



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b>  <b>C12Q 1/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/53101</b>  <b>(43) International Publication Date:</b> 21 October 1999 (21.10.99)
<b>(21) International Application Number:</b> PCT/US99/08268  <b>(22) International Filing Date:</b> 13 April 1999 (13.04.99)  <b>(30) Priority Data:</b> 60/081,483      13 April 1998 (13.04.98)      US 09/067,638      28 April 1998 (28.04.98)      US  <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US      09/067,638 (CIP) Filed on      28 April 1998 (28.04.98)  <b>(71) Applicant (for all designated States except US):</b> ISIS PHARMACEUTICALS, INC. [US/US]; 2292 Faraday Avenue, Carlsbad, CA 92008 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> COWSERT, Lex, M. [US/US]; 3008 Newshire Street, Carlsbad, CA 92008 (US). BAKER, Brenda, F. [US/US]; 2395-J Caringa Way, Carlsbad, CA 92009 (US). MCNEIL, John [US/US]; 427 Retaheim Way, La Jolla, CA 92037 (US). FREIER, Susan, M. [US/US]; 2946 Renault Street, San Diego, CA 92122 (US). SASMOR, Henri, M. [US/US]; 1751 Orange Blossom		Way, Encinitas, CA 92024 (US). BROOKS, Douglas, G. [US/US]; 1156 Caminito Amarillo, San Marcos, CA 92069 (US). OHASI, Cara [CA/US]; 2174 Cambridge Avenue, Cardiff, CA 92007 (US). WYATT, Jacqueline, R. [US/US]; 1065 Hymettus Avenue, Encinitas, CA 92024 (US). BORCHERS, Alexander, H. [US/US]; 733 Winding Way, Encinitas, CA 92024 (US). VICKERS, Timothy, A. [US/US]; 253 Luiseno Avenue, Oceanside, CA 92057 (US).  <b>(74) Agents:</b> CALDWELL, John, W. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).  <b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> IDENTIFICATION OF GENETIC TARGETS FOR MODULATION BY OLIGONUCLEOTIDES AND GENERATION OF OLIGONUCLEOTIDES FOR GENE MODULATION  <b>(57) Abstract</b>  Iterative, preferably computer based iterative processes for generating synthetic compounds with desired physical, chemical and/or bioactive properties, i.e., active compounds, are provided. During iterations of the processes, a target nucleic acid sequence is provided or selected, and a library of candidate nucleobase sequences is generated <i>in silico</i> according to defined criteria. A "virtual" oligonucleotide chemistry is chosen and a library of virtual oligonucleotide compounds having the selected nucleobase sequences is generated. These virtual compounds are reviewed and compounds predicted to have particular properties are selected. The selected compounds are robotically synthesized and are preferably robotically assayed for a desired physical, chemical or biological activity. Active compounds are thus generated and, at the same time, preferred sequences and regions of the target nucleic acid that are amenable to oligonucleotide or sequence-based modulation are identified.		

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**IDENTIFICATION OF GENETIC TARGETS FOR MODULATION  
BY OLIGONUCLEOTIDES AND GENERATION OF  
OLIGONUCLEOTIDES FOR GENE MODULATION**

**CROSS REFERENCE TO RELATED APPLICATIONS**

5           The present application is a continuation-in-part of U.S. Serial No. 09/067,638 filed April 28, 1998, which claims priority to provisional application Serial No. 60/081,483 filed April 13, 1998, each of which is incorporated herein by reference in its entirety.

**FIELD OF THE INVENTION**

10           The present invention relates generally to the generation and identification of synthetic compounds having defined physical, chemical or bioactive properties. More particularly, the present invention relates to the automated generation of oligonucleotide compounds targeted to a given nucleic acid sequence via computer-based, iterative robotic synthesis of synthetic oligonucleotide compounds and robotic or robot-assisted analysis of  
15           the activities of such compounds. Information gathered from assays of such compounds is used to identify nucleic acid sequences that are tractable to a variety of nucleotide sequence-based technologies, for example, antisense drug discovery and target validation.

## BACKGROUND OF THE INVENTION

### 1. Oligonucleotide Technology

Synthetic oligonucleotides of complementarity to targets are known to hybridize with particular, target nucleic acids in a sequence-specific manner. In one example, compounds complementary to the "sense" strand of nucleic acids that encode polypeptides, are referred to as "antisense oligonucleotides." A subset of such compounds may be capable of modulating the expression of a target nucleic acid; such synthetic compounds are described herein as "active oligonucleotide compounds."

Oligonucleotide compounds are commonly used *in vitro* as research reagents and diagnostic aids, and *in vivo* as therapeutic and bioactive agents. Oligonucleotide compounds can exert their effect by a variety of means. One such means takes advantage of an endogenous nuclease, such as RNase H in eukaryotes or RNase P in prokaryotes, to degrade the DNA/RNA hybrid formed between the oligonucleotide sequence and mRNA (Chiang *et al.*, *J. Biol. Chem.*, 1991, 266, 18162; Forster *et al.*, *Science*, 1990, 249, 783). Another means involves covalently linking of a synthetic moiety having nuclease activity to an oligonucleotide having an antisense sequence. This does not rely upon recruitment of an endogenous nuclease to modulate target activity. Synthetic moieties having nuclease activity include, but are not limited to, enzymatic RNAs, lanthanide ion complexes, and other reactive species. (Haseloff *et al.*, *Nature*, 1988, 334, 585; Baker *et al.*, *J. Am. Chem. Soc.*, 1997, 119, 8749).

Despite the advances made in utilizing antisense technology to date, it is still common to identify target sequences amenable to antisense technologies through an empirical approach (Szoka, *Nature Biotechnology*, 1997, 15, 509). Accordingly, the need exists for systems and methods for efficiently and effectively identifying target nucleotide sequences that are suitable for antisense modulation. The present disclosure answers this need by providing systems and methods for automatically identifying such sequences via *in silico*, robotic or other automated means.

### 2. Identification of Active Oligonucleotide Compounds

Traditionally, new chemical entities with useful properties are generated by (1) identifying a chemical compound (called a "lead compound") with some desirable property or activity, (2) creating variants of the lead compound, and (3) evaluating the



property and activity of such variant compounds. The process has been called "SAR," i.e., structure activity relationship. Although "SAR" and its handmaiden, rational drug design, has been utilized with some degree of success, there are a number of limitations to these approaches to lead compound generation, particularly as it pertains to the discovery of bioactive oligonucleotide compounds. In attempting to use SAR with oligonucleotides, it has been recognized that RNA structure can inhibit duplex formation with antisense compounds, so much so that "moving" the target nucleotide sequence even a few bases can drastically decrease the activity of such compounds (Lima *et al.*, *Biochemistry*, 1992, 31, 12055).

Heretofore, the preferred method of searching for lead antisense compounds has been the manual synthesis and analysis of such compounds. Consequently, a fundamental limitation of the conventional approach is its dependence upon the availability, number and cost of antisense compounds produced by manual, or at best semi-automated, means. Moreover, the assaying of such compounds has traditionally been performed by tedious manual techniques. Thus, the traditional approach to generating active antisense compounds is limited by the relatively high cost and long time required to synthesize and screen a relatively small number of candidate antisense compounds.

Accordingly, the need exists for systems and methods for efficiently and effectively generating new active antisense and other oligonucleotide compounds targeted to specific nucleic acid sequences. The present disclosure answers this need by providing systems and methods for automatically generating and screening active antisense compounds via robotic and other automated means.

### 3. Gene Function Analysis

Efforts such as the Human Genome Project are making an enormous amount of nucleotide sequence information available in a variety of forms, e.g., genomic sequences, cDNAs, expressed sequence tags (ESTs) and the like. This explosion of information has led one commentator to state that "genome scientists are producing more genes than they can put a function to" (Kahn, *Science*, 1995, 270, 369). Although some approaches to this problem have been suggested, no solution has yet emerged. For example, methods of looking at gene expression in different disease states or stages of development only provide, at best, an association between a gene and a disease or stage of development

(Nowak, *Science*, 1995, 270, 368). Another approach, looking at the proteins encoded by genes, is developing but “this approach is more complex and big obstacles remain” (Kahn, *Science*, 1995, 270, 369). Furthermore, neither of these approaches allows one to directly utilize nucleotide sequence information to perform gene function analysis.

5           In contrast, antisense technology does allow for the direct utilization of nucleotide sequence information for gene function analysis. Once a target nucleic acid sequence has been selected, antisense sequences hybridizable to the sequence can be generated using techniques known in the art. Typically, a large number of candidate antisense oligonucleotides (ASOs) are synthesized having sequences that are more-or-less randomly  
10           spaced across the length of the target nucleic acid sequence (e.g., a “gene walk”) and their ability to modulate the expression of the target nucleic acid is assayed. Cells or animals can then be treated with one or more active antisense oligonucleotides, and the resulting effects determined in order to determine the function(s) of the target gene. Although the practicality and value of this empirical approach to determining gene function has been  
15           acknowledged in the art, it has also been stated that this approach “is beyond the means of most laboratories and is not feasible when a new gene sequence is identified, but whose function and therapeutic potential are unknown” (Szoka, *Nature Biotechnology*, 1997, 15, 509).

          Accordingly, the need exists for systems and methods for efficiently and  
20           effectively determining the function of a gene that is uncharacterized except that its nucleotide sequence, or a portion thereof, is known. The present disclosure answers this need by providing systems and methods for automatically generating active antisense compounds to a target nucleotide sequence via robotic means. Such active antisense compounds are contacted with cells, cell-free extracts, tissues or animals capable of  
25           expressing the gene of interest and subsequent biochemical or biological parameters are measured. The results are compared to those obtained from a control cell culture, cell-free extract, tissue or animal which has not been contacted with an active antisense compound in order to determine the function of the gene of interest.

#### 4. Target Validation

30           Determining the nucleotide sequence of a gene is no longer an end unto itself; rather, it is “merely a means to an end. The critical next step is to validate the gene and its

[gene] product as a potential drug target” (Glasser, *Genetic Engineering News*, 1997, 17, 1). This process, i.e., confirming that modulation of a gene that is suspected of being involved in a disease or disorder actually results in an effect that is consistent with a causal relationship between the gene and the disease or disorder, is known as target validation.

5           Efforts such as the Human Genome Project are yielding a vast number of complete or partial nucleotide sequences, many of which might correspond to or encode targets useful for new drug discovery efforts. The challenge represented by this plethora of information is how to use such nucleotide sequences to identify and rank valid targets for drug discovery. Antisense technology provides one means by which this might be  
10           accomplished; however, the many manual, labor-intensive and costly steps involved in traditional methods of developing active antisense compounds has limited their use in target validation (Szoka, *Nature Biotechnology*, 1997, 15, 509). Nevertheless, the great target specificity that is characteristic of antisense compounds makes them ideal choices for target validation, especially when the functional roles of proteins that are highly related  
15           are being investigated (Albert *et al.*, *Trends in Pharm. Sci.*, 1994, 15, 250).

          Accordingly, the need exists for systems and methods for developing compounds efficiently and effectively that modulate a gene, wherein such compounds can be directly developed from nucleotide sequence information. Such compounds are needed to confirm that modulation of a gene that is thought to be involved in a disease or disorder will in fact  
20           cause an *in vitro* or *in vivo* effect indicative of the origin, development, spread or growth of the disease or disorder.

          The present disclosure answers this need by providing systems and methods for automatically generating active oligonucleotide and other compounds, especially antisense compounds, to a target nucleotide sequence via robotic or other automated means. Such  
25           active compounds are contacted with a cell culture, cell-free extract, tissue or animal capable of expressing the gene of interest, and subsequent biochemical or biological parameters indicative of the potential gene product function are measured. These results are compared to those obtained with a control cell system, cell-free extract, tissue or animal which has not been contacted with an active antisense compound in order to  
30           determine whether or not modulation of the gene of interest affects a specific cellular function. The resulting active antisense compounds may be used as positive controls when

other, non antisense-based agents directed to the same target nucleic acid, or to its gene product, are screened.

It should be noted that embodiments of the invention drawn to gene function analysis and target validation have parameters that are shared with other embodiments of the invention, but also have unique parameters. For example, antisense drug discovery naturally requires that the toxicity of the antisense compounds be manageable, whereas, for gene function analysis or target validation, overt toxicity resulting from the antisense compounds is acceptable unless it interferes with the assay being used to evaluate the effects of treatment with such compounds.

U.S. Patent 5,563,036 to Peterson *et al.* describes systems and methods of screening for compounds that inhibit the binding of a transcription factor to a nucleic acid. In a preferred embodiment, an assay portion of the process is stated to be performed by a computer controlled robot.

U.S. Patent 5,708,158 to Hoey describes systems and methods for identifying pharmacological agents stated to be useful for diagnosing or treating a disease associated with a gene the expression of which is modulated by a human nuclear factor of activated T cells. The methods are stated to be particularly suited to high-throughput screening wherein one or more steps of the process are performed by a computer controlled robot.

U.S. Patents 5,693,463 and 5,716,780 to Edwards *et al.* describe systems and methods for identifying non-oligonucleotide molecules that specifically bind to a DNA molecule based on their ability to compete with a DNA-binding protein that recognizes the DNA molecule.

U.S. Patents 5,463,564 and 5,684,711 to Agrafiotis *et al.* describe computer based iterative processes for generating chemical entities with defined physical, chemical and/or bioactive properties.

## SUMMARY OF THE INVENTION

The present invention is directed to automated systems and methods for defining sets of compounds that modulate the expression of target nucleic acid sequences, and generating sets of oligonucleotides that modulate the expression of target nucleic acid sequences. The present invention is also directed to identifying nucleic acid sequences

amenable to antisense binding of oligonucleotides to those nucleic acid sequences by the systems and methods of the invention. For purposes of illustration, the present invention is described herein with respect to the production and identification of active antisense oligonucleotides; however, the present invention is not limited to this embodiment.

5           The present invention is directed to iterative processes for defining chemical compounds with prescribed sets of physical, chemical and/or biological properties, and to systems for implementing these processes. During each iteration of a process as contemplated herein, a target nucleic acid sequence is provided or selected, and a library of (candidate) virtual compounds is generated *in silico* (that is in a computer manipulatable  
10           and reliable form) according to defined criteria. A library of virtual compounds is generated. These virtual compounds are reviewed and compounds predicted to have particular desired properties are selected. The selected compounds are synthesized, preferably in a robotic, batchwise manner; and then they are robotically assayed for a desired physical, chemical or biological activity in order to identify compounds with the  
15           desired properties. Active compounds are, thus, generated and, at the same time, preferred sequences and regions of the target nucleic acid that are amenable to modulation are identified. The preferred compounds of the invention are oligonucleotides that bind to a target nucleic acid sequence.

          In subsequent iterations of the process, second libraries of candidate compounds  
20           are generated and/or selected to give rise to a second virtual compound library. Through multiple iterations of the process, a library of target nucleic acid sequences that are tractable to modulation via binding of these compounds to the nucleic acid sequence are identified. Such modulation includes, but is not limited to, antisense technology, gene function analysis and target validation.

25           The present invention is also directed to processes for validating the function of a gene or the product of the gene comprising generating *in silico* a library of nucleobase sequences targeted to the gene and robotically assaying a plurality of synthetic compounds having at least some of the nucleobase sequences for effects on biological function.

          Further features and advantages of the present invention, as well as the structure  
30           and operation of various embodiments of the present invention, are described in detail below with reference to the accompanying drawings. In the drawings, like reference

numbers indicate identical or functionally similar elements.

## BRIEF DESCRIPTION OF THE DRAWINGS

The present invention will be described with reference to the accompanying drawings, wherein:

5            Figures 1 and 2 are a flow diagram of one method according to the present invention depicting the overall flow of data and materials among various elements of the invention.

            Figure 3 is a flow diagram depicting the flow of data and materials among elements of step **200** of Figure 1.

10           Figures 4 and 5 are a flow diagram depicting the flow of data and materials among elements of step **300** of Figure 1.

            Figure 6 is a flow diagram depicting the flow of data and materials among elements of step **306** of Figure 4.

15           Figure 7 is another flow diagram depicting the flow of data and materials among elements of step **306** of Figure 4.

            Figure 8 is a another flow diagram depicting the flow of data and materials among elements of step **306** of Figure 4.

            Figure 9 is a flow diagram depicting the flow of data and materials among elements of step **350** of Figure 5.

20           Figures 10 and 11 are flow diagrams depicting a logical analysis of data and materials among elements of step **400** of Figure 1.

            Figure 12 is a flow diagram depicting the flow of data and materials among the elements of step **400** of Figure 1.

25           Figures 13 and 14 are flow diagrams depicting the flow of data and materials among elements of step **500** of Figure 1.

            Figure 15 is a flow diagram depicting the flow of data and materials among elements of step **600** of Figure 1.

            Figure 16 is a flow diagram depicting the flow of data and materials among elements of step **700** of Figure 1.

30           Figure 17 is a flow diagram depicting the flow of data and materials among the

elements of step **1100** of Figure 2.

Figure 18 is a block diagram showing the interconnecting of certain devices utilized in conjunction with a preferred method of the invention;

Figure 19 is a flow diagram showing a representation of data storage in a relational database utilized in conjunction with one method of the invention;

Figure 20 is a flow diagram depicting the flow of data and materials in effecting a preferred embodiment of the invention as set forth in Example 14;

Figure 21 is a flow diagram depicting the flow of data and materials in effecting a preferred embodiment of the invention as set forth in Example 15;

Figure 22 is a flow diagram depicting the flow of data and materials in effecting a preferred embodiment of the invention as set forth in Example 2;

Figure 23 is a pictorial elevation view of a preferred apparatus used to robotically synthesize oligonucleotides; and

Figure 24 is a pictorial plan view of an apparatus used to robotically synthesize oligonucleotides.

## DETAILED DESCRIPTION OF THE INVENTION

Certain preferred methods of this invention are now described with reference to the flow diagram of Figures 1 and 2.

**1. Target Nucleic Acid Selection.** The target selection process, process step **100**, provides a target nucleotide sequence that is used to help guide subsequent steps of the process. It is generally desired to modulate the expression of the target nucleic acid for any of a variety of purposes, such as, e.g., drug discovery, target validation and/or gene function analysis.

One of the primary objectives of the target selection process, step **100**, is to identify molecular targets that represent significant therapeutic opportunities, to provide new and efficacious means of drug discovery and to determine the function of genes that are uncharacterized except for nucleotide sequence. To meet these objectives, genes are classified based upon specific sets of selection criteria.

One such set of selection criteria concerns the quantity and quality of target nucleotide sequence. There must be sufficient target nucleic acid sequence information

available for oligonucleotide design. Moreover, such information must be of sufficient quality to give rise to an acceptable level of confidence in the data to perform the methods described herein. Thus, the data must not contain too many missing or incorrect base entries. In the case of a target sequence that encodes a polypeptide, such errors can often  
5 be detected by virtually translating all three reading frames of the sense strand of the target sequence and confirming the presence of a continuous polypeptide sequence having predictable attributes, e.g., encoding a polypeptide of known size, or encoding a polypeptide that is about the same length as a homologous protein. In any event, only a very high frequency of sequence errors will frustrate the methods of the invention; most  
10 oligonucleotides to the target sequence will avoid such errors unless such errors occur frequently throughout the entire target sequence.

Another preferred criterion is that appropriate culturable cell lines or other source of reproducible genetic expression should be available. Such cell lines express, or can be induced to express, the gene comprising the target nucleic acid sequence. The  
15 oligonucleotide compounds generated by the process of the invention are assayed using such cell lines and, if such assaying is performed robotically, the cell line is preferably tractable to robotic manipulation such as by growth in 96 well plates. Those skilled in the art will recognize that if an appropriate cell line does not exist, it will nevertheless be possible to construct an appropriate cell line. For example, a cell line can be transfected  
20 with an expression vector comprising the target gene in order to generate an appropriate cell line for assay purposes.

For gene function analysis, it is possible to operate upon a genetic system having a lack of information regarding, or incomplete characterization of, the biological function(s) of the target nucleic acid or its gene product(s). This is a powerful agent of the invention.  
25 A target nucleic acid for gene function analysis might be absolutely uncharacterized, or might be thought to have a function based on minimal data or homology to another gene. By application of the process of the invention to such a target, active compounds that modulate the expression of the gene can be developed and applied to cells. The resulting cellular, biochemical or molecular biological responses are observed, and this information  
30 is used by those skilled in the art to elucidate the function of the target gene.

For target validation and drug discovery, another selection criterion is disease



association. Candidate target genes are placed into one of several broad categories of known or deduced disease association. Level 1 Targets are target nucleic acids for which there is a strong correlation with disease. This correlation can come from multiple scientific disciplines including, but not limited to, epidemiology, wherein frequencies of gene abnormalities are associated with disease incidence; molecular biology, wherein gene expression and function are associated with cellular events correlated with a disease; and biochemistry, wherein the *in vitro* activities of a gene product are associated with disease parameters. Because there is a strong therapeutic rationale for focusing on Level 1 Targets, these targets are most preferred for drug discovery and/or target validation.

Level 2 Targets are nucleic acid targets for which the combined epidemiological, molecular biological, and/or biochemical correlation with disease is not so clear as for Level 1. Level 3 Targets are targets for which there is little or no data to directly link the target with a disease process, but there is indirect evidence for such a link, i.e., homology with a Level 1 or Level 2 target nucleic acid sequence or with the gene product thereof. In order not to prejudice the target selection process, and to ensure that the maximum number of nucleic acids actually involved in the causation, potentiation, aggravation, spread, continuance or after-effects of disease states are investigated, it is preferred to examine a balanced mix of Level 1, 2 and 3 target nucleic acids.

In order to carry out drug discovery, experimental systems and reagents shall be available in order for one to evaluate the therapeutic potential of active compounds generated by the process of the invention. Such systems may be operable *in vitro* (e.g., *in vitro* models of cell:cell association) or *in vivo* (e.g., animal models of disease states). It is also desirable, but not obligatory, to have available animal model systems which can be used to evaluate drug pharmacology.

Candidate targets nucleic acids can also classified by biological processes. For example, programmed cell death ("apoptosis") has recently emerged as an important biological process that is perturbed in a wide variety of diseases. Accordingly, nucleic acids that encode factors that play a role in the apoptotic process are identified as candidate targets. Similarly, potential target nucleic acids can be classified as being involved in inflammation, autoimmune disorders, cancer, or other pathological or dysfunctional processes.

Moreover, genes can often be grouped into families based on sequence homology and biological function. Individual family members can act redundantly, or can provide specificity through diversity of interactions with downstream effectors, or through expression being restricted to specific cell types. When one member of a gene family is associated with a disease process then the rationale for targeting other members of the same family is reasonably strong. Therefore, members of such gene families are preferred target nucleic acids to which the methods and systems of the invention may be applied. Indeed, the potent specificity of antisense compounds for different gene family members makes the invention particularly suited for such targets (Albert *et al.*, *Trends Pharm. Sci.*, 1994, 15, 250). Those skilled in the art will recognize that a partial or complete nucleotide sequence of such family members can be obtained using the polymerase chain reaction (PCR) and “universal” primers, i.e., primers designed to be common to all members of a given gene family.

PCR products generated from universal primers can be cloned and sequenced or directly sequenced using techniques known in the art. Thus, although nucleotide sequences from cloned DNAs, or from complementary DNAs (cDNAs) derived from mRNAs, may be used in the process of the invention, there is no requirement that the target nucleotide sequence be isolated from a cloned nucleic acid. Any nucleotide sequence, no matter how determined, of any nucleic acid, isolated or prepared in any fashion, may be used as a target nucleic acid in the process of the invention.

Furthermore, although polypeptide-encoding nucleic acids provide the target nucleotide sequences in one embodiment of the invention, other nucleic acids may be targeted as well. Thus, for example, the nucleotide sequences of structural or enzymatic RNAs may be utilized for drug discovery and/or target validation when such RNAs are associated with a disease state, or for gene function analysis when their biological role is not known.

**2. Assembly of Target Nucleotide Sequence.** Figure 3 is a block diagram detailing the steps of the target nucleotide sequence assembly process, process step **200** in accordance with one embodiment of the invention. The oligonucleotide design process, process step **300**, is facilitated by the availability of accurate target sequence information. Because of limitations of automated genome sequencing technology, gene sequences are

often accumulated in fragments. Further, because individual genes are often being sequenced by independent laboratories using different sequencing strategies, sequence information corresponding to different fragments is often deposited in different databases. The target nucleic acid assembly process take advantage of computerized homology search algorithms and sequence fragment assembly algorithms to search available databases for related sequence information and incorporate available sequence information into the best possible representation of the target nucleic acid molecule, for example a RNA transcript. This representation is then used to design oligonucleotides, process step **300**, which can be tested for biological activity in process step **700**.

In the case of genes directing the synthesis of multiple transcripts, i.e. by alternative splicing, each distinct transcript is a unique target nucleic acid for purposes of step **300**. In one embodiment of the invention, if active compounds specific for a given transcript isoform are desired, the target nucleotide sequence is limited to those sequences that are unique to that transcript isoform. In another embodiment of the invention, if it is desired to modulate two or more transcript isoforms in concert, the target nucleotide sequence is limited to sequences that are shared between the two or more transcripts.

In the case of a polypeptide-encoding nucleic acid, it is generally preferred that full-length cDNA be used in the oligonucleotide design process step **300** (with full-length cDNA being defined as reading from the 5' cap to the poly A tail). Although full-length cDNA is preferred, it is possible to design oligonucleotides using partial sequence information. Therefore it is not necessary for the assembly process to generate a complete cDNA sequence. Further in some cases it may be desirable to design oligonucleotides targeting introns. In this case the process can be used to identify individual introns at process step **220**.

The process can be initiated by entering initial sequence information on a selected molecular target at process step **205**. In the case of a polypeptide-encoding nucleic acid, the full-length cDNA sequence is generally preferred for use in oligonucleotide design strategies at process step **300**. The first step is to determine if the initial sequence information represents the full-length cDNA, decision step **210**. In the case where the full-length cDNA sequence is available the process advances directly to the oligonucleotide design step **300**. When the full-length cDNA sequence is not available, databases are

searched at process step **212** for additional sequence information.

The algorithm preferably used in process steps **212** and **230** is BLAST (Altschul, *et al.*, *J. Mol. Biol.*, **1990**, 215, 403), or “Gapped BLAST” (Altschul *et al.*, *Nucl. Acids Res.*, 1997, 25, 3389). These are database search tools based on sequence homology used to identify related sequences in a sequence database. The BLAST search parameters are set to only identify closely related sequences. Some preferred databases searched by BLAST are a combination of public domain and proprietary databases. The databases, their contents, and sources are listed in Table 1.

**Table 1: Database Sources of Target Sequences**

	Database	Contents	Source
10	NR	All non-redundant GenBank, EMBL, DDBJ and PDB sequences	National Center for Biotechnology Information at the National Institutes of Health
	Month	All new or revised GenBank, EMBL, DDBJ and PDB sequences released in the last 30 days	National Center for Biotechnology Information at the National Institutes of Health
	Dbest	Non-redundant database of GenBank, EMBL, DDBJ and EST divisions	National Center for Biotechnology Information at the National Institutes of Health
	Dbsts	Non-redundant database of GenBank, EMBL, DDBJ and STS divisions	National Center for Biotechnology Information at the National Institutes of Health
15	Htgs	High throughput genomic sequences	National Center for Biotechnology Information at the National Institutes of Health

When genomic sequence information is available at decision step **215**, introns are removed and exons are assembled into continuous sequence representing the cDNA sequence in process step **220**. Exon assembly occurs using the Phragment Assembly Program “Phrap” (Copyright University of Washington Genome Center, Seattle, WA). The Phrap algorithm analyzes sets of overlapping sequences and assembles them into one

continuous sequence referred to as a "contig." The resulting contig is preferably used to search databases for additional sequence information at process step 230. When genomic information is not available, the results of process step 212 are analyzed for individual exons at decision step 225. Exons are frequently recorded individually in databases. If multiple complete exons are identified, they are preferably assembled into a contig using Phrap at process step 250. If multiple complete exons are not identified at decision step 225, then sequences can be analyzed for partial sequence information in decision step 228. ESTs identified in the database dbEST are examples of such partial sequence information. If additional partial information is not found, then the process is advanced to process step 230 at decision step 228. If partial sequence information is found in process 212 then that information is advanced to process step 230 via decision step 228.

Process step 230, decision step 240, decision step 260 and process step 250 define a loop designed to extend iteratively the amount of sequence information available for targeting. At the end of each iteration of this loop, the results are analyzed in decision steps 240 and 260. If no new information is found then the process advances at decision step 240 to process step 300. If there is an unexpectedly large amount of sequence information identified, suggesting that the process moved outside the boundary of the gene into repetitive genomic sequence, then the process is preferably cycled back one iteration and that sequence is advanced at decision step 240 to process step 300. If a small amount of new sequence information is identified, then the loop is iterated such as by taking the 100 most 5-prime (5') and 100 most 3-prime (3') bases and iterating them through the BLAST homology search at process step 230. New sequence information is added to the existing contig at process step 250.

### **3. *In Silico* Generation of a Set of Nucleobase Sequences and Virtual Oligonucleotides.**

For the following steps 300 and 400, they may be performed in the order described below, i.e., step 300 before step 400, or, in an alternative embodiment of the invention, step 400 before step 300. In this alternate embodiment, each oligonucleotide chemistry is first assigned to each oligonucleotide sequence. Then, each combination of oligonucleotide chemistry and sequence is evaluated according to the parameters of step 300. This embodiment has the desirable feature of taking into account the effect of

alternative oligonucleotide chemistries on such parameters. For example, substitution of 5-methyl cytosine (5MeC or m5c) for cytosine in an antisense compound may enhance the stability of a duplex formed between that compound and its target nucleic acid. Other oligonucleotide chemistries that enhance oligonucleotide:[target nucleic acid] duplexes are known in the art (see for example, Freier *et al.*, *Nucleic Acids Research*, 1997, 25, 4429). As will be appreciated by those skilled in the art, different oligonucleotide chemistries may be preferred for different target nucleic acids. That is, the optimal oligonucleotide chemistry for binding to a target DNA might be suboptimal for binding to a target RNA having the same nucleotide sequence.

In effecting the process of the invention in the order step 300 before step 400 as seen in Figure 1, from a target nucleic acid sequence assembled at step 200, a list of oligonucleotide sequences is generated as represented in the flowchart shown in Figures 4 and 5. In step 302, the desired oligonucleotide length is chosen. In a preferred embodiment, oligonucleotide length is between from about 8 to about 30, more preferably from about 12 to about 25, nucleotides. In step 304, all possible oligonucleotide sequences of the desired length capable of hybridizing to the target sequence obtained in step 200 are generated. In this step, a series of oligonucleotide sequences are generated, simply by determining the most 5' oligonucleotide possible and "walking" the target sequence in increments of one base until the 3' most oligonucleotide possible is reached.

In step 305, a virtual oligonucleotide chemistry is applied to the nucleobase sequences of step 304 in order to yield a set of virtual oligonucleotides that can be evaluated *in silico*. Default virtual oligonucleotide chemistries include those that are well-characterized in terms of their physical and chemical properties, e.g., 2'-deoxyribonucleic acid having naturally occurring bases (A, T, C and G), unmodified sugar residues and a phosphodiester backbone.

#### **4. *In Silico* Evaluation of Thermodynamic Properties of Virtual Oligonucleotides.**

In step 306, a series of thermodynamic, sequence, and homology scores are preferably calculated for each virtual oligonucleotide obtained from step 305.

Thermodynamic properties are calculated as represented in Figure 6. In step 308, the desired thermodynamic properties are selected. As many or as few as desired can be

selected; optionally, none will be selected. The desired properties will typically include step 309, calculation of the free energy of the target structure. If the oligonucleotide is a DNA molecule, then steps 310, 312, and 314 are performed. If the oligonucleotide is an RNA molecule, then steps 311, 313 and 315 are performed. In both cases, these steps  
5 correspond to calculation of the free energy of intramolecular oligonucleotide interactions, intermolecular interactions and duplex formation. In addition, a free energy of oligonucleotide-target binding is preferably calculated at step 316.

Other thermodynamic and kinetic properties may be calculated for oligonucleotides as represented at step 317. Such other thermodynamic and kinetic  
10 properties may include melting temperatures, association rates, dissociation rates, or any other physical property that may be predictive of oligonucleotide activity.

The free energy of the target structure is defined as the free energy needed to disrupt any secondary structure in the target binding site of the targeted nucleic acid. This region includes any intra-target nucleotide base pairs that need to be disrupted in order for  
15 an oligonucleotide to bind to its complementary sequence. The effect of this localized disruption of secondary structure is to provide accessibility by the oligonucleotide. Such structures will include double helices, terminal unpaired and mismatched nucleotides, loops, including hairpin loops, bulge loops, internal loops and multibranch loops (Serra *et al.*, *Methods in Enzymology*, 1995, 259, 242).

20 The intermolecular free energies refer to inherent energy due to the most stable structure formed by two oligonucleotides; such structures include dimer formation. Intermolecular free energies should also be taken into account when, for example, two or more oligonucleotides, of different sequence are to be administered to the same cell in an assay.

25 The intramolecular free energies refer to the energy needed to disrupt the most stable secondary structure within a single oligonucleotide. Such structures include, for example, hairpin loops, bulges and internal loops. The degree of intramolecular base pairing is indicative of the energy needed to disrupt such base pairing.

The free energy of duplex formation is the free energy of denatured  
30 oligonucleotide binding to its denatured target sequence. The oligonucleotide-target binding is the total binding involved, and includes the energies involved in opening up

intra- and inter- molecular oligonucleotide structures, opening up target structure, and duplex formation.

The most stable RNA structure is predicted based on nearest neighbor analysis (Xia, T., *et al.*, *Biochemistry*, 1998, 37, 14719-14735; Serra *et al.*, *Methods in Enzymology*, 1995, 259, 242). This analysis is based on the assumption that stability of a given base pair is determined by the adjacent base pairs. For each possible nearest neighbor combination, thermodynamic properties have been determined and are provided. For double helical regions, two additional factors need to be considered, an entropy change required to initiate a helix and a entropy change associated with self-complementary strands only. Thus, the free energy of a duplex can be calculated using the equation:

$$\Delta G^{\circ}_T = \Delta H^{\circ} - T\Delta S^{\circ}$$

where:

$\Delta G$  is the free energy of duplex formation,

$\Delta H$  is the enthalpy change for each nearest neighbor,

$\Delta S$  is the entropy change for each nearest neighbor, and  $T$  is temperature.

The  $\Delta H$  and  $\Delta S$  for each possible nearest neighbor combination have been experimentally determined. These letter values are often available in published tables. For terminal unpaired and mismatched nucleotides, enthalpy and entropy measurements for each possible nucleotide combination are also available in published tables. Such results are added directly to values determined for duplex formation. For loops, while the available data is not as complete or accurate as for base pairing, one known model determines the free energy of loop formation as the sum of free energy based on loop size, the closing base pair, the interactions between the first mismatch of the loop with the closing base pair, and additional factors including being closed by AU or UA or a first mismatch of GA or UU. Such equations may also be used for oligoribonucleotide-target RNA interactions.

The stability of DNA duplexes is used in the case of intra- or intermolecular oligodeoxyribonucleotide interactions. DNA duplex stability is calculated using similar equations as RNA stability, except experimentally determined values differ between nearest neighbors in DNA and RNA and helix initiation tends to be more favorable in



DNA than in RNA (SantaLucia *et al.*, *Biochemistry*, 1996, 35, 3555).

Additional thermodynamic parameters are used in the case of RNA/DNA hybrid duplexes. This would be the case for an RNA target and oligodeoxynucleotide. Such parameters were determined by Sugimoto *et al.* (*Biochemistry*, 1995, 34, 11211). In addition to values for nearest neighbors, differences were seen for values for enthalpy of helix initiation.

### 5. *In Silico* Evaluation of Target Accessibility

Target accessibility is believed to be an important consideration in selecting oligonucleotides. Such a target site will possess minimal secondary structure and thus, will require minimal energy to disrupt such structure. In addition, secondary structure in oligonucleotides, whether inter- or intra-molecular, is undesirable due to the energy required to disrupt such structures. Oligonucleotide-target binding is dependent on both these factors. It is desirable to minimize the contributions of secondary structure based on these factors. The other contribution to oligonucleotide-target binding is binding affinity. Favorable binding affinities based on tighter base pairing at the target site is desirable.

Following the calculation of thermodynamic properties ending at step 317, the desired sequence properties to be scored are selected at step 324. As many or as few as desired can be selected; optionally, none will be selected. These properties include the number of strings of four guanosine residues in a row at step 325 or three guanosine in a row at step 326, the length of the longest string of adenosines at step 327, cytidines at step 328 or uridines or thymidines at step 329, the length of the longest string of purines at step 330 or pyrimidine at step 331, the percent composition of adenosine at step 332, cytidine at step 333, guanosine at step 334 or uridines or thymidines at step 335, the percent composition of purines at step 336 or pyrimidines at step 337, the number of CG dinucleotide repeats at step 338, CA dinucleotide repeats at step 339 or UA or TA dinucleotide repeats at step 340. In addition, other sequence properties may be used as found to be relevant and predictive of antisense efficacy, as represented at step 341.

These sequence properties may be important in predicting oligonucleotide activity, or lack thereof. For example, U.S. Patent 5,523,389 discloses oligonucleotides containing stretches of three or four guanosine residues in a row. Oligonucleotides having such sequences may act in a sequence-independent manner. For an antisense approach, such a

mechanism is not usually desired. In addition, high numbers of dinucleotide repeats may be indicative of low complexity regions which may be present in large numbers of unrelated genes. Unequal base composition, for example, 90% adenosine, can also give non-specific effects. From a practical standpoint, it may be desirable to remove oligonucleotides that possess long stretches of other nucleotides due to synthesis considerations. Other sequences properties, either listed above or later found to be of predictive value may be used to select oligonucleotide sequences.

Following step 341, the homology scores to be calculated are selected in step 342. Homology to nucleic acids encoding protein isoforms of the target, as represented at step 343, may be desired. For example, oligonucleotides specific for an isoform of protein kinase C can be selected. Also, oligonucleotides can be selected to target multiple isoforms of such genes. Homology to analogous target sequences, as represented at step 344, may also be desired. For example, an oligonucleotide can be selected to a region common to both humans and mice to facilitate testing of the oligonucleotide in both species. Homology to splice variants of the target nucleic acid, as represented at step 345, may be desired. In addition, it may be desirable to determine homology to other sequence variants as necessary, as represented in step 346.

Following step 346, from which scores were obtained in each selected parameter, a desired range is selected to select the most promising oligonucleotides, as represented at step 347. Typically, only several parameters will be used to select oligonucleotide sequences. As structure prediction improves, additional parameters may be used. Once the desired score ranges are chosen, a list of all oligonucleotides having parameters falling within those ranges will be generated, as represented at step 348.

#### **6. Targeting Oligonucleotides to Functional Regions of a Nucleic Acid.**

It may be desirable to target oligonucleotide sequences to specific functional regions of the target nucleic acid. A decision is made whether to target such regions, as represented in decision step 349. If it is desired to target functional regions then process step 350 occurs as seen in greater detail in Figure 9. If it is not desired then the process proceeds to step 375.

In step 350, as seen in Figure 9, the desired functional regions are selected. Such regions include the transcription start site or 5' cap at step 353, the 5' untranslated region

at step 354, the start codon at step 355, the coding region at step 356, the stop codon at step 357, the 3' untranslated region at step 358, 5' splice sites at step 359 or 3' splice sites at step 360, specific exons at step 361 or specific introns at step 362, mRNA stabilization signal at step 363, mRNA destabilization signal at step 364, poly-adenylation signal at step 365, poly-A addition site at step 366, poly-A tail at step 367, or the gene sequence 5' of known pre-mRNA at step 368. In addition, additional functional sites may be selected, as represented at step 369.

Many functional regions are important to the proper processing of the gene and are attractive targets for antisense approaches. For example, the AUG start codon is commonly targeted because it is necessary to initiate translation. In addition, splice sites are thought to be attractive targets because these regions are important for processing of the mRNA. Other known sites may be more accessible because of interactions with protein factors or other regulatory molecules.

After the desired functional regions are selected and determined, then a subset of all previously selected oligonucleotides are selected based on hybridization to only those desired functional regions, as represented by step 370.

#### **7. Uniform Distribution of Oligonucleotides.**

Whether or not targeting functional sites is desired, a large number of oligonucleotide sequences may result from the process thus far. In order to reduce the number of oligonucleotide sequences to a manageable number, a decision is made whether to uniformly distribute selected oligonucleotides along the target, as represented in step 375. A uniform distribution of oligonucleotide sequences will aim to provide complete coverage throughout the complete target nucleic acid or the selected functional regions. A computer-based program is used to automate the distribution of sequences, as represented in step 380. Such a program factors in parameters such as length of the target nucleic acid, total number of oligonucleotide sequences desired, oligonucleotide sequences per unit length, number of oligonucleotide sequences per functional region. Manual selection of oligonucleotide sequences is also provided for by step 385. In some cases, it may be desirable to manually select oligonucleotide sequences. For example, it may be useful to determine the effect of small base shifts on activity. Once the desired number of oligonucleotide sequences is obtained either from step 380 or step 385, then these

oligonucleotide sequences are passed onto step 400 of the process, where oligonucleotide chemistries are assigned.

### 8. Assignment of Actual Oligonucleotide Chemistry.

Once a set of select nucleobase sequences has been generated according to the preceding process and decision steps, actual oligonucleotide chemistry is assigned to the sequences. An "actual oligonucleotide chemistry" or simply "chemistry" is a chemical motif that is common to a particular set of robotically synthesized oligonucleotide compounds. Preferred chemistries include, but are not limited to, oligonucleotides in which every linkage is a phosphorothioate linkage, and chimeric oligonucleotides in which a defined number of 5' and/or 3' terminal residues have a 2'-methoxyethoxy modification.

Chemistries can be assigned to the nucleobase sequences during general procedure step 400 (Figure 1). The logical basis for chemistry assignment is illustrated in Figures 10 and 11 and an iterative routine for stepping through an oligonucleotide nucleoside by nucleoside is illustrated in Figure 12. Chemistry assignment can be effected by assignment directly into a word processing program, via an interactive word processing program or via automated programs and devices. In each of these instances, the output file is selected to be in a format that can serve as an input file to automated synthesis devices.

### 9. Oligonucleotide Compounds.

In the context of this invention, in reference to oligonucleotides, the term "oligonucleotide" is used to refer to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. Thus this term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms, i.e., phosphodiester linked A, C, G, T and U nucleosides, because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

The oligonucleotide compounds in accordance with this invention can be of various lengths depending on various parameters, including but not limited to those discussed above in reference to the selection criteria of general procedure 300. For use as

antisense oligonucleotides compounds of the invention preferably are from about 8 to about 30 nucleobases in length (i.e. from about 8 to about 30 linked nucleosides).

Particularly preferred are antisense oligonucleotides comprising from about 12 to about 25 nucleobases. A discussion of antisense oligonucleotides and some desirable modifications can be found in De Mesmaeker *et al.*, *Acc. Chem. Res.*, 1995, 28, 366. Other lengths of oligonucleotides might be selected for non-antisense targeting strategies, for instance using the oligonucleotides as ribozymes. Such ribozymes normally require oligonucleotides of longer length as is known in the art.

A nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a normal (where normal is defined as being found in RNA and DNA) pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link adjacent nucleosides to one another to form a linear polymeric compound. In turn the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA and DNA is a 3' to 5' phosphodiester linkage.

Specific examples of preferred oligonucleotides useful in this invention include oligonucleotides containing modified backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

#### 10. Selection of Oligonucleotide Chemistries.

In a general logic scheme as illustrated in Figures 10 and 11, for each nucleoside position, the user or automated device is interrogated first for a base assignment, followed

by a sugar assignment, a linker assignment and finally a conjugate assignment. Thus for each nucleoside, at process step **410** a base is selected. In selecting the base, base chemistry 1 can be selected at process step **412** or one or more alternative bases are selected at process steps **414**, **416** and **418**. After base selection is effected, the sugar portion of the nucleoside is selected. Thus for each nucleoside, at process step **420** a sugar is selected that together with the select base will complete the nucleoside. In selecting the sugar, sugar chemistry 1 can be selected at process **422** or one or more alternative sugars are selected at process steps **424**, **426** and **428**. For each two adjacent nucleoside units, at process step **430**, the internucleoside linker is selected. The linker chemistry for the internucleoside linker can be linker chemistry 1 selected at process step **432** or one or more alternative internucleoside linker chemistries are selected at process steps **434**, **436** and **438**.

In addition to the base, sugar and internucleoside linkage, at each nucleoside position, one or more conjugate groups can be attached to the oligonucleotide via attachment to the nucleoside or attachment to the internucleoside linkage. The addition of a conjugate group is integrated at process step **440** and the assignment of the conjugate group is effected at process step **450**.

For illustrative purposes in Figures 10 and 11, for each of the bases, the sugars, the internucleoside linkers, or the conjugates, chemistries 1 through n are illustrated. As described in this specification, it is understood that the number of alternate chemistries between chemistry 1 and alternative chemistry n, for each of the bases, the sugars, the internucleoside linkages and the conjugates, is variable and includes, but is not limited to, each of the specific alternative bases, sugar, internucleoside linkers and conjugates identified in this specification as well as equivalents known in the art.

Utilizing the logic as described in conjunction with Figures 10 and 11, chemistry is assigned, as is shown in Figure 12, to the list of oligonucleotides from general procedure **300**. In assigning chemistries to the oligonucleotides in this list, a pointer can be set at process step **452** to the first oligonucleotide in the list and at step **453** to the first nucleotide of that first oligonucleotide. The base chemistry is selected at step **410**, as described above, the sugar chemistry is selected at step **420**, also as described above, followed by selection of the internucleoside linkage at step **430**, also as described above.

At decision 440, the process branches depending on whether a conjugate will be added at the current nucleotide position. If a conjugate is desired, the conjugate is selected at step 450, also as described above.

Whether or not a conjugate was added at decision step 440, an inquiry is made at decision step 454. This inquiry asks if the pointer resides at the last nucleotide in the current oligonucleotide. If the result at decision step 454 is "No," the pointer is moved to the next nucleotide in the current oligonucleotide and the loop including steps 410, 420, 430, 440 and 454 is repeated. This loop is reiterated until the result at decision step 454 is "Yes."

When the result at decision step 454 is "Yes," a query is made at decision step 460 concerning the location of the pointer in the list of oligonucleotides. If the pointer is not at the last oligonucleotide of the list, the "No" path of the decision step 460 is followed and the pointer is moved to the first nucleotide of the next oligonucleotide in the list at process step 458. With the pointer set to the next oligonucleotide in the list, the loop that starts at process steps 453 is reiterated. When the result at decision step 460 is "Yes," chemistry has been assigned to all of the nucleotides in the list of oligonucleotides.

### 11. Description of Oligonucleotide Chemistries.

As is illustrated in Figure 10, for each nucleoside of an oligonucleotide, chemistry selection includes selection of the base forming the nucleoside from a large palette of different base units available. These may be "modified" or "natural" bases (also reference herein as nucleobases) including the natural purine bases adenine (A) and guanine (G), and the natural pyrimidine bases thymine (T), cytosine (C) and uracil (U). They further can include modified nucleobases including other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo uracils and cytosines particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and

3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in the *Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch *et al.*, *Angewandte Chemie, International Edition*, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyl-adenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred for selection as the base. These are particularly useful when combined with a 2'-O-methoxyethyl sugar modifications, described below.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Patent 3,687,808, as well as U.S. Patents 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091; 5,614,617; and 5,681,941, each of which is incorporated herein by reference in its entirety. Reference is also made to allowed United States patent application 08/762,488, filed on December 10, 1996, commonly owned with the present application and which is incorporated herein by reference in its entirety.

In selecting the base for any particular nucleoside of an oligonucleotide, consideration is first given to the need of a base for a particular specificity for hybridization to an opposing strand of a particular target. Thus if an "A" base is required, adenine might be selected however other alternative bases that can effect hybridization in a manner mimicking an "A" base such as 2-aminoadenine might be selected should other consideration, e.g., stronger hybridization (relative to hybridization achieved with adenine), be desired.



As is illustrated in Figure 10, for each nucleoside of an oligonucleotide, chemistry selection includes selection of the sugar forming the nucleoside from a large palette of different sugar or sugar surrogate units available. These may be modified sugar groups, for instance sugars containing one or more substituent groups. Preferred substituent groups comprise the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; or O, S- or N-alkynyl; wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred substituent groups comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl), 2'-O-methoxyethyl, or 2'-MOE) (Martin *et al.*, *Helv. Chim. Acta*, 1995, 78, 486) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylamino oxyethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in co-owned United States patent application Serial Number 09/016,520, filed on January 30, 1998, which is incorporated herein by reference in its entirety.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the sugar group, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. The nucleosides of the oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar.

Representative United States patents that teach the preparation of such modified sugars structures include, but are not limited to, U.S. Patents 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811;

5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned with the present application, each of which is incorporated herein by reference in its entirety, together with allowed United States patent application 08/468,037, filed on June 5, 1995, which is  
5 commonly owned with the present application and which is incorporated herein by reference in its entirety.

As is illustrated in Figure 10, for each adjacent pair of nucleosides of an oligonucleotide, chemistry selection includes selection of the internucleoside linkage. These internucleoside linkages are also referred to as linkers, backbones or oligonucleotide  
10 backbones. For forming these nucleoside linkages, a palette of different internucleoside linkages or backbones is available. These include modified oligonucleotide backbones, for example, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates,  
15 phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

20 Representative United States patents that teach the preparation of the above phosphorus containing linkages include, but are not limited to, U.S. Patents 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,625,050; 25 and 5,697,248, certain of which are commonly owned with this application, each of which is incorporated herein by reference in its entirety.

Preferred internucleoside linkages for oligonucleotides that do not include a phosphorus atom therein, i.e., for oligonucleosides, have backbones that are formed by short chain alkyl or cycloalkyl intersugar linkages, mixed heteroatom and alkyl or  
30 cycloalkyl intersugar linkages, or one or more short chain heteroatomic or heterocyclic intersugar linkages. These include those having morpholino linkages (formed in part from

the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Patents 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, certain of which are commonly owned with this application, each of which is incorporated herein by reference in its entirety.

In other preferred oligonucleotides, i.e., oligonucleotide mimetics, both the sugar and the intersugar linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-phosphate backbone of an oligonucleotide is replaced with an amide-containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is incorporated herein by reference in its entirety. Further teaching of PNA compounds can be found in Nielsen *et al.*, *Science*, 1991, 254, 1497.

For the internucleoside linkages, the most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH<sub>2</sub>-NH-O-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-O-CH<sub>2</sub>- [known as a methylene (methylimino) or MMI backbone], -CH<sub>2</sub>-O-N(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-N(CH<sub>3</sub>)-CH<sub>2</sub>- and -O-N(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>2</sub>- (wherein the native phosphodiester backbone is represented as -O-P-O-CH<sub>2</sub>-) of the above referenced U.S. patent 5,489,677, and the amide

backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. Patent 5,034,506.

In attaching a conjugate group to one or more nucleosides or internucleoside linkages of an oligonucleotide, various properties of the oligonucleotide are modified. Thus modification of the oligonucleotides of the invention to chemically link one or more moieties or conjugates to the oligonucleotide are intended to enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety (Letsinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553), cholic acid (Manoharan *et al.*, *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan *et al.*, *Ann. N.Y. Acad. Sci.*, 1992, 660, 306; Manoharan *et al.*, *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765), a thiocholesterol (Oberhauser *et al.*, *Nucl. Acids Res.*, 1992, 20, 533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras *et al.*, *EMBO J.*, 1991, 10, 111; Kabanov *et al.*, *FEBS Lett.*, 1990, 259, 327; Svinarchuk *et al.*, *Biochimie*, 1993, 75, 49), a phospholipid, e.g., di-hexadecyl-*rac*-glycerol or triethylammonium 1,2-di-O-hexadecyl-*rac*-glycero-3-H-phosphonate (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36, 3651; Shea *et al.*, *Nucl. Acids Res.*, 1990, 18, 3777), a polyamine or a polyethylene glycol chain (Manoharan *et al.*, *Nucleosides & Nucleotides*, 1995, 14, 969), or adamantane acetic acid (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36, 3651), a palmityl moiety (Mishra *et al.*, *Biochim. Biophys. Acta*, 1995, 1264, 229), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, 277, 923).

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S. Patents 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923;

5,599,928 and 5,688,941, certain of which are commonly owned with the present application, and each of which is herein incorporated by reference in its entirety.

## 12. Chimeric Compounds.

It is not necessary for all positions in a given compound to be uniformly modified.

5 In fact, more than one of the aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes compounds which are chimeric compounds. "Chimeric" compounds or "chimeras," in the context of this invention, are compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one  
10 monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. An additional region of the oligonucleotide may serve as a substrate for  
15 enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids.

By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. Consequently, comparable results can often be obtained with shorter  
20 oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite  
25 structures representing the union of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as "hybrids" or "gapmers". Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S. Patents 5,013,830; 5,149,797; 5,220,007; 5,256,775;  
30 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the present application and each of

which is incorporated herein by reference in its entirety, together with commonly owned and allowed United States patent application serial number 08/465,880, filed on June 6, 1995, which is incorporated herein by reference in its entirety.

### 13. Description of Automated Oligonucleotide Synthesis.

5 In the next step of the overall process (illustrated in Figures 1 and 2), oligonucleotides are synthesized on an automated synthesizer. Although many devices may be employed, the synthesizer is preferably a variation of the synthesizer described in United States patents 5,472,672 and 5,529,756, each of which is incorporated herein by reference in its entirety. The synthesizer described in those patents is modified to include  
10 movement in along the Y axis in addition to movement along the X axis. As so modified, a 96-well array of compounds can be synthesized by the synthesizer. The synthesizer further includes temperature control and the ability to maintain an inert atmosphere during all phases of synthesis. The reagent array delivery format employs orthogonal X-axis motion of a matrix of reaction vessels and Y-axis motion of an array of reagents. Each  
15 reagent has its own dedicated plumbing system to eliminate the possibility of cross-contamination of reagents and line flushing and/or pipette washing. This in combined with a high delivery speed obtained with a reagent mapping system allows for the extremely rapid delivery of reagents. This further allows long and complex reaction sequences to be performed in an efficient and facile manner.

20 The software that operates the synthesizer allows the straightforward programming of the parallel synthesis of a large number of compounds. The software utilizes a general synthetic procedure in the form of a command (.cmd) file, which calls upon certain reagents to be added to certain wells *via* lookup in a sequence (.seq) file. The bottle position, flow rate, and concentration of each reagent is stored in a lookup table (.tab) file.  
25 Thus, once any synthetic method has been outlined, a plate of compounds is made by permutating a set of reagents, and writing the resulting output to a text file. The text file is input directly into the synthesizer and used for the synthesis of the plate of compounds. The synthesizer is interfaced with a relational database allowing data output related to the synthesized compounds to be registered in a highly efficient manner.

30 Building of the .seq, .cmd and .tab files is illustrated in Figure 13. Thus as a part of the general oligonucleotide synthesis procedure **500**, for each linker chemistry at

process step **502**, a synthesis file, i.e., a .cmd file, is built at process step **504**. This file can be built fresh to reflect a completely new set of machine commands reflecting a set of chemical synthesis steps or it can modify an existing file stored at process step **504** by editing that stored file in process step **508**. The .cmd files are built using a word processor and a command set of instructions as outlined below.

It will be appreciated that the preparation of control software and data files is within the routine skill of persons skilled in annotated nucleotide synthesis. The same will depend upon the hardware employed, the chemistries adopted and the design paradigm selected by the operator.

In a like manner to the building the .cmd files, .tab files are built to reflect the necessary reagents used in the automatic synthesizer for the particular chemistries that have been selected for the linkages, bases, sugars and conjugate chemistries. Thus for each of a set of these chemistries at process step **510**, a .tab file is built at process step **512** and stored at process step **514**. As with the .cmd files, an existing .tab file can be edited at process step **516**.

Both the .cmd files and the .tab files are linked together at process step **518** and stored for later retrieval in an appropriate sample database **520**. Linking can be as simple as using like file names to associate a .cmd file to its appropriate .tab file, e.g., synthesis\_1.cmd is linked to synthesis\_1.tab by use of the same preamble in their names.

The automated, multi-well parallel array synthesizer employs a reagent array delivery format, in which each reagent utilized has a dedicated plumbing system. As seen in Figures 23 and 24, an inert atmosphere **522** is maintained during all phases of a synthesis. Temperature is controlled *via* a thermal transfer plate **524**, which holds an injection molded reaction block **526**. The reaction plate assembly slides in the X-axis direction, while for example eight nozzle blocks (**528**, **530**, **532**, **534**, **536**, **538**, **540** and **542**) holding the reagent lines slide in the Y-axis direction, allowing for the extremely rapid delivery of any of 64 reagents to 96 wells. In addition, there are for example, six banks of fixed nozzle blocks (**544**, **546**, **548**, **550**, **552** and **554**) which deliver the same reagent or solvent to eight wells at once, for a total of 72 possible reagents.

In synthesizing oligonucleotides for screening, the target reaction vessels, a 96 well plate **556** (a 2-dimensional array), moves in one direction along the X axis, while the

series of independently controlled reagent delivery nozzles (528, 530, 532, 534, 536, 538, 540 and 542) move along the Y-axis relative to the reaction vessel 558. As the reaction plate 556 and reagent nozzles (528, 530, 532, 534, 536, 538, 540 and 542) can be moved independently at the same time, this arrangement facilitates the extremely rapid delivery of up to 72 reagents independently to each of the 96 reaction vessel wells.

The system software allows the straightforward programming of the synthesis of a large number of compounds by supplying the general synthetic procedure in the form of the command file to call upon certain reagents to be added to specific wells *via* lookup in the sequence file with the bottle position, flow rate, and concentration of each reagent being stored in the separate reagent table file. Compounds can be synthesized on various scales. For oligonucleotides, a 200 nmole scale is typically selected while for other compounds larger scales, as for example a 10  $\mu$ mole scale (3-5 mg), might be utilized. The resulting crude compounds are generally >80% pure, and are utilized directly for high throughput screening assays. Alternatively, prior to use the plates can be subjected to quality control (see general procedure 600 and Example 9) to ascertain their exact purity. Use of the synthesizer results in a very efficient means for the parallel synthesis of compounds for screening.

The software inputs accept tab delimited text files (as discussed above for file 504 and 512) from any text editor. A typical command file, a .cmd file, is shown in Example 3 at Table 2. Typical sequence files, .seq files, are shown in Example 3 at Tables 3 and 4 (.SEQ file), and a typical reagent file, a .tab file, is shown in Example 3 at Table 5. Table 3 illustrates the sequence file for an oligonucleotide having 2'-deoxy nucleotides at each position with a phosphorothioate backbone throughout. Table 4 illustrates the sequence file for an oligonucleotide, again having a phosphorothioate backbone throughout, however, certain modified nucleoside are utilized in portions of the oligonucleotide. As shown in this table, 2'-O-(2-methoxyethyl) modified nucleosides are utilized in a first region (a wing) of the oligonucleotide, followed by a second region (a gap) of 2'-deoxy nucleotides and finally a third region (a further wing) that has the same chemistry as the first region. Typically some of the wells of the 96 well plate 556 may be left empty (depending on the number of oligonucleotides to be made during an individual synthesis) or some of the wells may have oligonucleotides that will serve as standards for comparison



or analytical purposes.

Prior to loading reagents, moisture sensitive reagent lines are purged with argon at 522 for 20 minutes. Reagents are dissolved to appropriate concentrations and installed on the synthesizer. Large bottles, collectively identified as 558 in Figure 23 (containing 8 delivery lines) are used for wash solvents and the delivery of general activators, trityl group cleaving reagents and other reagents that may be used in multiple wells during any particular synthesis. Small septa bottles, collectively identified as 560 in Figure 23, are utilized to contain individual nucleotide amidite precursor compounds. This allows for anhydrous preparation and efficient installation of multiple reagents by using needles to pressurize the bottle, and as a delivery path. After all reagents are installed, the lines are primed with reagent, flow rates measured, then entered into the reagent table (.tab file). A dry resin loaded plate is removed from vacuum and installed in the machine for the synthesis.

The modified 96 well polypropylene plate 556 is utilized as the reaction vessel. The working volume in each well is approximately 700  $\mu$ l. The bottom of each well is provided with a pressed-fit 20  $\mu$ m polypropylene frit and a long capillary exit into a lower collection chamber as is illustrated in Figure 5 of the above referenced United States Patent 5,372,672. The solid support for use in holding the growing oligonucleotide during synthesis is loaded into the wells of the synthesis plate 556 by pipetting the desired volume of a balanced density slurry of the support suspended in an appropriate solvent, typically an acetonitrile-methylene chloride mixture. Reactions can be run on various scales as for instance the above noted 200 nmole and 10  $\mu$ mol scales. For oligonucleotide synthesis a CPG support is preferred, however other medium loading polystyrene-PEG supports such as TENTAGEL™ or ARGOGEL™ can also be used.

As seen in Figure 24, the synthesis plate is transported back and forth in the X-direction under an array of 8 moveable banks (530, 532, 534, 536, 538, 540, 542 and 544) of 8 nozzles (64 total) in the Y-direction, and 6 banks (544, 546, 548, 550, 552 and 554) of 48 fixed nozzles, so that each well can receive the appropriate amounts of reagents and/or solvents from any reservoir (large bottle or smaller septa bottle). A sliding balloon-type seal 562 surrounds this nozzle array and joins it to the reaction plate headspace 564. A slow sweep of nitrogen or argon 522 at ambient pressure across the plate headspace is used

to preserve an anhydrous environment.

The liquid contents in each well do not drip out until the headspace pressure exceeds the capillary forces on the liquid in the exit nozzle. A slight positive pressure in the lower collection chamber can be added to eliminate residual slow leakage from filled wells, or to effect agitation by bubbling inert gas through the suspension. In order to empty the wells, the headspace gas outlet valve is closed and the internal pressure raised to about 2 psi. Normally, liquid contents are blown directly to waste **566**. However, a 96 well microtiter plate can be inserted into the lower chamber beneath the synthesis plate in order to collect the individual well eluents for spectrophotometric monitoring (trityl, etc.) of reaction progress and yield.

The basic plumbing scheme for the machine is the gas-pressurized delivery of reagents. Each reagent is delivered to the synthesis plate through a dedicated supply line, collectively identified at **568**, solenoid valve collectively identified at **570** and nozzle, collectively identified at **572**. Reagents never cross paths until they reach the reaction well. Thus, no line needs to be washed or flushed prior to its next use and there is no possibility of cross-contamination of reagents. The liquid delivery velocity is sufficiently energetic to thoroughly mix the contents within a well to form a homogeneous solution, even when employing solutions having drastically different densities. With this mixing, once reactants are in homogeneous solution, diffusion carries the individual components into and out of the solid support matrix where the desired reaction takes place. Each reagent reservoir can be plumbed to either a single nozzle or any combination of up to 8 nozzles. Each nozzle is also provided with a concentric nozzle washer to wash the outside of the delivery nozzles in order to eliminate problems of crystallized reactant buildup due to slow evaporation of solvent at the tips of the nozzles. The nozzles and supply lines can be primed into a set of dummy wells directly to waste at any time.

The entire plumbing system is fabricated with teflon tubing, and reagent reservoirs are accessed *via* syringe needle/septa or direct connection into the higher capacity bottles. The septum vials **560** are held in removable 8-bottle racks to facilitate easy setup and cleaning. The priming volume for each line is about 350  $\mu$ l. The minimum delivery volume is about 2  $\mu$ l, and flow rate accuracy is  $\pm 5\%$ . The actual amount of material delivered depends on a timed flow of liquid. The flow rate for a particular solvent will

depend on its viscosity and wetting characteristics of the teflon tubing. The flow rate (typically 200-350  $\mu$ l per sec) is experimentally determined, and this information is contained in the reagent table setup file.

Heating and cooling of the reaction block **526** is effected utilizing a recirculating heat exchanger plate **524**, similar to that found in PCR thermocyclers, that nests with the polypropylene synthesis plate **556** to provide good thermal contact. The liquid contents in a well can be heated or cooled at about 10°C per minute over a range of +5 to +80°C, as polypropylene begins to soften and deform at about 80°C. For temperatures greater than this, a non-disposable synthesis plate machined from stainless steel or monel with replaceable frits can be utilized.

The hardware controller can be any of a wide variety, but conveniently can be designed around a set of three 1 MHz 86332 chips. This controller is used to drive the single X-axis and 8 Y-axis stepper motors as well as provide the timing functions for a total of 154 solenoid valves. Each chip has 16 bidirectional timer I/O and 8 interrupt channels in its timer processing unit (TPU). These are used to provide the step and direction signals, and to read 3 encoder inputs and 2 limit switches for controlling up to three motors per chip. Each 86332 chip also drives a serial chain of 8 UNC5891A darlington array chips to provide power to 64 valves with msec resolution. The controller communicates with the Windows software interface program running on a PC via a 19200 Hz serial channel, and uses an elementary instruction set to communicate valve\_number, time\_open, motor\_number and position\_data.

The three components of the software program that run the array synthesizer are the generalized procedure or command (.cmd) file which specifies the synthesis instructions to be performed, the sequence (.seq) file which specifies the scale of the reaction and the order in which variable groups will be added to the core synthon, and the reagent table (.tab) file which specifies the name of a chemical, its location (bottle number), flow rate, and concentration are utilized in conjunction with a basic set of command instructions.

One basic set of command instructions can be:

ADD

IF {block of instructions} END\_IF

## 38

```

REPEAT      {block of instructions}      END_REPEAT
PRIME, NOZZLE_WASH
WAIT, DRAIN
LOAD, REMOVE
5  NEXT_SEQUENCE
    LOOP_BEGIN, LOOP_END

```

The ADD instruction has two forms, and is intended to have the look and feel of a standard chemical equation. Reagents are specified to be added by a molar amount if the number proceeds the name identifier, or by an absolute volume in microliters if the number follows the identifier. The number of reagents to be added is a parsed list, separated by the “+” sign. For variable reagent identifiers, the key word, <seq>, means look in the sequence table for the identity of the reagent to be added, while the key word, <act>, means add the reagent which is associated with that particular <seq>. Reagents are delivered in the order specified in the list.

Thus:

ADD ACN 300

means: Add 300  $\mu$ l of the named reagent acetonitrile; ACN to each well of active synthesis

ADD <seq> 300

means: If the sequence pointer in the .seq file is to a reagent in the list of reagents, independent of scale, add 300  $\mu$ l of that particular reagent specified for that well.

ADD 1.1 PYR + 1.0 <seq> + 1.1 <act1>

means: If the sequence pointer in the .seq file is to a reagent in the list of acids in the Class ACIDS\_1, and PYR is the name of pyridine, and ethyl chloroformate is defined in the .tab file to activate the class, ACIDS\_1, then this instruction means:

Add 1.1 equiv. pyridine

1.0 equiv. of the acid specified for that well and

1.1 equiv. of the activator, ethyl chloroformate

The IF command allows one to test what type of reagent is specified in the <seq> variable

and process the succeeding block of commands accordingly.

Thus:

ACYLATION {the procedure name}

BEGIN

```
5          IF CLASS = ACIDS_1
            ADD 1.0 <seq> + 1.1 <act1> + 1.1 PYR
            WAIT 60
        ENDIF
        IF CLASS = ACIDS_2
10          ADD 1.0 <seq> + 1.2 <act1> + 1.2 TEA
        ENDIF
        WAIT 60
        DRAIN 10
    END
```

15 means: Operate on those wells for which reagents contained in the Acid\_1 class are specified, WAIT 60 sec, then operate on those wells for which reagents contained in the Acid\_2 class are specified, then WAIT 60 sec longer, then DRAIN the whole plate. Note that the Acid\_1 group has reacted for a total of 120 sec, while the Acid\_2 group has reacted for only 60 sec.

20 The REPEAT command is a simple way to execute the same block of commands multiple times.

Thus:

WASH\_1 {the procedure name}

BEGIN

```
25          REPEAT 3
            ADD ACN 300
            DRAIN 15
        END_REPEAT
    END
```

30 means: repeats the add acetonitrile and drain sequence for each well three times.

The PRIME command will operate either on specific named reagents or on nozzles

which will be used in the next associated <seq> operation. The  $\mu$ l amount dispensed into a prime port is a constant that can be specified in a config.dat file.

The NOZZLE\_WASH command for washing the outside of reaction nozzles free from residue due to evaporation of reagent solvent will operate either on specific named reagents or on nozzles which have been used in the preceding associated <seq> operation. The machine is plumbed such that if any nozzle in a block has been used, all the nozzles in that block will be washed into the prime port.

The WAIT and DRAIN commands are by seconds, with the drain command applying a gas pressure over the top surface of the plate in order to drain the wells.

The LOAD and REMOVE commands are instructions for the machine to pause for operator action.

The NEXT\_SEQUENCE command increments the sequence pointer to the next group of substituents to be added in the sequence file. The general form of a .seq file entry is the definition:

Well_No	Well_ID	Scale	Sequence
---------	---------	-------	----------

The sequence information is conveyed by a series of columns, each of which represents a variable reagent to be added at a particular position. The scale ( $\mu$ mole) variable is included so that reactions of different scale can be run at the same time if desired. The reagents are defined in a lookup table (the .tab file), which specifies the name of the reagent as referred to in the sequence and command files, its location (bottle number), flow rate, and concentration. This information is then used by the controller software and hardware to determine both the appropriate slider motion to position the plate and slider arms for delivery of a specific reagent, as well as the specific valve and time required to deliver the appropriate reagents. The adept classification of reagents allows the use of conditional IF loops from within a command file to perform addition of different reagents differently during a "single step" performed across 96 wells simultaneously. The special class ACTIVATORS defines certain reagents that always get added with a particular class of reagents (for example tetrazole during a phosphitylation reaction in adding the next nucleotide to a growing oligonucleotide).

The general form of the .tab file is the definition:

<b>Class</b>	<b>Bottle</b>	<b>Reagent Name</b>	<b>Flow_rate</b>	<b>Conc.</b>
--------------	---------------	---------------------	------------------	--------------

5           The LOOP\_BEGIN and LOOP\_END commands define the block of commands which will continue to operate until a NEXT\_SEQUENCE command points past the end of the longest list of reactants in any well.

          Not included in the command set is a MOVE command. For all of the above commands, if any plate or nozzle movement is required, this is automatically executed in order to perform the desired solvent or reagent delivery operation. This is accomplished by the controller software and hardware, which determines the correct nozzle(s) and well(s) required for a particular reagent addition, then synchronizes the position of the requisite nozzle and well prior to adding the reagent.

15           A MANUAL mode can also be utilized in which the synthesis plate and nozzle blocks can be "homed" or moved to any position by the operator, the nozzles primed or washed, the various reagent bottles depressurized or washed with solvent, the chamber pressurized, etc. The automatic COMMAND mode can be interrupted at any point, MANUAL commands executed, and then operation resumed at the appropriate location. The sequence pointer can be incremented to restart a synthesis anywhere within a command file.

20           In reference to Figure 14, the list of oligonucleotides for synthesis can be rearranged or grouped for optimization of synthesis. Thus at process step 574, the oligonucleotides are grouped according to a factor on which to base the optimization of synthesis. As illustrated in the Examples below, one such factor is the 3' most nucleoside of the oligonucleotide. Using the amidite approach for oligonucleotide synthesis, a nucleotide bearing a 3' phosphoramite is added to the 5' hydroxyl group of a growing nucleotide chain. The first nucleotide (at the 3' terminus of the oligonucleotide - the 3' most nucleoside) is first connected to a solid support. This is normally done batchwise on a large scale as is standard practice during oligonucleotide synthesis.

30           Such solid supports pre-loaded with a nucleoside are commercially available. In utilizing the multi well format for oligonucleotide synthesis, for each oligonucleotide to be

synthesized, an aliquot of a solid support bearing the proper nucleoside thereon is added to the well for synthesis. Prior to loading the sequence of oligonucleotides to be synthesized in the .seq file, they are sorted by the 3' terminal nucleotide. Based on that sorting, all of the oligonucleotide sequences having an "A" nucleoside at their 3' end are grouped together, those with a "C" nucleoside are grouped together as are those with "G" or "T" nucleosides. Thus in loading the nucleoside-bearing solid support into the synthesis wells, machine movements are conserved.

The oligonucleotides can be grouped by the above described parameter or other parameters that facilitate the synthesis of the oligonucleotides. Thus in Figure 14, sorting is noted as being effected by some parameter of type 1, as for instance the above described 3' most nucleoside, or other types of parameters from type 2 to type n at process steps 576, 578 and 580. Since synthesis will be from the 3' end of the oligonucleotides to the 5' end, the oligonucleotide sequences are reverse sorted to read 3' to 5'. The oligonucleotides are entered in the .seq file in this form, i.e., reading 3' to 5'.

Once sorted into types, the position of the oligonucleotides on the synthesis plates is specified at process step 582 by the creation of a .seq file as described above. The .seq file is associated with the respective .cmd and .tab files needed for synthesis of the particular chemistries specified for the oligonucleotides at process step 584 by retrieval of the .cmd and .tab files at process step 586 from the sample database 520. These files are then input into the multi well synthesizer at process step 588 for oligonucleotide synthesis. Once physically synthesized, the list of oligonucleotides again enters the general procedure flow as indicated in Figure 1. For shipping, storage or other handling purposes, the plates can be lyophilized at this point if desired. Upon lyophilization, each well contains the oligonucleotides located therein as a dry compound.

#### 14. Quality Control.

In an optional step, quality control is performed on the oligonucleotides at process step 600 after a decision is made (decision step 550) to perform quality control. Although optional, quality control may be desired when there is some reason to think that some aspect of the synthetic process step 500 has been compromised. Alternatively, samples of the oligonucleotides may be taken and stored in the event that the results of assays conducted using the oligonucleotides (process step 700) yield confusing results or



suboptimal data. In the latter event, for example, quality control might be performed after decision step **800** if no oligonucleotides with sufficient activity are identified. In either event, decision step **650** follows quality control step process **600**. If one or more of the oligonucleotides do not pass quality control, process step **500** can be repeated, i.e., the oligonucleotides are synthesized for a second time.

The operation of the quality control system general procedure **600** is detailed in steps **610-660** of Figure 15. Also referenced in the following discussion are the robotics and associated analytical instrumentation as shown in Figure 18.

During step **610** (Figure 15), sterile, double-distilled water is transferred by an automated liquid handler (**2040** of Figure 18) to each well of a multi-well plate containing a set of lyophilized antisense oligonucleotides. The automated liquid handler (**2040** of Figure 18) reads the barcode sticker on the multi-well plate to obtain the plate's identification number. Automated liquid handler **2040** then queries Sample Database **520** (which resides in Database Server **2002** of Figure 18) for the quality control assay instruction set for that plate and executes the appropriate steps. Three quality control processes are illustrated, however, it is understood that other quality control processes or steps maybe practiced in addition to or in place of the processes illustrated.

The first illustrative quality control process (steps **622** to **626**) quantitates the concentration of oligonucleotide in each well. If this quality control step is performed, an automated liquid handler (**2040** of Figure 18) is instructed to remove an aliquot from each well of the master plate and generate a replicate daughter plate for transfer to the UV spectrophotometer (**2016** of Figure 18). The UV spectrophotometer (**2016** of Figure 18) then measures the optical density of each well at a wavelength of 260 nanometers. Using standardized conversion factors, a microprocessor within UV spectrophotometer (**2016** of Figure 18) then calculates a concentration value from the measured absorbance value for each well and output the results to Sample Database **520**.

The second illustrative quality control process steps **632** to **636**) quantitates the percent of total oligonucleotide in each well that is full length. If this quality control step is performed, an automated liquid handler (**2040** of Figure 18) is instructed to remove an aliquot from each well of the master plate and generate a replicate daughter plate for transfer to the multichannel capillary gel electrophoresis apparatus (**2022** of Figure 18).

The apparatus electrophoretically resolves in capillary tube gels the oligonucleotide product in each well. As the product reaches the distal end of the tube gel during electrophoresis, a detection window dynamically measures the optical density of the product that passes by it. Following electrophoresis, the value of percent product that passed by the detection window with respect to time is utilized by a built in microprocessor to calculate the relative size distribution of oligonucleotide product in each well. These results are then output to the Sample Database (520).

The third illustrative quality control process steps 632 to 636 quantitates the mass of the oligonucleotide in each well that is full length. If this quality control step is performed, an automated liquid handler (2040 of Figure 18) is instructed to remove an aliquot from each well of the master plate and generate a replicate daughter plate for transfer to the multichannel liquid electrospray mass spectrometer (2018 of Figure 18). The apparatus then uses electrospray technology to inject the oligonucleotide product into the mass spectrometer. A built in microprocessor calculates the mass-to-charge ratio to arrive at the mass of oligonucleotide product in each well. The results are then output to Sample Database 520.

Following completion of the selected quality control processes, the output data is manually examined or is examined using an appropriate algorithm and a decision is made as to whether or not the plate receives "Pass" or "Fail" status. The current criteria for acceptance, for 18 mer oligonucleotides, is that at least 85% of the oligonucleotides in a multi-well plate must be 85% or greater full length product as measured by both capillary gel electrophoresis and mass spectrometry. An input (manual or automated) is then made into Sample Database 520 as to the pass/fail status of the plate. If a plate fails, the process cycles back to step 500, and a new plate of the same oligonucleotides is automatically placed in the plate synthesis request queue (process 554 of Figure 15). If a plate receives "Pass" status, an automated liquid handler (2040 of Figure 18) is instructed to remove appropriate aliquots from each well of the master plate and generate two replicate daughter plates in which the oligonucleotide in each well is at a concentration of 30 micromolar. The plate then moves on to process 700 for oligonucleotide activity evaluation.

**15. Cell Lines for Assaying Oligonucleotide Activity.** The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types

provided that the target nucleic acid, or its gene product, is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following four cell types are provided for illustrative purposes, but other cell types can be routinely used.

5                   **T-24 cells:** The transitional cell bladder carcinoma cell line T-24 is obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in complete McCoy's 5A basal media (Life Technologies, Gaithersburg, MD) supplemented with 10% fetal calf serum, penicillin 100 units per milliliter, and streptomycin 100 micrograms per milliliter (all from Life Technologies).  
10 Cells are routinely passaged by trypsinization and dilution when they reach 90% confluence. Cells are routinely seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis. For Northern blotting or other analysis, cells are seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

15                   **A549 cells:** The human lung carcinoma cell line A549 is obtained from the ATCC (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Life Technologies) supplemented with 10% fetal calf serum, penicillin 100 units per milliliter, and streptomycin 100 micrograms per milliliter (all from Life Technologies). Cells are routinely passaged by trypsinization and dilution when they reach 90% confluence.

20                   **NHDF cells:** Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville, MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corp.) as provided by the supplier. Cells are maintained for up to 10 passages as recommended by the supplier.

**HEK cells:** Human embryonic keratinocytes (HEK) were obtained from  
25 the Clonetics Corp. HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corp.) as provided by the supplier. Cell are routinely maintained for up to 10 passages as recommended by the supplier.

#### **16. Treatment of Cells with Candidate Compounds:**

                    When cells reach about 80% confluency, they are treated with oligonucleotide. For  
30 cells grown in 96-well plates, wells are washed once with 200  $\mu$ l OPTI-MEM-1<sup>TM</sup> reduced-serum medium (Life Technologies) and then treated with 130  $\mu$ l of OPTI-MEM-

1<sup>TM</sup> containing 3.75  $\mu$ g/ml LIPOFECTIN<sup>TM</sup> (Life Technologies) and the desired oligonucleotide at a final concentration of 150 nM. After 4 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16 hours after oligonucleotide treatment.

5 Alternatively, for cells resistant to cationic mediated transfection, oligonucleotides can be introduced by electroporation. Electroporation conditions must be optimized for every cell type. In general, oligonucleotide is added directly to complete growth media to a final concentration between 1 and 20 micromolar. An electronic pulse is delivered to the cells using a BTX T820 ELECTRO SQUARE PORATOR<sup>TM</sup> using a Multi-coaxial 96-  
10 well electrode (BT840) (BTX Corporation, San Diego, California). Following electroporation, the cells are returned to the incubator for 16 hours.

#### 17. Assaying Oligonucleotide Activity:

Oligonucleotide-mediated modulation of expression of a target nucleic acid can be assayed in a variety of ways known in the art. For example, target RNA levels can be  
15 quantitated by, e.g., Northern blot analysis, competitive PCR, or reverse transcriptase polymerase chain reaction (RT-PCR). RNA analysis can be performed on total cellular RNA or, preferably in the case of polypeptide-encoding nucleic acids, poly(A)+ mRNA. For RT-PCR, poly(A)+ mRNA is preferred. Methods of RNA isolation are taught in, for example, Ausubel *et al.* (*Short Protocols in Molecular Biology*, 2nd Ed., pp. 4-1 to 4-13,  
20 Greene Publishing Associates and John Wiley & Sons, New York, 1992). Northern blot analysis is routine in the art (*Id.*, pp. 4-14 to 4-29).

Alternatively, total RNA can be prepared from cultured cells or tissue using the QIAGEN RNeasy®-96 kit for the high throughput preparation of RNA (QIAGEN, Inc., Valencia, CA). Essentially, protocols are carried out according to the manufacturer's  
25 directions. Optionally, a DNase step is included to remove residual DNA prior to RT-PCR.

To improve efficiency and accuracy the repetitive pipeting steps and elution step have been automated using a QIAGEN Bio-Robot 9604. Essentially after lysing of the oligonucleotide treated cell cultures in situ, the plate is transferred to the robot deck where  
30 the pipeting, DNase treatment, and elution steps are carried out.

Reverse transcriptase polymerase chain reaction (RT-PCR) can be conveniently

accomplished using the commercially available ABI PRISM® 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to the manufacturer's instructions. Other methods of PCR are also known in the art.

5 Target protein levels can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis (immunoblotting), Enzyme-linked immunosorbent assay (ELISA) or fluorescence-activated cell sorting (FACS). Antibodies directed to a protein encoded by a target nucleic acid can be identified and obtained from a variety of sources, such as the MSRS catalog of antibodies, (Aerie Corporation, Birmingham, MI or via the internet at <http://www.ANTIBODIES-PROBES.com/>), or can  
10 be prepared via conventional antibody generation methods. Methods for preparation of polyclonal, monospecific ("antipeptide") and monoclonal antisera are taught by, for example, Ausubel *et al.* (*Short Protocols in Molecular Biology*, 2nd Ed., pp. 11-3 to 11-54, Greene Publishing Associates and John Wiley & Sons, New York, 1992).

Immunoprecipitation methods are standard in the art and are described by, for  
15 example, Ausubel *et al.* (*Id.*, pp. 10-57 to 10-63). Western blot (immunoblot) analysis is standard in the art (*Id.*, pp. 10-32 to 10-10-35). Enzyme-linked immunosorbent assays (ELISA) are standard in the art (*Id.*, pp. 11-5 to 11-17).

Because it is preferred to assay the compounds of the invention in a batchwise fashion, i.e., in parallel to the automated synthesis process described above, preferred  
20 means of assaying are suitable for use in 96-well plates and with robotic means. Accordingly, automated RT-PCR is preferred for assaying target nucleic acid levels, and automated ELISA is preferred for assaying target protein levels.

The assaying step, general procedure step **700**, is described in detail in Figure 16. After an appropriate cell line is selected at process step **710**, a decision is made at decision  
25 step **714** as to whether RT-PCR will be the only method by which the activity of the compounds is evaluated. In some instances, it is desirable to run alternative assay methods at process step **718**; for example, when it is desired to assess target polypeptide levels as well as target RNA levels, an immunoassay such as an ELISA is run in parallel with the RT-PCR assays. Preferably, such assays are tractable to semi-automated or robotic means.

30 When RT-PCR is used to evaluate the activities of the compounds, cells are plated into multi-well plates (typically, 96-well plates) in process step **720** and treated with test or

control oligonucleotides in process step 730. Then, the cells are harvested and lysed in process step 740 and the lysates are introduced into an apparatus where RT-PCR is carried out in process step 750. A raw data file is generated, and the data is downloaded and compiled at step 760. Spreadsheet files with data charts are generated at process step 770, and the experimental data is analyzed at process step 780. Based on the results, a decision is made at process step 785 as to whether it is necessary to repeat the assays and, if so, the process begins again with step 720. In any event, data from all the assays on each oligonucleotide are compiled and statistical parameters are automatically determined at process step 790.

#### **18. Classification of Compounds Based on Their Activity:**

Following assaying, general procedure step 700, oligonucleotide compounds are classified according to one or more desired properties. Typically, three classes of compounds are used: active compounds, marginally active (or "marginal") compounds and inactive compounds. To some degree, the selection criteria for these classes vary from target to target, and members of one or more classes may not be present for a given set of oligonucleotides.

However, some criteria are constant. For example, inactive compounds will typically comprise those compounds having 5% or less inhibition of target expression (relative to basal levels). Active compounds will typically cause at least 30% inhibition of target expression, although lower levels of inhibition are acceptable in some instances. Marginal compounds will have activities intermediate between active and inactive compounds, with preferred marginal compounds having activities more like those of active compounds.

#### **19. Optimization of Lead Compounds by Sequence.**

One means by which oligonucleotide compounds are optimized for activity is by varying their nucleobase sequences so that different regions of the target nucleic acid are targeted. Some such regions will be more accessible to oligonucleotide compounds than others, and "sliding" a nucleobase sequence along a target nucleic acid only a few bases can have significant effects on activity. Accordingly, varying or adjusting the nucleobase sequences of the compounds of the invention is one means by which suboptimal compounds can be made optimal, or by which new active compounds can be generated.

The operation of the gene walk process 1100 detailed in steps 1104-1112 of Figure 17 is detailed as follows. As used herein, the term "gene walk" is defined as the process by which a specified oligonucleotide sequence  $x$  that binds to a specified nucleic acid target  $y$  is used as a frame of reference around which a series of new oligonucleotides sequences capable of hybridizing to nucleic acid target  $y$  are generated that are sequence shifted increments of oligonucleotide sequence  $x$ . Gene walking can be done "downstream", "upstream" or in both directions from a specified oligonucleotide.

During step 1104 the user manually enters the identification number of the oligonucleotide sequence around which it is desired to execute gene walk process 1100 and the name of the corresponding target nucleic acid. The user then enters the scope of the gene walk at step 1104, by which is meant the number of oligonucleotide sequences that it is desired to generate. The user then enters in step 1108 a positive integer value for the sequence shift increment. Once this data is generated, the gene walk is effected. This causes a subroutine to be executed that automatically generates the desired list of sequences by walking along the target sequence. At that point, the user proceeds to process 400 to assign chemistries to the selected oligonucleotides.

Example 16 below, details a gene walk. In subsequent steps, this new set of nucleobase sequences generated by the gene walk is used to direct the automated synthesis at general procedure step 500 of a second set of candidate oligonucleotides. These compounds are then taken through subsequent process steps to yield active compounds or reiterated as necessary to optimize activity of the compounds.

## 20. Optimization of Lead Compounds by Chemistry.

Another means by which oligonucleotide compounds of the invention are optimized is by reiterating portions of the process of the invention using marginal or active compounds from the first iteration and selecting additional chemistries to the nucleobase sequences thereof.

Thus, for example, an oligonucleotide chemistry different from that of the first set of oligonucleotides is assigned at general procedure step 400. The nucleobase sequences of marginal compounds are used to direct the synthesis at general procedure step 500 of a second set of oligonucleotides having the second assigned chemistry. The resulting second set of oligonucleotide compounds is assayed in the same manner as the first set at

procedure process step **700** and the results are examined to determine if compounds having sufficient activity have been generated at decision step **800**.

### **21. Identification of Sites Amenable to Antisense Technologies.**

5 In a related process, a second oligonucleotide chemistry is assigned at procedure step **400** to the nucleobase sequences of all of the oligonucleotides (or, at least, all of the active and marginal compounds) and a second set of oligonucleotides is synthesized at procedure step **500** having the same nucleobase sequences as the first set of compounds. The resulting second set of oligonucleotide compounds is assayed in the same manner as the first set at procedure step **700** and active and marginal compounds are identified at  
10 procedure steps **800** and **1000**.

In order to identify sites on the target nucleic acid that are amenable to a variety of antisense technologies, the following mathematically simple steps are taken. The sequences of active and marginal compounds from two or more such automated syntheses/assays are compared and a set of nucleobase sequences that are active, or  
15 marginally so, in both sets of compounds is identified. The reverse complements of these nucleobase sequences corresponds to sequences of the target nucleic acid that are tractable to a variety of antisense and other sequence-based technologies. These antisense-sensitive sites are assembled into contiguous sequences (contigs) using the procedures described for assembling target nucleotide sequences (at procedure step **200**).

### **22. Systems for Executing Preferred Methods of the Invention.**

20 An embodiment of computer, network and instrument resources for effecting the methods of the invention is shown in Figure 18. In this embodiment, four computer servers are provided. First, a large database server **2002** stores all chemical structure, sample tracking and genomic, assay, quality control, and program status data. Further, this  
25 database server serves as the platform for a document management system. Second, a compute engine **2004** runs computational programs including RNA folding, oligonucleotide walking, and genomic searching. Third, a file server **2006** allows raw instrument output storage and sharing of robot instructions. Fourth, a groupware server **2008** enhances staff communication and process scheduling.

30 A redundant high-speed network system is provided between the main servers and the bridges **2026**, **2028** and **2030**. These bridges provide reliable network access to the



many workstations and instruments deployed for this process. The instruments selected to support this embodiment are all designed to sample directly from standard 96 well microtiter plates, and include an optical density reader **2016**, a combined liquid chromatography and mass spectroscopy instrument **2018**, a gel fluorescence and scintillation imaging system **2032** and **2042**, a capillary gel electrophoreses system **2022** and a real-time PCR system **2034**.

Most liquid handling is accomplished automatically using robots with individually controllable robotic pipetters **2038** and **2020** as well as a 96-well pipette system **2040** for duplicating plates. Windows NT or Macintosh workstations **2044**, **2024**, and **2036** are deployed for instrument control, analysis and productivity support.

### **23. Relational Database.**

Data is stored in an appropriate database. For use with the methods of the invention, a relational database is preferred. Figure 19 illustrates the data structure of a sample relational database. Various elements of data are segregated among linked storage elements of the database.

## **EXAMPLES**

The following examples illustrate the invention and are not intended to limit the same. Those skilled in the art will recognize, or be able to ascertain through routine experimentation, numerous equivalents to the specific procedures, materials and devices described herein. Such equivalents are considered to be within the scope of the present invention.

### **EXAMPLE 1: Selection of CD40 as a Target**

Cell-cell interactions are a feature of a variety of biological processes. In the activation of the immune response, for example, one of the earliest detectable events in a normal inflammatory response is adhesion of leukocytes to the vascular endothelium, followed by migration of leukocytes out of the vasculature to the site of infection or injury. The adhesion of leukocytes to vascular endothelium is an obligate step in their migration out of the vasculature (for a review, see Albelda *et al.*, *FASEB J.*, 1994, 8, 504). As is well known in the art, cell-cell interactions are also critical for propagation of both B-

lymphocytes and T-lymphocytes resulting in enhanced humoral and cellular immune responses, respectively (for a reviews, see Makgoba *et al.*, *Immunol. Today*, 1989, 10, 417; Janeway, *Sci. Amer.*, 1993, 269, 72).

CD40 was first characterized as a receptor expressed on B-lymphocytes. It was later found that engagement of B-cell CD40 with CD40L expressed on activated T-cells is essential for T-cell dependent B-cell activation (i.e. proliferation, immunoglobulin secretion, and class switching) (for a review, see Gruss *et al. Leuk. Lymphoma*, 1997, 24, 393). A full cDNA sequence for CD40 is available (GenBank accession number X60592, incorporated herein by reference as SEQ ID NO:85).

As interest in CD40 mounted, it was subsequently revealed that functional CD40 is expressed on a variety of cell types other than B-cells, including macrophages, dendritic cells, thymic epithelial cells, Langerhans cells, and endothelial cells (*Ibid.*). These studies have led to the current belief that CD40 plays a much broader role in immune regulation by mediating interactions of T-cells with cell types other than B-cells. In support of this notion, it has been shown that stimulation of CD40 in macrophages and dendritic results is required for T-cell activation during antigen presentation (*Id.*). Recent evidence points to a role for CD40 in tissue inflammation as well. Production of the inflammatory mediators IL-12 and nitric oxide by macrophages has been shown to be CD40 dependent (Buhlmann *et al.*, *J. Clin. Immunol.*, 1996, 16, 83). In endothelial cells, stimulation of CD40 by CD40L has been found to induce surface expression of E-selectin, ICAM-1, and VCAM-1, promoting adhesion of leukocytes to sites of inflammation (Buhlmann *et al.*, *J. Clin. Immunol.*, 1996, 16, 83; Gruss *et al.*, *Leuk Lymphoma*, 1997, 24, 393). Finally, a number of reports have documented overexpression of CD40 in epithelial and hematopoietic tumors as well as tumor infiltrating endothelial cells, indicating that CD40 may play a role in tumor growth and/or angiogenesis as well (Gruss *et al.*, *Leuk Lymphoma*, 1997, 24, 393-422; Kluth *et al. Cancer Res*, 1997, 57, 891).

Due to the pivotal role that CD40 plays in humoral immunity, the potential exists that therapeutic strategies aimed at downregulating CD40 may provide a novel class of agents useful in treating a number of immune associated disorders, including but not limited to graft versus host disease, graft rejection, and autoimmune diseases such as multiple sclerosis, systemic lupus erythematosus, and certain forms of arthritis. Inhibitors

of CD40 may also prove useful as an anti-inflammatory compound, and could therefore be useful as treatment for a variety of diseases with an inflammatory component such as asthma, rheumatoid arthritis, allograft rejections, inflammatory bowel disease, and various dermatological conditions, including psoriasis. Finally, as more is learned about the association between CD40 overexpression and tumor growth, inhibitors of CD40 may prove useful as anti-tumor agents as well.

Currently, there are no known therapeutic agents which effectively inhibit the synthesis of CD40. To date, strategies aimed at inhibiting CD40 function have involved the use of a variety of agents that disrupt CD40/CD40L binding. These include monoclonal antibodies directed against either CD40 or CD40L, soluble forms of CD40, and synthetic peptides derived from a second CD40 binding protein, A20. The use of neutralizing antibodies against CD40 and/or CD40L in animal models has provided evidence that inhibition of CD40 stimulation would have therapeutic benefit for GVHD, allograft rejection, rheumatoid arthritis, SLE, MS, and B-cell lymphoma (Buhlmann *et al.*, *J. Clin. Immunol.*, 1996, 16, 83). However, due to the expense, short half-life, and bioavailability problems associated with the use of large proteins as therapeutic agents, there is a long felt need for additional agents capable of effectively inhibiting CD40 function. Oligonucleotides compounds avoid many of the pitfalls of current agents used to block CD40/CD40L interactions and may therefore prove to be uniquely useful in a number of therapeutic applications.

## **EXAMPLE 2: Generation of Virtual Oligonucleotides Targeted to CD40**

The process of the invention was used to select oligonucleotides targeted to CD40, generating the list of oligonucleotide sequences with desired properties as shown in Figure 22. From the assembled CD40 sequence, the process began with determining the desired oligonucleotide length to be eighteen nucleotides, as represented in step 2500. All possible oligonucleotides of this length were generated by Oligo 5.0™, as represented in step 2504. Desired thermodynamic properties were selected in step 2508. The single parameter used was oligonucleotides of melting temperature less than or equal to 40°C were discarded. In step 2512, oligonucleotide melting temperatures were calculated by Oligo 5.0™. Oligonucleotide sequences possessing an undesirable score were discarded. It

is believed that oligonucleotides with melting temperatures near or below physiological and cell culture temperatures will bind poorly to target sequences. All oligonucleotide sequences remaining were exported into a spreadsheet. In step 2516, desired sequence properties are selected. These include discarding oligonucleotides with at least one stretch of four guanoses in a row and stretches of six of any other nucleotide in a row. In step 2520, a spreadsheet macro removed all oligonucleotides containing the text string "GGGG." In step 2524, another spreadsheet macro removed all oligonucleotides containing the text strings "AAAAAA" or "CCCCCC" or "TTTTTT." From the remaining oligonucleotide sequences, 84 sequences were selected manually with the criteria of having an uniform distribution of oligonucleotide sequences throughout the target sequence, as represented in step 2528. These oligonucleotide sequences were then passed to the next step in the process, assigning actual oligonucleotide chemistries to the sequences.

**EXAMPLE 3: Input Files For Automated Oligonucleotide Synthesis Command File (.cmd File)**

Table 2 is a command file for synthesis of an oligonucleotide having regions of 2'-O-(2-methoxyethyl) nucleosides and a central region of 2'-deoxy nucleosides each linked by phosphorothioate internucleotide linkages.

**Table 2**

```
SOLID_SUPPORT_SKIP
      BEGIN
                Next_Sequence
      END

INITIAL-WASH
      BEGIN
                Add ACN 300
                Drain 10
      END
```

55

```
LOOP-BEGIN
DEBLOCK
    BEGIN
        Prime TCA
        Load Tray
        Repeat 2
            Add TCA 150
            Wait 10
            Drain 8
        End_Repeat
        Remove Tray
        Add TCA 125
        Wait 10
        Drain 8
    END
END

WASH_AFTER_DEBLOCK
    BEGIN
        Repeat 3
            Add ACN 250 To_All
            Drain 10
        End_Repeat
    END

COUPLING
    BEGIN
        if class = DEOXY_THIOATE
            Nozzle wash <act1>
            prime <act1>
            prime <seq>
            Add <act1> 70 + <seq> 70
```

56

```
        Wait 40
            Drain 5
        end-if
    if class = MOE_THIOATE
5        Nozzle wash <act1>
            Prime <act1>
            prime <seq>
            Add <act1> 120 + <seq> 120
            Wait 230
10        Drain 5
        End_if
    END
```

```
WASH_AFTER_COUPLING
    BEGIN
15        Add ACN 200 To_All
        Drain 10
    END
```

```
OXIDIZE
20    BEGIN
        if class = DEOXY_THIOATE
        Add BEAU 180
        Wait 40
            Drain 7
25        end_if
        if class = MOE_THIOATE
        Add BEAU 200
            Wait 120
            Drain 7
30        end_if
```

57

END

CAP

BEGIN

5                   Add CAP\_B 80 + CAP\_A 80

Wait 20

Drain 7

END

10    WASH\_AFTER\_CAP

BEGIN

Add ACN 150 To\_All

Drain 5

Add ACN 250 To\_All

15                   Drain 11

END

BASE\_COUNTER

BEGIN

20                   Next\_Sequence

END

LOOP\_END

DEBLOCK\_FINAL

25                   BEGIN

Prime TCA

Load Tray

Repeat 2

Add TCA 150 To\_All

30                   Wait 10

58

```

        Drain 8
            End_Repeat
            Remove Tray
            Add TCA 125 To_All
5         Wait 10
            Drain 10
        END

FINAL_WASH
        BEGIN
10         Repeat 4
            Add ACN 300 to_All
            Drain_12
            End_Repeat
        END
15     ENDALL
        BEGIN
            Wait 3
        END

```

**Sequence files (.seq Files)**

20        Table 3 is a .seq file for oligonucleotides having 2'-deoxy nucleosides linked by phosphorothioate internucleotide linkages.

**Table 3**

Identity of columns: **Syn #, Well, Scale, Nucleotide at particular position** (identified using base identifier followed by backbone identifier where "s" is phosphorothioate).

25        Note the columns wrap around to next line when longer than one line.

1	A01	200	As	Cs	Cs	As	Gs	Gs	As	Cs	Gs
Gs	Cs	Gs	Gs	As	Cs	Cs	As	G			



## 59

5	2	A02	200	As	Cs	Gs	Gs	Cs	Gs	Gs	As	Cs
	Cs	As	Gs	As	Gs	Ts	Gs	Gs	A			
	3	A03	200	As	Cs	Cs	As	As	Gs	Cs	As	Gs
10				As	Cs	Gs	Gs	As	Cs	G		
	4	A04	200	As	Gs	Gs	As	Gs	As	Cs	Cs	Cs
	Cs	Gs	As	Cs	Gs	As	As	Cs	G			
15	5	A05	200	As	Cs	Cs	Cs	Cs	Gs	As	Cs	Gs
	As	As	Cs	Gs	As	Cs	Ts	Gs	G			
	6	A06	200	As	Cs	Gs	As	As	Cs	Gs	As	Cs
20				Ts	Gs	Gs	Cs	As	G			
	7	A07	200	As	Cs	Gs	As	Cs	Ts	Gs	Gs	Cs
	Gs	As	Cs	As	Gs	Gs	Ts	As	G			
25	8	A08	200	As	Cs	As	Gs	Gs	Ts	As	Gs	Gs
	Ts	Cs	Ts	Ts	Gs	Gs	Ts	Gs	G			
	9	A09	200	As	Gs	Gs	Ts	Cs	Ts	Ts	Gs	Gs
30				Ts	Gs	Gs	As	Cs	G			
	10	A10	200	As	Gs	Ts	Cs	As	Cs	Gs	As	Cs
	As	As	Gs	As	As	As	Cs	As	C			
30	11	A11	200	As	Cs	Gs	As	Cs	As	As	Gs	As
	As	As	Cs	As	Cs	Gs	Gs	Ts	C			
	12	A12	200	As	Gs	As	As	As	Cs	As	Cs	Gs
30				Gs	Ts	Cs	Gs	Gs	Ts	Cs	Cs	T
	13	B01	200	As	As	Cs	As	Cs	Gs	Gs	Ts	Cs
	Gs	Gs	Ts	Cs	Cs	Ts	Gs	Ts	C			
30	14	B02	200	As	Cs	Ts	Cs	As	Cs	Ts	Gs	As
	Cs	Gs	Ts	Gs	Ts	Cs	Ts	Cs	A			
	15	B03	200	As	Cs	Gs	Gs	As	As	Gs	Gs	As
30				As	Cs	Gs	Cs	Cs	As	Ts	T	
	16	B04	200	As	Ts	Cs	Ts	Gs	Ts	Gs	Gs	As
	Cs	Cs	Ts	Ts	Gs	Ts	Cs	Ts	C			
	17	B05	200	As	Cs	As	Cs	Ts	Ts	Cs	Ts	Ts

## 60

	Cs	Cs	Gs	As	Cs	Cs	Gs	Ts	G			
	18	B06	200	As	Cs	Ts	Cs	Ts	Cs	Gs	As	Cs
	As	Cs	As	Gs	Gs	As	Cs	Gs	T			
	19	B07	200	As	As	As	Cs	Cs	Cs	Cs	As	Gs
5	Ts	Ts	Cs	Gs	Ts	Cs	Ts	As	A			
	20	B08	200	As	Ts	Gs	Ts	Cs	Cs	Cs	Cs	As
	As	As	Gs	As	Cs	Ts	As	Ts	G			
	21	B09	200	As	Cs	Gs	Cs	Ts	Cs	Gs	Gs	Gs
	As	Cs	Gs	Gs	Gs	Ts	Cs	As	G			
10	22	B10	200	As	Gs	Cs	Cs	Gs	As	As	Gs	As
	As	Gs	As	Gs	Gs	Ts	Ts	As	C			
	23	B11	200	As	Cs	As	Cs	As	Gs	Ts	As	Gs
	As	Cs	Gs	As	As	As	Gs	Cs	T			
	24	B12	200	As	Cs	As	Cs	Ts	Cs	Ts	Gs	Gs
15	Ts	Ts	Ts	Cs	Ts	Gs	Gs	As	C			
	25	C01	200	As	Cs	Gs	As	Cs	Cs	As	Gs	As
	As	As	Ts	As	Gs	Ts	Ts	Ts	T			
	26	C02	200	As	Gs	Ts	Ts	As	As	As	As	Gs
	Gs	Gs	Cs	Ts	Gs	Cs	Ts	As	G			
20	27	C03	200	As	Gs	Gs	Ts	Ts	Gs	Ts	Gs	As
	Cs	Gs	As	Cs	Gs	As	Gs	Gs	T			
	28	C04	200	As	As	Ts	Gs	Ts	As	Cs	Cs	Ts
	As	Cs	Gs	Gs	Ts	Ts	Gs	Gs	C			
	29	C05	200	As	Gs	Ts	Cs	As	Cs	Gs	Ts	Cs
25	Cs	Ts	Cs	Ts	Cs	Ts	Gs	Ts	C			
	30	C06	200	Cs	Ts	Gs	Gs	Cs	Gs	As	Cs	As
	Gs	Gs	Ts	As	Gs	Gs	Ts	Cs	T			
	31	C07	200	Cs	Ts	Cs	Ts	Gs	Ts	Gs	Ts	Gs
	As	Cs	Gs	Gs	Ts	Gs	Gs	Ts	C			
30	32	C08	200	Cs	As	Gs	Gs	Ts	Cs	Gs	Ts	Cs
	Ts	Ts	Cs	Cs	Cs	Gs	Ts	Gs	G			

## 61

5	33	C09	200	Cs	Ts	Gs	Ts	Gs	Gs	Ts	As	Gs
	As	Cs	Gs	Ts	Gs	Gs	As	Cs	A			
	34	C10	200	Cs	Ts	As	As	Cs	Gs	As	Ts	Gs
	Ts	Cs	Cs	Cs	Cs	As	As	As	G			
	35	C11	200	Cs	Ts	Gs	Ts	Ts	Cs	Gs	As	Cs
10	As	Cs	Ts	Cs	Ts	Gs	Gs	Ts	T			
	36	C12	200	Cs	Ts	Gs	Gs	As	Cs	Cs	As	As
	Cs	As	Cs	Gs	Ts	Ts	Gs	Ts	C			
	37	D01	200	Cs	Cs	Gs	Ts	Cs	Cs	Gs	Ts	Gs
	Ts	Ts	Ts	Gs	Ts	Ts	Cs	Ts	G			
15	38	D02	200	Cs	Ts	Gs	As	Cs	Ts	As	Cs	As
	As	Cs	As	Gs	As	Cs	As	Cs	C			
	39	D03	200	Cs	As	As	Cs	As	Gs	As	Cs	As
	Cs	Cs	As	Gs	Gs	Gs	Gs	Ts	C			
	40	D04	200	Cs	As	Gs	Gs	Gs	Gs	Ts	Cs	Cs
20	Ts	As	Gs	Cs	Cs	Gs	As	Cs	T			
	41	D05	200	Cs	Ts	Cs	Ts	As	Gs	Ts	Ts	As
	As	As	As	Gs	Gs	Gs	Cs	Ts	G			
	42	D06	200	Cs	Ts	Gs	Cs	Ts	As	Gs	As	As
	Gs	Gs	As	Cs	Cs	Gs	As	Gs	G			
25	43	D07	200	Cs	Ts	Gs	As	As	As	Ts	Gs	Ts
	As	Cs	Cs	Ts	As	Cs	Gs	Gs	T			
	44	D08	200	Cs	As	Cs	Cs	Cs	Gs	Ts	Ts	Ts
	Gs	Ts	Cs	Cs	Gs	Ts	Cs	As	A			
	45	D09	200	Cs	Ts	Cs	Gs	As	Ts	As	Cs	Gs
30	Gs	Gs	Ts	Cs	As	Gs	Ts	Cs	A			
	46	D10	200	Gs	Gs	Ts	As	Gs	Gs	Ts	Cs	Ts
	Ts	Gs	Gs	Ts	Gs	Gs	Gs	Ts	G			
	47	D11	200	Gs	As	Cs	Ts	Ts	Ts	Gs	Cs	Cs
	Ts	Ts	As	Cs	Gs	Gs	As	As	G			
	48	D12	200	Gs	Ts	Gs	Gs	As	Gs	Ts	Cs	Ts

## 62

	Ts	Ts	Gs	Ts	Cs	Ts	Gs	Ts	G			
	49	E01	200	Gs	Gs	As	Gs	Ts	Cs	Ts	Ts	Ts
	Gs	Ts	Cs	Ts	Gs	Ts	Gs	Gs	T			
	50	E02	200	Gs	Gs	As	Cs	As	Cs	Ts	Cs	Ts
5	Cs	Gs	As	Cs	As	Cs	As	Gs	G			
	51	E03	200	Gs	As	Cs	As	Cs	As	Gs	Gs	As
	Cs	Gs	Ts	Gs	Gs	Cs	Gs	As	G			
	52	E04	200	Gs	As	Gs	Ts	As	Cs	Gs	As	Gs
	Cs	Gs	Gs	Gs	Cs	Cs	Gs	As	A			
10	53	E05	200	Gs	As	Cs	Ts	As	Ts	Gs	Gs	Ts
	As	Gs	As	Cs	Gs	Cs	Ts	Cs	G			
	54	E06	200	Gs	As	As	Gs	As	Gs	Gs	Ts	Ts
	As	Cs	As	Cs	As	Gs	Ts	As	G			
	55	E07	200	Gs	As	Gs	Gs	Ts	Ts	As	Cs	As
15	Cs	As	Gs	Ts	As	Gs	As	Cs	G			
	56	E08	200	Gs	Ts	Ts	Gs	Ts	Cs	Cs	Gs	Ts
	Cs	Cs	Gs	Ts	Gs	Ts	Ts	Ts	G			
	57	E09	200	Gs	As	Cs	Ts	Cs	Ts	Cs	Gs	Gs
	Gs	As	Cs	Cs	As	Cs	Cs	As	C			
20	58	E10	200	Gs	Ts	As	Gs	Gs	As	Gs	As	As
	Cs	Cs	As	Cs	Gs	As	Cs	Cs	A			
	59	E11	200	Gs	Gs	Ts	Ts	Cs	Ts	Ts	Cs	Gs
	Gs	Ts	Ts	Gs	Gs	Ts	Ts	As	T			
	60	E12	200	Gs	Ts	Gs	Gs	Gs	Gs	Ts	Ts	Cs
25	Gs	Ts	Cs	Cs	Ts	Ts	Gs	Gs	G			
	61	F01	200	Gs	Ts	Cs	As	Cs	Gs	Ts	Cs	Cs
	Ts	Cs	Ts	Gs	As	As	As	Ts	G			
	62	F02	200	Gs	Ts	Cs	Cs	Ts	Cs	Cs	Ts	As
	Cs	Cs	Gs	Ts	Ts	Ts	Cs	Ts	C			
30	63	F03	200	Gs	Ts	Cs	Cs	Cs	Cs	As	Cs	Gs
	Ts	Cs	Cs	Gs	Ts	Cs	Ts	Ts	C			

## 63

5	64	F04	200	Ts	Cs	As	Cs	Cs	As	Gs	Gs	As
	Cs	Gs	Gs	Cs	Gs	Gs	As	Cs	C			
	65	F05	200	Ts	As	Cs	Cs	As	As	Gs	Cs	As
	Gs	As	Cs	Gs	Gs	As	Gs	As	C			
	66	F06	200	Ts	Cs	Cs	Ts	Gs	Ts	Cs	Ts	Ts
10	Ts	Gs	As	Cs	Cs	As	Cs	Ts	C			
	67	F07	200	Ts	Gs	Ts	Cs	Ts	Ts	Ts	Gs	As
	Cs	Cs	As	Cs	Ts	Cs	As	Cs	T			
	68	F08	200	Ts	Gs	As	Cs	Cs	As	Cs	Ts	Cs
	As	Cs	Ts	Gs	As	Cs	Gs	Ts	G			
15	69	F09	200	Ts	Gs	As	Cs	Gs	Ts	Gs	Ts	Cs
	Ts	Cs	As	As	Gs	Ts	Gs	As	C			
	70	F10	200	Ts	Cs	As	As	Gs	Ts	Gs	As	Cs
	Ts	Ts	Ts	Gs	Cs	Cs	Ts	Ts	A			
	71	F11	200	Ts	Gs	Ts	Ts	Ts	As	Ts	Gs	As
20	Cs	Gs	Cs	Ts	Gs	Gs	Gs	Gs	T			
	72	F12	200	Ts	Ts	As	Ts	Gs	As	Cs	Gs	Cs
	Ts	Gs	Gs	Gs	Gs	Ts	Ts	Gs	G			
	73	G01	200	Ts	Gs	As	Cs	Gs	Cs	Ts	Gs	Gs
	Gs	Gs	Ts	Ts	Gs	Gs	As	Ts	C			
25	74	G02	200	Ts	Cs	Gs	Ts	Cs	Ts	Ts	Cs	Cs
	Cs	Gs	Ts	Gs	Gs	As	Gs	Ts	C			
	75	G03	200	Ts	Gs	Gs	Ts	As	Gs	As	Cs	Gs
	Ts	Gs	Gs	As	Cs	As	Cs	Ts	T			
	76	G04	200	Ts	Ts	Cs	Ts	Ts	Cs	Cs	Gs	As
30	Cs	Cs	Gs	Ts	Gs	As	Cs	As	T			
	77	G05	200	Ts	Gs	Gs	Ts	As	Gs	As	Cs	Gs
	Cs	Ts	Cs	Gs	Gs	Gs	As	Cs	G			
	78	G06	200	Ts	As	Gs	As	Cs	Gs	Cs	Ts	Cs
	Gs	Gs	Gs	As	Cs	Gs	Gs	Gs	T			
	79	G07	200	Ts	Ts	Ts	Ts	As	Cs	As	Gs	Ts

64

	Gs	Gs	Gs	As	As	Cs	Cs	Ts	G			
	80	G08	200	Ts	Gs	Gs	Gs	As	As	Cs	Cs	Ts
	Gs	Ts	Ts	Cs	Gs	As	Cs	As	C			
	81	G09	200	Ts	Cs	Gs	Gs	Gs	As	Cs	Cs	As
5	Cs	Cs	As	Cs	Ts	As	Gs	Gs	G			
	82	G10	200	Ts	As	Gs	Gs	As	Cs	As	As	As
	Cs	Gs	Gs	Ts	As	Gs	Gs	As	G			
	83	G11	200	Ts	Gs	Cs	Ts	As	Gs	As	As	Gs
	Gs	As	Cs	Cs	Gs	As	Gs	Gs	T			
10	84	G12	200	Ts	Cs	Ts	Gs	Ts	Cs	As	Cs	Ts
	Cs	Cs	Gs	As	Cs	Gs	Ts	Gs	G			

Table 4 is a .seq file for oligonucleotides having regions of 2'-O-(2-methoxyethyl)-nucleosides and a central region of 2'-deoxy nucleosides each linked by phosphorothioate internucleotide linkages.

15

**Table 4**

Identity of columns: **Syn #, Well, Scale, Nucleotide at particular position** (identified using base identifier followed by backbone identifier where "s" is phosphorothioate and "moe" indicated a 2'-O-(2-methoxyethyl) substituted nucleoside). The columns wrap around to next line when longer than one line.

20

	1	A01	200	moeAs	moeCs	moeCs	moeAs	Gs	Gs	As	Cs	Gs	Gs	Cs	Gs	Gs	As
				moeCs	moeCs	moeAs	moeG										
	2	A02	200	moeAs	moeCs	moeGs	moeGs	Cs	Gs	Gs	As	Cs	Cs	As	Gs	As	Gs
				moeTs	moeGs	moeGs	moeA										
25	3	A03	200	moeAs	moeCs	moeCs	moeAs	As	Gs	Cs	As	Gs	As	Cs	Gs	Gs	As
				moeGs	moeAs	moeCs	moeG										
	4	A04	200	moeAs	moeGs	moeGs	moeAs	Gs	As	Cs	Cs	Cs	Gs	As	Cs	Gs	
				moeAs	moeAs	moeCs	moeG										
	5	A05	200	moeAs	moeCs	moeCs	moeCs	Cs	Gs	As	Cs	Gs	As	As	Cs	Gs	As

moeCs moeTs moeGs moeG

6 A06 200 moeAs moeCs moeGs moeAs As Cs Gs As Cs Ts Gs Gs Cs Gs  
moeAs moeCs moeAs moeG

7 A07 200 moeAs moeCs moeGs moeAs Cs Ts Gs Gs Cs Gs As Cs As Gs  
5 moeGs moeTs moeAs moeG

8 A08 200 moeAs moeCs moeAs moeGs Gs Ts As Gs Gs Ts Cs Ts Ts Gs  
moeGs moeTs moeGs moeG

9 A09 200 moeAs moeGs moeGs moeTs Cs Ts Ts Gs Gs Ts Gs Gs Gs Ts  
moeGs moeAs moeCs moeG

10 10 A10 200 moeAs moeGs moeTs moeCs As Cs Gs As Cs As As Gs As As  
moeAs moeCs moeAs moeC

11 A11 200 moeAs moeCs moeGs moeAs Cs As As Gs As As As Cs As Cs  
moeGs moeGs moeTs moeC

12 A12 200 moeAs moeGs moeAs moeAs As Cs As Cs Gs Gs Ts Cs Gs Gs  
15 moeTs moeCs moeCs moeT

13 B01 200 moeAs moeAs moeCs moeAs Cs Gs Gs Ts Cs Gs Gs Ts Cs Cs  
moeTs moeGs moeTs moeC

14 B02 200 moeAs moeCs moeTs moeCs As Cs Ts Gs As Cs Gs Ts Gs Ts  
moeCs moeTs moeCs moeA

20 15 B03 200 moeAs moeCs moeGs moeGs As As Gs Gs As As Cs Gs Cs Cs  
moeAs moeCs moeTs moeT

16 B04 200 moeAs moeTs moeCs moeTs Gs Ts Gs Gs As Cs Cs Ts Ts Gs  
moeTs moeCs moeTs moeC

17 B05 200 moeAs moeCs moeAs moeCs Ts Ts Cs Ts Ts Cs Cs Gs As Cs  
25 moeCs moeGs moeTs moeG

18 B06 200 moeAs moeCs moeTs moeCs Ts Cs Gs As Cs As Cs As Gs Gs  
moeAs moeCs moeGs moeT

19 B07 200 moeAs moeAs moeAs moeCs Cs Cs Cs As Gs Ts Ts Cs Gs Ts  
moeCs moeTs moeAs moeA

30 20 B08 200 moeAs moeTs moeGs moeTs Cs Cs Cs Cs As As As Gs As Cs  
moeTs moeAs moeTs moeG

- 21 B09 200 moeAs moeCs moeGs moeCs Ts Cs Gs Gs Gs As Cs Gs Gs Gs  
moeTs moeCs moeAs moeG
- 22 B10 200 moeAs moeGs moeCs moeCs Gs As As Gs As As Gs As Gs Gs  
moeTs moeTs moeAs moeC
- 5 23 B11 200 moeAs moeCs moeAs moeCs As Gs Ts As Gs As Cs Gs As As  
moeAs moeGs moeCs moeT
- 24 B12 200 moeAs moeCs moeAs moeCs Ts Cs Ts Gs Gs Ts Ts Ts Cs Ts  
moeGs moeGs moeAs moeC
- 25 C01 200 moeAs moeCs moeGs moeAs Cs Cs As Gs As As As Ts As Gs  
10 moeTs moeTs moeTs moeT
- 26 C02 200 moeAs moeGs moeTs moeTs As As As As Gs Gs Gs Cs Ts Gs  
moeCs moeTs moeAs moeG
- 27 C03 200 moeAs moeGs moeGs moeTs Ts Gs Ts Gs As Cs Gs As Cs Gs  
moeAs moeGs moeGs moeT
- 15 28 C04 200 moeAs moeAs moeTs moeGs Ts As Cs Cs Ts As Cs Gs Gs Ts  
moeTs moeGs moeGs moeC
- 29 C05 200 moeAs moeGs moeTs moeCs As Cs Gs Ts Cs Cs Ts Cs Ts Cs  
moeTs moeGs moeTs moeC
- 30 C06 200 moeCs moeTs moeGs moeGs Cs Gs As Cs As Gs Gs Ts As Gs  
20 moeGs moeTs moeCs moeT
- 31 C07 200 moeCs moeTs moeCs moeTs Gs Ts Gs Ts Gs As Cs Gs Gs Ts  
moeGs moeGs moeTs moeC
- 32 C08 200 moeCs moeAs moeGs moeGs Ts Cs Gs Ts Cs Ts Ts Cs Cs Cs  
moeGs moeTs moeGs moeG
- 25 33 C09 200 moeCs moeTs moeGs moeTs Gs Gs Ts As Gs As Cs Gs Ts Gs  
moeGs moeAs moeCs moeA
- 34 C10 200 moeCs moeTs moeAs moeAs Cs Gs As Ts Gs Ts Cs Cs Cs Cs  
moeAs moeAs moeAs moeG
- 35 C11 200 moeCs moeTs moeGs moeTs Ts Cs Gs As Cs As Cs Ts Cs Ts  
30 moeGs moeGs moeTs moeT
- 36 C12 200 moeCs moeTs moeGs moeGs As Cs Cs As As Cs As Cs Gs Ts



moeTs moeGs moeTs moeC  
 37 D01 200 moeCs moeCs moeGs moeTs Cs Cs Gs Ts Gs Ts Ts Ts Gs Ts  
 moeTs moeCs moeTs moeG  
 38 D02 200 moeCs moeTs moeGs moeAs Cs Ts As Cs As As Cs As Gs As  
 5 moeCs moeAs moeCs moeC  
 39 D03 200 moeCs moeAs moeAs moeCs As Gs As Cs As Cs Cs As Gs Gs  
 moeGs moeGs moeTs moeC  
 40 D04 200 moeCs moeAs moeGs moeGs Gs Gs Ts Cs Cs Ts As Gs Cs Cs  
 moeGs moeAs moeCs moeT  
 10 41 D05 200 moeCs moeTs moeCs moeTs As Gs Ts Ts As As As As Gs Gs  
 moeGs moeCs moeTs moeG  
 42 D06 200 moeCs moeTs moeGs moeCs Ts As Gs As As Gs Gs As Cs Cs  
 moeGs moeAs moeGs moeG  
 43 D07 200 moeCs moeTs moeGs moeAs As As Ts Gs Ts As Cs Cs Ts As  
 15 moeCs moeGs moeGs moeT  
 44 D08 200 moeCs moeAs moeCs moeCs Cs Gs Ts Ts Ts Gs Ts Cs Cs Gs  
 moeTs moeCs moeAs moeA  
 45 D09 200 moeCs moeTs moeCs moeGs As Ts As Cs Gs Gs Gs Ts Cs As  
 moeGs moeTs moeCs moeA  
 20 46 D10 200 moeGs moeGs moeTs moeAs Gs Gs Ts Cs Ts Ts Gs Gs Ts Gs  
 moeGs moeGs moeTs moeG  
 47 D11 200 moeGs moeAs moeCs moeTs Ts Ts Gs Cs Cs Ts Ts As Cs Gs  
 moeGs moeAs moeAs moeG  
 48 D12 200 moeGs moeTs moeGs moeGs As Gs Ts Cs Ts Ts Ts Gs Ts Cs  
 25 moeTs moeGs moeTs moeG  
 49 E01 200 moeGs moeGs moeAs moeGs Ts Cs Ts Ts Ts Gs Ts Cs Ts Gs  
 moeTs moeGs moeGs moeT  
 50 E02 200 moeGs moeGs moeAs moeCs As Cs Ts Cs Ts Cs Gs As Cs As  
 moeCs moeAs moeGs moeG  
 30 51 E03 200 moeGs moeAs moeCs moeAs Cs As Gs Gs As Cs Gs Ts Gs Gs  
 moeCs moeGs moeAs moeG

- 52 E04 200 moeGs moeAs moeGs moeTs As Cs Gs As Gs Cs Gs Gs Gs Cs  
moeCs moeGs moeAs moeA
- 53 E05 200 moeGs moeAs moeCs moeTs As Ts Gs Gs Ts As Gs As Cs Gs  
moeCs moeTs moeCs moeG
- 5 54 E06 200 moeGs moeAs moeAs moeGs As Gs Gs Ts Ts As Cs As Cs As  
moeGs moeTs moeAs moeG
- 55 E07 200 moeGs moeAs moeGs moeGs Ts Ts As Cs As Cs As Gs Ts As  
moeGs moeAs moeCs moeG
- 56 E08 200 moeGs moeTs moeTs moeGs Ts Cs Cs Gs Ts Cs Cs Gs Ts Gs  
10 moeTs moeTs moeTs moeG
- 57 E09 200 moeGs moeAs moeCs moeTs Cs Ts Cs Gs Gs Gs As Cs Cs As  
moeCs moeCs moeAs moeC
- 58 E10 200 moeGs moeTs moeAs moeGs Gs As Gs As As Cs Cs As Cs Gs  
moeAs moeCs moeCs moeA
- 15 59 E11 200 moeGs moeGs moeTs moeTs Cs Ts Ts Cs Gs Gs Ts Ts Gs Gs  
moeTs moeTs moeAs moeT
- 60 E12 200 moeGs moeTs moeGs moeGs Gs Gs Ts Ts Cs Gs Ts Cs Cs Ts  
moeTs moeGs moeGs moeG
- 61 F01 200 moeGs moeTs moeCs moeAs Cs Gs Ts Cs Cs Ts Cs Ts Gs As  
20 moeAs moeAs moeTs moeG
- 62 F02 200 moeGs moeTs moeCs moeCs Ts Cs Cs Ts As Cs Cs Gs Ts Ts  
moeTs moeCs moeTs moeC
- 63 F03 200 moeGs moeTs moeCs moeCs Cs Cs As Cs Gs Ts Cs Cs Gs Ts  
moeCs moeTs moeTs moeC
- 25 64 F04 200 moeTs moeCs moeAs moeCs Cs As Gs Gs As Cs Gs Gs Cs Gs  
moeGs moeAs moeCs moeC
- 65 F05 200 moeTs moeAs moeCs moeCs As As Gs Cs As Gs As Cs Gs Gs  
moeAs moeGs moeAs moeC
- 66 F06 200 moeTs moeCs moeCs moeTs Gs Ts Cs Ts Ts Ts Gs As Cs Cs  
30 moeAs moeCs moeTs moeC
- 67 F07 200 moeTs moeGs moeTs moeCs Ts Ts Ts Gs As Cs Cs As Cs Ts

moeCs moeAs moeCs moeT  
 68 F08 200 moeTs moeGs moeAs moeCs Cs As Cs Ts Cs As Cs Ts Gs As  
 moeCs moeGs moeTs moeG  
 69 F09 200 moeTs moeGs moeAs moeCs Gs Ts Gs Ts Cs Ts Cs As As Gs  
 5 moeTs moeGs moeAs moeC  
 70 F10 200 moeTs moeCs moeAs moeAs Gs Ts Gs As Cs Ts Ts Ts Gs Cs  
 moeCs moeTs moeTs moeA  
 71 F11 200 moeTs moeGs moeTs moeTs Ts As Ts Gs As Cs Gs Cs Ts Gs  
 moeGs moeGs moeGs moeT  
 10 72 F12 200 moeTs moeTs moeAs moeTs Gs As Cs Gs Cs Ts Gs Gs Gs Gs  
 moeTs moeTs moeGs moeG  
 73 G01 200 moeTs moeGs moeAs moeCs Gs Cs Ts Gs Gs Gs Gs Ts Ts Gs  
 moeGs moeAs moeTs moeC  
 74 G02 200 moeTs moeCs moeGs moeTs Cs Ts Ts Cs Cs Cs Gs Ts Gs Gs  
 15 moeAs moeGs moeTs moeC  
 75 G03 200 moeTs moeGs moeGs moeTs As Gs As Cs Gs Ts Gs Gs As Cs  
 moeAs moeCs moeTs moeT  
 76 G04 200 moeTs moeTs moeCs moeTs Ts Cs Cs Gs As Cs Cs Gs Ts Gs  
 moeAs moeCs moeAs moeT  
 20 77 G05 200 moeTs moeGs moeGs moeTs As Gs As Cs Gs Cs Ts Cs Gs Gs  
 moeGs moeAs moeCs moeG  
 78 G06 200 moeTs moeAs moeGs moeAs Cs Gs Cs Ts Cs Gs Gs Gs As Cs  
 moeGs moeGs moeGs moeT  
 79 G07 200 moeTs moeTs moeTs moeTs As Cs As Gs Ts Gs Gs Gs As As  
 25 moeCs moeCs moeTs moeG  
 80 G08 200 moeTs moeGs moeGs moeGs As As Cs Cs Ts Gs Ts Ts Cs Gs  
 moeAs moeCs moeAs moeC  
 81 G09 200 moeTs moeCs moeGs moeGs Gs As Cs Cs As Cs Cs As Cs Ts  
 moeAs moeGs moeGs moeG  
 30 82 G10 200 moeTs moeAs moeGs moeGs As Cs As As As Cs Gs Gs Ts As  
 moeGs moeGs moeAs moeG

83 G11 200 moeTs moeGs moeCs moeTs As Gs As As Gs Gs As Cs Cs Gs  
 moeAs moeGs moeGs moeT  
 84 G12 200 moeTs moeCs moeTs moeGs Ts Cs As Cs Ts Cs Cs Gs As Cs  
 moeGs moeTs moeGs moeG

## 5 Reagent file (.tab File)

Table 5 is a .tab file for reagents necessary for synthesizing an oligonucleotides having both 2'-O-(2-methoxyethyl)nucleosides and 2'-deoxy nucleosides located therein.

**Table 5**

Identity of columns: **GroupName, Bottle ID, ReagentName, FlowRate, Concentration.**

10 Wherein reagent name is identified using base identifier, "moe" indicated a 2'-O-(2-methoxyethyl) substituted nucleoside and "cpg" indicates a control pore glass solid support medium. The columns wrap around to next line when longer than one line.

### SUPPORT

BEGIN

15           0       moeG       moeG       cpg   100   1  
             0       moe5meC   moe5meC   cpg   100   1  
             0       moeA       moeA       cpg   100   1  
             0       moeT       moeT       cpg   100   1  
             END

### 20 DEBLOCK

BEGIN

70           TCA       TCA                   100   1  
             END

### WASH

25           BEGIN  
             65       ACN       ACN           190   1

71

END

## OXIDIZERS

BEGIN

68	BEAU	BEAUCAGE	320	1
----	------	----------	-----	---

5

END

## CAPPING

BEGIN

66	CAP_B	CAP_B	220	1
----	-------	-------	-----	---

67	CAP_A	CAP_A	230	1
----	-------	-------	-----	---

10

END

## DEOXY THIOATE

BEGIN

31,32	Gs	deoxyG	270	1
-------	----	--------	-----	---

39,40	5meCs	5methyldeoxyC	270	1
-------	-------	---------------	-----	---

15

37,38	As	deoxyA	270	1
-------	----	--------	-----	---

29,30	Ts	deoxyT	270	1
-------	----	--------	-----	---

END

## MOE-THIOATE

20

BEGIN

15,16	moeGs	methoxyethoxyG	240	1
-------	-------	----------------	-----	---

23,24	moe5meCs	methoxyethoxyC	240	1
-------	----------	----------------	-----	---

21,22	moeAs	methoxyethoxyA	240	1
-------	-------	----------------	-----	---

13,14	moeTs	methoxyethoxyT	240	1
-------	-------	----------------	-----	---

25

END

## ACTIVATORS

BEGIN

5,6,7,8	SET	s-ethyl-tet	280	
---------	-----	-------------	-----	--

Activates

DEOXY\_THIOATE

MOE\_THIOATE

END

#### 5      **EXAMPLE 4: Oligonucleotide Synthesis - 96 Well Plate Format**

Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry using a multi well automated synthesizer utilizing input files as described in EXAMPLE 3 above. The oligonucleotides were synthesized by assembling 96 sequences simultaneously in a standard 96 well format. Phosphodiester internucleotide linkages were  
10      afforded by oxidation with aqueous iodine. Phosphorothioate internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE/ABI, Pharmacia). Non-standard nucleosides are synthesized as per known literature or patented  
15      methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

Following synthesis, oligonucleotides were cleaved from support and deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature (55-60°C) for 12-16 hours and the released product then dried *in vacuo*. The dried product was then re-suspended in sterile  
20      water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic pipettors.

#### **EXAMPLE 5: Alternative Oligonucleotide Synthesis**

Unsubstituted and substituted phosphodiester oligonucleotides are alternately synthesized on an automated DNA synthesizer (Applied Biosystems model 380B) using  
25      standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates are synthesized as per the phosphodiester oligonucleotides except the standard oxidation bottle was replaced by 0.2 M solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the stepwise thiation of the phosphite linkages. The thiation wait step was increased to 68 sec and was followed by the capping

step. After cleavage from the CPG column and deblocking in concentrated ammonium hydroxide at 55°C (18 hr), the oligonucleotides were purified by precipitating twice with 2.5 volumes of ethanol from a 0.5 M NaCl solution.

Phosphinate oligonucleotides are prepared as described in U.S. Patent 5,508,270,  
5 incorporated herein by reference in its entirety.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, incorporated herein by reference in its entirety.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are prepared as described in U.S. Patents 5,610,289 or 5,625,050, each of which is incorporated herein by reference in  
10 its entirety.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, incorporated herein by reference in its entirety.

Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093  
15 and WO 94/02499, respectively), each of which is incorporated herein by reference in its entirety.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are prepared as described in U.S. Patent 5,476,925, incorporated herein by reference in its entirety.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent  
20 5,023,243, incorporated herein by reference in its entirety.

Boranophosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, each of which is incorporated herein by reference in its entirety.

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as  
25 MDH linked oligonucleosides, and methylenecarbonylamino linked oligonucleosides, also identified as amide-3 linked oligonucleosides, and methyleneaminocarbonyl linked oligonucleosides, also identified as amide-4 linked oligonucleosides, as well as mixed backbone compounds having, for instance, alternating MMI and PO or PS linkages are prepared as described in U.S. Patents 5,378,825; 5,386,023; 5,489,677; 5,602,240 and 5,610,289, each  
30 of which is incorporated herein by reference in its entirety.

Formacetal and thioformacetal linked oligonucleosides are prepared as described in

U.S. Patents 5,264,562 and 5,264,564, each of which is incorporated herein by reference in its entirety.

Ethylene oxide linked oligonucleosides are prepared as described in U.S. Patent 5,223,618, incorporated herein by reference in its entirety.

#### 5      **EXAMPLE 6: PNA Synthesis**

Peptide nucleic acids (PNAs) are prepared in accordance with any of the various procedures referred to in Peptide Nucleic Acids (PNA): Synthesis, Properties and Potential Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5. They may also be prepared in accordance with U.S. Patents 5,539,082; 5,700,922, and 5,719,262, each of which is  
10      incorporated herein by reference in its entirety.

#### **EXAMPLE 7: Chimeric Oligonucleotide Synthesis**

Chimeric oligonucleotides, oligonucleosides or mixed oligonucleotides/oligonucleosides of the invention can be of several different types. These include a first type wherein the "gap" segment of linked nucleosides is positioned between  
15      5' and 3' "wing" segments of linked nucleosides and a second "open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound. Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers."

##### 20      **A.      [2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric Phosphorothioate Oligonucleotides**

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidites for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidites for 5' and 3' wings. The standard synthesis cycle is  
25      modified by increasing the wait step after the delivery of tetrazole and base to 600 s repeated four times for DNA and twice for 2'-O-methyl. The fully protected oligonucleotide was cleaved from the support and the phosphate group is deprotected in 3:1 Ammonia/Ethanol at room temperature overnight then lyophilized to dryness.



Treatment in methanolic ammonia for 24 hrs at room temperature is done to deprotect all bases and the samples are again lyophilized to dryness.

**B. [2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(2-Methoxyethyl)]**

**Chimeric Phosphorothioate Oligonucleotides**

5 [2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(2-methoxyethyl)] chimeric phosphorothioate oligonucleotides are prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-(2-methoxyethyl) amidites for the 2'-O-methyl amidites.

**C. [2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy Phosphorothioate]--[**

10 **[2'-O-(2-Methoxyethyl) Phosphodiester] Chimeric Oligonucleotide**

[2'-O-(2-methoxyethyl phosphodiester)]--[2'-deoxy phosphorothioate]--[2'-O-(2-methoxyethyl) phosphodiester] chimeric oligonucleotides are prepared as per the above procedure for the 2'-O-methyl chimeric oligonucleotide with the substitution of 2'-O-(2-methoxyethyl) amidites for the 2'-O-methyl amidites in the wing portions. Sulfurization utilizing 3,4-dihydro-2H-benzothio-3-one 1,1 dioxide (Beaucage Reagent) is used to generate the phosphorothioate internucleotide linkages within the wing portions of the chimeric structures. Oxidization with iodine is used to generate the phosphodiester internucleotide linkages for the center gap.

15 Other chimeric oligonucleotides, chimeric oligonucleosides and mixed chimeric oligonucleotides/oligonucleosides are synthesized according to United States Patent 5,623,065, which is incorporated herein by reference in its entirety.

**EXAMPLE 8: Output Oligonucleotides From Automated Oligonucleotide Synthesis**

Using the .seq files, the .cmd files and .tab file of Example 3, oligonucleotides were prepared as per the protocol of the 96 well format of Example 4. The  
25 oligonucleotides were prepared utilizing phosphorothioate chemistry to give in one instance a first library of phosphorothioate oligodeoxynucleotides. The oligonucleotides were prepared in a second instance as a second library of hybrid oligonucleotides having phosphorothioate backbones with a first and third "wing" region of 2'-O-(2-methoxyethyl)nucleotides on either side of a center gap region of 2'-deoxy nucleotides.  
30 The two libraries contained the same set of oligonucleotide sequences. Thus the two

libraries are redundant with respect to sequence but are unique with respect to the combination of sequence and chemistry. Because the sequences of the second library of compounds is the same as the first (however the chemistry is different), for brevity sake, the second library is not shown.

5 For illustrative purposes Tables 6-a and 6-b show the sequences of an initial first library, i.e., a library of phosphorothioate oligonucleotides targeted to a CD40 target. The compounds of Table 6-a shows the members of this library listed in compliance with the established rule for listing SEQ ID NO:, i.e., in numerical SEQ ID NO: order.

**Table 6-a**

10 **Sequences of Oligonucleotides Targeted to CD40 by SEQ ID NO.:**

	NUCLEOBASE SEQUENCE	SEQ ID NO.
	CCAGGCGGCAGGACCACT	1
	GACCAGGCGGCAGGACCA	2
	AGGTGAGACCAGGCGGCA	3
15	CAGAGGCAGACGAACCAT	4
	GCAGAGGCAGACGAACCA	5
	GCAAGCAGCCCCAGAGGA	6
	GGTCAGCAAGCAGCCCCA	7
	GACAGCGGTCAGCAAGCA	8
20	GATGGACAGCGGTCAGCA	9
	TCTGGATGGACAGCGGTC	10
	GGTGGTTCTGGATGGACA	11
	GTGGGTGGTTCTGGATGG	12
	GCAGTGGGTGGTTCTGGA	13
25	CACAAAGAACAGCACTGA	14
	CTGGCACAAAGAACAGCA	15
	TCCTGGCTGGCACAAAGA	16
	CTGTCCTGGCTGGCACAA	17
	CTCACCAGTTTCTGTCCT	18
30	TCACTCACCAGTTTCTGT	19

	GTGCAGTCACTCACCAGT	20
	ACTCTGTGCAGTCACTCA	21
	CAGTGAAGTCTGTGCAGT	22
	ATTCCGTTTCAGTGAAGT	23
5	GAAGGCATTCCGTTTCAG	24
	TTCACCGCAAGGAAGGCA	25
	CTCTGTTCCAGGTGTCTA	26
	CTGGTGGCAGTGTGTCTC	27
	TGGGGTCGCAGTATTTGT	28
10	GGTTGGGGTCGCAGTATT	29
	CTAGGTTGGGGTCGCAGT	30
	GGTGCCCTTCTGCTGGAC	31
	CTGAGGTGCCCTTCTGCT	32
	GTGTCTGTTTCTGAGGTG	33
15	TGGTGTCTGTTTCTGAGG	34
	ACAGGTGCAGATGGTGTC	35
	TTCACAGGTGCAGATGGT	36
	GTGCCAGCCTTCTTCACA	37
	TACAGTGCCAGCCTTCTT	38
20	GGACACAGCTCTCACAGG	39
	TGCAGGACACAGCTCTCA	40
	GAGCGGTGCAGGACACAG	41
	AAGCCGGGCGAGCATGAG	42
	AATCTGCTTGACCCCAAA	43
25	GAAACCCCTGTAGCAATC	44
	GTATCAGAAACCCCTGTA	45
	GCTCGCAGATGGTATCAG	46
	GCAGGGCTCGCAGATGGT	47
	TGGGCAGGGCTCGCAGAT	48
30	GACTGGGCAGGGCTCGCA	49
	CATTGGAGAAGAAGCCGA	50

	GATGACACATTGGAGAAG	51
	GCAGATGACACATTGGAG	52
	TCGAAAGCAGATGACACA	53
	GTCCAAGGGTGACATTTT	54
5	CACAGCTTGTCCAAGGGT	55
	TTGGTCTCACAGCTTGTC	56
	CAGGTCTTTGGTCTCACA	57
	CTGTTGCACAACCAGGTC	58
	GTTTGTGCCTGCCTGTTG	59
10	GTCTTGTTTGTGCCTGCC	60
	CCACAGACAACATCAGTC	61
	CTGGGGACCACAGACAAC	62
	TCAGCCGATCCTGGGGAC	63
	CACCACCAGGGCTCTCAG	64
15	GGGATCACCACCAGGGCT	65
	GAGGATGGCAAACAGGAT	66
	ACCAGCACCAAGAGGATG	67
	TTTTGATAAAGACCAGCA	68
	TATTGGTTGGCTTCTTGG	69
20	GGGTTCCCTGCTTGGGGTG	70
	GTCGGGAAAATTGATCTC	71
	GATCGTCGGGAAAATTGA	72
	GGAGCCAGGAAGATCGTC	73
	TGGAGCCAGGAAGATCGT	74
25	TGGAGCAGCAGTGTTGGA	75
	GTAAAGTCTCCTGCACTG	76
	TGGCATCCATGTAAAGTC	77
	CGGTTGGCATCCATGTAA	78
	CTCTTTGCCATCCTCCTG	79
30	CTGTCTCTCCTGCACTGA	80
	GGTGCAGCCTCACTGTCT	81

AACTGCCTGTTTGCCAC	82
CTTCTGCCTGCACCCCTG	83
ACTGACTGGGCATAGCTC	84

5           The sequences shown in Table 6-a, above, and Table 6-b, below, are in a 5' to 3' direction. This is reversed with respect to 3' to 5' direction shown in the .seq files of Example 3. For synthesis purposes, the .seq files are generated reading from 3' to 5'. This allows for aligning all of the 3' most "A" nucleosides together, all of the 3' most "G" nucleosides together, all of the 3' most "C" nucleosides together and all of the 3' most "T" nucleosides together. Thus when the first nucleoside of each particular oligonucleotide (attached to the solid support) is added to the wells on the plates, machine movement is reduced since an automatic pipette can move in a linear manner down one row and up another on the 96 well plate.

15           The location of the well holding each particular oligonucleotides is indicated by row and column. There are eight rows designated A to H and twelve columns designated 1 to 12 in a typical 96 well format plate. Any particular well location is indicated by its "Well No." which is indicated by the combination of the row and the column, e.g. A08 is the well at row A, column 8.

20           In Table 6-b below, the oligonucleotides of Table 6-a are shown reordered according to the Well No. on their synthesis plate. The order shown in Table 6-b is the actually order as synthesized on an automated synthesizer taking advantage of the preferred placement of the first nucleoside according to the above alignment criteria.

**Table 6-b:**

25           **Sequences of Oligonucleotides Targeted to CD40 Order by Synthesis Well No.**

Well No.		SEQ ID NO:
A01	GACCAGGCGGCAGGACCA	2
A02	AGGTGAGACCAGGCGGCA	3
A03	GCAGAGGCAGACGAACCA	5
A04	GCAAGCAGCCCCAGAGGA	6
A05	GGTCAGCAAGCAGCCCCA	7
A06	GACAGCGGTCAGCAAGCA	8
A07	GATGGACAGCGGTCAGCA	9
A08	GGTGGTTCTGGATGGACA	11

	A09	GCAGTGGGTGGTTCTGGA	13
	A10	CACAAAGAACAGCACTGA	14
	A11	CTGGCACAAAGAACAGCA	15
	A12	TCCTGGCTGGCACAAAGA	16
5	B01	CTGTCCTGGCTGGCACAA	17
	B02	ACTCTGTGCAGTCACTCA	21
	B03	TTCACCGCAAGGAAGGCA	25
	B04	CTCTGTTCCAGGTGTCTA	26
	B05	GTGCCAGCCTTCTTCACA	37
10	B06	TGCAGGACACAGCTCTCA	40
	B07	AATCTGCTTGACCCCAA	43
	B08	GTATCAGAAACCCCTGTA	45
	B09	GACTGGGCAGGGCTCGCA	49
	B10	CATTGGAGAAGAAGCCGA	50
15	B11	TCGAAAGCAGATGACACA	53
	B12	CAGGTCTTTGGTCTCACA	57
	C01	TTTTGATAAAGACCAGCA	68
	C02	GATCGTCGGGAAAATTGA	72
	C03	TGGAGCAGCAGTGTTGGA	75
20	C04	CGGTTGGCATCCATGTAA	78
	C05	CTGTCTCTCCTGCACTGA	80
	C06	TCTGGATGGACAGCGGTC	10
	C07	CTGGTGGCAGTGTGTCTC	27
	C08	GGTGCCCTTCTGCTGGAC	31
25	C09	ACAGGTGCAGATGGTGTC	35
	C10	GAAACCCCTGTAGCAATC	44
	C11	TTGGTCTCACAGCTTGTC	56
	C12	CTGTTGCACAACCAGGTC	58
	D01	GTCTTGTTTGTGCCTGCC	60
30	D02	CCACAGACAACATCAGTC	61
	D03	CTGGGGACCACAGACAAC	62
	D04	TCAGCCGATCCTGGGGAC	63
	D05	GTCGGGAAAATTGATCTC	71
	D06	GGAGCCAGGAAGATCGTC	73
35	D07	TGGCATCCATGTAAAGTC	77
	D08	AACTGCCTGTTTGCCAC	82
	D09	ACTGACTGGGCATAGCTC	84
	D10	GTGGGTGGTTCTGGATGG	12
	D11	GAAGGCATTCCGTTTCAG	24
40	D12	GTGTCTGTTTCTGAGGTG	33
	E01	TGGTGTCTGTTTCTGAGG	34
	E02	GGACACAGCTCTCACAGG	39
	E03	GAGCGGTGCAGGACACAG	41
	E04	AAGCCGGGCGAGCATGAG	42
45	E05	GCTCGCAGATGGTATCAG	46
	E06	GATGACACATTGGAGAAG	51
	E07	GCAGATGACACATTGGAG	52
	E08	GTTTGTGCCTGCCTGTTG	59
	E09	CACCACCAGGGCTCTCAG	64
50	E10	ACCAGCACCAAGAGGATG	67
	E11	TATTGGTTGGCTTCTTGG	69

	E12	GGGTCCTGCTTGGGGTG	70
	F01	GTAAAGTCTCCTGCACTG	76
	F02	CTCTTTGCCATCCTCCTG	79
	F03	CTTCTGCCTGCACCCCTG	83
5	F04	CCAGGCGGCAGGACCACT	1
	F05	CAGAGGCAGACGAACCAT	4
	F06	CTCACCAGTTTCTGTCCT	18
	F07	TCACTCACCAGTTTCTGT	19
	F08	GTGCAGTCACTCACCAGT	20
10	F09	CAGTGAACCTCTGTGCAGT	22
	F10	ATTCCGTTTCAGTGAAC	23
	F11	TGGGGTCGCAGTATTTGT	28
	F12	GGTTGGGGTCGCAGTATT	29
	G01	CTAGGTTGGGGTCGCAGT	30
15	G02	CTGAGGTGCCCTTCTGCT	32
	G03	TTCACAGGTGCAGATGGT	36
	G04	TACAGTGCCAGCCTTCTT	38
	G05	GCAGGGCTCGCAGATGGT	47
	G06	TGGGCAGGGCTCGCAGAT	48
20	G07	GTCCAAGGGTGACATTTT	54
	G08	CACAGCTTGTCCAAGGGT	55
	G09	GGGATCACCACCAGGGCT	65
	G10	GAGGATGGCAAACAGGAT	66
	G11	TGGAGCCAGGAAGATCGT	74
25	G12	GGTGCAGCCTCACTGTCT	81

### EXAMPLE 9: Oligonucleotide Analysis

#### A. Oligonucleotide Analysis - 96 Well Plate Format

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96 well format (Beckman MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors.

#### B. Alternative Oligonucleotide Analysis

After cleavage from the controlled pore glass support (Applied Biosystems) and deblocking in concentrated ammonium hydroxide at 55°C for 18 hours, the oligonucleotides or oligonucleosides are purified by precipitation twice out of 0.5 M NaCl with 2.5 volumes ethanol. Synthesized oligonucleotides are analyzed by polyacrylamide gel electrophoresis on denaturing gels. Oligonucleotide purity is checked by <sup>31</sup>P nuclear

magnetic resonance spectroscopy, and/or by HPLC, as described by Chiang *et al.*, *J. Biol. Chem.* 1991, 266, 18162.

#### EXAMPLE 10: Automated Assay of CD40 Oligonucleotide Activity

##### 5           A. Poly(A)+ mRNA isolation.

Poly(A)+ mRNA was isolated according to Miura *et al.* (*Clin. Chem.*, 1996, 42, 1758). Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ l cold PBS. 60  $\mu$ l lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was added to each well, the plate was gently agitated and then incubated at room temperature for five minutes. 55  $\mu$ l of lysate was transferred to Oligo d(T) coated 96 well plates (AGCT Inc., Irvine, CA). Plates were incubated for 60 minutes at room temperature, washed 3 times with 200  $\mu$ l of wash buffer (10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the final wash, the plate was blotted on paper towels to remove excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ l of elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C was added to each well, the plate was incubated on a 90°C plate for 5 minutes, and the eluate then transferred to a fresh 96-well plate. Cells grown on 100 mm or other standard plates may be treated similarly, using appropriate volumes of all solutions.

##### 20           B. Total RNA isolation

Total mRNA was isolated using an RNEASY 96-well kit and buffers purchased from Qiagen Inc. (Valencia CA) following the manufacturer's recommended procedures. Briefly, for cells grown on 96-well plates, growth medium was removed from the cells and each well was washed with 200  $\mu$ l cold PBS. 100  $\mu$ l Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 100  $\mu$ l of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96-well plate attached to a QIAVAC manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 15 seconds. 1  $\mu$ l of Buffer RW1 was added to each well of the RNEASY 96-well plate and the vacuum again applied for 15 seconds. 1  $\mu$ l of Buffer RPE was then added to each well of the RNEASY 96-well plate and the vacuum applied for a period of 15 seconds. The



Buffer RPE wash was then repeated and the vacuum was applied for an additional 10 minutes. The plate was then removed from the QIAVACÔ manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVACÔ manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by  
5 pipetting 60 mL water into each well, incubating 1 minute, and then applying the vacuum for 30 seconds. The elution step was repeated with an additional 60 mL water.

### C. RT-PCR Analysis of CD40 mRNA Levels

Quantitation of CD40 mRNA levels was determined by reverse transcriptase  
10 polymerase chain reaction (RT-PCR) using the ABI PRISM™\_7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in real-time.

15 As opposed to standard PCR, in which amplification products are quantitated after the PCR is completed, products in RT-PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., JOE or FAM, PE-Applied Biosystems, Foster City, CA) is  
20 attached to the 5' end of the probe and a quencher dye (e.g., TAMRA, PE-Applied Biosystems, Foster City, CA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension  
25 phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated.

With each cycle, additional reporter dye molecules are cleaved from their respective probes, and the fluorescence intensity is monitored at regular (six-second)  
30 intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from

untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

RT-PCR reagents were obtained from PE-Applied Biosystems, Foster City, CA. RT-PCR reactions were carried out by adding 25 ml PCR cocktail (1x TAQMAN™ buffer A, 5.5 mM MgCl<sub>2</sub>, 300 mM each of dATP, dCTP and dGTP, 600 mM of dUTP, 100 nM each of forward primer, reverse primer, and probe, 20 U RNase inhibitor, 1.25 units AMPLITAQ GOLD™, and 12.5 U MuLV reverse transcriptase) to 96 well plates containing 25 ml poly(A) mRNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. following a 10 minute incubation at 95°C to activate the AMPLITAQ GOLD™, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

For CD40, the PCR primers were:

forward: 5' CAGAGTTCACCTGAAACGGAATGC 3'  
(SEQ ID NO:86)  
reverse: 5' GGTGGCAGTGTGTCTCTCTGTTC 3' (SEQ ID NO:87), and  
PCR probe: 5' FAM-TTCCTTGCGGTGAAAGCGAATTCCT-TAMRA 3' (SEQ ID NO:88) where *FAM* (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye and *TAMRA* (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

For GAPDH, the PCR primers were:

forward: 5' GAAGGTGAAGGTCGGAGTC 3' (SEQ ID NO:89)  
reverse: 5' GAAGATGGTGATGGGATTTTC 3' (SEQ ID NO:90), and  
PCR probe: 5' JOE-CAAGCTTCCCGTTCTCAGCC-TAMRA 3' (SEQ ID NO. 91)  
where *JOE* (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye and *TAMRA* (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

#### **EXAMPLE 11: Inhibition of CD40 Expression by Phosphorothioate Oligodeoxynucleotides**

In accordance with the present invention, a series of oligonucleotides complementary to mRNA were designed to target different regions of the human CD40 mRNA, using published sequences (GenBank accession number X60592, incorporated herein by reference as SEQ ID NO: 85). The oligonucleotides are shown in Table 7.

Target sites are indicated by the beginning nucleotide numbers, as given in the sequence source reference (X60592), to which the oligonucleotide binds. All compounds in Table 7 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. Data are averages from three experiments.

5

Table 7:

## Inhibition of CD40 mRNA Levels by Phosphorothioate Oligodeoxynucleotides

TARGET		SEQ ID		
ISIS#	SITE	SEQUENCE	% INHIB.	NO.
10	18	CCAGGCGGCAGGACCA	30.71	1
	20	GACCAGGCGGCAGGAC	28.09	2
	26	AGGTGAGACCAGGCGG	21.89	3
	48	CAGAGGCAGACGAACC	0.00	4
	49	GCAGAGGCAGACGAAC	0.00	5
15	73	GCAAGCAGCCCCAGAG	0.00	6
	78	GGTCAGCAAGCAGCCC	29.96	7
	84	GACAGCGGTCAGCAAG	0.00	8
	88	GATGGACAGCGGTCAG	0.00	9
	92	TCTGGATGGACAGCGG	0.00	10
20	98	GGTGGTTCTGGATGGA	0.00	11
	101	GTGGGTGGTTCTGGAT	0.00	12
	104	GCAGTGGGTGGTTCTG	0.00	13
	152	CACAAAGAACAGCACT	0.00	14
	156	CTGGCACAAAGAACAG	0.00	15
25	162	TCCTGGCTGGCACAAA	0.00	16
	165	CTGTCTGGCTGGCAC	4.99	17
	176	CTCACCAGTTTCTGTCC	0.00	18
	179	TCACTCACCAGTTTCTG	0.00	19
	185	GTGCAGTCACTACCA	0.00	20
30	190	ACTCTGTGCAGTCACTC	0.00	21
	196	CAGTGAACCTGTGCA	5.30	22
	205	ATTCCGTTTCAGTGAAC	0.00	23
	211	GAAGGCATTCCGTTTC	9.00	24
	222	TTCACCGCAAGGAAGG	0.00	25
35	250	CTCTGTTCCAGGTGTCT	0.00	26
	267	CTGGTGGCAGTGTGTC	0.00	27
	286	TGGGGTCGCAGTATTT	0.00	28
	289	GGTTGGGGTCGCAGTA	0.00	29
	292	CTAGGTTGGGGTCGCA	0.00	30
40	318	GGTGCCCTTCTGCTGG	19.67	31
	322	CTGAGGTGCCCTTCTGC	15.63	32
	332	GTGTCTGTTTCTGAGGT	0.00	33
	334	TGGTGTCTGTTTCTGAG	0.00	34
	345	ACAGGTGCAGATGGTG	0.00	35
	348	TTCACAGGTGCAGATG	0.00	36

## 86

	18659	360	GTGCCAGCCTTCTTCAC	5.67	37
	18660	364	TACAGTGCCAGCCTTCT	7.80	38
	18661	391	GGACACAGCTCTCACA	0.00	39
	18662	395	TGCAGGACACAGCTCT	0.00	40
5	18663	401	GAGCGGTGCAGGACAC	0.00	41
	18664	416	AAGCCGGGCGAGCATG	0.00	42
	18665	432	AATCTGCTTGACCCCA	5.59	43
	18666	446	GAAACCCCTGTAGCAA	0.10	44
	18667	452	GTATCAGAAACCCCTG	0.00	45
10	18668	463	GCTCGCAGATGGTATC	0.00	46
	18669	468	GCAGGGCTCGCAGATG	34.05	47
	18670	471	TGGGCAGGGCTCGCAG	0.00	48
	18671	474	GACTGGGCAGGGCTCG	2.71	49
	18672	490	CATTGGAGAAGAAGCC	0.00	50
15	18673	497	GATGACACATTGGAGA	0.00	51
	18674	500	GCAGATGACACATTGG	0.00	52
	18675	506	TCGAAAGCAGATGACA	0.00	53
	18676	524	GTCCAAGGGTGACATT	8.01	54
	18677	532	CACAGCTTGTCCAAGG	0.00	55
20	18678	539	TTGGTCTCACAGCTTGT	0.00	56
	18679	546	CAGGTCTTTGGTCTCAC	6.98	57
	18680	558	CTGTTGCACAACCAGG	18.76	58
	18681	570	GTTTGTGCCTGCCTGTT	2.43	59
	18682	575	GTCTTGTTTGTGCCTGC	0.00	60
25	18683	590	CCACAGACAACATCAG	0.00	61
	18684	597	CTGGGGACCACAGACA	0.00	62
	18685	607	TCAGCCGATCCTGGGG	0.00	63
	18686	621	CACCACCAGGGCTCTC	23.31	64
	18687	626	GGGATCACCACCAGGG	0.00	65
30	18688	657	GAGGATGGCAAACAGG	0.00	66
	18689	668	ACCAGCACCAAGAGGA	0.00	67
	18690	679	TTTTGATAAAGACCAG	0.00	68
	18691	703	TATTGGTTGGCTTCTTG	0.00	69
	18692	729	GGGTTCTCTGCTTGGGG	0.00	70
35	18693	750	GTCGGGAAAATTGATC	0.00	71
	18694	754	GATCGTCGGGAAAATT	0.00	72
	18695	765	GGAGCCAGGAAGATCG	0.00	73
	18696	766	TGGAGCCAGGAAGATC	0.00	74
	18697	780	TGGAGCAGCAGTGTTG	0.00	75
40	18698	796	GTAAAGTCTCCTGCAC	0.00	76
	18699	806	TGGCATCCATGTAAAG	0.00	77
	18700	810	CGGTTGGCATCCATGT	0.00	78
	18701	834	CTCTTTGCCATCCTCCT	4.38	79
	18702	861	CTGTCTCTCCTGCACTG	0.00	80
45	18703	873	GGTGCAGCCTCACTGT	0.00	81
	18704	910	AACTGCCTGTTTGCCCA	33.89	82
	18705	954	CTTCTGCCTGCACCCCT	0.00	83
	18706	976	ACTGACTGGGCATAGC	0.00	84

As shown in Table 7, SEQ ID NOS: 1, 2, 7, 47 and 82 demonstrated at least 25% inhibition of CD40 expression and are therefore preferred compounds of the invention.

### EXAMPLE 12: Inhibition of CD40 Expression by Phosphorothioate 2'-MOE

#### Gapmer Oligonucleotides

- 5 In accordance with the present invention, a second series of oligonucleotides complementary to mRNA were designed to target different regions of the human CD40 mRNA, using published sequence X60592. The oligonucleotides are shown in Table 8. Target sites are indicated by the beginning or initial nucleotide numbers, as given in the sequence source reference (X60592), to which the oligonucleotide binds.
- 10 All compounds in Table 8 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings." The wings are composed of 2'-O-(2-methoxyethyl) (2'-MOE) nucleotides. The intersugar (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide.
- 15 Cytidine residues in the 2'-MOE wings are 5-methylcytidines. Data are averaged from three experiments.

**Table 8:**

#### Inhibition of CD40 mRNA Levels by Chimeric Phosphorothioate Oligonucleotides

	ISIS#	TARGET SEQUENCE	% Inhibition	SEQ ID
20	19211	18 CCAGGCGGCAGGACCA	75.71	1
	19212	20 GACCAGGCGGCAGGA	77.23	2
	19213	26 AGGTGAGACCAGGCG	80.82	3
	19214	48 CAGAGGCAGACGAAC	23.68	4
	19215	49 GCAGAGGCAGACGAA	45.97	5
25	19216	73 GCAAGCAGCCCCAGAG	65.80	6
	19217	78 GGTCAGCAAGCAGCCC	74.73	7
	19218	84 GACAGCGGTCAGCAAG	67.21	8
	19219	88 GATGGACAGCGGTCAG	65.14	9
	19220	92 TCTGGATGGACAGCGG	78.71	10
30	19221	98 GGTGGTTCTGGATGGA	81.33	11
	19222	101 GTGGGTGGTTCTGGAT	57.79	12
	19223	104 GCAGTGGGTGGTTCTG	73.70	13
	19224	152 CACAAAGAACAGCACT	40.25	14
	19225	156 CTGGCACAAAGAACAG	60.11	15

	19226	162	TCCTGGCTGGCACAAA	10.18	16
	19227	165	CTGTCCTGGCTGGCAC	24.37	17
	19228	176	CTCACCAGTTTCTGTC	22.30	18
	19229	179	TCACTCACCAGTTTCT	40.64	19
5	19230	185	GTGCAGTCACTCACCA	82.04	20
	19231	190	ACTCTGTGCAGTCACT	37.59	21
	19232	196	CAGTGA ACTCTGTGCA	40.26	22
	19233	205	ATTCCGTTTCAGTGAA	56.03	23
	19234	211	GAAGGCATTCCGTTTC	32.21	24
10	19235	222	TTCACCGCAAGGAAGG	61.03	25
	19236	250	CTCTGTTCCAGGTGTCT	62.19	26
	19237	267	CTGGTGGCAGTGTGTC	70.32	27
	19238	286	TGGGGTCGCAGTATTT	0.00	28
	19239	289	GGTTGGGGTCGCAGTA	19.40	29
15	19240	292	CTAGGTTGGGGTCGCA	36.32	30
	19241	318	GGTGCCCTTCTGCTGG	78.91	31
	19242	322	CTGAGGTGCCCTTCTG	69.84	32
	19243	332	GTGTCTGTTTCTGAGG	63.32	33
	19244	334	TGGTGTCTGTTTCTGA	42.83	34
20	19245	345	ACAGGTGCAGATGGTG	73.31	35
	19246	348	TTCACAGGTGCAGATG	47.72	36
	19247	360	GTGCCAGCCTTCTTCA	61.32	37
	19248	364	TACAGTGCCAGCCTTC	46.82	38
	19249	391	GGACACAGCTCTCACA	0.00	39
25	19250	395	TGCAGGACACAGCTCT	52.05	40
	19251	401	GAGCGGTGCAGGACAC	50.15	41
	19252	416	AAGCCGGGCGAGCATG	32.36	42
	19253	432	AATCTGCTTGACCCCA	0.00	43
	19254	446	GAAACCCCTGTAGCAA	0.00	44
30	19255	452	GTATCAGAAACCCCTG	36.13	45
	19256	463	GCTCGCAGATGGTATC	64.65	46
	19257	468	GCAGGGCTCGCAGATG	74.95	47
	19258	471	TGGGCAGGGCTCGCAG	0.00	48
	19259	474	GACTGGGCAGGGCTCG	82.00	49
35	19260	490	CATTGGAGAAGAAGCC	41.31	50
	19261	497	GATGACACATTGGAGA	13.81	51
	19262	500	GCAGATGACACATTGG	78.48	52
	19263	506	TCGAAAGCAGATGACA	59.28	53
	19264	524	GTCCAAGGGTGACATT	70.99	54
40	19265	532	CACAGCTTGTCCAAGG	0.00	55
	19266	539	TTGGTCTCACAGCTTG	45.92	56
	19267	546	CAGGTCTTTGGTCTCA	63.95	57
	19268	558	CTGTTGCACAACCAGG	82.32	58
	19269	570	GTTTGTGCCTGCCTGTT	70.10	59
45	19270	575	GTCTTGTTTGTGCCTGC	68.95	60
	19271	590	CCACAGACAACATCAG	11.22	61
	19272	597	CTGGGGACCACAGACA	9.04	62
	19273	607	TCAGCCGATCCTGGGG	0.00	63
	19274	621	CACCACCAGGGCTCTC	23.08	64
50	19275	626	GGGATCACCACCAGGG	57.94	65
	19276	657	GAGGATGGCAAACAG	49.14	66

	19277	668	ACCAGCACCAAGAGG	3.48	67
	19278	679	TTTGTATAAAGACCAG	30.58	68
	19279	703	TATTGGTTGGCTTCTTG	49.26	69
	19280	729	GGGTTCTGCTTGGGG	13.95	70
5	19281	750	GTCGGGAAAATTGATC	54.78	71
	19282	754	GATCGTCGGGAAAATT	0.00	72
	19283	765	GGAGCCAGGAAGATC	69.47	73
	19284	766	TGGAGCCAGGAAGATC	54.48	74
	19285	780	TGGAGCAGCAGTGTTG	15.17	75
10	19286	796	GTAAAGTCTCCTGCAC	30.62	76
	19287	806	TGGCATCCATGTAAAG	65.03	77
	19288	810	CGGTGGCATCCATGT	34.49	78
	19289	834	CTCTTGCCATCCTCCT	41.84	79
	19290	861	CTGTCTCTCCTGCACT	25.68	80
15	19291	873	GGTGCAGCCTCACTGT	76.27	81
	19292	910	AACTGCCTGTTTGCCC	63.34	82
	19293	954	CTTCTGCCTGCACCCC	0.00	83
	19294	976	ACTGACTGGGCATAGC	11.55	84

20 As shown in Table 8, SEQ ID NOS: 1, 2, 3, 6, 7, 8, 9, 10, 11, 12, 13, 15, 20, 23, 25, 26, 27, 31, 32, 33, 35, 37, 40, 41, 46, 47, 49, 52, 53, 54, 57, 58, 59, 60, 65, 71, 73, 74, 77, 81 and 82 demonstrated at least 50% inhibition of CD40 expression and are therefore preferred compounds of the invention.

### 25 **EXAMPLE 13: Oligonucleotide-Sensitive Sites of the CD40 Target Nucleic Acid**

As the data presented in the preceding two Examples shows, several sequences were present in preferred compounds of two distinct oligonucleotide chemistries. Specifically, compounds having SEQ ID NOS: 1, 2, 7, 47 and 82 are preferred in both instances. These compounds map to different regions of the CD40 transcript but  
30 nevertheless define accessible sites of the target nucleic acid.

For example, SEQ ID NOS: 1 and 2 overlap each other and both map to the 5'-untranslated region (5'-UTR) of CD40. Accordingly, this region of CD40 is particularly preferred for modulation via sequence-based technologies. Similarly, SEQ ID NOS: 7 and 47 map to the open reading frame of CD40, whereas SEQ ID NO: 82 maps to the 3'-  
35 untranslated region (3'-UTR). Thus, the ORF and 3'-UTR of CD40 may be targeted by sequence-based technologies as well.

The reverse complements of the active CD40 compounds are easily determined by those skilled in the art and may be assembled to yield nucleotide sequences corresponding

to accessible sites on the target nucleic acid. For example, the assembled reverse complement of SEQ ID NOS: 1 and 2 is represented below as SEQ ID NO:92:

5'- AGTGGTCCTGCCGCCTGGTC -3' SEQ ID NO:92

TCACCAGGACGGCGGACC -5' SEQ ID NO:1

5 ACCAGGACGGCGGACCAG -5' SEQ ID NO:2

10 Through multiple iterations of the process of the invention, more extensive “footprints” are generated. A library of this information is compiled and may be used by those skilled in the art in a variety of sequence-based technologies to study the molecular and biological functions of CD40 and to investigate or confirm its role in various diseases and disorders.

#### EXAMPLE 14: Site Selection Program

15 In a preferred embodiment of the invention, illustrated in Figure 20, an application is deployed which facilitates the selection process for determining the target positions of the oligos to be synthesized, or “sites.” This program is written using a three-tiered object-oriented approach. All aspects of the software described, therefore, are tightly integrated with the relational database. For this reason, explicit database read and write steps are not shown. It should be assumed that each step described includes database access. The description below illustrates one way the program can be used. The actual  
20 interface allows users to skip from process to process at will, in any order.

Before running the site picking program, the target must have all relevant properties computed as described previously and indicated in process step 2204. When the site picking program is launched at process step 2206 the user is presented with a panel showing targets which have previously been selected and had their properties calculated.  
25 The user selects one target to work with at process step 2208 and proceeds to decide if any derived properties will be needed at process step 2210. Derived properties are calculated by performing mathematical operations on combinations of pre-calculated properties as defined by the user at process step 2212.

30 The derived properties are made available as peers with all the pre-calculated properties. The user selects one of the properties to view plotted versus target position at process step 2214. This graph is shown above a linear representation of the target. The



horizontal or position axis of both the graph and target are linked and scalable by the user. The zoom range goes from showing the full target length to showing individual target bases as letters and individual property points. The user next selects a threshold value below or above which all sites will be eliminated from future consideration at process step **2216**. The user decides whether to eliminate more sites based on any other properties at process step **2218**. If they choose to eliminate more, they return to pick another property to display at process step **2214** and threshold at process step **2216**.

After eliminating sites, the user selects from the remaining list by choosing any property at process step **2220** and then choosing a manual or automatic selection technique at process step **2222**. In the automatic technique, the user decides whether they want to pick from maxima or minima and the number of maxima or minima to be selected as sites at process step **2224**. The software automatically finds and picks the points. When picking manually the user must decide if they wish to use automatic peak finding at process step **2226**. If the user selects automatic peak finding, then user must click on the graphed property with the mouse at process step **2236**. The nearest maxima or minima, depending on the modifier key held down, to the selected point will be picked as the site. Without the peak finding option, the user must pick a site at process step **2238** by clicking on its position on the linear representation of target.

Each time a site, or group of sites, is picked, a dynamic property is calculated for all possible sites (not yet eliminated) at process step **2230**. This property indicates the nearness of the site to a picked site allowing the user to pick sites in subsequent iterations based on target coverage. After new sites are picked, the user determines if the desired number of sites has been picked. If too few sites have been picked the user returns to pick more **2220**. If too many sites have been picked, the user may eliminate them by selecting and deleting them on the target display at process step **2234**. If the correct number of sites is picked, and the user is satisfied with the set of picked sites, the user registers these sites to the database along with their name, notebook number, and page number at process step **2238**. The database time stamps this registration event.

**EXAMPLE 15: Site Selection Program**

In a preferred embodiment of the invention, illustrated in Figure 21, an application is deployed which facilitates the assignment of specific chemical structure to the complement of the sequence of the sites previously picked and facilitates the registration and ordering of these now fully defined antisense compounds. This program is written using a three-tiered object-oriented approach. All aspects of the software described, therefore, are tightly integrated with the relational database. For this reason, explicit database read and write steps are not shown, it being understood that each step described also includes appropriate database read/write access.

To begin using the oligonucleotide chemistry assignment program, the user launches it at process step **2302**. The user then selects from the previously selected sets of oligonucleotides at process step **2304**, registered to the database in site picker's process step **2238**. Next, the user must decide whether to manually assign the chemistry a base at a time, or run the sites through a template at process step **2306**. If the user chooses to use a template, they must determine if a desired template is available at process step **2308**. If a template is not available with the desired chemistry modifications and the correct length, the user can define one at process step **2314**.

To define a template, the user must select the length of the oligonucleotide the template is to define. This oligonucleotide is then represented as a bar with selectable regions. The user sets the number of regions on the oligonucleotide, and the positions and lengths of these regions by dragging them back and forth on the bar. Each region is represented by a different color.

For each region, the user defines the chemistry modifications for the sugars, the linkers, and the heterocycles at each base position in the region. At least four heterocycle chemistries must be given, one for each of the four possible base types (A, G, C or T or U) in the site sequence the template will be applied to. A user interface is provided to select these chemistries which show the molecular structure of each component selected and its modification name. By pushing on a pop-up list next to each of the pictures, the user may choose from a list of structures and names, those possible to put in this place. For example, the heterocycle that represents the base type G is shown as a two dimensional structure diagram. If the user clicks on the pop-up list, a row of other possible structures

and names is shown. The user drags the mouse to the desired chemistry and releases the mouse. Now the newly selected molecule is displayed as the choice for G type heterocycle modifications.

Once the user has created a template, or selected an existing one, the software  
5 applies the template at process step **2312** to each of the complements of the sites in the list. When the templates are applied, it is possible that chemistries will be defined which are impossible to make with the chemical precursors presently used on the automatic synthesizer. To check this, a database is maintained of all precursors previously designed, and their availability for automated synthesis. When the templates are applied, the  
10 resulting molecules are tested at process step **2316** against this database to see if they are readily synthesized.

If a molecule is not readily synthesized, it is added to a list that the user inspects. At process step **2318**, the user decides whether to modify the chemistry to make it compatible with the currently recognized list of available chemistries or to ignore it. To  
15 modify a chemistry, the user must use the base at a time interface at process step **2322**. The user can also choose to go directly to this step, bypassing templates all together at process step **2306**.

The base at a time interface at process step **2322** is very similar to the template editor at process step **2314** except that instead of specifying chemistries for regions, they  
20 are defined one base at a time. This interface also differs in that it dynamically checks to see if the design is readily synthesized as the user makes selections. In other words, each choice made limits the choices the software makes available on the pop-up selection lists. To accommodate this function, an additional choice is made available on each pop-up of "not defined." For example, this allows the user to inhibit linker choice from restricting  
25 the sugar choices by first setting the linker to "not defined." The user would then pick the sugar, and then pick from the remaining linker choices available.

Once all of the sites on the list are assigned chemistries or dropped, they are registered at process step **2324** to a commercial chemical structure database. Registering to this database makes sure the structure is unique, assigns it a new identifier if it is  
30 unique, and allows future structure and substructure searching by creating various hash-tables. The compound definition is also stored at process step **2326** to various hash tables

referred to as chemistry/position tables. These allow antisense compound searching and categorization based on oligonucleotide chemistry modification sequences and equivalent base sequences.

The results of the registration are displayed at process step **2328** with the new IDs if they are new compounds and with the old IDs if they have been previously registered. The user next selects which of the compounds processed they wish to order for synthesis at process step **2330** and registers an order list at process step **2332** by including scientist name, notebook number and page number. The database time-stamps this entry. The user may then choose at process step **2334**, to quit the program at process step **2338**, go back to the beginning and choose a new site list to work with process step **2304**, or start the oligonucleotide ordering interface at process step **2336**.

#### EXAMPLE 16: Gene Walk to Optimize Oligonucleotide Sequence

A gene walk is executed using a CD40 antisense oligonucleotide having SEQ ID NO:15 (5'-CTGGCACAAAGAACAGCA-3'). In effecting this gene walk, the following parameters are used:

Gene Walk Parameter	Entered value
Oligonucleotide Sequence ID:	15
Name of Gene Target:	CD40
Scope of Gene Walk:	20
Sequence Shift Increment:	1

Entering these values and effecting the gene walk centered on SEQ ID NO: 15 automatically generates the following new oligonucleotides:

**Table 9:**

**Oligonucleotide Generated By Gene Walk**

SEQ ID	Sequence
93	GAACAGCACTGACTG
94	AGAACAGCACTGACT
95	AAGAACAGCACTGAC
96	AAAGAACAGCACTGA
97	CAAAGAACAGCACTG

5	98	ACAAAGAACAGCACT
	14	CACAAAGAACAGCAC
	100	GCACAAAGAACAGCA
	101	GGCACAAAGAACAGC
	102	TGGCACAAAGAACAG
	15	CTGGCACAAAGAACA
	103	GCTGGCACAAAGAAC
	104	GGCTGGCACAAAGAA
10	105	TGGCTGGCACAAAGA
	106	CTGGCTGGCACAAAG
	107	CCTGGCTGGCACAAA
	16	TCCTGGCTGGCACAA
	109	GTCCTGGCTGGCACA
	110	TGTCCTGGCTGGCACA
15	17	CTGTCCTGGCTGGCAC
	112	TCTGTCCTGGCTGGCA

The list shown above contains 20 oligonucleotide sequences directed against the CD40 nucleic acid sequence. They are ordered by the position along the CD40 sequence at which the 5' terminus of each oligonucleotide hybridizes. Thus, the first ten oligonucleotides are single-base frame shift sequences directed against the CD40 sequence upstream of compound SEQ ID NO: 15 and the latter ten are single-base frame shift sequences directed against the CD40 sequence downstream of compound SEQ ID NO: 15.

#### 25 **EXAMPLE 17: Automated Assay of RhoC Oligonucleotide Activity**

RhoC, a member of the Rho subfamily of small GTPases, is a protein that has been shown to be involved in a diverse set of signaling pathways including the ultimate regulation of the dynamic organization of the cytoskeleton.

30 Oligonucleotides were designed as described in Example 2, synthesized as described in Examples 3 through 8, analyzed as described in Examples 9 and assayed as described in Example 10 except for target specific primer and probes.

RhoC probes and primers were designed to hybridize to the human RhoC sequence, using published sequence information (GenBank accession number L25081, incorporated herein by reference as SEQ ID NO:113).

35 For RhoC the PCR primers were:

forward primer: TGATGTCATCCTCATGTGCTTCT (SEQ ID NO: 114)

reverse primer: CCAGGATGATGGGCACGTT (SEQ ID NO: 115) and the PCR probe was: FAM-CGACAGCCCTGACAGCCTGGAAA-TAMRA (SEQ ID NO: 116) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

5

**EXAMPLE 18: Antisense inhibition of RhoC expression- phosphorothioate oligodeoxynucleotides**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human RhoC RNA, using published sequences (GenBank accession number L25081, incorporated herein by reference as SEQ ID NO: 113). The oligonucleotides are shown in Table 10. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. L25081), to which the oligonucleotide binds. All compounds in Table 10 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds were analyzed for effect on RhoC mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from three experiments. If present, "N.D." indicates "no data".

**Table 10**

**Inhibition of RhoC mRNA levels by phosphorothioate oligodeoxynucleotides**

ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
25304	5' UTR	4	gagctgagatgaagtcaa	29	117
25305	5' UTR	44	gctgaagttcccaggctg	25	118
25306	5' UTR	47	ccggctgaagttcccagg	42	119
25307	Coding	104	ggcaccatccccaacgat	81	120
25308	Coding	105	aggcaccatccccaacga	81	121
25309	Coding	111	tcccacaggcaccatccc	70	122
25310	Coding	117	aggtcttcccacaggcac	40	123
25311	Coding	127	atgaggaggcaggtcttc	41	124
25312	Coding	139	ttgctgaagacgatgagg	23	125
25313	Coding	178	tcaaagacagtagggacg	0	126
25314	Coding	181	ttctcaaagacagtaggg	2	127
25315	Coding	183	agttctcaaagacagtag	38	128
25316	Coding	342	tgtttccaggctgtcag	59	129

## 97

	25317	Coding	433	tcgtcttcgctcaggtcc	79	130
	25318	Coding	439	gtgtgctcgtcttgctc	67	131
	25319	Coding	445	ctcctggtgtgctcgtct	67	132
	25320	Coding	483	cagaccgaacgggctcct	65	133
5	25321	Coding	488	ttcctcagaccgaacggg	57	134
	25322	Coding	534	actcaaggtagccaaagg	33	135
	25323	Coding	566	ctcccgactccctcctt	91	136
	25324	Coding	575	ctcaaacacctcccgac	34	137
	25325	Coding	581	ggccatctcaaacacctc	64	138
10	25326	Coding	614	cttggtcttgcgacctg	72	139
	25327	Coding	625	cccctccgacgtgttgc	66	140
	25328	3' UTR	737	gtatggagccctcaggag	60	141
	25329	3' UTR	746	gagccttcagtatggagc	63	142
	25330	3' UTR	753	gaaaatggagccttcagt	24	143
15	25331	3' UTR	759	ggaactgaaaatggagcc	2	144
	25332	3' UTR	763	ggagggaactgaaaatgg	13	145
	25333	3' UTR	766	gcaggagggaactgaaaa	27	146
	25334	3' UTR	851	agggcagggcataggcgt	31	147
	25335	3' UTR	854	ggaagggcagggcatagg	21	148
20	25336	3' UTR	859	catgaggaagggcagggc	0	149
	25337	3' UTR	920	taaagtgtgtgtgtga	39	150
	25338	3' UTR	939	cctgtgagccagaagtgt	69	151
	25339	3' UTR	941	ttcctgtgagccagaagt	69	152
	25340	3' UTR	945	cactttcctgtgagccag	82	153
25	25341	3' UTR	948	agacactttcctgtgagc	69	154
	25342	3' UTR	966	actctgggtccctactgc	20	155
	25343	3' UTR	992	tgcagaaacaactccagg	0	156

**Example 19: Antisense inhibition of RhoC expression- phosphorothioate 2'-MOE****30 gapmer oligonucleotides**

In accordance with the present invention, a second series of oligonucleotides targeted to human RhoC were synthesized. The oligonucleotide sequences are shown in Table 11. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession No. L25081), to which the oligonucleotide binds.

35 All compounds in Table 11 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE) nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the

oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from three experiments. If present, "N.D." indicates "no data".

5

Table 11

**Inhibition of RhoC mRNA levels by chimeric phosphorothioate oligonucleotides  
having 2'-MOE wings and a deoxy gap**

	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
10	25344	5' UTR	4	gagctgagatgaagtcaa	0	117
	25345	5' UTR	44	gctgaagttcccaggctg	35	118
	25346	5' UTR	47	ccggctgaagtcccagg	53	119
	25347	Coding	104	ggcaccatccccaacgat	50	120
	25348	Coding	105	aggcaccatccccaacga	56	121
15	25349	Coding	111	tcccacaggcaccatccc	4	122
	25350	Coding	117	aggtcttcccacaggcac	11	123
	25351	Coding	127	atgaggaggcaggcttc	6	124
	25352	Coding	139	ttgctgaagacgatgagg	15	125
	25353	Coding	178	tcaaagacagtagggacg	32	126
20	25354	Coding	181	ttctcaaagacagtaggg	7	127
	25355	Coding	183	agttctcaaagacagtag	39	128
	25356	Coding	342	tgtttccaggctgtcag	59	129
	25357	Coding	433	tcgtcttcctcaggctcc	48	130
	25358	Coding	439	gtgtgctcgtcttgctc	36	131
25	25359	Coding	445	ctcctggtgtgctcgtct	61	132
	25360	Coding	483	cagaccgaacgggctcct	50	133
	25361	Coding	488	ttcctcagaccgaacggg	14	134
	25362	Coding	534	actcaaggtagccaaagg	32	135
	25363	Coding	566	ctcccgcactccctcctt	21	136
30	25364	Coding	575	ctcaaacacctcccgcac	9	137
	25365	Coding	581	ggccatctcaaacacctc	66	138
	25366	Coding	614	ctgttcttgccgacctg	61	139
	25367	Coding	625	cccctccgacgcttgctc	0	140
	25368	3' UTR	737	gtatggagccctcaggag	28	141
35	25369	3' UTR	746	gagccttcagtatggagc	32	142
	25370	3' UTR	753	gaaaatggagccttcagt	0	143
	25371	3' UTR	759	ggaactgaaaatggagcc	40	144
	25372	3' UTR	763	ggagggaactgaaaatgg	45	145
	25373	3' UTR	766	gcaggagggaactgaaaa	35	146
40	25374	3' UTR	851	agggcagggcataggcgt	5	147
	25375	3' UTR	854	ggaagggcagggcataagg	0	148



25376	3' UTR	859	catgaggaagggcagggc	0	149
25377	3' UTR	920	taaagtgcgtggtgtgtga	20	150
25378	3' UTR	939	cctgtgagccagaagtgt	67	151
25379	3' UTR	941	ttcctgtgagccagaagt	61	152
5 25380	3' UTR	945	cactttcctgtgagccag	80	153
25381	3' UTR	948	agacactttcctgtgagc	0	154
25382	3' UTR	966	actctgggtccctactgc	0	155
25383	3' UTR	992	tcagaaacaactccagg	0	156

## 10 **EXAMPLE 20: Automated Assay of Cellular Inhibitor of Apoptosis-2 Expression Oligonucleotide Activity**

Cellular Inhibitor of Apoptosis-2 (also known as c-IAP-2, apoptosis inhibitor 2, API-2, hIAP-1, and MIHC) is a member of the inhibitor of apoptosis (IAP) family of anti-apoptotic proteins which interfere with the transmission of intracellular death signals.

15 Oligonucleotides were designed as described in Example 2, synthesized as described in Examples 3 through 8, analyzed as described in Examples 9 and assayed as described in Example 10 except for target specific primer and probes. Cellular Inhibitor of Apoptosis-2 probes and primers were designed to hybridize to the human Cellular Inhibitor of Apoptosis-2 sequence, using published sequence information (GenBank  
20 accession number U37546, incorporated herein by reference as SEQ ID NO:157).

For Cellular Inhibitor of Apoptosis-2 the PCR primers were:

forward primer: GGAATCAGGTGTTGGGAATCTG (SEQ ID NO: 158)

reverse primer: CAAGTACTCACACCTTGGAAACCA (SEQ ID NO: 159) and the PCR probe was: FAM-AGATGATCCATGGGTTCAACATGCCAA-TAMRA (SEQ ID NO:

25 160) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

## **EXAMPLE 21: Antisense inhibition of Cellular Inhibitor of Apoptosis-2 expression-phosphorothioate oligodeoxynucleotides**

30 In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human Cellular Inhibitor of Apoptosis-2 RNA, using published sequences (GenBank accession number U37546, incorporated herein by reference as SEQ ID NO: 157). The oligonucleotides are shown in Table 12. Target sites

are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. U37546), to which the oligonucleotide binds. All compounds in Table 12 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds were analyzed for effect on Cellular Inhibitor of Apoptosis-2 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from three experiments. If present, "N.D." indicates "no data".

Table 12

**Inhibition of Cellular Inhibitor of Apoptosis-2 mRNA levels by phosphorothioate oligodeoxynucleotides**

	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
15	23412	5' UTR	3	actgaagacattttgaat	62	161
	23413	5' UTR	37	cttagaggtagctaaaat	29	162
	23414	5' UTR	49	gcacttttatttcttaga	70	163
	23415	5' UTR	62	attttaattagaagcact	0	164
	23416	5' UTR	139	accatatttcactgattc	70	165
20	23417	5' UTR	167	ctaactcaaaggaggaaa	0	166
	23418	5' UTR	175	cacaagacctaactcaaa	27	167
	23419	5' UTR	268	gctctgctgtcaagtgtt	57	168
	23420	5' UTR	303	tgtgtgactcatgaagct	23	169
	23421	5' UTR	335	ttcagtggcattcaatca	23	170
25	23422	5' UTR	357	cttctccaggctactaga	50	171
	23423	5' UTR	363	ggtcaacttctccaggct	65	172
	23424	5' UTR	437	taaaacccttcacagaag	0	173
	23425	5' UTR	525	ttaaggaagaatacaca	0	174
	23426	5' UTR	651	gcatggctttgcttttat	0	175
30	23427	Coding	768	caaacgtgttggegttt	35	176
	23428	Coding	830	agcaggaaaagtgaata	0	177
	23429	Coding	1015	ttaacggaatttagactc	0	178
	23430	Coding	1064	atttggtactgaagaagg	0	179
	23431	Coding	1118	agagccacggaaatatcc	9	180
35	23432	Coding	1168	aatcttgatttgctctg	7	181
	23433	Coding	1231	gtaagtaacttgccattt	0	182
	23434	Coding	1323	agcaagccactctgtctc	50	183
	23435	Coding	1436	tgaagtgtcttgaagctg	0	184
	23436	Coding	1580	tttgacatcatcactgtt	0	185
	23437	Coding	1716	tggttgaaacttgacgga	0	186
	23438	Coding	1771	tcattctctgggctgtct	40	187

## 101

	23439	Coding	1861	gcagcattaatcacagga	0	188
	23440	Coding	2007	tttctctctctcttccc	10	189
	23441	Coding	2150	aacatcatgttctgttc	9	190
	23442	Coding	2273	atataacacagcttcagc	0	191
5	23443	Coding	2350	aattgttcttccactggt	0	192
	23444	Coding	2460	aagaaggagcacaatctt	70	193
	23445	3' UTR	2604	gaaaccaaattaggataa	12	194
	23446	3' UTR	2753	tgtagtgtctacctcttt	69	195
	23447	3' UTR	2779	ctgaaattttgattgaat	14	196
10	23448	3' UTR	2795	tacaatttcaataatgct	38	197
	23449	3' UTR	2920	gggtctcagtatgctgcc	21	198
	23450	3' UTR	3005	ccttcgatgtataggaca	0	199
	23451	3' UTR	3040	catgtccctaaaatgtca	0	200

15

**EXAMPLE 22: Antisense inhibition of Cellular Inhibitor of Apoptosis-2 expression-phosphorothioate 2'-MOE gapmer oligonucleotides**

In accordance with the present invention, a second series of oligonucleotides targeted to human Cellular Inhibitor of Apoptosis-2 were synthesized. The oligonucleotide sequences are shown in Table 13. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. U37546), to which the oligonucleotide binds.

All compounds in Table 13 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from three experiments. If present, "N.D." indicates "no data".

Table 13

**Inhibition of Cellular Inhibitor of Apoptosis-2 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap**

5	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
	23452	5' UTR	3	actgaagacattttgaat	35	161
	23453	5' UTR	37	cttagaggtagctaaaat	26	162
	23454	5' UTR	49	gcacttttatttcttaga	76	163
10	23455	5' UTR	62	attttaattagaagcact	0	164
	23456	5' UTR	139	accatatttactgattc	0	165
	23457	5' UTR	167	ctaactcaaaggaggaaa	5	166
	23458	5' UTR	175	cacaagacctaactcaaa	0	167
	23459	5' UTR	268	gctctgctgtcaagtgtt	57	168
15	23460	5' UTR	303	tgtgtgactcatgaagct	67	169
	23461	5' UTR	335	ttcagtggcattcaatca	59	170
	23462	5' UTR	357	cttctccaggctactaga	0	171
	23463	5' UTR	363	ggccaacttctccaggct	75	172
	23464	5' UTR	437	taaaacccttcacagaag	11	173
20	23465	5' UTR	525	ttaaggaagaaatacaca	0	174
	23466	5' UTR	651	gcatggctttgcttttat	46	175
	23467	Coding	768	caaactgtgtggcgcttt	47	176
	23468	Coding	830	agcaggaaaagtgaata	39	177
	23469	Coding	1015	ttaacggaatttagactc	12	178
25	23470	Coding	1064	atttggtactgaagaagg	34	179
	23471	Coding	1118	agagccacggaaatatacc	54	180
	23472	Coding	1168	aaatcttgatttgcctctg	34	181
	23473	Coding	1231	gtaagtaatctggcattt	0	182
	23474	Coding	1323	agcaagccactctgtctc	42	183
30	23475	Coding	1436	tgaagtgtcttgaagctg	0	184
	23476	Coding	1580	tttgacatcatcactgtt	57	185
	23477	Coding	1716	tggttggaacttgacgga	23	186
	23478	Coding	1771	tcctctctgggctgtct	66	187
	23479	Coding	1861	gcagcattaatcacagga	65	188
35	23480	Coding	2007	tttctctctctcttccc	0	189
	23481	Coding	2150	aacatcatgttcttgttc	13	190
	23482	Coding	2273	atataacacagcttcagc	0	191
	23483	Coding	2350	aattgttcttcactggt	60	192
	23484	Coding	2460	aagaaggagcacaaatctt	65	193
40	23485	3' UTR	2604	gaaaccaaattaggataa	0	194
	23486	3' UTR	2753	tgtagtgtacctctttt	73	195
	23487	3' UTR	2779	ctgaaatttgattgaat	4	196

23488	3' UTR	2795	tacaatttcaataatgct	0	197
23489	3' UTR	2920	gggtctcagtatgctgcc	42	198
23490	3' UTR	3005	ccttcgatgtataggaca	71	199
23491	3' UTR	3040	catgtccctaaaatgtca	45	200

5

**EXAMPLE 23: Automated Assay of ELK-1 Oligonucleotide Activity**

ELK-1 (also known as p62TCF) is a member of the ternary complex factor (TCF) subfamily of Ets domain proteins and utilizes a bipartite recognition mechanism mediated by both protein-DNA and protein-protein interactions. This results in gene regulation not only by direct DNA binding but also by indirect DNA binding through recruitment by other factors (Rao et al., *Science*, **1989**, 244, 66-70). The formation of ternary complexes with an array of proteins allows the differential regulation of many genes. The mechanism by which ELK-1 controls various signal transduction pathways involves regulating the activity of the Egr-1, pip92, nur77 and c-fos promoters by binding to the serum response element (SRE) in these promoters in response to extracellular stimuli such as growth factors, mitogens and oncogene products (Sharrocks et al., *Int. J. Biochem. Cell Biol.*, **1997**, 29, 1371-1387). ELK-1 has also been shown to mediate other functions within the cell including apoptosis.

Oligonucleotides were designed as described in Example 2, synthesized as described in Examples 3 through 8, analyzed as described in Examples 9 and assayed as described in Example 10 except for target specific primer and probes. ELK-1 probes and primers were designed to hybridize to the human ELK-1 sequence, using published sequence information (GenBank accession number M25269, incorporated herein by reference as SEQ ID NO:201).

For ELK-1 the PCR primers were:  
 forward primer: GCAAGGCAATGGCCACAT (SEQ ID NO: 202)  
 reverse primer: CTCCTCTGCATCCACCAGCTT (SEQ ID NO: 203) and the PCR probe was: FAM-TCTCCTGGACTTCACGGGATGGTGGT-TAMRA (SEQ ID NO: 204) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

30

**EXAMPLE 24: Antisense inhibition of ELK-1 expression-phosphorothioate oligodeoxynucleotides**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human ELK-1 RNA, using published sequences (GenBank accession number M25269, incorporated herein by reference as SEQ ID NO: 201). The oligonucleotides are shown in Table 14. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. M25269), to which the oligonucleotide binds. All compounds in Table 14 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds were analyzed for effect on ELK-1 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from three experiments. If present, "N.D." indicates "no data".

**Table 14**

**Inhibition of ELK-1 mRNA levels by phosphorothioate oligodeoxynucleotides**

ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
20	24752	5' UTR 11	cccctgcgtttccctaca	15	205
	24753	5' UTR 50	ggtggtggtggcggtggc	29	206
	24754	5' UTR 139	ggcgttggcaatgttggc	82	207
	24755	5' UTR 167	aagttgaggtgtgtgta	0	208
	24756	5' UTR 189	aggccacggacgggtctc	92	209
	24757	5' UTR 229	gattgattcgctacgatg	71	210
25	24758	5' UTR 255	gggatgcggaggagtgcg	74	211
	24759	5' UTR 289	agtgtcacgccatcca	22	212
	24760	Coding 328	aaactgccacagcgtcac	64	213
	24761	Coding 381	gaagtcaggagatgatg	62	214
	24762	Coding 395	caccaccatcccgtgaag	88	215
30	24763	Coding 455	tctgttcttgcgtagtc	62	216
	24764	Coding 512	tgttcttgcatagtagt	52	217
	24765	Coding 527	tcacctgcggatgatgt	57	218
	24766	Coding 582	gagcacctgcgacctca	72	219
	24767	Coding 600	ggcgggcagtcctcagtg	82	220
35	24768	Coding 787	ggtgaaggtggaatagag	58	221
	24769	Coding 993	tccgatttcaggtttggg	55	222
	24770	Coding 1110	ttggtggtttctggcaca	67	223

## 105

	24771	Coding	1132	tggagggacttctggctc	69	224
	24772	Coding	1376	gcgtaggaagcagggatg	34	225
	24773	Coding	1440	gtgctccagaagtgaatg	64	226
	24774	Coding	1498	actggatggaaactggaa	34	227
5	24775	Coding	1541	ggccatccacgctgatag	74	228
	24776	3' UTR	1701	ccaccacaatcagagcat	74	229
	24777	3' UTR	1711	gatccccaccccaccaca	16	230
	24778	3' UTR	1765	tgtttctgtggaggaga	48	231
	24779	3' UTR	1790	aaacagagaagttgtgga	11	232
10	24780	3' UTR	1802	gggactgacagaaaacag	0	233
	24781	3' UTR	1860	ataaataaataaacgcc	18	234
	24782	3' UTR	1894	gttaggtcaggctcatcc	56	235
	24783	3' UTR	1974	gttctcaagccagacctc	52	236
	24784	3' UTR	1992	aataaagaaagaaaggtc	41	237
15	24785	3' UTR	2006	agggcaggctgagaaata	29	238
	24786	3' UTR	2053	cttctactcacatccaaa	54	239
	24787	3' UTR	2068	caaaacaaactaactctt	24	240
	24788	3' UTR	2080	ggaataataaaacaaaac	40	241
	24789	3' UTR	2107	ttcttctggaccctga	93	242
20	24790	3' UTR	2161	ccaagggtgtgattcttc	81	243
	24791	3' UTR	2200	tgtctgagagaaaggttg	55	244

**EXAMPLE 25: Antisense inhibition of ELK-1 expression- phosphorothioate 2'-MOE gapmer oligonucleotides**

25 In accordance with the present invention, a second series of oligonucleotides targeted to human ELK-1 were synthesized. The oligonucleotide sequences are shown in Table 15. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. M25269), to which the oligonucleotide binds.

30 All compounds in Table 15 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

35 Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from three experiments. If present, "N.D." indicates "no data".

Table 15

**Inhibition of ELK-1 mRNA levels by chimeric phosphorothioate oligonucleotides  
having 2'-MOE wings and a deoxy gap**

	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
5	24792	5' UTR	11	cccctgcgtttccctaca	23	205
	24793	5' UTR	50	ggtggtggtggcggtggc	80	206
	24794	5' UTR	139	ggcgttggcaatgttggc	91	207
	24795	5' UTR	167	aagttgaggctgtgtga	27	208
10	24796	5' UTR	189	aggccacggacgggtctc	79	209
	24797	5' UTR	229	gattgattcgctacgatg	69	210
	24798	5' UTR	255	gggatgcggaggagtgcg	42	211
	24799	5' UTR	289	agtgtcacgccatccca	45	212
	24800	Coding	328	aaactgccacagcgtcac	57	213
15	24801	Coding	381	gaagtcaggagatgatg	55	214
	24802	Coding	395	caccaccatcccgtgaag	41	215
	24803	Coding	455	tcttgttctgcgtagtc	80	216
	24804	Coding	512	tgttctgtcatagtagt	65	217
	24805	Coding	527	tcaccttgcggatgatgt	70	218
20	24806	Coding	582	gagcaccctgcgacctca	64	219
	24807	Coding	600	ggcgggcagtcctcagtg	67	220
	24808	Coding	787	ggtgaagggtggaatagag	45	221
	24809	Coding	993	tccgatttcaggtttggg	75	222
	24810	Coding	1110	ttggtggtttctggcaca	82	223
25	24811	Coding	1132	tggaggggacttctggctc	60	224
	24812	Coding	1376	gcgtagggaagcagggatg	49	225
	24813	Coding	1440	gtgtccagaagtgaatg	71	226
	24814	Coding	1498	actggatggaaactggaa	62	227
	24815	Coding	1541	ggccatccacgtgatag	78	228
30	24816	3' UTR	1701	ccaccacaatcagagcat	54	229
	24817	3' UTR	1711	gatccccaccccaccaca	44	230
	24818	3' UTR	1765	tgtttctgtggaggaga	74	231
	24819	3' UTR	1790	aaacagagaagttgtgga	64	232
	24820	3' UTR	1802	gggactgacagaaaacag	16	233
35	24821	3' UTR	1860	ataaataaataaaccgcc	38	234
	24822	3' UTR	1894	gttaggtcaggctcatcc	59	235
	24823	3' UTR	1974	gttctcaagccagacctc	62	236
	24824	3' UTR	1992	aataaagaaagaaaggtc	35	237
	24825	3' UTR	2006	agggcagggtgagaaata	0	238
40	24826	3' UTR	2053	cttctactcacatccaaa	46	239
	24827	3' UTR	2068	caaaacaaactaactctt	38	240
	24828	3' UTR	2080	ggaataataaaacaaaac	37	241



107

24829	3' UTR	2107	ttcttcctggaccctga	71	242
24830	3' UTR	2161	ccaagggtgtgattcttc	88	243
24831	3' UTR	2200	tgtctgagagaaaggttg	65	244

5

**EXAMPLE 26: Automated Assay of Gi alpha proteins Oligonucleotide Activity**

G-alpha-11 is a member of the Gq subfamily of G proteins whose primary function is to activate PLC-b isoforms producing second messengers and affecting intracellular calcium stores.

10

Oligonucleotides were designed as described in Example 2, synthesized as described in Examples 3 through 8, analyzed as described in Examples 9 and assayed as described in Example 10 except for target specific primer and probes. G-alpha-11 probes and primers were designed to hybridize to the human G-alpha-11 sequence, using published sequence information (GenBank accession number AF011497, incorporated herein by reference as SEQ ID NO:245). For G-alpha-11 the PCR primers were: forward primer: TGACCACCTTCGAGCATCAG (SEQ ID NO: 246) reverse primer: CGGTCGTAGCATTCCTGGAT (SEQ ID NO: 247) and the PCR probe was: FAM-TCAGTGCCATCAAGACCCTGTGGGAG-TAMRA (SEQ ID NO: 248) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

15

20

**EXAMPLE 27: Antisense inhibition of G-alpha-11 expression- phosphorothioate oligodeoxynucleotides**

25

30

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human G-alpha-11 RNA, using published sequences (GenBank accession number AF011497, incorporated herein by reference as SEQ ID NO: 245). The oligonucleotides are shown in Table 16. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. AF011497), to which the oligonucleotide binds. All compounds in Table 16 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds were analyzed for effect on G-alpha-11 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from

three experiments. If present, "N.D." indicates "no data".

Table 16

## Inhibition of G-alpha-11 mRNA levels by phosphorothioate oligodeoxynucleotides

	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
5	20576	Coding	1	gatggactccagagtcac	0	249
	20577	Coding	6	gccatgatggactccaga	75	250
	20578	Coding	9	cacgccatgatggactcc	0	251
	20579	Coding	25	ctcatcgctcaggcaaca	61	252
10	20580	Coding	31	cttcacatcatcgctcag	20	253
	20581	Coding	36	gactccttcacatcatcg	15	254
	20582	Coding	45	atccgcttgactccttc	17	255
	20583	Coding	50	cgttgatccgcttgact	0	256
	20584	Coding	61	ctcgatctcgccgtgat	0	257
15	20585	Coding	77	cccgcgcagctgcttct	58	258
	20586	Coding	106	cttgagctcgccgggc	31	259
	20587	Coding	116	gcagcagcagcttgagct	0	260
	20588	Coding	127	gcccgtgccgagcagcag	0	261
	20589	Coding	146	acgtgctctcccgtctct	28	262
20	20590	Coding	159	atctgcttgatgaacgtg	0	263
	20591	Coding	162	cgcactgcttgatgaac	0	264
	20592	Coding	184	gtagccggcgccgtggat	1	265
	20593	Coding	197	tgtcctcctccagtagc	0	266
	20594	Coding	199	cttgctcctccagtagc	79	267
25	20595	Coding	207	aagccgcgcttgctctcc	56	268
	20596	Coding	222	tagacgagcttggtgaag	0	269
	20597	Coding	230	tgttctggtagacgagct	0	270
	20598	Coding	242	tggcggtagaatgttct	0	271
	20599	Coding	258	cggatcatggcctgcatg	1	272
30	20600	Coding	271	cgtctccatggcccggat	49	273
	20601	Coding	285	tagaggatcttgagcgtc	0	274
	20602	Coding	287	tgtagaggatcttgagcg	0	275
	20603	Coding	297	tgctcgtacttgtagagg	7	276
	20604	Coding	306	gccttgcttgctcgtac	25	277
35	20605	Coding	309	ttggccttgcttgctcg	0	278
	20606	Coding	319	caggagcgcattggcctt	0	279
	20607	Coding	340	ctccacgtccacctcccg	69	280
	20608	Coding	349	ggcaccttctccacgtc	27	281
	20609	Coding	362	gatgctcgaaggtggtca	33	282
40	20610	Coding	373	actgacgtactgatgctc	36	283
	20611	Coding	382	cttgatggcactgacgta	78	284
	20612	Coding	388	cagggtcttgatggcact	0	285

## 109

	20613	Coding	409	ctggatgccccgggtcctc	0	286
	20614	Coding	411	tcctggatgccccgggtcc	30	287
	20615	Coding	429	cgcctgcggtcgtagcat	0	288
	20616	Coding	440	gctggtactcgcgcctgc	41	289
5	20617	Coding	459	tacttggcagagtcggag	34	290
	20618	Coding	468	gtcaggtagtagtctggca	76	291
	20619	Coding	479	ggtcaacgtcggtcaggt	18	292
	20620	Coding	489	gtggcgatgcgggtcaacg	1	293
	20621	Coding	503	gcaggtagcccaaggtgg	20	294
10	20622	Coding	518	cgtcctgctgggtgggca	40	295
	20623	Coding	544	ggtggtgggcacgcggac	0	296
	20624	Coding	555	tcgatgatgccgggtggtg	0	297
	20625	Coding	572	ccaggtcgaaagggtact	0	298
	20626	Coding	578	tgttctccaggtcgaaag	33	299
15	20627	Coding	584	agatgatgttctccaggt	0	300
	20628	Coding	591	atccggaagatgatgttc	0	301
	20629	Coding	624	ctccgctccgaccgtgg	56	302
	20630	Coding	634	gatccacttctccgctc	59	303
	20631	Coding	655	tgtcacgttctcaaagca	0	304
20	20632	Coding	663	atgatggatgtcacgttc	0	305
	20633	Coding	671	cgagaaacatgatggatg	0	306
	20634	Coding	682	gctgagggcgacagagaaa	75	307
	20635	Coding	709	cgactccaccaggacttg	40	308
	20636	Coding	726	atccggttctcgttgcc	22	309
25	20637	Coding	728	ccatccggttctcgttgt	19	310
	20638	Coding	744	agggctttgctctctcc	77	311
	20639	Coding	754	ggtccggaacagggttt	26	312
	20640	Coding	766	gtaggtgatgatggtccg	0	313
	20641	Coding	787	ggaggagttctggaacca	64	314
30	20642	Coding	803	tgaggaaaggatgacgg	0	315
	20643	Coding	818	gcaggtccttctgttga	6	316
	20644	Coding	831	atcttgcctccagcagg	4	317
	20645	Coding	842	gcgagtacaggatcttgt	17	318
	20646	Coding	858	aagtagtcaccagggtgc	0	319
35	20647	Coding	910	gatgaactcccgcccgc	52	320
	20648	Coding	935	ggttcagggtccacgaaca	71	321
	20649	Coding	958	gtagatgatctgtcgct	0	322
	20650	Coding	972	cacgtgaagtgtgagtag	0	323
	20651	Coding	993	atgttctccgtgtcgggtg	0	324
40	20652	Coding	1014	acggccgcgaacacgaag	6	325
	20653	Coding	1027	gatggtgtccttcacggc	0	326
	20654	Coding	1043	tcaggttcagctgcagga	3	327
	20655	Coding	1059	accagattgtactccttc	0	328

**EXAMPLE 28: Antisense inhibition of G-alpha-11 expression- phosphorothioate 2'-MOE gapmer oligonucleotides**

In accordance with the present invention, a second series of oligonucleotides targeted to human G-alpha-11 were synthesized. The oligonucleotide sequences are shown in Table 17. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. AF011497), to which the oligonucleotide binds.

All compounds in Table 17 are chimeric oligonucleotides ("gapmers") 18 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

Data were obtained by real-time quantitative PCR as described in other examples herein and are averaged from three experiments. If present, "N.D." indicates "no data".

**Table 17**

**Inhibition of G-alpha-11 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap**

	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
20	20981	Coding	1	gatggactccagagtc	0	249
	20982	Coding	6	gccatgatggactccaga	0	250
	20983	Coding	9	cacgccatgatggactcc	0	251
25	20984	Coding	25	ctcatcgctcaggcaaca	0	252
	20985	Coding	31	cttcacctcatcgctcag	2	253
	20986	Coding	36	gactccttcacctcatcg	0	254
	20987	Coding	45	atccgcttggaactcttc	19	255
	20988	Coding	50	cgttgatccgcttggaact	15	256
30	20989	Coding	61	ctcgatctcggcggtgat	0	257
	20990	Coding	77	cccgccgcagctgcttct	41	258
	20991	Coding	106	cttgagctcgcgccgggc	19	259
	20992	Coding	116	gcagcagcagcttgagct	23	260
	20993	Coding	127	gcccgtgccgagcagcag	38	261
35	20994	Coding	146	acgtgctctcccgtct	34	262
	20995	Coding	159	atctgcttgatgaacgtg	56	263

## 111

	20996	Coding	162	cgcacatctgcttgatgaac	31	264
	20997	Coding	184	gtagccggcgccgtggat	0	265
	20998	Coding	197	tgctcctcctccagtagc	42	266
	20999	Coding	199	cttgctcctcctccagta	0	267
5	21000	Coding	207	aagccgcgcttgctcctcc	73	268
	21001	Coding	222	tagacgagcttggtgaag	0	269
	21002	Coding	230	tggtctggtagacgagct	61	270
	21003	Coding	242	tggcgggtgaagatgttct	14	271
	21004	Coding	258	cggatcatggcctgcatg	84	272
10	21005	Coding	271	cgctccatggcccgat	70	273
	21006	Coding	285	tagaggatcttgagcgtc	39	274
	21007	Coding	287	tgtagaggatcttgagcg	28	275
	21008	Coding	297	tgctcgtacttgtagagg	70	276
	21009	Coding	306	gccttggtctgctcgtac	76	277
15	21010	Coding	309	ttggccttggtctgctcg	0	278
	21011	Coding	319	caggagcgcattggcctt	87	279
	21012	Coding	340	ctccacgtccacctcccg	0	280
	21013	Coding	349	ggtcaccttctccacgtc	69	281
	21014	Coding	362	gatgctcgaaggtggtca	0	282
20	21015	Coding	373	actgacgtactgatgctc	69	283
	21016	Coding	382	cttgatggcactgacgta	32	284
	21017	Coding	388	cagggtcttgatggcact	19	285
	21018	Coding	409	ctggatgcccgggtcctc	63	286
	21019	Coding	411	tcctggatgcccgggtcc	56	287
25	21020	Coding	429	cgctcgcggtcgtagcat	73	288
	21021	Coding	440	gctggtactcgcgctgc	68	289
	21022	Coding	459	tacttggcagagtcggag	50	290
	21023	Coding	468	gtcaggtagtacttgga	13	291
	21024	Coding	479	ggtaacgcggtcaggt	64	292
30	21025	Coding	489	gtggcgatcggtcaacg	52	293
	21026	Coding	503	gcaggtagcccaaggtgg	52	294
	21027	Coding	518	cgctcgtggtgggca	0	295
	21028	Coding	544	gggtggtgggcacgcggac	81	296
	21029	Coding	555	tcgatgatccggtggtg	48	297
35	21030	Coding	572	ccaggtcgaaagggtact	61	298
	21031	Coding	578	tggtctccaggtcgaaag	0	299
	21032	Coding	584	agatgatgttctccaggt	0	300
	21033	Coding	591	atccggaagatgatgttc	0	301
	21034	Coding	624	ctccgctccgaccgctgg	59	302
40	21035	Coding	634	gatccacttctccgctc	17	303
	21036	Coding	655	tgtcacgttctcaaagca	9	304
	21037	Coding	663	atgatggatgtcacgttc	41	305
	21038	Coding	671	cgagaaacatgatggatg	0	306

## 112

	21039	Coding	682	gctgagggcgacgagaaa	11	307
	21040	Coding	709	cgactccaccaggacttg	0	308
	21041	Coding	726	atccggttctcgttgcc	67	309
	21042	Coding	728	ccatccggttctcgttgt	30	310
5	21043	Coding	744	agggttctcctctcc	61	311
	21044	Coding	754	ggtccggaacagggttt	72	312
	21045	Coding	766	gtaggtgatgatgtccg	68	313
	21046	Coding	787	ggaggagtcttgaacca	54	314
	21047	Coding	803	tgaggaaaggatgacgg	23	315
10	21048	Coding	818	gcaggtccttctgtga	0	316
	21049	Coding	831	atctgtctccagcagg	39	317
	21050	Coding	842	gcgagtacaggatcttgt	74	318
	21051	Coding	858	aagtagtccaccagggtgc	36	319
	21052	Coding	910	gatgaactcccgcgccgc	67	320
15	21053	Coding	935	ggttcagggtccacgaaca	37	321
	21054	Coding	958	gtagatgatctgtcgt	64	322
	21055	Coding	972	cacgtgaagtgtgagtag	37	323
	21056	Coding	993	atgttctccgtgtcgggtg	0	324
	21057	Coding	1014	acggccgcgaacacgaag	0	325
20	21058	Coding	1027	gatggtgtccttcacggc	69	326
	21059	Coding	1043	tcagggtcagctgcagga	0	327
	21060	Coding	1059	accagattgtactcttc	0	328

**EXAMPLE 29: Automated Assay of AKT-1 Oligonucleotide Activity**

25 Akt-1 (also known as PKB alpha and RAC-PK alpha) is a member of the AKT/PKB family of serine/threonine kinases and has been shown to be involved in a diverse set of signaling pathways.

Oligonucleotides were designed as described in Example 2, synthesized as described in Examples 3 through 8, analyzed as described in Examples 9 and assayed as described in Example 10 except for target specific primer and probes. AKT-1 probes and primers were designed to hybridize to the human AKT-1 sequence, using published sequence information (GenBank accession number M63167, incorporated herein by reference as SEQ ID NO:329). For Akt-1 the PCR primers were:

forward primer: CGTGACCATGAACGAGTTTGA (SEQ ID NO: 330)

35 reverse primer: CAGGATCACCTTGCCGAAA (SEQ ID NO: 331) and the PCR probe was: FAM-CTGAAGCTGCTGGGCAAGGGCA-TAMRA (SEQ ID NO: 332) where FAM (PE-Applied Biosystems, Foster City, CA) is the fluorescent reporter dye) and TAMRA (PE-Applied Biosystems, Foster City, CA) is the quencher dye.

**EXMAMPLE 30: Antisense inhibition of Akt-1 expression- phosphorothioate oligodeoxynucleotides**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human Akt-1 RNA, using published sequences (GenBank accession number M63167, incorporated herein by reference as SEQ ID NO: 329). The oligonucleotides are shown in Table 18. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. M63167), to which the oligonucleotide binds. All compounds in Table 18 are oligodeoxynucleotides with phosphorothioate backbones (internucleoside linkages) throughout. The compounds were analyzed for effect on Akt-1 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from three experiments. If present, "N.D." indicates "no data".

**Table 18**

**Inhibition of Akt-1 mRNA levels by phosphorothioate oligodeoxynucleotides**

ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
28880	5' UTR	4	ccctgtgccctgtcccag	55	333
28881	5' UTR	27	cctaagcccctgggtgaca	15	334
28882	5' UTR	62	cttgacttctttgaccc	68	335
28883	5' UTR	70	ggcagccccccttgacttc	53	336
28884	Coding	213	caaccctccttcacaata	24	337
28885	Coding	234	tactcccctcggttggtgc	0	338
28886	Coding	281	tgccatcattcttgagga	65	339
28887	Coding	293	agccaatgaaggtgccat	67	340
28888	Coding	352	cacagagaagttgttgag	22	341
28889	Coding	496	agtctggatggcggtgtg	49	342
28890	Coding	531	tectcctcctcctgcttc	9	343
28891	Coding	570	cctgagttgtcactgggt	49	344
28892	Coding	666	ccgaaagtgccttgccc	56	345
28893	Coding	744	gccacgatgacttccttc	60	346
28894	Coding	927	cggctctcggagaacaca	0	347
28895	Coding	990	acgttcttctcagtggtgc	30	348
28896	Coding	1116	gtgccgcaaaagggtcttc	66	349
28897	Coding	1125	tactcaggtgtgccgcaa	66	350
28898	Coding	1461	ggcttgaagggtgggctg	41	351
28899	Coding	1497	tcaaatacctggtgtca	51	352

## 114

	28900	Coding	1512	gccgtgaactcctcatca	56	353
	28901	Coding	1541	ggtcaggtgggtgatgg	0	354
	28902	Coding	1573	ctcgtgtccacacactc	61	355
	28903	3' UTR	1671	gcctctccatccctccaa	76	356
5	28904	3' UTR	1739	acagcgtggcttctctca	12	357
	28905	3' UTR	1814	ttttctccctaccccg	64	358
	28906	3' UTR	1819	gatagttttctccctac	0	359
	28907	3' UTR	1831	taaaacccgcaggatagt	74	360
	28908	3' UTR	1856	ggagaacaaactggatga	0	361
10	28909	3' UTR	1987	ctggctgacagagtgagg	59	362
	28910	3' UTR	1991	gcggctggctgacagagt	61	363
	28911	3' UTR	2031	cccagagagatgacagat	46	364
	28912	3' UTR	2127	gctgctgtgtgcctgcca	38	365
	28913	3' UTR	2264	cataatacacaataacaa	39	366
15	28914	3' UTR	2274	atttgaacaacataatac	11	367
	28915	3' UTR	2397	aagtgtctaccgtggagag	57	368
	28916	3' UTR	2407	cgaaaagggtcaagtgtcta	41	369
	28917	3' UTR	2453	cagggagtcagggaggggc	13	370
	28918	3' UTR	2545	aaagttgaatgttgtaaa	10	371
20	28919	3' UTR	2553	aaaataactaaagttgaat	25	372

**EXAMPLE 31: Antisense inhibition of Akt-1 expression- phosphorothioate 2'-MOE gapmer oligonucleotides**

In accordance with the present invention, a second series of oligonucleotides  
 25 targeted to human Akt-1 were synthesized. The oligonucleotide sequences are shown in Table 19. Target sites are indicated by nucleotide numbers, as given in the sequence source reference (Genbank accession no. M63167), to which the oligonucleotide binds.

All compounds in Table 19 are chimeric oligonucleotides ("gapmers") 18  
 nucleotides in length, composed of a central "gap" region consisting of ten 2'-  
 30 deoxynucleotides, which is flanked on both sides (5' and 3' directions) by four-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. Cytidine residues in the 2'-MOE wings are 5-methylcytidines.

Data were obtained by real-time quantitative PCR as described in other examples  
 35 herein and are averaged from three experiments. If present, "N.D." indicates "no data".



Table 19

**Inhibition of Akt-1 mRNA levels by chimeric phosphorothioate oligonucleotides  
having 2'-MOE wings and a deoxy gap**

	ISIS#	REGION	TARGET SITE	SEQUENCE	% Inhibition	SEQ ID NO.
5	28920	5' UTR	4	ccctgtgccctgtcccag	88	333
	28921	5' UTR	27	cctaagcccctggtgaca	44	334
	28922	5' UTR	62	ctttgacttctttgaccc	61	335
	28923	5' UTR	70	ggcagccccttgacttc	79	336
10	28924	Coding	213	caaccctccttcacaata	72	337
	28925	Coding	234	tactcccctcgtttgtgc	39	338
	28926	Coding	281	tgccatcattcttgagga	73	339
	28927	Coding	293	agccaatgaagggtccat	62	340
	28928	Coding	352	cacagagaagttgttgag	48	341
15	28929	Coding	496	agtctggatggcggttgt	43	342
	28930	Coding	531	tectcctcctcctgcttc	49	343
	28931	Coding	570	cctgagttgtcactgggt	71	344
	28932	Coding	666	ccgaaagtgcccttgccc	64	345
	28933	Coding	744	gccacgatgacttccttc	66	346
20	28934	Coding	927	cggtcctcggagaacaca	77	347
	28935	Coding	990	acgttcttctccgagtgc	89	348
	28936	Coding	1116	gtgccgcaaaaggtcttc	61	349
	28937	Coding	1125	tactcaggtgtgccgcaa	74	350
	28938	Coding	1461	ggcttgaagggtgggctg	54	351
25	28939	Coding	1497	tcaaaatacctggtgtca	78	352
	28940	Coding	1512	gccgtgaactcctcatca	88	353
	28941	Coding	1541	ggtcaggtggtgtgatgg	71	354
	28942	Coding	1573	ctcgtgtccacacactc	83	355
	28943	3' UTR	1671	gcctctccatccctccaa	86	356
30	28944	3' UTR	1739	acagcgtggcttctctca	73	357
	28945	3' UTR	1814	ttttcttcctaccccgc	77	358
	28946	3' UTR	1819	gatagttttcttcctac	43	359
	28947	3' UTR	1831	taaaacccgcaggatagt	64	360
	28948	3' UTR	1856	ggagaacaaactggatga	70	361
35	28949	3' UTR	1987	ctggctgacagagtgagg	90	362
	28950	3' UTR	1991	gcggctggctgacagagt	82	363
	28951	3' UTR	2031	cccagagagatgacagat	53	364
	28952	3' UTR	2127	gctgctgtgtgcctgcca	80	365
	28953	3' UTR	2264	cataatacacaataacaa	48	366
40	28954	3' UTR	2274	atttgaacaacataatac	39	367
	28955	3' UTR	2397	aagtgtctaccgtggagag	38	368
	28956	3' UTR	2407	cgaaaagggtcaagtgtcta	83	369

## 116

28957	3' UTR	2453	cagggagtcagggagggc	59	370
28958	3' UTR	2545	aaagttgaatgttgtaa	25	371
28959	3' UTR	2553	aaaatactaaagttgaat	45	372

**What is claimed is:**

1. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid sequence comprising generating a library of virtual compounds *in silico* according to defined criteria, and evaluating *in silico* the binding of said virtual compounds with said target nucleic acid according to defined criteria.

2. A method of defining a set of oligonucleotides that modulate the expression of a target nucleic acid sequence via binding of said oligonucleotides with said target nucleic acid sequence comprising generating *in silico* a plurality of virtual oligonucleotides according to defined criteria, and evaluating *in silico* the binding of said plurality of virtual oligonucleotides with said target nucleic acid according to defined criteria.

3. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid comprising, generating *in silico* a library of virtual compounds according to defined criteria wherein said virtual compounds modulate the expression of said target nucleic acid sequence, and robotically synthesizing synthetic compounds corresponding to at least some of said virtual compounds.

4. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid comprising generating *in silico* virtual compounds according to defined criteria wherein said virtual compounds modulate the expression of said target nucleic acid sequence, synthesizing synthetic compounds corresponding to at least some of said virtual compounds, and robotically assaying said synthetic compounds for one or more desired physical, chemical or biological properties.

5. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid sequence comprising generating *in silico* a library of nucleobase sequences according to

defined criteria and evaluating *in silico* a plurality of virtual oligonucleotides having said nucleobase sequences according to defined criteria.

6. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid sequence comprising evaluating *in silico* a plurality of virtual compounds according to defined criteria and robotically synthesizing a plurality of synthetic compounds corresponding to said plurality of virtual compounds.
7. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid sequence comprising evaluating *in silico* a plurality of virtual compounds according to defined criteria and robotically assaying a plurality of synthetic compounds corresponding to at least some of said virtual compounds for one or more desired physical, chemical or biological properties.
8. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid sequence comprising generating a library of nucleobase sequences *in silico* according to defined criteria and robotically synthesizing a plurality of synthetic compounds compounds having said nucleobase sequences.
9. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid sequence comprising robotically synthesizing a plurality of synthetic compounds and robotically assaying said plurality of synthetic compounds for one or more desired physical, chemical or biological properties.
10. A method of defining a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid sequence comprising generating a library of nucleobase sequences *in silico* according to

defined criteria and robotically assaying a plurality of synthetic compounds having at least some of said nucleobase sequences for one or more desired physical, chemical or biological properties.

5        11.     A method of generating a set of oligonucleotides that modulate the expression of a target nucleic acid sequence via binding of said oligonucleotides with said target nucleic acid sequence, comprising the steps of:

              (a) generating a library of nucleobase sequences *in silico* according to defined criteria;

10            (b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria; and

              (c) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides.

15        12.     A method of generating a set of oligonucleotides that modulate the expression of a target nucleic acid sequence via binding of said oligonucleotides with said target nucleic acid sequence, comprising the steps of:

              (a) generating a library of nucleobase sequences *in silico* according to defined criteria;

20            (b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria; and

              (c) robotically assaying a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides for one or more desired physical, chemical or biological properties.

25

13.     A method of generating a set of oligonucleotides that modulate the expression of a target nucleic acid sequence via binding of said oligonucleotides with said target nucleic acid sequence, comprising the steps of:

30            (a) generating a library of nucleobase sequences *in silico* according to defined criteria;

              (b) robotically synthesizing a plurality of synthetic oligonucleotides having at least

some of said nucleobase sequences; and

(c) robotically assaying said plurality of synthetic oligonucleotides for one or more desired physical, chemical or biological properties.

5 14. A method of generating a set of oligonucleotides that modulate the expression of a target nucleic acid sequence via binding of said oligonucleotides with said target nucleic acid sequence, comprising the steps of:

(a) evaluating *in silico* a plurality of virtual oligonucleotides according to defined criteria;

10 (b) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides; and

(c) robotically assaying said plurality of synthetic oligonucleotides for one or more desired physical, chemical or biological properties.

15 15. A method of generating a set of oligonucleotides that modulate the expression of a target nucleic acid sequence via binding of said oligonucleotides with said target nucleic acid sequence, comprising the steps of:

(a) generating a library of nucleobase sequences *in silico* according to defined criteria;

20 (b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria;

(c) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides; and

25 (d) robotically assaying said plurality of synthetic oligonucleotides for one or more desired physical, chemical or biological properties.

16. A method of generating a set of oligonucleotides that modulate the expression of a target nucleic acid sequence via binding of said oligonucleotides with said target nucleic acid sequence, comprising the steps of:

30 (a) generating a library of nucleobase sequences *in silico* according to defined criteria;

(b) choosing an oligonucleotide chemistry;

(c) robotically synthesizing a set of synthetic oligonucleotides having said nucleobase sequences of step (a) and said oligonucleotide chemistry of step (b);

5 (d) robotically assaying said set of synthetic oligonucleotides of step (c) for a physical, chemical or biological activity; and

(e) selecting a subset of said set of synthetic oligonucleotides of step (c) having a desired level of physical, chemical or biological activity in order to generate said set of compounds.

10 17. A method of generating a set of oligonucleotides that modulate the expression of a target nucleic acid sequence via binding of said oligonucleotides with said target nucleic acid sequence, comprising the steps of:

(a) generating a library of nucleobase sequences *in silico* according to defined criteria;

15 (b) choosing an oligonucleotide chemistry;

(c) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) and the oligonucleotide chemistry of (b) according to defined criteria, and selecting those having preferred characteristics, in order to generate a set of preferred nucleobase sequences;

20 (d) robotically synthesizing a set of synthetic oligonucleotides having said preferred nucleobase sequences of step (c) and said oligonucleotide chemistry of step (b);

(e) robotically assaying said set of synthetic oligonucleotides of step (d) for a physical, chemical or biological activity; and

25 (f) selecting a subset of said set of synthetic oligonucleotides of step (d) having a desired level of physical, chemical or biological activity in order to generate said set of oligonucleotides.

18. The method of claim 12, wherein said step of robotically assaying said plurality of synthetic oligonucleotide compounds is performed by computer-controlled real-time  
30 polymerase chain reaction or by computer-controlled enzyme-linked immunosorbent assay.

19. The method of claim 11, wherein said target nucleic acid sequence is that of a genomic DNA, a cDNA, a product of a polymerase chain reaction, an expressed sequence tag, an mRNA or a structural RNA.

5 20. The method of claim 11, wherein said target nucleic acid sequence is a human nucleic acid.

21. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences comprising generating a  
10 library of antisense nucleobase sequences *in silico* according to defined criteria.

22. A method of identifying a set of compounds that modulate the expression of a target nucleic acid sequence via binding of said compounds with said target nucleic acid sequence comprising evaluating *in silico* a plurality of virtual oligonucleotides according  
15 to defined criteria.

23. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an compound to said nucleic acid sequences comprising robotically synthesizing a plurality of synthetic antisense compounds.

20 24. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an compound to said nucleic acid sequences comprising robotically assaying a plurality of synthetic antisense compounds for one or more desired physical, chemical or biological properties.

25 25. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences comprising generating *in silico* a library of nucleobase sequences according to defined criteria and evaluating *in silico* a plurality of virtual oligonucleotides having said nucleobase sequences according to  
30 defined criteria.



26. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences comprising evaluating *in silico* a plurality of virtual oligonucleotides according to defined criteria and robotically synthesizing a plurality of synthetic oligonucleotides corresponding to least some of said virtual oligonucleotides.

27. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences comprising evaluating *in silico* a plurality of virtual oligonucleotides according to defined criteria and robotically assaying a plurality of synthetic oligonucleotides corresponding to least some of said virtual oligonucleotides for one or more desired physical, chemical or biological properties.

28. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences comprising generating a library of nucleobase sequences *in silico* according to defined criteria and robotically synthesizing a plurality of synthetic oligonucleotides having said nucleobase sequences.

29. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences comprising robotically synthesizing a plurality of synthetic oligonucleotides and robotically assaying said plurality of synthetic oligonucleotides for one or more desired physical, chemical or biological properties.

30. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences comprising generating a library of nucleobase sequences *in silico* according to defined criteria and robotically assaying a plurality of synthetic oligonucleotides having said nucleobase sequences for one or more desired physical, chemical or biological properties.

31. A method of identifying one or more nucleic acid sequences amenable to antisense

binding of an oligonucleotide to said nucleic acid sequences comprising the steps of:

(a) generating a library of nucleobase sequences *in silico* according to defined criteria;

5 (b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria; and

(c) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides.

32. A method of identifying one or more nucleic acid sequences amenable to antisense  
10 binding of an oligonucleotide to said nucleic acid sequences, comprising the steps of:

(a) generating a library of nucleobase sequences *in silico* according to defined criteria;

(b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria; and

15 (c) robotically assaying a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides for one or more desired physical, chemical or biological properties.

33. A method of identifying one or more nucleic acid sequences amenable to antisense  
20 binding of an oligonucleotide to said nucleic acid sequences, comprising the steps of:

(a) generating a library of nucleobase sequences *in silico* according to defined criteria;

(b) robotically synthesizing a plurality of synthetic oligonucleotides having at least some of said nucleobase sequences; and

25 (c) robotically assaying said plurality of synthetic oligonucleotides for one or more desired physical, chemical or biological properties.

34. A method of identifying one or more nucleic acid sequences amenable to antisense  
binding of an oligonucleotide to said nucleic acid sequences, comprising the steps of:

30 (a) evaluating *in silico* a plurality of virtual oligonucleotides according to defined criteria;

(b) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides; and

(c) robotically assaying said plurality of synthetic oligonucleotides for one or more desired physical, chemical or biological properties.

5

35. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences, comprising the steps of:

(a) generating a library of nucleobase sequences *in silico* according to defined criteria;

10

(b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria;

(c) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said plurality of virtual oligonucleotides; and

(d) robotically assaying said plurality of synthetic oligonucleotides for one or more desired physical, chemical or biological properties.

15

36. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences, comprising the steps of:

(a) generating a library of nucleobase sequences *in silico* according to defined criteria;

20

(b) choosing an oligonucleotide chemistry;

(c) robotically synthesizing a set of synthetic oligonucleotides having said nucleobase sequences of step (a) and said oligonucleotide chemistry of step (b);

(d) robotically assaying said set of synthetic oligonucleotides of step (c) for a physical, chemical or biological activity; and

25

(e) selecting a subset of said set of synthetic oligonucleotides of step (c) having a desired level of physical, chemical or biological activity.

37. A method of identifying one or more nucleic acid sequences amenable to antisense binding of an oligonucleotide to said nucleic acid sequences, comprising the steps of:

30

(a) generating a library of nucleobase sequences *in silico* according to defined

criteria;

(b) choosing an oligonucleotide chemistry;

(c) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria, and selecting those having preferred characteristics, in order to generate a set of preferred nucleobase sequences;

(d) robotically synthesizing a set of synthetic oligonucleotides having said preferred nucleobase sequences of step (b) and said oligonucleotide chemistry of step (c);

(e) robotically assaying said set of synthetic oligonucleotides of step (d) for a physical, chemical or biological activity; and

(f) selecting a subset of said set of oligonucleotides of step (d) having a desired level of physical, chemical or biological activity.

38. The method of claim 32, wherein said step of robotically assaying said plurality of synthetic antisense oligonucleotides is performed by computer-controlled real-time polymerase chain reaction or by computer-controlled enzyme-linked immunosorbent assay.

39. The method of claim 31, wherein said nucleic acid sequence is that of a genomic DNA, a cDNA, a product of a polymerase chain reaction, an expressed sequence tag, an mRNA or a structural RNA.

40. The method of claim 31, wherein said nucleic acid sequence is a human nucleic acid.

41. A computer formatted medium comprising computer readable instructions for identifying compounds that have one or more desired properties according to defined criteria and that bind to a genomic DNA, a cDNA, a product of a polymerase chain reaction, an expressed sequence tag, an mRNA or a structural RNA.

42. A computer formatted medium comprising computer readable instructions for performing the method of any one of claims 1 to 20.

43. A computer formatted medium comprising computer readable instructions for performing a method of identifying one or more nucleic acid sequences amenable to antisense binding of a compound to said nucleic acid sequences.

5 44. A computer formatted medium comprising computer readable instructions for performing the method of any one of claims 21 to 40.

45. A computer formatted medium comprising one or more nucleic acid sequences amenable to antisense binding of a compound to said nucleic acid sequences in computer  
10 readable form.

46. A computer formatted medium comprising one or more nucleic acid sequences amenable to antisense binding of a compound to said nucleic acid sequences in computer readable form, wherein said one or more nucleic acid sequences is identified according to  
15 the method of any one of claims 21-40.

47. A process for validating the function of a gene or the product of said gene comprising generating *in silico* a library of nucleobase sequences targeted to said gene and robotically assaying a plurality of synthetic compounds having at least some of said  
20 nucleobase sequences for effects on biological function.

48. A process for validating the function of a gene or the product of said gene, comprising the steps of:

25 (a) generating a library of nucleobase sequences *in silico* according to defined criteria;

(b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria; and

(c) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides.

30

49. A process for validating the function of a gene or the product of said gene,

comprising the steps of:

- (a) generating a library of nucleobase sequences *in silico* according to defined criteria;
- (b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) according to defined criteria; and
- (c) robotically assaying a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides for effects on biological function.

50. A process for validating the function of a gene or the product of said gene, comprising the steps of:

- (a) generating a library of nucleobase sequences *in silico* according to defined criteria;
- (b) robotically synthesizing a plurality of synthetic oligonucleotides having at least some of said nucleobase sequences; and
- (c) robotically assaying said plurality of synthetic oligonucleotides for effects on biological function.

51. A process for validating the function of a gene or the product of said gene, comprising the steps of:

- (a) evaluating *in silico* a plurality of virtual oligonucleotides according to defined criteria;
- (b) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides; and
- (c) robotically assaying said plurality of synthetic oligonucleotides for effects on biological function.

52. A process for validating the function of a gene or the product of said gene, comprising the steps of:

- (a) generating a library of nucleobase sequences *in silico* according to defined criteria;
- (b) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase

sequences of (a) according to defined criteria;

(c) robotically synthesizing a plurality of synthetic oligonucleotides corresponding to at least some of said virtual oligonucleotides; and

5 (d) robotically assaying said plurality of synthetic oligonucleotides for effects on biological function.

53. A process for validating the function of a gene or the product of said gene, comprising the steps of:

10 (a) generating a library of nucleobase sequences *in silico* according to defined criteria;

(b) choosing an oligonucleotide chemistry;

(c) robotically synthesizing a set of synthetic oligonucleotides having said nucleobase sequences of step (a) and said oligonucleotide chemistry of step (b);

15 (d) robotically assaying said set of synthetic oligonucleotides of step (c) for effects on biological function; and

(e) selecting a subset of said set of synthetic oligonucleotides of step (c) having a desired level of physical, chemical or biological activity in order to generate said set of compounds.

20 54. A process for validating the function of a gene or the product of said gene, comprising the steps of:

(a) generating a library of nucleobase sequences *in silico* according to defined criteria;

(b) choosing an oligonucleotide chemistry;

25 (c) evaluating *in silico* a plurality of virtual oligonucleotides having the nucleobase sequences of (a) and the oligonucleotide chemistry of (b) according to defined criteria, and selecting those having preferred characteristics, in order to generate a set of preferred nucleobase sequences;

30 (d) robotically synthesizing a set of synthetic oligonucleotides having said preferred nucleobase sequences of step (c) and said oligonucleotide chemistry of step (b);

(e) robotically assaying said set of synthetic oligonucleotides of step (d) for effects

on biological function; and

(f) selecting a subset of said set of synthetic oligonucleotides of step (d) having a desired level of physical, chemical or biological activity in order to generate said set of oligonucleotides.







Figure 1

2/24

**Figure 2**

3/24





Figure 3

4/24

*Figure 4*

5/24




Figure 5

6/24






Figure 6

7/24

From connector  
at step 317

**Figure 7**

8/24

*Figure 8*



9/24

350 →




Figure 9

10/24




Figure 10

11/24




Figure 11

12/24




Figure 12

13/24




Figure 13

14/24




Figure 14

15/24




Figure 15

16/24





Figure 16



17/24

*Figure 17*




Figure 18

19/24




Figure 19





Figure 20

21/24



22/24




Figure 22

23/24




FIG. 23




FIG. 24



## SEQUENCE LISTING

<110> ISIS Pharmaceuticals, Inc.

Cowsert, Lex M.

Baker, Brenda F.

McNeil, John

Freier, Susan M.

Sasmor, Henri M.

Brooks, Douglas G.

Ohashi, Cara

Wyatt, Jacqueline R.

Borchers, Alexander

Vickers, Timothy A.

<120> Identification of Genetic Targets for Modulation by  
Oligonucleotides and Generation of Oligonucleotides for Gene  
Modulation

<130> ISIS-3456

<140>

<141>

<150> US 09/067,638

<151> 1998-04-28

<150> US 60/081,483

<151> 1998-04-13

<160> 372

<210> 1

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 1

ccaggcggca ggaccact

18

<210> 2

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 2

gaccaggcgg caggacca

18

<210> 3

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 3

aggtgagacc agcggca

18

<210> 4

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 4

cagaggcaga cgaaccat

18

<210> 5

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 5

gcagaggcag acgaacca

18

<210> 6

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 6  
gcaagcagcc ccagagga 18

<210> 7  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 7  
ggtcagcaag cagcccca 18

<210> 8  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 8  
gacagcggtc agcaagca 18

<210> 9  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 9  
gatggacagc ggtcagca 18

<210> 10  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 10  
tctggatgga cagcggtc 18

<210> 11  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 11  
ggtggttctg gatggaca 18

<210> 12  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 12  
gtgggtggtt ctggatgg 18

<210> 13  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 13  
gcagtgggtg gttctgga 18

<210> 14  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 14  
cacaaagaac agcactga 18

<210> 15

<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 15  
ctggcacaaa gaacagca 18

<210> 16  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 16  
tcctggctgg cacaaga 18

<210> 17  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 17  
ctgtcctggc tggcaca 18

<210> 18  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 18  
ctcaccagtt tctgtcct 18

<210> 19  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 19

tcactcacca gtttctgt

18

<210> 20

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 20

gtgcagtcac tcaccagt

18

<210> 21

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 21

actctgtgca gtcactca

18

<210> 22

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 22

cagtgaactc tgtgcagt

18

<210> 23

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 23

attccgtttc agtgaact

18

<210> 24

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 24

gaaggcattc cgttcag

18

<210> 25

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 25

ttcaccgcaa ggaaggca

18

<210> 26

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 26

ctctgttcca ggtgtcta

18

<210> 27

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 27  
ctggtggcag tgtgtctc 18

<210> 28  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 28  
tggggtcgca gtatttgt 18

<210> 29  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 29  
ggttggggtc gcagtatt 18

<210> 30  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 30  
ctaggttggg gtcgcagt 18

<210> 31  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 31  
ggtgcccttc tgctggac 18



<210> 32  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 32  
ctgaggtgcc cttctgct 18

<210> 33  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 33  
gtgtctgttt ctgaggtg 18

<210> 34  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 34  
tgggtgtctgt ttctgagg 18

<210> 35  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 35  
acaggtgcag atggtgtc 18

<210> 36

<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 36  
ttcacagtg cagatggt 18

<210> 37  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 37  
gtgccagcct tcttcaca 18

<210> 38  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 38  
tacagtgccca gccttctt 18

<210> 39  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 39  
ggacacagct ctcacagg 18

<210> 40  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 40

tgcaggacac agctctca

18

<210> 41

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 41

gagcgggtgca ggacacag

18

<210> 42

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 42

aagccggggcg agcatgag

18

<210> 43

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 43

aatctgcttg accccaaa

18

<210> 44

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 44

gaaaccctg tagcaatc

18

<210> 45

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 45

gtatcagaaa cccctgta

18

<210> 46

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 46

gctcgcagat ggtatcag

18

<210> 47

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 47

gcagggctcg cagatggt

18

<210> 48

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 48  
tgggcagggc tcgcagat 18

<210> 49  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 49  
gactgggcag ggctcgca 18

<210> 50  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 50  
cattggagaa gaagccga 18

<210> 51  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 51  
gatgacacat tggagaag 18

<210> 52  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 52  
gcagatgaca cattggag 18

<210> 53  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 53  
tcgaaagcag atgacaca 18

<210> 54  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 54  
gtccaagggt gacatttt 18

<210> 55  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 55  
cacagcttgt ccaagggt 18

<210> 56  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 56  
ttggtctcac agcttgct 18

<210> 57

<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 57  
caggtctttg gtctcaca 18

<210> 58  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 58  
ctgttcaca accaggtc 18

<210> 59  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 59  
gtttgtgcct gcctgttg 18

<210> 60  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 60  
gtcttggttg tgcctgcc 18

<210> 61  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 61

ccacagacaa catcagtc

18

<210> 62

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 62

ctggggacca cagacaac

18

<210> 63

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 63

tcagccgatc ctggggac

18

<210> 64

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 64

caccaccagg gctctcag

18

<210> 65

<211> 18

<212> DNA

<213> Artificial Sequence



<223> Antisense Oligonucleotide

<400> 65

gggatcacca ccagggt

18

<210> 66

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 66

gaggatggca aacaggat

18

<210> 67

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 67

accagcacca agaggatg

18

<210> 68

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 68

tttgataaa gaccagca

18

<210> 69

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 69  
tattggttgg cttcttgg 18

<210> 70  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 70  
gggttcctgc ttggggtg 18

<210> 71  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 71  
gtcgggaaaa ttgatctc 18

<210> 72  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 72  
gatcgtcggg aaaattga 18

<210> 73  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 73  
ggagccagga agatcgtc 18

<210> 74  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 74  
tggagccagg aagatcgt 18

<210> 75  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 75  
tggagcagca gtgttgga 18

<210> 76  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 76  
gtaaagtctc ctgcactg 18

<210> 77  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 77  
tggcatccat gttaaagtc 18

<210> 78

<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 78  
cggttggcat ccatgtaa 18

<210> 79  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 79  
ctctttgccca tctcctg 18

<210> 80  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 80  
ctgtctctcc tgcactga 18

<210> 81  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 81  
ggtgcagcct cactgtct 18

<210> 82  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 82

aactgcctgt ttgccac

18

<210> 83

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 83

cttctgcctg caccctg

18

<210> 84

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 84

actgactggg catagctc

18

<210> 85

<211> 1004

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (48)..(881)

<400> 85

gcctcgctcg ggcgccagc ggtcctgccg cctggctcct cctcgcc atg gtt cgt 56

Met Val Arg

1

ctg cct ctg cag tgc gtc ctc tgg ggc tgc ttg ctg acc gct gtc cat 104

Leu Pro Leu Gln Cys Val Leu Trp Gly Cys Leu Leu Thr Ala Val His

5                    10                    15  
 cca gaa cca ccc act gca tgc aga gaa aaa cag tac cta ata aac agt    152  
 Pro Glu Pro Pro Thr Ala Cys Arg Glu Lys Gln Tyr Leu Ile Asn Ser  
 20                    25                    30                    35  
 cag tgc tgt tct ttg tgc cag cca gga cag aaa ctg gtg agt gac tgc    200  
 Gln Cys Cys Ser Leu Cys Gln Pro Gly Gln Lys Leu Val Ser Asp Cys  
 40                    45                    50  
 aca gag ttc act gaa acg gaa tgc ctt cct tgc ggt gaa agc gaa ttc    248  
 Thr Glu Phe Thr Glu Thr Glu Cys Leu Pro Cys Gly Glu Ser Glu Phe  
 55                    60                    65  
 cta gac acc tgg aac aga gag aca cac tgc cac cag cac aaa tac tgc    296  
 Leu Asp Thr Trp Asn Arg Glu Thr His Cys His Gln His Lys Tyr Cys  
 70                    75                    80  
 gac ccc aac cta ggg ctt cgg gtc cag cag aag ggc acc tca gaa aca    344  
 Asp Pro Asn Leu Gly Leu Arg Val Gln Gln Lys Gly Thr Ser Glu Thr  
 85                    90                    95  
 gac acc atc tgc acc tgt gaa gaa ggc tgg cac tgt acg agt gag gcc    392  
 Asp Thr Ile Cys Thr Cys Glu Glu Gly Trp His Cys Thr Ser Glu Ala  
 100                    105                    110                    115  
 tgt gag agc tgt gtc ctg cac cgc tca tgc tgc ccc ggc ttt ggg gtc    440  
 Cys Glu Ser Cys Val Leu His Arg Ser Cys Ser Pro Gly Phe Gly Val  
 120                    125                    130  
 aag cag att gct aca ggg gtt tct gat acc atc tgc gag ccc tgc cca    488  
 Lys Gln Ile Ala Thr Gly Val Ser Asp Thr Ile Cys Glu Pro Cys Pro  
 135                    140                    145  
 gtc ggc ttc ttc tcc aat gtg tca tct gct ttc gaa aaa tgt cac cct    536  
 Val Gly Phe Phe Ser Asn Val Ser Ser Ala Phe Glu Lys Cys His Pro  
 150                    155                    160  
 tgg aca agc tgt gag acc aaa gac ctg gtt gtg caa cag gca ggc aca    584  
 Trp Thr Ser Cys Glu Thr Lys Asp Leu Val Val Gln Gln Ala Gly Thr  
 165                    170                    175  
 aac aag act gat gtt gtc tgt ggt ccc cag gat cgg ctg aga gcc ctg    632  
 Asn Lys Thr Asp Val Val Cys Gly Pro Gln Asp Arg Leu Arg Ala Leu  
 180                    185                    190                    195

gtg gtg atc ccc atc atc ttc ggg atc ctg ttt gcc atc ctc ttg gtg 680  
 Val Val Ile Pro Ile Ile Phe Gly Ile Leu Phe Ala Ile Leu Leu Val  
 200 205 210

ctg gtc ttt atc aaa aag gtg gcc aag aag cca acc aat aag gcc ccc 728  
 Leu Val Phe Ile Lys Lys Val Ala Lys Lys Pro Thr Asn Lys Ala Pro  
 215 220 225

cac ccc aag cag gaa ccc cag gag atc aat ttt ccc gac gat ctt cct 776  
 His Pro Lys Gln Glu Pro Gln Glu Ile Asn Phe Pro Asp Asp Leu Pro  
 230 235 240

ggc tcc aac act gct gct cca gtg cag gag act tta cat gga tgc caa 824  
 Gly Ser Asn Thr Ala Ala Pro Val Gln Glu Thr Leu His Gly Cys Gln  
 245 250 255

ccg gtc acc cag gag gat ggc aaa gag agt cgc atc tca gtg cag gag 872  
 Pro Val Thr Gln Glu Asp Gly Lys Glu Ser Arg Ile Ser Val Gln Glu  
 260 265 270 275

aga cag tga ggctgcaccc acccaggagt gtggccacgt gggcaaacag 921  
 Arg Gln

gcagttggcc agagagcctg gtgctgctgc tgcaggggtg caggcagaag cggggagcta 981

tgcccagtca gtccagccc ctc 1004

<210> 86

<211> 23

<212> DNA

<213> Artificial Sequence

<223> PCR Primer

<400> 86

cagagttcac tgaaacggaa tgc 23

<210> 87

<211> 23

<212> DNA

<213> Artificial Sequence

<223> PCR Primer

<400> 87	
ggtggcagtg tgtctctctg ttc	23
<210> 88	
<211> 25	
<212> DNA	
<213> Artificial Sequence	
<223> PCR Probe	
<400> 88	
ttccttgccg tgaaagcgaa ttct	25
<210> 89	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<223> PCR Primer	
<400> 89	
gaaggtgaag gtcggagtc	19
<210> 90	
<211> 20	
<212> DNA	
<213> Artificial Sequence	
<223> PCR Primer	
<400> 90	
gaagatggtg atgggatttc	20
<210> 91	
<211> 20	
<212> DNA	
<213> Artificial Sequence	
<223> PCR Probe	
<400> 91	
caagcttccc gttctcagcc	20



<210> 92  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<223> Assembled Target Region

<400> 92  
agtggctctg ccgcctggtc 20

<210> 93  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 93  
gaacagcact gactgtt 18

<210> 94  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 94  
agaacagcac tgactgt 18

<210> 95  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 95  
aagaacagca ctgactgt 18

<210> 96

<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 96  
aaagaacagc actgactg 18

<210> 97  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 97  
caaagaacag cactgact 18

<210> 98  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 98  
acaaagaaca gcactgac 18

<210> 99

<400> 99  
000

<210> 100  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 100

gcacaaagaa cagcactg 18

<210> 101

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 101

ggcacaaga acagcact 18

<210> 102

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 102

tggcacaag aacagcac 18

<210> 103

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 103

gctggcaca agaacagc 18

<210> 104

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 104

ggctggcaca aagaacag 18

<210> 105  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 105  
tggctggcac aaagaaca 18

<210> 106  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 106  
ctggctggca caaagaac 18

<210> 107  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 107  
cctggctggc acaaagaa 18

<210> 108

<400> 108  
000

<210> 109  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

&lt;400&gt; 109

gtcctggctg gcacaaag

18

&lt;210&gt; 110

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 110

tgtcctggct ggcacaaa

18

&lt;210&gt; 111

&lt;400&gt; 111

000

&lt;210&gt; 112

&lt;211&gt; 18

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; Antisense Oligonucleotide

&lt;400&gt; 112

tctgtcctgg ctggcaca

18

&lt;210&gt; 113

&lt;211&gt; 1058

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (77)..(658)

&lt;400&gt; 113

gccttgactt catctcagct ccagagcccg cctctcttc ctgcagcctg ggaacttcag 60

ccggctggag cccacc atg gct gca atc cga aag aag ctg gtg atc gtt 109

Met Ala Ala Ile Arg Lys Lys Leu Val Ile Val  
 1 5 10

ggg gat ggt gcc tgt ggg aag acc tgc ctc ctc atc gtc ttc agc aag 157  
 Gly Asp Gly Ala Cys Gly Lys Thr Cys Leu Leu Ile Val Phe Ser Lys  
 15 20 25

gat cag ttt ccg gag gtc tac gtc cct act gtc ttt gag aac tat att 205  
 Asp Gln Phe Pro Glu Val Tyr Val Pro Thr Val Phe Glu Asn Tyr Ile  
 30 35 40

gcg gac att gag gtg gac ggc aag cag gtg gag ctg gct ctg tgg gac 253  
 Ala Asp Ile Glu Val Asp Gly Lys Gln Val Glu Leu Ala Leu Trp Asp  
 45 50 55

aca gca ggg cag gaa gac tat gat cga ctg cgg cct ctc tcc tac ccg 301  
 Thr Ala Gly Gln Glu Asp Tyr Asp Arg Leu Arg Pro Leu Ser Tyr Pro  
 60 65 70 75

gac act gat gtc atc ctc atg tgc ttc tcc atc gac agc cct gac agc 349  
 Asp Thr Asp Val Ile Leu Met Cys Phe Ser Ile Asp Ser Pro Asp Ser  
 80 85 90

ctg gaa aac att cct gag aag tgg acc cca gag gtg aag cac ttc tgc 397  
 Leu Glu Asn Ile Pro Glu Lys Trp Thr Pro Glu Val Lys His Phe Cys  
 95 100 105

ccc aac gtg ccc atc atc ctg gtg ggg aat aag aag gac ctg agg caa 445  
 Pro Asn Val Pro Ile Ile Leu Val Gly Asn Lys Lys Asp Leu Arg Gln  
 110 115 120

gac gag cac acc agg aga gag ctg gcc aag atg aag cag gag ccc gtt 493  
 Asp Glu His Thr Arg Arg Glu Leu Ala Lys Met Lys Gln Glu Pro Val  
 125 130 135

cgg tct gag gaa ggc cgg gac atg gcg aac cgg atc agt gcc ttt ggc 541  
 Arg Ser Glu Glu Gly Arg Asp Met Ala Asn Arg Ile Ser Ala Phe Gly  
 140 145 150 155

tac ctt gag tgc tca gcc aag acc aag gag gga gtg cgg gag gtg ttt 589  
 Tyr Leu Glu Cys Ser Ala Lys Thr Lys Glu Gly Val Arg Glu Val Phe  
 160 165 170

gag atg gcc act cgg gct ggc ctc cag gtc cgc aag aac aag cgt cgg 637  
 Glu Met Ala Thr Arg Ala Gly Leu Gln Val Arg Lys Asn Lys Arg Arg  
 175 180 185

agg ggc tgt ccc att ctc tga gatccccca aagggccctt ttctacatg 688  
 Arg Gly Cys Pro Ile Leu  
 190

ccccctccct tcacaggggt acagaaatta tccccctaca accccagcct cctgagggct 748

ccatactgaa ggetccattt tcagttccct cctgcccagg actgcattgt ttctagccc 808

cgaggtgtgg cacgggccct cctcccagc gctctgggag ccacgcctat gccctgccct 868

tcctcatggg cccctgggga tcttgccct ttgaccttc ccaaaggatg gtcacacacc 928

agcactttat acacttctgg ctcacaggaa agtgtctgca gtagggaccc agagtcccag 988

gccctggag ttgtttctgc aggggccttg tctctcactg catttggta ggggggcatg 1048

aataaaggct 1058

<210> 114

<211> 23

<212> DNA

<213> Artificial Sequence

<223> PCR Primer

<400> 114

tgatgtcatc ctcattgtct tct 23

<210> 115

<211> 19

<212> DNA

<213> Artificial Sequence

<223> PCR Primer

<400> 115

ccaggatgat gggcacgtt 19

<210> 116

<211> 23

<212> DNA

<213> Artificial Sequence

<223> PCR Probe

<400> 116

cgacagccct gacagcctgg aaa

23

<210> 117

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 117

gagctgagat gaagtcaa

18

<210> 118

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 118

gctgaagttc ccaggctg

18

<210> 119

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 119

ccggctgaag ttcccagg

18

<210> 120

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide



<400> 120  
ggcaccatcc ccaacgat 18

<210> 121  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 121  
aggcaccatc cccaacga 18

<210> 122  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 122  
tcccacaggc accatccc 18

<210> 123  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 123  
aggtcttccc acaggcac 18

<210> 124  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 124  
atgaggaggc aggtcttc 18

<210> 125  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 125  
ttgctgaaga cgatgagg 18

<210> 126  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 126  
tcaaagacag tagggacg 18

<210> 127  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 127  
ttctcaaaga cagtaggg 18

<210> 128  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 128  
agttctcaaa gacagtag 18

<210> 129

<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 129  
tgttttccag gctgtcag 18

<210> 130  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 130  
tcgtcttgcc tcaggtcc 18

<210> 131  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 131  
gtgtgctcgt cttgcctc 18

<210> 132  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 132  
ctcctggtgt gctcgtct 18

<210> 133  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 133

cagaccgaac gggctcct

18

<210> 134

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 134

ttcctcagac cgaacggg

18

<210> 135

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 135

actcaaggta gccaaagg

18

<210> 136

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 136

ctcccgcact ccctcctt

18

<210> 137

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 137

ctcaaacacc tccgcac

18

<210> 138

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 138

ggccatctca aacacctc

18

<210> 139

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 139

cttggtcttg cggacctg

18

<210> 140

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 140

cccctccgac gcttggtc

18

<210> 141

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 141  
gtatggagcc ctcaggag 18

<210> 142  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 142  
gagccttcag tatggagc 18

<210> 143  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 143  
gaaaatggag ccttcagt 18

<210> 144  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 144  
ggaactgaaa atggagcc 18

<210> 145  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 145  
ggaggggaact gaaaatgg 18

<210> 146  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 146  
gcaggaggga actgaaa 18

<210> 147  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 147  
agggcagggc ataggcgt 18

<210> 148  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 148  
ggaagggcag ggcatagg 18

<210> 149  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 149  
catgaggaag ggcagggc 18

<210> 150

<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 150  
taaagtgctg gtgtgtga 18

<210> 151  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 151  
cctgtgagcc agaagtgt 18

<210> 152  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 152  
ttcctgtgag ccagaagt 18

<210> 153  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 153  
cactttcctg tgagccag 18

<210> 154  
<211> 18  
<212> DNA



<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 154

agacactttc ctgtgagc 18

<210> 155

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 155

actctgggtc cctactgc 18

<210> 156

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 156

tgcagaaaca actccagg 18

<210> 157

<211> 3076

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (725)..(2539)

<400> 157

gaattcaaaa tgtcttcagt tgtaaacttt accattattt tacgtacctc taagaaataa 60

aagtgcctct aattaaaata tgatgtcatt aattatgaaa tacttcttga taacagaagt 120

tttaaaatag ccatcttaga atcagtgaatg tattattttc ctcctttgag 180  
 ttaggtcttg tgctttttt tctggccac taaatttcac aatttcaaaa aagcaaaaata 240  
 aacatattct gaatatattt gctgtgaaac acttgacagc agagctttcc accatgaaaa 300  
 gaagcttcat gagtcacaca ttacatcttt ggggttgattg aatgccactg aaacattcta 360  
 gtagcctgga gaagttgacc tacctgtgga gatgcctgcc attaaatggc atcctgatgg 420  
 cttataacac atcactcttc tgtgaagggt ttaattttc aacacagctt actctgtagc 480  
 atcatgttta cattgtatgt ataaagatta tacaaagggt caattgtgta ttcttcctt 540  
 aaaaagtatc agtataggat ttagaatctc catgttgaaa ctctaaatgc atagaaataa 600  
 aaataataaa aaattttca ttttgcttt tcagcctagt attaaaactg ataaaagcaa 660  
 agccatgcac aaaactacct cccatagaga aggctagtcc cttttctcc ccattcattt 720  
 catt atg aac ata gta gaa aac agc ata ttc tta tca aat ttg atg 766  
 Met Asn Ile Val Glu Asn Ser Ile Phe Leu Ser Asn Leu Met  
 1 5 10  
 aaa agc gcc aac acg ttt gaa ctg aaa tac gac ttg tca tgt gaa ctg 814  
 Lys Ser Ala Asn Thr Phe Glu Leu Lys Tyr Asp Leu Ser Cys Glu Leu  
 15 20 25 30  
 tac cga atg tct acg tat tcc act ttt cct gct ggg gtt cct gtc tca 862  
 Tyr Arg Met Ser Thr Tyr Ser Thr Phe Pro Ala Gly Val Pro Val Ser  
 35 40 45  
 gaa agg agt ctt gct cgt gct ggt ttc tat tac act ggt gtg aat gac 910  
 Glu Arg Ser Leu Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Val Asn Asp  
 50 55 60  
 aag gtc aaa tgc ttc tgt tgt ggc ctg atg ctg gat aac tgg aaa aga 958  
 Lys Val Lys Cys Phe Cys Cys Gly Leu Met Leu Asp Asn Trp Lys Arg  
 65 70 75  
 gga gac agt cct act gaa aag cat aaa aag ttg tat cct agc tgc aga 1006  
 Gly Asp Ser Pro Thr Glu Lys His Lys Lys Leu Tyr Pro Ser Cys Arg  
 80 85 90  
 ttc gtt cag agt cta aat tcc gtt aac aac ttg gaa gct acc tct cag 1054  
 Phe Val Gln Ser Leu Asn Ser Val Asn Asn Leu Glu Ala Thr Ser Gln

95            100            105            110  
 cct act ttt cct tct tca gta aca aat tcc aca cac tca tta ctt ccg 1102  
 Pro Thr Phe Pro Ser Ser Val Thr Asn Ser Thr His Ser Leu Leu Pro  
           115            120            125  
 ggt aca gaa aac agt gga tat ttc cgt ggc tct tat tca aac tct cca 1150  
 Gly Thr Glu Asn Ser Gly Tyr Phe Arg Gly Ser Tyr Ser Asn Ser Pro  
           130            135            140  
 tca aat cct gta aac tcc aga gca aat caa gat ttt tct gcc ttg atg 1198  
 Ser Asn Pro Val Asn Ser Arg Ala Asn Gln Asp Phe Ser Ala Leu Met  
           145            150            155  
 aga agt tcc tac cac tgt gca atg aat aac gaa aat gcc aga tta ctt 1246  
 Arg Ser Ser Tyr His Cys Ala Met Asn Asn Glu Asn Ala Arg Leu Leu  
           160            165            170  
 act ttt cag aca tgg cca ttg act ttt ctg tgc cca aca gat ctg gca 1294  
 Thr Phe Gln Thr Trp Pro Leu Thr Phe Leu Ser Pro Thr Asp Leu Ala  
 175            180            185            190  
 aaa gca ggc ttt tac tac ata gga cct gga gac aga gtg gct tgc ttt 1342  
 Lys Ala Gly Phe Tyr Tyr Ile Gly Pro Gly Asp Arg Val Ala Cys Phe  
           195            200            205  
 gcc tgt ggt gga aaa ttg agc aat tgg gaa ccg aag gat aat gct atg 1390  
 Ala Cys Gly Gly Lys Leu Ser Asn Trp Glu Pro Lys Asp Asn Ala Met  
           210            215            220  
 tca gaa cac ctg aga cat ttt ccc aaa tgc cca ttt ata gaa aat cag 1438  
 Ser Glu His Leu Arg His Phe Pro Lys Cys Pro Phe Ile Glu Asn Gln  
           225            230            235  
 ctt caa gac act tca aga tac aca gtt tct aat ctg agc atg cag aca 1486  
 Leu Gln Asp Thr Ser Arg Tyr Thr Val Ser Asn Leu Ser Met Gln Thr  
           240            245            250  
 cat gca gcc cgc ttt aaa aca ttc ttt aac tgg ccc tct agt gtt cta 1534  
 His Ala Ala Arg Phe Lys Thr Phe Phe Asn Trp Pro Ser Ser Val Leu  
 255            260            265            270  
 gtt aat cct gag cag ctt gca agt gcg ggt ttt tat tat gtg ggt aac 1582  
 Val Asn Pro Glu Gln Leu Ala Ser Ala Gly Phe Tyr Tyr Val Gly Asn  
           275            280            285

agt gat gat gtc aaa tgc ttt tgc tgt gat ggt gga ctc agg tgt tgg 1630  
 Ser Asp Asp Val Lys Cys Phe Cys Cys Asp Gly Gly Leu Arg Cys Trp  
           290              295              300

gaa tct gga gat gat cca tgg gtt caa cat gcc aag tgg ttt cca agg 1678  
 Glu Ser Gly Asp Asp Pro Trp Val Gln His Ala Lys Trp Phe Pro Arg  
           305              310              315

tgt gag tac ttg ata aga att aaa gga cag gag ttc atc cgt caa gtt 1726  
 Cys Glu Tyr Leu Ile Arg Ile Lys Gly Gln Glu Phe Ile Arg Gln Val  
           320              325              330

caa gcc agt tac cct cat cta ctt gaa cag ctg cta tcc aca tca gac 1774  
 Gln Ala Ser Tyr Pro His Leu Leu Glu Gln Leu Leu Ser Thr Ser Asp  
           335              340              345              350

agc cca gga gat gaa aat gca gag tca tca att atc cat ttt gaa cct 1822  
 Ser Pro Gly Asp Glu Asn Ala Glu Ser Ser Ile Ile His Phe Glu Pro  
           355              360              365

gga gaa gac cat tca gaa gat gca atc atg atg aat act cct gtg att 1870  
 Gly Glu Asp His Ser Glu Asp Ala Ile Met Met Asn Thr Pro Val Ile  
           370              375              380

aat gct gcc gtg gaa atg ggc ttt agt aga agc ctg gta aaa cag aca 1918  
 Asn Ala Ala Val Glu Met Gly Phe Ser Arg Ser Leu Val Lys Gln Thr  
           385              390              395

gtt caa aga aaa atc cta gca act gga gag aat tat aga cta gtc aat 1966  
 Val Gln Arg Lys Ile Leu Ala Thr Gly Glu Asn Tyr Arg Leu Val Asn  
           400              405              410

gat ctt gtg tta gac tta ctc aat gca gaa gat gaa ata agg gaa gag 2014  
 Asp Leu Val Leu Asp Leu Leu Asn Ala Glu Asp Glu Ile Arg Glu Glu  
           415              420              425              430

gag aga gaa aga gca act gag gaa aaa gaa tca aat gat tta tta tta 2062  
 Glu Arg Glu Arg Ala Thr Glu Glu Lys Glu Ser Asn Asp Leu Leu Leu  
           435              440              445

atc cgg aag aat aga atg gca ctt ttt caa cat ttg act tgt gta att 2110  
 Ile Arg Lys Asn Arg Met Ala Leu Phe Gln His Leu Thr Cys Val Ile  
           450              455              460

cca atc ctg gat agt cta cta act gcc gga att att aat gaa caa gaa 2158  
 Pro Ile Leu Asp Ser Leu Leu Thr Ala Gly Ile Ile Asn Glu Gln Glu

465                      470                      475  
 cat gat gtt att aaa cag aag aca cag acg tct tta caa gca aga gaa    2206  
 His Asp Val Ile Lys Gln Lys Thr Gln Thr Ser Leu Gln Ala Arg Glu  
 480                      485                      490  
 ctg att gat acg att tta gta aaa gga aat att gca gcc act gta ttc    2254  
 Leu Ile Asp Thr Ile Leu Val Lys Gly Asn Ile Ala Ala Thr Val Phe  
 495                      500                      505                      510  
 aga aac tct ctg caa gaa gct gaa gct gtg tta tat gag cat tta ttt    2302  
 Arg Asn Ser Leu Gln Glu Ala Glu Ala Val Leu Tyr Glu His Leu Phe  
 515                      520                      525  
 gtg caa cag gac ata aaa tat att ccc aca gaa gat gtt tca gat cta    2350  
 Val Gln Gln Asp Ile Lys Tyr Ile Pro Thr Glu Asp Val Ser Asp Leu  
 530                      535                      540  
 cca gtg gaa gaa caa ttg cgg aga cta caa gaa gaa aga aca tgt aaa    2398  
 Pro Val Glu Glu Gln Leu Arg Arg Leu Gln Glu Glu Arg Thr Cys Lys  
 545                      550                      555  
 gtg tgt atg gac aaa gaa gtg tcc ata gtg ttt att cct tgt ggt cat    2446  
 Val Cys Met Asp Lys Glu Val Ser Ile Val Phe Ile Pro Cys Gly His  
 560                      565                      570  
 cta gta gta tgc aaa gat tgt gct cct tct tta aga aag tgt cct att    2494  
 Leu Val Val Cys Lys Asp Cys Ala Pro Ser Leu Arg Lys Cys Pro Ile  
 575                      580                      585                      590  
 tgt agg agt aca atc aag ggt aca gtt cgt aca ttt ctt tca tga        2539  
 Cys Arg Ser Thr Ile Lys Gly Thr Val Arg Thr Phe Leu Ser  
 595                      600                      605  
 agaagaacca aaacatcatc taaactttag aattaattta ttaatgtat tataacttta 2599  
 acttttatcc taatttggtt tcctaaaat tttatttat ttacaactca aaaaacattg 2659  
 ttttggttaa catatttata tatgtatcta aaccatatga acatatattt ttagaaact 2719  
 aagagaatga taggcttttg ttcttatgaa cgaaaaagag gtagcactac aaacacaata 2779  
 ttcaatcaaa atttcagcat tattgaaatt gtaagtgaag taaaacttaa gatatttgag 2839  
 ttaaccttta agaattttta atatttggc attgtactaa tacctggttt ttttttgg 2899

ttgtttttt gtacagacag ggcagcatac tgagaccctg cctttaaaaa caaacagaac 2959

aaaaacaaaa caccaggac acatttctct gtctttttg atcagtgtcc tatacatcga 3019

aggtgtgcat atatgttgaa tgacatttta gggacatggg gttttataa agaattc 3076

<210> 158

<211> 22

<212> DNA

<213> Artificial Sequence

<223> PCR Primer

<400> 158

ggactcaggt gttgggaatc tg

22

<210> 159

<211> 24

<212> DNA

<213> Artificial Sequence

<223> PCR Primer

<400> 159

caagtactca caccttggaa acca

24

<210> 160

<211> 27

<212> DNA

<213> Artificial Sequence

<223> PCR Probe

<400> 160

agatgatcca tgggttcaac atgccaa

27

<210> 161

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 161  
actgaagaca ttttgaat 18

<210> 162  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 162  
cttagaggta cgtaaaat 18

<210> 163  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 163  
gcacttttat ttcttaga 18

<210> 164  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 164  
attttaatta gaagcact 18

<210> 165  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 165  
accatatttc actgattc 18

<210> 166  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 166  
ctaactcaaa ggaggaaa 18  
  
<210> 167  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 167  
cacaagacct aactcaaa 18  
  
<210> 168  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 168  
gctctgctgt caagtgtt 18  
  
<210> 169  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 169  
tgtgtgactc atgaagct 18  
  
<210> 170



<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 170  
ttcagtggca ttcaatca 18

<210> 171  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 171  
cttctccagg ctactaga 18

<210> 172  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 172  
ggtcaacttc tcaggct 18

<210> 173  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 173  
taaaaccctt cacagaag 18

<210> 174  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 174

ttaaggaaga aatacaca

18

<210> 175

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 175

gcatggcttt gcttttat

18

<210> 176

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 176

caaacgtgtt ggcgcttt

18

<210> 177

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 177

agcaggaaaa gtggaata

18

<210> 178

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 178

ttaacggaat ttagactc

18

<210> 179

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 179

atttggtact gaagaagg

18

<210> 180

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 180

agagccacgg aaatatcc

18

<210> 181

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 181

aaatcttgat ttgctctg

18

<210> 182

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 182 gtaagtaatc tggcattt	18
<210> 183 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 183 agcaagccac tctgtctc	18
<210> 184 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 184 tgaagtgtct tgaagctg	18
<210> 185 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 185 tttgacatca tcaactgtt	18
<210> 186 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 186 tggcttgaac ttgacgga	18

<210> 187  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 187  
tcatctcctg ggctgtct 18

<210> 188  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 188  
gcagcattaa tcacagga 18

<210> 189  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 189  
tttctctctc ctcttccc 18

<210> 190  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 190  
aacatcatgt tcttggtc 18

<210> 191

<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 191  
atataacaca gcttcagc 18

<210> 192  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 192  
aattgttctt ccactggc 18

<210> 193  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 193  
aagaaggagc acaatctt 18

<210> 194  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 194  
gaaaccaaatt taggataa 18

<210> 195  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 195

tgtagtgccta cctctttt

18

<210> 196

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 196

ctgaaatttt gattgaat

18

<210> 197

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 197

tacaatttca ataagtct

18

<210> 198

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 198

gggtctcagt atgctgcc

18

<210> 199

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 199

ccttcgatgt ataggaca

18

<210> 200

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 200

catgtcccta aaatgtca

18

<210> 201

<211> 2266

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (316)..(1602)

<400> 201

aattccgagc ttagggaaa cgcaggggcg gcttctaggt gctgccgccg ccaccgccac 60

caccacctcc accgccgct cggaaccag gcctgggggg cggtggggcc gcgtatggag 120

ccccgcccc ccggagctgc caacattgcc aacgccaccg ccacgtaca cacagcctca 180

acttcagga gaccgctcg tggccttatt tatccacc ttctgtaca tcgtagcgaa 240

tcaatccgtg gcgcgcact cctccgcac cctcttaac agtaccctg ggatggcgtg 300

agcactcccc cagcg atg gac cca tct gtg acg ctg tgg cag ttt ctg 348

Met Asp Pro Ser Val Thr Leu Trp Gln Phe Leu

1 5 10

ctg cag ctg ctg aga gag caa ggc aat ggc cac atc atc tcc tgg act 396

Leu Gln Leu Leu Arg Glu Gln Gly Asn Gly His Ile Ile Ser Trp Thr



15                      20                      25  
 tca cgg gat ggt ggt gaa ttc aag ctg gtg gat gca gag gag gtg gcc    444  
 Ser Arg Asp Gly Gly Glu Phe Lys Leu Val Asp Ala Glu Glu Val Ala  
      30                      35                      40  
 cgg ctg tgg gga cta cgc aag aac aag acc aac atg aat tac gac aag    492  
 Arg Leu Trp Gly Leu Arg Lys Asn Lys Thr Asn Met Asn Tyr Asp Lys  
      45                      50                      55  
 ctc agc cgg gcc ttg cgg tac tac tat gac aag aac atc atc cgc aag    540  
 Leu Ser Arg Ala Leu Arg Tyr Tyr Tyr Asp Lys Asn Ile Ile Arg Lys  
      60                      65                      70                      75  
 gtg agc ggc cag aag ttc gtc tac aag ttt gtg tcc tac cct gag gtc    588  
 Val Ser Gly Gln Lys Phe Val Tyr Lys Phe Val Ser Tyr Pro Glu Val  
              80                      85                      90  
 gca ggg tgc tcc act gag gac tgc ccg ccc cag cca gag gtg tct gtt    636  
 Ala Gly Cys Ser Thr Glu Asp Cys Pro Pro Gln Pro Glu Val Ser Val  
              95                      100                      105  
 acc tcc acc atg cca aat gtg gcc cct gct gct ata cat gcc gcc cca    684  
 Thr Ser Thr Met Pro Asn Val Ala Pro Ala Ala Ile His Ala Ala Pro  
              110                      115                      120  
 ggg gac act gtc tct gga aag cca ggc aca ccc aag ggt gca gga atg    732  
 Gly Asp Thr Val Ser Gly Lys Pro Gly Thr Pro Lys Gly Ala Gly Met  
              125                      130                      135  
 gca ggc cca ggc ggt ttg gca cgc agc agc cgg aac gag tac atg cgc    780  
 Ala Gly Pro Gly Gly Leu Ala Arg Ser Ser Arg Asn Glu Tyr Met Arg  
              140                      145                      150                      155  
 tcg ggc ctc tat tcc acc ttc acc atc cag tct ctg cag ccg cag cca    828  
 Ser Gly Leu Tyr Ser Thr Phe Thr Ile Gln Ser Leu Gln Pro Gln Pro  
              160                      165                      170  
 ccc cct cat cct cgg cct gct gtg gtg ctc ccc aat gca gct cct gca    876  
 Pro Pro His Pro Arg Pro Ala Val Val Leu Pro Asn Ala Ala Pro Ala  
              175                      180                      185  
 ggg gca gca gcg ccc ccc tcg ggg agc agg agc acc agt cca agc ccc    924  
 Gly Ala Ala Ala Pro Pro Ser Gly Ser Arg Ser Thr Ser Pro Ser Pro  
              190                      195                      200

ttg gag gcc tgt ctg gag gct gaa gag gcc ggc ttg cct ctg cag gtc 972  
 Leu Glu Ala Cys Leu Glu Ala Glu Glu Ala Gly Leu Pro Leu Gln Val  
 205 210 215

atc ctg acc ccg ccc gag gcc cca aac ctg aaa tcg gaa gag ctt aat 1020  
 Ile Leu Thr Pro Pro Glu Ala Pro Asn Leu Lys Ser Glu Glu Leu Asn  
 220 225 230 235

gtg gag ccg ggt ttg ggc cgg gct ttg ccc cca gaa gtg aaa gta gaa 1068  
 Val Glu Pro Gly Leu Gly Arg Ala Leu Pro Pro Glu Val Lys Val Glu  
 240 245 250

ggg ccc aag gaa gag ttg gaa gtt gcg ggg gag aga ggg ttt gtg cca 1116  
 Gly Pro Lys Glu Glu Leu Glu Val Ala Gly Glu Arg Gly Phe Val Pro  
 255 260 265

gaa acc acc aag gcc gag cca gaa gtc cct cca cag gag ggc gtg cca 1164  
 Glu Thr Thr Lys Ala Glu Pro Glu Val Pro Pro Gln Glu Gly Val Pro  
 270 275 280

gcc cgg ctg ccc gcg gtt gtt atg gac acc gca ggg cag gcg ggc ggc 1212  
 Ala Arg Leu Pro Ala Val Val Met Asp Thr Ala Gly Gln Ala Gly Gly  
 285 290 295

cat gcg gct tcc agc cct gag atc tcc cag ccg cag aag ggc cgg aag 1260  
 His Ala Ala Ser Ser Pro Glu Ile Ser Gln Pro Gln Lys Gly Arg Lys  
 300 305 310 315

ccc cgg gac cta gag ctt cca ctc agc ccg agc ctg cta ggt ggg ccg 1308  
 Pro Arg Asp Leu Glu Leu Pro Leu Ser Pro Ser Leu Leu Gly Gly Pro  
 320 325 330

gga ccc gaa cgg acc cca gga tcg gga agt ggc tcc ggc ctc cag gct 1356  
 Gly Pro Glu Arg Thr Pro Gly Ser Gly Ser Gly Ser Gly Leu Gln Ala  
 335 340 345

ccg ggg ccg gcg ctg acc cca tcc ctg ctt cct acg cat aca ttg acc 1404  
 Pro Gly Pro Ala Leu Thr Pro Ser Leu Leu Pro Thr His Thr Leu Thr  
 350 355 360

ccg gtg ctg ctg aca ccc agc tcg ctg cct cct agc att cac ttc tgg 1452  
 Pro Val Leu Leu Thr Pro Ser Ser Leu Pro Pro Ser Ile His Phe Trp  
 365 370 375

agc acc ctg agt ccc att gcg ccc cgt agc ccg gcc aag ctc tcc ttc 1500  
 Ser Thr Leu Ser Pro Ile Ala Pro Arg Ser Pro Ala Lys Leu Ser Phe

<400> 202  
gcaaggcaat ggccacat

<210> 203  
<211> 21  
<212> DNA  
<213> Artificial Sequence

<223> PCR Primer

<400> 203  
ctcctctgca tccaccagct t 21

<210> 204  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<223> PCR Probe

<400> 204  
tctcctggac ttcacgggat ggtggt 26

<210> 205  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 205  
cccctgcgtt tcctaca 18

<210> 206  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 206  
ggtggtggtg gcggtggc 18

<210> 207  
<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 207

ggcgttggca atgttggc

18

<210> 208

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 208

aagttgaggc tgtgtgta

18

<210> 209

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 209

aggccacgga cgggtctc

18

<210> 210

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 210

gattgattcg ctacgatg

18

<210> 211

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 211

gggatgcgga ggagtgcg

18

<210> 212

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 212

agtgtcacg ccatccca

18

<210> 213

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 213

aaactgccac agcgtcac

18

<210> 214

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 214

gaagtccagg agatgatg

18

<210> 215

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 215  
caccaccatc ccgtgaag 18

<210> 216  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 216  
tcttgttctt gcgtagtc 18

<210> 217  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 217  
tgttcttgtc atagtagt 18

<210> 218  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 218  
tcaccttgcg gatgatgt 18

<210> 219  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 219  
gagcaccctg cgacctca 18

<210> 220

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 220

ggcgggcagt cctcagtg

18

<210> 221

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 221

ggtgaagtg gaatagag

18

<210> 222

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 222

tccgattca ggtttggg

18

<210> 223

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 223

ttggtggtt ctggcaca

18

<210> 224



<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 224  
tggagggact tctggctc 18

<210> 225  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 225  
gcgtaggaag cagggatg 18

<210> 226  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 226  
gtgctccaga agtgaatg 18

<210> 227  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 227  
actggatgga aactggaa 18

<210> 228  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 228

ggccatccac gctgatag

18

<210> 229

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 229

ccaccacaat cagagcat

18

<210> 230

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 230

gatccccacc ccaccaca

18

<210> 231

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 231

tgttttctgt ggaggaga

18

<210> 232

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 232

aaacagagaa gttgtgga

18

<210> 233

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 233

gggactgaca gaaaacag

18

<210> 234

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 234

ataaataaat aaaccgcc

18

<210> 235

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 235

gttaggtcag gctcatcc

18

<210> 236

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 236 gtctcaagc cagacctc	18
<210> 237 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 237 aataaagaaa gaaaggtc	18
<210> 238 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 238 agggcaggct gagaaata	18
<210> 239 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 239 cttctactca catccaaa	18
<210> 240 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 240 caaaacaac taactctt	18

<210> 241  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 241  
ggaataataa aacaaac 18

<210> 242  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 242  
ttcttctgg acccctga 18

<210> 243  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 243  
ccaagggtgt gattcttc 18

<210> 244  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 244  
tgtctgagag aaaggttg 18

<210> 245

&lt;211&gt; 1080

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1080)

&lt;400&gt; 245

atg act ctg gag tcc atc atg gcg tgt tgc ctg agc gat gag gtg aag 48  
 Met Thr Leu Glu Ser Ile Met Ala Cys Cys Leu Ser Asp Glu Val Lys  
 1 5 10 15

gag tcc aag cgg atc aac gcc gag atc gag aag cag ctg cgg cgg gac 96  
 Glu Ser Lys Arg Ile Asn Ala Glu Ile Glu Lys Gln Leu Arg Arg Asp  
 20 25 30

aag cgc gac gcc cgg cgc gag ctc aag ctg ctg ctg ctc ggc acg ggc 144  
 Lys Arg Asp Ala Arg Arg Glu Leu Lys Leu Leu Leu Gly Thr Gly  
 35 40 45

gag agc ggg aag agc acg ttc atc aag cag atg cgc atc atc cac ggc 192  
 Glu Ser Gly Lys Ser Thr Phe Ile Lys Gln Met Arg Ile Ile His Gly  
 50 55 60

gcc ggc tac tgc gag gag gac aag cgc ggc ttc acc aag ctc gtc tac 240  
 Ala Gly Tyr Ser Glu Glu Asp Lys Arg Gly Phe Thr Lys Leu Val Tyr  
 65 70 75 80

cag aac atc ttc acc gcc atg cag gcc atg atc cgg gcc atg gag acg 288  
 Gln Asn Ile Phe Thr Ala Met Gln Ala Met Ile Arg Ala Met Glu Thr  
 85 90 95

ctc aag atc ctc tac aag tac gag cag aac aag gcc aat gcg ctc ctg 336  
 Leu Lys Ile Leu Tyr Lys Tyr Glu Gln Asn Lys Ala Asn Ala Leu Leu  
 100 105 110

atc cgg gag gtg gac gtg gag aag gtg acc acc ttc gag cat cag tac 384  
 Ile Arg Glu Val Asp Val Glu Lys Val Thr Thr Phe Glu His Gln Tyr  
 115 120 125

gtc agt gcc atc aag acc ctg tgg gag gac cgg ggc atc cag gaa tgc 432  
 Val Ser Ala Ile Lys Thr Leu Trp Glu Asp Pro Gly Ile Gln Glu Cys  
 130 135 140

tac gac cgc agg cgc gag tac cag ctc tcc gac tct gcc aag tac tac 480  
 Tyr Asp Arg Arg Arg Glu Tyr Gln Leu Ser Asp Ser Ala Lys Tyr Tyr  
 145 150 155 160

ctg acc gac gtt gac cgc atc gcc acc ttg ggc tac ctg ccc acc cag 528  
 Leu Thr Asp Val Asp Arg Ile Ala Thr Leu Gly Tyr Leu Pro Thr Gln  
 165 170 175

cag gac gtg ctg cgg gtc cgc gtg ccc acc acc ggc atc atc gag tac 576  
 Gln Asp Val Leu Arg Val Arg Val Pro Thr Thr Gly Ile Ile Glu Tyr  
 180 185 190

cct ttc gac ctg gag aac atc atc ttc cgg atg gtg gat gtg ggg ggc 624  
 Pro Phe Asp Leu Glu Asn Ile Ile Phe Arg Met Val Asp Val Gly Gly  
 195 200 205

cag cgg tcg gag cgg agg aag tgg atc cac tgc ttt gag aac gtg aca 672  
 Gln Arg Ser Glu Arg Arg Lys Trp Ile His Cys Phe Glu Asn Val Thr  
 210 215 220

tcc atc atg ttt ctc gtc gcc ctc agc gaa tac gac caa gtc ctg gtg 720  
 Ser Ile Met Phe Leu Val Ala Leu Ser Glu Tyr Asp Gln Val Leu Val  
 225 230 235 240

gag tcg gac aac gag aac cgg atg gag gag agc aaa gcc ctg ttc cgg 768  
 Glu Ser Asp Asn Glu Asn Arg Met Glu Glu Ser Lys Ala Leu Phe Arg  
 245 250 255

acc atc atc acc tac ccc tgg ttc cag aac tcc tcc gtc atc ctc ttc 816  
 Thr Ile Ile Thr Tyr Pro Trp Phe Gln Asn Ser Ser Val Ile Leu Phe  
 260 265 270

ctc aac aag aag gac ctg ctg gag gac aag atc ctg tac tcg cac ctg 864  
 Leu Asn Lys Lys Asp Leu Leu Glu Asp Lys Ile Leu Tyr Ser His Leu  
 275 280 285

gtg gac tac ttc ccc gag ttc gat ggt ccc cag cgg gac gcc cag gcg 912  
 Val Asp Tyr Phe Pro Glu Phe Asp Gly Pro Gln Arg Asp Ala Gln Ala  
 290 295 300

gcg cgg gag ttc atc ccg aag atg ttc gtg gac ctg aac ccc gac agc 960  
 Ala Arg Glu Phe Ile Pro Lys Met Phe Val Asp Leu Asn Pro Asp Ser  
 305 310 315 320

gac aag atc atc tac tca cac ttc acg tgt gcc acc gac acg gag aac 1008  
 Asp Lys Ile Ile Tyr Ser His Phe Thr Cys Ala Thr Asp Thr Glu Asn

325                      330                      335

atc cgc ttc gtg ttc gcg gcc gtg aag gac acc atc ctg cag ctg aac    1056  
 Ile Arg Phe Val Phe Ala Ala Val Lys Asp Thr Ile Leu Gln Leu Asn  
           340                      345                      350

ctg aag gag tac aat ctg gtc taa                                      1080  
 Leu Lys Glu Tyr Asn Leu Val  
           355

<210> 246  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<223> PCR Primer

<400> 246  
 tgaccacctt cgagcatcag                                      20

<210> 247  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<223> PCR Primer

<400> 247  
 cggtcgtagc attcctggat                                      20

<210> 248  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<223> PCR Probe

<400> 248  
 tcagtgccat caagaccctg tgggag                                      26

<210> 249  
 <211> 18



<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 249  
gatggactcc agagtcac 18

<210> 250  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 250  
gccatgatgg actccaga 18

<210> 251  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 251  
cacgccatga tggactcc 18

<210> 252  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 252  
ctcatcgctc aggcaaca 18

<210> 253  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 253

cttcacctca tcgctcag

18

<210> 254

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 254

gactccttca cctcatcg

18

<210> 255

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 255

atccgcttgg actccttc

18

<210> 256

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 256

cgttgatccg cttggact

18

<210> 257

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 257  
ctcgatctcg gcgttgat 18

<210> 258  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 258  
cccgccgcag ctgcttct 18

<210> 259  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 259  
cttgagctcg cgccgggc 18

<210> 260  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 260  
gcagcagcag ctgagct 18

<210> 261  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 261  
gcccgtgccg agcagcag 18

<210> 262

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 262

acgtgctctt cccgctct

18

<210> 263

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 263

atctgcttga tgaacgtg

18

<210> 264

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 264

cgcacgtgct tgatgaac

18

<210> 265

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 265

gtagccggcg ccgtggat

18

<210> 266

<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 266  
tgtcctctc cgagtagc 18

<210> 267  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 267  
cttgctctcc tccgagta 18

<210> 268  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 268  
aagccgcgct tgtctctc 18

<210> 269  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 269  
tagacgagct tggtagaag 18

<210> 270  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 270

tgttctggta gacgagct

18

<210> 271

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 271

tggcggtgaa gatgttct

18

<210> 272

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 272

cggatcatgg cctgcatg

18

<210> 273

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 273

cgtctccatg gcccgat

18

<210> 274

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 274

tagaggatct tgagcgtc 18

<210> 275

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 275

tgtagaggat cttagcgc 18

<210> 276

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 276

tgctcgtagc tgtagagg 18

<210> 277

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 277

gccttggtct gctcgtac 18

<210> 278

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 278 ttggccttgt tctgctcg	18
<210> 279 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 279 caggagcgca ttggcctt	18
<210> 280 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 280 ctccacgtcc acctcccg	18
<210> 281 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 281 ggtcaccttc tccacgtc	18
<210> 282 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 282 gatgctcgaa ggtggtca	18



<210> 283  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 283  
actgacgtac tgatgctc 18

<210> 284  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 284  
cttgatggca ctgacgta 18

<210> 285  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 285  
cagggtcttg atggcact 18

<210> 286  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 286  
ctggatgccc ggtcctc 18

<210> 287

<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 287  
tcctggatgc ccgggtcc 18

<210> 288  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 288  
cgctgcggt cgtagcat 18

<210> 289  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 289  
gctggctac gcgcctgc 18

<210> 290  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 290  
tactggcag agtcggag 18

<210> 291  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 291

gtcaggtagt acttgga

18

<210> 292

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 292

ggtcaacgtc ggtcaggt

18

<210> 293

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 293

gtggcgatgc ggtcaacg

18

<210> 294

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 294

gcaggtagcc caaggtgg

18

<210> 295

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 295

cgtcctgctg ggtgggca

18

<210> 296

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 296

ggtggtgggc acgcggac

18

<210> 297

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 297

tcgatgatgc cgttggtg

18

<210> 298

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 298

ccaggtcgaa aggttact

18

<210> 299

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 299 tgttctccag gtcgaaag	18
<210> 300 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 300 agatgatggt ctccaggt	18
<210> 301 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 301 atccggaaga tgatgttc	18
<210> 302 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 302 ctccgctccg accgctgg	18
<210> 303 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 303 gatccacttc ctccgctc	18

<210> 304

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 304

tgtcacgttc tcaaagca

18

<210> 305

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 305

atgatggatg tcacgttc

18

<210> 306

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 306

cgagaaacat gatggatg

18

<210> 307

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 307

gctgagggcg acgagaaa

18

<210> 308

<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 308  
cgactccacc aggacttg 18

<210> 309  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 309  
atccggttct cgtgtcc 18

<210> 310  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 310  
ccatccggtt ctcgttg 18

<210> 311  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 311  
agggtttgc tctctcc 18

<210> 312  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 312

ggtccggaac agggcttt

18

<210> 313

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 313

gtagtgatg atggtccg

18

<210> 314

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 314

ggaggagttc tggaacca

18

<210> 315

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 315

tgaggaagag gatgacgg

18

<210> 316

<211> 18

<212> DNA

<213> Artificial Sequence



<223> Antisense Oligonucleotide

<400> 316

gcaggtcctt cttgttga

18

<210> 317

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 317

atcttgctct ccagcagg

18

<210> 318

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 318

gcgagtacag gatcttgt

18

<210> 319

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 319

aagtagtcca ccaggtgc

18

<210> 320

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 320  
gatgaactcc cgcgccgc 18

<210> 321  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 321  
ggttcaggtc cacgaaca 18

<210> 322  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 322  
gtagatgac ttgtcgct 18

<210> 323  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 323  
cacgtgaagt gtgagtag 18

<210> 324  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 324  
atgttctccg tgcggtg 18

<210> 325

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 325

acggccgcga acacgaag

18

<210> 326

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 326

gatggtgtcc ttcacggc

18

<210> 327

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 327

tcaggttcag ctgcagga

18

<210> 328

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 328

accagattgt actccttc

18

<210> 329

&lt;211&gt; 2610

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (199)..(1641)

&lt;400&gt; 329

atcctgggac agggcacagg gccatctgtc accaggggct tagggaaggc cgagccagcc 60

tgggtcaaag aagtcaaagg ggctgcctgg aggaggcagc ctgtcagctg gtgcatcaga 120

ggctgtggcc aggccagctg ggctcgggga gcgccagcct gagaggagcg cgtgagcgtc 180

gcgggagcct cgggcacc atg agc gac gtg gct att gtg aag gag ggt tgg 231

Met Ser Asp Val Ala Ile Val Lys Glu Gly Trp

1 5 10

ctg cac aaa cga ggg gag tac atc aag acc tgg cgg cca cgc tac ttc 279

Leu His Lys Arg Gly Glu Tyr Ile Lys Thr Trp Arg Pro Arg Tyr Phe

15 20 25

ctc ctc aag aat gat ggc acc ttc att ggc tac aag gag cgg ccg cag 327

Leu Leu Lys Asn Asp Gly Thr Phe Ile Gly Tyr Lys Glu Arg Pro Gln

30 35 40

gat gtg gac caa cgt gag gct ccc ctc aac aac ttc tct gtg gcg cag 375

Asp Val Asp Gln Arg Glu Ala Pro Leu Asn Asn Phe Ser Val Ala Gln

45 50 55

tgc cag ctg atg aag acg gag cgg ccc cgg ccc aac acc ttc atc atc 423

Cys Gln Leu Met Lys Thr Glu Arg Pro Arg Pro Asn Thr Phe Ile Ile

60 65 70 75

cgc tgc ctg cag tgg acc act gtc atc gaa cgc acc ttc cat gtg gag 471

Arg Cys Leu Gln Trp Thr Thr Val Ile Glu Arg Thr Phe His Val Glu

80 85 90

act cct gag gag cgg gag gag tgg aca acc gcc atc cag act gtg gct 519

Thr Pro Glu Glu Arg Glu Glu Trp Thr Thr Ala Ile Gln Thr Val Ala

95 100 105

gac ggc ctc aag aag cag gag gag gag gag atg gac ttc cgg tcg ggc 567

Asp Gly Leu Lys Lys Gln Glu Glu Glu Glu Met Asp Phe Arg Ser Gly

110                      115                      120  
 tca ccc agt gac aac tca ggg gct gaa gag atg gag gtg tcc ctg gcc    615  
 Ser Pro Ser Asp Asn Ser Gly Ala Glu Glu Met Glu Val Ser Leu Ala  
 125                      130                      135  
 aag ccc aag cac cgc gtg acc atg aac gag ttt gag tac ctg aag ctg    663  
 Lys Pro Lys His Arg Val Thr Met Asn Glu Phe Glu Tyr Leu Lys Leu  
 140                      145                      150                      155  
 ctg ggc aag ggc act ttc ggc aag gtg atc ctg gtg aag gag aag gcc    711  
 Leu Gly Lys Gly Thr Phe Gly Lys Val Ile Leu Val Lys Glu Lys Ala  
 160                      165                      170  
 aca ggc cgc tac tac gcc atg aag atc ctc aag aag gaa gtc atc gtg    759  
 Thr Gly Arg Tyr Tyr Ala Met Lys Ile Leu Lys Lys Glu Val Ile Val  
 175                      180                      185  
 gcc aag gac gag gtg gcc cac aca ctc acc gag aac cgc gtc ctg cag    807  
 Ala Lys Asp Glu Val Ala His Thr Leu Thr Glu Asn Arg Val Leu Gln  
 190                      195                      200  
 aac tcc agg cac ccc ttc ctc aca gcc ctg aag tac tct ttc cag acc    855  
 Asn Ser Arg His Pro Phe Leu Thr Ala Leu Lys Tyr Ser Phe Gln Thr  
 205                      210                      215  
 cac gac cgc ctc tgc ttt gtc atg gag tac gcc aac ggg ggc gag ctg    903  
 His Asp Arg Leu Cys Phe Val Met Glu Tyr Ala Asn Gly Gly Glu Leu  
 220                      225                      230                      235  
 ttc ttc cac ctg tcc cgg gaa cgt gtg ttc tcc gag gac cgg gcc cgc    951  
 Phe Phe His Leu Ser Arg Glu Arg Val Phe Ser Glu Asp Arg Ala Arg  
 240                      245                      250  
 ttc tat ggc gct gag att gtg tca gcc ctg gac tac ctg cac tcg gag    999  
 Phe Tyr Gly Ala Glu Ile Val Ser Ala Leu Asp Tyr Leu His Ser Glu  
 255                      260                      265  
 aag aac gtg gtg tac cgg gac ctc aag ctg gag aac ctc atg ctg gac    1047  
 Lys Asn Val Val Tyr Arg Asp Leu Lys Leu Glu Asn Leu Met Leu Asp  
 270                      275                      280  
 aag gac ggg cac att aag atc aca gac ttc ggg ctg tgc aag gag ggg    1095  
 Lys Asp Gly His Ile Lys Ile Thr Asp Phe Gly Leu Cys Lys Glu Gly  
 285                      290                      295

atc aag gac ggt gcc acc atg aag acc ttt tgc ggc aca cct gag tac 1143  
 Ile Lys Asp Gly Ala Thr Met Lys Thr Phe Cys Gly Thr Pro Glu Tyr  
 300 305 310 315

ctg gcc ccc gag gtg ctg gag gac aat gac tac ggc cgt gca gtg gac 1191  
 Leu Ala Pro Glu Val Leu Glu Asp Asn Asp Tyr Gly Arg Ala Val Asp  
 320 325 330

tgg tgg ggg ctg ggc gtg gtc atg tac gag atg atg tgc ggt cgc ctg 1239  
 Trp Trp Gly Leu Gly Val Val Met Tyr Glu Met Met Cys Gly Arg Leu  
 335 340 345

ccc ttc tac aac cag gac cat gag aag ctt ttt gag ctc atc ctc atg 1287  
 Pro Phe Tyr Asn Gln Asp His Glu Lys Leu Phe Glu Leu Ile Leu Met  
 350 355 360

gag gag atc cgc ttc ccg cgc acg ctt ggt ccc gag gcc aag tcc ttg 1335  
 Glu Glu Ile Arg Phe Pro Arg Thr Leu Gly Pro Glu Ala Lys Ser Leu  
 365 370 375

ctt tca ggg ctg ctc aag aag gac ccc aag cag agg ctt ggc ggg ggc 1383  
 Leu Ser Gly Leu Leu Lys Lys Asp Pro Lys Gln Arg Leu Gly Gly Gly  
 380 385 390 395

tcc gag gac gcc aag gag atc atg cag cat cgc ttc ttt gcc ggt atc 1431  
 Ser Glu Asp Ala Lys Glu Ile Met Gln His Arg Phe Phe Ala Gly Ile  
 400 405 410

gtg tgg cag cac gtg tac gag aag aag ctc agc cca ccc ttc aag ccc 1479  
 Val Trp Gln His Val Tyr Glu Lys Lys Leu Ser Pro Pro Phe Lys Pro  
 415 420 425

cag gtc acg tcg gag act gac acc agg tat ttt gat gag gag ttc acg 1527  
 Gln Val Thr Ser Glu Thr Asp Thr Arg Tyr Phe Asp Glu Glu Phe Thr  
 430 435 440

gcc cag atg atc acc atc aca cca cct gac caa gat gac agc atg gag 1575  
 Ala Gln Met Ile Thr Ile Thr Pro Pro Asp Gln Asp Asp Ser Met Glu  
 445 450 455

tgt gtg gac agc gag cgc agg ccc cac ttc ccc cag ttc tcc tac tcg 1623  
 Cys Val Asp Ser Glu Arg Arg Pro His Phe Pro Gln Phe Ser Tyr Ser  
 460 465 470 475

gcc agc agc acg gcc tga ggcggcgggtg gactgcgctg gacgatagct 1671  
 Ala Ser Ser Thr Ala

480

tggagggatg gagaggcggc ctctgcat gatctgtatt taatggttt tattctcgg 1731  
 gtgcattga gagaagccac gctgtcctct cgagcccaga tggaaagacg ttttgtgct 1791  
 gtgggcagca cctcccccg cagcggggta gggaagaaaa ctatcctgcg ggtttaatt 1851  
 tattcatcc agttgttct ccgggtgtgg ctcagccct cagaacaac cgattcacgt 1911  
 agggaaatgt taaggacttc tacagctatg cgcaatgtgg cattgggggg ccgggcaggt 1971  
 cctgccatg tgtccctca ctctgcagc cagccgccct gggctgtctg tcaccagcta 2031  
 tctgtcatct ctctggggcc ctgggctca gttcaacctg gtggcaccag atgcaacctc 2091  
 actatggtat gctggccagc accctctct ggggggtggca ggcacacagc agccccccag 2151  
 cactaaggcc gtgtctctga ggacgtcatc ggaggctggg ccctgggat gggaccaggg 2211  
 atgggggatg ggccaggggt taccagtg gacagaggag caaggttaa attgttatt 2271  
 gtgtattatg ttgtcaaat gcattttggg ggttttaat cttgtgaca ggaaagccct 2331  
 ccccttccc cttctgtgc acagttcttg gtgactgtcc caccggagcc tccccctag 2391  
 atgatcttc caggttagca cttgacctt tcgacgcta accctccgc tgtcgccca 2451  
 ggccctcct gactccctgt ggggggtggc atccctgggc cctccacgc ctctggcca 2511  
 gacgtgccg ctgccgtgc accacggcgt tttttacaa cattcaact tagtatttt 2571  
 actattataa tataatatg aaccttcct ccaattct 2610

&lt;210&gt; 330

&lt;211&gt; 21

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;223&gt; PCR Primer

&lt;400&gt; 330

cgtgaccatg aacgagtttg a

21

<210> 331  
<211> 19  
<212> DNA  
<213> Artificial Sequence

<223> PCR Primer

<400> 331  
caggatcacc ttgccgaaa 19

<210> 332  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<223> PCR Probe

<400> 332  
ctgaagctgc tgggcaaggg ca 22

<210> 333  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 333  
ccctgtgcc tgtcccag 18

<210> 334  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 334  
cctaagcccc tggtgaca 18

<210> 335  
<211> 18



<212> DNA	
<213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 335	
ctttgacttc ttgaccc	18
<210> 336	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 336	
ggcagcccct ttgacttc	18
<210> 337	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 337	
caaccctcct tcacaata	18
<210> 338	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 338	
tactcccctc gtttgtgc	18
<210> 339	
<211> 18	
<212> DNA	
<213> Artificial Sequence	

<223> Antisense Oligonucleotide

<400> 339

tgccatcatt ctgagga 18

<210> 340

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 340

agccaatgaa ggtgcat 18

<210> 341

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 341

cacagagaag ttgtgag 18

<210> 342

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 342

agtctggatg gcggtgt 18

<210> 343

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 343 tcctcctcct cctgcttc	18
<210> 344 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 344 cctgagttgt cactgggt	18
<210> 345 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 345 ccgaaagtgc ccttgccc	18
<210> 346 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 346 gccacgatga cttccttc	18
<210> 347 <211> 18 <212> DNA <213> Artificial Sequence	
<223> Antisense Oligonucleotide	
<400> 347 cggtcctcgg agaacaca	18

<210> 348  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 348  
acgttcttct ccgagtgc 18

<210> 349  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 349  
gtgccgcaaa aggtcttc 18

<210> 350  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 350  
tactcaggtg tgccgcaa 18

<210> 351  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 351  
ggcttgaagg gtgggctg 18

<210> 352

<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 352  
tcaaaatacc tgggtca 18

<210> 353  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 353  
gccgtgaact cctcatca 18

<210> 354  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 354  
ggtcaggtgg tgtgatgg 18

<210> 355  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 355  
ctcgctgtcc acacactc 18

<210> 356  
<211> 18  
<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 356

gcctctccat ccctccaa

18

<210> 357

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 357

acagcgtggc ttctctca

18

<210> 358

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 358

ttttctccc taccgccg

18

<210> 359

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 359

gatagtttc ttcctac

18

<210> 360

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 360

taaaacccgc aggatagt

18

<210> 361

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 361

ggagaacaaa ctggatga

18

<210> 362

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 362

ctggctgaca gaggtagg

18

<210> 363

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 363

gcggctggct gacagagt

18

<210> 364

<211> 18

<212> DNA

<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 364  
cccagagaga tgacagat 18

<210> 365  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 365  
gctgctgtgt gcctgcca 18

<210> 366  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 366  
cataatacac aataacaa 18

<210> 367  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 367  
atttgaacaa cataatac 18

<210> 368  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<223> Antisense Oligonucleotide

<400> 368  
aagtgtacc gtggagag 18



<210> 369  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 369  
cgaaaaggctc aagtgcta 18

<210> 370  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 370  
cagggagtca gggagggc 18

<210> 371  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 371  
aaagttgaat gttgtaaa 18

<210> 372  
<211> 18  
<212> DNA  
<213> Artificial Sequence  
  
<223> Antisense Oligonucleotide  
  
<400> 372  
aaaatactaa agttgaat 18

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/08268

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) :C12Q 1/68

US CL :435/6

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 422/50,68.1; 435/5,6,7.1,7.2; 436/501; 514/2,44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,612,455 A (HOEY) 18 March 1997, see especially the abstract, summary of the invention in columns 1-2, and column 7, line 45, through column 12, line 4.	1-54
Y	US 4,806,463 A (GOODCHILD et al.) 21 February 1989, see entire disclosure.	1-54
Y	GHOSH et al. Evaluation of some properties of a phosphorodithioate oligodeoxyribonucleotide for antisense application. Nucleic Acids Research. 1993, Volume 21, Number 24, pages 5761-5766, see entire disclosure.	1-54

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

02 JULY 1999

Date of mailing of the international search report

16 AUG 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

ARDIN MARSCHER

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/08268

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MIRABELLI et al. In vitro and in vivo pharmacologic activities of antisense oligonucleotides. Anti-Cancer Drug Design. December 1991, Volume 6, pages 647-661, see entire disclosure.	1-54
Y	US 5,436,327 A (SOUTHERN) 25 July 1995, see entire disclosure.	1-54
Y	US 5,700,637 A (SOUTHERN) 23 December 1997, see entire disclosure.	1-54
Y	WO 89/10977 A1 (ISIS INNOVATION LIMITED) 16 November 1989, see entire disclosure.	1-54
Y	UHLMANN et al. Antisense Oligonucleotides: A New Therapeutic Principle. Chemical Reviews. June 1990, Volume 90, Number 4, pages 543-584, see entire disclosure.	1-54
P,Y	US 5,783,431 A (PETERSON et al.) 21 July 1998, see entire disclosure.	1-54
P,Y	US 5,824,485 A (THOMPSON et al.) 20 October 1998, see entire disclosure.	1-54

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/08268

### B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, EMBASE, WPI, BIOTECH ABS., CAS, MEDLINE using search terms: robot, screen, drug, antisense, silico, solid, support, substrate, library, array, matrix, and express