



(51) International Patent Classification:

A61K 48/00 (2006.01) A61K 47/46 (2006.01)

(21) International Application Number:

PCT/IB2017/056131

(22) International Filing Date:

04 October 2017 (04.10.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/405,074 06 October 2016 (06.10.2016) US

(71) Applicant: **ENGENEIC MOLECULAR DELIVERY PTY LTD** [AU/AU]; Building 2, 25 Sirius Road, Lane Cove West, Sydney, New South Wales 2066 (AU).

(72) Inventors: **BRAHMBHATT, Himanshu**; c/o Engeneic Molecular Delivery PTY LTD, Building 2, 25 Sirius Road, Lane Cove West, Sydney, New South Wales 2066 (AU). **MACDIARMID, Jennifer**; c/o Engeneic Molecular Delivery PTY LTD, Building 2, 25 Sirius Road, Lane Cove West, Sydney, New South Wales 2066 (AU).

(81) Designated States (unless otherwise indicated, for every kind of national protection available):


AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: BACTERIAL MINICELLS FOR DELIVERING NUCLEIC ACID ADJUVANTS AND METHODS OF USING THE SAME

Figure 1



(57) Abstract: This disclosure provides intact bacterially derived minicells containing nucleic acids adjuvants or plasmids encoding nucleic acids adjuvants that can produce a desired immune response in target cells. This disclosure further provides methods that employ minicells to deliver nucleic acids adjuvants for use in the treatment of diseases, including neoplastic disease and cancer.

WO 2018/065923 A1

**Declarations under Rule 4.17:**

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

**Published:**

- *with international search report (Art. 21(3))*

## **BACTERIAL MINICELLS FOR DELIVERING NUCLEIC ACID ADJUVANTS AND METHODS OF USING THE SAME**

### **CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims priority from U.S. Provisional Patent Application No. 62/405,074, filed October 6, 2016. The content of that application is incorporated herein by reference in its entirety.

### **Field of Invention**

[0002] The present disclosure relates generally to the field of nucleic acid-based therapy, and, in particular, to delivery of nucleic acid adjuvants to mammalian target cells (*e.g.*, T cells, dendritic cells) using bacterially-derived, intact minicells. The disclosure has particular utility for treating cancer, especially in the context of promoting anti-tumor immune responses, and infections (*e.g.*, viral infections).

### **Background of the Invention**

[0003] The following discussion is merely provided to aid the reader in understanding the disclosure and is not admitted to describe or constitute prior art thereto.

[0004] Recent advances have identified a number of nucleic acids that are sensed by receptors in mammalian cells. The engagement of nucleic acid receptors activates the innate immune system in multiple ways. For instance, the binding of certain nucleic acids (*i.e.*, nucleic acid adjuvants) to cognate receptors results in triggering Type I and Type II interferons (IFNs). These interferons can function as adjuvants in augmenting anti-tumor activity by the immune system (reviewed in Junt and Barchet, 2015).

[0005] Aside from triggering cell-intrinsic and IFN-mediated effector mechanisms, nucleic acid sensor agonists can activate dendritic cells (DCs), promoting cytokine secretion, maturation and antigen presentation. This, in turn, enhances and shapes the quality of adaptive immune responses. Owing to their immune-enhancing properties, oligonucleotide or small-molecule agonists of nucleic acid sensors are being used in clinical trials to boost the immune response against poorly immunogenic cancers, and as adjuvants in therapeutic immunizations against cancer or in prophylactic vaccines against infections.

[0006] A number of agonists of nucleic acid sensors have been discovered and tested as adjuvants to provoke anti-viral immunity or enhance anti-tumor activity. Despite the promise of these agonists of nucleic acid sensors as adjuvants, few have actually entered clinical development.

[0007] This may be due to several factors. In particular, nucleic acid agonists need to be delivered into specific cells, such as dendritic cells, but these molecules do not have homing properties to assist in locating or entering the target cells. Additionally, when administered *in vivo*, free nucleic acids are unstable in serum and can be rapidly degraded by nucleases. As a result, nucleic acid adjuvants and agonists of nucleic acid sensors have been relegated to topical use, or must be covalently linked to protein antigens.

[0008] Thus, there is a clear need in the art for a delivery system capable of delivering therapeutic nucleic acid adjuvants and agonists of nucleic acid sensors to target cells. The present disclosure provides such a delivery system.

### Summary of the Invention

[0009] Described herein are compositions and methods for treating diseases using bacterially derived, intact minicells comprising nucleic acid adjuvants or agonists of nucleic acid sensors. In general, the disclosed minicell delivery vectors function by transporting the nucleic acid adjuvants or agonists of nucleic acid sensors to target cells, such as T cells or dendritic cells, to elicit an immune response to help fight various diseases, including cancer and/or infections.

[0010] In one aspect, the present disclosure provides compositions comprising (a) an intact minicell that comprising at least one nucleic acid adjuvant molecule, a plasmid comprising a segment that encodes at least one nucleic acid adjuvant molecule, or an agonist of a nucleic acid sensor, and (b) a pharmaceutically acceptable carrier therefore, wherein the at least one nucleic acid adjuvant molecule or agonist of a nucleic acid receptor triggers an immune response from a target cell.

[0011] In another aspect, the present disclosure provides methods of delivering a nucleic acid adjuvant or an agonist of a nucleic acid sensor to a target cell, comprising contacting a target cell with an intact minicell comprising (i) at least one nucleic acid adjuvant molecule, (ii) a plasmid

comprised of a segment that encodes at least one nucleic acid adjuvant molecule, or (iii) at least one agonist of a nucleic acid sensor, wherein the target cell engulfs the minicell.

[0012] In yet another aspect, the present disclosure provides methods of treating a disease in a subject, comprising administering to a subject with a disease an intact minicell comprising (i) at least one nucleic acid adjuvant molecule, (ii) a plasmid comprising a segment that encodes at least one nucleic acid adjuvant molecule, or (iii) at least one agonist of a nucleic acid receptor, wherein the minicell is engulfed by a target cell following administration.

[0013] In some embodiments, the immune response produced in the target cells comprises the production of Type I interferon, including interferon- $\alpha$  and/or interferon- $\beta$ .

[0014] In some embodiments, the at least one nucleic acid adjuvant comprises a nucleic acid that binds to at least one of TLR3, TLR7, TLR8, TLR9, RIG-I, MDA5, AIM2, cGAS, or IFI16.

[0015] In some embodiment, the intact minicell comprises at least two nucleic acid adjuvants. In other embodiments, the at least one nucleic acid adjuvant comprises a sequence of at least about 40 nucleotides, for example, the at least one nucleic acid adjuvant may be a 40-mer or a 50-mer double stranded RNA or DNA.

[0016] In some embodiments, the intact minicell comprises a nucleic acid adjuvant and an agonist of a nucleic acid sensor, and in some embodiments, the at least one agonist of a nucleic acid sensor comprises a polynucleotide product of PNPase1, poly(I:C), poly-ICLC, imiquimod, imidazoquinoline resiquimod, CpG-ODNs or 2'3' cyclic GAMP (GMP-AMP), .

[0017] In some embodiments, the disclosed minicells may further comprise a bispecific ligand. In some embodiments, the bispecific ligand may comprise a first arm that carries specificity for a minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor. For example, the minicell surface structure may be an O-polysaccharide component of a lipopolysaccharide on the minicell surface and the mammalian cell surface receptor may be capable of activating receptor-mediated endocytosis or macropinocytosis of the minicell. In some embodiments, the bispecific ligand comprises an antibody or antibody fragment.

[0018] In some embodiments, the composition comprises fewer than about 1 contaminating parent bacterial cell per  $10^7$  minicells, per  $10^8$  minicells, per  $10^9$  minicells, per  $10^{10}$  minicells, or per  $10^{11}$  minicells.

[0019] In some embodiments, the plasmid encoding a nucleic acid adjuvant comprises a regulatory element operably linked to the segment that encodes at least one nucleic acid adjuvant, and in some embodiments, the plasmid encodes multiple nucleic acid adjuvant molecules.

[0020] In some embodiments, the target cell is a mammalian cell, such as a human immune cell. For example, the target cell can be a monocytic cell, a macrophage, a T cell, or a dendritic cell or a NK cell or a iNKT cell. In some embodiments, the target cell is phagocytosis- or endocytosis- or macropinocytosis competent.

[0021] In some embodiments, the contact between the minicell and the mammalian cell can occur *in vitro* or *in vivo*.

[0022] In some embodiments, the disease being treated can be cancer or an infection.

[0023] In some embodiments, the administration of the disclosed minicell can further comprise administering a drug for the treatment of the disease to the subject, such as a chemotherapeutic or an anti-viral or an anti-bacterial. In these embodiments, the drug may be packaged in an intact minicell as well. In some embodiments, treatment may further comprise radiation treatment.

[0024] The foregoing general description and following detailed description are exemplary and explanatory and not limiting of the disclosure.

#### **Brief Description of the Drawings**

[0025] Figure 1 shows endosomal and cytosolic nucleic acid-sensing pathways (adapted from Junt and Barchet, 2015). The figure indicates some of the major nucleic acid-sensing pathways described herein. Abbreviations include: cGAMP = cyclic GMP-AMP; cGAS = cGAMP synthase; DC = dendritic cell; dsRNA = double-stranded RNA; ER = endoplasmic reticulum; IL = interleukin; IRAK = IL-1 receptor-associated kinase; IRF = IFN-regulatory factor; MAVS = mitochondrial antiviral signaling protein; MDA5 = melanoma differentiation-associated protein

5; MYD88 = myeloid differentiation primary response protein 88; NF- $\kappa$ B = nuclear factor- $\kappa$ B; pDC = plasmacytoid DC; RIG-I = retinoic acid-inducible gene I; ssRNA = single-stranded RNA; STING = stimulator of IFN genes; TBK1 = TANK-binding kinase 1; TRIF = TIR domain-containing adaptor protein inducing IFN $\beta$ ; and XCR1 = XC-chemokine receptor 1.

[0026] Figure 2 shows a nucleic acid-sensing pathway in the nucleus (adapted from Diner, Lumm, and Christea, 2015). Following entry of double stranded foreign DNA into a host cell, it binds directly to DNA sensors IFI16 and IFIX (1). IFI16 signals to STING via a mechanism that has yet to be elucidated (2). Upon activation and dimerization of STING, TBK-1 is phosphorylated (3), resulting in the phosphorylation of IRF3 and NF- $\kappa$ B (4), which translocate back into the nucleus to induce the expression of cytokines (5).

[0027] Figure 3 shows the results of an *in vivo* study comparing the efficacy of drug-packaged minicells alone to a combination treatment of drug- (PNU159682) and nucleic acid adjuvant- (40-mer) packaged minicells in a mouse xenograft tumor model. As shown in the figure, mice receiving  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells<sub>40-mer</sub> (Group 3) showed highly significant tumor regression by day 44 after a total of 6 doses, while mice receiving  $^{EGFR}$ minicells<sub>PNU-159682</sub> alone (Group 2) were merely stable. Group 1 represents the control mice that received only saline.

[0028] Figure 4 shows results from a comparison of anti-tumor efficacy of nucleic acid adjuvant 40-mers and 50-mers in a mouse xenograft model. Group 1 (control) received sterile physiological saline. The other groups were administered as follows: Group 2  $^{EGFR}$ minicells<sub>PNU-159682</sub>, Group 3  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells<sub>40-mer</sub>, Group 4  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells<sub>50-mer</sub>, Group 5  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells.

### Detailed Description

[0029] The present disclosure provides novel minicell-based delivery vectors of nucleic acid adjuvants and/or agonists of nucleic acid sensors and methods of using the same. In particular, the disclosure has utility for enhancing a beneficial immune response to fight a disease in a patient, such as a neoplastic disease (*e.g.*, cancer) or an infection. For example, the disclosed minicell delivery vectors can enhance anti-tumor efficacy when administering the disclosed minicell delivery vectors either alone or in combination with a bispecific ligand targeted, cytotoxic drug or siRNA or miRNA-packaged minicells in the same patient.

[0030] Previously, the present inventors discovered that therapeutically significant concentrations of siRNA or miRNA (~23 nucleotides in length) could be successfully packaged in intact minicells. This was unexpected because these molecules are large (~ 14,000 daltons) and it was not known that such large molecules could enter into intact minicells via the double outer membrane. Further, once packaged, these siRNAs or miRNAs did not leak or diffuse out from the minicells *in vitro* or *in vivo*.

[0031] The present inventors made an even more surprising discovery that much larger double stranded oligonucleotides that are 40 and 50 nucleotides long (40-mer and 50-mer) can be transported into the intact minicells and these oligonucleotides can also be packaged in therapeutically effective concentrations that do not diffuse out of the minicells. The present disclosure provides a detailed description of these minicells and methods of using the same.

[0032] Throughout this disclosure, various publications, patents and published patent specifications are referenced by an identifying citation. The disclosures of these publications, patents and published patent specifications are hereby incorporated by reference into the present disclosure to more fully describe the state of the art to which this disclosure pertains.

#### 1. Therapeutic Minicell Compositions

[0033] The minicell compositions of the present disclosure are useful in delivering nucleic acid adjuvants to target cells in a mammal, in particular, a human cancer patient.

[0034] In the context of this disclosure, administering the disclosed minicell delivery vectors comprising nucleic acid adjuvants for treating a given disease (i.e., cancer) depends on several factors, in keeping with conventional medical practice. These factors include but are not limited to the patient's age, the status and progression of the disease, and whatever previous or current therapy the patient may have received or is receiving.

##### a. Minicells

[0035] As used herein, "minicell" refers to a derivative of a bacterial cell that is lacking in chromosomes ("chromosome-free") and is engendered by a disturbance in the coordination, during binary fission, of cell division with DNA segregation. Minicells are distinct from other small vesicles, such as so-called "membrane blebs" (~ 0.2µm or less in size), which are

generated and released spontaneously in certain situations but which are not due to specific genetic rearrangements or episomal gene expression. By the same token, intact minicells are distinct from bacterial ghosts, which are not generated due to specific genetic rearrangements or episomal gene expression. Bacterially derived minicells employed in this disclosure are fully intact and, thus, are distinguished from other chromosome-free forms of bacterial cellular derivatives characterized by an outer or defining membrane that is disrupted or degraded, even removed. See U.S. patent No. 7,183,105 at column 111, lines 54 *et seq.* The intact membrane that characterizes the minicells of the present disclosure allows retention of the therapeutic payload (*e.g.*, nucleic acid adjuvants or agonists of nucleic acid sensors) within the minicell until the payload is released, post-uptake, within a target cell.

[0036] Minicells of the invention can be anucleate forms of *E. coli* or other bacterial cells (*e.g.*, *S. typhimurium*). In *E. coli*, for example, mutation of *min* genes, such as *minCD*, can remove the inhibition of septum formation at the cell poles during cell division, resulting in production of a normal daughter cell and an anucleate minicell. See de Boer et al. (1992); Raskin & de Boer (1999); Hu & Lutkenhaus (1999); Harry (2001). For practicing the disclosed methods, in some embodiments it is desirable for minicells to have intact cell walls (“intact minicells”).

[0037] In addition to *min* operon mutations, anucleate minicells may also be generated using a range of other genetic rearrangements or mutations that affect septum formation, for example in the *divIVB1* in *B. subtilis*. See Reeve and Cornett (1975); Levin et al. (1992). Minicells also can be formed following a perturbation in the levels of gene expression of proteins involved in cell division/chromosome segregation. For example, overexpression of *minE* leads to polar division and production of minicells. Similarly, chromosome-less minicells may result from defects in chromosome segregation for example the *smc* mutation in *Bacillus subtilis* (Britton et al. (1998)), *spoOJ* deletion in *B. subtilis* (Ireton et al. (1994)), *mukB* mutation in *E. coli* (Hiraga et al. (1989)), and *parC* mutation in *E. coli* (Stewart and D’Ari (1992)). Gene products may be supplied *in trans*. When over-expressed from a high-copy number plasmid, for example, CafA may enhance the rate of cell division and/or inhibit chromosome partitioning after replication (Okada et al. (1994)), resulting in formation of chained cells and anucleate minicells (Wachi et al. (1989); Okada et al. (1993)).

[0038] Accordingly, minicells can be prepared for the present disclosure from any bacterial cell, be it of Gram-positive or Gram-negative origin. Furthermore, the minicells used in the disclosure may possess intact cell walls (*i.e.*, are “intact minicells”), as noted above, and may be distinguished from other small vesicles, such as membrane blebs, which are not attributable to specific genetic rearrangements or episomal gene expression.

[0039] Thus, in some embodiments, the parental (source) bacteria can comprise one or more bacteria selected from Terra-/Glidobacteria (BV1), Proteobacteria (BV2), and BV4 including Spirochaetes, Sphingobacteria, and Planctobacteria. In some embodiments, the bacteria can comprise one or more selected from Firmicutes (BV3), such as Bacilli, Clostridia or Tenericutes/Mollicutes, or Actinobacteria (BV5) such as Actinomycetales or Bifidobacteriales.

[0040] In some embodiments, the bacteria can comprise one or more selected from Eobacteria (Chloroflexi, Deinococcus-Thermus), Cyanobacteria, Thermodesulfobacteria, thermophiles (Aquificae, Thermotogae), Alpha, Beta, Gamma (Enterobacteriaceae), Delta or Epsilon Proteobacteria, Spirochaetes, Fibrobacteres, Chlorobi/Bacteroidetes, Chlamydiae/Verrucomicrobia, Planctomycetes, Acidobacteria, Chrysiogenetes, Deferribacteres, Fusobacteria, Gemmatimonadetes, Nitrospirae, Synergistetes, Dictyoglomi, Lentisphaerae Bacillales, Bacillaceae, Listeriaceae, Staphylococcaceae, Lactobacillales, Enterococcaceae, Lactobacillaceae, Leuconostocaceae, Streptococcaceae, Clostridiales, Halanaerobiales, Thermoanaerobacterales, Mycoplasmatales, Entomoplasmatales, Anaeroplasmatales, Acholeplasmatales, Haloplasmatales, Actinomycineae, Actinomycetaceae, Corynebacterineae, Mycobacteriaceae, Nocardiaceae, Corynebacteriaceae, Frankineae, Frankiaceae, Micrococcineae, Brevibacteriaceae, and Bifidobacteriaceae.

[0041] For pharmaceutical use, a composition of the disclosure can comprise minicells that are isolated as thoroughly as possible from immunogenic components and other toxic contaminants. Methodology for purifying bacterially derived minicells to remove free endotoxin and parent bacterial cells are described in further detail below, as well as in WO 2004/113507, which is incorporated by reference here in its entirety.

[0042] Another structural element of a minicell derived from Gram-negative bacteria is the O-polysaccharide component of lipopolysaccharide (LPS), which is embedded in the outer

membrane via the lipid A anchor. This component is a chain of repeat carbohydrate-residue units, with as many as 70 to 100 repeat units of four to five sugars per chain. Because these chains are not rigid, in a liquid environment, as *in vivo*, they can adopt a waving, flexible structure.

[0043] As discussed in further detail below, minicells of the present disclosure can comprise at least one nucleic acid adjuvant or a plasmid that encodes a nucleic acid adjuvant for which delivery is desired, as well as at least one agonist of nucleic acid sensors. Nucleic acid adjuvants and agonists of nucleic acid sensors of the present disclosure can bind cognate receptors and produce a desired immune response (*e.g.*, an IFN immune response). Both nucleic acid adjuvants and agonists of nucleic acid sensors of the present disclosure are discussed in more detail below.

[0044] Additionally, in some embodiments, the present disclosure provides methods of delivering at least one nucleic acid adjuvant or agonist of a nucleic acid sensor to a target cell, comprising contacting the target cell with an intact minicell comprising at least one nucleic acid adjuvant molecule, a plasmid comprising a segment that encodes at least one nucleic acid adjuvant molecule, or at least one agonist of a nucleic acid sensor, wherein the target cell engulfs the minicell. Following engulfment of the minicell by the target cell, the nucleic acid adjuvant molecule or agonist of a nucleic acid sensor is released into the cytoplasm of the target cell or expressed by the target cell. Minicells may be brought into contact with the target cells via bispecific ligands, as described in WO 2005/056749. Contact between the minicell and the target cell may be *in vitro* or *in vivo*.

b. Nucleic Acid Adjuvants

[0045] For the purposes of this disclosure, a “nucleic acid adjuvant” is a nucleic acid (*i.e.* polynucleotide or oligonucleotide) that binds to and/or activates a nucleic acid sensor or receptor (*i.e.* a cognate receptor) in a target cell and elicits a desirable immune response. A nucleic acid adjuvant may comprise DNA, RNA, and/or synthetic nucleotides. Further, a nucleic acid adjuvant may be single-stranded (ss) or double stranded (ds) and it may comprise more complex nucleic acid conformations such as hairpin loops, triplexes, and quadruplexes. For example, in

some embodiments, the disclosed nucleic acid adjuvant may be ssDNA, dsDNA, ssRNA, or dsRNA.

[0046] Further, the nucleic acid adjuvant of the disclosure may be varying lengths. In particular, the nucleic acid may be very small, *e.g.*, a dinucleotide. In other words, the nucleic acid adjuvant may be about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, about 50, about 51, about 52, about 53, about 54, about 55, about 56, about 57, about 58, about 59, or about 60 or more nucleotides in length.

[0047] Alternatively, the nucleic acid may have a size greater than about 27 nucleotides in length and up to about 4- about 5KB in a single intact minicell. In some embodiments of the invention, the nucleic acid present in the bacterial minicell can be about 27, about 28, about 29, about 30, about 35, about 40, about 50, about 60, about 75, about 100, about 200, about 300, about 400, about 500, about 600, about 700, about 800, about 900, about 1000, about 1500, about 2000, about 2500, about 3000, about 3500, about 4000, about 4500, or about 5000 nucleotides in length.

[0048] Recent advances have identified a number of nucleic acids that are sensed by receptors in mammalian cells. The binding of these nucleic acids to cognate receptors results in triggering Type I and Type II interferons. These interferons can function as adjuvants in augmenting anti-tumor efficacy (reviewed in Junt and Barchet, 2015).

[0049] For example, nucleic acid sensors of the Toll-like receptor (TLR) family are confined to endosomes (Blasius and Beutler, 2010) and are selectively expressed by a few cell types, mainly those in the innate immune system (*see* Figure 1). TLRs are transmembrane proteins consisting of amino-terminal leucine-rich repeats (LRRs) on the luminal side and a cytosolic Toll/IL-1R (TIR) domain.

[0050] In particular, *TLR3* is activated by double-stranded RNA (dsRNA). *TLR7* detects single-stranded RNA (ssRNA) and short dsRNA. *TLR8* binds short ssRNA and ssRNA breakdown products (Tanji et al. (2015)). *TLR9* instead senses DNA that contains unmethylated CpG motifs that are commonly found in bacterial DNA (Ohto et al. (2015)).

[0051] In TLR7, TLR8 and TLR9, the TIR domain initiates signaling by aggregating the adaptor protein myeloid differentiation primary response protein 88 (MYD88). Activation of the MYD88-dependent signaling cascade ultimately leads to the activation of nuclear factor- $\kappa$ B (NF- $\kappa$ B) and the transcription of genes encoding pro-inflammatory cytokines.

[0052] Exclusively in plasmacytoid dendritic cells (pDCs), the MYD88 signaling complex is able to induce ample transcription of genes encoding various interferon- $\alpha$  (IFN $\alpha$ ) subtypes via a direct activation of IFN-regulatory factor 7 (IRF7).

[0053] By contrast, TLR3 signals via TIR domain-containing adaptor protein inducing IFN $\beta$  (TRIF), which activates NF- $\kappa$ B, mitogen activated protein kinase (MAPK) and IRF3 signaling and results in the transcription of *IFNB* and pro-inflammatory cytokines (Hornung (2014)).

[0054] Thus, in some embodiments, the nucleic acid adjuvants of the disclosure may bind to and activate TLR7, TLR8, and/or TLR9. The nucleic acid adjuvants may bind to and activate these receptors in various cell types, including immune cells and non-immune cells. For instance, in some embodiments, the nucleic acid adjuvants may be targeted to immune cells such as T cells and/or monocytic cells, which can include but are not limited to, monocytes, macrophages, dendritic cells (*e.g.*, pDCs, conventional dendritic cells, or myeloid dendritic cells).

[0055] In some embodiments, the nucleic acid adjuvants of the disclosure may bind to and activate TLR3. In these embodiments, the nucleic acid adjuvants may also be targeted to immune cells or non-immune cells.

[0056] Another set of RNA and DNA sensors that may be activated by the disclosed nucleic acid adjuvants is broadly expressed in the cytosol of immune and non-immune cells (*see* Figure 1). The DExD/H-box helicases retinoic acid-inducible gene I (*RIG-I*) and melanoma differentiation-associated protein 5 (*MDA5*) detect complementary dsRNA structures. Both RIG-I and MDA5 consist of a carboxy-terminal ligand-binding domain, a central DEAxD/H-box helicase domain

and an N-terminal caspase activation and recruitment domain (CARD). These CARDS engage similar CARDS of the signalling adaptor mitochondrial antiviral signalling protein (MAVS) on the outer mitochondrial membrane, which leads to multimerization of MAVS and activation of the MAVS signalling complex (Hou et al. (2011)).

[0057] RIG-I is activated by 5' triphosphorylated or 5' diphosphorylated ends of short dsRNA (Hornung et al. (2006); Goubau et al. (2014)). MDA5 is thought to bind dsRNA and branched high-molecular RNA forms (Kato et al. (2008); Pichlmair et al. (2009)).

[0058] Thus, in some embodiments, the nucleic acid adjuvants of the disclosure may bind to and activate RIG-I and/or MDA5. In these embodiments, the nucleic acid adjuvants may also be targeted to immune cells or non-immune cells.

[0059] Another cytosolic sensor that may be activated by the disclosed nucleic acid adjuvants is cyclic GMP-AMP (*cGAMP*) synthase (cGAS) (Sun et al. (2013)), a receptor for dsDNA. Upon ligand binding, cGAS produces the non-canonically linked cyclic dinucleotide (CDN) [G(2',5')pA(3',5')p] (2'3'-cGAMP), which functions as a second messenger to activate the stimulator of IFN genes (STING) on the endoplasmic reticulum (ER) (Wu et al. (2013); Gao et al. (2013)).

[0060] Thus, in some embodiments, the nucleic acid adjuvants of the disclosure may bind to and activate cGAS. In these embodiments, the nucleic acid adjuvants may also be targeted to immune cells or non-immune cells.

[0061] By contrast, absent in melanoma 2 (AIM2), which is another cytosolic receptor for dsDNA that may be activated by the disclosed nucleic acid adjuvants, triggers the release not of IFNs but of IL-1 $\beta$  and IL-18.

[0062] Thus, in some embodiments, the nucleic acid adjuvants of the disclosure may bind to and activate AIM2. In these embodiments, the nucleic acid adjuvants may also be targeted to immune cells or non-immune cells.

[0063] Recent studies have established the existence of cellular DNA sensors that detect dsDNA within the nucleus to trigger immune signalling. Although the actions of these DNA sensors

have been predominantly studied in the cytosol and endosomes, emerging evidence is shifting focus to the nucleus.

**[0064]** IFI16 (interferon-inducible protein 16) is a DNA sensor (Dawson and Trapani (1995)) that belongs to the highly homologous HIN-200 (hemopoietic expression—interferon-inducibility—nuclear localization) protein family characterized by a 200 amino acid motif containing a DNA binding domain at the C-terminus and a PYRIN domain at the N-terminus, involved mainly in protein-protein interactions. IFI16 is the first viral DNA sensor shown to function within the nucleus and also in the cytoplasm (Dawson and Trapani (1995); Unterholzner et al. (2010)). IFI16 subcellular localization is influenced by the cell type, post-translational modification and cell treatment.

**[0065]** For example, pathogen invasion causes the formation of IFI16 foci in the cytoplasm and induces interferon  $\beta$  (IFN $\beta$ ) gene expression (Unterholzner et al. (2010)) and UV-light causes the transfer of IFI16 from the nucleus to the cytoplasm (Costa et al. (2011)).

**[0066]** The DNA sensing ability of IFI16 is related to the activation of interferon  $\beta$  expression through interaction with stimulator of interferon genes (Unterholzner et al. (2010)), and interferon  $\alpha$  expression (Thompson et al. (2014)). IFI16 binding to DNA is not sequence-specific or AT content-dependent, but is strongly DNA length-dependent (Unterholzner et al. (2010)).

**[0067]** Based on crystallographic studies, the IFI16 HIN-B—double stranded DNA interface is accomplished through electrostatic interactions between the negatively charged sugar-phosphate backbone and positively charged protein residues (Jin et al. (2012)). IFI16 binding to long plasmid DNA was studied and preferences for supercoiled over linear forms and for cruciform structure over double stranded DNA was observed (Brazda et al. (2012)).

**[0068]** IFI16 shows preferential binding to quadruplex DNA with positive effects on quadruplex DNA formation and stabilization (Haronikova et al. (2016)). Across non-immune cell types, IFI16 predominantly localizes to the nucleus (Diner et al. (2015); Li et al. (2012), (2013); Orzalli et al. (2012)).

[0069] Thus, in some embodiments, the nucleic acid adjuvants of the disclosure may bind to and activate IFI16. In these embodiments, the nucleic acid adjuvants may also be targeted to immune cells or non-immune cells.

[0070] In addition to those mentioned above, several other nucleic acid sensors have been described in the art. The skilled artisan will understand that any known nucleic acid sensor may be activated or targeted using the disclosed minicell delivery vectors comprising a nucleic acid adjuvant designed to bind to the cognate receptor (*i.e.* nucleic acid sensor). Accordingly, the nucleic acid sensors that may be targeted or activated by the disclosed compositions are not particularly limited.

[0071] For the purposes of this disclosure, in some embodiments, the minicell delivery vectors may comprise at least one nucleic acid adjuvant. For example, the minicell delivery vector may comprise at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 different nucleic acid adjuvants. In some embodiments, the disclosed minicell delivery vector may comprise both at least one nucleic acid adjuvant and at least one agonist of a nucleic acid sensor.

c. Nucleic Acid-Sensing Pathway Agonists

[0072] For the purposes of this disclosure, an “agonist of a nucleic acid sensor” is a compound—either natural or synthetic—that is capable of agonizing known nucleic acid sensors, including but not limited to those disclosed in the previous section, and can be used to elicit a desired immune response (*i.e.* an IFN response).

[0073] For example, polynucleotide products of the enzyme polynucleotide phosphorylase (PNPase 1) have been studied as synthetic inducers of IFN activity (Field et al. (1967)). Similarly, the dsRNA mimetic polyinosinic:polycytidylic acid (poly(I:C)), was shown to function as an agonist for both TLR3 and MDA5 (Alexopoulou et al. (2001); Gitlin et al. (2006)).

[0074] An analogue of poly(I:C), poly-ICLC, that is formulated with poly-L-lysine to increase RNase resistance, is currently being evaluated as a promising cancer vaccine adjuvant (reviewed in Ammi et al. (2014)).

[0075] Small molecule agonists of nucleic acid sensors have also been developed. As an example, imiquimod, an imidazoquinoline derivative, is an agonist of TLR7 (Hemmi et al. (2002)). It has been shown to be effective in the treatment of dermatological neoplasias, such as basal cell carcinoma and actinic keratosis. Local induction of pro-inflammatory cytokines, local recruitment of immune cells, and improved antigen presentation for the induction of T helper 1 (T<sub>H</sub>1) and CD8<sup>+</sup> T cell responses, all of which result from administration of imiquimod, are considered to be important for treatment success.

[0076] Agonists may also be able to bind to two or more nucleic acid sensors. For example, imidazoquinoline resiquimod is a dual agonist of TLR7 and TLR8. In humans, resiquimod therefore potently activates additional cell types and elicits a broader range of cytokines. A drawback may be that this probably contributes to the more frequent occurrence of systemic adverse effects observed in clinical trials (Huen and Rook (2014)).

[0077] Attempts are being made to develop agonists for human STING and these are being modelled on the physiological agonist 2'3'-cGAMP, which has recently been shown to have vaccine adjuvant properties (Li et al. (2013)).

[0078] Synthetic oligonucleotides can also be designed and used as agonists of nucleic acid sensors. For example, TLR9-stimulatory synthetic CpG oligodeoxynucleotides (CpG-ODNs) were designed based on the immune-stimulatory properties of bacterial DNA that, in contrast to human DNA, is rich in unmethylated CpG motifs (Krieg et al. (1995)). Optimization of sequence features and backbone modifications led to CpG-ODN subtypes that preferentially activate either B cells or pDCs. Multiple clinical trials are under way to determine antitumour activity of CpG-ODNs, in combination with chemotherapy or in therapeutic vaccines.

[0079] The combined use of different agonists may prove more beneficial, as recent data have indicated that co-engagement of multiple nucleic acid receptors elicited the most potent immune responses in preclinical trials of vaccine adjuvants (Goff et al. (2015); Temizoz et al. (2015)).

[0080] Thus, in some embodiments, the minicell delivery vectors may comprise at least one agonist of a nucleic acid sensor. For example, the minicell delivery vector may comprise at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10

different agonists of nucleic acid sensors. In some embodiments, the disclosed minicell delivery vector may comprise both at least one nucleic acid adjuvant and at least one agonist of a nucleic acid sensor.

d. *Nucleic Acids Encoding Nucleic Acid Adjuvants*

**[0081]** In some embodiments of the present disclosure, the disclosed minicell delivery vectors can comprise nucleic acids that encode at least one nucleic acid adjuvant. For example, a plasmid may encode a nucleic acid adjuvant that is expressed inside of a target cell (*i.e.* a mammalian cell or an immune cell). This makes possible endogenous delivery of nucleic acid adjuvants, which, for the treatment of some diseases, may be advantageous over the transient nature of exogenous delivery.

**[0082]** Thus, bacterially-derived intact minicells may carry plasmid DNA encoding one or more nucleic acid adjuvants, including but not limited to, double-stranded or single-stranded RNA or DNA that can bind to and/or activate TLR3, TLR7, TLR8, TLR9, RIG-I, MDA5, AIM2, cGAS, or IFI16. Using minicells that encode multiple nucleic acid adjuvants, it is possible to produce a robust immune response in a patient. For example, a first nucleic acid adjuvant can be expressed from an U6 promoter and a second nucleic acid adjuvant can be expressed from an H1 promoter. These multiple expression cassettes may be carried on a single plasmid, but may also be on different plasmids.

**[0083]** In some embodiments, different nucleic acid adjuvants can be expressed from a single promoter, where the recombinant plasmid carries an expression cassette comprised of multiple nucleic acid adjuvant sequences, which are linked together via non-coding polynucleotide sequences. A single gene transcription terminator can be placed downstream of the complete expression cassette.

**[0084]** In some embodiments, a plasmid may encode the sense and antisense strands of an dsRNA or dsDNA as two independent transcripts that, after expression within a target cell, hybridize to form a functional nucleic acid adjuvant. In some embodiments, a plasmid encodes one or more dsRNA or dsDNA that each are expressed as a single transcript that forms a short hairpin stem-loop structure. The hairpin structure may be processed by other enzymes in the cell to produce a functional nucleic acid adjuvant.

[0085] A nucleic acid encoding a nucleic acid adjuvant molecule to be introduced via the approach described in this section also can have the desired coding segment linked operatively to a regulatory element, such as a promoter, a terminator, an enhancer and/or a signal sequence. A suitable promoter can be tissue-specific or even cell-specific, as the therapeutic context dictates.

[0086] A signal sequence can be used, for the purposes of this disclosure, to effect secretion of an expression product or localization of an expression product to a particular cellular compartment. Thus, a nucleic acid (*i.e.* a plasmid) encoding a therapeutic nucleic acid adjuvant molecule that is delivered via intact minicells may include a signal sequence, in proper reading frame, such that the expression product of interest is secreted by an engulfing cell or its progeny, thereby to influence surrounding cells, in keeping with the chosen treatment paradigm. Illustrative signal sequences include the haemolysin C-terminal secretion sequence, described in U.S. Patent No. 5,143,830, the BAR1 secretion sequence, disclosed in U.S. Patent No. 5,037,743, and the signal sequence portion of the zsig32 polypeptide, described in U.S. Patent No. 6,025,197.

e. Reporter Elements

[0087] A nucleic acid adjuvant molecule or an agonist of a nucleic acid sensor delivered via the disclosed minicell delivery vectors may further comprise a reporter element. A reporter element confers on its target host cell a readily detectable phenotype or characteristic, typically by encoding a polypeptide, not otherwise produced by the host, that can be detected, upon expression, by histological or *in situ* analysis, such as by *in vivo* imaging techniques. For example, a reporter element delivered by an intact minicell, according to the present disclosure, could code for a protein that produces, in the engulfing host cell, a colorimetric or fluorometric change that is detectable by *in situ* analysis and that is a quantitative or semi-quantitative function of transcriptional activation. Illustrative of these proteins are esterases, phosphatases, proteases and other enzymes, the activity of which generates a detectable chromophore or fluorophore.

[0088] Preferred examples of reporter elements include, but are not limited to, *E. coli*  $\beta$ -galactosidase, which effects a color change via cleavage of an indigogenic substrate, indoly1- $\beta$ -D-galactoside, and a luciferase, which oxidizes a long-chain aldehyde (bacterial luciferase) or a

heterocyclic carboxylic acid (luciferin), with the concomitant release of light. Also useful in this context is a reporter element that encodes the green fluorescent protein (GFP) of the jellyfish, *Aequorea victoria*, as described by Prasher et al. (1995). The field of GFP-related technology is illustrated by two published PCT applications, WO 95/21191 (discloses a polynucleotide sequence encoding a 238 amino-acid GFP apoprotein, containing a chromophore formed from amino acids 65 through 67) and WO 95/21191 (discloses a modification of the cDNA for the apopeptide of *A. victoria* GFP, providing a peptide having altered fluorescent properties), and by a report of Heim et al. (1994) of a mutant GFP, characterized by a 4-to-6-fold improvement in excitation amplitude.

[0089] Other genes for use as a reporter element include those that can transform a target cell of the minicell delivery vector to express distinguishing cell-surface antigens, e.g., viral envelope proteins such as HIV gp120 or herpes gD, which are readily detectable by immunoassays.

f. Packaging Nucleic Acid Adjuvants in Minicells

[0090] Nucleic acid adjuvants that can be encoded by a nucleic acid can be introduced into minicells by transforming the encoding nucleic acid into the parental bacterial cell in a vector, such as a plasmid, that encodes the nucleic acid adjuvant. When a minicell is formed from the parental bacterial cell, the minicell retains certain copies of the plasmid and/or the expression product, the nucleic acid adjuvant. More details of packaging an expression product into a minicell is provided in WO 03/033519, the content of which is incorporated into the present disclosure in its entirety by reference.

[0091] Nucleic acid adjuvants also can be packaged into minicells directly. Thus, a nucleic acid adjuvant can be packaged directly into intact minicells by co-incubating a plurality of intact minicells with the nucleic acid adjuvant in a buffer. The buffer composition can be varied, as a function of conditions well known in this field, to optimize the loading of the nucleic acid adjuvant into the intact minicells. The buffer also may be varied in dependence on the nucleotide sequence and the length of the nucleic acid adjuvant to be loaded in the minicells. Once packaged, the nucleic acid remains inside the minicell and is protected from degradation. Prolonged incubation studies with siRNA-packaged minicells incubated in sterile saline showed, for example, no leakage of siRNAs.

[0092] In some embodiments, multiple nucleic acid adjuvants can be packaged in the same minicell. Such an approach can be used to produce a robust immune response in a patient. For example, cancer patients routinely present with tumors capable of evading the innate immune response. To combat this evasion, minicells can be packaged with therapeutically significant concentrations of nucleic acid adjuvants or agonists of nucleic acid sensors to immune cells (*e.g.*, T cells, dendritic cells, or monocytes) to stimulate the immune system to detect and destroy the cancer cells. Furthermore, packaging into the same minicell multiple nucleic acid adjuvants or agonists of nucleic acid sensors can enhance therapeutic success since many cognate receptors belong to distinct signaling pathways and can induce the production and secretion of various cytokines or chemokines including IFN $\alpha$  or IFN $\beta$ . More details of directly packaging a nucleic acid into a minicell is provided in WO 2009/027830, the contents of which are incorporated into the present disclosure in its entirety by reference.

[0093] Small molecule drugs, including agonists of nucleic acid sensors, whether hydrophilic or hydrophobic, can be packaged in minicells by creating a concentration gradient of the drug between an extracellular medium containing minicells and the minicell cytoplasm. When the extracellular medium contains a higher drug concentration than the minicell cytoplasm, the drug naturally moves down this concentration gradient, into the minicell cytoplasm. When the concentration gradient is reversed, however, the drug does not move out of the minicells.

g. Directing Minicells to Target Cells

[0094] For the purposes of this disclosure, the minicells of a composition, as described above, are directed to a target cell via a ligand. Target cells may include mammalian cells, and, in particular, human cells. Target cells may be immune cells or non-immune cells, but in preferred embodiments, the target cells are immune cells, such as T cell, monocytes, dendritic cells, and/or macrophages.

[0095] In some embodiments the ligand is “bispecific.” That is, the ligand displays a specificity for both minicell and target (*e.g.*, immune) cell components, such that it causes a given minicell to bind to the target cell, whereby the latter engulfs the former. Use of bispecific ligands to target a minicell to a tumor cell is further described in WO 05/056749 and WO 05/079854, the respective contents of which are incorporated here in the entirety by reference. Once such a

ligand is attached to a minicell, the unoccupied specificity (“monospecificity”) of the ligand pertains until it interacts with the target cell.

[0096] The ligand can be expressed from within the minicells or their parents and then is displayed on the minicells surface. Alternatively, the ligand can be attached to (“coated on”) the cell membrane of the minicells, *e.g.*, by virtue of ligand-receptor interaction. In either instance the ligand does not require a specificity to the minicell and only displays a specificity to a component that is characteristic of the target cells. That is, such component need not be unique to the target cells, *per se*, or even to the particular kind of immune cell being targeted, so long as the target cells present the component on their cell surface.

[0097] In some embodiments, minicell delivery vectors contained in an administered composition of the disclosure, may contact and bind to a targeted type of cell, eliciting their uptake into the cells, which then are affected by the therapeutic payload. That payload can be at least one nucleic acid adjuvant and/or at least one agonist of a nucleic acid sensor.

[0098] The inventors found that this targeted delivery approach is broadly applicable to a range of mammalian target cells, including cells that normally are refractory to specific adhesion and endocytosis of minicells. Thus, the type of cell being targeted is not particularly limited. Indeed, the binding of a minicell to a target cell precedes rapid endocytosis of the minicells by even non-phagocytic cells. However, in some embodiments, a suitable target cell for the present disclosure is characterized by expression of a cell surface receptor that, upon binding of a ligand, facilitates endocytosis or macropinocytosis.

[0099] The term “endocytosis” encompasses (1) phagocytosis and (2) pinocytosis, itself a category inclusive of (2a) macropinocytosis, all of which tend to access the late-endosome/lysosome pathway. The interaction between the ligand on a minicell and a mammalian cell surface receptor, the present inventors discovered, activates a particular endocytosis pathway, involving receptor mediated endocytosis (rME) to the late-endosomal/lysosomal compartment. By virtue of such an endocytosis pathway, the present inventors further discovered that the minicells were able to release their payload into the cytoplasm of the target mammalian cell. In the event the payload is an encoding nucleic acid, the nucleic acid not only

is not completely degraded in the late-endosomal/lysosomal compartment, but also is expressed in the target mammalian cell.

[0100] Ligands useful in the above-described targeted delivery approach, pursuant to this disclosure, include any agent that binds to a surface component on a target cell and to a surface component on a minicell. In some embodiments, the surface component on a target cell is a receptor. The ligands can comprise a polypeptide and/or carbohydrate component. In some embodiments, antibodies are preferred ligands.

[0101] For example, an antibody that carries specificity for a surface component, such as an immune cell marker, on the target mammalian cells can be used efficiently to target the minicells to the target cells to produce an immune response.

[0102] In some embodiments, preferred ligands comprise antibodies and/or antibody derivatives. In its present use, the term “antibody” encompasses an immunoglobulin molecule obtained by *in vitro* or *in vivo* generation of an immunogenic response. Accordingly, the “antibody” category includes monoclonal antibodies and humanized antibodies, as well as antibody derivatives, such as single-chain antibody fragments (scFv), bispecific antibodies, etc. A large number of different bispecific protein and antibody-based ligands are known, as evidenced by the review article of Caravella and Lugovskoy (2010), incorporated here by reference in its entirety. Antibodies and antibody derivatives useful in the present disclosure also can be obtained by recombinant DNA techniques.

[0103] This disclosure provides a composition comprising bispecific ligand targeted, nucleic acid adjuvant and/or agonist of nucleic acid receptor-packaged minicells that are able to deliver therapeutically significant concentrations of the nucleic acid adjuvant or agonist of a nucleic acid receptor into desired cells of the mammalian host such that a desired, beneficial immune response is elicited in the host.

## 2. **Methods of Treatment**

[0104] Provided herein are methods of treating or preventing various diseases or conditions with the disclosed minicell delivery vectors. In some embodiments, the disease to be treated or prevented comprises a tumor, cancer, malignant disease, or cancer cell proliferation. In some

embodiments, the disease to be treated or prevented is an infection, such as a viral infection. More specifically, the disclosure provides for methods of activating an IFN immune response, and, in particular, the production of interferon type 1. Such methods may comprise administering a therapeutically effective amount of the disclosed minicell delivery vectors containing at least one nucleic acid adjuvant and/or at least one agonist of a nucleic acid sensor.

**[0105]** The disclosed methods may be used to stimulate the immune system to recognize and/or destroy foreign bodies or pathogen cells (*i.e.* viruses, cells infected with viruses, cancer cells, etc.) by administering intact minicells that contain at least one nucleic acid adjuvant, a nucleic acid encoding at least one nucleic acid adjuvant, or an agonist of a nucleic acid sensor.

**[0106]** In some embodiments, administration of the disclosed minicells containing a nucleic acid adjuvant or an agonist of a nucleic acid sensor is not intended to directly kill a foreign body or pathogenic cell. Rather, the administration of such minicells is intended to adjuvate the therapeutic efficacy of another therapeutic agent.

**[0107]** For instance, in some embodiments, a subject may be administered a first dose of the disclosed minicells containing a nucleic acid adjuvant or an agonist of a nucleic acid sensor and a second dose of a minicell containing another therapeutic agent (*i.e.* siRNA or another biologic compound) or drug. In some embodiments, the second dose may comprise a cytotoxic drug, such as a chemotherapeutic, and in some embodiments, the first and second doses may be targeted to different cell types. For instance, a minicell for use in the first dose containing a nucleic acid adjuvant or an agonist of a nucleic acid sensor may be targeted to an immune cell, while a minicell for use in the second dose containing a chemotherapeutic drug may be targeted to a tumor cell. Chemotherapeutics and drugs that are co-administered or administered before or after administration of the disclosed minicells comprising a nucleic acid adjuvant or an agonist of a nucleic acid sensor may be packaged in another minicell or another type of targeted particle, or the chemotherapeutics or drugs may be administered in an unencapsulated, free form.

**[0108]** Additionally, the disclosed minicells comprising a nucleic acid adjuvant or an agonist of a nucleic acid sensor may be administered to a subject that is receiving, has received, or will receive radiation therapy for the treatment of cancer. Indeed, the disclosed minicells comprising a nucleic acid adjuvant or an agonist of a nucleic acid sensor can augment any known

form of cancer treatment by, for example, increasing a subject's innate anti-tumor immune defense.

[0109] Indeed, the disclosed methods allow for the administration of oligonucleotides that are greater than about 27 nucleotides in length and up to about 4- about 5KB in a single intact minicell. In other embodiments, the nucleic acid can be very small, with a size of at least about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, about 31, about 32, about 33, about 34, about 35, about 36, about 37, about 38, about 39, about 40, about 41, about 42, about 43, about 44, about 45, about 46, about 47, about 48, about 49, about 50, about 51, about 52, about 53, about 54, about 55, about 56, about 57, about 58, about 59, or about 60 or more nucleotides in length.

[0110] These oligonucleotides can trigger immunomodulation in the form of production of Type I interferon (IFN $\alpha$  and IFN $\beta$ ). Production of Type I interferon has been shown to produce an adjuvant effect and assist in treating cancer and/or infections. Thus, in some embodiments, administering the disclosed minicells containing a nucleic acid adjuvant or an agonist of a nucleic acid sensor in connection with another therapeutic agent or drug will result in higher antitumor efficacy than administering the other therapeutic agent or drug alone.

[0111] The disclosed methods are a dramatic improvement over the prior art, as it is known to be difficult to administer Type I interferons directly. Interferons have a short half-life in circulation and IV administration of interferons generally results in severe side effects. Thus, the disclosed methods provide a novel means of promoting the production of a therapeutically effective level of interferons in a manner that is physiologically acceptable.

[0112] The disclosed minicells may be targeted directly to specific immune cells, or they may be non-targeted. In some embodiments, non-targeted intact minicells are phagocytosed in macrophages or other monocytic cells, and thereby elicit the desired immunomodulation in these cells.

[0113] Without being bound by theory, it is believed that the disclosed methods work by activating interferon type 1 production, which in turn acts to stimulate immune cells that are capable of killing cancer cells or infected cells. It is known that the immune system is often suppressed in cancer patient, and the disclosed methods may provide a means of “waking up” the cancer patient’s immune system.

[0114] Thus, once the cancer patient’s immune system is active, the success of cytotoxic frontline therapies will be more efficacious. Pairing the disclosed minicells containing a nucleic acid adjuvant or an agonist of a nucleic acid sensor with intact minicells comprising a cytotoxic drug provides superior efficacy over the treatment of the drug alone, as shown in Examples 1 and 2.

[0115] As discussed in more detail below, formulations of the invention can be administered via various routes and to various sites in a mammalian body, to achieve the therapeutic effect(s) desired, either locally or systemically. Delivery may be accomplished, for example, by oral administration, by application of the formulation to a body cavity, by inhalation or insufflation, or by parenteral, intramuscular, intravenous, intraportal, intrahepatic, peritoneal, subcutaneous, intratumoral, or intradermal administration. The mode and site of administration is dependent on the location of the target cells.

a. Purity

[0116] Minicells of the invention are substantially free from contaminating parent bacterial cells. Thus, minicell-containing formulations of the invention preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^7$  minicells, more preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^8$  minicells, even more preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^9$  minicells, still more preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^{10}$  minicells and most preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^{11}$  minicells.

[0117] Methods of purifying minicells are known in the art and described in PCT/IB02/04632, WO 2004/113507, and U.S. patent No. 8,591,862. One such method combines cross-flow filtration (feed flow is parallel to a membrane surface; Forbes, 1987) and dead-end filtration (feed flow is perpendicular to the membrane surface). Optionally, the filtration combination can

be preceded by a differential centrifugation, at low centrifugal force, to remove some portion of the bacterial cells and thereby enrich the supernatant for minicells.

**[0118]** Another purification method employs density gradient centrifugation in a biologically compatible medium. After centrifugation, a minicell band is collected from the gradient, and, optionally, the minicells are subjected to further rounds of density gradient centrifugation to maximize purity. The method may further include a preliminary step of performing differential centrifugation on the minicell-containing sample. When performed at low centrifugal force, differential centrifugation will remove some portion of parent bacterial cells, thereby enriching the supernatant for minicells.

**[0119]** Particularly effective purification methods exploit bacterial filamentation to increase minicell purity. Thus a minicell purification method can include the steps of (a) subjecting a sample containing minicells to a condition that induces parent bacterial cells to adopt a filamentous form, followed by (b) filtering the sample to obtain a purified minicell preparation.

**[0120]** Known minicell purification methods also can be combined. For example, one highly effective combination of methods is as follows:

**[0121]** Step A: Differential centrifugation of a minicell producing bacterial cell culture. This step, which may be performed at 2000g for about 20 minutes, removes most parent bacterial cells, while leaving minicells in the supernatant.

**[0122]** Step B: Density gradient centrifugation using an isotonic and non-toxic density gradient medium. This step separates minicells from many contaminants, including parent bacterial cells, with minimal loss of minicells. Preferably, this step is repeated within a purification method.

**[0123]** Step C: Cross-flow filtration through a 0.45  $\mu\text{m}$  filter to further reduce parent bacterial cell contamination.

**[0124]** Step D: Stress-induced filamentation of residual parent bacterial cells. This may be accomplished by subjecting the minicell suspension to any of several stress-inducing environmental conditions.

[0125] Step E: Antibiotic treatment to kill parent bacterial cells.

[0126] Step F: Cross-flow filtration to remove small contaminants, such as membrane blebs, membrane fragments, bacterial debris, nucleic acids, media components and so forth, and to concentrate the minicells. A 0.2  $\mu\text{m}$  filter may be employed to separate minicells from small contaminants, and a 0.1  $\mu\text{m}$  filter may be employed to concentrate minicells.

[0127] Step G: Dead-end filtration to eliminate filamentous dead bacterial cells. A 0.45  $\mu\text{m}$  filter may be employed for this step.

[0128] Step H: Removal of endotoxin from the minicell preparation. Anti-Lipid A coated magnetic beads may be employed for this step.

[0129] In some embodiments, the purification process achieves removal of (a) smaller vesicles, such as membrane blebs, which are generally smaller than about 0.2  $\mu\text{m}$  in size, (b) free endotoxins released from cell membranes, and (c) parental bacteria, whether live or dead, and their debris, which are sources of free endotoxins, too. Such removal can be implemented with, *inter alia*, a 0.2  $\mu\text{m}$  filter to remove smaller vesicles and cell debris, a 0.45  $\mu\text{m}$  filter to remove parental cells following induction of the parental cells to form filaments, antibiotics to kill live bacterial cells, and antibodies against free endotoxins.

[0130] Underlying the disclosed purification procedure is a discovery by the present inventors that, despite the difference of their bacterial sources, all intact minicells are approximately 400 nm in size, *i.e.*, larger than membrane blebs and other smaller vesicles and yet smaller than parental bacteria. Size determination for minicells can be accomplished by using solid-state, such as electron microscopy, or by liquid-based techniques, *e.g.*, dynamic light scattering. The size value yielded by each such technique can have an error range, and the values can differ somewhat between techniques. Thus, the size of minicells in a dried state can be measured via electron microscopy as approximately 400 nm  $\pm$  50 nm. On the other hand, dynamic light scattering can measure the same minicells to be approximately 500 nm  $\pm$  50 nm in size. Also, drug-packaged, ligand-targeted minicells can be measured, again using dynamic light scattering, to be approximately 500 nm  $\pm$  50 nm.

[0131] This scatter of size values is readily accommodated in practice, *e.g.*, for purposes of isolating minicells from immunogenic components and other toxic contaminants, as described above. That is, an intact, bacterially derived minicell is characterized by cytoplasm surrounded by a rigid membrane, which gives the minicell a rigid, spherical structure. This structure is evident in transmission-electron micrographs, in which minicell diameter is measured, across the minicell, between the outer limits of the rigid membrane. This measurement provides the above-mentioned size value of 400 nm  $\pm$  50 nm.

[0132] Another structural element of a minicell derived from Gram-negative bacteria is the O-polysaccharide component of lipopolysaccharide (LPS), which is embedded in the outer membrane via the lipid A anchor. The component is a chain of repeat carbohydrate-residue units, with as many as 70 to 100 repeat units of four to five sugars per chain. Because these chains are not rigid, in a liquid environment, as *in vivo*, they can adopt a waving, flexible structure that gives the general appearance of seaweed in a coral sea environment; *i.e.*, the chains move with the liquid while remaining anchored to the minicell membrane.

[0133] Influenced by the O-polysaccharide component, dynamic light scattering can provide a value for minicell size of about 500 nm as noted above. Nevertheless, minicells from Gram-negative and Gram-positive bacteria alike readily pass through a 0.45  $\mu$ m filter, which substantiates an effective minicell size of 400 nm  $\pm$  50 nm. The above-mentioned scatter in sizes is encompassed by the present invention and, in particular, is denoted by the qualifier “approximately” in the phrase “approximately 400 nm in size” and the like.

[0134] In relation to toxic contaminants, a composition of the disclosure can contain less than about 350 EU free endotoxin. Illustrative in this regard are levels of free endotoxin of about 250 EU, about 200 EU, about 150 EU, about 100 EU, about 90 EU, about 80 EU, about 70 EU, about 60 EU, about 50 EU, about 40 EU, about 30 EU, about 20 EU, about 15 EU, about 10 EU, about 9 EU, about 8 EU, about 7 EU, about 6 EU, about 5 EU, about 4 EU, about 3 EU, about 2 EU, about 1 EU, about 0.9 EU, about 0.8 EU, about 0.7 EU, about 0.6 EU, about 0.5 EU, about 0.4 EU, about 0.3 EU, about 0.2 EU, about 0.1 EU, about 0.05 EU, and about 0.01 EU, respectively.

[0135] A composition of the disclosure also can contain at least about  $10^8$  minicells, *e.g.*, at least about  $5 \times 10^8$ . Alternatively, the composition can contain on the order of about  $10^9$  or about  $10^{10}$

vesicles, *e.g.*, about  $5 \times 10^9$ , about  $1 \times 10^{10}$  or about  $5 \times 10^{10}$  vesicles. Amongst any such number of minicells, moreover, a composition of the disclosure can contain fewer than about 10 contaminating live/parent bacterial cells, *e.g.*, fewer than about 9, about 8, about 7, about 6, about 5, about 4, about 3, about 2, or about 1 live/parent bacterial cells.

b. Formulations

[0136] Pharmaceutical compositions suitable for use in the methods described herein can include the disclosed minicell delivery vectors and a pharmaceutically acceptable carrier or diluent.

[0137] The invention includes within its scope compositions, or formulations, comprising (a) an intact minicell and (b) a pharmaceutically acceptable carrier therefor, where the minicell contains at least one nucleic acid adjuvant molecule, a plasmid comprising a segment that encodes at least one nucleic acid adjuvant molecule, and/or at least one agonist of a nucleic acid sensor. The at least one nucleic acid adjuvant or agonist of a nucleic acid sensor may comprise any of the nucleic acid adjuvants or agonists of nucleic acid sensors described herein, or a combination thereof.

[0138] In some embodiments, the formulation may also further comprise a drug, as described herein. Preferably, the minicell of the formulation contains the drug. Alternatively, the minicell may contain a nucleic acid molecule, such as a plasmid, that encodes the drug.

[0139] The minicell-containing formulations preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^7$  minicells, more preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^8$  minicells, even more preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^9$  minicells, still more preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^{10}$  minicells and most preferably contain fewer than about 1 contaminating parent bacterial cell per  $10^{11}$  minicells.

[0140] The formulations also optionally contain a bispecific ligand for targeting the minicell to a target cell. The minicell and ligand may be any of those described herein. Thus, in some embodiments the minicell contains at least one nucleic acid adjuvant, a nucleic acid encoding a nucleic acid adjuvant, and/or at least one agonist of a nucleic acid sensor and a bispecific ligand that is capable of binding to a surface component of the minicell and to a surface component of a target mammalian cell.

[0141] A formulation comprising minicells and, optionally drugs and bispecific ligands, of the present disclosure can be formulated in conventional manner, using one or more pharmaceutically acceptable carriers or excipients.

[0142] Formulations may be presented in unit dosage form, *e.g.*, in ampules or vials, or in multi-dose containers, with or without an added preservative. The formulation can be a solution, a suspension, or an emulsion in oily or aqueous vehicles, and may contain formulatory agents, such as suspending, stabilizing and/or dispersing agents. A suitable solution is isotonic with the blood of the recipient and is illustrated by physiological saline, Ringer's solution, and dextrose solution. Alternatively, formulations may be in lyophilized powder form, for reconstitution with a suitable vehicle, *e.g.*, sterile, pyrogen-free water or physiological saline. The formulations also may be in the form of a depot preparation. Such long-acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection.

[0143] In some embodiments, the disclosed minicell delivery vectors may be formulated to be administered concurrently with another therapeutic agent or drug. In some embodiments, the disclosed minicell delivery vectors may be formulated to be administered in sequence with another therapeutic agent or drug. For example, in some embodiments, the disclosed minicell delivery vectors may be administered either before or after the patient has received a treatment of chemotherapy for the treatment of cancer. Alternatively, in some embodiments the disclosed minicell delivery vectors may be administered either before or after the patient has received a vaccine to prevent an infection.

c. Administration Schedules

[0144] In general, the formulations disclosed herein may be used at appropriate dosages defined by routine testing, to obtain optimal physiological effect, while minimizing any potential toxicity. The dosage regimen may be selected in accordance with a variety of factors including age, weight, sex, medical condition of the patient; the severity of the condition to be treated, the route of administration, and the renal and hepatic function of the patient.

[0145] Optimal precision in achieving concentrations of minicell within the range that yields maximum efficacy with minimal side effects may require a regimen based on the kinetics of the minicell availability to target sites and target cells. Distribution, equilibrium, and elimination of

a minicell or drug may be considered when determining the optimal concentration for a treatment regimen. The dosages of the minicells and drugs may be adjusted when used in combination, to achieve desired effects.

**[0146]** Moreover, the dosage administration of the formulations may be optimized using a pharmacokinetic/pharmacodynamic modeling system. For example, one or more dosage regimens may be chosen and a pharmacokinetic/pharmacodynamic model may be used to determine the pharmacokinetic/pharmacodynamic profile of one or more dosage regimens. Next, one of the dosage regimens for administration may be selected which achieves the desired pharmacokinetic/pharmacodynamic response based on the particular pharmacokinetic/pharmacodynamic profile. *See, e.g., WO 00/67776.*

**[0147]** Specifically, the formulations may be administered at least once a week over the course of several weeks. In some embodiment, the formulations are administered at least once a week over several weeks to several months. In some embodiment, the formulations are administered at least twice a week over several weeks to several months. In some embodiment, the formulations are administered at least three times a week over several weeks to several months. In some embodiment, the formulations are administered at least four times a week over several weeks to several months. In some embodiment, the formulations are administered at least five times a week over several weeks to several months. In some embodiment, the formulations are administered at least six times a week over several weeks to several months.

**[0148]** More specifically, the formulations may be administered at least once a day for about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30 or about 31 days. Alternatively, the formulations may be administered about once every day, about once every 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or 31 days or more.

**[0149]** The formulations may alternatively be administered about once every week, about once every 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 weeks or more. Alternatively, the formulations may be administered at least once a week for about 2, about 3,

about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19 or about 20 weeks or more.

[0150] Alternatively, the formulations may be administered about once every month, about once every 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 months or more.

[0151] The formulations may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

[0152] In a method in which minicells are administered before another therapeutic agent or drug, administration of the therapeutic agent or drug may occur anytime from several minutes to several hours before or after administration of the minicells. The other therapeutic agent or drug may alternatively be administered anytime from several hours to several days, possibly several weeks up to several months before or after the minicells.

[0153] More specifically, the minicells may be administered at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23 or about 24 hours before or after another therapeutic agent or drug. Moreover, the minicells may be administered at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, about 23, about 24, about 25, about 26, about 27, about 28, about 29, about 30, or about 31 days before or after the administration of another therapeutic agent or drug. In yet another embodiment, the minicells may be administered at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19 or about 20 weeks or more before or after the administration of another therapeutic agent or drug. In a further embodiment, the minicells may be administered at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11 or about 12 months before or after the administration of another therapeutic agent or drug.

[0154] Dosage regimens can be adjusted to provide the optimum desired response (*e.g.*, a therapeutic response like tumor regression or remission). For example, in some embodiments, a

single bolus of minicell delivery vectors may be administered, while in some embodiments, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the situation. For example, in some embodiments the disclosed minicell delivery vectors may be administered once or twice weekly by subcutaneous, intramuscular, or intravenous injection. In some embodiments, the disclosed minicell delivery vectors may be administered once or twice monthly by subcutaneous, intramuscular, or intravenous injection. In some embodiments, the disclosed minicell delivery vectors may be administered once every week, once every other week, once every three weeks, once every four weeks, once every other month, once every three months, once every four months, once every five months, or once every six months.

[0155] Particular treatment regimens may be evaluated according to whether they will improve a given patient's outcome, meaning the treatment will reduce the risk of recurrence or increase the likelihood of progression-free survival of the given cancer or infection.

[0156] Thus, for the purposes of this disclosure, a subject is treated if one or more beneficial or desired results, including desirable clinical results, are obtained. For example, beneficial or desired clinical results include, but are not limited to, one or more of the following: decreasing one or more symptoms resulting from the disease, increasing the quality of life of those suffering from the disease, decreasing the dose of other medications required to treat the disease, delaying the progression of the disease, and/or prolonging survival of individuals.

[0157] Furthermore, while the subject of the methods is generally a patient with some type of cancer or infection, the age of the patient is not limited. The disclosed methods are useful for treating tumors, cancer, malignant disease, or cancer cell proliferation with various recurrence and prognostic outcomes across all age groups and cohorts. The disclosed methods are also useful for treating or preventing various infections, including both bacterial and viral, either before or throughout the course of the infection. Further, in some embodiments, the subject may be a pediatric subject, while in other embodiments, the subject may be an adult subject.

### 3. **Definitions**

[0158] Unless defined otherwise, technical and scientific terms used herein have the meanings commonly understood by one of ordinary skill in the art, unless otherwise defined. Any suitable

materials and/or methodologies known to those of ordinary skill in the art can be utilized in carrying out the methods described herein.

[0159] For convenience, the meaning of certain terms and phrases employed in the specification, examples, and appended claims are provided below. Other terms and phrases are defined throughout the specification.

[0160] The singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise.

[0161] As used herein, the term “about” will be understood by persons of ordinary skill in the art and will vary to some extent depending upon the context in which it is used. If there are uses of the term which are not clear to persons of ordinary skill in the art given the context in which it is used, “about” will mean up to plus or minus 10% of the particular term.

[0162] As used herein, the term “comprising” is intended to mean that the compositions and methods include the recited elements, but not excluding others. “Consisting essentially of” when used to define compositions and methods, shall mean excluding other elements of any essential significance to the composition or method. “Consisting of” shall mean excluding more than trace elements of other ingredients for claimed compositions and substantial method steps. Embodiments defined by each of these transition terms are within the scope of this disclosure. Accordingly, it is intended that the methods and compositions can include additional steps and components (comprising) or alternatively including steps and compositions of no significance (consisting essentially of) or alternatively, intending only the stated method steps or compositions (consisting of).

[0163] As used herein, “sequence” refers to all or a portion of a polynucleotide or polypeptide sequence.

[0164] “Cancer,” “neoplasm,” “tumor,” “malignancy” and “carcinoma,” used interchangeably herein, refer to cells or tissues that exhibit an aberrant growth phenotype characterized by a significant loss of control of cell proliferation. The methods and compositions of this invention particularly apply to precancerous, malignant, pre-metastatic, metastatic, and non-metastatic cells.

[0165] “Complementary” refers to the topological compatibility or matching together of the interacting surfaces of two molecules, such as base pairing in complementary nucleotide sequences. Thus, molecules can be described as complementary, and furthermore, the contact surface characteristics are complementary to each other.

[0166] “Cytokine” is a generic term for proteins released by one cell population that act on another cell population as intercellular mediators.

[0167] “Drug” refers to any physiologically or pharmacologically active substance that produces a local or systemic effect in animals, particularly mammals and humans.

[0168] “Expression” generally refers to the process by which a polynucleotide sequence undergoes successful transcription and translation such that detectable levels of the amino acid sequence or protein are expressed. In certain contexts herein, expression refers to the production of a nucleic acid sequence (*i.e.* a nucleic acid adjuvant), rather than the expression of a protein or amino acids. In other contexts, expression refers to the production of protein.

[0169] “Host cell” refers to a cell that may be, or has been, used as a recipient for a recombinant vector or other transfer of polynucleotides, and includes the progeny of the original cell that has been transfected. The progeny of a single cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent due to natural, accidental, or deliberate mutation.

[0170] “Hybridization” refers to any process by which a polynucleotide sequence binds to a complementary sequence through base pairing.

[0171] As used herein, the phrases “therapeutically effective amount” means that a dose of the disclosed bead particles provides the specific pharmacological effect for which the drug is administered in a subject in need of such treatment, *e.g.*, to reduce, ameliorate, or eliminate cancer/tumor growth, progression, or recurrence. It is emphasized that a therapeutically effective amount of a minicell containing a nucleic acid adjuvant or agonist of a nucleic acid sensor will not always be effective in treating the cancer/tumors of every individual subject, even though such dosage is deemed to be a therapeutically effective amount by those of skill in the art. Those skilled in the art can adjust what is deemed to be a therapeutically effective amount in

accordance with standard practices as needed to treat a specific subject and/or specific type of disease, *i.e.*, a given cancer or infection. The therapeutically effective amount may vary based on the route of administration and dosage form, the age and weight of the subject, and/or the subject's condition, including the progression, stage, and/or class of the disease at the time of treatment.

**[0172]** The terms "treatment" or "treating" refer to obtaining a desired pharmacological and/or physiologic effect on an active disease. For instance, the effect may be completely or partially reducing, ameliorating or eliminating a disease, for instance, inhibiting cancer/tumor growth and/or progression, or causing cancer/tumor cell death. "Treatment" covers any treatment of a disease in a mammal, particularly a human, and includes: (1) inhibiting a disease symptom, *i.e.*, arresting its development; or (2) relieving the disease symptom, *i.e.*, causing regression of the disease or symptom.

**[0173]** The terms "prevent" or "preventing" as used herein refer to preventing the disease or symptom from occurring in a subject, which may be predisposed to the disease or symptom and/or has not yet been diagnosed as having it. "Preventing" a disease can include stopping the formation of cancer/tumor cells or inhibiting the recurrence of cancer/tumor growth or prophylactically stopping an infection from taking hold in an individual.

**[0174]** "Individual," "subject," "host," and "patient," are used interchangeably herein, and refer to any mammalian subject for whom diagnosis, treatment, or therapy is desired. In one preferred embodiment, the individual, subject, host, or patient is a human. Other subjects may include, but are not limited to, cattle, horses, dogs, cats, guinea pigs, rabbits, rats, primates, and mice.

**[0175]** "Oligonucleotide" refers to a polynucleotide comprising, for example, from about 10 nucleotides (nt) to about 1000 nt. Oligonucleotides for use in the invention are preferably from about 10 nt to about 150 nt. The oligonucleotide may be a naturally occurring oligonucleotide or a synthetic oligonucleotide. Oligonucleotides may be modified.

**[0176]** "Minicell" refers to anucleate forms of bacterial cells, engendered by a disturbance in the coordination, during binary fission, of cell division with DNA segregation. Minicells are distinct from other small vesicles that are generated and released spontaneously in certain situations and,

in contrast to minicells, are not due to specific genetic rearrangements or episomal gene expression. For practicing the present invention, it is desirable for minicells to have intact cell walls (“intact minicells”).

**[0177]** “Modified oligonucleotide” and “Modified polynucleotide” refer to oligonucleotides or polynucleotides with one or more chemical modifications at the molecular level of the natural molecular structures of all or any of the bases, sugar moieties, internucleoside phosphate linkages, as well as to molecules having added substitutions or a combination of modifications at these sites. The internucleoside phosphate linkages may be phosphodiester, phosphotriester, phosphoramidate, siloxane, carbonate, carboxymethylester, acetamidate, carbamate, thioether, bridged phosphoramidate, bridged methylene phosphonate, phosphorothioate, methylphosphonate, phosphorodithioate, bridged phosphorothioate or sulfone internucleotide linkages, or 3'-3', 5'-3', or 5'-5' linkages, and combinations of such similar linkages. The phosphodiester linkage may be replaced with a substitute linkage, such as phosphorothioate, methylamino, methylphosphonate, phosphoramidate, and guanidine, and the ribose subunit of the polynucleotides may also be substituted (*e.g.*, hexose phosphodiester; peptide nucleic acids). The modifications may be internal (single or repeated) or at the end(s) of the oligonucleotide molecule, and may include additions to the molecule of the internucleoside phosphate linkages, such as deoxyribose and phosphate modifications which cleave or crosslink to the opposite chains or to associated enzymes or other proteins. The terms “modified oligonucleotides” and “modified polynucleotides” also include oligonucleotides or polynucleotides comprising modifications to the sugar moieties (*e.g.*, 3'-substituted ribonucleotides or deoxyribonucleotide monomers), any of which are bound together via 5' to 3' linkages.

**[0178]** The phrase “nucleic acid molecules” and the term “polynucleotides” denote polymeric forms of nucleotides of any length, either ribonucleotides or deoxynucleotides. They include single-, double-, or multi-stranded DNA or RNA, genomic DNA, cDNA, DNA-RNA hybrids, or a polymer comprising purine and pyrimidine bases or other natural, chemically or biochemically modified, non-natural, or derivatized nucleotide bases. The backbone of a polynucleotide can comprise sugars and phosphate groups (as may typically be found in RNA or DNA), or modified or substituted sugar or phosphate groups. Alternatively, the backbone of the polynucleotide can comprise a polymer of synthetic subunits such as phosphoramidites and thus can be an

oligodeoxynucleoside phosphoramidate or a mixed phosphoramidate-phosphodiester oligomer. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs, uracyl, other sugars, and linking groups such as fluororibose and thioate, and nucleotide branches. A polynucleotide may be further modified, such as by conjugation with a labeling component. Other types of modifications include caps, substitution of one or more of the naturally occurring nucleotides with an analog, and introduction of means for attaching the polynucleotide to proteins, metal ions, labeling components, other polynucleotides, or a solid support.

**[0179]** “Pharmaceutically acceptable” refers to physiological compatibility. A pharmaceutically acceptable carrier or excipient does not abrogate biological activity of the composition being administered, is chemically inert and is not toxic to the organism in which it is administered.

**[0180]** “Polypeptide” and “protein,” used interchangeably herein, refer to a polymeric form of amino acids of any length, which may include translated, untranslated, chemically modified, biochemically modified, and derivatized amino acids. A polypeptide or protein may be naturally occurring, recombinant, or synthetic, or any combination of these. Moreover, a polypeptide or protein may comprise a fragment of a naturally occurring protein or peptide. A polypeptide or protein may be a single molecule or may be a multi-molecular complex. In addition, such polypeptides or proteins may have modified peptide backbones. The terms include fusion proteins, including fusion proteins with a heterologous amino acid sequence, fusions with heterologous and homologous leader sequences, with or without N-terminal methionine residues, immunologically tagged proteins, and the like.

**[0181]** “Purified” refers to a compound that is removed from its natural environment and is at least about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, about 99.9%, or about 99.99% free from other components with which it is naturally associated.

**[0182]** “Sequence Identity” refers to a degree of similarity or complementarity. There may be partial identity or complete identity. A partially complementary sequence is one that at least partially inhibits an identical sequence from hybridizing to a target polynucleotide; it is referred

to using the functional term “substantially identical.” The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization, and the like) under conditions of low stringency. A substantially identical sequence or probe will compete for and inhibit the binding (*i.e.*, the hybridization) of a completely identical sequence or probe to the target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (*i.e.*, selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (*e.g.*, less than about 30% identity); in the absence of non-specific binding, the probe will not hybridize to the second non-complementary target sequence.

**[0183]** Another way of viewing sequence identity, in the context to two nucleic acid or polypeptide sequences, entails referencing residues in the two sequences that are the same when aligned for maximum correspondence over a specified region. As used here, “percentage of sequence identity” means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

**[0184]** The compositions and methods of the disclosure may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms “comprising”, “including,” “containing”, etc. shall be read expansively and without limitation. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof,

but it is recognized that various modifications are possible within the scope of the disclosure claimed.

### *References*

- [0185] Alexopoulou, L., Holt, A. C., Medzhitov, R. & Flavell, R. A., Recognition of double-stranded RNA and activation of NF- $\kappa$ B by Toll-like receptor 3. *Nature* 413, 732–738 (2001).
- [0186] Ammi, R. et al., Poly(I:C) as cancer vaccine adjuvant: knocking on the door of medical breakthroughs. *Pharmacol. Ther.* 140, 120–131 (2014).
- [0187] Blasius, A. L. & Beutler, B., Intracellular toll-like receptors. *Immunity* 32, 305–315 (2010).
- [0188] Brazda V, Coufal J, Liao JC, Arrowsmith CH., Preferential binding of IFI16 protein to cruciform structure and superhelical DNA. *Biochem Biophys Res Commun.* 422, 716–720 (2012).
- [0189] Costa S, Borgogna C, Mondini M, De Andrea M, Meroni PL, Berti E, et al., Redistribution of the nuclear protein IFI16 into the cytoplasm of ultraviolet B-exposed keratinocytes as a mechanism of autoantigen processing. *Br J Dermatol.* 164, 282–90 (2011).
- [0190] Dawson MJ, Trapani JA., The interferon-inducible autoantigen, IFI 16: localization to the nucleolus and identification of a DNA-binding domain. *Biochem Biophys Res Commun.* 214, 152–62 (1995).
- [0191] Diner, B. A., Li, T., Greco, T. M., Crow, M. S., Fuesler, J. A., Wang, J., and Cristea, I. M., The functional interactome of PYHIN immune regulators reveals IFIX is a sensor of viral DNA. *Mol. Syst. Biol.* 11, 787 (2015).
- [0192] Diner, BA., Lum KK., Cristea, IM., The emerging role of nuclear viral DNA sensors. *J. Biol. Chem.* 290, 26412-26421 (2015).

- [0193] Field, A. K., Tytell, A. A., Lampson, G. P. & Hilleman, M. R., Inducers of interferon and host resistance. II. Multistranded synthetic polynucleotide complexes. *Proc. Natl Acad. Sci. USA* 58, 1004–1010 (1967).
- [0194] Gao, P. et al., Cyclic [G(2',5')pA(3',5')p] is the metazoan second messenger produced by DNA-activated cyclic GMP-AMP synthase. *Cell* 153, 1094–1107 (2013).
- [0195] Gitlin, L. et al., Essential role of mda-5 in type I IFN responses to polyriboinosinic:polyribocytidylic acid and encephalomyocarditis picornavirus. *Proc. Natl Acad. Sci. USA* 103, 8459–8464 (2006).
- [0196] Goff, P. H. et al., Synthetic TLR4 and TLR7 ligands as influenza virus vaccine adjuvants induce rapid, sustained and broadly protective responses. *J. Virol.* 89, 3221–3335 (2015).
- [0197] Goubau, D. et al., Antiviral immunity via RIG-I-mediated recognition of RNA bearing 5'-diphosphates. *Nature* 514, 372–375 (2014).
- [0198] Haronikova L, Coufal J, Kejnovska I, Jagelska EB, Fojta M et al., IFI16 Preferentially Binds to DNA with Quadruplex Structure and Enhances DNA Quadruplex Formation. *PLoS One* 11, 1-19 (2016).
- [0199] Hemmi, H. et al., Small anti-viral compounds activate immune cells via the TLR7 MyD88-dependent signaling pathway. *Nat. Immunol.* 3, 196–200 (2002).
- [0200] Hornung, V. et al., 5'-triphosphate RNA is the ligand for RIG-I. *Science* 314, 994–997 (2006).
- [0201] Hornung, V., SnapShot: nucleic acid immune sensors, part 1. *Immunity* 41, 868–868.e1 (2014).
- [0202] Hou, F. et al., MAVS forms functional prion-like aggregates to activate and propagate antiviral innate immune response. *Cell* 146, 448–461 (2011).
- [0203] Huen, A. O. & Rook, A. H., Toll receptor agonist therapy of skin cancer and cutaneous T-cell lymphoma. *Curr. Opin. Oncol.* 26, 237–244 (2014).

- [0204] Jin T, Perry A, Jiang J, Smith P, Curry JA, Unterholzner L, et al., Structures of the HIN domain:DNA complexes reveal ligand binding and activation mechanisms of the AIM2 inflammasome and IFI16 receptor. *Immunity*. 36, 561–571 (2012).
- [0205] Junt, T., Barchet, W., Translating nucleic acid-sensing pathways into therapies. *Nat Rev Immunol*. 15, 529-544 (2015).
- [0206] Kato, H. et al., Length-dependent recognition of double-stranded ribonucleic acids by retinoic acid-inducible gene-I and melanoma differentiation-associated gene 5. *J. Exp. Med.* 205, 1601–1610 (2008).
- [0207] Krieg, A. M. et al., CpG motifs in bacterial DNA trigger direct B-cell activation. *Nature* 374, 546–549 (1995).
- [0208] Li, X.-D. et al., Pivotal roles of cGAS–cGAMP signaling in antiviral defense and immune adjuvant effects. *Science* 341, 1390–1394 (2013).
- [0209] Li, T., Chen, J., and Cristea, I. M., Human cytomegalovirus tegument protein pUL83 inhibits IFI16-mediated DNA sensing for immune evasion. *Cell Host Microbe* 14, 591–599 (2013).
- [0210] Li, T., Diner, B. A., Chen, J., and Cristea, I. M., Acetylation modulates cellular distribution and DNA sensing ability of interferon-inducible protein IFI16. *Proc. Natl. Acad. Sci. U.S.A.* 109, 10558–10563 (2012).
- [0211] Ohto, U. et al., Structural basis of CpG and inhibitory DNA recognition by Toll-like receptor 9. *Nature* 520, 702–705 (2015).
- [0212] Orzalli, M. H., DeLuca, N. A., and Knipe, D. M., Nuclear IFI16 induction of IRF-3 signaling during herpesviral infection and degradation of IFI16 by the viral ICP0 protein. *Proc. Natl. Acad. Sci. U.S.A.* 109, E3008–E3017 (2012).
- [0213] Pichlmair, A. et al., Activation of MDA5 requires higher-order RNA structures generated during virus infection. *J. Virol.* 83, 10761–10769 (2009).

[0214] Quintieri, L., Geroni, C., Fantin, M., Battaglia, R., Rosato, A., Speed, W., Zanovello, P., Floreani, M., Formation and Antitumor Activity of PNU-159682, A Major Metabolite of Nemorubicin in Human Liver Microsomes. *Clin. Cancer Res.* 11, 1608-1617 (2005).

[0215] Sun, L., Wu, J., Du, F., Chen, X. & Chen, Z. J., Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* 339, 786–791 (2013).

[0216] Tanji, H. et al., Toll-like receptor 8 senses degradation products of single-stranded RNA. *Nat. Struct. Mol. Biol.* 22, 109–115 (2015).

[0217] Temizoz, B. et al., TLR9 and STING agonists synergistically induce innate and adaptive type II IFN. *Eur. J. Immunol.* 45, 1159–1169 (2015).

[0218] Thompson MR, Sharma S, Atianand M, Jensen SB, Carpenter S, Knipe DM, et al., Interferon gamma inducible protein (IFI) 16 transcriptionally regulates type I interferons and other interferon-stimulated genes and controls the interferon response to both DNA and RNA viruses. *J Biol Chem.* 289, 23568–81 (2014).

[0219] Unterholzner L, Keating SE, Baran M, Horan KA, Jensen SB, Sharma S, et al., IFI16 is an innate

[0220] Immune sensor for intracellular DNA. *Nat Immunol.* 11, 997–1004 (2010).

[0221] Wu, J. et al., Cyclic GMP–AMP is an endogenous second messenger in innate immune signaling by cytosolic DNA. *Science* 339, 826–830 (2013).

[0222] The following examples are given to illustrate the present disclosure. It should be understood that the invention is not to be limited to the specific conditions or details described in these examples.

## Examples

### ***Example 1 – Significantly Enhanced Tumor Regression Following Treatment With EGFR minicells<sub>PNU-159682</sub> and minicells<sub>40-mer</sub>***

[0223] This example demonstrates that combined treatment of mouse xenografts with bispecific ligand-targeted and PNU-159682-packaged intact minicells with non-targeted, 40-mer-packaged intact minicells can enhance anti-tumor efficacy as compared to treatment with the former alone.

[0224] Minicells were obtained from an *S. typhimurium minCDE*- mutant strain and were purified using a gradient centrifugation / filamentation / filtration / endotoxin removal procedure as previously described (MacDiarmid et al. (2007)). The purified minicells were packaged with the chemotherapeutic drug doxorubicin-derivative, PNU-159682 (Quintieri et al. (2005)) or double stranded 40 nucleotide long RNA (40-mer) as described (MacDiarmid et al. (2007), (2009)).

[0225] The bispecific antibody (BsAb) used for targeting the minicells was a single polypeptide containing binding specificity for both the *S. typhimurium* O-polysaccharide (present on minicells) and for human EGFR (overexpressed on alveolar adenocarcinoma cells A549). The O-polysaccharide specificity was derived from a mouse monoclonal antibody, for which the variable regions were isolated from the hybridoma cell line 1H10 and presented as a single-chain variable fragment (scFv). The EGFR specificity, also presented as a scFv, was derived from the commercial antibody VECTIBIX<sup>®</sup> (Amgen, USA). The two scFv components were separated by flexible linker, and a 6xHis tag incorporated at the N-terminus to facilitate purification by immobilised metal affinity chromatography, and a c-myc tag at the C-terminus to aid in additional detection. The DNA vector encoding the BsAb contained a hCMV promoter for high level expression and a signal peptide for the secretion of the BsAb into the cell culture medium. The expression vector encoding the BsAb was stably transfected into suspension adapted Chinese hamster ovary (CHO) cells in chemically defined, protein and animal origin free medium and the protein is expressed over 10 days in culture.

[0226] The two chromatographic columns were used to purify the antibody and are an immobilized metal ion affinity chromatography (IMAC- HisTrap Excel column) and hydroxyapatite chromatography (BioRad CHT I column), which provided final antibody purity

of >98%. For viral safety of product, the antibody went through a solvent/detergent inactivation (using TNBP/Tween) and a viral filtration. The final yield of antibody was 10 mg from 1 L of cell culture supernatant.

[0227] The mice (6 week old female athymic nude mice) used in this example were purchased from Animal Resources Centre, Perth, WA, Australia, and all animal experiments were performed in compliance with the guide of care and use of laboratory animals and with Animal Ethics Committee approval. The experiments were performed in the NSW Agriculture accredited small animal facility at EnGeneIC Pty Ltd (Sydney, NSW, Australia). Human alveolar adenocarcinoma cells (A549, ATCC) were grown in tissue culture to full confluency in T-75 flasks in RPMI 1640 medium supplemented with 5% Bovine Calf Serum (GIBCO-BRL Life Technologies, Invitrogen Corporation, Carlsbad, CA, USA) and glutamine (Invitrogen) in a humidified atmosphere of 95% air and 5% CO<sub>2</sub> at 37°C.

[0228]  $1 \times 10^6$  cells in 50  $\mu$ L serum-free media together with 50  $\mu$ L growth factor reduced matrigel (BD Biosciences, Franklin Lakes, NJ, USA) were injected subcutaneously between the shoulder blades of each mouse using a 23-gauge needle. The tumors were measured twice a week using an electronic digital caliper (Mitutoyo, Japan, precision to 0.001) and mean tumor volume was calculated using the formula, length (mm)  $\times$  width<sup>2</sup> (mm)  $\times$  0.5 = volume (mm<sup>3</sup>). The treatments were commenced when the tumors reached a mean of  $\sim 285$  mm<sup>3</sup> and mice were randomized to four different groups of 7 mice per group. All treatments were administered intravenously (i.v.) in a total volume of 100  $\mu$ l. All minicell doses contained  $1 \times 10^9$  minicells of the respective type.

[0229] The experiment was designed as follows. Group 1 (control) received no sterile physiological saline. Group 2 <sup>EGFR</sup>minicells<sub>PNU-159682</sub>, Group 3 <sup>EGFR</sup>minicells<sub>PNU-159682</sub> + minicells<sub>40-mer</sub>.

[0230] The results revealed (Figure 3) that mice treated with <sup>EGFR</sup>minicells<sub>PNU-159682</sub> (Group 2) achieved tumor stabilization. In contrast, mice treated with <sup>EGFR</sup>minicells<sub>PNU-159682</sub> + minicells<sub>40-mer</sub> (Group 3) showed *highly significant tumor regression* by day 44 after a total of 6 doses.

[0231] These results demonstrate the effectiveness of the nucleic acid adjuvants according to the invention, as the therapeutic results were *significantly* better with use of the adjuvant as compared to treatment with non-adjuvanted drug.

***Example 2 – Significantly Enhanced Tumor Regression Following Treatment With  $^{EGFR}$  minicells<sub>PNU-159682</sub> and minicells<sub>40-mer</sub> or minicells<sub>50-mer</sub>***

[0232] This example demonstrates that combined treatment of mouse xenografts with bispecific ligand-targeted and PNU-159682-packaged intact minicells with non-targeted, 40-mer- or 50-mer-packaged intact minicells can enhance anti-tumor efficacy as compared to treatment with the former alone.

[0233] The various minicells were prepared as described in Example 1. Seven mice per group were prepared for the A549 xenograft as described in Example 1.

[0234] The experiment was designed as follows. Group 1 (control) received no sterile physiological saline. Group 2  $^{EGFR}$ minicells<sub>PNU-159682</sub>, Group 3  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells<sub>40-mer</sub>, Group 4  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells<sub>50-mer</sub>, Group 5  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells.

[0235] The results revealed (Figure 4) that mice treated with  $^{EGFR}$ minicells<sub>PNU-159682</sub> (Group 2) and  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells (Group 5) achieved tumor stabilization. In contrast, mice treated with  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells<sub>40-mer</sub> (Group 3) or with  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells<sub>50-mer</sub> (Group 4) showed *significant enhancement of tumor regression* by day 38 after a total of 6 doses. Of further interest, when the saline treated mice had developed large tumor volumes ( $\sim 600\text{mm}^3$ ) by day 36 and the treatment was changed to  $^{EGFR}$ minicells<sub>PNU-159682</sub> + minicells<sub>40-mer</sub>, the tumor volumes plummeted to  $\sim 300\text{mm}^3$  by day 38. This is a dramatic 50% decrease in tumor volume over a 2 day period.

[0236] These results demonstrate the effectiveness of the nucleic acid adjuvants according to the invention, as the therapeutic results were *significantly* better with use of the adjuvant as compared to treatment with non-adjuvanted drug.

***Example 3 – Prophetic Human Treatment***

[0237] This example illustrates methods using the disclosed minicell delivery vectors in the treatment of cancer.

[0238] A patient known to have or suspected of having cancer is administered a therapeutically effective amount of a bacterially-derived minicell delivery vector comprising at least one nucleic acid adjuvant or agonist of a nucleic acid sensor, by intravenous, intramuscular, or subcutaneous injection. The patient is evaluated for the presence and/or severity of signs and symptoms associated with cancer, including, but not limited to, pain, weakness, tumor size, etc., and the patient is treated until one or more signs/symptoms are reduced, ameliorated, or eliminated.

[0239] Optionally, samples may be taken from the patient to monitor cancer progression following treatment. Optionally, another dose of the bacterially-derived minicell delivery vector comprising at least one nucleic acid adjuvant or agonist of a nucleic acid sensor is administered if signs/symptoms persist and/or if the cancer progresses or recurs.

\* \* \* \* \*

[0240] One skilled in the art readily appreciates that the present disclosure is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. Modifications therein and other uses will occur to those skilled in the art. These modifications are encompassed within the spirit of the disclosure and are defined by the scope of the claims, which set forth non-limiting embodiments of the disclosure. All public documents referenced herein, including but not limited to published patent documents, are specifically incorporated by reference.

**What is claimed:**

1. A composition comprising:
  - (a) an intact bacterial minicell comprising:
    - (i) at least one nucleic acid adjuvant molecule,
    - (ii) at least one a plasmid comprising a segment that encodes at least one nucleic acid adjuvant molecule, or
    - (iii) at least one an agonist of a nucleic acid sensor, and
  - (b) a pharmaceutically acceptable carrier therefore,  
wherein the nucleic acid adjuvant molecule or agonist of a nucleic acid sensor produces an immune response from a target cell.
2. The composition of claim 1, wherein:
  - (a) the immune response from the target cell comprises the production of Type I interferon; and/or
  - (b) the immune response from the target cell comprises the production of Type I interferon, wherein Type I interferon comprises interferon- $\alpha$  or interferon- $\beta$ .
3. The composition of claim 1 or 2, wherein the at least one nucleic acid adjuvant comprises a nucleic acid that binds to at least one of TLR3, TLR7, TLR8, TLR9, RIG-I, MDA5, AIM2, cGAS, or IFI16.
4. The composition of any one of claims 1-3, wherein:
  - (a) the intact minicell comprises at least two nucleic acid adjuvant molecules; and/or
  - (b) the intact minicell comprises a nucleic acid adjuvant molecule and an agonist of a nucleic acid sensor.
5. The composition of any one of claims 1-4, wherein:
  - (a) the at least one nucleic acid adjuvant molecule comprises a sequence of at least 40 nucleotides;
  - (b) the at least one nucleic acid adjuvant molecule is a 40-mer; or
  - (c) the at least one nucleic acid adjuvant molecule is a 50-mer.

6. The composition of any one of claims 1-5, wherein the at least one agonist of a nucleic acid sensor comprises a polynucleotide product of PNPase1, poly(I:C), poly-ICLC, imiquimod, imidazoquinoline resiquimod, or CpG-ODNs.
7. The composition of any one of claims 1-6, further comprising a bispecific ligand.
8. The composition of claim 7, wherein:
- (a) the bispecific ligand comprises a first arm that carries specificity for a minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor; and/or
  - (b) the bispecific ligand comprises a first arm that carries specificity for a minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor, wherein the minicell surface structure is an O-polysaccharide component of a lipopolysaccharide on the minicell surface; and/or
  - (c) the bispecific ligand comprises a first arm that carries specificity for a minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor, wherein the mammalian cell surface receptor is capable of activating receptor-mediated endocytosis of the minicell; and/or
  - (d) the bispecific ligand comprises an antibody or antibody fragment.
9. The composition of any one of claims 1-8, wherein:
- (a) the composition comprises fewer than about 1 contaminating parent bacterial cell per  $10^7$  minicells; or
  - (b) the composition comprises fewer than about 1 contaminating parent bacterial cell per  $10^8$  minicells; or
  - (c) the composition comprises fewer than about 1 contaminating parent bacterial cell per  $10^9$  minicells; or
  - (d) the composition comprises fewer than about 1 contaminating parent bacterial cell per  $10^{10}$  minicells; or
  - (e) the composition comprises fewer than about 1 contaminating parent bacterial cell per  $10^{11}$  minicells.

10. The composition of any one of claims 1-9, wherein:
- (a) the plasmid comprises a regulatory element operably linked to the segment that encodes at least one nucleic acid adjuvant molecule; and/or
  - (b) the plasmid encodes more than one nucleic acid adjuvant molecule.
11. A method of delivering a nucleic acid adjuvant to a target cell, comprising contacting a target cell with an intact minicell comprising:
- (a) at least one nucleic acid adjuvant molecule,
  - (b) a plasmid comprised of a segment that encodes at least one nucleic acid adjuvant molecule, or
  - (c) at least one agonist of a nucleic acid receptor,
- wherein the target cell engulfs the minicell.
12. The method of claim 11, wherein the target cell is a:
- (a) mammalian cell; or
  - (b) human cell; or
  - (c) human immune cell; or
  - (d) human immune cell which is a monocytic cell, a macrophage, a T cell, or a dendritic cell.
13. The method of claim 11 or 12, wherein:
- (a) the at least one nucleic acid adjuvant binds a receptor on the target cell; and/or
  - (b) the at least one nucleic acid adjuvant comprises a nucleic acid that binds to at least one of TLR3, TLR7, TLR8, TLR9, RIG-I, MDA5, AIM2, cGAS, or IFI16.
14. The method of any one of claims 11-13, wherein:
- (a) the intact minicell comprises at least two nucleic acid adjuvant molecules; and/or
  - (b) the intact minicell comprises a nucleic acid adjuvant molecule and an agonist of a nucleic acid sensor.
15. The method of any one of claims 11-14, wherein:
- (a) the at least one nucleic acid adjuvant molecule comprises a sequence of at least 40 nucleotides;

- (b) the at least one nucleic acid adjuvant molecule is a 40-mer; or
- (c) the at least one nucleic acid adjuvant molecule is a 50-mer.

16. The method of any one of claims 11-15, wherein the at least one agonist of a nucleic acid sensor comprises a polynucleotide product of PNPase1, poly(I:C), poly-ICLC, imiquimod, imidazoquinoline resiquimod, or CpG-ODNs.

17. The method of any one of claims 11-16, wherein:

- (a) contact between the minicell and the mammalian cell occurs *in vitro*; or
- (b) contact between the minicell and the mammalian cell occurs *in vivo*.

18. The method of any one of claims 11-17, wherein:

- (a) the plasmid comprises a regulatory element operably linked to the segment that encodes at least one nucleic acid adjuvant molecule; and/or
- (b) the plasmid encodes multiple nucleic acid adjuvant molecules.

19. The method of any one of claims 11-18, further comprising a bispecific ligand.

20. The method of claim 19, wherein:

(a) the bispecific ligand comprises a first arm that carries specificity for a minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor; and/or

(b) the bispecific ligand comprises a first arm that carries specificity for a minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor, wherein the minicell surface structure is an O-polysaccharide component of a lipopolysaccharide on the minicell surface; and/or

(c) the bispecific ligand comprises a first arm that carries specificity for a minicell surface structure and a second arm that carries specificity for a non-phagocytic mammalian cell surface receptor, wherein the mammalian cell surface receptor is capable of activating receptor-mediated endocytosis of the minicell; and/or

- (d) the bispecific ligand comprises an antibody or antibody fragment.

21. The method of any one of claims 11-20, wherein the target cell is phagocytosis- or endocytosis-competent.

22. A method of treating a disease in a subject, comprising administering to a subject with a disease an intact minicell composition according to any one of claims 1-10, wherein the minicell is engulfed by a target cell following administration.

23. Use of a composition according to any one of claims 1-10 for treating a disease, wherein the minicell is engulfed by a target cell following administration.

24. The method of claim 22 or the use of claim 23, wherein:

- (a) the disease is cancer; or
- (b) the disease is an infection.

25. The method of claim 22 or 24, or the use of claim 23 or 24, further comprising:

- (a) administering a drug for the treatment of the disease to the subject; and/or
- (b) administering a drug for the treatment of the disease to the subject, wherein the drug is a chemotherapeutic; and/or
- (c) administering a drug for the treatment of the disease to the subject, wherein the drug is an anti-viral; and/or
- (d) administering a drug for the treatment of the disease to the subject, wherein the drug is packaged in an intact minicell.

Figure 1




Figure 2




Figure 3





Figure 4



**A. CLASSIFICATION OF SUBJECT MATTER****A61K 48/00(2006.01)i, A61K 47/46(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**Minimum documentation searched (classification system followed by classification symbols)  
A61K 48/00; A61K 39/44; A61K 31/7088; C12N 15/00; A61K 38/21; C12N 1/20; A61K 47/46Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
Korean utility models and applications for utility models  
Japanese utility models and applications for utility modelsElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
eKOMPASS(KIPO internal) & Keywords: minicell, nucleic acid adjuvant, agonist, nucleic acid sensor, immune response, Type I interferon**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2007-0298056 A1 (BRAHMBHATT, HIMANSHU et al.) 27 December 2007 See abstract; paragraph [0134]; claims 1, 30, 37.	1-3,11-13
Y	GLAS, MARTIN et al., 'Targeting the cytosolic innate immune receptors RIG-I and MDA5 effectively counteracts cancer cell heterogeneity in glioblastoma' Stem Cells, 2013, Vol.31, pages 1064-1074 See abstract; pages 1070, 1072; Figure 1C.	1-3,11-13
Y	US 2015-0098897 A1 (ENGINEIC MOLECULAR DELIVERY PTY LTD.) 09 April 2015 See abstract; claims 1-16.	1-3,11-13
Y	US 2012-0142079 A1 (SABBADINI, ROGER A. et al.) 07 June 2012 See abstract; claims 1-13.	1-3,11-13
Y	GIACALONE, MATTHEW J. et al., 'The use of bacterial minicells to transfer plasmid DNA to eukaryotic cells' Cellular Microbiology, 2006, Vol.8, No.10, pages 1624-1633 See abstract; pages 1629-1630.	1-3,11-13

 Further documents are listed in the continuation of Box C. See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

28 December 2017 (28.12.2017)

Date of mailing of the international search report

**28 December 2017 (28.12.2017)**

Name and mailing address of the ISA/KR


International Application Division  
Korean Intellectual Property Office  
189 Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea

Facsimile No. +82-42-481-8578

Authorized officer

KAM, YOO LIM

Telephone No. +82-42-481-3516



**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: 22,24-25  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claims 22 and 24-25 pertain to methods for treatment of the human body by therapy, and thus relate to a subject matter which this International Searching Authority is not required, under PCT Article 17(2)(a)(i) and PCT Rule 39.1(iv), to search.
2.  Claims Nos.: 8,20  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
Claims 8 and 20 refer to one of unsearchable claims which do not comply with PCT Rule 6.4(a).
3.  Claims Nos.: 4-7,9-10,14-19,21-25  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of any additional fees.
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

**Remark on Protest**

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

**PCT/IB2017/056131**

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2007-0298056 A1	27/12/2007	AU 2005-276145 A1	02/03/2006
		AU 2005-276145 B2	27/05/2010
		AU 2005-276145 C1	06/01/2011
		AU 2008-291833 A1	05/03/2009
		AU 2008-291833 B2	08/11/2012
		AU 2010-212520 A1	16/09/2010
		AU 2010-212520 B2	31/01/2013
		AU 2013-203202 A1	02/05/2013
		AU 2013-203202 B2	16/02/2017
		AU 2017-203234 A1	08/06/2017
		CA 2577938 A1	02/03/2006
		CA 2577938 C	16/04/2013
		CA 2682704 A1	05/03/2009
		CA 2682704 C	08/07/2014
		CA 2803995 A1	02/03/2006
		CA 2803995 C	10/06/2014
		CA 2844647 A1	05/03/2009
		CA 2844647 C	26/07/2016
		CA 2933978 A1	05/03/2009
		CN 101072876 A	14/11/2007
		CN 101072876 B	28/11/2012
		CN 101715489 A	26/05/2010
		CN 101715489 B	25/05/2016
		CN 102921019 A	13/02/2013
		CN 102921019 B	01/04/2015
		CN 105854029 A	17/08/2016
		EP 1791959 A2	06/06/2007
		EP 2145002 A2	20/01/2010
		EP 2145002 B1	07/01/2015
		EP 2386640 A2	16/11/2011
		EP 2386640 A3	27/06/2012
		EP 2386640 A9	28/12/2011
		EP 2386640 B1	28/01/2015
		EP 2532746 A2	12/12/2012
		EP 2532746 A3	14/08/2013
		EP 2865755 A2	29/04/2015
		EP 2865755 A3	06/05/2015
		EP 2865755 B1	01/02/2017
		EP 3205724 A1	16/08/2017
		JP 2008-510794 A	10/04/2008
		JP 2010-523487 A	15/07/2010
		JP 4965447 B2	04/07/2012
		JP 5167338 B2	21/03/2013
		US 2008-0299084 A1	04/12/2008
		US 2011-0111041 A1	12/05/2011
		US 2013-0261170 A1	03/10/2013
		US 2014-0221459 A1	07/08/2014
		US 2014-0315976 A1	23/10/2014
		US 2016-0051687 A1	25/02/2016

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/IB2017/056131

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		US 2016-0113883 A1	28/04/2016
		US 2017-258734 A1	14/09/2017
		US 8669101 B2	11/03/2014
		US 8691963 B2	08/04/2014
		US 8735566 B2	27/05/2014
		US 8956864 B2	17/02/2015
		US 9066982 B2	30/06/2015
		US 9242007 B2	26/01/2016
		US 9730897 B2	15/08/2017
		WO 2006-021894 A2	02/03/2006
		WO 2006-021894 A3	04/05/2006
		WO 2009-027830 A2	05/03/2009
		WO 2009-027830 A3	14/05/2009
US 2015-0098897 A1	09/04/2015	AU 2014-330895 A1	14/04/2016
		CA 2926161 A1	09/04/2015
		CN 105658233 A	08/06/2016
		EP 3052122 A1	10/08/2016
		JP 2016-532639 A	20/10/2016
		KR 10-2016-0058885 A	25/05/2016
		US 2017-0326235 A1	16/11/2017
		US 9731011 B2	15/08/2017
		WO 2015-049589 A1	09/04/2015
US 2012-0142079 A1	07/06/2012	AU 2002-318168 A1	09/09/2003
		AU 2002-318168 B2	13/12/2007
		AU 2008-201082 B2	15/09/2011
		AU 2008-201082 C1	26/06/2014
		CA 2517027 A1	04/09/2003
		EP 1487965 A2	22/12/2004
		EP 2272946 A2	12/01/2011
		EP 2272946 A3	27/07/2011
		EP 2272946 B1	08/04/2015
		EP 2272946 B9	24/06/2015
		EP 2390308 A1	30/11/2011
		US 2003-0166099 A1	04/09/2003
		US 2003-0166279 A1	04/09/2003
		US 2003-0190601 A1	09/10/2003
		US 2003-0190683 A1	09/10/2003
		US 2003-0190749 A1	09/10/2003
		US 2003-0194714 A1	16/10/2003
		US 2003-0194798 A1	16/10/2003
		US 2003-0198995 A1	23/10/2003
		US 2003-0198996 A1	23/10/2003
		US 2003-0199005 A1	23/10/2003
		US 2003-0199088 A1	23/10/2003
		US 2003-0199089 A1	23/10/2003
		US 2003-0202937 A1	30/10/2003
		US 2003-0203411 A1	30/10/2003
		US 2003-0203481 A1	30/10/2003

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No.

**PCT/IB2017/056131**

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		US 2003-0207833 A1	06/11/2003
		US 2003-0211086 A1	13/11/2003
		US 2003-0211599 A1	13/11/2003
		US 2003-0219408 A1	27/11/2003
		US 2003-0219888 A1	27/11/2003
		US 2003-0224369 A1	04/12/2003
		US 2003-0224444 A1	04/12/2003
		US 2003-0232335 A1	18/12/2003
		US 2005-0147590 A1	07/07/2005
		US 2007-0122881 A1	31/05/2007
		US 2009-0011490 A1	08/01/2009
		US 2011-0281330 A1	17/11/2011
		US 2012-0135503 A1	31/05/2012
		US 2012-0252099 A1	04/10/2012
		US 2013-0337545 A1	19/12/2013
		US 2015-0218254 A1	06/08/2015
		US 6930182 B1	16/08/2005
		US 7183105 B2	27/02/2007
		US 7396822 B2	08/07/2008
		US 7871815 B2	18/01/2011
		US 8101396 B2	24/01/2012
		US 8129166 B2	06/03/2012
		US 8524484 B2	03/09/2013
		US 9017986 B2	28/04/2015
		US 9670270 B2	06/06/2017
		WO 03-072014 A2	04/09/2003