

Description**BACKGROUND OF THE INVENTION**

[0001] The present invention relates generally to methods for manufacturing downstream products of 4-hydroxybutyrate or 1,4-butanediol.

[0002] 1,4-Butanediol (BDO) is a four carbon dialcohol that currently is manufactured exclusively through various petrochemical routes. BDO is part of a large volume family of solvents and polymer intermediates that includes gamma-butyrolactone (GBL), tetrahydrofuran (THF), pyrrolidone, N-methylpyrrolidone (NMP), and N-vinyl-pyrrolidone. The overall market opportunity for this family exceeds \$4.0 B.

[0003] Approximately 2.5B lb BDO is produced globally per year with 4-5% annual growth and a recent selling price ranging from \$1.00-1.20/lb. The demand for BDO stems largely from its use as an intermediate for polybutylene terephthalate (PBT) plastic resins, polyurethane thermoplastics and co-polyester ethers. BDO also serves as a primary precursor to THF, which is employed as an intermediate for poly(tetramethylene glycol) PTMEG copolymers required for lycra and spandex production. Approximately 0.7 B lb of THF is produced globally per year with an annual growth rate over 6%. A significant percentage of growth (>30%) for both BDO and THF is occurring in Asia (China and India). GBL currently is a smaller volume (0.4 B lb/year) product which has numerous applications as a solvent, as an additive for inks, paints, and dyes, as well as the primary precursor to pyrrolidone derivatives such as NMP.

[0004] Conventional processes for the synthesis of BDO use petrochemical feedstocks for their starting materials. For example, acetylene is reacted with 2 molecules of formaldehyde in the Reppe synthesis reaction (Kroschwitz and Grant, Encyclopedia of Chem. Tech., John Wiley and Sons, Inc., New York (1999)), followed by catalytic hydrogenation to form 1,4-butanediol. It has been estimated that 90% of the acetylene produced in the U.S. is consumed for butanediol production. Alternatively, it can be formed by esterification and catalytic hydrogenation of maleic anhydride, which is derived from butane. Downstream, butanediol can be further transformed; for example, by oxidation to γ -butyrolactone, which can be further converted to pyrrolidone and N-methyl-pyrrolidone, or hydrogenolysis to tetrahydrofuran (Figure 1). These compounds have varied uses as polymer intermediates, solvents, and additives, and have a combined market of nearly 2 billion lb/year.

[0005] The conventional hydrocarbon feedstock-based approach utilizes methane to produce formaldehyde. Thus, a large percentage of the commercial production of BDO relies on methane as a starting material. The production of acetylene also relies on petroleum-based starting material (see Figure 1). Therefore, the costs of BDO production fluctuate with the price of petroleum and natural gas.

[0006] JPS62285779 (A) relates to a bacterial strain belonging to the *Bacillus* genus (C-105 strain (FERM P-8788)) and production of 1,4-butanediol using the same.

[0007] It is desirable to develop a method for production of these chemicals by alternative means that not only substitute renewable for petroleum-based feedstocks, and also use less energy- and capital-intensive processes. The Department of Energy has proposed 1,4-diacids, and particularly succinic acid, as key biologically-produced intermediates for the manufacture of the butanediol family of products (DOE Report, "Top Value-Added Chemicals from Biomass", 2004). However, succinic acid is costly to isolate and purify and requires high temperatures and pressures for catalytic reduction to butanediol.

[0008] Thus, there exists a need for alternative means for effectively producing commercial quantities of 1,4-butanediol and its chemical precursors. The present invention satisfies this need and provides related advantages as well.

SUMMARY OF INVENTION

[0009] The invention is defined by the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

Figure 1 is a schematic diagram showing an entry point of 4-hydroxybutanoic acid (4-HB) into the product pipeline of the 1,4-butanediol (BDO) family of chemicals, and comparison with chemical synthesis routes from petrochemical feedstocks. Solid black arrows show chemical synthesis routes; dashed blue arrows show a biosynthetic route to 4-HB and subsequent conversion steps to BDO family chemicals.

Figure 2 is a schematic diagram showing biochemical pathways to 4-hydroxybutyrate (4-HB) and γ -butyrolactone (GBL) production. Enzymes catalyzing the 4-HB biosynthetic reactions are: (1) CoA-independent succinic semial-

selected from the set of genes comprising *adhE*, *ldhA*, *pflAB*, *mdh*, and *aspA*.

43. The method of item 34, wherein said set of metabolic modifications comprise disruption of the genes *adhE*, *ldhA*, *pflAB*, *mdh*, and *aspA*.

44. The method of item 34, wherein said non-naturally occurring microorganism further comprises a 1,4-butanediol (BDO) biosynthetic pathway, said pathway comprising at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or alcohol dehydrogenase, wherein said exogenous nucleic acid is expressed in sufficient amounts to produce 1,4-butanediol (BDO).

Claims

1. A method for manufacturing a downstream product of 4-hydroxybutyrate or 1,4-butanediol, comprising:

- (a) culturing under exponential growth phase in a sufficient amount of nutrients and media a non-naturally occurring microorganism comprising disruption of the genes *adhE*, *ldhA*, *pflAB* and *mdh*, wherein said microorganism expresses at least one exogenous nucleic acid encoding an enzyme in a 4-hydroxybutyrate or 1,4-butanediol biosynthetic pathway in sufficient amounts to produce 4-hydroxybutyrate or 1,4-butanediol, said at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or alcohol dehydrogenase, and wherein said microorganism produces 4-hydroxybutyrate or 1,4-butanediol;
- (b) optionally isolating 4-hydroxybutyrate or 1,4-butanediol; and
- (c) chemically converting 4-hydroxybutyrate or 1,4-butanediol to the downstream product.

2. The method of claim 1, wherein the microorganism exhibits stable growth-coupled production of 1,4-butanediol.

3. The method of claim 1 or 2, wherein the downstream product is gamma-butyrolactone.

4. The method of claim 3, further comprising the step of isolating gamma-butyrolactone.

5. The method of claim 3 or 4, wherein the gamma-butyrolactone is chemically converted from 4-hydroxybutyrate.

6. The method of claim 3 or 4, wherein the gamma-butyrolactone is chemically converted from 1,4-butanediol.

7. The method of claim 1 or 2, wherein the downstream product is tetrahydrofuran.

8. The method of claim 1 or 2, wherein the downstream product is pyrrolidone.

9. The method of any one of claims 1-8, wherein the 1,4-butanediol, 4-hydroxybutyrate or gamma-butyrolactone is isolated from a fermentation culture.

10. The method of any one of claims 1-9, wherein the 1,4-butanediol or gamma-butyrolactone is purified by distillation.

Patentansprüche

1. Verfahren zur Herstellung eines nachgelagerten ("downstream") Produkts von 4-Hydroxybutyrat oder 1,4-Butandiol, umfassend:

- (a) das Kultivieren eines nicht natürlich vorkommenden Mikroorganismus, der eine Störung der Gene *adhE*, *ldhA*, *pflAB* und *mdh* umfasst, in exponentieller Wachstumsphase in einer ausreichenden Menge von Nährstoffen und Medien, wobei der Mikroorganismus mindestens eine exogene Nukleinsäure, die ein Enzym in einem 4-

Description**BACKGROUND OF THE INVENTION**

[0001] The present invention relates generally to methods for manufacturing downstream products of 4-hydroxybutyrate or 1,4-butanediol.

[0002] 1,4-Butanediol (BDO) is a four carbon dialcohol that currently is manufactured exclusively through various petrochemical routes. BDO is part of a large volume family of solvents and polymer intermediates that includes gamma-butyrolactone (GBL), tetrahydrofuran (THF), pyrrolidone, N-methylpyrrolidone (NMP), and N-vinyl-pyrrolidone. The overall market opportunity for this family exceeds \$4.0 B.

[0003] Approximately 2.5B lb BDO is produced globally per year with 4-5% annual growth and a recent selling price ranging from \$1.00-1.20/lb. The demand for BDO stems largely from its use as an intermediate for polybutylene terephthalate (PBT) plastic resins, polyurethane thermoplastics and co-polyester ethers. BDO also serves as a primary precursor to THF, which is employed as an intermediate for poly(tetramethylene glycol) PTMEG copolymers required for lycra and spandex production. Approximately 0.7 B lb of THF is produced globally per year with an annual growth rate over 6%. A significant percentage of growth (>30%) for both BDO and THF is occurring in Asia (China and India). GBL currently is a smaller volume (0.4 B lb/year) product which has numerous applications as a solvent, as an additive for inks, paints, and dyes, as well as the primary precursor to pyrrolidone derivatives such as NMP.

[0004] Conventional processes for the synthesis of BDO use petrochemical feedstocks for their starting materials. For example, acetylene is reacted with 2 molecules of formaldehyde in the Reppe synthesis reaction (Kroschwitz and Grant, Encyclopedia of Chem. Tech., John Wiley and Sons, Inc., New York (1999)), followed by catalytic hydrogenation to form 1,4-butanediol. It has been estimated that 90% of the acetylene produced in the U.S. is consumed for butanediol production. Alternatively, it can be formed by esterification and catalytic hydrogenation of maleic anhydride, which is derived from butane. Downstream, butanediol can be further transformed; for example, by oxidation to γ -butyrolactone, which can be further converted to pyrrolidone and N-methyl-pyrrolidone, or hydrogenolysis to tetrahydrofuran (Figure 1). These compounds have varied uses as polymer intermediates, solvents, and additives, and have a combined market of nearly 2 billion lb/year.

[0005] The conventional hydrocarbon feedstock-based approach utilizes methane to produce formaldehyde. Thus, a large percentage of the commercial production of BDO relies on methane as a starting material. The production of acetylene also relies on petroleum-based starting material (see Figure 1). Therefore, the costs of BDO production fluctuate with the price of petroleum and natural gas.

[0006] JPS62285779 (A) relates to a bacterial strain belonging to the *Bacillus* genus (C-105 strain (FERM P-8788)) and production of 1,4-butanediol using the same.

[0007] It is desirable to develop a method for production of these chemicals by alternative means that not only substitute renewable for petroleum-based feedstocks, and also use less energy- and capital-intensive processes. The Department of Energy has proposed 1,4-diacids, and particularly succinic acid, as key biologically-produced intermediates for the manufacture of the butanediol family of products (DOE Report, "Top Value-Added Chemicals from Biomass", 2004). However, succinic acid is costly to isolate and purify and requires high temperatures and pressures for catalytic reduction to butanediol.

[0008] Thus, there exists a need for alternative means for effectively producing commercial quantities of 1,4-butanediol and its chemical precursors. The present invention satisfies this need and provides related advantages as well.

SUMMARY OF INVENTION

[0009] The invention is defined by the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawings will be provided by the Office upon request and payment of the necessary fee.

Figure 1 is a schematic diagram showing an entry point of 4-hydroxybutanoic acid (4-HB) into the product pipeline of the 1,4-butanediol (BDO) family of chemicals, and comparison with chemical synthesis routes from petrochemical feedstocks. Solid black arrows show chemical synthesis routes; dashed blue arrows show a biosynthetic route to 4-HB and subsequent conversion steps to BDO family chemicals.

Figure 2 is a schematic diagram showing biochemical pathways to 4-hydroxybutyrate (4-HB) and γ -butyrolactone (GBL) production. Enzymes catalyzing the 4-HB biosynthetic reactions are: (1) CoA-independent succinic semial-

dehyde dehydrogenase; (2) succinyl-CoA synthetase; (3) CoA-dependent succinic semialdehyde dehydrogenase; (4) glutamate:succinic semialdehyde transaminase; (5) glutamate decarboxylase; (6) 4-hydroxybutanoate dehydrogenase. Conversion (7) corresponds to a spontaneous, non-enzymatic reaction which converts 4-HB to GBL.

Figure 3 is a schematic diagram showing the chemical synthesis of 1,4-butanediol (BDO) and downstream products γ -butyrolactone (GBL), tetrahydrofuran (THF) and several pyrrolidones.

Figure 4 is a schematic process flow diagram of bioprocesses for the production of γ -butyrolactone. Panel (a) illustrates fed-batch fermentation with batch separation and panel (b) illustrates fed-batch fermentation with continuous separation.

Figure 5 shows the hypothetical production envelopes of an OptKnock-designed strain contrasted against a typical non-growth-coupled production strain. Note that the potential evolutionary trajectories of the OptKnock strain are fundamentally different in that they lead to a high producing phenotype.

Figure 6 shows biochemical pathways to 1,4-butanediol. 1) CoA-independent succinic semialdehyde dehydrogenase; 2) succinyl-CoA synthetase; 3) CoA-dependent succinic semialdehyde dehydrogenase; 4) glutamate:succinate semialdehyde transaminase; 5) glutamate decarboxylase; 6) 4-hydroxybutanoate dehydrogenase; 7) 4-hydroxybutyryl CoA:acetyl-CoA transferase; 8) aldehyde dehydrogenase; 9) alcohol dehydrogenase.

Figure 7 shows the anaerobic growth rate versus BDO yield solution boundaries for an *E. coli* strain possessing the BDO production pathways shown and OptKnock predicted knockouts in Figure 5 assuming a) PEP carboxykinase irreversibility and b) PEP carboxykinase reversibility. A basis glucose uptake rate of 20 mmol/gDW/hr is assumed along with a non-growth associated ATP maintenance requirement of 7.6 mmol/gDW/hr.

Figure 8 shows a pictorial representation of *E. coli* central metabolism.

Figure 9 shows the anaerobic growth rate versus BDO yield solution boundaries for an *E. coli* strain possessing the BDO production pathways shown and OptKnock predicted knockouts in Figure 5 assuming PEP carboxykinase reversibility. A basis glucose uptake rate of 20 mmol/gDW/hr is assumed along with a non-growth associated ATP maintenance requirement of 7.6 mmol/gDW/hr.

DETAILED DESCRIPTION OF THE INVENTION

[0011] This invention is directed to the design and production of non-naturally occurring microorganisms having bio-synthetic production capabilities for 4-hydroxybutanoic acid (4-HB), γ -butyrolactone and 1,4-butanediol. The invention utilizes *in silico* stoichiometric models of *Escherichia coli* metabolism that identify metabolic designs for biosynthetic production of 4-hydroxybutanoic acid (4-HB) and 1,4-butanediol (BDO). The results described herein indicate that metabolic pathways can be designed and recombinantly engineered to achieve the biosynthesis of 4-HB and downstream products such as 1,4-butanediol in *Escherichia coli* and other cells or organisms. Biosynthetic production of 4-HB, for example, for the *in silico* designs can be confirmed by construction of strains having the designed metabolic genotype. These metabolically engineered cells or organisms also can be subjected to adaptive evolution to further augment 4-HB biosynthesis, including under conditions approaching theoretical maximum growth.

[0012] The invention is further directed to metabolic engineering strategies for attaining high yields of 1,4-butanediol (BDO) in *Escherichia coli* (see Examples V-VII). As disclosed herein, a genome-scale stoichiometric model of *E. coli* metabolism was employed using the bilevel optimization framework OptKnock to identify *in silico* strategies with multiple knockouts. The deletions are placed such that the redundancy in the network is reduced with the ultimate effect of coupling growth to the production of BDO in the network. The growth-coupled BDO production characteristic of the designed strains make them genetically stable and amenable to continuous bioprocesses. Strain design strategies were identified assuming the addition of non-native reaction capabilities into *E. coli* leading to a metabolic pathway from succinate semialdehyde to BDO. Out of the hundreds of strategies identified by OptKnock, one design emerged as satisfying multiple criteria. This design, utilizing the removal of *adhE*, *ldhA*, *mdh*, *aspA*, and *pflAB*, 1) led to a high predicted BDO yield at maximum growth, 2) required a reasonable number of knockouts, 3) had no detrimental effect on the maximum theoretical BDO yield, 4) brought about a tight coupling of BDO production with cell growth, and 5) was robust with respect to the assumed irreversibility or reversibility of PEP carboxykinase. Also disclosed herein are methods for the experimental testing of the strain designs and their evolution towards the theoretical maximum growth.

[0013] In certain embodiments, the 4-HB biosynthesis characteristics of the designed strains make them genetically stable and particularly useful in continuous bioprocesses. Separate strain design strategies were identified with incor-

poration of different non-native or heterologous reaction capabilities into *E. coli* leading to 4-HB and 1,4-butanediol producing metabolic pathways from either CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase and CoA-dependent succinic semialdehyde dehydrogenase, or glutamate:succinic semialdehyde transaminase. *In silico* metabolic designs were identified that resulted in the biosynthesis of 4-HB in both *E. coli* and yeast species from each of these metabolic pathways. The 1,4-butanediol intermediate γ -butyrolactone can be generated in culture by spontaneous cyclization under conditions at pH<7.5, particularly under acidic conditions, such as below pH 5.5, for example, pH<7, pH<6.5, pH<6, and particularly at pH<5.5 or lower.

[0014] Strains identified via the computational component of the platform can be put into actual production by genetically engineering any of the predicted metabolic alterations which lead to the biosynthetic production of 4-HB, 1,4-butanediol or other intermediate and/or downstream products. In yet a further embodiment, strains exhibiting biosynthetic production of these compounds can be further subjected to adaptive evolution to further augment product biosynthesis. The levels of product biosynthesis yield following adaptive evolution also can be predicted by the computational component of the system.

[0015] As used herein, the term "non-naturally occurring" when used in reference to a microbial organism or microorganism of the disclosure is intended to mean that the microbial organism has at least one genetic alteration not normally found in a naturally occurring strain of the referenced species, including wild-type strains of the referenced species. Genetic alterations include, for example, modifications introducing expressible nucleic acids encoding metabolic polypeptides, other nucleic acid additions, nucleic acid deletions and/or other functional disruption of the microbial genetic material. Such modification include, for example, coding regions and functional fragments thereof, for heterologous, homologous or both heterologous and homologous polypeptides for the referenced species. Additional modifications include, for example, non-coding regulatory regions in which the modifications alter expression of a gene or operon. Exemplary metabolic polypeptides include enzymes within a 4-HB biosynthetic pathway and enzymes within a biosynthetic pathway for a BDO family of compounds.

[0016] A metabolic modification refers to a biochemical reaction that is altered from its naturally occurring state. Therefore, non-naturally occurring microorganisms having genetic modifications to nucleic acids encoding metabolic polypeptides or, functional fragments thereof. Exemplary metabolic modifications are described further below for both *E. coli* and yeast microbial organisms.

[0017] As used herein, the term "isolated" when used in reference to a microbial organism is intended to mean an organism that is substantially free of at least one component as the referenced microbial organism is found in nature. The term includes a microbial organism that is removed from some or all components as it is found in its natural environment. The term also includes a microbial organism that is removed from some or all components as the microbial organism is found in non-naturally occurring environments. Therefore, an isolated microbial organism is partly or completely separated from other substances as it is found in nature or as it is grown, stored or subsisted in non-naturally occurring environments. Specific examples of isolated microbial organisms include partially pure microbes, substantially pure microbes and microbes cultured in a medium that is non-naturally occurring.

[0018] As used herein, the terms "microbial," "microbial organism" or "microorganism" is intended to mean any organism that exists as a microscopic cell that is included within the domains of archaea, bacteria or eukarya. Therefore, the term is intended to encompass prokaryotic or eukaryotic cells or organisms having a microscopic size and includes bacteria, archaea and eubacteria of all species as well as eukaryotic microorganisms such as yeast and fungi. The term also includes cell cultures of any species that can be cultured for the production of a biochemical.

[0019] As used herein, the term "4-hydroxybutanoic acid" is intended to mean a 4-hydroxy derivative of butyric acid having the chemical formula $C_4H_8O_3$ and a molecular mass of 104.11 g/mol (126.09 g/mol for its sodium salt). The chemical compound 4-hydroxybutanoic acid also is known in the art as 4-HB, 4-hydroxybutyrate, gamma-hydroxybutyric acid or GHB. The term as it is used herein is intended to include any of the compound's various salt forms and include, for example, 4-hydroxybutanoate and 4-hydroxybutyrate. Specific examples of salt forms for 4-HB include sodium 4-HB and potassium 4-HB. Therefore, the terms 4-hydroxybutanoic acid, 4-HB, 4-hydroxybutyrate, 4-hydroxybutanoate, gamma-hydroxybutyric acid and GHB as well as other art recognized names are used synonymously herein.

[0020] As used herein, the term "monomeric" when used in reference to 4-HB is intended to mean 4-HB in a non-polymeric or underivatized form. Specific examples of polymeric 4-HB include poly-4-hydroxybutanoic acid and copolymers of, for example, 4-HB and 3-HB. A specific example of a derivatized form of 4-HB is 4-HB-CoA. Other polymeric 4-HB forms and other derivatized forms of 4-HB also are known in the art.

[0021] As used herein, the term " γ -butyrolactone" is intended to mean a lactone having the chemical formula $C_4H_6O_2$ and a molecular mass of 86.089 g/mol. The chemical compound γ -butyrolactone also is known in the art as GBL, butyrolactone, 1,4-lactone, 4-butyrolactone, 4-hydroxybutyric acid lactone, and gamma-hydroxybutyric acid lactone. The term as it is used herein is intended to include any of the compound's various salt forms.

[0022] As used herein, the term "1,4 butanediol" is intended to mean an alcohol derivative of the alkane butane, carrying two hydroxyl groups which has the chemical formula $C_4H_{10}O_2$ and a molecular mass of 90.12 g/mol. The chemical compound 1,4 butanediol also is known in the art as BDO and is a chemical intermediate or precursor for a

family of compounds referred to herein as BDO family of compounds, some of which are exemplified in Figure 1.

[0023] As used herein, the term "tetrahydrofuran" is intended to mean a heterocyclic organic compound corresponding to the fully hydrogenated analog of the aromatic compound furan which has the chemical formula C_4H_8O and a molecular mass of 72.11 g/mol. The chemical compound tetrahydrofuran also is known in the art as THF, tetrahydrofuran, 1,4-epoxybutane, butylene oxide, cyclotetramethylene oxide, oxacyclopentane, diethylene oxide, oxolane, furanidine, hydrofuran, tetra-methylene oxide. The term as it is used herein is intended to include any of the compound's various salt forms.

[0024] As used herein, the term "CoA" or "coenzyme A" is intended to mean an organic cofactor or prosthetic group (nonprotein portion of an enzyme) whose presence is required for the activity of many enzymes (the apoenzyme) to form an active enzyme system. Coenzyme A functions in certain condensing enzymes, acts in acetyl or other acyl group transfer and in fatty acid synthesis and oxidation, pyruvate oxidation and in other acetylation.

[0025] As used herein, the term "substantially anaerobic" when used in reference to a culture or growth condition is intended to mean that the amount of oxygen is less than about 10% of saturation for dissolved oxygen in liquid media. The term also is intended to include sealed chambers of liquid or solid medium maintained with an atmosphere of less than about 1% oxygen.

[0026] The non-naturally occurring microbial organisms as disclosed can contain stable genetic alterations, which refers to microorganisms that can be cultured for greater than five generations without loss of the alteration. Generally, stable genetic alterations include modifications that persist greater than 10 generations, particularly stable modifications will persist more than about 25 generations, and more particularly, stable genetic modifications will be greater than 50 generations, including indefinitely.

[0027] Those skilled in the art will understand that the genetic alterations, including metabolic modifications exemplified herein are described with reference to *E. coli* and yeast genes and their corresponding metabolic reactions. However, given the complete genome sequencing of a wide variety of organisms and the high level of skill in the area of genomics, those skilled in the art will readily be able to apply the teachings and guidance provided herein to essentially all other organisms. For example, the *E. coli* metabolic alterations exemplified herein can readily be applied to other species by incorporating the same or analogous encoding nucleic acid from species other than the referenced species. Such genetic alterations include, for example, genetic alterations of species homologs, in general, and in particular, orthologs, paralogs or nonorthologous gene displacements.

[0028] An ortholog is a gene or genes that are related by vertical descent and are responsible for substantially the same or identical functions in different organisms. For example, mouse epoxide hydrolase and human epoxide hydrolase can be considered orthologs for the biological function of hydrolysis of epoxides. Genes are related by vertical descent when, for example, they share sequence similarity of sufficient amount to indicate they are homologous, or related by evolution from a common ancestor. Genes can also be considered orthologs if they share three-dimensional structure but not necessarily sequence similarity, of a sufficient amount to indicate that they have evolved from a common ancestor to the extent that the primary sequence similarity is not identifiable. Genes that are orthologous can encode proteins with sequence similarity of about 25% to 100% amino acid sequence identity. Genes encoding proteins sharing an amino acid similarity less than 25% can also be considered to have arisen by vertical descent if their three-dimensional structure also shows similarities. Members of the serine protease family of enzymes, including tissue plasminogen activator and elastase, are considered to have arisen by vertical descent from a common ancestor.

[0029] Orthologs include genes or their encoded gene products that through, for example, evolution, have diverged in structure or overall activity. For example, where one species encodes a gene product exhibiting two functions and where such functions have been separated into distinct genes in a second species, the three genes and their corresponding products are considered to be orthologs. For the growth-coupled production of a biochemical product, those skilled in the art will understand that the orthologous gene harboring the metabolic activity to be disrupted is to be chosen for construction of the non-naturally occurring microorganism. An example of orthologs exhibiting separable activities is where distinct activities have been separated into distinct gene products between two or more species or within a single species. A specific example is the separation of elastase proteolysis and plasminogen proteolysis, two types of serine protease activity, into distinct molecules as plasminogen activator and elastase. A second example is the separation of mycoplasma 5'-3' exonuclease and *Drosophila* DNA polymerase III activity. The DNA polymerase from the first species can be considered an ortholog to either or both of the exonuclease or the polymerase from the second species and vice versa.

[0030] In contrast, paralogs are homologs related by, for example, duplication followed by evolutionary divergence and have similar or common, but not identical functions. Paralogs can originate or derive from, for example, the same species or from a different species. For example, microsomal epoxide hydrolase (epoxide hydrolase I) and soluble epoxide hydrolase (epoxide hydrolase II) can be considered paralogs because they represent two distinct enzymes, co-evolved from a common ancestor, that catalyze distinct reactions and have distinct functions in the same species. Paralogs are proteins from the same species with significant sequence similarity to each other suggesting that they are homologous, or related through co-evolution from a common ancestor. Groups of paralogous protein families include

HipA homologs, luciferase genes, peptidases, and others.

[0031] A nonorthologous gene displacement is a nonorthologous gene from one species that can substitute for a referenced gene function in a different species. Substitution includes, for example, being able to perform substantially the same or a similar function in the species of origin compared to the referenced function in the different species. Although generally, a nonorthologous gene displacement will be identifiable as structurally related to a known gene encoding the referenced function, less structurally related but functionally similar genes and their corresponding gene products nevertheless will still fall within the meaning of the term as it is used herein. Functional similarity requires, for example, at least some structural similarity in the active site or binding region of a nonorthologous gene compared to a gene encoding the function sought to be substituted. Therefore, a nonorthologous gene includes, for example, a paralog or an unrelated gene.

[0032] Therefore, in identifying and constructing the non-naturally occurring microbial organisms having 4-HB, GBL and/or BDO biosynthetic capability, those skilled in the art will understand with applying the teaching and guidance provided herein to a particular species that the identification of metabolic modifications can include identification and inclusion or inactivation of orthologs. To the extent that paralogs and/or nonorthologous gene displacements are present in the referenced microorganism that encode an enzyme catalyzing a similar or substantially similar metabolic reaction, those skilled in the art also can utilize these evolutionally related genes.

[0033] Orthologs, paralogs and nonorthologous gene displacements can be determined by methods well known to those skilled in the art. For example, inspection of nucleic acid or amino acid sequences for two polypeptides will reveal sequence identity and similarities between the compared sequences. Based on such similarities, one skilled in the art can determine if the similarity is sufficiently high to indicate the proteins are related through evolution from a common ancestor. Algorithms well known to those skilled in the art, such as Align, BLAST, Clustal W and others compare and determine a raw sequence similarity or identity, and also determine the presence or significance of gaps in the sequence which can be assigned a weight or score. Such algorithms also are known in the art and are similarly applicable for determining nucleotide sequence similarity or identity. Parameters for sufficient similarity to determine relatedness are computed based on well known methods for calculating statistical similarity, or the chance of finding a similar match in a random polypeptide, and the significance of the match determined. A computer comparison of two or more sequences can, if desired, also be optimized visually by those skilled in the art. Related gene products or proteins can be expected to have a high similarity, for example, 25% to 100% sequence identity. Proteins that are unrelated can have an identity which is essentially the same as would be expected to occur by chance, if a database of sufficient size is scanned (about 5%). Sequences between 5% and 24% may or may not represent sufficient homology to conclude that the compared sequences are related. Additional statistical analysis to determine the significance of such matches given the size of the data set can be carried out to determine the relevance of these sequences.

[0034] Exemplary parameters for determining relatedness of two or more sequences using the BLAST algorithm, for example, can be as set forth below. Briefly, amino acid sequence alignments can be performed using BLASTP version 2.0.8 (Jan-05-1999) and the following parameters: Matrix: 0 BLOSUM62; gap open: 11; gap extension: 1; x_dropoff: 50; expect: 10.0; wordsize: 3; filter: on. Nucleic acid sequence alignments can be performed using BLASTN version 2.0.6 (Sept-16-1998) and the following parameters: Match: 1; mismatch: -2; gap open: 5; gap extension: 2; x_dropoff: 50; expect: 10.0; wordsize: 11; filter: off. Those skilled in the art will know what modifications can be made to the above parameters to either increase or decrease the stringency of the comparison, for example, and determine the relatedness of two or more sequences.

[0035] The disclosure provides a non-naturally occurring microbial biocatalyst including a microbial organism having a 4-hydroxybutanoic acid (4-HB) biosynthetic pathway that includes at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, glutamate:succinic semialdehyde transaminase or glutamate decarboxylase, wherein the exogenous nucleic acid is expressed in sufficient amounts to produce monomeric 4-hydroxybutanoic acid (4-HB). Succinyl-CoA synthetase is also referred to as succinyl-CoA synthase or succinyl CoA ligase.

[0036] The non-naturally occurring microbial biocatalysts of the disclosure include microbial organisms that employ combinations of metabolic reactions for biosynthetically producing the compounds of the disclosure. Exemplary compounds produced by the non-naturally occurring microorganisms include, for example, 4-hydroxybutanoic acid, 1,4-butanediol and γ -butyrolactone. The relationships of these exemplary compounds with respect to chemical synthesis or biosynthesis are exemplified in Figure 1.

[0037] In one embodiment, a non-naturally occurring microbial organism is engineered to produce 4-HB. This compound is one useful entry point into the 1,4-butanediol family of compounds. The biochemical reactions for formation of 4-HB from succinate, from succinate through succinyl-CoA or from α -ketoglutarate are shown in Figure 2.

[0038] The invention is described herein with general reference to the metabolic reaction, reactant or product thereof, or with specific reference to one or more nucleic acids or genes encoding an enzyme associated with or catalyzing the referenced metabolic reaction, reactant or product. Unless otherwise expressly stated herein, those skilled in the art will understand that reference to a reaction also constitutes reference to the reactants and products of the reaction. Similarly,

unless otherwise expressly stated herein, reference to a reactant or product also references the reaction and that reference to any of these metabolic constituents also references the gene or genes encoding the enzymes that catalyze the referenced reaction, reactant or product. Likewise, given the well known fields of metabolic biochemistry, enzymology and genomics, reference herein to a gene or encoding nucleic acid also constitutes a reference to the corresponding encoded enzyme and the reaction it catalyzes as well as the reactants and products of the reaction.

[0039] The production of 4-HB via biosynthetic modes using the microbial organisms of the disclosure is particularly useful because it results in monomeric 4-HB. The non-naturally occurring microbial organisms of the disclosure and their biosynthesis of 4-HB and BDO family compounds also is particularly useful because the 4-HB product (1) is secreted; (2) is devoid of any derivatizations such as Coenzyme A; (3) avoids thermodynamic changes during biosynthesis, and (4) allows for the spontaneous chemical conversion of 4-HB to γ -butyrolactone (GBL) in acidic pH medium. This latter characteristic is exemplified as step 7 in Figure 2 and also is particularly useful for efficient chemical synthesis or biosynthesis of BDO family compounds such as 1,4-butanediol and/or tetrahydrofuran (THF), for example.

[0040] Microbial organisms generally lack the capacity to synthesize 4-HB and therefore, any of the compounds shown in Figure 1 are known to be within the 1,4-butanediol family of compounds or known by those in the art to be within the 1,4-butanediol family of compounds. Moreover, organisms having all of the requisite metabolic enzymatic capabilities are not known to produce 4-HB from the enzymes described and biochemical pathways exemplified herein. Rather, with the possible exception of a few anaerobic microorganisms described further below, the microorganisms having the enzymatic capability use 4-HB as a substrate to produce, for example, succinate. In contrast, the non-naturally occurring microbial organisms of the disclosure generate 4-HB as a product. As described above, the biosynthesis of 4-HB in its monomeric form is not only particularly useful in chemical synthesis of BDO family of compounds, it also allows for the further biosynthesis of BDO family compounds and avoids altogether chemical synthesis procedures.

[0041] The non-naturally occurring microbial organisms of the disclosure that can produce monomeric 4-HB are produced by ensuring that a host microbial organism includes functional capabilities for the complete biochemical synthesis of at least one 4-HB biosynthetic pathway of the disclosure. Ensuring at least one requisite 4-HB biosynthetic pathway confers 4-HB biosynthesis capability onto the host microbial organism.

[0042] Three requisite 4-HB biosynthetic pathways are exemplified herein and shown for purposes of illustration in Figure 2. One requisite 4-HB biosynthetic pathway includes the biosynthesis of 4-HB from succinate. The enzymes participating in this 4-HB pathway include CoA-independent succinic semialdehyde dehydrogenase and 4-hydroxybutanoate dehydrogenase. Another requisite 4-HB biosynthetic pathway includes the biosynthesis from succinate through succinyl-CoA. The enzymes participating in this 4-HB pathway include succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase and 4-hydroxybutanoate dehydrogenase. A third requisite 4-HB biosynthetic pathway includes the biosynthesis of 4-HB from α -ketoglutarate. The enzymes participating in this 4-HB biosynthetic pathway include glutamate:succinic semialdehyde transaminase, glutamate decarboxylase and 4-hydroxybutanoate dehydrogenase. Each of these 4-HB biosynthetic pathways, their substrates, reactants and products are described further below in the Examples.

[0043] The non-naturally occurring microbial organisms of the disclosure can be produced by introducing expressible nucleic acids encoding one or more of the enzymes participating in one or more 4-HB biosynthetic pathways. Depending on the host microbial organism chosen for biosynthesis, nucleic acids for some or all of a particular 4-HB biosynthetic pathway can be expressed. For example, if a chosen host is deficient in both enzymes in the succinate to 4-HB pathway (the succinate pathway) and this pathway is selected for 4-HB biosynthesis, then expressible nucleic acids for both CoA-independent succinic semialdehyde dehydrogenase and 4-hydroxybutanoate dehydrogenase are introduced into the host for subsequent exogenous expression. Alternatively, if the chosen host is deficient in 4-hydroxybutanoate dehydrogenase then an encoding nucleic acid is needed for this enzyme to achieve 4-HB biosynthesis.

[0044] In like fashion, where 4-HB biosynthesis is selected to occur through the succinate to succinyl-CoA pathway (the succinyl-CoA pathway), encoding nucleic acids for host deficiencies in the enzymes succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase and/or 4-hydroxybutanoate dehydrogenase are to be exogenously expressed in the recipient host. Selection of 4-HB biosynthesis through the α -ketoglutarate to succinic semialdehyde pathway (the α -ketoglutarate pathway) will utilize exogenous expression for host deficiencies in one or more of the enzymes for glutamate:succinic semialdehyde transaminase, glutamate decarboxylase and/or 4-hydroxybutanoate dehydrogenase.

[0045] Depending on the 4-HB biosynthetic pathway constituents of a selected host microbial organism, the non-naturally occurring microbial 4-HB biocatalysts of the invention will include at least one exogenously expressed 4-HB pathway-encoding nucleic acid and up to all six 4-HB pathway encoding nucleic acids. For example, 4-HB biosynthesis can be established from all three pathways in a host deficient in 4-hydroxybutanoate dehydrogenase through exogenous expression of a 4-hydroxybutanoate dehydrogenase encoding nucleic acid. In contrast, 4-HB biosynthesis can be established from all three pathways in a host deficient in all six enzymes through exogenous expression of all six of CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase and 4-hydroxybutanoate

dehydrogenase.

[0046] Given the teachings and guidance provided herein, those skilled in the art will understand that the number of encoding nucleic acids to introduce in an expressible form will parallel the 4-HB pathway deficiencies of the selected host microbial organism. Therefore, a non-naturally occurring microbial organism of the disclosure can have one, two, three, four, five or six encoding nucleic acids encoding the above enzymes constituting the 4-HB biosynthetic pathways. In some embodiments, the non-naturally occurring microbial organisms also can include other genetic modifications that facilitate or optimize 4-HB biosynthesis or that confer other useful functions onto the host microbial organism. One such other functionality can include, for example, augmentation of the synthesis of one or more of the 4-HB pathway precursors such as succinate, succinyl-CoA and/or α -ketoglutarate.

[0047] In some embodiments, a non-naturally occurring microbial organism of the disclosure is generated from a host that contains the enzymatic capability to synthesize 4-HB. In this specific embodiment it can be useful to increase the synthesis or accumulation of a 4-HB pathway product to, for example, drive 4-HB pathway reactions toward 4-HB production. Increased synthesis or accumulation can be accomplished by, for example, overexpression of nucleic acids encoding one or more of the above-described 4-HB pathway enzymes. Over expression of the 4-HB pathway enzyme or enzymes can occur, for example, through exogenous expression of the endogenous gene or genes, or through exogenous expression of the heterologous gene or genes. Therefore, naturally occurring organisms can be readily generated to be non-naturally 4-HB producing microbial organisms of the disclosure through overexpression of one, two, three, four, five or all six nucleic acids encoding 4-HB biosynthetic pathway enzymes. In addition, a non-naturally occurring organism can be generated by mutagenesis of an endogenous gene that results in an increase in activity of an enzyme in the 4-HB biosynthetic pathway.

[0048] In particularly useful embodiments, exogenous expression of the encoding nucleic acids is employed. Exogenous expression confers the ability to custom tailor the expression and/or regulatory elements to the host and application to achieve a desired expression level that is controlled by the user. However, endogenous expression also can be utilized in other embodiments such as by removing a negative regulatory effector or induction of the gene's promoter when linked to an inducible promoter or other regulatory element. Thus, an endogenous gene having a naturally occurring inducible promoter can be up-regulated by providing the appropriate inducing agent, or the regulatory region of an endogenous gene can be engineered to incorporate an inducible regulatory element, thereby allowing the regulation of increased expression of an endogenous gene at a desired time. Similarly, an inducible promoter can be included as a regulatory element for an exogenous gene introduced into a non-naturally occurring microbial organism (see Example II).

[0049] "Exogenous" as it is used herein is intended to mean that the referenced molecule or the referenced activity is introduced into the host microbial organism. Therefore, the term as it is used in reference to expression of an encoding nucleic acid refers to introduction of the encoding nucleic acid in an expressible form into the microbial organism. When used in reference to a biosynthetic activity, the term refers to an activity that is introduced into the host reference organism. The source can be, for example, a homologous or heterologous encoding nucleic acids that expresses the referenced activity following introduction into the host microbial organism. Therefore, the term "endogenous" refers to a referenced molecule or activity that is present in the host. Similarly, the term when used in reference to expression of an encoding nucleic acid refers to expression of an encoding nucleic acid contained within the microbial organism. The term "heterologous" refers to a molecule or activity derived from a source other than the referenced species whereas "homologous" refers to a molecule or activity derived from the host microbial organism. Accordingly, exogenous expression of an encoding nucleic acid can utilize either or both a heterologous or homologous encoding nucleic acid.

[0050] Sources of encoding nucleic acids for a 4-HB pathway enzyme can include, for example, any species where the encoded gene product is capable of catalyzing the referenced reaction. Such species include both prokaryotic and eukaryotic organisms including, but not limited to, bacteria, including archaea and eubacteria, and eukaryotes, including yeast, plant, insect, animal, and mammal, including human. For example, the microbial organisms having 4-HB biosynthetic production are exemplified herein with reference to *E. coli* and yeast hosts. However, with the complete genome sequence available for now more than 550 species (with more than half of these available on public databases such as the NCBI), including 395 microorganism genomes and a variety of yeast, fungi, plant, and mammalian genomes, the identification of genes encoding the requisite 4-HB biosynthetic activity for one or more genes in related or distant species, including for example, homologues, orthologs, paralogs and nonorthologous gene displacements of known genes, and the interchange of genetic alterations between organisms is routine and well known in the art. Accordingly, the metabolic alterations enabling biosynthesis of 4-HB and other compounds of the invention described herein with reference to a particular organism such as *E. coli* or yeast can be readily applied to other microorganisms, including prokaryotic and eukaryotic organisms alike. Given the teachings and guidance provided herein, those skilled in the art will know that a metabolic alteration exemplified in one organism can be applied equally to other organisms.

[0051] In some instances, such as when an alternative 4-HB biosynthetic pathway exists in an unrelated species, 4-HB biosynthesis can be conferred onto the host species by, for example, exogenous expression of a paralog or paralogs from the unrelated species that catalyzes a similar, yet non-identical metabolic reaction to replace the referenced reaction. Because certain differences among metabolic networks exist between different organisms, those skilled in the art will

understand that the actual genes usage between different organisms may differ. However, given the teachings and guidance provided herein, those skilled in the art also will understand that the teachings and methods of the invention can be applied to all microbial organisms using the cognate metabolic alterations to those exemplified herein to construct a microbial organism in a species of interest that will synthesize monomeric 4-HB.

[0052] Host microbial organisms can be selected from, and the non-naturally occurring microbial organisms generated in, for example, bacteria, yeast, fungus or any of a variety of other microorganisms applicable to fermentation processes. Exemplary bacteria include species selected from *E. coli*, *Klebsiella oxytoca*, *Anaerobiospirillum succiniciproducens*, *Actinobacillus succinogenes*, *Mannheimia succiniciproducens*, *Rhizobium etli*, *Bacillus subtilis*, *Corynebacterium glutamicum*, *Gluconobacter oxydans*, *Zymomonas mobilis*, *Lactococcus lactis*, *Lactobacillus plantarum*, *Streptomyces coelicolor*, *Clostridium acetobutylicum*, *Pseudomonas fluorescens*, and *Pseudomonas putida*. Exemplary yeasts or fungi include species selected from *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces marxianus*, *Aspergillus terreus*, *Aspergillus niger* and *Pichia pastoris*.

[0053] Methods for constructing and testing the expression levels of a non-naturally occurring 4-HB-producing host can be performed, for example, by recombinant and detection methods well known in the art. Such methods can be found described in, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Third Ed., Cold Spring Harbor Laboratory, New York (2001); Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, MD (1999). 4-HB and GBL can be separated by, for example, HPLC using a Spherisorb 5 ODS1 column and a mobile phase of 70% 10 mM phosphate buffer (pH=7) and 30% methanol, and detected using a UV detector at 215 nm (Hennessy et al. *J. Forensic Sci.* 46(6):1-9 (2004)). BDO is detected by gas chromatography or by HPLC and refractive index detector using an Aminex HPX-87H column and a mobile phase of 0.5 mM sulfuric acid (Gonzalez-Pajuelo et al., *Met. Eng.* 7:329-336 (2005)).

[0054] The non-naturally occurring microbial organisms of the disclosure are constructed using methods well known in the art as exemplified above to exogenously express at least one nucleic acid encoding a 4-HB pathway enzyme in sufficient amounts to produce monomeric 4-HB. Exemplary levels of expression for 4-HB enzymes in each pathway are described further below in the Examples. Following the teachings and guidance provided herein, the non-naturally occurring microbial organisms of the disclosure can achieve biosynthesis of monomeric 4-HB resulting in intracellular concentrations between about 0.1-25 mM or more. Generally, the intracellular concentration of monomeric 4-HB is between about 3-20mM, particularly between about 5-15 mM and more particularly between about 8-12 mM, including about 10 mM or more. Intracellular concentrations between and above each of these exemplary ranges also can be achieved from the non-naturally occurring microbial organisms of the disclosure. As described further below, one exemplary growth condition for achieving biosynthesis of 4-HB includes anaerobic culture or fermentation conditions. In certain embodiments, the non-naturally occurring microbial organisms of the disclosure can be sustained, cultured or fermented under anaerobic or substantially anaerobic conditions. Briefly, anaerobic conditions refers to an environment devoid of oxygen. Substantially anaerobic conditions include, for example, a culture, batch fermentation or continuous fermentation such that the dissolved oxygen concentration in the medium remains between 0 and 10% of saturation. Substantially anaerobic conditions also includes growing or resting cells in liquid medium or on solid agar inside a sealed chamber maintained with an atmosphere of less than 1 % oxygen. The percent of oxygen can be maintained by, for example, sparging the culture with an N₂/CO₂ mixture or other suitable non-oxygen gas or gases.

[0055] The disclosure also provides a non-naturally occurring microbial biocatalyst including a microbial organism having 4-hydroxybutanoic acid (4-HB) and 1,4-butanediol (BDO) biosynthetic pathways that include at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or alcohol dehydrogenase, wherein the exogenous nucleic acid is expressed in sufficient amounts to produce 1,4-butanediol (BDO).

[0056] Non-naturally occurring microbial organisms also can be generated which biosynthesize BDO. Following the teachings and guidance provided previously for the construction of microbial organisms that synthesize 4-HB, additional BDO pathways can be incorporated into the 4-HB producing microbial organisms to generate organisms that also synthesize BDO and other BDO family compounds. The chemical synthesis of BDO and its downstream products are illustrated in Figure 3. The non-naturally occurring microbial organisms of the disclosure capable of BDO biosynthesis circumvent these chemical synthesis using 4-HB as an entry point as illustrated in Figure 2. As described further below, the 4-HB producers can be used to chemically convert 4-HB to GBL and then to BDO or THF, for example. Alternatively, the 4-HB producers can be further modified to include biosynthetic capabilities for conversion of 4-HB and/or GBL to BDO.

[0057] The additional BDO pathways to introduce into 4-HB producers include, for example, the exogenous expression in a host deficient background or the overexpression of a CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or an alcohol dehydrogenase. In the absence of endogenous acyl-CoA synthetase capable of modifying 4-HB, the non-naturally occurring BDO producing microbial organisms can further include an exogenous acyl-CoA synthetase selective for 4-HB. Exemplary alcohol and aldehyde dehydrogenases that can be used for these

in vivo conversions from 4-HB to BDO are listed below in Table 1.

Table 1. Alcohol and Aldehyde Dehydrogenases for Conversion of 4-HB to BDO.

	ALCOHOL	DEHYDROGENASES
5	ec:1.1.1.1	alcohol dehydrogenase
	ec:1.1.1.2	alcohol dehydrogenase (NADP+)
	ec:1.1.1.4	(R,R)-butanediol dehydrogenase
	ec:1.1.1.5	acetoin dehydrogenase
10	ec:1.1.1.6	glycerol dehydrogenase
	ec:1.1.1.7	propanediol-phosphate dehydrogenase
	ec:1.1.1.8	glycerol-3-phosphate dehydrogenase (NAD+)
	ec:1.1.1.11	D-arabinitol 4-dehydrogenase
	ec:1.1.1.12	L-arabinitol 4-dehydrogenase
15	ec:1.1.1.13	L-arabinitol 2-dehydrogenase
	ec:1.1.1.14	L-identol 2-dehydrogenase
	ec:1.1.1.15	D-identol 2-dehydrogenase
	ec:1.1.1.16	galactitol 2-dehydrogenase
20	ec:1.1.1.17	mannitol-1-phosphate 5-dehydrogenase
	ec:1.1.1.18	inositol 2-dehydrogenase
	ec:1.1.1.21	aldehyde reductase
	ec:1.1.1.23	histidinol dehydrogenase
	ec:1.1.1.26	glyoxylate reductase
25	ec:1.1.1.27	L-lactate dehydrogenase
	ec:1.1.1.28	D-lactate dehydrogenase
	ec:1.1.1.29	glycerate dehydrogenase
	ec:1.1.1.30	3-hydroxybutyrate dehydrogenase
30	ec:1.1.1.31	3-hydroxyisobutyrate dehydrogenase
	ec:1.1.1.35	3-hydroxyacyl-CoA dehydrogenase
	ec:1.1.1.36	acetoacetyl-CoA reductase
	ec:1.1.1.37	malate dehydrogenase
	ec:1.1.1.38	malate dehydrogenase (oxaloacetate-decarboxylating)
35	ec:1.1.1.39	malate dehydrogenase (decarboxylating)
	ec:1.1.1.40	malate dehydrogenase (oxaloacetate-decarboxylating) (NADP+)
	ec:1.1.1.41	isocitrate dehydrogenase (NAD+)
	ec:1.1.1.42	isocitrate dehydrogenase (NADP+)
40	ec:1.1.1.54	allyl-alcohol dehydrogenase
	ec:1.1.1.55	lactaldehyde reductase (NADPH)
	ec:1.1.1.56	ribitol 2-dehydrogenase
	ec:1.1.1.59	3-hydroxypropionate dehydrogenase
	ec:1.1.1.60	2-hydroxy-3-oxopropionate reductase
45	ec:1.1.1.61	4-hydroxybutyrate dehydrogenase
	ec:1.1.1.66	omega-hydroxydecanoate dehydrogenase
	ec:1.1.1.67	mannitol 2-dehydrogenase
	ec:1.1.1.71	alcohol dehydrogenase [NAD(P)+]
50	ec:1.1.1.72	glycerol dehydrogenase (NADP+)
	ec:1.1.1.73	octanol dehydrogenase
	ec:1.1.1.75	(R)-aminopropanol dehydrogenase
	ec:1.1.1.76	(S,S)-butanediol dehydrogenase
	ec:1.1.1.77	lactaldehyde reductase
55	ec:1.1.1.78	methylglyoxal reductase (NADH-dependent)
	ec:1.1.1.79	glyoxylate reductase (NADP+)
	ec:1.1.1.80	isopropanol dehydrogenase (NADP+)

(continued)

ALCOHOL**DEHYDROGENASES**

	ec:1.1.1.81	hydroxypyruvate reductase
5	ec:1.1.1.82	malate dehydrogenase (NADP+)
	ec:1.1.1.83	D-malate dehydrogenase (decarboxylating)
	ec:1.1.1.84	dimethylmalate dehydrogenase
	ec:1.1.1.85	3-isopropylmalate dehydrogenase
10	ec:1.1.1.86	ketol-acid reductoisomerase
	ec:1.1.1.87	homoisocitrate dehydrogenase
	ec:1.1.1.88	hydroxymethylglutaryl-CoA reductase
	ec:1.1.1.90	aryl-alcohol dehydrogenase
	ec:1.1.1.91	aryl-alcohol dehydrogenase (NADP+)
15	ec:1.1.1.92	oxaloglycolate reductase (decarboxylating)
	ec:1.1.1.94	glycerol-3-phosphate dehydrogenase [NAD(P)+]
	ec:1.1.1.95	phosphoglycerate dehydrogenase
	ec:1.1.1.97	3-hydroxybenzyl-alcohol dehydrogenase
20	ec:1.1.1.101	acylglycerone-phosphate reductase
	ec:1.1.1.103	L-threonine 3-dehydrogenase
	ec:1.1.1.104	4-oxoproline reductase
	ec:1.1.1.105	retinol dehydrogenase
	ec:1.1.1.110	indolelactate dehydrogenase
25	ec:1.1.1.112	indanol dehydrogenase
	ec:1.1.1.113	L-xylose 1-dehydrogenase
	ec:1.1.1.129	L-threonate 3-dehydrogenase
	ec:1.1.1.137	ribitol-5-phosphate 2-dehydrogenase
	ec:1.1.1.138	mannitol 2-dehydrogenase (NADP+)
30	ec:1.1.1.140	sorbitol-6-phosphate 2-dehydrogenase
	ec:1.1.1.142	D-pinitol dehydrogenase
	ec:1.1.1.143	sequoyitol dehydrogenase
	ec:1.1.1.144	perillyl-alcohol dehydrogenase
35	ec:1.1.1.156	glycerol 2-dehydrogenase (NADP+)
	ec:1.1.1.157	3-hydroxybutyryl-CoA dehydrogenase
	ec:1.1.1.163	cyclopentanol dehydrogenase
	ec:1.1.1.164	hexadecanol dehydrogenase
	ec:1.1.1.165	2-alkyn-1-ol dehydrogenase
40	ec:1.1.1.166	hydroxycyclohexanecarboxylate dehydrogenase
	ec:1.1.1.167	hydroxymalonate dehydrogenase
	ec:1.1.1.174	cyclohexane-1,2-diol dehydrogenase
	ec:1.1.1.177	glycerol-3-phosphate 1-dehydrogenase (NADP+)
45	ec:1.1.1.178	3-hydroxy-2-methylbutyryl-CoA dehydrogenase
	ec:1.1.1.185	L-glycol dehydrogenase
	ec:1.1.1.190	indole-3-acetaldehyde reductase (NADH)
	ec:1.1.1.191	indole-3-acetaldehyde reductase (NADPH)
	ec:1.1.1.192	long-chain-alcohol dehydrogenase
50	ec:1.1.1.194	coniferyl-alcohol dehydrogenase
	ec:1.1.1.195	cinnamyl-alcohol dehydrogenase
	ec:1.1.1.198	(+)-borneol dehydrogenase
	ec:1.1.1.202	1,3-propanediol dehydrogenase
55	ec:1.1.1.207	(-)-menthol dehydrogenase
	ec:1.1.1.208	(+)-neomenthol dehydrogenase
	ec:1.1.1.216	farnesol dehydrogenase
	ec:1.1.1.217	benzyl-2-methyl-hydroxybutyrate dehydrogenase

(continued)

ALCOHOL

ec:1.1.1.222
 5 ec:1.1.1.223
 ec:1.1.1.226
 ec:1.1.1.229
 ec:1.1.1.237
 10 ec:1.1.1.244
 ec:1.1.1.245
 ec:1.1.1.250
 ec:1.1.1.251
 ec:1.1.1.255
 15 ec:1.1.1.256
 ec:1.1.1.257
 ec:1.1.1.258
 ec:1.1.1.259
 ec:1.1.1.261
 20 ec:1.1.1.265
 ec:1.1.1.283
 ec:1.1.1.286
 ec:1.1.1.287

ALDEHYDE

ec:1.2.1.2
 ec:1.2.1.3
 ec:1.2.1.4
 30 ec:1.2.1.5
 ec:1.2.1.7
 ec:1.2.1.8
 ec:1.2.1.9
 35 ec:1.2.1.10
 ec:1.2.1.11
 ec:1.2.1.12
 ec:1.2.1.13
 ec:1.2.1.15
 40 ec:1.2.1.16
 ec:1.2.1.17
 ec:1.2.1.18
 ec:1.2.1.19
 45 ec:1.2.1.20
 ec:1.2.1.21
 ec:1.2.1.22
 ec:1.2.1.23
 ec:1.2.1.24
 50 ec:1.2.1.25
 ec:1.2.1.26
 ec:1.2.1.27
 ec:1.2.1.28
 55 ec:1.2.1.29
 ec:1.2.1.30
 ec:1.2.1.31
 ec:1.2.1.32

DEHYDROGENASES

(R)-4-hydroxyphenyllactate dehydrogenase
 isopiperitenol dehydrogenase
 4-hydroxycyclohexanecarboxylate dehydrogenase
 diethyl 2-methyl-3-oxosuccinate reductase
 hydroxyphenylpyruvate reductase
 methanol dehydrogenase
 cyclohexanol dehydrogenase
 D-arabinitol 2-dehydrogenase
 galactitol 1-phosphate 5-dehydrogenase
 mannitol dehydrogenase
 15 fluoren-9-ol dehydrogenase
 4-(hydroxymethyl)benzenesulfonate dehydrogenase
 6-hydroxyhexanoate dehydrogenase
 3-hydroxypimeloyl-CoA dehydrogenase
 glycerol-1-phosphate dehydrogenase [NAD(P)+]
 20 3-methylbutanal reductase
 methylglyoxal reductase (NADPH-dependent)
 isocitrate-homoisocitrate dehydrogenase
 D-arabinitol dehydrogenase (NADP+) butanol dehydrogenase

DEHYDROGENASES

formate dehydrogenase
 aldehyde dehydrogenase (NAD+)
 aldehyde dehydrogenase (NADP+)
 30 aldehyde dehydrogenase [NAD(P)+]
 benzaldehyde dehydrogenase (NADP+)
 betaine-aldehyde dehydrogenase
 glyceraldehyde-3-phosphate dehydrogenase (NADP+)
 35 acetaldehyde dehydrogenase (acetylating)
 aspartate-semialdehyde dehydrogenase
 glyceraldehyde-3-phosphate dehydrogenase (phosphorylating)
 glyceraldehyde-3-phosphate dehydrogenase (NADP+) (phosphorylating)
 malonate-semialdehyde dehydrogenase
 40 succinate-semialdehyde dehydrogenase [NAD(P)+]
 glyoxylate dehydrogenase (acylating)
 malonate-semialdehyde dehydrogenase (acetylating)
 aminobutyraldehyde dehydrogenase
 45 glutarate-semialdehyde dehydrogenase
 glycolaldehyde dehydrogenase
 lactaldehyde dehydrogenase
 2-oxoaldehyde dehydrogenase (NAD+)
 succinate-semialdehyde dehydrogenase
 50 2-oxoisovalerate dehydrogenase (acylating)
 2,5-dioxovalerate dehydrogenase
 methylmalonate-semialdehyde dehydrogenase (acylating)
 benzaldehyde dehydrogenase (NAD+)
 aryl-aldehyde dehydrogenase
 55 aryl-aldehyde dehydrogenase (NADP+)
 L-aminoadipate-semialdehyde dehydrogenase
 aminomuconate-semialdehyde dehydrogenase

(continued)

ALDEHYDE**DEHYDROGENASES**

	ec:1.2.1.36	retinal dehydrogenase
5	ec:1.2.1.39	phenylacetaldehyde dehydrogenase
	ec:1.2.1.41	glutamate-5-semialdehyde dehydrogenase
	ec:1.2.1.42	hexadecanal dehydrogenase (acylating)
	ec:1.2.1.43	formate dehydrogenase (NADP+)
10	ec:1.2.1.45	4-carboxy-2-hydroxymuconate-6-semialdehyde dehydrogenase
	ec:1.2.1.46	formaldehyde dehydrogenase
	ec:1.2.1.47	4-trimethylammonibutyraldehyde dehydrogenase
	ec:1.2.1.48	long-chain-aldehyde dehydrogenase
	ec:1.2.1.49	2-oxoaldehyde dehydrogenase (NADP+)
15	ec:1.2.1.51	pyruvate dehydrogenase (NADP+)
	ec:1.2.1.52	oxoglutarate dehydrogenase (NADP+)
	ec:1.2.1.53	4-hydroxyphenylacetaldehyde dehydrogenase
	ec:1.2.1.57	butanal dehydrogenase
	ec:1.2.1.58	phenylglyoxylate dehydrogenase (acylating)
20	ec:1.2.1.59	glyceraldehyde-3-phosphate dehydrogenase (NAD(P)+) (phosphorylating)
	ec:1.2.1.62	4-formylbenzenesulfonate dehydrogenase
	ec:1.2.1.63	6-oxohexanoate dehydrogenase
	ec:1.2.1.64	4-hydroxybenzaldehyde dehydrogenase
25	ec:1.2.1.65	salicylaldehyde dehydrogenase
	ec:1.2.1.66	mycothiol-dependent formaldehyde dehydrogenase
	ec:1.2.1.67	vanillin dehydrogenase
	ec:1.2.1.68	coniferyl-aldehyde dehydrogenase
	ec:1.2.1.69	fluoroacetaldehyde dehydrogenase
30	ec:1.2.1.71	succinylglutamate-semialdehyde dehydrogenase

[0058] Therefore, in addition to any of the various modifications exemplified previously for establishing 4-HB biosynthesis in a selected host, the BDO producing microbial organisms can include any of the previous combinations and permutations of 4-HB pathway metabolic modifications as well as any combination of expression for CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or an alcohol dehydrogenase to generate biosynthetic pathways for BDO. Therefore, the BDO producers of the disclosure can have exogenous expression of, for example, one, two, three, four, five, six, seven, eight, nine or all 10 enzymes corresponding to any of the six 4-HB pathway and/or any of the 4 BDO pathway enzymes.

[0059] Design and construction of the genetically modified microbial organisms is carried out using methods well known in the art to achieve sufficient amounts of expression to produce BDO. In particular, the non-naturally occurring microbial organisms of the disclosure can achieve biosynthesis of BDO resulting in intracellular concentrations between about 0.1-25 mM or more. Generally, the intracellular concentration of BDO is between about 3-20mM, particularly between about 5-15 mM and more particularly between about 8-12 mM, including about 10 mM or more. Intracellular concentrations between and above each of these exemplary ranges also can be achieved from the non-naturally occurring microbial organisms of the disclosure. As with the 4-HB producers, the BDO producers also can be sustained, cultured or fermented under anaerobic conditions.

[0060] The disclosure further provides a method for the production of 4-HB. The method includes culturing a non-naturally occurring microbial organism having a 4-hydroxybutanoic acid (4-HB) biosynthetic pathway comprising at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, glutamate:succinic semialdehyde transaminase or glutamate decarboxylase under substantially anaerobic conditions for a sufficient period of time to produce monomeric 4-hydroxybutanoic acid (4-HB). The method can additionally include chemical conversion of 4-HB to GBL and to BDO or THF, for example.

[0061] It is understood that, in methods of the disclosure, any of the one or more exogenous nucleic acids can be combined in a non-naturally occurring microbial organism of the disclosure so long as the desired product is produced, for example, 4-HB, BDO, THF or GBL. For example, a non-naturally occurring microbial organism having a 4-HB biosynthetic pathway can comprise at least two exogenous nucleic acids encoding desired enzymes, such as the combination

of 4-hydroxybutanoate dehydrogenase and CoA-independent succinic semialdehyde dehydrogenase; 4-hydroxybutanoate dehydrogenase and CoA-dependent succinic semialdehyde dehydrogenase; CoA-dependent succinic semialdehyde dehydrogenase and succinyl-CoA synthetase; succinyl-CoA synthetase and glutamate decarboxylase, and the like. Thus, it is understood that any combination of two or more enzymes of a biosynthetic pathway can be included in a non-naturally occurring microbial organism of the disclosure. Similarly, it is understood that any combination of three or more enzymes of a biosynthetic pathway can be included in a non-naturally occurring microbial organism of the disclosure, for example, 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase and succinyl-CoA synthetase; 4-hydroxybutanoate dehydrogenase, CoA-dependent succinic semialdehyde dehydrogenase and glutamate:succinic semialdehyde transaminase, and so forth, as desired, so long as the combination of enzymes of the desired biosynthetic pathway results in production of the corresponding desired product.

[0062] Any of the non-naturally occurring microbial organisms described previously can be cultured to produce the biosynthetic products of the disclosure. For example, the 4-HB producers can be cultured for the biosynthetic production of 4-HB. The 4-HB can be isolated or be treated as described below to generate GBL, THF and/or BDO. Similarly, the BDO producers can be cultured for the biosynthetic production of BDO. The BDO can be isolated or subjected to further treatments for the chemical synthesis of BDO family compounds such as those downstream compounds exemplified in Figure 3.

[0063] In some embodiments, culture conditions include anaerobic or substantially anaerobic growth or maintenance conditions. Exemplary anaerobic conditions have been described previously and are well known in the art. Exemplary anaerobic conditions for fermentation processes are described below in the Examples. Any of these conditions can be employed with the non-naturally occurring microbial organisms as well as other anaerobic conditions well known in the art. Under such anaerobic conditions, the 4-HB and BDO producers can synthesize monomeric 4-HB and BDO, respectively, at intracellular concentrations of 5-10 mM or more as well as all other concentrations exemplified previously.

[0064] A number of downstream compounds also can be generated for the 4-HB and BDO producing non-naturally occurring microbial organisms of the disclosure. With respect to the 4-HB producing microbial organisms of the disclosure, monomeric 4-HB and GBL exist in equilibrium in the culture medium. The conversion of 4-HB to GBL can be efficiently accomplished by, for example, culturing the microbial organisms in acid pH medium. A pH less than or equal to 7.5, in particular at or below pH 5.5, spontaneously converts 4-HB to GBL as illustrated in Figure 1.

[0065] The resultant GBL can be separated from 4-HB and other components in the culture using a variety of methods well known in the art. Such separation methods include, for example, the extraction procedures exemplified in the Examples as well as methods which include continuous liquid-liquid extraction, pervaporation, membrane filtration, membrane separation, reverse osmosis, electrodialysis, distillation, crystallization, centrifugation, extractive filtration, ion exchange chromatography, size exclusion chromatography, adsorption chromatography, and ultrafiltration. All of the above methods are well known in the art. Separated GBL can be further purified by, for example, distillation.

[0066] Another downstream compound that can be produced from the 4-HB producing non-naturally occurring microbial organisms of the disclosure includes, for example, BDO. This compound can be synthesized by, for example, chemical hydrogenation of GBL. Chemical hydrogenation reactions are well known in the art. One exemplary procedure includes the chemical reduction of 4-HB and/or GBL or a mixture of these two components deriving from the culture using a heterogeneous or homogeneous hydrogenation catalyst together with hydrogen, or a hydride-based reducing agent used stoichiometrically or catalytically, to produce 1,4-butanediol.

[0067] Other procedures well known in the art are equally applicable for the above chemical reaction and include, for example, WO No. 82/03854 (Bradley, et al.), which describes the hydrogenolysis of gamma-butyrolactone in the vapor phase over a copper oxide and zinc oxide catalyst. British Pat. No. 1,230,276, which describes the hydrogenation of gamma-butyrolactone using a copper oxide-chromium oxide catalyst. The hydrogenation is carried out in the liquid phase. Batch reactions also are exemplified having high total reactor pressures. Reactant and product partial pressures in the reactors are well above the respective dew points. British Pat. No. 1,314,126, which describes the hydrogenation of gamma-butyrolactone in the liquid phase over a nickel-cobalt-thorium oxide catalyst. Batch reactions are exemplified as having high total pressures and component partial pressures well above respective component dew points. British Pat. No. 1,344,557, which describes the hydrogenation of gamma-butyrolactone in the liquid phase over a copper oxide-chromium oxide catalyst. A vapor phase or vapor-containing mixed phase is indicated as suitable in some instances. A continuous flow tubular reactor is exemplified using high total reactor pressures. British Pat. No. 1,512,751, which describes the hydrogenation of gamma-butyrolactone to 1,4-butanediol in the liquid phase over a copper oxide-chromium oxide catalyst. Batch reactions are exemplified with high total reactor pressures and, where determinable, reactant and product partial pressures well above the respective dew points. U.S. Pat. No. 4,301,077, which describes the hydrogenation to 1,4-butanediol of gamma-butyrolactone over a Ru-Ni-Co-Zn catalyst. The reaction can be conducted in the liquid or gas phase or in a mixed liquid-gas phase. Exemplified are continuous flow liquid phase reactions at high total reactor pressures and relatively low reactor productivities. U.S. Pat. No. 4,048,196, which describes the production of 1,4-butanediol by the liquid phase hydrogenation of gamma-butyrolactone over a copper oxide-zinc oxide catalyst. Further exemplified is a continuous flow tubular reactor operating at high total reactor pressures and high reactant and product

partial pressures. And U.S. Patent No. 4,652,685, which describes the hydrogenation of lactones to glycols.

[0068] A further downstream compound that can be produced from the 4-HB producing microbial organisms of the disclosure includes, for example, THF. This compound can be synthesized by, for example, chemical hydrogenation of GBL. One exemplary procedure well known in the art applicable for the conversion of GBL to THF includes, for example, chemical reduction of 4-HB and/or GBL or a mixture of these two components deriving from the culture using a heterogeneous or homogeneous hydrogenation catalyst together with hydrogen, or a hydride-based reducing agent used stoichiometrically or catalytically, to produce tetrahydrofuran. Other procedures well known in the art are equally applicable for the above chemical reaction and include, for example, U.S. Patent No. 6,686,310, which describes high surface area sol-gel route prepared hydrogenation catalysts. Processes for the reduction of gamma butyrolactone to tetrahydrofuran and 1,4-butanediol also are described.

[0069] The culture conditions can include, for example, liquid culture procedures as well as fermentation and other large scale culture procedures. As described further below in the Examples, particularly useful yields of the biosynthetic products of the disclosure can be obtained under anaerobic or substantially anaerobic culture conditions.

[0070] The disclosure further provides a method of manufacturing 4-HB. The method includes fermenting a non-naturally occurring microbial organism having a 4-hydroxybutanoic acid (4-HB) biosynthetic pathway comprising at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, glutamate:succinic semialdehyde transaminase or glutamate decarboxylase under substantially anaerobic conditions for a sufficient period of time to produce monomeric 4-hydroxybutanoic acid (4-HB), the process comprising fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation.

[0071] The culture and chemical hydrogenations described above also can be scaled up and grown continuously for manufacturing of 4-HB, GBL, BDO and/or THF. Exemplary growth procedures include, for example, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation. All of these processes are well known in the art. Employing the 4-HB producers allows for simultaneous 4-HB biosynthesis and chemical conversion to GBL, BDO and/or THF by employing the above hydrogenation procedures simultaneous with continuous cultures methods such as fermentation. Other hydrogenation procedures also are well known in the art and can be equally applied to the methods of the invention.

[0072] Fermentation procedures are particularly useful for the biosynthetic production of commercial quantities of 4-HB and/or BDO. Generally, and as with non-continuous culture procedures, the continuous and/or near-continuous production of 4-HB or BDO will include culturing a non-naturally occurring 4-HB or BDO producing organism of the invention in sufficient nutrients and medium to sustain and/or nearly sustain growth in an exponential phase. Continuous culture under such conditions can include, for example, 1 day, 2, 3, 4, 5, 6 or 7 days or more. Additionally, continuous culture can include 1 week, 2, 3, 4 or 5 or more weeks and up to several months. Alternatively, microorganisms of the disclosure can be cultured for hours, if suitable for a particular application. It is to be understood that the continuous and/or near-continuous culture conditions also can include all time intervals in between these exemplary periods.

[0073] Fermentation procedures are well known in the art. Briefly, fermentation for the biosynthetic production of 4-HB, BDO or other 4-HB derived products of the disclosure can be utilized in, for example, fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation. Examples of batch and continuous fermentation procedures well known in the art are exemplified further below in the Examples.

[0074] In addition, to the above fermentation procedures using the 4-HB or BDO producers of the disclosure for continuous production of substantial quantities of monomeric 4-HB and BDO, respectively, the 4-HB producers also can be, for example, simultaneously subjected to chemical synthesis procedures as described previously for the chemical conversion of monomeric 4-HB to, for example, GBL, BDO and/or THF. The BDO producers can similarly be, for example, simultaneously subjected to chemical synthesis procedures as described previously for the chemical conversion of BDO to, for example, THF, GBL, pyrrolidones and/or other BDO family compounds. In addition, the products of the 4-HB and BDO producers can be separated from the fermentation culture and sequentially subjected to chemical conversion, as disclosed herein.

[0075] Briefly, hydrogenation of GBL in the fermentation broth can be performed as described by Frost et al., *Biotechnology Progress* 18: 201-211 (2002). Another procedure for hydrogenation during fermentation include, for example, the methods described in, for example, U.S. Patent No. 5,478,952. This method is further exemplified in the Examples below.

[0076] Therefore, the invention additionally provides a method of manufacturing γ -butyrolactone (GBL), tetrahydrofuran (THF) or 1,4-butanediol (BDO). The method includes fermenting a non-naturally occurring microbial organism having 4-hydroxybutanoic acid (4-HB) and 1,4-butanediol (BDO) biosynthetic pathways, the pathways comprise at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent 1,4-butanediol

diol semialdehyde dehydrogenase, CoA-dependent 1,4-butanediol semialdehyde dehydrogenase, 1,4-butanediol alcohol dehydrogenase, either CoA-dependent or independent, under substantially anaerobic conditions for a sufficient period of time to produce 1,4-butanediol (BDO), GBL or THF, the fermenting comprising fed-batch fermentation and batch separation; fed-batch fermentation and continuous separation, or continuous fermentation and continuous separation.

[0077] In addition to the biosynthesis of 4-HB, BDO and other products of the disclosure as described herein, the non-naturally occurring microbial organisms and methods of the disclosure also can be utilized in various combinations with each other and with other microbial organisms and methods well known in the art to achieve product biosynthesis by other routes. For example, one alternative to produce BDO other than use of the 4-HB producers and chemical steps or other than use of the BDO producer directly is through addition of another microbial organism capable of converting 4-HB or a 4-HB product exemplified herein to BDO.

[0078] One such procedure includes, for example, the fermentation of a 4-HB producing microbial organism of the disclosure to produce 4-HB, as described above and below. The 4-HB can then be used as a substrate for a second microbial organism that converts 4-HB to, for example, BDO, GBL and/or THF. The 4-HB can be added directly to another culture of the second organism or the original culture of 4-HB producers can be depleted of these microbial organisms by, for example, cell separation, and then subsequent addition of the second organism to the fermentation broth can be utilized to produce the final product without intermediate purification steps. One exemplary second organism having the capacity to biochemically utilize 4-HB as a substrate for conversion to BDO, for example, is *Clostridium acetobutylicum* (see, for example, Jewell et al., Current Microbiology, 13:215-19 (1986)).

[0079] In other disclosure, the non-naturally occurring microbial organisms and methods of the disclosure can be assembled in a wide variety of subpathways to achieve biosynthesis of, for example, 4-HB and/or BDO as described. In these disclosure, biosynthetic pathways for a desired product can be segregated into different microbial organisms and the different microbial organisms can be co-cultured to produce the final product. In such a biosynthetic scheme, the product of one microbial organism is the substrate for a second microbial organism until the final product is synthesized. For example, the biosynthesis of BDO can be accomplished as described previously by constructing a microbial organism that contains biosynthetic pathways for conversion of a substrate such as endogenous succinate through 4-HB to the final product BDO. Alternatively, BDO also can be biosynthetically produced from microbial organisms through co-culture or co-fermentation using two organisms in the same vessel. A first microbial organism being a 4-HB producer with genes to produce 4-HB from succinic acid, and a second microbial organism being a BDO producer with genes to convert 4-HB to BDO.

[0080] Given the teachings and guidance provided herein, those skilled in the art will understand that a wide variety of combinations and permutations exist for the non-naturally occurring microbial organisms and methods of the disclosure together with other microbial organisms, with the co-culture of other non-naturally occurring microbial organisms having subpathways and with combinations of other chemical and/or biochemical procedures well known in the art to produce 4-HB, BDO, GBL and THF products.

[0081] One computational method for identifying and designing metabolic alterations favoring biosynthesis of a product is the OptKnock computational framework, Burgard et al., Biotechnol Bioeng, 84: 647-57 (2003). OptKnock is a metabolic modeling and simulation program that suggests gene deletion strategies that result in genetically stable microorganisms which overproduce the target product. Specifically, the framework examines the complete metabolic and/or biochemical network of a microorganism in order to suggest genetic manipulations that force the desired biochemical to become an obligatory byproduct of cell growth. By coupling biochemical production with cell growth through strategically placed gene deletions or other functional gene disruption, the growth selection pressures imposed on the engineered strains after long periods of time in a bioreactor lead to improvements in performance as a result of the compulsory growth-coupled biochemical production. Lastly, when gene deletions are constructed there is a negligible possibility of the designed strains reverting to their wild-type states because the genes selected by OptKnock are to be completely removed from the genome. Therefore, this computational methodology can be used to either identify alternative pathways that lead to biosynthesis of 4-HB and/or BDO or used in connection with the non-naturally occurring microbial organisms for further optimization of 4-HB and/or BDO biosynthesis.

[0082] Briefly, OptKnock is a term used herein to refer to a computational method and system for modeling cellular metabolism. The OptKnock program relates to a framework of models and methods that incorporate particular constraints into flux balance analysis (FBA) models. These constraints include, for example, qualitative kinetic information, qualitative regulatory information, and/or DNA microarray experimental data. OptKnock also computes solutions to various metabolic problems by, for example, tightening the flux boundaries derived through flux balance models and subsequently probing the performance limits of metabolic networks in the presence of gene additions or deletions. OptKnock computational framework allows the construction of model formulations that enable an effective query of the performance limits of metabolic networks and provides methods for solving the resulting mixed-integer linear programming problems. The metabolic modeling and simulation methods referred to herein as OptKnock are described in, for example, U.S. Patent Application Serial No. 10/043,440, filed January 10, 2002, and in International Patent No. PCT/US02/00660, filed January

10, 2002.

[0083] Another computational method for identifying and designing metabolic alterations favoring biosynthetic production of a product is metabolic modeling and simulation system termed SimPheny®. This computational method and system is described in, for example, U.S. Patent Application Serial No. 10/173,547, filed June 14, 2002, and in International Patent Application No. PCT/US03/18838, filed June 13, 2003.

[0084] SimPheny® is a computational system that can be used to produce a network model *in silico* and to simulate the flux of mass, energy or charge through the chemical reactions of a biological system to define a solution space that contains any and all possible functionalities of the chemical reactions in the system, thereby determining a range of allowed activities for the biological system. This approach is referred to as constraints-based modeling because the solution space is defined by constraints such as the known stoichiometry of the included reactions as well as reaction thermodynamic and capacity constraints associated with maximum fluxes through reactions. The space defined by these constraints can be interrogated to determine the phenotypic capabilities and behavior of the biological system or of its biochemical components. Analysis methods such as convex analysis, linear programming and the calculation of extreme pathways as described, for example, in Schilling et al., J. Theor. Biol. 203:229-248 (2000); Schilling et al., Biotech. Bioeng. 71:286-306 (2000) and Schilling et al., Biotech. Prog. 15:288-295 (1999), can be used to determine such phenotypic capabilities. As described in the Examples below, this computation methodology was used to identify and analyze the feasible as well as the optimal 4-HB biosynthetic pathways in 4-HB non-producing microbial organisms.

[0085] As described above, one constraints-based method used in the computational programs applicable to the invention is flux balance analysis. Flux balance analysis is based on flux balancing in a steady state condition and can be performed as described in, for example, Varma and Palsson, Biotech. Bioeng. 12:994-998 (1994). Flux balance approaches have been applied to reaction networks to simulate or predict systemic properties of, for example, adipocyte metabolism as described in Fell and Small, J. Biochem. 138:781-786 (1986), acetate secretion from *E. coli* under ATP maximization conditions as described in Majewski and Domach, Biotech. Bioeng. 35:732-738 (1990) or ethanol secretion by yeast as described in Vanrolleghem et al., Biotech. Prog. 12:434-448 (1996). Additionally, this approach can be used to predict or simulate the growth of *E. coli* on a variety of single-carbon sources as well as the metabolism of *H. influenzae* as described in Edwards and Palsson, Proc. Natl. Acad. Sci. USA 97:5528-5533 (2000), Edwards and Palsson, J. Biol. Chem. 274:17410-17416 (1999) and Edwards et al., Nature Biotech. 19:125-130 (2001).

[0086] Once the solution space has been defined, it can be analyzed to determine possible solutions under various conditions. This computational approach is consistent with biological realities because biological systems are flexible and can reach the same result in many different ways. Biological systems are designed through evolutionary mechanisms that have been restricted by fundamental constraints that all living systems must face. Therefore, constraints-based modeling strategy embraces these general realities. Further, the ability to continuously impose further restrictions on a network model via the tightening of constraints results in a reduction in the size of the solution space, thereby enhancing the precision with which physiological performance or phenotype can be predicted.

[0087] Given the teachings and guidance provided herein, those skilled in the art will be able to apply various computational frameworks for metabolic modeling and simulation to design and implement biosynthesis of 4-HB, BDO, GBL, THF and other BDO family compounds in host microbial organisms other than *E. coli* and yeast. Such metabolic modeling and simulation methods include, for example, the computational systems exemplified above as SimPheny® and OptKnock. For illustration of the invention, some methods are described herein with reference to the OptKnock computation framework for modeling and simulation. Those skilled in the art will know how to apply the identification, design and implementation of the metabolic alterations using OptKnock to any of such other metabolic modeling and simulation computational frameworks and methods well known in the art.

[0088] The ability of a cell or organism to biosynthetically produce a biochemical product can be illustrated in the context of the biochemical production limits of a typical metabolic network calculated using an *in silico* model. These limits are obtained by fixing the uptake rate(s) of the limiting substrate(s) to their experimentally measured value(s) and calculating the maximum and minimum rates of biochemical production at each attainable level of growth. The production of a desired biochemical generally is in direct competition with biomass formation for intracellular resources. Under these circumstances, enhanced rates of biochemical production will necessarily result in sub-maximal growth rates. The knock-outs suggested by the above metabolic modeling and simulation programs such as OptKnock are designed to restrict the allowable solution boundaries forcing a change in metabolic behavior from the wild-type strain. Although the actual solution boundaries for a given strain will expand or contract as the substrate uptake rate(s) increase or decrease, each experimental point will lie within its calculated solution boundary. Plots such as these enable accurate predictions of how close the designed strains are to their performance limits which also indicates how much room is available for improvement.

[0089] The OptKnock mathematical framework is exemplified herein for pinpointing gene deletions leading to product biosynthesis and, particularly, growth-coupled product biosynthesis. The procedure builds upon constraint-based metabolic modeling which narrows the range of possible phenotypes that a cellular system can display through the successive imposition of governing physico-chemical constraints, Price et al., Nat. Rev. Microbiol., 2: 886-97 (2004). As described

above, constraint-based models and simulations are well known in the art and generally invoke the optimization of a particular cellular objective, subject to network stoichiometry, to suggest a likely flux distribution.

[0090] Briefly, the maximization of a cellular objective quantified as an aggregate reaction flux for a steady state metabolic network comprising a set $N = \{1, \dots, N\}$ of metabolites and a set $M = \{1, \dots, M\}$ of metabolic reactions is expressed mathematically as follows:

$$\begin{aligned}
 &\text{maximize} && v_{\text{cellular objective}} \\
 &\text{subject to} && \sum_{j=1}^M S_{ij} v_j = 0, && \forall i \in N \\
 & && v_{\text{substrate}} = v_{\text{substrate_uptake}} \text{ mmol/gDW}\cdot\text{hr} && \forall i \in \{\text{limiting substrate(s)}\} \\
 & && v_{\text{atp}} \geq v_{\text{atp_main}} \text{ mmol/gDW}\cdot\text{hr} \\
 & && v_j \geq 0, && \forall j \in \{\text{irrev. reactions}\}
 \end{aligned}$$

where S_{ij} is the stoichiometric coefficient of metabolite i in reaction j , v_j is the flux of reaction j , $v_{\text{substrate_uptake}}$ represents the assumed or measured uptake rate(s) of the limiting substrate(s), and $v_{\text{atp_main}}$ is the non-growth associated ATP maintenance requirement. The vector v includes both internal and external fluxes. In this study, the cellular objective is often assumed to be a drain of biosynthetic precursors in the ratios required for biomass formation, Neidhardt, F.C. et al., *Escherichia coli* and *Salmonella*: Cellular and Molecular Biology, 2nd ed. 1996, Washington, D.C.: ASM Press. 2 v. (xx, 2822, lxxvi). The fluxes are generally reported per 1 gDW·hr (gram of dry weight times hour) such that biomass formation is expressed as g biomass produced/gDW·hr or 1/hr.

[0091] The modeling of gene deletions, and thus reaction elimination, first employs the incorporation of binary variables into the constraint-based approach framework, Burgard et al., *Biotechnol. Bioeng.*, 74: 364-375 (2001), Burgard et al., *Biotechnol. Prog.*, 17: 791-797 (2001). These binary variables,

$$y_j = \begin{cases} 1, & \text{if reaction flux } v_j \text{ is active} \\ 0, & \text{if reaction flux } v_j \text{ is not active} \end{cases}, \forall j \in M$$

assume a value of 1 if reaction j is active and a value of 0 if it is inactive. The following constraint,

$$v_j^{\min} \cdot y_j \leq v_j \leq v_j^{\max} \cdot y_j, \quad \forall j \in M$$

ensures that reaction flux v_j is set to zero only if variable y_j is equal to zero. Alternatively, when y_j is equal to one, v_j is

free to assume any value between a lower $v_j^{\min}$ and an upper $v_j^{\max}$ bound. Here, $v_j^{\min}$ and $v_j^{\max}$ are identified by minimizing and maximizing, respectively, every reaction flux subject to the network constraints described above, Mahadevan et al., *Metab. Eng.*, 5: 264-76 (2003).

[0092] Optimal gene/reaction knockouts are identified by solving a bilevel optimization problem that chooses the set of active reactions ($y_j = 1$) such that an optimal growth solution for the resulting network overproduces the chemical of interest. Mathematically, this bilevel optimization problem is expressed as the following bilevel mixed-integer optimization problem:

$$\text{maximize } y_j \quad v_{\text{chemical}} \quad (\text{OptKnock})$$

(continued)

$$\begin{array}{ll}
 \text{subject to} & \text{maximize} \quad v_{biomass} \\
 v_j & \\
 \text{subject to} & \sum_{j=1}^M S_{ij} v_j = 0, \quad \forall i \in N \\
 & v_{substrate} = v_{substrate_uptake} \quad \forall i \in \{\text{limiting substrate(s)}\} \\
 & v_{atp} \geq v_{atp_min} \\
 & \\
 & v_{biomass} \geq v_{biomass}^{target} \\
 & v_j^{min} \cdot y_j \leq v_j \leq v_j^{max} \cdot y_j, \quad \forall j \in M \\
 & \sum_{j \in M^{forward}} (1 - y_j) = K \\
 y_j \in \{0, 1\}, & \forall j \in M
 \end{array}$$

where $v_{chemical}$ is the production of the desired target product, for example succinate or other biochemical product, and K is the number of allowable knockouts. Note that setting K equal to zero returns the maximum biomass solution of the complete network, while setting K equal to one identifies the single gene/reaction knockout ($y_j = 0$) such that the resulting network involves the maximum overproduction given its maximum biomass yield. The final constraint ensures that the resulting network meets a minimum biomass yield. Burgard et al., *Biotechnol. Bioeng.*, 84: 647-57 (2003), provide a more detailed description of the model formulation and solution procedure. Problems containing hundreds of binary variables can be solved in the order of minutes to hours using CPLEX 8.0, *GAMS: The Solver Manuals*. 2003: GAMS Development Corporation, accessed via the GAMS, Brooke et al., GAMS Development Corporation (1998), modeling environment on an IBM RS6000-270 workstation. The OptKnock framework has already been able to identify promising gene deletion strategies for biochemical overproduction, Burgard et al., *Biotechnol. Bioeng.*, 84: 647-57 (2003), Pharkya et al., *Biotechnol. Bioeng.*, 84: 887-899 (2003), and establishes a systematic framework that will naturally encompass future improvements in metabolic and regulatory modeling frameworks.

[0093] Any solution of the above described bilevel OptKnock problem will provide one set of metabolic reactions to disrupt. Elimination of each reaction within the set or metabolic modification can result in 4-HB or BDO as an obligatory product during the growth phase of the organism. Because the reactions are known, a solution to the bilevel OptKnock problem also will provide the associated gene or genes encoding one or more enzymes that catalyze each reaction within the set of reactions. Identification of a set of reactions and their corresponding genes encoding the enzymes participating in each reaction is generally an automated process, accomplished through correlation of the reactions with a reaction database having a relationship between enzymes and encoding genes.

[0094] Once identified, the set of reactions that are to be disrupted in order to achieve 4-HB or BDO production are implemented in the target cell or organism by functional disruption of at least one gene encoding each metabolic reaction within the set. One particularly useful means to achieve functional disruption of the reaction set is by deletion of each encoding gene. However, in some instances, it can be beneficial to disrupt the reaction by other genetic aberrations including, for example, mutation, deletion of regulatory regions such as promoters or *cis* binding sites for regulatory factors, or by truncation of the coding sequence at any of a number of locations. These latter aberrations, resulting in less than total deletion of the gene set can be useful, for example, when rapid assessments of the succinate coupling are desired or when genetic reversion is less likely to occur.

[0095] To identify additional productive solutions to the above described bilevel OptKnock problem which lead to further sets of reactions to disrupt or metabolic modifications that can result in the biosynthesis, including growth-coupled biosynthesis of 4-HB or other biochemical product, an optimization method, termed integer cuts, can be implemented. This method proceeds by iteratively solving the OptKnock problem exemplified above with the incorporation of an additional constraint referred to as an integer cut at each iteration. Integer cut constraints effectively prevent the solution procedure from choosing the exact same set of reactions identified in any previous iteration that obligatory couples product biosynthesis to growth. For example, if a previously identified growth-coupled metabolic modification specifies

reactions 1, 2, and 3 for disruption, then the following constraint prevents the same reactions from being simultaneously considered in subsequent solutions: $y_1 + y_2 + y_3 \geq 1$. The integer cut method is well known in the art and can be found described in, for example, reference, Burgard et al., *Biotechnol. Prog.*, 17: 791-797 (2001). As with all methods described herein with reference to their use in combination with the OptKnock computational framework for metabolic modeling and simulation, the integer cut method of reducing redundancy in iterative computational analysis also can be applied

with other computational frameworks well known in the art including, for example, SimPheny®. [0096] Constraints of the above form preclude identification of larger reaction sets that include previously identified sets. For example, employing the integer cut optimization method above in a further iteration would preclude identifying a quadruple reaction set that specified reactions 1, 2, and 3 for disruption since these reactions had been previously identified. To ensure identification of all possible reaction sets leading to biosynthetic production of a product, a modification of the integer cut method can be employed.

[0097] Briefly, the modified integer cut procedure begins with iteration 'zero' which calculates the maximum production of the desired biochemical at optimal growth for a wild-type network. This calculation corresponds to an OptKnock solution with K equaling 0.

[0098] Next, single knockouts are considered and the two parameter sets, $objstore_{iter}$ and $ystore_{iter,j}$, are introduced to store the objective function ($v_{chemical}$) and reaction on-off information (y_j), respectively, at each iteration, $iter$. The following constraints are then successively added to the OptKnock formulation at each iteration.

$$v_{chemical} \geq objstore_{iter} + \varepsilon - M \cdot \sum_{j \in ystore_{iter,j} = 0} y_j$$

[0099] In the above equation, ε and M are a small and a large numbers, respectively. In general, ε can be set at about 0.01 and M can be set at about 1000. However, numbers smaller and/or larger than these numbers also can be used. M ensures that the constraint can be binding only for previously identified knockout strategies, while ε ensures that adding knockouts to a previously identified strategy must lead to an increase of at least ε in biochemical production at optimal growth. The approach moves onto double deletions whenever a single deletion strategy fails to improve upon the wild-type strain. Triple deletions are then considered when no double deletion strategy improves upon the wild-type strain, and so on. The end result is a ranked list, represented as desired biochemical production at optimal growth, of distinct deletion strategies that differ from each other by at least one knockout. This optimization procedure as well as the identification of a wide variety of reaction sets that, when disrupted, lead to the biosynthesis, including growth-coupled production, of a biochemical product. Given the teachings and guidance provided herein, those skilled in the art will understand that the methods and metabolic engineering designs exemplified herein are equally applicable to identify new biosynthetic pathways and/or to the obligatory coupling of cell or microorganism growth to any biochemical product.

[0100] The methods exemplified above and further illustrated in the Examples below enable the construction of cells and organisms that biosynthetically produce, including obligatory couple production of a target biochemical product to growth of the cell or organism engineered to harbor the identified genetic alterations. In this regard, metabolic alterations have been identified that result in the biosynthesis of 4-HB and 1,4-butanediol. Microorganism strains constructed with the identified metabolic alterations produce elevated levels of 4-HB or BDO compared to unmodified microbial organisms. These strains can be beneficially used for the commercial production of 4-HB, BDO, THF and GBL, for example, in continuous fermentation process without being subjected to the negative selective pressures.

[0101] Therefore, the computational methods described herein enable the identification and implementation of metabolic modifications that are identified by an *in silico* method selected from OptKnock or SimPheny. The set of metabolic modifications can include, for example, addition of one or more biosynthetic pathway enzymes and/or functional disruption of one or more metabolic reactions including, for example, disruption by gene deletion.

[0102] Application of the OptKnock method to identify pathways suitable for growth-coupled production of 1,4-butanediol (BDO) is described in Examples V-VII. As discussed above, the OptKnock methodology was developed on the premise that mutant microbial networks can be evolved towards their computationally predicted maximum-growth phenotypes when subjected to long periods of growth selection. In other words, the approach leverages an organism's ability to self-optimize under selective pressures. The OptKnock framework allows for the exhaustive enumeration of gene deletion combinations that force a coupling between biochemical production and cell growth based on network stoichiometry. The identification of optimal gene/reaction knockouts requires the solution of a bilevel optimization problem that chooses the set of active reactions such that an optimal growth solution for the resulting network overproduces the biochemical of interest (Burgard et al. *Biotechnol. Bioeng.* 84:647-657 (2003)).

[0103] Growth-coupled biochemical production can be visualized in the context of the biochemical production limits of a typical metabolic network. These limits are obtained from an *in silico* metabolic model by fixing the uptake rate(s) of the limiting substrate(s) to their experimentally measured value(s) and calculating the maximum and minimum rates of biochemical production at each attainable level of growth. The production envelopes essentially bracket what is

possible by encompassing all potential biological phenotypes available to a given strain. Although exceptions exist, typically the production of a desired biochemical is in direct competition with biomass formation for intracellular resources (see Figure 5, gray region). Thus increased biochemical yields will necessarily result in sub-maximal growth. Furthermore, the application of growth selective pressures may drive the performance of a non-growth-coupled production strain towards a low producing phenotype (point A, Figure 5), regardless of its initial starting point. The knockouts suggested by OptKnock are designed to restrict the allowable solution boundary, forcing a change in metabolic behavior as depicted in Figure 5 (cyan region). Evolutionary engineering approaches can thus be applied to drive the performance of an OptKnock designed strain to the rightmost boundary. This will result in a high producing strain that should be inherently stable to further growth selective pressures (point B, Figure 5).

[0104] An *in silico* stoichiometric model of *E. coli* metabolism was employed to identify essential genes for metabolic pathways as exemplified previously and described in, for example, U.S. patent publications US 2002/0012939, US 2003/0224363, US 2004/0029149, US 2004/0072723, US 2003/0059792, US 2002/0168654 and US 2004/0009466, and in U.S. Patent No. 7,127,379. As disclosed herein, the OptKnock mathematical framework was applied to pinpoint gene deletions leading to the growth-coupled production of BDO (see Examples V-VII). The strategies were initially determined by employing a reduced model of the *E. coli* metabolic network. This "small" model captures the key metabolic functionalities in the network, thus eliminating the redundancy associated with the genome-scale metabolic networks. Care was taken to ensure that the model was not reduced to the point where potentially active pathways possessing viable targets were neglected. Overall, the reduced model contained 262 reactions and its implementation reduced OptKnock CPU times approximately ten-fold when compared to the application of OptKnock to the genome-scale *E. coli* model (Reed et al., *Genome Biol.* 4:R54 (2003)).

[0105] Further, the solution of the bilevel OptKnock problem provides only one set of deletions. To enumerate all meaningful solutions, that is, all sets of knockouts leading to growth-coupled production formation, an optimization technique, termed integer cuts, can be implemented. This entails iteratively solving the OptKnock problem with the incorporation of an additional constraint referred to as an integer cut at each iteration, as discussed above.

[0106] For biochemical pathways to 1,4-butanediol, the conversion of glucose to BDO in *E. coli* is expected to derive from a total of three intracellular reduction steps from succinate semialdehyde (see Figure 6). Succinate semialdehyde is natively produced by *E. coli* through the TCA cycle intermediate, alpha-ketoglutarate, via the action of two enzymes, glutamate:succinic semialdehyde transaminase and glutamate decarboxylase. An alternative pathway, used by the obligate anaerobe *Clostridium kluyveri* to degrade succinate, activates succinate to succinyl-CoA, and then converts succinyl-CoA to succinic semialdehyde using a CoA-dependant succinic semialdehyde dehydrogenase (Sohling and Gottschalk, *Eur. J. Biochem.* 212:121-127 (1993)). The conversion of succinate semialdehyde to BDO first requires the activity of 4-hydroxybutanoate (4-HB) dehydrogenase, an enzyme which is not native to *E. coli* or yeast but is found in various bacteria such as *C. kluyveri* and *Ralstonia eutropha* (Lutke-Eversloh and Steinbuchel, *FEMS Microbiol. Lett.* 181:63-71 (1999); Sohling and Gottschalk, *J. Bacteriol.* 178:871-880 (1996); Valentin et al., *Eur. J. Biochem.* 227:43-60 (1995); Wolff and Kenealy, *Protein Expr. Purif.* 6:206-212 (1995)). Precedent for the 4-HB to BDO conversion has been demonstrated in the strict anaerobe, *Clostridium acetobutylicum*, which when fed 4-HB, was shown to quantitatively produce BDO (Jewell et al., *Current Microbiology* 13:215-219 (1986)). The required biotransformations from 4-HB to BDO are assumed to be similar to those of the butyric acid to butanol conversion common to *Clostridia* species which proceeds via a CoA-derivative (Girbal et al., *FEMS -Microbiology Reviews* 17:287-297 (1995)).

[0107] Further disclosed is a non-naturally occurring microorganism comprising one or more gene disruptions, the one or more gene disruptions occurring in genes encoding an enzyme obligatory to coupling 1,4-butanediol production to growth of the microorganism when the gene disruption reduces an activity of the enzyme, whereby the one or more gene disruptions confers stable growth-coupled production of 1,4-butanediol onto the non-naturally occurring microorganism. The one or more gene disruptions can be, for example, those disclosed in Table 6 or 7. The one or more gene disruptions can comprise a deletion of the one or more genes, such as those genes disclosed in Table 6 or 7.

[0108] As disclosed herein, the non-naturally occurring microorganism can be a bacterium, yeast or fungus. For example, the non-naturally occurring microorganism can be a bacterium such as *Escherichia coli*, *Klebsiella oxytoca*, *Anaerobiospirillum succiniciproducens*, *Actinobacillus succinogenes*, *Mannheimia succiniciproducens*, *Rhizobium etli*, *Bacillus subtilis*, *Corynebacterium glutamicum*, *Gluconobacter oxydans*, *Zymomonas mobilis*, *Lactococcus lactis*, *Lactobacillus plantarum*, *Streptomyces coelicolor*, *Clostridium acetobutylicum*, *Pseudomonas fluorescens*, and *Pseudomonas putida*. The non-naturally occurring microorganism can also be a yeast such as *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces marxianus*, *Aspergillus terreus*, *Aspergillus niger*, and *Pichia pastoris*.

[0109] A non-naturally occurring microorganism of the disclosure can comprise a set of metabolic modifications obligatory coupling 1,4-butanediol production to growth of the microorganism, the set of metabolic modifications comprising disruption of one or more genes selected from the set of genes comprising *adhE*, *ldhA*, *pflAB*; *adhE*, *ldhA*, *pflAB*, *mdh*; *adhE*, *ldhA*, *pflAB*, *mdh*, *mgo*; *adhE*, *ldhA*, *pflAB*, *mdh*, *aspA*; *adhE*, *mdh*, *ldhA*, *pflAB*, *sfcA*; *adhE*, *mdh*, *ldhA*, *pflAB*, *maeB*; *adhE*, *mdh*, *ldhA*, *pflAB*, *sfcA*, *maeB*; *adhE*, *ldhA*, *pflAB*, *mdh*, *pntAB*; *adhE*, *ldhA*, *pflAB*, *mdh*, *gdhA*; *adhE*, *ldhA*,

pflAB, *mdh*, *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE*, *ycdW*; and *adhE*, *ldhA*, *pflAB*, *mdh*, *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE*, *ycdW*, *prpC*, *gsk*, or an ortholog thereof, wherein the microorganism exhibits stable growth-coupled production of 1,4-butanediol. The disclosure provides disclosure a non-naturally occurring microorganism comprising a set of metabolic modifications obligatory to coupling 1,4-butanediol production to growth of the microorganism, the set of metabolic modifications comprising disruption of one or more genes, or an ortholog thereof, wherein the set of metabolic modifications comprises disruption of *adhE* and *ldhA*, wherein the microorganism exhibits stable growth-coupled production of 1,4-butanediol. The set of metabolic modifications can further comprise disruption of *mdh*. The set of metabolic modifications can further comprise disruption of one or more genes selected from the set of genes comprising *mgo*, *aspA*, *sfcA*, *maeB*, *pntAB*, and *gdhA* and can include, for example, disruption of *sfcA* and *maeB*. The set of metabolic modifications can further comprise disruption of one or more genes selected from the set of genes comprising *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE*, *ycdW*, *prpC*, and *gsk*, including disruption of all of *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE* and *ycdW*, and can further comprise disruption of *prpC* and *gsk*. Any of the above-described set of metabolic modifications can further comprise disruption of *pflAB*. In a particular disclosure the set of metabolic modifications comprise disruption of one or more genes selected from the set of genes comprising *adhE*, *ldhA*, *pflAB*, *mdh*, and *aspA*, including up to all of genes *adhE*, *ldhA*, *pflAB*, *mdh*, and *aspA*.

[0110] A non-naturally occurring microorganism of the disclosure can further comprise a 1,4-butanediol (BDO) biosynthetic pathway comprising at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or alcohol dehydrogenase, wherein the exogenous nucleic acid is expressed in sufficient amounts to produce 1,4-butanediol (BDO), as disclosed herein.

[0111] The disclosure additionally provides a method of producing a non-naturally occurring microorganism having stable growth-coupled production of 1,4-butanediol by identifying *in silico* a set of metabolic modifications requiring 1,4-butanediol production during exponential growth under a defined set of conditions, and genetically modifying a microorganism to contain the set of metabolic modifications requiring 1,4-butanediol production. Such methods can additionally include the addition of exogenous genes expressing desired enzyme activities to a microorganism. Such a set of metabolic modifications can be identified by an *in silico* method selected from OptKnock or SimPheny (see above and Examples V-VII).

[0112] Furthermore, the invention provides a method of producing 1,4-butanediol coupled to the growth of a microorganism. The method can include the steps of culturing under exponential growth phase in a sufficient amount of nutrients and media a non-naturally occurring microorganism comprising a set of metabolic modifications obligatorily coupling 1,4-butanediol production to growth of the microorganism, wherein the microorganism exhibits stable growth-coupled production of 1,4-butanediol, and isolating 1,4-butanediol produced from the non-naturally occurring microorganism. The set of metabolic modifications comprising disruption of one or more genes can be selected from the set of genes comprising *adhE*, *ldhA*, *pflAB*; *adhE*, *ldhA*, *pflAB*, *mdh*; *adhE*, *ldhA*, *pflAB*, *mdh*, *mgo*; *adhE*, *ldhA*, *pflAB*, *mdh*, *aspA*; *adhE*, *mdh*, *ldhA*, *pflAB*, *sfcA*; *adhE*, *mdh*, *ldhA*, *pflAB*, *maeB*; *adhE*, *mdh*, *ldhA*, *pflAB*, *sfcA*, *maeB*; *adhE*, *ldhA*, *pflAB*, *mdh*, *pntAB*; *adhE*, *ldhA*, *pflAB*, *mdh*, *gdhA*; *adhE*, *ldhA*, *pflAB*, *mdh*, *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE*, *ycdW*; and *adhE*, *ldhA*, *pflAB*, *mdh*, *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE*, *ycdW*, *prpC*, *gsk*, or an ortholog thereof.

[0113] In one embodiment, the disclosure provides a method of producing 1,4-butanediol coupled to the growth of a microorganism. The method can include the steps of culturing under exponential growth phase in a sufficient amount of nutrients and media a non-naturally occurring microorganism comprising a set of metabolic modifications obligatorily coupling 1,4-butanediol production to growth of the microorganism, the set of metabolic modifications comprising disruption of one or more genes, or an ortholog thereof, wherein the set of metabolic modifications comprises disruption of *adhE* and *ldhA*, wherein the microorganism exhibits stable growth-coupled production of 1,4-butanediol; and isolating 1,4-butanediol produced from the non-naturally occurring microorganism. In an additional disclosure the set of metabolic modifications can further comprise disruption of *mdh*. In another disclosure, the set of metabolic modifications can further comprise disruption of one or more genes selected from the set of genes comprising *mgo*, *aspA*, *sfcA*, *maeB*, *pntAB*, and *gdhA* and can include, for example, disruption of *sfcA* and *maeB*. In still another disclosure, the set of metabolic modifications can further comprise disruption of one or more genes selected from the set of genes comprising *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE*, *ycdW*, *prpC*, and *gsk*, including disruption of all of *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE* and *ycdW*, and can further comprise disruption of *prpC* and *gsk*. Any of the above-described set of metabolic modifications can further comprise disruption of *pflAB*.

[0114] In a method of producing BDO, the non-naturally occurring microorganism can further comprise a 1,4-butanediol (BDO) biosynthetic pathway comprising at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or

alcohol dehydrogenase, wherein the exogenous nucleic acid is expressed in sufficient amounts to produce 1,4-butanediol (BDO).

[0115] It is understood that modifications which do not substantially affect the activity of the various embodiments of this invention are also included within the definition of the invention provided herein. Accordingly, the following examples are intended to illustrate but not limit the present invention.

EXAMPLE I

Biosynthesis of 4-Hydroxybutanoic Acid

[0116] This Example describes the biochemical pathways for 4-HB production.

[0117] Previous reports of 4-HB synthesis in microbes have focused on this compound as an intermediate in production of the biodegradable plastic poly-hydroxyalkanoate (PHA) (U.S. Patent No. 6,117,658). The use of 4-HB/3-HB copolymers can be advantageous over the more traditional poly-3-hydroxybutyrate polymer (PHB) because the resulting plastic is less brittle (Saito and Doi, Intl. J. Biol. Macromol. 16:99-104 (1994)). The production of monomeric 4-HB described herein is a fundamentally different process for several reasons: 1) The product is secreted, as opposed to PHA which is produced intracellularly and remains in the cell; 2) for organisms that produce hydroxybutanoate polymers, free 4-HB is not produced, but rather the Coenzyme A derivative is used by the polyhydroxyalkanoate synthase; 3) in the case of the polymer, formation of the granular product changes thermodynamics; and 4) extracellular pH is not an issue for production of the polymer, whereas it will affect whether 4-HB is present in the free acid or conjugate base state, and also the equilibrium between 4-HB and GBL.

[0118] 4-HB can be produced in two enzymatic reduction steps from succinate, a central metabolite of the TCA cycle, with succinic semialdehyde as the intermediate (Figure 2). The first of these enzymes, succinic semialdehyde dehydrogenase, is native to many organisms including *E. coli*, in which both NADH- and NADPH-dependent enzymes have been found (Donnelly and Cooper, Eur. J. Biochem. 113:555-561 (1981); Donnelly and Cooper, J. Bacteriol. 145:1425-1427 (1981); Marek and Henson, J. Bacteriol. 170:991-994 (1988)). There is also evidence supporting succinic semialdehyde dehydrogenase activity in *S. cerevisiae* (Ramos et al., Eur. J. Biochem. 149:401-404 (1985)), and a putative gene has been identified by sequence homology. However, most reports indicate that this enzyme proceeds in the direction of succinate synthesis, opposite to that shown in Figure 2 (Donnelly and Cooper, supra; Lutke-Eversloh and Steinbuchel, FEMS Microbiol. Lett. 181:63-71 (1999)), participating in the degradation pathway of 4-HB and gamma-aminobutyrate. An alternative pathway, used by the obligate anaerobe *Clostridium kluyveri* to degrade succinate, activates succinate to succinyl-CoA, then converts succinyl-CoA to succinic semialdehyde using an alternative succinic semialdehyde dehydrogenase which is known to function in this direction (Sohling and Gottschalk, Eur. J. Biochem. 212:121-127 (1993)). However, this route has the energetic cost of ATP required to convert succinate to succinyl-CoA.

[0119] The second enzyme of the pathway, 4-hydroxybutanoate dehydrogenase, is not native to *E. coli* or yeast but is found in various bacteria such as *C. kluyveri* and *Ralstonia eutropha* (Lutke-Eversloh and Steinbuchel, supra; Sohling and Gottschalk, J. Bacteriol. 178:871-880 (1996); Valentin et al., Eur. J. Biochem. 227:43-60 (1995); Wolff and Kenealy, Protein Expr. Purif. 6:206-212 (1995)). These enzymes are known to be NADH-dependent, though NADPH-dependent forms can exist. An additional pathway to 4-HB from alpha-ketoglutarate was demonstrated in *E. coli* resulting in the accumulation of poly(4-hydroxybutyric acid) (Song et al., Wei Sheng Wu Xue Bao. 45:382-386 (2005)). The recombinant strain required the overexpression of three heterologous genes, PHA synthase (*R. eutropha*), 4-hydroxybutyrate dehydrogenase (*R. eutropha*) and 4-hydroxybutyrate:CoA transferase (*C. kluyveri*), along with two native *E. coli* genes: glutamate:succinic semialdehyde transaminase and glutamate decarboxylase. Steps 4 and 5 in Figure 2 can alternatively be carried out by an alpha-ketoglutarate decarboxylase such as the one identified in *Euglena gracilis* (Shigeoka et al., Biochem. J. 282(Pt2):319-323 (1992); Shigeoka and Nakano, Arch. Biochem. Biophys. 288:22-28 (1991); Shigeoka and Nakano, Biochem. J. 292(Pt 2):463-467 (1993)). However, this enzyme has not yet been applied to impact the production of 4-HB or related polymers in any organism.

[0120] The reported directionality of succinic semialdehyde dehydrogenase led to the investigation of the thermodynamics of 4-HB metabolism. Specifically, this study investigated whether or not the reactions involved in the conversion of succinate or succinyl-CoA to 4-HB are thermodynamically favorable (i.e., $\Delta G_r < 0$) under the typical physiological conditions present in *E. coli* and *S. cerevisiae*. All oxidation/reduction reactions were assumed to utilize NADH, although the results for assuming NADPH utilization would be similar. Standard Gibbs free energies of formation (ΔG_f°) were calculated for each compound in Figure 2 based on the group contribution method (Mavrouniotis, M.L., J. Biol. Chem. 266:14440-14445 (1991)). Each standard Gibbs energy of formation was then transformed in order to obtain a criterion of spontaneous change at specified pressure, temperature, pH, and ionic strength (Alberty, R.A., Biochem. Biophys. Acta 1207:1-11 (1994)) (equation 1).

$$\Delta G_f'(I, pH) = \Delta G_f^\circ(I=0) + N_H RT \ln(10^{pH}) - 2.915 \sqrt{I} \frac{z^2 - N_H}{1 + B\sqrt{I}} \quad (1)$$

[0121] Where ΔG_f° is the standard Gibbs energy of formation, N_H is the number of hydrogen atoms in the compound, R is the universal gas constant, T is constant at 298K, z is the charge of the molecule at the pH of interest, I is the ionic strength in M, and B is a constant equal to $1.6 \text{ L}^{0.5}/\text{mol}^{0.5}$.

[0122] Equation 1 reveals that both intracellular pH and ionic strength play a role in determining thermodynamic feasibility. Normally, intracellular pH of cells is very well regulated, even when there are large variations in the culture pH. The intracellular pH of *E. coli* and *S. cerevisiae* have both been reported in the literature. *E. coli* maintains an intracellular pH of 7.4-7.7 during typical growth conditions in neutral buffers, but can drop to 7.2 in pH 6 medium, and even go as low as 6.9 for external pH of 5 (Riondet et al., Biotechnology Tech. 11:735-738 (1997)). However, growth of *E. coli* is severely inhibited at external pH below 6. Yeast pH exhibits more variation. During exponential growth phase, *S. cerevisiae* internal pH has been measured to be in the range of 6.7-7.0 with external pH controlled at 5.0 (Dombek and Ingram, Appl. Environ. Microbiol. 53:1286-1291 (1987)). On the other hand, in resting cells the internal pH drops to below 6 when the external pH is 6 or less (Imai and Ohno, J. Biotechnol. 38:165-172 (1995)). This analysis assumes an intracellular pH of 7.4 for *E. coli* and 6.8 for *S. cerevisiae*. An ionic strength of 0.15 also was assumed (Valenti et al., *supra*).

[0123] Transformed Gibbs energies of formation were calculated at the standard state (pH = 7.0, $I = 0$) and at physiological states of *E. coli* (pH = 7.4, $I = 0.15$) and *S. cerevisiae* (pH = 6.8, $I = 0.15$). Transformed Gibbs energies of reaction ($\Delta G_r'$) were then calculated by taking the difference in $\Delta G_f'$ between the products and reactants. The transformed Gibbs energies of the reactions necessary to convert succinate or succinyl-CoA to 4-HB are provided in Table 2. Note that the standard error, $U_{f,est}$, on ΔG_f calculated by the group contribution theory is 4 kcal/mol. The uncertainty in ΔG_r , $U_{r,est}$, can be calculated as the Euclidean norm of the uncertainty for ΔG_f of each compound (Equation).

$$U_{r,est} = \sqrt{\sum_{i=1}^n n_i^2 * U_{f,est}^2} = \sqrt{\sum_{i=1}^n 16n_i^2} \quad (2)$$

Where n is the stoichiometric coefficient and i is the compound. For the examined reactions, this uncertainty is on the order of 8 kcal/mol.

Table 2. Gibbs free energy of reaction (kcal/mole) at different pH and ionic strength values. The first column is under standard conditions, while the others are adjusted according to equation 1. Temperature is constant at 298 K. Error bars for these values are on the order of 8 kcal/mol, as calculated by equation 2. Abbreviations: suc, succinate; sucса, succinic semialdehyde; succoa, succinyl-CoA; Pi, inorganic phosphate.

Reaction	$\Delta G_r^{\circ'}$ pH = 7.0 IS = 0	$\Delta G_r'$ pH = 7.4 IS = 0.15 M	$\Delta G_r'$ pH = 6.8 IS = 0.15 M
succ + NADH + 2 H ⁺ → sucса + NAD + h ₂ o	12.0	14.4	12.8
succ + coa + ATP → succoa + ADP + Pi	0.30	-0.03	-0.03
succoa + NADH + H ⁺ → sucса + NAD + coa	4.4	7.0	6.2
sucса + NADH + H ⁺ → 4hb + NAD	-5.0	-3.8	-4.6

[0124] Table 2 reveals that the reaction most likely to encounter a thermodynamic barrier after considering potential uncertainty in our calculations is succinic semialdehyde dehydrogenase (step 1 in Figure 2). Whether this reaction can be driven closer to thermodynamic feasibility by varying the assumed concentrations of the participating metabolites also was studied. For example, the standard Gibbs energies assume concentrations of 1 M for all participating compounds (except water). In an anaerobic environment, NADH will be present at a several-fold higher concentration than NAD. Assuming $[\text{NADH}] = 5 \times [\text{NAD}]$, we calculated the effect on $\Delta G_r'$ using the equation

$$\Delta G_f' = \Delta G_f^{0'} + RT \ln \frac{\prod [prod]}{\prod [react]} \quad (3)$$

[0125] This change results in a difference of only about 1 kcal/mol in the delta G values for succinic semialdehyde dehydrogenase. Equation 3 was also used to calculate other effects on ΔG_r , such as high succinate concentration to drive the reactions. A 1000-fold difference in the concentrations of succinate and succinic semialdehyde will contribute about 5 kcal/mol to delta G. Taken together with an assumed uncertainty of 8 kcal/mol, the possibility that succinic semialdehyde dehydrogenase will operate in the direction towards succinic semialdehyde under some set of physiological conditions cannot be eliminated. Thus we still consider the direct route from succinate to 4-HB in our subsequent theoretical analysis.

[0126] The microbial production capabilities of 4-hydroxybutyrate were explored in two microbes, *Escherichia coli* and *Saccharomyces cerevisiae*, using *in silico* metabolic models of each organism. Potential pathways to 4-HB proceed via a succinate, succinyl-CoA, or alpha-ketoglutarate intermediate as shown in Figure 2.

[0127] The first step in the 4-HB production pathway from succinate involves the conversion of succinate to succinic semialdehyde via an NADH- or NADPH-dependant succinic semialdehyde dehydrogenase. In *E. coli*, *gabD* is an NADP-dependant succinic semialdehyde dehydrogenase and is part of a gene cluster involved in 4-aminobutyrate uptake and degradation (Niegemann et al., Arch. Microbiol. 160:454-460 (1993); Schneider et al., J. Bacteriol. 184:6976-6986 (2002)). *sad* is believed to encode the enzyme for NAD-dependant succinic semialdehyde dehydrogenase activity (Marek and Henson, *supra*). *S. cerevisiae* contains only the NADPH-dependant succinic semialdehyde dehydrogenase, putatively assigned to UGA2, which localizes to the cytosol (Huh et al., Nature 425:686-691 (2003)). The maximum yield calculations assuming the succinate pathway to 4-HB in both *E. coli* and *S. cerevisiae* require only the assumption that a non-native 4-HB dehydrogenase has been added to their metabolic networks.

[0128] The pathway from succinyl-CoA to 4-hydroxybutyrate was described in U.S1 Patent No. 6,117,658 as part of a process for making polyhydroxyalkanoates comprising 4-hydroxybutyrate monomer units. *Clostridium kluyveri* is one example organism known to possess CoA-dependant succinic semialdehyde dehydrogenase activity (Sohling and Gottschalk, *supra*; Sohling and Gottschalk, *supra*). In this study, it is assumed that this enzyme, from *C. kluyveri* or another organism, is expressed in *E. coli* or *S. cerevisiae* along with a non-native or heterologous 4-HB dehydrogenase to complete the pathway from succinyl-CoA to 4-HB. The pathway from alpha-ketoglutarate to 4-HB was demonstrated in *E. coli* resulting in the accumulation of poly(4-hydroxybutyric acid) to 30% of dry cell weight (Song et al., *supra*). As *E. coli* and *S. cerevisiae* natively or endogenously possess both glutamate:succinic semialdehyde transaminase and glutamate decarboxylase (Coleman et al., J. Biol. Chem. 276:244-250 (2001)), the pathway from AKG to 4-HB can be completed in both organisms by assuming only that a non-native 4-HB dehydrogenase is present.

EXAMPLE II

Production of 4-Hydroxybutanoic Acid in *E. coli*

[0129] This Example describes the biosynthetic yields for 4-hydroxybutanoic acid resulting from each biochemical pathway.

[0130] In this section, the maximum theoretical yields of 4-HB from glucose are calculated assuming that each of the three metabolic pathways depicted in Figure 2 are functional in *E. coli*. A genome-scale metabolic model of *E. coli*, similar to the one described in Reed et al., Genome Biol. 4:R54 (2003), was used as the basis for the analysis. The energetic gain, in terms of ATP molecules produced, of each maximum yielding pathway is calculated assuming anaerobic conditions, unless otherwise stated. 4-Hydroxybutyrate is assumed to exit in *E. coli* via proton symport, as is the case with most organic acids. It is also possible that GBL is secreted by simple diffusion, and in this case the energetics would be more favorable than in the case considered here. The impact of cofactor specificity (i.e., NADH or NADPH-dependence) of the participating enzymes on the maximum yield and energetics of each pathway also was investigated.

[0131] The results from the analysis are shown in Tables 3 A-C. From an energetic and yield standpoint, the succinate to 4-HB pathway is the most promising provided that the thermodynamic concerns raised in Example I can be overcome. Specifically, the calculations reveal that the maximum theoretical yield of 4-HB from glucose is 1.33 mol/mol (0.77 g/g; 0.89 Cmol/Cmol) assuming the succinate to 4-HB pathway is functional. In addition, the anaerobic production of 4-HB via succinate would result in the net production of either 1.8, 1.5, or 1.1 mol of ATP per glucose depending upon the assumed cofactor specificity of the participating enzymes. These energetic yields are comparable to the 2.0 ATP per glucose that can be obtained via substrate level phosphorylation by the production of ethanol or lactate suggesting the potential for anaerobic homo-4-HB production in *E. coli*.

[0132] The succinyl-CoA route to 4-HB is the second most promising pathway when considering maximum yield and

energetics. A 1.33 mol/mol yield of 4-HB is achievable in *E. coli* if at least one of the pathway steps is assumed NADH-dependant. However, because this pathway requires the formation of succinyl-CoA, its energetic yield is considerably lower than that of the succinate pathway. An oxygen requirement is anticipated at high 4-HB yields if both the CoA-dependant succinic semialdehyde dehydrogenase and 4-HB dehydrogenase steps are assumed NADPH-dependant. In this case, the production of 4-HB at the maximum yield would result in no net ATP gain and could not support the energetic maintenance demands needed for *E. coli* survival. Thus, some energy would have to originate from oxidative phosphorylation to enable homo-fermentative 4-HB production. The alpha-ketoglutarate pathway toward 4-HB is the least favorable of the three potential routes with a maximum achievable yield of 1.0 mol 4-HB per mol of glucose. In addition to the lower maximum yield, this pathway requires the utilization of 1.5 moles of oxygen per mol of glucose converted to 4-HB. The energetics of this pathway are unaffected by the assumed cofactor specificity of 4-HB dehydrogenase.

Table 3. The overall substrate conversion stoichiometry to 4-HB assuming the A) succinate, B) succinyl-CoA, or C) alpha-ketoglutarate production routes are functional in *E. coli*. Glucose and oxygen are taken up while all other molecules are produced.

A) Succinate Pathway				
Cofactor Specificity	2 NADH steps	1 NADH step 1 NADPH step	2 NADPH steps	
Glucose	-1.000	-1.000	-1.000	
Oxygen	0.000	0.000	0.000	
Protons	1.333	1.333	1.333	
4HB	1.333	1.333	1.333	
CO2	0.667	0.667	0.667	
H2O	0.667	0.667	0.667	
ATP	1.800	1.510	1.097	
B) Succinyl-CoA Pathway				
Cofactor Specificity	2 NADH steps	1 NADH step 1 NADPH step	2 NADPH steps	2 NADPH steps
Glucose	-1.000	-1. 000	-1.000	-1.000
Oxygen	0.000	0.000	-0.036	0.000
Protons	1.333	1.333	1.325	1.294
4HB	1.333	1.333	1.325	1.294
CO2	0.667	0.667	0.698	0.082
H2O	0.667	0.667	0.698	0.470
ATP	0.467	0.177	0.000	0.000
C) Alpha-ketoglutarate Pathway				
Cofactor Specificity	1 NADH step		1 NADPH step	
Glucose	-1.000		-1.000	
Oxygen	-1.500		-1.500	
Protons	1.000		1.000	
4HB	1.000		1.000	
CO2	2.000		2.000	
H2O	2.000		2.000	

(continued)

C) Alpha-ketoglutarate Pathway

Cofactor Specificity	1 NADH step	1 NADPH step
ATP	5.500	5.500

[0133] In order to corroborate the computational predictions proposed in this report, the strains expressing a complete pathway to 4-HB can be constructed and tested. Corroboration is performed with both *E. coli* (Example II) and *S. cerevisiae* (Example III). In *E. coli*, the relevant genes are expressed in a synthetic operon behind an inducible promoter on a medium- or high-copy plasmid; for example the P_{BAD} promoter which is induced by arabinose, on a plasmid of the pBAD series (Guzman et al., J. Bacteriol. 177:4121-4130 (1995)). In *S. cerevisiae*, genes are integrated into the chromosome behind the PDC1 promoter, replacing the native pyruvate carboxylase gene. It has been reported that this results in higher expression of foreign genes than from a plasmid (Ishida et al., Appl. Environ. Microbiol. 71:1964-1970 (2005)), and will also ensure expression during anaerobic conditions.

[0134] Cells containing the relevant constructs are grown in minimal media containing glucose, with addition of arabinose in the case of *E. coli* containing genes expressed under the P_{BAD} promoter. Periodic samples are taken for both gene expression and enzyme activity analysis. Enzyme activity assays are performed on crude cell extracts using procedures well known in the art. Alternatively, assays based on the oxidation of NAD(P)H, which is produced in all dehydrogenase reaction steps and detectable by spectrophotometry can be utilized. In addition, antibodies can be used to detect the level of particular enzymes. In lieu of or in addition to enzyme activity measurements, RNA can be isolated from parallel samples and transcript of the gene of interest measured by reverse transcriptase PCR. Any constructs lacking detectable transcript expression are reanalyzed to ensure the encoding nucleic acids are harbored in an expressible form. Where transcripts are detected, this result indicates either a lack of translation or production of inactive enzyme. A variety of methods well known in the art can additionally be employed, such as codon optimization, engineering a strong ribosome binding site, use of a gene from a different species, and prevention of N-glycosylation (for expression of bacterial enzymes in yeast) by conversion of Asn residues to Asp. Once all required enzyme activities are detected, the next step is to measure the production of 4-HP *in vivo*. Triplicate shake flask cultures are grown either anaerobically or microaerobically, depending on the conditions required (see above), and periodic samples taken. Organic acids present in the culture supernatants are analyzed by HPLC using the Aminex AH-87X column. The elution time of 4-HB will be determined using a standard purchased from a chemical supplier.

[0135] The CoA-independent pathway can be implemented and tested for corroboration. In this case, the genes overexpressed are the native succinic semialdehyde dehydrogenase from each organism, and the 4-hydroxybutanoate dehydrogenase from *Ralstonia eutropha*. Once both enzyme activities are detected as discussed above, the strains are tested for 4-HB production. Corroboration also can be obtained from implementing the CoA-dependent pathway. The CoA-dependent succinic semialdehyde dehydrogenase and the 4-hydroxybutanoate dehydrogenase from *Clostridium kluyveri* are expressed as described above. In addition, overexpression of the native succinyl-CoA synthetase also can be performed, to funnel more succinate into the heterologous pathway. Finally, if 4-HB production is unfavorable, different culture conditions can be tested, such as a change in oxygenation status which can manipulate the NAD(P)H/NAD(P) ratio.

EXAMPLE III**Production of 4-Hydroxybutanoic Acid in Yeast**

[0136] This Example describes the biosynthetic yields for 4-hydroxybutanoic acid resulting from each biochemical pathway in *S. cerevisiae*.

[0137] In this section, the maximum theoretical yields of 4-HB from glucose are calculated assuming that each of the three metabolic pathways depicted in Figure 2 are functional in *S. cerevisiae*. A genome-scale metabolic model of *S. cerevisiae*, similar to the one described in Forster et al. (Genome Res. 13:244-253 (2003)) was used as the basis for the analysis. The energetic gain of each maximum yielding pathway is calculated assuming anaerobic conditions unless otherwise stated. 4-hydroxybutyrate is assumed to exit *S. cerevisiae* via proton symport, as is the case with most organic acids. The impact of cofactor specificity (i.e., NADH or NADPH-dependence) of the participating enzymes on the maximum yield and energetics of each pathway was also investigated.

[0138] The results from the analysis are shown in Tables 4 A-C. As with *E. coli*, the succinate to 4-HB pathway is the most promising provided that the thermodynamic concerns raised in Example I can be overcome. The calculations reveal

that the maximum theoretical yield of 4-HB from glucose is 1.33 mol/mol (0.77 g/g; 0.89 Cmol/Cmol) in *S. cerevisiae*. In addition, the anaerobic production of 4-HB via succinate would result in the net production of either 1.4, 1.1, or 0.5 mol of ATP per glucose depending upon the assumed cofactor specificity of the participating enzymes.

[0139] The succinyl-CoA route to 4-HB is the second most favorable pathway. A maximum yield of 1.33 mol 4-HB/mol glucose is achievable in *S. cerevisiae* regardless of cofactor specificity. However, net energy generation at the maximum theoretical yield is possible only if both the CoA-dependant succinic semialdehyde dehydrogenase and 4-HB dehydrogenase steps are assumed to be NADH-dependant. If either step is NADPH-dependant, no net ATP will be gained from anaerobic 4-HB production and an alternate energy source (e.g., oxidative phosphorylation) would be required to support cell growth and maintenance. The alpha-ketoglutarate route toward 4-HB is the least favorable of the three potential pathways in *S. cerevisiae* although the maximum yield of 1.1-1.2 mol 4-HB per mol glucose is slightly higher than was found in *E. coli*. Nevertheless, this pathway requires an oxygen uptake of 0.8-0.9 mol oxygen per mol glucose to become energetically neutral.

Table 4. The overall substrate conversion stoichiometry to 4-HB in *S. cerevisiae*., assuming the A) succinate, B) succinyl-CoA, or C) alpha-ketoglutarate production routes are functional in *S. cerevisiae*. Glucose and oxygen are taken up while all other molecules are produced.

A) Succinate Pathway			
Cofactor Specificity	2 NADH steps	1 NADH step 1 NADPH step	2 NADPH steps
Glucose	-1.000	-1.000	-1.000
Oxygen	0.000	0.000	0.000
Protons	1.333	1.333	1.333
4HB	1.333	1.333	1.333
CO₂	0.667	0.667	0.667
H₂O	0.667	0.667	0.667
ATP	1.444	1.067	0.533
B) Succinyl-CoA Pathway			
Cofactor Specificity	2 NADH steps	1 NADH step 1 NADPH step	2 NADPH steps
Glucose	-1.000	-1.000	-1.000
Oxygen	0.000	0.000	0.000
Protons	1.333	1.333	1.333
4HB	1.333	1.333	1.333
CO₂	0.667	0.667	0.667
H₂O	0.667	0.667	0.667
ATP	0.533	0.000	0.000
C) Alpha-ketoglutarate Pathway			
Cofactor Specificity	1 NADH step	1 NADPH step	
Glucose	-1.000	-1.000	
Oxygen	-0.785	-0.879	
Protons	1.159	1.138	
4HB	1.159	1.138	
CO₂	1.364	1.448	
H₂O	1.364	1.448	

(continued)

C) Alpha-ketoglutarate Pathway

Cofactor Specificity	1 NADH step	1 NADPH step
ATP	0.000	0.000

EXAMPLE IV

Biosynthesis of 4-Hydroxybutanoic Acid, γ -Butyrolactone and 1,4-Butanediol

[0140] This Example describes the biosynthetic production of 4-hydroxybutanoic acid, γ -butyrolactone and 1,4-butanediol using fermentation and other bioprocesses.

[0141] Methods for the integration of the 4-HB fermentation step into a complete process for the production of purified GBL, 1,4-butanediol (BDO) and tetrahydrofuran (THF) are described below. Since 4-HB and GBL are in equilibrium, the fermentation broth will contain both compounds. At low pH this equilibrium is shifted to favor GBL. Therefore, the fermentation can operate at pH 7.5 or less. After removal of biomass, the product stream enters into a separation step in which GBL is removed and the remaining stream enriched in 4-HB is recycled. Finally, GBL is distilled to remove any impurities. The process operates in one of three ways: 1) fed-batch fermentation and batch separation; 2) fed-batch fermentation and continuous separation; 3) continuous fermentation and continuous separation. The first two of these modes are shown schematically in Figure 4. The integrated fermentation procedures described below also are used for the BDO producing cells of the invention for biosynthesis of BDO and subsequent BDO family products.

[0142] Fermentation protocol to produce 4-HB/GBL (batch): The production organism is grown in a 10L bioreactor sparged with an N_2/CO_2 mixture, using 5 L broth containing 5 g/L potassium phosphate, 2.5 g/L ammonium chloride, 0.5 g/L magnesium sulfate, and 30 g/L corn steep liquor, and an initial glucose concentration of 20 g/L. As the cells grow and utilize the glucose, additional 70% glucose is fed into the bioreactor at a rate approximately balancing glucose consumption. The temperature of the bioreactor is maintained at 30 degrees C. Growth continues for approximately 24 hours, until 4-HB reaches a concentration of between 20-200 g/L, with the cell density being between 5 and 10 g/L. The pH is not controlled, and will typically decrease to pH 3-6 by the end of the run. Upon completion of the cultivation period, the fermenter contents are passed through a cell separation unit (e.g., centrifuge) to remove cells and cell debris, and the fermentation broth is transferred to a product separations unit. Isolation of 4-HB and/or GBL would take place by standard separations procedures employed in the art to separate organic products from dilute aqueous solutions, such as liquid-liquid extraction using a water immiscible organic solvent (e.g., toluene) to provide an organic solution of 4-HB/GBL. The resulting solution is then subjected to standard distillation methods to remove and recycle the organic solvent and to provide GBL (boiling point 204-205°C) which is isolated as a purified liquid.

[0143] Fermentation protocol to produce 4-HB/GBL (fully continuous): The production organism is first grown up in batch mode using the apparatus and medium composition described above, except that the initial glucose concentration is 30-50 g/L. When glucose is exhausted, feed medium of the same composition is supplied continuously at a rate between 0.5 L/hr and 1 L/hr, and liquid is withdrawn at the same rate. The 4-HB concentration in the bioreactor remains constant at 30-40 g/L, and the cell density remains constant between 3-5 g/L. Temperature is maintained at 30 degrees C, and the pH is maintained at 4.5 using concentrated NaOH and HCl, as required. The bioreactor is operated continuously for one month, with samples taken every day to assure consistency of 4-HB concentration. In continuous mode, fermenter contents are constantly removed as new feed medium is supplied. The exit stream, containing cells, medium, and products 4-HB and/or GBL, is then subjected to a continuous product separations procedure, with or without removing cells and cell debris, and would take place by standard continuous separations methods employed in the art to separate organic products from dilute aqueous solutions, such as continuous liquid-liquid extraction using a water immiscible organic solvent (e.g., toluene) to provide an organic solution of 4-HB/GBL. The resulting solution is subsequently subjected to standard continuous distillation methods to remove and recycle the organic solvent and to provide GBL (boiling point 204-205°C) which is isolated as a purified liquid.

[0144] GBL Reduction Protocol: Once GBL is isolated and purified as described above, it will then be subjected to reduction protocols such as those well known in the art (references cited) to produce 1,4-butanediol or tetrahydrofuran (THF) or a mixture thereof. Heterogeneous or homogeneous hydrogenation catalysts combined with GBL under hydrogen pressure are well known to provide the products 1,4-butanediol or tetrahydrofuran (THF) or a mixture thereof. It is important to note that the 4-HB/GBL product mixture that is separated from the fermentation broth, as described above, may be subjected directly, prior to GBL isolation and purification, to these same reduction protocols to provide the products 1,4-butanediol or tetrahydrofuran or a mixture thereof. The resulting products, 1,4-butanediol and THF are then

isolated and purified by procedures well known in the art.

[0145] Fermentation and hydrogenation protocol to produce BDO or THF directly (batch): Cells are grown in a 10L bioreactor sparged with an N_2/CO_2 mixture, using 5 L broth containing 5 g/L potassium phosphate, 2.5 g/L ammonium chloride, 0.5 g/L magnesium sulfate, and 30 g/L corn steep liquor, and an initial glucose concentration of 20 g/L. As the cells grow and utilize the glucose, additional 70% glucose is fed into the bioreactor at a rate approximately balancing glucose consumption. The temperature of the bioreactor is maintained at 30 degrees C. Growth continues for approximately 24 hours, until 4-HB reaches a concentration of between 20-200 g/L, with the cell density being between 5 and 10 g/L. The pH is not controlled, and will typically decrease to pH 3-6 by the end of the run. Upon completion of the cultivation period, the fermenter contents are passed through a cell separation unit (e.g., centrifuge) to remove cells and cell debris, and the fermentation broth is transferred to a reduction unit (e.g., hydrogenation vessel), where the mixture 4-HB/GBL is directly reduced to either 1,4-butanediol or THF or a mixture thereof. Following completion of the reduction procedure, the reactor contents are transferred to a product separations unit. Isolation of 1,4-butanediol and/or THF would take place by standard separations procedures employed in the art to separate organic products from dilute aqueous solutions, such as liquid-liquid extraction using a water immiscible organic solvent (e.g., toluene) to provide an organic solution of 1,4-butanediol and/or THF. The resulting solution is then subjected to standard distillation methods to remove and recycle the organic solvent and to provide 1,4-butanediol and/or THF which are isolated as a purified liquids.

[0146] Fermentation and hydrogenation protocol to produce BDO or THF directly (fully continuous): The cells are first grown up in batch mode using the apparatus and medium composition described above, except that the initial glucose concentration is 30-50 g/L. When glucose is exhausted, feed medium of the same composition is supplied continuously at a rate between 0.5 L/hr and 1 L/hr, and liquid is withdrawn at the same rate. The 4-HB concentration in the bioreactor remains constant at 30-40 g/L, and the cell density remains constant between 3-5 g/L. Temperature is maintained at 30 degrees C, and the pH is maintained at 4.5 using concentrated NaOH and HCl, as required. The bioreactor is operated continuously for one month, with samples taken every day to assure consistency of 4-HB concentration. In continuous mode, fermenter contents are constantly removed as new feed medium is supplied. The exit stream, containing cells, medium, and products 4-HB and/or GBL, is then passed through a cell separation unit (e.g., centrifuge) to remove cells and cell debris, and the fermentation broth is transferred to a continuous reduction unit (e.g., hydrogenation vessel), where the mixture 4-HB/GBL is directly reduced to either 1,4-butanediol or THF or a mixture thereof. Following completion of the reduction procedure, the reactor contents are transferred to a continuous product separations unit. Isolation of 1,4-butanediol and/or THF would take place by standard continuous separations procedures employed in the art to separate organic products from dilute aqueous solutions, such as liquid-liquid extraction using a water immiscible organic solvent (e.g., toluene) to provide an organic solution of 1,4-butanediol and/or THF. The resulting solution is then subjected to standard continuous distillation methods to remove and recycle the organic solvent and to provide 1,4-butanediol and/or THF which are isolated as a purified liquids.

[0147] Fermentation protocol to produce BDO directly (batch): The production organism is grown in a 10L bioreactor sparged with an N_2/CO_2 mixture, using 5 L broth containing 5 g/L potassium phosphate, 2.5 g/L ammonium chloride, 0.5 g/L magnesium sulfate, and 30 g/L corn steep liquor, and an initial glucose concentration of 20 g/L. As the cells grow and utilize the glucose, additional 70% glucose is fed into the bioreactor at a rate approximately balancing glucose consumption. The temperature of the bioreactor is maintained at 30 degrees C. Growth continues for approximately 24 hours, until BDO reaches a concentration of between 20-200 g/L, with the cell density generally being between 5 and 10 g/L. Upon completion of the cultivation period, the fermenter contents are passed through a cell separation unit (e.g., centrifuge) to remove cells and cell debris, and the fermentation broth is transferred to a product separations unit. Isolation of BDO would take place by standard separations procedures employed in the art to separate organic products from dilute aqueous solutions, such as liquid-liquid extraction using a water immiscible organic solvent (e.g., toluene) to provide an organic solution of BDO. The resulting solution is then subjected to standard distillation methods to remove and recycle the organic solvent and to provide BDO (boiling point 228-229°C) which is isolated as a purified liquid.

[0148] Fermentation protocol to produce BDO directly (fully continuous): The production organism is first grown up in batch mode using the apparatus and medium composition described above, except that the initial glucose concentration is 30-50 g/L. When glucose is exhausted, feed medium of the same composition is supplied continuously at a rate between 0.5 L/hr and 1 L/hr, and liquid is withdrawn at the same rate. The BDO concentration in the bioreactor remains constant at 30-40 g/L, and the cell density remains constant between 3-5 g/L. Temperature is maintained at 30 degrees C, and the pH is maintained at 4.5 using concentrated NaOH and HCl, as required. The bioreactor is operated continuously for one month, with samples taken every day to assure consistency of BDO concentration. In continuous mode, fermenter contents are constantly removed as new feed medium is supplied. The exit stream, containing cells, medium, and the product BDO, is then subjected to a continuous product separations procedure, with or without removing cells and cell debris, and would take place by standard continuous separations methods employed in the art to separate organic products from dilute aqueous solutions, such as continuous liquid-liquid extraction using a water immiscible organic solvent (e.g., toluene) to provide an organic solution of BDO. The resulting solution is subsequently subjected to standard continuous distillation methods to remove and recycle the organic solvent and to provide BDO (boiling point

228-229°C) which is isolated as a purified liquid (mpt 20°C).

EXAMPLE V

In silico Derived Knockout Strategies for Growth-Coupled Production of 1,4-Butanediol in *Escherichia coli*

[0149] This example describes the *in silico* design of knockout strategies to generate growth-coupled production of 1,4-butanediol in *E. coli*.

[0150] The strategy for generating growth-coupled production of BDO first involves the demonstration of a functional pathway which is subsequently optimized through adaptive evolution and targeted gene insertions, deletions, and over-expressions. Described below in more detail are strain-engineering strategies identified via OptKnock for generating growth-coupled BDO production strains of *Escherichia coli*. All implementations of OptKnock assume that sufficient activities of all enzymes outlined in Figure 6 are available to *E. coli* under all conditions. In addition, all reduction steps in this pathway are assumed to be NADH-dependent although many of the designs are expected to be applicable regardless of cofactor specificity.

[0151] The BDO yield potential of the biochemical pathways to 1,4-butanediol in *E. coli* is detailed below. Three conditions were used, anaerobic, anaerobic + nitrate addition, and aerobic. The maximum theoretical yields for each scenario assuming that the production of BDO must be energetically neutral are shown in Table 5. Table 5 shows the maximum theoretical yields of 1,4-butanediol (BDO) for five sets of environmental conditions: 1) aerobic respiration, 2) anaerobic fermentation with acetate co-production, 3) anaerobic fermentation with ethanol co-production, 4) nitrate respiration leading to nitrite formation, and 5) nitrate respiration leading to ammonia formation. Negative values indicate metabolites taken up; positive values indicate metabolites secreted. Molar units are assumed along with pathway energetic neutrality. The maximum theoretical yield without assuming energetic neutrality is 1.091 mol/mol (0.545 g/g) glucose for all cases. The highest yield is obtained under aerobic conditions where enough ATP to render the pathway energetically neutral can be generated with minimal loss of carbon through respiration. In the anaerobic case, the yield drops as ATP is made via substrate level phosphorylation and either acetate or ethanol is made as a byproduct. Controlled nitrate addition can provide nearly the same yield as oxygen, the exact amount depending upon whether further reduction of nitrite to ammonia occurs.

Table 5. Maximum theoretical yields of 1,4-butanediol (BDO) for five sets of environmental conditions.

	Aerobic	Anerobic			
		Acetate Coproduction	Ethanol Coproduction	Nitrate to Nitrite	Nitrate to Ammonia
Glucose	-1.000	-1.000	-1.000	-1.000	1.000
Oxygen	-0.068				
Nitrate				-0.147	-0.092
H ⁺		0.144			-0.184
H ₂ O	0.607	0.519	0.498	0.612	0.621
CO ₂	1.686	1.558	1.668	1.690	1.770
Nitrite				0.147	
Ammonia					0.092
Acetate		0.144			
Ethanol			0.173		
BDO	1.079	1.039	0.997	1.078	1.058

[0152] If phosphoenolpyruvate (PEP) carboxykinase is assumed to operate in the direction of phosphoenolpyruvate to oxaloacetate, BDO production becomes an energy-generating endeavor and the maximum theoretical yield for all scenarios becomes 0.545 g/g with CO₂ as the only byproduct. PEP carboxykinase is known to produce oxaloacetate from PEP in rumen bacteria such as *Mannheimia succiniciproducens* (Hong et al., Nat. Biotechnol. 22:1275-1281 (2004)). However, the role of PEP carboxykinase in producing oxaloacetate in *E. coli* is believed to be minor as compared to

PEP carboxylase possibly due to the higher K_m for bicarbonate of PEP carboxykinase (Kim et al., Appl. Environ. Microbiol. 70:1238-1241 (2004)). Nevertheless, activity of the native *E. coli* PEP carboxykinase from PEP towards oxaloacetate has been recently demonstrated in *ppc* mutants of K-12 (Kwon et al., J. Microbiol. Biotechnol. 16:1448-1452 (2006)).

[0153] In more detail, the designs identified for increasing BDO production in *E. coli* are described below. A non-growth associated energetic maintenance requirement of 7.6 mmol/gDW/hr was assumed along with a maximum specific glucose uptake rate of 20 mmol/gDW/hr. BDO was assumed to be exported via diffusion. Knockout strategies were identified assuming that 1) PEP carboxykinase is irreversible and functions only to convert oxaloacetate to PEP and 2) PEP carboxykinase is reversible. For both cases, the OptKnock code described in Burgard et al. (Biotechnol. Bioeng. 84:647-657 (2003)) was modified to allow for limited, unlimited or no nitrate respiration. Adding the possibility of nitrate respiration serves to increase the maximum theoretical yield of BDO when PEP carboxykinase is assumed irreversible and also, in some cases, enables the selection of knockout strategies that otherwise would not be selected under anaerobic conditions due to unfavorable energetics. Specifically, six nitrate uptake reactions were added to the *E. coli* network with lower bounds of 0, -2, -5, -10, -20, or -1000 mmol/gDW/hr (the negative values signify metabolite uptake in the stoichiometric model). The dual constraints of OptKnock were adjusted accordingly and an additional constraint was added that allowed only one nitrate uptake reaction to be active at a time. OptKnock selects the optimum amount of nitrate respiration for each identified knockout strategy. Finally, all simulations assume that the reaction catalyzed by *adhE* in *E. coli* (that is, acetyl-CoA + NADH $\rightarrow$ ethanol + NAD) has been removed. Because acetyl-CoA is a necessary intermediate for BDO production, nearly all OptKnock designs would include this deletion anyway to prevent ethanol production from competing with BDO production. Including this deletion *a priori* was done to lower the CPU time of the computational procedure.

[0154] The knockout strategies derived by OptKnock are given in Tables 6 and 7 assuming PEP carboxykinase to be irreversible and reversible, respectively. In this description, the knockout strategies are listed by reaction abbreviations in all capital letters for simplicity. The corresponding genes that would have to be knocked out to prevent a particular reaction from occurring in *E. coli* are provided in Table 8 along with the reaction stoichiometry. The metabolite names corresponding to Table 8 are listed in Table 9.

Table 6. Knockout strategies derived by OptKnock, assuming PEP carboxykinase to be irreversible.

	Metabolic Transformations Targeted for Removal $\uparrow, \downarrow, \#$	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
1	ADHEr	5.67	0.57	25.42	0.00	-5.74	28.42	-20.00	0.00	57.91
2	ADHEr, PFLi	9.95	0.50	22.31	0.00	14.90	0.00	-20.00	0.00	25.85
3	ADHEr, NADH6	6.92	0.52	25.03	0.00	-0.07	20.82	-20.00	0.00	49.54
4	ADHEr, THD2	6.55	0.56	23.75	0.00	-3.54	26.67	-20.00	0.00	54.38
5	ADHEr, PGI	6.21	0.30	28.99	0.00	-5.98	30.57	-20.00	0.00	61.72
6	ADHEr, PFK	6.13	0.31	29.10	0.00	-6.17	30.70	-20.00	0.00	61.98
7	ADHEr, FBA	6.13	0.31	29.10	0.00	-6.17	30.70	-20.00	0.00	61.98
8	ADHEr, ATPS4r	6.15	0.61	23.38	0.00	-3.10	26.55	-20.00	0.00	54.24
9	ADHEr, TPI	6.13	0.31	29.10	0.00	-6.17	30.70	-20.00	0.00	61.98
10	ADHEr, SUCD4	5.85	0.47	26.89	0.00	-5.92	29.33	-20.00	0.00	59.54
11	ADHEr, RPE	5.78	0.57	25.20	0.00	-5.45	28.19	-20.00	0.00	57.44
12	ADHEr, GLCpts	5.78	0.51	26.33	0.00	-5.85	28.99	-20.00	0.00	58.92
13	ADHEr, GLUDy	5.77	0.52	26.20	0.00	-5.84	28.90	-20.00	0.00	58.78
14	ADHEr, TAL	5.73	0.57	25.30	0.00	-5.59	28.30	-20.00	0.00	57.67
15	ADHEr, MDH	5.73	0.54	25.90	0.00	-5.80	28.72	-20.00	0.00	58.44
16	ADHEr, FUM	5.73	0.54	25.90	0.00	-5.80	28.72	-20.00	0.00	58.44
17	ADHEr, CBMK2	5.68	0.57	25.48	0.00	-5.75	28.46	-20.00	0.00	57.99
18	ADHEr, HEX1, PGI	11.82	0.22	18.07	0.00	8.10	19.23	-20.00	0.00	38.88

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5	19	ADHEr, EDA, PGI	11.82	0.22	18.05	0.00	8.10	19.22	-20.00	0.00	38.86
	20	ADHEr, PFLi, PGI	10.87	0.20	25.96	0.00	16.29	0.00	-20.00	0.00	27.38
10	21	ADHEr, FBA, PFLi	10.83	0.20	26.02	0.00	16.23	0.00	-20.00	0.00	27.46
	22	ADHEr, PFLi, TPI	10.83	0.20	26.02	0.00	16.23	0.00	-20.00	0.00	27.46
	23	ADHEr, PFK, PFLi	10.83	0.20	26.02	0.00	16.23	0.00	-20.00	0.00	27.46
15	24	ADHEr, PFLi, THD2	10.49	0.49	21.02	0.00	15.71	0.00	-20.00	0.00	24.49
	25	ADHEr, GLCpts, PFLi	10.15	0.43	23.16	0.00	15.20	0.00	-20.00	0.00	26.22
	26	ADHEr, GLUDy, PFLi	10.10	0.45	22.94	0.00	15.12	0.00	-20.00	0.00	26.12
	27	ADHEr, PFLi, RPE	10.02	0.50	22.14	0.00	15.00	0.00	-20.00	0.00	25.67
20	28	ADHEr, PFLi, TAL	9.99	0.50	22.22	0.00	14.95	0.00	-20.00	0.00	25.76
	29	ADHEr, CBMK2, PFLi	9.96	0.49	22.36	0.00	14.92	0.00	-20.00	0.00	25.87
	30	ADHEr, ATPS4r, PGI	8.97	0.26	23.60	0.00	0.95	24.98	-20.00	0.00	50.46
25	31	ADHEr, ATPS4r, NADH6	8.62	0.48	23.04	0.00	6.28	13.23	-20.00	0.00	39.68
	32	ADHEr, ATPS4r, TPI	8.26	0.26	27.70	0.00	3.99	16.78	-20.00	0.00	46.33
	33	ADHEr, ATPS4r, PFK	8.26	0.26	27.70	0.00	3.99	16.78	-20.00	0.00	46.33
30	34	ADHEr, ATPS4r, FBA	8.26	0.26	27.70	0.00	3.99	16.78	-20.00	0.00	46.33
	35	ADHEr, NADH6, THD2	7.88	0.50	23.49	0.00	2.67	18.21	-20.00	0.00	45.24
	36	ADHEr, ATPS4r, FUM	7.86	0.49	22.44	0.00	0.04	23.43	-20.00	0.00	49.33
	37	ADHEr, ATPS4r, MDH	7.86	0.49	22.44	0.00	0.04	23.43	-20.00	0.00	49.33
35	38	ADHEr, NADH6, PGI	7.58	0.23	28.71	0.00	0.18	22.36	-20.00	0.00	52.73
	39	ADHEr, FUM, THD2	7.55	0.49	22.66	0.00	-1.33	25.25	-20.00	0.00	51.43
	40	ADHEr, MDH, THD2	7.55	0.49	22.66	0.00	-1.33	25.25	-20.00	0.00	51.43
40	41	ADHEr, NADH6, PFK	7.50	0.24	28.80	0.00	-0.03	22.54	-20.00	0.00	53.03
	42	ADHEr, FBA, NADH6	7.50	0.24	28.80	0.00	-0.03	22.54	-20.00	0.00	53.03
	43	ADHEr, NADH6, TPI	7.50	0.24	28.80	0.00	-0.03	22.54	-20.00	0.00	53.03
45	44	ADHEr, GLCpts, NADH6	7.06	0.45	25.94	0.00	-0.06	21.24	-20.00	0.00	50.38
	45	ADHEr, NADH6, RPE	7.05	0.52	24.82	0.00	0.30	20.47	-20.00	0.00	48.96
	46	ADHEr, GLUDy, NADH6	7.02	0.47	25.73	0.00	-0.06	21.14	-20.00	0.00	50.18
50	47	ADHEr, FLTM, NADH6	7.01	0.47	25.63	0.00	-0.06	21.10	-20.00	0.00	50.09
	48	ADHEr, MDH, NADH6	7.01	0.47	25.63	0.00	-0.06	21.10	-20.00	0.00	50.09
	49	ADHEr, NADH6, TAL	6.98	0.52	24.92	0.00	0.12	20.64	-20.00	0.00	49.23
55	50	ADHEr, GLCpts, TIID2	6.56	0.49	24.86	0.00	-3.91	27.44	-20.00	0.00	55.80
	51	ADHEr, GLUDy, THD2	6.56	0.50	24.67	0.00	-3.84	27.31	-20.00	0.00	55.57

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
52	ADHEr, PGI, SUCD4	6.33	0.22	30.11	0.00	-6.16	31.28	-20.00	0.00	62.99
53	ADHEr, GLCpts, PGI	6.28	0.26	29.59	0.00	-6.08	30.96	-20.00	0.00	62.41
54	ADHEr, HEX1, PFK	6.28	0.22	30.24	0.00	-6.31	31.41	-20.00	0.00	63.24
55	ADHEr, FBA, HEX1	6.28	0.22	30.24	0.00	-6.31	31.41	-20.00	0.00	63.24
56	ADHEr, HEX1, TPI	6.28	0.22	30.24	0.00	-6.31	31.41	-20.00	0.00	63.24
57	ADHEr, PFK, SUCD4	6.27	0.23	30.17	0.00	-6.30	31.37	-20.00	0.00	63.17
58	ADHEr, FBA, SUCD4	6.27	0.23	30.17	0.00	-6.30	31.37	-20.00	0.00	63.17
59	ADHEr, SUCD4, TPI	6.27	0.23	30.17	0.00	-6.30	31.37	-20.00	0.00	63.17
60	ADHEr, GLUDy, PGI	6.26	0.27	29.44	0.00	-6.06	30.86	-20.00	0.00	62.23
61	ADHEr, GLCpts, TPI	6.21	0.26	29.69	0.00	-6.24	31.07	-20.00	0.00	62.64
62	ADHEr, GLCpts, PFK	6.21	0.26	29.69	0.00	-6.24	31.07	-20.00	0.00	62.64
63	ADHEr, FBA, GLCpts	6.21	0.26	29.69	0.00	-6.24	31.07	-20.00	0.00	62.64
64	ADHEr, PFK, RPE	6.20	0.30	29.01	0.00	-6.02	30.60	-20.00	0.00	61.77
65	ADHEr, FBA, RPE	6.20	0.30	29.01	0.00	-6.02	30.60	-20.00	0.00	61.77
66	ADHEr, RPE, TPI	6.20	0.30	29.01	0.00	-6.02	30.60	-20.00	0.00	61.77
67	ADHEr, GLUDy, TPI	6.19	0.27	29.54	0.00	-6.23	30.98	-20.00	0.00	62.47
68	ADHEr, FBA, GLUD	6.19	0.27	29.54	0.00	-6.23	30.98	-20.00	0.00	62.47
69	ADHEr, GLUDy, PFK	6.19	0.27	29.54	0.00	-6.23	30.98	-20.00	0.00	62.47
70	ADHEr, TAL, TPI	6.17	0.31	29.05	0.00	-6.09	30.65	-20.00	0.00	61.87
71	ADHEr, PFK, TAL	6.17	0.31	29.05	0.00	-6.09	30.65	-20.00	0.00	61.87
72	ADHEr, FBA, TAL	6.17	0.31	29.05	0.00	-6.09	30.65	-20.00	0.00	61.87
73	ADHEr, ATPS4r, GLUDy	6.17	0.55	24.36	0.00	-3.40	27.23	-20.00	0.00	55.50
74	ADHEr, PYK, SUCD4	6.00	0.38	28.04	0.00	-6.05	30.04	-20.00	0.00	60.81
75	ADHEr, GLCpts, SUCD4	5.97	0.40	27.84	0.00	-6.03	29.92	-20.00	0.00	60.59
76	ADHEr, RPE, SUCD4	5.95	0.46	26.74	0.00	-5.68	29.16	-20.00	0.00	59.19
77	ADHEr, FUM, GLUDy	5.94	0.42	27.58	0.00	-6.00	29.76	-20.00	0.00	60.30
78	ADHEr, GLUDy, MDH	5.94	0.42	27.58	0.00	-6.00	29.76	-20.00	0.00	60.30
79	ADHEr, GLUDy, SUCD4	5.94	0.42	27.54	0.00	-5.99	29.74	-20.00	0.00	60.27
80	ADHEr, SUCD4, TAL	5.90	0.46	26.81	0.00	-5.79	29.24	-20.00	0.00	59.36
81	ADHEr, GLCpts, RPE	5.89	0.51	26.14	0.00	-5.59	28.78	-20.00	0.00	58.51
82	ADHEr, GLCpts, GLUDy	5.87	0.46	27.02	0.00	-5.93	29.42	-20.00	0.00	59.69
83	ADHEr, GLCpts, MDH	5.85	0.47	26.83	0.00	-5.91	29.29	-20.00	0.00	59.47
84	ADHEr, FUM, GLCpts	5.85	0.47	26.83	0.00	-5.91	29.29	-20.00	0.00	59.47

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+	
5		85	ADHEr, GLUDy, RPE	5.87	0.52	26.00	0.00	-5.57	28.69	-20.00	0.00	58.35
		86	ADHEr, GLCpts, TAL	5.84	0.51	26.23	0.00	-5.72	28.88	-20.00	0.00	58.71
10		87	ADHEr, FUM, RPE	5.84	0.54	25.70	0.00	-5.53	28.50	-20.00	0.00	58.02
		88	ADHEr, MDH, RPE	5.84	0.54	25.70	0.00	-5.53	28.50	-20.00	0.00	58.02
		89	ADHEr, MDH, PYK	5.82	0.47	26.84	0.00	-5.98	29.28	-20.00	0.00	59.61
15		90	ADHEr, FUM, PYK	5.82	0.47	26.84	0.00	-5.98	29.28	-20.00	0.00	59.61
		91	ADHEr, MDH, TAL	5.79	0.54	25.79	0.00	-5.66	28.61	-20.00	0.00	58.22
		92	ADHEr, FUM, TAL	5.79	0.54	25.79	0.00	-5.66	28.61	-20.00	0.00	58.22
		93	ADHEr, GLUDy, TAL	5.82	0.52	26.09	0.00	-5.70	28.79	-20.00	0.00	58.55
20		94	ADHEr, CBMK2, GLU5K	5.68	0.57	25.51	0.00	-5.76	28.48	-20.00	0.00	58.02
		95	ADHEr, CBMK2, G5SD	5.68	0.57	25.51	0.00	-5.76	28.48	-20.00	0.00	58.02
25		96	ADHEr, ASNS2, CBMK2	5.68	0.57	25.51	0.00	-5.76	28.48	-20.00	0.00	58.02
		97	ADHEr, CBMK2, SO4t2	5.68	0.57	25.50	0.00	-5.75	28.47	-20.00	0.00	58.00
		98	ADHEr, CBMK2, HEX1	5.68	0.57	25.48	0.00	-5.75	28.46	-20.00	0.00	57.99
30		99	ADHEr, EDA, PFLi, PGI	14.76	0.16	16.11	0.00	22.13	0.00	-20.00	0.00	17.24
		100	ADHEr, EDA, NADH6, PGI	14.39	0.13	16.96	0.00	21.96	1.24	-20.00	0.00	19.11
35		101	ADHEr, FRD2, GLUDy, LDH_D	12.91	0.12	12.29	0.00	-0.02	38.75	-20.00	0.00	51.90
		102	ADHEr, FRD2, LDH_D, THD2	12.89	0.13	12.25	0.00	-0.02	38.70	-20.00	0.00	51.85
40		103	ADHEr, ACKr, ACS, PPC	12.62	0.11	12.68	0.00	0.33	39.20	-20.00	0.00	52.67
		104	ADHEr, GLUDy, LDH_ D, PPC	12.60	0.16	11.81	0.00	0.47	38.84	-20.00	0.00	51.79
45		105	ADHEr, LDH_D, PPC, THD2	12.60	0.16	11.72	0.00	0.49	38.81	-20.00	0.00	51.71
		106	ADHEr, ATPS4r, EDA, PGI	11.95	0.15	19.06	0.00	8.00	19.85	-20.00	0.00	39.98
50		107	ADHEr, EDA, GLCpts, PGI	11.95	0.15	19.03	0.00	8.00	19.83	-20.00	0.00	39.94
		108	ADHEr, EDA, GLUDy, PGI	11.86	0.20	18.37	0.00	8.07	19.42	-20.00	0.00	39.21
55		109	ADHEr, GLUDy, HEX1, PGI	11.85	0.20	18.38	0.00	8.05	19.43	-20.00	0.00	39.24
		110	ADHEr, ATPS4r, FRD2, LDH_D	11.72	0.28	7.78	0.00	9.70	25.16	-20.00	0.00	37.14

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+	
5		111	ADHEr, ACKr, NADH6, PYK	11.48	0.56	11.79	0.00	27.19	0.00	-20.00	0.00	15.78
10		112	ADHEr, ACKr, LDH_D, NADH6	11.02	0.64	11.37	0.00	26.49	0.00	-20.00	0.00	15.94
		113	ADHEr, GLCpts, PFLi, PGI	10.95	0.17	26.34	0.00	16.41	0.00	-20.00	0.00	27.55
15		114	ADHEr, GLUDy, PFLi, PGI	10.93	0.18	26.24	0.00	16.38	0.00	-20.00	0.00	27.51
		115	ADHEr, FBA, GLCpts, PFLi	10.91	0.17	26.39	0.00	16.36	0.00	-20.00	0.00	27.62
20		116	ADHEr, GLCpts, PFK, PFLi	10.91	0.17	26.39	0.00	16.36	0.00	-20.00	0.00	27.62
		117	ADHEr, GLCpts, PFLi, TPI	10.91	0.17	26.39	0.00	16.36	0.00	-20.00	0.00	27.62
25		118	ADHEr, FBA, GLUDy, PFLi	10.89	0.18	26.30	0.00	16.33	0.00	-20.00	0.00	27.58
		119	ADHEr, GLUDy, PFK, PFLi	10.89	0.18	26.30	0.00	16.33	0.00	-20.00	0.00	27.58
30		120	ADHEr, GLUDy, PFLi, TPI	10.89	0.18	26.30	0.00	16.33	0.00	-20.00	0.00	27.58
		121	ADHEr, FBA, PFLi, RPE	10.86	0.20	25.97	0.00	16.28	0.00	-20.00	0.00	27.40
35		122	ADHEr, PFLi, RPE, TPI	10.86	0.20	25.97	0.00	16.28	0.00	-20.00	0.00	27.40
		123	ADHEr, PFK, PFLi, RPE	10.86	0.20	25.97	0.00	16.28	0.00	-20.00	0.00	27.40
		124	ADHEr, PFK, PFLi, TAL	10.85	0.20	26.00	0.00	16.25	0.00	-20.00	0.00	27.43
40		125	ADHEr, PFLi, TAL, TPI	10.85	0.20	26.00	0.00	16.25	0.00	-20.00	0.00	27.43
		126	ADHEr, FBA, PFLi, TAL	10.85	0.20	26.00	0.00	16.25	0.00	-20.00	0.00	27.43
		127	ADHEr, ATPS4r, NADH6, PGI	10.64	0.17	23.71	0.00	8.96	13.97	-20.00	0.00	38.90
45		128	ADHEr, GLCpts, PFLi, THD2	10.62	0.42	22.05	0.00	15.90	0.00	-20.00	0.00	25.05
		129	ADHEr, GLUDy, PFLi, THD2	10.59	0.44	21.77	0.00	15.85	0.00	-20.00	0.00	24.89
50		130	ADHEr, LDH_D, PFLi, SUCD4	10.53	0.30	24.78	0.00	15.78	0.00	-20.00	0.00	26.92
		131	ADHEr, LDH_D, NADH6, PFLi	10.31	0.38	23.85	0.00	15.45	0.00	-20.00	0.00	26.52
55		132	ADHEr, ATPS4r, LDH_D, SUCD4	10.30	0.35	21.88	0.00	12.39	8.06	-20.00	0.00	32.44

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+	
5		133	ADHEr, FUM, PFLi, PYK	10.29	0.38	23.75	0.00	15.41	0.00	-20.00	0.00	26.47
10		134	ADHEr, MDH, PFLi, PYK	10.29	0.38	23.75	0.00	15.41	0.00	-20.00	0.00	26.47
		135	ADHEr, GLCpts, GLUDy, PFLi	10.28	0.39	23.71	0.00	15.40	0.00	-20.00	0.00	26.46
15		136	ADHEr, GLCpts, PFLi, RPE	10.21	0.43	23.02	0.00	15.29	0.00	-20.00	0.00	26.07
		137	ADHEr, GLCpts, PFLi, TAL	10.18	0.43	23.09	0.00	15.25	0.00	-20.00	0.00	26.14
20		138	ADHEr, GLUDy, PFLi, RPE	10.16	0.45	22.79	0.00	15.22	0.00	-20.00	0.00	25.96
		139	ADHEr, CBMK2, GLCpts, PFLi	10.16	0.43	23.21	0.00	15.22	0.00	-20.00	0.00	26.24
25		140	ADHEr, LDH_D, MDH, PFLi	10.15	0.43	23.14	0.00	15.19	0.00	-20.00	0.00	26.21
		141	ADHEr, FUM, LDH_D, PFLi	10.15	0.43	23.14	0.00	15.19	0.00	-20.00	0.00	26.21
30		142	ADHEr, GLUDy, PFLi, TAL	10.13	0.45	22.86	0.00	15.17	0.00	-20.00	0.00	26.04
		143	ADHEr, CBMK2, GLUDy, PFLi	10.11	0.44	22.98	0.00	15.14	0.00	-20.00	0.00	26.14
35		144	ADHEr, CBMK2, PFLi, RPE	10.04	0.49	22.20	0.00	15.02	0.00	-20.00	0.00	25.69
		145	ADHEr, CBMK2, PFLi, TAL	10.00	0.49	22.28	0.00	14.97	0.00	-20.00	0.00	25.78
40		146	ADHEr, ASNS2, G5SD, PFLi	9.96	0.49	22.35	0.00	14.91	0.00	-20.00	0.00	25.87
		147	ADHEr, ASNS2, GLU5K, PFLi	9.96	0.49	22.35	0.00	14.91	0.00	-20.00	0.00	25.87
45		148	ADHEr, ATPS4r, GLCpts, MDH	9.96	0.37	21.98	0.00	9.04	11.74	-20.00	0.00	36.36
		149	ADHEr, ATPS4r, FUM, GLCpts	9.96	0.37	21.98	0.00	9.04	11.74	-20.00	0.00	36.36
50		150	ADHEr, ATPS4r, FBA, NADH6	9.87	0.17	27.44	0.00	11.15	7.29	-20.00	0.00	35.96
		151	ADHEr, ATPS4r, NADH6, TPI	9.87	0.17	27.44	0.00	11.15	7.29	-20.00	0.00	35.96
55		152	ADHEr, ATPS4r, NADH6, PFK	9.87	0.17	27.44	0.00	11.15	7.29	-20.00	0.00	35.96
		153	ADHEr, ATPS4r, FUM, PGL	9.82	0.44	23.35	0.00	13.64	2.12	-20.00	0.00	28.60

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5		154 ADHEr, ATPS4r, MDH, PGDH	9.82	0.44	23.35	0.00	13.64	2.12	-20.00	0.00	28.60
10		155 ADHEr, ATPS4r, FUM, G6PDHv	9.82	0.44	23.35	0.00	13.64	2.12	-20.00	0.00	28.60
		156 ADHEr, ATPS4r, FUM, PGDH	9.82	0.44	23.35	0.00	13.64	2.12	-20.00	0.00	28.60
15		157 ADHEr, ATPS4r, MDH, PGL	9.82	0.44	23.35	0.00	13.64	2.12	-20.00	0.00	28.60
		158 ADHEr, ATPS4r, G6PDHy, MDH	9.82	0.44	23.35	0.00	13.64	2.12	-20.00	0.00	28.60
20		159 ADHEr, ATPS4r, LDH_ D, PPC	9.81	0.44	12.15	0.00	1.32	36.78	-20.00	0.00	52.07
		160 ADHEr, ATPS4r, FUM, TAL	9.77	0.44	23.32	0.00	13.29	2.67	-20.00	0.00	29.13
25		161 ADHEr, ATPS4r, MDH, TAL	9.77	0.44	23.32	0.00	13.29	2.67	-20.00	0.00	29.13
		162 ADHEr, ATPS4r, GLCpts, NADH6	9.76	0.42	22.90	0.00	13.62	4.00	-20.00	0.00	29.90
30		163 ADHEr, ATPS4r, MDH, RPE	9.72	0.44	23.30	0.00	12.97	3.18	-20.00	0.00	29.62
		164 ADHEr, ATPS4r, FUM, RPE	9.72	0.44	23.30	0.00	12.97	3.18	-20.00	0.00	29.62
35		165 ADHEr, ATPS4r, NADH6, PGDH	9.39	0.46	23.42	0.00	11.71	4.69	-20.00	0.00	31.40
		166 ADHEr, ATPS4r, NADH6, PGL	9.39	0.46	23.42	0.00	11.71	4.69	-20.00	0.00	31.40
40		167 ADHEr, ATPS4r, G6PDHy, NADH6	9.39	0.46	23.42	0.00	11.71	4.69	-20.00	0.00	31.40
		168 ADHEr, ATPS4r, NADH6, TAL	9.34	0.46	23.40	0.00	11.35	5.25	-20.00	0.00	31.94
45		169 ADHEr, ATPS4r, NADH6, RPE	9.29	0.46	23.37	0.00	11.02	5.76	-20.00	0.00	32.44
		170 ADHEr, G6PDHy, ME2, THD2	9.08	0.43	19.50	0.00	0.35	26.49	-20.00	0.00	49.05
50		171 ADHEr, ME2, PGL, THD2	9.08	0.43	19.50	0.00	0.35	26.49	-20.00	0.00	49.05
		172 ADHEr, G6PDHy, PPC, THD2	8.99	0.47	0.00	0.00	6.58	33.74	-20.00	0.00	45.14
55		173 ADHEr, PGL, PPC, THD2	8.99	0.47	0.00	0.00	6.58	33.74	-20.00	0.00	45.14
		174 ADHEr, ATPS4r, GLUDy, NADH6	8.65	0.43	23.82	0.00	6.11	13.67	-20.00	0.00	40.57

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
175	ADHEr, ACKr, FRD2, LDH_D	8.46	0.64	8.81	0.00	9.05	20.63	-20.00	0.00	37.61
176	ADHEr, ATPS4r, FBA, RPE	8.29	0.26	27.65	0.00	3.98	16.88	-20.00	0.00	46.36
177	ADHEr, ATPS4r, RPE, TPI	8.29	0.26	27.65	0.00	3.98	16.88	-20.00	0.00	46.36
178	ADHEr, ATPS4r, PFK, RPE	8.29	0.26	27.65	0.00	3.98	16.88	-20.00	0.00	46.36
179	ADHEr, ATPS4r, GLUDy, PFK	8.28	0.23	28.10	0.00	3.79	17.22	-20.00	0.00	46.98
180	ADHEr, ATPS4r, GLUDy, TPI	8.28	0.23	28.10	0.00	3.79	17.22	-20.00	0.00	46.98
181	ADHEr, ATPS4r, FBA, GLUDy	8.28	0.23	28.10	0.00	3.79	17.22	-20.00	0.00	46.98
182	ADHEr, ATPS4r, PFK, TAL	8.28	0.26	27.67	0.00	3.98	16.83	-20.00	0.00	46.35
183	ADHEr, ATPS4r, TAL, TPI	8.28	0.26	27.67	0.00	3.98	16.83	-20.00	0.00	46.35
184	ADHEr, ATPS4r, FBA, TAL	8.28	0.26	27.67	0.00	3.98	16.83	-20.00	0.00	46.35
185	ADHEr, ASPT, MDH, PYK	8.16	0.28	23.57	0.00	-4.24	32.94	-20.00	0.00	58.48
186	ADHEr, MDH, PGL, THD2	8.00	0.71	13.36	0.00	7.12	24.68	-20.00	0.00	43.08
187	ADHEr, G6PDHy, MDH, THD2	8.00	0.71	13.36	0.00	7.12	24.68	-20.00	0.00	43.08
188	ADHEr, GLCpts, NADH6, THD2	7.89	0.43	24.60	0.00	2.33	18.97	-20.00	0.00	46.64
189	ADHEr, GLUDy, NADH6, THD2	7.89	0.45	24.31	0.00	2.42	18.77	-20.00	0.00	46.27
190	ADHEr,ASPT,LDH_D, MDH	7.71	0.45	21.65	0.00	-4.12	31.31	-20.00	0.00	56.15
191	ADHEr, GLCpts, NADH6, PGI	7.65	0.19	29.27	0.00	0.15	22.63	-20.00	0.00	53.28
192	ADHEr, LDH_D, SUCD4, THD2	7.64	0.40	24.07	0.00	-1.65	26.18	-20.00	0.00	53.11
193	ADHEr, GLUDy, NADH6, PGI	7.62	0.21	29.06	0.00	0.16	22.53	-20.00	0.00	53.08
194	ADHEr, ACKr, FUM, LDH_D	7.62	0.32	7.79	0.00	-2.23	9.47	-20.00	0.00	37.34
195	ADHEr, FBA, GLCpts, NADH6	7.59	0.20	29.34	0.00	-0.03	22.79	-20.00	0.00	53.53

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
196	ADHEr, GLCpts, NADH6, TPI	7.59	0.20	29.34	0.00	-0.03	22.79	-20.00	0.00	53.53
197	ADHEr, ACKr, LDH_D, MDH	7.62	0.32	7.79	0.00	-2.23	9.47	-20.00	0.00	37.34
198	ADHEr, GLCpts, NADH6, PFK	7.59	0.20	29.34	0.00	-0.03	22.79	-20.00	0.00	53.53
199	ADHEr, GLCpts, MDH, THD2	7.58	0.43	23.76	0.00	-1.67	26.01	-20.00	0.00	52.82
200	ADHEr, FUM, GLCpts, THD2	7.58	0.43	23.76	0.00	-1.67	26.01	-20.00	0.00	52.82
201	ADHEr, NADH6, PFK, RPE	7.57	0.23	28.73	0.00	0.14	22.39	-20.00	0.00	52.79
202	ADHEr, FBA, NADH6, RPE	7.57	0.23	28.73	0.00	0.14	22.39	-20.00	0.00	52.79
203	ADHEr, NADH6, RPE, TPI	7.57	0.23	28.73	0.00	0.14	22.39	-20.00	0.00	52.79
204	ADHEr, GLUDy, NADH6, TPI	7.56	0.21	29.14	0.00	-0.03	22.70	-20.00	0.00	53.35
205	ADHEr, GLUDy, NADH6, PFK	7.56	0.21	29.14	0.00	-0.03	22.70	-20.00	0.00	53.35
206	ADHEr, FBA, GLUDy, NADH6	7.56	0.21	29.14	0.00	-0.03	22.70	-20.00	0.00	53.35
207	ADHEr, NADH6, PFK, TAL	7.54	0.24	28.76	0.00	0.06	22.46	-20.00	0.00	52.90
208	ADHEr, FBA, NADH6, TAL	7.54	0.24	28.76	0.00	0.06	22.46	-20.00	0.00	52.90
209	ADHEr, NADH6, TAL, TPI	7.54	0.24	28.76	0.00	0.06	22.46	-20.00	0.00	52.90
210	ADHEr, ACKr, AKGD, ATPS4r	7.24	0.58	7.55	0.00	52.00	0.00	-20.00	0.00	11.65
211	ADHEr, GLCpts, NADH6, RPE	7.17	0.45	25.76	0.00	0.26	20.93	-20.00	0.00	49.87
212	ADHEr, FUM, GLCpts, NADH6	7.15	0.40	26.55	0.00	-0.06	21.52	-20.00	0.00	50.95
213	ADHEr, GLCpts, MDH, NADH6	7.15	0.40	26.55	0.00	-0.06	21.52	-20.00	0.00	50.95
214	ADHEr, GLCpts, GLUDy, NADH6	7.15	0.41	26.54	0.00	-0.06	21.51	-20.00	0.00	50.94
215	ADHEr, FUM, NADH6, PYK	7.15	0.41	26.54	0.00	-0.06	21.51	-20.00	0.00	50.94
216	ADHEr, MDH, NADH6, PYK	7.15	0.41	26.54	0.00	-0.06	21.51	-20.00	0.00	50.94

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5											
	217	ADHEr, ACKr, ATPS4r, SUCOAS	7.14	0.58	7.46	0.00	16.55	35.80	-20.00	0.00	47.39
10	218	ADHEr, GLUDy, NADH6, RPE	7.14	0.46	25.54	0.00	0.27	20.82	-20.00	0.00	49.66
	219	ADHEr, MDH, NADH6, RPE	7.13	0.47	25.45	0.00	0.27	20.78	-20.00	0.00	49.57
15	220	ADHEr, FUM, NADH6, RPE	7.13	0.47	25.45	0.00	0.27	20.78	-20.00	0.00	49.57
	221	ADHEr, GLCpts, NADH6, TAL	7.12	0.45	25.84	0.00	0.11	21.08	-20.00	0.00	50.11
20	222	ADHEr, GLUDy, NADH6, TAL	7.09	0.46	25.63	0.00	0.11	20.98	-20.00	0.00	49.91
	223	ADHEr, FUM, NADH6, TAL	7.07	0.47	25.53	0.00	0.11	20.93	-20.00	0.00	49.82
25	224	ADHEr, MDH, NADH6, TAL	7.07	0.47	25.53	0.00	0.11	20.93	-20.00	0.00	49.82
	225	ADHEr, CBMK2, GLU5K, NADH6	6.93	0.51	25.11	0.00	-0.07	20.86	-20.00	0.00	49.62
30	226	ADHEr, CBMK2, G5SD, NADH6	6.93	0.51	25.11	0.00	-0.07	20.86	-20.00	0.00	49.62
	227	ADHEr, CBMK2, NADH6, SO4t2	6.93	0.51	25.10	0.00	-0.07	20.86	-20.00	0.00	49.60
35	228	ADHEr, ASNS2, CBMK2, NADH6	6.93	0.51	25.11	0.00	-0.07	20.86	-20.00	0.00	49.61
	229	ADHEr, ATPS4r, PYK, SUCD4	6.69	0.36	19.85	0.00	7.01	7.99	-20.00	0.00	38.27
40	230	ADHEr, GLCpts, PGI, SUCD4	6.40	0.18	30.74	0.00	-6.26	31.69	-20.00	0.00	63.71
	231	ADHEr, FUM, GLUDy, PGI	6.38	0.19	30.56	0.00	-6.23	31.57	-20.00	0.00	63.51
45	232	ADHEr, GLUDy, MDH, PGI	6.38	0.19	30.56	0.00	-6.23	31.57	-20.00	0.00	63.51
	233	ADHEr, GLUDy, PGI, SUCD4	6.37	0.20	30.45	0.00	-6.21	31.50	-20.00	0.00	63.38
50	234	ADHEr, GLCpts, SUCD4, TPI	6.35	0.18	30.80	0.00	-6.37	31.76	-20.00	0.00	63.86
	235	ADHEr, GLCpts, PFK, SUCD4	6.35	0.18	30.80	0.00	-6.37	31.76	-20.00	0.00	63.86
55	236	ADHEr, FBA, GLCpts, SUCD4	6.35	0.18	30.80	0.00	-6.37	31.76	-20.00	0.00	63.86
	237	ADHEr, HEX 1, RPE, TPI	6.32	0.22	30.18	0.00	-6.20	31.34	-20.00	0.00	63.09

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+	
5		238	ADHEr, FBA, HEX1, RPE	6.32	0.22	30.18	0.00	-6.20	31.34	-20.00	0.00	63.09
10		239	ADHEr, HEX1, PFK, RPE	6.32	0.22	30.18	0.00	-6.20	31.34	-20.00	0.00	63.09
		240	ADHEr, FUM, GLUDy, TPI	6.32	0.20	30.62	0.00	-6.35	31.65	-20.00	0.00	63.67
15		241	ADHEr, FBA, GLUDy, MDH	6.32	0.20	30.62	0.00	-6.35	31.65	-20.00	0.00	63.67
		242	ADHEr, FLTM, GLUDy, PFK	6.32	0.20	30.62	0.00	-6.35	31.65	-20.00	0.00	63.67
20		243	ADHEr, GLUDy, MDH, PFK	6.32	0.20	30.62	0.00	-6.35	31.65	-20.00	0.00	63.67
		244	ADHEr, GLUDy, MDH, TPI	6.32	0.20	30.62	0.00	-6.35	31.65	-20.00	0.00	63.67
25		245	ADHEr, FBA, FUM, GLUDy	6.32	0.20	30.62	0.00	-6.35	31.65	-20.00	0.00	63.67
		246	ADHEr, GLCpts, GLUDy, PGI	6.32	0.23	29.99	0.00	-6.14	31.21	-20.00	0.00	62.85
30		247	ADHEr, RPE, SUCD4, TPI	6.32	0.23	30.12	0.00	-6.19	31.30	-20.00	0.00	63.02
		248	ADHEr, FBA, RPE, SUCD4	6.32	0.23	30.12	0.00	-6.19	31.30	-20.00	0.00	63.02
35		249	ADHEr, PFK, RPE, SUCD4	6.32	0.23	30.12	0.00	-6.19	31.30	-20.00	0.00	63.02
		250	ADHEr, GLUDy, HEX1, PFK	6.31	0.20	30.55	0.00	-6.34	31.60	-20.00	0.00	63.59
40		251	ADHEr, GLUDy, HEX1, TPI	6.31	0.20	30.55	0.00	-6.34	31.60	-20.00	0.00	63.59
		252	ADHEr, FBA, GLUDy, HEX1	6.31	0.20	30.55	0.00	-6.34	31.60	-20.00	0.00	63.59
45		253	ADHEr, FBA, GLUDy, SUCD4	6.31	0.20	30.51	0.00	-6.34	31.58	-20.00	0.00	63.55
		254	ADHEr, GLUDy, SUCD4, TPI	6.31	0.20	30.51	0.00	-6.34	31.58	-20.00	0.00	63.55
50		255	ADHEr, GLUDy, PFK, SUCD4	6.31	0.20	30.51	0.00	-6.34	31.58	-20.00	0.00	63.55
		256	ADHEr, FBA, HEX1, TAL	6.30	0.22	30.21	0.00	-6.25	31.37	-20.00	0.00	63.16
55		257	ADHEr, HEX1, PFK, TAL	6.30	0.22	30.21	0.00	-6.25	31.37	-20.00	0.00	63.16
		258	ADHEr, HEX1, TAL, TPI	6.30	0.22	30.21	0.00	-6.25	31.37	-20.00	0.00	63.16

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5											
	259	ADHEr, PFK, SUCD4, TAL	6.29	0.23	30.14	0.00	-6.24	31.33	-20.00	0.00	63.09
10	260	ADHEr, SUCD4, TAL, TPI	6.29	0.23	30.14	0.00	-6.24	31.33	-20.00	0.00	63.09
	261	ADHEr, FBA, SUCD4, TAL	6.29	0.23	30.14	0.00	-6.24	31.33	-20.00	0.00	63.09
15	262	ADHEr, ACKr, LDH_D, SUCD4	6.28	0.22	6.40	0.00	-6.31	7.56	-20.00	0.00	39.41
	263	ADHEr, GLCpts, RPE, TPI	6.26	0.26	29.61	0.00	-6.11	30.98	-20.00	0.00	62.45
20	264	ADHEr, GLCpts, PFK, RPE	6.26	0.26	29.61	0.00	-6.11	30.98	-20.00	0.00	62.45
	265	ADHEr, FBA, GLCpts, RPE	6.26	0.26	29.61	0.00	-6.11	30.98	-20.00	0.00	62.45
25	266	ADHEr, ACT6, LDH_D, MDH	6.26	0.23	0.00	0.00	-6.29	1.22	-20.00	0.00	32.99
	267	ADHEr, GLCpts, GLUDy, TPI	6.26	0.24	30.08	0.00	-6.29	31.31	-20.00	0.00	63.07
30	268	ADHEr, ACT6, FUM, LDH_D	6.26	0.23	0.00	0.00	-6.29	1.22	-20.00	0.00	32.99
	269	ADHEr, FBA, GLCpts, GLUDy	6.26	0.24	30.08	0.00	-6.29	31.31	-20.00	0.00	63.07
35	270	ADHEr, GLCpts, GLUDy, PFK	6.26	0.24	30.08	0.00	-6.29	31.31	-20.00	0.00	63.07
	271	ADHEr, GLUDy, RPE, TPI	6.25	0.27	29.46	0.00	-6.09	30.88	-20.00	0.00	62.28
40	272	ADHEr, FBA, GLUDy, RPE	6.25	0.27	29.46	0.00	-6.09	30.88	-20.00	0.00	62.28
	273	ADHEr, GLUDy, PFK, RPE	6.25	0.27	29.46	0.00	-6.09	30.88	-20.00	0.00	62.28
45	274	ADHEr, GLCpts, TAL, TPI	6.24	0.26	29.65	0.00	-6.17	31.02	-20.00	0.00	62.54
	275	ADHEr, FBA, GLCpts, TAL	6.24	0.26	29.65	0.00	-6.17	31.02	-20.00	0.00	62.54
50	276	ADHEr, GLCpts, PFK, TAL	6.24	0.26	29.65	0.00	-6.17	31.02	-20.00	0.00	62.54
	277	ADHEr, GLUDy, PFK TAL	6.22	0.27	29.50	0.00	-6.15	30.93	-20.00	0.00	62.37
55	278	ADHEr, FBA, GLUDy, TAL	6.22	0.27	29.50	0.00	-6.15	30.93	-20.00	0.00	62.37
	279	ADHEr, GLUDy, TAL, TPI	6.22	0.27	29.50	0.00	-6.15	30.93	-20.00	0.00	62.37

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5		280 ADHEr, GLUDy, MDH, PYK	6.09	0.33	28.74	0.00	-6.13	30.48	-20.00	0.00	61.59
10		281 ADHEr, FUM, GLUDy, PYK	6.09	0.33	28.74	0.00	-6.13	30.48	-20.00	0.00	61.59
		282 ADHEr, PYK, RPE, SUCD4	6.08	0.38	27.93	0.00	-5.86	29.92	-20.00	0.00	60.54
15		283 ADHEr, GLUDy, PYK, SUCD4	6.08	0.34	28.65	0.00	-6.12	30.43	-20.00	0.00	61.49
		284 ADHEr, MDH, PYK, SUCD4	6.06	0.35	28.51	0.00	-6.10	30.34	-20.00	0.00	61.33
20		285 ADHEr, FUM, PYK, SUCD4	6.06	0.35	28.51	0.00	-6.10	30.34	-20.00	0.00	61.33
		286 ADHEr, GLCpts, RPE, SUCD4	6.06	0.40	27.70	0.00	-5.83	29.77	-20.00	0.00	60.29
25		287 ADHEr, GLCpts, GLUDy, MDH	6.05	0.35	28.44	0.00	-6.10	30.30	-20.00	0.00	61.26
		288 ADHEr, FUM, GLCpts, GLUDy	6.05	0.35	28.44	0.00	-6.10	30.30	-20.00	0.00	61.26
30		289 ADHEr, GLCpts, GLUDy, SUCD4	6.04	0.36	28.39	0.00	-6.09	30.27	-20.00	0.00	61.21
		290 ADHEr, PYK, SUCD4, TAL	6.04	0.38	27.98	0.00	-5.95	29.98	-20.00	0.00	60.67
35		291 ADHEr, GLUDy, MDH, RPE	6.03	0.41	27.44	0.00	-5.79	29.60	-20.00	0.00	59.98
		292 ADHEr, FUM, GLUDy, RPE	6.03	0.41	27.44	0.00	-5.79	29.60	-20.00	0.00	59.98
40		293 ADHEr, GLUDy, RPE, SUCD4	6.02	0.42	27.40	0.00	-5.78	29.58	-20.00	0.00	59.94
		294 ADHEr, GLCpts, SUCD4, TAL	6.02	0.40	27.77	0.00	-5.92	29.84	-20.00	0.00	60.43
45		295 ADHEr, GLUDy, MDH, TAL	5.99	0.42	27.50	0.00	-5.89	29.68	-20.00	0.00	60.14
		296 ADHEr, FUM, GLUDy, TAL	5.99	0.42	27.50	0.00	-5.89	29.68	-20.00	0.00	60.14
50		297 ADHEr, GLUDy, SUCD4, TAL	5.98	0.42	27.47	0.00	-5.88	29.66	-20.00	0.00	60.10
		298 ADHEr, GLCpts, GLUDy, RPE	5.96	0.46	26.84	0.00	-5.70	29.23	-20.00	0.00	59.31
55		299 ADHEr, FUM, GLCpts, RPE	5.94	0.47	26.66	0.00	-5.67	29.11	-20.00	0.00	59.10
		300 ADHEr, GLCpts, MDH, RPE	5.94	0.47	26.66	0.00	-5.67	29.11	-20.00	0.00	59.10

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5		301 ADHEr, FUM, LDH_D, SUCD4	5.92	0.43	27.41	0.00	-5.98	29.66	-20.00	0.00	60.12
10		302 ADHEr, LDH_D, MDH, SUCD4	5.92	0.43	27.41	0.00	-5.98	29.66	-20.00	0.00	60.12
		303 ADHEr, GLCpts, GLUDy, TAL	5.92	0.46	26.93	0.00	-5.81	29.32	-20.00	0.00	59.49
15		304 ADHEr, FUM, GLCpts, TAL	5.90	0.47	26.74	0.00	-5.78	29.20	-20.00	0.00	59.28
		305 ADHEr, GLCpts, MDH, TAL	5.90	0.47	26.74	0.00	-5.78	29.20	-20.00	0.00	59.28
20		306 ADHEr, CBMK2, GLU5K, SUCD4	5.86	0.46	26.97	0.00	-5.93	29.38	-20.00	0.00	59.63
		307 ADHEr, CBMK2, G5SD, SUCD4	5.86	0.46	26.97	0.00	-5.93	29.38	-20.00	0.00	59.63
25		308 ADHEr, CBMK2, GLU5K, RPE	5.79	0.56	25.29	0.00	-5.47	28.25	-20.00	0.00	57.55
		309 ADHEr, CBMK2, G5SD, RPE	5.79	0.56	25.29	0.00	-5.47	28.25	-20.00	0.00	57.55
30		310 ADHEr, CBMK2, GLCpts, GLU5K	5.79	0.50	26.42	0.00	-5.86	29.04	-20.00	0.00	59.02
		311 ADHEr, ASNS2, CBMK2, RPE	5.79	0.56	25.29	0.00	-5.47	28.25	-20.00	0.00	57.55
35		312 ADHEr, CBMK2, G5SD, GLCpts	5.79	0.50	26.42	0.00	-5.86	29.04	-20.00	0.00	59.02
		313 ADHEr, ASNS2, CBMK2, GLCpts	5.79	0.50	26.42	0.00	-5.86	29.04	-20.00	0.00	59.02
40		314 ADHEr, CBMK2, GLU5K, TAL	5.74	0.57	25.40	0.00	-5.60	28.36	-20.00	0.00	57.77
		315 ADHEr, CBMK2, G5SD, TAL	5.74	0.57	25.40	0.00	-5.60	28.36	-20.00	0.00	57.77
45		316 ADHEr, ASNS2, CBMK2, TAL	5.74	0.57	25.40	0.00	-5.60	28.36	-20.00	0.00	57.77
		317 ADHEr, CBMK2, GLU5K, MDH	5.74	0.53	25.99	0.00	-5.81	28.77	-20.00	0.00	58.54
50		318 ADHEr, CBMK2, FUM, G5SD	5.74	0.53	25.99	0.00	-5.81	28.77	-20.00	0.00	58.54
		319 ADHEr, CBMK2, G5SD, MDH	5.74	0.53	25.99	0.00	-5.81	28.77	-20.00	0.00	58.54
55		320 ADHEr, CBMK2, FUM, GLU5K	5.74	0.53	25.99	0.00	-5.81	28.77	-20.00	0.00	58.54
		321 ADHEr, ASNS2, CBMK2, MDH	5.74	0.53	25.99	0.00	-5.81	28.77	-20.00	0.00	58.54

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
322	ADHEr, ASNS2, CBMK2, FUM	5.74	0.53	25.99	0.00	-5.81	28.77	-20.00	0.00	58.54
323	ADHEr, ASNS2, GLU5K, SO4t2	5.68	0.57	25.48	0.00	-5.75	28.46	-20.00	0.00	57.99
324	ADHEr, ASPT, LDH_D, MDH, PFLi	14.96	0.23	14.07	0.00	22.42	0.00	-20.00	0.00	15.71
325	ADHEr, EDA, GLUDy, PFLi, PGI	14.81	0.14	16.32	0.00	22.21	0.00	-20.00	0.00	17.33
326	ADHEr, ATPS4r, G6PDHy, GLCpts, MDH	14.63	0.18	15.95	0.00	21.94	0.00	-20.00	0.00	17.25
327	ADHEr, ATPS4r, GLCpts, MDH, PGL	14.63	0.18	15.95	0.00	21.94	0.00	-20.00	0.00	17.25
328	ADHEr, EDA, GLUDy, NADH6, PGI	14.42	0.11	17.14	0.00	21.96	1.32	-20.00	0.00	19.28
329	ADHEr, G6PDHy, LDH_D, PPC, THD2	13.73	0.10	9.55	0.00	1.40	40.38	-20.00	0.00	50.65
330	ADHEr, LDH_D, PGL, PPC, THD2	13.73	0.10	9.55	0.00	1.40	40.38	-20.00	0.00	50.65
331	ADHEr, ATPS4r, FRD2, LDH_D, NADH6	13.37	0.17	4.27	6.42	13.23	18.61	-20.00	0.00	30.48
332	ADHEr, EDA, LDH_D, PPC, THD2	13.09	0.16	10.61	0.00	1.33	38.58	-20.00	0.00	50.31
333	ADHEr, FRD2, GLUDy, LDH_D, RPE	12.96	0.12	12.19	0.00	0.07	38.74	-20.00	0.00	51.78
334	ADHEr, FRD2, LDH_D, RPE, THD2	12.95	0.12	12.13	0.00	0.07	38.68	-20.00	0.00	51.70
335	ADHEr, FRD2, GLUDy, LDH_D, THD2	12.94	0.11	12.36	0.00	-0.02	38.83	-20.00	0.00	52.00
336	ADHEr, FRD2, GLUDy, LDH_D, TAL	12.94	0.12	12.24	0.00	0.03	38.74	-20.00	0.00	51.84
337	ADHEr, FRD2, LDH_D, TAL, THD2	12.92	0.13	12.19	0.00	0.03	38.69	-20.00	0.00	51.77
338	ADHEr, ATPS4r, LDH_D, NADH6, PPC	12.83	0.28	9.13	0.00	9.66	29.15	-20.00	0.00	40.26
339	ADHEr, ASPT, FUM, GLUDy, LDH_D	12.77	0.13	12.57	0.00	-0.13	38.55	-20.00	0.00	52.04
340	ADHEr, ASPT, FUM, LDH_D, THD2	12.75	0.13	12.54	0.00	-0.14	38.50	-20.00	0.00	51.99
341	ADHEr, ME2, PFLi, PGL, THD2	12.69	0.38	17.31	0.00	19.01	0.00	-20.00	0.00	19.99

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5										
	342 ADHEr, G6PDHy, ME2, PFLi, THD2	12.69	0.38	17.31	0.00	19.01	0.00	-20.00	0.00	19.99
10	343 ADHEr, ACKr, ACS, PPC, RPE	12.67	0.10	12.72	0.00	0.38	39.23	-20.00	0.00	52.69
	344 ADHEr, GLUDy, LDH_ D, PPC, RPE	12.67	0.16	11.69	0.00	0.58	38.82	-20.00	0.00	51.63
15	345 ADHEr, LDH_D, PPC, RPE, THD2	12.67	0.16	11.57	0.00	0.60	38.78	-20.00	0.00	51.52
	346 ADHEr, GLUDy, LDH_ D, PPC, TAL	12.64	0.16	11.75	0.00	0.53	38.83	-20.00	0.00	51.71
20	347 ADHEr, LDH_D, PPC, TAL, THD2	12.63	0.16	11.65	0.00	0.55	38.79	-20.00	0.00	51.61
	348 ADHEr, GLCpts, GLUDy, LDH_D, PPC	12.62	0.10	12.80	0.00	0.31	39.24	-20.00	0.00	52.79
25	349 ADHEr, GLCpts, LDH_ D, PPC, THD2	12.62	0.11	12.74	0.00	0.32	39.22	-20.00	0.00	52.73
	350 ADHEr, ACKr, GLUDy, LDH_D, PGI	12.38	0.13	12.45	0.00	9.92	10.79	-20.00	0.00	30.65
30	351 ADHEr, ASPT, ATPS4r, GLCpts, MDH	12.20	0.20	10.35	0.00	18.28	0.00	-20.00	0.00	19.76
	352 ADHEr, LDH_D, PFLi, SUCD4, THD2	12.11	0.25	21.60	0.00	18.15	0.00	-20.00	0.00	23.35
35	353 ADHEr, ACKr, AKGD, ATPS4r, PYK	12.05	0.24	12.18	0.00	12.02	25.48	-20.00	0.00	39.36
	354 ADHEr, ACKr, ATPS4r, PYK, SUCOAS	12.00	0.24	12.13	0.00	12.20	25.51	-20.00	0.00	39.36
40	355 ADHEr, ATPS4r, EDA, GLUDy, PGI	11.98	0.13	19.29	0.00	7.97	19.98	-20.00	0.00	40.22
	356 ADHEr, EDA, GLCpts, GLUDy, PGI	11.98	0.14	19.25	0.00	7.98	19.96	-20.00	0.00	40.18
45	357 ADHEr, ACKr, LDH_D, MDH, SUCD4	11.91	0.20	12.01	3.31	5.46	24.78	-20.00	0.00	41.50
	358 ADHEr, ACKr, GLUDy, NADH6, PYK	11.90	0.49	12.16	0.00	27.81	0.00	-20.00	0.00	15.63
50	359 ADHEr, FUM, LDH_D, PFLi, THD2	11.76	0.39	19.51	0.00	17.62	0.00	-20.00	0.00	22.30
	360 ADHEr, LDH_D, MDH, PFLi, THD2	11.76	0.39	19.51	0.00	17.62	0.00	-20.00	0.00	22.30
55	361 ADHEr, Act6, ATPS4r, LDH_D, PPC	11.67	0.14	0.00	0.00	1.15	24.89	-19.46	0.00	38.65
	362 ADHEr, ACKr, LDH_D, PYRt2, SUCD4	11.66	0.21	11.78	0.00	8.63	21.96	-20.00	0.00	37.33

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5	363 ADHEr, ATPS4r, FUM, LDH_D, SUCD4	11.59	0.23	20.54	0.00	13.84	9.06	-20.00	0.00	31.25
10	364 ADHEr, ATPS4r, LDH_ D, MDH, SUCD4	11.59	0.23	20.54	0.00	13.84	9.06	-20.00	0.00	31.25
	365 ADHEr, ACKr, CBMK2, NADH6, PYK	11.52	0.55	11.82	0.00	27.24	0.00	-20.00	0.00	15.76
15	366 ADHEr, ACKr, NADH6, PYK, RPE	11.50	0.56	11.81	0.00	27.22	0.00	-20.00	0.00	15.77
	367 ADHEr, ACKr, ASNS2, NADH6, PYK	11.50	0.56	11.80	0.00	27.21	0.00	-20.00	0.00	15.77
20	368 ADHEr, ACKr, NADH6, PYK, TAL	11.49	0.56	11.80	0.00	27.20	0.00	-20.00	0.00	15.77
	369 ADHEr, ASPT, ATPS4r, FUM, LDH_D	11.46	0.30	8.96	0.00	8.56	25.90	-20.00	0.00	38.85
25	370 ADHEr, ASPT, ATPS4r, LDH_D, MDH	11.43	0.34	17.13	0.00	8.27	17.71	-20.00	0.00	37.24
	371 ADHEr, MDH, PFLi, PYK, THD2	11.38	0.38	20.90	0.00	17.05	0.00	-20.00	0.00	23.58
30	372 ADHEr, FUM, PFLi, PYK, THD2	11.38	0.38	20.90	0.00	17.05	0.00	-20.00	0.00	23.58
	373 ADHEr, ACKr, FUM, LDH_D, NADH6	11.25	0.53	11.54	0.00	23.26	0.00	-20.00	0.00	17.50
35	374 ADHEr, ATPS4r, G6PDHy, LDH_D, SUCD4	11.24	0.27	20.72	0.00	13.49	8.70	-20.00	0.00	31.38
	375 ADHEr, ATPS4r, LDH_ D, PGL, SUCD4	11.24	0.27	20.72	0.00	13.49	8.70	-20.00	0.00	31.38
40	376 ADHEr, ATPS4r, LDH_ D, PGDH, SUCD4	11.24	0.27	20.72	0.00	13.49	8.70	-20.00	0.00	31.38
	377 ADHEr, GLYCL, PGL, PPC, THD2	11.24	0.14	2.95	0.00	3.54	28.60	-20.00	0.02	42.95
45	378 ADHEr, G6PDHy, GLYCL, PPC, THD2	11.24	0.14	2.95	0.00	3.54	28.60	-20.00	0.02	42.95
	379 ADHEr, FTHFD, G6PDHy, PPC, THD2	11.22	0.14	2.94	0.00	3.54	28.55	-20.00	0.00	42.92
50	380 ADHEr, FTHFD, PGL, PPC, THD2	11.22	0.14	2.94	0.00	3.54	28.55	-20.00	0.00	42.92
	381 ADHEr, MTHFC, PGL, PPC, THD2	11.22	0.14	2.94	0.00	3.54	28.55	-20.00	0.00	42.92
55	382 ADHEr, G6PDHy, MTHFC, PPC, THD2	11.22	0.14	2.94	0.00	3.54	28.55	-20.00	0.00	42.92

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
5	383 ADHEr, ATPS4r, LDH_ D, PFLi, SUCD4	11.20	0.32	22.07	0.00	17.78	0.00	-20.00	0.00	24.34
10	384 ADHEr, ATPS4r, LDH_ D, SUCD4, TAL	11.18	0.28	20.84	0.00	13.41	8.69	-20.00	0.00	31.49
	385 ADHEr, ATPS4r, LDH_ D, RPE, SUCD4	11.13	0.28	23.11	0.00	17.68	0.00	-20.00	0.00	25.09
15	386 ADHEr, ATPS4r, FUM, GLCpts, NADH6	11.13	0.27	19.49	0.00	16.68	0.00	-20.00	0.00	24.27
	387 ADHEr, ATPS4r, GLCpts, MDH, NADH6	11.13	0.27	19.49	0.00	16.68	0.00	-20.00	0.00	24.27
20	388 ADHEr, ATPS4r, LDH_ D, NADH6, PFLi	11.12	0.34	22.41	0.00	16.65	0.00	-20.00	0.00	24.82
	389 ADHEr, ACKr, MAL5, NADH12, NADH6	11.06	0.63	11.41	0.00	26.56	0.00	-20.00	0.00	15.92
25	390 ADHEr, ACKr, ICL, NADH12, NADH6	11.06	0.63	11.41	0.00	26.56	0.00	-20.00	0.00	15.92
	391 ADHEr, ACKr, CBMK2, LDH_D, NADH6	11.06	0.64	11.40	0.00	26.54	0.00	-20.00	0.00	15.93
30	392 ADHEr, ATPS4r, GLCpts, MDH, PFLi	11.04	0.35	22.48	0.00	16.54	0.00	-20.00	0.00	24.94
	393 ADHEr, ATPS4r, FUM, GLCpts, PFLi	11.04	0.35	22.48	0.00	16.54	0.00	-20.00	0.00	24.94
35	394 ADHEr, ACKr, ASNS2, LDH_D, NADH6	11.03	0.64	11.38	0.00	26.51	0.00	-20.00	0.00	15.93
	395 ADHEr, ACKr, ATPS4r, LDH_D, THD2	11.00	0.38	11.21	0.00	5.04	32.88	-20.00	0.00	46.77
40	396 ADHEr, GLCpts, GLUDy, PFLi, PGI	11.00	0.15	26.58	0.00	16.49	0.00	-20.00	0.00	27.66
	397 ADHEr, GLCpts, GLUDy, PFK, PFLi	10.97	0.15	26.63	0.00	16.45	0.00	-20.00	0.00	27.73
45	398 ADHEr, FBA, GLCpts, GLUDy, PFLi	10.97	0.15	26.63	0.00	16.45	0.00	-20.00	0.00	27.73
	399 ADHEr, GLCpts, GLUDy, PFLi, TPI	10.97	0.15	26.63	0.00	16.45	0.00	-20.00	0.00	27.73
50	400 ADHEr, ATPS4r, GLCpts, NADH6, PGI	10.97	0.11	27.23	0.00	15.95	0.99	-20.00	0.00	29.02
	401 ADHEr, ATPS4r, GLCpts, NADH6, TPI	10.95	0.12	27.26	0.00	15.95	0.94	-20.00	0.00	29.02
55	402 ADHEr, ATPS4r, GLCpts, NADH6, PFK	10.95	0.12	27.26	0.00	15.95	0.94	-20.00	0.00	29.02
	403 ADHEr, ATPS4r, FBA, GLCpts, NADH6	10.95	0.12	27.26	0.00	15.95	0.94	-20.00	0.00	29.02

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	BDO	BIO	AC	ALA	CO2	FOR	GLC	GLY	H+
404	ADHEr, GLCpts, PFLi, RPE, TPI	10.94	0.17	26.35	0.00	16.40	0.00	-20.00	0.00	27.57
405	ADHEr, GLCpts, PFK, PFLi, RPE	10.94	0.17	26.35	0.00	16.40	0.00	-20.00	0.00	27.57
406	ADHEr, FBA, GLCpts, PFLi, RPE	10.94	0.17	26.35	0.00	16.40	0.00	-20.00	0.00	27.57
407	ADHEr, ATPS4r, NADH6, PFLi, PYK	10.93	0.28	24.13	0.00	16.38	0.00	-20.00	0.00	26.12
408	ADHEr, GLCpts, PFK, PFLi, TAL	10.93	0.17	26.37	0.00	16.38	0.00	-20.00	0.00	27.59
409	ADHEr, FBA, GLCpts, PFLi, TAL	10.93	0.17	26.37	0.00	16.38	0.00	-20.00	0.00	27.59
410	ADHEr, GLCpts, PFLi, TAL, TPI	10.93	0.17	26.37	0.00	16.38	0.00	-20.00	0.00	27.59
411	ADHEr, FBA, GLUDy, PFLi, RPE	10.92	0.18	26.25	0.00	16.37	0.00	-20.00	0.00	27.52
412	ADHEr, GLUDy, PFK, PFLi, RPE	10.92	0.18	26.25	0.00	16.37	0.00	-20.00	0.00	27.52
413	ADHEr, GLUDy, PFLi, RPE, TPI	10.92	0.18	26.25	0.00	16.37	0.00	-20.00	0.00	27.52
414	ADHEr, GLUDy, PFLi, TAL, TPI	10.91	0.18	26.27	0.00	16.35	0.00	-20.00	0.00	27.55
415	ADHEr, GLUDy, PFK, PFLi, TAL	10.91	0.18	26.27	0.00	16.35	0.00	-20.00	0.00	27.55
416	ADHEr, FBA, GLUDy, PFLi, TAL	10.91	0.18	26.27	0.00	16.35	0.00	-20.00	0.00	27.55
417	ADHEr, ATPS4r, LDH_D, NADH6, SUCD4	10.79	0.32	21.77	0.00	14.53	5.29	-20.00	0.00	29.37
418	ADHEr, FUM, LDH_D, PFLi, SUCD4	10.76	0.23	25.74	0.00	16.13	0.00	-20.00	0.00	27.34
419	ADHEr, LDH_D, MDH, PFLi, SUCD4	10.76	0.23	25.74	0.00	16.13	0.00	-20.00	0.00	27.34
	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
1	ADHEr	-0.90	0.00	-4.96	0.00	-0.61	0.00	-0.10	0.00	0.00
2	ADHEr, PFLi	14.07	0.00	-4.31	0.00	-0.53	0.00	-0.09	0.00	0.00
3	ADHEr, NADH6	2.51	0.00	-4.48	0.00	-0.55	0.00	-0.09	0.00	0.00
4	ADHEr, THD2	0.13	0.00	-4.83	0.00	-0.60	0.00	-0.10	0.00	0.00
5	ADHEr, PGI	-6.65	0.00	-2.62	0.00	-0.32	0.00	-0.05	0.00	0.00

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	6	ADHEr, PFK	-6.68	0.00	-2.66	0.00	-0.33	0.00	-0.05	0.00	0.00
	7	ADHEr, FBA	-6.68	0.00	-2.66	0.00	-0.33	0.00	-0.05	0.00	0.00
10	8	ADHEr, ATPS4r	1.88	0.00	-5.25	-2.00	-0.65	0.00	-0.11	0.00	0.00
	9	ADHEr, TPI	-6.68	0.00	-2.66	0.00	-0.33	0.00	-0.05	0.00	0.00
	10	ADHEr, SUCD4	-3.22	0.00	-4.04	0.00	-0.50	0.00	-0.08	0.00	0.00
15	11	ADHEr, RPE	-0.76	0.00	-4.95	0.00	-0.61	0.00	-0.10	0.00	0.00
	12	ADHEr, GLCpts	-2.34	0.00	-4.39	0.00	-0.54	0.00	-0.09	0.00	0.00
	13	ADHEr, GLUDy	-2.13	0.00	-4.47	0.00	-0.55	0.00	-0.09	0.00	0.00
	14	ADHEr, TAL	-0.83	0.00	-4.95	0.00	-0.61	0.00	-0.10	0.00	0.00
20	15	ADHEr, MDH	-1.66	0.00	-4.66	0.00	-0.58	0.00	-0.09	0.00	0.00
	16	ADHEr, FUM	-1.66	0.00	-4.66	0.00	-0.58	0.00	-0.09	0.00	0.00
	17	ADHEr, CBMK2	-1.01	0.00	-4.92	0.00	-0.61	0.00	-0.10	0.00	0.00
25	18	ADHEr, HEX1, PGI	0.36	0.00	-1.93	0.00	-0.24	0.00	-0.04	0.00	0.00
	19	ADHEr, EDA, PGI	0.38	0.00	-1.94	0.00	-0.24	0.00	-0.04	0.00	0.00
	20	ADHEr, PFLi, PGI	9.09	0.00	-1.73	0.00	-0.21	0.00	-0.03	0.00	0.00
	21	ADHEr, FBA, PFLi	9.11	0.00	-1.75	0.00	-0.22	0.00	-0.04	0.00	0.00
30	22	ADHEr, PFLi, TPI	9.11	0.00	-1.75	0.00	-0.22	0.00	-0.04	0.00	0.00
	23	ADHEr, PFK, PFLi	9.11	0.00	-1.75	0.00	-0.22	0.00	-0.04	0.00	0.00
	24	ADHEr, PFLi, THD2	14.16	0.00	-4.23	0.00	-0.52	0.00	-0.09	0.00	0.00
35	25	ADHEr, GLCpts, PFLi	12.93	0.00	-3.72	0.00	-0.46	0.00	-0.08	0.00	0.00
	26	ADHEr, GLUDy, PFLi	13.23	0.00	-3.87	0.00	-0.48	0.00	-0.08	0.00	0.00
	27	ADHEr, PFLi, RPE	14.08	0.00	-4.30	0.00	-0.53	0.00	-0.09	0.00	0.00
	28	ADHEr, PFLi, TAL	14.07	0.00	-4.30	0.00	-0.53	0.00	-0.09	0.00	0.00
40	29	ADHEr, CBMK2, PFLi	13.99	0.00	-4.27	0.00	-0.53	0.00	-0.09	0.00	0.00
	30	ADHEr, ATPS4r, PGI	-3.19	0.00	-2.28	0.00	-0.28	0.00	-0.05	0.00	0.00
	31	ADHEr, ATPS4r, NADH6	6.46	0.00	-4.15	0.00	-0.51	0.00	-0.08	0.00	0.00
45	32	ADHEr, ATPS4r, TPI	0.48	0.00	-2.25	0.00	-0.28	0.00	-0.05	0.00	0.00
	33	ADHEr, ATPS4r, PFK	0.48	0.00	-2.25	0.00	-0.28	0.00	-0.05	0.00	0.00
	34	ADHEr, ATPS4r, FBA	0.48	0.00	-2.25	0.00	-0.28	0.00	-0.05	0.00	0.00
50	35	ADHEr, NADH6, THD2	3.90	0.00	-4.30	0.00	-0.53	0.00	-0.09	0.00	0.00
	36	ADHEr, ATPS4r, FUM	1.11	0.00	-4.22	0.00	-0.52	0.00	-0.09	0.00	0.00
	37	ADHEr, ATPS4r, MDH	1.11	0.00	-4.22	0.00	-0.52	0.00	-0.09	0.00	0.00
55	38	ADHEr, NADH6, PGI	-3.12	0.00	-2.02	0.00	-0.25	0.00	-0.04	0.00	0.00
	39	ADHEr, FUM, THD2	0.18	0.00	-4.28	0.00	-0.53	0.00	-0.09	0.00	0.00
	40	ADHEr, MDH, THD2	0.18	0.00	-4.28	0.00	-0.53	0.00	-0.09	0.00	0.00

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	41	ADHEr, NADH6, PFK	-3.17	0.00	-2.06	0.00	-0.25	0.00	-0.04	0.00	0.00
	42	ADHEr, FBA, NADH6	-3.17	0.00	-2.06	0.00	-0.25	0.00	-0.04	0.00	0.00
10	43	ADHEr, NADH6, TPI	-3.17	0.00	-2.06	0.00	-0.25	0.00	-0.04	0.00	0.00
	44	ADHEr, GLCpts, NADH6	1.14	0.00	-3.90	0.00	-0.48	0.00	-0.08	0.00	0.00
	45	ADHEr, NADH6, RPE	2.70	0.00	-4.46	0.00	-0.55	0.00	-0.09	0.00	0.00
15	46	ADHEr, GLUDy, NADH6	1.46	0.00	-4.03	0.00	-0.50	0.00	-0.08	0.00	0.00
	47	ADHEr, FUM, NADH6	1.61	0.00	-4.10	0.00	-0.51	0.00	-0.08	0.00	0.00
	48	ADHEr, MDH, NADH6	1.61	0.00	-4.10	0.00	-0.51	0.00	-0.08	0.00	0.00
20	49	ADHEr, NADH6, TAL	2.61	0.00	-4.47	0.00	-0.55	0.00	-0.09	0.00	0.00
	50	ADHEr, GLCpts, THD2	-1.43	0.00	-4.27	0.00	-0.53	0.00	-0.09	0.00	0.00
	51	ADHEr, GLUDy, THD2	-1.17	0.00	-4.36	0.00	-0.54	0.00	-0.09	0.00	0.00
25	52	ADHEr, PGI, SUCD4	-8.37	0.00	-1.95	0.00	-0.24	0.00	-0.04	0.00	0.00
	53	ADHEr, GLCpts, PGI	-7.58	0.00	-2.25	0.00	-0.28	0.00	-0.05	0.00	0.00
	54	ADHEr, HEX1, PFK	-8.47	0.00	-1.94	0.00	-0.24	0.00	-0.04	0.00	0.00
30	55	ADHEr, FBA, HEX1	-8.47	0.00	-1.94	0.00	-0.24	0.00	-0.04	0.00	0.00
	56	ADHEr, HEX1, TPI	-8.47	0.00	-1.94	0.00	-0.24	0.00	-0.04	0.00	0.00
	57	ADHEr, PFK, SUCD4	-8.37	0.00	-1.98	0.00	-0.25	0.00	-0.04	0.00	0.00
	58	ADHEr, FBA, SUCD4	-8.37	0.00	-1.98	0.00	-0.25	0.00	-0.04	0.00	0.00
35	59	ADHEr, SUCD4, TPI	-8.37	0.00	-1.98	0.00	-0.25	0.00	-0.04	0.00	0.00
	60	ADHEr, GLUDy, PGI	-7.35	0.00	-2.35	0.00	-0.29	0.00	-0.05	0.00	0.00
	61	ADHEr, GLCpts, TPI	-7.61	0.00	-2.29	0.00	-0.28	0.00	-0.05	0.00	0.00
40	62	ADHEr, GLCpts, PFK	-7.61	0.00	-2.29	0.00	-0.28	0.00	-0.05	0.00	0.00
	63	ADHEr, FBA, GLCpts	-7.61	0.00	-2.29	0.00	-0.28	0.00	-0.05	0.00	0.00
	64	ADHEr, PFK, RPE	-6.66	0.00	-2.63	0.00	-0.32	0.00	-0.05	0.00	0.00
	65	ADHEr, FBA, RPE	-6.66	0.00	-2.63	0.00	-0.32	0.00	-0.05	0.00	0.00
45	66	ADHEr, RPE, TPI	-6.66	0.00	-2.63	0.00	-0.32	0.00	-0.05	0.00	0.00
	67	ADHEr, GLUDy, TPI	-7.38	0.00	-2.38	0.00	-0.29	0.00	-0.05	0.00	0.00
	68	ADHEr, FBA, GLUDy	-7.38	0.00	-2.38	0.00	-0.29	0.00	-0.05	0.00	0.00
50	69	ADHEr, GLUDy, PFK	-7.38	0.00	-2.38	0.00	-0.29	0.00	-0.05	0.00	0.00
	70	ADHEr, TAL, TPI	-6.67	0.00	-2.64	0.00	-0.33	0.00	-0.05	0.00	0.00
	71	ADHEr, PFK, TAL	-6.67	0.00	-2.64	0.00	-0.33	0.00	-0.05	0.00	0.00
	72	ADHEr, FBA, TAL	-6.67	0.00	-2.64	0.00	-0.33	0.00	-0.05	0.00	0.00
55	73	ADHEr, ATPS4r, GLUDy	0.48	0.00	-4.75	-2.00	-0.59	0.00	-0.10	0.00	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
74	ADHEr, PYK, SUCD4	-5.01	0.00	-3.32	0.00	-0.41	0.00	-0.07	0.00	0.00
75	ADHEr, GLCpts, SUCD4	-4.70	0.00	-3.45	0.00	-0.43	0.00	-0.07	0.00	0.00
76	ADHEr, RPE, SUCD4	-3.14	0.00	-4.01	0.00	-0.50	0.00	-0.08	0.00	0.00
77	ADHEr, FUM, GLUDy	-4.29	0.00	-3.61	0.00	-0.45	0.00	-0.07	0.00	0.00
78	ADHEr, GLUDy, MDH	-4.29	0.00	-3.61	0.00	-0.45	0.00	-0.07	0.00	0.00
79	ADHEr, GLUDy, SUCD4	-4.24	0.00	-3.63	0.00	-0.45	0.00	-0.07	0.00	0.00
80	ADHEr, SUCD4, TAL	-3.18	0.00	-4.02	0.00	-0.50	0.00	-0.08	0.00	0.00
81	ADHEr, GLCpts, RPE	-2.22	0.00	-4.37	0.00	-0.54	0.00	-0.09	0.00	0.00
82	ADHEr, GLCpts, GLUDy	-3.42	0.00	-3.96	0.00	-0.49	0.00	-0.08	0.00	0.00
83	ADHEr, GLCpts, MDH	-3.12	0.00	-4.08	0.00	-0.50	0.00	-0.08	0.00	0.00
84	ADHEr, FUM, GLCpts	-3.12	0.00	-4.08	0.00	-0.50	0.00	-0.08	0.00	0.00
85	ADHEr, GLUDy, RPE	-2.00	0.00	-4.46	0.00	-0.55	0.00	-0.09	0.00	0.00
86	ADHEr, GLCpts, TAL	-2.28	0.00	-4.38	0.00	-0.54	0.00	-0.09	0.00	0.00
87	ADHEr, FUM, RPE	-1.54	0.00	-4.64	0.00	-0.57	0.00	-0.09	0.00	0.00
88	ADHEr, MDH, RPE	-1.54	0.00	-4.64	0.00	-0.57	0.00	-0.09	0.00	0.00
89	ADHEr, MDH, PYK	-3.17	0.00	-4.03	0.00	-0.50	0.00	-0.08	0.09	0.00
90	ADHEr, FUM, PYK	-3.17	0.00	-4.03	0.00	-0.50	0.00	-0.08	0.09	0.00
91	ADHEr, MDH, TAL	-1.60	0.00	-4.65	0.00	-0.58	0.00	-0.09	0.00	0.00
92	ADHEr, FUM, TAL	-1.60	0.00	-4.65	0.00	-0.58	0.00	-0.09	0.00	0.00
93	ADHEr, GLUDy, TAL	-2.06	0.00	-4.47	0.00	-0.55	0.00	-0.09	0.00	0.00
94	ADHEr, CBMK2, GLU5K	-1.05	0.00	-4.90	0.00	-0.61	0.00	-0.10	0.00	0.00
95	ADHEr, CBMK2, G5SD	-1.05	0.00	-4.90	0.00	-0.61	0.00	-0.10	0.00	0.00
96	ADHEr, ASNS2, CBMK2	-1.05	0.00	-4.90	0.00	-0.61	0.00	-0.10	0.00	0.00
97	ADHEr, CBMK2, SO4t2	-1.03	0.00	-4.91	0.00	-0.61	0.00	-0.10	0.00	0.00
98	ADHEr, CBMK2, HEX1	-1.01	0.00	-4.92	0.00	-0.61	0.00	-0.10	0.00	0.00
99	ADHEr, EDA, PFLi, PGI	10.28	0.00	-1.38	0.00	-0.17	0.00	-0.03	0.00	0.00
100	ADHEr, EDA, NADH6, PGI	9.91	0.00	-1.11	-2.00	-0.14	0.00	-0.02	0.00	0.00
101	ADHEr, FRD2, GLUDy, LDH_D	-10.70	0.00	-1.05	0.00	-0.13	0.00	-0.02	0.00	0.00
102	ADHEr, FRD2, LDH_D, THD2	-10.60	0.00	-1.09	0.00	-0.13	0.00	-0.02	0.00	0.00

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5											
	103	ADHEr, ACKr, ACS, PPC	-10.25	0.00	-0.96	-2.00	-0.12	0.00	-0.02	0.00	0.00
10	104	ADHEr, GLUDy, LDH_ D, PPC	-9.20	0.00	-1.38	-2.00	-0.17	0.00	-0.03	0.00	0.00
	105	ADHEr, LDH_D, PPC, THD2	-9.09	0.00	-1.43	-2.00	-0.18	0.00	-0.03	0.00	0.00
15	106	ADHEr, ATPS4r, EDA, PGI	-1.21	0.00	-1.30	0.00	-0.16	0.00	-0.03	0.00	0.00
	107	ADHEr, EDA, GLCpts, PGI	-1.16	0.00	-1.32	0.00	-0.16	0.00	-0.03	0.00	0.00
20	108	ADHEr, EDA, GLUDy, PGI	-0.12	0.00	-1.73	0.00	-0.21	0.00	-0.04	0.00	0.00
	109	ADHEr, GLUDy, HEX1, PGI	-0.13	0.00	-1.74	0.00	-0.21	0.00	-0.04	0.00	0.00
25	110	ADHEr, ATPS4r, FRD2, LDH_D	5.32	0.00	-4.64	-5.00	-0.30	0.00	-0.05	0.00	2.22
	111	ADHEr, ACKr, NADH6, PYK	25.99	0.00	-4.86	-20.00	-0.60	0.00	-0.10	0.00	0.00
30	112	ADHEr, ACKr, LDH_D, NADH6	27.24	0.00	-5.56	-20.00	-0.69	0.00	-0.11	0.00	0.00
	113	ADHEr, GLCpts, PFLi, PGI	8.60	0.00	-1.48	0.00	-0.18	0.00	-0.03	0.00	0.00
35	114	ADHEr, GLUDy, PFLi, PGI	8.73	0.00	-1.55	0.00	-0.19	0.00	-0.03	0.00	0.00
	115	ADHEr, FBA, GLCpts, PFLi	8.62	0.00	-1.50	0.00	-0.19	0.00	-0.03	0.00	0.00
40	116	ADHEr, GLCpts, PFK, PFLi	8.62	0.00	-1.50	0.00	-0.19	0.00	-0.03	0.00	0.00
	117	ADHEr, GLCpts, PFLi, TPI	8.62	0.00	-1.50	0.00	-0.19	0.00	-0.03	0.00	0.00
45	118	ADHEr, FBA, GLUDy, PFLi	8.75	0.00	-1.56	0.00	-0.19	0.00	-0.03	0.00	0.00
	119	ADHEr, GLUDy, PFK, PFLi	8.75	0.00	-1.56	0.00	-0.19	0.00	-0.03	0.00	0.00
50	120	ADHEr, GLUDy, PFLi, TPI	8.75	0.00	-1.56	0.00	-0.19	0.00	-0.03	0.00	0.00
	121	ADHEr, FBA, PFLi, RPE	9.09	0.00	-1.74	0.00	-0.21	0.00	-0.04	0.00	0.00
55	122	ADHEr, PFLi, RPE, TPI	9.09	0.00	-1.74	0.00	-0.21	0.00	-0.04	0.00	0.00
	123	ADHEr, PFK, PFLi, RPE	9.09	0.00	-1.74	0.00	-0.21	0.00	-0.04	0.00	0.00
	124	ADHEr, PFK, PFLi, TAL	9.10	0.00	-1.74	0.00	-0.22	0.00	-0.04	0.00	0.00

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL	
5		125	ADHEr, PFLi, TAL, TPI	9.10	0.00	-1.74	0.00	-0.22	0.00	-0.04	0.00	0.00
		126	ADHEr, FBA, PFLi, TAL	9.10	0.00	-1.74	0.00	-0.22	0.00	-0.04	0.00	0.00
10		127	ADHEr, ATPS4r, NADH6, PGI	1.46	0.00	-1.48	0.00	-0.18	0.00	-0.03	0.00	0.00
		128	ADHEr, GLCpts, PFLi, THD2	13.01	0.00	-3.65	0.00	-0.45	0.00	-0.07	0.00	0.00
15		129	ADHEr, GLUDy, PFLi, THD2	13.33	0.00	-3.81	0.00	-0.47	0.00	-0.08	0.00	0.00
		130	ADHEr, LDH_D, PFLi, SUCD4	10.77	0.00	-2.61	0.00	-0.32	0.00	-0.05	0.00	0.00
20		131	ADHEr, LDH_D, NADH6, PFLi	12.01	0.00	-3.25	0.00	-0.40	0.00	-0.07	0.00	0.00
		132	ADHEr, ATPS4r, LDH_ D, SUCD4	8.53	0.00	-3.04	-2.00	-0.38	0.00	-0.06	0.00	0.00
25		133	ADHEr, FUM, PFLi, PYK	12.15	0.00	-3.32	0.00	-0.41	0.00	-0.07	0.00	0.00
		134	ADHEr, MDH, PFLi, PYK	12.15	0.00	-3.32	0.00	-0.41	0.00	-0.07	0.00	0.00
30		135	ADHEr, GLCpts, GLUDy, PFLi	12.20	0.00	-3.35	0.00	-0.41	0.00	-0.07	0.00	0.00
		136	ADHEr, GLCpts, PFLi, RPE	12.94	0.00	-3.71	0.00	-0.46	0.00	-0.08	0.00	0.00
35		137	ADHEr, GLCpts, PFLi, TAL	12.93	0.00	-3.72	0.00	-0.46	0.00	-0.08	0.00	0.00
		138	ADHEr, GLUDy, PFLi, RPE	13.24	0.00	-3.87	0.00	-0.48	0.00	-0.08	0.00	0.00
40		139	ADHEr, CBMK2, GLCpts, PFLi	12.87	0.00	-3.69	0.00	-0.46	0.00	-0.07	0.00	0.00
		140	ADHEr, LDH_D, MDH, PFLi	12.95	0.00	-3.73	0.00	-0.46	0.00	-0.08	0.00	0.00
45		141	ADHEr, FUM, LDH_D, PFLi	12.95	0.00	-3.73	0.00	-0.46	0.00	-0.08	0.00	0.00
		142	ADHEr, GLUDy, PFLi, TAL	13.23	0.00	-3.87	0.00	-0.48	0.00	-0.08	0.00	0.00
50		143	ADHEr, CBMK2, GLUDy, PFLi	13.17	0.00	-3.84	0.00	-0.48	0.00	-0.08	0.00	0.00
		144	ADHEr, CBMK2, PFLi, RPE	14.01	0.00	-4.26	0.00	-0.53	0.00	-0.09	0.00	0.00
55		145	ADHEr, CBMK2, PFLi, TAL	14.00	0.00	-4.26	0.00	-0.53	0.00	-0.09	0.00	0.00
		146	ADHEr, ASNS2, G5SD, PFLi	14.01	0.00	-4.28	0.00	-0.53	0.00	-0.09	0.00	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	147	ADHEr, ASNS2, GLU5K, PFLi	14.01	0.00	-4.28	0.00	-0.53	0.00	-0.09	0.00
10	148	ADHEr, ATPS4r, GLCpts, MDH	5.89	0.00	-3.22	0.00	-0.40	0.00	-0.07	0.00
	149	ADHEr, ATPS4r, FUM, GLCpts	5.89	0.00	-3.22	0.00	-0.40	0.00	-0.07	0.00
15	150	ADHEr, ATPS4r, FBA, NADH6	4.45	0.00	-1.50	0.00	-0.19	0.00	-0.03	0.00
	151	ADHEr, ATPS4r, NADH6, TPI	4.45	0.00	-1.50	0.00	-0.19	0.00	-0.03	0.00
20	152	ADHEr, ATPS4r, NADH6, PFK	4.45	0.00	-1.50	0.00	-0.19	0.00	-0.03	0.00
	153	ADHEr, ATPS4r, FUM, PGL	11.87	0.00	-3.80	0.00	-0.47	0.00	-0.08	0.00
25	154	ADHEr, ATPS4r, MDH, PGDH	11.87	0.00	-3.80	0.00	-0.47	0.00	-0.08	0.00
	155	ADHEr, ATPS4r, FUM, G6PDH	11.87	0.00	-3.80	0.00	-0.47	0.00	-0.08	0.00
30	156	ADHEr, ATPS4r, FUM, PGDH	11.87	0.00	-3.80	0.00	-0.47	0.00	-0.08	0.00
	157	ADHEr, ATPS4r, MDH, PGL	11.87	0.00	-3.80	0.00	-0.47	0.00	-0.08	0.00
35	158	ADHEr, ATPS4r, G6PDHy, MDH	11.87	0.00	-3.80	0.00	-0.47	0.00	-0.08	0.00
	159	ADHEr, ATPS4r, LDH_ D, PPC	-0.41	0.00	-3.83	-10.00	-0.47	0.00	-0.08	0.02
40	160	ADHEr, ATPS4r, FUM, TAL	11.60	0.00	-3.81	0.00	-0.47	0.00	-0.08	0.00
	161	ADHEr, ATPS4r, MDH, TAL	11.60	0.00	-3.81	0.00	-0.47	0.00	-0.08	0.00
45	162	ADHEr, ATPS4r, GLCpts, NADH6	11.59	0.00	-3.65	-2.00	-0.45	0.00	-0.07	0.00
	163	ADHEr, ATPS4r, MDH, RPE	11.34	0.00	-3.82	0.00	-0.47	0.00	-0.08	0.00
50	164	ADHEr, ATPS4r, FUM, RPE	11.34	0.00	-3.82	0.00	-0.47	0.00	-0.08	0.00
	165	ADHEr, ATPS4r, NADH6, PGDH	10.80	0.00	-4.00	0.00	-0.50	0.00	-0.08	0.00
55	166	ADHEr, ATPS4r, NADH6, PGL	10.80	0.00	-4.00	0.00	-0.50	0.00	-0.08	0.00
	167	ADHEr, ATPS4r, G6PDHy, NADH6	10.80	0.00	-4.00	0.00	-0.50	0.00	-0.08	0.00

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL	
5		168	ADHEr, ATPS4r, NADH6, TAL	10.51	0.00	-4.01	0.00	-0.50	0.00	-0.08	0.00	0.00
10		169	ADHEr, ATPS4r, NADH6, RPE	10.25	0.00	-4.02	0.00	-0.50	0.00	-0.08	0.00	0.00
		170	ADHEr, G6PDHy, ME2, THD2	-0.86	0.00	-3.72	0.00	-0.46	0.00	-0.08	0.00	0.00
15		171	ADHEr, ME2, PGL, THD2	-0.86	0.00	-3.72	0.00	-0.46	0.00	-0.08	0.00	0.00
		172	ADHEr, G6PDHy, PPC, THD2	6.22	8.06	-4.07	-20.00	-0.50	0.00	-0.08	0.00	0.00
20		173	ADHEr, PGL, PPC, THD2	6.22	8.06	-4.07	-20.00	-0.50	0.00	-0.08	0.00	0.00
		174	ADHEr, ATPS4r, GLUDy, NADH6	5.40	0.00	-3.75	0.00	-0.46	0.00	-0.08	0.00	0.00
25		175	ADHEr, ACKr, FRD2, LDH_D	15.97	0.00	-6.24	-15.00	-0.69	2.92	-0.11	0.00	0.68
		176	ADHEr, ATPS4r, FBA, RPE	0.40	0.00	-2.23	0.00	-0.28	0.00	-0.05	0.00	0.00
30		177	ADHEr, ATPS4r, RPE, TPI	0.40	0.00	-2.23	0.00	-0.28	0.00	-0.05	0.00	0.00
		178	ADHEr, ATPS4r, PFK, RPE	0.40	0.00	-2.23	0.00	-0.28	0.00	-0.05	0.00	0.00
35		179	ADHEr, ATPS4r, GLUDy, PFK	-0.22	0.00	-2.02	0.00	-0.25	0.00	-0.04	0.00	0.00
		180	ADHEr, ATPS4r, GLUDy, TPI	-0.22	0.00	-2.02	0.00	-0.25	0.00	-0.04	0.00	0.00
40		181	ADHEr, ATPS4r, FBA, GLUDy	-0.22	0.00	-2.02	0.00	-0.25	0.00	-0.04	0.00	0.00
		182	ADHEr, ATPS4r, PFK, TAL	0.44	0.00	-2.24	0.00	-0.28	0.00	-0.05	0.00	0.00
45		183	ADHEr, ATPS4r, TAL, TPI	0.44	0.00	-2.24	0.00	-0.28	0.00	-0.05	0.00	0.00
		184	ADHEr, ATPS4r, FBA, TAL	0.44	0.00	-2.24	0.00	-0.28	0.00	-0.05	0.00	0.00
50		185	ADHEr, ASPT, MDH, PYK	-7.34	0.00	-2.39	0.00	-0.30	0.00	-0.05	0.00	0.00
		186	ADHEr, MDH, PGL, THD2	12.12	0.00	-6.14	-15.00	-0.76	0.00	-0.12	0.00	0.00
55		187	ADHEr, G6PDHy, MDH, THD2	12.12	0.00	-6.14	-15.00	-0.76	0.00	-0.12	0.00	0.00
		188	ADHEr, GLCpts, NADH6, THD2	2.35	0.00	-3.74	0.00	-0.46	0.00	-0.08	0.00	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	189 ADHEr, GLUDy, NADH6, THD2	2.75	0.00	-3.88	0.00	-0.48	0.00	-0.08	0.00	0.00
10	190 ADHEr, ASPT, LDH_D, MDH	-3.60	0.00	-3.89	0.00	-0.48	0.00	-0.08	0.00	0.00
	191 ADHEr, GLCpts, NADH6, PGI	-3.94	0.00	-1.68	0.00	-0.21	0.00	-0.03	0.00	0.00
15	192 ADHEr, LDH_D, SUCD4, THD2	-1.90	0.00	-3.49	0.00	-0.43	0.00	-0.07	0.00	0.00
	193 ADHEr, GLUDy, NADH6, PGI	-3.63	0.00	-1.81	0.00	-0.22	0.00	-0.04	0.00	0.00
20	194 ADHEr, ACKr, FUM, LDH_D	13.82	0.00	-2.77	0.00	-0.34	17.80	-0.06	0.00	0.00
	195 ADHEr, FBA, GLCpts, NADH6	-3.99	0.00	-1.71	0.00	-0.21	0.00	-0.03	0.00	0.00
25	196 ADHEr, GLCpts, NADH6, TPI	-3.99	0.00	-1.71	0.00	-0.21	0.00	-0.03	0.00	0.00
	197 ADHEr, ACKr, LDH_D, MDH	13.82	0.00	-2.77	0.00	-0.34	17.80	-0.06	0.00	0.00
30	198 ADHEr, GLCpts, NADH6, PFK	-3.99	0.00	-1.71	0.00	-0.21	0.00	-0.03	0.00	0.00
	199 ADHEr, GLCpts, MDH, THD2	-1.38	0.00	-3.71	0.00	-0.46	0.00	-0.08	0.00	0.00
35	200 ADHEr, FUM, GLCpts, THD2	-1.38	0.00	-3.71	0.00	-0.46	0.00	-0.08	0.00	0.00
	201 ADHEr, NADH6, PFK, RPE	-3.13	0.00	-2.03	0.00	-0.25	0.00	-0.04	0.00	0.00
40	202 ADHEr, FBA, NADH6, RPE	-3.13	0.00	-2.03	0.00	-0.25	0.00	-0.04	0.00	0.00
	203 ADHEr, NADH6, RPE, TPI	-3.13	0.00	-2.03	0.00	-0.25	0.00	-0.04	0.00	0.00
45	204 ADHEr, GLUDy, NADH6, TPI	-3.69	0.00	-1.84	0.00	-0.23	0.00	-0.04	0.00	0.00
	205 ADHEr, GLUDy, NADH6, PFK	-3.69	0.00	-1.84	0.00	-0.23	0.00	-0.04	0.00	0.00
50	206 ADHEr, FBA, GLUDy, NADH6	-3.69	0.00	-1.84	0.00	-0.23	0.00	-0.04	0.00	0.00
	207 ADHEr, NADH6, PFK, TAL	-3.15	0.00	-2.04	0.00	-0.25	0.00	-0.04	0.00	0.00
55	208 ADHEr, FBA, NADH6, TAL	-3.15	0.00	-2.04	0.00	-0.25	0.00	-0.04	0.00	0.00
	209 ADHEr, NADH6, TAL, TPI	-3.15	0.00	-2.04	0.00	-0.25	0.00	-0.04	0.00	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	210 ADHEr, ACKr, AKGD, ATPS4r	55.34	0.00	-4.99	-82.37	-0.62	0.00	-0.10	0.00	0.00
10	211 ADHEr, GLCpts, NADH6, RPE	1.30	0.00	-3.88	0.00	-0.48	0.00	-0.08	0.00	0.00
	212 ADHEr, FUM, GLCpts, NADH6	0.21	0.00	-3.50	0.00	-0.43	0.00	-0.07	0.00	0.00
15	213 ADHEr, GLCpts, MDH, NADH6	0.21	0.00	-3.50	0.00	-0.43	0.00	-0.07	0.00	0.00
	214 ADHEr, GLCpts, GLUDy, NADH6	0.22	0.00	-3.51	0.00	-0.43	0.00	-0.07	0.00	0.00
20	215 ADHEr, FUM, NADH6, PYK	0.23	0.00	-3.51	0.00	-0.43	0.00	-0.07	0.00	0.00
	216 ADHEr, MDH, NADH6, PYK	0.23	0.00	-3.51	0.00	-0.43	0.00	-0.07	0.00	0.00
25	217 ADHEr, ACKr, ATPS4r, SUCOAS	20.07	0.00	-5.03	-47.55	-0.62	0.00	-0.10	0.00	0.00
	218 ADHEr, GLUDy, NADH6, RPE	1.63	0.00	-4.01	0.00	-0.50	0.00	-0.08	0.00	0.00
30	219 ADHEr, MDH, NADH6, RPE	1.76	0.00	-4.07	0.00	-0.50	0.00	-0.08	0.00	0.00
	220 ADHEr, FUM, NADH6, RPE	1.76	0.00	-4.07	0.00	-0.50	0.00	-0.08	0.00	0.00
35	221 ADHEr, GLCpts, NADH6, TAL	1.23	0.00	-3.89	0.00	-0.48	0.00	-0.08	0.00	0.00
	222 ADHEr, GLUDy, NADH6, TAL	1.55	0.00	-4.02	0.00	-0.50	0.00	-0.08	0.00	0.00
40	223 ADHEr, FUM, NADH6, TAL	1.69	0.00	-4.08	0.00	-0.51	0.00	-0.08	0.00	0.00
	224 ADHEr, MDH, NADH6, TAL	1.69	0.00	-4.08	0.00	-0.51	0.00	-0.08	0.00	0.00
45	225 ADHEr, CBMK2, GLU5K, NADH6	2.38	0.00	-4.43	0.00	-0.55	0.00	-0.09	0.00	0.00
	226 ADHEr, CBMK2, G5SD, NADH6	2.38	0.00	-4.43	0.00	-0.55	0.00	-0.09	0.00	0.00
50	227 ADHEr, CBMK2, NADH6, SO4t2	2.40	0.00	-4.44	0.00	-0.55	0.00	-0.09	0.00	0.00
	228 ADHEr, ASNS2, CBMK2, NADH6	2.38	0.00	-4.43	0.00	-0.55	0.00	-0.09	0.00	0.00
55	229 ADHEr, ATPS4r, PYK, SUCD4	6.92	7.87	-3.12	-2.00	-0.39	0.00	-0.06	0.00	0.00
	230 ADHEr, GLCpts, PGI, SUCD4	-9.35	0.00	-1.56	0.00	-0.19	0.00	-0.03	0.00	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	231	ADHEr, FUM, GLUDy, PGI	-9.07	0.00	-1.67	0.00	-0.21	0.00	-0.03	0.00
10	232	ADHEr, GLUDy, MDH, PGI	-9.07	0.00	-1.67	0.00	-0.21	0.00	-0.03	0.00
	233	ADHEr, GLUDy, PGI, SUCD4	-8.89	0.00	-1.74	0.00	-0.22	0.00	-0.04	0.00
15	234	ADHEr, GLCpts, SUCD4, TPI	-9.35	0.00	-1.59	0.00	-0.20	0.00	-0.03	0.00
	235	ADHEr, GLCpts, PFK, SUCD4	-9.35	0.00	-1.59	0.00	-0.20	0.00	-0.03	0.00
20	236	ADHEr, FBA, GLCpts, SUCD4	-9.35	0.00	-1.59	0.00	-0.20	0.00	-0.03	0.00
	237	ADHEr, HEX1, RPE, TPI	-8.46	0.00	-1.92	0.00	-0.24	0.00	-0.04	0.00
25	238	ADHEr, FBA, HEX1, RPE	-8.46	0.00	-1.92	0.00	-0.24	0.00	-0.04	0.00
	239	ADHEr, HEX1, PFK, RPE	-8.46	0.00	-1.92	0.00	-0.24	0.00	-0.04	0.00
30	240	ADHEr, FUM, GLUDy, TPI	-9.08	0.00	-1.70	0.00	-0.21	0.00	-0.03	0.00
	241	ADHEr, FBA, GLUDy, MDH	-9.08	0.00	-1.70	0.00	-0.21	0.00	-0.03	0.00
35	242	ADHEr, FUM, GLUDy, PFK	-9.08	0.00	-1.70	0.00	-0.21	0.00	-0.03	0.00
	243	ADHEr, GLUDy, MDH, PFK	-9.08	0.00	-1.70	0.00	-0.21	0.00	-0.03	0.00
40	244	ADHEr, GLUDy, MDH, TPI	-9.08	0.00	-1.70	0.00	-0.21	0.00	-0.03	0.00
	245	ADHEr, FBA, FUM, GLUDy	-9.08	0.00	-1.70	0.00	-0.21	0.00	-0.03	0.00
45	246	ADHEr, GLCpts, GLUDy, PGI	-8.19	0.00	-2.02	0.00	-0.25	0.00	-0.04	0.00
	247	ADHEr, RPE, SUCD4, TPI	-8.37	0.00	-1.95	0.00	-0.24	0.00	-0.04	0.00
50	248	ADHEr, FBA, RPE, SUCD4	-8.37	0.00	-1.95	0.00	-0.24	0.00	-0.04	0.00
	249	ADHEr, PFK, RPE, SUCD4	-8.37	0.00	-1.95	0.00	-0.24	0.00	-0.04	0.00
55	250	ADHEr, GLUDy, HEX1, PFK	-8.96	0.00	-1.75	0.00	-0.22	0.00	-0.04	0.00
	251	ADHEr, GLUDy, HEX1, TPI	-8.96	0.00	-1.75	0.00	-0.22	0.00	-0.04	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	252	ADHEr, FBA, GLUDy, HEX1	-8.96	0.00	-1.75	0.00	-0.22	0.00	-0.04	0.00
10	253	ADHEr, FBA, GLUDy, SUCD4	-8.90	0.00	-1.77	0.00	-0.22	0.00	-0.04	0.00
	254	ADHEr, GLUDy, SUCD4, TPI	-8.90	0.00	-1.77	0.00	-0.22	0.00	-0.04	0.00
15	255	ADHEr, GLUDy, PFK, SUCD4	-8.90	0.00	-1.77	0.00	-0.22	0.00	-0.04	0.00
	256	ADHEr, FBA, HEX1, TAL	-8.47	0.00	-1.93	0.00	-0.24	0.00	-0.04	0.00
20	257	ADHEr, HEX1, PFK, TAL	-8.47	0.00	-1.93	0.00	-0.24	0.00	-0.04	0.00
	258	ADHEr, HEX1, TAL, TPI	-8.47	0.00	-1.93	0.00	-0.24	0.00	-0.04	0.00
25	259	ADHEr, PFK, SUCD4, TAL	-8.37	0.00	-1.97	0.00	-0.24	0.00	-0.04	0.00
	260	ADHEr, SUCD4, TAL, TPI	-8.37	0.00	-1.97	0.00	-0.24	0.00	-0.04	0.00
30	261	ADHEr, FBA, SUCD4, TAL	-8.37	0.00	-1.97	0.00	-0.24	0.00	-0.04	0.00
	262	ADHEr, ACKr, LDH_D, SUCD4	15.34	0.00	-1.92	0.00	-0.24	23.88	-0.04	0.00
35	263	ADHEr, GLCpts, RPE, TPI	-7.59	0.00	-2.26	0.00	-0.28	0.00	-0.05	0.00
	264	ADHEr, GLCpts, PFK, RPE	-7.59	0.00	-2.26	0.00	-0.28	0.00	-0.05	0.00
40	265	ADHEr, FBA, GLCpts, RPE	-7.59	0.00	-2.26	0.00	-0.28	0.00	-0.05	0.00
	266	ADHEr, Act6, LDH_D, MDH	21.84	0.00	-2.02	0.00	-0.25	30.11	-0.04	0.00
45	267	ADHEr, GLCpts, GLUDy, TPI	-8.22	0.00	-2.04	0.00	-0.25	0.00	-0.04	0.00
	268	ADHEr, Act6, FUM, LDH_D	21.84	0.00	-2.02	0.00	-0.25	30.11	-0.04	0.00
50	269	ADHEr, FBA, GLCpts, GLUDy	-8.22	0.00	-2.04	0.00	-0.25	0.00	-0.04	0.00
	270	ADHEr, GLCpts, GLUDy, PFK	-8.22	0.00	-2.04	0.00	-0.25	0.00	-0.04	0.00
55	271	ADHEr, GLUDy, RPE, TPI	-7.35	0.00	-2.35	0.00	-0.29	0.00	-0.05	0.00
	272	ADHEr, FBA, GLUDy, RPE	-7.35	0.00	-2.35	0.00	-0.29	0.00	-0.05	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,§	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	273	ADHEr, GLUDy, PFK, RPE	-7.35	0.00	-2.35	0.00	-0.29	0.00	-0.05	0.00
10	274	ADHEr, GLCpts, TAL, TPI	-7.60	0.00	-2.27	0.00	-0.28	0.00	-0.05	0.00
	275	ADHEr, FBA, GLCpts, TAL	-7.60	0.00	-2.27	0.00	-0.28	0.00	-0.05	0.00
15	276	ADHEr, GLCpts, PFK, TAL	-7.60	0.00	-2.27	0.00	-0.28	0.00	-0.05	0.00
	277	ADHEr, GLUDy, PFK, TAL	-7.36	0.00	-2.36	0.00	-0.29	0.00	-0.05	0.00
20	278	ADHEr, FBA, GLUDy, TAL	-7.36	0.00	-2.36	0.00	-0.29	0.00	-0.05	0.00
	279	ADHEr, GLUDy, TAL, TPI	-7.36	0.00	-2.36	0.00	-0.29	0.00	-0.05	0.00
25	280	ADHEr, GLUDy, MDH, PYK	-6.12	0.00	-2.88	0.00	-0.36	0.00	-0.06	0.00
	281	ADHEr, FUM, GLUDy, PYK	-6.12	0.00	-2.88	0.00	-0.36	0.00	-0.06	0.00
30	282	ADHEr, PYK, RPE, SUCD4	-4.99	0.00	-3.28	0.00	-0.41	0.00	-0.07	0.00
	283	ADHEr, GLUDy, PYK, SUCD4	-5.98	0.00	-2.94	0.00	-0.36	0.00	-0.06	0.00
35	284	ADHEr, MDH, PYK, SUCD4	-5.76	0.00	-3.02	0.00	-0.37	0.00	-0.06	0.00
	285	ADHEr, FUM, PYK, SUCD4	-5.76	0.00	-3.02	0.00	-0.37	0.00	-0.06	0.00
40	286	ADHEr, GLCpts, RPE, SUCD4	-4.64	0.00	-3.42	0.00	-0.42	0.00	-0.07	0.00
	287	ADHEr, GLCpts, GLUDy, MDH	-5.65	0.00	-3.07	0.00	-0.38	0.00	-0.06	0.00
45	288	ADHEr, FUM, GLCpts, GLUDy	-5.65	0.00	-3.07	0.00	-0.38	0.00	-0.06	0.00
	289	ADHEr, GLCpts, GLUDy, SUCD4	-5.58	0.00	-3.10	0.00	-0.38	0.00	-0.06	0.00
50	290	ADHEr, PYK, SUCD4, TAL	-5.00	0.00	-3.30	0.00	-0.41	0.00	-0.07	0.00
	291	ADHEr, GLUDy, MDH, RPE	-4.23	0.00	-3.58	0.00	-0.44	0.00	-0.07	0.00
55	292	ADHEr, FUM, GLUDy, RPE	-4.23	0.00	-3.58	0.00	-0.44	0.00	-0.07	0.00
	293	ADHEr, GLUDy, RPE, SUCD4	-4.17	0.00	-3.60	0.00	-0.45	0.00	-0.07	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,§	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5										
	294 ADHEr, GLCpts, SUCD4, TAL	-4.67	0.00	-3.43	0.00	-0.42	0.00	-0.07	0.00	0.00
10	295 ADHEr, GLUDy, MDH, TAL	-4.26	0.00	-3.60	0.00	-0.44	0.00	-0.07	0.00	0.00
	296 ADHEr, FUM, GLUDy, TAL	-4.26	0.00	-3.60	0.00	-0.44	0.00	-0.07	0.00	0.00
15	297 ADHEr, GLUDy, SUCD4, TAL	-4.21	0.00	-3.62	0.00	-0.45	0.00	-0.07	0.00	0.00
	298 ADHEr, GLCpts, GLUDy, RPE	-3.31	0.00	-3.94	0.00	-0.49	0.00	-0.08	0.00	0.00
20	299 ADHEr, FUM, GLCpts, RPE	-3.02	0.00	-4.06	0.00	-0.50	0.00	-0.08	0.00	0.00
	300 ADHEr, GLCpts, MDH, RPE	-3.02	0.00	-4.06	0.00	-0.50	0.00	-0.08	0.00	0.00
25	301 ADHEr, FUM, LDH_D, SUCD4	-4.04	0.00	-3.71	0.00	-0.46	0.00	-0.08	0.00	0.00
	302 ADHEr, LDH_D, MDH, SUCD4	-4.04	0.00	-3.71	0.00	-0.46	0.00	-0.08	0.00	0.00
30	303 ADHEr, GLCpts, GLUDy, TAL	-3.36	0.00	-3.95	0.00	-0.49	0.00	-0.08	0.00	0.00
	304 ADHEr, FUM, GLCpts, TAL	-3.07	0.00	-4.07	0.00	-0.50	0.00	-0.08	0.00	0.00
35	305 ADHEr, GLCpts, MDH, TAL	-3.07	0.00	-4.07	0.00	-0.50	0.00	-0.08	0.00	0.00
	306 ADHEr, CBMK2, GLU5K, SUCD4	-3.34	0.00	-3.99	0.00	-0.49	0.00	-0.08	0.00	0.00
40	307 ADHEr, CBMK2, G5SD, SUCD4	-3.34	0.00	-3.99	0.00	-0.49	0.00	-0.08	0.00	0.00
	308 ADHEr, CBMK2, GLU5K, RPE	-0.91	0.00	-4.89	0.00	-0.60	0.00	-0.10	0.00	0.00
45	309 ADHEr, CBMK2, G5SD, RPE	-0.91	0.00	-4.89	0.00	-0.60	0.00	-0.10	0.00	0.00
	310 ADHEr, CBMK2, GLCpts, GLU5K	-2.47	0.00	-4.34	0.00	-0.54	0.00	-0.09	0.00	0.00
50	311 ADHEr, ASNS2, CBMK2, RPE	-0.91	0.00	-4.89	0.00	-0.60	0.00	-0.10	0.00	0.00
	312 ADHEr, CBMK2, G5SD, GLCpts	-2.47	0.00	-4.34	0.00	-0.54	0.00	-0.09	0.00	0.00
55	313 ADHEr, ASNS2, CBMK2, GLCpts	-2.47	0.00	-4.34	0.00	-0.54	0.00	-0.09	0.00	0.00
	314 ADHEr, CBMK2, GLU5K, TAL	-0.98	0.00	-4.89	0.00	-0.61	0.00	-0.10	0.00	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	315 ADHEr, CBMK2, G5SD, TAL	-0.98	0.00	-4.89	0.00	-0.61	0.00	-0.10	0.00	0.00
10	316 ADHEr, ASNS2, CBMK2, TAL	-0.98	0.00	-4.90	0.00	-0.61	0.00	-0.10	0.00	0.00
	317 ADHEr, CBMK2, GLU5K, MDH	-1.80	0.00	-4.61	0.00	-0.57	0.00	-0.09	0.00	0.00
15	318 ADHEr, CBMK2, FUM, G5SD	-1.80	0.00	-4.61	0.00	-0.57	0.00	-0.09	0.00	0.00
	319 ADHEr, CBMK2, G5SD, MDH	-1.80	0.00	-4.61	0.00	-0.57	0.00	-0.09	0.00	0.00
20	320 ADHEr, CBMK2, FUM, GLU5K	-1.80	0.00	-4.61	0.00	-0.57	0.00	-0.09	0.00	0.00
	321 ADHEr, ASNS2, CBMK2, MDH	-1.80	0.00	-4.61	0.00	-0.57	0.00	-0.09	0.00	0.00
25	322 ADHEr, ASNS2, CBMK2, FUM	-1.80	0.00	-4.61	0.00	-0.57	0.00	-0.09	0.00	0.00
	323 ADHEr, ASNS2, GLU5K, SO4t2	-1.01	0.00	-4.92	0.00	-0.61	0.00	-0.10	0.00	0.00
30	324 ADHEr, ASPT, LDH_D, MDH, PFLi	11.71	0.00	-2.00	0.00	-0.25	0.00	-0.04	0.00	0.00
	325 ADHEr, EDA, GLUDy, PFLi, PGI	10.00	0.00	-1.23	0.00	-0.15	0.00	-0.02	0.00	0.00
35	326 ADHEr, ATPS4r, G6PDHy, GLCpts, MDH	10.67	0.00	-1.59	0.00	-0.20	0.00	-0.03	0.00	0.00
	327 ADHEr, ATPS4r, GLCpts, MDH, PGL	10.67	0.00	-1.59	0.00	-0.20	0.00	-0.03	0.00	0.00
40	328 ADHEr, EDA, GLUDy, NADH6, PGI	9.64	0.00	-0.99	-2.00	-0.12	0.00	-0.02	0.00	0.00
	329 ADHEr, G6PDHy, LDH_D, PPC, THD2	-10.49	0.00	-0.87	-2.00	-0.11	0.00	-0.02	0.00	0.00
45	330 ADHEr, LDH_D, PGL, PPC, THD2	-10.49	0.00	-0.87	-2.00	-0.11	0.00	-0.02	0.00	0.00
	331 ADHEr, ATPS4r, FRD2, LDH_D, NADH6	9.33	0.00	-7.86	-5.00	-0.18	0.00	-0.03	0.00	0.00
50	332 ADHEr, EDA, LDH_D, PPC, THD2	-8.88	0.00	-1.36	-2.00	-0.17	0.00	-0.03	0.00	0.00
	333 ADHEr, FRD2, GLUDy, LDH_D, RPE	-10.71	0.00	-1.03	0.00	-0.13	0.00	-0.02	0.00	0.00
55	334 ADHEr, FRD2, LDH_D, RPE, THD2	-10.58	0.00	-1.08	0.00	-0.13	0.00	-0.02	0.00	0.00

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5											
	335	ADHEr, FRD2, GLUDy, LDH_D, THD2	-10.88	0.00	-0.98	0.00	-0.12	0.00	-0.02	0.00	0.00
10											
	336	ADHEr, FRD2, GLUDy, LDH_D, TAL	-10.71	0.00	-1.04	0.00	-0.13	0.00	-0.02	0.00	0.00
	337	ADHEr, FRD2, LDH_D, TAL, THD2	-10.59	0.00	-1.09	0.00	-0.13	0.00	-0.02	0.00	0.00
15											
	338	ADHEr, ATPS4r, LDH_D, NADH6, PPC	1.95	0.00	-2.42	-10.00	-0.30	0.00	-0.05	0.00	0.00
	339	ADHEr, ASPT, FUM, GLUDy, LDH_D	-10.53	0.00	-1.12	0.00	-0.14	0.00	-0.02	0.00	0.00
20											
	340	ADHEr, ASPT, FUM, LDH_D, THD2	-10.42	0.00	-1.16	0.00	-0.14	0.00	-0.02	0.00	0.00
	341	ADHEr, ME2, PFLi, PGL, THD2	13.21	0.00	-3.25	0.00	-0.40	0.00	-0.07	0.00	0.00
25											
	342	ADHEr, G6PDHy, ME2, PFLi, THD2	13.21	0.00	-3.25	0.00	-0.40	0.00	-0.07	0.00	0.00
	343	ADHEr, ACKr, ACS, PPC, RPE	-10.40	0.00	-0.89	-2.00	-0.11	0.00	-0.02	0.00	0.00
30											
	344	ADHEr, GLUDy, LDH_D, PPC, RPE	-9.20	0.00	-1.37	-2.00	-0.17	0.00	-0.03	0.00	0.00
	345	ADHEr, LDH_D, PPC, RPE, THD2	-9.06	0.00	-1.42	-2.00	-0.18	0.00	-0.03	0.00	0.00
35											
	346	ADHEr, GLUDy, LDH_D, PPC, TAL	-9.20	0.00	-1.37	-2.00	-0.17	0.00	-0.03	0.00	0.00
	347	ADHEr, LDH_D, PPC, TAL, THD2	-9.08	0.00	-1.42	-2.00	-0.18	0.00	-0.03	0.00	0.00
40											
	348	ADHEr, GLCpts, GLUDy, LDH_D, PPC	-10.40	0.00	-0.91	-2.00	-0.11	0.00	-0.02	0.00	0.00
	349	ADHEr, GLCpts, LDH_D, PPC, THD2	-10.32	0.00	-0.93	-2.00	-0.12	0.00	-0.02	0.00	0.00
45											
	350	ADHEr, ACKr, GLUDy, LDH_D, PGI	6.43	0.00	-1.13	0.00	-0.14	6.48	-0.02	0.00	0.00
	351	ADHEr, ASPT, ATPS4r, GLCpts, MDH	9.70	8.01	-1.71	0.00	-0.21	0.00	-0.03	0.00	0.00
50											
	352	ADHEr, LDH_D, PFLi, SUCD4, THD2	10.54	0.00	-2.12	0.00	-0.26	0.00	-0.04	0.00	0.00
	353	ADHEr, ACKr, AKGD, ATPS4r, PYK	4.35	0.00	-2.07	-13.40	-0.26	0.00	-0.04	0.00	0.00
55											
	354	ADHEr, ACKr, ATPS4r, PYK, SUCOAS	4.62	0.00	-2.09	-13.93	-0.26	0.00	-0.04	0.00	0.00
	355	ADHEr, ATPS4r, EDA, GLUDy, PGI	-1.56	0.00	-1.15	0.00	-0.14	0.00	-0.02	0.00	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,§	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	356	ADHEr, EDA, GLCpts, GLUDy, PGI	-1.51	0.00	-1.18	0.00	-0.15	0.00	-0.02	0.00
10	357	ADHEr, ACKr, LDH_D, MDH, SUCD4	0.47	0.00	-5.01	0.00	-0.21	0.00	-0.03	0.00
	358	ADHEr, ACKr, GLUDy, NADH6, PYK	24.87	0.00	-4.23	-20.00	-0.52	0.00	-0.09	0.00
15	359	ADHEr, FUM, LDH_D, PFLi, THD2	13.05	0.00	-3.40	0.00	-0.42	0.00	-0.07	0.00
	360	ADHEr, LDH_D, MDH, PFLi, THD2	13.05	0.00	-3.40	0.00	-0.42	0.00	-0.07	0.00
20	361	ADHEr, Act6, ATPS4r, LDH_D, PPC	4.77	0.00	-1.18	-5.00	-0.15	12.79	-0.02	0.00
	362	ADHEr, ACKr, LDH_D, PYRt2, SUCD4	2.88	0.00	-3.91	0.00	-0.22	0.00	-0.04	0.00
25	363	ADHEr, ATPS4r, FUM, LDH_D, SUCD4	6.51	0.00	-2.01	-2.00	-0.25	0.00	-0.04	0.00
	364	ADHEr, ATPS4r, LDH_ D, MDH, SUCD4	6.51	0.00	-2.01	-2.00	-0.25	0.00	-0.04	0.00
30	365	ADHEr, ACKr, CBMK2, NADH6, PYK	25.89	0.00	-4.80	-20.00	-0.59	0.00	-0.10	0.00
	366	ADHEr, ACKr, NADH6, PYK, RPE	25.93	0.00	-4.82	-20.00	-0.60	0.00	-0.10	0.00
35	367	ADHEr, ACKr, ASNS2, NADH6, PYK	25.95	0.00	-4.83	-20.00	-0.60	0.00	-0.10	0.00
	368	ADHEr, ACKr, NADH6, PYK, TAL	25.96	0.00	-4.84	-20.00	-0.60	0.00	-0.10	0.00
40	369	ADHEr, ASPT, ATPS4r, FUM, LDH_D	4.50	0.00	-4.47	-5.00	-0.33	0.00	-0.05	0.00
	370	ADHEr, ASPT, ATPS4r, LDH_D, MDH	3.02	0.00	-2.92	0.00	-0.36	0.00	-0.06	0.00
45	371	ADHEr, MDH, PFLi, PYK, THD2	12.56	0.00	-3.25	0.00	-0.40	0.00	-0.07	0.00
	372	ADHEr, FUM, PFLi, PYK, THD2	12.56	0.00	-3.25	0.00	-0.40	0.00	-0.07	0.00
50	373	ADHEr, ACKr, FUM, LDH_D, NADH6	23.94	0.00	-4.61	-15.00	-0.57	2.16	-0.09	0.00
	374	ADHEr, ATPS4r, G6PDHy, LDH_D, SUCD4	7.29	0.00	-2.38	-2.00	-0.29	0.00	-0.05	0.00
55	375	ADHEr, ATPS4r, LDH_ D, PGL, SUCD4	7.29	0.00	-2.38	-2.00	-0.29	0.00	-0.05	0.00

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5											
	376	ADHEr, ATPS4r, LDH_D, PGDH, SUCD4	7.29	0.00	-2.38	-2.00	-0.29	0.00	-0.05	0.00	0.00
10	377	ADHEr, GLYCL, PGL, PPC, THD2	-5.10	10.37	-1.24	-2.00	-0.15	0.00	-0.02	0.00	0.00
	378	ADHEr, G6PDHy, GLYCL, PPC, THD2	-5.10	10.37	-1.24	-2.00	-0.15	0.00	-0.02	0.00	0.00
15	379	ADHEr, FTHFD, G6PDHy, PPC, THD2	-5.08	10.43	-1.22	-2.00	-0.15	0.00	-0.02	0.00	0.00
	380	ADHEr, FTHFD, PGL, PPC, THD2	-5.08	10.43	-1.22	-2.00	-0.15	0.00	-0.02	0.00	0.00
20	381	ADHEr, MTHFC, PGL, PPC, THD2	-5.08	10.43	-1.22	-2.00	-0.15	0.00	-0.02	0.00	0.00
	382	ADHEr, G6PDHy, MTHFC, PPC, THD2	-5.08	10.43	-1.22	-2.00	-0.15	0.00	-0.02	0.00	0.00
25	383	ADHEr, ATPS4r, LDH_D, PFLi, SUCD4	12.44	0.00	-2.76	-2.00	-0.34	0.00	-0.06	0.00	0.00
	384	ADHEr, ATPS4r, LDH_D, SUCD4, TAL	7.30	0.00	-2.40	-2.00	-0.30	0.00	-0.05	0.00	0.00
30	385	ADHEr, ATPS4r, LDH_D, RPE, SUCD4	11.65	0.00	-2.41	-2.00	-0.30	0.00	-0.05	0.00	0.00
	386	ADHEr, ATPS4r, FUM, GLCpts, NADH6	10.51	2.86	-2.34	0.00	-0.29	0.00	-0.05	0.00	0.00
35	387	ADHEr, ATPS4r, GLCpts, MDH, NADH6	10.51	2.86	-2.34	0.00	-0.29	0.00	-0.05	0.00	0.00
	388	ADHEr, ATPS4r, LDH_D, NADH6, PFLi	11.74	0.00	-2.93	0.00	-0.36	0.00	-0.06	0.00	0.00
40	389	ADHEr, ACKr, MALS, NADH12, NADH6	27.12	0.00	-5.49	-20.00	-0.68	0.00	-0.11	0.00	0.00
	390	ADHEr, ACKr, ICL, NADH12, NADH6	27.12	0.00	-5.49	-20.00	-0.68	0.00	-0.11	0.00	0.00
45	391	ADHEr, ACKr, CBMK2, LDH_D, NADH6	27.15	0.00	-5.51	-20.00	-0.68	0.00	-0.11	0.00	0.00
	392	ADHEr, ATPS4r, GLCpts, MDH, PFLi	11.83	0.00	-2.99	0.00	-0.37	0.00	-0.06	0.00	0.00
50	393	ADHEr, ATPS4r, FUM, GLCpts, PFLi	11.83	0.00	-2.99	0.00	-0.37	0.00	-0.06	0.00	0.00
	394	ADHEr, ACKr, ASNS2, LDH_D, NADH6	27.21	0.00	-5.54	-20.00	-0.69	0.00	-0.11	0.00	0.00
55	395	ADHEr, ACKr, ATPS4r, LDH_D, THD2	0.94	0.00	-3.26	-10.00	-0.40	0.00	-0.07	0.00	0.00
	396	ADHEr, GLCpts, GLUDy, PFLi, PGI	8.29	0.00	-1.32	0.00	-0.16	0.00	-0.03	0.00	0.00

EP 2 679 676 B1

(continued)

		Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5	397	ADHEr, GLCpts, GLUDy, PFK, PFLi	8.31	0.00	-1.34	0.00	-0.17	0.00	-0.03	0.00	0.00
10	398	ADHEr, FBA, GLCpts, GLUDy, PFLi	8.31	0.00	-1.34	0.00	-0.17	0.00	-0.03	0.00	0.00
	399	ADHEr, GLCpts, GLUDy, PFLi, TPI	8.31	0.00	-1.34	0.00	-0.17	0.00	-0.03	0.00	0.00
15	400	ADHEr, ATPS4r, GLCpts, NADH6, PGI	7.07	0.00	-0.99	0.00	-0.12	0.00	-0.02	0.00	0.00
	401	ADHEr, ATPS4r, GLCpts, NADH6, TPI	7.11	0.00	-1.00	0.00	-0.12	0.00	-0.02	0.00	0.00
20	402	ADHEr, ATPS4r, GLCpts, NADH6, PFK	7.11	0.00	-1.00	0.00	-0.12	0.00	-0.02	0.00	0.00
	403	ADHEr, ATPS4r, FBA, GLCpts, NADH6	7.11	0.00	-1.00	0.00	-0.12	0.00	-0.02	0.00	0.00
25	404	ADHEr, GLCpts, PFLi, RPE, TPI	8.61	0.00	-1.49	0.00	-0.18	0.00	-0.03	0.00	0.00
	405	ADHEr, GLCpts, PFK, PFLi, RPE	8.61	0.00	-1.49	0.00	-0.18	0.00	-0.03	0.00	0.00
30	406	ADHEr, FBA, GLCpts, PFLi, RPE	8.61	0.00	-1.49	0.00	-0.18	0.00	-0.03	0.00	0.00
	407	ADHEr, ATPS4r, NADH6, PFLi, PYK	10.57	0.00	-2.42	0.00	-0.30	0.00	-0.05	0.00	0.00
35	408	ADHEr, GLCpts, PFK, PFLi, TAL	8.62	0.00	-1.49	0.00	-0.18	0.00	-0.03	0.00	0.00
	409	ADHEr, FBA, GLCpts, PFLi, TAL	8.62	0.00	-1.49	0.00	-0.18	0.00	-0.03	0.00	0.00
40	410	ADHEr, GLCpts, PFLi, TAL, TPI	8.62	0.00	-1.49	0.00	-0.18	0.00	-0.03	0.00	0.00
	411	ADHEr, FBA, GLUD, PFLi, RPE	8.73	0.00	-1.55	0.00	-0.19	0.00	-0.03	0.00	0.00
45	412	ADHEr, GLUDy, PFK, PFLi, RPE	8.73	0.00	-1.55	0.00	-0.19	0.00	-0.03	0.00	0.00
	413	ADHEr, GLUDy, PFLi, RPE, TPI	8.73	0.00	-1.55	0.00	-0.19	0.00	-0.03	0.00	0.00
50	414	ADHEr, GLUDy, PFLi, TAL, TPI	8.74	0.00	-1.56	0.00	-0.19	0.00	-0.03	0.00	0.00
	415	ADHEr, GLUDy, PFK, PFLi, TAL	8.74	0.00	-1.56	0.00	-0.19	0.00	-0.03	0.00	0.00
55	416	ADHEr, FBA, GLUDy, PFLi, TAL	8.74	0.00	-1.56	0.00	-0.19	0.00	-0.03	0.00	0.00
	417	ADHEr, ATPS4r, LDH_ D, NADH6, SUCD4	9.67	0.00	-2.81	-2.00	-0.35	0.00	-0.06	0.00	0.00

EP 2 679 676 B1

(continued)

	Metabolic Transformations Targeted for Removal †,‡,#	H2O	LAC	NH4	NO3	PI	PYR	SO4	SUC	VAL
5										
	418 ADHEr, FUM, LDH_D, PFLi, SUCD4	9.50	0.00	-1.95	0.00	-0.24	0.00	-0.04	0.00	0.00
10	419 ADHEr, LDH_D, MDH, PFLi, SUCD4	9.50	0.00	-1.95	0.00	-0.24	0.00	-0.04	0.00	0.00

15

20

25

30

35

40

45

50

55

Table 7. Knockout strategies derived by OptKnock assuming FLP carboxylase to be reversible.

	Metabolic Trans- formations Targeted For Removal + #	BDO	BIO	AC	ALA	CO2	FOR	FUM	GLC	GLY	H+
1	ADHEC, NADH6	6.43	0.72	22.08	0.00	1.90	27.29	0.00	20.00	0.00	44.88
2	ADHEC, FNO	6.29	0.03	33.29	0.00	6.29	33.43	0.00	20.00	0.00	66.92
3	ADHEC, PGM	6.29	0.03	33.29	0.00	6.29	33.43	0.00	20.00	0.00	66.92
4	ADHEC, PPK	5.67	0.57	35.42	0.00	5.74	28.42	0.00	20.00	0.00	57.91
5	ADHEC, SURTS	8.54	0.65	24.41	0.00	5.65	37.79	0.00	20.00	0.00	56.89
6	ADHEC, ADPSR	4.41	0.82	32.86	0.00	4.52	37.15	0.00	20.00	0.00	55.83
7	ADHEC, PGI	1.81	0.52	0.00	0.00	11.12	2.74	0.00	20.00	0.00	56.18
8	ADHEC, FUM	1.32	0.74	14.95	0.00	13.58	18.85	0.00	20.00	0.00	63.42
9	ADHEC, HENI	0.82	0.77	13.74	0.00	14.47	17.74	0.00	20.00	0.00	64.01
10	ADHEC, MDH	0.72	0.67	14.49	0.00	15.13	18.02	0.00	20.00	0.00	65.93
11	ADHEC, PP	0.33	0.49	15.60	0.00	16.86	18.16	0.00	20.00	0.00	70.17
12	ADHEC, PBA	0.33	0.49	15.60	0.00	16.86	18.16	0.00	20.00	0.00	70.17
13	ADHEC, PPK	0.33	0.49	15.60	0.00	16.86	18.16	0.00	20.00	0.00	70.17
14	ADHEC, HENI, PGI	11.52	0.39	15.75	0.00	8.35	17.80	0.00	20.00	0.00	36.34
15	ADHEC, PPI, PPK	9.95	0.50	22.31	0.00	14.90	0.00	0.00	20.00	0.00	25.85
16	ADHEC, PPI, SURTS	9.81	0.55	21.71	0.00	14.68	0.00	0.00	20.00	0.00	25.59
17	ADHEC, ACCK, NADH6	9.81	0.86	10.27	0.00	24.66	0.00	0.00	20.00	0.00	16.36
18	ADHEC, NADH6, PPI	9.71	0.58	17.27	0.00	14.52	0.00	0.00	20.00	0.00	25.40
19	ADHEC, NADH6, PGM	7.82	0.08	30.86	0.00	9.99	13.48	0.00	20.00	0.00	44.95
20	ADHEC, FNO, NADH6	7.82	0.08	30.86	0.00	9.99	13.48	0.00	20.00	0.00	44.95

5
10
15
20
25
30
35
40
45
50
55

21	ADH1c, ASPT, MDH	20.76	0.00	-4.06	30.56	0.00	-20.00	0.00	58.08
22	ADH1c, NADH6, PGI	22.03	0.00	0.27	21.51	0.00	-20.00	0.00	51.08
23	ADH1c, NADH6, TPI	22.27	0.00	0.05	21.80	0.00	-20.00	0.00	51.52
24	ADH1c, FBA, NADH6	22.27	0.00	-0.05	21.80	0.00	-20.00	0.00	51.52
25	ADH1c, NADH6, PFK	22.27	0.00	0.05	21.80	0.00	-20.00	0.00	51.52
26	ADH1c, NADH6, PCK	28.03	0.00	-0.67	20.82	0.00	-20.00	0.00	49.54
27	ADH1c, MDH, NADH6	24.04	0.00	-0.08	30.57	0.00	-20.00	0.00	48.62
28	ADH1c, FUM, NADH6	23.01	0.00	-0.09	29.90	0.00	-20.00	0.00	47.66
29	ADH1c, PCK, THD2	23.25	0.00	3.54	26.67	0.00	-20.00	0.00	54.58
30	ADH1c, NADH6, RPE	21.20	0.00	5.45	23.37	0.00	-20.00	0.00	40.40
31	ADH1c, NADH6, TAT	21.41	0.00	5.15	23.97	0.00	-20.00	0.00	40.88
32	ADH1c, PGI, PCK	28.99	0.00	-5.08	30.57	0.00	-20.00	0.00	61.72
33	ADH1c, PGI, SUCD4	28.63	0.00	5.93	30.55	0.00	-20.00	0.00	61.51
34	ADH1c, ATPS4c, PCK	25.38	0.00	3.70	26.55	0.00	-20.00	0.00	54.24
35	ADH1c, PFK, PCK	29.10	0.00	-6.17	30.70	0.00	-20.00	0.00	61.98
36	ADH1c, FBA, PCK	29.10	0.00	-6.17	30.70	0.00	-20.00	0.00	61.98
37	ADH1c, PCK, TPI	29.20	0.00	6.17	30.70	0.00	-20.00	0.00	61.98
38	ADH1c, FBA, FEN1	28.91	0.00	-6.15	30.59	0.00	-20.00	0.00	61.78
39	ADH1c, FEN1, PFK	28.91	0.00	-6.15	30.59	0.00	-20.00	0.00	61.78
40	ADH1c, FEN1, TPI	28.91	0.00	-6.15	30.59	0.00	-20.00	0.00	61.78
41	ADH1c, MDH, THD2	14.90	0.00	-3.07	18.16	0.00	-20.00	0.00	49.89
42	ADH1c, SUCD4, TPI	28.75	0.00	6.13	30.48	0.00	-20.00	0.00	61.59
43	ADH1c, FBA, SUCD4	28.75	0.00	-6.13	30.48	0.00	-20.00	0.00	61.59
44	ADH1c, PFK, SUCD4	28.75	0.00	-6.13	30.48	0.00	-20.00	0.00	61.59
45	ADH1c, FUM, PFK	15.28	0.00	-0.93	0.00	0.00	-20.00	0.00	36.10
46	ADH1c, PCK, SUCD4	26.89	0.00	-5.92	29.33	0.00	-20.00	0.00	59.54
47	ADH1c, PCK, RPE	25.20	0.00	-5.45	28.29	0.00	-20.00	0.00	57.44
48	ADH1c, G.Caps, PCK	26.33	0.00	5.85	28.99	0.00	-20.00	0.00	58.92
49	ADH1c, G.Caps, MDH	26.30	0.00	-5.85	28.97	0.00	-20.00	0.00	58.89

5
10
15
20
25
30
35
40
45
50
55

50	ADHEr, GLUDy, PCK	5,77	0,52	26,20	0,00	5,84	28,90	4,00	20,00	0,00	58,78
51	ADHEr, MDL, SUC'D4	5,74	0,53	25,90	0,00	5,81	28,78	4,00	20,00	0,00	58,55
52	ADHEr, PCK, TAY	5,73	0,57	25,30	0,00	5,89	28,30	4,00	20,00	0,00	57,57
53	ADHEr, FUM, PCK	5,73	0,54	25,90	0,00	5,80	28,72	4,00	20,00	0,00	58,44
54	ADHEr, MDL, PCK	5,73	0,54	25,90	0,00	5,80	28,73	4,00	20,00	0,00	58,44
55	ADHEr, RPE, SUC'D4	5,68	0,64	24,22	0,00	5,31	27,57	4,00	20,00	0,00	56,34
56	ADHEr, MDL, SUC'D4	5,68	0,57	25,49	0,00	5,75	28,46	4,00	20,00	0,00	57,99
57	ADHEr, FUM, GLUDy	5,66	0,58	25,35	0,00	5,74	28,38	4,00	20,00	0,00	57,84
58	ADHEr, GLUDy, SUC'D4	5,66	0,58	25,34	0,00	5,74	28,37	4,00	20,00	0,00	57,83
59	ADHEr, GLCPs, SUC'D4	5,65	0,58	25,28	0,00	5,73	28,34	4,00	20,00	0,00	57,77
60	ADHEr, SUC'D4, TAY	5,61	0,64	23,31	0,00	5,46	27,67	4,00	20,00	0,00	56,86
61	ADHEr, FUM, SUC'D4	5,57	0,63	24,65	0,00	5,66	27,94	4,00	20,00	0,00	57,07
62	ADHEr, HEN1, SUC'D4	5,56	0,64	23,55	0,00	5,64	27,88	4,00	20,00	0,00	56,95
63	ADHEr, CBMK2, SUC'D4	5,55	0,64	23,49	0,00	5,64	27,84	4,00	20,00	0,00	56,89
64	ADHEr, FUM, HEN1	5,44	0,70	23,61	0,00	5,53	27,29	4,00	20,00	0,00	55,91
65	ADHEr, HEN1, PPL	5,24	0,72	14,09	0,00	3,70	0,00	0,00	20,00	0,00	48,10
66	ADHEr, ATPS4r, PGI	5,00	0,52	26,72	0,00	4,66	29,44	0,00	20,00	0,00	59,85
67	ADHEr, ATPS4r, PBA	4,97	0,53	26,89	0,00	4,99	29,65	0,00	20,00	0,00	60,29
68	ADHEr, ATPS4r, PTK	4,97	0,53	26,89	0,00	4,99	29,65	0,00	20,00	0,00	60,29
69	ADHEr, ATPS4r, LPI	4,97	0,53	26,89	0,00	4,99	29,65	0,00	20,00	0,00	60,29
70	ADHEr, PPL, LPI	4,88	0,44	16,04	0,00	1,18	0,00	0,00	20,00	0,00	43,62
71	ADHEr, PCK, PPL	4,88	0,44	16,04	0,00	1,18	0,00	0,00	20,00	0,00	43,62
72	ADHEr, PBA, PPL	4,88	0,44	16,04	0,00	1,18	0,00	0,00	20,00	0,00	43,62
73	ADHEr, HEN1, LUD2	4,87	0,74	23,17	0,00	6,49	25,99	0,00	20,00	0,00	56,39
74	ADHEr, ATPS4r, RPE	4,59	0,81	22,69	0,00	4,12	26,91	0,00	20,00	0,00	55,35
75	ADHEr, ATPS4r, GLUDy	4,56	0,73	24,10	0,00	4,66	27,92	0,00	20,00	0,00	57,20
76	ADHEr, ATPS4r, TAY	4,50	0,81	23,77	0,00	4,37	27,02	0,00	20,00	0,00	55,57
77	ADHEr, ATPS4r, CBMK2	4,47	0,81	22,97	0,00	4,53	27,22	0,00	20,00	0,00	55,95
78	ADHEr, EDNA, PGI	4,21	0,51	0,00	0,00	5,69	2,64	0,00	20,00	0,00	48,84

5
10
15
20
25
30
35
40
45
50
55

79	ADHET, PEG, PGL	2.48	0.52	0.00	0.00	-8.45	0.00	0.00	-20.00	0.00	52.22
80	ADHET, NDEL, PCK	1.36	0.47	17.85	0.00	-14.86	20.31	0.00	-20.00	0.00	68.38
81	ADHET, NDEL, TPI	1.36	0.47	17.85	0.00	-14.86	20.31	0.00	-20.00	0.00	68.38
82	ADHET, FBA, MDH	1.36	0.47	17.85	0.00	-14.86	20.31	0.00	-20.00	0.00	68.38
83	ADHET, HEXN, RPE	1.13	0.36	13.77	0.00	-13.77	17.76	0.00	-20.00	0.00	63.00
84	ADHET, NDEL, RPE	1.00	0.57	14.57	0.00	-14.57	18.02	0.00	-20.00	0.00	65.11
85	ADHET, HEXN, TAI	0.98	0.76	13.76	0.00	-14.10	17.75	0.00	-20.00	0.00	63.53
86	ADHET, NDEL, TAI	0.86	0.67	14.80	0.00	-14.81	18.02	0.00	-20.00	0.00	65.57
87	ADHET, RPE, TPI	0.47	0.48	15.53	0.00	-16.54	18.07	0.00	-20.00	0.00	69.75
88	ADHET, PCK, RPE	0.47	0.48	15.53	0.00	-16.54	18.07	0.00	-20.00	0.00	69.75
89	ADHET, FBA, RPE	0.47	0.48	15.53	0.00	-16.54	18.07	0.00	-20.00	0.00	69.75
90	ADHET, PCK, TAI	0.40	0.49	15.57	0.00	-16.69	18.11	0.00	-20.00	0.00	69.95
91	ADHET, FBA, TAI	0.40	0.49	15.57	0.00	-16.69	18.11	0.00	-20.00	0.00	69.95
92	ADHET, HEXN, TPI, PGL	14.20	0.34	13.87	0.00	21.29	0.00	0.00	-20.00	0.00	16.25
93	ADHET, HEXN, NADH6, PGL	14.07	0.28	14.98	0.00	19.90	2.36	0.00	-20.00	0.00	19.37
94	ADHET, FBA, NADH6, PGL	14.00	0.31	14.58	0.00	19.89	2.18	0.00	-20.00	0.00	18.93
95	ADHET, ACK, NADH6, PGL	13.92	0.34	14.10	0.00	21.85	0.00	0.00	-20.00	0.00	16.55
96	ADHET, PRD2, FDEL, D, MDH	13.31	0.25	5.13	0.00	13.27	19.78	0.00	-20.00	0.00	25.86
97	ADHET, ATPS-4, PGL, SUCD4	13.14	0.27	5.98	0.00	17.23	5.93	0.00	-20.00	0.00	24.84
98	ADHET, ATPS-6, FDEL2, NADH6	12.33	0.49	18.46	0.00	19.46	0.00	0.00	-20.00	0.00	18.94
99	ADHET, ACK, NADH6, TPI	12.03	0.46	12.29	0.00	28.02	0.00	0.00	-20.00	0.00	15.58
100	ADHET, ACK, FBA, NADH6	12.03	0.46	12.29	0.00	28.02	0.00	0.00	-20.00	0.00	15.58
101	ADHET, ACK, NADH6, PCK	12.03	0.46	12.29	0.00	28.02	0.00	0.00	-20.00	0.00	15.58
102	ADHET, PRD2, FDEL, D, MDH	11.95	0.26	5.09	0.00	7.62	17.42	4.40	-20.00	0.00	33.78
103	ADHET, HEXN, PGL, PCK	11.82	0.22	18.07	0.00	8.10	19.23	0.00	-20.00	0.00	38.88
104	ADHET, FBA, PGL, PCK	11.82	0.22	18.05	0.00	8.10	19.22	0.00	-20.00	0.00	38.86
105	ADHET, HEXN, PGL, SUCD4	11.77	0.25	17.60	0.00	8.15	18.98	0.00	-20.00	0.00	38.43
106	ADHET, FBA, PGL, SUCD4	11.76	0.26	17.62	0.00	8.15	18.95	0.00	-20.00	0.00	38.39
107	ADHET, ATPS-6, FBA, PGL	11.62	0.34	16.57	0.00	8.27	18.28	0.00	-20.00	0.00	37.18

5
10
15
20
25
30
35
40
45
50
55

108	ADHEr, GLUTD ₂ , HEXI, PGI	11.59	0.35	15.29	0.00	8.29	18.13	0.00	20.00	0.00	30.23
109	ADHEr, NADH, PGI, THD2	11.18	0.48	13.34	0.00	4.81	23.87	0.00	20.00	0.00	30.67
110	ADHEr, G6PDH ₂ , MDH, THD2	11.18	0.48	13.34	0.00	4.81	23.87	0.00	20.00	0.00	30.62
111	ADHEr, PFI, PGI, PPCK	10.87	0.20	25.96	0.00	16.29	0.00	0.00	-20.00	0.00	27.38
112	ADHEr, PFI, PPCK, TPI	10.83	0.20	26.02	0.00	16.23	0.00	0.00	20.00	0.00	27.46
113	ADHEr, FBA, PFI, PPCK	10.83	0.20	26.02	0.00	16.23	0.00	0.00	-20.00	0.00	27.46
114	ADHEr, PFK, PFI, PPCK	10.83	0.20	26.02	0.00	16.23	0.00	0.00	20.00	0.00	27.46
115	ADHEr, ACK, MDH, NADH ₆	10.80	0.34	11.20	0.00	23.65	0.00	0.00	20.00	0.00	16.47
116	ADHEr, NADH ₆ , PFI, PGI	10.79	0.23	25.57	0.00	16.16	0.00	0.00	-20.00	0.00	27.21
117	ADHEr, PFI, PGI, SUCD4	10.79	0.23	25.57	0.00	16.16	0.00	0.00	20.00	0.00	27.21
118	ADHEr, FBA, PFI, SUCD4	10.74	0.23	25.64	0.00	16.09	0.00	0.00	-20.00	0.00	27.30
119	ADHEr, PFK, PFI, SUCD4	10.74	0.23	25.64	0.00	16.09	0.00	0.00	20.00	0.00	27.30
120	ADHEr, NADH ₆ , PFK, PFI	10.74	0.23	25.64	0.00	16.09	0.00	0.00	-20.00	0.00	27.30
121	ADHEr, FBA, NADH ₆ , PFI	10.74	0.23	25.64	0.00	16.09	0.00	0.00	20.00	0.00	27.30
122	ADHEr, PFI, SUCD4, TPI	10.74	0.23	25.64	0.00	16.09	0.00	0.00	-20.00	0.00	27.30
123	ADHEr, NADH ₆ , PFI, TPI	10.74	0.23	25.64	0.00	16.08	0.00	0.00	20.00	0.00	27.28
124	ADHEr, HEXI, PFK, PFI	10.73	0.24	25.61	0.00	16.08	0.00	0.00	-20.00	0.00	27.28
125	ADHEr, FBA, HEXI, PFI	10.73	0.24	25.61	0.00	16.08	0.00	0.00	20.00	0.00	27.28
126	ADHEr, HEXI, PFI, TPI	10.73	0.24	25.61	0.00	16.08	0.00	0.00	-20.00	0.00	27.28
127	ADHEr, PFI, PPCK, THD2	10.49	0.49	21.02	0.00	15.71	0.00	0.00	20.00	0.00	24.49
128	ADHEr, ACK, GLUTD ₂ , NADH ₆	10.46	0.75	10.81	0.00	25.55	0.00	0.00	-20.00	0.00	16.16
129	ADHEr, ACK, GLCPs, NADH ₆	10.28	0.77	10.70	0.00	25.38	0.00	0.00	20.00	0.00	16.26
130	ADHEr, ACK, AKGId, ATPS4r	10.24	0.58	10.55	0.00	11.67	22.40	0.00	-20.00	0.00	36.95
131	ADHEr, ATPS4r, NADH ₆ , PFI	10.17	0.53	19.34	0.00	15.23	0.00	0.00	20.00	0.00	24.37
132	ADHEr, GLCPs, PFI, PPCK	10.15	0.43	23.16	0.00	15.20	0.00	0.00	-20.00	0.00	26.22
133	ADHEr, ACK, ATPS4r, SUCOAS	10.12	0.58	10.44	0.00	10.60	23.88	0.00	-20.00	0.00	38.45
134	ADHEr, ACK, ME2, NADH ₆	10.11	0.77	10.53	0.00	22.03	0.00	0.00	20.00	0.00	18.19
135	ADHEr, GLUTD ₂ , PFI, PPCK	10.10	0.45	22.94	0.00	15.12	0.00	0.00	-20.00	0.00	26.12
136	ADHEr, ME2, PFI, SUCD4	10.05	0.47	22.71	0.00	15.03	0.00	0.00	20.00	0.00	26.02

5
10
15
20
25
30
35
40
45
50
55

137	ADHEr, MDL, NADH6, PFL	10.04	0.47	22.67	0.00	15.02	0.00	0.00	0.00	20.00	0.00	26.01
138	ADHEr, PFL, PCK, RPE	10.02	0.50	22.14	0.00	15.00	0.00	0.00	0.00	20.00	0.00	25.50
139	ADHEr, PFL, PCK, TAI	9.99	0.50	22.22	0.00	14.95	0.00	0.00	0.00	20.00	0.00	25.76
140	ADHEr, GLADy, PFL, SUCD4	9.98	0.49	22.45	0.00	14.95	0.00	0.00	0.00	20.00	0.00	25.91
141	ADHEr, CBMK2, PFL, PCK	9.96	0.49	22.36	0.00	14.92	0.00	0.00	0.00	20.00	0.00	25.87
142	ADHEr, ATPS4r, MDL, SUCD4	9.92	0.57	20.45	0.00	15.84	0.00	0.00	0.00	20.00	0.00	24.48
143	ADHEr, PFL, RPE, SUCD4	9.90	0.54	21.57	0.00	14.81	0.00	0.00	0.00	20.00	0.00	25.47
144	ADHEr, AC Kt, CBMK2, NADH6	9.86	0.85	10.32	0.00	24.74	0.00	0.00	0.00	20.00	0.00	16.34
145	ADHEr, PFL, SUCD4, TAI	9.86	0.54	21.64	0.00	14.75	0.00	0.00	0.00	20.00	0.00	25.50
146	ADHEr, AC Kt, NADH6, RPE	9.86	0.85	10.32	0.00	24.73	0.00	0.00	0.00	20.00	0.00	16.34
147	ADHEr, AC Kt, TUM, NADH6	9.84	0.85	10.30	0.00	24.70	0.00	0.00	0.00	20.00	0.00	16.35
148	ADHEr, AC Kt, NADH6, TAI	9.85	0.85	10.30	0.00	24.70	0.00	0.00	0.00	20.00	0.00	16.35
149	ADHEr, AC Kt, ASSS2, NADH6	9.85	0.85	10.29	0.00	24.69	0.00	0.00	0.00	20.00	0.00	16.35
150	ADHEr, CBMK2, PFL, SUCD4	9.83	0.54	21.78	0.00	14.70	0.00	0.00	0.00	20.00	0.00	25.62
151	ADHEr, AC Kt, NADH6, NADH6	9.83	0.85	10.29	0.00	24.68	0.00	0.00	0.00	20.00	0.00	16.36
152	ADHEr, AC Kt, NADH6, S04r2	9.82	0.85	10.28	0.00	24.67	0.00	0.00	0.00	20.00	0.00	16.36
153	ADHEr, NADH6, NADH6, PFL	9.81	0.55	21.71	0.00	14.65	0.00	0.00	0.00	20.00	0.00	25.59
154	ADHEr, TUM, NADH6, PFL	9.80	0.55	21.69	0.00	14.67	0.00	0.00	0.00	20.00	0.00	25.58
155	ADHEr, AC Kt, PGL, SUCD4	9.80	0.22	9.92	0.00	12.33	0.00	0.00	0.00	20.00	0.00	30.56
156	ADHEr, NADH6, PFL, TAI	9.75	0.58	21.19	0.00	14.60	0.00	0.00	0.00	20.00	0.00	25.30
157	ADHEr, CBMK2, NADH6, PFL	9.72	0.58	21.33	0.00	14.55	0.00	0.00	0.00	20.00	0.00	25.13
158	ADHEr, TUM, HAXL, PFL	9.55	0.53	20.52	0.00	14.29	0.00	0.00	0.00	20.00	0.00	25.11
159	ADHEr, MDL, NADH6, THD2	9.34	0.55	19.65	0.00	11.38	0.00	0.00	0.00	20.00	0.00	36.73
160	ADHEr, ATPS4r, MDL, NADH6	9.28	0.55	19.75	0.00	7.20	0.00	0.00	0.00	20.00	0.00	37.01
161	ADHEr, ATPS4r, TUM, NADH6	9.03	0.63	21.23	0.00	11.62	0.00	0.00	0.00	20.00	0.00	29.45
162	ADHEr, ATPS4r, PGL, PCK	8.97	0.26	23.60	0.00	0.95	0.00	0.00	0.00	20.00	0.00	50.46
163	ADHEr, ASPT, MDL, NADH6	8.65	0.48	19.82	0.00	0.06	0.00	0.00	0.00	20.00	0.00	49.25
164	ADHEr, ATPS4r, NADH6, PCK	8.62	0.48	23.04	0.00	6.28	0.00	0.00	0.00	20.00	0.00	39.68
165	ADHEr, ASPT, MDL, THD2	8.60	0.50	28.60	0.00	11.82	0.00	0.00	0.00	20.00	0.00	51.55

5
10
15
20
25
30
35
40
45
50
55

166	ADHEr, ATPS4r, GL Cps, SUCD4	8.49	0.65	23.41	0.00	13.16	2.07	0.00	-20.00	0.00	28.11
167	ADHEr, ASP1, MDH, PGd	8.36	0.23	23.87	0.00	-4.10	33.24	0.00	-20.00	0.00	58.77
168	ADHEr, ASP1, FBA, MDH	8.27	0.24	24.01	0.00	4.27	33.31	0.00	-20.00	0.00	59.01
169	ADHEr, ASP1, MDH, PCK	8.27	0.73	24.01	0.00	4.27	33.31	0.00	-20.00	0.00	59.01
170	ADHEr, ASP1, MDH, TPI	8.27	0.24	24.01	0.00	-4.27	33.31	0.00	-20.00	0.00	59.01
171	ADHEr, ATPS4r, PCK, PCK	8.26	0.76	27.70	0.00	3.99	16.78	0.00	-20.00	0.00	46.33
172	ADHEr, ATPS4r, FBA, PCK	8.26	0.76	27.70	0.00	3.99	16.78	0.00	-20.00	0.00	46.33
173	ADHEr, ATPS4r, PCK, TPI	8.26	0.26	27.70	0.00	3.99	16.78	0.00	-20.00	0.00	46.33
174	ADHEr, ACKr, EDN, PGd	8.12	0.45	8.36	0.00	1.85	16.77	0.00	-20.00	0.00	42.04
175	ADHEr, ATPS4r, HENr, NADH6	7.99	0.67	23.56	0.00	6.75	16.40	0.00	-20.00	0.00	36.71
176	ADHEr, NADH6, PCK, THD2	7.88	0.50	23.49	0.00	2.67	18.21	0.00	-20.00	0.00	45.24
177	ADHEr, ATPS4r, GL Cps, MDH	7.87	0.16	22.87	0.00	10.07	23.70	0.00	-20.00	0.00	49.84
178	ADHEr, ATPS4r, MDH, PCK	7.86	0.49	22.44	0.00	10.04	23.43	0.00	-20.00	0.00	49.33
179	ADHEr, ATPS4r, GCM, PCK	7.86	0.49	23.44	0.00	0.04	23.43	0.00	-20.00	0.00	49.33
180	ADHEr, ENOr, NADH6, RPE	7.84	0.08	30.83	0.00	10.05	13.42	0.00	-20.00	0.00	44.85
181	ADHEr, NADH6, PGM, RPE	7.83	0.08	30.83	0.00	10.05	13.42	0.00	-20.00	0.00	44.85
182	ADHEr, NADH6, PGM, TAC	7.83	0.08	30.84	0.00	10.02	13.45	0.00	-20.00	0.00	44.90
183	ADHEr, ENOr, NADH6, TAC	7.83	0.08	30.84	0.00	10.02	13.45	0.00	-20.00	0.00	44.90
184	ADHEr, ASP1, GL Cps, MDH	7.76	0.46	21.57	0.00	-4.11	31.25	0.00	-20.00	0.00	56.07
185	ADHEr, ASP1, MDH, RPE	7.65	0.52	20.47	0.00	3.75	30.40	0.00	-20.00	0.00	54.60
186	ADHEr, ASP1, GL Cps, MDH	7.65	0.47	21.39	0.00	4.10	31.16	0.00	-20.00	0.00	55.85
187	ADHEr, ME2, NADH6, THD2	7.62	0.87	13.24	0.00	18.87	0.00	0.00	-20.00	0.00	23.43
188	ADHEr, ME2, SUCD4, THD2	7.61	0.52	22.02	0.00	-0.99	24.76	0.00	-20.00	0.00	50.51
189	ADHEr, ASP1, MDH, TAC	7.58	0.53	20.61	0.00	3.90	30.48	0.00	-20.00	0.00	54.83
190	ADHEr, NADH6, PGL, PCK	7.58	0.23	28.71	0.00	0.18	22.36	0.00	-20.00	0.00	52.73
191	ADHEr, TUNr, PCK, THD2	7.55	0.49	22.66	0.00	-1.33	25.25	0.00	-20.00	0.00	57.43
192	ADHEr, MDH, PCK, THD2	7.55	0.29	23.66	0.00	-1.33	25.25	0.00	-20.00	0.00	57.43
193	ADHEr, GL Cps, MDH, THD2	7.56	0.37	23.10	0.00	-1.46	25.55	0.00	-20.00	0.00	57.99
194	ADHEr, ASP1, CBNK3, MDH	7.52	0.52	20.82	0.00	-4.06	30.61	0.00	-20.00	0.00	55.15

195	ADHEr, ATPS4r, FBA, SUCD4	7.51	0.43	25.99	0.00	6.86	13.76	0.00	0.00	0.00	42.72
196	ADHEr, ATPS4r, PEK, SUCD4	7.51	0.42	25.99	0.00	6.86	13.76	0.00	0.00	0.00	42.72
197	ADHEr, ATPS4r, SUCD4, TPI	7.51	0.43	25.99	0.00	6.86	13.76	0.00	0.00	0.00	42.72
198	ADHEr, FBA, NADH6, PEK	7.50	0.24	28.80	0.00	6.03	22.54	0.00	0.00	0.00	53.03
199	ADHEr, NADH6, PEK, PECK	7.50	0.24	28.80	0.00	6.03	22.54	0.00	0.00	0.00	53.03
200	ADHEr, NADH6, PECK, TPI	7.50	0.24	28.80	0.00	6.03	22.54	0.00	0.00	0.00	53.03
201	ADHEr, HENT, PECK, THD2	7.47	0.69	14.32	0.00	8.03	0.00	0.00	0.00	0.00	31.52
202	ADHEr, HENT, NADH6, PEK	7.46	0.26	28.53	0.00	6.04	22.42	0.00	0.00	0.00	52.78
203	ADHEr, FBA, HENT, NADH6	7.46	0.26	28.53	0.00	6.03	22.42	0.00	0.00	0.00	52.78
204	ADHEr, HENT, NADH6, TPI	7.46	0.26	28.53	0.00	6.04	22.42	0.00	0.00	0.00	52.78
205	ADHEr, ATPS4r, G6PDHr, MDH	7.44	0.58	13.04	0.00	6.96	0.00	0.00	0.00	0.00	33.76
206	ADHEr, ATPS4r, MDH, PGI	7.44	0.58	13.04	0.00	6.96	0.00	0.00	0.00	0.00	33.76
207	ADHEr, GLUTDy, NADH6, PGI	7.43	0.57	27.55	0.00	6.24	17.78	0.00	0.00	0.00	51.58
208	ADHEr, ACKr, FRD2, TPI, D	7.42	0.97	7.95	0.00	15.46	13.33	0.00	0.00	0.00	29.24
209	ADHEr, ACKr, TPI, D, SUCD4	7.42	0.97	7.95	2.12	13.46	13.33	0.00	0.00	0.00	30.30
210	ADHEr, ATPS4r, FLM, GLUTDy	7.38	0.54	24.75	0.00	2.27	17.14	0.00	0.00	0.00	45.22
211	ADHEr, NADH6, PEK, RPE	7.34	0.36	27.05	0.00	6.21	21.57	0.00	0.00	0.00	51.14
212	ADHEr, FBA, NADH6, RPE	7.34	0.36	27.05	0.00	6.21	21.57	0.00	0.00	0.00	51.14
213	ADHEr, NADH6, RPE, TPI	7.34	0.36	27.05	0.00	6.21	21.57	0.00	0.00	0.00	51.14
214	ADHEr, FBA, GLUTDy, NADH6	7.33	0.32	27.69	0.00	0.04	22.04	0.00	0.00	0.00	52.00
215	ADHEr, GLUTDy, NADH6, PEK	7.33	0.32	27.69	0.00	0.04	22.04	0.00	0.00	0.00	52.00
216	ADHEr, GLUTDy, NADH6, TPI	7.34	0.32	27.69	0.00	0.04	22.04	0.00	0.00	0.00	52.00
217	ADHEr, ATPS4r, FLM, HENT	7.31	0.66	21.12	0.00	0.90	20.04	0.00	0.00	0.00	45.88
218	ADHEr, NADH6, TPI, TPI	7.30	0.36	27.10	0.00	0.09	21.68	0.00	0.00	0.00	51.32
219	ADHEr, NADH6, PEK, TPI	7.30	0.36	27.10	0.00	0.09	21.68	0.00	0.00	0.00	51.32
220	ADHEr, FBA, NADH6, TPI	7.30	0.36	27.10	0.00	0.09	21.68	0.00	0.00	0.00	51.32
221	ADHEr, ATPS4r, MDH, THD2	7.27	0.62	19.64	0.00	1.10	22.88	0.00	0.00	0.00	49.00
222	ADHEr, GLUTDy, MDH, NADH6	7.06	0.45	15.98	0.00	0.06	21.26	0.00	0.00	0.00	50.42
223	ADHEr, G6Pds, NADH6, PECK	7.06	0.45	15.94	0.00	0.06	21.24	0.00	0.00	0.00	50.38

5
10
15
20
25
30
35
40
45
50
55

224	ADHEP, NADH6, PPCK, RPE	7.05	0.52	24.82	0.00	0.30	20.47	0.00	-20.00	0.00	-8.06
225	ADHEP, GLUTD, NADH6, PPCK	7.02	0.47	25.73	0.00	-0.06	21.14	0.00	-20.00	0.00	50.18
226	ADHEP, FUM, NADH6, PPCK	7.01	0.37	25.63	0.00	-0.09	21.10	0.00	-20.00	0.00	50.09
227	ADHEP, MDL, NADH6, PPCK	7.01	0.37	25.63	0.00	-0.06	21.10	0.00	-20.00	0.00	50.09
228	ADHEP, ATPS4, FRO2, THD2	7.00	0.32	0.00	0.00	17.78	8.35	0.00	-20.00	0.00	21.33
229	ADHEP, NADH6, PPCK, FAL	6.98	0.52	24.92	0.00	0.12	20.63	0.00	-20.00	0.00	49.23
230	ADHEP, FUM, GLUTD, NADH6	6.92	0.52	25.04	0.00	-0.07	20.83	0.00	-20.00	0.00	49.55
231	ADHEP, GLCP6, MDL, NADH6	6.91	0.52	24.95	0.00	0.07	20.89	0.00	-20.00	0.00	49.49
232	ADHEP, MDL, NADH6, RPE	6.91	0.59	23.79	0.00	0.34	19.96	0.00	-20.00	0.00	47.94
233	ADHEP, MDL, NADH6, FAL	6.84	0.59	23.91	0.00	0.34	20.16	0.00	-20.00	0.00	48.26
234	ADHEP, HEN, NADH6, THD2	6.83	0.92	18.08	0.00	17.46	1.07	0.00	-20.00	0.00	28.73
235	ADHEP, FUM, NADH6, RPE	6.77	0.66	22.78	0.00	0.38	19.37	0.00	-20.00	0.00	46.96
236	ADHEP, CBMK2, MDL, NADH6	6.77	0.59	24.11	0.00	0.08	20.49	0.00	-20.00	0.00	48.68
237	ADHEP, FUM, ME2, NADH6	6.76	0.59	24.04	0.00	0.08	20.37	0.00	-20.00	0.00	48.62
238	ADHEP, FUM, NADH6, FAL	6.69	0.66	22.89	0.00	0.16	19.68	0.00	-20.00	0.00	47.29
239	ADHEP, ATPS4, MDL, PGD1	6.64	0.60	16.99	0.00	6.32	0.00	0.00	-20.00	0.00	35.28
240	ADHEP, FUM, HEN, NADH6	6.64	0.65	23.23	0.00	0.09	20.67	0.00	-20.00	0.00	47.87
241	ADHEP, CBMK2, FUM, NADH6	6.61	0.66	23.09	0.00	-0.09	19.94	0.00	-20.00	0.00	47.74
242	ADHEP, GLCP6, PPCK, THD2	6.56	0.39	24.86	0.00	-3.91	27.44	0.00	-20.00	0.00	55.80
243	ADHEP, GLUTD, PPCK, THD2	6.56	0.50	24.67	0.00	3.84	27.31	0.00	-20.00	0.00	55.57
244	ADHEP, CBMK2, NADH6, FAL	6.49	0.77	21.51	0.00	5.18	24.02	0.00	-20.00	0.00	30.97
245	ADHEP, ATPS4, MDL, FAL	6.48	0.61	16.56	0.00	5.95	0.00	0.00	-20.00	0.00	35.75
246	ADHEP, ATPS4, GLUTD, NADH6	6.45	0.65	22.52	0.00	1.91	16.38	0.00	-20.00	0.00	45.54
247	ADHEP, PGL, PPCK, SC94	6.43	0.22	30.11	0.00	-5.16	31.28	0.00	-20.00	0.00	62.99
248	ADHEP, ATPS4, MDL, RPE	6.33	0.61	16.77	0.00	5.52	0.00	0.00	-20.00	0.00	36.25
249	ADHEP, FBP, PGM, THD2	6.33	0.03	33.24	0.00	-6.19	33.35	0.00	-20.00	0.00	66.75
250	ADHEP, FNO, FBP, THD2	6.33	0.03	33.21	0.00	-6.19	33.35	0.00	-20.00	0.00	66.75
251	ADHEP, FBA, PGM, THD2	6.33	0.03	33.21	0.00	-6.19	33.35	0.00	-20.00	0.00	66.75
252	ADHEP, FNO, FBA, THD2	6.33	0.03	33.21	0.00	-6.19	33.35	0.00	-20.00	0.00	66.75

5
10
15
20
25
30
35
40
45
50
55

253	ADHET, PNO, THD2, TPI	6.33	0.03	33.23	0.00	-6.19	33.35	0.00	-20.00	0.00	66.75
254	ADHET, PGM, THD2, TPI	6.33	0.03	33.23	0.00	-6.19	33.35	0.00	-20.00	0.00	66.75
255	ADHET, GLCps, PGL, PPCK	6.28	0.26	29.59	0.00	-6.08	30.96	0.00	-20.00	0.00	62.47
256	ADHET, PBA, HEXT, PPCK	6.38	0.23	30.23	0.00	6.37	31.47	0.00	-20.00	0.00	63.24
257	ADHET, HEXT, PFK, PPCK	6.28	0.22	30.24	0.00	-6.37	31.47	0.00	-20.00	0.00	63.34
258	ADHET, HEXT, PPCK, TPI	6.28	0.22	30.24	0.00	6.37	31.47	0.00	-20.00	0.00	63.24
259	ADHET, PPCK, SUCD4, TPI	6.27	0.23	30.17	0.00	-6.30	31.37	0.00	-20.00	0.00	63.17
260	ADHET, PFK, PPCK, SUCD4	6.27	0.23	30.17	0.00	-6.30	31.37	0.00	-20.00	0.00	63.17
261	ADHET, PBA, PPCK, SUCD4	6.37	0.23	30.17	0.00	6.30	31.37	0.00	-20.00	0.00	63.17
262	ADHET, GLUDy, PGL, PPCK	6.26	0.27	29.44	0.00	6.06	30.86	0.00	-20.00	0.00	62.25
263	ADHET, GLCps, PGL, SUCD4	6.24	0.28	29.26	0.00	-6.03	30.74	0.00	-20.00	0.00	62.02
264	ADHET, PUN, GLUDy, PGL	6.24	0.29	29.24	0.00	-6.02	30.73	0.00	-20.00	0.00	62.03
265	ADHET, GLUDy, MDL, PGL	6.24	0.29	29.24	0.00	6.02	30.73	0.00	-20.00	0.00	62.00
266	ADHET, PBA, HEXT, SUCD4	6.24	0.28	29.95	0.00	-6.27	31.23	0.00	-20.00	0.00	62.93
267	ADHET, HEXT, SUCD4, TPI	6.24	0.28	29.95	0.00	6.27	31.23	0.00	-20.00	0.00	62.93
268	ADHET, HEXT, PFK, SUCD4	6.24	0.25	29.95	0.00	-6.27	31.25	0.00	-20.00	0.00	62.93
269	ADHET, GLUDy, PGL, SUCD4	6.23	0.29	29.14	0.00	-6.07	30.67	0.00	-20.00	0.00	61.88
270	ADHET, PUN, HEXT, THD2	6.21	0.59	27.25	0.00	3.64	25.85	0.00	-20.00	0.00	52.95
271	ADHET, PBA, GLCps, PPCK	6.21	0.26	29.69	0.00	-6.24	31.07	0.00	-20.00	0.00	62.64
272	ADHET, GLCps, PFK, PPCK	6.21	0.26	29.69	0.00	6.24	31.07	0.00	-20.00	0.00	62.64
273	ADHET, GLCps, PPCK, TPI	6.21	0.26	29.69	0.00	6.24	31.07	0.00	-20.00	0.00	62.64
274	ADHET, PFK, PPCK, RPE	6.20	0.30	29.03	0.00	6.02	30.60	0.00	-20.00	0.00	61.77
275	ADHET, PPCK, RPE, TPI	6.20	0.30	29.01	0.00	-6.02	30.60	0.00	-20.00	0.00	61.77
276	ADHET, PBA, PPCK, RPE	6.20	0.30	29.01	0.00	-6.02	30.60	0.00	-20.00	0.00	61.77
277	ADHET, GLUDy, PFK, PPCK	6.19	0.27	29.54	0.00	-6.23	30.98	0.00	-20.00	0.00	62.47
278	ADHET, GLUDy, PPCK, TPI	6.19	0.27	29.54	0.00	-6.23	30.98	0.00	-20.00	0.00	62.47
279	ADHET, PBA, GLUDy, PPCK	6.19	0.27	29.54	0.00	6.23	30.98	0.00	-20.00	0.00	62.47
280	ADHET, ADPS-g, M12, THD2	6.19	0.77	21.39	0.00	-2.37	25.20	0.00	-20.00	0.00	57.73
281	ADHET, HEXT, PFK, RPE	6.18	0.32	28.82	0.00	-5.99	30.48	0.00	-20.00	0.00	61.55

5
10
15
20
25
30
35
40
45
50
55

282	ADHEr, HEXT, RPE, TPI	6.18	0.32	28.82	0.00	-5.99	30.48	0.00	-20.00	0.00	61.55
283	ADHEr, FBA, HEXT, RPE	6.18	0.32	28.82	0.00	-5.99	30.48	0.00	-20.00	0.00	61.55
284	ADHEr, FBA, PPK, TAL	6.17	0.31	29.05	0.00	6.09	30.65	0.00	-20.00	0.00	61.87
285	ADHEr, PPK, TAL, TPI	6.17	0.31	29.05	0.00	-6.09	30.65	0.00	-20.00	0.00	61.87
286	ADHEr, PPK, PPK, TAL	6.17	0.31	29.05	0.00	-6.09	30.65	0.00	-20.00	0.00	61.87
287	ADHEr, ATPSr, GLUDy, PPK	6.17	0.55	24.36	0.00	-5.40	27.23	0.00	-20.00	0.00	55.50
288	ADHEr, FBA, GLUDy, HEXT	6.17	0.29	29.36	0.00	-6.20	30.87	0.00	-20.00	0.00	62.28
289	ADHEr, GLUDy, HEXT, TPI	6.17	0.29	29.36	0.00	-6.20	30.87	0.00	-20.00	0.00	62.28
290	ADHEr, GLUDy, HEXT, PPK	6.17	0.29	29.36	0.00	6.20	30.87	0.00	-20.00	0.00	62.28
291	ADHEr, GLCps, SUCD4, TPI	6.16	0.29	29.36	0.00	-6.20	30.86	0.00	-20.00	0.00	62.27
292	ADHEr, FBA, GLCps, SUCD4	6.16	0.29	29.36	0.00	-6.20	30.86	0.00	-20.00	0.00	62.27
293	ADHEr, GLCps, PPK, SUCD4	6.16	0.29	29.36	0.00	6.20	30.85	0.00	-20.00	0.00	62.25
294	ADHEr, GLUDy, MDH, PPK	6.16	0.29	29.36	0.00	6.20	30.85	0.00	-20.00	0.00	62.25
295	ADHEr, FBA, GLUDy, MDH	6.16	0.29	29.36	0.00	6.20	30.85	0.00	-20.00	0.00	62.25
296	ADHEr, FBA, FUM, GLUDy	6.16	0.29	29.36	0.00	6.20	30.85	0.00	-20.00	0.00	62.25
297	ADHEr, FUM, GLUDy, PPK	6.16	0.29	29.36	0.00	-6.20	30.85	0.00	-20.00	0.00	62.25
298	ADHEr, GLUDy, MDH, TPI	6.16	0.29	29.36	0.00	-6.20	30.85	0.00	-20.00	0.00	62.25
299	ADHEr, FUM, GLUDy, TPI	6.16	0.29	29.36	0.00	6.20	30.85	0.00	-20.00	0.00	62.25
300	ADHEr, RPE, SUCD4, TPI	6.16	0.33	28.65	0.00	-5.97	30.57	0.00	-20.00	0.00	61.36
301	ADHEr, PPK, RPE, SUCD4	6.16	0.33	28.65	0.00	5.97	30.57	0.00	-20.00	0.00	61.36
302	ADHEr, FBA, RPE, SUCD4	6.16	0.33	28.65	0.00	5.97	30.57	0.00	-20.00	0.00	61.36
303	ADHEr, GLUDy, PPK, SUCD4	6.15	0.30	29.25	0.00	-6.19	30.80	0.00	-20.00	0.00	62.15
304	ADHEr, FBA, GLUDy, SUCD4	6.15	0.30	29.25	0.00	6.19	30.80	0.00	-20.00	0.00	62.15
305	ADHEr, GLUDy, SUCD4, TPI	6.15	0.30	29.25	0.00	-6.19	30.80	0.00	-20.00	0.00	62.15
306	ADHEr, HEXT, PPK, TAL	6.14	0.32	28.86	0.00	6.07	30.53	0.00	-20.00	0.00	61.65
307	ADHEr, FBA, HEXT, TAL	6.14	0.32	28.86	0.00	6.07	30.53	0.00	-20.00	0.00	61.65
308	ADHEr, HEXT, TAL, TPI	6.14	0.32	28.86	0.00	-6.07	30.53	0.00	-20.00	0.00	61.65
309	ADHEr, PPK, SUCD4, TAL	6.13	0.33	28.70	0.00	-6.05	30.43	0.00	-20.00	0.00	61.47
310	ADHEr, FBA, SUCD4, TAL	6.13	0.33	28.70	0.00	6.05	30.43	0.00	-20.00	0.00	61.47

5
10
15
20
25
30
35
40
45
50
55

311	ADHET, SUCD4, IAL, IPI	6.13	0.33	28.70	0.00	-6.05	30.43	0.00	20.00	0.00	61.47
312	ADHET, FUM, ME2, IHD2	6.10	0.62	14.90	0.00	3.07	18.16	0.00	20.00	0.00	59.89
313	ADHET, AIPSE2, ME2, SUCD4	6.04	0.65	18.68	0.00	0.48	22.08	0.00	20.00	0.00	48.37
314	ADHET, PPCK, PYK, SUCD4	6.00	0.38	28.14	0.00	-6.05	30.04	0.00	20.00	0.00	64.81
315	ADHET, GLCps, PPCK, SUCD4	5.97	0.40	27.84	0.00	6.03	30.92	0.00	20.00	0.00	64.50
316	ADHET, PPCK, RPE, SUCD4	5.95	0.46	26.74	0.00	-5.68	29.16	0.00	20.00	0.00	59.19
317	ADHET, FUM, GLUTDy, PPCK	5.94	0.42	27.58	0.00	6.00	29.76	0.00	20.00	0.00	64.30
318	ADHET, GLUTDy, MDH, PPCK	5.94	0.43	27.58	0.00	6.00	29.76	0.00	20.00	0.00	64.40
319	ADHET, GLUTDy, PPCK, SUCD4	5.94	0.42	27.54	0.00	-5.99	29.74	0.00	20.00	0.00	64.27
320	ADHET, PPCK, SUCD4, IAL	5.90	0.46	26.81	0.00	5.79	29.24	0.00	20.00	0.00	59.36
321	ADHET, GLCps, GLUTDy, MDH	5.89	0.45	27.19	0.00	5.95	29.52	0.00	20.00	0.00	59.87
322	ADHET, GLCps, PPCK, RPE	5.89	0.51	26.14	0.00	-5.89	28.78	0.00	20.00	0.00	58.51
323	ADHET, GLUTDy, MDH, RPE	5.88	0.51	26.17	0.00	5.59	28.77	0.00	20.00	0.00	58.49
324	ADHET, GLUTDy, PPCK, RPE	5.87	0.52	26.00	0.00	5.57	28.69	0.00	20.00	0.00	58.35
325	ADHET, GLCps, GLUTDy, PPCK	5.87	0.46	27.02	0.00	5.94	29.42	0.00	20.00	0.00	59.99
326	ADHET, GLCps, MDH, SUCD4	5.86	0.46	26.97	0.00	-5.93	29.38	0.00	20.00	0.00	59.93
327	ADHET, MDH, RPE, SUCD4	5.85	0.53	25.81	0.00	5.54	28.57	0.00	20.00	0.00	58.14
328	ADHET, GLCps, MDH, PPCK	5.85	0.47	26.84	0.00	-5.91	29.29	0.00	20.00	0.00	59.47
329	ADHET, FUM, GLCps, PPCK	5.85	0.47	26.83	0.00	5.91	29.29	0.00	20.00	0.00	59.47
330	ADHET, MDH, PPCK, RPE	5.84	0.54	25.70	0.00	-5.53	28.50	0.00	20.00	0.00	58.02
331	ADHET, FUM, PPCK, RPE	5.84	0.54	25.70	0.00	-5.53	28.50	0.00	20.00	0.00	58.02
332	ADHET, GLCps, PPCK, IAL	5.84	0.51	26.23	0.00	5.72	28.88	0.00	20.00	0.00	58.71
333	ADHET, GLUTDy, MDH, IAL	5.83	0.51	26.21	0.00	-5.71	28.86	0.00	20.00	0.00	58.68
334	ADHET, MDH, PPCK, PYK	5.82	0.47	26.84	0.00	-5.98	29.28	0.00	20.00	0.00	59.61
335	ADHET, FUM, PPCK, PYK	5.82	0.47	26.84	0.00	5.98	29.28	0.00	20.00	0.00	59.61
336	ADHET, GLUTDy, PPCK, IAL	5.82	0.52	26.09	0.00	-5.70	28.79	0.00	20.00	0.00	58.55
337	ADHET, FUM, GLUTDy, SUCD4	5.81	0.49	26.54	0.00	-5.87	29.10	0.00	20.00	0.00	59.17
338	ADHET, MDH, SUCD4, IAL	5.80	0.53	25.90	0.00	5.67	28.67	0.00	20.00	0.00	58.43
339	ADHET, GLCps, ME2, SUCD4	5.80	0.50	26.45	0.00	-5.86	29.06	0.00	20.00	0.00	59.05

5
10
15
20
25
30
35
40
45
50
55

340	ADHEP, MF2, RPE, SUCD4	5.79	0.57	25.28	0.00	-5.46	28.24	0.00	-20.00	0.00	57.53
341	ADHEP, FUM, PPCK, FAL	5.79	0.54	25.79	0.00	-5.66	28.61	0.00	-20.00	0.00	58.22
342	ADHEP, MDL, PPCK, FAL	5.79	0.54	25.79	0.00	-5.66	28.61	0.00	-20.00	0.00	58.22
343	ADHEP, GLUDy, PROIZ, SUCD4	5.78	0.51	26.35	0.00	-5.85	29.00	0.00	-20.00	0.00	58.95
344	ADHEP, FUM, GLUDy, RPE	5.78	0.57	25.18	0.00	-5.45	28.18	0.00	-20.00	0.00	57.43
345	ADHEP, GLUDy, RPE, SUCD4	5.78	0.57	25.17	0.00	-5.45	28.17	0.00	-20.00	0.00	57.41
346	ADHEP, GLCPis, RPE, SUCD4	5.78	0.57	25.15	0.00	-5.44	28.15	0.00	-20.00	0.00	57.39
347	ADHEP, FUM, GLUDy, MF2	5.78	0.51	26.40	0.00	-5.85	28.97	0.00	-20.00	0.00	58.89
348	ADHEP, GLUDy, MD2, SUCD4	5.78	0.51	26.30	0.00	-5.85	28.96	0.00	-20.00	0.00	58.88
349	ADHEP, FUM, GLCPis, GLUDy	5.77	0.52	26.32	0.00	-5.84	28.91	0.00	-20.00	0.00	58.80
350	ADHEP, GLCPis, GLUDy, SUCD4	5.76	0.52	26.17	0.00	-5.83	28.88	0.00	-20.00	0.00	58.74
351	ADHEP, FUM, MF2, SUCD4	5.74	0.55	25.99	0.00	-5.81	28.78	0.00	-20.00	0.00	58.55
352	ADHEP, MF2, SUCD4, FAL	5.74	0.57	25.38	0.00	-5.60	28.34	0.00	-20.00	0.00	57.75
353	ADHEP, FUM, GLUDy, FAL	5.72	0.58	25.26	0.00	-5.59	28.27	0.00	-20.00	0.00	57.62
354	ADHEP, GLUDy, SUCD4, FAL	5.72	0.58	25.35	0.00	-5.58	28.26	0.00	-20.00	0.00	57.61
355	ADHEP, GLCPis, SUCD4, FAL	5.72	0.58	25.21	0.00	-5.58	28.24	0.00	-20.00	0.00	57.57
356	ADHEP, MD2, PGGL, IHD2	5.71	0.95	0.00	0.00	6.29	15.87	0.00	20.00	0.00	40.76
357	ADHEP, G6PDy, MF2, IHD2	5.71	0.95	0.00	0.00	6.29	15.82	0.00	-20.00	0.00	40.76
358	ADHEP, FUM, RPE, SUCD4	5.71	0.62	24.48	0.00	-5.35	27.73	0.00	-20.00	0.00	56.63
359	ADHEP, IHDy, RPE, SUCD4	5.69	0.63	24.34	0.00	-5.32	27.64	0.00	-20.00	0.00	56.47
360	ADHEP, CBMK2, GLUSK, PPCK	5.68	0.57	25.81	0.00	-5.76	28.48	0.00	-20.00	0.00	58.07
361	ADHEP, CBMK2, GSSD, PPCK	5.68	0.57	25.51	0.00	-5.76	28.48	0.00	-20.00	0.00	58.02
362	ADHEP, ASSS, CBMK2, PPCK	5.68	0.57	25.51	0.00	-5.76	28.48	0.00	-20.00	0.00	58.02
363	ADHEP, CBMK2, PPCK, SO4L2	5.68	0.57	25.50	0.00	-5.75	28.47	0.00	-20.00	0.00	58.01
364	ADHEP, GLUDy, IHDy, SUCD4	5.67	0.57	25.42	0.00	-5.75	28.42	0.00	-20.00	0.00	57.92
365	ADHEP, FUM, SUCD4, FAL	5.64	0.63	24.56	0.00	-5.49	27.83	0.00	-20.00	0.00	56.84
366	ADHEP, IHDy, SUCD4, FAL	5.63	0.63	24.44	0.00	-5.48	27.78	0.00	-20.00	0.00	56.79
367	ADHEP, FUM, IHDy, SUCD4	5.62	0.60	25.62	0.00	-5.70	28.77	0.00	-20.00	0.00	57.47
368	ADHEP, FUM, IHDy, RPE	5.58	0.70	24.35	0.00	-5.18	27.62	0.00	-20.00	0.00	55.35

5
10
15
20
25
30
35
40
45
50
55

369	ADHER. CBMK2, FUM, SUC, D5	5.58	0.62	24.75	2.00	-5.67	27.99	0.00	-20.00	0.00	57.15
370	ADHER. HENI, PELL, RPE	5.54	0.72	14.12	0.00	3.77	0.00	0.00	20.00	0.00	37.22
371	ADHER. FUM, HENI, TAI	5.51	0.70	23.47	0.00	5.35	27.15	0.00	-20.00	0.00	58.62
372	ADHER. CBMK2, FUM, HENI	5.45	0.70	23.69	0.00	-5.54	27.34	0.00	20.00	0.00	56.03
373	ADHER. HENI, PELL, TAI	5.40	0.72	14.11	0.00	3.45	0.00	0.00	-20.00	0.00	37.64
374	ADHER. GLYCL, HE NI, PELL	5.25	0.72	14.16	0.00	3.09	0.00	0.00	-20.00	0.00	38.10
375	ADHER. ATPS4r, GLUDY, PGI	5.14	0.46	23.54	0.00	-4.59	29.95	0.00	20.00	0.00	60.78
376	ADHER. PELL, PGDI, PGI	5.08	0.43	16.39	0.00	1.66	0.00	0.00	20.00	0.00	43.14
377	ADHER. PELL, PGI, TAI	5.05	0.43	16.21	0.00	1.57	0.00	0.00	20.00	0.00	43.21
378	ADHER. ATPS4r, PEK, RPE	5.03	0.52	26.76	0.00	-4.75	29.48	0.00	-20.00	0.00	59.94
379	ADHER. ATPS4r, FBA, RPE	5.03	0.52	26.76	0.00	-4.75	29.48	0.00	-20.00	0.00	59.94
380	ADHER. ATPS4r, RPE, TPI	5.03	0.52	26.76	0.00	-4.75	29.48	0.00	-20.00	0.00	59.94
381	ADHER. PELL, PGI, RPE	5.02	0.43	16.04	0.00	1.47	0.00	0.00	20.00	0.00	43.27
382	ADHER. ATPS4r, GLUDY, TPI	5.02	0.47	27.70	0.00	5.08	30.15	0.00	20.00	0.00	61.18
383	ADHER. ATPS4r, FBA, GLUDY	5.02	0.47	27.70	0.00	-5.08	30.15	0.00	-20.00	0.00	61.18
384	ADHER. ATPS4r, GLUDY, PEK	5.02	0.47	27.70	0.00	-5.08	30.15	0.00	-20.00	0.00	61.18
385	ADHER. FBA, PELL, PGI	5.01	0.44	15.95	0.00	1.43	0.00	0.00	20.00	0.00	43.31
386	ADHER. PEK, PELL, PGI	5.01	0.44	15.95	0.00	1.43	0.00	0.00	-20.00	0.00	43.31
387	ADHER. PELL, PGI, TPI	5.01	0.44	15.95	0.00	1.43	0.00	0.00	-20.00	0.00	43.31
388	ADHER. PELL, RPE, TPI	4.99	0.44	15.97	0.00	1.38	0.00	0.00	-20.00	0.00	43.37
389	ADHER. PEK, PELL, RPE	4.99	0.44	15.97	0.00	1.38	0.00	0.00	-20.00	0.00	43.37
390	ADHER. FBA, PELL, RPE	4.99	0.44	15.97	0.00	1.38	0.00	0.00	20.00	0.00	43.37
391	ADHER. ATPS4r, PEK, TAI	4.98	0.52	26.82	0.00	-4.85	29.56	0.00	-20.00	0.00	60.10
392	ADHER. ATPS4r, FBA, TAI	4.98	0.52	26.82	0.00	-4.85	29.56	0.00	-20.00	0.00	60.10
393	ADHER. ATPS4r, TAI, TPI	4.98	0.52	26.82	0.00	-4.85	29.56	0.00	20.00	0.00	60.10
394	ADHER. FBA, PELL, TAI	4.94	0.44	16.00	0.00	1.28	0.00	0.00	-20.00	0.00	43.49
395	ADHER. PELL, TAI, TPI	4.94	0.44	16.00	0.00	1.28	0.00	0.00	-20.00	0.00	43.49
396	ADHER. PEK, PELL, TAI	4.94	0.44	16.00	0.00	1.28	0.00	0.00	20.00	0.00	43.49
397	ADHER. GLYCL, HE NI, THERP	4.90	0.74	22.16	0.00	-6.57	26.62	0.00	-20.00	0.00	56.37

398	ADHET, GLUDY, PFK, PFI1	4.89	0.39	16.45	0.00	1.06	0.00	0.00	-20.00	0.00	44.29
399	ADHET, GLUDY, PFI1, TPI	4.89	0.39	16.45	0.00	1.06	0.00	0.00	20.00	0.00	44.29
400	ADHET, PBA, GLUDY, PFI1	4.89	0.39	16.45	0.00	1.06	0.00	0.00	20.00	0.00	44.29
401	ADHET, PBA, PFI1, PGI	4.83	0.50	0.00	0.00	-3.17	0.00	0.00	-20.00	0.00	45.11
402	ADHET, ATPS4r, GLUDY, RPI	4.72	0.72	23.95	0.00	4.53	27.69	0.00	20.00	0.00	56.73
403	ADHET, ATPS4r, GLUDY, TAI	4.65	0.72	24.01	0.00	-4.47	27.80	0.00	-20.00	0.00	56.96
404	ADHET, ATPS4r, CBMK2, RPI	4.60	0.80	22.80	0.00	-4.14	26.97	0.00	-20.00	0.00	55.45
405	ADHET, ATPS4r, CBMK2, GLUDY	4.57	0.72	24.19	0.00	-4.67	27.97	0.00	-20.00	0.00	57.30
406	ADHET, ATPS4r, CBMK2, TAI	4.51	0.80	22.85	0.00	-4.32	27.69	0.00	-20.00	0.00	55.69
407	ADHET, ASNS2, ATPS4r, GLU5K	4.47	0.81	23.95	0.00	4.53	27.20	0.00	20.00	0.00	55.95
408	ADHET, ASNS2, ATPS4r, GSND	4.42	0.81	22.95	0.00	-4.53	27.20	0.00	-20.00	0.00	55.95
409	ADHET, AC Kc, PFI1, PGI	3.90	0.50	3.27	0.00	-6.43	0.00	0.00	-20.00	0.00	50.40
410	ADHET, AC Kc, AKGD, PRD2	2.76	1.03	3.32	0.00	14.67	0.00	0.00	-20.00	0.00	25.87
411	ADHET, AC Kc, PRD2, SUCOAS	3.91	1.03	2.47	0.00	12.55	0.00	0.00	-20.00	0.00	27.45
412	ADHET, FUM, G6PDH, TAI	3.40	0.71	14.98	0.00	12.86	18.72	0.00	20.00	0.00	62.51
413	ADHET, FUM, PGDH, TAI	3.40	0.71	14.98	0.00	-12.86	18.72	0.00	-20.00	0.00	62.51
414	ADHET, FUM, PGL, TAI	3.40	0.71	14.98	0.00	-12.86	18.72	0.00	-20.00	0.00	62.51
415	ADHET, FUM, ME2, TPI	3.36	0.47	17.83	0.00	13.86	20.51	0.00	20.00	0.00	68.38
416	ADHET, FUM, ME2, PFK	3.36	0.47	17.83	0.00	-13.86	20.51	0.00	-20.00	0.00	68.38
417	ADHET, PBA, FUM, ME2	3.36	0.47	17.83	0.00	13.86	20.51	0.00	20.00	0.00	68.38
418	ADHET, PRD2, GLUDY, LIDI D	3.27	0.24	0.00	0.00	33.89	3.68	53.86	20.00	0.00	73.09
419	ADHET, PRD2, LIDI D, TPI	3.15	0.40	0.00	0.00	15.41	4.39	0.00	-20.00	0.00	22.97
420	ADHET, G6PDH, HEAN, TAI	3.01	0.73	14.02	0.00	-13.53	17.85	0.00	-20.00	0.00	62.75
421	ADHET, HEAN, PGDH, TAI	3.01	0.73	14.02	0.00	13.53	17.85	0.00	20.00	0.00	62.75
422	ADHET, HEAN, PGI, TAI	3.04	0.73	14.02	0.00	-13.53	17.85	0.00	-20.00	0.00	62.75
423	ADHET, MDH, PGDH, TAI	0.89	0.65	14.71	0.00	14.28	18.10	0.00	20.00	0.00	64.72
424	ADHET, G6PDH, MDH, TAI	0.89	0.65	14.71	0.00	-14.28	18.10	0.00	-20.00	0.00	64.72
425	ADHET, MDH, PGI, TAI	0.89	0.65	14.71	0.00	-14.28	18.10	0.00	-20.00	0.00	64.72
426	ADHET, PGDH, TAI, TPI	0.43	0.47	15.75	0.00	-16.25	18.18	0.00	20.00	0.00	69.35

5
10
15
20
25
30
35
40
45
50
55

427	ADH1c, FBA, PGDH, FAL	0.43	0.47	15.73	0.00	-16.28	18.18	0.00	20.00	0.00	69.35
428	ADH1c, PK, PGDH, FAL	0.43	0.47	15.73	0.00	16.28	18.18	0.00	20.00	0.00	69.35
429	ADH1c, GLXCL, FAL, TPI	0.41	0.49	15.57	0.00	16.70	18.12	0.00	-20.00	0.00	69.95
430	ADH1c, TAN, THDS, TPI	0.40	0.49	15.57	0.00	-16.69	18.11	0.00	-20.00	0.00	69.95
431	ADH1c, LDH D, FAL, TPI	0.40	0.49	15.57	0.00	16.69	18.11	0.00	20.00	0.00	69.95
432	ADH1c, ASPT, EDA, MDH, PGI	16.17	0.05	8.39	0.00	13.88	23.36	0.00	-20.00	0.00	33.47
433	ADH1c, APTS4, FRD2, LDH D, ME7	15.11	0.23	0.00	0.00	23.25	9.12	0.00	20.00	0.00	3.86
434	ADH1c, EDA, PPI, PGI, PPCK	14.76	0.16	16.11	0.00	22.13	0.00	0.00	20.00	0.00	17.24
435	ADH1c, APTS4, FRD2, LDH D, MDH	15.02	0.23	0.00	0.00	23.19	9.11	0.00	-20.00	0.00	3.93
436	ADH1c, EDA, PPI, PGI, STCD4	14.65	0.19	15.67	0.00	21.97	0.00	0.00	20.00	0.00	17.05
437	ADH1c, EDA, NADH6, PPI, PGI	14.65	0.19	15.67	0.00	21.97	0.00	0.00	-20.00	0.00	17.05
438	ADH1c, EDA, NADH6, PGI, PPCK	14.47	0.09	17.44	0.00	19.97	3.46	0.00	-20.00	0.00	21.55
439	ADH1c, ASPT, LDH D, MDH, PPI	14.38	0.36	13.01	0.00	21.55	0.00	0.00	20.00	0.00	15.56
440	ADH1c, GLDy, HE XI, PPI, PGI	14.31	0.31	14.25	0.00	21.45	0.00	0.00	-20.00	0.00	16.44
441	ADH1c, ACK, GLCps, NADH6, PGI	14.24	0.27	14.48	0.00	22.22	0.00	0.00	20.00	0.00	16.72
442	ADH1c, GLDy, HE XI, NADH6, PGI	14.13	0.25	15.36	0.00	19.91	2.53	0.00	20.00	0.00	19.65
443	ADH1c, EDA, GLDy, NADH6, PGI	14.07	0.27	15.01	0.00	19.91	2.48	0.00	-20.00	0.00	19.24
444	ADH1c, ACK, PPI, PGI, STCD4	14.04	0.19	14.14	0.00	21.04	0.00	0.00	20.00	0.00	17.69
445	ADH1c, ACK, NADH6, PPI, PGI	14.04	0.19	14.14	0.00	22.14	0.00	0.00	20.00	0.00	16.50
446	ADH1c, ACK, GLDy, NADH6, PGI	14.03	0.30	14.19	0.00	22.02	0.00	0.00	20.00	0.00	16.65
447	ADH1c, EDA, GLCps, NADH6, PGI	13.92	0.28	14.76	0.00	21.90	0.00	0.00	20.00	0.00	17.17
448	ADH1c, ACK, CBMK2, NADH6, PGI	13.94	0.34	14.12	0.00	21.89	0.00	0.00	-20.00	0.00	16.84
449	ADH1c, APTS4, FRD2, NADH6, PGI	13.86	0.22	14.56	0.00	21.75	0.00	0.00	20.00	0.00	17.62
450	ADH1c, APTS4, NADH6, PPI, PGI	13.86	0.22	14.56	0.00	21.75	0.00	0.00	-20.00	0.00	17.62
451	ADH1c, APTS4, GLCps, NADH6, PPI	13.80	0.27	16.36	0.00	20.68	0.00	0.00	20.00	0.00	18.31
452	ADH1c, APTS4, MDH, NADH6, PGI	13.69	0.43	13.35	0.00	20.51	0.00	0.00	20.00	0.00	16.43
453	ADH1c, APTS4, G6PDH, MDH, NADH6	13.69	0.43	13.35	0.00	20.51	0.00	0.00	20.00	0.00	16.43
454	ADH1c, ACK, PPI, GLDy, LDH D	13.68	0.36	13.87	0.00	18.43	4.12	0.00	-20.00	0.00	20.56
455	ADH1c, APTS4, NADH6, PGI, STCD4	13.66	0.25	15.79	0.00	19.61	3.75	0.00	20.00	0.00	21.47

5
10
15
20
25
30
35
40
45
50
55

456	ADHEC, ACK, GLUDy, LDIH_D, SUCD4	13.56	0.37	13.76	0.00	13.71	5.22	0.00	-20.00	0.00	21.50
457	ADHEC, AIPS4c, G6PDHy, MDH, THD2	13.53	0.44	10.63	0.00	14.18	12.16	0.00	-20.00	0.00	25.92
458	ADHEC, AIPS4c, MDH, PGL, THD2	13.53	0.44	10.63	0.00	14.18	17.16	0.00	20.00	0.00	25.92
459	ADHEC, ASPT, G6PDHy, MDH, PYK	13.44	0.26	11.28	0.00	7.24	25.83	0.00	-20.00	0.00	38.92
460	ADHEC, ASPT, FBA, MDH, PYK	13.44	0.26	11.28	0.00	7.24	25.83	0.00	20.00	0.00	38.92
461	ADHEC, ASPT, MDH, PGL, PYK	13.44	0.26	11.28	0.00	7.24	25.83	0.00	-20.00	0.00	38.92
462	ADHEC, FRD2, LDIH_D, MDH, SUCOAS	13.35	0.25	4.81	0.00	13.43	19.89	0.00	-20.00	0.00	28.60
463	ADHEC, ASPT, LDIH_D, MDH, SUCOAS	13.27	0.26	4.78	0.00	13.14	19.81	0.00	20.00	0.00	28.92
464	ADHEC, AC6, LDIH_D, MDH, SUCD4	13.22	0.26	0.00	0.00	17.54	14.32	0.00	-20.00	0.00	23.06
465	ADHEC, AIPS4c, GLUDy, PGL, SUCD4	13.17	0.24	16.40	0.00	17.06	7.36	0.00	-20.00	0.00	25.48
466	ADHEC, ASPT, FUM, LDIH_D, MDH	13.15	0.27	5.76	0.00	12.63	20.09	0.00	20.00	0.00	29.76
467	ADHEC, ASPT, LDIH_D, MAL'S, MDH	13.15	0.27	5.76	0.00	12.63	20.09	0.00	20.00	0.00	29.76
468	ADHEC, ASPT, ICT, LDIH_D, MDH	13.15	0.27	5.76	0.00	12.63	20.09	0.00	20.00	0.00	29.76
469	ADHEC, AC6, LDIH_D, MDH, NADH6	13.07	0.51	0.00	0.00	23.45	0.00	0.00	-20.00	0.00	13.57
470	ADHEC, FRD2, GLUDy, LDIH_D, PPC	12.91	0.12	12.29	0.00	0.02	38.75	0.00	20.00	0.00	51.90
471	ADHEC, FRD2, LDIH_D, PPC, THD2	12.89	0.14	12.25	0.00	-0.02	38.70	0.00	-20.00	0.00	51.85
472	ADHEC, ACK, AIPS4c, LDIH_D, SUCD4	12.73	0.47	12.99	0.00	16.72	6.70	0.00	20.00	0.00	24.06
473	ADHEC, ACK, ACS, PPC, PPC	12.62	0.11	12.68	0.00	0.34	39.20	0.00	-20.00	0.00	52.57
474	ADHEC, GLUDy, LDIH_D, PPC, PPC	12.60	0.16	11.81	0.00	0.47	38.84	0.00	-20.00	0.00	51.79
475	ADHEC, AIPS4c, FRD2, NADH6, SUTabc	12.60	0.48	14.92	0.00	19.87	0.00	0.00	20.00	0.00	18.32
476	ADHEC, LDIH_D, PPC, PPC, THD2	12.60	0.16	11.72	0.00	0.49	38.82	0.00	-20.00	0.00	51.71
477	ADHEC, ASPT, AIPS4c, GLCps, MDH	12.57	0.29	9.83	0.00	18.83	0.00	0.00	20.00	0.00	18.28
478	ADHEC, G6PDHy, MDH, NADH6, THD2	12.57	0.58	7.32	0.00	24.57	3.87	0.00	-20.00	0.00	15.05
479	ADHEC, MDH, NADH6, PGL, THD2	12.57	0.58	7.32	0.00	24.57	2.87	0.00	-20.00	0.00	15.05
480	ADHEC, ACK, FBA, GLUDy, NADH6	12.36	0.41	2.58	0.00	28.57	0.00	0.00	20.00	0.00	15.47
481	ADHEC, ACK, GLUDy, NADH6, PTK	12.36	0.41	2.58	0.00	28.57	0.00	0.00	-20.00	0.00	15.47
482	ADHEC, ACK, GLUDy, NADH6, TPI	12.36	0.41	2.58	0.00	28.57	0.00	0.00	20.00	0.00	18.94
483	ADHEC, AIPS4c, MDH, NADH6, PFK	12.33	0.49	15.46	0.00	19.46	0.00	0.00	-20.00	0.00	18.94
484	ADHEC, AIPS4c, FRD2, NADH6, PFK	12.33	0.49	15.46	0.00	19.46	0.00	0.00	-20.00	0.00	18.94

5
10
15
20
25
30
35
40
45
50
55

485	ADH1c, ATPS4c, G6PDHc, GLCP1c, MDH1	12.30	0.34	0.00	0.00	12.96	0.00	0.00	-20.00	0.00	24.29
486	ADH1c, ATPS4c, GLCP1c, MDH1, PGI	12.30	0.34	0.00	0.00	12.96	0.00	0.00	20.00	0.00	24.29
487	ADH1c, ACKc, FBA, GLCP1c, NADH6	12.15	0.44	12.39	0.00	28.20	0.00	0.00	20.00	0.00	15.54
488	ADH1c, ACKc, GLCP1c, NADH6, TPI	12.15	0.44	12.39	0.00	28.20	0.00	0.00	20.00	0.00	15.54
489	ADH1c, ACKc, GLCP1c, NADH6, PFK	12.15	0.44	12.39	0.00	28.20	0.00	0.00	20.00	0.00	15.54
490	ADH1c, ACKc, LDH1D, MDH1, SUC1D4	12.15	0.40	12.37	0.00	6.33	23.73	0.00	20.00	0.00	38.94
491	ADH1c, ACKc, AKG1D, ATPS4c, FBA	12.15	0.22	12.15	0.00	47.66	0.00	0.00	20.00	0.00	13.85
492	ADH1c, ACKc, AKG1D, ATPS4c, PFK	12.15	0.22	12.15	0.00	47.66	0.00	0.00	20.00	0.00	13.85
493	ADH1c, ACKc, AKG1D, ATPS4c, TPI	12.15	0.22	12.15	0.00	47.66	0.00	0.00	20.00	0.00	13.85
494	ADH1c, FBA, PGI, PCK, SUC1D4	12.09	0.37	10.13	0.00	7.88	20.50	0.00	20.00	0.00	41.14
495	ADH1c, ACKc, ATPS4c, FBA, SUC1D4	12.09	0.23	12.21	0.00	12.27	25.58	0.00	20.00	0.00	39.40
496	ADH1c, ACKc, ATPS4c, PFK, SUC1D4	12.09	0.23	12.21	0.00	12.27	25.58	0.00	20.00	0.00	39.40
497	ADH1c, ACKc, ATPS4c, SUC1D4, TPI	12.09	0.23	12.21	0.00	57.85	0.00	0.00	20.00	0.00	13.82
498	ADH1c, FRD2, LDH1D, ME2, SUC1D4	12.08	0.26	3.82	0.00	8.33	11.37	4.02	20.00	0.00	33.19
499	ADH1c, ACKc, CBMK2, FBA, NADH6	12.06	0.46	12.31	0.00	28.05	0.00	0.00	20.00	0.00	15.57
500	ADH1c, ACKc, CBMK2, NADH6, PFK	12.06	0.46	12.31	0.00	28.05	0.00	0.00	20.00	0.00	15.57
501	ADH1c, ACKc, CBMK2, NADH6, TPI	12.06	0.46	12.31	0.00	28.05	0.00	0.00	20.00	0.00	15.57
502	ADH1c, ACKc, NADH6, RPE, TPI	12.05	0.46	12.30	0.00	28.05	0.00	0.00	20.00	0.00	15.58
503	ADH1c, ACKc, NADH6, PFK, RPE	12.05	0.46	12.30	0.00	28.05	0.00	0.00	20.00	0.00	15.58
504	ADH1c, ACKc, FBA, NADH6, RPE	12.05	0.46	12.30	0.00	28.05	0.00	0.00	20.00	0.00	15.58
505	ADH1c, ACKc, ASSN1, FBA, NADH6	12.05	0.46	12.30	0.00	28.05	0.00	0.00	20.00	0.00	15.58

Table 7 (cont'd)

		H2O	ILE	LAC	NH4	NO3	PHE	PI	SO4	SUC	THR	VAL
	Metabolic Trans- formations Targeted For Removal val, #											
1	ADHEr, NADHe6	8.96	0.00	0.00	6.26	3.00	0.00	0.78	-0.13	0.23	0.00	0.00
2	ADHEr, FNO	12.07	0.00	0.00	-0.24	-2.00	0.00	0.03	0.00	0.00	0.00	0.00
3	ADHEr, PGNr	-12.07	0.00	0.00	0.24	2.00	0.00	0.03	0.00	0.00	0.00	0.00
4	ADHEr, PPK	0.90	0.00	0.00	-4.96	0.00	0.00	-0.61	-0.10	0.00	0.00	0.00
5	ADHEr, SUCD4	0.68	0.00	0.00	-8.69	0.00	0.00	-0.69	0.11	0.00	0.00	0.00
6	ADHEr, ATPSHr	6.08	0.00	0.00	-7.09	-5.00	0.00	0.88	-0.14	0.00	0.00	0.00
7	ADHEr, PGI	21.51	0.00	0.00	-4.52	0.00	0.00	-0.86	-0.09	24.86	0.00	0.00
8	ADHEr, FUNr	10.90	0.00	0.00	-6.44	0.00	0.00	-0.80	0.13	12.16	0.00	0.00
9	ADHEr, HEAT	12.79	0.00	0.00	6.62	0.00	0.00	0.82	-0.13	13.54	0.00	0.00
10	ADHEr, MDH	10.81	0.00	0.00	-5.83	0.00	0.00	-0.72	0.12	14.32	0.00	0.00
11	ADHEr, TPI	8.25	0.00	0.00	4.23	0.00	0.00	0.51	0.09	16.46	0.00	0.00
12	ADHEr, PBA	8.25	0.00	0.00	4.23	0.00	0.00	0.51	-0.09	16.46	0.00	0.00
13	ADHEr, PPK	8.25	0.00	0.00	-4.23	0.00	0.00	-0.52	-0.09	16.46	0.00	0.00
14	ADHEr, HXN1, PGI	4.02	0.00	0.00	-3.36	0.00	0.00	0.42	0.07	0.00	0.00	0.00
15	ADHEr, PFL, PPK	14.07	0.00	0.00	4.31	0.00	0.00	0.53	0.09	0.00	0.00	0.00
16	ADHEr, PFL, SUCD4	14.87	0.00	0.00	-4.72	0.00	0.00	-0.58	0.10	0.00	0.00	0.00
17	ADHEr, ACKr, NADHe6	30.54	0.00	0.00	7.41	20.00	0.00	0.92	0.15	0.00	0.00	0.00
18	ADHEr, NADHe6, PFL	15.46	0.00	0.00	-5.03	0.00	0.00	-0.62	-0.10	0.00	0.00	0.00
19	ADHEr, NADHe6, PGM	3.71	0.00	0.00	-0.73	-10.00	0.00	-0.09	0.01	0.00	0.00	0.00
20	ADHEr, FNO, NADHe6	3.71	0.00	0.00	-0.73	-10.00	0.00	-0.09	0.01	0.00	0.00	0.00
21	ADHEr, ASP1, MDH	-1.88	0.00	0.00	-1.57	0.00	0.00	-0.57	-0.09	0.00	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

22	ADDER, NADDE6, PGL	0.60	0.00	0.00	-3.07	0.00	0.00	-0.38	0.06	0.00	0.00
23	ADDER, NADDE6, TPI	0.77	0.00	0.00	-3.11	0.00	0.00	0.38	0.06	0.00	0.00
24	ADDER, FBA, NADDE6	-0.77	0.00	0.00	-2.11	0.00	0.00	-0.38	-0.06	0.00	0.00
25	ADDER, NADDE6, PPK	0.77	0.00	0.00	3.11	0.00	0.00	-0.38	-0.06	0.00	0.00
26	ADDER, NADDE6, PPK	2.51	0.00	0.00	-4.48	0.00	0.00	0.55	0.09	0.00	0.00
27	ADDER, MDH, NADDE6	4.03	0.00	0.00	5.12	0.00	0.00	-0.63	-0.10	0.00	0.00
28	ADDER, GUT, NADDE6	5.55	0.00	0.00	5.78	0.00	0.00	0.72	0.12	0.00	0.00
29	ADDER, PPK, THD2	0.13	0.00	0.00	-4.53	0.00	0.00	0.60	-0.10	0.00	0.00
30	ADDER, NADDE6, RPE	13.24	0.00	0.00	6.66	-5.00	0.00	-0.82	-0.13	0.00	0.00
31	ADDER, NADDE6, TAC	12.89	0.00	0.00	6.69	5.00	0.00	-0.83	0.12	0.00	0.00
32	ADDER, PGL, PPK	-6.65	0.00	0.00	-2.62	0.00	0.00	0.37	0.05	0.00	0.00
33	ADDER, PGL, SUCD4	-6.10	0.00	0.00	-2.84	0.00	0.00	-0.35	-0.06	0.00	0.00
34	ADDER, ATP54r, PPK	1.88	0.00	0.00	-5.25	0.00	0.00	0.65	0.11	0.00	0.00
35	ADDER, PPK, PPK	-6.68	0.00	0.00	-2.66	0.00	0.00	0.33	-0.05	0.00	0.00
36	ADDER, FBA, PPK	-6.68	0.00	0.00	-2.66	0.00	0.00	-0.33	-0.05	0.00	0.00
37	ADDER, PPK, TPI	6.68	0.00	0.00	-2.66	0.00	0.00	-0.33	0.05	0.00	0.00
38	ADDER, FBA, HEX1	-6.39	0.00	0.00	-2.77	0.00	0.00	-0.34	-0.06	0.00	0.00
39	ADDER, HEX1, PPK	-6.39	0.00	0.00	-2.77	0.00	0.00	-0.34	-0.06	0.00	0.00
40	ADDER, HEX1, TPI	6.39	0.00	0.00	-2.77	0.00	0.00	0.34	0.06	0.00	0.00
41	ADDER, MDH, THD2	8.45	0.00	0.00	5.39	0.00	0.00	-0.67	-0.11	0.00	0.00
42	ADDER, SUCD4, TPI	-6.13	0.00	0.00	7.88	0.00	0.00	-0.36	0.06	0.00	0.00
43	ADDER, FBA, SUCD4	-6.13	0.00	0.00	-2.88	0.00	0.00	0.36	0.06	0.00	0.00
44	ADDER, PPK, SUCD4	-6.13	0.00	0.00	2.88	0.00	0.00	-0.36	-0.06	0.00	0.00
45	ADDER, GUT, PGL	19.69	0.00	0.00	6.02	0.00	0.00	-0.75	-0.17	0.00	0.00
46	ADDER, PPK, SUCD4	3.22	0.00	0.00	-3.04	0.00	0.00	0.50	-0.08	0.00	0.00
47	ADDER, PPK, RPE	-0.16	0.00	0.00	-4.28	0.00	0.00	-0.61	-0.10	0.00	0.00
48	ADDER, GUT, PPK	-2.34	0.00	0.00	4.39	0.00	0.00	-0.54	-0.09	0.00	0.00
49	ADDER, GUT, MDH	2.39	0.00	0.00	-0.31	0.00	0.00	0.55	0.09	0.00	0.00
50	ADDER, GUT, PPK	-2.13	0.00	0.00	-4.47	0.00	0.00	-0.55	-0.09	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

51	ADHEG, MDH, SUCTD4	1.81	0.00	0.00	-4.62	0.00	-0.57	-0.09	0.00	0.00	0.00
52	ADHEG, PCK, FAL	-0.83	0.00	0.00	-4.95	0.00	-0.61	-0.10	0.00	0.00	0.00
53	ADHEG, FAL, PCK	-1.66	0.00	0.00	-4.66	0.00	0.58	0.09	0.00	0.00	0.00
54	ADHEG, MDH, PCK	1.66	0.00	0.00	-4.66	0.00	0.58	-0.09	0.00	0.00	0.00
55	ADHEG, RPE, SUCTD4	0.74	0.00	0.00	-5.54	0.00	-0.69	-0.11	0.00	0.00	0.00
56	ADHEG, ME2, SUCTD4	1.01	0.00	0.00	4.92	0.00	0.61	0.10	0.00	0.00	0.00
57	ADHEG, FAL, GLUDY	0.80	0.00	0.00	5.00	0.00	0.57	0.10	0.00	0.00	0.00
58	ADHEG, GLUDY, SUCTD4	-0.78	0.00	0.00	-5.00	0.00	-0.52	-0.10	0.00	0.00	0.00
59	ADHEG, GLCPS, SUCTD4	-0.69	0.00	0.00	-5.05	0.00	-0.52	-0.10	0.00	0.00	0.00
60	ADHEG, SUCTD4, FAL	0.71	0.00	0.00	-5.56	0.00	0.69	-0.11	0.00	0.00	0.00
61	ADHEG, FAL, SUCTD4	0.50	0.00	0.00	-5.44	0.00	-0.67	-0.11	0.00	0.00	0.00
62	ADHEG, BEN1, SUCTD4	0.47	0.00	0.00	5.51	0.00	0.68	0.11	0.00	0.00	0.00
63	ADHEG, CBMK2, SUCTD4	0.56	0.00	0.00	-5.55	0.00	-0.69	-0.11	0.00	0.00	0.00
64	ADHEG, FAL, BEN1	1.94	0.00	0.00	-6.10	0.00	-0.75	0.12	0.00	0.00	0.00
65	ADHEG, BEN1, PFL1	20.54	0.00	0.00	6.25	0.00	0.77	0.13	0.44	0.00	0.00
66	ADHEG, ATPS4, PGI	0.23	0.00	0.00	-4.49	0.00	-0.56	-0.09	0.00	0.00	0.00
67	ADHEG, ATPS4, FBA	-0.28	0.00	0.00	4.56	0.00	0.56	0.09	0.00	0.00	0.00
68	ADHEG, ATPS4, PFK	-0.25	0.00	0.00	4.56	0.00	0.56	0.09	0.00	0.00	0.00
69	ADHEG, ATPS4, TPI	0.25	0.00	0.00	-4.56	0.00	-0.56	-0.09	0.00	0.00	0.00
70	ADHEG, PFL1, TPI	16.61	0.00	0.00	3.82	0.00	0.47	0.08	12.23	0.00	0.00
71	ADHEG, PFK, PFL1	16.61	0.00	0.00	-3.82	0.00	-0.47	0.08	12.23	0.00	0.00
72	ADHEG, FBA, PFL1	16.61	0.00	0.00	3.82	0.00	0.47	0.08	12.23	0.00	0.00
73	ADHEG, BEN1, FIDM	3.30	0.00	0.00	-6.40	0.00	-0.49	-0.13	1.81	0.00	0.00
74	ADHEG, ATPS4, RPE	6.06	0.00	0.00	-6.97	0.00	-0.86	0.14	0.00	0.00	0.00
75	ADHEG, ATPS4, GLUDY	4.14	0.00	0.00	-6.31	0.00	-0.75	-0.13	0.00	0.00	0.00
76	ADHEG, ATPS4, FAL	6.07	0.00	0.00	7.03	0.00	-0.87	-0.14	0.00	0.00	0.00
77	ADHEG, ATPS4, CBMK2	5.91	0.00	0.00	7.02	0.00	0.87	0.14	0.00	0.00	0.00
78	ADHEG, FIDM, PGI	23.66	0.00	0.00	-4.37	0.00	-0.54	-0.09	21.30	0.00	0.00
79	ADHEG, PFL1, PGI	22.80	0.00	0.00	-4.46	0.00	-0.55	-0.09	24.28	0.00	0.00

80	ADHER. MEDL. PPK	5.91	0.00	0.00	-2.11	0.00	0.00	-0.51	-0.08	13.43	0.00	0.00
81	ADHER. MEDL. PPI	5.91	0.00	0.00	-2.11	0.00	0.00	-0.51	-0.08	13.43	0.00	0.00
82	ADHER. FBA, NDIH	5.91	0.00	0.00	-1.11	0.00	0.00	-0.51	-0.08	13.43	0.00	0.00
83	ADHER. HENL, RPE	12.13	0.00	0.00	6.89	0.00	0.00	-0.52	-0.13	13.08	0.00	0.00
84	ADHER. MEDL. RPE	10.68	0.00	0.00	-5.80	0.00	0.00	-0.72	-0.13	13.90	0.00	0.00
85	ADHER. HENL, TAL	12.21	0.00	0.00	6.61	0.00	0.00	-0.82	-0.13	13.50	0.00	0.00
86	ADHER. MEDL, TAL	10.33	0.00	0.00	-5.81	0.00	0.00	-0.73	-0.12	14.10	0.00	0.00
87	ADHER. RPE, PPI	8.73	0.00	0.00	-1.19	0.00	0.00	-0.52	-0.08	16.35	0.00	0.00
88	ADHER. PPK, RPE	8.23	0.00	0.00	-2.19	0.00	0.00	-0.52	-0.08	16.35	0.00	0.00
89	ADHER. FBA, RPE	8.73	0.00	0.00	1.19	0.00	0.00	-0.52	-0.08	16.35	0.00	0.00
90	ADHER. PPK, TAL	8.24	0.00	0.00	-1.21	0.00	0.00	-0.52	-0.09	16.34	0.00	0.00
91	ADHER. FBA, TAL	8.74	0.00	0.00	1.21	0.00	0.00	-0.52	-0.09	16.34	0.00	0.00
92	ADHER. HENL, PPL, PGI	15.38	0.00	0.00	2.97	0.00	0.00	-0.37	-0.06	0.00	0.00	0.00
93	ADHER. HENL, NADH6, PGI	10.91	0.00	0.00	2.40	0.00	0.00	-0.30	-0.05	0.00	0.00	0.00
94	ADHER. FBA, NADH6, PGI	11.52	0.00	0.00	-2.66	0.00	0.00	-0.35	-0.05	0.00	0.00	0.00
95	ADHER. ACK, NADH6, PGI	13.73	0.00	0.00	2.97	0.00	0.00	-0.37	-0.06	0.00	0.00	0.00
96	ADHER. FRD2, LDH, D, M, D1	6.75	2.14	0.00	-2.35	0.00	0.00	-0.27	-0.04	0.00	0.00	0.00
97	ADHER. ATPS4c, PGI, SUC104	9.08	0.00	0.00	-2.36	0.00	0.00	-0.29	-0.05	0.00	0.00	0.00
98	ADHER. ATPS4c, LDH2, NADH6	16.09	0.00	0.00	-2.23	0.00	0.00	-0.52	-0.09	0.00	0.00	0.00
99	ADHER. ACK, NADH6, PPI	23.49	0.00	0.00	-2.03	0.00	0.00	-0.50	-0.08	0.00	0.00	0.00
100	ADHER. ACK, FBA, NADH6	24.49	0.00	0.00	-2.01	0.00	0.00	-0.50	-0.08	0.00	0.00	0.00
101	ADHER. ACK, NADH6, PPK	25.49	0.00	0.00	-3.01	0.00	0.00	-0.50	-0.08	0.00	0.00	0.00
102	ADHER. FRD2, LDH, D, M, D2	10.90	1.81	0.00	-3.06	0.00	0.00	-0.28	-0.05	0.00	0.00	0.00
103	ADHER. HENL, PGI, PPK	0.36	0.00	0.00	1.93	0.00	0.00	0.24	-0.03	0.00	0.00	0.00
104	ADHER. FBA, PGI, PPK	0.38	0.00	0.00	1.94	0.00	0.00	-0.24	-0.04	0.00	0.00	0.00
105	ADHER. HENL, PGI, SUC104	1.01	0.00	0.00	-2.19	0.00	0.00	-0.27	-0.04	0.00	0.00	0.00
106	ADHER. FBA, PGI, SUC104	1.02	0.00	0.00	2.21	0.00	0.00	-0.27	-0.04	0.00	0.00	0.00
107	ADHER. ATPS4c, FBA, PGI	2.80	0.00	0.00	2.91	0.00	0.00	-0.36	-0.06	0.00	0.00	0.00
108	ADHER. GLUDx, HENL, PGI	3.13	0.00	0.00	3.05	0.00	0.00	-0.38	-0.06	0.00	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

109	ADHEr, MDL, PGL, THD2	2.41	0.00	0.00	-4.15	0.00	0.00	0.00	-0.51	-0.08	0.00	0.00
110	ADHEr, G6PD45, MDH, THD2	2.41	0.00	0.00	-4.15	0.00	0.00	0.00	-0.51	-0.08	0.00	0.00
111	ADHEr, PFL, PGL, PPCK	9.09	0.00	0.00	1.75	0.00	0.00	0.00	0.21	0.03	0.00	0.00
112	ADHEr, PFL, PPCK, TPI	9.11	0.00	0.00	-1.75	0.00	0.00	0.00	-0.22	-0.04	0.00	0.00
113	ADHEr, FBA, PFL, PPCK	9.11	0.00	0.00	1.75	0.00	0.00	0.00	0.22	0.04	0.00	0.00
114	ADHEr, PK, PFL, PPCK	9.11	0.00	0.00	-1.75	0.00	0.00	0.00	-0.22	-0.04	0.00	0.00
115	ADHEr, ACK, MDH, NADH6	26.42	0.00	0.00	6.41	-15.00	0.00	0.00	-0.59	0.13	0.00	0.00
116	ADHEr, NADH6, PFL, PGL	9.59	0.00	0.00	-1.99	0.00	0.00	0.00	0.25	-0.04	0.00	0.00
117	ADHEr, PFL, PGL, SUCD4	9.59	0.00	0.00	-1.99	0.00	0.00	0.00	-0.25	-0.04	0.00	0.00
118	ADHEr, FBA, PFL, SUCD4	9.62	0.00	0.00	-2.01	0.00	0.00	0.00	0.25	-0.04	0.00	0.00
119	ADHEr, PK, PFL, SUCD4	9.62	0.00	0.00	-2.01	0.00	0.00	0.00	-0.25	-0.04	0.00	0.00
120	ADHEr, NADH6, PK, PFL	9.62	0.00	0.00	2.01	0.00	0.00	0.00	0.25	0.04	0.00	0.00
121	ADHEr, FBA, NADH6, PFL	9.62	0.00	0.00	2.01	0.00	0.00	0.00	0.25	0.04	0.00	0.00
122	ADHEr, PFL, SUCD4, TPI	9.62	0.00	0.00	-2.01	0.00	0.00	0.00	-0.25	-0.04	0.00	0.00
123	ADHEr, NADH6, PFL, TPI	9.62	0.00	0.00	2.01	0.00	0.00	0.00	0.25	0.04	0.00	0.00
124	ADHEr, HEN, PK, PFL	9.63	0.00	0.00	-2.03	0.00	0.00	0.00	-0.25	-0.04	0.00	0.00
125	ADHEr, FBA, HEN, PFL	9.63	0.00	0.00	2.03	0.00	0.00	0.00	0.25	0.04	0.00	0.00
126	ADHEr, HEN, PFL, TPI	9.63	0.00	0.00	-2.02	0.00	0.00	0.00	-0.25	-0.04	0.00	0.00
127	ADHEr, PFL, PPCK, THD2	14.16	0.00	0.00	-3.23	0.00	0.00	0.00	-0.52	-0.09	0.00	0.00
128	ADHEr, ACK, GLUD5, NADH6	28.93	0.00	0.00	6.51	-20.00	0.00	0.00	-0.81	0.13	0.00	0.00
129	ADHEr, ACK, GLCps, NADH6	29.25	0.00	0.00	6.68	-20.00	0.00	0.00	-0.83	0.14	0.00	0.00
130	ADHEr, ACK, AKG2, ATPS4	12.00	0.00	0.00	-4.99	-15.00	0.00	0.00	-0.62	-0.10	0.00	0.00
131	ADHEr, ATPS4, NADH6, PFL	14.83	0.00	0.00	-4.62	0.00	0.00	0.00	-0.57	-0.09	0.00	0.00
132	ADHEr, GLCps, PFL, PPCK	12.93	0.00	0.00	3.72	0.00	0.00	0.00	0.46	-0.08	0.00	0.00
133	ADHEr, ACK, ATPS4, SUCOAS	11.13	0.00	0.00	-5.03	-14.78	0.00	0.00	-0.62	0.10	0.00	0.00
134	ADHEr, ACK, ME2, NADH6	27.22	0.00	0.00	6.69	-15.00	0.00	0.00	-0.83	0.14	0.00	0.00
135	ADHEr, GLUD5, PFL, PPCK	13.23	0.00	0.00	-3.87	0.00	0.00	0.00	-0.48	-0.08	0.00	0.00
136	ADHEr, ME2, PFL, SUCD4	13.54	0.00	0.00	4.03	0.00	0.00	0.00	-0.50	0.08	0.00	0.00
137	ADHEr, MDH, NADH6, PFL	13.59	0.00	0.00	-3.06	0.00	0.00	0.00	-0.50	-0.08	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

138	ADHEr, PLL, PPCK, RPE	14.08	0.00	0.00	-4.30	0.00	-0.53	-0.09	0.00	0.00	0.00
139	ADHEr, PLL, PPCK, TAL	14.07	0.00	0.00	-4.30	0.00	-0.53	-0.09	0.00	0.00	0.00
140	ADHEr, GLUTS, PELL, SUCD4	13.88	0.00	0.00	4.21	0.00	-0.52	-0.09	0.00	0.00	0.00
141	ADHEr, CBMK2, PLL, PPCK	13.99	0.00	0.00	4.37	0.00	0.53	0.09	0.00	0.00	0.00
142	ADHEr, ATPS-g, LIDH2, SUCD4	15.53	0.00	0.00	-4.91	2.00	-0.64	-0.10	0.00	0.00	0.00
143	ADHEr, PLL, RPE, SUCD4	14.82	0.00	0.00	4.67	0.00	0.58	0.09	0.00	0.00	0.00
144	ADHEr, ACKr, CBMK2, NADH6	30.40	0.00	0.00	-7.33	20.00	0.00	0.91	0.15	0.00	0.00
145	ADHEr, PLL, SUCD4, TAL	14.84	0.00	0.00	-4.70	0.00	-0.58	-0.09	0.00	0.00	0.00
146	ADHEr, ACKr, NADH6, RPE	30.41	0.00	0.00	7.34	20.00	0.00	0.91	0.15	0.00	0.00
147	ADHEr, ACKr, LUM, NADH6	30.45	0.00	0.00	-1.36	20.00	0.00	-0.91	-0.15	0.00	0.00
148	ADHEr, ACKr, NADH6, TAL	30.47	0.00	0.00	7.37	20.00	0.00	-0.91	-0.15	0.00	0.00
149	ADHEr, ACKr, ANS2, NADH6	30.48	0.00	0.00	7.38	20.00	0.00	0.91	0.15	0.00	0.00
150	ADHEr, CBMK2, PLL, SUCD4	14.78	0.00	0.00	-4.68	0.00	0.00	-0.58	-0.09	0.00	0.00
151	ADHEr, ACKr, NADH2, NADH6	30.49	0.00	0.00	-7.38	20.00	0.00	-0.91	-0.15	0.00	0.00
152	ADHEr, ACKr, NADH6, SOM2	30.51	0.00	0.00	7.39	20.00	0.00	0.91	0.15	0.00	0.00
153	ADHEr, NADH2, NADH6, PLL	14.87	0.00	0.00	-4.72	0.00	0.00	-0.58	-0.10	0.00	0.00
154	ADHEr, EUN, NADH6, PLL	14.89	0.00	0.00	4.73	0.00	0.00	-0.59	-0.10	0.00	0.00
155	ADHEr, ACKr, PGL, SUCD4	12.62	0.00	0.00	-6.04	0.00	0.00	0.23	0.04	1.94	4.75
156	ADHEr, NADH6, PLL, TAL	15.43	0.00	0.00	-5.00	0.00	0.00	-0.62	-0.10	0.00	0.00
157	ADHEr, CBMK2, NADH6, PPL	15.57	0.00	0.00	4.98	0.00	0.00	0.62	0.10	0.00	0.00
158	ADHEr, EUN, HEN1, PPL	15.52	0.00	0.00	5.47	0.00	0.00	0.68	0.11	0.00	0.00
159	ADHEr, MDL, NADH6, THD2	8.10	0.00	0.00	-4.75	0.00	0.00	-0.59	-0.10	0.00	0.00
160	ADHEr, ATPS-g, MDL, NADH6	8.00	0.00	0.00	4.76	0.00	0.00	0.59	0.10	0.00	0.00
161	ADHEr, ATPS-g, EUM, NADH6	14.08	0.00	0.00	-5.43	0.00	0.00	-0.67	-0.11	0.00	0.00
162	ADHEr, ATPS-g, PGL, PPCK	3.19	0.00	0.00	2.28	0.00	0.00	-0.28	0.05	0.00	0.00
163	ADHEr, ASPT, MDL, NADH6	0.03	0.00	0.00	-4.14	0.00	0.00	0.51	0.08	0.00	0.00
164	ADHEr, ATPS-g, NADH6, PPCK	6.46	0.00	0.00	-4.15	0.00	0.00	-0.51	-0.08	0.00	0.00
165	ADHEr, ASPT, MDL, THD2	-1.24	0.00	0.00	4.34	0.00	0.00	-0.54	-0.09	0.00	0.00
166	ADHEr, ATPS-g, GLCPs, SUCD4	17.57	0.00	0.00	5.67	5.00	0.00	0.70	0.11	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

167	ADHET, ASP1, MDH, PGI	-8.18	0.00	0.00	-2.02	0.00	0.00	-0.25	-0.04	-0.03	0.00
168	ADHET, ASP1, FBA, MDH	-8.17	0.00	0.00	-2.06	0.00	0.00	-0.25	-0.04	0.00	0.00
169	ADHET, ASP1, MDH, PER	8.17	0.00	0.00	2.06	0.00	0.00	0.25	-0.04	0.00	0.00
170	ADHET, ASP1, MDH, TPI	-8.17	0.00	0.00	-2.06	0.00	0.00	-0.25	-0.04	0.00	0.00
171	ADHET, ATPS4E, P, K, PPK	0.48	0.00	0.00	2.25	0.00	0.00	0.28	0.05	0.00	0.00
172	ADHET, ATPS4E, FBA, PPK	0.48	0.00	0.00	-2.25	0.00	0.00	-0.28	-0.05	0.00	0.00
173	ADHET, ATPS4E, PPK, TPI	0.48	0.00	0.00	2.25	0.00	0.00	0.28	-0.05	0.00	0.00
174	ADHET, ACK, LIDA, PGI	11.87	0.00	0.00	3.90	0.00	0.00	0.48	0.08	0.88	0.00
175	ADHET, ATPS4E, HEN1, NADH6	11.00	0.00	0.00	-5.78	0.00	0.00	-0.72	-0.12	0.00	0.00
176	ADHET, NADH6, PPK, THD2	3.90	0.00	0.00	-4.30	0.00	0.00	-0.53	-0.09	0.00	0.00
177	ADHET, ATPS4E, GLUC, MDH	0.51	0.00	0.00	3.99	0.00	0.00	0.49	0.08	0.00	0.00
178	ADHET, ATPS4E, MDH, PPK	1.11	0.00	0.00	4.22	0.00	0.00	-0.52	-0.09	0.00	0.00
179	ADHET, ATPS4E, P, K, PPK	1.11	0.00	0.00	4.22	0.00	0.00	0.52	0.09	0.00	0.00
180	ADHET, FNO, NADH6, RPE	3.75	0.00	0.00	0.75	0.00	0.00	0.09	0.01	0.00	0.00
181	ADHET, NADH6, PGM, RPE	3.75	0.00	0.00	-0.75	0.00	0.00	-0.09	-0.01	0.00	0.00
182	ADHET, NADH6, PGM, FAL	4.75	0.00	0.00	-0.75	-0.00	0.00	-0.09	-0.01	0.00	0.00
183	ADHET, FNO, NADH6, TPI	3.75	0.00	0.00	0.75	0.00	0.00	0.09	0.01	0.00	0.00
184	ADHET, ASP1, GLUC, MDH	3.56	0.00	0.00	-3.94	0.00	0.00	-0.49	-0.08	0.00	0.00
185	ADHET, ASP1, MDH, RPE	-1.29	0.00	0.00	4.52	0.00	0.00	-0.56	0.09	0.00	0.00
186	ADHET, ASP1, GLUTD, MDH	3.71	0.00	0.00	-4.08	0.00	0.00	-0.51	0.08	0.00	0.00
187	ADHET, ME2, NADH6, THD2	27.21	0.00	0.00	-7.53	-15.00	0.00	-0.93	-0.18	0.00	0.00
188	ADHET, ME2, SUCD4, THD2	0.98	0.00	0.00	-4.53	0.00	0.00	0.56	0.09	0.00	0.00
189	ADHET, ASP1, MDH, FAL	1.83	0.00	0.00	-4.56	0.00	0.00	-0.56	-0.09	0.00	0.00
190	ADHET, NADH6, PGI, PPK	3.12	0.00	0.00	2.02	0.00	0.00	-0.25	-0.04	0.00	0.00
191	ADHET, FEN, PPK, THD2	0.18	0.00	0.00	4.28	0.00	0.00	0.53	0.09	0.00	0.00
192	ADHET, MDH, PPK, THD2	0.18	0.00	0.00	-4.28	0.00	0.00	-0.53	-0.09	0.00	0.00
193	ADHET, GLUTD, MDH, THD2	0.33	0.00	0.00	-4.05	0.00	0.00	-0.50	-0.08	0.00	0.00
194	ADHET, ASP1, CBMK2, MDH	-7.99	0.00	0.00	4.53	0.00	0.00	0.56	0.09	0.00	0.00
195	ADHET, ATPS4E, FBA, SUCD4	7.01	0.00	0.00	-3.62	-5.00	0.00	-0.55	-0.07	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

196	ADHEr, ATPS4r, PFK, SUCD4	7.01	0.00	0.00	-3.62	-5.09	0.00	-0.45	-0.07	0.00	0.00	0.00
197	ADHEr, ATPS4r, SUCD4, TP1	7.01	0.00	0.00	-3.62	-5.09	0.00	-0.45	-0.07	0.00	0.00	0.00
198	ADHEr, FBA, NADH6, PFK	-3.17	0.00	0.00	-2.06	0.00	0.00	-0.25	-0.04	0.00	0.00	0.00
199	ADHEr, NADH6, PFK, PFK	-3.17	0.00	0.00	-2.06	0.00	0.00	-0.25	-0.04	0.00	0.00	0.00
200	ADHEr, NADH6, PFK, TP1	-3.17	0.00	0.00	-2.06	0.00	0.00	-0.25	-0.04	0.00	0.00	0.00
201	ADHEr, HEN1, PFK, THD2	19.49	0.00	0.00	6.04	0.00	0.00	0.74	0.17	6.13	0.00	0.00
202	ADHEr, HEN1, NADH6, PFK	-2.76	0.00	0.00	2.23	0.00	0.00	0.28	0.05	0.00	0.00	0.00
203	ADHEr, FBA, HEN1, NADH6	-2.76	0.00	0.00	-2.23	0.00	0.00	-0.28	-0.05	0.00	0.00	0.00
204	ADHEr, HEN1, NADH6, TP1	-2.76	0.00	0.00	-2.23	0.00	0.00	-0.28	-0.05	0.00	0.00	0.00
205	ADHEr, ATPS4r, G6PDH6, MDH	18.41	0.00	0.00	3.09	0.00	0.00	0.67	-0.20	8.31	0.00	0.00
206	ADHEr, ATPS4r, MDH, PFK	18.41	0.00	0.00	-4.09	0.00	0.00	-0.62	-0.20	8.31	0.00	0.00
207	ADHEr, GLUTD, NADH6, PFK	-1.39	0.00	0.00	2.74	0.00	0.00	-0.34	0.06	0.00	0.00	0.00
208	ADHEr, ACKr, FRD2, THD2	26.90	0.00	0.00	-9.46	-20.00	0.00	-1.04	-0.17	0.00	0.00	0.00
209	ADHEr, ACKr, LDH D, SUCD4	26.90	0.00	0.00	-10.53	-20.00	0.00	-1.04	-0.17	0.00	0.00	0.00
210	ADHEr, ATPS4r, FUM, GLUTD	5.95	0.00	0.00	-4.66	0.00	0.00	-0.58	0.09	0.00	0.00	0.00
211	ADHEr, NADH6, PFK, RPE	0.62	0.00	0.00	3.08	0.00	0.00	0.38	-0.06	0.00	0.00	0.00
212	ADHEr, FBA, NADH6, RPE	-0.62	0.00	0.00	-3.08	0.00	0.00	-0.38	-0.06	0.00	0.00	0.00
213	ADHEr, NADH6, RPE, TP1	-0.62	0.00	0.00	3.08	0.00	0.00	0.38	0.06	0.00	0.00	0.00
214	ADHEr, FBA, GLUTD, NADH6	1.50	0.00	0.00	2.77	0.00	0.00	-0.34	-0.05	0.00	0.00	0.00
215	ADHEr, GLUTD, NADH6, PFK	-1.50	0.00	0.00	-2.77	0.00	0.00	0.34	0.05	0.00	0.00	0.00
216	ADHEr, GLUTD, NADH6, TP1	-1.50	0.00	0.00	2.77	0.00	0.00	-0.34	-0.05	0.00	0.00	0.00
217	ADHEr, ATPS4r, FUM, HEN1	5.75	0.00	0.00	-5.74	0.00	0.00	-0.77	-0.12	0.00	0.00	0.00
218	ADHEr, NADH6, TP1, TP1	-0.66	0.00	0.00	-3.09	0.00	0.00	-0.38	0.06	0.00	0.00	0.00
219	ADHEr, NADH6, PFK, TP1	0.66	0.00	0.00	3.09	0.00	0.00	0.38	-0.06	0.00	0.00	0.00
220	ADHEr, FBA, NADH6, TP1	-0.66	0.00	0.00	-3.09	0.00	0.00	-0.38	0.06	0.00	0.00	0.00
221	ADHEr, ATPS4r, MDH, THD2	4.07	0.00	0.00	5.55	0.00	0.00	-0.66	-0.17	1.04	0.00	0.00
222	ADHEr, GLUTD, MDH, NADH6	-4.07	0.00	0.00	-3.57	0.00	0.00	-0.38	-0.08	0.00	0.00	0.00
223	ADHEr, GLUTD, NADH6, PFK	1.14	0.00	0.00	-3.90	0.00	0.00	-0.48	0.08	0.00	0.00	0.00
224	ADHEr, NADH6, PFK, RPE	2.70	0.00	0.00	4.46	0.00	0.00	0.55	-0.09	0.00	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

225	ADH1G, GLUD3, NADH6, PPK	1.46	0.00	0.00	4.03	0.00	0.00	0.50	0.08	0.00	0.00	0.00
226	ADH1G, FUM, NADH6, PPK	1.51	0.00	0.00	4.10	0.00	0.00	0.51	0.08	0.00	0.00	0.00
227	ADH1G, MDH, NADH6, PPK	1.61	0.00	0.00	4.10	0.00	0.00	-0.51	-0.08	0.00	0.00	0.00
228	ADH1G, ATPS4L, ERD2, THD2	27.21	0.00	0.00	13.79	2.00	0.00	0.55	0.06	0.00	0.00	10.48
229	ADH1G, NADH6, PPK, TAI	2.51	0.00	0.00	-4.47	0.00	0.00	-0.55	-0.09	0.00	0.00	0.00
230	ADH1G, FUM, GLUD3, NADH6	2.48	0.00	0.00	4.47	0.00	0.00	0.55	0.09	0.00	0.00	0.00
231	ADH1G, GLUD3, MDH, NADH6	2.59	0.00	0.00	4.52	0.00	0.00	-0.56	-0.09	0.00	0.00	0.00
232	ADH1G, MDH, NADH6, RPE	4.24	0.00	0.00	-5.10	0.00	0.00	-0.63	-0.10	0.00	0.00	0.00
233	ADH1G, MDH, NADH6, TAI	4.13	0.00	0.00	5.11	0.00	0.00	-0.63	0.10	0.00	0.00	0.00
234	ADH1G, IHEX1, NADH6, THD2	27.21	0.00	0.00	7.98	15.00	0.00	0.99	0.76	0.00	0.00	0.00
235	ADH1G, FUM, NADH6, RPE	5.73	0.00	0.00	-5.72	0.00	0.00	0.71	-0.12	0.00	0.00	0.00
236	ADH1G, CBMK2, MDH, NADH6	3.00	0.00	0.00	-5.08	0.00	0.00	-0.63	-0.10	0.00	0.00	0.00
237	ADH1G, FUM, MDH, NADH6	4.00	0.00	0.00	-5.12	0.00	0.00	-0.63	-0.10	0.00	0.00	0.00
238	ADH1G, FUM, NADH6, TAI	5.65	0.00	0.00	-5.75	0.00	0.00	0.71	-0.12	0.00	0.00	0.00
239	ADH1G, ATPS4L, MDH, PGD4	17.83	0.00	0.00	5.22	0.00	0.00	-0.65	0.11	7.01	0.00	0.00
240	ADH1G, FUM, IHEX1, NADH6	5.22	0.00	0.00	5.63	0.00	0.00	0.70	0.11	0.00	0.00	0.00
241	ADH1G, CBMK2, FUM, NADH6	5.13	0.00	0.00	-5.73	0.00	0.00	-0.71	-0.12	0.00	0.00	0.00
242	ADH1G, GLUD3, PPK, THD2	11.43	0.00	0.00	4.27	0.00	0.00	0.53	0.09	0.00	0.00	0.00
243	ADH1G, GLUD3, PPK, THD2	1.17	0.00	0.00	3.36	0.00	0.00	0.54	0.09	0.00	0.00	0.00
244	ADH1G, CBMK2, NADH6, TAI	12.72	0.00	0.00	6.63	-5.00	0.00	-0.82	0.13	0.00	0.00	0.00
245	ADH1G, ATPS4L, MDH, TAI	18.04	0.00	0.00	5.25	0.00	0.00	-0.65	0.11	7.45	0.00	0.00
246	ADH1G, ATPS4L, GLUD3, NADH6	8.28	0.00	0.00	5.60	2.00	0.00	0.69	-0.11	0.92	0.00	0.00
247	ADH1G, PGL, PPK, SUCD4	-8.37	0.00	0.00	-1.95	0.00	0.00	-0.24	-0.04	0.00	0.00	0.00
248	ADH1G, ATPS4L, MDH, RPE	18.24	0.00	0.00	5.28	0.00	0.00	-0.65	0.11	7.87	0.00	0.00
249	ADH1G, FBR, PGH, THD2	12.02	0.00	0.00	-0.23	-2.00	0.00	-0.03	0.00	0.00	0.00	0.00
250	ADH1G, ENO, FBR, THD2	-12.02	0.00	0.00	-0.23	-2.00	0.00	-0.03	0.00	0.00	0.00	0.00
251	ADH1G, FBA, PGH, THD2	-12.02	0.00	0.00	-0.23	-2.00	0.00	-0.03	0.00	0.00	0.00	0.00
252	ADH1G, ENO, FBA, THD2	-12.02	0.00	0.00	0.23	2.00	0.00	0.03	0.00	0.00	0.00	0.00
253	ADH1G, ENO, THD2, TPI	-12.02	0.00	0.00	-0.23	-2.00	0.00	-0.03	0.00	0.00	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

254	ADHET, PGM, HE02, TPI	7.07	0.00	0.00	0.25	2.00	0.00	0.03	0.00	0.00	0.00
255	ADHET, GL Cps, PGL, PPCK	7.58	0.00	0.00	2.25	0.00	0.00	0.28	0.05	0.00	0.00
256	ADHET, FBA, HEX1, PPCK	8.47	0.00	0.00	9.4	0.00	0.00	0.24	0.02	0.00	0.00
257	ADHET, HEX1, PFK, PPCK	8.47	0.00	0.00	9.4	0.00	0.00	0.24	0.02	0.00	0.00
258	ADHET, HEX1, PPCK, TPI	8.47	0.00	0.00	9.4	0.00	0.00	0.24	0.02	0.00	0.00
259	ADHET, PPCK, SUCD4, TPI	8.37	0.00	0.00	7.98	0.00	0.00	0.25	0.02	0.00	0.00
260	ADHET, PFK, PPCK, SUCD4	8.37	0.00	0.00	7.98	0.00	0.00	0.25	0.02	0.00	0.00
261	ADHET, FBA, PPCK, SUCD4	8.47	0.00	0.00	9.4	0.00	0.00	0.24	0.02	0.00	0.00
262	ADHET, GLUTD, PGL, PPCK	7.35	0.00	0.00	2.35	0.00	0.00	0.29	0.05	0.00	0.00
263	ADHET, GL Cps, PGL, SUCD4	7.06	0.00	0.00	2.46	0.00	0.00	0.30	0.05	0.00	0.00
264	ADHET, FUM, GLUTD, PGL	7.03	0.00	0.00	2.47	0.00	0.00	0.31	0.05	0.00	0.00
265	ADHET, GLUTD, M3H, PGL	7.03	0.00	0.00	2.47	0.00	0.00	0.31	0.05	0.00	0.00
266	ADHET, FBA, HEX1, SUCD4	8.02	0.00	0.00	2.12	0.00	0.00	0.26	0.02	0.00	0.00
267	ADHET, HEX1, SUCD4, TPI	8.02	0.00	0.00	2.12	0.00	0.00	0.26	0.02	0.00	0.00
268	ADHET, HEX1, PFK, SUCD4	8.02	0.00	0.00	2.12	0.00	0.00	0.26	0.02	0.00	0.00
269	ADHET, GLUTD, PGL, SUCD4	6.88	0.00	0.00	2.53	0.00	0.00	0.31	0.05	0.00	0.00
270	ADHET, FUM, HEX1, HE02	2.71	0.00	0.00	5.93	0.00	0.00	0.73	0.12	0.00	0.00
271	ADHET, FBA, GL Cps, PPCK	7.67	0.00	0.00	2.29	0.00	0.00	0.28	0.05	0.00	0.00
272	ADHET, GL Cps, PFK, PPCK	7.67	0.00	0.00	2.29	0.00	0.00	0.28	0.05	0.00	0.00
273	ADHET, GL Cps, PPCK, TPI	7.67	0.00	0.00	2.29	0.00	0.00	0.28	0.05	0.00	0.00
274	ADHET, PFK, PPCK, RPF	6.66	0.00	0.00	2.63	0.00	0.00	0.32	0.05	0.00	0.00
275	ADHET, PPCK, RPF, TPI	6.66	0.00	0.00	2.63	0.00	0.00	0.32	0.05	0.00	0.00
276	ADHET, FBA, PPCK, RPF	6.66	0.00	0.00	2.63	0.00	0.00	0.32	0.05	0.00	0.00
277	ADHET, GLUTD, PFK, PPCK	7.38	0.00	0.00	2.38	0.00	0.00	0.29	0.05	0.00	0.00
278	ADHET, GLUTD, PPCK, TPI	7.38	0.00	0.00	2.38	0.00	0.00	0.29	0.05	0.00	0.00
279	ADHET, FBA, GLUTD, PPCK	7.38	0.00	0.00	2.38	0.00	0.00	0.29	0.05	0.00	0.00
280	ADHET, ATPS4, HE02, HE02	4.44	0.00	0.00	6.13	2.00	0.00	0.75	0.12	0.00	0.00
281	ADHET, HEX1, PFK, RPF	6.36	0.00	0.00	2.74	0.00	0.00	0.34	0.06	0.00	0.00
282	ADHET, HEX1, RPF, TPI	6.36	0.00	0.00	2.74	0.00	0.00	0.34	0.06	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

283	ADHEr, FBA, HEX1, RPE	6.36	0.00	0.00	-2.73	0.00	0.00	0.34	0.00	0.00	0.00	0.00
284	ADHEr, FBA, PECK, IAI	6.67	0.00	0.00	2.64	0.00	0.00	0.34	0.00	0.00	0.00	0.00
285	ADHEr, PECK, IAI, TPI	-6.67	0.00	0.00	-2.64	0.00	0.00	-0.33	0.00	0.00	0.00	0.00
286	ADHEr, PEK, PECK, IAI	6.67	0.00	0.00	2.64	0.00	0.00	0.33	0.00	0.00	0.00	0.00
287	ADHEr, ADPSr, GLUTy, PECK	6.48	0.00	0.00	-3.75	-2.00	0.00	-0.59	0.00	0.00	0.00	0.00
288	ADHEr, FBA, GLUTy, HEX1	-7.10	0.00	0.00	2.49	0.00	0.00	0.31	0.00	0.00	0.00	0.00
289	ADHEr, GLUTy, HEX1, TPI	7.10	0.00	0.00	-2.49	0.00	0.00	-0.31	0.00	0.00	0.00	0.00
290	ADHEr, GLUTy, HEX1, PEK	-7.10	0.00	0.00	2.49	0.00	0.00	-0.31	0.00	0.00	0.00	0.00
291	ADHEr, GLCps, SUCTD4, TPI	-7.09	0.00	0.00	2.49	0.00	0.00	-0.31	0.00	0.00	0.00	0.00
292	ADHEr, FBA, GLCps, SUCTD4	7.09	0.00	0.00	-2.49	0.00	0.00	0.31	0.00	0.00	0.00	0.00
293	ADHEr, GLCps, PEK, SUCTD4	-7.09	0.00	0.00	2.49	0.00	0.00	-0.31	0.00	0.00	0.00	0.00
294	ADHEr, GLUTy, MDH, PEK	7.06	0.00	0.00	-2.50	0.00	0.00	0.31	0.00	0.00	0.00	0.00
295	ADHEr, FBA, GLUTy, MDH	-7.06	0.00	0.00	2.50	0.00	0.00	-0.31	0.00	0.00	0.00	0.00
296	ADHEr, FBA, FUM, GLUTy	7.06	0.00	0.00	-2.50	0.00	0.00	0.31	0.00	0.00	0.00	0.00
297	ADHEr, FUM, GLUTy, PEK	-7.06	0.00	0.00	2.50	0.00	0.00	-0.31	0.00	0.00	0.00	0.00
298	ADHEr, GLUTy, MDH, TPI	-7.06	0.00	0.00	2.50	0.00	0.00	0.31	0.00	0.00	0.00	0.00
299	ADHEr, FUM, GLUTy, TPI	7.06	0.00	0.00	-2.50	0.00	0.00	-0.31	0.00	0.00	0.00	0.00
300	ADHEr, RPE, SUCTD4, TPI	6.17	0.00	0.00	-2.83	0.00	0.00	-0.35	0.00	0.00	0.00	0.00
301	ADHEr, PEK, RPE, SUCTD4	-6.17	0.00	0.00	2.83	0.00	0.00	0.35	0.00	0.00	0.00	0.00
302	ADHEr, FBA, RPE, SUCTD4	6.17	0.00	0.00	-2.83	0.00	0.00	-0.35	0.00	0.00	0.00	0.00
303	ADHEr, GLUTy, PEK, SUCTD4	6.92	0.00	0.00	-2.56	0.00	0.00	-0.32	0.00	0.00	0.00	0.00
304	ADHEr, FBA, GLUTy, SUCTD4	-6.92	0.00	0.00	2.56	0.00	0.00	0.32	0.00	0.00	0.00	0.00
305	ADHEr, GLUTy, SUCTD4, TPI	-6.92	0.00	0.00	2.56	0.00	0.00	0.32	0.00	0.00	0.00	0.00
306	ADHEr, HEX1, PEK, IAI	6.37	0.00	0.00	-2.76	0.00	0.00	-0.34	0.00	0.00	0.00	0.00
307	ADHEr, FBA, HEX1, IAI	-6.37	0.00	0.00	2.76	0.00	0.00	0.34	0.00	0.00	0.00	0.00
308	ADHEr, HEX1, IAI, TPI	6.37	0.00	0.00	-2.76	0.00	0.00	-0.34	0.00	0.00	0.00	0.00
309	ADHEr, PEK, SUCTD4, IAI	6.12	0.00	0.00	-2.86	0.00	0.00	-0.35	0.00	0.00	0.00	0.00
310	ADHEr, FBA, SUCTD4, IAI	-6.12	0.00	0.00	2.86	0.00	0.00	0.35	0.00	0.00	0.00	0.00
311	ADHEr, SUCTD4, IAI, TPI	-6.12	0.00	0.00	2.86	0.00	0.00	0.35	0.00	0.00	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

312	ADHEP, FUM, ME2, TMD4	8.35	0.00	-5.39	0.00	-0.67	-0.11	6.20	0.00	0.00
313	ADHEP, ADP5R, ME2, SUCD4	6.35	0.00	-5.62	0.00	-0.69	-0.11	0.00	0.00	0.00
314	ADHEP, PPCK, PYK, SUCD4	-5.01	0.00	-5.32	0.00	-0.41	-0.07	0.00	0.00	0.00
315	ADHEP, GLCPs, PPCK, SUCD4	4.70	0.00	5.35	0.00	0.43	0.07	0.00	0.00	0.00
316	ADHEP, PPCK, RPE, SUCD4	-3.14	0.00	-4.01	0.00	-0.50	-0.08	0.00	0.00	0.00
317	ADHEP, FUM, GLUDy, PPCK	-4.29	0.00	5.67	0.00	0.45	0.07	0.00	0.00	0.00
318	ADHEP, GLUDy, MDH, PPCK	4.29	0.00	5.67	0.00	0.45	0.07	0.00	0.00	0.00
319	ADHEP, GLUDy, PPCK, SUCD4	-4.24	0.00	-5.63	0.00	-0.45	-0.07	0.00	0.00	0.00
320	ADHEP, PPCK, SUCD4, TAI	3.15	0.00	-4.02	0.00	-0.50	-0.08	0.00	0.00	0.00
321	ADHEP, GLCPs, GLUDy, MDH	3.68	0.00	5.85	0.00	0.48	0.08	0.00	0.00	0.00
322	ADHEP, GLCPs, PPCK, RPE	-2.32	0.00	-4.37	0.00	-0.54	-0.09	0.00	0.00	0.00
323	ADHEP, GLUDy, MDH, RPE	2.19	0.00	4.39	0.00	0.54	0.09	0.00	0.00	0.00
324	ADHEP, GLUDy, PPCK, RPE	-2.00	0.00	-4.46	0.00	-0.55	-0.09	0.00	0.00	0.00
325	ADHEP, GLCPs, GLUDy, PPCK	3.42	0.00	5.96	0.00	0.49	0.08	0.00	0.00	0.00
326	ADHEP, GLCPs, MDH, SUCD4	3.35	0.00	5.99	0.00	0.49	0.08	0.00	0.00	0.00
327	ADHEP, MDH, RPE, SUCD4	1.71	0.00	-3.57	0.00	-0.57	-0.09	0.00	0.00	0.00
328	ADHEP, GLCPs, MDH, PPCK	-3.12	0.00	-4.08	0.00	-0.50	-0.08	0.00	0.00	0.00
329	ADHEP, FUM, GLCPs, PPCK	3.17	0.00	4.08	0.00	0.50	0.08	0.00	0.00	0.00
330	ADHEP, MDH, PPCK, RPE	-1.54	0.00	-4.64	0.00	-0.57	-0.09	0.00	0.00	0.00
331	ADHEP, FUM, PPCK, RPE	1.54	0.00	4.64	0.00	0.57	0.09	0.00	0.00	0.00
332	ADHEP, GLCPs, PPCK, TAI	2.28	0.00	3.38	0.00	0.53	0.09	0.00	0.00	0.00
333	ADHEP, GLUDy, MDH, TAI	-2.34	0.00	-4.40	0.00	-0.54	-0.09	0.00	0.00	0.00
334	ADHEP, MDH, PPCK, PYK	-3.17	0.00	-4.03	0.00	-0.50	-0.08	0.00	0.00	0.00
335	ADHEP, FUM, PPCK, PYK	3.17	0.00	4.03	0.00	0.50	0.08	0.00	0.00	0.00
336	ADHEP, GLUDy, PPCK, TAI	-2.06	0.00	-4.47	0.00	-0.55	-0.09	0.00	0.00	0.00
337	ADHEP, FUM, GLUDy, SUCD4	-2.62	0.00	-4.78	0.00	-0.53	-0.09	0.00	0.00	0.00
338	ADHEP, MDH, SUCD4, TAI	1.76	0.00	4.89	0.00	0.57	0.09	0.00	0.00	0.00
339	ADHEP, GLCPs, ME2, SUCD4	-2.52	0.00	-4.32	0.00	-0.53	-0.09	0.00	0.00	0.00
340	ADHEP, ME2, RPE, SUCD4	0.89	0.00	4.90	0.00	0.61	0.10	0.00	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

341	ADHET, FUM, PPCK, TAL	-1.60	0.00	0.00	-4.65	0.00	0.00	-0.58	-0.09	0.00	0.00
342	ADHET, MDL, PPCK, TAL	-1.60	0.00	0.00	-4.65	0.00	0.00	-0.58	-0.09	0.00	0.00
343	ADHET, GLUDY, PROIZ, SUCD4	2.37	0.00	0.00	4.38	0.00	0.00	-0.54	-0.09	0.00	0.00
344	ADHET, FUM, GLUDY, RPE	-0.34	0.00	0.00	4.95	0.00	0.00	0.61	0.10	0.00	0.00
345	ADHET, GLUDY, RPE, SUCD5	-0.72	0.00	0.00	4.96	0.00	0.00	-0.61	-0.10	0.00	0.00
346	ADHET, GLCps, RPE, SUCD5	-0.69	0.00	0.00	4.98	0.00	0.00	0.62	0.10	0.00	0.00
347	ADHET, FUM, GLUDY, ME2	-2.29	0.00	0.00	-4.41	0.00	0.00	-0.55	-0.09	0.00	0.00
348	ADHET, GLUDY, ME2, SUCD4	2.28	0.00	0.00	4.41	0.00	0.00	-0.55	-0.09	0.00	0.00
349	ADHET, FUM, GLCps, GLUDY	2.16	0.00	0.00	4.46	0.00	0.00	0.55	0.09	0.00	0.00
350	ADHET, GLCps, GLUDY, SUCD4	-2.08	0.00	0.00	-4.49	0.00	0.00	-0.56	-0.09	0.00	0.00
351	ADHET, FUM, ME2, SUCD4	1.81	0.00	0.00	-4.60	0.00	0.00	-0.57	-0.09	0.00	0.00
352	ADHET, ME2, SUCD4, TAL	0.95	0.00	0.00	4.91	0.00	0.00	0.61	0.10	0.00	0.00
353	ADHET, FUM, GLUDY, TAL	-0.77	0.00	0.00	4.98	0.00	0.00	-0.62	-0.10	0.00	0.00
354	ADHET, GLUDY, SUCD4, TAL	-0.75	0.00	0.00	4.99	0.00	0.00	-0.62	-0.10	0.00	0.00
355	ADHET, GLCps, SUCD4, TAL	0.69	0.00	0.00	5.01	0.00	0.00	0.62	0.10	0.00	0.00
356	ADHET, ME2, PGL, HED2	26.98	0.00	0.00	8.25	-20.00	0.00	-1.02	-0.17	8.93	0.00
357	ADHET, G6PDH, ME2, HED2	26.98	0.00	0.00	8.25	30.00	0.00	-1.02	-0.17	8.93	0.00
358	ADHET, FUM, RPE, SUCD4	0.34	0.00	0.00	-5.38	0.00	0.00	0.67	0.11	0.00	0.00
359	ADHET, HEXT, RPE, SUCD4	0.56	0.00	0.00	-5.47	0.00	0.00	-0.68	-0.11	0.00	0.00
360	ADHET, CBMK2, GLUSK, PPCK	1.05	0.00	0.00	4.90	0.00	0.00	0.61	0.10	0.00	0.00
361	ADHET, CBMK2, G5SD, PPCK	1.05	0.00	0.00	-4.90	0.00	0.00	0.61	0.10	0.00	0.00
362	ADHET, ASNS2, CBMK2, PPCK	-1.05	0.00	0.00	-4.90	0.00	0.00	-0.61	-0.10	0.00	0.00
363	ADHET, CBMK2, PPCK, SUC42	1.03	0.00	0.00	4.91	0.00	0.00	0.61	0.10	0.00	0.00
364	ADHET, GLUDY, HEXT, SUCD4	-0.91	0.00	0.00	-4.96	0.00	0.00	0.61	0.10	0.00	0.00
365	ADHET, FUM, SUCD5, TAL	0.32	0.00	0.00	5.41	0.00	0.00	-0.67	-0.11	0.00	0.00
366	ADHET, HEXT, SUCD4, TAL	0.52	0.00	0.00	5.49	0.00	0.00	0.68	0.11	0.00	0.00
367	ADHET, FUM, HEXT, SUCD4	-0.27	0.00	0.00	-5.21	0.00	0.00	-0.65	-0.11	0.00	0.00
368	ADHET, FUM, HEXT, RPE	2.09	0.00	0.00	6.07	0.00	0.00	-0.75	-0.12	0.00	0.00
369	ADHET, CBMK2, FUM, SUCD4	0.18	0.00	0.00	5.39	0.00	0.00	0.67	0.11	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

370	ADHEr, HEX1, PET, RPE	0.00	0.00	-6.22	0.00	0.00	-0.37	-0.13	8.99	0.00	0.00
371	ADHEr, FUM, HEX1, TAL	0.00	0.00	-6.08	0.00	0.00	-0.35	-0.12	0.00	0.00	0.00
372	ADHEr, CBNK2, FUM, HEX1	0.00	0.00	6.05	0.00	0.00	0.75	0.12	0.00	0.00	0.00
373	ADHEr, HEX1, PET, TAL	0.00	0.00	-6.24	0.00	0.00	-0.37	-0.13	9.20	0.00	0.00
374	ADHEr, GLYCL, HEX1, PET	0.00	0.00	6.33	0.00	0.00	0.77	0.13	9.39	0.00	0.00
375	ADHEr, ATPS4r, GLUDy, PGL	0.00	0.00	-4.00	-5.00	0.00	-0.49	-0.08	0.00	0.00	0.00
376	ADHEr, PET, PGL, PGL, PGL	0.00	0.00	3.77	0.00	0.00	0.46	0.08	1.82	0.00	0.00
377	ADHEr, PET, PGL, TAL	0.00	0.00	-3.75	0.00	0.00	-0.46	-0.08	1.96	0.00	0.00
378	ADHEr, ATPS4r, PEK, RPE	0.00	0.00	4.50	5.00	0.00	0.86	0.09	0.00	0.00	0.00
379	ADHEr, ATPS4r, FBA, RPE	0.00	0.00	-4.50	-5.00	0.00	-0.86	-0.09	0.00	0.00	0.00
380	ADHEr, ATPS4r, RPE, PPI	0.00	0.00	4.50	5.00	0.00	0.86	0.09	0.00	0.00	0.00
381	ADHEr, PET, PGL, RPE	0.00	0.00	-3.75	0.00	0.00	-0.47	-0.08	12.07	0.00	0.00
382	ADHEr, ATPS4r, GLUDy, PPI	0.00	0.00	-4.05	-5.00	0.00	0.50	-0.08	0.00	0.00	0.00
383	ADHEr, ATPS4r, FBA, GLUDy	0.00	0.00	-4.05	-5.00	0.00	-0.50	-0.08	0.00	0.00	0.00
384	ADHEr, ATPS4r, GLUDy, PEK	0.00	0.00	4.05	5.00	0.00	0.50	0.08	0.00	0.00	0.00
385	ADHEr, FBA, PET, PGL	0.00	0.00	-3.77	0.00	0.00	0.47	-0.08	12.13	0.00	0.00
386	ADHEr, PEK, PET, PGL	0.00	0.00	-3.77	0.00	0.00	-0.47	-0.08	12.13	0.00	0.00
387	ADHEr, PET, PGL, PPI	0.00	0.00	3.77	0.00	0.00	0.47	0.08	12.13	0.00	0.00
388	ADHEr, PET, RPE, PPI	0.00	0.00	-3.78	0.00	0.00	-0.47	-0.08	12.15	0.00	0.00
389	ADHEr, PEK, PET, RPE	0.00	0.00	3.78	0.00	0.00	0.47	0.08	12.15	0.00	0.00
390	ADHEr, FBA, PET, RPE	0.00	0.00	-3.78	0.00	0.00	-0.47	-0.08	12.15	0.00	0.00
391	ADHEr, ATPS4r, PEK, TAL	0.00	0.00	4.53	5.00	0.00	0.56	0.09	0.00	0.00	0.00
392	ADHEr, ATPS4r, FBA, TAL	0.00	0.00	-4.53	-5.00	0.00	-0.56	-0.09	0.00	0.00	0.00
393	ADHEr, ATPS4r, TAL, PPI	0.00	0.00	4.53	5.00	0.00	0.56	0.09	0.00	0.00	0.00
394	ADHEr, FBA, PET, TAL	0.00	0.00	-3.86	0.00	0.00	0.47	0.08	12.18	0.00	0.00
395	ADHEr, PET, TAL, PPI	0.00	0.00	-3.86	0.00	0.00	-0.47	-0.08	12.18	0.00	0.00
396	ADHEr, PEK, PET, TAL	0.00	0.00	-3.86	0.00	0.00	-0.47	-0.08	12.18	0.00	0.00
397	ADHEr, GLYCL, HEX1, TAL	0.00	0.00	6.47	0.00	0.00	0.79	-0.13	1.43	0.00	0.00
398	ADHEr, GLUDy, PEK, PPI	0.00	0.00	-3.42	0.00	0.00	-0.42	-0.07	12.52	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

399	ADHEr, GLUDy, PEEL, TPI	15.92	0.00	0.00	3.42	0.00	0.00	-0.42	-0.07	12.52	0.00	0.00
400	ADHEr, FRA, GLUDy, PEEL	15.92	0.00	0.00	3.42	0.00	0.00	0.42	-0.07	12.52	0.00	0.00
401	ADHEr, EDA, PEEL, PGL	21.92	0.00	0.00	-4.32	0.00	0.00	-0.53	-0.09	30.78	0.00	0.00
402	ADHEr, ATPS2r, GLUDy, RPE	4.15	0.00	0.00	6.22	0.00	0.00	0.77	-0.13	0.00	0.00	0.00
403	ADHEr, ATPS4r, GLUDy, TAI	4.15	0.00	0.00	-6.26	0.00	0.00	-0.77	-0.13	0.00	0.00	0.00
404	ADHEr, ATPS4r, CBNK2, RPE	5.89	0.00	0.00	6.91	0.00	0.00	-0.85	0.14	0.00	0.00	0.00
405	ADHEr, ATPS4r, CBNK2, GLUDy	4.00	0.00	0.00	6.26	0.00	0.00	0.77	-0.13	0.00	0.00	0.00
406	ADHEr, ATPS4r, CBNK2, TAI	5.90	0.00	0.00	-6.96	0.00	0.00	-0.85	-0.14	0.00	0.00	0.00
407	ADHEr, ASNS2r, ATPS4r, GJL SK	5.95	0.00	0.00	-7.03	0.00	0.00	0.87	0.14	0.00	0.00	0.00
408	ADHEr, ASNS2r, ATPS4r, G5SD	5.95	0.00	0.00	7.03	0.00	0.00	-0.87	0.14	0.00	0.00	0.00
409	ADHEr, ACK2, PEEL, PGL	21.51	0.00	0.00	-4.32	0.00	0.00	0.53	-0.09	17.79	0.00	0.00
410	ADHEr, ACK2, AKGLD, FRD2	30.75	0.00	15.19	-8.96	0.00	0.00	1.19	-0.18	0.00	0.00	0.00
411	ADHEr, ACK2, FRD2, ST COAS	30.09	0.00	16.64	-8.95	0.00	0.00	1.11	-0.18	0.49	0.00	0.00
412	ADHEr, FUM, G6PDHy, TAI	11.80	0.00	0.00	-6.43	0.00	0.27	0.76	-0.12	17.63	0.00	0.00
413	ADHEr, FUM, PGDHL, TAI	11.80	0.00	0.00	-6.43	0.00	0.27	0.76	-0.12	17.63	0.00	0.00
414	ADHEr, FUM, PGL, TAI	11.80	0.00	0.00	6.44	0.00	0.27	-0.76	-0.12	17.63	0.00	0.00
415	ADHEr, FUM, ME2, TPI	5.91	0.00	0.00	-4.11	0.00	0.00	0.51	0.08	13.43	0.00	0.00
416	ADHEr, FUM, ME2, PPK	5.91	0.00	0.00	-4.11	0.00	0.00	-0.51	-0.08	13.43	0.00	0.00
417	ADHEr, FRA, FUM, ME2	5.91	0.00	0.00	-4.11	0.00	0.00	-0.51	-0.08	13.43	0.00	0.00
418	ADHEr, FRD2, GLUDy, UDEL D	36.99	0.00	0.00	-2.07	0.00	0.00	0.26	-0.04	0.00	0.00	0.00
419	ADHEr, FRD2, COH D, THD2	37.30	0.29	0.00	-19.20	0.00	0.00	0.43	-0.07	0.00	0.00	15.46
420	ADHEr, G6PDHy, HEN1, TAI	12.92	0.00	0.00	-6.60	0.00	0.27	-0.78	-0.13	12.69	0.00	0.00
421	ADHEr, HEN1, PGDHL, TAI	12.92	0.00	0.00	6.60	0.00	0.27	0.78	-0.13	12.69	0.00	0.00
422	ADHEr, HEN1, PGL, TAI	12.92	0.00	0.00	-6.60	0.00	0.27	0.78	-0.13	12.69	0.00	0.00
423	ADHEr, MDL, PGDHL, TAI	11.43	0.00	0.00	-5.84	0.00	0.24	-0.69	-0.17	13.54	0.00	0.00
424	ADHEr, G6PDHy, MDL, TAI	11.43	0.00	0.00	5.84	0.00	0.24	0.69	-0.17	13.54	0.00	0.00
425	ADHEr, MDL, PGL, TAI	11.43	0.00	0.00	-5.84	0.00	0.24	-0.69	-0.17	13.54	0.00	0.00
426	ADHEr, PGDHL, TAI, TPI	8.73	0.00	0.00	-4.23	0.00	0.18	-0.50	0.08	15.97	0.00	0.00
427	ADHEr, FRA, PGDHL, TAI	8.73	0.00	0.00	-4.23	0.00	0.18	-0.50	0.08	15.97	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

428	ADHET, PEK, PGDEL, TAC	8.23	0.00	0.00	-4.23	0.00	0.18	-0.50	0.08	15.97	0.00	0.00
429	ADHET, GLYC, TAU, TPI	8.24	0.00	0.00	-4.26	0.00	0.00	-0.52	-0.09	15.38	0.00	0.00
430	ADHET, TAU, THDS, TPI	8.25	0.00	0.00	-4.21	0.00	0.00	0.52	0.09	16.41	0.00	0.00
431	ADHET, LDIH D, TAU, TPI	8.24	0.00	0.00	-4.21	0.00	0.00	-0.52	-0.09	16.41	0.00	0.00
432	ADHET, ASP, EDA, MDL, PGI	8.26	0.00	0.00	-4.39	0.00	0.00	0.05	0.01	0.00	0.00	0.00
433	ADHET, ATPS-g, FRD2, LDIH D, MFI	15.46	2.12	0.00	5.10	-2.00	0.00	-0.25	-0.04	0.00	0.00	0.00
434	ADHET, EDA, PET, PGI, PCK	10.28	0.00	0.00	1.38	0.00	0.00	0.17	0.03	0.00	0.00	0.00
435	ADHET, ATPS-g, FRD2, LDIH D, MDI	15.60	2.10	0.00	-5.18	-2.00	0.00	-0.25	-0.06	0.00	0.00	0.00
436	ADHET, EDA, PET, PGI, SUCD4	10.88	0.00	0.00	1.68	0.00	0.00	0.21	0.03	0.00	0.00	0.00
437	ADHET, EDA, NADH6, PGI, PGI	10.88	0.00	0.00	-1.68	0.00	0.00	-0.21	0.03	0.00	0.00	0.00
438	ADHET, EDA, NADH6, PGI, PCK	7.19	0.00	0.00	0.80	0.00	0.00	-0.10	-0.02	0.00	0.00	0.00
439	ADHET, ASP, LDIH D, MDL, PET	13.74	0.00	0.00	3.10	0.00	0.00	0.58	0.06	0.00	0.00	0.00
440	ADHET, GLUDY, HEXT, PET, PGI	12.79	0.00	0.00	2.67	0.00	0.00	-0.31	-0.05	0.00	0.00	0.00
441	ADHET, ACK, C, Cps, NADH6, PGI	13.18	0.00	0.00	3.35	2.00	0.00	0.29	0.05	0.20	0.00	0.00
442	ADHET, GLUDY, HEXT, NADH6, PGI	10.34	0.00	0.00	2.15	0.00	0.00	0.27	0.04	0.00	0.00	0.00
443	ADHET, EDA, GLUDY, NADH6, PGI	10.88	0.00	0.00	-2.37	0.00	0.00	-0.29	-0.05	0.00	0.00	0.00
444	ADHET, ACK, PET, PGI, SUCD4	10.52	0.00	2.18	1.66	0.00	0.00	0.21	0.03	0.00	0.00	0.00
445	ADHET, ACK, NADH6, PET, PGI	12.71	0.00	0.00	-2.25	0.00	0.00	-0.21	-0.04	0.00	0.00	0.00
446	ADHET, ACK, GLUDY, NADH6, PGI	13.53	0.00	0.31	1.61	2.00	0.00	0.52	0.05	0.00	0.00	0.00
447	ADHET, EDA, GLCps, NADH6, PGI	13.25	0.00	0.00	2.43	2.00	0.00	-0.30	-0.05	0.21	0.00	0.00
448	ADHET, ACK, CBMK2, NADH6, PGI	4.17	0.00	0.00	1.94	2.00	0.00	0.36	0.06	0.00	0.00	0.00
449	ADHET, ATPS-g, FRD2, NADH6, PGI	11.98	0.00	1.48	-1.92	-2.00	0.00	-0.24	-0.04	0.00	0.00	0.00
450	ADHET, ATPS-g, NADH6, PET, PGI	11.98	0.00	1.48	1.92	2.00	0.00	0.24	0.04	0.00	0.00	0.00
451	ADHET, ATPS-g, GLCps, NADH6, PET	11.92	0.00	0.00	-2.38	0.00	0.00	0.29	0.05	0.00	0.00	0.00
452	ADHET, ATPS-g, MDL, NADH6, PGI	14.76	0.00	0.00	3.75	0.00	0.00	0.46	0.08	0.00	0.00	0.00
453	ADHET, ATPS-g, GLCps, MDL, NADH6	13.76	0.00	0.00	-3.75	0.00	0.00	-0.46	-0.08	0.00	0.00	0.00
454	ADHET, ACK, EDA, GLUDY, LDIH D	11.37	0.00	0.00	3.12	0.00	0.00	-0.39	-0.06	0.00	0.00	0.00
455	ADHET, ATPS-g, NADH6, PGI, SUCD4	10.53	0.00	0.00	3.17	2.00	0.00	0.27	0.04	0.00	0.00	0.00
456	ADHET, ACK, GLUDY, LDIH D, SUCD4	10.89	0.00	0.00	-3.19	0.00	0.00	-0.39	-0.06	0.00	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

457	ADHET, ATPS4r, G6PDHs, MDH, THD2	8.72	0.00	0.00	3.81	0.00	0.00	0.47	0.08	0.00	0.00	0.00
458	ADHET, ATPS4r, MDH, PGL, THD2	8.72	0.00	0.00	-3.81	0.00	0.00	0.47	0.08	0.00	0.00	0.00
459	ADHET, ASPT, G6PDHs, MDH, PYK	-1.53	0.00	0.00	-2.21	0.00	0.00	-0.27	-0.04	0.00	0.00	0.00
460	ADHET, ASPT, FBA, MDH, PYK	-1.53	0.00	0.00	3.21	0.00	0.00	0.27	0.04	0.00	0.00	0.00
461	ADHET, ASPT, MDH, PGL, PYK	-1.53	0.00	0.00	-2.21	0.00	0.00	-0.27	-0.04	0.00	0.00	0.00
462	ADHET, FRLD2, LDH D, MDH, SUCOA5	6.97	2.29	0.00	4.35	0.00	0.00	-0.27	-0.04	0.00	0.00	0.00
463	ADHET, ASPT, LDH D, MDH, SUCOA5	7.01	2.11	0.00	4.33	0.00	0.00	-0.27	-0.04	0.00	0.00	0.00
464	ADHET, AC36, LDH D, MDH, SUCD4	13.99	0.00	0.00	-7.34	0.00	0.00	0.28	0.05	0.00	0.00	4.89
465	ADHET, ATPS4r, GLUDs, PGL, SUCD4	8.35	0.00	0.00	-2.11	-2.00	0.00	0.26	0.04	0.00	0.00	0.00
466	ADHET, ASPT, FUM, LDH D, MDH	6.45	1.98	0.00	4.33	0.00	0.00	-0.29	-0.05	0.00	0.00	0.00
467	ADHET, ASPT, LDH D, MAL S, MDH	6.45	1.98	0.00	-4.33	0.00	0.00	0.29	0.05	0.00	0.00	0.00
468	ADHET, ASPT, ICL, LDH D, MDH	6.45	1.98	0.00	-4.33	0.00	0.00	0.29	0.05	0.00	0.00	0.00
469	ADHET, AC36, LDH D, MDH, NADH6	25.20	0.00	0.00	5.48	10.00	0.00	0.54	0.09	4.43	0.00	1.10
470	ADHET, FRLD2, GLUDs, LDH D, PPC	10.70	0.00	0.00	-1.05	0.00	0.00	0.13	0.02	0.00	0.00	0.00
471	ADHET, FRLD2, LDH D, PPC, THD2	-10.60	0.00	0.00	-1.09	0.00	0.00	0.13	-0.02	0.00	0.00	0.00
472	ADHET, ACK3, ATPS4r, LDH D, SUCD4	12.66	0.00	0.00	4.09	2.00	0.00	0.51	0.08	0.00	0.00	0.00
473	ADHET, ACK3, ACS, PPC, PPC	10.35	0.00	0.00	0.96	2.00	0.00	0.12	0.02	0.00	0.00	0.00
474	ADHET, GLUDs, LDH D, PPC, PPC	-9.20	0.00	0.00	-1.38	-2.00	0.00	0.17	0.05	0.00	0.00	0.00
475	ADHET, ATPS4r, FRLD2, NADH6, SUCOA5	16.04	0.00	0.00	-4.14	-2.00	0.00	-0.51	-0.08	0.00	0.00	0.00
476	ADHET, LDH D, PPC, PPC, THD2	9.09	0.00	0.00	-1.43	2.00	0.00	0.18	0.03	0.00	0.00	0.00
477	ADHET, ASPT, ATPS4r, GLCps, MDH	11.52	0.00	6.37	-2.53	0.00	0.00	0.41	0.05	0.00	0.00	0.00
478	ADHET, G6PDHs, MDH, NADH6, THD2	24.76	0.00	0.00	-5.93	-15.00	0.00	-0.73	-0.12	0.00	0.00	0.00
479	ADHET, MDH, NADHs, PGL, THD2	24.76	0.00	0.00	5.93	15.00	0.00	-0.73	0.12	0.00	0.00	0.00
480	ADHET, ACK3, FBA, GLUDs, NADH6	23.51	0.00	0.00	3.52	20.00	0.00	0.44	-0.07	0.00	0.00	0.00
481	ADHET, ACK3, GLUDs, NADH6, PFK	23.51	0.00	0.00	-3.52	-20.00	0.00	-0.44	-0.07	0.00	0.00	0.00
482	ADHET, ACK3, GLUDs, NADH6, TPI	23.51	0.00	0.00	3.52	20.00	0.00	0.44	0.07	0.00	0.00	0.00
483	ADHET, ATPS4r, MEHC, NADH6, PGL	16.09	0.00	0.00	-3.23	-2.00	0.00	0.52	0.00	0.00	0.00	0.00
484	ADHET, ATPS4r, FRLD2, NADH6, PGL	16.09	0.00	0.00	-3.23	-2.00	0.00	-0.52	-0.00	0.00	0.00	0.00
485	ADHET, ATPS4r, G6PDHs, GLCps, MDH	17.81	0.00	0.00	2.93	0.00	0.00	0.36	0.06	10.94	0.00	0.00

5
10
15
20
25
30
35
40
45
50
55

486	ADHEC, ATPS4c, GL Cpos, MDEH, PGL	17.8	0.00	0.00	3.93	0.00	0.00	0.56	0.00	10.94	0.00	0.00
487	ADHEC, ACKc, FBA, GL Cpos, NADH6	24.17	0.00	0.00	3.83	30.00	0.00	0.47	0.00	0.00	0.00	0.00
488	ADHEC, ACKc, GL Cpos, NADH6, TPI	24.17	0.00	0.00	3.83	20.00	0.00	0.47	0.00	0.00	0.00	0.00
489	ADHEC, ACKc, GL Cpos, NADH6, PFK	24.17	0.00	0.00	3.83	20.00	0.00	0.47	0.00	0.00	0.00	0.00
490	ADHEC, ACKc, TDIH D, MDEH, SUCD4	1.57	0.00	0.00	3.46	0.00	0.00	0.43	0.00	0.00	0.00	0.00
491	ADHEC, ACKc, AKG2, ATPS4c, FBA	29.63	0.00	0.00	1.94	38.95	0.00	0.24	0.00	0.00	0.00	0.00
492	ADHEC, ACKc, AKG2, ATPS4c, PFK	29.63	0.00	0.00	1.94	38.95	0.00	0.24	0.00	0.00	0.00	0.00
493	ADHEC, ACKc, AKG2, ATPS4c, TPI	4.07	0.00	0.00	1.94	13.49	0.00	0.24	0.00	0.00	0.00	0.00
494	ADHEC, FLDX, PGL, PFK, SUCD4	2.89	0.00	0.00	0.62	0.00	0.00	0.05	0.00	0.00	0.00	0.00
495	ADHEC, ACKc, ATPS4c, FBA, SUCOAS	4.32	0.00	0.00	1.95	13.90	0.00	0.24	0.00	0.00	0.00	0.00
496	ADHEC, ACKc, ATPS4c, PFK, SUCOAS	4.32	0.00	0.00	1.95	13.90	0.00	0.24	0.00	0.00	0.00	0.00
497	ADHEC, ACKc, ATPS4c, SUCOAS, TPI	29.91	0.00	0.00	1.95	39.48	0.00	0.24	0.00	0.00	0.00	0.00
498	ADHEC, FRD2, TDIH D, ME2, SUCOAS	10.83	1.87	0.00	4.10	0.00	0.00	0.28	0.00	0.12	0.00	0.00
499	ADHEC, ACKc, CBMK2, FBA, NADH6	24.41	0.00	0.00	3.97	20.00	0.00	0.49	0.00	0.00	0.00	0.00
500	ADHEC, ACKc, CBMK2, NADH6, PFK	24.41	0.00	0.00	3.97	20.00	0.00	0.49	0.00	0.00	0.00	0.00
501	ADHEC, ACKc, CBMK2, NADH6, TPI	24.41	0.00	0.00	3.97	20.00	0.00	0.49	0.00	0.00	0.00	0.00
502	ADHEC, ACKc, NADH6, RPE, TPI	24.45	0.00	0.00	3.99	20.00	0.00	0.49	0.00	0.00	0.00	0.00
503	ADHEC, ACKc, NADH6, PFK, RPE	24.45	0.00	0.00	3.99	20.00	0.00	0.49	0.00	0.00	0.00	0.00
504	ADHEC, ACKc, FBA, NADH6, RPT	24.45	0.00	0.00	3.99	20.00	0.00	0.49	0.00	0.00	0.00	0.00
505	ADHEC, ACKc, ASNS2, FBA, NADH6	24.46	0.00	0.00	4.00	20.00	0.00	0.49	0.00	0.00	0.00	0.00

Table 8. Corresponding genes to be knocked out to prevent a particular reaction from occurring in *E. coli*.

Reaction Abbreviation	Reaction Stoichiometry*	Genes Encoding the Enzyme(s) Catalyzing Each Reaction&
ACKr	[c] : ac + atp <==> actp + adp	(b3115 or b2296 or b1849))
ACS	[c] : ac + atp + coa --> accoa + amp + ppi	b4069
ACT6	ac[p] + h[p] <==> ac[c] + h[c]	Non-gene associated
ADHEr	[c] : etoh + nad <==> acald + h + nadh [c] : acald + coa + nad <==> accoa + h + nadh	(b0356 or b1478 or b1241) (b1241 or b0351)
AKGD	[c] : akg + coa + nad --> co2 + nadh + succoa	(b0116 and b0726 and b0727)
ASNS2	[c] : asp-L + atp + nh4 --> amp + asn-L + h + ppi	b3744
ASPT	[c] : asp-L --> fum + nh4	b4139
ATPS4r	adp[c] + (4) h[p] + pi[c] <==> atp[c] + (3) h[c] + h2o[c]	((b3736 and b3737 and b3738) and (b3731 and b3732 and b3733 and b3734 and b3735)) or ((b3736 and b3737 and b3738) and (b3731 and b3732 and b3733 and b3734 and b3735) and b3739))
CBMK2	[c] : atp + co2 + nh4 <==> adp + cbp + (2) h	(b0521 or b0323 or b2874)
EDA	[c] : 2ddg6p --> g3p + pyr	b1850
ENO	[c] : 2pg <==> h2o + pep	b2779
FBA	[c] : fdp <==> dhap + g3p	(b2097 or b2925 or b1773)
FBP	[c] : fdp + h2o --> f6p + pi	(b4232 or b3925)
FDH2	for[p] + (2) h[c] + q8[c] --> co2 [c] + h[p] + q8h2[c] for[p] + (2) h[c] + mqn8[c] --> co2[c] + h[p] + mql8[c]	((b3892 and b3893 and b3894) or (b1474 and b1475 and b1476))
FRD2	[c] : fum + mql8 --> mqn8 + succ [c] : 2dmmql8 + fum --> 2dmmq8 + succ	(b4151 and b4152 and b4153 and b4154)
FTHFD	[c] : 10fthf + h2o --> for + h + thf	b1232
FUM	[c] : fum + h2o <==> mal-L	(b1612 or b4122 or b1611)
G5SD	[c] : glu5p + h + nadph --> glu5sa + nadp + pi	b0243
G6PDHy	[c] : g6p + nadp <==> 6pgl + h + nadph	b1852
GLCpts	glc-D[p] + pep[c] --> g6p[c] + pyr[c]	((b2417 and b1101 and b2415 and b2416) or (b1817 and b1818 and b1819 and b2415 and b2416) or (b2417 and b1621 and b2415 and b2416))
GLU5K	[c] : atp + glu-L --> adp + glu5p	b0242
GLUDy	[c] : glu-L + h2o + nadp <==> akg + h + nadph + nh4	b1761

EP 2 679 676 B1

(continued)

	Reaction Abbreviation	Reaction Stoichiometry*	Genes Encoding the Enzyme(s) Catalyzing Each Reaction&
5	GLYCL	[c] : gly + nad + thf --> co2 + mlthf + nadh + nh4	(b2904 and b2903 and b2905 and b0116)
	HEX1	[c] : atp + glc-D --> adp + g6p + h	b2388
10	ICL	[c] : icit --> glx + succ	b4015
	LDH_D	[c] : lac-D + nad <==> h + nadh + pyr	(b2133 or b1380)
15	MALS	[c] : accoa + glx + h2o --> coa + h + mal-L	(b4014 or b2976)
	MDH	[c] : mal-L + nad <==> h + nadh + oaa	b3236
20	ME2	[c] : mal-L + nadp --> co2 + nadph + pyr	b2463
	MTHFC	[c] : h2o + methf <==> 10fthf + h	b0529
25	NADH12	[c] : h + mqn8 + nadh --> mql8 + nad [c] : h + nadh + q8 --> nad + q8h2 [c] : 2dmmq8 + h + nadh --> 2dmmql8 + nad	b1109
30	NADH6	(4) h[c] + nadh[c] + q8[c] --> (3) h[p] + nad[c] + q8h2[c] (4) h[c] + mqn8[c] + nadh[c] --> (3) h[p] + mql8[c] + nad[c] 2dmmq8[c] + (4) h[c] + nadh[c] --> 2dmmql8[c] + (3) h[p] + nad[c]	(b2276 and b2277 and b2278 and b2279 and b2280 and b2281 and b2282 and b2283 and b2284 and b2285 and b2286 and b2287 and b2288)
35	PFK	[c] : atp + f6p --> adp + fdp + h	(b3916 or b1723)
40	PFLi	[c] : coa + pyr --> accoa + for	((b0902 and b0903) and b2579) or (b0902 and b0903) or (b0902 and b3114) or (b3951 and b3952))
	PGDH	[c] : 6pgc + nadp --> co2 + nadph + ru5p-D	b2029
	PGI	[c] : g6p <==> f6p	b4025
45	PGL	[c] : 6pgl + h2o --> 6pgc + h	b0767
	PGM	[c] : 2pg <==> 3pg	(b3612 or b4395 or b0755)
	PPC	[c] : co2 + h2o + pep --> h + oaa + pi	b3956
50	PPCK	[c] : atp + oaa --> adp + co2 + pep	b3403
	PRO1z	[c] : fad + pro-L --> 1pyr5c + fadh2 + h	b1014
55	PYK	[c] : adp + h + pep --> atp + pyr	b1854 or b1676)
	PYRt2	h[p] + pyr[p] <==> h[c] + pyr[c]	Non-gene associated

(continued)

Reaction Abbreviation	Reaction Stoichiometry*	Genes Encoding the Enzyme(s) Catalyzing Each Reaction&
RPE	[c] : ru5p-D <==> xu5p-D	(b4301 or b3386)
SO4t2	so4[e] <==> so4[p]	(b0241 or b0929 or b1377 or b2215)
SUCD4	[c] : q8 + succ --> fum + q8h2	(b0721 and b0722 and b0723 and b0724)
SUCOAS	[c] : atp + coa + succ <==> adp + pi + succoa	(b0728 and b0729)
SULabc	atp[c] + h2o[c] + so4[p] --> adp [c] + h[c] + pi[c] + so4[c]	((b2422 and b2425 and b2424 and b2423) or (b0763 and b0764 and b0765) or (b2422 and b2424 and b2423 and b3917))
TAL	[c] : g3p + s7p <==> e4p + f6p	(b2464 or b0008)
THD2	(2) h[p] + nadh[c] + nadp[c] --> (2) h[c] + nad[c] + nadph[c]	(b1602 and b1603)
THD5	[c] : nad + nadph --> nadh + nadp	(b3962 or (b1602 and b1603))
TPI	[c] : dhap <==> g3p	b3919

Table 9. Metabolite names corresponding to abbreviations used in Table 8.

Metabolite Abbreviation	Metabolite Name
10fthf	10-Formyltetrahydrofolate
1pyr5c	1-Pyrroline-5-carboxylate
2ddg6p	2-Dehydro-3-deoxy-D-gluconate 6-phosphate
2dmmq8	2-Demethylmenaquinone 8
2dmmql8	2-Demethylmenaquinol 8
2pg	D-Glycerate 2-phosphate
3pg	3-Phospho-D-glycerate
6c	6-Phospho-D-gluconate
6pgl	6-phospho-D-glucono-1,5-lactone
ac	Acetate
acald	Acetaldehyde
accoa	Acetyl-CoA
actp	Acetyl phosphate
adp	ADP
akg	2-Oxolutarate
amp	AMP
asn-L	L-Asparagine
asp-L	L-Aspartate
atp	ATP
cbp	Carbamoyl phosphate
co2	CO2

EP 2 679 676 B1

(continued)

Metabolite Abbreviation	Metabolite Name
coa	Coenzyme A
dhap	Dihydroxyacetone phosphate
e4p	D-Erythrose 4-phosphate
etoh	Ethanol
f6p	D-Fructose 6-phosphate
fad	Flavin adenine dinucleotide oxidized
fadh2	Flavin adenine dinucleotide reduced
fdp	D-Fructose 1,6-bisphosphate
for	Formate
fum	Fumarate
g3p	Glyceraldehyde 3-phosphate
g6p	D-Glucose 6-phosphate
glc-D	D-Glucose
glu5p	L-Glutamate 5-phosphate
glu5sa	L-Glutamate 5-semialdehyde
glu-L	L-Glutamate
glx	Glyoxylate
gly	Glycine
h	H ⁺
h2o	H ₂ O
icit	Isocitrate
lac-D	D-Lactate
mal-L	L-Malate
methf	5,10-Methenyltetrahydrofolate
mlthf	5,10-Methylenetetrahydrofolate
mql8	Menaquinol 8
mqn8	Menaquinone 8
nad	Nicotinamide adenine dinucleotide
nadh	Nicotinamide adenine dinucleotide - reduced
nadp	Nicotinamide adenine dinucleotide phosphate
nadph	Nicotinamide adenine dinucleotide phosphate-reduced
nh4	Ammonium
oaa	Oxaloacetate
pep	Phosphoenolpyruvate
pi	Phosphate
ppi	Diphosphate
pro-L	L-Proline
pyr	Pyruvate

(continued)

Metabolite Abbreviation	Metabolite Name
q8	Ubiquinone-8
q8h2	Ubiquinol-8
ru5p-D	D-Ribulose 5-phosphate
s7p	Sedoheptulose 7-phosphate
so4	Sulfate
succ	Succinate
succoa	Succinyl-CoA
thf	5,6,7,8-Tetrahydrofolate
xu5p-D	D-Xylulose 5-phosphate

[0155] A number of criteria were applied to select the most practical sets of genes to target for removal. First, the designs were limited to include only knockouts that would not significantly (that is, > 5%) reduce the maximum theoretical yield of BDO under anaerobic conditions with or without the presence of nitrate as an electron acceptor. Such knockouts would create an artificial ceiling on any future metabolic engineering efforts and are thus undesirable. To this end, a series of linear programming (LP) problems were solved that maximized the BDO yield for the wild-type *E. coli* metabolic network assuming every reaction was individually deleted from the network. As used herein, reference to the wild-type *E. coli* network assumes that the BDO pathway is available. The term "wild-type" is thus a surrogate name for the undeleted *E. coli* network. Reactions whose deletion negatively affects the maximum BDO yield assuming PEP carboxykinase to be irreversible or reversible are shown in Tables 10 and 11, respectively. Table 10 shows reactions which, when deleted, reduce the maximum theoretical BDO yield under anaerobic conditions with or without the presence of nitrate, assuming that PEP carboxykinase cannot be used to produce oxaloacetate. Strain AB3 contains deletions in ADHE, LDH_D, and PFLi. 'Inf' indicates that the non-growth associated energetic requirements cannot be satisfied.

Table 10. Reactions which, when deleted, reduce the maximum theoretical BDO yield under anaerobic conditions with or without the presence of nitrate, assuming that PEP carboxykinase cannot be used to produce oxaloacetate.

		WT, Anaerobic		WT, Nitrate		AB3, Anaerobic	
	MAXIMUM MASS YIELD	0.477g/g		0.528 g/g		0.477 g/g	
Abbreviation	Reaction Name	BDO Yield	% of Max Yield	BDOYield	% of Max Yield	BDO Yield	% of Max Yield
4HBACT	4-hydroxybutyrate acetyl-CoA transferase	0.00	0%	0.00	0%	0.00	0%
4HBDH	4-hydroxybutyrate dehydrogenase	0.00	0%	0.00	0%	0.00	0%
4HBALDDH	4-hydroxybutyraldehyde dehydrogenase	0.00	0%	0.00	0%	0.00	0%
ACKr	acetate kinase	0.44	93%	0.50	95%	0.44	93%
ACONT	aconitase	0.41	86%	0.48	91%	0.41	86%
ACT6	acetate transport in/out via proton symport	0.42	89%	0.53	100%	0.40	83%
ATPS4r	ATP synthase (four protons for one ATP)	0.45	95%	0.45	86%	0.45	95%
BTDP2	1,4 butanediol dehydrogenase	0.00	0%	0.00	0%	0.00	0%
BTDt1	1,4-butanediol transport (Diffusion)	0.00	0%	0.00	0%	0.00	0%

EP 2 679 676 B1

(continued)

Abbreviation	Reaction Name	BDO Yield	% of Max Yield	BDOYield	% of Max Yield	BDO Yield	% of Max Yield
CO2t	CO2 transport out via diffusion	0.36	75%	0.36	68%	0.27	57%
CS	citrate synthase	0.41	86%	0.48	91%	0.41	86%
ENO	enolase	0.03	7%	0.46	88%	Inf	Inf
FBA	fructose-bisphosphate aldolase	0.47	98%	0.53	100%	0.47	98%
FRD3	fumarate reductase	0.47	99%	0.53	100%	0.47	99%
FUM	fumarase	0.48	100%	0.53	100%	0.48	100%
GAPD	glyceraldehyde-3-phosphate dehydrogenase (NAD)	Inf	Inf	0.44	82%	Inf	Inf
GLCpts	D-glucose transport via PEP:Pyr PTS	0.45	94%	0.52	99%	0.45	94%
H2Ot5	H2O transport via diffusion	0.44	91%	0.47	88%	0.39	81%
ICDH _y	isocitrate dehydrogenase (NADP)	0.48	100%	0.53	100%	0.48	100%
ICL	Isocitrate lyase	0.48	100%	0.53	100%	0.48	100%
MALS	malate synthase	0.48	100%	0.53	100%	0.48	100%
NADH6	NADH dehydrogenase (ubiquinone-8 & 3.5 protons)	0.48	100%	0.53	100%	0.48	100%
NADH8	NADH dehydrogenase (demethylmenaquinone-8 & 2.8 protons)	0.47	99%	0.53	100%	0.47	99%
NO3R1	Nitrate reductase (Ubiquinol-8)	0.48	100%	0.52	99%	0.48	100%
NO3t7	nitrate transport in via nitrite antiport	0.48	100%	0.48	90%	0.48	100%
PDH	pyruvate dehydrogenase	0.43	89%	0.51	97%	0.29	60%
PFK	phosphofructokinase	0.47	98%	0.53	100%	0.47	98%
PGI	glucose-6-phosphate isomerase	0.43	90%	0.52	98%	0.43	90%
PGK	phosphoglycerate kinase	Inf	Inf	0.44	82%	Inf	Inf
PGM	phosphoglycerate mutase	0.03	7%	0.46	88%	Inf	Inf
PPC	phosphoenolpyruvate carboxylase	0.40	84%	0.52	98%	0.14	30%
PTAr	phosphotransacetylase	0.44	93%	0.50	95%	0.44	93%
SSALcoax	CoA-dependant succinate semialdehyde dehydrogenase	0.40	83%	0.52	98%	0.31	64%
TPI	triose-phosphate isomerase	0.35	74%	0.50	95%	0.35	74%
		AB3, Nitrate		AB3 MDH ASPT, Anaerobic		AB3 MDHASPT, Nitrate	
	MAXIMUM MASS YIELD	0.528 g/g		0.477 g/g		0.528 g/g	

EP 2 679 676 B1

(continued)

	Abbreviation	Reaction Name	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield
5	4HBACT	4-hydroxybutyrate acetyl-CoA transferase	0.00	0%	0.00	0%	0.00	0%
10	4HBDH	4-hydroxybutyrate dehydrogenase	0.00	0%	0.00	0%	0.00	0%
	4HBTALDDH	4-hydroxybutyraldehyde dehydrogenase	0.00	0%	0.00	0%	0.00	0%
15	ACKr	acetate kinase	0.50	95%	0.44	93%	0.50	95%
	ACONT	aconitase	0.48	91%	0.32	67%	0.43	81%
	ACT6	acetate transport in/out via proton symport	0.53	100%	0.39	83%	0.53	100%
20	ATPS4r	ATP synthase (four protons for one ATP)	0.45	86%	0.45	95%	0.45	86%
	BTDP2	1,4 butanediol dehydrogenase	0.00	0%	0.00	0%	0.00	0%
25	BTDIl	1,4-butanediol transport (Diffusion)	0.00	0%	0.00	0%	0.00	0%
	CO2t	CO2 transport out via diffusion	0.27	52%	0.26	54%	0.26	49%
	CS	citrate synthase	0.48	91%	0.32	67%	0.43	81%
	ENO	enolase	0.46	88%	Inf	Inf	0.45	85%
30	FBA	fructose-bisphosphate aldolase	0.53	100%	0.46	96%	0.52	99%
	FRD3	fumarate reductase	0.53	100%	0.47	99%	0.53	100%
	FUM	fumarase	0.53	100%	0.40	84%	0.52	99%
35	GAPD	glyceraldehyde-3-phosphate dehydrogenase (NAD)	0.44	82%	Inf	Inf	0.42	79%
	GLCpts	D-glucose transport via PEP:Pyr PTS	0.52	99%	0.45	94%	0.52	99%
40	H2Ot5	H2O transport via diffusion	0.39	73%	0.36	76%	0.37	70%
	ICDH _{Hy}	isocitrate dehydrogenase (NADP)	0.53	100%	0.46	96%	0.51	97%
	ICL	Isocitrate lyase	0.53	100%	0.37	78%	0.52	99%
45	MALS	malate synthase	0.53	100%	0.40	84%	0.52	99%
	NADH6	NADH dehydrogenase (ubiquinone-8 & 3.5 protons)	0.53	100%	0.48	100%	0.53	100%
50	NADH8	NADH dehydrogenase (demethylmenaquinone-8 & 2.8 protons)	0.53	100%	0.47	99%	0.53	100%
	NO3R1	Nitrate reductase (Ubiquinol-8)	0.52	99%	0.48	100%	0.52	99%
	NO3t7	nitrate transport in via nitrite antiport	0.48	90%	0.48	100%	0.48	90%
55	PDH	pyruvate dehydrogenase	0.48	91%	0.22	46%	0.47	90%
	PFK	phosphofructokinase	0.53	100%	0.46	96%	0.52	99%

(continued)

Abbreviation	Reaction Name	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield
PGI	glucose-6-phosphate isomerase	0.52	98%	0.24	51%	0.52	98%
PGK	phosphoglycerate kinase	0.44	82%	Inf	Inf	0.42	79%
PGM	phosphoglycerate mutase	0.46	88%	Inf	Inf	0.45	85%
PPC	phosphoenolpyruvate carboxylase	0.52	98%	0.00	0%	0.00	0%
PTAr	phosphotransacetylase	0.50	95%	0.44	93%	0.50	95%
SSALcoax	CoA-dependant succinate semialdehyde dehydrogenase	0.52	98%	0.19	39%	0.52	98%
TPI	triose-phosphate isomerase	0.50	95%	0.28	58%	0.49	93%

[0156] Table 11 shows reactions which, when deleted, reduce the maximum theoretical BDO yield under anaerobic conditions with or without the presence of nitrate, assuming that PEP carboxykinase can be used to produce oxaloacetate. 'Inf' indicates that the non-growth associated energetic requirements cannot be satisfied.

Table 11. Reactions which, when deleted, reduce the maximum theoretical BDO yield under anaerobic conditions with or without the presence of nitrate, assuming that PEP carboxykinase can be used to produce oxaloacetate.

		WT, Anaerobic		WT, Nitrate		AB3, Anaerobic	
	MAXIMUM MASS YIELD	0.545 g/g		0.545 g/g		0.545 g/g	
Abbreviation	Reaction Name	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield
4HBACT	4-hydroxybutyrate acetyl-CoA transferase	0.00	0%	0.00	0%	0.00	0%
4HBDH	4-hydroxybutyrate dehydrogenase	0.00	0%	0.00	0%	0.00	0%
4HBTALDDH	4-hydroxybutyralde-hyde dehydrogenase	0.00	0%	0.00	0%	0.00	0%
ACKr	acetate kinase	0.50	92%	0.53	97%	0.50	92%
ACONT	aconitase	0.47	86%	0.51	94%	0.47	86%
BTDP2	1,4 butanediol dehydrogenase	0.00	0%	0.00	0%	0.00	0%
BTDtl	1,4-butanediol transport (Diffusion)	0.00	0%	0.00	0%	0.00	0%
CO2t	CO2 transport out via diffusion	0.36	67%	0.36	67%	0.33	61%
CS	citrate synthase	0.47	86%	0.51	94%	0.47	86%
ENO	enolase	0.07	14%	0.46	85%	Inf	Inf
GAPD	glyceraldehyde-3-phosphate dehydrogenase (NAD)	Inf	Inf	0.44	80%	Inf	Inf
H2Ot5	H2O transport via diffusion	0.50	92%	0.50	92%	0.46	84%

EP 2 679 676 B1

(continued)

Abbreviation	Reaction Name	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield
ICDH _y	isocitrate dehydrogenase (NADP)	0.50	92%	0.53	97%	0.50	92%
PDH	pyruvate dehydrogenase	0.53	97%	0.54	99%	0.39	71%
PGI	glucose-6-phosphate isomerase	0.54	98%	0.54	100%	0.54	98%
PGK	phosphoglycerate kinase	Inf	Inf	0.44	80%	Inf	Inf
PGM	phosphoglycerate mutase	0.07	14%	0.46	85%	Inf	Inf
PPCK	Phosphoenol-pyruvate carboxykinase	0.48	88%	0.53	97%	0.48	88%
PTAr	Phosphotrans-acetylase	0.50	92%	0.53	97%	0.50	92%
SSALcoax	CoA-dependant succinate semialdehyde dehydrogenase	0.52	95%	0.54	98%	0.48	89%
TPI	triose-phosphate isomerase	0.44	81%	0.52	96%	0.44	81%
		AB3, Nitrate		AB3 MDH ASPT, Anaerobic		AB3 MDH ASPT, Nitrate	
	MAXIMUM MASS YIELD	0.545 g/g		0.545 g/g		0.545 g/g	
Abbreviation	Reaction Name	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield
4HBACT	4-hydroxybutyrate acetyl-CoA transferase	0.00	0%	0.00	0%	0.00	0%
4HBDH	4-hydroxybutyrate dehydrogenase	0.00	0%	0.00	0%	0.00	0%
4HBTALDDH	4-hydroxybutyralde-hyde dehydrogenase	0.00	0%	0.00	0%	0.00	0%
ACKr	acetate kinase	0.53	97%	0.48	89%	0.53	96%
ACONT	aconitase	0.51	94%	0.36	66%	0.45	83%
BTDP2	1,4 butanediol dehydrogenase	0.00	0%	0.00	0%	0.00	0%
BTDtl	1,4-butanediol transport (Diffusion)	0.00	0%	0.00	0%	0.00	0%
CO2t	CO2 transport out via diffusion	0.33	61%	0.29	54%	0.30	54%
CS	citrate synthase	0.51	94%	0.36	66%	0.45	83%
ENO	enolase	0.46	85%	0.00	0%	0.46	85%
GAPD	glyceraldehyde-3-phosphate dehydrogenase (NAD)	0.44	80%	0.00	0%	0.43	80%
H2Ot5	H2O transport via diffusion	0.46	84%	0.44	81%	0.44	81%

(continued)

Abbreviation	Reaction Name	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield	BDO Yield	% of Max Yield
ICDH _y	isocitrate dehydrogenase (NADP)	0.53	97%	0.50	92%	0.53	97%
PDH	pyruvate dehydrogenase	0.51	94%	0.31	57%	0.50	92%
PGI	glucose-6-phosphate isomerase	0.54	100%	0.40	73%	0.53	98%
PGK	phosphoglycerate kinase	0.44	80%	0.00	0%	0.43	80%
PGM	phosphoglycerate mutase	0.46	85%	0.00	0%	0.46	85%
PPCK	Phosphoenol-pyruvate carboxykinase	0.53	97%	0.48	88%	0.53	97%
PTAr	Phosphotrans-acetylase	0.53	97%	0.48	89%	0.53	96%
SSALcoax	CoA-dependant succinate semialdehyde dehydrogenase	0.54	98%	0.46	84%	0.54	98%
TPI	triose-phosphate isomerase	0.52	96%	0.40	73%	0.52	95%

[0157] The above-described analysis led to three critical observations. One critical observation was that acetate kinase and phosphotransacetylase are required to achieve the maximum BDO yields under all conditions by regenerating acetyl-CoA from the acetate produced by 4-hydroxybutyrate:acetyl-CoA transferase. This finding strongly suggests that eliminating acetate formation by deleting *ackA-pta* may not be a viable option. Thus a successful strain will likely have to provide an intracellular environment where converting acetate to acetyl-CoA is beneficial and thermodynamically feasible. Otherwise, a set of enzymes capable of performing the required BDO reductions without passing through a CoA derivative would have to be found or the co-production of 1 mol of acetate per mol of BDO would have to be accepted.

[0158] A second critical observation was that the TCA cycle enzymes citrate synthase (CS), aconitase (ACONT), and isocitrate dehydrogenase (ICDH_y) are required to achieve the maximum BDO yields under all conditions. This indicates that the reverse TCA cycle flux from oxaloacetate to succinate to succinyl-CoA must be complemented to some extent by CS, ACONT, and ICDH_y for maximum production.

[0159] A third critical observation was that supplanting PEP carboxylase with PEP carboxykinase in *E. coli* can positively impact the BDO program. The maximum BDO yield under anaerobic conditions with and without the presence of nitrate is 3% and 12% lower, respectively, if PEP carboxylase carries out the PEP to oxaloacetate conversion as compared to if PEP carboxykinase carries out the conversion. Furthermore, under anaerobic conditions without nitrate addition, PEP carboxykinase can lessen the requirement for pyruvate dehydrogenase activity for maximum BDO production. Specifically, the maximum BDO yield drops 11% if this typically aerobic enzyme has no activity if PEP carboxykinase is assumed irreversible as compared to a 3% reduction if PEP carboxykinase can catalyze the production of oxaloacetate. An alternative to pyruvate dehydrogenase activity can be utilized by coupling non-native formate dehydrogenase, capable of catalyzing the reduction of formate to carbon dioxide, to pyruvate formate lyase activity.

[0160] The next two criteria applied to evaluate the OptKnock designs were the number of required knockouts and the predicted BDO yield at maximum growth. Analysis of the one, two, and three reaction deletion strategies in Table 6 (that is, PEP carboxykinase assumed irreversible) revealed that so few knockouts were insufficient to prevent high acetate yields. The predicted acetate/BDO ratios for all one, two, or three deletion designs, was at least 1.5. Even allowing for four deletions led to only two designs, #99 and #100, with predicted BDO yields above 0.35 g/g, and those designs suggest the removal of the glycolysis gene, *pgi*, which encodes phosphoglucose isomerase. Given the anticipated importance of glycolysis in the fermentation of *E. coli*, it was decided to pursue such high-risk designs only as a last resort. Furthermore, the suggested deletions lower the maximum theoretical yield in designs #99 and #100 by 8% and 15%, respectively. The highest producing four deletion strategy that did not negatively impact the maximum theoretical yield was design #129, which had a predicted BDO yield at maximum growth of only 0.26 g/g.

[0161] Only one five deletion design in Table 6 satisfies all criteria for a successful design. This knockout strategy involves the removal of ADHEr (alcohol dehydrogenase), PFLi (pyruvate formate lyase), LDH_D (lactate dehydrogenase), MDH (malate dehydrogenase), and ASPT (aspartate transaminase). The suggested knockouts do not reduce the max-

imum theoretical BDO yield under any of the conditions examined. A strain engineered with these knockouts is predicted to achieve a BDO yield at maximum growth of 0.37 g/g assuming anaerobic conditions and PEP carboxykinase irreversibility. This design has several desirable properties. Most notably, it prevents the network from producing high yields of the natural fermentation products, ethanol, formate, lactate, and succinate. The prevention of homosuccinate production via the MDH deletion as opposed to removing PEP carboxylase, fumarase, or fumarate reductase, is particularly intriguing because it blocks the energy-yielding fermentation pathway from oxaloacetate to succinate that could arise if PEP carboxykinase is assumed reversible without negatively impacting the maximum BDO yield. In this design, succinate semialdehyde can be made via succinyl-CoA or alpha-ketoglutarate. The succinyl-CoA is formed from succinate via succinyl-CoA synthetase. The succinate can be formed from both the reverse TCA cycle reactions (PEP carboxylase, PEP carboxykinase, fumarase, fumarate reductase) and the glyoxylate shunt (malate synthase, isocitrate lyase).

[0162] The BDO versus biomass solution boundaries for the ADHER, PFLi, LDH_D, MDH, ASPT knockout strategy are shown in Figure 7A and 7B, assuming PEP carboxykinase irreversibility or reversibility, respectively. Note that the solution boundaries are obtained using a genome-scale model of *E. coli* metabolism as opposed to the reduced model because these calculations are not CPU-intensive. The solution boundaries reveal that the growth coupling of BDO is robust with respect to the assumption of PEP carboxykinase reversibility. However, the deletion of the proton-pumping transhydrogenase (THD2) or glutamate dehydrogenase (GLUDy) may be necessary to achieve an obligatory coupling of cell growth with BDO production. The only negative aspect of the design is that the MDH and ASPT deletions drop the maximum ATP yield of BDO production slightly (~20%) if PEP carboxykinase reversibility is assumed. This causes the optimal growth solution of the design strategy to drop below the black BDO vs. biomass line of the wild-type network in Figure 7B. However, this finding suggests the possibility of engineering an optimal balance of MDH activity where enough is present to ensure efficient BDO production while also being limited enough to prevent succinate from becoming the major fermentation product. Note that succinate is predicted to be the major fermentation product of the wild-type network if PEP carboxykinase reversibility is assumed.

[0163] Tables 10 and 11 list reactions whose deletion negatively impacts the maximum BDO yield in an intermediate strain, referred to as AB3, which lacks ADHER, LDH_D, and PFLi as well as a strain lacking ASPT and MDH in addition to the AB3 deletions. Note that the suggested deletions place a very high importance on obtaining pyruvate dehydrogenase, citrate synthase, and aconitase activity under completely anaerobic conditions. If sufficient pyruvate dehydrogenase activity cannot be attained, an alternative is to leave PFLi intact and supplement its activity with a non-native formate dehydrogenase that can capture one reducing equivalent while converting formate to carbon dioxide.

[0164] Figure 8 and Table 12 depict the flux ranges that the *E. coli* network can attain while reaching either the maximum BDO yield (cases 1-4) or the maximum biomass yield (case 5) under anaerobic conditions. Cases 1 and 2 assume that no gene deletions have taken place. Case 2 exhibits tighter flux ranges than case 1 due to the fact that an additional constraint enforcing the maximum ATP yield at the maximum BDO yield is imposed. Cases 3 and 4 are analogous to cases 1 and 2 except that the fluxes encoding the reactions for ADHER, ASPT, LDH_D, MDH, and PFLi have been set to zero. The flux ranges assuming biomass yield maximization in the presence of the ADHER, ASPT, LDH_D, MDH, and PFLi knockouts are shown in case 5.

[0165] Table 12 shows achievable ranges of central metabolic fluxes under anaerobic conditions assuming PEP carboxykinase to be reversible. Bold flux values were set as constraints on the system. Five cases are considered: Case 1, maximum BDO yield of the wild-type network; Case 2, maximum ATP yield assuming the maximum BDO yield of the wild-type network; Case3, maximum BDO yield of the network with fluxes through ADHER, ASPT, LDH_D, MDH, and PFLi set to zero; Case 4, maximum ATP yield assuming the maximum BDO yield of the network with fluxes through ADHER, ASPT, LDH_D, MDH, and PFLi set to zero; and Case 5, maximum biomass yield of the network with fluxes through ADHER, ASPT, LDH_D, MDH, and PFLi set to zero.

Table 12. Achievable ranges of central metabolic fluxes under anaerobic conditions assuming PEP carboxykinase to be reversible. Reactions that are assumed inactive in cases 3, 4, and 5 are indicated with bold font.

Reaction Abbreviation	CASE 1		CASE 2		CASE 3		CASE 4		CASE 5	
	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX
	mmol/gDW/hr									
GLCpts	0.0	20.0	18.2	18.2	0.0	20.0	20.0	20.0	20.0	20.0
HEX 1	0.0	20.0	1.8	1.8	0.0	20.0	0.0	0.0	0.0	0.0
G6PDHy	0.0	27.3	0.0	0.0	0.0	15.3	0.0	0.0	0.0	0.0
PGL	0.0	27.3	0.0	0.0	0.0	15.3	0.0	0.0	0.0	0.0
PGDH	0.0	27.3	0.0	0.0	0.0	15.3	0.0	0.0	0.0	0.0

EP 2 679 676 B1

(continued)

	CASE 1		CASE 2		CASE 3		CASE 4		CASE 5	
Reaction Abbreviation	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX
	mmol/gDW/hr									
RPE	-6.4	18.2	0.0	0.0	-6.2	10.2	0.0	0.0	-0.2	-0.2
RPI	-9.1	0.0	0.0	0.0	-6.2	0.0	0.0	0.0	-0.2	-0.2
TKT1	-3.2	9.1	0.0	0.0	-3.1	5.1	0.0	0.0	-0.1	-0.1
TKT2	-3.2	9.1	0.0	0.0	-3.1	5.1	0.0	0.0	-0.2	-0.2
TAL	-3.2	9.1	0.0	0.0	-3.1	5.1	0.0	0.0	-0.1	-0.1
EDA	0.0	12.7	0.0	0.0	0.0	10.5	0.0	0.0	0.0	0.0
PGDHY	0.0	12.7	0.0	0.0	0.0	10.5	0.0	0.0	0.0	0.0
PGI	-7.3	20.0	20.0	20.0	4.7	20.0	20.0	20.0	20.0	20.0
FBP	0.0	15.9	0.0	0.0	0.0	12.7	0.0	0.0	0.0	0.0
PFK	7.3	35.9	20.0	20.0	9.5	32.7	20.0	20.0	19.7	19.7
FBA	7.3	20.0	20.0	20.0	9.5	20.0	20.0	20.0	19.7	19.7
TPI	7.3	20.0	20.0	20.0	9.5	20.0	20.0	20.0	19.7	19.7
GAPD	27.3	40.0	40.0	40.0	29.5	40.0	40.0	40.0	39.3	39.3
PGK	27.3	40.0	40.0	40.0	29.5	40.0	40.0	40.0	39.3	39.3
PGM	27.3	43.6	40.0	40.0	29.5	43.6	40.0	40.0	39.0	39.0
ENO	27.3	43.6	40.0	40.0	29.5	43.6	40.0	40.0	39.0	39.0
PYK	0.0	29.5	0.0	0.0	0.0	29.5	1.8	1.8	10.7	10.7
PDH	0.0	34.5	14.5	18.2	12.0	34.5	21.8	21.8	30.1	30.1
PFLi	0.0	11.9	0.0	3.6	0.0	0.0	0.0	0.0	0.0	0.0
PPC	0.0	15.9	0.0	0.0	0.0	12.7	0.0	0.0	0.0	0.0
PPCK	5.5	51.4	21.8	21.8	5.5	30.9	18.2	18.2	8.1	8.1
CS	9.1	34.1	18.2	18.2	13.1	32.7	18.2	18.2	7.5	7.5
CITL	0.0	15.9	0.0	0.0	0.0	12.7	0.0	0.0	0.0	0.0
ACONT	9.1	21.8	18.2	18.2	13.1	21.8	18.2	18.2	7.5	7.5
ICDH _y	1.8	21.4	18.2	18.2	1.8	21.3	14.5	14.5	0.2	0.2
AKGD	0.0	21.4	0.0	18.2	0.0	21.3	0.0	14.5	0.0	0.0
SUCOAS	0.5	20.0	3.6	3.6	0.6	20.0	7.3	7.3	14.9	14.9
FRD	0.0	12.7	3.6	3.6	0.0	8.7	3.6	3.6	7.5	7.5
FUM	-55.5	12.7	-14.5	3.6	-5.1	8.7	3.6	3.6	7.2	7.2
MDH	-50.9	33.2	-14.5	3.6	0.0	0.0	0.0	0.0	0.0	0.0
ICL	0.0	18.2	0.0	0.0	0.0	18.2	3.6	3.6	7.2	7.2
MALS	0.0	16.4	0.0	0.0	0.0	16.4	3.6	3.6	7.2	7.2
ME	0.0	29.5	0.0	0.0	0.0	12.7	0.0	0.0	0.0	0.0
ASPTA	0.0	66.7	0.0	18.2	0.0	7.5	0.0	0.0	0.6	0.6
ASPT	0.0	66.7	0.0	18.2	0.0	0.0	0.0	0.0	0.0	0.0
LDH _D	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

(continued)

	CASE 1		CASE 2		CASE 3		CASE 4		CASE 5	
Reaction Abbreviation	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX
	mmol/gDW/hr									
ADHEr	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
PTAr	5.9	37.7	21.8	21.8	9.1	34.5	21.8	21.8	0.1	0.1
ACKr	5.9	37.7	21.8	21.8	9.1	34.5	21.8	21.8	0.1	0.1
GLUDc	0.0	21.4	0.0	18.2	0.0	21.3	0.0	14.5	0.0	0.0
ABTA	0.0	21.4	0.0	18.2	0.0	21.2	0.0	14.5	0.0	0.0
SSAL_coa	0.5	37.7	3.6	21.8	0.6	34.5	7.3	21.8	14.8	14.8
ATPM	0.0	15.9	15.9	15.9	0.0	12.7	12.7	12.7	7.6	7.6
BDOsyn	21.8	21.8	21.8	21.8	21.8	21.8	21.8	21.8	14.8	14.8
	1/hr									
BIOMASS	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.21	0.21

[0166] The first two four deletion designs (#92 and #93) listed in Table 7 (that is, PEP carboxykinase assumed reversible) were considered next due to their relatively high predicted BDO yields. Design #92 (ADHEr, HEX1, PFLi, and PGI) would need additional knockouts that eliminate succinate and lactate production and the maximum BDO yield of design #93 (ADHEr, EDA, NADH6, PGI) was only 90% of the theoretical maximum of the wild-type network. Both designs called for the removal of PGI which, as mentioned above, was ruled undesirable due to the anticipated importance of glycolysis on fermentation. Design #98 (ADHEr, ATPS4r, FDH2, NADH6) was the first four deletion design to require the removal of ATP synthase without additionally requiring the PGI knockout. However, this design would also require the deletion of genes to prevent lactate and succinate production in order to raise the predicted BDO yield at maximum growth. Upon further analysis, it was found that nearly all promising designs in Table 7 could be improved by ensuring that the deletions (i.e., ADHEr, LDH_D, MDH, ASPT, and PFLi) were also implemented if not already specified.

[0167] Efforts were next focused on identifying reaction deletions that could supplement the core set (that is, ADHEr, LDH_D, MDH, ASPT, PFLi). Table 13 shows known *E. coli* genes responsible for catalyzing the reactions targeted for removal. The designation "[c]" refers to cytosolic. The genes encoding these reactions along with the reactions of the core set are provided in Table 13. The effect of the additional deletions on the BDO versus biomass solution boundaries is shown in Figure 9. Notably, the coupling between BDO and biomass production becomes more and more pronounced as additional deletions are accumulated. The predicted BDO yield at maximum growth after accumulating all deletions is 0.46 g/g. Lastly, none of the deletions negatively impact the maximum theoretical BDO yield.

Table 13. Known *E. coli* genes responsible for catalyzing the reactions targeted for removal.

Reaction Abbreviation	Reaction Stoichiometry	Genes Encoding the Enzyme(s) Catalyzing Each Reaction &
ADHEr	[c] : etoh + nad <=> acald + h + nadh [c] : acald + coa + nad <==> accoa + h + nadh	(b0356 or b1478 or b1241) (b1241 or b0351)
PFLi	[c] : coa + pyr --> accoa + for	((b0902 and b0903) and b2579) or (b0902 and b0903) or (b0902 and b3114) or (b3951 and b3952))
MDH	[c] : mal-L + nad <==> h + nadh + oaa	b3236
ASPT	[c] : asp-L --> fum + nh4	b4139
LDH_D	[c] : lac-D + nad <==> h + nadh + pyr	(b2133 or b1380)
DHAPT	[c] : dha + pep --> dhap + pyr	(b1200 and b1199 and b1198 and b2415 and b2416)

(continued)

Reaction Abbreviation	Reaction Stoichiometry	Genes Encoding the Enzyme(s) Catalyzing Each Reaction &
DRPA	[c] : 2dr5p --> acald + g3p	b4381
PYK	[c] : adp + h + pep --> atp + pyr	(b1854 or b1676)
EDD	[c] : 6pgc --> 2ddg6p + h2o	b1851
GLYCLTDx GLYCLTDy	[c] glx + h + nadh --> glyclt + nad [c] glx + h + nadph --> glyclt + nadp	(b3553 or b1033)
MCITS	[c] : h2o + oaa + ppcoa --> 2mcit + coa + h	b0333
INSK	[c] : atp + ins --> adp + h + im	b0477

[0168] In the results shown in Table 13, OptKnock identifies reactions to be eliminated from an organism to enhance biochemical production. Any combination (that is, at least one and at most all) of the listed gene deletions could conceivably have the desired effect of ensuring that the corresponding reaction is non-functional in *E. coli*. The most practical experimental strategy for eliminating the reactions targeted for removal must be determined on a case-by-case basis.

EXAMPLE VI

Generation of Engineered Strains

[0169] In order to validate the computational predictions of Example V, the strains are constructed, evolved, and tested. *Escherichia coli* K-12 MG1655 serves as the wild-type strain into which the deletions are introduced. The strains are constructed by incorporating in-frame deletions using homologous recombination via the λ Red recombinase system of (Datsenko and Wanner, Proc. Natl. Acad. Sci. USA 97:6640-6645 (2000)). The approach involves replacing a chromosomal sequence (that is, the gene targeted for removal) with a selectable antibiotic resistance gene, which itself is later removed. The knockouts are integrated one by one into the recipient strain. No drug resistance markers or scars will remain after each deletion, allowing accumulation of multiple mutations in each target strain. The deletion technology completely removes the gene targeted for removal so as to substantially reduce the possibility of the constructed mutants reverting back to the wild-type. During the initial stages of strain development, non-native genes enabling BDO production are expressed in a synthetic operon behind an inducible promoter on a medium- or high-copy plasmid; for example the PBAD promoter which is induced by arabinose, on a plasmid of the pBAD series (Guzman et al., J. Bacteriol. 177:4121-4130 (1995)). This promoter is known to be very easily titratable, allowing expression to be fine tuned over a 1000-fold range of arabinose concentrations. If BDO production is successful, these genes are then be integrated into the chromosome to promote stability.

[0170] The engineered strains are characterized by measuring growth rate, substrate uptake rate, and product/by-product secretion rate. These strains are initially anticipated to exhibit suboptimal growth rates until their metabolic networks have adjusted to their missing functionalities. To enable this adjustment, the strains are adaptively evolved. By subjecting the strains to adaptive evolution, cellular growth rate becomes the primary selection pressure and the mutant cells are compelled to reallocate their metabolic fluxes in order to enhance their rates of growth. This reprogramming of metabolism has been recently demonstrated for several *E. coli* mutants that had been adaptively evolved on various substrates to reach the growth rates predicted *a priori* by an *in silico* model (Fong and Palsson, Nat. Genet. 36:1056-1058 (2004)). Should the OptKnock predictions prove successful, the growth improvements brought about by adaptive evolution are accompanied by enhanced rates of BDO production. Adaptive evolution is performed in triplicate (running in parallel) due to differences in the evolutionary patterns witnessed previously in *E. coli* (Fong and Palsson, *supra*, 2004; Fong et al., J. Bacteriol. 185:6400-6408 (2003); Ibarra et al., Nature 420:186-189 (2002)) that could potentially result in one strain having superior production qualities over the others. Evolutions are run for a period of 2-6 weeks, depending on the rate of growth improvement obtained. In general, evolutions are stopped once a stable growth phenotype is obtained.

[0171] Following the adaptive evolution process, the new strains are again characterized by measuring growth rate, substrate uptake rate, and product/byproduct secretion rate. These results are compared to the OptKnock predictions by plotting actual growth and production yields along side the production described above. The most successful OptKnock design/evolution combinations are chosen to pursue further, and are characterized in lab-scale batch and continuous fermentations. The growth-coupled biochemical production concept behind the OptKnock approach also results in the

generation of genetically stable overproducers. Thus, the cultures are maintained in continuous mode for one month to evaluate long-term stability. Periodic samples are taken to ensure that yield and productivity are maintained throughout the process.

EXAMPLE VII

OptKnock Strains for Production of BDO

[0172] As described in Examples V and VI, the application of the OptKnock methodology has been applied for generating promising deletion targets to generate BDO producing strains. OptKnock identifies reactions to be eliminated from an organism to couple the biochemical production and biomass yields. The designs provide a list of the metabolic reactions to be targeted for removal by OptKnock. The *E. coli* genes known to encode the enzymes that catalyze each reaction were also provided to describe which genetic modifications must be implemented to realize the predicted growth-coupled production phenotypes. Obviously, if new discoveries reveal that additional genes in the *E. coli* genome can confer one or more of the reaction functionalities targeted for removal in a given design, then these genes should be removed as well as the ones described herein. Note that preventing the activity of only a subset (that is, at least one and at most all) of the reactions in each of the designs may sometimes be sufficient to confer a growth-coupled producing phenotype. For example, if a design calls for the removal of a particular reaction whose activity *in vivo* is not sufficient to uncouple growth from BDO production, then the genes encoding the enzymes that catalyze this reaction can be left intact. In addition, any combination (that is, at least one and at most all) of the listed gene deletions for a given reaction could conceivably have the desired effect of ensuring that the reaction is non-functional in *E. coli*.

[0173] Multiple deletion strategies are listed in Table 6 and 7 for enhancing the coupling between 1,4-butanediol production and *E. coli* growth assuming PEP carboxykinase to be irreversible and reversible, respectively. One design (that is, ADHER, ASPT, MDH, LDH_D, PFLi) emerged as the most promising upon satisfying multiple criteria. The suggested deletions 1) led to a high predicted BDO yield at maximum growth, 2) required a reasonable number of knockouts, 3) had no detrimental effect on the maximum theoretical BDO yield, 4) brought about a tight coupling of BDO production with cell growth, and 5) was robust with respect to the irreversibility/reversibility of PEP carboxykinase. The following list specifies the minimal set of required gene deletions predicted to render BDO the major fermentation product of *E. coli*:

adhE (b1421), *ldhA* (b1380).

pflAB (b0902, b0903) is not included in the minimal set because its deletion forces a reliance on pyruvate dehydrogenase to provide sufficient acetyl-CoA for cell growth and one reducing equivalent from pyruvate. As pyruvate dehydrogenase activity is low under anaerobic conditions and inhibited by high NADH concentrations, a plausible alternative to the *pflAB* deletion is to add a non-native formate dehydrogenase to *E. coli* that can capture the reducing power that is otherwise lost via formate secretion. Nevertheless, adding *pflAB* to the minimal deletion set yields:

adhE (b1421), *ldhA* (b1380), *pflAB* (b0902, b0903).

mdh (b3236) is not included in the minimal set because there are multiple deletions capable of preventing succinate from becoming the major fermentation product of *E. coli* as opposed to BDO. Examples include the genes encoding fumarase and/or fumarate reductase. However, eliminating malate dehydrogenase appears to be the most logical choice to attenuate succinate production as it leaves intact a pathway for the conversion of the glyoxylate shunt product, malate, to BDO. Adding the malate dehydrogenase deletion to the minimal set above yields:

adhE (b1421), *ldhA* (b1380), *pflAB* (b0902, b0903), *mdh* (b3236).

[0174] The gene, *mgo*, which encodes a malate:quinone-oxidoreductase, is believed to catalyze the oxidation of malate to oxaloacetate (van der Rest et al., J. Bacteriol. 182:6892-6899 (2000)). However, if it is shown to also catalyze the formation of malate from oxaloacetate, its removal will be necessary to ensure that it does not circumvent the *mdh* deletion. This leads to the deletion set:

adhE (b1421), *ldhA* (b1380), *pflAB* (b0902, b0903), *mdh* (b3236), *mgo* (b2210).

aspA is left out of the minimal set as it is questionable whether or not aspartate deaminase can carry enough flux to circumvent the malate dehydrogenase deletion. However, if this scenario is indeed possible, then the minimal list of required deletions becomes:

adhE (b1421), *ldhA* (b1380), *pflAB* (b0902, b0903), *mdh* (b3236), *aspA* (b4139).

[0175] For the calculations above, NADH and NADPH-dependent malic enzymes of *E. coli* were assumed to operate irreversibly catalyzing only the conversion of malate to carbon dioxide and pyruvate. If these enzymes can also catalyze the formation of malate from pyruvate and carbon dioxide, the genes encoding one or both malic enzymes will have to be removed to prevent succinate from becoming the major fermentation product. This leads to the following sets of deletions:

adhE (b1421), *mdh* (b3236), *ldhA* (b1380), *pflAB* (b0902, b0903), *sfcA* (b1479)

adhE (b1421), *mdh* (b3236), *ldhA* (b1380), *pflAB* (b0902, b0903), *maeB* (b2463)

adhE (b1421), *mdh* (b3236), *ldhA* (b1380), *pflAB* (b0902, b0903), *sfcA* (b1479), *maeB* (b2463)

[0176] The minimal set of deletions can be supplemented with additional deletions aimed at tightening the coupling of BDO production to cell growth. These sets of the deletions are listed below.

adhE (b1421), *ldhA* (b1380), *pflAB* (b0902, b0903), *mdh* (b3236), *pntAB* (b1602, b1603)

adhE (b1421), *ldhA* (b1380), *pflAB* (b0902, b0903), *mdh* (b3236), *gdhA* (b1761)

adhE (b1421), *ldhA* (b1380), *pflAB* (b0902, b0903), *mdh* (b3236), *pykA* (b1854), *pykF* (b1676), *dhaKLM* (b1198, b1199, b1200), *deoC* (b4381), *edd* (b1851), *yiaE* (b3553), *ycdW* (b1033)

adhE (b1421), *ldhA* (b1380), *pflAB* (b0902, b0903), *mdh* (b3236), *pykA* (b1854), *pykF* (b1676), *dhaKLM* (b1198, b1199, b1200), *deoC* (b4381), *edd* (b1851), *yiaE* (b3553), *ycdW* (b1033), *prpC* (b0333), *gsk* (b0477)

[0177] Strains possessing the deletions listed in this Example can be supplemented with additional deletions if it is found that the suggested deletions do not reduce the activity of their corresponding reactions to the extent required to attain growth coupled BDO production or if native *E. coli* genes, through adaptive evolution or mutagenesis, attain mutations conferring activities capable of circumventing the proposed design strategies.

[0178] Disclosed items:

1. A non-naturally occurring microorganism comprising one or more gene disruptions, said one or more gene disruptions occurring in genes encoding an enzyme obligatory to coupling 1,4-butanediol production to growth of said microorganism when said gene disruption reduces an activity of said enzyme, whereby said one or more gene disruptions confers stable growth-coupled production of 1,4-butanediol onto said non-naturally occurring microorganism.

2. The non-naturally occurring microorganism of item 1, wherein said one or more gene disruptions comprise a metabolic modification listed in Table 6 or 7.

3. The non-naturally occurring microorganism of item 1, wherein said one or more gene disruptions comprise a deletion of said one or more genes.

4. The non-naturally occurring microorganism of item 1, wherein said non-naturally occurring microorganism is selected from the group of microorganisms having a metabolic modification listed in Table 6 or 7.

5. The non-naturally occurring microorganism of item 1, wherein said microorganism comprises a bacterium, yeast or fungus.

6. The non-naturally occurring microorganism of item 5, wherein said bacteria comprises a species selected from *Escherichia coli*, *Klebsiella oxytoca*, *Anaerobiospirillum succiniciproducens*, *Actinobacillus succinogenes*, *Mannheimia succiniciproducens*, *Rhizobium etli*, *Bacillus subtilis*, *Corynebacterium glutamicum*, *Gluconobacter oxydans*, *Zymomonas mobilis*, *Lactococcus lactis*, *Lactobacillus plantarum*, *Streptomyces coelicolor*, *Clostridium acetobutylicum*, *Pseudomonas fluorescens*, and *Pseudomonas putida*.

7. The non-naturally occurring microorganism of item 5, wherein said yeast comprises a species selected from *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces marxianus*, *Aspergillus terreus*, *Aspergillus niger*, and *Pichia pastoris*.

8. A non-naturally occurring microorganism comprising a set of metabolic modifications obligatory to coupling 1,4-butanediol production to growth of said microorganism, said set of metabolic modifications comprising disruption of

one or more genes, or an ortholog thereof, wherein said set of metabolic modifications comprises disruption of *adhE* and *ldhA*, wherein said microorganism exhibits stable growth-coupled production of 1,4-butanediol.

9. The non-naturally occurring microorganism of item 8, wherein said set of metabolic modifications further comprise disruption of *mdh*.

10. The non-naturally occurring microorganism of item 9, wherein said set of metabolic modifications further comprise disruption of one or more genes selected from the set of genes comprising *mgo*, *aspA*, *sfcA*, *maeB*, *pntAB*, and *gdhA*.

11. The non-naturally occurring microorganism of item 10, wherein said set of metabolic modifications comprise disruption of *sfcA* and *maeB*.

12. The non-naturally occurring microorganism of item 9, wherein said set of metabolic modifications further comprise disruption of one or more genes selected from the set of genes comprising *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE*, *ycdW*, *prpC*, and *gsk*.

13. The non-naturally occurring microorganism of item 12, wherein said set of metabolic modifications comprise disruption of *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE* and *ycdW*.

14. The non-naturally occurring microorganism of item 13, wherein said set of metabolic modifications further comprise disruption of *prpC* and *gsk*.

15. The non-naturally occurring microorganism of any of items 9-14, wherein said set of metabolic modifications further comprise disruption of *pflAB*.

16. The non-naturally occurring microorganism of item 8, wherein said set of metabolic modifications comprise disruption of one or more genes selected from the set of genes comprising *adhE*, *ldhA*, *pflAB*, *mdh*, and *aspA*.

17. The method of item 8, wherein said set of metabolic modifications comprise disruption of the genes *adhE*, *ldhA*, *pflAB*, *mdh*, and *aspA*.

18. The non-naturally occurring microorganism of item 8, wherein said microorganism further comprises a 1,4-butanediol (BDO) biosynthetic pathway, said pathway comprising at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or alcohol dehydrogenase, wherein said exogenous nucleic acid is expressed in sufficient amounts to produce 1,4-butanediol (BDO).

19. The non-naturally occurring microorganism of item 8, wherein said disruption of one or more genes comprises a deletion.

20. The non-naturally occurring microorganism of item 8, wherein said microorganism comprises a bacterium, yeast or fungus.

21. The non-naturally occurring microorganism of item 20, wherein said bacteria comprises a species selected from *E. coli*, *K. oxytoca*, *A. succiniciproducens*, *A. succinogenes*, *M. succiniciproducens*, *R. etli*, *B. subtilis*, *C. glutamicum*, *G. oxydans*, *Z. mobilis*, *L. lactis*, *L. plantarum*, *S. coelicolor*, *C. acetobutylicum*, *P. fluorescens*, and *P. putida*.

22. The non-naturally occurring microorganism of item 20, wherein said yeast comprises a species selected from *S. cerevisiae*, *S. pombe*, *K. lactis*, *K. marxianus*, *A. terreus*, *A. niger*, and *P. pastoris*.

23. A method of producing a non-naturally occurring microorganism having stable growth-coupled production of 1,4-butanediol, comprising:

- (a) identifying *in silico* a set of metabolic modifications requiring 1,4-butanediol production during exponential growth, and
- (b) genetically modifying a microorganism to contain said set of metabolic modifications requiring 1,4-butanediol

production.

24. The method of item 23, wherein said set of metabolic modifications are identified by an *in silico* method selected from OptKnock or SimPheny.

25. The method of item 23, wherein said set of metabolic modifications comprise functional disruption of one or more metabolic reactions.

26. The method of item 25, wherein said metabolic modifications are disrupted by gene deletion.

27. The method of item 25, wherein said non-naturally occurring microorganism comprises a microorganism having a metabolic modification selected from the set of metabolic modifications listed in Table 6 or 7.

28. The method of item 23, further comprising culturing said genetically modified microorganism.

29. The method of item 28, further comprising adaptively evolving said genetically modified microorganism under conditions requiring 1,4-butanediol production.

30. The method of item 23, wherein said non-naturally occurring microorganism comprises a bacterium, yeast or fungus.

31. The method of item 30, wherein said bacteria comprises a species selected from *E. coli*, *K. oxytoca*, *A. succiniciproducens*, *A. succinogenes*, *M. succiniciproducens*, *R. etli*, *B. subtilis*, *C. glutamicum*, *G. oxydans*, *Z. mobilis*, *L. lactis*, *L. plantarum*, *S. coelicolor*, *C. acetobutylicum*, *P. fluorescens*, and *P. putida*.

32. The non-naturally occurring microorganism of item 30, wherein said yeast comprises a species selected from *S. cerevisiae*, *S. pombe*, *K. lactis*, *K. marxianus*, *A. terreus*, *A. niger*, and *P. pastoris*.

33. A microorganism produced by the method of item 23 or 29.

34. A method of producing 1,4-butanediol coupled to the growth of a microorganism, comprising:

(a) culturing under exponential growth phase in a sufficient amount of nutrients and media a non-naturally occurring microorganism comprising a set of metabolic modifications obligatorily coupling 1,4-butanediol production to growth of said microorganism, said set of metabolic modifications comprising disruption of one or more genes, or an ortholog thereof, wherein said set of metabolic modifications comprises disruption of *adhE* and *ldhA*, wherein said microorganism exhibits stable growth-coupled production of 1,4-butanediol; and

(b) isolating 1,4-butanediol produced from said non-naturally occurring microorganism.

35. The method of item 34, wherein said set of metabolic modifications further comprise disruption of *mdh*.

36. The method of item 35, wherein said set of metabolic modifications further comprise disruption of one or more genes selected from the set of genes comprising *mgo*, *aspA*, *sfcA*, *maeB*, *pntAB*, and *gdhA*.

37. The method of item 36, wherein said set of metabolic modifications comprise disruption of *sfcA* and *maeB*.

38. The method of item 35, wherein said set of metabolic modifications further comprise disruption of one or more genes selected from the set of genes comprising *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE*, *ycdW*, *prpC*, and *gsk*.

39. The method of item 38, wherein said set of metabolic modifications comprise disruption of *pykA*, *pykF*, *dhaKLM*, *deoC*, *edd*, *yiaE* and *ycdW*.

40. The method of item 39, wherein said set of metabolic modifications further comprise disruption of *prpC* and *gsk*.

41. The method of any of items 34-40, wherein said set of metabolic modifications further comprise disruption of *pflAB*.

42. The method of item 34, wherein said set of metabolic modifications comprise disruption of one or more genes

selected from the set of genes comprising *adhE*, *ldhA*, *pflAB*, *mdh*, and *aspA*.

43. The method of item 34, wherein said set of metabolic modifications comprise disruption of the genes *adhE*, *ldhA*, *pflAB*, *mdh*, and *aspA*.

44. The method of item 34, wherein said non-naturally occurring microorganism further comprises a 1,4-butanediol (BDO) biosynthetic pathway, said pathway comprising at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or alcohol dehydrogenase, wherein said exogenous nucleic acid is expressed in sufficient amounts to produce 1,4-butanediol (BDO).

Claims

1. A method for manufacturing a downstream product of 4-hydroxybutyrate or 1,4-butanediol, comprising:

- (a) culturing under exponential growth phase in a sufficient amount of nutrients and media a non-naturally occurring microorganism comprising disruption of the genes *adhE*, *ldhA*, *pflAB* and *mdh*, wherein said microorganism expresses at least one exogenous nucleic acid encoding an enzyme in a 4-hydroxybutyrate or 1,4-butanediol biosynthetic pathway in sufficient amounts to produce 4-hydroxybutyrate or 1,4-butanediol, said at least one exogenous nucleic acid encoding 4-hydroxybutanoate dehydrogenase, CoA-independent succinic semialdehyde dehydrogenase, succinyl-CoA synthetase, CoA-dependent succinic semialdehyde dehydrogenase, 4-hydroxybutyrate:CoA transferase, glutamate:succinic semialdehyde transaminase, glutamate decarboxylase, CoA-independent aldehyde dehydrogenase, CoA-dependent aldehyde dehydrogenase or alcohol dehydrogenase, and wherein said microorganism produces 4-hydroxybutyrate or 1,4-butanediol;
- (b) optionally isolating 4-hydroxybutyrate or 1,4-butanediol; and
- (c) chemically converting 4-hydroxybutyrate or 1,4-butanediol to the downstream product.

2. The method of claim 1, wherein the microorganism exhibits stable growth-coupled production of 1,4-butanediol.

3. The method of claim 1 or 2, wherein the downstream product is gamma-butyrolactone.

4. The method of claim 3, further comprising the step of isolating gamma-butyrolactone.

5. The method of claim 3 or 4, wherein the gamma-butyrolactone is chemically converted from 4-hydroxybutyrate.

6. The method of claim 3 or 4, wherein the gamma-butyrolactone is chemically converted from 1,4-butanediol.

7. The method of claim 1 or 2, wherein the downstream product is tetrahydrofuran.

8. The method of claim 1 or 2, wherein the downstream product is pyrrolidone.

9. The method of any one of claims 1-8, wherein the 1,4-butanediol, 4-hydroxybutyrate or gamma-butyrolactone is isolated from a fermentation culture.

10. The method of any one of claims 1-9, wherein the 1,4-butanediol or gamma-butyrolactone is purified by distillation.

Patentansprüche

1. Verfahren zur Herstellung eines nachgelagerten ("downstream") Produkts von 4-Hydroxybutyrat oder 1,4-Butandiol, umfassend:

- (a) das Kultivieren eines nicht natürlich vorkommenden Mikroorganismus, der eine Störung der Gene *adhE*, *ldhA*, *pflAB* und *mdh* umfasst, in exponentieller Wachstumsphase in einer ausreichenden Menge von Nährstoffen und Medien, wobei der Mikroorganismus mindestens eine exogene Nukleinsäure, die ein Enzym in einem 4-

EPO - Munich
64

11. März 2013




FIGURE 1




FIGURE 2




FIGURE 3





FIGURE 4



Biochemical production envelopes of a:

- ☒ Typical Production Strain
- ☐ OptKnock-Designed Strain

Potential phenotypes:

- ☐ Prior to Growth Selection
- ☒ Following Growth Selection

FIGURE 5




FIGURE 6





FIGURE 7




FIGURE 8




FIGURE 9