



(51) International Patent Classification:

C07D 491/04 (2006.01) *A61P 31/14* (2006.01)
A61K 31/7068 (2006.01)

(21) International Application Number:

PCT/US2015/021638

(22) International Filing Date:

20 March 2015 (20.03.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/968,763 21 March 2014 (21.03.2014) US

(71) Applicant: BRISTOL-MYERS SQUIBB COMPANY
[US/US]; Route 206 and Province Line Road, Princeton,
New Jersey 08543 (US).(72) Inventors: WANG, Tao; c/o Bristol-Myers Squibb Com-
pany, 5 Research Parkway, Wallingford, Connecticut
06492 (US). EASTMAN, Kyle J.; c/o Bristol-Myers
Squibb Company, 5 Research Parkway, Wallingford, Con-
necticut 06492 (US). ZHANG, Zhongxing; c/o Brist-
ol-Myers Squibb Company, 5 Research Parkway, Walling-
ford, Connecticut 06492 (US). PARCELLA, Kyle E.; c/o
Bristol-Myers Squibb Company, 5 Research Parkway,
Wallingford, Connecticut 06492 (US). YIN, Zhiwei; c/o
Bristol-Myers Squibb Company, 5 Research Parkway,
Wallingford, Connecticut 06492 (US). KADOW, John F.;
c/o Bristol-Myers Squibb Company, 5 Research Parkway,
Wallingford, Connecticut 06492 (US).(74) Agents: LEVIS, John F. et al.; Bristol-Myers Squibb
Company, Route 206 and Province Line Road, Princeton,
New Jersey 08543 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) Title: CYANO CONTAINING AZABENZOFURAN COMPOUNDS FOR THE TREATMENT OF HEPATITIS C



I

(57) Abstract: Compounds of Formula I, including their salts, as well as compositions and methods of using the compounds are set forth. The compounds have activity against hepatitis C virus (HCV) and may be useful in treating those infected with HCV: (I)

CYANO CONTAINING AZABENZOFURAN COMPOUNDS FOR THE
TREATMENT OF HEPATITIS C

CROSS REFERENCE TO RELATED APPLICATION

- 5 This application claims the priority of U.S. Provisional Application Serial No. 61/968,763 filed March 21, 2014, which is hereby incorporated by reference in its entirety.

FIELD OF THE INVENTION

- 10 The invention relates to novel compounds, including their salts, which have activity against hepatitis C virus (HCV) and which are useful in treating those infected with HCV. The invention also relates to compositions and methods of making and using these compounds.

15 BACKGROUND OF THE INVENTION

- Hepatitis C virus (HCV) is a major human pathogen, infecting an estimated 170 million persons worldwide - roughly five times the number infected by human immunodeficiency virus type 1. A substantial fraction of these HCV infected individuals develop serious progressive liver disease, including cirrhosis and hepatocellular carcinoma (Lauer, G. M.; Walker, B. D. *N. Engl. J. Med.* **2001**, 345, 41-52).
- 20

- HCV is a positive-stranded RNA virus. Based on a comparison of the deduced amino acid sequence and the extensive similarity in the 5'-untranslated region, HCV has been classified as a separate genus in the Flaviviridae family. All members of the Flaviviridae family have enveloped virions that contain a positive stranded RNA genome encoding all known virus-specific proteins via translation of a single, uninterrupted, open reading frame.\
- 25

- Considerable heterogeneity is found within the nucleotide and encoded amino acid sequence throughout the HCV genome. At least six major genotypes have been characterized, and more than 50 subtypes have been described. The major genotypes of HCV differ in their distribution worldwide, and the clinical significance of the genetic
- 30

heterogeneity of HCV remains elusive despite numerous studies of the possible effect of genotypes on pathogenesis and therapy.

The single strand HCV RNA genome is approximately 9500 nucleotides in length
5 and has a single open reading frame (ORF) encoding a single large polyprotein of about
3000 amino acids. In infected cells, this polyprotein is cleaved at multiple sites by
cellular and viral proteases to produce the structural and non-structural (NS) proteins. In
the case of HCV, the generation of mature non-structural proteins (NS2, NS3, NS4A,
NS4B, NS5A, and NS5B) is effected by two viral proteases. The first one is believed to
10 be a metalloprotease and cleaves at the NS2-NS3 junction; the second one is a serine
protease contained within the N-terminal region of NS3 (also referred to as NS3 protease)
and mediates all the subsequent cleavages downstream of NS3, both in cis, at the NS3-
NS4A cleavage site, and in trans, for the remaining NS4A-NS4B, NS4B-NS5A, NS5A-
NS5B sites. The NS4A protein appears to serve multiple functions, acting as a cofactor
15 for the NS3 protease and possibly assisting in the membrane localization of NS3 and
other viral replicase components. The complex formation of the NS3 protein with NS4A
seems necessary to the processing events, enhancing the proteolytic efficiency at all of the
sites. The NS3 protein also exhibits nucleoside triphosphatase and RNA helicase
activities. NS5B (also referred to as HCV polymerase) is a RNA-dependent RNA
20 polymerase that is involved in the replication of HCV. The HCV NS5B protein is
described in “Structural Analysis of the Hepatitis C Virus RNA Polymerase in Complex
with Ribonucleotides (Bressanelli; S. et al., *Journal of Virology* **2002**, 3482-3492; and
Defrancesco and Rice, *Clinics in Liver Disease* **2003**, 7, 211-242).


25 Currently, the most effective HCV therapy employs a combination of alpha-
interferon and ribavirin, leading to sustained efficacy in 40% of patients (Poynard, T. et
al. *Lancet* **1998**, 352, 1426-1432). Recent clinical results demonstrate that pegylated
alpha-interferon is superior to unmodified alpha-interferon as monotherapy (Zeuzem, S.
et al. *N. Engl. J. Med.* **2000**, 343, 1666-1672). However, even with experimental
30 therapeutic regimens involving combinations of pegylated alpha-interferon and ribavirin,
a substantial fraction of patients do not have a sustained reduction in viral load. Thus,
there is a clear and important need to develop effective therapeutics for treatment of HCV
infection.

HCV-796, an HCV NS5B inhibitor, has shown an ability to reduce HCV RNA levels in patients. The viral RNA levels decreased transiently and then rebounded during dosing when treatment was with the compound as a single agent but levels dropped more robustly when combined with the standard of care which is a form of interferon and ribavirin. The development of this compound was suspended due to hepatic toxicity observed during extended dosing of the combination regimens. US patent 7,265,152 and the corresponding PCT patent application WO2004/041201 describe compounds of the HCV-796 class. Other compounds have been disclosed; see for example, WO2009/101022, as well as WO 2012/058125.

What is therefore needed in the art are additional compounds which are novel and effective against hepatitis C. Additionally, these compounds should provide advantages for pharmaceutical uses, for example, with regard to one or more of their mechanism of action, binding, inhibition efficacy, target selectivity, solubility, safety profiles, or bioavailability. Also needed are new formulations and methods of treatment which utilize these compounds.

SUMMARY OF THE INVENTION

One aspect of the invention is a compound of Formula I, including pharmaceutically acceptable salts thereof:



I

wherein

Z is C-R⁵ or N;

R⁰ is hydrogen;

R¹ is methyl;

R² is phenyl that is independently substituted with 0-2 halo or methoxy, or is para substituted with W-Ar;

5

W is -O- or -NH-;

Ar is phenyl or para-halophenyl;

10 R³ is hydrogen, fluoro, or chloro;

R⁴, R⁵, and R⁶ are independently selected from the group of hydrogen, halo, alkyl, haloalkyl, alkoxy, and perdeuteroalkoxy;

15 R^{7a}, R^{7b} are each independently selected from the group of hydrogen, alkyl, cycloalkyl, and Ar¹, or together

R^{7a} and R^{7b} form a 3-7 membered carbocyclic ring;

20 R⁸ is hydrogen,

Ar¹ is phenyl, a 5-membered heteroaromatic ring or a 6-membered heteroaromatic ring;

R⁹ is selected from the group of hydrogen, halo, R²⁰¹, OR²⁰², and NR²⁰³R²⁰⁴;

25

R²⁰¹ is alkyl, alkenyl, or C₁-C₄ alkyl with between one to all of the hydrogens replaced by fluoro;

R²⁰² is C₁-C₃ alkyl, or C₁-C₃ alkyl with between one to all of the hydrogens replaced by
30 fluoro;

R²⁰³ is hydrogen; and

R²⁰⁴ is C₁-C₃ alkyl, C₁-C₃ hydroxyalkyl, or is C₁-C₃ alkyl with between one to all of the hydrogens replaced by fluoro.

The invention also relates to pharmaceutical compositions comprising a
5 compound of Formula I, including pharmaceutically acceptable salts thereof, and a pharmaceutically acceptable carrier, excipient, and/or diluent.

In addition, the invention provides one or more methods of treating hepatitis C infection comprising administering a therapeutically effective amount of a compound of
10 Formula I to a patient.

Also provided as part of the invention are one or more methods for making the compounds of Formula I.

15 The present invention is directed to these, as well as other important ends, hereinafter described.

DETAILED DESCRIPTION OF THE EMBODIMENTS

The singular forms “a”, “an”, and “the” include plural reference unless the context
20 clearly dictates otherwise.

Unless otherwise specifically set forth elsewhere in the application, the following terms may be used herein and shall have the following meanings: “Hydrogen” or “H” refers to hydrogen, including its isotopes, such as deuterium which may be represented herein by the letter “D”. “Halo” means fluoro, chloro, bromo, or iodo. “Alkyl” means a
25 straight or branched alkyl group composed of 1 to 6 carbons. “Alkenyl” means a straight or branched alkyl group composed of 2 to 6 carbons with at least one double bond. “Cycloalkyl” means a monocyclic ring system composed of 3 to 7 carbons. “Hydroxyalkyl,” “alkoxy” and other terms with a substituted alkyl moiety include straight and branched isomers composed of 1 to 6 carbon atoms for the alkyl moiety. “Halo”
30 includes all halogenated isomers from monohalo substituted to perhalo substituted in substituents defined with halo, for example, “Haloalkyl” and “haloalkoxy”, “halophenyl”, “halophenoxy.” “Aryl” means a monocyclic or bicyclic aromatic hydrocarbon groups having 6 to 12 carbon atoms, or a bicyclic fused ring system wherein one or both of the

rings is a phenyl group. Bicyclic fused ring systems consist of a phenyl group fused to a four- to six-membered aromatic or non-aromatic carbocyclic ring. Representative examples of aryl groups include, but are not limited to, indanyl, indenyl, naphthyl, phenyl, and tetrahydronaphthyl. "Heteroaryl" means a 5 to 7 membered monocyclic or 8
5 to 11 membered bicyclic aromatic ring system with 1-5 heteroatoms independently selected from nitrogen, oxygen, and sulfur. Parenthetical and multiparenthetical terms are intended to clarify bonding relationships to those skilled in the art. For example, a term such as ((R)alkyl) means an alkyl substituent further substituted with the substituent R. Substituents which are illustrated by chemical drawing to bond at variable positions on a
10 multiple ring system (for example a bicyclic ring system) are intended to bond to the ring where they are drawn to append.


The invention includes all pharmaceutically acceptable salt forms of the compounds. Pharmaceutically acceptable salts are those in which the counter ions do not contribute significantly to the physiological activity or toxicity of the compounds and as
15 such function as pharmacological equivalents. These salts can be made according to common organic techniques employing commercially available reagents. Some anionic salt forms include acetate, acistrate, besylate, bromide, camsylate, chloride, citrate, fumarate, glucuronate, hydrobromide, hydrochloride, hydroiodide, iodide, lactate, maleate, mesylate, nitrate, pamoate, phosphate, succinate, sulfate, tartrate, tosylate, and
20 xinofoate. Some cationic salt forms include ammonium, aluminum, benzathine, bismuth, calcium, choline, diethylamine, diethanolamine, lithium, magnesium, meglumine, 4-phenylcyclohexylamine, piperazine, potassium, sodium, tromethamine, and zinc.

Some of the compounds of the invention possess asymmetric carbon atoms. The invention includes all stereoisomeric forms, including enantiomers and diastereomers as
25 well as mixtures of stereoisomers such as racemates. Some stereoisomers can be made using methods known in the art. Stereoisomeric mixtures of the compounds and related intermediates can be separated into individual isomers according to methods commonly known in the art. The use of wedges or hashes in the depictions of molecular structures in the following schemes and tables is intended only to indicate relative stereochemistry,
30 and should not be interpreted as implying absolute stereochemical assignments.

The invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen

include deuterium and tritium. Isotopes of carbon include ^{13}C and ^{14}C . Isotopically-labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed. Such compounds may have a variety of potential uses, for example as standards and reagents in determining biological activity. In the case of stable isotopes, such compounds may have the potential to favorably modify biological, pharmacological, or pharmacokinetic properties.

As set forth above, the invention is directed to one or more compounds of Formula I, including pharmaceutically acceptable salts thereof:



I

wherein

Z is C- R^5 or N;

R^0 is hydrogen;

R^1 is methyl;

R^2 is phenyl that is independently substituted with 0-2 halo or methoxy, or is para substituted with W-Ar;

W is -O- or -NH-;

Ar is phenyl or para-halophenyl;

R^3 is hydrogen, fluoro, or chloro;

R^4 , R^5 , and R^6 are independently selected from the group of hydrogen, halo, alkyl, haloalkyl, alkoxy, and perdeuteroalkoxy;

R^{7a} , R^{7b} are each independently selected from the group of hydrogen, alkyl, cycloalkyl, and Ar^1 , or together

R^{7a} and R^{7b} form a 3-7 membered carbocyclic ring;

R^8 is hydrogen;

Ar^1 is phenyl, a 5-membered heteroaromatic ring or a 6-membered heteroaromatic ring;

R^9 is selected from the group of hydrogen, halo, R^{201} , OR^{202} , and $NR^{203}R^{204}$;

R^{201} is alkyl, alkenyl, or C_1 - C_4 alkyl with between one to all of the hydrogens replaced by fluoro.

R^{202} is C_1 - C_3 alkyl, or C_1 - C_3 alkyl with between one to all of the hydrogens replaced by fluoro;

R^{203} is hydrogen; and

R^{204} is C_1 - C_3 alkyl, C_1 - C_3 hydroxyalkyl, or is C_1 - C_3 alkyl with between one to all of the hydrogens replaced by fluoro.

It is preferred in the compound of Formula I above that R^2 is para-fluorophenyl.

It is also preferred that R^3 is hydrogen.

In addition, it is preferred that R^4 , R^5 , and R^6 are each independently selected from the group of hydrogen, fluoro, $-OCH_3$, and $-OCD_3$.

It is further preferred that R^{7a} is selected from the group of hydrogen, methyl, fluoromethyl, and cyclopropyl.

It is also preferred that R^{7b} is selected from the group of hydrogen, methyl, fluoromethyl, cyclopropyl, and Ar^1 .

In certain embodiments, it is also preferred that together R^{7a} and R^{7b} form a cyclopropyl or cyclobutyl ring.

It is further preferred that Ar^1 is phenyl or pyrimidyl.

It is also preferred that R^9 is R^{201} or $NR^{203}R^{204}$.

In addition, it is preferred that R^{201} is $-CH_2CH_2CF_3$ or vinyl.

It is also preferred that R^{204} is $-CH_2CF_3$, $-CH_2CF_2CF_3$ or $-CH_2CH_2OH$.

In a further embodiment of the invention, it is preferred that R^2 is para-fluorophenyl, R^3 is hydrogen, R^4 , R^5 , and R^6 are each independently selected from the group of hydrogen, fluoro, $-OCH_3$, and $-OCD_3$,

R^{7a} is selected from the group of hydrogen, methyl, fluoromethyl, and cyclopropyl,

R^{7b} is selected from the group of hydrogen, methyl, fluoromethyl, cyclopropyl, and Ar^1 ,

or together R^{7a} and R^{7b} form a cyclopropyl or cyclobutyl ring;

Ar^1 is phenyl or pyrimidyl,

R^9 is R^{201} or $NR^{203}R^{204}$;

R^{201} is $-CH_2CH_2CF_3$ or vinyl; and

R^{204} is $-CH_2CF_3$, $-CH_2CF_2CF_3$ or $-CH_2CH_2OH$.

In a further embodiment of the invention, it is preferred that R^4 is hydrogen, R^5 is hydrogen or fluoro, R^{7b} is selected from the group of hydrogen, methyl, fluoromethyl, and cyclopropyl, or together R^{7a} and R^{7b} form a cyclopropyl or cyclobutyl ring; and R^{201} is $-\text{CH}_2\text{CH}_2\text{CF}_3$.

In certain embodiments of the compound of Formula I above, it is preferred that Z is N.

Also preferred are compounds wherein Z is N, R^4 is hydrogen, and R^6 is $-\text{OCD}_3$.

In certain embodiments of the compound of Formula I above, it is preferred that Z is CR^5 .

Other preferred compounds include those wherein Z is CR^5 , R^4 is hydrogen, R^5 is hydrogen or fluoro, R^6 is hydrogen, fluoro, or $-\text{OCH}_3$,

R^{7a} is selected from hydrogen, methyl, fluoromethyl, or cyclopropyl

R^{7b} is selected from hydrogen, methyl, fluoromethyl, or cyclopropyl

or together R^{7a} and R^{7b} form a cyclopropyl or cyclobutyl ring; and

R^{201} is $-\text{CH}_2\text{CH}_2\text{CF}_3$.

Also preferred are compounds of Formula I wherein R^5 is hydrogen, R^6 is fluoro, R^{7a} is methyl, R^{7b} is cyclopropyl, and R^9 is R^{201} .


Other preferred compounds include those wherein the compound exists as a single enantiomer which is: (R)-5-(3-((1-cyano-1-cyclopropylethyl)carbamoyl)-4-fluorophenyl)-2-(4-fluorophenyl)-N-methyl-6-(3,3,3-trifluoropropyl)furo[2,3-b]pyridine-3-carboxamide, or the other single enantiomer which is: (S)-5-(3-((1-cyano-1-


cyclopropylethyl)carbamoyl)-4-fluorophenyl)-2-(4-fluorophenyl)-N-methyl-6-(3,3,3-trifluoropropyl)furo[2,3-b]pyridine-3-carboxamide.

- Even more preferred is the compound from the two pure enantiomers above
 5 wherein the compound exists as the single enantiomer that exhibits a minus rotation when optical rotation is measured via standard methods in a polarimeter.


Preferred compounds of the invention, including pharmaceutically acceptable salts thereof, are selected from the group of:

10







,




,




,




,




,




,




,




,




,




,




,




,




,




,




,









,















,




,












,




,




,



,










,




,









,




,




,




,




,




,





,




,












Also preferred are compounds, including pharmaceutically acceptable salts thereof, which are selected from the group of:







,




,




,









,




,




,







,




,




,



,




,




,







,




,




,









,








,




,






and




second eluting isomer


Further preferred are compounds, including pharmaceutically acceptable salts thereof, which are selected from the group of:




,




,















,




,












,




,




,




,




,




,




,




,








,




,




,







,




,








,












,




,









and




More preferred compounds, including pharmaceutically acceptable salts thereof, are selected from the group of:







,




,




,




,











,




,











and



Other more preferred compounds, including pharmaceutically acceptable salts thereof, are selected from the group of:




, and




.

Further preferred is the compound, including pharmaceutically acceptable salts thereof, which is identified as:



5

In addition, the compound, including pharmaceutically acceptable salts thereof, which is identified as:



Pharmaceutical Compositions and Methods of Treatment

5 The compounds according to the various embodiments herein set forth demonstrate activity against HCV NS5B, and can be useful in treating HCV and HCV infection. Therefore, another aspect of the invention is a composition comprising a compound of Formula I, including pharmaceutically acceptable salts thereof, and a pharmaceutically acceptable carrier, excipient and/or diluent.

10

Another aspect of the invention is a composition further comprising an additional compound having anti-HCV activity.

15 Another aspect of the invention is a composition wherein the additional compound having anti-HCV activity is an interferon or a ribavirin. Another aspect of the invention is wherein the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, interferon lambda, and lymphoblastoid interferon tau.

20

Another aspect of the invention is a composition wherein the compound having additional anti-HCV activity is a cyclosporin. Another aspect of the invention is where the cyclosporin is cyclosporin A.

25 Another aspect of the invention is a composition wherein the additional compound having anti-HCV activity is selected from the group consisting of interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiquimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

Another aspect of the invention is a composition wherein the additional compound having anti-HCV activity is effective to inhibit the function of a target selected from HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A protein, IMPDH, and a
5 nucleoside analog for the treatment of an HCV infection.

Another aspect of the invention is a composition comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, a pharmaceutically acceptable carrier, an interferon and ribavirin.

10

Another aspect of the invention is a method of inhibiting the function of the HCV replicon comprising contacting the HCV replicon with a compound of Formula I or a pharmaceutically acceptable salt thereof.

15

Another aspect of the invention is a method of inhibiting the function of the HCV NS5B protein comprising contacting the HCV NS5B protein with a compound of Formula I or a pharmaceutically acceptable salt thereof.

Another aspect of the invention is a method of treating an HCV infection in a
20 patient comprising administering to the patient a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt thereof. In another embodiment the compound is effective to inhibit the function of the HCV replicon. In another embodiment the compound is effective to inhibit the function of the HCV NS5B protein.

25

Another aspect of the invention is a method of treating an HCV infection in a patient comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, in conjunction with (prior to, after, or concurrently) another compound having anti-HCV activity.

30

Another aspect of the invention is the method wherein the other compound having anti-HCV activity is an interferon or a ribavirin.

Another aspect of the invention is the method where the interferon is selected from interferon alpha 2B, pegylated interferon alpha, consensus interferon, interferon alpha 2A, interferon lambda, and lymphoblastoid interferon tau.

5 Another aspect of the invention is the method where the other compound having anti-HCV activity is a cyclosporin.

Another aspect of the invention is the method where the cyclosporin is cyclosporin A.

10

Another aspect of the invention is the method where the other compound having anti-HCV activity is selected from interleukin 2, interleukin 6, interleukin 12, a compound that enhances the development of a type 1 helper T cell response, interfering RNA, anti-sense RNA, Imiqimod, ribavirin, an inosine 5'-monophosphate dehydrogenase inhibitor, amantadine, and rimantadine.

15

Another aspect of the invention is the method wherein the other compound having anti-HCV activity is effective to inhibit the function of a target selected from the group consisting of HCV metalloprotease, HCV serine protease, HCV polymerase, HCV helicase, HCV NS4B protein, HCV entry, HCV assembly, HCV egress, HCV NS5A protein, IMPDH, and a nucleoside analog for the treatment of an HCV infection.

20

Another aspect of the invention is the method wherein the other compound having anti-HCV activity is effective to inhibit the function of target in the HCV life cycle other than the HCV NS5B protein.

25

“Therapeutically effective” means the amount of agent required to provide a meaningful patient benefit, e.g., inhibiting, ameliorating, or healing of acute conditions caused by HCV infection and/or inhibiting, ameliorating, or healing of the HCV infection itself, when applied to an individual so infected, as understood by practitioners in the field of hepatitis and HCV infection.

30

“Patient” means a person infected with the HCV virus and suitable for therapy as understood by practitioners in the field of hepatitis and HCV infection.

“Treatment,” “therapy,” “regimen,” “HCV infection,” and related terms are used as understood by practitioners in the field of hepatitis and HCV infection.

5 The compounds of this invention are generally given as pharmaceutical compositions comprised of a therapeutically effective amount of a compound or its pharmaceutically acceptable salt and a pharmaceutically acceptable carrier and may contain conventional excipients. Pharmaceutically acceptable carriers are those conventionally known carriers having acceptable safety profiles. Compositions
10 encompass all common solid and liquid forms including for example capsules, tablets, lozenges, and powders as well as liquid suspensions, syrups, elixers, and solutions. Compositions are made using common formulation techniques, and conventional excipients (such as binding and wetting agents) and vehicles (such as water and alcohols) are generally used for compositions. See, for example, *Remington's Pharmaceutical*
15 *Sciences*, Mack Publishing Company, Easton, PA, 17th edition, 1985.

Solid compositions are normally formulated in dosage units and compositions providing from about 1 to 1000 mg of the active ingredient per dose are preferred. Some examples of dosages are 1 mg, 10 mg, 100 mg, 250 mg, 500 mg, and 1000 mg.
20 Generally, other agents will be present in a unit range similar to agents of that class used clinically. Typically, this is 0.25-1000 mg/unit.

Liquid compositions are usually in dosage unit ranges. Generally, the liquid composition will be in a unit dosage range of 1-100 mg/mL. Some examples of dosages
25 are 1 mg/mL, 10 mg/mL, 25 mg/mL, 50 mg/mL, and 100 mg/mL. Generally, other agents will be present in a unit range similar to agents of that class used clinically. Typically, this is 1-100 mg/mL.

The invention encompasses all conventional modes of administration; oral and
30 parenteral methods are preferred. Generally, the dosing regimen will be similar to other agents used clinically. Typically, the daily dose will be 1-100 mg/kg body weight daily. Generally, more compound is required orally and less parenterally. The specific dosing regimen, however, will be determined by a physician using sound medical judgment.

The invention also encompasses methods where the compound is given in combination therapy. That is, the compound can be used in conjunction with, but separately from, other agents useful in treating hepatitis and HCV infection. In these combination methods, the compound will generally be given in a daily dose of 1-100 mg/kg body weight daily in conjunction with other agents. The other agents generally will be given in the amounts used therapeutically. The specific dosing regimen, however, will be determined by a physician using sound medical judgment.

Some examples of compounds suitable for compositions and methods are listed in Table 1.

Table 1

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
NIM811		Cyclophilin Inhibitor	Novartis
Zadaxin		Immuno-modulator	Sciclone
Suvus		Methylene blue	Bioenvision
Actilon (CPG10101)		TLR9 agonist	Coley
Batabulin (T67)	Anticancer	β -tubulin inhibitor	Tularik Inc., South San Francisco, CA
ISIS 14803	Antiviral	antisense	ISIS Pharmaceuticals Inc, Carlsbad, CA/Elan Pharmaceuticals Inc., New York, NY
Summetrel	Antiviral	antiviral	Endo Pharmaceuticals Holdings Inc., Chadds Ford, PA
GS-9132 (ACH-806)	Antiviral	HCV Inhibitor	Achillion / Gilead
Pyrazolopyrimidine compounds and salts From WO-2005047288 26 May 2005	Antiviral	HCV Inhibitors	Arrow Therapeutics Ltd.
Levovirin	Antiviral	IMPDH inhibitor	Ribapharm Inc., Costa Mesa, CA
Merimepodib (VX-497)	Antiviral	IMPDH inhibitor	Vertex Pharmaceuticals Inc., Cambridge, MA

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
XTL-6865 (XTL-002)	Antiviral	monoclonal antibody	XTL Biopharmaceuticals Ltd., Rehovot, Israel
Telaprevir (VX-950, LY-570310)	Antiviral	NS3 serine protease inhibitor	Vertex Pharmaceuticals Inc., Cambridge, MA/ Eli Lilly and Co. Inc., Indianapolis, IN
HCV-796	Antiviral	NS5B Replicase Inhibitor	Wyeth / Viropharma
NM-283	Antiviral	NS5B Replicase Inhibitor	Idenix / Novartis
GL-59728	Antiviral	NS5B Replicase Inhibitor	Gene Labs / Novartis
GL-60667	Antiviral	NS5B Replicase Inhibitor	Gene Labs / Novartis
2'C MeA	Antiviral	NS5B Replicase Inhibitor	Gilead
PSI 6130	Antiviral	NS5B Replicase Inhibitor	Roche
R1626	Antiviral	NS5B Replicase Inhibitor	Roche
2'C Methyl adenosine	Antiviral	NS5B Replicase Inhibitor	Merck
JTK-003	Antiviral	RdRp inhibitor	Japan Tobacco Inc., Tokyo, Japan
Levovirin	Antiviral	ribavirin	ICN Pharmaceuticals, Costa Mesa, CA
Ribavirin	Antiviral	ribavirin	Schering-Plough Corporation, Kenilworth, NJ
Viramidine	Antiviral	Ribavirin Prodrug	Ribapharm Inc., Costa Mesa, CA
Heptazyme	Antiviral	ribozyme	Ribozyme Pharmaceuticals Inc., Boulder, CO
BILN-2061	Antiviral	serine protease inhibitor	Boehringer Ingelheim Pharma KG, Ingelheim, Germany
SCH 503034	Antiviral	serine protease inhibitor	Schering Plough
Zadazim	Immune modulator	Immune modulator	SciClone Pharmaceuticals Inc., San Mateo, CA
Ceplene	Immunomodulator	immune modulator	Maxim Pharmaceuticals Inc., San Diego, CA

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
CellCept	Immunosuppressant	HCV IgG immuno-suppressant	F. Hoffmann-La Roche LTD, Basel, Switzerland
Civacir	Immunosuppressant	HCV IgG immuno-suppressant	Nabi Biopharmaceuticals Inc., Boca Raton, FL
Albuferon - α	Interferon	albumin IFN- α 2b	Human Genome Sciences Inc., Rockville, MD
Infergen A	Interferon	IFN alfacon-1	InterMune Pharmaceuticals Inc., Brisbane, CA
Omega IFN	Interferon	IFN- ω	Intarcia Therapeutics
IFN- β and EMZ701	Interferon	IFN- β and EMZ701	Transition Therapeutics Inc., Ontario, Canada
Rebif	Interferon	IFN- β 1a	Serono, Geneva, Switzerland
Roferon A	Interferon	IFN- α 2a	F. Hoffmann-La Roche LTD, Basel, Switzerland
Intron A	Interferon	IFN- α 2b	Schering-Plough Corporation, Kenilworth, NJ
Intron A and Zadaxin	Interferon	IFN- α 2b/ α 1-thymosin	RegeneRx Biopharma. Inc., Bethesda, MD/ SciClone Pharmaceuticals Inc, San Mateo, CA
Rebetron	Interferon	IFN- α 2b/ribavirin	Schering-Plough Corporation, Kenilworth, NJ
Actimmune	Interferon	INF- γ	InterMune Inc., Brisbane, CA
Interferon- β	Interferon	Interferon- β -1a	Serono
Multiferon	Interferon	Long lasting IFN	Viragen/ Valentis
Wellferon	Interferon	Lympho-blastoid IFN- α n1	GlaxoSmithKline plc, Uxbridge, UK
Omniferon	Interferon	natural IFN- α	Viragen Inc., Plantation, FL
Pegasys	Interferon	PEGylated IFN- α 2a	F. Hoffmann-La Roche LTD, Basel, Switzerland
Pegasys and Ceplene	Interferon	PEGylated IFN- α 2a/ immune modulator	Maxim Pharmaceuticals Inc., San Diego, CA

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
Pegasys and Ribavirin	Interferon	PEGylated IFN- α 2a/ribavirin	F. Hoffmann-La Roche LTD, Basel, Switzerland
PEG-Intron	Interferon	PEGylated IFN- α 2b	Schering-Plough Corporation, Kenilworth, NJ
PEG-Intron / Ribavirin	Interferon	PEGylated IFN- α 2b/ribavirin	Schering-Plough Corporation, Kenilworth, NJ
IP-501	Liver protection	antifibrotic	Indevus Pharmaceuticals Inc., Lexington, MA
IDN-6556	Liver protection	caspase inhibitor	Idun Pharmaceuticals Inc., San Diego, CA
ITMN-191 (R-7227)	Antiviral	serine protease inhibitor	InterMune Pharmaceuticals Inc., Brisbane, CA
GL-59728	Antiviral	NS5B Replicase Inhibitor	Genelabs
ANA-971	Antiviral	TLR-7 agonist	Anadys
Boceprevir	Antiviral	serine protease inhibitor	Schering Plough
TMS-435	Antiviral	serine protease inhibitor	Tibotec BVBA, Mechelen, Belgium
BI-201335	Antiviral	serine protease inhibitor	Boehringer Ingelheim Pharma KG, Ingelheim, Germany
MK-7009	Antiviral	serine protease inhibitor	Merck
PF-00868554	Antiviral	replicase inhibitor	Pfizer
ANA598	Antiviral	Non-Nucleoside NS5B Polymerase Inhibitor	Anadys Pharmaceuticals, Inc., San Diego, CA, USA
IDX375	Antiviral	Non-Nucleoside Replicase Inhibitor	Idenix Pharmaceuticals, Cambridge, MA, USA
BILB 1941	Antiviral	NS5B Polymerase Inhibitor	Boehringer Ingelheim Canada Ltd R&D, Laval, QC, Canada
PSI-7851	Antiviral	Nucleoside Polymerase Inhibitor	Pharmasset, Princeton, NJ, USA
PSI-7977	Antiviral	Nucleotide NS5B Polymerase Inhibitor	Pharmasset, Princeton, NJ, USA
VCH-759	Antiviral	NS5B Polymerase Inhibitor	ViroChem Pharma

<i>Brand Name</i>	<i>Physiological Class</i>	<i>Type of Inhibitor or Target</i>	<i>Source Company</i>
VCH-916	Antiviral	NS5B Polymerase Inhibitor	ViroChem Pharma
GS-9190	Antiviral	NS5B Polymerase Inhibitor	Gilead
Peg-interferon lambda	Antiviral	Interferon	ZymoGenetics/Bristol-Myers Squibb

Synthesis Methods

5 The compounds may be made by methods available in the art, as well as those described below. Some reagents and intermediates are available in the art. Other reagents and intermediates can be made by methods available in the art using commercially available materials. The variables (e.g. numbered “R” substituents) used to describe the synthesis of the compounds are intended only to illustrate how to make and
10 are not to be confused with variables used in the claims or in other sections of the specification. Abbreviations used within the schemes generally follow conventions used in the art.

Abbreviations used in the schemes generally follow conventions used in the art.

15 Chemical abbreviations used in the specification and examples are defined as follows: “NaHMDS” for sodium bis(trimethylsilyl)amide; “DMF” for N,N-dimethylformamide; “MeOH” for methanol; “NBS” for N-bromosuccinimide; “Ar” for aryl; “TFA” for trifluoroacetic acid; “LAH” for lithium aluminum hydride; “DMSO” for dimethylsulfoxide; “h” for hours; “rt” for room temperature or retention time (context
20 will dictate); “min” for minutes; “EtOAc” for ethyl acetate; “THF” for tetrahydrofuran; “EDTA” for ethylenediaminetetraacetic acid; “Et₂O” for diethyl ether; “DMAP” for 4-dimethylaminopyridine; “DCE” for 1,2-dichloroethane; “ACN” for acetonitrile; “DME” for 1,2-dimethoxyethane; “HOBt” for 1-hydroxybenzotriazole hydrate; “DIEA” for diisopropylethylamine.

25


For the section of compounds in the 0000 series all Liquid Chromatography (LC) data were recorded on a Shimadzu LC-10AS or LC-20AS liquid chromatograph using a

SPD-10AV or SPD-20A UV-Vis detector and Mass Spectrometry (MS) data were determined with a Micromass Platform for LC in electrospray mode.

- HPLC Method (i.e., compound isolation).* Compounds purified by preparative HPLC were diluted in methanol (1.2 mL) and purified using a Shimadzu LC-8A or LC-10A or Dionex APS-3000 or Waters Acquity™ automated preparative HPLC system.

Examples:

10 Preparation of Compounds 10001:



- Step 1: To a mixture of Compound 1 (5 g), 5-borono-2-methoxybenzoic acid (3.07 g) and Cs_2CO_3 (8.49 g) in dioxane (120 mL) and water (20 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (1.51 g). The mixture was flushed with nitrogen and then heated at 85 °C for 16 hours. The mixture was diluted with water and acidified with 1N HCl to pH ~ 3 and then extracted with EtOAc (2 x 150 mL). The organic layers were combined, washed with brine, dried over MgSO_4 and concentrated under vacuum. The residue was purified by titration with EtOAc to give Compound 2.

Compound 2	
MS (MHZ) ⁺ Calcd.	455.1
MS (MHZ) ⁺ Observ.	454.9
Retention Time	1.84 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	30
Final % B	100

Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um


- Step 2: To a solution of Compound 2 (300 mg), 2-amino-2-methylpropanenitrile (66.6 mg) and HATU (376 mg) in DMF (5 mL) was added iPr_2NEt (0.46 mL). The mixture was stirred at room temperature for 4 hours. The mixture was diluted with EtOAc (200 mL), washed water (50 mL), brine (50 mL), dried over $MgSO_4$ and concentrated under vacuum. The residue was purified by titration with EtOAc to give Compound 10001.

10001	
MS (MHZ) ⁺ Calcd.	521.1
MS (MHZ) ⁺ Observ.	521.2
Retention Time	1.85 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	30
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compounds 10002:

10

Compound 10002 was prepared via the same procedure towards compound 10001, using 1-aminocyclobutanecarbonitrile as the starting material.




10002

MS (MHZ) ⁺ Calcd.	533.1
MS (MHZ) ⁺ Observ.	533.1
Retention Time	1.90 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	30
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compounds 10003:

Compound 10003 was prepared via the same procedure towards compound 10001, using
 5 1-aminocyclopentanecarbonitrile hydrochloride as the starting material.



10003


MS (MHZ) ⁺ Calcd.	547.2
MS (MHZ) ⁺ Observ.	547.1
Retention Time	1.65 min

LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compounds 10004:

Compound 10004 was prepared via the same procedure towards compound 10001 from


- 5 Compound 1, using 3-borono-benzoic acid as the starting material in the Step 1.

 10004	
MS (MHZ) ⁺ Calcd.	491.1
MS (MHZ) ⁺ Observ.	491.1
Retention Time	1.85 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	30
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA

Column	PHENOMENEX-LUNA 2.0 x 30mm 3um
--------	--------------------------------


Preparation of Compounds 10005:

- Compound 10005 was prepared via the same procedure towards compound 10002 from
 5 Compound 1, using 3-borono-benzoic acid as the starting material in the Step 1.


 10005	
MS (MHZ) ⁺ Calcd.	503.1
MS (MHZ) ⁺ Observ.	503.1
Retention Time	1.89 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	30
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compounds 10006:

- 10 Compound 10006 was prepared via the same procedure towards compound 10003 from
 Compound 1, using 3-borono-benzoic acid as the starting material in the Step 1.

 10006	
MS (MHZ) ⁺ Calcd.	517.1
MS (MHZ) ⁺ Observ.	517.1
Retention Time	1.60 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compound 11001:



5


A mixture of Compound 10004 (30 mg), CF₃CH₂CH₂BF₃K (43.6 mg), cesium carbonate (59.7 mg), dicyclohexyl(2',6'-diisopropoxy-[1,1'-biphenyl]-2-yl)phosphine (11.41 mg) and diacetoxypalladium (2.74 mg) in toluene (3 mL) and water (0.3 mL) was heated at 80 °C for 16 hours. The mixture was diluted with EtOAc (20 mL), washed with water (20 mL), brine (20 mL), dried over MgSO₄ and concentrated under vacuum. The residue was purified by preparative HPLC system.

11001	
MS (MHZ) ⁺ Calcd.	553.2
MS (MHZ) ⁺ Observ.	553.2
Retention Time	1.79 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

Preparation of Compound 11002:

5


Compound 11002 was prepared via the same procedure towards compound 11001, using Compound 10005 as the starting material.

 11002	
MS (MHZ) ⁺ Calcd.	565.2
MS (MHZ) ⁺ Observ.	565.3
Retention Time	1.82 min
LC Condition	


Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

Preparation of Compound 11003:

- Compound 11003 was prepared via the same procedure towards compound 11001, using
 5 Compound 10006 as the starting material.

 11003	
MS (MHZ) ⁺ Calcd.	579.2
MS (MHZ) ⁺ Observ.	579.2
Retention Time	1.82 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220

Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Intermediate 5:

- 5 Step 1: To a mixture of Compound 1 (100 mg), (3-(methoxycarbonyl)phenyl)boronic acid (46.9 mg) and Cs_2CO_3 (170 mg) in dioxane (4 mL) and water (1 mL) was added $\text{Pd}(\text{PPh}_3)_4$ (30.1 mg). The mixture was flushed with nitrogen and then heated at 85 °C for 4 hours. The mixture was diluted with water and extracted with EtOAc (2 x 10 mL). The organic layers were combined, washed with brine (2 x 10 mL), dried over MgSO_4 and concentrated under vacuum. The residue was purified by titration with EtOAc to give Compound 3.
- 10 concentrated under vacuum. The residue was purified by titration with EtOAc to give Compound 3.

Compound 3	
MS (MHZ) ⁺ Calcd.	439.1
MS (MHZ) ⁺ Observ.	439.0
Retention Time	1.76 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220

Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

- Step 2: A mixture of Compound 3 (1g), $\text{CF}_3\text{CH}_2\text{CH}_2\text{BF}_3\text{K}$ (1.63 g), Cs_2CO_3 (2.23 g), dicyclohexyl(2',6'-diisopropoxy-[1,1'-biphenyl]-2-yl)phosphine (0.43 g) and diacetoxypalladium (0.10 g) in toluene (50 mL) and water (5.0 mL) was heated at 90°C for 16 hours. The mixture was diluted with EtOAc (250 mL), washed with water (100 mL), brine (100 mL), dried over MgSO_4 and concentrated under vacuum. The residue was purified by silica gel column (hexanes : EtOAc = 1:1 to 1:2) to give Compound 4.

Compound 4	
MS (MHZ) ⁺ Calcd.	501.1
MS (MHZ) ⁺ Observ.	501.1
Retention Time	1.88 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

- Step 3: A mixture of Compound 4 (400 mg) and NaOH (4.0 mL, 1N) in THF (30 mL) and water (15 mL) was heated at 80°C for 6 hours. The mixture was acidified by 1N HCl to pH ~ 5 and extracted with EtOAc (2 x 50mL). The organic layers were combined, washed with brine (2 x 50 mL), dried over MgSO_4 and concentrated under vacuum to give Compound 5 which was used as was.

Compound 5

MS (MHZ) ⁺ Calcd.	487.1
MS (MHZ) ⁺ Observ.	487.0
Retention Time	1.64 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compounds 11004, 11005, 11008, 11011, 11012 and 11013:

- 5 iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 5 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.


LC Condition A	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

LC Condition B	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

Cmpd #	LC Method	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
11004	A		581.2	581.2	1.88
11005	B		579.2	579.3	1.92
11008	A		539.2	539.1	1.67

Cmpd #	LC Method	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
11011	A		617.2	617.3	1.50
11012	B		571.2	571.4	1.83
11013	B		589.2	589.4	1.86

Preparation of Compounds 11006 and 11007:




5

Compounds 11006 and 11007 were two enantiomers separated from sample 11005. The absolute stereochemistry is not determined.

Preparation of Compounds 11009 and 11010:

10



Compounds 11009 and 11010 were two enantiomers separated from sample 11008. The absolute stereochemistry is not determined.

5

Preparation of Compound 12001:


Compound 12001 was prepared via the same procedure towards compound 11001, using Compound 10001 as the starting material.

10

12001	
MS (MHZ) ⁺ Calcd.	583.2
MS (MHZ) ⁺ Observ.	583.1
Retention Time	1.69 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um


Preparation of Compound 12002:

- Compound 12002 was prepared via the same procedure towards compound 11001, using
- 5 Compound 10002 as the starting material.


 12002	
MS (MHZ) ⁺ Calcd.	595.2
MS (MHZ) ⁺ Observ.	595.3
Retention Time	1.93 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

Preparation of Compound 12003:

- 10 Compound 12003 was prepared via the same procedure towards compound 11001, using
- Compound 10003 as the starting material.

 12003	
MS (M-H) ⁺ Calcd.	609.2
MS (M-H) ⁺ Observ.	609.3
Retention Time	1.95 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

Preparation of Intermediate 6:



- 5 Intermediate 6 was prepared via the same procedure towards compound 11001, using Intermediate 2 as the starting material.

Compound 6	
MS (MHZ) ⁺ Calcd.	517.1

MS (MHZ) ⁺ Observ.	517.0
Retention Time	1.63 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um


Preparation of Compounds 12004 and 12005:

- 5 iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 6 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.

LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Cmpd #	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
12004		611.2	611.2	1.88
12005		609.2	609.1	1.82

Preparation of Intermediate 9:



5

Step 1: To a mixture of Compound 1 (1 g), (4-fluoro-3-(methoxycarbonyl)phenyl)boronic acid (0.62 g) and Cs₂CO₃ (1.70 g) in dioxane (40 mL) and water (4 mL) was added Pd(PPh₃)₄ (0.30 g). The mixture was flushed with nitrogen and then heated at 85°C for 16 hours. The mixture was diluted with water and then extracted with EtOAc (2 x 100 mL). The organic layers were combined, washed with brine (100 mL), dried over MgSO₄ and concentrated under vacuum. The residue was purified by titration with EtOAc to give Compound 7.

Compound 7	
MS (MHZ) ⁺ Calcd.	457.1
MS (MHZ) ⁺ Observ.	457.0
Retention Time	1.76 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

- Step 2: A mixture of Compound 7 (270 mg), potassium trifluoro(3,3,3-trifluoropropyl)borate (422 mg), cesium carbonate (578 mg), dicyclohexyl(2',6'-diisopropoxy-[1,1'-biphenyl]-2-yl)phosphine (110 mg) and
- 5 diacetoxypalladium (26.5 mg) in toluene (10 mL) and water (1.0 mL) was heated at 80°C for 16 hours. The mixture was diluted with EtOAc (20 mL), washed with water (20 mL), brine (20 mL), dried over MgSO₄ and concentrated under vacuum to give Compound 8 which was used as was.

Compound 8	
MS (MHZ) ⁺ Calcd.	519.1
MS (MHZ) ⁺ Observ.	519.1
Retention Time	1.94 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100

Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

- Step 3: To a suspension of Compound 8 (50 mg) in acetone (3 mL) and water (1 mL) was added NaOH (1.93 mL, 1N). The mixture was heated at 80°C for 4 hours. The mixture was acidified by 1N HCl to pH ~ 3. The precipitate was
- 5 collected by filtration to give Compound 9 which was used as was.

Compound 9	
MS (MHZ) ⁺ Calcd.	505.1
MS (MHZ) ⁺ Observ.	505.0
Retention Time	1.64 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um




Preparation of Compounds 11004, 11005, 11008, 11011, 11012 and 11013:

- 10 iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 9 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.

LC Condition A	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um


LC Condition B	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

TABLE 1

Cmpd #	LC Method	Structure	Coupling / Base Agent Used	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
13001	B		HATU / iPr ₂ NEt	597.2	597.3	1.94
13002	B		HATU / iPr ₂ NEt	599.2	599.3	1.96
13003	A		HATU / iPr ₂ NEt	557.2	557.0	1.57

12276 WO 2015

Cmpd #	LC Method	Structure	Coupling / Base Agent Used	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
13006	B		HATU / iPr ₂ NEt	597.2	597.4	1.91


Preparation of Compounds 13004 and 13005:

5

Compounds 13004 and 13005 were two enantiomers separated from sample 13003. The absolute stereochemistry is not determined.

Preparation of Compounds 13007 and 13008:

10



Compounds 13007 and 13008 were two enantiomers separated from sample 13006. The absolute stereochemistry is not determined.

15

Chiral separation and purity analysis of Compounds 13007 and 13008:

Chiralcel OD-H preparative column, 30 x 250mm, 5 μ m

Mobile Phase: 10% MeOH (0.1%DEA) in CO₂, 150bar

20 Temp: 35°C

Flow rate: 70.0 mL/min. for 22 min.

UV monitored @ 316nm

Injection: 0.5ml of ~40mg/mL solution in 1:1 MeOH:CHCl₃

Retention Time: 15.25 minutes (Compound 13007) and 17.68 minutes

25 (Compound 13008)

Two analytical LC/MS injections were used to determine the final purity. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0%B, 0-100% B over 3 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0%B, 0-100% B over 3 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm.

Compound 13308 was further recrystallized from a mixed solution of EtOH and H₂O to give a white solid.

Characterization of Compound 13308:

a. Optical rotation

Result: $[\alpha]_d^{20^\circ\text{C}} = -33.48^\circ$ (3.435mg/ml, CHCl₃)

b. Mass spectrum:

Mass Range: m/z 120-1200
Ionization and Mode: Electrospray ionization, positive ion mode
Result: $\text{ES}^+ = 597.3$

c. Elemental Analysis: %composition difference (Δ) = experimental - theoretical (Acceptance criterion: $\Delta \leq \pm 0.4$)

Result: C₃₁H₂₅F₅N₄O₃•0.06 H₂O: $\Delta\text{C} = -0.09\%$; $\Delta\text{H} = -0.19\%$;
 $\Delta\text{N} = +0.06\%$

d. Proton Spectrum :

Experimental: 7.3 mg sample dissolved in 600 μ l DMSO- d_6 , 4 scans, 32K points at room temperature. The ^1H chemical shifts are referenced to TMS at 0.0 ppm.

5 ^1H NMR (500MHz, DMSO- d_6) δ 9.06 (s, 1H), 8.54 (q, $J=4.4$ Hz, 1H), 8.07 (dd, $J=9.0, 5.5$ Hz, 2H), 7.97 (s, 1H), 7.69 (m, 1H), 7.66 (m, 1H), 7.48 (t, $J=8.5$ Hz, 1H), 7.43 (dd, $J=8.8, 7.3$ Hz, 2H), 3.03 (dd, $J=9.1, 6.6$ Hz, 2H), 2.83 (d, $J=4.7$ Hz, 3H), 2.79 (m, 2H), 1.67 (s, 3H), 1.50 (m, 1H), 0.64 (m, 3H), 0.53 (m, 1H)

10 e. Carbon Spectrum:

Experimental: 7.3 mg sample dissolved in 600 μ l DMSO- d_6 . Carbon resonance frequency is 125.73 MHz. 1024 scans, 32K points at room temperature. The ^{13}C chemical shifts are referenced to TMS at 0.0 ppm.

15 ^{13}C NMR (125.73MHz, DMSO- d_6) δ 163.4, 162.4, 163.0 (d, $J=248.9$ Hz), 158.8, 158.7 (d, $J=250.7$ Hz), 152.0, 151.4, 135.0 (d, $J=3.6$ Hz), 133.6 (d, $J=8.2$ Hz), 131.9, 131.8, 130.8, 129.9 (d, $J=9.1$ Hz), 127.5 (q, $J=277.0$ Hz), 125.2 (d, $J=2.7$ Hz), 123.9 (d, $J=15.4$ Hz), 118.8, 117.5, 116.4 (d, $J=22.7$ Hz), 116.1 (d, $J=22.7$ Hz), 112.6, 52.2, 31.3 (q, $J=27.3$ Hz), 27.1, 26.2, 23.8, 18.6, 2.9, 1.7.

20


f. Fluorine Spectrum

Experimental: 7.3 mg sample dissolved in 600 μ l DMSO- d_6 , fluorine resonance frequency is 470.45 MHz, 16 scans, 64K points at room temperature. The ^{19}F chemical shifts are referenced to CFCl_3 at 0.0 ppm.

25

^{19}F NMR (470.45 MHz, DMSO- d_6) δ -64.74, -109.85, -115.68.

30

Preparation of Intermediate 10:



Intermediate 10 was prepared via the same procedure towards Intermediate 9, using
 5 methyl 3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate as the starting material in Step 1.

Preparation of Compounds 14001 and 14002:


- 10 $i\text{Pr}_2\text{NEt}$ or Et_3N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 10 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.

LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

15

Cmpd #	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
14001		571.2	571.1	1.89
14002		597.2	597.1	1.98

Preparation of Intermediate 14:



5

Step 1: To a mixture of Compound 1 (500 mg), 3-borono-4-fluorobenzoic acid (264 mg) and Cs₂CO₃ (849 mg) in DMF (15 mL) and water (1.5 mL) was added Pd(PPh₃)₄ (151 mg). The mixture was flushed with nitrogen and then heated at 85°C for 6 hours. The mixture was diluted with water and then extracted with EtOAc (2 x 50 mL). The organic layers were combined, washed with brine (50 mL) and concentrated under vacuum. The residue was purified by titration with EtOAc to give Compound 11.

Compound 11	
MS (MHZ) ⁺ Calcd.	443.1
MS (MHZ) ⁺ Observ.	442.9
Retention Time	1.67 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Step 2: A mixture of Compound 2 (70 mg), iodomethane (0.049 mL) and Cs₂CO₃ (103 mg) in MeOH (3 mL) in a sealed tube was heated at 80°C for 6 hours. The mixture was diluted with MeOH. The solid was removed by filtration. The filtrate was concentrated

5 to give a residue which was purified by preparative HPLC system.

Compound 12	
MS (MHZ) ⁺ Calcd.	457.1
MS (MHZ) ⁺ Observ.	456.9
Retention Time	1.91 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220

Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Step 3: A mixture of Compound 12 (25 mg), potassium trifluoro(3,3,3-trifluoropropyl)borate (39.1 mg), cesium carbonate (53.5 mg), dicyclohexyl(2',6'-diisopropoxy-[1,1'-biphenyl]-2-yl)phosphine (10.21 mg) and diacetoxypalladium (2.46 mg) in toluene (2 mL) / water (0.2 mL) was degassed and heated at 80°C for 16 hours. The mixture was diluted with EtOAc (10 mL), then washed with water (10 mL) and brine (10 mL). The organic layer was separated and concentrated under vacuum. The residue was purified by titration with EtOAc to Compound 13.

Compound 13	
MS (MHZ) ⁺ Calcd.	519.1
MS (MHZ) ⁺ Observ.	519.0
Retention Time	2.09 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

10

Step 4: To a suspension of Compound 13 (15 mg) in THF (3 mL) and water (0.3 mL) was added NaOH (0.5 mL, 1N). The mixture was heated at 80°C for 4 hours. The mixture was acidified with 1N HCl to pH ~ 3. All solvents were removed under vacuum to give Compound 14 which was used as was.



15

Compound 14	
MS (MHZ) ⁺ Calcd.	505.1
MS (MHZ) ⁺ Observ.	505.0
Retention Time	1.98 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um


Preparation of Compounds 15001 and 15002:

- 5 iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 14 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.

LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Cmpd #	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
15001		597.2	597.1	2.00
15002		571.2	571.1	1.71

5 Preparation of Intermediate 15:




Intermediate 15 was prepared via the same procedure towards Intermediate 9, using methyl 2,3-dichloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate as the starting material in Step 1.

Compound 15	
MS (MHZ) ⁺ Calcd.	555.0
MS (MHZ) ⁺ Observ.	555.0

Retention Time	1.97 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	60
Final % B	90
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compound 16001:



5


To a solution of Compound 15 (8 mg), 2-amino-2-methylpropanenitrile (2.424 mg, 0.029 mmol) and HATU (8.22 mg, 0.022 mmol) in DMF (1 mL) was added $i\text{Pr}_2\text{NEt}$ (10.06 μl). The mixture was stirred at room temperature for 4 hours. The product was isolated by preparative HPLC system.

10

Compound 16001	
MS (MHZ) ⁺ Calcd.	621.1
MS (MHZ) ⁺ Observ.	621.1
Retention Time	1.59 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA

Start % B	60
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Intermediate 16:




5

Intermediate 16 was prepared via the same procedure towards Intermediate 9, using (3,4-difluoro-5-(methoxycarbonyl)phenyl)boronic acid as the starting material in Step 1.


Compound 16	
MS (MHZ) ⁺ Calcd.	523.1
MS (MHZ) ⁺ Observ.	523.4
Retention Time	1.41 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate

Column	Phenomenex LUNA C18, 30x2, 3u
--------	-------------------------------

Preparation of Compound 17001:

5

Compound 18001	
MS (MHZ) ⁺ Calcd.	615.2
MS (MHZ) ⁺ Observ.	615.1
Retention Time	1.84 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compound 18001:


10

To a solution of Compound 17 (15 mg), 2-amino-2-cyclopropylpropanenitrile (9.52 mg) and HATU (16.44 mg) in DMF (1 mL) was added $i\text{Pr}_2\text{NEt}$ (0.02 mL). The mixture was stirred at room temperature for 4 hours. The product was isolated by preparative HPLC system.

5

Compound 18001	
MS (MHZ) ⁺ Calcd.	613.2
MS (MHZ) ⁺ Observ.	613.2
Retention Time	2.00 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Intermediate 19:



10

Step 1: A mixture of Compound 3 (800 mg) and NaOH (9.12 mL, 1N) in THF (30 mL) and water (15 mL) was heated at 80°C for 6 hours. The mixture was acidified by 1N HCl to pH ~ 5 and extracted with EtOAc (2 x 50 mL). The organic layers were combined, washed with brine (50 mL), dried over MgSO_4

15 and concentrated under vacuum to give Compound 18.

Compound 18	
MS (MHZ) ⁺ Calcd.	425.1
MS (MHZ) ⁺ Observ.	425.0
Retention Time	1.49 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

- Step 2: A mixture of Compound 18 (175 mg), chloro[2-(dicyclohexylphosphino)-3,6-dimethoxy-2',4',6'-tri-*i*-propyl-1,1'-biphenyl][2-(2-aminoethyl)phenyl]palladium (II) (28.2 mg) and sodium 2-methylbutan-2-olate (194 mg) in dioxane (10 mL) was heated at 90°C for 30 minutes. The mixture was diluted with EtOAc (20 mL), washed with water (20 mL), brine (20 mL), dried over MgSO₄ and concentrated under vacuum. The residue was purified by preparative HPLC system.

Compound 19	
MS (MHZ) ⁺ Calcd.	488.1
MS (MHZ) ⁺ Observ.	488.0
Retention Time	1.91 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	30
Final % B	100

Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compounds 20001, 20002, 20005 and 20006:




- 5 iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 19 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.


LC Condition A	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um


LC Condition B	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220

Solvent Pair	ACN: Water: Ammonium Acetate
Column	Phenomenex LUNA C18, 30x2, 3u

12278WOPCT

#	LC Method	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
20001	B		554.2	554.3	1.85
20002	A		540.2	540.1	1.59
20005	A		602.2	602.2	1.77

Cmpd #	LC Method	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
20006	A		580.2	580.1	1.71


Preparation of Compounds 20003 and 20004:

5

Compounds 20003 and 20004 were two enantiomers separated from sample 20002. The absolute stereochemistry is not determined.


Preparation of Compounds 20007 and 20008:

10



Compounds 20007 and 20008 were two enantiomers separated from sample 20006. The absolute stereochemistry is not determined.


15

Preparation of Intermediate 20:

20 Intermediate 20 was prepared via the same procedure towards Intermediate 19 from Compound 18, using Compound 2 as the starting material.

Compound 20	
MS (MHZ) ⁺ Calcd.	518.1
MS (MHZ) ⁺ Observ.	518.0
Retention Time	1.89 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	30
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compound 21001:




- 5 To a solution of Compound 20 (20 mg), 2-amino-2-methylpropanenitrile hydrochloride (9.32 mg, 0.077 mmol) and HATU (22.05 mg) in DMF (2 mL) was added iPr₂NEt (0.027 mL). The mixture was stirred at room temperature for 4 hours. The product was isolated by preparative HPLC system.

Compound 21001	
MS (MHZ) ⁺ Calcd.	584.2
MS (MHZ) ⁺ Observ.	584.3
Retention Time	1.87 min
LC Condition	

Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

Preparation of Intermediate 21:



5

Intermediate 21 was prepared via the same procedure towards Intermediate 19, using Compound 7 as the starting material at step 1.


Compound 21	
MS (MHZ) ⁺ Calcd.	506.1
MS (MHZ) ⁺ Observ.	506.0
Retention Time	1.60 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220

Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compounds 22001 and 22004:


- 5 iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 21 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.

LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

Cmpd #	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
22001		558.2	558.3	1.77

Cmpd #	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
22004		598.2	598.4	1.86

Preparation of Compounds 22002 and 22003:




5

Compounds 22002 and 22003 were two enantiomers separated from sample 22001. The absolute stereochemistry is not determined.

Preparation of Compounds 22005 and 22006:


10



Compounds 22005 and 22006 were two enantiomers separated from sample 22004. The absolute stereochemistry is not determined.

15

Preparation of Intermediate 23:



Step 1: Intermediate 22 was prepared via the same procedure towards Intermediate 7, using methyl 3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate as the starting material.

Compound 22	
MS (MHZ) ⁺ Calcd.	457.1
MS (MHZ) ⁺ Observ.	456.9
Retention Time	2.01 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um



Step 2: A mixture of Compound 22 (50 mg), 2,2,2-trifluoroethanamine (54.2 mg), chloro[2-(dicyclohexylphosphino)-3,6-dimethoxy-2',4',6'-tri-*i*-propyl-1,1'-biphenyl][2-(2-aminoethyl)phenyl]palladium (II) (17.49 mg) and sodium 2-methylbutan-2-olate (48.2 mg) in dioxane (3 mL) was heated at 65°C for 20 minutes. The mixture was diluted with EtOAc (20 mL), washed with 1N HCl (20 mL) and brine (20 mL). The organic layer was dried over MgSO₄ and concentrated under vacuum. The residue was purified by preparative HPLC system to Compound 23.

Compound 23	
MS (MHZ) ⁺ Calcd.	506.1
MS (MHZ) ⁺ Observ.	506.0
Retention Time	1.60 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um


Preparation of Compounds 23001 and 23002:

- 5 iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 23 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.

LC Condition	
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um
Column	Phenomenex LUNA C18, 30x2, 3u

Cmpd #	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
23001		598.2	598.1	1.73
23002		572.2	572.1	1.60

Preparation of Intermediate 26:



Step 1: Intermediate 24 was prepared via the same procedure towards Intermediate 7, using methyl 2,3-dichloro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate as the starting material.

10

Compound 24	
MS (MHZ) ⁺ Calcd.	507.0

MS (MHZ) ⁺ Observ.	507.0
Retention Time	2.07 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Step 2: To a suspension of Compound 24 (40 mg) in THF (3 mL) and water (1 mL) was added NaOH (1 mL, 1 N). The mixture was heated at 80°C for 4 hours. The mixture was acidified with 1N HCl to pH ~ 3. The solid was

5 collected by filtration to give Compound 25.


Compound 25	
MS (MHZ) ⁺ Calcd.	493.0
MS (MHZ) ⁺ Observ.	493.2
Retention Time	1.38 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

- Step 3: A mixture of Compound 25 (37 mg), 2,2,2-trifluoroethanamine (37.1 mg), chloro[2-(dicyclohexylphosphino)-3,6-dimethoxy-2',4',6'-tri-*i*-propyl-1,1'-biphenyl][2-(2-aminoethyl)phenyl]palladium (II) (11.97 mg) and sodium 2-methylbutan-2-olate (33.0 mg) in dioxane (5 mL) was heated at 80°C for 20 minutes. The mixture was diluted with EtOAc (20 mL), washed with 1N HCl (20 mL) and brine (20 mL). The organic layer was dried over MgSO₄ and concentrated under vacuum. The residue was purified by preparative HPLC system to Compound 26.

10


Compound 26	
MS (MHZ) ⁺ Calcd.	619.1
MS (MHZ) ⁺ Observ.	619.1
Retention Time	1.69 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

15

Preparation of Compound 24001:

- 5 To a solution of Compound 26 (15 mg), 2-amino-2-methylpropanenitrile (4.08 mg) and HATU (13.82 mg) in DMF (2 mL) was added $i\text{Pr}_2\text{NEt}$ (0.017 mL). The mixture was stirred at room temperature for 4 hours. The product was isolated by preparative HPLC system.

Compound 24001	
MS (MHZ) ⁺ Calcd.	685.2
MS (MHZ) ⁺ Observ.	685.3
Retention Time	1.89 min
LC Condition	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

Preparation of Intermediate 27:

- 5 A mixture of Compound 2 (460 mg), 2,2,3,3,3-pentafluoropropan-1-amine (781 mg), chloro[2-(dicyclohexylphosphino)-3,6-dimethoxy-2',4',6'-tri-*i*-propyl-1,1'-biphenyl][2-(2-aminoethyl)phenyl]palladium (II) (84 mg) and sodium 2-methylbutan-2-olate (577 mg) in dioxane (25 mL) was heated at 85°C for 30 minutes. The mixture was diluted with EtOAc (100 mL), washed
- 10 with water (100 mL) and brine (100 mL), dried over MgSO₄ and concentrated under vacuum to give Compound 27.

Compound 27	
MS (MHZ) ⁺ Calcd.	538.1
MS (MHZ) ⁺ Observ.	538.0
Retention Time	1.79 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um




Preparation of Compounds 30001-30003:


iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 27 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.

LC Condition A	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles

LC Condition B	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Phenomenex LUNA C18, 30x2, 3u

12278WOPCT

Cmpd #	LC Method	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
30001	B		604.2	604.3	1.84
30002	A		632.2	632.4	3.29
30003	A		630.2	630.4	3.24

Preparation of Intermediate 28:

5

Intermediate 28 was prepared via the same procedure towards Intermediate 20 from Compound 2, using 2,2,3,3,3-pentafluoropropan-1-amine as the starting material.

Compound 28	
MS (MHZ) ⁺ Calcd.	568.1
MS (MHZ) ⁺ Observ.	568.1
Retention Time	1.74 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

10 **Preparation of Compounds 31001 and 31002:**

iPr₂NEt or Et₃N (2 eq.) and HATU or HCTU or DEBPT (1.3 eq.) were added into a solution of Compound 28 (1 eq.) and amine (1.3 eq.) in DMF or THF. The reaction was



stirred at room temperature or 85°C for 30 minutes to 72 hours. The desired product was isolated by preparative HPLC system.

LC Condition A	
Solvent A	5 % ACN: 95% Water : 10mM Ammonium Actetate
Solvent B	95 % ACN: 5% Water : 10mM Ammonium Actetate
Start % B	0
Final % B	100
Gradient Time	0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	ACN: Water: Ammonium Actetate
Column	Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles


LC Condition B	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

5

10

Cmpd #	LC Method	Structure	MS (MHZ) ⁺ Calcd.	MS (MHZ) ⁺ Observ.	Retention Time (min)
31001	B		662.2	662.2	1.97
31002	A		660.2	660.3	3.39

Preparation of Intermediate 29:




5

Intermediate 29 was prepared via the same procedure towards Intermediate 19 from Compound 18, using 2-aminoethanol as the starting material.

Compound 29	
MS (MHZ) ⁺ Calcd.	450.1
MS (MHZ) ⁺ Observ.	450.1
Retention Time	1.88 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA


Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compound 40001:

- 5 To a solution of Compound 29 (10 mg), 2-amino-2-methylpropanenitrile (3.74 mg) and HATU (12.69 mg) in DMF (1.5 mL) was added $i\text{Pr}_2\text{NEt}$ (0.016 mL). The mixture was stirred at room temperature for 4 hours. The product was isolated by preparative HPLC system.

Compound 40001	
MS (MHZ) ⁺ Calcd.	516.2
MS (MHZ) ⁺ Observ.	516.1
Retention Time	1.87 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	0
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min

Wavelength	220
Solvent Pair	Water - Methanol- TFA
Column	PHENOMENEX-LUNA 2.0 x 30mm 3um

Preparation of Compound 50001:

5

A mixture of Compound 10004 (80 mg), potassium vinyltrifluoroborate (76 mg), cesium carbonate (159 mg), dicyclohexyl(2',6'-diisopropoxy-[1,1'-biphenyl]-2-yl)phosphine (30.4 mg) and diacetoxypalladium (7.32 mg) in toluene (8 mL) and water (0.8 mL) was heated at 80°C for 16 hours. The mixture was diluted with EtOAc (20 mL), washed with water (20 mL) and brine (20 mL), dried over MgSO₄ and concentrated under vacuum. The residue purified by preparative HPLC.


Compound 50001	
MS (MHZ) ⁺ Calcd.	483.2
MS (MHZ) ⁺ Observ.	483.2
Retention Time	2.49 min
LC Condition	
Solvent A	90% Water -10% Methanol-0.1% TFA
Solvent B	10% Water -90% Methanol-0.1% TFA
Start % B	50
Final % B	100
Gradient Time	2 min
Flow Rate	1 mL/min
Wavelength	220
Solvent Pair	Water - Methanol- TFA

Column	PHENOMENEX-LUNA 2.0 x 30mm 3um
--------	--------------------------------

General procedure for the preparation of compounds K1001-K1003


*i*Pr₂NEt (3 eq.) and HATU (1.5 eq.) were added into a solution of 2-fluoro-5-(2-(4-fluorophenyl)-3-(methylcarbamoyl)-6-(3,3,3-trifluoropropyl)furo[2,3-b]pyridin-5-yl)benzoic acid (1 eq.) and amine (2 eq.) in DMF. The reaction mixture was stirred at room temperature for 1 hour. The entire reaction mixture was purified via preparative HPLC to obtain the desired product.

10 Compound K1001




Amine nucleophile = 1-aminocyclopropanecarbonitrile hydrochloride. LCMS: Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 2.90 min, (M+H)⁺: 569. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 3.91 min, (M+H)⁺: 569.

25 Compound K1002



- Amine nucleophile = 1-aminocyclobutanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 2.11 min, (M+H)⁺: 583. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 4.01 min, (M-H)⁺: 581. ¹H NMR (500MHz, DMSO-d₆) δ 9.43 (s, 1H), 8.54 - 8.48 (m, 1H), 8.06 (dd, J=8.9, 5.5 Hz, 2H), 7.97 (s, 1H), 7.74 - 7.65 (m, 2H), 7.49 (t, J=9.3 Hz, 1H), 7.41 (t, J=8.9 Hz, 2H), 3.05 - 2.97 (m, 2H), 2.85 - 2.64 (m, 7H), 2.47 (d, J=11.3 Hz, 2H), 2.11 - 2.00 (m, 2H).

Compound K1003



20

Amine nucleophile = 2-amino-2-methylpropanenitrile hydrochloride. Injection 1 conditions: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles;

Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.11 mL/min.

Retention time: 4.06 min, (M+H)⁺: 571. Injection 2 conditions: Column: Waters Acquity


- 5 UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 0.05% TFA; Mobile Phase B: 95:5 acetonitrile:water with 0.1% TFA; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.11 mL/min. Retention time: 3.08 min, (M+H)⁺: 571. ¹H NMR (500MHz, DMSO-d₆) δ 8.98 (s, 1H), 8.54 - 8.45 (m, 1H), 8.06 (dd, J=8.4, 5.6 Hz, 2H), 7.96 (s, 1H), 7.69 - 7.63 (m, 2H), 7.47 (s, 1H), 7.41 (t, J=8.7 Hz, 2H), 3.06 - 2.98 (m, 2H), 2.85 - 2.72 (m, 5H), 1.73 - 1.65 (m, 6H).
- 10

General procedure for the preparation of compounds K2001-K2005:

- 15 *i*Pr₂NEt (8 eq.) and HATU (1.5 eq.) were added into a solution of 2-fluoro-5-(2-(4-fluorophenyl)-3-(methylcarbamoyl)-6-((2,2,2-trifluoroethyl)amino)furo[2,3-b]pyridin-5-yl)benzoic acid (1 eq.) and amine (1 eq.) in DMF. The reaction mixture was stirred at room temperature for 1 hour. The entire reaction mixture was purified via preparative HPLC to obtain the desired product.

20


Compound K2001



- 25 Amine nucleophile = 2-amino-2-methylpropanenitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min


hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 2.92 min, (M+H)⁺: 572. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.86 min, (M+H)⁺: 572.

Compound K2002



Amine nucleophile = 1-aminocyclopropanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 2.86 min, (M+H)⁺: 570. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.81 min, (M+H)⁺: 570.

Compound K2003



Amine nucleophile = 1-aminocyclobutanecarbonitrile hydrochloride. Injection 1


conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

- 5 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 2.96 min, (M+H)⁺: 584. Injection 2

conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

- 10 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.90 min, (M+H)⁺: 584.

15 **Compound K2004**



Amine nucleophile = 1-aminocyclopentanecarbonitrile hydrochloride. Injection 1


- 20 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5

acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min

hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 3.05 min, (M+H)⁺: 598. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.98 min, (M+H)⁺: 598.

Compound K2005




Amine nucleophile = 2-amino-2,3-dimethylbutanenitrile. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 3.10 min, (M+H)⁺: 600. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 4.02 min, (M+H)⁺: 600.

General procedure for the preparation of compound K3001:

*i*Pr₂NEt (3 eq.) and HATU (1.5 eq.) were added into a solution of 3-(2-(4-fluorophenyl)-3-(methylcarbamoyl)-6-(3,3,3-trifluoropropyl)furo[2,3-*b*]pyridin-5-yl)benzoic acid (1 eq.) and amine (2 eq.) in DMF. The reaction mixture was stirred at room temperature for 1 hour. The entire reaction mixture was purified via preparative HPLC to obtain the desired product.

Compound K3001




Amine nucleophile = 1-aminocyclopropanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 2.87 min, (M+H)⁺: 551. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 3.93 min, (M+H)⁺: 551. ¹H NMR (500MHz, DMSO-*d*₆) δ 9.43 (s, 1H), 8.54 - 8.44 (m, 1H), 8.10 - 8.00 (m, 2H), 7.97 (s, 2H), 7.90 (s, 1H), 7.74 - 7.60 (m, 2H), 7.42 (s, 2H), 3.05 - 2.96 (m, 2H), 2.84 - 2.69 (m, 5H), 1.63 - 1.52 (m, 2H), 1.34 - 1.26 (m, 2H).

General procedure for the preparation of compounds K4001-K4003:

*i*Pr₂NEt (8 eq.) and HATU (1.5 eq.) were added into a solution of 3-(2-(4-fluorophenyl)-3-(methylcarbamoyl)-6-((2,2,2-trifluoroethyl)amino)furo[2,3-*b*]pyridin-5-yl)benzoic acid

(1 eq.) and amine (1 eq.) in DMF. The reaction mixture was stirred at room temperature for 1 hour. The entire reaction mixture was purified via preparative HPLC to obtain the desired product.

5 Compound K4001




Amine nucleophile = 1-aminocyclopentanecarbonitrile hydrochloride. Injection 1

- 10 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 2.94 min, (M+H)⁺: 580. Injection 2
- 15 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.98 min, (M+H)⁺: 580.

20

Compound K4002



1-aminocyclobutanecarbonitrile hydrochloride was the amine used. Injection 1


conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

5 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5
acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min
hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1
mL/min; Detection: UV at 220 nm. Retention time: 2.86 min, (M+H)⁺: 566. Injection 2

conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

10 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5
methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min
hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5
mL/min; Detection: UV at 220 nm. Retention time: 3.91 min, (M+H)⁺: 566.

15 **Compound K4003**



Amine nucleophile = 1-aminocyclopropanecarbonitrile hydrochloride. Injection 1


20 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5

acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min

hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 2.99 min, (M+H)⁺: 552. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.78 min, (M+H)⁺: 552.

Compound K4004




Amine nucleophile = 2-amino-2,3-dimethylbutanenitrile. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 3.18 min, (M+H)⁺: 582. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.97 min, (M+H)⁺: 582.

25 General procedure for the preparation of compounds K5001-K5004:

*i*Pr₂NEt (8 eq.) and HATU (1.5 eq.) were added into a solution of 5-(2-(4-fluorophenyl)-3-(methylcarbamoyl)-6-((2,2,2-trifluoroethyl)amino)furo[2,3-b]pyridin-5-yl)-2-methoxybenzoic acid (1 eq.) and amine (1 eq.) in DMF. The reaction mixture was stirred at room temperature for 1 hour. The entire reaction mixture was purified via preparative HPLC to obtain the desired product.

Compound K5001




10

Amine nucleophile = 1-aminocyclobutanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 3.15 min, (M+H)⁺: 596. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.88 min, (M+H)⁺: 596.

20

Compound K5002




Amine nucleophile = 1-aminocyclopropanecarbonitrile hydrochloride. Injection 1

conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

- 5 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 2.91 min, (M+H)⁺: 582. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:
- 10 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.81 min, (M+H)⁺: 582.

15 **Compound K5003**




Amine nucleophile = 1-aminocyclopentanecarbonitrile hydrochloride. Injection 1

- 20 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min

hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 3.01 min, (M+H)⁺: 610. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 3.98 min, (M+H)⁺: 610.

Compound K5004




Amine nucleophile = 2-amino-2,3-dimethylbutanenitrile. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 3.20 min, (M+H)⁺: 612. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 4.03 min, (M+H)⁺: 612.

25 General procedure for the preparation of compounds K6001:

*i*Pr₂NEt (8 eq.) and HATU (1.5 eq.) were added into a solution of 5-(2-(4-fluorophenyl)-3-(methylcarbamoyl)-6-(3,3,3-trifluoropropyl)furo[2,3-*b*]pyridin-5-yl)-2-methoxybenzoic acid (1 eq.) and amine (1 eq.) in DMF. The reaction mixture was stirred at room temperature for 1 hour. The entire reaction mixture was purified via preparative HPLC to obtain the desired product.

Compound K6001



10

Amine nucleophile = 1-aminocyclopropanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 2.95 min, (M+H)⁺: 581. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 3.93 min, (M+H)⁺: 581.

20

General procedure for the preparation of compounds K7001-K7005:


*i*Pr₂NEt (8 eq.) and HATU (1.5 eq.) were added into a solution of 5-(2-(4-fluorophenyl)-3-(methylcarbamoyl)-6-(3,3,3-trifluoropropyl)furo[2,3-*b*]pyridin-5-yl)-2-methoxynicotinic acid (1 eq.) and amine (1 eq.) in DMF. The reaction mixture was

25

stirred at room temperature for 1 hour. The entire reaction mixture was purified via preparative HPLC to obtain the desired product.

Compound K7001

5




Amine nucleophile = 2-amino-2-methylpropanenitrile hydrochloride. Injection 1

conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

- 10 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 3.17 min, (M+H)⁺: 584. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 4.09 min, (M+H)⁺: 584.

20 Compound K7002




Amine nucleophile = 1-aminocyclobutanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5

- 5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 3.20 min, (M+H)⁺: 596. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 4.11 min, (M+H)⁺: 596.

Compound K7003

15




Amine nucleophile = 1-aminocyclopropanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5

- 20 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 2.94 min, (M+H)⁺: 582. Injection 2 conditions:Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; MobilePhase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes,
- 25

then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 3.99 min, (M+H)⁺: 582.


Compound K7004

5



Amine nucleophile = 1-aminocyclopentanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 3.18 min, (M+H)⁺: 610. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 4.16 min, (M+H)⁺: 610.

20 Compound K7005




Amine nucleophile = 2-amino-2,3-dimethylbutanenitrile. Injection 1 conditions: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.11 mL/min. Retention time: 3.33 min, (M+H)⁺: 612. Injection 2 conditions: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 0.05% TFA; Mobile Phase B: 95:5 acetonitrile:water with 0.1% TFA; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.11 mL/min. Retention time: 4.31 min, (M+H)⁺: 612.

General procedure for the preparation of compounds K8001-K8005:

*i*Pr₂NEt (8 eq.) and HATU (1.5 eq.) were added into a solution of 5-(2-(4-fluorophenyl)-3-(methoxycarbonyl)-6-((2,2,2-trifluoroethyl)amino)furo[2,3-b]pyridin-5-yl)-2-methoxynicotinic acid (1 eq.) and amine (1 eq.) in DMF. The reaction mixture was stirred at room temperature for 1 hour. The entire reaction mixture was purified via preparative HPLC to obtain the desired product.


Compound K8001



Amine nucleophile = 2-amino-2-methylpropanenitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

- 5 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 2.92 min, (M+H)⁺: 585. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; MobilePhase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 3.88 min, (M+H)⁺: 585.

15 Compound K8002




- Amine nucleophile = 1-aminocyclobutanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min

hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 3.05 min, (M+H)⁺: 597. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 3.87 min, (M+H)⁺: 597.

Compound K8003


10



Amine nucleophile = 1-aminocyclopropanecarbonitrile hydrochloride. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 2.82 min, (M+H)⁺: 583. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 3.83 min, (M+H)⁺: 583.

Compound K8004

25



Amine nucleophile = 1-aminocyclopentanecarbonitrile hydrochloride. Injection 1

conditions: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles;


- 5 Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.11 mL/min.

Retention time: 3.14 min, $(M+H)^+$: 611. Injection 2 conditions: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 0.05% TFA; Mobile Phase B: 95:5 acetonitrile:water with 0.1% TFA; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.11 mL/min. Retention time: 4.15 min, $(M+H)^+$: 611.

10

Compound K8005

15




Amine nucleophile = 2-amino-2,3-dimethylbutanenitrile was. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A:

- 20 5:95acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min

hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min. Retention time: 3.11 min, (M+H)⁺: 613. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min. Retention time: 4.04 min, (M+H)⁺: 613.

Preparation of compound K9001:


10



A mixture of Chloro[2-(dicyclohexylphosphino)-3,6-dimethoxy-2',4', 6'-triisopropyl-1,1'-biphenyl][2-(2-aminoethyl)phenyl]palladium(II) (4.5 mg, 5.6 μmol), 5-(3-((2-(1,2,4-oxadiazol-3-yl)propan-2-yl)carbamoyl)phenyl)-6-chloro-2-(4-fluorophenyl)-N-methylfuro[2,3-b]pyridine-3-carboxamide (30 mg, 0.056 mmol), di-tert-butyl(2',4',6'-triisopropyl-3-methoxy-6-methyl-[1,1'-biphenyl]-2-yl)phosphine (2.6 mg, 5.6 μmol), sodium 2-methylbutan-2-olate (30 mg, 0.28 mmol) were combined in trifluoroethanol and heated to 65 °C for 2 hours and then at 90 °C for 16 hours. The reaction mixture was purified by preparative reverse phase HPLC on a C18 column using a suitably buffered H₂O/CH₃CN gradient, and concentrated. The minor product is consistent with : 5-(3-((2-cyanopropan-2-yl)carbamoyl)phenyl)-2-(4-fluorophenyl)-N-methyl-6-(2,2,2-trifluoroethoxy)furo[2,3-b]pyridine-3-carboxamide (1.0 mg, 1.7 μmol, 3.0 % yield) by LCMS and NMR. ¹H NMR (400MHz, CDCl₃) δ 8.31 - 8.28 (m, 1H), 7.96 - 7.89 (m, 4H), 7.77 - 7.73 (m, 1H), 7.61 - 7.55 (m, 1H), 7.26 - 7.21 (m, 2H), 6.22 - 6.17 (m, 1H), 5.88 - 5.80 (m, 1H), 4.95 - 4.86 (m, 2H), 3.01 (d, J=5.0 Hz, 3H), 1.85 (s, 6H). LC-MS retention time: 1.76 min; m/z (M+H)⁺: 555. LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a Phenomenex-Luna 3u C18 2.0x30mm

column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nM. The elution conditions employed a flow rate of 1 mL/min, a gradient of 100% solvent A / 0% solvent B to 0% solvent A / 100% solvent B, a gradient time of 2 min, a hold time of 1 min, and an analysis time of 3 min where solvent A was 10% acetonitrile / 90% H₂O / 0.1% trifluoroacetic acid and solvent B was 10% H₂O / 90% acetonitrile / 0.1% trifluoroacetic acid. MS data was determined using a Micromass Platform for LC in electrospray mode.

Preparation of compound K10001:




Pd/C (9.0 mg, 8.5 μ mol) was added to a stirring solution of (E)-5-(3-((2-(1,2,4-oxadiazol-3-yl)propan-2-yl)carbamoyl)phenyl)-2-(4-fluorophenyl)-N-methyl-6-(prop-1-en-1-yl)furo[2,3-b]pyridine-3-carboxamide (23 mg, 0.043 mmol) in MeOH (853 μ l) at room temperature. The reaction mixture was placed in a Parr bomb and charged with 25 PSI of H₂ (g) and the reaction mixture was allowed to stir for 4 hours. LCMS indicated no conversion. Pd/C (9.0 mg, 8.5 μ mol) was added and the reaction mixture was placed in a Parr bomb and charged with 50 PSI of H₂ (g) and the reaction mixture was allowed to stir for 16 hours. The reaction mixture was filtered and purified by preparative reverse phase HPLC on a C18 column using a suitably buffered H₂O/CH₃CN gradient, and concentrated to give 5-(3-((2-cyanopropan-2-yl)carbamoyl)phenyl)-2-(4-fluorophenyl)-N-methyl-6-propylfuro[2,3-b]pyridine-3-carboxamide (2.5 mg, 4.7 μ mol, 11 % yield) consistent by LCMS and NMR. ¹H NMR (400MHz, CDCl₃) δ 8.01 - 7.95 (m, 2H), 7.94 (s, 1H), 7.82 (d, J=7.5 Hz, 1H), 7.70 (s, 1H), 7.56 - 7.45 (m, 2H), 7.22 (t, J=8.7 Hz, 2H), 6.58 (s, 1H), 6.09 - 6.01 (m, 1H), 2.99 (d, J=5.0 Hz, 3H), 2.74 - 2.67 (m, 2H), 1.86 (s, 6H), 1.71 - 1.65 (m, 2H), 0.85 (t, J=7.4 Hz, 3H). LC-MS retention time: 2.05 min; m/z (M+H)⁺: 499. LC data was recorded on a Shimadzu LC-10AS liquid chromatograph equipped with a

Phenomenex-Luna 3u C18 2.0x30mm column using a SPD-10AV UV-Vis detector at a detector wave length of 220 nM. The elution conditions employed a flow rate of 1 mL/min, a gradient of 100% solvent A / 0% solvent B to 0% solvent A / 100% solvent B, a gradient time of 2 min, a hold time of 1 min, and an analysis time of 3 min where
 5 solvent A was 10% methanol / 90% H₂O / 0.1% trifluoroacetic acid and solvent B was 10% H₂O / 90% methanol / 0.1% trifluoroacetic acid. MS data was determined using a Micromass Platform for LC in electrospray mode.

Preparation of compounds K11001 and K11002:

10



2-(3H-[1,2,3]triazolo[4,5-b]pyridin-3-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate(V) (45 mg, 0.12 mmol) was added to stirring solution of 3-(6-(sec-butyl)-2-(4-fluorophenyl)-3-(methylcarbamoyl)furo[2,3-b]pyridin-5-yl)benzoic acid (35
 15 mg, 0.078 mmol), N-ethyl-N-isopropylpropan-2-amine (41 μ l, 0.24 mmol) and 2-amino-2-methylpropanenitrile hydrochloride (11 mg, 0.094 mmol) in DMF (0.8 μ l) at room temperature. The mixture was allowed to stir at room temperature for 30 minutes. The entire reaction mixture was purified via preparative LC/MS with the following
 20 conditions: Column: XBridge C18, 19 x 200 mm, 5- μ m particles; Mobile Phase A: 5:95 acetonitrile: water with 10-mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile: water with 10-mM ammonium acetate; Gradient: 40-80% B over 20 minutes, then a 7-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired product were combined and dried via centrifugal evaporation. The material was further purified through
 25 chiral separation.

First eluting isomer: The yield of the product was 8.9 mg, and its purity was 100%. Two analytical LC/MS injections were used to determine the final purity. Injection 1

- conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 2.99 min, (M+H)⁺: 513
- 5 Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 4.04 min, (M+H)⁺: 513. ¹H NMR (500MHz, DMSO-d₆) δ 8.86 (s, 1H), 8.55 - 8.47 (m, 1H), 8.10 - 8.02 (m, 2H), 7.98 - 7.94 (m, 1H), 7.90 (s, 2H), 7.66 - 7.61 (m, 1H), 7.59 - 7.55 (m, 1H), 7.43 - 7.36 (m, 1H), 2.93 - 2.86 (m, 1H), 2.81 (d, J=4.3 Hz, 3H), 1.83 - 1.74 (m, 1H), 1.71 (s, 6H), 1.58 - 1.47 (m, 1H), 1.19 (d, J=6.7 Hz, 3H), 0.65 (s, 3H).
- 10 Second eluting isomer: The yield of the product was 9.5 mg, and its purity was 100%. Two analytical LC/MS injections were used to determine the final purity. Injection 1 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 1 mL/min; Detection: UV at 220 nm. Retention time: 2.99 min, (M+H)⁺: 513. Injection 2 conditions: Column: Waters BEH C18, 2.0 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 methanol:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 methanol:water with 10 mM ammonium acetate; Temperature: 40 °C; Gradient: 0.5 min hold at 0%B, 0-100% B over 4 minutes, then a 0.5-minute hold at 100% B; Flow: 0.5 mL/min; Detection: UV at 220 nm. Retention time: 4.04 min, (M+H)⁺: 513. ¹H NMR (500MHz, DMSO-d₆) δ 8.82 (s, 1H), 8.50 - 8.45 (m, J=4.3 Hz, 1H), 8.06 (dd, J=8.9, 5.5 Hz, 2H), 7.96 (d, J=7.9 Hz, 1H), 7.92 - 7.85 (m, 2H), 7.67 - 7.60 (m, 1H), 7.60 - 7.54 (m, 1H), 7.41 (t, J=8.9 Hz, 2H), 2.93 - 2.86 (m, 1H), 2.81 (d, J=4.6 Hz, 2H), 1.84 - 1.75 (m, 1H), 1.71 (s, 6H), 1.58 - 1.49 (m, 1H), 1.20 (d, J=6.7 Hz, 3H), 0.66 (t, J=7.3 Hz, 3H).
- 15
- 20
- 25
- 30

Biological Methods

The compound demonstrated activity against HCV NS5B as determined in the following HCV RdRp assays.

5

HCV NS5B RdRp cloning, expression, and purification. The cDNA encoding NS5B proteins of HCV genotype 1b (Con1), a genotype 1b variant with amino acid 316 mutated from cysteine to asparagine, and genotype 2a (JFH-1), were cloned into the pET21a expression vector. Each untagged protein was expressed with an 18 amino acid C-terminal truncation to enhance the solubility. The E. coli competent cell line BL21(DE3) was used for expression of the protein. Cultures were grown at 37 °C for ~ 4 hours until the cultures reached an optical density of 2.0 at 600 nm. The cultures were cooled to 20 °C and induced with 1 mM IPTG. Fresh ampicillin was added to a final concentration of 50 µg/mL and the cells were grown overnight at 20 °C.

15

Cell pellets (3L) were lysed for purification to yield 15-24 mgs of purified NS5B. The lysis buffer consisted of 20 mM Tris-HCl, pH 7.4, 500 mM NaCl, 0.5% triton X-100, 1 mM DTT, 1mM EDTA, 20% glycerol, 0.5 mg/mL lysozyme, 10 mM MgCl₂, 15 ug/mL deoxyribonuclease I, and Complete TM protease inhibitor tablets (Roche). After addition of the lysis buffer, frozen cell pellets were resuspended using a tissue homogenizer. To reduce the viscosity of the sample, aliquots of the lysate were sonicated on ice using a microtip attached to a Branson sonicator. The sonicated lysate was centrifuged at 100,000 x g for 30 minutes at 4 °C and filtered through a 0.2 µm filter unit (Corning).

20

25

The protein was purified using two sequential chromatography steps: Heparin sepharose CL-6B and polyU sepharose 4B. The chromatography buffers were identical to the lysis buffer but contained no lysozyme, deoxyribonuclease I, MgCl₂ or protease inhibitor and the NaCl concentration of the buffer was adjusted according to the requirements for charging the protein onto the column. Each column was eluted with a NaCl gradient which varied in length from 5-50 column volumes depending on the column type. After the final chromatography step, the resulting purity of the enzyme is >90% based on SDS-PAGE analysis. The enzyme was aliquoted and stored at -80 °C.

30

HCV NS5B RdRp enzyme assay. An on-bead solid phase homogeneous assay was used in a 384-well format to assess NS5B inhibitors (WangY-K, Rigat K, Roberts S, and Gao M (2006) Anal Biochem, 359: 106-111). The biotinylated oligo dT₁₂ primer was captured on streptavidin-coupled imaging beads (GE, RPNQ0261) by mixing primer and beads in 1X buffer and incubating at room temperature for three hours. Unbound primer was removed after centrifugation. The primer-bound beads were resuspended in 3X reaction mix (20 mM Hepes buffer, pH 7.5, dT primer coupled beads, poly A template, ³H-UTP, and RNase inhibitor (Promega N2515)). Compounds were serially diluted 1:3 in DMSO and aliquoted into assay plates. Equal volumes (5 µL) of water, 3X reaction mix, and enzyme in 3X assay buffer (60 mM Hepes buffer, pH 7.5, 7.5 mM MgCl₂, 7.5 mM KCl, 3 mM DTT, 0.03 mg/mL BSA, 6 % glycerol) were added to the diluted compound on the assay plate. Final concentration of components in 384-well assay: 0.36 nM template, 15 nM primer, 0.29 µM ³H-UTP (0.3 µCi), 1.6 U/µL RNase inhibitor, 7 nM NS5B enzyme, 0.01 mg/mL BSA, 1 mM DTT, and 0.33 µg/µL beads, 20 mM Hepes buffer, pH 7.5, 2.5 mM MgCl₂, 2.5 mM KCl, and 0.1 % DMSO.

Reactions were allowed to proceed for 24 hours at 30° C and terminated by the addition of 50 mM EDTA (5 µL). After incubating for at least 15 minutes, plates were read on an Amersham LEADseeker multimodality imaging system.

IC₅₀ values for compounds were determined using ten different [I]. IC₅₀ values were calculated from the inhibition using the four-parameter logistic formula $y = A + ((B - A) / (1 + ((C/x)^D)))$, where A and B denote minimal and maximal % inhibition, respectively, C is the IC₅₀, D is hill slope and x represents compound concentration.

Cell lines. The cell lines used to evaluate compounds consist of a human hepatocyte derived cell line (Huh-7) that constitutively expresses a genotype 1b (Con-1) HCV replicon or a genotype 1b (Con-1) HCV replicon with an asparagine replacing the cysteine at amino acid 316, or a genotype 2a (JFH-1) replicon, containing a Renilla luciferase reporter gene. These cells were maintained in Dulbecco's modified Eagle medium (DMEM) containing 10% FBS, 100 U/mL penicillin/streptomycin and 1.0 mg/mL G418.

HCV replicon luciferase assay. To evaluate compound efficacy, titrated compounds were transferred to sterile 384-well tissue culture treated plates, and the plates were seeded with HCV replicon cells (50 μ L at a density of 2.4×10^3 cells/well) in DMEM containing 4 % FBS (final DMSO concentration at 0.5 %). After 3 days

5 incubation at 37°C, cells were analyzed for Renilla Luciferase activity using the EnduRen substrate (Promega cat #E6485) according to the manufacturer's directions. Briefly, the EnduRen substrate was diluted in DMEM and then added to the plates to a final concentration of 7.5 μ M. The plates were incubated for at least 1 h at 37°C then read on a Viewlux Imager (PerkinElmer) using a luminescence program. The 50% effective

10 concentration (EC_{50}) was calculated using the four-parameter logistic formula noted above.

To assess cytotoxicity of compounds, Cell Titer-Blue (Promega) was added to the EnduRen-containing plates and incubated for at least 4 hrs at 37°C. The fluorescence

15 signal from each well was read using a Viewlux Imager. All CC_{50} values were calculated using the four-parameter logistic formula.







Compound EC_{50} data is expressed as A: < 100 nM; B = 100-1000 nM; C > 1000 nM). Representative data for compounds are reported in Table 2.

Table 2






20







Cmpd#	Structure	EC_{50} (μ M) 1b
10001		0.0738 A
10002		A






Cmpd#	Structure	EC ₅₀ (uM) 1b
10003		A
10004		0.1051 B
10005		A
10006		0.0760 A
11001		A
11002		A
11003		0.0046 A






Cmpd#	Structure	EC ₅₀ (uM) 1b
11004		A
11005		A
11006 Chiral		0.0022 A
11007 Chiral		0.0027 A
11008		A
11009 Chiral		A







Cmpd#	Structure	EC ₅₀ (uM) 1b
11010 Chiral		A
11011		0.0075 A
11012		A
11013		N/A
12001		0.0025 A







Cmpd#	Structure	EC ₅₀ (uM) 1b
12002		A
12003		A
12004		0.0033 A
12005		N/A
13001		N/A






Cmpd#	Structure	EC ₅₀ (uM) 1b
13002		0.0024 A
13003		A
13004 Chiral		A
13005 Chiral		A
13006		0.0036 A
13007 Chiral		A






Cmpd#	Structure	EC ₅₀ (uM) 1b
13008 Chiral		A
14001		0.0022 A
14002		A
15001		A
15002		0.0042 A






Cmpd#	Structure	EC ₅₀ (uM) 1b
16001		A
17001		0.0018 A
18001		0.0037 A
20001		A
20002		A





Cmpd#	Structure	EC ₅₀ (uM) 1b
20003 Chiral		A
20004 Chiral		A
20005		0.0140 A
20006		0.0031 A
20007 Chiral		A
20008 Chiral		A





Cmpd#	Structure	EC ₅₀ (uM) 1b
21001		A
22001		0.0055 A
22002 Chiral		A
22003 Chiral		A
22004		0.0040 A
22005 Chiral		A





Cmpd#	Structure	EC ₅₀ (uM) 1b
22006 Chiral		A
23001		A
23002		0.0072 A
24001		0.0539 A
30001		A





Cmpd#	Structure	EC ₅₀ (uM) 1b
30002		0.0394 A
30003		A
31001		A
31002		0.0061 A
40001		A





Cmpd#	Structure	EC ₅₀ (uM) 1b
50001		0.0107 A
K1001		0.0061 A
K1002		A
K1003		A
K2001		A





Cmpd#	Structure	EC ₅₀ (uM) 1b
K2002		0.0034 A
K2003		A
K2004		A
K2005		A





Cmpd#	Structure	EC ₅₀ (uM) 1b
K3001		A
K4001		A
K4002		A
K4003		0.0043 A

Cmpd#	Structure	EC ₅₀ (uM) 1b
K4004		0.0040 A
K5001		A
K5002		A
K5003		A

Cmpd#	Structure	EC ₅₀ (uM) 1b
K5004		0.0028 A
K6001		A
K7001		A
K7002		A

Cmpd#	Structure	EC ₅₀ (uM) 1b
K7003		A
K7004		A
K7005		0.0028 A
K8001		A

Cmpd#	Structure	EC ₅₀ (uM) 1b
K8002		A
K8003		A
K8004		A
K8005		0.0036 A

Cmpd#	Structure	EC ₅₀ (uM) 1b
K9001		0.0177 uM (EC ₅₀ for 1a instead) A
K10001		A
K11001	 first eluting isomer	0.0133 A
K11002	 second eluting isomer	0.0044 A

It will be evident to one skilled in the art that the present disclosure is not limited to the foregoing illustrative examples, and that it can be embodied in other specific forms without departing from the essential attributes thereof. It is therefore desired that the examples be considered in all respects as illustrative and not restrictive, reference being


5 made to the appended claims, rather than to the foregoing examples, and all changes

which come within the meaning and range of equivalency of the claims are therefore intended to be embraced therein.

CLAIMS

We claim:

1. A compound of Formula I, including pharmaceutically acceptable salts thereof:



I

wherein

Z is C-R⁵ or N;

R⁰ is hydrogen;

R¹ is methyl;

R² is phenyl that is independently substituted with 0-2 halo or methoxy, or is para substituted with W-Ar;

W is -O- or -NH-;

Ar is phenyl or para-halophenyl;

R³ is hydrogen, fluoro, or chloro;

R⁴, R⁵, and R⁶ are independently selected from the group of hydrogen, halo, alkyl, haloalkyl, alkoxy, and perdeuteroalkoxy;

R^{7a} , R^{7b} are each independently selected from the group of hydrogen, alkyl, cycloalkyl, and Ar^1 , or together

R^{7a} and R^{7b} form a 3-7 membered carbocyclic ring;

R^8 is hydrogen,

Ar^1 is phenyl, a 5-membered heteroaromatic ring or a 6-membered heteroaromatic ring;

R^9 is hydrogen, halo, R^{201} , OR^{202} , $NR^{203}R^{204}$;

R^{201} is alkyl, alkenyl, or C_1 - C_4 alkyl with between one to all of the hydrogens replaced by fluoro;

R^{202} is C_1 - C_3 alkyl, or C_1 - C_3 alkyl with between one to all of the hydrogens replaced by fluoro;

R^{203} is hydrogen; and

R^{204} is C_1 - C_3 alkyl, or C_1 - C_3 hydroxyalkyl, or is C_1 - C_3 alkyl with between one to all of the hydrogens replaced by fluoro.

2. A compound of claim 1 wherein:

R^2 is para fluorophenyl, R^3 is hydrogen, R^4 , R^5 , and R^6 are each independently selected from hydrogen, fluoro, $-OCH_3$, and $-OCD_3$,

R^{7a} is selected from the group of hydrogen, methyl, fluoromethyl, and cyclopropyl;

R^{7b} is selected from the group of hydrogen, methyl, fluoromethyl, cyclopropyl, and Ar^1 ,

or together R^{7a} and R^{7b} form a cyclopropyl or cyclobutyl ring;

Ar¹ is phenyl or pyrimidyl;

R⁹ is R²⁰¹ or NR²⁰³R²⁰⁴;

R²⁰¹ is -CH₂CH₂CF₃ or vinyl; and

R²⁰⁴ is -CH₂CF₃, -CH₂CF₂CF₃ or -CH₂CH₂OH.

3. A compound of claim 2 wherein R⁴ is hydrogen, R⁵ is hydrogen or fluoro, R^{7b} is selected from the group of hydrogen, methyl, fluoromethyl, and cyclopropyl, or together R^{7a} and R^{7b} form a cyclopropyl or cyclobutyl ring; and

R²⁰¹ is -CH₂CH₂CF₃.

4. A compound of claim 2 wherein Z is N.

5. A compound of claim 2 wherein Z is CR⁵.

6. A compound of claim 4 wherein R⁴ is hydrogen and R⁶ is -OCD₃.


7. A compound of claim 1 wherein R⁴ is hydrogen, R⁵ is hydrogen or fluoro, R⁶ is hydrogen, fluoro, or -OCH₃, R^{7a} is selected from the group of hydrogen, methyl, fluoromethyl, and cyclopropyl, R^{7b} is selected from hydrogen, methyl, fluoromethyl, or cyclopropyl, or together R^{7a} and R^{7b} form a cyclopropyl or cyclobutyl ring; and








R²⁰¹ is -CH₂CH₂CF₃.


8. A compound of claim 7 wherein R⁵ is hydrogen, R⁶ is fluoro, R^{7a} is methyl, R^{7b} is cyclopropyl, and R⁹ is R²⁰¹.

9. The compound existing as a single enantiomer which is: (R)-5-(3-((1-cyano-1-cyclopropylethyl)carbamoyl)-4-fluorophenyl)-2-(4-fluorophenyl)-N-methyl-6-(3,3,3-trifluoropropyl)furo[2,3-b]pyridine-3-carboxamide.


10. The compound existing as a single enantiomer which is: (S)-5-(3-((1-cyano-1-cyclopropylethyl)carbamoyl)-4-fluorophenyl)-2-(4-fluorophenyl)-N-methyl-6-(3,3,3-trifluoropropyl)furo[2,3-b]pyridine-3-carboxamide.
11. A compound of claim 9 or claim 10, wherein said compound exists as the single enantiomer that exhibits a minus rotation when optical rotation is measured via standard methods in a polarimeter.
12. A compound, including pharmaceutically acceptable salts thereof, which is selected from the group of:









,




,




,




,




,




,




,




,




,



,















,




,












,




,




,









,















,









,





































13. A compound, including pharmaceutically acceptable salts thereof, which is selected from the group of:

























,




,




,









,




,




,







2




3



3




2







first eluting isomer







and




second eluting isomer


14. A compound, including pharmaceutically acceptable salts thereof, which is selected from the group of:









,




,




,









,










,




,









,




,








,













,












,
























and






15. A compound, including pharmaceutically acceptable salts thereof, which is selected from the group of:














and




16. A compound, including pharmaceutically acceptable salts thereof, which is selected from the group of:



17. A compound, including pharmaceutically acceptable salts thereof, which is selected from the group of:





18. A composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier, excipient and/or diluent.
19. A composition comprising a compound of claim 3 and a pharmaceutically acceptable carrier, excipient and/or diluent.
20. A composition comprising a compound of claim 12 and a pharmaceutically acceptable carrier, excipient and/or diluent.
21. A composition comprising a compound of claim 16 and a pharmaceutically acceptable carrier, excipient and/or diluent.
22. A composition comprising a compound of claim 8 and a pharmaceutically acceptable carrier, excipient and/or diluent.
23. A composition comprising a compound of claim 17 and a pharmaceutically acceptable carrier, excipient and/or diluents.
24. A method of treating hepatitis C infection comprising administering a therapeutically effective amount of a compound of claim 1 to a patient.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2015/021638

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D491/04 A61K31/7068 A61P31/14
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2011/112186 A1 (SQUIBB BRISTOL MYERS CO [US]; PRACITTO RICHARD [US]; KADOW JOHN F [US]) 15 September 2011 (2011-09-15) page 40, 3rd and 6th compound; page 41, 6th compound; page 43, 3rd and 6th compound; page 50, 3rd compound; page 57, 3rd compound; page 1, line 5 - page 1, line 8 -----	1-24



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

7 May 2015

Date of mailing of the international search report

26/05/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Schmid, Arnold

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2015/021638

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2011112186 A1	15-09-2011	CN 102906089 A	30-01-2013
		EP 2545050 A1	16-01-2013
		JP 5572725 B2	13-08-2014
		JP 2013522191 A	13-06-2013
		WO 2011112186 A1	15-09-2011
