



(51) International Patent Classification:

C07D 401/04 (2006.01) A61P 25/34 (2006.01)
A61K 31/465 (2006.01)

(21) International Application Number:

PCT/EP2018/085444

(22) International Filing Date:

18 December 2018 (18.12.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

17210181.8 22 December 2017 (22.12.2017) EP

(71) Applicants: **SIEGFRIED AG** [CH/CH]; Untere
Brühlstrasse 4, 4800 Zofingen (CH). **CONTRAF-NICO-
TEX-TOBACCO GMBH** [DE/DE]; Herbststraße 8, 74072
Heilbronn (DE).

(72) Inventors: **WEBER, Beat**; Wiesenstrasse 4, 4800 Zofin-
gen (CH). **PAN, Ben**; Hongyang Building 4 Room 1403,
East Youth Road, Nantong Chongchuan Area, Jiangsu
Province, 226001 (CN).

(74) Agent: **HARMS, Guido**; Siegfried Hameln Services
GmbH, Langes Feld 13, 31789 Hameln (DE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))


Published:

— with international search report (Art. 21(3))


(54) Title: ENANTIOMERIC SEPARATION OF RACEMIC NICOTINE BY ADDITION OF AN O,O'-DISUBSTITUTED TAR-
TARIC ACID ENANTIOMER



Formula I-a
mixture of (R)- and (S)-Nicotine



Formula I-b
(S)-Nicotine




Formula I-c
(R)-Nicotine

(57) Abstract: The present invention relates to a method of separating racemic nicotine of Formula (I-a) as a mixture of the (R)- and (S)-enantiomers into the enantiomerically pure (S)- and (R)-nicotine represented by Formula (I-b) and (I-c), by adding a mixture of the L- and the D-enantiomer of a O,O'-disubstituted tartaric acid, wherein the molar ratio of the L- to the D-enantiomer is from 80:20 to 95:5, and obtaining the (S)-nicotine of formula (I-b), or by adding O,O'-dibenzoyl-D-tartaric acid and obtaining the (R)-nicotine of formula (I-c).


ENANTIOMERIC SEPARATION OF RACEMIC NICOTINE BY ADDITION OF AN O,O'-DISUBSTITUTED TARTARIC ACID ENANTIOMER

Field of the invention


- 5 The present invention relates to a method of separating nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomers into the enantiomerically pure substances represented by Formula I-b and Formula I-c.



10 Formula I-a
mixture of (R)- and (S)-Nicotine



Formula I-b
(S)-Nicotine



Formula I-c
(R)-Nicotine

The invention further relates to a method of preparing enantiomerically pure (R)-nicotine and enantiomerically pure (S)-nicotine, where the enantiomers are generally difficult to separate.

15

Background of the Invention

Nicotine is a naturally occurring alkaloid used in various applications. Especially (S)-nicotine is used as active pharmaceutical ingredient to treat nicotine abuse and nicotine dependency. Success has
20 further been reported when treating Tourette's syndrome, Alzheimer's disease, schizophrenia and other diseases related to disorders of the nervous system. Common ways of administration are gums, creams, transdermal patches, tablets, nasal sprays, and electric cigarettes.

Considerable quantities of nicotine are also used in agriculture as plant protectant or pesticide
25 against aphids.

Natural nicotine is extracted from tobacco plants, a process that requires efficient purification steps to remove undesired, harmful impurities. The increasing demand of nicotine creates a need

to offer Ecological and economical ways to prepare synthetic nicotine in, particularly chemically and stereochemically, very pure form.

State of the art

5

Nicotine ((S)-3-(1-methylpyrrolidin-2-yl)pyridine) and its enantiomers have been prepared for many years by various, not satisfying methods. Known syntheses are usually expensive and use agents that harm the environment.

10 Pictet A. reported in 1904 already a synthesis for nicotine, including the use of tartaric acid to separate the enantiomers (Berichte der deutschen chemischen Gesellschaft, vol. 37, 1904, pages 1225-1235). Tartaric acid has then been used afterwards for decades (see for example: Aceto M. D., et al. (J. Med. Chem., 1979, vol. 22, 174-177)).

15 More recently Chavdarian C. G. et al. disclosed more modern ideas on the synthesis of optically active nicotinoids (J. Org. Chem., 1982, vol. 41, 1069-1073).

Katsuyama A. et al. reported a way of synthesizing nicotine using potassium *tert*-butanolate for the racemization of nicotine to prepare the starting material for further separation of the
20 enantiomers (Bull. Spec. CORESTA Symposium, Winston-Salem, 1982, p. 15, S05, ISSN.0525-6240).

Further, EP 4 487 172 discloses a synthesis route over 5 different steps, offering a net yield of 37.7%.

25 In US 2016/0326134 a synthesis of nicotine comprising the condensation of 1-methylpyrrolidin-2-one and methyl nicotinate in the presence of a strong base (as K *tert*-butoxide) at reflux to the intermediate potassium 1-methyl-3-nicotinoyl-4,5-dihydro-1H-pyrrol-2-olate is described, which then can be converted into the racemic mixture of R/S nicotine. Di-*para*-toluoyl-L-tartaric acid serves as resolution agent.

30

EP 2 484 673 (US 8,378,111) relies on well-known routes of synthesis and discloses D-DBTA (D-dibenzoyl ester of tartaric acid) as agent to separate the enantiomers.

WO 2016/065209 (EP 3 209 653, US 9,556,142) discloses a preparative way comprising 3 steps to the intermediate myosmine, including the condensation of N-vinylegous-pyrrolidinone and nicotinate ester in the presence of a metal hydride.

- 5 Bowman E. R. et al. describe the preparation of d-nicotine di-(p-toluoyl)-L-tartrate from D,L-nicotine using di-(p-toluoyl)-L-tartaric acid monohydrate as separation agent giving an yield of 50% (Synthetic Comm., 1982, vol. 22, 11, 871-879) .

- 10 Aceto M. D. et al. describe a multi-step resolution process to obtain optically high pure nicotine salts, each step using either pure di-p-toluoyl-D-tartrate or pure di-p-toluoyl-L-tartrate in solvents chosen from methanol, acetone or mixtures thereof (J. Med. Chem, 1979, vol. 2, 22, 174-177).


- 15 Glassco W. et al. describe the synthesis and optical resolution of (+)- and (-)-cis-2,3,3a,4,5,9b-hexahydro-1-methyl-1-H-pyrrolo-[3,2-b]isoquinoline (J. Med. Chem. 1993, 36, 22, 3381-3385).

- During the last years development was mainly focused on purification and optimization of the resolution step of the optically active enantiomers. But there is still a need for further improvement in this step, particularly towards more efficient, more ecologic synthesis of enantiomerically enriched, essentially pure or even pure (R)- or (S)-nicotine and the use of
20 environmental friendly agents and solvents.

Summary of the invention

- 25 The current invention offers a novel method for preparing essentially enantiomeric pure (R)- or (S)-nicotine, including a specific process to separate the enantiomers. At the same time increased yield and a high purity of the final product has been found. Overall the novel method is economically and ecologically superior compared to methods known in the art. The inventors found that racemic nicotine can be separated by an inventive method using economically and ecologically advantageous agents. Furthermore, they found that separation of the racemic
30 mixture is also efficient when using separation agents that are not enantiomerically pure.


In a first aspect the present invention provides a method of preparing a compound of Formula I-b, comprising



(I-b)

providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer

5




(I-a);

separating enantiomers of the compound of Formula I-a by addition of a chiral O,O'-disubstituted tartaric acid, wherein the chiral O,O'-disubstituted tartaric acid comprises the L-enantiomer; and
 obtaining the compound of Formula I-b, wherein the O,O'-disubstituted tartaric acid is a mixture of the L-enantiomer and the D-enantiomer, wherein the molar ratio of the L-enantiomer to the D-enantiomer is preferably 80:20 or more, further preferably 90:10 or more, even further preferably 95:5 or more.

10

Furthermore disclosed is a method of preparing a compound of Formula I-c, comprising


15



(I-c)

providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer

20




(I-a);

separating enantiomers of the compound of Formula I-a by addition of a chiral O,O'-disubstituted tartaric acid, wherein the chiral O,O'-disubstituted tartaric acid is the D-enantiomer; and

- 5 obtaining the compound of Formula I-c, wherein the chiral O,O'-disubstituted tartaric acid is O,O'-dibenzoyl-D-tartaric acid.


Further described is a method of preparing a compound of Formula I-b, comprising



10

(I-b)

providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer



(I-a);

15

separating enantiomers of the compound of Formula I-a by addition of at least a mixture of two enantiomeric separating agents, wherein one enantiomeric separating agent is preferably used at a molar ratio of 50:50 or more, further preferably 80:20 or more, even further preferably 90:10 or more, even more preferably 95:5 or more, compared to the other enantiomeric separating agent;

20 and

obtaining the compound of Formula I-b.

Further embodiments are disclosed in the dependent claims and can be taken from the following description and examples, without being limited thereto.

5 Detailed description of the present invention


If not defined otherwise technical and scientific terms have the same meaning as is generally understood by a skilled person in the field of the invention.

- 10 All ranges disclosed herein are to be considered to be supplemented by the term “about”, unless clearly defined to the contrary or otherwise clear from the context.

All numbers or percentages relating to amounts of a substance within this application are given in wt.%, unless clearly defined to the contrary or otherwise clear from the context.

15


Described is a method of preparing a compound of Formula I-b, i.e. (S)-nicotine, is disclosed, comprising



(I-b)

20


providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer



(I-a);


separating enantiomers of the compound of Formula I-a by addition of a chiral O,O'-disubstituted tartaric acid, wherein the chiral O,O'-disubstituted tartaric acid comprises the L-enantiomer; and obtaining the compound of Formula I-b. In this method the O,O'-disubstituted tartaric acid can consist of the L-enantiomer. In this method the chiral O,O'-disubstituted tartaric acid can further
5 be chosen from O,O'-dibenzoyl tartaric acid (DBTA) and O,O'-ditoluoyl tartaric acid (DTTA), and for example the chiral O,O'-disubstituted tartaric acid can be O,O'-dibenzoyl-L-tartaric acid.

In a first aspect a method of preparing a compound of Formula I-b, i.e. (S)-nicotine, is disclosed, comprising



(I-b)

providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer




(I-a);

separating enantiomers of the compound of Formula I-a by addition of a chiral O,O'-disubstituted tartaric acid, wherein the chiral O,O'-disubstituted tartaric acid comprises the L-enantiomer; and obtaining the compound of Formula I-b, wherein the O,O'-disubstituted tartaric acid is a mixture
20 of the L-enantiomer and the D-enantiomer, wherein the molar ratio of the L-enantiomer to the D-enantiomer is preferably 80:20 or more, further preferably 90:10 or more, even further preferably 95:5 or more.

The nicotine of Formula I-a is not particularly restricted and can be obtained in any suitable way,
25 e.g. by extraction from a natural product or by a synthetic method. It is a mixture of the (R)- and (S)-enantiomer of nicotine that is not particularly limited and which can comprise the two

enantiomers at any ratio, as long as both enantiomers are contained. It can be a racemic mixture, i.e. a mixture with a molar ratio of 50:50, but it can also be a mixture with a ratio of the (S)-enantiomer to the (R)-enantiomer in the range of e.g. 1:99 to 99:1, e.g. 10:90 to 90:10, e.g. 20:80 to 80:20, e.g. 30:70 to 70:30, e.g. 40:60 to 60:40, e.g. 45:55 to 55:45, or any other ratio in between these ratios. The present method of the first aspect allows the separation of the (S)-enantiomer out of this mixture.

An exemplary synthesis of nicotine of Formula I-a as a racemate ((rac)-nicotine) follows the scheme as presented hereunder in steps 1a, 1b, 1c, and 1d, as also can be found in literature:



A mixture of (R)- and (S)-nicotine, e.g. racemic nicotine, can be e.g. synthesized, e.g. in a one-pot synthesis with a telescoping method, and generally with a method starting with a condensation of ethyl nicotinate and 1-vinyl-2-pyrrolidone in presence of a base. In the presence of a strong acid the amide nitrogen is deprotected and decarboxylation and ring closure to myosmine takes place. Reduction of the pyrrole ring to a pyrrolidine ring is performed, followed by methylation to

nicotine. The mixture of nicotine can be resolved with a resolving agent/separating agent in the method of the first aspect, e.g. L-DBTA, to obtain the target product (S)-nicotine, as shown in step 2. The obtained (R)-nicotine rich side stream can further be recycled by racemization, as shown in step 3, and undergo a further separating/resolving step in line with the method of the first aspect.

In the method of the first aspect the chiral O,O'-disubstituted tartaric acid is not particularly limited as long as it is chiral, i.e. optically active. The two substituents on the oxygen of the hydroxy groups are not particularly limited and can be the same or different. According to certain embodiments they are chosen from alkyl groups with 1 to 20 C-atoms, alkene and/or alkyne groups with 2 to 20 C-atoms, aryl groups with 6 to 20 C-atoms; and/or alkyl aryl and/or aryl alkyl groups with 7 to 20 C-atoms that all can be substituted or unsubstituted by functional groups like halogen groups, nitro groups, amine groups, ester groups, amide groups, etc., and which all are preferably unsubstituted. Preferred substituents in the chiral O,O'-disubstituted tartaric acid are aryl groups with 6 to 20 C-atoms; and/or alkyl aryl and/or aryl alkyl groups with 7 to 20 C-atoms that are not substituted.

According to certain embodiments, the chiral O,O'-disubstituted tartaric acid is chosen from O,O'-dibenzoyl tartaric acid (DBTA) and O,O'-ditoluoyl tartaric acid, e.g. O,O'-di-o-toluoyl tartaric acid, O,O'-di-m-toluoyl tartaric acid and/or O,O'-di-p-toluoyl tartaric acid (DTTA), and or mixtures thereof, preferably O,O'-dibenzoyl tartaric acid. According to certain embodiments, it is added in ethanol as solvent.

In the present method the chiral O,O'-disubstituted tartaric acid comprises the L-enantiomer. The O,O'-disubstituted tartaric acid comprises the L-enantiomer and the D-enantiomer as a mixture. It is preferable that the L-enantiomer is contained in excess of the D-enantiomer, i.e. at a molar ratio of bigger than 50:50, e.g. at a molar ratio of L-enantiomer to D-enantiomer of at least 80:20, preferably at least 90:10, further preferably 95:5 or more, wherein this ratio can also be described as enantiomeric excess (ee) of at least 60%, preferably at least 80%, further preferably at least 90%.

Described is that the chiral O,O'-disubstituted tartaric acid is O,O'-dibenzoyl-L-tartaric acid, i.e. has an ee of 100%.

According to certain embodiments, the O,O'-disubstituted tartaric acid is added in ethanol as solvent.

5 While it has been found that the separation of the (R)- and (S)-nicotine can be achieved with a separating agent that is a pure enantiomer, i.e. the L-enantiomer of the chiral O,O'-disubstituted tartaric acid if the (S)-nicotine is to be obtained, it has also surprisingly been found that the same is also accomplished if not a pure separating agent, also termed resolution agent, is used, but also
10 when a mixture of enantiomers of the chiral O,O'-disubstituted tartaric acid is used, which surprisingly achieved the separation effect. Even though pure separating agents / resolution agents are available, there is an advantage in using the economically and ecologically easier accessible agents as mixtures with an excess of one enantiomer, e.g. the L-enantiomer if the (S)-nicotine (compound of Formula I-b) is to be obtained.

15 The O,O'-disubstituted tartaric acid is a mixture of the L-enantiomer and the D-enantiomer, wherein the L-enantiomer is contained in excess over the D-enantiomer, preferably wherein the molar ratio of the L-enantiomer to the D-enantiomer is 80:20 or more, preferably 90:10 or more, further preferably 95:5 or more. According to certain embodiments, it is added in ethanol as solvent.


20 According to certain embodiments, the O,O'-disubstituted tartaric acid is a mixture of O,O'-dibenzoyl-L-tartaric acid (L-DBTA) and O,O'-dibenzoyl-D-tartaric acid (D-DBTA) with a molar ratio of L-DBTA to D-DBTA of more than 50:50, preferably 80:20 or more, further preferably 90:10 or more, even further preferably 95:5 or more. According to certain embodiments, it is added in
25 ethanol as solvent.

In the present method, the solvent used for adding the O,O'-disubstituted tartaric acid is not particularly restricted and can be any suitable solvent in which the O,O'-disubstituted tartaric acid can be dissolved. According to certain embodiments, the solvent is ethanol. For the separation,
30 the mixture obtained by adding the O,O'-disubstituted tartaric acid to the compound of Formula I-a can be e.g. refluxed over a certain time period to react the mixture and/or can be stirred under heating, e.g. to a temperature of at least 40°C, preferably 50°C or more, further preferably 60°C or more, preferably for at least 15 min, further preferably 30 minutes or more. After such heating a

cooling can take place, e.g. to a temperature below 20°C, preferably below 10°C, further preferably below 5°C, preferably for at least 8 hours, preferably at least 10 hours, e.g. 12 hours or more. Also two or more of the reflux and/or heating and optionally cooling steps can be carried out, wherein between each step also seeding with a salt of the S-nicotine and the L-O,O'-disubstituted tartaric acid can take place.


The obtaining of the compound of Formula I-b is not particularly restricted and can be carried out by suitable methods, e.g. hydrolyzing the obtained salt of the (S)-nicotine with the separating agent with water in alkaline medium, extracting with an organic solvent like toluene, and distilling of the solvent. For obtaining the salt of the (S)-nicotine with the separating agent, it can be precipitated beforehand, filtered, and optionally washed, e.g. with ethanol. The steps of precipitating, filtering and washing therein can be carried out repeatedly, e.g. two, three, four or more times.

In a second aspect the present invention relates to a method of preparing a compound of Formula I-c, comprising



(I-c)

providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer



(I-a);

separating enantiomers of the compound of Formula I-a by addition of a chiral O,O'-disubstituted tartaric acid, wherein the chiral O,O'-disubstituted tartaric acid is the D-enantiomer; and obtaining the compound of Formula I-c, wherein the chiral O,O'-disubstituted tartaric acid is O,O'-dibenzoyl-D-tartaric acid.

5

Again, the nicotine of Formula I-a is not particularly restricted and can be obtained in any suitable way, e.g. by extraction from a natural product or by a synthetic method, as e.g. shown above. It is a mixture of the (R)- and (S)-enantiomer of nicotine that is not particularly limited and which can comprise the two enantiomers at any ratio, as long as both enantiomers are contained. It can be a racemic mixture, i.e. a mixture with a molar ratio of 50:50, but it can also be a mixture with a ratio of the (S)-enantiomer to the (R)-enantiomer in the range of e.g. 1:99 to 99:1, e.g. 10:90 to 90:10, e.g. 20:80 to 80:20, e.g. 30:70 to 70:30, e.g. 40:60 to 60:40, e.g. 45:55 to 55:45, or any other ratio in between these ratios. The present method of the second aspect allows the separation of the (R)-enantiomer out of this mixture.

15

Described is that the O,O'-disubstituted tartaric acid is chiral, i.e. optically active and contains the D-enantiomer. The two substituents on the oxygen of the hydroxy groups can be the same or different. They can be chosen from alkyl groups with 1 to 20 C-atoms, alkene and/or alkyne groups with 2 to 20 C-atoms, aryl groups with 6 to 20 C-atoms; and/or alkyl aryl and/or aryl alkyl groups with 7 to 20 C-atoms that all can be substituted or unsubstituted by functional groups like halogen groups, nitro groups, amine groups, ester groups, amide groups, etc., and which all are preferably unsubstituted. Preferred substituents in the chiral O,O'-disubstituted tartaric acid are aryl groups with 6 to 20 C-atoms; and/or alkyl aryl and/or aryl alkyl groups with 7 to 20 C-atoms that are not substituted.

25

Described is that the chiral O,O'-disubstituted tartaric acid can be chosen from O,O'-dibenzoyl-D-tartaric acid and O,O'-ditoluoyl-D-tartaric acid, e.g. O,O'-di-o-toluoyl-D-tartaric acid, O,O'-di-m-toluoyl-D-tartaric acid and/or O,O'-di-p-toluoyl-D-tartaric acid, and or mixtures thereof, preferably O,O'-dibenzoyl-D-tartaric acid.

30

The chiral O,O'-disubstituted tartaric acid can be added in ethanol as solvent.

In the second aspect the chiral O,O'-disubstituted tartaric acid is O,O'-dibenzoyl-D-tartaric acid, i.e. has an ee of 100%. According to certain embodiments, it is added in ethanol as solvent.


It has been found that the separation of the (R)- and (S)-nicotine can be achieved with a separating agent that is a pure enantiomer, i.e. the D-enantiomer of the chiral O,O'-disubstituted tartaric acid if the (R)-nicotine is to be obtained.

In the present method, the solvent used for adding the O,O'-disubstituted D-tartaric acid is not particularly restricted and can be any suitable solvent in which the O,O'-disubstituted D-tartaric acid can be dissolved. According to certain embodiments, the solvent is ethanol. For the separation, the mixture obtained by adding the O,O'-disubstituted tartaric acid to the compound of Formula I-a can be e.g. refluxed over a certain time period to achieve a clear solution and/or can be stirred under heating, e.g. to a temperature of at least 40°C, preferably 50°C or more, further preferably 60°C or more, preferably for at least 15 min, further preferably 30 minutes or more. After such heating a cooling can take place, e.g. to a temperature below 20°C, preferably below 10°C, further preferably below 5°C, preferably for at least 8 hours, preferably at least 10 hours, e.g. 12 hours or more. Also two or more of the reflux and/or heating and optionally cooling steps can be carried out, wherein between each step also seeding with a salt of the R-nicotine and the D-O,O'-disubstituted tartaric acid, i.e. O,O'-dibenzoyl-D-tartaric acid, can take place.

The obtaining of the compound of Formula I-c is not particularly restricted and can be carried out by suitable methods, e.g. hydrolyzing the obtained salt of the (R)-nicotine with the separating agent with water in alkaline medium, extracting with an organic solvent like toluene, and distilling of the solvent. For obtaining the salt of the (R)-nicotine with the separating agent, it can be precipitated beforehand, filtered, and optionally washed, e.g. with ethanol. The steps of precipitating, filtering and washing therein can be carried out repeatedly, e.g. two, three, four or more times.


According to certain embodiments of the second aspect, a seeding with the pure enantiomer, i.e. the (R)-enantiomer, and/or its salt with the O,O'-disubstituted tartaric acid, is carried out, preferably at an increased temperature, further preferably at between 30 and 50°C, particularly preferably at about 40°C, to obtain an increased yield and purity.

Further described is a method of preparing a compound of Formula I-b, comprising



(I-b)

- 5 providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer



(I-a);

- 10 separating enantiomers of the compound of Formula I-a by addition of at least a mixture of two enantiomeric separating agents, wherein one enantiomeric separating agent is used at a molar ratio of 50:50 or more, preferably more than 50:50; further preferably 80:20 or more, even further preferably 90:10 or more, particularly further preferably 95:5 or more, compared to the other enantiomeric separating agent; and obtaining the compound of Formula I-b.

15

- Again, the nicotine of Formula I-a is not particularly restricted and can be obtained in any suitable way, e.g. by extraction from a natural product or by a synthetic method, as e.g. shown above. It is a mixture of the (R)- and (S)-enantiomer of nicotine that is not particularly limited and which can comprise the two enantiomers at any ratio, as long as both enantiomers are contained. It can be a racemic mixture, i.e. a mixture with a molar ratio of 50:50, but it can also be a mixture with a ratio of the (S)-enantiomer to the (R)-enantiomer in the range of e.g. 1:99 to 99:1, e.g. 10:90 to 90:10, e.g. 20:80 to 80:20, e.g. 30:70 to 70:30, e.g. 40:60 to 60:40, e.g. 45:55 to 55:45, or any other ratio in between these ratios. The present method of the third aspect allows the separation of the (S)-enantiomer out of this mixture.
- 20

The addition of at least the mixture of two enantiomeric separating agents can be carried out with ethanol as solvent. For the separation, at least the mixture of the two enantiomeric separating agents and the compound of Formula I-a can be e.g. refluxed over a certain time
5 period to react the mixture and/or can be stirred under heating, e.g. to a temperature of at least 40°C, preferably 50°C or more, further preferably 60°C or more, preferably for at least 15 min, further preferably 30 minutes or more. After such heating a cooling can take place, e.g. to a temperature below 20°C, preferably below 10°C, further preferably below 5°C, preferably for at least 8 hours, preferably at least 10 hours, e.g. 12 hours or more. Also two or more of the reflux
10 and/or heating and optionally cooling steps can be carried out, wherein between each step also seeding with a salt of the nicotine enantiomer to be obtained and the corresponding O,O'-disubstituted tartaric acid for resolution can take place.

The at least mixture of the two enantiomeric separating agents is not particularly restricted, as
15 long as it contains at least two enantiomers and one enantiomeric separating agent is used at a molar ratio of 50:50 or more, e.g. more than 50:50, preferably 80:20 or more, further preferably 90:10 or more, even further preferably 95:5 or more, compared to the other enantiomeric separating agent. Preferably the two enantiomeric separating agents in the at least a mixture of two enantiomeric separating agents are optical antipodes, i.e. two enantiomers, i.e. two
20 stereoisomers that are mirror images of each other that are non-superposable. The mixture can also contain more than one pair of enantiomeric separating agents, e.g. a mixture of two enantiomers each of two enantiomeric separating agents (e.g. DBTA and DTTA), a mixture of two enantiomers each of three enantiomeric separating agents, etc. with the molar ratios for at least one pair, e.g. at least two pairs, e.g. at least three pairs, etc. of 50:50 or more, e.g. more than
25 50:50, preferably 80:20 or more, further preferably 90:10 or more, even further preferably 95:5 or more. If two pairs of enantiomeric separating agents are contained, they can be contained in any suitable ratio, e.g. at a molar ratio between 1:4 and 4:1, e.g. between 1:2 and 2:1, e.g. 1:1. Accordingly, mixtures with three pairs of enantiomeric separating agents or more can be contained in any suitable ratio, e.g. at a molar ratio of between 4:1:1, 1:4:1 and 1:1:4, e.g. between
30 2:1:1, 1:2:1 and 1:1:2, e.g. 1:1:1 for three pairs, etc. Any mixture of suitable enantiomeric separating agents can be applied, e.g. chiral O,O'-disubstituted tartaric acids as described with regard to the first and second aspect of the invention, e.g. chiral O,O'-disubstituted tartaric acids with two acyl substituents, two benzoyl substituents, two toluoyl substituents, with two different substituents,

etc., but also others, e.g. D- and L-tartaric acid, as well as mixtures thereof, as indicated above. Preferably, the enantiomeric separating agents are enantiomeric separating agents with a D- and an L-configuration, wherein preferably the L-configuration is used in excess, i.e. at a molar ratio of more than 50:50, preferably 80:20 or more, further preferably 90:10 or more, even further preferably 95:5 or more, compared to the other enantiomeric separating agent in D-configuration.

It has surprisingly been found that a separation of (S)-nicotine is accomplished if not a pure separating agent, also termed resolution agent, is used, but also when a mixture of separating agents is used with an ee given as above, particularly in ethanol as solvent, which surprisingly achieved the separation effect. Even though pure separating agents / resolution agents are available, there is an advantage in using the economically and ecologically easier accessible agents as mixtures with an excess of one enantiomer, e.g. the L-enantiomer if the (S)-nicotine (compound of Formula I-b) is to be obtained.

The obtaining of the compound of Formula I-b is not particularly restricted and can be carried out by suitable methods, e.g. hydrolyzing the obtained salt of the (S)-nicotine with the separating agent with water in alkaline medium, extracting with an organic solvent like toluene, and distilling of the solvent. For obtaining the salt of the (S)-nicotine with the separating agent, it can be precipitated beforehand, filtered, and optionally washed, e.g. with ethanol. The steps of precipitating, filtering and washing therein can be carried out repeatedly, e.g. two, three, four or more times.

In this method a seeding with the pure enantiomer and/or its salt with the enantiomeric separating agent(s) can be carried out, preferably at an increased temperature, further preferably at between 30 and 50°C, particularly preferably at about 40°C, to obtain an increased yield and purity.

The above embodiments can be combined arbitrarily, if appropriate. Further embodiments and implementations of the invention comprise also not explicitly cited combinations of features mentioned beforehand or hereinafter with regard to examples of the invention. Particularly, a skilled person will also add single aspects as improvements or supplements to the respective basic form of the present invention.

Examples

The present invention will now be described in detail with reference to several examples thereof.

5 However, these examples are illustrative and do not limit the scope of the invention.

Reference Example: General Procedure to prepare racemic nicotine

1.0 eq. of ethyl nicotinate, toluene, 1.6 eq. of sodium ethoxide, and 1.2 eq. of 1-vinyl-2-pyrrolidone (NVP) were charged into a flask at room temperature of about 20°C under anhydrous conditions.

10 Then the reaction was run at 100°C for 3 hours. The reaction was completed, and after the mixture had been cooled down to 30°C, 695.0 g HCl (36% by weight in water) was added dropwise. The low boiling components like acetaldehyde, ethanol and the gaseous CO₂ were removed by distillation, together with parts of the toluene and the water. When the reaction temperature reached 105°C, distillation was stopped and the reaction mixture was stirred at a
15 temperature between 100°C and 105°C overnight. After completion of the reaction, the pH was adjusted to a value between 9.5 to 10.5 using NaOH (30% by weight in water). 360.0 g iso-propanol and 1.0 eq. of NaBH₄ (in respect of the ethyl nicotinate) were charged into the reaction vessel. The reaction was run at room temperature (about 20°C) for more than 3 hours (at this point the content of myosmine was below 3.0% by weight). 2.0 eq. of formic acid (HCOOH, in
20 respect of the ethyl nicotinate) and 1.2 eq. paraformaldehyde ((HCHO)_n, in respect of the ethyl nicotinate) were added and the mixture was stirred at 65°C for at least 3 hours. After the reaction was finished (at this point the content of myosmine was below 0.5% by weight), the pH of the mixture was adjusted to a value of 13 to 14 using NaOH (30% by weight in water). Water was added until all inorganic solids were dissolved. The mixture was extracted twice with toluene. The
25 combined organic phases were concentrated to obtain the crude product. By distillation of the crude product, racemic nicotine was obtained as a colorless oil with a purity of 98.9% (yield was 66.0%) as sample 1.

Further procedures according to the above with different amounts of reactants and solvents that
30 achieve similar results are shown in Table 1 below:

Table 1: Results of a series of reference examples with varying amounts of formic acid and paraformaldehyde

ethyl nicotinate g	HCl g	i-prop. g	NaBH ₄ g	HCOOH g	(HCHO) _n g	Yield %	purity %
11.4 (1.0 eq)	36.0	20.0	2.86 (1.0 eq)	10.3 (3.0 eq)	4.5 (2.0 eq)	46.0	99.7
11.4 (1.0 eq)	36.0	20.0	2.86 (1.0 eq)	10.3 (3.0 eq)	4.5 (2.0 eq)	46.0	98.9
68.4 (1.0 eq)	216.0	48.0	17.2 (1.0 eq)	61.8 (3.0 eq)	13.5 (1.0 eq)	51.8	99.8
68.4 (1.0 eq)	216.0	48.0	17.2 (1.0 eq)	61.8 (3.0 eq)	13.5 (1.0 eq)	49.6	98.7
68.4 (1.0 eq)	216.0	48.0	17.2 (1.0 eq)	31.0 (1.5 eq)	16.2 (1.2 eq)	45.5	99.2
205.2 (1.0 eq)	648.0	360.0	51.6 (1.0 eq)	124.0 (2.0 eq)	48.6 (1.2 eq)	47.0	95.6
205.2 (1.0 eq)	695.0	360.0	51.6 (1.0 eq)	124.0 (2.0 eq)	48.6 (1.2 eq)	66.0	98.9
205.2 (1.0 eq)	840.0	360.0	51.6 (1.0 eq)	124.0 (2.0 eq)	48.6 (1.2 eq)	47.0	99.1
1436.4 (1.0 eq)	4125.0	2500.0	432.0 (1.2 eq)	1312.0 (3.0 eq)	285.0 (1.0 eq)	56.0	99.1
1437 (1.0 eq)	4125.0	2500.0	432.0 (1.2 eq)	875.0 (2.0 eq)	313.0 (1.1 eq)	59.7	97.5

In all cases exactly 1.2 eq NVP related to ethyl nicotinate, exactly 1.6 eq EtONa related to ethyl nicotinate, and 60.0 g toluene per 11.4 g ethyl nicotinate were used.

5 Example 1: Resolution Step

1.0 g racemic nicotine, as obtained by the previous Reference Example, sample 1, was mixed at room temperature with 10 g ethanol (1) and 2.2 g dibenzoyl-L-tartaric acid (L-DBTA) (1 equivalent). The mixture was refluxed for a few minutes and cooled down to room temperature (about 20°C). Precipitation started, and the mixture was stirred overnight (10 to 12 hours) at 20°C. A precipitate that formed was filtered, washed with 2.5 g ethanol (2). The crude product was dissolved in 5.0 g ethanol (3). The mixture was refluxed for a few minutes and cooled down to room temperature. Precipitation started, and the mixture was stirred overnight (10 to 12 hours) at 20°C. The precipitate was filtered and washed with 2.5 g ethanol (4). The product was dried and the pure product was obtained. Using different amounts as given in Table 2 (samples 2 to 5), similar resolution/separation experiments were carried out. Furthermore, it was found that an increased yield and purity could be obtained with seeding at 40°C.

Example 2:

3.2 g nicotine-L-DBTA as produced in Example 1, sample 1, were suspended in 7.2 g water and 7.2 g toluene. Aqueous ammonia (25% by weight) was added until the pH was between 9.8 and 10.4. The phases were separated, and the aqueous phase was extracted twice with 2.4 g toluene. The toluene phases were combined, and toluene was removed by distillation. The residue was distilled under vacuum, yielding 0.93 g pure (S)-nicotine. Enantiopurity was determined by chiral HPLC. The same experiments were carried out with samples 2 to 5 of Example 1, and the results are presented in Table 2.

Table 2: samples of Example 1 using various amounts of solvents in steps 1, 2, 3 and 4

	DBTA 1 g	Ethanol (1) g	Ethanol (2) g	Ethanol (3) g	Ethanol (4) g	Yield	Chiral purity
1	100% L-DBTA	10.0	2.5	5.0	2.5	67.6%	99.8% S-isomer
2	100% L-DBTA	10.0	2.5	10.0	2.5	70.8%	99.6% S-isomer
3	98% L-DTBA 2% D-DTBA	7.5	2.5	5.0	2.5	69.8%	98.2% S-isomer
4	95% L-DTBA 5% D-DTBA	7.5	2.5	5.0	2.5	60.0%	99.6% S-isomer
5	90% L-DTBA 10% D-DTBA	7.5	2.5	5.0	2.5	65.3%	99.6% S-isomer

Molar equivalents of resolution agent and racemic nicotine have been used. The amount of ethanol is chosen as weight multiple of the racemic nicotine.

Example 3: Alternative Resolution Step

1 g racemic nicotine, as obtained in sample 1 of the Reference example, was mixed at room temperature with 10.0 g ethanol (1) and 2.2 g dibenzoyl-L-tartaric acid (L-DBTA) (1 equivalent). The mixture was heated for a few minutes to 70°C and stirred for not less than 0.5 hour at 60°C, seeded with (S)-nicotine L-DBTA salt and slowly cooled to 2°C to 3°C and stirred overnight. The precipitate formed was filtered and the crude product was dissolved in ethanol (2). The mixture was heated to 75°C for a few minutes to obtain a clear solution, cooled to 60°C, seeded with (S)-nicotine L-DBTA salt and stirring for 0.5 hours. The obtained suspension is slowly cooled to 2°C to 3°C and stirred overnight. The precipitate is filtered off, washed with pre-cooled ethanol (3) and dried under vacuum. (S)-nicotine was produced as in Example 2. The results of different samples 6 to 8 with different amounts of ethanol are shown in Table 3 below.

Table 3: samples 6 to 9 of Example 3 using various amounts of solvents in steps 1, 2, 3 and 4

	DBTA 1 g	Ethanol (1) g	Ethanol (2) g	Ethanol (3) g	Yield	Chiral purity
6	95% L_DTBA 5% D-DTBA	4.1	3.6	1.1	67.8%	99.6% S-isomer
7	95% L_DTBA 5% D-DTBA	4.1	3.0	1.1	67.1%	99.4% S-isomer
8	95% L_DTBA 5% D-DTBA	3.6	3.6	1.0	66.2%	99.6% S-isomer

Example 4:


Equivalent amounts of dibenzoyl-D-tartaric acid (23.2 g) and racemic nicotine (10 g) were dissolved in ethanol and stirred for 1 hour, refluxed for 15 minutes, cooled to room temperature and stirred for another hour. (R)-nicotine dibenzoyl-D-tartrate was obtained. (R)-nicotine was produced as in Example 2. After recrystallization in an iso-propanol-methanol mixture (1.0: 0.3), (R)-nicotine was obtained. The results are given in the following table 4.

Table 4: amounts for samples in Example 4

DBTA	Ethanol	Seeds	Yield	Chiral purity
100% D-DBTA	10.0	(R)- nicotine-D- DBTA	60.0%	87.9% (R)-isomer


Claims:

1. A method of preparing a compound of Formula I-b, comprising



(I-b)

providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer




(I-a);

separating enantiomers of the compound of Formula I-a by addition of a chiral O,O'-disubstituted tartaric acid, wherein the chiral O,O'-disubstituted tartaric acid comprises the L-enantiomer; and obtaining the compound of Formula I-b, wherein the O,O'-disubstituted tartaric acid is a mixture of the L-enantiomer and the D-enantiomer, wherein the molar ratio of the L-enantiomer to the D-enantiomer is preferably 80:20 or more, further preferably 90:10 or more, even further preferably 95:5 or more.


2. The method of claim 1, wherein the chiral O,O'-disubstituted tartaric acid is chosen from O,O'-dibenzoyl tartaric acid (DBTA) and O,O'-ditoluoyl tartaric acid (DTTA).
3. The method of claim 1 or 2, wherein the chiral O,O'-disubstituted tartaric acid is O,O'-dibenzoyl-L-tartaric acid (L-DBTA).

4. The method of any one of claims 1 to 3, wherein the O,O'-disubstituted tartaric acid is a mixture of O,O'-dibenzoyl-L-tartaric acid (L-DBTA) and O,O'-dibenzoyl-D-tartaric acid (D-DBTA) with a molar ratio of L-DBTA to D-DBTA of 80:20 or more, preferably 90:10 or more, further preferably 95:5 or more.
5. The method of any one of claims 1 to 4, wherein the addition of the chiral O,O'-disubstituted tartaric acid is carried out with ethanol as solvent.
6. A method of preparing a compound of Formula I-c, comprising



(I-c)

providing nicotine of Formula I-a as a mixture of the (R)- and (S)-enantiomer



(I-a);

separating enantiomers of the compound of Formula I-a by addition of a chiral O,O'-disubstituted tartaric acid, wherein the chiral O,O'-disubstituted tartaric acid is the D-enantiomer; and
obtaining the compound of Formula I-c, wherein the chiral O,O'-disubstituted tartaric acid is O,O'-dibenzoyl-D-tartaric acid.

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/085444

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D401/04 A61K31/465 A61P25/34
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2016/326134 A1 (WILLIS BERT [US] ET AL) 10 November 2016 (2016-11-10) cited in the application	1-5
A	page 2, right-hand column, paragraph [0018] - page 3, left-hand column, paragraph [0019] page 5, left-hand column, paragraph [0033] - left-hand column, paragraph [0039]; example 2 page 6 - page 8; table 1 claims 13, 14 ----- -/--	6



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

11 February 2019

Date of mailing of the international search report

20/02/2019

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Cortés Suárez, José

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2018/085444

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 2 484 673 A2 (DIVIS LAB LTD [IN]) 8 August 2012 (2012-08-08) cited in the application page 4, paragraph [0015] - page 5, paragraph [0019]; tables 1, 2 page 6 - page 7; examples 1-5, comparative example 1 -----	1-6
X	EDWARD R BOWMAN ET AL: "A CONVENIENT METHOD FOR THE PREPARATION OF RACEMIC NICOTINE1", SYNTHETIC COMMUNICAT, TAYLOR & FRANCIS INC, PHILADELPHIA, PA; US, vol. 12, no. 11, January 1982 (1982-01), pages 871-879, XP008169935, ISSN: 0039-7911, DOI: 10.1080/00397918208065965 [retrieved on 2006-12-06]	1-5
A	page 874, paragraph 2 - page 875, paragraph 3 -----	6

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2018/085444

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2016326134 A1	10-11-2016	US 2016326134 A1	10-11-2016
		US 2018030028 A1	01-02-2018

EP 2484673 A2	08-08-2012	EP 2484673 A2	08-08-2012
		ES 2625537 T3	19-07-2017
		US 2012197022 A1	02-08-2012
