

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
14 April 2005 (14.04.2005)

PCT

(10) International Publication Number
WO 2005/033245 A1

(51) International Patent Classification⁷: **C09K 11/06**,
C07D 209/12, G01N 33/58

(21) International Application Number:
PCT/GB2004/004167

(22) International Filing Date:
30 September 2004 (30.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0323171.9 3 October 2003 (03.10.2003) GB

(71) Applicant (for all designated States except US): **COVENTRY UNIVERSITY** [GB/GB]; Priory Street, Coventry CV1 5FB (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LYNCH, Daniel, Eric** [GB/GB]; Coventry University, Priory Street, Coventry CV1 5FB (GB). **HEPTINSTALL, John** [AU/GB]; c/o Coventry University, Priory Street, Coventry CV1 5FB (GB).

(74) Agents: **STAGG, Diana, C.** et al.; Marks & Clerk, 144 New Walk, Leicester LE1 7JA (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:


— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FLUORESCENT COMPOUND




(57) Abstract: A water-soluble photoluminescent compound including a symmetrical skeletal structure of the formula (A) in which x may represent any integer, and in which the phenyl rings may be substituted, is disclosed. A protein detector comprising such a compound and a method of measuring the total dissolved protein content of a fluid sample are also disclosed.

Fluorescent Compound

The present invention relates to novel photoluminescent compounds, to a novel protein
5 detector and to a novel method of detecting the presence of protein in a fluid, in particular
water, or bodily fluids.

Photoluminescent compounds are known, and known photoluminescent compounds include
the class of dyes known as squaraine dyes, in particular the indolenine series of squaraine
10 dyes, which are substituted and unsubstituted compounds of the general formula:



Known indolenine squaraine dyes include both symmetrical and asymmetrical compounds,
15 and are described, for example, in an article by E. Terpetschnig et al. in Anal. Chim. Acta
282 (1993) pages 633 - 641.

These dyes are insoluble in water and are used, inter alia, for the qualitative detection of the presence of protein in fluids. In order to use these dyes for the detection of the presence of proteins, which are themselves water-soluble in a fluid, in particular an aqueous fluid, the dye
5 has to be dissolved in a solvent comprising a mixture of water and an alcohol such as methanol. When protein is added to a water/methanol solution of an indolenine squaraine dye, there is an increase in fluorescence. The analytical method disclosed in the article cited above is a purely qualitative method, simply recording the presence or absence of protein in the test fluid sample, and cannot therefore be used to determine the amount of protein
10 present.

Current methods for the quantitative analysis of protein in fluids and semiquantitatively / quantitatively as stains for proteins in gels include the Bradford [M.M. Bradford in Anal. Biochem. 72 (1976) 248], Lowry [O.H. Lowry et al. in J. Biol. Chem. 193 (1951) 265]
15 methods and bicinchoninic acid (BCA) [P.K. Smith et al. in Anal. Biochem. 150 (1985) 76] methods for fluids and silver stain [C.R. Merrill in Meth. Enzymol. 182 (1990) 477] and Coomassie Blue [S. Fazekas de St. Groth et al. in Biochim. Biophys. Acta 71 (1963) 377] for gels. The Bradford, Lowry and BCA methods are colourimetric techniques (ie. colour change) and are suitable for solution protein ranges of 8 – 2000 µg/mL, require multiple
20 reagents and incubation times and suffer from interference from numerous common reagents. Coomassie Blue, used in the Bradford method, takes 20 – 30 mins and is not hindered by tris buffers. Silver stain is a very sensitive (10 – 100 ng/mL) chromatogenic-technique and is used as a gel stain but care must be taken in the handling of the gel and the whole technique


takes 2 – 3 hours. Coomassie Blue is also a gel stain and can be used instead of silver stain if the protein range is in the chromatogenic μg range. Other commercial techniques for the detection of protein concentration include the NanoOrangeTM Protein Quantitation Kit and the CBQCATM Protein Quantitation Kit, both from Molecular Probes, which are ultra-
5 sensitive solution techniques reliant on changes in fluorescence but are not linear over the ranges given for the squaraine derivative. These two techniques also require 30 mins preparation time.

It is an object of the present invention to provide a novel photoluminescent compound, in
10 particular a novel photoluminescent compound that is soluble in water, or water containing solvent mixtures, and retains its photoluminescent properties in these media.

It is a further object of the present invention to provide a method for detecting the presence of protein in fluids, or as a gel stain, in which the above time constraints are reduced or
15 substantially obviated and at a greater sensitivity.

The present invention provides water-soluble photoluminescent compounds including a symmetrical skeletal structure of the formula

20



in which x may represent any integer, and in which the phenyl rings may be substituted.

5

Particularly preferred compounds according to the invention include those wherein $x = 3$.


Further preferred compounds according to the invention include those which are unsubstituted or in which one or both phenyl rings are substituted in the 5-position with alkyl

10 (including ⁱpropyl and ^tbutyl) or halogen groups.

Particularly preferred compounds according to the invention include 2,4-bis(1-(propan-3-sulfonic acid)-3,3-trimethyl-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate or any metal or quaternary nitrogen (ie. ammonium, mono-, di- or trialkylammonium, pyridinium

15 etc) salt of the sulfonic acid.

The present invention further provides a protein detector, which comprises a compound including a symmetrical skeletal structure of the general formula




in which x may represent any integer, and in which the phenyl rings may be substituted, in
 5 solution in water or a water containing solvent mixture, at concentrations from 1×10^{-9} to 1
 moles per litre.

The present invention further provides a method for measuring the total dissolved protein
 content of a fluid sample, which method includes the steps of:

10


- (a) dissolving a compound including a symmetrical skeletal structure of the
 general formula;



in which x may represent any integer, and in which the phenyl rings may be substituted, in
 5 solution in water or a water containing solvent mixture, at concentrations from 1×10^{-10} to 1
 moles per litre.

- (b) admixing the solution of step (a) with a test fluid sample;
 - 10 (c) measuring the fluorescence of the sample; and
 - (d) comparing the fluorescence with a standard value or values (ie. calibration
 plot), to obtain a value for the total protein content.
- 15 The present invention further provides a method for detecting and / or quantifying
 proteins separated electrophoretically in a supporting matrix, for example polyacrylamide,
 agrose or starch, either in the presence or absence of sodium dodecylsulfate (SDS), that
 has been fixed in an aqueous / organic / acid mixture comprising aqueous methanol and


acetic acid, wherein the matrix is subsequently stained with a photoluminescent compound including a symmetrical skeletal structure of the general formula;



5

in which x may represent any integer, and in which the phenyl rings may be substituted, in solution in 10% aqueous methanol or aqueous acetic acid, at a concentration of from 1×10^{-10} to 1 moles per litre, and destained to visualise bands.

- 10 In order to establish standard values for the fluorescence of aqueous solutions of proteins over the required detection ranges, the fluorescence of solutions containing known concentrations of the protein BSA, was measured using a 5×10^{-8} M aqueous solution of the compound including a symmetrical skeletal structure of formula:



where $x = 3$.

Dye 1

5

at an excitation wavelength of 624 nm ($\epsilon = 1.0 \times 10^5 \text{ mol}^{-1} \text{ cm}^{-1}$) and a detection wavelength of 638 nm.



The results, for two separate calibration plots over the range 0-100 ng/mL BSA are shown
 10 in Figure 1; the results for three separate calibration plots, over the detection range of 0-500 ng/mL BSA are shown in Figure 2 and the results, for three separate calibration plots, over the detection range of 0-10 $\mu\text{g/mL}$ BSA are shown in Figure 3.


It is particularly striking to note the linear response over these detection ranges.


15

It is also an advantage of the photo luminescent compounds of the present invention, that once a calibration curve has been constructed using BSA then any number of protein concentrations can be determined instantaneously. The fluorescence is also at a high

wavelength and is not masked by any diffuse lower wavelength organic fluorescence.

Examples**Figure 1****Figure 2**

**Figure 3**

Experimental**Starting materials**

2,3,3-trimethyl-1-(propan-3-sulfonyl)indolenine was prepared using the literature method of E. Havinga et al. in *Synthetic Metals* 69 (1995) pages 581-582 by heating 2,3,3-trimethylindolenine in excess 1,3-propanesultone and toluene, using a Dean and Stark apparatus. Upon cooling the excess solvent was decanted off and the product washed repeatedly with petroleum ether to yield a red oil. Crystals of 2,3,3-trimethyl-1-(propan-3-sulfonyl)indolenine formed from the oil upon standing for several weeks and the single crystal x-ray structure is shown in Figure 4.





Figure 4

Molecular conformation and atom-naming scheme for the structure of 2,3,3-trimethyl-1-(propan-3-sulfonyl)indolenine.

5

Preparation of Dye 1


2,4-bis(1-(propan-3-sulfonic acid)-3,3-dimethyl-2-indolinyldenemethyl)cyclobutenebis(ylium)-1,3-diolate was prepared using the literature method of Sprenger and Ziegenbein in Agnew. Chem. Int. Ed. 6 (1967) page 533 by refluxing 2:1 molar amounts of 2,3,3-trimethyl-1-(propan-3-sulfonyl)indolenine and squaric acid with catalytic amounts of quinaline in 50/50 toluene/butan-1-ol using a Dean and Stark apparatus. The product was collected in vacuo after removal of the reaction solvents and repeated heated washings with petroleum ether.

15

Claims

1. A water-soluble photoluminescent compound including a symmetrical skeletal structure of the formula

5



in which x may represent any integer, and in which the phenyl rings may be substituted.

10


- 2 A photoluminescent compound according to claim 1 wherein $x = 3$.

- 3 A photoluminescent compound according to claim 1 or claim 2 which is unsubstituted.

15

- 4 A photoluminescent compound according to claim 1 or claim 2 in which one or both phenyl rings are substituted in the 5-position with alkyl, including ⁱpropyl and ^tbutyl, or halogen groups.

5. A photoluminescent compound according to any of claims 1 to 4 which is 2,4-bis(1-(propan-3-sulfonic acid)-3,3-trimethyl-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate or any metal or quaternary nitrogen (ie. ammonium, mono-, di- or trialkylammonium, pyridinium etc) salt of the sulfonic acid.
6. A photoluminescent compound substantially as herein described and with reference to the Examples.
7. A protein detector, which comprises a photoluminescent compound including a symmetrical skeletal structure of the general formula




- in which x may represent any integer, and in which the phenyl rings may be substituted, in solution in water, at concentrations from 1×10^{-10} to 1 moles per litre.
8. A method for measuring the total dissolved protein content of a fluid sample,

which method includes the steps of:

- (a) dissolving a photoluminescent compound including a symmetrical skeletal structure of the general formula;

5



in which x may represent any integer, and in which the phenyl rings may be substituted, in solution in water, at concentrations from 1×10^{-10} to 1 moles per litre.


10

- (b) admixing the solution of step (a) with a test fluid sample;

- (c) measuring the fluorescence of the sample; and

- 15 (d) comparing the fluorescence with a standard value, to obtain a value for the total dissolved protein content.

9. A method for detecting and / or quantifying proteins separated electrophoretically in a supporting matrix, either in the presence or absence of sodium dodecylsulfate (SDS), that has been fixed in an aqueous / organic / acid mixture comprising aqueous methanol and acetic acid, wherein the matrix is subsequently stained with a photoluminescent compound including a symmetrical skeletal structure of the general formula;
- 5 compound including a symmetrical skeletal structure of the general formula;



- in which x may represent any integer, and in which the phenyl rings may be substituted, in solution in 10% aqueous methanol or aqueous acetic acid, at a concentration of from 1×10^{-10} to 1 moles per litre, and destained to visualise bands.
- 10

10. A method according to claim 9 wherein the supporting matrix is polyacrylamide, agrose or starch.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB2004/004167

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C09K11/06 C07D209/12 G01N33/58

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C09K C07D G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01/11370 A (DEROOVER GEERT ; MISSFELDT MICHAEL (DE); SIMON LYDIA (DE); BAYER AG (D) 15 February 2001 (2001-02-15) * page 51 through page 57, page 85, claims *	1-10
A	WO 01/02374 A (SURROMED INC) 11 January 2001 (2001-01-11) the whole document ----- -/--	1-10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

29 November 2004

Date of mailing of the international search report

21/12/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Nemes, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2004/004167

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TERPETSCHNIG E ET AL: "SYNTHESIS, SPECTRAL PROPERTIES AND PHOTOSTABILITIES OF SYMMETRICAL AND UNSYMMETRICAL SQUARAINES; A NEW CLASS OF FLUOROPHORES WITH LONG-WAVELENGTH EXCITATION AND EMISSION" ANALYTICA CHIMICA ACTA, ELSEVIER, AMSTERDAM, NL, vol. 282, no. 3, 29 October 1993 (1993-10-29), pages 633-641, XP000563718 ISSN: 0003-2670 cited in the application the whole document	1-10
A	US 2002/147354 A1 (THEODOROPULOS SPYROS) 10 October 2002 (2002-10-10) the whole document	1-10

INTERNATIONAL SEARCH REPORT
information on patent family members

I International Application No
PCT/GB2004/004167

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0111370	A	15-02-2001	DE 19937024 A1	08-02-2001
			AU 6986100 A	05-03-2001
			CA 2381088 A1	15-02-2001
			CN 1369062 T	11-09-2002
			WO 0111370 A1	15-02-2001
			EP 1206703 A1	22-05-2002
			JP 2003506567 T	18-02-2003
WO 0102374	A	11-01-2001	AU 6206800 A	22-01-2001
			CA 2376955 A1	11-01-2001
			EP 1196398 A1	17-04-2002
			WO 0102374 A1	11-01-2001
			US 6403807 B1	11-06-2002
US 2002147354	A1	10-10-2002	NONE	