



(86) Date de dépôt PCT/PCT Filing Date: 2004/12/01
(87) Date publication PCT/PCT Publication Date: 2005/09/09
(45) Date de délivrance/Issue Date: 2011/07/19
(85) Entrée phase nationale/National Entry: 2006/11/30
(86) N° demande PCT/PCT Application No.: US 2004/040103
(87) N° publication PCT/PCT Publication No.: 2005/082325
(30) Priorité/Priority: 2004/02/24 (US10/785,271)

(51) Cl.Int./Int.Cl. *A61K 8/85* (2006.01),
A61K 8/31 (2006.01), *A61K 8/35* (2006.01),
A61K 8/37 (2006.01), *A61K 8/40* (2006.01),
A61Q 17/04 (2006.01)
(72) Inventeurs/Inventors:
BONDA, CRAIG A., US;
NEUDAHL, GARY A., US
(73) Propriétaire/Owner:
HALLSTAR INNOVATIONS CORP., US
(74) Agent: SMART & BIGGAR

(54) Titre : PHOTOSTABILISATION D'UNE COMPOSITION D'ECRAN SOLAIRE COMPRENANT UNE COMBINAISON
D'UN COMPOSE α -CYANO- β,β -DIPHENYLACRYLATE ET D'UN DIALKYLE NAPHTHALATE
(54) Title: PHOTOSTABILIZATION OF A SUNSCREEN COMPOSITION WITH A COMBINATION OF AN α -CYANO- β,β -
DIPHENYLACRYLATE COMPOUND AND A DIALKYL NAPHTHALATE

(57) **Abrégé/Abstract:**

Sunscreen compositions including a dibenzoylmethane derivative, such as avobenzone, that are made more stable by the addition of (a) an α -cyano- β,β -diphenylacrylate compound, e.g., octocrylene and (b) a diester or polyester of naphthalene dicarboxylic acid (diethylhexyl 2,6-naphthalate), having a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0. More particularly, the invention relates to sunscreen compositions which utilize a combination of octocrylene and diethylhexyl 2,6-naphthalate to stabilize other photoactive compounds present in a sunscreen composition and, in particular, to stabilize dibenzoylmethane derivatives, without, or with levels less than 0.5% by weight of, a methoxy-substituted benzophenone derivative, such as benzophenone-3.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
9 September 2005 (09.09.2005)

PCT

(10) International Publication Number
WO 2005/082325 A1

(51) International Patent Classification⁷: **A61K 7/42**

(21) International Application Number:
PCT/US2004/040103

(22) International Filing Date: 1 December 2004 (01.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/785,271 24 February 2004 (24.02.2004) US

(71) Applicant (for all designated States except US): **CPH INNOVATIONS CORP.** [US/US]; 311 South Wacker Drive, Suite 4700, Chicago, IL 60606-6622 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BONDA, Craig, A.** [US/US]; 0S515 Summit Drive, Winfield, IL 60190 (US). **NEUDAHL, Gary, A.** [US/US]; 559 Ada Street, Cary, IL 60013 (US).

(74) Agent: **ANDERSON, Richard, H.**; Marshall, Gerstein & Borun LLP, 233 S. Wacker Drive, Suite 6300, Sears Tower, Chicago, IL 60606-6357 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PHOTOSTABILIZATION OF A SUNSCREEN COMPOSITION WITH A COMBINATION OF AN α -CYANO- β,β -DIPHENYLACRYLATE COMPOUND AND A DIALKYL NAPHTHANATE

(57) Abstract: Sunscreen compositions including a dibenzoylmethane derivative, such as avobenzene, that are made more stable by the addition of (a) an α -cyano- β,β -diphenylacrylate compound, e.g., octocrylene and (b) a diester or polyester of naphthalene dicarboxylic acid (diethylhexyl 2,6-naphthalate), having a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0. More particularly, the invention relates to sunscreen compositions which utilize a combination of octocrylene and diethylhexyl 2,6-naphthalate to stabilize other photoactive compounds present in a sunscreen composition and, in particular, to stabilize dibenzoylmethane derivatives, without, or with levels less than 0.5% by weight of, a methoxy-substituted benzophenone derivative, such as benzophenone-3.

WO 2005/082325 A1

64267-1444

**PHOTOSTABILIZATION OF A SUNSCREEN COMPOSITION WITH A
COMBINATION OF AN α -CYANO- β,β -DIPHENYLACRYLATE COMPOUND
AND A DIALKYL NAPHTHALATE**

Background Of The Invention

Field of the Invention

[0002] The invention relates to sunscreen compositions including a dibenzoylmethane derivative, such as avobenzene, that are made more stable by the addition of (a) an α -cyano- β,β -diphenylacrylate compound, e.g., octocrylene, and (b) a diester or polyester of naphthalene dicarboxylic acid (e.g., diethylhexyl 2,6-naphthalate), having a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0. More particularly, the invention relates to sunscreen compositions which utilize a combination of octocrylene and diethylhexyl 2,6-naphthalate to stabilize other photoactive compounds present in a sunscreen composition and, in particular, to stabilize dibenzoylmethane derivatives, without, or with levels less than 0.5% by weight of, a methoxy-substituted benzophenone derivative, such as benzophenone-3.

Brief Description of Related Technology

[0003] It is well known that ultraviolet radiation (light) having a wavelength from about 280 nm or 290 nm to about 320 nm (UV-B) is harmful to human skin, causing burns that are detrimental to the development of a good sun tan. UV-A radiation (about 320 nm to about 400 nm), while producing tanning of the skin, also can cause damage, particularly to very lightly-colored or sensitive skin, leading to reduction of skin elasticity and to wrinkles. Therefore, a sunscreen composition for use on human skin preferably includes both a UV-A and a UV-B filter to prevent most of the sunlight within the full range of about 280 nm or 290 nm to about 400 nm from damaging human skin.

[0004] Ultraviolet radiation from the sun or artificial sources can also cause harm to coatings containing photoactive substances, such as photoactive pigments and dyes, by breaking down chemical bonds in the structure of a component such as a polymer, a pigment, or a dye. This photodegradation can lead to color fading, loss of gloss, and loss of physical

64267-1444

and protective properties of a coating. Photodegradation can take place in several steps which include one or more components of a coating absorbing UV radiation. The absorbed radiation can excite the absorbing molecules and raise them to a higher energy level, which can be very reactive. If the molecule cannot be relaxed, bond cleavage and the formation of free radicals will occur. These free radicals can attack one or more color molecules and/or a polymer backbone and form more free radicals. UV-A and UV-B filters can also be used to accept UV radiation to protect a pigmented coating.

[0005] The UV-B filters that are most widely used in the U.S. in commercial sunscreen compositions are paramethoxycinnamic acid esters, such as 2-ethylhexyl paramethoxycinnamate, commonly referred to as octyl methoxycinnamate or PARSOL[®] MCX, octyl salicylate, and oxybenzone.

[0006] The organic UV-A filters most commonly used in commercial sunscreen compositions are the dibenzoylmethane derivatives, particularly 4-(1,1-dimethylethyl)-4'-methoxydibenzoylmethane (also called avobenzone, sold under the brand name PARSOL[®] 1789). Other dibenzoylmethane derivatives described as UV-A filters are disclosed in U.S. Patent Nos. 4,489,057, 4,387,089 and 4,562,067.

It is also well known that the above.

described UV-A filters, particularly the dibenzoylmethane derivatives, can suffer from rapid photochemical degradation, when used alone or when combined with the above-described most commercially used UV-B filters.

[0007] The performance of a photoactive compound or a combination of photoactive compounds in a sunscreen composition has been extremely difficult to predict based on the levels of photoactive compounds in the formulation, particularly when the formulation includes one or more photoactive compounds that suffer from relatively rapid photodegradation, such as avobenzone. Because of this, each formulation has required expensive laboratory testing to determine the UV absorbance, as a function of time (quantity) of exposure of the formulation to UV radiation. Moreover, a particularly difficult problem is presented when one photoactive compound in a sunscreen composition acts to increase the rate of photodegradation of another photoactive compound in the composition. This can be accomplished in a number of ways, including a bimolecular reaction between two photoactive compounds and a lowering of the threshold energy need to raise a photoactive compound to its excited state. For example, when avobenzone is combined with octyl

methoxycinnamate, a bimolecular pathway leads to the rapid photodegradation of both the dibenzoylmethane derivative and the octyl methoxycinnamate.

[0008] Methods and compositions for stabilizing photoactive compounds, such as dibenzoylmethane derivatives, e.g., avobenzene, with diesters and/or polyesters of naphthalene dicarboxylic acid are described in U.S. Patent Nos. 5,993,789, 6,284,916 and Gers-Barlag, et al. U.S. Patent No. 6,491,901 ('901). Alternatively, Deflandre, et al., U.S. Patent No. 5,576,354 and Gonzenbach et al., U.S. Patent No. 6,033,649 describe the use of an α -cyano- β , β -diphenylacrylate compound, such as octocrylene, to stabilize a sunscreen composition including a dibenzoylmethane derivative. Thus, Deflandre et al. teach a minimum 1% by weight of octocrylene and Gonzenbach et al. teach a minimum of 0.5% by weight of octocrylene to stabilize a dibenzoylmethane derivative, such as avobenzene, in a sunscreen composition. Gers-Barlag, et al. U.S. Patent No. 6,491,901 ('901) discloses sunscreen compositions containing a dibenzoylmethane derivative with a stabilizing combination of octocrylene and diesters or polyesters of naphthalene dicarboxylic acid wherein the examples have a weight ratio of octocrylene to the diester or polyester of naphthalene dicarboxylic acid in the range of 0.16 to 0.725, and the claims call for a weight ratio between about 0.03 and about 0.9.

[0009] In accordance with the '901 patent, the combination of octocrylene with one or more dialkyl naphthalates, according to the '901 invention, acts synergistically to prevent decomposition of avobenzene (see col. 4, lines 21-35). The combination of octocrylene and dialkyl naphthalates of the '901 patent, in the example formulations, disclose 1% to 2.9% octocrylene and 4% to 8% dialkyl naphthalate combinations.

[0010] In accordance with the compositions and methods described herein, it has been found that weight ratios of an α -cyano- β , β -diphenylacrylate, e.g., octocrylene, to dialkyl naphthalate of at least 0.95, preferably at least about 1.0, provide surprisingly better results for photostabilizing a dibenzoylmethane derivative, such as avobenzene, than lower weight ratios, contrary to the teachings of the '901 patent.

[0011] In accordance with another embodiment of the compositions and methods described herein, the formulations contain 0% to about 10%, by weight, to less than .05% by weight benzophenone-3, more preferably about 0.1 to about 0.49% by weight benzophenone-3. By including benzophenone-3 in the formulation, a PA+++ rating UVA protection grade, as defined by the Technical Committee of the Japan Cosmetic Association (JCIA) in the

64267-1444

Japanese Persistent Pigment Darkening Protocol, attached as an appendix to this application, is obtained. It should be understood that benzophenone-3 in the formulations represents a separate, preferred embodiment, and is not necessary to provide excellent and surprising photostabilization of one or more dibenzoylmethane derivatives so long as the octocrylene/dialkyl naphthalate weight ratio is at least 0.95, preferably at least about 1.0, as shown in the examples.

Summary

[0012] One aspect of the compositions and methods described herein is a composition including a mixture of a dibenzoylmethane derivative, and a combination of (a) an α -cyano- β,β -diphenylacrylate compound, such as octocrylene, and (b) a diester or polyester of naphthalene dicarboxylic acid, wherein the weight ration of (a)/(b) is at least 0.95, preferably at least about 1.0.

[0013] Another aspect of another embodiment of the compositions and methods described herein is a composition including a mixture of a dibenzoylmethane derivative, such as avobenzene, together with a combination of (a) an α -cyano- β,β -diphenylacrylate compound, such as octocrylene, (b) a diester or polyester of naphthalene dicarboxylic acid, wherein the weight ratio of (a) to (b) is at least 0.95, preferably at least about 1.0; and (c) 0 to about 3.0% by weight of benzophenone-3, preferably less than about 0.5% by weight benzophenone-3, more preferably about 0.1% by weight to about 0.49% by weight benzophenone-3.

[0014] Another aspect of another embodiment of the compositions and method described herein is a composition including a mixture of a dibenzoylmethane derivative, such as avobenzene, together with a combination of (a) an α -cyano- β,β -diphenylacrylate compound, such as octocrylene, (b) a diester or polyester of naphthalene dicarboxylic acid, wherein the weight ratio of (a) to (b) is at least 0.95, preferably at least about 1.0 and (c) a compound capable of raising the dielectric constant of an oil phase of the composition to a level of at least about 7.0 preferably at least about 8.0, such as dimethyl capramide and/or diethylhexyl malate.

[0015] Another aspect of the compositions and methods described herein is a composition including a mixture of (a) a dibenzoylmethane derivative, such as avobenzene, (b) an α -cyano- β,β -diphenylacrylate compound, such as oxybenzone, (c) a diester or polyester of naphthalene dicarboxylic acid, wherein the weight ratio of (b) to (c) is at least 0.95,

64267-1444

preferably at least about 1.0; (d) 0 to about 3.0% by weight of benzophenone-3, preferably less than about 0.5% by weight benzophenone-3, more preferably about 0.1% by weight to about 0.49% by weight benzophenone-3; and (e) a compound
 5 capable of raising the dielectric constant of an oil phase of the composition to a level of at least about 7.0 preferably at least about 8.0, such as dimethyl capramide and/or diethylhexyl malate.



[0016] Yet another aspect of the invention is a
 10 composition including a mixture of a dibenzoylmethane derivative, such as avobenzene, an α -cyano- β,β -diphenylacrylate compound, such as octocrylene, a diester or polyester of naphthalene dicarboxylic acid, and less than 0.5% benzophenone-3 capable of receiving a
 15 UVA protection grade of PA+++ (has a PFA - Protection Factor of UVA - of at least 8.0) when tested in accordance with the present Japanese Persistent Darkening Protocol.

According to one aspect of the present invention, there is provided a sunscreen composition, comprising a mixture
 20 of a dibenzoylmethane derivative selected from the group consisting of 2-methyldibenzoylmethane;
 4-methyldibenzoylmethane; 4-isopropyldibenzoylmethane;
 4-tert-butyldibenzoylmethane; 2,4-dimethyldibenzoylmethane;
 2,5-dimethyldibenzoylmethane; 4,4'-diisopropyldibenzoylmethane;
 25 4,4'-dimethoxydibenzoylmethane; 4-tert-butyl-4'-methoxydibenzoylmethane; 2-methyl-5-isopropyl-4'-methoxydibenzoylmethane; 2-methyl-5-tert-butyl-4'-methoxydibenzoylmethane; 2,4-dimethyl-4'-methoxydibenzoylmethane; 2,6-dimethyl-4-tert-butyl-
 30 4'-methoxydibenzoylmethane, and combinations thereof, with:

64267-1444

(a) octocrylene; and

(b) a diester or polyester of naphthalene dicarboxylic acid selected from the group consisting of formula (I), formula (II), and combinations thereof:



5

wherein:

R^1 and R^2 are the same or different and each is selected from the group consisting of a C_1 - C_{22} alkyl group, a residue of a diol having the structure $\text{HO}-\text{R}^3-\text{OH}$, and a
 10 residue of a polyglycol having the structure $\text{HO}-\text{R}^4-(\text{-O}-\text{R}^3-\text{-})_n-\text{OH}$;

each R^3 and R^4 is the same or different and each is selected from the group consisting of a C_1 - C_6 straight or branched chain alkylene group;

15 m and n are each in a range of 1 to 100 and p is in a range of 0 to 100; and

the weight ratio of (a)/(b) is at least 0.95.

According to other aspects of the present invention the composition described herein may be used for
 20 reduction of contact of UV radiation with human skin, for example, wherein the composition is formulation for application to the human skin.

64267-1444

Detailed Description Of The Preferred Embodiments

[0017] Sunscreen compositions containing a dibenzoylmethane derivative UV-A filter compound, such as 4-(1,1-dimethylethyl)-4'-methoxydibenzoylmethane (PARSOL® 1789) are photostabilized by containing (a) an α -cyano- β , β -diphenylacrylate compound, such as octocrylene; and (b) a diester or polyester of naphthalene dicarboxylic acid, wherein the weight ratio of (a)/(b) is at least 0.95, preferably at least about 1.0, more preferably from about 1/1 to about 2/1.

[0018] One aspect of the sunscreen compositions and methods described herein is to photostabilize a sunscreen composition containing a dibenzoylmethane derivative, such as avobenzene, by including in the composition a combination of (a) an α -cyano- β , β -diphenylacrylate compound, such as octocrylene, in an amount in the range of about 0.5% to about 8.0% by weight, preferably about 2% to about 7% by weight, more preferably about 1.5% to about 5% by weight, and (b) a diester or polyester of naphthalene dicarboxylic acid (DEHN), in an amount in the range of about 0.5% to about 8.0% by weight, preferably about 2% to about 7% by weight, more preferably about 1.5% to about 5% by weight, wherein the weight ratio of (a) to (b) is at least 0.95, preferably at least about 1.0 to achieve a stable sunscreen composition.

[0019] In accordance with another embodiment of the sunscreen compositions and methods described herein, an α -cyano- β , β -diphenylacrylate compound is combined with a highly polar solvent or blend of solvents with a high polarity to enhance the photostabilizing effect

of the α -cyano- β,β -diphenylacrylate compound, and thereby enhance the stability of the sunscreen composition.

[0020] In accordance with another embodiment of the compositions and methods described herein the photostabilizing effect of the diester or polyester of naphthalene dicarboxylic acid on the dibenzoylmethane derivative is enhanced by adding benzophenone-3, preferably at a level less than about 0.5% by weight.


[0021] A photoactive compound can be considered stable when, for example, after 30 MED irradiation the photoactive compound has retained at least 90% of its original absorbance at a wavelength or a range of wavelengths of interest (*e.g.*, the wavelength at which or near a photoactive compound has a peak absorbance, such as 350-360 nm for avobenzene). Likewise, a sunscreen composition can include a plurality of photoactive compounds and a sunscreen composition, as a whole, can be considered stable when, for example, after 30 MED irradiation the sunscreen composition has retained at least 90% of its original absorbance at one or more wavelengths of interest (*e.g.*, at or near the peak absorbance wavelength of the primary photoactive compounds).

[0022] It has surprisingly been found that the addition of (a) an α -cyano- β,β -diphenylacrylate compound, and (b) a diester or polyester of naphthalene dicarboxylic acid significantly increases the photostability of the sunscreen composition. Without intending to be limited to any particular mechanism of achieving this increase in stability, it is believed that a diester or polyester of naphthalene dicarboxylic acid stabilizes a dibenzoylmethane derivative by accepting the triplet energy of the dibenzoylmethane derivative once the dibenzoylmethane derivative has reached an excited state as a result of the absorption of ultra-violet light. Once a dibenzoylmethane derivative is excited, it is prone to degrade according to a number of pathways, however, the degradation of the dibenzoylmethane derivative can be substantially reduced or prevented by the use of a diester or polyester of naphthalene dicarboxylic acid to quench (accept) the triplet excited state energy present in an excited dibenzoylmethane molecule. Thus, in one pathway of degradation, a dibenzoylmethane derivative is excited to its triplet state and the excited state triplet energy is released in a bond breaking step, thereby preventing the dibenzoylmethane derivative from further accepting ultra-violet radiation. A diester or polyester of naphthalene dicarboxylic acid may stabilize a dibenzoylmethane derivative by accepting the triplet state (excited state) energy of the excited dibenzoylmethane derivative in such a way as to convert the excited

dibenzoylmethane derivative back to a non-excited state that is capable of reaccepting ultra-violet radiation (energy transfer).

[0023] For this process to work continuously, the diester or polyester of naphthalene dicarboxylic acid must transfer or convert the energy that was accepted from the excited dibenzoylmethane derivative. Without intending to be limited to a particular mechanism, it is believed that when a diester or polyester of naphthalene dicarboxylic acid is excited to its triplet state, the triplet excited state energy is dissipated through vibrations (*i.e.*, heat), which in this group of molecules is a slow mode of dissipating energy. It has been found, quite surprisingly, that by the addition of (a) an α -cyano- β,β -diphenylacrylate compound, together with (b) a diester or polyester of naphthalene dicarboxylic acid, in a weight ratio of (a)/(b) of at least 0.95, preferably at least 1.0, provides the α -cyano- β,β -diphenylacrylate compound with the capacity to accept triplet excited state energy from an excited diester or polyester of naphthalene dicarboxylic acid. Thus, according to one possible mechanism, the efficiency of the dissipation of the excited state energy in an excited diester or polyester of naphthalene dicarboxylic acid is greatly improved by a transfer of energy from an excited diester or polyester of naphthalene dicarboxylic acid to an α -cyano- β,β -diphenylacrylate compound.

[0024] Without intending to be limited to any particular mechanism by which an α -cyano- β,β -diphenylacrylate compound is able to quench the excited state of a diester or polyester of naphthalene dicarboxylic acid, it is believed that the α -cyano- β,β -diphenylacrylate compound accepts the excited state energy and dissipates the energy kinetically in the form of rapid isomerizations. This process is shown below:



wherein the α -cyano- β,β -diphenylacrylate compound (octocrylene shown above as A), accepts the triplet excited state energy and forms a diradical (shown above as A*) at the α and β positions of the acrylate, which converts the double bond into a single bond and allows for the free rotation of the phenyl groups. This rotation occurs rapidly and efficiently to dissipate any excited state energy accepted by the α -cyano- β,β -diphenylacrylate compound. In

64267-1444

solution (*e.g.*, a sunscreen composition), a key limitation on one compound's ability to stabilize another is the ability of the two compounds to come into contact with one another. Thus, according to this mechanism of stabilization, it is preferred to have at least as much of the α -cyano- β,β -diphenylacrylate compound as the diester and/or polyester of naphthalene dicarboxylic acid so that the α -cyano- β,β -diphenylacrylate compound is abundant enough to quickly come into contact with an excited diester or polyester of naphthalene dicarboxylic acid.

[0025] Commonly-assigned U.S. Patent No. 6,485,713 and U.S. Patent No. 6,537,529, describe compositions and methods for increasing the stability of photoactive compounds in a sunscreen composition, *e.g.*, by the addition of polar solvents to the oil phase of a composition. It has been found, quite surprisingly, that by increasing the polarity of the oil phase of a sunscreen composition including (a) an α -cyano- β,β -diphenylacrylate compound, and (b) a diester or polyester of naphthalene dicarboxylic acid, in a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0, the stability of the sunscreen composition is increased. Now knowing that the polarity of the solution affects the stability, one might expect that the more polar the solution is, the greater the stability it will impart to the photoactive compound. In contrast, and even more surprisingly, it has been found that as the polarity of a solvent system including a dissolved, rapidly-photodegradable compound is increased, the rate of photodecay initially decreases -- but then increases again as the polarity is further increased. Thus, a photodegradable compound in solution will degrade as a second-order function of the overall polarity of the solution. Currently accepted photochemical theory provides the possibility that the mechanism by which a photodegradable compound is stabilized is the transfer of a photonicallly-excited electron to a nearby molecule of the same or different species (*see, e.g.*, N.J. Turro, Modern Molecular Photochemistry, Chapter 9, Benjamin/Cummings Publ. Co., Menlo Park, California (1991)), however photochemical theory does not describe the observed phenomena. Though not intending to be bound by such a belief, the observed phenomena are believed to coincide with the electron transfer theory of Professor Rudolph A. Marcus of the California Institute of Technology, for which he received the 1992 Nobel Prize in Chemistry.

[0026] The dielectric constant of a solvent system is a preferred measure of polarity of a solvent system, for example because the dielectric constant is a measure of both inherent and inducible dipole moments. Other measures of polarity include, but are not limited to, the

64267-1444

induced and/or inherent (permanent) dipole moment (*e.g.*, in Debye units), the Dimroth-Reichardt E_T parameter, and ionizing power. See generally, C. Reichardt, "Solvents and Solvent Effects in Organic Chemistry" 2nd ed., Chap. 7: Empirical Parameters of Solvent Polarity, VCH Publishers, New York, New York, (1988). Moreover, a more detailed description of these methods of measuring the polarity of the compound or a series of compounds can be found in commonly assigned U.S. Patents 6,537,529 and 6,485,713. Examples of polar solvents include, but are not limited to diethylhexyl malate, dimethyl capramide, or a combination thereof.

[0027] Mathematically, photodegradation can be described by an exponential function. Thus, $Q(a)$, the absorbance after a radiation dose (*i.e.*, exposure to a quantity of radiation), can be described by the general equation (i),

$$Q(a) = Ae^{-kr} \quad (i)$$

wherein A is the original (pre-exposure) absorbance, e is the natural logarithm base, k is the rate constant of the photodecay, and r is the cumulative dose (*e.g.*, in MED units). Because the absorbance decreases as the cumulative dose increases (photodecay), the overall term $-k$ will be negative, and the greater the value of $-k$ (*i.e.*, closer to zero) and, thus, the lower the rate constant of photodecay, the lower is the rate of photodecay. For example, when $Q(a)$ is plotted on a log scale versus r on a linear scale, the function forms a straight line with a slope equal to $-k$.

[0028] Furthermore, it has been found that, for a set of photoactive compounds that includes a photodegradable compound (*e.g.* avobenzene), the rate constant of photodecay of the set of photoactive compounds can be described as a second-order function of the polarity, preferably the dielectric constant (*i.e.*, relative permittivity) of the filter set dissolved in the solvent system. Thus, for example, the rate constant of photodecay of a filter set that include one or more of a photoactive compound, can be described by the general equation (ii),

$$k = -(x\varepsilon^2 + y\varepsilon + z) \quad (ii)$$

wherein x , y , and z can be empirically determined. The dielectric constant at the theoretical minimum rate constant of photodecay $-k_{\min}$ described by formula (iii),

$$\varepsilon_{k \min} = \frac{-y}{2x} \quad (iii)$$

wherein x and y are defined as above.

[0029] The phenomena described above, coupled with the knowledge that, heretofore, sunscreen compositions have been formulated without specific regard to the relationship between polarity and photostability and, in newly-discovered fact, have had non-optimal polarities, forms the basis for at least one aspect of the compositions described herein.

[0030] A photoactive compound is one that responds to light photoelectrically. In the compositions disclosed herein, a photoactive compound is one that responds to UV radiation photoelectrically. For example, photoactive compounds that respond to UV radiation photoelectrically by rapid photo degradation can benefit highly from the compositions disclosed herein, even though the benefits of the compositions disclosed herein are not limited to such compounds. Photostability is a potential problem with all UV filters because they are deliberately selected as UV-absorbing molecules. In other applications, a photoactive compound may be a pigment or a dye (*e.g.*, a hydrophobic dye).

[0031] UV filters include compounds selected from the following categories (with specific examples) including: p-aminobenzoic acid, its salts and its derivatives (ethyl, isobutyl, glyceryl esters; p-dimethylaminobenzoic acid); anthranilates (o-aminobenzoates; methyl, menthyl, phenyl, benzyl, phenylethyl, linalyl, terpinyl, and cyclohexenyl esters); salicylates (octyl, amyl, phenyl, benzyl, menthyl (homosalate), glyceryl, and dipropyleneglycol esters); cinnamic acid derivatives (menthyl and benzyl esters, alpha-phenyl cinnamionitrile; butyl cinnamoyl pyruvate); dihydroxycinnamic acid derivatives (umbelliferone, methylumbelliferone, methylaceto-umbelliferone); camphor derivatives (3-benzylidene, 4-methylbenzylidene, polyacrylamidomethyl benzylidene, benzalkonium methosulfate, benzylidene camphor sulfonic acid, and terephthalylidene dicamphor sulfonic acid); trihydroxycinnamic acid derivatives (esculetin, methylesculetin, daphnetin, and the glucosides, esculin and daphnin); hydrocarbons (diphenylbutadiene, stilbene); dibenzalacetone; benzalacetophenone; naphtholsulfonates (sodium salts of 2-naphthol-3,6-disulfonic and of 2-naphthol-6,8-disulfonic acids); dihydroxy-naphthoic acid and its salts; o- and p-hydroxydiphenyldisulfonates; coumarin derivatives (7-hydroxy, 7-methyl, 3-phenyl); diazoles (2-acetyl-3-bromoindazole, phenyl benzoxazole, methyl naphthoxazole, various aryl benzothiazoles); quinine salts (bisulfate, sulfate, chloride, oleate, and tannate); quinoline derivatives (8-hydroxyquinoline salts, 2-phenylquinoline); hydroxy- or methoxy-substituted benzophenones; uric acid derivatives; vilouric acid derivatives; tannic acid and its derivatives; hydroquinone; and benzophenones (oxybenzone, sulisobenzene, dioxybenzone, benzoescorcinol, 2,2',4,4'-tetrahydroxybenzophenone, 2,2'-dihydroxy-4,4'-

dimethoxybenzophenone, octabenzene, 4-isopropylidibenzoylmethane, butylmethoxydibenzoylmethane, etocrylene, and 4-isopropyl-dibenzoylmethane).

[0032] Particularly useful are: 2-ethylhexyl *p*-methoxycinnamate, 4,4'-*t*-butyl methoxydibenzoylmethane, 2-hydroxy-4-methoxybenzophenone, octyldimethyl *p*-aminobenzoic acid, digalloyltriolate, 2,2-dihydroxy-4-methoxybenzophenone, ethyl 4-[bis(hydroxypropyl)]aminobenzoate, 2-ethylhexyl-2-cyano-3,3-diphenylacrylate, 2-ethylhexylsalicylate, glycerol *p*-aminobenzoate, 3,3,5-trimethylcyclohexylsalicylate, methylanthranilate, *p*-dimethylaminobenzoic acid or aminobenzoate, 2-ethylhexyl *p*-dimethylaminobenzoate, 2-phenylbenzimidazole-5-sulfonic acid, 2-(*p*-dimethylaminophenyl-5-sulfoniobenzoxazoic acid, and combinations thereof.

[0033] A sunscreen composition disclosed herein includes a dibenzoylmethane derivative. Preferred dibenzoylmethane derivatives include 2-methyldibenzoylmethane; 4-methyldibenzoylmethane; 4-isopropylidibenzoylmethane; 4-*tert*-butyldibenzoylmethane; 2,4-dimethyldibenzoylmethane; 2,5-dimethyldibenzoylmethane; 4,4'-diisopropylidibenzoylmethane; 4,4'-dimethoxydibenzoylmethane; 4-*tert*-butyl-4'-methoxydibenzoylmethane; 2-methyl-5-isopropyl-4'-methoxydibenzoylmethane; 2-methyl-5-*tert*-butyl-4'-methoxydibenzoylmethane; 2,4-dimethyl-4'-methoxydibenzoylmethane; 2,6-dimethyl-4-*tert*-butyl-4'-methoxydibenzoylmethane, and combinations thereof. The compositions disclosed herein preferably include a dibenzoylmethane derivative in a range of about 0.1% to about 25% by weight of the total weight of the composition.

[0034] In any embodiment of a sunscreen composition an additional photoactive compound can be added to the composition. Additional photoactive compounds can be selected from any of the UV-A filters, UV-B filters, and combinations thereof. In a cosmetically-acceptable sunscreen embodiment for use on human skin, a photoactive compound preferably is selected from approved (if regulated), cosmetically-acceptable UV-A filters, UV-B filters, and combinations thereof.

[0035] For example, for a product marketed in the United States, preferred cosmetically-acceptable photoactive compounds and concentrations (reported as a percentage by weight of the total cosmetic sunscreen composition) include: aminobenzoic acid (also called para-aminobenzoic acid and PABA; 15% or less), avobenzene (also called butyl methoxy dibenzoylmethane; 3% or less), cinoxate (also called 2-ethoxyethyl *p*-methoxycinnamate; 3% or less), dioxybenzone (also called benzophenone-8; 3% or less),

64267-1444

homosalate (15% or less), menthyl anthranilate (also called menthyl 2-aminobenzoate ; 5% or less), octocrylene (also called 2-ethylhexyl-2-cyano-3,3 diphenylacrylate; 10% or less), octyl methoxycinnamate (7.5% or less), octyl salicylate (also called 2-ethylhexyl salicylate ; 5% or less), oxybenzone (also called benzophenone-3; 6% or less), padimate O (also called octyl dimethyl PABA; 8% or less), phenylbenzimidazole sulfonic acid (water soluble; 4% or less), sulisobenzone (also called benzophenone-4; 10% or less), titanium dioxide (2.5% or less), triethanolamine salicylate (also called triethanolamine salicylate; 12% or less), and zinc oxide (25% or less).

[0036] Other preferred cosmetically-acceptable photoactive compounds and concentrations (percent by weight of the total cosmetic sunscreen composition) include diethanolamine methoxycinnamate (10% or less), ethyl-[bis(hydroxypropyl)] aminobenzoate (5% or less), glyceryl aminobenzoate (3% or less), 4-isopropyl dibenzoylmethane (5% or less), 4-methylbenzylidene camphor (6% or less), terephthalylidene dicamphor sulfonic acid (10% or less), and sulisobenzone (also called benzophenone-4, 10% or less).

[0037] For a product marketed in the European Union, preferred cosmetically-acceptable photoactive compounds and concentrations (reported as a percentage by weight of the total cosmetic sunscreen composition) include: PABA (5% or less), camphor benzalkonium methosulfate (6% or less), homosalate (10% or less), benzophenone-3 (10% or less), phenylbenzimidazole sulfonic acid (8% or less, expressed as acid), terephthalylidene dicamphor sulfonic acid (10% or less, expressed as acid), butyl methoxydibenzoylmethane (5% or less), benzylidene camphor sulfonic acid (6% or less, expressed as acid), octocrylene (10% or less, expressed as acid), polyacrylamidomethyl benzylidene camphor (6% or less), ethylhexyl methoxycinnamate (10% or less), PEG-25 PABA (10% or less), isooamyl p-methoxycinnamate (10% or less), ethylhexyl triazone (5% or less), drometrizole trielloxane (15% or less), diethylhexyl butamido triazone (10% or less), 4-methylbenzylidene camphor (4% or less), 3-benzylidene camphor (2% or less), ethylhexyl salicylate (5% or less), ethylhexyl dimethyl PABA (8% or less), benzophenone-4 (5%, expressed as acid), methylene bis-benzotriazolyl tetramethylbutylphenol (10% or less), disodium phenyl dibenzimidazole tetrasulfonate (10% or less, expressed as acid), bis-ethylhexyloxyphenol methoxyphenyl triazine (10% or less), methylene bisbenzotriazolyl tetramethylbutylphenol (10% or less, also called TINOSORB* M), and bisethylhexyloxyphenol methoxyphenyl triazine (10% or less, also called TINOSORB* S).

*Trade-mark

64267-1444

[0038] All of the above-described UV filters are commercially available. For example, suitable commercially-available organic UV filters are identified by trade name and supplier in Table 1 below:

TABLE I

CTFA Name	Trade Name	Supplier
benzophenone-3	UVINUL* M-40	BASF Chemical Co.
benzophenone-4	UVINUL* MS-40	BASF Chemical Co.
benzophenone-8	SPECTRA-SORB* UV-24	American Cyanamid
DEA-methoxycinnamate	BERNEL HYDRO	Bernel Chemical
ethyl dihydroxypropyl-PABA	AMERSCREEN* P	Amerchol Corp.
glyceryl PABA	NIPA G.M.P.A.	Nipa Labs.
homosalate	KEMESTER* HMS	Humko Chemical
menthyl anthranilate	SUNAROME UVA	Felton Worldwide
octocrylene	UVINUL* N-539	BASF Chemical Co.
octyl dimethyl PABA	AMERSCOL*	Amerchol Corp.
octyl methoxycinnamate	PARSOL* MCX	Bernel Chemical
PABA	PABA	National Starch
2-phenylbenzimidazole-5-sulphonic acid	EUSOLEX* 6300	EM Industries
TEA salicylate	SUNAROME* W	Felton Worldwide
2-(4-methylbenzildene)-camphor	EUSOLEX* 6300	EM Industries
benzophenone-1	UVINUL* 400	BASF Chemical Co.
benzophenone-2	UVINUL* D-50	BASF Chemical Co.
benzophenone-6	UVINUL* D-49	BASF Chemical Co.
benzophenone-12	UVINUL* 408	BASF Chemical Co.
4-isopropyl dibenzoyl methane	EUSOLEX* 8020	EM Industries
butyl methoxy dibenzoyl methane	PARSOL* 1789	Givaudan Corp.
etocrylene	UVINUL* N-35	BASF Chemical Co.
methylene bisbenzotriazolyl tetramethylbutylphenol	TINOSORB* M	Ciba Specialty Chemicals
bisethylhexyloxyphenol methoxyphenyl triazine.	TINOSORB* S	Ciba Specialty Chemicals

[0039] As used herein, the term "alkyl" includes straight chained and branched hydrocarbon groups containing the indicated number of carbon atoms, typically methyl, ethyl, propyl, and butyl groups. The term "alkyl" also includes "bridged alkyl," *e.g.*, a C₄-C₁₆ bicyclic or polycyclic hydrocarbon group, for example, norbornyl, adamantyl, bicyclo[2.2.2]octyl, bicyclo[2.2.1]heptyl, bicyclo[3.2.1]octyl, or decahydronaphthyl. The term "cycloalkyl" is defined as a cyclic hydrocarbon group, *e.g.*, cyclopropyl, cyclobutyl, cyclohexyl, and cyclopentyl.

*Trade-mark

[0040] As used herein, the term "alkenyl" is defined identically as "alkyl," except for containing a carbon-carbon double bond. The term "cycloalkenyl" is identical to "cycloalkyl" except containing a carbon-carbon double bond, *e.g.*, cyclopropyl, cyclobutyl, cyclohexyl, and cyclopentyl.

[0041] As used herein, the term "aryl," alone or in combination, is defined herein as a monocyclic or polycyclic aromatic group, preferably a monocyclic or bicyclic aromatic group, *e.g.*, phenyl or naphthyl.

[0042] A sunscreen composition disclosed herein may include a variety of photoactive compounds, including one or more UV-A photoactive compounds and one or more UV-B photoactive compounds. Preferably, a sunscreen composition includes a photoactive compound selected from the group consisting of p-aminobenzoic acid and salts and derivatives thereof; anthranilate and derivatives thereof; dibenzoylmethane and derivatives thereof; salicylate and derivatives thereof; cinnamic acid and derivatives thereof; dihydroxycinnamic acid and derivatives thereof; camphor and salts and derivatives thereof; trihydroxycinnamic acid and derivatives thereof; dibenzalacetone naphtholsulfonate and salts and derivatives thereof; benzalacetophenone naphtholsulfonate and salts and derivatives thereof; dihydroxy-naphthoic acid and salts thereof; o-hydroxydiphenyldisulfonate and salts and derivatives thereof; p-hydroxydiphenyldisulfonate and salts and derivatives thereof; coumarin and derivatives thereof; diazole derivatives; quinine derivatives and salts thereof; quinoline derivatives; hydroxy-substituted benzophenone derivatives; methoxy-substituted benzophenone derivatives; uric acid derivatives; vilouric acid derivatives; tannic acid and derivatives thereof; hydroquinone; benzophenone derivatives; 1,3,5-triazine derivatives, phenyldibenzimidazole tetrasulfonate and salts and derivatives thereof; terephthalylidene dicamphor sulfonic acid and salts and derivatives thereof; methylene bis-benzotriazolyl tetramethylbutylphenol and salts and derivatives thereof; bis-ethylhexyloxyphenol methoxyphenyl triazine and salts and derivatives thereof; diethylamino hydroxybenzoyl hexyl benzoate and salts and derivatives thereof; and combinations of the foregoing.

[0043] A preferred combination of photoactive compounds in a sunscreen composition includes a UV-A and a UV-B photoactive compound. However, when 2-ethylhexyl-*p*-methoxycinnamate is included in a mixture with a dibenzoylmethane derivative, the dibenzoylmethane derivative becomes particularly unstable. Without intending to be limited to any particular mechanism, it is believed that the cinnamate ester reacts with an excited-



state dibenzoylmethane derivative in a bimolecular pathway that renders both the dibenzoylmethane derivative and the cinnamate ester incapable of absorbing UV radiation.

[0044] It has been found, quite surprisingly, that a combination of (a) an α -cyano- β,β -diphenylacrylate compound and (b) a diester or polyester of naphthalene dicarboxylic acid in a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0, increases the stability of a sunscreen composition including 2-ethylhexyl-*p*-methoxycinnamate and a dibenzoylmethane derivative. Thus, one embodiment of a sunscreen composition includes the photoactive compound 2-ethylhexyl-*p*-methoxycinnamate, a dibenzoylmethane derivative, and a combination of (a) and (b) in weight ratios previously defined.

[0045] It has been found, quite surprisingly, that the addition of a methoxy-substituted benzophenone derivative to a sunscreen composition including a dibenzoylmethane derivative and a diester or polyester of naphthalene dicarboxylic acid results in an increase in the stability of the dibenzoylmethane derivative present in the composition. A methoxy-substituted benzophenone derivative has dual purposes in the sunscreen composition, both to act as a photoactive compound, and to increase the photostability (lower the rate constant of photodecay) of one or more photoactive compounds present in the sunscreen composition. Without intending to be limited to any particular mechanism, it is believed that a methoxy-substituted benzophenone derivative quenches (accepts) the singlet excited state of the diester or polyester of naphthalene dicarboxylic acid, and thereby prevents the excited diester or polyester from reaching the triplet excited state. Preferably, a sunscreen composition disclosed herein includes a methoxy-substituted benzophenone derivative such as benzophenone-3. The methoxy-substituted benzophenone derivative preferably is present in a sunscreen composition in an amount of 0.5% or less by weight of the total weight of the composition.

[0046] One embodiment of a sunscreen composition disclosed herein includes a mixture of a dibenzoylmethane derivative, and a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0, wherein (a) is an α -cyano- β,β -diphenylacrylate compound, and (b) is a diester or polyester of naphthalene dicarboxylic acid selected from the group consisting of formula (I), formula (II), and combinations thereof:

64267-1444



[0047] wherein R^1 and R^2 are the same or different and selected from the group consisting of a C_1 - C_{22} alkyl groups, residues of diols having the structure $\text{HO}-\text{R}^3-\text{OH}$, and residues of polyglycols having the structure $\text{HO}-\text{R}^4-(\text{O}-\text{R}^3)_n-\text{OH}$; wherein each R^3 and R^4 is the same or different and selected from the group consisting of C_1 - C_6 straight or branched chain alkylene groups; and wherein m and n are each in a range of 1 to 100 and p is in a range of 0 to 100. Although any α -cyano- β,β -diphenylacrylate compound may be used in this embodiment, preferably, the α -cyano- β,β -diphenylacrylate compound is 2-ethylhexyl-2-cyano-3,3-diphenylacrylate (also known as octocrylene).

[0048] The method of preparation of particularly useful diesters and polyesters of naphthalene dicarboxylic acid and the use of diesters and polyesters of naphthalene dicarboxylic acid in a sunscreen composition are described in U.S. Patent Nos. 5,993,789 and 6,284,916. Preferably, a composition of this embodiment includes a diester of formula (II) wherein R^1 and R^2 are 2-ethylhexane and p is 0. Preferably, the compositions disclosed herein include a diester or polyester of naphthalene dicarboxylic acid in a range of about 0.1% to about 15% by weight of the total weight of the composition.

[0049] As described above, the stability of photoactive compounds present in a sunscreen composition can be increased by controlling the polarity of the oil phase of the composition. Because prior sunscreen formulations have typically had lower than optimal polarities, adding a high-polarity component to the oil phase to raise the oil phase polarity improves the photostability of the photoactive compounds. Thus, preferably, a sunscreen composition includes an oil phase comprising a dibenzoylmethane derivative, (a) an α -cyano- β,β -diphenylacrylate compound, (b) a diester or polyester of naphthalene dicarboxylic acid, in a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0, and a solvent system, wherein the solvent system includes an effective amount of a polar solvent, or a blend

64267-1444

solvents with a high polarity, to increase the photostability of the dibenzoylmethane derivative or other photoactive compounds present in the sunscreen composition. Suitable polar solvents for use in a sunscreen composition are disclosed in commonly assigned U.S. Patent Publication No. 2003-18690 and U.S. Patent No. 6,485,713. A composition of this embodiment preferably has a dielectric constant of at least 7, preferably at least about 8.

[0050] Another embodiment of the sunscreen compositions disclosed herein includes a mixture of a dibenzoylmethane derivative, (a) an α -cyano- β,β -diphenylacrylate compound, (b) a diester or polyester of naphthalene dicarboxylic acid, in a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0, and (c) benzophenone-3 in an amount of about 0.1 to 10% by weight, preferably less than about 0.5 % by weight. Although any α -cyano- β,β -diphenylacrylate compound may be used according to this embodiment, preferably, the α -cyano- β,β -diphenylacrylate compound is 2-ethylhexyl-2-cyano-3,3-diphenylacrylate (also known as octocrylene).

[0051] Another embodiment of a sunscreen composition disclosed herein includes a mixture of a dibenzoylmethane derivative, (a) an α -cyano- β,β -diphenylacrylate compound, and (b) a diester or polyester of naphthalene dicarboxylic acid, in a weight ratio of (a)/(b) of at least 0.95, preferably at least about 1.0, wherein said sunscreen composition has a dielectric constant of at least about 8.

EXAMPLES

[0052] The following examples are provided to illustrate the invention but are not intended to limit the scope of the invention.

[0053] Two sunscreen compositions were produced by mixing the ingredients and concentrations (formulations) shown in the following two examples:

64267-1444

Example 1***In vivo* PFA 8.63**

Phase	INCI/CTFA Chemical Name	Trade Name, Supplier	% w/w
A.	Octisalate	(USP, RTD*HALL STAR)	5.00
	Homosalate	(NeoHeliopan* HMS, Symrise)	7.50
	Avobenzene	(Parsol 1789, Roche)	3.00
	Octocrylene	(NeoHeliopan* 303, Symrise)	2.50
	Diethylhexyl 2,6-naphthalate	(Symrise*)	2.50
	Dimethyl capramide	(Spectrasolv* DMDA, RTD*HALLSTAR)	1.00
	Diethylhexyl malate	(Spectrasolv* 16, RTD*HALLSTAR)	2.01
	Benzophenone-3	(NeoHeliopan* BB, Symrise)	0.49
B.	Polyglyceryl-3 distearate	(Cremorphor* GS-32, BASF)	3.00
	Sorbitan isostearate	(Crill* 6, Croda)	4.00
	Stearic acid	(V-1655, RTD*HALLSTAR)	3.05
	PVP/Eicosene copolymer	(Ganex* V-220, ISP)	2.00
	Dimethicone (100 cSt)	(Mirasil* DM100, RTD*HALLSTAR)	0.40
	Silica	(R972, Degussa)	0.25
C.	Deionized water	Water	Q.S.
	Disodium EDTA	Disodium EDTA	0.05
	Carbomer	(Carbopol* Ultrez 10, B.F. Goodrich)	0.05
D.	Methylpropanediol	(MPDiol, Lyondell)	2.00
	Glycerin	Glycerin	3.00
	Phenoxyethanol(methyl- paraben(ethylparaben(propyl- paraben(butylparaben	(CoSept* PEP, RTD*HALLSTAR)	0.60
	Triethanolamine (TEA)	Triethanolamine	1.74

Procedure

1. Oil Phase: Blend "A" liquid additives. Heat to 80°C, stirring to dissolve Oxybenzone, Avobenzene. Add "B" additives with stirring until clear and homogeneous. NOTE: Addition of Dimethicone will turn oil phase turbid. Add silica and stir to wet and disperse thoroughly.
2. Water Phase: Dissolve EDTA salt in water. Sprinkle carbomer on water and allow to wet completely. With stirring, heat to 75°C. Preblend preservative, Methylpropanediol, Glycerin, and TEA, and add to water.
3. With stirring, add oil ("A" + "B") to water ("C" + "D"). A crude yellowish emulsion will form. Remove from heat. Continue stirring until temperature falls below 75°C. Homogenize for three minutes to form a white, fine emulsion. Switch to sweep. Stir while cooling. Package when temperature drops below 35°C.
4. Don't forget to add back water lost during processing.

[0054] The following in-vivo PFA values were obtained from using the composition of Example 1 on five individuals, in accordance with the Japanese Pigment Darkening Protocol (Appendix):

*Trade-mark

64267-1444

Table II
Individual PFA Values

Subject	CPTC#	Skin Type	Age/Sex	<u>180 Minutes</u>	
				Standard	CAB4-269
1) JA	38973	IV	54/M	5.86	9.38
2) BB	38974	III	54/F	4.68	7.50
3) RA	388912	III	26/M	5.86	9.38
4) TF	40813	III	19/M	4.69	7.50
5) NF	12775	III	56/M	4.69	9.38
Average PFA (N=5)				5.16	8.63

Example 2

[0055] The following composition was produced and tested for photostability in the same manner described with reference to Example 1, above. The addition of 0.20% by weight diethylhexyl malate to the control was to maintain identical oil phase volumes in all formulations tested. The sunscreen compositions included very low ratios of octocrylene (OC) to DEHN (diester or polyester of naphthalene dicarboxylic acid) of 0.06 and 0.026.

***In vivo* PFA 7.27**

Phase	INCI/CTFA Chemical Name	Trade Name, Supplier	% w/w
A.	Octisalate	(USP, RTD*HALL STAR)	5.00
	Homosalate	(NeoHeliopan* HMS, Symrise)	7.25
	Avobenzone	(NeoHeliopan* 357, Symrise)	3.00
	Octocrylene	(NeoHeliopan* 303, Symrise)	0.26
	Diethylhexyl 2,6-naphthalate	(Symrise*)	10.0
	Dimethyl capramide	(Spectrasolv [®] DMDA, RTD*HALLSTAR	0.49
B.	Stearyl alcohol	(Alfol* 18, Sasol)	1.00
	Polyglyceryl-3 methyl glucose distearate	(TegoCare* 450, Goldschmidt)	3.00
	Steareth-21	(Brij* 721, Uniqema)	0.31
	Steareth-2	(Brij* 721, Uniqema)	0.19
	C30-38 Olefin/Isopropyl maleate/MA copolymer	(Performa* V 1608, New Phase)	2.00
C.	Deionized water	Water	Q.S.
	Disodium EDTA	Disodium EDTA	0.05
D.	Glycerin	Glycerin	4.00
	Phenoxyethanol(methylparaben(ethylparaben(propylparaben(butylparaben	(Phenonip*, Clariant)	0.60
E.	Carbomer	(Carbopol* Ultrez 10, B.F. Goodrich)	0.20
F.	Sodium hydroxide (25%)	Sodium hydroxide	0.28

*Trade-mark

Procedure

1. Oil Phase: In secondary vessel, blend "A" liquid additives. Heat to 80°C, stirring to dissolve Oxybenzone, Avobenzone. Increase heat to 90°C. Add "B" additives with stirring until clear and homogeneous.
2. Water Phase: In small vessel, pre-disperse carbomer ("E") in water (15-20 parts water to 1 part carbomer; allow to sit for 15-20 minutes) and set aside. Charge primary vessel with water (less amount of water used to pre-disperse carbomer). Dissolve Disodium EDTA in water. Heat to 85°C. Pre-blend preservative, Glycerin and add to water. Stir to disperse.
3. With homogenization, add oil ("A" + "B") to water ("C" + "D"). While continuing to homogenize, add pre-dispersed carbomer ("E"). Maintain heat at 80-90°C, homogenize for 2 minutes.
4. Remove from heat. Stir with sweep while cooling. When temperature is below 40°C slowly neutralize with Sodium hydroxide ("F") solution to pH 6-6.5. Continue stirring to smooth, homogenous lotion. NOTE: Be sure to add back any water lost during processing.

APPENDIX 3: JCIA/PERSISTENT PIGMENT DARKENING PROTOCOL

WO 2005/082325 **Japan Cosmetic Industry Association** PCT/US2004/040103
Measurement Standards for UVA Protection Efficacy

Introduction

The damaging effects of UV rays on the skin have become widely recognized by consumers, and there have been reports in the media warning of increases in the level of UV rays due to environmental pollution. This has resulted in the appearance of many "Cosmetics with UV Protection" on the market. These products protect the skin by reducing or blocking the effects of UV rays.

Cosmetics with UV Protection can be roughly divided into two groups. One is "Suntan Cosmetics" which are used for the purpose of obtaining a beautifully suntanned complexion while limiting the affects of UV rays on the skin to a minimum, the other is "Sunscreen Cosmetic" which are used for the purpose of preventing Sunburn and Suntan.

UV rays that reach the surface of the earth can be divided into the A region of UV light (UVA: 320-400 nm) and the B region of UV light (UVB: 280-320 nm), and these two types of UV rays have different effects on the skin. UVB causes erythema of the skin several hours after exposure, and several days after exposure to UVB may lead to increased pigmentation, dryness and scale. UVA causes darkening of the skin immediately after exposure (immediate pigment darkening) and in the event of exposure to large amounts of UVA, this darkening appears to be transformed to delayed pigment darkening. There are also reports that UVA increases sensitivity of the skin to UVB. In addition to these acute responses, UV rays contribute to skin cancer and to aging of the skin typified by blotches and wrinkles. The relative contributions of UVB and UVA to these various reactions are not known, but the effects of the deep penetration of UVA rays cannot be ignored.

Under these circumstances, the expression "UV Protection" (in product claims) is not always adequate, so there is a need to clarify whether a product protects against UVA or UVB, and to what extent it protects against each.

SPF is an index of the protection against sunburn provided by cosmetics with UV protection. The SPF value is determined in accordance with the Japan Cosmetic Industry Association SPF Measurement Standards (Effective from January 1992) in Japan, in accordance with COLIPA regulations (October 1994) in Europe, and in accordance with FDA regulations (Tentative Final Monograph, May 1993) in the US. Because these methods are quite similar, their SPF values are roughly comparable even though a uniform method is not employed worldwide. Throughout the world the SPF value acts as an index that consumers use in product selection.

With respect to an index or measurement methods for UVA protection, however, a uniform measurement method has not yet been established on a national or industry-wide level although several papers on the subject have been published, and studies are underway in various countries. Throughout the world there are products displaying numerical values, etc., for UVA protection, but because there is particular concern that a uniform measurement method has not been established in Japan and these numerical values may cause confusion among consumers in their product selection, it has been decided to employ some types of index to list the level of UVA protection on cosmetic products.

Therefore, for the purpose of establishing a method for measuring UVA protection, the Technical Committee of the Japan Cosmetic Industry Association reorganized its previous SPF task force in November 1992 and established the Ultra Violet task force. This task force has handled the basic research project on UV protection approved and project on UV protection approved and

sponsored by WO 2005/082325 Human Science Foundation and has compiled its results in this document. PCT/US2004/040103

The fundamental principles toward the standard are described below:

(1) The standard is intended to provide uniform measurement method of WA (Protection Factor of WA) values and labeling method for the grade of UVA protection on sunscreen and suntan cosmetics enable consumers to select products which meet consumers desired UV light protection efficacy.

(2) The standard shall go into effect on January 1, 1996.

(3) The standard shall be reviewed when new technological findings warrant it

The standard consist of "I. Measurement Method of UVA Protection efficacy," listing itemized measurement conditions and "U. Annotation" providing practical points catty out tests using this method.

I. PFA Measurement Method

1. Selection of Test Subjects and Test Sites (Annotation I)

(1) Subjects must be healthy males and females at least 18 years old and belong to Skin Type II, III or IV mentioned below.

Subjects must be asked of their physical conditions and must be excluded if they have photodermatitis or take medicine (such as anti-inflammatory agent, anti-hypertensive agent etc.) relating to skin's photosensitivity.

Skin Type:	I.	Always bums easily; never tans
	II.	Always burns easily; tans minimally
	III.	Bums moderately; tans gradually
	IV.	Burns minimally; always tans well
	V.	Rarely bums; tans profusely
	VI.	Never bums; deeply pigmented

The skin types are classified based on the typical skin reactions to 30 to 45 minutes sun bathing after a winter season of no sun exposure.

(2) Test site is the back, and the skin must have almost uniform color without pigmentation, nevus and so

2. Number of Subjects (Annotation 2)

Each test must be performed with at least 10 subjects and the standard error for measuring PFA shall not exceed 10% of obtained PFA value.

The standard sample shall be prepared according to the formula described below. Measurement of the standard sample shall be performed concurrently with the measurement of the test sample. Formula and Preparation method of the Standard sample (a cream containing 5% 4-tert-Butyl-4'-methoxydibenzoylmethane and 3% 2-Ethylhexyl p-methoxycinnamate).

		%by Weight
A1	Purified Water (JSCI)	57.13
A2	Dipropylene Glycol (JSCI)	5.00
A3	Potassium Hydroxide (JSCI)	0.12
A4	Trisodium Edetate (JSC1)	0.05
A5	Phenoxyethanol (JScQ	0.3
B1	Stearic Acid (JSCI)	3.0
B2	(3)glyceryl Monostearate, Self-emulsifying (JSCI)	3.0
B3	Catostearyl Alcohol (ISCI)	5.0
B4	Petrolatum (JSCJ~	3.0
B5	Glyceryl Tri-Z-ethylhexanoate (JCIC)	15.0
B6	2-Ethylhexyl p-Methoxycinnamate (JSCI)	3.0
B7	4-tert-Butyl-4'-Metttoxydibenzoylmethane (JSCI)	5.0
B8	Ethyl Parahydroxybenzoate (JSCI)	0.2
B9	Methyl Parahydroxybenzoate (JSC1)	0.2

Weigh out each of the ingredients in A, dissolve them in the purified water, and heat the solution to 70° C. Weigh out each of the ingredients in B and heat them to 70°C so that they dissolve completely. Add B to A, emulsify the mixture, and adjust the size of the emulsified particles with a homogenizer, etc. Cool the emulsion to obtain the standard sample.

4. Amount of the Samples to be Applied (Annotation 4)

The amount of the samples to be applied shall be 2 mg/cm² or 2 ul/cm² each.

5. Area of the Samples to be Applied (Annotation 5)

The area (for applying samples) shall be at least 20 cm².

6. Time from Application to Exposure

Radiation exposure shall begin at lest 15 minutes after the samples are applied.

An artificial light shall be used as a source of light, which must satisfy following conditions...

(1) The UV light emission is UVA range shall have a continuous spectrum similar to sun light. Moreover, the ratio of UVA I (340-400 nm) and UVA II (320-340 nm) shall be close to that of sunlight (UVA II/UVA-8-20%).

(2) To avoid extreme sunburn, UV ray shorter than 320 nm shall be excluded through the use of an appropriate filter. Monitoring and maintenance shall be performed to insure That the above conditions are always maintained.

8. Radiation Field (Annotation 7)

A single radiation field shall be at least 0.5 cm². The radiation fields of the untreated area shall be equivalent to the radiation field of the treated area.

9. Progression of UV Dose (Annotation 8)

A UV dose progresses geometrically and the increment shall be 25% maximum.

10. MPPD (Minimal persistent Pigment Darkening Dose) (Annotation 9)

The MPPD is defined as the minimum dose of UV rays that produces slight darkening over essentially the whole radiation field within 2 to 4 hours after exposure. Determination of MPPD shall be conducted in a room with sufficient lighting at a fixed time within 2 to 4 hours after the end of exposure. At least two trained evaluators are desired to read MPPD).

11. Calculation Method of the PFA Value (Annotation 10)

PFA value shall be obtained from the following equation by using MPPDs at sites untreated and treated by a test sample.

$$\text{PFA value} = \text{MPPD in protected skin} / \text{MPPD in unprotected skin}$$

PFA value of a test sample is defined as the arithmetic mean of each subject's PFA values obtained from the above equation.

12. Method for Expressing UVA Protection (Annotation 11)

For labeling PFA values in UV protecting products, the figures to the right of the decimal point shall be discarded from the PFA value of the sample that has been calculated according to the above method to make it an integer. Then, when the value is not less than 2, it shall be classified according to the following PA (Protection grade of UVA), and this classification shall be expressed on the label. PA shall be placed together with the SPF value.

2 or more but less than 4	PA+
4 or more but less than 8	PA++
8 or more	PA+++

II. Annotations**(Annotation 1) Selection of Subjects and Test Sites**

For the PFA value measurement to be an index of the persistent pigment darkening induced by UVA, it must be assumed that a stable darkening occurs from exposure to UV rays. If the skin color is dark, the determination of this reaction is very difficult. Moreover, as shown in Figure 1, no significant differences were found between PFA values obtained from skin types II, III, and IV. Therefore, skin types II, III, and IV have been stipulated. Results of a skin type survey of Japanese people conducted by the Japan Cosmetic Industry Association show that approximately 74% of Japanese belong to skin types II, III, and IV.

The subjects shall be selected from applicants who understand the objective of the test by interview questions in the "Questionnaire for subject Selection". Examples of drugs that contribute to photosensitivity include the following:

Hypotensive agents, psychotropic agents, tranquilizers, antihistamines, oral hypoglycemic agents, and tetracycline antibiotics,

Because it is necessary for the test subjects to evaluate their own skin types and agree to being a test subject, the minimum age for test subjects was set at 10 and above. It is also recommended that subjects not be older than 60 years old.



64267-1444

CLAIMS:

1. A sunscreen composition, comprising a mixture of a dibenzoylmethane derivative selected from the group consisting of 2-methyldibenzoylmethane; 4-methyldibenzoylmethane;
 5 4-isopropyldibenzoylmethane; 4-tert-butyldibenzoylmethane; 2,4-dimethyldibenzoylmethane; 2,5-dimethyldibenzoylmethane; 4,4'-diisopropyldibenzoylmethane; 4,4'-dimethoxydibenzoylmethane; 4-tert-butyl-4'-methoxydibenzoylmethane; 2-methyl-5-isopropyl-
 10 4'-methoxydibenzoylmethane; 2-methyl-5-tert-butyl-4'-methoxydibenzoylmethane; 2,4-dimethyl-4'-methoxydibenzoylmethane; 2,6-dimethyl-4-tert-butyl-4'-methoxydibenzoylmethane, and combinations thereof, with:

(a) octocrylene; and

- 15 (b) a diester or polyester of naphthalene dicarboxylic acid selected from the group consisting of formula (I), formula (II), and combinations thereof:



wherein:

- 20 R^1 and R^2 are the same or different and each is selected from the group consisting of a C_1 - C_{22} alkyl group, a residue of a diol having the structure $\text{HO}-\text{R}^3-\text{OH}$, and a

64267-1444

residue of a polyglycol having the structure
 $\text{HO-R}^4\text{-(O-R}^3\text{)}_n\text{-OH}$;

each R^3 and R^4 is the same or different and each is
selected from the group consisting of a $\text{C}_1\text{-C}_6$ straight or
5 branched chain alkylene group;

m and n are each in a range of 1 to 100 and p is
in a range of 0 to 100; and

the weight ratio of (a)/(b) is at least 0.95.

2. The composition of claim 1, wherein the weight
10 ratio of (a) to (b) is at least about 1.0.

3. The composition of claim 1 or 2, further
comprising benzophenone-3.

4. The composition of claim 3, wherein the
benzophenone-3 is contained in an amount of about 0.1% by
15 weight to about 10% by weight of the sunscreen composition.

5. The composition of claim 3, wherein the
benzophenone-3 is present in an amount of 0.5% or less by
weight of the total weight of the composition.

6. The composition of claim 3, wherein the
20 benzophenone-3 is included in the sunscreen composition in
an amount less than about 0.5% by weight.

7. The composition of any one of claims 1 to 6,
wherein the dibenzoylmethane derivative is present in an
amount of about 0.1% to about 25% by weight of the total
25 weight of the composition.

64267-1444

8. The composition of any one of claims 1 to 7, wherein the octocrylene is present in an amount of at least about 0.5% by weight of the total weight of the composition.

9. The composition of any one of claims 1 to 8,
5 wherein the octocrylene is present in an amount of about 1.0% to about 8% by weight of the total weight of the composition.

10. The composition of any one of claims 1 to 9, comprising the diester of formula (II) wherein R^1 and R^2 are
10 2-ethylhexane and p is 0.

11. The composition of any one of claims 1 to 10, wherein the diester or polyester of naphthalene dicarboxylic acid is present in a range of about 0.1% to about 15% by weight of the total weight of the composition.

15 12. The composition of any one of claims 1 to 11, wherein the weight ratio of (a) to (b) is in the range of about 1/1 to about 2/1.

13. The composition of any one of claims 1 to 12, further comprising a photoactive compound selected from the
20 group consisting of p-aminobenzoic acid, salts thereof; anthranilate; salicylate; cinnamic acid; dihydroxycinnamic acid; camphor, derivatives thereof; trihydroxycinnamic acid; dibenzalacetone; naphtholsulfonate, salts thereof; benzalacetophenone; dihydroxy-naphthoic acid, salts thereof;
25 o-hydroxydiphenyldisulfonate, salts thereof; p-hydroxydiphenyldisulfonate, salts thereof; coumarin; tannic acid; hydroquinone; phenyldibenzimidazole tetrasulfonate, salts thereof; terephthalylidene dicamphor sulfonic acid, salts thereof; methylene bis-benzotriazolyl
30 tetramethylbutylphenol, salts thereof;

64267-1444

bis-ethylhexyloxyphenol methoxyphenyl triazine, salts thereof; diethylamino hydroxybenzoyl hexyl benzoate, salts thereof; and combinations of the foregoing.

14. The composition of claim 13, wherein the
5 photoactive compound comprises 2-ethylhexyl-
p-methoxycinnamate.

15. A use of the composition as defined in any one of
claims 1 to 14 in the reduction of contact of UV radiation
with human skin.

10 16. The use of claim 15, wherein the composition is
formulated for application to the human skin.