

ORGANISATION AFRICAINE DE LA PROPRIETE INTELLECTUELLE
(O.A.P.I.)



(19)

(11) N°

11538

(51) Inter. Cl.⁷

A61K 31/00

BREVET D'INVENTION

(21) Numéro de dépôt : 1200000279

(22) Date de dépôt : 13.04.1999

(30) Priorité(s) : DE
14.04.1998 N° 19816196.4
09.06.1998 N° 19825585.3
22.09.1998 N° 19843222.4
22.09.1998 N° 19843223.2

(24) Délivré le : 13.08.2001

(45) Publié le : 24 MAI 2004

(73) Titulaire(s) :

Hassan JOMAA
Breslauer Strasse 24
D-35398 GIESSEN
(DE)


(72) Inventeur(s) : (Le Titulaire)

(74) Mandataire : CABINET J. EKEME
B.P. 6370
YAOUNDE - Cameroun

(54) Titre : Use of organophosphoric compounds for the therapeutic and preventative treatment of infections.

(57) Abrégé :

Use of organophosphorus compounds of general Formula (I)



(I)

for the therapeutic and prophylactic treatment of infections in humans and animals caused by viruses, fungi and parasites.

Use of organophosphorus compounds for the therapeutic and prophylactic treatment of infections

The invention relates to the use of organophosphorus compounds and their salts, esters and amides for the therapeutic and prophylactic treatment of infections which are caused by viruses, fungi and parasites in humans and animals. According to the invention the organophosphorus compounds comprise phosphonic acid derivatives, phosphinoyl derivatives and phosphinic acid derivatives.

The suitability of aminohydrocarbyl phosphonic acid derivatives and some of their esters and salts for use in pharmaceutical compositions is already known. Up until now however only their antimicrobial efficacy against bacteria in humans and animals and against fungi in plants has been described (DE 27 33 658 A1, US 4 143 135, US 4 182 758 and US 4 206 156, US 4 994 447, US 4 888 330, US 4 210 635, US 3 955 958, US 4 196 193, US 4 268 503, US 4 330 529, US 5 189 030, US 3 764 677, US 3 764 676). Furthermore substances from this group have been described as herbicides (US 4 693 742, US 5 002 602, US 4 131 448, US 3 977 860, US 4 062 669), as algaecides (US 3 887 353), as means for regulating plant growth (US 4 127 401, US 4 120 688, US 3 961 934, US 4 431 438, US 3 853 530, US 4 205 977, US 4 025 332, US 3 894 861) and as reagents in dye production (US 4 051 175). In DE 27 33 658 A1 the use of aminohydrocarbyl phosphonic acid derivatives for the treatment of bacterial diseases has been described. Whilst the document speaks of a microbial efficacy against pathogenic microorganisms in the introduction to the description, from the overall context however it is clear that the invention relates exclusively to bacteria. Thus on page 16, second paragraph "antimicrobial efficacy" is defined as "antibacterial efficacy".

There is a serious need for the provision of preparations to enhance the treatment of humans and animals and the protection of plants, which preparations not only possess a strong efficacy but in contrast to other pharmaceutical compositions and plant protective agents, contain reduced side effects and therefore represent a lower risk to health for humans.

The object of the present invention is therefore to provide a substance which fulfils the conditions given above in the case of infections caused by viruses, fungi and parasites in humans and animals.

This object is achieved in a completely surprising manner by the group of substances defined in claim 1. This group of substances demonstrates an anti-infectious effect against viruses, fungi and unicellular and multicellular parasites. In the context of this invention the strictly scientific definition of parasites is to be used. This means that unicellular parasites are understood to mean protozoa exclusively.

The organophosphorus compounds used according to the invention correspond to the general Formula (I):



in which R_1 and R_2 are the same or different and are selected from the group which consists of hydrogen, substituted and unsubstituted alkyl, substituted and unsubstituted hydroxyalkyl, substituted and unsubstituted alkenyl, substituted and unsubstituted alkynyl, substituted and unsubstituted aryl, substituted and unsubstituted acyl, substituted and unsubstituted cycloalkyl, substituted and

unsubstituted aralkyl, substituted and unsubstituted heterocyclic radical, halogen, OX_1 and OX_2 , wherein X_1 and X_2 can be the same or different and are selected from the group which consists of hydrogen, substituted and unsubstituted alkyl, substituted and unsubstituted hydroxyalkyl, substituted and unsubstituted alkenyl, substituted and unsubstituted alkynyl, substituted and unsubstituted aryl, substituted and unsubstituted acyl, substituted and unsubstituted cycloalkyl, substituted and unsubstituted aralkyl, substituted and unsubstituted heterocyclic radicals,

A is selected from the group which consists of an alkylene radical, an alkenylene radical and a hydroxyalkylene radical,

R_3 and R_4 are selected independently from the group which consists of hydrogen, substituted and unsubstituted C_{1-26} -alkyl, substituted and unsubstituted hydroxy- C_{1-26} -alkyl, substituted and unsubstituted aryl, substituted and unsubstituted acyl, substituted and unsubstituted aralkyl, substituted and unsubstituted C_{1-26} -alkenyl, substituted and unsubstituted C_{1-26} -alkynyl, substituted and unsubstituted cycloalkyl, substituted and unsubstituted heterocyclic radical, halogen, OX_3 and OX_4 ,

wherein X_3 and X_4 are selected independently from the group which consists of hydrogen, substituted and unsubstituted C_{1-26} -alkyl, substituted and unsubstituted hydroxyl- C_{1-26} -alkyl, substituted and unsubstituted aryl, substituted and unsubstituted aralkyl, substituted and unsubstituted C_{1-26} -alkenyl, substituted and unsubstituted C_{1-26} -alkynyl, substituted and unsubstituted cycloalkyl, substituted and unsubstituted heterocyclic radical, a silyl, a cation of an organic and inorganic base, in particular a metal of the first, second or third main group of the periodic system, ammonium,

substituted ammonium and ammonium compounds which derive from ethylene diamine or amino acids,

and their pharmaceutically acceptable salts, esters and amides and salts of esters.

- 5 Phosphonic acid derivatives are particularly preferred.

In particular compounds which contain the following Formula (II) are suitable:



wherein,

- 10 X_1 is selected from the group which consists of hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkyl, substituted and unsubstituted heterocyclic radical;

R_2 , R_3 , R_4 and A contain the same meaning as in Formula (I).

- 15 A is particularly preferably a chain of three carbon atoms which bind the nitrogen atom to the phosphorus atom.

- 20 Compounds of Formula (II) are particularly preferred in which R_2 = acyl, in particular an acetyl, R_3 = hydrogen, methyl or ethyl, R_4 = hydrogen, methyl, ethyl or OX_4 with X_4 = hydrogen, sodium, potassium, methyl, ethyl, X_1 = H and A = alkylene, alkenylene or hydroxyalkylene. Particularly good results are achieved with R_2 = formyl or acetyl and A = propylene, propenylene or hydroxypropylene.

Furthermore compounds are particularly preferred in which R_1 is an alkyl, hydroxyalkyl, alkynyl, or alkenyl group with 16 or 18 carbon atoms or OX_3 , wherein X_3 is an alkyl, alkynyl, hydroxyalkyl or alkenyl group with 16 or 18 carbon atoms, R_4 is an alkyl, alkynyl, hydroxyalkyl or alkenyl group with 16 or 18 carbon atoms or OX_4 , wherein X_4 is an alkyl, alkynyl, hydroxyalkyl or alkenyl group with 16 or 18 carbon atoms.

Special features of the above definitions and suitable examples thereof are given below:

- 10 "Acyl" is a substituent which originates from an acid such as from an organic carboxylic acid, carbonic acid, carbamic acid or the thioacid or imidic acid corresponding to the individually present acids, or from an organic sulphonic acid, wherein these acids comprise in each case aliphatic, aromatic and/or heterocyclic groups in the molecule and carbamoyl or carbamimidoyl.

Suitable examples of these acyl groups will be given below.

Acyl radicals originating from an aliphatic acid are designated as aliphatic acyl groups and include:

- 20 Alkanoyl (for example formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl etc.);

Alkenoyl (for example acryloyl, methacryloyl, crotonoyl etc.);

Alkylthioalkanoyl (for example methylthioacetyl, ethylthioacetyl etc.);

- 25 Alkane sulphonyl (for example mesyl, ethane sulphonyl, propane sulphonyl etc.);

011538

Alkoxy carbonyl (for example methoxy carbonyl, ethoxy carbonyl, propoxy carbonyl, isopropoxy carbonyl, butoxy carbonyl, isobutoxy carbonyl etc.);

Alkyl carbamoyl (for example methyl carbamoyl etc.);

- 5 (N-alkyl)-thiocarbamoyl (for example (N-methyl)-thiocarbamoyl etc.);

Alkyl carbamimidoyl (for example methyl carbamimidoyl etc.);

Oxalo;

- 10 Alkoxalyl (for example methoxalyl, ethoxalyl, propoxalyl etc.).

In the above examples of aliphatic acyl groups the aliphatic hydrocarbon part, in particular the alkyl group and the alkane radical can optionally contain one or more suitable substituents such as amino, halogen (for example fluorine, chlorine, bromine etc.), hydroxy, hydroxyimino, carboxy, alkoxy (for example methoxy, ethoxy, propoxy etc.), alkoxy carbonyl, acylamino (for example benzyloxy carbonylamino etc.), acyloxy (for example acetoxy, benzoyloxy etc.) and the like. Preferred aliphatic acyl radicals with such substitutes are for example alkanoyls substituted with amino, carboxy, amino and carboxy, halogen, acylamino or the like.

25 Acyl residues originate from an acid with substituted or unsubstituted aryl groups, wherein the aryl group can comprise phenyl, toluyl, xylyl, naphthyl, and the like are designated as aromatic acyl radicals. Suitable examples are given below:

Aroyl (for example benzoyl, toluoyl, xyloyl, naphthoyl, phthaloyl etc.);

Aralkanoyl (for example phenylacetyl etc.);

Aralkenoyl (for example cinnamoyl etc.);

Aryloxyalkanoyl (for example phenoxyacetyl etc.);

Arylthioalkanoyl (for example phenylthioacetyl etc.);

Arylaminoalkanoyl (for example N-phenylglycyl, etc.);

- 5 Arene sulphonyl (for example benzene sulphonyl, tosyl toluene sulphonyl, naphthalene sulphonyl etc.);

Aryloxycarbonyl (for example phenoxycarbonyl, naphthyl-oxy carbonyl etc.);

Aralkoxycarbonyl (for example benzyloxycarbonyl etc.);

- 10 Arylcarbamoyl (for example phenylcarbamoyl, naphthylcarbamoyl etc.);

Arylglyoxyloyl (for example phenylglyoxyloyl etc.).

- 15 In the above-mentioned examples of aromatic acyl radicals the aromatic hydrocarbon part (in particular the aryl radical) and/or the aliphatic hydrocarbon part (in particular the alkane radical) can optionally contain one or more suitable substituents such as those which were already mentioned as suitable substituents for the alkyl group and the alkane radical. In particular and as an example of preferred aromatic acyl radicals with particular substituents, aroyl substituted by halogen and hydroxy or by halogen and acyloxy and aralkanoyl substituted by hydroxy, hydroxyimino, dihalogenalkanoyloxyimino are mentioned as well as
- 20

Arylthiocarbamoyl (for example phenylthiocarbamoyl etc.);

- 25 Arylcarbamimidoyl (for example phenylcarbamimidoyl etc.).

011538

A heterocyclic acyl radical is understood to be an acyl radical which originates from an acid with hererocyclic group. These include:

5 Heterocyclic carbonyl, in which the heterocyclic radical is an aromatic or aliphatic 5 to 6 membered heterocycle with at least one heteroatom from the group nitrogen, oxygen and sulphur (for example thiophenyl, furoyl, pyrrolcarbonyl, nicotinoyl etc.);

10 Heterocyclic alkanoyl in which the heterocyclic radical is 5 to 6 membered and has at least one heteroatom from the group nitrogen, oxygen and sulphur (for example thiophenylacetyl, furylacetyl, imidazolylpropionyl, tetrazolylacetyl, 2-(2-amino-4-thiazolyl)-2-methoxyiminoacetyl etc.) and the like.

15 In the above examples of heterocyclic acyl radicals the heterocycles and/or the aliphatic hydrocarbon part can optionally contain one or more suitable substituents, such as the same as those which were mentioned as suitable for alkyl and alkane groups.

20 "Alkyl" is a straight- or branched-chain alkyl radical with up to 9 carbon atoms, unless defined otherwise, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl, hexyl and the like.

25 "Hydroxylalkyl" is a straight- or branched-chain alkyl radical with up to 9 carbon atoms, unless defined otherwise, which contains at least one hydroxyl group, preferably one or two hydroxyl groups.

"Alkenyl" includes straight- or branched-chain alkenyl groups with up to 9 carbon atoms, unless defined otherwise, such as, for example, vinyl, propenyl, (for example 1-propenyl, 2-


011538

propenyl), 1-methylpropenyl, 2-methylpropenyl, butenyl, 2-ethylpropenyl, pentenyl, hexenyl.

"Alkynyl" includes straight- or branched-chain alkynyl groups with up to 9 carbon atoms, unless defined otherwise.

- 5 Cycloalkyl preferably represents an optionally substituted C3 to C7 cycloalkyl. Alkyl, alkenyl, alkynyl, alkoxy (for example methoxy, ethoxy etc.), halogen (for example fluorine, chlorine, bromine etc.), nitro and the like are *inter alia* suitable as possible substituents.
- 10+ Aryl is an aromatic hydrocarbon radical such as phenyl naphthyl etc., which can optionally contain one or more suitable substituents such as alkyl, alkenyl, alkynyl, alkoxy (for example methoxy, ethoxy etc.), halogen (for example fluorine, chlorine, bromine etc.), nitro and the like.
- 15 "Aralkyl" includes mono-, di-, triphenylalkyls such as benzyl, phenethyl, benzhydryl, trityl and the like, wherein the aromatic part can optionally contain one or more suitable substituents such as alkoxy (for example methoxy, ethoxy etc.), halogen (for example fluorine, chlorine, bromine etc.), nitro and the like.
- 20


"Alkylene" includes straight- or branched-chain alkylene groups which contain up to 9 carbon atoms and can be reproduced by the Formula



- 25 in which n is an integer from 1 to 9, such as methylene, ethylene, trimethylene, methylethylene, tetramethylene, 1-methyltrimethylene, 2-ethylethylene, pentamethylene, 2-methyltetramethylene, isopropylethylene, hexamethylene, and the like. Preferred alkylene radicals contain up to 4 carbon


atoms and radicals with 3 carbon atoms such as, for example, trimethylene are particularly preferred. The hydrogen atoms can be replaced by other substituents such as, for example, halogen radicals.

- 5 "Alkenylene" includes straight- or branched-chain alkenylene groups with up to 9 carbon atoms which can be reproduced by the Formula



- 10 in which n is an integer from 2 to 9, such as, for example, vinylene, propenylene (for example 1-propenylene, 2-propenylene), 1-methylpropenylene, 2-methylpropenylene, butenylene, 2-ethylpropenylene, pentenylene, hexenylene and the like. The alkenylene radical can particularly preferably contain up to 5 carbon atoms and in particular 3 carbon atoms
15 such as, for example, 1-propenylene. The hydrogen atoms can be replaced by other substituents such as, for example, halogen radicals.

- 20 "Hydroxyalkylene" can include straight- or branched-chain alkylene radicals which contain up to 9 carbon atoms, wherein at least one selected carbon atom is substituted by a hydroxy group. These radicals can be reproduced by the Formula



- 25 in which n is an integer from 1 to 9 and z is an integer, to which $1 \leq z \leq n$ applies. Suitable examples of such hydroxyalkylene groups include hydroxymethylene, hydroxyethylene (for example 1-hydroxyethylene and 2-hydroxyethylene), hydroxytrimethylene (for example 1-hydroxytrimethylene, 2-hydroxytrimethylene and 3-hydroxytrimethylene), hydroxy-tetramethylene (for example
30 2-hydroxytetramethylene), 2-hydroxy-2-methyltrimethylene,

hydroxypentamethylene (for example 2-hydroxypentamethylene), hydroxyhexamethylene (for example 2-hydroxyhexamethylene) and the like. A lower hydroxyalkylene with up to 4 carbon atoms is particularly preferred and in particular one with 3 carbon atoms such as, for example, 2-hydroxytrimethylene. The hydrogen atoms can be replaced by other substituents such as, for example, halogen radicals.

The radicals X_3 and X_4 can preferably be selected such that esters form on the phosphinic group or phosphonic group. Suitable examples of such esters in accordance with the Formulae (I) and (II) include alkyl esters (for example methyl ester, ethyl ester, propyl ester, isopropyl ester, butyl ester, isobutyl ester, hexyl ester, hexadecanyl ester, octadecanyl ester etc.);

Aralkyl esters (benzyl ester, phenethyl ester, benzohydryl ester, trityl ester etc.);

Aryl esters (for example phenyl ester, tolyl ester, naphthyl ester etc.);

Aroylalkyl esters (for example phenacyl ester etc.); and silyl esters (for example from trialkylhalogensilyl, dialkyldihalogensilyl, alkyltrihalogensilyl, dialkylarylhalogensilyl, trialkoxyhalogensilyl, dialkylaralkylhalogensilyl, dialkoxydihalogensilyl, trialkoxyhalogensilyl etc.) and the like.

With the above esters the alkane and/or arene part can contain at least one suitable substituent as desired such as halogen, alkoxy, hydroxy, nitro or the like.

X_3 and X_4 are preferably a metal from the first, second or third main group of the periodic system, ammonium, substituted ammonium or ammonium compounds which derive from

011538

ethylenediamine or amino acids. In other words the salt compounds of organophosphorus compounds are formed with organic or inorganic bases (for example sodium salt, potassium salt, calcium salt, aluminium salt, ammonium salt, magnesium salt, triethylamine salt, ethanolamine salt, dicyclohexylamine salt, ethylenediamine salt, N,N'-dibenzylethylenediamine salt etc.) and salts with amino acids (for example arginine salt, aspartic acid salt, glutamic acid salt etc.) and the like.

The compounds used according to the invention in accordance with the Formulae (I) or (II) can be present in their protonised form as ammonium salt of organic or inorganic acids such as hydrochloric acid, hydrobromic acid, sulphuric acid, nitric acid, methane sulphonc acid, p-toluene sulphonc acid, acetic acid, lactic acid, maleic acid, fumaric acid, oxalic acid, tartaric acid, benzoic acid etc.

The compounds used according to the invention of Formulas (I) or (II) permit for example the emergence of spatial isomers for groups containing double bonds or chiral groups R_1 , R_2 , R_3 , R_4 , X_1 , X_2 , X_3 , X_4 or A. The use of compounds according to the invention consists of all spatial isomers both as pure substances and in the form of their mixtures.

The organophosphorus compounds are suitable in particular for the therapeutic and prophylactic treatment of infections in humans and animals which are caused by viruses, unicellular and multicellular parasites and fungi.

The compounds are effective against unicellular parasites (protozoa), in particular against pathogens of malaria and sleeping sickness as well as Chagas' disease, toxoplasmosis, amoebic dysentery, leishmaniasis, trichomoniasis, pneumocystosis, balantidiasis, cryptosporidiasis, sarcocystosis, acanthamebiasis, naegleriasis, coccidiosis, giardiasis and lambliosis.

011538

They are therefore particularly suitable as malaria prophylactics and as prophylactics for sleeping sickness as well as Chagas' disease, toxoplasmosis, amoebic dysentery, leishmaniasis, trichomoniasis, pneumocystosis, balantidiasis, cryptosporidiasis, sarcocystosis, acanthamebiasis, naegleriasis, coccidiosis, giardiasis and lambliosis.

Combinations with an antibiotic can also be used to treat the above-mentioned diseases. Isoniazid, rifampicin, ethambutol, pyrazineamide, streptomycin, protionamide and dapsone are particularly suitable for combination with other anti-infectives for the treatment of tuberculosis.

The agents according to the invention can furthermore be used in particular in infections with the following viruses:

- Parvoviridae: parvo viruses, dependo viruses, denso viruses,
- 15 Adenoviridae: adeno viruses, mastadeno viruses, aviadeno viruses,
- Papovaviridae: papova viruses, in particular papilloma viruses (so called wart viruses), polyoma viruses, in particular JC virus, BK virus and miopapova viruses,
- 20 Herpesviridae: all herpes viruses, in particular herpes simplex viruses, varicella-zoster viruses, human cytomegalo virus, Epstein-Barr viruses, all human herpes viruses, human herpes virus 6, human herpes virus 7, human herpes virus 8,
- Poxviridae: pox viruses, orthopox, parapox, molluscum
- 25 contagiosum virus, avipox viruses, capripox viruses, leporipox viruses, all primary hepatotropic viruses, hepatitis viruses: hepatitis A viruses, hepatitis B viruses, hepatitis C viruses, hepatitis D viruses, hepatitis E viruses, hepatitis F viruses, hepatitis G viruses,
- 30 Hepadna viruses: all hepatitis viruses, hepatitis B virus, hepatitis D viruses,

011538

Picornaviridae: picorna viruses, all entero viruses, all polio viruses, all coxsackie viruses, all echo viruses, all rhino viruses, hepatitis A virus, aphtho viruses,

Calciviridae: hepatitis E viruses,

5 Reoviridae: reo viruses, orbi viruses, rota viruses,

Togaviridae: toga viruses, alpha viruses, rubi viruses, pestiviruses, rubella virus,

Flaviviridae: flavi viruses, ESME virus, hepatitis C virus,

Orthomyxoviridae: all influenza viruses,

10 Paramyxoviridae: paramyxo viruses, morbilli virus, pneumo virus, measles virus, mumps virus

Rhabdoviridae: rhabdo viruses, rabies virus, lyssa virus, viscula stomatitis virus,

Coronaviridae: corona viruses,

15 Bunyaviridae: bunya viruses, nairo virus, phlebo virus, uuku virus, hanta virus, hantaan virus,

Arenaviridae: arena viruses, lymphocytic choriomeningitis virus,

20 Retroviridae: retro viruses, all HTL viruses, human T-cell leukaemia virus, oncorna viruses, spuma viruses, lenti viruses, all HI viruses,

Filoviridae: Marburg and Ebola virus,

Slow virus infections, prions,

Oncoviruses and leukaemia viruses.

25 The organophosphorus compounds used according to the invention are therefore suitable for fighting the following viral infections:

30 Eradication of papilloma viruses to prevent tumours, in particular tumours in the sexual organs caused by papilloma viruses in humans, eradication of JC viruses and BK viruses, eradication of herpes viruses, eradication of human herpes viruses 8 for the treatment of Kaposi's sarcoma, eradication of cytomegalo viruses before transplants, eradication of Epstein-Barr viruses before transplants and to prevent tumours

associated with Epstein-Barr viruses, eradication of hepatitis viruses for the treatment of chronic liver diseases and for the prevention of tumours of the liver and cirrhosis of the liver, eradication of coxsackie viruses in patients with

5 cardiomyopathy, eradication of coxsackie viruses in diabetes mellitus patients, eradication of immune system debilitating viruses in humans and animals, treatment of secondary infections in AIDS patients, treatment of inflammations of viral origin of the respiratory tract (larynx papillomas,

10 hyperplasias, rhinitis, pharyngitis, bronchitis, pneumonias), of the sensory organs (keratoconjunctivitis), of the nervous system (poliomyelitis, meningoencephalitis, encephalitis, subacute sclerosing panencephalitis SSPE, progressive multifocal leukoencephalopathy, lymphocytic choriomeningitis), of the

15 gastro-intestinal tract (stomatitis, gingivostomatitis, oesophagitis, gastritis, gastroenteritis, diarrhoea-causing diseases), of the liver and gall bladder system (hepatitis, cholangitis, hepatocellular carcinoma), of the lymphatic tissue (mononucleosis, lymphadenitis), of the haematopoietic system, of

20 the sexual organs (mumpsorchitis), of the skin (warts, dermatitis, herpes labialis, heat rash, herpes zoster, shingles), of the mucous membranes (papillomas, conjunctiva papillomas, hyperplasias, dysplasias), of the heart/blood vessel system (arteritis, myocarditis, endocarditis,

25 pericarditis), of the kidney/urinary tract system, of the sexual organs (anogenital lesions, warts, genital warts, acute condylomas, displasias, papillomas, cervix dysplasias, condylomata acuminata, epidermodysplasia verruci formis), of the organs of motion (myositis, myalgias), treatment of foot

30 and mouth diseases in cloven-hoofed animals, of Colorado tick fever, of Dengue syndrome, of haemorrhagic fever, of early summer meningoencephalitis (ESME) and of yellow fever.

The described compounds, i.e. the organophosphorus compounds according to Formulae (I) and (II) and esters and amides

35 thereof on the phosphinic group or phosphonic group and salts

thereof have a strongly cytotoxic efficacy against unicellular and multicellular parasites, in particular against pathogens of malaria and sleeping sickness. Accordingly the compounds used according to the invention can be used for the treatment of infectious diseases which are caused by viruses, parasites and fungi in humans and animals. The compounds are also suitable for use in the prevention of diseases which are caused by viruses, parasites and fungi, in particular as malaria prophylactics and as sleeping sickness prophylactics.

The organophosphorus compounds used according to the invention, which generally include pharmaceutically acceptable salts, amides, esters, a salt of such an ester or else compounds which upon application provide the compounds used according to the invention as metabolic products or decomposition products, also called "prodrugs", can all be prepared for administration like known anti-infectious agents in any suitable manner (mixed with a non-toxic pharmaceutically acceptable carrier).

Pharmaceutically acceptable salts of the compounds include salts which form the compounds used according to the invention of Formulae (I) and (II) in their protonised form as an ammonium salt of inorganic or organic acids, such as hydrochloric acid, sulphuric acid, citric acid, maleic acid, fumaric acid, tartaric acid, p-toluene sulphonic acid.

Salts which are formed by suitable selection of X_3 and X_4 , such as sodium salt, potassium salt, calcium salt, ammonium salt, ethanolamine salt, triethylamine salt, dicyclohexylamine salt and salts of an amino acid such as arginine salt, aspartic acid salt, glutamine acid salt are particularly suitable pharmaceutically.

The activity of substances is determined in a test system. This system is based on the measuring of the inhibition of growth of parasites, viruses, fungi or plants *in vitro*. To this end, test

procedures are used, some of which are known to the person skilled in the art.

5 To determine the anti-malaria activity, for example, the inhibition of the growth of malaria parasites in blood cultures is determined.

The determining of antiviral activity is based on inhibition of the formation of viral elements in cell cultures.

10 The determining of fungicidal activity is based on the inhibition of the growth of fungi on culture media and in liquid cultures.

Some of the microorganisms which should be investigated can only be investigated in animal models. In this case we will use the corresponding models.

15 Substances which demonstrate an efficacy in the *in vitro* measuring systems will be further investigated in *in vivo* models.

The anti-parasitic, antiviral or fungicide activity will be further evaluated in the appropriate animal models.

20 The pharmaceutically effective preparations can be prepared in the form of pharmaceutical preparations in dispensing units. This means that the preparation can be present in the form of individual parts, for example tablets, dragees, capsules, pills, suppositories and ampoules, the active ingredient content of which corresponds to a fraction or a multiple of a
25 single dose. The dispensing units can, for example, contain 1, 2, 3 or 4 single doses or $1/2$, $1/3$ or $1/4$ of a single dose. A single dose preferably contains the quantity of active ingredient which is administered during one application and

which usually corresponds to a whole, a half or a third or a quarter of a daily dose.

Non-toxic, inert pharmaceutically suitable excipients are understood to mean solid, semi-solid or liquid diluents,
5 fillers and formulation auxiliary agents of all kinds.

Tablets, dragees, capsules, pills, granules, suppositories, solutions, suspensions and emulsions, pastes, ointments, gels, creams, lotions, powders and sprays are mentioned as suitable pharmaceutical preparations. Tablets, dragees, capsules, pills
10 and granules can contain in addition to the conventional excipients, the active ingredient or active ingredients such as (a) fillers and diluents, for example starches, lactose, cane sugar, glucose, mannitol and silicic acid, (b) binders, for example carboxymethylcellulose, alginates, gelatines,
15 polyvinylpyrrolidone, (c) moisture-retaining agents, for example glycerol, (d) dispersing agents, for example agar-agar, calcium carbonate and sodium carbonate, (e) solution retarders, for example paraffin and (f) resorption accelerators, for example quaternary ammonium compounds, (g) wetting agents, for
20 example cetyl alcohol, glycerol monostearate, (h) adsorption agents, for example kaolin and bentonite and (i) lubricants, for example talcum, calcium and magnesium stearate and solid polyethylene glycols or mixtures of the substances listed under (a) to (i).

25 The tablets, dragees, capsules, pills and granular materials can be provided with the conventional coatings and casings optionally containing opaquing agents and can also be put together so that they release the active ingredient or active ingredients only or preferably in a specific part of the
30 intestinal tract optionally with sustain release, wherein polymer substances and waxes for example can be used as embedding compounds.

The active ingredient or active ingredients can optionally also be present in microencapsulated form with one or more of the above-mentioned excipients.

5 In addition to the active ingredient or active ingredients suppositories can also contain the conventional water soluble or water insoluble excipients, for example polyethylene glycols, fats, for example cocoa fat and higher esters (for example C14-alcohol with C16-fatty acid) or mixtures of these substances.

10 In addition to the active ingredient or active ingredients, ointments, pastes, creams and gels can contain the conventional excipients, for example animal and vegetable fats, waxes, paraffins, starches, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc
15 and zinc oxide or mixtures of these substances.

In addition to the active ingredient or active ingredients, powders and sprays can contain the conventional excipients, for example lactose, talc, silicic acid, aluminium hydroxide, calcium silicate, and polyamide powder or mixtures of these
20 substances. Sprays can additionally contain the conventional blowing agents, for example chlorofluorohydrocarbons.

In addition to the active ingredient or active ingredients, solutions and emulsions can contain the conventional excipients such as solvents, solubilisers and emulsifiers, for example
25 water, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils, in particular cotton seed oil, peanut oil, corn oil, olive oil, castor oil and sesame oil, glycerol, glycerol formal, tetrahydrofurfuryl
30 alcohol, polyethylene glycols and fatty acid esters of sorbitan or mixtures of these substances.

The solutions and emulsions can also be present in sterile and blood isotonic form for parenteral application.

In addition to the active ingredient or active ingredients, suspensions can contain conventional excipients such as liquid diluents, for example water, ethyl alcohol, propylene glycol, suspending agents, for example ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminium metahydroxide, bentonite, agar-agar and tragacanth or mixtures of these substances.

The above-mentioned formulations can also contain dyes, preservatives and odour and flavour improving additives, for example peppermint oil and eucalyptus oil and sweeteners, for example saccharine.

The agents of Formulae (I) and (II) should be present in the above-listed pharmaceutical preparations preferably in a concentration of approximately 0.1 to 99.5% by weight, preferably of approximately 0.5 to 95% by weight of the total mixture.

In addition to the compounds of Formulae (I) and (II) the pharmaceutical preparations can also contain further pharmaceutical agents.

The compounds can be used with hitherto described substances with antibacterial, antiviral, antifungal and antiparasitic properties. Compounds which have already found application in treatment or are still being used belong to this group. Substances which are listed in the Red List or in Simon/Stille, Antibiotika-Therapie in Klinik und Praxis, 9th Edition 1998 Schattauer Verlag or under

<http://www.customs.treas.gv/imp-exp/rulings/harmoniz/hrm129.html>

on the Internet are particularly suitable for this purpose. In particular derivatives with penicillins, benzyl penicillin (Penicillin G), phenoxy penicillins, isoxazolyl penicillins, amino penicillins, ampicillin, amoxicillin, bacampicillin, 5 carboxy penicillin, ticarcillin, temocillin, acyalamino pencillins, azlocillin, mezlocillin, piperacillin, apalcillin, mecillinam, cephalosporins, cefazolin group, cefuroxime group, cefoxitin group, cefoxitin, cefotetan, cefmetazole, latamoxef, flomoxef, cefotaxime group, ceftazidime, ceftazidime group, 10 ceftazidime, cefpirom, cefepim, other cephalosporins, cefsulodine, cefoperazone, oralcephalosporins of the cefalexine group, loracarbef, cefprozil, new oralcephalosporins with expanded spectrum, cefixime, cefpodoxim-proxetil, cefuroxime-axetil, cefetamet, cefotiam hexetil, cefdinir, ceftibutene, 15 other β -lactam antibiotics, carbapenem, imipenem/cilastatin, meropenem, biapenem, aztreonam, β -lactamase inhibitors, clavulanic acid/amoxicillin, clavulanic acid/ticarcillin, sulbactam/ampicillin, tazobactam/piperacillin, tetracyclines, oxytetracycline, rolitetracycline, doxycycline, minocycline, 20 chloramphenicol, aminoglycosides, gentamicin, tobramycin, netilmicin, amikacin, spectinomycin, macrolides, erythromycin, clarithromycin, roxithromycin, azithromycin, dirithromycin, spiramycin, josamycin, lincosamides, clindamycin, fusidic acid, glycopeptide antibiotics, vancomycin, tecoplanin, pristinaamycin 25 derivatives, fosfomycin, antimicrobial folic acid antagonists, sulphonamides, co-trimoxazole, trimethoprim, other diaminopyrimidine-sulphonamide combinations, nitrofurans, nitrofurantoin, nitrofurazone, Gyrase inhibitors (quinolones), norfloxacin, ciprofloxacin, ofloxacin, sparfloxacin, enoxacin, 30 fleroxacin, pefloxacin, lomefloxacin, Bay Y3118, nitroimidazoles, antimycobacterial agents, isoniazid, rifampicin, rifabutin, ethambutol, pyrazinamide, streptomycin, capreomycin, prothionamide, terizidon, dapsone, clofazimine, topical antibiotics, bacitracin, tyrothricin, polymyxins,

neomycin, kanamycin, paromomycin, mupirocin, antiviral agents, acyclovir, ganciclovir, azidothymidine, didanosin, zalcitabin, thiacytidine, stavudine, ribavirin, idoxuridine, trifluridine, foscarnet, amantadine, interferons, tibol derivatives,
5 proteinase inhibitors, antifungal agents, polyenes, amphotericin B, nystatin, natamycin, azoles, azoles for septic treatment, miconazole, ketoconazole, itraconazole, fluconazole, UK-109.496, azoles for topical application, clotrimazole, econazole, isoconazole, oxiconazole, bifonazole, flucytosine,
10 griseofulvin, ciclopiroxolamine, tolnaftate, naftifine, terbinafine, amorolfine, anthraquinones, betulinic acid, semianthraquinones, xanthenes, naphthoquinones, aryaminoalcohols, quinine, quinidines, mefloquine, halofantrine, chloroquine, amodiaquine, acridine,
15 benzonaphthylidine, mepacrine, pyronaridine, dapsone, sulphonamide, sulfadoxine, sulfalenes, trimethoprim, proguanil, chlorproguanil, diaminopyrimidine, pyrimethamine, primaquine, aminoquinolines, WR 238,605, tetracycline, doxycycline, clindamycin, norfloxacin, ciprofloxacin, ofloxacin,
20 artemisinin, dihydroartemisinin, 10b arte methether, arte ether, arte sunate, atovaquon, suramin, melarsoprol, nifurtimox, stibogluconate-sodium, pentamidine, amphotericin B, metronidazole, clioquinol, mebendazole, niclosamide, praziquantel, pyrantel, tiabendazole, diethylcarbamazine,
25 ivermectin, bithionol, oxamniquine, metrifonate, piperazine, embonate can be [used].

The organophosphorus compounds can furthermore be present in the pharmaceutical preparations in combination with
sulphonamide, sulfadoxin, artemisinin, atovaquon, quinine,
30 chloroquine, hydroxychloroquine, mefloquine, halofantrine, pyrimethamine, artesunate, tetracyclines, doxycycline, proguanil, metronidazole, praziquantel, niclosamide, mebendazole, pyrantel, tiabendazole, diethylcarbamazine, piperazine, pyrivinum, metrifonate, oxamniquine, bithionol or suramin or
35 several of these substances.

The above-listed pharmaceutical preparations are produced in the conventional manner by known methods, for example by mixing the active ingredient or active ingredients with the excipient or excipients.

5 The above-mentioned preparations can be used in humans and animals either orally, rectally, parenterally, (intravenously, intramuscularly, subcutaneously), intracisternally, intra-
10 vaginally, intraperitoneally, topically (powder, ointment, drops) and for the treatment of infections in cavities, orifices. Suitable preparations are injection solutions, solutions and suspensions for oral treatment, gels, infusions, emulsions, ointments or drops. Ophthalmological and dermatological formulations, silver and other salts, eardrops, eye ointments, powders or solutions can be used for topical
15 treatment. With animals the absorption can occur via the food or drinking water in suitable formulations. Furthermore gels, powders, tablets, sustain release tablets, premixes, concentrates, granules, pellets, tablets, boli, capsules, aerosols, sprays, inhalers can be used with humans and animals.
20 The compounds used according to the invention can furthermore be incorporated into other carrier materials such as, for example, plastics materials (plastics chains for topical treatment), collagen or bone cement.

25 In general it has proved advantageous both in human and veterinary medicine to administer the active ingredient or active ingredients of Formulae (I) and (II) in total quantities of approximately 0.05 to approximately 600, preferably 0.5 to 200 mg/kg body weight every 24 hours, optionally in the form of several individual doses in order to achieve the desired
30 results. An individual dose contains the active ingredient or active ingredients preferably in quantities from approximately 1 to approximately 200, in particular 1 to 60 mg/kg body weight. It may however be necessary to deviate from the above-mentioned dosages and this is dependent on the nature and the

body weight of the patient to be treated, the nature and the severity of the disease, the nature of the preparation and the application of the pharmaceutical composition as well as the time scale or interval within which the administration takes place.

Thus in some cases it may be sufficient to get by with less than the above-mentioned quantity of active ingredient whilst in other cases the above-stated quantity of active ingredient must be exceeded. The person skilled in the art can determine the optimum dosage and method of application of the active ingredient required in each case by virtue of their expert knowledge.

The compounds according to the invention can be given in the conventional concentrations and preparations to animals together with the feedstuff or the feedstuff preparations or with the drinking water.

Example 1

Test of the efficacy of substances against malaria in vivo

The various derivatives were tested by the modified Peters' test. The substances were applied here in a quarter of the median lethal dose (LD50). In the test batch ten mice were infected with *Plasmodium vinckei*, the pathogen of mouse malaria. After confirmation of the infection by blood testing four mice were treated. Six mice, which were not treated, served as a control group. The treatment with 1,000 mg/kg/d, 3-(N-formyl-N-hydroxylamino)-propylphosphonic acid monosodium salt over 3 days led to a destruction of the parasites in the mice's blood. After just one day the treated group was free of living parasites. The control mice had to be killed on day 5 after infection with parasitaemia of > 80%. The treated mice were still free from parasites 8 weeks after the treatment had stopped. Further experiments showed an efficacy of 50 mg/kg/d

3-(N-formyl-N-hydroxylamino)-propylphosphonic acid monosodium salt in mice with a parasitaemia of 80%. These mice were also free of living parasites after 1 day.

Example 2

5 Protection from malaria in a test with infected mice

The efficacy of compounds *in vivo* against malaria was tested using male mice weighing 20 to 25 g (BALB/c-strain). Four mice were treated intraperitoneally with 50 mg/kg 3-(N-formyl-N-hydroxylamino)-propylphosphonic acid monosodium salt one day
10 before infection. The mice were then infected with *Plasmodium vinckei*. Mice which were not pretreated with the substance served as a control group. No infection could be detected in the treated mice, whilst the control mice were killed after 5 days with a parasitaemia of 80%. The treated mice were still
15 free of parasites 8 weeks after infection.

Example 3

In vitro cytotoxicity against malaria parasites

On the principle of IC₅₀ determination according to Vial et al (the concentration at which the vitality of parasites is
20 reduced by a half)

To determine the IC₅₀ value according to Vial et al malaria parasites are initially cultivated for a complete 48 hour cycle in the presence of inhibitors, in the subsequent 24 hours the survival rate is measured by [³H]-hypoxanthine insertion.

25 Carrying out the test

A dilution series of 3-(N-formyl-N-hydroxylamino)-propylphosphonic acid monosodium salt is placed in 20 µl aliquots concentrated by 10 on a microtitre plate. Then 180 µl parasite suspension in culture medium is added to each well.
30 Asynchronous cultures with a parasitaemia of approximately 0.4%

and 2% haematocrite are used. The microtitre plates are subsequently incubated for 48 hours. Then 30 μ l [3 H]-hypoxanthine are added to each well. After 24 hours of incubation the cells were harvested and the incorporated radioactivity was measured. In Fig. 1 the results with the strains HB3 and Dd2 are shown with known resistances against other malaria pharmaceutical compositions. In both strains there is a IC50 value of approximately 100 μ g/l. The resistances of these strains are:

Plasmodium falciparum HB3 (Honduras) is resistant to pyrimethamine.

Plasmodium falciparum Dd2 (Indo-China) is resistant to cloroquine, quinine, pyrimethamine, cycloguanil and sulfadoxine.

No cross resistances with antimalarial agents were found.

Example 4

Preparation of 3-bromopropylphosphonic acid diethylester

471 g (238 ml, 2.33 mol) 1,3-dibromopropane and 77.6 g (81 ml, 0.467 mol) triethylphosphite were introduced into a 500 ml flask and heated for 30 minutes to 155-160°C. 20 ml ethyl bromide (boiling point: 40°C) were distilled off under normal pressure via a reflux condenser and a distilling apparatus. Concentration of the solution under vacuum (8 torr ($1.07 \cdot 10^3$ Pa)) resulted in 380 g (191 ml, 1.863 mol) 1,3 dibromopropane (surplus educt). 88.1 g (0.34 mol) could be distilled as a colourless liquid (boiling point: 96°C, 0.1 torr (13.33 Pa)) from the remaining yellow oil. This corresponds to a yield of 73%. (Hewitt, Teese, Aust.J.Chem. 1984, 37, 205-10 US-Patent 4 206 156).

$^1\text{H-NMR}$ (CDCl_3) δ = 4.08 (quintet, $J=7$ Hz, 4H), 1.33 (t, $J=7$ Hz, 6H)

$^{13}\text{C-NMR}$ (CDCl_3) δ = 61.2 (OCH_2CH_3), 33.10 ($J=18$, 3 Hz), 25.6 ($J=4$, 4 Hz), 24.14 ($J=120$, 6 Hz) 16.04 (OCH_2CH_3)

Example 5

Preparation of 3-(N-hydroxyamino)-propylphosphonic acid diethylester

Firstly 32.0 g (0.8 mol) of sodium hydroxide dissolved in 75 ml water then 75 ml methanol, and finally 25.5 g (0.098 mol) 3-bromopropyl phosphonic acid diethylester were added drop by drop to a solution of 55.6 g (0.8 mol) hydroxylamine hydrochloride in 100 ml water with cooling with ice. This led to a clouding of the solution. After 3 hours stirring at a temperature of 40 to 45°C methanol was removed under reduced pressure, the resulting aqueous solution was saturated with NaHCO_3 (pH=8), shaken out three times with 60 ml toluene in each case (the toluene phase was discarded) and then shaken out with chloroform (1 x with 90 ml, 2 x with 50 ml in each case). The slightly yellowy chloroform phase was dried over MgSO_4 . After filtering of the dehydrating agent the solution was concentrated under reduced pressure. 15.43 g (0.0728 mol) 3-(N-hydroxyamino)-propylphosphonic acid diethylester were obtained as an almost colourless oil. This corresponds to a yield of 74.3% (DE-A-27 33 658).

$^1\text{H-NMR}$ (CDCl_3) δ = 5.94 (wide s, 2H), 4.13 (quintet, $J=7$ Hz, 4H) 2.90 (t, $J=7$ Hz, 2H) 1.5-2.2 (m, 4H), 1.33 (t, $J=7$ Hz, 6H)

$^{13}\text{C-NMR}$ (CDCl_3) δ = 61.23 (OCH_2CH_3), 53.34 (NCH_2 , $J=15$, 9 Hz), 22.75 ($J=141$, 9 Hz), 19.77 Hz, 16.08 (OCH_2CH_3).

Example 6**Preparation of 3-(N-hydroxyamino)-propylphosphonic acid**

12.9 g (0.0608 mol) 3-(N-hydroxylamino)-propylphosphonic acid diethylester and 130 ml concentrated HCl were heated for 6
5 hours under reflux (oil bath temperature: 150°C). The resulting yellow/orange solution is concentrated under reduced pressure. The remaining oil is taken up in 30 ml water, stirred with 3 spoonfuls of activated carbon for 30 minutes, filtered out from the activated carbon and the colourless solution is
10 concentrated in the entire diaphragm pump vacuum. After take-up in 30 ml water a pH of 4.0 to 4.5 is adjusted with approximately 4.7 g (0.056 mol) NaHCO₃ (from pH = 1.5 the product precipitates). Suction filtering of the white solid resulted in 5.83 g 3-(hydroxyamino)-propylphosphonic acid
15 (melting point: 160°C, decomposition). This corresponds to a yield of 61.8%. (DE-A-27 33 658, Öhler Synthesis 1995, 539-543).

¹H-NMR (CDCl₃) δ = 3.49 (t, J=7, 4 Hz, 2H), 2.1 (m, 2H), 1.82 (m, PCH₂, 2H)

20 ¹³C-NMR (CDCl₃) δ = 56.26 (NCH₂, J=15 Hz), 29.61 (PC, J=134 Hz), 22.37 (C-2, J=3, 8 Hz).

Example 7

Preparation of 3-(N-formylhydroxyamino)-propylphosphonic acid diethylester

1.38 (0.030 mol) formic acid were added drop by drop at room temperature to 2.04 g (0.020 mol) acetic anhydride and stirred at the same temperature. This solution was added with cooling with ice to 2.8 g (0.013 mol) 3-(N-hydroxyamino)-propylphosphonic acid diethylester dissolved in chloroform. The reaction mixture is stirred for 30 minutes at 0-5°C and for a further 1.5 hours at room temperature. After concentration under reduced pressure until an oily residue is obtained, this oily residue is taken up in 15 ml methanol and 5 ml water, adjusted to pH = 8 with 2 n NaOH and stirred for a further 1.5 hours at room temperature. Methanol is removed under reduced pressure and the resultant aqueous solution is adjusted to pH = 5 with concentrated HCl. The yellow solution is extracted with chloroform (1 x 30 ml, 2 x 10 ml in each case) and the CHCl₃ phases are dried over MgSO₄. After concentration of the solution in the entire diaphragm pump vacuum 3 g of a yellow oil are obtained. After removal of volatile constituents in the entire diaphragm pump vacuum, chromatography on 60 g SiO₂ with chloroform shows: methanol in the ratio 25:1 2.65 g product as yellow oil (DE-A-27 33 658).

¹H-NMR (CDCl₃) δ = 8.4 (CHO, 0.5 H), 7.94 (CHO, 0.5 H), 4.1 (quintet, 4 H), 3.68 (t, 2 H), 1.7-2.19 (m, 4 H), 1.36 (t, J=7 Hz, 6 H)

¹³C-NMR (CDCl₃) δ = 162.65 (CHO), 156.96 (CHO), 61.72 (OCH₂CH₃), 46.31 (NCH₂, J=15, 9 Hz), 22.15 (PC, J=142, 0 Hz), 19.13 (C-2), 16.08 (OCH₂CH₃).

011538

Example 8**Preparation of 3-(N-acetylhydroxyamino)-propylphosphonic acid diethylester**

2.8 g (0.013 mol) 3, (N-hydroxyamino)-propylphosphonic acid
5 diethylester are dissolved in 30 ml methylene chloride and
added drop by drop with cooling with ice to 2.65 g (0.026 mol)
acetic anhydride. The reaction mixture is stirred for 30
minutes at 0-5°C and for a further 1.5 hours at room
temperature. After concentration under reduced pressure until a
10 yellow oily radical is achieved, this oily residue is taken up
in 15 ml methanol and 5 ml water, adjusted to pH 8 with 2 n
NaOH and stirred for a further 1.5 hours at room temperature.
Methanol is removed under reduced pressure and the resultant
solution is adjusted to pH = 5 with concentrated HCl. The
15 yellow solution is extracted repeatedly with methylene chloride
(1 x 30 ml, 2 x 10 ml in each case), the combined CH₂Cl₂ phases
are dried over MgSO₄ and the solvent is removed at room
temperature under reduced pressure. 3.7 g of a yellow oil are
obtained which are freed from adhering volatile substances in
20 the entire diaphragm pump vacuum. 2.78 g yellow oil remain.

¹³C-NMR (CDCl₃) δ = 171.96 (C=O), 61.62 (OCH₂CH₃), 47.44 (J=15,
49 Hz), 22.13 (PC, J=141, 8 Hz), 19.3, 15.9
(OCH₂CH₃).

Example 9**Preparation of 3-(N-formylhydroxyamino)-propylphosphonic acid monosodium salt**

2 ml formic acid are added drop by drop to 4 ml acetamide at
0-5°C. The solution is stirred at this temperature for 10
minutes and for a further 15 minutes at room temperature,
30 subsequently cooled to 0°C again and 3.28 g (0.021 mol) 3-(N-
hydroxyamino)-propylphosphonic acid dissolved in 6 ml formic

acid are added drop by drop at 0-5°C. After 1 hour of stirring at room temperature the solution is condensed in a rotary evaporator under reduced pressure, the oil is dissolved in 50 ml methanol, heated to 60°C and mixed with 10 ml ethanol.

5 The resulting oily separated material is separated without stirring by decanting. To precipitate white crystals the methanolic solution is mixed with a further 50 ml ethanol, boiled up and the white residue filtered out. This residue is taken up in 80 ml methanol and 100 ml ethanol are added with
10 stirring. The mixture is stirred further overnight at room temperature. A solid is obtained which is filtered out.

(DE-A-27 33 658)

Example 10

Preparation of 3-(N-acetylhydroxyamino)-propylphosphonic acid 15 monosodium salt

A suspension of 3.8 g (0.02 mol) 3-(N-hydroxyamino)-propylphosphonic acid is introduced into 20 ml water and 4.51 g (0.044 mol) acetic anhydride are added drop by drop at room temperature. After the solution has been stirred for 1.5 hours
20 at room temperature a pH of 2.5 is adjusted with 0.2 n NaOH, the solution is concentrated in the entire diaphragm pump vacuum, taken up twice in 40 ml water in each case, which are again removed by concentration and the oil is washed twice with 30 ml ether in each case, taken up in 50 ml water and a pH of
25 6.5 is adjusted with 1.6 g NaHCO₃. After removal of volatile constituents under vacuum, 20 ml n-butanol are added to remove the residual water and are also removed under reduced pressure. The oil is boiled up twice with isopropanol, the isopropanol phase is discarded and the remaining glassy resin is pulverised
30 with a spatula to a yellow solid (5.65 g). In order to recrystallise it, it is taken up in only a little methanol, filtered from the undissolved component and acetone is added dropwise to the filtrate. A first filtering results in 1 g product with a melting point of 175°C. Recrystallisation as

described above is carried out again for further purification.
(DE-A-27. 33 658)

Example 11

Preparation of antiparasitically effective agents

5 Preparation for injections:

(1) The required quantity of sterile antiparasitically effective agent, 3-(N-formyl-N-hydroxylamino)-propylphosphonic acid monosodium salt was divided into phials or ampoules which then contained 500 mg active ingredient. The phials were
10 hermetically sealed to exclude bacteria. For injections 2 ml sterile water were added to the phials in each case and the content was administered.

In substantially the same manner as described under (1) further injectable preparations of the antiparasitically effective
15 agent were prepared as described below:

(2) 250 mg 3-(N-formyl-N-hydroxylamino)-propylphosphonic acid monosodium salt were used as active ingredient for the injections.

(3) 250 mg 3-(N-formyl-N-hydroxylamino)-trans-1-propenyl-phosphonic acid monosodium salt were used as active ingredient
20 for the injections.

(4) 500 mg 3-(N-acetyl-N-hydroxylamino)-2-hydroxypropyl-phosphonic acid monosodium salt were used as active ingredient for the injections.

(5) 250 mg 3-(N-formyl-N-hydroxylamino)-2-hydroxypropyl-phosphonic acid monopotassium salt were used as active
25 ingredient for the injections.

Preparation of tablets:

A suitable tablet recipe is formed by the following mixture:

	3- (N-formyl-N-hydroxylamino) -propylphosphonic acid monosodium salt	200 mg
	Mannitol	400 mg
5	Starch	50 mg
	Magnesium stearate	10 mg

Preparation of capsules:

	3- (N-formyl-N-hydroxylamino) -propylphosphonic acid monopotassium salt	300 mg
10	Magnesium stearate	15 mg


The present components were mixed and then introduced into a hard gelatine capsule in the conventional manner.

Preparation of an oily suspension:

	3- (N-acetyl-N-hydroxylamino) -propylphosphonic acid monosodium salt	200 mg
15	Lanette-Wax SX®	50 mg
	Soft paraffin	100 mg
	Brilliant blue FCF	25 mg

The above components were mixed with liquid paraffin for a total quantity of 3 g to achieve an infusion preparation.

Examples of methods of synthesis for substances with the following structure:



where $\text{R}_1=\text{R}_2=\text{C}_{18}\text{H}_{37}$

12

where $\text{R}_1=\text{H}$ or Na^+ and $\text{R}_2=\text{C}_{18}\text{H}_{37}$

13

where $\text{R}_1=\text{H}$ or Na^+ and $\text{R}_2=\text{C}_{16}\text{H}_{33}$

14

Example 12:3-(N-formyl-N-hydroxylamino)-propylphosphonic acid dioctadecyl-ester 12

1 equivalent fosmidomycin (FR-31564) and 6 equivalents
tris(octadecyl)-orthoformic acid were heated under reduced
5 pressure and vigorous stirring for 2 hours under reflux. Then
methanol and formic acid octadecylester were distilled off
- also under reduced pressure - wherein the temperature must be
kept below the decomposition temperature of the product.
Further volatile secondary products are removed in the oil pump
10 vacuum in order finally to obtain 12 as a highly viscous oil.
(On the carrying out of the procedure cf.: D.A. Nicholson, W.A.
Cilley, O.T. Quimby, J Org Chem. 1970, 35, 3149-50)

The monoesters can be provided starting from both fosmidomycin
and di-octadecylester 12.

15 Example 13:3-(N-formyl-N-hydroxylamino)-propylphosphonic acid
mono-octadecyl-ester 13

First proposal:

0.21 mmol fosmidomycin (phosphonic acid) and 0.2 mmol
20 n-octadecanol are dissolved in 1-2 ml dry pyridine and
0.67 mmol trichloroacetonitrile is added dropwise thereto. The
reaction mixture is heated for 16 hours to 80°C and then
condensed under vacuum. After take up in water (for the
diaphragm filtering of undissolved components) the solution is
25 condensed again under reduced pressure and the product is
chromatographed onto silica gel with ethyl acetate, ethanol and
water as eluent. The product 13 is produced in this case as a
viscous rubber-like to glass-like compound.

(On the carrying out of the procedure cf.: G.B. Brookes, D.
30 Edwards, J.D.I. Hatto, T.C. Smale, R. Southgate, Tetrahedron
1995, 51, 7999-814)

Second proposal:

0.2 mmol. of the diester 3-(N-formyl-N-hydroxylamino)-
propylphosphonic acid dioctadecylester 13 dissolved in ethanol
is added to 1 equivalent KOH (ethanolic solution) and boiled
5 for 10 hours under reflux. Owing to the introduction of CO₂,
potassium salts can precipitate as carbonates and can be
removed by filtration. The filtrate is condensed to dryness,
the oil is dried over P₂O₅, washed with petrol ether and the
product can finally be recrystallised from absolute ethanol by
10 the addition of isopropanol.

(On the carrying out of the procedure cf.: V. Jagodic, Chem Ber
1960, 93, 2308-13)

Example 14:

3-(N-formyl-N-hydroxylamino)-propylphosphonic acid
15 monohexadecyl-ester 14

14 can be synthesised in the same way as 13.

011538

CLAIMS :

1. Use of organophosphorus compounds of the general formula
(1)



in which R_1 and R_2 are the same or different and are selected from the group which consists of hydrogen, substituted and unsubstituted alkyl, substituted and unsubstituted hydroxyalkyl, substituted and unsubstituted alkenyl, substituted and unsubstituted alkynyl, substituted and unsubstituted aryl, substituted and unsubstituted alkyl, substituted and unsubstituted cycloalkyl, substituted and unsubstituted aralkyl, substituted and unsubstituted heterocyclic radical, halogen, OX_1 and OX_2 ,

wherein X_1 and X_2 may be the same or different and are selected from the group which consists of hydrogen, substituted and unsubstituted alkyl, substituted and unsubstituted hydroxyalkyl, substituted and unsubstituted alkenyl, substituted and unsubstituted alkynyl, substituted and unsubstituted aryl, substituted and unsubstituted acyl, substituted and unsubstituted cycloalkyl, substituted and unsubstituted aralkyl, substituted and unsubstituted heterocyclic radical,

A is selected from the group which consists of an alkylene radical, an alkenylene radical and a hydroxyalkylene radical,

R_3 and R_4 are selected independently from the group which consists of hydrogen, substituted and unsubstituted C_{1-26} -alkyl, substituted and unsubstituted hydroxy- C_{1-26} -alkyl,

011538

substituted and unsubstituted aryl, substituted and unsubstituted acyl, substituted and unsubstituted aralkyl, substituted and unsubstituted C₁₋₂₆-alkenyl, substituted and unsubstituted C₁₋₂₆-alkinyl, substituted and unsubstituted cycloalkyl, substituted and unsubstituted heterocyclic radical, halogen, OX₃ and OX₄,

wherein X₃ and X₄ are selected independently from the group which consists of hydrogen, substituted and unsubstituted C₁₋₂₆-alkyl, substituted and unsubstituted hydroxyl-C₁₋₂₆-alkyl, substituted and unsubstituted aryl, substituted and unsubstituted aralkyl, substituted and unsubstituted C₁₋₂₆-alkenyl, substituted and unsubstituted C₁₋₂₆-alkinyl, substituted and unsubstituted cycloalkyl, substituted and unsubstituted heterocyclic radical, a silyl, a cation of an organic and inorganic base, in particular a metal of the first, second or third main group of the periodic system, ammonium, substituted ammonium and ammonium compounds which derive from ethylene diamine or amino acids, and their pharmaceutically acceptable salts, esters and salts of esters in order to produce a pharmaceutical composition for therapeutic and prophylactic treatment of infections in humans and animals caused by parasites, fungi and viruses.

2. Use according to claim 1, characterised in that the organophosphorus compounds correspond to Formula (II),



wherein,

X₁ is selected from the group which consists of hydrogen, substituted or unsubstituted acyl, substituted or unsubstituted alkyl, substituted or unsubstituted aryl, substituted or unsubstituted aralkyl, substituted or unsubstituted cycloalkyl, substituted or unsubstituted heterocyclic radical.

3. Use according to claim 2, characterised in that R_2 is an acyl radical, preferably an alkanoyl radical and particularly preferably a formyl or acetyl radical.
4. Use according to one of the preceding claims, R_3 and R_4 are selected independently from the group which consists of hydrogen, methyl, ethyl, or R_4 is OX_4 , wherein X_4 is selected from the group which consists of hydrogen, sodium, potassium, methyl and ethyl.
5. Use according to one of the preceding claims, characterised in that between the phosphorus atom and the nitrogen atom A forms a chain of three carbon atoms, preferably a propylene, propenylene or hydroxypropylene chain.
6. Use according to claim 1 or 2, characterised in that R_4 represents OX_4 and X_4 is selected from the group which consists of hydrogen, ammonium and metals of the first and second main group of the periodic system, preferably sodium, potassium, calcium or magnesium, ammonium compounds which derive from ethylenediamine or amino acids, preferably ethanolamine, ethylenediamine, N,N-dibenzylethylenediamine and arginine.
7. Use according to one of the preceding claims for the treatment of infections which are caused by viruses selected from the group which consists of viruses of the genus parvoviridae, in particular parvo viruses, dependo viruses, denso viruses, viruses of the adenoviridae genus, in particular adeno viruses, mastadeno viruses, aviadeno viruses, viruses of the genus papovaviridae, in particular papova viruses, in particular papilloma viruses (so called wart viruses), polyoma viruses, in

particular JC-virus, BK-virus and miopapova viruses,
viruses of the genus herpesviridae, in particular herpes
simplex viruses, of the varicella-zoster viruses, human
cytomegalo virus, Epstein-Barr viruses, human herpes
5 virus 6, human herpes virus 7, human herpes virus 8,
viruses of the genus poxviridae, in particular pox
viruses, orthopox, parapox, molluscum contagiosum virus,
avipox viruses, capripox viruses, leporipox viruses,
primary hepatotropic viruses, in particular hepatitis
10 viruses, such as hepatitis A viruses, hepatitis B
viruses, hepatitis C viruses, hepatitis D viruses,
hepatitis E viruses, hepatitis F viruses, hepatitis G
viruses, hepadna viruses, in particular all hepatitis
viruses, such as hepatitis B virus, hepatitis D viruses,
15 viruses of the genus picornaviridae, in particular
picorna viruses, all entero viruses, all polio viruses,
all coxsackie viruses, all echo viruses, all rhino
viruses, hepatitis A virus, aphtho viruses, viruses of
the genus calciviridae, in particular hepatitis E
20 viruses, viruses of the genus reoviridae, in particular
reo viruses, orbi viruses, rota viruses, viruses of the
genus togaviridae, in particular toga viruses, alpha
viruses, rubi viruses, pestiviruses, rubella virus,
viruses of the genus flaviviridae, in particular flavi
25 viruses, ESME virus, hepatitis C virus, viruses of the
genus orthomyxoviridae, in particular influenza viruses,
viruses of the genus paramyxoviridae, in particular
paramyxo viruses, morbilli virus, pneumo virus, measles
virus, mumps virus, viruses of the genus rhabdoviridae,
30 in particular rhabdo viruses, rabies virus, lyssa virus,
visceral stomatitis virus, viruses of the genus
coronaviridae, in particular corona viruses, viruses of
the genus bunyaviridae, in particular bunya viruses,
nairo virus, phlebo virus, uuku virus, hanta virus,
35 hantaan virus, viruses of the genus arenaviridae, in
particular arena viruses, lymphocytic choriomeningitis
virus, viruses of the genus retroviridae, in particular
retro viruses, all HTL viruses, human T-cell leukaemia
virus, oncornaviruses, spuma viruses, lenti viruses, all
40 HI viruses, viruses of the genus filoviridae in
particular Marburg and Ebola virus, slow viruses,
oncoviruses and leukaemia viruses.

011538

8. Use according to one of the preceding claims for the prevention and treatment of infections caused by unicellular parasites namely pathogens of malaria, sleeping sickness, Chagas' disease, toxoplasmosis, amoebic dysentery, leishmaniasis, trichomoniasis, pneumocystosis, balantidiasis, cryptosporidiasis, sarcocystosis, acanthamebiasis, naegleriasis, coccidiosis, giardiasis and lambliosis.

9. Use according to one of claims 1 to 8 in a pharmaceutical preparation which has an active content of at least one organophosphorus compound and a pharmaceutically acceptable carrier.

10. Use according to claim 9, characterised in that the pharmaceutical preparation has at least one further pharmaceutical active ingredient, in particular sulphonamide, sulfadoxine, artemisinin, atovaquon, quinine, chloroquine, hydroxychloroquine, mefloquine, halofantrine, pyrimethamine, artesin, tetracyclines, doxycycline, proguanil, metronidazole, praziquantel, niclosamide, mebendazol, pyrantel, tiabendazol, diethylcarbazine, piperazine, pyrivinum, metrifonate, oxamni-
quine, bithionol or suramin.

11. Use according to claim 10, characterised by one or more components of the group which consists of benzyl penicillin (Penicillin G), phenoxy penicillins, isoxazolyl penicillins, amino penicillins, ampicillin, amoxicillin, bacampicillin, carboxy penicillin, ticarcillin, temocillin, acyalamino penicillins, azlocillin, mezlocillin, piperacillin, apalcillin, mecillinam, cephalosporins, cefazolin group, cefuroxime group, cefoxitin group, cefoxitin, cefotetan, cefmetazole, latamoxef, flomoxef, cefotaxime group, ceftazidime, ceftazidime groups, ceftazidime, cefpirom, cefepim, other cephalosporins, cefsulodine, cefoperazone, oralcephalosporins of the cefalexine group, laracarbef, cefprozil, new oralcephalosporins with expanded spectrum, cefixime, cefpodoxim proxetil, cefuroxime

axetil, cefetamet, cefotiam hexetil, cefdinir, ceftibutene,
other β -lactam antibiotics, carbapenem, imipenem/cilastatin,
meropenem, biapenem, aztreonam, β -lactamase inhibitors,
calvulanic acid/ amoxicillin, calvulanic acid/ticarcillin,
5 sulbactam/ampicillin, tazobactam/piperacillin, tetracyclines,
oxytetracycline, roli-tetracycline, doxycycline, minocycline,
chloramphenicol, amino-glycosides, gentamicin, tobramycin,
netilmicin, amikacin, spectinomycin, macrolides, erythromycin,
clarithromycin, roxithromycin, azithromycin, dirithromycin,
10 spiramycin, josamycin, lincosamides, clindamycin, fusidic acid,
glycopeptide antibiotics, vancomycin, tecoplanin, pristinamycin
derivates, fosfomycin, antimicrobial folic acid antagonists,
sulphonamides, co-trimoxazole, trimethoprim, other
diaminopyrimidine-sulphonamide combinations, nitrofurans,
15 nitrofurantoin, nitrofurazone, Gyrase inhibitors (quinolones),
norflaxacin, ciproflaxacin, ofloxacin, sparflaxacin, enoxacin,
fleroxacin, pefloxacin, lomefloxacin, Bay Y3118,
nitroimidazoles, antimycobacterial agents, isoniazid,
rifampicin, rifabutin, ethambutol, pyrazinamide, streptomycin,
20 capreomycin, prothionamide, terizidon, dapsone, clofazimine,
topical antibiotics, bacitracin, tyrothricin, polymyxins,
neomycin, kanamycin, paromomycin, mupirocin, antiviral agents,
acyclovir, ganciclovir, azidothymidine, didanosin, zalcitabin,
thiacytidine, stavudine, ribavirin, idoxuridine, trifluridine,
25 foscarnet, amantadine, interferons, tibol derivatives,
proteinase inhibitors, antifungal agents, polyenes,
amphotericin B, nystatin, natamycin, azoles, azoles for septic
treatment, miconazole, ketoconazole, itraconazole, fluconazole,
UK-109.496, azoles for topical application, clotrimazole,
30 econazole, isoconazole, oxiconazole, bifonazole, flucytosine,
griseofulvin, ciclopiroxolamine, tolnaftate, naftifine,
terbinafine, amorolfine, anthraquinones, betulinic acid,
semianthraquinones, xanthones, naphthoquinones,
aryaminoalcohols, quinine, quini-dines, mefloquine,
35 halofantrine, chloroquine, amodiaquine, acridine,

benzonaphthyridine, mepacrine, pyronaridine, dapsone,
sulphonamide, sulfadoxine, sulfalenes, trimethoprim, proguanil,
chlorproguanil, diaminopyrimidine, pyrimethamine, primaquine,
aminoquinolines, WR 238,605, tetracycline, doxycycline, clinda-
5 mycin, norfloxacin, ciprofloxacin, ofloxacin, artemisinin,
dihydroartemisinin, 10b arte methether, arte ether, arte sunate,
atovaquon, suramin, melarsoprol, nifurtimox, stibogluconate-
sodium, pentamidine, amphotericin B, metronidazole, clioquinol,
mebendazole, niclosamide, praziquantel, pyrantel, tiabendazole,
10 diethylcarbamazine, ivermectin, bithionol, oxamniquine, metri-
fonate, piperazine, embonate.

