

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 March 2004 (11.03.2004)

PCT

(10) International Publication Number
WO 2004/019991 A2

(51) International Patent Classification⁷: **A61K 47/48**

(21) International Application Number:
PCT/EP2003/009545

(22) International Filing Date: 28 August 2003 (28.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02019137.5 30 August 2002 (30.08.2002) EP

(71) Applicant: **F. HOFFMANN-LA ROCHE AG** [CH/CH];
124 Grenzacherstrasse, CH-4070 Basle (CH).

(72) Inventors: **BRANDT, Michael**; Faltergatter 29, 82393 If-
feldorf (DE). **BROCKMANN, Marc**; Glockengiesserstr.
87, 23552 Luebeck (DE). **LAMSZUS, Katrin**; Am Ree
52a, 22459 Hamburg (DE). **PAPADIMITRIOU, Apollon**;
Bachstrasse 38a, 83673 Bichl (DE). **SCHUELL, Chris-
tine**; Waldst. 1, 82377 Penzberg (DE).

(74) Agent: **SCHREINER, Siegfried**; Roche Diagnostics
GmbH, Patent Department (TR-E), P.O. Box 11 52, 82372
Penzberg (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,
UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: SCATTER FACTOR/HEPATOCYTE GROWTH FACTOR ANTAGONIST NK4 FOR THE TREATMENT OF GLIOMA

(57) Abstract: A pharmaceutical composition for the treatment of a tumor derived from glia cells of the central nervous system (CNS) of a patient, characterized in that the composition contains a pharmaceutically effective amount of a fragment of the hepatocyte growth factor, said fragment consisting of the N-terminal hairpin domain and the four kringle domains of the hepatocyte growth factor α -chain.



WO 2004/019991 A2

Scatter Factor/Hepatocyte Growth Factor antagonist NK4 for the treatment of glioma

The present invention relates to the treatment of glioma or tumor derived from glia cells of the central nervous system (CNS) by use of the scatter factor/hepatocyte growth factor antagonist NK4 and to a composition for treatment or inhibition of such disorders as well as the use of NK4 for the manufacturing of such a pharmaceutical composition.

Background of the Invention

Gliomas or tumors derived from glia cells of the central nervous system (CNS) are the most frequent malignant glioma account for approximately 30% of all primary brain tumors in adults and having a median survival rate of about 12 months, glioma are usually treated by surgery, chemotherapy, radiotherapy and/or fractionated stereotyped radio surgery.

Gliomas are heterogeneous in the cellular content and divided into groups of astrocytoma, anaplastic astrocytoma and glioblastoma multiforme. Glioma treatment is reviewed by Cher, L.M., Med. J. Aust. 175 (2001) 277-282.

Hepatocyte growth factor (HGF/SF) is a polypeptide identified and purified by Nakamura, T., et al., Biochem. Biophys. Res. Comm. 22 (1984) 1450-1459. It was further found that hepatocyte growth factor is identical to scatter factor (SF), Weidner, K.M., et al., Proc. Natl. Acad. Sci. USA 88 (1991) 7001-7005. HGF is a glycoprotein with a molecular weight of about 100 kDa involved in the development of a number of cellular phenotypes including proliferation, mitogenesis, formation of branching tubules and, in the case of tumor cells, invasion and metastasis. For a status review, see Stuart, K.A., et al., Int. J. Exp. Pathol. 81 (2000) 17-30. Both rat HGF and human HGF have been sequenced and cloned (Miyazawa, K. et al., Biochem. Biophys. Res. Comm. 163 (1989) 967-973; Nakamura, T., et al., Nature 342 (1989) 440-443; Seki, T., et al., Biochem. Biophys. Res. Comm. 172 (1990) 321-327; Tashiro, K., et al., Proc. Natl. Acad. Sci. USA 87 (1990) 3200-3204; Okajima, A., et al., Eur. J. Biochem. 193 (1990) 375-381). The pharmacokinetics and pharmacological effects of an HGF lacking the first five N-

terminal amino acids (dHGF) were investigated by Uematsu, Y., et al., J. Pharm. Sciences 88 (1999) 131-135. It was found that the serum concentration of dHGF decreased rapidly and therefore infusion would be preferred against bolus injection as administration route.

5 U.S. Patent No. 5,977,310 describes PEG-modified HGF. Such PEG-modified HGF has a prolonged clearance in vivo and has the same physiological activity as HGF. However, according to U.S. Patent No. 5,977,310, it is only possible to prolong the half life of HGF from 59.2 minutes to 76.7 minutes or 95.6 minutes, respectively (see Example 5 of U.S. Patent No. 5,977,310). It is further suggested in this patent
10 that the molar amount of the PEG reagent may be selected from the range of from 5 to 100 times of the molar weight of HGF. In the case of modifying an amino group of lysine or the N-terminus of protein, a preferred molar range of the PEG reagent is of from 10 to 25 times of the molar weight of HGF. The molecular weight of the attached PEG chain was about 10 kDa. Methods for the synthesis of conjugates
15 consisting of PEG and polypeptides such as HGF are also described in WO 94/13322. These conjugates are linked together at predefined positions as random conjugation leads, according to the authors, to the introduction of polymeric moieties into domains of the molecule that mediate the therapeutically or diagnostically desirable activities. Consequently, the molecules may acquire a
20 prolonged half-life in vivo and, in the case of heterologous proteins, reduced immunogenicity, but at the expense of a significant or complete loss of the desired biological activities (see, e.g., Kitamura, K., et al., Cancer Res. 51 (1991) 4310-4315 and Maiti, P.K., et al., Int. J. Cancer Suppl. 3 (1988) 17-22). PEGylated IFN- α shows, for example, only 7% of the potency compared to non-PEGylated IFN- α
25 (Bailon, P., Bioconjugate Chem. 12 (2001) 195-202).

It was further known from Cao, B., et al., Proc. Natl. Acad. Sci. USA 98 (2001) 7443-7448 that the growth of human glioblastoma multiforme xenograft expression Met and HGF/SF where markedly reduced in the presence of neutralizing monoclonal antibodies against HGF.

30 It was further found that an HGF/SF fragment, termed NK4, consisting of the N-terminal hairpin domain and the four kringle domains of HGF/SF has pharmacological properties that are completely different from those of HGF/SF, and is an antagonist to the influence of HGF/SF on the motility and the invasion of colon cancer cells, and is, in addition, an angiogenesis inhibitor that suppresses

tumor growth and metastasis (WO 93/23541; Parr, C., et al., *Int. J. Cancer* 85 (2000) 563-570; Kuba, K., et al., *Cancer Res.* 60 (2000) 6737-6743; Date, K., et al., *FEBS Letters* 420 (1997) 1-6; Date, K., et al., *Oncogene* 17 (1989) 3045-3054; Tomioka, D., et al., *Cancer Res.* 61 (2001) 7518-7524).

- 5 As emerges from Kuba, K., et al., *Cancer Res.* 60 (2000) 6737-6743, in animal experiments, for detecting an effect of NK4 on lung metastases, NK4 had to be infused continuously over a period of two weeks.

It is known that the attachment of polymers to certain polypeptides may increase the serum half life of such polypeptides. This was found, for example, for
10 PEGylated Interleukin-6 (EP 0 442 724) or Interleukin-2 (WO 90/07938) and erythropoietin (WO 01/02017). However, the attachment of polyethylene glycol and other polymers did not necessarily lead to prolongation of their serum half lives. It is known, for example, that the conjugation of different polyethylene glycols to Interleukin-8, G-CSF and other interleukins results in the production of
15 molecules with impaired properties (Mehvar, R., *J. Pharm. Pharmaceut. Sci.* 3 (2000) 125-136). Thus, the outcome of a PEGylation of a polypeptide is highly unpredictable. Gaertner, H.F., and Offord, R.E., *Bioconjugate Chem.* 7 (1996) 38-44 describes the site-specific attachment of PEG to the amino terminus of proteins. Gaertner et al. state (as already mentioned in WO 94/13222, see above) that
20 PEGylation presents a big problem if the attachment sites cannot be precisely controlled, as this might have important implications for protein stability and function.

Francis, G.E., et al., *Int. J. Hematol.* 68 (1998) 1-18 present an overview of PEGylation of cytokines and other therapeutic proteins. Francis et al. state that with
25 the majority of methods of PEGylation, substantial reduction of bioactivity has been reported (typically, 20-95%). According to Francis et al., PEGylation of proteins is always based on trial and error and virtually all parameters of such a PEGylation can have a surprising and very profound effect on the functionality of the product. Tsutsumi, Y., et al., *Thromb. Haemost.* 77 (1997) 168-173 describes
30 the PEGylation of Interleukin-6. According to Tsutsumi et al. about 54% of the lysine amino groups of IL-6 were coupled with PEG with a molecular weight of 5 kDa per PEG group. Tsutsumi, Y., et al., in *Proc. Natl. Acad. Sci. USA* 97 (2000) 8548-8553, describe the chemical modification of an immunotoxin by PEG. As random PEGylation was accompanied by a significant loss of specific cytotoxic

activity, Tsutsumi performs a site-specific PEGylation by using an immunotoxin mutant with one or two additional cysteins which are used for PEG coupling. Heinzerling, L., et al., Dermatol. 201 (2000) 154-157 describes the coupling of PEG to Interferon- α with a molecular weight of 5 kDa. Tsutsumi, Y., et al., in J. Pharmacol. Exp. Ther. 278 (1996) 1006-1011, describe the PEG modification of TNF- α , whereby the molecular weight of the PEG groups used is again 5 kDa. As the PEGylated TNF- α applied has a molecular weight of at least 84 kDa (by a molecular weight of 17 kDa of TNF- α) there are at least 13 5-kDa PEG groups attached to TNF- α .

PEGylation of proteins and its pharmacological effects are also reviewed by Reddy, K.R., Ann. Pharmacotherapy 34 (2000) 915-923. Again it is stated that PEGylation of therapeutic proteins must be carefully evaluated. Each protein is, according to Reddy et al., different, requires different optimization chemistry and therefore the influence of PEGylation cannot be predicted.

PEGylated NK4 and DNA and polypeptide sequences of NK4 are described in the International Patent Application No. PCT/EP02/01837.

It was an object of the present invention to provide a method for the treatment of glioma and tumors derived from glia cells and methods for manufacturing of such pharmaceutical compositions.

Summary of the Invention

It was surprisingly found that NK4 and preferably PEGylated NK4 have a strong inhibitory effect on glioma proliferative activity, invasion and angiogenesis. The present invention provides therefore the use of NK4, preferably NK4 conjugates, consisting of NK4 being covalently linked to one polyethyleneglycol (PEG) group of about 20 to 40 kDa (monoPEGylated NK4) preferably via an ϵ -amino group of NK4 lysine or the N-terminal amino group for glioma treatment.

The current invention comprises a pharmaceutical composition for the treatment of a tumor derived from glia cells of the central nervous system (CNS) of a patient, characterized in that the composition contains a pharmaceutically effective amount of a fragment of the hepatocyte growth factor, said fragment consisting of the N-

terminal hairpin domain and the four Kringle domains of the hepatocyte growth factor α -chain (NK4) and preferably monoPEGylated NK4.

5 The current invention comprises use of a fragment of the hepatocyte growth factor, said fragment consisting of the N-terminal hairpin domain and the four Kringle domains of the hepatocyte growth factor α -chain (NK4) for the manufacture of a pharmaceutical agent containing said NK4 in a pharmaceutically active amount for the treatment of a tumor derived from glia cells in the central nervous system (CNS).

10 The current invention further comprises a method for the treatment of human brain cancer (in a patient) further characterized in that a pharmaceutically effective amount of NK4, preferably monoPEGylated NK4, is administered to a patient and therefore a method for the treatment of a patient having a tumor derived from glia cells in the central nervous system (CNS), comprising administering to said patient
15 a fragment of the hepatocyte growth factor, said fragment consisting of the N-terminal hairpin domain and the four Kringle domains of the hepatocyte growth factor α -chain (NK4) in an amount effective for treatment of such tumors in such patients.

It was surprisingly found that NK4 and preferably PEGylated NK4 have superior effects in the treatment of gliomas in comparison to the treatment known in the
20 state of the art.

Detailed Description of the Invention

Human HGF is a disulfide-linked heterodimer, which can be cleaved in an α -subunit of 463 amino acids and a β -subunit of 234 amino acids, by cleavage between amino acids R494 and V495. The N-terminus of the α -chain is preceded
25 by 31 amino acids started with a methionine group. This segment includes a signal sequence of 31 amino acids. The α -chain starts at amino acid 32 and contains four kringle domains. The so-called "hairpin domain" consists of amino acids 70-96. The kringle 1 domain consists of amino acids 128-206. The kringle 2 domain consists of amino acids 211-288, the kringle 3 domain consists of amino acids 305-383, and the
30 kringle 4 domain consists of amino acids 391-469 of the α -chain, approximately. There exist variations of these sequences, essentially not affecting the biological properties of NK4 (especially not affecting its activities antagonistic to HGF and its

antiangiogenic activities), which variations are described, for example, in WO 93/23541. Also the length of NK4 can vary within a few amino acids as long as its biological properties are not affected.

5 NK4 is composed of the N-terminal 447 amino acids of the HGF/SF α -chain, which includes the above-mentioned hairpin domain and the four kringle domains. It can be produced recombinantly, either by the production of recombinant human HGF/SF and digestion with elastase (Date, K., FEBS Letters 420 (1997) 1-6) or by recombinant expression of an NK4 encoding nucleic acid in appropriate host cells, as described below. NK4 glycoprotein has a molecular weight of about 57 kDa (52
10 kDa for the polypeptide part alone) and has the in vivo biological activity of causing inhibition of tumor growth, angiogenesis and/or metastasis.

"PEGylated NK4" and "monoPEGylated NK4" as used herein therefore means that NK4 has attached covalently - and preferably one - polyethylene glycol group with a molecular weight of 20 to 40 kDa. The group can be attached, preferably randomly,
15 at different sites of the NK4 molecule, preferably, however, at the most reactive sites, e.g., the lysine side chains and the N-terminal amino group. The monoPEGylated NK4 (which therefore preferably is a mixture of monoPEGylated NK4 molecules, PEGylated at different sites which are the ϵ -amino groups of NK4 lysine and the N-terminal amino group) is at least 90% of the preparation, and
20 most preferably, the monoPEGylated NK4 is 92%, or more, of the preparation (the remaining part of the preparation is nonPEGylated NK4 and/or multiPEGylated NK4). The monoPEGylated NK4 preparations according to the invention are therefore substantially homogeneous enough to display the advantages of a homogeneous preparation, e.g., in a pharmaceutical application.


25 The PEG polymer molecules used according to the invention have a molecular weight of about 20 to 40 kDa, whereby PEG polymers with about 20, 30 or 40 kDa are preferred (by "molecular weight" as used here there is to be understood the mean molecular weight of the PEG; the term "about" indicates that in said PEG preparations, some molecules will weigh more and some less than the stated
30 molecular weight).

"PEG or PEG group" according to the invention means a residue containing poly(ethylene glycol) as an essential part. Such a PEG can contain further chemical groups which are necessary for binding reactions; which results from the chemical

synthesis of the molecule; or which is a spacer for optimal distance of parts of the molecule. In addition, such a PEG can consist of one or more PEG side-chains which are linked together. PEGs with more than one PEG chain are called multiarmed or branched PEGs. Branched PEGs can be prepared, for example, by the addition of polyethylene oxide to various polyols, including glycerol, pentaerythriol, and sorbitol. For example, a four-armed branched PEG can be prepared from pentaerythriol and ethylene oxide. Branched PEG are described in, for example, EP-A 0 473 084 and US Patent No. 5,932,462. Especially preferred are PEGs with two PEG side-chains (PEG2) linked via the primary amino groups of a lysine (Monfardini, C., et al., *Bioconjugate Chem.* 6 (1995) 62-69). As PEG polymers with a molecular weight of 20-30 kDa linear PEG molecules are preferred and as PEG polymers with a molecular weight of more than 30 kDa, especially with 40 kDa, branched PEGs are preferred. As PEG 40 kDa a two-armed PEG (PEG2) is particularly preferred.

According to the invention a method is provided for the production of a substantially homogeneous monoPEGylated NK4. PEGylation of NK4 can be performed according to the methods of the state of the art, for example by reaction of NK4 with electrophilically activated PEGs (supplier: Shearwater Corp., USA, www.shearwatercorp.com). Preferred PEG reagents are, e.g., N-hydroxysuccinimidyl propionates (PEG-SPA) or butanoates (PEG-SBA) or branched N-hydroxysuccinimides such as mPEG2-NHS (Monfardini, C., et al., *supra*). Such methods result in an NK4 polypeptide which is randomly PEGylated at an ϵ -amino group of an NK4 lysine or the N-terminal amino group. Not randomly, N-terminally PEGylated NK4 can be produced according to WO 94/01451.


In a preferred embodiment of the invention, said NK4 is covalently linked to one poly(ethylene glycol) group of the formula



with the $-\text{CO}$ (i.e. carbonyl) of the poly(ethylene glycol) group forming an amide bond with one of the amino groups of NK4; R being lower alkyl; x being 2 or 3; m being from about 450 to about 950; and n and m being chosen so that the molecular weight of the conjugate minus the NK4 protein is from about 20 to


40 kDa. As amino group of NK4 the ϵ -amino group of NK4 lysine is the available (free) amino group.

More specifically, the above conjugates may be represented by formula (I)



5 wherein P is the group of an NK4 protein as described herein, (i.e. without the amino group or amino groups which form an amide linkage with the carbonyl shown in formula (I); and wherein R is lower alkyl; x is 2 or 3; m is from about 450 to about 950 and is chosen so that the molecular weight of the conjugate minus the NK4 protein is from about 20 to 40 kDa. As used herein, the given ranges of "m" merely have an orientational meaning. The ranges of "m" are determined in any case, and exactly, by the molecular weight of the PEG group.

In a further preferred embodiment of the invention, said NK4 is covalently linked to one poly(ethylene glycol) group of the formula




15 wherein y is 1 to 4, preferably 4, n and p together are chosen such that the molecular weight of the conjugate minus the NK4 protein is from about 20 to 40 kDa, preferably 40 kDa, and n and p differ by not more than 25%, preferably by not more than 10%, and most preferably are identical, and R is lower alkyl.

20 As used herein, "lower alkyl" means a linear or branched alkyl group having from one to six carbon atoms (C_1 - C_6) alkyl). Examples of lower alkyl groups include methyl, ethyl and isopropyl. In accordance with this invention, R is any lower alkyl. Conjugates in which R is methyl are preferred.

25 The symbol "m" represents the number of ethylene oxide groups (OCH_2CH_2) in the poly(ethylene oxide) group. A single PEG sub-unit of ethylene oxide has a molecular weight of about 44 daltons. Thus, the molecular weight of the conjugate

(excluding the molecular weight of the NK4) depends on the number "m". In the conjugates of this invention "m" is from about 450 to about 950 (corresponding to a molecular weight of about 20 kDa to about 40 kDa). The number m is selected such that the resulting conjugate of this invention has a physiological activity comparable to unmodified NK4, which activity may represent the same as, more than, or a fraction of the corresponding activity of unmodified NK4. A molecular weight of "about" a certain number means that it is within a reasonable range of that number as determined by conventional analytical techniques. The number "m" is selected so that the molecular weight of each poly(ethylene glycol) group covalently linked to the NK4 protein is from about 20 kDa to about 40 kDa.

The compound of formula (I) can be prepared, for example, from a known activated polymeric material:



in which R and m are as described above, by condensing the compound of Formula II with the NK4 protein. Compounds of formula (II) in which x is 3 are alpha-lower alkoxybutyric acid succinimidyl esters of poly(ethylene glycol) (lower alkoxy-PEG-SBA). Compounds of formula (II) in which x is 2 are alpha-lower alkoxypropionic acid succinimidyl esters of poly(ethylene glycol) (lower alkoxy-PEG-SPA). Any conventional method of reacting an activated ester with an amine to form an amide can be utilized. In the reaction described above, the exemplified succinimidyl ester is a leaving group causing the amide formation. The use of succinimidyl esters such as the compounds of formula II to produce conjugates with proteins are disclosed in U.S. Patent No. 5,672,662, issued September 30, 1997 (Harris, et al.).

Human NK4 contains 30 free ε-amino groups of 30 lysine residues. When the PEGylation reagent was combined with a SBA compound of Formula II, it has been found that at a protein concentration of about 5 to 10 mg/ml, at a pH of about 7.0 to 8.0, a protein:PEG ratio of about 1:3 and a reaction temperature of from 20-25°C, a mixture of mono-, di-, and trace amounts of the tri-PEGylated species were produced. When the protein:PEG ratio was about 1:1 or 1:2 (for example,


preferably about 1:2 for 30 kDa PEG-SBA and about 1:5 for 40 kDa PEG2-NHS), primarily the monoPEGylated species is produced. By manipulating the reaction conditions (e.g., ratio of reagents, pH, temperature, protein concentration, time of reaction etc.), the relative amounts of the different monoPEGylated species can be optimized.

Typical, but not limiting, conditions are about 8 to 12 mg/ml NK4, 0.3 M potassium phosphate, pH 8, 25° C, reaction time of 1 h. Under such conditions using 30 kDa PEG-SBA (1:2, protein:PEG), the yield is about 38% monoPEGylated NK4.

10 Monopegylated NK4 can also be produced according to the methods described in WO 94/01451. WO 94/01451 describes a method for preparing a recombinant polypeptide with a modified terminal amino acid alpha-carbon reactive group. The steps of the method involve forming the recombinant polypeptide and protecting it with one or more biologically added protecting groups at the N-terminal alpha-amine and C-terminal alpha- carboxyl. The polypeptide can then be reacted with
15 chemical protecting agents to selectively protect reactive side chain groups and thereby prevent side chain groups from being modified. The polypeptide is then cleaved with a cleavage reagent specific for the biological protecting group to form an unprotected terminal amino acid alpha-carbon reactive group. The unprotected
20 terminal amino acid alpha-carbon reactive group is modified with a chemical modifying agent. The side chain protected terminally modified single copy polypeptide is then de-protected at the side chain groups to form a terminally modified recombinant single copy polypeptide. The number and sequence of steps in the method can be varied to achieve selective modification at the N- and/or C-
25 terminal amino acid of the polypeptide.

Further preferred conjugates according to the invention consist of NK4 protein being covalently linked to a lower-alkoxy poly(ethylene glycol) group via a linker of the formula $-C(O)-X-S-Y-$ with the C(O) of the linker forming an amide bond with an amino group of NK4 (as mentioned above, the ϵ -amino group of lysine residues is available), X is $-(CH_2)_k-$ or $-CH_2(O-CH_2-CH_2)_k-$, k is from 1 to 10, Y is
30

- 11 -




the average molecular weight of the poly(ethylene glycol) moiety is from about 20 kDa to about 40 kDa and the molecular weight of the conjugate is from about 72 kDa to about 92 kDa at a molecular weight of 52 kDa for NK4 polypeptide, or
 5 from about 77 kDa to about 97 kDa at a molecular weight of 57 kDa for NK4 glycoprotein.

This NK4 species may also be represented by formula (III)




wherein R may be any lower alkyl, by which is meant a linear or branched alkyl
 10 group having from one to six carbon atoms such as methyl, ethyl, isopropyl, etc. A preferred alkyl is methyl. X may be $-(CH_2)_k-$ or $-CH_2(O-CH_2-CH_2)_k-$, wherein k is from 1 to about 10. Preferably, k is from 1 to about 4, more preferably, k is 1 or 2. Most preferably, X is $-(CH_2)$.


In formula III, Y is



preferably




5 more preferably




In formula (III), the number m is selected such that the resulting conjugate of formula (III) has a physiological activity comparable to unmodified NK4, which activity may represent the same as, more than, or a fraction of the corresponding activity of unmodified NK4. m represents the number of ethylene oxide chains in the PEG unit. A single PEG subunit of $-(OCH_2CH_2)-$ has a molecular weight of about 44 daltons. Thus, the molecular weight of the conjugate (excluding the molecular weight of the NK4) depends on the number m . A molecular weight of "about" a certain number means that it is within a reasonable range of that number as determined by conventional analytical techniques. m is therefore an integer ranging from about 450 to about 950 (corresponding to a molecular weight of from about 20 to 40 kDA).


Preferred NK4 proteins of formula (III) are represented by the formulae:



and



Most preferred NK4 protein products are represented by the formula:




These NK4 proteins may be prepared by

- (a) covalently reacting a free amino group, preferably an ϵ -amino group of a lysine amino acid of an NK4 protein or the N-terminal amino group represented by the formula, P-NH₂, with a bi-functional reagent represented by the formula, Z-CO-X-S-Q, to form an intermediate with an amide linkage represented by the formula:



wherein P is an NK4 protein less the amino group which forms an amide linkage; Z is a reactive group, e.g. a carboxylic-NHS ester; X is $-(CH_2)_k-$ or $-CH_2(O-CH_2-CH_2)_k-$, wherein k is from 1 to about 10; and Q is a protecting group, like alkanoyl, e.g. acetyl.

- (b) covalently reacting the intermediate with an amide linkage from step (a) with an activated polyethylene glycol derivative represented by the formula, W-[OCH₂CH₂]_m-OR, to form an NK4 protein product represented by the formula:




wherein W is a sulfhydryl reactive form of Y; m is an integer ranging from about 450 to about 950; R is lower alkyl; and Y is as defined above.

In this embodiment, the bi-functional reagent is preferably N-succinimidyl-S-acetylthiopropionate or N-succinimidyl-S-acetylthioacetate, Z is preferably N-hydroxy-succinimide, and the activated polyethylene glycol derivative W-


$[\text{OCH}_2\text{CH}_2]_m\text{-OR}$ is preferably selected from the group consisting of iodo-acetyl-methoxy-PEG, methoxy-PEG-vinylsulfone, and methoxy-PEG-maleimide.

5 In more detail, the NK4 proteins of formula (III) may be prepared by covalent linking of a thiol group to NK4 ("activation") and coupling the resulting activated NK4 with a poly(ethylene glycol) (PEG) derivative. The first step for the preparation of monoPEGylated NK4 according to the present invention comprises covalent linking of a thiol group via NH_2 -groups of NK4. This activation of NK4 is performed with bi-functional reagents which carry a protected thiol group and an additional reactive group, such as active esters (e.g., a succinimidylester),
10 anhydrides, esters of sulphonic acids, halogenides of carboxylic acids and sulphonic acids, respectively. The thiol group is protected by groups known in the art, e.g., acetyl groups. These bi-functional reagents are able to react with the amino groups by forming an amide linkage.

15 In a preferred embodiment the activation of the amino groups is performed by reaction with bi-functional reagents having a succinimidyl moiety. The bi-functional reagents may carry different spacer species, e.g. $-(\text{CH}_2)_k-$ or $-\text{CH}_2-(\text{O}-\text{CH}_2-\text{CH}_2-)_k-$ moieties, wherein k is from 1 to about 10, preferably from 1 to about 4, and more preferably 1 or 2, and most preferably 1. Examples of these reagents are N-succinimidyl-S-acetylthiopropionate (SATP) and N-succinimidyl-S-acetylthioacetate (SATA)
20




Acetylthioalkyl-carboxylic-NHS-ester, like




SATP

or



SATA

2-(Acetylthio)-(ethoxy)_k-acetic-acid-NHS-ester

with k as defined above.

The preparation of the bi-functional reagents is known in the art. Precursors of 2-(acetylthio)-(ethoxy)_k-acetic-acid-NHS-esters are described in DE-3924705, while the derivatization to the acetylthio compound is described by March, J., Advanced Organic Chemistry (1977) 375-376. SATA is commercially available (Molecular Probes, Eugene, OR, USA and Pierce, Rockford, IL).


The addition of only one thiol group to an NK4 molecule can be selected by adjusting the reaction parameters, i.e., the protein (NK4) concentration and the protein/bi-functional reagent ratio.

The reaction is carried out, for example, in an aqueous buffer solution, pH 6.5–8.0, e.g., in 10 or 100 mM potassium phosphate, with or without 300 mM NaCl, pH 7.3. The bi-functional reagent may be added in DMSO. After completion of the reaction, preferably after 30 minutes, the reaction is stopped by addition of lysine. Excess bifunctional reagent may be separated by methods known in the art, e.g., by dialysis or column filtration. The average number of thiol groups added to NK4

can be determined by photometric methods described in, for example, Grasetti, D.R., and Murray, J.F. in *J. Appl. Biochem. Biotechnol.* 119 (1967) 41-49.


The above reaction is followed by covalent coupling of an activated polyethylene glycol (PEG) derivative. Suitable PEG derivatives are activated PEG molecules with an average molecular weight of from about 20 to about 40 kDa.

Activated PEG derivatives are known in the art and are described in, for example, Morpurgo, M., et al., *Bioconjugate Chem.* 7 (1996) 363-368 for PEG-vinylsulfone. Linear chain and branched chain PEG species are suitable for the preparation of the compounds of Formula 1. Examples of reactive PEG reagents are iodo-acetyl-methoxy-PEG and methoxy-PEG-vinylsulfone:




The use of these iodo-activated substances is known in the art and described e.g. by Hermanson, G.T., in *Bioconjugate Techniques*, Academic Press, San Diego (1996) p. 147-148.

Most preferably, the PEG species are activated by maleimide using (alkoxy-PEG-maleimide), such as methoxy-PEG-maleimide (MW 20,000 to 40,000; Shearwater Polymers, Inc.). The structure of alkoxy-PEG-maleimide is as follows:



with R and m are as defined above, preferably



The coupling reaction with alkoxy-PEG-maleimide takes place after in situ cleavage of the thiol protecting group in an aqueous buffer solution, e.g. 10 mM potassium phosphate, 300 mM NaCl, 2 mM EDTA, pH 6.2. The cleavage of the protecting group may be performed, for example, with hydroxylamine in DMSO at 25°C, pH 6.2 for about 90 minutes. For the PEG modification the molar ratio of activated NK4/alkoxy-PEG-maleimide should be from about 1:1 to about 1:6. The reaction may be stopped by addition of cysteine and reaction of the remaining thiol (-SH) groups with N-methylmaleimide or other appropriate compounds capable of forming disulfide bonds. Because of the reaction of any remaining active thiol groups with a protecting group such as N-methylmaleimide or other suitable protecting group, the NK4 proteins in the conjugates of this invention may contain such protecting groups. Generally the procedure described herein will produce a mixture of molecules having varying numbers of thiols protected by different numbers of the protecting group, depending on the number of activated thiol groups on the protein that were not conjugated to PEG-maleimide.

Whereas N-methylmaleimide forms the same type of covalent bond when used to block the remaining thiol-groups on the PEGylated protein, disulfide compounds will lead in an intermolecular sulfide/disulfide exchange reaction to a disulfide bridged coupling of the blocking reagent. Preferred blocking reagents for that type of blocking reaction are oxidized glutathione (GSSG), cysteine and cystamine. Whereas with cysteine no additional net charge is introduced into the PEGylated protein, the use of the blocking reagents GSSG or cystamine results in an additional negative or positive charge.

25 The further purification of the compounds of formula (III), including the separation of mono- from di-, tri- and multi-PEGylated NK4 species, may be done by methods known in the art, e.g., column chromatography. The percentage of mono-PEG conjugates can be controlled by pooling broader fractions around the elution peak to decrease the percentage of mono-PEG or narrower fractions to
30 increase the percentage of mono-PEG in the composition. About ninety percent

mono-PEG conjugates is a good balance of yield and activity. Compositions in which, for example, at least ninety-two percent or at least ninety-six percent of the conjugates are mono-PEG species may be desired. In an embodiment of this invention the percentage of mono-PEG conjugates is from ninety percent to ninety-six percent.

Pharmaceutical formulations

The compounds of the present invention can be formulated according to methods for the preparation of pharmaceutical compositions which methods are known to the person skilled in the art. For the production of such compositions, NK4 or monoPEGylated NK4 according to the invention is combined in a mixture with a pharmaceutically acceptable carrier. Such acceptable carriers are described, for example, in Remington's Pharmaceutical Sciences, 18th edition, 1990, Mack Publishing Company, edited by Oslo et al. (e.g. pp. 1435-1712). Typical compositions contain an effective amount of the substance according to the invention, for example from about 0.1 to 100 mg/ml, together with a suitable amount of a carrier. The compositions may be administered parenterally.

This invention further provides pharmaceutical compositions containing conjugates described herein in which the percentage of mono-PEG conjugates is preferably at least ninety percent, more preferably at least ninety-two percent.

The pharmaceutical formulations according to the invention can be prepared according to known methods in the art. Usually, solutions of monoPEGylated NK4 are dialyzed against the buffer intended to be used in the pharmaceutical composition and the desired final protein concentration is adjusted by concentration or dilution.

Such pharmaceutical compositions may be used for administration for injection and contain an effective amount of the monoPEGylated NK4 together with pharmaceutically acceptable diluents, preservatives, solubilizers, emulsifiers, adjuvants and/or carriers. Such compositions include diluents of various buffer contents (e.g. arginine, acetate, phosphate), pH and ionic strength, additives such as detergents and solubilizing agents (e.g. Tween 80/polysorbate, pluronic F68, sodium chloride, sodium sulfate), antioxidants (e.g. ascorbic acid, L-methionine), preservatives (Timersol, benzyl alcohol) and bulking substances (e.g. saccharose,

mannitol), incorporation of the material into particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc. or into liposomes. Such compositions may influence the physical state stability rate of release and clearance of the monoPEGylated NK4 according to the invention.

5 **Dosages and drug concentrations**

Typically, in a standard glioma treatment regimen, patients are treated with dosages in the range between 1 to 30 mg of NK4 or monoPEGylated NK4 per kg per day over a certain period of time, lasting from one day to about 30 days or even longer. Drug is applied as a single daily subcutaneous or i.v. bolus injection of a
10 pharmaceutical formulation containing 0.1 to 100 mg monoPEGylated NK4 per ml.

The following examples and references are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

15 **Example 1**

Recombinant production of NK4

NK4 for therapeutic uses may be produced by recombinant means using bacterial or eukaryotic expression systems. Suitable eukaryotic expression systems are for example engineered HeLa, BHK or preferably CHO cells. Cells engineered for NK4
20 production are cultivated in a suitable medium. Typically, a 1 to 5 liter cell culture is used as inoculum for a 10 liter fermenter. After 3 to 5 days, the culture in the 10 liter fermenter can be used as inoculum for the 100 liter fermenter. After additional 3 to 5 days of fermentation, this culture can be used as inoculum for the 1000 liter production fermenter. After 3 to 4 days cells are removed by filtration or
25 centrifugation and discarded. The NK4 containing supernatant is filtered, collected and processed during purification. The purification process is described in the following example.

Example 2**Purification**

Heparin-Sepharose consists of Sepharose beads to the surface of which heparin is covalently bound. Since NK4 shows a high affinity to heparin it is retained on this column and can be eluted with high salt concentrations, whereas protein contaminants and other impurities either do not bind or elute at lower salt concentrations. NK4 containing fractions, eluting at about 0.7 to 1.1 M NaCl in 50 mM Hepes pH 7.5 are collected and loaded onto a hydroxyapatite column. NK4 elutes with about 0.4 to 0.7 M potassium phosphate, pH 7.5. The resulting fractions are substantially free of contaminating proteins and can be further purified by S-sepharose chromatography.

Example 3**Production of monoPEGylated NK4**

NK4 purified in accordance with the above mentioned procedure was used for PEGylation reactions. Three of the above-mentioned suitable methods are exemplarily described.

a) PEGylation of NK4 with mPEG-SBA

Aliquots of NK4 were reacted with methoxy-PEG-SBA (5 kDa for comparison, 20 kDa and 30 kDa, respectively; Shearwater Polymers, Inc., Huntsville Alabama). Reaction was carried out at a protein to reagent ratio between 1:1 and 1:5 for about 2 h at room temperature (a ratio of 1:2 is preferred when using 20 and 30 kDa PEG). The reaction was stopped by the addition of 10 mM Tris-buffer or arginine HCl, pH 8, and samples were analyzed by SDS-PAGE or size exclusion chromatography on a Superose 6 column (Pharmacia) using as buffer solution 500 mmol/l potassium phosphate, pH 6.8, for equilibration and elution. The reaction was optimized by varying protein to reagent ratio, pH, time and temperature, in order to obtain predominantly monoPEGylated NK4.

Such conditions are, for example:

	Concentration NK4:	8-12 mg/ml
	Buffer system/pH:	0.3 M potassium phosphate, pH 8
	Temperature:	25°C
5	Reaction time:	1 h
	Molar ratios (protein:reagent):	1:2
	Yield:	
	MonoPEGylated NK4:	38%
10	DiPEGylated NK4:	17%
	UnPEGylated NK4:	45%

b) PEGylation of NK4 with mPEG-SPA

15 Aliquots of NK4 (protein concentration 8 to 12 mg/ml in 0.3 M potassium phosphate, pH 8) were reacted with methoxy-PEG-SPA (5 kDa for comparison and 20 kDa, respectively; Shearwater Polymers, Inc., Huntsville Alabama). Reaction was carried out at a protein to reagent ratio of 1:2 for about 2 h at room temperature. The reaction was stopped by the addition of 10 mM Tris-buffer or arginine HCl and samples were analyzed by SDS-PAGE, reversed phase HPLC or size exclusion
20 chromatography on a Superose 6 column (Pharmacia) using as buffer solution 500 mmol/l potassium phosphate, pH 6.8, for equilibration and elution. The reaction was optimized by varying protein to reagent ratio, pH, time and temperature, in order to obtain predominantly monoPEGylated NK4, compared to di- and tri-PEGylated NK4.

25 **c) PEGylation of NK4 with mPEG2-NHS**

This PEGylation was performed as described in Example 3c with the exception that instead of PEG-SPA mPEG2-NHS (40 kDa PEG, branched via a lysine linker) was used at a molar ratio of 1:5 (protein:PEG reagent).

Example 4**Isolation of monoPEGylated NK4**

5 MonoPEGylated NK4 can be separated from unPEGylated, di- and tri-PEGylated NK4 by running a preparative size exclusion chromatography (e.g. Superose 6 or Superdex 200; Pharmacia) using as buffer solution 500 mmol/l K-phosphate pH 6.8, for equilibration and elution, or by ion exchange chromatography. The purified protein contains predominantly the monoPEGylated species. Fractions were collected and analyzed by SDS-PAGE and reversed phase chromatography.

Example 5**10 Molecular characterization of monoPEGylated NK4****a) Size exclusion chromatography**

15 The monoPEGylated species elutes earlier in size exclusion chromatography (e.g. Superose 6 or Superdex 200; Pharmacia; using as buffer solution 500 mmol/l K-phosphate pH 6.8, for equilibration and elution) as compared to the unmodified form. This is due to an increased hydrodynamic radius of the molecule.

b) SDS-PAGE

20 In SDS-PAGE proteins are separated according to their molecular weight. Due to an increase in molecular weight by PEGylation, the monoPEGylated NK4 shows a shorter migration distance as compared to the unmodified NK4. The migration distance is inversely correlated with the chain length of the PEG moiety and the number of PEG groups attached per NK4 molecule.

c) Peptide mapping

25 Digestion of monoPEGylated NK4 with sequence-specific endo-proteinases (e.g. LysC or trypsin) results in a characteristic peptide map. The resulting peptides can be separated by reversed phase chromatography and analyzed by mass spectrometry and/or N-terminal sequencing. This allows for a determination of the PEG-modified groups within the NK4 molecule.

d) Reversed phase chromatography

MonoPEGylated NK4 can also be characterized by reversed phase chromatography. PEGylation of NK4 results in a change in retention time as compared to unmodified NK4.

5 Example 6

Inhibition of intracerebral glioblastoma growth by intratumoral injections of the Scatter Factor/Hepatocyte growth factor (SF/HGF)-antagonist NK4 using a guide screw for mice

Methods:

- 10 Guide screws were implanted into the cranial vault of 16 nude mice. U87 glioblastoma cells (1.7×10^6) were injected intracerebrally through the screw, after which the mice received daily intratumoral injections of either NK4 (25 μ g in 2.5 μ l PBS) or PBS over 19 days. Animals were sacrificed and tumor volume, vessel density, proliferative activity and apoptosis were determined histologically. The effects of NK4 on glioma and endothelial cell migration and proliferation were analyzed in vitro using modified Boyden chamber and proliferation assays.
- 15

Results:

- Tumor volume was reduced by 61 % in animals treated with NK4 compared to PBS controls ($p < 0.01$). Microvessel density was reduced by 65 % in NK4-treated mice compared to controls ($p < 0.01$), and proliferative activity of the tumor cells was reduced by 33 % compared to controls ($p < 0.05$). In vitro, NK4 showed a dose-dependent inhibition of SF/HGF-induced directional glioma cell migration (up to 100%) and of SF/HGF-induced endothelial cell proliferation.
- 20

List of References

- Bailon, P., *Bioconjugate Chem.* 12 (2001) 195-202
- Cao, B., et al., *Proc. Natl. Acad. Sci. USA* 98 (2001) 7443-7448
- 5 Cher, L. M., *Med. J. Aust.* 175 (2001) 277-282
- Date, K., et al., *FEBS Letters* 420 (1997) 1-6
- Date, K., et al., *Oncogene* 17 (1989) 3045-3054
- EP 0 473 084
- Francis, G.E., et al., *Int. J. Hematol.* 68 (1998) 1-18
- 10 Gaertner, H.F., and Offord, R.E., *Bioconjugate Chem.* 7 (1996) 38-44
- Grasetti, D.R., and Murray, J.F., *J. Appl. Biochem. Biotechnol.* 119 (1967) 41-49
- Heinzerling, L., et al., *Dermatol.* 201 (2000) 154-157
- Hermanson, G.T., in *Bioconjugate Techniques*, Academic Press, San Diego (1996)
147-148
- 15 International Patent Application No. PCT/EP02/01837
- Kitamura, K., et al., *Cancer Res.* 51 (1991) 4310-4315
- Kuba, K., et al., *Cancer Res.* 60 (2000) 6737-6743
- Maiti, P.K., et al., *Int. J. Cancer Suppl.* 3 (1988) 17-22
- March, J., *Advanced Organic Chemistry* (1977) 375-376
- 20 Mehvar, R., *J. Pharm. Pharmaceut. Sci.* 3 (2000) 125-136
- Miyazawa, K. et al., *Biochem. Biophys. Res. Comm.* 163 (1989) 967-973
- Monfardini, C., et al., *Bioconjugate Chem.* 6 (1995) 62-69
- Morpurgo, M., et al., *Bioconjugate Chem.* 7 (1996) 363-368
- Nakamura, T., et al., *Biochem. Biophys. Res. Comm.* 22 (1984) 1450-1459
- 25 Nakamura, T., et al., *Nature* 342 (1989) 440-443
- Okajima, A., et al., *Eur. J. Biochem.* 193 (1990) 375-381
- Parr, C., et al., *Int. J. Cancer* 85 (2000) 563-570
- Reddy, K.R., *Ann. Pharmacotherapy* 34 (2000) 915-923
- Remington's *Pharmaceutical Sciences*, 18th edition, 1990, Mack Publishing
30 Company, edited by Oslo et al. (e.g. pp. 1435-1712)
- Seki, T., et al., *Biochem. Biophys. Res. Comm.* 172 (1990) 321-327
- Stuart, K.A., et al., *Int. J. Exp. Pathol.* 81 (2000) 17-30
- Tashiro, K., et al., *Proc. Natl. Acad. Sci. USA* 87 (1990) 3200-3204
- Tomioka, D., et al., *Cancer Res.* 61 (2001) 7518-7524
- 35 Tsutsumi, Y, et al., *Proc. Natl. Acad. Sci. USA* 97 (2000) 8548-8553
- Tsutsumi, Y., et al., *J. Pharmacol. Exp. Ther.* 278 (1996) 1006-1011
- Tsutsumi, Y., et al., *Thromb. Haemost.* 77 (1997) 168-173

Uematsu, Y., et al., J. Pharm. Sciences 88 (1999) 131-135

US 5,672,662

US 5,932,462

US 5,977,310

5 Weidner, K.M., et al., Proc. Natl. Acad. Sci. USA 88 (1991) 7001-7005

WO 01/02017

WO 90/07938

WO 93/23541

WO 94/01451

10 WO 94/13222

Patent Claims

1. A pharmaceutical composition for the treatment of a tumor derived from glia cells of the central nervous system (CNS) of a patient, characterized in that the composition contains a pharmaceutically effective amount of a fragment of the hepatocyte growth factor, said fragment consisting of the N-terminal hairpin domain and the four kringle domains of the hepatocyte growth factor α -chain (NK4).
5
2. A pharmaceutical composition of claim 1, characterized in that it contains as NK4 monoPEGylated NK4.
- 10 3. Use of a fragment of the hepatocyte growth factor, said fragment consisting of the N-terminal hairpin domain and the four kringle domains of the hepatocyte growth factor α -chain (NK4) for the manufacture of a pharmaceutical agent containing said NK4 in a pharmaceutically active amount for the treatment of a tumor derived from glia cells in the central nervous system (CNS).
15
4. Use according to claim 3, characterized in that as NK4 monoPEGylated NK4 is used.
5. Method for the treatment of a patient having a tumor derived from glia cells in the central nervous system (CNS), comprising administering to said patient a fragment of the hepatocyte growth factor, said fragment consisting of the N-terminal hairpin domain and the four kringle domains of the hepatocyte growth factor α -chain (NK4) in an amount effective for treatment of such tumors in such patients.
20
6. Method according to claim 5, characterized in that as NK4 monoPEGylated NK4 is used.
25