(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
24 November 2022 (24.11.2022)

WIPO | PCT

(10) International Publication Number
WO 2022/245500 A1

(51) International Patent Classification:
A61K 38/20 (2006.01) *C07K 16/46* (2006.01)
A61K 47/68 (2017.01) *A61K 39/395* (2006.01)
C07K 14/54 (2006.01) *C07K 19/00* (2006.01)
C07K 16/28 (2006.01)

(21) International Application Number:
PCT/US2022/026584

(22) International Filing Date:
27 April 2022 (27.04.2022)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
63/190,669 19 May 2021 (19.05.2021) US
63/223,684 20 July 2021 (20.07.2021) US
PCT/US2021/056312 22 October 2021 (22.10.2021) US
PCT/US2021/062485 08 December 2021 (08.12.2021) US
63/297,631 07 January 2022 (07.01.2022) US

(71) Applicant: **ASHER BIOTHERAPEUTICS, INC.** [US/US]; 650 Gateway Boulevard, Suite 100, South San Francisco, California 94080 (US).

(72) Inventors: **YEUNG, Yik Andy**; 650 Gateway Boulevard, Suite 100, South San Francisco, California 94080 (US). **GREER, Renee L.**; 650 Gateway Boulevard, Suite 100, South San Francisco, California 94080 (US). **NGUYEN, Henry C.**; 650 Gateway Boulevard, Suite 100, South San Francisco, California 94080 (US). **LIU, David**; 650 Gateway Boulevard, Suite 100, South San Francisco, California 94080 (US). **KANG, Byong**; 650 Gateway Boulevard, Suite 100, South San Francisco, California 94080 (US). **DJURETIC, Ivana**; 650 Gateway Boulevard, Suite 100, South San Francisco, California 94080 (US).

(74) Agent: **HUANG, Wei**; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, California 94304 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(54) Title: IL-21 POLYPEPTIDES AND TARGETED CONSTRUCTS

FIG. 6



(57) Abstract: The present disclosure provides methods and compositions comprising IL-21 polypeptides, targeted cytokine construct that selectively activates target immune cells (e.g., CD8+ T cells) over other immune cell types. The cytokine of the present disclosure can further comprise mutations that alter its charge distribution to improve its half-life within the blood.

[Continued on next page]

WO 2022/245500 A1

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

IL-21 POLYPEPTIDES AND TARGETED CONSTRUCTS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to: US Provisional Application No. 63/190,669, filed May 19, 2021; US Provisional Application No. 63/223,684, filed July 20, 2021; PCT/US2021/056312, filed October 22, 2021; PCT/US2021/062458, filed December 8, 2021; and US Provisional Application No. 63/297,631, filed January 7, 2022. The entire contents of each application is incorporated herein by this reference.

BACKGROUND

[0002] Interleukin-21 (IL-21) is a T cell derived pleiotropic cytokine that regulates the activity of both innate and adaptive immune cells. IL-21 can augment T cell survival and effector function. Interleukin-21 is a type I cytokine and a member of the common cytokine receptor gamma-chain (γc) family that has emerged as a promising immune therapeutic for the treatment of cancer. In some cases, IL-21 that is produced by activated CD4+ T cells and natural killer T (NKT) cells signals via a heterodimeric receptor complex comprised of a discrete IL-21 receptor (IL-21R) subunit together with the γc. In some cases, activation of the IL-21R complex leads to the activation of the JAK/STAT signaling pathway. IL-21R is broadly expressed in hematopoietic cells including T and B lymphocytes, natural killer (NK) cells and myeloid cells. IL-21 is a potent mitogen and survival factor for both NK cells and activated T cells. IL-21 can support the differentiation of CD4 + T helper 17 (Th17) as well as follicular helper T cells (Tfh) and can antagonize regulatory T cell (Treg) differentiation. Additionally, IL-21 can augment the survival of CD8+ T cells resulting in a less activated but more persistent T cell phenotype that leads to enhanced tumor and viral control. A challenge of cytokine immunotherapy is that in some cases, while activating immune cells to potentiate immune responses, the same cytokine can also activate counter-regulatory pathways as exemplified by IL-2 and IFNγ. These counter-regulatory pathways can activate regulatory T cell responses and inhibitory pathways. Because it plays a key role in anti-tumor and anti-viral responses, in addition to exerting major effects on inflammatory responses that lead to the development of autoimmune diseases and inflammatory diseases, IL-21 has been an attractive target for several therapies.

[0003] There remains a need for treatment modalities utilizing IL-21, including modalities combining IL-21 moieties that direct IL-21 to specific cell types.

SUMMARY

[0004] Provided herein, in one embodiment of the disclosure is an IL-21 polypeptide or a functional fragment or a variant thereof comprising a polypeptide sequence having at least 80% sequence identity to SEQ ID NO: 1, wherein the IL-21 polypeptide has an isoelectric point that is at least about 0.6 units to about 5 units lower, compared to that of a wild-type IL-21 protein having a sequence of SEQ ID NO: 1. In some embodiments, the isoelectric point of SEQ ID NO: 1 is about 9.42. In some embodiments, the IL-21 polypeptide has an isoelectric point of about 7.12 to about 8.72. In some embodiments, the polypeptide provides an improved exposure compared to the wild-type IL-21 protein, as measured by at least about 1.5 times greater under the curve (AUC) for the polypeptide, relative to that of the wild-type IL-21, when administered to a subject, at equivalent concentrations. In some embodiments, the IL-21 polypeptide comprises at least one amino acid substitution that reduces the isoelectric point of the IL-21 polypeptide by about 0.6 units to about 5 units relative to the human IL-21 polypeptide without the amino acid substitution. In some embodiments, the IL-21 polypeptide comprises at least four amino acid substitutions that reduce the isoelectric point of the IL-21 polypeptide about 0.6 units to about 5 units relative to the human IL-21 polypeptide without the amino acid substitutions. In some embodiments, the IL-21 polypeptide comprises up to 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 amino acid substitutions that reduce the isoelectric point.

[0005] In some embodiments, the disclosure provides an IL-21 polypeptide or a functional fragment or variant thereof, wherein relative to a region of a human IL-21 polypeptide that comprises about 2 to 20 positively charged amino acid residues, the IL-21 polypeptide comprises at least one amino acid substitution of the about 2 to 20 positively charged amino acid residues. In some embodiments, the region of the human IL-21 polypeptide does not include amino acid residues that bind a human IL-21 receptor. In some embodiments, the region of the human IL-21 polypeptide comprises amino acid residues S80 to T92 of the human IL-21 polypeptide comprising SEQ ID NO: 1. In some embodiments, the disclosure provides an IL-21 polypeptide or a functional fragment or variant thereof, wherein a human IL-21 polypeptide comprises at least one positively charged amino acid residue on the surface of the human IL-21 polypeptide in a three-dimensional structure of the human IL-21 polypeptide which does not directly interact with or bind a human IL-21 receptor, and wherein the IL-21 polypeptide comprises at least one amino acid substitution of the at least one positively charged amino acid residue. In some embodiments, the disclosure provides an IL-21 polypeptide that does not comprise an amino acid substitution at G84 of the human IL-21 polypeptide comprising SEQ ID NO: 1.

[0006] In some embodiments, the IL-21 polypeptide comprises a mutation at one or more positions selected from the group consisting of: K56, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide comprises a mutation (e.g., an amino acid substitution) at one or more positions selected from the group consisting of: S80, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide comprises 4, 5, 6, 7, 8, 9, 10, 11, or 12 amino acid substitutions at positions selected from S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, or T92 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide comprises 4, 5, 6, 7, 8, 9, 10, 11, or 12 amino acid substitutions selected from S80G, T81G, N82G, N82E, A82G, A83E, A83S, R85G, R85E, R85S, R86G, R86E, Q87G, Q87E, Q87S, K88G, H89G, H89S, R90G, R90S, R90E, R90A, L91G, L91S, T92G, T92S. In some embodiments, the IL-21 polypeptide comprises (a) R85G, R86G, K88G and R90E, or (b) S80G, T81G, N82E, A83G, R85G, R86G, Q87G, K88G, H89G, R90E, L91G, and T92G. In some embodiments, the disclosure provides an IL-21 polypeptide that does not comprise an amino acid substitution at G84 of the human IL-21 polypeptide comprising SEQ ID NO: 1.

[0007] In some embodiments, the disclosure provides an IL-21 polypeptide or functional fragment or variant thereof comprising the amino acid sequence: QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKAQLKSANT GNNERIINVSIKKLKRKPPX$_1$X$_2$X$_3$X$_4$GX$_5$X$_6$X$_7$X$_8$X$_9$X$_{10}$X$_{11}$X$_{12}$CPSCDSYEKKPPKEFLERFKSL LQKMIHQHLSSRTHGSEDS (SEQ ID NO: 380), wherein X$_1$ = G, S; X$_2$ = G, T; X$_3$ = G, E, N; X$_4$ = G, S, E, A; X$_5$ = G, E, S, R; X$_6$ = G, E, R; X$_7$ = S, G, E, Q; X$_8$ = G, K; X$_9$ = G, S, H; X$_{10}$ = A, E, S, G, R; X$_{11}$ = S, G, L; and X$_{12}$ = G, S, T, provided at least one of X$_5$, X$_6$, X$_8$, X$_9$, and X$_{10}$ is not the amino acid residue at the identical position set forth in SEQ ID NO: 1, optionally wherein:

(i) X$_5$ = G, X$_6$ = G, X$_8$ = G, X$_{10}$ = A;

(ii) X$_5$ = G, X$_6$ = G, X$_8$ = G, X$_{10}$ = E;

(iii) X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = E, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = S;

(iv) X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = E;

(v) X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = E;

(vi) X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = S;

(vii) X$_5$ = G, X$_6$ = G, X$_7$ = G, X$_8$ = G, X$_9$ = G, X$_{10}$ = E;

(viii) X$_5$ = G, X$_6$ = G, X$_7$ = G, X$_8$ = G, X$_9$ = G, X$_{10}$ = G;

(ix) X$_5$ = G, X$_6$ = G, X$_7$ = E, X$_8$ = G, X$_9$ = G, X$_{10}$ = G;

(x) X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = G, X$_{11}$ = S, X$_{12}$ = G;

(xi) X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = G, X_{11} = G, X_{12} = G;

(xii) X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = G, X_6 = G, X_7 = E, X_8 = G, X_9 = G, X_{10} = G, X_{11} = S, X_{12} = G;

(xiii) X_1 = G, X_2 = G, X_3 = G, X_4 = E, X_5 = G, X_6 = G, X_7 = E, X_8 = G, X_9 = G, X_{10} = G, X_{11} = S, X_{12} = G;

(xiv) X_3 = G, X_4 = G, X_5 = S, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = G, X_{11} = G, X_{12} = S;

(xv) X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = G, X_{11} = G, X_{12} = S;

(xvi) X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = G, X_{11} = G, X_{12} = G;

(xvii) X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = G, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xviii) X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xix) X_3 = G, X_4 = G, X_5 = S, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = E, X_{11} = G, X_{12} = S;

(xx) X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xxi) X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxii) X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxiii) X_3 = G, X_4 = G, X_5 = G, X_6 = E, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxiv) X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxv) X_1 = G, X_2 = G, X_3 = G, X_4 = E, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G; or

(xxvi) X_1 = G, X_2 = G, X_3 = E, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G. In some embodiments, the disclosure provides an IL-21 polypeptide that does not comprise an amino acid substitution at G84 of the human IL-21 polypeptide comprising SEQ ID NO: 1.

[0008] In some embodiments, the disclosure provides an IL-21 polypeptide or functional fragment or variant thereof comprising an amino acid sequence at least 90%, about 95%, about 96%, about 97%, about 98%, or about 99% identical to an amino acid sequence selected from SEQ ID NOs: 2, 4-15, and 23-40. In some embodiments, the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15, and 23-40. In some embodiments, the IL-21 polypeptide comprises the amino acid sequence of SEQ ID NO: 40.

[0009] In some embodiments, the disclosure provides an IL-21 polypeptide or functional fragment or variant thereof comprising at least one amino acid substitution that reduces binding to a

human IL-21 receptor compared to binding by the human IL-21 polypeptide. In some embodiments, the at least one amino acid substitution is at one or more amino acid residues at positions R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, or K117, of SEQ ID NO: 1. In some embodiments, the at least one amino acid substitution is selected from: R5F, R5A, R5E, R5S, R5T, R5N, R5Q, R5V, R5I, R5L, R5Y, I8E, R9A, R9D, R9E, R9H, R9S, R9T, R9N, R9G, R9V, R9I, R9L, R9Y, R11D, R11E, L13F, L13R, I14D, I16A, I16S, I16R, V17I, V17A, D18A, K72A, K72E, K73A, K73E, K75A, K75E, L74I, L74F, L74M, L74V, R76E, R76F, R76A, R76N, R76D, R76S, R76T, R76Q, R76V, R76I, R76L, R76Y, R76M, K77A, K77E, and K117A. In some embodiments, the at least one amino acid substitution is R76E or R76Q. In some embodiments, the IL-21 polypeptide comprises a mutation at a position selected from the group consisting of: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to the sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, or SEQ ID NO: 15. In some embodiments, the IL-21 polypeptide comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to the sequence of SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, or SEQ ID NO: 21. In some embodiments, the IL-21 polypeptide comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from SEQ ID NOs: 2, 4-15, and 23-40. In some embodiments, the IL-21 polypeptide comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from SEQ ID NOs: 16-21, 41-98 and 374-379. In some embodiments, the IL-21 polypeptide comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from SEQ ID NOs: 16-21, 41-93 and 374-379. In some embodiments, the IL-21 polypeptide comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from SEQ ID NOs: 94-98.

[0010] Provided herein, in one embodiment of the disclosure, is a targeted cytokine construct comprising an IL-21 polypeptide or a functional fragment or a variant thereof according to the present disclosure and an antibody or an antigen binding fragment thereof. In some

embodiments, the antibody or antigen binding fragment thereof specifically binds to a CD8+ T cell. In some embodiments, the antibody or antigen binding fragment thereof specifically binds to at least one of: CD8α, CD8αα, or CD8αβ. In some embodiments, the antibody or antigen binding fragment thereof specifically binds to CD8β.

[0011] Provided herein, in one embodiment of the disclosure, is a targeted cytokine construct comprising: a) an IL-21 polypeptide or a functional fragment or a variant thereof comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1; and b) an antibody or an antigen binding fragment thereof that specifically binds to at least one of CD8α, CD8αα, or CD8αβ.

[0012] Provided herein, in one embodiment of the disclosure, is a targeted cytokine construct comprising: a) an IL-21 polypeptide or a functional fragment or a variant thereof comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1; and b) an antibody or an antigen binding fragment thereof that specifically binds to CD8β.

[0013] In some embodiments, the IL-21 polypeptide comprises a mutation at one or more amino acid positions of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide comprises a mutation at one or more positions selected from the group consisting of: K56, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1.In some embodiments, the IL-21 polypeptide comprises mutation at a position selected from the group consisting of: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to the sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, or SEQ ID NO: 15.

[0014] In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to the sequence of SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, or SEQ ID NO: 21. In some embodiments, the construct activates CD8+ T cells with at least about 10-fold or greater potency as compared to activation of NK or CD4+ T cells. In some embodiments, the construct activates CD8+ T cells with a potency that is at least about 10-fold to about 100,000-fold greater as compared to activation of NK or CD4+ T cells. In some embodiments, the antibody or antigen binding fragment thereof comprises: i) a first polypeptide, arranged from N-to-C terminus,

comprising: a variable light chain (VL) amino acid sequence, and a light chain constant region amino acid sequence (CL1); ii) a second polypeptide, arranged from N-to-C terminus, comprising: a variable heavy chain (VH) amino acid sequence, a heavy chain CH1 constant region amino acid sequence, a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence; iii) a third polypeptide, arranged from N-to-C terminus, comprising: a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence, wherein the CH2 and CH3 domains of each of the second and third polypeptides form an Fc domain. In some embodiments, the third polypeptide, arranged from N-to-C terminus, comprises: a variable heavy chain (VH) amino acid sequence, a heavy chain CH1 constant region amino acid sequence, a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, a heavy chain CH3 constant region amino acid sequence, wherein the antibody or antigen binding fragment thereof further comprises: iv) a fourth polypeptide, arranged from N-to-C terminus, comprising: a variable light chain (VL) amino acid sequence, and a light chain constant amino acid sequence (CL1). In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof and the antibody or antigen binding fragment thereof are operably linked to each other. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof is linked to the N-terminus or C-terminus of the antibody or antigen binding fragment thereof. In some embodiments, the IL-21 polypeptide or a functional fragment or variant thereof is conjugated to the C-terminus of the second or the third polypeptide. In some embodiments, the targeted cytokine construct comprises at least one molecule of the IL-21 polypeptide. In some embodiments, the Fc domain is a human IgG Fc domain. In some embodiments, the Fc domain is an IgG1, IgG2, IgG3, or IgG4 Fc domain. In some embodiments, the Fc domain comprises one or more modifications that promote heterodimerization. In some embodiments, the second polypeptide comprises a knob modification in the CH2 or the CH3 domain, and the third polypeptide comprises a hole modification in the CH2 or the CH3 domain; or wherein the third polypeptide comprises a knob modification in the CH2 or the CH3 domain and the second polypeptide comprises a hole modification in the CH2 or the CH3. In some embodiments, at least one of the second and the third polypeptide comprises the following mutations: L234A, L235A, and G237A, numbering according to the EU index.

[0015] Provided herein, in one embodiments of the disclosure, is a targeted cytokine construct comprising: a) an IL-21 polypeptide or a functional fragment or variant thereof comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1; and b) an antibody or an antigen binding fragment thereof comprising a first antigen binding arm and a

second antigen binding arm, wherein the first and the second antigen binding arms bind to two different antigens, wherein at least one of the first and the second antigen binding arm specifically binds to CD8α, CD8αα, CD8αβ.

[0016] Provided herein, in one embodiments of the disclosure, is a targeted cytokine construct comprising: a) an IL-21 polypeptide or a functional fragment or variant thereof comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1; and b) an antibody or an antigen binding fragment thereof comprising a first antigen binding arm and a second antigen binding arm, wherein the first and the second antigen binding arms bind to two different antigens, wherein at least one of the first and the second antigen binding arm specifically binds to CD8β. In some embodiments, the targeted cytokine construct activates CD8+ T cells with at least about 10-fold or greater potency as compared to activation of NK cells or CD4+ T cells. In some embodiments, targeted cytokine construct activates CD8+ T cells a potency that is at least about 10-fold to about 100,000-fold greater as compared to activation of NK cells or CD4+ T cells. In some embodiments, the IL-21 polypeptide or a functional fragment thereof or a variant thereof comprises a mutation at a position selected from the group consisting of: K56, T81, N82, A83, R85, G84, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide or a functional fragment thereof or a variant thereof comprises a mutation at a position selected from the group consisting of: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to the sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, or SEQ ID NO: 15. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to the sequence of SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, or SEQ ID NO: 21.

[0017] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and

(ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β. In some embodiments, the IL-21 polypeptide does not comprise a G84 substitution. In some embodiments, the at least one amino acid substitution providing reduced binding is R76E or R76Q.

[0018] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least four amino acid substitutions are at residues R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0019] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least four amino acid substitutions are at residues R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at a residue selected from R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, K117, and a combination thereof; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0020] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least four amino acid substitutions are at residues R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution selected from: (a) R11D; (b) R11E; (c) I14D, D18A and K117A; (d) R76E; (e) R5F; (f) R76F; (g) I8E; (h) R5A; (i) R5E; (j) R5S; (k) R5T; (l) R5N; (m) R5Q; (n) R5V; (o) R5I; (p) R5L; (q) R5Y; (r) R76A; (s) R76N; (t) R76D; (u) R76S; (v) R76T; (w) R76Q; (x) R76V; (y) R76I; (z) R76L; (aa) R76Y; (bb) K77A; (cc) K77E; (dd) K72A; (ee) K72E;

(ff) K75A; (gg) K75E; (hh) K73A; (ii) K73E; (jj) R5F and K77A; (kk) R5F and K77E; (ll) R5F and K72A; (mm) R5F and K72E; (nn) R5F and K76A; (oo) R5F and K76E; (pp) K73A and K76F; (qq) K73E and K76F; (rr) R9A; (ss) R9D; (tt) R9E; (uu) R9H; (vv) R9S; (ww) R9T; (xx) R9N; (zz) R9G; (aaa) R9V; (bbb) R9I; (ccc) R9L; (ddd) R9Y; (eee) K72A and R76F; (fff) K75A and R76F; (ggg) R76F and K77A; (hhh) K75E and R76F; (iii) V17I and L74I; (jjj) I16A and L74F; (kkk) I16S, V17I, and L74V; (lll) I16R, V17I, and L74I; (mmm) L13F, I16A, V17A and L74M; and (nnn) L13R, I16A, V17I and L74I; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0021] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are at residues S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, T92, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at a residue selected from R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, K117, and a combination thereof; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0022] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are at residues S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, T92, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at residue R76; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0023] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0024] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at a residue selected from R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, K117, and a combination thereof; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0025] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at residue R76; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0026] In some embodiments, the IL-21 polypeptide and the antibody or antigen binding fragment thereof are linked to each other via a linker. In some embodiments, the IL-21 polypeptide is linked to the N-terminus or C-terminus of the antibody or antigen binding fragment thereof. In some embodiments, the IL-21 polypeptide is linked to the C-terminus of the antibody or antigen binding fragment thereof.

[0027] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, and wherein the HC constant domain comprises a knob modification;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0028] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0029] Provided herein, in one embodiment of the disclosure, is a polynucleotide encoding an IL-21 polypeptide or a functional fragment or a variant thereof according to any one of claims 1-8.

[0030] Provided herein, in one embodiment of the disclosure, is a polynucleotide that codes for a targeted cytokine construct comprising an IL-21 polypeptide or a functional fragment or a variant thereof and an antibody or an antigen binding fragment thereof, according to the present disclosure, the polynucleotide comprising a coding sequence for the IL-21 polypeptide and a coding sequence for the antibody or an antigen binding fragment thereof.

[0031] Provided herein, in one embodiment of the disclosure, is a vector comprising the polynucleotide the present disclosure.

[0032] Provided herein, in one embodiment of the disclosure, is a host cell comprising the polynucleotide or the vector of the present disclosure.

[0033] Provided herein, in one embodiment of the disclosure, is a pharmaceutical composition comprising an IL-21 polypeptide or a functional fragment or variant thereof according to the present disclosure and a pharmaceutically acceptable carrier.

[0034] Provided herein, in one embodiment of the disclosure, is a pharmaceutical composition comprising a targeted cytokine construct according to the present disclosure and a pharmaceutically acceptable carrier.

[0035] Provided herein, in one embodiment of the disclosure, is a method for selective activation of CD8+ T cells, wherein the method comprises contacting a population of cells comprising CD8+ T cells, CD4+ T cells, and NK cells, with a targeted cytokine construct according to the present disclosure. In some embodiments, the selective activation comprises activation of CD8+ T cells with at least about 10-fold or greater potency, as compared to activation of NK cells or CD4+ T cells in the population of cells. In some embodiments, the selective activation comprises activation of CD8+ T cells with a potency that is at least about 10-fold to about 100,000-fold greater as compared to activation of NK cells or CD4+ T cells in the population of cells. In some embodiments, the selective activation of CD8+ T cells results in increased STAT3 phosphorylation of the CD8+ T cells compared to the STAT3 phosphorylation of the NK cells or CD4+ T cells in the population of cells.

[0036] Provided herein, in one embodiment of the disclosure, is a method of treating a disease in a subject, the method comprising administering an IL-21 polypeptide or a functional fragment or a variant thereof according to the present disclosure, a targeted cytokine construct described herein, or a pharmaceutical composition according to the present disclosure. In some embodiments, the method further comprises an additional therapeutic agent. In some embodiments, the disease comprises a cancer or a chronic infection. In some embodiments, the disease comprises the cancer and wherein the cancer is acute lymphoblastic leukemia (ALL) (including non T cell ALL), acute myeloid leukemia, B cell prolymphocytic leukemia, B cell acute lymphoid leukemia ("BALL"), blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloid leukemia, chronic or acute leukemia, diffuse large B cell lymphoma (DLBCL), follicular lymphoma (FL), hairy cell leukemia, Hodgkin's Disease, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, monoclonal gammopathy of undetermined significance (MGUS), multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin's lymphoma (NHL), plasma cell proliferative disorder (including asymptomatic myeloma

(smoldering multiple myeloma or indolent myeloma), plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, plasmacytomas (including plasma cell dyscrasia; solitary myeloma; solitary plasmacytoma; extramedullary plasmacytoma; and multiple plasmacytoma), POEMS syndrome (also known as Crow-Fukase syndrome; Takatsuki disease; and PEP syndrome), primary mediastinal large B cell lymphoma (PMBC), small cell- or a large cell-follicular lymphoma, splenic marginal zone lymphoma (SMZL), systemic amyloid light chain amyloidosis, T cell acute lymphoid leukemia ("TALL"), T cell lymphoma, transformed follicular lymphoma, or Waldenstrom macroglobulinemia, Mantle cell lymphoma (MCL), Transformed follicular lymphoma (TFL), Primary mediastinal B cell lymphoma (PMBCL), Multiple myeloma, Hairy cell lymphoma/leukemia, lung cancer, small-cell lung cancer, non-small cell lung (NSCL) cancer, bronchioloalveolar cell lung cancer, squamous cell cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, head and neck cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, thyroid cancer, uterine cancer, , gastrointestinal cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, endometrial carcinoma, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the cervix, carcinoma of the vagina, vulval cancer, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, bladder cancer, liver cancer, hepatoma, hepatocellular cancer, cervical cancer, salivary gland carcinoma, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwannomas, ependymomas, medulloblastomas, meningiomas, squamous cell carcinomas, pituitary adenoma and Ewing's sarcoma, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers.

[0037] In some embodiments, the antibody or antigen binding fragment comprises a heavy chain variable (VH) domain and a light chain variable (VL) domain; and wherein: (a) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:137, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:138, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:139; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:140, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:141, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:142; (b) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:149, a CDR-H2

comprising the amino acid sequence of SEQ ID NO:150, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:151; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:154; (c) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:155, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:156, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:157; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:158, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:159, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:160; (d) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:161, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:162, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:163; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:164, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:165, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:166; (e) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:167, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:168, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:169; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:170, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:171, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:172; (f) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:173, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:174, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:175; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; (g) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:179, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:180, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:181; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:182, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:183, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:184; (h) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:143, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:144, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:145; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:146, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:147, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:148; (i) the VH

domain comprises a CDR-H1 comprising the amino acid sequence of X1X2AIS, wherein X1 is S, K, G, N, R, D, T, or G, and wherein X2 is Y, L, H, or F (SEQ ID NO:185), a CDR-H2 comprising the amino acid sequence of X1X2X3PX4X5X6X7X8X9YX10QKFX11G, wherein X1 is G or H, X2 is I or F, X3 is I, N, or M, X4 is G, N, H, S, R, I, or A, X5 is A, N, H, S, T, F, or Y, X6 is A, D, or G, X7 is T, E, K, V, Q, or A, X8 is A or T, X9 is N or K, X10 is A or N, and X11 is Q or T (SEQ ID NO:186), and a CDR-H3 comprising the amino acid sequence of X1X2X3GX4X5LFX6X7, wherein X1 is D or A, X2 is A, G, E, R, Y, K, N, Q, L, or F, X3 is A, L, P, or Y, X4 is I or L, X5 is R, A, Q, or S, X6 is A or D, and X7 is D, E, A, or S (SEQ ID NO:187); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of X1X2SX3X4IX5GX6LN, wherein X1 is R or G, X2 is A or T, X3 is Q or E, X4 is E, N, T, S, A, K, D, G, R, or Q, X5 is Y or S, and X6 is A or V (SEQ ID NO:188), a CDR-L2 comprising the amino acid sequence of GX1X2X3LX4X5, wherein X1 is A or S, X2 is T, S, E, Q, or D, X3 is N, R, A, E, or H, X4 is Q or A, and X5 is S or D (SEQ ID NO:189), and a CDR-L3 comprising the amino acid sequence of QX1X2X3X4X5PWT, wherein X1 is S, N, D, Q, A, or E, X2 is T, I, or S, X3 is Y, L, or F, X4 is D, G, T, E, Q, A, or Y, and X5 is A, T, R, S, K, or Y (SEQ ID NO:190); (j) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:200, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:201; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202; (k) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207; (l) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202; (m) the VH domain comprises a CDR-H1 comprising the amino acid sequence of X1YX2MS, wherein X1 is S, D, E, A, or Q and X2 is A, G, or T (SEQ ID NO:208), a CDR-H2 comprising the amino acid sequence of DIX1X2X3GX4X5TX6YADSVKG, wherein X1 is T, N, S, Q, E, H, R, or A, X2 is Y, W, F, or H, X3 is A, S, Q, E, or T, X4 is G or E, X5 is S or I, and X6 is A or G (SEQ ID NO:209),

and a CDR-H3 comprising the amino acid sequence of X1X2X3YX4WX5X6AX7DX8, wherein X1 is S or A, X2 is N, H, A, D, L, Q, Y, or R, X3 is A, N, S, or G, X4 is A, V, R, E, or S, X5 is D or S, X6 is D, N, Q, E, S, T, or L, X7 is L, F, or M, and X8 is I, Y, or V (SEQ ID NO:210); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); (n) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); (o) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:260, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); or (p) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:252, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:253, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:254; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:255, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:256, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:257.

[0038] In some embodiments, the antibody or antigen binding fragment comprises a heavy chain variable (VH) domain and a light chain variable (VL) domain; and wherein: (a) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:226, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:151; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:154; (b) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:228, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:157; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:158, a CDR-L2 comprising the amino acid sequence of SEQ

ID NO:159, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:160; (c) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:163; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:164, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:165, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:166; (d) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:229, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:169; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:170, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:171, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:172; (e) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:230, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:231, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:175; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; (f) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:230, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:232, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:181; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:182, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:183, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:184; (g) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:224, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:225; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:140, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:141, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:142; (h) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:233, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:234, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:145; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:146, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:147, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:148; (i) the VH domain comprises a CDR-H1 comprising the amino acid sequence of GX1X2FX3X4X5, wherein X1 is G, Y, S, or A, X2 is T, S, G, R, N, or H, X3 is S, T, R, H, Y, G, or P, X4 is S, K, G, N, R, D, T, or G, and X5 is Y, L, H, or F (SEQ ID NO:235), a CDR-H2 comprising the amino acid sequence

of X1PX2X3X4X5, wherein X1 is I, N, or M, X2 is G, N, H, S, R, I, or A, X3 is A, N, H, S, T, F, or Y, X4 is A, D, or G, and X5 is T, E, K, V, Q, or A (SEQ ID NO:236), and a CDR-H3 comprising the amino acid sequence of X1X2X3GX4X5LFX6X7, wherein X1 is D or A, X2 is A, G, E, R, Y, K, N, Q, L, or F, X3 is A, L, P, or Y, X4 is I or L, X5 is R, A, Q, or S, X6 is A or D, and X7 is D, E, A, or S (SEQ ID NO:237); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of X1X2SX3X4IX5GX6LN, wherein X1 is R or G, X2 is A or T, X3 is Q or E, X4 is E, N, T, S, A, K, D, G, R, or Q, X5 is Y or S, and X6 is A or V (SEQ ID NO:188), a CDR-L2 comprising the amino acid sequence of GX1X2X3LX4X5, wherein X1 is A or S, X2 is T, S, E, Q, or D, X3 is N, R, A, E, or H, X4 is Q or A, and X5 is S or D (SEQ ID NO:189), and a CDR-L3 comprising the amino acid sequence of QX1X2X3X4X5PWT, wherein X1 is S, N, D, Q, A, or E, X2 is T, I, or S, X3 is Y, L, or F, X4 is D, G, T, E, Q, A, or Y, and X5 is A, T, R, S, K, or Y (SEQ ID NO:190); (j) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:242, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202; (k) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207; (l) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202; (m) the VH domain comprises a CDR-H1 comprising the amino acid sequence of GFTFX1X2Y, wherein X1 is S, D, E, Q, S, or A and X2 is S, D, E, A, or Q (SEQ ID NO:244), a CDR-H2 comprising the amino acid sequence of X1X2X3GX4X5, wherein X1 is T, N, S, Q, E, H, R or A, X2 is Y, W, F, or H, X3 is A, S, Q, E, or T, X4 is G or E, and X5 is S or I (SEQ ID NO:245), and a CDR-H3 comprising the amino acid sequence of X1X2X3YX4WX5X6AX7DX8, wherein X1 is S or A, X2 is N, H, A, D, L, Q, Y, or R, X3 is A, N, S, or G, X4 is A, V, R, E, or S, X5 is D or S, X6 is D, N, Q, E, S, T, or L, X7 is L, F, or M, and X8 is I, Y, or V (SEQ ID NO:246); and wherein the VL domain comprises a CDR-L1 comprising

the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); (n) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); (o) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:261, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); or (p) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:224, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:284; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:285, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:286, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:287.

[0039] In some embodiments, (a) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:109, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:110; (b) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:111, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:112; (c) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:113, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:114; (d) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:115, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:116; (e) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the

sequence of SEQ ID NO:117, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:118; (f) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:119, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:120; (g) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:123; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:124; (h) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:129, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:130; (i) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:131; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:132; (j) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:125; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:126; (k) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:127, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:128; (l) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:133; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:134; (m) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:135; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:136; (n) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:107; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:108; (o) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:121; and wherein the VL domain comprises

an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:122; or (p) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:258; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:259.

[0040] In some embodiments, (a) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(b) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(c) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(d) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

[0041] In some embodiments, (a) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:129, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:130; or (b) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the

sequence of SEQ ID NO:127, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:128.

[0042] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least four amino acid substitutions are at residues R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β, wherein the antibody or antigen binding fragment thereof comprises:

(i) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(ii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(iii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(iv) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid

sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

[0043] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least four amino acid substitutions are at residues R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at a residue selected from R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, K117, and a combination thereof; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β, wherein the antibody or antigen binding fragment thereof comprises:

(i) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(ii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(iii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(iv) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3

comprising the amino acid sequence of SEQ ID NO:288; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

[0044] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least four amino acid substitutions are at residues R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution selected from: (a) R11D; (b) R11E; (c) I14D, D18A and K117A; (d) R76E; (e) R5F; (f) R76F; (g) I8E; (h) R5A; (i) R5E; (j) R5S; (k) R5T; (l) R5N; (m) R5Q; (n) R5V; (o) R5I; (p) R5L; (q) R5Y; (r) R76A; (s) R76N; (t) R76D; (u) R76S; (v) R76T; (w) R76Q; (x) R76V; (y) R76I; (z) R76L; (aa) R76Y; (bb) K77A; (cc) K77E; (dd) K72A; (ee) K72E; (ff) K75A; (gg) K75E; (hh) K73A; (ii) K73E; (jj) R5F and K77A; (kk) R5F and K77E; (ll) R5F and K72A; (mm) R5F and K72E; (nn) R5F and K76A; (oo) R5F and K76E; (pp) K73A and K76F; (qq) K73E and K76F; (rr) R9A; (ss) R9D; (tt) R9E; (uu) R9H; (vv) R9S; (ww) R9T; (xx) R9N; (zz) R9G; (aaa) R9V; (bbb) R9I; (ccc) R9L; (ddd) R9Y; (eee) K72A and R76F; (fff) K75A and R76F; (ggg) R76F and K77A; (hhh) K75E and R76F; (iii) V17I and L74I; (jjj) I16A and L74F; (kkk) I16S, V17I, and L74V; (lll) I16R, V17I, and L74I; (mmm) L13F, I16A, V17A and L74M; and (nnn) L13R, I16A, V17I and L74I; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β, wherein the antibody or antigen binding fragment thereof comprises:

(i) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(ii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(iii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(iv) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

[0045] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are at residues S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, T92, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at a residue selected from R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, K117, and a combination thereof; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β, wherein the antibody or antigen binding fragment thereof comprises:

(i) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(ii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino

acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(iii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(iv) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

[0046] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are at residues S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, T92, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at residue R76; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

[0047] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β, wherein the antibody or antigen binding fragment thereof comprises:

(i) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid

sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(ii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(iii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(iv) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

[0048] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at a residue selected from R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, K117, and a combination thereof; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β, wherein the antibody or antigen binding fragment thereof comprises:

(i) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(ii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(iii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(iv) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

[0049] In some embodiments, the disclosure provides a fusion protein comprising:

a) an IL-21 polypeptide comprising (i) an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide, wherein the at least one amino acid substitution is at residue R76; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β, wherein the antibody or antigen binding fragment thereof comprises:

(i) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(ii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and a VL domain comprising comprises a

CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(iii) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(iv) a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

[0050] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein

at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0051] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0052] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid

sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0053] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination

thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0054] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0055] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC

constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0056] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0057] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid

sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0058] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

[0059] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

[0060] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

[0061] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2

comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

[0062] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein

at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0063] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0064] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid

sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0065] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination

thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0066] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0067] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC

constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0068] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0069] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid

sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15 and 23-40, and at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

[0070] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

[0071] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

[0072] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

[0073] In some embodiments, the disclosure provides a fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

[0074] In some embodiments, the fusion protein comprises four polypeptide chains, wherein:

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:262, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:263, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:264, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:262;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:266, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:267, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:268, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:266;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:270, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:271, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:272, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 270;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:275, the third

polypeptide chain comprises the amino acid sequence of SEQ ID NO:276, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274; or

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:278, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:279, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:280, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 278.

[0075] In some embodiments, the fusion protein comprises four polypeptide chains, wherein:

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:262, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:263, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:265, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:262;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:266, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:267, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:269, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:266;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:270, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:271, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:273, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 270;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:275, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:277, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274; or

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:278, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:279, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:281, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 278.

[0076] In some embodiments, the fusion protein comprises four polypeptide chains, wherein:

(a) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:297, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:298, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:299, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:297;

(b) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:301, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:302, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:303 and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:301;

(c) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:305, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:306, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:307, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 305; or

(d) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:310, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:311, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309.

[0077] In some embodiments, the fusion protein comprises four polypeptide chains, wherein:

(a) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:297, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:298, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:300, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:297;

(b) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:301, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:302, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:304 and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:301;

(c) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:305, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:306, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:308, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 305; or

(d) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:310, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:312, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309.

INCORPORATION BY REFERENCE

[0078] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0079] The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0080] **FIGS. 1A-1C** show the amino acid sequences of the following polypeptides: mature IL-21 (SEQ ID NO: 1) (**FIG. 1A**), IL-21R (SEQ ID NO: 381) (**FIG. 1B**), and the common gamma chain (SEQ ID NO: 382) (**FIG. 1C**).

[0081] **FIG. 2** shows the amino acid sequences of the mature wild-type IL-21 polypeptide (SEQ ID NO: 1) and the modifications to generate charge variants. "X" denotes the amino acid position substituted in the sequence of wild-type IL-21 polypeptide to generate charge variants.

[0082] **FIGS. 3A -3F** show ion exchange chromatography traces of fusion proteins of anti-CD8 antibodies and either wild type IL-21 or IL-21 charge variants after Protein A purification. Fusion proteins of xhCD8 and either wild type IL-21 or exemplary IL-21 charge variants v1 through v6 are shown in **FIG. 3A**, as measured by HPLC. Fusion proteins of xhCD8.1 and exemplary IL-21 charge variants v2, or variants v7 through v31 are shown in **FIGS. 3B** through **3E**, and fusion protein of xhCD8.1 and wild type IL-21 is shown in **FIG. 3F**, as measured by FPLC. In all cases, the peak(s) on the left are undesired products, primarily homodimer, whereas the peak on the right is the desired heterodimer fusion protein, and the percentages of total of each denoted region are shown.

[0083] **FIGS. 4A-4D** show mouse pharmacokinetic data for fusion proteins. Serum concentrations are shown for 1mg/kg of a control antibody (xCtrl) with either the wild-type IL-21 or exemplary charge variant IL-21 polypeptide IL-21 v2 dosed either intravenously **(FIG. 4A)** or subcutaneously (**FIG. 4B**). In a separate experiment, serum concentrations are additionally shown for 1mg/kg wild type IL-21, or exemplary charge variant IL-21 polypeptides, IL-21 v2 or IL-21 v31 dosed subcutaneously (**FIG. 4C**) or intravenously (**FIG. 4D**).

[0084] **FIGS. 5A-5B** show the activation of STAT3 by wild-type IL-21 in CD8+ T cells , CD4+ T cells , NK cells and B cells from human peripheral blood mononuclear cells (PBMCs)

(**FIG. 5A**) or from whole blood (**FIG. 5B**). STAT3 activation was measured by flow cytometry, and all cell types were potently activated by wild-type IL-21.

[0085] **FIG. 6** is a schematic showing the general mechanism for how targeted fusions of mutant IL-21 polypeptides with CD8 antigen binding molecules and untargeted fusions with mutant IL-21 polypeptides work to stimulate cells expressing or not expressing CD8 antigens.

[0086] **FIG. 7** shows activation of STAT3 in human CD8+ T cells by fusion proteins of anti CD8 antibody with either wild-type IL-21, exemplary IL-21 polypeptide charge variants IL-21 v1, IL-21 v2 or IL-21 v6.

[0087] **FIGS. 8A-8F** show activation of STAT3 in human PBMCs by fusion proteins of an anti-human CD8 antibody, xhCD8, with various attenuated versions of IL-21 v2. STAT3 activation is shown for CD8+ T cells, CD4+ T cells, NK cells and B cells. Fusion proteins of xhCD8 with exemplary IL-21 polypeptides comprising attenuation mutations, IL-21 v2.1 (**FIG. 8A**), IL-21 v2.2 (**FIG. 8B**), IL-21 v2.3 (**FIG. 8C**), IL-21 v2.4 (**FIG. 8D**), IL-21 v2.5 (**FIG. 8E**) and IL-21 v2.6 (**FIG. 8F**), all show preferential STAT3 activation of CD8+ T cells over NK, B and CD4+ T cells, as measured by EC_{50} as well as maximal STAT3 activation.

[0088] **FIGS. 9A-9H** show activation of STAT3 in human PBMCs by fusion proteins of an anti-human CD8 antibody, xhCD8.1, with additional attenuated versions of IL-21 v2 or untargeted Fc fusion to IL-21 v2 as a control. STAT3 activation is shown in CD8+ T cells in **FIGS. 9A, 9C, 9E** and **9G**, and in CD4+ T cells in **FIGS. 9B, 9D, 9F,** and **9H.**

[0089] **FIGS. 10A-10F** show activation of STAT3 in human whole blood by fusion proteins of an anti-human CD8 antibody, xhCD8.1, with various attenuated versions of IL-21 v31. STAT3 activation is shown for CD8+ T cells, CD4+ T cells , NK cells, B cells and myeloid cells. Fusion proteins of xhCD8.1 with exemplary IL-21 polypeptides comprising attenuation mutations, IL-21 v31.4 (**FIG. 10A**), IL-21 v31.6 (**FIG. 10B**), IL-21 v31.23 (**FIG. 10C**), IL-21 v31.48 (**FIG. 10D**), and IL-21 v31.51 (**FIG. 10E**), all show preferential STAT3 activation of CD8+ T cells over NK, B and CD4+ T cells, as measured by EC_{50} as well as maximal STAT3 activation. In the case of xhCD8.1-hIL21 v31.6, data from two separate experiments are shown (**FIG. 10B**). Untargeted, unattenuated Fc-hIL21 v2 is shown in **FIG. 10F,** with activation observed in all cell types shown.

[0090] **FIGS. 11A-11C** show activation of STAT3 in mouse splenocytes. STAT3 activation is shown for CD8+ T cells, CD4+ T cells, NK cells and B cells. Activation profiles are shown for recombinant wild type mouse IL-21 (**FIG. 11A**); fusion protein of untargeted control antibody, xCtrl, and mouse IL-21 charge variant mIL-21 v1 (**FIG. 11B**); and fusion protein of anti-mouse CD8 antibody, xmCD8, and attenuated mouse IL-21 charge variant, mIL21 v1.1 (**FIG. 11C**). For both recombinant wild type mouse IL-21 and xCtrl-mIL21 v1, all cell types are

activated, whereas for xmCD8-mIL21 v1.1, there is preferential STAT3 activation of CD8+ T cells over NK, B, and CD4+ T cells, as measured by as measured by EC_{50} as well as maximal STAT3 activation.

[0091] **FIG. 12** shows tumor growth curves in an MC38 syngeneic tumor model treated with a single dose of either PBS; 0.3 or 1 mg/kg (mpk) of xCtrl-mIL21 v1, a fusion protein of untargeted control antibody, xCtrl, and unattenuated mouse IL-21 charge variant, IL-21 v1; or 0.1, 0.3 or 1mpk of xmCD8-mIL21 v1.1, a fusion protein of anti-mouse CD8 antibody, xmCD8, and attenuated mouse IL-21 charge variant, IL21 v1.1. Each subpanel shows tumor growth curves from each individual animal, the fraction showing complete responses (CR), and an arrow denoting the treatment time.

[0092] **FIGS. 13A-13D** show the impact of FTY720 on the anti-tumor potency of xmCD8-mIL21 v1.1 in an MC38 tumor model. **FIG. 13A** shows the dosing scheme for FTY720 and xmCD8-mIL21 v1.1, and tumor growth curves for 0.1 mpk (**FIG. 13B**), 0.3 mpk (**FIG. 13C**), or 1 mpk (**FIG. 13D**) xmCD8-mIL21 v1.1. Plotted values are the mean tumor volume and error bars show the standard error of the mean, with each group comprising 8 animals.

[0093] **FIGS. 14A-14F** show activation of STAT3 in human whole blood from additional donors by fusion proteins of an anti-human CD8 antibody, xhCD8.1, with various attenuated versions of IL-21 v31. STAT3 activation is shown for CD8+ T cells, CD4+ T cells, NK cells, and B cells. Fusion proteins of xhCD8.1 with exemplary IL-21 polypeptides comprising attenuation mutations, IL-21 v31.4 (**FIG. 14A**), IL-21 v31.6 (**FIG. 14B**), IL-21 v31.23 (**FIG. 14C**), IL-21 v31.48 (**FIG. 14D**), and IL-21 v31.51 (**FIG. 14E**), all show preferential STAT3 activation of CD8+ T cells over NK, B and CD4+ T cells, as measured by EC_{50} . In the case of xhCD8.1-hIL21 v31.4 and xhCD8.1-hIL21 v31.23, data from two different donors analyzed in two separate experiments are shown (**FIGS. 14A & 14C**). Untargeted, unattenuated Fc-hIL21 v31 is shown in **FIG. 14F,** with activation observed in all cell types shown.

[0094] **FIGS. 15A-15D** show activation of STAT3 in human whole blood by fusion proteins of an anti-human CD8 antibody, xhCD8v11, with various attenuated versions of IL-21 v31. STAT3 activation is shown for CD8+ T cells, CD4+ T cells, NK cells, and B cells. Fusion proteins of xhCD8v11 with exemplary IL-21 polypeptides comprising attenuation mutations, IL-21 v31.4 (**FIG. 15A**), IL-21 v31.6 (**FIG. 15B**), IL-21 v31.23 (**FIG. 15C**), and IL-21 v31.51 (**FIG. 15D**), all show preferential STAT3 activation of CD8+ T cells over NK, B and CD4+ T cells, as measured by EC_{50}. In the case of xhCD8v11-hIL21 v31.23, data from two different donors analyzed in two separate experiments are shown (**FIG. 15C**).

[0095] **FIGS. 16A-16C** show activation of STAT3 in human whole blood by fusion proteins of an anti-human CD8 antibody, xhCD8.1, with various versions of IL-21 v2 charge variant that also contained additional mutations. STAT3 activation is shown for CD8+ T cells, CD4+ T cells, NK cells, and B cells. Fusion proteins of xhCD8v.1 with exemplary IL-21 polypeptides comprising attenuation mutations, IL-21 v2.63 (**FIG. 16A**), IL-21 v2.64 (**FIG. 16B**), and IL-21 v2.65 (**FIG. 16C**) all show preferential STAT3 activation of CD8+ T cells over NK, B and CD4+ T cells, as measured by EC_{50}.

[0096] **FIGS. 17A-17C** show ion exchange chromatography traces of fusion proteins of anti-CD8 antibodies and IL-21R attenuated IL-21 charge variants after Protein A purification. Traces are shown for xhCD8v11-hIL21 v31.23 (**FIG. 17A**), xhCD8.1-hIL21 v31.23 (**FIG. 17B**), and xhCD8.1-hIL21 v31.4 (**FIG. 17C**), as measured by FPLC. In all cases, the peak(s) to the left of the major peak are undesired products, primarily homodimer, whereas the major peak in the middle is the desired heterodimer fusion protein, and the percentages of total of the desired peak are shown.

[0097] **FIG. 18** shows the pharmacokinetic data for fusion proteins in wild-type C57BL6 mice. Serum concentrations are shown for fusion proteins of anti-human CD8 antibodies, xhCD8.1 or xhCD8v11, with either hIL-21 v31.4, hIL-21 v31.23, or hIL-21 v0.4 dosed intravenously. The anti-human CD8 antibodies, xhCD8.1 and xhCD8v11 are not cross reactive to mouse CD8. Exposures are shown for xhCD8.1-hIL21 v31.4, xhCD8v11-hIL21 v31.23, xhCD8.1-hIL21 v31.23 and xhCD8.1-hIL21 v0.4.

[0098] **FIGS. 19A & 19B** show the characterization data of recombinant cytokine variants based on wild-type IL-21 or the IL-21 v31 charge variant. SDS-PAGE of material purified by IMAC followed by preparative SEC is shown in **FIG. 19A.** Analytical SEC of material purified by IMAC followed by preparative SEC is shown in **FIG. 19B.** Retention times and peak widths are also reported to evaluate column interactions. In summary, recombinant cytokine variants based on the IL-21 v31 charge variant have lower retention times and narrower peak widths compared to variants based on wild-type IL-21, suggesting fewer column interactions and more favorable biophysical properties with IL-21 v31 based constructs.

[0099] **FIG. 20** shows the exposure of recombinant (non-fusion) wildtype human IL-21 and IL-21 v31 charge variant in mouse. IL-21 v31 shows increased in vivo exposure compared to wildtype IL-21.

[0100] **FIGS. 21A and 21B** each provide a structural schematic showing the interaction between wild-type IL-21 (SEQ ID NO: 1) and IL-21R (SEQ ID NO: 381). Residues S80-T92 of SEQ ID NO: 1 form a largely unstructured, flexible loop (circled). **FIG. 21A** indicates residues P78

and P79 (Pro-Pro), and C93 and P94 (Cys-Pro) surrounding the flexible loop. **FIG. 21B** shows the charge of wild-type IL-21 indicating the flexible loop is highly charged.

[0101] **FIG. 22** shows the activities of IL-21 fusion molecules in cultures of alloreactive T cells and activated dendritic cells isolated from two different human PBMC donors. Molecules tested include two CD8+ T cell targeted fusions, xhCD8.1-hIL21 v31.4 and xhCD8.1-hIL21 v31.23, and an untargeted, unattenuated Fc-hIL21 v31. IL-2 release in culture was measured to determine the level of alloreactive T cell activation. In summary, Fc-hIL21 v31 suppressed activation of alloreactive T cells, while the CD8+ T cell targeted fusions did not.

[0102] **FIGS. 23A-23D** show in vitro activation of STAT3 in cynomolgus monkey whole blood by fusion proteins of anti-human CD8 antibodies, xhCD8.1 or xhCD8v11, with various attenuated versions of IL-21 v31. STAT3 activation is shown for CD8+ T cells, CD4+ T cells, and B cells. CD8+ T cell targeted fusion proteins, xhCD8.1-IL-21 v31.4 (**FIG. 23A**), xhCD8.1-IL-21 v31.23 (**FIG. 23B**) and xhCD8v11-IL-21 v31.23 (**FIG. 23C**), all show preferential STAT3 activation of CD8+ T cells over B and CD4+ T cells, as measured by EC_{50}. Untargeted, unattenuated Fc-hIL21 v31 is shown in **FIG. 23D,** with activation observed in all cell types shown.

[0103] **FIG. 24** shows in vivo activation of STAT3 in cynomolgus monkey by xhCD8.1-IL-21 v31.4. STAT3 activation levels in CD8+ T cells, CD4+ T cells, NK cells and B cells were assessed 20 min post-dose. Results show that xhCD8.1-IL-21 v31.4 preferentially activates STAT3 in CD8+ T cells over CD4+ T cells, NK cells and B cells, as measured by STAT3 MFI.

DETAILED DESCRIPTION OF THE INVENTION

[0104] This disclosure, in some embodiments, provides IL-21 polypeptides or functional fragments or variants thereof that comprise improved properties relative to a wild-type IL-21 (SEQ ID NO: 1). Examples of improved properties include, but are not limited to: (i) improved systemic exposure (*e.g.*, as measured by an increased area under the curve (AUC) upon administering the IL-21 polypeptide or a functional fragment or variant thereof to a subject, relative to that following administration of a wild-type IL-21 protein); (ii) attenuated binding to IL-21 receptor; (iii) reduced cytotoxicity, or any combination thereof.

[0105] As described herein, the present disclosure provides IL-21 polypeptides or functional fragments or variants thereof with reduced charge (*i.e.,* isoelectric point) and reduced binding affinity to human IL-21 receptor relative to a human IL-21 polypeptide (SEQ ID NO: 1). In particular, the disclosure provides IL-21 polypeptides with one or more amino acid substitutions in a positively charged region of located within residues S80-T92 of human IL-21. As shown in **FIG. 21B**, residues within S80-T92 of human IL-21 form a flexible loop that does not directly

interact with the human IL-21 receptor. Without wishing to be bound by theory, it is believed that residues P78, P79, C93 and P94 of human IL-21 provide structural rigidity surrounding the loop. The present disclosure provides IL-21 polypeptides having one or more amino acid substitutions within the flexible loop with a reduced isoelectric point and improved bioavailability and biological activity. As described herein, the IL-21 polypeptides of the disclosure having reduced isoelectric points relative to the human IL-21 polypeptide demonstrate increased circulatory half-life and reduced polyreactivity (*e.g.,* with positively charged proteins abundant in circulation). As shown in **FIGs. 4A-4D,** fusion proteins with IL-21 polypeptides having reduced isoelectric points relative to human IL-21 have improved half-life. The disclosure further provides fusion proteins of IL-21 polypeptides to anti-CD8 antibodies which selectively activate CD8+ T cells over CD4+ T cells, NK cells and B cells, including *in vivo* in non-human primates (*see* **FIG. 24**). Without being bound by theory, fusion proteins of the IL-21 polypeptides with antigen binding domains that specifically bind CD8ab or CD8b selectively activate CD8+ T cells relative to NK cells which express CD8aa. In addition, the IL-21 polypeptides with reduced isoelectric points have lower binding affinities to proteins abundant in circulation, such as heparin and hemoglobin (*see* **Table 5**). Further, without being bound by theory, IL-21 polypeptides of disclosure having glycine and/or serine substitutions in the flexible loop may have reduced immunogenicity relative to unmodified human IL-21 following administration to a subject.

[0106] The disclosure also provides IL-21 polypeptides having reduced binding to human IL-21 receptors. As shown herein, attenuated binding of the IL-21 polypeptide of the disclosure to its receptor reduces IL-21 mediated suppressive effects on alloreactive T cells and dendritic cells (*see* **FIG. 22**). Without wishing to be bound by theory, attenuated binding of the IL-21 polypeptide of the disclosure to its receptor is expected to avoid the sink effect of binding to B cells and NK cells. Also demonstrated herein is the therapeutic efficacy of fusion proteins comprising the IL-21 polypeptides of the disclosure and anti-CD8 targeting antibodies. Without being bound by theory, linking of the IL-21 polypeptide with attenuated binding to an anti-CD8 targeting antibody provides selective restoration of the attenuated IL-21 polypeptide. Specifically, **FIG. 12** shows such fusion proteins provide anti-tumor efficacy *in vivo*. This efficacy results, at least in part, from peripheral activation of CD8+ T cells as blocking lymphocyte trafficking reduces the curative response (*see* **FIGs. 13A-13D**). Without being bound by theory, T cells in the tumor microenvironment may be dysfunctional and express inhibitory molecules, and thus peripheral activation of CD8+ T cells by the fusion protein provides the anti-tumor immune response. Accordingly, the present disclosure provides novel IL-21 polypeptides and fusion proteins

comprising the IL-21 polypeptides having improved biological properties, as well as methods of using such polypeptides to treat diseases in subjects.

[0107] Various embodiments of the present disclosure provide methods and compositions involving the use of cytokines (*e.g.*, interleukin 21 (IL-21) or IL-21 polypeptides or functional fragments or variants thereof comprising one or more mutated amino acid residues relative to a wild-type IL-21 polypeptide sequence) that are directed or targeted to activate specific cell subsets (*e.g.*, CD8+ T cells), hereby referred to as targeted cytokine constructs. Without wishing to be bound by a certain theory, it is contemplated that being able to selectively activate certain cell types may enable a more robust response as well as the ability to limit the activation of other cell types that would be counter-productive to a successful immune response. In some cases, the methods and compositions disclosed herein allow for targeted delivery of IL-21 polypeptides to desired cell types and for administration of IL-21 at relatively low concentrations to elicit selective activation of CD8+ T cells. In some cases, use of targeted cytokine constructs disclosed herein provide more selective cell (such as a CD8+ T cells) activation compared to activation of other cell types, such as natural killer (NK) cells, CD4+ T cells or B cells.

Definitions

[0108] Unless otherwise defined herein, scientific and technical terms used in connection with this disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. The meaning and scope of the terms should be clear, however, in the event of any latent ambiguity, definitions provided herein take precedent over any dictionary or extrinsic definition. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. In this application, the use of "or" means "and/or" unless stated otherwise. Furthermore, the use of the term "including", as well as other forms, such as "includes" and "included", is not limiting.

[0109] Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well-known and commonly used in the art. The methods and techniques of the present disclosure are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclatures used in connection with, and the laboratory procedures and techniques of, analytical

chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well-known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

[0110] As used in the specification and claims, the singular forms "a," "an," and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a chimeric transmembrane receptor polypeptide" includes a plurality of chimeric transmembrane receptor polypeptides.

[0111] The term "about" or "approximately" means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. For example, "about" can mean within 1 or more than 1 standard deviation, per the practice in the art. Alternatively, "about" can mean a range of up to 20%, up to 10%, up to 5%, or up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value. Where particular values are described in the application and claims, unless otherwise stated, the term "about" meaning within an acceptable error range for the particular value should be assumed.

[0112] As used herein, "area under the curve" or "AUC" is a pharmacokinetic parameter that refers to the area under the plot of plasma concentration of a drug (*e.g.*, IL-21 polypeptide or fusion protein described herein) versus time after dosage. In some embodiments, the AUC measures a patient's exposure to a drug and depends on dose, bioavailability, and clearance. Methods for determining the AUC are known to those of skill in the art and include, but is not limited to, the Rectangle Method, the Trapezoidal Rule, and Simpson's Rule.

[0113] A "cytokine" is a form of immunomodulatory polypeptide that can mediate cross-talk between initiating/primary cells and target/effector cells. It can function as a soluble form or cell-surface associated to bind the "cytokine receptor" on target immune cells to activate signaling. "Cytokine receptor" as used here is the polypeptide on the cell surface that activates intracellular signaling upon binding the cytokine on the extracellular cell surface. Cytokines can include, but are not limited to, chemokines, interferons, interleukins, lymphokines, and tumor necrosis factors. Cytokines are produced by a wide range of cells, including immune cells, endothelial cells, fibroblasts, and stromal cells. A given cytokine may be produced by more than one cell type. Cytokines are pleiotropic; since the receptors are expressed on multiple immune cell subsets, one cytokine can activate the signaling pathway in multiple cells. However, depending on the cell type,

the signaling events for a cytokine can result in different downstream cellular events such as activation, proliferation, survival, apoptosis, effector function and secretion of other immunomodulatory proteins. A given cytokine, in some embodiments, is a wild-type cytokine polypeptide, a fragment thereof, or a variant thereof, such as a mutated cytokine polypeptide (also referred to herein as a mutein cytokine, *e.g.*, a mutein IL-21 or an IL-21 mutein).

[0114] The term "antigen," as used herein, refers to a molecule or a fragment thereof capable of being bound by a selective binding agent. As an example, an antigen can be a ligand that can be bound by a selective binding agent such as a receptor. As another example, an antigen can be an antigenic molecule that can be bound by a selective binding agent such as an immunological protein (*e.g.*, an antibody). An antigen can also refer to a molecule or fragment thereof capable of being used in an animal to produce antibodies capable of binding to that antigen.

[0115] The term "antibody", as used herein, means any antigen-binding molecule comprising at least one complementarity determining region (CDR) that specifically binds to or interacts with a particular antigen. The term "antibody" includes immunoglobulin molecules comprising four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, as well as multimers thereof (e.g., IgM). Each heavy chain comprises a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region comprises three domains, CH1, CH2 and CH3. Each light chain comprises a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region comprises one domain (CL1). The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. In different embodiments of the disclosure, the FRs of an antibody (or antigen-binding portion thereof) may be identical to the human germline sequences, or may be naturally or artificially modified. An amino acid consensus sequence may be defined based on a side-by-side analysis of two or more CDRs.

[0116] The term "intact antibody" refers to an antibody comprising four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. In one embodiment, the antibody is an intact antibody. In one embodiment, the intact antibody is an intact human IgG1, IgG2 or IgG4 isotype. In certain embodiments, the antibody, or antigen-binding fragment thereof, is a human IgG1, IgG2, or IgG4 isotype.

[0117] The terms "antigen-binding portion" of an antibody, "antigen-binding fragment," or "antibody-fragment," of an antibody, and the like, as used herein, include any naturally occurring, enzymatically obtainable, synthetic, or genetically engineered polypeptide or glycoprotein that specifically binds an antigen to form a complex. Antigen-binding fragments of an antibody may be derived, e.g., from intact antibody molecules using any suitable standard techniques such as proteolytic digestion or recombinant genetic engineering techniques involving the manipulation and expression of DNA encoding antibody variable and optionally constant domains. Such DNA is known and/or is readily available from, e.g., commercial sources, DNA libraries (including, e.g., phage-antibody libraries), or can be synthesized. The DNA may be sequenced and manipulated chemically or by using molecular biology techniques, for example, to arrange one or more variable and/or constant domains into a suitable configuration, or to introduce codons, create cysteine residues, modify, add or delete amino acids, etc.

[0118] Non-limiting examples of antigen-binding fragments include: (i) Fab fragments; (ii) F(ab')2 fragments; (iii) Fd fragments; (iv) Fv fragments; (v) single-chain Fv (scFv) molecules; (vi) dAb fragments; and (vii) minimal recognition units consisting of the amino acid residues that mimic the hypervariable region of an antibody (e.g., an isolated complementarity determining region (CDR) such as a CDR3 peptide), or a constrained FR3-CDR3-FR4 peptide.

[0119] The term "variable region" or "variable domain" of an antibody, or fragment thereof, as used herein refers to the portions of the light and heavy chains of antibody molecules that include amino acid sequences of complementarity determining regions (CDRs; i.e., CDR-1, CDR-2, and CDR-3), and framework regions (FRs). VH refers to the variable domain of the heavy chain. VL refers to the variable domain of the light chain. According to the methods used in this disclosure, the amino acid positions assigned to CDRs and FRs may be defined according to Kabat (Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md., 1987 and 1991)). Amino acid numbering of antibodies or antigen binding fragments is also according to that of Kabat.

[0120] The term "complementarity determining regions" or "CDRs" as used herein refers to the complementarity determining region within antibody variable sequences. There are three CDRs in each of the variable regions of the heavy chain and the light chain, which are designated CDR1, CDR2 and CDR3, for each of the variable regions. The term "CDR set" as used herein refers to a group of three CDRs that occur in a single variable region capable of binding the antigen. The exact boundaries of these CDRs have been defined differently according to different systems. The system described by Kabat (Kabat et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987) and (1991)) not only provides an unambiguous residue

numbering system applicable to any variable region of an antibody, but also provides precise residue boundaries defining the three CDRs. These CDRs may be referred to as Kabat CDRs. Chothia and coworkers (Chothia et al., J. Mol. Biol. 196:901-917 (1987) and Chothia et al., Nature 342:877-883 (1989)) found that certain sub- portions within Kabat CDRs adopt nearly identical peptide backbone conformations, despite having great diversity at the level of amino acid sequence. These sub-portions were designated as L1, L2 and L3 or H1, H2 and H3 where the "L" and the "H" designates the light chain and the heavy chains regions, respectively. These regions may be referred to as Chothia CDRs, which have boundaries that overlap with Kabat CDRs. Other boundaries defining CDRs overlapping with the Kabat CDRs have been described by Padlan (FASEB J. 9:133-139 (1995)) and MacCallum (J Mol Biol 262 (5):732-45 (1996)). Still other CDR boundary definitions may not strictly follow one of the above systems, but will nonetheless overlap with the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. The methods used herein may utilize CDRs defined according to any of these systems, although preferred embodiments use Kabat or Chothia defined CDRs.

[0121] The term "framework regions" (hereinafter FR) as used herein refers to those variable domain residues other than the CDR residues. Each variable domain typically has four FRs identified as FR1, FR2, FR3 and FR4. Common structural features among the variable regions of antibodies, or functional fragments thereof, are well known in the art. The DNA sequence encoding a particular antibody can generally be found following well known methods such as those described in Kabat, et al. 1987 Sequence of Proteins of Immunological Interest, U.S. Department of Health and Human Services, Bethesda MD, which is incorporated herein as a reference. In addition, a general method for cloning functional variable regions from antibodies can be found in Chaudhary, V.K., et al., 1990 Proc. Natl. Acad. Sci. USA 87:1066, which is incorporated herein as a reference.

[0122] A "variant Fc region" comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one "amino acid modification" as herein defined. A variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, e.g., from about one to about ten amino acid substitutions, from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. In one embodiment, a variant Fc region herein can have a sequence that has at least about 80% identity, at least about 85% identity, at least about 90% identity, at least about 95% identity or at least about 99% identity with a native sequence Fc region.

According to another embodiment, the variant Fc region herein can have a sequence that has at least about 80% identity, at least about 85% identity, at least about 90% identity, at least about 95% identity or at least about 99% identity with an Fc region of a parent polypeptide.

[0123] The terms "Fc receptor" or "FcR," as used herein, generally refers to a receptor, or any derivative, variant or fragment thereof, that can bind to the Fc region of an antibody. In certain embodiments, the FcR is one which binds an IgG antibody (a gamma receptor, Fcgamma R) and includes receptors of the Fcgamma RI (CD64), Fcgamma RII (CD32), and Fcgamma RIII (CD16) subclasses, including allelic variants and alternatively spliced forms of these receptors. Fcgamma RII receptors include Fcgamma RIIA (an "activating receptor") and Fcgamma RIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. The term "FcR" also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus.

[0124] By "effector function" as used herein is meant a biochemical event that results from the interaction of an antibody Fc region with an Fc receptor or ligand, which vary with the antibody isotype. Effector functions include but are not limited to antibody-dependent cell-mediated cytotoxicity (ADCC), antibody-dependent cell-mediated phagocytosis (ADCP), complement-dependent cytotoxicity (CDC), cytokine secretion, immune complex-mediated antigen uptake by antigen presenting cells, down regulation of cell surface receptors (*e.g.* B cell receptor), and B cell activation. "Antibody-dependent cell-mediated cytotoxicity" or "ADCC" refers to a cell-mediated reaction in which nonspecific cytotoxic cells that express FcRs (such as Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. ADCC is correlated with binding to FcγRIIIa; increased binding to FcγRIIIa leads to an increase in ADCC activity. To assess ADCC activity of a molecule of interest, an in vitro ADCC assay such as that described in U.S. Pat. No. 5,500,362 or 5,821,337 may be performed. "ADCP" or antibody dependent cell-mediated phagocytosis as used herein is meant the cell-mediated reaction wherein nonspecific cytotoxic cells that express FcγRs recognize bound antibody on a target cell and subsequently cause phagocytosis of the target cell.

[0125] "Fc null" and "Fc null variant" are used interchangeably and used herein to describe a modified Fc which have reduced or abolished effector functions. Such Fc null or Fc null variant have reduced or abolished to FcγRs and/or complement receptors. Preferably, such Fc null or Fc null variant has abolished effector functions. Exemplary methods for the modification include but not limited to chemical alteration, amino acid residue substitution, insertion and deletions. Exemplary amino acid positions on Fc molecules where one or more modifications were introduced to decrease effector function of the resulting variant (numbering based on the EU numbering

scheme) at position i) IgG1: C220, C226, C229, E233, L234, L235, G237, P238, S239 D265, S267, N297, L328, P331, K322, A327 and P329, ii) IgG2: V234, G237, D265, H268, N297, V309, A330, A331, K322 and iii) IgG4: L235, G237, D265 and E318. Exemplary Fc molecules having decreased effector function include those having one or more of the following substitutions: i) IgG1: N297A, N297Q, D265A/N297A, D265A/N297Q, C220S/C226S/C229S/P238S, S267E/L328F, C226S/C229S/E233P/L234V/L235A, L234F/L235E/P331S, L234A/L235A, L234A/L235A/G237A, L234A/L235A/G237A/K322A, L234A/L235A/G237A/A330S/A331S, L234A/L235A/P329G,E233P/L234V/L235A/G236del/S239K, E233P/L234V/L235A/G236del/S267K, E233P/L234V/L235A/G236del/S239K/A327G, E233P/L234V/L235A/G236del/S267K/A327G and E233P/L234V/L235A/G236del, L234A/L235A/G237deleted; ii) IgG2: A330S/A331S, V234A/G237A, V234A/G237A/D265A, D265A/A330S/A331S, V234A/G237A/D265A/A330S/A331S, and H268Q/V309L/A330S/A331S; iii) IgG4: L235A/G237A/E318A, D265A, L235A/G237A/D265A and L235A/G237A/D265A/E318A.

[0126] "Epitope" as used herein refers to a determinant capable of specific binding to the variable region of an antibody molecule known as a paratope. Epitopes are groupings of molecules such as amino acids or sugar side chains and usually have specific structural characteristics, as well as specific charge characteristics. A single antigen may have more than one epitope. The epitope may comprise amino acid residues directly involved in the binding and other amino acid residues, which are not directly involved in the binding, such as amino acid residues which are effectively blocked by the antigen binding peptide (in other words, the amino acid residue is within the footprint of the antigen binding peptide). Epitopes may be either conformational or linear. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids. Antibodies that recognize the same epitope can be verified in a simple immunoassay showing the ability of one antibody to block the binding of another antibody to a target antigen, for example "binning".

[0127] "Linker" as used herein refers to a molecule that connect two polypeptide chains. Linker can be a polypeptide linker or a synthetic chemical linker (for example, see disclosed in Protein Engineering, 9(3), 299-305, 1996). The length and sequence of the polypeptide linkers is not particularly limited and can be selected according to the purpose by those skilled in the art. Polypeptide linker comprises one or more amino acids. Preferably, polypeptide linker is a peptide with a length of at least 5 amino acids, preferably with a length of 5 to 100, more preferably of 10 to 50 amino acids. In one embodiment, said peptide linker is G, S, GS, SG, SGG, GGS, and GSG (with G=glycine and S=serine). In another embodiment, said peptide linker is $(GGGS)_xG_n$ (SEQ ID NO: 294) or $(GGGGS)_xG_n$ (SEQ ID NO: 295), $(GGGGGS)_xG_n$ (SEQ ID NO: 296), S(GGGS)xGn

(SEQ ID NO:289), S(GGGGS)xGn (SEQ ID NO:290), or S(GGGGGS)xGn (SEQ ID NO:291), with x=1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 and n=0, 1, 2 or 3. In some embodiments, the linker is $(GGGGS)_xG_n$ with x=2,3, or 4 and n=0 (SEQ ID NO: 372); in some embodiments the linker is $(GGGGS)_xG_n$ with x=3 and n=0 (SEQ ID NO: 373). In some embodiments, the linker comprises the sequence SGGGGSGGGGSGGGGS (SEQ ID NO:292), or SGGGGSGGGGSGGGG (SEQ ID NO:293). Synthetic chemical linkers include crosslinking agents that are routinely used to crosslink peptides, for example, N-hydroxy succinimide (NHS), disuccinimidyl suberate (DSS), bis(succinimidyl) suberate (BS3), dithiobis(succinimidyl propionate) (DSP), dithiobis(succinimidyl propionate) (DTSSP), ethylene glycol bis(succinimidyl succinate) (EGS), ethylene glycol bis(sulfosuccinimidyl succinate) (sulfo-EGS), disuccinimidyl tartrate (DST), disulfosuccinimidyl tartrate (sulfo-DST), bis[2-(succinimidoxycarbonyloxy)ethyl] sulfone (BSOCOES), and bis[2-(succinimidoxycarbonyloxy)ethyl] sulfone (sulfo-BSOCOES).

[0128] The term "nucleotide," as used herein, generally refers to a base-sugar-phosphate combination. A nucleotide can comprise a synthetic nucleotide. A nucleotide can comprise a synthetic nucleotide analog. Nucleotides can be monomeric units of a nucleic acid sequence (*e.g.*, deoxyribonucleic acid (DNA) and ribonucleic acid (RNA)). The term nucleotide can include ribonucleoside triphosphates adenosine triphosphate (ATP), uridine triphosphate (UTP), cytosine triphosphate (CTP), guanosine triphosphate (GTP) and deoxyribonucleoside triphosphates such as dATP, dCTP, dITP, dUTP, dGTP, dTTP, or derivatives thereof. Such derivatives can include, for example, [αS]dATP, 7-deaza-dGTP and 7-deaza-dATP, and nucleotide derivatives that confer nuclease resistance on the nucleic acid molecule containing them. The term nucleotide as used herein, in some examples, refers to dideoxyribonucleoside triphosphates (ddNTPs) and their derivatives. Illustrative examples of dideoxyribonucleoside triphosphates can include, but are not limited to, ddATP, ddCTP, ddGTP, ddITP, and ddTTP. A nucleotide can be unlabeled or detectably labeled by well-known techniques. Labeling can also be carried out with quantum dots. Detectable labels can include, for example, radioactive isotopes, fluorescent labels, chemiluminescent labels, bioluminescent labels and enzyme labels. Fluorescent labels of nucleotides can include but are not limited fluorescein, 5-carboxyfluorescein (FAM), 2′7′-dimethoxy-4′5-dichloro-6-carboxyfluorescein (JOE), rhodamine, 6-carboxyrhodamine (R6G), N,N,N′,N′-tetramethyl-6-carboxyrhodamine (TAMRA), 6-carboxy-X-rhodamine (ROX), 4-(4′dimethylaminophenylazo) benzoic acid (DABCYL), Cascade Blue, Oregon Green, Texas Red, Cyanine and 5-(2′-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS). Specific examples of fluorescently labeled nucleotides can include [R6G]dUTP, [TAMRA]dUTP, [R110]dCTP, [R6G]dCTP, [TAMRA]dCTP, [JOE]ddATP, [R6G]ddATP, [FAM]ddCTP, [R110]ddCTP,

[TAMRA]ddGTP, [ROX]ddTTP, [dR6G]ddATP, [dR110]ddCTP, [dTAMRA]ddGTP, and [dROX]ddTTP available from Perkin Elmer, Foster City, Calif; FluoroLink DeoxyNucleotides, FluoroLink Cy3-dCTP, FluoroLink Cy5-dCTP, FluoroLink Fluor X-dCTP, FluoroLink Cy3-dUTP, and FluoroLink Cy5-dUTP available from Amersham, Arlington Heights, Ill.; Fluorescein-15-dATP, Fluorescein-12-dUTP, Tetramethyl-rodamine-6-dUTP, IR770-9-dATP, Fluorescein-12-ddUTP, Fluorescein-12-UTP, and Fluorescein-15-2′-dATP available from Boehringer Mannheim, Indianapolis, Ind.; and Chromosome Labeled Nucleotides, BODIPY-FL-14-UTP, BODIPY-FL-4-UTP, BODIPY-TMR-14-UTP, BODIPY-TMR-14-dUTP, BODIPY-TR-14-UTP, BODIPY-TR-14-dUTP, Cascade Blue-7-UTP, Cascade Blue-7-dUTP, fluorescein-12-UTP, fluorescein-12-dUTP, Oregon Green 488-5-dUTP, Rhodamine Green-5-UTP, Rhodamine Green-5-dUTP, tetramethylrhodamine-6-UTP, tetramethylrhodamine-6-dUTP, Texas Red-5-UTP, Texas Red-5-dUTP, and Texas Red-12-dUTP available from Molecular Probes, Eugene, Oreg. Nucleotides can also be labeled or marked by chemical modification. A chemically-modified single nucleotide can be biotin-dNTP. Some non-limiting examples of biotinylated dNTPs can include, biotin-dATP (*e.g.*, bio-N6-ddATP, biotin-14-dATP), biotin-dCTP (*e.g.*, biotin-11-dCTP, biotin-14-dCTP), and biotin-dUTP (*e.g.*, biotin-11-dUTP, biotin-16-dUTP, biotin-20-dUTP).

[0129] The terms “polynucleotide,” “oligonucleotide,” and “nucleic acid” are used interchangeably to refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof, either in single-, double-, or multi-stranded form. A polynucleotide can be exogenous or endogenous to a cell. A polynucleotide can exist in a cell-free environment. A polynucleotide can be a gene or fragment thereof. A polynucleotide can be DNA. A polynucleotide can be RNA. A polynucleotide can have any three-dimensional structure, and can perform any function, known or unknown. A polynucleotide can comprise one or more analogs (*e.g.*, altered backbone, sugar, or nucleobase). If present, modifications to the nucleotide structure can be imparted before or after assembly of the polymer. Some non-limiting examples of analogs include: 5-bromouracil, peptide nucleic acid, xeno nucleic acid, morpholinos, locked nucleic acids, glycol nucleic acids, threose nucleic acids, dideoxynucleotides, cordycepin, 7-deaza-GTP, fluorophores (*e.g.*, rhodamine or fluorescein linked to the sugar), thiol containing nucleotides, biotin linked nucleotides, fluorescent base analogs, CpG islands, methyl-7-guanosine, methylated nucleotides, inosine, thiouridine, pseudourdine, dihydrouridine, queuosine, and wyosine. Non-limiting examples of polynucleotides include coding or non-coding regions of a gene or gene fragment, loci (locus) defined from linkage analysis, exons, introns, messenger RNA (mRNA), transfer RNA (tRNA), ribosomal RNA (rRNA), short interfering RNA (siRNA), short-hairpin RNA (shRNA), micro-RNA (miRNA), ribozymes, cDNA,

recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, cell-free polynucleotides including cell-free DNA (cfDNA) and cell-free RNA (cfRNA), nucleic acid probes, and primers. The sequence of nucleotides can be interrupted by non-nucleotide components.

[0130] The term "gene," as used herein, refers to a nucleic acid (*e.g.*, DNA such as genomic DNA and cDNA) and its corresponding nucleotide sequence that is involved in encoding an RNA transcript. The term as used herein with reference to genomic DNA includes intervening, non-coding regions as well as regulatory regions and can include 5' and 3' ends. In some uses, the term encompasses the transcribed sequences, including 5' and 3' untranslated regions (5'-UTR and 3'-UTR), exons and introns. In some genes, the transcribed region will contain "open reading frames" that encode polypeptides. In some uses of the term, a "gene" comprises only the coding sequences (*e.g.*, an "open reading frame" or "coding region") necessary for encoding a polypeptide. In some cases, genes do not encode a polypeptide, for example, ribosomal RNA genes (rRNA) and transfer RNA (tRNA) genes. In some cases, the term "gene" includes not only the transcribed sequences, but in addition, also includes non-transcribed regions including upstream and downstream regulatory regions, enhancers and promoters. A gene, in some cases, refers to an "endogenous gene" or a native gene in its natural location in the genome of an organism. A gene, in some cases, refers to an "exogenous gene" or a non-native gene. A non-native gene, in some cases, refers to a gene not normally found in the host organism but which is introduced into the host organism by gene transfer. A non-native gene , in some cases, refers to a gene not in its natural location in the genome of an organism. A non-native gene, in some cases, also refers to a naturally occurring nucleic acid or polypeptide sequence that comprises mutations, insertions and/or deletions (*e.g.*, non-native sequence).

[0131] The term "regulating" with reference to expression or activity, as used herein, refers to altering the level of expression or activity. Regulation can occur at the transcription level and/or translation level.

[0132] The terms "peptide," "polypeptide," and "protein" are used interchangeably herein to refer to a polymer of at least two amino acid residues joined by peptide bond(s). This term does not connote a specific length of polymer, nor is it intended to imply or distinguish whether the peptide is produced using recombinant techniques, chemical or enzymatic synthesis, or is naturally occurring. The terms apply to naturally occurring amino acid polymers as well as amino acid polymers comprising at least one modified amino acid. In some cases, the polymer can be interrupted by non-amino acids. The terms include amino acid chains of any length, including full length proteins, and proteins with or without secondary and/or tertiary structure (*e.g.*, domains).

The terms also encompass an amino acid polymer that has been modified, for example, by disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, oxidation, and any other manipulation such as conjugation with a labeling component. The terms "amino acid" and "amino acids," as used herein, generally refer to natural and non-natural amino acids, including, but not limited to, modified amino acids and amino acid analogues. Modified amino acids can include natural amino acids and non-natural amino acids, which have been chemically modified to include a group or a chemical moiety not naturally present on the amino acid. Amino acid analogues can refer to amino acid derivatives. The term "amino acid" includes both D-amino acids and L-amino acids.

[0133] The terms "derivative," "variant," and "fragment," when used herein with reference to a polypeptide, refers to a polypeptide related to a wild type polypeptide, for example either by amino acid sequence, structure (*e.g.*, secondary and/or tertiary), activity (*e.g.*, enzymatic activity) and/or function. Derivatives, variants and fragments of a polypeptide can comprise one or more amino acid variations (*e.g.*, mutations, insertions, and deletions), truncations, modifications, or combinations thereof compared to a wild type polypeptide.

[0134] The term "residue" as used herein means a position in a protein and its associated amino acid identity. For example, Leu 234 (also referred to as Leu234 or L234) is a residue at position 234 in the human antibody IgG1.

[0135] The term "wild-type" as used herein means an amino acid sequence or a nucleotide sequence that is found in nature, including allelic variations. A wild-type protein has an amino acid sequence or a nucleotide sequence that has not been intentionally modified.

[0136] The terms "substitution" or "mutation" refers to a change to the polypeptide backbone wherein an amino acid occurring naturally in the wild-type sequence of a polypeptide is substituted to another amino acid not naturally occurring at the same position in the said polypeptide. In some cases, a mutation or mutations is introduced to modify a polypeptide's affinity to its receptor thereby altering its activity such that it becomes different from the affinity and activity of the wild-type cognate polypeptide. Mutations can also improve a polypeptide's biophysical properties. Amino acid mutations can be generated using genetic or chemical methods well known in the art. Genetic methods may include site-directed mutagenesis, PCR, gene synthesis and the like. It is contemplated that methods of altering the side chain group of an amino acid by methods other than genetic engineering, such as chemical modification, may also be useful.

[0137] The terms "affinity" or "binding affinity" refer to the strength of the sum total of non-covalent interactions between a single binding site of a molecule (*e.g.*, an antibody) and its binding partner (*e.g.*, an antigen). Unless indicated otherwise, as used herein, "binding affinity"

refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (*e.g.*, antibody and antigen). The affinity can generally be represented by the dissociation constant (K_D), which is the ratio of dissociation and association rate constants (koff and kon, respectively). Thus, equivalent affinities may comprise different rate constants, as long as the ratio of the rate constants remains the same. Affinity can be measured by common methods known in the art, such as enzyme-linked immunosorbent assay (ELISA), surface plasmon resonance (SPR) technologies (*e.g.*, BIAcore), BioLayer Interferometry (BLI) technologies (*e.g.* Octet) and other traditional binding assays (Heeley, Endocr Res 28, 217-229 (2002). In some examples, binding affinity is determined by Scatchard analysis, which comprises generating a Scatchard plot, which is a plot of the ratio of concentrations of bound ligand to unbound ligand versus the bound ligand concentration.

[0138] The terms "binding" or "specific binding" as used here, can refer to the ability of a polypeptide or an antigen binding domain to selectively interact with the receptor for the polypeptide or target antigen, respectively, and this specific interaction can be distinguished from non-targeted or undesired or non-specific interactions. Examples of specific binding can include but are not limited to IL-21 cytokine binding to its specific receptors (*e.g.*, IL-21R and common gamma chain) and an antigen binding domain binding to a specific antigen (*e.g.*, CD8 or PD-1).

[0139] The terms "subject," "individual," and "patient" are used interchangeably herein to refer to a vertebrate, preferably a mammal such as a human. Mammals include, but are not limited to, murines, simians, humans, farm animals, sport animals, and pets. Tissues, cells and their progeny of a biological entity obtained *in vivo* or cultured *in vitro* are also encompassed.

[0140] The terms "treatment" and "treating," as used herein, refer to an approach for obtaining beneficial or desired results including but not limited to a therapeutic benefit and/or a prophylactic benefit. For example, a treatment can comprise administering a system or cell population disclosed herein. By therapeutic benefit is meant any therapeutically relevant improvement in or effect on one or more diseases, conditions, or symptoms under treatment. For prophylactic benefit, a composition can be administered to a subject at risk of developing a particular disease, condition, or symptom, or to a subject reporting one or more of the physiological symptoms of a disease, even though the disease, condition, or symptom may not have yet been manifested.

[0141] The terms "effective amount," or "therapeutically effective amount," or "effective dose," or "effective dosage," refer to the quantity of a composition, for example a composition comprising immune cells such as lymphocytes (*e.g.*, T lymphocytes and/or NK cells), which can be combined with a targeted cytokine construct of the present disclosure, that is sufficient to result in a

desired activity upon administration to a subject in need thereof. Within the context of the present disclosure, the term "therapeutically effective" refers to that quantity of a composition that is sufficient to delay the manifestation, arrest the progression, relieve or alleviate at least one symptom of a disorder treated by the methods of the present disclosure.

[0142] As used herein, the term "Percent (%) amino acid sequence identity" with respect to a sequence is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as EMBOSS MATCHER, EMBOSS WATER, EMBOSS STRETCHER, EMBOSS NEEDLE, EMBOSS LALIGN, BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. Alignment for purposes of determining percent amino acid sequence identity can for example be achieved using publicly available sequence comparison computer program ALIGN-2. The source code for the ALIGN-2 sequence comparison computer program is available with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program can be compiled for use on a UNIX operating system, such as a digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

[0143] For all amino acid positions discussed in the present disclosure, in the context of antibodies or antigen binding fragments thereof, numbering is according to the EU index. The "EU index" or "EU index as in Kabat et al." or "EU numbering scheme" refers to the numbering of the EU antibody (See Edelman et al., 1969; Kabat et al., 1991).

Interleukin-21 Polypeptides

[0144] Interleukin-21 (IL-21) is a cytokine that can be expressed by T cells, B cells, NK cells and myeloid cells, and regulates the activity of both innate and adaptive immune cells and improves T cell survival and effector function. Several Phase I and II clinical trials include IL-21 as the investigational product for the treatment of cancers, inflammatory diseases, and autoimmune diseases, including, melanoma, renal cell carcinoma, acute myeloid leukemia, non-Hodgkin's

lymphoma, ovarian cancer, colorectal cancer, systemic lupus erythematosus, Crohn's disease and rheumatoid arthritis.

[0145] IL-21 has a four-helix bundle structure and exists as a monomer. In humans, two isoforms of IL-21 are known, each of which are derived from a precursor molecule. The first IL-21 isoform comprises 162 amino acids (aa), the first 29 of which make up the signal peptide; and the second IL-21 isoform comprises 153 aa, the first 29 of which make up the signal peptide as in the first isoform.

[0146] IL-21 binds to the heterodimeric IL-21 receptor complex, comprising of an IL-21 receptor (IL-21R) and common gamma chain (γc). IL-21 receptor complex is expressed on the surface of T, B, and NK cells.IL-21 receptor complex is similar in structure to the IL-2 receptor complex, in that each of these cytokine receptor complex comprises a γc.

[0147] When IL-21 binds to IL-21 receptor complex, the JAK/STAT signaling pathway is activated to activate target genes. While IL-21-induced signaling may be therapeutically desirable, careful consideration of the timing and the location of the signaling is needed, given IL-21's broad expression profile and due to the fact that IL-21 has the ability to potentiate CD8+ T cell responses as well as to suppress antigen presentation and T cell priming.

[0148] The present disclosure provides, in some embodiments, IL-21 polypeptides or functional fragments or variants thereof, comprising at least one amino acid substitution, relative to the wild-type IL-21 amino acid sequence, which is provided herein as SEQ ID NO: 1. Unless noted otherwise, the terms "wild-type IL-21," "wild-type IL-21 polypeptide," "human IL-21," and "human IL-21 polypeptide" are used interchangeably and can refer to the amino acid sequence of SEQ ID NO: 1. Such IL-21 polypeptides comprising at least one amino acid substitution relative to SEQ ID NO: 1 are also referred to herein as IL-21 muteins. In exemplary aspects, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises at least one and not more than X amino acid substitutions, wherein X is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 or greater. In some embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises at least 35 amino acid substitutions compared to SEQ ID NO: 1. In exemplary embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by 10 amino acids, 15 amino acids, 20 amino acids, or 25 amino acids. In some embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by no more than 16 amino acids. In some embodiments, an IL-21 polypeptide or

functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 amino acids. In exemplary embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by no more than 7 amino acids or no more than 5 amino acids. In some embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by 10, 11, 12, 13, 14, 15 or 16 amino acids. In exemplary embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by 3, 4, 5, or 6 amino acids. In some embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by 5 to 16. In exemplary embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by 3 to 6 amino acids or 1 to 5 amino acids. In exemplary embodiments, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, comprises an amino acid sequence which differs from the amino acid sequence of human IL-21 (SEQ ID NO: 1) by one or two amino acids.

Altered Isoelectric Point or Charge Distribution

[0149] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least one mutation that alters the isoelectric point with or without affecting its binding to IL-21R. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant therefor comprises at least one mutation that alters the isoelectric point without affecting binding to IL-21R. An altered isoelectric point of a protein can change the surface charge distribution, thereby affecting protein yield, non-specific binding, and the blood or circulatory half-life for a given protein. In some embodiments, the mutations in the IL-21 polypeptide or a functional fragment or a variant thereof decreases the theoretical isoelectric point by a given unit measured by a publicly available database ProtParam (SwissProt), which is hereby incorporated by reference.

[0150] In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof has an isoelectric point difference of about 0.6 to about 5.0 units compared to the isoelectric point of SEQ ID NO: 1, which has an isoelectric point of about 9.42. In some embodiments, the IL-

21 polypeptide or a functional fragment or a variant thereof has an isoelectric point of about 6.0 to about 9.0. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a theoretical isoelectric point of about 6 to about 9. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a theoretical isoelectric point of at least about 6. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a theoretical isoelectric point of at most about 9. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a theoretical isoelectric point of about 9 to about 8.9, about 9 to about 8.8, about 9 to about 8.7, about 9 to about 8.5, about 9 to about 8.3, about 9 to about 8.1, about 9 to about 7.9, about 9 to about 7.7, about 9 to about 7.5, about 9 to about 7.3, about 9 to about 7, about 9 to about 6.5, about 9 to about 6.0, about 8.9 to about 8.8, about 8.9 to about 8.7, about 8.9 to about 8.5, about 8.9 to about 8.3, about 8.9 to about 8.1, about 8.9 to about 7.9, about 8.9 to about 7.7, about 8.9 to about 7.5, about 8.9 to about 7.3, about 8.9 to about 7, about 8.9 to about 6.5, about 8.9 to about 6.0, about 8.8 to about 8.7, about 8.8 to about 8.5, about 8.8 to about 8.3, about 8.8 to about 8.1, about 8.8 to about 7.9, about 8.8 to about 7.7, about 8.8 to about 7.5, about 8.8 to about 7.3, about 8.8 to about 7, about 8.8 to about 6.5, about 8.8 to about 6.0, about 8.7 to about 8.6, about 8.7 to about 8.5, about 8.7 to about 8.3, about 8.7 to about 8.1, about 8.7 to about 7.9, about 8.7 to about 7.7, about 8.7 to about 7.5, about 8.7 to about 7.3, about 8.7 to about 7, about 8.7 to about 6.5, about 8.7 to about 6.0, about 8.5 to about 8.3, about 8.5 to about 8.4, about 8.5 to about 8.39, about 8.5 to about 8.1, about 8.5 to about 7.9, about 8.5 to about 7.7, about 8.5 to about 7.5, about 8.5 to about 7.3, about 8.5 to about 7, about 8.5 to about 6.5, about 8.5 to about 6.0, about 8.39, about 8.3 to about 8.1, about 8.3 to about 7.9, about 8.3 to about 7.7, about 8.3 to about 7.5, about 8.3 to about 7.3, about 8.3 to about 7, about 8.3 to about 6.5, about 8.3 to about 6.0, about 8.1 to about 7.9, about 8.1 to about 7.7, about 8.1 to about 7.5, about 8.1 to about 7.3, about 8.1 to about 7, about 8.1 to about 6.5, about 8.1 to about 6.0, about 7.9 to about 7.7, about 7.9 to about 7.5, about 7.9 to about 7.3, about 7.9 to about 7, about 7.9 to about 6.5, about 7.9 to about 6.0, about 7.7 to about 7.5, about 7.7 to about 7.3, about 7.7 to about 7, about 7.5 to about 7.3, about 7.5 to about 7, or about 7.3 to about 7, about 7.3 to about 6.5, about 7.3 to about 6.0, about 7.0 to about 6.5, about 7.0 to about 6.0, or about 6.5 to about 6.0. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a theoretical isoelectric point of about 9, about 8.9, about 8.8, about 8.79, about 8.7, about 8.5, about 8.39, about 8.3, about 8.1, about 7.9, about 7.7, about 7.5, about 7.3, about 7, about 6.5 or about 6.0. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof has a theoretical isoelectric point of less than 9.0. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof has a theoretical isoelectric point of less

than 8.9, less than 8.8, less than 8.7, less than 8.5, less than 8.3, less than 8.1, less than 7.9, less than 7.7, less than 7.5, less than 7.3, less than 7.2, less than 7.0, less than 6.5 or less than 6.2.

[0151] In some instances, an altered isoelectric point can increase the protein yield of IL-21 polypeptide or a functional fragment or a variant thereof during the purification compared to wild-type IL-21. In some embodiments, the increased protein yield is at least 5% greater than wild-type IL-21. In some embodiments, the increased protein yield is at least 5%, 10%, 15%, 20%, 25%, 50%, 75%, 100%, 125%, 150%, 175%, 200%, 225%, 250%, 275%, or 300% greater than wild-type IL-21. In some embodiments, the increased protein yield is at least 5%, 10%, 15%, 20%, 25%, 50%, 75%, 100%, 125%, 150%, 175%, 200%, 225%, 250%, 275%, or 300% greater than wild-type IL-21. In some embodiments, the increased protein yield is 5% to 10 %, 5 % to 15 %, 5 % to 20 %, 5 % to 25 %, 5 % to 50 %, 5 % to 75 %, 5 % to 100 %, 5 % to 125 %, 5 % to 150 %, 5 % to 175 %, 5 % to 200 %, 5 % to 225%, 5 % to 250 %, 5 % to 275 %, 5 % to 300 %, 10 % to 15 %, 10 % to 20 %, 10 % to 25 %, 10 % to 50 %, 10 % to 75 %, 10 % to 100 %, 10 % to 125 %, 10 % to 150 %, 10 % to 175 %, 10 % to 200 %, 10 % to 225%, 10 % to 250 %, 10 % to 275 %, 10 % to 300 %, 15 % to 20 %, 15 % to 25 %, 15 % to 50 %, 15 % to 75 %, 15 % to 100 %, 15 % to 125 %, 15 % to 150 %, 15 % to 175 %, 15 % to 200 %, 15 % to 225%, 15 % to 250 %, 15 % to 275 %, 15 % to 300 %, 20 % to 25 %, 20 % to 50 %, 20 % to 75 %, 20 % to 100 %, 20 % to 125 %, 20 % to 150 %, 20 % to 175 %, 20 % to 200 %, 20 % to 225%, 20 % to 250 %, 20 % to 275 %, 20 % to 300 %, 25 % to 50 %, 25 % to 75 %, 25 % to 100 %, 25 % to 125 %, 25 % to 150 %, 25 % to 175 %, 25 % to 200 %, 25 % to 225%, 25 % to 250 %, 25 % to 275 %, 25 % to 300 %, 50 % to 75 %, 50 % to 100 %, 50 % to 125 %, 50 % to 150 %, 50 % to 175 %, 50 % to 200 %, 75 % to 100 %, 75 % to 125 %, 75 % to 150 %, 75 % to 175 %, 75 % to 200 %, 75 % to 225%, 75 % to 250 %, 75 % to 275 %, 75 % to 300 %, 100 % to 125 %, 100 % to 150 %, 100 % to 175 %, 100 % to 200 %, 100 % to 225%, 100 % to 250 %, 100 % to 275 %, 100 % to 300 %, 125 % to 150 %, 125 % to 175 %, 125 % to 200 %, 125 % to 225%, 125 % to 250 %, 125 % to 275 %, 125 % to 300 %, 150 % to 175 %, 150 % to 200 %, 175 % to 200 %, 175 % to 225%, 175 % to 250 %, 175 % to 275 %, 175 % to 300 %, 200 % to 225%, 200 % to 250 %, 200 % to 275 %, 200 % to 300 %, 225 % to 250 %, 225 % to 275 %, 225 % to 300 %, 250 % to 275 %, 250 % to 300 %, or 275 % to 300 %. In some embodiments, the increased protein yield is about 5 %, 10 %, 15 %, 20 %, 25 %, 50 %, 75 %, 100 %, 125 %, 150 %, 175 %, 200 %, 225%, 250%, 275%, or 300%.

[0152] In some embodiments, relative to a region of wild-type IL-21 polypeptide that comprises one or more positively charged amino acid residues, the IL-21 polypeptide or a functional fragment or a variant thereof comprises at least one amino acid substitution of the one or more positively charged amino acid residues. In some cases, the region of wild-type IL-21

comprises 2-20 positively charged amino acid residues. In some embodiments, the region in wild-type IL-21 comprises 2-3, 2-4, 2-5, 2-7. 2-8, 2-9, 2-10, 2-11, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 2-20, 3-4, 3-5, 3-6, 3-7, 3-8, 3-9, 3-10, 3-11, 3-12, 3-13, 3-14, 3-15, 3-16, 3-17, 3-18, 3-19, 3-20, 4-5, 4-6, 4-7, 4-8, 4-9, 4-10, 4-11, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 4-20, 5-6, 5-7, 5-8, 5-9, 5-10, 5-11, 5-12, 5-13, 5-14, 5-15, 5-16, 5-17, 5-18, 5-19, 5-20, 6-7, 6-8, 6-9, 6-10, 6-11, 6-12, 6-13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 6-20, 7-8, 7-9, 7-10, 7-11, 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 7-20, 8-9, 8-10, 8-11, 8-12, 8-13, 8-14, 8-15, 8-16, 8-17, 8-18, 8-19, 8-20, 9-10, 9-11, 9-12, 9-13, 9-14, 9-15, 9-16, 9-17, 9-18, 9-19, 9-20, 10-11, 10-12, 10-13, 10-14, 10-15, 10-16, 10-17, 10-18, 10-19, 10-20, 11-12,11-13, 11-14, 11-15, 11-16, 11-17, 11-18, 11-19, 11-20, 12-13, 12-14, 12-15, 12-16, 12-17, 12-18, 12-19, 12-20, 13-14, 13-15, 13-16, 13-17, 13-18, 13-19, 13-20, 14-15, 14-16, 14-17, 3-18, 14-19, 14-20, 15-16, 15-17, 15-18, 15-19, 15-20, 16-17, 16-18, 16-19, 16-20, 17-18, 17-19, 17-20 18-19, 18-20, or 19-20 positively charged amino acid residues. In some embodiments, the region of wild-type IL-21 comprises 12 positively charged amino acid residues. In some embodiments, the region of wild-type IL-21 comprises 2-20 amino acid residues, of which at least one amino acid is positively charged. In some embodiments, the region of wild-type IL-21 comprises 2-3, 2-4, 2-5, 2-7. 2-8, 2-9, 2-10, 2-11, 2-12, 2-13, 2-14, 2-15, 2-16, 2-17, 2-18, 2-19, 2-20, 3-4, 3-5, 3-6, 3-7, 3-8, 3-9, 3-10, 3-11, 3-12, 3-13, 3-14, 3-15, 3-16, 3-17, 3-18, 3-19, 3-20, 4-5, 4-6, 4-7, 4-8, 4-9, 4-10, 4-11, 4-12, 4-13, 4-14, 4-15, 4-16, 4-17, 4-18, 4-19, 4-20, 5-6, 5-7, 5-8, 5-9, 5-10, 5-11, 5-12, 5-13, 5-14, 5-15, 5-16, 5-17, 5-18, 5-19, 5-20, 6-7, 6-8, 6-9, 6-10, 6-11, 6-12, 6-13, 6-14, 6-15, 6-16, 6-17, 6-18, 6-19, 6-20, 7-8, 7-9, 7-10, 7-11, 7-12, 7-13, 7-14, 7-15, 7-16, 7-17, 7-18, 7-19, 7-20, 8-9, 8-10, 8-11, 8-12, 8-13, 8-14, 8-15, 8-16, 8-17, 8-18, 8-19, 8-20, 9-10, 9-11, 9-12, 9-13, 9-14, 9-15, 9-16, 9-17, 9-18, 9-19, 9-20, 10-11, 10-12, 10-13, 10-14, 10-15, 10-16, 10-17, 10-18, 10-19, 10-20, 11-12,11-13, 11-14, 11-15, 11-16, 11-17, 11-18, 11-19, 11-20, 12-13, 12-14, 12-15, 12-16, 12-17, 12-18, 12-19, 12-20, 13-14, 13-15, 13-16, 13-17, 13-18, 13-19, 13-20, 14-15, 14-16, 14-17, 3-18, 14-19, 14-20, 15-16, 15-17, 15-18, 15-19, 15-20, 16-17, 16-18, 16-19, 16-20, 17-18, 17-19, 17-20 18-19, 18-20, or 19-20 amino acid residues with at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 positively charged amino acid residues. In some embodiments, the region of wild-type IL-21 comprises 12 amino acid residues, of which 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 amino acid residues are positively charged. In some embodiments, the region of wild-type IL-21 comprises 12 amino acid residues, of which 2, 3, 4, 5, or 6 amino acid residues are positively charged.

[0153] In some embodiments, amino acids within the region of wild-type IL-21 do not bind a human IL-21 receptor. In some embodiments, the region of wild-type IL-21 comprises amino acid residues S80-T92 of SEQ ID NO: 1. In some embodiments, the region of wild-type IL-21

comprises positively charged amino acid residues R85, R86, K88, H89, and R90. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises at least one amino acid substitution in the region of wild-type IL-21. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises at least one amino acid substitution of S80-T92. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises at least four amino acid substitutions of S80-T92. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises at least four amino acid substitutions of S80-T92, provided G84 is not substituted. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises at least one amino acid substitution at R85, R86, K88, H89 or R90. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid substitution at one or more of S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91 and T92. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises amino acid substitutions at S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91 and T92.

[0154] In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof that comprises an altered isoelectric point relative to wild-type IL-21 (SEQ ID NO: 1) comprises a mutation in a position selected from the group consisting of K56, S80, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof that comprises an altered isoelectric point relative to wild-type IL-21 (SEQ ID NO: 1) comprises a mutation in a position selected from the group consisting of K56, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof that comprises an altered isoelectric point relative to wild-type IL-21 (SEQ ID NO: 1) comprises a mutation in a position selected from the group consisting of S80, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof that comprises an altered isoelectric point relative to wild-type IL-21 (SEQ ID NO: 1) comprises a mutation in a position selected from the group consisting of S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1.

[0155] In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position K56 of SEQ ID NO: 1. In some instances, the mutation comprises K56G, K56S, K56E, K56D, or K56A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position S80. In some embodiments, the mutation comprises S80G, S80A, S80D or S80E. In some embodiments, the IL-

21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position T81 of SEQ ID NO: 1. In some instances, the mutation comprises T81G, T81S, T81E, T81D, or T81A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position N82 of SEQ ID NO: 1. In some instances, the mutation comprises N82G, N82S, N82E, N82D, or N82A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position A83 of SEQ ID NO: 1. In some instances, the mutation comprises A83G, A83S, A83E, or A83D. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position G84 of SEQ ID NO: 1. In some instances, the mutation comprises G84A, G84S, G84E, or G84D. In some embodiments, G84 is not mutated. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position R85 of SEQ ID NO: 1. In some instances, the mutation comprises R85G, R85S, R85E, R85D, or R85A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position R86 of SEQ ID NO: 1. In some instances, the mutation comprises R86G, R86S, R86E, R86D, or R86A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position Q87 of SEQ ID NO: 1. In some instances, the mutation comprises Q87G, Q87S, Q87E, Q87D, or Q87A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position K88 of SEQ ID NO: 1. In some instances, the mutation comprises K88G, K88S, K88E, K88D, or K88A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position H89 of SEQ ID NO: 1. In some instances, the mutation comprises H89G, H89S, H89E, H89D, or H89A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position R90 of SEQ ID NO: 1. In some instances, the mutation comprises R90G, R90S, R90E, R90D, or R90A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position L91 of SEQ ID NO: 1. In some instances, the mutation comprises L91G, L91S, L91E, L91D, or L91A. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof comprises a mutation at position T92 of SEQ ID NO: 1. In some instances, the mutation comprises T92G, T92S, T92E, T92D, or T92A.

[0156] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof that has an altered isoelectric point relative to the IL-21 of SEQ ID NO: 1 comprises an amino acid sequence that is at least 75% identical to the sequence of SEQ ID NO: 1 and comprises a mutation in at least one position selected from the group consisting of positions K56 and R90 of SEQ ID NO: 1. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant

thereof that has an altered isoelectric point relative to the IL-21 of SEQ ID NO: 1 comprises an amino acid sequence that is at least 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% identical to the sequence of SEQ ID NO: 1 and comprises a mutation in at least one position selected from the group consisting of positions K56 and R90 of SEQ ID NO: 1. The mutation at position K56, in some instances, is K56G, K56S, K56E, K56D, or K56A; the mutation at position R90, in some instances, is R90G, R90S, R90E, R90D, or R90A. The altered isoelectric point, in some examples, is a reduced isoelectric point compared to the IL-21 of SEQ ID NO: 1 (which is about 9.42), *e.g.*, reduced by at least about 0.6 units.

[0157] In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids selected from the group consisting of G85, G86, G88, and A90; G85, G86, G88, and E90; A56, A75, G85, G86, G88, and E90; A56, E75, G85, G86, G88, and A90; E56, A75, G85, G86, G88, and A90; G80, G81, G82, S83, E85, G86, S87, G88, G89, and S90; G80, G81, G82, S83, G85, G86, S87, G88, G89, S90; G85, G86, S87, G88, G89, and E90; G85, G86, S87, G88, G89, and S90; G85, G86, G87, G88, G89, and E90; G85, G86, G87, G88, G89, and G90; or G85, G86, E87, G88, G89, and G90, wherein the position numbers correspond to SEQ ID NO: 1.

[0158] In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, G88, and E90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, G88, and E90.In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids A56, A75, G85, G86, G88, and E90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids A56, A75, G85, G86, G88, and E90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: A56, E75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids A56, E75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: A56, A75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide comprises at

least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids A56, A75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: E56, A75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids E56, A75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, S83, E85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, S83, E85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, S83, G85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, S83, G85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, S83, G85, G86, S87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, S83, G85, G86, S87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G85, G86, S87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, S87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G85, G86, G87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, G87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G85, G86, G87, G88, G89, and G90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises

amino acids G85, G86, G87, G88, G89, and G90. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G85, G86, E87, G88, G89, and G90. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, E87, G88, G89, and G90. Provided in some embodiments are charge variant IL-21 polypeptides or functional fragments or variants thereof that comprise a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a sequence selected from the group consisting of SEQ ID NOs: 2, 4, 5, 6, 7, 8, 9, 10, 11, 12,13, 14 and 15.

[0159] In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, S83, G85, G86, S87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, S83, G85, G86, S87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, G83, G85, G86, G87, G88, G89, G90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, G83, G85, G86, G87, G88, G89, G90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, S83, G85, G86, E87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, S83, G85, G86, E87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, E83, G85, G86, E87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, E83, G85, G86, E87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G82, G83, S85, G86, G87, G88, S89, G90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G82, G83, S85, G86, G87, G88, S89, G90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G82, G83, E85, G86, G87, G88, S89, G90, G91 and S92. In

some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G82, G83, E85, G86, G87, G88, S89, G90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, G83, E85, G86, G87, G88, G89, G90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, G83, E85, G86, G87, G88, G89, G90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, S83, G85, G86, S87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, S83, G85, G86, S87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G82, G83, S85, G86, G87, G88, S89, E90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G82, G83, S85, G86, G87, G88, S89, E90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G82, G83, G85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G82, G83, G85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide

comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G82, G83, G85, E86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G82, G83, G85, E86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G82, G83, E85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G82, G83, E85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, G83, E84, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, G83, E84, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, G82, E83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, G82, E83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 1 and comprises amino acids: G80, G81, E82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 1 and comprises amino acids G80, G81, E82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. Provided in some embodiments are charge variant IL-21 polypeptides or functional fragments or variants thereof that comprise a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a sequence selected from the group consisting of SEQ ID NOs. 23-40. Provided in some embodiments are charge variant IL-21 polypeptides or functional fragments or variants thereof that comprise a sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12,13, 14 15, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and 40.

[0160] In some embodiments, an IL-21 polypeptide comprises about 75 % sequence identity to about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 and 15. In some embodiments, an IL-21 polypeptide comprises at least about 75 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 and 15. In some embodiments, an IL-21

polypeptide comprises at most about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 and 15. In some embodiments, an IL-21 polypeptide comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 and 15. In some embodiments, an

IL-21 polypeptide comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 and 15.

[0161] In some embodiments, an IL-21 polypeptide comprises about 75 % sequence identity to about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 23-40. In some embodiments, an IL-21 polypeptide comprises at least about 75 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 23-40. In some embodiments, an IL-21 polypeptide comprises at most about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 23-40. In some embodiments, an IL-21 polypeptide comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 %

sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 23-40. In some embodiments, an IL-21 polypeptide comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 23-40.

[0162] In some embodiments, an IL-21 polypeptide comprises about 75 % sequence identity to about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12,13, 14 15, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and 40. In some embodiments, an IL-21 polypeptide comprises at least about 75 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12,13, 14 15, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and 40. In some embodiments, an IL-21 polypeptide comprises at most about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12,13, 14 15, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and 40. In some embodiments, an IL-21 polypeptide comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 %

sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12,13, 14 15, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and 40. In some embodiments, an IL-21 polypeptide comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to a sequence selected from the group consisting of SEQ ID NOs. 2, 4, 5, 6, 7, 8, 9, 10, 11, 12,13, 14 15, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and 40.

[0163] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 4 and comprises amino acids G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 4 and comprises amino acids G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 2 and comprises amino acids G85, G86, G88, and E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at

least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 2 and comprises amino acids G85, G86, G88, and E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 5 and comprises amino acids A56, A75, G85, G86, G88, and E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 5 and comprises amino acids A56, A75, G85, G86, G88, and E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 6 and comprises amino acids: A56, E75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 6 and comprises amino acids A56, E75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 7 and comprises amino acids: A56, A75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 7 and comprises amino acids A56, A75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 8 and comprises amino acids: E56, A75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 8 and comprises amino acids E56, A75, G85, G86, G88, and A90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 9 and comprises amino acids: G80, G81, G82, S83, E85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 9 and comprises amino acids G80, G81, G82, S83, E85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 10 and comprises amino acids: G80, G81, G82, S83, G85, G86, S87, G88, G89, S90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 10 and comprises amino acids G80, G81, G82, S83, G85, G86, S87, G88, G89, S90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 10 and comprises amino acids: G80, G81, G82, S83, G85, G86, S87, G88, G89, E90. In some

embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 10 and comprises amino acids G80, G81, G82, S83, G85, G86, S87, G88, G89, E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 11 and comprises amino acids: G85, G86, S87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 11 and comprises amino acids G85, G86, S87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 12 and comprises amino acids: G85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 12 and comprises amino acids G85, G86, S87, G88, G89, and S90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 13 and comprises amino acids: G85, G86, G87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 13 and comprises amino acids G85, G86, G87, G88, G89, and E90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 14 and comprises amino acids: G85, G86, G87, G88, G89, and G90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 14 and comprises amino acids G85, G86, G87, G88, G89, and G90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 80% sequence identity to SEQ ID NO: 15 and comprises amino acids: G85, G86, E87, G88, G89, and G90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 15 and comprises amino acids G85, G86, E87, G88, G89, and G90.

[0164] In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 23 and comprises amino acids: G80, G81, G82, S83, G85, G86, S87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 23 and comprises amino acids G80, G81, G82, S83, G85, G86, S87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 24 and comprises

amino acids: G80, G81, G82, G83, G85, G86, G87, G88, G89, G90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 24 and comprises amino acids G80, G81, G82, G83, G85, G86, G87, G88, G89, G90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 25 and comprises amino acids: G80, G81, G82, S83, G85, G86, E87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 25 and comprises amino acids G80, G81, G82, S83, G85, G86, E87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 26 and comprises amino acids: G80, G81, G82, E83, G85, G86, E87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 26 and comprises amino acids G80, G81, G82, E83, G85, G86, E87, G88, G89, G90, S91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 27 and comprises amino acids: G82, G83, S85, G86, G87, G88, S89, G90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 27 and comprises amino acids G82, G83, S85, G86, G87, G88, S89, G90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 28 and comprises amino acids: G82, G83, E85, G86, G87, G88, S89, G90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 28 and comprises amino acids G82, G83, E85, G86, G87, G88, S89, G90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 29 and comprises amino acids: G80, G81, G82, G83, E85, G86, G87, G88, G89, G90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 29 and comprises amino acids G80, G81, G82, G83, E85, G86, G87, G88, G89, G90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 30 and comprises amino acids: G80, G81, G82, S83, G85, G86, S87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 30 and comprises amino acids G80, G81, G82, S83, G85, G86, S87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 31 and comprises amino acids: G80, G81, G82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence

identity to SEQ ID NO: 31 and comprises amino acids G80, G81, G82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 32 and comprises amino acids: G82, G83, S85, G86, G87, G88, S89, E90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 32 and comprises amino acids G82, G83, S85, G86, G87, G88, S89, E90, G91 and S92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 33 and comprises amino acids: G82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 33 and comprises amino acids G82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 34 and comprises amino acids: G85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 34 and comprises amino acids G85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 35 and comprises amino acids: G82, G83, G85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 35 and comprises amino acids G82, G83, G85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 36 and comprises amino acids: G82, G83, G85, E86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 36 and comprises amino acids G82, G83, G85, E86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 37 and comprises amino acids: G82, G83, E85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 37 and comprises amino acids G82, G83, E85, G86, G87, G88, G89, E90, and G91. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 38 and comprises amino acids: G80, G81, G82, G83, E84, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 38 and comprises amino acids G80, G81, G82, G83, E84, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 39 and comprises amino acids: G80, G81, G82, E83, G85, G86, G87, G88, G89, E90, G91 and G92. In some

embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 39 and comprises amino acids G80, G81, G82, E83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 80% sequence identity to SEQ ID NO: 40 and comprises amino acids: G80, G81, E82, G83, G85, G86, G87, G88, G89, E90, G91 and G92. In some embodiments, an IL-21 polypeptide comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity to SEQ ID NO: 40 and comprises amino acids G80, G81, E82, G83, G85, G86, G87, G88, G89, E90, G91 and G92.

[0165] In some embodiments, an IL-21 polypeptide comprises QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKAQLX$_1$SAN TGNNERIINVSIKKLX$_2$RKPPX$_3$X$_4$X$_5$X$_6$X$_7$X$_8$X$_9$X$_{10}$X$_{11}$X$_{12}$X$_{13}$X$_{14}$X$_{15}$CPSCDSYEKKPPKEFLER FKSLLQKMIHQHLSSRTHGSEDS (SEQ ID NO: 383), wherein X$_1$ = A, E, K; X$_2$ = A, E, K; X$_3$ = G, S; X$_4$ = G, T; X$_5$ = G, E, N; X$_6$ = G, S, E, A; X$_7$ = E, G; X$_8$ = G, E, S, R; X$_9$ = G, E, R; X$_{10}$ = S, G, E, Q; X$_{11}$ = G, K; X$_{12}$ = G, S, H; X$_{13}$ = A, E, S, G, R; X$_{14}$ = S, G, L; X$_{15}$ = G, S, T; and wherein at least one amino acid residue is not the amino acid residue at the identical position set forth in SEQ ID NO: 1. In some embodiments, at least one of X$_8$, X$_9$, X$_{11}$, X$_{12}$, and X$_{13}$ is not the amino acid residue at the identical position set forth in SEQ ID NO: 1.

[0166] In some embodiments, the IL-21 polypeptide comprises the amino acid sequence: QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKAQLKSANT GNNERIINVSIKKLKRKPPX$_1$X$_2$X$_3$X$_4$GX$_5$X$_6$X$_7$X$_8$X$_9$X$_{10}$X$_{11}$X$_{12}$CPSCDSYEKKPPKEFLERFKSL LQKMIHQHLSSRTHGSEDS (SEQ ID NO: 380), wherein X$_1$ = G, S; X$_2$ = G, T; X$_3$ = G, E, N; X$_4$ = G, S, E, A; X$_5$ = G, E, S, R; X$_6$ = G, E, R; X$_7$ = S, G, E, Q; X$_8$ = G, K; X$_9$ = G, S, H; X$_{10}$ = A, E, S, G, R; X$_{11}$ = S, G, L; and X$_{12}$ = G, S, T, provided at least one of X$_5$, X$_6$, X$_8$, X$_9$, and X$_{10}$ is not the amino acid residue at the identical position set forth in SEQ ID NO: 1. In some embodiments, X$_5$ = G, X$_6$ = G, X$_8$ = G, X$_{10}$ = A. In some embodiments, X$_5$ = G, X$_6$ = G, X$_8$ = G, X$_{10}$ = E. In some embodiments, X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = E, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = S. In some embodiments, X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = E. In some embodiments, X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = E. In some embodiments, X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = S. In some embodiments, X$_5$ = G, X$_6$ = G, X$_7$ = G, X$_8$ = G, X$_9$ = G, X$_{10}$ = E. In some embodiments, X$_5$ = G, X$_6$ = G, X$_7$ = G, X$_8$ = G, X$_9$ = G, X$_{10}$ = G. In some embodiments, X$_5$ = G, X$_6$ = G, X$_7$ = E, X$_8$ = G, X$_9$ = G, X$_{10}$ = G. In some embodiments, X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = G, X$_{11}$ = S, X$_{12}$ = G. In some embodiments, X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = G, X$_5$ = G, X$_6$ = G, X$_7$ = G, X$_8$ = G, X$_9$ = G, X$_{10}$ = G, X$_{11}$ = G, X$_{12}$ = G. In some embodiments, X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = G, X$_6$ = G, X$_7$ =

E, X_8 = G, X_9 = G, X_{10} = G, X_{11} = S, X_{12} = G. In some embodiments, X_1 = G, X_2 = G, X_3 = G, X_4 = E, X_5 = G, X_6 = G, X_7 = E, X_8 = G, X_9 = G, X_{10} = G, X_{11} = S, X_{12} = G. In some embodiments, X_3 = G, X_4 = G, X_5 = S, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = G, X_{11} = G, X_{12} = S. In some embodiments, X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = G, X_{11} = G, X_{12} = S. In some embodiments, X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = G, X_{11} = G, X_{12} = G. In some embodiments, X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = G, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G. In some embodiments, X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G. In some embodiments, X_3 = G, X_4 = G, X_5 = S, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = E, X_{11} = G, X_{12} = S. In some embodiments, X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G. In some embodiments, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G. In some embodiments, X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G. In some embodiments, X_3 = G, X_4 = G, X_5 = G, X_6 = E, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G. In some embodiments, X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G. In some embodiments, X_1 = G, X_2 = G, X_3 = G, X_4 = E, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G. In some embodiments, X_1 = G, X_2 = G, X_3 = E, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G.

[0167] In some embodiments, the IL-21 polypeptide comprises an amino acid sequence starting at position 78 as defined by SEQ ID NO: 1, wherein the amino acid sequence is $PPX_1X_2X_3X_4GX_5X_6X_7X_8X_9X_{10}X_{11}X_{12}CP$ (SEQ ID NO: 386), wherein X_1 = G, S; X_2 = G, T; X_3 = G, E, N; X_4 = G, S, E, A; X_5 = G, E, S, R; X_6 = G, E, R; X_7 = S, G, E, Q; X_8 = G, K; X_9 = G, S, H; X_{10} = A, E, S, G, R; X_{11} = S, G, L; and X_{12} = G, S, T, provided at least one of X_5, X_6, X_8, X_9, and X_{10} is not the amino acid residue at the identical position set forth in SEQ ID NO: 1.

[0168] In some embodiments, the IL-21 polypeptide comprises an amino acid sequence starting at position 80 as defined by SEQ ID NO: 1, wherein the amino acid sequence is $X_1X_2X_3X_4GX_5X_6X_7X_8X_9X_{10}X_{11}X_{12}$, wherein X_1 = G, S; X_2 = G, T; X_3 = G, E, N; X_4 = G, S, E, A; X_5 = G, E, S, R; X_6 = G, E, R; X_7 = S, G, E, Q; X_8 = G, K; X_9 = G, S, H; X_{10} = A, E, S, G, R; X_{11} = S, G, L; and X_{12} = G, S, T, provided at least one of X_5, X_6, X_8, X_9, and X_{10} is not the amino acid residue at the identical position set forth in SEQ ID NO: 1.

Reduced IL-21R binding

[0169] In some embodiments, the disclosure provides an IL-21 polypeptide or a functional fragment or a variant thereof with reduced binding affinity to IL-21R. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least one mutation that

reduces its binding affinity to IL-21R. Such mutations are in some embodiments in one or more positions selected from the group consisting of: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117, wherein the position numbering is number according to the amino acid sequence of SEQ ID NO: 1. In some embodiments, such mutations are in one or more positions selected from the group consisting of: R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, and K117, wherein the position numbering is number according to the amino acid sequence of SEQ ID NO: 1.In some embodiments, the mutation at position R5 comprises an amino acid substitution selected from the group consisting of A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, the mutation at position I8 comprises an amino acid substitution selected from the group consisting of Q, H, E. In some embodiments, the mutation at position R9 comprises an amino acid substitution selected from the group consisting of A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, the mutation at position R11 comprises an amino acid substitution selected from the group consisting of D or E. In some embodiments, the mutation at position Q12 comprises an amino acid substitution selected from the group consisting of L, I, or Y. In some embodiments, the mutation at position L13 comprises an amino acid substitution selected from the group consisting of F or R. In some embodiments, the mutation at position I14 comprises an amino acid substitution selected from the group consisting of D or E. In some embodiments, the mutation at position D15 comprises an amino acid substitution selected from the group consisting of R, K, H, L, Y, or F. In some embodiments, the mutation at position I16 comprises an amino acid substitution selected from the group consisting of A, S or R. In some embodiments, the mutation at position V17 comprises an amino acid substitution selected from the group consisting of I or A. In some embodiments, the mutation at position D18 comprises an amino acid substitution selected from the group consisting of A, K, or R. In some embodiments, the mutation at position Q19 comprises an amino acid substitution selected from the group consisting of L, or Y. In some embodiments, the mutation at position Y23 comprises an amino acid substitution of E. In some embodiments, the mutation at position R65 comprises an amino acid substitution selected from the group consisting of G, S, E, D or A. In some embodiments, the mutation at position S70 comprises an amino acid substitution selected from the group consisting of H, Y, L, V or F. In some embodiments, the mutation at position K72 comprises an amino acid substitution selected from the group consisting of G, S, E, D or A. In some embodiments, the mutation at position K73 comprises an amino acid substitution selected from the group consisting of A, Y, L, F, G, S, T, E, or D. In some embodiments, the mutation at position L74 comprises an amino acid substitution selected from the group consisting of I, F, V, or M. In some embodiments, the mutation at position K75 comprises an amino acid substitution selected from the group

consisting of G, S, E, D or A. In some embodiments, the mutation at position R76 comprises an amino acid substitution selected from the group consisting of A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, the mutation at position K77 comprises an amino acid substitution selected from the group consisting of G, S, E, D or A. In some embodiments, the mutation at position S80 comprises an amino acid substitution of H, A, G, E, or D. In some embodiments, the mutation at position Q116 comprises an amino acid substitution is Y. In some embodiments, the mutation at position K117 comprises an amino acid substitution selected from the group consisting of A, D, or E.

[0170] In some cases, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least one amino acid substitution selected from: R5F, R5A, R5E, R5S, R5T, R5N, R5Q, R5V, R5I, R5L, R5Y, I8E, R9A, R9D, R9E, R9H, R9S, R9T, R9N, R9G, R9V, R9I, R9L, R9Y, R11D, R11E, L13F, L13R, I14D, I16A, I16S, I16R, V17I, V17A, D18A, K72A, K72E, K73A, K73E, K75A, K75E, L74I, L74F, L74M, L74V, R76E, R76F, R76A, R76N, R76D, R76S, R76T, R76Q, R76V, R76I, R76L, R76Y, R76M, K77A, K77E, and K117A. In some cases, an IL-21 polypeptide or a functional fragment or a variant thereof comprises amino acid substitution is R76E or R76Q.

[0171] An example sequence for an IL-21 polypeptide or a functional fragment or a variant thereof of this disclosure is provided as follows:
QGQDX$_1$HMX$_2$X$_3$MX$_4$X$_5$LX$_6$X$_7$IVX$_8$X$_9$LKNX$_{10}$VNDLVPEFLPAPEDVETNCEWSAFSCFQKA QLKSANTGNNEX$_{11}$IINVX$_{12}$IX$_{13}$X$_{14}$LX$_{15}$X$_{16}$X$_{17}$PPX$_{18}$TNAGRRQKHRLTCPSCDSYEKKPPKE FLERFKSLLX$_{19}$X$_{20}$MIHQHLSSRTHGSEDS (SEQ ID NO: 22). In some embodiments, X$_1$ = R, A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, X$_2$ = I, Q, H, E. In some embodiments, X$_3$= R, A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, X$_4$ = R, D or E. In some embodiments, X$_5$ = Q, L, I, or Y. In some embodiments, X$_6$ = I, D or E. In some embodiments, X$_7$ = D, R, K, H, L, Y, or F. In some embodiments, X$_8$ = D, A, K, or R. In some embodiments, X$_9$ = Q, L, or Y. In some embodiments, X$_{10}$ = Y or E. In some embodiments, X$_{11}$ = R, G, S, E, D, or A. In some embodiments, X$_{12}$= S, H, Y, L, V, or F. In some embodiments, X$_{13}$ = K, G, S, E, D, or A. In some embodiments, X$_{14}$ = K, A, Y, L, F, G, S, T, E, A, or D. In some embodiments, X$_{15}$ = K, G, S, E, D, or A. In some embodiments, X$_{16}$ = R, A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, X$_{17}$ = K, G, S, E, D, or A. In some embodiments, X$_{18}$ = S, H, A, G, E, or D. In some embodiments, X$_{19}$ = Q or Y. In some embodiments, X$_{20}$ = K, A, D, or E.

[0172] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof with reduced binding to IL-21R comprises the sequence:
QGQDX$_1$HMX$_2$X$_3$MX$_4$QX$_5$X$_6$DX$_7$X$_8$X$_9$QLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKA

QLKSANTGNNERIINVSIX$_{10}$X$_{11}$X$_{12}$X$_{13}$X$_{14}$X$_{15}$PPSTNAGRRQKHRLTCPSCDSYEKKPPKEFLE RFKSLLQ X$_{16}$MIHQHLSSRTHGSEDS (SEQ ID NO: 384), wherein X$_1$ = F, A, E, S, T, N, Q, V, I, L, Y, R; X$_2$ = E, I; X$_3$ = A, D, E, H, S, T, N, G, V, I, L, Y, R; X$_4$ = D, E, R; X$_5$ = F, R, L; X$_6$ = D, I; X$_7$ = A, S, R, I; X$_8$ = I, A, V; X$_9$ = A, D; X$_{10}$ = A, E, K; X$_{11}$ = A, E, K; X$_{12}$ = I, F, M, L; X$_{13}$ = A, K, E; X$_{14}$ = E, F, A, N, D, S, T, Q, V, I, L, Y, M, R; X$_{15}$ = A, E, K; X$_{16}$ = A, K; provided at least one of X$_1$-X$_{16}$ is not the amino acid residue at the identical position set forth in SEQ ID NO: 1.

[0173] In some embodiments, the disclosure provides an IL-21 polypeptide comprising an amino acid sequence at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to an amino acid sequence selected from SEQ ID NOs: 313-371. In some embodiments, the disclosure provides an IL-21 polypeptide comprising an amino acid sequence selected from SEQ ID NOs: 313-371.

Combination of Mutations

[0174] An IL-21 polypeptide or a functional fragment of a variant thereof, as described herein, comprises, in some embodiments, a combination of (a) mutations that alter its isoelectric point relative to a wild-type IL-21 (SEQ ID NO: 1) and (b) mutations that attenuate its binding to IL-21R, relative to the wild-type IL-21. In some embodiments, a mutation in group **(a)** is at a position selected from positions: K56, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92; and a mutation in group **(b)** is at a position selected from the group consisting of: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117, wherein the position numbering is according to the amino acid sequence of SEQ ID NO: 1. In some embodiments, a mutation in group **(a)** is at a position selected from positions: S80, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92; and a mutation in group **(b)** is at a position selected from the group consisting of: R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, and K117, wherein the position numbering is according to the amino acid sequence of SEQ ID NO: 1. In some embodiments, a mutation in group **(a)** is at a position selected from positions: S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, and T92; and a mutation in group **(b)** is at a position selected from the group consisting of: R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, and K117, wherein the position numbering is according to the amino acid sequence of SEQ ID NO: 1. In some embodiments, a mutation in group **(a)** is at a position selected from positions: S80, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92; and a mutation in group **(b)** is at position R76, wherein the position numbering is according to the amino acid sequence of SEQ ID NO: 1.

[0175] In some embodiments, an IL-21 polypeptide comprises a consensus sequence as follows:

QGQDX$_1$HMX$_2$X$_3$MX$_4$X$_5$LX$_6$X$_7$IVX$_8$X$_9$LKNX$_{10}$VNDLVPEFLPAPEDVETNCEWSAFSCFQKA QLX$_{11}$SANTGNNEX$_{12}$IINVX$_{13}$IX$_{14}$X$_{15}$LX$_{16}$X$_{17}$X$_{18}$PPX$_{19}$X$_{20}$X$_{21}$X$_{22}$X$_{23}$X$_{24}$X$_{25}$X$_{26}$X$_{27}$X$_{28}$X$_{29}$X$_{30}$X$_{31}$CPSCDSYEKKPPKEFLERFKSLLX$_{32}$X$_{33}$MIHQHLSSRTHGSEDS (SEQ ID NO: 3). In some embodiments, X$_1$ = R, A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, X$_2$ = I, Q, H, E. In some embodiments, X$_3$= R, A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, X$_4$ = R, D or E. In some embodiments, X$_5$ = Q, L, I, or Y. In some embodiments, X$_6$ = I or D or E. In some embodiments, X$_7$ = D, R, K, H, L, Y, or F. In some embodiments, X$_8$ = D, A, K, or R. In some embodiments, X$_9$ = Q, L, or Y. In some embodiments, X$_{10}$ = Y or E. X$_{11}$ = G, S, E, D, or A. In some embodiments, X$_{12}$ = R, G, S, E, D, or A. In some embodiments, X$_{13}$= S, H, Y, L, V, or F. In some embodiments, X$_{14}$ = K, G, S, E, D, or A. In some embodiments, X$_{15}$ = K, A, Y, L, F, G, S, T, E, A, or D. In some embodiments, X$_{16}$ = K, G, S, E, D, or A. In some embodiments, X$_{17}$ = R, A, D, E, S, T, N, Q, V, I, L, Y, or F. In some embodiments, X$_{18}$ = K, G, S, E, D, or A. In some embodiments, X$_{19}$ = S, H, A, G, E, or D. In some embodiments, X$_{20}$ = G, S, E, D, or A. In some embodiments, X$_{21}$ = G, S, E, D, or A. In some embodiments, X$_{22}$ = G, S, E, or D. In some embodiments, X$_{23}$ = A, S, E, or D. In some embodiments, X$_{24}$ = G, S, E, D, or A. In some embodiments, X$_{25}$ = G, S, E, D, or A. In some embodiments, X$_{26}$ = G, S, E, D, or A. In some embodiments, X$_{27}$ = G, S, E, D, or A. In some embodiments, X$_{28}$ = G, S, E, D, or A. In some embodiments, X$_{29}$ = G, S, E, D, or A. In some embodiments, X$_{30}$ = G, S, E, D, or A. In some embodiments, X$_{31}$ = G, S, E, D, or A. In some embodiments, X$_{32}$ = Y. In some embodiments, X$_{33}$ = A, D, or E.

[0176] In some embodiments, an IL-21 polypeptide comprises a consensus sequence as follows:

QGQDX$_1$HMX$_2$X$_3$MX$_4$QX$_5$X$_6$DX$_7$X$_8$X$_9$QLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKA QLX$_{10}$SANTGNNERIINVSIX$_{11}$X$_{12}$X$_{13}$X$_{14}$X$_{15}$X$_{16}$PPX$_{17}$X$_{18}$X$_{19}$X$_{20}$X$_{21}$X$_{22}$X$_{23}$X$_{24}$X$_{25}$X$_{26}$X$_{27}$X$_{28}$X$_{29}$C PSCDSYEKKPPKEFLERFKSLLQ X$_{30}$MIHQHLSSRTHGSEDS (SEQ ID NO: 385), wherein X$_1$ = F, A, E, S, T, N, Q, V, I, L, Y, R; X$_2$ = E, I; X$_3$ = A, D, E, H, S, T, N, G, V, I, L, Y, R; X$_4$ = D, E, R; X$_5$ = F, R, L; X$_6$ = D, I; X$_7$ = A, S, R, I; X$_8$ = I, A, V; X$_9$ = A, D; X$_{10}$ = A, E, K; X$_{11}$ = A, E, K; X$_{12}$ = A, E, K; X$_{13}$ = I, F, M, L; X$_{14}$ = A, K, E; X$_{15}$ = E, F, A, N, D, S, T, Q, V, I, L, Y, M, R; X$_{16}$ = A, E, K; X$_{17}$ = G, S; X$_{18}$ = G, T; X$_{19}$ = G, E, N; X$_{20}$ = G, S, E, A; X$_{21}$ = E, G; X$_{22}$ = G, E, S, R; X$_{23}$ = G, E, R; X$_{24}$ = S, G, E, Q; X$_{25}$ = G, K; X$_{26}$ = G, S, H; X$_{27}$ = A, E, S, G, R; X$_{28}$ = S, G, L; X$_{29}$ = G, S, T; X$_{30}$ = A, K, provided at least one of X$_1$-X$_9$ X$_{11}$-X$_{16}$ and X$_{30}$ is not the amino acid residue at the identical position set forth in SEQ ID NO: 1 and reduces binding to IL-21R, and provided at least

one of X_{10} and X_{17}-X_{29} is not the amino acid residue at the identical position set forth in SEQ ID NO: 1 and reduces the isoelectric point relative to human IL-21 (SEQ ID NO: 1).

[0177] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprising a combination of mutations comprises at least one mutation in a position of SEQ ID NO: 2. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprising a combination of mutations comprises at least one mutation in a position selected from the group consisting of positions: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117, wherein the position numbering is according to SEQ ID NO: 2. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprising a combination of mutations comprises at least one mutation in a position selected from the group consisting of positions: R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, and K117, wherein the position numbering is according to SEQ ID NO: 2.

[0178] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprising a combination of mutations comprises at least one mutation in a position of SEQ ID NO: 40. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprising a combination of mutations comprises at least one mutation in a position selected from the group consisting of positions: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117, wherein the position numbering is according to SEQ ID NO: 40. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprising a combination of mutations comprises at least one mutation in a position selected from the group consisting of positions: R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, and K117, wherein the position numbering is according to SEQ ID NO: 40.

[0179] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises (i) four or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the four or more amino acid substitutions are within S80 to T92 of SEQ ID NO: 1, wherein at least four amino acid substitutions are at residues R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution selected from (a) R11D; (b) R11E; (c) I14D, D18A and K117A; (d) R76E; (e) R5F; (f) R76F; (g) I8E; (h) R5A; (i) R5E; (j) R5S; (k) R5T; (l) R5N; (m) R5Q; (n) R5V; (o) R5I; (p) R5L; (q) R5Y; (r) R76A; (s) R76N; (t) R76D; (u) R76S; (v) R76T; (w) R76Q; (x) R76V; (y) R76I; (z) R76L; (aa) R76Y; (bb) K77A; (cc) K77E; (dd) K72A; (ee) K72E; (ff) K75A; (gg) K75E; (hh) K73A; (ii) K73E; (jj) R5F and K77A; (kk) R5F and K77E; (ll) R5F and K72A;

(mm) R5F and K72E; (nn) R5F and K76A; (oo) R5F and K76E; (pp) K73A and K76F; (qq) K73E and K76F; (rr) R9A; (ss) R9D; (tt) R9E; (uu) R9H; (vv) R9S; (ww) R9T; (xx) R9N; (zz) R9G; (aaa) R9V; (bbb) R9I; (ccc) R9L; (ddd) R9Y; (eee) K72A and R76F; (fff) K75A and R76F; (ggg) R76F and K77A; (hhh) K75E and R76F; (iii) V17I and L74I; (jjj) I16A and L74F; (kkk) I16S, V17I, and L74V; (lll) I16R, V17I, and L74I; (mmm) L13F, I16A, V17A and L74M; and (nnn) L13R, I16A, V17I and L74I.

[0180] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises (i) an amino acid sequence selected from SEQ ID Nos: 2, 4-15 and 23-40, and (ii) at least one amino acid substitution selected from (a) R11D; (b) R11E; (c) I14D, D18A and K117A; (d) R76E; (e) R5F; (f) R76F; (g) I8E; (h) R5A; (i) R5E; (j) R5S; (k) R5T; (l) R5N; (m) R5Q; (n) R5V; (o) R5I; (p) R5L; (q) R5Y; (r) R76A; (s) R76N; (t) R76D; (u) R76S; (v) R76T; (w) R76Q; (x) R76V; (y) R76I; (z) R76L; (aa) R76Y; (bb) K77A; (cc) K77E; (dd) K72A; (ee) K72E; (ff) K75A; (gg) K75E; (hh) K73A; (ii) K73E; (jj) R5F and K77A; (kk) R5F and K77E; (ll) R5F and K72A; (mm) R5F and K72E; (nn) R5F and K76A; (oo) R5F and K76E; (pp) K73A and K76F; (qq) K73E and K76F; (rr) R9A; (ss) R9D; (tt) R9E; (uu) R9H; (vv) R9S; (ww) R9T; (xx) R9N; (zz) R9G; (aaa) R9V; (bbb) R9I; (ccc) R9L; (ddd) R9Y; (eee) K72A and R76F; (fff) K75A and R76F; (ggg) R76F and K77A; (hhh) K75E and R76F; (iii) V17I and L74I; (jjj) I16A and L74F; (kkk) I16S, V17I, and L74V; (lll) I16R, V17I, and L74I; (mmm) L13F, I16A, V17A and L74M; and (nnn) L13R, I16A, V17I and L74I.

[0181] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11D mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 16. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11D mutation and comprises at least about 75 % sequence identity SEQ ID NO: 16. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11D mutation and comprises at most about 100 % sequence identity SEQ ID NO: 16. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11D mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence

identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 16. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11D mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 16.

[0182] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 17. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 17. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11E mutation and comprises at most about 100 %

sequence identity SEQ ID NO: 17. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11E mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 17. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R11E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about

85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 17.

[0183] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof or a functional fragment or a variant thereof comprises a I14D, D18A and K117A mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 18. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof or a functional fragment or a variant thereof comprises a I14D, D18A and K117A mutation and comprises at least about 75 % sequence identity SEQ ID NO: 18. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof or a functional fragment or a variant thereof comprises a I14D, D18A and K117A mutation and comprises at most about 100 % sequence identity SEQ ID NO: 18. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof or a functional fragment or a variant thereof comprises a I14D, D18A and K117A mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 %

sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity to SEQ ID NO: 18. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof or a functional fragment or a variant thereof comprises a I14D, D18A and K117A mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 18.

[0184] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 19. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 19. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 19. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence

identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 19. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 19.

[0185] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R5F mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 20. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R5F mutation and comprises at least about 75 % sequence identity SEQ ID NO: 20. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at most about 100 % sequence identity SEQ ID NO: 20. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R5F mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90

% sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 29. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R5F mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 20.

[0186] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 21. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises at least about 75 % sequence identity SEQ ID NO: 21. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises at most about 100 % sequence identity SEQ ID NO: 21. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence

identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity to SEQ ID NO: 21. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 21.

[0187] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an I8E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 41. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an I8E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 41. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an I8E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 41. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an I8E mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence

identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 41. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an I8E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 41.

[0188] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5A mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 42. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5A mutation and comprises at least about 75 % sequence identity SEQ ID NO: 42. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5A mutation and comprises at most about 100 % sequence identity SEQ ID NO: 42. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5A mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence

identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 42. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5A mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 42.

[0189] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 43. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 43. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 43. In some embodiments, an IL-21 polypeptide or a functional

fragment or a variant thereof comprises an R5E mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 43. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 %

sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 43.

[0190] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5S mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 44. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5S mutation and comprises at least about 75 % sequence identity SEQ ID NO: 44. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5S mutation and comprises at most about 100 % sequence identity SEQ ID NO: 44. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5S mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence

identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 44. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5S mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 44.

[0191] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5T mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 45. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5T mutation and comprises at least about 75 % sequence identity SEQ ID NO: 45. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5T mutation and comprises at most about 100 % sequence identity SEQ ID NO: 45. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5T mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence

identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 45. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5T mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 45.

[0192] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5N mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 46. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5N mutation and comprises at least about 75 % sequence identity SEQ ID NO: 46. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5N mutation and comprises at most about 100 % sequence identity SEQ ID NO: 46. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5N mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence

identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 46. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5N mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 46.

[0193] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5Q mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 47. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5Q mutation and comprises at least about 75 % sequence identity SEQ ID NO: 47. In some embodiments, an IL-21 polypeptide or a functional

fragment or a variant thereof comprises an R5Q mutation and comprises at most about 100 % sequence identity SEQ ID NO: 47. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5Q mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 47. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises

an R5Q mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 47.

[0194] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5V mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 48. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5V mutation and comprises at least about 75 % sequence identity SEQ ID NO: 48. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5V mutation and comprises at most about 100 % sequence identity SEQ ID NO: 48. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5V mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence

identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 48. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5V mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 48.

[0195] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5I mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 49. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5I mutation and comprises at least about 75 % sequence identity SEQ ID NO: 49. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5I mutation and comprises at most about 100 % sequence identity SEQ ID NO: 49. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5I mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence

identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 49. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5I mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 49.

[0196] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5L mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 50. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5L mutation and comprises at least about 75 % sequence identity SEQ ID NO: 50. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5L mutation and comprises at most about 100 % sequence identity SEQ ID NO: 50. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5L mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence

identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 50. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5L mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 50.

[0197] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5Y mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 51. In some embodiments, an IL-21 polypeptide or a functional

fragment or a variant thereof comprises an R5Y mutation and comprises at least about 75 % sequence identity SEQ ID NO: 51. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5Y mutation and comprises at most about 100 % sequence identity SEQ ID NO: 51. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5Y mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence

identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 51. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R5Y mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 51.

[0198] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76A mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 52. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76A mutation and comprises at least about 75 % sequence identity SEQ ID NO: 52. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76A mutation and comprises at most about 100 % sequence identity SEQ ID NO: 52. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76A mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence

identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 52. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76A mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 52.

[0199] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76N mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 53. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76N mutation and comprises at least about 75 % sequence identity SEQ ID NO: 53. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76N mutation and comprises at most about 100 % sequence identity SEQ ID NO: 53. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76N mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence

identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 53. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76N mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 53.

[0200] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76D mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 54. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76D mutation and comprises at least about 75 % sequence identity SEQ ID NO: 54. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76D mutation and comprises at most about 100 % sequence identity SEQ ID NO: 54. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76D mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence

identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 54. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76D mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 54.

[0201] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76S mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 55. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76S mutation and comprises at least about 75 % sequence identity SEQ ID NO: 55. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76S mutation and comprises at most about 100 % sequence identity SEQ ID NO: 55. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76S mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence

identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 55. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76S mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 55.

[0202] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76T mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 56. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76T mutation and comprises at least about 75 % sequence identity SEQ ID NO: 56. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76T mutation and comprises at most about 100 % sequence identity SEQ ID NO: 56. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76T mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence

identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 6 In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76T mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 56.

[0203] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Q mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 57. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Q mutation and comprises at least about 75 % sequence identity SEQ ID NO: 57. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Q mutation and comprises at most about 100 % sequence identity SEQ ID NO: 57. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Q mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence

identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 57. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Q mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 57.

[0204] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76V mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 58. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76V mutation and comprises at least about 75 % sequence identity SEQ ID NO: 58. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76V mutation and comprises at most about 100 % sequence identity SEQ ID NO: 58. In some embodiments, an IL-21 polypeptide or a functional

fragment or a variant thereof comprises an R76V mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 58. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76V mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about

96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 58.

[0205] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76I mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 59. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76I mutation and comprises at least about 75 % sequence identity SEQ ID NO: 59. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76I mutation and comprises at most about 100 % sequence identity SEQ ID NO: 59. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76I mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence

identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 59. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76I mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 59.

[0206] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76L mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 60. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76L mutation and comprises at least about 75 % sequence identity SEQ ID NO: 60. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76L mutation and comprises at most about 100 % sequence identity SEQ ID NO: 60. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76L mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence

identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 60. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76L mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 60.

[0207] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Y mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 61. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Y mutation and comprises at least about 75 % sequence identity SEQ ID NO: 61. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Y mutation and comprises at most about 100 % sequence identity SEQ ID NO: 61. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Y mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence

identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 61. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R76Y mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 61.

[0208] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K77A mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 62. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K77A mutation and comprises at least about 75 % sequence identity SEQ ID NO: 62. In some embodiments, an IL-21 polypeptide or a functional

fragment or a variant thereof comprises a K77A mutation and comprises at most about 100 % sequence identity SEQ ID NO: 62. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K77A mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 62. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a

K77A mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 62.

[0209] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K77E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 63. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K77E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 63. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K77E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 63. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K77E mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence

identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 63. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K77E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 63.

[0210] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72A mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 64. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72A mutation and comprises at least about 75 % sequence identity SEQ ID NO: 64. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72A mutation and comprises at most about 100 % sequence identity SEQ ID NO: 64. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72A mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence

identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 64. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72A mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 64.

[0211] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 65. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 65. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 65. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72E mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence

identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 65. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K72E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 65.

[0212] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75A mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 66. In some embodiments, an IL-21 polypeptide or a functional

fragment or a variant thereof comprises a K75A mutation and comprises at least about 75 % sequence identity SEQ ID NO: 66. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75A mutation and comprises at most about 100 % sequence identity SEQ ID NO: 66. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75A mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence

identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 66. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75A mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 66.

[0213] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 67. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 67. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 67. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75E mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence

identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 67. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K75E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 67.

[0214] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73A mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 68. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73A mutation and comprises at least about 75 % sequence identity SEQ ID NO: 68. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73A mutation and comprises at most about 100 % sequence identity SEQ ID NO: 68. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73A mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence

identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 68. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73A mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 68.

[0215] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 69. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 69. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 69. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73E mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence

identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 69. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a K73E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 69.

[0216] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77A mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 70. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77A mutations and comprises at least about 75 % sequence identity SEQ ID NO: 70. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77A mutations and comprises at most about 100 % sequence identity SEQ ID NO: 70. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77A mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 %

sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 70. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77A mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 70.

[0217] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77E mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 71. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77E mutations and comprises at least about 75 % sequence identity SEQ ID NO: 71. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77E mutations and comprises at most about 100 % sequence identity SEQ ID NO: 71. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77E mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 %

sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 71. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K77E mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 71.

[0218] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72A mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 72. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72A mutations and comprises at least about 75 % sequence identity SEQ ID NO: 72. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72A mutations and comprises at most about 100 % sequence identity SEQ ID NO: 72. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72A mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 %

sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 72. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72A mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 72.

[0219] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72E mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 73. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72E mutations and comprises at least

about 75 % sequence identity SEQ ID NO: 73. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72E mutations and comprises at most about 100 % sequence identity SEQ ID NO: 73. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72E mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID

NO: 73. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K72E mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 73.

[0220] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76A mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 74. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76A mutations and comprises at least about 75 % sequence identity SEQ ID NO: 74. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76A mutations and comprises at most about 100 % sequence identity SEQ ID NO: 74. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76A mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 %

sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 74. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76A mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 74.

[0221] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76E mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 75. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76E mutations and comprises at least about 75 % sequence identity SEQ ID NO: 75. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76E mutations and comprises at most about 100 % sequence identity SEQ ID NO: 75. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76E mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 %

sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 75. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R5F and K76E mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 75.

[0222] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73A and K76F mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 76. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73A and K76F mutations and comprises at least about 75 % sequence identity SEQ ID NO: 76. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73A and K76F mutations and comprises at most about 100 % sequence identity SEQ ID NO: 76. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73A and K76F mutations and comprises

about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 76. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73A and K76F mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 %

sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 76.

[0223] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73E and K76F mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 77. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73E and K76F mutations and comprises at least about 75 % sequence identity SEQ ID NO: 77. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73E and K76F mutations and comprises at most about 100 % sequence identity SEQ ID NO: 77. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73E and K76F mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 %

sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 77. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K73E and K76F mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 77.

[0224] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9A mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 78. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9A mutation and comprises at least about 75 % sequence identity SEQ ID NO: 78. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9A mutation and comprises at most about 100 % sequence identity SEQ ID NO: 78. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9A mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence

identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 78. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9A mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 78.

[0225] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9D mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 79. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9D mutation and comprises at least about 75 % sequence identity SEQ ID NO: 79. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9D mutation and comprises at most about 100 % sequence identity SEQ ID NO: 79. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9D mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence

identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 79. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9D mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 79.

[0226] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 80. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9E mutation and comprises at least about 75 %

sequence identity SEQ ID NO: 80. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 80. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9E mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 80. In

some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 80.

[0227] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9H mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 81. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9H mutation and comprises at least about 75 % sequence identity SEQ ID NO: 81. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9HE mutation and comprises at most about 100 % sequence identity SEQ ID NO: 81. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9H mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence

identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 81. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9H mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 81.

[0228] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 82. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises at least about 75 % sequence identity SEQ ID NO: 82. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises at most about 100 % sequence identity SEQ ID NO: 82. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence

identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 82. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 82.

[0229] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9T mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 83. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9T mutation and comprises at least about 75 % sequence identity SEQ ID NO: 83. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9T mutation and comprises at most about 100 % sequence identity SEQ ID NO: 83. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9T mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence

identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 83. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9T mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 83.

[0230] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9N mutation and comprises about 75 % sequence identity to about 100 %

sequence identity to SEQ ID NO: 84. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9N mutation and comprises at least about 75 % sequence identity SEQ ID NO: 84. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9N mutation and comprises at most about 100 % sequence identity SEQ ID NO: 84. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9N mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence

identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 84. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9N mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 84.

[0231] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 85. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises at least about 75 % sequence identity SEQ ID NO: 85. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises at most about 100 % sequence identity SEQ ID NO: 85. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence

identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 85. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 85.

[0232] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9V mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 86. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9V mutation and comprises at least about 75 % sequence identity SEQ ID NO: 86. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9V mutation and comprises at most about 100 % sequence identity SEQ ID NO: 86. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9V mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence

identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 86. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9V mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 86.

[0233] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9I mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 87. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9I mutation and comprises at least about 75 % sequence identity SEQ ID NO: 87. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9I mutation and comprises at most about 100 % sequence identity SEQ ID NO: 87. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9I mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence

identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 87. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9I mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 87.

[0234] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9L mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 88. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9L mutation and comprises at least about 75 % sequence identity SEQ ID NO: 88. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9L mutation and comprises at most about 100 % sequence identity SEQ ID NO: 88. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9L mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence

identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 88. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9L mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 88.

[0235] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9Y mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 89. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9Y mutation and comprises at least about 75 % sequence identity SEQ ID NO: 89. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9Y mutation and comprises at most about 100 % sequence identity SEQ ID NO: 89. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9Y mutation and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence

identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 89. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9Y mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 89.

[0236] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K72A and R76F mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K72A and R76F mutations and comprises at least about 75 % sequence identity SEQ ID NO:90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K72A and R76F mutations and comprises at most about 100 % sequence identity SEQ ID NO: 90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K72A and R76F mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 %

sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 90. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K72A and R76F mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 90.

[0237] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75A and R76F mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 91. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75A and R76F mutations and comprises at least about 75 % sequence identity SEQ ID NO:91. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75A and R76F mutations and comprises at

most about 100 % sequence identity SEQ ID NO: 91. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75A and R76F mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 91. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75A and R76F mutations and comprises about 75 % sequence identity, about 80 %

sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 91.

[0238] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R76F and K77A mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 92. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R76F and K77A mutations and comprises at least about 75 % sequence identity SEQ ID NO:92. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R76F and K77A mutations and comprises at most about 100 % sequence identity SEQ ID NO: 92. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R76F and K77A mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 %

sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 92. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises R76F and K77A mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 92.

[0239] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75E and R76F mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 93. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75E and R76F mutations and comprises at least about 75 % sequence identity SEQ ID NO:93. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75E and R76F mutations and comprises at most about 100 % sequence identity SEQ ID NO: 93. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75E and R76F mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 %

sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 93. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises K75E and R76F mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 93.

[0240] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises V17I and L74I mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 374. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises V17I and L74I mutations and comprises at least about 75 % sequence identity SEQ ID NO:374. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises V17I and L74I mutations and comprises at most about 100 % sequence identity SEQ ID NO: 374. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises V17I and L74I mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 %

sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 374. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises V17I and L74I mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 374.

[0241] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16A and L74F mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 375. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16A and L74F mutations and comprises at least about 75 % sequence identity SEQ ID NO:375. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16A and L74F mutations and comprises at most about 100 % sequence identity SEQ ID NO: 375. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16A and L74F mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 %

sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 375. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16A and L74F mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 375.

[0242] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16S, V17I, and L74V mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 376. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16S, V17I, and L74V mutations and comprises at least about 75 % sequence identity SEQ ID NO:376. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16S, V17I, and L74V mutations and comprises at most about 100 % sequence identity SEQ ID NO: 376. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16S, V17I, and L74V mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence

identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 376. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16S, V17I, and L74V mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 376.

[0243] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16R, V17I, and L74I mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 377. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16R, V17I, and L74I mutations and comprises at least about 75 % sequence identity SEQ ID NO:377. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16R, V17I, and L74I mutations and comprises at most about 100 % sequence identity SEQ ID NO: 377. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16R, V17I, and L74I mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 %

sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 377. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises I16R, V17I, and L74I mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 377.

[0244] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13F, I16A, V17A and L74M mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 378. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13F, I16A, V17A and L74M

mutations and comprises at least about 75 % sequence identity SEQ ID NO:378. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13F, I16A, V17A and L74M mutations and comprises at most about 100 % sequence identity SEQ ID NO: 378. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13F, I16A, V17A and L74M mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 %

sequence identity to about 100 % sequence identity, to SEQ ID NO: 378. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13F, I16A, V17A and L74M mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 378.

[0245] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13R, I16A, V17I and L74I mutations and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 379. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13R, I16A, V17I and L74I mutations and comprises at least about 75 % sequence identity SEQ ID NO:379. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13R, I16A, V17I and L74I mutations and comprises at most about 100 % sequence identity SEQ ID NO: 379. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13R, I16A, V17I and L74I mutations and comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 %

sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 379. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises L13R, I16A, V17I and L74I mutations and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 379.

[0246] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76E mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 94. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76E mutation and comprises at least about 75 % sequence identity SEQ ID NO: 94. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76E mutation and comprises at most about 100 % sequence identity SEQ ID NO: 94. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence

identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 94. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76E mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 94.

[0247] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 95. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises at least about 75 % sequence identity SEQ ID NO: 95. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises at most about 100 % sequence identity SEQ ID NO: 95. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence

identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 95. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76F mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 95.

[0248] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76Q mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 96. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76Q mutation and comprises at least about 75 % sequence identity SEQ ID NO: 96. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76Q mutation and comprises at most about 100 % sequence identity SEQ ID NO: 96. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence

identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 96. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises a R76Q mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 96.

[0249] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 97. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises at least about 75 % sequence identity SEQ ID NO: 97. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises at most about 100 % sequence identity SEQ ID NO: 97. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence

identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 97. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9S mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 97.

[0250] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises about 75 % sequence identity to about 100 % sequence identity to SEQ ID NO: 98. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises at least about 75 % sequence identity SEQ ID NO: 98. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises at most about 100 % sequence identity SEQ ID NO: 98. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises about 75 % sequence identity to about 80 % sequence identity, about 75 % sequence identity to about 85 % sequence identity, about 75 % sequence identity to about 90 % sequence identity, about 75 % sequence identity to about 95 % sequence identity, about 75 % sequence identity to about 96 % sequence identity, about 75 % sequence identity to about 97 % sequence identity, about 75 % sequence identity to about 98 % sequence identity, about 75 % sequence identity to about 99 % sequence identity, about 75 % sequence identity to about 100 % sequence identity, about 80 % sequence identity to about 85 % sequence identity, about 80 % sequence identity to about 90 % sequence identity, about 80 % sequence identity to about 95 % sequence identity, about 80 % sequence identity to about 96 % sequence

identity, about 80 % sequence identity to about 97 % sequence identity, about 80 % sequence identity to about 98 % sequence identity, about 80 % sequence identity to about 99 % sequence identity, about 80 % sequence identity to about 100 % sequence identity, about 85 % sequence identity to about 90 % sequence identity, about 85 % sequence identity to about 95 % sequence identity, about 85 % sequence identity to about 96 % sequence identity, about 85 % sequence identity to about 97 % sequence identity, about 85 % sequence identity to about 98 % sequence identity, about 85 % sequence identity to about 99 % sequence identity, about 85 % sequence identity to about 100 % sequence identity, about 90 % sequence identity to about 95 % sequence identity, about 90 % sequence identity to about 96 % sequence identity, about 90 % sequence identity to about 97 % sequence identity, about 90 % sequence identity to about 98 % sequence identity, about 90 % sequence identity to about 99 % sequence identity, about 90 % sequence identity to about 100 % sequence identity, about 95 % sequence identity to about 96 % sequence identity, about 95 % sequence identity to about 97 % sequence identity, about 95 % sequence identity to about 98 % sequence identity, about 95 % sequence identity to about 99 % sequence identity, about 95 % sequence identity to about 100 % sequence identity, about 96 % sequence identity to about 97 % sequence identity, about 96 % sequence identity to about 98 % sequence identity, about 96 % sequence identity to about 99 % sequence identity, about 96 % sequence identity to about 100 % sequence identity, about 97 % sequence identity to about 98 % sequence identity, about 97 % sequence identity to about 99 % sequence identity, about 97 % sequence identity to about 100 % sequence identity, about 98 % sequence identity to about 99 % sequence identity, about 98 % sequence identity to about 100 % sequence identity, or about 99 % sequence identity to about 100 % sequence identity, to SEQ ID NO: 98. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises an R9G mutation and comprises about 75 % sequence identity, about 80 % sequence identity, about 85 % sequence identity, about 90 % sequence identity, about 95 % sequence identity, about 96 % sequence identity, about 97 % sequence identity, about 98 % sequence identity, about 99 % sequence identity, or about 100 % sequence identity to SEQ ID NO: 98.

[0251] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NOs: 16-21. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NOs: 16-21. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NOs: 16-21. In some embodiments, an IL-21 polypeptide comprises a sequence selected from the group consisting of: SEQ ID NOs: 16-21.

[0252] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NOs: 16-21, 41-93, and 374-379. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NOs: 16-21, 41-93, and 374-379. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NOs: 16-21, 41-93, and 374-379. In some embodiments, an IL-21 polypeptide comprises a sequence selected from the group consisting of: SEQ ID NOs: 16-21, 41-93, and 374-379.

[0253] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NOs: 94-98. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NOs: 94-98. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NOs: 94-98. In some embodiments, an IL-21 polypeptide comprises a sequence selected from the group consisting of: SEQ ID NOs: 97-98.

[0254] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NOs: 16-21, 41-98, and 374-379. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NOs: 16-21, 41-98, and 374-379. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NOs: 16-21, 41-98, and 374-379. In some embodiments, an IL-21 polypeptide comprises a sequence selected from the group consisting of: SEQ ID NOs: 16-21, 41-98, and 374-379.

[0255] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 16 In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 16. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 16. In some embodiments, an IL-21 polypeptide comprises SEQ ID NO: 16.

[0256] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 17. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 17. In some

embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 17. In some embodiments, an IL-21 polypeptide comprises SEQ ID NO: 17.

[0257] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 18. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 18. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 18. In some embodiments, an IL-21 polypeptide comprises SEQ ID NO: 18.

[0258] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 19. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 19. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 19. In some embodiments, an IL-21 polypeptide comprises SEQ ID NO: 19.

[0259] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 20. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 20. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 20. In some embodiments, an IL-21 polypeptide comprises SEQ ID NO: 20.

[0260] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 21. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 21. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 21. In some embodiments, an IL-21 polypeptide SEQ ID NO: 21.

[0261] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 41. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 41. In some

embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 41. In some embodiments, an IL-21 polypeptide SEQ ID NO: 41.

[0262] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 42. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 42. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 42. In some embodiments, an IL-21 polypeptide SEQ ID NO: 42.

[0263] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 43. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 43. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 43. In some embodiments, an IL-21 polypeptide SEQ ID NO: 43.

[0264] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 44. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 44. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 44. In some embodiments, an IL-21 polypeptide SEQ ID NO: 44.

[0265] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 45. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 45. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 45. In some embodiments, an IL-21 polypeptide SEQ ID NO: 45.

[0266] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 46 In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 46. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 46. In some embodiments, an IL-21 polypeptide SEQ ID NO: 46.

[0267] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 47. In some embodiments, an IL-

21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 47. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 47. In some embodiments, an IL-21 polypeptide SEQ ID NO: 47.

[0268] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 48. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 48. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 48. In some embodiments, an IL-21 polypeptide SEQ ID NO: 48.

[0269] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 49. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 49. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 49. In some embodiments, an IL-21 polypeptide SEQ ID NO: 49.

[0270] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 50. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 50. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 50. In some embodiments, an IL-21 polypeptide SEQ ID NO: 50.

[0271] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 51. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 51. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO:5. In some embodiments, an IL-21 polypeptide SEQ ID NO: 51.

[0272] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 52. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 52. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 52. In some embodiments, an IL-21 polypeptide SEQ ID NO: 52.

[0273] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 53. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 53. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 53. In some embodiments, an IL-21 polypeptide SEQ ID NO: 53.

[0274] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 54. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 54. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 54. In some embodiments, an IL-21 polypeptide SEQ ID NO: 54.

[0275] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 55. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 55. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 55. In some embodiments, an IL-21 polypeptide SEQ ID NO: 55.

[0276] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 56. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 56. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 56. In some embodiments, an IL-21 polypeptide SEQ ID NO: 56.

[0277] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 57. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 57. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 57. In some embodiments, an IL-21 polypeptide SEQ ID NO: 57.

[0278] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 58. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 58. In some

embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 58. In some embodiments, an IL-21 polypeptide SEQ ID NO: 58.

[0279] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 59. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 59. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 59. In some embodiments, an IL-21 polypeptide SEQ ID NO: 59.

[0280] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 60. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 60. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 60. In some embodiments, an IL-21 polypeptide SEQ ID NO: 60.

[0281] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 61. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 61. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 61. In some embodiments, an IL-21 polypeptide SEQ ID NO: 61.

[0282] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 62. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 62. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 62. In some embodiments, an IL-21 polypeptide SEQ ID NO: 62.

[0283] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 63. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 63. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 63. In some embodiments, an IL-21 polypeptide SEQ ID NO: 63.

[0284] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 64. In some embodiments, an IL-

21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 64. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 64. In some embodiments, an IL-21 polypeptide SEQ ID NO: 64.

[0285] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 65. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 65. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 65. In some embodiments, an IL-21 polypeptide SEQ ID NO: 65.

[0286] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 66. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 66. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 66. In some embodiments, an IL-21 polypeptide SEQ ID NO: 66.

[0287] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 67. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 67. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 67. In some embodiments, an IL-21 polypeptide SEQ ID NO: 67.

[0288] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 68. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 68. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 68. In some embodiments, an IL-21 polypeptide SEQ ID NO: 68.

[0289] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 69. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 69. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 69. In some embodiments, an IL-21 polypeptide SEQ ID NO: 69.

[0290] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 70. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 70. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 70. In some embodiments, an IL-21 polypeptide SEQ ID NO: 70.

[0291] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 71. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 71. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 71. In some embodiments, an IL-21 polypeptide SEQ ID NO: 71.

[0292] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 72. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 72. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 72. In some embodiments, an IL-21 polypeptide SEQ ID NO: 72.

[0293] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 73. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 73. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 73. In some embodiments, an IL-21 polypeptide SEQ ID NO: 73.

[0294] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 74. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 74. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 74. In some embodiments, an IL-21 polypeptide SEQ ID NO: 74.

[0295] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 75. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 75. In some

embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 75. In some embodiments, an IL-21 polypeptide SEQ ID NO: 75.

[0296] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 76. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 76. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 76. In some embodiments, an IL-21 polypeptide SEQ ID NO: 76.

[0297] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 77. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 77. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 77. In some embodiments, an IL-21 polypeptide SEQ ID NO: 77.

[0298] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 78. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 78. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 78. In some embodiments, an IL-21 polypeptide SEQ ID NO: 78.

[0299] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 79. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 79. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 79. In some embodiments, an IL-21 polypeptide SEQ ID NO: 79.

[0300] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 80. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 80. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 80. In some embodiments, an IL-21 polypeptide SEQ ID NO: 80.

[0301] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 81. In some embodiments, an IL-

21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 81. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 81. In some embodiments, an IL-21 polypeptide SEQ ID NO: 81.

[0302] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 82. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 82. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 82. In some embodiments, an IL-21 polypeptide SEQ ID NO: 82.

[0303] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 83. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 83. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 83. In some embodiments, an IL-21 polypeptide SEQ ID NO: 83.

[0304] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 84. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 84. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 84. In some embodiments, an IL-21 polypeptide SEQ ID NO: 84.

[0305] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 85. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 85. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 85. In some embodiments, an IL-21 polypeptide SEQ ID NO: 85.

[0306] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 86. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 86. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 86. In some embodiments, an IL-21 polypeptide SEQ ID NO: 86.

[0307] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 87. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 87. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 87. In some embodiments, an IL-21 polypeptide SEQ ID NO: 87.

[0308] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 88. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 88. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 88. In some embodiments, an IL-21 polypeptide SEQ ID NO: 88.

[0309] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 89. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 89. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 89. In some embodiments, an IL-21 polypeptide SEQ ID NO: 89.

[0310] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 90. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 90. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 90. In some embodiments, an IL-21 polypeptide SEQ ID NO: 90.

[0311] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 91. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 91. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 91. In some embodiments, an IL-21 polypeptide SEQ ID NO: 91.

[0312] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 92. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 92. In some

embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 92. In some embodiments, an IL-21 polypeptide SEQ ID NO: 92.

[0313] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 93. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 93. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 93. In some embodiments, an IL-21 polypeptide SEQ ID NO: 93.

[0314] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 94. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 94. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 94. In some embodiments, an IL-21 polypeptide SEQ ID NO: 94.

[0315] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 95. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 95. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 95. In some embodiments, an IL-21 polypeptide SEQ ID NO: 95.

[0316] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 96. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 96. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 96. In some embodiments, an IL-21 polypeptide SEQ ID NO: 96.

[0317] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 97. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 97. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 97. In some embodiments, an IL-21 polypeptide SEQ ID NO: 97.

[0318] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 98. In some embodiments, an IL-

21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 98. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 98. In some embodiments, an IL-21 polypeptide SEQ ID NO: 98.

[0319] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 374. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 374. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 374. In some embodiments, an IL-21 polypeptide SEQ ID NO: 374.

[0320] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 375. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 375. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 375. In some embodiments, an IL-21 polypeptide SEQ ID NO: 375.

[0321] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 376. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 376. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 376. In some embodiments, an IL-21 polypeptide SEQ ID NO: 376.

[0322] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 377. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 377. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 377. In some embodiments, an IL-21 polypeptide SEQ ID NO: 377.

[0323] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 378. In some embodiments, an IL-

21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 378. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 378. In some embodiments, an IL-21 polypeptide SEQ ID NO: 378.

[0324] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof comprises at least 75% sequence identity to SEQ ID NO: 379. In some embodiments, an IL-21 polypeptide comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to SEQ ID NO: 379. In some embodiments, an IL-21 polypeptide comprises 75-80%, 80-85%, 85-90%, 90-95%, 95-99% sequence identity to SEQ ID NO: 379. In some embodiments, an IL-21 polypeptide SEQ ID NO: 379.

Antigen Binding Protein or Functional Fragments thereof

[0325] Some embodiments of this disclosure provides a targeted cytokine construct comprising an antigen binding protein (*e.g.*, an antibody or an antigen binding fragment thereof) as a targeting protein and an IL-21 polypeptide as disclosed herein. The antigen binding protein or a functional fragment thereof can, in some embodiments, specifically bind to CD8+ T cells and comprise specificity to at least one antigen selected from the group consisting of CD8-alpha (CD8α or CD8a), CD8-beta (CD8-β), CD8- $\alpha\alpha$, or CD8-αβ. As provided herein, the term "CD8-alpha" can be used interchangeably with "CD8α" or "CD8a". As provided herein, the term "CD8-beta" can be used interchangeably with the terms "CD8-β" or "CD8b". As provided herein the term "CD8- $\alpha\alpha$" can be used interchangeably with "CD8aa". As provided herein the term "CD8-αβ" can be used interchangeably with "CD8ab". In some embodiments, the antibody or antigen binding fragment thereof may specifically bind to a different antigen, such as an immune checkpoint protein.

[0326] In some embodiments, the antigen binding protein or functional fragment thereof comprises an antigen binding domain selected from the group consisting of an scFv, a single domain antibody, VHH, VNAR, sdAbs, nanobody, or a functional fragment thereof.

[0327] In some embodiments, the antigen binding protein or a functional fragment thereof comprises a full-length intact antibody. The full-length intact antibody can comprise: a first polypeptide, arranged from N-to-C terminus, comprising: variable light chain (VL) amino acid sequence, and a light chain constant region amino acid sequence (CL1); a second polypeptide, arranged from N-to-C terminus, comprising: a variable heavy chain (VH) amino acid sequence, a

heavy chain CH1 constant region amino acid sequence, a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence; a third polypeptide, arranged from N-to-C terminus, comprising: a variable heavy chain (VH) amino acid sequence, a heavy chain CH1 constant region amino acid sequence, a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence; and a fourth polypeptide, arranged from N-to-C terminus, comprising: a variable light chain (VL) amino acid sequence, and a light chain constant amino acid sequence (CL1), wherein the CH2 and CH3 domains of each of the second and third polypeptides form an Fc domain.

[0328] In some embodiments, the antigen binding protein or a functional fragment thereof comprises: i) a first polypeptide, arranged from N-to-C terminus, comprising: a variable light chain (VL) amino acid sequence; and a light chain constant region amino acid sequence (CL1) domain; ii) a second polypeptide, arranged from N-to-C terminus, comprising: a variable heavy chain (VH) amino acid sequence, a heavy chain CH1 constant region amino acid sequence, a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence; and iii) a third polypeptide, arranged from N-to-C terminus, comprising: a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence, wherein the CH2 and CH3 domains of each of the second and third polypeptides form an Fc domain.

[0329] In some embodiments, the Fc domain is a human IgG Fc domain. In some embodiments, the Fc domain is an IgG1, IgG2, IgG3, or IgG4 Fc domain. In some embodiments, the Fc domain comprises one or more modifications that promote heterodimerization. In some embodiments, the second polypeptide comprises a knob modification in the CH2 or the CH3 domain, and the third polypeptide comprises a hole modification in the CH2 or the CH3 domain; or wherein the third polypeptide comprises a knob modification in the CH2 or the CH3 domain and the second polypeptide comprises a hole modification in the CH2 or the CH3 domain. In some embodiments, the second polypeptide comprises a knob modification in the CH3 domain, and the third polypeptide comprises a hole modification in the CH3 domain; or wherein the third polypeptide comprises a knob modification in the CH3 domain and the second polypeptide comprises a hole modification in the CH3 domain.

[0330] In some embodiments, the antigen binding protein or functional fragment thereof is a monovalent or bivalent antibody. In some embodiments, the bivalent antibody is monospecific or bispecific. In some embodiments, the antigen binding protein or functional fragment thereof is a humanized antibody or a functional fragment thereof.

[0331] A bispecific antibody can comprise two antigen binding arms, wherein the two antigen binding arms recognize different antigens. In some embodiments, the first or the second antigen binding arm independently binds to an immune checkpoint protein. In some embodiments, the immune checkpoint protein is selected from the group consisting of PD-1, CD28, CTLA-4, ICOS, TMIGD2, 4-1BB, BTLA, CD160, LIGHT, LAG3, OX40, CD27, CD39, CD73, CD40L, GITR, DNAM-1, TIGIT, CD96, 2B4, TIM-3, CEACAM1, SIRPα, DC-SIGN, CD200R, DR3, PD-L1, PD-L2, CD80, CD86, ICOS ligand, B7-H3, B7-H4, VISTA, B7-H7(HHLA2), 4-1BBL, HVEM, OX40L, CD70, CD40, GITRL, CD155, CD48, Galectin-9, Adenosine, IDO, TDO, CD47, BTN2A1, CD200, LAYN, LAIR1, CD276, VTCN1, KIR, A2AR, MHC class I, MHC class II, GALS, TGFR, TREM1, TREM2, HLA-G, CCR4, CCR8, CSF1R, MICA/B, LILRB4, SIGLEC-15, arginase and TL1A. In some embodiments, the first antigen binding arm binds to PD-1. In some embodiments, the first or the second antigen binding arm independently binds to CD8$\boldsymbol{\alpha}$, CD8$\boldsymbol{\alpha\alpha}$, CD8$\boldsymbol{\alpha\beta}$ or CD8$\boldsymbol{\beta}$.

[0332] In some embodiments, the antigen binding molecules of the present disclosure bind to an epitope on CD8a wherein the binding of the antigen binding molecule to CD8a does not block the interaction of CD8aa or CD8ab with MHC class I molecules on target cells or antigen presenting cells. In some embodiments, the antigen binding molecule of the present disclosure binds to an epitope on CD8b wherein the binding of the antigen binding molecule to CD8b does not block the interaction of CD8ab with MHC class I molecules on target cells or antigen presenting cells.

[0333] In some embodiments, the fusion protein binds human CD8, and the binding of the fusion protein to CD8 does not block the interaction of CD8 with MHC class I. In some embodiments, the antigen binding molecule of the present disclosure binds to an epitope on CD8ab wherein the binding of the antigen binding molecule to CD8ab does not block the interaction of CD8aa or CD8ab with MHC class I molecules on target cells or antigen presenting cells. In some embodiments, the antigen binding molecule of the present disclosure binds to an epitope on CD8a wherein the binding of the antigen binding molecule to CD8a does not block the interaction of CD8ab with MHC class I molecules on target cells or antigen presenting cells.

[0334] In some embodiments, the fusion protein binds human CD8, and the binding of the fusion protein to CD8 does not block the interaction of CD8 with MHC class I. In some embodiments, the antigen binding molecule of the present disclosure binds to an epitope on CD8α wherein the binding of the antigen binding molecule to CD8α does not block the interaction of CD8αα or CD8αβ with MHC class I molecules on target cells or antigen presenting cells. In some embodiments, the antigen binding molecule of the present disclosure binds to an epitope on CD8β

wherein the binding of the antigen binding molecule to CD8β does not block the interaction of CD8αβ with MHC class I molecules on target cells or antigen presenting cells. In some embodiments, whether an anti-CD8 antibody or fusion protein of the present disclosure blocks the interaction of CD8 with MHC class I can be assayed, *e.g.*, by assaying activation status of CD8+ T cells (*e.g.*, upon antigen stimulation) in the presence or absence of the anti-CD8 antibody or fusion protein.

[0335] In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of EVQLVESGGGLVQPGRSLKLSCAASGFTFSNYYMAWVRQAPTKGLEWVAYINTGGGTTY YRDSVKGRFTISRDDAKSTLYLQMDSLRSEDTATYYCTTAIGYYFDYWGQGVMVTVSS (SEQ ID NO:99) and a VL domain comprising the sequence of DIQLTQSPASLSASLGETVSIECLASEDIYSYLAWYQQKPGKSPQVLIYAANRLQDGVPSRF SGSGSGTQYSLKISGMQPEDEGDYFCLQGSKFPYTFGAGTKLELK (SEQ ID NO:100). In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of EVKLQESGPSLVQPSQTLSLTCSVSGFSLISDSVHWVRQPPGKGLEWMGGIWADGSTDYNS ALKSRLSISRDTSKSQGFLKMNSLQTDDTAIYFCTSNRESYYFDYWGQGTMVTVSS (SEQ ID NO:101) and a VL domain comprising the sequence of DIQMTQSPASLSASLGDKVTITCQASQNIDKYIAWYQQKPGKAPRQLIHYTSTLVSGTPSRF SGSGSGRDYSFSISSVESEDIASYYCLQYDTLYTFGAGTKLELK (SEQ ID NO:102). In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of EVKLQESGPSLVQPSQTLSLTCSVSGFSLISDSVHWVRQPPGKGLEWMGGIWADGSTDYNS ALKSRLSISRDTSKSQGFLKMNSLQTDDTAIYFCTSARESYYFDYWGQGTMVTVSS (SEQ ID NO:103) and a VL domain comprising the sequence of DIQMTQSPASLSASLGDKVTITCQASQNIDKYIAWYQQKPGKAPRQLIHYTSTLVSGTPSRF SGSGSGRDYSFSISSVESEDIASYYCLQYATLYTFGAGTKLELK (SEQ ID NO:104). In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of EVQLVESGGALVQPGRSLKLSCAASGLTFSDCYMAWVRQTPTKGLEWVSYISSDGGSTYY GDSVKGRFTISRDNAKSTLYLQMNSLRSEDMATYYCACATDLSSYWSFDFWGPGTMVTV SS (SEQ ID NO:105) and a VL domain comprising the sequence of DIQMTQSPSSLPVSLGERVTISCRASQGISNNLNWYQQKPDGTIKPLIYHTSNLQSGVPSRFS GSGSGTDYSLTISSLEPEDFAMYYCQQDATFPLTFGSGTKLEIK (SEQ ID NO:106).

[0336] In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:107 and a VL domain comprising the sequence of SEQ ID NO:108. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:109 and a VL domain comprising the sequence of SEQ ID NO:110. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:111 and a VL domain comprising the sequence of SEQ ID NO:112. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:113 and a VL domain comprising the sequence of SEQ ID NO:114. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:115 and a VL domain comprising the sequence of SEQ ID NO:116. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:117 and a VL domain comprising the sequence of SEQ ID NO:118. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:119 and a VL domain comprising the sequence of SEQ ID NO:120. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:121 and a VL domain comprising the sequence of SEQ ID NO:122. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:123 and a VL domain comprising the sequence of SEQ ID NO:124. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:125 and a VL domain comprising the sequence of SEQ ID NO:126. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:127 and a VL domain comprising the sequence of SEQ ID NO:128. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:129 and a VL domain comprising the sequence of SEQ ID NO:130. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:131 and a VL domain comprising the sequence of SEQ ID NO:132. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:133 and a VL domain comprising the sequence of SEQ ID NO:134. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:135 and a VL

domain comprising the sequence of SEQ ID NO:136. In some embodiments, an anti-CD8 antibody or fusion protein of the present disclosure comprises a VH domain comprising the sequence of SEQ ID NO:258 and a VL domain comprising the sequence of SEQ ID NO:259.

[0337] In some embodiments, the antigen binding molecules (and fusion proteins) of the present disclosure specifically bind human CD8b and/or human CD8ab.

[0338] In some embodiments, the anti-CD8 antibody of the present disclosure is a human antibody or antibody fragment. In some embodiments, the anti-CD8 antibody of the present disclosure is a humanized antibody or antibody fragment.

[0339] In some embodiments, the anti-CD8 antibody of the present disclosure specifically binds human CD8b and/or human CD8ab with at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, at least 60-fold, at least 70-fold, at least 80-fold, at least 90-fold, at least 100-fold, or at least 200-fold higher affinity than its binding to human CD8a and/or human CD8aa, *e.g.*, as expressed on natural killer (NK) cells (*e.g.*, human NK cells). In some embodiments, the anti-CD8 antibody of the present disclosure specifically binds human CD8b and/or human CD8ab with at least 10-fold higher affinity than its binding to human CD8a and/or human CD8aa, *e.g.*, as expressed on natural killer (NK) cells. In some embodiments, the human CD8b and/or human CD8ab are expressed on the surface of a human cell, *e.g.*, a human T cell.

[0340] In some embodiments, the anti-CD8 antibody of the present disclosure specifically binds to a cell expressing a human CD8ab heterodimer on its surface (*e.g.*, a human T cell) with an EC50 that is less than 1000nM. In some embodiments, the anti-CD8 antibody of the present disclosure specifically binds to human CD8+ T cells.

[0341] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:137, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:138, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:139 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:140, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:141, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:142. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized.

[0342] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:143, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:144, and a CDR-H3 comprising the

amino acid sequence of SEQ ID NO:145 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:146, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:147, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:148. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v8 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v8 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized.

[0343] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:252, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:253, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:254 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:255, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:256, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:257. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:258 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:259. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v1 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v1 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized.

[0344] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:149, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:150, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:151 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:154. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:109 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:110. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v2 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v2 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized.

[0345] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:155, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:156, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:157 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:158, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:159, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:160. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:111 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:112. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v3 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v3 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized.

[0346] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:161, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:162, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:163 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:164, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:165, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:166. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:113 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:114. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v4 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v4 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized.

[0347] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:167, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:168, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:169 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:170, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:171, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:172. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:115 and/or the VL domain

comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:116. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v5 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v5 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized.

[0348] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:173, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:174, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:175 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:117 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:118. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v6 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v6 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is human.

[0349] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:179, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:180, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:181 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:182, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:183, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:184. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:119 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:120. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v7 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v7 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is human.

[0350] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of X_1X_2AIS, wherein X_1 is S, K, G, N, R, D, T, or G, and wherein X_2 is Y, L, H, or F (SEQ ID NO:185), a CDR-H2

comprising the amino acid sequence of $X_1X_2X_3PX_4X_5X_6X_7X_8X_9YX_{10}QKFX_{11}G$, wherein X_1 is G or H, X_2 is I or F, X_3 is I, N, or M, X_4 is G, N, H, S, R, I, or A, X_5 is A, N, H, S, T, F, or Y, X_6 is A, D, or G, X_7 is T, E, K, V, Q, or A, X_8 is A or T, X_9 is N or K, X_{10} is A or N, and X_{11} is Q or T (SEQ ID NO:186), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5LFX_6X_7$, wherein X_1 is D or A, X_2 is A, G, E, R, Y, K, N, Q, L, or F, X_3 is A, L, P, or Y, X_4 is I or L, X_5 is R, A, Q, or S, X_6 is A or D, and X_7 is D, E, A, or S (SEQ ID NO:187) and a VL domain comprising a CDR-L1 comprising the amino acid sequence of $X_1X_2SX_3X_4IX_5GX_6LN$, wherein X_1 is R or G, X_2 is A or T, X_3 is Q or E, X_4 is E, N, T, S, A, K, D, G, R, or Q, X_5 is Y or S, and X_6 is A or V (SEQ ID NO:188), a CDR-L2 comprising the amino acid sequence of $GX_1X_2X_3LX_4X_5$, wherein X_1 is A or S, X_2 is T, S, E, Q, or D, X_3 is N, R, A, E, or H, X_4 is Q or A, and X_5 is S or D (SEQ ID NO:189), and a CDR-L3 comprising the amino acid sequence of $QX_1X_2X_3X_4X_5PWT$, wherein X_1 is S, N, D, Q, A, or E, X_2 is T, I, or S, X_3 is Y, L, or F, X_4 is D, G, T, E, Q, A, or Y, and X_5 is A, T, R, S, K, or Y (SEQ ID NO:190). In some embodiments, the VH domain further comprises a FW-1 comprising the sequence QVQLVQSGAEVKKPGSSVKVSCKASGGTFS (SEQ ID NO:191), a FW-2 comprising the sequence WVRQAPGQGLEWMG (SEQ ID NO:192), a FW-3 comprising the sequence RVTITADESTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO:193), and/or a FW-4 comprising the sequence WGQGTLVTVSS (SEQ ID NO:194). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:195), a FW-2 comprising the sequence WYQQKPGKAPKLLIY (SEQ ID NO:196), a FW-3 comprising the sequence GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:197), and/or a FW-4 comprising the sequence FGGGTKVEIK (SEQ ID NO:198).

[0351] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:200, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:201 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:123 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:124. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v9 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v9 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized. In some embodiments, an anti-CD8

antibody of the present disclosure comprises a VH domain comprising a CDR-H1, CDR-H2, and CDR-H3 from the sequence of SEQ ID NO:123 and a VL domain comprising a CDR-L1, CDR-L2, and CDR-L3 from the sequence of SEQ ID NO:124. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence QVQLVQSGAEVKKPGSSVKVSCKASGGTFS (SEQ ID NO:191), a FW-2 comprising the sequence WVRQAPGQGLEWMG (SEQ ID NO:192), a FW-3 comprising the sequence RVTITADESTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO:193), and/or a FW-4 comprising the sequence WGQGTLVTVSS (SEQ ID NO:194). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:195), a FW-2 comprising the sequence WYQQKPGKAPKLLIY (SEQ ID NO:196), a FW-3 comprising the sequence GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:197), and/or a FW-4 comprising the sequence FGGGTKVEIK (SEQ ID NO:198).

[0352] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:129 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:130. In some embodiments, the VH domain comprises the amino acid sequence of SEQ ID NO:129; and the VL domain comprises the amino acid sequence of SEQ ID NO:130. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v12 (xhCD8.1) (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v12 (xhCD8.1) (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1, CDR-H2, and CDR-H3 from the sequence of SEQ ID NO:129 and a VL domain comprising a CDR-L1, CDR-L2, and CDR-L3 from the sequence of SEQ ID NO:130. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence QVQLVQSGAEVKKPGSSVKVSCKASGGTFS (SEQ ID NO:191), a FW-2 comprising the sequence WVRQAPGQGLEWMG (SEQ ID NO:192), a FW-3 comprising the sequence RVTITADESTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO:193), and/or a FW-4 comprising

the sequence WGQGTLVTVSS (SEQ ID NO:194). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:195), a FW-2 comprising the sequence WYQQKPGKAPKLLIY (SEQ ID NO:196), a FW-3 comprising the sequence GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:197), and/or a FW-4 comprising the sequence FGGGTKVEIK (SEQ ID NO:198).

[0353] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:131 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:132. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v13 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v13 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1, CDR-H2, and CDR-H3 from the sequence of SEQ ID NO:131 and a VL domain comprising a CDR-L1, CDR-L2, and CDR-L3 from the sequence of SEQ ID NO:132. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence QVQLVQSGAEVKKPGSSVKVSCKASGGTFS (SEQ ID NO:191), a FW-2 comprising the sequence WVRQAPGQGLEWMG (SEQ ID NO:192), a FW-3 comprising the sequence RVTITADESTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO:193), and/or a FW-4 comprising the sequence WGQGTLVTVSS (SEQ ID NO:194). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:195), a FW-2 comprising the sequence WYQQKPGKAPKLLIY (SEQ ID NO:196), a FW-3 comprising the sequence GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:197), and/or a FW-4 comprising the sequence FGGGTKVEIK (SEQ ID NO:198).

[0354] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of X_1YX_2MS, wherein X_1 is S, D, E, A, or Q and X_2 is A, G, or T (SEQ ID NO:208), a CDR-H2 comprising the amino acid sequence of $DIX_1X_2X_3GX_4X_5TX_6YADSVKG$, wherein X_1 is T, N, S, Q, E, H, R, or A, X_2 is Y, W,

F, or H, X_3 is A, S, Q, E, or T, X_4 is G or E, X_5 is S or I, and X_6 is A or G (SEQ ID NO:209), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3YX_4WX_5X_6AX_7DX_8$, wherein X_1 is S or A, X_2 is N, H, A, D, L, Q, Y, or R, X_3 is A, N, S, or G, X_4 is A, V, R, E, or S, X_5 is D or S, X_6 is D, N, Q, E, S, T, or L, X_7 is L, F, or M, and X_8 is I, Y, or V (SEQ ID NO:210) and a VL domain comprising a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178). In some embodiments, the VH domain further comprises a FW-1 comprising the sequence EVQLVESGGGLVQPGGSLRLSCAASGFTFS (SEQ ID NO:211), a FW-2 comprising the sequence WVRQAPGKGLEWVS (SEQ ID NO:212), a FW-3 comprising the sequence RFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR (SEQ ID NO:213), and/or a FW-4 comprising the sequence WGQGTMVTVSS (SEQ ID NO:214) or WGQGTLVTVSS (SEQ ID NO:215). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence EIVLTQSPGTLSLSPGERATLSC (SEQ ID NO:216), a FW-2 comprising the sequence WYQQKPGQAPRLLIY (SEQ ID NO:217), a FW-3 comprising the sequence GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC (SEQ ID NO:218), and/or a FW-4 comprising the sequence FGQGTKVEIK (SEQ ID NO:219).

[0355] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:125 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:126. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v10 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v10 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1, CDR-H2, and CDR-H3 from the sequence of SEQ ID NO:125 and a VL domain comprising a CDR-L1, CDR-L2, and CDR-L3 from the sequence of SEQ ID NO:126. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence EVQLVESGGGLVQPGGSLRLSCAASGFTFS (SEQ

ID NO:211), a FW-2 comprising the sequence WVRQAPGKGLEWVS (SEQ ID NO:212), a FW-3 comprising the sequence RFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR (SEQ ID NO:213), and/or a FW-4 comprising the sequence WGQGTMVTVSS (SEQ ID NO:214) or WGQGTLVTVSS (SEQ ID NO:215). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence EIVLTQSPGTLSLSPGERATLSC (SEQ ID NO:216), a FW-2 comprising the sequence WYQQKPGQAPRLLIY (SEQ ID NO:217), a FW-3 comprising the sequence GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC (SEQ ID NO:218), and/or a FW-4 comprising the sequence FGQGTKVEIK (SEQ ID NO:219).

[0356] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:127 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:128. In some embodiments, the VH domain comprises the amino acid sequence of SEQ ID NO:127; and the VL domain comprises the amino acid sequence of SEQ ID NO:128. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v11 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v11 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1, CDR-H2, and CDR-H3 from the sequence of SEQ ID NO:127 and a VL domain comprising a CDR-L1, CDR-L2, and CDR-L3 from the sequence of SEQ ID NO:128. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence EVQLVESGGGLVQPGGSLRLSCAASGFTFS (SEQ ID NO:211), a FW-2 comprising the sequence WVRQAPGKGLEWVS (SEQ ID NO:212), a FW-3 comprising the sequence RFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR (SEQ ID NO:213), and/or a FW-4 comprising the sequence WGQGTMVTVSS (SEQ ID NO:214) or WGQGTLVTVSS (SEQ ID NO:215). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence EIVLTQSPGTLSLSPGERATLSC (SEQ ID NO:216), a FW-2 comprising the sequence WYQQKPGQAPRLLIY (SEQ ID NO:217), a FW-3 comprising the

sequence GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC (SEQ ID NO:218), and/or a FW-4 comprising the sequence FGQGTKVEIK (SEQ ID NO:219).

[0357] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:260, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:133 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:134. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v14 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v14 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1, CDR-H2, and CDR-H3 from the sequence of SEQ ID NO:133 and a VL domain comprising a CDR-L1, CDR-L2, and CDR-L3 from the sequence of SEQ ID NO:134. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence EVQLVESGGGLVQPGGSLRLSCAASGFTFS (SEQ ID NO:211), a FW-2 comprising the sequence WVRQAPGKGLEWVS (SEQ ID NO:212), a FW-3 comprising the sequence RFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR (SEQ ID NO:213), and/or a FW-4 comprising the sequence WGQGTMVTVSS (SEQ ID NO:214) or WGQGTLVTVSS (SEQ ID NO:215). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence EIVLTQSPGTLSLSPGERATLSC (SEQ ID NO:216), a FW-2 comprising the sequence WYQQKPGQAPRLLIY (SEQ ID NO:217), a FW-3 comprising the sequence GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC (SEQ ID NO:218), and/or a FW-4 comprising the sequence FGQGTKVEIK (SEQ ID NO:219).

[0358] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:260, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178. In some embodiments, the VH domain comprises an amino acid sequence that is at least 90%, at least 95%,

at least 99%, or 100% identical to the sequence of SEQ ID NO:135 and/or the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:136. In some embodiments, an anti-CD8 antibody of the present disclosure comprises 1, 2, or 3 heavy chain CDRs of antibody xhCD8v15 (*e.g.*, as shown in Tables 10-12) and/or 1, 2, or 3 light chain CDRs of antibody xhCD8v15 (*e.g.*, as shown in Tables 10-12). In some embodiments, the antibody is humanized. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1, CDR-H2, and CDR-H3 from the sequence of SEQ ID NO:135 and a VL domain comprising a CDR-L1, CDR-L2, and CDR-L3 from the sequence of SEQ ID NO:136. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence EVQLVESGGGLVQPGGSLRLSCAASGFTFS (SEQ ID NO:211), a FW-2 comprising the sequence WVRQAPGKGLEWVS (SEQ ID NO:212), a FW-3 comprising the sequence RFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR (SEQ ID NO:213), and/or a FW-4 comprising the sequence WGQGTMVTVSS (SEQ ID NO:214) or WGQGTLVTVSS (SEQ ID NO:215). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence EIVLTQSPGTLSLSPGERATLSC (SEQ ID NO:216), a FW-2 comprising the sequence WYQQKPGQAPRLLIY (SEQ ID NO:217), a FW-3 comprising the sequence GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC (SEQ ID NO:218), and/or a FW-4 comprising the sequence FGQGTKVEIK (SEQ ID NO:219).

[0359] Multiple definitions for the CDR sequences of antibody variable domains are known in the art. Unless otherwise specified, CDR sequences are described herein according to the definition of Kabat (*see, e.g.,* Kabat et al., *Sequences of Proteins of Immunological Interest*, Fifth Edition, NIH Publication 91-3242, Bethesda MD (1991), vols. 1-3). However, other definitions are known and contemplated for use. For example, in some embodiments, CDR sequences can be described by the definition of Chothia (*see, e.g.,* Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987).

[0360] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:224, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:225 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:140, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:141, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:142. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:226, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of

SEQ ID NO:151 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:154. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:228, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:157 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:158, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:159, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:160. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:163 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:164, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:165, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:166. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:229, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:169 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:170, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:171, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:172. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:230, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:231, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:175 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:230, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:232, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:181 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:182, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:183, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:184. In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:233, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:234, and a CDR-

H3 comprising the amino acid sequence of SEQ ID NO:145 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:146, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:147, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:148.

[0361] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of $GX_1X_2FX_3X_4X_5$, wherein X_1 is G, Y, S, or A, X_2 is T, S, G, R, N, or H, X_3 is S, T, R, H, Y, G, or P, X_4 is S, K, G, N, R, D, T, or G, and X_5 is Y, L, H, or F (SEQ ID NO:235), a CDR-H2 comprising the amino acid sequence of $X_1PX_2X_3X_4X_5$, wherein X_1 is I, N, or M, X_2 is G, N, H, S, R, I, or A, X_3 is A, N, H, S, T, F, or Y, X_4 is A, D, or G, and X_5 is T, E, K, V, Q, or A (SEQ ID NO:236), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5LFX_6X_7$, wherein X_1 is D or A, X_2 is A, G, E, R, Y, K, N, Q, L, or F, X_3 is A, L, P, or Y, X_4 is I or L, X_5 is R, A, Q, or S, X_6 is A or D, and X_7 is D, E, A, or S (SEQ ID NO:237) and a VL domain comprising a CDR-L1 comprising the amino acid sequence of $X_1X_2SX_3X_4IX_5GX_6LN$, wherein X_1 is R or G, X_2 is A or T, X_3 is Q or E, X_4 is E, N, T, S, A, K, D, G, R, or Q, X_5 is Y or S, and X_6 is A or V (SEQ ID NO:188), a CDR-L2 comprising the amino acid sequence of $GX_1X_2X_3LX_4X_5$, wherein X_1 is A or S, X_2 is T, S, E, Q, or D, X_3 is N, R, A, E, or H, X_4 is Q or A, and X_5 is S or D (SEQ ID NO:189), and a CDR-L3 comprising the amino acid sequence of $QX_1X_2X_3X_4X_5PWT$, wherein X_1 is S, N, D, Q, A, or E, X_2 is T, I, or S, X_3 is Y, L, or F, X_4 is D, G, T, E, Q, A, or Y, and X_5 is A, T, R, S, K, or Y (SEQ ID NO:190). In some embodiments, the VH domain further comprises a FW-1 comprising the sequence QVQLVQSGAEVKKPGSSVKVSCKAS (SEQ ID NO:238), a FW-2 comprising the sequence AISWVRQAPGQGLEWMGGI (SEQ ID NO:239), a FW-3 comprising the sequence ANYAQKFQGRVTITADESTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO:240), and/or a FW-4 comprising the sequence WGQGTLVTVSS (SEQ ID NO:194). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:195), a FW-2 comprising the sequence WYQQKPGKAPKLLIY (SEQ ID NO:196), a FW-3 comprising the sequence GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:197), and/or a FW-4 comprising the sequence FGGGTKVEIK (SEQ ID NO:198).

[0362] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:242, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202. In some

embodiments, the VH domain further comprises a FW-1 comprising the sequence QVQLVQSGAEVKKPGSSVKVSCKAS (SEQ ID NO:238), a FW-2 comprising the sequence AISWVRQAPGQGLEWMGGI (SEQ ID NO:239), a FW-3 comprising the sequence ANYAQKFQGRVTITADESTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO:240), and/or a FW-4 comprising the sequence WGQGTLVTVSS (SEQ ID NO:194). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:195), a FW-2 comprising the sequence WYQQKPGKAPKLLIY (SEQ ID NO:196), a FW-3 comprising the sequence GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:197), and/or a FW-4 comprising the sequence FGGGTKVEIK (SEQ ID NO:198).

[0363] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence QVQLVQSGAEVKKPGSSVKVSCKAS (SEQ ID NO:238), a FW-2 comprising the sequence AISWVRQAPGQGLEWMGGI (SEQ ID NO:239), a FW-3 comprising the sequence ANYAQKFQGRVTITADESTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO:240), and/or a FW-4 comprising the sequence WGQGTLVTVSS (SEQ ID NO:194). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:195), a FW-2 comprising the sequence WYQQKPGKAPKLLIY (SEQ ID NO:196), a FW-3 comprising the sequence GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:197), and/or a FW-4 comprising the sequence FGGGTKVEIK (SEQ ID NO:198).

[0364] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence QVQLVQSGAEVKKPGSSVKVSCKAS (SEQ ID NO:238), a FW-2 comprising the sequence AISWVRQAPGQGLEWMGGI (SEQ ID NO:239), a FW-3 comprising the sequence ANYAQKFQGRVTITADESTSTAYMELSSLRSEDTAVYYCAR (SEQ ID NO:240), and/or a

FW-4 comprising the sequence WGQGTLVTVSS (SEQ ID NO:194). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:195), a FW-2 comprising the sequence WYQQKPGKAPKLLIY (SEQ ID NO:196), a FW-3 comprising the sequence GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:197), and/or a FW-4 comprising the sequence FGGGTKVEIK (SEQ ID NO:198).

[0365] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of $GFTFX_1X_2Y$, wherein X_1 is S, D, E, Q, S, or A and X_2 is S, D, E, A, or Q (SEQ ID NO:244), a CDR-H2 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5$, wherein X_1 is T, N, S, Q, E, H, R or A, X_2 is Y, W, F, or H, X_3 is A, S, Q, E, or T, X_4 is G or E, and X_5 is S or I (SEQ ID NO:245), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3YX_4WX_5X_6AX_7DX_8$, wherein X_1 is S or A, X_2 is N, H, A, D, L, Q, Y, or R, X_3 is A, N, S, or G, X_4 is A, V, R, E, or S, X_5 is D or S, X_6 is D, N, Q, E, S, T, or L, X_7 is L, F, or M, and X_8 is I, Y, or V (SEQ ID NO:246) and a VL domain comprising a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178). In some embodiments, the VH domain further comprises a FW-1 comprising the sequence EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:247), a FW-2 comprising the sequence AMSWVRQAPGKGLEWVSDI (SEQ ID NO:248), a FW-3 comprising the sequence TAYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR (SEQ ID NO:249), and/or a FW-4 comprising the sequence WGQGTMVTVSS (SEQ ID NO:214) or WGQGTLVTVSS (SEQ ID NO:215). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence EIVLTQSPGTLSLSPGERATLSC (SEQ ID NO:216), a FW-2 comprising the sequence WYQQKPGQAPRLLIY (SEQ ID NO:217), a FW-3 comprising the sequence GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC (SEQ ID NO:218), and/or a FW-4 comprising the sequence FGQGTKVEIK (SEQ ID NO:219).

[0366] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:247), a FW-2 comprising the sequence

AMSWVRQAPGKGLEWVSDI (SEQ ID NO:248), a FW-3 comprising the sequence TAYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR (SEQ ID NO:249), and/or a FW-4 comprising the sequence WGQGTMVTVSS (SEQ ID NO:214) or WGQGTLVTVSS (SEQ ID NO:215). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence EIVLTQSPGTLSLSPGERATLSC (SEQ ID NO:216), a FW-2 comprising the sequence WYQQKPGQAPRLLIY (SEQ ID NO:217), a FW-3 comprising the sequence GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC (SEQ ID NO:218), and/or a FW-4 comprising the sequence FGQGTKVEIK (SEQ ID NO:219).

[0367] In some embodiments, an anti-CD8 antibody of the present disclosure comprises a VH domain comprising a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:261, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288 and a VL domain comprising a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178. In some embodiments, the VH domain further comprises a FW-1 comprising the sequence EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:247), a FW-2 comprising the sequence AMSWVRQAPGKGLEWVSDI (SEQ ID NO:248), a FW-3 comprising the sequence TAYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAR (SEQ ID NO:249), and/or a FW-4 comprising the sequence WGQGTMVTVSS (SEQ ID NO:214) or WGQGTLVTVSS (SEQ ID NO:215). In some embodiments, the VL domain further comprises a FW-1 comprising the sequence EIVLTQSPGTLSLSPGERATLSC (SEQ ID NO:216), a FW-2 comprising the sequence WYQQKPGQAPRLLIY (SEQ ID NO:217), a FW-3 comprising the sequence GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC (SEQ ID NO:218), and/or a FW-4 comprising the sequence FGQGTKVEIK (SEQ ID NO:219). In some embodiments, the present disclosure provides an anti-CD8 antibody comprising a VH domain comprising CDR-H1, CDR-H2, and CDR-H3 sequences of a single antibody listed in Table 10 and a VL domain comprising CDR-L1, CDR-L2, and CDR-L3 sequences of the single antibody listed in Table 10. For example, the anti-CD8 antibody comprises the six CDRs of antibody xhCD8, xhCD8v1, xhCD8v2, xhCD8v3, xhCD8v4, xhCD8v5, xhCD8v6, xhCD8v7, xhCD8v8, xhCD8v9, xhCD8v10, xhCD8v11, xhCD8v12 (xhCD8.1), xhCD8v13, xhCD8v14, xhCD8v15, V9 family, or V11 family shown in Table 10. In some embodiments, the present disclosure provides an anti-CD8 antibody comprising a VH domain comprising CDR-H1, CDR-H2, and CDR-H3 sequences of a single antibody listed in Table 10 and a VL domain comprising CDR-L1, CDR-L2, and CDR-L3 sequences of the single antibody listed in Table 10. For example, the anti-CD8 antibody comprises the six CDRs of

antibody xhCD8, xhCD8v1, xhCD8v2, xhCD8v3, xhCD8v4, xhCD8v5, xhCD8v6, xhCD8v7, xhCD8v8, xhCD8v9, xhCD8v10, xhCD8v11, xhCD8v12 (xhCD8.1), xhCD8v13, xhCD8v14, xhCD8v15, V9 family, or V11 family shown in Table 10. In some embodiments, the present disclosure provides a fusion protein comprising an anti-CD8 antibody comprising a VH domain comprising CDR-H1, CDR-H2, and CDR-H3 sequences of the single antibody listed in Table 10 and a VL domain comprising CDR-L1, CDR-L2, and CDR-L3 sequences of a single antibody listed in Table 10. For example, the anti-CD8 antibody of the fusion protein comprises the six CDRs of antibody xhCD8, xhCD8v1, xhCD8v2, xhCD8v3, xhCD8v4, xhCD8v5, xhCD8v6, xhCD8v7, xhCD8v8, xhCD8v9, xhCD8v10, xhCD8v11, xhCD8v12 (xhCD8.1), xhCD8v13, xhCD8v14, xhCD8v15, V9 family, or V11 family shown in Table 10. In some embodiments, the present disclosure provides a fusion protein comprising an anti-CD8 antibody comprising a VH domain comprising CDR-H1, CDR-H2, and CDR-H3 sequences of a single antibody listed in Table 11 and a VL domain comprising CDR-L1, CDR-L2, and CDR-L3 sequences of the single antibody listed in Table 11. For example, the anti-CD8 antibody of the fusion protein comprises the six CDRs of antibody xhCD8, xhCD8v1, xhCD8v2, xhCD8v3, xhCD8v4, xhCD8v5, xhCD8v6, xhCD8v7, xhCD8v8, xhCD8v9, xhCD8v10, xhCD8v11, xhCD8v12 (xhCD8.1), xhCD8v13, xhCD8v14, xhCD8v15, V9 family, or V11 family shown in Table 11. In some embodiments, the present disclosure provides an anti-CD8 antibody comprising a VH domain comprising CDR-H1, CDR-H2, and CDR-H3 sequences of a VH domain listed in Table 12 and a VL domain comprising CDR-L1, CDR-L2, and CDR-L3 sequences of a VL domain listed in Table 12 (in some embodiments, the VH and VL domains are from the same single antibody listed in Table 12). For example, the anti-CD8 antibody comprises the VH and VL of antibody xhCD8, xhCD8v1, xhCD8v2, xhCD8v3, xhCD8v4, xhCD8v5, xhCD8v6, xhCD8v7, xhCD8v8, xhCD8v9, xhCD8v10, xhCD8v11, xhCD8v12 (xhCD8.1), xhCD8v13, xhCD8v14, or xhCD8v15 shown in Table 12. In some embodiments, the present disclosure provides a fusion protein comprising an anti-CD8 antibody comprising a VH domain comprising CDR-H1, CDR-H2, and CDR-H3 sequences of a VH domain listed in Table 12 and a VL domain comprising CDR-L1, CDR-L2, and CDR-L3 sequences of a VL domain listed in Table 12 (in some embodiments, the VH and VL domains are from the same single antibody listed in Table 12). In some embodiments, the present disclosure provides an anti-CD8 antibody comprising a VH domain sequence and a VL domain sequence for a single antibody as listed in Table 12. In some embodiments, the present disclosure provides a fusion protein comprising an anti-CD8 antibody comprising a VH domain sequence and a VL domain sequence for a single antibody as listed in Table 12. For example, the anti-CD8 antibody of the fusion protein comprises the VH and VL of antibody xhCD8, xhCD8v1, xhCD8v2,

xhCD8v3, xhCD8v4, xhCD8v5, xhCD8v6, xhCD8v7, xhCD8v8, xhCD8v9, xhCD8v10, xhCD8v11, xhCD8v12 (xhCD8.1), xhCD8v13, xhCD8v14, or xhCD8v15 shown in Table 12.

Linkers

[0368] In some embodiments, the targeted cytokine construct comprises a linker between the antigen binding protein or a functional fragment thereof (*e.g.*, an antibody or an antigen binding fragment thereof as described above) and the IL-21 polypeptide or a functional fragment or a variant thereof. Many different linker polypeptides are known in the art and may be used in the context of an targeted cytokine construct. In some embodiments, the targeted cytokine construct comprises one or more copies of a peptide comprising the sequence of: GGGGS (SEQ ID NO:23), GGNGT (SEQ ID NO: 24), or YGNGT (SEQ ID NO: 25) between the antigen binding protein and the IL-21 polypeptide or a functional fragment or a variant thereof. In some embodiments, the polypeptide region between the antigen binding protein and the IL-21 polypeptide or a functional fragment or a variant thereof comprises a single copy of GGGGS (SEQ ID NO: 23), GGNGT (SEQ ID NO: 24), or YGNGT (SEQ ID NO: 25).

Targeted Cytokine Protein

[0369] The present disclosure provides in some embodiments a fusion protein comprising an antigen binding protein or a functional fragment and an IL-21 polypeptide or a functional fragment or a variant thereof. In some embodiments, the fusion protein can comprise the antigen binding protein or a functional fragment that is operably linked to an IL-21 polypeptide or a functional fragment or a variant thereof. In some embodiments, the antigen binding protein or functional fragment thereof binds a receptor on a T cell. In some embodiments, the T cell is a CD8+ T cell. In some embodiments, the T cell is not a Treg cell (*e.g.,* T cell expressing FoxP3). In some embodiments, the antigen binding protein or functional fragment thereof is an anti-CD8 antibody or antigen binding fragment thereof as described herein.

[0370] In some embodiments, the targeted cytokine construct comprises an antibody or antigen binding fragment having the structure: i) a first polypeptide, arranged from N-to-C terminus, comprising: a variable light chain (VL) amino acid sequence, and a light chain constant region amino acid sequence (CL1); ii) a second polypeptide, arranged from N-to-C terminus, comprising: a variable heavy chain (VH) amino acid sequence, a heavy chain CH1 constant region amino acid sequence, a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence; iii) a third polypeptide, arranged from N-to-C terminus, comprising : a hinge region amino acid sequence, a heavy chain

CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence, wherein the CH2 and CH3 domains of each of the second and third polypeptides form an Fc domain.

[0371] In some embodiments, iii) the third polypeptide comprises, arranged from N-to-C terminus, comprising: a variable heavy chain (VH) amino acid sequence, a heavy chain CH1 constant region amino acid sequence, a hinge region amino acid sequence, a heavy chain CH2 constant region amino acid sequence, and a heavy chain CH3 constant region amino acid sequence; wherein the antibody or antigen binding fragment thereof further comprises: iv) a fourth polypeptide, arranged from N-to-C terminus, comprising: a variable light chain (VL) amino acid sequence, and a light chain constant amino acid sequence (CL1).

[0372] In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof and the antibody or antigen binding fragment thereof are operably linked to each other. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof is linked to the N-terminus or C-terminus of the antibody or antigen binding fragment thereof. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof is conjugated to the C-terminus of the first, second, or third polypeptide. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof is conjugated to the N-terminus of the first, second, or third polypeptide. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof is conjugated to the C-terminus of the first, second, third, or fourth polypeptide. In some embodiments, the IL-21 polypeptide or a functional fragment or a variant thereof is conjugated to the hinge region of the second or third polypeptide. In some embodiments, the targeted cytokine construct comprises at least one molecule of the IL-21 polypeptide or a functional fragment or a variant thereof. In some embodiments, the targeted cytokine construct comprises at least 1, 2, 3, 4, or 5 molecules of the IL-21 polypeptide, such from 1 to 2, 1 to 3, 1 to 4, 1 to 5, 2 to 3, 2 to 4, 2 to 5, 3 to 4, 3 to 5, or 4 to 5 molecules of the IL-21 polypeptide or a functional fragment or a variant thereof.

[0373] In some embodiments, the Fc domain comprises a human IgG Fc domain, such as IgG1, IgG2, IgG3, or IgG4 Fc domain. In some embodiments, the Fc domain comprises one or more modifications that promote heterodimerization. In some embodiments, the second polypeptide comprises a knob modification in the CH2 or the CH3 domain, and the third polypeptide comprises a hole modification in the CH2 or the CH3 domain; or wherein the third polypeptide comprises a knob modification in the CH2 or the CH3 domain and the second polypeptide comprises a hole modification in the CH2 or the CH3. For example, an IgG Fc domain of the targeted cytokine construct can comprise a knob-into-hole modification in the CH3 domains. The “hole” heavy chain

may be connected to an IL-21 polypeptide and carry the S354C, T366S, L368A and Y407V mutations in the CH3 domain, whereas the unfused "knob" heavy chain can carry the Y349C and T366W mutations in the CH3 domain (EU numbering). Alternately, the "hole" heavy chain may be unfused and carry the S354C, T366S, L368A and Y407V mutations in the CH3 domain, whereas the "knob" heavy chain may be connected to the IL-21 polypeptide and can carry the Y349C and T366W mutations in the CH3 domain (EU numbering). In some embodiments, at least one of said first and second Fc domains comprise the following Fc mutations according to EU numbering: L234A, L235A, and G237A.

[0374] In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof is conjugated to at least one of: the N-terminus and the C-terminus of the second or the third polypeptide. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof is conjugated to the C-terminus of the second or the third polypeptide. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof is conjugated to at least one of: the N-terminus or the C-terminus of the second or the third polypeptide. In some embodiments, an IL-21 polypeptide or a functional fragment or a variant thereof is conjugated to the N-terminus or the C-terminus of first or the fourth polypeptide. In some embodiments, an IL-21 polypeptide is conjugated to at least one of: the hinge region of the second or the third polypeptide.

[0375] In some embodiments, a fusion protein of the present disclosure displays one or more of the following: binds human CD8 and does not block an interaction of CD8 with MHC class I; and activates CD8+ T cells with at least 10-fold, 25-fold, 50-fold, 100-fold, 250-fold, 500-fold, or 1000-fold greater potency, *e.g.*, as compared to activation of NK cells, CD4+ T cells or B cells. In some embodiments, whether an anti-CD8 antibody or fusion protein of the present disclosure blocks the interaction of CD8 with MHC class I can be assayed, *e.g.*, by assaying activation status of CD8+ T cells (*e.g.*, upon antigen stimulation) in the presence or absence of the anti-CD8 antibody or fusion protein. In some embodiments, activation of CD8+ T cells, CD4+ T cells, and/or NK cells can be measured, *e.g.*, by assaying STAT3 phosphorylation. For an exemplary assay and conditions, *see, e.g.*, **Example 3.**

[0376] A targeted cytokine construct comprising and IL-21 polypeptide or a functional fragment or a variant thereof of the disclosure, in certain embodiments, the IL-21 polypeptide binds to IL-21R with K_D that is greater than about 1 nM. In some embodiments, the IL-21 polypeptide binds to IL-21R with K_D that is greater than about 10 nM. In some embodiments, the IL-21 polypeptide binds to IL-21R with K_D that is greater than about 100 nM. In some embodiments, the IL-21 polypeptide binds to IL-21R with K_D that is greater than about 1μM. In some embodiments, the IL-21 polypeptide binds to IL-21R with K_D that is greater than about 10μM. In some embodiments, the

IL-21 polypeptide binds to IL-21R with K_D that is greater than about 100μM. In some embodiments, the IL-21 polypeptide binds to In some embodiments the IL-21 polypeptide or functional fragment thereof has a K_D of about 1 nM, about 5 nM, about 10 nM, about 20 nM, about 25 nM, about 50 nM, about 100 nM, about 200 nM, about 300 nM, about 400 nM, about 500 nM, about 600 nM, about 700 nM, about 800 nM, about 900 nM, about 1.0 μM, about 1.5 μM, about 2.0 μM, about 2.5 μM, about 3.0 μM, about 3.5 μM, about 4.0 μM, about 4.5 μM, about 5.0 μM about 10.0 μM, about 20.0 μM, about 30.0 μM, about 40.0 μM, about 50.0 μM, about 100 μM or about 1000 μM . In some embodiments, the IL-21 polypeptide has a K_D of about 1 nM to about 5 nM, about 1 nM to about 10 nM, about 1 nM to about 20 nM, about 1 nM to about 25 nM, about 1 nM to about 50 nM, about 1 nM to about 100 nM, about 1 nM to about 200 nM, about 1 nM to about 300 nM, about 1 nM to about 400 nM, about 1 nM to about 500 nM, about 1 nM to about 600 nM, about 1 nM to about 700 nM, about 1 nM to about 800 nM, about 1 nM to about 900 nM, about 1 nM to about 1.0 μM, about 1 nM to about 1.5 μM, about 1 nM to about 2.0 μM, about 1 nM to about 2.5 μM, about 1 nM to about 3.0 μM, about 1 nM to about 3.5 μM, about 1 nM to about 4.0 μM, about 1 nM to about 4.5 μM, about 1 nM to about 5.0 μM, about 5 nM to about 10 nM, about 5 nM to about 20 nM, about 5 nM to about 25 nM, about 5 nM to about 50 nM, about 5 nM to about 100 nM, about 5 nM to about 200 nM, about 5 nM to about 300 nM, about 5 nM to about 400 nM, about 5 nM to about 500 nM, about 5 nM to about 600 nM, about 5 nM to about 700 nM, about 5 nM to about 800 nM, about 5 nM to about 900 nM, about 5 nM to about 1.0 μM, about 5 nM to about 1.5 μM, about 5 nM to about 2.0 μM, about 5 nM to about 2.5 μM, about 5 nM to about 3.0 μM, about 5 nM to about 3.5 μM, about 5 nM to about 4.0 μM, about 5 nM to about 4.5 μM, about 5 nM to about 5.0 μM, 10 nM to about 20 nM, about 10 nM to about 25 nM, about 10 nM to about 50 nM, about 10 nM to about 100 nM, about 10 nM to about 200 nM, about 10 nM to about 300 nM, about 10 nM to about 400 nM, about 10 nM to about 500 nM, about 10 nM to about 600 nM, about 10 nM to about 700 nM, about 10 nM to about 800 nM, about 10 nM to about 900 nM, about 10 nM to about 1.0 μM, about 10 nM to about 1.5 μM, about 10 nM to about 2.0 μM, about 10 nM to about 2.5 μM, about 10 nM to about 3.0 μM, about 10 nM to about 3.5 μM, about 10 nM to about 4.0 μM, about 10 nM to about 4.5 μM, about 10 nM to about 5.0 μM, about 20 nM to about 25 nM, about 20 nM to about 50 nM, about 20 nM to about 100 nM, about 20 nM to about 200 nM, about 20 nM to about 300 nM, about 20 nM to about 400 nM, about 20 nM to about 500 nM, about 20 nM to about 600 nM, about 20 nM to about 700 nM, about 20 nM to about 800 nM, about 20 nM to about 900 nM, about 20 nM to about 1.0 μM, about 20 nM to about 1.5 μM, about 20 nM to about 2.0 μM, about 20 nM to about 2.5 μM, about 20 nM to about 3.0 μM, about 20 nM to about 3.5 μM, about 20 nM to about 4.0 μM, about 20 nM to about 4.5 μM,

about 20 nM to about 5.0 μM, about 25 nM to about 50 nM, about 25 nM to about 100 nM, about 25 nM to about 200 nM, about 25 nM to about 300 nM, about 25 nM to about 400 nM, about 25 nM to about 500 nM, about 25 nM to about 600 nM, about 25 nM to about 700 nM, about 25 nM to about 800 nM, about 25 nM to about 900 nM, about 25 nM to about 1.0 μM, about 25 nM to about 1.5 μM, about 25 nM to about 2.0 μM, about 25 nM to about 2.5 μM, about 25 nM to about 3.0 μM, about 25 nM to about 3.5 μM, about 25 nM to about 4.0 μM, about 25 nM to about 4.5 μM, about 25 nM to about 5.0 μM, about 50 nM to about 100 nM, about 50 nM to about 200 nM, about 50 nM to about 300 nM, about 50 nM to about 400 nM, about 50 nM to about 500 nM, about 50 nM to about 600 nM, about 50 nM to about 700 nM, about 50 nM to about 800 nM, about 50 nM to about 900 nM, about 50 nM to about 1.0 μM, about 50 nM to about 1.5 μM, about 50 nM to about 2.0 μM, about 50 nM to about 2.5 μM, about 50 nM to about 3.0 μM, about 50 nM to about 3.5 μM, about 50 nM to about 4.0 μM, about 50 nM to about 4.5 μM, about 50 nM to about 5.0 μM, about 100 nM to about 200 nM, about 100 nM to about 300 nM, about 100 nM to about 400 nM, about 100 nM to about 500 nM, about 100 nM to about 600 nM, about 100 nM to about 700 nM, about 100 nM to about 800 nM, about 100 nM to about 900 nM, about 100 nM to about 1.0 μM, about 100 nM to about 1.5 μM, about 100 nM to about 2.0 μM, about 100 nM to about 2.5 μM, about 100 nM to about 3.0 μM, about 100 nM to about 3.5 μM, about 100 nM to about 4.0 μM, about 100 nM to about 4.5 μM, about 100 nM to about 5.0 μM, about 200 nM to about 300 nM, about 200 nM to about 400 nM, about 200 nM to about 500 nM, about 200 nM to about 600 nM, about 200 nM to about 700 nM, about 200 nM to about 800 nM, about 200 nM to about 900 nM, about 200 nM to about 1.0 μM, about 200 nM to about 1.5 μM, about 200 nM to about 2.0 μM, about 200 nM to about 2.5 μM, about 200 nM to about 3.0 μM, about 200 nM to about 3.5 μM, about 200 nM to about 4.0 μM, about 200 nM to about 4.5 μM, about 200 nM to about 5.0 μM, about 300 nM to about 400 nM, about 300 nM to about 500 nM, about 300 nM to about 600 nM, about 300 nM to about 700 nM, about 300 nM to about 800 nM, about 300 nM to about 900 nM, about 300 nM to about 1.0 μM, about 300 nM to about 1.5 μM, about 300 nM to about 2.0 μM, about 300 nM to about 2.5 μM, about 300 nM to about 3.0 μM, about 300 nM to about 3.5 μM, about 300 nM to about 4.0 μM, about 300 nM to about 4.5 μM, about 300 nM to about 5.0 μM, about 400 nM to about 500 nM, about 400 nM to about 600 nM, about 400 nM to about 700 nM, about 400 nM to about 800 nM, about 400 nM to about 900 nM, about 400 nM to about 1.0 μM, about 400 nM to about 1.5 μM, about 400 nM to about 2.0 μM, about 400 nM to about 2.5 μM, about 400 nM to about 3.0 μM, about 400 nM to about 3.5 μM, about 400 nM to about 4.0 μM, about 400 nM to about 4.5 μM, about 400 nM to about 5.0 μM , about 500 nM to about 600 nM, about 500 nM to about 700 nM, about 500 nM to about 800 nM, about 500 nM to about 900 nM,

about 500 nM to about 1.0 μM, about 500 nM to about 1.5 μM, about 500 nM to about 2.0 μM, about 500 nM to about 2.5 μM, about 500 nM to about 3.0 μM, about 500 nM to about 3.5 μM, about 500 nM to about 4.0 μM, about 500 nM to about 4.5 μM, about 500 nM to about 5.0 μM, about 600 nM to about 700 nM, about 600 nM to about 800 nM, about 600 nM to about 900 nM, about 600 nM to about 1.0 μM, about 600 nM to about 1.5 μM, about 600 nM to about 2.0 μM, about 600 nM to about 2.5 μM, about 600 nM to about 3.0 μM, about 600 nM to about 3.5 μM, about 600 nM to about 4.0 μM, about 600 nM to about 4.5 μM, about 600 nM to about 5.0 μM, about 700 nM to about 800 nM, about 700 nM to about 900 nM, about 700 nM to about 1.0 μM, about 700 nM to about 1.5 μM, about 700 nM to about 2.0 μM, about 700 nM to about 2.5 μM, about 700 nM to about 3.0 μM, about 700 nM to about 3.5 μM, about 700 nM to about 4.0 μM, about 700 nM to about 4.5 μM, about 700 nM to about 5.0 μM, about 800 nM to about 900 nM, about 800 nM to about 1.0 μM, about 800 nM to about 1.5 μM, about 800 nM to about 2.0 μM, about 800 nM to about 2.5 μM, about 800 nM to about 3.0 μM, about 800 nM to about 3.5 μM, about 800 nM to about 4.0 μM, about 800 nM to about 4.5 μM, about 800 nM to about 5.0 μM, about 900 nM to about 1.0 μM, about 900 nM to about 1.5 μM, about 900 nM to about 2.0 μM, about 900 nM to about 2.5 μM, about 900 nM to about 3.0 μM, about 900 nM to about 3.5 μM, about 900 nM to about 4.0 μM, about 900 nM to about 4.5 μM, about 900 nM to about 5.0 μM, about 1.0 μM to about 1.5 μM, about 1.0 μM to about 2.0 μM, about 1.0 μM to about 2.5 μM, about 1.0 μM to about 3.0 μM, about 1.0 μM to about 3.5 μM, about 1.0 μM to about 4.0 μM, about 1.0 μM to about 4.5 μM, about 1.0 μM to about 5.0 μM, about 1.5 μM to about 2.0 μM, about 1.5 μM to about 2.5 μM, about 1.5 μM to about 3.0 μM, about 1.5 μM to about 3.5 μM, about 1.5 μM to about 4.0 μM, about 1.5 μM to about 4.5 μM, about 1.5 μM to about 5.0 μM, about 2.0 μM to about 2.5 μM, about 2.0 μM to about 3.0 μM, about 2.0 μM to about 3.5 μM, about 2.0 μM to about 4.0 μM, about 2.0 μM to about 4.5 μM, about 2.0 μM to about 5.0 μM, about 2.5 μM to about 3.0 μM, about 2.5 μM to about 3.5 μM, about 2.5 μM to about 4.0 μM, about 2.5 μM to about 4.5 μM, about 2.5 μM to about 5.0 μM, about 3.0 μM to about 3.5 μM, about 3.0 μM to about 4.0 μM, about 3.0 μM to about 4.5 μM, about 3.0 μM to about 5.0 μM, about 3.5 μM to about 4.0 μM, about 3.5 μM to about 4.5 μM, about 3.5 μM to about 5.0 μM, about 4.0 μM to about 4.5 μM, about 4.0 μM to about 5.0 μM, about 4.5 μM to about 5.0 μM, about 10 μM to about 100 μM, or about 100 μM to about 1000 μM. In some embodiments, the K_D is determined by surface plasmon resonance or using a BIAcore instrument.

[0377] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:262, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:263, a third polypeptide chain comprises

the amino acid sequence of SEQ ID NO:264, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:262.

[0378] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:266, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:267, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:268, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:266.

[0379] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:270, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:271, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:272, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 270.

[0380] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:275, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:276, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274.

[0381] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:278, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:279, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:280, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 278.

[0382] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:262, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:263, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:265, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:262.

[0383] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:266, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:267, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:269, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:266.

[0384] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:270, a second polypeptide

chain comprises the amino acid sequence of SEQ ID NO:271, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:273, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 270.

[0385] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:275, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:277, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274.

[0386] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:278, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:279, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:281, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 278.

[0387] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:297, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:298, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:299, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:297.

[0388] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:301, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:302, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:303 and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:301.

[0389] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:305, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:306, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:307, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 305.

[0390] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:310, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:311, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309.

[0391] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:297, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:298, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:300, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:297.

[0392] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:301, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:302, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:304 and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:301.

[0393] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO:305, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:306, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:308, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 305.

[0394] In some embodiments, the disclosure provides a fusion protein comprising: a first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309, a second polypeptide chain comprises the amino acid sequence of SEQ ID NO:310, a third polypeptide chain comprises the amino acid sequence of SEQ ID NO:312, and a fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309.

[0395] Also provided, in some embodiments, are at least one polynucleotide encoding a targeted cytokine construct, antibody or functional fragment thereof, or a cytokine or functional fragment thereof. In some embodiments, the polynucleotide molecules are provided as a DNA construct. In other embodiments, the polynucleotide molecules are provided as a messenger RNA transcript.

[0396] In some embodiments are provided nucleic acid sequences that are codon optimized for expression in a host cell, *e.g.*, a bacterium, such as *E. coli*, or a eukaryotic cell, such as a CHO cell. In some examples, the nucleic acid sequences are codon optimized for expression in CHO cells.

[0397] The polynucleotide molecules are constructed by known methods such as by incorporating the genes encoding the binding proteins into a genetic construct linked to a suitable promoter, and optionally a suitable transcription terminator, and expressing it in bacteria or other appropriate expression system such as, for example CHO cells. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. The promoter is selected such that it drives the expression of the polynucleotide in the respective host cell.

[0398] In some embodiments, a polynucleotide as described herein is inserted into a vector, preferably an expression vector, which represents a further embodiment. This recombinant vector can be constructed according to known methods. Vectors of particular interest include plasmids, phagemids, phage derivatives, virii (*e.g.*, retroviruses, adenoviruses, adeno-associated viruses, herpes viruses, lentiviruses, and the like), and cosmids.

[0399] A variety of expression vector/host systems may be utilized to contain and express the polynucleotide encoding the polypeptide of the described herein. Examples of expression vectors for expression in E.coli are pSKK (Le Gall *et al.*, J Immunol Methods. (2004) 285(1):111-27) or pcDNA5 (Invitrogen) for expression in mammalian cells.

[0400] Thus, the targeted cytokine constructs as described herein, in some embodiments, are produced by introducing at least one vector encoding the protein as described above into a host cell and culturing said host cell under conditions whereby the protein domains are expressed, may be isolated and, optionally, further purified.

Method of Use

[0401] The disclosure further provides a method for selectively activating CD8+ T cells over other cells present in PBMC populations. These other cells can include, but are not limited to, CD4+, NK cells, B cells, dendritic cells, and innate lymphoid cells. In certain embodiments, the method of the disclosure can also be used to for the treatment of cancers, infectious diseases, chronic infections.

[0402] In some embodiments the disease or disorder can be selected for the group consisting of an autoimmune disease, inflammation, autoimmune disease, atopic diseases, paraneoplastic autoimmune diseases, cartilage inflammation, arthritis, rheumatoid arthritis, juvenile arthritis, juvenile rheumatoid arthritis, pauciarticular juvenile rheumatoid arthritis, polyarticular juvenile rheumatoid arthritis, systemic onset juvenile rheumatoid arthritis, juvenile ankylosing spondylitis, juvenile enteropathic arthritis, juvenile reactive arthritis, juvenile Reiter's Syndrome, SEA Syndrome (Seronegativity, Enthesopathy, Arthropathy Syndrome), juvenile dermatomyositis, juvenile psoriatic arthritis, juvenile scleroderma, juvenile systemic lupus erythematosus, juvenile vasculitis, pauciarticular rheumatoid arthritis, polyarticular rheumatoid arthritis, systemic onset rheumatoid arthritis, ankylosing spondylitis, enteropathic arthritis, reactive arthritis, Reiter's Syndrome, SEA Syndrome (Seronegativity, Enthesopathy, Arthropathy Syndrome), dermatomyositis, psoriatic arthritis, scleroderma, vasculitis, myelitis, polymyositis , dermatomyositis, polyarteritis nodosa, Wegener's granulomatosis, arteritis, esophagitis rheumatica, sarcoidosis, sclerosis, primary biliary sclerosis, sclerosing cholangitis, Sjogren's syndrome, psoriasis, plaque psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis, erythrodermic

psoriasis, dermatitis, atopic dermatitis, atherosclerosis, lupus, Still's disease, Systemic Lupus Erythematosus (SLE), myasthenia gravis, inflammatory bowel disease (IBD), Crohn's disease, ulcerative colitis, celiac disease, multiple sclerosis (MS), asthma, COPD, rhinosinusitis, rhinosinusitis with polyps, eosinophilic esophagitis, eosinophilic bronchitis, Guillain-Barre disease, Type I diabetes mellitus, thyroiditis (e.g., Graves' disease), Addison's disease, Raynaud's phenomenon, autoimmune hepatitis, GVHD, transplantation rejection, kidney damage, hepatitis C-induced vasculitis, or spontaneous loss of pregnancy.

[0403] In some embodiments, the disease is the cancer. In some embodiments, the cancer is acute lymphoblastic leukemia (ALL) (including non T cell ALL), acute myeloid leukemia, B cell prolymphocytic leukemia, B cell acute lymphoid leukemia ("BALL"), blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloid leukemia, chronic or acute leukemia, diffuse large B cell lymphoma (DLBCL), follicular lymphoma (FL), hairy cell leukemia, Hodgkin's Disease, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, monoclonal gammopathy of undetermined significance (MGUS), multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin's lymphoma (NHL), plasma cell proliferative disorder (including asymptomatic myeloma, smoldering multiple myeloma or indolent myeloma), plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, plasmacytomas (including plasma cell dyscrasia; solitary myeloma; solitary plasmacytoma; extramedullary plasmacytoma; and multiple plasmacytoma), POEMS syndrome (also known as Crow-Fukase syndrome; Takatsuki disease; and PEP syndrome), primary mediastinal large B cell lymphoma (PMBC), small cell- or a large cell-follicular lymphoma, splenic marginal zone lymphoma (SMZL), systemic amyloid light chain amyloidosis, T cell acute lymphoid leukemia ("TALL"), T cell lymphoma, transformed follicular lymphoma, or Waldenstrom macroglobulinemia, Mantle cell lymphoma (MCL), Transformed follicular lymphoma (TFL), Primary mediastinal B cell lymphoma (PMBCL), Multiple myeloma, Hairy cell lymphoma/leukemia, lung cancer, small-cell lung cancer, non-small cell lung (NSCL) cancer, bronchioloalveolar cell lung cancer, squamous cell cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, head and neck cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, thyroid cancer, uterine cancer, gastrointestinal cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, endometrial carcinoma, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the cervix, carcinoma of the vagina, vulval cancer, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine

system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, bladder cancer, liver cancer, hepatoma, hepatocellular cancer, cervical cancer, salivary gland carcinoma, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytoma, schwannomas, ependymomas, medulloblastomas, meningiomas, squamous cell carcinomas, pituitary adenoma and Ewing's sarcoma, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers.

[0404] In some embodiments, the disclosure provides a method of enhancing and/or restoring T cell activity in an aging subject, comprising administering to the subject a composition described herein. In some embodiments, the disclosure provides a method of enhancing efficacy of a vaccine in an aging subject, comprising administering to the subject a composition described herein prior to receiving the vaccine. In some embodiments, the disclosure provides a method of treating a disease, disorder, or condition, associated with aging of immune cells, comprising administering to the subject a composition described herein.

[0405] The targeted cytokine of the present disclosure provides selective activation of T cells, *e.g.*, CD8+ T cells, over other non-CD8+ cell types, such as CD4+ T cells or NK cells. IL-21 generally acts on a broad class of cells, such as regulatory T cells. Stimulation and subsequent activation of cells via IL-21 can be measured by phosphorylation of STAT3. Cellular activation by IL-21 can be reported as an EC50 of STAT3 phosphorylation (pSTAT3). In some embodiments, the contacting results in a selective increase in cytotoxic function of CD8+ T cells in the culture of cells, as compared to the cytotoxic function of the CD8+ T cells, upon contacting the culture of cells with a reference construct that comprises the same IL-21 polypeptide or a functional fragment or variant thereof but does not comprise the antigen binding protein or functional fragment thereof.

[0406] In some embodiments, potency of activation of CD8+ T cell and other cell types is measured by EC50 of cell activation as assessed by STAT3 phosphorylation. In some embodiments, the targeted cytokine construct has an EC50 of STAT3 phosphorylation for CD8+ T cells lower than other cell types. In some embodiments, the other cells are non-CD8+ cells. The non-CD8+ cells can be, but are not limited to B cells, CD4+ T cells, or NK cells. In some embodiments, the selective activation of CD8+ T cells is measured by an increase in STAT3 phosphorylation of the CD8+ T cells compared to the STAT3 phosphorylation of non-CD8+ cells in the culture of cells. In some embodiments, the increase in STAT3 phosphorylation is at least 5X, at least10X or at least 100x more for the CD8+ T cells compared to the STAT3 phosphorylation of

non-CD8+ cells in the culture of cells. In some embodiments, the targeted cytokine construct has a lower EC50 of STAT3 phosphorylation for CD8+ T cells than other cell types by at least 5x, at least 10x, at least 50x, at least 100x, at least 1,000x, at least 10,000x, or at least 100,000x.

[0407] In some embodiments, the targeted cytokine construct activates CD8+ T cells with at least about 10-fold or greater potency as compared to activation of NK cells. In some embodiments, the construct activates CD8+ T cells a potency that is at least about 10-fold to about 50-fold greater as compared to activation of NK cells. In some embodiments, the construct activates CD8+ T cells a potency that is at least about 10-fold to about 15-fold, 15-fold to about 20-fold, 20-fold to about 25-fold, 25-fold to about 30-fold, 30-fold to about 35-fold, 35-fold to about 40-fold, 40-fold to about 45-fold, 45-fold to about 50-fold, 50-fold to about 100-fold, 100-fold to about 1,000-fold, 1,000-fold to about 10,000-fold, 10,000-fold to about 100,000-fold, about 10-fold, 15-fold, 20-fold, 25-fold, 30-fold, 35-fold, 40-fold, 45-fold, 50-fold, 100-fold, 1,000-fold, 10,000-fold or about 100,000-fold greater as compared to activation of NK cells.

[0408] In some embodiments, the targeted cytokine construct activates CD8+ T cells with at least about 10-fold or greater potency as compared to activation of CD4+ T cells. In some embodiments, the construct activates CD8+ T cells a potency that is at least about 10-fold to about 50-fold greater as compared to activation of CD4+ cells. In some embodiments, the construct activates CD4+ T cells a potency that is at least about 10-fold to about 15-fold, 15-fold to about 20-fold, 20-fold to about 25-fold, 25-fold to about 30-fold, 30-fold to about 35-fold, 35-fold to about 40-fold, 40-fold to about 45-fold, 45-fold to about 50-fold, 50-fold to about 100-fold, 100-fold to about 1,000-fold, 1,000-fold to about 10,000-fold, 10,000-fold to about 100,000-fold, about 10-fold, 15-fold, 20-fold, 25-fold, 30-fold, 35-fold, 40-fold, 45-fold, 50-fold, 100-fold, 1,000-fold, 10,000-fold or about 100,000-fold greater as compared to activation of CD4+ T cells.

Host Cells

[0409] Provided herein are host cells comprising a nucleic acid or vector of the present disclosure. As used herein, the term "host cell" refers to any type of cell that can contain the presently disclosed vector and is capable of producing an expression product encoded by the nucleic acid (e.g., mRNA, protein). The host cell in some aspects is an adherent cell or a suspended cell, i.e., a cell that grows in suspension. The host cell in exemplary aspects is a cultured cell or a primary cell which is isolated directly from an organism, *e.g.*, a human. The host cell can be of any cell type, can originate from any type of tissue, and can be of any developmental stage.

[0410] In exemplary aspects, the cell is a eukaryotic cell, including, but not limited to, a yeast cell, filamentous fungi cell, protozoa cell, algae cell, insect cell, or mammalian cell. Such host cells are

described in the art. *See, e.g.*, Frenzel, et al., Front Immunol 4: 217 (2013). In exemplary aspects, the eukaryotic cells are mammalian cells. In exemplary aspects, the mammalian cells are non-human mammalian cells. In some aspects, the cells are Chinese Hamster Ovary (CHO) cells and derivatives thereof (e.g., CHO-K1, CHO pro-3, CS9), mouse myeloma cells (e.g., NS0, GS-NS0, Sp2/0), cells engineered to be deficient in dihydrofolate reductase (DHFR) activity (e.g., DUKX-X11, DG44), human embryonic kidney 293 (HEK293) cells or derivatives thereof (e.g., HEK293T, HEK293-EBNA), green African monkey kidney cells (e.g., COS cells, VERO cells), human cervical cancer cells (e.g., HeLa), human bone osteosarcoma epithelial cells U2-OS, adenocarcinomic human alveolar basal epithelial cells A549, human fibrosarcoma cells HT1080, mouse brain tumor cells CAD, embryonic carcinoma cells P19, mouse embryo fibroblast cells NIH 3T3, mouse fibroblast cells L929, mouse neuroblastoma cells N2a, human breast cancer cells MCF-7, retinoblastoma cells Y79, human retinoblastoma cells SO-Rb50, human liver cancer cells Hep G2, mouse B myeloma cells J558L, or baby hamster kidney (BHK) cells (Gaillet et al. 2007; Khan, Adv Pharm Bull 3(2): 257-263 (2013)). In a particular embodiment, the host cell is CS9 (a CHO cell line).

[0411] For purposes of amplifying or replicating the vector, the host cell is in some aspects is a prokaryotic cell, e.g., a bacterial cell.

[0412] Also provided by the present disclosure is a population of cells comprising at least one host cell described herein. The population of cells in some aspects is a heterogeneous population comprising the host cell comprising vectors described, in addition to at least one other cell, which does not comprise any of the vectors. Alternatively, in some aspects, the population of cells is a substantially homogeneous population, in which the population comprises mainly host cells (e.g., consisting essentially of) comprising the vector. The population in some aspects is a clonal population of cells, in which all cells of the population are clones of a single host cell comprising a vector, such that all cells of the population comprise the vector. In exemplary embodiments of the present disclosure, the population of cells is a clonal population comprising host cells comprising a vector as described herein.

Pharmaceutical Compositions

[0413] Compositions comprising an IL-21 polypeptide, a conjugate comprising the IL-21 polypeptide, a fusion protein comprising the IL-21 polypeptide and an antigen-binding protein, a conjugate comprising the antigen-binding protein, a fusion protein comprising the antigen-binding protein of the disclosure, a nucleic acid, vector, or host cell, of the present disclosure, or a combination thereof, are provided herein. The compositions in some aspects comprise an IL-21

polypeptide, an antigen-binding protein, a conjugate, fusion protein, nucleic acid, vector, or host cell of the present disclosure, or a combination thereof, in isolated and/or purified form. In some aspects, the composition comprises a single type (*e.g.*, structure/combination of mutations relative to wild-type IL-21 polypeptide) of an IL-21 polypeptide, antigen-binding protein, a conjugate, fusion protein, nucleic acid, vector, or host cell of the present disclosure, or comprises a combination of two or more different types of IL-21 polypeptides, antigen-binding proteins, conjugates, fusion proteins, nucleic acids, vectors or host cells of the present disclosure.

[0414] In some embodiments, the composition comprises agents which enhance the chemico-physico features of an IL-21 polypeptide, antigen-binding protein, a conjugate, fusion protein, nucleic acid, vector, or host cell, *e.g.*, *via* stabilizing, for example, the IL-21 polypeptide or fusion protein at certain temperatures (*e.g.*, room temperature), increasing shelf life, reducing degradation, *e.g.*, oxidation protease mediated degradation, increasing half-life of, for example, the IL-21 mutein or fusion protein, etc. In some embodiments, the composition comprises any of the agents disclosed herein as a heterologous moiety or conjugate moiety, optionally, in admixture with an IL-21 polypeptide, conjugates, fusion proteins, nucleic acids, vectors, or host cells of the present disclosure.

[0415] In some embodiments, the composition additionally comprises a pharmaceutically acceptable carrier, diluents, or excipient. In some embodiments, an IL-21 polypeptides, antigen-binding proteins, conjugates, fusion proteins, nucleic acids, vectors, or host cells as presently disclosed (hereinafter referred to as "active agents"), or combinations thereof, are formulated into a pharmaceutical composition comprising the active agent, along with a pharmaceutically acceptable carrier, diluent, or excipient. In this regard, the present disclosure further provides pharmaceutical compositions comprising an active agent (any of the IL-21 polypeptides, antigen-binding proteins, conjugates, fusion proteins, nucleic acids, vectors, or host cells of the present disclosure), which pharmaceutical composition is suitable for administration to a subject, *e.g.*, a mammal.

[0416] In some embodiments, the active agent is present in the pharmaceutical composition at a purity level suitable for administration to a subject. In some embodiments, the active agent has a purity level of at least about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98% or about 99%, and a pharmaceutically acceptable diluent, carrier or excipient. In some embodiments, the compositions contain an active agent at a concentration of about 0.001 to about 200.0 mg/ml.

[0417] In exemplary aspects, the pharmaceutical compositions comprise a pharmaceutically acceptable carrier. As used herein, the term "pharmaceutically acceptable carriers" include but are not limited to any of the standard pharmaceutical carriers, such as a phosphate buffered saline

solution, water, emulsions such as an oil/water or water/oil emulsion, and various types of wetting agents. The term also encompasses any of the agents approved by a regulatory agency of the US Federal government or listed in the US Pharmacopeia for use in animals, including humans.

[0418] The pharmaceutical composition can comprise any pharmaceutically acceptable ingredients, including, for example, acidifying agents, additives, adsorbents, aerosol propellants, air displacement agents, alkalizing agents, anticaking agents, anticoagulants, antimicrobial preservatives, antioxidants, antiseptics, bases, binders, buffering agents, chelating agents, coating agents, coloring agents, desiccants, detergents, diluents, disinfectants, disintegrants, dispersing agents, dissolution enhancing agents, dyes, emollients, emulsifying agents, emulsion stabilizers, fillers, film forming agents, flavor enhancers, flavoring agents, flow enhancers, gelling agents, granulating agents, humectants, lubricants, mucoadhesive, ointment bases, ointments, oleaginous vehicles, organic bases, pastille bases, pigments, plasticizers, polishing agents, preservatives, sequestering agents, skin penetrants, solubilizing agents, solvents, stabilizing agents, suppository bases, surface active agents, surfactants, suspending agents, sweetening agents, therapeutic agents, thickening agents, tonicity agents, toxicity agents, viscosity-increasing agents, water-absorbing agents, water-miscible cosolvents, water softeners, or wetting agents. *See, e.g.*, the Handbook of Pharmaceutical Excipients, Third Edition, A. H. Kibbe (Pharmaceutical Press, London, U K, 2000), which is incorporated by reference in its entirety. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980), which is incorporated by reference in its entirety.

[0419] In some embodiments, the pharmaceutical composition comprises formulation materials that are nontoxic to recipients at the dosages and concentrations employed. In specific embodiments, pharmaceutical compositions comprising an active agent and one or more pharmaceutically acceptable salts; polyols; surfactants; osmotic balancing agents; tonicity agents; anti-oxidants; antibiotics; antimycotics; bulking agents; lyoprotectants; anti-foaming agents; chelating agents; preservatives; colorants; analgesics; or additional pharmaceutical agents. In exemplary aspects, the pharmaceutical composition comprises one or more polyols and/or one or more surfactants, optionally, in addition to one or more excipients, including but not limited to, pharmaceutically acceptable salts; osmotic balancing agents (tonicity agents); anti-oxidants; antibiotics; antimycotics; bulking agents; cytoprotectants; anti-foaming agents; chelating agents; preservatives; colorants; and analgesics.

[0420] In some embodiments, the pharmaceutical composition can contain formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the

composition. In such embodiments, suitable formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine); antimicrobials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetra acetic acid (EDTA)); complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrin); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents; hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate, triton, tromethamine, lecithin, cholesterol, tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants. See, REMINGTON'S PHARMACEUTICAL SCIENCES, 18″ Edition, (A. R. Genrmo, ed.), 1990, Mack Publishing Company.

[0421] The pharmaceutical compositions can be formulated to achieve a physiologically compatible pH. In some embodiments, the pH of the pharmaceutical composition can be for example between about 4 or about 5 and about 8.0 or about 4.5 and about 7.5 or about 5.0 to about 7.5. In exemplary embodiments, the pH of the pharmaceutical composition is between 5.5 and 7.5.

Routes of Administration

[0422] With regard to the present disclosure, the active agent, or pharmaceutical composition comprising the same, can be administered to the subject via any suitable route of administration. For example, the active agent can be administered to a subject via parenteral, nasal, oral, pulmonary, topical, vaginal, or rectal administration. The following discussion on routes of administration is merely provided to illustrate exemplary embodiments and should not be construed as limiting the scope in any way.

[0423] Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and

solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The term, "parenteral" means not through the alimentary canal but by some other route such as subcutaneous, intramuscular, intraspinal, or intravenous. The active agent of the present disclosure can be administered with a physiologically acceptable diluent in a pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an alcohol, such as ethanol or hexadecyl alcohol, a glycol, such as propylene glycol or polyethylene glycol, dimethyl sulfoxide, glycerol, ketals such as 2,2-dimethyl-153-dioxolane-4-methanol, ethers, poly(ethylene glycol) 400, oils, fatty acids, fatty acid esters or glycerides, or acetylated fatty acid glycerides with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers, methylcellulose, hydroxypropyl methylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

[0424] Oils, which can be used in parenteral formulations include petroleum, animal, vegetable, or synthetic oils. Specific examples of oils include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral. Suitable fatty acids for use in parenteral formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters.

[0425] Suitable soaps for use in parenteral formulations include fatty alkali metal, ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylenepolypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl-β-aminopropionates, and 2-alkyl-imidazoline quaternary ammonium salts, and I mixtures thereof.

[0426] The parenteral formulations in some embodiments contain from about 0.5% to about 25% by weight of the active agent of the present disclosure in solution. Preservatives and buffers can be used. In order to minimize or eliminate irritation at the site of injection, such compositions can contain one or more nonionic surfactants having a hydrophile-lipophile balance (HLB) of from about 12 to about 17. The quantity of surfactant in such formulations will typically range from about 5% to about 15% by weight. Suitable surfactants include polyethylene glycol sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol.

The parenteral formulations in some aspects are presented in unit-dose or multi-dose sealed containers, such as ampoules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid excipient, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions in some aspects are prepared from sterile powders, granules, and tablets of the kind previously described.

[0427] Injectable formulations are in accordance with the present disclosure. The requirements for effective pharmaceutical carriers for injectable compositions are well-known to those of ordinary skill in the art (see, e.g., Pharmaceutics and Pharmacy Practice, J. B. Lippincott Company, Philadelphia, Pa., Banker and Chalmers, eds., pages 238-250 (1982), and ASHP Handbook on Injectable Drugs, Toissel, 4th ed., pages 622-630 (1986)).

Dosages

[0428] The active agents of the disclosure are believed to be useful in methods more specifically activating CD8+ T cells through IL-21 signaling, as described herein, and are thus believed to be useful in methods of treating or preventing one or more diseases, *e.g.*, cancer. For purposes of the disclosure, the amount or dose of the active agent administered should be sufficient to effect, *e.g.*, a therapeutic or prophylactic response, in the subject or animal over a reasonable time frame. For example, the dose of the active agent of the present disclosure should be sufficient to treat cancer as described herein in a period of from about 1 to 4 minutes, 1 to 4 hours or 1 to 4 weeks or longer, e.g., 5 to 20 or more weeks, from the time of administration. In certain embodiments, the time period could be even longer. The dose will be determined by the efficacy of the particular active agent and the condition of the animal (*e.g.*, human), as well as the body weight of the animal (*e.g.*, human) to be treated.

[0429] Many assays for determining an administered dose are known in the art. For purposes herein, an assay, which comprises comparing the extent to which cancer is treated upon administration of a given dose of the active agent of the present disclosure to a mammal among a set of mammals, each set of which is given a different dose of the active agent, could be used to determine a starting dose to be administered to a mammal. The extent to which cancer is treated upon administration of a certain dose can be represented by, for example, the cytotoxicity of the active agent or the extent of tumor regression achieved with the active agent in a mouse xenograft model. Methods of measuring cytotoxicity of the fusion proteins and methods of assaying tumor regression are known in the art.

[0430] The dose of the active agent of the present disclosure also will be determined by the existence, nature and extent of any adverse side effects that might accompany the administration of

a particular active agent of the present disclosure. Typically, the attending physician will decide the dosage of the active agent of the present disclosure with which to treat each individual patient, taking into consideration a variety of factors, such as age, body weight, general health, diet, sex, active agent of the present disclosure to be administered, route of administration, and the severity of the condition being treated. By way of example and not intending to limit the present disclosure, the dose of the active agent of the present disclosure can be about 0.000001 to about 1 g/kg body weight of the subject being treated/day, from about 0.000001 to about 0.001 g/kg body weight/day, or about 0.01 mg to about 1 g/kg body weight/day.

Controlled Release Formulations

[0431] In some embodiments, the active agents described herein can be modified into a depot form, such that the manner in which the active agent of the present disclosure is released into the body to which it is administered is controlled with respect to time and location within the body (see, for example, U.S. Pat. No. 4,450,150). Depot forms of active agents of the present disclosure can be, for example, an implantable composition comprising the active agents and a porous or non-porous material, such as a polymer, wherein the active agent is encapsulated by or diffused throughout the material and/or degradation of the non-porous material. The depot is then implanted into the desired location within the body of the subject and the active agent is released from the implant at a predetermined rate.

[0432] The pharmaceutical composition comprising the active agent in certain aspects is modified to have any type of in vivo release profile. In some aspects, the pharmaceutical composition is an immediate release, controlled release, sustained release, extended release, delayed release, or bi-phasic release formulation. Methods of formulating peptides for controlled release are known in the art. See, for example, Qian et al., J Pharm 374: 46-52 (2009) and International Patent Application Publication Nos. WO 2008/130158, WO2004/033036; WO2000/032218; and WO 1999/040942.

[0433] The instant compositions can further comprise, for example, micelles or liposomes, or some other encapsulated form, or can be administered in an extended release form to provide a prolonged storage and/or delivery effect.

Combinations

[0434] In some embodiments, the IL-21 polypeptide or a targeted cytokine construct composition is administered in combination with a cell therapy (*e.g.*, an adoptive cell therapy, a T-cell therapy, such as a CAR-T cell therapy, a TCR-T cell therapy, a tumor infiltrating lymphocyte (TIL) therapy), cancer vaccine, anti-angiogenesis agent, antibody drug conjugate, chemotherapeutic

agent, immune stimulating agent (e.g. IL-2), immune priming agent (e.g. IL-2) or immune checkpoint inhibitor (ICI). In some embodiments, the chemotherapeutic agent is a kinase inhibitor, antimetabolite, cytotoxin or cytostatic agent, anti-hormonal agent, platinum-based chemotherapeutic agent, methyltransferase inhibitor, antibody, or anti-cancer peptide.

[0435] In some cases, the IL-21 polypeptide disclosed herein is used to treat, or enhance the efficacy of treatment associated with tumor types not responsive to a therapy using IL-2 (*e.g.*, a therapy that promotes IL-2 signaling in one or more cells of the subject). In some cases, the IL-21 polypeptide enhances the efficacy of treatment associated with tumor types not responsive to a therapy that targets PD-1 (*e.g.*, a therapy that inhibits PD-1 signaling in one or more cells of the subject). In some cases, the IL-21 polypeptide enhances the efficacy of treatment associated with tumor types not responsive to both a therapy comprising IL-2 and a therapy that targets PD-1. For example, PD-1 resistant tumor are associated with reduced memory T cell numbers and increased exhausted T cell numbers, suggesting use of the IL-21 polypeptide of the present disclosure can improve the outcome of the treatment of these tumors.

[0436] In some embodiments, the immune checkpoint inhibitor targets PD-L1, PD-1, CTLA-4, CEACAM, LAIR1, CD160, 2B4, CD80, CD86, CD276, VTCN1, HVEM, KIR, A2AR, MHC class I, MHC class II, GALS, adenosine, TGFR, OX40, CD137, CD40, CD47, TREM1, TREM2, HLA-G, CCR4, CCR8, CD39, CD73, IDO, CSF1R, TIM-3, BTLA, VISTA, LAG-3, TIGIT, IDO, MICA/B, LILRB4, SIGLEC-15, or arginase, including without limitation an inhibitor of PD-1 (*e.g.*, an anti-PD-1 antibody), PD-L1 (*e.g.*, an anti-PD-L1 antibody), or CTLA-4 (*e.g.*, an anti-CTLA-4 antibody).

[0437] Examples of anti-PD-1 antibodies include, without limitation, pembrolizumab, nivolumab, cemiplimab, zimberelimab (Arcus), sasanlimab (Pfizer), JTX-4014, spartalizumab (PDR001; Novartis), camrelizumab (SHR1210; Jiangsu HengRui Medicine), sintilimab (IBI308; Innovent and Eli Lilly), tislelizumab (BGB-A317), toripalimab (JS 001), dostarlimab (TSR-042, WBP-285), INCMGA00012 (MGA012), AMP-224, and AMP-514 (MEDI0680). Examples of anti-PD-L1 antibodies include, without limitation, atezolizumab, avelumab, durvalumab, KN035, and CK-301 (Checkpoint Therapeutics). Examples of PD-L1 inhibitors (non-antibody based) include, without limitation, AUNP12, CA-170, and BMS-986189. Examples of anti-CTLA-4 antibodies include, without limitation, ipilimumab, tremelimumab, BMS-986218, BMS-986249, BMS-986288, HBM4003, ONC-392, KN044, ADG116, ADU-1604, AGEN1181, AGEN1884, MK-1308, and REGN4659.

[0438] In some embodiments, an IL-21 polypeptide or fusion protein thereof is administered in combination with a tumor targeting antibody. In some embodiments, a tumor targeting antibody is

specific for a cancer antigen or a tumor antigen. In some embodiments, the tumor targeting antibody is Rituxan or cetuximab.

[0439] The IL-21 polypeptide of the present disclosure can be administered in combination with another cytokine therapy. In some embodiments, the cytokine therapy comprises IL-2 or any functional variant thereof, or a target IL-2 construct comprising IL-2 or functional variant thereof (for example, such as those described in international patent application nos. PCT/US20/36454 and PCT/US21/56312, each of which is hereby incorporated by reference in its entirety). In some embodiments, the IL-21 polypeptide or target cytokine construct composition is administered in combination with IL-2 or functional variant thereof, a target IL-2 construct comprising IL-2 or functional variant thereof, such as those described in international patent application nos. PCT/US20/36454 and PCT/US21/56312.

Kits

[0440] The present disclosure additionally provides kits comprising an IL-21 polypeptide or functional fragment or variant thereof, an antigen-binding protein, a conjugate, fusion protein, nucleic acid, vector, or host cell of the present disclosure, or a combination thereof. The kit in exemplary aspects comprises at least one IL-21 polypeptide or a functional fragment or a variant thereof, antigen-binding protein, a conjugate, fusion protein, nucleic acid, vector, or host cell of the present disclosure, or a combination thereof, in a container. In exemplary aspects, the at least one IL-21 polypeptide or a functional fragment or a variant thereof, antigen-binding protein, a conjugate, fusion protein, nucleic acid, vector, or host cell of the present disclosure, is provided in the kit as a unit dose. For purposes herein "unit dose" refers to a discrete amount dispersed in a suitable carrier. In exemplary aspects, the unit dose is the amount sufficient to provide a subject with a desired effect, e.g., treatment of cancer. In exemplary aspects, the kit comprises several unit doses, e.g., a week or month supply of unit doses, optionally, each of which is individually packaged or otherwise separated from other unit doses. In some embodiments, the components of the kit/unit dose are packaged with instructions for administration to a patient. In some embodiments, the kit comprises one or more devices for administration to a patient, e.g., a needle and syringe, and the like. In some aspects, the at least one IL-21 polypeptide or a functional fragment or a variant thereof, antigen-binding protein, a conjugate, fusion protein, nucleic acid, vector, or host cell of the present disclosure, or a combination thereof, is/are pre-packaged in a ready to use form, *e.g.*, a syringe, an intravenous bag, etc. In exemplary aspects, the ready to use form is for a single use. In exemplary aspects, the kit comprises multiple single use, ready to use forms of the at least one IL-21 polypeptide or a functional fragment or a variant thereof, antigen-

binding protein, a conjugate, fusion protein, nucleic acid, vector, or host cell of the present disclosure. In some aspects, the kit further comprises other therapeutic or diagnostic agents or pharmaceutically acceptable carriers (*e.g.*, solvents, buffers, diluents, etc.), including any of those described herein.

Methods of Manufacture

[0441] The IL-21 polypeptide or functional fragment or variant thereof, as described herein, may be obtained by methods known in the art. Suitable methods of *de novo* synthesizing polypeptides are described in, for example, Chan et al., Fmoc Solid Phase Peptide Synthesis, Oxford University Press, Oxford, United Kingdom, 2005; Peptide and Protein Drug Analysis, ed. Reid, R., Marcel Dekker, Inc., 2000; Epitope Mapping, ed. Westwood et al., Oxford University Press, Oxford, United Kingdom, 2000; and U.S. Pat. No. 5,449,752. Additional exemplary methods of making the peptides of the invention are set forth herein.

[0442] In some embodiments, the IL-21 polypeptide or functional fragment or variant thereof, as described herein, can be commercially synthesized by companies, such as Synpep (Dublin, Calif.), Peptide Technologies Corp. (Gaithersburg, Md.), Multiple Peptide Systems (San Diego, Calif.), Peptide 2.0 Inc. (Chantilly, Va.), and American Peptide Co. (Sunnyvale, Calif.). In this respect, the IL-21 muteins can be synthetic, recombinant, isolated, and/or purified.

[0443] Also, in some aspects, an IL-21 polypeptide or functional fragment or variant thereof, as described herein, can be recombinantly produced using a nucleic acid encoding the amino acid sequence of the peptide using standard recombinant methods. See, for instance, Sambrook *et al.*, Molecular Cloning: A Laboratory Manual· 3rd ed., Cold Spring Harbor Press, Cold Spring Harbor, N.Y. 2001; and Ausubel et al., Current Protocols in Molecular Biology, Greene Publishing Associates and John Wiley & Sons, N Y, 1994.

[0444] Methods of making an IL-21 polypeptide or functional fragment or variant thereof, as described herein, or a targeted cytokine construct comprising an IL-21 polypeptide or functional fragment or variant thereof, as described herein, are provided herein. The method, in exemplary embodiments, comprises culturing a host cell of the present disclosure to express the IL-21 polypeptide or functional fragment or variant thereof, as described herein, or a targeted cytokine construct, and harvesting the expressed IL-21 polypeptide or functional fragment or variant thereof.

[0445] Methods of making fusion protein comprising an IL-21 polypeptide or a functional fragment or a variant thereof or a targeted cytokine construct as described herein are also provided herein. The method, in exemplary embodiments, comprises culturing a host cell of the present disclosure to express the fusion protein and harvesting the expressed fusion protein.

[0446] In exemplary embodiments, the method comprises culturing a host cell comprising a nucleic acid encoding the IL-21 polypeptide or a functional fragment or a variant thereof or fusion protein as described herein so as to express the IL-21 polypeptide or a functional fragment or a variant thereof or fusion protein. The host cell can be any of the host cells described herein. In exemplary aspects, the host cell is selected from the group consisting of: CHO cells, NS0 cells, COS cells, VERO cells, and BHK cells. In exemplary aspects, the step of culturing a host cell comprises culturing the host cell in a growth medium to support the growth and expansion of the host cell. In exemplary aspects, the growth medium increases cell density, culture viability and productivity in a timely manner. In exemplary aspects, the growth medium comprises amino acids, vitamins, inorganic salts, glucose, and serum as a source of growth factors, hormones, and attachment factors. In exemplary aspects, the growth medium is a fully chemically defined media consisting of amino acids, vitamins, trace elements, inorganic salts, lipids and insulin or insulin-like growth factors. In addition to nutrients, the growth medium also helps maintain pH and osmolality. Several growth media are commercially available and are described in the art. See, e.g., Arora, "Cell Culture Media: A Review" MATER METHODS 3:175 (2013).

[0447] In exemplary aspects, the method of making an IL-21 polypeptide or a functional fragment or a variant thereof or fusion protein of the present disclosure comprises culturing the host cell in a feed medium. In exemplary aspects, the method comprises culturing in a feed medium in a fed-batch mode. Methods of recombinant protein production are known in the art. See, e.g., Li et al., "Cell culture processes for monoclonal antibody production" MAbs 2(5): 466-477 (2010).

[0448] The method making an IL-21 polypeptide or a functional fragment or a variant thereof or fusion protein can comprise one or more steps for purifying the polypeptide from a cell culture or the supernatant thereof and preferably recovering the purified protein. In exemplary aspects, the method comprises one or more chromatography steps, *e.g.*, affinity chromatography (e.g., protein A affinity chromatography), ion exchange chromatography, hydrophobic interaction chromatography. In exemplary aspects, the method comprises purifying the protein using a Protein A affinity chromatography resin.

[0449] In exemplary embodiments, the method further comprises steps for formulating the purified protein, etc., thereby obtaining a formulation comprising the purified protein. Such steps are described in Formulation and Process Development Strategies for Manufacturing, eds. Jameel and Hershenson, John Wiley & Sons, Inc. (Hoboken, N.J.), 2010.

Other Embodiments

[0450] The disclosure relates to the following embodiments. Throughout this section, the term "embodiment" is abbreviated as "E" followed by an ordinal. For example, "E1" is equivalent to Embodiment 1.

[0451] E1. An IL-21 polypeptide or a functional fragment or a variant thereof comprising a polypeptide sequence having at least 80% sequence identity to SEQ ID NO: 1, wherein the IL-21 polypeptide has an isoelectric point that is at least about 0.6 units to about 5 units lower, compared to that of a wild-type IL-21 protein having a sequence of SEQ ID NO: 1.

[0452] E2. The IL-21 polypeptide of embodiment 1, wherein the isoelectric point of SEQ ID NO: 1 is about 9.42.

[0453] E3. The IL-21 polypeptide of embodiment 1 or 2, wherein the IL-21 polypeptide has an isoelectric point of about 7.12 to about 8.72.

[0454] E4. The IL-21 polypeptide of any one of embodiments 1-3, wherein the polypeptide provides an improved exposure compared to the wild-type IL-21 protein, as measured by at least about 1.5 times greater under the curve (AUC) for the polypeptide, relative to that of the wild-type IL-21, when administered to a subject, at equivalent concentrations.

[0455] E5. The IL-21 polypeptide of any one of embodiments 1-4, wherein the IL-21 polypeptide comprises a mutation at one or more positions selected from the group consisting of: K56, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1.

[0456] E6. The IL-21 polypeptide of embodiment 5, comprising a mutation at a position selected from the group consisting of: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117 of SEQ ID NO: 1.

[0457] E7. The IL-21 polypeptide of any one of embodiments 1-6, comprising an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, and SEQ ID NO: 23 – SEQ ID NO: 40.

[0458] E8. The IL-21 polypeptide of any one of embodiments 1-7, comprising an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, and SEQ ID NO: 41 - SEQ ID NO: 93.

[0459] E9. A targeted cytokine construct comprising an IL-21 polypeptide or a functional fragment or a variant thereof according to any one of embodiments 1-8 and an antibody or an antigen binding fragment thereof.

[0460] E10. The targeted cytokine construct of embodiment 9, wherein the antibody or antigen binding fragment thereof specifically binds to a CD8+ T cell.

[0461] E11. The targeted cytokine construct of embodiment 9 or 10, wherein the antibody or antigen binding fragment thereof specifically binds to at least one of: CD8α, CD8αα, or CD8αβ.

[0462] E12. The targeted cytokine construct of embodiment 9 or 10, wherein the antibody or antigen binding fragment thereof specifically binds to CD8β.

[0463] E13. A targeted cytokine construct comprising:

a) an IL-21 polypeptide or a functional fragment or a variant thereof comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1; and

b) an antibody or an antigen binding fragment thereof that specifically binds to at least one of CD8α, CD8αα, or CD8αβ.

[0464] E14. A targeted cytokine construct comprising:

a) an IL-21 polypeptide or a functional fragment or a variant thereof comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8β.

[0465] E15. The targeted cytokine construct of embodiment 13 or 14, wherein the IL-21 polypeptide comprises a mutation at one or more amino acid positions of SEQ ID NO: 1.

[0466] E16. The targeted cytokine construct of any one of embodiments 13-15, wherein the IL-21 polypeptide comprises a mutation at one or more positions selected from the group consisting of: K56, T81, N82, A83, G84, R85, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1.

[0467] E17. The targeted cytokine construct of any one of embodiments 13-16, wherein the IL-21 polypeptide comprises mutation at a position selected from the group consisting of: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117 of SEQ ID NO: 1.

[0468] E18. The targeted cytokine construct of any one of embodiments 13-17, wherein the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, or SEQ ID NO: 15, and SEQ ID NO: 23 – SEQ ID NO: 40.

[0469] E19. The targeted cytokine construct of any one of embodiments 13-18, wherein the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, and SEQ ID NO: 41 - SEQ ID NO: 93.

[0470] E20. The targeted cytokine construct of any one of embodiments 13-19, wherein the construct activates CD8+ T cells with at least about 10-fold or greater potency as compared to activation of NK or CD4+ T cells.

[0471] E21. The targeted cytokine construct of any one of embodiments 13-20, wherein the construct activates CD8+ T cells with a potency that is at least about 10-fold to about 100,000-fold greater as compared to activation of NK or CD4+ T cells.

[0472] E22. The targeted cytokine construct of any one of embodiments 13-21, wherein the antibody or antigen binding fragment thereof comprises:

i) a first polypeptide, arranged from N-to-C terminus, comprising:

a variable light chain (VL) amino acid sequence, and

a) a light chain constant region amino acid sequence (CL1);

ii) a second polypeptide, arranged from N-to-C terminus, comprising:

a) a variable heavy chain (VH) amino acid sequence,

b) a heavy chain CH1 constant region amino acid sequence,

c) a hinge region amino acid sequence,

d) a heavy chain CH2 constant region amino acid sequence, and

e) a heavy chain CH3 constant region amino acid sequence; and

iii) a third polypeptide, arranged from N-to-C terminus, comprising:

a) a hinge region amino acid sequence,

b) a heavy chain CH2 constant region amino acid sequence, and

c) a heavy chain CH3 constant region amino acid sequence,

wherein the CH2 and CH3 domains of each of the second and third polypeptides form an Fc domain.

[0473] E23. The targeted cytokine construct of embodiment 22, wherein the third polypeptide, arranged from N-to-C terminus, comprises:

a) a variable heavy chain (VH) amino acid sequence,

b) a heavy chain CH1 constant region amino acid sequence,

c) a hinge region amino acid sequence,

d) a heavy chain CH2 constant region amino acid sequence,

e) a heavy chain CH3 constant region amino acid sequence,

wherein the antibody or antigen binding fragment thereof further comprises:

iv) a fourth polypeptide, arranged from N-to-C terminus, comprising:

a) a variable light chain (VL) amino acid sequence, and

b) a light chain constant amino acid sequence (CL1).

[0474] E24. The targeted cytokine construct of embodiment 22 or 23, wherein the IL-21 polypeptide or a functional fragment or a variant thereof and the antibody or antigen binding fragment thereof are operably linked to each other.

[0475] E25. The targeted cytokine construct of embodiment 24, wherein the IL-21 polypeptide or a functional fragment or a variant thereof is linked to the N-terminus or C-terminus of the antibody or antigen binding fragment thereof.

[0476] E26. The targeted cytokine construct of any one of embodiments 22-25, wherein the IL-21 polypeptide or a functional fragment or variant thereof is conjugated to the C-terminus of the second or the third polypeptide.

[0477] E27. The targeted cytokine construct of any one of embodiments 9-26, comprising at least one molecule of the IL-21 polypeptide.

[0478] E28. The targeted cytokine construct of any one of embodiments 22-27, wherein the Fc domain is a human IgG Fc domain.

[0479] E29. The targeted cytokine construct of any one of embodiments 22-28, wherein the Fc domain is an IgG1, IgG2, IgG3, or IgG4 Fc domain.

[0480] E30. The targeted cytokine construct of embodiment 28 or 29, wherein the Fc domain comprises one or more modifications that promote heterodimerization.

[0481] E31. The targeted cytokine construct of any one of embodiments 22-30, wherein the second polypeptide comprises a knob modification in the CH2 or the CH3 domain, and the third polypeptide comprises a hole modification in the CH2 or the CH3 domain; or wherein the third polypeptide comprises a knob modification in the CH2 or the CH3 domain and the second polypeptide comprises a hole modification in the CH2 or the CH3.

[0482] E32. The fusion protein of any one of embodiments 22-31, wherein at least one of the second and the third polypeptide comprises the following mutations: L234A, L235A, and G237A, numbering according to the EU index.

[0483] E33. A targeted cytokine construct comprising:

a) an IL-21 polypeptide or a functional fragment or variant thereof comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1; and

b) an antibody or an antigen binding fragment thereof comprising a first antigen binding arm and a second antigen binding arm, wherein the first and the second antigen binding arms bind to two different antigens, wherein at least one of the first and the second antigen binding arm specifically binds to CD8α, CD8αα, CD8αβ.

[0484] E34. A targeted cytokine construct comprising:

a) an IL-21 polypeptide or a functional fragment or variant thereof comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1; and

b) an antibody or an antigen binding fragment thereof comprising a first antigen binding arm and a second antigen binding arm, wherein the first and the second antigen binding arms bind to two different antigens, wherein at least one of the first and the second antigen binding arm specifically binds to CD8β.

[0485] E35. The targeted cytokine construct of embodiment 33 or 34, wherein the construct activates CD8+ T cells with at least about 10-fold or greater potency as compared to activation of NK cells or CD4+ T cells.

[0486] E36. The targeted cytokine construct of embodiment 34 or 35, wherein the construct activates CD8+ T cells a potency that is at least about 10-fold to about 100,000-fold greater as compared to activation of NK cells or CD4+ T cells.

[0487] E37. The targeted cytokine construct of any one of embodiments 34-36, wherein the IL-21 polypeptide or a functional fragment thereof or a variant thereof comprises a mutation at a position selected from the group consisting of: K56, T81, N82, A83, R85, G84, R86, Q87, K88, H89, R90, L91, and T92 of SEQ ID NO: 1.

[0488] E38. The targeted cytokine construct of any one of embodiments 34-37, wherein the IL-21 polypeptide or a functional fragment thereof or a variant thereof comprises a mutation at a position selected from the group consisting of: R5, I8, R9, R11, Q12, I14, D15, D18, Q19, Y23, R65, S70, K72, K73, K75, R76, K77, S80, Q116, and K117 of SEQ ID NO: 1.

[0489] E39. The targeted cytokine construct of any one of embodiments 34-38, wherein the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid sequence that is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 14, SEQ ID NO: 15, and SEQ ID NO: 23 – SEQ ID NO: 40.

[0490] E40. The targeted cytokine construct of any one of embodiments 34-39, wherein the IL-21 polypeptide or a functional fragment or a variant thereof comprises an amino acid sequence that

is at least about 85%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to a sequence selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 21, and SEQ ID NO: 41 – SEQ ID NO:93.

[0491] E41. A polynucleotide encoding an IL-21 polypeptide or a functional fragment or a variant thereof according to any one of embodiments 1-8.

[0492] E42. A polynucleotide that codes for a targeted cytokine construct comprising an IL-21 polypeptide or a functional fragment or a variant thereof and an antibody or an antigen binding fragment thereof, according to any one of embodiments 9-40, the polynucleotide comprising a coding sequence for the IL-21 polypeptide and a coding sequence for the antibody or an antigen binding fragment thereof.

[0493] E43. A vector comprising the polynucleotide of embodiment 41 or 42

[0494] E44. A host cell comprising the polynucleotide of embodiment 41 or 42 or the vector of embodiment 43.

[0495] E45. A pharmaceutical composition comprising an IL-21 polypeptide or a functional fragment or variant thereof according to any one of embodiments 1-8 and a pharmaceutically acceptable carrier.

[0496] E46. A pharmaceutical composition comprising a targeted cytokine construct according to any one of embodiments 9-40 and a pharmaceutically acceptable carrier.

[0497] E47. A method for selective activation of CD8+ T cells, wherein the method comprises contacting a population of cells comprising CD8+ T cells, CD4+ T cells, and NK cells, with a targeted cytokine construct according to any one of embodiments 9-40.

[0498] E48. The method of embodiment 47, wherein selective activation comprises activation of CD8+ T cells with at least about 10-fold or greater potency, as compared to activation of NK cells or CD4+ T cells in the population of cells.

[0499] E49. The method of embodiment 47 or 48, wherein selective activation comprises activation of CD8+ T cells with a potency that is at least about 10-fold to about 100,000-fold greater as compared to activation of NK cells or CD4+ T cells in the population of cells.

[0500] E50. The method of any one of embodiments 47-49, wherein the selective activation of CD8+ T cells results in increased STAT3 phosphorylation of the CD8+ T cells compared to the STAT3 phosphorylation of the NK cells or CD4+ T cells in the population of cells.

[0501] E51. A method of treating a disease in a subject, the method comprising administering an IL-21 polypeptide or a functional fragment or a variant thereof according to any one of

embodiments 1-8, a targeted cytokine construct according to any one of embodiments 9-40, or a pharmaceutical composition according to embodiment 45 or 46.

[0502] E52. The method of embodiment 51, further comprising an additional therapeutic agent.

[0503] E53. The method of embodiment 51 or 52, wherein the disease comprises a cancer or a chronic infection.

[0504] E54. The method of any one of embodiments 51-53, wherein the disease comprises the cancer and wherein the cancer is acute lymphoblastic leukemia (ALL) (including non T cell ALL), acute myeloid leukemia, B cell prolymphocytic leukemia, B cell acute lymphoid leukemia ("BALL"), blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloid leukemia, chronic or acute leukemia, diffuse large B cell lymphoma (DLBCL), follicular lymphoma (FL), hairy cell leukemia, Hodgkin's Disease, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, monoclonal gammopathy of undetermined significance (MGUS), multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin's lymphoma (NHL), plasma cell proliferative disorder (including asymptomatic myeloma (smoldering multiple myeloma or indolent myeloma), plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, plasmacytomas (including plasma cell dyscrasia; solitary myeloma; solitary plasmacytoma; extramedullary plasmacytoma; and multiple plasmacytoma), POEMS syndrome (also known as Crow-Fukase syndrome; Takatsuki disease; and PEP syndrome), primary mediastinal large B cell lymphoma (PMBC), small cell- or a large cell-follicular lymphoma, splenic marginal zone lymphoma (SMZL), systemic amyloid light chain amyloidosis, T cell acute lymphoid leukemia ("TALL"), T cell lymphoma, transformed follicular lymphoma, or Waldenstrom macroglobulinemia, Mantle cell lymphoma (MCL), Transformed follicular lymphoma (TFL), Primary mediastinal B cell lymphoma (PMBCL), Multiple myeloma, Hairy cell lymphoma/leukemia, lung cancer, small-cell lung cancer, non-small cell lung (NSCL) cancer, bronchioloalveolar cell lung cancer, squamous cell cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, head and neck cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, thyroid cancer, uterine cancer, , gastrointestinal cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, endometrial carcinoma, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the cervix, carcinoma of the vagina, vulval cancer, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the

bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, bladder cancer, liver cancer, hepatoma, hepatocellular cancer, cervical cancer, salivary gland carcinoma, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwannomas, ependymomas, medulloblastomas, meningiomas, squamous cell carcinomas, pituitary adenoma and Ewing's sarcoma, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers.

[0505] E55. The targeted cytokine construct of any one of embodiments 13-40, wherein the antibody or antigen binding fragment comprises a heavy chain variable (VH) domain and a light chain variable (VL) domain; and wherein:

(a) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:137, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:138, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:139; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:140, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:141, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:142;

(b) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:149, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:150, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:151; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:154;

(c) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:155, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:156, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:157; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:158, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:159, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:160;

(d) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:161, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:162, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:163; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:164, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:165, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:166;

(e) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:167, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:168, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:169; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:170, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:171, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:172;

(f) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:173, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:174, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:175; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

(g) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:179, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:180, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:181; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:182, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:183, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:184;

(h) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:143, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:144, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:145; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:146, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:147, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:148;

(i) the VH domain comprises a CDR-H1 comprising the amino acid sequence of X_1X_2AIS, wherein X_1 is S, K, G, N, R, D, T, or G, and wherein X_2 is Y, L, H, or F (SEQ ID NO:185), a CDR-H2 comprising the amino acid sequence of $X_1X_2X_3PX_4X_5X_6X_7X_8X_9YX_{10}QKFX_{11}G$, wherein X_1 is G or H, X_2 is I or F, X_3 is I, N, or M, X_4 is G, N, H, S, R, I, or A, X_5 is A, N, H, S, T, F, or Y, X_6 is A, D, or G, X_7 is T, E, K, V, Q, or A, X_8 is A or T, X_9 is N or K, X_{10} is A or N, and X_{11} is Q or T (SEQ ID NO:186), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5LFX_6X_7$, wherein X_1 is D or A, X_2 is A, G, E, R, Y, K, N, Q, L, or F, X_3 is A, L, P, or Y, X_4 is I or L, X_5 is R, A, Q, or S, X_6 is A or D, and X_7 is D, E, A, or S (SEQ ID NO:187); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of $X_1X_2SX_3X_4IX_5GX_6LN$, wherein X_1 is R or G, X_2 is A or T, X_3 is Q or E, X_4 is E, N, T, S, A, K, D, G, R, or Q, X_5 is Y or S, and X_6 is A

or V (SEQ ID NO:188), a CDR-L2 comprising the amino acid sequence of $GX_1X_2X_3LX_4X_5$, wherein X_1 is A or S, X_2 is T, S, E, Q, or D, X_3 is N, R, A, E, or H, X_4 is Q or A, and X_5 is S or D (SEQ ID NO:189), and a CDR-L3 comprising the amino acid sequence of $QX_1X_2X_3X_4X_5PWT$, wherein X_1 is S, N, D, Q, A, or E, X_2 is T, I, or S, X_3 is Y, L, or F, X_4 is D, G, T, E, Q, A, or Y, and X_5 is A, T, R, S, K, or Y (SEQ ID NO:190);

(j) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:200, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:201; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202;

(k) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(l) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202;

(m) the VH domain comprises a CDR-H1 comprising the amino acid sequence of X_1YX_2MS, wherein X_1 is S, D, E, A, or Q and X_2 is A, G, or T (SEQ ID NO:208), a CDR-H2 comprising the amino acid sequence of $DIX_1X_2X_3GX_4X_5TX_6YADSVKG$, wherein X_1 is T, N, S, Q, E, H, R, or A, X_2 is Y, W, F, or H, X_3 is A, S, Q, E, or T, X_4 is G or E, X_5 is S or I, and X_6 is A or G (SEQ ID NO:209), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3YX_4WX_5X_6AX_7DX_8$, wherein X_1 is S or A, X_2 is N, H, A, D, L, Q, Y, or R, X_3 is A, N, S, or G, X_4 is A, V, R, E, or S, X_5 is D or S, X_6 is D, N, Q, E, S, T, or L, X_7 is L, F, or M, and X_8 is I, Y, or V (SEQ ID NO:210); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178);

(n) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178);

(o) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:260, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); or

(p) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:252, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:253, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:254; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:255, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:256, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:257.

[0506] E56. The targeted cytokine construct of any one of embodiments 13- 40, wherein the antibody or antigen binding fragment comprises a heavy chain variable (VH) domain and a light chain variable (VL) domain; and wherein:

(a) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:226, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:151; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:154;

(b) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:228, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:157; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:158, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:159, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:160;

(c) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:163; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:164, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:165, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:166;

(d) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:229, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:169; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:170, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:171, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:172;

(e) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:230, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:231, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:175; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

(f) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:230, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:232, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:181; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:182, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:183, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:184;

(g) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:224, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:225; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:140, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:141, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:142;

(h) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:233, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:234, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:145; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:146, a CDR-L2 comprising the amino

acid sequence of SEQ ID NO:147, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:148;

(i) the VH domain comprises a CDR-H1 comprising the amino acid sequence of $GX_1X_2FX_3X_4X_5$, wherein X_1 is G, Y, S, or A, X_2 is T, S, G, R, N, or H, X_3 is S, T, R, H, Y, G, or P, X_4 is S, K, G, N, R, D, T, or G, and X_5 is Y, L, H, or F (SEQ ID NO:235), a CDR-H2 comprising the amino acid sequence of $X_1PX_2X_3X_4X_5$, wherein X_1 is I, N, or M, X_2 is G, N, H, S, R, I, or A, X_3 is A, N, H, S, T, F, or Y, X_4 is A, D, or G, and X_5 is T, E, K, V, Q, or A (SEQ ID NO:236), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5LFX_6X_7$, wherein X_1 is D or A, X_2 is A, G, E, R, Y, K, N, Q, L, or F, X_3 is A, L, P, or Y, X_4 is I or L, X_5 is R, A, Q, or S, X_6 is A or D, and X_7 is D, E, A, or S (SEQ ID NO:237); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of $X_1X_2SX_3X_4IX_5GX_6LN$, wherein X_1 is R or G, X_2 is A or T, X_3 is Q or E, X_4 is E, N, T, S, A, K, D, G, R, or Q, X_5 is Y or S, and X_6 is A or V (SEQ ID NO:188), a CDR-L2 comprising the amino acid sequence of $GX_1X_2X_3LX_4X_5$, wherein X_1 is A or S, X_2 is T, S, E, Q, or D, X_3 is N, R, A, E, or H, X_4 is Q or A, and X_5 is S or D (SEQ ID NO:189), and a CDR-L3 comprising the amino acid sequence of $QX_1X_2X_3X_4X_5PWT$, wherein X_1 is S, N, D, Q, A, or E, X_2 is T, I, or S, X_3 is Y, L, or F, X_4 is D, G, T, E, Q, A, or Y, and X_5 is A, T, R, S, K, or Y (SEQ ID NO:190);

(j) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:242, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202;

(k) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(l) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino

acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202;

(m) the VH domain comprises a CDR-H1 comprising the amino acid sequence of $GFTFX_1X_2Y$, wherein X_1 is S, D, E, Q, S, or A and X_2 is S, D, E, A, or Q (SEQ ID NO:244), a CDR-H2 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5$, wherein X_1 is T, N, S, Q, E, H, R or A, X_2 is Y, W, F, or H, X_3 is A, S, Q, E, or T, X_4 is G or E, and X_5 is S or I (SEQ ID NO:245), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3YX_4WX_5X_6AX_7DX_8$, wherein X_1 is S or A, X_2 is N, H, A, D, L, Q, Y, or R, X_3 is A, N, S, or G, X_4 is A, V, R, E, or S, X_5 is D or S, X_6 is D, N, Q, E, S, T, or L, X_7 is L, F, or M, and X_8 is I, Y, or V (SEQ ID NO:246); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178);

(n) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178);

(o) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:261, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); or

(p) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:224, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:284; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:285, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:286, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:287.

[0507] E57. The targeted cytokine construct of embodiment 55 or embodiment 56, wherein:

(a) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:109, and wherein the VL domain

comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:110;

(b) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:111, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:112;

(c) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:113, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:114;

(d) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:115, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:116;

(e) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:117, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:118;

(f) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:119, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:120;

(g) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:123; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:124;

(h) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:129, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:130;

(i) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:131; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:132;

(j) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:125; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:126;

(k) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:127, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:128;

(l) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:133; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:134;

(m) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:135; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:136;

(n) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:107; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:108; or

(o) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:121; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:122; or

(p) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:258; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:259.

[0508] E58. The targeted cytokine construct of any one of embodiments 13- 40, and 55-57, wherein the targeted cytokine construct comprises four polypeptide chains, wherein:

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:262, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:263, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:264, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:262;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:266, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:267, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:268, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:266;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:270, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:271, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:272, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 270;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:275, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:276, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274; or

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:278, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:279, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:280, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 278.

[0509] E59. The targeted cytokine construct of any one of embodiments 13- 40, and 55-57, wherein the targeted cytokine construct comprises four polypeptide chains, wherein:

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:262, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:263, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:265, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:262;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:266, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:267, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:269, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:266;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:270, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:271, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:273, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 270;

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:275, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:277, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274; or

the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:278, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:279, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:281, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 278.

EXAMPLES

Example 1: Design and biophysical characterization of exemplary IL-21 polypeptides and fusion proteins comprising the same

Materials and Methods

Recombinant DNA techniques

[0510] Techniques involving recombinant DNA manipulation, used for this study, were previously described in Sambrook *et al.*, Molecular cloning: A laboratory manual; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989. All reagents were used according to the manufacturer's instructions. DNA sequences were determined by double strand sequencing.

Gene synthesis

[0511] Desired gene segments were either generated by PCR using appropriate templates or synthesized at Genewiz (South Plainfield, NJ), Integrated DNA Technologies (Coralville, IA) or GeneScript (Piscataway, NJ) from synthetic oligonucleotides. The gene segments were cloned into the expression vectors using either Gibson assembly® method or using restriction digest followed by ligation. DNA was purified from transformed bacteria and concentration was determined by UV visible spectroscopy. DNA sequencing was used to confirm the DNA sequences of the subcloned gene fragments.

Isolation of antibody genes

[0512] Antibodies binding to CD8 antigens were generated using either humanization of mouse antibodies or *in vitro* phage display system.

[0513] For humanization, complementarity-determining regions (CDRs) of mouse residues were grafted into selected human framework(s) which exhibit close sequence similarity to the parental mouse framework and good stability. The resulting CDR-grafted antibodies were further humanized to remove any unnecessary non-human mutations.

[0514] For the *in vitro* display method, a non-immune human single chain Fv phage library generated from naïve B cells was panned for 5 to 6 rounds to isolate antibodies against the CD8 antigens. After the panning, individual phage clones that exhibited specific binding to target

antigen over non-specific antigens in ELISA were identified. DNA fragments of heavy and light chain V-domain of the specific binders were subsequently cloned and sequenced.

Cloning of antibody constructs

[0515] General information regarding the nucleotide sequences of human immunoglobulins light and heavy chains can be found in: IMGT® (the international ImMunoGeneTics information system®) from Lefranc et al. IMGT®, the international ImMunoGeneTics information system® 25 years on. Nucleic Acids Res. 2015 Jan;43. The DNA fragments of heavy and light chain V-domains were inserted in frame into the human IgG1 and Kappa light chain containing mammalian expression vector.

Cloning of fusion constructs

[0516] The IL-21 portions of the constructs were cloned in frame with the heavy chain using a $(G_4S)_3$ 15-mer linker between the C-terminus of the IgG heavy chain and the N-terminus of IL-21. The C-terminal lysine residue of the IgG heavy chain was eliminated after fusing the IL-21 portion. To generate the constructs with asymmetric geometry, knob-into-hole modification was introduced into the CH3 domains of the Fc region to facilitate heterodimerization. Specifically, the "hole" domain carried the S354C, T366S, L368A and Y407V mutations in the CH3 domain, whereas the "knob" domain carried the Y349C and T366W mutations in the CH3 domain (EU numbering). To abolish FcγR binding/effector function and prevent FcR co-activation, mutations L234A/L235A/G237A (EU numbering) were introduced into the CH2 domain of each of the IgG heavy chains or the Fc region. The expression of the antibody-IL-21 fusion constructs was driven by an CMV promoter and transcription terminated by a synthetic polyA signal sequence located downstream of the coding sequence.

Purification of fusion proteins with IL-21 polypeptides

[0517] Constructs encoding fusion proteins with IL-21 polypeptides as used in the examples were produced by co-transfecting exponentially growing Expi293 cells with the mammalian expression vectors using polyethylenimine (PEI). Supernatants were collected after 5 days of culture. IL-21 fusion constructs were first purified by affinity chromatography using a protein A matrix. The protein A column was equilibrated and washed in phosphate-buffered saline (PBS). The fusion constructs were eluted with 100 mM sodium citrate, pH 3.6 directly into 1 M Tris-HCl, pH 9 neutralization buffer. The eluted fractions were pooled and diluted into 20 mM MES, 25 mM sodium chloride pH 5.5 for further purification by ion-exchange chromatography (Mono-S, Cytiva)

to separate heterodimers from homodimers. After loading the protein, the column was washed with 20 mM MES, pH 5.5. The protein was then eluted with a gradient of sodium chloride from 25 mM NaCl to 500 mM NaCl in 20 mM MES, pH 5.5 buffer. The major eluent peak corresponding to the heterodimer was collected and analyzed by analytical size exclusion chromatography (Superdex 200 Increase 5/150 GL, Cytiva). For *in vitro* studies, the protein was dialyzed into PBS if the % monomer was ≥98%. Otherwise, and for all *in vivo* studies, the ion exchange eluent peak was polished by size exclusion chromatography (HiLoad Superdex 200 pg 16/600 or 26/600, Cytiva) in PBS.

[0518] The protein concentration of purified IL-21 fusion constructs was determined by measuring the optical density (OD) at 280 nm, using the molar extinction coefficient calculated on the basis of the amino acid sequence. Purity, integrity and monomeric state of the fusion constructs were analyzed by SDS-PAGE in the presence and absence of a reducing agent (5 mM 1,4-dithiothreitol) and stained with Coomassie blue (SimpleBlue™ SafeStain, Invitrogen). The NuPAGE® Pre-Cast gel system (Invitrogen) was used according to the manufacturer's instructions (4-20% Tris-glycine gels or 3-12% Bis-Tris). The aggregate content of immunoconjugate samples was analyzed using a Superdex 200 5/150 GL analytical size-exclusion column (Cytiva).

Binding affinity determination by surface plasmon resonance (SPR) for IL-21 fusion proteins with human IL-21R

[0519] Kinetic rate constants (k_a and k_d) as well as affinity (K_D) of unattenuated IL-21 fusion proteins with human IL-21R were measured by surface plasmon resonance (SPR) using a BIAcore® T200 (Cytiva) at 37° C. Briefly, to determine the affinities, IL-21 fusion proteins were diluted to 10 µg/mL and captured at 10 µL/min *via* their Fc onto a CM4 sensor chip by a covalently immobilized anti-human Fc antibody (Southern Biotech, Catalog No. 2014-01). Protein was not captured on flow cell 1 to serve as a reference surface. Human IL-21R (Biolegend, Catalog No. 773104), diluted in HBS-P+ buffer supplemented with 1 g/L BSA, at concentrations of 0.12, 0.37, 1.1, 3.3 and 10 nM, or buffer was flowed over the surface-captured fusion protein for 2 minutes at 30 µL/min. Dissociation was monitored for 10 minutes, and the anti-hIgG-Fc surface was regenerated with three 60-seconds injections of 75 mM phosphoric acid before recapturing fusion protein in each subsequent cycle. All interactions were measured in triplicate. Binding data were double-referenced and analyzed by Biacore® Evaluation Software version 3.2 using a 1:1 Langmuir with mass transport model to determine k_a, k_d and K_D using Biacore Evaluation Software.

[0520] Binding affinity (K_D) of attenuated IL-21 fusion proteins with human IL-21R were measured by surface plasmon resonance (SPR) using a BIAcore® T200 (Cytiva) at 37° C.

Recombinant human IL-21R was generated in house. Briefly, recombinant human IL-21R, comprised of the extracellular domain of IL-21R fused to 8× Histidine tag, were expressed in secreted form from HEK293 cells and purified by IMAC and size exclusion column chromatography. To determine the affinities, IL-21 fusion proteins were diluted to 10 µg/mL and captured at 10 µL/min *via* their Fc onto a CM4 sensor chip by a covalently immobilized anti-human Fc antibody (Southern Biotech). Protein was not captured on flow cell 1 to serve as a reference surface. A serial dilution was generated for human IL-21R, in HBS-P+ buffer supplemented with 1 g/L BSA, starting at nominal concentrations between 20-5 µM, followed by six to seven 5-fold serial dilutions, and flowed over the surface-captured fusion protein for 2 minutes at 30 µL/min. Buffer was also flowed over the surface to serve as a blank subtraction. Dissociation was monitored for 5-10 minutes, and the anti-hIgG-Fc surface was regenerated with three 30 to 60-seconds injections of 75 mM phosphoric acid before recapturing fusion protein in each subsequent binding cycle. Binding data were double-referenced and analyzed by Biacore® Evaluation Software version 3.2 using steady-state affinity for very weak interactions, or when possible, a 1:1 Langmuir with mass transport model to determine kinetic rate constants (k_a, k_d) and K_D.

Polyreactivity assessment by ELISA

[0521] In order to measure polyreactivity of candidate fusion proteins, an ELISA assay was used to check for binding to a panel of irrelevant antigens. The following were used as antigens and purchased from Sigma: keyhole limpet hemocyanin, hemoglobin, and heparin biotin sodium salt.

[0522] Antigens were diluted in PBS to a concentration of 5 µg/mL and coated onto a 384-well Nunc MaxiSorp plate (Thermo Fisher Scientific) at a volume of 25 µL per well. The plates were incubated overnight at 4°C. The antigens were removed, and the plate was washed with milli-Q water (Millipore). Wells were filled with PBS supplemented with 0.05% Tween and 1 mM EDTA (assay buffer) and then incubated at room temperature for 1 hour. The assay buffer was removed, and the wells washed with milli-Q water. 25 µL of 10 µg/mL of assay buffer only (negative control), bococizumab (positive control for polyreactivity), or fusion proteins, diluted in assay buffer were added and incubated at room temperature for 1 hour. Samples were removed and the plate was washed with milli-Q water. 25 µl of the detection antibody, 1:25000 diluted horseradish peroxidase conjugated goat anti-human IgG (Jackson ImmunoResearch), was added and allowed to incubate for 1 hour at room temperature. The reagent was removed, and wells were washed with milli-Q water. Wells were developed using 25 µL of KPL SureBlue TMB Microwell Substrate (SeraCare) for 5-7 mins and quenched with 25 µL of 0.1 M HCl. The absorbance at 450 nm was

recorded on a SpectraMax iD5 plate reader (Molecular Devices) and normalized against the assay buffer only control wells.

Results

Design of IL-21 charge variant polypeptides

[0523] Design of various exemplary IL-21 polypeptides is depicted in **FIG. 2**, which shows the amino acid sequence of wild-type IL-21 with "X" denoting the amino acid substituted in the sequence of wild-type IL-21 for another amino acid to generate an exemplary IL-21 polypeptide as described herein. The IL-21 polypeptides were charge variants with altered (*e.g.*, reduced) isoelectric point) relative to that of wild-type IL-21.

[0524] Increased surface charge or pI in protein-based molecules have been shown to correlate with increased systemic clearance, and decreased subcutaneous bioavailability (Boswell, CA *et al.* Bioconjug Chem. 21: 2153-2163, 2010; Herve, F *et al.* AAPS J. 10: 455-472, 2008; Zheng Y. MAbs 4: 243-255, 2012). Additionally, positive surface charge has been shown to correlate with decreased soluble expression and increased aggregation (Niwa, T, *et al.* PNAS, 106: 4201, 2009; Chan, P et al. Sci Reports, 3: 3333, 2013; Dudgeon K. et al, PNAS 109: 10879-84, 2012). In order to evaluate the impact of reducing positive surface charge on wild type IL-21, which has a theoretical pI of about 9.42, several exemplary charge variants were designed to remove positively charged surface residues at several locations predicted not to interfere with binding to the IL-21 receptor or common gamma chain. Specifically, several exemplary charge variant designs were made (Exemplary IL-21 polypeptide 1 to 31, also referred to as IL-21 v1 through IL-21 v31), amino acid substitutions to either glycine, serine, aspartic acid, alanine or glutamic acid, with sequences shown in **Table 2**. Theoretical pI values of the modified cytokine variants are also listed, as calculated by the ExPASy ProtParam tool.

Expression of IL-21 charge variant polypeptides as antibody fusions

[0525] Protein expression characteristics of fusion proteins of an exemplary anti-CD8 antibody, either xhCD8 or xhCD8.1, with the exemplary IL-21 charge variant polypeptides, exemplary IL-21 polypeptide 1 to 6, were compared to fusion proteins with wild type IL-21, and the results are summarized in **Table 3**. Yields are reported after an intermediate step, after Protein A, as well as after two-steps of purification, Protein A and ion exchange chromatography, for selected protein fusions. In the case of fusion proteins with exemplary IL-21 polypeptide 1, exemplary IL-21 polypeptide 2, and exemplary IL-21 polypeptides 6 through 31, yields were higher than for fusion

proteins with wild type IL-21 at both the Protein A intermediate stage, as well as the fully purified stage, after Protein A and ion exchange chromatography.

[0526] Ion exchange chromatography traces also show improvements for fusion proteins of all charge variants, exemplary IL-21 polypeptide, compared to fusion proteins of wild type IL-21, as shown in **FIG. 3A - 3E**. In all traces, two peaks were observed, one representing a homodimer, an undesired product comprising the antibody only without cytokine, and one representing a heterodimer, the desired antibody-cytokine fusion product. The elution of the heterodimer peak was observed to be earlier for all exemplary charge variant fusion proteins compared to the wild type IL-21 fusion protein, consistent with a decrease in isoelectric point and surface charge. **FIG. 3A** shows ion exchange chromatography traces for fusion proteins of xhCD8 and either wild type IL-21 or exemplary IL-21 polypeptides 1 through 6. Here, the portion of total protein that was the desired product was about 32% in the case of the wild type IL-21 fusion protein, and higher for fusion proteins of all IL-21 charge variants, ranging from about 40% to about 85%. In **FIGS. 3B** through **3E**, ion exchange chromatography traces are shown for fusion proteins of xhCD8.1 and exemplary IL-21 polypeptides 2 or 7 through 31. Here, the portion of total protein that was the desired product ranged from 77% to 88% for the charge-modified exemplary IL-21 polypeptides, which is higher than the portion of total protein of about 34% in the case of the fusion protein of xhCD8.1 and wild type IL-21 (**FIG. 3F).**

Kinetics characterization of wild type IL-21 compared to selected charge variants

[0527] Binding kinetics to IL-21R at 37°C were compared for wild type IL-21 (Fc-hIL21 WT) and selected charge variants exemplary IL-21 polypeptide 1 (xhCD8-hIL21 v1), exemplary IL-21 polypeptide 2 (xhCD8-Fc-hIL21 v2), and exemplary IL-21 polypeptide 6 (xhCD8-hIL21 v6) and exemplary IL-21 polypeptide 27 through 31 (xhCD8.1-hIL21 v27-31). For wild type IL-21, a fusion protein to the human Fc was used, whereas, for the charge variants, a fusion protein to an anti-CD8 antibody was used. Kinetic and affinity parameters derived from averaging data from 3 replicate experiments are shown in **Table 4**. The IL-21R binding affinities of exemplary IL-21 polypeptide 1, exemplary IL-21 polypeptide 2, exemplary IL-21 polypeptide 27 through 31, are within 2-fold of that of wild type IL-21, indicating that there was no effect on IL-21R affinity by the charge modifications in exemplary IL-21 polypeptide. In the case of exemplary IL-21 polypeptide 6, there was approximately a 3-fold change in affinity.

Polyreactivity of fusion proteins with wild type IL-21 versus charge variants

[0528] The results of a polyreactivity ELISA assay that measures binding of IL-21 fusion proteins to a panel of irrelevant proteins are shown in **Table 5**. Bococizumab, which has been shown to be polyreactive, is also included as a positive control. The polyreactivity of fusion protein with wild type IL-21, xhCD8.1-hIL21, was compared to that of fusion proteins comprising the same antibody, xhCD8.1, and IL-21 charge variants. Notably, binding to irrelevant antigens was reduced for all fusion proteins with the IL-21 charge variants compared to xhCD8.1-hIL21, indicating an improvement in polyreactivity for the charge variants.

Example 2: Pharmacokinetics of exemplary IL-21 charge variant polypeptide containing fusion proteins in mice

Materials and Methods

Mouse pharmacokinetics evaluation

[0529] Female C57BL/6 mice at 8 weeks of age were used for pharmacokinetic evaluation of fusion proteins containing exemplary IL-21 polypeptide 2 (xCtrl-hIL21 v2) or wild-type IL-21 (xCtrl-hIL21) polypeptide. The fusion proteins were: xCtrl -hIL21 and xCtrl-hIL21 v2, fusion proteins of a control (Ctrl) antibody that is non-binding to mouse proteins, with either wild type IL-21 or exemplary IL-21 polypeptide 2, respectively. The control antibody was xhCD8.1 which does not bind mouse CD8b. All animals were housed and handled in accordance with the Institutional Animal Care and Use Committee of Explora BioLabs. Proteins were administered either intravenously or subcutaneously at a dose of 1 mg/kg in groups of 2 mice. Tail vein bleeds were performed at 30 minutes, 1, 2 5, and optionally 7 days post-dose for the mice dosed intravenously, and at 2 hours, 1, 2 and 4 days post-dose for the mice dosed subcutaneously. At each time point, approximately 25 µL blood and serum was collected. Serum was stored at -20 °C until they were used for analysis.

ELISA evaluation of serum samples

[0530] Mouse serum concentration of the test molecules was measured by ELISA, using a standard curve generated by the proteins used for treatment. Briefly, 100 µL per well of 2 µg/mL of AffiniPure F(ab')$_2$ Fragment Donkey Anti-Human IgG, Fcγ fragment specific (Jackson ImmunoResearch) in PBS was coated onto a 96-well Nunc MaxiSorp plate (Thermo Fisher Scientific) overnight at 4 °C. Coating solution was removed, and the plates were blocked with 250 µL per well of PBS + 5% skim milk for 2 hours at room temperature. Plates were washed with PBS + 0.05% Tween-20, and either protein standard or diluted serum was added. For the protein standard, a serial dilution was generated starting at 0.03 µg/mL, followed by seven 3-fold serial

dilutions. Serum samples were diluted between 100-fold and 5000-fold, in order to keep the final readout within the range of the standard curve. All dilutions were in PBS + 0.5% BSA + 0.05% Tween-20, and 100 µL was used per/well. Serum samples and the protein standard were incubated at room temperature for 1-2 hours, followed by washing with PBS + 0.05% Tween-20. Next, 100 µL per well of detection antibody (Peroxidase AffiniPure F'ab')$_2$ Fragment Goat Anti-Human IgG, Fcγ fragment specific) at 1:3000 dilution in PBS + 0.5% BSA + 0.05% Tween-20, and the plates were incubated at room temperature for 45 minutes. The plates were developed using 100 µL of 1-step Ultra TMB-ELISA substrate solution (Thermo Fisher Scientific) for 1-3 minutes, and stopped with 100 µL of 2 M sulfuric acid. The absorbance at 450 nm was read on a SpectraMax iD5 plate reader (Molecular Devices). A nonlinear regression was used to fit the standard curve, and absorbance at 450 nm of diluted serum samples were interpolated to the standard curve using Prism software (GraphPad). All serum concentrations were then converted to nanomolar concentration.

Results

[0531] The pharmacokinetics results are shown in **FIGS. 4A-4D**. With both the intravenous dosing and the subcutaneous dosing, the fusion protein comprising the charge variant exemplary IL-21 polypeptide sequence exhibited greater exposure than fusion protein comprising the wild type IL-21 sequence. Based on area under the curve (AUC) calculations using the trapezoidal rule, there is about 2.5-fold improvement in AUC by IV dosing and 2.3 to 2.8-fold improvement in AUC by subcutaneous dosing for the IL-21 charge variant exemplary polypeptide 2 (xCtrl-hIL21 v2) over wild type IL-21 (xCtrl-hIL21). Additionally, there is about 3.5-fold improvement in AUC by subcutaneous dosing for the IL-21 charge variant exemplary polypeptide 31 (xCtrl-hIL21 v31) over wild type IL-21 (xCtrl-hIL21). When comparing the intravenously-dosed exposure of xCtrl-hIL21 v31 and xCtrl-hIL21, there is an AUC ratio of 4.2 (**FIG. 4D**), indicating a large improvement of the hIL21 v31 charge modifications over wild type hIL21.

[0532] Based on the kinetics reported in **Example 1**, there is negligible difference in the binding kinetics of wild-type IL-21 and either exemplary IL-21 polypeptide 2 or exemplary IL-21 polypeptide 31 to IL-21R. Therefore, the observed differences in exposure cannot be attributed to any differences in target-mediated drug disposition of fusion proteins comprising exemplary IL-21 polypeptide 2 or exemplary IL-21 polypeptide 31 and wild type IL-21. Additionally, the targeting (Ctrl) antibody in the two fusion proteins described in each of **FIGs. 4A-4D** is the same and does not bind any mouse protein. The data suggest that the reduction in positive surface charge in exemplary IL-21 polypeptide 2 or exemplary IL-21 polypeptide 31 is responsible for the observed improvements in exposure over wild type IL-21.

Example 3: Evaluation of STAT3 activation by IL-21 and fusion proteins

Materials and Methods

Evaluation of STAT3 activation in human primary cells

[0533] The ability of IL-21 to activate various immune cell subsets was determined in an assay measuring the phosphorylation of STAT3 by flow cytometry in either human PBMCs or human whole blood.

[0534] PBMCs were isolated from blood of healthy donors using Ficoll-Paque Plus (Cytiva) and red blood cells were lysed using ACK lysis buffer. PBMCs were resuspended in serum-free RPMI1640 media at 40 x 10^6 cells/mL and aliquoted into 96-well U-bottom plates (50 µL per well). IL-21 fusion proteins and control proteins including recombinant IL-21 and control fusion proteins, were diluted to desired concentrations and added to wells (50 µL added as 2x stimulus). Incubation was typically performed for 10-20 min at 37 °C. Antibodies that could not be applied following methanol permeabilization - CD8a (SK1, Biolegend), CD4 (RPA-T4, Biolegend), CD62L (DREG-56, Biolegend), and CD19 (HIB19, Biolegend) – were added directly to the wells immediately following stimulation and incubated on ice for 10 min. Staining was stopped with 100 µL ice cold 8% PFA (4% final) for 10 min on ice. Cells were washed 2x with wash buffer (2% FBS in PBS). Cells were permeabilized in 100 µL pre-chilled Phosflow Perm buffer III (BD Biosciences) for 45 minutes on ice. Cells were washed 2x with wash buffer and stained for 30-45min at 4°C with antibodies against: CD3 (UCHT1, BD Biosciences), CD45RO (UCHL1, Biolegend), HLA-DR (L243, Biolegend), perforin (dG9, BD Biosciences), and pSTAT3 pY705 (4/P-STAT3, BD Biosciences) diluted in permeabilization buffer (eBiosciences). Cells were then analyzed on a flow cytometer. Cell populations were defined as follows: CD8+ T cells: CD3+CD8+, CD4+ T cells: CD3+CD4+, B cells: CD19+, NK cells: CD3-CD19-HLA-DR-Perforin+. Data were expressed as percent pSTAT3 positive, and in some cases as pSTAT3 mean fluorescence intensity (MFI), and imported into GraphPad Prism.

[0535] For assays in human whole blood, 90 µL human blood was used per well in a 1mL 96 well deep plate well and prewarmed for 10 minutes at 37°C. IL-21 fusion proteins and control proteins, such as wild-type IL-21, and control fusion proteins were prepared and pre-warmed to 37°C at 10x strength. 10 µL of prewarmed 10x stimuli was added to each well, creating 100 µL total volume at 1x stimuli concentration. Incubation was typically performed for 25 min at 37°C. The stimulation was quenched by adding pre-fix antibody staining cocktail, vortexing briefly and incubating on ice for 10 minutes in the dark. The pre-fix staining cocktail contained TruStain FcX (Biolegend) and antibodies against: CD4 (RPA-T4, Biolegend), CD19 (HIB19, BD), CD56 (NCAM16.2, BD), CD16 (3G8, Biolegend), and CD8 (SK1, Biolegend). 900 µL pre-warmed Lyse Fix (BD) was

added to the sample wells and incubated at 37°C for 10 minutes. Cells were washed in pre-chilled wash buffer containing PBS + 0.5% bovine serum albumin and 2mM EDTA. Pre-chilled Perm Buffer III (BD) was added and incubated for 60 minutes at -20°C, followed by two washes in wash buffer and one wash in TFP Perm/Wash (BD). Cells were resuspended in 25 µL "post-methanol" staining cocktail prepared in TFP Perm/Wash containing antibodies against: CD3 (UCHT1, BD), CD14 (MΦP9, BD), CD11c (B-ly6, BD), HLA-DR (L243, Biolegend), and pSTAT3 pY705 (4/P-STAT3, BD Biosciences). Cells were incubated for 30 min at 4°C in the dark, then washed in TFP Perm/Wash buffer, followed by fixation in 100 µL 4% PFA for 10 minutes at room temperature. Cells were washed in wash buffer and analyzed on a flow cytometer. Data were expressed as percent pSTAT3 positive and imported into GraphPad Prism. Cell populations were defined as follows: CD8+ T cells: CD3+CD8+, CD4+ T cells: CD3+CD4+, B cells: CD19+, NK cells: CD3-CD19-CD16+CD56dim, myeloid cells: CD11c+ HLA-DR+.

Results

Evaluation of STAT3 activation by wild type IL-21

[0536] STAT3 activation profiles for different cell types in human PBMCs or whole blood by recombinant wild type human IL-21 are shown in **FIGs. 5A-5B**, respectively. CD8+ T cells, CD4+ T cells, NK and B cells are potently activated by recombinant IL-21 with similar EC_{50}s in the pM range and high maximal STAT3 activation.

Selective activation of CD8+ T cells by CD8-targeted IL-21charge variants

[0537] The concept of cis-targeting for selective activation of CD8+ T cells with CD8-targeted attenuated IL-21 is shown **FIG. 6**. Affinity attenuation of an IL-21 charge variant reduces activity on all cells expressing IL-21R. CD8-targeting enables the binding of the targeted fusion to CD8+ T cells and restores the IL-21 activity specifically on CD8+ T cells. In contrast, untargeted IL-21 mutein fusion showed weak to no activity on all IL-21R+ cell types, including CD8+ T cells.

[0538] STAT3 activation by fusion proteins comprising anti-CD8 antibody and either wild type IL-21 or exemplary IL-21 charge variant polypeptides were evaluated on CD8+ T cells. Data are shown in **FIG. 7**. Fusion proteins comprising anti-CD8 antibody and exemplary IL-21 polypeptide 1 (xhCD8-hIL21 v1), exemplary IL-21 polypeptide 2 (xhCD8-hIL21 v2) or exemplary IL-21 polypeptide 6 (xhCD8-hIL21 v6) was seen to mediate STAT3 activation with equivalent potency to that of fusion protein comprising anti-CD8 antibody and wild type IL-21 (xhCD8-hIL21).

[0539] An initial round of IL-21 muteins were first generated on the exemplary IL-21 polypeptide sequence 2 (SEQ ID NO: 2) as a background and tested for STAT3 activation in human PBMCs.

Table 6 summarizes the sequences of these muteins, along with the measured affinity constants using the methods described in **Example 1**. STAT3 activation profiles are shown for fusion proteins comprising anti-CD8 antibody, xhCD8, and IL-21 v2.1 (**FIG. 8A**), IL-21 v2.2 (**FIG. 8B**), IL-21 v2.3 (**FIG. 8C**), IL-21 v2.4 (**FIG. 8D**), IL-21 v2.5 (**FIG. 8E**), and IL-21 v2.6 (**FIG. 8F**). For all muteins, preferential STAT3 activation of CD8+ T cells was observed, as measured by EC_{50} as well as maximal STAT3 activation.

[0540] Additional mutations that attenuate binding IL-21R were designed on the background of exemplary IL-21 polypeptide sequence 2, resulting in IL-21 muteins, IL-21 v2.7 through IL-21 v2.59. These were constructed as fusion proteins with an alternate anti-CD8 antibody, xhCD8.1, which shares the same epitope as xhCD8. Sequence information and binding affinities of the muteins are described in **Table 6**. STAT3 activation profiles are shown for CD8 T cells in **FIGS. 9A, 9C, 9E,** and **9G,** and for CD4 T cells in **FIGS. 9B, 9D, 9F,** and **9H**.

[0541] Selected IL-21 muteins that attenuate IL-21R binding were then generated on the exemplary IL-21 polypeptide sequence 31 (SEQ ID NO:94 - SEQ ID NO:98) as a background and tested for STAT3 activation in whole human blood. **Table 7** summarizes the sequences of these IL-21 muteins, while **Table 8** summarizes example sequences for full length antibody-cytokine fusion proteins. STAT3 activation profiles in whole blood are shown for fusion proteins comprising anti-CD8 antibody, xhCD8.1, and IL-21 v31.4 (**FIG. 10A**), IL-21 v31.6 (**FIG. 10B**), IL-21 v31.23 (**FIG. 10C**), IL-21 v31.48 (**FIG. 10D**), IL-21 v31.51 (**FIG. 10E**). As a positive control, STAT3 activation of untargeted, unattenuated IL-21 fusion, Fc-IL-21v2 is shown in **FIG. 10F.** Additional data from whole blood of separate donors are shown in **FIGS. 14A-14F**, and reveals a consistent pattern of pSTAT3 activation across multiple donors.

[0542] Fusion proteins of exemplary IL-21 polypeptide sequences with an alternate anti-CD8 antibody, xhCD8v11, were constructed and evaluated in a whole human blood STAT3 activation assay. The results of these are shown in **FIGS. 15A-15D**, and reveals that all xhCD8v11 fusions selectively activate pSTAT3 signaling in CD8 T cells over other cell types.

Example 4: Construction and in vitro characterization of mouse surrogates of cis-targeted IL-21

Materials and Methods

Cloning of murine fusion constructs

[0543] Murine IL-21 portions of the constructs were cloned in frame with the heavy chain using a $(G_4S)_3$ 15-mer linker between the C-terminus of the mouse IgG2a heavy chain and the N-terminus of IL-21. The C-terminal lysine residue of the IgG heavy chain was eliminated after fusing the IL-

21 portion. To generate the constructs with asymmetric geometry, electrostatic modifications were introduced into the CH3 domains of the Fc region to facilitate heterodimerization. Specifically, the positively charged modifications carried the E356K and D399K mutations in the CH3 domain, whereas the negatively charged modifications carried the K409E and K439D mutations in the CH3 domain (EU numbering). To abolish FcγR binding/effector function and prevent FcR co-activation, mutations L234A/L235A/P329G (EU numbering) were introduced into the CH2 domain of each of the IgG heavy chains or the Fc region. The expression of the antibody-IL-21 fusion constructs was driven by an CMV promoter and transcription terminated by a synthetic polyA signal sequence located downstream of the coding sequence.

Evaluation of STAT3 activation in mouse splenocytes

[0544] The ability of IL-21 to activate various immune cell subsets was determined in an assay measuring the phosphorylation of STAT3 by flow cytometry in mouse splenocytes
Mouse spleens were mechanically dissociated in RPMI-1640 containing 10% FBS using the OctoMACS dissociation system (Miltenyi Biotech). Splenocytes were resuspended at 1-5 x 10^6 cells/ml in plain RPMI and rested for 1 hour at 37 °C with gentle agitation. Following rest, cells were resuspended in serum-free RPMI1640 media at 40 x 10^6 cells/mL and aliquoted into 96-well U-bottom plates (50 µL per well). IL-21 fusion proteins and control proteins including recombinant IL-21 (Biolegend) and control fusion proteins, were diluted to desired concentrations and added to wells (50 µL added as 2x stimulus) and incubated for 15 min at 37 °C. Antibodies that could not be applied following methanol permeabilization – Nkp46 (29A1.4, Biolegend), B220 (RA3-6B2, Biolegend) – were added directly to the wells immediately following stimulation and incubated on ice for 10 min. Cells were washed 3x in pre-chilled wash buffer containing PBS + 0.5% bovine serum albumin and 2mM EDTA and fixed with 100uL BD Fix/Perm buffer for 30 minutes at 4C in the dark. Cells were washed 3x in pre-chilled wash buffer and resuspended in 100 µL pre-chilled Phosflow Perm buffer III (BD Biosciences) for 60 minutes at -20°C. Cells were washed 2x with wash buffer and 1x with TFP permeabilization buffer (BD Biosciences) and stained for 30-45min at 4°C with antibodies against: CD4 (RM4-5, BD Biosciences), CD3 (17A2, Biolegend), CD8 (53-6.7, Biolegend), CD11b (M1/70, Biolegend), CD11c (N418, Biolegend) and pSTAT3 pY705 (4/P-STAT3, BD Biosciences) diluted in TFP permeabilization buffer. Cell populations were defined as follows: CD8+ T cells: CD3+CD8+, CD4+ T cells: CD3+CD4+, B cells: B220+, NK cells: CD3-Nkp4.6+. Data were expressed as percent pSTAT3 positive, and in some cases as pSTAT3 mean fluorescence intensity (MFI), and imported into GraphPad Prism.

Results

Design of engineered murine IL-21 muteins

[0545] Using the methods described in **Example 1**, initial attempts to express and purify fusion proteins with wild type murine IL-21 resulted in yields that were too low to generate usable material, less than 0.05 mg/L. Using the same design principles as for human IL-21, charge variants of murine IL-21 were designed to eliminate or reverse positive charge on a region of murine IL-21 that is expected to not affect binding to IL-21R or the gamma common chain. A fusion protein of xCtrl, a control antibody that does not bind any mouse proteins, and mIL-21 v1, a charge variant of murine IL-21 with sequence shown in **Table 9**, was successfully expressed and purified with final yield of approximately 8.6 mg/L after Protein A and ion exchange chromatography.

[0546] In order to create a CD8 cis-targeted murine surrogate, the same design principles were used as described in **Example 3** and **FIG. 6**. Mutations that attenuated binding affinity to mouse IL-21R were introduced to mIL-21v1, resulting in the murine IL-21 mutein, mIL-21 v1.1, with sequence shown in **Table 9**. The fusion protein of xmCD8, a mouse CD8 specific antibody, and mIL-21 v1.1 was expressed and purified with final yield of 36.9 mg/L after Protein A and ion exchange chromatography.

Selective activation of CD8+ T cells by murine CD8-targeted IL-21 charge variants

[0547] STAT3 activation profiles in mouse splenocytes are shown for wild type murine IL-21 (Biolegend) (**FIG. 11A**), xCtrl-mIL21 v1 (**FIG. 11B**), and xmCD8-mIL21 v1.1 (**FIG. 11C**). Wild type murine IL-21 and xCtrl-mIL21 v1 exhibit negligible selectivity of CD8+ T cell activation, whereas xmCD8-mIL21 v1.1 shows potent activation and >1000-fold selectivity on CD8+ T cells.

Example 5: In vivo efficacy of murine surrogates of cis-targeted IL-21

Materials and Methods

MC38 mouse tumor model

[0548] C57BL6 female mice (Jackson Labs) at 8-10 weeks of age were housed and acclimated at the vivarium facility. Cultured MC38 cells were harvested and resuspended in serum free media (1x DMEM, Sigma D6429) at $1.5x10^7$ cells/mL for implantation. Mice were shaved and $1.5x10^6$ cells (100 μL) are implanted subcutaneously above rear hind leg. Tumors were measured until tumor volume of 60-120mm3 was reached at approximately 8 days post implantation (TV=width*width*length*0.5). Mice were then randomized into groups by tumor volume. Where indicated, 1 mg/kg FTY720 (Sigma) was dosed intraperitoneally 3x weekly beginning at 6 days post-implantation (~2 days prior to therapeutic agent dosing). Following randomization, each

mouse was weighed and dosed subcutaneously with therapeutic agents under the scruff of the neck. Tumor volumes and body weights were measured every 3-4 days until either end of study (30-40 days post initial dose) or max tumor volume (2000 mm^3) was reached. At end of study mice were euthanized by CO_2 with appropriate secondary euthanasia.

Results

[0549] To the best of our knowledge, there has not been any reported single-agent efficacy of systemically administered untargeted mouse IL-21 in an MC38 mouse model. For example, three doses of approximately 10 mg/kg mouse IL-21 yielded 0/10 complete responses (CRs) in a previously reported study (Lewis et al. Oncoimmunology, Vol 7(1): 2018). Extension of cytokine half-life by fusion to an untargeted antibody also failed to control MC38 tumor growth with 0/5 CRs observed from three doses of approximately 2mg/kg antibody-IL-21 fusion (Deng et al. J Clin Invest Insight, Vol 5(7): 2020). **FIG. 12** shows potent efficacy with xCtrl-mIL21 v1, with 1/8 CRs and 5/8 CRs observed with a single dose of 0.3 mg/kg or 1 mg/kg, respectively. xCtrl-mIL21 v1 is differentiated from the construct used by Deng et al in that it is charge optimized. Based on the trends observed with human IL-21 charge variants in **Example 2**, it is speculated that xCtrl-mIL21 v1 would have improved bioavailability and exposure compared to the Deng et al construct, and these improved properties may be responsible for the observed efficacy in the MC38 tumor model.

[0550] CD8-targeted IL-21, xmCD8-mIL21 v1.1, also shows potent efficacy with 5/8 CRs, 7/8 CRs, and 8/8 CRs observed with single doses of 0.1, 0.3 and 1 mg/kg, respectively. When comparing efficacy in groups treated with dose-matched xCtrl-mIL21 v1, it is observed that xmCD8-IL21 v1.1 outperforms xmCtrl-mIL21 v1 in terms of the frequency of CRs as well as the degree of tumor growth inhibition. This suggests that there is an efficacy benefit to specifically targeting CD8+ T cells, and that pleiotropic IL-21 activation of cells other than CD8+ T cells by untargeted xCtrl-mIL21 v1 may inhibit the anti-tumor efficacy from IL-21 activity on CD8+ T cells.

[0551] FTY720 is a potent small molecule agonist of sphingosine 1-phosphate receptor molecule that blocks lymphocyte trafficking (Brinkmann et al Nat Rev Drug Disc, 2010). The effect of FTY720 on the efficacy of a single dose of xmCD8-mIL21 v1.1 was evaluated in the MC38 tumor model, with dosing schedule described in **FIG. 13A**. The tumor growth curves of 0.1, 0.3 or 1 mg/kg of xmCD8-mIL21 v1.1 with and without FTY720 treatment are shown in **FIGS. 13B, 13C** and **13D**, respectively. At the 0.1 mg/kg xmCD8-mIL21 v1.1 dose, less tumor growth inhibition and fewer CRs are observed when lymphocyte trafficking is blocked by FTY720 (**FIG. 13B**),

indicating that the curative response of 0.1 mg/kg xmCD8-mIL21 v1.1 is contributed by peripheral activation of CD8+ T cells.

Example 6: Evaluation of CD8-targeted IL-21 in PD1-resistant tumors and/or in combination with CD8-targeted IL2

[0552] The efficacy of CD8-targeted IL21 is evaluated in additional syngeneic mouse tumor models, using the methods previously described in **Example 5.**

[0553] In order to evaluate the efficacy of CD8-targeted IL-21 in tumor models that are PD-1 treatment resistant, efficacy is tested in a panel of syngeneic tumor models that are resistant to anti PD-1 treatment. Some examples of tumor models that are PD-1 resistant include B16F10, 4T1 and Pan02. CD8-targeted IL21 is administered either intravenously or subcutaneously with a single dose ranging from 0.01 to 5mg/kg and tumor growth will be measured.

[0554] In order to evaluate the efficacy of CD8-targeted IL-21 in combination with CD8-targeted IL2 (as described in international patent application nos. PCT/US20/36454 and PCT/US21/56312, each of which is hereby incorporated by reference in its entirety), efficacy is tested in panels of tumor models where CD8-targeted IL2 as an single agent is both efficacious and not efficacious. In tumor models where CD8-targeted IL2 is efficacious, the dose of CD8-targeted IL2 is reduced to suboptimal doses to give moderate to low single-agent efficacy. Combinations of the suboptimal doses of CD8-targeted IL2 are combined with various doses of CD8-targeted IL21 ranging from 0.01 to 5 mg/kg, and tumor growth is measured. In tumor models where CD8-targeted IL2 is not efficacious as a single agent, a moderate to high dose of CD8-targeted IL2 (e.g., 0.1-5 mg/kg) is used in combination with various doses of CD8-targeted IL21 ranging from 0.1 to 5 mg/kg. Control groups with PBS-treatment as well as single agent dosing is used for comparison. Simultaneous and staggered dosing schemes for CD8-targeted IL2 and CD8-targeted IL21 are also evaluated.

Example 7: Construction and evaluation of additional IL-21 variants with attenuated binding to IL-21R

Materials and Methods

Library construction and selection

[0555] Based on the published crystal structure of IL-21 cytokine in complex with IL-21R, amino acid residues on IL-21 were identified that do not make direct contacts with IL-21R but that may affect the structure of IL-21 in a manner that would attenuate binding affinity to IL-21R. Four different combinations of four residues were selected for diversification using NNK degenerate

codons to generate four independent libraries. Diversification was incorporated onto the IL-21 v2 background at the following sites: library 1 contained NNK diversity at positions V17, L20, L55, and L74; library 2 contained NNK diversity at positions I16, L20, L55 and L74; library 3 contained NNK diversity at positions I16, V17, L20 and L74; and library 4 contained NNK diversity at positions L13, I16, V17, and L74.

[0556] Using the IL-21 v2 sequence as a template, NNK-diversified libraries were constructed using PCR with primers containing degenerate nucleotides (Integrated DNA Technologies). PCR fragments were designed with flanking regions that match a linearized yeast display plasmid. Libraries were constructed according to published protocols for selection by yeast display (for example, Chao,G *et al.* Nat Protocols. 1: 755-768, 2006).

[0557] The four NNK libraries were displayed on yeast and sorted to isolate clones with desired affinity to IL21R. Controls hIL-21 v2.4 and hIL-21 v2.6 were also expressed clonally in a yeast display format. Overnight-induced yeast cells were labeled with mouse anti-human IL-21 (R&D Systems) and a recombinant human Fc fusion of heterodimeric IL-21R and common gamma chain (IL-21R-gc-Fc). After washing, cells were labeled with anti-mouse IgG Alexa488 and anti-human IgG Fc DyLight650 conjugates (Thermo). Two rounds of sorting were performed to enrich yeast populations with binding to IL-21R-gc-Fc between that of hIL-21 v2.4 and hIL-21 v2.6. After two rounds of sorting, plasmids were recovered from yeast by miniprep and transformed into E. coli to sequence individual clones.

Results

Characterization of additional IL-21 variants

[0558] Nine sequences of interest were identified and expressed as fusion proteins to xhCD8.1. IL-21 variant sequences are listed in **Table 6** as IL-21 v2.60 through IL-21 v2.65. The K_D's of binding to IL-21R of these variants were measured as described in **Example 1** and are listed in **Table 6**. Results shows that IL-21v2.61, IL-21v2.62, IL-21v2.63, IL-21v2.64, and IL-21v2.65, exhibit at least 50-fold reduction in binding to IL-21 receptors. The data here also shows that affinity attenuation to IL-21 receptors can be achieved by modifying non-contact residues.

As described in **Example 3**, variants with IL-21R affinity attenuation exhibit reduced activity on cells expressing IL-21R, whereas fusing the variants to a CD8-targeting domain enables preferential binding of the targeted fusions to CD8+ T cells and thereby restores the IL-21 activity specifically on CD8+ T cells. STAT3 activation in human whole blood was measured for selected fusion proteins of xhCD8.1 to additional IL-21 variants, hIL21 v2.63 (**FIG. 16A**), hIL21 v2.64

(**FIG. 16B**), and hIL21 v2.65 (**FIG. 16C**). In all cases shown, STAT3 is activated on CD8+ T cells with >100-fold selectively over non-CD8 T cells.

Example 8: Characterization of selected hIL-21 fusion proteins

[0559] Several hIL-21 fusion proteins were selected for additional characterization, namely xhCD8.1-hIL21 v31.4, xhCD8.1-hIL-21 v31.23, and xhCD8v11-hIL21 v31.23.

[0560] The yields of selected fusion proteins are shown in **Table 13** after ProA and after ProA+IEX purification. These yields are mostly higher than the charge-modified unattenuated fusion proteins reported in **Table 3**, which shows increased yield over the fusion with wild-type IL-21. Ion-exchange chromatography traces are shown in **FIGs. 17A-17C** with percentage of total signal specified that represents the desired heterodimer product. These results indicate that combination of IL-21R attenuation modifications and the charge variant modifications, v31, along with different targeting arms, xhCD8v11 or xhCD8.1, can maintain good expression properties with a high percentage of material being the correctly formed heterodimer.

[0561] Polyreactivity data for selected fusion proteins are shown in **Table 15** and reveal that these fusions exhibit favorable polyreactivity profiles as compared to positive control Bococizumab.

Example 9: Comparison of IL-21R attenuation muteins on wild type or v31 charge variant background

[0562] As shown in **Example 2,** the modification of charge in IL-21 reduces polyreactivity and improves bioavailability and exposure in mouse. Additionally, selected charge variants that were tested and reported in **Table 4** do not affect binding affinity to IL-21R. However, some of the mutations used for IL-21R attenuation either eliminates or reverses a positively charged residue. The effect of the charge reduction from an exemplary IL-21R attenuation mutein was evaluated by comparing the IL-21R attenuation mutein on either the wild type IL21 background, or the IL21 v31 charge variant background. Specifically, fusion proteins of hIL-21 v31.4 were compared to hIL-21 v0.4, which contains the same arginine to glutamic acid mutation for IL-21R attenuation as hIL-21 v31.4 but on the wild type IL-21 background. The sequence of hIL-21 v0.4 can be found in **Table 14.**

[0563] Fusion proteins of xhCD8.1 with either hIL-21 v0.4 or hIL-21 v31.4 were evaluated for polyreactivity. As shown in **Table 15** and **Table 5**, xhCD8.1-hIL21 v0.4 exhibits an improved polyreactivity profile compared to xhCD8.1-hIL21 (wild type IL-21 fusion), but the polyreactivity profile of xhCD8.1-hIL21 v31.4 is further improved with the incorporation of charge modifications (v31).

[0564] Highly polyreactive proteins exhibit reduced in vivo exposure due to non-specific interaction with cellular components. A pharmacokinetic study in mouse was set up to evaluate the impact of polyreactivity on the in vivo exposures of targeted fusions with and without v31 charge modification. Targeted fusions were generated using antibodies targeting human CD8, such as xhCD8.1 and xhCD8v11, which do not cross react with mouse CD8, therefore any observed exposure difference is primarily driven by the cytokine part of the fusion. Results in **FIG. 18** shows that xhCD8.1-hIL21 v31.4 exhibits 1.9-fold increase in area-under-the-curve over xhCD8.1-hIL21 v0.4, which lacks v31 charge modification and exhibit higher polyreactivity profile (**Table 15**). Such results suggest that charge modification such as v31can reduce polyreactivity and in turn enhance the in vivo exposure of IL-21 variants. Similarly, charge-modified xhCD8.1-hIL-21 v31.23, and xhCD8v11-hIL21 v31.23 show extended exposure in vivo over xhCD8.1-hIL21 v0.4, demonstrating the importance of the v31 charge modification for improved bioavailability and exposure.

Example 10: Characterization of recombinant (non-fusion) hIL-21 and hIL-21 variants

Materials and Methods

Expression and purification of non-fusion polypeptides

[0565] Recombinant (non-fusion) hIL-21 and hIL-21 variants were cloned into expression vectors and expressed as described in **Example 1**. A Gly-Gly-Ala-Ser linker, followed by 8His-tag was inserted downstream of the cytokine molecule for purification. Supernatants were collected after 5 days of expression in Expi293 cells (Thermo Fisher), and were used either directly for a Biacore assay or subjected to purification. IL-21 polypeptides were first purified by affinity chromatography using Ni Sepharose resin (Cytiva). The column was equilibrated and washed with PBS supplemented with 0.5 M NaCl. Proteins were eluted with PBS supplemented with 0.5 M NaCl and 0.25 M imidazole. The eluted fractions were polished by size exclusion chromatography (HiLoad Superdex 200 pg 16/600, Cytiva) in PBS.

Polyreactivity assessment of non-fusion polypeptides

[0566] For polyreactivity measurements of recombinant (non-fusion) hIL-21 and hIL-21 variants, an ELISA assay was performed as described in **Example 1** with the exception that detection was performed with 1:1000 diluted horseradish peroxidase conjugated mouse anti-His tag antibody (BioLegend).

Biacore analysis of non-fusion IL-21 variants

[0567] Kinetic rate constants (ka and kd) as well as binding affinity (KD) of non-fusion IL-21 muteins with human IL-21R were measured by surface plasmon resonance (SPR) using a BIAcore® 8K (Cytiva) at 25° C. To determine the affinities, a human IL-21 antibody (R&D Systems, Catalog No. MAB15001) diluted to 10 µg/mL was first captured at 10 µL/min for 2 minutes on flow cell 1 and 2 onto a CM5 sensor chip that was covalently immobilized with anti-mouse Fc antibody (Cytiva, Catalog No. BR100838). Next, IL-21 WT muteins supernatants were captured for 2 minutes at 10 µL/min on flow cell 2 with the human IL-21 antibody. No protein was captured on flow cell 1 to serve as a reference surface. A serial dilution was generated for the recombinant human IL-21R, that was generated in house as described earlier, in HBS-P+ buffer supplemented with 1 g/L BSA, starting at nominal concentrations of 20 µM, followed by five 5-fold serial dilutions, and flowed over the surface for 2 minutes at 30 µL/min to measure association. Buffer was also flowed over the surface to serve as a blank subtraction. Dissociation was monitored for 5 minutes, and the anti-mouse IgG-Fc surface was regenerated with 180 seconds injections of 10 mM glycine-HCl pH 1.7 before recapturing human IL-21 antibody/hIL-21 WT mutein protein in each subsequent binding cycle. Binding data were double-referenced and analyzed by Biacore® Insight Evaluation Software version 4.0.8 using steady-state affinity for very weak interactions, or when possible, a 1:1 Langmuir with mass transport model to determine kinetic rate constants (ka, kd) and KD.

Results

[0568] IL-21R attenuating mutations were incorporated on either the wild type hIL-21 or hIL-21 v31 background. The sequence of the cytokine portion, not inclusive of the downstream linker and 8His-tag, are shown in **Table 14** with affinities reported as well where measured.

[0569] Selected IL-21R muteins were selected for full purification. After preparative SEC, the fully-purified materials were re-analyzed by SDS-PAGE (**FIG. 19A**) and analytical SEC (**FIG. 19B**). Based on SDS-PAGE analysis of the fully-purified protein, all non-fusion hIL-21 and hIL-21 variants appear to be monomeric and pure, with no contaminating bands observed. Analytical SEC reveals an increase in peak width and retention time for all wild-type hIL-21 based constructs (hIL-21, hIL-21 v0.4, hIL-21 v0.23 & hIL-21 v0.51) compared to the hIL-21 v31 based constructs (hIL-21 v31.4, hIL-21 v31.23 & hIL-21 v31.51). This indicates that wild-type hIL-21 based constructs have more column interactions than hIL-21 v31 based constructs, and that the hIL-21 v31 based constructs possess better developability properties (Kohli et al., MAbs, 7(4): 752-758, 2015).

[0570] Polyreactivity was also measured for these fully purified constructs. The results are shown in **Table 16.** Overall, there was less polyreactivity observed with hIL-21 v31 compared to any of

the hIL-21 wild-type (v0) based variants, which suggests that the v31 charge modification alone improves polyreactivity more than any IL-21R attenuating mutation on the wild-type background.

Example 11: Exposure of recombinant (non-fusion) hIL-21 and hIL-21 variants

Materials and Methods

ELISA evaluation of serum samples

[0571] Mouse serum concentration of the test molecules was measured by ELISA, using a standard curve generated using known concentrations of test molecules. Briefly, 100 µL per well of 2 µg/mL of anti-human IL-21 antibody (R&D Systems, Catalog No. AF15001) in PBS was coated onto a 96-well Nunc MaxiSorp plate (Thermo Fisher Scientific) overnight at 4 °C. Coating solution was removed, and the plates were blocked with 250 µL per well of Blocker Casein in PBS (ThermoFisher Scientific) for 2 hours at room temperature. Plates were washed with PBS + 0.1% Tween-20, and either protein standard or diluted serum was added. For the protein standard, a serial dilution was generated starting at 0.00333 µg/mL, followed by four 3-fold serial dilutions. Serum samples were diluted between 50-fold and 2000-fold, in order to keep the final readout within the range of the standard curve. All dilutions were in PBS + 0.5% BSA + 0.05% Tween-20, and 100 µL was used per/well. Serum samples and the protein standard were incubated at room temperature for 1-2 hours, followed by washing with PBS + 0.1% Tween-20. Next, 100 µL per well of His Tag Biotinylated Antibody (R&D Systems, Catalog No. BAM050) at 10 nM was added and the plates were incubated at room temperature for 1 hour. Plates were washed with PBS + 0.1% Tween-20. Then, 100 µL per well of Peroxidase Streptavidin (Jackson ImmunoResearch, Laboratories Inc., Catalog no. 016-030-084) at 1:2000 dilution in PBS + 0.5% BSA + 0.05% Tween-20 was added, and the plates were incubated at room temperature for 45 minutes. The plates were developed using 100 µL of 1-step Ultra TMB-ELISA substrate solution (Thermo Fisher Scientific) for 1-3 minutes and stopped with 100 µL of 2 M sulfuric acid. The absorbance at 450 nm was read on a SpectraMax iD5 plate reader (Molecular Devices). A nonlinear regression was used to fit the standard curve, and absorbance at 450 nm of diluted serum samples were converted to concentrations by interpolating the standard curve.

Results

[0572] The exposure of human recombinant (non-fusion) wildtype IL-21 and IL-21 charge variant, v31, were evaluated in mouse. Due to the lack of significant difference in molecular weight between the two IL-21 molecules (~16kDa), renal filtration mediated clearance is not expected to contribute to any meaningful difference in exposure. Instead, the major difference in exposures

between wildtype IL-21 and IL-21v31 is mediated by the non-specific clearance, as v31 modification do not alter the IL-21R affinities (Table 14). Results in **FIG. 20** shows that v31 modification improves exposure compared to wild type IL-21, and a 4.3x increase in AUC ratio is observed. Overall, charge modification increases the bioavailability of recombinant IL-21 in non-fusion format, and this is consistent with the improvement observed in antibody fusion format (**FIG. 4D**).

Example 12: Measurement of suppression of T cell activation

Materials and Methods

[0573] *Activation of allogeneic T cells assay*

[0574] Immature human dendritic cells (DCs) were purchased (HemaCare - Charles River Laboratories) and activated for 3 days in GM-CSF, IL-4, and TNFa according to kit protocol (R&D Systems). Total T cells were isolated and enriched by a negative selection magnetic bead kit (Miltenyi Biotech). Activated DCs were harvested and plated at a ratio of 1:10 with T cells from mis-matched donors in the presence or absence of IL-21 molecules for 3 days. Molecules tested include two CD8+ T cell targeted fusions, xhCD8.1-hIL21 v31.4 and xhCD8.1-hIL21 v31.23, and an untargeted, unattenuated Fc-hIL21 v31. All three IL-21 molecules were tested at respective concentrations that give EC_{95} response in CD8 T cells in an in vitro pSTAT3 assay. Following completion of co-culture, supernatants were harvested and assessed for IL-2 secretion by ELISA following the manufacturing protocol (IL-2 instant ELISA, ThermoFisher Scientific).

Results

[0575] IL-21 can mediate suppressive effects on dendritic cells (DCs), inhibiting their maturation and inducing apoptosis of conventional DCs. Additionally, IL-21 can inhibit T cell priming in mixed lymphocyte cultures (Wan et al, Immunity, (2013) 38:514–27). The ability of CD8+ T cell targeted fusions to avoid the IL-21 mediated suppressive effect on dendritic cells was evaluated in co-cultures of DCs and T cells from mismatched human donors. In the absence of IL-21, incubation of alloreactive T cells and DCs induces the release of IL-2 in culture. **FIG. 22** shows that unattenuated Fc-hIL21 v31 inhibits IL-2 release, and therefore suppresses T cell activation. In contrast, both CD8+ T cell targeted fusions, xhCD8.1-hIL21 v31.4 and xhCD8.1-hIL21 v31.23, maintain similar level of IL-2 release as control, supporting that CD8+ T cell targeted IL-21 fusions do not interfere with the alloresponse of mismatched T cells and dendritic cells.

Example 13: Evaluation of CD8-targeted IL-21 fusions in cynomolgus monkeys

Materials and Methods

Evaluation of in vitro STAT3 activation in cynomolgus monkey blood cells

[0576] The activation of STAT3 in cynomolgus monkey whole blood by IL-21 molecules was similarly performed as described in Example 3, with the exception that cynomolgus monkey specific antibodies were used to stain various cell types in blood. Specifically, antibodies used were CD3 (SP34.2, BD), CD4 (L200, BD), CD8a (SK1, Biolegend), CD14 (M5E2, Biolegend), CD20 (2H7, Biolegend), HLA-DR (L243, Biolegend), NKG2A (REA110, Miltenyi) and pStat3 pY705 (4/P-STAT3, BD). Cell populations were defined as follows: CD8+ T cells: CD3+CD8+, CD4+ T cells: CD3+CD4+, B cells: CD20+, NK cells: CD3-CD19-NKG2A+.

Evaluation of in vivo STAT3 activation in cynomolgus monkey

[0577] Two naïve cynomolgus non-human primates (Macaca fascicularis) were dosed intravenously of xhCD8.1-IL-21 v31.4. Blood samples were collected 20 min post dose. STAT3 activation in various blood cell types was determined following the STAT3 protocol described in Example 3. The following set of antibodies were used to stain various cell types and phosphorylated STAT3: CD3 (SP34.2, BD), CD4 (L200, BD), CD8a (SK1, Biolegend), CD14 (M5E2, Biolegend), CD20 (2H7, Biolegend), HLA-DR (L243, Biolegend), NKG2A (REA110, Miltenyi) and pStat3 pY705 (4/P-STAT3, BD).

Results

[0578] Cynomolgus monkey blood were stimulated with various IL-21 molecules in vitro and STAT3 activation in various blood cell types were determined. Similar to the preferential activities demonstrated on human CD8+ T cells (**FIG. 14** and **15**), CD8+ T cell targeted IL-21 fusion proteins xhCD8.1-IL-21 v31.4 (**FIG. 23A**), xhCD8.1-IL-21 v31.23 (**FIG. 23B**) and xhCD8v11-IL-21 v31.23 (**FIG. 23C**) all show preferential STAT3 activation of CD8+ T cells over B cells and CD4+ T cells in cynomolgus monkey blood, as measured by EC_{50}. In contrast, untargeted, unattenuated Fc-hIL21 v31 (**FIG. 23D**) indiscriminately activates STAT3 in all cell types.

[0579] The in vivo selectivity of a CD8+T cell targeted fusion, xhCD8.1-IL-21 v31.4, was assessed. Cynomolgus monkeys were dosed with xhCD8.1-IL-21 v31.4 and in vivo STAT3 activation in various blood cell types was determined by directly staining for the phosphorylated STAT3 levels in collected blood samples. Similar to observed in vitro selectivity, **FIG. 24** shows that xhCD8.1-IL-21 v31.4 selectively and preferentially activate STAT3 in CD8+T cells over NK cells, B cells and CD4+ T cells in vivo.

[0580] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

Table 1: IL-21 Sequences

SEQ ID NO:	Name	Sequence
1	Wild-Type IL-21	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVET NCEWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPS TNAGRRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIH QHLSSRTHGSEDS
2	Exemplary IL-21 polypeptide sequence	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVET NCEWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPS TNAGGGQGHELTCPSCDSYEKKPPKEFLERFKSLLQKMIH QHLSSRTHGSEDS
3	Exemplary IL-21 scaffold sequence	QGQDX$_1$HMX$_2$X$_3$MX$_4$X$_5$LX$_6$X$_7$IVX$_8$X$_9$LKNX$_{10}$VNDLVPEFLPAPED VETNCEWSAFSCFQKAQLX$_{11}$SANTGNNEX$_{12}$IINVX$_{13}$IX$_{14}$X$_{15}$LX$_{16}$X$_{17}$X$_{18}$PPX$_{19}$X$_{20}$X$_{21}$X$_{22}$X$_{23}$X$_{24}$X$_{25}$X$_{26}$X$_{27}$X$_{28}$X$_{29}$X$_{30}$X$_{31}$CPSCDSYEKKP PKEFLERFKSLLX$_{32}$X$_{33}$MIHQHLSSRTHGSEDS
22	Exemplary IL-21 scaffold sequence	QGQDX$_1$HMX$_2$X$_3$MX$_4$X$_5$LX$_6$X$_7$IVX$_8$X$_9$LKNX$_{10}$VNDLVPEFLPAPED VETNCEWSAFSCFQKAQLKSANTGNNEX$_{11}$IINVX$_{12}$IX$_{13}$X$_{14}$LX$_{15}$X$_{16}$ X$_{17}$PPX$_{18}$TNAGRRQKHRLTCPSCDSYEKKPPKEFLERFKSLLX$_{19}$X$_{20}$ MIHQHLSSRTHGSEDS

Table 2: IL-21 Charge Variants

IL-21 Variant	Amino Acid Sequence	Theoretical pI	SEQ ID NO
Exemplary IL-21 polypeptide sequence 1	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTNAGGGQGHALTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.72	4
Exemplary IL-21 polypeptide sequence 2	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTNAGGGQGHELTCPSCDSYEKKPPK EFLERFKSLLQ KMIHQHLSSRTHGSEDS	8.39	2
Exemplary IL-21 polypeptide sequence 3	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLASANTGNNERIINVS IKKLARKPPSTNAGGGQGHALTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.87	5
Exemplary IL-21 polypeptide sequence 4	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLESANTGNNERIINVS IKKLARKPPSTNAGGGQGHALTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.12	6

IL-21 Variant	Amino Acid Sequence	Theoretical pI	SEQ ID NO
Exemplary IL-21 polypeptide sequence 5	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLASANTGNNERIINVS IKKLERKPPSTNAGGGQGHALTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.12	7
Exemplary IL-21 polypeptide sequence 6	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLASANTGNNERIINVS IKKLARKPPSTNAGGGQGHELTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.12	8
Exemplary IL-21 polypeptide sequence 7	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGSGEGSGGSLTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	9
Exemplary IL-21 polypeptide sequence 8	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGSGGGSGGELTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	10
Exemplary IL-21 polypeptide sequence 9	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTNAGGGSGGELTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	11
Exemplary IL-21 polypeptide sequence 10	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTNAGGGSGGSLTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.72	12
Exemplary IL-21 polypeptide sequence 11	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTNAGGGGGGELTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	13
Exemplary IL-21 polypeptide sequence 12	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTNAGGGGGGGLTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.72	14
Exemplary IL-21 polypeptide sequence 13	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTNAGGGEGGGLTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	15
Exemplary IL-21 polypeptide sequence 14	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGSGGGSGGGSGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.72	23
Exemplary IL-21 polypeptide sequence 15	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGGGGGGGGGGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.72	24
Exemplary IL-21 polypeptide sequence 16	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGSGGGEGGGSGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	25

IL-21 Variant	Amino Acid Sequence	Theoretical pI	SEQ ID NO
Exemplary IL-21 polypeptide sequence 17	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGEGGGEGGGSGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.85	26
Exemplary IL-21 polypeptide sequence 18	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTGGGSGGGSGGSCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.72	27
Exemplary IL-21 polypeptide sequence 19	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTGGGEGGGSGGSCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	28
Exemplary IL-21 polypeptide sequence 20	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGGGEGGGGGGGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	29
Exemplary IL-21 polypeptide sequence 21	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGSGGGSGGEGGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	30
Exemplary IL-21 polypeptide sequence 22	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGGGGGGGGEGGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	31
Exemplary IL-21 polypeptide sequence 23	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTGGGSGGGSEGSCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	32
Exemplary IL-21 polypeptide sequence 24	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTGGGGGGGEGGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	33
Exemplary IL-21 polypeptide sequence 25	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTNAGGGGGEGTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	34
Exemplary IL-21 polypeptide sequence 26	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTGGGGGGGEGTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	8.39	35
Exemplary IL-21 polypeptide sequence 27	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTGGGGEGGGEGTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.85	36
Exemplary IL-21 polypeptide sequence 28	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPSTGGGEGGGGEGTCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.85	37

IL-21 Variant	Amino Acid Sequence	Theoretical pI	SEQ ID NO
Exemplary IL-21 polypeptide sequence 29	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGGEGGGGGEGGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.85	38
Exemplary IL-21 polypeptide sequence 30	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGGEGGGGGGEGGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.85	39
Exemplary IL-21 polypeptide sequence 31	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAP EDVETNCEWSAFSCFQKAQLKSANTGNNERIINVS IKKLKRKPPGGEGGGGGGGEGGCPSCDSYEKKPPK EFLERFKSLLQKMIHQHLSSRTHGSEDS	7.85	40

Table 3: Intermediate and final yields for fusion proteins of IL-21 charge variants

IL-21 Mutein	Antibody fusion	Yield after Protein A (mg/L)	Yield after Protein A+IEX (mg/L)
WT	xhCD8	36	8
v1	xhCD8	61	21
v2	xhCD8	112	23
v3	xhCD8	35	ND*
v4	xhCD8	31	ND
v5	xhCD8	30	ND
v6	xhCD8	40	9
WT	xhCD8.1	32	5
v2	xhCD8.1	129	69
v7	xhCD8.1	132	67
v8	xhCD8.1	114	46
v9	xhCD8.1	120	65
v10	xhCD8.1	111	46
v11	xhCD8.1	126	59
v12	xhCD8.1	117	63
v13	xhCD8.1	153	78
v14	xhCD8.1	78	29
v15	xhCD8.1	87	22
v16	xhCD8.1	111	46
v17	xhCD8.1	127	57
v18	xhCD8.1	81	26
v19	xhCD8.1	97	33
v20	xhCD8.1	99	32
v21	xhCD8.1	90	34
v22	xhCD8.1	97	34
v23	xhCD8.1	130	40
v24	xhCD8.1	110	49
v25	xhCD8.1	119	43

v26	xhCD8.1	99	37
v27	xhCD8.1	123	43
v28	xhCD8.1	107	36
v29	xhCD8.1	97	39
v30	xhCD8.1	136	59
v31	xhCD8.1	116	59

* ND indicates that the measurement was not determined

Table 4: IL-21R binding kinetics of IL-21 charge variants

Construct	k_a ($x10^6$ M^{-1} s^{-1})	k_d ($x10^{-4}$ s^{-1})	K_D (pM)
Fc-hIL21 WT	5.0 ± 0.1	7.6 ± 0.3	154 ± 7
xhCD8-hIL21 v1	8.3 ± 0.5	9.7 ± 2.7	117 ± 33
xhCD8-hIL21 v2	8.9 ± 0.4	11 ± 1	122 ± 14
xhCD8-hIL21 v6	5.2 ± 0.2	25 ± 2	478 ± 53
xhCD8.1-hIL21 v27	13.1 ± 0.5	13.9 ± 0.2	106 ± 4
xhCD8.1-hIL21 v28	13.1 ± 0.6	13.9 ± 0.2	106 ± 5
xhCD8.1-hIL21 v29	12.7 ± 0.2	11.9 ± 0.2	94 ± 2
xhCD8.1-hIL21 v30	11.6 ± 0.2	10.7 ± 0.2	93 ± 2
xhCD8.1-hIL21 v31	12.0 ± 0.3	11.3 ± 0.2	94 ± 3

Table 5: Polyreactivity ELISA of IL-21 fusion proteins

Construct	Binding to KLH	Binding to heparin	Binding to hemoglobin
Bococi-zumab	47.13	15.58	29.12
xhCD8.1-hIL21	30.11	32.85	10.90
xhCD8.1-hIL21 v2	9.68	11.75	4.95
xhCD8.1-hIL21 v14	13.20	12.37	4.50
xhCD8.1-hIL21 v15	12.01	11.28	5.36
xhCD8.1-hIL21 v16	11.22	10.26	5.20
xhCD8.1-hIL21 v17	9.90	9.47	5.02
xhCD8.1-hIL21 v18	11.49	11.85	5.09
xhCD8.1-hIL21 v19	10.47	12.34	5.13

xhCD8.1-hIL21 v20	11.27	11.20	5.03
xhCD8.1-hIL21 v21	12.22	12.87	6.08
xhCD8.1-hIL21 v22	11.28	12.72	5.37
xhCD8.1-hIL21 v23	11.00	11.18	5.72
xhCD8.1-hIL21 v24	9.24	9.49	5.05
xhCD8.1-hIL21 v25	10.87	11.69	5.77
xhCD8.1-hIL21 v26	9.89	11.14	4.85
xhCD8.1-hIL21 v27	9.49	9.93	4.74
xhCD8.1-hIL21 v28	9.76	9.13	4.26
xhCD8.1-hIL21 v29	10.57	9.08	5.38
xhCD8.1-hIL21 v30	11.36	10.46	5.69
xhCD8.1-hIL21 v31	10.50	8.80	5.29

Table 6: IL-21 charge variants with reduced affinity to IL-21R

IL-21 Mutein	Amino Acid Sequence	K_D at 37°C (nM)	SEQ ID NO
IL-21 v2.1	QGQDRHMIRMDQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	ND	16
IL-21 v2.2	QGQDRHMIRMEQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	ND	17
IL-21 v2.3	QGQDRHMIRMRQLDDIVAQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQAMIHQHLSSRTHGSEDS	ND	18
IL-21 v2.4	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKEKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	>5000	19
IL-21 v2.5	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	ND	20
IL-21 v2.6	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKFKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	1700	21
IL-21 v2.7	QGQDRHMERMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	156	41

IL-21 v2.8	QGQDAHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	18.9	42
IL-21 v2.9	QGQDEHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	420	43
IL-21 v2.10	QGQDSHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	24.4	44
IL-21 v2.11	QGQDTHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	9.4	45
IL-21 v2.12	QGQDNHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	54.6	46
IL-21 v2.13	QGQDQHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	15.2	47
IL-21 v2.14	QGQDVHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	6.5	48
IL-21 v2.15	QGQDIHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	6.2	49
IL-21 v2.16	QGQDLHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	2.1	50
IL-21 v2.17	QGQDYHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	9.8	51
IL-21 v2.18	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKAKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	3180	52
IL-21 v2.19	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKNKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	2310	53
IL-21 v2.20	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKDKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	NB	54
IL-21 v2.21	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKSKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	3800	55
IL-21 v2.22	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKTKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	3030	56
IL-21 v2.23	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKQKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	2410	57
IL-21 v2.24	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKVKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	5180	58

IL-21 v2.25	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKIKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	2990	59
IL-21 v2.26	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKLKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	651	60
IL-21 v2.27	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKYKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	2150	61
IL-21 v2.28	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRAPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	0.693	62
IL-21 v2.29	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKREPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	2.8	63
IL-21 v2.30	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIAKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	0.419	64
IL-21 v2.31	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIEKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	3.4	65
IL-21 v2.32	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLARKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	0.236	66
IL-21 v2.33	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLERKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	1.3	67
IL-21 v2.34	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKALKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	2.0	68
IL-21 v2.35	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKELKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	449.0	69
IL-21 v2.36	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRAPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	125.0	70
IL-21 v2.37	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKREPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	601.0	71
IL-21 v2.38	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIAKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	66.9	72
IL-21 v2.39	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIEKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	614.0	73
IL-21 v2.40	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLARKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	54.7	74
IL-21 v2.41	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLERKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	104.0	75

IL-21 v2.42	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKALKFKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	NB	76
IL-21 v2.43	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKELKFKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	NB	77
IL-21 v2.44	QGQDRHMIAMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	3250	78
IL-21 v2.45	QGQDRHMIDMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	NB	79
IL-21 v2.46	QGQDRHMIEMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	NB	80
IL-21 v2.47	QGQDRHMIHMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	16.5	81
IL-21 v2.48	QGQDRHMISMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	4760	82
IL-21 v2.49	QGQDRHMITMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	>5000	83
IL-21 v2.50	QGQDRHMINMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	2880	84
IL-21 v2.51	QGQDRHMIGMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	4840	85
IL-21 v2.52	QGQDRHMIVMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	>5000	86
IL-21 v2.53	QGQDRHMIIMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	> 5000	87
IL-21 v2.54	QGQDRHMILMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	4690	88
IL-21 v2.55	QGQDRHMIYMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	394	89
IL-21 v2.56	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIAKLKFKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	156	90
IL-21 v2.57	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLAFKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	18.9	91
IL-21 v2.58	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKFAPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	420	92

IL-21 v2.59	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLEFKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	24.4	93
IL-21 v2.60	QGQDRHMIRMRQLIDIIDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKIKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	0.15	374
IL-21 v2.61	QGQDRHMIRMRQLIDAVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKFKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	11.9	375
IL-21 v2.62	QGQDRHMIRMRQLIDSIDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKVKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	13.6	376
IL-21 v2.63	QGQDRHMIRMRQLIDRIDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKIKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	1400	377
IL-21 v2.64	QGQDRHMIRMRQFIDAADQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKMKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	1050	378
IL-21 v2.65	QGQDRHMIRMRQRIDAIDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKIKRKPPSTNAGGGQGHE LTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	5910	379

ND indicates that the affinity was not determined
NB indicates no binding detected at highest concentration tested

Table 7: Additional IL-21 charge variants with reduced affinity to IL-21R

IL-21 Mutein	Amino Acid Sequence	SEQ ID NO
IL-21 v31.4	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKEKPPGGEGGGGGGGE GGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	94
IL-21 v31.6	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKFKPPGGEGGGGGGGE GGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	95
IL-21 v31.23	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKQKPPGGEGGGGGGGE GGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	96
IL-21 v31.48	QGQDRHMISMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPGGEGGGGGGGE GGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	97
IL-21 v31.51	QGQDRHMIGMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWS AFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPGGEGGGGGGGE GGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS	98

Table 8: Example antibody-cytokine fusion protein sequences

Name	*Light chain sequence*	*Heavy chain sequence (with IL-21 fusion)*	*Heavy chain sequence (without IL-21 fusion)*	*Heavy chain sequence (without IL-21 fusion) plus C-term lysine*
xhCD8.1-hIL21 v31.4	DIQMTQSPSSLS ASVGDRVTITCR ASQSIYGALNWY QQKPGKAPKLLI YGASNLQSGVPS RFSGSGSGTDFT LTISSLQPEDFA TYYCQSTYTAPW TFGGGTKVEIKR TVAAPSVFIFPP SDEQLKSGTASV VCLLNNFYPREA KVQWKVDNALQS GNSQESVTEQDS KDSTYSLSSTLT LSKADYEKHKVY ACEVTHQGLSSP VTKSFNRGEC (SEQ ID NO:262)	QVQLVQSGAEVKKPGSSV KVSCKASGGTFSSYAISW VRQAPGQGLEWMGGIIPG YATANYAQKFQGRVTITA DESTSTAYMELSSLRSED TAVYYCARDAAGIRLFAD WGQGTLVTVSSASTKGPS VFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSG LYSLSSVVTVPSSSLGTQ TYICNVNHKPSNTKVDKK VEPKSCDKTHTCPPCPAP EAAGAPSVFLFPPKPKDT LMISRTPEVTCVVVDVSH EDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAK GQPREPQVYTLPPCREEM TKNQVSLSCAVKGFYPSD IAVEWESNGQPENNYKTT PPVLDSDGSFFLVSKLTV DKSRWQQGNVFSCSVMHE ALHNHYTQKSLSLSPGSG GGGSGGGGSGGGGSQGQD RHMIRMRQLIDIVDQLKN YVNDLVPEFLPAPEDVET NCEWSAFSCFQKAQLKSA NTGNNERIINVSIKKLKE KPPGGEGGGGGGGEGGCP SCDSYEKKPPKEFLERFK SLLQKMIHQHLSSRTHGS EDS (SEQ ID NO: 263)	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPG (SEQ ID NO:264)	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPGK (SEQ ID NO:265)
xhCD8.1-hIL21 v31.6	DIQMTQSPSSLS ASVGDRVTITCR ASQSIYGALNWY QQKPGKAPKLLI YGASNLQSGVPS RFSGSGSGTDFT LTISSLQPEDFA TYYCQSTYTAPW TFGGGTKVEIKR TVAAPSVFIFPP SDEQLKSGTASV VCLLNNFYPREA KVQWKVDNALQS GNSQESVTEQDS KDSTYSLSSTLT LSKADYEKHKVY ACEVTHQGLSSP VTKSFNRGEC (SEQ ID NO:266)	QVQLVQSGAEVKKPGSSV KVSCKASGGTFSSYAISW VRQAPGQGLEWMGGIIPG YATANYAQKFQGRVTITA DESTSTAYMELSSLRSED TAVYYCARDAAGIRLFAD WGQGTLVTVSSASTKGPS VFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSG LYSLSSVVTVPSSSLGTQ TYICNVNHKPSNTKVDKK VEPKSCDKTHTCPPCPAP EAAGAPSVFLFPPKPKDT LMISRTPEVTCVVVDVSH EDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAK GQPREPQVYTLPPCREEM TKNQVSLSCAVKGFYPSD	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW

		IAVEWESNGQPENNYKTT PPVLDSDGSFFLVSKLTV DKSRWQQGNVFSCSVMHE ALHNHYTQKSLSLSPGSG GGGSGGGGSGGGGSQGQD RHMIRMRQLIDIVDQLKN YVNDLVPEFLPAPEDVET NCEWSAFSCFQKAQLKSA NTGNNERIINVSIKKLKF KPPGGEGGGGGGGEGGCP SCDSYEKKPPKEFLERFK SLLQKMIHQHLSSRTHGS EDS (SEQ ID NO:267)	LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPG (SEQ ID NO:268)	LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPGK (SEQ ID NO:269)
xhCD8.1-hIL21 v31.23	DIQMTQSPSSLS ASVGDRVTITCR ASQSIYGALNWY QQKPGKAPKLLI YGASNLQSGVPS RFSGSGSGTDFT LTISSLQPEDFA TYYCQSTYTAPW TFGGGTKVEIKR TVAAPSVFIFPP SDEQLKSGTASV VCLLNNFYPREA KVQWKVDNALQS GNSQESVTEQDS KDSTYSLSSTLT LSKADYEKHKVY ACEVTHQGLSSP VTKSFNRGEC (SEQ ID NO:270)	QVQLVQSGAEVKKPGSSV KVSCKASGGTFSSYAISW VRQAPGQGLEWMGGIIPG YATANYAQKFQGRVTITA DESTSTAYMELSSLRSED TAVYYCARDAAGIRLFAD WGQGTLVTVSSASTKGPS VFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSG LYSLSSVVTVPSSSLGTQ TYICNVNHKPSNTKVDKK VEPKSCDKTHTCPPCPAP EAAGAPSVFLFPPKPKDT LMISRTPEVTCVVVDVSH EDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAK GQPREPQVYTLPPCREEM TKNQVSLSCAVKGFYPSD IAVEWESNGQPENNYKTT PPVLDSDGSFFLVSKLTV DKSRWQQGNVFSCSVMHE ALHNHYTQKSLSLSPGSG GGGSGGGGSGGGGSQGQD RHMIRMRQLIDIVDQLKN YVNDLVPEFLPAPEDVET NCEWSAFSCFQKAQLKSA NTGNNERIINVSIKKLKQ KPPGGEGGGGGGGEGGCP SCDSYEKKPPKEFLERFK SLLQKMIHQHLSSRTHGS EDS (SEQ ID NO:271)	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPG (SEQ ID NO:272)	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPGK (SEQ ID NO:273)
xhCD8.1-hIL21 v31.48	DIQMTQSPSSLS ASVGDRVTITCR ASQSIYGALNWY QQKPGKAPKLLI YGASNLQSGVPS RFSGSGSGTDFT LTISSLQPEDFA TYYCQSTYTAPW TFGGGTKVEIKR TVAAPSVFIFPP SDEQLKSGTASV VCLLNNFYPREA	QVQLVQSGAEVKKPGSSV KVSCKASGGTFSSYAISW VRQAPGQGLEWMGGIIPG YATANYAQKFQGRVTITA DESTSTAYMELSSLRSED TAVYYCARDAAGIRLFAD WGQGTLVTVSSASTKGPS VFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSG LYSLSSVVTVPSSSLGTQ TYICNVNHKPSNTKVDKK	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG

	KVQWKVDNALQS GNSQESVTEQDS KDSTYSLSSTLT LSKADYEKHKVY ACEVTHQGLSSP VTKSFNRGEC (SEQ ID NO:274)	VEPKSCDKTHTCPPCPAP EAAGAPSVFLFPPKPKDT LMISRTPEVTCVVVDVSH EDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAK GQPREPQVYTLPPCREEM TKNQVSLSCAVKGFYPSD IAVEWESNGQPENNYKTT PPVLDSDGSFFLVSKLTV DKSRWQQGNVFSCSVMHE ALHNHYTQKSLSLSPGSG GGGSGGGGSGGGGSQGQD RHMISMRQLIDIVDQLKN YVNDLVPEFLPAPEDVET NCEWSAFSCFQKAQLKSA NTGNNERIINVSIKKLKR KPPGGEGGGGGGGEGGCP SCDSYEKKPPKEFLERFK SLLQKMIHQHLSSRTHGS EDS (SEQ ID NO:275)	LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPG (SEQ ID NO:276)	LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPGK (SEQ ID NO:277)
xhCD8.1-hIL21 v31.51	DIQMTQSPSSLS ASVGDRVTITCR ASQSIYGALNWY QQKPGKAPKLLI YGASNLQSGVPS RFSGSGSGTDFT LTISSLQPEDFA TYYCQSTYTAPW TFGGGTKVEIKR TVAAPSVFIFPP SDEQLKSGTASV VCLLNNFYPREA KVQWKVDNALQS GNSQESVTEQDS KDSTYSLSSTLT LSKADYEKHKVY ACEVTHQGLSSP VTKSFNRGEC (SEQ ID NO:278)	QVQLVQSGAEVKKPGSSV KVSCKASGGTFSSYAISW VRQAPGQGLEWMGGIIPG YATANYAQKFQGRVTITA DESTSTAYMELSSLRSED TAVYYCARDAAGIRLFAD WGQGTLVTVSSASTKGPS VFPLAPSSKSTSGGTAAL GCLVKDYFPEPVTVSWNS GALTSGVHTFPAVLQSSG LYSLSSVVTVPSSSLGTQ TYICNVNHKPSNTKVDKK VEPKSCDKTHTCPPCPAP EAAGAPSVFLFPPKPKDT LMISRTPEVTCVVVDVSH EDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAK GQPREPQVYTLPPCREEM TKNQVSLSCAVKGFYPSD IAVEWESNGQPENNYKTT PPVLDSDGSFFLVSKLTV DKSRWQQGNVFSCSVMHE ALHNHYTQKSLSLSPGSG GGGSGGGGSGGGGSQGQD RHMIGMRQLIDIVDQLKN YVNDLVPEFLPAPEDVET NCEWSAFSCFQKAQLKSA NTGNNERIINVSIKKLKR KPPGGEGGGGGGGEGGCP SCDSYEKKPPKEFLERFK SLLQKMIHQHLSSRTHGS EDS (SEQ ID NO:279)	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPG (SEQ ID NO:280)	QVQLVQSGAEVKKPG SSVKVSCKASGGTFS SYAISWVRQAPGQGL EWMGGIIPGYATANY AQKFQGRVTITADES TSTAYMELSSLRSED TAVYYCARDAAGIRL FADWGQGTLVTVSSA STKGPSVFPLAPSSK STSGGTAALGCLVKD YFPEPVTVSWNSGAL TSGVHTFPAVLQSSG LYSLSSVVTVPSSSL GTQTYICNVNHKPSN TKVDKKVEPKSCDKT HTCPPCPAPEAAGAP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSH EDPEVKFNWYVDGVE VHNAKTKPREEQYNS TYRVVSVLTVLHQDW LNGKEYKCKVSNKAL PAPIEKTISKAKGQP REPQVCTLPPSREEM TKNQVSLWCLVKGFY PSDIAVEWESNGQPE NNYKTTPPVLDSDGS FFLYSKLTVDKSRWQ QGNVFSCSVMHEALH NHYTQKSLSLSPGK (SEQ ID NO:281)
xhCD8v11-hIL21 v31.4	EIVLTQSPGTLS LSPGERATLSCR ASQSVSSNLAWY	EVQLVESGGGLVQPGGSL RLSCAASGFTFSSYAMSW VRQAPGKGLEWVSDITYA	EVQLVESGGGLVQPG GSLRLSCAASGFTFS SYAMSWVRQAPGKGL	EVQLVESGGGLVQPG GSLRLSCAASGFTFS SYAMSWVRQAPGKGL

	QQKPGQAPRLLI YGASSRATGIPD RFSGSGSGTDFT LTISRLEPEDFA VYYCQQYGSSPP VTFGQGTKVEIK RTVAAPSVFIFP PSDEQLKSGTAS VVCLLNNFYPRE AKVQWKVDNALQ SGNSQESVTEQD SKDSTYSLSSTL TLSKADYEKHKV YACEVTHQGLSS PVTKSFNRGEC (SEQ ID NO: 297)	GGSTAYADSVKGRFTISR DNAKNSLYLQMNSLRAED TAVYYCARSNAYAWDDAL DIWGQGTLVTVSSASTKG PSVFPLAPSSKSTSGGTA ALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQS SGLYSLSSVVTVPSSSLG TQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCP APEAAGAPSVFLFPPKPK DTLMISRTPEVTCVVVDV SHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRV VSVLTVLHQDWLNGKEYK CKVSNKALPAPIEKTISK AKGQPREPQVYTLPPCRE EMTKNQVSLSCAVKGFYP SDIAVEWESNGQPENNYK TTPPVLDSDGSFFLVSKL TVDKSRWQQGNVFSCSVM HEALHNHYTQKSLSLSPG SGGGGSGGGGSGGGGSQG QDRHMIRMRQLIDIVDQL KNYVNDLVPEFLPAPEDV ETNCEWSAFSCFQKAQLK SANTGNNERIINVSIKKL KEKPPGGEGGGGGGGEGG CPSCDSYEKKPPKEFLER FKSLLQKMIHQHLSSRTH GSEDS (SEQ ID NO: 298)	EWVSDITYAGGSTAY ADSVKGRFTISRDNA KNSLYLQMNSLRAED TAVYYCARSNAYAWD DALDIWGQGTLVTVS SASTKGPSVFPLAPS SKSTSGGTAALGCLV KDYFPEPVTVSWNSG ALTSGVHTFPAVLQS SGLYSLSSVVTVPSS SLGTQTYICNVNHKP SNTKVDKKVEPKSCD KTHTCPPCPAPEAAG APSVFLFPPKPKDTL MISRTPEVTCVVVDV SHEDPEVKFNWYVDG VEVHNAKTKPREEQY NSTYRVVSVLTVLHQ DWLNGKEYKCKVSNK ALPAPIEKTISKAKG QPREPQVCTLPPSRE EMTKNQVSLWCLVKG FYPSDIAVEWESNGQ PENNYKTTPPVLDSD GSFFLYSKLTVDKSR WQQGNVFSCSVMHEA LHNHYTQKSLSLSPG (SEQ ID NO: 299)	EWVSDITYAGGSTAY ADSVKGRFTISRDNA KNSLYLQMNSLRAED TAVYYCARSNAYAWD DALDIWGQGTLVTVS SASTKGPSVFPLAPS SKSTSGGTAALGCLV KDYFPEPVTVSWNSG ALTSGVHTFPAVLQS SGLYSLSSVVTVPSS SLGTQTYICNVNHKP SNTKVDKKVEPKSCD KTHTCPPCPAPEAAG APSVFLFPPKPKDTL MISRTPEVTCVVVDV SHEDPEVKFNWYVDG VEVHNAKTKPREEQY NSTYRVVSVLTVLHQ DWLNGKEYKCKVSNK ALPAPIEKTISKAKG QPREPQVCTLPPSRE EMTKNQVSLWCLVKG FYPSDIAVEWESNGQ PENNYKTTPPVLDSD GSFFLYSKLTVDKSR WQQGNVFSCSVMHEA LHNHYTQKSLSLSPG K (SEQ ID NO: 300)
xhCD8v11-hIL21 v31.6	EIVLTQSPGTLS LSPGERATLSCR ASQSVSSNLAWY QQKPGQAPRLLI YGASSRATGIPD RFSGSGSGTDFT LTISRLEPEDFA VYYCQQYGSSPP VTFGQGTKVEIK RTVAAPSVFIFP PSDEQLKSGTAS VVCLLNNFYPRE AKVQWKVDNALQ SGNSQESVTEQD SKDSTYSLSSTL TLSKADYEKHKV YACEVTHQGLSS PVTKSFNRGEC (SEQ ID NO: 301)	EVQLVESGGGLVQPGGSL RLSCAASGFTFSSYAMSW VRQAPGKGLEWVSDITYA GGSTAYADSVKGRFTISR DNAKNSLYLQMNSLRAED TAVYYCARSNAYAWDDAL DIWGQGTLVTVSSASTKG PSVFPLAPSSKSTSGGTA ALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQS SGLYSLSSVVTVPSSSLG TQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCP APEAAGAPSVFLFPPKPK DTLMISRTPEVTCVVVDV SHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRV VSVLTVLHQDWLNGKEYK CKVSNKALPAPIEKTISK AKGQPREPQVYTLPPCRE EMTKNQVSLSCAVKGFYP SDIAVEWESNGQPENNYK TTPPVLDSDGSFFLVSKL TVDKSRWQQGNVFSCSVM HEALHNHYTQKSLSLSPG SGGGGSGGGGSGGGGSQG QDRHMIRMRQLIDIVDQL KNYVNDLVPEFLPAPEDV ETNCEWSAFSCFQKAQLK	EVQLVESGGGLVQPG GSLRLSCAASGFTFS SYAMSWVRQAPGKGL EWVSDITYAGGSTAY ADSVKGRFTISRDNA KNSLYLQMNSLRAED TAVYYCARSNAYAWD DALDIWGQGTLVTVS SASTKGPSVFPLAPS SKSTSGGTAALGCLV KDYFPEPVTVSWNSG ALTSGVHTFPAVLQS SGLYSLSSVVTVPSS SLGTQTYICNVNHKP SNTKVDKKVEPKSCD KTHTCPPCPAPEAAG APSVFLFPPKPKDTL MISRTPEVTCVVVDV SHEDPEVKFNWYVDG VEVHNAKTKPREEQY NSTYRVVSVLTVLHQ DWLNGKEYKCKVSNK ALPAPIEKTISKAKG QPREPQVCTLPPSRE EMTKNQVSLWCLVKG FYPSDIAVEWESNGQ PENNYKTTPPVLDSD GSFFLYSKLTVDKSR	EVQLVESGGGLVQPG GSLRLSCAASGFTFS SYAMSWVRQAPGKGL EWVSDITYAGGSTAY ADSVKGRFTISRDNA KNSLYLQMNSLRAED TAVYYCARSNAYAWD DALDIWGQGTLVTVS SASTKGPSVFPLAPS SKSTSGGTAALGCLV KDYFPEPVTVSWNSG ALTSGVHTFPAVLQS SGLYSLSSVVTVPSS SLGTQTYICNVNHKP SNTKVDKKVEPKSCD KTHTCPPCPAPEAAG APSVFLFPPKPKDTL MISRTPEVTCVVVDV SHEDPEVKFNWYVDG VEVHNAKTKPREEQY NSTYRVVSVLTVLHQ DWLNGKEYKCKVSNK ALPAPIEKTISKAKG QPREPQVCTLPPSRE EMTKNQVSLWCLVKG FYPSDIAVEWESNGQ PENNYKTTPPVLDSD GSFFLYSKLTVDKSR WQQGNVFSCSVMHEA

		SANTGNNERIINVSIKKL KFKPPGGEGGGGGGGEGG CPSCDSYEKKPPKEFLER FKSLLQKMIHQHLSSRTH GSEDS (SEQ ID NO: 302)	WQQGNVFSCSVMHEA LHNHYTQKSLSLSPG (SEQ ID NO: 303)	LHNHYTQKSLSLSPG K (SEQ ID NO: 304)
xhCD8v11-hIL21 v31.23	EIVLTQSPGTLS LSPGERATLSCR ASQSVSSNLAWY QQKPGQAPRLLI YGASSRATGIPD RFSGSGSGTDFT LTISRLEPEDFA VYYCQQYGSSPP VTFGQGTKVEIK RTVAAPSVFIFP PSDEQLKSGTAS VVCLLNNFYPRE AKVQWKVDNALQ SGNSQESVTEQD SKDSTYSLSSTL TLSKADYEKHKV YACEVTHQGLSS PVTKSFNRGEC (SEQ ID NO: 305)	EVQLVESGGGLVQPGGSL RLSCAASGFTFSSYAMSW VRQAPGKGLEWVSDITYA GGSTAYADSVKGRFTISR DNAKNSLYLQMNSLRAED TAVYYCARSNAYAWDDAL DIWGQGTLVTVSSASTKG PSVFPLAPSSKSTSGGTA ALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQS SGLYSLSSVVTVPSSSLG TQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCP APEAAGAPSVFLFPPKPK DTLMISRTPEVTCVVVDV SHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRV VSVLTVLHQDWLNGKEYK CKVSNKALPAPIEKTISK AKGQPREPQVYTLPPCRE EMTKNQVSLSCAVKGFYP SDIAVEWESNGQPENNYK TTPPVLDSDGSFFLVSKL TVDKSRWQQGNVFSCSVM HEALHNHYTQKSLSLSPG SGGGGSGGGGSGGGGSQG QDRHMIRMRQLIDIVDQL KNYVNDLVPEFLPAPEDV ETNCEWSAFSCFQKAQLK SANTGNNERIINVSIKKL KQKPPGGEGGGGGGGEGG CPSCDSYEKKPPKEFLER FKSLLQKMIHQHLSSRTH GSEDS (SEQ ID NO: 306)	EVQLVESGGGLVQPG GSLRLSCAASGFTFS SYAMSWVRQAPGKGL EWVSDITYAGGSTAY ADSVKGRFTISRDNA KNSLYLQMNSLRAED TAVYYCARSNAYAWD DALDIWGQGTLVTVS SASTKGPSVFPLAPS SKSTSGGTAALGCLV KDYFPEPVTVSWNSG ALTSGVHTFPAVLQS SGLYSLSSVVTVPSS SLGTQTYICNVNHKP SNTKVDKKVEPKSCD KTHTCPPCPAPEAAG APSVFLFPPKPKDTL MISRTPEVTCVVVDV SHEDPEVKFNWYVDG VEVHNAKTKPREEQY NSTYRVVSVLTVLHQ DWLNGKEYKCKVSNK ALPAPIEKTISKAKG QPREPQVCTLPPSRE EMTKNQVSLWCLVKG FYPSDIAVEWESNGQ PENNYKTTPPVLDSD GSFFLYSKLTVDKSR WQQGNVFSCSVMHEA LHNHYTQKSLSLSPG (SEQ ID NO: 307)	EVQLVESGGGLVQPG GSLRLSCAASGFTFS SYAMSWVRQAPGKGL EWVSDITYAGGSTAY ADSVKGRFTISRDNA KNSLYLQMNSLRAED TAVYYCARSNAYAWD DALDIWGQGTLVTVS SASTKGPSVFPLAPS SKSTSGGTAALGCLV KDYFPEPVTVSWNSG ALTSGVHTFPAVLQS SGLYSLSSVVTVPSS SLGTQTYICNVNHKP SNTKVDKKVEPKSCD KTHTCPPCPAPEAAG APSVFLFPPKPKDTL MISRTPEVTCVVVDV SHEDPEVKFNWYVDG VEVHNAKTKPREEQY NSTYRVVSVLTVLHQ DWLNGKEYKCKVSNK ALPAPIEKTISKAKG QPREPQVCTLPPSRE EMTKNQVSLWCLVKG FYPSDIAVEWESNGQ PENNYKTTPPVLDSD GSFFLYSKLTVDKSR WQQGNVFSCSVMHEA LHNHYTQKSLSLSPG K (SEQ ID NO: 308)
xhCD8v11-hIL21 v31.51	EIVLTQSPGTLS LSPGERATLSCR ASQSVSSNLAWY QQKPGQAPRLLI YGASSRATGIPD RFSGSGSGTDFT LTISRLEPEDFA VYYCQQYGSSPP VTFGQGTKVEIK RTVAAPSVFIFP PSDEQLKSGTAS VVCLLNNFYPRE AKVQWKVDNALQ SGNSQESVTEQD SKDSTYSLSSTL TLSKADYEKHKV YACEVTHQGLSS PVTKSFNRGEC	EVQLVESGGGLVQPGGSL RLSCAASGFTFSSYAMSW VRQAPGKGLEWVSDITYA GGSTAYADSVKGRFTISR DNAKNSLYLQMNSLRAED TAVYYCARSNAYAWDDAL DIWGQGTLVTVSSASTKG PSVFPLAPSSKSTSGGTA ALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQS SGLYSLSSVVTVPSSSLG TQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCP APEAAGAPSVFLFPPKPK DTLMISRTPEVTCVVVDV SHEDPEVKFNWYVDGVEV HNAKTKPREEQYNSTYRV VSVLTVLHQDWLNGKEYK CKVSNKALPAPIEKTISK AKGQPREPQVYTLPPCRE	EVQLVESGGGLVQPG GSLRLSCAASGFTFS SYAMSWVRQAPGKGL EWVSDITYAGGSTAY ADSVKGRFTISRDNA KNSLYLQMNSLRAED TAVYYCARSNAYAWD DALDIWGQGTLVTVS SASTKGPSVFPLAPS SKSTSGGTAALGCLV KDYFPEPVTVSWNSG ALTSGVHTFPAVLQS SGLYSLSSVVTVPSS SLGTQTYICNVNHKP SNTKVDKKVEPKSCD KTHTCPPCPAPEAAG APSVFLFPPKPKDTL MISRTPEVTCVVVDV SHEDPEVKFNWYVDG	EVQLVESGGGLVQPG GSLRLSCAASGFTFS SYAMSWVRQAPGKGL EWVSDITYAGGSTAY ADSVKGRFTISRDNA KNSLYLQMNSLRAED TAVYYCARSNAYAWD DALDIWGQGTLVTVS SASTKGPSVFPLAPS SKSTSGGTAALGCLV KDYFPEPVTVSWNSG ALTSGVHTFPAVLQS SGLYSLSSVVTVPSS SLGTQTYICNVNHKP SNTKVDKKVEPKSCD KTHTCPPCPAPEAAG APSVFLFPPKPKDTL MISRTPEVTCVVVDV SHEDPEVKFNWYVDG

	(SEQ ID NO: 309)	EMTKNQVSLSCAVKGFYP SDIAVEWESNGQPENNYK TTPPVLDSDGSFFLVSKL TVDKSRWQQGNVFSCSVM HEALHNHYTQKSLSLSPG SGGGGSGGGGSGGGGSQG QDRHMIGMRQLIDIVDQL KNYVNDLVPEFLPAPEDV ETNCEWSAFSCFQKAQLK SANTGNNERIINVSIKKL KRKPPGGEGGGGGGGEGG CPSCDSYEKKPPKEFLER FKSLLQKMIHQHLSSRTH GSEDS (SEQ ID NO: 310)	VEVHNAKTKPREEQY NSTYRVVSVLTVLHQ DWLNGKEYKCKVSNK ALPAPIEKTISKAKG QPREPQVCTLPPSRE EMTKNQVSLWCLVKG FYPSDIAVEWESNGQ PENNYKTTPPVLDSD GSFFLYSKLTVDKSR WQQGNVFSCSVMHEA LHNHYTQKSLSLSPG (SEQ ID NO: 311)	VEVHNAKTKPREEQY NSTYRVVSVLTVLHQ DWLNGKEYKCKVSNK ALPAPIEKTISKAKG QPREPQVCTLPPSRE EMTKNQVSLWCLVKG FYPSDIAVEWESNGQ PENNYKTTPPVLDSD GSFFLYSKLTVDKSR WQQGNVFSCSVMHEA LHNHYTQKSLSLSPG K (SEQ ID NO: 312)

Table 9: Engineered mouse IL-21 muteins

Mouse IL-21 Mutein	Amino Acid Sequence	SEQ ID NO
mIL-21 v1	QGPDRLLIRLRHLIDIVEQLKIYENDLDPELLSAPQDVKGHCEHAAFACFQKA KLKPSNPGNNKTFIIDLVAQLERRLPAGEGGEGQEHIAKCPSCDSYEKRTPKE FLERLKWLLQKMIHQHLS	282
mIL-21 v1.1	QGPDRLLIRLRHLIDIVEQLKIYENDLDPELLSAPQDVKGHCEHAAFACFQKA KLKPSNPGNNKTFIIDLVAQLEERLPAGEGGEGQEHIAKCPSCDSYEKRTPKE FLERLKWLLQKMIHQHLS	283

Table 10. Anti-CD8 antibody CDRs (Kabat)

Name	CDR-H1	CDR-H2	CDR-H3	CDR-L1	CDR-L2	CDR-L3
xhCD8	KYTMH (SEQ ID NO:137)	HFNPNNDETKYNQKFTG (SEQ ID NO:138)	DGLGLRLFAD (SEQ ID NO:139)	GASENIYGALN (SEQ ID NO:140)	GATNLAD (SEQ ID NO:141)	QNILDTPWT (SEQ ID NO:142)
xhCD8v1	KYAIS (SEQ ID NO:252)	HFNPNNDETKYNQKFQG (SEQ ID NO:253)	DGLGLRLFAD (SEQ ID NO:254)	RASENIYGALN (SEQ ID NO:255)	GATNLAD (SEQ ID NO:256)	QNILDTPWT (SEQ ID NO:257)
xhCD8v2	NFAIS (SEQ ID NO:149)	GIIPGHAKANYAQKFQG (SEQ ID NO:150)	DGLGIRLFAD (SEQ ID NO:151)	RASQEIYGALN (SEQ ID NO:152)	GATNLQS (SEQ ID NO:153)	QDIYDAPWT (SEQ ID NO:154)
xhCD8v3	KFAIS (SEQ ID NO:155)	GIIPGHAKANYAQKFQG (SEQ ID NO:156)	DGLGIRLFAD (SEQ ID NO:157)	RASQEIYGALN (SEQ ID NO:158)	GATNLQS (SEQ ID NO:159)	QDIYDAPWT (SEQ ID NO:160)
xhCD8v4	KYAIS	GIIPGHAKANYAQKFQG	DGLGIRLFAD	RASQKIYGALN	GATNLQS	QNTYDTPWT

	(SEQ ID NO:161)	(SEQ ID NO:162)	(SEQ ID NO:163)	(SEQ ID NO:164)	(SEQ ID NO:165)	(SEQ ID NO:166)
xhCD8v5	GHAIS (SEQ ID NO:167)	GIIPGHAKAN YAQKFQG (SEQ ID NO:168)	DGLGIRLFAD (SEQ ID NO:169)	RASQKIYGAL N (SEQ ID NO:170)	GATNLQS (SEQ ID NO:171)	QNTYDTPW T (SEQ ID NO:172)
xhCD8v6	DYGMS (SEQ ID NO:173)	DINWSGEIT AYADSVKG (SEQ ID NO:174)	SNSYRWDD ALDI (SEQ ID NO:175)	RASQSVSSN LA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T (SEQ ID NO:178)
xhCD8v7	DYAMH (SEQ ID NO:179)	VISYDGSNKY YADSVKG (SEQ ID NO:180)	DRIGWYDYD AFDI (SEQ ID NO:181)	RASHSVGSN LA (SEQ ID NO:182)	DASNRAT (SEQ ID NO:183)	QQRSNWPP T (SEQ ID NO:184)
xhCD8v8	SYWMN (SEQ ID NO:143)	QIYPGDGDT NYNGKFKG (SEQ ID NO:144)	SGAAFSSYYA MDY (SEQ ID NO:145)	RASENIYSNL A (SEQ ID NO:146)	AATNLAD (SEQ ID NO:147)	QHFWGTPW T (SEQ ID NO:148)
xhCD8v9	SYAIS (SEQ ID NO:199)	GIIPGAATAN YAQKFQG (SEQ ID NO:200)	DAAGIRLFA D (SEQ ID NO:201)	RASQEIYGAL N (SEQ ID NO:152)	GATNLQS (SEQ ID NO:153)	QSTYDAPWT (SEQ ID NO:202)
xhCD8v10	SYAMS (SEQ ID NO:220)	DITYAGGSTA YADSVKG (SEQ ID NO:221)	SNAYAWDD ALDI (SEQ ID NO:222)	RASQSVSSN LA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T (SEQ ID NO:178)
xhCD8v11	SYAMS (SEQ ID NO:220)	DITYAGGSTA YADSVKG (SEQ ID NO:221)	SNAYAWDD ALDI (SEQ ID NO:222)	RASQSVSSN LA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T (SEQ ID NO:178)
xhCD8v12 (xhCD8.1)	SYAIS (SEQ ID NO:199)	GIIPGYATAN YAQKFQG (SEQ ID NO:203)	DAAGIRLFA D (SEQ ID NO:204)	RASQSIYGAL N (SEQ ID NO:205)	GASNLQS (SEQ ID NO:206)	QSTYTAPWT (SEQ ID NO:207)
xhCD8v13	SYAIS (SEQ ID NO:199)	GIIPGYATAN YAQKFQG (SEQ ID NO:203)	DAAGIRLFA D (SEQ ID NO:204)	RASQEIYGAL N (SEQ ID NO:152)	GATNLQS (SEQ ID NO:153)	QSTYDAPWT (SEQ ID NO:202)
xhCD8v14	SYAMS (SEQ ID NO:220)	DISYAGGSTA YADSVKG (SEQ ID NO:260)	SNAYAWDD ALDI (SEQ ID NO:222)	RASQSVSSN LA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T (SEQ ID NO:178)
xhCD8v15	SYAMS (SEQ ID NO:220)	DISYAGGSTA YADSVKG	SNAYAWDD ALDI (SEQ ID NO:222)	RASQSVSSN LA	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T

		(SEQ ID NO:260)		(SEQ ID NO:176)		(SEQ ID NO:178)
V9 family	X_1X_2AIS X_1 is S, K, G, N, R, D, T, or G X_2 is Y, L, H, or F (SEQ ID NO:185)	$X_1X_2X_3PX_4X_5X_6X_7X_8X_9YX_{10}QKFX_{11}G$ X_1 is G or H, X_2 is I or F, X_3 is I, N, or M, X_4 is G, N, H, S, R, I, or A, X_5 is A, N, H, S, T, F, or Y, X_6 is A, D, or G, X_7 is T, E, K, V, Q, or A, X_8 is A or T, X_9 is N or K, X_{10} is A or N, X_{11} is Q or T (SEQ ID NO:186)	$X_1X_2X_3GX_4X_5LFX_6X_7$ X_1 is D or A, X_2 is A, G, E, R, Y, K, N, Q, L, or F, X_3 is A, L, P, or Y, X_4 is I or L, X_5 is R, A, Q, or S, X_6 is A or D, X_7 is D, E, A, or S (SEQ ID NO:187)	$X_1X_2SX_3X_4IX_5GX_6LN$ X_1 is R or G, X_2 is A or T, X_3 is Q or E, X_4 is E, N, T, S, A, K, D, G, R, or Q, X_5 is Y or S, X_6 is A or V (SEQ ID NO:188)	$GX_1X_2X_3LX_4X_5$ X_1 is A or S, X_2 is T, S, E, Q, or D, X_3 is N, R, A, E, or H, X_4 is Q or A, X_5 is S or D (SEQ ID NO:189)	$QX_1X_2X_3X_4X_5PWT$ X_1 is S, N, D, Q, A, or E, X_2 is T, I, or S, X_3 is Y, L, or F, X_4 is D, G, T, E, Q, A, or Y, X_5 is A, T, R, S, K, or Y (SEQ ID NO:190)
V11 family	X_1YX_2MS X_1 is S, D, E, A, or Q X_2 is A, G, or T (SEQ ID NO:208)	$DIX_1X_2X_3GX_4X_5TX_6YADSVKG$ X_1 is T, N, S, Q, E, H, R, or A, X_2 is Y, W, F, or H, X_3 is A, S, Q, E, or T, X_4 is G or E, X_5 is S or I, X_6 is A or G (SEQ ID NO:209)	$X_1X_2X_3YX_4WX_5X_6AX_7DX_8$ X_1 is S or A, X_2 is N, H, A, D, L, Q, Y, or R, X_3 is A, N, S, or G, X_4 is A, V, R, E, or S, X_5 is D or S, X_6 is D, N, Q, E, S, T, or L, X_7 is L, F, or M, X_8 is I, Y, or V (SEQ ID NO:210)	RASQSVSSNLA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPVT (SEQ ID NO:178)

Table 11. Anti-CD8 antibody CDRs (Chothia)

Name	CDR-H1	CDR-H2	CDR-H3	CDR-L1	CDR-L2	CDR-L3
xhCD8	GYTFTKY (SEQ ID NO:223)	NPNNDE (SEQ ID NO:224)	DGLGLRLFAD (SEQ ID NO:225)	GASENIYGALN (SEQ ID NO:140)	GATNLAD (SEQ ID NO:141)	QNILDTPWT (SEQ ID NO:142)
xhCD8v1	GYTFTKY (SEQ ID NO:223)	NPNNDE (SEQ ID NO:224)	DGLGLRLFAD (SEQ ID NO:284)	RASENIYGALN (SEQ ID NO:285)	GATNLAD (SEQ ID NO:286)	QNILDTPWT (SEQ ID NO:287)
xhCD8v2	GYRFHNF (SEQ ID NO:226)	IPGHAK (SEQ ID NO:227)	DGLGIRLFAD (SEQ ID NO:151)	RASQEIYGALN (SEQ ID NO:152)	GATNLQS (SEQ ID NO:153)	QDIYDAPWT (SEQ ID NO:154)
xhCD8v3	GSRFYKF (SEQ ID NO:228)	IPGHAK (SEQ ID NO:227)	DGLGIRLFAD (SEQ ID NO:157)	RASQEIYGALN (SEQ ID NO:158)	GATNLQS (SEQ ID NO:159)	QDIYDAPWT (SEQ ID NO:160)
xhCD8v4	GYTFTKY (SEQ ID NO:223)	IPGHAK (SEQ ID NO:227)	DGLGIRLFAD (SEQ ID NO:163)	RASQKIYGALN (SEQ ID NO:164)	GATNLQS (SEQ ID NO:165)	QNTYDTPWT (SEQ ID NO:166)
xhCD8v5	GSGFRGH (SEQ ID NO:229)	IPGHAK (SEQ ID NO:227)	DGLGIRLFAD (SEQ ID NO:169)	RASQKIYGALN (SEQ ID NO:170)	GATNLQS (SEQ ID NO:171)	QNTYDTPWT (SEQ ID NO:172)
xhCD8v6	GFTFDDY (SEQ ID NO:230)	NWSGEI (SEQ ID NO:231)	SNSYRWDDALDI (SEQ ID NO:175)	RASQSVSSNLA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPVT (SEQ ID NO:178)
xhCD8v7	GFTFDDY (SEQ ID NO:230)	SYDGSN (SEQ ID NO:232)	DRIGWYDYDAFDI (SEQ ID NO:181)	RASHSVGSNLA (SEQ ID NO:182)	DASNRAT (SEQ ID NO:183)	QQRSNWPPT (SEQ ID NO:184)
xhCD8v8	GYAFSSY (SEQ ID NO:233)	YPGDGD (SEQ ID NO:234)	SGAAFSSYYAMDY (SEQ ID NO:145)	RASENIYSNLA (SEQ ID NO:146)	AATNLAD (SEQ ID NO:147)	QHFWGTPWT (SEQ ID NO:148)
xhCD8v9	GGTFSSY (SEQ ID NO:241)	IPGAAT (SEQ ID NO:242)	DAAGIRLFAD (SEQ ID NO:204)	RASQEIYGALN (SEQ ID NO:152)	GATNLQS (SEQ ID NO:153)	QSTYDAPWT (SEQ ID NO:202)
xhCD8v10	GFTFSSY (SEQ ID NO:250)	TYAGGS (SEQ ID NO:251)	SNAYAWDDALDI (SEQ ID NO:288)	RASQSVSSNLA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPVT (SEQ ID NO:178)

xhCD8v11	GFTFSSY (SEQ ID NO:250)	TYAGGS (SEQ ID NO:251)	SNAYAWDD ALDI (SEQ ID NO:288)	RASQSVSSN LA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T (SEQ ID NO:178)
xhCD8v12 (xhCD8.1)	GGTFSSY (SEQ ID NO:241)	IPGYAT (SEQ ID NO:243)	DAAGIRLFA D (SEQ ID NO:204)	RASQSIYGAL N (SEQ ID NO:205)	GASNLQS (SEQ ID NO:206)	QSTYTAPWT (SEQ ID NO:207)
xhCD8v13	GGTFSSY (SEQ ID NO:241)	IPGYAT (SEQ ID NO:243)	DAAGIRLFA D (SEQ ID NO:204)	RASQEIYGAL N (SEQ ID NO:152)	GATNLQS (SEQ ID NO:153)	QSTYDAPWT (SEQ ID NO:202)
xhCD8v14	GFTFSSY (SEQ ID NO:250)	SYAGGS (SEQ ID NO:261)	SNAYAWDD ALDI (SEQ ID NO:288)	RASQSVSSN LA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T (SEQ ID NO:178)
xhCD8v15	GFTFSSY (SEQ ID NO:250)	SYAGGS (SEQ ID NO:261)	SNAYAWDD ALDI (SEQ ID NO:288)	RASQSVSSN LA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T (SEQ ID NO:178)
V9 family	$GX_1X_2FX_3X_4X_5$ X_1 is G, Y, S, or A, X_2 is T, S, G, R, N, or H, X_3 is S, T, R, H, Y, G, or P, X_4 is S, K, G, N, R, D, T, or G, X_5 is Y, L, H, or F (SEQ ID NO:235)	$X_1PX_2X_3X_4X_5$ X_1 is I, N, or M, X_2 is G, N, H, S, R, I, or A, X_3 is A, N, H, S, T, F, or Y, X_4 is A, D, or G, X_5 is T, E, K, V, Q, or A (SEQ ID NO:236)	$X_1X_2X_3GX_4X_5$ LFX_6X_7 X_1 is D or A, X_2 is A, G, E, R, Y, K, N, Q, L, or F, X_3 is A, L, P, or Y, X_4 is I or L, X_5 is R, A, Q, or S, X_6 is A or D, X_7 is D, E, A, or S (SEQ ID NO:237)	$X_1X_2SX_3X_4IX_5$ GX_6LN X_1 is R or G, X_2 is A or T, X_3 is Q or E, X_4 is E, N, T, S, A, K, D, G, R, or Q, X_5 is Y or S, X_6 is A or V (SEQ ID NO:188)	$GX_1X_2X_3LX_4X_5$ X_1 is A or S, X_2 is T, S, E, Q, or D, X_3 is N, R, A, E, or H, X_4 is Q or A, X_5 is S or D (SEQ ID NO:189)	$QX_1X_2X_3X_4X_5$ PWT X_1 is S, N, D, Q, A, or E, X_2 is T, I, or S, X_3 is Y, L, or F, X_4 is D, G, T, E, Q, A, or Y, X_5 is A, T, R, S, K, or Y (SEQ ID NO:190)
V11 family	$GFTFX_1X_2Y$ X_1 is S, D, E, Q, S, or A X_2 is S, D, E, A, or Q (SEQ ID NO:244)	$X_1X_2X_3GX_4X_5$ X_1 is T, N, S, Q, E, H, R or A, X_2 is Y, W, F, or H, X_3 is A, S, Q, E, or T, X_4 is G or E, X_5 is S or I (SEQ ID NO:245)	$X_1X_2X_3YX_4W$ $X_5X_6AX_7DX_8$ X_1 is S or A, X_2 is N, H, A, D, L, Q, Y, or R, X_3 is A, N, S, or G, X_4 is A, V, R, E, or S, X_5 is D or S, X_6 is	RASQSVSSN LA (SEQ ID NO:176)	GASSRAT (SEQ ID NO:177)	QQYGSSPPV T (SEQ ID NO:178)

			D, N, Q, E, S, T, or L, X_7 is L, F, or M, X_8 is I, Y, or V (SEQ ID NO:246)			

Table 12. Anti-CD8 antibody variable domain sequences

Name	VH	VL
xhCD8	QVHLQQSGPELVKPGASVKMSCKTSGYTFTKY TMHWVKQGHEESLEWIGHFNPNNDETKYNQ KFTGKATLTVDKSSTTAYMELRSLTSDDSALYY CARDGLGLRLFADWGQGTLITVSA (SEQ ID NO:107)	DIQMTQSPASLSASVGETVTITCGASENIYGAL NWYQRKQGKSPQLLIFGATNLADGVSSRFSGS GSDRQYSLKISSLHPDDVATYYCQNILDTPWTF GGGTKLEIK (SEQ ID NO:108)
xhCD8v1	QVQLVQSGAEVKKPGSSVKVSCKASGYTFTKY AISWVRQAPGQGLEWMGHFNPNNDETKYN QKFQGRVTITADESTSTAYMELSSLRSEDTAVY YCARDGLGLRLFADWGQGT LVTVSS (SEQ ID NO:258)	DIQMTQSPSSLSASVGDRVTITCRASENIYGAL NWYQQKPGKAPKLLIYGATNLADGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQNILDTPWTF GGGTKLEIK (SEQ ID NO:259)
xhCD8v2	QVQLVQSGAEVKKPGSSVKVSCKASGYRFHNF AISWVRQAPGQGLEWMGGIIPGHAKANYAQ KFQGRVTITADESTSTAYMELSSLRSEDTAVYY CARDGLGIRLFADWGQGTLVTVSS (SEQ ID NO:109)	DIQMTQSPSSLSASVGDRVTITCRASQEIYGAL NWYQQKPGKAPKLLIYGATNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQDIYDAPWTF GGGTKVEIK (SEQ ID NO:110)
xhCD8v3	QVQLVQSGAEVKKPGSSVKVSCKASGSRFYKF AISWVRQAPGQGLEWMGGIIPGHAKANYAQ KFQGRVTITADESTSTAYMELSSLRSEDTAVYY CARDGLGIRLFADWGQGTLVTVSS (SEQ ID NO:111)	DIQMTQSPSSLSASVGDRVTITCRASQEIYGAL NWYQQKPGKAPKLLIYGATNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQDIYDAPWTF GGGTKVEIK (SEQ ID NO:112)
xhCD8v4	QVQLVQSGAEVKKPGSSVKVSCKASGYTFTKY AISWVRQAPGQGLEWMGGIIPGHAKANYAQ KFQGRVTITADESTSTAYMELSSLRSEDTAVYY CARDGLGIRLFADWGQGTLVTVSS (SEQ ID NO:113)	DIQMTQSPSSLSASVGDRVTITCRASQKIYGAL NWYQQKPGKAPKLLIYGATNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQNTYDTPWTF GGGTKVEIK (SEQ ID NO:114)
xhCD8v5	QVQLVQSGAEVKKPGSSVKVSCKASGSGFRG HAISWVRQAPGQGLEWMGGIIPGHAKANYA QKFQGRVTITADESTSTAYMELSSLRSEDTAVY YCARDGLGIRLFADWGQGTLVTVSS (SEQ ID NO:115)	DIQMTQSPSSLSASVGDRVTITCRASQKIYGAL NWYQQKPGKAPKLLIYGATNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQNTYDTPWTF GGGTKVEIK (SEQ ID NO:116)
xhCD8v6	EVQLVESGGGAVRPGGSLRLSCAASGFTFDDY GMSWVRQAPGKGLEWVSDINWSGEITAYAD SVKGRFTISRDNAKNSLYLQMNSLRAEDTAVY YCARSNSYRWDDALDIWGQGTMVTVSS (SEQ ID NO:117)	EIVLTQSPATLSVSPGERATLSCRASQSVSSNLA WYQQKPGQAPRLLIYGASSRATGIPDRFSGSG SGTDFTLTISRLEPEDFAVYYCQQYGSSPPVTF GQGTKVEIK (SEQ ID NO:118)

xhCD8v7	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDY AMHWVRQAPGKGLEWVAVISYDGSNKYYAD SVKGRFTISRDNSKNTLYLQMNSLRAEDTAVY YCAKDRIGWYDYDAFDIWGQGTMVTVSS (SEQ ID NO:119)	EIVLTQSPATLSVTPGEGATLSCRASHSVGSNL AWYQQKPGQAPRLLIYDASNRATGIPARFSGS GSGTDFTLTISSLEPEDLAVYYCQQRSNWPPTF GQGTRLEIK (SEQ ID NO:120)
xhCD8v8	QVQLQQSGAELVRPGSSVKISCKASGYAFSSY WMNWVKQRPGQGLEWIGQIYPGDGDTNYN GKFKGKATLTADKSSSTAYMQLSSLTSEDSAVY FCARSGAAFSSYYAMDYWGQGTSVTVSS (SEQ ID NO:121)	DIQMTQSPASLSVSVGETVTITCRASENIYSNL AWYQQKQGKSPQLLVYAATNLADGVPSRFSG SGSGTQYSLKINSLQSEDFGSYYCQHFWGTPW TFGGGTKLEIK (SEQ ID NO:122)
xhCD8v9	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSY AISWVRQAPGQGLEWMGGIIPGAATANYAQ KFQGRVTITADESTSTAYMELSSLRSEDTAVYY CARDAAGIRLFADWGQGTLVTVSS (SEQ ID NO:123)	DIQMTQSPSSLSASVGDRVTITCRASQEIYGAL NWYQQKPGKAPKLLIYGATNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQSTYDAPWTF GGGTKVEIK (SEQ ID NO:124)
xhCD8v10	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSY AMSWVRQAPGKGLEWVSDITYAGGSTAYAD SVKGRFTISRDNAKNSLYLQMNSLRAEDTAVY YCARSNAYAWDDALDIWGQGTMVTVSS (SEQ ID NO:125)	EIVLTQSPGTLSLSPGERATLSCRASQSVSSNLA WYQQKPGQAPRLLIYGASSRATGIPDRFSGSG SGTDFTLTISRLEPEDFAVYYCQQYGSSPPVTF GQGTKVEIK (SEQ ID NO:126)
xhCD8v11	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSY AMSWVRQAPGKGLEWVSDITYAGGSTAYAD SVKGRFTISRDNAKNSLYLQMNSLRAEDTAVY YCARSNAYAWDDALDIWGQGTLVTVSS (SEQ ID NO:127)	EIVLTQSPGTLSLSPGERATLSCRASQSVSSNLA WYQQKPGQAPRLLIYGASSRATGIPDRFSGSG SGTDFTLTISRLEPEDFAVYYCQQYGSSPPVTF GQGTKVEIK (SEQ ID NO:128)
xhCD8v12 (xhCD8.1)	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSY AISWVRQAPGQGLEWMGGIIPGYATANYAQ KFQGRVTITADESTSTAYMELSSLRSEDTAVYY CARDAAGIRLFADWGQGTLVTVSS (SEQ ID NO:129)	DIQMTQSPSSLSASVGDRVTITCRASQSIYGAL NWYQQKPGKAPKLLIYGASNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQSTYTAPWTF GGGTKVEIK (SEQ ID NO:130)
xhCD8v13	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSY AISWVRQAPGQGLEWMGGIIPGYATANYAQ KFQGRVTITADESTSTAYMELSSLRSEDTAVYY CARDAAGIRLFADWGQGTLVTVSS (SEQ ID NO:131)	DIQMTQSPSSLSASVGDRVTITCRASQEIYGAL NWYQQKPGKAPKLLIYGATNLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQSTYDAPWTF GGGTKVEIK (SEQ ID NO:132)
xhCD8v14	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSY AMSWVRQAPGKGLEWVSDISYAGGSTAYAD SVKGRFTISRDNAKNSLYLQMNSLRAEDTAVY YCARSNAYAWDDALDIWGQGTMVTVSS (SEQ ID NO:133)	EIVLTQSPGTLSLSPGERATLSCRASQSVSSNLA WYQQKPGQAPRLLIYGASSRATGIPDRFSGSG SGTDFTLTISRLEPEDFAVYYCQQYGSSPPVTF GQGTKVEIK (SEQ ID NO:134)
xhCD8v15	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSY AMSWVRQAPGKGLEWVSDISYAGGSTAYAD SVKGRFTISRDNAKNSLYLQMNSLRAEDTAVY YCARSNAYAWDDALDIWGQGTLVTVSS (SEQ ID NO:135)	EIVLTQSPGTLSLSPGERATLSCRASQSVSSNLA WYQQKPGQAPRLLIYGASSRATGIPDRFSGSG SGTDFTLTISRLEPEDFAVYYCQQYGSSPPVTF GQGTKVEIK (SEQ ID NO:136)

Table 13: Intermediate and final yields for fusion proteins of IL-21 charge variants

Fusion protein	Yield after Protein A (mg/L)	Yield after Protein A+IEX (mg/L)
xhCD8.1-hIL21 v31.4	198	89
xhCD8.1-hIL21 v31.23	170	83
xhCD8v11-hIL21 v31.23	158	77

Table 14: Wild type IL-21 variants

IL-21 Mutein	Amino Acid Sequence	K_D at 25°C (nM)	SEQ ID NO
IL-21	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	0.25	1
IL-21 v0.1	QGQDRHMIRMDQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	0.54	313
IL-21 v0.2	QGQDRHMIRMEQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	0.37	314
IL-21 v0.3	QGQDRHMIRMRQLDDIVAQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQAMIHQHLSSR THGSEDS	NB	315
IL-21 v0.4	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKEKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	> 10000	316
IL-21 v0.5	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	5.5	317
IL-21 v0.6	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKFKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	2205	318
IL-21 v0.7	QGQDRHMERMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	171.9	319
IL-21 v0.8	QGQDAHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG	12.4	320

	RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS		
IL-21 v0.9	QGQDEHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	184.6	321
IL-21 v0.10	QGQDSHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	20.8	322
IL-21 v0.11	QGQDTHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	6.6	323
IL-21 v0.12	QGQDNHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	23.4	324
IL-21 v0.13	QGQDQHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	7.7	325
IL-21 v0.14	QGQDVHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	4.6	326
IL-21 v0.15	QGQDIHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	6.2	327
IL-21 v0.16	QGQDLHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	2.5	328
IL-21 v0.17	QGQDYHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	7.2	329
IL-21 v0.18	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKAKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	2976	330
IL-21 v0.19	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKNKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	3981	331
IL-21 v0.20	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKDKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	> 10000	332

IL-21 v0.21	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKSKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	3671	333
IL-21 v0.22	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKTKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	4085	334
IL-21 v0.23	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKQKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	3375	335
IL-21 v0.24	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKVKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	4351	336
IL-21 v0.25	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKIKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	3639	337
IL-21 v0.26	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKLKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	934.1	338
IL-21 v0.27	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKYKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	1353	339
IL-21 v0.28	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRAPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	0.77	340
IL-21 v0.29	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKREPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	0.69	341
IL-21 v0.30	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIAKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	0.72	342
IL-21 v0.31	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIEKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	2.3	343
IL-21 v0.32	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLARKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	0.78	344
IL-21 v0.33	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLERKPPSTNAG	1.2	345

	RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS		
IL-21 v0.34	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKALKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	1.81	346
IL-21 v0.35	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKELKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	511.4	347
IL-21 v0.36	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRAPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	40.2	348
IL-21 v0.37	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKREPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	46.4	349
IL-21 v0.38	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIAKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	35.0	350
IL-21 v0.39	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIEKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	265.1	351
IL-21 v0.40	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLARKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	31.9	352
IL-21 v0.41	QGQDFHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLERKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	86.9	353
IL-21 v0.42	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKALKFKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	>10000	354
IL-21 v0.43	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKELKFKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	NB	355
IL-21 v0.44	QGQDRHMIAMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	4982	356
IL-21 v0.45	QGQDRHMIDMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	>10000	357

IL-21 v0.46	QGQDRHMIEMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	> 10000	358
IL-21 v0.47	QGQDRHMIHMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	11.9	359
IL-21 v0.48	QGQDRHMISMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	3932	360
IL-21 v0.49	QGQDRHMITMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	11664	361
IL-21 v0.50	QGQDRHMINMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	4159	362
IL-21 v0.51	QGQDRHMIGMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	7677	363
IL-21 v0.52	QGQDRHMIVMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	11082	364
IL-21 v0.53	QGQDRHMIIMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	>10000	365
IL-21 v0.54	QGQDRHMILMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	7168	366
IL-21 v0.55	QGQDRHMIYMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	675.7	367
IL-21 v0.56	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIAKLKFKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	3431	368
IL-21 v0.57	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLAFKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	5041	369
IL-21 v0.58	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKFAPPSTNAG	4760	370

	RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS		
IL-21 v0.59	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLEFKPPSTNAG RRQKHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	NB	371
hIL21 v31	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPGGEGG GGGGGEGGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	0.24	40
hIL21 v31.4	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKEKPPGGEGG GGGGGEGGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	>10000	94
hIL21 v31.23	QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKQKPPGGEGG GGGGGEGGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	6059	96
hIL21 v31.51	QGQDRHMIGMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPGGEGG GGGGGEGGCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	7730	98
IL-21 v2.63	QGQDRHMIRMRQLIDRIDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKIKRKPPSTNAG GGQGHELTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	1186	377
IL-21 v2.64	QGQDRHMIRMRQFIDAADQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKMKRKPPSTNAG GGQGHELTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	2823	378
IL-21 v2.65	QGQDRHMIRMRQRIDAIDQLKNYVNDLVPEFLPAPEDVETNC EWSAFSCFQKAQLKSANTGNNERIINVSIKKIKRKPPSTNAG GGQGHELTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSR THGSEDS	10471	379

Table 15: Polyreactivity ELISA of IL-21R-attenuated fusion proteins

Construct	Binding to KLH	Binding to heparin	Binding to hemoglobin
Bococizumab	24.35	10.20	25.37
xhCD8.1-hIL21 v0.4	5.67	7.39	6.59
xhCD8.1-hIL21 v31.4	2.98	2.92	3.94
xhCD8.1-hIL21 v31.23	4.66	3.61	5.45
xhCD8v11-hIL21 v31.23	3.06	3.37	2.10

Table 16: Polyreactivity ELISA of non-fusion IL-21 and IL-21 variants

Construct	Binding to KLH	Binding to heparin	Binding to hemoglobin
hIL21	24.06	51.09	31.42
hIL21 v0.51	23.54	48.90	30.58
hIL21 v0.4	22.93	44.65	28.60
hIL21 v0.23	23.76	51.02	30.94
hIL21 v31	20.97	44.73	19.99
hIL21 v31.51	15.67	31.45	12.22
hIL21 v31.4	6.44	12.22	3.40
hIL21 v31.23	12.78	26.22	9.81

CLAIMS

1. A fusion protein comprising:

a) an IL-21 polypeptide or a functional fragment or a variant thereof comprising an amino acid sequence that is at least 80% identical to a human IL-21 polypeptide comprising SEQ ID NO: 1; and

b) an antibody or an antigen binding fragment thereof that specifically binds to at least one of CD8α, CD8αα, or CD8αβ.

2. The fusion protein of claim 1, wherein the IL-21 polypeptide has an isoelectric point that is at least about 0.6 units to about 5 units lower, compared to that of the human IL-21 polypeptide.

3. The fusion protein of claim 2, wherein the IL-21 polypeptide comprises at least one amino acid substitution that reduces the isoelectric point of the IL-21 polypeptide by about 0.6 units to about 5 units relative to the human IL-21 polypeptide without the amino acid substitution.

4. The fusion protein of claim 2, wherein the IL-21 polypeptide comprises at least four amino acid substitutions that reduce the isoelectric point of the IL-21 polypeptide about 0.6 units to about 5 units relative to the human IL-21 polypeptide without the amino acid substitutions.

5. The fusion protein of claim 3 or 4, wherein the IL-21 polypeptide comprises up to 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 amino acid substitutions that reduce the isoelectric point.

6. The fusion protein of any one of claims 1-5, wherein the human IL-21 polypeptide has an isoelectric point of about 9.42.

7. The fusion protein of any one of claims 1-6, wherein the IL-21 polypeptide has an isoelectric point of about 7.12 to about 8.72.

8. The fusion protein of any one of claims 1-7, wherein the IL-21 polypeptide results in an improved exposure following administration to a subject relative to the human IL-21 polypeptide, as measured by at least about 1.5 times greater under the curve (AUC) for the IL-21 polypeptide.

9. The fusion protein of any one of claims 2-8, wherein relative to a region of the human IL-21 polypeptide that comprises about 2 to 20 positively charged amino acid residues, the IL-21

polypeptide comprises at least one amino acid substitution of the about 2 to 20 positively charged amino acid residues.

10. The fusion protein of claim 9, wherein the region of the human IL-21 polypeptide does not include amino acid residues that bind an IL-21 receptor.

11. The fusion protein of claim 9 or 10, wherein the region of the human IL-21 polypeptide comprises amino acid residues S80 to T92 of the human IL-21 polypeptide comprising SEQ ID NO: 1.

12. The fusion protein of any one of claims 9-11, wherein the human IL-21 polypeptide comprises at least one positively charged amino acid residue on the surface of the human IL-21 polypeptide in a three-dimensional structure of the human IL-21 polypeptide and does not bind an IL-21 receptor, and wherein the IL-21 polypeptide comprises at least one amino acid substitution of the at least one positively charged amino acid residue.

13. The fusion protein of any one of claims 11-12, wherein the IL-21 polypeptide does not comprise an amino acid substitution at G84 of the human IL-21 polypeptide comprising SEQ ID NO: 1.

14. The fusion protein of any one of claims 2-13, wherein the IL-21 polypeptide comprises 4, 5, 6, 7, 8, 9, 10, 11, or 12 amino acid substitutions at positions selected from S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, or T92 of SEQ ID NO: 1.

15. The fusion protein of claim 14, wherein the IL-21 polypeptide comprises 4, 5, 6, 7, 8, 9, 10, 11, or 12 amino acid substitutions selected from S80G, T81G, N82G, N82E, A82G, A83E, A83S, R85G, R85E, R85S, R86G, R86E, Q87G, Q87E, Q87S, K88G, H89G, H89S, R90G, R90S, R90E, R90A, L91G, L91S, T92G, T92S.

16. The fusion protein of claim 14, wherein the IL-21 polypeptide comprises (a) R85G, R86G, K88G and R90E, or (b) S80G, T81G, N82E, A83G, R85G, R86G, Q87G, K88G, H89G, R90E, L91G, and T92G.

17. The fusion protein of any one of claims 1-16, wherein the IL-21 polypeptide comprises the amino acid sequence:

QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKAQLKSANT GNNERIINVSIKKLKRKPPX$_1$X$_2$X$_3$X$_4$GX$_5$X$_6$X$_7$X$_8$X$_9$X$_{10}$X$_{11}$X$_{12}$CPSCDSYEKKPPKEFLERFKSL LQKMIHQHLSSRTHGSEDS (SEQ ID NO: 380), wherein X_1 = G, S; X_2 = G, T; X_3 = G, E, N; X_4 = G, S, E, A; X_5 = G, E, S, R; X_6 = G, E, R; X_7 = S, G, E, Q; X_8 = G, K; X_9 = G, S, H; X_{10} = A, E, S, G, R; X_{11} = S, G, L; and X_{12} = G, S, T, provided at least one of X_5, X_6, X_8, X_9, and X_{10} is not the amino acid residue at the identical position set forth in SEQ ID NO: 1, optionally wherein:

(i) X_5 = G, X_6 = G, X_8 = G, X_{10} = A;

(ii) X_5 = G, X_6 = G, X_8 = G, X_{10} = E;

(iii) X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = E, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = S;

(iv) X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = G, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = E;

(v) X_5 = G, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = E;

(vi) X_5 = G, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = S;

(vii) X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E;

(viii) X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = G;

(ix) X_5 = G, X_6 = G, X_7 = E, X_8 = G, X_9 = G, X_{10} = G;

(x) X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = G, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = G, X_{11} = S, X_{12} = G;

(xi) X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = G, X_{11} = G, X_{12} = G;

(xii) X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = G, X_6 = G, X_7 = E, X_8 = G, X_9 = G, X_{10} = G, X_{11} = S, X_{12} = G;

(xiii) X_1 = G, X_2 = G, X_3 = G, X_4 = E, X_5 = G, X_6 = G, X_7 = E, X_8 = G, X_9 = G, X_{10} = G, X_{11} = S, X_{12} = G;

(xiv) X_3 = G, X_4 = G, X_5 = S, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = G, X_{11} = G, X_{12} = S;

(xv) X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = G, X_{11} = G, X_{12} = S;

(xvi) X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = G, X_{11} = G, X_{12} = G;

(xvii) X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = G, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xviii) X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xix) X_3 = G, X_4 = G, X_5 = S, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = E, X_{11} = G, X_{12} = S;

(xx) X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xxi) X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxii) X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxiii) X_3 = G, X_4 = G, X_5 = G, X_6 = E, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxiv) X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxv) X_1 = G, X_2 = G, X_3 = G, X_4 = E, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G; or

(xxvi) X_1 = G, X_2 = G, X_3 = E, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G.

18. The fusion protein of any one of claims 1-17, wherein the IL-12 polypeptide comprises an amino acid sequence at least 90% identical to an amino acid sequence selected from SEQ ID NOs: 2, 4-15, and 23-40.

19. The fusion protein of claim 18, wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 2, 4-15, and 23-40.

20. The fusion protein of claim 19, wherein the IL-21 polypeptide comprises the amino acid sequence of SEQ ID NO: 40.

21. The fusion protein of any one of claims 1-20, wherein the IL-21 polypeptide comprises at least one amino acid substitution that reduces binding to an IL-21 receptor compared to binding by the human IL-21 polypeptide.

22. The fusion protein of claim 21, wherein the at least one amino acid substitution is at one or more amino acid residues at positions R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, or K117, of SEQ ID NO: 1.

23. The fusion protein of claim 22, wherein the at least one amino acid substitution is selected from: R5F, R5A, R5E, R5S, R5T, R5N, R5Q, R5V, R5I, R5L, R5Y, I8E, R9A, R9D, R9E, R9H, R9S, R9T, R9N, R9G, R9V, R9I, R9L, R9Y, R11D, R11E, L13F, L13R, I14D, I16A, I16S, I16R, V17I, V17A, D18A, K72A, K72E, K73A, K73E, K75A, K75E, L74I, L74F, L74M, L74V, R76E, R76F, R76A, R76N, R76D, R76S, R76T, R76Q, R76V, R76I, R76L, R76Y, R76M, K77A, K77E, and K117A.

24. The fusion protein of claim 22 or 23, wherein the at least one amino acid substitution is R76E or R76Q.

25. A fusion protein comprising:

a) an IL-21 polypeptide comprising (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions is within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide; and

b) an antibody or an antigen binding fragment thereof that specifically binds to CD8αβ or CD8β.

26. The fusion protein of claim 25, wherein the IL-21 polypeptide does not comprise a G84 substitution.

27. The fusion protein of claim 25 or 26, wherein the at least one amino acid substitution providing reduced binding is R76E or R76Q.

28. The fusion protein of any one of claims 1-27, wherein the IL-21 polypeptide comprises an amino acid sequence at least 90% identical to an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

29. The fusion protein of claim 28, wherein the IL-21 polypeptide comprises an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

30. The fusion protein of any one of claims 1-29, wherein the IL-21 polypeptide and the antibody or antigen binding fragment thereof are linked to each other via a linker.

31. The fusion protein of anyone of claims 1-30, wherein the IL-21 polypeptide is linked to the N-terminus or C-terminus of the antibody or antigen binding fragment thereof.

32. The fusion protein of claim 31, wherein the IL-21 polypeptide is linked to the C-terminus of the antibody or antigen binding fragment thereof.

33. The fusion protein of any one of claims 1-32, wherein the antibody or antigen binding fragment thereof comprises:

i) a first polypeptide comprising a variable light chain (VL) amino acid sequence, and a light chain constant region (CL1) amino acid sequence;

ii) a second polypeptide comprising a variable heavy chain (VH) amino acid sequence, and a heavy chain (HC) constant region amino acid sequence; and

iii) a third polypeptide, comprising a HC constant region amino acid sequence,

wherein the heavy chain constant regions of the second and third polypeptides form an Fc domain.

34. The fusion protein of claim 33, wherein the third polypeptide comprises a VH amino acid sequence.

35. The fusion protein of claim 33 or 34, comprising a fourth polypeptide comprising a VL amino acid sequence and a CL1 amino acid sequence.

36. The fusion protein of any one of claims 1-32, wherein the antibody or antigen binding fragment thereof comprises

i) a first polypeptide, arranged from N-to-C terminus, comprising:

a) a VL amino acid sequence, and

b) a CL1 amino acid sequence;

ii) a second polypeptide, arranged from N-to-C terminus, comprising:

a) a VH amino acid sequence,

b) a heavy chain CH1 constant region amino acid sequence,

c) a hinge region amino acid sequence,

d) a heavy chain CH2 constant region amino acid sequence, and

e) a heavy chain CH3 constant region amino acid sequence; and

iii) a third polypeptide, arranged from N-to-C terminus, comprising:

a) a hinge region amino acid sequence,

b) a heavy chain CH2 constant region amino acid sequence, and

c) a heavy chain CH3 constant region amino acid sequence,

wherein the CH2 and CH3 domains of each of the second and third polypeptides form an Fc domain.

37. The fusion protein of claim 36, wherein the third polypeptide, arranged from N-to-C terminus, comprises:

a) a VH amino acid sequence,
b) a heavy chain CH1 constant region amino acid sequence,
c) the hinge region amino acid sequence,
d) the heavy chain CH2 constant region amino acid sequence,
e) the heavy chain CH3 constant region amino acid sequence.

38. The fusion protein of claim 36 or 37, wherein the antibody or antigen binding fragment thereof comprises:

iv) a fourth polypeptide, arranged from N-to-C terminus, comprising:

a) a VL amino acid sequence, and
b) a CL1 amino acid sequence.

39. The fusion protein of any one of claims 33-38, wherein the IL-21 polypeptide is linked to the first polypeptide, the second polypeptide, the third polypeptide, or the fourth polypeptide, provided the IL-21 polypeptide does not interfere with binding of the antibody or antigen binding fragment thereof.

40. The fusion protein of any one of claims 33-35, wherein the IL-21 polypeptide is linked to the heavy chain constant region of the second polypeptide or the third polypeptide.

41. The fusion protein of claim 40, wherein the IL-21 polypeptide is linked to the C-terminus of the heavy chain constant region of the second polypeptide or the third polypeptide.

42. The fusion protein of any one of claims 36-38, wherein the IL-21 polypeptide is linked to the CH1, CH2 or CH3 regions of the second polypeptide or the third polypeptide.

43. The fusion protein of claim 42, wherein the IL-21 polypeptide is linked to the CH3 region of the second polypeptide or the third polypeptide.

44. The fusion protein of any one of claims 33-43, wherein the Fc domain is a human IgG Fc domain.

45. The fusion protein of claim 44, wherein the Fc domain is an IgG1, IgG2, IgG3, or IgG4 Fc domain.

46. The fusion protein of any one of claims 33-45, wherein the Fc domain comprises one or more modifications that promote heterodimerization.

47. The fusion protein of any one of claims 33-35, 39-41 and 44-46, wherein the second polypeptide comprises a knob modification in the HC constant region, and the third polypeptide comprises a hole modification in the HC constant region; or wherein the third polypeptide comprises a knob modification in HC constant region and the second polypeptide comprises a hole modification in the HC constant region.

48. The fusion protein of any one of claims 36-39 and 42-46, wherein the second polypeptide comprises a knob modification in the CH3 domain, and the third polypeptide comprises a hole modification in the CH3 domain; or wherein the third polypeptide comprises a knob modification in the CH3 domain and the second polypeptide comprises a hole modification in the CH3.

49. The fusion protein of claim 47 or 48, wherein the knob modification comprises amino acid substitutions Y349C and T366W according to EU numbering, and wherein the hole modification comprises amino acid substitutions S354C, T366S, L368A and Y407V according to EU numbering.

50. The fusion protein of any one of claims 33-49, wherein the Fc domain comprises an amino acid mutation that reduces binding of the Fc domain to an Fc gamma receptor.

51. The fusion protein of claim 50, wherein the amino acid mutation in the Fc domain is selected from L234A, L235A, and G237A, according to EU numbering.

52. The fusion protein of any one of claims 33-50, wherein the Fc domain comprises amino acid substitutions L234A, L235A, and G237A, according to EU numbering.

53. The fusion protein of any one of claims 1-52, wherein the antibody or antigen binding fragment thereof specifically binds to CD8αβ or CD8β.

54. A fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, and wherein the HC constant domain comprises a knob modification;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions is within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

55. A fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions is within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

56. The fusion protein of claim 54 or 55, wherein the at least one amino acid substitution providing reduced binding is R76E or R76Q.

57. The fusion protein of any one of claims 1-56, wherein the antibody or antigen binding fragment thereof comprises a heavy chain variable (VH) domain and a light chain variable (VL) domain; and wherein:

(a) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:137, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:138, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:139; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:140, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:141, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:142;

(b) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:149, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:150, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:151; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:154;

(c) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:155, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:156, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:157; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:158, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:159, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:160;

(d) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:161, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:162, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:163; and wherein the VL domain comprises a

CDR-L1 comprising the amino acid sequence of SEQ ID NO:164, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:165, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:166;

(e) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:167, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:168, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:169; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:170, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:171, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:172;

(f) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:173, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:174, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:175; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

(g) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:179, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:180, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:181; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:182, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:183, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:184;

(h) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:143, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:144, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:145; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:146, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:147, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:148;

(i) the VH domain comprises a CDR-H1 comprising the amino acid sequence of X_1X_2AIS, wherein X_1 is S, K, G, N, R, D, T, or G, and wherein X_2 is Y, L, H, or F (SEQ ID NO:185), a CDR-H2 comprising the amino acid sequence of $X_1X_2X_3PX_4X_5X_6X_7X_8X_9YX_{10}QKFX_{11}G$, wherein X_1 is G or H, X_2 is I or F, X_3 is I, N, or M, X_4 is G, N, H, S, R, I, or A, X_5 is A, N, H, S, T, F, or Y, X_6 is A, D, or G, X_7 is T, E, K, V, Q, or A, X_8 is A or T, X_9 is N or K, X_{10} is A or N, and X_{11} is Q or T (SEQ ID NO:186), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5LFX_6X_7$, wherein X_1 is D or A, X_2 is A, G, E, R, Y, K, N, Q, L, or F, X_3 is A, L, P, or Y, X_4 is I or L, X_5 is

R, A, Q, or S, X_6 is A or D, and X_7 is D, E, A, or S (SEQ ID NO:187); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of $X_1X_2SX_3X_4IX_5GX_6LN$, wherein X_1 is R or G, X_2 is A or T, X_3 is Q or E, X_4 is E, N, T, S, A, K, D, G, R, or Q, X_5 is Y or S, and X_6 is A or V (SEQ ID NO:188), a CDR-L2 comprising the amino acid sequence of $GX_1X_2X_3LX_4X_5$, wherein X_1 is A or S, X_2 is T, S, E, Q, or D, X_3 is N, R, A, E, or H, X_4 is Q or A, and X_5 is S or D (SEQ ID NO:189), and a CDR-L3 comprising the amino acid sequence of $QX_1X_2X_3X_4X_5PWT$, wherein X_1 is S, N, D, Q, A, or E, X_2 is T, I, or S, X_3 is Y, L, or F, X_4 is D, G, T, E, Q, A, or Y, and X_5 is A, T, R, S, K, or Y (SEQ ID NO:190);

(j) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:200, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:201; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202;

(k) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(l) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202;

(m) the VH domain comprises a CDR-H1 comprising the amino acid sequence of X_1YX_2MS, wherein X_1 is S, D, E, A, or Q and X_2 is A, G, or T (SEQ ID NO:208), a CDR-H2 comprising the amino acid sequence of $DIX_1X_2X_3GX_4X_5TX_6YADSVKG$, wherein X_1 is T, N, S, Q, E, H, R, or A, X_2 is Y, W, F, or H, X_3 is A, S, Q, E, or T, X_4 is G or E, X_5 is S or I, and X_6 is A or G (SEQ ID NO:209), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3YX_4WX_5X_6AX_7DX_8$, wherein X_1 is S or A, X_2 is N, H, A, D, L, Q, Y, or R, X_3 is A, N, S, or G, X_4 is A, V, R, E, or S, X_5 is D or S, X_6 is D, N, Q, E, S, T, or L, X_7 is L, F, or M, and X_8 is I, Y, or V (SEQ ID NO:210); and wherein the VL domain comprises a CDR-L1 comprising the

amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178);

(n) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178);

(o) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:260, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); or

(p) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:252, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:253, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:254; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:255, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:256, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:257.

58. The fusion protein of any one of claims 1-56, wherein the antibody or antigen binding fragment thereof comprises a VH domain and a VL domain; and wherein:

(a) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:226, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:151; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:154;

(b) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:228, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:157; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:158, a CDR-L2 comprising the amino

acid sequence of SEQ ID NO:159, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:160;

(c) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:163; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:164, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:165, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:166;

(d) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:229, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:227, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:169; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:170, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:171, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:172;

(e) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:230, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:231, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:175; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

(f) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:230, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:232, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:181; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:182, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:183, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:184;

(g) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:224, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:225; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:140, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:141, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:142;

(h) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:233, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:234, and a CDR-H3

comprising the amino acid sequence of SEQ ID NO:145; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:146, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:147, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:148;

(i) the VH domain comprises a CDR-H1 comprising the amino acid sequence of $GX_1X_2FX_3X_4X_5$, wherein X_1 is G, Y, S, or A, X_2 is T, S, G, R, N, or H, X_3 is S, T, R, H, Y, G, or P, X_4 is S, K, G, N, R, D, T, or G, and X_5 is Y, L, H, or F (SEQ ID NO:235), a CDR-H2 comprising the amino acid sequence of $X_1PX_2X_3X_4X_5$, wherein X_1 is I, N, or M, X_2 is G, N, H, S, R, I, or A, X_3 is A, N, H, S, T, F, or Y, X_4 is A, D, or G, and X_5 is T, E, K, V, Q, or A (SEQ ID NO:236), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5LFX_6X_7$, wherein X_1 is D or A, X_2 is A, G, E, R, Y, K, N, Q, L, or F, X_3 is A, L, P, or Y, X_4 is I or L, X_5 is R, A, Q, or S, X_6 is A or D, and X_7 is D, E, A, or S (SEQ ID NO:237); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of $X_1X_2SX_3X_4IX_5GX_6LN$, wherein X_1 is R or G, X_2 is A or T, X_3 is Q or E, X_4 is E, N, T, S, A, K, D, G, R, or Q, X_5 is Y or S, and X_6 is A or V (SEQ ID NO:188), a CDR-L2 comprising the amino acid sequence of $GX_1X_2X_3LX_4X_5$, wherein X_1 is A or S, X_2 is T, S, E, Q, or D, X_3 is N, R, A, E, or H, X_4 is Q or A, and X_5 is S or D (SEQ ID NO:189), and a CDR-L3 comprising the amino acid sequence of $QX_1X_2X_3X_4X_5PWT$, wherein X_1 is S, N, D, Q, A, or E, X_2 is T, I, or S, X_3 is Y, L, or F, X_4 is D, G, T, E, Q, A, or Y, and X_5 is A, T, R, S, K, or Y (SEQ ID NO:190);

(j) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:242, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202;

(k) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(l) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a

CDR-L1 comprising the amino acid sequence of SEQ ID NO:152, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:153, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:202;

(m) the VH domain comprises a CDR-H1 comprising the amino acid sequence of $GFTFX_1X_2Y$, wherein X_1 is S, D, E, Q, S, or A and X_2 is S, D, E, A, or Q (SEQ ID NO:244), a CDR-H2 comprising the amino acid sequence of $X_1X_2X_3GX_4X_5$, wherein X_1 is T, N, S, Q, E, H, R or A, X_2 is Y, W, F, or H, X_3 is A, S, Q, E, or T, X_4 is G or E, and X_5 is S or I (SEQ ID NO:245), and a CDR-H3 comprising the amino acid sequence of $X_1X_2X_3YX_4WX_5X_6AX_7DX_8$, wherein X_1 is S or A, X_2 is N, H, A, D, L, Q, Y, or R, X_3 is A, N, S, or G, X_4 is A, V, R, E, or S, X_5 is D or S, X_6 is D, N, Q, E, S, T, or L, X_7 is L, F, or M, and X_8 is I, Y, or V (SEQ ID NO:246); and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178);

(n) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178);

(o) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:261, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of RASQSVSSNLA (SEQ ID NO:176), a CDR-L2 comprising the amino acid sequence of GASSRAT (SEQ ID NO:177), and a CDR-L3 comprising the amino acid sequence of QQYGSSPPVT (SEQ ID NO:178); or

(p) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:223, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:224, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:284; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:285, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:286, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:287.

59. The fusion protein of any one of claims 1-56, wherein the antibody or antigen binding fragment thereof comprises a VH domain and a VL domain; and wherein:

(a) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:109, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:110;

(b) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:111, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:112;

(c) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:113, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:114;

(d) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:115, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:116;

(e) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:117, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:118;

(f) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:119, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:120;

(g) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:123; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:124;

(h) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:129, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:130;

(i) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:131; and wherein the VL domain

comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:132;

(j) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:125; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:126;

(k) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:127, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:128;

(l) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:133; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:134;

(m) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:135; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:136;

(n) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:107; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:108; or

(o) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:121; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:122; or

(p) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:258; and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:259.

60. The fusion protein of any one of claims 1-56, wherein the antibody or antigen binding fragment thereof comprises a VH domain and a VL domain; and wherein:

(a) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(b) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:241, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:243, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

(c) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178; or

(d) the VH domain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:250, a CDR-H2 comprising the amino acid sequence of SEQ ID NO: 251, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:288; and wherein the VL domain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178.

61. The fusion protein of any one of claims 1-56, wherein the antibody or antigen binding fragment thereof comprises a VH domain and a VL domain; and wherein:

(a) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:129, and wherein the VL domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:130; or

(b) the VH domain comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:127, and wherein the VL domain

comprises an amino acid sequence that is at least 90%, at least 95%, at least 99%, or 100% identical to the sequence of SEQ ID NO:128.

62. A fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions is within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

63. A fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:205, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:206, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:207;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:199, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:203, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:204;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions is within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

64. A fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions is within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

65. A fusion protein comprising:

a) a first polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain comprises a CDR-L1 comprising the amino acid sequence of SEQ ID NO:176, a CDR-L2 comprising the amino acid sequence of SEQ ID NO:177, and a CDR-L3 comprising the amino acid sequence of SEQ ID NO:178;

b) a second polypeptide comprising a VH chain and a HC constant domain, wherein the HC constant domain comprises a knob modification, and wherein the VH chain comprises a CDR-H1 comprising the amino acid sequence of SEQ ID NO:220, a CDR-H2 comprising the amino acid sequence of SEQ ID NO:221, and a CDR-H3 comprising the amino acid sequence of SEQ ID NO:222;

c) a third polypeptide comprising a VH chain and a HC constant domain, wherein an IL-21 polypeptide is operably linked to the C-terminus of the HC constant domain, wherein the HC constant domain comprises a hole modification, and wherein the VH chain of the second polypeptide and the third polypeptide are the same; and

d) a fourth polypeptide comprising a VL chain and a CL1 chain, wherein the VL chain of the first polypeptide and the fourth polypeptide are the same, and

wherein the VH chains and VL chains comprise CDRs that specifically bind to CD8αβ or CD8β,

wherein the IL-21 polypeptide comprises (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions is within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination

thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

66. The fusion protein of any one of claims 62-65, wherein the at least one amino acid substitution providing reduced binding is R76E or R76Q.

67. The fusion protein of any one of claims 1-8, comprising a first polypeptide chain, a second polypeptide chain, a third polypeptide chain, and a fourth polypeptide chain, wherein:

(a) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:262, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:263, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:264, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:262;

(b) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:266, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:267, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:268, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:266;

(c) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:270, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:271, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:272, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 270;

(d) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:275, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:276, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274; or

(e) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:278, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:279, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:280, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 278.

68. The fusion protein of any one of claims 1-8, comprising a first polypeptide chain, a second polypeptide chain, a third polypeptide chain, and a fourth polypeptide chain, wherein:

(a) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:262, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:263, the third

polypeptide chain comprises the amino acid sequence of SEQ ID NO:265, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:262;

(b) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:266, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:267, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:269, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:266;

(c) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:270, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:271, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:273, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 270;

(d) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:275, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:277, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 274; or

(e) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:278, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:279, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:281, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 278.

69. The fusion protein of any one of claims 1-8, comprising a first polypeptide chain, a second polypeptide chain, a third polypeptide chain, and a fourth polypeptide chain, wherein:

(a) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:297, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:298, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:299, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:297;

(b) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:301, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:302, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:303 and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:301;

(c) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:305, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:306, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:307, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 305; or

(d) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:310, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:311, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309.

70. The fusion protein of any one of claims 1-8, comprising a first polypeptide chain, a second polypeptide chain, a third polypeptide chain, and a fourth polypeptide chain, wherein:

(a) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:297, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:298, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:300, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:297;

(b) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:301, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:302, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:304 and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:301;

(c) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO:305, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:306, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:308, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 305; or

(d) the first polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309, the second polypeptide chain comprises the amino acid sequence of SEQ ID NO:310, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:312, and the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO: 309.

71. A polynucleotide encoding the fusion protein of any one of claims 1-70.

72. The polynucleotide of claim 71, comprising a nucleotide sequence encoding the IL-21 polypeptide or the mutant IL-21 polypeptide, and at least one nucleotide sequence encoding the antibody or antigen binding fragment thereof.

73. A vector comprising the polynucleotide of claim 71 or 72.

74. A host cell comprising the vector of claim 73.

75. A pharmaceutical composition comprising the fusion protein of any one of claims 1-70.

76. A method of activating a population of CD8+ T cells, comprising contacting the population of CD8+ T cells with the fusion protein of any one of claims 1-70 or the pharmaceutical composition of claim 75.

77. A method of selective activation of CD8+ T cells, comprising contacting a population of cells comprising CD8+ T cells, CD4+ T cells and NK cells, with the fusion protein of any one of claims 1-70 or the pharmaceutical composition of claim 75.

78. The method of claim 76 or 77, wherein the CD8+ T cells are activated at least about 10-fold higher compared to activation of the NK cells or the CD4+ T cells by the fusion protein or pharmaceutical composition.

79. The method of any one of claims 77-78, wherein the CD8+ T cells are activated with a potency that is at least about 10-fold to about 100,000-fold greater as compared to activation of the NK cells or the CD4+ T cells in the population of cells.

80. The method of any one of claims 76-79, wherein activation of CD8+ T cells results in increased STAT3 phosphorylation of the CD8+ T cells compared to the STAT3 phosphorylation of the NK cells or the CD4+ T cells.

81. A method of treating a disease in a subject, the method comprising administering to the subject an effective amount of the fusion protein of any one of claims 1-70 or the pharmaceutical composition of claim 75, thereby treating the disease.

82. The method of claim 81, comprising administering an additional therapeutic agent.

83. The method of claim 81 or 82, wherein the disease comprises a cancer or a chronic infection.

84. The method of claim 81 or 82, wherein the disease comprises a cancer, and wherein the cancer is selected from acute lymphoblastic leukemia (ALL) (including non T cell ALL), acute myeloid leukemia, B cell prolymphocytic leukemia, B cell acute lymphoid leukemia ("BALL"),

blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloid leukemia, chronic or acute leukemia, diffuse large B cell lymphoma (DLBCL), follicular lymphoma (FL), hairy cell leukemia, Hodgkin's Disease, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, monoclonal gammopathy of undetermined significance (MGUS), multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin's lymphoma (NHL), plasma cell proliferative disorder (including asymptomatic myeloma (smoldering multiple myeloma or indolent myeloma), plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, plasmacytomas (including plasma cell dyscrasia; solitary myeloma; solitary plasmacytoma; extramedullary plasmacytoma; and multiple plasmacytoma), POEMS syndrome (also known as Crow-Fukase syndrome; Takatsuki disease; and PEP syndrome), primary mediastinal large B cell lymphoma (PMBC), small cell- or a large cell-follicular lymphoma, splenic marginal zone lymphoma (SMZL), systemic amyloid light chain amyloidosis, T cell acute lymphoid leukemia ("TALL"), T cell lymphoma, transformed follicular lymphoma, or Waldenstrom macroglobulinemia, Mantle cell lymphoma (MCL), Transformed follicular lymphoma (TFL), Primary mediastinal B cell lymphoma (PMBCL), Multiple myeloma, Hairy cell lymphoma/leukemia, lung cancer, small-cell lung cancer, non-small cell lung (NSCL) cancer, bronchioloalveolar cell lung cancer, squamous cell cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, head and neck cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, thyroid cancer, uterine cancer, , gastrointestinal cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, endometrial carcinoma, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the cervix, carcinoma of the vagina, vulval cancer, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, bladder cancer, liver cancer, hepatoma, hepatocellular cancer, cervical cancer, salivary gland carcinoma, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwannomas, ependymomas, medulloblastomas, meningiomas, squamous cell carcinomas, pituitary adenoma and Ewing's sarcoma, including refractory versions of any of the above cancers, and any combination of one or more of the above cancers.

85. Use of the fusion protein of any one of claims 1-70 for selectively activating CD8+ T cells, comprising contacting a population of cells comprising CD8+ T cells, CD4+ T cells, and NK cells with the fusion protein, wherein the CD8+ T cells are selectively activated compared to activation of the CD4+ T cells or the NK cells in the population of cells.

86. The fusion protein of any one of claims 1-70 for use in a method of selectively activating CD8+ T cells, comprising contacting a population of cells comprising CD8+ T cells, CD4+ T cells, and NK cells with the fusion protein, wherein the CD8+ T cells are selectively activated compared to activation of the CD4+ T cells or the NK cells in the population of cells.

87. Use of the fusion protein of any one of claims 1-70 for treating a disease in a subject, comprising administering the fusion protein to the subject.

88. The fusion protein of any one of claims 1-70 for use in a method of treating a disease in a subject, comprising administering the fusion protein to the subject.

89. A kit comprising a container comprising the fusion protein of any one of claims 1-70, and optionally a pharmaceutically acceptable carrier, and instructions for administering the fusion protein to a subject with a disease.

90. The use of claim 87, the fusion protein of claim 88, or the kit of claim 89, wherein the disease is a cancer or a chronic infection.

91. An IL-21 polypeptide or a functional fragment or a variant thereof comprising a polypeptide sequence having at least 80% sequence identity to SEQ ID NO: 1, wherein the IL-21 polypeptide has an isoelectric point that is at least about 0.6 units to about 5 units lower, compared to that of a human IL-21 polypeptide having a sequence of SEQ ID NO: 1.

92. The IL-21 polypeptide of claim 91, wherein the isoelectric point of the human IL-21 polypeptide is about 9.42.

93. The IL-21 polypeptide of claim 91 or 92, comprising an isoelectric point of about 7.12 to about 8.72.

94. The IL-21 polypeptide of anyone of claims 91-93, comprising at least one amino acid substitution that reduces the isoelectric point of the IL-21 polypeptide by about 0.6 units to about 5 units relative to the human IL-21 polypeptide without the amino acid substitution.

95. The IL-21 polypeptide of claim 94, comprising up to 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 amino acid substitutions that reduce the isoelectric point.

96. The IL-21 polypeptide of anyone of claims 91-95, wherein the IL-21 polypeptide results in an improved exposure following administration to a subject relative to the human IL-21 polypeptide, as measured by at least about 1.5 times greater under the curve (AUC) for the IL-21 polypeptide.

97. The IL-21 polypeptide of anyone of claims 91-96, wherein relative to a region of the human IL-21 polypeptide that comprises about 2 to 20 positively charged amino acid residues, the IL-21 polypeptide comprises at least one amino acid substitution of the about 2 to 20 positively charged amino acid residues.

98. The IL-21 polypeptide of claim 97, wherein the region of the human IL-21 polypeptide does not include amino acid residues that bind an IL-21 receptor.

99. The IL-21 polypeptide of claim 97 or 98, wherein the region of the human IL-21 polypeptide comprises amino acid residues S80 to T92 of the human IL-21 polypeptide comprising SEQ ID NO: 1.

100. The IL-21 polypeptide of any one of claims 97-99, wherein the human IL-21 polypeptide comprises at least one positively charged amino acid residue located in spatial proximity of the region in a three-dimensional structure of the human IL-21 polypeptide, and wherein the IL-21 polypeptide comprises at least one amino acid substitution of at least one positively charged amino acid residue located in spatial proximity of the region.

101. The IL-21 polypeptide of any one of claims 99-100, wherein the IL-21 polypeptide does not comprise an amino acid substitution at G84 of the human IL-21 polypeptide comprising SEQ ID NO: 1.

102. The IL-21 polypeptide of any one of claims 91-101, comprising at least four amino acid substitutions at positions selected from S80, T81, N82, A83, R85, R86, Q87, K88, H89, R90, L91, or T92, of SEQ ID NO: 1.

103. The IL-21 polypeptide of claim 102, comprising at least four amino acid substitutions selected from S80G, T81G, N82G, N82E, A82G, A83E, A83S, R85G, R85E, R85S, R86G, R86E, Q87G, Q87E, Q87S, K88G, H89G, H89S, R90G, R90S, R90E, R90A, L91G, L91S, T92G, T92S.

104. The IL-21 polypeptide of claim 102, comprising 4-12 amino acid substitutions selected from S80G, T81G, N82G, N82E, A82G, A83E, A83S, R85G, R85E, R85S, R86G, R86E, Q87G, Q87E, Q87S, K88G, H89G, H89S, R90G, R90S, R90E, R90A, L91G, L91S, T92G, T92S.

105. The IL-21 polypeptide of any one of claims 91-104, comprising the amino acid sequence: QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKAQLKSANT GNNERIINVSIKKLKRKPPX$_1$X$_2$X$_3$X$_4$GX$_5$X$_6$X$_7$X$_8$X$_9$X$_{10}$X$_{11}$X$_{12}$CPSCDSYEKKPPKEFLERFKSL LQKMIHQHLSSRTHGSEDS (SEQ ID NO: 380), wherein X$_1$ = G, S; X$_2$ = G, T; X$_3$ = G, E, N; X$_4$ = G, S, E, A; X$_5$ = G, E, S, R; X$_6$ = G, E, R; X$_7$ = S, G, E, Q; X$_8$ = G, K; X$_9$ = G, S, H; X$_{10}$ = A, E, S, G, R; X$_{11}$ = S, G, L; and X$_{12}$ = G, S, T, provided at least one of X$_5$, X$_6$, X$_8$, X$_9$, and X$_{10}$ is not the amino acid residue at the identical position set forth in SEQ ID NO: 1, optionally wherein:

(i) X$_5$ = G, X$_6$ = G, X$_8$ = G, X$_{10}$ = A;

(ii) X$_5$ = G, X$_6$ = G, X$_8$ = G, X$_{10}$ = E;

(iii) X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = E, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = S;

(iv) X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = E;

(v) X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = E;

(vi) X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = S;

(vii) X$_5$ = G, X$_6$ = G, X$_7$ = G, X$_8$ = G, X$_9$ = G, X$_{10}$ = E;

(viii) X$_5$ = G, X$_6$ = G, X$_7$ = G, X$_8$ = G, X$_9$ = G, X$_{10}$ = G;

(ix) X$_5$ = G, X$_6$ = G, X$_7$ = E, X$_8$ = G, X$_9$ = G, X$_{10}$ = G;

(x) X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = G, X$_6$ = G, X$_7$ = S, X$_8$ = G, X$_9$ = G, X$_{10}$ = G, X$_{11}$ = S, X$_{12}$ = G;

(xi) X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = G, X$_5$ = G, X$_6$ = G, X$_7$ = G, X$_8$ = G, X$_9$ = G, X$_{10}$ = G, X$_{11}$ = G, X$_{12}$ = G;

(xii) X$_1$ = G, X$_2$ = G, X$_3$ = G, X$_4$ = S, X$_5$ = G, X$_6$ = G, X$_7$ = E, X$_8$ = G, X$_9$ = G, X$_{10}$ = G, X$_{11}$ = S, X$_{12}$ = G;

(xiii) X_1 = G, X_2 = G, X_3 = G, X_4 = E, X_5 = G, X_6 = G, X_7 = E, X_8 = G, X_9 = G, X_{10} = G, X_{11} = S, X_{12} = G;

(xiv) X_3 = G, X_4 = G, X_5 = S, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = G, X_{11} = G, X_{12} = S;

(xv) X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = G, X_{11} = G, X_{12} = S;

(xvi) X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = G, X_{11} = G, X_{12} = G;

(xvii) X_1 = G, X_2 = G, X_3 = G, X_4 = S, X_5 = G, X_6 = G, X_7 = S, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xviii) X_1 = G, X_2 = G, X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xix) X_3 = G, X_4 = G, X_5 = S, X_6 = G, X_7 = G, X_8 = G, X_9 = S, X_{10} = E, X_{11} = G, X_{12} = S;

(xx) X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G;

(xxi) X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxii) X_3 = G, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxiii) X_3 = G, X_4 = G, X_5 = G, X_6 = E, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxiv) X_3 = G, X_4 = G, X_5 = E, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G;

(xxv) X_1 = G, X_2 = G, X_3 = G, X_4 = E, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G; or

(xxvi) X_1 = G, X_2 = G, X_3 = E, X_4 = G, X_5 = G, X_6 = G, X_7 = G, X_8 = G, X_9 = G, X_{10} = E, X_{11} = G, X_{12} = G.

106. The IL-21 polypeptide of any one of claims 91-105, comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from SEQ ID NOs: 2, 4-15, and 23-40.

107. The IL-21 polypeptide of claim 106, comprising an amino acid sequence selected from SEQ ID NOs: 2, 4-15, and 23-40.

108. The IL-21 polypeptide of claim 107, comprising the amino acid sequence of SEQ ID NO: 40.

109. The IL-21 polypeptide of any one of claims 91-108, comprising at least one amino acid substitution that reduces binding to an IL-21 receptor compared to binding by the human IL-21 polypeptide.

110. The IL-21 polypeptide of claim 109, wherein the at least one amino acid substitution is at one or more amino acid residues at positions R5, I8, R9, R11, L13, I14, I16, V17, D18, K72, K73, L74, K75, R76, K77, or K117, or SEQ ID NO: 1.

111. The IL-21 polypeptide of claim 110, wherein the at least one amino acid substitution is selected from: R5F, R5A, R5E, R5S, R5T, R5N, R5Q, R5V, R5I, R5L, R5Y, I8E, R9A, R9D, R9E, R9H, R9S, R9T, R9N, R9G, R9V, R9I, R9L, R9Y, R11D, R11E, L13F, L13R, I14D, I16A, I16S, I16R, V17I, V17A, D18A, K72A, K72E, K73A, K73E, K75A, K75E, L74I, L74F, L74M, L74V, R76E, R76F, R76A, R76N, R76D, R76S, R76T, R76Q, R76V, R76I, R76L, R76Y, R76M, K77A, K77E, and K117A.

112. The IL-21 polypeptide of claim 110 or 111, wherein the at least one amino acid substitution is R76E or R76Q.

113. The IL-21 polypeptide of any one of claims 91-112, comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

114. The IL-21 polypeptide of any one of claims 91-112, comprising an amino acid sequence selected from SEQ ID NOs: 16-21, 41-98, and 374-379.

115. An IL-21 polypeptide comprising, (i) one or more amino acid substitutions providing a reduced isoelectric point relative to a human IL-21 polypeptide comprising SEQ ID NO: 1, wherein the one or more amino acid substitutions is within S80 to T92 of SEQ ID NO: 1, wherein at least one amino acid substitution is at residue R85, R86, K88, H89, R90, or a combination thereof, and (ii) at least one amino acid substitution providing reduced binding to a human IL-21 receptor relative to binding by the human IL-21 polypeptide.

116. A fusion protein comprising (i) the IL-21 polypeptide of any one of claim 91-115; and (ii) a polypeptide of interest.

117. The fusion protein of claim 116, wherein the polypeptide of interest is antibody or antigen binding fragment thereof.

118. The fusion protein of claim 117, wherein the antibody or antigen binding fragment thereof binds a receptor on T cells.

119. The fusion protein of claim 118, wherein the receptor is on CD8+ T cells.

120. The fusion protein of claim 118, wherein the receptor is on T cells expressing an IL-21 receptor.

121. The fusion protein of any one of claims 118-120, wherein the receptor is not expressed on Treg cells.

122. A polynucleotide encoding the IL-21 polypeptide of any one of claims 91-115 or the fusion protein of any one of claims 116-121.

123. A vector comprising the polynucleotide of claim 122.

124. A host cell comprising the vector of claim 123.

125. A pharmaceutical composition comprising the IL-21 polypeptide or fusion protein of any one of claims 91-121.

126. A method of activating T cells, comprising contacting the T cells with the IL-21 polypeptide of any one of claims 91-115, the fusion protein of any one of claims 116-121, or the pharmaceutical composition of claim 125.

127. A method of treating a disease in a subject, comprising administering to the subject the IL-21 polypeptide of any one of claims 91-115, the fusion protein of any one of claims 116-121, or the pharmaceutical composition of claim 125.

128. The method of claim 127, comprising administering an additional therapeutic agent.

129. The method of claim 127 or 128, wherein the disease comprises a cancer or a chronic infection.

130. The method of claim 127 or 128, wherein the disease comprises a cancer, and wherein the cancer is selected from acute lymphoblastic leukemia (ALL) (including non T cell ALL), acute myeloid leukemia, B cell prolymphocytic leukemia, B cell acute lymphoid leukemia ("BALL"), blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloid leukemia, chronic or acute leukemia, diffuse large B cell lymphoma (DLBCL), follicular lymphoma (FL), hairy cell leukemia, Hodgkin's Disease, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, monoclonal gammopathy of undetermined significance (MGUS), multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin's lymphoma (NHL), plasma cell proliferative disorder (including asymptomatic myeloma (smoldering multiple myeloma or indolent myeloma), plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, plasmacytomas (including plasma cell dyscrasia; solitary myeloma; solitary plasmacytoma; extramedullary plasmacytoma; and multiple plasmacytoma), POEMS syndrome (also known as Crow-Fukase syndrome; Takatsuki disease; and PEP syndrome), primary mediastinal large B cell lymphoma (PMBC), small cell- or a large cell-follicular lymphoma, splenic marginal zone lymphoma (SMZL), systemic amyloid light chain amyloidosis, T cell acute lymphoid leukemia ("TALL"), T cell lymphoma, transformed follicular lymphoma, or Waldenstrom macroglobulinemia, Mantle cell lymphoma (MCL), Transformed follicular lymphoma (TFL), Primary mediastinal B cell lymphoma (PMBCL), Multiple myeloma, Hairy cell lymphoma/leukemia, lung cancer, small-cell lung cancer, non-small cell lung (NSCL) cancer, bronchioloalveolar cell lung cancer, squamous cell cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, head and neck cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, thyroid cancer, uterine cancer, , gastrointestinal cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, endometrial carcinoma, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the cervix, carcinoma of the vagina, vulval cancer, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, bladder cancer, liver cancer, hepatoma, hepatocellular cancer, cervical cancer, salivary gland carcinoma, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwannomas, ependymomas, medulloblastomas, meningiomas, squamous cell carcinomas, pituitary adenoma and

Ewing's sarcoma, including refractory versions of any of the above cancers, and any combination of one or more of the above cancers.

131. Use of the IL-21 polypeptide of any one of claims 91-115 or the fusion protein of any one of claims 116-121 for activating T cells, comprising contacting T cells with the IL-21 polypeptide or fusion protein.

132. The IL-21 polypeptide of any one of claims 91-115 or the fusion protein of any one of claims 116-121for use in a method of activating T cells, comprising contacting T cells with the IL-21 polypeptide or the fusion protein.

133. Use of the IL-21 polypeptide of any one of claims 91-115 or the fusion protein of any one of claims 116-121, for treating a disease in a subject, comprising administering the fusion protein to the subject.

134. The IL-21 polypeptide of any one of claims 91-115 or the fusion protein of any one of claims 116-121 for use in a method of treating a disease in a subject, comprising administering the fusion protein to the subject.

135. A kit comprising a container comprising the IL-21 polypeptide of any one of claims 91-115 or the fusion protein of any one of claims 116-121, and optionally a pharmaceutically acceptable carrier, and instructions for administering the fusion protein to a subject with a disease.

136. The use of claim 133, the fusion protein of claim 134, or the kit of claim 135, wherein the disease is a cancer or a chronic infection.

FIG. 1A

Amino acid sequence of wild type IL-21

QGQDRHMIRMRQLIDIVDQLKNYVNDLVPEFLPAPEDVETNCEWSAFSCFQKAQLKSANTGNNERIINVSIKKLKRKPPSTNAGRRQ KHRLTCPSCDSYEKKPPKEFLERFKSLLQKMIHQHLSSRTHGSEDS (SEQ ID NO: 1)

FIG. 1B

Amino acid sequence of IL-21R

CPDLVCYTDYLQTVICILEMWNLHPSTLTLTWQDQYEELKDEATSCSLHRSAHNATHATYTCHMDVFHFMADDIFSVNITDQSGNYS QECGSFLLAESIKPAPPFNVTVTFSGQYNISWRSDYEDPAFYMLKGKLQYELQYRNRGDPWAVSPRRKLISVDSRSVSLLPLEFRKDSSY ELQVRAGPMPGSSYQGTWSEWSDPVIFQTQSEELKEGWNPHLLLLLLLVIVFIPAFWSLKTHPLWRLWKKIWAVPSPERFFMPLYKG CSGDFKKWVGAPFTGSSLELGPWSPEVPSTLEVYSCHPPRSPAKRLQLTELQEPAELVESDGVPKPSFWPTAQNSGGSAYSEERDRPY GLVSIDTVTVLDAEGPCTWPCSCEDDGYPALDLDAGLEPSPGLEDPLLDAGTTVLSCGCVSAGSPGLGGPLGSLLDRLKPPLADGED WAGGLPWGGRSPGGVSESEAGSPLAGLDMDTFDSGFVGSDCSSPVECDFTSPGDEGPPRSYLRQWVVIPPPLSSPGPQAS (SEQ ID NO: 381)

FIG. 1C

Amino acid sequence of gamma common receptor

LNTTILTPNGNEDTTADFFLTTMPTDSLSVSTLPLPEVQCFVFNVEYMNCTWNSSSEPQPTNLTLHYWYKNSDNDKVQKCSHYLFSEE ITSGCQLQKKEIHLYQTFVVQLQDPREPRRQATQMLKLQNLVIPWAPENLTLHKLSESQLELNWNNRFLNHCLEHLVQYRTDWDHS WTEQSVDYRHKFSLPSVDGQKRYTFRVRSRFNPLCGSAQHWSEWSHPIHWGSNTSKENPFLFALEAVVISVGSMGLIISLLCVYFWL ERTMPRIPTLKNLEDLVTEYHGNFSAWSGVSKGLAESLQPDYSERLCLVSEIPPKGGALGEGPGASPCNQHSPYWAPPCYTLKPET (SEQ ID NO: 382)

Amino acid number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Wild-type residue	Q	G	Q	D	R	H	M	I	R	M	R	Q	L	I	D	I	V	D	Q	L
Mutation position																				
Amino acid number	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40
Wild-type residue	K	N	Y	V	N	D	L	V	P	E	F	L	P	A	P	E	D	V	E	T
Mutation position																				
Amino acid number	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60
Wild-type residue	N	C	E	W	S	A	F	S	C	F	Q	K	A	Q	L	K	S	A	M	T
Mutation position																X				
Amino acid number	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
Wild-type residue	G	N	N	E	R	I	I	N	V	S	I	K	K	L	K	R	K	P	P	S
Mutation position																				X
Amino acid number	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
Wild-type residue	T	N	A	G	R	R	Q	K	H	R	L	T	C	P	S	C	D	S	Y	E
Mutation position	X	X	X	X	X	X	X	X	X	X	X	X								
Amino acid number	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120
Wild-type residue	K	K	P	P	K	E	F	L	E	R	F	K	S	L	L	Q	K	M	I	H
Mutation position																				
Amino acid number	121	122	123	124	125	126	127	128	129	130	131	132	133							
Wild-type residue	Q	H	L	S	S	R	T	H	G	S	E	D	S							
Mutation position																				

FIG. 2


A.U.
homodimer
heterodimer
v6
v5
v4
v3
v2
v1
Wild Type
40%
57%
56%
60%
15%
20%
68%
60%
43%
44%
40%
85%
80%
32%
time (min)
Higher pI

FIG. 3A

FIG. 3B


homodimer
heterodimer
mAU
2000
1500
1000
500
0
8
18
28
38
time (min)
v12
v11
v10
v9
v8
v7
v2
83%
84%
80%
83%
88%
76%
85%

FIG. 3C


homodimer
heterodimer
mAU
2000
1500
1000
500
0
8
18
28
38
time (min)
v19
v18
v17
v16
v15
v14
v13
88%
86%
85%
81%
77%
82%
82%

FIG. 3D


homodimer
heterodimer
mAU
2000
1500
1000
500
0
6
18
28
38
time (min)
v20
v21
v22
v23
v24
v25
85%
88%
88%
84%
88%
84%

FIG. 3E


homodimer
heterodimer
mAU
2000
1500
1000
500
0
8
18
28
38
time (min)
v26
v27
v28
v29
v30
v31
86%
86%
85%
87%
78%
83%

FIG. 3F


WT
34%
heterodimer
undesired products
mAU
200
150
100
50
0
0
10
20
30
40
time (min)

FIG. 4B


Mouse PK (1mg/kg SubQ)
xCtrl-hIL21 v2
xCtrl-hIL21
Serum Concentration (nM)
100
10
1
0
24
48
72
96
120
Time after dosing (hr)

FIG. 4A


Mouse PK (1mg/kg IV)
xCtrl-hIL21 v2
xCtrl-hIL21
Serum Concentration (nM)
100
10
1
0
24
48
72
96
120
Time after dosing (hr)

FIG. 4D


Mouse PK (1mg/kg IV)
xCtrl-hIL21 v31
xCtrl-hIL21
AUC Ratio: 4.2
Serum Concentration (nM)
1000
100
10
1
0.1
0
48
96
144
192
Time after dosing (hr)

FIG. 4C


Mouse PK (1mg/kg SubQ)
xCtrl-hIL21 v2
xCtrl-hIL21 v31
xCtrl-hIL21
Serum Concentration (nM)
100
10
1
0
48
96
144
192
Time after dosing (hr)

FIG. 5A


Wild type IL-21
(PBMC)
CD8+ T cells
CD4+ T cells
NK cells
B cells
%STAT3+
100
80
60
40
20
0
0.0001
0.01
1
100
10000
Concentration (nM)

FIG. 5B


Wild type IL-21
(Whole Blood)
CD8+ T cells
CD4+ T cells
NK cells
B cells
% STAT3+
100
80
60
40
20
0
0.0001
0.01
1
100
10000
Concentration (nM)

FIG. 6


IL-21 with WT potency
Targeted IL-21 fusion protein
Untargeted IL-21 fusion protein
Binding/ activation
Weak to no binding/activity
Binding/ activation
Weak to no binding
no binding
All IL-21R+ cells (CD8+ and CD8-)
Undesired IL-21R+ cells (CD8-)
Desired IL-21R+ cells (CD8+)
Desired IL-21R+ cells (CD8+)
Activation: + - + -
IL-21 receptor
Target antigen (i.e. CD8)
Anti-antigen (i.e. CD8) antibody
Activation
IL-21 polypeptide with wild-type potency
IL-21 polypeptide with attenuated potency
Control antibody

FIG. 7


CD8 T cell activation
% pStat3+
50
40
30
20
10
0
0.00010.001 0.01 0.1 1 10 100 1000
Concentration (nM)
xhCD8-hIL21
xhCD8-hIL21 v1
xhCD8-hIL21 v2
xhCD8-hIL21 v6

FIG. 8A


xhCD8-IL21 v2.1
CD8+
CD4+
B cells
NK cells
% pStat3+
Concentration (nM)
0
20
40
60
80
100
0.0001
0.001
0.01
0.1
1
10
100
1000

FIG. 8B


xhCD8-IL21 v2.2
CD8+
CD4+
B cells
NK cells
% pStat3+
Concentration (nM)
0
20
40
60
80
100
0.0001
0.001
0.01
0.1
1
10
100
1000

FIG. 8C


xhCD8-IL21 v2.3
CD8+
CD4+
B cells
NK cells
% pStat3+
Concentration (nM)
0
20
40
60
80
100
0.0001
0.001
0.01
0.1
1
10
100
1000

FIG. 8D


xhCD8-IL21 v2.4
CD8+
CD4+
B cells
NK cells
% pStat3+
Concentration (nM)
0
20
40
60
80
100
0.0001
0.001
0.01
0.1
1
10
100
1000

FIG. 8E


xhCD8-IL21 v2.5
CD8+
CD4+
B cells
NK cells
% pStat3+
Concentration (nM)
0
20
40
60
80
100
0.0001
0.001
0.01
0.1
1
10
100
1000

FIG. 8F


xhCD8-IL21 v2.6
CD8
CD4
B Cells
NK
% pStat3+
nM
0
20
40
60
80
100
0.0001
0.001
0.01
0.1
1
10
100
1000

FIG. 9A


CD8 T cells
%pStat3+
0
20
40
60
80
0.00010.001 0.01 0.1 1 10 100 1000
Concentration (nM)
Fc-hIL21 v2
xhCD8.1-hIL21 v2.8
xhCD8.1-hIL21 v2.9
xhCD8.1-hIL21 v2.10
xhCD8.1-hIL21 v2.11
xhCD8.1-hIL21 v2.12
xhCD8.1-hIL21 v2.14
xhCD8.1-hIL21 v2.16

FIG. 9B


CD4 T cells
%pStat3+
0
20
40
60
80
0.00010.001 0.01 0.1 1 10 100 1000
Concentration (nM)
Fc-hIL21 v2
xhCD8.1-hIL21 v2.8
xhCD8.1-hIL21 v2.9
xhCD8.1-hIL21 v2.10
xhCD8.1-hIL21 v2.11
xhCD8.1-hIL21 v2.12
xhCD8.1-hIL21 v2.14
xhCD8.1-hIL21 v2.16

FIG. 9C


CD8 T cells
%pStat3+
Concentration (nM)
0.00010.001 0.01 0.1 1 10 100 1000
0 20 40 60 80
Fc-hIL21 v2
xhCD8.1-hIL21 v2.4
xhCD8.1-hIL21 v2.18
xhCD8.1-hIL21 v2.19
xhCD8.1-hIL21 v2.20
xhCD8.1-hIL21 v2.23
xhCD8.1-hIL21 v2.24

FIG. 9D


CD4 T cells
%pStat3+
Concentration (nM)
0.00010.001 0.01 0.1 1 10 100 1000
0 20 40 60
Fc-hIL21 v2
xhCD8.1-hIL21 v2.4
xhCD8.1-hIL21 v2.18
xhCD8.1-hIL21 v2.19
xhCD8.1-hIL21 v2.20
xhCD8.1-hIL21 v2.23
xhCD8.1-hIL21 v2.24

FIG. 9E


CD8 T cells
%pStat3+
Concentration (nM)
0.0001
0.001
0.01
0.1
1
10
100
1000
0
20
40
60
80
Fc-hIL21 v2
xhCD8.1-hIL21 v2.44
xhCD8.1-hIL21 v2.48
xhCD8.1-hIL21 v2.49
xhCD8.1-hIL21 v2.50
xhCD8.1-hIL21 v2.51
xhCD8.1-hIL21 v2.52
xhCD8.1-hIL21 v2.53
xhCD8.1-hIL21 v2.54

FIG. 9F


CD4 T cells
%pStat3+
Concentration (nM)
0.0001
0.001
0.01
0.1
1
10
100
1000
0
20
40
60
80
100
Fc-hIL21 v2
xhCD8.1-hIL21 v2.44
xhCD8.1-hIL21 v2.48
xhCD8.1-hIL21 v2.49
xhCD8.1-hIL21 v2.50
xhCD8.1-hIL21 v2.51
xhCD8.1-hIL21 v2.52
xhCD8.1-hIL21 v2.53
xhCD8.1-hIL21 v2.54

FIG. 9G


CD8 T cells
%pStat3+
100
80
60
40
20
0
0.00010.001 0.01 0.1 1 10 100 1000
Concentration (nM)
Fc-hIL21 v2
xhCD8.1-hIL21 v2.4
xhCD8.1-hIL21 v2.5
xhCD8.1-hIL21 v2.51
xhCD8.1-hIL21 v2.56
xhCD8.1-hIL21 v2.57
xhCD8.1-hIL21 v2.58
xhCD8.1-hIL21 v2.59

FIG. 9H


CD4 T cells
%pStat3+
100
80
60
40
20
0
0.00010.001 0.01 0.1 1 10 100 1000
Concentration (nM)
Fc-hIL21 v2
xhCD8.1-hIL21 v2.4
xhCD8.1-hIL21 v2.5
xhCD8.1-hIL21 v2.51
xhCD8.1-hIL21 v2.56
xhCD8.1-hIL21 v2.57
xhCD8.1-hIL21 v2.58
xhCD8.1-hIL21 v2.59

FIG. 10B



FIG. 10A



FIG. 10D



FIG. 10C



FIG. 10E



FIG. 10F





FIG. 11A



FIG. 11C



FIG. 11B

FIG. 12


MC38 Tumor Growth Curves
PBS
0/8 CR
0.3mpk xCtrl-mIL21 v1
1/8 CR
1mpk xCtrl-mIL21 v1
5/8 CR
0.1mpk xmCD8-mIL21 v1.1
5/8 CR
0.3mpk xmCD8-mIL21 v1.1
7/8 CR
1mpk xmCD8-mIL21 v1.1
8/8 CR
Tumor Volume mm^3
Days Post Treatment
0
500
1000
1500
2000
2500
10
20
30

FIG. 13A



FIG. 13B



FIG. 13C



FIG. 13D



FIG. 14A



FIG. 14B



FIG. 14C



FIG. 14D



FIG. 14F



FIG. 14E



FIG. 15B



FIG. 15A



FIG. 15C



FIG. 15D



FIG. 16B



FIG. 16A



FIG. 16C


xhCD8.1-hIL21 v2.65
CD8
CD4
B
NK
% pSTAT3+
100
80
60
40
20
0
0.00010.001 0.01 0.1 1 10 100 1000
Concentration (nM)

FIG. 17A



FIG. 17B



FIG. 17C



FIG. 18


Mouse PK (1mg/kg IV)
xhCD8.1-hIL21 v31.4
xhCD8v11-hIL21 v31.23
xhCD8.1-hIL21 v31.23
xhCD8.1-hIL21 v0.4
Serum Concentration (nM)
1000
100
10
1
0
48
96
144
192
Time after dosing (hr)


hIL21
hIL21 vO.51
hIL21 vO.4
hIL21 vO.23
hIL21 v31
hIL21 v31.51
hIL21 v31.23
hIL21 31.23
250
150
100
75
50
37
25
20
15
10

FIG. 19A

FIG. 19B

Construct	Ret time (min)	Peak width (min)
hIL21 v31.23	4.43	0.22
hIL21 v31.4	4.43	0.22
hIL21 v31.51	4.44	0.21
hIL21 v31	4.44	0.21
hIL21 v0.23	4.58	0.35
hIL21 v0.4	4.52	0.22
hIL21 v0.51	4.52	0.35
hIL21	4.60	0.48



FIG. 20


Mouse PK (1mg/kg IV)
IL-21
IL-21 v31
LLOQ
AUC Ratio: 4.3
Concentration (nM)
100
10
1
0.1
0.01
50
100
150
200
Time post-dose (min)

FIG. 21A





FIG. 21B

FIG. 22


no IL-21
Fc-hIL21v31
xhCD8.1-hIL21 v31.4
xhCD8.1-hIL21 v31.23
IL-2 release (pg/ml)
150
100
50
0

FIG. 23B



FIG. 23A



FIG. 23D



FIG. 23C



FIG. 24


In vivo pSTAT3 in cynomolgus monkeys
xhCD8.1-hIL21 v31.4
pStat3 MFI
1.5
1.0
0.5
0.0
CD8+ T cells
CD4+ T cells
B cells
NK cells

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US22/26584

A. CLASSIFICATION OF SUBJECT MATTER

IPC - INV. A61K 38/20; A61K 47/68; C07K 14/54; C07K 16/28; C07K 16/46; A61K 39/395; C07K 19/00 (2022.01)
ADD.

CPC - INV. A61K 38/20; A61K 47/68; C07K 14/54; C07K 16/2815; C07K 16/46; C07K 19/00; A61K 39/395
ADD. C07K 2319/00; C12N 2501/2321

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y --- A	(SHEN, S et al.) Engineered IL-21 Cytokine Muteins Fused to Anti-PD-1 Antibodies Can Improve CD8+ T Cell Function and Anti-tumor Immunity. Frontiers in Immunology. Epub 8 May 2020, Vol. 11, No. 832; pages 1-15; abstract; page 3, 2nd column, 1st paragraph; page 6, 1st column, 2nd paragraph; supplementary figure 2A; DOI: 10.3389/fimmu.2020.00832	1 --- 2-5, 25-27, 54-56, 62-66, 91-93, 115
Y --- A	(BONDENSGAARD, K et al.) The Existence of Multiple Conformers of Interleukin-21 Directs Engineering of a Superpotent Analogue. The Journal of Biological Chemistry. 10 August 2007, Epub 12 June 2007, Vol. 282, No. 32; pages 23326-23336; fig. 6A; DOI: 10.1074/jbc.M701313200	1 --- 2-5, 25-27, 54-56, 62-66, 91-93
Y --- A	WO 2020/247843 A2 (ASHER BIOTHERAPEUTICS, INC.) 10 December 2020; abstract; paragraphs [0013], [0067]; fig. 4A	1 --- 2-5, 25-27, 54-56, 62-66, 91-93, 115
A	US 2018/0142027 A1 (CHUGAI SEIYAKU KABUSHIKI KAISHA) 24 May 2018; paragraphs [0044], [0048], [0049], embodiment 31	2-5, 25-27, 54-56, 62-66, 91-93, 115
A	(YOUNG, PA et al.) Antibody-Cytokine Fusion Proteins for Treatment of Cancer: Engineering Cytokines for Improved Efficacy and Safety. Seminars in Oncology. October 2014, Epub 12 August 2014, Vol. 41, No. 5; pages 1-19; entire document; DOI: 10.1053/j.seminoncol.2014.08.002	1-5, 25-27, 54-56, 62-66, 91-93, 115

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:
"A" document defining the general state of the art which is not considered to be of particular relevance
"D" document cited by the applicant in the international application
"E" earlier application or patent but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed
"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
18 August 2022 (18.08.2022)	SEP 09 2022
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-8300	Authorized officer Shane Thomas Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US22/26584

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:

   a. ☐ forming part of the international application as filed:

      ☐ in the form of an Annex C/ST.25 text file.

      ☐ on paper or in the form of an image file.

   b. ☐ furnished together with the international application under PCT Rule 13*ter*.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.

   c. ☒ furnished subsequent to the international filing date for the purposes of international search only:

      ☒ in the form of an Annex C/ST.25 text file (Rule 13*ter*.1(a)).

      ☐ on paper or in the form of an image file (Rule 13*ter*.1(b) and Administrative Instructions, Section 713).

2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US22/26584

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 6-24, 28-53, 57-61, 67-90, 94-114, 116-136
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

☐ No protest accompanied the payment of additional search fees.