



(86) Date de dépôt PCT/PCT Filing Date: 2007/11/22
(87) Date publication PCT/PCT Publication Date: 2008/07/10
(85) Entrée phase nationale/National Entry: 2009/06/19
(86) N° demande PCT/PCT Application No.: EP 2007/010122
(87) N° publication PCT/PCT Publication No.: 2008/080455
(30) Priorité/Priority: 2006/12/21 (DE10 2006 060 598.5)


(51) Cl.Int./Int.Cl. *C07D 231/54* (2006.01),
A61K 31/416 (2006.01), *A61K 31/423* (2006.01),
A61K 31/454 (2006.01), *A61K 31/5377* (2006.01),
A61P 35/00 (2006.01), *C07D 261/20* (2006.01),
C07D 413/04 (2006.01), *C07D 413/06* (2006.01),
C07D 413/12 (2006.01), *C07D 413/14* (2006.01)

(71) Demandeur/Applicant:
MERCK PATENT GESELLSCHAFT MIT
BESCHRAENKTER HAFTUNG, DE

(72) Inventeurs/Inventors:
SCHIEMANN, KAI, DE;
FINSINGER, DIRK, DE;
ZENKE, FRANK, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : TETRAHYDROBENZO-ISOXAZOLE ET DERIVE DE TETRAHYDRO-INDAZOLE COMME MODULATEURS
DE LA PROTEINE MOTEUR MITOTIQUE
(54) Title: TETRAHYDROBENZOISOXAZOLE AND TETRAHYDROINDAZOLE DERIVATIVES AS MODULATORS OF
THE MITOTIC MOTOR PROTEIN



(57) **Abrégé/Abstract:**

Compounds of Formula (I), in which A¹, A², R¹, X¹, X², X³, Y, R², Cy, and n have the meanings set forth in claim 1, can be used, among other things, for the treatment of tumors.



ABSTRACT

Compounds of Formula (I), in which A^1 , A^2 , R^1 , X^1 , X^2 , X^3 , Y , R^2 , Cy , and n have the meanings set forth in claim 1, can be used, among other things, for the treatment of tumors.

TETRAHYDROBENZOXISOXAZOLE AND TETRAHYDROINDAZOLE DERIVATIVES AS MODULATORS
OF THE MITOTIC MOTOR PROTEIN

BACKGROUND OF THE INVENTION

5 The invention had the object of finding novel compounds having valuable properties, in particular those which can be used for the preparation of medicaments.

10 The present invention relates to compounds of the formula I and to the use thereof for the treatment and prophylaxis of diseases in which the inhibition, regulation and/or modulation of mitotic motor proteins, in particular the mitotic motor protein Eg5, plays a role, furthermore to pharmaceutical compositions which comprise these compounds.

15 In detail, the present invention relates to compounds of the formula I which which preferably inhibit, regulate and/or modulate one or more mitotic motor proteins, to compositions which comprise these compounds, and to methods for the use thereof for the treatment of diseases and complaints such as angiogenesis, cancer, tumour formation, growth and propagation, arterio-
20 sclerosis, ocular diseases, choroidal neovascularisation and diabetic retinopathy, inflammatory diseases, arthritis, neurodegeneration, restenosis, wound healing or transplant rejection. In particular, the compounds according to the invention are suitable for the therapy or prophylaxis of cancer diseases.

25 During mitosis, various kinesins regulate the formation and dynamics of the spindle apparatus, which is responsible for correct and coordinated alignment and separation of the chromosomes. It has been observed that specific inhibition of a mitotic motor protein – Eg5 – results in collapse of the spindle
30 fibres. The result of this is that the chromosomes can no longer be distributed correctly over the daughter cells. This results in mitotic arrest and can thus cause cell death. Upregulation of the motor protein Eg5 has been described, for example, in tissue from breast lung and colon tumours. Since

Eg5 takes on a mitosis-specific function, it is principally rapidly dividing cells and not fully differentiated cells that are affected by Eg5 inhibition. In addition, Eg5 regulates exclusively the movement of mitotic microtubuli (spindle apparatus) and not that of the cytoskeleton. This is crucial for the side-effect profile of the compounds according to the invention since, for example, neuropathies, as observed in the case of Taxol, do not occur or only do so to a weakened extent. The inhibition of Eg5 by the compounds according to the invention is therefore a relevant therapy concept for the treatment of malignant tumours.

In general, all solid and non-solid tumours can be treated with the compounds of the formula I, such as, for example, monocytic leukaemia, brain, urogenital, lymphatic system, stomach, laryngeal and lung carcinoma, including lung adenocarcinoma and small-cell lung carcinoma. Further examples include prostate, pancreatic and breast carcinoma.

Surprisingly, it has been found that the compounds according to the invention effect specific inhibition of mitotic motor proteins, in particular Eg5. The compounds according to the invention preferably exhibit an advantageous biological activity which can easily be detected in the assays described herein, for example. In such assays, the compounds according to the invention preferably exhibit and cause an inhibiting effect, which is usually documented by IC_{50} values in a suitable range, preferably in the micromolar range and more preferably in the nanomolar range.

As discussed herein, effects of the compound according to the invention are relevant to various diseases. Accordingly, the compounds according to the invention are useful in the prophylaxis and/or treatment of diseases which are influenced by inhibition of one or more mitotic motor proteins, in particular Eg5.

The present invention therefore relates to compounds according to the invention as medicaments and/or medicament active ingredients in the treatment and/or prophylaxis of the said diseases and to the use of compounds

according to the invention for the preparation of a pharmaceutical for the treatment and/or prophylaxis of the said diseases, and also to a method for the treatment of the said diseases comprising the administration of one or more compounds according to the invention to a patient in need of such an administration.

It can be shown that the compounds according to the invention have an advantageous effect in a xenotransplant tumour model.

The host or patient can belong to any mammal species, for example a primate species, particularly humans; rodents, including mice, rats and hamsters; rabbits; horses, cattle, dogs, cats, etc. Animal models are of interest for experimental investigations, providing a model for the treatment of a human disease.


The susceptibility of a certain cell to treatment with the compounds according to the invention can be determined by testing in vitro. Typically, a culture of the cell is combined with a compound according to the invention at various concentrations for a period of time which is sufficient to enable the active ingredients to induce cell death or inhibit migration, usually between approximately one hour and one week. For testing in vitro, cultivated cells from a biopsy sample can be used. The viable cells remaining after the treatment are then counted.

The dose varies depending on the specific compound used, the specific disease, the patient status, etc. Typically, a therapeutic dose is sufficient considerably to reduce the undesired cell population in the target tissue, while the viability of the patient is maintained. The treatment is generally continued until a considerable reduction has occurred, for example at least about a 50% reduction in the cell burden, and can be continued until essentially no undesired cells are detected in the body.

SUMMARY OF THE INVENTION

Compounds of the formula I

5




in which

10

A^1, A^2 , independently of one another, denote N, O or S,


X^1, X^2, X^3 , independently of one another, denote a single bond, NR^3-NR^3 , NR^3 , O, S, or one of the following groups:

15



20

Y denotes $C=O$, SO , SO_2 , $(CR^1_2)_n$,



25

Cy denotes H, a carbocyclic or heterocyclic saturated, unsaturated or aromatic radical, which may be unsubstituted or mono- or polysubstituted by alkyl, Hal, CN, OH; OR, OCF_3 , CF_3 , COOR or by a $(CR^1_2)_n-Y-X^1-(CR^1_2)_n-Q$ group,

30

Q denotes H, alkyl, cycloalkyl, aryl or heteroaryl

R, R^1 , denote H, alkyl, Hal, alkoxy, OH, alkenyl, alkoxyalkyl, hydroxyalkyl,

R^2, R^3 $(CH_2)_n-Q$, $(CH_2)_n-Cy$ or $(CH_2)_n-NR_2$,

Hal denotes F, Br or Cl

n denotes 0, 1, 2, 3, 4, 5, 6, 7, or 8,

m denotes 1 or 2

and

p denotes 0, 1 or 2,

and pharmaceutically usable derivatives, solvates, tautomers, salts and stereoisomers thereof, including mixtures thereof in all ratios.

The invention also relates to the optically active forms, the enantiomers, the racemates, the diastereomers and the hydrates and solvates of these compounds. The term solvates of the compounds is taken to mean adductions of inert solvent molecules onto the compounds of the formula I which form owing to their mutual attractive force. Solvates are, for example, mono- or dihydrates or alkoxides.

The term pharmaceutically usable derivatives is taken to mean, for example, the salts of the compounds according to the invention and also so-called prodrug compounds.

The term prodrug derivatives is taken to mean compounds of the formula I which have been modified by means of, for example, alkyl or acyl groups, sugars or oligopeptides and which are rapidly cleaved in the organism to form the effective compounds according to the invention.

These also include biodegradable polymer derivatives of the compounds according to the invention, as described, for example, in Int. J. Pharm. 115, 61-67 (1995).

Similar compounds are described, for example, in Tetrahedron Lett. **1988**, 29, 5855-5858, Tetrahedron Lett. **2003**, 44, 217-219, J. Org. Chem. **1997**, 62, 4880-4882, J. Org. Chem. **1999**, 64, 6462-6467, Chem. Lett. **1995**,

423-424, J. Org. Chem. **2000**, 65, 5009-5013, Chem. Lett. **2003**, 32, 222-223, US2003149069A1, but are not mentioned in connection with cancer treatments and/or do not contain the features essential to the invention.

5 The expression "effective amount" denotes the amount of a medicament or of a pharmaceutical active ingredient which causes in a tissue, system, animal or human a biological or medical response which is sought or desired, for example, by a researcher or physician.

In addition, the expression "therapeutically effective amount" denotes an
10 amount which causes at least one of the following effects in a human or another mammal (compared with a subject who has not received this amount):

improvement in the treatment, healing, prevention or elimination of a disease, syndrome, condition, complaint, disorder or side-effects or also the
15 reduction in the progress of a disease, complaint or disorder.


The term "therapeutically effective amount" also encompasses the amounts which are effective for increasing or enhancing normal physiological function.

The invention also relates to the use of mixtures of the compounds of the
20 formula I, for example mixtures of two diastereomers, for example in the ratio 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:100 or 1:1000.

These are particularly preferably mixtures of stereoisomeric compounds.

The invention relates to the compounds of the formula I and salts thereof
25 and to a process for the preparation of compounds of the formula I according to the patent claims and pharmaceutically usable derivatives, salts, solvates and stereoisomers thereof, characterised in that a compound of the formula II


30



in which A^1 , A^2 , R^1 , and X^1 have the meanings indicated above,

is reacted with a compound of the formula III

5



in which

Y , X^2 , R^2 , X^3 and Cy have the meaning indicated above,

10

preferably in the presence of an activating reagent', such as, for example, N-(3-dialkylaminoalkyl)-N'-alkylcarbodiimide, in particular N-(3dimethylamino-propyl)-N'-ethylcarbodiimide.

15

The mixtures of diastereomers and enantiomers of the compounds of the formula I which may be obtained by the process described above are preferably separated by chromatography or crystallisation.

20

If desired, the bases and acids of the formula I obtained by the process described above are converted into their salts.

25

Above and below, the radicals Hal , R , R^1 , R^2 , R^3 , X^1 , X^2 , X^3 , Y , Q , Cy , m , n and p have the meanings indicated for the formula I, unless expressly indicated otherwise. If individual radicals occur a number of times within a compound, the radicals adopt the meanings indicated, independently of one another.

30

Alkyl is preferably unbranched (linear) or branched, and has 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 C atoms. Alkyl preferably denotes methyl, furthermore ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, furthermore also pentyl, 1-, 2- or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-, 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethyl-

butyl, 1- or 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2- or 1,2,2-trimethylpropyl, further preferably, for example, trifluoromethyl.

Alkyl very particularly preferably denotes alkyl having 1, 2, 3, 4, 5 or 6 C atoms, preferably methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, trifluoromethyl, pentafluoroethyl or 1,1,1-trifluoroethyl. Alkyl also denotes cycloalkyl.


Cycloalkyl preferably denotes cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl, but in particular cyclopentyl.

R , R^1 , R^2 , preferably denotes H, alkyl, CF_3 , OCF_3 , SCN, CN, OH, O alkyl -OCOalkyl, OCOH, Hal, SCF_3 , preferably also t-butyl, -CH(CH₃)CH₂CH₃, isopropyl, ethyl or methyl. In particular, R^1 denotes t-butyl, isopropyl, ethyl, CF_3 , methyl, Br, Cl, SCF_3 , CH(CH₃)CH₂CH₃, n-propyl, OCH₃, SCH₃, n-butyl, -SCN, CH₂CN. R^1 particularly preferably denotes t-butyl, isopropyl, ethyl or CF_3 .

A^1 , A^2 preferably denote, independently of one another, O or N. A^1 and A^2 are particularly preferably not identical.


R^3 preferably denote H, alkyl, hydroxyalkyl, alkoxyalkyl, $(CH_2)_n$ Q or $(CH_2)_nN(R^1)_2$.

X^1 , X^2 preferably denote NR^3 , O or the of the following group




X preferably denotes a single bond or $(CH_2)_n$

Y preferably denotes C=O or



5 Cy preferably denotes substituted or unsubstituted cyclopentyl, aryl or heteroaryl. In particular, Cy denotes one of the following groups

10



in which q denotes 1 or 2

15

Preference is given to compounds of the formula I in which the $Y-X^2-(CR^2)_n-X$ group represents a single bond.

Q preferably denotes aryl or heteroaryl

20

u preferably denotes 0, 1 or 2.

m preferably denotes 1.

p preferably denotes 1.

25

Aryl preferably denotes phenyl, naphthyl or biphenyl, each of which is unsubstituted or mono-, di- or trisubstituted by Hal, A, OH, OA, NH₂, NO₂, CN, COOH, COOA, CONH₂, NHCOA, NHCONH₂, NHSO₂A, CHO, COA, SO₂NH₂, SO₂A, -CH₂-COOH or -OCH₂-COOH.

30

Aryl preferably denotes phenyl, o-, m- or p-tolyl, o-, m- or p-ethylphenyl, o-, m- or p-propylphenyl, o-, m- or p-isopropylphenyl, o-, m- or p-tert-butylphenyl, o-, m- or p-hydroxyphenyl, o-, m- or p-methoxyphenyl, o-, m- or p-nitrophenyl, o-, m- or p-aminophenyl, o-, m- or p-(N-methylamino)phenyl, o-, m- or p-(N-methylaminocarbonyl)phenyl, o-, m- or p-acetamidophenyl, o-, m- or p-methoxyphenyl, o-, m- or p-ethoxyphenyl, o-, m- or p-ethoxycarbonylphenyl, o-, m- or p-(N,N-dimethylamino)phenyl, o-, m- or p-(N,N-dimethyl-

aminocarbonyl)phenyl, o-, m- or p-(N-ethylamino)phenyl, o-, m- or p-(N,N-diethylamino)phenyl, o-, m- or p-fluorophenyl, o-, m- or p-bromophenyl, o-, m- or p-chlorophenyl, o-, m- or p-(methylsulfonamido)phenyl, o-, m- or p-(methylsulfonyl)phenyl, further preferably 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-difluorophenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dichlorophenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dibromophenyl, 2,4- or 2,5-dinitrophenyl, 2,5- or 3,4-dimethoxyphenyl, 3-nitro-4-chlorophenyl, 3-amino-4-chloro-, 2-amino-3-chloro-, 2-amino-4-chloro-, 2-amino-5-chloro- or 2-amino-6-chlorophenyl, 2-nitro-4-N,N-dimethylamino- or 3-nitro-4-N,N-dimethylaminophenyl, 2,3-diaminophenyl, 2,3,4-, 2,3,5-, 2,3,6-, 2,4,6- or 3,4,5-trichlorophenyl, 2,4,6-trimethoxyphenyl, 2-hydroxy-3,5-dichlorophenyl, p-iodophenyl, 3,6-dichloro-4-aminophenyl, 4-fluoro-3-chlorophenyl, 2-fluoro-4-bromophenyl, 2,5-difluoro-4-bromophenyl, 3-bromo-6-methoxyphenyl, 3-chloro-6-methoxyphenyl, 3-chloro-4-acetamidophenyl, 3-fluoro-4-methoxyphenyl, 3-amino-6-methylphenyl, 3-chloro-4-acetamidophenyl or 2,5-dimethyl-4-chlorophenyl.

Heteroaryl preferably denotes a mono- or bicyclic aromatic heterocycle having one or more N, O and/or S atoms which is unsubstituted or mono-, di- or trisubstituted by Hal, A, NO₂, NHA, NA₂, OA, COOA or CN.

Heteroaryl particularly preferably denotes a monocyclic saturated or aromatic heterocycle having one N, S or O atom, which may be unsubstituted or mono-, di- or trisubstituted by Hal, A, NHA, NA₂, NO₂, COOA or benzyl.

Irrespective of further substitutions, unsubstituted heteroaryl denotes, for example, 2- or 3-furyl, 2- or 3-thienyl, 1-, 2- or 3-pyrrolyl, 1-, 2, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or 5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 3- or 4-pyridazinyl, pyrazinyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 4- or 5-isoindolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyra-

zoyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or 7- benzisoxazolyl, 2-, 4-,
 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-, 5-, 6- or 7-
 benz-2,1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolyl, 1-, 3-, 4-, 5-, 6-, 7-
 or 8-isoquinolyl, 3-, 4-, 5-, 6-, 7- or 8-cinnolinyl, 2-, 4-, 5-, 6-, 7- or 8-quinazo-
 5 linyl, 5- or 6-quinoxaliny, 2-, 3-, 5-, 6-, 7- or 8-2H-benzo-1,4-oxazinyl, further
 preferably 1,3-benzodioxol-5-yl, 1,4-benzodioxan-6-yl, 2,1,3-benzothiadiazo-
 4- or -5-yl or 2,1,3-benzoxadiazol-5-yl.

Hal preferably denotes F, Cl or Br particularly preferably F or Cl.


10

Throughout the invention, all radicals which occur more than once may be
 identical or different, i.e. are independent of one another.

The compounds of the formula I can have one or more chiral centres and
 15 therefore exist in various stereoisomeric forms. The formula I encompasses
 all these forms.


Particularly preferred compounds of the formula I are those of the sub-
 formulae IA to IC:

20




IA

25



IB

30



IC

5

in which

R^1 , X^1 , X^2 , X^3 , Y , R^2 and Cy have the meanings indicated above

10

The compounds of the formula I and also the starting materials for their preparation are, in addition, prepared by methods known per se, as described in the literature (for example in the standard works, such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart), to be precise under reaction conditions which are known and suitable for the said reactions. Use may also be made here of variants known per se which are not mentioned here in greater detail.

15

20

If desired, the starting materials may also be formed in situ so that they are not isolated from the reaction mixture, but instead are immediately converted further into the compounds of the formula I.

25

The reaction is generally carried out in an inert solvent, preferably in the presence of an activating reagent' such as N-(3-alkylaminoalkyl)-N'-alkylcarbodiimide, in particular N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide. Depending on the conditions used, the reaction time is between a few minutes and 14 days, the reaction temperature is between about 0° and 180°, normally between 0° and 100°, particularly preferably between 0°C and 70°C.

30

Suitable inert solvents are, for example, hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichloroethylene, 1,2-dichloroethane, carbon tetrachloride, chloroform or dichloromethane, or mixtures of the said solvents.

If desired, a functionally modified amino and/or hydroxyl group in a compound of the formula I can be liberated by solvolysis or hydrogenolysis by conventional methods. This can be carried out, for example, using NaOH or KOH in water, water/THF or water/dioxane at temperatures between 0 and 100°.

The reduction of an ester to the aldehyde or alcohol or the reduction of a nitrile to the aldehyde or amine is carried out by methods as are known to the person skilled in the art and are described in standard works of organic chemistry.

The said compounds according to the invention can be used in their final non-salt form. On the other hand, the present invention also encompasses the use of these compounds in the form of their pharmaceutically acceptable salts, which can be derived from various organic and inorganic acids and bases by procedures known in the art. Pharmaceutically acceptable salt forms of the compounds of the formula I are for the most part prepared by conventional methods. If the compound of the formula I contains a carboxyl group, one of its suitable salts can be formed by reacting the compound with a suitable base to give the corresponding base-addition salt. Such bases are, for example, alkali metal hydroxides, including potassium hydroxide, sodium hydroxide and lithium hydroxide; alkaline earth metal hydroxides, such as barium hydroxide and calcium hydroxide; alkali metal alkoxides, for example potassium ethoxide and sodium propoxide; and various organic bases, such as piperidine, diethanolamine and N-methylglutamine. The aluminium salts of the compounds of the formula I are likewise included. In the case of certain compounds of the formula I, acid-addition salts can be formed by treating these compounds with pharmaceutically acceptable organic and inorganic acids, for example hydrogen halides, such as hydrogen chloride, hydrogen bromide or hydrogen iodide, other mineral acids and corresponding salts thereof, such as sulfate, nitrate or phosphate and the like, and alkyl- and monoarylsulfonates, such as ethanesulfonate, toluene-

sulfonate and benzenesulfonate, and other organic acids and corresponding salts thereof, such as acetate, trifluoroacetate, tartrate, maleate, succinate, citrate, benzoate, salicylate, ascorbate and the like. Accordingly, pharmaceutically acceptable acid-addition salts of the compounds of the formula I include the following: acetate, adipate, alginate, arginate, aspartate, benzoate, benzenesulfonate (besylate), bisulfate, bisulfite, bromide, butyrate, camphorate, camphorsulfonate, caprylate, chloride, chlorobenzoate, citrate, cyclopentanepropionate, digluconate, dihydrogenphosphate, dinitrobenzoate, dodecylsulfate, ethanesulfonate, fumarate, galacterate (from mucic acid), galacturonate, glucoheptanoate, gluconate, glutamate, glycerophosphate, hemisuccinate, hemisulfate, heptanoate, hexanoate, hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, iodide, isethionate, isobutyrate, lactate, lactobionate, malate, maleate, malonate, mandelate, metaphosphate, methanesulfonate, methylbenzoate, monohydrogenphosphate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, oleate, pamoate, pectinate, persulfate, phenylacetate, 3-phenylpropionate, phosphate, phosphonate, phthalate, but this does not represent a restriction.

Furthermore, the base salts of the compounds according to the invention include aluminium, ammonium, calcium, copper, iron(III), iron(II), lithium, magnesium, manganese(III), manganese(II), potassium, sodium and zinc salts, but this is not intended to represent a restriction. Of the above-mentioned salts, preference is given to ammonium; the alkali metal salts sodium and potassium, and the alkaline earth metal salts calcium and magnesium. Salts of the compounds of the formula I which are derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary and tertiary amines, substituted amines, also including naturally occurring substituted amines, cyclic amines, and basic ion exchanger resins, for example arginine, betaine, caffeine, chlorprocaine, choline, N,N'-dibenzylethylenediamine (benzathine), dicyclohexylamine, diethanolamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lidocaine, lysine, meglumine,

N-methyl-D-glucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethanolamine, triethylamine, trimethylamine, tripropylamine and tris(hydroxymethyl)methylamine (tromethamine), but this is not intended to represent a restriction.

5

Compounds of the present invention which contain basic nitrogen-containing groups can be quaternised using agents such as (C₁-C₄)alkyl halides, for example methyl, ethyl, isopropyl and tert-butyl chloride, bromide and iodide; di(C₁-C₄)alkyl sulfates, for example dimethyl, diethyl and diamyl sulfate; 10 (C₁₀-C₁₈)alkyl halides, for example decyl, dodecyl, lauryl, myristyl and stearyl chloride, bromide and iodide; and aryl(C₁-C₄)alkyl halides, for example benzyl chloride and phenethyl bromide. Both water- and oil-soluble compounds according to the invention can be prepared using such salts.

15

The above-mentioned pharmaceutical salts which are preferred include acetate, trifluoroacetate, besylate, citrate, fumarate, gluconate, hemisuccinate, hippurate, hydrochloride, hydrobromide, isethionate, mandelate, meglumine, nitrate, oleate, phosphonate, pivalate, sodium phosphate, stearate, sulfate, sulfosalicylate, tartrate, thiomalate, tosylate and trometh- 20 amine, but this is not intended to represent a restriction.

25

The acid-addition salts of basic compounds of the formula I are prepared by bringing the free base form into contact with a sufficient amount of the desired acid, causing the formation of the salt in a conventional manner. The free base can be regenerated by bringing the salt form into contact with a base and isolating the free base in a conventional manner. The free base forms differ in a certain respect from the corresponding salt forms thereof with respect to certain physical properties, such as solubility in polar solvents; for the purposes of the invention, however, the salts otherwise correspond to the respective free base forms thereof. 30

As mentioned, the pharmaceutically acceptable base-addition salts of the compounds of the formula I are formed with metals or amines, such as alkali

metals and alkaline earth metals or organic amines. Preferred metals are sodium, potassium, magnesium and calcium. Preferred organic amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, N-methyl-D-glucamine and procaine.

5

The base-addition salts of acidic compounds according to the invention are prepared by bringing the free acid form into contact with a sufficient amount of the desired base, causing the formation of the salt in a conventional manner. The free acid can be regenerated by bringing the salt form into contact with an acid and isolating the free acid in a conventional manner. The free acid forms differ in a certain respect from the corresponding salt forms thereof with respect to certain physical properties, such as solubility in polar solvents; for the purposes of the invention, however, the salts otherwise correspond to the respective free acid forms thereof.

15

If a compound according to the invention contains more than one group which is capable of forming pharmaceutically acceptable salts of this type, the invention also encompasses multiple salts. Typical multiple salt forms include, for example, bitartrate, diacetate, difumarate, dimeglumine, diphosphate, disodium and trihydrochloride, but this is not intended to represent a restriction.

20

With regard to that stated above, it can be seen that the expression "pharmaceutically acceptable salt" in the present connection is taken to mean an active ingredient which comprises a compound of the formula I in the form of one of its salts, in particular if this salt form imparts improved pharmacokinetic properties on the active ingredient compared with the free form of the active ingredient or any other salt form of the active ingredient used earlier. The pharmaceutically acceptable salt form of the active ingredient can also provide this active ingredient for the first time with a desired pharmacokinetic property which it did not have earlier and can even have a positive influence on the pharmacodynamics of this active ingredient with respect to its therapeutic efficacy in the body.

25

30

5 The invention furthermore relates to medicaments comprising at least one compound of the formula I and/or pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios, and optionally excipients and/or adjuvants.

Pharmaceutical formulations can be administered in the form of dosage units which comprise a predetermined amount of active ingredient per dosage unit. Such a unit can comprise, for example, 0.5 mg to 1 g, preferably 1 mg to 700 mg, particularly preferably 5 mg to 100 mg, of a compound according to the invention, depending on the condition treated, the method of administration and the age, weight and condition of the patient, or pharmaceutical formulations can be administered in the form of dosage units which comprise a predetermined amount of active ingredient per dosage unit. Preferred dosage unit formulations are those which comprise a daily dose or part-dose, as indicated above, or a corresponding fraction thereof of an active ingredient. Furthermore, pharmaceutical formulations of this type can be prepared using one of the processes which are generally known in the pharmaceutical art.

20 Pharmaceutical formulations can be adapted for administration via any desired suitable method, for example by oral (including buccal or sublingual), rectal, nasal, topical (including buccal, sublingual or transdermal), vaginal or parenteral (including subcutaneous, intramuscular, intravenous or intradermal) methods. Such formulations can be prepared using all processes known in the pharmaceutical art by, for example, combining the active ingredient with the excipient(s) or adjuvant(s).

Pharmaceutical formulations adapted for oral administration can be administered as separate units, such as, for example, capsules or tablets; powders or granules; solutions or suspensions in aqueous or non-aqueous liquids; edible foams or foam foods; or oil-in-water liquid emulsions or water-in-oil liquid emulsions.

Thus, for example, in the case of oral administration in the form of a tablet or capsule, the active-ingredient component can be combined with an oral, non-toxic and pharmaceutically acceptable inert excipient, such as, for example, ethanol, glycerol, water and the like. Powders are prepared by comminuting the compound to a suitable fine size and mixing it with a pharmaceutical excipient comminuted in a similar manner, such as, for example, an edible carbohydrate, such as, for example, starch or mannitol. A flavour, preservative, dispersant and dye may likewise be present.

Capsules are produced by preparing a powder mixture as described above and filling shaped gelatine shells therewith. Glidants and lubricants, such as, for example, highly disperse silicic acid, talc, magnesium stearate, calcium stearate or polyethylene glycol in solid form, can be added to the powder mixture before the filling operation. A disintegrant or solubiliser, such as, for example, agar-agar, calcium carbonate or sodium carbonate, may likewise be added in order to improve the availability of the medicament after the capsule has been taken.

In addition, if desired or necessary, suitable binders, lubricants and disintegrants as well as dyes can likewise be incorporated into the mixture. Suitable binders include starch, gelatine, natural sugars, such as, for example, glucose or beta-lactose, sweeteners made from maize, natural and synthetic rubber, such as, for example, acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. The lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. The disintegrants include, without being restricted thereto, starch, methylcellulose, agar, bentonite, xanthan gum and the like. The tablets are formulated by, for example, preparing a powder mixture, granulating or dry-pressing the mixture, adding a lubricant and a disintegrant and pressing the entire mixture to give tablets. A powder mixture is prepared by mixing the compound comminuted in a suitable manner with a diluent or a base, as described above, and optionally with a binder, such as, for example, car-

boxymethylcellulose, an alginate, gelatine or polyvinylpyrrolidone, a dissolution retardant, such as, for example, paraffin, an absorption accelerator, such as, for example, a quaternary salt, and/or an absorbant, such as, for example, bentonite, kaolin or dicalcium phosphate. The powder mixture can be
5 granulated by wetting it with a binder, such as, for example, syrup, starch paste, acacia mucilage or solutions of cellulose or polymer materials and pressing it through a sieve. As an alternative to granulation, the powder mixture can be run through a tableting machine, giving lumps of non-uniform shape which are broken up to form granules. The granules can be lubricated
10 by addition of stearic acid, a stearate salt, talc or mineral oil in order to prevent sticking to the tablet casting moulds. The lubricated mixture is then pressed to give tablets. The compounds according to the invention can also be combined with a free-flowing inert excipient and then pressed directly to give tablets without carrying out the granulation or dry-pressing steps. A
15 transparent or opaque protective layer consisting of a shellac sealing layer, a layer of sugar or polymer material and a gloss layer of wax may be present. Dyes can be added to these coatings in order to be able to differentiate between different dosage units.

20 Oral liquids, such as, for example, solution, syrups and elixirs, can be prepared in the form of dosage units so that a given quantity comprises a pre-specified amount of the compound. Syrups can be prepared by dissolving the compound in an aqueous solution with a suitable flavour, while elixirs are prepared using a non-toxic alcoholic vehicle. Suspensions can be formulated
25 by dispersion of the compound in a non-toxic vehicle. Solubilisers and emulsifiers, such as, for example, ethoxylated isostearyl alcohols and polyoxyethylene sorbitol ethers, preservatives, flavour additives, such as, for example, peppermint oil or natural sweeteners or saccharin, or other artificial sweeteners and the like, can likewise be added.

30

The dosage unit formulations for oral administration can, if desired, be encapsulated in microcapsules. The formulation can also be prepared in such a

way that the release is extended or retarded, such as, for example, by coating or embedding of particulate material in polymers, wax and the like.

5 The compounds of the formula I and salts, solvates and physiologically functional derivatives thereof can also be administered in the form of liposome delivery systems, such as, for example, small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from various phospholipids, such as, for example, cholesterol, stearylamine or phosphatidylcholines.

10

The compounds of the formula I and the salts, solvates and physiologically functional derivatives thereof can also be delivered using monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds can also be coupled to soluble polymers as targeted medicament carriers. Such polymers may encompass polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidophenol, polyhydroxyethyl-aspartamidophenol or polyethylene oxide polylysine, substituted by palmitoyl radicals. The compounds may furthermore be coupled to a class of biodegradable polymers which are suitable for achieving controlled release of a medicament, for example polylactic acid, poly-epsilon-caprolactone, polyhydroxybutyric acid, polyorthoesters, polyacetals, polydihydroxypyranes, polycyanoacrylates and crosslinked or amphipathic block copolymers of hydrogels.

20

25 Pharmaceutical formulations adapted for transdermal administration can be administered as independent plasters for extended, close contact with the epidermis of the recipient. Thus, for example, the active ingredient can be delivered from the plaster by iontophoresis, as described in general terms in Pharmaceutical Research, 3(6), 318 (1986).

30

Pharmaceutical compounds adapted for topical administration can be formulated as ointments, creams, suspensions, lotions, powders, solutions, pastes, gels, sprays, aerosols or oils.

For the treatment of the eye or other external tissue, for example mouth and skin, the formulations are preferably applied as topical ointment or cream. In the case of formulation to give an ointment, the active ingredient can be employed either with a paraffinic or a water-miscible cream base. Alternatively, the active ingredient can be formulated to give a cream with an oil-in-water cream base or a water-in-oil base.

Pharmaceutical formulations adapted for topical application to the eye include eye drops, in which the active ingredient is dissolved or suspended in a suitable carrier, in particular an aqueous solvent.

Pharmaceutical formulations adapted for topical application in the mouth encompass lozenges, pastilles and mouthwashes.

Pharmaceutical formulations adapted for rectal administration can be administered in the form of suppositories or enemas.

Pharmaceutical formulations adapted for nasal administration in which the carrier substance is a solid comprise a coarse powder having a particle size, for example, in the range 20-500 microns, which is administered in the manner in which snuff is taken, i.e. by rapid inhalation via the nasal passages from a container containing the powder held close to the nose. Suitable formulations for administration as nasal spray or nose drops with a liquid as carrier substance encompass active-ingredient solutions in water or oil.

Pharmaceutical formulations adapted for administration by inhalation encompass finely particulate dusts or mists, which can be generated by various types of pressurised dispensers with aerosols, nebulisers or insufflators.

Pharmaceutical formulations adapted for vaginal administration can be administered as pessaries, tampons, creams, gels, pastes, foams or spray formulations.

Pharmaceutical formulations adapted for parenteral administration include aqueous and non-aqueous sterile injection solutions comprising antioxidants, buffers, bacteriostatics and solutes, by means of which the formulation is rendered isotonic with the blood of the recipient to be treated; and aqueous and non-aqueous sterile suspensions, which may comprise suspension media and thickeners. The formulations can be administered in single-dose or multidose containers, for example sealed ampoules and vials, and stored in freeze-dried (lyophilised) state, so that only the addition of the sterile carrier liquid, for example water for injection purposes, immediately before use is necessary. Injection solutions and suspensions prepared in accordance with the recipe can be prepared from sterile powders, granules and tablets.

It goes without saying that, in addition to the above particularly mentioned constituents, the formulations may also comprise other agents usual in the art with respect to the particular type of formulation; thus, for example, formulations which are suitable for oral administration may comprise flavours.

A therapeutically effective amount of a compound of the formula I depends on a number of factors, including, for example, the age and weight of the animal, the precise condition which requires treatment, and its severity, the nature of the formulation and the method of administration, and is ultimately determined by the treating doctor or vet. However, an effective amount of a compound according to the invention for the treatment of neoplastic growth, for example colon or breast carcinoma, is generally in the range from 0.1 to 100 mg/kg of body weight of the recipient (mammal) per day and particularly typically in the range from 1 to 10 mg/kg of body weight per day. Thus, the actual amount per day for an adult mammal weighing 70 kg is usually between 70 and 700 mg, where this amount can be administered as a single dose per day or more usually in a series of part-doses (such as, for example, two, three, four, five or six) per day, so that the total daily dose is the same. An effective amount of a salt or solvate or of a physiologically functional derivative thereof can be determined as the fraction of the effective amount

of the compound according to the invention *per se*. It can be assumed that similar doses are suitable for the treatment of other conditions mentioned above.

- 5 The invention furthermore relates to medicaments comprising at least one compound of the formula I and/or pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios, and at least one further medicament active ingredient.
- 10 The invention also relates to a set (kit) consisting of separate packs of
- (a) an effective amount of a compound of the formula I and/or pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios,
- and
- 15 (b) an effective amount of a further medicament active ingredient.

The set comprises suitable containers, such as boxes, individual bottles, bags or ampoules. The set may, for example, comprise separate ampoules, each containing an effective amount of a compound of the formula I and/or

20 pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios,

and an effective amount of a further medicament active ingredient in dissolved or lyophilised form.

- 25 The medicaments from Table 1 are preferably, but not exclusively, combined with the compounds of the formula I. A combination of the formula I and medicaments from Table 1 can also be combined with compounds of the formula V.

5

10

15

20

25

30

Table 1.		
Alkylating agents	Cyclophosphamide Busulfan Ifosfamide Melphalan Hexamethylmelamine Thiotepa Chloroambucil Dacarbazine Carmustine	Lomustine Procarbazine Altretamine Estramustine phosphate Mechloroethamine Streptozocin Temozolomide Semustine
Platinum agents	Cisplatin Oxaliplatin Spiroplatin Carboxyphthalatoplatinum Tetraplatin Ormiplatin Iproplatin	Carboplatin ZD-0473 (AnorMED) Lobaplatin (Aetema) Satraplatin (Johnson Matthey) BBR-3464 (Hoffmann- La Roche) SM-11355 (Sumitomo) AP-5280 (Access)
Antimetabolites	Azacytidine Gemcitabine Capecitabine 5-Fluorouracil Floxuridine 2-Chlorodeoxyadenosine 6-Mercaptopurine 6-Thioguanine Cytarabine 2-Fluorodeoxycytidine Methotrexate Idatrexate	Tomudex Trimetrexate Deoxycoformycin Fludarabine Pentostatin Raltitrexed Hydroxyurea Decitabine (SuperGen) Clofarabine (Bioenvision) Irofulven (MGI Pharma) DMDC (Hoffmann-La Roche) Ethynylcytidine (Taiho)
Topoisomerase inhibitors	Amsacrine Epirubicin Etoposide Teniposide or mitoxantrone Irinotecan (CPT-11) 7-Ethyl-10- hydroxycamptothecin Topotecan Dexrazoxanet (TopoTarget) Pixantrone (Novuspharma) Rebeccamycin analogue (Exelixis)	Rubitecan (SuperGen) Exatecan mesylate (Daiichi) Quinamed (ChemGenex) Gimatecan (Sigma- Tau) Diflomotecan (Beaufour- Ipsen) TAS-103 (Taiho) Elsamitrucin (Spectrum) J-107088 (Merck & Co) BNP-1350 (BioNumerik) CKD-602 (Chong Kun Dang) KW-2170 (Kyowa Hakko)

	BBR-3576 (Novuspharma)	
5	Antitumour antibiotics	Dactinomycin (Actinomycin D) Doxorubicin (Adriamycin) Deoxyrubicin Valrubicin Daunorubicin (Daunomycin) Epirubicin Therarubicin Idarubicin Rubidazon Plicamycin Porfiromycin Cyanomorpholinodoxo-rubicin Mitoxantrone (Novantron)
10		Amonafide Azonafide Anthrapyrazole Oxantrazole Losoxantrone Bleomycin sulfate (Blenoxan) Bleomycinic acid Bleomycin A Bleomycin B Mitomycin C MEN-10755 (Menarini) GPX-100 (Gem Pharmaceuticals)
15	Antimitotic agents	Paclitaxel Docetaxel Colchicine Vinblastine Vincristine Vinorelbine Vindesine Dolastatin 10 (NCI) Rhizoxin (Fujisawa) Mivobulin (Warner-Lambert) Cemadotin (BASF) RPR 109881A (Aventis) TXD 258 (Aventis) Epothilone B (Novartis) T 900607 (Tularik) T 138067 (Tularik) Cryptophycin 52 (Eli Lilly) Vinflunine (Fabre) Auristatin PE (Teikoku Hormone) BMS 247550 (BMS) BMS 184476 (BMS) BMS 188797 (BMS) Taxoprexin (Protarga)
20		SB 408075 (GlaxoSmithKline) E7010 (Abbott) PG-TXL (Cell Therapeutics) IDN 5109 (Bayer) A 105972 (Abbott) A 204197 (Abbott) LU 223651 (BASF) D 24851 (ASTA Medica) ER-86526 (Eisai) Combretastatin A4 (BMS) Isohomohalichondrin-B (PharmaMar) ZD 6126 (AstraZeneca) PEG-Paclitaxel (Enzon) AZ10992 (Asahi) IDN-5109 (Indena) AVLB (Prescient NeuroPharma) Azaepothilone B (BMS) BNP- 7787 (BioNumerik) CA-4-Prodrug (OXiGENE) Dolastatin-10 (NrH) CA-4 (OXiGENE)
25		
30	Aromatase inhibitors	Aminoglutethimide Letrozole Anastrozole Formestan Exemestan Atamestan (BioMedicines) YM-511 (Yamanouchi)

	Thymidylate synthase inhibitors	Pemetrexed (Eli Lilly) ZD-9331 (BTG)	Nolatrexed (Eximias) CoFactor™ (BioKeys)
5	DNA antagonists	Trabectedin (PharmaMar) Glufosfamide (Baxter International) Albumin + 32P (Isotope Solutions) Thymectacin (NewBiotics) Edotreotid (Novartis)	Mafosfamide (Baxter International) Apaziquone (Spectrum Pharmaceuticals) O6-Benzylguanine (Paligent)
10	Farnesyl transferase inhibitors	Arglabin (NuOncology Labs) Ionafernib (Schering-Plough) BAY-43-9006 (Bayer)	Tipifarnib (Johnson & Johnson) Perillyl alcohol (DOR BioPharma)
15	Pump inhibitors	CBT-1 (CBA Pharma) Tariquidar (Xenova) MS-209 (Schering AG)	Zosuquidar trihydrochloride (Eli Lilly) Biricodar dicitrate (Vertex)
	Histone acetyl transferase inhibitors	Tacedinaline (Pfizer) SAHA (Aton Pharma) MS-275 (Schering AG)	Pivaloyloxymethyl butyrate (Titan) Depsipeptide (Fujisawa)
20	Metalloproteinase inhibitors Ribonucleoside reductase inhibitors	Neovastat (Aeterna Laboratories) Marimastat (British Biotech) Gallium maltolate (Titan) Triapin (Vion)	CMT -3 (CollaGenex) BMS-275291 (Celltech) Tezacitabine (Aventis) Didox (Molecules for Health)
25	TNF-alpha agonists/ antagonists	Virulizin (Lorus Therapeutics) CDC-394 (Celgene)	Revimid (Celgene)
	Endothelin-A receptor antagonists	Atrasentan (Abbot) ZD-4054 (AstraZeneca)	YM-598 (Yamanouchi)
30	Retinoic acid receptor agonists	Fenretinide (Johnson & Johnson) LGD-1550 (Ligand)	Alitretinoin (Ligand)
	Immuno-modulators	Interferon Oncophage (Antigenics)	Dexosome therapy (Anosys)

5		GMK (Progenics) Adenocarcinoma vaccine (Biomira) CTP-37 (AVI BioPharma) JRX-2 (Immuno-Rx) PEP-005 (Peplin Biotech) Synchrovax vaccines (CTL Immuno) Melanoma vaccine (CTL Immuno) p21-RAS vaccine (GemVax)	Pentrix (Australian Cancer Technology) JSF-154 (Tragen) Cancer vaccine (Intercell) Norelin (Biostar) BLP-25 (Biomira) MGV (Progenics) !3-Alethin (Dovetail) CLL-Thera (Vasogen)
10	Hormonal and antihormonal agents	Oestrogens Conjugated oestrogens Ethinylloestradiol Chlorotrianisene Idenestrol Hydroxyprogesterone caproate Medroxyprogesterone Testosterone Testosterone propionate Fluoxymesterone Methyltestosterone Diethylstilbestrol Megestrol Tamoxifen Toremofin Dexamethasone	Prednisone Methylprednisolone Prednisolone Aminoglutethimide Leuprolide Goserelin Leuporelin Bicalutamide Flutamide Octreotide Nilutamide Mitotan P-04 (Novogen) 2-Methoxyoestradiol (EntreMed) Arzoxifen (Eli Lilly)
15			
20			
25	Photodynamic agents	Talaporfin (Light Sciences) Theralux (Theratechnologies) Motexafin-Gadolinium (Pharmacyclics)	Pd bacteriopheophorbide (Yeda) Lutetium-Texaphyrin (Pharmacyclics) Hypericin
30	Tyrosine kinase inhibitors	Imatinib (Novartis) Leflunomide (Sugen/Pharmacia) ZDI839 (AstraZeneca) Erlotinib (Oncogene Science) Canertinib (Pfizer) Squalamine (Genaera) SU5416 (Pharmacia) SU6668 (Pharmacia) ZD4190 (AstraZeneca) ZD6474 (AstraZeneca)	Kahalide F (PharmaMar) CEP- 701 (Cephalon) CEP-751 (Cephalon) MLN518 (Millenium) PKC412 (Novartis) Phenoxodiol O Trastuzumab (Genentech) C225 (ImClone) rhu-Mab (Genentech) MDX-H210 (Medarex) 2C4 (Genentech) MDX-447 (Medarex)

		Vatalanib (Novartis) PKI166 (Novartis) GW2016 (GlaxoSmithKline) EKB-509 (Wyeth) EKB-569 (Wyeth)	ABX-EGF (Abgenix) IMC-1C11 (ImClone)
5	Various agents	SR-27897 (CCK-A inhibitor, Sanofi- Synthelabo) Tocladesine (cyclic AMP agonist, Ribapharm) Alvocidib (CDK inhibitor, Aventis) CV-247 (COX-2 inhibitor, Ivy Medical) P54 (COX-2 inhibitor, Phytopharm) CapCell™ (CYP450 stimulant, Bavarian Nordic) GCS-100 (gal3 antagonist, GlycoGenesys) G17DT immunogen (gastrin inhibitor, Aphton) Efaproxiral (oxygenator, Allos Therapeutics) PI-88 (heparanase inhibitor, Progen) Tesmifen (histamine antagonist, YM BioSciences) Histamine (histamine H2 receptor agonist, Maxim) Tiazofurin (IMPDH inhibitor, Ribapharm) Cilengitide (integrin antagonist, Merck KGaA) SR-31747 (IL-1 antagonist, Sanofi-Synthelabo) CCI-779 (mTOR kinase inhibitor, Wyeth) Exisulind (PDE-V inhibitor, Cell Pathways) CP-461 (PDE-V inhibitor, Cell Pathways) AG-2037 (GART inhibitor, Pfizer) WX-UK1 (plasminogen activator inhibitor, Willex) PBI-1402 (PMN stimulant,	BCX-1777 (PNP inhibitor, BioCryst) Ranpirnase (ribonuclease stimulant, Alfacell) Galarubicin (RNA synthesis inhibitor, Dong- A) Tirapazamine (reducing agent, SRI International) N-Acetylcysteine (reducing agent, Zambon) R-Flurbiprofen (NF-kappaB inhibitor, Encore) 3CPA (NF-kappaB inhibitor, Active Biotech) Seocalcitol (vitamin D receptor agonist, Leo) 131-I-TM-601 (DNA antagonist, TransMolecular) Eflornithine (ODC inhibitor, ILEX Oncology) Minodronic acid (osteoclast inhibitor, Yamanouchi) Indisulam (p53 stimulant, Eisai) Aplidine (PPT inhibitor, PharmaMar) Rituximab (CD20 antibody, Genentech) Gemtuzumab (CD33 antibody, Wyeth Ayerst) PG2 (haematopoiesis promoter, Pharmagenesis) Immunol™ (triclosan mouthwash, Endo) Triacetyluridine (uridine prodrug, Wellstat) SN-4071 (sarcoma agent, Signature BioScience) TransMID-107™ (immunotoxin, KS
10			
15			
20			
25			
30			

5

10

15

ProMetic LifeSciences)	Biomedix)
Bortezomib (proteasome inhibitor, Millennium)	PCK-3145 (apoptosis promoter, Procyon)
SRL-172 (T-cell stimulant, SR Pharma)	Doranidazole (apoptosis promoter, Pola)
TLK-286 (glutathione-S transferase inhibitor, Telik)	CHS-828 (cytotoxic agent, Leo)
PT-100 (growth factor agonist, Point Therapeutics)	Trans-retinoic acid (differentiator, NIH)
Midostaurin (PKC inhibitor, Novartis)	MX6 (apoptosis promoter, MAXIA)
Bryostatin-1 (PKC stimulant, GPC Biotech)	Apomine (apoptosis promoter, ILEX Oncology)
CDA-II (apoptosis promoter, Everlife)	Urocidin (apoptosis promoter, Bioniche)
SDX-101 (apoptosis promoter, Salmedix)	Ro-31-7453 (apoptosis promoter, La Roche)
Ceflatonin (apoptosis promoter, ChemGenex)	Brostallicin (apoptosis promoter, Pharmacia)

The compounds of the formula I are preferably combined with known anti-cancer agents.

These known anti-cancer agents include the following: oestrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors and other angiogenesis inhibitors. The present compounds are particularly suitable for administration at the same time as radiotherapy. The synergistic effects of inhibition of VEGF in combination with radiotherapy have been described by specialists (see WO 00/61186).

"Oestrogen receptor modulators" refers to compounds which interfere with or inhibit the binding of oestrogen to the receptor, regardless of mechanism. Examples of oestrogen receptor modulators include, but are not limited to, tamoxifen, raloxifene, idoxifene, LY353381, LY 117081, toremifene, fulvestrant, 4-[7-(2,2-dimethyl-1-oxopropoxy-4-methyl-2-[4-[2-(1-piperidinyl)-ethoxy]phenyl]-2H-1-benzopyran-3-yl]phenyl 2,2-dimethylpropanoate, 4,4'-dihydroxybenzophenone-2,4-dinitrophenylhydrazone and SH646.

"Androgen receptor modulators" refers to compounds which interfere with or inhibit the binding of androgens to the receptor, regardless of mechanism.

Examples of androgen receptor modulators include finasteride and other 5 α -reductase inhibitors, nilutamide, flutamide, bicalutamide, liarozole and abiraterone acetate.

"Retinoid receptor modulators" refers to compounds which interfere with or inhibit the binding of retinoids to the receptor, regardless of mechanism.

Examples of such retinoid receptor modulators include bexarotene, tretinoin, 13-cis-retinoic acid, 9-cis-retinoic acid, α -difluoromethylornithine, ILX23-7553, trans-N-(4'-hydroxyphenyl)retinamide and N-4-carboxyphenylretinamide.

"Cytotoxic agents" refers to compounds which result in cell death primarily through direct action on the cellular function or inhibit or interfere with cell myosis, including alkylating agents, tumour necrosis factors, intercalators, microtubulin inhibitors and topoisomerase inhibitors.

Examples of cytotoxic agents include, but are not limited to, tirapazimine, sertenef, cachectin, ifosfamide, tasonermin, lonidamine, carboplatin, altretamine, prednimustine, dibromodulcitol, ranimustine, fotemustine, nedaplatin, oxaliplatin, temozolomide, heptaplatin, estramustine, improsulfan tosylate, trofosfamide, nimustine, dibrospidium chloride, pumitepa, lobaplatin, satraplatin, profiromycin, cisplatin, irofulven, dexifosfamide, cis-aminedichloro(2-methylpyridine)platinum, benzylguanine, glufosfamide, GPX100, (trans,trans,trans)bis-mu-(hexane-1,6-diamine)-mu-[diamineplatinum(II)]bis-[diamine(chloro)platinum(II)] tetrachloride, diarizidinylspermine, arsenic trioxide, 1-(11-dodecylamino-10-hydroxyundecyl)-3,7-dimethylxanthine, zorubicin, idarubicin, daunorubicin, bisantrene, mitoxantrone, pirarubicin, pinafide, valrubicin, amrubicin, antineoplastone, 3'-deamino-3'-morpholino-13-deoxo-10-hydroxycarminomycin, annamycin, galarubicin, elinafide, MEN10755 and 4-demethoxy-3-deamino-3-aziridiny-4-methylsulfonyldaunorubicin (see WO 00/50032).

Examples of microtubulin inhibitors include paclitaxel, vindesine sulfate, 3',4'-didehydro-4'-deoxy-8'-norvincal leukoblastine, docetaxol, rhizoxin, dolastatin, mivobulin isethionate, auristatin, cemadotin, RPR109881, BMS184476, vin-

flunine, cryptophycin, 2,3,4,5,6-pentafluoro-N-(3-fluoro-4-methoxyphenyl)-benzenesulfonamide, anhydrovinblastine, N,N-dimethyl-L-valyl-L-valyl-N-methyl-L-valyl-L-prolyl-L-proline-t-butylamide, TDX258 and BMS188797. Some examples of topoisomerase inhibitors are topotecan, hycaptamine, irinotecan, rubitecan, 6-ethoxypropionyl-3',4'-O-exobenzylidenechartreusin, 9-methoxy-N,N-dimethyl-5-nitropyrazolo[3,4,5-kl]acridine-2-(6H)propan-amine, 1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4':b,7]indolizino[1,2b]quinoline-10,13(9H,15H)-dione, lurtotecan, 7-[2-(N-isopropylamino)ethyl]-(20S)camptothecin, BNP1350, BNPI1100, BN80915, BN80942, etoposide phosphate, teniposide, sobuzoxane, 2'-dimethylamino-2'-deoxyetoposide, GL331, N-[2-(dimethylamino)ethyl]-9-hydroxy-5,6-dimethyl-6H-pyrido[4,3-b]carbazole-1-carboxamide, asulacrine, (5a,5aB,8aa,9b)-9-[2-[N-[2-(dimethylamino)ethyl]-N-methylamino]ethyl]-5-[4-hydroxy-3,5-dimethoxyphenyl]-5,5a,6,8,8a,9-hexahydro-furo(3',4':6,7)naphtho(2,3-d)-1,3-dioxol-6-one, 2,3-(methylenedioxy)-5-methyl-7-hydroxy-8-methoxybenzo[c]phenanthridinium, 6,9-bis[(2-aminoethyl)amino]benzo[g]isoquinoline-5,10-dione, 5-(3-aminopropylamino)-7,10-dihydroxy-2-(2-hydroxyethylaminomethyl)-6H-pyrazolo[4,5,1-de]acridin-6-one, N-[1-[2(diethylamino)ethylamino]-7-methoxy-9-oxo-9H-thioxanthen-4-ylmethyl]formamide, N-(2-(dimethylamino)ethyl)acridine-4-carboxamide, 6-[[2-(dimethylamino)ethyl]amino]-3-hydroxy-7H-indeno[2,1-c]quinolin-7-one and dimesna.

"Antiproliferative agents" include antisense RNA and DNA oligonucleotides such as G3139, ODN698, RVASKRAS, GEM231 and INX3001 and anti-metabolites such as enocitabine, carmofur, tegafur, pentostatin, doxifluridine, trimetrexate, fludarabine, capecitabine, galocitabine, cytarabine ocfosfate, fosteabine sodium hydrate, raltitrexed, paltitrexid, emitefur, tiazofurin, decitabine, nolatrexed, pemetrexed, nelzarabine, 2'-deoxy-2'-methylidenecytidine, 2'-fluoromethylene-2'-deoxycytidine, N-[5-(2,3-dihydrobenzofuryl)sulfonyl]-N'-(3,4-dichlorophenyl)urea, N6-[4-deoxy-4-[N2-[2(E),4(E)-tetradecadienoyl]-glycylamino]-L-glycero-B-L-mannoheptopyranosyl]adenine, aplidine, ecteinascidin, troxacitabine, 4-[2-amino-4-oxo-4,6,7,8-tetrahydro-3H-pyrimidino-[5,4-b]-1,4-thiazin-6-yl-(S)-ethyl]-2,5-thienoyl-L-glutamic acid, aminopterin,

5-fluorouracil, alanosine, 11-acetyl-8-(carbamoyloxymethyl)-4-formyl-6-methoxy-14-oxa-1,11-diazatetracyclo(7.4.1.0.0)tetradeca-2,4,6-trien-9-yl-acetic acid ester, swainsonine, lometrexol, dexrazoxane, methioninase, 2'-cyano-2'-deoxy-N4-palmitoyl-1-B-D-arabinofuranosyl cytosine and 3-amino-pyridine-2-carboxaldehyde thiosemicarbazone. "Antiproliferative agents" also include monoclonal antibodies to growth factors other than those listed under "angiogenesis inhibitors", such as trastuzumab, and tumour suppressor genes, such as p53, which can be released via recombinant virus-mediated gene transfer (see US Patent No. 6,069,134, for example).

10

Particular preference is given to the use of the compound according to the invention for the treatment and prophylaxis of tumour diseases.

15

The tumour is preferably selected from the group of tumours of the squamous epithelium, the bladder, the stomach, the kidneys, of head and neck, the oesophagus, the cervix, the thyroid, the intestine, the liver, the brain, the prostate, the urogenital tract, the lymphatic system, the stomach, the larynx and/or the lung.

20

The tumour is furthermore preferably selected from the group lung adenocarcinoma, small-cell lung carcinomas, pancreatic cancer, glioblastomas, colon carcinoma and breast carcinoma.

25


Preference is furthermore given to the use for the treatment of a tumour of the blood and immune system, preferably for the treatment of a tumour selected from the group of acute myeloid leukaemia, chronic myeloid leukaemia, acute lymphatic leukaemia and/or chronic lymphatic leukaemia.

30

The invention also encompasses a method for the treatment of a patient who has a neoplasm, such as a cancer, by administration of

a) one or more of the compounds of the formula I:

b) and one or more of the compounds of the formula V or acid-addition salts thereof, in particular hydrochlorides:




10

in which Y' and Z' each, independently of one another, denote O or N, R⁶ and R⁷ each, independently of one another, denote H, OH, halogen, OC1-10-alkyl, OCF₃, NO₂ or NH₂, S denotes an integer between 2 and 6, in each case inclusive, and R⁸ and R⁹ are each, independently of one another, preferably in the meta- or para-position and are selected from the group:

15

20




where the first and second compounds are administered simultaneously or within 14 days of one another in amounts which are sufficient to inhibit the growth of the neoplasm.

30


The combination of the compounds of the formula I with the compounds of the formula V and other pentamedine analogues results in a synergistic effect in the inhibition of neoplasias. Combinations comprising the compounds of the formula V are mentioned, for example, in WO 02058684.

The mechanism of action of pentamidine or derivatives thereof has not been clearly explained at present: pentamidine or derivatives thereof appears to have pleiotropic actions since it results in a decrease in DNA, RNA and protein synthesis. It was recently described that pentamidine is a capable inhibitor of PRL1, -2 and 3 phosphatases (Pathak et al., 2002) and tyrosine phosphatases, and overexpression thereof is accompanied by neoplastic malignant tumours in humans. On the other hand, it has been described that pentamidine is a medicament which binds to the DNA minor groove (Puckowska et al., 2004) and is able to exert its action via disturbance of gene expression and/or DNA synthesis.

Other suitable pentamidine analogues include stilbamidine (G-1) and hydroxystilbamidine (G-2) and indole analogues thereof (for example G-3):




(G-1)



(G-2)

and

35



5

(G-3)

10

Each amidine unit may be replaced, independently of one another, by one of the units defined above for R^8 and R^{11} . As in the case of benzimidazoles and pentamidines, salts of stilbamidine, hydroxystilbamidine and indole derivatives thereof are also suitable for the process according to the invention. Preferred salts include, for example, dihydrochloride and methanesulfonate salts.

15

20

25

30

Still other analogues are those which fall under a formula which are provided in one of the US patents No. 5,428,051, 5,521,189, 5,602,172, 5,643,935, 5,723,495, 5,843,980, 6,172,104 and 6,326,395 or the US patent application with the publication no. US 2002/0019437 A1, each of which is incorporated in its entirety by way of reference. Illustrative analogues include 1,5-bis(4'-(N-hydroxyamidino)phenoxy)pentane, 1,3-bis(4'-(N-hydroxyamidino)phenoxy)propane, 1,3-bis(2'-methoxy-4'-(N-hydroxyamidino)phenoxy)propane, 1,4-bis(4'-(N-hydroxyamidino)phenoxy)butane, 1,5-bis(4'-(N-hydroxyamidino)phenoxy)pentane, 1,4-bis(4'-(N-hydroxyamidino)phenoxy)butane, 1,3-bis(4'-(4-hydroxyamidino)phenoxy)propane, 1,3-bis(2'-methoxy-4'-(N-hydroxyamidino)phenoxy)propane, 2,5-bis[4-amidinophenyl]furan, 2,5-bis[4-amidinophenyl]furan bisamide oxime, 2,5-bis[4-amidinophenyl]furan bis-O-methylamide oxime, 2,5-bis[4-amidinophenyl]furan bis-O-ethylamide oxime, 2,8-diamidinodibenzothiophene, 2,8-bis(N-isopropylamidino)carbazole, 2,8-bis(N-hydroxyamidino)carbazole, 2,8-bis(2-imidazoliny)l)dibenzothiophene, 2,8-bis(2-imidazoliny)l)-5,5-dioxodibenzothiophene, 3,7-diamidinodibenzothiophene, 3,7-bis(N-isopropylamidino)dibenzothiophene, 3,7-bis(N-hydroxyamidino)dibenzothiophene, 3,7-diaminodibenzothiophene, 3,7-dibromodibenzothiophene, 3,7-dicyanodibenzothiophene, 2,8-diamidinodibenzo-


furan, 2,8-di(2-imidazoliny)ldibenzofuran, 2,8-di(N-isopropylamidino)dibenzo-
furan, 2,8-di(N-hydroxylamidino)dibenzofuran, 3,7-di(2-imidazoliny)ldibenzo-
furan, 3,7-di(isopropylamidino)dibenzofuran, 3,7-di(A-hydroxylamidino)-
dibenzofuran, 2,8-dicyanodibenzofuran, 4,4'-dibromo-2,2'-dinitrobiphenyl,
5 2-methoxy-2'-nitro-4,4'-dibromobiphenyl, 2-methoxy-2'-amino-4,4'-dibromo-
biphenyl, 3,7-dibromodibenzofuran, 3,7-dicyanodibenzofuran, 2,5-bis(5-
amidino-2-benzimidazolyl)pyrrole, 2,5-bis[5-(2-imidazoliny)l-2-benzimida-
zolyl]pyrrole, 2,6-bis[5-(2-imidazoliny)l-2-benzimidazolyl]pyridine, 1-methyl-
2,5-bis(5-amidino-2-benzimidazolyl)pyrrole, 1-methyl-2,5-bis[5-(2-imidazolyl)-
10 2-benzimidazolyl]pyrrole, 1-methyl-2,5-bis[5-(1,4,5,6-tetrahydro-2-pyrimi-
diny)l-2-benzimidazolyl]pyrrole, 2,6-bis(5-amidino-2-benzimidazolyl)pyridine,
2,6-bis[5-(1,4,5,6-tetrahydro-2-pyrimidiny)l-2-benzimidazolyl]pyridine, 2,5-bis-
(5-amidino-2-benzimidazolyl)furan, 2,5-bis[5-(2-imidazoliny)l-2-benzimidazo-
lyl]furan, 2,5-bis(5-N-isopropylamidino-2-benzimidazolyl)furan, 2,5-bis(4-
15 guanylphenyl)furan, 2,5-bis(4-guanylphenyl)-3,4-dimethylfuran, 2,5-di-p-[2-
(3,4,5,6-tetrahydropyrimidyl)phenyl]furan, 2,5-bis[4-(2-imidazoliny)phenyl]-
furan, 2,5-[bis{4-(2-tetrahydropyrimidiny)}phenyl]-p-(tolylloxy)furan, 2,5-[bis-
{4-(2-imidazoliny)}phenyl]-3-p-(tolylloxy)furan, 2,5-bis{4-[5-(N-2-aminoethyl-
amido)benzimidazol-2-yl]phenyl}furan, 2,5-bis[4-(3a,4,5,6,7,7a-hexahydro-
20 1H-benzimidazol-2-yl)phenyl]furan, 2,5-bis[4-(4,5,6,7-tetrahydro-1H-1,3-
diazepin-2-yl)phenyl]furan, 2,5-bis(4-N,N-dimethylcarboxhydrazidophenyl)-
furan, 2,5-bis{4-[2-(N-2-hydroxyethyl)imidazoliny]phenyl}furan, 2,5-bis[4-(N-
isopropylamidino)phenyl]furan, 2,5-bis{4-[3-(dimethylaminopropyl)amidino]-
phenyl}furan, 2,5-bis{4-[N-(3-aminopropyl)amidino]phenyl}furan, 2,5-bis[2-
25 (imidzaoliny)phenyl]-3,4-bis(methoxymethyl)furan, 2,5-bis[4-N-(dimethyl-
aminoethyl)guanyl]phenylfuran, 2,5-bis{4-[(N-2-hydroxyethyl)guanyl]phenyl}-
furan, 2,5-bis[4-N-(cyclopropylguanyl)phenyl]furan, 2,5-bis[4-(N,N-diethyl-
aminopropyl)guanyl]phenylfuran, 2,5-bis{4-[2-(N-ethylimidazoliny)]phenyl}-
furan, 2,5-bis{4-[N-(3-pentylguanyl)]}phenylfuran, 2,5-bis[4-(2-imidazoliny)-
30 phenyl]-3-methoxyfuran, 2,5-bis[4-(N-isopropylamidino)phenyl]-3-methyl-
furan, bis[5-amidino-2-benzimidazolyl]methane, bis[5-(2-imidazolyl)-2-benz-
imidazolyl]methane, 1,2-bis[5-amidino-2-benzimidazolyl]ethane, 1,2-bis[5-(2-
imidazolyl)-2-benzimidazolyl]ethane, 1,3-bis[5-amidino-2-benzimidazolyl]-

propane, 1,3-bis[5-(2-imidazolyl)-2-benzimidazolyl]propane, 1,4-bis[5-amidino-2-benzimidazolyl]propane, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]butane, 1,8-bis[5-amidino-2-benzimidazolyl]octane, trans-1,2-bis[5-amidino-2-benzimidazolyl]ethene, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]-1-butene, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]-2-butene, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]-1-methylbutane, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]-2-ethylbutane, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]-1-methyl-1-butene, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]-2,3-diethyl-2-butene, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]-1,3-butadiene, 1,4-bis[5-(2-imidazolyl)-2-benzimidazolyl]-2-methyl-1,3-butadiene, bis[5-(2-pyrimidyl)-2-benzimidazolyl]methane, 1,2-bis[5-(2-pyrimidyl)-2-benzimidazolyl]ethane, 1,3-bis[5-amidino-2-benzimidazolyl]propane, 1,3-bis[5-(2-pyrimidyl)-2-benzimidazolyl]propane, 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]butane, 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]-1-butene, 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]-2-butene, 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]-1-methylbutane, 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]-2-ethylbutane, 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]-1-methyl-1-butene, 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]-2,3-diethyl-2-butene, 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]-1,3-butadiene and 1,4-bis[5-(2-pyrimidyl)-2-benzimidazolyl]-2-methyl-1,3-butadiene, 2,4-bis(4-guanylphenyl)pyrimidine, 2,4-bis(4-imidazolin-2-yl)pyrimidine, 2,4-bis[(tetrahydropyrimidinyl-2-yl)phenyl]pyrimidine, 2-(4-[N-i-propylguanyl]phenyl)-4-(2-methoxy-4-[N-i-propylguanyl]phenyl)pyrimidine, 4-(N-cyclopentylamidino)-1,2-phenylenediamine, 2,5-bis[2-(5-amidino)benzimidazolyl]furan, 2,5-bis[2-{5-(2-imidazolino)}benzimidazolyl]furan, 2,5-bis[2-(5-N-isopropylamidino)benzimidazolyl]furan, 2,5-bis[2-(5-N-cyclopentylamidino)benzimidazolyl]furan, 2,5-bis[2-(5-amidino)benzimidazolyl]pyrrole, 2,5-bis[2-{5-(2-imidazolino)}benzimidazolyl]pyrrole, 2,5-bis[2-(5-N-isopropylamidino)benzimidazolyl]pyrrole, 2,5-bis[2-(5-N-cyclopentylamidino)benzimidazolyl]pyrrole, 1-methyl-2,5-bis[2-(5-amidino)benzimidazolyl]pyrrole, 2,5-bis[2-{5-(2-imidazolino)}benzimidazolyl]-1-methylpyrrole, 2,5-bis[2-(5-N-cyclopentylamidino)benzimidazolyl]-1-methylpyrrole, 2,5-bis[2-(5-N-isopropylamidino)benzimidazolyl]thiophene, 2,6-bis[2-{5-(2-imidazolino)}benzimidazolyl]pyridine, 2,6-bis[2-(5-amidino)benzimidazolyl]pyridine, 4,4'-bis-

[2-(5-N-isopropylamidino)benzimidazolyl]-1,2-diphenylethane, 4,4'-bis[2-(5-N-cyclopentylamidino)benzimidazolyl]-2,5-diphenylfuran, 2,5-bis[2-(5-amidino)-benzimidazolyl]benzo[b]furan, 2,5-bis[2-(5-N-cyclopentylamidino)benzimidazolyl]benzo[b]furan, 2,7-bis[2-(5-N-isopropylamidino)benzimidazolyl]-fluorine, 2,5-bis[4-(3-(N-morpholinopropyl)carbamoyl)phenyl]furan, 2,5-bis[4-(2-N,N-dimethylaminoethylcarbamoyl)phenyl]furan, 2,5-bis[4-(3-N,N-dimethylaminopropylcarbamoyl)phenyl]furan, 2,5-bis[4-(3-N-methyl-3-N-phenylaminopropylcarbamoyl)phenyl]furan, 2,5-bis[4-(3-N,N8,N11-trimethylaminopropylcarbamoyl)phenyl]furan, 2,5-bis[3-amidinophenyl]furan, 2,5-bis[3-(N-isopropylamidino)amidinophenyl]furan, 2,5-bis[3-[(N-(2-dimethylaminoethyl)amidino]phenyl]furan, 2,5-bis[4-(N-2,2,2-trichloroethoxycarbonyl)-amidinophenyl]furan, 2,5-bis[4-(N-thioethylcarbonyl)amidinophenyl]furan, 2,5-bis[4-(N-benzyloxycarbonyl)amidinophenyl]furan, 2,5-bis[4-(N-phenoxy-carbonyl)amidinophenyl]furan, 2,5-bis[4-(N-(4-fluoro)phenoxy-carbonyl)-amidinophenyl]furan, 2,5-bis[4-(N-(4-methoxy)phenoxy-carbonyl)amidinophenyl]furan, 2,5-bis[4-(1-acetoxyethoxycarbonyl)amidinophenyl]furan and 2,5-bis[4-(N-(3-fluoro)phenoxy-carbonyl)amidinophenyl]furan. Processes for the preparation of one of the above compounds are described in US patents No. 5,428,051, 5,521,189, 5,602,172, 5,643,935, 5,723,495, 5,843,980, 6,172,104 and 6,326,395 or the US patent application with the publication no. US 2002/0019437 A1.

Pentamidine metabolites are likewise suitable in the antiproliferative combination according to the invention. Pentamidine is rapidly metabolised in the body to at least seven primary metabolites. Some of these metabolites have one or more effects in common with pentamidine pentamidine metabolites have antiproliferative action when combined with a benzimidazole or an analogue thereof.

Seven pentamidine analogues are shown below.



The combinations according to the invention of compounds of the formula I and formula V or analogues thereof and metabolites thereof are suitable for the treatment of neoplasms. Combination therapy can be carried out alone or in combination with another therapy (for example operation, irradiation, chemotherapy, biological therapy). In addition, a person whose risk of developing a neoplasm is greater (for example someone who is genetically predisposed or someone who previously had a neoplasm) can be given prophylactic treatment in order to inhibit or delay neoplasm formation.

30

The invention likewise relates to the combination of kinesin ATPase Eg5/KSP with the compounds of the formula V, pentamidine, analogues thereof and/or metabolites thereof.

The dosage and frequency of administration of each compound in the combination can be controlled independently. For example, one compound may be administered orally three times daily, while the second compound may be administered intramuscularly once per day. The compounds may also be formulated together, leading to administration of both compounds.

The antiproliferative combinations according to the invention can also be provided as components of a pharmaceutical package. The two medicaments can be formulated together or separately and in individual dosage amounts.

Under another aspect, the invention encompasses a for the treatment of a patient who has a neoplasm, such as a cancer, by administration of a compound of the formula (I) and (V) in combination with an antiproliferative agent. Suitable antiproliferative agents encompass those provided in Table 1.


Above and below, all temperatures are indicated in °C. In the following examples, "conventional work-up" means: if necessary, water is added, the pH is adjusted, if necessary, to values between 2 and 10, depending on the constitution of the end product, the mixture is extracted with ethyl acetate or dichloromethane, the phases are separated, the organic phase is dried over sodium sulfate and evaporated, and the product is purified by chromatography on silica gel and/or by crystallisation. R_f values on silica gel; eluent: ethyl acetate/methanol 9:1.

Mass spectrometry (MS): EI (electron impact ionisation) M⁺
 FAB (fast atom bombardment) (M+H)⁺
 ESI (electrospray ionisation) (M+H)⁺
APCI-MS (atmospheric pressure chemical ionisation – mass spectrometry)
(M+H)⁺

Example 1

Synthesis of N-(3-hydroxybenzyl)-4,5,6,7-tetrahydrobenzo[d]isoxazole-3-carboxamide

5



10

The commercially available acid **1** (100 mg, 0.60 mmol), the amine **2** (73.6 mg, 0.60 mmol) and *N*-methylmorpholine (0.07 ml, 0.60 mmol) were dissolved in DMF (5 ml), and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (115 mg, 0.06 mmol) and 1-hydroxybenzotriazole (80.0 mg, 0.06 mmol) were added successively. The mixture was stirred at RT for 15 h and precipitated using water. The residue was filtered off and purified by column chromatography (ethyl acetate/cyclohexane), giving the anide **3** as colourless solid.

15


20

Example 2

Synthesis of N-[2-(2-dimethylaminoethylcarbamoyl)-1-(3-hydroxyphenyl)-ethyl]-4,5,6,7-tetrahydrobenzo[d]isoxazole-3-carboxamide

25

30




b. Compound **4** was obtained analogously to procedure a. from the acid **1** and methyl 3-amino-3-(3-hydroxyphenyl)propionate.

Compound **4** (40 mg, 0.12 mmol) and *N,N*-dimethylethylenediamine (0.5 ml) were stirred at 100°C for 12 h in a pressure flask. Ethyl acetate was added to the cooled solution, the mixture was washed with water, dried, filtered and evaporated to dryness. The residue was recrystallised from EtOH/water, giving a colourless solid, which was identified as compound **5**.

Example 3

Synthesis of *N*-((1*S*,2*S*)-2-methylaminoindan-1-yl)-4,5,6,7-tetrahydrobenzo-[d]isoxazole-3-carboxamide



c. Compound **6** was obtained analogously to procedure a. from the acid **1** and *cis*-1-amino-2-indanol.


Compound **6** (158 mg, 0.53 mmol) was initially introduced in dichloromethane (5 ml), triethylamine (0.06 ml, 0.80 mmol) was added, and methanesulfonyl chloride (0.15 ml, 10.06 mmol, dissolved in 1 ml of DCM) was added dropwise at 0°C. The mixture was subsequently stirred at RT for 12 h. The mixture was evaporated to dryness, the residue was taken up in ethyl acetate and washed with water. The org. phase was dried, filtered and evaporated to dryness. The residue (about 140 mg of crude substance) was employed in the next reaction without further purification.

Half of the crude substance (70 mg) was taken up in methylamine (33% solution in EtOH, 1 ml) and stirred at 100°C for 8 h in a pressure flask. The solution

was evaporated to dryness and purified directly by column chromatography (ethyl acetate/cyclohexane), giving 42 mg of a colourless solid (**7**).

5 Example 4

Synthesis of 3-(4-phenyl-4,5-dihydrooxazol-2-yl)-4,5,6,7-tetrahydrobenzo[d]-isoxazole



d. Compound **8** was obtained analogously to procedure a. from the acid **1** and 2-phenylglycinol.

Compound **8** (106 mg, 0.37 mmol) was dissolved in dichloromethane (5 ml), thionyl chloride (0.06 ml, 0.89 mmol) was added, and the mixture was stirred at 70°C for 2 h in a pressure flask. The batch was allowed to cool, and sat. NaHCO₃ solution was added to the mixture. The org. phase was separated off, dried over Na₂SO₄, filtered and evaporated to dryness. This residue was dissolved in MeOH (5 ml), NaOH (about 15 mg, 0.37 mmol) was added, and the mixture was left to stir at 70°C for a further 2 h in a pressure flask. The batch was allowed to cool and was evaporated to dryness. The residue was taken up in DCM (5 ml) and extracted twice with sat. NaHCO₃ solution. The org. phase was dried over Na₂SO₄, filtered and evaporated to dryness. The product was subsequently crystallised from ethyl acetate/cyclohexane, giving compound **9** as colourless solid.

Example 5


Hydrazines can also be converted into the corresponding carbohydrazides analogously to a. using the acid **1**.

5

Example 6

Synthesis of N-(2-methylbenzyl)-4,5,6,7-tetrahydro-1H-indazole-3-carboxamide

10



15


e. Analogously to a., the commercial product **10** (50 mg, 0.28 mmol) was reacted with 2-methylbenzylamine (34 l, 0.28 mmol) using *N*-methymorpholine (1 eq), *N*-(3-dimethylaminopropyl)-*N*'-ethylcarbodiimide hydrochloride (1 eq) and 1-hydroxybenzotriazole (1 eq). The product crystallised cleanly out of the reaction solution after addition of water, giving the amide **11** as colourless solid.

20

Example 7

Synthesis 5-ethyl-4,5,6,7-tetrahydrobenzo[c]isoxazole-3-carboxylic acid

25



30

f. Sodium ethoxide (14 ml of a 20% solution in EtOH) was initially introduced with ice-cooling, and a solution of 4-ethylcyclohexanone (5.00 ml, 36.3 mmol) and diethyl oxalate (4.91 ml, 36.3 mmol) was slowly added dropwise with stirring and ice-cooling. During the dropwise addition, the temperature was kept below 5°C. The reaction mixture was subsequently allowed to come slowly to

RT and was stirred for a further 15 h. The reaction mixture was poured into a mixture of ice and 10 ml of conc. H_2SO_4 and extracted twice with DCM. The org. phase was washed again with sat. NaHCO_3 soln., dried, filtered and evaporated to dryness. The crude substance was reacted further without further purification (see g.).

g. The crude substance 14 (6.8 g, 30.1 mmol) was dissolved in acetic acid (10 ml), hydroxylammonium chloride (2.09 g, 30.1 mmol), dissolved in 5 ml of water, was slowly added dropwise with ice-cooling. The mixture was subsequently stirred under reflux overnight. After cooling to RT, the reaction mixture was poured into ice/water and neutralised using sat. NaHCO_3 soln. The mixture was then extracted twice with dichloromethane and once with ethyl acetate. The combined org. phases were dried over Na_2SO_4 , filtered and evaporated to dryness. The residue consisted of a small amount of the desired acid and the corresponding ethyl ester, which was discarded.


The aqueous phase was re-acidified using 2 N HCl, extracted a number of times with ethyl acetate. The combined org. phases were dried over Na_2SO_4 , filtered and evaporated, giving compound 15 as yellowish solid.

Example 8

Analogously to f. and g., cyclohexanone was converted into 4,5,6,7-tetrahydrobenzo[c]isoxazole-3-carboxylic acid. Both acids were reacted further analogously to a., b., c. and d.

Example 9

Synthesis of 1-(4,5,6,7-tetrahydrobenzo[d]isoxazol-3-yl)-3-o-tolylurea and N-(4,5,6,7-tetrahydrobenzo[d]isoxazol-3-yl)-2-o-tolylacetamide



h. 2-Oxocyclohexanecarbonitrile **16** (1.00 g, 8.12 mmol) and hydroxylammonium chloride (0.56 g, 8.12 mmol) was taken up in acetic acid (1 ml) and stirred at 60°C for 15 h in a pressure flask. The mixture was evaporated to dryness and purified directly by column chromatography (ethyl acetate/cyclohexane). Compound **17** was isolated as colourless solid.

i. Compound **17** (130 mg, 0.94 mmol) was dissolved in DCM (2 ml), and o-tolyl isocyanate (138 mg, 1.04 mmol) was added at RT. After 12 h at RT, the mixture was evaporated to dryness and purified by column chromatography, giving compound **18**.

j. The amine **17** (130 mg, 0.94 mmol) was taken up in DCM (2 ml), triethylamine was added at RT (0.16 ml, 1.13 mmol), and the mixture was cooled to 0°C. o-Tolylacetyl chloride (166 mg, 0.99 mmol), dissolved in DCM (1 ml), was added dropwise to this mixture and stirred for a further 12 h at RT. The reaction mixture was washed 2x with water, dried, filtered and evaporated to dryness. Work-up by column chromatography (ethyl acetate/cyclohexane) gave the amide **19** as colourless solid.

The following compounds according to the invention are obtained analogously using or corresponding precursors.

5

10

15

20


25

30

48


MOLECULAR STRUCTURES

5




10

10




11

15




12

20



13


25



14


30

49




5

15




10

16




15

17




20

18



25

19




30

20


50

5




21

10



22


15



23


20

24



25

25




30

26


51

5




27

10




28

15




29

20



30

25




31

30


52

5




32

10




33

15




34

20




35

25




36

30



37


5



10


38

15




39

20




40

25



41


30



42


54

5




43

10




44

15




45

20



46

25




47

30


55

5




48

10



49


15



50


20

25



51


30



52


56

5




53

10




54

15




20

55



25


56




30

57


57




58




59



60



61




62




58

5




63

10




64

15




20

65



25


66




30

67


59




68




69




70



71




72



73


60

5




74

10




75

15




76

20



77

25




78

30


61

5




79

10



80


15



20


81

25



82


30



83


62

5




10

84




15

85



20

86




25

87

30


63

5



88


10



Chiral

89


15



Chiral


90

20



91


25



92


30

64




5

93




10

94




15

95




20

96



25

97




30

98

65


5

99




10

100




15

101




20

102



25


103



30


66

5




104

10




15

105




20

106



25

107




30

108


67

5




109

10




110

15




111

20




112

25




113

30




114

68




5

115



10


116



15


20

117



25


118



30


119

69




5

120




10

121




15

122




20

123



25

124




30

125

70


5



10

126


15



20


127

25




30

128




129


71




130




131




132



133




134




135

72



136



137

The following examples relate to medicaments:

Example C: Injection vials

A solution of 100 g of an active ingredient of the formula I and 5 g of di-sodium hydrogenphosphate in 3 l of bidistilled water is adjusted to pH 6.5 using 2 N hydrochloric acid, sterile filtered, transferred into injection vials, lyophilised under sterile conditions and sealed under sterile conditions. Each injection vial contains 5 mg of active ingredient.

Example D: Suppositories

A mixture of 20 g of an active ingredient of the formula I with 100 g of soya lecithin and 1400 g of cocoa butter is melted, poured into moulds and allowed to cool. Each suppository contains 20 mg of active ingredient.

Example E: Solution

A solution is prepared from 1 g of an active ingredient of the formula I, 9.38 g of $\text{NaH}_2\text{PO}_4 \cdot 2 \text{H}_2\text{O}$, 28.48 g of $\text{Na}_2\text{HPO}_4 \cdot 12 \text{H}_2\text{O}$ and 0.1 g of benzalkonium chloride in 940 ml of bidistilled water. The pH is adjusted to 6.8,

and the solution is made up to 1 l and sterilised by irradiation. This solution can be used in the form of eye drops.

Example F: Ointment

5

500 mg of an active ingredient of the formula I are mixed with 99.5 g of Vaseline under aseptic conditions.

Example G: Tablets

10

A mixture of 1 kg of active ingredient of the formula I, 4 kg of lactose, 1.2 kg of potato starch, 0.2 kg of talc and 0.1 kg of magnesium stearate is pressed in a conventional manner to give tablets in such a way that each tablet contains 10 mg of active ingredient.

15

Example H: Dragees

Tablets are pressed analogously to Example E and subsequently coated in a conventional manner with a coating of sucrose, potato starch, talc, tragacanth and dye.

20

Example I: Capsules

2 kg of active ingredient of the formula I are introduced into hard gelatine capsules in a conventional manner in such a way that each capsule contains 20 mg of the active ingredient.

25


Example J: Ampoules

A solution of 1 kg of active ingredient of the formula I in 60 l of bidistilled water is sterile filtered, transferred into ampoules, lyophilised under sterile conditions and sealed under sterile conditions. Each ampoule contains 10 mg of active ingredient.

30

Patent Claims


1. Compounds of the formula I




in which

10 A^1, A^2 , independently of one another, denote N, O or S,

X^1, X^2, X^3 , independently of one another, denote a single bond, NR^3 , NR^3 , NR^3 , O, S, or one of the following groups:



20 Y denotes $C=O$, SO , SO_2 , $(CR^1_2)_n$,



25 Cy denotes H, a carbocyclic or heterocyclic saturated, unsaturated or aromatic radical, which may be unsubstituted or mono- or polysubstituted by alkyl, Hal, CN, OH; OR, OCF_3 , CF_3 , COOR or by a $(CR^1_2)_n$ -Y- X^1 -(CR^1_2) $_n$ -Q group,

30 Q denotes H, alkyl, cycloalkyl, aryl or heteroaryl

R, R^1, R^2, R^3 denote H, alkyl, Hal, alkoxy, OH, alkenyl, alkoxyalkyl, hydroxyalkyl, $(CH_2)_n-Q$, $(CH_2)_n-Cy$ or $(CH_2)_nNR_2$,

Hal denotes F, Br or Cl

5

n denotes 0, 1, 2, 3, 4, 5, 6, 7, or 8,

m denotes 1 or 2

and

10

p denotes 0, 1 or 2,

and pharmaceutically usable derivatives, solvates, tautomers, salts and stereoisomers thereof, including mixtures thereof in all ratios.

15

2. Compounds according to Claim 1 in which A^1, A^2 denote O and/or N.


3. Compounds according to Claim 1 or 2 in which R^1 and R^2 denotes H, alkyl, CF_3 , OCF_3 , $OCOH$, Hal or SCF_3 .

20

4. Compounds according to one or more of Claims 1-3 in which R^3 , denotes H, alkyl, hydroxyalkyl, alkoxyalkyl, $(CH_2)_n Q$ or $(CH_2)_n NR_2$, in which Q, R^1 and n have the meaning indicated in Claim 1.

25

5. Compounds according to one or more of Claims 1-4 in which X^1, X^2 , denotes NR^3 , O or the following group:




30

and R^3 , m and p have the meaning indicated in Claim 1.

6. Compounds according to one or more of Claims 1-5 in which X^3 denotes a single bond or $(CH^2)_n$ and n has the meaning indicated in Claim 1.

- 5 7. Compounds according to one or more of Claims 1-6 in which

Y


 $C=O$, SO , SO_2 , $(CR^1_2)_n$,

- 10 8. Compounds according to one or more of Claims 1-7 in which Cy denotes substituted or unsubstituted cyclopentyl cyclohexyl, aryl or heteroaryl.


- 15 9. Compounds according to one or more of Claims 1-8 in which Q denotes aryl or heteroaryl.

10. Compounds according to one or more of Claims 1-9 in which the $Y-X^2-(CR^2_2)_n-X$ group denotes a single bond.


- 20 11. Compounds of the sub-formulae IA to IC:



IA



IB



IC


5

in which R^1 , X^1 , X^2 , X^3 , Y , R^2 and Cy have the meaning indicated in Claim 1.

10

12. Process for the preparation of compounds of the formula I according to Claims 1-11 and pharmaceutically usable derivatives, salts, solvates, tautomers and stereoisomers thereof, characterised in that a compound of the formula II


15



20

in which A^1 , A^2 and X^1 have the meanings indicated in Claim 1,

with a compound of the formula III



25

in which

Y , X^2 , R^2 , X^3 and Cy have the meaning indicated in Claim 1,

and/or optionally


a base or acid of the formula I is converted into one of its salts.

30

13. Medicaments comprising at least one compound of the formula I according to Claim 1 to 11 and/or pharmaceutically usable derivatives,


salts, solvates, tautomers and stereoisomers thereof, including mixtures thereof in all ratios, and optionally excipients and/or adjuvants.

14. Mixture comprise one or more compounds of the formula I and amount of one or more compounds of the formula V, analogues thereof and/or metabolites thereof,



in which


- Y' and Z' each, independently of one another, denote O or N, R⁹ and R¹⁰ each, independently of one another, denote H, OH, halogen, OC1-10-alkyl, OCF₃, NO₂ or NH₂, s denotes an integer between 2 and 6, in each case inclusive, and R⁸ and R¹¹ are each, independently of one another, in the meta- or para-position and are selected from the group:



15. Use according to Claim 14, where the compound of the formula V used is pentamidine or salts thereof.

16. Use of compounds according to Claim 1 to 11 and pharmaceutically usable derivatives, salts, solvates, tautomers and stereoisomers thereof, including mixtures thereof in all ratios, or of the mixture according to Claim 14 for the preparation of a medicament for the treatment of diseases which can be influenced by the inhibition, regulation and/or modulation of the mitotic motor protein Eg5.
17. Use of compound according to Claim 1 to 11 or of the mixture according to Claim 14 for the preparation of a medicament for the treatment and prophylaxis of cancer diseases.
18. Use according to Claim 17, where the cancer diseases are accompanied by a tumour from the group of tumours of the squamous epithelium, bladder, stomach, kidneys, head and neck, oesophagus, cervix, thyroid, intestine, liver, brain, prostate, urogenital tract, lymphatic system, stomach, larynx and/or lung.
19. Use according to Claim 18, where the tumour originates from the group monocytic leukaemia, lung adenocarcinoma, small-cell lung carcinomas, pancreatic cancer, glioblastomas and breast carcinoma and colorectal carcinoma.
20. Use according to Claim 19, where the cancer disease to be treated is a tumour of the blood and immune system.
21. Use according to Claim 20, where the tumour originates from the group of acute myeloid leukaemia, chronic myeloid leukaemia, acute lymphatic leukaemia and/or chronic lymphatic leukaemia.
22. Use of compounds of the formula I according to Claim 1 to 11 and/or physiologically acceptable salts and solvates thereof for the preparation of a medicament for the treatment of tumours in combination with a


therapeutically effective amount of one or more compounds of the formula V, analogues thereof and/or metabolites thereof,



in which

10 Y' and Z' each, independently of one another, denote O or N, R⁹ and R¹⁰ each, independently of one another, denote H, OH, halogen, OC1-10-alkyl, OCF₃, NO₂ or NH₂, s denotes an integer between 2 and 6, in each case inclusive, and R⁸ and R¹¹ are each, independently of one another, in the meta- or para-position and are selected from the

15 group:



25 the compounds of the formula I and the compounds of the formula V, analogues thereof and/or metabolites thereof, are administered simultaneously or within 14 days of one another in amounts which are sufficient to inhibit the growth of a tumour or of other hyperproliferative cells.

30 23. Use according to Claim 22, where the compound of the formula V used is pentamidine or salts thereof.

24. Use of compounds of the formula I according to Claim 1 to 11 and/or physiologically acceptable salts and solvates thereof for the preparation of a medicament for the treatment of tumours where a therapeutically effective amount of a compound of the formula I is administered in combination with radiotherapy and a compound from the group 1) oestrogen receptor modulator, 2) androgen receptor modulator, 3) retinoid receptor modulator, 4) cytotoxic agent, 5) antiproliferative agent, 6) prenyl-protein transferase inhibitor, 7) HMG-CoA reductase inhibitor, 8) HIV protease inhibitor, 9) reverse transcriptase inhibitor and 10) further angiogenesis inhibitors.

15

20

25

30

Fetherstonhaugh
Ottawa, Canada
Patent Agents

