



Office de la Propriété

Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2482598 C 2012/09/25

(11)(21) **2 482 598**

(12) **BREVET CANADIEN**  
**CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2003/04/22  
(87) Date publication PCT/PCT Publication Date: 2003/10/30  
(45) Date de délivrance/Issue Date: 2012/09/25  
(85) Entrée phase nationale/National Entry: 2004/10/14  
(86) N° demande PCT/PCT Application No.: US 2003/012541  
(87) N° publication PCT/PCT Publication No.: 2003/089456  
(30) Priorité/Priority: 2002/04/22 (US60/374,530)

(51) Cl.Int./Int.Cl. *C12N 9/12*(2006.01),  
*A61K 38/00*(2006.01), *A61K 38/08*(2006.01),  
*A61K 38/10*(2006.01), *A61K 38/45*(2006.01),  
*C07K 7/06*(2006.01), *C07K 7/08*(2006.01)

(72) Inventeur/Inventor:  
MOCHLY-ROSEN, DARIA, US

(73) Propriétaire/Owner:  
THE BOARD OF TRUSTEES OF THE LEELAND  
STANFORD JUNIOR UNIVERSITY, US

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : INHIBITEURS PEPTIDIQUES DE LA PROTEINE KINASE C

(54) Title: PEPTIDE INHIBITORS OF PROTEIN KINASE C

(57) Abrégé/Abstract:

PKC V5 isozyme-specific peptides are described. The sequences and compositions comprising the sequences are useful for treating diseases states associated with the PKC isozyme from which they are respectively derived. Methods of treatment, pharmaceutical formulations and methods of identifying compounds that mimic the activity of the peptides are also described.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 October 2003 (30.10.2003)

PCT

(10) International Publication Number  
**WO 2003/089456 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 38/00**

(21) International Application Number:  
PCT/US2003/012541

(22) International Filing Date: 22 April 2003 (22.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/374,530 22 April 2002 (22.04.2002) US

(71) Applicant (for all designated States except US): **THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY [US/US]**; 900 Welch Road, Suite 350, Palo Alto, CA 94304 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **MOCHLY-ROSEN, Daria** [US/US]; 900 Welch Road, Suite 350, Palo Alto, CA 94304 (US).

(74) Agent: **LOWIN, David, A.**; P.O. Box 620535, Woodside, CA 94062-0535 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report: 12 February 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2003/089456 A3

(54) Title: PEPTIDE INHIBITORS OF PROTEIN KINASE C

(57) Abstract: PKC V5 isozyme-specific peptides are described. The sequences and compositions comprising the sequences are useful for treating diseases states associated with the PKC isozyme from which they are respectively derived. Methods of treatment, pharmaceutical formulations and methods of identifying compounds that mimic the activity of the peptides are also described.

## PEPTIDE INHIBITORS OF PROTEIN KINASE C

5

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of co-pending provisional application 60/374,530, filed April 22, 2002, incorporated herein by reference.

10

15

### FIELD OF THE INVENTION

The present invention relates to peptides effective for the isozyme-specific modulation of protein kinase C.

20

### BACKGROUND OF THE INVENTION

Protein kinase C ("PKC") is a key enzyme in signal transduction involved in a variety of cellular functions, including cell growth, regulation of gene expression, and ion channel activity. The PKC family of isozymes includes at least 11 different protein kinases that can be divided into at least three subfamilies based on their homology and sensitivity to activators. Each isozyme includes a number of homologous ("conserved" or "C") domains interspersed with isozyme-unique ("variable" or "V") domains. Members of the "classical" or "cPKC" subfamily,  $\alpha$ ,  $\beta_I$ ,  $\beta_{II}$  and  $\gamma$ PKC, contain four homologous domains (C1, C2, C3 and C4) and require calcium, phosphatidylserine, and diacylglycerol or phorbol esters for activation. Members of the "novel" or "nPKC" subfamily,  $\delta$ ,  $\epsilon$ ,  $\eta$  and  $\theta$ PKC, lack the C2 homologous domain and do not require calcium for activation. Finally, members of the "atypical" or "aPKC" subfamily,  $\zeta$  and  $\lambda$ PKC, lack both the C2 and one-half of the C1 homologous domains and are insensitive to diacylglycerol, phorbol esters and calcium.

Studies on the subcellular distribution of PKC isozymes demonstrate that activation of PKC results in its redistribution in the cells (also termed translocation), such that activated PKC isozymes associate with the plasma membrane, cytoskeletal elements, nuclei, and other subcellular compartments (Saito, N. *et al.*, 5 *Proc. Natl. Acad. Sci. USA* **86**:3409-3413 (1989); Papadopoulos, V. and Hall, P.F. *J. Cell Biol.* **108**:553-567 (1989); Mochly-Rosen, D., *et al.*, *Molec. Biol. Cell* (formerly *Cell Reg.*) **1**:693-706, (1990)). The unique cellular functions of different PKC isozymes are determined by their subcellular location. For example, activated  $\beta_1$ PKC is found inside the nucleus, whereas activated  $\beta_{II}$ PKC is found at 10 the perinucleus and cell periphery of cardiac myocytes (Disatnik, M.H., *et al.*, *Exp. Cell Res.* **210**:287-297 (1994)).  $\epsilon$ PKC, a member of the novel PKC family independent from calcium but requiring phospholipids for activation, is found in primary afferent neurons both in the dorsal root ganglia as well as in the superficial layers of the dorsal spinal cord.

15 The localization of different PKC isozymes to different areas of the cell in turn appears due to binding of the activated isozymes to specific anchoring molecules termed Receptors for Activated C-Kinase ("RACKs"). RACKs are thought to function by selectively anchoring activated PKC isozymes to their respective subcellular sites. RACKs bind only fully activated PKC and are not necessarily substrates of the enzyme. Nor is the binding to RACKs mediated via the catalytic 20 domain of the kinase (Mochly-Rosen, D., *et al.*, *Proc. Natl. Acad. Sci. USA* **88**:3997-4000 (1991)). Translocation of PKC reflects binding of the activated enzyme to RACKs anchored to the cell particulate fraction and the binding to RACKs is required for PKC to produce its cellular responses (Mochly-Rosen, D., *et al.*, *Science* **268**:247-251 (1995)). Inhibition of PKC binding to RACKs *in vivo* 25 inhibits PKC translocation and PKC-mediated function (Johnson, J.A., *et al.*, *J. Biol. Chem.*, **271**:24962-24966 (1996a); Ron, D., *et al.*, *Proc. Natl. Acad. Sci. USA* **92**:492-496 (1995); Smith, B.L. and Mochly-Rosen, D., *Biochem. Biophys. Res. Commun.*, **188**:1235-1240 (1992)).

30 In general, translocation of PKC is required for proper function of PKC isozymes. Peptides that mimic either the PKC-binding site on RACKs (Mochly-Rosen, D., *et al.*, *J. Biol. Chem.*, **226**:1466-1468 (1991a); Mochly-Rosen, D., *et al.*, 1995) or the RACK-binding site on PKC (Ron, *et al.*, 1995; Johnson, J.A., *et al.*,

1996a) are isozyme-specific translocation inhibitors of PKC that selectively inhibit the function of the enzyme *in vivo*.

Individual isozymes of PKC have been implicated in the mechanisms of various disease states, including the following: cancer (alpha and delta PKC); 5 cardiac hypertrophy and heart failure (beta I and beta II PKC) nociception (gamma and epsilon PKC); ischemia including myocardial infarction (delta PKC); immune response, particularly T-cell mediated (theta PKC); and fibroblast growth and memory (zeta PKC). Various PKC isozyme- and variable region-specific peptides have been previously described (see, for example, U.S. Patent No. 5,783,405). 10 The role of  $\epsilon$ PKC in pain perception has recently been reported (WO 00/01415; U.S. Patent No. 6,376,467) including therapeutic use of the  $\epsilon$ V1-2 peptide (a selective inhibitor of  $\epsilon$ PKC first described in the above-referenced '405 patent). The binding specificity for RACK1, a selective anchoring protein for  $\beta_{II}$ PKC, has 15 recently been reported to reside in the V5 region of  $\beta_{II}$ PKC (Stebbins, E. *et al.*, *J. Biol. Chem.* **271**:29644-29650 (2001)), including the testing of certain N-, middle, and C-terminus peptides alone, in combination and together with a mixture of three peptides from the  $\beta$ C2 domain.

Notwithstanding such reported advances, new, selective agents and methods for the treatment of disease, including alternatives to known PKC 20 isozyme- and variable region-specific peptides have and continue to be desired.

### SUMMARY OF THE INVENTION

The present invention provides PKC V5 isozyme-specific peptides, pharmaceutical compositions and methods for the treatment diseases modulated 25 by a PKC isozyme. The peptides are selective modulators of the corresponding PKC isozymes.

In one aspect the invention pertains to a PKC V5 isozyme-specific peptide.

In another aspect of the invention, the peptide is selected from 6 to 12 of the N-terminal 15 amino acids of the V5 domain.

30 In still another aspect the peptide does not include the N-terminal 2 amino acids of the V5 domain.

In yet another aspect, the peptide has 6 to 8 amino acids.

Still another aspect of the invention pertains to an above-described peptide

conjugated to a carrier peptide, preferably Cys-Cys bonded to a carrier peptide selected from poly-Arg, Tat, or the *Drosophila* Antennapedia homeodomain.

The invention further provides pharmaceutical formulations including a pharmaceutically acceptable excipient and an above-described peptide or peptide/carrier conjugate.

The above-referenced peptides, peptide conjugates and pharmaceutical formulations include all modifications, derivations, fragments, combinations, or hybrids thereof that retain the desired activity. Excluded from the peptides of the invention are: SEQ ID NO:5, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:31, SEQ ID NO:37, SEQ ID NO:43, SEQ ID NO:48, SEQ ID NO:57, SEQ ID NO:63 and SEQ ID NO:64.

Also provided in the present invention is a method of treatment for a disease state modulated by a PKC isozyme comprising administering a therapeutically effective amount of an above-described PKC V5 isozyme-specific peptide, peptide/carrier conjugate (or a modification, derivation, fragment, combination, or hybrid thereof that retains the desired activity) or a pharmaceutical formulation thereof to a mammal in need of such treatment. In one embodiment, excluded from the methods of treatment of the invention are those methods employing a peptide of SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:19, SEQ ID NO:20, or SEQ ID NO:21. In another such embodiment, excluded from the methods of treatment of the invention are those methods employing a peptide of SEQ ID NO:5, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:31, SEQ ID NO:37, SEQ ID NO:43, SEQ ID NO:48, SEQ ID NO:57, SEQ ID NO:63 or SEQ ID NO:64.

In another aspect, the invention includes the use of the peptides described herein to identify compounds that treat disease.

An additional aspect includes the use of a peptides or peptide/carrier conjugate in the preparation of a medicament for use in the treatment of disease.

These and other objects and features of the invention will be more fully appreciated when the following detailed description of the invention is read in conjunction with the accompanying drawings.

**BRIEF DESCRIPTION OF THE SEQUENCES**

SEQ ID NO:1 represents the V5 domain of the human  $\alpha$ PKC isozyme.  
SEQ ID NO:2 is a fragment derived from SEQ ID NO:1.  
5 SEQ ID NO:3 is a modified fragment derived from SEQ ID NO:1.  
SEQ ID NO:4 is a modified fragment derived from SEQ ID NO:1.  
SEQ ID NO:5 is a fragment derived from SEQ ID NO:1.  
SEQ ID NO:6 represents the V5 domain of the human  $\beta$ <sub>I</sub>PKC isozyme.  
SEQ ID NO:7 is a fragment derived from SEQ ID NO:6.  
10 SEQ ID NO:8 is a fragment derived from SEQ ID NO:6.  
SEQ ID NO:9 is a modified fragment derived from SEQ ID NO:6.  
SEQ ID NO:10 is a modified fragment derived from SEQ ID NO:6.  
SEQ ID NO:11 is a fragment derived from SEQ ID NO:6.  
SEQ ID NO:12 is a fragment derived from SEQ ID NO:6.  
15 SEQ ID NO:13 is a fragment derived from SEQ ID NO:6.  
SEQ ID NO:14 represents the V5 domain of the human  $\beta$ <sub>II</sub>PKC isozyme.  
SEQ ID NO:15 is a fragment derived from SEQ ID NO:14.  
SEQ ID NO:16 is a fragment derived from SEQ ID NO:14.  
SEQ ID NO:17 is a fragment derived from SEQ ID NO:14.  
20 SEQ ID NO:18 is a modified fragment derived from SEQ ID NO:14.  
SEQ ID NO:19 is a fragment derived from SEQ ID NO:14.  
SEQ ID NO:20 is a fragment derived from SEQ ID NO:14.  
SEQ ID NO:21 is a fragment derived from SEQ ID NO:14.  
SEQ ID NO:22 represents the V5 domain of the human  $\gamma$ PKC isozyme.  
25 SEQ ID NO:23 is a fragment derived from SEQ ID NO:22.  
SEQ ID NO:24 is a fragment derived from SEQ ID NO:22.  
SEQ ID NO:25 is a modified fragment derived from SEQ ID NO:22.  
SEQ ID NO:26 represents the V5 domain of the human  $\delta$ PKC isozyme.  
SEQ ID NO:27 is a fragment derived from SEQ ID NO:26.  
30 SEQ ID NO:28 is a fragment derived from SEQ ID NO:26.  
SEQ ID NO:29 is a modified fragment derived from SEQ ID NO:26.

SEQ ID NO:30 is a modified fragment derived from SEQ ID NO:26.

SEQ ID NO:31 is a fragment derived from SEQ ID NO:26.

SEQ ID NO:32 represents the V5 domain of the human  $\epsilon$ PKC isozyme.

SEQ ID NO:33 is a fragment derived from SEQ ID NO:32.

5 SEQ ID NO:34 is a fragment derived from SEQ ID NO:32.

SEQ ID NO:35 is a modified fragment derived from SEQ ID NO:32.

SEQ ID NO:36 is a modified fragment derived from SEQ ID NO:32.

SEQ ID NO:37 is a fragment derived from SEQ ID NO:32.

SEQ ID NO:38 represents the V5 domain of the human  $\eta$ PKC isozyme.

10 SEQ ID NO:39 is a fragment derived from SEQ ID NO:38.

SEQ ID NO:40 is a fragment derived from SEQ ID NO:38.

SEQ ID NO:41 is a modified fragment derived from SEQ ID NO:38.

SEQ ID NO:42 is a modified fragment derived from SEQ ID NO:38.

SEQ ID NO:43 is a fragment derived from SEQ ID NO:38.

15 SEQ ID NO:44 represents the V5 domain of the human  $\eta$ PKC isozyme.

SEQ ID NO:45 is a fragment derived from SEQ ID NO:44.

SEQ ID NO:46 is a fragment derived from SEQ ID NO:44.

SEQ ID NO:47 is a modified fragment derived from SEQ ID NO:44.

SEQ ID NO:48 is a fragment derived from SEQ ID NO:44.

20 SEQ ID NO:49 represents the V5 domain of the human  $\mu$ PKC isozyme.

SEQ ID NO:50 is a fragment derived from SEQ ID NO:49.

SEQ ID NO:51 is a fragment derived from SEQ ID NO:49.

SEQ ID NO:52 is a modified fragment derived from SEQ ID NO:49.

SEQ ID NO:53 represents the V5 domain of the human  $\theta$ PKC isozyme.

25 SEQ ID NO:54 is a fragment derived from SEQ ID NO:53.

SEQ ID NO:55 is a fragment derived from SEQ ID NO:53.

SEQ ID NO:56 is a modified fragment derived from SEQ ID NO:53.

SEQ ID NO:57 is a fragment derived from SEQ ID NO:53.

SEQ ID NO:58 represents the V5 domain of the human  $\xi$ PKC isozyme.

30 SEQ ID NO:59 is a fragment derived from SEQ ID NO:58.

SEQ ID NO:60 is a fragment derived from SEQ ID NO:58.

SEQ ID NO:61 is a modified fragment derived from SEQ ID NO:58.

SEQ ID NO:62 is a modified fragment derived from SEQ ID NO:58.

SEQ ID NO:63 is a fragment derived from SEQ ID NO:58.

SEQ ID NO:64 represents a fragment derived from the V5 domain of the  
5 human  $\mu$ PKC isozyme.

SEQ ID NO:65 is a Tat-derived carrier peptide (Tat 47-57): Tyr Gly Lys Lys  
Arg Arg Gln Arg Arg Arg.

SEQ ID NO:66 is the *Drosophila* Antennapedia homeodomain-derived  
carrier peptide: Cys Arg Gln Ile Lys Ile Trp Phe Gln Asn Arg Arg Met Lys Trp Lys  
10 Lys.

## DETAILED DESCRIPTION OF THE INVENTION

### DEFINITIONS

Unless otherwise indicated, all terms herein have the same meaning as they  
would to one skilled in the art of the present invention. Practitioners are particularly  
15 directed to *Current Protocols in Molecular Biology* (Ausubel, F. M. et al., John Wiley  
and Sons, Inc., Media Pa.) for definitions and terms of the art.

Abbreviations for amino acid residues are the standard 3-letter and/or 1-letter  
codes used in the art to refer to one of the 20 common L-amino acids.

20 A "conserved set" of amino acids refers to a contiguous sequence of amino  
acids that is identical or closely homologous (e.g., having only conservative amino  
acid substitutions) between members of a group of proteins. A conserved set may  
be anywhere from two to over 50 amino acid residues in length. Typically, a  
conserved set is between two and ten contiguous residues in length. For example,  
25 for the two peptides CGRNE(SEQ ID NO:15) and ACGRNAE(SEQ ID NO:19),  
there are 6 identical positions (CGRNAE) that form the conserved set of amino  
acids for these two sequences.

30 "Conservative amino acid substitutions" are substitutions that do not result  
in a significant change in the activity or tertiary structure of a selected polypeptide  
or protein. Such substitutions typically involve replacing a selected amino acid  
residue with a different residue having similar physico-chemical properties. For  
example, substitution of Glu for Asp is considered a conservative substitution since  
both are similarly-sized negatively-charged amino acids. Groupings of amino

acids by physico-chemical properties are known to those of skill in the art.

"Domain" and "region" are used interchangeably herein and refer to a contiguous sequence of amino acids within a PKC isozyme, typically characterized by being either conserved or variable.

5 "Peptide" and "polypeptide" are used interchangeably herein and refer to a compound made up of a chain of amino acid residues linked by peptide bonds. Unless otherwise indicated, the sequence for peptides is given in the order from the "N" (or amino) terminus to the "C" (or carboxyl) terminus.

10 Two amino acid sequences or two nucleotide sequences are considered "homologous" (as this term is preferably used in this specification) if they have an alignment score of >5 (in standard deviation units) using the program ALIGN with the mutation gap matrix and a gap penalty of 6 or greater (Dayhoff, M. O., in ATLAS OF PROTEIN SEQUENCE AND STRUCTURE (1972) Vol. 5, National Biomedical Research Foundation, pp. 101-110, and Supplement 2 to this volume, pp. 1-10.) The two 15 sequences (or parts thereof) are more preferably homologous if their amino acids are greater than or equal to 50%, more preferably 70%, still more preferably 80%, identical when optimally aligned using the ALIGN program mentioned above.

20 A peptide or peptide fragment is "derived from" a parent peptide or polypeptide if it has an amino acid sequence that is homologous to the amino acid sequence of, or is a conserved fragment from, the parent peptide or polypeptide.

"Modulate" intends a lessening, an increase, or some other measurable change in PKC activation.

"Management," for example in the context of treating pain, intends both a lessening of pain and/or induction of analgesia.

25 The term "treatment" or "treating" means any treatment of disease in a mammal, including: (a) preventing or protecting against the disease, that is, causing the clinical symptoms not to develop; (b) inhibiting the disease, that is, arresting or suppressing the development of clinical symptoms; and/or (c) relieving the disease, that is, causing the regression of clinical symptoms. It will be 30 understood by those skilled in the art that in human medicine, it is not always possible to distinguish between "preventing" and "suppressing" since the ultimate inductive event or events may be unknown, latent, or the patient is not ascertained until well after the occurrence of the event or events. Therefore, as used herein

the term "prophylaxis" is intended as an element of "treatment" to encompass both "preventing" and "suppressing" as defined herein. The term "protection," as used herein, is meant to include "prophylaxis."

5 The term "effective amount" means a dosage sufficient to provide treatment for the disorder or disease state being treated. This will vary depending on the patient, the disease and the treatment being effected.

10 The term "pharmaceutically acceptable carrier" or "pharmaceutically acceptable excipient" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

15

#### PKC PEPTIDES OF THE INVENTION

Prior reports have focused on PKC isozyme- and variable region-specific peptides from the V1, V3 and V5 domains, but the therapeutic and related potentials of peptides from the V5 domain remain largely unidentified. Moreover, 20 to the extent that peptides from the V5 domain have been described, such peptides have been predominantly selected from the approximate center of the V5 domain, starting about 25 amino acids from the N-terminus of the V5 domain.

Generally, the invention contemplates use of a peptide derived from a V5 region of any protein kinase C isozyme for use in treating and/or managing 25 disease. The present invention pertains to such PKC isozyme-specific V5 peptides, including methods of use and treatment, compositions of matter, and pharmaceutical formulations thereof. In particular, preferred are peptides selected from about 6 to 12 of the N-terminal 15 amino acids of the V5 domain, or a conservative modification or juxtaposition thereof. Further preferred are those peptides that do not include the N-terminal 2 amino acids of the V5 domain, especially peptides having 6 to 8 amino acids. The peptides can be used in native 30 form or modified by conjugation to a carrier, for example via a disulfide bond between a Cys on the carrier and a Cys within or added to the peptide, such as

those described below.

It will be appreciated that peptides homologous to the native sequences and peptides having conservative amino acid substitutions and/or juxtapositions, as well as fragments that retain activity, are within the scope of peptides contemplated. For example, one or more amino acids (preferably no more than two) can be substituted, changing between R and K; between V, L, I, R and D; and/or between G, A, P and N. Thus, the term "a PKC V5 peptide" contemplates the native sequence and all modifications, derivations, fragments, combinations, and hybrids thereof that retain the desired activity.

The following sequences correspond to the V5 domain of various PKC isozymes and to exemplary fragments derived therefrom. Exemplary modified peptides are also described below, where the substitution(s) are indicated in lower case. In all cases, it is appreciated that sequences derived from and homologous to those expressly indicated herein (e.g., closely homologous sequences from other species) are contemplated. All peptides described herein can be prepared by chemical synthesis using either automated or manual solid phase synthetic technologies, known in the art. The peptides can also be prepared recombinantly, using techniques known in the art.

The V5 domain of the  $\alpha$ PKC isozyme has the amino acid sequence: "PKVCG KGAENFDKFF TRGQPVLTPP DQLVIANIDQ SDFEGFSYVN PQFVHPILQS AV" (SEQ ID NO:1), taken from amino acid residue 616 et seq. of  $\alpha$ PKC. Exemplary peptides include GKGAEN (SEQ ID NO:2), corresponding to amino acid residues 620-625, and modified peptides "arGAEN" (SEQ ID NO:3) and cGKGAEN (SEQ ID NO:4). Excluded is the peptide QLVIAN (SEQ ID NO:5).

The V5 domain of the  $\beta$ PKC isozyme has the amino acid sequence: "PK ARDKRDTSNF DKEFTRQPVE LTPTDKLFIM NLDQNEFAGF SYTNPEFVIN V" (SEQ ID NO:6). Exemplary peptides include RDKRDTS (SEQ ID NO:7) and ARDKRDTSNF DK (SEQ ID NO:8), and modified peptides cARDKRDTS (SEQ ID NO:9) and gRDKRDTS (SEQ ID NO:10). Excluded are the peptides ARDKRDTS (SEQ ID NO:11), KLFIMN (SEQ ID NO:12) and AGFSYTNPEF VINV (SEQ ID NO:13).

The V5 domain of the  $\beta$ II-PKC isozyme has the amino acid sequence:  
"PK ACGRNAENFD RFFTRHPPVL TPPDQE VIRN IDQSEFEGFS FVNSEFLKPE  
VKS" (SEQ ID NO:14). Exemplary peptides include CGRNAE (SEQ ID NO:15),  
KACGRNAE (SEQ ID NO:16) and CGRNAEN (SEQ ID NO:17) and modified  
5 peptide ACGkNAE (SEQ ID NO:18). Excluded are the peptides ACGRNAE  
(SEQ ID NO:19) QE VIRN (SEQ ID NO:20) and SFVNSEFLKP EVKS (SEQ  
ID NO:21).

The V5 domain of the  $\gamma$ PKC isozyme has the amino acid sequence:  
"PRPCGRSG ENFDKFFTRA APALT TPPDRL VLASIDQADF QGFTYVNPDF  
10 VHPDARSPTS PVPVPVM" (SEQ ID NO:22) taken from amino acid residue 633 et  
seq. of  $\gamma$ PKC. Exemplary are the peptides GRSGEN (SEQ ID NO:23) and  
PCGRSGEN (SEQ ID NO:24), and modified peptide GkSGEN (SEQ ID NO:25).

The V5 domain of the  $\delta$ PKC isozyme has the amino acid sequence:  
"PKVKSPRDY SNFDQEFLNE KARLSYS DKN LIDSMDQSAF AGFSFVNPKF  
15 EHLLED" (SEQ ID NO:26). Exemplary peptides include VKSPRDYS (SEQ ID  
NO:27) taken from amino acid residues 624-631, PKVKSPRDY SN (SEQ ID  
NO:28), and modified peptides VKSPcRDYS (SEQ ID NO:29) and iKSPR $\gamma$ YS  
(SEQ ID NO:30). Excluded is the peptide KNLIDS (SEQ ID NO:31).

The V5 domain of the  $\epsilon$ PKC isozyme has the amino acid sequence:  
"PRIK TKRDVNNFDQ DFTREEPVLT LVDEAIVKQI NQEEFKGFSY FGEDLMP"  
20 (SEQ ID NO:32). Exemplary peptides include IKT KRDV (SEQ ID NO:33) taken  
from amino acid residues 689-695, and TKRDVNNFDQ (SEQ ID NO:34), and  
modified peptides cEAIVKQ (SEQ ID NO:35) and IKT K $\gamma$  (SEQ ID NO:36).  
Excluded is the peptide EAIVKQ (SEQ ID NO:37).

The V5 domain of the  $\eta$ PKC isozyme has the amino acid sequence:  
"PRIKSREDV SNFDPDFIKE EPVLTPIDEGL HLP MINQDEF RNFSYV SPEL QP"  
25 (SEQ ID NO:38). Exemplary peptides include IKSREDVS (SEQ ID NO:39)  
taken from amino acid residues 634-641, and PRIKSREDV (SEQ ID NO:40), and  
modified peptides vrSREDVS (SEQ ID NO:41) and EGHdPM (SEQ ID  
NO:42). Excluded is the peptide EGH LPM (SEQ ID NO:43).

The V5 domain of the  $\iota$ PKC isozyme has the amino acid sequence:  
"PNISGEFGL DNFDSQFTNE PVQLTPDDDD IVRKIDQSEF EGFEYINPLL

MSAEECV" (SEQ ID NO:44). Exemplary peptides include ISGEFGLD (SEQ ID NO:45) taken from amino acid resides 534-541 and DDDIVRK (SEQ ID NO:46), and modified peptide cSGEFGLD (SEQ ID NO:47). Excluded is the peptide DDIVRK (SEQ ID NO:48).

5 The V5 domain of the  $\mu$ PKC isozyme has the amino acid sequence:  
"PQVKLCDFGF ARIIGEKSFR RSVVGTPAYL APEVLRNKGY NRSLDMWSVG  
VIIYVSLSGT FPFNEDEDIH DQIQNAAFMY PPNPWKEISH EAIDLINNLL  
QVKMRKRYSV DKTLSHPWLQ DYQTWLDLRE LECKIGERYI THESDDLWWE  
KYAGEQRLQY PTHLINPSAS HSDTPETEET EMKALGERVS IL" (SEQ ID  
10 NO:49). Exemplary peptides include VKLCDFGF (SEQ ID NO:50) taken from amino acid resides 723-730, and QVKLCDFGFA (SEQ ID NO:51), and modified peptide irLCDFaF (SEQ ID NO:52).

The V5 domain of the  $\theta$ PKC isozyme has the amino acid sequence:  
"PKVKSPFD CSNFDKEFLN EKPRLSFADR ALINSMDQNM FRNFSFMNPG  
15 MERLIS" (SEQ ID NO:53). Exemplary peptides include VKSPFDCS (SEQ ID NO:54) taken from amino acid resides 655-662, and DRALINS (SEQ ID NO:55), and modified peptides VrSPFDCS (SEQ ID NO:56). Excluded is the peptide RALINS (SEQ ID NO:57).

The V5 domain of the  $\xi$ PKC isozyme has the amino acid sequence:  
20 "PQIT DDYGLDNFDT QFTSEPVQLT PDDEDAIKRI DQSEFEGFEY INPLLSTEE  
SV" (SEQ ID NO:58). Exemplary peptides include ITDDYGLD (SEQ ID NO:59) taken from amino acid resides 539-546, and DDYGLDN (SEQ ID NO:60), and modified peptides ITDDYGd $\beta$  (SEQ ID NO:61) and EDAIR (SEQ ID NO:62). Excluded is the peptide EDAIKR (SEQ ID NO:63).

25 Also excluded is the peptide SDSPEA (SEQ ID NO:64), identified as  $\mu$ PKC V5 peptide " $\mu$ V5-1" in U.S. Patent No. 5,783,405.

In all of the exemplary fragments recited above, conservative modifications and other modifications that do not appreciably alter the activity can be made and fall within the contemplated peptides.

30 Preferred are the peptides of the group: SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:23,

SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:50, 5 SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61 and SEQ ID NO:62, optionally including all modifications, derivations, fragments, combinations, and hybrids thereof that retain the desired activity.

More preferred are the peptides of the group: SEQ ID NO:2, SEQ ID NO:7, 10 SEQ ID NO:8, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:59 and SEQ ID NO:60, optionally including all modifications, 15 derivations, fragments, combinations, and hybrids thereof that retain the desired activity.

Still more preferred are the peptides of the group: SEQ ID NO:2, SEQ ID NO:7, SEQ ID NO:15, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:33, SEQ ID NO:39, SEQ ID NO:45, SEQ ID NO:50, SEQ ID NO:54 and SEQ ID 20 NO:59, optionally including all modifications, derivations, fragments, combinations, and hybrids thereof that retain the desired activity.

Similarly preferred are the peptides of the group: SEQ ID NO:2, SEQ ID NO:23, SEQ ID NO:27, SEQ ID NO:33, SEQ ID NO:39, SEQ ID NO:45, SEQ ID NO:50, SEQ ID NO:54 and SEQ ID NO:59, optionally including all 25 modifications, derivations, fragments, combinations, and hybrids thereof that retain the desired activity.

## UTILITY

### Therapeutic Indications

30 The PKC isozyme-specific V5 peptides of the present invention are useful as modulators of PKC, being selective for the isozyme from which they are derived. In that regard, the peptides can be used in the treatment of mammalian (especially

human) disease states associated with particular PKC isozymes, including:  $\alpha$ PKC (hyperproliferative cellular diseases, such as cancer);  $\beta_I$ PKC and  $\beta_{II}$ PKC (cardiac hypertrophy and heart failure);  $\gamma$ PKC (pain management);  $\delta$ PKC (protecting tissue from damage due to an ischemic or hypoxic event, such as myocardial infarction and stroke, or apoptosis induced by UV irradiation, and for inhibiting fibroblast growth to promote scarless wound healing);  $\epsilon$ PKC (pain management, myocardial dysfunction);  $\theta$ PKC (immune system modulation, particularly involving T-cell mediated responses); and  $\zeta$ PKC (memory and stimulating fibroblast growth).

By way of example, pain is a basic clinical symptom seen by physicians and is often categorized as mild, moderate, or severe. The  $\gamma$ PKC and  $\epsilon$ PKC peptides described herein are suitable for treatment of pain in any of these categories. For example, cancer and post-operative surgical pain are often described as being in the moderate-to-severe category. Tumor infiltration of bone, nerve, soft tissue, or viscera are common causes of cancer pain. Various factors influence the prevalence of cancer pain in patients, such as the tumor type, state, and site, as well as patient variables. With respect to post-operative pain, the severity of the pain is often dependent on location and extent of intervention.

More particularly, the  $\gamma$ PKC and  $\epsilon$ PKC peptides are suited to treatment of acute or chronic pain caused, for example, by neuropathic or inflammatory conditions. Exemplary inflammatory conditions contemplated for treatment include, but are not limited to, sunburn, osteoarthritis, colitis, carditis, dermatitis, myostis, neuritis, and rheumatoid arthritis, lupus and other collagen vascular diseases, as well as post-operative surgical pain. Conditions associated with neuropathic pain include, but are not limited to, trauma, surgery, amputation, abscess, demyelinating diseases, trigeminal neuralgia, cancer, chronic alcoholism, stroke, thalamic pain syndrome, diabetes, herpes infections, and the like.

Inflammation and nerve damage can induce hyperalgesia, where a noxious stimulus is perceived as intensely painful due to a lowering of pain threshold. Accordingly, in its embodiments addressed to the treatment of pain, the invention contemplates a composition and a method for treating hyperalgesia in a patient. Additionally, the invention contemplate compositions and methods for treating allodynia in a subject; that is, treating the pain associated with a normally non-noxious stimulus.

Use in Methods of Identification

Another aspect the usefulness of the invention is a method of identifying compounds that modulate pain, for example, by using the peptides described herein as research tools for identification of compounds that mimic the activity of the peptides. The invention also contemplates use of the peptides in assays to detect the site of action of the peptides or in studies on the mechanism of action of the peptides. In identifying compounds that mimic the activity of the peptides, compounds that are able to bind to cellular receptors to which the peptides bind or otherwise act in the same or a similar physiological manner as the peptides can be identified by several techniques. For example, one method comprises adding a test compound to a biological assay that determines the activity of a PKC peptide of the invention and detecting the activity of the test compound. Alternatively, test compounds that modulate the activity of a PKC isozyme can be determined with an assay and then tested for a corresponding therapeutic activity.

For example, a competitive binding screening assay can be used to identify compounds that mimic the activity of a PKC isozyme by adding a test compound and a detectably labeled peptide of the invention to mammalian cells, tissue, or the suitable RACK under conditions that allow binding of the peptide and comparing the results against binding of the labeled peptide (without test compound) to the cell, tissue or RACK. Compounds that mimic the activity of the peptide can compete with the peptide for binding to the cell, tissue or RACK. Consequently, a smaller amount of RACK-bound labeled peptide (or a larger amount of RACK-unbound labeled peptide) will be measured when the test compound mimics the activity of the peptide by binding to the receptor (as compared to the amounts of free and RACK-bound labeled peptide when a test compound does not mimic the activity of the peptide, does not bind to the receptor, or does so with less affinity).

In general, identification of compounds that mimic the activity of PKC isozymes are identified by measuring the ability of a test compound to inhibit, enhance, or modulate the activity of the corresponding PKC isozyme. The activity of the PKC isozyme in a selected assay is measured in the presence and absence of the test compound. The assay can be a competitive binding assay (e.g., as described above) or a cellular assay the monitors modulation of a second messenger

production, changes in cellular metabolism, or effects on enzymatic activity. Compounds identified as mimicking or modulating the activity of the PKC isozyme are then tested for therapeutic activity using a corresponding *in vivo* disease model.

5 **TESTING**

Activity of the peptides of the invention can be determined using any of the established *in vitro* and *in vivo* assays of efficacy in treatment of the above-mentioned indications. Ultimately, of course, safety and efficacy are determined in controlled human clinical studies.

10 Representative testing protocols are summarized below in the context of determining activity for the treatment of pain, and for the treatment of ischemia. Testing protocols for other PKC V5 peptide indications will be well known to those skilled in the art.

15 **V5 Domain Activity as Illustrated by PKC Peptides for Pain Management**

The effect of a PKC peptide to modulate nociception is investigated using models of acute inflammatory pain induced by capsaicin or by formalin. These models, and others, having long-term increases of sensitivity to noxious stimuli can be useful in modeling certain human pathological pain. The capsaicin model of 20 inflammation, together with a low rate thermal test, mimics central sensitization and hyperalgesia resulting from chronic pain. Application of capsaicin to the skin produces a robust, hours-long, C fiber selective hyperalgesia indicated by significant lowering of paw withdrawal latencies during low heating rate thermal tests. Capsaicin is the active ingredient in spicy "hot" foods. The receptor for 25 capsaicin (VR-1 vanilloid receptor found on C fibers) has been recently cloned. It is a ligand-gated, non-selective cation channel. In addition to responding to capsaicin, VR-1 also responds to thermal stimuli (approximately 43°C) (Kidd B.L., *et al.*, *Br. J. Anaesth.*, 87(1):3-11 (2001)) and to protons, suggesting that its activity 30 is enhanced during inflammation. Capsaicin has been shown to selectively activate and sensitize C fibers, and not A $\delta$ . Therefore, A $\delta$  latency measurements are used as controls for animal wellbeing during the studies.

The formalin model in rodents has been validated as a predictive test of treating injury-induced pain in humans (Dennis, S.G. and Melzack, R. in Advances

in Pain Research and Therapy, Vol. 3, 747, Eds. J.J. Bonica et al., Raven Press, New York, 1979; Tjolsen, A., et al., *Pain*, 51:5-17 (1992)). The model produces a bi-phasic response, where the initial phase is triggered by a primary afferent barrage, similar in character to that described for the acute phasic tests except that chemical nociceptors are the mediators. The second phase is considered to be the hyperalgesic spontaneous activity that results from the initial tissue damage and reflects the lowering of nociceptive threshold plus the priming or "wind up" of the corresponding spinal circuitry. Thus, both peripheral and central neuronal circuits and mediators are required to induce and sustain this painful tissue-injury condition.

When tested as described above,  $\gamma$ PKC V5 peptides and  $\varepsilon$ PKC V5 peptides confer a substantial reduction in pain.

#### V5 Domain Activity as Illustrated by PKC Peptides for Ischemia

Administration to isolated rat cardiac myocytes can be employed to determine the activity of  $\delta$ PKC peptides in protection from ischemia. A peptide or carrier-peptide conjugate is introduced into isolated adult rat cardiac myocytes ten minutes prior to prolonged ischemia. Cell damage is assessed using an osmotic fragility test by measuring uptake of trypan blue.

Administration to whole hearts *ex vivo* can be employed to determine if the peptides have activity when introduced extracellularly to a whole organ. Peptides optionally conjugated to a carrier peptide, a Tat-derived peptide are delivered into Langendorff perfused rat hearts prior to induction of an ischemic period. After perfusion with the peptides, global ischemia is effected for 30 minutes. After the 30 minute ischemic period, the amount of creatine phosphokinase (CPK) released is monitored during a 30 minute reperfusion period.

Another study determines if the peptides can be delivered to an intact organ to provide protection after an ischemic insult. In this study, the rat heart model described above is used and the hemodynamic parameters are measured during the 20 minutes of global ischemia and the 20 minutes of reperfusion. Test peptides are delivered at a concentration of 500 nM, only during the reperfusion period.

Protecting tissue from damage due to an ischemic or hypoxic event can be

evaluated by administering the peptide *in vivo* to adult female pigs, preferably during the last 10 minutes of a 30 minute ischemic insult. Five days later, the hearts are analyzed for tissue damage.

In another study, left ventriculogram is performed in pigs (n=5) at three time points: (1) before occlusion of left anterior descending artery by balloon catheter (pre ischemia); (2) immediately after reperfusion with 2.5  $\mu$ M/10 mL of  $\delta$ V1-1 (post ischemia); and (3) before sacrifice five days later (5 days post ischemia), using 6 Fr. of pig-tail catheter. LVG is recorded by 2 views, right anterior oblique and left anterior oblique. Ejection fraction (EF), the percent of blood ejected in a beat, during maximum contraction, of the total maximum present in the left ventricle, is analyzed by the software, Plus Plus (Sanders Data Systems), and the averages of two views are evaluated. Ejection fractions are calculated based on left ventricle dimensions. Ejection fraction is a measure of how well the heart is functioning, with a higher ejection fraction indicative of a better functioning heart. An ejection fraction of less than 50% in a short period of time can suggest progression into a state of heart failure.

Inhibition of damage to the brain as a result of stroke can be examined in a rat cerebral ischemia model. Ischemia is induced using an intraluminal suture to occlude the ostium of the middle cerebral artery. Test peptide, optionally conjugated to a Tat peptide or the Tat peptide alone are injected into the carotid artery before and after a two hour occlusion period. The brain from each animal is harvested 24 hours later, stained, and examined.

When tested as described above,  $\delta$ PKC V5 peptides, when delivered before, during, or after ischemia, confer a substantial reduction of damage to the heart and brain induced by ischemia.

#### ADMINISTRATION

The peptides are prepared for administration by combining with a pharmaceutically-acceptable carrier or diluent. Thus, a further aspect of the invention provides pharmaceutical compositions comprising a peptide of the invention in a dosage form for administration to a subject. Such a dosage form includes, but is not limited to, tablets, capsules, suspensions, syrups for oral administration, where suitable pharmaceutical carriers include starch, lactose, talc,

magnesium stearate, aqueous solutions, oil-water emulsions, and the like. Other dosage forms include intrathecal, intravenous, intramuscular, subcutaneous, where suitable pharmaceutical carriers include buffered-aqueous or non-aqueous media. The peptides can be locally administered (e.g., near a site of inflammation or 5 peripheral nerve damage) for example, by topical application, intradermal injection or drug delivery catheter.

The amount of the peptide in the composition can be varied so that a suitable dose is obtained and a therapeutic effect is achieved. The dosage will depend on a 10 number of factors such as the route of administration, the duration of treatment, the size and physical condition of the patient, the potency of the peptide and the patient's response. Effective amounts of the peptide can be determined by testing the peptide in one or more models known in the art, including those described herein.

The peptides can be administered as needed, hourly, several times per day, daily, or as often as the person experiencing the pain or that person's physician 15 deems appropriate. The peptides can be administered on an on-going basis for management of chronic indications, or can be administered on a short term basis prior to after an acute indications.

The peptides of the invention can be administered alone or linked to a carrier 20 peptide, such as a Tat carrier peptide (of which the peptide having the sequence identified as SEQ ID NO:65 is exemplary). Other suitable carrier peptides are known and contemplated, such as the *Drosophila* Antennapedia homeodomain (SEQ ID NO:66; Théodore, L., *et al.* *J. Neurosci.* **15**:7158 (1995); Johnson, J. A., *et al.*, *Circ. Res.* **79**:1086 (1996b)), where the PKC peptide is cross-linked via an 25 N-terminal Cys-Cys bond to the Antennapedia carrier. Polyarginine is another exemplary carrier peptide (Mitchell *et al.*, *J. Peptide Res.*, **56**:318-325 (2000); Rolbard *et al.*, *Nature Med.*, **6**:1253-1257 (2000)).

### EXAMPLES

The following examples further illustrate the invention described herein and are 30 in no way intended to limit the scope of the invention.

**Example 1****Effect of PKC Peptides on Capsaicin-Induced Nociception**

Adult male Sprague-Dawley rats weighing between 200-250 g are lightly anaesthetized with urethane (800 mg/kg, i.p.). The dorsal surface of each animal is 5 painted with India ink to ensure heat is applied evenly to the dorsal surface. Baseline measurements of all animals (n=10/test group) are taken for 45 minutes for both C-fibers (0.9 °C/sec heating rate) and A $\delta$ -fibers (6.5 °C/sec heating rate). The peptides administered are a positive control  $\epsilon$ PKC antagonist having the amino acid sequence EAIVKQ (SEQ ID NO:37) that had previously been identified active 10 in the assay, the  $\gamma$ PKC peptides identified as SEQ ID NO:23, SEQ ID NO:24 and SEQ ID NO:25 and the  $\epsilon$ PKC peptides identified as SEQ ID NO:33, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35 and SEQ ID NO:36. All peptides are linked 15 to a Tat carrier peptide, of which the sequenced identified as SEQ ID NO:65 is exemplary. A second series of the peptides is tested without a carrier. Control and test peptides are delivered intrathecally via direct lumbar puncture (10  $\mu$ M peptide in 20  $\mu$ L) 15 minutes prior to topical application of capsaicin on the left hind paw (100  $\mu$ L of 3% capsaicin). As negative controls, saline and Tat-carrier peptide 20 are also administered to two separate groups of test animals. A latency measurement is taken post peptide but prior to capsaicin application to control for direct peptide effect. Twenty minutes after the application of capsaicin, the ink is re-applied, and the dorsal surface of the hind paws is subjected to low rate heating for a maximum of 20 seconds. Foot withdrawal latencies are measured at 15 minute intervals. The  $\epsilon$ PKC and  $\gamma$ PKC test peptides result in a decreased pain score relative to the negative control pups.

25

**Example 2****Effect of PKC Peptides on Pre-Existing Capsaicin-Induced Nociception**

Testing of the  $\gamma$ PKC and  $\epsilon$ PKC peptides for treatment of pre-existing chronic pain is done as follows. The ability of the peptides to reverse established capsaicin-induced thermal hyperalgesia is determined using the procedure described in 30 Example 1, except the test peptides are administered at various concentrations of test peptides (1  $\mu$ M, 50  $\mu$ M, and 100  $\mu$ M) over a 10-minute period commencing 25 minutes post capsaicin treatment (i.e., capsaicin is administered after the baseline

measurements). Thermal testing is then done as described in Example 1. Thirty minutes after capsaicin application, paw withdrawal latency was measured at regular intervals for 75 minutes. The  $\epsilon$ PKC and  $\gamma$ PKC test peptides result in a decreased pain score relative to the negative control pups.

5

### Example 3

#### Inhibition of $\delta$ PKC Translocation

##### A. Peptide Preparation

$\delta$ V5 PKC peptides (SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29 and SEQ ID NO:30) are synthesized and purified. The peptides are modified with a carrier peptide by cross-linking via an N-terminal Cys-Cys bond to the *Drosophila* Antennapedia homeodomain (Théodore, L., et al.; Johnson, J. A. et al., 1996b) or a Tat-derived peptide.

##### B. Peptide Delivery Into Cells

Primary cardiac myocyte cell cultures (90-95% pure) are prepared from newborn rats (Gray, M.O. et al.; Disatnik M.-H. et al.). The peptides are introduced into cells at an applied concentration of 500 nM in the presence and absence of phorbol 12-myristate 13-acetate (PMA) at concentrations of 3 nm and 10 nm, respectively, for 10-20 minutes. In a third set of cells, the peptides are applied at a concentration of 500 nM in the presence and absence of 500 nM  $\psi$  $\delta$ RACK.

Translocation of  $\delta$ PKC isozyme is assessed by using  $\delta$ PKC isozyme-specific antibodies in Western blot analysis (Santa Cruz Biotechnology). Western blot analysis of cytosolic and particulate fractions of treated cells is carried out as described by Johnson et al., 1995. Subcellular localization of delta PKC isozymes is assessed by chemiluminescence of blots probed with anti- $\delta$ PKC, anti- $\alpha$ PKC and anti- $\epsilon$ PKC antibodies. Amounts of PKC isozymes in each fraction are quantitated using a scanner and translocation is expressed as the amount of isozymes in the particulate fraction over the amount of isozymes in non-treated cells. Changes in translocation of  $\delta$ PKC isozyme are also determined by immunofluorescence study of treated and fixed cells (Gray et al., 1997). Translocation is determined by counting over 100 cells/treatment in a blinded fashion.

When tested as described above, the  $\delta$ V5 PKC peptides are active inhibitors of delta PKC translocation.

**Example 4****Peptide Administration to Isolated Cardiac Myocytes**

The peptides are prepared as described in Example 3.

5 Adult male Wistar rat cardiomyocytes are prepared on a Langendorff apparatus (van der Heide, R.S. *et al.*) by collagenase treatment (Armstrong, S. *et al.*). The cells are treated with peptides at concentrations of 10 nM, 100 nM, 500 nM, and 1  $\mu$ M in the presence or absence of 1  $\mu$ M  $\psi\delta$ RACK.  $\beta$ PKC-selective activator was used as a control.

10 For stimulated ischemia, adult myocytes treated in microcentrifuge tubes with  $\delta$ V1-1 and/or  $\psi\delta$ RACK peptides conjugated to the carrier are washed twice with degassed glucose-free incubation buffer and pelleted. On top of a minimal amount of buffer, the cell pellets are overlaid with either a micro-balloon (Sig Manufacturing, Montezuma, IA) or with degassed buffer saturated with nitrogen, and sealed with an airtight top. Tubes are then incubated at 37°C for either 180 minutes or 90 minutes.

15 Cell damage is assessed by an osmotic fragility test by measuring the uptake of trypan blue added in a hypotonic (85 mosM) solution. Similar results are also obtained by using a live-dead kit (Molecular Probes) or measuring the release of lactose dehydrogenase to the medium using a kit (Sigma) as previously described (Chen, *et al.*, 1999; Gray *et al.*, 1997; Mackay *et al.*, 1999).

20 The  $\delta$ V5 PKC peptides inhibit ischemic damage when tested as described above.

**Example 5****Ex vivo Peptide Administration to Whole Hearts and Effect on Cell Damage**

25 Adult, male rats are anesthetized with i.p. avertin, and their hearts are rapidly removed and cannulated via the aorta for perfusion as described in the art (Colbert, M. C. *et al.*) using Langendorff set-up. Care is taken to have the hearts perfused within 90 seconds of removal. The hearts are perfused with oxygenated Krebs-Henseleit solution comprised of, in nmol/L, NaCl 120; KCl 5.8; NaHCO<sub>3</sub> 25; NaH<sub>2</sub>O<sub>4</sub> 1.2; MgSO<sub>4</sub> 1.2; CaCl<sub>2</sub> 1.0; and dextrose 10, pH 7.4 at 37°C.

30 After a 10-20 minute equilibration period, the hearts are perfused with  $\delta$ V5 PKC peptides, prepared as described in Example 1 but conjugated to a Tat-derived

peptide, for 20 minutes. Perfusion is maintained at a constant flow of 10 mL/min with Krebs-Hanseleit solution containing 0.5  $\mu$ M of the appropriate peptide. The Langendorff method employed uses retrograde flow from the ventricle to the aorta and into the coronary arteries, bypassing the pulmonary arteries.

5 To induce global ischemia, flow is interrupted for 30 minutes. After the ischemic event, the hearts are re-perfused for 30-60 minutes. During reperfusion, ischemia-induced cell damage is determined by measuring the activity of creatine phosphokinase (CPK) (absorbance at 520 nm) in the perfusate using a Sigma kit. As controls, some *ex vivo* hearts are left untreated, or maintained under normoxia 10 conditions, or treated with the Tat-carrier peptide alone, or treated with Tat-carrier peptide conjugated to a previously identified inactive peptide.

The  $\delta$ V5 PKC peptides inhibit ischemic damage when tested as described above.

15

### Example 6

#### *In vivo Administration of $\delta$ V1-1 After Ischemia*

Adult female pigs, 35-40kg in weight, are anesthetized and a catheter is introduced through the carotid artery into the heart. Using conventional intervention cardiology techniques, a wire is placed through a catheter and into the 20 left anterior descending artery. A balloon is run over this wire to a site of occlusion where it is then inflated to block blood flow for 30 minutes. During the last 10 minutes of the 30-minute occlusion, either a control comprised of a carrier peptide alone or a test peptide (optionally conjugated to a carrier Tat peptide as described above) is delivered by slow diffusion (1mL/min) directly downstream of the 25 occlusion. Approximately 20 $\mu$ g of test peptide (~400ng per kg body weight) is administered.

After 30 minutes of occlusion, the balloon is removed and pigs are left to recover from surgery. Five days later, the pigs are euthanized and hearts are harvested. After heart removal, the LAD is occluded. With the occlusion in place, 30 Evans Blue dye, which stains all areas not at risk of infarct in blue while leaving all areas with no access to blood flow red, is infused. Hearts are then cut into slices and stained with a tetrazolium red dye which stains all live areas red and infarcted dead tissue in white. Each heart has multiple tissue slices with distinctive areas

marking the area at risk for ischemia and the infarcted area. From this, the percent infarct per area at risk for each slice and for the entire heart is determined.

The  $\delta$ V5 PKC peptides inhibit ischemic damage when tested as described above.

5

### Example 7

#### In vivo Administration to Rats for Stroke Damage Protection

##### A. Cerebral Ischemia Model

Adult male Sprague-Dawley rats weighing between 280-320 g are used.  
10 Animals are maintained under isofluorane anesthesia during all surgical procedures. Physiological parameters are monitored and maintained in the normal range. Rectal temperature is also measured. At the completion of the experiment, the animals are euthanized with a barbiturate overdose and prepared for histological analysis.  
15

##### B. Focal model

Ischemia is induced using an occluding intraluminal suture. An uncoated 30 mm long segment of 3-0 nylon monofilament suture with the tip rounded by flame is inserted into the stump of the common carotid artery and advanced into the internal carotid artery approximately 19-20 mm from the bifurcation in order to  
20 occlude the ostium of the middle cerebral artery. Sham control animals undergo similar anesthesia and surgical manipulation, but do not experience ischemia. At the end of a 2 hour ischemic period, the suture is removed and the animal allowed to recover. Brains are harvested after 24 hrs of reperfusion.

##### C. Peptide delivery

25 Test peptide optionally conjugated to Tat peptide, or Tat carrier control peptide (50 $\mu$ L of 10 $\mu$ M solution in saline) are injected into the carotid artery either immediately before or before and after the 2 hours occlusion. The final blood concentration of test peptide is 1  $\mu$ M.

##### D. Histology

30 Animals are perfused with heparinized saline and brains removed and sectioned into 2 mm thick slices. To assess ischemic injury, brain sections are stained with cresyl violet or with triphenyl tetrazolium chloride, a live tissue stain to indicate the regions of infarct. Areas of infarction (white) are then measured using

an image analysis system previously described (Yenari, M.A. *et al.*, 1998; Maier, C. *et al.*, 1998).

The  $\delta$ V5 PKC peptides inhibit ischemic damage when tested as described above.

5

REMAINDER OF PAGE INTENTIONALLY LEFT BLANK

WO 03/089456

PCT/US03/12541

While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many 5 modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

10

## SEQUENCE LISTING

<110> The Board of Trustees of the Leland Stanford Junior University

<120> Peptide inhibitors of protein kinase C

<130> 08901452CA

<140>

<141> 2003-04-22

<150> US 60/374,530

<151> 2002-04-22

<160> 66

<170> PatentIn version 3.1

<210> 1

<211> 57

<212> PRT

<213> Artificial

<220>

<223> V5 domain of the human alphaPKC isozyme

<400> 1

Pro Lys Val Cys Gly Lys Gly Ala Glu Asn Phe Asp Lys Phe Phe Thr  
1 5 10 15

Arg Gly Gln Pro Val Leu Thr Pro Pro Asp Gln Leu Val Ile Ala Asn  
20 25 30

Ile Asp Gln Ser Asp Phe Glu Gly Phe Ser Tyr Val Asn Pro Gln Phe  
35 40 45

Val His Pro Ile Leu Gln Ser Ala Val  
50 55

<210> 2

<211> 6

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:1

<400> 2

Gly Lys Gly Ala Glu Asn  
1 5

<210> 3  
<211> 6  
<212> PRT  
<213> Artificial

<220>  
<223> modified fragment derived from SEQ ID NO:1

<400> 3

Ala Arg Gly Ala Glu Asn  
1 5

<210> 4  
<211> 7  
<212> PRT  
<213> Artificial

<220>  
<223> modified fragment derived from SEQ ID NO:1

<400> 4

Cys Gly Lys Gly Ala Glu Asn  
1 5

<210> 5  
<211> 6  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:1

<400> 5

Gln Leu Val Ile Ala Asn  
1 5

<210> 6  
<211> 53  
<212> PRT  
<213> Artificial

<220>  
<223> V5 domain of the human beta1PKC isozyme

<400> 6

Pro Lys Ala Arg Asp Lys Arg Asp Thr Ser Asn Phe Asp Lys Glu Phe  
1 5 10 15

Thr Arg Gln Pro Val Glu Leu Thr Pro Thr Asp Lys Leu Phe Ile Met  
20 25 30

Asn Leu Asp Gln Asn Glu Phe Ala Gly Phe Ser Tyr Thr Asn Pro Glu  
35 40 45

Phe Val Ile Asn Val  
50

<210> 7

<211> 7

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:6

<400> 7

Arg Asp Lys Arg Asp Thr Ser  
1 5

<210> 8

<211> 12

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:6

<400> 8

Ala Arg Asp Lys Arg Asp Thr Ser Asn Phe Asp Lys  
1 5 10

<210> 9

<211> 9

<212> PRT

<213> Artificial

<220>

<223> modified fragment derived from SEQ ID NO:6

<400> 9

Cys Ala Arg Asp Lys Arg Asp Thr Ser  
1 5

<210> 10  
<211> 8  
<212> PRT  
<213> Artificial

<220>  
<223> modified fragment derived from SEQ ID NO:6

<400> 10

Gly Arg Asp Lys Arg Asp Thr Ser  
1 5

<210> 11  
<211> 8  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:6

<400> 11

Ala Arg Asp Lys Arg Asp Thr Ser  
1 5

<210> 12  
<211> 6  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:6

<400> 12

Lys Leu Phe Ile Met Asn  
1 5

<210> 13  
<211> 14  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:6

<400> 13

Ala Gly Phe Ser Tyr Thr Asn Pro Glu Phe Val Ile Asn Val  
1 5 10

<210> 14  
<211> 55  
<212> PRT  
<213> Artificial

&lt;220&gt;

&lt;223&gt; V5 domain of the human beta2PKC isozyme

&lt;400&gt; 14

Pro Lys Ala Cys Gly Arg Asn Ala Glu Asn Phe Asp Arg Phe Phe Thr  
1 5 10 15

Arg His Pro Pro Val Leu Thr Pro Pro Asp Gln Glu Val Ile Arg Asn  
20 25 30

Ile Asp Gln Ser Glu Phe Glu Gly Phe Ser Phe Val Asn Ser Glu Phe  
35 40 45

Leu Lys Pro Glu Val Lys Ser  
50 55

&lt;210&gt; 15

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; fragment derived from SEQ ID NO:14

&lt;400&gt; 15

Cys Gly Arg Asn Ala Glu  
1 5

&lt;210&gt; 16

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; fragment derived from SEQ ID NO:14

&lt;400&gt; 16

Lys Ala Cys Gly Arg Asn Ala Glu  
1 5

&lt;210&gt; 17

&lt;211&gt; 7

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; fragment derived from SEQ ID NO:14

<400> 17

Cys Gly Arg Asn Ala Glu Asn  
1 5

<210> 18

<211> 7

<212> PRT

<213> Artificial

<220>

<223> modified fragment derived from SEQ ID NO:14

<400> 18

Ala Cys Gly Lys Asn Ala Glu  
1 5

<210> 19

<211> 7

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:14

<400> 19

Ala Cys Gly Arg Asn Ala Glu  
1 5

<210> 20

<211> 6

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:14

<400> 20

Gln Glu Val Ile Arg Asn  
1 5

<210> 21

<211> 14

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:14

<400> 21

Ser Phe Val Asn Ser Glu Phe Leu Lys Pro Glu Val Lys Ser  
 1 5 10

<210> 22  
 <211> 65  
 <212> PRT  
 <213> Artificial

<220>  
 <223> V5 domain of the human gammaPKC isozyme

<400> 22

Pro Arg Pro Cys Gly Arg Ser Gly Glu Asn Phe Asp Lys Phe Phe Thr  
 1 5 10 15

Arg Ala Ala Pro Ala Leu Thr Pro Pro Asp Arg Leu Val Leu Ala Ser  
 20 25 30

Ile Asp Gln Ala Asp Phe Gln Gly Phe Thr Tyr Val Asn Pro Asp Phe  
 35 40 45

Val His Pro Asp Ala Arg Ser Pro Thr Ser Pro Val Pro Val Pro Val  
 50 55 60

Met  
 65

<210> 23  
 <211> 6  
 <212> PRT  
 <213> Artificial

<220>  
 <223> fragment derived from SEQ ID NO:22

<400> 23

Gly Arg Ser Gly Glu Asn  
 1 5

<210> 24  
 <211> 8  
 <212> PRT  
 <213> Artificial

<220>  
 <223> fragment derived from SEQ ID NO:22

<400> 24

Pro Cys Gly Arg Ser Gly Glu Asn  
1 5

<210> 25  
<211> 6  
<212> PRT  
<213> Artificial  
  
<220>  
<223> modified fragment derived from SEQ ID NO:22  
  
<400> 25

Gly Lys Ser Gly Glu Asn  
1 5

<210> 26  
<211> 55  
<212> PRT  
<213> Artificial  
  
<220>  
<223> V5 domain of the human deltaPKC isozyme  
  
<400> 26

Pro Lys Val Lys Ser Pro Arg Asp Tyr Ser Asn Phe Asp Gln Glu Phe  
1 5 10 15

Leu Asn Glu Lys Ala Arg Leu Ser Tyr Ser Asp Lys Asn Leu Ile Asp  
20 25 30

Ser Met Asp Gln Ser Ala Phe Ala Gly Phe Ser Phe Val Asn Pro Lys  
35 40 45

Phe Glu His Leu Leu Glu Asp  
50 55

<210> 27  
<211> 8  
<212> PRT  
<213> Artificial  
  
<220>  
<223> fragment derived from SEQ ID NO:26  
  
<400> 27

Val Lys Ser Pro Arg Asp Tyr Ser  
1 5

<210> 28  
<211> 11  
<212> PRT  
<213> Artificial  
  
<220>  
<223> fragment derived from SEQ ID NO:26  
  
<400> 28

Pro Lys Val Lys Ser Pro Arg Asp Tyr Ser Asn  
1 5 10

<210> 29  
<211> 9  
<212> PRT  
<213> Artificial  
  
<220>  
<223> modified fragment derived from SEQ ID NO:26  
  
<400> 29

Val Lys Ser Pro Cys Arg Asp Tyr Ser  
1 5

<210> 30  
<211> 8  
<212> PRT  
<213> Artificial  
  
<220>  
<223> modified fragment derived from SEQ ID NO:26  
  
<400> 30

Ile Lys Ser Pro Arg Leu Tyr Ser  
1 5

<210> 31  
<211> 6  
<212> PRT  
<213> Artificial  
  
<220>  
<223> fragment derived from SEQ ID NO:26  
  
<400> 31

Lys Asn Leu Ile Asp Ser  
1 5

<210> 32  
<211> 51

<212> PRT  
<213> Artificial

<220>  
<223> V5 domain of the human epsilonPKC isozyme

<400> 32

Pro Arg Ile Lys Thr Lys Arg Asp Val Asn Asn Phe Asp Gln Asp Phe  
1 5 10 15

Thr Arg Glu Glu Pro Val Leu Thr Leu Val Asp Glu Ala Ile Val Lys  
20 25 30

Gln Ile Asn Gln Glu Glu Phe Lys Gly Phe Ser Tyr Phe Gly Glu Asp  
35 40 45

Leu Met Pro  
50

<210> 33  
<211> 7  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:32

<400> 33

Ile Lys Thr Lys Arg Asp Val  
1 5

<210> 34  
<211> 10  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:32

<400> 34

Thr Lys Arg Asp Val Asn Asn Phe Asp Gln  
1 5 10

<210> 35  
<211> 7  
<212> PRT  
<213> Artificial

<220>

<223> modified fragment derived from SEQ ID NO:32

<400> 35

Cys Glu Ala Ile Val Lys Gln  
1 5

<210> 36

<211> 7

<212> PRT

<213> Artificial

<220>

<223> modified fragment derived from SEQ ID NO:32

<400> 36

Ile Lys Thr Lys Arg Leu Ile  
1 5

<210> 37

<211> 6

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:32

<400> 37

Glu Ala Ile Val Lys Gln  
1 5

<210> 38

<211> 51

<212> PRT

<213> Artificial

<220>

<223> V5 domain of the human etaPKC isozyme

<400> 38

Pro Arg Ile Lys Ser Arg Glu Asp Val Ser Asn Phe Asp Pro Asp Phe  
1 5 10 15

Ile Lys Glu Glu Pro Val Leu Thr Pro Ile Asp Glu Gly His Leu Pro  
20 25 30

Met Ile Asn Gln Asp Glu Phe Arg Asn Phe Ser Tyr Val Ser Pro Glu  
35 40 45

Leu Gln Pro  
50

<210> 39  
<211> 8  
<212> PRT  
<213> Artificial  
  
<220>  
<223> fragment derived from SEQ ID NO:38  
  
<400> 39

Ile Lys Ser Arg Glu Asp Val Ser  
1 5

<210> 40  
<211> 9  
<212> PRT  
<213> Artificial  
  
<220>  
<223> fragment derived from SEQ ID NO:38  
  
<400> 40

Pro Arg Ile Lys Ser Arg Glu Asp Val  
1 5

<210> 41  
<211> 8  
<212> PRT  
<213> Artificial  
  
<220>  
<223> modified fragment derived from SEQ ID NO:38  
  
<400> 41

Val Arg Ser Arg Glu Asp Val Ser  
1 5

<210> 42  
<211> 6  
<212> PRT  
<213> Artificial  
  
<220>  
<223> modified fragment derived from SEQ ID NO:38  
  
<400> 42

Glu Gly His Asp Pro Met  
1 5

<210> 43  
<211> 6  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:38

<400> 43

Glu Gly His Leu Pro Met  
1 5

<210> 44  
<211> 56  
<212> PRT  
<213> Artificial

<220>  
<223> V5 domain of the human iotaPKC isozyme

<400> 44

Pro Asn Ile Ser Gly Glu Phe Gly Leu Asp Asn Phe Asp Ser Gln Phe  
1 5 10 15

Thr Asn Glu Pro Val Gln Leu Thr Pro Asp Asp Asp Asp Ile Val Arg  
20 25 30

Lys Ile Asp Gln Ser Glu Phe Glu Gly Phe Glu Tyr Ile Asn Pro Leu  
35 40 45

Leu Met Ser Ala Glu Glu Cys Val  
50 55

<210> 45  
<211> 8  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:44

<400> 45

Ile Ser Gly Glu Phe Gly Leu Asp  
1 5

<210> 46  
<211> 7

<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:44

<400> 46

Asp Asp Asp Ile Val Arg Lys  
1 5

<210> 47  
<211> 8  
<212> PRT  
<213> Artificial

<220>  
<223> modified fragment derived from SEQ ID NO:44

<400> 47

Cys Ser Gly Glu Phe Gly Leu Asp  
1 5

<210> 48  
<211> 6  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:44

<400> 48

Asp Asp Ile Val Arg Lys  
1 5

<210> 49  
<211> 192  
<212> PRT  
<213> Artificial

<220>  
<223> V5 domain of the human muPKC isozyme

<400> 49

Pro Gln Val Lys Leu Cys Asp Phe Gly Phe Ala Arg Ile Ile Gly Glu  
1 5 10 15

Lys Ser Phe Arg Arg Ser Val Val Gly Thr Pro Ala Tyr Leu Ala Pro  
20 25 30

Glu Val Leu Arg Asn Lys Gly Tyr Asn Arg Ser Leu Asp Met Trp Ser  
 35 40 45

Val Gly Val Ile Ile Tyr Val Ser Leu Ser Gly Thr Phe Pro Phe Asn  
 50 55 60

Glu Asp Glu Asp Ile His Asp Gln Ile Gln Asn Ala Ala Phe Met Tyr  
 65 70 75 80

Pro Pro Asn Pro Trp Lys Glu Ile Ser His Glu Ala Ile Asp Leu Ile  
 85 90 95

Asn Asn Leu Leu Gln Val Lys Met Arg Lys Arg Tyr Ser Val Asp Lys  
 100 105 110

Thr Leu Ser His Pro Trp Leu Gln Asp Tyr Gln Thr Trp Leu Asp Leu  
 115 120 125

Arg Glu Leu Glu Cys Lys Ile Gly Glu Arg Tyr Ile Thr His Glu Ser  
 130 135 140

Asp Asp Leu Arg Trp Glu Lys Tyr Ala Gly Glu Gln Arg Leu Gln Tyr  
 145 150 155 160

Pro Thr His Leu Ile Asn Pro Ser Ala Ser His Ser Asp Thr Pro Glu  
 165 170 175

Thr Glu Glu Thr Glu Met Lys Ala Leu Gly Glu Arg Val Ser Ile Leu  
 180 185 190

<210> 50  
 <211> 8  
 <212> PRT  
 <213> Artificial

<220>  
 <223> fragment derived from SEQ ID NO:49

<400> 50

Val Lys Leu Cys Asp Phe Gly Phe  
 1 5

<210> 51  
 <211> 10  
 <212> PRT  
 <213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:49

<400> 51

Gln Val Lys Leu Cys Asp Phe Gly Phe Ala  
1 5 10

<210> 52

<211> 8

<212> PRT

<213> Artificial

<220>

<223> modified fragment derived from SEQ ID NO:49

<400> 52

Ile Arg Leu Cys Asp Phe Ala Phe  
1 5

<210> 53

<211> 54

<212> PRT

<213> Artificial

<220>

<223> V5 domain of the human thetaPKC isozyme

<400> 53

Pro Lys Val Lys Ser Pro Phe Asp Cys Ser Asn Phe Asp Lys Glu Phe  
1 5 10 15

Leu Asn Glu Lys Pro Arg Leu Ser Phe Ala Asp Arg Ala Leu Ile Asn  
20 25 30

Ser Met Asp Gln Asn Met Phe Arg Asn Phe Ser Phe Met Asn Pro Gly  
35 40 45

Met Glu Arg Leu Ile Ser  
50

<210> 54

<211> 8

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:53

<400> 54

Val Lys Ser Pro Phe Asp Cys Ser  
1 5

<210> 55

<211> 7

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:53

<400> 55

Asp Arg Ala Leu Ile Asn Ser  
1 5

<210> 56

<211> 8

<212> PRT

<213> Artificial

<220>

<223> modified fragment derived from SEQ ID NO:53

<400> 56

Val Arg Ser Pro Phe Asp Cys Ser  
1 5

<210> 57

<211> 6

<212> PRT

<213> Artificial

<220>

<223> fragment derived from SEQ ID NO:53

<400> 57

Arg Ala Leu Ile Asn Ser  
1 5

<210> 58

<211> 56

<212> PRT

<213> Artificial

<220>

<223> V5 domain of the human xiPKC isozyme

<400> 58

Pro Gln Ile Thr Asp Asp Tyr Gly Leu Asp Asn Phe Asp Thr Gln Phe  
1 5 10 15

Thr Ser Glu Pro Val Gln Leu Thr Pro Asp Asp Glu Asp Ala Ile Lys  
20 25 30

Arg Ile Asp Gln Ser Glu Phe Glu Gly Phe Glu Tyr Ile Asn Pro Leu  
35 40 45

Leu Leu Ser Thr Glu Glu Ser Val  
50 55

<210> 59  
<211> 8  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:58

<400> 59

Ile Thr Asp Asp Tyr Gly Leu Asp  
1 5

<210> 60  
<211> 7  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:58

<400> 60

Asp Asp Tyr Gly Leu Asp Asn  
1 5

<210> 61  
<211> 8  
<212> PRT  
<213> Artificial

<220>  
<223> modified fragment derived from SEQ ID NO:58

<400> 61

Ile Thr Asp Asp Tyr Gly Asp Leu  
1 5

<210> 62  
<211> 5  
<212> PRT  
<213> Artificial

<220>  
<223> modified fragment derived from SEQ ID NO:58

<400> 62

Glu Asp Ala Ile Arg  
1 5

<210> 63  
<211> 6  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from SEQ ID NO:58

<400> 63

Glu Asp Ala Ile Lys Arg  
1 5

<210> 64  
<211> 6  
<212> PRT  
<213> Artificial

<220>  
<223> fragment derived from the V5 domain of the human muPKC isozyme

<400> 64

Ser Asp Ser Pro Glu Ala  
1 5

<210> 65  
<211> 10  
<212> PRT  
<213> Artificial

<220>  
<223> Tat-derived carrier peptide (Tat 47-57)

<400> 65

Tyr Gly Lys Lys Arg Arg Gln Arg Arg Arg  
1 5 10

<210> 66  
<211> 17

<212> PRT  
<213> Artificial

<220>

<223> Drosophila Antennapedia homeodomain-derived carrier peptide

<400> 66

Cys Arg Gln Ile Lys Ile Trp Phe Gln Asn Arg Arg Met Lys Trp Lys  
1 5 10 15

Lys

**Claims**

1. A peptide consisting of a sequence of six to twelve amino acid residues that is at least 70% identical to a contiguous sequence of six to twelve amino acid residues from the variable 5 (V5) domain of delta protein kinase C (PKC) identified as SEQ ID NO:26, provided that the peptide is not SEQ ID NO:31.
2. The peptide of claim 1 wherein said peptide is selected from the N-terminal 15 amino acids of said V5 domain.
3. The peptide of claim 1 or claim 2 wherein said peptide does not include the N-terminal 2 amino acids of said V5 domain.
4. The peptide of any one of claims 1-3 wherein said peptide has 6 to 8 amino acids.
5. The peptide of any one of claims 1-4, wherein said peptide is linked to a carrier peptide.
6. The peptide of claim 5 wherein said peptide is linked to a carrier peptide selected from poly-Arg, a Tat-derived peptide, or the Drosophila Antennapedia homeodomain.
7. The peptide of claim 1 selected from the group consisting of SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, and SEQ ID NO:30.
8. A pharmaceutical formulation comprising a pharmaceutically acceptable excipient and the peptide of any one of Claims 1-7.
9. A use of a therapeutically effective amount of a peptide consisting of a sequence of six to twelve amino acid residues that is at least 70% identical to a contiguous sequence of six to twelve amino acid residues from the variable 5 (V5) domain of delta

protein kinase C (PKC) identified as SEQ ID NO:26, provided that the peptide is not SEQ ID NO:31, for reducing tissue damage due to ischemia.

10. A use of a therapeutically effective amount of a peptide consisting of a contiguous sequence of six to twelve amino acid residues that is at least 70% identical to a contiguous sequence of six to twelve amino acid residues from the variable 5 (v5) domain of delta protein kinase C (PKC) identified as SEQ ID NO:26, provided that the peptide is not SEQ ID NO:31, for the production of a medicament for reducing tissue damage due to ischemia.
11. The use of claim 9 or claim 10 wherein said peptide is selected from 6 to 12 of the N-terminal 15 amino acids of said V5 domain.
12. The use of claim 10 or claim 11 wherein said peptide does not include the N-terminal 2 amino acids of said V5 domain.
13. The use of claim 10 or claim 11 wherein said peptide has 6 to 8 amino acids.
14. The use of any one of claims 10-13, wherein said peptide is linked to a carrier peptide.
15. The use of claim 14 wherein said peptide is linked to a carrier peptide selected from poly-Arg, a Tat-derived peptide, or the Drosophila Antennapedia homeodomain.
16. The use of claim 10 or claim 11 wherein said peptide is selected from the group consisting of SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, and SEQ ID NO:30.
17. The use of any one of claims 10-16, wherein the tissue damage is due to cardiac ischemia.

18. The peptide of claim 5, wherein said peptide is linked to the carrier peptide via a disulfide bond.
19. The peptide of claim 18, wherein the carrier peptide is the Tat-derived peptide consisting of SEQ ID NO:65.
20. The peptide of claim 7, wherein said peptide is linked to a carrier peptide.
21. The peptide of claim 20, wherein said peptide is linked to the carrier peptide via a disulfide bond.
22. The peptide of claim 20, wherein the carrier peptide is the Tat-derived peptide consisting of SEQ ID NO:65.