



(22) Date de dépôt/Filing Date: 2006/10/18  
 (41) Mise à la disp. pub./Open to Public Insp.: 2007/04/26  
 (62) Demande originale/Original Application: 2 626 262  
 (30) Priorité/Priority: 2005/10/18 (US60/727,512)

(51) Cl.Int./Int.Cl. *C12N 9/22* (2006.01),  
*A61K 38/46* (2006.01), *A61K 48/00* (2006.01),  
*A61P 31/00* (2006.01), *C12N 15/09* (2006.01),  
*C12N 15/55* (2006.01), *C12N 15/79* (2006.01),  
*C12N 15/90* (2006.01), *C12Q 1/68* (2006.01),  
*G06F 19/16* (2011.01)

(71) Demandeur/Applicant:  
DUKE UNIVERSITY, US

(72) Inventeurs/Inventors:  
HELLINGA, HOMME W., US;  
SMITH, JAMES JEFFERSON, US;  
JANTZ, DEREK, US

(74) Agent: MBM INTELLECTUAL PROPERTY LAW LLP

(54) Titre : MEGANUCLEASES CONCUES RATIONNELLEMENT POSSEDANT UNE SPECIFICITE SEQUENCE MODIFIEE ET UNE AFFINITE DE LIAISON POUR L'ADN

(54) Title: RATIONALLY-DESIGNED MEGANUCLEASES WITH ALTERED SEQUENCE SPECIFICITY AND DNA-BINDING AFFINITY

(57) **Abrégé/Abstract:**

Rationally-designed LAGLIDADG meganucleases and methods of making such meganucleases are provided. In addition, methods are provided for using the meganucleases to generate recombinant cells and organisms having a desired DNA sequence inserted into a limited number of loci within the genome, as well as methods of gene therapy, for treatment of pathogenic infections, and for in vitro applications in diagnostics and research.



## **ABSTRACT**

Rationally-designed LAGLIDADG meganucleases and methods of making such meganucleases are provided. In addition, methods are provided for using the meganucleases to generate recombinant cells and organisms having a desired DNA sequence inserted into a limited number of loci within the genome, as well as methods of gene therapy, for treatment of pathogenic infections, and for in vitro applications in diagnostics and research.

## **DEMANDES OU BREVETS VOLUMINEUX**

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS  
COMPREND PLUS D'UN TOME.**

**CECI EST LE TOME \_\_\_\_ DE \_\_\_\_**

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

## **JUMBO APPLICATIONS / PATENTS**

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE  
THAN ONE VOLUME.**

**THIS IS VOLUME 1 OF 2**

NOTE: For additional volumes please contact the Canadian Patent Office.

**RATIONALLY-DESIGNED MEGANUCLEASES  
WITH ALTERED SEQUENCE SPECIFICITY AND DNA-BINDING AFFINITY**

[0001]

GOVERNMENT SUPPORT

[0002] The invention was supported in part by grants 2R01-GM-0498712, 5F32-GM072322 and 5 DP1 OD000122 from the National Institute of General Medical Sciences of National Institutes of Health of the United States of America. Therefore, the U.S. government may have certain rights in the invention.

FIELD OF THE INVENTION

[0003] The invention relates to the field of molecular biology and recombinant nucleic acid technology. In particular, the invention relates to rationally-designed, non-naturally-occurring meganucleases with altered DNA recognition sequence specificity and/or altered affinity. The invention also relates to methods of producing such meganucleases, and methods of producing recombinant nucleic acids and organisms using such meganucleases.

BACKGROUND OF THE INVENTION

[0004] Genome engineering requires the ability to insert, delete, substitute and otherwise manipulate specific genetic sequences within a genome, and has numerous therapeutic and biotechnological applications. The development of effective means for genome modification remains a major goal in gene therapy, agrotechnology, and synthetic biology (Porteus *et al.* (2005), *Nat. Biotechnol.* 23: 967-73; Tzfira *et al.* (2005), *Trends Biotechnol.* 23: 567-9; McDaniel *et al.* (2005), *Curr. Opin. Biotechnol.* 16: 476-83). A common method for inserting or modifying a DNA sequence involves introducing a transgenic DNA sequence flanked by sequences homologous to the genomic target and

selecting or screening for a successful homologous recombination event. Recombination with the transgenic DNA occurs rarely but can be stimulated by a double-stranded break in the genomic DNA at the target site. Numerous methods have been employed to create DNA double-stranded breaks, including irradiation and chemical treatments. Although these methods efficiently stimulate recombination, the double-stranded breaks are randomly dispersed in the genome, which can be highly mutagenic and toxic. At present, the inability to target gene modifications to unique sites within a chromosomal background is a major impediment to successful genome engineering.

[0005] One approach to achieving this goal is stimulating homologous recombination at a double-stranded break in a target locus using a nuclease with specificity for a sequence that is sufficiently large to be present at only a single site within the genome (see, e.g., Porteus *et al.* (2005), *Nat. Biotechnol.* 23: 967-73). The effectiveness of this strategy has been demonstrated in a variety of organisms using chimeric fusions between an engineered zinc finger DNA-binding domain and the non-specific nuclease domain of the FokI restriction enzyme (Porteus (2006), *Mol Ther* 13: 438-46; Wright *et al.* (2005), *Plant J.* 44: 693-705; Urnov *et al.* (2005), *Nature* 435: 646-51). Although these artificial zinc finger nucleases stimulate site-specific recombination, they retain residual non-specific cleavage activity resulting from under-regulation of the nuclease domain and frequently cleave at unintended sites (Smith *et al.* (2000), *Nucleic Acids Res.* 28: 3361-9). Such unintended cleavage can cause mutations and toxicity in the treated organism (Porteus *et al.* (2005), *Nat. Biotechnol.* 23: 967-73).

[0006] A group of naturally-occurring nucleases which recognize 15-40 base-pair cleavage sites commonly found in the genomes of plants and fungi may provide a less toxic genome engineering alternative. Such "meganucleases" or "homing endonucleases" are frequently associated with parasitic DNA elements, such as group 1 self-splicing introns and inteins. They naturally promote homologous recombination or gene insertion at specific locations in the host genome by producing a double-stranded break in the chromosome, which recruits the cellular DNA-repair machinery (Stoddard (2006), *Q. Rev. Biophys.* 38: 49-95). Meganucleases are commonly grouped into four families: the LAGLIDADG family, the GIY-YIG family, the His-Cys box family and the HNH family. These families are characterized by structural motifs, which affect catalytic activity and

recognition sequence. For instance, members of the LAGLIDADG family are characterized by having either one or two copies of the conserved LAGLIDADG motif (see Chevalier *et al.* (2001), *Nucleic Acids Res.* 29(18): 3757-3774). The LAGLIDADG meganucleases with a single copy of the LAGLIDADG motif form homodimers, whereas members with two copies of the LAGLIDADG motif are found as monomers. Similarly, the GIY-YIG family members have a GIY-YIG module, which is 70–100 residues long and includes four or five conserved sequence motifs with four invariant residues, two of which are required for activity (see Van Roey *et al.* (2002), *Nature Struct. Biol.* 9: 806-811). The His-Cys box meganucleases are characterized by a highly conserved series of histidines and cysteines over a region encompassing several hundred amino acid residues (see Chevalier *et al.* (2001), *Nucleic Acids Res.* 29(18): 3757-3774). In the case of the NHN family, the members are defined by motifs containing two pairs of conserved histidines surrounded by asparagine residues (see Chevalier *et al.* (2001), *Nucleic Acids Res.* 29(18): 3757-3774). The four families of meganucleases are widely separated from one another with respect to conserved structural elements and, consequently, DNA recognition sequence specificity and catalytic activity.

[0007] Natural meganucleases, primarily from the LAGLIDADG family, have been used to effectively promote site-specific genome modification in plants, yeast, *Drosophila*, mammalian cells and mice, but this approach has been limited to the modification of either homologous genes that conserve the meganuclease recognition sequence (Monnat *et al.* (1999), *Biochem. Biophys. Res. Commun.* 255: 88-93) or to pre-engineered genomes into which a recognition sequence has been introduced (Rouet *et al.* (1994), *Mol. Cell. Biol.* 14: 8096-106; Chilton *et al.* (2003), *Plant Physiol.* 133: 956-65; Puchta *et al.* (1996), *Proc. Natl. Acad. Sci. USA* 93: 5055-60; Rong *et al.* (2002), *Genes Dev.* 16: 1568-81; Gouble *et al.* (2006), *J. Gene Med.* 8(5):616-622).

[0008] Systematic implementation of nuclease-stimulated gene modification requires the use of engineered enzymes with customized specificities to target DNA breaks to existing sites in a genome and, therefore, there has been great interest in adapting meganucleases to promote gene modifications at medically or biotechnologically relevant sites (Porteus *et al.* (2005), *Nat. Biotechnol.* 23: 967-73;

Sussman *et al.* (2004), *J. Mol. Biol.* 342: 31-41; Epinat *et al.* (2003), *Nucleic Acids Res.* 31: 2952-62).

[0009] The meganuclease I-CreI from *Chlamydomonas reinhardtii* is a member of the LAGLIDADG family which recognizes and cuts a 22 base-pair recognition sequence in the chloroplast chromosome, and which presents an attractive target for meganuclease redesign. The wild-type enzyme is a homodimer in which each monomer makes direct contacts with 9 base pairs in the full-length recognition sequence. Genetic selection techniques have been used to identify mutations in I-CreI that alter base preference at a single position in this recognition sequence (Sussman *et al.* (2004), *J. Mol. Biol.* 342: 31-41; Chames *et al.* (2005), *Nucleic Acids Res.* 33: e178; Seligman *et al.* (2002), *Nucleic Acids Res.* 30: 3870-9) or, more recently, at three positions in the recognition sequence (Arnould *et al.* (2006), *J. Mol. Biol.* 355: 443-58). The I-CreI protein-DNA interface contains nine amino acids that contact the DNA bases directly and at least an additional five positions that can form potential contacts in modified interfaces. The size of this interface imposes a combinatorial complexity that is unlikely to be sampled adequately in sequence libraries constructed to select for enzymes with drastically altered cleavage sites.

[0010] There remains a need for nucleases that will facilitate precise modification of a genome. In addition, there remains a need for techniques for generating nucleases with pre-determined, rationally-designed recognition sequences that will allow manipulation of genetic sequences at specific genetic loci and for techniques utilizing such nucleases to genetically engineer organisms with precise sequence modifications.

#### SUMMARY OF THE INVENTION

[0011] The present invention is based, in part, upon the identification and characterization of specific amino acid residues in the LAGLIDADG family of meganucleases that make contacts with DNA bases and the DNA backbone when the meganucleases associate with a double-stranded DNA recognition sequence, and thereby affect the specificity and activity of the enzymes. This discovery has been used, as described in detail below, to identify amino acid substitutions which can alter the recognition sequence specificity and/or DNA-binding affinity of the meganucleases, and

to rationally design and develop meganucleases that can recognize a desired DNA sequence that naturally-occurring meganucleases do not recognize. The invention also provides methods that use such meganucleases to produce recombinant nucleic acids and organisms by utilizing the meganucleases to cause recombination of a desired genetic sequence at a limited number of loci within the genome of the organism, for gene therapy, for treatment of pathogenic infections, and for *in vitro* applications in diagnostics and research.

[0012] Thus, in some embodiments, the invention provides recombinant meganucleases having altered specificity for at least one recognition sequence half-site relative to a wild-type I-CreI meganuclease, in which the meganuclease includes a polypeptide having at least 85% sequence similarity to residues 2-153 of the wild-type I-CreI meganuclease of SEQ ID NO: 1, but in which the recombinant meganuclease has specificity for a recognition sequence half-site which differs by at least one base pair from a half-site within an I-CreI meganuclease recognition sequence selected from SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and SEQ ID NO: 5, and in which the recombinant meganuclease includes at least one modification listed in Table 1 which is not an excluded modification found in the prior art.

[0013] In other embodiments, the invention provides recombinant meganucleases having altered specificity for at least one recognition sequence half-site relative to a wild-type I-MsoI meganuclease, in which the meganuclease includes a polypeptide having at least 85% sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6, but in which the recombinant meganuclease has specificity for a recognition sequence half-site which differs by at least one base pair from a half-site within an I-MsoI meganuclease recognition sequence selected from SEQ ID NO: 7 and SEQ ID NO: 8, and in which the recombinant meganuclease includes at least one modification listed in Table 2 which is not an excluded modification found in the prior art.

[0014] In other embodiments, the invention provides recombinant meganucleases having altered specificity for a recognition sequence relative to a wild-type I-SceI meganuclease, in which the meganuclease includes a polypeptide having at least 85% sequence similarity to residues 3-186 of the I-SceI meganuclease of SEQ ID NO: 9, but in which the recombinant meganuclease has specificity for a recognition sequence which



differs by at least one base pair from an I-SceI meganuclease recognition sequence of SEQ ID NO: 10 and SEQ ID NO: 11, and in which the recombinant meganuclease includes at least one modification listed in Table 3 which is not an excluded modification found in the prior art.

[0015] In other embodiments, the invention provides recombinant meganucleases having altered specificity for at least one recognition sequence half-site relative to a wild-type I-CeuI meganuclease, in which the meganuclease includes a polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12, but in which the recombinant meganuclease has specificity for a recognition sequence half-site which differs by at least one base pair from a half-site within an I-CeuI meganuclease recognition sequence selected from SEQ ID NO: 13 and SEQ ID NO: 14, and in which the recombinant meganuclease includes at least one modification listed in Table 4 which is not an excluded modification found in the prior art.

[0016] The meganucleases of the invention can include one, two, three or more of the modifications which have been disclosed herein in order to affect the sequence specificity of the recombinant meganucleases at one, two, three or more positions within the recognition sequence. The meganucleases can include only the novel modifications disclosed herein, or can include the novel modifications disclosed herein in combination with modifications found in the prior art. Specifically excluded, however, are recombinant meganucleases comprising only the modifications of the prior art.

[0017] In another aspect, the invention provides for recombinant meganucleases with altered binding affinity for double-stranded DNA which is not sequence-specific. This is accomplished by modifications of the meganuclease residues which make contacts with the backbone of the double-stranded DNA recognition sequence. The modifications can increase or decrease the binding affinity and, consequently, can increase or decrease the overall activity of the enzyme. Moreover, increases/decreases in binding and activity have been found to cause decreases/increases in sequence specificity. Thus, the invention provides a means for altering sequence specificity generally by altering DNA-binding affinity.

[0018] Thus, in some embodiments, the invention provides for recombinant meganucleases having altered binding affinity for double-stranded DNA relative to a

wild-type I-CreI meganuclease, in which the meganuclease includes a polypeptide having at least 85% sequence similarity to residues 2-153 of the I-CreI meganuclease of SEQ ID NO: 1, and in which the DNA-binding affinity has been either (1) increased by at least one modification corresponding to a substitution selected from (a) substitution of E80, D137, I81, L112, P29, V64 or Y66 with H, N, Q, S, T, K or R, or (b) substitution of T46, T140 or T143 with K or R; or, conversely, (2) decreased by at least one modification corresponding to a substitution selected from (a) substitution of K34, K48, R51, K82, K116 or K139 with H, N, Q, S, T, D or E, or (b) substitution of I81, L112, P29, V64, Y66, T46, T140 or T143 with D or E.

[0019] In other embodiments, the invention provides for recombinant meganucleases having altered binding affinity for double-stranded DNA relative to a wild-type I-MsoI meganuclease, in which the meganuclease includes a polypeptide having at least 85% sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6, and in which the DNA-binding affinity has been either (1) increased by at least one modification corresponding to a substitution selected from (a) substitution of E147, I85, G86 or Y118 with H, N, Q, S, T, K or R, or (b) substitution of Q41, N70, S87, T88, H89, Q122, Q139, S150 or N152 with K or R; or, conversely, (2) decreased by at least one modification corresponding to a substitution selected from (a) substitution of K36, R51, K123, K143 or R144 with H, N, Q, S, T, D or E, or (b) substitution of I85, G86, Y118, Q41, N70, S87, T88, H89, Q122, Q139, S150 or N152 with D or E.

[0020] In other embodiments, the invention provides for recombinant meganucleases having altered binding affinity for double-stranded DNA relative to a wild-type I-SceI meganuclease, in which the meganuclease includes a polypeptide having at least 85% sequence similarity to residues 3-186 of the I-SceI meganuclease of SEQ ID NO: 9, and in which the DNA-binding affinity has been either (1) increased by at least one modification corresponding to a substitution selected from (a) substitution of D201, L19, L80, L92, Y151, Y188, I191, Y199 or Y222 with H, N, Q, S, T, K or R, or (b) substitution of N15, N17, S81, H84, N94, N120, T156, N157, S159, N163, Q165, S166, N194 or S202 with K or R; or, conversely, (2) decreased by at least one modification corresponding to a substitution selected from (a) substitution of K20, K23, K63, K122, K148, K153, K190, K193, K195 or K223 with H, N, Q, S, T, D or E, or (b) substitution

of L19, L80, L92, Y151, Y188, I191, Y199, Y222, N15, N17, S81, H84, N94, N120, T156, N157, S159, N163, Q165, S166, N194 or S202 with D or E.

**[0021]** In other embodiments, the invention provides for recombinant meganucleases having altered binding affinity for double-stranded DNA relative to a wild-type I-CeuI meganuclease, in which the meganuclease includes a polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12, and in which the DNA-binding affinity has been either (1) increased by at least one modification corresponding to a substitution selected from (a) substitution of D25 or D128 with H, N, Q, S, T, K or R, or (b) substitution of S68, N70, H94, S117, N120, N129 or H172 with K or R; or, conversely, (2) decreased by at least one modification corresponding to a substitution selected from (a) substitution of K21, K28, K31, R112, R114 or R130 with H, N, Q, S, T, D or E, or (b) substitution of S68, N70, H94, S117, N120, N129 or H172 with D or E.

**[0022]** The meganucleases of the invention can include one, two, three or more of the modifications of backbone contact residues which have been disclosed herein in order to affect DNA-binding affinity. In addition, these modifications affecting DNA-binding affinity can be combined with one or more of the novel modifications of the base contact residues described above which alter the sequence specificity of the recombinant meganucleases at specific positions within the recognition sequence, or with the prior art modifications described above, or with a combination of the novel modifications and prior art modifications. In particular, by combining backbone contact modifications and base contact modifications, recombinant meganucleases can be rationally-designed with desired specificity and activity. For example, increases in DNA-binding affinity can be designed which may offset losses in affinity resulting from designed changes to base contact residues, or decreases in affinity can be designed which may also decrease sequence specificity and broaden the set of recognition sequences for an enzyme.

**[0023]** In another aspect, the invention provides for rationally-designed meganuclease monomers with altered affinity for homo- or heterodimer formation. The affinity for dimer formation can be measured with the same monomer (*i.e.*, homodimer formation) or with a different monomer (*i.e.*, heterodimer formation) such as a reference wild-type meganuclease. These recombinant meganucleases have modifications to the

amino acid residues which are present at the protein-protein interface between monomers in a meganuclease dimer. The modifications can be used to promote heterodimer formation and create meganucleases with non-palindromic recognition sequences.

[0024] Thus, in some embodiments, the invention provides recombinant meganuclease monomers having altered affinity for dimer formation with a reference meganuclease monomer, in which the recombinant monomer includes a polypeptide having at least 85% sequence similarity to residues 2-153 of the I-CreI meganuclease of SEQ ID NO: 1, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (a) substitution of K7, K57 or K96 with D or E, or (b) substitution of E8 or E61 with K or R. Based upon such recombinant monomers, the invention also provides recombinant meganuclease heterodimers including (1) a first polypeptide having at least 85% sequence similarity to residues 2-153 of the I-CreI meganuclease of SEQ ID NO: 1, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (a) substitution of K7, K57 or K96 with D or E, and (2) a second polypeptide having at least 85% sequence similarity to residues 2-153 of the I-CreI meganuclease of SEQ ID NO: 1, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (b) substitution of E8 or E61 with K or R.

[0025] In other embodiments, the invention provides recombinant meganuclease monomers having altered affinity for dimer formation with a reference meganuclease monomer, in which the recombinant monomer includes a polypeptide having at least 85% sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (a) substitution of R302 with D or E, or (b) substitution of D20, E11 or Q64 with K or R. Based upon such recombinant monomers, the invention also provides recombinant meganuclease heterodimers including (1) a first polypeptide having at least 85% sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (a) substitution of R302 with D or E, and (2) a second polypeptide having at least 85%

sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (b) substitution of D20, E11 or Q64 with K or R.

[0026] In other embodiments, the invention provides recombinant meganuclease monomers having altered affinity for dimer formation with a reference meganuclease monomer, in which the recombinant monomer includes a polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (a) substitution of R93 with D or E, or (b) substitution of E152 with K or R. Based upon such recombinant monomers, the invention also provides recombinant meganuclease heterodimers including (1) a first polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (a) substitution of R93 with D or E, and (2) a second polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12, but in which affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from (b) substitution of E152 with K or R.

[0027] The recombinant meganuclease monomers or heterodimers with altered affinity for dimer formation can also include one, two, three or more of the modifications of base contact residues described above; one, two, three or more of the modifications of backbone contact residues described above; or combinations of both. Thus, for example, the base contacts of a monomer can be modified to alter sequence specificity, the backbone contacts of a monomer can be modified to alter DNA-binding affinity, and the protein-protein interface can be modified to affect dimer formation. Such a recombinant monomer can be combined with a similarly modified monomer to produce a rationally-designed meganuclease heterodimer with desired sequence specificity and activity.

[0028] In another aspect, the invention provides for various methods of use for the rationally-designed meganucleases described and enabled herein. These methods include producing genetically-modified cells and organisms, treating diseases by gene therapy,

treating pathogen infections, and using the recombinant meganucleases for *in vitro* applications for diagnostics and research.

[0029] Thus, in one aspect, the invention provides methods for producing a genetically-modified eukaryotic cell including an exogenous sequence of interest inserted in a chromosome, by transfecting the cell with (i) a first nucleic acid sequence encoding a meganuclease of the invention, and (ii) a second nucleic acid sequence including said sequence of interest, wherein the meganuclease produces a cleavage site in the chromosome and the sequence of interest is inserted into the chromosome at the cleavage site either by homologous recombination or non-homologous end-joining.

[0030] Alternatively, in another aspect, the invention provides methods for producing a genetically-modified eukaryotic cell including an exogenous sequence of interest inserted in a chromosome, by introducing a meganuclease protein of the invention into the cell, and transfecting the cell with a nucleic acid including the sequence of interest, wherein the meganuclease produces a cleavage site in the chromosome and the sequence of interest is inserted into the chromosome at the cleavage site either by homologous recombination or non-homologous end-joining.

[0031] In another aspect, the invention provides methods for producing a genetically-modified eukaryotic cell by disrupting a target sequence in a chromosome, by transfecting the cell with a nucleic acid encoding a meganuclease of the invention, wherein the meganuclease produces a cleavage site in the chromosome and the target sequence is disrupted by non-homologous end-joining at the cleavage site.

[0032] In another aspect, the invention provides methods of producing a genetically-modified organism by producing a genetically-modified eukaryotic cell according to the methods described above, and growing the genetically-modified eukaryotic cell to produce the genetically-modified organism. In these embodiments, the eukaryotic cell can be selected from a gamete, a zygote, a blastocyst cell, an embryonic stem cell, and a protoplast cell.

[0033] In another aspect, the invention provides methods for treating a disease by gene therapy in a eukaryote, by transfecting at least one cell of the eukaryote with one or more nucleic acids including (i) a first nucleic acid sequence encoding a meganuclease of the invention, and (ii) a second nucleic acid sequence including a sequence of interest,

wherein the meganuclease produces a cleavage site in the chromosome and the sequence of interest is inserted into the chromosome by homologous recombination or non-homologous end-joining, and insertion of the sequence of interest provides gene therapy for the disease.

[0034] Alternatively, in another aspect, the invention provides methods for treating a disease by gene therapy in a eukaryote, by introducing a meganuclease protein of the invention into at least one cell of the eukaryote, and transfecting the cell with a nucleic acid including a sequence of interest, wherein the meganuclease produces a cleavage site in the chromosome and the sequence of interest is inserted into the chromosome at the cleavage site by homologous recombination or non-homologous end-joining, and insertion of the sequence of interest provides gene therapy for the disease.

[0035] In another aspect, the invention provides methods for treating a disease by gene therapy in a eukaryote by disrupting a target sequence in a chromosome of the eukaryotic, by transfecting at least one cell of the eukaryote with a nucleic acid encoding a meganuclease of the invention, wherein the meganuclease produces a cleavage site in the chromosome and the target sequence is disrupted by non-homologous end-joining at the cleavage site, wherein disruption of the target sequence provides the gene therapy for the disease.

[0036] In another aspect, the invention provides methods for treating a viral or prokaryotic pathogen infection in a eukaryotic host by disrupting a target sequence in a genome of the pathogen, by transfecting at least one infected cell of the host with a nucleic acid encoding a meganuclease of the invention, wherein the meganuclease produces a cleavage site in the genome and the target sequence is disrupted by either (1) non-homologous end-joining at the cleavage site or (2) by homologous recombination with a second nucleic acid, and wherein disruption of the target sequence provides treatment for the infection.

[0037] More generally, in another aspect, the invention provides methods for rationally-designing recombinant meganucleases having altered specificity for at least one base position of a recognition sequence, by (1) determining at least a portion of a three-dimensional structure of a reference meganuclease-DNA complex; (2) identifying amino acid residues forming a base contact surface at the base position; (3) determining a

distance between a  $\beta$ -carbon of at least a first residue of the contact surface and at least a first base at the base position; and (4) identifying an amino acid substitution to promote the desired change by either (a) for a first residue which is  $< 6 \text{ \AA}$  from the first base, selecting a substitution from Group 1 and/or Group 2 which is a member of an appropriate one of Group G, Group C, Group T or Group A; or (b) for a first residue which is  $> 6 \text{ \AA}$  from said first base, selecting a substitution from Group 2 and/or Group 3 which is a member of an appropriate one of Group G, Group C, Group T or Group A, where each of the Groups is defined herein. This method may be repeated for additional contact residues for the same base, and for contact residues for the other base at the same position, as well as for additional positions.

[0038] In addition, in another general aspect, the invention provides methods for rationally-designing a recombinant meganuclease having increased DNA-binding affinity, by (1) determining at least a portion of a three-dimensional structure of a reference meganuclease-DNA complex; (2) identifying amino acid contact residues forming a backbone contact surface; and (3) identifying an amino acid substitution to increase the DNA-binding affinity by (a) for a contact residue having a negatively-charged or hydrophobic side chain, selecting a substitution having an uncharged/polar or positively-charged side chain; or (b) for a contact residue having an uncharged/polar side chain, selecting a substitution having a positively-charged side chain. Conversely, the invention also provides methods for rationally-designing a recombinant meganuclease having decreased DNA-binding affinity, by (1) determining at least a portion of a three-dimensional structure of a reference meganuclease-DNA complex; (2) identifying amino acid contact residues forming a backbone contact surface; (3) identifying an amino acid substitution to decrease the DNA-binding affinity by (a) for a contact residue having a positively-charged side chain, selecting a substitution having an uncharged/polar or negatively-charged side chain; or (b) for a contact residue having an hydrophobic or uncharged/polar side chain, selecting a substitution having a negatively-charged side chain.

[0039] These and other aspects and embodiments of the invention will be apparent to one of ordinary skill in the art based upon the following detailed description of the invention.



### BRIEF DESCRIPTION OF THE FIGURES

[0040] Figure 1(A) illustrates the interactions between the I-CreI homodimer and its naturally-occurring double-stranded recognition sequence, based upon crystallographic data. This schematic representation depicts the recognition sequence (SEQ ID NO: 2 and SEQ ID NO: 3), shown as unwound for illustration purposes only, bound by the homodimer, shown as two ovals. The bases of each DNA half-site are numbered -1 through -9, and the amino acid residues of I-CreI which form the recognition surface are indicated by one-letter amino acid designations and numbers indicating residue position. Solid black lines: hydrogen bonds to DNA bases. Dashed lines: amino acid positions that form additional contacts in enzyme designs but do not contact the DNA in the wild-type complex. Arrows: residues that interact with the DNA backbone and influence cleavage activity.

[0041] Figure 1(B) illustrates the wild-type contacts between the A-T base pair at position -4 of the cleavage half-site on the right side of Figure 1(A). Specifically, the residue Q26 is shown to interact with the A base. Residue I77 is in proximity to the base pair but not specifically interacting.

[0042] Figure 1(C) illustrates the interactions between a rationally-designed variant of the I-CreI meganuclease in which residue I77 has been modified to E77. As a result of this change, a G-C base pair is preferred at position -4. The interaction between Q26 and the G base is mediated by a water molecule, as has been observed crystallographically for the cleavage half-site on the left side of Figure 1(A).

[0043] Figure 1(D) illustrates the interactions between a rationally-designed variant of the I-CreI meganuclease in which residue Q26 has been modified to E26 and residue I77 has been modified to R77. As a result of this change, a C-G base pair is preferred at position -4.

[0044] Figure 1(E) illustrates the interactions between a rationally-designed variant of the I-CreI meganuclease in which residue Q26 has been modified to A26 and residue I77 has been modified to Q77. As a result of this change, a T-A base pair is preferred at position -4.

[0045] Figure 2(A) shows a comparison of one recognition sequence for each of the wild type I-CreI meganuclease (WT) and 11 rationally-designed meganuclease heterodimers of the invention. Bases that are conserved relative to the WT recognition sequence are shaded. The 9 bp half-sites are bolded. WT: wild-type (SEQ ID NO: 4); CF:  $\Delta$ F508 allele of the human CFTR gene responsible for most cases of cystic fibrosis (SEQ ID NO: 25); MYD: the human DM kinase gene associated with myotonic dystrophy (SEQ ID NO: 27); CCR: the human CCR5 gene (a major HIV co-receptor) (SEQ ID NO: 26); ACH: the human FGFR3 gene correlated with achondroplasia (SEQ ID NO: 23); TAT: the HIV-1 TAT/REV gene (SEQ ID NO: 15); HSV: the HSV-1 UL36 gene (SEQ ID NO: 28); LAM: the bacteriophage  $\lambda$  p05 gene (SEQ ID NO: 22); POX: the Variola (smallpox) virus gp009 gene (SEQ ID NO: 30); URA: the *Saccharomyces cerevisiae* URA3 gene (SEQ ID NO: 36); GLA: the *Arabidopsis thaliana* GL2 gene (SEQ ID NO: 32); BRP: the *Arabidopsis thaliana* BP-1 gene (SEQ ID NO: 33).

[0046] Figure 2(B) illustrates the results of incubation of each of wild-type I-CreI (WT) and 11 rationally-designed meganuclease heterodimers with plasmids harboring the recognition sites for all 12 enzymes for 6 hours at 37°C. Percent cleavage is indicated in each box.

[0047] Figure 3 illustrates cleavage patterns of wild-type and rationally-designed I-CreI homodimers. (A) wild type I-CreI. (B) I-CreI K116D. (C-L) rationally-designed meganucleases of the invention. Enzymes were incubated with a set of plasmids harboring palindromes of the intended cleavage half-site the 27 corresponding single-base pair variations. Bar graphs show fractional cleavage (F) in 4 hours at 37°C. Black bars: expected cleavage patterns based on Table 1. Gray bars: DNA sites that deviate from expected cleavage patterns. White circles indicate bases in the intended recognition site. Also shown are cleavage time-courses over two hours. The open circle time-course plots in C and L correspond to cleavage by the CCR1 and BRP2 enzymes lacking the E80Q mutation. The cleavage sites correspond to the 5' (left column) and 3' (right column) half-sites for the heterodimeric enzymes described in Fig. 2(A).

## DETAILED DESCRIPTION OF THE INVENTION

### 1.1 Introduction

[0048] The present invention is based, in part, upon the identification and characterization of specific amino acids in the LAGLIDADG family of meganucleases that make specific contacts with DNA bases and non-specific contacts with the DNA backbone when the meganucleases associate with a double-stranded DNA recognition sequence, and which thereby affect the recognition sequence specificity and DNA-binding affinity of the enzymes. This discovery has been used, as described in detail below, to identify amino acid substitutions in the meganucleases that can alter the specificity and/or affinity of the enzymes, and to rationally design and develop meganucleases that can recognize a desired DNA sequence that naturally-occurring meganucleases do not recognize, and/or that have increased or decreased specificity and/or affinity relative to the naturally-occurring meganucleases. Furthermore, because DNA-binding affinity affects enzyme activity as well as sequence-specificity, the invention provides rationally-designed meganucleases with altered activity relative to naturally-occurring meganucleases. In addition, the invention provides rationally-designed meganucleases in which residues at the interface between the monomers associated to form a dimer have been modified in order to promote heterodimer formation. Finally, the invention provides uses for the rationally-designed meganucleases in the production of recombinant cells and organisms, as well as in gene therapy, anti-pathogen, anti-cancer, and *in vitro* applications, as disclosed herein.

[0049] As a general matter, the invention provides methods for generating rationally-designed LAGLIDADG meganucleases containing altered amino acid residues at sites within the meganuclease that are responsible for (1) sequence-specific binding to individual bases in the double-stranded DNA recognition sequence, or (2) non-sequence-specific binding to the phosphodiester backbone of a double-stranded DNA molecule. Because enzyme activity is correlated to DNA-binding affinity, however, altering the amino acids involved in binding to the DNA recognition sequence can alter not only the specificity of the meganuclease through specific base pair interactions, but also the activity of the meganuclease by increasing or decreasing overall binding affinity for the

double-stranded DNA. Similarly, altering the amino acids involved in binding to the DNA backbone can alter not only the activity of the enzyme, but also the degree of specificity or degeneracy of binding to the recognition sequence by increasing or decreasing overall binding affinity for the double-stranded DNA.

[0050] As described in detail below, the methods of rationally-designing meganucleases include the identification of the amino acids responsible for DNA recognition/binding, and the application of a series of rules for selecting appropriate amino acid changes. With respect to meganuclease sequence specificity, the rules include both steric considerations relating to the distances in a meganuclease-DNA complex between the amino acid side chains of the meganuclease and the bases in the sense and anti-sense strands of the DNA, and considerations relating to the non-covalent chemical interactions between functional groups of the amino acid side chains and the desired DNA base at the relevant position.

[0051] Finally, a majority of natural meganucleases that bind DNA as homodimers recognize pseudo- or completely palindromic recognition sequences. Because lengthy palindromes are expected to be rare, the likelihood of encountering a palindromic sequence at a genomic site of interest is exceedingly low. Consequently, if these enzymes are to be redesigned to recognize genomic sites of interest, it is necessary to design two enzyme monomers recognizing different half-sites that can heterodimerize to cleave the non-palindromic hybrid recognition sequence. Therefore, in some aspects, the invention provides rationally-designed meganucleases in which monomers differing by at least one amino acid position are dimerized to form heterodimers. In some cases, both monomers are rationally-designed to form a heterodimer which recognizes a non-palindromic recognition sequence. A mixture of two different monomers can result in up to three active forms of meganuclease dimer: the two homodimers and the heterodimer. In addition or alternatively, in some cases, amino acid residues are altered at the interfaces at which monomers can interact to form dimers, in order to increase or decrease the likelihood of formation of homodimers or heterodimers.

[0052] Thus, in one aspect, the invention provide methods for rationally designing LAGLIDADG meganucleases containing amino acid changes that alter the specificity and/or activity of the enzymes. In another aspect, the invention provides the rationally-

designed meganucleases resulting from these methods. In another aspect, the invention provides methods that use such rationally-designed meganucleases to produce recombinant nucleic acids and organisms in which a desired DNA sequence or genetic locus within the genome of an organism is modified by the insertion, deletion, substitution or other manipulation of DNA sequences. In another aspect, the invention provides methods for reducing the survival of pathogens or cancer cells using rationally-designed meganucleases which have pathogen-specific or cancer-specific recognition sequences.

## 1.2 References and Definitions

[0053] The patent and scientific literature referred to herein establishes knowledge that is available to those of skill in the art.

[0054] As used herein, the term "meganuclease" refers to an endonuclease that binds double-stranded DNA at a recognition sequence that is greater than 12 base pairs. Naturally-occurring meganucleases can be monomeric (*e.g.*, I-SceI) or dimeric (*e.g.*, I-CreI). The term meganuclease, as used herein, can be used to refer to monomeric meganucleases, dimeric meganucleases, or to the monomers which associate to form a dimeric meganuclease. The term "homing endonuclease" is synonymous with the term "meganuclease."

[0055] As used herein, the term "LAGLIDADG meganuclease" refers either to meganucleases including a single LAGLIDADG motif, which are naturally dimeric, or to meganucleases including two LAGLIDADG motifs, which are naturally monomeric. The term "mono-LAGLIDADG meganuclease" is used herein to refer to meganucleases including a single LAGLIDADG motif, and the term "di-LAGLIDADG meganuclease" is used herein to refer to meganucleases including two LAGLIDADG motifs, when it is necessary to distinguish between the two. Each of the two structural domains of a di-LAGLIDADG meganuclease which includes a LAGLIDADG motif can be referred to as a LAGLIDADG subunit.

[0056] As used herein, the term "rationally-designed" means non-naturally occurring and/or genetically engineered. The rationally-designed meganucleases of the invention differ from wild-type or naturally-occurring meganucleases in their amino acid sequence or primary structure, and may also differ in their secondary, tertiary or quaternary structure. In addition, the rationally-designed meganucleases of the invention also differ from wild-type or naturally-occurring meganucleases in recognition sequence-specificity and/or activity.

[0057] As used herein, with respect to a protein, the term "recombinant" means having an altered amino acid sequence as a result of the application of genetic engineering techniques to nucleic acids which encode the protein, and cells or organisms which express the protein. With respect to a nucleic acid, the term "recombinant" means having an altered nucleic acid sequence as a result of the application of genetic engineering techniques. Genetic engineering techniques include, but are not limited to, PCR and DNA cloning technologies; transfection, transformation and other gene transfer technologies; homologous recombination; site-directed mutagenesis; and gene fusion. In accordance with this definition, a protein having an amino acid sequence identical to a naturally-occurring protein, but produced by cloning and expression in a heterologous host, is not considered recombinant.

[0058] As used herein with respect to recombinant proteins, the term "modification" means any insertion, deletion or substitution of an amino acid residue in the recombinant sequence relative to a reference sequence (*e.g.*, a wild-type).

[0059] As used herein, the term "genetically-modified" refers to a cell or organism in which, or in an ancestor of which, a genomic DNA sequence has been deliberately modified by recombinant technology. As used herein, the term "genetically-modified" encompasses the term "transgenic."

[0060] As used herein, the term "wild-type" refers to any naturally-occurring form of a meganuclease. The term "wild-type" is not intended to mean the most common allelic variant of the enzyme in nature but, rather, any allelic variant found in nature. Wild-type meganucleases are distinguished from recombinant or non-naturally-occurring meganucleases.

[0061] As used herein, the term "recognition sequence half-site" or simply "half site" means a nucleic acid sequence in a double-stranded DNA molecule which is recognized by a monomer of a mono-LAGLIDADG meganuclease or by one LAGLIDADG subunit of a di-LAGLIDADG meganuclease.

[0062] As used herein, the term "recognition sequence" refers to a pair of half-sites which is bound and cleaved by either a mono-LAGLIDADG meganuclease dimer or a di-LAGLIDADG meganuclease monomer. The two half-sites may or may not be separated by base pairs that are not specifically recognized by the enzyme. In the cases of I-CreI, I-MsoI and I-CeuI, the recognition sequence half-site of each monomer spans 9 base pairs, and the two half-sites are separated by four base pairs which are not recognized specifically but which constitute the actual cleavage site (which has a 4 base pair overhang). Thus, the combined recognition sequences of the I-CreI, I-MsoI and I-CeuI meganuclease dimers normally span 22 base pairs, including two 9 base pair half-sites flanking a 4 base pair cleavage site. The base pairs of each half-site are designated -9 through -1, with the -9 position being most distal from the cleavage site and the -1 position being adjacent to the 4 central base pairs, which are designated N<sub>1</sub>-N<sub>4</sub>. The strand of each half-site which is oriented 5' to 3' in the direction from -9 to -1 (*i.e.*, towards the cleavage site), is designated the "sense" strand and the opposite strand is designated the "antisense strand", although neither strand may encode protein. Thus, the "sense" strand of one half-site is the antisense strand of the other half-site. See, for example, Figure 1(A). In the case of the I-SceI meganuclease, which is a di-LAGLIDADG meganuclease monomer, the recognition sequence is an approximately 18 bp non-palindromic sequence, and there are no central base pairs which are not specifically recognized. By convention, one of the two strands is referred to as the "sense" strand and the other the "antisense" strand, although neither strand may encode protein.

[0063] As used herein, the term "specificity" means the ability of a meganuclease to recognize and cleave double-stranded DNA molecules only at a particular sequence of base pairs referred to as the recognition sequence, or only at a particular set of recognition sequences. The set of recognition sequences will share certain conserved positions or sequence motifs, but may be degenerate at one or more positions. A highly-specific

meganuclease is capable of cleaving only one or a very few recognition sequences. Specificity can be determined in a cleavage assay as described in Example 1. As used herein, a meganuclease has "altered" specificity if it binds to and cleaves a recognition sequence which is not bound to and cleaved by a reference meganuclease (*e.g.*, a wild-type) or if the rate of cleavage of a recognition sequence is increased or decreased by a statistically significant ( $p < 0.05$ ) amount relative to a reference meganuclease.

[0064] As used herein, the term "degeneracy" means the opposite of "specificity." A highly-degenerate meganuclease is capable of cleaving a large number of divergent recognition sequences. A meganuclease can have sequence degeneracy at a single position within a half-site or at multiple, even all, positions within a half-site. Such sequence degeneracy can result from (i) the inability of any amino acid in the DNA-binding domain of a meganuclease to make a specific contact with any base at one or more positions in the recognition sequence, (ii) the ability of one or more amino acids in the DNA-binding domain of a meganuclease to make specific contacts with more than one base at one or more positions in the recognition sequence, and/or (iii) sufficient non-specific DNA binding affinity for activity. A "completely" degenerate position can be occupied by any of the four bases and can be designated with an "N" in a half-site. A "partially" degenerate position can be occupied by two or three of the four bases (*e.g.*, either purine (Pu), either pyrimidine (Py), or not G).

[0065] As used herein with respect to meganucleases, the term "DNA-binding affinity" or "binding affinity" means the tendency of a meganuclease to non-covalently associate with a reference DNA molecule (*e.g.*, a recognition sequence or an arbitrary sequence). Binding affinity is measured by a dissociation constant,  $K_D$  (*e.g.*, the  $K_D$  of I-CreI for the WT recognition sequence is approximately 0.1 nM). As used herein, a meganuclease has "altered" binding affinity if the  $K_D$  of the recombinant meganuclease for a reference recognition sequence is increased or decreased by a statistically significant ( $p < 0.05$ ) amount relative to a reference meganuclease.

[0066] As used herein with respect to meganuclease monomers, the term "affinity for dimer formation" means the tendency of a meganuclease monomer to non-covalently associate with a reference meganuclease monomer. The affinity for dimer formation can be measured with the same monomer (*i.e.*, homodimer formation) or with a different



monomer (*i.e.*, heterodimer formation) such as a reference wild-type meganuclease. Binding affinity is measured by a dissociation constant,  $K_D$ . As used herein, a meganuclease has "altered" affinity for dimer formation if the  $K_D$  of the recombinant meganuclease monomer for a reference meganuclease monomer is increased or decreased by a statistically significant ( $p < 0.05$ ) amount relative to a reference meganuclease monomer.

[0067] As used herein, the term "palindromic" refers to a recognition sequence consisting of inverted repeats of identical half-sites. In this case, however, the palindromic sequence need not be palindromic with respect to the four central base pairs, which are not contacted by the enzyme. In the case of dimeric meganucleases, palindromic DNA sequences are recognized by homodimers in which the two monomers make contacts with identical half-sites.

[0068] As used herein, the term "pseudo-palindromic" refers to a recognition sequence consisting of inverted repeats of non-identical or imperfectly palindromic half-sites. In this case, the pseudo-palindromic sequence not only need not be palindromic with respect to the four central base pairs, but also can deviate from a palindromic sequence between the two half-sites. Pseudo-palindromic DNA sequences are typical of the natural DNA sites recognized by wild-type homodimeric meganucleases in which two identical enzyme monomers make contacts with different half-sites.

[0069] As used herein, the term "non-palindromic" refers to a recognition sequence composed of two unrelated half-sites of a meganuclease. In this case, the non-palindromic sequence need not be palindromic with respect to either the four central base pairs or the two monomer half-sites. Non-palindromic DNA sequences are recognized by either di-LAGLIDADG meganucleases, highly degenerate mono-LAGLIDADG meganucleases (*e.g.*, I-CeuI) or by heterodimers of mono-LAGLIDADG meganuclease monomers that recognize non-identical half-sites.

[0070] As used herein, the term "activity" refers to the rate at which a meganuclease of the invention cleaves a particular recognition sequence. Such activity is a measurable enzymatic reaction, involving the hydrolysis of phosphodiester bonds of double-stranded DNA. The activity of a meganuclease acting on a particular DNA substrate is affected by the affinity or avidity of the meganuclease for that particular DNA

substrate which is, in turn, affected by both sequence-specific and non-sequence-specific interactions with the DNA.

[0071] As used herein, the term "homologous recombination" refers to the natural, cellular process in which a double-stranded DNA-break is repaired using a homologous DNA sequence as the repair template (see, *e.g.* Cahill *et al.* (2006), *Front. Biosci.* 11:1958-1976). The homologous DNA sequence may be an endogenous chromosomal sequence or an exogenous nucleic acid that was delivered to the cell. Thus, in some embodiments, a rationally-designed meganuclease is used to cleave a recognition sequence within a target sequence and an exogenous nucleic acid with homology to or substantial sequence similarity with the target sequence is delivered into the cell and used as a template for repair by homologous recombination. The DNA sequence of the exogenous nucleic acid, which may differ significantly from the target sequence, is thereby incorporated into the chromosomal sequence. The process of homologous recombination occurs primarily in eukaryotic organisms. The term "homology" is used herein as equivalent to "sequence similarity" and is not intended to require identity by descent or phylogenetic relatedness.

[0072] As used herein, the term "non-homologous end-joining" refers to the natural, cellular process in which a double-stranded DNA-break is repaired by the direct joining of two non-homologous DNA segments (see, *e.g.* Cahill *et al.* (2006), *Front. Biosci.* 11:1958-1976). DNA repair by non-homologous end-joining is error-prone and frequently results in the untemplated addition or deletion of DNA sequences at the site of repair. Thus, in certain embodiments, a rationally-designed meganuclease can be used to produce a double-stranded break at a meganuclease recognition sequence within a target sequence to disrupt a gene (*e.g.*, by introducing base insertions, base deletions, or frameshift mutations) by non-homologous end-joining. In other embodiments, an exogenous nucleic acid lacking homology to or substantial sequence similarity with the target sequence may be captured at the site of a meganuclease-stimulated double-stranded DNA break by non-homologous end-joining (see, *e.g.* Salomon, *et al.* (1998), *EMBO J.* 17:6086-6095). The process of non-homologous end-joining occurs in both eukaryotes and prokaryotes such as bacteria.

[0073] As used herein, the term "sequence of interest" means any nucleic acid sequence, whether it codes for a protein, RNA, or regulatory element (*e.g.*, an enhancer, silencer, or promoter sequence), that can be inserted into a genome or used to replace a genomic DNA sequence using a meganuclease protein. Sequences of interest can have heterologous DNA sequences that allow for tagging a protein or RNA that is expressed from the sequence of interest. For instance, a protein can be tagged with tags including, but not limited to, an epitope (*e.g.*, c-myc, FLAG) or other ligand (*e.g.*, poly-His). Furthermore, a sequence of interest can encode a fusion protein, according to techniques known in the art (see, *e.g.*, Ausubel *et al.*, Current Protocols in Molecular Biology, Wiley 1999). In some embodiments, the sequence of interest is flanked by a DNA sequence that is recognized by the recombinant meganuclease for cleavage. Thus, the flanking sequences are cleaved allowing for proper insertion of the sequence of interest into genomic recognition sequences cleaved by the recombinant meganuclease. In some embodiments, the entire sequence of interest is homologous to or has substantial sequence similarity with the a target sequence in the genome such that homologous recombination effectively replaces the target sequence with the sequence of interest. In other embodiments, the sequence of interest is flanked by DNA sequences with homology to or substantial sequence similarity with the target sequence such that homologous recombination inserts the sequence of interest within the genome at the locus of the target sequence. In some embodiments, the sequence of interest is substantially identical to the target sequence except for mutations or other modifications in the meganuclease recognition sequence such that the meganuclease can not cleave the target sequence after it has been modified by the sequence of interest.

[0074] As used herein with respect to both amino acid sequences and nucleic acid sequences, the terms "percentage similarity" and "sequence similarity" refer to a measure of the degree of similarity of two sequences based upon an alignment of the sequences which maximizes similarity between aligned amino acid residues or nucleotides, and which is a function of the number of identical or similar residues or nucleotides, the number of total residues or nucleotides, and the presence and length of gaps in the sequence alignment. A variety of algorithms and computer programs are available for determining sequence similarity using standard parameters. As used herein, sequence

similarity is measured using the BLASTp program for amino acid sequences and the BLASTn program for nucleic acid sequences, both of which are available through the National Center for Biotechnology Information ([www.ncbi.nlm.nih.gov/](http://www.ncbi.nlm.nih.gov/)), and are described in, for example, Altschul *et al.* (1990), *J. Mol. Biol.* 215:403-410; Gish and States (1993), *Nature Genet.* 3:266-272; Madden *et al.* (1996), *Meth. Enzymol.* 266:131-141; Altschul *et al.* (1997), *Nucleic Acids Res.* 25:3389-3402; Zhang *et al.* (2000), *J. Comput. Biol.* 7(1-2):203-14. As used herein, percent similarity of two amino acid sequences is the score based upon the following parameters for the BLASTp algorithm: word size = 3; gap opening penalty = -11; gap extension penalty = -1; and scoring matrix = BLOSUM62. As used herein, percent similarity of two nucleic acid sequences is the score based upon the following parameters for the BLASTn algorithm: word size = 11; gap opening penalty = -5; gap extension penalty = -2; match reward = 1; and mismatch penalty = -3.

[0075] As used herein with respect to modifications of two proteins or amino acid sequences, the term "corresponding to" is used to indicate that a specified modification in the first protein is a substitution of the same amino acid residue as in the modification in the second protein, and that the amino acid position of the modification in the first proteins corresponds to or aligns with the amino acid position of the modification in the second protein when the two proteins are subjected to standard sequence alignments (*e.g.*, using the BLASTp program). Thus, the modification of residue "X" to amino acid "A" in the first protein will correspond to the modification of residue "Y" to amino acid "A" in the second protein if residues X and Y correspond to each other in a sequence alignment, and despite the fact that X and Y may be different numbers.

[0076] As used herein, the recitation of a numerical range for a variable is intended to convey that the invention may be practiced with the variable equal to any of the values within that range. Thus, for a variable which is inherently discrete, the variable can be equal to any integer value within the numerical range, including the end-points of the range. Similarly, for a variable which is inherently continuous, the variable can be equal to any real value within the numerical range, including the end-points of the range. As an example, and without limitation, a variable which is described as having values between 0 and 2 can take the values 0, 1 or 2 if the variable is inherently discrete, and can

take the values 0.0, 0.1, 0.01, 0.001, or any other real values  $\geq 0$  and  $\leq 2$  if the variable is inherently continuous.

[0077] As used herein, unless specifically indicated otherwise, the word "or" is used in the inclusive sense of "and/or" and not the exclusive sense of "either/or."

#### 2.1 Rationally-Designed Meganucleases with Altered Sequence-Specificity

[0078] In one aspect of the invention, methods for rationally designing recombinant LAGLIDADG family meganucleases are provided. In this aspect, recombinant meganucleases are rationally-designed by first predicting amino acid substitutions that can alter base preference at each position in the half-site. These substitutions can be experimentally validated individually or in combinations to produce meganucleases with the desired cleavage specificity.

[0079] In accordance with the invention, amino acid substitutions that can cause a desired change in base preference are predicted by determining the amino acid side chains of a reference meganuclease (*e.g.*, a wild-type meganuclease, or a non-naturally-occurring reference meganuclease) that are able to participate in making contacts with the nucleic acid bases of the meganuclease's DNA recognition sequence and the DNA phosphodiester backbone, and the spatial and chemical nature of those contacts. These amino acids include but are not limited to side chains involved in contacting the reference DNA half-site. Generally, this determination requires having knowledge of the structure of the complex between the meganuclease and its double-stranded DNA recognition sequence, or knowledge of the structure of a highly similar complex (*e.g.*, between the same meganuclease and an alternative DNA recognition sequence, or between an allelic or phylogenetic variant of the meganuclease and its DNA recognition sequence).

[0080] Three-dimensional structures, as described by atomic coordinates data, of a polypeptide or complex of two or more polypeptides can be obtained in several ways. For example, protein structure determinations can be made using techniques including, but not limited to, X-ray crystallography, NMR, and mass spectrometry. Another approach is to analyze databases of existing structural co-ordinates for the meganuclease of interest or a related meganuclease. Such structural data is often available from

databases in the form of three-dimensional coordinates. Often this data is accessible through online databases (*e.g.*, the RCSB Protein Data Bank at [www.rcsb.org/pdb](http://www.rcsb.org/pdb)).

[0081] Structural information can be obtained experimentally by analyzing the diffraction patterns of, for example, X-rays or electrons, created by regular two- or three-dimensional arrays (*e.g.*, crystals) of proteins or protein complexes. Computational methods are used to transform the diffraction data into three-dimensional atomic coordinates in space. For example, the field of X-ray crystallography has been used to generate three-dimensional structural information on many protein-DNA complexes, including meganucleases (see, *e.g.*, Chevalier *et al.* (2001), *Nucleic Acids Res.* 29(18): 3757-3774).

[0082] Nuclear Magnetic Resonance (NMR) also has been used to determine inter-atomic distances of molecules in solution. Multi-dimensional NMR methods combined with computational methods have succeeded in determining the atomic coordinates of polypeptides of increasing size (see, *e.g.*, Tzakos *et al.* (2006), *Annu. Rev. Biophys. Biomol. Struct.* 35:19-42.).

[0083] Alternatively, computational modeling can be used by applying algorithms based on the known primary structures and, when available, secondary, tertiary and/or quaternary structures of the protein/DNA, as well as the known physiochemical nature of the amino acid side chains, nucleic acid bases, and bond interactions. Such methods can optionally include iterative approaches, or experimentally-derived constraints. An example of such computational software is the CNS program described in Adams *et al.* (1999), *Acta Crystallogr. D. Biol. Crystallogr.* 55 (Pt 1): 181-90. A variety of other computational programs have been developed that predict the spatial arrangement of amino acids in a protein structure and predict the interaction of the amino acid side chains of the protein with various target molecules (see, *e.g.*, U.S. Pat. No. 6,988,041).

[0084] Thus, in some embodiments of the invention, computational models are used to identify specific amino acid residues that specifically interact with DNA nucleic acid bases and/or facilitate non-specific phosphodiester backbone interactions. For instance, computer models of the totality of the potential meganuclease-DNA interaction can be produced using a suitable software program, including, but not limited to, MOLSCRIPT™ 2.0 (Avatar Software AB, Stockholm, Sweden), the graphical display

program O (Jones *et al.* (1991), *Acta Crystallography*, A47: 110), the graphical display program GRASP™ (Nicholls *et al.* (1991), *PROTEINS, Structure, Function and Genetics* 11(4): 281ff), or the graphical display program INSIGHT™ (TSI, Inc., Shoreview, MN). Computer hardware suitable for producing, viewing and manipulating three-dimensional structural representations of protein-DNA complexes are commercially available and well known in the art (*e.g.*, Silicon Graphics Workstation, Silicon Graphics, Inc., Mountainview, CA).

[0085] Specifically, interactions between a meganuclease and its double-stranded DNA recognition sequences can be resolved using methods known in the art. For example, a representation, or model, of the three dimensional structure of a multi-component complex structure, for which a crystal has been produced, can be determined using techniques which include molecular replacement or SIR/MIR (single/multiple isomorphous replacement) (*see, e.g.*, Brunger (1997), *Meth. Enzym.* 276: 558-580; Navaza and Saludjian (1997), *Meth. Enzym.* 276: 581-594; Tong and Rossmann (1997), *Meth. Enzym.* 276: 594-611; and Bentley (1997), *Meth. Enzym.* 276: 611-619) and can be performed using a software program, such as AMoRe/Mosflm (Navaza (1994), *Acta Cryst.* A50: 157-163; CCP4 (1994), *Acta Cryst.* D50: 760-763) or XPLOR (*see, Brünger et al.* (1992), X-PLOR Version 3.1. A System for X-ray Crystallography and NMR, Yale University Press, New Haven, CT).

[0086] The determination of protein structure and potential meganuclease-DNA interaction allows for rational choices concerning the amino acids that can be changed to affect enzyme activity and specificity. Decisions are based on several factors regarding amino acid side chain interactions with a particular base or DNA phosphodiester backbone. Chemical interactions used to determine appropriate amino acid substitutions include, but are not limited to, van der Waals forces, steric hindrance, ionic bonding, hydrogen bonding, and hydrophobic interactions. Amino acid substitutions can be selected which either favor or disfavor specific interactions of the meganuclease with a particular base in a potential recognition sequence half-site in order to increase or decrease specificity for that sequence and, to some degree, overall binding affinity and activity. In addition, amino acid substitutions can be selected which either increase or decrease binding affinity for the phosphodiester backbone of double-stranded DNA in

order to increase or decrease overall activity and, to some degree, to decrease or increase specificity.

[0087] Thus, in specific embodiments, a three-dimensional structure of a meganuclease-DNA complex is determined and a "contact surface" is defined for each base-pair in a DNA recognition sequence half-site. In some embodiments, the contact surface comprises those amino acids in the enzyme with  $\beta$ -carbons less than 9.0 Å from a major groove hydrogen-bond donor or acceptor on either base in the pair, and with side chains oriented toward the DNA, irrespective of whether the residues make base contacts in the wild-type meganuclease-DNA complex. In other embodiments, residues can be excluded if the residues do not make contact in the wild-type meganuclease-DNA complex, or residues can be included or excluded at the discretion of the designer to alter the number or identity of the residues considered. In one example, as described below, for base positions -2, -7, -8, and -9 of the wild-type I-CreI half-site, the contact surfaces were limited to the amino acid positions that actually interact in the wild-type enzyme-DNA complex. For positions -1, -3, -4, -5, and -6, however, the contact surfaces were defined to contain additional amino acid positions that are not involved in wild-type contacts but which could potentially contact a base if substituted with a different amino acid.

[0088] It should be noted that, although a recognition sequence half-site is typically represented with respect to only one strand of DNA, meganucleases bind in the major groove of double-stranded DNA, and make contact with nucleic acid bases on both strands. In addition, the designations of "sense" and "antisense" strands are completely arbitrary with respect to meganuclease binding and recognition. Sequence specificity at a position can be achieved either through interactions with one member of a base pair, or by a combination of interactions with both members of a base-pair. Thus, for example, in order to favor the presence of an A/T base pair at position X, where the A base is on the "sense" strand and the T base is on the "antisense" strand, residues are selected which are sufficiently close to contact the sense strand at position X and which favor the presence of an A, and/or residues are selected which are sufficiently close to contact the antisense strand at position X and which favor the presence of a T. In accordance with the



invention, a residue is considered sufficiently close if the  $\beta$ -carbon of the residue is within 9Å of the closest atom of the relevant base.

[0089] Thus, for example, an amino acid with a  $\beta$ -carbon within 9Å of the DNA sense strand but greater than 9Å from the antisense strand is considered for potential interactions with only the sense strand. Similarly, an amino acid with a  $\beta$ -carbon within 9Å of the DNA antisense strand but greater than 9Å from the sense strand is considered for potential interactions with only the antisense strand. Amino acids with  $\beta$ -carbons that are within 9Å of both DNA strands are considered for potential interactions with either strand.

[0090] For each contact surface, potential amino acid substitutions are selected based on their predicted ability to interact favorably with one or more of the four DNA bases. The selection process is based upon two primary criteria: (i) the size of the amino acid side chains, which will affect their steric interactions with different nucleic acid bases, and (ii) the chemical nature of the amino acid side chains, which will affect their electrostatic and bonding interactions with the different nucleic acid bases.

[0091] With respect to the size of side chains, amino acids with shorter and/or smaller side chains can be selected if an amino acid  $\beta$ -carbon in a contact surface is  $<6$  Å from a base, and amino acids with longer and/or larger side chains can be selected if an amino acid  $\beta$ -carbon in a contact surface is  $>6$  Å from a base. Amino acids with side chains that are intermediate in size can be selected if an amino acid  $\beta$ -carbon in a contact surface is 5-8 Å from a base.

[0092] The amino acids with relatively shorter and smaller side chains can be assigned to Group 1, including glycine (G), alanine (A), serine (S), threonine (T), cysteine (C), valine (V), leucine (L), isoleucine (I), aspartate (D), asparagine (N) and proline (P). Proline, however, is expected to be used less frequently because of its relative inflexibility. In addition, glycine is expected to be used less frequently because it introduces unwanted flexibility in the peptide backbone and its very small size reduces the likelihood of effective contacts when it replaces a larger residue. On the other hand, glycine can be used in some instances for promoting a degenerate position. The amino acids with side chains of relatively intermediate length and size can be assigned to Group 2, including lysine (K), methionine (M), arginine (R), glutamate (E) and glutamine (Q).

The amino acids with relatively longer and/or larger side chains can be assigned to Group 3, including lysine (K), methionine (M), arginine (R), histidine (H), phenylalanine (F), tyrosine (Y), and tryptophan (W). Tryptophan, however, is expected to be used less frequently because of its relative inflexibility. In addition, the side chain flexibility of lysine, arginine, and methionine allow these amino acids to make base contacts from long or intermediate distances, warranting their inclusion in both Groups 2 and 3. These groups are also shown in tabular form below:

Group 1	Group 2	Group 3
glycine (G)	glutamine (Q)	arginine (R)
alanine (A)	glutamate (E)	histidine (H)
serine (S)	lysine (K)	phenylalanine (F)
threonine (T)	methionine (M)	tyrosine (Y)
cysteine (C)	arginine (R)	tryptophan (W)
valine (V)		lysine (K)
leucine (L)		methionine (M)
isoleucine (I)		
aspartate (D)		
asparagine (N)		
proline (P)		

[0093] With respect to the chemical nature of the side chains, the different amino acids are evaluated for their potential interactions with the different nucleic acid bases (*e.g.*, van der Waals forces, ionic bonding, hydrogen bonding, and hydrophobic interactions) and residues are selected which either favor or disfavor specific interactions of the meganuclease with a particular base at a particular position in the double-stranded DNA recognition sequence half-site. In some instances, it may be desired to create a half-site with one or more complete or partial degenerate positions. In such cases, one

may choose residues which favor the presence of two or more bases, or residues which disfavor one or more bases. For example, partial degenerate base recognition can be achieved by sterically hindering a pyrimidine at a sense or antisense position.

**[0094]** Recognition of guanine (G) bases is achieved using amino acids with basic side chains that form hydrogen bonds to N7 and O6 of the base. Cytosine (C) specificity is conferred by negatively-charged side chains which interact unfavorably with the major groove electronegative groups present on all bases except C. Thymine (T) recognition is rationally-designed using hydrophobic and van der Waals interactions between hydrophobic side chains and the major groove methyl group on the base. Finally, adenine (A) bases are recognized using the carboxamide side chains Asn and Gln or the hydroxyl side chain of Tyr through a pair of hydrogen bonds to N7 and N6 of the base. Lastly, His can be used to confer specificity for a purine base (A or G) by donating a hydrogen bond to N7. These straightforward rules for DNA recognition can be applied to predict contact surfaces in which one or both of the bases at a particular base-pair position are recognized through a rationally-designed contact.

**[0095]** Thus, based on their binding interactions with the different nucleic acid bases, and the bases which they favor at a position with which they make contact, each amino acid residue can be assigned to one or more different groups corresponding to the different bases they favor (i.e., G, C, T or A). Thus, Group G includes arginine (R), lysine (K) and histidine (H); Group C includes aspartate (D) and glutamate (E); Group T includes alanine (A), valine (V), leucine (L), isoleucine (I), cysteine (C), threonine (T), methionine (M) and phenylalanine (F); and Group A includes asparagine (N), glutamine (N), tyrosine (Y) and histidine (H). Note that histidine appears in both Group G and Group A; that serine (S) is not included in any group but may be used to favor a degenerate position; and that proline, glycine, and tryptophan are not included in any particular group because of predominant steric considerations. These groups are also shown in tabular form below:

Group G	Group C	Group T	Group A
arginine (R)	aspartate (D)	alanine (A)	asparagine (N)
lysine (K)	glutamate (E)	valine (V)	glutamine (Q)
histidine (H)		leucine (L)	tyrosine (Y)
		isoleucine (I)	histidine (H)
		cysteine (C)	
		threonine (T)	
		methionine (M)	
		phenylalanine (F)	

[0096] Thus, in accordance with the invention, in order to effect a desired change in the recognition sequence half-site of a meganuclease at a given position X, (1) determine at least the relevant portion of the three-dimensional structure of the wild-type or reference meganuclease-DNA complex and the amino acid residue side chains which define the contact surface at position X; (2) determine the distance between the  $\beta$ -carbon of at least one residue comprising the contact surface and at least one base of the base pair at position X; and (3)(a) for a residue which is  $< 6 \text{ \AA}$  from the base, select a residue from Group 1 and/or Group 2 which is a member of the appropriate one of Group G, Group C, Group T or Group A to promote the desired change, and/or (b) for a residue which is  $> 6 \text{ \AA}$  from the base, select a residue from Group 2 and/or Group 3 which is a member of the appropriate one of Group G, Group C, Group T or Group A to promote the desired change. More than one such residue comprising the contact surface can be selected for analysis and modification and, in some embodiments, each such residue is analyzed and multiple residues are modified. Similarly, the distance between the  $\beta$ -carbon of a residue included in the contact surface and each of the two bases of the base pair at position X

can be determined and, if the residue is within 9Å of both bases, then different substitutions can be made to affect the two bases of the pair (e.g., a residue from Group 1 to affect a proximal base on one strand, or a residue from Group 3 to affect a distal base on the other strand). Alternatively, a combination of residue substitutions capable of interacting with both bases in a pair can affect the specificity (e.g., a residue from the T Group contacting the sense strand combined with a residue from the A Group contacting the antisense strand to select for T/A). Finally, multiple alternative modifications of the residues can be validated either empirically (e.g., by producing the recombinant meganuclease and testing its sequence recognition) or computationally (e.g., by computer modeling of the meganuclease-DNA complex of the modified enzyme) to choose amongst alternatives.

[0097] Once one or more desired amino acid modifications of the wild-type or reference meganuclease are selected, the rationally-designed meganuclease can be produced by recombinant methods and techniques well known in the art. In some embodiments, non-random or site-directed mutagenesis techniques are used to create specific sequence modifications. Non-limiting examples of non-random mutagenesis techniques include overlapping primer PCR (see, e.g., Wang *et al.* (2006), *Nucleic Acids Res.* 34(2): 517–527), site-directed mutagenesis (see, e.g., U.S. Pat. No. 7,041,814), cassette mutagenesis (see, e.g., U.S. Pat. No. 7,041,814), and the manufacturer's protocol for the Altered Sites® II Mutagenesis Systems kit commercially available from Promega Biosciences, Inc. (San Luis Obispo, CA).

[0098] The recognition and cleavage of a specific DNA sequence by a rationally-designed meganuclease can be assayed by any method known by one skilled in the art (see, e.g., U.S. Pat. Pub. No. 2006/0078552). In certain embodiments, the determination of meganuclease cleavage is determined by *in vitro* cleavage assays. Such assays use *in vitro* cleavage of a polynucleotide substrate comprising the intended recognition sequence of the assayed meganuclease and, in certain embodiments, variations of the intended recognition sequence in which one or more bases in one or both half-sites have been changed to a different base. Typically, the polynucleotide substrate is a double-stranded DNA molecule comprising a target site which has been synthesized and cloned into a vector. The polynucleotide substrate can be linear or circular, and typically comprises

only one recognition sequence. The meganuclease is incubated with the polynucleotide substrate under appropriate conditions, and the resulting polynucleotides are analyzed by known methods for identifying cleavage products (*e.g.*, electrophoresis or chromatography). If there is a single recognition sequence in a linear, double-strand DNA substrate, the meganuclease activity is detected by the appearance of two bands (products) and the disappearance of the initial full-length substrate band. In one embodiment, meganuclease activity can be assayed as described in, for example, Wang *et al.* (1997), *Nucleic Acid Res.*, 25: 3767-3776.

[0099] In other embodiments, the cleavage pattern of the meganuclease is determined using *in vivo* cleavage assays (see, *e.g.*, U.S. Pat. Pub. No. 2006/0078552). In particular embodiments, the *in vivo* test is a single-strand annealing recombination test (SSA). This kind of test is known to those of skill in the art (Rudin *et al.* (1989), *Genetics* 122: 519-534; Fishman-Lobell *et al.* (1992), *Science* 258: 480-4).

[0100] As will be apparent to one of skill in the art, additional amino acid substitutions, insertions or deletions can be made to domains of the meganuclease enzymes other than those involved in DNA recognition and binding without complete loss of activity. Substitutions can be conservative substitutions of similar amino acid residues at structurally or functionally constrained positions, or can be non-conservative substitutions at positions which are less structurally or functionally constrained. Such substitutions, insertions and deletions can be identified by one of ordinary skill in the art by routine experimentation without undue effort. Thus, in some embodiments, the recombinant meganucleases of the invention include proteins having anywhere from 85% to 99% sequence similarity (*e.g.*, 85%, 87.5%, 90%, 92.5%, 95%, 97.5%, 99%) to a reference meganuclease sequence. With respect to each of the wild-type I-CreI, I-MsoI, I-SceI and I-CeuI proteins, the most N-terminal and C-terminal sequences are not clearly visible in X-ray crystallography studies, suggesting that these positions are not structurally or functionally constrained. Therefore, these residues can be excluded from calculation of sequence similarity, and the following reference meganuclease sequences can be used: residues 2-153 of SEQ ID NO: 1 for I-CreI, residues 6-160 of SEQ ID NO: 6 for I-MsoI, residues 3-186 of SEQ ID NO: 9 for I-SceI, and residues 5-211 of SEQ ID NO: 12 for I-CeuI.

## 2.2 LAGLIDADG Family Meganucleases

[0101] The LAGLIDADG meganuclease family is composed of more than 200 members from a diverse phylogenetic group of host organisms. All members of this family have one or two copies of a highly conserved LAGLIDADG motif along with other structural motifs involved in cleavage of specific DNA sequences. Enzymes that have a single copy of the LAGLIDADG motif (*i.e.*, mono-LAGLIDADG meganucleases) function as dimers, whereas the enzymes that have two copies of this motif (*i.e.*, di-LAGLIDADG meganucleases) function as monomers.

[0102] All LAGLIDADG family members recognize and cleave relatively long sequences (> 12bp), leaving four nucleotide 3' overhangs. These enzymes also share a number of structural motifs in addition to the LAGLIDADG motif, including a similar arrangement of anti-parallel  $\beta$ -strands at the protein-DNA interface. Amino acids within these conserved structural motifs are responsible for interacting with the DNA bases to confer recognition sequence specificity. The overall structural similarity between some members of the family (*e.g.*, I-CreI, I-MsoI, I-SceI and I-CeuI) has been elucidated by X-ray crystallography. Accordingly, the members of this family can be modified at particular amino acids within such structural motifs to change the over-all activity or sequence-specificity of the enzymes, and corresponding modifications can reasonable be expected to have similar results in other family members. See, generally, Chevalier *et al.* (2001), *Nucleic Acid Res.* 29(18): 3757-3774).

### 2.2.1 Meganucleases Derived from I-CreI

[0103] In one aspect, the present invention relates to rationally-designed meganucleases which are based upon or derived from the I-CreI meganuclease of *Chlamydomonas reinhardtii*. The wild-type amino acid sequence of the I-CreI meganuclease is shown in SEQ ID NO: 1, which corresponds to Genbank Accession # PO5725. Two recognition sequence half sites of the wild-type I-CreI meganuclease from crystal structure PDB # 1BP7 are shown below:

```

Position   -9-8-7-6-5-4-3-2-1
           5'-G A A A C T G T C T C A C G A C G T T T T G-3' SEQ ID NO: 2
           3'-C T T T G A C A G A G T G C T G C A A A A C-5' SEQ ID NO: 3
Position                                     -1-2-3-4-5-6-7-8-9

```

Note that this natural recognition sequence is not perfectly palindromic, even outside the central four base pairs. The two recognition sequence half-sites are shown in bold on their respective sense strands.

[0104] Wild-type I-CreI also recognizes and cuts the following perfectly palindromic (except for the central N<sub>1</sub>-N<sub>4</sub> bases) sequence:

```

Position   -9-8-7-6-5-4-3-2-1
           5'-C A A A C T G T C G T G A G A C A G T T T G-3' SEQ ID NO: 4
           3'-G T T T G A C A G C A C T C T G T C A A A C-5' SEQ ID NO: 5
Position                                     -1-2-3-4-5-6-7-8-9

```

[0105] The palindromic sequence of SEQ ID NO: 4 and SEQ ID NO: 5 is considered to be a better substrate for the wild-type I-CreI because the enzyme binds this site with higher affinity and cleaves it more efficiently than the natural DNA sequence. For the purposes of the following disclosure, and with particular regard to the experimental results presented herein, this palindromic sequence cleaved by wild-type I-CreI is referred to as "WT" (see, e.g., Figure 2(A)). The two recognition sequence half-sites are shown in bold on their respective sense strands.

[0106] Figure 1(A) depicts the interactions of a wild-type I-CreI meganuclease homodimer with a double-stranded DNA recognition sequence, Figures 1(B) shows the specific interactions between amino acid residues of the enzyme and bases at the -4 position of one half-site for a wild-type enzyme and one wild-type recognition sequence, and Figures 1(C)-(E) show the specific interactions between amino acid residues of the enzyme and bases at the -4 position of one half-site for three rationally-designed meganucleases of the invention with altered specificity at position -4 of the half-site.

[0107] Thus, the base preference at any specified base position of the half-site can be rationally altered to each of the other three base pairs using the methods disclosed herein. First, the wild type recognition surface at the specified base position is determined (e.g., by analyzing meganuclease-DNA complex co-crystal structures; or by



computer modeling of the meganuclease-DNA complexes). Second, existing and potential contact residues are determined based on the distances between the  $\beta$ -carbons of the surrounding amino acid positions and the nucleic acid bases on each DNA strand at the specified base position. For example, and without limitation, as shown in Figure 1(A), the I-CreI wild type meganuclease-DNA contact residues at position -4 involve a glutamine at position 26 which hydrogen bonds to an A base on the antisense DNA strand. Residue 77 was also identified as potentially being able to contact the -4 base on the DNA sense strand. The  $\beta$ -carbon of residue 26 is 5.9 Å away from N7 of the A base on the antisense DNA strand, and the  $\beta$ -carbon of residue 77 is 7.15 Å away from the C5-methyl of the T on the sense strand. According to the distance and base chemistry rules described herein, a C on the sense strand could hydrogen bond with a glutamic acid at position 77 and a G on the antisense strand could bond with glutamine at position 26 (mediated by a water molecule, as observed in the wild-type I-CreI crystal structure) (see Fig 1(C)); a G on the sense strand could hydrogen bond with an arginine at position 77 and a C on the antisense strand could hydrogen bond with a glutamic acid at position 26 (see Fig 1(D)); an A on the sense strand could hydrogen bond with a glutamine at position 77 and a T on the antisense strand could form hydrophobic contacts with an alanine at position 26 (see Fig. 1(E)). If the base specific contact is provided by position 77, then the wild-type contact, Q26, can be substituted (*e.g.*, with a serine residue) to reduce or remove its influence on specificity. Alternatively, complementary mutations at positions 26 and 77 can be combined to specify a particular base pair (*e.g.*, A26 specifies a T on the antisense strand and Q77 specifies an A on the sense strand (Fig. 1(E)). These predicted residue substitutions have all been validated experimentally.

[0108] Thus, in accordance with the invention, a substantial number of amino acid modifications to the DNA recognition domain of the I-CreI meganuclease have been identified which, singly or in combination, result in recombinant meganucleases with specificities altered at individual bases within the DNA recognition sequence half-site, such that these rationally-designed meganucleases have half-sites different from the wild-type enzyme. The amino acid modifications of I-CreI and the resulting change in recognition sequence half-site specificity are shown in Table 1:

TABLE 1

Posn.	Favored Sense-Strand Base										A/C/G/T	
	A	C	G	T	A/T	A/C	A/G	C/T	G/T	A/G/T		
-1	Y75 L75* C75* Y139* C46* A46*	R70* H75* R75* H46* K46* R46*	K70 E70* E75* E46* D46*	Q70* C70 L70 Y75* Q75* H75* H139 Q46* H46*				T46*				G70 A70 S70 G46*
-2	Q70 T44* A44* V44* I44* L44* N44*	E70 D70 K44* R44*	H70 D44* E44*	Q44*	C44*							
-3	Q68 C24* I24*	E68 F68 K24* R24*	R68	M68 C68 L68 F68		H68		Y68	K68			
-4	A26* Q77	E77 K26*	R77 E26*					S77 Q26*				S26*
-5		E42	R42			K28*	C28* Q42					M66 K66

-6	Q40 C28*	E40 R28*	R40	C40 I40 V40 C79 I79 V79 Q28*	A40 A79 A28* H28*						S40 S28*
-7	N30* Q38	E38 K30* R30*	K38 R38 E30*	I38 L38		C38					H38 N38 Q30*
-8	F33 Y33	E33 D33	F33 H33	L33 V33 I33 F33 C33		R32*	R33				
-9		E32	R32 K32	L32 V32 A32 C32				D32 I32			S32 N32 H32 Q32 T32

Bold entries are wild-type contact residues and do not constitute "modifications" as used herein.

An asterisk indicates that the residue contacts the base on the antisense strand.

### 2.2.2 Meganucleases Derived from I-MsoI

[0109] In another aspect, the present invention relates to rationally-designed meganucleases which are based upon or derived from the I-MsoI meganuclease of *Monomastix sp.* The wild-type amino acid sequence of the I-MsoI meganuclease is shown in SEQ ID NO: 6, which corresponds to Genbank Accession # AAL34387. Two recognition sequence half-sites of the wild-type I-MsoI meganuclease from crystal structure PDB # 1M5X are shown below:

Position	-9-8-7-6-5-4-3-2-1	
	5'- <b>C A G A A C G T C G T G A G A C A G T T C C</b> -3'	SEQ ID NO: 7
	3'- <b>G T C T T G C A G C A C T C T G T C A A G G</b> -5'	SEQ ID NO: 8
Position		-1-2-3-4-5-6-7-8-9

Note that the recognition sequence is not perfectly palindromic, even outside the central four base pairs. The two recognition sequence half-sites are shown in bold on their respective sense strands.

[0110] In accordance with the invention, a substantial number of amino acid modifications to the DNA recognition domain of the I-MsoI meganuclease have been identified which, singly or in combination, can result in recombinant meganucleases with specificities altered at individual bases within the DNA recognition sequence half-sites, such that these rationally-designed meganucleases have recognition sequences different from the wild-type enzyme. Amino acid modifications of I-MsoI and the predicted change in recognition sequence half-site specificity are shown in Table 2:

TABLE 2

Position	Favored Sense-Strand Base			
	A	C	G	T
-1	K75* Q77 A49* C49* K79*	D77 E77 K49* R75* K75* R79* K79*	K77 R77 E49* E79*	C77 L77 Q79*
-2	Q75 K81 C47* I47* L47*	E75 D75 R47* K47* K81* R81*	K75 E47* E81*	A75 C75 V75 I75 T75 Q47* Q81*
-3	Q72 C26* L26* V26* A26* I26*	E72 Y72 H26* K26* R26*	R72 K72 Y26* F26*	K72 Y72 H26*
-4	K28 Q83	K28* R28* E83	R83 K83	K28 K83 Q28*
-5	K28 C28* L28* I28*	K28* R28*	R45 E28*	Q28*
-6	I30* V30* S30* L30* Q43	E43 E85 K30* R30*	R43 K43 K85 R85 E30* D30*	K43 I85 V85 L85 Q30*
-7	Q41	E32 E41	R32 R41 K41	K32 M41 L41 I41
-8	Y35 K35	E32	R32 K32 K35 R35	K32 K35
-9	N34 H34	D34 E34 S34	K34 R34 H34	S34 C34 V34 T34 A34

Bold entries are represent wild-type contact residues and do not constitute "modifications" as used herein. An asterisk indicates that the residue contacts the base on the antisense strand.

### 2.2.3 Meganucleases Derived from I-SceI

[0111] In another aspect, the present invention relates to rationally-designed meganucleases which are based upon or derived from the I-SceI meganuclease of *Saccharomyces cerevisiae*. The wild-type amino acid sequence of the I-SceI meganuclease is shown in SEQ ID NO: 9, which corresponds to Genbank Accession # CAA09843. The recognition sequence of the wild-type I-SceI meganuclease from crystal structure PDB # 1R7M is shown below:

Sense	5'	T	T	A	C	C	C	T	G	T	T	A	T	C	C	C	T	A	G	-3'	SEQ ID NO:	10
Antisense	3'	A	A	T	G	G	G	A	C	A	A	T	A	G	G	G	A	T	C	-5'	SEQ ID NO:	11
Position		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18			

Note that the recognition sequence is non-palindromic and there are not four base pairs separating half-sites.

[0112] In accordance with the invention, a substantial number of amino acid modifications to the DNA recognition domain of the I-SceI meganuclease have been identified which, singly or in combination, can result in recombinant meganucleases with specificities altered at individual bases within the DNA recognition sequence, such that these rationally-designed meganucleases have recognition sequences different from the wild-type enzyme. The amino acid modifications of I-SceI and the predicted change in recognition sequence specificity are shown in Table 3:

TABLE 3

Position	Favored Sense-Strand Base			
	A	C	G	T
4	K50	R50* K50* E57	E50* R57 K57	K57 M57 Q50*
5	K48 Q102	R48* K48* E102 E59	E48* K102 R102	Q48* C102 L102 V102
6	K59	R59* K59*	K84 E59*	Q59* Y46
7	C46* L46* V46*	R46* K46* E86	K86 R86 E46*	K68 C88 L86 Q46*
8	K61* S61* V61* A61* L61*	E88 R61* H61*	E61* R88 K88	K88 Q61* H61*
9	T98* C98* V98* L98*	R98* K98*	E98* D98*	Q98*
10	V96* C96* A96*	K96* R96*	D96* E96*	Q96*
11	C90* L90*	K90* R90*	E90*	Q90*
12	Q193	E165 E193 D193	K165 R165	C165 L165 C193 V193 A193 T193 S193
13	C193* L193*	K193* R193* D192	E193* D193* K163 R192	Q193* C163 L163
14	L192* C192*	E161 R192* K192*	K147 K161 R161 R197 D192* E192*	K161 Q192*

15		E151	K151	C151 L151 K151
17	<b>N152*</b> <b>S152*</b> <b>C150*</b> <b>L150*</b> <b>V150*</b> <b>T150*</b>	<b>K152*</b> <b>K150*</b>	<b>N152*</b> <b>S152*</b> <b>D152*</b> <b>D150*</b> <b>E150*</b>	<b>Q152*</b> <b>Q150*</b>
18	<b>K155*</b> <b>C155*</b>	<b>R155*</b> <b>K155*</b>	<b>E155*</b>	<b>H155*</b> <b>Y155*</b>

Bold entries are wild-type contact residues and do not constitute "modifications" as used herein.

An asterisk indicates that the residue contacts the base on the antisense strand.

#### 2.2.4 Meganucleases Derived from I-CeuI

[0113] In another aspect, the present invention relates to rationally-designed meganucleases which are based upon or derived from the I-CeuI meganuclease of *Chlamydomonas eugametos*. The wild-type amino acid sequence of the I-CeuI meganuclease is shown in SEQ ID NO: 12, which corresponds to Genbank Accession # P32761. Two recognition sequence half sites of the wild-type I-CeuI meganuclease from crystal structure PDB # 2EX5 are shown below:

```

Position      -9-8-7-6-5-4-3-2-1
5'-A T A A C G G T C C T A A G G T A G C G A A-3' SEQ ID NO:
13
3'-T A T T G C C A G G A T T C C A T C G C T T-5' SEQ ID NO:
14
Position      -1-2-3-4-5-6-7-8-9

```

Note that the recognition sequence is non-palindromic, even outside the central four base pairs, despite the fact that I-CeuI is a homodimer, due to the natural degeneracy in the I-CeuI recognition interface (Spiegel *et al.* (2006), *Structure* 14:869-80). The two recognition sequence half-sites are shown in bold on their respective sense strands.

[0114] In accordance with the invention, a substantial number of amino acid modifications to the DNA recognition domain of the I-CeuI meganuclease have been identified which, singly or in combination, result in recombinant meganucleases with specificities altered at individual bases within the DNA recognition sequence half-site,



such that these rationally-designed meganucleases can have recognition sequences different from the wild-type enzyme. The amino acid modifications of I-CeuI and the predicted change in recognition sequence specificity are shown in Table 4:

TABLE 4

Position	Favored Sense-Strand Base			
	A	C	G	T
-1	C92* A92* V92*	K116* R116* D116* K92*	E116* E92*	Q116* Q92*
-2	Q117 C90* L90* V90*	E117 D117 R174* K124* K90* R90* K68*	K117 R124 K124 E124* E90* D90*	C117 V117 T117 Q90*
-3	C70* V70* T70* L70* K70*	K70*	E70* E88*	Q70*
-4	Q126 N126 K88* L88* C88* C72* L72* V72*	E126 D126 R88* K88* K72*	R126 K126 E88* D88*	K126 L126 Q88*
-5	C74* L74* V74* T74*	K74*	E74* K128 R128 E128	C128 L128 V128 T128
-6	Q86	D86 E86 R84* K84*	K128 R128 R86 K86 E84*	K86 C86 L86
-7	L76* C76* K76*	R76* K76* H76*	E76* R84	H76* Q76*
-8	Y79 R79 Q76	D79 E79 D76	R79 K79 K76	C79 L79 V79

		<b>E76</b>	<b>R76</b>	<b>L76</b>
<b>-9</b>	<b>Q78</b>	<b>D78</b>	<b>R78</b>	<b>K78</b>
	N78	E78	K78	V78
	H78		H78	L78
	K78			C78
				T78

Bold entries are wild-type contact residues and do not constitute "modifications" as used herein.

An asterisk indicates that the residue contacts the base on the antisense strand.

### 2.2.5 Specifically-Excluded Recombinant Meganucleases

**[0115]** The present invention is not intended to embrace certain recombinant meganucleases which have been described in the prior art, and which have been developed by alternative methods. These excluded meganucleases include those described by Arnould *et al.* (2006), *J. Mol. Biol.* 355: 443-58; Sussman *et al.* (2004), *J. Mol. Biol.* 342: 31-41; Chames *et al.* (2005), *Nucleic Acids Res.* 33: e178; Seligman *et al.* (2002), *Nucleic Acids Res.* 30: 3870-9; and Ashworth *et al.* (2006), *Nature* 441(7093):656-659;

including recombinant meganucleases based on I-CreI with single substitutions selected from C33, R33, A44, H33, K32, F33, R32, A28, A70, E33, V33, A26, and R66. Also excluded are recombinant meganucleases based on I-CreI with three substitutions selected from A68/N70/N75 and D44/D70/N75, or with four substitutions selected from K44/T68/G60/N75 and R44/A68/T70/N75. Lastly, specifically excluded is the recombinant meganuclease based on I-MsoI with the pair of substitutions L28 and R83. These substitutions or combinations of substitutions are referred to herein as the "excluded modifications."

### 2.2.6 Meganucleases with Multiple Changes in the Recognition Sequence Half-Site

**[0116]** In another aspect, the present invention relates to rationally-designed meganucleases which are produced by combining two or more amino acid modifications as described in sections 2.2.1-2.2.4 above, in order to alter half-site preference at two or more positions in a DNA recognition sequence half-site. For example, without limitation,

and as more fully described below, the enzyme DJ1 was derived from I-CreI by incorporating the modifications R30/E38 (which favor C at position -7), R40 (which favors G at position -6), R42 (which favors G at position -5), and N32 (which favors complete degeneracy at position -9). The rationally-designed DJ1 meganuclease invariably recognizes C<sub>-7</sub> G<sub>-6</sub> G<sub>-5</sub> compared to the wild-type preference for A<sub>-7</sub> A<sub>-6</sub> C<sub>-5</sub>, and has increased tolerance for A at position -9.

[0117] The ability to combine residue substitutions that affect different base positions is due in part to the modular nature of the LAGLIDADG meganucleases. A majority of the base contacts in the LAGLIDADG recognition interfaces are made by individual amino acid side chains, and the interface is relatively free of interconnectivity or hydrogen bonding networks between side chains that interact with adjacent bases. This generally allows manipulation of residues that interact with one base position without affecting side chain interactions at adjacent bases. The additive nature of the mutations listed in sections 2.2.1-2.2.4 above is also a direct result of the method used to identify these mutations. The method predicts side chain substitutions that interact directly with a single base. Interconnectivity or hydrogen bonding networks between side chains is generally avoided to maintain the independence of the substitutions within the recognition interface.

[0118] Certain combinations of side chain substitutions are completely or partially incompatible with one another. When an incompatible pair or set of amino acids are incorporated into a rationally-designed meganuclease, the resulting enzyme will have reduced or eliminated catalytic activity. Typically, these incompatibilities are due to steric interference between the side chains of the introduced amino acids and activity can be restored by identifying and removing this interference. Specifically, when two amino acids with large side chains (*e.g.*, amino acids from group 2 or 3) are incorporated at amino acid positions that are adjacent to one another in the meganuclease structure (*e.g.*, positions 32 and 33, 28 and 40, 28 and 42, 42 and 77, or 68 and 77 in the case of meganucleases derived from I-CreI), it is likely that these two amino acids will interfere with one another and reduce enzyme activity. This interference can be eliminated by substituting one or both incompatible amino acids to an amino acid with a smaller side chain (*e.g.*, group 1 or group 2). For example, in rationally-designed meganucleases

derived from I-CreI, K28 interferes with both R40 and R42. To maximize enzyme activity, R40 and R42 can be combined with a serine or aspartic acid at position 28.

[0119] Combinations of amino substitutions, identified as described herein, can be used to rationally alter the specificity of a wild-type meganuclease (or a previously modified meganuclease) from an original recognition sequence to a desired recognition sequence which may be present in a nucleic acid of interest (*e.g.*, a genome). Figure 2A, for example, shows the "sense" strand of the I-CreI meganuclease recognition sequence WT (SEQ ID NO: 4) as well as a number of other sequences for which a rationally-designed meganuclease would be useful. Conserved bases between the WT recognition sequence and the desired recognition sequence are shaded. In accordance with the invention, recombinant meganucleases based on the I-CreI meganuclease can be rationally-designed for each of these desired recognition sequences, as well as any others, by suitable amino acid substitutions as described herein.

### 3. Rationally-Designed Meganucleases with Altered DNA-Binding Affinity

[0120] As described above, the DNA-binding affinity of the recombinant meganucleases of the invention can be modulated by altering certain amino acids that form the contact surface with the phosphodiester backbone of DNA. The contact surface comprises those amino acids in the enzyme with  $\beta$ -carbons less than 9 Å from the DNA backbone, and with side chains oriented toward the DNA, irrespective of whether the residues make contacts with the DNA backbone in the wild-type meganuclease-DNA complex. Because DNA-binding is a necessary precursor to enzyme activity, increases/decreases in DNA-binding affinity have been shown to cause increases/decreases, respectively, in enzyme activity. However, increases/decreases in DNA-binding affinity also have been shown to cause decreases/increases in the meganuclease sequence-specificity. Therefore, both activity and specificity can be modulated by modifying the phosphodiester backbone contacts.

[0121] Specifically, to increase enzyme activity/decrease enzyme specificity:

[0122] (i) Remove electrostatic repulsion between the enzyme and DNA backbone. If an identified amino acid has a negatively-charged side chain (*e.g.*, aspartic acid, glutamic acid) which would be expected to repulse the negatively-charged DNA

backbone, the repulsion can be eliminated by substituting an amino acid with an uncharged or positively-charged side chain, subject to effects of steric interference. An experimentally verified example is the mutation of glutamic acid 80 in I-CreI to glutamine.

[0123] (ii) Introduce electrostatic attraction interaction between the enzyme and the DNA backbone. At any of the positions of the contact surface, the introduction of an amino acid with a positively-charged side chain (*e.g.*, lysine or arginine) is expected to increase binding affinity, subject to effects of steric interference.

[0124] (iii) Introduce a hydrogen-bond between the enzyme and the DNA backbone. If an amino acid of the contact surface does not make a hydrogen bond with the DNA backbone because it lacks an appropriate hydrogen-bonding functionality or has a side chain that is too short, too long, and/or too inflexible to interact with the DNA backbone, a polar amino acid capable of donating a hydrogen bond (*e.g.*, serine, threonine, tyrosine, histidine, glutamine, asparagine, lysine, cysteine, or arginine) with the appropriate length and flexibility can be introduced, subject to effects of steric interference.

[0125] Specifically, to decrease enzyme activity/increase enzyme specificity:

[0126] (i) Introduce electrostatic repulsion between the enzyme and the DNA backbone. At any of the positions of the contact surface, the introduction of an amino acid with a negatively-charged side chain (*e.g.*, glutamic acid, aspartic acid) is expected to decrease binding affinity, subject to effects of steric interference.

[0127] (ii) Remove electrostatic attraction between the enzyme and DNA. If any amino acid of the contact surface has a positively-charged side chain (*e.g.*, lysine or arginine) that interacts with the negatively-charged DNA backbone, this favorable interaction can be eliminated by substituting an amino acid with an uncharged or negatively-charged side chain, subject to effects of steric interference. An experimentally verified example is the mutation of lysine 116 in I-CreI to aspartic acid.

[0128] (iii) Remove a hydrogen-bond between the enzyme and the DNA backbone. If any amino acid of the contact surface makes a hydrogen bond with the DNA backbone, it can be substituted to an amino acid that would not be expected to make

a similar hydrogen bond because its side chain is not appropriately functionalized or it lacks the necessary length/flexibility characteristics.

[0129] For example, in some recombinant meganucleases based on I-CreI, the glutamic acid at position 80 in the I-CreI meganuclease is altered to either a lysine or a glutamine to increase activity. In another embodiment, the tyrosine at position 66 of I-CreI is changed to arginine or lysine, which increases the activity of the meganuclease. In yet another embodiment, enzyme activity is decreased by changing the lysine at position 34 of I-CreI to aspartic acid, changing the tyrosine at position 66 to aspartic acid, and/or changing the lysine at position 116 to aspartic acid.

[0130] The activities of the recombinant meganucleases can be modulated such that the recombinant enzyme has anywhere from no activity to very high activity with respect to a particular recognition sequence. For example, the DJ1 recombinant meganuclease when carrying glutamic acid mutation at position 26 loses activity completely. However, the combination of the glutamic acid substitution at position 26 and a glutamine substitution at position 80 creates a recombinant meganuclease with high specificity and activity toward a guanine at -4 within the recognition sequence half-site (see Figure 1(D)).

[0131] In accordance with the invention, amino acids at various positions in proximity to the phosphodiester DNA backbone can be changed to simultaneously affect both meganuclease activity and specificity. This "tuning" of the enzyme specificity and activity is accomplished by increasing or decreasing the number of contacts made by amino acids with the phosphodiester backbone. A variety of contacts with the phosphodiester backbone can be facilitated by amino acid side chains. In some embodiments, ionic bonds, salt bridges, hydrogen bonds, and steric hindrance affect the association of amino acid side chains with the phosphodiester backbone. For example, for the I-CreI meganuclease, alteration of the lysine at position 116 to an aspartic acid removes a salt bridge between nucleic acid base pairs at positions -8 and -9, reducing the rate of enzyme cleavage but increasing the specificity.

[0132] The residues forming the backbone contact surface of each of the wild-type I-CreI (SEQ ID NO: 1), I-MsoI (SEQ ID NO: 6), I-SceI (SEQ ID NO: 9) and I-CeuI (SEQ ID NO: 12) meganucleases are identified in Table 5 below:

TABLE 5

I-CreI	I-MsoI	I-SceI	I-CeuI
P29, K34, T46, K48, R51, V64, Y66, E80, I81, K82, L112, K116, D137, K139, T140, T143	K36, Q41, R51, N70, I85, G86, S87, T88, H89, Y118, Q122, K123, Q139, K143, R144, E147, S150, N152	N15, N17, L19, K20, K23, K63, L80, S81, H84, L92, N94, N120, K122, K148, Y151, K153, T156, N157, S159, N163, Q165, S166, Y188, K190, I191, K193, N194, K195, Y199, D201, S202, Y222, K223	K21, D25, K28, K31, S68, N70, H94, R112, R114, S117, N120, D128, N129, R130, H172

**[0133]** To increase the affinity of an enzyme and thereby make it more active/less specific:

- (1) Select an amino acid from Table 5 for the corresponding enzyme that is either negatively-charged (D or E), hydrophobic (A, C, F, G, I, L, M, P, V, W, Y), or uncharged/polar (H, N, Q, S, T).
- (2) If the amino acid is negatively-charged or hydrophobic, mutate it to uncharged/polar (less effect) or positively-charged (K or R, more effect).
- (3) If the amino acid is uncharged/polar, mutate it to positively-charged.

**[0134]** To decrease the affinity of an enzyme and thereby make it less active/more specific:

- (1) Select an amino acid from Table 5 for the corresponding enzyme that is either positively-charged (K or R), hydrophobic (A, C, F, G, I, L, M, P, V, W, Y), or uncharged/polar (H, N, Q, S, T).

- (2) If the amino acid is positively-charged, mutate it to uncharged/polar (less effect) or negatively-charged (more effect).
- (3) If the amino acid is hydrophobic or uncharged/polar, mutate it to negatively-charged.

#### 4. Heterodimeric Meganucleases

[0135] In another aspect, the invention provides meganucleases which are heterodimers formed by the association of two monomers, one of which may be a wild-type and one or both of which may be a non-naturally-occurring or recombinant form. For example, wild-type I-CreI meganuclease is normally a homodimer composed of two monomers that each bind to one half-site in the pseudo-palindromic recognition sequence. A heterodimeric recombinant meganuclease can be produced by combining two meganucleases that recognize different half-sites, for example by co-expressing the two meganucleases in a cell or by mixing two meganucleases in solution. The formation of heterodimers can be favored over the formation of homodimers by altering amino acids on each of the two monomers that affect their association into dimers. In particular embodiments, certain amino acids at the interface of the two monomers are altered from negatively-charged amino acids (D or E) to positively-charged amino acids (K or R) on a first monomer and from positively-charged amino acids to negatively-charged amino acids on a second monomer (Table 6). For example, in the case of meganucleases derived from I-CreI, lysines at positions 7 and 57 are mutated to glutamic acids in the first monomer and glutamic acids at positions 8 and 61 are mutated to lysines in the second monomer. The result of this process is a pair of monomers in which the first monomer has an excess of positively-charged residues at the dimer interface and the second monomer has an excess of negatively-charged residues at the dimer interface. The first and second monomer will, therefore, associate preferentially over their identical monomer pairs due to the electrostatic interactions between the altered amino acids at the interface.

TABLE 6

<b>I-CreI: First Monomer Substitutions</b>	<b>I-CreI: Second Monomer Substitutions</b>
--	---



K7 to E7 or D7 K57 to E57 or D57 K96 to E96 or D96	E8 to K8 or R8 E61 to K61 or R61
<b>I-MsoI: First Monomer Substitutions</b>	<b>I-MsoI: Second Monomer Substitutions</b>
R302 to E302 or D302	D20 to K60 or R60 E11 to K11 or R11 Q64 to K64 or R64
<b>I-CeuI: First Monomer Substitutions</b>	<b>I-CeuI: Second Monomer Substitutions</b>
R93 to E93 or D93	E152 to K152 or R152

**[0136]** Alternatively, or in addition, certain amino acids at the interface of the two monomers can be altered to sterically hinder homodimer formation. Specifically, amino acids in the dimer interface of one monomer are substituted with larger or bulkier residues that will sterically prevent the homodimer. Amino acids in the dimer interface of the second monomer optionally can be substituted with smaller residues to compensate for the bulkier residues in the first monomer and remove any clashes in the heterodimer, or can be unmodified.

**[0137]** In another alternative or additional embodiment, an ionic bridge or hydrogen bond can be buried in the hydrophobic core of a heterodimeric interface. Specifically, a hydrophobic residue on one monomer at the core of the interface can be substituted with a positively charged residue. In addition, a hydrophobic residue on the second monomer, that interacts in the wild type homodimer with the hydrophobic residue substituted in the first monomer, can be substituted with a negatively charged residue. Thus, the two substituted residues can form an ionic bridge or hydrogen bond. At the same time, the electrostatic repulsion of an unsatisfied charge buried in a hydrophobic interface should disfavor homodimer formation.

[0138] Finally, as noted above, each monomer of the heterodimer can have different amino acids substituted in the DNA recognition region such that each has a different DNA half-site and the combined dimeric DNA recognition sequence is non-palindromic.

##### 5. Methods of Producing Recombinant Cells and Organisms

[0139] Aspects of the present invention further provide methods for producing recombinant, transgenic or otherwise genetically-modified cells and organisms using rationally-designed meganucleases. Thus, in certain embodiments, recombinant meganucleases are developed to specifically cause a double-stranded break at a single site or at relatively few sites in the genomic DNA of a cell or an organism to allow for precise insertion(s) of a sequence of interest by homologous recombination. In other embodiments, recombinant meganucleases are developed to specifically cause a double-stranded break at a single site or at relatively few sites in the genomic DNA of a cell or an organism to either (a) allow for rare insertion(s) of a sequence of interest by non-homologous end-joining or (b) allow for the disruption of the target sequence by non-homologous end-joining. As used herein with respect to homologous recombination or non-homologous end-joining of sequences of interest, the term "insertion" means the ligation of a sequence of interest into a chromosome such that the sequence of interest is integrated into the chromosome. In the case of homologous recombination, an inserted sequence can replace an endogenous sequence, such that the original DNA is replaced by exogenous DNA of equal length, but with an altered nucleotide sequence. Alternatively, an inserted sequence can include more or fewer bases than the sequence it replaces.

[0140] Therefore, in accordance with this aspect of the invention, the recombinant organisms include, but are not limited to, monocot plant species such as rice, wheat, corn (maize) and rye, and dicot species such as legumes (*e.g.*, kidney beans, soybeans, lentils, peanuts, peas), alfalfa, clover, tobacco and *Arabidopsis* species. In addition, the recombinant organisms can include, but are not limited to, animals such as humans and non-human primates, horses, cows, goats, pigs, sheep, dogs, cats, guinea pigs, rats, mice, lizards, fish and insects such as *Drosophila* species. In other embodiments, the organism is a fungus such as a *Candida*, *Neurospora* or *Saccharomyces* species.

[0141] In some embodiments, the methods of the invention involve the introduction of a sequence of interest into a cell such as a germ cell or stem cell that can become a mature recombinant organism or allow the resultant genetically-modified organism to give rise to progeny carrying the inserted sequence of interest in its genome.

[0142] Meganuclease proteins can be delivered into cells to cleave genomic DNA, which allows for homologous recombination or non-homologous end-joining at the cleavage site with a sequence of interest, by a variety of different mechanisms known in the art. For example, the recombinant meganuclease protein can be introduced into a cell by techniques including, but not limited to, microinjection or liposome transfections (see, e.g., Lipofectamine™, Invitrogen Corp., Carlsbad, CA). The liposome formulation can be used to facilitate lipid bilayer fusion with a target cell, thereby allowing the contents of the liposome or proteins associated with its surface to be brought into the cell.

Alternatively, the enzyme can be fused to an appropriate uptake peptide such as that from the HIV TAT protein to direct cellular uptake (see, e.g., Hudecz *et al.* (2005), *Med. Res. Rev.* 25: 679-736).

[0143] Alternatively, gene sequences encoding the meganuclease protein are inserted into a vector and transfected into a eukaryotic cell using techniques known in the art (see, e.g., Ausubel *et al.*, Current Protocols in Molecular Biology, Wiley 1999). The sequence of interest can be introduced in the same vector, a different vector, or by other means known in the art.

[0144] Non-limiting examples of vectors for DNA transfection include virus vectors, plasmids, cosmids, and YAC vectors. Transfection of DNA sequences can be accomplished by a variety of methods known to those of skill in the art. For instance, liposomes and immunoliposomes are used to deliver DNA sequences to cells (see, e.g., Lasic *et al.* (1995), *Science* 267: 1275-76). In addition, viruses can be utilized to introduce vectors into cells (see, e.g., U.S. Pat. No. 7,037,492). Alternatively, transfection strategies can be utilized such that the vectors are introduced as naked DNA (see, e.g., Rui *et al.* (2002), *Life Sci.* 71(15): 1771-8).

[0145] General methods for delivering nucleic acids into cells include: (1) chemical methods (Graham *et al.* (1973), *Virology* 54(2):536-539; Zatloukal *et al.* (1992), *Ann. N.Y. Acad. Sci.*, 660:136-153; (2) physical methods such as microinjection

(Capecchi (1980), *Cell* 22(2):479-488, electroporation (Wong *et al.* (1982), *Biochim. Biophys. Res. Commun.* 107(2):584-587; Fromm *et al.* (1985), *Proc. Nat'l Acad. Sci. USA* 82(17):5824-5828; U.S. Pat. No. 5,384,253) and ballistic injection (Johnston *et al.* (1994), *Methods Cell. Biol.* 43(A): 353-365; Fynan *et al.* (1993), *Proc. Nat'l Acad. Sci. USA* 90(24): 11478-11482); (3) viral vectors (Clapp (1993), *Clin. Perinatol.* 20(1): 155-168; Lu *et al.* (1993), *J. Exp. Med.* 178(6):2089-2096; Eglitis *et al.* (1988), *Adv. Exp. Med. Biol.* 241:19-27; Eglitis *et al.* (1988), *Biotechniques* 6(7):608-614); and (4) receptor-mediated mechanisms (Curiel *et al.* (1991), *Proc. Nat'l Acad. Sci. USA* 88(19):8850-8854; Curiel *et al.* (1992), *Hum. Gen. Ther.* 3(2):147-154; Wagner *et al.* (1992), *Proc. Nat'l Acad. Sci. USA* 89 (13):6099-6103).

[0146] In certain embodiments, a genetically-modified plant is produced, which contains the sequence of interest inserted into the genome. In certain embodiments, the genetically-modified plant is produced by transfecting the plant cell with DNA sequences corresponding to the recombinant meganuclease and the sequence of interest, which may or may not be flanked by the meganuclease recognition sequences and/or sequences substantially identical to the target sequence. In other embodiments, the genetically-modified plant is produced by transfecting the plant cell with DNA sequences corresponding to the recombinant meganuclease only, such that cleavage promotes non-homologous end-joining and disrupts the target sequence containing the recognition sequence. In such embodiments, the meganuclease sequences are under the control of regulatory sequences that allow for expression of the meganuclease in the host plant cells. These regulatory sequences include, but are not limited to, constitutive plant promoters such as the NOS promoter, chemically-inducible gene promoters such as the dexamethasone-inducible promoter (see, *e.g.*, Gremillon *et al.* (2004), *Plant J.* 37:218-228), and plant tissue specific promoters such as the LGC1 promoter (see, *e.g.*, Singh *et al.* (2003), *FEBS Lett.* 542:47-52).

[0147] Suitable methods for introducing DNA into plant cells include virtually any method by which DNA can be introduced into a cell, including but not limited to *Agrobacterium* infection, PEG-mediated transformation of protoplasts (Omirulleh *et al.* (1993), *Plant Molecular Biology*, 21:415-428), desiccation/inhibition-mediated DNA

uptake, electroporation, agitation with silicon carbide fibers, ballistic injection or microprojectile bombardment, and the like.

[0148] In other embodiments, a genetically-modified animal is produced using a recombinant meganuclease. As with plant cells, the nucleic acid sequences can be introduced into a germ cell or a cell that will eventually become a transgenic organism. In some embodiments, the cell is a fertilized egg, and exogenous DNA molecules can be injected into the pro-nucleus of the fertilized egg. The micro-injected eggs are then transferred into the oviducts of pseudopregnant foster mothers and allowed to develop. The recombinant meganuclease is expressed in the fertilized egg (*e.g.*, under the control of a constitutive promoter, such as 3-phosphoglycerate kinase), and facilitates homologous recombination of the sequence of interest into one or a few discrete sites in the genome. Alternatively, the genetically-modified animals can be obtained by utilizing recombinant embryonic stem ("ES") cells for the generation of the transgenics, as described by Gossler *et al.* (1986), *Proc. Natl. Acad. Sci. USA* 83:9065-9069.

[0149] In certain embodiments, a recombinant mammalian expression vector is capable of directing tissue-specific expression of the nucleic acid preferentially in a particular cell type. Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987), *Genes Dev.* 1: 268-277), lymphoid-specific promoters (Calame and Eaton (1988), *Adv. Immunol.* 43: 235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989), *EMBO J.* 8: 729-733) and immunoglobulins (Banerji *et al.* (1983), *Cell* 33: 729-740; Queen and Baltimore (1983), *Cell* 33: 741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle (1989), *Proc. Natl. Acad. Sci. USA* 86: 5473-5477), pancreas-specific promoters (Edlund *et al.* (1985), *Science* 230: 912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Pat. No. 4,873,316 and European Pat. Pub. EP 0 264 166). Developmentally-regulated promoters are also encompassed, *e.g.*, the murine hox promoters (Kessel and Gruss (1990), *Science* 249: 374-379) and the  $\alpha$ -fetoprotein promoter (Campes and Tilghman (1989), *Genes Dev.* 3: 537-546).

[0150] In certain embodiments, a rationally-designed meganuclease may be tagged with a peptide epitope (*e.g.*, an HA, FLAG, or Myc epitope) to monitor expression

levels or localization. In some embodiments, the meganuclease may be fused to a sub-cellular localization signal such as a nuclear-localization signal (*e.g.*, the nuclear localization signal from SV40) or chloroplast or mitochondrial localization signals. In other embodiments, the meganuclease may be fused to a nuclear export signal to localize it to the cytoplasm. The meganuclease may also be fused to an unrelated protein or protein domain such as a protein that stimulates DNA-repair or homologous recombination (*e.g.*, *recA*, *RAD51*, *RAD52*, *RAD54*, *RAD57* or *BRCA2*).

#### 6. Methods for Gene Therapy

**[0151]** Aspects of the invention allow for the use of recombinant meganuclease for gene therapy. As used herein, “gene therapy” means therapeutic treatments that comprise introducing into a patient a functional copy of at least one gene, or gene regulatory sequence such as a promoter, enhancer, or silencer to replace a gene or gene regulatory region that is defective in its structure and/or function. The term “gene therapy” can also refer to modifications made to a deleterious gene or regulatory element (*e.g.*, oncogenes) that reduce or eliminate expression of the gene. Gene therapy can be performed to treat congenital conditions, conditions resulting from mutations or damage to specific genetic loci over the life of the patient, or conditions resulting from infectious organisms.

**[0152]** In some aspects of the invention, dysfunctional genes are replaced or disabled by the insertion of exogenous nucleic acid sequences into a region of the genome affecting gene expression. In certain embodiments, the recombinant meganuclease is targeted to a particular sequence in the region of the genome to be modified so as to alleviate the condition. The sequence can be a region within an exon, intron, promoter, or other regulatory region that is causing dysfunctional expression of the gene. As used herein, the term “dysfunctional expression” means aberrant expression of a gene product either by the cell producing too little of the gene product, too much of the gene product, or producing a gene product that has a different function such as lacking the necessary function or having more than the necessary function.

**[0153]** Exogenous nucleic acid sequences inserted into the modified region can be used to provide “repaired” sequences that normalize the gene. Gene repair can be

accomplished by the introduction of proper gene sequences into the gene allowing for proper function to be reestablished. In these embodiments, the nucleic acid sequence to be inserted can be the entire coding sequence for a protein or, in certain embodiments, a fragment of the gene comprising only the region to be repaired. In other embodiments the nucleic acid sequence to be inserted comprises a promoter sequence or other regulatory elements such that mutations causing abnormal expression or regulation are repaired. In other embodiments, the nucleic acid sequence to be inserted contains the appropriate translation stop codon lacking in a mutated gene. The nucleic acid sequence can also have sequences for stopping transcription in a recombinant gene lacking appropriate transcriptional stop signals.

[0154] Alternatively, the nucleic acid sequences can eliminate gene function altogether by disrupting the regulatory sequence of the gene or providing a silencer to eliminate gene function. In some embodiments, the exogenous nucleic acid sequence provides a translation stop codon to prevent expression of the gene product. In other embodiments, the exogenous nucleic acid sequences provide transcription stop element to prevent expression of a full length RNA molecule. In still other embodiments, gene function is disrupted directly by the meganuclease by introducing base insertions, base deletions, and/or frameshift mutations through non-homologous end-joining.

[0155] In many instances, it is desirable to direct the proper genetic sequences to a target cell or population of cells that is the cause of the disease condition. Such targeting of therapeutics prevents healthy cells from being targeted by the therapeutics. This increases the efficacy of the treatment, while decreasing the potentially adverse effects that the treatment could have on healthy cells.

[0156] Delivery of recombinant meganuclease genes and the sequence of interest to be inserted into the genome to the cells of interest can be accomplished by a variety of mechanisms. In some embodiments, the nucleic acids are delivered to the cells by way of viruses with particular viral genes inactivated to prevent reproduction of the virus. Thus, a virus can be altered so that it is capable only of delivery and maintenance within a target cell, but does not retain the ability to replicate within the target cell or tissue. One or more DNA sequences can be introduced to the altered viral genome, so as to produce a viral genome that acts like a vector, and may or may not be inserted into a host genome

and subsequently expressed. More specifically, certain embodiments include employing a retroviral vector such as, but not limited to, the MFG or pLJ vectors. An MFG vector is a simplified Moloney murine leukemia virus vector (MoMLV) in which the DNA sequences encoding the pol and env proteins have been deleted to render it replication defective. A pLJ retroviral vector is also a form of the MoMLV (see, *e.g.*, Korman *et al.* (1987), *Proc. Nat'l Acad. Sci.*, 84:2150-2154). In other embodiments, a recombinant adenovirus or adeno-associated virus can be used as a delivery vector.

[0157] In other embodiments, the delivery of recombinant meganuclease protein and/or recombinant meganuclease gene sequences to a target cell is accomplished by the use of liposomes. The production of liposomes containing nucleic acid and/or protein cargo is known in the art (see, *e.g.*, Lasic *et al.* (1995), *Science* 267: 1275-76). Immunoliposomes incorporate antibodies against cell-associated antigens into liposomes, and can delivery DNA sequences for the meganuclease or the meganuclease itself to specific cell types (see, *e.g.*, Lasic *et al.* (1995), *Science* 267: 1275-76; Young *et al.* (2005), *J. Calif. Dent. Assoc.* 33(12): 967-71; Pfeiffer *et al.* (2006), *J. Vasc. Surg.* 43(5):1021-7). Methods for producing and using liposome formulations are well known in the art, (see, *e.g.*, U.S. Pat. No. 6,316,024, U.S. Pat. No. 6,379,699, U.S. Pat. No. 6,387,397, U.S. Pat. No. 6,511,676 and U.S. Pat. No. 6,593,308, and references cited therein). In some embodiments, liposomes are used to deliver the sequences of interest as well as the recombinant meganuclease protein or recombinant meganuclease gene sequences.

#### 7. Methods for Treating Pathogen Infection.

[0158] Aspects of the invention also provide methods of treating infection by a pathogen. Pathogenic organisms include viruses such as, but not limited to, herpes simplex virus 1, herpes simplex virus 2, human immunodeficiency virus 1, human immunodeficiency virus 2, variola virus, polio virus, Epstein-Barr virus, and human papilloma virus and bacterial organisms such as, but not limited to, *Bacillus anthracis*, *Haemophilus* species, *Pneumococcus* species, *Staphylococcus aureus*, *Streptococcus* species, methicillin-resistant *Staphylococcus aureus*, and *Mycoplasma tuberculosis*.



Pathogenic organisms also include fungal organisms such as, but not limited to, *Candida*, *Blastomyces*, *Cryptococcus*, and *Histoplasma* species.

[0159] In some embodiments, a rationally-designed meganuclease can be targeted to a recognition sequence within the pathogen genome, *e.g.*, to a gene or regulatory element that is essential for growth, reproduction, or toxicity of the pathogen. In certain embodiments, the recognition sequence may be in a bacterial plasmid. Meganuclease-mediated cleavage of a recognition sequence in a pathogen genome can stimulate mutation within a targeted, essential gene in the form of an insertion, deletion or frameshift, by stimulating non-homologous end-joining. Alternatively, cleavage of a bacterial plasmid can result in loss of the plasmid along with any genes encoded on it, such as toxin genes (*e.g.*, *B. anthracis* Lethal Factor gene) or antibiotic resistance genes. As noted above, the meganuclease may be delivered to the infected patient, animal, or plant in either protein or nucleic acid form using techniques that are common in the art. In certain embodiments, the meganuclease gene may be incorporated into a bacteriophage genome for delivery to pathogenic bacteria.

[0160] Aspects of the invention also provide therapeutics for the treatment of certain forms of cancer. Because human viruses are often associated with tumor formation (*e.g.*, Epstein-Barr Virus and nasopharyngeal carcinomas; Human Papilloma Virus and cervical cancer) inactivation of these viral pathogens may prevent cancer development or progression. Alternatively, double-stranded breaks targeted to the genomes of these tumor-associated viruses using rationally-designed meganucleases may be used to trigger apoptosis through the DNA damage response pathway. In this manner, it may be possible to selectively induce apoptosis in tumor cells harboring the viral genome.

#### 8. Methods for Genotyping and Pathogen Identification

[0161] Aspects of the invention also provide tools for *in vitro* molecular biology research and development. It is common in the art to use site-specific endonucleases (*e.g.*, restriction enzymes) for the isolation, cloning, and manipulation of nucleic acids such as plasmids, PCR products, BAC sequences, YAC sequences, viruses, and genomic sequences from eukaryotic and prokaryotic organisms (see, *e.g.*, Ausubel *et al.*, Current

Protocols in Molecular Biology, Wiley 1999). Thus, in some embodiments, a rationally-designed meganuclease may be used to manipulate nucleic acid sequences *in vitro*. For example, rationally-designed meganucleases recognizing a pair of recognition sequences within the same DNA molecule can be used to isolate the intervening DNA segment for subsequent manipulation such as ligation into a bacterial plasmid, BAC, or YAC.

[0162] In another aspect, this invention provides tools for the identification of pathogenic genes and organisms. In one embodiment, rationally-designed meganucleases can be used to cleave recognition sites corresponding to polymorphic genetic regions correlated to disease to distinguish disease-causing alleles from healthy alleles (*e.g.*, a rationally-designed meganuclease which recognizes the  $\Delta F$ -508 allele of the human CFTR gene, see example 4). In this embodiment, DNA sequences isolated from a human patient or other organism are digested with a rationally-designed meganuclease, possibly in conjunction with additional site-specific nucleases, and the resulting DNA fragment pattern is analyzed by gel electrophoresis, capillary electrophoresis, mass spectrometry, or other methods known in the art. This fragmentation pattern and, specifically, the presence or absence of cleavage by the rationally-designed meganuclease, indicates the genotype of the organism by revealing whether or not the recognition sequence is present in the genome. In another embodiment, a rationally-designed meganuclease is targeted to a polymorphic region in the genome of a pathogenic virus, fungus, or bacterium and used to identify the organism. In this embodiment, the rationally-designed meganuclease cleaves a recognition sequence that is unique to the pathogen (*e.g.*, the spacer region between the 16S and 23S rRNA genes in a bacterium; see, *e.g.*, van der Giessen *et al.* (1994), *Microbiology* 140:1103-1108) and can be used to distinguish the pathogen from other closely-related organisms following endonuclease digest of the genome and subsequent analysis of the fragmentation pattern by electrophoresis, mass spectrometry, or other methods known in the art.

#### 9. Methods for the Production of Custom DNA-binding Domains.

[0163] In another aspect, the invention provides rationally-designed DNA-binding proteins that lack endonuclease cleavage activity. The catalytic activity of a rationally-designed meganuclease can be eliminated by mutating amino acids involved in catalysis

(*e.g.*, the mutation of Q47 to E in I-CreI, see Chevalier *et al.* (2001), *Biochemistry*. 43:14015-14026); the mutation of D44 or D145 to N in I-SceI; the mutation of E66 to Q in I-CeuI; the mutation of D22 to N in I-MsoI). The inactivated meganuclease can then be fused to an effector domain from another protein including, but not limited to, a transcription activator (*e.g.*, the GAL4 transactivation domain or the VP16 transactivation domain), a transcription repressor (*e.g.*, the KRAB domain from the Kruppel protein), a DNA methylase domain (*e.g.*, M.CviPI or M.SssI), or a histone acetyltransferase domain (*e.g.*, HDAC1 or HDAC2). Chimeric proteins consisting of an engineered DNA-binding domain, most notably an engineered zinc finger domain, and an effector domain are known in the art (see, *e.g.*, Papworth *et al.* (2006), *Gene* 366:27-38).

#### EXAMPLES

[0164] This invention is further illustrated by the following examples, which should not be construed as limiting. Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific substances and procedures described herein. Such equivalents are intended to be encompassed in the scope of the claims that follow the examples below. Examples 1-4 below refer specifically to rationally-designed meganucleases based on I-CreI, but rationally-designed meganucleases based on I-SceI, I-MsoI, I-CeuI, and other LAGLIDADG meganucleases can be similarly produced and used, as described herein.

#### EXAMPLE 1

##### Rational Design of Meganucleases Recognizing the HIV-1 TAT Gene

##### 1. Meganuclease Design.

[0165] A pair of meganucleases were designed to recognize and cleave the DNA site 5'-GAAGAGCTCATCAGAACAGTCA-3' (SEQ ID NO: 15) found in the HIV-1 TAT Gene. In accordance with Table 1, two meganucleases, TAT1 and TAT2, were designed to bind the half-sites 5'-GAAGAGCTC-3' (SEQ ID NO: 16) and 5'-TGACTGTTC-3' (SEQ ID NO: 17), respectively, using the following base contacts (non-WT contacts are in bold):

## TAT1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>G</u>	<u>A</u>	<u>A</u>	<u>G</u>	<u>A</u>	<u>G</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>Y33</u>	<u>N30/</u> <u>Q38</u>	<u>R40</u>	<u>K28</u>	<u>S26/</u> <u>R77</u>	<u>K24/</u> <u>Y68</u>	<u>Q44</u>	<u>R70</u>

## TAT2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>G</u>	<u>A</u>	<u>C</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>C32</u>	<u>R33</u>	<u>N30/</u> <u>Q38</u>	<u>R28/</u> <u>E40</u>	<u>M66</u>	<u>S26/</u> <u>R77</u>	<u>Y68</u>	<u>Q44</u>	<u>R70</u>

[0166] The two enzymes were cloned, expressed in *E. coli*, and assayed for enzyme activity against the corresponding DNA recognition sequence as described below. In both cases, the rationally-designed meganucleases were found to be inactive. A second generation of each was then produced in which E80 was mutated to Q to improve contacts with the DNA backbone. The second generation TAT2 enzyme was found to be active against its intended recognition sequence while the second generation TAT1 enzyme remained inactive. Visual inspection of the wild-type I-CreI co-crystal structure suggested that TAT1 was inactive due to a steric clash between R40 and K28. To alleviate this clash, TAT1 variants were produced in which K28 was mutated to an amino acid with a smaller side chain (A, S, T, or C) while maintaining the Q80 mutation. When these enzymes were produced in *E. coli* and assayed, the TAT1 variants with S28 and T28 were both found to be active against the intended recognition sequence while maintaining the desired base preference at position -7.

## 2. Construction of Recombinant Meganucleases.

[0167] Mutations for the redesigned I-CreI enzymes were introduced using mutagenic primers in an overlapping PCR strategy. Recombinant DNA fragments of I-CreI generated in a primary PCR were joined in a secondary PCR to produce full-length recombinant nucleic acids. All recombinant I-CreI constructs were cloned into pET21a vectors with a six histidine tag fused at the 3' end of the gene for purification (Novagen

Corp., San Diego, CA). All nucleic acid sequences were confirmed using Sanger Dideoxynucleotide sequencing (see Sanger *et al.* (1977), *Proc. Natl. Acad. Sci. USA.* 74(12): 5463-7).

[0168] Wild-type I-CreI and all engineered meganucleases were expressed and purified using the following method. The constructs cloned into a pET21a vector were transformed into chemically competent BL21 (DE3) pLysS, and plated on standard 2xYT plates containing 200 µg/ml carbanicillin. Following overnight growth, transformed bacterial colonics were scraped from the plates and used to inoculate 50 ml of 2XYT broth. Cells were grown at 37°C with shaking until they reached an optical density of 0.9 at a wavelength of 600 nm. The growth temperature was then reduced from 37°C to 22°C. Protein expression was induced by the addition of 1mM IPTG, and the cells were incubated with agitation for two and a half hours. Cells were then pelleted by centrifugation for 10 min. at 6000 xg. Pellets were resuspended in 1ml binding buffer (20 mM Tris-HCL, pH 8.0, 500 mM NaCl, 10 mM imidazole) by vortexing. The cells were then disrupted with 12 pulses of sonication at 50% power and the cell debris was pelleted by centrifugation for 15 min. at 14,000 xg. Cell supernatants were diluted in 4 ml binding buffer and loaded onto a 200 µl nickel-charged metal-chelating Sepharose<sup>TM</sup> column (Pharmacia).

[0169] The column was subsequently washed with 4 ml wash buffer (20 mM Tris-HCl, pH 8.0, 500 mM NaCl, 60 mM imidazole) and with 0.2 ml elution buffer (20 mM Tris-HCl, pH 8.0, 500 mM NaCl, 400 mM imidazole). Meganuclease enzymes were eluted with an additional 0.6 ml of elution buffer and concentrated to 50-130 µl using Vivospin disposable concentrators (ISC, Inc., Kaysville, UT). The enzymes were exchanged into SA buffer (25 mM Tris-HCL, pH 8.0, 100 mM NaCl, 5 mM MgCl<sub>2</sub>, 5mM EDTA) for assays and storage using Zeba spin desalting columns (Pierce Biotechnology, Inc., Rockford, IL). The enzyme concentration was determined by absorbance at 280 nm using an extinction coefficient of 23,590 M<sup>-1</sup>cm<sup>-1</sup>. Purity and molecular weight of the enzymes was then confirmed by MALDI-TOF mass spectrometry.

[0170] Heterodimeric enzymes were produced either by purifying the two proteins independently, and mixing them *in vitro* or by constructing an artificial operon for tandem expression of the two proteins in *E. coli*. In the former case, the purified

meganucleases were mixed 1:1 in solution and pre-incubated at 42°C for 20 minutes prior to the addition of DNA substrate. In the latter case, the two genes were cloned sequentially into the pET-21a expression vector using *NdeI/EcoRI* and *EcoRI/HindIII*. The first gene in the operon ends with two stop codons to prevent read-through errors during transcription. A 12-base pair nucleic acid spacer and a Shine-Dalgarno sequence from the pET21 vector separated the first and second genes in the artificial operon.

### 3. Cleavage Assays.

[0171] All enzymes purified as described above were assayed for activity by incubation with linear, double-stranded DNA substrates containing the meganuclease recognition sequence. Synthetic oligonucleotides corresponding to both sense and antisense strands of the recognition sequence were annealed and were cloned into the *SmaI* site of the pUC19 plasmid by blunt-end ligation. The sequences of the cloned binding sites were confirmed by Sanger dideoxynucleotide sequencing. All plasmid substrates were linearized with *XmnI*, *ScaI* or *BpmI* concurrently with the meganuclease digest. The enzyme digests contained 5 µl 0.05 µM DNA substrate, 2.5 µl 5 µM recombinant I-CreI meganuclease, 9.5 µl SA buffer, and 0.5 µl *XmnI*, *ScaI*, or *BpmI*. Digests were incubated at either 37°C, or 42°C for certain meganuclease enzymes, for four hours. Digests were stopped by adding 0.3 mg/ml Proteinase K and 0.5% SDS, and incubated for one hour at 37°C. Digests were analyzed on 1.5% agarose and visualized by ethidium bromide staining.

[0172] To evaluate meganuclease half-site preference, rationally-designed meganucleases were incubated with a set of DNA substrates corresponding to a perfect palindrome of the intended half-site as well as each of the 27 possible single-base-pair substitutions in the half-site. In this manner, it was possible to determine how tolerant each enzyme is to deviations from its intended half-site.

### 4. Recognition Sequence-Specificity.

[0173] Purified recombinant TAT1 and TAT2 meganucleases recognized DNA sequences that were distinct from the wild-type meganuclease recognition sequence (Fig. 2(B)). The wild-type I-CreI meganuclease cleaves the WT recognition sequence, but cuts

neither the intended sequence for TAT1 nor the intended sequence for TAT2. TAT1 and TAT2, likewise, cut their intended recognition sequences but not the wild-type sequence. The meganucleases were then evaluated for half-site preference and overall specificity (Fig 3). Wild-type I-CreI was found to be highly tolerant of single-base-pair substitutions in its natural half-site. In contrast, TAT1 and TAT2 were found to be highly-specific and completely intolerant of base substitutions at positions -1, -2, -3, -6, and -8 in the case of TAT1, and positions -1, -2, and -6 in the case of TAT2.

## EXAMPLE 2

### Rational Design of Meganucleases with Altered DNA-Binding Affinity

#### 1. Meganucleases with increased affinity and increased activity.

[0174] The meganucleases CCR1 and BRP2 were designed to cleave the half-sites 5'-AACCCTCTC-3' (SEQ ID NO: 18) and 5'-CTCCGGGTC-3' (SEQ ID NO: 19), respectively. These enzymes were produced in accordance with Table 1 as in Example 1:

#### CCR1:

Position	-9	-8	-7	-6	-5	-4	-3	-2	-1
Base	A	A	C	C	C	T	C	T	C
Contact Residues	N32	Y33	R30/ E38	R28/ E40	E42	Q26	K24/ Y68	Q44	R70

#### BRP2:

Position	-9	-8	-7	-6	-5	-4	-3	-2	-1
Base	C	T	C	C	G	G	G	T	C
Contact Residues	S32	C33	R30/ E38	R28/ E40	R42	S26/ R77	R68	Q44	R70

[0175] Both enzymes were expressed in *E. coli*, purified, and assayed as in Example 1. Both first generation enzymes were found to cleave their intended recognition sequences with rates that were considerably below that of wild-type I-CreI with its natural recognition sequence. To alleviate this loss in activity, the DNA-binding affinity of CCR1 and BRP2 was increased by mutating E80 to Q in both enzymes. These

second-generation versions of CCR1 and BRP2 were found to cleave their intended recognition sequences with substantially increased catalytic rates.

2. Meganucleases with decreased DNA-binding affinity and decreased activity but increased specificity.

[0176] Wild-type I-CreI was found to be highly-tolerant of substitutions to its half-site (Fig. 3(A)). In an effort to make the enzyme more specific, the lysine at position 116 of the enzyme, which normally makes a salt-bridge with a phosphate in the DNA backbone, was mutated to aspartic acid to reduce DNA-binding affinity. This rationally-designed enzyme was found to cleave the wild-type recognition sequence with substantially reduced activity but the recombinant enzyme was considerably more specific than wild-type. The half-site preference of the K116D variant was evaluated as in Example 1 and the enzyme was found to be entirely intolerant of deviation from its natural half-site at positions -1, -2, and -3, and displayed at least partial base preference at the remaining 6 positions in the half-site (Fig. 3(B)).

### EXAMPLE 3

#### Rationally-Designed Meganuclease Heterodimers

1. Cleavage of non-palindromic DNA sites by meganuclease heterodimers formed in solution.

[0177] Two meganucleases, LAM1 and LAM2, were designed to cleave the half-sites 5'-TGCGGTGTC-3' (SEQ ID NO: 20) and 5'-CAGGCTGTC-3' (SEQ ID NO: 21), respectively. The heterodimer of these two enzymes was expected to recognize the DNA sequence 5'-TGCGGTGTCCGGCGACAGCCTG-3' (SEQ ID NO: 22) found in the bacteriophage  $\lambda$  p05 gene.



## LAM1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>G</u>	<u>C</u>	<u>G</u>	<u>G</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>C32</u>	<u>R33</u>	<u>R30/ E38</u>	<u>D28/ R40</u>	<u>R42</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

## LAM2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>A</u>	<u>G</u>	<u>G</u>	<u>C</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>Y33</u>	<u>E30/ R38</u>	<u>R40</u>	<u>K28/ E42</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

[0178] LAM1 and LAM 2 were cloned, expressed in *E. coli*, and purified individually as described in Example 1. The two enzymes were then mixed 1:1 and incubated at 42°C for 20 minutes to allow them to exchange subunits and re-equilibrate. The resulting enzyme solution, expected to be a mixture of LAM1 homodimer, LAM2 homodimer, and LAM1/LAM2 heterodimer, was incubated with three different recognition sequences corresponding to the perfect palindrome of the LAM1 half-site, the perfect palindrome of the LAM2 half-site, and the non-palindromic hybrid site found in the bacteriophage  $\lambda$  genome. The purified LAM1 enzyme alone cuts the LAM1 palindromic site, but neither the LAM2 palindromic site, nor the LAM1/LAM2 hybrid site. Likewise, the purified LAM2 enzyme alone cuts the LAM2 palindromic site but neither the LAM1 palindromic site nor the LAM1/LAM2 hybrid site. The 1:1 mixture of LAM1 and LAM2, however, cleaves all three DNA sites. Cleavage of the LAM1/LAM2 hybrid site indicates that two distinct redesigned meganucleases can be mixed in solution to form a heterodimeric enzyme capable of cleaving a non-palindromic DNA site.

2. Cleavage of non-palindromic DNA sites by meganuclease heterodimers formed by co-expression.

[0179] Genes encoding the LAM1 and LAM2 enzymes described above were arranged into an operon for simultaneous expression in *E. coli* as described in Example 1. The co-expressed enzymes were purified as in Example 1 and the enzyme mixture

incubated with the three potential recognition sequences described above. The co-expressed enzyme mixture was found to cleave all three sites, including the LAM1/LAM2 hybrid site, indicating that two distinct rationally-designed meganucleases can be co-expressed to form a heterodimeric enzyme capable of cleaving a non-palindromic DNA site.

3. Preferential cleavage of non-palindromic DNA sites by meganuclease heterodimers with modified protein-protein interfaces.

[0180] For applications requiring the cleavage of non-palindromic DNA sites, it is desirable to promote the formation of enzyme heterodimers while minimizing the formation of homodimers that recognize and cleave different (palindromic) DNA sites. To this end, variants of the LAM1 enzyme were produced in which lysines at positions 7, 57, and 96 were changed to glutamic acids. This enzyme was then co-expressed and purified as in above with a variant of LAM2 in which glutamic acids at positions 8 and 61 were changed to lysine. In this case, formation of the LAM1 homodimer was expected to be reduced due to electrostatic repulsion between E7, E57, and E96 in one monomer and E8 and E61 in the other monomer. Likewise, formation of the LAM2 homodimer was expected to be reduced due to electrostatic repulsion between K7, K57, and K96 on one monomer and K8 and K61 on the other monomer. Conversely, the LAM1/LAM2 heterodimer was expected to be favored due to electrostatic attraction between E7, E57, and E96 in LAM1 and K8 and K61 in LAM2. When the two meganucleases with modified interfaces were co-expressed and assayed as described above, the LAM1/LAM2 hybrid site was found to be cleaved preferentially over the two palindromic sites, indicating that substitutions in the meganuclease protein-protein interface can drive the preferential formation of heterodimers.

**EXAMPLE 4**

Additional Meganuclease Heterodimers Which Cleave Physiologic DNA Sequences

1. Meganuclease heterodimers which cleave DNA sequences relevant to gene therapy.

[0181] A rationally-designed meganuclease heterodimer (ACH1/ACH2) can be produced that cleaves the sequence 5'-CTGGGAGTCTCAGGACAGCCTG-3' (SEQ ID NO: 23) in the human FGFR3 gene, mutations in which cause achondroplasia. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

ACH1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>T</u>	<u>G</u>	<u>G</u>	<u>G</u>	<u>A</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>D32</u>	<u>C33</u>	<u>E30/</u> <u>R38</u>	<u>R40/</u> <u>D28</u>	<u>R42</u>	<u>A26/</u> <u>Q77</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

ACH2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>A</u>	<u>G</u>	<u>G</u>	<u>C</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>D32</u>	<u>Y33</u>	<u>E30/</u> <u>R38</u>	<u>R40</u>	<u>K28/</u> <u>E42</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

[0182] A rationally-designed meganuclease heterodimer (HGH1/HGH2) can be produced that cleaves the sequence 5'-CCAGGTGTCTCTGGACTCCTCC-3' (SEQ ID NO: 24) in the promoter of the Human Growth Hormone gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

HGH1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>C</u>	<u>A</u>	<u>G</u>	<u>G</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>D32</u>	<u>C33</u>	<u>N30/</u> <u>Q38</u>	<u>R40/</u> <u>D28</u>	<u>R42</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

HGH2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>G</u>	<u>G</u>	<u>A</u>	<u>G</u>	<u>G</u>	<u>A</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>K32</u>	<u>R33</u>	<u>N30/ Q38</u>	<u>R40/ D28</u>	<u>R42</u>	<u>A26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

[0183] A rationally-designed meganuclease heterodimer (CF1/CF2) can be produced that cleaves the sequence 5'-GAAAATATCATTGGTGTTCCT-3' (SEQ ID NO: 25) in the  $\Delta$ F508 allele of the human CFTR gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

CF1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>G</u>	<u>A</u>	<u>A</u>	<u>A</u>	<u>A</u>	<u>T</u>	<u>A</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>Y33</u>	<u>N30/ Q38</u>	<u>Q40</u>	<u>K28</u>	<u>Q26</u>	<u>H68/ C24</u>	<u>Q44</u>	<u>R70</u>

CF2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>A</u>	<u>G</u>	<u>G</u>	<u>A</u>	<u>A</u>	<u>A</u>	<u>C</u>	<u>A</u>	<u>C</u>
<u>Contact Residues</u>	<u>N32</u>	<u>R33</u>	<u>E30/ R38</u>	<u>Q40</u>	<u>K28</u>	<u>A26</u>	<u>Y68/ C24</u>	<u>T44</u>	<u>R70</u>

[0184] A rationally-designed meganuclease heterodimer (CCR1/CCR2) can be produced that cleaves the sequence 5'-AACCTCTCCAGTGAGATGCCT-3' (SEQ ID NO: 26) in the human CCR5 gene (an HIV co-receptor). For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

CCR1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>A</u>	<u>A</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>T</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>N32</u>	<u>Y33</u>	<u>R30/</u> <u>E38</u>	<u>E40/</u> <u>R28</u>	<u>E42</u>	<u>Q26</u>	<u>Y68/</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

CCR2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>A</u>	<u>G</u>	<u>G</u>	<u>C</u>	<u>A</u>	<u>T</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>N32</u>	<u>R33</u>	<u>E30/</u> <u>R38</u>	<u>E40</u>	<u>K28</u>	<u>Q26</u>	<u>Y68/</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

[0185] A rationally-designed meganuclease heterodimer (MYD1/MYD2) can be produced that cleaves the sequence 5'-GACCTCGTCCTCCGACTCGCTG-3' (SEQ ID NO: 27) in the 3' untranslated region of the human DM kinase gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

MYD1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>G</u>	<u>A</u>	<u>C</u>	<u>C</u>	<u>T</u>	<u>C</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>Y33</u>	<u>R30/</u> <u>E38</u>	<u>E40/</u> <u>R28</u>	<u>K66</u>	<u>Q26/</u> <u>E77</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

MYD1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>A</u>	<u>G</u>	<u>C</u>	<u>G</u>	<u>A</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>Y33</u>	<u>E30/</u> <u>R38</u>	<u>E40/</u> <u>R28</u>	<u>R42</u>	<u>A26</u> <u>Q77</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

## 2. Meganuclease heterodimers which cleave DNA sequences in pathogen genomes.

[0186] A rationally-designed meganuclease heterodimer (HSV1/HSV2) can be produced that cleaves the sequence 5'-CTCGATGTCGGACGACACGGCA-3' (SEQ ID NO: 28) in the UL36 gene of Herpes Simplex Virus-1 and Herpes Simplex Virus-2. For

example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

## HSV1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>T</u>	<u>C</u>	<u>G</u>	<u>A</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>C33</u>	<u>R30/</u> <u>E38</u>	<u>R40/</u>	<u>Q42/</u> <u>K28</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

## HSV2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>G</u>	<u>C</u>	<u>C</u>	<u>G</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>C32</u>	<u>R33</u>	<u>R30/</u> <u>E38</u>	<u>E40/</u> <u>R28</u>	<u>R42</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

[0187] A rationally-designed meganuclease heterodimer (ANT1/ANT2) can be produced that cleaves the sequence 5'-ACAAGTGTCTATGGACAGTTTA-3' (SEQ ID NO: 29) in the *Bacillus anthracis* genome. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

## ANT1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>A</u>	<u>C</u>	<u>A</u>	<u>A</u>	<u>G</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>N32</u>	<u>C33</u>	<u>N30/</u> <u>Q38</u>	<u>Q40/</u> <u>A28</u>	<u>R42</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

## ANT2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>A</u>	<u>A</u>	<u>A</u>	<u>C</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>C32</u>	<u>Y33</u>	<u>N30/</u> <u>Q38</u>	<u>Q40</u>	<u>E42</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

[0188] A rationally-designed meganuclease heterodimer (POX1/POX2) can be produced that cleaves the sequence 5'-AAAAGTGTCAAATGACATCGCA-3' (SEQ ID NO: 30) in the Variola (smallpox) virus gp009 gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

POX1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>A</u>	<u>A</u>	<u>A</u>	<u>A</u>	<u>C</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>N32</u>	<u>C33</u>	<u>N30/ Q38</u>	<u>Q40</u>	<u>K28</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

POX2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>G</u>	<u>C</u>	<u>G</u>	<u>A</u>	<u>T</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>C32</u>	<u>R33</u>	<u>R30/ E38</u>	<u>R40</u>	<u>C28/ Q42</u>	<u>Q26</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

[0189] A rationally-designed meganuclease homodimer (EBB1/EBB1) can be produced that cleaves the pseudo-palindromic sequence 5'-CGGGGTCTCGTGCGAGGCCTCC-3' (SEQ ID NO: 31) in the Epstein-Barr Virus *BALF2* gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

EBB1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	<u>T</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>R33</u>	<u>D30/ Q38</u>	<u>R40/ D28</u>	<u>R42</u>	<u>Q26</u>	<u>Y68/ K24</u>	<u>Q44</u>	<u>R70</u>

EBB1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>G</u>	<u>G</u>	<u>A</u>	<u>G</u>	<u>G</u>	<u>C</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>R33</u>	<u>D30/</u> <u>Q38</u>	<u>R40/</u> <u>D28</u>	<u>R42</u>	<u>Q26</u>	<u>Y68/</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

### 3. Meganuclease heterodimers which cleave DNA sequences in plant genomes.

[0190] A rationally-designed meganuclease heterodimer (GLA1/GLA2) can be produced that cleaves the sequence 5'-CACTAACTCGTATGAGTCGGTG-3' (SEQ ID NO: 32) in the *Arabidopsis thaliana* GL2 gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

GLA1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>A</u>	<u>C</u>	<u>T</u>	<u>A</u>	<u>A</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>Y33</u>	<u>R30/</u> <u>E38</u>	<u>S40/</u> <u>C79</u>	<u>K28</u>	<u>A26/</u> <u>Q77</u>	<u>Y68/</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

GLA2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>A</u>	<u>C</u>	<u>C</u>	<u>G</u>	<u>A</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>Y33</u>	<u>R30/</u> <u>E38</u>	<u>E40/</u> <u>R28</u>	<u>R42</u>	<u>A26</u> <u>Q77</u>	<u>Y68/</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

[0191] A rationally-designed meganuclease heterodimer (BRP1/BRP2) can be produced that cleaves the sequence 5'-TGCCTCCTCTAGAGACCCGGAG-3' (SEQ ID NO: 33) in the *Arabidopsis thaliana* BP1 gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

BRP1:



<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>G</u>	<u>C</u>	<u>C</u>	<u>T</u>	<u>C</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>C32</u>	<u>R33</u>	<u>R30/</u> <u>E38</u>	<u>R28/</u> <u>E40</u>	<u>K66</u>	<u>Q26/</u> <u>E77</u>	<u>Y68/</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

## BRP2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>T</u>	<u>C</u>	<u>C</u>	<u>G</u>	<u>G</u>	<u>G</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>C33</u>	<u>R30/</u> <u>E38</u>	<u>E40/</u> <u>R28</u>	<u>R42</u>	<u>S26</u> <u>R77</u>	<u>R68</u>	<u>Q44</u>	<u>R70</u>

[0192] A rationally-designed meganuclease heterodimer (MGC1/MGC2) can be produced that cleaves the sequence 5'-TAAAATCTCTAAGGTCTGTGCA-3' (SEQ ID NO: 34) in the *Nicotiana tabacum* Magnesium Chelatase gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

## MGC1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>A</u>	<u>A</u>	<u>A</u>	<u>A</u>	<u>T</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>C32</u>	<u>Y33</u>	<u>N30/</u> <u>Q38</u>	<u>Q40/</u>	<u>K28</u>	<u>Q26</u>	<u>Y68/</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

## MGC2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>G</u>	<u>C</u>	<u>A</u>	<u>C</u>	<u>A</u>	<u>G</u>	<u>A</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>R33</u>	<u>R30/</u> <u>E38</u>	<u>Q40</u>	<u>K28</u>	<u>A26</u> <u>Q77</u>	<u>R68</u>	<u>T44</u>	<u>R70</u>

[0193] A rationally-designed meganuclease heterodimer (CYP/HGH2) can be produced that cleaves the sequence 5'-CAAGAATTCAAGCGAGCATTAA-3' (SEQ ID NO: 35) in the *Nicotiana tabacum* CYP82E4 gene. For example, a meganuclease was

designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

CYP:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>C</u>	<u>A</u>	<u>A</u>	<u>G</u>	<u>A</u>	<u>A</u>	<u>T</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>D32</u>	<u>Y33</u>	<u>N30/</u> <u>Q38</u>	<u>R40/</u>	<u>K28</u>	<u>Q77/</u> <u>A26</u>	<u>Y68</u>	<u>Q44</u>	<u>R70</u>

HGH2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>T</u>	<u>A</u>	<u>A</u>	<u>T</u>	<u>G</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>C33</u>	<u>N30/</u> <u>Q38</u>	<u>Q40</u>	<u>K66</u>	<u>R77/</u> <u>S26</u>	<u>Y68</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

4. Meganuclease heterodimers which cleave DNA sequences in yeast genomes.

[0194] A rationally-designed meganuclease heterodimer (URA1/URA2) can be produced that cleaves the sequence 5'-TTAGATGACAAGGGAGACGCAT-3' (SEQ ID NO: 36) in the *Saccharomyces cerevisiae* URA3 gene. For example, a meganuclease was designed based on the I-CreI meganuclease, as described above, with the following contact residues and recognition sequence half-sites:

URA1:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>T</u>	<u>T</u>	<u>A</u>	<u>G</u>	<u>A</u>	<u>T</u>	<u>G</u>	<u>A</u>	<u>C</u>
<u>Contact Residues</u>	<u>S32</u>	<u>C33</u>	<u>N30/</u> <u>Q38</u>	<u>R40</u>	<u>K28</u>	<u>Q26</u>	<u>R68</u>	<u>T44</u>	<u>R70</u>

URA2:

<u>Position</u>	<u>-9</u>	<u>-8</u>	<u>-7</u>	<u>-6</u>	<u>-5</u>	<u>-4</u>	<u>-3</u>	<u>-2</u>	<u>-1</u>
<u>Base</u>	<u>A</u>	<u>T</u>	<u>G</u>	<u>C</u>	<u>G</u>	<u>T</u>	<u>C</u>	<u>T</u>	<u>C</u>
<u>Contact Residues</u>	<u>N32</u>	<u>C33</u>	<u>E30/</u> <u>R38</u>	<u>E40/</u> <u>R28</u>	<u>R42</u>	<u>Q26</u>	<u>Y68/</u> <u>K24</u>	<u>Q44</u>	<u>R70</u>

#### 5. Recognition Sequence Specificity.

[0195] The rationally-designed meganucleases outlined above in this Example were cloned, expressed in *E. coli*, and purified as in Example 1. Each purified meganuclease was then mixed 1:1 with its corresponding heterodimerization partner (e.g., ACH1 with ACH2, HGH1 with HGH2, etc.) and incubated with a linearized DNA substrate containing the intended non-palindromic DNA recognition sequence for each meganuclease heterodimer. As shown in Figure 3, each rationally-designed meganuclease heterodimer cleaves its intended DNA site.

## **DEMANDES OU BREVETS VOLUMINEUX**

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS  
COMPREND PLUS D'UN TOME.**

**CECI EST LE TOME \_\_\_\_\_ DE \_\_\_\_\_**

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

## **JUMBO APPLICATIONS / PATENTS**

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE  
THAN ONE VOLUME.**

**THIS IS VOLUME   1   OF   2**

NOTE: For additional volumes please contact the Canadian Patent Office.

**THE EMBODIMENTS OF THE INVENTION FOR WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:**

1. A recombinant meganuclease having altered specificity for at least one recognition sequence half-site relative to a wild-type I-MsoI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6; and having specificity for a recognition sequence half-site which differs by at least one base pair from a half-site within an I-MsoI meganuclease recognition sequence selected from the group consisting of SEQ ID NO: 7 and SEQ ID NO: 8; wherein said recombinant meganuclease comprises at least one modification of Table 2 which is not an excluded modification.
2. A recombinant meganuclease having altered specificity for a recognition sequence relative to a wild-type I-SceI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 3-186 of the I- SceI meganuclease of SEQ ID NO: 9; and having specificity for a recognition sequence which differs by at least one base pair from an I-SceI meganuclease recognition sequence of SEQ ID NO: 10 and SEQ ID NO: 11; wherein said recombinant meganuclease comprises at least one modification of Table 3 which is not an excluded modification.
3. A recombinant meganuclease having altered specificity for at least one recognition sequence half-site relative to a wild-type I-CeuI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12; and having specificity for a recognition sequence half-site which differs by at least one base pair from a half-site within an I-CeuI meganuclease recognition sequence selected from the group consisting of SEQ ID NO: 13 and SEQ ID NO: 14; wherein said recombinant meganuclease comprises at least one modification of Table 4 which is not an excluded modification.
4. A recombinant meganuclease having altered specificity for at least one recognition sequence half-site relative to a wild-type I-MsoI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 6-160 of the I- MsoI meganuclease of SEQ ID NO: 6; and having specificity for a recognition sequence half-site which differs by at least one base pair from a half-site within an I-MsoI meganuclease recognition sequence selected from the group

consisting of SEQ ID NO: 7 and SEQ ID NO: 8; wherein:

(1) specificity at position -1 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of K75, Q77, A49, C49 and K79;

(b) to a T on a sense strand by a modification selected from the group consisting of C77, L77 and Q79; or

(c) to a G on a sense strand by a modification selected from the group consisting of K77, R77, E49 and E79; and/or

(2) specificity at position -2 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of Q75, K81, C47, I47 and L47;

(b) to a C on a sense strand by a modification selected from the group consisting of E75, D75, R47, K47, K81 and R81; or

(c) to a G on a sense strand by a modification selected from the group consisting of K75, E47 and E81; and/or

(3) specificity at position -3 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of Q72, C26, L26, V26, A26 and I26;

(b) to a C on a sense strand by a modification selected from the group consisting of E72, Y72, H26, K26 and R26; or

(c) to a T on a sense strand by a modification selected from the group consisting of K72, Y72 and H26; and/or

(4) specificity at position -4 has been altered:

(a) to a T on a sense strand by a modification selected from the group consisting of K28, K83 and Q28;

(b) to a G on a sense strand by a modification selected from the group consisting of R83 and K83; or

(c) to an A on a sense strand by a modification selected from the group consisting of K28 and Q83; and/or

(5) specificity at position -5 has been altered:

(a) to a G on a sense strand by a modification selected from the group consisting of R45 and

E28;

(b) to a T on a sense strand by a modification comprising Q28; or

(c) to a C on a sense strand by a modification comprising R28; and/or

(6) specificity at position -6 has been altered:

(a) to a T on a sense strand by a modification selected from the group consisting of K43, V85, L85 and Q30;

(b) to a C on a sense strand by a modification selected from the group consisting of E43, E85, K30 and R30; or

(c) to a G on a sense strand by a modification selected from the group consisting of R43, K43, K85, R85, E30 and D30; and/or

(7) specificity at position -7 has been altered:

(a) to a C on a sense strand by a modification selected from the group consisting of E32 and E41;

(b) to a G on a sense strand by a modification selected from the group consisting of R32, R41 and K41;

(c) to a T on a sense strand by a modification selected from the group consisting of K32, M41, L41 and I41; and/or

(8) specificity at position -8 has been altered:

(a) to a T on a sense strand by a modification selected from the group consisting of K32 and K35;

(b) to a C on a sense strand by a modification comprising E32; or

(c) to a G on a sense strand by a modification consisting of K32, K35 and R35; and/or

(9) specificity at position -9 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of N34 and H34;

(b) to a T on a sense strand by a modification selected from the group consisting of S34, C34, V34, T34 and A34; or

(c) to a G on a sense strand by a modification selected from the group consisting of K34, R34 and H34.

5. A recombinant meganuclease having altered specificity for a recognition sequence relative to a wild-type I-SceI meganuclease, comprising: a polypeptide having at least 85% sequence

similarity to residues 3-186 of the I-SceI meganuclease of SEQ ID NO: 9; and having specificity for a recognition sequence which differs by at least one base pair from an I-SceI meganuclease recognition sequence of SEQ ID NO: 10 and SEQ ID NO: 11; wherein:

(1) specificity at position 4 has been altered:

(a) to an A on a sense strand by a modification comprising K50;

(b) to a T on a sense strand by a modification selected from the group consisting of K57, M57 and Q50; or

(c) to a G on a sense strand by a modification selected from the group consisting of E50, R57 and K57; and/or

(2) specificity at position 5 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of K48, Q102;

(b) to a G on a sense strand by a modification selected from the group consisting of E48, K102 and R102; or

(c) to a T on a sense strand by a modification selected from the group consisting of Q48, C102, L102 and V102; and/or

(3) specificity at position 6 has been altered:

(a) to an A on a sense strand by a modification comprising K59;

(b) to a C on a sense strand by a modification selected from the group consisting of R59 and K59; or

(b) to a G on a sense strand by a modification selected from the group consisting of K84 and E59; and/or

(4) specificity at position 7 has been altered:

(a) to a C on a sense strand by a modification selected from the group consisting of R46, K46 and E86;

(b) to a G on a sense strand by a modification selected from the group consisting of K86, R86 and E46; or

(c) to an A on a sense strand by a modification selected from the group consisting of C46, L46 and V46; and/or

(5) specificity at position 8 has been altered:

(a) to a C on a sense strand by a modification selected from the group consisting of E88, R61 and H61;



- (b) to a T on a sense strand by a modification selected from the group consisting of K88, Q61 and H61; or
- (c) to an A on a sense strand by a modification selected from the group consisting of K61, S61 , V61 , A61 and L61; and/or
- (6) specificity at position 9 has been altered:
- (a) to an A on a sense strand by a modification selected from the group consisting of C98, V98 and L98;
- (b) to a C on a sense strand by a modification selected from the group consisting of R98 and K98; or
- (c) to a G on a sense strand by a modification selected from the group consisting of E98 and D98; and/or
- (7) specificity at position 10 has been altered:
- (a) to a C on a sense strand by a modification selected from the group consisting of K96 and R96;
- (b) to a G on a sense strand by a modification selected from the group consisting of D96 and E96; or
- (c) to an A on a sense strand by a modification selected from the group consisting of C96 and A96; and/or
- (8) specificity at position 11 has been altered:
- (a) to a T on a sense strand by a modification comprising Q90;
- (b) to a C on a sense strand by a modification selected from the group consisting of K90 and R90; or
- (c) to a G on a sense strand by a modification comprising E90; and/or
- (9) specificity at position 12 has been altered:
- (a) to an A on a sense strand by a modification comprising Q193;
- (b) to a C on a sense strand by a modification selected from the group consisting of E165, E193 and D193; or
- (c) to a G on a sense strand by a modification selected from the group consisting of K 165 and R165; and/or
- (10) specificity at position 13 has been altered:
- (a) to a T on a sense strand by a modification selected from the group consisting of Q193, C163

and L163;

(b) to a G on a sense strand by a modification selected from the group consisting of E193, D193, K163 and R192; or

(c) to an A on a sense strand by a modification selected from the group consisting of C193 and L193; and/or

(11) specificity at position 14 has been altered:

(a) to a T on a sense strand by a modification selected from the group consisting of K161 and Q192; (b) to an A on a sense strand by a modification selected from the group consisting of L192 and C192;

(c) to a G on a sense strand by a modification selected from the group consisting of K147, K161, R161, R197, D192 and E192; or

(d) to a T on a sense strand by a modification selected from the group consisting of K161 and Q192; and/or

(12) specificity at position 15 has been altered:

(a) to a T on a sense strand by a modification selected from the group consisting of C151, L151 and K151;

(b) to a G on a sense strand by a modification comprising K151 ; or

(c) to a C on a sense strand by a modification comprising E151; and/or

(13) specificity at position 17 has been altered:

(a) to a T on a sense strand by a modification selected from the group consisting of G152 and Q150;

(b) to a C on a sense strand by a modification selected from the group consisting of K152 and K150; or

(c) to a G on a sense strand by a modification selected from the group consisting of N152, S152, D152, D150 and E150; and/or

(14) specificity at position 18 has been altered:

(a) to a T on a sense strand by a modification selected from the group consisting of H155 and Y155;

(b) to a C on a sense strand by a modification selected from the group consisting of R155 and K155; or

(c) to an A on a sense strand by a modification selected from the group consisting of K155 and

C155.

6. A recombinant meganuclease having altered specificity for at least one recognition sequence half-site relative to a wild-type I-CeuI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12; and having specificity for a recognition sequence half-site which differs by at least one base pair from a half-site within an I-CeuI meganuclease recognition sequence selected from the group consisting of SEQ ID NO: 13 and SEQ ID NO: 14; wherein:

(1) specificity at position -1 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of C92, A92 and V92;

(b) to a T on a sense strand by a modification selected from the group consisting of Q116 and Q92; or

(c) to a G on a sense strand by a modification selected from the group consisting of E116 and E92; and/or

(2) specificity at position -2 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of Q117, C90, L90 and V90;

(b) to a G on a sense strand by a modification selected from the group consisting of K117, R124, K124, E124, E90 and D90; or

(c) to a C on a sense strand by a modification selected from the group consisting of E117, D117, R174, K124, K90, R90 and K68; and/or

(3) specificity at position -3 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of C70, V70, T70, L70 and K70;

(b) to a T on a sense strand by a modification comprising Q70;

(b) to a C on a sense strand by a modification consisting of K70; and/or

(4) specificity at position -4 has been altered:

(a) to a C on a sense strand by a modification selected from the group consisting of E126, D126, R88, K88 and K72;

(b) to a T on a sense strand by a modification selected from the group consisting of K126, L126

and Q88; or

(c) to an A on a sense strand by a modification selected from the group consisting of Q126, N126, K88, L88, C88, C72, L72 and V72; and/or

(5) specificity at position -5 has been altered:

(a) to a G on a sense strand by a modification selected from the group consisting of E74, K128, R128 and E128;

(b) to a T on a sense strand by a modification selected from the group consisting of C128, L128, V128 and T128; or

(c) to an A on a sense strand by a modification selected from the group consisting of C74, L74, V74 and T74; and/or

(6) specificity at position -6 has been altered:

(a) to a T on a sense strand by a modification selected from the group consisting of K86, C86 and L86;

(b) to a C on a sense strand by a modification selected from the group consisting of D86, E86, R84 and K84; or

(c) to a G on a sense strand by a modification selected from the group consisting of K128, R128, R86, K86 and E84; and/or

(7) specificity at position -7 has been altered:

(a) to a C on a sense strand by a modification selected from the group consisting of R76, K76 and H76;

(b) to a G on a sense strand by a modification selected from the group consisting of E76 and R84; or

(c) to a T on a sense strand by a modification consisting of H76 and Q76; and/or

(8) specificity at position -8 has been altered:

(a) to an A on a sense strand by a modification selected from the group consisting of Y79, R79 and Q76;

(b) to a C on a sense strand by a modification selected from the group consisting of D79, E79, D76 and E76; or

(c) to a G on a sense strand by a modification selected from the group consisting of R79, K79, K76 and R76; and/or

(9) specificity at position -9 has been altered:

- (a) to a T on a sense strand by a modification selected from the group consisting of K78, V78, L78, C78 and T78;
- (b) to a C on a sense strand by a modification selected from the group consisting of D78 and E78; or
- (c) to a G on a sense strand by a modification selected from the group consisting of R78, K78 and H78.

7. A recombinant meganuclease having altered binding affinity for double-stranded DNA relative to a wild-type I-MsoI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 6-160 of the I- MsoI meganuclease of SEQ ID NO: 6; wherein DNA-binding affinity has been increased by at least one modification corresponding to a substitution selected from the group consisting of:

- (a) substitution of E147, 185, G86 or Y118 with H, N, Q, S, T, K or R; or
- (b) substitution of Q41, N70, S87, T88, H89, Q122, Q139, S150 or N152 with K or R.

8. A recombinant meganuclease having altered binding affinity for double-stranded DNA relative to a wild-type I-MsoI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 6-160 of the I- MsoI meganuclease of SEQ ID NO: 6; wherein DNA-binding affinity has been decreased by at least one modification corresponding to a substitution selected from the group consisting of:

- (a) substitution of K36, R51, K123, K143 or R144 with H, N, Q, S, T, D or E; or
- (b) substitution of I85, G86, Y118, Q41, N70, S87, T88, H89, Q122, Q139, S150 or N152 with D or E.

9. A recombinant meganuclease having altered binding affinity for double-stranded DNA relative to a wild-type I-SceI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 3-186 of the I-SceI meganuclease of SEQ ID NO: 9; wherein DNA-binding affinity has been increased by at least one modification corresponding to a substitution selected from the group consisting of:

- (a) substitution of D201, L19, L80, L92, Y151, Y188, I191, Y199 or Y222 with H, N, Q, S, T, K or R; or

(b) substitution of N15, N17, S81, H84, N94, N120, T156, N157, S159, N163, Q165, S166, N194 or S202 with K or R.

10. A recombinant meganuclease having altered binding affinity for double-stranded DNA relative to a wild-type I-SceI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 3-186 of the I- SceI meganuclease of SEQ ID NO: 9; wherein DNA-binding affinity has been decreased by at least one modification corresponding to a substitution selected from the group consisting of:

(a) substitution of K20, K23, K63, K122, K148, K153, K190, K193, K195 or K223 with H, N, Q, S, T, D or E; or (b) substitution of L19, L80, L92, Y151, Y188, I191, Y199, Y222, N15, N17, S81, H84, N94, N120, T156, N157, S159, N163, Q165, S166, N194 or S202 with D or E.

11. A recombinant meganuclease having altered binding affinity for double-stranded DNA relative to a wild-type I-CeuI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12; wherein DNA-binding affinity has been increased by at least one modification corresponding to a substitution selected from the group consisting of:

(a) substitution of D25 or D128 with H, N, Q, S, T, K or R; or  
(b) substitution of S68, N70, H94, S117, N120, N129 or H172 with K or R.

12. A recombinant meganuclease having altered binding affinity for double-stranded DNA relative to a wild-type I-CeuI meganuclease, comprising: a polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12; wherein DNA-binding affinity has been decreased by at least one modification corresponding to a substitution selected from the group consisting of:

(a) substitution of K21, K28, K31, R112, R114 or R130 with H, N, Q, S, T, D or E; or  
(b) substitution of S68, N70, H94, S117, N120, N129 or H172 with D or E.

13. A recombinant meganuclease monomer having altered affinity for dimer formation with a reference meganuclease monomer, comprising: a polypeptide having at least 85% sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6; wherein affinity for

dimer formation has been altered by at least one modification corresponding to a substitution selected from the group consisting of:

- (a) substitution of R302 with D or E; or
- (b) substitution of D20, E11 or Q64 with K or R.

14. A recombinant meganuclease heterodimer comprising: a first polypeptide having at least 85% sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6; wherein affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from the group consisting of:

- (a) substitution of R302 with D or E; and a second polypeptide having at least 85% sequence similarity to residues 6-160 of the I-MsoI meganuclease of SEQ ID NO: 6; wherein affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from the group consisting of: (b) substitution of D20, E11 or Q64 with K or R.

15. A recombinant meganuclease monomer having altered affinity for dimer formation with a reference meganuclease monomer, comprising: a polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12; wherein affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from the group consisting of:

- (a) substitution of R93 with D or E; or
- (b) substitution of E152 with K or R.

16. A recombinant meganuclease heterodimer comprising: a first polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12; wherein affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from the group consisting of:

- (a) substitution of R93 with D or E; and a second polypeptide having at least 85% sequence similarity to residues 5-211 of the I-CeuI meganuclease of SEQ ID NO: 12; wherein affinity for dimer formation has been altered by at least one modification corresponding to a substitution selected from the group consisting of:
- (b) substitution of E 152 with K or R.

17. A recombinant meganuclease monomer or heterodimer of claim 13 or claim 14, further comprising: at least one modification selected from Table 2; wherein said modification is not an excluded modification.

18. A recombinant meganuclease monomer or heterodimer of claim 15 or claim 16, further comprising: at least one modification selected from Table 4; wherein said modification is not an excluded modification.

19. A method for producing a genetically-modified eukaryotic cell including an exogenous sequence of interest inserted in a chromosome of said eukaryotic cell, comprising: transfecting a eukaryotic cell with one or more nucleic acids including (i) a first nucleic acid sequence encoding a meganuclease, and (ii) a second nucleic acid sequence including said sequence of interest; wherein said meganuclease produces a cleavage site in said chromosome and said sequence of interest is inserted into said chromosome at said cleavage site; and wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18.

20. A method as in claim 19 wherein: said second nucleic acid further comprises sequences homologous to sequences flanking said cleavage site and said sequence of interest is inserted at said cleavage site by homologous recombination.

21. A method as in claim 19 wherein: said second nucleic acid lacks substantial homology to said cleavage site and said sequence of interest is inserted into said chromosome by non-homologous end-joining.

22. A method for producing a genetically-modified eukaryotic cell including an exogenous sequence of interest inserted in a chromosome of said eukaryotic cell, comprising: introducing a meganuclease protein into a eukaryotic cell; and transfecting said eukaryotic cell with a nucleic acid including said sequence of interest; wherein said meganuclease produces a cleavage site in said chromosome and said sequence of interest is inserted into said chromosome at said cleavage site; and wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18.



23. A method as in claim 22 wherein: said nucleic acid further comprises sequences homologous to sequences flanking said cleavage site and said sequence of interest is inserted at said cleavage site by homologous recombination.

24. A method as in claim 22 wherein: said nucleic acid lacks substantial homology to said cleavage site and said sequence of interest is inserted into said chromosome by non-homologous end-joining.

25. A method for producing a genetically-modified eukaryotic cell by disrupting a target sequence in a chromosome of said eukaryotic cell, comprising: transfecting a eukaryotic cell with a nucleic acid encoding a meganuclease; wherein said meganuclease produces a cleavage site in said chromosome and said target sequence is disrupted by non-homologous end-joining at said cleavage site; and wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18.

26. A method of producing a genetically-modified organism comprising: producing a genetically-modified eukaryotic cell according to the method of any one of claims 19-25; and growing said genetically-modified eukaryotic cell to produce said genetically-modified organism.

27. A method as in claim 26 wherein: said eukaryotic cell is selected from the group consisting of a gamete, a zygote, a blastocyst cell, an embryonic stem cell, and a protoplast cell.

28. Use of one or more nucleic acids to treat a disease by gene therapy in a eukaryote, wherein said one or more nucleic acids includes  
(i) a first nucleic acid sequence encoding a meganuclease, and (ii) a second nucleic acid sequence including a sequence of interest; wherein said meganuclease produces a cleavage site in said chromosome and said sequence of interest is inserted into said chromosome at said cleavage site; wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18; and wherein insertion of said sequence of interest provides said gene therapy for said disease.

29. The use according to claim 28 wherein: said second nucleic acid sequence further comprises sequences homologous to sequences flanking said cleavage site and said sequence of interest is inserted at said cleavage site by homologous recombination.

30. The use according to claim 28 wherein: said second nucleic acid sequence lacks substantial homology to said cleavage site and said sequence of interest is inserted into said chromosome by non-homologous end-joining.

31. Use of a meganuclease protein and a nucleic acid including a sequence of interest to treat a disease by gene therapy in a eukaryote; wherein said meganuclease produces a cleavage site in said chromosome and said sequence of interest is inserted into said chromosome at said cleavage site; wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18; and wherein insertion of said sequence of interest provides said gene therapy for said disease.

32. The use according to claim 31 wherein: said nucleic acid further comprises sequences homologous to sequences flanking said cleavage site and said sequence of interest is inserted at said cleavage site by homologous recombination.

33. The use according to claim 31 wherein: said nucleic acid lacks substantial homology to said cleavage site and said sequence of interest is inserted into said chromosome by non-homologous end-joining.

34. Use of a nucleic acid encoding a meganuclease to treat a disease by gene therapy in a eukaryote by disrupting a target sequence in a chromosome of said eukaryotic cell; wherein said meganuclease produces a cleavage site in said chromosome and said target sequence is disrupted by non-homologous end-joining at said cleavage site; wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18; and wherein disruption of said target sequence provides said gene therapy for said disease.

35. Use of a nucleic acid encoding a meganuclease to treat a viral pathogen infection in a

eukaryotic host by disrupting a target sequence in a genome of said viral pathogen; wherein said meganuclease produces a cleavage site in said viral genome and said target sequence is disrupted by non-homologous end-joining at said cleavage site; wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18; and wherein disruption of said target sequence provides treatment for said infection.

36. Use of a first nucleic acid encoding a meganuclease and a second nucleic acid to treat a viral pathogen infection in a eukaryotic host by disrupting a target sequence in a genome of said viral pathogen; wherein said meganuclease produces a cleavage site in said viral genome and said target sequence is disrupted by homologous recombination of said viral genome and said second nucleic acid at said cleavage site; wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18; wherein said second nucleic acid comprises sequences homologous to sequences flanking said cleavage site; and wherein disruption of said target sequence provides treatment for said infection.

37. Use of a nucleic acid encoding a meganuclease to treat a prokaryotic pathogen infection in a eukaryotic host by disrupting a target sequence in a genome of said prokaryotic pathogen; wherein said meganuclease produces a cleavage site in said prokaryotic genome and said target sequence is disrupted by non-homologous end-joining at said cleavage site; wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18; and wherein disruption of said target sequence provides treatment for said infection.

38. Use of a first nucleic acid encoding a meganuclease and a second nucleic acid to treat a prokaryotic pathogen infection in a eukaryotic host by disrupting a target sequence in a genome of said prokaryotic pathogen; wherein said meganuclease produces a cleavage site in said prokaryotic genome and said target sequence is disrupted by homologous recombination of said prokaryotic genome and said second nucleic acid at said cleavage site; wherein said meganuclease is a recombinant meganuclease of any one of claims 1-18; wherein said second nucleic acid comprises sequences homologous to sequences flanking said cleavage site; and wherein disruption of said target sequence provides treatment for said infection.

39. A method for rationally-designing a recombinant meganuclease having altered specificity for at least one base position of a recognition sequence, said altered specificity including at least one desired change, the method comprising:

- (1) determining at least a portion of a three-dimensional structure of a reference meganuclease-DNA complex;
- (2) identifying amino acid residues forming a base contact surface at said base position;
- (3) determining a distance between a  $\beta$ -carbon of at least a first residue comprising said contact surface and at least a first base at said base position;
- (4) identifying an amino acid substitution to promote said desired change by
  - (a) for a first residue which is  $< 6$  Å from said first base, selecting a substitution from Group 1 and/or Group 2 which is a member of an appropriate one of Group G, Group C, Group T or Group A to promote said desired change; or
  - (b) for a first residue which is  $> 6$  Å from said first base, selecting a substitution from Group 2 and/or Group 3 which is a member of an appropriate one of Group G, Group C, Group T or Group A to promote said desired change.

40. A method for rationally-designing a recombinant meganuclease having increased DNA-binding affinity, comprising:

- (1) determining at least a portion of a three-dimensional structure of a reference meganuclease-DNA complex;
- (2) identifying amino acid contact residues forming a backbone contact surface;
- (3) identifying an amino acid substitution to increase said DNA-binding affinity by
  - (a) for a contact residue having a negatively-charged or hydrophobic side chain, selecting a substitution having an uncharged/polar or positively-charged side chain; or
  - (b) for a contact residue having an uncharged/polar side chain, selecting a substitution having a positively-charged side chain.

41. A method for rationally-designing a recombinant meganuclease having decreased DNA-binding affinity, comprising:

- ( 1 ) determining at least a portion of a three-dimensional structure of a reference meganuclease-DNA complex;

- (2) identifying amino acid contact residues forming a backbone contact surface;
- (3) identifying an amino acid substitution to decrease said DNA-binding affinity by
  - (a) for a contact residue having a positively-charged side chain, selecting a substitution having an uncharged/polar or negatively-charged side chain; or
  - (b) for a contact residue having an hydrophobic or uncharged/polar side chain, selecting a substitution having a negatively-charged side chain.




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 1E

2/4

WT **CAA**ACTGTCGTGAGACAGTTG

CF1/CF2 G**AAA**A**TAT**CA T T G**GTG**T**TT**CCT

MYD1/MYD2 G**ACC**T**CGT**CCT**CC**G**AC**T**CG**C**TG**

CCR1/CCR2 A**ACC**C**CTC**CA**GT**G**AG**A**TG**CCT

ACH1/ACH2 **CTG**GGAG**GTC**TCAG**GAC**AG**CCT**G

TAT1/TAT2 G**AG**AG**CTC**A**TC**AG**AC**AG**TCA**

HSV1/HSV2 **CTC**GA**TGT**CG**GAC**G**ACA**CGGCA

LAM1/LAM2 T**GCG**G**TGT**C**CG**G**C**G**AC**AG**CCT**G

POX1/POX2 A**AA**ACTGTC**AA**AT**GAC**A**TG**CGCA

URA1/URA2 T**TAG**A**TG**AC**AA**GG**GAG**A**CG**CAT

GLA1/GLA2 **CAC**T**AA**CT**CGT**AT**GAG**T**CG**GG**TG**

BRP1/BRP2 T**GC**CT**CC****TC**TA**GAG**AC**CC**GG**AG**

FIG. 2A

Enzyme Site													
	WT	CF	MYD	CCR	ACH	TAT	HSV	LAM	POX	URA	GLA	BRP	
WT	100	0	0	0	0	0	47	0	100	100	0	0	Wild Type
CF1/CF2	0	63	0	0	0	0	0	0	0	0	0	0	Cystic Fibrosis
MYD1/MYD2	0	0	94	0	0	0	16	0	0	0	0	0	Myotonic Dystrophy
CCR1/CCR2	0	0	0	100	0	0	0	0	0	0	0	0	CCR5
ACH1/ACH2	0	0	0	0	87	0	0	0	0	0	0	0	Achondroplasia
TAT1/TAT2	0	0	0	0	0	81	0	0	0	0	0	0	HIV-1
HSV1/HSV2	0	0	0	0	0	0	67	0	0	0	0	0	Herpes Simplex 1
LAM1/LAM2	0	0	0	0	0	0	0	87	0	0	0	0	Lambda Phage
POX1/POX2	100	0	0	0	0	0	0	0	100	0	0	0	Variola Virus
URA1/URA2	0	0	0	0	0	0	0	0	0	70	0	0	<i>S. cerevisiae</i> URA3
GLA1/GLA2	0	0	0	0	0	0	0	0	0	0	88	0	<i>A. thaliana</i> GL2
BRP1/BRP2	0	0	0	0	0	0	0	0	0	0	0	100	<i>A. thaliana</i> BP1

FIG. 2B

FIG. 3A  
FIG. 3B

FIG. 3




FIG. 3A






FIG. 3B

SUBSTITUTE SHEET (RULE 26)