



US00PP32312P2

(12) **United States Plant Patent**  
**Hofmann**

(10) **Patent No.:** **US PP32,312 P2**

(45) **Date of Patent:** **Oct. 13, 2020**

(54) **MANDEVILLA PLANT NAMED**  
**'INMANBEGSC'**

(50) Latin Name: *Mandevilla sanderi*  
Varietal Denomination: **Inmanbegsc**

(71) Applicant: **Birgit Hofmann**, Rudesheim am Rhein  
(DE)

(72) Inventor: **Birgit Hofmann**, Rudesheim am Rhein  
(DE)

(73) Assignee: **INNOVAPLANT ZIERPFLANZEN**  
**GmbH & Co. KG**, Gensingen (DE)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **16/602,286**

(22) Filed: **Sep. 6, 2019**

(51) **Int. Cl.**  
*A01H 5/02* (2018.01)  
*A01H 6/08* (2018.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./232**  
CPC ..... *A01H 6/088* (2018.05)

(58) **Field of Classification Search**  
USPC ..... Plt./232  
CPC ..... *A01H 6/088*  
See application file for complete search history.

*Primary Examiner* — Anne Marie Grunberg  
(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Mandevilla* plant named  
'Inmanbegsc', characterized by its broadly upright and vin-  
ing plant habit; moderately vigorous growth habit; freely  
branching habit; glossy dark green-colored leaves; early and  
freely flowering habit; large deep scarlet red-colored flowers  
with slightly undulate margins and orange red-colored centers  
that resist fading; and relative tolerance to low tempera-  
tures.

**2 Drawing Sheets**

**1**

Botanical designation: *Mandevilla sanderi*.  
Cultivar denomination: 'INMANBEGSC'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of *Mandevilla* plant, botanically known as *Mandevilla san-*  
*deri* and hereinafter referred to by the name 'Inmanbegsc'.

The new *Mandevilla* plant is a product of a planned  
breeding program conducted by the Inventor in Johannes-  
burg, South Africa and Gensingen, Germany. The objective  
of the breeding program is to create new freely branching  
*Mandevilla* plants that flower early and freely and have  
attractive flowers that resist fading and sun scald.

The new *Mandevilla* plant originated from a cross-pollina-  
tion conducted by the Inventor in Johannesburg, South  
Africa in January, 2014 of a proprietary selection of *Man-*  
*devilla sanderi* identified as code number D11-2524-1, not  
patented, as the female, or seed parent with a proprietary  
selection of *Mandevilla sanderi* identified as code number  
D11-2102-1, not patented, as the male, or pollen, parent. The  
new *Mandevilla* plant was discovered and selected by the  
Inventor as a single flowering plant from within the progeny  
of the stated cross-pollination in a controlled greenhouse  
environment in Gensingen, Germany in June, 2015.

Asexual reproduction of the new *Mandevilla* plant by  
vegetative cuttings in a controlled greenhouse environment  
in Gensingen, Germany since August, 2015 has shown that  
the unique features of this new *Mandevilla* plant are stable  
and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

Plants of the new *Mandevilla* have not been observed  
under all possible combinations of environmental conditions

**2**

and cultural practices. The phenotype may vary somewhat  
with variations in environmental conditions such as tem-  
perature and light intensity without, however, any variance  
in genotype.

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of 'Inman-  
begsc'. These characteristics in combination distinguish  
'Inmanbegsc' as a new and distinct *Mandevilla* plant:

1. Broadly upright and vining plant habit.
2. Moderately vigorous growth habit.
3. Freely branching habit.
4. Glossy dark green-colored leaves.
5. Early and freely flowering habit.
6. Large deep scarlet red-colored flowers with slightly  
undulate margins and orange red-colored centers that  
resist fading.
7. Relatively tolerant to low temperatures.

Plants of the new *Mandevilla* can be compared to plants  
of the female parent selection. Plants of the new *Mandevilla*  
differ primarily from plants of the female parent selection in  
the following characteristics:

1. Plants of the new *Mandevilla* are more freely branching  
and bushier than plants of the female parent selection.
2. Plants of the new *Mandevilla* flower later than plants of  
the female parent selection.
3. Plants of the new *Mandevilla* have deep scarlet red-  
colored flowers with orange red-colored centers  
whereas plants of the female parent selection have  
slightly deep red-colored flowers with white-colored  
centers.

Plants of the new *Mandevilla* can be compared to plants  
of the male parent selection. Plants of the new *Mandevilla*  
differ primarily from plants of the male parent selection in  
the following characteristics:

1. Plants of the new *Mandevilla* have stronger roots than plants of the male parent selection.
2. Plants of the new *Mandevilla* are more upright than plants of the male parent selection.
3. Leaves of plants of the new *Mandevilla* are smoother and flatter than leaves of plants of the male parent selection.
4. Flower margins of plants of the new *Mandevilla* are undulate whereas flower margins of plants of the male parent selection are not undulate.

Plants of the new *Mandevilla* can also be compared to plants of *Mandevilla hybrida* 'Inmanredeep', disclosed in U.S. Plant Pat. No. 29,096. In side-by-side comparisons, plants of the new *Mandevilla* differ primarily from plants 'Inmanredeep' in the following characteristics:

1. Plants of the new *Mandevilla* are larger than plants of 'Inmanredeep'.
2. Plants of the new *Mandevilla* have larger flowers than plants of 'Inmanredeep'.
3. Flowers of plants of the new *Mandevilla* are brighter red in color than flowers of plants of 'Inmanredeep'.
4. Flower margins of plants of the new *Mandevilla* are undulate whereas flower margins of plants of 'Inmanredeep' are not undulate.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Mandevilla* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the actual colors of the new *Mandevilla* plant.

The photograph on the first sheet (FIG. 1 of 2) comprises a side perspective view of a typical flowering plant of 'Inmanbegsc' grown in a container.

The photograph on the second sheet (FIG. 2 of 2) is a close-up view of a typical flower of 'Inmanbegsc'.

#### DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown during the late spring and early summer in 11-cm containers in a glass-covered greenhouse in Heidesheim, Germany and under cultural practices typical of commercial *Mandevilla* production. During the production of the plants, day temperatures ranged from 12° C. to 35° C. and night temperatures ranged from 12° C. to 18° C. Plants were seven months old when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2015 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Mandevilla sanderi* 'Inmanbegsc'. Parentage:

*Female, or seed, parent.*—Proprietary selection of *Mandevilla sanderi* identified as code number D11-2524-1.

*Male, or pollen, parent.*—Proprietary selection of *Mandevilla sanderi* identified as code number D11-2102-1.

Propagation:

*Type.*—By vegetative cuttings.

*Time to initiate roots, summer.*—About three weeks at temperatures ranging from 20° C. to 35° C.

*Time to initiate roots, winter.*—About three weeks at temperatures ranging from 20° C. to 25° C.

*Time to produce a rooted young plant, summer.*—About 20 to 25 days at temperatures ranging from 20° C. to 35° C.

*Time to produce a rooted young plant, winter.*—About 25 to 30 days at temperatures ranging from 20° C. to 25° C.

*Root description.*—Thick, fleshy; typically white in color, actual color of the roots is dependent on substrate composition, water quality, fertilizers, substrate temperature and physiological age of roots.

*Rooting habit.*—Low branching; medium density.

Plant description:

*Plant and growth habit.*—Broadly upright and vining plant habit; roughly broadly obovate in overall shape; moderately vigorous growth habit and moderate growth rate.

*Plant height, soil level to top of foliar plane.*—About 35.3 cm.

*Plant height, soil level to top of floral plane.*—About 37 cm.

*Plant diameter (spread).*—About 31.3 cm.

*Lateral branch description.*—Branching habit: Freely branching habit, typically about three primary lateral branches each with about two secondary lateral branches. Length: About 16.3 cm. Diameter: About 2.5 mm. Internode length: About 2.5 cm. Aspect: Primary lateral branches, erect to about 10° from vertical; secondary lateral branches, about 20° from primary branch axis. Strength: Strong. Texture and luster: Smooth, glabrous; glossy; becoming woody with development. Color, developing: Close to 144A. Color, developed: Close to between 144A and 146D; when woody, close to N199A.

Leaf description:

*Arrangement.*—Opposite, simple.

*Length.*—About 6.8 cm.

*Width.*—About 5.1 cm.

*Shape.*—Broadly elliptic-oblong, broadly ovate to broadly obovate.

*Apex.*—Short apiculate.

*Base.*—Obtuse to truncate.

*Margin.*—Entire; not undulate.

*Texture and luster, upper and lower surfaces.*—Smooth, glabrous; moderately coriaceous; moderately glossy.

*Venation pattern.*—Pinnate.

*Color.*—Developing leaves, upper surface: Close to 146A. Developing leaves, lower surface: Close to 146B. Full expanded leaves, upper surface: Darker than between 139A and 147A; venation, close to 143A. Fully expanded leaves, lower surface: Close to between 146A and 147B; venation, close to 144C.

*Petioles.*—Length: About 1 cm. Diameter: About 2 mm by 2 mm. Strength: Moderately strong. Texture and luster, upper and lower surfaces: Smooth, glabrous; moderately glossy. Color, upper surface: Close to 144A. Color, lower surface: Close to 144B to 144C.

## Flower description:

*Flower type and flowering habit.*—Single salverform flowers arranged in axillary cymes; flowers face outwardly to somewhat upright; freely flowering habit with about five flowers per inflorescence and about 45 flower buds and open flowers developing per plant during the flowering season.

*Natural flowering season.*—Plants flower continuously from spring into the autumn in Germany; plants begin flowering about 16 weeks after propagation (dependent on light level).

*Flower longevity on the plant.*—Individual flowers last about ten days; flowers not persistent.

*Fragrance.*—None detected.

*Inflorescence height.*—About 11 cm.

*Inflorescence diameter.*—About 10.8 cm.

*Flower buds.*—Length: About 6.1 cm. Diameter: About 1.1 cm. Shape: Narrowly oblanceolate. Texture and luster: Smooth, glabrous; slightly glossy. Color: Distally, close to 53A; towards the base, close to 53B to 53C; at the base, close to 145A, 145B and 145C.

*Flowers.*—Appearance: Flared trumpet, corolla fused and five-parted. Diameter: Large, about 8.6 cm. Depth (length): About 5.9 cm. Throat diameter: About 1.8 cm. Tube length: About 5.1 cm. Tube diameter: Proximally, about 4 mm; distally, about 2.1 cm.

*Corolla.*—Quantity and arrangement: Five petals arranged in a single whorl; lower 58% portion of the petals are fused into a funnellform tube. Petal length: About 8.8 cm. Petal width: About 3.6 cm. Petal shape and appearance: Roughly unequal spatulate. Petal apex: Apiculate. Petal margin: Entire; moderately undulate. Petal texture and luster, upper surface: Smooth, glabrous; velvety; matte. Petal texture and luster, lower surface: Smooth, glabrous; slightly velvety; moderately glossy. Throat texture: Smooth, glabrous; velvety. Tube texture: Smooth, glabrous; slightly velvety. Color: Petal, when opening, upper surface: More intense than between 53A and 187C. Petal, when opening, lower surface: Close to 53A. Petal, fully opened, upper surface: Slightly more intense than 53A; venation, similar to lamina; color does not change with development. Petal, fully opened, lower surface: Close to 53A; venation, similar to lamina; color becoming closer to 53B with development. Throat: Close to 31A to 31B flushed with close to 15A proximally; at the base, close to N144B; venation, similar to lamina. Tube: Close to

47A; proximally, close to 145C and at the base, close to 145B; venation, similar to lamina.

*Calyx.*—Quantity and arrangement: Five sepals arranged in a single whorl. Calyx length: About 7 mm. Calyx diameter: About 5 mm. Sepal length: About 6 mm. Sepal width: About 1.5 mm. Sepal shape: Lanceolate. Sepal apex: Narrowly acuminate. Sepal base: Broadly cuneate. Sepal margin: Entire. Sepal texture and luster, upper and lower surfaces: Smooth, glabrous; moderately glossy. Sepal color: When developing, upper and lower surfaces: Close to N144A; fading towards the base to close to 144A; at the apex, close to 179B. Fully developed, upper and lower surfaces: Close to N144A; fading towards the base to close to 144A; at the apex, close to 179B.

*Peduncles.*—Length: About 6.5 cm. Diameter: About 2 mm. Strength: Strong. Aspect: About 30° from lateral branch axis. Texture and luster: Smooth, glabrous; glossy. Color: Close to 144B.

*Pedicels.*—Length: About 1.3 cm. Diameter: About 2 mm. Strength: Strong. Aspect: About 15° to 40° from peduncle axis. Texture and luster: Smooth, glabrous; moderately glossy. Color: Close to 143C.

*Reproductive organs.*—Stamens: Quantity and arrangement: Typically five; basifixed; anthers connivent. Filament length: About 3 mm. Filament color: Close to 153D. Anther shape: Narrowly oblong. Anther size: About 9 mm by 1.5 mm. Anther color: Close to 161D. Pollen amount: None observed. Pistils: Quantity: Typically one. Pistil length: About 2.7 cm. Style length: About 2.4 cm. Style color: Close to 145D. Stigma diameter: About 2 mm. Stigma shape: Club-shaped, pointed. Stigma color: Close to 148C. Ovary color: Close to 143C.

*Seeds and fruits.*—To date, seed and fruit production have not been observed on plants of the new *Mandevilla*.

*Pathogen & pest resistance:* To date, plants of the new *Mandevilla* have not been noted to be resistant to pathogens and pests common to *Mandevilla* plants.

*Temperature tolerance:* Plants of the new *Mandevilla* have been observed to tolerate temperatures ranging from about 5° C. to about 40° C. and to be suitable for USDA Hardiness Zones 9 to 13.

It is claimed:

1. A new and distinct *Mandevilla* plant named 'Inmanbegs' as illustrated and described.

\* \* \* \* \*

FIG. 1



FIG. 2

