



(51) International Patent Classification:

C07D 487/04 (2006.01) *A61K 31/519* (2006.01)
C07D 519/00 (2006.01) *A61P 29/00* (2006.01)

(21) International Application Number:

PCT/EP2012/001737

(22) International Filing Date:

23 April 2012 (23.04.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/517,582 21 April 2011 (21.04.2011) US

(71) Applicant (for all designated States except US): **ORIGENIS GMBH** [DE/DE]; Am Klopferspitz 19a, 52152 Martinsried (DE).

(72) Inventors; and

(75) **Inventors/Applicants (for US only): ALMSTETTER, Michael** [DE/DE]; Am Klopferspitz 19a, 52152 Martinsried (DE). **THORMANN, Michael** [DE/DE]; Am Klopferspitz 19a, 52152 Martinsried (DE). **TREML, Andreas** [DE/DE]; Am Klopferspitz 19a, 52152 Martinsried (DE). **TRAUBE, Nadine** [DE/DE]; Am Klopferspitz 19a, 52152 Martinsried (DE).

(74) **Agents: FORSTMAYER, Dietmar** et al.; Boeters & Lieck, Oberanger 32, 80331 München (DE).


(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: PYRAZOLO [4, 3-D] PYRIMIDINES USEFUL AS KINASE INHIBITORS



(I)

(57) **Abstract:** The present invention relates to novel compounds of formula (I) that are capable of inhibiting one or more kinases, especially SYK (Spleen Tyrosine Kinase), LRRK2 (Leucine-rich repeat kinase 2) and/or MYLK (Myosin light chain kinase) or mutants thereof. The compounds find applications in the treatment of a variety of diseases. These diseases include autoimmune diseases, inflammatory diseases, bone diseases, metabolic diseases, neurological and neurodegenerative diseases, cancer, cardiovascular diseases, allergies, asthma, alzheimer's disease, parkinson's disease, skin disorders, eye diseases, infectious diseases and hormone-related diseases.

PYRAZOLO [4 , 3-D] PYRIMIDINES USEFUL AS KINASE INHIBITORS

The present invention relates to novel compounds that are capable of inhibiting one or more kinases, especially SYK (Spleen Tyrosine Kinase), LRRK2 (Leucine-rich repeat kinase 2) and/or MYLK (Myosin light chain kinase) or mutants thereof. The compounds find applications in the treatment of a variety of diseases. These diseases include autoimmune diseases, inflammatory diseases, bone diseases, metabolic diseases, neurological and neurodegenerative diseases, cancer, cardiovascular diseases, allergies, asthma, alzheimer's disease, parkinson's disease, skin disorders, eye diseases, infectious diseases and hormone-related diseases.

BACKGROUND OF THE INVENTION

Protein kinases constitute a large family of structurally related enzymes that are responsible for the control of a variety of signal transduction processes within cells (see, e.g., Hardie and Hanks, The Protein Kinase Facts Book, I and II, Academic Press, San Diego, Calif., 1995). Protein kinases are thought to have evolved from a common ancestral gene due to the conservation of their structure and catalytic function. Almost all kinases contain a similar 250-300 amino acid catalytic domain. The kinases can be categorized into families by the substrates they phosphorylate (e.g., protein-tyrosine, protein-serine/threonine, lipids, etc.). Sequence motifs have been identified that generally correspond to each of these families (see, e.g., Hanks & Hunter, (1995), FASEB J. 9:576-596; Knighton et al., (1991), Science 253:407-414; Hiles et al., (1992), Cell 70:419-429; Kunz et al., (1993), Cell 73:585-596; Garcia-Bustos et al., (1994), EMBO J. 13:2352-2361).

Many diseases are associated with abnormal cellular responses triggered by protein kinase-mediated events. These diseases include autoimmune diseases, inflammatory diseases, bone diseases, metabolic diseases, neurological and neurodegenerative diseases, cancer, cardiovascular diseases, allergies, asthma, alzheimer's disease, parkinson's disease skin disorders, infectious diseases and hormone-related diseases. As a consequence, there has been substantial effort in medicinal chemistry to find inhibitors of protein kinases for use as therapeutic agents.

SYK - Spleen Tyrosine Kinase

Syk is known to play an essential role in adaptive immune response and immune cell signaling. Recent findings impressively demonstrate a variety of further biological functions as cellular adhesion, innate immune recognition, osteoclast maturation, platelet activation and vascular development (Moscai, A. et al., Nat Rev Immunol, 10:387 - 402, 2010). Syk associates with a variety of receptors of immune cells (mast cells, B cells, macrophages and neutrophils) and non-immune cells (osteoclasts, breast cancer cells) and orchestrates various different cellular processes including cytokine production, bone resorption and phagocytosis. Due to the interaction with immunoreceptors and G-coupled receptors Syk not only functions as a protein kinase but also as a true protein adaptor and therefore became a central paradigm in immune cell signaling.

Immunoreceptor tyrosine activation motif (ITAM)-mediated signaling has emerged as a primary event in signaling pathways responsible for human pathologies. ITAM-mediated and hemITAM-mediated signaling is responsible for relaying activation

signals initiated at classical immune receptors such as T-cell receptors, B-cell receptors, Fc receptors in immune cells and at GPVI and FcγRIIa in platelets to downstream intracellular molecules such as Syk and ZAP-70 (Underhill, D. M and Goodridge, H. S., Trends Immunol., 28:66-73, 2007) but also furthermore with hemITAM-containing factors as CLEC7A and other C-type lectins.

The binding of a ligand to an ITAM-containing receptor triggers signaling events which allows for the recruitment of proteins from a family of nonreceptor tyrosine kinases called the Src family. These kinases phosphorylate tyrosine residues within the ITAM sequence, a region with which the tandem SH2 domains on either Syk or ZAP-70 interact.

Syk, along with Zap-70, is a member of the Syk family of protein tyrosine kinases. The interaction of Syk or ZAP-70 with diphosphorylated ITAM sequences induces a conformation change in the kinases that allows for tyrosine phosphorylation of the kinase itself. Phosphorylated Syk family members activate a multitude of downstream signaling pathway proteins which include Src homology 2 (SH2) domains. Direct binding partners of Syk are VAV family members, phospholipase C gamma (PLCγ, PLCγ 2), phosphoinositide 3-kinases (PI3Ks), SH2 domain-containing leukocyte protein family members (SLP-76 or SLP-65). Other signaling intermediates are p38, Janus kinase (JNK), RAS homologue (RHO) family, Ca⁺⁺, diacylglycerol DAG, TEC family, caspase-recruitment domain - B cell lymphoma 10 - mucosa-associated lymphoid tissue lymphoma translocation protein 1 (CARD-BCL-10-MALT1) complex, protein tyrosine kinase 2 (PYK2), nuclear factor of activated T cells (NFAT), protein kinase C (PKC), RAS guanyl-releasing protein (RASGRP), extracellular signal-regulated kinase (ERK), AKT, NLR family,

pyrin domain-containing 3 (NLRP3) inflammasome, NLR family and nuclear factor kappaB (NFkappaB) and factors in the canonical and non-canonical signaling pathways. These contribute to a variety of cellular responses as cytoskeletal changes, ROS production, differentiation, proliferation, survival of cells and cytokine release.

Syk as a key mediator of immunoreceptor and non immunoreceptor signaling in a host of inflammatory cells is identified as a key player in the pathogenesis of a variety of diseases and disorders attributed to dysfunctional signaling including autoimmune diseases such as rheumatoid arthritis, systemic lupus, multiple sclerosis, hemolytic anemia, immune-thrombocytopenia purpura, and heparin-induced thrombocytopenia, functional gastrointestinal disorders, asthma, allergic disorders, anaphylactic shock and arteriosclerosis (Riccaboni, M. et al., DDT, 15:517 - 529, 2010). Interestingly, many of the above mentioned diseases are thought to occur through crosslinking of Fc receptors by antibodies which, via Syk, activate a signaling cascade in mast, basophil and other immune cells that result in the release of cell mediators responsible for inflammatory reactions. The release of mediators and the production of cytokines in IgE stimulation-dependent allergic and inflammatory reactions from mast cells and basophiles can be controlled by inhibiting the kinase activity of Syk (Rossi, A. B. et al., J Allergy Clin Immunol., 118:749-755, 2006). In immune-thrombocytopenia, antibody bound platelets are cleared by the spleen by an Fc receptor/ITAM/Syk-mediated process (Crow, A. R. et al., Blood, 106:abstract 2165, 2005). Drug-induced thrombocytopenia, caused by heparin-platelet factor 4 immune complexes that activate platelet FcgammaRIIa, also

involve Syk signaling downstream of receptor engagement (Reilly, M. P., Blood, 98:2442-2447, 2001).

Syk has also been shown to mediate signaling by classes of receptors that do not contain conventional ITAM motifs as integrins and lectins (Kerrigan, A. M. et al., Immunol. Rev., 234:335-352, 2010). Furthermore Syk plays an important role in pathogen recognition like fungi, bacteria and viruses (Hughes, C. E., et al., Blood, 115:2947-2955, 2010; Geijtenbeek, T. B. et al., Nat Rev Immunol, 9:465-479, 2009). The mechanism of Syk activation by integrins-mediated Platelet agonists induce inside-out integrin signaling resulting in fibrinogen binding and platelet aggregation. This initiates outside-in signaling which produces further stimulation of platelets. Syk is activated during both phases of integrin signaling, and inhibition of Syk is shown to inhibit platelet adhesion to immobilized proteins (Law, D. A. et al., Blood, 93:2645-2652, 1999). Release of arachidonic acid and serotonin and platelet aggregation induced by collagen are markedly inhibited in platelets derived from Syk deficient mouse (Poole, A. et al., EMBO J., 16:2333-2341, 1997). Thus Syk inhibitors may also possess anticoagulation action.

Because of the role Syk plays in Ig-induced platelet activations, it is of interest in arteriosclerosis and restenosis. Arteriosclerosis is a class of diseases characterized by the thickening and hardening of the arterial walls of blood vessels. Although all blood vessels are susceptible to this serious degenerative condition, the aorta and the coronary arteries serving the heart are most often affected. Arteriosclerosis is of profound clinical importance since it can increase the risk of heart attacks, myocardial infarctions, strokes, and aneurysms.

The traditional treatment for arteriosclerosis includes vascular recanalization procedures for less-serious blockages and coronary bypass surgery for major blockages. A serious shortcoming of intravascular procedures is that, in a significant number of treated individuals, some or all of the treated vessels restenose (i.e., re-narrow). While the exact hormonal and cellular processes promoting restenosis have not been determined, restenosis is thought to be due in part to mechanical injury to the walls of the blood vessels caused by the balloon catheter or other intravascular device. In response to this injury, adhering platelets, infiltrating macrophages, leukocytes, or the smooth muscle cells themselves release cell-derived growth factors such as platelet-derived growth factor (PDGF), with subsequent proliferation and migration of medial smooth muscle cells (SMCs) through the internal elastic lamina to the area of the vessel intima. Further proliferation and hyperplasia of intimal SMCs and, most significantly, production of large amounts of extracellular matrix over a period of three to six months results in the filling in and narrowing of the vascular space sufficient to significantly obstruct blood flow.

In addition to the role Syk plays in Ig-induced platelet activations, Syk plays a very important role in collagen-mediated signaling. The primary adhesive protein responsible for platelet adhesion and activation is collagen. Collagen is a filamentous protein contained within the fibrotic caps of atheromas which becomes exposed to blood during plaque rupture. Collagen functions initially by binding von Willebrand factor which tethers platelets through binding platelet membrane GPIb. Collagen functions secondarily by engaging the two collagen receptors on platelets, GPVI and integrin $\alpha_2\beta_1$.

GPVI exists in platelet membranes as a complex with FcRgamma, an interaction required for the expression of GPVI. Activation of FcgammaRIIa on platelets results in platelet shape change, secretion and thrombosis. Signaling by the GPVI/FcRgamma complex is initiated by tyrosine phosphorylation of the ITAM domain of FcRgamma followed by the recruitment of Syk. Activation of GPVI leads to induction of multiple platelet functions including: activation of integrins alpha2beta1 to achieve firm platelet adhesion, and GP IIb-IIIa which mediates platelet aggregation and thrombosis growth; platelet secretion, allowing for the delivery of inflammatory proteins such as CD40L, RANTES and TGFbeta to the vessel wall; and the expression of P-selectin which allows for the recruitment of leukocytes. Therefore, it is believed that Syk inhibitors can inhibit thrombotic events mediated by platelet adhesion, activation and aggregation.

It has been reported that the tyrosine phosphorylation of intracellular protein (activation) induced by stimulation of a receptor for IgG antibody, FcgammaR, and the phagocytosis mediated by FcgammaR are considerably inhibited in macrophages derived from Syk deficient mouse (Crowley, M. T. et al., J. Exp. Med., 186:1027-1039, 1997). This suggests that Syk has a markedly important role in the FcgammaR-mediated phagocytosis of macrophages.

It has also been reported that an antisense oligonucleotide of Syk suppresses the apoptosis inhibition of eosinophils induced by GM-CSF (Yousefi, S. et al., J. E. Med., 183:1407-1414, 1996), showing that Syk is essential for the life extending signal of eosinophils caused by GM-CSF and the like. Since life extension of eosinophils is closely related to the transition of diseases into a chronic state in allergic

disorders, such as asthma, Syk inhibitors can also serve as therapeutic agents for chronic eosinophilic inflammation.

Syk is important for the activation of B-cells via a B-cell antigen receptor and is involved in the phosphatidylinositol metabolism and increase in the intracellular calcium concentration caused by the antigen receptor stimulation (Hutchcroft, J E. et al., J. Biol. Chem., 267:8613-8619, 1992; and Takata, M. et al., EMBO J., 13:1341-1349, 1994). Thus, Syk inhibitors may be used to control the function of B-cells and are, therefore, expected to serve as therapeutic agents for antibody-related diseases.

Syk binds to a T-cell antigen receptor, quickly undergoes tyrosine phosphorylation through crosslinking of the receptor and synergistically acts upon intracellular signals mediated by Src tyrosine kinases such as Lck (Couture, C. et al., Proc. Natl. Acad. Sci. USA, 91:5301-5305, 1994; and Couture, C. et al., Mol. Cell. Biol., 14:5249-5258, 1994). Syk is present in mature T-cell populations, such as intraepithelial gammadelta T-cells and naive alphabeta T-cells, and has been reported to be capable of phosphorylation of multiple components of the TCR signaling cascade (Latour, S. et. al., Mol Cell Biol., 17:4434-4441, 1997). As a consequence, Syk inhibitors may serve as agents for inhibiting cellular immunity mediated by T-cell antigen receptor.

Recent comparative genomic hybridization studies have identified Syk as another gene important in the pathogenesis of Mantle Cell Lymphoma (MCL) (Chen, R. et al. Journal of Clinical Oncology, 2007 ASCO Annual Meeting Proceedings (Post-Meeting Edition), Vol 25, No 18S (June 20 Supplement), 2007: 8056). MCL represents 5-10% of all non-Hodgkins lymphomas and

it is a difficult form of lymphoma to treat. It has the worst prognosis among the B cell lymphomas with median survival of three years. It has been reported that Syk is overexpressed in MCL (Rinaldi, A, et. al, Br. J. Haematol, 2006; 132:303-316) and that Syk mediates mTOR (mammalian target of Rapamycin) survival signals in follicular, mantel cell, Burkitt's, and diffuse large B-cell non-Hodgkin's lymphomas (Leseux, L., et. al, Blood, 2006; 108:4156-4162).

Several lines of evidence suggest that many B-cell lymphomas depend upon B-cell receptor (BCR)-mediated survival signals. BCR signaling induces receptor oligomerization and phosphorylation of Igalpha and beta immunoreceptor tyrosine-based activated motifs by SRC family kinases. ITAM phosphorylation results in the recruitment and activation of Syk that initiates downstream events and amplifies the original BCR signal. Given the role of tonic BCR signaling in normal B cell and Syk-dependent survival of non-Hodgkins lymphoma cell lines in vitro (Chen, L., et. al, Blood, 2006; 108:3428-3433), Syk inhibition is a promising rational treatment target for certain B-cell lymphomas and chronic lymphocytic leukemia (CLL) (Stefania Gobessi, Luca Laurenti, Pablo Longo, Laura Carsetti, Giuseppe Leone, Dimitar G. Efremov, Constitutive activation of the protein tyrosine kinase Syk in Chronic Lymphocytic Leukemia B-cells, Blood, 2007, 110, Abstract 1123). Recent data shows that administration of a multikinase inhibitor which inhibits Syk, may have significant clinical activity in CLL patients (Friedberg J W et al, Blood 2008; 112(11), Abstract 3).

The oncogenic potential of Syk has been described in a number of different settings. Clinically, Syk over-expression is reported in Mantle Cell Lymphoma (Rinaldi, A, et. al, Br. J.

Haematol., 2006; 132:303-316) and the TEL-Syk fusion protein (Translocated ETS Leukemia) generated by a chromosomal translocation (t(9;12)(q22;p12)) leads to increased Syk activity and is associated with myelodysplastic syndrome (Kuno, Y., et. al, Blood, 2001; 97:1050-1055). Leukemia is induced in mice by adoptively transferring bone marrow cells that express human TEL-Syk (Wossning, T., JEM, 2006; 203:2829-2840). Further, in mouse primary bone marrow cells, over-expression of Syk results in IL-7 independent growth in culture (Wossning, T., et. al, JEM, 2006; 203:2829-2840).

Interestingly, Syk signaling appears to be required for B-cell development and survival in humans and mouse. Inducible loss of the B-cell receptor (Lam, K., et. al, Cell, 1997; 90:1073-1083) or Igalpha (Kraus, M., et. al, Cell, 2004; 117:787-800) results in loss of peripheral B-cells in mice. Over-expression of the protein tyrosine phosphatase PTP-RO, which is known to negatively regulate Syk activity, inhibits proliferation and induces apoptosis in cell lines derived from non-Hodgkin's lymphomas (Chen, L., et. al, Blood, 2006; 108:3428-3433). Finally, B-cell lymphomas rarely exhibit loss of BCR expression, and anti-idiotypic therapy rarely leads to resistance (Kuppers, R. Nat Rev Cancer, 2005; 5:251-262).

Engagement of the antigen-specific B cell receptor (BCR) activates multiple signaling pathways that ultimately regulate the cells activation status, promoting survival and clonal expansion. Signaling through the BCR is made possible by its association with two other members of the immunoglobulin super-family; Igalpha and Igbeta, each bearing an immunotyrosine based activation motif (ITAM) (Jumaa, Hendriks et al. Annu Rev Immunol 23: 415-45 (2005)). The ITAM domain is directly phosphorylated by Src family kinases in response to

BCR engagement. Syk docks with and phosphorylates the ITAM, a process that enhances its kinase activity, resulting in Syk autophosphorylation and tyrosine phosphorylation of multiple downstream substrates (Rolli, Gallwitz et al. Mol Cell 10(5): 1057-69 (2002). This signaling pathway is active in B cells beginning at the transition from pro- to pre-B cell stage of development, when the newly formed pre-BCR is expressed. In fact, B cell development arrests at the pro-B cell stage in Syk knockout mice (Cheng, Rowley et al. 1995; Turner, Mee et al. Nature 378(6554): 303-6 (1995). Inducible loss of the B cell receptor (Lam, Kuhn et al. Cell 90(6): 1073-83 (1997) or Igalpha (Kraus, Alimzhanov et al. Cell 117(6): 787-800 (2004) results in loss of peripheral B cells in mice. Human B cells also appear to require Syk for proliferation and survival. Over-expression of the protein tyrosine phosphatase PTP-RO, a negative regulator of Syk activity, inhibits proliferation and induces apoptosis in cell lines derived from non-Hodgkin's lymphomas (NHL) (Chen, Juszczynski et al. Blood 108(10): 3428-33 (2006). Knock down of Syk by siRNA in the NHL line SUDHL-4 led to a block in the G1/S transition of the cell cycle (Gururaj an, Dasu et al. J Immunol 178(1): 111-21 (2007). Together, these data suggest that Syk signaling is required for the development, proliferation, and even survival of human and mouse B cells.

Conversely, the oncogenic potential of Syk has been described in a number of different settings. Clinically, Syk over-expression is reported in Mantle Cell Lymphoma (Rinaldi, Kwee et al. Br J Haematol 132(3): 303-16 (2006) and the TEL-Syk fusion protein (Translocated ETS Leukemia) generated by a chromosomal translocation (t(9;12)(q22;p12)) leads to increased Syk activity and is associated with myelodysplastic syndrome (Kuno, Abe et al. Blood 97(4): 1050-5 (2001).

Leukemia is induced in mice by the adoptive transfer of bone marrow cells that express human TEL-Syk (Wossning, Herzog et al. J Exp Med 203(13): 2829-40 (2006). Further, in mouse primary bone marrow cells, over-expression of Syk results in IL-7 independent growth in culture (Wossning, Herzog et al. 2006). Consistently, Syk was reported to mediate mTOR (mammalian target of Rapamycin) survival signals in follicular, mantle cell, Burkitt's, and diffuse large B-cell NHL (Leseux, Hamdi et al. Blood 108(13): 4156-62 (2006). Additional recent studies also suggest that Syk-dependant survival signals may play a role in B-cell malignancies, including DLBCL, mantle cell lymphoma and follicular lymphoma (Gururajan, Jennings et al. 2006; Irish, Czerwinski et al. J Immunol 176(10): 5715-9 (2006)). Given the role of tonic BCR signaling in normal B cells and Syk-dependent survival of NHL cell lines in vitro, the specific inhibition of Syk may prove promising for the treatment of certain B-cell lymphomas.

Recently, R406 (Rigel Pharmaceuticals) was reported to inhibit ITAM signaling in response to various stimuli, including FcepsilonR1 and BCR induced Syk activation (Brasemann, Taylor et al. J Pharmacol Exp Ther 319(3): 998-1008 (2006). Interestingly, this ATP-competitive inhibitor of Syk was also active against Flt3, cKit, and JAK kinases, but not against Src kinase (Brasemann, Taylor et al. 2006). Activating mutations to Flt3 are associated with AML and inhibition of this kinase is currently under clinical development (Burnett and Knapper Hematology Am Soc Hematol Educ Program 2007: 429-34 (2007). Over-activation of the tyrosine kinase cKit is also associated with hematologic malignancies, and a target for cancer therapy (Heinrich, Griffith et al. Blood 96(3): 925-32 (2000). Similarly, JAK3 signaling is implicated in leukemias and lymphomas, and is currently exploited as a potential

therapeutic target (Heinrich, Griffith et al. 2000). Importantly, the multi-kinase inhibitory activity of R406 attenuates BCR signaling in lymphoma cell lines and primary human lymphoma samples, resulting in apoptosis of the former (Chen, Monti et al. Blood 111(4): 2230-7 (2008). Further, a phase II clinical trial reported favorable results by this compound in refractory NHL and chronic lymphocytic leukemia (Friedberg J W et al, Blood 2008; 112(11), Abstract 3). Although the precise mechanism of action is unclear for R406, the data suggest that inhibition of kinases that mediate survival signaling in lymphocytes is clinically beneficial.

Additional recent studies also suggest that Syk-dependant survival signals may play a role in B-cell malignancies, including DLBCL, mantle cell lymphoma and follicular lymphoma (see e.g., S. Linfengshen et al. Blood, February 2008; 111: 2230-2237; J. M. Irish et al. Blood, 2006; 108: 3135-3142; A. Renaldi et al. Brit J. Haematology, 2006; 132: 303-316; M. Guruoajan et al. J. Immunol, 2006; 176: 5715-5719; L. Laseux et al. Blood, 2006; 108: 4156-4162.

A recent publication summarizes the frequent finding of eye involvement with rheumatoid arthritis and other autoimmune diseases. Scleritis, episcleritis and keratoconjunctivitis sicca may represent the leading clinical manifestation of these autoimmune diseases. All components of the visual organ might be affected. Autoimmune reactions based on the patient's genetic predisposition are assumed to be of significance in pathogenesis of eye diseases.

This manifests Syk as relevant therapeutic target in ocular diseases. (Feist, E., Pleyer, U., Z Rheumatol, 69: 403 - 410, 2010).

Furthermore SYK is also a relevant target in the treatment of fungal, viral and bacterial infections of the eye e.g. fungal keratitis. Dectin-1 mediated activation of p-Syk, and further factors as p-IkB or NFkB lead to the production of IL-1b and CXCL1/KC that are important for neutrophil and mononuclear cell recruitment to the corneal stroma. Leal, S. M., Cowden, S., Hsia, Y.-C., Ghannoum, M. A., Momany, M., & Pearlman, E. (2010). Distinct roles for Dectin-1 and TLR4 in the pathogenesis of *Aspergillus fumigatus* keratitis. *PLOS Pathogens*, 6.

In general recent evidence shows that SYK is a essential target for treatment of PRR and CLR mediated adaptive immune response. Kingeter, L. M., & Lin, X. (2012). C-type lectin receptor-induced NF- κ B activation in innate immune and inflammatory responses. *Cellular & molecular immunology*, 9(2), 105-112. Drummond, R. A., Saijo, S., Iwakura, Y., & Brown, G. D. (2011). The role of Syk/CARD9 coupled C-type lectins in antifungal immunity. *European journal of immunology*, 41(2), 276-281. Lee, H.-M., Yuk, J.-M., Kim, K.-H., Jang, J., Kang, G., Park, J. B., Son, J.-W., et al. (2011). *Mycobacterium abscessus* activates the NLRP3 inflammasome via Dectin-1-Syk and p62/SQSTM1. *Immunology and cell biology*.

According to one embodiment, the present invention provides compounds that are capable of inhibiting one or more kinases, more particularly SYK and mutants thereof.

LRRK2 - Leucine-rich repeat kinase 2

There has been much interest raised by the discovery that different autosomal dominant point mutations within the gene

encoding for LRRK2 predispose humans to develop late-onset Parkinson's disease (PD), with a clinical appearance indistinguishable from idiopathic PD (see Paisan-Ruiz, C., Jain, S., Evans, E. W., Gilks, W. P., Simon, J., van der Brug, M., Lopez de Munain, A., Aparicio, S., Gille A. M., Khan, N., Johnson, J., Martinez, J. R., Nicholl, D., Carrera, I. M., Pena, A. S., de Silva, R., Lees, A., Marti-Masso, J. F., Perez-Tur, J., Wood, N. W. and Singleton, A. B. (2004) Cloning of the gene containing mutations that cause PARK8-linked Parkinson's disease. *Neuron*. 44, 595-600; Mata, I. F., Wedemeyer, W. J., Farrer, M. J., Taylor, J. P. and Gallo, K. A. (2006) LRRK2 in Parkinson's disease: protein domains and functional insights. *Trends Neurosci*. 29, 286-293; Taylor, J. P., Mata, I. F. and Farrer, M. J. (2006) LRRK2: a common pathway for parkinsonism, pathogenesis and prevention? *Trends Mol Med*. 12, 76-82). The genetic analysis undertaken to date indicates that mutations in LRRK2 are relatively frequent, not only accounting for 5-10% of familial PD, but also being found in a significant proportion of sporadic PD cases (see Farrer, M., Stone, J., Mata, I. F., Lincoln, S., Kachergus, J., Hulihan, M., Strain, K. J. and Maraganore, D. M. (2005) LRRK2 mutations in Parkinson disease. *Neurology*. 65, 738-740; Zabetian, C. P., Samii, A., Mosley, A. D., Roberts, J. W., Leis, B. C, Yearout, D., Raskind, W. H. and Griffith, A. (2005) A clinic-based study of the LRRK2 gene in Parkinson disease yields new mutations. *Neurology*. 65, 741-744. Little is known about how LRRK2 is regulated in cells, what its physiological substrates are and how mutations cause or increase risk of PD.

Genomewide-wide association studies show a possible involvement in further neurodegenerative diseases like Alzheimer furthermore leprosy but also revealed a higher probability of cancer occurrence for carriers of LRRK2 mutants

and may indicate an involvement of this kinase and mutants in cancer development. Inzelberg, et al. The LRRK2 G2019S mutation is associated with Parkinson disease and concomitant non-skin cancers. *Neurology*, 2012, 78, 781-786. Zhao, Y., Ho, P., Yih, Y., Chen, C., Lee, W. L., & Tan, E. K. (2011). LRRK2 variant associated with Alzheimer's disease. *Neurobiology of aging*, 32(11), 1990-1993. Lewis, P. A., & Manzoni, C. (2012). LRRK2 and Human Disease: A Complicated Question or a Question of Complexes? *Science Signaling*, 5(207).

An unexpected finding was the involvement of LRRK2 as a major susceptibility gene for Crohn's disease (CD) and other related inflammatory diseases. LRRK2 deficiency in mice confers enhanced susceptibility to experimental colitis. The complex nature of the multidomain LRRK2 protein makes it plausible that LRRK2 may also regulate different pathways in immune reactions through its involvement in NFAT1 regulation in participating in the NRON complex in immune cells. Liu, Z., & Lenardo, M. J. (2012) "The role of LRRK2 in inflammatory bowel disease", *Cell research*; "LRRK2 as a negative regulator of NFAT: implications for the pathogenesis of inflammatory bowel disease" Puja Vora, Dermot PB McGovern. *Expert Review of Clinical Immunology*, Mar 2012, Vol. 8, No. 3, Pages 227-229.

According to one embodiment, the present invention provides compounds that are capable of inhibiting one or more kinases, more particularly, LRRK, even more preferably LRRK2.

Myosin Light Chain Kinase (MLCK or MYLK)

Inhibitors of MYLK (or MLCK) are of interest in the treatment and/or prevention of any disorder where tissue barrier dysfunction or changes in cell motility are part of the

disease mechanism or progression of pathophysiology. These include a large number of diseases in a variety of categories, including but not limited to skin disorders: including ichthyosis vulgaris, atopic dermatitis, psoriasis, eczema, allergic skin disease, and hypersensitivity reactions; intestinal disorders: including inflammatory bowel disease, Crohn's disease, ulcers, bacterial infections hemorrhagic shock, diarrhea, colitis, viral and alcoholic liver disease, pancreatitis; lung disorders: including acute lung injury after infection, mechanical ventilation-induced injury, sepsis, thrombin-induced lung injury, lung injury after reperfusion; interstitial cystitis of the bladder; coronary disease after ischemia-reperfusion injury, flow-induced injury, aortic aneurysm, hypertension; burn-induced injury; chorioretinal vascular disease; neurologic disorders: including multiple sclerosis, Alzheimer's disease, vascular dementia, traumatic brain injury, ALS, Parkinson's disease, stroke, meningoencephalitis, cerebral hemorrhage, Guillain-Barre syndrome, vasogenic brain edema, hypoxia-induced injury and blood brain barrier compromise after ethanol toxicity; and cancers, including metastatic cancers such as non-small cell lung cancers, pancreatic cancer, adenocarcinoma and prostate cancer. See, e.g., Behanna H A, Watterson D M and Ralay Ranaivo H (2006) Development of a novel bioavailable inhibitor of the calmodulin-regulated protein kinase MLCK: a lead compound that attenuates vascular leak. *Biochim Biophys Acta* 1763: 1266-1274; Behanna H A, Bergan R and Watterson D M (2007), unpublished observations; Bratcher J M and Korelitz B I (2006) Toxicity of infliximab in the course of Crohn's disease. *Expert Opin Drug Saf* 5: 9-16; Clayburgh D R, Shen L and Turner J R (2004) A porous defense: the leaky epithelial barrier in intestinal disease. *Lab Invest* 84: 282-291; Clayburgh D R, Barrett T A, Tang Y, Meddings J B, Van Eldik L


J, Watterson D M, Clarke L L, Mrsny R J and Turner J R (2005) Epithelial myosin light chain kinase-dependent barrier dysfunction mediates T cell activation-induced diarrhea in vivo. *J Clin Invest* 115: 2702-2715; Demling R H (2005) The burn edema process: current concepts. *J Burn Care Rehabil* 26: 207-227; Dreyfuss D and Saumon G (1998) Ventilator-induced lung injury: lessons from experimental studies. *Am J Respir Crit. Care Med* 157: 294-323; Haorah J, Heilman D, Knipe B, Chrastil J, Leibhart J, Ghorpade A, Miller D W and Persidsky Y (2005) Ethanol-induced activation of myosin light chain kinase leads to dysfunction of tight junctions and blood-brain barrier compromise. *Alcohol Clin Exp Res* 29: 999-1009; Huang Q, Xu W, Ustinova E, Wu M, Childs E, Hunter F and Yuan S (2003) Myosin light chain kinase-dependent microvascular hyperpermeability in thermal injury. *Shock* 20: 363-368; Kaneko K, Satoh K, Masamune A, Satoh A and Shimosegawa T (2002) Myosin light chain kinase inhibitors can block invasion and adhesion of human pancreatic cancer cell lines. *Pancreas* 24: 34-41; Ma T Y, Boivin M A, Ye D, Pedram A and Said H M (2005) Mechanism of TNF α modulation of Caco-2 intestinal epithelial tight junction barrier: role of myosin light-chain kinase protein expression. *Am J Physiol Gastrointest Liver Physiol* 288: G422-G430; Minamiya Y, Nakagawa T, Saito H, Matsuzaki I, Taguchi K, Ito M and Ogawa J (2005) Increased expression of myosin light chain kinase mRNA is related to metastasis in non-small cell lung cancer. *Tumour Biol* 26: 153-157; Ralay Ranaivo H, Carusio N, Wangenstein R, Ohlmann P, Loichot C, Tesse A, Chalupsky K, Lobysheva I, Haiech J, Watterson D M and Andriantsitohaina R (2007) Protection against endotoxic shock as a consequence of reduced nitrosative stress in MLCK210-null mice. *Am J Pathol* 170:439-446; Reynoso R, Perrin R M, Breslin J W, Daines D A, Watson K D, Watterson D M, Wu M H and Yuan S A role for long chain

myosin light chain kinase (MLCK-210) in microvascular hyperpermeability during severe burns. Shock, June 14 epub; Rossi J, Bayram M, Udelson J E, Lloyd-Jones D, Adams K F, Oconnor C M, Stough W G, Ouyang J, Shin D D, Orlandi C and Gheorghiade M (2007) Improvement in hyponatremia during hospitalization for worsening heart failure is associated with improved outcomes: insights from the Acute and Chronic Therapeutic Impact of a Vasopressin Antagonist in Chronic Heart Failure (ACTIV in CHF) trial. Acute Card Care 9:82-86; Scott K G, Meddings J B, Kirk D R, Lees-Miller S P and Buret A G (2002) Intestinal infection with Giardia spp. reduces epithelial barrier function in a myosin light chain kinase-dependent fashion. Gastroenterology 123: 1179-1190; Tohtong R, Phattarasakul K, Jiraviriyakul A and Sutthiphongchai T (2003) Dependence of metastatic cancer cell invasion on MLCK-catalyzed phosphorylation of myosin regulatory light chain. Prostate Cancer Prostatic Dis 6: 212-216; Yuan S Y (2002) Protein kinase signaling in the modulation of microvascular permeability. Vascul Pharmacol 39: 213-223; Yuan S Y, Wu M H, Ustinova E E, Guo M, Tinsley J H, De Lanerolle P and Xu W (2002) Myosin light chain phosphorylation in neutrophil-stimulated coronary microvascular leakage. Circ Res 90: 1214-1221; Zolotarevsky Y, Hecht G, Koutsouris A, Gonzalez D E, Quan C, Tom J, Mrsny R J and Turner JR (2002) A membrane-permeant peptide that inhibits MLC kinase restores barrier function in in vitro models of intestinal disease. Gastroenterology 123 (2002) 163-172. Role of myosin light chain kinase in regulation of basal blood pressure and maintenance of salt-induced hypertension. (2011). Role of myosin light chain kinase in regulation of basal blood pressure and maintenance of salt-induced hypertension. American journal of physiology. Heart and circulatory physiology, 301(2).

According to one embodiment, the present invention provides compounds that are capable of inhibiting one or more kinases, especially MYLK (or MLCK).

STATEMENT OF INVENTION

The present invention provides one or more compounds of formula (I)



(I)

wherein

A is O, S, C=O, NR⁴ or CR⁵R⁶ (especially NH);

R¹ is an optionally substituted alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl group;

R² is an optionally substituted alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or hetero-aralkyl group, wherein R² is preferably bound to the pyrimidine ring of formula (I) via a carbon-carbon bond;

R³ is a hydrogen atom, a halogen atom, NO₂, N₃, OH, SH, NH₂ or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group;

R⁴ is a hydrogen atom or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group;

R⁵ is a hydrogen atom, NO₂, N₃, OH, SH, NH₂ or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group; and

R⁶ is a hydrogen atom, NO₂, N₃, OH, SH, NH₂ or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group;

or a pharmaceutically acceptable salt, ester, solvate or hydrate or a pharmaceutically acceptable formulation thereof.

The expression alkyl refers to a saturated, straight-chain or branched hydrocarbon group that contains from 1 to 20 carbon atoms, preferably from 1 to 12 carbon atoms, especially from 1 to 6 (e.g. 1, 2, 3 or 4) carbon atoms, for example a methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, tert-butyl, n-pentyl, iso-pentyl, n-hexyl, 2,2-dimethylbutyl or n-octyl group. Furthermore, the term alkyl refers to groups in which one or more hydrogen atoms have been replaced by a

halogen atom (preferably F or Cl) such as, for example, a 2,2,2-trichloroethyl or a trifluoromethyl group.

The expressions alkenyl and alkynyl refer to at least partially unsaturated, straight-chain or branched hydrocarbon groups that contain from 2 to 20 carbon atoms, preferably from 2 to 12 carbon atoms, especially from 2 to 6 (e.g. 2, 3 or 4) carbon atoms, for example an ethenyl (vinyl), propenyl (allyl), iso-propenyl, butenyl, ethinyl, propinyl, butinyl, acetylenyl, propargyl, isoprenyl or hex-2-enyl group. Preferably, alkenyl groups have one or two (especially preferably one) double bond(s), and alkynyl groups have one or two (especially preferably one) triple bond(s). Furthermore, the terms alkenyl and alkynyl refer to groups in which one or more hydrogen atoms have been replaced by a halogen atom (preferably F or Cl).

The expression heteroalkyl refers to an alkyl, alkenyl or alkynyl group in which one or more (preferably 1, 2 or 3) carbon atoms have been replaced by an oxygen, nitrogen, phosphorus, boron, selenium, silicon or sulfur atom (preferably by an oxygen, sulfur or nitrogen atom). The expression heteroalkyl furthermore refers to a carboxylic acid or to a group derived from a carboxylic acid, such as, for example, acyl, acylalkyl, alkoxycarbonyl, acyloxy, acyloxyalkyl, carboxyalkylamide or alkoxycarbonyloxy.

Preferably, a heteroalkyl group contains from 1 to 12 carbon atoms and from 1 to 4 hetero atoms selected from oxygen, nitrogen and sulphur (especially oxygen and nitrogen). Especially preferably, a heteroalkyl group contains from 1 to 6 (e.g. 1, 2, 3 or 4) carbon atoms and 1, 2 or 3 (especially 1 or 2) hetero atoms selected from oxygen, nitrogen and sulphur

(especially oxygen and nitrogen). The term C₁-C₆ heteroalkyl refers to a heteroalkyl group containing from 1 to 6 carbon atoms and 1, 2 or 3 heteroatoms selected from O, S and/or N (especially O and/or N). The term C₁-C₄ heteroalkyl refers to a heteroalkyl group containing from 1 to 4 carbon atoms and 1, 2 or 3 heteroatoms selected from O, S and/or N (especially O and/or N). Furthermore, the term heteroalkyl refers to groups in which one or more hydrogen atoms have been replaced by a halogen atom (preferably F or Cl).

Examples of heteroalkyl groups are groups of formulae:
 R^a-O-Y^a-, R^a-S-Y^a-, R^a-N(R^b)-Y^a-, R^a-CO-Y^a-, R^a-O-CO-Y^a-,
 R^a-CO-O-Y^a-, R^a-CO-N(R^b)-Y^a-, R^a-N(R^b)-CO-Y^a-, R^a-O-CO-N(R^b)-Y^a-,
 R^a-N(R^b)-CO-O-Y^a-, R^a-N(R^b)-CO-N(R^c)-Y^a-, R^a-O-CO-O-Y^a-,
 R^a-N(R^b)-C(=NR^d)-N(R^c)-Y^a-, R^a-CS-Y^a-, R^a-O-CS-Y^a-, R^a-CS-O-Y^a-,
 R^a-CS-N(R^b)-Y^a-, R^a-N(R^b)-CS-Y^a-, R^a-O-CS-N(R^b)-Y^a-,
 R^a-N(R^b)-CS-O-Y^a-, R^a-N(R^b)-CS-N(R^c)-Y^a-, R^a-O-CS-O-Y^a-,
 R^a-S-CO-Y^a-, R^a-CO-S-Y^a-, R^a-S-CO-N(R^b)-Y^a-, R^a-N(R^b)-CO-S-Y^a-,
 R^a-S-CO-O-Y^a-, R^a-O-CO-S-Y^a-, R^a-S-CO-S-Y^a-, R^a-S-CS-Y^a-,
 R^a-CS-S-Y^a-, R^a-S-CS-N(R^b)-Y^a-, R^a-N(R^b)-CS-S-Y^a-, R^a-S-CS-O-Y^a-,
 R^a-O-CS-S-Y^a-, wherein R^a being a hydrogen atom, a C₁-C₆ alkyl, a C₂-C₆ alkenyl or a C₂-C₆ alkynyl group; R^b being a hydrogen atom, a C₁-C₆ alkyl, a C₂-C₆ alkenyl or a C₂-C₆ alkynyl group; R^c being a hydrogen atom, a C₁-C₆ alkyl, a C₂-C₆ alkenyl or a C₂-C₆ alkynyl group; R^d being a hydrogen atom, a C₁-C₆ alkyl, a C₂-C₆ alkenyl or a C₂-C₆ alkynyl group and Y^a being a direct bond, a C₁-C₆ alkylene, a C₂-C₆ alkenylene or a C₂-C₆ alkynylene group, wherein each heteroalkyl group contains at least one carbon atom and one or more hydrogen atoms may be replaced by fluorine or chlorine atoms.

Specific examples of heteroalkyl groups are methoxy, trifluoromethoxy, ethoxy, n-propyloxy, isopropyloxy, butoxy,

tert-butyloxy, methoxymethyl, ethoxymethyl, $-\text{CH}_2\text{CH}_2\text{OH}$, $-\text{CH}_2\text{OH}$, methoxyethyl, 1-methoxyethyl, 1-ethoxyethyl, 2-methoxyethyl or 2-ethoxyethyl, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino, diethylamino, isopropylethylamino, methylamino methyl, ethylamino methyl, diisopropylamino ethyl, methylthio, ethylthio, isopropylthio, enol ether, dimethylamino methyl, dimethylamino ethyl, acetyl, propionyl, butyryloxy, acetyloxy, methoxycarbonyl, ethoxycarbonyl, propionyloxy, acetylamino or propionylamino, carboxymethyl, carboxyethyl or carboxypropyl, N-ethyl-N-methylcarbamoyl or N-methylcarbamoyl. Further examples of heteroalkyl groups are nitrile, isonitrile, cyanate, thiocyanate, isocyanate, isothiocyanate and alkyl nitrile groups.

The expression cycloalkyl refers to a saturated or partially unsaturated (for example, a cycloalkenyl group) cyclic group that contains one or more rings (preferably 1 or 2), and contains from 3 to 14 ring carbon atoms, preferably from 3 to 10 (especially 3, 4, 5, 6 or 7) ring carbon atoms. The expression cycloalkyl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH_2 , =NH, N_3 or NO_2 groups, thus, for example, cyclic ketones such as, for example, cyclohexanone, 2-cyclohexenone or cyclopentanone. Further specific examples of cycloalkyl groups are a cyclopropyl, cyclobutyl, cyclopentyl, spiro[4,5]decanyl, norbornyl, cyclohexyl, cyclopentenyl, cyclohexadienyl, decalanyl, bicyclo[4.3.0]nonyl, tetraline, cyclopentylcyclohexyl, fluorocyclohexyl or cyclohex-2-enyl group.

The expression heterocycloalkyl refers to a cycloalkyl group as defined above in which one or more (preferably 1, 2 or 3)

ring carbon atoms have been replaced by an oxygen, nitrogen, silicon, selenium, phosphorus or sulfur atom (preferably by an oxygen, sulfur or nitrogen atom). A heterocycloalkyl group has preferably 1 or 2 ring(s) containing from 3 to 10 (especially 3, 4, 5, 6 or 7) ring atoms (preferably selected from C, O, N and S). The expression heterocycloalkyl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups. Examples are a piperidyl, prolinyl, imidazolidinyl, piperazinyl, morpholinyl, urotropinyl, pyrrolidinyl, tetrahydrothiophenyl, tetrahydropyranyl, tetrahydrofuryl or 2-pyrazolinyl group and also lactames, lactones, cyclic imides and cyclic anhydrides.

The expression alkylcycloalkyl refers to groups that contain both cycloalkyl and also alkyl, alkenyl or alkynyl groups in accordance with the above definitions, for example alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkenyl, alkenylcycloalkyl and alkynylcycloalkyl groups. An alkylcycloalkyl group preferably contains a cycloalkyl group that contains one or two rings having from 3 to 10 (especially 3, 4, 5, 6 or 7) ring carbon atoms, and one or two alkyl, alkenyl or alkynyl groups (especially alkyl groups) having 1 or 2 to 6 carbon atoms.

The expression heteroalkylcycloalkyl refers to alkylcycloalkyl groups as defined above in which one or more (preferably 1, 2 or 3) carbon atoms have been replaced by an oxygen, nitrogen, silicon, selenium, phosphorus or sulfur atom (preferably by an oxygen, sulfur or nitrogen atom). A heteroalkylcycloalkyl group preferably contains 1 or 2 rings having from 3 to 10 (especially 3, 4, 5, 6 or 7) ring atoms, and one or two alkyl, alkenyl, alkynyl or heteroalkyl groups (especially alkyl or

heteroalkyl groups) having from 1 or 2 to 6 carbon atoms. Examples of such groups are alkylheterocycloalkyl, alkylheterocycloalkenyl, alkenylheterocycloalkyl, alkynylheterocycloalkyl, heteroalkylcycloalkyl, heteroalkylheterocycloalkyl and heteroalkylheterocycloalkenyl, the cyclic groups being saturated or mono-, di- or tri-unsaturated.

The expression aryl refers to an aromatic group that contains one or more rings containing from 6 to 14 ring carbon atoms, preferably from 6 to 10 (especially 6) ring carbon atoms. The expression aryl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, SH, NH₂, N₃ or NO₂ groups. Examples are the phenyl, naphthyl, biphenyl, 2-fluorophenyl, anilinyll, 3-nitrophenyl or 4-hydroxyphenyl group.

The expression heteroaryl refers to an aromatic group that contains one or more rings containing from 5 to 14 ring atoms, preferably from 5 to 10 (especially 5 or 6 or 9 or 10) ring atoms, and contains one or more (preferably 1, 2, 3 or 4) oxygen, nitrogen, phosphorus or sulfur ring atoms (preferably O, S or N). The expression heteroaryl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, SH, N₃, NH₂ or NO₂ groups. Examples are pyridyl (e.g. 4-pyridyl), imidazolyl (e.g. 2-imidazolyl), phenylpyrrolyl (e.g. 3-phenylpyrrolyl), thiazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxadiazolyl, thiadiazolyl, indolyl, indazolyl, tetrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, isoxazolyl, triazolyl, tetrazolyl, isoxazolyl, indazolyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzthiazolyl, pyridazinyl, quinolinyl, isoquinolinyl, pyrrolyl, purinyl, carbazolyl, acridinyl, pyrimidyl, 2,3'-

bifuryl, pyrazolyl (e.g. 3-pyrazolyl) and isoquinolinyl groups.

The expression aralkyl refers to groups containing both aryl and also alkyl, alkenyl, alkynyl and/or cycloalkyl groups in accordance with the above definitions, such as, for example, arylalkyl, arylalkenyl, arylalkynyl, arylcycloalkyl, arylcycloalkenyl, alkylaryl cycloalkyl and alkylaryl cycloalkenyl groups. Specific examples of aralkyls are toluene, xylene, mesitylene, styrene, benzyl chloride, o-fluorotoluene, 1H-indene, tetraline, dihydronaphthalene, indanone, phenylcyclopentyl, cumene, cyclohexylphenyl, fluorene and indane. An aralkyl group preferably contains one or two aromatic ring systems (1 or 2 rings) containing from 6 to 10 carbon atoms and one or two alkyl, alkenyl and/or alkynyl groups containing from 1 or 2 to 6 carbon atoms and/or a cycloalkyl group containing 5 or 6 ring carbon atoms.

The expression heteroaralkyl refers to an aralkyl group as defined above in which one or more (preferably 1, 2, 3 or 4) carbon atoms have been replaced by an oxygen, nitrogen, silicon, selenium, phosphorus, boron or sulfur atom (preferably oxygen, sulfur or nitrogen), that is to say to groups containing both aryl or heteroaryl, respectively, and also alkyl, alkenyl, alkynyl and/or heteroalkyl and/or cycloalkyl and/or heterocycloalkyl groups in accordance with the above definitions. A heteroaralkyl group preferably contains one or two aromatic ring systems (1 or 2 rings) containing from 5 or 6 to 10 ring carbon atoms and one or two alkyl, alkenyl and/or alkynyl groups containing 1 or 2 to 6 carbon atoms and/or a cycloalkyl group containing 5 or 6 ring carbon atoms, wherein 1, 2, 3 or 4 of these carbon atoms have been replaced by oxygen, sulfur or nitrogen atoms.

Examples are arylheteroalkyl, arylheterocycloalkyl, arylheterocycloalkenyl, arylalkylheterocycloalkyl, arylalkenylheterocycloalkyl, arylalkynylheterocycloalkyl, arylalkylheterocycloalkenyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, heteroarylheteroalkyl, heteroarylcycloalkyl, heteroarylcycloalkenyl, heteroarylheterocycloalkyl, heteroarylheterocycloalkenyl, heteroarylalkylcycloalkyl, heteroarylalkylheterocycloalkenyl, heteroarylheteroalkylcycloalkyl, heteroarylheteroalkylcycloalkenyl and heteroarylheteroalkylheterocycloalkyl groups, the cyclic groups being saturated or mono-, di- or tri-unsaturated. Specific examples are a tetrahydroisoquinolinyl, benzoyl, 2- or 3-ethylindolyl, 4-methylpyridino, 2-, 3- or 4-methoxyphenyl, 4-ethoxyphenyl, 2-, 3- or 4-carboxyphenylalkyl group.

The expression "optionally substituted" especially refers to groups in which one, two, three or more hydrogen atoms may have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups. This expression refers furthermore to groups that may be substituted by one, two, three or more unsubstituted C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ heteroalkyl, C₃-C₁₈ cycloalkyl, C₂-C₁₇ heterocycloalkyl, C₄-C₂₀ alkylcycloalkyl, C₂-C₁₉ heteroalkylcycloalkyl, C₆-C₁₈ aryl, C₁-C₁₇ heteroaryl, C₇-C₂₀ aralkyl or C₂-C₁₉ heteroaralkyl groups. This expression refers furthermore especially to groups that may be substituted by one, two, three or more unsubstituted C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ heteroalkyl, C₃-C₁₀ cycloalkyl, C₂-C₉ heterocycloalkyl, C₇-C₁₂ alkylcycloalkyl, C₂-C₁₁ heteroalkylcycloalkyl, C₆-C₁₀ aryl, C₁-C₉ heteroaryl, C₇-C₁₂ aralkyl or C₂-C₁₁ heteroaralkyl groups.

Preferred substituents are F, Cl, Br, OH, =O, NH₂, C₁₋₄ alkyl (e.g. methyl, ethyl, t-butyl), NMe₂, CONH₂, CH₂NMe₂, NHSO₂Me, C(CH₃)₂CN, COMe, OMe, SMe, COOMe, COOEt, CH₂COOH, OCH₂COOH, COOH, SMe, SO₂Me, cyclopropyl, SO₂NH₂, SO₂NHMe, SO₂CH₂CH₂OH, SF₅, SO₂NMe₂, OCF₃, SO₂CF₃, COMe, CN or CF₃.

Especially preferred substituents are F, Cl, Br, OH, NH₂, Me, Ethyl, NMe₂, CONH₂, OMe, CN or CF₃.

According to a preferred embodiment, all alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aralkyl and heteroaralkyl groups described herein may optionally be substituted.

When an aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group contains more than one ring, these rings may be bonded to each other via a single or double bond or these rings may be annulated.

Further preferred are compounds of formula (I) or (Ia) wherein R¹ is an optionally substituted aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aralkyl or heteroaralkyl group.

Moreover preferred are compounds of formula (I) or (Ia) wherein R¹ is an optionally substituted phenyl or naphthyl group or an optionally substituted heteroaryl group having one or two rings containing 5, 6, 7, 8, 9 or 10 ring atoms, or an optionally substituted arylheterocycloalkyl or heteroaryl-heterocycloalkyl group containing two or three (especially


two) rings and from 9 to 20 (especially 9 or 10) ring atoms. Preferably, the heteroatoms are selected from S, O and N, especially from N and O. Further preferably, the number of heteroatoms is from 1 to 6 (especially 1, 2, 3 or 4).

Preferred are compounds of formula (I) wherein A is NH.

Further preferred are compounds of formula (I) wherein R³ is H, F, Cl, CH₃, CF₃, NO₂, cyclopropyl, CN, N₃, OH, SH, OMe, SMe, NHMe, NMe₂ or NH₂.

Moreover preferred are compounds of formula (I) wherein R³ is a hydrogen atom.

Especially preferred are compounds of formula (I) wherein A is NH and R³ is H, i.e. compounds of formula (Ia):



wherein R¹ and R² are as defined above.


Further preferred are compounds of formula (I) or (Ia) wherein R² is an optionally substituted aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aralkyl or heteroaralkyl group.


Moreover preferred are compounds of formula (I) or (Ia) wherein R^2 is an optionally substituted phenyl or naphthyl group or an optionally substituted heteroaryl group having one or two rings containing 5, 6, 7, 8, 9 or 10 ring atoms, or an optionally substituted arylheterocycloalkyl, heteroaryl-cycloalkyl or heteroarylheterocycloalkyl group containing two or three (especially two) rings and from 9 to 20 (especially 9 or 10) ring atoms. Preferably, the heteroatoms are selected from S, O and N. Further preferably, the number of heteroatoms is from 1 to 6 (especially 1, 2, 3 or 4).


As already mentioned above, according to a preferred embodiment of the present invention, R^2 is bound to the pyrimidine ring of formula (I) or (Ia) via a carbon-carbon bond. In other words, R^2 is bound to the pyrimidine moiety of formula (I) or (Ia) via a carbon atom, i.e. a carbon atom of group R^2 is bound to the pyrimidine moiety.


Especially preferred are compounds of formula (I) or (Ia) wherein, R^2 is selected from the following groups: phenyl; 3-anisyl; 4-anisyl; 2-anisyl; 4,5 dimethoxyphenyl; 6-hydroxyphenyl; 4-pyridyl; 6-pyrrolidinyl-3-pyridinyl; 3-pyrid-3-ylphenyl; 3-morpholinophenyl; 1-methyl-4-piperidyl; 2-thienyl; 3-thienyl; 3-pyrazolyl; 2-benzthiazolyl; imidazo[1,2-a]pyridin-2-yl; 3,6 dimethoxyphenyl; benzimidaz-2-yl and 1-methyl-2-benzimidazolyl.

Especially preferably, R^2 is selected from the following groups:










Moreover preferably, R^2 is derived from the following aldehydes: benzaldehyde; isonicotinaldehyde; 6-(pyrrolidin-1-yl)nicotinaldehyde; 3,4-dimethoxybenzaldehyde; 1-methylpiperidine-4-carbaldehyde; 1H-pyrazole-3-carbaldehyde; 4-methoxybenzaldehyde; 3-methoxybenzaldehyde; thiophene-2-carbaldehyde; thiophene-3-carbaldehyde; benzo[d]thiazole-2-carbaldehyde; 3-(pyridin-3-yl)benzaldehyde; 2-hydroxybenzaldehyde; 3-morpholinobenzaldehyde; 2-methoxybenzaldehyde; 2,5-dimethoxybenzaldehyde; N-(2-aminoethyl)-3-formylbenzamide; imidazo[1,2-a]pyridine-2-carbaldehyde; 1H-benzo[d]imidazole-2-carbaldehyde; 1-methyl-1H-benzo[d]imidazole-2-carbaldehyde; 1H-imidazole-5-carbaldehyde; 1H-imidazole-2-carbaldehyde; 3-(2H-tetrazol-5-yl)benzaldehyde; 2-oxo-2-phenylacetaldehyde; 3-formylbenzoic acid; 2-(3-formylphenoxy)acetic acid; 5-formylthiophene-3-carboxylic acid; 2-(3-formylphenyl)acetic acid; 6-formylpicolinic acid; thiazole-2-carbaldehyde; 3-(((4-methylbenzyl)amino)methyl)benzaldehyde; 3-(3-(dimethylamino)propoxy)benzaldehyde; 3-formylbenzenesulfonamide; N-(3-formylphenyl)methanesulfonamide; thiazole-4-carbaldehyde; oxazole-4-carbaldehyde; oxazole-2-carbaldehyde; methyl 3-

formylbenzoate; methyl 4-formylbenzoate; 3-(pentafluorosulfanyl)benzaldehyde; 3-(methylthio)benzaldehyde; 3-(trifluoromethoxy)benzaldehyde; 3-(trifluoromethyl)benzaldehyde; 3-chlorobenzaldehyde; 2-fluorobenzaldehyde; 3-fluorobenzaldehyde; 4-formylbenzoic acid; 3-(methylsulfonyl)benzaldehyde; 1H-indazole-6-carbaldehyde; 1H-pyrrolo[3,2-b]pyridine-6-carbaldehyde; 4-fluorobenzaldehyde; 3,4-difluorobenzaldehyde; 3,5-difluorobenzaldehyde; 2,6-difluorobenzaldehyde; 2,4-difluorobenzaldehyde; 3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazine-6-carbaldehyde;


Further preferred are compounds of formula (Ib):



(Ib)

wherein R^8 and R^9 together are part of an optionally substituted cycloalkyl, heterocycloalkyl, aryl or heteroaryl group and R^1 is as defined above.

Further preferred are compounds of formula (Ic):



(Ic)

wherein R¹⁰ and R¹¹ together are part of an optionally substituted cycloalkyl or heterocycloalkyl group and R¹ is as defined above.

Further preferred are compounds of formula (I), (Ia), (Ib) and (Ic) wherein R¹ is an aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aralkyl or heteroaralkyl group, all of which may optionally be substituted.

Further preferred are compounds of formula (I), (Ia), (Ib) and (Ic) wherein R¹ is an aryl, heteroaryl, CH₂-aryl or CH₂-heteroaryl group, all of which may optionally be substituted.

Moreover preferred are compounds of formula (I), (Ia), (Ib) and (Ic) wherein R¹ is an optionally substituted phenyl or naphthyl group or an optionally substituted heteroaryl group having one or two rings containing 5, 6, 7, 8, 9 or 10 ring atoms, or an optionally substituted arylheterocycloalkyl, heteroarylcycloalkyl or heteroarylheterocycloalkyl group containing two or three (especially two) rings and from 9 to 20 (especially 9 or 10) ring atoms. Preferably, the heteroatoms are selected from S, O and N, especially from N

and O. Further preferably, the number of heteroatoms is from 1 to 6 (especially 1, 2, 3 or 4).

Especially preferably, R^1 is an optionally substituted phenyl group or an optionally substituted heteroaryl group having one ring containing 5 or 6 ring atoms. Preferably this phenyl or heteroaryl group is substituted by one or more (especially one) C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_1 - C_{10} heteroalkyl, C_3 - C_{18} cycloalkyl, C_2 - C_{17} heterocycloalkyl, C_4 - C_{20} alkylcycloalkyl, C_2 - C_{19} heteroalkylcycloalkyl, C_6 - C_{18} aryl, C_1 - C_{17} heteroaryl, C_7 - C_{20} aralkyl or C_2 - C_{19} heteroaralkyl group(s). Preferably, the heteroatoms are selected from S, O and N, especially from N and O. Especially preferably the phenyl or heteroaryl group is substituted by one or more (especially one) C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_6 heteroalkyl, C_3 - C_{10} cycloalkyl, C_2 - C_9 heterocycloalkyl, C_7 - C_{12} alkylcycloalkyl, C_2 - C_{11} heteroalkylcycloalkyl, C_6 - C_{10} aryl, C_1 - C_9 heteroaryl, C_7 - C_{12} aralkyl or C_2 - C_{11} heteroaralkyl group(s). Preferably, the heteroatoms are selected from S, O and N, especially from N and O.

Further preferred are compounds of formula (I), (Ia), (Ib) and (Ic) wherein R^1 is a group of formula $-CH_2-Ar$ wherein Ar is an optionally substituted phenyl or naphthyl group or an optionally substituted heteroaryl group having one or two rings containing 5, 6, 7, 8, 9 or 10 ring atoms, or an optionally substituted arylheterocycloalkyl, heteroarylalkyl or heteroarylheterocycloalkyl group containing two or three (especially two) rings and from 9 to 20 (especially 9 or 10) ring atoms. Preferably, the heteroatoms are selected from S, O and N, especially from N and O. Further preferably, the number of heteroatoms is from 1 to 6 (especially 1, 2, 3 or 4).

Especially preferably, Ar is an optionally substituted phenyl group or an optionally substituted heteroaryl group having one ring containing 5 or 6 ring atoms. Preferably this phenyl or heteroaryl group is substituted by one or more (especially one) C₁-C₁₀ alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ heteroalkyl, C₃-C₁₈ cycloalkyl, C₂-C₁₇ heterocycloalkyl, C₄-C₂₀ alkylcycloalkyl, C₂-C₁₉ heteroalkylcycloalkyl, C₆-C₁₈ aryl, C₁-C₁₇ heteroaryl, C₇-C₂₀ aralkyl or C₂-C₁₉ heteroaralkyl group(s). Preferably, the heteroatoms are selected from S, O and N, especially from N and O. Especially preferably the phenyl or heteroaryl group is substituted by one or more (especially one) C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ heteroalkyl, C₃-C₁₀ cycloalkyl, C₂-C₉ heterocycloalkyl, C₇-C₁₂ alkylcycloalkyl, C₂-C₁₁ heteroalkylcycloalkyl, C₆-C₁₀ aryl, C₁-C₉ heteroaryl, C₇-C₁₂ aralkyl or C₂-C₁₁ heteroaralkyl group(s). Preferably, the heteroatoms are selected from S, O and N, especially from N and O.

Further preferably, R¹ is a group of formula X¹-L¹-Y¹ or a group of formula X¹-L¹-Y¹-L²-Z¹ wherein

X¹ is an optionally substituted phenyl group or an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N;

L¹ is a bond or a group of formula -CH₂-, -C(=O)-, -SO-, -SO₂-, -NH-C(=O)-, -C(=O)-NH-, -C(=O)-O-, -O-C(=O)-, -NH-C(=O)-O-, -O-C(=O)-NH-, -NH-SO₂-NH-, -CH₂-NH-CH₂-, -NH-SO₂-, -SO₂-NH- or -NH-C(=O)-NH- (preferably, L¹ is a bond or a group of formula -CH₂-, -C(=O)-, -SO₂- or -NH-C(=O)-NH-);

Y¹ is an optionally substituted phenyl group, an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N, an optionally substituted C₃-C₇ cycloalkyl group or an optionally

substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N (preferably, Y^1 is an optionally substituted C_3 - C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N);

L^2 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO-$, $-SO_2-$, $-NH-C(=O)-$, $-C(=O)-NH-$; $-C(=O)-O-$, $-O-C(=O)-$, $-NH-C(=O)-O-$, $-O-C(=O)-NH-$, $-NH-SO_2-NH-$, $-CH_2-NH-CH_2-$, $-NH-SO_2-$, $-SO_2-NH-$ or $-NH-C(=O)-NH-$ (preferably, L^2 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO_2-$ or $-NH-C(=O)-NH-$; especially preferably, L^2 is a bond); and

Z^1 is an optionally substituted phenyl group, an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N, an optionally substituted C_3 - C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N (preferably, Z^1 is an optionally substituted C_3 - C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N).

Further preferably, R^2 is a group of formula $X^2-L^3-Y^2$ or a group of formula $X^2-L^3-Y^2-L^4-Z^2$ wherein

X^2 is an optionally substituted phenyl group or an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N;

L^3 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO-$, $-SO_2-$, $-NH-C(=O)-$, $-C(=O)-NH-$; $-C(=O)-O-$, $-O-C(=O)-$, $-NH-C(=O)-O-$, $-O-C(=O)-NH-$, $-NH-SO_2-NH-$, $-CH_2-NH-CH_2-$, $-NH-SO_2-$, $-SO_2-NH-$ or $-NH-C(=O)-NH-$ (preferably, L^3 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO_2-$ or $-NH-C(=O)-NH-$);

Y^2 is an optionally substituted phenyl group, an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N, an optionally substituted C_3 - C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N (preferably, Y^2 is an optionally substituted C_3 - C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N);


L^4 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO-$, $-SO_2-$, $-NH-C(=O)-$, $-C(=O)-NH-$; $-C(=O)-O-$, $-O-C(=O)-$, $-NH-C(=O)-O-$, $-O-C(=O)-NH-$, $-NH-SO_2-NH-$, $-CH_2-NH-CH_2-$, $-NH-SO_2-$, $-SO_2-NH-$ or $-NH-C(=O)-NH-$ (preferably, L^4 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO_2-$ or $-NH-C(=O)-NH-$; especially preferably, L^4 is a bond); and


Z^2 is an optionally substituted phenyl group, an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N, an optionally substituted C_3 - C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N (preferably, Z^2 is an optionally substituted C_3 - C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N).


Especially preferred are compounds of formula (I), (Ia), (Ib) and/or (Ic) wherein, R^1 is selected from the following groups: 4-morpholinophenyl; 4-(4-methylpiperazino)phenyl; 6-(4-methylpiperazino)-3-pyridyl; 4-piperidinophenyl; 4-pyrrolidinophenyl; 4-carboxyphenyl; 4-(di-methylamino)phenyl; p-amino-carbonylphenyl; 4-cyanophenyl; 4,5 dimethoxyphenyl; 3-


hydroxyphenyl; 3-aminophenyl; 3-anisyl; 1,6-benzomorpholinylene; 6-indazolyl; 2H-1,4-benzoxazin-3-on-6-yl; 2-methyl-benzimidazol-6-yl; 6-benzotriazolyl; 5-cyclopropylpyrazol-3-yl; 3-pyrazolyl; 2-aminocyclohexyl; 4-chlorophenyl; 3-dimethylaminophenyl; 3-(aminocarbonyl)phenyl; 2-(aminocarbonyl)phenyl; 4-(N-ethylpiperazino)phenyl; 4-(4-cyclopropyl-1-piperazinyl)phenyl; 4-dimethylaminomethylphenyl; 4-(4-methylpiperazinomethyl) phenyl and 4-pyrrolidinomethyl-phenyl.

Especially preferably, R^1 is selected from the following groups:









Especially preferred are compounds of formula (I), (Ia), (Ib) and/or (Ic) wherein, R¹ is derived from the following amines:

m-toluidine; 3-(trifluoromethyl)aniline; 3,4,5-trimethoxyaniline; 1H-indazol-5-amine; aniline; 1H-indazol-6-amine; 3-chloroaniline; 7-methyl-1H-indazol-5-amine; 2-methoxyethan-1-amine; thiophen-2-ylmethanamine; 6-methyl-1H-indazol-5-amine; 2H-indazol-6-amine; methyl 4-aminobenzoate; 1H-benzo[d]imidazol-5-amine; 2H-indazol-7-amine; (1-methyl-1H-pyrrol-2-yl)methanamine; benzo[d][1,3]dioxol-5-amine; pyridin-3-amine; 1-methyl-1H-indazol-6-amine; 6-methoxypyridin-3-amine; 4-(4-methylpiperazin-1-yl)aniline; 4-(4-methyl-1,4-diazepan-1-yl)aniline; pyridin-2-amine; 5-bromopyridin-2-amine; isoquinolin-3-amine; 4-methylpyridin-2-amine; 4,6-dimethylpyridin-2-amine; 1H-indazol-7-amine; benzene-1,3-diamine; 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one; 1H-benzo[d][1,2,3]triazol-5-amine; 3-aminobenzimidamide; 4-(piperidin-1-yl)aniline; N1,N1-dimethylbenzene-1,4-diamine; 3-aminobenzamide; 3,4-dimethoxyaniline; 4-morpholinoaniline; 2-methyl-1H-benzo[d]imidazol-6-amine; 4-aminobenzoic acid; 4-aminobenzamide; 4-aminobenzonitrile; 3-methoxyaniline; 4-methoxyaniline; 3-aminobenzonitrile; benzo[c][1,2,5]thiadiazol-5-amine; 3-aminopyridin-2(1H)-one; 2-ethoxyaniline; 1H-pyrazol-3-amine; 5-amino-1H-pyrazole-4-carboxamide; 2-phenoxyaniline; 3-phenoxyaniline; 5-amino-1H-benzo[d]imidazol-2(3H)-one; 1H-indol-5-amine; 4-(aminomethyl)aniline; 1H-indol-6-amine; N1,N1-dimethylbenzene-1,3-diamine; 3-phenyl-1H-pyrazol-5-amine; N1,N1-diethylbenzene-1,4-diamine; 4-(pyrrolidin-1-yl)aniline; 4H-1,2,4-triazole-3,5-diamine; 3-morpholinoaniline; 3-cyclobutyl-1H-pyrazol-5-amine; 4-(4,5-dihydro-1H-imidazol-2-yl)aniline; 4-(4-aminophenyl)morpholin-3-one; 2,2-dimethyl-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-amine; 7-amino-3,4-dihydroquinolin-2(1H)-one; 6-amino-2H-benzo[b][1,4]thiazin-3(4H)-one; 5-(tert-

butyl)-1H-pyrazol-3-amine; 3-methyl-1H-pyrazol-5-amine; 5-cyclopropyl-1H-pyrazol-3-amine; 4-(1H-tetrazol-5-yl)aniline; 2,3-dihydrobenzo[b][1,4]dioxin-6-amine; 4-(1-methylpiperidin-4-yl)aniline; 6-morpholinopyridin-3-amine; 4-(2-methoxyethoxy)aniline; 4-ethoxy-3-methoxyaniline; 1-(4-aminophenyl)pyrrolidin-2-one; 4-thiomorpholinoaniline; 5-aminobenzo[d]oxazol-2(3H)-one; 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 7-aminoquinazolin-4-ol; 4-(4-aminophenyl)thiomorpholine 1,1-dioxide; 2-(4-aminophenyl)acetamide; 3-aminophenol; 3,4-diethoxyaniline; 6-amino-1H-benzo[d][1,3]oxazine-2,4-dione; 5-amino-2-methoxyphenol; 3-methoxy-N-methylaniline; N-(3-aminophenyl)acetamide; 1H-pyrazol-4-amine; 4-fluoro-3-methoxyaniline; 3-fluoro-4-methoxyaniline; 1-methyl-1H-benzo[d]imidazol-5-amine; 1-(3-aminophenyl)ethan-1-one; N-(4-aminophenyl)acetamide; 1H-pyrrolo[2,3-b]pyridin-6-amine; 3-aminobenzenesulfonamide; 4-aminobenzenesulfonamide; pyridine-2,6-diamine; 1,2,3-trimethyl-1H-indol-5-amine; pyrimidine-2,4-diamine; 5-(methylthio)-4H-1,2,4-triazol-3-amine; 5-cyclopropyl-4H-1,2,4-triazol-3-amine; N-(5-amino-2-methoxyphenyl)acetamide; 1H-benzo[d]imidazol-2-amine; 1H-imidazol-2-amine; 1-(4-aminophenyl)ethan-1-one; 4H-benzo[d][1,3]dioxin-6-amine; 1,3-dihydroisobenzofuran-5-amine; 1-methyl-1H-benzo[d]imidazol-6-amine; 4,5-dimethylthiazol-2-amine; 2-methyl-4-(4-methylpiperazin-1-yl)aniline; 6-methylpyridin-2-amine; 4-methylthiazol-2-amine; 4,5,6,7-tetrahydrobenzo[d]thiazol-2-amine; 4-phenoxyaniline; 2-methyl-1,2,3,4-tetrahydrobenzo[4,5]imidazo[1,2-a]pyrazin-8-amine; 4-(pyridin-4-ylmethyl)aniline; 4-aminobenzene-1,2-diol; 4-((1-methylpiperidin-4-yl)oxy)aniline; 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one; 6-(4-methylpiperazin-1-yl)pyridin-3-amine; N1,N1,2-trimethylbenzene-1,4-diamine; 4-(4-cyclopropylpiperazin-1-yl)aniline; ammonia; 3-fluoro-4-

morpholinoaniline; 7-aminoquinoxalin-2(1H)-one; 3-methyl-4-(4-methylpiperazin-1-yl)aniline; 4-(piperazin-1-yl)aniline; 4-((dimethylamino)methyl)aniline; 2-fluoro-4-morpholinoaniline; 4-(4-ethylpiperazin-1-yl)aniline; 8-amino-4,5-dihydro-1H-benzo[b]azepin-2(3H)-one; 5-amino-1,3-dimethyl-1H-benzo[d]imidazol-2(3H)-one; 4-benzylaniline; 2-methyl-4-morpholinoaniline; N1-methyl-N1-(1-methylpiperidin-4-yl)benzene-1,4-diamine; 4-(2-morpholinoethyl)aniline; 3-chloro-4-(4-methylpiperazin-1-yl)aniline; 1,2,3,4-tetrahydroquinolin-7-amine; cyclohexane-1,2-diamine; pyridin-4-amine; 2-(4-aminophenyl)-N-(4-methoxyphenethyl)acetamide; 3-(piperazin-1-yl)aniline; 4-amino-N-(2-(diethylamino)ethyl)benzamide; 2-(4-methylpiperazin-1-yl)pyrimidin-5-amine; 7-amino-2H-benzo[b][1,4]oxazin-3(4H)-one; 3,4-dihydro-2H-benzo[b][1,4]oxazin-7-amine; 3-(4-methylpiperazin-1-yl)aniline; 3-(2-(piperazin-1-yl)ethoxy)aniline; 6-methyl-5,6,7,8-tetrahydro-1,6-naphthyridin-3-amine; (4-aminophenyl)(pyrrolidin-1-yl)methanone; (4-aminophenyl)(morpholino)methanone; 4-(pyrrolidin-1-ylmethyl)aniline; (4-aminophenyl)(4-methylpiperazin-1-yl)methanone; N2-(2-(dimethylamino)ethyl)pyrimidine-2,5-diamine; 4-(morpholinomethyl)aniline; 4-((4-methylpiperazin-1-yl)methyl)aniline; 4-(4-ethylpiperazin-1-yl)-3-fluoroaniline; 4-(2-(4-benzylpiperidin-1-yl)ethyl)aniline; 4-((4-benzylpiperidin-1-yl)methyl)aniline; p-toluidine; 6-(2-(dimethylamino)ethoxy)pyridin-3-amine; 2-methyl-1H-benzo[d]imidazol-5-amine; N2-(3-(dimethylamino)propyl)pyridine-2,5-diamine; N2-(2-(dimethylamino)ethyl)pyridine-2,5-diamine; 6-((1-methylpiperidin-4-yl)oxy)pyridin-3-amine; 4-(4-cyclopentylpiperazin-1-yl)aniline; 4-(4-isobutylpiperazin-1-yl)aniline; 4-(4-isopropylpiperazin-1-yl)aniline; 4-(4-

(cyclopropylmethyl)piperazin-1-yl)aniline; 4-(4-(tert-butyl)piperazin-1-yl)aniline; 2-(4-(4-aminophenyl)piperazin-1-yl)acetic acid; 2-(4-amino-2-methoxyphenoxy)acetic acid; (4-aminophenyl)methanol; 4-(4-(2-(dimethylamino)ethyl)piperazin-1-yl)aniline; 2-(4-aminophenyl)acetic acid; 6-amino-2-naphthoic acid; 3-aminobenzoic acid; 4'-amino-[1,1'-biphenyl]-4-carboxylic acid; 1-(4-aminophenyl)-3-(m-tolyl)urea; 2-(4-aminophenoxy)acetic acid; 2-methylisoindolin-5-amine; (4-amino-3-methoxyphenyl)(4-(4-methylpiperazin-1-yl)piperidin-1-yl)methanone; 4-amino-N-(2-(dimethylamino)ethyl)-N-methylbenzamide; (4-aminophenyl)(2-(methoxymethyl)pyrrolidin-1-yl)methanone; (4-aminophenyl)(azetidin-1-yl)methanone; 4-amino-N,N-dimethylbenzamide; (4-aminophenyl)(4-methyl-1,4-diazepan-1-yl)methanone; 1-(4-aminobenzoyl)piperidin-4-one; (4-aminophenyl)(1,4-dioxo-8-azaspiro[4.5]decan-8-yl)methanone; (4-aminophenyl)(3-(dimethylamino)pyrrolidin-1-yl)methanone; 1-methyl-1,2,3,4-tetrahydroquinolin-6-amine; 3-aminophenylsulphur pentafluoride; 4-fluoroaniline; 3,4-difluoroaniline; N-(4-aminophenyl)-2,2,2-trifluoroacetamide; 3-((6-amino-2H-benzo[b][1,4]oxazin-3-yl)amino)propan-1-ol; N3-phenethyl-2H-benzo[b][1,4]oxazine-3,6-diamine; 3,5-difluoroaniline; 3-fluoro-4-methylaniline; 3,4,5-trifluoroaniline; 4-nitroaniline; 3-methoxy-4-morpholinoaniline; 3-(methylsulfonyl)aniline; 2-(4-aminophenyl)-1,1,1,3,3,3-hexafluoropropan-2-ol; 4-(difluoromethoxy)-3-methoxyaniline; 3-fluoro-4-(trifluoromethyl)aniline; 3-fluoro-4-(trifluoromethoxy)aniline; 2,3-dimethoxyaniline; 2,4-dimethoxyaniline; 3,5-dimethoxyaniline; 4-amino-N,N-dimethylbenzenesulfonamide; 3-amino-N-cyclopropylbenzenesulfonamide; 4-(2H-1,2,3-triazol-2-yl)aniline; 3-(methylsulfinyl)aniline; 3-(2H-1,2,3-triazol-2-yl)aniline; 3-amino-N-methylbenzenesulfonamide; 3-

(morpholinosulfonyl) aniline; 3-
 ((trifluoromethyl) sulfonyl) aniline; 2-((3-
 aminophenyl) sulfonyl) ethan-1-ol; N-(4-aminophenyl)-4-
 fluorobenzamide; 4-morpholino-3-nitroaniline; 2,4-
 difluoroaniline; 2-aminobenzamide; 4-chloroaniline; N1,N1-
 dimethylethane-1,2-diamine; (1-methylpiperidin-4-
 yl) methanamine; 1-methyl-1,2,3,4-tetrahydroquinolin-7-amine;
 2-(4-aminophenyl)-2-methylpropanenitrile; 4-aminophenylsulphur
 pentafluoride; 3-amino-N,N-dimethylbenzenesulfonamide; 2-
 (methylsulfonyl) aniline; 4-(4-(4-methylpiperazin-1-
 yl)piperidin-1-yl) aniline; 3-((dimethylamino)methyl) aniline;
 (4-aminophenyl) (4-(4-methylpiperazin-1-yl)piperidin-1-
 yl) methanone.

Furthermore R¹ can be derived from the following amines:

formamide; 2-aminoethan-1-ol; prop-2-yn-1-amine; N1-
 methylethane-1,2-diamine; 2-aminoacetonitrile; 3-aminopropan-
 1-ol; butan-1-amine; cyclopropanamine; propan-2-amine; 3-
 aminopropanenitrile; 4-aminobutan-1-ol; cyclobutanamine; 2-
 aminopropan-1-ol; acetamide; cyclopropylmethanamine; 5-
 aminopentan-1-ol; 2-aminoacetamide; isoxazol-3-amine; thiazol-
 2-amine; 3-aminopropane-1,2-diol; cyclopentanamine; piperidin-
 4-amine; piperidin-3-amine; pyrimidin-2-amine; 2-
 aminocyclopentanol; 3-aminopropanamide; tetrahydro-2H-pyran-4-
 amine; 2-methylpropan-2-amine; o-toluidine; 2,2,2-
 trifluoroethan-1-amine; phenylmethanamine; piperidin-4-
 ylmethanamine; 2-aminocyclohexanol; 4-aminobutanamide;
 piperidin-3-ylmethanamine; 1-methyl-1H-pyrazol-4-amine; 2-
 methoxyaniline; 2-chloroaniline; 2-aminopropanamide; 4-
 methylthiophen-2-amine; 2-phenylethan-1-amine; 1H-pyrazol-5-
 amine; 5-methylisoxazol-3-amine; 2-morpholinoethan-1-amine; 1-
 (aminomethyl)-N-methylcyclopropanamine; 1-methyl-1H-pyrrol-3-
 amine; 5-methylthiazol-2-amine; 5-methylthiophen-2-amine; 4-

aminophenol; 3-fluoroaniline; 3,5-dimethylisoxazol-4-amine; 3-morpholinopropan-1-amine; 2-aminobutanamide; 4-iodoaniline; (3-aminophenyl)methanol; 2-aminothiazole-4-carbaldehyde; 3-bromoaniline; 2,6-dimethylaniline; 4-ethylaniline; 3-amino-2-methylphenol; 4-(methylthio)aniline; 3-ethylaniline; 1-phenylethan-1-amine; 2-(4-aminophenyl)ethan-1-ol; 5-aminonicotinaldehyde; 6-aminonicotinaldehyde; 4-aminobenzaldehyde; 3-aminobenzaldehyde; indolin-6-amine; 4-amino-2-methoxyphenol; 2-aminopyrimidine-5-carbaldehyde; 5-aminopyrazine-2-carbaldehyde; 5-aminopicolinaldehyde; 3-methoxy-4-methylaniline; 6-aminopyrazine-2-carbaldehyde; N1,6-dimethylbenzene-1,3-diamine; 5-methyl-1H-pyrazol-3-amine; 4-ethoxyaniline; 2,3-dihydrobenzofuran-5-amine; 3-ethoxyaniline; benzo[d]thiazol-5-amine; benzo[d]thiazol-6-amine; piperidine-3-carboxamide; imidazo[1,2-a]pyridin-6-amine; piperidine-4-carboxamide; benzo[d]thiazol-7-amine; benzo[d]isoxazol-5-amine; 4-methoxy-3-methylaniline; benzo[d]thiazol-2-amine; 4-vinyylaniline; benzo[c][1,2,5]thiadiazol-4-amine; 1-aminocyclopropanecarboxamide; 2-phenylcyclopropanamine; 2-aminocyclopentanecarboxamide; 3-vinyylaniline; (5-amino-2-methoxyphenyl)methanol; 2-(4-aminophenoxy)ethan-1-ol; 1,2,3,4-tetrahydroisoquinolin-6-amine; (4-amino-2-methoxyphenyl)methanol; 2-amino-4-methylpyrimidine-5-carbaldehyde; 6-amino-4-methylnicotinaldehyde; 2-isopropoxyaniline; 6-amino-2-methylnicotinaldehyde; 4-amino-2-methylphenol; 5-amino-2-methylphenol; 3-chloro-4-methoxyaniline; 3,5-dimethylaniline; N-(3-aminophenyl)formamide; 2-(3-aminophenoxy)ethan-1-ol; N-(6-aminopyridin-2-yl)formamide; 4-amino-2-fluorophenol; 5-amino-2-hydroxybenzonitrile; 4-amino-3-fluorophenol; N-(4-aminophenyl)formamide; 2,4-dimethylaniline; 3,4-dimethylaniline; 2-fluoro-5-methylaniline; 2,5-dimethylaniline; quinoxalin-6-amine; quinolin-6-amine; 2-

amino-3-methylbutanamide; quinoxalin-5-amine; naphthalen-1-amine; naphthalen-2-amine; 4-fluoro-3-methylaniline; quinolin-5-amine; quinolin-8-amine; 2,6-dimethylpyrimidin-4-amine; 1-(4-aminophenyl)ethan-1-ol; 2,3,4,5-tetrahydrobenzo[b][1,4]oxazepin-7-amine; 3-methoxy-4-(methoxymethyl)aniline; 2-fluoro-4-methoxyaniline; 5-amino-6-methoxypyrazine-2-carbaldehyde; 2-amino-4-methoxypyrimidine-5-carbaldehyde; 6-amino-5-methoxynicotinaldehyde; 3-chloro-4-fluoroaniline; 4H-benzo[b][1,4]oxazin-6-amine; 4-isopropylaniline; 4-amino-2,5-dimethylphenol; 4-amino-2-chlorophenol; 4H-benzo[b][1,4]oxazin-7-amine; 3-(2-methoxyethoxy)aniline; 4-methoxy-2-methylaniline; 5-methoxy-2-methylaniline; 3-(2-(methylamino)ethoxy)aniline; 3-isopropylaniline; 4-amino-2,3-dimethylphenol; N-(5-amino-2-methylphenyl)formamide; 2-amino-4-methylpentanamide; 4-chloro-3-methylaniline; 3-aminocyclopentanecarboxamide; 2-chloro-5-fluoropyrimidin-4-amine; 3,4-dihydroquinolin-6-amine; 2-amino-4-methylpentanethioamide; 2-(isopentyloxy)aniline; 6-amino-5-methylnicotinaldehyde; 5-amino-6-methylpyrazine-2-carbaldehyde; 2-amino-6-methylpyrimidine-4-carbaldehyde; 2-methyl-2H-indazol-6-amine; 5-amino-6-methylpicolinaldehyde; 5-amino-4-methylpicolinaldehyde; 4-isopropoxyaniline; 1-methyl-1H-indazol-5-amine; 3,5-dichloroaniline; 3,4-dichloroaniline; [1,1'-biphenyl]-2-amine; 2,6-dimethoxypyridin-3-amine; 4-methoxy-3,5-dimethylaniline; 2-methyl-2H-indazol-5-amine; 3-(ethyl(hydroxy)amino)aniline; 3-isopropoxyaniline; N1-isopropylbenzene-1,3-diamine; 4-amino-5-chloro-2-methylphenol; 1-methyl-1H-indol-4-amine; 1H-indazol-4-amine; 1-methyl-1H-indazol-4-amine; 2-methyl-2H-indazol-4-amine; 1H-indol-4-amine; 1H-benzo[d]imidazol-6-amine; 1H-benzo[d][1,2,3]triazol-6-amine; 2-methylbenzo[d]thiazol-5-amine; 4-(methylsulfinyl)aniline; 1-methyl-1H-indol-5-amine; 3-(2-aminophenyl)propanamide; 2-((2-aminocyclohexyl)amino)acetic

acid; (2,3,6-trifluorophenyl)methanamine; 5-bromo-2-chloropyrimidin-4-amine; 1-methyl-1H-indazol-7-amine; 1-methyl-1H-benzo[d]imidazol-4-amine; 2-methyl-2H-indazol-7-amine; 2-methylbenzo[d]oxazol-7-amine; 4-methyl-3,4-dihydro-2H-benzo[b][1,4]oxazin-7-amine; 4-methyl-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 4-(4H-1,2,4-triazol-4-yl)aniline; 4-(1H-imidazol-1-yl)aniline; 4-(1H-pyrazol-1-yl)aniline; 5-aminoindolin-2-one; 6-aminoindolin-2-one; 3-methoxy-4-(2-methoxyethoxy)aniline; 3-(4-amino-2-methoxyphenoxy)propan-1-ol; 2-((4-aminophenyl)(methylamino)ethan-1-ol; 2-methyl-1,2,3,4-tetrahydroisoquinolin-6-amine; 4-amino-N-methylbenzamide; 3-amino-N-methylbenzamide; 1-(piperidin-4-yl)-1H-pyrazol-4-amine; 4-(oxazol-4-yl)aniline; 4-(pyrrolidin-3-yl)aniline; 2-(trifluoromethoxy)aniline; 3-chloro-4-methoxy-5-methylaniline; 2-((2-aminophenyl)imino)acetic acid; 3-(oxazol-5-yl)aniline; (5-aminobenzofuran-2-yl)methanol; 3,4,5-trimethylaniline; N-(5-amino-2-fluorophenyl)formamide; methyl 4-amino-1-methyl-1H-pyrrole-2-carboxylate; methyl 3-aminobenzoate; N-(3-amino-4-ethoxyphenyl)formamide; 2-((3-aminophenyl)amino)propan-1-ol; 2-methyl-2,3-dihydrobenzo[b][1,4]dioxin-6-amine; 4-(1H-1,2,4-triazol-1-yl)aniline; 3-(1H-pyrazol-1-yl)aniline; 2-amino-2-phenylacetamide; 4-(thiazol-4-yl)aniline; 1,2-dimethyl-1H-indol-4-amine; 4-(oxazol-5-yl)aniline; 1-ethyl-1H-indol-4-amine; 3-(thiazol-2-yl)aniline; 4-(1,2,3-thiadiazol-4-yl)aniline; 3-(isoxazol-3-yl)aniline; 4-(isoxazol-3-yl)aniline; 4-(isoxazol-5-yl)aniline; 4-(thiophen-2-yl)aniline; 3-(1H-tetrazol-1-yl)aniline; 4-(1H-tetrazol-1-yl)aniline; 3-(1H-imidazol-1-yl)aniline; 5-aminobenzofuran-2(3H)-one; 8-methylquinolin-4-amine; 2-amino-2-(pyridin-3-yl)acetamide; 1-phenyl-1H-pyrazol-4-amine; 1-phenyl-1H-pyrrol-3-amine; 3-(2H-tetrazol-2-yl)aniline; 3-(1H-1,2,4-triazol-1-yl)aniline; 3-(1H-1,2,3-triazol-1-yl)aniline; 4-(1H-1,2,3-

triazol-1-yl)aniline; 3-(pyrrolidin-1-yl)aniline; 3-(1H-pyrrol-1-yl)aniline; 4-(1H-pyrrol-1-yl)aniline; 4-(1,3,4-oxadiazol-2-yl)aniline; 4-(thiazol-2-yl)aniline; 3-(thiazol-4-yl)aniline; 3-(oxazol-4-yl)aniline; 3-(thiazol-5-yl)aniline; 4-(thiazol-5-yl)aniline; 6-fluoronaphthalen-2-amine; methyl 2-((2-aminocyclohexyl)amino)acetate; 4-isobutoxyaniline; 2-methylquinolin-6-amine; 2-methylquinolin-8-amine; 3-methylcinnolin-5-amine; 2-(4-aminophenyl)propan-2-ol; 2-((4-aminophenyl)(ethyl)amino)ethan-1-ol; 4-(2-(dimethylamino)ethoxy)aniline; 4-(tetrahydro-2H-pyran-4-yl)aniline; 3-methoxy-4-((2-methoxyethoxy)methyl)aniline; N1-(2-methoxyethyl)-N1-methylbenzene-1,4-diamine; 4-isopropoxy-3-methoxyaniline; 4-amino-2-methoxybenzoic acid; 4-(piperidin-4-yl)aniline; 4-amino-N,2-dimethylbenzamide; 6-(tetrahydro-2H-pyran-4-yl)pyridin-3-amine; 6-(piperidin-4-yl)pyridin-3-amine; 4-(3-(dimethylamino)propyl)aniline; 4-(pyridin-3-yl)aniline; 4-(piperidin-3-yl)aniline; 2-ethyl-1,2,3,4-tetrahydroisoquinolin-6-amine; 3,5-dichloro-4-methoxyaniline; 4-amino-2-chloro-6-methylphenol; 3-(4-aminophenoxy)propane-1,2-diol; 3-(tert-butyl)aniline; 2-(5-amino-1H-indazol-1-yl)ethan-1-ol; 2-(6-amino-1H-indazol-1-yl)ethan-1-ol; 4-chloro-2,5-dimethoxyaniline; ethyl 3-aminobenzoate; 4-(tert-butyl)aniline; 4-chloro-3,5-dimethylaniline; N-(3-amino-5-chlorophenyl)formamide; 4-(trifluoromethyl)aniline; [1,1'-biphenyl]-3-amine; 6-amino-2H-chromen-2-one; 7-amino-2H-chromen-2-one; methyl 2-(4-aminophenyl)acetate; methyl 2-(3-aminophenyl)acetate; methyl 5-amino-2-hydroxybenzoate; methyl 4-amino-2-hydroxybenzoate; 5-amino-2-methoxybenzoic acid; 3-(2-(dimethylamino)ethoxy)aniline; 4-methyl-4H-benzo[b][1,4]oxazin-7-amine; 3-amino-4-isopropylphenol; 5-methyl-3-phenylisoxazol-4-amine; 3-(pyrimidin-2-yl)aniline; 3-(pyrimidin-5-yl)aniline; 2,3-dihydro-1H-pyrrolo[1,2-a]indol-8-amine; 4-(pyrimidin-2-yl)aniline; 3-(pyridin-3-yl)aniline; 4-

(pyridin-2-yl)aniline; 6-methoxynaphthalen-2-amine; 2-methyl-2H-indol-4-amine; 4-chloronaphthalen-1-amine; 3-(pyridin-4-yl)aniline; 4-amino-2-methoxybenzamide; 3-(5-methyl-1H-tetrazol-1-yl)aniline; 2-fluoro-4-(1H-pyrazol-1-yl)aniline; 2-fluoro-4-(thiazol-4-yl)aniline; 4-(pyrimidin-5-yl)aniline; 3-(pyrazin-2-yl)aniline; 4-(pyrazin-2-yl)aniline; 3-(tetrahydro-2H-pyran-4-yl)aniline; 3-(pyridazin-4-yl)aniline; 4-(pyridazin-4-yl)aniline; 4-(pyridin-4-yl)aniline; 3-(pyridin-2-yl)aniline; [1,1'-biphenyl]-4-amine; 7-chloro-1H-indazol-6-amine; 6-bromonaphthalen-2-amine; 3-(1-methyl-1H-tetrazol-5-yl)aniline; 4-(4-methyl-4H-1,2,4-triazol-3-yl)aniline; 4-(1-methyl-1H-imidazol-2-yl)aniline; 3-(1-methyl-1H-imidazol-2-yl)aniline; 4-(2-methyl-1H-imidazol-1-yl)aniline; 3-(2-methyl-1H-imidazol-1-yl)aniline; 2-(4-aminophenyl)-2-methylpropan-1-ol; 1-(4-aminophenyl)azetidin-3-ol; 2-aminoquinazoline-6-carbaldehyde; 1-(4-aminophenyl)-2-methylpropan-2-ol; 2-(4-aminophenoxy)-N-methylacetamide; 4-(1,4-oxazepan-4-yl)aniline; 3-methoxy-4-(pyrrolidin-1-yl)aniline; 4-amino-N-propylbenzamide; 3-aminoquinoline-6-carbaldehyde; 4-((tetrahydrofuran-2-yl)methoxy)aniline; 4-((tetrahydro-2H-pyran-4-yl)oxy)aniline; 4-(pyridin-4-yloxy)aniline; 4-(3-fluoroazetidin-1-yl)aniline; 4-amino-N-(2-hydroxyethyl)benzamide; 1-(4-aminophenyl)cyclobutanol; 2-aminoquinoline-6-carbaldehyde; 4-(2-methoxypropan-2-yl)aniline; 2-((4-amino-2-methoxyphenyl)(methyl)amino)ethan-1-ol; 4-methoxy-3-(pyrrolidin-1-yl)aniline; 4-(3-methylazetidin-1-yl)aniline; 2,3-dimethyl-2H-indazol-6-amine; 4-(trifluoromethoxy)aniline; 3-methyl-1H-indazol-6-amine; 1-(2-morpholinoethyl)-1H-pyrazol-4-amine; 3-(trifluoromethoxy)aniline; 4-amino-N-ethoxybenzamide; 3-amino-N-propylbenzamide; 4-((2-methyl-1H-imidazol-1-yl)methyl)aniline; 3-(5-amino-1H-indazol-1-yl)propan-1-ol; 4-amino-2,6-dichlorophenol; 3-(6-amino-1H-indazol-1-yl)propan-1-

ol; 2-((3-aminophenyl)imino)acetamide; methyl 5-amino-2-methoxybenzoate; ethyl 2-((2-aminophenyl)imino)acetate; 4-((trifluoromethyl)thio)aniline; 5-amino-2-hydroxybenzoic acid; 4-amino-2-hydroxybenzoic acid; 2-((3-aminophenyl)imino)acetic acid; 2,2'-((3-aminophenyl)azanediyl)bis(ethan-1-ol); 2,2-difluorobenzo[d][1,3]dioxol-5-amine; 2-((methylamino)methylene)-2,3-dihydrobenzofuran-5-amine; 3-aminophenyl ethylcarbamate; 1-(4-aminophenyl)-3-ethylurea; 1-(3-aminophenyl)-3-ethylurea; 6-amino-2-methyl-2H-benzo[b][1,4]oxazin-3(4H)-one; 4-methyl-[1,1'-biphenyl]-3-amine; 2-methyl-1H-indol-4-amine; 3-(4-aminopiperidin-1-yl)-3-oxopropanenitrile; 4-(pyridin-3-yloxy)aniline; 1-(3-aminophenyl)pyrrolidin-2-one; 4-(5,6-dihydro-4H-1,3-oxazin-2-yl)aniline; 4-(3,6-dihydro-2H-pyran-4-yl)aniline; 6-aminoquinoline-2-carbonitrile; 2-chloro-5-cyclopropylpyrimidin-4-amine; 3-(3-aminopiperidin-1-yl)-3-oxopropanenitrile; 3-((tetrahydro-2H-pyran-4-yl)oxy)aniline; 3-(3,6-dihydro-2H-pyran-4-yl)aniline; dibenzo[b,d]furan-2-amine; 3-methoxy-4-(oxazol-5-yl)aniline; 4-methoxy-3-(2H-1,2,3-triazol-2-yl)aniline; 1-(4-aminophenyl)pyrrolidin-3-ol; 1-(4-aminophenoxy)-2-methylpropan-2-ol; 4-((tetrahydro-2H-pyran-4-yl)methoxy)aniline; (4-(4-aminophenyl)morpholin-3-yl)methanol; 4-(2H-tetrazol-5-yl)aniline; 2-methoxy-N1-(2-methoxyethyl)-N1-methylbenzene-1,4-diamine; 4-(1-methoxy-2-methylpropan-2-yl)aniline; 5-amino-1-methyl-6-oxo-1,6-dihydropyridine-3-carbaldehyde; 2-((4-amino-2-ethoxyphenyl)(methyl)amino)ethan-1-ol; 4-(2-(pyrrolidin-1-yl)ethoxy)aniline; 4-(1-methyl-1H-pyrazol-4-yl)aniline; 4-(1H-imidazol-4-yl)aniline; 4-(methylsulfonyl)aniline; 2-methoxy-[1,1'-biphenyl]-4-amine; 4-(1-methylpyrrolidin-3-yl)aniline; 5-amino-2-methylisoindolin-1-one; 6-amino-2-methylisoindolin-1-one; 3-(2-(pyrrolidin-1-yl)ethoxy)aniline; 2-(2-methoxyethyl)-1,2,3,4-tetrahydroisoquinolin-6-amine; 4-

methoxy-3-(trifluoromethyl)aniline; 2-((3-aminophenyl)imino)-N-methylacetamide; 3-(1H-tetrazol-5-yl)aniline; N-(4-aminophenyl)-N-methylacetamide; 3-(benzyloxy)aniline; 3-(1H-pyrazol-3-yl)aniline; 2-amino-7-oxabicyclo[4.2.0]octa-1,3,5-triene-8-carboxylic acid; (5-amino-1H-indol-2-yl)methanol; 6-methoxy-[1,1'-biphenyl]-3-amine; 4-methoxy-[1,1'-biphenyl]-3-amine; ethyl (4-amino-2-hydroxyphenyl)carbamate; 3-fluoro-4-(thiazol-4-yl)aniline; 3-fluoro-4-(1H-pyrazol-1-yl)aniline; 4-amino-N-(3-hydroxypropyl)benzamide; 3-fluoro-4-(1H-imidazol-1-yl)aniline; 1-(4-aminophenyl)pyridin-2(1H)-one; 1-(4-aminophenyl)-1-methylurea; butyl 4-aminobenzoate; 4-(5-methyl-1,2,4-oxadiazol-3-yl)aniline; 3-(2-methylthiazol-4-yl)aniline; 4-(3-methyl-1H-pyrazol-1-yl)aniline; 3-methyl-5-(2H-1,2,3-triazol-2-yl)aniline; 4-methyl-3-(2H-1,2,3-triazol-2-yl)aniline; 2-amino-2-(3-hydroxyphenyl)acetamide; 2-amino-2-(3-fluorophenyl)acetamide; 3-fluoro-5-(2H-1,2,3-triazol-2-yl)aniline; 4-fluoro-3-(2H-1,2,3-triazol-2-yl)aniline; 4-amino-2-(2H-1,2,3-triazol-2-yl)benzonitrile; 3-fluoro-4-(2H-1,2,3-triazol-2-yl)aniline; 3-(2H-tetrazol-5-yl)aniline; 3-chloro-1H-indazol-5-amine; 4-(methylthio)-7H-pyrrolo[2,3-d]pyrimidin-2-amine; 4-(2-(methoxymethyl)pyrrolidin-1-yl)aniline; 2'-methoxy-[1,1'-biphenyl]-3-amine; 4-(1-methyl-1H-pyrazol-3-yl)aniline; 3-(1-methyl-1H-pyrazol-3-yl)aniline; 3-(1-methyl-1H-pyrazol-4-yl)aniline; 4-(5-methyl-1,3,4-thiadiazol-2-yl)aniline; 3-(5-methyl-1,2,4-oxadiazol-3-yl)aniline; 3-(4-methyl-2H-1,2,3-triazol-2-yl)aniline; 3-(4-methyl-1H-1,2,3-triazol-1-yl)aniline; 3-(5-methylisoxazol-3-yl)aniline; 3-methyl-5-(2H-tetrazol-2-yl)aniline; 3-methyl-4-(2H-1,2,3-triazol-2-yl)aniline; 3-methyl-4-(1H-pyrazol-1-yl)aniline; 3-(5-methylfuran-2-yl)aniline; 2-amino-2-(m-tolyl)acetamide; 2-amino-2-(p-tolyl)acetamide; 3-amino-1H-indazole-6-carbaldehyde; 3-ethoxy-4-morpholinoaniline; 3-amino-1H-indazole-5-carbaldehyde; 1-(4-aminophenyl)piperidin-

3-ol; 3-methoxy-4-((tetrahydro-2H-pyran-4-yl)oxy)aniline; 2-(4-amino-2-methoxyphenoxy)-N-methylacetamide; 2-amino-1H-benzo[d]imidazole-6-carbaldehyde; 2-(4-aminophenoxy)-N-(2-hydroxyethyl)acetamide; 5-amino-2-morpholinobenzonitrile; 1-(4-aminophenyl)piperidin-4-ol; (1-(4-aminophenyl)pyrrolidin-3-yl)methanol; 4-(4-fluoropiperidin-1-yl)aniline; 3-(methoxymethyl)-4-morpholinoaniline; 3-methoxy-4-((tetrahydrofuran-2-yl)methoxy)aniline; 4-(3-(dimethylamino)propoxy)-3-methoxyaniline; 1-(4-amino-2-methoxyphenyl)azetidin-3-ol; 4-amino-N-(2-hydroxyethyl)-2-methoxybenzamide; 1-(4-amino-2-methoxyphenyl)-2-methylpropan-2-ol; 3-(4-aminophenoxy)-2,2-dimethylpropan-1-ol; 4-(2-methylmorpholino)aniline; 6-(2-methylmorpholino)pyridin-3-amine; 3-methyl-4-(piperidin-4-yl)aniline; 4-(2-morpholinoethoxy)aniline; 3-(2-morpholinoethoxy)aniline; 3-methyl-4-morpholinoaniline; (1-(4-aminophenethyl)pyrrolidin-2-yl)methanol; 4-(2-methylpyridin-4-yl)aniline; 6-(1-methylpiperidin-4-yl)pyridin-3-amine; 2-(2-methylmorpholino)pyrimidin-5-amine; 3-methyl-4-(tetrahydro-2H-pyran-4-yl)aniline; 4-amino-N-cyclopropylbenzamide; 4-(1-methylpiperidin-3-yl)aniline; 4-(1-ethylpyrrolidin-3-yl)aniline; 5-methyl-6-morpholinopyridin-3-amine; 4-methyl-3-(trifluoromethyl)aniline; 5-aminobenzofuran-2-carboxylic acid; 2-((dimethylamino)methyl)benzofuran-5-amine; ethyl 2-((3-aminophenyl)imino)acetate; 3-fluoro-5-(trifluoromethyl)aniline; 4-amino-2-(trifluoromethyl)phenol; 5-amino-2,3-dihydrobenzofuran-2-carboxylic acid; N-(4-aminophenyl)methanesulfonamide; 7-amino-4-methyl-2H-benzo[b][1,4]oxazin-3(4H)-one; 6-amino-4-methyl-2H-benzo[b][1,4]oxazin-3(4H)-one; 3-methyl-5-(trifluoromethyl)aniline; 6-amino-1-methyl-3,4-dihydroquinolin-2(1H)-one; methyl (4-aminophenyl)(methyl)carbamate; N-(4-aminophenyl)-2-hydroxy-N-

methylacetamide; 6-aminoquinolin-2(1H)-one; 6-amino-1-methylquinolin-2(1H)-one; 4-(ethylsulfonyl)aniline; N-(4-aminophenyl)-N-methylpropionamide; 4-(3,5-dimethyl-1H-pyrazol-1-yl)aniline; 2-amino-2-(3-chlorophenyl)acetamide; 5-fluoro-4-(piperazin-1-yl)pyrimidin-2-amine; 4-(2-(piperidin-1-yl)ethoxy)aniline; 3-chloro-5-(2H-1,2,3-triazol-2-yl)aniline; 4-chloro-3-(2H-1,2,3-triazol-2-yl)aniline; (1-(4-aminophenyl)-1H-1,2,3-triazol-4-yl)methanol; 3-(5-fluoropyrimidin-2-yl)aniline; 4'-fluoro-[1,1'-biphenyl]-3-amine; 3'-fluoro-[1,1'-biphenyl]-3-amine; 4-bromo-3-(2H-1,2,3-triazol-2-yl)aniline; dibenzo[b,d]furan-3-amine; 3-bromo-5-(2H-1,2,3-triazol-2-yl)aniline; 3-methoxy-5-(2H-1,2,3-triazol-2-yl)aniline; 3-methoxy-5-(1H-tetrazol-1-yl)aniline; 2-amino-2-(4-methoxyphenyl)acetamide; 1-(4-aminophenyl)-3-methylazetidin-3-ol; (4-(4-aminophenyl)morpholin-2-yl)methanol; 1-(4-amino-2-methoxyphenyl)pyrrolidin-3-ol; 2-(5-amino-2-morpholinophenoxy)ethan-1-ol; (1-(4-aminophenyl)piperidin-4-yl)methanol; 4-(4-aminophenoxy)cyclohexanol; 4-(4-aminophenyl)piperazin-2-one; 4-(3,3-difluoroazetidin-1-yl)aniline; 2-(1-(4-aminophenyl)pyrrolidin-3-yl)ethan-1-ol; 4-amino-N-(oxetan-3-yl)benzamide; 1-(4-aminophenyl)-2,2,2-trifluoroethan-1-ol; 1-(4-amino-2-methoxyphenoxy)-2-methylpropan-2-ol; 3-methoxy-4-((tetrahydro-2H-pyran-4-yl)methoxy)aniline; 4-amino-N-(2-hydroxyethyl)-N-methylbenzamide; 2-(4-aminophenoxy)-N,N-dimethylacetamide; 4-(3-fluoro-3-methylazetidin-1-yl)aniline; 3-methyl-4-(2-methylmorpholino)aniline; 4-(1-ethylpiperidin-4-yl)aniline; 3-fluoro-4-(2-methylmorpholino)aniline; 3-methyl-4-(1-methylpiperidin-4-yl)aniline; 4-(2,3-dihydroimidazo[2,1-b]thiazol-6-yl)aniline; 3-fluoro-4-(1-methylpiperidin-4-yl)aniline; 5-methyl-6-(2-methylmorpholino)pyridin-3-amine; 3-fluoro-4-(piperazin-1-yl)methyl)aniline; 4'-methoxy-[1,1'-biphenyl]-4-amine; 1-(6-amino-3,4-dihydroisoquinolin-2(1H)-

yl)ethan-1-one; 4-amino-N,N-diethylbenzamide; 5-(4-ethylpiperazin-1-yl)pyridin-2-amine; 2-(isopropylsulfonyl)aniline; 3-methyl-4-(1,2,3,6-tetrahydropyridin-4-yl)aniline; 2-methyl-5-(2-morpholinoethoxy)aniline; methyl 5-aminobenzofuran-2-carboxylate; 5-amino-N-methyl-2,3-dihydrobenzofuran-2-carboxamide; methyl 5-amino-2,3-dihydrobenzofuran-2-carboxylate; 3-methoxy-5-(trifluoromethyl)aniline; 5-amino-N-methylbenzofuran-2-carboxamide; 2-((3-aminophenyl)imino)-N-(2-hydroxyethyl)acetamide; 4-chloro-3-(trifluoromethyl)aniline; 3-nitroaniline; methyl 4-amino-2,3-dihydrobenzofuran-7-carboxylate; 6-amino-N-methyl-1H-indole-1-carboxamide; ethyl 2-amino-7-oxabicyclo[4.2.0]octa-1(6),2,4-triene-8-carboxylate; 3-aminophenyl isopropylcarbamate; 3-chloro-4-morpholinoaniline; 2-(6-amino-1H-indazol-1-yl)acetamide; 1-(4-aminophenyl)azetidine-2-carboxamide; 6-amino-2-naphthamide; 3-amino-5-(2H-1,2,3-triazol-2-yl)benzonitrile; 3-amino-5-(1H-1,2,3-triazol-1-yl)benzonitrile; 4-amino-N-cyclobutylbenzamide; 3-(4-methoxypyrimidin-2-yl)aniline; 4-(6-methoxypyridin-3-yl)aniline; 3-(6-methoxypyridin-2-yl)aniline; 3-(6-methoxypyridin-3-yl)aniline; 3'-methoxy-[1,1'-biphenyl]-3-amine; 4'-methoxy-[1,1'-biphenyl]-3-amine; N-(4-aminophenyl)-3-hydroxy-N-methylpropanamide; 9-methyl-9H-carbazol-3-amine; 2-(1-(4-aminophenyl)piperidin-4-yl)ethan-1-ol; 1-(4-amino-2-methoxyphenyl)piperidin-3-ol; 1-(4-aminophenyl)-3-methylpyrrolidin-3-ol; 2-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-ol; 4-(3,3-difluoropyrrolidin-1-yl)aniline; 4-(2-oxa-6-azaspiro[3.3]heptan-6-yl)aniline; 3-(2-methoxyethoxy)-4-morpholinoaniline; 3-(4-amino-2-methoxyphenoxy)-2,2-dimethylpropan-1-ol; 1-(4-amino-2-methoxyphenyl)piperidin-4-ol; (1-(4-amino-2-methoxyphenyl)pyrrolidin-3-yl)methanol; 4-(3-methoxy-3-methylazetidin-1-yl)aniline; 1-(4-amino-2-

ethoxyphenoxy)-2-methylpropan-2-ol; 1-(4-amino-2-ethoxyphenyl)pyrrolidin-3-ol; 4-amino-N-ethyl-N-(2-hydroxyethyl)benzamide; 4-((4-ethylpiperazin-1-yl)methyl)aniline; 1-(4-aminophenyl)-3-ethylazetidin-3-ol; 3-methyl-4-(2-(pyrrolidin-1-yl)ethoxy)aniline; 4-methyl-3-(2-(pyrrolidin-1-yl)ethoxy)aniline; 1-(4-aminophenethyl)piperidin-4-ol; 2-(4-(4-aminophenyl)piperidin-1-yl)ethan-1-ol; 3-methyl-4-((1-methylpiperidin-4-yl)oxy)aniline; 2-methoxy-5-methyl-4-(piperidin-4-yl)aniline; 4-(1-(2-methoxyethyl)-1H-pyrazol-4-yl)aniline; 3-fluoro-4-(2-(pyrrolidin-1-yl)ethoxy)aniline; 4-amino-N-cyclopentylbenzamide; 4-(1-(2-methoxyethyl)pyrrolidin-3-yl)aniline; 2-methoxy-4-(4-methylpiperazin-1-yl)aniline; 2-(4-(4-amino-1H-pyrazol-1-yl)piperidin-1-yl)acetonitrile; (4-aminophenyl)(piperazin-1-yl)methanone; 1-(1-(2-methoxyethyl)piperidin-4-yl)-1H-pyrazol-4-amine; 4-(2-(4-methylpiperazin-1-yl)ethyl)aniline; 4-(1,2-dimethylpiperidin-4-yl)-3-fluoroaniline; methyl 2-(5-amino-1H-indazol-1-yl)acetate; methyl 2-(6-amino-1H-indazol-1-yl)acetate; 2-(5-amino-1H-indazol-1-yl)-N-methylacetamide; 3-chloro-4-(trifluoromethoxy)aniline; 4-(4,5-dichloro-1H-imidazol-1-yl)aniline; 2-(6-amino-1H-indazol-1-yl)-N-methylacetamide; (3-aminophenyl)(phenyl)methanone; methyl 3-amino-5-formamidobenzoate; 5-methoxy-2-methyl-[1,1'-biphenyl]-4-amine; 2-((3-aminophenyl)imino)-N,N-dimethylacetamide; 2-((3-aminophenyl)imino)-N-(2-(methylamino)ethyl)acetamide; ethyl 5-aminobenzofuran-2-carboxylate; 1-(4-aminophenyl)pyrrolidine-2-carboxamide; (1-(2-amino-5-fluoropyrimidin-4-yl)piperidin-4-yl)methanol; 4-amino-N-(2-hydroxyethyl)benzenesulfonamide; 4-(2-amino-5-fluoropyrimidin-4-yl)piperazin-2-one; 3-(1H-benzo[d][1,2,3]triazol-1-yl)aniline; 3-(1H-indazol-1-yl)aniline; 3-(2H-benzo[d][1,2,3]triazol-2-yl)aniline; 3-(1H-benzo[d]imidazol-1-yl)aniline; 3-(2H-indazol-2-yl)aniline; 3-

(imidazo[1,2-a]pyridin-2-yl)aniline; 3-(4-aminophenyl)pyridin-2(1H)-one; 3-(benzo[d][1,3]dioxol-4-yl)aniline; 3-(benzo[d][1,3]dioxol-5-yl)aniline; 3-(2,3-dihydrobenzofuran-5-yl)aniline; 3-(imidazo[1,2-a]pyridin-6-yl)aniline; 4-(imidazo[1,2-a]pyridin-6-yl)aniline; 6-amino-N-methyl-2-naphthamide; 3-fluoro-4-(4-methyl-1H-pyrazol-1-yl)aniline; 1-(4-aminophenyl)-4-methylpiperidin-4-ol; 1-(4-amino-2-methoxyphenyl)-3-methylazetidin-3-ol; (4-(4-amino-2-methoxyphenyl)morpholin-2-yl)methanol; 2-(4-(4-aminophenyl)piperazin-1-yl)acetaldehyde; 4-(4-(3-fluoropropyl)piperazin-1-yl)aniline; 4-(4,4-difluoropiperidin-1-yl)aniline; 4-(3,3-difluoropiperidin-1-yl)aniline; 4-(8-oxa-3-azabicyclo[3.2.1]octan-3-yl)aniline; 4-amino-N-(tetrahydro-2H-pyran-4-yl)benzamide; (4-aminophenyl)(3-hydroxyazetidin-1-yl)methanone; 3-(4-amino-2-ethoxyphenoxy)-2,2-dimethylpropan-1-ol; 4-(4-ethylpiperazin-1-yl)-3-methoxyaniline; 4-(2,6-dimethylmorpholino)aniline; 1-(4-aminophenyl)-3-methylpiperidin-3-ol; 2-(4-aminophenyl)-N,2-dimethylpropanamide; 4-(1-(2-methoxyethyl)piperidin-4-yl)aniline; 4-(1-(3-fluoropropyl)piperidin-4-yl)aniline; 4-(2-(4-ethylpiperazin-1-yl)ethyl)aniline; 4-amino-N-phenylbenzamide; 3-((dimethylamino)methyl)-1H-indazol-6-amine; 1-(4-aminophenyl)-N,N-dimethylpyrrolidin-3-amine; 4-(4-ethoxypiperidin-1-yl)-3-fluoroaniline; 6-(2,6-dimethylmorpholino)pyridin-3-amine; 6-(1-(2-methoxyethyl)piperidin-4-yl)pyridin-3-amine; 4-methyl-3-(2-morpholinoethoxy)aniline; 6-amino-2,2-dimethyl-2H-benzo[b][1,4]oxazin-3(4H)-one; 3-(5-amino-1H-indazol-1-yl)-N-methylpropanamide; 3-(6-amino-1H-indazol-1-yl)-N-methylpropanamide; 5-amino-N-(2-hydroxyethyl)benzofuran-2-carboxamide; 3-(5-amino-2H-indazol-2-yl)-N-methylpropanamide; N-(3-amino-5-(trifluoromethyl)phenyl)formamide; 4-(benzyloxy)-3-chloroaniline; 6-amino-2,2-difluoro-2H-benzo[b][1,4]oxazin-

3(4H)-one; 5-amino-N-(2-hydroxyethyl)-2,3-dihydrobenzofuran-2-carboxamide; 5-amino-1H-indole-2-carboxylic acid; methyl 5-amino-3-oxo-2,3-dihydrobenzofuran-2-carboxylate; methyl 2-amino-8-methyl-7-oxabicyclo[4.2.0]octa-1,3,5-triene-5-carboxylate; 2-methoxy-5-nitroaniline; N-(3-aminophenyl)pivalamide; N-(4-aminophenyl)-N-methylcyclopropanecarboxamide; N-(4-amino-2-chlorophenyl)-N-methylacetamide; 3-((4-aminophenyl)sulfonyl)propanenitrile; 2-morpholinoquinolin-6-amine; 4-amino-N-(2-methoxyethyl)benzenesulfonamide; 4-amino-N-cyclopropyl-N-methylbenzamide; tert-butyl 4-aminopiperidine-1-carboxylate; 1-(4-aminophenyl)piperidine-2-carboxamide; 3,5-difluoro-4-morpholinoaniline; 4-(4-aminophenyl)thiomorpholine-2,3-dione; 3-(2H-benzo[b][1,4]oxazin-4(3H)-yl)aniline; 3-(quinolin-3-yl)aniline; 3-(quinolin-4-yl)aniline; 3',4'-difluoro-[1,1'-biphenyl]-3-amine; 3-(quinolin-5-yl)aniline; 3-(quinolin-8-yl)aniline; 3-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)aniline; 4-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)aniline; 3-(quinolin-6-yl)aniline; 4-(methylsulfinyl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine; 4-(4-(2-methoxyethyl)piperazin-1-yl)aniline; 2',4'-dimethoxy-[1,1'-biphenyl]-3-amine; 2',3'-dimethoxy-[1,1'-biphenyl]-3-amine; 3',4'-dimethoxy-[1,1'-biphenyl]-3-amine; 1-(4-aminophenyl)-N-methylpyrrolidine-2-carboxamide; tert-butyl 3-aminopiperidine-1-carboxylate; 2-(4-(4-amino-2-methoxyphenyl)piperazin-1-yl)ethan-1-ol; 1-(4-amino-2-ethoxyphenyl)-3-methylazetidin-3-ol; 4-(2-oxa-7-azaspiro[3.5]nonan-7-yl)aniline; 4-(7-oxa-2-azaspiro[3.5]nonan-2-yl)aniline; 1-(4-amino-2-methoxyphenyl)-3-methylpyrrolidin-3-ol; 1-(4-aminophenyl)-N,N-dimethylpiperidin-4-amine; 2-(4-aminophenyl)-1,1,1-trifluoropropan-2-ol; 1-(4-amino-2-fluorophenyl)-3-methylazetidin-3-ol; 4-(1-cyclopropylpiperidin-4-yl)aniline; 4-(1-(2-methoxyethyl)piperidin-4-yl)-3-methylaniline; 3-(4-(4-

aminophenyl)piperidin-1-yl)propanenitrile; 4-amino-N-(3-methoxypropyl)benzenesulfonamide; 2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)aniline; 4-(1-isopropylpiperidin-4-yl)aniline; 2-(3-aminophenoxy)-1-morpholinoethan-1-one; 2-(4-(4-amino-2-methylphenyl)piperidin-1-yl)acetonitrile; 4-(1-(3-fluoropropyl)piperidin-4-yl)-3-methylaniline; ethyl 3-(5-amino-1H-indazol-1-yl)propanoate; ethyl 3-(6-amino-1H-indazol-1-yl)propanoate; 5-amino-N,N-dimethyl-2,3-dihydrobenzofuran-2-carboxamide; ethyl 2-(4-aminophenyl)-2-methylpropanoate; ethyl 3-(5-amino-2H-indazol-2-yl)propanoate; 4-fluoro-3-nitroaniline; 2-fluoro-5-nitroaniline; methyl 5-amino-1H-indole-2-carboxylate; tert-butyl (4-aminophenyl)carbamate; tert-butyl (3-aminophenyl)carbamate; 4-methyl-3-nitroaniline; 2-methyl-5-nitroaniline; 1-(4-aminophenyl)-N-methylpiperidine-2-carboxamide; tert-butyl 4-(aminomethyl)piperidine-1-carboxylate; isopropyl (4-aminophenyl)(methyl)carbamate; 2-(morpholinomethyl)quinolin-6-amine; 4-(pyrrolidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine; benzyl 4-aminocyclohexanecarboxylate; 4-amino-N-cyclobutyl-N-methylbenzamide; 1-(4-aminophenyl)-1H-1,2,3-triazole-4-carboxamide; 2-(4-aminophenoxy)-1-morpholinoethan-1-one; 4-amino-N-cyclopropylbenzenesulfonamide; 2'-(pyrrolidin-3-yl)-[1,1'-biphenyl]-3-amine; 1-(methylsulfonyl)-1H-indazol-6-amine; N-(4-aminophenyl)-2-(dimethylamino)-N-methylacetamide; 5-(4-aminophenyl)-N,N-dimethylpyridin-2-amine; 4-(2-methyl-1-morpholinopropan-2-yl)aniline; 1-(4-amino-2-methoxyphenyl)-4-methylpiperidin-4-ol; 5-amino-2-morpholinobenzamide; (4-aminophenyl)(4-hydroxypiperidin-1-yl)methanone; 3-methoxy-4-(4-(2-methoxyethyl)piperazin-1-yl)aniline; 1-(4-amino-2-methoxyphenyl)-3-methylpiperidin-3-ol; 3-(4-aminophenyl)-1,4-dimethylpiperazin-2-one; 1-(4-(4-aminophenyl)piperidin-1-yl)ethan-1-one; 4-(2-(4-(2-methoxyethyl)piperazin-1-yl)ethyl)aniline; 4-amino-N-(2-morpholinoethyl)benzamide; 2-

(methylsulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-amine; 1-(4-aminophenethyl)-N,N-dimethylpyrrolidin-3-amine; 3-methyl-4-(3-(4-methylpiperazin-1-yl)propoxy)aniline; 3-(4-(4-amino-2-methylphenyl)piperidin-1-yl)propanenitrile; 3-chloro-4-(2,6-dimethylmorpholino)aniline; ethyl 5-amino-1H-indole-2-carboxylate; 2-((3-aminophenyl)imino)-1-morpholinoethan-1-one; 2-((3-aminophenyl)imino)-N-(2,3-dihydroxypropyl)acetamide; 2-((3-aminophenyl)imino)-1-(piperazin-1-yl)ethan-1-one; 4-chloro-3-nitroaniline; 4-(pyrrolidin-1-ylsulfonyl)aniline; 2,2,3,3-tetrafluoro-2,3-dihydrobenzo[b][1,4]dioxin-6-amine; methyl 5-amino-2-(trifluoromethoxy)benzoate; ethyl 7-amino-1H-indole-2-carboxylate; 5-amino-N-isopropyl-2,3-dihydrobenzofuran-2-carboxamide; 1-(4-(5-aminopyridin-2-yl)piperazin-1-yl)ethan-1-one; benzyl 3-(aminomethyl)piperidine-1-carboxylate; 6-amino-N,N-dimethyl-2-naphthamide; 4-(4-aminophenyl)piperazine-1-carboxamide; 1-(4-aminophenyl)piperidine-3-carboxamide; 4-(piperidin-1-yl)-7H-pyrrolo[2,3-d]pyrimidin-2-amine; 4-amino-N-cyclobutylbenzenesulfonamide; 1-(4-aminophenyl)piperidine-4-carboxylic acid; 1-(4-aminophenyl)-4-hydroxypyrrolidine-2-carboxamide; 1-(4-aminophenyl)piperidine-4-carboxamide; 2'-(piperidin-4-yl)-[1,1'-biphenyl]-3-amine; 2'-(piperidin-3-yl)-[1,1'-biphenyl]-3-amine; 2-phenyl-1H-indol-4-amine; 2',5'-dimethoxy-[1,1'-biphenyl]-3-amine; 1-(4-aminophenyl)-N-methylpyrrolidine-3-carboxamide; tert-butyl 2-(4-aminophenoxy)acetate; (4-aminophenyl)(2-(hydroxymethyl)morpholino)methanone; 4-amino-N-methyl-N-(tetrahydro-2H-pyran-4-yl)benzamide; 1-(4-amino-2-ethoxyphenyl)-4-methylpiperidin-4-ol; (4-aminophenyl)(3-hydroxy-3-methylazetidin-1-yl)methanone; 3-(4-(4-aminophenyl)piperidin-1-yl)propane-1,2-diol; 1-(4-aminophenethyl)-N,N-dimethylpiperidin-4-amine; 4-(2,2-dimethylmorpholino)-3-methylaniline; 4-amino-N-(1-

methylpiperidin-4-yl) benzamide; 4-(2-(tetrahydro-2H-pyran-4-yl)thiazol-4-yl) aniline; 4-(2-(piperidin-3-yl)thiazol-4-yl) aniline; 4-(2-(pyridin-3-yl)thiazol-4-yl) aniline; 4-(1-(piperidin-4-yl)-1H-pyrazol-4-yl) aniline; 4-(1-cyclopentylpiperidin-4-yl) aniline; 3-(1,2,3,6-tetrahydropyridin-4-yl)-1H-indazol-6-amine; 4-(1-ethylpiperidin-4-yl)-2-methoxy-5-methylaniline; methyl 4-(4-aminophenyl)piperazine-1-carboxylate; tert-butyl 2-((3-aminophenyl)imino)acetate; (5-aminobenzofuran-2-yl)(pyrrolidin-1-yl)methanone; methyl 3-amino-5-(trifluoromethyl)benzoate; 2-((3-aminophenyl)imino)-N-(3-(dimethylamino)propyl)acetamide; ethyl 6-amino-4H-benzo[b]imidazo[1,5-d][1,4]oxazine-3-carboxylate; 1-(2-amino-5-fluoropyrimidin-4-yl)piperidine-3-carboxamide; 1-(4-(4-aminophenyl)piperazin-1-yl)propan-1-one; 2-amino-4-(m-tolyl)pyrimidine-5-carboxamide; 2-amino-4-(2-(2-hydroxyethyl)piperidin-1-yl)pyrimidine-5-carboxamide; 4-(morpholinosulfonyl) aniline; 4-((thiazol-4-ylmethyl)sulfonyl) aniline; 4-((tetrahydro-2H-pyran-4-yl)sulfonyl) aniline; 3-(2H-1,2,3-triazol-2-yl)-5-(trifluoromethyl) aniline; 1-(4-aminophenyl)-N-methylpiperidine-3-carboxamide; 1-(4-aminophenyl)-N-methylpiperidine-4-carboxamide; 6-amino-2,2,4-trimethyl-2H-benzo[b][1,4]oxazin-3(4H)-one; 1-(4-(4-aminophenyl)-3-methylpiperazin-1-yl)ethan-1-one; 2-(4-aminophenylsulfonamido)acetamide; 4-(phenylsulfonyl) aniline; 3-(2H-1,2,3-triazol-2-yl)-4-(trifluoromethyl) aniline; 3-morpholino-4-(1H-pyrazol-1-yl) aniline; 3-(1H-1,2,3-triazol-1-yl)-4-(trifluoromethyl) aniline; 4-(3-(trifluoromethyl)-1H-pyrazol-1-yl) aniline; 2-(6-amino-2-oxo-3,4-dihydroquinolin-1(2H)-yl)acetic acid; 1-(4-aminophenyl)-4-hydroxy-N-methylpyrrolidine-2-carboxamide; 1-(4-aminophenyl)-N,N-dimethylpyrrolidine-2-carboxamide; 4-(4-

((dimethylamino)methyl)piperidin-1-yl)-5-fluoropyrimidin-2-amine; 7-amino-2,2,4-trimethyl-2H-benzo[b][1,4]oxazin-3(4H)-one; 3-(4-methyl-3,4-dihydro-2H-benzo[b][1,4]oxazin-7-yl)aniline; N-(1-(4-aminophenyl)piperidin-4-yl)acetamide; 1-(4-(4-aminophenyl)-1,4-diazepan-1-yl)ethan-1-one; 1-(4-(4-aminophenyl)-5,6-dihydropyridin-1(2H)-yl)ethan-1-one; 1-(4-(4-aminophenyl)-1,2,3,6-tetrahydropyridin-2-yl)ethan-1-one; 2-(1-(4-aminophenyl)piperidin-4-yl)propan-2-ol; 4-(piperidin-4-yl)-3-(trifluoromethyl)aniline; 4-(1-cyclopropylpiperidin-4-yl)-3-methylaniline; 4-(piperazin-1-yl)-3-(trifluoromethyl)aniline; 1-(4-aminophenethyl)piperidine-3-carboxylic acid; 1-(4-aminophenethyl)piperidine-4-carboxylic acid; 4-morpholino-3-(trifluoromethyl)aniline; 4-amino-N-(4-chlorophenyl)benzamide; 4-(4-(4-aminophenyl)piperidin-1-yl)butan-2-one; 4-(4-aminophenyl)-N-ethylpiperidine-1-carboxamide; 4-(1-isopropylpiperidin-4-yl)-3-methylaniline; 2-(4-(4-amino-2-methylphenyl)piperidin-1-yl)acetamide; (5-aminobenzofuran-2-yl)(morpholino)methanone; 5-amino-N-(2,3-dihydroxypropyl)-2,3-dihydrobenzofuran-2-carboxamide; 5-amino-N-(2,3-dihydroxypropyl)benzofuran-2-carboxamide; 5-amino-N-(1,3-dihydroxypropan-2-yl)benzofuran-2-carboxamide; 2-amino-4-(3-methoxyphenyl)pyrimidine-5-carboxamide; 3-amino-N-methoxy-N-phenylbenzamide; 1-(4-aminophenyl)-N,N-dimethylpiperidine-2-carboxamide; 2-amino-4-(3-ethylphenyl)pyrimidine-5-carboxamide; 1-(1-(2-amino-5-fluoropyrimidin-4-yl)piperidin-4-yl)urea; 2-(1-(2-amino-5-fluoropyrimidin-4-yl)piperidin-4-yl)acetamide; 2-amino-4-(3-(hydroxymethyl)piperidin-1-yl)pyrimidine-5-carboxamide; 4-(4-(pyridin-2-yl)piperazin-1-yl)aniline; 3-(4-phenylpiperazin-1-yl)aniline; 3'-morpholino-[1,1'-biphenyl]-3-amine; 4'-morpholino-[1,1'-biphenyl]-3-amine; 3'-morpholino-[1,1'-biphenyl]-4-amine; 4'-morpholino-[1,1'-biphenyl]-4-amine; 2'-(methylsulfonyl)-[1,1'-biphenyl]-3-amine; 1-(4-(4-aminophenyl)piperazin-1-yl)-2-methoxyethan-1-

one; 2',3',4'-trimethoxy-[1,1'-biphenyl]-3-amine; 4-acetyl-1-(4-aminophenyl)piperazin-2-one; (2-aminocyclohexyl)(tert-butyl)carbamate; 4-(4-(1-methylcyclopropyl)piperazin-1-yl)aniline; 2-(4-aminophenoxy)-1-(3-hydroxy-3-methylazetidin-1-yl)ethan-1-one; (4-aminophenyl)(4-hydroxy-4-methylpiperidin-1-yl)methanone; 4-(2-(4-(pyrrolidin-1-yl)piperidin-1-yl)ethyl)aniline; 1-(4-(4-amino-2-methylphenyl)piperidin-1-yl)ethan-1-one; 4-(1-(methylsulfonyl)pyrrolidin-3-yl)aniline; 4-(1-(2-morpholinoethyl)-1H-pyrazol-4-yl)aniline; 4-(1-methylpiperidin-4-yl)-3-(trifluoromethyl)aniline; 1-(4-(4-aminophenyl)piperidin-1-yl)-2-(ethylamino)ethan-1-one; 4-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)aniline; 4-(4-aminophenyl)piperidine-1-carboxylate; 1-(4-(4-amino-2-methylphenyl)piperidin-1-yl)-3-methoxypropan-2-ol; 2-(4-(4-amino-2-methylphenyl)piperidin-1-yl)-N-methylacetamide; 4-(4-benzylpiperazin-1-yl)aniline; 2-((3-aminophenyl)imino)-1-(4-methylpiperazin-1-yl)ethan-1-one; 2-((3-aminophenyl)imino)-N-(2-morpholinoethyl)acetamide; (5-aminobenzofuran-2-yl)(1,4-diazepan-1-yl)methanone; (6-aminonaphthalen-2-yl)(morpholino)methanone; 1-(4-(4-amino-3-methylphenyl)piperazin-1-yl)ethan-1-one; 1-((1-(2-amino-5-fluoropyrimidin-4-yl)piperidin-4-yl)methyl)urea; 4-(2-amino-5-fluoropyrimidin-4-yl)piperazine-1-carboxamide; 1-(4-(4-amino-2-fluorophenyl)piperazin-1-yl)ethan-1-one; 1-(4-(4-aminophenyl)-2-methylpiperazin-1-yl)ethan-1-one; 4-(1-(methylsulfonyl)piperidin-4-yl)aniline; 4-(2-(4-morpholinopiperidin-1-yl)ethyl)aniline; 3-methyl-4-(1-(piperidin-4-yl)-1H-pyrazol-4-yl)aniline; 4-(1-(methylsulfonyl)piperidin-3-yl)aniline; 1-(4-(4-amino-2-methylphenyl)piperidin-1-yl)-2-(ethylamino)ethan-1-one; 1-(3-(4-aminophenyl)pyrrolidin-1-yl)-2-(dimethylamino)ethan-1-one; ethyl 3-(4-(4-aminophenyl)piperidin-1-yl)propanoate; tert-butyl 6-amino-3,4-dihydroisoquinoline-2(1H)-carboxylate; 4-

(benzyloxy)-3-(trifluoromethyl)aniline; 5-amino-N-(1-hydroxy-2-methylpropan-2-yl)benzofuran-2-carboxamide; 1-((4-aminophenyl)sulfonyl)piperidin-4-ol; 4-(4-(methylsulfonyl)piperazin-1-yl)aniline; 1-(4-aminophenyl)-N,N-dimethylpiperidine-4-carboxamide; 2-amino-4-(4-(2-hydroxyethyl)-1,4-diazepan-1-yl)pyrimidine-5-carboxamide; 4-(4-aminophenyl)-N,N-dimethylpiperazine-1-carboxamide; 1-(4-(4-amino-2-chlorophenyl)piperazin-1-yl)ethan-1-one; 3-(2-amino-5-nitrophenyl)propanamide; 3-(4-(2-amino-5-fluoropyrimidin-4-yl)piperazin-1-yl)-3-oxopropanenitrile; 3-fluoro-4-(3-(trifluoromethyl)-1H-pyrazol-1-yl)aniline; (4-(4-aminophenyl)piperazin-1-yl)(cyclopropyl)methanone; 1-(3'-amino-[1,1'-biphenyl]-4-yl)piperidin-2-one; 1-(3'-amino-[1,1'-biphenyl]-3-yl)pyridin-2(1H)-one; 1-(3'-amino-[1,1'-biphenyl]-4-yl)pyridin-2(1H)-one; 4-(methylsulfonyl)-3-morpholinoaniline; 3'-(methylsulfonyl)-[1,1'-biphenyl]-4-amine; 4'-(methylsulfonyl)-[1,1'-biphenyl]-4-amine; 3'-(methylsulfonyl)-[1,1'-biphenyl]-3-amine; 4'-(methylsulfonyl)-[1,1'-biphenyl]-3-amine; methyl 4-(4-aminophenyl)-3-methylpiperazine-1-carboxylate; N-(4-aminophenyl)-2-(benzyloxy)-N-methylacetamide; 2-(4-amino-N-methylphenylsulfonamido)acetic acid; 1-(4-aminophenyl)-4-hydroxy-N,N-dimethylpyrrolidine-2-carboxamide; 1-(4-aminophenyl)-N,N-dimethylpiperidine-3-carboxamide; N-(1-(4-aminophenyl)piperidin-4-yl)-N-methylacetamide; 1-(4-(4-aminophenyl)piperidin-1-yl)-2-(dimethylamino)ethan-1-one; 4-(1-(4,4,4-trifluorobutyl)piperidin-4-yl)aniline; 3-methyl-4-(6-(piperazin-1-yl)pyridin-3-yl)aniline; 1-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)-1H-pyrazol-4-amine; 1-(4-(5-aminopyridin-2-yl)piperidin-1-yl)-2-(dimethylamino)ethan-1-one; 4-(1-(ethylsulfonyl)piperidin-4-yl)aniline; 2-((3-aminophenyl)imino)-N-(2-(benzylamino)ethyl)acetamide; 1-(4-(4-aminophenyl)-3-

methylypiperazin-1-yl)-2-methoxyethan-1-one; N-(1-(2-amino-5-fluoropyrimidin-4-yl)piperidin-4-yl)-2-cyanoacetamide; 4-((4-(methylsulfonyl)piperazin-1-yl)methyl)aniline; 4-(4-(methylsulfonyl)-1,4-diazepan-1-yl)aniline; 4-(1-(methylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)aniline; 4-(2-(methylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)aniline; 2-amino-4-(methyl(1-methylpiperidin-4-yl)amino)pyrimidine-5-carboxamide; 4-(4-(ethylsulfonyl)piperazin-1-yl)aniline; 1-(4-(4-aminobenzoyl)piperazin-1-yl)ethan-1-one; 4-(2-methyl-4-(methylsulfonyl)piperazin-1-yl)aniline; 2,6-diisopropyl-4-phenoxyaniline; 2-methyl-4-(4-(methylsulfonyl)piperazin-1-yl)aniline; 3-methyl-4-(1-(2-morpholinoethyl)-1H-pyrazol-4-yl)aniline; 4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)aniline; 1-(4-(4-amino-2-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethan-1-one; 5-amino-2-(morpholine-4-carbonyl)benzofuran-3(2H)-one; tert-butyl (6-amino-4H-chromen-4-yl)carbamate; 2-amino-4-(3,5-dimethylphenyl)pyrimidine-5-carboxamide; (1-(4-aminophenyl)piperidin-4-yl)(pyrrolidin-1-yl)methanone; 4-acetyl-1-(4-aminophenyl)piperazine-2-carboxamide; 3-(4-(4-aminophenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol; tert-butyl 3-(4-aminophenyl)pyrrolidine-1-carboxylate; ethyl 1-(3-aminobenzoyl)piperidine-4-carboxylate; (1-(4-aminophenyl)piperidin-4-yl)(morpholino)methanone; (1-(4-aminophenyl)piperidin-4-yl)(piperidin-1-yl)methanone; 4-acetyl-1-(4-aminophenyl)-N-methylpiperazine-2-carboxamide; tert-butyl 4-(4-aminophenyl)piperidine-1-carboxylate; butyl 2-(4-amino-N-methylphenylsulfonamido)acetate; 4-(4-(cyclopropylsulfonyl)piperazin-1-yl)aniline; N-(1-(4-aminophenyl)piperidin-4-yl)-N-methylmethanesulfonamide; 4-(4-(ethylsulfonyl)-2-methylpiperazin-1-yl)aniline; 3-methyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)aniline; (4-aminophenyl)(4-(methylsulfonyl)piperazin-1-yl)methanone; 4-(4-

(cyclopropylsulfonyl)-2-methylpiperazin-1-yl)aniline; 1-(4-(4-aminophenyl)piperidin-1-yl)-2-(methylsulfonyl)ethan-1-one; methyl 4-((6-amino-2H-indazol-2-yl)methyl)-3-methoxybenzoate; methyl 4-((6-amino-1H-indazol-1-yl)methyl)-3-methoxybenzoate; 4-((6-amino-1H-indazol-1-yl)methyl)-3-methoxy-N-methylbenzamide; 2-isopropyl-5-methylcyclohexyl 5-amino-2,3-dihydrobenzofuran-2-carboxylate; ethyl 4-amino-3-(2-amino-5-nitrobenzyl)-4-oxobutanoate; 3,4-bis(3-(trifluoromethyl)-1H-pyrazol-1-yl)aniline.

Especially preferred compounds of formula (I) are:

N-(1H-indazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-benzo[d][1,2,3]triazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-phenyl-N-(4-(piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; 3-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzamide; N-(3,4-dimethoxyphenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-methoxyphenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzoic acid; 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzamide; 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzonitrile; N-(1H-indazol-5-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-5-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 2-((5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzamide; N-(1H-indazol-6-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-

yl)benzene-1,4-diamine; N-(3,4-dimethoxyphenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; 5-(3,4-dimethoxyphenyl)-N-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1-(5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine; N,5-bis(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; N-(1H-indazol-6-yl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; N-(4-chlorophenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-6-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; N-(3,4-dimethoxyphenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-6-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; 5-phenyl-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-phenyl-N-(4-(pyrrolidin-1-yl)phenyl)-

1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(1H-indazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-pyrazol-3-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(pyridin-4-yl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N,5-di(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; N-(1H-pyrazol-3-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(pyrrolidin-1-yl)phenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-5-yl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(1-methylpiperidin-4-yl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; 5-(1-methylpiperidin-4-yl)-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(1-methylpiperidin-4-yl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(4-methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-

methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1-(5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine; N-(5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 3-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; 5-(4-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-6-yl)-5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; N-(3,4-dimethoxyphenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; N-(1H-indazol-6-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1-(5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine; N-(1H-indazol-6-yl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 3-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; 3-((5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; N-(1H-indazol-6-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol;

N- (5- (thiophen-3-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) -3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 3-((5- (thiophen-3-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) phenol; N- (5-cyclopropyl-1H-pyrazol-3-yl) -5- (4-methoxyphenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (5- (4-methoxyphenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) -3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N- (5-cyclopropyl-1H-pyrazol-3-yl) -5- (3-methoxyphenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (5- (3-methoxyphenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) -3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N- (5-cyclopropyl-1H-pyrazol-3-yl) -5- (thiophen-2-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (5- (thiophen-2-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) -3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N- (5-cyclopropyl-1H-pyrazol-3-yl) -5- (1-methylpiperidin-4-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (5- (1-methylpiperidin-4-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) -3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 3-((5- (1-methylpiperidin-4-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) phenol; 5- (benzo[d]thiazol-2-yl) -N- (4- (4-methylpiperazin-1-yl) phenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5- (benzo[d]thiazol-2-yl) -N- (3,4-dimethoxyphenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (5- (benzo[d]thiazol-2-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) -3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 5- (benzo[d]thiazol-2-yl) -N- (1H-indazol-6-yl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (4- (4-methylpiperazin-1-yl) phenyl) -5- (3- (pyridin-3-yl) phenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (5-cyclopropyl-1H-pyrazol-3-yl) -5- (3- (pyridin-3-yl) phenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (3,4-dimethoxyphenyl) -5- (3- (pyridin-3-yl) phenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (4-morpholinophenyl) -5- (3- (pyridin-3-yl) phenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-amine; N- (5- (3- (pyridin-3-yl) phenyl) -1H-pyrazolo[4,3-d]pyrimidin-7-yl) -3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N- (1H-indazol-6-yl) -5- (3-

(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 2-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N-(1H-indazol-6-yl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N3-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,3-diamine; N1-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)cyclohexane-1,2-diamine; N1-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,3-diamine; N-(6-(4-methylpiperazin-1-yl)pyridin-3-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-ethylpiperazin-1-yl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-((dimethylamino)methyl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,5-dimethoxyphenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,5-dimethoxyphenyl)-N-(4-(4-ethylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2,5-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-aminoethyl)-3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzamide; N-(3,4-dimethoxyphenyl)-5-(imidazo[1,2-a]pyridin-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-ethylpiperazin-1-yl)phenyl)-5-(imidazo[1,2-a]pyridin-2-yl)-1H-

pyrazolo[4,3-d]pyrimidin-7-amine; 5-(imidazo[1,2-a]pyridin-2-yl)-N-(1H-indazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(imidazo[1,2-a]pyridin-2-yl)-N-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 4-((5-(imidazo[1,2-a]pyridin-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzoic acid; 5-(1H-benzo[d]imidazol-2-yl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)methyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-methoxyphenyl)-N-(4-(pyrrolidin-1-ylmethyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)-3-(m-tolyl)urea; (4-methylpiperazin-1-yl)(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; pyrrolidin-1-yl(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; N-(4-thiomorpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 4-(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)thiomorpholine 1,1-dioxide; 1-(4-(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; (3-methoxy-4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(4-(4-methylpiperazin-1-yl)piperidin-1-yl)methanone; 5-(1H-imidazol-5-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(1H-imidazol-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; pyrrolidin-1-yl(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; 4-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)thiomorpholine 1,1-dioxide; 1-(4-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; N-

(4-thiomorpholinophenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; (4-methylpiperazin-1-yl)(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; 2,2-dimethyl-N-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-amine; 5-(3-(2H-tetrazol-5-yl)phenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; (7-((2-(dimethylamino)ethyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; (7-((1-methylpiperidin-4-yl)methyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; (7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; (7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; (4-((5-benzoyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(pyrrolidin-1-yl)methanone; 7-((5-benzoyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; morpholino(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; 6-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]thiazin-3(4H)-one; 7-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; (4-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(pyrrolidin-1-yl)methanone; N-(4-(4-(tert-butyl)piperazin-1-yl)phenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; N-(4-(4-(tert-butyl)piperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 8-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-4,5-

dihydro-1H-benzo[b]azepin-2(3H)-one; 6-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]thiazin-3(4H)-one; 1-methyl-N-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-1,2,3,4-tetrahydroquinolin-7-amine; 2-methyl-2-(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)propanenitrile; 5-(4-methoxyphenyl)-N-(4-thiomorpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 4-(4-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)thiomorpholine 1,1-dioxide; 1-(4-(4-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; 6-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; N-(4-(4-(tert-butyl)piperazin-1-yl)phenyl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 2,2-dimethyl-N-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-amine; 2-(3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)acetic acid; 5-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)thiophene-3-carboxylic acid; 2-(3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)acetic acid; 6-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)picolinic acid; N-(4-(pentafluorosulfanyl)phenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-(((4-methylbenzyl)amino)methyl)phenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-(3-(dimethylamino)propoxy)phenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 3-(7-((3,4-

dimethoxyphenyl) amino) -1H-pyrazolo[4,3-d]pyrimidin-5-yl) benzenesulfonamide; 1-(4-(4-((5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) phenyl) piperazin-1-yl) ethan-1-one; N-(3,4-dimethoxyphenyl)-5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-3,4-dihydroquinolin-2(1H)-one; N-(3-(7-((4-(4-methylpiperazin-1-yl) phenyl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl) phenyl) methanesulfonamide; N-(3-(7-((4-(piperazin-1-yl) phenyl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl) phenyl) methanesulfonamide; N-(3-(7-((3,4-dimethoxyphenyl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl) phenyl) methanesulfonamide; N-(3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl) phenyl) methanesulfonamide; N-(3-(7-((4-(4-acetylpiperazin-1-yl) phenyl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl) phenyl) methanesulfonamide; 1-(4-(4-((5-(3-(2H-tetrazol-5-yl) phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) phenyl) piperazin-1-yl) ethan-1-one; 5-(3-(2H-tetrazol-5-yl) phenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 3-(7-((3,4-dimethoxyphenyl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl) benzoic acid; 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl) benzoic acid; 7-((5-(3-(2H-tetrazol-5-yl) phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-3,4-dihydroquinolin-2(1H)-one; N-(4-(4-methylpiperazin-1-yl) phenyl)-5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-(4-((5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) phenyl) piperazin-1-yl) ethan-1-one; N-(3,4-dimethoxyphenyl)-5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl) phenyl)-5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-(4-((5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) phenyl) piperazin-1-yl) ethan-1-one; N-(3,4-dimethoxyphenyl)-5-(oxazol-4-yl)-1H-

pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-(4-((5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; N-(3,4-dimethoxyphenyl)-5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; 7-((5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; 7-((5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; N-(4-(1-methylpiperidin-4-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; methyl 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; methyl 4-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((2-oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-8-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 3-(7-((2-oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-8-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; N-(3,4-dimethoxyphenyl)-5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-(pentafluorosulfanyl)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-

amine; 7-((5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; N-(3,4-dimethoxyphenyl)-5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-(methylthio)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-(trifluoromethyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(3-(trifluoromethyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-chlorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-chlorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; 3-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 3-(7-((3-methoxy-4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 4-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; methyl 4-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-


d]pyrimidin-5-yl)benzoate; 4-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; 4-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-(4-((5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; 5-(1H-indazol-6-yl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(1H-indazol-6-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; 5-(3-(methylsulfonyl)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N,5-bis(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-(methylsulfonyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 2-((3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)sulfonyl)ethan-1-ol; 1-(4-(4-((5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; 7-((5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; N-(4-morpholinophenyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N,N-dimethyl-4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide; N-cyclopropyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide; N-(3-(2H-1,2,3-triazol-2-yl)phenyl)-5-(thiophen-2-yl)-1H-

pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(2H-1,2,3-triazol-2-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N,N-dimethyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide; N-methyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide; N-(3-(morpholinosulfonyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(thiophen-2-yl)-N-(3-((trifluoromethyl)sulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-(methylsulfinyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; N-(3,4-dimethoxyphenyl)-5-(4-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(4-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,5-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(2-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,6-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,6-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2,6-difluorophenyl)-

1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,6-difluorophenyl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,4-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,4-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 6-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 6-(7-((2-methyl-1H-benzo[d]imidazol-5-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-(methylsulfonyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-((dimethylamino)methyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; morpholino(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; N-(4-(morpholinomethyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-fluoro-4-morpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; (4-(4-methylpiperazin-1-yl)piperidin-1-yl)(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; 6-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 6-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(4-morpholinophenyl)-1H-

pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(2-methyl-1H-benzo[d]imidazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-methylisoindolin-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1,3-dihydroisobenzofuran-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-(tert-butyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; methyl 3-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((1H-indazol-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 5-(3-aminophenyl)-N-(4-((dimethylamino)methyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(2-methylisoindolin-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-((dimethylamino)methyl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-((dimethylamino)methyl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine.

Preferably, the following compounds are excluded from the scope of the present application:



Especially preferably, the compounds disclosed in WO 98/29413 are excluded from the scope of the present application.

The present invention further provides pharmaceutical compositions comprising one or more compounds of formula (I), (Ia), (Ib) and/or (Ic) as defined herein or a pharmaceutically acceptable ester, prodrug, hydrate, solvate or salt thereof, optionally in combination with a pharmaceutically acceptable carrier.

The pharmaceutical composition optionally comprises one or more of the following compounds or is administered in combination with one or more of these compounds:

Chlorhexidine; polynoxylin; domiphen; oxyquinoline; neomycin; miconazole; natamycin; various; hexetidine; tetracycline; mepartricin; metronidazole; clotrimazole; chlortetracycline; doxycycline; minocycline; triamcinolone; dexamethasone; hydrocortisone; epinephrine; benzydamine; adrenalone; amlexanox; becaplermin; algeldrate; aloglutamol; magaldrate; almagate; hydrotalcite; almasilate; cimetidine; ranitidine; famotidine; nizatidine; niperotidine; roxatidine; lafutidine; misoprostol; enprostil; omeprazole; pantoprazole; lansoprazole; rabeprazole; esomeprazole; carbenoxolone; sucralfate; pirenzepine; proglumide; gefarnate; sulglicotide; acetoxolone; zolimidine; troxipide; oxyphencyclimine; camylofin; mebeverine; trimebutine; rociverine; dicycloverine; dihexyverine; difemerine; piperidolate; benzilone; glycopyrronium; oxyphenonium; penthienate; propantheline; methantheline; tridihexethyl; isopropamide; hexocyclium; poldine; mepenzolate; bevonium; pipenzolate; diphemanil; fenpiverinium; dimethylaminopropionylphenothiazine; nicofetamide; tiropramide; papaverine; drotaverine; moxaverine; alosetron; tegaserod; cilansetron; prucalopride; fenpiprane; diisopromine; chlorbenzoxamine; pinaverium; fenoverine; idanpramine; proxazole; alverine; trepibutone; isometheptene; caroverine; phloroglucinol; silicones; trimethyldiphenylpropylamine; atropine; hyoscyamine; butylscopolamine; methylatropine; methylscopolamine; fentonium; metoclopramide; cisapride; domperidone; bromopride; alizapride; clebopride; ondansetron; granisetron; tropisetron; dolasetron; palonosetron; scopolamine; chlorobutanol; metopimazine; dronabinol; nabilone; aprepitant; casopitant; piprozolin; hymecromone; cyclobutyrol; silymarin; citiolone; epomediol; oxyphenisatine; bisacodyl; dantron; phenolphthalein; cascara; bisoxatin; ethulose; sterculia; linseed; methylcellulose; lactulose; lactitol;

pentaerithrityl; macrogol; mannitol; sorbitol; glycerol; oil; alvimopan; lubiprostone; nystatin; streptomycin; paromomycin; kanamycin; vancomycin; colistin; rifaximin; phthalylsulfathiazole; sulfaguanidine; succinylsulfathiazole; broxyquinoline; acetarsol; nifuroxazide; nifurzide; pectin; kaolin; crosopovidone; attapulgit; diosmectite; diphenoxylate; opium; loperamide; difenoxin; prednisolone; prednisone; betamethasone; tixocortol; budesonide; beclometasone; sulfasalazine; mesalazine; olsalazine; balsalazide; ceratonia; racecadotril; phentermine; fenfluramine; amfepramone; dexfenfluramine; mazindol; etilamfetamine; cathine; clobenzorex; mefenorex; sibutramine; orlistat; rimonabant; diastase; pepsin; tilactase; phenformin; metformin; buformin; glibenclamide; chlorpropamide; tolbutamide; glibornuride; tolazamide; carbutamide; glipizide; gliquidone; gliclazide; metahexamide; glisoxepide; glimepiride; acetohexamide; glymidine; acarbose; miglitol; voglibose; troglitazone; rosiglitazone; pioglitazone; sitagliptin; vildagliptin; saxagliptin; alogliptin; repaglinide; nateglinide; exenatide; pramlintide; benfluorex; liraglutide; mitiglinide; tolrestat; betacarotene; ergocalciferol; dihydrotachysterol; alfacalcidol; calcitriol; colecalciferol; calcifediol; sulbutiamine; benfotiamine; nicotinamide; biotin; inositol; tocofersolan; dexpanthenol; pantethine; androstanolone; stanozolol; metandienone; metenolone; oxymetholone; quinbolone; prasterone; oxandrolone; norethandrolone; nandrolone; ethylestrenol; levocarnitine; ademetonine; glutamine; mercaptamine; betaine; alglucerase; imiglucerase; laronidase; sacrosidase; galsulfase; idursulfase; nitisinone; miglustat; sapropterin; dicoumarol; phenindione; warfarin; phenprocoumon; acenocoumarol; clorindione; diphenadione; tiocloamarol; heparin; dalteparin; enoxaparin; nadroparin; parnaparin; reviparin; danaparoid; tinzaparin; sulodexide;

bemiparin; ditazole; cloricromen; picotamide; clopidogrel; ticlopidine; dipyridamole; epoprostenol; indobufen; iloprost; abciximab; aloxiprin; eptifibatide; tirofiban; triflusal; beraprost; treprostinil; prasugrel; streptokinase; alteplase; anistreplase; urokinase; fibrinolysin; brinase; reteplase; saruplase; ancrod; tenecteplase; desirudin; lepirudin; argatroban; melagatran; ximelagatran; bivalirudin; defibrotide; fondaparinux; rivaroxaban; camostat; phytomenadione; menadione; thrombin; collagen; etamsylate; carbazochrome; batroxobin; romiplostim; eltrombopag; dextriferron; cyanocobalamin; hydroxocobalamin; cobamamide; mecobalamin; erythropoietin; albumin; dextran; hydroxyethylstarch; erythrocytes; thrombocytes; carbohydrates; electrolytes; trometamol; carbamide; cetylpyridinium; nitrofurazone; sulfamethizole; taurolidine; noxytiolin; glucose; glycine; lysine; hyaluronidase; chymotrypsin; trypsin; desoxyribonuclease; bromelains; hematin; acetyldigitoxin; acetyldigoxin; digitoxin; digoxin; deslanoside; metildigoxin; gitoformate; proscillaridin; g-strophanthin; cymarin; peruvoside; quinidine; procainamide; disopyramide; sparteine; ajmaline; prajmaline; lorajmine; lidocaine; mexiletine; tocainide; aprindine; propafenone; flecainide; lorcainide; encainide; amiodarone; bunaftine; dofetilide; ibutilide; tedisamil; moracizine; cibenzoline; etilefrine; isoprenaline; norepinephrine; dopamine; norfenefrine; phenylephrine; dobutamine; oxedrine; metaraminol; methoxamine; mephentermine; dimetofrine; prenalterol; dopexamine; gepefrine; ibopamine; midodrine; octopamine; fenoldopam; cafedrine; arbutamine; theodrenaline; amrinone; milrinone; enoximone; bucladesine; angiotensinamide; xamoterol; levosimendan; propatyl nitrate; trol nitrate; tenitramine; flosequinan; prenylamine; oxyfedrine; benziodarone; carbocromen; hexobendine; etafenone; heptaminol; imolamine; dilazep; trapidil; molsidomine;

efloxate; cinepazet; cloridarol; nicorandil; linsidomine;
nesiritide; alprostadil; camphora; indometacin;
creatinolfosfate; fosfocreatine; ubidecarenone; adenosine;
tiracizine; acadesine; trimetazidine; ibuprofen; ivabradine;
ranolazine; icatibant; regadenoson; rescinnamine; reserpine;
deserpidine; methoserpidine; bietaserpine; clonidine;
guanfacine; tolondine; moxonidine; rilmenidine; trimetaphan;
mecamylamine; prazosin; indoramin; trimazosin; doxazosin;
urapidil; betanidine; guanethidine; guanoxan; debrisoquine;
guanoclor; guanazodine; guanoxabenz; diazoxide; dihydralazine;
hydralazine; endralazine; cadralazine; minoxidil;
nitroprusside; pinacidil; veratrum; metirosine; pargyline;
ketanserin; bosentan; ambrisentan; sitaxentan;
bendroflumethiazide; hydroflumethiazide; hydrochlorothiazide;
chlorothiazide; polythiazide; trichlormethiazide;
cyclopenthiazide; methyclothiazide; cyclothiazide; mebutizide;
quinethazone; clopamide; chlortalidone; mefruside;
clofenamide; metolazone; meticrane; xipamide; indapamide;
clorexolone; fenquizone; mersalyl; theobromine; cicletanine;
furosemide; bumetanide; piretanide; torasemide; muzolimine;
etozolin; spironolactone; canrenone; eplerenone; amiloride;
triamterene; tolvaptan; conivaptan; isoxsuprine; buphenine;
bamethan; phentolamine; tolazoline; ciclonicate; pentifylline;
pentoxifylline; nicergoline; dihydroergocristine;
kallidinogenase; cyclandelate; phenoxybenzamine; vincamine;
moxisylyte; bencyclane; vinburnine; suloctidil; buflomedil;
naftidrofuryl; butalamine; visnadine; cetiedil; cinepazide;
ifenprodil; azapetine; fasudil; fluorometholone;
fluocortolone; fluocinonide; tetracaine; benzocaine;
cinchocaine; procaine; oxetacaine; pramocaine; tribenoside;
organo-heparinoid; polidocanol; phenol; rutoside; monoxerutin;
diosmin; troxerutin; hidrosmin; alprenolol; oxprenolol;
pindolol; propranolol; timolol; sotalol; nadolol; mepindolol;

carteolol; tertatolol; bopindolol; bupranolol; penbutolol; cloranolol; practolol; metoprolol; atenolol; acebutolol; betaxolol; bevantolol; bisoprolol; celiprolol; esmolol; epanolol; s-atenolol; nebivolol; talinolol; labetalol; carvedilol; amlodipine; felodipine; isradipine; nicardipine; nifedipine; nimodipine; nisoldipine; nitrendipine; lacidipine; nilvadipine; manidipine; barnidipine; lercanidipine; cilnidipine; benidipine; mibefradil; verapamil; gallopamil; diltiazem; fendiline; bepridil; lidoflazine; perhexiline; captopril; enalapril; lisinopril; perindopril; ramipril; quinapril; benazepril; cilazapril; fosinopril; trandolapril; spirapril; delapril; moexipril; temocapril; zofenopril; imidapril; losartan; eprosartan; valsartan; irbesartan; tasosartan; candesartan; telmisartan; remikiren; aliskiren; simvastatin; lovastatin; pravastatin; fluvastatin; atorvastatin; cerivastatin; rosuvastatin; pitavastatin; clofibrate; bezafibrate; gemfibrozil; fenofibrate; simfibrate; ronifibrate; ciprofibrate; etofibrate; clofibrade; colestyramine; colestipol; colextran; colesevelam; niceritrol; nicofuranose; acipimox; dextrothyroxine; probucol; tiadenol; meglutol; policosanol; ezetimibe; hachimycin; pecilocin; pyrrolnitrin; griseofulvin; econazole; chlormidazole; isoconazole; tiabendazole; tioconazole; ketoconazole; sulconazole; bifonazole; oxiconazole; fenticonazole; omoconazole; sertaconazole; fluconazole; flutrimazole; bromochlorosalicylanilide; methylrosaniline; tribromometacresol; chlorphenesin; ticlatone; sulbentine; haloprogin; ciclopirox; terbinafine; amorolfine; dimazole; tolnaftate; tolclate; flucytosine; naftifine; butenafine; octinoxate; dextranomer; crilanomer; enoxolone; collagenase; thonzylamine; mepyramine; thenalidine; tripelennamine; chloropyramine; promethazine; tolpropamine; dimetindene; clemastine; bamipine; isothipendyl; diphenhydramine;

chlorphenoxamine; oxybuprocaine; quinisocaine; dithranol;
trioxysalen; methoxsalen; calcipotriol; tacalcitol;
tazarotene; bergapten; etretinate; acitretin; demeclocycline;
oxytetracycline; chloramphenicol; bacitracin; gentamicin;
tyrothricin; mupirocin; virginiamycin; amikacin; retapamulin;
sulfathiazole; mafenide; sulfanilamide; sulfamerazine;
idoxuridine; tromantadine; aciclovir; podophyllotoxin;
inosine; penciclovir; lysozyme; ibacitabine; edoxudine;
imiquimod; docosanol; methylprednisolone; clobetasone;
flumetasone; fluocortin; fluperolone; fluprednidene; desonide;
alclometasone; clocortolone; flucolorolone; desoximetasone;
diflucortolone; fludroxycortide; diflorasone; amcinonide;
halometasone; mometasone; fluticasone; prednicarbate;
difluprednate; ulobetasol; clobetasol; halcinonide;
aminoacridine; euflavine; dibrompropamidine; propamidine;
hexamidine; polihexanide; hexachlorophene; policresulen;
triclosan; chloroxylenol; biphenylol;
iodine/octylphenoxypolyglycolether; povidone-iodine; iodine;
diiodohydroxypropane; dequalinium; chlorquinaldol; clioquinol;
benzalkonium; cetrimonium; cetrime; mercurochrome;
thiomersal; silver; eosin; propanol; isopropanol; ethanol;
framycetin; benzododecinium; iodoform; bithionol; sulfur;
tioxolone; mesulfen; tretinoin; retinol; adapalene;
isotretinoin; motretinide; clindamycin; erythromycin;
meclocycline; resorcinol; dapsone; ichtasol; xenysalate;
others; tacrolimus; pimecrolimus; mequinol; tiratricol;
oxaceprol; finasteride; hydroquinone; monobenzene;
eflornithine; diclofenac; alitretinoin; candicidin;
carfecillin; pentamycin; diiodohydroxyquinoline; Sulfonamides:
sulfatolamide; ornidazole; azanidazole; propenidazole;
butoconazole; terconazole; clodantoin; nifuratel;
furazolidone; protiofate; methylergometrine; ergometrine;
dinoprost; dinoprostone; gemeprost; carboprost; sulprostone;

ritodrine; fenoterol; bromocriptine; lisuride; cabergoline; quinagolide; metergoline; terguride; naproxen; flunoxaprofen; atosiban; norethisterone; lynestrenol; levonorgestrel; quingestanol; megestrol; medroxyprogesterone; norgestrienone; etonogestrel; desogestrel; fluoxymesterone; methyltestosterone; testosterone; mesterolone; ethinylestradiol; estradiol; estriol; chlorotrianisene; estrone; promestriene; dienestrol; diethylstilbestrol; methallenestril; moxestrol; tibolone; gestonorone; hydroxyprogesterone; progesterone; dydrogesterone; medrogestone; nomegestrol; demegestone; chlormadinone; promegestone; allylestrenol; ethisterone; etynodiol; methylestrenolone; urofollitropin; cyclofenil; clomifene; epimestrol; cyproterone; danazol; gestrinone; mifepristone; raloxifene; bazedoxifene; emepronium; flavoxate; meladrazine; oxybutynin; terodiline; propiverine; tolterodine; solifenacin; trospium; darifenacin; fesoterodine; sildenafil; yohimbine; apomorphine; tadalafil; vardenafil; phenazopyridine; succinimide; dapoxetine; alfuzosin; tamsulosin; terazosin; silodosin; dutasteride; corticotropin; tetracosactide; thyrotropin; somatropin; somatrem; mecasermin; sermorelin; pegvisomant; vasopressin; desmopressin; lypressin; terlipressin; ornipressin; argipressin; demoxytocin; oxytocin; carbetocin; gonadorelin; nafarelin; histrelin; somatostatin; octreotide; lanreotide; vapreotide; ganirelix; cetrorelix; aldosterone; fludrocortisone; desoxycortone; paramethasone; cortisone; prednylidene; rimexolone; deflazacort; cloprednol; meprednisone; cortivazol; trilostane; methylthiouracil; propylthiouracil; benzylthiouracil; carbimazole; thiamazole; diiodotyrosine; dibromotyrosine; glucagon; teriparatide; elcatonin; cinacalcet; paricalcitol; doxercalciferol; lymecycline; metacycline; rolitettracycline; penimepicycline; clomocycline; tigecycline; thiamphenicol; ampicillin;

pivampicillin; carbenicillin; amoxicillin; carindacillin;
bacampicillin; epicillin; pivmecillinam; azlocillin;
mezlocillin; mecillinam; piperacillin; ticarcillin;
metampicillin; talampicillin; sulbenicillin; temocillin;
hetacillin; benzylpenicillin; phenoxymethylpenicillin;
propicillin; azidocillin; pheneticillin; penamecillin;
clometocillin; dicloxacillin; cloxacillin; meticillin;
oxacillin; flucloxacillin; sulbactam; tazobactam;
sultamicillin; cefalexin; cefaloridine; cefalotin; cefazolin;
cefadroxil; cefazedone; cefatrizine; cefapirin; cefradine;
cefacetrile; cefroxadine; ceftezole; cefoxitin; cefuroxime;
cefamandole; cefaclor; cefotetan; cefonicid; cefotiam;
loracarbef; cefmetazole; cefprozil; ceforanide; cefotaxime;
ceftazidime; cefsulodin; ceftriaxone; cefmenoxime; latamoxef;
ceftizoxime; cefixime; cefodizime; cefetamet; cefpiramide;
cefoperazone; cefpodoxime; ceftibuten; cefdinir; cefditoren;
cefcapene; cefepime; cefpirome; cefozopran; meropenem;
ertapenem; doripenem; biapenem; trimethoprim; brodimoprim;
iclaprim; sulfaisodimidine; sulfadimidine; sulfapyridine;
sulfafurazole; sulfathiourea; sulfamethoxazole; sulfadiazine;
sulfamoxole; sulfadimethoxine; sulfalene; sulfametomidine;
sulfametoxydiazine; sulfamethoxypyridazine; sulfaperin;
sulfaphenazole; sulfamazone; spiramycin; midecamycin;
oleandomycin; roxithromycin; josamycin; troleandomycin;
clarithromycin; azithromycin; miocamycin; rokitamycin;
dirithromycin; flurithromycin; telithromycin; lincomycin;
pristinamycin; quinupristin/dalfopristin; streptoduocin;
tobramycin; netilmicin; sisomicin; dibekacin; ribostamycin;
isepamicin; arbekacin; ofloxacin; ciprofloxacin; pefloxacin;
enoxacin; temafloxacin; norfloxacin; lomefloxacin; fleroxacin;
sparfloxacin; rufloxacin; grepafloxacin; levofloxacin;
trovafloxacin; moxifloxacin; gemifloxacin; gatifloxacin;
prulifloxacin; pazufloxacin; garenoxacin; rosoxacin;

cinoxacin; flumequine; teicoplanin; telavancin; dalbavancin; oritavancin; tinidazole; nitrofurantoin; nifurtimol; fosfomicin; xibornol; clofoctol; spectinomycin; methenamine; nitroxoline; linezolid; daptomycin; itraconazole; voriconazole; posaconazole; caspofungin; micafungin; anidulafungin; cycloserine; rifampicin; rifamycin; rifabutin; rifapentine; capreomycin; isoniazid; protionamide; tiocarlide; ethionamide; pyrazinamide; ethambutol; terizidone; morinamide; clofazimine; metisazone; vidarabine; ribavirin; ganciclovir; famciclovir; valaciclovir; cidofovir; valganciclovir; brivudine; rimantadine; foscarnet; fosfonet; saquinavir; indinavir; ritonavir; nelfinavir; amprenavir; lopinavir; fosamprenavir; atazanavir; tipranavir; darunavir; zidovudine; didanosine; zalcitabine; stavudine; lamivudine; abacavir; emtricitabine; entecavir; telbivudine; clevidine; nevirapine; delavirdine; efavirenz; etravirine; zanamivir; oseltamivir; moroxydine; pleconaril; enfuvirtide; raltegravir; maraviroc; maribavir; palivizumab; nebacumab; diphtheria-poliomyelitis-tetanus; diphtheria-pertussis-poliomyelitis-tetanus; diphtheria-rubella-tetanus; cyclophosphamide; chlorambucil; melphalan; chlormethine; ifosfamide; trofosfamide; prednimustine; bendamustine; busulfan; treosulfan; mannosulfan; thiotepa; triaziquone; carboquone; carmustine; lomustine; semustine; streptozocin; fotemustine; nimustine; ranimustine; etoglucid; mitobronitol; pipobroman; temozolomide; dacarbazine; methotrexate; raltitrexed; pemetrexed; pralatrexate; mercaptopurine; tioguanine; cladribine; fludarabine; clofarabine; nelarabine; cytarabine; fluorouracil; tegafur; carmofur; gemcitabine; capecitabine; azacitidine; decitabine; vinblastine; vincristine; vindesine; vinorelbine; vinflunine; etoposide; teniposide; demecolcine; paclitaxel; docetaxel; trabectedin; dactinomycin; doxorubicin; daunorubicin; epirubicin; aclarubicin; zorubicin; idarubicin;

mitoxantrone; pirarubicin; valrubicin; bleomycin; plicamycin; mitomycin; ixabepilone; cisplatin; carboplatin; oxaliplatin; satraplatin; procarbazine; edrecolomab; rituximab; trastuzumab; alemtuzumab; gemtuzumab; cetuximab; bevacizumab; panitumumab; catumaxomab; ofatumumab; temoporfin; efaproxiral; imatinib; gefitinib; erlotinib; sunitinib; sorafenib; dasatinib; lapatinib; nilotinib; temsirolimus; everolimus; pazopanib; amsacrine; asparaginase; altretamine; hydroxycarbamide; lonidamine; pentostatin; miltefosine; masoprocold; estramustine; mitoguazone; topotecan; tiazofurine; irinotecan; mitotane; pegaspargase; bexarotene; bortezomib; celecoxib; anagrelide; oblimersen; vorinostat; romidepsin; fosfestrol; buserelin; leuprorelin; goserelin; triptorelin; tamoxifen; toremifene; fulvestrant; flutamide; nilutamide; bicalutamide; aminoglutethimide; formestane; anastrozole; letrozole; vorozole; exemestane; abarelix; degarelix; filgrastim; molgramostim; sargramostim; lenograstim; ancestim; pegfilgrastim; aldesleukin; oprelvekin; lentinan; roquinimex; pegademase; pidotimod; thymopentin; immunocyanin; tasonermin; mifamurtide; plerixafor; muromonab-CD3; sirolimus; leflunomide; alefacept; gusperimus; efalizumab; abetimus; natalizumab; abatacept; eculizumab; etanercept; infliximab; afelimomab; adalimumab; golimumab; daclizumab; basiliximab; anakinra; rilonacept; ustekinumab; mepolizumab; tocilizumab; canakinumab; ciclosporin; azathioprine; thalidomide; lenalidomide; phenylbutazone; mofebutazone; oxyphenbutazone; clofezone; kebuzone; sulindac; tolmetin; zomepirac; alclofenac; bumadizone; etodolac; lonazolac; fentiazac; acetaminophen; difenpiramide; oxametacin; proglumetacin; ketorolac; aceclofenac; bufexamac; piroxicam; tenoxicam; droxicam; lornoxicam; meloxicam; ketoprofen; fenoprofen; fenbufen; benoxaprofen; suprofen; piroprofen; flurbiprofen; indoprofen; oxaprozin; ibuprofen; dexibuprofen; alminoprofen;

dexketoprofen; rofecoxib; valdecoxib; parecoxib; etoricoxib;
lumiracoxib; nabumetone; azapropazone; glucosamine;
proquazone; orgotein; nimesulide; feprazone; diacerein;
morniflumate; tenidap; oxycinchophen; auranofin;
aurothioglucose; aurotioprol; penicillamine; bucillamine;
etofenamate; felbinac; bendazac; suxibuzone; nifenazone;
capsaicin; zucapsaicin; alcuronium; tubocurarine;
dimethyltubocurarine; suxamethonium; pancuronium; gallamine;
vecuronium; atracurium; hexafluronium; cisatracurium;
phenprobamate; carisoprodol; methocarbamol; styramate;
febarbamate; chlormezanone; chlorzoxazone; baclofen;
tizanidine; pridinol; tolperisone; thiocolchicoside;
mephenesin; tetrazepam; cyclobenzaprine; eperisone;
fenyramidol; dantrolene; allopurinol; tisopurine; febuxostat;
probenecid; sulfinpyrazone; benzbromarone; isobromindione;
colchicine; cinchophen; pegloticase; ipriflavone; denosumab;
hydroquinine; chymopapain; halothane; chloroform; enflurane;
trichloroethylene; isoflurane; desflurane; sevoflurane;
methohexital; hexobarbital; thiopental; narcobarbital;
fentanyl; alfentanil; sufentanil; phenoperidine; anileridine;
remifentanil; droperidol; ketamine; propanidid; alfaxalone;
etomidate; propofol; esketamine; xenon; metabutethamine;
chloroprocaine; bupivacaine; mepivacaine; prilocaine;
butanilicaine; etidocaine; articaine; ropivacaine;
levobupivacaine; cocaine; dyclonine; morphine; hydromorphone;
nicomorphine; oxycodone; dihydrocodeine; diamorphine;
papaveretum; ketobemidone; pethidine; dextromoramide;
piritramide; dextropropoxyphene; bezitramide; pentazocine;
phenazocine; buprenorphine; butorphanol; nalbuphine; tilidine;
tramadol; dezocine; meptazinol; tapentadol; salicylamide;
salsalate; ethenzamide; dipyrrocetyl; benorilate; diflunisal;
guacetisal; phenazone; aminophenazone; propyphenazone;
paracetamol; phenacetin; bucetin; propacetamol; rimazolium;

glafenine; floctafenine; viminol; nefopam; flupirtine;
ziconotide; methoxyflurane; nabiximols; dihydroergotamine;
ergotamine; methysergide; flumedroxone; sumatriptan;
naratriptan; zolmitriptan; rizatriptan; almotriptan;
eletriptan; frovatriptan; pizotifen; ipرازochrome;
dimetotiazine; oxetorone; methylphenobarbital; phenobarbital;
primidone; barbexaclone; metharbital; ethotoin; phenytoin;
mephenytoin; fosphenytoin; paramethadione; trimethadione;
ethadione; ethosuximide; phensuximide; mesuximide; clonazepam;
carbamazepine; oxcarbazepine; rufinamide; eslicarbazepine;
valpromide; vigabatrin; progabide; tiagabine; sultiame;
phenacemide; lamotrigine; felbamate; topiramate; gabapentin;
pheneturide; levetiracetam; zonisamide; pregabalin;
stiripentol; lacosamide; carisbamate; beclamide;
trihexyphenidyl; biperiden; metixene; procyclidine;
profenamine; dexetimide; phenglutarimide; mazaticol;
bornaprine; tropatepine; etanautine; benzatropine;
etybenzatropine; levodopa; melevodopa; amantadine; pergolide;
ropinirole; pramipexole; piribedil; rotigotine; selegiline;
rasagiline; tolcapone; entacapone; budipine; chlorpromazine;
levomepromazine; promazine; acepromazine; triflupromazine;
cyamemazine; chlorproethazine; dixyrazine; fluphenazine;
perphenazine; prochlorperazine; thiopropazate;
trifluoperazine; acetophenazine; thioproperazine;
butaperazine; perazine; periciazine; thioridazine;
mesoridazine; pipotiazine; haloperidol; trifluoperidol;
melperone; moperone; pipamperone; bromperidol; benperidol;
fluanisone; oxypertine; molindone; sertindole; ziprasidone;
flupentixol; clopenthixol; chlorprothixene; tiotixene;
zuclopenthixol; fluspirilene; pimozone; penfluridol; loxapine;
clozapine; olanzapine; quetiapine; asenapine; clotiapine;
#sulpiride; sultopride; tiapride; remoxipride; amisulpride;
veralipride; levosulpiride; lithium; prothipendyl;

risperidone; mosapramine; zotepine; aripiprazole;
paliperidone; diazepam; chlordiazepoxide; medazepam; oxazepam;
lorazepam; adinazolam; bromazepam; clobazam; ketazolam;
prazepam; alprazolam; halazepam; pinazepam; camazepam;
nordazepam; fludiazepam; etizolam; clotiazepam; cloxazolam;
tofisopam; hydroxyzine; captodiame; meprobamate; emylcamate;
mebutamate; benzoctamine; buspirone; mephenoxalone;
gedocarnil; etifoxine; pentobarbital; amobarbital;
butobarbital; barbital; aprobarbital; secobarbital; talbutal;
vinylbital; vinbarbital; cyclobarbital; heptabarbital;
reposal; etallobarbital; allobarbital; proxibarbal;
chloralodol; dichloralphenazone; paraldehyde; flurazepam;
nitrazepam; flunitrazepam; estazolam; triazolam; lormetazepam;
temazepam; midazolam; brotizolam; quazepam; loprazolam;
doxefazepam; cinolazepam; glutethimide; methyprylon;
pyrithyldione; zopiclone; zolpidem; zaleplon; eszopiclone;
melatonin; ramelteon; methaqualone; clomethiazole; bromisoval;
carbromal; propiomazine; triclofos; ethchlorvynol; valerian;
hexapropymate; bromides; apronal; valnoctamide;
methylpentynol; niaprazine; dexmedetomidine; desipramine;
imipramine; clomipramine; opipramol; trimipramine;
lofepramine; dibenzepin; amitriptyline; nortriptyline;
protriptyline; doxepin; iprindole; melitracen; butriptyline;
dosulepin; amoxapine; dimetacrine; amineptine; maprotiline;
quinupramine; zimeldine; fluoxetine; citalopram; paroxetine;
sertraline; alaproclate; fluvoxamine; etoperidone;
escitalopram; isocarboxazid; nialamide; phenelzine;
tranylcypromine; iproniazide; iproclozide; moclobemide;
toloxatone; oxitriptan; tryptophan; mianserin; nomifensine;
trazodone; nefazodone; minaprine; bifemelane; viloxazine;
oxaflozane; mirtazapine; bupropion; medifoxamine; tianeptine;
pivagabine; venlafaxine; milnacipran; reboxetine; gepirone;

duloxetine; agomelatine; desvenlafaxine; amfetamine; dexamfetamine; metamfetamine; methylphenidate; pemoline; fencamfamin; modafinil; fenozolone; atomoxetine; fenetylline; dexmethylphenidate; caffeine; propentofylline; meclofenoxate; pyritinol; piracetam; deanol; fipexide; citicoline; oxiracetam; pirisudanol; linopirdine; nifedipine; aniracetam; acetylcarnitine; idebenone; prolintane; pipradrol; pramiracetam; adrafinil; vinpocetine; tacrine; donepezil; rivastigmine; galantamine; memantine; neostigmine; pyridostigmine; distigmine; ambenonium; carbachol; bethanechol; pilocarpine; cevimeline; nicotine; varenicline; disulfiram; acamprosate; naltrexone; methadone; levacetylmethadol; lofexidine; betahistine; cinnarizine; flunarizine; acetylleucine; tirilazad; riluzole; xaliproden; amifampridine; tetrabenazine; tilbroquinol; nimorazole; secnidazole; diloxanide; clefamide; etofamide; teclozan; arsthinol; difetarstone; glycobarsol; chiniofon; emetine; phanquinone; mepacrine; atovaquone; trimetrexate; tenonitroazole; dihydroemetine; fumagillin; nitazoxanide; chloroquine; hydroxychloroquine; primaquine; amodiaquine; proguanil; quinine; mefloquine; pyrimethamine; artemisinin; artemether; artesunate; artemotil; artenimol; halofantrine; benznidazole; nifurtimox; melarsoprol; praziquantel; oxamniquine; metrifonate; niridazole; stibophen; triclabendazole; mebendazole; albendazole; cicloheximide; flubendazole; fenbendazole; piperazine; diethylcarbamazine; pyrantel; oxantel; levamisole; ivermectin; pyrvinium; buphenium; niclosamide; desaspidin; dichlorophen; dioxanthogen; thiram; clofenotane; lindane; pyrethrum; bioallethrin; phenothrin; permethrin; malathion; quassia; cyfluthrin; cypermethrin; decamethrin; tetramethrin; diethyltoluamide; dimethylphthalate; dibutylphthalate; dibutylsuccinate; dimethylcarbamate; etohexadiol; cyclopentamine; ephedrine;

oxymetazoline; tetryzoline; xylometazoline; naphazoline;
tramazoline; metizoline; tuaminoheptane; fenoxazoline;
tymazoline; levocabastine; azelastine; antazoline; nedocromil;
olopatadine; flunisolide; ritiometan; phenylpropanolamine;
pseudoephedrine; ambazone; benzethonium; myristyl-
benzalkonium; hexylresorcinol; fusafungine; gramicidin;
orciprenaline; salbutamol; terbutaline; rimiterol;
hexoprenaline; isoetarine; pirbuterol; tretoquinol;
carbuterol; tulobuterol; salmeterol; formoterol; clenbuterol;
reproterol; procaterol; bitolterol; indacaterol; ciclesonide;
fenspiride; methoxyphenamine; bambuterol; diprophylline;
proxiphylline; theophylline; aminophylline; etamiphylline;
bamifylline; bufylline; doxofylline; zafirlukast; pranlukast;
montelukast; ibudilast; eprozinol; omalizumab; seratrodast;
roflumilast; tyloxapol; guaifenesin; ipecacuanha; senega;
creosote; guaiacolsulfonate; levoverbenone; acetylcysteine;
bromhexine; carbocisteine; eprazinone; mesna; ambroxol;
sobrerol; domiodol; letosteine; stepronin; tiopronin;
neltexine; erdosteine; ethylmorphine; hydrocodone; codeine;
normethadone; noscapine; pholcodine; dextromethorphan;
thebacon; dimemorfan; acetyldihydrocodeine; benzonatate;
benproperine; clobutinol; isoaminile; pentoxyverine;
oxolamine; oxeladin; clofedanol; pipazetate; butamirate;
fedrilate; zipeprol; dibunate; droxypropine; prenoxdiazine;
dropropizine; cloperastine; meprotixol; piperidione;
tipepidine; morclofone; nepinalone; levodropropizine;
dimethoxanate; bromazine; diphenylpyraline; carbinoxamine;
doxylamine; brompheniramine; dexchlorpheniramine;
chlorphenamine; pheniramine; dexbrompheniramine; talastine;
histapyrrodine; methapyrilene; alimemazine; thiethylperazine;
methdilazine; hydroxyethylpromethazine; thiazinam;
mequitazine; oxomemazine; buclizine; cyclizine;
chlorcyclizine; meclozine; oxatomide; cetirizine;

levocetirizine; cyproheptadine; phenindamine; triprolidine; pyrrobutamine; azatadine; astemizole; terfenadine; loratadine; mebhydrolin; dehtropine; ketotifen; acrivastine; tritoqualine; ebastine; pimethixene; epinastine; mizolastine; fexofenadine; desloratadine; rupatadine; doxapram; nikethamide; pentetrazol; etamivan; bemegride; prethcamide; almitrine; dimeflin; mepixanox; dihydrostreptomycin; micronomicin; azidamfenicol; sulfadicramide; sulfacetamide; sulfafenazol; trifluridine; interferon; fomivirsen; bibrocathol; picloxydine; medrysone; formocortal; loteprednol; pranoprofen; nepafenac; bromfenac; dipivefrine; apraclonidine; brimonidine; ecothiopate; demecarium; physostigmine; fluostigmine; aceclidine; acetylcholine; paraoxon; acetazolamide; diclofenamide; dorzolamide; brinzolamide; methazolamide; levobunolol; metipranolol; befunolol; latanoprost; unoprostone; bimatoprost; travoprost; tafluprost; dapiprazole; cyclopentolate; homatropine; tropicamide; lodoxamide; emedastine; proxymetacaine; fluorescein; hypromellose; verteporfin; anecortave; pegaptanib; ranibizumab; guaiazulen; alum; iodoheparinate; nalorphine; edetates; pralidoxime; thiosulfate; dimercaprol; obidoxime; protamine; naloxone; flumazenil; methionine; cholinesterase; glutathione; fomepizole; sugammadex; deferoxamine; deferiprone; deferasirox; sevelamer; dexrazoxane; amifostine; rasburicase; palifermin; glucarpidase; oxygen; helium; nitrogen; nalfurafine; sincalide; ceruletide; metyrapone; corticorelin; somatostatin; galactose; sulfobromophthalein; tuberculin; betazole; pentagastrin; phenolsulfonphthalein; alsactide; protirelin; secretin; bentiromide; iodamide; methiodal; diiodone; metrizamide; iothexol; iopamidol; iopromide; iotrolan; ioversol; iopentol; iodixanol; iomeprol; iobitridol; ioxilan; adapiodone; iopydol; propyliodone; iofendylate; gadodiamide; gadoteridol; mangafodipir; gadoversetamide; gadobutrol;

gadofosveset; ferumoxsil; ferristene; perflubron;
perflenapent.

It is a further object of the present invention to provide a compound of formula (I), (Ia), (Ib) and/or (Ic) as defined herein or a pharmaceutical composition as defined herein for the preparation of a medicament for the treatment of one or more diseases mentioned herein.

Preferably the compounds of the present invention may be used for the treatment and/or prevention of the following conditions:

respiratory tract/obstructive airways diseases and disorders including:

rhinorrhea, tracheal constriction, airway contraction, acute-, allergic, atrophic rhinitis or chronic rhinitis (such as rhinitis caseosa, hypertrophic rhinitis, rhinitis purulenta, rhinitis sicca), rhinitis medicamentosa, membranous rhinitis (including croupous, fibrinous and pseudomembranous rhinitis), scrofulous rhinitis, perennial allergic rhinitis, seasonal rhinitis (including rhinitis nervosa (hay fever) and vasomotor rhinitis), pollinosis, asthma (such as bronchial, atopic, allergic, intrinsic, extrinsic, exercise-induced, cold air-induced, occupational, bacterial infection-induced, and dust asthma particularly chronic or inveterate asthma (e.g. late asthma and airways hyper-responsiveness)), bronchitis (including chronic, acute, arachidic, catarrhal, croupus, phthinoid and eosinophilic bronchitis), cardiobronchitis, pneumoconiosis, chronic inflammatory disease of the lung which result in interstitial fibrosis, such as interstitial lung disease (ILD) (e.g., idiopathic pulmonary fibrosis, or ILD associated with rheumatoid arthritis, or other autoimmune

conditions), acute lung injury (ALI), adult respiratory distress syndrome (ARDS), chronic obstructive pulmonary, airways or lung disease (CORD, COAD, COLD or COPD, such as irreversible COPD), chronic sinusitis, conjunctivitis (e.g. allergic conjunctivitis), cystic fibrosis, extrinsic allergic alveolitis (like farmer's lung and related diseases), fibroid lung, hypersensitivity lung diseases, hypersensitivity pneumonitis, idiopathic interstitial pneumonia, nasal congestion, nasal polyposis, otitis media, and cough (chronic cough associated with inflammation or iatrogenic induced), pleurisy, pulmonary congestion, emphysema, bronchiectasis, sarcoidosis, lung fibrosis, including cryptogenic fibrosing alveolitis, fibrosis complicating anti-neoplastic therapy and chronic infection, including tuberculosis and aspergillosis and other fungal infections, vasculitic and thrombotic disorders of the lung vasculature, and pulmonary hypertension, acute viral infection including the common cold, and infection due to respiratory syncytial virus, influenza, coronavirus (including SARS) and adenovirus, allergic bronchopulmonary mycosis, emphysema, diffuse panbronchiolitis, systemic anaphylaxis or hypersensitivity responses, drug allergies (e.g., to penicillin, cephalosporins), insect sting allergies, and food related allergies which may have effects remote from the gut (such as migraine, rhinitis and eczema), anaphylactic shock, vascular spasms;

bone and joint related diseases and disorders including:

osteoporosis, arthritis (including rheumatic, infectious, autoimmune, chronic, malignant), seronegative spondyloarthropathies (such as ankylosing spondylitis, rheumatoid spondylitis, psoriatic arthritis, enthesopathy, Bechet's disease, Marie-Strümpell arthritis, arthritis of inflammatory bowel disease, and Reiter's disease), systemic

sclerosis, osteoarthritis, osteoarthrosis, both primary and secondary to e.g. congenital hip dysplasia, cervical and lumbar spondylitis, and low back and neck pain, Still's disease, reactive arthritis and undifferentiated spondarthropathy, septic arthritis and other infection-related arthropathies and bone disorders such as tuberculosis, including Pott's disease and Poncet's syndrome, acute and chronic crystal-induced synovitis including urate gout, calcium pyrophosphate deposition disease, and calcium apatite related tendon, bursar and synovial inflammation, primary and secondary Sjogren's syndrome, systemic sclerosis and limited scleroderma, mixed connective tissue disease, and undifferentiated connective tissue disease, inflammatory myopathies including, polymyalgia rheumatica, juvenile arthritis including idiopathic inflammatory arthritides of whatever joint distribution and associated syndromes, other joint disease (such as intervertebral disc degeneration or temporomandibular joint degeneration), rheumatic fever and its systemic complications, vasculitides including giant cell arteritis, Takayasu's arteritis, polyarteritis nodosa, microscopic polyarteritis, and vasculitides to associated with viral infection, hypersensitivity reactions, cryoglobulins, paraproteins, low back pain, Familial Mediterranean fever, Muckle-Wells syndrome, and Familial Hibernian Fever, Kikuchi disease, drug-induced arthralgias, tendonitides, polychondritis, and myopathies, osteoporosis, osteomalacia like osteoporosis, osteopenia, osteogenesis imperfecta, osteopetrosis, osteofibrosis, osteonecrosis, Paget's disease of bone, hypophosphatemia, Felty's syndrome, Still's disease, slack of artificial joint implant, sprain or strain of muscle or joint, tendinitis, fasciitis, periarthritides humeroscapularis, cervico-omo-brachial syndrome, tenosynovitis;

Skin and eye related diseases and disorders including:
glaucoma, ocular hypertension, cataract, retinal detachment, psoriasis (including psoriasis vulgaris, pustular psoriasis, arthritic psoriasis, erythroderma psoriaticum), palmoplantar pustulosis, xeroderma, eczematous diseases (like atopic dermatitis, ultraviolet radiation dermatitis, contact dermatitis, and seborrheic dermatitis), phytodermatitis, photodermatitis, cutaneous eosinophilias, chronic skin ulcers, cutaneous lupus erythematosus, contact hypersensitivity/allergic contact dermatitis (including sensitivity to poison ivy, sumac, or oak), and eosinophilic folliculitis (Ofuji's disease), pruritus, drug eruptions, urticaria (acute or chronic, allergic or non-allergic), acne, erythema, dermatitis herpetiformis, scleroderma, vitiligo, lichen planus, lichen sclerosus et atrophica, pyodermis gangrenosum, skin sarcoid, pemphigus, ocular pemphigus, pemphigoid, epidermolysis bullosa, angioedema, vasculitides, toxic erythemas, cutaneous eosinophilias, alopecia areata, male-pattern baldness, Sweet's syndrome, Stevens-Johnson syndrome, Weber-Christian syndrome, erythema multiforme, cellulitis, both infective and non infective, panniculitis, cutaneous Lymphomas, non-melanoma skin cancer and other dysplastic lesions, blepharitis, iritis, anterior and posterior uveitis, choroiditis, autoimmune, degenerative or inflammatory disorders affecting the retina, ophthalmitis including sympathetic ophthalmitis, sarcoidosis, xerosis infections including viral, fungal, and bacterial, allergic conjunctivitis, increased fibrosis, keloids, keloplasty, post-surgical scars, epidermolysis bullosa, dry eye, ocular inflammation, allergic conjunctivitis, vernal conjunctivitis, vernal keratoconjunctivitis, and giant papillary conjunctivitis, ocular angiogenesis, cornea damage and scar,

all forms of macular degeneration, macular edema, macular dystrophy, abnormal wound healing, scleritis, episcleritis, pachydermia, peripheral ulcerative keratitis, fungal keratitis, herpetic keratitis, invasive aspergillosis; conical cornea, dystorpha epithelialis corneae, severe intraocular inflammation;

gastrointestinal tract and abdominal related diseases and disorders including:

celiac/coeliac disease (e.g. celiac sprue), cholecystitis, enteritis (including infectious, ischemic, radiation, drug-induced, and eosinophilic gastroenteritis), eosinophilic esophagitis, eosinophilic gastrointestinal inflammation, allergen induced diarrhea, enteropathy associated with seronegative arthropathies, gastritis, autoimmune atrophic gastritis, ischemic bowel disease, inflammatory bowel disease (Crohn's disease and ulcerative colitis), colitis, Mooren's ulcer, irritable bowel syndrome, necrotizing enterocolitis, gut ischemia, glossitis, gingivitis, periodontitis, oesophagitis, including reflux, proctitis, fibrosis and cirrhosis of the liver, pancreatitis, both acute and chronic, pancreatic fibrosis, pancreatic sclerosis, pancreatolithiasis, hepatic cirrhosis, hepatitis (congestive, autoimmune, acute, fulminant, chronic, drug-induced, alcoholic, lupoid, steatohepatitis and chronic viral), fatty liver, primary biliary cirrhosis, hepatic porphyria, and gastrointestinal related allergic disorders, spastic colon, diverticulitis, gastroenteric bleeding, Behcet's disease; partial liver resection, acute liver necrosis (e.g. necrosis caused by toxins, viral hepatitis, shock or anoxia), hemolytic uremic syndrome;

hematological disorders including:

anemias, coagulation, myeloproliferative disorders, hemorrhagic disorders, leukopenia, eosinophilic disorders,

leukemias (e.g. myelogenous, lymphomas, plasma cell dyscrasias, disorders of the spleen, Banti's disease, hemophilia, purpura (including idiopathic thrombocytopenic purpura), Wiskott-Aldrich syndrome;

metabolic disorders including:

obesity, amyloidosis, disturbances of the amino and acid metabolism like branched chain disease, hyperaminoacidemia, hyperaminoaciduria, disturbances of the metabolism of urea, hyperammonemia, mucopolysaccharidoses e.g. Maroteaux-Lamy syndrome, storage disease like glycogen storage diseases and lipid storage diseases, glycogenosis I diseases like Cori's disease, malabsorption diseases like intestinal carbohydrate malabsorption, oligosaccharidase deficiency like maltase-, lactase-, sucrase-insufficiency, disorders of the metabolism of fructose, disorders of the metabolism of galactose, galactosaemia, disturbances of carbohydrate utilization like diabetes, hypoglycemia, disturbances of pyruvate metabolism, hypolipidemia, hypolipoproteinemia, hyperlipidemia, hyperlipoproteinemia, carnitine or carnitine acyltransferase deficiency, disturbances of the porphyrin metabolism, porphyrins, disturbances of the purine metabolism, lysosomal diseases, metabolic diseases of nerves and nervous systems like gangliosidoses, sphingolipidoses, sulfatidoses, leucodystrophies, Lesch-Nyhan syndrome;

cerebellar dysfunction, disturbances of brain metabolism like: dementia, Alzheimer's disease, Huntington's chorea, Parkinson's disease, Pick's disease, toxic encephalopathy, demyelinating neuropathies like inflammatory neuropathy, Guillain-Barre syndrome; Meniere's disease and radiculopathy, primary and secondary metabolic disorders associated with hormonal defects like any disorder stemming from either an hyperfunction or hypofunction of some hormone-secreting

endocrine gland and any combination thereof. Sipple's syndrome, pituitary gland dysfunction and its effects on other endocrine glands, such as the thyroid, adrenals, ovaries, and testes, acromegaly, hyper- and hypothyroidism, euthyroid goiter, euthyroid sick syndrome, thyroiditis, and thyroid cancer, over or underproduction of the adrenal steroid hormones, adrenogenital syndrome, Cushing's syndrome, Addison's disease of the adrenal cortex, Addison's pernicious anemia, primary and secondary aldosteronism, diabetes insipidus, diabetes mellitus, carcinoid syndrome, disturbances caused by the dysfunction of the parathyroid glands, pancreatic islet cell dysfunction, diabetes, disturbances of the endocrine system of the female like estrogen deficiency, resistant ovary syndrome; muscle weakness, myotonia. Duchenne's and other muscular dystrophies, dystrophia myotonica of Steinert, mitochondrial myopathies like I disturbances of the catabolic metabolism in the muscle, carbohydrate and lipid storage myopathies, glycogenoses, myoglobinuria, malignant hyperthermia, polymyalgia rheumatica, dermatomyositis, multiple myositis, primary myocardial disease, cardiomyopathy; disorders of the ectoderm, neurofibromatosis, scleroderma and polyarthritis, Louis-Bar syndrome, von Hippel-Lindau disease, Sturge-Weber syndrome, tuberous sclerosis, amyloidosis, porphyria; sexual dysfunction of the male and female; confused states and seizures due to inappropriate secretion of antidiuretic hormone from the pituitary gland, Liddle's syndrome, Bartter's syndrome, Fanconi's I syndrome, and renal electrolyte wasting; transplant rejection related conditions including: acute and chronic allograft rejection following solid organ transplant, for example, transplantation of kidney, heart, liver, lung, and cornea, chronic graft versus host disease, skin graft rejection, and bone marrow transplant rejection,

immunosuppression;

genitourinary related conditions including:

nephritis (interstitial, acute interstitial (allergic), and glomerulonephritis), nephrotic syndrome, cystitis including acute and chronic (interstitial) cystitis and Hunner's ulcer, acute and chronic urethritis, prostatitis, epididymitis, oophoritis, salpingitis, vulvo vaginitis, vulvovaginal candidiasis, Peyronie's disease, and erectile dysfunction, renal disease, renal fibrosis, nephropylitis, secondary contracted kidney, steroid dependent and steroid-resistant nephrosis, Goodpasture's syndrome;

CNS related diseases and disorders including:

neurodegenerative diseases, Alzheimer's disease and other cementing disorders including CJD and nvCJD, amyloidosis, and other demyelinating syndromes, cerebral atherosclerosis and vasculitis, temporal arteritis, myasthenia gravis, acute and chronic so pain (acute, intermittent or persistent, whether of central or peripheral origin) including post-operative, visceral pain, headache, migraine, neuralgia (including trigeminal), atypical facial pain, joint and bone pain, pain arising from cancer and tumor invasion, neuropathic pain syndromes including diabetic, post-herpetic, and HIV-associated neuropathies, neurosarcoidosis, to brain injuries, cerebrovascular diseases and their consequences, Parkinson's disease, corticobasal degeneration, motor neuron disease, dementia, including ALS (Amyotrophic lateral sclerosis), multiple sclerosis, traumatic brain injury, stroke, post-stroke, post-traumatic brain injury, and small-vessel cerebrovascular disease, dementias, vascular dementia, dementia with Lewy bodies, frontotemporal dementia and Parkinsonism linked 1 to chromosome 17, frontotemporal dementias, including Pick's disease, progressive supranuclear palsy, corticobasal degeneration, Huntington's disease,

thalamic degeneration, HIV dementia, schizophrenia with dementia, and Korsakoff's psychosis, within the meaning of the definition are also considered to be CNS disorders central and peripheral nervous system complications of malignant, infectious or autoimmune processes, algesia, cerebral infarction, attack, cerebral ischemia, head injury, spinal cord injury, myelopathic muscular atrophy, Shy-Drager syndrome, Reye's syndrome, progressive multifocal leukoencephalopathy, normal pressure hydrocephalus, sclerosing panencephalitis, frontal lobe type dementia, acute anterior poliomyelitis (poliomyelitis), poliomyelitis neurosis, viral encephalitis, allergic encephalomyelitis, epileptic encephalopathies, Creutzfeldt-Jakob disease, Kuru disease, bovine spongiform encephalopathy (mad cow disease), scrapie, epilepsy, cerebral amyloid angiopathy, depression, mania, manic-depressive psychosis, hereditary cerebellar ataxia, peripheral neuropathy, Nasu-Hakola syndrome, Machado-Joseph disease;

inflammatory or immunological diseases or disorders including: general inflammation (of the ocular, nasal, pulmonary, and gastrointestinal passages), mastocytosis/mast cell disorders (cutaneous, systemic, mast cell activation syndrome, and pediatric mast cell diseases), mastitis (mammary gland), vaginitis, vasculitis (e.g., necrotizing, cutaneous, and hypersensitivity vasculitis), Wegener granulomatosis, myositis (including polymyositis, dermatomyositis), basophil related diseases including basophilic leukemia and basophilic leukocytosis, and eosinophil related diseases such as Churg-Strauss syndrome, eosinophilic granuloma, lupus erythematosus (such as, systemic lupus erythematosus, subacute cutaneous lupus erythematosus, and discoid lupus erythematosus), chronic thyroiditis, Hashimoto's thyroiditis, Grave's disease, type I diabetes, complications arising from diabetes mellitus, other

immune disorders, eosinophilia fasciitis, hyper IgE syndrome, Addison's disease, antiphospholipid syndrome, immunodeficiency disease, acquired immune deficiency syndrome (AIDS), leprosy, Sezary syndrome, paraneoplastic syndromes, and other autoimmune disorders, ferveescence, myositis, nervous diseases selected from multiple myositis, bursitis, Evans syndrome, leukotriene B4-mediated diseases, idiopathic hypoparathyroidism, nephrotic syndrome lupus, immunosuppression;

cardiovascular diseases and disorders including:

congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertension, cerebral trauma, occlusive vascular disease, stroke, cerebrovascular disorder, atherosclerosis, restenosis, affecting the coronary and peripheral is circulation, pericarditis, myocarditis, inflammatory and auto-immune cardiomyopathies including myocardial sarcoid, endocarditis, valvulitis, and aortitis including infective (e.g. syphilitic), hypertensive vascular diseases, peripheral vascular diseases, and atherosclerosis, vasculitides, disorders of the proximal and peripheral veins including phlebitis and thrombosis, including deep vein thrombosis and complications of varicose veins, aortic aneurism, periarteritis nodosa, cardiac fibrosis, post-myocardial infarction, idiopathic cardiomyopathy; angioplasty;

oncological diseases and disorders including:

common cancers (prostrate, breast, lung, ovarian, pancreatic, bowel and colon, abdomen, stomach (and any other digestive system cancers), liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head, neck, nervous system (central and peripheral), lymphatic system, blood, pelvic, skin, bone, soft

tissue, spleen, thoracic, urogenital, and brain tumors), breast cancer, genitourinary cancer, lung cancer, gastrointestinal cancer, epidermoid cancer, melanoma, ovarian cancer, pancreas cancer, neuroblastoma, malignancies affecting the bone marrow (including the leukaemias) and lymphoproliferative systems, such as Hodgkin's and non-Hodgkin's lymphoma, B-cell lymphoma, follicular lymphoma, metastatic disease and tumour recurrences, and paraneoplastic syndromes, as well as hypergammaglobulinemia, lymphoproliferative diseases, disorders, and/or conditions, paraproteinemias, purpura (including idiopathic thrombocytopenic purpura), Waldenström's Macroglobulinemia, Gaucher's Disease, histiocytosis, retinoblastoma and any other hyperproliferative disease, sarcomata, cachexia, tumor growth, tumor invasion, metastasis, AIDS-related lymphomas, malignant immunoproliferative diseases, multiple myeloma and malignant plasma cell neoplasms, lymphoid leukemia, acute or chronic myeloid leukemia, acute or chronic lymphocytic leukemia, monocytic leukemia, other leukemias of specified cell type, leukemia of unspecified cell type, other and unspecified malignant neoplasms of lymphoid, haematopoietic and related tissues, for example diffuse large cell lymphoma, T-cell lymphoma or cutaneous T-cell lymphoma). Myeloid cancer includes e.g. acute or chronic myeloid leukaemia, keratoleukoma and

other diseases and disorders including:

pain, migraine, sleep disorders, fever, sepsis, idiopathic thrombocytopenia pupura, post-operative adhesions, flushing, ischemic/reperfusion injury in the heart, brain, peripheral limbs, bacterial infection, viral infection, fungal infection, thrombosis, endotoxin shock, septic shock, thermal regulation including fever, Raynaud's disease, gangrene, diseases requiring anti-coagulation therapy, congestive heart failure,

mucus secretion disorders, pulmonary hypotension, prostanoid-induced smooth muscle contract associated with dysmenorrhea and premature labor, premature delivery, reperfusion injury, burn, thermal injury, hemorrhage or traumatic shock, menstrual pain, menstrual cramp, dysmenorrhea, periodontosis, rickettsial infectious disease, protozoal disease, reproduction disease, toothache, pain after tooth extraction, Herpes zoster, Herpes simplex, retroperitoneal fibrosis, various radiation injuries and the like.

A therapeutically effective amount of a compound in accordance with this invention means an amount of compound that is effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated. Determination of a therapeutically effective amount is within the skill in the art.

The therapeutically effective amount or dosage of a compound according to this invention can vary within wide limits and may be determined in a manner known in the art. Such dosage may be adjusted to the individual requirements in each particular case including the specific compound being administered, the route of administration, the condition being treated, as well as the patient being treated.

Examples of pharmacologically acceptable salts of sufficiently basic compounds of formula (I), (Ia), (Ib) and (Ic) are salts of physiologically acceptable mineral acids like hydrochloric, hydrobromic, sulfuric and phosphoric acid; or salts of organic acids like methanesulfonic, p-toluenesulfonic, lactic, acetic, trifluoroacetic, citric, succinic, fumaric, maleic and salicylic acid. Further, a sufficiently acidic compound of formula (I), (Ia), (Ib) and (Ic) may form alkali or earth

alkali metal salts, for example sodium, potassium, lithium, calcium or magnesium salts; ammonium salts; or organic base salts, for example methylamine, dimethylamine, trimethylamine, triethylamine, ethylenediamine, ethanolamine, choline hydroxide, meglumin, piperidine, morpholine, tris-(2-hydroxyethyl)amine, lysine or arginine salts; all of which are also further examples of salts of formula (I), (Ia), (Ib) and (Ic). Compounds of formula (I), (Ia), (Ib) and (Ic) may be solvated, especially hydrated. The hydratization/hydration may occur during the process of production or as a consequence of the hygroscopic nature of the initially water free compounds of formula (I), (Ia), (Ib) and (Ic). The solvates and/or hydrates may e.g. be present in solid or liquid form.

It should be appreciated that certain compounds of formula (I), (Ia), (Ib) and (Ic) may have tautomeric forms from which only one might be specifically mentioned or depicted in the following description, different geometrical isomers (which are usually denoted as cis/trans isomers or more generally as (E) and (Z) isomers) or different optical isomers as a result of one or more chiral carbon atoms (which are usually nomenclatured under the Cahn-Ingold-Prelog or R/S system). All these tautomeric forms, geometrical or optical isomers (as well as racemates and diastereomers) and polymorphous forms are included in the invention. Since the compounds of formula (I), (Ia), (Ib) and (Ic) may contain asymmetric C-atoms, they may be present either as achiral compounds, mixtures of diastereomers, mixtures of enantiomers or as optically pure compounds. The present invention comprises both all pure enantiomers and all pure diastereomers, and also the mixtures thereof in any mixing ratio.

According to a further embodiment of the present invention, one or more hydrogen atoms of the compounds of the present invention may be replaced by deuterium. Deuterium modification improves the metabolic properties of a drug with little or no change in its intrinsic pharmacology. Deuterium substitution at specific molecular positions improves metabolic stability, reduces formation of toxic metabolites and/or increases the formation of desired active metabolites. Accordingly, the present invention also encompasses the partially and fully deuterated compounds of formula (I), (Ia), (Ib) and (Ic). The term hydrogen also encompasses deuterium.

The therapeutic use of compounds according to formula (I), (Ia), (Ib) and (Ic), their pharmacologically acceptable salts, solvates and hydrates, respectively, as well as formulations and pharmaceutical compositions also lie within the scope of the present invention.

The pharmaceutical compositions according to the present invention comprise at least one compound of formula (I), (Ia), (Ib) and (Ic) as an active ingredient and, optionally, carrier substances and/or adjuvants.

The present invention also relates to pro-drugs which are composed of a compound of formula (I), (Ia), (Ib) and/or (Ic) and at least one pharmacologically acceptable protective group which will be cleaved off under physiological conditions, such as an alkoxy-, arylalkyloxy-, acyl-, acyloxymethyl group (e.g. pivaloyloxymethyl), an 2-alkyl-, 2-aryl- or 2-arylalkyloxycarbonyl-2-alkylidene ethyl group or an acyloxy group as defined herein, e.g. ethoxy, benzyloxy, acetyl or acetyloxy or, especially for a compound of formula (I), (Ia), (Ib) and/or (Ic), carrying a hydroxy group (-OH): a sulfate, a

phosphate ($-\text{OPO}_3$ or $-\text{OCH}_2\text{OPO}_3$) or an ester of an amino acid. Especially preferred are pro-drugs of the hydroxy group of a compound of formula (I), (Ia), (Ib) and/or (Ic).

Preferably, the present invention also relates to a prodrug, a biohydrolyzable ester, a biohydrolyzable amide, a polymorph, tautomer, stereoisomer, metabolite, N-oxide, biohydrolyzable carbamate, biohydrolyzable ether, physiologically functional derivative, atropisomer, or in vivo-hydrolysable precursor, diastereomer or mixture of diastereomers, chemically protected form, affinity reagent, complex, chelate and a stereoisomer of the compounds of formula (I), (Ia), (Ib) and/or (Ic).

As used herein, the term pharmaceutically acceptable ester especially refers to esters which hydrolyze in vivo and include those that break down readily in the human body to leave the parent compound or a salt thereof. Suitable ester groups include, for example, those derived from pharmaceutically acceptable aliphatic carboxylic acids, particularly alkanoic, alkenoic, cycloalkanoic and alkanedioic acids, in which each alkyl or alkenyl moiety advantageously has not more than 6 carbon atoms. Examples of particular esters include, but are not limited to, formates, acetates, propionates, butyrates, acrylates and ethylsuccinates.


As mentioned above, therapeutically useful agents that contain compounds of formula (I), (Ia), (Ib) and/or (Ic), their solvates, salts or formulations are also comprised in the scope of the present invention. In general, compounds of formula (I), (Ia), (Ib) and/or (Ic) will be administered by using the known and acceptable modes known in the art, either alone or in combination with any other therapeutic agent.

For oral administration such therapeutically useful agents can be administered by one of the following routes: oral, e.g. as tablets, dragees, coated tablets, pills, semisolids, soft or hard capsules, for example soft and hard gelatine capsules, aqueous or oily solutions, emulsions, suspensions or syrups, parenteral including intravenous, intramuscular and subcutaneous injection, e.g. as an injectable solution or suspension, rectal as suppositories, by inhalation or insufflation, e.g. as a powder formulation, as microcrystals or as a spray (e.g. liquid aerosol), transdermal, for example via an transdermal delivery system (TDS) such as a plaster containing the active ingredient or intranasal. For the production of such tablets, pills, semisolids, coated tablets, dragees and hard, e.g. gelatine, capsules the therapeutically useful product may be mixed with pharmaceutically inert, inorganic or organic excipients as are e.g. lactose, sucrose, glucose, gelatine, malt, silica gel, starch or derivatives thereof, talc, stearinic acid or their salts, dried skim milk, and the like. For the production of soft capsules one may use excipients as are e.g. vegetable, petroleum, animal or synthetic oils, wax, fat, polyols. For the production of liquid solutions, emulsions or suspensions or syrups one may use as excipients e.g. water, alcohols, aqueous saline, aqueous dextrose, polyols, glycerin, lipids, phospholipids, cyclodextrins, vegetable, petroleum, animal or synthetic oils. Especially preferred are lipids and more preferred are phospholipids (preferred of natural origin; especially preferred with a particle size between 300 to 350 nm) preferred in phosphate buffered saline (pH = 7 to 8, preferred 7.4). For suppositories one may use excipients as are e.g. vegetable, petroleum, animal or synthetic oils, wax, fat and polyols. For aerosol formulations one may use compressed gases suitable for this purpose, as are e.g. oxygen, nitrogen and


carbon dioxide. The pharmaceutically useful agents may also contain additives for conservation, stabilization, e.g. UV stabilizers, emulsifiers, sweetener, aromatizers, salts to change the osmotic pressure, buffers, coating additives and antioxidants.

In general, in the case of oral or parenteral administration to adult humans weighing approximately 80 kg, a daily dosage of about 10 mg to about 10,000 mg, preferably from about 20 mg to about 1,000 mg, should be appropriate, although the upper limit may be exceeded when indicated. The daily dosage can be administered as a single dose or in divided doses, or for parenteral administration, it may be given as continuous infusion or subcutaneous injection.


The present invention refers furthermore to compounds of formulas (II), (IIIa) and/or (IIIb) wherein R^2 and R^3 are defined as above and PG is a protecting group.



(II)



(IIIa)



(IIIb)

Protecting groups are known to a person skilled in the art and e.g. described in P. J. Kocienski, *Protecting Groups*, Georg Thieme Verlag, Stuttgart, 1994 and in T. W. Greene, P. G. M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, New York, 1999.

Preferably, PG is a 4-methoxy benzyl group or a carboxybenzyl (Cbz or Z) group; especially preferably, PG is a 4-methoxy benzyl group.

Preferred compounds of formula (II) are:

7-chloro-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine;
7-chloro-5-(3,4-dimethoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine;
7-chloro-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-

d]pyrimidine; 7-chloro-5-(1H-pyrazol-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole; 7-chloro-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol; 4-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)morpholine; 7-chloro-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,5-dimethoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; N-(2-aminoethyl)-3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzamide; 7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(1H-benzo[d]imidazol-2-yl)-7-chloro-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-5-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2H-pyrazolo[4,3-d]pyrimidine; (7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; 3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; 2-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)acetic acid; 5-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiophene-3-carboxylic acid; 2-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)acetic acid; 6-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)picolinic acid; 2-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; N-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzyl)-1-(p-tolyl)methanamine; 3-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)-N,N-dimethylpropan-1-amine; 3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; N-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; 4-(7-

chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; 4-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; 2-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; methyl 3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-(trifluoromethoxy)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-(trifluoromethyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-chlorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2-fluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-fluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 4-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; 7-chloro-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-indazol-6-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(4-fluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-difluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,5-difluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,6-difluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,4-difluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 6-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one.

Preferred compounds of formula (IIIa) and (IIIb) are:

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-

d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrazol-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole; 7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol; 4-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)morpholine; 7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,5-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; N-(2-aminoethyl)-3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzamide; 7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(1H-benzo[d]imidazol-2-yl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-5-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; (7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; 2-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)acetic acid; 5-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiophene-3-carboxylic acid; 2-(3-(7-chloro-2-(4-methoxybenzyl)-2H-


pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)acetic acid; 6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)picolinic acid; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzyl)-1-(p-tolyl)methanamine; 3-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)-N,N-dimethylpropan-1-amine; 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethoxy)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-chlorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; 7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-indazol-6-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-

pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(4-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,5-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,6-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one.

GENERAL SYNTHESIS


The procedures shown in the following synthesis schemes may be used to provide chloride intermediates. The chloride intermediates may then be used in an amination reaction with various amines and a subsequent deprotection to provide the final products.

Step 1:




4-Amino-2H-pyrazole-3-carboxylic acid amide is reacted with an aldehyde in the presence of an oxidant esp. 2,3-Dichloro-5,6-dicyano-1,4-benzoquinone to result in the formation of 5-substituted 1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one.

Step 2:




The pyrazole is protected e.g. with a para-methoxy-benzyl protection group. This step results in the formation of two isomers - 5-substituted 1-(4-Methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one and 5-substituted 2-(4-Methoxy-benzyl)-2,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one. Both isomers can be used in the following steps.

Step 3:




5-substituted 1-(4-Methoxy-benzyl)-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one and 5-substituted 2-(4-Methoxy-benzyl)-2,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one are chlorinated e.g. with POCl₃, resulting in the formation of 5-substituted 7-Chloro-1-(4-methoxy-benzyl)-1H-pyrazolo[4,3-d]pyrimidine and 5-substituted 7-Chloro-2-(4-methoxy-benzyl)-2H-pyrazolo[4,3-d]pyrimidine.

Step 4:



5-substituted 7-Chloro-1-(4-methoxy-benzyl)-1H-pyrazolo[4,3-d]pyrimidine and 5-substituted 7-Chloro-2-(4-methoxy-benzyl)-2H-pyrazolo[4,3-d]pyrimidine are reacted e.g. with the corresponding amine to result in the formation of the protected final products.

Step 5:



In a final step the para-methoxy-benzyl protection group is removed e.g. with trifluoroacetic acid to provide the final product for biological characterisation.

EXAMPLES

Materials and Methods

BIOLOGICAL ASSAYS - Protein Kinase Assays

SYK ASSAY

In the assay OMNIA® KINASE ASSAY by Invitrogen Corporation (Carlsbad) the effect of invention of a compound on the phosphorylation is determined by measurement of fluorescence intensity of a chelation-enhanced fluorophore called SOX. Upon phosphorylation of the peptide by the kinase of interest, Mg²⁺ is chelated to form a bridge between the SOX moiety and the phosphate group that is transferred to the specific tyrosine on the peptide. The fluorescence intensity is directly proportional to the amount of peptide phosphorylation.

To the wells of an 384 well small volume plate (Greiner, Frickenhausen) are added (i) the compound under test in 5 % DMSO/distilled water (2 µl), (ii) 16 µl of the master mix containing ATP, DTT, Kinase Reaction Buffer, Omina Peptide Substrate Tyr 7 resulting in a final concentration of 1 mM ATP, 0.2 mM DTT and 10 µM Peptide Substrate.

The Master Mix and the assay plate was incubated to reaction temperature before the measurement (30° C). The reaction was started with addition of (iii) 2 µl 4µg/ml SYK kinase (Invitrogen, Carlsbad). During measurement fluorescence intensity readings were collected using a TECAN M1000 at a wavelength of λ_{ex} 360/ λ_{em} 485 nm every 30 s for 30 minutes. The reaction velocity was plotted versus the inhibitor concentration to determine the IC₅₀ using XLFit 5.0 (IDBS, Guildford) to fit to a sigmoidal dose response curve, and the apparent K_i values were calculated from the IC₅₀ using the Cheng-Prusoff equation (Cheng, Y.; Prusoff, W. H. Biochem. Pharmacol. 22, 3099-3108, 1973).

The ATP dependency was determined according to Lai C-J-, Wu JC A Simple Kinetic Method for Rapid Mechanistic Analysis of Reversible Enzyme Inhibitors. Assays and Drug Dev. Technologies. 2003;1(4):527-535. To demonstrate a competition effect of the test compounds towards ATP the corresponding test compound was used at the 50% inhibitory concentration. Assay conditions as described previously were maintained. ATP concentrations used was 1000, 333, 100, 33.3, 10, 3.3, 1 μ M.

LRRK2 - LRRK2 G2019S - ASSAY

In the assay LanthaScreen™ Eu Kinase Binding Assay by Invitrogen Corporation (Carlsbad) the effect of invention of a compound on the phosphorylation is determined by measurement of fluorescence intensity emission ratio based on the binding and displacement of a proprietary, Alexa Fluor® 647- labeled, ATP-competitive kinase inhibitor scaffold (kinase tracer) to the kinase of interest. Binding of the tracer to the kinase is detected using a europium-labeled anti-tag antibody, which binds to the kinase of interest. Simultaneous binding of both the tracer and antibody to the kinase results in a high degree of FRET (fluorescence resonance energy transfer) from the europium (Eu) donor fluorophore to the Alexa Fluor® 647 acceptor fluorophore on the kinase tracer. Binding of an inhibitor to the kinase competes for binding with the tracer, resulting in a loss of FRET. The fluorescence intensity ratio is directly proportional to the amount of peptide phosphorylation.

To the wells of an 384 well small volume plate (Greiner, Frickenhausen) are added (i) the compound under test in 5 % DMSO/distilled water (5 μ l), (ii) 5 μ l of the kinase antibody mixture resulting in a final concentration 5 nM LRRK2 or their mutants, 2 nM EU-Anti-GST antibody in 1x kinase buffer A. The reaction was started with addition of (iii) 5 μ l resulting in

a final concentration of 10 nM tracer 236. The assay plate was incubated at RT for 1 h and fluorescence intensity readings were collected using a TECAN M1000 at two wavelengths of λ_{ex} 340/ λ_{em} 615 nm and λ_{ex} 340/ λ_{em} 665 nm with a delay time of 100 μ s and an integration time of 200 μ s after 60 minutes. The emission ratio was calculated by division of the acceptor/tracer emission (665 nm) by the antibody/donor emission (615 nm). The inhibitor concentration was plotted versus the emission ratio to determine the IC₅₀ using XLFit 5.0 (IDBS, Guildford) to fit to a sigmoidal dose response curve with a variable slope.

MYLK ASSAY

In the assay LanthaScreen™ Eu Kinase Binding Assay by Invitrogen Corporation (Carlsbad) the effect of invention of a compound on the phosphorylation is determined by measurement of fluorescence intensity emission ratio based on the binding and displacement of a proprietary, Alexa Fluor® 647- labeled, ATP-competitive kinase inhibitor scaffold (kinase tracer) to the kinase of interest as described above.

To the wells of an 384 well small volume plate (Greiner, Frickenhausen) are added (i) the compound under test in 5 % DMSO/distilled water (5 μ l), (ii) 5 μ l of the kinase antibody mixture resulting in a final concentration 5 nM MYLK, 2 nM EU-Anti-GST antibody in 1x kinase buffer A. The reaction was started with addition of (iii) 5 μ l resulting in a final concentration of 30 nM tracer 236. The assay plate was incubated at RT for 1 h and fluorescence intensity readings were collected using a TECAN M1000 at two wavelengths of λ_{ex} 340/ λ_{em} 615 nm and λ_{ex} 340/ λ_{em} 665 nm with a delay time of 100 μ s and an integration time of 200 μ s after 60 minutes. The emission ratio was calculated by division of the

acceptor/tracer emission (665 nM) by the antibody/donor emission (615 nM). The inhibitor concentration was plotted versus the emission ratio to determine the IC50 using XLFit 5.0 (IDBS, Guildford) to fit to a sigmoidal dose response curve with a variable slope.

BIOLOGICAL ASSAYS - Cellular Assay

LAD2 Assay

The inhibition of Syk may also be determined by examining IgE mediated release of histamine in vitro using human conjunctival tissue mast cells or in LAD2 mast cells (Leuk Res. 2003 Aug. 27(8):677-82). Human conjunctival tissue mast cells (HCTMCs) are obtained using the methodology outlined in U.S. Patent No. 5,360,720, for example. Briefly, HCTMCs are enzymatically released from human conjunctival tissues and then partially enriched by density centrifugation over a PERCOLL(R) cushion. A monodispersed cell suspension is obtained from the resulting pellet, and these cells are used for a histamine release assay. Cells are exposed to drug or control prior to stimulation with anti-human IgE, which triggers mast cell degranulation and histamine release to the supernatant. Histamine is then measured in the supernatant by EIA (Beckman Coulter), RIA (Beckman Coulter) or other method known to one of skill in the art. A decrease in histamine release drug treated cells indicates that the compound has potential for further investigation.

The LAD2 mast cell line is used in much the same way with the exception that the cells are passively sensitized with human IgE myeloma prior to stimulation with anti-human IgE to cross-link receptor bound IgE and trigger degranulation.

GENERAL PROCEDURES FOR SYNTHESIS OF COMPOUNDS
CHROMATOGRAPHY

The compound verification via analytical HPLC-MS was done after purification using the following instrumentation, column and method:

Analytical method for compound purity

Instrumentation:

Agilent MSD 1100

Analytical Methods:

Solvents:

A: acetonitrile

B: H₂O

C: 2% HCOOH in acetonitrile

D: 0.1% NEt₃ in acetonitrile

The following analytical methods were used:

Method A

Column SunFire C18 from Waters 2.1x50mm 2.5µm particle size, thermostated @ 40°

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	0.6
2.5	10	5	0	0.6
4	10	5	0	0.6
4.5	90	5	0	0.6
6	90	5	0	0.6

Stop time @ 7min

MS: ESI positive, Mass scan from 100 to 800
gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method B

Column ODS-AQ from YMC 4.0x50mm 2.5µm particle size,
thermostated @ RT

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	1.3
2.5	10	5	0	1.3
4	10	5	0	1.3
4.5	90	5	0	1.3
6	90	5	0	1.3

Stop time @ 7min

MS: ESI positive, Mass scan from 100 to 800
gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method C

Column ODS-AQ from YMC 2.1x50mm 3µm particle size,
thermostated @ 40°C

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	0.6
2.5	10	5	0	0.6
4	10	5	0	0.6
4.5	90	5	0	0.6
6	90	5	0	0.6

Stop time @ 7min

MS: ESI positive, Mass scan from 100 to 800

gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method D

Column ODS-AQ from YMC 2.1x50mm 3µm particle size, incl.
GuardCol 2.1x10mm, 3µm particle size
thermostated @ 40°C

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	0.6
10.0	10	5	0	0.6
13.0	10	5	0	0.6
14.0	90	5	0	0.6

Stop time @ 15min

MS: ESI positive, Mass scan from 100 to 800

gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method E

Column ODS-AQ from YMC 2.1x50mm 3µm particle size,
thermostated @ 40°C

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	0.6
5.0	10	5	0	0.6
7.0	10	5	0	0.6
8.0	90	5	0	0.6

Stop time @ 10min

MS: ESI positive, Mass scan from 100 to 800

gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method F

Column SunFire C18 from Waters 2.1x50mm 2.5µm particle size,
thermostated @ 40°

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	1.2
7.0	10	5	0	1.2
8.5	10	5	0	1.2
9.0	90	5	0	1.2
11.0	90	5	0	1.2

Stop time @ 12min

MS: ESI positive, Mass scan from 100 to 800

gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method G

Column YMC C8 OS 4.0x50mm 4µm particle size,

thermostated @ 40°

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	1.5
2.5	10	5	0	1.5
4.0	10	5	0	1.5
4.5	90	5	0	1.5
6.0	90	5	0	1.5

Stop time @ 6min

MS: ESI positive, Mass scan from 100 to 800

gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method H

Column Waters XBridge C18 2.1x50mm 2.5µm particle size,

thermostated @ 40°

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]

0	90	5	0	0.55
2.5	10	5	0	0.55
4.0	10	5	0	0.55
4.5	90	5	0	0.55
6.0	90	5	0	0.55

Stop time @ 6 min

MS: ESI positive, Mass scan from 100 to 800

gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method I

Column Waters XBridge C18 2.1x50mm 2.5µm particle size,
thermostated @ 40°

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	0	5	0.55
2.5	10	0	5	0.55
4.0	10	0	5	0.55
4.5	90	0	5	0.55
6.0	90	0	5	0.55

Stop time @ 6 min

MS: ESI positive, Mass scan from 100 to 800

gradient fragmentation: 50 to 125V

UV: detection @ 220 and 254nm

Method J

Column Waters SunFire C18 2.1x50mm, 2.5µm particle size
thermostated@40°C

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	0.6
10	10	5	0	0.6
13	10	5	0	0.6
14	90	5	0	0.6

Stoptime@15min

MS:ESIpositive,Massscanfrom100to800

gradientfragmentation:50to125V

UV: detection @ 220 and 254nm

Method K

Column YMC ODS-AQ 2.1x50mm, 3µm particle size
incl. GuardCol 2.1x10mm, 3µm particle size
thermostated@40°C

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	99	5	0	0.6
2.5	10	5	0	0.6
4	10	5	0	0.6
4.5	99	5	0	0.6
6	99	5	0	0.6

Stoptime@7min

MS:ESIpositive,Massscanfrom100to800

gradientfragmentation:50to125V

UV: detection @ 220 and 254nm

Method L

Column YMC TriArt C18 2.0x50mm, 1.9µm, #TA12SP90502WT
thermostated@40°C

Gradient:

Time [min]	%B	%C	%D	Flow [mL/min]
0	90	5	0	0.45
0.5	90	5	0	0.45
4.5	10	5	0	0.45
5.5	10	5	0	0.45
5.6	90	5	0	0.45

Stoptime@10min

MS:ESIpositive,Massscanfrom100to800

gradientfragmentation:50to125V

UV: detection @ 220 and 254nm

Purification and characterisation:

The resulting crude reaction products were purified in an automatic process using a semi-preparative HPLC-MS with mass-triggered sampling of the desired peak:

Purification via semi-preparative HPLC-MS

Instrumentation:

2x Varian PrepStar SD-1
1x Dionex P580 Pump 1 Channel (MakeUP I)
1x Dionex AXP-MS (MakeUP II)
1x Dionex MSQ
1x Dionex UVD 340V - Prep Flow Cell
Gilson 215 Liquid Handler

Column:

SunFire Prep C18 OBD 5 μ m 19x50mm

Method:

Column Flow: 30 ml/min
Solvent A: methanol, 0,3% acetic acid
Solvent B: water, 0,3 % acetic acid

Time table for gradient:

Time (min)	Solv. A	Solv. B
0.0	30.00	70.00
10.0	100.00	0.00
14.0	100.00	0.00
14.4	30.00	70.00
16.4	30.00	70.00

Detection:

UV 254nm, Mass Spectrometer Detector (API-ES, positive)

COMPOUND PREPARATION

Where the preparation of starting materials is not described, these are commercially available, known in the literature, or readily obtainable by those skilled in the art using standard procedures. Where it is stated that compounds were prepared analogously to earlier examples or intermediates, it will be appreciated by the skilled person that the reaction time, number of equivalents of reagents and temperature can be modified for each specific reaction and that it may be necessary or desirable to employ different workup or purification techniques. Where reactions are carried out using microwave irradiation, the microwave used is an Initiator 60 supplied by Biotage. The actual power supplied varies during the course of the reaction in order to maintain a constant temperature.

Abbreviations

DCM = Dichloromethane

DMF = N,N-Dimethylformamide

DDQ = 2,3-Dichloro-5,6-dicyano-1,4-benzoquinone

THF = Tetrahydrofuran

MeOH = Methanol

TFA = Trifluoroacetic acid

TEA = Triethylamine

rm = Reaction mixture

rt = Room temperature

AcOH = Acetic acid

MeCN = Acetonitrile

EtOH = Ethanol

EtOAc = Ethyl Acetate

LCMS = Mass spectrometry directed high pressure liquid chromatography

UV = Ultraviolet

DMSO = Dimethylsulphoxide

INTERMEDIATES

Preparation of

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine

Step1: 5-Phenyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one

4-Amino-2H-pyrazole-3-carboxylic acid amide (5mmol) and benzaldehyde (5mmol) were dissolved in acetic acid (14ml). Subsequently DDQ (3.75mmol) were added and the reaction mixture is heated in the microwave to 180°C for 30 minutes. The precipitate was filtered and washed with Et2O.

Mexact = 212,079

Yield 80%

HPLC-MS method: A

Retention Time: 2.1 min

mass found (m+H): 213.1

Step2: Protection of Pyrazole with 1-Chloromethyl-4-methoxybenzene

5-Phenyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one (3.9 mmol) was dissolved in 14ml acetonitrile. 1-Chloromethyl-4-methoxybenzene (4.7 mmol), 1.2 eq triethylamine (4.7 mmol) and sodiumiodide (0.39mmol) was added subsequently. The reaction mixture was heated in the microwave for 30min at 160°C. After cooling down the reaction the resulting precipitate was filtered and washed with Et2O.

The reaction is leading to two isomers (1-(4-Methoxy-benzyl)-5-phenyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one and 2-(4-Methoxy-benzyl)-5-phenyl-2,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one).

Both isomers can be used for the next steps.

Yield: 47%

Mexact = 332,149

HPLC-MS method: A

Retention time: 2.96 min / 3.2 min (Isomers)

Mass found (m+H): 333.1

Step3: Chlorination

The mixture of 1-(4-Methoxy-benzyl)-5-phenyl-1,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one and 2-(4-Methoxy-benzyl)-5-phenyl-2,6-dihydro-pyrazolo[4,3-d]pyrimidin-7-one (0.18 mmol) was suspended in POCl₃. The mixture was heated to 100°C in a reaction vessel. After 0,5h - 24h LC-MS showed reaction completion. The mixture was cooled to rt. The mixture was concentrated to dryness, cooled to 0°C, and quenched with ice/water. The mixture was extracted with DCM. The organic layer was washed with NaHCO₃ sat. aq. and water, dried (Na₂SO₄) filtered and concentrated.

The product was used without further purification.

The product is an isomeric mixture of 7-Chloro-1-(4-methoxy-benzyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidine and 7-Chloro-2-(4-methoxy-benzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine

Both isomers can be used for the next steps.

Mexact = 350,113

HPLC-MS method: A

Retention time: 3.98 min / 4.3 min

Mass found (m+H): 351.1

The following Chlorides and Isomers were synthesized according to the above mentioned procedure:

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrazol-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole; 7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol; 4-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)morpholine; 7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,5-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; N-(2-aminoethyl)-3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzamide; 7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(1H-benzo[d]imidazol-2-yl)-7-chloro-2-(4-methoxybenzyl)-2H-

pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-5-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; (7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester; 2-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)acetic acid methyl ester; 5-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiophene-3-carboxylic acid methyl ester; 2-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)acetic acid methyl ester; 6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)picolinic acid methyl ester; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzyl)-1-(p-tolyl)methanamine; 3-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)-N,N-dimethylpropan-1-amine; 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-

methoxybenzyl)-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethoxy)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-chlorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester; 7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-indazol-6-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(4-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,5-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,6-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-Chloro-2-(4-methoxybenzyl)-5-(3-nitrophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one;

Example # 1

Preparation of N-(1H-indazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 327.1386 g/mol

HPLC-MS: analytical method A

rt: 2.194 min - found mass: 328 (m/z+H)

Example # 2

Preparation of N-(1H-benzo[d][1,2,3]triazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-benzo[d][1,2,3]triazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 328.1316 g/mol

HPLC-MS: analytical method A

rt: 2.336 min - found mass: 329 (m/z+H)

Example # 3

Preparation of 5-phenyl-N-(4-(piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(piperidin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 370.2241 g/mol

HPLC-MS: analytical method A

rt: 2.245 min - found mass: 371 (m/z+H)

Example # 4

Preparation of N1,N1-dimethyl-N4-(5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 330.1856 g/mol

HPLC-MS: analytical method I

rt: 3.175 min - found mass: 331 (m/z+H)

Example # 5

Preparation of 3-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzamide

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-aminobenzamide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 330.1405 g/mol

HPLC-MS: analytical method A

rt: 2.290 min - found mass: 331 (m/z+H)

Example # 6

Preparation of N-(3,4-dimethoxyphenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL).

The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 347.1618 g/mol

HPLC-MS: analytical method A

rt: 2.544 min - found mass: 348 (m/z+H)

Example # 7

Preparation of N-(4-morpholinophenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 372.199 g/mol

HPLC-MS: analytical method A

rt: 2.546 min - found mass: 373 (m/z+H)

Example # 8

Preparation of N-(3-methoxyphenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-methoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 317.1479 g/mol

HPLC-MS: analytical method A

rt: 2.867 min - found mass: 318 (m/z+H)

Example # 9

Preparation of N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-methyl-1H-benzo[d]imidazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 341.1581 g/mol

HPLC-MS: analytical method I

rt: 2.576 min - found mass: 342 (m/z+H)

Example # 10

Preparation of 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzoic-acid

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-aminobenzoic-acid (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at

140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 331.1224 g/mol

HPLC-MS: analytical method A

rt: 2.581 min - found mass: 332 (m/z+H)

Example # 11

Preparation of 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzamide

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-aminobenzamide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 330.1405 g/mol

HPLC-MS: analytical method A

rt: 2.287 min - found mass: 331 (m/z+H)

Example # 12

Preparation of 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzonitrile

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-aminobenzonitrile (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 312.126 g/mol

HPLC-MS: analytical method A

rt: 3.081 min - found mass: 313 (m/z+H)

Example # 13

Preparation of N-(1H-indazol-5-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 328.1316 g/mol

HPLC-MS: analytical method A

rt: 1.850 min - found mass: 329 (m/z+H)

Example # 14

Preparation of N-(1H-indazol-5-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was

added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 397.2021 g/mol

HPLC-MS: analytical method A

rt: 2.011 min - found mass: 398 (m/z+H)

Example # 15

Preparation of 2-((5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzamide

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-aminobenzamide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 390.1682 g/mol

HPLC-MS: analytical method A

rt: 1.960 min - found mass: 391 (m/z+H)

Example # 16

Preparation of N-(1H-indazol-6-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 328.1316 g/mol

HPLC-MS: analytical method A

rt: 1.935 min - found mass: 329 (m/z+H)

Example # 17

Preparation of N1,N1-dimethyl-N4-(5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 331.1786 g/mol

HPLC-MS: analytical method A

rt: 0.678 min - found mass: 332 (m/z+H)

Example # 18

Preparation of N-(3,4-dimethoxyphenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial

(2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 348.1548 g/mol

HPLC-MS: analytical method A

rt: 2.013 min - found mass: 349 (m/z+H)

Example # 19

Preparation of N-(4-morpholinophenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 373.192 g/mol

HPLC-MS: analytical method A

rt: 1.974 min - found mass: 374 (m/z+H)

Example # 20

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 386.2296 g/mol

HPLC-MS: analytical method A

rt: 0.504 min - found mass: 387 (m/z+H)

Example # 21

Preparation of 6-((5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 359.1274 g/mol

HPLC-MS: analytical method A

rt: 1.930 min - found mass: 360 (m/z+H)

Example # 22

Preparation of 5-(3,4-dimethoxyphenyl)-N-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine

(0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 387.1664 g/mol

HPLC-MS: analytical method A

rt: 2.233 min - found mass: 388 (m/z+H)

Example # 23

Preparation of N1-(5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 390.2133 g/mol

HPLC-MS: analytical method A

rt: 2.173 min - found mass: 391 (m/z+H)

Example # 24

Preparation of N,5-bis(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 407.1896 g/mol

HPLC-MS: analytical method A

rt: 2.303 min - found mass: 408 (m/z+H)

Example # 25

Preparation of 5-(3,4-dimethoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 432.2267 g/mol

HPLC-MS: analytical method A

rt: 2.282 min - found mass: 433 (m/z+H)

Example # 26

Preparation of 6-((5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 418.1621 g/mol

HPLC-MS: analytical method A

rt: 2.218 min - found mass: 419 (m/z+H)

Example # 27

Preparation of N-(1H-indazol-6-yl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 348.211 g/mol

HPLC-MS: analytical method A

rt: 0.486 min - found mass: 349 (m/z+H)

Example # 28

Preparation of N1,N1-dimethyl-N4-(5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 351.258 g/mol

HPLC-MS: analytical method A

rt: 0.293 min - found mass: 352 (m/z+H)

Example # 29

Preparation of N-(4-chlorophenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-chloroaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 342.1648 g/mol

HPLC-MS: analytical method A

rt: 0.935 min - found mass: 343 (m/z+H)

Example # 30

Preparation of N-(1H-indazol-6-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 397.2021 g/mol

HPLC-MS: analytical method A

rt: 2.060 min - found mass: 398 (m/z+H)

Example # 31

Preparation of N1,N1-dimethyl-N4-(5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction

mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 400.2491 g/mol

HPLC-MS: analytical method A

rt: 1.890 min - found mass: 401 (m/z+H)

Example # 32

Preparation of N-(3,4-dimethoxyphenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 417.2253 g/mol

HPLC-MS: analytical method I

rt: 2.551 min - found mass: 418 (m/z+H)

Example # 33

Preparation of N-(4-morpholinophenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 442.2624 g/mol

HPLC-MS: analytical method I

rt: 2.553 min - found mass: 443 (m/z+H)

Example # 34

Preparation of 5-(3,4-dimethoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 445.2643 g/mol

HPLC-MS: analytical method A

rt: 1.761 min - found mass: 446 (m/z+H)

Example # 35

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was

added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 385.2366 g/mol

HPLC-MS: analytical method I

rt: 2.028 min - found mass: 386 (m/z+H)

Example # 36

Preparation of N-(1H-indazol-6-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 327.1386 g/mol

HPLC-MS: analytical method A

rt: 2.428 min - found mass: 328 (m/z+H)

Example # 37

Preparation of 6-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 358.1344 g/mol

HPLC-MS: analytical method A

rt: 2.390 min - found mass: 359 (m/z+H)

Example # 38

Preparation of 5-phenyl-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 277.1205 g/mol

HPLC-MS: analytical method A

rt: 2.202 min - found mass: 278 (m/z+H)

Example # 39

Preparation of 5-phenyl-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(pyrrolidin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave

vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 356.2046 g/mol

HPLC-MS: analytical method A

rt: 2.714 min - found mass: 357 (m/z+H)

Example # 40

Preparation of 5-(3,4-dimethoxyphenyl)-N-(1H-indazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 387.1664 g/mol

HPLC-MS: analytical method A

rt: 2.102 min - found mass: 388 (m/z+H)

Example # 41

Preparation of 5-(3,4-dimethoxyphenyl)-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 337.1483 g/mol

HPLC-MS: analytical method A

rt: 2.046 min - found mass: 338 (m/z+H)

Example # 42

Preparation of 5-(3,4-dimethoxyphenyl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(pyrrolidin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 416.2323 g/mol

HPLC-MS: analytical method A

rt: 2.560 min - found mass: 417 (m/z+H)

Example # 43

Preparation of N-(1H-pyrazol-3-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-pyrazol-3-amine (0.3 mmol 2

eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 278.1135 g/mol

HPLC-MS: analytical method A

rt: 0.823 min - found mass: 279 (m/z+H)

Example # 44

Preparation of 5-(pyridin-4-yl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(pyrrolidin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was

concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 357.1976 g/mol

HPLC-MS: analytical method A

rt: 2.138 min - found mass: 358 (m/z+H)

Example # 45

Preparation of N,5-di(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrazol-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 267.1069 g/mol

HPLC-MS: analytical method A

rt: 0.817 min - found mass: 268 (m/z+H)

Example # 46

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 455.3001 g/mol

HPLC-MS: analytical method A

rt: 1.038 min - found mass: 456 (m/z+H)

Example # 47

Preparation of 6-((5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 428.1979 g/mol

HPLC-MS: analytical method A

rt: 2.047 min - found mass: 429 (m/z+H)

Example # 48

Preparation of N-(1H-pyrazol-3-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was

added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 347.184 g/mol

HPLC-MS: analytical method A

rt: 1.867 min - found mass: 348 (m/z+H)

Example # 49

Preparation of N-(4-(pyrrolidin-1-yl)phenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(pyrrolidin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 426.2681 g/mol

HPLC-MS: analytical method A

rt: 2.326 min - found mass: 427 (m/z+H)

Example # 50

Preparation of N-(1H-indazol-5-yl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 348.211 g/mol

HPLC-MS: analytical method I

rt: 0.465 min - found mass: 349 (m/z+H)

Example # 51

Preparation of N-(3,4-dimethoxyphenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 368.2343 g/mol

HPLC-MS: analytical method I

rt: 0.331 min - found mass: 369 (m/z+H)

Example # 52

Preparation of 5-(1-methylpiperidin-4-yl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a

microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 393.2714 g/mol

HPLC-MS: analytical method I

rt: 0.331 min - found mass: 394 (m/z+H)

Example # 53

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 406.309 g/mol

HPLC-MS: analytical method I

rt: 0.326 min - found mass: 407 (m/z+H)

Example # 54

Preparation of 6-((5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 379.2068 g/mol

HPLC-MS: analytical method I

rt: 0.331 min - found mass: 380 (m/z+H)

Example # 55

Preparation of 5-(1-methylpiperidin-4-yl)-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 298.193 g/mol

HPLC-MS: analytical method I

rt: 0.344 min - found mass: 299 (m/z+H)

Example # 56

Preparation of 5-(1-methylpiperidin-4-yl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(pyrrolidin-1-

yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 377.277 g/mol.

HPLC-MS: analytical method I

rt: 2.282 min - found mass: 378 (m/z+H)

Example # 57

Preparation of N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 5-cyclopropyl-1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 317.159 g/mol

HPLC-MS: analytical method A

rt: 2.540 min - found mass: 318 (m/z+H)

Example # 58

Preparation of N-(3,4-dimethoxyphenyl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 377.1757 g/mol

HPLC-MS: analytical method A

rt: 2.386 min - found mass: 378 (m/z+H)

Example # 59

Preparation of 5-(4-methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 402.2128 g/mol

HPLC-MS: analytical method A

rt: 2.336 min - found mass: 403 (m/z+H)

Example # 60

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 377.1757 g/mol

HPLC-MS: analytical method A

rt: 2.561 min - found mass: 378 (m/z+H)

Example # 61

Preparation of 5-(3-methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 402.2128 g/mol

HPLC-MS: analytical method A

rt: 2.487 min - found mass: 403 (m/z+H)

Example # 62

Preparation of 5-(3-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 415.2504 g/mol

HPLC-MS: analytical method B

rt: 1.631 min - found mass: 416 (m/z+H)

Example # 63

Preparation of N1-(5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 360.1994 g/mol

HPLC-MS: analytical method A

rt: 2.312 min - found mass: 361 (m/z+H)

Example # 64

Preparation of N-(5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 344.16 g/mol

HPLC-MS: analytical method A

rt: 2.358 min - found mass: 345 (m/z+H)

Example # 65

Preparation of 3-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-aminophenol (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The

reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 303.1284 g/mol

HPLC-MS: analytical method A

rt: 2.408 min - found mass: 304 (m/z+H)

Example # 66

Preparation of 5-(4-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 415.2504 g/mol

HPLC-MS: analytical method B

rt: 1.540 min - found mass: 416 (m/z+H)

Example # 67

Preparation of N-(1H-indazol-6-yl)-5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 357.1524 g/mol

HPLC-MS: analytical method A

rt: 2.485 min - found mass: 358 (m/z+H)

Example # 68

Preparation of N-(3,4-dimethoxyphenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 353.1143 g/mol

HPLC-MS: analytical method A

rt: 2.541 min - found mass: 354 (m/z+H)

Example # 69

Preparation of N-(4-morpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave

reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 378.1515 g/mol

HPLC-MS: analytical method A

rt: 2.497 min - found mass: 379 (m/z+H)

Example # 70

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 391.1891 g/mol

HPLC-MS: analytical method B

rt: 1.587 min - found mass: 392 (m/z+H)

Example # 71

Preparation of N1,N1-dimethyl-N4-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 336.138 g/mol

HPLC-MS: analytical method A

rt: 2.230 min - found mass: 337 (m/z+H)

Example # 72

Preparation of N-(3,4-dimethoxyphenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 353.1143 g/mol

HPLC-MS: analytical method A

rt: 2.319 min - found mass: 354 (m/z+H)

Example # 73

Preparation of N-(4-morpholinophenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was

added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 378.1515 g/mol

HPLC-MS: analytical method A

rt: 2.288 min - found mass: 379 (m/z+H)

Example # 74

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 391.1891 g/mol

HPLC-MS: analytical method B

rt: 1.478 min - found mass: 392 (m/z+H)

Example # 75

Preparation of N1,N1-dimethyl-N4-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 336.138 g/mol

HPLC-MS: analytical method A

rt: 2.121 min - found mass: 337 (m/z+H)

Example # 76

Preparation of N-(1H-indazol-6-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 333.091 g/mol

HPLC-MS: analytical method A

rt: 2.242 min - found mass: 334 (m/z+H)

Example # 77

Preparation of N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 5-cyclopropyl-1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry,

3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 323.1115 g/mol

HPLC-MS: analytical method A

rt: 2.379 min - found mass: 324 (m/z+H)

Example # 78

Preparation of N1-(5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,4-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 360.1994 g/mol

HPLC-MS: analytical method A

rt: 2.244 min - found mass: 361 (m/z+H)

Example # 79

Preparation of N-(1H-indazol-6-yl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 357.1524 g/mol

HPLC-MS: analytical method A

rt: 2.294 min - found mass: 358 (m/z+H)

Example # 80

Preparation of 3-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-aminophenol (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 333.1423 g/mol

HPLC-MS: analytical method A

rt: 2.253 min - found mass: 334 (m/z+H)

Example # 81

Preparation of 3-((5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-aminophenol (0.3

mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 333.1423 g/mol

HPLC-MS: analytical method A

rt: 2.486 min - found mass: 334 (m/z+H)

Example # 82

Preparation of N-(1H-indazol-6-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 333.091 g/mol

HPLC-MS: analytical method A

rt: 2.463 min - found mass: 334 (m/z+H)

Example # 83

Preparation of 3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-aminophenol (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 309.0809 g/mol

HPLC-MS: analytical method A

rt: 2.441 min - found mass: 310 (m/z+H)

Example # 84

Preparation of N-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 350.1124 g/mol

HPLC-MS: analytical method A

rt: 2.255 min - found mass: 351 (m/z+H)

Example # 85

Preparation of 3-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-aminophenol (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 309.0809 g/mol

HPLC-MS: analytical method A

rt: 2.255 min - found mass: 310 (m/z+H)

Example # 86

Preparation of N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 5-cyclopropyl-1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 347.1729 g/mol

HPLC-MS: analytical method A

rt: 2.412 min - found mass: 348 (m/z+H)

Example # 87

Preparation of N-(5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 374.1738 g/mol

HPLC-MS: analytical method A

rt: 2.296 min - found mass: 375 (m/z+H)

Example # 88

Preparation of N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 5-cyclopropyl-1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 347.1729 g/mol

HPLC-MS: analytical method A

rt: 2.604 min - found mass: 348 (m/z+H)

Example # 89

Preparation of N-(5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 374.1738 g/mol

HPLC-MS: analytical method A

rt: 2.434 min - found mass: 375 (m/z+H)

Example # 90

Preparation of N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 5-cyclopropyl-1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue

was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 323.1115 g/mol

HPLC-MS: analytical method F

rt: 5.010 min - found mass: 324 (m/z+H)

Example # 91

Preparation of N-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 350.1124 g/mol

HPLC-MS: analytical method A

rt: 2.425 min - found mass: 351 (m/z+H)

Example # 92

Preparation of N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 5-cyclopropyl-1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 338.2315 g/mol

HPLC-MS: analytical method A

rt: 0.330 min - found mass: 339 (m/z+H)

Example # 93

Preparation of N-(5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 365.2325 g/mol

HPLC-MS: analytical method A

rt: 0.321 min - found mass: 366 (m/z+H)

Example # 94

Preparation of 3-((5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol

7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-aminophenol (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5

min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 324.2009 g/mol

HPLC-MS: analytical method A

rt: 0.308 min - found mass: 325 (m/z+H)

Example # 95

Preparation of 5-(benzo[d]thiazol-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 442.2001 g/mol

HPLC-MS: analytical method A

rt: 2.111 min - found mass: 443 (m/z+H)

Example # 96

Preparation of 5-(benzo[d]thiazol-2-yl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 404.1254 g/mol

HPLC-MS: analytical method A

rt: 3.058 min - found mass: 405 (m/z+H)

Example # 97

Preparation of N-(5-(benzo[d]thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 401.1235 g/mol

HPLC-MS: analytical method A

rt: 2.909 min - found mass: 402 (m/z+H)

Example # 98

Preparation of 5-(benzo[d]thiazol-2-yl)-N-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave

vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 384.1021 g/mol

HPLC-MS: analytical method C

rt: 2.360 min - found mass: 385 (m/z+H)

Example # 99

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 462.2667 g/mol

HPLC-MS: analytical method A

rt: 1.861 min - found mass: 463 (m/z+H)

Example # 100

Preparation of N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 5-cyclopropyl-1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 394.1891 g/mol

HPLC-MS: analytical method A

rt: 2.318 min - found mass: 395 (m/z+H)

Example # 101

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 424.1919 g/mol

HPLC-MS: analytical method A

rt: 2.349 min - found mass: 425 (m/z+H)

Example # 102

Preparation of N-(4-morpholinophenyl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline

(0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 449.2291 g/mol

HPLC-MS: analytical method A

rt: 2.299 min - found mass: 450 (m/z+H)

Example # 103

Preparation of N-(5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 421.1901 g/mol

HPLC-MS: analytical method A

rt: 2.218 min - found mass: 422 (m/z+H)

Example # 104

Preparation of N-(1H-indazol-6-yl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 404.1687 g/mol

HPLC-MS: analytical method A

rt: 2.238 min - found mass: 405 (m/z+H)

Example # 105

Preparation of 2-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 401.231 g/mol

HPLC-MS: analytical method A

rt: 2.011 min - found mass: 402 (m/z+H)

Example # 106

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

4-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)morpholine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 470.3015 g/mol

HPLC-MS: analytical method A

rt: 1.793 min - found mass: 471 (m/z+H)

Example # 107

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

4-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)morpholine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 432.2267 g/mol

HPLC-MS: analytical method A

rt: 2.410 min - found mass: 433 (m/z+H)

Example # 108

Preparation of 5-(2-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 415.2504 g/mol

HPLC-MS: analytical method A

rt: 1.338 min - found mass: 416 (m/z+H)

Example # 109

Preparation of N-(3,4-dimethoxyphenyl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 377.1757 g/mol

HPLC-MS: analytical method A

rt: 2.215 min - found mass: 378 (m/z+H)

Example # 110

Preparation of N-(5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 374.1738 g/mol

HPLC-MS: analytical method A

rt: 2.227 min - found mass: 375 (m/z+H)

Example # 111

Preparation of N-(1H-indazol-6-yl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue

was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 357.1524 g/mol

HPLC-MS: analytical method A

rt: 2.125 min - found mass: 358 (m/z+H)

Example # 112

Preparation of 5-(2-methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 402.2128 g/mol

HPLC-MS: analytical method A

rt: 2.252 min - found mass: 403 (m/z+H)

Example # 113

Preparation of N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 5-cyclopropyl-1H-pyrazol-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 347.1729 g/mol

HPLC-MS: analytical method A

rt: 2.242 min - found mass: 348 (m/z+H)

Example # 114

Preparation of N1,N1-dimethyl-N3-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,3-diamine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and N1,N1-dimethylbenzene-1,3-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 336.138 g/mol

HPLC-MS: analytical method C

rt: 2.085 min - found mass: 337 (m/z+H)

Example # 115

Preparation of (1R,2R)-N-(5-Thiophen-3-yl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-cyclohexane-1,2-diamine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and ((1R,2R)-2-Amino-cyclohexyl)-carbamic acid tert-butyl ester (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture

was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 314.159 g/mol

HPLC-MS: analytical method C

rt: 1.411 min - found mass: 315 (m/z+H)

Example # 116

Preparation of (1S,2R)-N-(5-Thiophen-3-yl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-cyclohexane-1,2-diamine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and ((1S,2R)-2-Amino-cyclohexyl)-carbamic acid tert-butyl ester (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 314.159 g/mol

HPLC-MS: analytical method C

rt: 1.388 min - found mass: 315 (m/z+H)

Example # 117

Preparation of N1-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,3-diamine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and benzene-1,3-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 308.099 g/mol

HPLC-MS: analytical method A

rt: 1.937 min - found mass: 309 (m/z+H)

Example # 118

Preparation of N-(6-(4-methylpiperazin-1-yl)pyridin-3-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-(4-methylpiperazin-1-yl)pyridin-3-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 392.1821 g/mol

HPLC-MS: analytical method C

rt: 1.428 min - found mass: 393 (m/z+H)

Example # 119

Preparation of N-(4-(4-ethylpiperazin-1-yl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-ethylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 399.2561 g/mol

HPLC-MS: analytical method C

rt: 1.586 min - found mass: 400 (m/z+H)

Example # 120

Preparation of N-(4-((dimethylamino)methyl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-((dimethylamino)methyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 344.2051 g/mol

HPLC-MS: analytical method C

rt: 1.601 min - found mass: 345 (m/z+H)

Example # 121

Preparation of 5-(2,5-dimethoxyphenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,5-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 407.1896 g/mol

HPLC-MS: analytical method A

rt: 2.363 min - found mass: 408 (m/z+H)

Example # 122

Preparation of 5-(2,5-dimethoxyphenyl)-N-(4-(4-ethylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,5-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-ethylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 459.2839 g/mol

HPLC-MS: analytical method A

rt: 1.848 min - found mass: 460 (m/z+H)

Example # 123

Preparation of N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2,5-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,5-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-cyclopropylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 471.2833 g/mol

HPLC-MS: analytical method A

rt: 1.976 min - found mass: 472 (m/z+H)

Example # 124

Preparation of N-(3,4-dimethoxyphenyl)-5-(imidazo[1,2-a]pyridin-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was

evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 387.1664 g/mol

HPLC-MS: analytical method A

rt: 1.962 min - found mass: 388 (m/z+H)

Example # 125

Preparation of N-(4-(4-ethylpiperazin-1-yl)phenyl)-5-(imidazo[1,2-a]pyridin-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-ethylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 439.2607 g/mol

HPLC-MS: analytical method C

rt: 1.787 min - found mass: 440 (m/z+H)

Example # 126

Preparation of 5-(imidazo[1,2-a]pyridin-2-yl)-N-(1H-indazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 367.1431 g/mol

HPLC-MS: analytical method A

rt: 1.835 min - found mass: 368 (m/z+H)

Example # 127

Preparation of 5-(imidazo[1,2-a]pyridin-2-yl)-N-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 367.1431 g/mol

HPLC-MS: analytical method A

rt: 1.900 min - found mass: 368 (m/z+H)

Example # 128

Preparation of 4-((5-(imidazo[1,2-a]pyridin-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzoic-acid

7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-aminobenzoic-acid (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was

added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 371.1269 g/mol

HPLC-MS: analytical method C

rt: 2.104 min - found mass: 372 (m/z+H)

Example # 129

Preparation of 5-(1H-benzo[d]imidazol-2-yl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

5-(1H-benzo[d]imidazol-2-yl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 387.1664 g/mol

HPLC-MS: analytical method A

rt: 2.255 min - found mass: 388 (m/z+H)

Example # 130

Preparation of N-(3,4-dimethoxyphenyl)-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 401.1859 g/mol

HPLC-MS: analytical method A

rt: 2.267 min - found mass: 402 (m/z+H)

Example # 131

Preparation of 5-(3-methoxyphenyl)-N-(4-((4-methylpiperazin-1-yl)methyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-Amino-phenyl)-(4-methyl-piperazin-1-yl)-methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and dissolved in THF (dry) LiAlH₄ powder was added (excess, 2 by 2 eq) until completion of reaction is observed (by LCMS). The reaction was quenched with water (1mL per gram LiAlH₄), then NaOH (ca. 15% aq., 1mL per g LiAlH₄), water (3mL per gram LiAlH₄). The mixture was filtered, washed with THF, MeOH, MeCN (ca. 10ML each). The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 429.27 g/mol

HPLC-MS: analytical method A

rt: 2.239 min - found mass: 430 (m/z+H)

Example # 132

Preparation of 5-(3-methoxyphenyl)-N-(4-(pyrrolidin-1-ylmethoxy)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-Amino-phenyl)-pyrrolidin-1-yl-methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and dissolved in THF (dry) LiAlH₄ powder was added (excess, 2 by 2 eq) until completion of reaction is observed (by LCMS). The reaction was quenched with water (1mL per gram LiAlH₄), then NaOH (ca. 15% aq., 1mL per g LiAlH₄), water (3mL per gram LiAlH₄). The mixture was filtered, washed with THF, MeOH, MeCN (ca. 10ML each). The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 400.238 g/mol

HPLC-MS: analytical method A

rt: 2.021 min - found mass: 401 (m/z+H)

Example # 133

Preparation of 1-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)-3-(m-tolyl)urea

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1-(4-aminophenyl)-3-(m-tolyl)urea (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 441.1620 g/mol

HPLC-MS: analytical method C

rt: 2.74 min - found mass: 442 (m/z+H)

Example # 134

Preparation of (4-methylpiperazin-1-yl)(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-aminophenyl)(4-methylpiperazin-1-yl)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The

reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 419.1830 g/mol

HPLC-MS: analytical method C

rt: 1.97 min - found mass: 420 (m/z+H)

Example # 135

Preparation of pyrrolidin-1-yl(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-aminophenyl)(pyrrolidin-1-yl)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 390.1510 g/mol

HPLC-MS: analytical method J

rt: 5.14 min - found mass: 391 (m/z+H)

Example # 136

Preparation of N-(4-thiomorpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-thiomorpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 394.1286 g/mol

HPLC-MS: analytical method C

rt: 2.81 min - found mass: 395 (m/z+H)

Example # 137

Preparation of 4-(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)thiomorpholine 1,1-dioxide

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-aminophenyl)thiomorpholine 1,1-dioxide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 426.1174 g/mol

HPLC-MS: analytical method A

rt: 2.32 min - found mass: 427 (m/z+H)

Example # 138

Preparation of 1-(4-(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl

in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 419.1830 g/mol

HPLC-MS: analytical method J

rt: 4.21 min - found mass: 420 (m/z+H)

Example # 139

Preparation of (3-methoxy-4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(4-(4-methylpiperazin-1-yl)piperidin-1-yl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-amino-3-methoxyphenyl)(4-(4-methylpiperazin-1-yl)piperidin-1-yl)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 532.2869 g/mol

HPLC-MS: analytical method A

rt: 1.91 min - found mass: 533 (m/z+H)

Example # 140

Preparation of 5-(1H-imidazol-5-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(1H-imidazol-5-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 375.2231 g/mol

HPLC-MS: analytical method A

rt: 0.47 min - found mass: 376 (m/z+H)

Example # 141

Preparation of 5-(1H-imidazol-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(1H-imidazol-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 375.2231 g/mol

HPLC-MS: analytical method A

rt: 0.41 min - found mass: 376 (m/z+H)

Example # 142

Preparation of pyrrolidin-1-yl(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-aminophenyl)(pyrrolidin-1-yl)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 390.1510 g/mol

HPLC-MS: analytical method A

rt: 2.40 min - found mass: 391 (m/z+H)

Example # 143

Preparation of 4-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)thiomorpholine 1,1-dioxide

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-aminophenyl)thiomorpholine 1,1-dioxide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further

purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 426.1174 g/mol

HPLC-MS: analytical method A

rt: 2.14 min - found mass: 427 (m/z+H)

Example # 144

Preparation of 1-(4-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 419.1830 g/mol

HPLC-MS: analytical method A

rt: 2.11 min - found mass: 420 (m/z+H)

Example # 145

Preparation of N-(4-thiomorpholinophenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-thiomorpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 394.1286 g/mol

HPLC-MS: analytical method A

rt: 2.50 min - found mass: 395 (m/z+H)

Example # 146

Preparation of 6-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 364.0869 g/mol

HPLC-MS: analytical method A

rt: 2.17 min - found mass: 365 (m/z+H)

Example # 147

Preparation of (4-methylpiperazin-1-yl)(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-aminophenyl)(4-methylpiperazin-1-yl)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl

in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 419.1830 g/mol

HPLC-MS: analytical method A

rt: 0.29 min - found mass: 420 (m/z+H)

Example # 148

Preparation of 2,2-dimethyl-N-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2,2-dimethyl-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and

purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 394.1286 g/mol

HPLC-MS: analytical method C

rt: 2.95 min - found mass: 395 (m/z+H)

Example # 149

Preparation of 5-(3-(2H-tetrazol-5-yl)phenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 453.2462 g/mol

HPLC-MS: analytical method L

rt: 3.29 min - found mass: 454 (m/z+H)

Example # 150

Preparation of (7-((2-(dimethylamino)ethyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone

(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone (0.16 mmol) and N1,N1-dimethylethane-1,2-diamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 310.1815 g/mol

HPLC-MS: analytical method L

rt: 3.17 min - found mass: 311 (m/z+H)

Example # 151

Preparation of (7-(((1-methylpiperidin-4-yl)methyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone

(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone (0.16 mmol) and (1-methylpiperidin-4-yl)methanamine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 350.2200 g/mol

HPLC-MS: analytical method L

rt: 2.92 min - found mass: 351 (m/z+H)

Example # 152

Preparation of (7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone

(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue

was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 413.2305 g/mol

HPLC-MS: analytical method L

rt: 3.50 min - found mass: 414 (m/z+H)

Example # 153

Preparation of (7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone

(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 375.1558 g/mol

HPLC-MS: analytical method L

rt: 4.51 min - found mass: 376 (m/z+H)

Example # 154

Preparation of (4-((5-benzoyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(pyrrolidin-1-yl)methanone

(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone (0.16 mmol) and (4-aminophenyl)(pyrrolidin-1-yl)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 412.1924 g/mol

HPLC-MS: analytical method L

rt: 4.42 min - found mass: 413 (m/z+H)

Example # 155

Preparation of 7-((5-benzoyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 384.1534 g/mol

HPLC-MS: analytical method L

rt: 4.22 min - found mass: 385 (m/z+H)

Example # 156

Preparation of 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid

Methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave

reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 416.1872 g/mol

HPLC-MS: analytical method A

rt: 2.39 min - found mass: 417 (m/z+H)

Example # 157

Preparation of morpholino(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-aminophenyl)(morpholino)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The

reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 406.1454 g/mol

HPLC-MS: analytical method A

rt: 2.44 min - found mass: 407 (m/z+H)

Example # 158

Preparation of 6-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]thiazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]thiazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 380.0640 g/mol

HPLC-MS: analytical method A

rt: 2.58 min - found mass: 381 (m/z+H)

Example # 159

Preparation of 7-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 386.1733 g/mol

HPLC-MS: analytical method A

rt: 2.30 min - found mass: 387 (m/z+H)

Example # 160

Preparation of (4-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(pyrrolidin-1-yl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-aminophenyl)(pyrrolidin-1-yl)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 414.2123 g/mol

HPLC-MS: analytical method A

rt: 2.44 min - found mass: 415 (m/z+H)

Example # 161

Preparation of N-(4-(4-(tert-butyl)piperazin-1-yl)phenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-(tert-butyl)piperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane

(4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 433.2476 g/mol

HPLC-MS: analytical method L

rt: 3.23 min - found mass: 434 (m/z+H)

Example # 162

Preparation of 6-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 364.0869 g/mol

HPLC-MS: analytical method A

rt: 2.40 min - found mass: 365 (m/z+H)

Example # 163

Preparation of N-(4-(4-(tert-butyl)piperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-(tert-butyl)piperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 433.2476 g/mol

HPLC-MS: analytical method A

rt: 1.94 min - found mass: 434 (m/z+H)

Example # 164

Preparation of 8-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-4,5-dihydro-1H-benzo[b]azepin-2(3H)-one

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 8-amino-4,5-dihydro-1H-benzo[b]azepin-2(3H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 400.1928 g/mol

HPLC-MS: analytical method A

rt: 2.33 min - found mass: 401 (m/z+H)

Example # 165

Preparation of 6-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]thiazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-

benzo[b][1,4]thiazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 380.0640 g/mol

HPLC-MS: analytical method A

rt: 2.63 min - found mass: 381 (m/z+H)

Example # 166

Preparation of 1-methyl-N-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-1,2,3,4-tetrahydroquinolin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1-methyl-1,2,3,4-tetrahydroquinolin-7-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 362.1571 g/mol

HPLC-MS: analytical method L

rt: 4.78 min - found mass: 363 (m/z+H)

Example # 167

Preparation of 2-methyl-2-(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)propanenitrile

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-(4-aminophenyl)-2-methylpropanenitrile (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 360.1371 g/mol

HPLC-MS: analytical method A

rt: 3.12 min - found mass: 361 (m/z+H)

Example # 168

Preparation of 5-(4-methoxyphenyl)-N-(4-thiomorpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-thiomorpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 418.1899 g/mol

HPLC-MS: analytical method A

rt: 2.57 min - found mass: 419 (m/z+H)

Example # 169

Preparation of 4-(4-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)thiomorpholine 1,1-dioxide

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-aminophenyl)thiomorpholine 1,1-dioxide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 450.1787 g/mol

HPLC-MS: analytical method A

rt: 2.23 min - found mass: 451 (m/z+H)

Example # 170

Preparation of 1-(4-(4-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further

purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 443.2443 g/mol

HPLC-MS: analytical method A

rt: 2.21 min - found mass: 444 (m/z+H)

Example # 171

Preparation of 6-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 388.1483 g/mol

HPLC-MS: analytical method A

rt: 2.26 min - found mass: 389 (m/z+H)

Example # 172

Preparation of N-(4-(4-(tert-butyl)piperazin-1-yl)phenyl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-(tert-butyl)piperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 457.3090 g/mol

HPLC-MS: analytical method C

rt: 2.13 min - found mass: 458 (m/z+H)

Example # 173

Preparation of 2,2-dimethyl-N-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2,2-dimethyl-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 394.1286 g/mol

HPLC-MS: analytical method A

rt: 3.18 min - found mass: 395 (m/z+H)

Example # 174

Preparation of 2-(3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)acetic acid

2-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)acetic acid methyl ester (0.16 mmol)

and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 459.2388 g/mol

HPLC-MS: analytical method L

rt: 3.04 min - found mass: 460 (m/z+H)

Example # 175

Preparation of 5-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)thiophene-3-carboxylic acid

5-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiophene-3-carboxylic acid methyl ester (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 435.1774 g/mol

HPLC-MS: analytical method L

rt: 3.19 min - found mass: 436 (m/z+H)

Example # 176

Preparation of 2-(3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)acetic acid

2-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)acetic acid methyl ester (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 443.2444 g/mol

HPLC-MS: analytical method L

rt: 3.06 min - found mass: 444 (m/z+H)

Example # 177

Preparation of 6-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)picolinic acid

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)picolinic acid methyl ester (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 430.2179 g/mol

HPLC-MS: analytical method L

rt: 2.70 min - found mass: 431 (m/z+H)

Example # 178

Preparation of N-(4-(pentafluorosulfanyl)phenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-Aminophenylsulphur pentafluoride (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 419.0376 g/mol

HPLC-MS: analytical method C

rt: 3.40 min - found mass: 420 (m/z+H)

Example # 179

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 392.1821 g/mol

HPLC-MS: analytical method D

rt: 2.67 min - found mass: 393 (m/z+H)

Example # 180

Preparation of 5-(3-(((4-methylbenzyl)amino)methyl)phenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzyl)-1-(p-tolyl)methanamine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 518.3447 g/mol

HPLC-MS: analytical method L

rt: 3.20 min - found mass: 519 (m/z+H)

Example # 181

Preparation of 5-(3-(3-(dimethylamino)propoxy)phenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

3-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)-N,N-dimethylpropan-1-amine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 486.3410 g/mol

HPLC-MS: analytical method L

rt: 3.21 min - found mass: 487 (m/z+H)

Example # 182

Preparation of 3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue

was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 464.2094 g/mol

HPLC-MS: analytical method L

rt: 0.46 min - found mass: 465 (m/z+H)

Example # 183

Preparation of 3-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 426.1346 g/mol

HPLC-MS: analytical method L

rt: 3.91 min - found mass: 427 (m/z+H)

Example # 184

Preparation of 1-(4-(4-((5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 420.1760 g/mol

HPLC-MS: analytical method L

rt: 3.66 min - found mass: 421 (m/z+H)

Example # 185

Preparation of N-(3,4-dimethoxyphenyl)-5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 354.1073 g/mol

HPLC-MS: analytical method L

rt: 3.93 min - found mass: 355 (m/z+H)

Example # 186

Preparation of 7-((5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave

reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 363.1049 g/mol

HPLC-MS: analytical method A

rt: 2.30 min - found mass: 364 (m/z+H)

Example # 187

Preparation of N-(3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide

N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 478.2289 g/mol

HPLC-MS: analytical method D

rt: 3.27 min - found mass: 479 (m/z+H)

Example # 188

Preparation of N-(3-(7-((4-(piperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide

N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide (0.16 mmol) and 4-(piperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 464.2094 g/mol

HPLC-MS: analytical method L

rt: 2.90 min - found mass: 465 (m/z+H)

Example # 189

Preparation of N-(3-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide

N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 440.1541 g/mol

HPLC-MS: analytical method L

rt: 4.06 min - found mass: 441 (m/z+H)

Example # 190

Preparation of N-(3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide

N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 449.1518 g/mol

HPLC-MS: analytical method D

rt: 4.65 min - found mass: 450 (m/z+H)

Example # 191

Preparation of N-(3-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide

N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 506.2228 g/mol

HPLC-MS: analytical method L

rt: 3.75 min - found mass: 507 (m/z+H)

Example # 192

Preparation of 1-(4-(4-((5-(3-(2H-tetrazol-5-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 481.2401 g/mol

HPLC-MS: analytical method A

rt: 2.34 min - found mass: 482 (m/z+H)

Example # 193

Preparation of 5-(3-(2H-tetrazol-5-yl)phenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 415.1714 g/mol

HPLC-MS: analytical method A

rt: 2.49 min - found mass: 416 (m/z+H)

Example # 194

Preparation of 3-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 391.1501 g/mol

HPLC-MS: analytical method L

rt: 4.04 min - found mass: 392 (m/z+H)

Example # 195

Preparation of 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 400.1478 g/mol

HPLC-MS: analytical method L

rt: 4.02 min - found mass: 401 (m/z+H)

Example # 196

Preparation of 7-((5-(3-(2H-tetrazol-5-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 424.1690 g/mol

HPLC-MS: analytical method L

rt: 3.97 min - found mass: 425 (m/z+H)

Example # 197

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue

was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 392.1821 g/mol

HPLC-MS: analytical method CP

rt: 2.13 min - found mass: 393 (m/z+H)

Example # 198

Preparation of 1-(4-(4-((5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 420.1760 g/mol

HPLC-MS: analytical method L

rt: 3.32 min - found mass: 421 (m/z+H)

Example # 199

Preparation of N-(3,4-dimethoxyphenyl)-5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 354.1073 g/mol

HPLC-MS: analytical method L

rt: 3.47 min - found mass: 355 (m/z+H)

Example # 200

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 376.2050 g/mol

HPLC-MS: analytical method CP

rt: 2.18 min - found mass: 377 (m/z+H)

Example # 201

Preparation of 1-(4-(4-((5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave

reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 404.1989 g/mol

HPLC-MS: analytical method A

rt: 2.01 min - found mass: 405 (m/z+H)

Example # 202

Preparation of N-(3,4-dimethoxyphenyl)-5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 338.1302 g/mol

HPLC-MS: analytical method A

rt: 2.13 min - found mass: 339 (m/z+H)

Example # 203

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 376.2050 g/mol

HPLC-MS: analytical method CP

rt: 2.23 min - found mass: 377 (m/z+H)

Example # 204

Preparation of 1-(4-(4-((5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 404.1989 g/mol

HPLC-MS: analytical method L

rt: 3.59 min - found mass: 405 (m/z+H)

Example # 205

Preparation of N-(3,4-dimethoxyphenyl)-5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction

mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 338.1302 g/mol

HPLC-MS: analytical method L

rt: 3.88 min - found mass: 339 (m/z+H)

Example # 206

Preparation of 7-((5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 363.1049 g/mol

HPLC-MS: analytical method L

rt: 3.42 min - found mass: 364 (m/z+H)

Example # 207

Preparation of 7-((5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 347.1278 g/mol

HPLC-MS: analytical method L

rt: 3.47 min - found mass: 348 (m/z+H)

Example # 208

Preparation of 7-((5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 347.1278 g/mol

HPLC-MS: analytical method L

rt: 3.68 min - found mass: 348 (m/z+H)

Example # 209

Preparation of N-(4-(1-methylpiperidin-4-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(1-methylpiperidin-4-yl)aniline (0.3 mmol 2 eq.,) were suspended

in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 390.1961 g/mol

HPLC-MS: analytical method L

rt: 4.22 min - found mass: 391 (m/z+H)

Example # 210

Preparation of methyl 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 414.1673 g/mol

HPLC-MS: analytical method L

rt: 4.53 min - found mass: 415 (m/z+H)

Example # 211

Preparation of methyl 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 416.1422 g/mol

HPLC-MS: analytical method L

rt: 4.50 min - found mass: 417 (m/z+H)

Example # 212

Preparation of 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester (0.16 mmol) and 6-amino-2H-benzo[b][1,4]oxazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 402.1227 g/mol

HPLC-MS: analytical method D

rt: 4.69 min - found mass: 403 (m/z+H)

Example # 213

Preparation of methyl 4-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 443.2443 g/mol

HPLC-MS: analytical method L

rt: 3.54 min - found mass: 444 (m/z+H)

Example # 214

Preparation of methyl 4-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a

microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 405.1697 g/mol

HPLC-MS: analytical method L

rt: 4.81 min - found mass: 406 (m/z+H)

Example # 215

Preparation of methyl 4-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 414.1673 g/mol

HPLC-MS: analytical method L

rt: 4.54 min - found mass: 415 (m/z+H)

Example # 216

Preparation of methyl 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 6-amino-2H-benzo[b][1,4]thiazin-3(4H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 432.1193 g/mol

HPLC-MS: analytical method L

rt: 4.81 min - found mass: 433 (m/z+H)

Example # 217

Preparation of methyl 3-(7-((2-oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-8-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 8-amino-4,5-dihydro-1H-benzo[b]azepin-2(3H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 428.1868 g/mol

HPLC-MS: analytical method L

rt: 4.65 min - found mass: 429 (m/z+H)

Example # 218

Preparation of 3-(7-((2-oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-8-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester (0.16 mmol) and 8-amino-4,5-dihydro-1H-benzo[b]azepin-2(3H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 414.1673 g/mol

HPLC-MS: analytical method L

rt: 4.04 min - found mass: 415 (m/z+H)

Example # 219

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 473.1128 g/mol

HPLC-MS: analytical method D

rt: 7.77 min - found mass: 474 (m/z+H)

Example # 220

Preparation of 5-(3-(pentafluorosulfanyl)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3

drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 498.1500 g/mol

HPLC-MS: analytical method D

rt: 7.74 min - found mass: 499 (m/z+H)

Example # 221

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and

purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 511.1876 g/mol

HPLC-MS: analytical method D

rt: 5.24 min - found mass: 512 (m/z+H)

Example # 222

Preparation of 7-((5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 482.1105 g/mol

HPLC-MS: analytical method D

rt: 7.25 min - found mass: 483 (m/z+H)

Example # 223

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 393.1528 g/mol

HPLC-MS: analytical method D

rt: 6.18 min - found mass: 394 (m/z+H)

Example # 224

Preparation of 5-(3-(methylthio)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 418.1899 g/mol

HPLC-MS: analytical method D

rt: 6.11 min - found mass: 419 (m/z+H)

Example # 225

Preparation of N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated

in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 431.2275 g/mol

HPLC-MS: analytical method D

rt: 3.80 min - found mass: 432 (m/z+H)

Example # 226

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethoxy)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 431.1394 g/mol

HPLC-MS: analytical method A

rt: 3.35 min - found mass: 432 (m/z+H)

Example # 227

Preparation of N-(4-morpholinophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethoxy)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 456.1765 g/mol

HPLC-MS: analytical method A

rt: 3.31 min - found mass: 457 (m/z+H)

Example # 228

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-(trifluoromethyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 415.1450 g/mol

HPLC-MS: analytical method A

rt: 3.30 min - found mass: 416 (m/z+H)

Example # 229

Preparation of N-(4-morpholinophenyl)-5-(3-(trifluoromethyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was

added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 440.1821 g/mol

HPLC-MS: analytical method A

rt: 3.28 min - found mass: 441 (m/z+H)

Example # 230

Preparation of 5-(3-chlorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3-chlorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 381.1201 g/mol

HPLC-MS: analytical method A

rt: 3.23 min - found mass: 382 (m/z+H)

Example # 231

Preparation of 5-(3-chlorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3-chlorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 406.1572 g/mol

HPLC-MS: analytical method A

rt: 3.20 min - found mass: 407 (m/z+H)

Example # 232

Preparation of N-(3,4-dimethoxyphenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 365.1497 g/mol

HPLC-MS: analytical method A

rt: 2.64 min - found mass: 366 (m/z+H)

Example # 233

Preparation of 5-(2-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a

microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 390.1868 g/mol

HPLC-MS: analytical method A

rt: 2.59 min - found mass: 391 (m/z+H)

Example # 234

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 365.1497 g/mol

HPLC-MS: analytical method A

rt: 3.01 min - found mass: 366 (m/z+H)

Example # 235

Preparation of 5-(3-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 390.1868 g/mol

HPLC-MS: analytical method A

rt: 2.97 min - found mass: 391 (m/z+H)

Example # 236

Preparation of 7-((5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 402.1505 g/mol

HPLC-MS: analytical method D

rt: 5.89 min - found mass: 403 (m/z+H)

Example # 237

Preparation of 3-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide (0.16 mmol) and 1-(4-(4-

aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 492.2033 g/mol

HPLC-MS: analytical method L

rt: 3.83 min - found mass: 493 (m/z+H)

Example # 238

Preparation of 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 435.1323 g/mol

HPLC-MS: analytical method L

rt: 3.91 min - found mass: 436 (m/z+H)

Example # 239

Preparation of 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 451.1718 g/mol

HPLC-MS: analytical method L

rt: 4.03 min - found mass: 452 (m/z+H)

Example # 240

Preparation of 3-(7-((3-methoxy-4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide

3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide (0.16 mmol) and 3-methoxy-4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 481.1856 g/mol

HPLC-MS: analytical method L

rt: 3.99 min - found mass: 482 (m/z+H)

Example # 241

Preparation of 4-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 429.2249 g/mol

HPLC-MS: analytical method L

rt: 3.12 min - found mass: 430 (m/z+H)

Example # 242

Preparation of methyl 4-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a

microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 430.2067 g/mol

HPLC-MS: analytical method L

rt: 4.83 min - found mass: 431 (m/z+H)

Example # 243

Preparation of 4-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved

in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 416.1872 g/mol

HPLC-MS: analytical method L

rt: 4.07 min - found mass: 417 (m/z+H)

Example # 244

Preparation of 4-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid

4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. For the saponification the residue was dissolved in methanol. 2M aqueous NaOH-solution was added to ensure basic conditions. The mixture was irradiated in a microwave reactor for 10 min at 120°C. The reaction mixture was

concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 391.1501 g/mol

HPLC-MS: analytical method L

rt: 4.14 min - found mass: 392 (m/z+H)

Example # 245

Preparation of N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-methyl-1H-benzo[d]imidazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 347.1105 g/mol

HPLC-MS: analytical method D

rt: 3.80 min - found mass: 348 (m/z+H)

Example # 246

Preparation of 1-(4-(4-((5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 491.2103 g/mol

HPLC-MS: analytical method L

rt: 4.09 min - found mass: 492 (m/z+H)

Example # 247

Preparation of 5-(1H-indazol-6-yl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(1H-indazol-6-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 412.2036 g/mol

HPLC-MS: analytical method D

rt: 4.88 min - found mass: 413 (m/z+H)

Example # 248

Preparation of 5-(1H-indazol-6-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(1H-indazol-6-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated

in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 425.2411 g/mol

HPLC-MS: analytical method D

rt: 3.53 min - found mass: 426 (m/z+H)

Example # 249

Preparation of N-(3,4-dimethoxyphenyl)-5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 425.1416 g/mol

HPLC-MS: analytical method L

rt: 4.38 min - found mass: 426 (m/z+H)

Example # 250

Preparation of 7-((5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 434.1393 g/mol

HPLC-MS: analytical method L

rt: 4.04 min - found mass: 435 (m/z+H)

Example # 251

Preparation of 5-(3-(methylsulfonyl)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 450.1787 g/mol

HPLC-MS: analytical method L

rt: 4.35 min - found mass: 451 (m/z+H)

Example # 252

Preparation of N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-methyl-1H-benzo[d]imidazol-5-amine (0.3 mmol 2 eq.,) were suspended in

MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 419.1378 g/mol

HPLC-MS: analytical method L

rt: 3.21 min - found mass: 420 (m/z+H)

Example # 253

Preparation of N,5-bis(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-(methylsulfonyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 443.0937 g/mol

HPLC-MS: analytical method L

rt: 4.29 min - found mass: 444 (m/z+H)

Example # 254

Preparation of N-(3-(methylsulfonyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-(methylsulfonyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 371.0664 g/mol

HPLC-MS: analytical method L

rt: 4.43 min - found mass: 372 (m/z+H)

Example # 255

Preparation of 2-((3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)sulfonyl)ethan-1-ol

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-((3-aminophenyl)sulfonyl)ethan-1-ol (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 401.0803 g/mol

HPLC-MS: analytical method L

rt: 4.12 min - found mass: 402 (m/z+H)

Example # 256

Preparation of 1-(4-(4-((5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one

7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 453.2351 g/mol

HPLC-MS: analytical method L

rt: 3.32 min - found mass: 454 (m/z+H)

Example # 257

Preparation of 7-((5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction

mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 396.1640 g/mol

HPLC-MS: analytical method CP

rt: 2.46 min - found mass: 397 (m/z+H)

Example # 258

Preparation of N-(4-morpholinophenyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 412.2036 g/mol

HPLC-MS: analytical method L

rt: 3.40 min - found mass: 413 (m/z+H)

Example # 259

Preparation of N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-methyl-1H-benzo[d]imidazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 381.1626 g/mol

HPLC-MS: analytical method L

rt: 3.20 min - found mass: 382 (m/z+H)

Example # 260

Preparation of N,N-dimethyl-4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-amino-N,N-dimethylbenzenesulfonamide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 400.0984 g/mol

HPLC-MS: analytical method L

rt: 4.90 min - found mass: 401 (m/z+H)

Example # 261

Preparation of N-cyclopropyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-amino-N-cyclopropylbenzenesulfonamide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane

(4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 412.0979 g/mol

HPLC-MS: analytical method L

rt: 4.74 min - found mass: 413 (m/z+H)

Example # 262

Preparation of N-(3-(2H-1,2,3-triazol-2-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-(2H-1,2,3-triazol-2-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 360.1031 g/mol

HPLC-MS: analytical method D

rt: 6.78 min - found mass: 361 (m/z+H)

Example # 263

Preparation of N-(4-(2H-1,2,3-triazol-2-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(2H-1,2,3-triazol-2-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 360.1031 g/mol

HPLC-MS: analytical method L

rt: 5.10 min - found mass: 361 (m/z+H)

Example # 264

Preparation of N,N-dimethyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-amino-N,N-dimethylbenzenesulfonamide (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 400.0984 g/mol

HPLC-MS: analytical method A

rt: 2.93 min - found mass: 401 (m/z+H)

Example # 265

Preparation of N-methyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-amino-N-methylbenzenesulfonamide (0.3 mmol 2 eq.,) were suspended in

MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 386.0789 g/mol

HPLC-MS: analytical method A

rt: 2.68 min - found mass: 387 (m/z+H)

Example # 266

Preparation of N-(3-(morpholinosulfonyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-(morpholinosulfonyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 442.1118 g/mol

HPLC-MS: analytical method A

rt: 2.87 min - found mass: 443 (m/z+H)

Example # 267

Preparation of 5-(thiophen-2-yl)-N-(3-((trifluoromethyl)sulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-((trifluoromethyl)sulfonyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 425.0301 g/mol

HPLC-MS: analytical method A

rt: 3.37 min - found mass: 426 (m/z+H)

Example # 268

Preparation of N-(3-(methylsulfinyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-(methylsulfinyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 355.0720 g/mol

HPLC-MS: analytical method A

rt: 2.51 min - found mass: 356 (m/z+H)

Example # 269

Preparation of 7-((5-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one

7-chloro-5-(1H-indazol-6-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 396.1640 g/mol

HPLC-MS: analytical method D

rt: 4.76 min - found mass: 397 (m/z+H)

Example # 270

Preparation of N-(3,4-dimethoxyphenyl)-5-(4-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(4-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 365.1497 g/mol

HPLC-MS: analytical method A

rt: 2.88 min - found mass: 366 (m/z+H)

Example # 271

Preparation of 5-(4-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(4-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 390.1868 g/mol

HPLC-MS: analytical method A

rt: 2.83 min - found mass: 391 (m/z+H)

Example # 272

Preparation of 5-(3,4-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 383.1376 g/mol

HPLC-MS: analytical method A

rt: 3.15 min - found mass: 384 (m/z+H)

Example # 273

Preparation of 5-(3,4-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 408.1747 g/mol

HPLC-MS: analytical method A

rt: 3.13 min - found mass: 409 (m/z+H)

Example # 274

Preparation of 5-(3,5-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(3,5-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue

was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 383.1376 g/mol

HPLC-MS: analytical method A

rt: 3.24 min - found mass: 384 (m/z+H)

Example # 275

Preparation of 5-(2-fluorophenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-methylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 403.2245 g/mol

HPLC-MS: analytical method A

rt: 1.84 min - found mass: 404 (m/z+H)

Example # 276

Preparation of 5-(2-fluorophenyl)-N-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-(methylsulfonyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 383.1018 g/mol

HPLC-MS: analytical method A

rt: 2.63 min - found mass: 384 (m/z+H)

Example # 277

Preparation of 5-(2-fluorophenyl)-N-(2-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-(methylsulfonyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 383.1018 g/mol

HPLC-MS: analytical method A

rt: 2.89 min - found mass: 384 (m/z+H)

Example # 278

Preparation of 5-(2-fluorophenyl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL),

HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 486.3146 g/mol

HPLC-MS: analytical method A

rt: 2.00 min - found mass: 487 (m/z+H)

Example # 279

Preparation of N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-cyclopropylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and

purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 429.2434 g/mol

HPLC-MS: analytical method A

rt: 2.03 min - found mass: 430 (m/z+H)

Example # 280

Preparation of N-(3,4-dimethoxyphenyl)-5-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(1H-indazol-6-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 387.1664 g/mol

HPLC-MS: analytical method D

rt: 4.93 min - found mass: 388 (m/z+H)

Example # 281

Preparation of 5-(2,6-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,6-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 383.1377 g/mol

HPLC-MS: analytical method A

rt: 2.71 min - found mass: 384 (m/z+H)

Example # 282

Preparation of 5-(2,6-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,6-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 408.1747 g/mol

HPLC-MS: analytical method A

rt: 2.71 min - found mass: 409 (m/z+H)

Example # 283

Preparation of N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,6-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-cyclopropylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The

reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 447.2313 g/mol

HPLC-MS: analytical method A

rt: 2.07 min - found mass: 448 (m/z+H)

Example # 284

Preparation of 5-(2,6-difluorophenyl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,6-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 504.3025 g/mol

HPLC-MS: analytical method A

rt: 1.97 min - found mass: 505 (m/z+H)

Example # 285

Preparation of 5-(2,4-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 383.1376 g/mol

HPLC-MS: analytical method A

rt: 2.83 min - found mass: 384 (m/z+H)

Example # 286

Preparation of 5-(2,4-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 408.1747 g/mol

HPLC-MS: analytical method A

rt: 2.79 min - found mass: 409 (m/z+H)

Example # 287

Preparation of 6-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one (0.16 mmol) and 7-amino-3,4-dihydroquinolin-2(1H)-one (0.3 mmol 2 eq.,) were suspended

in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 427.1598 g/mol

HPLC-MS: analytical method D

rt: 4.82 min - found mass: 428 (m/z+H)

Example # 288

Preparation of 6-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The

reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 443.1992 g/mol

HPLC-MS: analytical method D

rt: 4.96 min - found mass: 444 (m/z+H)

Example # 289

Preparation of 6-(7-((2-methyl-1H-benzo[d]imidazol-5-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one (0.16 mmol) and 2-methyl-1H-benzo[d]imidazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 412.1584 g/mol

HPLC-MS: analytical method D

rt: 3.81 min - found mass: 413 (m/z+H)

Example # 290

Preparation of N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 474.2791 g/mol

HPLC-MS: analytical method L

rt: 3.08 min - found mass: 475 (m/z+H)

Example # 291

Preparation of N-(2-(methylsulfonyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-(methylsulfonyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 371.0663 g/mol

HPLC-MS: analytical method L

rt: 4.94 min - found mass: 372 (m/z+H)

Example # 292

Preparation of N-(3-((dimethylamino)methyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-((dimethylamino)methyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 350.1576 g/mol

HPLC-MS: analytical method L

rt: 3.33 min - found mass: 351 (m/z+H)

Example # 293

Preparation of N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-(4-cyclopropylpiperazin-1-yl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 417.2081 g/mol

HPLC-MS: analytical method L

rt: 3.40 min - found mass: 418 (m/z+H)

Example # 294

Preparation of morpholino(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-aminophenyl)(morpholino)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 406.1454 g/mol

HPLC-MS: analytical method L

rt: 4.29 min - found mass: 407 (m/z+H)

Example # 295

Preparation of N-(4-(morpholinomethyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-Amino-phenyl)-morpholin-4-yl-methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and dissolved in THF (dry) LiAlH₄ powder was added (excess, 2 by 2 eq) until completion of reaction is observed (by LCMS). The reaction was quenched with water (1mL per gram LiAlH₄), then NaOH (ca. 15% aq., 1mL per g LiAlH₄), water (3mL per gram LiAlH₄). The mixture was filtered, washed with THF, MeOH, MeCN (ca. 10ML each). The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 392.1710 g/mol

HPLC-MS: analytical method L

rt: 3.20 min - found mass: 393 (m/z+H)

Example # 296

Preparation of N-(3-fluoro-4-morpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-fluoro-4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 396.1393 g/mol

HPLC-MS: analytical method L

rt: 4.90 min - found mass: 397 (m/z+H)

Example # 297

Preparation of (4-(4-methylpiperazin-1-yl)piperidin-1-yl)(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and (4-aminophenyl)(4-(4-methylpiperazin-1-yl)piperidin-1-yl)methanone (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 502.2731 g/mol

HPLC-MS: analytical method L

rt: 3.25 min - found mass: 503 (m/z+H)

Example # 298

Preparation of 6-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one (0.16 mmol) and 1-(4-(4-

aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 484.2308 g/mol

HPLC-MS: analytical method A

rt: 2.22 min - found mass: 485 (m/z+H)

Example # 299

Preparation of 6-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was

irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 418.1621 g/mol

HPLC-MS: analytical method A

rt: 2.41 min - found mass: 419 (m/z+H)

Example # 300

Preparation of 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one (0.16 mmol) and 3,4-dimethoxyaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The reaction mixture was dissolved in THF (dry) LiAlH₄ powder was added (excess, 2 by 2 eq) until completion of reaction is observed (by LCMS). The reaction was quenched with water (1mL per gram LiAlH₄), then NaOH (ca. 15% aq., 1mL per g LiAlH₄), water (3mL per gram LiAlH₄). The mixture was filtered, washed with THF, MeOH, MeCN (ca. 10ML each). The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was

concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 404.1877 g/mol

HPLC-MS: analytical method L

rt: 3.84 min - found mass: 405 (m/z+H)

Example # 301

Preparation of 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The reaction mixture was dissolved in THF (dry) LiAlH₄ powder was added (excess, 2 by 2 eq) until completion of reaction is observed (by LCMS). The reaction was quenched with water (1mL per gram LiAlH₄), then NaOH (ca. 15% aq., 1mL per g LiAlH₄), water (3mL per gram LiAlH₄). The mixture was filtered, washed with THF, MeOH, MeCN (ca. 10ML each). The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 429.2248 g/mol

HPLC-MS: analytical method L

rt: 3.81 min - found mass: 430 (m/z+H)

Example # 302

Preparation of 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(2-methyl-1H-benzo[d]imidazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one (0.16 mmol) and 2-methyl-1H-benzo[d]imidazol-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The reaction mixture was dissolved in THF (dry) LiAlH₄ powder was added (excess, 2 by 2 eq) until completion of reaction is observed (by LCMS). The reaction was quenched with water (1mL per gram LiAlH₄), then NaOH (ca. 15% aq., 1mL per g LiAlH₄), water (3mL per gram LiAlH₄). The mixture was filtered, washed with THF, MeOH, MeCN (ca. 10ML each). The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 398.1840 g/mol

HPLC-MS: analytical method L

rt: 3.09 min - found mass: 399 (m/z+H)

Example # 303

Preparation of N-(2-methylisoindolin-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-methylisoindolin-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 348.1375 g/mol

HPLC-MS: analytical method L

rt: 3.03 min - found mass: 349 (m/z+H)

Example # 304

Preparation of N-(1,3-dihydroisobenzofuran-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 1,3-dihydroisobenzofuran-5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 335.0999 g/mol

HPLC-MS: analytical method L

rt: 4.61 min - found mass: 336 (m/z+H)

Example # 305

Preparation of N-(3-(tert-butyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-(tert-butyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry,

3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 349.1646 g/mol

HPLC-MS: analytical method L

rt: 5.75 min - found mass: 350 (m/z+H)

Example # 306

Preparation of methyl 3-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 1-(4-(4-aminophenyl)piperazin-1-yl)ethan-1-one (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and

purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 471.2382 g/mol

HPLC-MS: analytical method D

rt: 5.83 min - found mass: 482 (m/z+H)

Example # 307

Preparation of methyl 3-(7-((1H-indazol-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 1H-indazol-6-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 385.1464 g/mol

HPLC-MS: analytical method D

rt: 5.91 min - found mass: 386 (m/z+H)

Example # 308

Preparation of methyl 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate

methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate (0.16 mmol) and 4-morpholinoaniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 430.2067 g/mol

HPLC-MS: analytical method D

rt: 6.26 min - found mass: 431 (m/z+H)

Example # 309

Preparation of 5-(3-aminophenyl)-N-(4-((dimethylamino)methyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-Chloro-2-(4-methoxy-benzyl)-5-(3-nitro-phenyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-((dimethylamino)methyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The reaction mixture was dissolved in methanol (3mL). Pd/C was added (20mg) and the reaction was stirred overnight under N₂ at room temperature. After filtration with celite the reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 359.2176 g/mol

HPLC-MS: analytical method D

rt: 0.63 min - found mass: 360 (m/z+H)

Example # 310

Preparation of 5-(2-fluorophenyl)-N-(2-methylisoindolin-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 2-methylisoindolin-

5-amine (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 360.1729 g/mol

HPLC-MS: analytical method C

rt: 2.00 min - found mass: 409 (m/z+H)

Example # 311

Preparation of N-(4-((dimethylamino)methyl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 4-((dimethylamino)methyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and

purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 362.1930 g/mol

HPLC-MS: analytical method C

rt: 2.17 min - found mass: 361 (m/z+H)

Example # 312

Preparation of N-(3-((dimethylamino)methyl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine

7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine (0.16 mmol) and 3-((dimethylamino)methyl)aniline (0.3 mmol 2 eq.,) were suspended in MeOH (dry, 3mL) in a microwave vial (2-5mL), HCl in dioxane (4M, 3 drops) was added. The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was evaporated and used without further purification. The residue was dissolved in TFA (3mL). The reaction mixture was irradiated in a microwave reactor for 5 min at 140°C. The reaction mixture was concentrated and purified by semi-preparative HPLC-MS and freeze dried from water/t-BuOH 4/1.

exact mass: 362.1930 g/mol

HPLC-MS: analytical method C

rt: 2.18 min - found mass: 363 (m/z+H)

BIOLOGICAL DATA

SYK Activities:

Ki lower than 10nM:

Example #1; Example #2; Example #4; Example #5; Example #6;
Example #7; Example #10; Example #11; Example #16; Example
#20; Example #23; Example #25; Example #30; Example #31;
Example #33; Example #34; Example #35; Example #36; Example
#46; Example #54; Example #58; Example #59; Example #60;
Example #61; Example #62; Example #63; Example #64; Example
#66; Example #67; Example #68; Example #69; Example #70;
Example #71; Example #72; Example #73; Example #74; Example
#75; Example #76; Example #78; Example #79; Example #82;
Example #84; Example #87; Example #91; Example #95; Example
#98; Example #99; Example #117; Example #118; Example #119;
Example #128; Example #131; Example #132; Example #134;
Example #137; Example #138; Example #143; Example #144;
Example #149; Example #151; Example #160; Example #161;
Example #162; Example #163; Example #164; Example #169;
Example #170; Example #172; Example #175; Example #179;
Example #182; Example #183; Example #184; Example #186;
Example #187; Example #188; Example #189; Example #190;
Example #191; Example #192; Example #193; Example #194;
Example #195; Example #196; Example #197; Example #198;
Example #200; Example #201; Example #203; Example #204;
Example #205; Example #206; Example #207; Example #208;
Example #213; Example #225; Example #232; Example #233;
Example #234; Example #235; Example #237; Example #238;
Example #239; Example #240; Example #241; Example #243;
Example #244; Example #245; Example #246; Example #247;

Example #248; Example #251; Example #256; Example #257;
Example #269; Example #275; Example #278; Example #280;
Example #281; Example #285; Example #290; Example #293;
Example #295; Example #296; Example #297; Example #298;
Example #300; Example #302; Example #303; Example #304;

Ki between 10nM and 100nM:

Example #3; Example #8; Example #9; Example #12; Example #14;
Example #17; Example #18; Example #19; Example #22; Example
#24; Example #26; Example #27; Example #28; Example #29;
Example #32; Example #37; Example #38; Example #40; Example
#47; Example #50; Example #51; Example #52; Example #53;
Example #57; Example #65; Example #77; Example #80; Example
#81; Example #83; Example #85; Example #86; Example #89;
Example #90; Example #92; Example #93; Example #94; Example
#96; Example #97; Example #100; Example #101; Example #102;
Example #103; Example #104; Example #105; Example #106;
Example #107; Example #108; Example #109; Example #110;
Example #111; Example #112; Example #114; Example #116;
Example #120; Example #122; Example #123; Example #124;
Example #125; Example #126; Example #127; Example #129;
Example #135; Example #136; Example #139; Example #140;
Example #141; Example #142; Example #145; Example #146;
Example #147; Example #152; Example #156; Example #157;
Example #158; Example #159; Example #165; Example #166;
Example #167; Example #168; Example #171; Example #173;
Example #176; Example #177; Example #180; Example #181;
Example #185; Example #199; Example #202; Example #209;
Example #210; Example #211; Example #212; Example #216;
Example #218; Example #220; Example #221; Example #223;
Example #226; Example #227; Example #228; Example #229;
Example #230; Example #231; Example #236; Example #242;
Example #249; Example #252; Example #253; Example #254;

Example #258; Example #260; Example #268; Example #270;
Example #271; Example #273; Example #276; Example #279;
Example #282; Example #283; Example #284; Example #286;
Example #287; Example #288; Example #289; Example #291;
Example #294; Example #299; Example #301; Example #306;
Example #309; Example #310; Example #311;

Ki between 100nM and 1000nM:

Example #13; Example #21; Example #39; Example #45; Example
#48; Example #55; Example #56; Example #88; Example #113;
Example #115; Example #121; Example #130; Example #133;
Example #148; Example #150; Example #153; Example #174;
Example #178; Example #215; Example #217; Example #219;
Example #222; Example #224; Example #255; Example #259;
Example #262; Example #263; Example #264; Example #265;
Example #266; Example #267; Example #272; Example #274;
Example #277; Example #292; Example #305; Example #307;
Example #312;

LRRK2 Activites:

IC50 lower than 10nM:

Example #36; Example #82; Example #135; Example #144; Example
#247; Example #269;

IC50 between 10nM and 100nM:

Example #23; Example #26; Example #37; Example #46; Example
#59; Example #62; Example #64; Example #69; Example #70;
Example #71; Example #72; Example #73; Example #75; Example
#76; Example #77; Example #79; Example #84; Example #87;

Example #89; Example #91; Example #95; Example #117; Example #118; Example #136; Example #137; Example #139; Example #143; Example #145; Example #147; Example #158; Example #161; Example #164; Example #166; Example #182; Example #183; Example #184; Example #185; Example #190; Example #195; Example #199; Example #206; Example #235; Example #238; Example #280; Example #297;

IC50 between 100nM and 1000nM:

Example #1; Example #2; Example #4; Example #5; Example #6; Example #7; Example #9; Example #10; Example #14; Example #16; Example #17; Example #18; Example #20; Example #21; Example #22; Example #24; Example #25; Example #27; Example #28; Example #32; Example #33; Example #34; Example #35; Example #40; Example #41; Example #47; Example #48; Example #50; Example #51; Example #58; Example #60; Example #61; Example #63; Example #65; Example #66; Example #67; Example #68; Example #74; Example #78; Example #80; Example #81; Example #83; Example #85; Example #86; Example #88; Example #93; Example #96; Example #97; Example #99; Example #101; Example #108; Example #109; Example #110; Example #111; Example #114; Example #119; Example #120; Example #126; Example #127; Example #131; Example #132; Example #134; Example #138; Example #140; Example #142; Example #146; Example #156; Example #157; Example #159; Example #162; Example #163; Example #169; Example #170; Example #171; Example #172; Example #173; Example #179; Example #186; Example #187; Example #188; Example #189; Example #191; Example #193; Example #194; Example #196; Example #197; Example #198; Example #200; Example #201; Example #202; Example #203; Example #204; Example #205; Example #207; Example #208; Example #209; Example #218; Example #225; Example #232; Example #233; Example #234; Example #237; Example #239;

Example #240; Example #248; Example #251; Example #254;
Example #257; Example #261; Example #265; Example #268;
Example #271; Example #275; Example #278; Example #279;
Example #281; Example #282; Example #283; Example #284;
Example #290; Example #293; Example #294; Example #295;
Example #296; Example #300; Example #301; Example #302;
Example #303; Example #304; Example #310; Example #311;

MYLK Activites:


IC50 lower than 5000nM:

Example #5; Example #8; Example #20; Example #21; Example #34;
Example #35; Example #45; Example #46; Example #57; Example
#58; Example #62; Example #68; Example #70; Example #74;
Example #83; Example #91; Example #95; Example #97; Example
#99; Example #114; Example #117; Example #118; Example #120;
Example #127; Example #131; Example #132; Example #134;
Example #138; Example #139; Example #143; Example #144;
Example #147; Example #160; Example #161; Example #163;
Example #172; Example #175; Example #179; Example #182;
Example #187; Example #188; Example #189; Example #190;
Example #191; Example #225; Example #238; Example #240;
Example #241; Example #247; Example #248; Example #252;
Example #254; Example #255; Example #268; Example #275;
Example #276; Example #277; Example #279; Example #280;
Example #284; Example #286; Example #288; Example #289;
Example #290; Example #293; Example #294; Example #295;

Figure 1 shows the LAD2-SYK Correlation

CLAIMS

1. A compound of formula (I)



(I)

wherein

A is NH, O, S, C=O, NR⁴ or CR⁵R⁶;

R¹ is an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group;

R² is an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group, wherein R² is bound to the pyrimidine ring of formula (I) via a carbon-carbon bond;

R³ is a hydrogen atom, a halogen atom, NO₂, N₃, OH, SH, NH₂ or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group;

R⁴ is an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group;

R⁵ is a hydrogen atom, NO₂, N₃, OH, SH, NH₂ or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group; and

R⁶ is a hydrogen atom, NO₂, N₃, OH, SH, NH₂ or an alkyl, alkenyl, alkynyl, heteroalkyl, aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group;

or a pharmaceutically acceptable salt, ester, solvate or hydrate or a pharmaceutically acceptable formulation thereof.


2. A compound according to claim 1, wherein A is NH.
3. A compound according to claim 1 or claim 2 wherein R³ is H, F, Cl, CH₃, CF₃, NO₂, cyclopropyl, CN, N₃, OH, SH, OMe, SMe, NHMe, NMe₂ or NH₂.
4. A compound according to any one of the preceding claims, wherein R³ is a hydrogen atom.
5. A compound according to any one of the preceding claims, wherein A is NH and R³ is H.







6. A compound according to any one of the preceding claims, wherein R^1 is an optionally substituted aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aralkyl or heteroaralkyl group.
7. A compound according to any one of the preceding claims, wherein R^1 is an optionally substituted phenyl or naphthyl group or an optionally substituted heteroaryl group having one or two rings containing 5, 6, 7, 8, 9 or 10 ring atoms, or an optionally substituted arylheterocycloalkyl, heteroarylcycloalkyl or heteroaryl-heterocycloalkyl group containing two or three rings and 9 to 20 ring atoms.
8. A compound according to any one of the preceding claims, wherein R^1 is a group of formula $X^1-L^1-Y^1$ or a group of formula $X^1-L^1-Y^1-L^2-Z^1$ wherein
 X^1 is an optionally substituted phenyl group or an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N;
 L^1 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO-$, $-SO_2-$, $-NH-C(=O)-$, $-C(=O)-NH-$; $-C(=O)-O-$, $-O-C(=O)-$, $-NH-C(=O)-O-$, $-O-C(=O)-NH-$, $-NH-SO_2-NH-$, $-CH_2-NH-CH_2-$, $-NH-SO_2-$, $-SO_2-NH-$ or $-NH-C(=O)-NH-$;
 Y^1 is an optionally substituted phenyl group, an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N, an optionally substituted C_3-C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N;






L^2 is a bond or a group of formula $-\text{CH}_2-$, $-\text{C}(=\text{O})-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NH}-\text{C}(=\text{O})-$, $-\text{C}(=\text{O})-\text{NH}-$, $-\text{C}(=\text{O})-\text{O}-$, $-\text{O}-\text{C}(=\text{O})-$, $-\text{NH}-\text{C}(=\text{O})-\text{O}-$, $-\text{O}-\text{C}(=\text{O})-\text{NH}-$, $-\text{NH}-\text{SO}_2-\text{NH}-$, $-\text{CH}_2-\text{NH}-\text{CH}_2-$, $-\text{NH}-\text{SO}_2-$, $-\text{SO}_2-\text{NH}-$ or $-\text{NH}-\text{C}(=\text{O})-\text{NH}-$; and






Z^1 is an optionally substituted phenyl group, an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N, an optionally substituted C_3 - C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N.

9. A compound according to any one of the preceding claims 1 to 5, wherein R^1 is selected from the following groups:











10. A compound according to any one of the preceding claims, wherein R^2 is an optionally substituted aryl, heteroaryl, cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aralkyl or heteroaralkyl group.
11. A compound according to any one of the preceding claims, wherein R^2 is an optionally substituted phenyl or naphthyl group or an optionally substituted heteroaryl group having one or two rings containing 5, 6, 7, 8, 9 or 10 ring atoms, or an optionally substituted arylheterocycloalkyl or heteroarylheterocycloalkyl group containing two or three rings and 9 to 20 ring atoms.
12. A compound according to any one of the preceding claims, wherein R^2 is a group of formula $X^2-L^3-Y^2$ or a group of formula $X^2-L^3-Y^2-L^4-Z^2$ wherein
 X^2 is an optionally substituted phenyl group or an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N;
 L^3 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO-$, $-SO_2-$, $-NH-C(=O)-$, $-C(=O)-NH-$; $-C(=O)-O-$, $-O-C(=O)-$, $-NH-C(=O)-O-$, $-O-C(=O)-NH-$, $-NH-SO_2-NH-$, $-CH_2-NH-CH_2-$, $-NH-SO_2-$, $-SO_2-NH-$ or $-NH-C(=O)-NH-$;
 Y^2 is an optionally substituted phenyl group, an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N, an optionally substituted C_3-C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N;
 L^4 is a bond or a group of formula $-CH_2-$, $-C(=O)-$, $-SO-$, $-SO_2-$, $-NH-C(=O)-$, $-C(=O)-NH-$; $-C(=O)-O-$, $-O-C(=O)-$, $-NH-$


$C(=O)-O-$, $-O-C(=O)-NH-$, $-NH-SO_2-NH-$, $-CH_2-NH-CH_2-$, $-NH-SO_2-$, $-SO_2-NH-$ or $-NH-C(=O)-NH-$; and

Z^2 is an optionally substituted phenyl group, an optionally substituted heteroaryl group containing 5 or 6 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N, an optionally substituted C_3-C_7 cycloalkyl group or an optionally substituted heterocycloalkyl group containing 3, 4, 5, 6 or 7 ring atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and N.

13. A compound according to any one of the preceding claims 1 to 10, wherein R^2 is selected from the following groups:







14. A compound according to claim 1 exemplified as follows:
 N-(1H-indazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine;
 N-(1H-benzo[d][1,2,3]triazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine;
 5-phenyl-N-(4-(piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine;
 N1,N1-dimethyl-N4-(5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine;
 3-((5-phenyl-1H-

pyrazolo[4,3-d]pyrimidin-7-yl) amino) benzamide; N-(3,4-dimethoxyphenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-methoxyphenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) benzoic acid; 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) benzamide; 4-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) benzonitrile; N-(1H-indazol-5-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-5-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 2-((5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) benzamide; N-(1H-indazol-6-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) benzene-1,4-diamine; N-(3,4-dimethoxyphenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; 5-(3,4-dimethoxyphenyl)-N-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1-(5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine; N,5-bis(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; N-(1H-indazol-6-yl)-5-

(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; N-(4-chlorophenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-6-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; N-(3,4-dimethoxyphenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-6-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; 5-phenyl-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-phenyl-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(1H-indazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dimethoxyphenyl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-pyrazol-3-yl)-5-(pyridin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(pyridin-4-yl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N,5-di(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one;

N-(1H-pyrazol-3-yl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(pyrrolidin-1-yl)phenyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-5-yl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(1-methylpiperidin-4-yl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; 5-(1-methylpiperidin-4-yl)-N-(1H-pyrazol-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(1-methylpiperidin-4-yl)-N-(4-(pyrrolidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(4-methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1-(5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine; N-(5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 3-((5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; 5-(4-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1H-indazol-6-yl)-5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-

d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; N-(3,4-dimethoxyphenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N4-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,4-diamine; N-(1H-indazol-6-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1-(5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-N4,N4-dimethylbenzene-1,4-diamine; N-(1H-indazol-6-yl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 3-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; 3-((5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; N-(1H-indazol-6-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; N-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 3-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-(3-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N-(5-

cyclopropyl-1H-pyrazol-3-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 3-((5-(1-methylpiperidin-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenol; 5-(benzo[d]thiazol-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(benzo[d]thiazol-2-yl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-(benzo[d]thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; 5-(benzo[d]thiazol-2-yl)-N-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N-(1H-indazol-6-yl)-5-(3-(pyridin-3-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 2-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-methoxyphenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-

pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-amine; N-(1H-indazol-6-yl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-methoxyphenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(5-cyclopropyl-1H-pyrazol-3-yl)-5-(2-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N1,N1-dimethyl-N3-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,3-diamine; N1-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)cyclohexane-1,2-diamine; N1-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)benzene-1,3-diamine; N-(6-(4-methylpiperazin-1-yl)pyridin-3-yl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-ethylpiperazin-1-yl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-((dimethylamino)methyl)phenyl)-5-phenyl-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,5-dimethoxyphenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,5-dimethoxyphenyl)-N-(4-(4-ethylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2,5-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-aminoethyl)-3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzamide; N-(3,4-dimethoxyphenyl)-5-(imidazo[1,2-a]pyridin-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-ethylpiperazin-1-yl)phenyl)-5-(imidazo[1,2-a]pyridin-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(imidazo[1,2-a]pyridin-2-yl)-N-(1H-indazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(imidazo[1,2-a]pyridin-2-yl)-N-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 4-((5-(imidazo[1,2-a]pyridin-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-

7-yl)amino)benzoic acid; 5-(1H-benzo[d]imidazol-2-yl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-methoxyphenyl)-N-(4-((4-methylpiperazin-1-yl)methyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-methoxyphenyl)-N-(4-(pyrrolidin-1-ylmethyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)-3-(m-tolyl)urea; (4-methylpiperazin-1-yl)(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; pyrrolidin-1-yl(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; N-(4-thiomorpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 4-(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)thiomorpholine 1,1-dioxide; 1-(4-(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; (3-methoxy-4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(4-(4-methylpiperazin-1-yl)piperidin-1-yl)methanone; 5-(1H-imidazol-5-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(1H-imidazol-2-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; pyrrolidin-1-yl(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; 4-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)thiomorpholine 1,1-dioxide; 1-(4-(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; N-(4-thiomorpholinophenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(thiophen-3-yl)-1H-

pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; (4-methylpiperazin-1-yl)(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; 2,2-dimethyl-N-(5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-amine; 5-(3-(2H-tetrazol-5-yl)phenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; (7-((2-(dimethylamino)ethyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; (7-(((1-methylpiperidin-4-yl)methyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; (7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; (7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; (4-((5-benzoyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(pyrrolidin-1-yl)methanone; 7-((5-benzoyl-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; morpholino(4-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; 6-((5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]thiazin-3(4H)-one; 7-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; (4-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)(pyrrolidin-1-yl)methanone; N-(4-(4-(tert-butyl)piperazin-1-yl)phenyl)-5-(thiophen-3-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-2H-benzo[b][1,4]oxazin-3(4H)-one; N-(4-(4-(tert-butyl)piperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 8-((5-(4-

methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-
4,5-dihydro-1H-benzo[b]azepin-2(3H)-one; 6-((5-(thiophen-
2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-2H-
benzo[b][1,4]thiazin-3(4H)-one; 1-methyl-N-(5-(thiophen-2-
yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-1,2,3,4-
tetrahydroquinolin-7-amine; 2-methyl-2-(4-((5-(thiophen-2-
yl)-1H-pyrazolo[4,3-d]pyrimidin-7-
yl) amino)phenyl)propanenitrile; 5-(4-methoxyphenyl)-N-(4-
thiomorpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine;
4-(4-((5-(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-
yl) amino)phenyl)thiomorpholine 1,1-dioxide; 1-(4-(4-((5-
(4-methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-
yl) amino)phenyl)piperazin-1-yl)ethan-1-one; 6-((5-(4-
methoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-2H-
benzo[b][1,4]oxazin-3(4H)-one; N-(4-(4-(tert-
butyl)piperazin-1-yl)phenyl)-5-(4-methoxyphenyl)-1H-
pyrazolo[4,3-d]pyrimidin-7-amine; 2,2-dimethyl-N-(5-
(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)-3,4-
dihydro-2H-benzo[b][1,4]thiazin-6-amine; 2-(3-(7-((4-(4-
methylpiperazin-1-yl)phenyl) amino)-1H-pyrazolo[4,3-
d]pyrimidin-5-yl)phenoxy)acetic acid; 5-(7-((4-(4-
methylpiperazin-1-yl)phenyl) amino)-1H-pyrazolo[4,3-
d]pyrimidin-5-yl)thiophene-3-carboxylic acid; 2-(3-(7-((4-
(4-methylpiperazin-1-yl)phenyl) amino)-1H-pyrazolo[4,3-
d]pyrimidin-5-yl)phenyl)acetic acid; 6-(7-((4-(4-
methylpiperazin-1-yl)phenyl) amino)-1H-pyrazolo[4,3-
d]pyrimidin-5-yl)picolinic acid; N-(4-
(pentafluorosulfanyl)phenyl)-5-(thiophen-3-yl)-1H-
pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-
1-yl)phenyl)-5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-
7-amine; 5-(3-(((4-methylbenzyl) amino)methyl)phenyl)-N-(4-
(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-
d]pyrimidin-7-amine; 5-(3-(3-

(dimethylamino)propoxy)phenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 3-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 1-(4-(4-((5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; N-(3,4-dimethoxyphenyl)-5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(thiazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; N-(3-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; N-(3-(7-((4-(piperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; N-(3-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; N-(3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; N-(3-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; 1-(4-(4-((5-(3-(2H-tetrazol-5-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; 5-(3-(2H-tetrazol-5-yl)phenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 3-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; 7-((5-(3-(2H-tetrazol-5-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-(4-((5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-

yl) amino) phenyl) piperazin-1-yl) ethan-1-one; N-(3,4-dimethoxyphenyl)-5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-(4-((5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) phenyl) piperazin-1-yl) ethan-1-one; N-(3,4-dimethoxyphenyl)-5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-(4-((5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino) phenyl) piperazin-1-yl) ethan-1-one; N-(3,4-dimethoxyphenyl)-5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(thiazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-3,4-dihydroquinolin-2(1H)-one; 7-((5-(oxazol-4-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-3,4-dihydroquinolin-2(1H)-one; 7-((5-(oxazol-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-3,4-dihydroquinolin-2(1H)-one; N-(4-(1-methylpiperidin-4-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; methyl 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; methyl 4-(7-((4-(4-methylpiperazin-1-yl)phenyl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-((3,4-dimethoxyphenyl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((3-oxo-3,4-dihydro-2H-benzo[b][1,4]thiazin-6-yl) amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((2-

oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-8-yl) amino) -1H-pyrazolo[4,3-d]pyrimidin-5-yl) benzoate; 3-(7-((2-oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-8-yl) amino) -1H-pyrazolo[4,3-d]pyrimidin-5-yl) benzoic acid; N-(3,4-dimethoxyphenyl)-5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-(pentafluorosulfanyl)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(3-(pentafluorosulfanyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl) amino)-3,4-dihydroquinolin-2(1H)-one; N-(3,4-dimethoxyphenyl)-5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-(methylthio)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-methylpiperazin-1-yl)phenyl)-5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(3-(trifluoromethoxy)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-(trifluoromethyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-morpholinophenyl)-5-(3-(trifluoromethyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-chlorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-chlorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-

((5-(3-(methylthio)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; 3-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 3-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 3-(7-((3-methoxy-4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; 4-(7-((4-(4-methylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; methyl 4-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 4-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; 4-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid; N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 1-(4-(4-((5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; 5-(1H-indazol-6-yl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(1H-indazol-6-yl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; 5-(3-(methylsulfonyl)phenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N,5-bis(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-(methylsulfonyl)phenyl)-5-(thiophen-2-yl)-1H-

pyrazolo[4,3-d]pyrimidin-7-amine; 2-((3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)sulfonyl)ethan-1-ol; 1-(4-(4-((5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)piperazin-1-yl)ethan-1-one; 7-((5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; N-(4-morpholinophenyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-methyl-1H-benzo[d]imidazol-5-yl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N,N-dimethyl-4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide; N-cyclopropyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide; N-(3-(2H-1,2,3-triazol-2-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(2H-1,2,3-triazol-2-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N,N-dimethyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide; N-methyl-3-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)benzenesulfonamide; N-(3-(morpholinosulfonyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(thiophen-2-yl)-N-(3-((trifluoromethyl)sulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-(methylsulfinyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 7-((5-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)-3,4-dihydroquinolin-2(1H)-one; N-(3,4-dimethoxyphenyl)-5-(4-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(4-fluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-

pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,5-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(4-(4-methylpiperazin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(3-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(2-(methylsulfonyl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3,4-dimethoxyphenyl)-5-(1H-indazol-6-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,6-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,6-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(2,6-difluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,6-difluorophenyl)-N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,4-difluorophenyl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(2,4-difluorophenyl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 6-(7-((2-oxo-1,2,3,4-tetrahydroquinolin-7-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 6-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 6-(7-((2-methyl-1H-benzo[d]imidazol-5-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; N-(4-(4-(4-methylpiperazin-1-yl)piperidin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-

(methylsulfonyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-((dimethylamino)methyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-(4-cyclopropylpiperazin-1-yl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; morpholino(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; N-(4-(morpholinomethyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-fluoro-4-morpholinophenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; (4-(4-methylpiperazin-1-yl)piperidin-1-yl)(4-((5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-yl)amino)phenyl)methanone; 6-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 6-(7-((3,4-dimethoxyphenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(3,4-dimethoxyphenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(4-morpholinophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; 5-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-N-(2-methyl-1H-benzo[d]imidazol-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(2-methylisoindolin-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(1,3-dihydroisobenzofuran-5-yl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-(tert-butyl)phenyl)-5-(thiophen-2-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; methyl 3-(7-((4-(4-acetylpiperazin-1-yl)phenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((1H-indazol-6-yl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 3-(7-((4-morpholinophenyl)amino)-1H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 5-(3-aminophenyl)-N-(4-((dimethylamino)methyl)phenyl)-1H-pyrazolo[4,3-

d]pyrimidin-7-amine; 5-(2-fluorophenyl)-N-(2-methylisoindolin-5-yl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(4-((dimethylamino)methyl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine; N-(3-((dimethylamino)methyl)phenyl)-5-(2-fluorophenyl)-1H-pyrazolo[4,3-d]pyrimidin-7-amine.


15. A pharmaceutical composition comprising a compound according to anyone of the preceding claims or a pharmaceutically acceptable ester, prodrug, hydrate, solvate or salt thereof, optionally in combination with a pharmaceutically acceptable carrier.
16. Use of a compound or a pharmaceutical composition according to anyone of the preceding claims for the preparation of a medicament for the treatment of respiratory tract/obstructive airways diseases and disorders, systemic anaphylaxis or hypersensitivity responses, drug allergies, bone and joint related diseases and disorders, skin and eye related diseases and disorders, gastrointestinal tract and abdominal related diseases and disorders, hematological disorders, metabolic disorders, cerebellar dysfunction, disturbances of brain metabolism, transplant rejection related conditions, genitourinary related conditions, CNS related diseases and disorders like parkinson's disease, inflammatory or immunological diseases or disorders, cardiovascular diseases and disorders, oncological diseases and disorders.
17. A compound or a pharmaceutical composition according to anyone of claims 1 to 15 for use in the treatment of respiratory tract/obstructive airways diseases and

disorders, systemic anaphylaxis or hypersensitivity responses, drug allergies, bone and joint related diseases and disorders, skin and eye related diseases and disorders, gastrointestinal tract and abdominal related diseases and disorders, hematological disorders, metabolic disorders, cerebellar dysfunction, disturbances of brain metabolism, transplant rejection related conditions, genitourinary related conditions, CNS related diseases and disorders like parkinson's disease, inflammatory or immunological diseases or disorders, cardiovascular diseases and disorders, oncological diseases and disorders.


18. A method of treatment of a disease mediated by kinase-activity comprising the administration, to a patient in need thereof, of a therapeutically effective amount of a compound according to any one of claims 1 to 14 or a pharmaceutical composition according to claim 15.
19. The method according to claim 18, wherein the kinase is selected from SYK (Spleen Tyrosine Kinase), LRRK2 (Leucine-rich repeat kinase 2) and/or Myosin light chain kinase (MYLK or MLCK) or mutants thereof.
20. The method according to claim 19, wherein said treatment is systemic.
21. The method according to claim 19, wherein said administration is topical.
22. The method according to claim 19, wherein said disease is selected from the group consisting of an inflammatory

disease, an autoimmune disease, an allergic disorder, and an ocular disorder.


23. The method according to claim 18 or 19, wherein the disease is selected from the group consisting of pruritus, eczema, asthma, rhinitis, dry eye, ocular inflammation, allergic conjunctivitis, vernal conjunctivitis, vernal keratoconjunctivitis, giant papillary conjunctivitis, fungal keratitis and uveitis.
24. The method according to claim 21, wherein said topical administration is to the skin.
25. The method according to claim 21, wherein said topical administration is to the eye.
26. The method according to claim 21, wherein said topical administration is intranasal or by inhalation.
27. A compound of formula (II), (IIIa) or (IIIb) wherein R^2 and R^3 are defined as in anyone of the preceding claims 1 to 13 and PG is a protecting group:



(II)



(IIIa)



(IIIb)

28. A compound according to claim 27 wherein PG is a 4-methoxybenzyl group or a carboxybenzyl group.
29. A compound according to claim 27 which is selected from the following compounds:
- 7-chloro-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-dimethoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-pyrazol-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-

chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole;
7-chloro-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol; 4-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)morpholine; 7-chloro-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,5-dimethoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; N-(2-aminoethyl)-3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzamide; 7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(1H-benzo[d]imidazol-2-yl)-7-chloro-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-5-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2H-pyrazolo[4,3-d]pyrimidine; (7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; 3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester; 2-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)acetic acid methyl ester; 5-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiophene-3-carboxylic acid methyl ester; 2-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)acetic acid methyl ester; 6-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)picolinic acid methyl ester; 2-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; N-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzyl)-1-(p-tolyl)methanamine; 3-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)-N,N-dimethylpropan-1-amine; 3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; N-(3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; 4-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; 4-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; 2-(7-chloro-2H-

pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; methyl 3-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-(trifluoromethoxy)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-(trifluoromethyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-chlorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2-fluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-fluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 4-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester; 7-chloro-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-indazol-6-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(4-fluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-difluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,5-difluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,6-difluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,4-difluorophenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-Chloro-5-(3-nitro-phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 6-(7-chloro-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one; 7-chloro-2-(4-methoxybenzyl)-5-phenyl-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(pyridin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(6-(pyrrolidin-1-yl)pyridin-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1-methylpiperidin-4-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-

chloro-2-(4-methoxybenzyl)-5-(1H-pyrazol-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(4-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(thiophen-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(thiophen-3-yl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzo[d]thiazole; 7-chloro-2-(4-methoxybenzyl)-5-(3-(pyridin-3-yl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenol; 4-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)morpholine; 7-chloro-2-(4-methoxybenzyl)-5-(2-methoxyphenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,5-dimethoxyphenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; N-(2-aminoethyl)-3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzamide; 7-chloro-5-(imidazo[1,2-a]pyridin-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(1H-benzo[d]imidazol-2-yl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1-methyl-1H-benzo[d]imidazol-2-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-5-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-imidazol-2-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 5-(3-(2H-tetrazol-5-yl)phenyl)-7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; (7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methanone; 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester; 2-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-

yl)phenoxy)acetic acid methyl ester; 5-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiophene-3-carboxylic acid methyl ester; 2-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)acetic acid methyl ester; 6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)picolinic acid methyl ester; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzyl)-1-(p-tolyl)methanamine; 3-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenoxy)-N,N-dimethylpropan-1-amine; 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzenesulfonamide; N-(3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)phenyl)methanesulfonamide; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)thiazole; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; 2-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)oxazole; methyl 3-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; methyl 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoate; 7-chloro-5-(3-(pentafluorosulfanyl)phenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-(methylthio)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethoxy)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(3-(trifluoromethyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-chlorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3-fluorophenyl)-2-(4-

methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 4-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)benzoic acid methyl ester; 7-chloro-2-(4-methoxybenzyl)-5-(3-(methylsulfonyl)phenyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(1H-indazol-6-yl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-2-(4-methoxybenzyl)-5-(1H-pyrrolo[3,2-b]pyridin-6-yl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(4-fluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(3,5-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-chloro-5-(2,6-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 7-Chloro-2-(4-methoxybenzyl)-5-(3-nitrophenyl)-2H-pyrazolo[4,3-d]pyrimidin; 7-chloro-5-(2,4-difluorophenyl)-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidine; 6-(7-chloro-2-(4-methoxybenzyl)-2H-pyrazolo[4,3-d]pyrimidin-5-yl)-2H-benzo[b][1,4]oxazin-3(4H)-one.




Fig. 1

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/001737

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D487/04 C07D519/00 A61K31/519 A61P29/00 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98/29413 A1 (YUAN JUN [US]) 9 July 1998 (1998-07-09) example 2A -----	1,2
X	EP 1 348 707 A1 (USTAV EX BOTAN AV CR I OF EX B [CZ] USTAV EX BOTAN AV CR V V I I O [CZ] 1 October 2003 (2003-10-01) example 7 -----	1,2,6
X	WO 02/074774 A1 (PFIZER LTD [GB]; DENTON STEPHEN MARTIN [GB]; KEMP MARK IAN [GB]; NEWMA) 26 September 2002 (2002-09-26) * preparations 71, 72 * -----	1,10,11, 27
X	US 6 531 475 B1 (HADDACH MUSTAPHA [US] ET AL) 11 March 2003 (2003-03-11) column 54; compounds 2', 3', 4', 5' ----- -/-	1,2,27
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search	Date of mailing of the international search report	
26 June 2012	02/07/2012	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Johnson, Claire	

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/001737

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JUN YUAN: "3-Aryl pyrazolo[4,3-d]pyrimidine derivatives: non-peptide CRF-1 antagonists", BIOORGANIC MEDICINAL CHEMISTRY LETTERS, vol. 12, 1 January 2002 (2002-01-01), pages 2133-2136, XP55030588, compound 16a -----	1,2, 15-17
A	EP 2 123 654 A1 (TAISHO PHARMA CO LTD [JP]; NISSAN CHEMICAL IND LTD [JP]) 25 November 2009 (2009-11-25) claims 1,12 -----	1-29
A	JEREMY NICHOLS R ET AL: "Substrate specificity and inhibitors of LRRK2, a protein kinase mutated in Parkinson's disease", BIOCHEMICAL JOURNAL, THE BIOCHEMICAL SOCIETY, LONDON, GB, vol. 424, 9 September 2009 (2009-09-09), pages 47-60, XP002654035, ISSN: 0264-6021, DOI: 10.1042/BJ20091035 figure 4 -----	1-29

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/001737

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9829413	A1	09-07-1998	AT 214701 T 15-04-2002
		AU 738324 B2 13-09-2001	
		AU 5811098 A 31-07-1998	
		BR 9713808 A 25-01-2000	
		CA 2276610 A1 09-07-1998	
		CN 1246860 A 08-03-2000	
		DE 69711241 D1 25-04-2002	
		DE 69711241 T2 12-12-2002	
		EP 0960110 A1 01-12-1999	
		ES 2172830 T3 01-10-2002	
		HK 1025958 A1 09-05-2003	
		JP 2001507705 A 12-06-2001	
		NO 993240 A 27-08-1999	
		NZ 336529 A 30-03-2001	
		US 5723608 A 03-03-1998	
		US 6211187 B1 03-04-2001	
		WO 9829413 A1 09-07-1998	
EP 1348707	A1	01-10-2003	AT 478872 T 15-09-2010
		AU 2003226731 A1 13-10-2003	
		CA 2480409 A1 09-10-2003	
		DK 1348707 T3 13-12-2010	
		EP 1348707 A1 01-10-2003	
		JP 2005527565 A 15-09-2005	
		US 2005080097 A1 14-04-2005	
		WO 03082872 A1 09-10-2003	
WO 02074774	A1	26-09-2002	AT 283856 T 15-12-2004
		BR 0208154 A 02-03-2004	
		CA 2439528 A1 26-09-2002	
		DE 60202134 D1 05-01-2005	
		DE 60202134 T2 24-11-2005	
		EP 1368352 A1 10-12-2003	
		ES 2231667 T3 16-05-2005	
		JP 2004523585 A 05-08-2004	
		MX PA03005937 A 10-09-2003	
		WO 02074774 A1 26-09-2002	
US 6531475	B1	11-03-2003	AU 6465701 A 26-11-2001
		US 6531475 B1 11-03-2003	
		WO 0187885 A1 22-11-2001	
EP 2123654	A1	25-11-2009	CA 2673965 A1 10-07-2008
		CN 101611035 A 23-12-2009	
		EP 2123654 A1 25-11-2009	
		US 2010298557 A1 25-11-2010	
		WO 2008081928 A1 10-07-2008	