



(51) International Patent Classification:

C07D 487/04 (2006.01) *A61P 1/16* (2006.01)
A61K 31/407 (2006.01) *A61P 25/00* (2006.01)
A61P 13/12 (2006.01) *A61P 37/06* (2006.01)

(21) International Application Number:

PCT/EP2013/069679

(22) International Filing Date:

23 September 2013 (23.09.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12185941.7 25 September 2012 (25.09.2012) EP

(71) *Applicant (for all designated States except SA, US):* F. **HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).

(71) *Applicant (for US only):* **HOFFMANN-LA ROCHE INC.** [US/US]; 340 Kingsland Street, Nutley, New Jersey 07110 (US).

(72) *Inventors:* **HERT, Jérôme**; Roettelerstrasse 15, CH-4058 Basel (CH). **HUNZIKER, Daniel**; Buerklistrasse 14, CH-4313 Mochlin (CH). **MATTEI, Patrizio**; Doernliweg 33, CH-4125 Riehen (CH). **MAUSER, Harald**; Inzlingerstrasse 1, CH-4125 Riehen (CH). **TANG, Guozhi**; Unit 34, No. 528 East Pailou Road, Pudong New District, Shanghai, 201314 (CN). **WANG, Lisha**; Thannerstrasse 80, 1st floor, CH-4054 Basel (CH).


(74) *Agent:* **SALUD, Carlos**; Grenzacherstrasse 124, CH-4070 Basel (CH).

(81) *Designated States (unless otherwise indicated, for every kind of national protection available):* AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) *Designated States (unless otherwise indicated, for every kind of regional protection available):* ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:— *of inventorship (Rule 4.17(iv))***Published:**— *with international search report (Art. 21(3))*

(54) Title: NEW BICYCLIC DERIVATIVES




(57) **Abstract:** The invention provides novel compounds having the general formula (I) wherein R^1 , Y, A, W, R^2 , m, n, p and q are as described herein, compositions including the compounds and methods of using the compounds.

New bicyclic derivatives

The present invention relates to organic compounds useful for therapy or prophylaxis in a mammal, and in particular to autotaxin (ATX) inhibitors which are inhibitors of lysophosphatidic acid (LPA) production and thus modulators of LPA levels and associated signaling, for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, conditions of the respiratory system, vascular and cardiovascular conditions, fibrotic diseases, cancer, ocular conditions, metabolic conditions, cholestatic and other forms of chronic pruritus and acute and chronic organ transplant rejection.

The present invention provides novel compounds of formula (I)





wherein

R^1 is alkyl, haloalkyl, substituted cycloalkyl, substituted cycloalkylalkyl, substituted piperazinyl, substituted piperidinyl, substituted indanyloxyalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted phenylalkynyl, substituted pyridinyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted pyridinylalkynyl, substituted thiophenyl, substituted thiophenylalkyl, substituted thiophenylalkenyl, substituted thiophenylalkynyl, naphthyl, substituted naphthyl, quinolyl, substituted quinolynyl, isoquinolyl, substituted isoquinolynyl, substituted 2,3-dihydro-1H-isoindol-2-yl, substituted 1H-indol-2-yl or substituted benzofuran-2-yl wherein substituted cycloalkyl, substituted cycloalkylalkyl, substituted piperazinyl, substituted piperidinyl, substituted indanyloxyalkyl, substituted phenyl, substituted phenylalkyl, substituted phenylalkynyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted pyridinylalkynyl, substituted thiophenyl, substituted thiophenylalkyl, substituted thiophenylalkenyl, substituted thiophenylalkynyl,

-2-

substituted naphthyl, substituted quinoliny, substituted isoquinoliny, substituted 2,3-dihydro-1H-isoindol-2-yl, substituted 1H-indol-2-yl and substituted benzofuran-2-yl are substituted with R^8 , R^9 and R^{10} ;

Y is $-OC(O)-$, $-NR^7C(O)-$, $-C(O)-$, $-S(O)_2-$,  or  ;

5 A is $-N-$ or CR^5- ;

W is $-O-$, $-S-$, $-NR^6-$, $-C(O)-$, $-S(O)_2-$, $-C(O)-NR^6-$ or $-CR^3R^4-$;


R^3 and R^4 are independently selected from H, halogen, alkyl and cycloalkyl;

R^5 , R^6 and R^7 are independently selected from H, alkyl and cycloalkyl;


10 R^8 , R^9 and R^{10} are independently selected from H, alkyl, hydroxyalkyl, haloalkyl, hydroxyhaloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, alkoxyhaloalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, phenyl, substituted phenyl, pyridinyl, substituted pyridinyl, pyrrolyl, substituted pyrrolyl, pyrrolydinyl, substituted pyrrolydinyl, tetrahydrofuranyl, substituted tetrahydrofuranyl, halogen, hydroxy, 15 cyano, alkylsulfanyl, haloalkylsulfanyl, cycloalkylsulfanyl, alkylsulfinyl, haloalkylsulfinyl, cycloalkylsulfinyl, alkylcarbonyl, haloalkylcarbonyl, cycloalkylcarbonyl, alkylsulfonyl, haloalkylsulfonyl, cycloalkylsulfonyl, substituted aminosulfonyl, substituted amino and substituted aminoalkyl, wherein substituted aminosulfonyl, substituted amino and substituted aminoalkyl are substituted on the 20 nitrogen atom with one to two substituents independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, alkylcarbonyl and cycloalkylcarbonyl, and wherein substituted phenyl, substituted pyrrolyl, substituted pyrrolydinyl, substituted tetrahydrofuranyl, and substituted pyridinyl are substituted with one to three substituents independently selected from alkyl, halogen, haloalkyl, 25 alkoxy and haloalkoxy;

m, n, p and q are independently selected from 1 or 2;


-3-

**B**


,

**C**


,

**D**


,

**E**


,

**F**


,

**G**


,

**H**


,

**I**


,

**J**


,

**K**


,

**L**


,

**M**


,

**N**


,

**O**


,

**P**


,

**Q**

,


**R**

,


**S**

,


-4-

**T**


,

**U**


,

**V**


,

**X**


,

**Z**


,

**AA**


,

**AB**


,

**AC**


,

**AD**


,

**AE**


,

**AF**


,

**AG**


,

**AH**

,

**AI**

,

**AJ**

;

R¹¹ is H, alkyl, haloalkyl or cycloalkyl;

R¹² is alkyl, halogen, haloalkyl and alkoxy;

R² is selected from the ring systems B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, X, Z, AA, AB, AC, AD, AE, AF, AG, AH, AI and AJ;

and pharmaceutically acceptable salts.

Autotaxin (ATX) is a secreted enzyme also called ectonucleotide pyrophosphatase / phosphodiesterase 2 or lysophospholipase D that is important for converting lysophosphatidyl choline (LPC) to the bioactive signaling molecule lysophosphatidic acid (LPA). It has been shown that plasma LPA levels are well correlated with ATX activity and hence ATX is believed to be an important source of extracellular LPA. Early experiments with a prototype ATX inhibitor have shown that such a compound is able to inhibit the LPA synthesizing activity in mouse plasma. Work conducted in the 1970s and early 1980s has demonstrated that LPA can elicit a wide range of cellular responses; including smooth muscle cell contraction, platelet activation, cell proliferation, chemotaxis and others. LPA mediates its effects via signaling to several G protein coupled receptors (GPCRs); the first members were originally denoted Edg (endothelial cell differentiation gene) receptors or ventricular zone gene-1(vzg-1) but are now called LPA receptors. The prototypic group now consists of LPA1/Edg-2/VZG-1, LPA2/Edg-4, and LPA3/Edg-7. Recently, three additional LPA receptors LPA4/p2y9/GPR23, LPA5/GPR92 and LPA6/p2Y5 have been described that are more closely related to nucleotide-selective purinergic receptors than to the prototypic LPA1-3 receptors. The ATX-LPA signaling axis is involved in a large range of physiological and pathophysiological functions, including, for example, nervous system function, vascular development, cardiovascular physiology, reproduction, immune system function, chronic inflammation, tumor metastasis and progression, organ fibrosis as well as obesity and/or other metabolic diseases such as diabetes mellitus. Therefore, increased activity of ATX and/or increased levels of LPA, altered LPA receptor expression and altered responses to LPA may contribute to the initiation, progression and/or outcome of a number of different pathophysiological conditions related to the ATX/LPA axis.

In accordance with the invention, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis of diseases, disorders or conditions that are associated with the activity of autotaxin and/or the biological activity of lysophosphatidic acid (LPA).

The compounds of formula (I) or their pharmaceutically acceptable salts and esters herein inhibit autotaxin activity and therefore inhibit LPA production and modulate LPA levels and associated signaling. Autotaxin inhibitors described herein are useful as agents for the treatment or prevention of diseases or conditions in which ATX activity and/or LPA signaling participates, is involved in the etiology or pathology of the disease, or is otherwise associated with at least

one symptom of the disease. The ATX-LPA axis has been implicated for example in angiogenesis, chronic inflammation, autoimmune diseases, fibrotic diseases, cancer and tumor metastasis and progression, ocular conditions, metabolic conditions such as obesity and/or diabetes mellitus, conditions such as cholestatic or other forms of chronic pruritus as well as acute and chronic organ transplant rejection.

Objects of the present invention are the compounds of formula (I) and their aforementioned salts and esters and their use as therapeutically active substances, a process for the manufacture of the said compounds, intermediates, pharmaceutical compositions, medicaments containing the said compounds, their pharmaceutically acceptable salts or esters, the use of the said compounds, salts or esters for the treatment or prophylaxis of disorders or conditions that are associated with the activity of ATX and/or the biological activity of lysophosphatidic acid (LPA), particularly in the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, conditions of the respiratory system, vascular and cardiovascular conditions, fibrotic diseases, cancer, ocular conditions, metabolic conditions, cholestatic and other forms of chronic pruritus and acute and chronic organ transplant rejection, and the use of the said compounds, salts or esters for the production of medicaments for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, conditions of the respiratory system, vascular and cardiovascular conditions, fibrotic diseases, cancer, ocular conditions, metabolic conditions, cholestatic and other forms of chronic pruritus and acute and chronic organ transplant rejection.

The term “alkenyl” denotes a monovalent linear or branched hydrocarbon group of 2 to 7 carbon atoms with at least one double bond. In particular embodiments, alkenyl has 2 to 4 carbon atoms with at least one double bond. Examples of alkenyl include ethenyl, propenyl, prop-2-enyl, isopropenyl, n-butenyl and iso-butenyl. Particular alkenyl group is ethenyl.

The term “alkoxy” denotes a group of the formula -O-R', wherein R' is an alkyl group. Examples of alkoxy group include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy and tert-butoxy. Particular alkoxy group include methoxy.

The term “alkoxyalkoxy” denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group has been replaced by another alkoxy group. Examples of alkoxyalkoxy group include methoxymethoxy, ethoxymethoxy, methoxyethoxy, ethoxyethoxy,

methoxypropoxy and ethoxypropoxy. Particular alkoxyalkoxy groups include methoxymethoxy and methoxyethoxy.

The term “alkoxyalkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by an alkoxyalkoxy group. Examples of
5 alkoxyalkoxyalkyl group include methoxymethoxymethyl, ethoxymethoxymethyl, methoxyethoxymethyl, ethoxyethoxymethyl, methoxypropoxymethyl, ethoxypropoxymethyl, methoxymethoxyethyl, ethoxymethoxyethyl, methoxyethoxyethyl, ethoxyethoxyethyl, methoxypropoxyethyl and ethoxypropoxyethyl.

The term “alkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen
10 atoms of the alkyl group has been replaced by an alkoxy group. Exemplary alkoxyalkyl groups include methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methoxypropyl, ethoxypropyl and isopropoxymethyl. Particular alkoxyalkyl group include include methoxymethyl, methoxyethyl and isopropoxymethyl.

The term “alkoxyhaloalkyl” denotes a haloalkyl group wherein at least one of the
15 hydrogen atoms of the haloalkyl group has been replaced by an alkoxy group. Exemplary alkoxyhaloalkyl groups include methoxytrifluoroethyl, ethoxytrifluoroethyl, methoxytrifluoropropyl, ethoxytrifluoropropyl and isopropoxytrifluoroethyl. Particular alkoxyhaloalkyl group include include methoxytrifluoroethyl.

The term “alkyl” denotes a monovalent linear or branched saturated hydrocarbon group of
20 1 to 12 carbon atoms. In particular embodiments, alkyl has 1 to 7 carbon atoms, and in more particular embodiments 1 to 4 carbon atoms. Examples of alkyl include methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl and sec-butyl, pentyl. Particular alkyl groups include methyl, ethyl, propyl and isopropyl. More particular alkyl groups are methyl and isopropyl.

The term “alkylcarbonyl” denotes a group of the formula $-C(O)-R'$, wherein R' is an alkyl
25 group. Examples of alkylcarbonyl groups include groups of the formula $-C(O)-R'$, wherein R' is methyl or ethyl. Particular alkylcarbonyl groups include groups of the formula $-C(O)-R'$, wherein R' is methyl.

The term “alkylsulfanyl” denotes a group of the formula $-S-R'$, wherein R' is an alkyl group. Examples of alkylsulfanyl groups include groups of the formula

-S-R', wherein R' is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl or tert-butyl. Particular alkylsulfanyl groups include group of the formula -S-R', wherein R' is methyl.

The term "alkylsulfinyl" denotes a group of the formula -S(O)-R', wherein R' is an alkyl group. Examples of alkylsulfinyl groups include groups of the formula

5 -S(O)-R', wherein R' is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl or tert-butyl.

Particular alkylsulfinyl groups include group of the formula -S(O)-R', wherein R' is methyl.

The term "alkylsulfonyl" denotes a group of the formula -S(O)₂-R', wherein R' is an alkyl group. Examples of alkylsulfonyl groups include groups of the formula

-S(O)₂-R', wherein R' is methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl or tert-butyl.

10 Particular alkylsulfonyl groups include group of the formula -S(O)₂-R', wherein R' is methyl.

The term "alkynyl" denotes a monovalent linear or branched saturated hydrocarbon group of 2 to 7 carbon atoms comprising one, two or three triple bonds. In particular embodiments alkynyl has from 2 to 4 carbon atoms comprising one or two triple bonds. Examples of alkynyl include ethynyl, propynyl, prop-2-ynyl, isopropynyl and n-butylnyl.

15 The term "amino" denotes a -NH₂ group.

The term "aminoalkyl" denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by an aminogroup. Examples of aminoalkyl include aminomethyl, aminoethyl, amino-1-methyl-ethyl, aminopropyl, aminomethylpropyl and aminopropyl. Particular examples are aminomethyl and haminoethyl.

20 The term "aminosulfonyl" denotes a -S(O)₂-NH₂ group.

The term "carbonyl" denotes a -C(O)- group.

The term "cyano" denotes a -C≡N group.

The term "cycloalkoxy" denotes a group of the formula -O-R', wherein R' is a cycloalkyl group. Examples of cycloalkoxy group include cyclopropoxy, cyclobutoxy, cyclopentyloxy,

25 cyclohexyloxy, cycloheptyloxy and cyclooctyloxy. Particular cycloalkoxy group is cyclopropoxy.

The term "cycloalkoxyalkyl" denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a cycloalkoxy group. Examples of cycloalkoxyalkyl groups include cyclopropoxymethyl, cyclopropoxyethyl, cyclobutoxymethyl,

cyclobutoxyethyl, cyclopentyloxymethyl, cyclopentyloxyethyl, cyclohexyloxymethyl, cyclohexyloxyethyl, cycloheptyloxymethyl, cycloheptyloxyethyl, cyclooctyloxymethyl and cyclooctyloxyethyl.

The term “cycloalkyl” denotes a monovalent saturated monocyclic or bicyclic hydrocarbon group of 3 to 10 ring carbon atoms. In particular embodiments, cycloalkyl denotes a monovalent saturated monocyclic hydrocarbon group of 3 to 8 ring carbon atoms. Bicyclic means a ring system consisting of two saturated carbocycles having two carbon atoms in common. Examples for monocyclic cycloalkyl are cyclopropyl, cyclobutanyl, cyclopentyl, cyclohexyl or cycloheptyl. Examples for bicyclic cycloalkyl are bicyclo[2.2.1]heptanyl or bicyclo[2.2.2]octanyl. Particular monocyclic cycloalkyl groups are cyclopropyl, cyclobutanyl, cyclopentyl and cyclohexyl. More particular monocyclic cycloalkyl group is cyclopropyl.

The term “cycloalkylalkoxy” denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group is replaced by a cycloalkyl group. Examples of cycloalkylalkoxy include cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy, cycloheptylmethoxy and cyclooctylmethoxy.

The term “cycloalkylalkoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a cycloalkylalkoxy group. Examples of cycloalkylalkoxyalkyl include cyclopropylmethoxymethyl, cyclopropylmethoxyethyl, cyclobutylmethoxymethyl, cyclobutylmethoxyethyl, cyclopentylmethoxyethyl, cyclopentylmethoxyethyl, cyclohexylmethoxymethyl, cyclohexylmethoxyethyl, cycloheptylmethoxymethyl, cycloheptylmethoxyethyl, cyclooctylmethoxymethyl and cyclooctylmethoxyethyl.

The term “cycloalkylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group is replaced by a cycloalkyl group. Examples of cycloalkylalkyl include cyclopropylmethyl, cyclopropylethyl, cyclopropylbutyl, cyclobutylpropyl, 2-cyclopropylbutyl, cyclopentylbutyl, cyclohexylmethyl, cyclohexylethyl, bicyclo[4.1.0]heptanymethyl, bicyclo[4.1.0]heptanylethyl, bicyclo[2.2.2]octanymethyl, bicyclo[2.2.2]octanylethyl, adamantanymethyl and adamantanylethyl. Particular examples of cycloalkylalkyl are cyclohexylmethyl, cyclohexylethyl, bicyclo[4.1.0]heptanymethyl, bicyclo[4.1.0]heptanylethyl, bicyclo[2.2.2]octanymethyl, bicyclo[2.2.2]octanylethyl, adamantanymethyl and adamantanylethyl. Further particular examples cycloalkylalkyl are cyclohexylmethyl,

cyclohexylethyl, bicyclo[4.1.0]heptanylmethyl, bicyclo[2.2.2]octanylmethyl, adamantanylmethyl and adamantanylethyl.

The term “cycloalkylcarbonyl” of the formula $-C(O)-R'$, wherein R' is a cycloalkyl group. Examples of cycloalkylcarbonyl groups include groups of the formula $-C(O)-R'$, wherein R' is
5 cyclopropyl.

The term “cycloalkylsulfanyl” denotes a group of the formula $-S-R'$, wherein R' is a cycloalkyl group. Examples of cycloalkylsulfanyl groups include groups of the formula $-S-R'$, wherein R' is cyclopropyl.

The term “cycloalkylsulfinyl” denotes a group of the formula $-S(O)-R'$, wherein R' is a
10 cycloalkyl group. Examples of cycloalkylsulfinyl groups include groups of the formula $-S(O)-R'$, wherein R' is cyclopropyl.

The term “cycloalkylsulfonyl” denotes a group of the formula $-S(O)_2-R'$, wherein R' is a cycloalkyl group. Examples of cycloalkylsulfonyl groups include groups of the formula $-S(O)_2-R'$, wherein R' is cyclopropyl.

15 The term “haloalkoxy” denotes an alkoxy group wherein at least one of the hydrogen atoms of the alkoxy group has been replaced by the same or different halogen atoms. The term “perhaloalkoxy” denotes an alkoxy group where all hydrogen atoms of the alkoxy group have been replaced by the same or different halogen atoms. Examples of haloalkoxy include fluoromethoxy, difluoromethoxy, trifluoromethoxy, trifluoroethoxy, trifluoromethylethoxy,
20 trifluorodimethylethoxy and pentafluoroethoxy. Particular haloalkoxy group is trifluoromethoxy.

The term “haloalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by the same or different halogen atoms. The term “perhaloalkyl” denotes an alkyl group where all hydrogen atoms of the alkyl group have been replaced by the same or different halogen atoms. Examples of haloalkyl include fluoromethyl,
25 difluoromethyl, trifluoromethyl, trifluoroethyl, trifluoromethylethyl and pentafluoroethyl. Particular haloalkyl group is trifluoromethyl.

The term “haloalkylsulfanyl” denotes a group of the formula $-S-R'$, wherein R' is a haloalkyl group. Examples of haloalkylsulfanyl groups include groups of the formula $-S-R'$, wherein R' is trifluoromethyl.

The term “haloalkylsulfinyl” denotes a group of the formula $-S(O)-R'$, wherein R' is a haloalkyl group. Examples of haloalkylsulfinyl groups include groups of the formula $-S(O)-R'$, wherein R' is trifluoromethyl.

5 The term “haloalkylsulfonyl” denotes a group of the formula $-S(O)_2-R'$, wherein R' is a haloalkyl group. Examples of haloalkylsulfonyl groups include groups of the formula $-S(O)_2-R'$, wherein R' is trifluoromethyl.

The term “halogen” and “halo” are used interchangeably herein and denote fluoro, chloro, bromo, or iodo. Particular halogens are chloro, fluoro and bromo. More particular halogens are chloro and fluoro.

10 The term “hydroxy” denotes an $-OH$ group.

The term “hydroxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a hydroxy group. Examples of hydroxyalkyl include hydroxymethyl, hydroxyethyl, hydroxy-1-methyl-ethyl, hydroxypropyl, hydroxymethylpropyl and dihydroxypropyl. Particular examples are hydroxymethyl and
15 hydroxyethyl.

The term “hydroxyhaloalkyl” denotes a haloalkyl group wherein at least one of the hydrogen atoms of the haloalkyl group has been replaced by an hydroxy group. Exemplary hydroxyhaloalkyl groups include hydroxytrifluoroethyl and hydroxytrifluoropropyl. Particular hydroxyhaloalkyl groups include hydroxytrifluoroethyl.

20 The term “indanyloxy” denotes a group of the formula $-O-R'$, wherein R' is an indanyl.

The term “indanyloxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a indanyloxy group. Exemplary indanyloxyalkyl groups include indanyloxymethyl, indanyloxyethyl and indanyloxypropyl. Particular indanyloxyalkyl group is indanyloxymethyl.

25 The term “phenoxy” denotes a group of the formula $-O-R'$, wherein R' is a phenyl.

The term “phenoxyalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced by a phenoxy group. Exemplary phenoxyalkyl groups

include phenoxyethyl, phenoxypropyl. Particular phenoxyalkyl group is phenoxyethyl.

The term “phenylalkenyl” denotes an alkenyl group wherein at least one of the hydrogen atoms of the alkenyl group has been replaced a phenyl. Particular phenylalkenyl group is phenylethenyl.

The term “phenylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced a phenyl. Particular phenylalkyl groups are benzyl, phenethyl and phenylpropyl. More particular phenylalkyl groups are benzyl and phenethyl. Further particular phenylalkyl group is benzyl.

The term “phenylalkynyl” denotes an alkynyl group wherein at least one of the hydrogen atoms of the alkynyl group has been replaced a phenyl. Particular phenylalkynyl group is phenylethynyl.

The term “phenylcycloalkyl” denotes a cycloalkyl group wherein at least one of the hydrogen atoms of the cycloalkyl group has been replaced a phenyl. Particular phenylcycloalkyl group is phenylcyclopropyl.

The term “pyridinylalkenyl” denotes an alkenyl group wherein at least one of the hydrogen atoms of the alkenyl group has been replaced a pyridinyl. Particular pyridinylalkenyl group is pyridinylethenyl.

The term “pyridinylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced a pyridinyl. Particular pyridinylalkyl groups are pyridinylmethyl, pyridinylethyl and pyridinylpropyl. More particular pyridinylalkyl group is pyridinylethyl.

The term “pyridinylalkynyl” denotes an alkynyl group wherein at least one of the hydrogen atoms of the alkynyl group has been replaced a pyridinyl. Particular pyridinylalkynyl group is pyridinylethynyl. The term “thiophenylalkenyl” denotes an alkenyl group wherein at least one of the hydrogen atoms of the alkenyl group has been replaced a thiophenyl. Particular thiophenylalkenyl group is thiophenylethenyl.

The term “thiophenylalkyl” denotes an alkyl group wherein at least one of the hydrogen atoms of the alkyl group has been replaced a thiophenyl. Particular thiophenylalkyl groups are

thiophenylmethyl, thiophenylethyl and thiophenylpropyl. More particular thiophenylalkyl group is thiophenylmethyl.

The term "thiophenylalkynyl" denotes an alkynyl group wherein at least one of the hydrogen atoms of the alkynyl group has been replaced a thiophenyl. Particular

5 thiophenylalkynyl group is thiophenylethynyl.

The term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. The salts are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, in particular

10 hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, N-acetylcystein and the like. In addition, these salts may be prepared by addition of an inorganic base or an organic base to the free acid. Salts derived from
15 an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium salts and the like. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine,
20 lysine, arginine, N-ethylpiperidine, piperidine, polyimine resins and the like. Particular pharmaceutically acceptable salts of compounds of formula (I) are the hydrochloride salts, methanesulfonic acid salts and citric acid salts.

"Pharmaceutically acceptable esters" means that compounds of general formula (I) may be derivatised at functional groups to provide derivatives which are capable of conversion back to
25 the parent compounds in vivo. Examples of such compounds include physiologically acceptable and metabolically labile ester derivatives, such as methoxymethyl esters, methylthiomethyl esters and pivaloyloxymethyl esters. Additionally, any physiologically acceptable equivalents of the compounds of general formula (I), similar to the metabolically labile esters, which are capable of producing the parent compounds of general formula (I) in vivo, are within the scope
30 of this invention.

The term "protecting group" (PG) denotes a group which selectively blocks a reactive site in a multifunctional compound such that a chemical reaction can be carried out selectively at another unprotected reactive site in the meaning conventionally associated with it in synthetic chemistry. Protecting groups can be removed at the appropriate point. Exemplary protecting groups are amino-protecting groups, carboxy-protecting groups or hydroxy-protecting groups. Particular protecting groups are the tert-butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), fluorenylmethoxycarbonyl (Fmoc) and benzyl (Bn) groups. Further particular protecting groups are the tert-butoxycarbonyl (Boc) and the fluorenylmethoxycarbonyl (Fmoc) groups. More particular protecting group is the tert-butoxycarbonyl (Boc) group.

10 The abbreviation uM means microMolar and is equivalent to the symbol μM .

The abbreviation uL means microliter and is equivalent to the symbol μL .

The abbreviation ug means microgram and is equivalent to the symbol μg .

The compounds of formula (I) can contain several asymmetric centers and can be present in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates.



According to the Cahn-Ingold-Prelog convention the asymmetric carbon atom can be of the "R" or "S" configuration.

Also an embodiment of the present invention are compounds according to formula (I) as described herein and pharmaceutically acceptable salts or esters thereof, in particular compounds according to formula (I) as described herein and pharmaceutically acceptable salts thereof, more particularly compounds according to formula (I) as described herein.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein

25 R^1 is alkyl, haloalkyl, substituted cycloalkyl, substituted cycloalkylalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted phenylalkynyl, substituted pyridinyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted pyridinylalkynyl, substituted thiophenyl, substituted thiophenylalkyl, substituted

thiophenylalkenyl, substituted thiophenylalkynyl, substituted 2,3-dihydro-1H-isoin-
 2-yl, substituted 1H-indol-2-yl or substituted benzofuran-2-yl wherein substituted
 cycloalkyl, substituted cycloalkylalkyl, substituted phenyl, substituted phenylalkyl,
 substituted phenylalkynyl, substituted phenoxyalkyl, substituted phenylcycloalkyl,
 substituted phenylalkenyl, substituted pyridinyl, substituted pyridinylalkyl, substituted
 pyridinylalkenyl, substituted pyridinylalkynyl, substituted thiophenyl, substituted
 thiophenylalkyl, substituted thiophenylalkenyl, substituted thiophenylalkynyl,
 substituted 2,3-dihydro-1H-isoin-2-yl, substituted 1H-indol-2-yl and substituted
 benzofuran-2-yl are substituted with R^8 , R^9 and R^{10} ;

Y is $-OC(O)-$, $-NR^7C(O)-$, $-C(O)-$, $-S(O)_2-$,  or  ;

A is $-N-$ or CR^5- ;

W is $-O-$, $-S-$, $-NR^6-$, $-C(O)-$, $-S(O)_2-$, $-C(O)-NR^6-$ or $-CR^3R^4-$;

R^3 and R^4 are independently selected from H, halogen, alkyl and cycloalkyl;


R^5 , R^6 and R^7 are independently selected from H, alkyl and cycloalkyl;

R^8 , R^9 and R^{10} are independently selected from H, alkyl, hydroxyalkyl, haloalkyl,
 hydroxyhaloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy,
 cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy,
 alkoxyhaloalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, phenyl, substituted phenyl,
 pyridinyl, substituted pyridinyl, halogen, hydroxy, cyano, alkylsulfanyl,
 haloalkylsulfanyl, cycloalkylsulfanyl, alkylsulfinyl, haloalkylsulfinyl,
 cycloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, cycloalkylsulfonyl, substituted
 aminosulfonyl, substituted amino and substituted aminoalkyl, wherein substituted
 aminosulfonyl, substituted amino and substituted aminoalkyl are substituted on the
 nitrogen atom with one to two substituents independently selected from H, alkyl,
 cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, alkylcarbonyl and
 cycloalkylcarbonyl, and wherein substituted phenyl and substituted pyridinyl are
 optionally substituted with one to three substituents independently selected from alkyl,
 halogen, haloalkyl, alkoxy and haloalkoxy;


-16-

m, n, p and q are independently selected from 1 or 2;


-17-

**B**


,

**C**


,

**D**


,

**E**


,

**F**


,

**G**


,

**H**


,

**I**


,

**J**


,

**K**


,

**L**


,

**M**


,

**N**


,

**O**


,

**P**


,

**Q**

,


**R**

,


**S**

,


-18-

**T**


,

**U**


,

**V**


,

**X**


,

**Z**


,

**AA**


,

**AB**


,

**AC**

,

**AD**

,

**AE**

;

R^{11} is H, alkyl, haloalkyl or cycloalkyl;

R^2 is selected from the ring systems B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, X, Z, AA, AB, AC, AD and AE;

5 and pharmaceutically acceptable salts.

A further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is substituted cycloalkylalkyl, substituted piperazinyl, substituted piperidinyl, substituted indanyloxyalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted

10 pyridinylalkyl, substituted pyridinylalkenyl, naphthyl, substituted naphthyl, substituted quinolinyl,


substituted isoquinolinyl, or substituted 1H-indol-2-yl, wherein substituted cycloalkylalkyl, substituted piperazinyl, substituted piperidinyl, substituted indanyloxyalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted naphthyl, substituted quinolinyl, substituted isoquinolinyl and substituted 1H-indol-2-yl are substituted with R^8 , R^9 and R^{10} .

Another further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is substituted cycloalkylalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinylalkyl, substituted pyridinylalkenyl or substituted 1H-indol-2-yl, wherein substituted cycloalkylalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, and substituted 1H-indol-2-yl are substituted with R^8 , R^9 and R^{10} .

A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is substituted phenylalkyl or substituted phenylalkenyl, wherein substituted phenylalkyl and substituted phenylalkenyl are substituted with R^8 , R^9 and R^{10} .


In a further embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^1 is phenylalkyl substituted with R^8 , R^9 and R^{10} .

The present invention also relates to compounds according to formula (I) as described

herein, wherein Y is $-\text{OC}(\text{O})-$, $-\text{C}(\text{O})-$, $-\text{S}(\text{O})_2-$ or .

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein Y is $-\text{OC}(\text{O})-$ or $-\text{C}(\text{O})-$.

Another further embodiment of the present invention are compounds according to formula (I) as described herein, wherein Y is $-\text{OC}(\text{O})-$ and of formula (In).



Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein Y is -C(O)-.

A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein A is -N-.

5 Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein W is -O-, -NR⁶-, -C(O)-, -S(O)₂-, -C(O)-NR⁶- or -CR³R⁴-.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein W is -C(O)-, -C(O)-NR⁶- or -CR³R⁴-.

10 Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein W is -C(O)-.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R² is selected from the ring systems B, C, D, E, G, H, M, O, P, R, S, T, U, V, X, Z, AA, AB, AC, AD, AE, AF, AG, AH and AI.

15 Another further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R² is selected from the ring systems B, C, D, E, G, H, M, O, P, R, S, T, U, V, X, Z, AA, AB, AC, AD and AE.

Also a further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R² is selected from the ring systems B, D, H, M, O, R and AJ.

20 A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R² is selected from the ring systems B, D, H, O and R.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R² is selected from the ring systems B and D.

25 A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R² is the ring system D.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R³ and R⁴ are H.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁵ is H.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁶ is H or alkyl.

5 A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁸, R⁹ and R¹⁰ are independently selected from H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen, cyano, haloalkylsulfanyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, pyrrolyl substituted with one alkyl, pyrrolydiny, tetrahydrofuranyl, alkylcarbonyl, and aminosulfonyl substituted on
10 the nitrogen atom with one to two substituents independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, alkylcarbonyl and cycloalkylcarbonyl.

Also a further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁸, R⁹ and R¹⁰ are independently selected from H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen,
15 cyano, haloalkylsulfanyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl and aminosulfonyl substituted on the nitrogen atom with one to two substituents independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, alkylcarbonyl and cycloalkylcarbonyl.

A further particular embodiment of the present invention are compounds according to
20 formula (I) as described herein, wherein R⁸, R⁹ and R¹⁰ are independently selected from H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen, cyano, haloalkylsulfanyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl and aminosulfonyl substituted on the nitrogen atom with two alkyl.

A further particular embodiment of the present invention are compounds according to
25 formula (I) as described herein, wherein R⁸, R⁹ and R¹⁰ are independently selected from H, alkyl, haloalkyl, haloalkoxy, halogen and alkylsulfonyl.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R⁸ is H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen, cyano, haloalkylsulfanyl,
30 haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, pyrrolyl substituted with one alkyl,

pyrrolydiny, tetrahydrofuranyl, alkylcarbonyl, or aminosulfonyl substituted on the nitrogen atom with two alkyl.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^8 is H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen, cyano, haloalkylsulfanyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl or aminosulfonyl substituted on the nitrogen atom with two alkyl.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^8 is haloalkyl, haloalkoxy, halogen or alkylsulfonyl.

10 A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^8 is haloalkoxy or halogen.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^8 is halogen.

15 The present invention also relates to compounds according to formula (I) as described herein, wherein R^9 is H, alkyl, haloalkyl, cycloalkyl, cycloalkoxy, alkoxy, haloalkoxy, alkoxyalkoxy, cyano or halogen.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^9 is H, alkyl, haloalkyl, alkoxy or halogen.

20 Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^9 is H, alkyl or halogen.

The present invention also relates to compounds according to formula (I) as described herein, wherein R^8 and R^9 are halogen.

25 Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^9 is H, alkyl, haloalkyl, cycloalkyl, cycloalkoxy, alkoxy, haloalkoxy, alkoxyalkoxy, cyano or halogen.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{10} is H or alkyl.

Another embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{10} is H.

Also an embodiment of the present invention are compounds according to formula (I) as described herein, wherein R^{11} is haloalkyl.

5 A particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein m is 1.

A further particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein n is 1.

10 A more particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein m and n are 1.

Also a particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein p and q are 1.

The present invention also relates to compounds according to formula (I) as described herein, wherein m, n, p and q are 1.

15 A furthermore particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein

R^1 is substituted phenylalkyl or substituted phenylalkenyl, wherein substituted phenylalkyl and substituted phenylalkenyl are substituted with R^8 , R^9 and R^{10} ;

Y is -OC(O)- or -C(O)-;

20 A is -N-;

W is -C(O)-;

R^8 is haloalkyl, haloalkoxy, halogen or alkylsulfonyl;

R^9 is H, alkyl or halogen;

R^{10} is H or alkyl;

25 m and n are 1;

-24-

p and q are independently selected from 1 or 2;

R^2 is selected from the ring systems B, D, H, M, O, R and AJ;

R^{12} is halogen

and pharmaceutically acceptable salts.

5 Also a furthermore particular embodiment of the present invention are compounds according to formula (I) as described herein, wherein

R^1 is substituted phenylalkyl or substituted phenylalkenyl, wherein substituted phenylalkyl and substituted phenylalkenyl are substituted with R^8 , R^9 and R^{10} ;

Y is -OC(O)- or -C(O)-;

10 A is -N-;

W is -C(O)-;

R^8 is haloalkyl, haloalkoxy, halogen or alkylsulfonyl;

R^9 is H, alkyl or halogen;

R^{10} is H or alkyl;

15 m and n are 1;

p and q are independently selected from 1 or 2;

R^2 is selected from the ring systems B and D;

and pharmaceutically acceptable salts.

Particular examples of compounds of formula (I) as described herein are selected from

20 (E)-1-[(3aS,8aR)-2-(4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)propan-1-one;

6-((3aR,6aS)-5-(3-(3,5-dichlorophenyl)propanoyl)octahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzo[d]oxazol-2(3H)-one;

(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

5 (3aR,6aS)-3,5-dichlorobenzyl 5-(9H-pyrido[3,4-b]indole-3-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-indole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

10 (3aR,6aS)-3,5-dichlorobenzyl 5-(9H-carbazole-3-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-indazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-benzo[d]imidazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

15 trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

cis-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

20 (3aR,8aS)-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate;

(1H-benzotriazol-5-yl)-{(3aS,6aR)-5-[2-(3-chloro-phenyl)-ethanesulfonyl]-hexahydropyrrolo[3,4-c]pyrrol-2-yl}-methanone;

(3aR,6aS)-3,5-dichlorobenzyl 5-(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

25 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chlorophenyl)-2,2-dimethylpropan-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(3aSR,6SR,7aSR)-6-((3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl)-hexahydro-benzooxazol-2-one;

5 (E)-1-[(3aS,8aR)-2-(benzo[c][1,2,5]oxadiazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-methyl-4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

10 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-methoxyphenyl)prop-2-en-1-one;

(E)-1-[(3aS,8aR)-2-((S)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one

15 (E)-1-[(3aS,8aR)-2-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one (E)-1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(2-isopropyl-phenyl)-prop-2-ene-1-one;

trans-3,5-dichlorobenzyl 2-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

20 (3aR,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

6-((3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl)-3H-benzooxazol-2-one;

(3aR,5s,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yloxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate;

25 (3aR,5r,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yloxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate;

(3aS,6aS)-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-2-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester;

5 (3aR,6aR)-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

10 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester;

(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

15 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester;

cis-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate;

20 (3aS,7aR)-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

trans-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate;

(3aR,8aS)-3,5-dichlorobenzyl 6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate;

25 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 1-(3-chloro-phenyl)-cyclopropyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid bicyclo[4.1.0]hept-7-ylmethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid adamantan-2-ylmethyl ester;

5 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 1-fluoro-cyclohexylmethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-adamantan-2-yl-ethyl ester;

10 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-adamantan-1-yl-ethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid adamantan-1-ylmethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid cyclohexylmethyl ester;

15 cis-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2,2-trifluoro-1-methoxy-ethyl)-benzyl ester;

cis-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2,2-trifluoro-1-hydroxy-ethyl)-benzyl ester;

20 (3aR,6aS)-2-cyclohexylethyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-5-trifluoromethoxy-benzyl ester;

(3aR,6aS)-3-chloro-5-cyanobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

25 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-trifluoromethoxy-benzyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-5-trifluoromethyl-benzyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-trifluoromethoxy-benzyl ester;

5 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-fluoro-3-trifluoromethoxy-benzyl ester;

(3aR,6aS)-3-cyano-5-fluorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

10 (3aR,6aS)-3-chloro-5-methoxybenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1S,4R)-3-methyl-bicyclo[2.2.1]hept-2-ylmethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1R,4S)-1-bicyclo[2.2.1]hept-2-ylmethyl ester;

15 (3aR,5S,6aS)-5-[(3H-[1,2,3]triazol-4-ylmethyl)-carbamoyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid 3,5-dichloro-benzyl ester;

(3aS,6aS)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

20 (3aS,6aR)-5-[(1H-[1,2,3]triazol-4-ylmethyl)-carbamoyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3,5-dichloro-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aR,5r,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-ylamino)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate;

25 (3aR,6aS)-3,5-dichlorobenzyl 5-((1H-benzo[d]imidazol-5-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

1-((3aR,6aS)-5-((1H-benzo[d][1,2,3]triazol-5-yl)methyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)propan-1-one;

(3aR,6aS)-3,5-dichlorobenzyl 5-((1H-indazol-5-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

5 (3aR,6aS)-3,5-dichlorobenzyl 5-((2-oxo-2,3-dihydrobenzo[d]oxazol-6-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

6-(((3aR,6aS)-5-(3-(3,5-dichlorophenyl)propanoyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methyl)benzo[d]oxazol-2(3H)-one;

10 4-{(E)-3-[(3aS,8aR)-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-oxo-propenyl}-benzonitrile;

(E)-1-[(3aS,8aR)-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(3aR,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-ylsulfonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

15 (3aR,6aS)-3,5-dichlorobenzyl 5-(4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

20 (E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

cis-5-((3aR,8aS)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)decahydropyrrolo[3,4-d]azepine-2-carbonyl)hexahydrooxazolo[5,4-c]pyridin-2(1H)-one;

6-{(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl}-5,6,7,8-tetrahydro-1H-[1,6]naphthyridin-2-one;

25 (3aR,7aR)-5-{(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl}-hexahydro-oxazolo[5,4-c]pyridin-2-one;

(E)-1-[(3aS,8aR)-2-(7,8-Dihydro-5H-[1,6]naphthyridine-6-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-3-(4-trifluoromethoxy-phenyl)-1-[(3aS,8aR)-2-(2-trifluoromethyl-5,6-dihydro-8H-[1,2,4]triazolo[1,5-a]pyrazine-7-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-prop-2-ene-1-one;

(3aS,8aR)-6-[(E)-3-(3-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid (1H-[1,2,3]triazol-4-ylmethyl)-amide;

(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

(3aS,6aR)-5-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1H-[1,2,3]triazol-4-ylmethyl)-amide;

(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid (4H-[1,2,4]triazol-3-ylmethyl)-amide;

(E)-1-[(3aS,8aR)-2-(6,7-dihydro-4H-[1,2,3]triazolo[1,5-a]pyrazine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-imidazo[4,5-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

(3aS,8aR)-6-[3-(3-chloro-phenyl)-2,2-dimethyl-propionyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid (3H-[1,2,3]triazol-4-ylmethyl)-amide;

(3aR,8aS)-N-(2-(1H-1,2,3-triazol-5-yl)ethyl)-6-((E)-3-(4-

(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

(3aR,7aS)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3,5-dichloro-benzyl ester;

(3aS,7aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3,5-dichloro-benzyl ester;

5 (+)-trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

(-)-trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

10 (-)-trans-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate;

(+)-trans-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate;

(E)-1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(3,5-dichloro-phenyl)-prop-2-en-1-one;

15 (1H-benzotriazol-5-yl)-{trans-2-[5-(4-chloro-phenyl)-[1,3,4]oxadiazol-2-yl]-octahydro-pyrrolo[3,4-c]pyridin-5-yl}-methanone;

(E)-1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

20 (1H-benzotriazol-5-yl)-[(3aR,6aS)-5-(5-chloro-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

(E)-1-[(3aR,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-5-trifluoromethyl-phenyl)-prop-2-ene-1-one;

1-[(3aR,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-5-trifluoromethyl-phenyl)-propan-1-one;

25 (1H-benzotriazol-5-yl)-[(3aR,6aS)-5-(6-chloro-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylsulfonyl)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-chlorophenyl)prop-2-en-1-one;

5 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-p-tolylprop-2-en-1-one;

4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)-N,N-dimethylbenzenesulfonamide;

10 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxyphenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)benzonitrile;

15 (E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one;

20 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluorophenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-phenylprop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-2-yl)prop-2-en-1-one;

25 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-3-yl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chlorophenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(4-chlorophenyl)prop-2-en-1-one;

5 (E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(difluoromethoxy)phenyl)prop-2-en-1-one;

10 (E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one;

4-((E)-3-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-oxoprop-1-enyl)benzonitrile;

(E)-1-((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

15 (-)-(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(+)-(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

20 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3,5-dichlorophenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-4-yl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2,4-difluorophenyl)prop-2-en-1-one;

25 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2,4-dichlorophenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3,4-dichlorophenyl)prop-2-en-1-one;

(E)-1-[(3aS,7aS)-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-difluoromethoxy-phenyl)-prop-2-ene-1-one;

5 4-{(E)-3-[(3aS,7aS)-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-oxo-propenyl}-benzonitrile;

4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)-3-fluorobenzonitrile;

10 4-((E)-3-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-oxoprop-1-enyl)-3-fluorobenzonitrile;

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(difluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-[cis-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

15 3-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)benzonitrile;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

20 (E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-chloro-2-fluorophenyl)prop-2-en-1-one;

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)prop-2-en-1-one;

(E)-1-((3aR,6aS)-5-(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

5 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

(E)-1-[(3aS,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-methanesulfonyl-phenyl)-prop-2-ene-1-one;

10 (E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3,5-dimethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

15 (E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-methoxy-phenyl)-prop-2-ene-1-one;

3-{(E)-3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-oxo-propenyl}-5-chloro-benzonitrile;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-methoxy-5-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

20 (E)-1-[(3aR,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

1-[(3aR,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

25 (3aR,7aR)-5-[(3aR,6aR)-5-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carbonyl]-hexahydro-oxazolo[5,4-c]pyridin-2-one;

(3aR,7aR)-5-[(3aR,6aR)-5-[3-(4-trifluoromethoxy-phenyl)-propionyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carbonyl]-hexahydro-oxazolo[5,4-c]pyridin-2-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-phenyl-prop-2-ene-1-one;

5 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-phenyl-propan-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-prop-2-ene-1-one;

10 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-propan-1-one;

(3aR,6aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-5-(3-(4-(trifluoromethoxy)phenyl)propanoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide;

15 (3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acryloyl)-N-methyloctahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-trifluoromethyl-pyridin-3-yl)-propan-1-one;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-trifluoromethoxy-phenoxy)-ethanone;

20 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-biphenyl-4-yl-propan-1-one;

25 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluoro-2-(trifluoromethyl)phenyl)prop-2-en-1-one;

1-[(3aS,6aR)-5-(1H-Benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(methylsulfonyl)phenyl)prop-2-en-1-one;

5 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylthio)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

10 1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-2-(3-(trifluoromethoxy)phenoxy)ethanone;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3-trifluoromethoxy-phenyl)-prop-2-en-1-one;

15 (E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one;

(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3-chloro-5-trifluoromethoxy-phenyl)-prop-2-en-1-one;

20 (E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3,5-dichloro-phenyl)-prop-2-en-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-phenyl-pyridin-3-yl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-trifluoromethyl-pyridin-2-yl)-prop-2-ene-1-one;

25 (E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-4-yl-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-3-yl-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-2-yl-phenyl)-prop-2-ene-1-one;

5 1-[(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-3-methyl-phenoxy)-ethanone;

1-[(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-methyl-phenoxy)-ethanone;

10 (E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-phenyl-pyridin-2-yl)-prop-2-ene-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylsulfinyl)phenyl)prop-2-en-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)propan-1-one;

15 (3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-6-(3-(4-(trifluoromethoxy)phenyl)propanoyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-(3-(3-fluoro-4-(trifluoromethoxy)phenyl)propanoyl)-N-methyloctahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

20 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-difluoromethoxy-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(2-fluoro-4-trifluoromethoxy-phenyl)-propan-1-one;

25 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluoro-2-(trifluoromethyl)phenyl)propan-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-methyl-4-(trifluoromethoxy)phenyl)propan-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-methoxyphenyl)propan-1-one;

5 1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(2-isopropyl-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-trifluoromethyl-pyridin-2-yl)-propan-1-one;

10 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-phenyl-pyridin-2-yl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-4-yl-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-3-yl-phenyl)-propan-1-one;

15 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-2-yl-phenyl)-propan-1-one;

(3aS,8aR)-6-[3-(4-trifluoromethoxy-phenyl)-propionyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid [2-(3H-[1,2,3]triazol-4-yl)-ethyl]-amide;

20 (E)-3-[4-(trifluoro-methoxy)-phenyl]-1-[(3aS,8aR)-2-((S)-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridine-6-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-prop-2-ene-1-one hydrochloride;

and pharmaceutically acceptable salts thereof.

Also particular examples of compounds of formula (I) as described herein are selected from

25 trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid
4-trifluoromethoxy-benzyl ester;

1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-2-(4-
trifluoromethoxy-phenoxy)-ethanone;

5 (E)-1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-
trifluoromethoxy-phenyl)-propenone;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-
d]azepin-6(7H)-yl)-3-(3-chloro-5-(trifluoromethyl)phenyl)prop-2-en-1-one;

10 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-
d]azepin-6(7H)-yl)-3-(4-methoxy-2-(trifluoromethyl)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-
d]azepin-6(7H)-yl)-3-(2-cyclopropylphenyl)prop-2-en-1-one;

trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid
4-fluoro-2-trifluoromethyl-benzyl ester;

15 trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid
2-cyclopropyl-4-trifluoromethyl-benzyl ester;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(2-
trifluoromethoxy-phenoxy)-ethanone;

20 trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid
2-methoxy-4-trifluoromethoxy-benzyl ester;

4-{2-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-
oxo-ethoxy}-3-trifluoromethyl-benzonitrile;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(4-
chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

25 1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-[4-
methyl-2-(1-methyl-pyrrolidin-3-yl)-phenoxy]-ethanone;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(2-chloro-4-fluoro-phenoxy)-ethanone;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(2-chloro-4-(trifluoromethyl)phenoxy)ethanone;

5 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(6-isopropyl-3,3-dimethyl-2,3-dihydro-1H-inden-5-yloxy)ethanone;

(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester;

10 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(5-chloro-2-(trifluoromethyl)phenoxy)ethanone;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-2-(2-tert-butyl-4-methoxyphenoxy)ethanone;

4-[2-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-2-oxoethoxy]-3-propan-2-ylbenzonitrile;

15 1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-3-[3-fluoro-4-(2,2,2-trifluoroethoxy)phenyl]propan-1-one;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-3-[2-fluoro-4-(2,2,2-trifluoroethoxy)phenyl]propan-1-one;

20 (3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-fluoro-4-(2,2,2-trifluoro-ethoxy)-benzyl ester;

(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 2-fluoro-4-(2,2,2-trifluoro-ethoxy)-benzyl ester;

(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 4-(2,2,2-trifluoro-ethoxy)-benzyl ester;

25 (3aS,6aS)-5-(3H-[1,2,3]triazolo[4,5-b]pyridine-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

1-[(3aR,6aR)-5-(1H-triazolo[4,5-b]pyridine-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[4-(trifluoromethoxy)phenyl]propan-1-one;

(3aS,6aS)-5-(3H-[1,2,3]triazolo[4,5-c]pyridine-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

5 (3aS,6aS)-5-(4-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(7-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

10 (3aS,6aS)-5-(6-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(4-chloro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(6-trifluoromethyl-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

15 (3aS,6aS)-5-(4-methyl-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(6-methyl-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

20 1-[(3aR,6aR)-5-(4-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-[4-(trifluoromethoxy)phenyl]propan-1-one;

(4-ethoxyquinolin-2-yl)((3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone;

(4-ethoxyquinolin-2-yl)((3aS,6aS)-5-(4-fluoro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone;

25 6-[(3aR,6aR)-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrole-5-carbonyl]-3H-1,3-benzoxazol-2-one;

(3aS,6aS)-5-(2-oxo-2,3-dihydro-benzooxazole-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-benzyl ester;

5 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-(1,1,2,2-tetrafluoro-ethoxy)-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-difluoromethoxy-3-fluoro-benzyl ester;

10 (3aS,6aS)-5-(1H-[1,2,3]triazolo[4,5-b]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-difluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-[1,2,3]triazolo[4,5-b]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

15 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-(2,2,2-trifluoro-ethoxy)-benzyl ester;

20 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 5-trifluoromethoxy-pyridin-2-ylmethyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-benzyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-5-methyl-benzyl ester;

25 (3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-ethoxy-benzyl ester;

5 (3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-benzyl ester;

10 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-5-methyl-benzyl ester;

(3aS,6aS)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

15 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethyl-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethyl-benzyl ester;

20 (3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethyl-benzyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

25 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-methanesulfonyl-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-ethoxy-5-fluoro-benzyl ester;

(3aS,6aS)-5-(4-methoxy-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

5 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-cyclobutoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropoxy-benzyl ester;

10 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-(2,2,2-trifluoro-ethoxy)-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-chloro-2-ethoxy-5-fluoro-benzyl ester;

(E)-1-[trans-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propenone;

15 (3aS,6aS)-5-(4-sulfamoyl-piperidine-1-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-phenyl-pyridin-3-yl)-propan-1-one;

20 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-isopropyl-phenoxy)-ethanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-trifluoromethyl-phenoxy)-ethanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(biphenyl-2-yloxy)-ethanone;

25 (E)-1-[(3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propenone;

1-((3aR,6aR)-5-((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-chloro-4-trifluoromethoxy-phenoxy)-ethanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-pyrrol-1-yl-phenoxy)-ethanone;

4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-methoxy-benzonitrile;

4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-benzonitrile;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-phenoxy-ethanone;

2-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-5-trifluoromethoxy-benzonitrile;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(2-isopropyl-5-methylphenoxy)ethanone;

(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(6-trifluoromethoxy-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(5-trifluoromethoxy-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

1-[trans-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chloro-5-(trifluoromethyl)phenyl)propan-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxy-2-(trifluoromethyl)phenyl)propan-1-one;

5 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-cyclopropylphenyl)propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[3-methoxy-5-(trifluoromethoxy)phenyl]propan-1-one;

10 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-isopropyl-5-methyl-phenoxy)-ethanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-bromo-4-trifluoromethoxy-phenoxy)-ethanone;

(1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(4'-chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

15 4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-isopropyl-benzonitrile;

2-(2-Acetyl-phenoxy)-1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-ethanone;

20 4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-5-isopropyl-2-methyl-benzonitrile;

(1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(4-methoxy-naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

25 4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-ethoxy-benzonitrile;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-4-trifluoromethoxy-phenyl)-propan-1-one;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-phenoxy)-ethanone;

5 [(3aS,6aS)-5-(4'-Chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-(R)-4,5,6,7-tetrahydro-1H-benzotriazol-5-yl-methanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

10 (1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(4'-chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-[2-(tetrahydro-furan-2-yl)-phenoxy]-ethanone;

(1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(4-methoxy-naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

15 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-tert-butyl-phenoxy)-ethanone;

[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-[trans-4-(4-chloro-phenyl)-cyclohexyl]-methanone;

20 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-4-trifluoromethyl-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(2-fluoro-4-trifluoromethyl-phenyl)-propan-1-one;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(2-pyridin-3-ylphenoxy)ethanone;

25 4-[3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-oxopropyl]-2-methyl-5-propan-2-ylbenzonitrile;

4-[3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-oxopropyl]-3-propan-2-ylbenzonitrile;

[(3aR,6aR)-5-[1-(4-chlorophenyl)piperidine-4-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(1H-benzotriazol-5-yl)methanone;

5 [(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(4-propan-2-yloxynaphthalen-2-yl)methanone;

[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(4-propan-2-yloxyquinolin-2-yl)methanone;

10 1-[(3aR,6aR)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[2-fluoro-4-(trifluoromethoxy)phenyl]propan-1-one;

4-[2-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-2-methyl-5-propan-2-ylbenzonitrile;

15 [(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-[1-(2,2,2-trifluoroethoxy)isoquinolin-3-yl]methanone;

1-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(4-bromo-2-tert-butylphenoxy)ethanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(4-bromo-2-tert-butylphenoxy)ethanone;

20 4-[2-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-3-tert-butylbenzonitrile;

4-[2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-3-tert-butylbenzonitrile;

25 [(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-[1-methyl-5-(trifluoromethoxy)indol-2-yl]methanone;

1-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-[4-(trifluoromethoxy)phenoxy]ethanone;

[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(1-ethoxyisoquinolin-3-yl)methanone;

5 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(2-tert-butyl-4-methoxyphenoxy)ethanone;

((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(4-ethoxyquinolin-2-yl)methanone;

10 ((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(4-(2,2,2-trifluoroethoxy)quinolin-2-yl)methanone;

((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(6-cyclobutoxy-5-(trifluoromethyl)pyridin-3-yl)methanone;

((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-bromo-6-(2-methoxyethoxy)pyridin-3-yl)methanone;

15 ((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-bromo-6-(cyclopropylmethoxy)pyridin-3-yl)methanone;

((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-cyclopropyl-6-(2,2,2-trifluoroethoxy)pyridin-3-yl)methanone;

20 ((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(6-(2,2,2-trifluoroethoxy)-5-(trifluoromethyl)pyridin-3-yl)methanone;

(1H-benzotriazol-5-yl)-{(3aS,6aS)-5-[4-(4-chloro-phenyl)-piperidine-1-carbonyl]-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl}-methanone;

(1H-benzotriazol-5-yl)-{(3aS,6aS)-5-[4-(4-chloro-phenyl)-piperazine-1-carbonyl]-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl}-methanone;

25 and pharmaceutically acceptable salts thereof.

Further particular examples of compounds of formula (I) as described herein are selected from

trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

5 trans-3,5-dichlorobenzyl 2-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

10 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one;

15 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-propan-1-one;

20 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

and pharmaceutically acceptable salts thereof.

Also further particular examples of compounds of formula (I) as described herein are selected from

25 trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

trans-3,5-dichlorobenzyl 2-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

5 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

10 1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-propan-1-one;


15 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

and pharmaceutically acceptable salts thereof.

Processes for the manufacture of compounds of formula (I) as described herein are an object of the invention.

20 The preparation of compounds of formula (I) of the present invention may be carried out in sequential or convergent synthetic routes. Syntheses of the invention are shown in the following general schemes. The skills required for carrying out the reactions and purifications of the resulting products are known to those persons skilled in the art. In case a mixture of enantiomers or diastereoisomers is produced during a reaction, these enantiomers or diastereoisomers can be
25 separated by methods described herein or known to the man skilled in the art such as e.g. (chiral) chromatography or crystallization. The substituents and indices used in the following description of the processes have the significance given herein.

Compounds of general formula (I) can be synthesised from amine precursor **1** and appropriate reagents, using methods well known in the art.



For instance, amine **1** is reacted with a suitable chloroformate ester of formula $R^1-O-C(O)Cl$ (**2**), or with an imidazole-1-carboxylate ester of formula (**3A**), or with a succinimidyl carbonate derivative of formula (**3B**), leading to a compound of formula (I) wherein Y is $-OC(O)-$.



The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence or not of a base, e. g., triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.

Chloroformate esters **2** are commercially available or can be synthesised from the corresponding alcohol of formula R^1-OH , by reaction with phosgene or a phosgene equivalent (e. g., diphosgene, triphosgene), as described in the literature.

Imidazole-1-carboxylate esters **3A** are synthesised from the corresponding alcohols of formula R^1-OH , by reaction with 1,1'-carbonyldiimidazole. The reaction is performed at room temperature, in a solvent such as dichloromethane, tetrahydrofuran or acetonitrile. The imidazole-1-carboxylate esters **3A** are typically not isolated but directly reacted with amines **1** as described above.

Succinimidyl carbonate derivatives **3B** are synthesised from the corresponding alcohols of formula R^1-OH , by reaction with N,N'-disuccinimidyl carbonate. The reaction is performed at room temperature, in a solvent such as dichloromethane, tetrahydrofuran, or acetonitrile,

optionally in the presence of a base, e. g., triethylamine. The succinimidyl carbonate derivatives **3B** are typically not isolated but directly reacted with amines **1** as described above.

Alcohols of formula R^1-OH are commercially available or can be produced by methods described herein or known in the art.

- 5 Alternatively, amine **1** is reacted with a suitable N-(chlorocarbonyl)amine of formula $R^1-N(R^7)-C(O)-Cl$ (**4**), or, in the case where R^7 is H, with an isocyanate of formula R^1-NCO (**5**), leading to compounds of formula (I) wherein Y is $-NR^7C(O)-$.

N-(Chlorocarbonyl)amines (**4**) are synthesised from the corresponding amines of formula $R^1-N(R^7)H$ by reaction with phosgene or a phosgene equivalent, as described in the literature.

- 10 Isocyanates **5** are commercially available or can be prepared from the corresponding amines of formula R^1-NH_2 , by reaction with phosgene or a phosgene equivalent (e. g., diphosgene, triphosgene, 1,1'-carbonyldiimidazole), as described in the literature.

- Alternatively, amine **1** is reacted with a suitable carboxylic acid of formula R^1-COOH (**6**) leading to a compound of formula (I), wherein Y is $-C(O)-$. The reaction is performed in the
15 presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium
20 hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between $-40^\circ C$ and $80^\circ C$ in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.




- Amine **1** can also be reacted with suitable acylating reagents such as acyl chlorides of formula R^1-COCl (**7**) to lead to compounds of formula (I) wherein Y is $-C(O)-$. The reaction is
25 performed in a solvent such as dichloromethane, tetrahydrofuran, or N,N-dimethylformamide, in the presence of a base such as triethylamine or 4-methylmorpholine, at temperatures between $0^\circ C$ and $80^\circ C$.

Carboxylic acids (**6**) and acyl halides (**7**) are commercially available or can be prepared as described herein or in the literature.

Alternatively, amine **1** is reacted with a suitable sulfonyl chloride of formula R^1-SO_2Cl (**8**), leading to compounds of formula (I) wherein Y is $-S(O_2)-$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g., triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.

Sulfonyl chlorides (**8**) are commercially available or can be synthesised as described herein or in the literature.

Alternatively, amine **1** is reacted with a suitable chloro-oxadiazole reagent of general formula **9**, or with oxadiazolone reagent **10**, leading to a compound of formula (I), wherein Y is

**9****10**

In the case where compounds of formula (I) are produced from amine **1** and chloro-oxadiazole **9**, the reaction is performed in the presence of a base, e. g., potassium carbonate, triethylamine, or 1,8-diazabicyclo[5.4.0]undec-7-ene, in a solvent such as toluene, ethanol, N,N-dimethylformamide, or 1,4-dioxane at temperatures between 20°C and 150°C.


In the case where compounds of formula (I) are produced from amine **1** and oxadiazolone **10**, the reaction is performed in the presence of a coupling agent, e. g. benzotriazol-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate and a base, e. g., diisopropylethylamine or 4-methylmorpholine, in a solvent such as N,N-dimethylformamide, at temperatures between 20°C and 100°C as described in the literature.



Oxadiazolones **10** are commercially available or can be produced as described in the literature.

-57-

Chloro-oxadiazoles **9** are commercially available or can be produced from the corresponding oxadiazolones, by reaction with a suitable halogenating reagent, e. g. phosphorus oxychloride and/or phosphorus pentachloride, at temperatures between 60°C and 120°C.

Alternatively, amine **1** is reacted with a suitable halo-thiadiazole reagent of general formula **11** (X = Cl or Br), or with thiadiazolethione reagent **12**, leading to compounds of (I)

wherein Y is 

**11****12**


In the case where compounds of formula (I) are produced from amine **1** and halo-thiadiazole **11**, the reaction is performed in the presence of a base, e. g. potassium carbonate, triethylamine or 1,8-diazabicyclo[5.4.0]undec-7-ene, in a solvent such as toluene, ethanol, N,N-dimethylformamide or 1,4-dioxane, at temperatures between 20°C and 150°C.

In the case where compounds of formula (I) are produced from amine **1** and thiadiazolethione **12**, the reaction is performed in a solvent such as ethanol or N,N-dimethylformamide at temperatures between 20°C and 100°C as described in the literature.

Thiadiazolethiones **12** are commercially available or can be produced as described in the literature.

Halo-thiadiazoles **11** are commercially available or can be produced as described in the literature.

Amines of general formula **1** are synthesised from suitably protected precursors **13**.


**13**

Suitable protective groups (PG) are tert-butoxycarbonyl, benzyloxycarbonyl and substituted benzyloxycarbonyl such as 3,5-dichloro benzyloxycarbonyl. The deprotection of intermediates **13** can be performed using methods and reagents known in the art.

For instance, in the case where PG is optionally substituted benzyloxycarbonyl, the deprotection may be performed by hydrogenation at pressures between 1 bar and 100 bar, in the presence of a suitable catalyst such as palladium on activated charcoal, at temperatures between 20°C and 150°C in solvents such as methanol or ethanol.


Alternatively, in the case where PG is tert-butoxycarbonyl, the deprotection may be performed in the presence of a suitable acid, e. g., hydrochloric acid or trifluoroacetic acid, in a solvent such as water, 2-propanol, dichloromethane, or 1,4-dioxane at temperatures between 0°C and 30°C.

Carbamates **13**, wherein A is N are represented by general structure **13A**.



PG is a suitable protective group, e. g., tert-butoxycarbonyl, benzyloxycarbonyl and substituted benzyloxycarbonyl such as 3,5-dichloro benzyloxycarbonyl.

Carbamates **13A** can be produced from amine precursors of general formula **14** by reaction with appropriate reagents, using methods known in the art.



For instance, **14** is reacted with alkylating agents of general formula $X-CR^3R^4-R^2$ (**15**) where X is a leaving group such as Cl, Br, I, or OSO_2CH_3 , leading to **13A**, wherein W is $-CR^3R^4-$. This reaction is performed in a solvent such as tetrahydrofuran or N,N-dimethylformamide, in the presence of a base, e. g. triethylamine or potassium carbonate, at temperatures between 0°C and 100°C.

Alternatively, for compounds of formula **13A**, wherein W is $-\text{CR}^3\text{R}^4-$, R^4 is hydrogen, alkyl or cycloalkyl, and R^3 is H, amine **14** is reacted with aldehydes or ketones of general formula $\text{R}^4-\text{C}(\text{O})-\text{R}^2$ (**16**) in a reductive amination reaction, leading to **13A**. This reaction is performed in the presence of a suitable reducing agent, e. g., sodium borohydride or sodium triacetoxyborohydride, in a solvent such as methanol, acetic acid, tetrahydrofuran, 1,2-dichloroethane or mixtures thereof, at temperatures between 0°C and 50°C.


Alternatively, amine **14** is reacted with a suitable carboxylic acid of formula R^2-COOH (**17**), leading to compounds of formula **13A**, wherein W is $-\text{C}(\text{O})-$. The reaction is performed in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

Alternatively, amine **14** is reacted with a suitable sulfonyl chloride of formula $\text{R}^2-\text{SO}_2\text{Cl}$ (**18**), leading to compounds of formula **13A**, wherein W is $-\text{S}(\text{O}_2)-$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g. triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.

Alternatively, amine **14** is reacted with a suitable N-(chlorocarbonyl)amine of formula $\text{R}^2-\text{N}(\text{R}^6)-\text{C}(\text{O})-\text{Cl}$ (**19**) leading to compounds of formula **13A**, wherein W is $-\text{C}(\text{O})-\text{NR}^6-$, or with an isocyanate of formula R^2-NCO (**20**), leading to compounds of formula **13A**, wherein W is $-\text{C}(\text{O})-\text{NR}^6-$ and R^6 is H.

Alternatively, amine **14** is reacted with phosgene or phosgene equivalent (diphosgene, triphosgene) in the presence of a base (e. g., pyridine, triethylamine) in a solvent such as dichloromethane or tetrahydrofuran, to provide the corresponding N-(chlorocarbonyl)amine of formula **21**, which is then reacted with amine of formula $\text{HN}(\text{R}^6)\text{R}^2$ (**22**), in the presence of a base such as triethylamine or diisopropylethylamine, in a solvent such as dichloromethane,

tetrahydrofuran, or N,N-dimethylformamide, leading to compounds of formula **13A**, wherein W is $-\text{C}(\text{O})-\text{NR}^6-$.




Alternatively, amine **14** is reacted with phosgene or a phosgene equivalent (diphosgene, triphosgene) in the presence of a base (e. g., pyridine, triethylamine), in a solvent such as dichloromethane or tetrahydrofuran, to the corresponding N-(chlorocarbonyl)amine of formula **21**, which is then reacted with amines of formula H-O, H-P, H-Q, H-R, H-T, H-U, H-V, H-X, H-AA or H-AF, in the presence of a base such as triethylamine or diisopropylethylamine, in a solvent such as dichloromethane, tetrahydrofuran, or N,N-dimethylformamide, leading to compounds of formula **13A**, wherein W is $-\text{C}(\text{O})-$ and R^2 is O, P, Q, R, T, U, V, X, AA or AF.

N-(Chlorocarbonyl)amines **19** are synthesised from the corresponding amines **22** by reaction with phosgene or a phosgene equivalent (diphosgene, triphosgene) as described in the literature.

Isocyanates **20** are commercially available or can be prepared from the corresponding amines of formula R^2-NH_2 , by reaction with phosgene or a phosgene equivalent (e. g., diphosgene, triphosgene, 1,1'-carbonyldiimidazole) as described in the literature.


Amines **14**, alkylating agents **15**, aldehydes/ketones **16**, carboxylic acids **17**, sulfonyl chlorides **18**, and amines **22** are commercially available or can be synthesised as described in the literature.

Carbamates **13** wherein A is CR^5 and R^5 is H are represented by general formula **13B**, wherein PG is a suitable protective group, e. g. tert-butoxycarbonyl, benzyloxycarbonyl and substituted benzyloxycarbonyl such as 3,5-dichloro benzyloxycarbonyl.




Compound **13B**, wherein W is $-\text{NR}^6-$, is produced from ketone **23** by reaction with an amine of formula $\text{HN}(\text{R}^6)\text{R}^2$ (**22**) in the presence of a suitable reducing agent, e. g. sodium

borohydride or sodium triacetoxyborohydride, in a solvent such as methanol, acetic acid, tetrahydrofuran, 1,2-dichloroethane, or mixtures thereof, at temperatures between 0°C and 50°C.



Ketones **23** and amines **22** are commercially available or can be prepared as described in the literature.

Compound **13B**, wherein W is –O– or –S–, is produced from alcohol **24** using methods and reagents known in the art.



For instance, alcohol **24** is reacted at room temperature with phenol HO–R² or thiophenol HS–R² in the presence of triphenylphosphine and an dialkylazodicarboxylate, e. g. diisopropylazodicarboxylate or diethylazodicarboxylate, in a solvent such as toluene, dichloromethane, or tetrahydrofuran, leading to **13B**, wherein W is –O– or –S–.

In the case wherein W is –O– or and R² is **B**, compounds of formula **13B** can also be produced from alcohol **24** in a three-step sequence. Therefore, **24** is reacted in step 1 at room temperature with 2,4-dihydroxybenzoic acid alkyl ester in the presence of triphenylphosphine and an dialkylazodicarboxylate, e. g. diisopropylazodicarboxylate or diethylazodicarboxylate, in a solvent such as toluene, dichloromethane or tetrahydrofuran, converting the hydroxy group into a 3-hydroxy-4-(alkoxycarbonyl)-phenyl ether substituent. This is hydrolysed in step 2 to the corresponding 3-hydroxy-4-carboxyphenyl group, using a base such as sodium hydroxide in water and in the presence of co-solvents such as tetrahydrofuran and/or methanol or ethanol. In step 3, the 3-hydroxy-4-carboxyphenyl ether intermediate can be subjected to a Curtius rearrangement as described in the literature, e. g. by using diphenylphosphoryl azide, in a solvent such as toluene, in the presence of a base, e. g., triethylamine, at temperatures between 60°C and 110°C, leading to the corresponding 2-oxo-2,3-dihydro-benzooxazol-6-yl ether **13B**, wherein W is –O– and R² is **B**.


-62-

Alternatively, conversion of alcohol **24** to the corresponding methanesulfonate using methanesulfonyl chloride in the presence of a base, e. g. triethylamine, in a solvent such as dichloromethane or tetrahydrofuran, at temperatures between -20°C and $+30^{\circ}\text{C}$, and treatment of the methanesulfonate intermediate with phenol $\text{HO}-\text{R}^2$ or thiophenol $\text{HS}-\text{R}^2$ in the presence of a base, e. g., potassium carbonate, in a solvent such as N,N-dimethylformamide or acetonitrile, at temperatures between 20°C and 100°C , leads to **13B**, wherein W is $-\text{O}-$ or $-\text{S}-$.


Compound **13B**, wherein W is $-\text{SO}_2-$, is produced from compound **13B**, wherein W is $-\text{S}-$ by oxidation with a suitable reagent, e. g., hydrogen peroxide or 3-chloroperbenzoic acid, in a solvent such as formic acid, acetic acid, or dichloromethane, at temperatures between 0°C and 50°C .

Alcohols **24** are produced from ketones **23** using a suitable reducing agent, e. g., sodium borohydride, in a solvent such as methanol, at temperatures between 0°C and 50°C .

Carbamates **13** wherein A is CR^5 , R^5 is H, and W is $-\text{C}(\text{O})-\text{N}(\text{R}^6)-$ are represented by general formula **13C**, wherein R^{12} is $\text{N}(\text{R}^6)\text{R}^2$, **O**, **P**, **Q**, **R**, **T**, **U**, **V**, **X**, **AA** or **AF**.



Amide **13C** is produced from carboxylic acid **25** by coupling reaction with an amine of formula $\text{HN}(\text{R}^6)\text{R}^2$ (**22**), **H-O**, **H-P**, **H-Q**, **H-R**, **H-T**, **H-U**, **H-V**, **H-X**, **H-AA** or **H-AF**.




The reaction is performed in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and

mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

Carboxylic acids **25** are commercially available or can be produced as described in the literature.

Compounds of formula (I), wherein A is N can be produced from amine precursors of general formula **26** by reaction with appropriate reagents, using methods known in the art.



For instance, an amine of formula **26** is reacted with alkylating agents of general formula $\text{X}-\text{CR}^3\text{R}^4-\text{R}^2$ (**15**) where X is a leaving group such as Cl, Br, I, or OSO_2CH_3 , leading to compounds of formula (I), wherein A is N and W is $-\text{CR}^3\text{R}^4-$. This reaction is performed in a solvent such as tetrahydrofuran or N,N-dimethylformamide, in the presence of a base, e. g., triethylamine or potassium carbonate, at temperatures between 0°C and 100°C .

Alternatively, an amine of formula **26** is reacted with aldehydes or ketones of general formula $\text{R}^4-\text{C}(\text{O})-\text{R}^2$ (**16**) in a reductive amination reaction, leading to compounds of formula (I) wherein A is N, W is $-\text{CR}^3\text{R}^4-$, R^4 is hydrogen, alkyl or cycloalkyl, and R^3 is H. This reaction is performed in the presence of a suitable reducing agent, e. g. sodium borohydride or sodium triacetoxyborohydride, in a solvent such as methanol, acetic acid, tetrahydrofuran, 1,2-dichloroethane or mixtures thereof, at temperatures between 0°C and 50°C .

Alternatively, amine **26** is reacted with a suitable carboxylic acid of formula R^2-COOH (**17**), leading to compounds of formula (I) wherein A is N and W is $-\text{C}(\text{O})-$. The reaction is performed in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran,


N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

Alternatively, amine **26** is reacted with a suitable sulfonyl chloride of formula

- 5 R^2-SO_2Cl (**18**), leading to (I) wherein A is N and W is $-S(O_2)-$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g. triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.


- 10 Alternatively, amine of formula **26** is reacted with a suitable N-(chlorocarbonyl)amine of formula $R^2-N(R^6)-C(O)-Cl$ (**19**) leading to compounds of formula (I), wherein A is N and W is $C(O)-NR^6$, or with isocyanate R^2-NCO (**20**), leading to compounds of formula (I), wherein A is N, W is $-C(O)-NR^6-$ and R^6 is H.

- Alternatively, amine **26** is reacted with phosgene or a phosgene equivalent (diphosgene, triphosgene) in the presence of a base (e. g., pyridine, triethylamine), in a solvent such as dichloromethane or tetrahydrofuran, to the corresponding N-(chlorocarbonyl)amine of formula **27**, which is then reacted with an amine of formula H-O, H-P, H-Q, H-R, H-T, H-U, H-V, H-X, H-AA or H-AF in the presence of a base such as triethylamine or diisopropylethylamine, in a solvent such as dichloromethane, tetrahydrofuran, or N,N-dimethylformamide, leading to
- 20 compounds of formula (I), wherein A is N, W is $-C(O)-$ and R^2 is O, P, Q, R, T, U, V, X, AA or AF.


**27**

- Amines **26** can be synthesised from their tert-butyl carbamate derivatives of formula **28** by carbamate deprotection. The deprotection may be performed in the presence of a suitable acid, e. g, hydrochloric acid or trifluoroacetic acid, in a solvent such as water, 2-propanol, dichloromethane, or 1,4-dioxane, at temperatures between 0°C and 30°C.
- 25

-65-



tert-Butyl carbamates **28** can be synthesised from amine precursors of formula **29** and appropriate reagents, using methods well known in the art.



For instance, an amine of formula **29** is reacted with a suitable chloroformate ester of formula $R^1-O-C(O)-Cl$ (**2**), or with an imidazole-1-carboxylate ester of formula (**3A**) or with a succinimidyl carbonate derivative of formula (**3B**), leading to compounds of formula **28**, wherein Y is $-OC(O)-$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence or not of a base, e. g. triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.


Alternatively, an amine of formula **29** is reacted with a suitable N-(chlorocarbonyl)amine of formula $R^1-N(R^7)-C(O)-Cl$ (**4**) leading to compounds of formula **28**, wherein Y is $-NR^7C(O)-$, or with an isocyanate of formula R^1-NCO (**5**) leading to compounds of formula **28**, wherein Y is $-NR^7C(O)-$ and R^7 is H.

Alternatively, amine **29** is reacted with a suitable carboxylic acid of formula R^1-COOH (**6**) leading to compounds of formula **28**, wherein Y is $-C(O)-$. The reaction is performed in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

Amine **29** can also be reacted with suitable acylating reagents, such as acyl chlorides of formula $R^1\text{-COCl}$ (**7**) to provide compounds of formula **28**, wherein Y is -C(O)- . The reaction is performed in a solvent such as dichloromethane, tetrahydrofuran, or N,N-dimethylformamide, in the presence of a base such as triethylamine or 4-methylmorpholine, at temperatures between 0°C and 80°C.

Alternatively, amine **29** is reacted with a suitable sulfonyl chloride, of formula $R^1\text{-SO}_2\text{Cl}$ (**8**), leading to compounds of formula **28**, wherein Y is $\text{-S(O}_2\text{)-}$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g. triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.


Alternatively, amine **29** is reacted with a suitable chloro-oxadiazole reagent of general formula **9**, or with oxadiazolone reagent **10**, leading to compounds of formula **28**, wherein Y is



In the case where **28** is produced from amine **29** and chloro-oxadiazole **9**, the reaction is performed in the presence of a base, e. g. potassium carbonate, triethylamine, or 1,8-diazabicyclo[5.4.0]undec-7-ene, in a solvent such as toluene, ethanol, N,N-dimethylformamide, or 1,4-dioxane, at temperatures between 20°C and 150°C.

In the case where **28** is produced from amine **29** and oxadiazolone **10**, the reaction is performed in the presence of a coupling agent, e. g., benzotriazol-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate, and a base, e. g. diisopropylethylamine or 4-methylmorpholine, in a solvent such as N,N-dimethylformamide, at temperatures between 20°C and 100°C, as described in the literature).

Alternatively, amine **29** is reacted with a suitable halo-thiadiazole reagent of general formula **11** (X is Cl or Br), or with thiadiazolethione reagent **12**, leading to compounds of



formula **28**, wherein Y is


In the case where **28** is produced from amine **29** and halo-thiadiazole **11**, the reaction is performed in the presence of a base, e. g. potassium carbonate, triethylamine, or 1,8-diazabicyclo[5.4.0]undec-7-ene, in a solvent such as toluene, ethanol, N,N-dimethylformamide, or 1,4-dioxane, at temperatures between 20°C and 150°C.

- 5 In the case where **28** is produced from amine **29** and thiadiazolethione **12**, the reaction is performed in a solvent such as ethanol or N,N-dimethylformamide, at temperatures between 20°C and 100°C, as described in the literature.

- Alternatively, amine **29** is acylated with a haloalkanoyl halide, e. g., bromoacetyl chloride, in the presence of a base, e. g. triethylamine, in a solvent such as dichloromethane or
10 tetrahydrofuran, at temperatures between -78°C and +20°C, leading to the corresponding haloalkanamide intermediate, which in the presence of a base, e. g. potassium carbonate or caesium carbonate, in a solvent such as N,N-dimethylformamide undergoes a nucleophilic substitution reaction with a substituted phenol, leading to compounds of formula **28**, wherein Y is -C(O)- and R¹ is substituted phenoxyalkyl.

- 15 Amines of formula **29** are commercially available or can be produced as described in the literature.


- Compounds of formula (I), wherein A is CR⁵ and W is -C(O)-N(R⁶)- can be produced from carboxylic acid precursors of general formula **30** by reaction with appropriate amine reagents of general formula HN(R⁶)R², H-O, H-P, H-Q, H-R, H-T, H-U, H-V, H-X, H-AA or
20 H-AF using methods known in the art.




- For instance, this reaction is performed in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-
25 phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base

such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

Carboxylic acids **30** can be produced from the corresponding ester precursors **31**, wherein R^a is lower alkyl, e. g. methyl or ethyl, using methods and reagents known in the art. For instance, the reaction is performed in the presence of a base, e. g., potassium hydroxide, sodium hydroxide, or lithium hydroxide, in solvents such as water, methanol, ethanol, tetrahydrofuran, or mixtures thereof, at temperatures between 20°C and 100°C.



Compounds of formula **31** can be synthesised from amine precursors of formula **32** and appropriate reagents, using methods well known in the art.



For instance, an amine of formula **32** is reacted with a suitable chloroformate ester of formula $R^1-O-C(O)-Cl$ (**2**), or with an imidazole-1-carboxylate ester of formula **3**, leading to compounds of formula **31**, wherein Y is $-OC(O)-$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g. triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.

Alternatively, an amine of formula **32** is reacted with a suitable N-(chlorocarbonyl)amine of formula $R^1-N(R^7)-C(O)-Cl$ (**4**) leading to compounds of formula **31**, wherein Y is $-NR^7C(O)-$, or with an isocyanate of formula R^1-NCO (**5**) leading to compounds of formula **31**, wherein Y is $-NR^7C(O)-$ and R^7 is H.


Alternatively, amine **32** is reacted with a suitable carboxylic acid of formula R^1-COOH (**6**) leading to compounds of formula **31**, wherein Y is $-C(O)-$. The reaction is performed in the presence of a coupling agent such as 1,1'-carbonyldiimidazole, N,N'-dicyclohexylcarbodiimide,

1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride, O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate or bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, in aprotic solvents such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, N-methylpyrrolidinone and mixtures thereof at temperatures between -40°C and 80°C in the presence or absence of a base such as triethylamine, diisopropylethylamine, 4-methylmorpholine and/or 4-(dimethylamino)pyridine.

Amine **32** can also be reacted with suitable acylating reagents, such as acyl chlorides of formula R^1-COCl (**7**) to lead to compounds of formula **31**, wherein Y is $-C(O)-$. The reaction is performed in a solvent such as dichloromethane, tetrahydrofuran, or N,N-dimethylformamide, in the presence of a base such as triethylamine or 4-methylmorpholine, at temperatures between 0°C and 80°C.

Alternatively, amine **32** is reacted with a suitable sulfonyl chloride of formula R^1-SO_2Cl (**8**), leading to compounds of formula **31**, wherein Y is $-S(O_2)-$. The reaction is performed in a suitable solvent such as dichloromethane, tetrahydrofuran, N,N-dimethylformamide, acetonitrile, acetone, water, or mixtures thereof, in the presence of a base, e. g. triethylamine, diisopropylethylamine, pyridine, potassium hydrogencarbonate, potassium carbonate, at temperatures between 0°C and the boiling point of the solvent or solvent mixture.

Alternatively, amine **32** is reacted with a suitable chloro-oxadiazole reagent of general formula **9**, or with oxadiazolone reagent **10**, leading to compounds of formula **31**, wherein Y is




In the case where **31** is produced from amine **32** and chloro-oxadiazole **9**, the reaction is performed in the presence of a base, e. g. potassium carbonate, triethylamine, or 1,8-diazabicyclo[5.4.0]undec-7-ene, in a solvent such as toluene, ethanol, N,N-dimethylformamide, or 1,4-dioxane, at temperatures between 20°C and 150°C.

In the case where **31** is produced from amine **32** and oxadiazolone **10**, the reaction is performed in the presence of a coupling agent, e. g. benzotriazol-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate, and a base, e. g. diisopropylethylamine or 4-

methylmorpholine, in a solvent such as N,N-dimethylformamide, at temperatures between 20°C and 100°C, as described in the literature.

Alternatively, amine **32** is reacted with a suitable halo-thiadiazole reagent of general formula **11** (X is Cl or Br), or with thiadiazolethione reagent **12**, leading to compounds of




5 formula **31**, wherein Y is

In the case where **31** is produced from amine **32** and halo-thiadiazole **11**, the reaction is performed in the presence of a base, e. g. potassium carbonate, triethylamine, or 1,8-diazabicyclo[5.4.0]undec-7-ene, in a solvent such as toluene, ethanol, N,N-dimethylformamide or 1,4-dioxane, at temperatures between 20°C and 150°C.

10 In the case where **31** is produced from amine **32** and thiadiazolethione **12**, the reaction is performed in a solvent such as ethanol or N,N-dimethylformamide, at temperatures between 20°C and 100°C, as described in the literature.

Amines of general formula **32** are synthesised from suitably protected precursors **33**.




33

15 Suitable protective groups (PG) are tert-butoxycarbonyl or benzyloxycarbonyl. The deprotection of intermediates **33** can be performed using methods and reagents known in the art.

For instance, in the case where PG is benzyloxycarbonyl, the deprotection may be performed by hydrogenation at pressures between 1 bar and 100 bar, in the presence of a suitable catalyst such as palladium on activated charcoal, at temperatures between 20°C and 150°C, in
20 solvents such as methanol or ethanol.

Alternatively, in the case where PG is tert-butoxycarbonyl, the deprotection may be performed in the presence of a suitable acid, e. g. hydrochloric acid or trifluoroacetic acid, in a solvent such as water, 2-propanol, dichloromethane, or 1,4-dioxane, at temperatures between 0°C and 30°C.


-71-



Substituents R^5 may be introduced starting from ester precursor **34**, using suitable reagents and methods known in the art. For instance, **34** is reacted with alkylating agents of general formula R^5-X wherein R^5 is alkyl or cycloalkyl and X is a leaving group such as Cl, Br, I, or OSO_2CH_3 , leading to **33**, wherein R^5 is alkyl or cycloalkyl. This reaction is performed in the presence of a suitable base, e. g., sodium hydride, lithium diisopropylamide, lithium bis(trimethylsilyl)amide, or lithium pyrrolidide, in a solvent such as tetrahydrofuran or toluene, at temperatures between $-78^\circ C$ and $+50^\circ C$.

Esters **34**, wherein R^a is methyl or ethyl, are produced from carboxylic acids **25**, using methods and reagents known in the art. For instance, **25** alkylated with methyl iodide or ethyl bromide, in the presence of a base, e. g., potassium carbonate, in a solvent such as N,N-dimethylformamide, at $-20^\circ C$ and $+30^\circ C$, leading to the methyl or ethyl ester **34**, respectively.

Also an embodiment of the present invention is a process to prepare a compound of formula (I) as defined above comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III);



wherein R^1 , R^2 , A, W, m, n, p and q are as defined above, Y is $-OC(O)-$.

In particular, in the presence of a coupling agent such as O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate, in a solvent such as N,N-dimethylformamide, in the presence of a base such as 4-methylmorpholine and at a temperature comprised between $-78^\circ C$ and reflux, particularly between $-10^\circ C$ and room temperature.

Also an object of the present invention is a compound according to formula (I) as described herein for use as a therapeutically active substance.

Likewise an object of the present invention is a pharmaceutical composition comprising a compound according to formula (I) as described herein and a therapeutically inert carrier.

5 An object of the invention is the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, conditions of the respiratory system, vascular and cardiovascular conditions, fibrotic diseases, cancer, ocular conditions, metabolic conditions, cholestatic and other forms of chronic pruritus and acute and chronic organ transplant rejection.

10 Renal conditions include, but are not limited to, acute kidney injury and chronic renal disease with and without proteinuria including end-stage renal disease (ESRD). In more detail, this includes decreased creatinine clearance and decreased glomerular filtration rate, micro-albuminuria, albuminuria and proteinuria, glomerulosclerosis with expansion of reticulated mesangial matrix with or without significant hypercellularity (particularly diabetic nephropathy
15 and amyloidosis), focal thrombosis of glomerular capillaries (particularly thrombotic microangiopathies), global fibrinoid necrosis, ischemic lesions, malignant nephrosclerosis (such as ischemic retraction, reduced renal blood flow and renal arteriopathy), swelling and proliferation of intracapillary (endothelial and mesangial) and/or extracapillary cells (crescents) like in glomerular nephritis entities, focal segmental glomerular sclerosis, IgA nephropathy,
20 vasculitides / systemic diseases as well as acute and chronic kidney transplant rejection.

Liver conditions include, but are not limited to, liver cirrhosis, hepatic congestion, cholestatic liver disease including pruritus, nonalcoholic steatohepatitis and acute and chronic liver transplant rejection.

25 Inflammatory conditions include, but are not limited to, arthritis, osteoarthritis, multiple sclerosis, systemic lupus erythematoses, inflammatory bowel disease, abnormal evacuation disorder and the like as well as inflammatory airways diseases such as idiopathic pulmonary fibrosis (IPF), chronic obstructive pulmonary disease (COPD) or chronic asthma bronchiale.

30 Further conditions of the respiratory system include, but are not limited to, other diffuse parenchymal lung diseases of different etiologies including iatrogenic drug-induced fibrosis, occupational and/or environmental induced fibrosis, systemic diseases and vasculitides,

granulomatous diseases (sarcoidosis, hypersensitivity pneumonia), collagen vascular disease, alveolar proteinosis, Langerhans cell granulomatosis, lymphangioleiomyomatosis, inherited diseases (Hermansky-Pudlak Syndrome, tuberous sclerosis, neurofibromatosis, metabolic storage disorders, familial interstitial lung disease), radiation induced fibrosis, silicosis, asbestos induced pulmonary fibrosis or acute respiratory distress syndrome (ARDS).

Conditions of the nervous system include, but are not limited to, neuropathic pain, schizophrenia, neuro-inflammation (e.g. astrogliosis), peripheral and/or autonomic (diabetic) neuropathies and the like.

Vascular conditions include, but are not limited to, atherosclerosis, thrombotic vascular disease as well as thrombotic microangiopathies, proliferative arteriopathy (such as swollen myointimal cells surrounded by mucinous extracellular matrix and nodular thickening), atherosclerosis, decreased vascular compliance (such as stiffness, reduced ventricular compliance and reduced vascular compliance), endothelial dysfunction and the like.

Cardiovascular conditions include, but are not limited to, acute coronary syndrome, coronary heart disease, myocardial infarction, arterial and pulmonary hypertension, cardiac arrhythmia such as atrial fibrillation, stroke and other vascular damage.

Fibrotic diseases include, but are not limited to myocardial and vascular fibrosis, renal fibrosis, liver fibrosis, pulmonary fibrosis, skin fibrosis, scleroderma and encapsulating peritonitis.

In a particular embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis of organ or skin fibrosis.

In another embodiment, the fibrotic disease is renal tubulo-interstitial fibrosis or glomerulosclerosis.

In another embodiment, the fibrotic disease is non-alcoholic liver steatosis, liver fibrosis or liver cirrhosis.

In another embodiment, the fibrotic disease is idiopathic pulmonary fibrosis.

Cancer and cancer metastasis include, but are not limited to, breast cancer, ovarian cancer, lung cancer, prostate cancer, mesothelioma, glioma, hepatic carcinoma, gastrointestinal cancers and progression and metastatic aggressiveness thereof.

5 Ocular conditions include, but are not limited to, proliferative and non-proliferative (diabetic) retinopathy, dry and wet age-related macular degeneration (AMD), macular edema, central arterial /venous occlusion, traumatic injury, glaucoma and the like.

Metabolic conditions include, but are not limited to, obesity and diabetes.

10 In another embodiment, the compounds of formula (I) or their pharmaceutically acceptable salts and esters can be used for the treatment or prophylaxis of cholestatic or non-cholestatic chronic pruritus.

The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, fibrotic diseases and acute and chronic organ transplant rejection.

15 The present invention also relates to the use of a compound according to formula (I) as described herein for the treatment or prophylaxis of renal conditions, liver conditions and fibrotic diseases.

20 A particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, fibrotic diseases and acute and chronic organ transplant rejection.

A particular embodiment of the present invention is a compound according to formula (I) as described herein for the treatment or prophylaxis of renal conditions, liver conditions and fibrotic diseases.

25 The present invention also relates to the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, fibrotic diseases and acute and chronic organ transplant rejection.

The present invention also relates to the use of a compound according to formula (I) as described herein for the preparation of a medicament for the treatment or prophylaxis of renal conditions, liver conditions and fibrotic diseases.

5 Also an object of the invention is a method for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, fibrotic diseases and acute and chronic organ transplant rejection, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

10 Also an object of the invention is a method for the treatment or prophylaxis of renal conditions, liver conditions and fibrotic diseases, which method comprises administering an effective amount of a compound according to formula (I) as described herein.

In a particular embodiment, the renal condition is selected from the group consisting of acute kidney injury, chronic kidney disease, diabetic nephropathy, acute kidney transplant rejection and chronic allograft nephropathy.

In another particular embodiment, the renal condition is acute kidney injury.

15 In another particular embodiment, the renal condition is chronic kidney disease.

In a further particular embodiment, the renal condition is diabetic nephropathy.

In another particular embodiment, the renal condition is acute kidney transplant rejection.

In another particular embodiment, the renal condition is chronic allograft nephropathy.

20 In a particular embodiment, the liver condition is acute and chronic liver transplant rejection

In a particular embodiment, the inflammatory condition is arthritis.

In a particular embodiment, the condition of the nervous system is neuropathic pain.

In another embodiment, the fibrotic disease is encapsulating peritonitis.

In another embodiment, the fibrotic disease is idiopathic pulmonary fibrosis.

Also an embodiment of the present invention are compounds of formula (I) as described herein, when manufactured according to any one of the described processes.

PRODUCTION OF HUMAN FULL LENGTH ATX, WITH AND WITHOUT HIS TAG

Autotaxin (ATX - ENPP2) cloning: cDNA was prepared from commercial human hematopoietic cells total RNA and used as template in overlapping PCR to generate a full length human ENPP2 ORF with or without a 3'-6xHis tag. These full length inserts were cloned into the pcDNA3.1V5-His TOPO (Invitrogen) vector. The DNA sequences of several single clones were verified. The DNA from a correct full length clone was used to transfect Hek293 cells for verification of protein expression. The sequence of the encoded ENPP2 conforms to Swissprot entry Q13822, with or without the additional C-terminal 6xHis tag.

ATX Fermentation: Recombinant protein was produced by large-scale transient transfection in 20 L controlled stirred tank bioreactors (Sartorius). During cell growth and transfection, temperature, stirrer speed, pH and dissolved oxygen concentration were maintained at 37°C, 120 rpm, 7.1 and 30% DO, respectively. FreeStyle 293-F cells (Invitrogen) were cultivated in suspension in FreeStyle 293 medium (Invitrogen) and transfected at ca. $1\text{--}1.5 \times 10^6$ cells/mL with above plasmid DNAs using X-tremeGENE Ro-1539 (commercial product, Roche Diagnostics) as complexing agent. Cells were fed a concentrated nutrient solution (J Immunol Methods 194 (1996), 19, 1-199 (page 193)) and induced by sodium butyrate (2 mM) at 72 h post-transfection and harvested at 96 h post-transfection. Expression was analyzed by Western Blot, enzymatic assay and/or analytical IMAC chromatography. After cooling the cell suspension to 4°C in a flow-through heat exchanger, cell separation and sterile filtration of supernatant was performed by filtration through Zeta Plus 60M02 E16 (Cuno) and Sartopore 2 XLG (Sartorius) filter units. The supernatant was stored at 4°C prior to purification.

ATX Purification: 20 liter of culture supernatant were conditioned for ultrafiltration by adding Brij 35 to a final concentration of 0.02% and by adjusting the pH to 7.0 using 1 M HCl. Then the supernatant was first microfiltered through a 0.2 μ m Ultracel-Pilot Open Channel PES filter (Whatman) and afterwards concentrated to 1 liter through an Ultracel-Pilot Screen Channel PES

-77-

filter with 30 kDa MWCO (Whatman). Prior to IMAC chromatography, NiSO₄ was added to a final concentration of 1 mM. The cleared supernatant was then applied to a HisTrap column (GE Healthcare) previously equilibrated in 50 mM Na₂HPO₄ pH 7.0, 0.5 M NaCl, 10% glycerol, 0.3% CHAPS, 0.02% NaN₃. The column was washed stepwise with the same buffer containing
5 20 mM, 40 mM and 50 mM imidazole, respectively. The protein was subsequently eluted using a linear gradient to 0.5 M imidazole in 15 column volumes. ATX containing fractions were pooled and concentrated using an Amicon cell equipped with a 30 kDa PES filter membrane. The protein was further purified by size exclusion chromatography on Superdex S-200 prep grade (XK 26/100) (GE Healthcare) in 20 mM BICINE pH 8.5, 0.15 M NaCl, 10% glycerol,
10 0.3% CHAPS, 0.02% NaN₃. Final yield of protein after purification was 5-10 mg ATX per liter of culture supernatant. The protein was stored at -80°C.

HUMAN ATX ENZYME INHIBITION ASSAY

ATX inhibition was measured by a fluorescence quenching assay using a specifically labeled substrate analogue (MR121 substrate). To obtain this MR121 substrate, BOC and TBS protected
15 6-amino-hexanoic acid (R)-3-({2-[3-(2-{2-[2-(2-amino-ethoxy)-ethoxy]-ethoxy}-ethoxy)-propionylamino]-ethoxy}-hydroxy-phosphoryloxy)-2-hydroxy-propyl ester (Ferguson et al., Org Lett 2006, 8 (10), 2023) was labeled with MR121 fluorophore (CAS 185308-24-1, 1-(3-carboxypropyl)-11-ethyl-1,2,3,4,8,9,10,11-octahydro-dipyrido[3,2-b:2',3'-i]phenoxazin-13-ium) on the free amine of the ethanolamine side and then, after deprotection, subsequently with
20 tryptophan on the side of the aminohexanoic acid.

Assay working solutions were made as follows:

Assay buffer (50 mM Tris-HCl, 140 mM NaCl, 5 mM KCl, 1 mM CaCl₂, 1 mM MgCl₂, 0.01% Triton-X-100, pH 8.0;

ATX solution: ATX (human His-tagged) stock solution (1.08 mg/mL in 20mM bicine, pH 8.5,
25 0.15 M NaCl, 10% glycerol, 0.3% CHAPS, 0.02% NaN₃), diluted to 1.4 – 2.5x final concentration in assay buffer;

MR121 substrate solution: MR121 substrate stock solution (800 µM MR121 substrate in DMSO), diluted to 2 – 5x final concentration in assay buffer.

Test compounds (10 mM stock in DMSO, 8 µL) were obtained in 384 well sample plates
30 (Corning Costar #3655) and diluted with 8 µL DMSO. Row-wise serial dilutions were made by transferring 8 µL cpd solution to the next row up to row O. The compound and control solutions

were mixed five times and 2 μL were transferred to 384 well assay plates (Corning Costar # 3702). Then, 15 μL of 41.7 nM ATX solution was added (30 nM final concentration), mixed five times and then incubated for 15 minutes at 30°C. 10 μL of MR121 substrate solution was added (1 μM final concentration), mixed 30 times and then incubated for 15 minutes at 30 °C.

- 5 Fluorescence was then measured every 2 minutes for 1 hour (Perkin Elmer plate: vision multimode reader); light intensity: 2.5%; exp. time: 1.4 sec, Filter: Fluo_630/690 nm) and IC_{50} values were calculated from these readouts.

Example	IC50 (μM)
1	0.008
1.01	0.035
1.02	0.077
1.03	0.025
1.04	0.127
1.05	0.09
1.06	0.255
1.07	1.034
1.08	0.503
1.09	0.009
1.10	0.078
1.11	0.004
1.12	9.26
1.13	0.226
1.14	0.81
1.15	0.001
1.16	0.382
1.17	0.651

Example	IC50 (μM)
1.18	0.01
1.19	0.012
1.20	0.083
1.21	0.005
1.22	0.009
2	0.01
2.01	0.005
2.02	0.006
3	0.036
3.01	1.093
4	0.008
4.01	0.041
4.02	0.011
4.03	0.011
4.04	0.046
4.05	0.011
4.06	0.012
4.07	0.016

Example	IC50 (μM)
4.08	0.047
4.09	0.049
4.10	0.004
4.11	0.007
4.12	1.476
4.13	2.907
4.14	4.194
4.15	17.63
4.16	0.398
4.17	0.799
4.18	5.617
4.19	1.583
4.20	0.722
4.21	3.48
4.22	1.381
4.23	0.017
4.24	0.056
4.25	0.122

Example	IC50 (μM)
4.26	0.051
4.27	0.004
4.28	0.123
4.29	0.643
4.30	0.05
4.31	1.817
4.32	3.854
4.33	0.036
4.34	0.019
4.35	0.98
4.36	0.009
5	0.248
6	10.42
6.01	0.077
6.02	2.809
6.03	0.531
6.04	0.282
6.05	0.179
6.06	0.008
7	0.022
8	0.108
8.01	0.002
8.02	0.186
8.03	0.911
8.04	0.192

Example	IC50 (μM)
8.05	0.149
8.06	0.192
8.07	1.845
9	0.709
9.01	0.038
9.02	1.6
9.03	0.042
9.04	0.713
9.05	0.406
9.06	0.247
9.07	0.074
9.08	3.402
9.09	0.162
10A	0.036
10B	0.118
11A	0.022
11B	0.009
12A	0.011
12B	0.005
13	0.057
14	0.926
15	0.026
15.01	1.615
15.02	0.096
15.03	0.362

Example	IC50 (μM)
15.04	2.019
15.05	0.015
15.06	0.012
15.07	0.021
15.08	0.977
15.09	0.042
15.10	0.034
15.11	0.013
15.12	0.006
15.13	0.114
15.14	0.062
15.15	0.156
15.16	5.786
15.17	3.953
15.18	0.02
15.19	0.009
15.20	0.049
15.21	0.005
15.22	0.053
15.23	0.032
15.24	0.01
15.25	0.044
15.26	0.008
15.27	0.002
15.28	4.42

Example	IC50 (μM)
15.29	0.034
15.30	0.014
15.31	0.038
15.32	0.019
15.33	0.328
15.34	0.003
15.35	0.031
15.36	0.003
15.37	0.093
15.38	0.05
15.39	0.005
15.40	0.005
15.41	0.017
15.42	0.006
15.43	0.13
15.44	0.006
15.45	0.039
15.46	0.018
15.47	0.096
15.48	0.006
15.49	0.016
15.50	0.006
15.51	0.006
15.52	0.003
15.53	0.007

Example	IC50 (μM)
15.54	0.28
15.55	0.478
15.56	0.655
15.57	0.494
15.58	0.005
15.59	0.011
15.60	1.037
15.61	0.054
15.62	0.068
15.63	0.012
15.64	0.005
15.65	0.003
15.66	0.013
15.67	0.014
15.68	0.171
15.69	0.008
15.70	0.007
15.71	1.365
15.72	0.015
15.73	0.116
15.74	0.012
15.75	0.005
15.76	0.013
15.77	0.008
15.78	0.009

Example	IC50 (μM)
15.79	0.135
15.80	0.06
15.81	0.019
15.82	1.995
15.83	6.859
15.84	0.012
16	0.015
17	0.004
17.01	0.085
17.02	0.101
17.03	0.016
17.04	0.008
17.05	0.006
17.06	0.01
17.07	0.043
17.08	0.016
17.09	0.058
17.10	0.021
17.11	0.028
17.12	0.032
17.13	0.031
17.14	0.464
18	6.735
1.23	0.058
1.24	0.013

Example	IC50 (μM)
1.25	0.114
1.26	0.009
1.27	0.008
1.28	0.006
1.29	0.008
1.30	0.02
1.31	0.021
1.32	0.04
1.33	0.021
1.34	0.013
1.35	0.01
1.36	7.455
1.37	0.025
1.38	0.01
1.39	0.007
1.40	0.007
1.41	0.002
1.42	0.007
1.43	0.014
1.44	0.003
1.45	0.007
1.46	0.002
1.47	0.002
1.48	0.0075
1.49	0.218

Example	IC50 (μM)
1.50	0.017
1.51	0.041
1.52	0.016
1.53	0.052
1.54	0.022
1.55	0.014
1.56	0.2
1.57	0.025
1.58	0.057
1.59	0.006
1.60	0.012
1.61	0.021
2.03	0.005
2.04	0.008
4.37	0.132
4.38	0.024
4.39	0.006
4.40	0.016
4.41	0.009
4.42	0.024
4.43	0.004
4.44	0.006
4.45	0.007
4.46	0.005
4.47	0.004

Example	IC50 (μM)
4.48	0.003
4.49	0.003
4.50	0.006
4.51	0.002
4.52	0.01
4.53	0.005
4.54	0.014
4.55	0.005
4.56	0.009
4.57	0.011
4.58	0.0052
4.59	0.009
4.60	0.005
4.61	0.239
4.62	0.012
4.63	0.037
4.64	0.019
4.65	0.014
4.66	0.04
4.67	0.008
6.07	0.207
9.10	0.007
15.85	0.029
15.86	0.019
15.87	0.103

Example	IC ₅₀ (μM)
15.88	0.027
15.89	0.007
15.90	0.007
15.91	0.012
15.92	0.249
15.93	0.921
15.94	1.018
15.95	1.896
15.96	0.074
15.97	0.009
15.98	0.013
15.99	0.012
17.15	0.047
17.16	0.309
17.17	0.004
17.18	0.014
17.19	0.012
17.20	0.002
19	0.007
19.01	0.012
19.02	0.007
19.03	0.008

Example	IC ₅₀ (μM)
19.04	1.708
19.05	0.005
19.06	0.25
19.07	0.024
19.08	0.036
19.09	0.01
19.10	0.006
19.11	0.008
19.12	0.01
19.13	0.01
19.14	0.273
19.15	0.02
19.16	0.031
19.17	0.01
19.18	0.011
19.19	0.005
19.20	0.459
19.21	0.003
19.22	0.002
19.23	0.025
19.24	0.007
19.25	0.015

Example	IC ₅₀ (μM)
19.26	0.005
19.27	0.006
19.28	0.002
19.29	0.001
19.30	0.005
19.31	0.002
19.32	0.002
19.33	0.006
19.34	0.003
19.35	0.002
19.36	0.005
19.37	0.015
19.38	0.005
19.39	0.007
19.40	0.1
19.41	0.685
19.42	0.01
19.43	0.014
20	0.029
20.01	0.305

Compounds of formula (I) and their pharmaceutically acceptable salts or esters thereof as described herein have IC₅₀ values between 0.00001 μM and 1000 μM, particular compounds

have IC₅₀ values between 0.0005 μM and 500 μM, further particular compounds have IC₅₀ values between 0.0005 μM and 50 μM, more particular compounds have IC₅₀ values between 0.0005 μM and 5 μM. These results have been obtained by using the enzymatic assay described above.

5 The compounds of formula (I) and their pharmaceutically acceptable salts can be used as medicaments (e.g. in the form of pharmaceutical preparations). The pharmaceutical preparations can be administered internally, such as orally (e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatin capsules, solutions, emulsions or suspensions), nasally (e.g. in the form of nasal sprays) or rectally (e.g. in the form of suppositories). However, the administration can also
10 be effected parenterally, such as intramuscularly or intravenously (e.g. in the form of injection solutions).

 The compounds of formula (I) and their pharmaceutically acceptable salts can be processed with pharmaceutically inert, inorganic or organic adjuvants for the production of tablets, coated tablets, dragées and hard gelatin capsules. Lactose, corn starch or derivatives thereof, talc, stearic
15 acid or its salts etc. can be used, for example, as such adjuvants for tablets, dragées and hard gelatin capsules.

 Suitable adjuvants for soft gelatin capsules, are, for example, vegetable oils, waxes, fats, semi-solid substances and liquid polyols, etc.

 Suitable adjuvants for the production of solutions and syrups are, for example, water,
20 polyols, saccharose, invert sugar, glucose, etc.

 Suitable adjuvants for injection solutions are, for example, water, alcohols, polyols, glycerol, vegetable oils, etc.

 Suitable adjuvants for suppositories are, for example, natural or hardened oils, waxes, fats, semi-solid or liquid polyols, etc.

25 Moreover, the pharmaceutical preparations can contain preservatives, solubilizers, viscosity-increasing substances, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They can also contain still other therapeutically valuable substances.

The dosage can vary in wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of about 0.1 mg to 20 mg per kg body weight, preferably about 0.5 mg to 4 mg per kg body weight (e.g. about 300 mg per person), divided into preferably 1-3 individual doses, which can
5 consist, for example, of the same amounts, should be appropriate. It will, however, be clear that the upper limit given herein can be exceeded when this is shown to be indicated.

The invention is illustrated hereinafter by Examples, which have no limiting character.

In case the preparative examples are obtained as a mixture of enantiomers, the pure enantiomers can be obtained by methods described herein or by methods known to those skilled
10 in the art, such as e.g. chiral chromatography or crystallization.

Examples


All examples and intermediates were prepared under nitrogen atmosphere if not specified otherwise.

Abbreviations:

- 5 aq. = aqueous; CAS-RN = Chemical Abstracts Service Registry Number; e.r. = enantiomeric ratio; HPLC = high performance liquid chromatography; MS = mass spectrum; sat. = saturated

Example 1

(E)-1-[(3aS,8aR)-2-(4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one






10




- To a solution of (E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one hydrochloride (intermediate 5; 40 mg, 102 μmol) in N,N-dimethylformamide (2 mL) were added N-methylmorpholine (51.8 mg, 512 μmol), 4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid (CAS-RN 33062-47-4; 17.1 mg, 102 μmol) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (42.8 mg, 113 μmol) at room temperature, then after 16 h the reaction mixture was partitioned between ethyl acetate and sat. aq. ammonium chloride solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (41 mg, 80%). Colourless gum, MS: 504.7 (M+H)⁺.




20




The following examples were prepared according to example 1, replacing (E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one hydrochloride and 4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid by the appropriate amine and carboxylic acid reagents, respectively.




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.01	1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)propan-1-one 	3-(3,5-dichlorophenyl)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)propan-1-one hydrochloride (intermediate 3)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	458.3 (M+H) ⁺
1.02	6-((3aR,6aS)-5-(3-(3,5-dichlorophenyl)propanoyl)octahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzo[d]oxazol-2(3H)-one 	3-(3,5-dichlorophenyl)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)propan-1-one hydrochloride (intermediate 3)	2-oxo-2,3-dihydro-benzo[d]-oxazole-6-carboxylic acid	474.1 (M+H) ⁺
1.03	(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	460.3 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.04	(3aR,6aS)-3,5-dichlorobenzyl 5-(9H-pyrido[3,4-b]indole-3-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	9H-pyrido[3,4-b]indole-3-carboxylic acid (CAS-RN 74214-63-4)	509.2 (M+H) ⁺
1.05	(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-indole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	1H-indole-5-carboxylic acid	458.3 (M+H) ⁺
1.06	(3aR,6aS)-3,5-dichlorobenzyl 5-(9H-carbazole-3-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	9H-carbazole-3-carboxylic acid (CAS-RN 51035-17-7)	508.3 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.07	(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-indazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	1H-indazole-5-carboxylic acid	459.2 (M+H) ⁺
1.08	(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-benzo[d]imidazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	1H-benzo[d]-imidazole-5-carboxylic acid	459.3 (M+H) ⁺
1.09	trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate 	trans-3,5-dichlorobenzyl hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate hydrochloride (intermediate 1.1)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	474.4 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.10	cis-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate 	cis-3,5-dichlorobenzyl hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate hydrochloride (intermediate 1.2)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	474.4 (M+H)⁺
1.11	(3aR,8aS)-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate 	(3aR,8aS)-3,5-dichlorobenzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate hydrochloride (intermediate 1.3)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	488.4 (M+H)⁺
1.12	(1H-benzotriazol-5-yl)-{(3aS,6aR)-5-[2-(3-chloro-phenyl)-ethanesulfonyl]-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl}-methanone 	(3aR,6aS)-2-(3-chlorophenethyl-sulfonyl)octahydro-pyrrolo[3,4-c]pyrrole (intermediate 21)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	460.5 (M+H)⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.13	(3aR,6aS)-3,5-dichlorobenzyl 5-(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	4,5,6,7-tetrahydro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (CAS-RN 33062-47-4)	464.5 (M+H) ⁺
1.14	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chlorophenyl)-2,2-dimethylpropan-1-one 	3-(3-chlorophenyl)-2,2-dimethyl-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)propan-1-one (intermediate 26.02)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	480.6 (M+H) ⁺
1.15	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.03)	1H-benzo[d]-imidazole-5-carboxylic acid	518.6 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.16	(3aSR,6SR,7aSR)-6-((3aR,8aS)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl)-hexahydro-benzooxazol-2-one 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	(3aSR,6SR,7aSR)-2-oxooctahydro-benzo[d]oxazole-6-carboxylic acid (intermediate 29)	522.7 (M+H) ⁺
1.17	(E)-1-((3aS,8aR)-2-(benzo[c][1,2,5]oxadiazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl)-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	benzo[c]-[1,2,5]oxadiazole-5-carboxylic acid	501.5 (M+H) ⁺
1.18	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-methyl-4-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(E)-3-(2-methyl-4-(trifluoromethoxy)-phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.05)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	514.7 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.19	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-methoxyphenyl)prop-2-en-1-one 	(E)-3-(3-fluoro-4-methoxyphenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.06)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	464.4 (M+H) ⁺
1.20	(E)-1-[(3aS,8aR)-2-((S)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxyphenyl)-prop-2-en-1-one 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	(-)-(S)-4,5,6,7-tetrahydro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	504.6 (M+H) ⁺
1.21	(E)-1-[(3aS,8aR)-2-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxyphenyl)-prop-2-en-1-one 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	(+)-(R)-4,5,6,7-tetrahydro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	504.6 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.22	(E)-1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(2-isopropyl-phenyl)-prop-2-en-1-one 	(E)-3-(2-isopropylphenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.07)	(+)-(R)-4,5,6,7-tetrahydro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	458.3 (M+H) ⁺
1.23	trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester 	trans-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester hydrochloride (intermediate 1.5)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	518.6 (M+H) ⁺
1.24	trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 4-trifluoromethoxy-benzyl ester 	trans-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 4-trifluoromethoxy-benzyl ester hydrochloride (intermediate 1.4)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	490.6 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.25	1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-2-(4-trifluoromethoxy-phenoxy)-ethanone 	1-(trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl)-2-(4-(trifluoromethoxy)phenoxy)-ethanone hydrochloride (intermediate 5.4)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	490.6 (M+H) ⁺
1.26	(E)-1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propenone 	(E)-1-(trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one hydrochloride (intermediate 5.3)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	486.7 (M+H) ⁺
1.27	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chloro-5-(trifluoromethyl)phenyl)prop-2-en-1-one 	(E)-3-(3-chloro-5-(trifluoromethyl)-phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.10)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	518.6 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.28	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxy-2-(trifluoromethyl)phenyl)prop-2-en-1-one 	(E)-3-(4-methoxy-2-(trifluoromethyl)-phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.09)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	514.7 (M+H) ⁺
1.29	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-cyclopropylphenyl)prop-2-en-1-one 	(E)-3-(2-cyclopropylphenyl)-1-((trans-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.08)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	456.7 (M+H) ⁺
1.30	trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 4-fluoro-2-trifluoromethyl-benzyl ester 	trans-(4-fluoro-2-(trifluoromethyl)-benzyl) hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate (intermediate 32.3)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	492.6 (M+H) ⁺

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.31	trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 2-cyclopropyl-4-trifluoromethyl-benzyl ester 	trans-(2-cyclopropyl-4-(trifluoromethyl)-benzyl) hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate (intermediate 32.2)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	514.6 (M+H) ⁺
1.32	1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(2-trifluoromethoxy-phenoxy)-ethanone 	1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(2-(trifluoromethoxy)-phenoxy)ethanone (intermediate 36.07)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	504.2 (M+H) ⁺
1.33	trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 2-methoxy-4-trifluoromethoxy-benzyl ester 	trans-(2-methoxy-4-(trifluoromethoxy)-benzyl) hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate (intermediate 32.1)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	520.6 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.34	4-{2-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-oxo-ethoxy}-3-trifluoromethyl-benzonitrile 	4-(2-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-oxoethoxy)-3-(trifluoromethyl)-benzonitrile (intermediate 36.05)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	511.1 (M-H) ⁻
1.35	1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone 	2-(4-chloro-2-isopropyl-5-methylphenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone (intermediate 36.06)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	510.2 (M+H) ⁺
1.36	1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-[4-methyl-2-(1-methyl-pyrrolidin-3-yl)-phenoxy]-ethanone 	2-(4-methyl-2-(1-methylpyrrolidin-3-yl)phenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone (intermediate 36.04)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	517.3 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.37	1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(2-chloro-4-fluorophenoxy)-ethanone 	2-(2-chloro-4-fluorophenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone (intermediate 36.03)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	470.1 (M-H) ⁻
1.38	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(2-chloro-4-(trifluoromethyl)phenoxy)ethanone 	2-(2-chloro-4-(trifluoromethyl)phenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone (intermediate 36.10)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	522.6 (M+H) ⁺
1.39	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(6-isopropyl-3,3-dimethyl-2,3-dihydro-1H-inden-5-yloxy)ethanone 	2-(6-isopropyl-3,3-dimethyl-2,3-dihydro-1H-inden-5-yloxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone (intermediate 36.01)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	530.7 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.40	(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester 	2-(2-fluoro-4-(trifluoromethoxy)-phenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	520.6 (M-H) ⁻
1.41	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(5-chloro-2-(trifluoromethyl)phenoxy)ethanone 	2-(5-chloro-2-(trifluoromethyl)-phenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone (intermediate 36)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	522.5 (M+H) ⁺
1.42	1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-2-(2-tert-butyl-4-methoxyphenoxy)ethanone 	2-(2-tert-butyl-4-methoxyphenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone (intermediate 36.08)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	506.6 (M+H) ⁺




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.43	4-[2-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-2-oxoethoxy]-3-propan-2-ylbenzonitrile 	3-isopropyl-4-(2-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-oxoethoxy)benzonitrile (intermediate 36.09)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	487.6 (M+H) ⁺
1.44	1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-3-[3-fluoro-4-(2,2,2-trifluoroethoxy)phenyl]propan-1-one 	3-(3-fluoro-4-(2,2,2-trifluoroethoxy)-phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)propan-1-one (intermediate 35.2)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	532.2 (M+H) ⁺
1.45	1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-3-[2-fluoro-4-(2,2,2-trifluoroethoxy)phenyl]propan-1-one 	3-(2-fluoro-4-(2,2,2-trifluoroethoxy)-phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)propan-1-one (intermediate 26.12)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	534.2 (M+H) ⁺


No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.46	(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-fluoro-4-(2,2,2-trifluoro-ethoxy)-benzyl ester	(3aR,8aS)-3-fluoro-4-(2,2,2-trifluoroethoxy)-benzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate (intermediate 32.4)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	536.2 (M+H) ⁺
1.47	(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 2-fluoro-4-(2,2,2-trifluoro-ethoxy)-benzyl ester	(3aR,8aS)-2-fluoro-4-(2,2,2-trifluoroethoxy)-benzyl octahydro-pyrrolo[3,4-d]azepine-6(7H)-carboxylate hydrochloride (intermediate 1.7)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	536.6 (M+H) ⁺
1.48	(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 4-(2,2,2-trifluoro-ethoxy)-benzyl ester	(3aR,8aS)-4-(2,2,2-trifluoroethoxy)benzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate hydrochloride (intermediate 1.6)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	518.6 (M+H) ⁺


No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.49	(3aS,6aS)-5-(3H-[1,2,3]triazolo[4,5-b]pyridine-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoromethoxy)benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	3H-1,2,3-triazolo[4,5-b]pyridine-6-carboxylic acid (M-H) ⁻ (CAS-RN 1260385-82-7)	475.5
1.50	1-[(3aR,6aR)-5-(1H-triazolo[4,5-b]pyridine-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[4-(trifluoromethoxy)phenyl]propan-1-one 	1-((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one dihydrochloride (intermediate 5.5)	1H-[1,2,3]triazolo[4,5-b]pyridine-5-carboxylic acid (M+H) ⁺ (CAS-RN 1216149-55-1)	475.4
1.51	(3aS,6aS)-5-(3H-[1,2,3]triazolo[4,5-c]pyridine-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoromethoxy)benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	3H-[1,2,3]triazolo[4,5-c]pyridine-6-carboxylic acid (M+H) ⁺ (intermediate 45)	477.4

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.52	(3aS,6aS)-5-(4-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoro-methoxy)benzyl hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	4-fluoro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 43.2)	494.4 (M+H) ⁺
1.53	(3aS,6aS)-5-(7-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoro-methoxy)benzyl hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	7-fluoro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 43.1)	494.6 (M+H) ⁺
1.54	(3aS,6aS)-5-(6-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoromethoxy)-benzyl hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	6-fluoro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 43)	494.5 (M+H) ⁺

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.55	(3aS,6aS)-5-(4-chloro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoromethoxy)benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	4-chloro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 43.5)	510.4 (M+H) ⁺
1.56	(3aS,6aS)-5-(6-trifluoromethyl-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoromethoxy)benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	6-(trifluoromethyl)-1H-benzo[d][1,2,3]triazole-5-carboxylic acid (intermediate 43.4)	542.4 (M-H) ⁻
1.57	(3aS,6aS)-5-(4-methyl-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoromethoxy)benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	4-methyl-1H-benzo[d][1,2,3]triazole-5-carboxylic acid (intermediate 43.3)	488.2 (M-H) ⁻




No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.58	(3aS,6aS)-5-(6-methyl-1H-benzotriazole-5-carbonyl)-hexahydropyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoromethoxy)benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	6-methyl-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 43.6)	488.2 (M-H) ⁻
1.59	1-[(3aR,6aR)-5-(4-fluoro-1H-benzotriazole-5-carbonyl)-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one 	1-((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)-phenyl)propan-1-one dihydrochloride (intermediate 5.5)	4-fluoro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 43.2)	492.2 (M+H) ⁺
1.60	(4-ethoxyquinolin-2-yl)((3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone 	(4-ethoxyquinolin-2-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 5.6)	(R)-4,5,6,7-tetrahydro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 30A)	461.3 (M+H) ⁺


No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
1.61	(4-ethoxyquinolin-2-yl)((3aS,6aS)-5-(4-fluoro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone 	(4-ethoxyquinolin-2-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 5.6)	4-fluoro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 43.2)	475.3 (M+H) ⁺

Example 2**trans-3,5-Dichlorobenzyl 2-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate**

To a solution of trans-3,5-dichlorobenzyl hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate hydrochloride (intermediate 1.1; 50 mg, 137 μ mol) in N,N-dimethylformamide (1 mL) were added 4-methylmorpholine (69.1 mg, 684 μ mol), 4-amino-3-hydroxybenzoic acid (20.9 mg, 137 μ mol) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (62.4 mg, 164 μ mol) at room temperature, then after 18 h 1,1'-carbonyldiimidazole (50.3 mg, 301 μ mol) was added. After 1 h, the reaction mixture was partitioned between ethyl acetate and 1 M aq. hydrochloric acid solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered and evaporated in vacuo. Chromatography (silica gel, gradient dichloromethane dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (35 mg, 52%). Light yellow gum, MS: 490.5 (M+H)⁺.


The following examples were prepared according to example 2, replacing trans-3,5-dichlorobenzyl hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate hydrochloride by the appropriate starting material.

Ex.	Systematic Name	Starting material	MS, m/e
2.01	(3aR,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	476.2 (M+H) ⁺
2.02	6-((3aS,8aR)-6-[(E)-3-(4-trifluoromethoxyphenyl)acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl)-3H-benzoxazol-2-one 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	514.6 (M-H) ⁻
2.03	6-((3aR,6aR)-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrole-5-carbonyl)-3H-1,3-benzoxazol-2-one 	1-((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one dihydrochloride (intermediate 5.5)	490.4 (M+H) ⁺

2.04	(3aS,6aS)-5-(2-oxo-2,3-dihydro-benzooxazole-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(3aS,6aS)-4-(trifluoromethoxy)benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	492.6 (M+H) ⁺
------	--	---	--------------------------

Example 3

(3aR,5s,6aS)-3,5-Dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yloxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate




5

A mixture of 4-((3aR,5s,6aS)-2-((3,5-dichlorobenzoyloxy)carbonyl)octahydrocyclopenta[c]pyrrol-5-yloxy)-2-hydroxybenzoic acid (intermediate 12.1; 63 mg, 135 μ mol), triethylamine (13.7 mg, 135 μ mol) and diphenylphosphoryl azide (37.2 mg, 135 μ mol) in toluene (2 mL) was heated at 110°C for 20 h. After evaporation of the solvent, the residue was purified by chromatography (silica gel; heptane – ethyl acetate gradient) to produce the title compound (19 mg, 30%). White solid, MS: 463.2 (M+H)⁺.

10


Example 3.01

(3aR,5r,6aS)-3,5-Dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yloxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate



15

The title compound was produced in analogy to example 3 from 4-((3aR,5r,6aS)-2-((3,5-dichlorobenzoyloxy)carbonyl)octahydrocyclopenta[c]pyrrol-5-yloxy)-2-hydroxybenzoic acid (intermediate 12). White solid, MS: 463.2 (M+H)⁺.




Example 4**(3aS,6aS)-3,5-Dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate**




- 5 To a solution of (3,5-dichlorophenyl)methanol (21.4 mg, 121 μmol) in acetonitrile (5 mL) was added N,N'-carbonyldiimidazole (20.6 mg, 127 μmol) at room temperature, then after 3 h triethylamine (61.3 mg, 606 μmol) and (1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4; 40 mg, 121 μmol) were added and the reaction mixture was heated at reflux. After 16 h the reaction
- 10 mixture was partitioned between ethyl acetate and sat. aq. ammonium chloride, the organic layer was washed with sat. aq. sodium hydrogen carbonate solution and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (38 mg, 68%). Light yellow foam, MS: 460.4 (M+H)⁺.




15




The following compounds were produced in analogy to example 4, replacing (1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride and (3,5-dichlorophenyl)methanol by the appropriate amine and alcohol precursors, respectively.




No.	Systematic Name	Amine	Alcohol	MS, m/e




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.01	trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-2-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester 	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride (intermediate 2.1)	(3-(methylsulfonyl)-5-(trifluoromethyl)phenyl)-methanol (CAS-RN 1003843-94-4)	552.5 (M+H)⁺
4.02	(3aR,6aR)-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 8)	(3,5-dichlorophenyl)-methanol	460.5 (M+H)⁺
4.03	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(3-chloro-5-(methylsulfonyl)-phenyl)-methanol (intermediate 17)	504.4 (M+H)⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.04	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(3-chloro-5-(methylsulfonyl)-phenyl)-methanol (intermediate 17)	504.4 (M+H) ⁺
4.05	(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(3-(methylsulfonyl)-5-(trifluoromethyl)-phenyl)-methanol (CAS-RN 1003843-94-4)	566.2 (M+H) ⁺
4.06	(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(3-chloro-5-(methylsulfonyl)-phenyl)-methanol (intermediate 17)	532.3 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.07	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(3-(methylsulfonyl)-5-(trifluoromethyl)-phenyl)-methanol (CAS-RN 1003843-94-4)	538.4 (M+H)⁺
4.08	cis-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate 	(1H-benzo[d][1,2,3]triazol-5-yl)(cis-tetrahydro-1H-pyrrolo[3,4-c]pyridin-5(6H,7H,7aH)-yl)methanone hydrochloride (intermediate 2)	(3,5-dichlorophenyl)-methanol	474.5 (M+H)⁺
4.09	(3aS,7aR)-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester 	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone hydrochloride (intermediate 2.1)	(3-chloro-5-(methylsulfonyl)-phenyl)-methanol (intermediate 17)	518.4 (M+H)⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.10	trans-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate 	(1H-benzo[d][1,2,3]triazol-5-yl)(trans-tetrahydro-1H-pyrrolo[3,4-c]pyridin-5(6H,7H,7aH)-yl)methanone hydrochloride (intermediate 2.1)	(3,5-dichlorophenyl)-methanol	474.4 (M+H)⁺
4.11	(3aR,8aS)-3,5-dichlorobenzyl 6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)methanone hydrochloride (intermediate 2.3)	(3,5-dichlorophenyl)-methanol	486.4 (M-H)⁻
4.12	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 1-(3-chloro-phenyl)-cyclopropyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	1-(3-chlorophenyl)-cyclopropanol (CAS-RN 43187-67-3)	452.5 (M+H)⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.13	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid bicyclo[4.1.0]hept-7-ylmethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	bicyclo[4.1.0]-heptan-7-ylmethanol	410.5 (M+H) ⁺
4.14	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid adamantan-2-ylmethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	2-adamantane-methanol (CAS-RN 22635-61-6)	450.5 (M+H) ⁺
4.15	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 1-fluorocyclohexylmethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(1-fluorocyclohexyl)-methanol (CAS-RN 117169-30-9)	414.4 (M-H) ⁻




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.16	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-adamantan-2-yl-ethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	2-(2-adamantyl)-ethanol (CAS-NR 39555-28-7)	464.5 (M+H) ⁺
4.17	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-adamantan-1-yl-ethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	1-adamantane-ethanol	464.5 (M+H) ⁺
4.18	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid adamantan-1-ylmethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	1-adamantane-methanol	450.5 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.19	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid cyclohexylmethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	cyclohexyl-methanol	398.5 (M+H) ⁺
4.20	cis-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2,2-trifluoro-1-methoxy-ethyl)-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(3-(2,2,2-trifluoro-1-methoxy-ethyl)phenyl)-methanol (intermediate 19)	504.5 (M+H) ⁺
4.21	cis-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2,2-trifluoro-1-hydroxy-ethyl)-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	2,2,2-trifluoro-1-(3-(hydroxy-methyl)-phenyl)ethanol (intermediate 18)	490.5 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.22	(3aR,6aS)-2-cyclohexylethyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	2-cyclohexylethanol	412.5 (M+H) ⁺
4.23	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-5-trifluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(3-fluoro-5-(trifluoromethoxy)-phenyl)-methanol	494.4 (M+H) ⁺
4.24	(3aR,6aS)-3-chloro-5-cyanobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	3-chloro-5-(hydroxymethyl)-benzonitrile (CAS-RN 1021871-35-1)	451.4 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.25	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-trifluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(3-(trifluoromethoxy)-phenyl)-methanol	476.4 (M+H) ⁺
4.26	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-5-trifluoromethyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(3-fluoro-5-(trifluoromethyl)-phenyl)-methanol	478.5 (M+H) ⁺
4.27	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-trifluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(3-chloro-5-(trifluoromethoxy)-phenyl)-methanol	510.4 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.28	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-fluoro-3-trifluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)(trans-hexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(4-fluoro-3-(trifluoromethoxy)-phenyl)-methanol (CAS-RN 86256-18-0)	494.5 (M+H) ⁺
4.29	(3aR,6aS)-3-cyano-5-fluorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahdropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(1H-benzo[d][1,2,3]triazol-5-yl)(trans-hexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	3-fluoro-5-(hydroxymethyl)-benzonitrile (CAS-RN 1021871-34-0)	435.4 (M+H) ⁺
4.30	(3aR,6aS)-3-chloro-5-methoxybenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahdropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(1H-benzo[d][1,2,3]triazol-5-yl)(trans-hexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(3-chloro-5-methoxy-phenyl)-methanol (CAS-RN 82477-68-7)	456.5 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.31	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1S,4R)-3-methyl-bicyclo[2.2.1]hept-2-ylmethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)(trans-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	2-norbornane-methanol (CAS-RN 6968-75-8)	423.2 (M+H) ⁺
4.32	(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1R,4S)-1-bicyclo[2.2.1]hept-2-ylmethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)(trans-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(1R,4S)-bicyclo[2.2.1]heptan-2-ylmethanol	410.2 (M+H) ⁺
4.33	(3aR,5s,6aS)-5-[(3H-[1,2,3]triazol-4-ylmethyl)-carbamoyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid 3,5-dichloro-benzyl ester 	(3aR,5s,6aS)-N-((1H-1,2,3-triazol-4-yl)methyl)octahydro-cyclopenta[c]pyrrole-5-carboxamide 2,2,2-trifluoroacetate (intermediate 4)	(3,5-dichloro-phenyl)-methanol	438.4 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.34	(3aS,6aS)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester 	(6,7-dihydro-1H-[1,2,3]triazolo[4,5-c]pyridin-5(4H)-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 25)	(3-chloro-5-(methylsulfonyl)-phenyl)-methanol (intermediate 17)	509.5 (M+H) ⁺
4.35	(3aS,6aR)-5-[(1H-[1,2,3]triazol-4-ylmethyl)-carbamoyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3,5-dichloro-benzyl ester 	(3aR,6aS)-N-((1H-1,2,3-triazol-4-yl)methyl)hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxamide 2,2,2-trifluoroacetate (intermediate 20)	(3,5-dichloro-phenyl)-methanol	440.3 (M+H) ⁺
4.36	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(4-(trifluoromethoxy)-phenyl)-methanol	476.5 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.37	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-(hydroxymethyl)-benzonitrile	415.5 (M-H) ⁻
4.38	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-(1,1,2,2-tetrafluoro-ethoxy)-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(4-(1,1,2,2-tetrafluoroethoxy)-phenyl)-methanol (CAS-RN 773868-39-6)	508.4 (M+H) ⁺
4.39	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-difluoromethoxy-3-fluoro-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(4-(difluoromethoxy)-3-fluorophenyl)-methanol (CAS-RN 1242252-59-0)	476.4 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.40	(3aS,6aS)-5-(1H-[1,2,3]triazolo[4,5-b]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester 	(1H-[1,2,3]triazolo[4,5-b]pyridin-5-yl)((3aR,6aR)-hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.6)	(3-fluoro-4-(trifluoromethoxy)-phenyl)-methanol (CAS-RN 886498-99-3)	495.3 (M+H) ⁺
4.41	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-difluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(4-(difluoromethoxy)-phenyl)-methanol (CAS-RN 170924-50-2)	458.6 (M+H) ⁺
4.42	(3aS,6aS)-5-(1H-[1,2,3]triazolo[4,5-b]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(1H-[1,2,3]triazolo[4,5-b]pyridin-5-yl)((3aR,6aR)-hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.6)	(4-(trifluoromethoxy)-phenyl)-methanol	477.6 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.43	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(3-fluoro-4-(trifluoromethoxy)-phenyl)-methanol (CAS-RN 886498-99-3)	494.4 (M+H) ⁺
4.44	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-(2,2,2-trifluoro-ethoxy)-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(4-(2,2,2-trifluoroethoxy)-phenyl)-methanol (CAS-RN 1020949-12-5)	490.4 (M+H) ⁺
4.45	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 5-trifluoromethoxy-pyridin-2-ylmethyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(5-(trifluoromethoxy)-pyridin-2-yl)methanol (CAS-RN 31181-85-8)	477.4 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.46	(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-benzyl ester 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	4-(hydroxymethyl)-3-isopropyl-benzonitrile (intermediate 41)	463.5 (M+H) ⁺
4.47	(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-5-methyl-benzyl ester 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	4-(hydroxymethyl)-5-isopropyl-2-methyl-benzonitrile (intermediate 41.1)	477.5 (M+H) ⁺
4.48	(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	(2-fluoro-4-(trifluoromethoxy)-phenyl)-methanol (CAS-RN 1240257-07-1)	498.4 (M+H) ⁺




No.	Systematic Name	Amine	Alcohol	MS, m/e
4.49	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(2-fluoro-4-(trifluoromethoxy)-phenyl)-methanol (CAS-RN 1240257-07-1)	494.6 (M+H) ⁺
4.50	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-ethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-ethoxy-4-(hydroxymethyl)-benzonitrile (intermediate 41.2)	459.6 (M+H) ⁺
4.51	(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	(3-fluoro-4-(trifluoromethoxy)-phenyl)-methanol (CAS-RN 886498-99-3)	496.4 (M-H) ⁻

No.	Systematic Name	Amine	Alcohol	MS, m/e
4.52	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-(hydroxymethyl)-3-isopropylbenzonitrile (intermediate 41)	459.5 (M+H) ⁺
4.53	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-5-methyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-(hydroxymethyl)-5-isopropyl-2-methylbenzonitrile (intermediate 41.1)	473.5 (M+H) ⁺
4.54	(3aS,6aS)-5-(1,4,6,7-tetrahydro[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(6,7-dihydro-1H-[1,2,3]triazolo[4,5-c]pyridin-5(4H)-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 25)	(4-(trifluoromethoxy)phenyl)methanol	481.6 (M+H) ⁺

No.	Systematic Name	Amine	Alcohol	MS, m/e
4.55	(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 8)	(4-(trifluoromethoxy)-phenyl)-methanol	476.4 (M+H) ⁺
4.56	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(2-fluoro-4-(trifluoromethyl)-phenyl)-methanol	478.4 (M+H) ⁺
4.57	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(4-(trifluoromethyl)-phenyl)-methanol	460.5 (M+H) ⁺

No.	Systematic Name	Amine	Alcohol	MS, m/e
4.58	(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	(4-(trifluoromethoxy)-phenyl)-methanol	480.5 (M+H) ⁺
4.59	(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethyl-benzyl ester 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	(4-(trifluoromethyl)-phenyl)-methanol	464.5 (M+H) ⁺
4.60	(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	(3-chloro-5-(methylsulfonyl)-phenyl)-methanol (intermediate 17)	508.4 (M+H) ⁺


No.	Systematic Name	Amine	Alcohol	MS, m/e
4.61	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-methanesulfonyl-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-(hydroxymethyl)-3-(methylsulfonyl)benzonitrile (intermediate 48)	493.2 (M-H) ⁻
4.62	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-ethoxy-5-fluoro-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	5-ethoxy-2-fluoro-4-(hydroxymethyl)-benzonitrile (intermediate 49)	477.3 (M-H) ⁻
4.63	(3aS,6aS)-5-(4-methoxy-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzylester 	(3aS,6aS)-4-(trifluoromethoxy)-benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	4-methoxy-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 45.1)	504.3 (M-H) ⁻

No.	Systematic Name	Amine	Alcohol	MS, m/e
4.64	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-cyclobutoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-cyclobutoxy-4-(hydroxymethyl)-benzonitrile (intermediate 46)	487.3 (M+H) ⁺
4.65	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropoxy-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-(hydroxymethyl)-3-isopropoxy-benzonitrile (intermediate 46.1)	475.3 (M+H) ⁺
4.66	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-(2,2,2-trifluoro-ethoxy)-benzyl ester 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-(hydroxymethyl)-3-(2,2,2-trifluoroethoxy)-benzonitrile (intermediate 47.1)	515.3 (M+H) ⁺

No.	Systematic Name	Amine	Alcohol	MS, m/e
4.67	(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-chloro-2-ethoxy-5-fluoro-benzyl ester	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(4-chloro-2-ethoxy-5-fluorophenyl) methanol (intermediate 47)	488.2 (M+H) ⁺

Example 5

(3aR,5r,6aS)-3,5-Dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-ylamino)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate




5

Sodium triacetoxyborohydride (121 mg, 556 μ mol,) was added at room temperature to a solution of (3aR,6aS)-3,5-dichlorobenzyl 5-oxohexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (intermediate 13; 128 mg, 371 μ mol), 6-aminobenzo[d]oxazol-2(3H)-one (CAS-RN 22876-17-1; 57.3 mg, 371 μ mol) and acetic acid (134 mg, 2.22 mmol) in 1,2-dichloroethane (2.5 mL), then after 16 h another portion of sodium triacetoxyborohydride (39.3 mg, 185 μ mol) was added. After another 6 h the reaction mixture was partitioned between ethyl acetate and sat. aq. sodium hydrogen carbonate solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 95:5:0.25), followed by HPLC chromatography (Reprosil Chiral-NR, heptane/ethanol 3:2) produced the title compound (70 mg, 41%). White solid, MS: 462.2 (M+H)⁺.

Example 6




(3aR,6aS)-3,5-Dichlorobenzyl 5-((1H-benzo[d]imidazol-5-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate






To a white suspension of (3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1; 35 mg, 99.5 μmol) and 1H-benzo[d]imidazole-5-carbaldehyde (15.0 mg, 99.5 μmol) in tetrahydrofuran (2 mL) were added sodium triacetoxyborohydride (31.6 mg, 149 μmol) and acetic acid (9.0 mg, 150 μmol), then after 3 h the reaction mixture was partitioned between ethyl acetate and sat. aq. sodium hydrogen carbonate solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (42 mg, 95%). White foam, MS: 445.3 (M+H)⁺.

The following compounds were produced in analogy to example 6, replacing (3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride and 1H-benzo[d]imidazole-5-carbaldehyde by the appropriate amine and aldehyde reagents, respectively.


No.	Systematic Name	Amine	Aldehyde	MS, m/e
6.01	1-((3aR,6aS)-5-((1H-benzo[d][1,2,3]triazol-5-yl)methyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)propan-1-one	3-(3,5-dichlorophenyl)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)propan-1-one hydrochloride (intermediate 3)	1H-benzo[d]-[1,2,3]triazole-5-carbaldehyde (CAS-RN 70938-42-0)	444.5 (M+H) ⁺

No.	Systematic Name	Amine	Aldehyde	MS, m/e
6.02	(3aR,6aS)-3,5-dichlorobenzyl 5-((1H-indazol-5-yl)methyl)hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	1H-indazole-5-carbaldehyde	445.2 (M+H) ⁺
6.03	(3aR,6aS)-3,5-dichlorobenzyl 5-((2-oxo-2,3-dihydrobenzo[d]oxazol-6-yl)methyl)hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate 	(3aR,6aS)-3,5-dichlorobenzyl hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1)	2-oxo-2,3-dihydro-benzo[d]ox-azole-6-carbaldehyde (CAS-RN 54903-15-0)	462.2 (M+H) ⁺
6.04	6-(((3aR,6aS)-5-(3-(3,5-dichlorophenyl)propanoyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)methyl)benzo[d]oxazol-2(3H)-one 	3-(3,5-dichloro-phenyl)-1-((3aR,6aS)-hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)propan-1-one hydrochloride (intermediate 3)	2-oxo-2,3-dihydro-benzo[d]ox-azole-6-carbaldehyde (CAS-RN 54903-15-0)	460.3 (M+H) ⁺

No.	Systematic Name	Amine	Aldehyde	MS, m/e
6.05	4-{(E)-3-[(3aS,8aR)-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-oxo-propenyl}-benzonitrile 	4-((E)-3-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)benzonitrile hydrochloride (intermediate 5.1)	1H-benzo[d]-[1,2,3]triazole-5-carbaldehyde (CAS-RN 70938-42-0)	427.6 (M+H) ⁺
6.06	(E)-1-[(3aS,8aR)-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	1H-benzo[d]-[1,2,3]triazole-5-carbaldehyde (CAS-RN 70938-42-0)	485.5 (M+H) ⁺
6.07	(E)-1-[trans-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propenone 	(E)-1-(trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride (intermediate 5.3)	1H-benzo[d]-[1,2,3]triazole-5-carbaldehyde (CAS-RN 70938-42-0)	472.7 (M+H) ⁺

Example 7


(3aR,6aS)-3,5-Dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-ylsulfonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate



To a suspension of (3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-
 5 carboxylate hydrochloride (intermediate 1; 40 mg, 114 μ mol) and pyridine (45.0 mg, 569 μ mol)
 in acetone (2 mL) was added 2-oxo-2,3-dihydrobenzo[d]oxazole-6-sulfonyl chloride (25.2 mg,
 108 μ mol) at room temperature, then after 40 h the reaction mixture was partitioned between sat.
 aq. sodium hydrogencarbonate solution and ethyl acetate. The organic layer was washed with
 brine, dried over magnesium sulfate, filtered, and evaporated. The residue was dissolved in ethyl
 10 acetate, then after 40 min the suspension formed was treated with ethyl acetate/heptane 1:1 and
 the precipitate was collected by filtration to afford the title compound (24 mg, 41%). White solid,
 MS: 512.2 (M+H)⁺.




Example 8




**(3aR,6aS)-3,5-Dichlorobenzyl 5-(4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine-5-
 15 carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate**





To a suspension of 4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine (CAS-RN 706757-05-3;
 52.5 mg, 423 μ mol) in dichloromethane (8 mL) was added a solution of (3aR,6aS)-3,5-
 dichlorobenzyl 5-(chlorocarbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate
 20 (intermediate 9; 168 mg, 423 μ mol) in dichloromethane (2 mL) dropwise at room temperature,
 then after 1 h, N,N-dimethylformamide (1 mL) was added. After 96 h, the reaction mixture was
 washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate,
 filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to
 dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound
 25 (136 mg, 69%). Colourless oil, MS: 465.5 (M+H)⁺.

The following examples were produced in analogy to example 8, replacing (3aR,6aS)-3,5-dichlorobenzyl 5-(chlorocarbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate by (3aR,8aS)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carbonyl chloride (intermediate 9.1) and 4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine by the appropriate amine precursor.

Ex.	Systematic Name	Amine	MS, m/e
8.01	(E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine (CAS-RN 706757-05-3)	505.7 (M+H) ⁺
8.02	(E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	4,5,6,7-tetrahydro-1H-pyrazolo[4,3-c]pyridine dihydrochloride (CAS-RN 157327-44-1)	504.7 (M+H) ⁺
8.03	cis-5-((3aR,8aS)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)decahydro-pyrrolo[3,4-d]azepine-2-carbonyl)-hexahydrooxazolo[5,4-c]pyridin-2(1H)-one 	cis-hexahydro-oxazolo[5,4-c]pyridin-2(1H)-one hydrochloride (intermediate 28)	523.5 (M+H) ⁺

Ex.	Systematic Name	Amine	MS, m/e
8.04	6-{(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl}-5,6,7,8-tetrahydro-1H-[1,6]naphthyridin-2-one 	5,6,7,8-tetrahydro-1,6-naphthyridin-2(1H)-one hydrochloride (CAS-RN 1211505-91-7)	531.6 (M+H) ⁺
8.05	(3aR,7aR)-5-{(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl}-hexahydro-oxazolo[5,4-c]pyridin-2-one 	(3aR,7aR)-hexahydro-oxazolo[5,4-c]pyridin-2(1H)-one hydrochloride (intermediate 27)	523.6 (M+H) ⁺
8.06	(E)-1-[(3aS,8aR)-2-(7,8-dihydro-5H-[1,6]naphthyridine-6-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	5,6,7,8-tetrahydro-1,6-naphthyridine dihydrochloride (CAS-RN 348623-30-3)	515.5 (M+H) ⁺



Ex.	Systematic Name	Amine	MS, m/e
8.07	(E)-3-(4-trifluoromethoxy-phenyl)-1-[(3aS,8aR)-2-(2-trifluoromethyl-5,6-dihydro-8H-[1,2,4]triazolo[1,5-a]pyrazine-7-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-prop-2-en-1-one 	2-(trifluoromethyl)-5,6,7,8-tetrahydro-[1,2,4]triazolo[1,5-a]pyrazine (CAS-RN 681249-57-0)	573.7 (M+H) ⁺




Example 9**(3aR,8aS)-N-((1H-1,2,3-Triazol-5-yl)methyl)-6-((E)-3-(4-(trifluoromethoxy)phenyl)-acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide**




5

A solution of bis(trichloromethyl)-carbonate (73.7 mg, 248 μ mol) in ethyl acetate (10 mL) was added dropwise at 0°C over a period of 5 minutes to a solution of (E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one (intermediate 5; 176 mg, 497 μ mol) in tetrahydrofuran (5 mL). The ice bath was removed, then after 30 min the reaction mixture was heated at reflux for 2 h. After concentration under vacuum, the residue was dissolved in tetrahydrofuran (7 mL), then after addition of (1H-1,2,3-triazol-4-yl)methanamine hydrochloride (66.8 mg, 497 μ mol) and triethylamine (251 mg, 2.48 mmol) the reaction mixture was stirred at room temperature for 17 hours and then partitioned between sat. aq. sodium hydrogen carbonate solution and ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane/ethyl acetate 4:1, then dichloromethane/methanol 95:5) afforded the title compound (68 mg, 28%). White solid, MS: 479.5 (M+H)⁺.

The following examples were produced in analogy to example 8, replacing (E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one by the appropriate starting material and (1H-1,2,3-triazol-4-yl)methanamine hydrochloride by the appropriate amine reagent.

Ex.	Systematic Name	Starting material	Amine reagent	MS, m/e
9.01	(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)-octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one (intermediate 5)	(1H-1,2,3-triazol-4-yl)-methanamine hydrochloride	479.5 (M+H) ⁺
9.02	(3aS,6aR)-5-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1H-[1,2,3]triazol-4-ylmethyl)-amide 	(E)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one (intermediate 26.01)	(1H-1,2,3-triazol-4-yl)-methanamine hydrochloride	451.5 (M+H) ⁺


Ex.	Systematic Name	Starting material	Amine reagent	MS, m/e
9.03	(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acryloyl)-octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide 	(E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.03)	(1H-1,2,3-triazol-4-yl)-methanamine hydrochloride	497.6 (M+H) ⁺
9.04	(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid (4H-[1,2,4]triazol-3-ylmethyl)-amide 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	1H-1,2,4-triazole-5-methanamine dihydrochloride (CAS-RN 859791-21-2)	479.2 (M+H) ⁺
9.05	(E)-1-[(3aS,8aR)-2-(6,7-dihydro-4H-[1,2,3]triazolo[1,5-a]pyrazine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	4,5,6,7-tetrahydro-[1,2,3]triazolo[1,5-a]pyrazine hydrochloride (CAS-RN 123308-28-1)	549.2 (M+HCOO) ⁻

Ex.	Systematic Name	Starting material	Amine reagent	MS, m/e
9.06	(E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-imidazo[4,5-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridine (CAS-RN 6882-74-2)	504.2 (M+H) ⁺
9.07	(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)-octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide 	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one (intermediate 5)	N-methyl-1H-1,2,3-triazole-5-methanamine (CAS-RN 1248059-33-7)	493.7 (M+H) ⁺
9.08	(3aS,8aR)-6-[3-(3-chloro-phenyl)-2,2-dimethyl-propionyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid (3H-[1,2,3]triazol-4-ylmethyl)-amide 	3-(3-chlorophenyl)-2,2-dimethyl-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)propan-1-one (intermediate 26.03)	(1H-1,2,3-triazol-4-yl)-methanamine hydrochloride	459.5 (M+H) ⁺

Ex.	Systematic Name	Starting material	Amine reagent	MS, m/e
9.09	(3aR,8aS)-N-(2-(1H-1,2,3-triazol-5-yl)ethyl)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)-octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one (intermediate 5)	2-(1H-1,2,3-triazol-5-yl)-ethanamine (CAS-RN 52845-67-7)	493.6 (M+H) ⁺
9.10	(3aS,6aS)-5-(4-sulfamoyl-piperidine-1-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester	(3aS,6aS)-4-(trifluoromethoxy)-benzyl hexahydro-pyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1.8)	piperidine-4-sulfonamide hydrochloride (CAS-RN 1251923-46-2)	521.6 (M+H) ⁺

Examples 10A and 10B

(3aR,7aS)-2-(1H-Benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3,5-dichloro-benzyl ester and (3aS,7aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3,5-dichloro-benzyl ester



Racemic cis-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate (example 1.02; 616 mg, 1.30 mmol) was separated by preparative HPLC using a Reprosil Chiral-NR column as the stationary phase and


heptane/ethanol 3:2 as the mobile phase. This produced the faster eluting enantiomer (example 10A; 227 mg, 37%; orange foam, MS: 474.5 (M+H)⁺), and the slower eluting enantiomer (example 10B; 211 mg, 34%; orange foam, MS: 474.5 (M+H)⁺).

- 5 The following examples were prepared in analogy to examples 10A and 10B by HPLC separation of their racemates:

No.	Starting material	Optical rotation sign	MS, m/e
11A	trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)-	(+)	474.4 (M+H) ⁺
11B	hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate (example 1.08)	(-)	474.4 (M+H) ⁺
12A	trans-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)-	(+)	474.5 (M+H) ⁺
12B	hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (example 4.10)	(-)	474.5 (M+H) ⁺

Example 13

- 10 (E)-1-[trans-2-(1H-Benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(3,5-dichloro-phenyl)-prop-2-en-1-one




- A solution of trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate (example 1.08; 105 mg, 221 μmol) in ethanol (2 mL) was stirred for 72 h under a hydrogen atmosphere (1 bar) in the presence of palladium (10% on carbon, 100 mg, 94 μmol), then insoluble material was removed by filtration through
 15 diatomaceous earth. The filtrate was evaporated, taken up in hydrochloric acid solution (5-6 M

-145-

in 2-propanol, 1 mL), then after 1 h concentrated in vacuo and the residue was triturated in ethyl acetate to produce an off-white solid (32 mg). This material was dissolved in N,N-dimethylformamide (1 mL), then 4-methylmorpholine (44.8 mg, 443 μmol), 3,5-dichlorocinnamic acid (19.2 mg, 88.5 μmol) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (50.5 mg, 133 μmol) were added at room temperature, then after 18 h the reaction mixture was partitioned between sat. aq. ammonium chloride solution and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel, gradient dichlormethane to dichlormethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (9 mg, 9%). Colourless gum, MS: 470.5 (M+H)⁺.

Example 14


(1H-Benzotriazol-5-yl)-{trans-2-[5-(4-chloro-phenyl)-[1,3,4]oxadiazol-2-yl]-octahydro-pyrrolo[3,4-c]pyridin-5-yl}-methanone



To a solution of 5-(4-chlorophenyl)-1,3,4-oxadiazol-2(3H)-one (CAS-RN 1711-61-1; 30 mg, 153 μmol) and N,N-diisopropylethylamine (98.6 mg, 763 μmol) in N,N-dimethylformamide (3.00 mL) was added (1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone hydrochloride (intermediate 2.1; 51.7 mg, 168 μmol) at room temperature, then after 10 min benzotriazol-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate (75.8 mg, 168 μmol) was added. After 16 h the reaction mixture was partitioned between water and ethyl acetate. The organic layer was washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (42 mg, 61%). White solid, MS: 450.4 (M+H)⁺.




Example 15




(E)-1-((3aR,6aS)-5-(1H-Benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one









- To a solution of (1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2; 30 mg, 102 μ mol), 4-methylmorpholine (51.6 mg, 511 μ mol) and (E)-3-(4-(trifluoromethoxy)phenyl)acrylic acid (23.7 mg, 102 μ mol) in N,N-dimethylformamide (1.5 mL) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (38.8 mg, 102 μ mol) at 0°C, then the reaction mixture was allowed to reach room temperature over a period of 16 h. After partitioning between ethyl acetate and sat. aq. sodium hydrogen carbonate solution the organic layer was washed with water and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (38 mg, 79%). White foam, MS: 472.4 (M+H)⁺.
- 15 The following compounds were produced in analogy to example 15, replacing (1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride and (E)-3-(4-(trifluoromethoxy)phenyl)acrylic acid by the appropriate amine and carboxylic acid, respectively.




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.01	(1H-benzotriazol-5-yl)-[(3aR,6aS)-5-(5-chloro-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	5-chloro-1H-indole-2-carboxylic acid	435.5 (M+H) ⁺
15.02	(E)-1-[(3aR,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-5-trifluoromethyl-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	(E)-3-(3-fluoro-5-(trifluoromethyl)-phenyl)acrylic acid	472.5 (M-H) ⁻
15.03	1-[(3aR,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-5-trifluoromethyl-phenyl)-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	3-(3-fluoro-5-(trifluoromethyl)-phenyl)-propanoic acid	474.5 (M-H) ⁻




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.04	(1H-benzotriazol-5-yl)-[(3aR,6aS)-5-(6-chloro-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	6-chloro-1H-indole-2-carboxylic acid	435.5 (M+H) ⁺
15.05	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylsulfonyl)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-(trifluoromethylsulfonyl)-phenyl)acrylic acid (CAS-RN 910654-44-3)	548.4 (M+H) ⁺
15.06	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-chlorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-chlorophenyl)-acrylic acid	450.4 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.07	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-p-tolylprop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-p-tolylacrylic acid	430.5 (M+H) ⁺
15.08	4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)-N,N-dimethylbenzenesulfonamide 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-(N,N-dimethylsulfamoyl)-phenyl)acrylic acid	523.4 (M+H) ⁺
15.09	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxyphenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-methoxyphenyl)acrylic acid	446.6 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.10	(E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)methanone hydrochloride (intermediate 2.3)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	500.4 (M+H) ⁺
15.11	4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)benzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-cyanophenyl)-acrylic acid	441.5 (M+H) ⁺
15.12	(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	472.5 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.13	1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	3-(4-(trifluoromethoxy)-phenyl)-propanoic acid	474.4 (M+H) ⁺
15.14	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-fluorophenyl)-acrylic acid	434.4 (M+H) ⁺
15.15	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-phenylprop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	cinnamic acid	416.5 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.16	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-2-yl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(pyridin-2-yl)acrylic acid	417.5 (M+H) ⁺
15.17	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-3-yl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(pyridin-3-yl)acrylic acid	417.5 (M+H) ⁺
15.18	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chlorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(3-chlorophenyl)-acrylic acid	448.4 (M-H) ⁻




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.19	(E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(4-chlorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)methanone hydrochloride (intermediate 2.3)	(E)-3-(4-chlorophenyl)-acrylic acid	450.4 (M+H) ⁺
15.20	(E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)methanone hydrochloride (intermediate 2.3)	(E)-3-(3-(trifluoromethoxy)-phenyl)acrylic acid	500.4 (M+H) ⁺
15.21	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(difluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	3-(4-(difluoromethoxy)-phenyl)acrylic acid	482.4 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.22	(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	470.6 (M-H) ⁻
15.23	4-((E)-3-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-oxoprop-1-enyl)benzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(4-cyanophenyl)-acrylic acid	413.6 (M+H) ⁺
15.24	(E)-1-((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 8)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	472.5 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.25	(-)-(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)(trans-tetrahydro-1H-pyrrolo[3,4-c]pyridin-5(6H,7H,7aH)-yl)methanone hydrochloride, enantiomer A (intermediate 7A)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	486.4 (M+H) ⁺
15.26	(+)-(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aS,7aS)-tetrahydro-1H-pyrrolo[3,4-c]pyridin-5(6H,7H,7aH)-yl)methanone hydrochloride, enantiomer B (intermediate 7B)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	486.4 (M+H) ⁺
15.27	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3,5-dichlorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(3,5-dichloro-phenyl)acrylic acid	484.5 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.28	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-4-yl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(pyridin-4-yl)acrylic acid	417.5 (M+H) ⁺
15.29	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2,4-difluorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(2,4-difluorophenyl)acrylic acid	452.6 (M+H) ⁺
15.30	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2,4-dichlorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(2,4-dichlorophenyl)acrylic acid	484.5 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.31	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3,4-dichlorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(3,4-dichlorophenyl)acrylic acid	484.3 (M+H) ⁺
15.32	(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-difluoromethoxyphenyl)-prop-2-en-1-one, enantiomer B 	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride, enantiomer B (intermediate 7B)	(E)-3-(4-(difluoromethoxy)-phenyl)acrylic acid	468.4 (M+H) ⁺
15.33	4- {(E)-3-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-oxo-propenyl} - benzonitrile, enantiomer B 	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride, enantiomer B (intermediate 7B)	(E)-3-(4-cyanophenyl)-acrylic acid	427.6 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.34	4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)-3-fluorobenzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-cyano-2-fluorophenyl)-acrylic acid (CAS-RN 669002-88-4)	459.6 (M+H) ⁺
15.35	4-((E)-3-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-oxoprop-1-enyl)-3-fluorobenzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-cyano-2-fluorophenyl)-acrylic acid (CAS-RN 669002-88-4)	431.5 (M+H) ⁺
15.36	(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(difluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-(difluoromethoxy)-phenyl)acrylic acid	454.6 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.37	(E)-1-[cis-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(1H-benzotriazol-5-yl)-cis-octahydro-pyrrolo[3,4-c]pyridin-2-yl-methanone (intermediate 6)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	486.3 (M+H) ⁺
15.38	3-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)benzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(3-cyanophenyl)-acrylic acid	441.6 (M+H) ⁺
15.39	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(2-fluoro-4-(trifluoromethoxy)-phenyl)acrylic acid (CAS-RN 1240261-81-7)	518.5 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.40	(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(2-fluoro-4-(trifluoromethoxy)-phenyl)acrylic acid (CAS-RN 1240261-81-7)	490.2 (M+H) ⁺
15.41	(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-chloro-2-fluorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(4-chloro-2-fluorophenyl)-acrylic acid	440.1 (M+H) ⁺
15.42	(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(3,5-dichloro-phenyl)acrylic acid	456.1 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.43	(E)-1-((3aR,6aS)-5-(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one 	((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone (intermediate 6.3)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	476.4 (M+H) ⁺
15.44	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(4-(trifluoromethoxy)-phenyl)-propanoic acid	474.6 (M+H) ⁺
15.45	(E)-1-[(3aS,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(6,7-dihydro-1H-[1,2,3]triazolo[4,5-c]pyridin-5(4H)-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone (intermediate 6.2)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	477.5 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.46	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-methanesulfonyl-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(3-chloro-5-(methylsulfonyl)-phenyl)acrylic acid (intermediate 23)	500.5 (M+H) ⁺
15.47	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3,5-dimethoxy-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(3,5-dimethoxy-phenyl)acrylic acid	448.5 (M+H) ⁺
15.48	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(3-chloro-5-(trifluoromethoxy)-phenyl)acrylic acid	506.4 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.49	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-methoxy-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(3-chloro-5-methoxy-phenyl)acrylic acid (intermediate 23.1)	452.5 (M+H) ⁺
15.50	3-{(E)-3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-oxo-propenyl}-5-chloro-benzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(3-chloro-5-cyanophenyl)-acrylic acid (intermediate 24.1)	447.5 (M+H) ⁺
15.51	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-methoxy-5-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(3-methoxy-5-(trifluoromethoxy)-phenyl)acrylic acid (intermediate 24)	502.5 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.52	(E)-1-[(3aR,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(6,7-dihydro-1H-[1,2,3]triazolo[4,5-c]pyridin-5(4H)-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 25)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	477.6 (M+H) ⁺
15.53	1-[(3aR,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one 	(6,7-dihydro-1H-[1,2,3]triazolo[4,5-c]pyridin-5(4H)-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 25)	3-(4-(trifluoromethoxy)-phenyl)-propanoic acid	479.6 (M+H) ⁺
15.54	(3aR,7aR)-5-[(3aR,6aR)-5-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carbonyl]-hexahydro-oxazolo[5,4-c]pyridin-2-one 	(3aR,7aR)-5-(((3aR,6aR)-octahydropyrrolo[3,4-c]pyrrole-2-carbonyl)hexahydro-oxazolo[5,4-c]pyridin-2(1H)-one hydrochloride (intermediate 25.1)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	495.6 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.55	(3aR,7aR)-5-[(3aR,6aR)-5-[3-(4-trifluoromethoxy-phenyl)-propionyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carbonyl]-hexahydro-oxazolo[5,4-c]pyridin-2-one 	(3aR,7aR)-5-[(3aR,6aR)-octahydropyrrolo[3,4-c]pyrrole-2-carbonyl]hexahydro-oxazolo[5,4-c]pyridin-2(1H)-one hydrochloride (intermediate 25.1)	3-(4-(trifluoromethoxy)-phenyl)-propanoic acid	497.7 (M+H) ⁺
15.56	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-phenyl-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	cinnamic acid	388.6 (M+H) ⁺
15.57	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-phenyl-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-phenyl-propanoic acid	390.6 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.58	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(4-(trifluoro-methyl)-phenyl)acrylic acid	456.7 (M+H) ⁺
15.59	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahdropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(4-(trifluoro-methyl)-phenyl)-propanoic acid	458.7 (M+H) ⁺
15.60	(3aR,6aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-5-(3-(4-(trifluoromethoxy)phenyl)propanoyl)hexahdropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide 	(3aR,6aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methylhexahdropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide 2,2,2-trifluoroacetate (intermediate 20.1)	3-(4-(trifluoro-methoxy)-phenyl)-propanoic acid	467.6 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.61	(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acryloyl)-N-methyloctahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide	(E)-3-(3-fluoro-4-(trifluoromethoxy)-phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.03)	N-methyl-1-(1H-1,2,3-triazol-4-yl)methan-amine (CAS-RN 1248059-33-7)	511.6 (M+H) ⁺
15.62	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-trifluoromethyl-pyridin-3-yl)-propan-1-one	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(6-(trifluoromethyl)-pyridin-3-yl)propanoic acid (CAS-RN 539855-70-4)	459.5 (M+H) ⁺
15.63	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-trifluoromethoxy-phenoxy)-ethanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-(trifluoromethoxy)-phenoxy)acetic acid	476.5 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.64	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-chloro-2-isopropyl-5-methylphenoxy)acetic acid (CAS-RN 5411-11-0)	482.6 (M+H) ⁺
15.65	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-biphenyl-4-ylpropan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(biphenyl-4-yl)propanoic acid	466.6 (M+H) ⁺
15.66	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluoro-2-(trifluoromethyl)phenyl)prop-2-en-1-one 	(E)-3-(4-fluoro-2-(trifluoromethyl)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one (intermediate 26.04)	1H-benzo[d][1,2,3]triazole-5-carboxylic acid	502.6 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.67	1-[(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone 	2-(4-chloro-2-isopropyl-5-methylphenoxy)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)ethanone hydrochloride (intermediate 5.2)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	482.7 (M+H) ⁺
15.68	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(methylsulfonyl)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-(methylsulfonyl)-phenyl)acrylic acid	494.4 (M+H) ⁺
15.69	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylthio)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-(trifluoromethylthio)-phenyl)acrylic acid	516.4 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.70	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	500.4 (M+H) ⁺
15.71	1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-2-(3-(trifluoromethoxy)phenoxy)ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)(trans-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.2)	2-(3-(trifluoromethoxy)-phenoxy)acetic acid (CAS-RN 836-33-9)	476.5 (M+H) ⁺
15.72	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone (intermediate 6.1)	(E)-3-(3-(trifluoromethoxy)-phenyl)acrylic acid	500.4 (M+H) ⁺



Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.73	(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride (intermediate 2.1)	(E)-3-(3-(trifluoromethoxy)-phenyl)acrylic acid	486.4 (M+H) ⁺
15.74	(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride (intermediate 2.1)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	486.5 (M+H) ⁺
15.75	(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3-chloro-5-trifluoromethoxy-phenyl)-prop-2-en-1-one 	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride (intermediate 2.1)	(E)-3-(3-chloro-5-(trifluoromethoxy)-phenyl)acrylic acid	520.4 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.76	(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3,5-dichloro-phenyl)-prop-2-en-1-one 	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride (intermediate 2.1)	(E)-3-(3,5-dichloro-phenyl)acrylic acid	470.4 (M+H) ⁺
15.77	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-phenyl-pyridin-3-yl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.04)	(E)-3-(6-phenylpyridin-3-yl)acrylic acid (intermediate 23.05)	465.5 (M+H) ⁺
15.78	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-trifluoromethyl-pyridin-2-yl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.04)	(E)-3-(5-(trifluoromethyl)-pyridin-2-yl)acrylic acid (CAS-RN 910654-24-9)	457.5 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.79	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-4-yl-phenyl)-prop-2-en-1-one	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(4-(pyridin-4-yl)phenyl)-acrylic acid (intermediate 23.07)	465.6 (M+H) ⁺
15.80	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-3-yl-phenyl)-prop-2-en-1-one	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(4-(pyridin-3-yl)phenyl)-acrylic acid (intermediate 23.08)	465.6 (M+H) ⁺
15.81	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-2-yl-phenyl)-prop-2-en-1-one	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(4-(pyridin-2-yl)phenyl)-acrylic acid (intermediate 23.09)	465.5 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.82	1-[(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-3-methylphenoxy)-ethanone 	2-(4-chloro-3-methylphenoxy)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)ethanone hydrochloride (intermediate 31.1)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	440.7 (M+H) ⁺
15.83	1-[(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-methylphenoxy)-ethanone 	2-(4-chloro-2-methylphenoxy)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)ethanone hydrochloride (intermediate 31)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	440.7 (M+H) ⁺
15.84	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-phenyl-pyridin-2-yl)-prop-2-en-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	(E)-3-(5-phenylpyridin-2-yl)acrylic acid (intermediate 23.06)	465.5 (M+H) ⁺


Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.85	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-phenyl-pyridin-3-yl)-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(6-phenylpyridin-3-yl)propanoic acid (intermediate 35.1)	467.6 (M+H) ⁺
15.86	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-isopropyl-phenoxy)-ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-isopropyl-phenoxy)acetic acid	434.6 (M+H) ⁺
15.87	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-trifluoromethyl-phenoxy)-ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-(trifluoromethyl)-phenoxy)acetic acid	460.5 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.88	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(biphenyl-2-yloxy)-ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(biphenyl-2-yloxy)acetic acid	468.5 (M+H) ⁺
15.89	(E)-1-[(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propenone 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	476.6 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.90	1-((3aR,6aR)-5-((R)-4,5,6,7-tetrahydro-1H-benzo[d]-[1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	3-(4-(trifluoromethoxy)-phenyl)-propanoic acid	478.5 (M+H) ⁺
15.91	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-chloro-4-trifluoromethoxy-phenoxy)-ethanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-chloro-4-(trifluoromethoxy)-phenoxy)acetic acid (intermediate 33.3)	510.6 (M+H) ⁺
15.92	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-pyrrol-1-yl-phenoxy)-ethanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-(1H-pyrrol-1-yl)phenoxy)-acetic acid (intermediate 33.2)	457.6 (M+H) ⁺


Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.93	4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-methoxy-benzonitrile	(1H-benzo[d]-[1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-cyano-2-methoxy-phenoxy)acetic acid (CAS-RN 115109-49-4)	447.5 (M+H) ⁺
15.94	4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-benzonitrile	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-cyano-phenoxy)acetic acid	417.5 (M+H) ⁺
15.95	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-phenoxy-ethanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-phenoxyacetic acid	392.6 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.96	2-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-5-trifluoromethoxy-benzonitrile	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-cyano-4-(trifluoromethoxy)-phenoxy)acetic acid (intermediate 34.2)	501.5 (M+H) ⁺
15.97	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(2-isopropyl-5-methylphenoxy)ethanone	2-(2-isopropyl-5-methylphenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone (intermediate 36.02)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	476.7 (M+H) ⁺
15.98	(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(6-trifluoromethoxy-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	6-(trifluoromethoxy)-1H-indole-2-carboxylic acid	485.4 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
15.99	(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(5-trifluoromethoxy-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	5-(trifluoromethoxy)-1H-indole-2-carboxylic acid	485.4 (M+H) ⁺

Example 16

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylsulfinyl)phenyl)prop-2-en-1-one



5


To solution of (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylthio)phenyl)prop-2-en-1-one (example 15.69; 25 mg, 48.5 μmol) in acetic acid (1 mL) was added 35% aq. hydrogen peroxide solution (9.43 mg, 97.0 μmol). The reaction mixture was heated at 95°C, then after 3 h another portion of 35% aq.

10 hydrogen peroxide solution (18.9 mg, 194 μmol) was added. After another 15 h the reaction mixture was partitioned between ethyl acetate and sat. aq. sodium hydrogencarbonate solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (11 mg, 43%). White foam, MS:

15 532.4 (M+H)⁺.


Example 17




1-((3aR,8aS)-2-(1H-Benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)propan-1-one










A solution of (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one (example 1.15; 63 mg, 122 μ mol) in methanol (4 mL) was stirred at room temperature under a hydrogen atmosphere (1 bar) in the presence of palladium (10% on activated charcoal, 39 mg, 0.37 mmol), then after 18 h insoluble material was removed by filtration through diatomaceous earth. The filtrate was evaporated and purified by chromatography (silica gel; heptane – ethyl acetate gradient) to produce the title compound (48 mg, 74%). White foam, MS: 520.7 (M+H)⁺.





The following examples were produced in analogy to example 17, replacing of (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one by the appropriate starting material.




Ex.	Systematic Name	Starting material	MS
17.01	(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-6-(3-(4-(trifluoromethoxy)phenyl)propanoyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide 	(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide (example 9.07)	495.6 (M+H) ⁺



Ex.	Systematic Name	Starting material	MS
17.02	(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-(3-(3-fluoro-4-(trifluoromethoxy)phenyl)-propanoyl)-N-methyloctahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide 	(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)-acryloyl)-N-methyloctahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide (example 15.61)	513.6 (M+H)⁺
17.03	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-difluoromethoxy-phenyl)-propan-1-one 	(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(difluoromethoxy)phenyl)-prop-2-en-1-one (example 15.36)	456.5 (M+H)⁺
17.04	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(2-fluoro-4-trifluoromethoxy-phenyl)-propan-1-one 	(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)-prop-2-en-1-one (example 15.40)	492.7 (M+H)⁺

Ex.	Systematic Name	Starting material	MS
17.05	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluoro-2-(trifluoromethyl)phenyl)propan-1-one 	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydro-pyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluoro-2-(trifluoromethyl)phenyl)-prop-2-en-1-one (example 15.66)	504.6 (M+H) ⁺
17.06	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-methyl-4-(trifluoromethoxy)phenyl)-propan-1-one 	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydro-pyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-methyl-4-(trifluoromethoxy)phenyl)-prop-2-en-1-one (example 1.18)	516.7 (M+H) ⁺
17.07	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-methoxyphenyl)propan-1-one 	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydro-pyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-methoxyphenyl)prop-2-en-1-one (example 1.19)	466.7 (M+H) ⁺

Ex.	Systematic Name	Starting material	MS
17.08	1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(2-isopropyl-phenyl)-propan-1-one 	(E)-1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(2-isopropyl-phenyl)-prop-2-en-1-one (example 1.22)	460.7 (M+H) ⁺
17.09	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-trifluoromethyl-pyridin-2-yl)-propan-1-one 	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-trifluoromethyl-pyridin-2-yl)-prop-2-en-1-one (example 15.78)	459.5 (M+H) ⁺
17.10	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-phenyl-pyridin-2-yl)-propan-1-one 	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-phenyl-pyridin-2-yl)-prop-2-en-1-one (example 15.84)	467.6 (M+H) ⁺
17.11	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-4-yl-phenyl)-propan-1-one 	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-4-yl-phenyl)-prop-2-en-1-one (example 15.79)	467.5 (M+H) ⁺


Ex.	Systematic Name	Starting material	MS
17.12	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-3-yl-phenyl)-propan-1-one 	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-3-yl-phenyl)-prop-2-en-1-one (example 15.80)	467.6 (M+H) ⁺
17.13	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-2-yl-phenyl)-propan-1-one 	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-2-yl-phenyl)-prop-2-en-1-one (example 15.81)	467.5 (M+H) ⁺
17.14	(3aS,8aR)-6-[3-(4-trifluoromethoxy-phenyl)-propionyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid [2-(3H-[1,2,3]triazol-4-yl)-ethyl]-amide 	(3aR,8aS)-N-(2-(1H-1,2,3-triazol-5-yl)ethyl)-6-((E)-3-(4-(trifluoromethoxy)-phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide (example 9.09)	495.2 (M+H) ⁺
17.15	1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one 	(E)-1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propenone (example 1.26)	488.7 (M+H) ⁺

Ex.	Systematic Name	Starting material	MS
17.16	1-[trans-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one 	(E)-1-[trans-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propenone (example 6.07)	474.4 (M+H) ⁺
17.17	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chloro-5-(trifluoromethyl)phenyl)propan-1-one 	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chloro-5-(trifluoromethyl)phenyl)prop-2-en-1-one (example 1.27)	520.6 (M+H) ⁺
17.18	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxy-2-(trifluoromethyl)phenyl)propan-1-one 	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxy-2-(trifluoromethyl)phenyl)prop-2-en-1-one (example 1.28)	516.7 (M+H) ⁺

Ex.	Systematic Name	Starting material	MS
17.19	1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-cyclopropylphenyl)propan-1-one 	(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-cyclopropylphenyl)prop-2-en-1-one (example 1.29)	458.7 (M+H) ⁺
17.20	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[3-methoxy-5-(trifluoromethoxy)phenyl]propan-1-one 	(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-methoxy-5-trifluoromethoxy-phenyl)-prop-2-ene-1-one (example 15.51)	504.4 (M+H) ⁺

Example 18

(E)-3-[4-(Trifluoro-methoxy)-phenyl]-1-[(3aS,8aR)-2-((S)-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridine-6-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-prop-2-en-1-one hydrochloride



Step 1: (S)-Di-tert-butyl 6-((3aR,8aS)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)-decahydropyrrolo[3,4-d]azepine-2-carbonyl)-6,7-dihydro-1H-imidazo[4,5-c]pyridine-1,5(4H)-dicarboxylate

- 10 The title compound was produced in analogy to example 15 from (E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one

-188-

hydrochloride (intermediate 5) and (S)-1,5-bis(tert-butoxycarbonyl)-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridine-6-carboxylic acid (CAS-RN 175289-42-6). White foam, MS: 704.3 (M+H)⁺.

Step 2: (E)-3-[4-(Trifluoro-methoxy)-phenyl]-1-[(3aS,8aR)-2-((S)-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridine-6-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-prop-2-en-1-one hydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from (S)-di-tert-butyl 6-(((3aR,8aS)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)decahydropyrrolo[3,4-d]azepine-2-carbonyl)-6,7-dihydro-1H-imidazo[4,5-c]pyridine-1,5(4H)-dicarboxylate. White solid, MS: 504.2 (M+H)⁺.




Example 19

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-isopropyl-5-methyl-phenoxy)-ethanone




To a suspension of (1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.04; 25 mg, 85.1 μmol, Eq: 1.00), N-methylmorpholine (43.0 mg, 426 μmol) and 2-(2-isopropyl-5-methylphenoxy)acetic acid (17.7 mg, 85.1 μmol) in N,N-dimethylformamide (4 mL) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (32.4 mg, 85.1 μmol) at 0°C, then the reaction mixture was allowed to reach room temperature over a period of 16 h. After partitioning between ethyl acetate and sat. aq. sodium hydrogen carbonate solution the organic layer was washed with water and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (37 mg, 97%). White foam, MS: 448.6 (M+H)⁺.




The following compounds were produced in analogy to example 19, replacing (1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone




hydrochloride and 2-(2-isopropyl-5-methylphenoxy)acetic acid by the appropriate amine and carboxylic acid, respectively.




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.01	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-bromo-4-trifluoromethoxy-phenoxy)-ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-bromo-4-(trifluoromethoxy)-phenoxy)acetic acid (intermediate 33.1)	554.3 (M+H) ⁺
19.02	(1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(4'-chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4'-chlorobi-phenyl-4-carboxylic acid	472.5 (M+H) ⁺
19.03	4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-isopropyl-benzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-cyano-2-isopropyl-phenoxy)acetic acid (intermediate 34.1)	459.6 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.04	2-(2-acetyl-phenoxy)-1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-ethanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-acetyl-phenoxy)acetic acid (CAS-RN 1878-62-2)	434.6 (M+H) ⁺
19.05	4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-5-isopropyl-2-methyl-benzonitrile	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-cyano-2-isopropyl-5-methyl-phenoxy)acetic acid (intermediate 34)	473.7 (M+H) ⁺
19.06	(1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-naphthoic acid	410.6 (M-H) ⁻



Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.07	(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(4-methoxy-naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-methoxy-2-naphthoic acid (CAS-RN 5773-93-3)	442.6 (M+H) ⁺
19.08	4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-ethoxy-benzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-cyano-2-ethoxy-phenoxy)acetic acid (CAS-RN 835888-68-1)	461.7 (M+H) ⁺
19.09	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-4-trifluoromethoxy-phenyl)-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(3-fluoro-4-(trifluoro-methoxy)-phenyl)-propanoic acid (intermediate 35)	492.4 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.10	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-phenoxy)-ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-chloro-2-isopropyl-phenoxy)acetic acid (CAS-RN 109042-01-5)	468.5 (M+H) ⁺
19.11	[(3aS,6aS)-5-(4'-chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-(R)-4,5,6,7-tetrahydro-1H-benzotriazol-5-yl-methanone 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	4'-chlorobi-phenyl-4-carboxylic acid	476.5 (M+H) ⁺
19.12	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 8)	3-(4-(trifluoromethoxy)-phenyl)-propanoic acid	474.4 (M+H) ⁺



Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.13	(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(4'-chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H-yl)methanone dihydrochloride (intermediate 8)	4'-chloro-biphenyl-4-carboxylic acid	472.6 (M+H) ⁺
19.14	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-[2-(tetrahydro-furan-2-yl)-phenoxy]-ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-(tetrahydro-furan-2-yl)-phenoxy)acetic acid (intermediate 33)	462.5 (M+H) ⁺
19.15	(1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(4-methoxy-naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H-yl)methanone dihydrochloride (intermediate 8)	4-methoxy-2-naphthoic acid (CAS-RN 5773-93-3)	442.6 (M+H) ⁺



Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.16	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-tert-butyl-phenoxy)-ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-tert-butyl-phenoxy)acetic acid (CAS-RN 19271-90-0)	448.5 (M+H) ⁺
19.17	[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-[trans-4-(4-chlorophenyl)-cyclohexyl]-methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	trans-4-(4-chlorophenyl)-cyclohexane-carboxylic acid	478.2 (M+H) ⁺
19.18	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-4-trifluoromethyl-phenyl)-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(3-fluoro-4-(trifluoromethyl)-phenyl)-propanoic acid	474.5 (M-H) ⁻




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.19	1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(2-fluoro-4-trifluoromethyl-phenyl)-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H-yl)methanone dihydrochloride (intermediate 2.4)	3-(2-fluoro-4-(trifluoromethyl)-phenyl)-propanoic acid	474.5 (M-H) ⁻
19.20	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(2-pyridin-3-ylphenoxy)ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-(pyridin-3-yl)phenoxy)-acetic acid (intermediate 34.3)	467.6 (M-H) ⁻
19.21	4-[3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-oxopropyl]-2-methyl-5-propan-2-ylbenzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H-yl)methanone dihydrochloride (intermediate 2.4)	3-(4-cyano-2-isopropyl-5-methyl-phenyl)-propanoic acid (intermediate 42.1)	471.7 (M+H) ⁺



Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.22	4-[3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-oxopropyl]-3-propan-2-ylbenzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(4-cyano-2-isopropylphenyl)-propanoic acid (intermediate 42)	457.7 (M+H) ⁺
19.23	[(3aR,6aR)-5-[1-(4-chlorophenyl)piperidine-4-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(1H-benzotriazole-5-yl)methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	1-(4-chlorophenyl)-piperidine-4-carboxylic acid (CAS-RN 845645-46-7)	479.6 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.24	[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(4-propan-2-yloxynaphthalen-2-yl)methanone	(((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	4-isopropoxy-2-naphthoic acid (CAS-RN 1368865-02-4)	474.5 (M+H) ⁺
19.25	[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(4-propan-2-yloxyquinolin-2-yl)methanone	(((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	4-isopropoxy-quinoline-2-carboxylic acid (CAS-RN 1406553-19-2)	475.5 (M+H) ⁺



Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.26	1-[(3aR,6aR)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[2-fluoro-4-(trifluoromethoxy)phenyl]-propan-1-one 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	3-(2-fluoro-4-(trifluoromethoxy)-phenyl)-propanoic acid (intermediate 35.4)	496.4 (M+H) ⁺
19.27	4-[2-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-2-methyl-5-propan-2-ylbenzonitrile 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	2-(4-cyano-2-isopropyl-5-methyl-phenoxy)acetic acid (intermediate 34)	477.5 (M+H) ⁺




Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.28	[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-[1-(2,2,2-trifluoroethoxy)isoquinolin-3-yl]methanone 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	1-(2,2,2-trifluoroethoxy)-isoquinoline-3-carboxylic acid (CAS-RN 1096982-79-4)	515.4 (M+H)⁺
19.29	1-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(4-bromo-2-tert-butylphenoxy)ethanone 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	2-(4-bromo-2-tert-butylphenoxy)acetic acid (CAS-RN 425372-86-7)	530.4 (M+H)⁺


Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.30	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(4-bromo-2-tert-butylphenoxy)ethanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(4-bromo-2-tert-butylphenoxy)acetic acid (CAS-RN 425372-86-7)	526.6 (M+H) ⁺
19.31	4-[2-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-3-tert-butylbenzonitrile 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	2-(2-tert-butyl-4-cyano-phenoxy)acetic acid (intermediate 40)	477.7 (M+H) ⁺
19.32	4-[2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-3-tert-butylbenzonitrile 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-tert-butyl-4-cyano-phenoxy)acetic acid (intermediate 40)	471.7 (M-H) ⁻

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.33	[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-[1-methyl-5-(trifluoromethoxy)indol-2-yl]methanone 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	1-methyl-5-(trifluoromethoxy)-1H-indole-2-carboxylic acid (CAS-RN 1257122-42-1)	501.5 (M-H)⁻
19.34	1-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-[4-(trifluoromethoxy)phenoxy]ethanone 	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	2-(4-(trifluoromethoxy)-phenoxy)acetic acid	478.5 (M-H)⁻

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.35	[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(1-ethoxyisoquinolin-3-yl)methanone	[(3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone hydrochloride (intermediate 2.5)	1-ethoxy-isoquinoline-3-carboxylic acid (CAS-RN 1094758-39-0)	461.5 (M+H) ⁺
19.36	1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(2-tert-butyl-4-methoxyphenoxy)ethanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	2-(2-tert-butyl-4-methoxyphenoxy)acetic acid	476.5 (M+H) ⁺
19.37	[(3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl](4-ethoxyquinolin-2-yl)methanone	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-ethoxy-quinoline-2-carboxylic acid (CAS-RN 40609-78-7)	457.6 (M+H) ⁺


Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.38	((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)(4-(2,2,2-trifluoroethoxy)quinolin-2-yl)methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	4-(2,2,2-trifluoroethoxy)-quinoline-2-carboxylic acid (CAS-RN 1281584-65-3)	511.6 (M+H) ⁺
19.39	((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)(6-cyclobutoxy-5-(trifluoromethyl)-pyridin-3-yl)methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	6-cyclobutoxy-5-(trifluoromethyl)-nicotinic acid	501.2 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.40	((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-bromo-6-(2-methoxyethoxy)pyridin-3-yl)methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	5-bromo-6-(2-methoxyethoxy)-nicotinic acid (CAS-RN 912454-34-3)	515.2 (M+H) ⁺
19.41	((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-bromo-6-(cyclopropylmethoxy)pyridin-3-yl)methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	5-bromo-6-(cyclopropylmethoxy)-nicotinic acid (CAS-RN 912454-38-7)	511.1 (M+H) ⁺
19.42	((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-cyclopropyl-6-(2,2,2-trifluoroethoxy)pyridin-3-yl)methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	5-cyclopropyl-6-(2,2,2-trifluoroethoxy)-nicotinic acid (CAS-RN 1427064-90-1)	501.5 (M+H) ⁺

Ex.	Systematic Name	Amine	Carboxylic acid	MS, m/e
19.43	((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)(6-(2,2,2-trifluoroethoxy)-5-(trifluoromethyl)pyridin-3-yl)methanone 	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride (intermediate 2.4)	6-(2,2,2-trifluoroethoxy)-5-(trifluoromethyl)-nicotinic acid	529.5 (M+H) ⁺

Example 20

(1H-Benzotriazol-5-yl)-{(3aS,6aS)-5-[4-(4-chloro-phenyl)-piperidine-1-carbonyl]-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl}-methanone




To a white suspension of (1H-benzo[d][1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride (intermediate 2.4; 40 mg, 136 μmol) and triethylamine (68.9 mg, 681 μmol) in dichloromethane (4 mL) was added a solution of 4-(4-chlorophenyl)piperidine-1-carbonyl chloride (intermediate 9.3; 44.9 mg, 163 μmol) in

10 dichloromethane (2 mL) at room temperature, then after 19 the reaction mixture was partitioned between sat. aq. sodium hydrogencarbonate solution and dichloromethane. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated.

Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (50 mg, 77%). White foam, MS:

15 479.6 (M+H)⁺.

The following example was produced in analogy to example 20, replacing 4-(4-chlorophenyl)piperidine-1-carbonyl chloride by the appropriate reagent.

Ex.	Systematic Name	Reagent	MS
20.01	(1H-benzotriazol-5-yl)-{(3aS,6aS)-5-[4-(4-chloro-phenyl)-piperazine-1-carbonyl]-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl}-methanone 	4-(4-chlorophenyl)piperazine-1-carbonyl chloride (CAS-RN 64985-84-8)	480.4 (M+H) ⁺

5

Intermediates

Intermediate 1

(3aR,6aS)-3,5-Dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride

- 10 Step 1: (3aR,6aS)-2-tert-Butyl 5-(3,5-dichlorobenzyl) tetrahydropyrrolo[3,4-c]pyrrole-2,5(1H,3H)-dicarboxylate

To a light brown solution of (3,5-dichlorophenyl)methanol (425 mg, 2.35 mmol) in dichloromethane (7 mL) was added N,N'-carbonyldiimidazole (401 mg, 2.47 mmol). The solution was stirred at room temperature for 3 h, then (3aR,6aS)-tert-butyl

- 15 hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (CAS-RN 250275-15-1; 526 mg, 2.35 mmol) was added, then after 15 h the reaction mixture was partitioned between 1 M aq. hydrochloric acid solution and dichloromethane. The organic layer was washed with brine, dried over magnesium sulfate, filtered and evaporated to afford the title compound (972 mg, 99%). Light brown viscous oil, MS: 359.2 (M+H-isobutene)⁺.

Step 2: (3aR,6aS)-3,5-Dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride

To a solution of (3aR,6aS)-2-tert-butyl 5-(3,5-dichlorobenzyl) tetrahydropyrrolo[3,4-c]pyrrole-2,5(1H,3H)-dicarboxylate (962 mg, 2.32 mmol) in 2-propanol (4 mL) was added hydrochloric acid (5–6 M in 2-propanol) (11.6 mL, 57.9 mmol), then after 3 h the reaction mixture was evaporated. The residue was taken up in ethyl acetate and a few drops of ethanol, then the precipitate was collected by filtration to produce the title compound (738 mg, 91%). White solid, MS: 315.3 (M+H)⁺.

- 10 The following intermediates were prepared according to intermediate 1, replacing (3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate and (3,5-dichlorophenyl)methanol by the appropriate amine and alcohol, respectively.

No.	Systematic Name	Amine	Alcohol	MS, m/e
1.1	(3aS,7aS)-3,5-dichlorobenzyl hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate hydrochloride	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	(3,5-dichlorophenyl)methanol	329.4 (M+H) ⁺
1.2	cis-3,5-dichlorobenzyl hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate hydrochloride	cis-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (intermediate 14)	(3,5-dichlorophenyl)methanol	329.4 (M+H) ⁺

No.	Systematic Name	Amine	Alcohol	MS, m/e
1.3	(3aR,8aS)-3,5-dichlorobenzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate hydrochloride	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester (CAS-RN 1251013-07-6)	(3,5-dichlorophenyl)methanol	343.4 (M+H) ⁺
1.4	trans-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 4-trifluoromethoxy-benzyl ester hydrochloride	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	(4-(trifluoromethoxy)phenyl)-methanol	345.6 (M+H) ⁺
1.5	trans-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester hydrochloride	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	(3-chloro-5-(methanesulfonyl)-phenyl)methanol (intermediate 17)	373.6 (M+H) ⁺
1.6	(3aR,8aS)-4-(2,2,2-trifluoroethoxy)benzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate hydrochloride	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester (CAS-RN 1251013-07-6)	(4-(2,2,2-trifluoroethoxy)-phenyl)methanol (CAS-RN 1020949-12-5)	n.a.

No.	Systematic Name	Amine	Alcohol	MS, m/e
1.7	(3aR,8aS)-2-fluoro-4-(2,2,2-trifluoroethoxy)benzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate hydrochloride	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester (CAS-RN 1251013-07-6)	(2-fluoro-4-(2,2,2-trifluoroethoxy)-phenyl)methanol (intermediate 38)	n.a.
1.8	(3aS,6aS)-4-(trifluoromethoxy)benzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride	(3aR,6aR)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 15.1)	(4-(trifluoromethoxy)-phenyl)methanol	331.5 (M+H) ⁺

Intermediate 2

(1H-Benzotriazol-5-yl)-cis-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone hydrochloride

Step 1: cis-tert-Butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate

To a solution of cis-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (intermediate 14; 50 mg, 210 μ mol) in N,N-dimethylformamide (1 mL) were added 4-methylmorpholine (84.9 mg, 840 μ mol), 1H-benzo[d][1,2,3]triazole-5-carboxylic acid (36.0 mg, 220 μ mol) and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (120 mg, 315 μ mol) at room temperature, then after 16 h the reaction mixture was partitioned between ethyl acetate and sat. aq. ammonium chloride solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (52 mg, 67%). Light yellow gum, MS: 370.5 (M-H)⁻.

Step 2: (1H-Benzotriazol-5-yl)-cis-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from (3aS,7aS)-tert-butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate. White solid, MS: 272.5 (M+H)⁺.

- 5 The following intermediates were prepared according to intermediate 2, replacing (3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate and 1H-benzo[d][1,2,3]triazole-5-carboxylic acid by the appropriate amine and carboxylic acid, respectively.

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
2.1	(1H-benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone; hydrochloride	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	272.5 (M+H) ⁺
2.2	(1H-benzo[d]-[1,2,3]triazol-5-yl)(trans-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride	(3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (CAS-RN 250275-15-1)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	258.5 (M+H) ⁺
2.3	(1H-benzo[d]-[1,2,3]triazol-5-yl)-((3aR,8aS)-octahydro-pyrrolo[3,4-d]azepin-6(7H)-yl)methanone hydrochloride	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester (CAS-RN 1251013-07-6)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	286.5 (M+H) ⁺

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
2.4	(1H-benzo[d]-[1,2,3]triazol-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride	(3aS,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 15)	1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid	258.5 (M+H) ⁺
2.5	((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)((R)-4,5,6,7-tetrahydro-1H-benzo[d]-[1,2,3]triazol-5-yl)-methanone hydrochloride	(3aS,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 15)	(+)-(R)-4,5,6,7-tetrahydro-1H-benzo[d]-[1,2,3]triazole-5-carboxylic acid (intermediate 30A)	262.6 (M+H) ⁺
2.6	(1H-[1,2,3]triazolo[4,5-b]pyridin-5-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride	(3aS,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 15)	1H-[1,2,3]triazolo[4,5-b]pyridine-5-carboxylic acid (CAS-RN 1216149-55-1)	257.5 (M-H) ⁻

Intermediate 3

3-(3,5-Dichlorophenyl)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)propan-1-one hydrochloride

- 5 Step 1: (3aR,6aS)-tert-Butyl 5-(3-(3,5-dichlorophenyl)propanoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

To a solution of 3-(3,5-dichlorophenyl)propanoic acid (103 mg, 470 μ mol), (3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (CAS-RN 250275-15-1; 100 mg, 471 μ mol) and 4-methylmorpholine (238 mg, 2.35 mmol) in N,N-dimethylformamide (1 mL) was added O-

(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (268 mg, 705 μ mol) at room temperature, then after 16 h the reaction mixture was partitioned between ethyl acetate and sat. aq. sodium hydrogencarbonate solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; 5 heptane – ethyl acetate gradient) afforded the title compound (161 mg, 83%). Colourless oil, MS: 357.1 (M–isobutene+H)⁺.

Step 2: 3-(3,5-Dichlorophenyl)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)propan-1-one hydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from (3aR,6aS)-tert-butyl 5-(3-(3,5-dichlorophenyl)propanoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate. White 10 solid, MS: 313.1 (M+H)⁺.

Intermediate 4

(3aR,5s,6aS)-N-((1H-1,2,3-Triazol-4-yl)methyl)octahydrocyclopenta[c]pyrrole-5-carboxamide 2,2,2-trifluoroacetate

15 Step 1: (3aR,5s,6aS)-tert-butyl 5-((1H-1,2,3-triazol-4-yl)methylcarbamoyl)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

To a solution of (3aR,5s,6aS)-2-(tert-butoxycarbonyl)octahydrocyclopenta[c]pyrrole-5-carboxylic acid (WuXi AppTec (Wuhan) Co., Ltd.; catalogue No. WX110047; 100 mg, 392 μ mol) and (1H-1,2,3-triazol-4-yl)methanamine hydrochloride (52.7 mg, 392 μ mol) in 20 dichloromethane (2 mL) were added at 0°C diisopropylethylamine (127 mg, 979 μ mol) and benzotriazol-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate (182 mg, 411 μ mol), then after 15 h at room temperature the reaction mixture was evaporated and the residue partitioned between ethyl acetate and water. The organic layer was dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel, ethyl acetate–methanol gradient) 25 produced the title compound (100 mg, 75%). White foam, MS: 334.5 (M–H)[–].

Step 2: (3aR,5s,6aS)-N-((1H-1,2,3-Triazol-4-yl)methyl)octahydrocyclopenta[c]pyrrole-5-carboxamide 2,2,2-trifluoroacetate

Trifluoroacetic acid (340 mg, 2.98 mmol) was added at room temperature to a solution of (3aR,5s,6aS)-tert-butyl 5-((1H-1,2,3-triazol-4-yl)methylcarbamoyl)hexahydrocyclo-

penta[c]pyrrole-2(1H)-carboxylate (100 mg, 298 μ mol) in dichloromethane, then after 4 h the reaction mixture was evaporated to produce the title compound, which was used directly in the next step. Light yellow oil, MS: 236.5 (M+H)⁺.

Intermediate 5

5 (E)-1-((3aR,8aS)-Octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride

Step 1: (3aR,8aS)-tert-Butyl 6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydro-pyrrolo[3,4-d]azepine-2(1H)-carboxylate

To a solution of (3aR,8aS)-tert-butyl octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate
10 hydrochloride (CAS-RN 1251013-07-6; 1.50 g, 5.42 mmol), 4-methylmorpholine (2.19 g, 21.7 mmol) and (E)-3-(4-(trifluoromethoxy)phenyl)acrylic acid (1.26 g, 5.42 mmol) in N,N-dimethylformamide (30 mL) was added added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (2.06 g, 5.42 mmol) at 0°C. After 60 min the ice bath
15 aq. sodium hydrogencarbonate solution. The organic layer was washed with sat. aq. ammonium chloride solution, water, and brine, dried over magnesium sulfate, filtered, and evaporated. The residue was triturated in heptane/ethyl acetate 9:1 to produce the title compound (2.20 g, 89%). White solid, MS: 399.5 (M+H-isobutene)⁺.

Step 2: (E)-1-((3aR,8aS)-Octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)-
20 phenyl)prop-2-en-1-one hydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from (3aR,8aS)-tert-butyl 6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate. White solid, MS: 355.5 (M+H)⁺.

25 The following intermediates were produced according to intermediate 5, replacing (3aR,8aS)-tert-butyl octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate hydrochloride and (E)-3-(4-(trifluoromethoxy)phenyl)acrylic acid by the appropriate amine and carboxylic acid precursors, respectively.

No.	Systematic Name	Amine	Carboxylic acid	MS
5.1	4-((E)-3-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)benzonitrile hydrochloride	(3aR,8aS)-tert-butyl octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(4-cyanophenyl)-acrylic acid	296.5 (M+H) ⁺
5.2	2-(4-chloro-2-isopropyl-5-methylphenoxy)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)ethanone hydrochloride	(3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (CAS-RN 250275-15-1)	2-[4-chloro-5-methyl-2-(1-methylethyl)-phenoxy]-acetic acid (CAS-RN 5411-11-0)	337.6 (M+H) ⁺
5.3	(E)-1-(trans-octahydropyrrolo[3,4-c]pyridin-5-yl)-3-(4-(trifluoromethoxy)-phenyl)prop-2-en-1-one hydrochloride	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	341.6 (M+H) ⁺
5.4	1-(trans-octahydropyrrolo[3,4-c]pyridin-5-yl)-2-(4-(trifluoromethoxy)-phenoxy)ethanone hydrochloride	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	2-(4-(trifluoromethoxy)-phenoxy)acetic acid	345.6 (M+H) ⁺

No.	Systematic Name	Amine	Carboxylic acid	MS
5.5	1-((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)-phenyl)propan-1-one dihydrochloride	(3aR,6aR)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 15.1)	3-(4-(trifluoromethoxy)-phenyl)propanoic acid	329.5 (M+H) ⁺
5.6	(4-ethoxyquinolin-2-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride	(3aR,6aR)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 15.1)	4-ethoxy-quinoline-2-carboxylic acid (CAS-RN 40609-78-7)	329.5 (M+H) ⁺

Intermediate 6

cis-3,5-Dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate

- 5 A solution of cis-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate (example 1.02; 100 mg, 211 μ mol) in methanol (3 mL) was stirred for 20 h at 100°C under a hydrogen atmosphere (10 bar) in the presence of palladium (10% on carbon, 10 mg), then insoluble material was removed by filtration through diatomaceous earth and the filtrate was evaporated to produce the title compound (62 mg), which
- 10 was used directly in the next step. Light yellow foam, MS: 272.5 (M+H)⁺.

The following intermediates were prepared according to intermediate 6, replacing cis-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate by the appropriate starting material.

No.	Systematic Name	Starting material	MS, m/e
6.1	(1H-benzo[d][1,2,3]triazol-5-yl)((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-2(1H)-yl)methanone	(3aR,8aS)-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate (example 1.04)	286.5 (M+H) ⁺
6.2	(6,7-dihydro-1H-[1,2,3]triazolo[4,5-c]pyridin-5(4H)-yl)((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone	(3aR,6aS)-3,5-dichlorobenzyl 5-(4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (example 8)	263.5 (M+H) ⁺
6.3	((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazol-5-yl)methanone	(3aR,6aS)-3,5-dichlorobenzyl 5-(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (example 1.01)	262.4 (M+H) ⁺

Intermediate 7A

(1H-Benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone hydrochloride, enantiomer A

- 5 Step 1: trans-tert-Butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate

The title compound was produced in analogy to intermediate 2, step 1 from trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5) and 1H-benzo[d][1,2,3]triazole-5-carboxylic acid. Light yellow foam, MS: 370.4 (M-H)⁻.

Step 2: (-)-trans-tert-Butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate and (+)-trans-tert-butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate

HPLC separation of racemic trans-tert-butyl 5-(1H-benzo[d][1,2,3]triazole-5-

- 5 carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (668 mg, 1.80 mmol) using a Reprosil Chiral-NR column as the stationary phase and heptane/ethanol 3:2 as the eluent produced the faster eluting (-)-enantiomer (251 mg, 37%; colourless gum, MS: 370.6 (M-H)⁻), followed by the slower eluting (+)-enantiomer (212 mg, 32%; colourless gum, MS: 370.6 (M-H)⁻).

- 10 Step 3: (1H-Benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone hydrochloride enantiomer A

The title compound was produced in analogy to intermediate 1, step 2 from (-)-trans-tert-butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate. White solid, MS: 272.5 (M+H)⁺.

15 **Intermediate 7B**

(1H-Benzotriazol-5-yl)-trans-octahydro-pyrrolo[3,4-c]pyridin-5-yl-methanone hydrochloride, enantiomer B

- The following intermediate was prepared according to intermediate 7A, step 3 from (+)-trans-tert-butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (intermediate 7A, step 2). White solid, MS: 272.5 (M+H)⁺.

Intermediate 8

(1H-Benzo[d][1,2,3]triazol-5-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride

- 25 Step 1: trans-tert-Butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

The title compound was produced in analogy to intermediate 2, step 1 from trans-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 16) and 1H-benzo[d][1,2,3]triazole-5-carboxylic acid. Light yellow foam, MS: 358.5 (M+H)⁺.

Step 2: (+)-(3aR,6aR)-tert-Butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate and (-)-(3aS,6aS)-tert-butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

HPLC separation of racemic trans-tert-butyl 5-(1H-benzo[d][1,2,3]triazole-5-

- 5 carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (790 mg, 2.21 mmol) using a Chiralpak AD column as the stationary phase and heptane/ethanol 7:3 as the eluent produced the faster eluting (+)-(R,R)-enantiomer (350 mg, 44%; light yellow foam, MS: 358.5 (M+H)⁺; e. r. 100:0), followed by the slower eluting (-)-(S,S)-enantiomer (388 mg, 49%; light yellow foam, MS: 358.5 (M+H)⁺; e. r. 4:96).

- 10 Step 3: (1H-Benzo[d][1,2,3]triazol-5-yl)((3aS,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone dihydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from (3aR,6aR)-tert-butyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate. White solid, MS: 258.5 (M+H)⁺.

15 **Intermediate 9**

(3aR,6aS)-3,5-Dichlorobenzyl 5-(chlorocarbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

- To a colourless solution of (3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride (intermediate 1; 152 mg, 432 μmol) and pyridine (106 mg, 1.34 mmol)
20 in dichloromethane (2 mL) was added dropwise a solution of triphosgene (57.7 mg, 195 μmol) in dichloromethane (2 mL) at 0°C, then after 30 min the ice bath was removed. After 16 h the reaction mixture was partitioned between 1 M aq. hydrochloric acid solution and dichloromethane. The organic layer was washed with water and brine, dried over magnesium sulfate, filtered and evaporated to afford the title compound (172 mg), which was used directly in
25 the next step. Colourless oil, MS: 401.3 (M+Na)⁺.

The following intermediates were prepared according to intermediate 9, replacing (3aR,6aS)-3,5-dichlorobenzyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate hydrochloride by the appropriate starting material.

No.	Systematic Name	Starting material	MS, m/e
9.1	(3aR,8aS)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carbonyl chloride	(E)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one hydrochloride (intermediate 5)	416 (M) ⁺
9.2	(3aR,6aR)-tert-butyl 5-(chlorocarbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate	(3aS,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 15)	274 (M) ⁺
9.3	4-(4-chlorophenyl)piperidine-1-carbonyl chloride	4-(4-chlorophenyl)piperidine hydrochloride	257.0 (M) ⁺

Intermediate 10

(3aR,5r,6aS)-3,5-Dichlorobenzyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

5 Step 1: (3aR,5r,6aS)-tert-Butyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

To a solution of (3aR,6aS)-tert-butyl 5-oxohexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (CAS-RN 146231-54-1; 4.3 g, 19.1 mmol) in methanol (100 mL) was added sodium borohydride (1.44 g, 38.2 mmol) at 0°C, then after 1 h the reaction mixture was treated with ice water and the methanol was removed under reduced pressure. The reaction mixture was partitioned between ethyl acetate and brine. The organic layer was dried over magnesium sulfate, filtered and evaporated to produce the title compound (4.38 g, 100%). Off-white solid, MS: 228.3 (M+H)⁺.

Step 2: (3aR,5r,6aS)-Octahydrocyclopenta[c]pyrrol-5-ol hydrochloride

(3aR,5r,6aS)-tert-butyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (4.37 g, 19.0 mmol) was combined with hydrochloric acid solution (5–6 M in 2-propanol) (49 mL, 245

mmol), then after 2 h the reaction mixture was evaporated und the residue was triturated in ethyl acetate to afford the title compound as an off-white solid (2.84 g, 91%).

Step 3: (3aR,5r,6aS)-3,5-Dichlorobenzyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

- 5 To a solution of (3,5-dichlorophenyl)methanol (541 mg, 3.06 mmol) in dichloromethane (10 mL) was added 1,1'-carbonyldiimidazole (520 mg, 3.21 mmol) at room temperature. Then after 3 h (3aR,5r,6aS)-octahydrocyclopenta[c]pyrrol-5-ol hydrochloride (500 mg, 3.06 mmol) and triethylamine (309 mg, 3.06 mmol) were added, then after 18 h the reaction mixture was partitioned between dichloromethane and water. The organic layer was washed with brine, dried
10 over magnesium sulfate, filtered and evaporated. Chromatography (silica gel; heptane – ethyl acetate gradient) produced the title compound (847 mg, 84%). Colourless oil, MS: 330.1 (M+H)⁺.

Intermediate 11

- 15 **(3aR,5s,6aS)-3,5-Dichlorobenzyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate**

Step 1: (3aR,5s,6aS)-3,5-Dichlorobenzyl 5-(4-nitrobenzoyloxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

- Diethyl azodicarboxylate solution (40% in toluene; 168 µL, 424 µmol) was added at room temperature to a solution of (3aR,5r,6aS)-3,5-dichlorobenzyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (intermediate 10; 140 mg, 424 µmol), 4-nitrobenzoic acid
20 (85.0 mg, 509 µmol), and triphenylphosphine (111 mg, 424 µmol) in toluene (10 mL), then after 16 h another portion of triphenylphosphine (33.4 mg, 127 µmol) and diethyl azodicarboxylate solution (40% in toluene; 50 µL, 127 µmol) was added. The reaction mixture was stirred for another 5 h at room temperature, evaporated and the residue was purified by chromatography
25 (silica gel; heptane – ethyl acetate gradient) to produce the title compound (167mg, 82%). Colourless oil, MS 479.0 (M+H)⁺.

Step 2: (3aR,5s,6aS)-3,5-Dichlorobenzyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

A mixture of (3aR,5s,6aS)-3,5-dichlorobenzyl 5-(4-nitrobenzoyloxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (163 mg, 340 μ mol) and 2 M aq. sodium hydroxide solution (1 mL, 2 mmol) in tetrahydrofuran (2 mL) was stirred at room temperature for 6 h. After evaporation of volatile material the residue was partitioned between 1 M aq. hydrochloric acid solution and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered and evaporated in vacuo to give the title compound (109 mg, 97%). Colourless oil, MS: 330.1 (M+H)⁺.

Intermediate 12

4-((3aR,5r,6aS)-2-((3,5-Dichlorobenzoyloxy)carbonyl)octahydrocyclopenta[c]pyrrol-5-yloxy)-2-hydroxybenzoic acid

Step 1: (3aR,5r,6aS)-3,5-Dichlorobenzyl 5-(3-hydroxy-4-(methoxycarbonyl)phenoxy)-hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

Diethyl azodicarboxylate solution (40% in toluene, 157 μ L, 398 μ mol) was added at room temperature to a solution of (3aR,5s,6aS)-3,5-dichlorobenzyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (101 mg, 306 μ mol), methyl 2,4-dihydroxybenzoate (68.9 mg, 398 μ mol), and triphenylphosphine (104 mg, 398 μ mol) in toluene (2 mL). Then after 18 h the reaction mixture was concentrated and the residue purified by chromatography (silica gel; heptane – ethyl acetate gradient) to produce the title compound (144 mg, 98%). Colourless gum, MS: 480.1 (M+H)⁺.

Step 2: 4-((3aR,5r,6aS)-2-((3,5-Dichlorobenzoyloxy)carbonyl)octahydrocyclopenta[c]pyrrol-5-yloxy)-2-hydroxybenzoic acid

The title compound was produced in analogy to intermediate 11, step 2 from (3aR,5r,6aS)-3,5-dichlorobenzyl 5-(3-hydroxy-4-(methoxycarbonyl)phenoxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate. Colourless gum, MS: 466.2 (M+H)⁺.

Intermediate 12.1

4-((3aR,5s,6aS)-2-((3,5-Dichlorobenzoyloxy)carbonyl)octahydrocyclopenta[c]pyrrol-5-yloxy)-2-hydroxybenzoic acid

The title compound was produced in analogy to intermediate 12 from (3aR,5r,6aS)-3,5-dichlorobenzyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (intermediate 10). Colourless gum, MS: 466.2 (M+H)⁺.

5 **Intermediate 13**

(3aR,6aS)-3,5-Dichlorobenzyl 5-oxohexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

To a stirred solution of (3aR,5r,6aS)-3,5-dichlorobenzyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (intermediate 10; 144 mg, 436 μmol) in dichloromethane (3 mL) was added 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one solution (15% in dichloromethane; 1.48 g, 523 μmol) at 0°C. Then after 2.5 h solid sodium bicarbonate (256 mg, 3.05 mmol) was added and the reaction mixture was stirred for another 5 min and was then filtered through diatomaceous earth. The filtrate was evaporated and purified by chromatography (silica gel; heptane – ethyl acetate gradient) to produce the title compound (132 mg, 88%). Colourless oil, MS⁺: 328.2 (M+H)⁺.

15 **Intermediate 14**

cis-tert-Butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate

Step 1: cis-tert-Butyl 6-oxohexahydropyrano[3,4-c]pyrrole-2(1H)-carboxylate

(3aR,6aS)-tert-Butyl 5-oxohexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (CAS-RN 146231-54-1; 5.00 g, 17.8 mmol) was dissolved in dry dichloromethane (250 mL) and then disodium hydrogen phosphate (63.0 g, 444 mmol) and 3-chloroperbenzoic acid (10.9 g, 44.4 mmol) were added. The suspension was stirred at room temperature for 72 h, then 2 M aq. sodium sulfite solution (200 mL) was added. After stirring for another 30 min, the organic layer was washed with sat. aq. sodium hydrogencarbonate solution, dried over magnesium sulfate, filtered, and evaporated to produce a mixture (4.66 g) containing the title compound (MS: 264.5 (M+Na)⁺) as the main product, along with some unreacted starting material.

Step 2: cis-tert-Butyl 3-(2-hydroxyethyl)-4-(hydroxymethyl)pyrrolidine-1-carboxylate

cis-tert-Butyl 6-oxohexahydropyrano[3,4-c]pyrrole-2(1H)-carboxylate (crude mixture from step 1; 4.46 g) was dissolved in ethanol (300 mL), then freshly powdered anhydrous calcium chloride

(3.69 g, 33.3 mmol) was added at once, then sodium borohydride (2.52 g, 66.5 mmol) was added portionwise at room temperature. After 1 h, the reaction was poured onto ice water (800 mL) and ethyl acetate (800 mL). The two phases were saturated with solid sodium chloride and stirred for 15 min, then the organic layer was separated, washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane – ethyl acetate gradient) produced the title compound (2.57 g, ca. 60% yield over 2 steps). Colourless oil, MS: 268.5 (M+Na)⁺.

Step 3: cis-tert-Butyl 3-(2-(methylsulfonyloxy)ethyl)-4-((methylsulfonyloxy)methyl)pyrrolidine-1-carboxylate

A solution of methanesulfonyl chloride (3.60 g, 31.4 mmol) in dichloromethane (5 mL) was added at 0°C to a solution of cis-tert-butyl 3-(2-hydroxyethyl)-4-(hydroxymethyl)pyrrolidine-1-carboxylate (2.57 g, 10.5 mmol) and N,N-diisopropylethylamine (8.12 g, 62.9 mmol) in dichloromethane (70 mL), then after 1 h the mixture was partitioned between ethyl acetate and sat. aq. ammonium chloride solution. The organic layer was washed with sat. aq. sodium hydrogencarbonate solution, dried over magnesium sulfate, filtered, and evaporated.

Chromatography (silica gel, heptane – ethyl acetate gradient) afforded the title compound (3.47 g, 83%). Light yellow oil, MS: 424.5 (M+Na)⁺.

Step 4: cis-tert-Butyl 5-benzylhexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate

To a solution of cis-tert-butyl 3-(2-(methylsulfonyloxy)ethyl)-4-((methylsulfonyloxy)methyl)pyrrolidine-1-carboxylate (3.05 g, 7.60 mmol) in acetonitrile (100 mL) were added phenylmethanamine (2.44 g, 22.8 mmol) and potassium carbonate (5.25 g, 38.0 mmol). The reaction mixture was heated at 95 °C for 22 h and was then partitioned between ethyl acetate and water. The organic layer was washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel, gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 95:5:0.25) produced the title compound (1.63 g, 64%). Light yellow oil, MS: 317.6 (M+H)⁺.

Step 5: cis-tert-Butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate

To a solution of cis-tert-butyl 5-benzylhexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (1.63 g, 4.89 mmol) in methanol (33 mL) was added palladium (10% on carbon; 260 mg, 245 μmol), and the reaction mixture was stirred under a hydrogen atmosphere (1 bar) at room temperature for 24 h, then insoluble material was removed by filtration through diatomaceous

earth. The filtrate was concentrated and the residue was chromatographed (silica gel, gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 95:5:0.25) to produce the title compound (895 mg, 81%). Light yellow oil, MS: 227.5 (M+H)⁺.

Intermediate 15

5 (3aS,6aS)-tert-Butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

Step 1: (3R,4R)-tert-Butyl 3,4-bis((methylsulfonyloxy)methyl)pyrrolidine-1-carboxylate

The title compound was produced in analogy to intermediate 14, step 3 from (3R,4R)-3,4-bis-hydroxymethyl-pyrrolidine-1-carboxylic acid tert-butyl ester (CAS-RN 895245-32-6). Light yellow oil, MS: 332.4 (M-isobutene+H)⁺.

10 Step 2: (3aS,6aS)-tert-Butyl 5-benzylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

The title compound was produced in analogy to intermediate 14, step 4 from (3R,4R)-tert-butyl 3,4-bis((methylsulfonyloxy)methyl)pyrrolidine-1-carboxylate. Light yellow solid, MS: 303.5 (M+H)⁺.

Step 3: (3aS,6aS)-tert-Butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

- 15** To a solution of (3aS,6aS)-tert-butyl 5-benzylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (2.22 g, 7.34 mmol) in methanol (20 mL) was added palladium (10% on carbon, 220 mg, 7.34 mmol), and the reaction mixture was stirred under a hydrogen atmosphere (1 bar) at room temperature for 24 h, then insoluble material was removed by filtration through diatomaceous earth. The filtrate was concentrated to produce the title compound (1.60 g, 100%). White waxy
- 20** solid, MS: 213.5 (M+H)⁺.

Intermediate 15.1

(3aR,6aR)-tert-Butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

- The title compound was produced in analogy to intermediate 15, replacing (3R,4R)-3,4-bis-hydroxymethyl-pyrrolidine-1-carboxylic acid tert-butyl ester by (3S,4S)-3,4-bis-hydroxymethyl-
- 25** pyrrolidine-1-carboxylic acid tert-butyl ester (CAS-RN 895245-30-4). White waxy solid, MS: 213.3 (M+H)⁺.

Intermediate 16**trans-tert-Butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate**

The title compound was produced in analogy to intermediate 15 from trans-3,4-bis(hydroxymethyl)pyrrolidine-1-carboxylic acid tert-butyl ester (CAS-RN 895245-31-5).

5 White waxy solid, MS: 213.5 (M+H)⁺.

Intermediate 17**(3-Chloro-5-(methylsulfonyl)phenyl)methanol**

To a solution of 3-chloro-5-(methylsulfonyl)benzoic acid (CAS-RN 151104-63-1; 500 mg, 2.13 mmol) in tetrahydrofuran (5 mL) was added slowly borane-tetrahydrofuran complex solution (1
10 M solution in tetrahydrofuran, 5.33 mL, 5.33 mmol) at 0°C, then after 3 h the ice-bath was removed and the reaction mixture was stirred at room temperature overnight. The mixture was then carefully treated with methanol (3 mL) and evaporated. The residue was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane – ethyl acetate
15 gradient afforded the title compound (428 mg, 91%). White solid, MS: 221.3 (M+H)⁺.

The following intermediate was produced according to intermediate 17, replacing 3-chloro-5-(methylsulfonyl)benzoic acid with the appropriate carboxylic acid

No.	Systematic Name	Starting material	MS, m/e
17.1	(2-cyclopropyl-4-(trifluoromethyl)phenyl)methanol	2-cyclopropyl-4-trifluoromethylbenzoic acid (CAS-RN 1236303-04-0)	216.0 (M) ⁺

20 **Intermediate 18****2,2,2-Trifluoro-1-(3-(hydroxymethyl)phenyl)ethanol**

Lithium borohydride solution (2 M in tetrahydrofuran, 1.15 mL, 2.31 mmol) was added dropwise at 0°C to a solution of methyl 3-(2,2,2-trifluoro-1-hydroxyethyl)benzoate (CAS-RN 1188323-28-5; 180 mg, 769 µmol) in tetrahydrofuran (6 mL), then after 15 min the ice bath was removed and the reaction mixture was heated at reflux for 21 h. Then another portion of lithium borohydride solution (2 M in tetrahydrofuran, 0.77 mL, 1.54 mmol) was added and the reaction mixture was heated at reflux for another 5 h. After cooling, the reaction mixture was partitioned between 1 M aq. hydrochloric acid solution and ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane – ethyl acetate gradient) afforded the title compound (86 mg, 53%). White solid, MS: 206.0 (M⁺).

10 **Intermediate 19**

(3-(2,2,2-Trifluoro-1-methoxyethyl)phenyl)methanol

Step 1: Methyl 3-(2,2,2-trifluoro-1-methoxyethyl)benzoate

Sodium hydride dispersion (60% in mineral oil, 93.2 mg, 2.33 mmol) was added at –5°C to a solution of methyl 3-(2,2,2-trifluoro-1-hydroxyethyl)benzoate (303 mg, 1.29 mmol) in tetrahydrofuran (8 mL), then after 30 min iodomethane (643 mg, 4.53 mmol) was added dropwise over a period of 5 min. After 1 h, the ice bath was removed and the reaction mixture was stirred for another 90 min. The reaction mixture was then partitioned between ethyl acetate and water, the organic layer was washed with brine, dried over sodium sulfate, filtered, and evaporated to produce the title compound (purity ca. 90%; 263 mg, 74%). Light yellow liquid, MS: 248.0 (M⁺).

Step 2: (3-(2,2,2-Trifluoro-1-methoxyethyl)phenyl)methanol

The title compound was produced in analogy to intermediate 18 from methyl 3-(2,2,2-trifluoro-1-methoxyethyl)benzoate. Colourless liquid, MS: 220.0 (M⁺).

Intermediate 20

25 **(3aR,6aS)-N-((1H-1,2,3-Triazol-4-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide 2,2,2-trifluoroacetate**

Step 1: (3aR,6aS)-tert-Butyl 5-((1H-1,2,3-triazol-4-yl)methylcarbamoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

-227-

A solution of triphosgene (140 mg, , 471 μmol) in ethyl acetate (12 mL) was added at 0°C to a solution of (3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (CAS-RN 250275-15-1; 200 mg, 942 μmol) in tetrahydrofuran (5 mL). The ice bath was removed, then after 30 min the reaction mixture was heated at reflux for 2 h and was then concentrated in vacuo. The residue was taken up in tetrahydrofuran (20 mL), then after the addition of triethylamine (286 mg, 2.83 mmol) and (1H-1,2,3-triazol-4-yl)methanamine hydrochloride (127 mg, 942 μmol) the reaction mixture was stirred at room temperature for 15 h. After partitioning between water and ethyl acetate, the organic layer was dried over sodium sulfate, filtered, and evaporated. Chromatography (silica gel; dichloromethane – methanol gradient) produced the title compound (110 mg, 34%). White foam, MS: MS: 337.5 (M+H)⁺.

Step 2: (3aR,6aS)-N-((1H-1,2,3-Triazol-4-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide 2,2,2-trifluoroacetate

The title compound was produced in analogy to intermediate 4, step 2 from (3aR,6aS)-tert-butyl 5-((1H-1,2,3-triazol-4-yl)methylcarbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate. Light yellow oil, MS: 237.5 (M+H)⁺.

The following intermediate was produced according to intermediate 20, replacing (1H-1,2,3-triazol-4-yl)methanamine hydrochloride by the appropriate amine.

No.	Systematic Name	Starting material	MS, m/e
20.1	(3aR,6aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methylhexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide 2,2,2-trifluoroacetate	N-methyl-1H-1,2,3-triazole-5-methanamine (CAS-RN 1248059-33-7)	251.5 (M+H) ⁺

20 Intermediate 21

(3aR,6aS)-2-(3-Chlorophenethylsulfonyl)octahydropyrrolo[3,4-c]pyrrole

Step 1: (3aR,6aS)-tert-Butyl 5-(3-chlorophenethylsulfonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

To a solution of (3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (CAS-RN 250275-15-1; 300 mg, 1.34 mmol) in N,N-dimethylformamide (5 mL) was added 2-(3-chlorophenyl)ethanesulfonyl chloride (321 mg, 1.34 mmol) triethylamine (291 mg, 2.87 mmol), and 4-(dimethylamino)pyridine, then after 16 h the reaction mixture was concentrated in vacuo. The residue was partitioned between sat. aq. sodium hydrogencarbonate solution and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated to produce the title compound (520 mg, 93%), which was directly used in the next step.

Step 2: (3aR,6aS)-2-(3-Chlorophenethylsulfonyl)octahydropyrrolo[3,4-c]pyrrole

To a solution of (3aR,6aS)-tert-butyl 5-(3-chlorophenethylsulfonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (520 mg, 1.25 mmol) in methanol (10 mL) was added hydrogen chloride solution (4 M in 1,4-dioxane, 6.3 mL), then after 1 h a few drops of 37% aq. hydrochloric acid solution were added, then after another 2 h the reaction mixture was concentrated to 1/3 of the volume. This was basified to pH 7 with 2 M aq. sodium hydroxide solution and was extracted with ethyl acetate. The organic layer was washed with brine and was dried over magnesium sulfate to produce the title compound (320 mg, 81%). Colourless oil, MS: 315.4 (M+H)⁺.

Intermediate 22

3-Chloro-5-(methylsulfonyl)benzaldehyde

To a clear colourless solution of (3-chloro-5-(methylsulfonyl)phenyl)methanol (intermediate 17; 505 mg, 2.29 mmol) in dichloromethane (10 mL) was added 1,1,1-triacetoxy-1,1-dihydro-1,2-benziodoxol-3(1H)-one solution (15% in dichloromethane; 5.7 mL, 2.75 mmol) at 0°C, then after 1 h the reaction mixture was allowed to reach room temperature over 1 h. After partitioning between dichloromethane and 1 M aq. sodium thiosulfate solution, the organic layer was washed with water and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; dichloromethane) afforded the title compound (376 mg, 75%). White solid, MS: 218 (M⁺).

The following intermediate was produced according to intermediate 20, replacing (1H-1,2,3-triazol-4-yl)methanamine hydrochloride by the appropriate amine.

No.	Systematic Name	Starting material	MS, m/e
22.1	4-formyl-3-isopropylbenzonitrile	4-(hydroxymethyl)-3-isopropylbenzonitrile (intermediate 41)	173 (M) ⁺
22.2	4-formyl-5-isopropyl-2-methylbenzonitrile	4-(hydroxymethyl)-5-isopropyl-2-methylbenzonitrile (intermediate 41.1)	187 (M) ⁺

5 **Intermediate 23**

(E)-3-(3-chloro-5-(methylsulfonyl)phenyl)acrylic acid

To a clear yellow solution of 3-chloro-5-(methylsulfonyl)benzaldehyde (intermediate 22; 370 mg, 1.69 mmol) and malonic acid (352 mg, 3.38 mmol) in pyridine (3 mL) was added piperidine (28.8 mg, 338 μ mol) and the reaction mixture was heated at reflux for 2 h. After cooling, the reaction mixture was treated with 4 M aq. hydrochloric acid solution. The precipitate was collected by filtration, washed with water, and dried to afford the title compound (352 mg, 78%). White solid, MS: 259.5 (M-H)⁻.

The following intermediates were prepared according to intermediate 23, replacing 3-chloro-5-(methylsulfonyl)benzaldehyde by the appropriate aldehyde.

No.	Systematic Name	Aldehyde	MS, m/e
23.01	(E)-3-(3-chloro-5-methoxyphenyl)acrylic acid	3-chloro-5-methoxybenzaldehyde (CAS-RN 164650-68-4)	211.1 (M+H) ⁺

No.	Systematic Name	Aldehyde	MS, m/e
23.02	(E)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)acrylic acid	2-fluoro-4-(trifluoromethoxy)benzaldehyde (CAS-RN 1227628-83-2)	249.1 (M-H) ⁻
23.03	(E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acrylic acid	3-fluoro-4-(trifluoromethoxy)benzaldehyde (CAS-RN 473917-15-6)	249.5 (M-H) ⁻
23.04	(E)-3-(4-fluoro-2-(trifluoromethyl)phenyl)acrylic acid	4-fluoro-2-(trifluoromethyl)benzaldehyde	224.3 (M-H) ⁻
23.05	(E)-3-(6-phenylpyridin-3-yl)acrylic acid	6-phenylnicotinaldehyde	224.3 (M-H) ⁻
23.06	(E)-3-(5-phenylpyridin-2-yl)acrylic acid	5-phenylpicolinaldehyde	224.3 (M-H) ⁻
23.07	(E)-3-(4-(pyridin-4-yl)phenyl)acrylic acid	4-(pyridin-4-yl)benzaldehyde	224.3 (M-H) ⁻
23.08	(E)-3-(4-(pyridin-3-yl)phenyl)acrylic acid	4-(pyridin-3-yl)benzaldehyde	224.3 (M-H) ⁻
23.09	(E)-3-(4-(pyridin-2-yl)phenyl)acrylic acid	4-(pyridin-2-yl)benzaldehyde	224.3 (M-H) ⁻
23.10	(E)-3-(2-cyclopropylphenyl)acrylic acid	2-cyclopropylbenzaldehyde	187.4 (M-H) ⁻
23.11	(E)-3-(2-fluoro-4-(2,2,2-trifluoroethoxy)phenyl)acrylic acid	2-fluoro-4-(2,2,2-trifluoroethoxy)benzaldehyde	263.0 (M-H) ⁻

(E)-3-(3-Methoxy-5-(trifluoromethoxy)phenyl)acrylic acidStep 1 (E)-tert-Butyl 3-(3-methoxy-5-(trifluoromethoxy)phenyl)acrylate

To a colourless solution of 1-bromo-3-methoxy-5-(trifluoromethoxy)benzene (CAS-RN 1330750-28-1; 1.00 g, 3.62 mmol) in N,N-dimethylformamide (10 mL) was added triethylamine (1.1 g, 10.8 mmol), tert-butyl acrylate (567 mg, 4.34 mmol), palladium(II) acetate (16.2 mg, 72.3 μ mol) and tri-o-tolylphosphine (88 mg, 289 μ mol). The light yellow reaction mixture was evacuated and backfilled with argon, three times. The reaction mixture was heated at 120°C, then after 16 h partitioned between ethyl acetate and sat. aq. sodium hydrogencarbonate solution. The organic layer was washed with water, sat. aq. ammonium chloride solution, and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane – ethyl acetate gradient) afforded the title compound (984 mg, 85%). Colourless oil, MS: 318 (M^+).

Step 2: (E)-3-(3-Methoxy-5-(trifluoromethoxy)phenyl)acrylic acid

To a solution of (E)-tert-butyl 3-(3-methoxy-5-(trifluoromethoxy)phenyl)acrylate (966 mg, 3.03 mmol) in dichloromethane (9 mL) was added trifluoroacetic acid (3.5 mL), then after 2 h the reaction mixture was concentrated in vacuo. The residue was triturated in heptane to afford the title compound (752 mg, 95%). White solid, MS: 261.2 ($M-H$)⁻.

The following intermediate was prepared according to intermediate 24, replacing 1-bromo-3-methoxy-5-(trifluoromethoxy)benzene by the appropriate starting material.

No.	Systematic Name	Starting material	MS, m/e
24.1	(E)-3-(3-chloro-5-methoxyphenyl)acrylic acid	3-chloro-5-iodobenzonitrile (CAS-RN 289039-30-1)	206.1 ($M-H$) ⁻

Intermediate 25

(6,7-Dihydro-1H-[1,2,3]triazolo[4,5-c]pyridin-5(4H)-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride

-232-

Step 1: (3aS,6aS)-tert-Butyl 5-(4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

To a colourless solution of 4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine (CAS-RN 706757-05-3; 102 mg, 819 μ mol) and N,N-diisopropylethylamine (222 mg, 1.72 mmol) in N,N-dimethylformamide (4 mL) was added a solution of (3aR,6aR)-tert-butyl 5-(chlorocarbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (intermediate 9.2; 225 mg, 819 μ mol) in dichloromethane (8 mL), then after 70 h the reaction mixture was partitioned between dichloromethane and sat. aq. ammonium chloride solution. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 90:10:0.25) produced the title compound (225 mg, 76%). White foam, MS: 363.6 (M+H)⁺.

Step 2: (6,7-Dihydro-1H-[1,2,3]triazolo[4,5-c]pyridin-5(4H)-yl)((3aR,6aR)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone hydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from (3aS,6aS)-tert-butyl 5-(4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate. White solid, MS: 263.5 (M+H)⁺.

The following intermediate was produced according to intermediate 25, replacing 4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine by the appropriate amine:

No.	Systematic Name	Amine	MS, m/e
25.1	(3aR,7aR)-5-((3aR,6aR)-octahydropyrrolo[3,4-c]pyrrole-2-carbonyl)hexahydrooxazolo[5,4-c]pyridin-2(1H)-one hydrochloride	(3aR,7aR)-hexahydrooxazolo[5,4-c]pyridin-2(1H)-one hydrochloride (intermediate 27)	281.1 (M+H) ⁺

Intermediate 26

(E)-1-((3aR,8aS)-Octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-(trifluoromethoxy)-phenyl)prop-2-en-1-one

-233-

Step 1: from (3aR,8aS)-tert-butyl 6-((E)-3-(3-(trifluoromethoxy)phenyl)acryloyl)-octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate

To a solution of (3aR,8aS)-tert-butyl octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate (CAS-RN 1251013-07-6; 353 mg, 1.47 mmol) and (E)-3-(3-(trifluoromethoxy)phenyl)acrylic acid (341 mg, 1.47 mmol) in N,N-dimethylformamide (10 mL) were added N-ethyldiisopropylamine (570 mg, 4.41 mmol) dropwise over a period of 2 minutes at room temperature under an argon atmosphere. The mixture was cooled down to 0°C and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluoro-phosphate (563 mg, 1.47 mmol) was added, then after 1 h the ice bath was removed. The reaction mixture was stirred for 16 h at room temperature, then partitioned between ethyl acetate and sat. aq. sodium hydrogencarbonate solution. The organic layer was washed with brine, dried over sodium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane – ethyl acetate gradient) produced the title compound (768 mg, 87%). White foam, MS: 399.5 (M+H–isobutene)⁺.

Step 2: (E)-1-((3aR,8aS)-Octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one

Trifluoroacetic acid (1.1 mL, 15 mmol) was added over 5 min to a solution of (3aR,8aS)-tert-butyl 6-((E)-3-(3-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate (768 mg, 1.49 mmol) in dichloromethane (12 mL) at room temperature, then after 5 h the reaction mixture was poured onto ice water, basified to pH 10 with 2 M aq. sodium hydroxide solution, and extracted with chloroform. The organic layer was washed with brine, dried over sodium sulfate, filtered, and evaporated. Chromatography (silica gel; dichloromethane/methanol 9:1) produced the title compound (526 mg, 95%). Yellow gum, MS: 355.5 (M+H)⁺.

The following intermediate was prepared according to intermediate 26, replacing (3aR,8aS)-tert-butyl octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate by the appropriate amine and (E)-3-(3-(trifluoromethoxy)phenyl)acrylic acid by the appropriate carboxylic acid.

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
26.01	(E)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one	(3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (CAS-RN 250275-15-1)	(E)-3-(4-(trifluoromethoxy)-phenyl)acrylic acid	327.5 (M+H) ⁺
26.02	3-(3-chlorophenyl)-2,2-dimethyl-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)propan-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	3-(3-chlorophenyl)-2,2-dimethylpropanoic acid (CAS-RN 1225505-29-2)	335.6 (M+H) ⁺
26.03	(E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(3-fluoro-4-(trifluoromethoxy)-phenyl)acrylic acid (intermediate 23.3)	373.6 (M+H) ⁺
26.04	(E)-3-(4-fluoro-2-(trifluoromethyl)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(4-fluoro-2-(trifluoromethyl)-phenyl)acrylic acid (intermediate 23.4)	357.6 (M+H) ⁺

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
26.05	(E)-3-(2-methyl-4-(trifluoromethoxy)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(2-methyl-4-(trifluoromethoxy)phenyl)-acrylic acid (CAS-RN 1262012-31-6)	369.6 (M+H) ⁺
26.06	(E)-3-(3-fluoro-4-methoxyphenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(3-fluoro-4-methoxyphenyl)-acrylic acid (CAS-RN 147906-08-9)	319.6 (M+H) ⁺
26.07	(E)-3-(2-isopropyl-phenyl)-1-(3aS,8aR)-octahydro-pyrrolo[3,4-d]azepin-6-yl-prop-2-en-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(2-isopropyl-phenyl)-acrylic acid (CAS-RN 1379383-70-6)	313.2 (M+H) ⁺
26.08	(E)-3-(2-cyclopropylphenyl)-1-(trans-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(2-cyclopropyl-phenyl)acrylic acid (intermediate 23.10)	311.6 (M+H) ⁺

No.	Systematic Name	Amine	Carboxylic acid	MS, m/e
26.09	(E)-3-(4-methoxy-2-(trifluoromethyl)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(4-methoxy-2-(trifluoromethyl)phenyl)-acrylic acid (CAS-RN 773131-66-1)	369.6 (M+H) ⁺
26.10	(E)-3-(3-chloro-5-(trifluoromethyl)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)prop-2-en-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	(E)-3-(3-chloro-5-(trifluoromethyl)phenyl)acrylic acid (CAS-RN 886761-69-9)	373.6 (M+H) ⁺
26.11	3-(3-fluoro-4-(2,2,2-trifluoroethoxy)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)propan-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6)	3-(3-fluoro-4-(2,2,2-trifluoroethoxy)phenyl)-propanoic acid	389.2 (M+H) ⁺
26.12	3-(2-fluoro-4-(2,2,2-trifluoroethoxy)phenyl)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)propan-1-one	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester (CAS-RN 1251013-07-6)	3-(2-fluoro-4-(2,2,2-trifluoroethoxy)-phenyl)propanoic acid (intermediate 35.3)	389.2 (M+H) ⁺

Intermediate 27**(3aR,7aR)-Hexahydrooxazolo[5,4-c]pyridin-2(1H)-one hydrochloride**

Step 1: (3aR,7aR)-tert-butyl 2-oxohexahydrooxazolo[5,4-c]pyridine-5(6H)-carboxylate

To a solution of (3R,4R)-tert-butyl 4-amino-3-hydroxypiperidine-1-carboxylate (CAS-RN 1007596-95-3; 500 mg, 2.31 mmol) in N,N-dimethylformamide (5.00 mL) was added imidazole (157 mg, 2.31 mmol) and 1,1'-carbonyldiimidazole (375 mg, 2.31 mmol) at room temperature, then after 18 h the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over magnesium sulfate, filtered and evaporated. The residue was chromatographed (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 95:5:0.25) to produce the title compound (401 mg, 72%) as a white solid.

10 Step 2: (3aR,7aR)-Hexahydrooxazolo[5,4-c]pyridin-2(1H)-one hydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from (3aR,7aR)-tert-butyl 2-oxohexahydrooxazolo[5,4-c]pyridine-5(6H)-carboxylate. White solid, MS: 143.2 (M+H)⁺.

Intermediate 28**cis-Hexahydrooxazolo[5,4-c]pyridin-2(1H)-one hydrochloride**15 Step 1: cis-tert-butyl 2-oxohexahydrooxazolo[5,4-c]pyridine-5(6H)-carboxylate

To a light yellow solution of 1-(tert-butoxycarbonyl)-3-hydroxypiperidine-4-carboxylic acid (CAS-RN 1260876-51-4; 196 mg, 799 µmol) in toluene (2 mL) was added triethylamine (97 mg, 0.96 mmol) and diphenylphosphoryl azide (269 mg, 959 µmol). The reaction mixture was heated at reflux for 18 h and was then partitioned between ethyl acetate and sat. aq. sodium hydrogen carbonate solution. The organic layer was washed with water and brine, dried over magnesium sulfate, filtered and evaporated. Chromatography (silica gel; ethyl acetate–methanol gradient) produced the title compound (66 mg, 34%). White solid, MS: 241.4 (M–H)[–].

Step 2: cis-Hexahydrooxazolo[5,4-c]pyridin-2(1H)-one hydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from cis-tert-butyl 2-oxohexahydrooxazolo[5,4-c]pyridine-5(6H)-carboxylate. White solid, MS: 142.1 (M⁺).

Intermediate 29**(3aSR,6SR,7aSR)-2-Oxoctahydrobenzo[d]oxazole-6-carboxylic acid**

Step 1: (1SR,3SR,4RS)-Methyl 4-bromo-3-(phenoxy carbonyloxy)cyclohexanecarboxylate

To a solution of (1SR,3SR,4RS)-methyl 4-bromo-3-hydroxycyclohexanecarboxylate (CAS 38361-11-4; 500 mg, 2.11 mmol) and pyridine (175 mg, 2.21 mmol) in dichloromethane (8 mL) was added a solution of phenyl carbonochloridate (347 mg, 2.21 mmol) in dichloromethane (1 mL) at -5°C , then after 1 h the reaction mixture was partitioned between dichloromethane and water. The organic layer was washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane – ethyl acetate gradient) produced the title compound (545 mg, 72%). Colourless oil, MS: 357.4 $(\text{M}+\text{H})^{+}$.

10 Step 2: (1SR,3SR,4SR)-Methyl 4-azido-3-(phenoxy carbonyloxy)cyclohexanecarboxylate

To a solution of (1SR,3SR,4RS)-methyl 4-bromo-3-(phenoxy carbonyloxy)cyclohexanecarboxylate (533 mg, 1.49 mmol) and 15-crown-5 (23.0 mg, 104 μmol) in N,N-dimethylformamide (4 mL) was added sodium azide (437 mg, 6.71 mmol). The reaction mixture was heated at 75°C for 72 h and was then partitioned between water and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; dichloromethane) produced the title compound (60 mg, 13%). Colourless oil, MS: 337.2 $(\text{M}+\text{NH}_4)^{+}$.

Step 3: (3aSR,6SR,7aSR)-Methyl 2-oxooctahydrobenzo[d]oxazole-6-carboxylate

To a colourless solution of (1R,3R,4R)-methyl 4-azido-3-

(phenoxy carbonyloxy)cyclohexanecarboxylate (56 mg, 175 μmol) in tetrahydrofuran (2 mL) and water (50 μL) was added triphenylphosphine (138 mg, 526 μmol). The solution was heated to 50°C for 3 h and was then partitioned between ethyl acetate and brine. The organic layer was dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 95:5:0.25) produced the title compound (49 mg), which contained triphenylphosphine oxide as an inseparable impurity.

Step 4: (3aSR,6SR,7aSR)-2-Oxooctahydrobenzo[d]oxazole-6-carboxylic acid

The title compound was produced in analogy to intermediate 11, step 2 from (3aSR,6SR,7aSR)-methyl 2-oxooctahydrobenzo[d]oxazole-6-carboxylate. White solid, MS: 184.3 $(\text{M}-\text{H})^{-}$.

Intermediate 30A and 30B**(+)-(R)-4,5,6,7-Tetrahydro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid and (-)-(S)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid**

Racemic 4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid (CAS-RN 33062-47-4;

- 5 1.10 g, 6.58 mmol) was separated by preparative HPLC using a Chiralpak AD column as the stationary phase and heptane/ethanol 3:2 as the mobile phase. This produced the faster eluting (+)-(R)-enantiomer (452 mg, 41%), followed by the slower eluting (-)-(S)-enantiomer (381 mg, 35%).

Intermediate 31

- 10 **2-(4-Chloro-2-methylphenoxy)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrole-2(1H)-yl)ethanone hydrochloride**

Step 1: (3aR,6aS)-tert-Butyl 5-(2-bromoacetyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

- 15 To a solution of (3aR,6aS)-tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (CAS-RN 250275-15-1; 400 mg, 1.88 mmol) and triethylamine (191 mg, 1.88 mmol) was added 2-bromoacetyl chloride (297 mg, 1.88 mmol) at -40°C, then after 2 h the reaction mixture was washed with water at 0°C. The organic layer was dried over magnesium sulfate, filtered, and evaporated to produce the title compound (628 mg, 100%), which was directly used in the next step.

- 20 Step 2: (3aR,6aS)-tert-Butyl 5-(2-(4-chloro-2-methylphenoxy)acetyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate

- To a solution of (3aR,6aS)-tert-butyl 5-(2-bromoacetyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate (100.6 mg, 302 µmol) and 4-chloro-2-methylphenol (64.6 mg, 453 µmol,) in N,N-dimethylformamide (5 mL) was added caesium carbonate (197 mg, 604 µmol) at room
25 temperature, then after 16 h the reaction mixture was partitioned between water and ethyl acetate. The organic layer was washed with brine, dried over sodium sulfate, filtered, and concentrated in vacuo. Chromatography (silica gel; heptane – ethyl acetate gradient) produced the title compound (57 mg, 48%). Colourless oil, MS: 339.5 (M+H-isobutene)⁺.

-240-

Step 3: 2-(4-Chloro-2-methylphenoxy)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)ethanone hydrochloride

The title compound was produced in analogy to intermediate 1, step 2 from (3aR,6aS)-tert-butyl 5-(2-(4-chloro-2-methylphenoxy)acetyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate.

- 5 Light brown solid, MS: 295.5 (M+H)⁺.

The following intermediate was produced in analogy to intermediate 31, replacing 4-chloro-2-methylphenol in step 2 by the appropriate phenol.

No.	Systematic Name	Phenol	MS, m/e
31.1	2-(4-chloro-3-methylphenoxy)-1-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)ethanone hydrochloride	4-chloro-3-methylphenol	295.5 (M+H) ⁺

10 Intermediate 32

(3aR,8aS)-2-Fluoro-4-(trifluoromethoxy)benzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate

Step 1: (3aR,8aS)-2-tert-Butyl 6-(2-fluoro-4-(trifluoromethoxy)benzyl) hexahydropyrrolo[3,4-d]azepine-2,6(1H,7H)-dicarboxylate

- 15 The title compound was produced in analogy to intermediate 1, step 1 from cis-octahydropyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride (CAS-RN 1251013-07-6) and (2-fluoro-4-(trifluoromethoxy)phenyl)methanol (CAS-RN 1240257-07-1).

Step 2: (3aR,8aS)-2-Fluoro-4-(trifluoromethoxy)benzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate

- 20 The title compound was produced in analogy to intermediate 4, step 2 from (3aR,8aS)-2-tert-butyl 6-(2-fluoro-4-(trifluoromethoxy)benzyl) hexahydropyrrolo[3,4-d]azepine-2,6(1H,7H)-dicarboxylate. Light brown gum, MS: 377.6 (M+H)⁺.

The following intermediates were produced according to intermediate 32, replacing cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester hydrochloride and (2-fluoro-4-(trifluoromethoxy)phenyl)methanol by the appropriate amine and alcohol, respectively.

No.	Systematic Name	Amine	Alcohol	MS, m/e
32.1	trans-(2-methoxy-4-(trifluoromethoxy)benzyl) hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	(2-methoxy-4-(trifluoromethoxy)phenyl)-methanol (CAS-RN 886500-30-7)	375.6 (M+H) ⁺
32.2	trans-(2-cyclopropyl-4-(trifluoromethyl)benzyl) hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	(2-cyclopropyl-4-(trifluoromethyl)-phenyl)methanol (intermediate 17.1)	369.3 (M+H) ⁺
32.3	trans-(4-fluoro-2-(trifluoromethyl)benzyl) hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate	trans-tert-butyl hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (CAS-RN 1251014-37-5)	(4-fluoro-2-(trifluoromethyl)-phenyl)methanol	347.6 (M+H) ⁺
32.4	(3aR,8aS)-3-fluoro-4-(2,2,2-trifluoroethoxy)benzyl octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate	cis-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid tert-butyl ester (CAS-RN 1251013-07-6)	[3-fluoro-4-(2,2,2-trifluoroethoxy)-phenyl]methanol (CAS-RN 1039931-47-9)	390.2 (M+)

Intermediate 33

2-(2-(Tetrahydrofuran-2-yl)phenoxy)acetic acidStep 1: Ethyl 2-(2-(tetrahydrofuran-2-yl)phenoxy)acetate

To a solution of 2-(tetrahydrofuran-2-yl)phenol (CAS-RN 40324-49-0; 510 mg, 3.11 mmol) in acetone (4 mL) were added potassium carbonate (859 mg, 6.21 mmol) and ethyl 2-bromoacetate (545 mg, 3.26 mmol) at room temperature, then after 3 h the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated to produce the title compound (754 mg, 97%). Light yellow oil, MS:251.5 (M+H)⁺.

Step 2: 2-(2-(Tetrahydrofuran-2-yl)phenoxy)acetic acid

To a solution of ethyl 2-(2-(tetrahydrofuran-2-yl)phenoxy)acetate (754mg, 3.01 mmol) in methanol (2.5 mL) and tetrahydrofuran (12 mL) was added 1 M aq. lithium hydroxide solution (5.12 mL, 5.12 mmol), then after 90 min the reaction mixture was acidified with 2 M aq. hydrochloric acid solution and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated to produce the title compound (650 mg, 97%). White solid, MS: 221.2 (M-H)⁻.

The following intermediates were produced according to intermediate 33, replacing 2-(tetrahydrofuran-2-yl)phenol by the appropriate phenol.

No.	Systematic Name	Phenol	MS, m/e
33.1	2-(2-bromo-4-(trifluoromethoxy)-phenoxy)acetic acid	2-bromo-4-(trifluoromethoxy)-phenol (CAS-RN 200956-13-4)	313.3 (M-H) ⁻
33.2	2-(2-(1H-pyrrol-1-yl)phenoxy)acetic acid	2-(1H-pyrrol-1-yl)phenol (CAS-RN 32277-91-1)	216.5 (M-H) ⁻
33.3	2-(2-chloro-4-(trifluoromethoxy)phenoxy)acetic acid	2-chloro-4-(trifluoromethoxy)phenol (CAS-RN 70783-75-4)	269.5 (M-H) ⁻

Intermediate 34**2-(4-Cyano-2-isopropyl-5-methylphenoxy)acetic acid**Step 1: tert-Butyl 2-(4-cyano-2-isopropyl-5-methylphenoxy)acetate

To a solution of 4-hydroxy-5-isopropyl-2-methylbenzonitrile (CAS-RN 858026-56-9; 156 mg, 890 μmol) in acetone (4 mL) were added potassium carbonate (246 mg, 1.78 mmol) and tert-butyl 2-bromoacetate (188 mg, 935 μmol) at room temperature, then after 3 h the reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated to produce the title compound (253 mg 98%). White solid, MS: 290.5 (M+H)⁺.

10 Step 2: 2-(4-Cyano-2-isopropyl-5-methylphenoxy)acetic acid

To a colourless solution of tert-butyl 2-(4-cyano-2-isopropyl-5-methylphenoxy)acetate (248 mg, 857 μmol) in dichloromethane (3 mL) was added trifluoroacetic acid (1.95 g, 17.1 mmol), then after 3 h the reaction mixture was concentrated and the residue triturated in heptane. The crude product was purified by chromatography (silica gel; dichloromethane–methanol gradient) to afford the title compound (179 mg, 90%). White solid, MS: 232.5 (M–H)[–].

The following intermediates were produced according to intermediate 34, replacing 4-hydroxy-5-isopropyl-2-methylbenzonitrile by the appropriate phenol.

No.	Systematic Name	Phenol	MS, m/e
34.1	2-(4-cyano-2-isopropylphenoxy)acetic acid	3-isopropyl-4-hydroxy-benzonitrile (CAS-RN 46057-54-9)	218.3 (M–H) [–]
34.2	2-(2-cyano-4-(trifluoromethoxy)phenoxy)acetic acid	2-hydroxy-5-(trifluoromethoxy)-benzonitrile	260.5 (M–H) [–]
34.3	2-(2-(pyridin-3-yl)phenoxy)acetic acid	2-(3-pyridinyl)-phenol (CAS-RN 54168-07-9)	230.2 (M+H) ⁺

Intermediate 35**3-(3-Fluoro-4-(trifluoromethoxy)phenyl)propanoic acid**

A solution of (E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acrylic acid (intermediate 23.3; 500 mg, 2.00 mmol) in methanol (7 mL) was stirred under a hydrogen atmosphere (1 bar) in the presence of palladium (10% on activated charcoal; 50 mg), then after 20 h insoluble material was removed by filtration through diatomaceous earth. The filtrate was concentrated to produce the title compound (485 mg, 96%). White solid, MS: 251.2 (M-H)⁻.

- 10 The following intermediates were produced according to intermediate 35, replacing (E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acrylic acid by the appropriate starting material.

No.	Systematic Name	Starting material	MS, m/e
35.1	3-(6-phenylpyridin-3-yl)propanoic acid	(E)-3-(6-phenylpyridin-3-yl)acrylic acid (intermediate 23.05)	226.3 (M-H) ⁻
35.2	3-(3-fluoro-4-(2,2,2-trifluoroethoxy)phenyl)propanoic acid	(E)-3-(3-fluoro-4-(2,2,2-trifluoroethoxy)phenyl)acrylic acid (CAS-RN 1087780-94-6)	265.1 (M-H) ⁻
35.3	3-(2-fluoro-4-(2,2,2-trifluoroethoxy)phenyl)propanoic acid	(E)-3-(2-fluoro-4-(2,2,2-trifluoroethoxy)phenyl)acrylic acid (intermediate 23.11)	265.1 (M-H) ⁻
35.4	3-(2-fluoro-4-(trifluoromethoxy)phenyl)propanoic acid	(E)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)acrylic acid (intermediate 23.02)	251.2 (M-H) ⁻

Intermediate 36

2-(5-Chloro-2-(trifluoromethyl)phenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone

Step 1: (3aR,8aS)-tert-butyl 6-(2-(5-chloro-2-(trifluoromethyl)phenoxy)acetyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate.

- 5 Caesium carbonate (132 mg, 404 μmol) was added to a solution of 5-chloro-2-(trifluoromethyl)phenol (47.7 mg, 242 μmol) and (3aR,8aS)-tert-butyl 6-(2-bromoacetyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate (intermediate 37; 73 mg, 202 μmol) in N,N-dimethylformamide (5 mL) at room temperature, then after 18 h the reaction mixture was partitioned between ice water and ethyl acetate. The organic layer was washed with
- 10 brine, dried over sodium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane–ethyl acetate gradient) produced the title compound (82 mg, 83%). White foam, MS: 421.5 (M+H–isobutene)⁺.

Step 2: 2-(5-Chloro-2-(trifluoromethyl)phenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone

- 15 The title compound was produced in analogy to intermediate 26, step 2 from (3aR,8aS)-tert-butyl 6-(2-(5-chloro-2-(trifluoromethyl)phenoxy)acetyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate. Light yellow foam, MS: 377.5 (M+H)⁺.

- The following intermediates were produced according to intermediate 36, replacing 5-chloro-2-
- 20 (trifluoromethyl)phenol by the appropriate phenol.

No.	Systematic Name	Phenol	MS, m/e
36.01	2-(6-isopropyl-3,3-dimethyl-2,3-dihydro-1H-inden-5-yloxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone	6-isopropyl-3,3-dimethyl-2,3-dihydro-1H-inden-5-ol	385.7 (M+H) ⁺

No.	Systematic Name	Phenol	MS, m/e
36.02	2-(2-isopropyl-5-methylphenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone	2-isopropyl-5-methylphenol	331.6 (M+H) ⁺
36.03	2-(2-chloro-4-fluorophenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone	2-chloro-4-fluorophenol	327.1 (M+H) ⁺
36.04	2-(4-methyl-2-(1-methylpyrrolidin-3-yl)phenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone	4-methyl-2-(1-methylpyrrolidin-3-yl)phenol	372.3 (M+H) ⁺
36.05	4-(2-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-oxoethoxy)-3-(trifluoromethyl)benzonitrile	4-hydroxy-3-(trifluoromethyl)benzonitrile	368.1 (M+H) ⁺
36.06	2-(4-chloro-2-isopropyl-5-methylphenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone	4-chloro-2-isopropyl-5-methylphenol	365.2 (M+H) ⁺
36.07	1-(3aS,8aR)-octahydro-pyrrolo[3,4-d]azepin-6-yl-2-(2-trifluoromethoxy-phenoxy)-ethanone	2-(trifluoromethoxy)phenol	359.6 (M+H) ⁺
36.08	2-(2-tert-butyl-4-methoxyphenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone	2-tert-butyl-4-methoxyphenol	361.6 (M+H) ⁺
36.09	3-isopropyl-4-(2-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-oxoethoxy)benzonitrile	4-hydroxy-3-isopropylbenzonitrile (CAS-RN CAS-RN 46057-54-9)	342.6 (M+H) ⁺

No.	Systematic Name	Phenol	MS, m/e
36.10	2-(2-chloro-4-(trifluoromethyl)phenoxy)-1-((3aR,8aS)-octahydropyrrolo[3,4-d]azepin-6(7H)-yl)ethanone	2-chloro-4-(trifluoromethyl)phenol	n.a.

Intermediate 37**(3aR,8aS)-tert-Butyl 6-(2-bromoacetyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate**

To a suspension of (3aR,8aS)-tert-butyl octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate hydrochloride (CAS-RN 1251013-07-6; 600 mg, 2.17 mmol) and triethylamine (439 mg, 4.34 mmol) in dichloromethane (40 mL) was added 2-bromoacetyl chloride (341 mg, 2.17 mmol) dropwise at -40°C, then the reaction mixture was allowed to reach room temperature over 2 h and partitioned between water and dichloromethane. The organic layer was dried over sodium sulfate, filtered, and evaporated to produce the title compound (600 mg, 69%; brown oil), which was directly used in the next step.

Intermediate 38**[2-Fluoro-4-(2,2,2-trifluoroethoxy)phenyl]methanol**

Sodium borohydride was added portionwise at 0°C to a solution of 2-fluoro-4-(2,2,2-trifluoroethoxy)benzaldehyde (intermediate 39; 500 mg, 2.25 mmol) in methanol (3 mL), then the reaction mixture was allowed to reach room temperature over 3 h. After addition of water (5 mL) and evaporation of methanol, the reaction mixture was partitioned between ethyl acetate and water. The organic layer was dried over sodium sulfate, filtered, and evaporated to produce the title compound (450 mg, 87%). Colourless oil, MS: 224 (M⁺).

Intermediate 39

2-Fluoro-4-(2,2,2-trifluoroethoxy)benzaldehyde

To a solution of 2-fluoro-4-hydroxybenzaldehyde (1.00 g, 7.14 mmol) in N,N-dimethylformamide (6 mL) were added potassium carbonate (1.48 g, 10.7 mmol) and 2,2,2-trifluoroethyl trifluoromethanesulfonate (1.99 g, 8.56 mmol) at room temperature. The reaction mixture was heated at 50°C for 2 h, then partitioned between ethyl acetate and water. The organic layer was dried over sodium sulfate, filtered, and evaporated to produce the title compound (1.30 g, 80%). White solid, MS: 222 (M^+).

Intermediate 40**2-(2-tert-Butyl-4-cyanophenoxy)acetic acid**

A mixture of 2-(4-bromo-2-tert-butylphenoxy)acetic acid (CAS-RN 425372-86-7; 453 mg, 1.58 mmol), 1,1'-bis(diphenylphosphino)ferrocene (26.2 mg, 47.3 μ mol), tris(dibenzylideneacetone)dipalladium(0) (14.4 mg, 15.8 μ mol), zinc cyanide (102 mg, 868 μ mol), zinc powder (4.13 mg, 63.1 μ mol) and zinc acetate (11.6 mg, 63.1 μ mol), N,N-dimethylformamide (4.5 mL) and water (45 μ L) was heated at 180°C for 15 min under microwave irradiation, then evaporated under vacuum. The residue was taken up in ethyl acetate, then insoluble material was removed by filtration through diatomaceous earth. The filtrate was evaporated and the residue purified by chromatography (silica gel; ethyl acetate-methanol gradient, then dichloromethane-methanol gradient) to produce the title compound (64 mg, 22%). Dark brown solid. MS: 232.5 ($M-H$)⁻.

Intermediate 41**4-(Hydroxymethyl)-3-isopropylbenzonitrile****Step 1: 4-Cyano-2-isopropylphenyl trifluoromethanesulfonate**

To a solution of pyridine (895 mg, 11.3 mmol) in dichloromethane (70 mL) was added trifluoromethanesulfonic anhydride (2.93 g, 10.4 mmol) at 0°C, then after 10 min a solution of 4-hydroxy-3-isopropylbenzonitrile (CAS-RN 46057-54-9; 1.52 g, 9.43 mmol) in dichloromethane (40 mL) was added dropwise to the white suspension that had formed. The ice bath was

-249-

removed, then after 75 min the reaction mixture was partitioned between dichloromethane and water. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane–dichloromethane gradient) produced the title compound (2.63 g, 95%). Yellow liquid, MS: 292.1 (M–H)[–].

5 Step 2: Methyl 4-cyano-2-isopropylbenzoate

A solution of 4-cyano-2-isopropylphenyl trifluoromethanesulfonate (2.62 g, 8.93 mmol), triethylamine (2.26 g, 22.3 mmol), and bis(diphenylphosphino)ferrocene-palladium(II) dichloride dichloromethane complex (365 mg, 447 μ mol) was stirred for 20 h under a carbon monoxide atmosphere (50 bar) at 110°C. After cooling the reaction mixture was evaporated and
10 the residue was purified by chromatography (silica gel; dichloromethane/heptane 1:1) to produce the title compound (1.40 g, 77%). Light yellow oil, MS: 218.5 (M+H)⁺.

Step 3: 4-(Hydroxymethyl)-3-isopropylbenzonitrile

Lithium borohydride solution (2 M in tetrahydrofuran, 9.06 mL, 18.1 mmol) was added at room temperature to a solution of methyl 4-cyano-2-isopropylbenzoate (1.227 g, 6.04 mmol) in
15 tetrahydrofuran (15 mL). The reaction mixture was heated at reflux for 1 h, then partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; gradient dichloromethane to dichloromethane/methanol/25% aq. ammonia solution 95:5:0.25) afforded the title compound (802 mg, 76%). Light yellow oil, MS: 176.2 (M+H)⁺.

20

The following intermediates were produced according to intermediate 41, replacing 4-hydroxy-3-isopropylbenzonitrile by the appropriate phenol.

No.	Systematic Name	Phenol	MS, m/e
41.1	4-(hydroxymethyl)-5-isopropyl-2-methylbenzonitrile	4-hydroxy-5-isopropyl-2-methylbenzonitrile (CAS-RN 858026-56-9)	190.3 (M+H) ⁺

No.	Systematic Name	Phenol	MS, m/e
41.2	3-ethoxy-4-(hydroxymethyl)benzonitrile	3-ethoxy-4-hydroxy-benzonitrile (CAS-RN 60758-79-4)	177 (M) ⁺

Intermediate 42

3-(4-Cyano-2-isopropylphenyl)propanoic acid

Triethylamine (1.22 g, 12.1 mmol) was added dropwise to formic acid (1.36 g, 29.6 mmol) at 0°C. This was added to 4-formyl-3-isopropylbenzonitrile (190 mg, 1.1 mmol) and 2,2-dimethyl-1,3-dioxane-4,6-dione (158 mg, 1.1 mmol). The solution was stirred at room temperature for 3 h, then poured upon ice water, acidified with 4 M aq. hydrochloric acid solution, and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated to afford crude 4-[(2,2-dimethyl-4,6-dioxo-1,3-dioxan-5-yl)methyl]-3-isopropylbenzonitrile (334 mg) as a light yellow foam, MS: 302.4 (M+H)⁺. This was dissolved in acetonitrile/water 100:1 (2 mL) and heated at 100°C for 30 min under microwave irradiation, then concentrated in vacuo. The residue was partitioned between 2 M hydrochloric acid solution and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated to afford the title compound (240 mg), which was used directly in the next step. Yellow oil, MS: 216.3 (M+H)⁺.

The following intermediate was produced according to intermediate 42, replacing 4-formyl-3-isopropylbenzonitrile by the appropriate aldehyde.

No.	Systematic Name	Phenol	MS, m/e

No.	Systematic Name	Phenol	MS, m/e
42.1	3-(4-cyano-2-isopropyl-5-methylphenyl)propanoic acid	4-formyl-5-isopropyl-2-methyl-benzonitrile (intermediate 22.2)	230.2 (M-H) ⁻

Intermediate 43

6-Fluoro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid

Step 1: Methyl 6-fluoro-1H-benzo[d][1,2,3]triazole-5-carboxylate

- 5 A solution of 5-bromo-6-fluoro-1H-benzo[d][1,2,3]triazole (CAS-RN 1242336-69-1; 492 mg, 2.28 mmol), triethylamine (576 mg, 5.69 mmol), and 1,1'-bis(diphenylphosphino)ferrocene-palladium(II) dichloride dichloromethane complex (74.4 mg, 91.1 μ mol), in methanol (6 mL) was stirred at 110°C under a carbon monoxide atmosphere (70 bar) for 18 h. After cooling insoluble material was separated by filtration through diatomaceous earth. The filtrate was
- 10 evaporated and purified by chromatography (silica gel; dichloromethane–methanol gradient) to produce the title compound (281 mg, 63%). Light red solid, MS: 194.2 (M-H)⁻.

Step 2: 6-Fluoro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid

- To a solution of methyl 6-fluoro-1H-benzo[d][1,2,3]triazole-5-carboxylate (276 mg, 1.41 mmol) in tetrahydrofuran (3 mL) and methanol (1.5 mL) was added 1 M aq. lithium hydroxide solution
- 15 1 M in water (4.24 mL, 4.24 mmol). After 5 h most of the organic solvents were removed by concentration under reduced pressure, then the aqueous solution was acidified with 1 M aq. hydrochloric acid solution. The precipitate was collected by filtration and dried to afford the title compound (266 mg, 100 %). Off-white solid, MS: 180.2 (M-H)⁻.
- 20 The following intermediates were produced according to intermediate 43, replacing 5-bromo-6-fluoro-1H-benzo[d][1,2,3]triazole by the appropriate starting material.

No.	Systematic Name	Starting material	MS, m/e
43.1	7-fluoro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid	5-bromo-7-fluoro-1H-benzo[d][1,2,3]triazole (intermediate 44)	180.2 (M-H) ⁻
43.2	4-fluoro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid	5-bromo-4-fluoro-1H-benzo[d][1,2,3]triazole (intermediate 44.1)	180.2 (M-H) ⁻
43.3	4-methyl-1H-benzo[d][1,2,3]triazole-5-carboxylic acid	5-bromo-4-methyl-1H-benzo[d][1,2,3]triazole (CAS-RN 1372795-26-0)	176.2 (M-H) ⁻
43.4	6-(trifluoromethyl)-1H-benzo[d][1,2,3]triazole-5-carboxylic acid	5-bromo-6-(trifluoromethyl)-1H-benzo[d][1,2,3]triazole (CAS-RN 157590-65-3)	230.1 (M-H) ⁻
43.5	4-chloro-1H-benzo[d][1,2,3]triazole-5-carboxylic acid	5-bromo-4-chloro-1H-benzo[d][1,2,3]triazole (CAS-RN 1388044-33-4)	196.2 (M-H) ⁻
43.6	6-methyl-1H-benzo[d][1,2,3]triazole-5-carboxylic acid	5-bromo-6-methyl-1H-benzo[d][1,2,3]triazole (CAS-RN 1388070-91-4)	176.4 (M-H) ⁻

Example 44**5-bromo-7-fluoro-1H-benzo[d][1,2,3]triazole**

A solution of sodium nitrite (185 mg, 2.68 mmol) in water (0.5 mL) was added dropwise at room temperature to a solution of 5-bromo-3-fluorobenzene-1,2-diamine (500 mg, 2.44 mmol) in water (5 mL) and acetic acid (1.8 mL). The reaction mixture was stirred at room temperature for 1 h, then heated at 85°C for another hour, then partitioned between water and ethyl acetate. The

-253-

organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated to produce the title compound (498 mg, 94%). Light brown solid, MS: 214.1 (M-H)⁻.

The following intermediate was produced according to intermediate 44, replacing 5-bromo-6-fluoro-1H-benzo[d][1,2,3]triazole by the appropriate starting material.

No.	Systematic Name	Starting material	MS, m/e
44.1	5-bromo-4-fluoro-1H-benzo[d][1,2,3]triazole	4-bromo-3-fluorobenzene-1,2-diamine	214.1 (M-H) ⁻

Intermediate 45

3H-[1,2,3]triazolo[4,5-c]pyridine-6-carboxylic acid

Step 1: Methyl 3H-[1,2,3]triazolo[4,5-c]pyridine-6-carboxylate

- 10 A solution of sodium nitrite (413 mg, 5.99 mmol) in water (1.5 mL) was added dropwise at 0°C to a solution of methyl 4,5-diaminopicolinate (CAS-RN 850689-13-3; 910 mg, 5.44 mmol) in water (10 mL) and acetic acid (2 mL) at 0°C, then after 1 h acetic acid (2 mL) was added to the suspension formed. The reaction mixture was stirred for 1 h at 85°C and filtered hot. The precipitate was triturated in methanol to produce the title compound (758 mg, 78%). Red solid,
- 15 MS: 177.2 (M-H)⁻.

Step 2: 3H-[1,2,3]Triazolo[4,5-c]pyridine-6-carboxylic acid

The title compound was produced in analogy to intermediate 43, step 2 from ethyl 3H-[1,2,3]triazolo[4,5-c]pyridine-6-carboxylate. Light brown solid, MS: 163.1 (M-H)⁻.

- 20 The following intermediate was produced according to intermediate 45, replacing methyl 4,5-diaminopicolinate by the appropriate starting material.

No.	Systematic Name	Starting material	MS, m/e
45.1	4-methoxy-1H-benzo[d][1,2,3]triazole-5-carboxylic acid	methyl 3,4-diamino-2-methoxybenzoate (CAS-RN 538372-37-1)	192.3 (M-H) ⁻

Intermediate 46

3-Cyclobutoxy-4-(hydroxymethyl)benzonitrile

5 Step 1: 4-Cyano-2-cyclobutoxybenzoic acid

To a suspension of sodium hydride (60% dispersion in mineral oil, 545 mg, 13.6 mmol) in N,N-dimethylformamide (20 mL) was added cyclobutanol (1.05 g, 13.9 mmol) dropwise below 30°C. The clear solution obtained was stirred for 2 h, then a solution of 4-cyano-2-fluorobenzoic acid (1.00 g, 6.06 mmol) in N,N-dimethylformamide (15 mL) was added dropwise below 35°C. The

10 light yellow suspension was stirred for 66 h at room temperature, then partitioned between water and heptane. The aqueous layer was separated and acidified to pH 2.5 with 3 M aq. hydrochloric acid solution to pH 2.5. The precipitate was collected by filtration, washed with water and dried to afford the title compound (1.03 g, 79%). White foam, MS: 216.2 (M-H)⁻.

Step 2: 3-Cyclobutoxy-4-(hydroxymethyl)benzonitrile

15 Borane dimethyl sulfide complex (429 mg, 5.64 mmol) was added at 0°C to a solution of 4-cyano-2-cyclobutoxybenzoic acid (613 mg, 2.82 mmol) in tetrahydrofuran (8 mL). After 30 min the ice bath was removed, then after 3 h the reaction was stopped by careful addition of water. The reaction mixture was extracted with ethyl acetate, the organic layer was washed with sat. aq. ammonium chloride solution and brine, dried over magnesium sulfate, filtered, and evaporated.

20 Chromatography (silica gel; dichloromethane) afforded the title compound (452 mg, 79%). White solid, MS: 203 (M⁺).

The following intermediate was produced according to intermediate 46, replacing cyclobutanol by the appropriate alcohol.

No.	Systematic Name	Alcohol	MS, m/e
46.1	4-(hydroxymethyl)-3-isopropoxybenzonitrile	2-propanol	191 (M^+)

Intermediate 47

5 (4-Chloro-2-ethoxy-5-fluorophenyl)methanol

Step 1: Ethyl 4-chloro-2-ethoxy-5-fluorobenzoate

To a solution of methyl 4-chloro-2,5-difluorobenzoate (CAS-RN 1214361-01-9; 848 mg, 4.11 mmol) in N,N-dimethylformamide (8 mL) was added a freshly prepared sodium ethoxide solution (94.4 mg/4.11 mmol sodium in 2 mL ethanol) at 0 °C. The reaction mixture was
 10 allowed to reach room temperature over 30 min, then partitioned between ethyl acetate and 1 M hydrochloric acid solution. The organic layer was washed with water and brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel; heptane–ethyl acetate gradient) afforded the title compound (484 mg, 48%) and methyl 4-chloro-2-ethoxy-5-fluorobenzoate (192 mg, 20%). White solid, MS: 247.2 ($M+H$)⁺.

15 Step 2: (4-Chloro-2-ethoxy-5-fluorophenyl)methanol

The title compound was produced in analogy to intermediate 41, step 3 from ethyl 4-chloro-2-ethoxy-5-fluorobenzoate. White solid, MS: 204 (M^+).

The following intermediate was produced according to intermediate 47, replacing methyl 4-chloro-2,5-difluorobenzoate tert-butyl ester hydrochloride and ethanol by the appropriate ester
 20 and alcohol, respectively.

No.	Systematic Name	Ester	Alcohol	MS, m/e

No.	Systematic Name	Ester	Alcohol	MS, m/e
47.1	4-(hydroxymethyl)-3-(2,2,2-trifluoroethoxy)benzonitrile	methyl 4-cyano-2-fluorobenzoate	2,2,2-trifluoroethanol	231 (M ⁺)

Intermediate 48

4-(Hydroxymethyl)-3-(methylsulfonyl)benzonitrile

Step 1: 4-Cyano-2-(methylthio)benzoate

- 5 To a solution of methyl 4-cyano-2-fluorobenzoate (500 mg, 2.79 mmol) in N,N-dimethylformamide (5 mL) was added sodium thiomethoxide (293 mg, 4.19 mmol, Eq: 1.5) at 0°C. After 2 h the reaction mixture was partitioned between sat. aq sodium hydrogen carbonate solution and ethyl acetate 3 times. The organic layer was washed with water and brine, dried over magnesium sulfate, filtered and evaporated to afford the title compound (434 mg, 75%).
- 10 White solid, MS: 207 (M⁺).

Step 2: 4-Cyano-2-(methylsulfonyl)benzoate

- To a of methyl 4-cyano-2-(methylthio)benzoate (420 mg, 2.03 mmol) in dichloromethane (10 mL) was added a suspension of 3-chloroperoxybenzoic acid (1.82 g, 8.11 mmol) in dichloromethane (15 mL) portionwise at 0°C. The reaction mixture was stirred at 0°C for 45 min
- 15 and at room temperature for 45 min, then partitioned between dichloromethane and 1 M sodium sulfite solution. The organic layer was washed with sat. aq. sodium hydrogen carbonate solution and brine, dried over magnesium sulfate, filtered, and evaporated. The residue was triturated in heptane/ethyl acetate 7:3 to afford the title compound (453 g, 93%). White solid, MS: 239 (M⁺).

Step 3: 4-(Hydroxymethyl)-3-(methylsulfonyl)benzonitrile

- 20 A solution of calcium chloride (390 mg, 3.51 mmo) in ethanol (10 mL) was added at room temperature to a solution of methyl 4-cyano-2-(methylsulfonyl)benzoate (420 mg, 1.76 mmol) in tetrahydrofuran (10 mL). Then sodium borohydride (266 mg, 7.02 mmol) was added portionwise over 20 min. After 2 h the reaction mixture was partitioned between sat aq. ammonium chloride

-257-

solution and ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate, filtered, and evaporated. Chromatography (silica gel, dichloromethane/ethanol 19:1) afforded the title compound (123 mg, 33%). White solid, MS: 211 (M^+).

5 Intermediate 49

5-Ethoxy-2-fluoro-4-(hydroxymethyl)benzonitrile

Step 1: Ethyl 4-bromo-2-ethoxy-5-fluorobenzoate

The title compound was produced in analogy to intermediate 47, step 1 from methyl 4-bromo-2,5-difluorobenzoate (CAS-RN 1193162-21-8). White solid, MS: 290 (M^+).

10 Step 2: Ethyl 4-cyano-2-ethoxy-5-fluorobenzoate

A mixture of ethyl 4-bromo-2-ethoxy-5-fluorobenzoate (50mg, 172 μmol), zinc cyanide (11.1 mg, 94.5 μmol), zinc powder (0.4 mg, 7 μmol , Eq: .04), zinc acetate (1.3 mg, 7 μmol , Eq: .04) 1,1'-bis(diphenylphosphino)ferrocene (2.9 mg, 5.2 μmol), and tris(dibenzylideneacetone)dipalladium(0) (1.6 mg, 1.8 μmol) in N,N-dimethylformamide (500 μl) and water (5 μl) was heated at 120°C for 15 min under microwave irradiation, then insoluble material was removed by filtration through diatomaceous earth and the filtrate was evaporated. The residue was purified by chromatography (silica gel; heptane–dichloromethane gradient) to produce the title compound (41 mg, quant.). White solid. MS: 237 (M^+).

Step 3: 5-Ethoxy-2-fluoro-4-(hydroxymethyl)benzonitrile

20 The title compound was produced in analogy to intermediate 41, step 3 from ethyl 4-cyano-2-ethoxy-5-fluorobenzoate. White solid. MS: 195 (M^+).

Example A

A compound of formula (I) can be used in a manner known per se as the active ingredient
25 for the production of tablets of the following composition:

Per tablet

-258-

	Active ingredient	200 mg
	Microcrystalline cellulose	155 mg
	Corn starch	25 mg
	Talc	25 mg
5	Hydroxypropylmethylcellulose	<u>20 mg</u>
		425 mg


Example B

A compound of formula (I) can be used in a manner known per se as the active ingredient for the production of capsules of the following composition:

10		<u>Per capsule</u>
	Active ingredient	100.0 mg
	Corn starch	20.0 mg
	Lactose	95.0 mg
	Talc	4.5 mg
15	Magnesium stearate	<u>0.5 mg</u>
		220.0 mg

Claims

1. Compounds of formula (I)




5 wherein

R^1 is alkyl, haloalkyl, substituted cycloalkyl, substituted cycloalkylalkyl, substituted piperazinyl, substituted piperidinyl, substituted indanyloxyalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted phenylalkynyl, substituted pyridinyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted pyridinylalkynyl substituted thiophenyl, substituted thiophenylalkyl, substituted thiophenylalkenyl, substituted thiophenylalkynyl, naphthyl, substituted naphthyl, quinolyl, substituted quinolinyl, isoquinolyl, substituted isoquinolinyl, substituted 2,3-dihydro-1H-isoindol-2-yl, substituted 1H-indol-2-yl or substituted benzofuran-2-yl wherein substituted

10 cycloalkyl, substituted cycloalkylalkyl, substituted piperazinyl, substituted piperidinyl, substituted indanyloxyalkyl, substituted phenyl, substituted phenylalkyl, substituted phenylalkynyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted pyridinylalkynyl, substituted thiophenyl, substituted thiophenylalkyl, substituted thiophenylalkenyl, substituted thiophenylalkynyl, substituted naphthyl, substituted quinolinyl, substituted isoquinolinyl, substituted 2,3-dihydro-1H-isoindol-2-yl, substituted 1H-indol-2-yl and substituted benzofuran-2-yl are substituted with R^8 , R^9 and R^{10} ;

15

20



25 A is $-\text{N}-$ or CR^5- ;

-260-

W is -O-, -S-, -NR⁶-, -C(O)-, -S(O)₂-, -C(O)-NR⁶- or -CR³R⁴-;


R³ and R⁴ are independently selected from H, halogen, alkyl and cycloalkyl;

R⁵, R⁶ and R⁷ are independently selected from H, alkyl and cycloalkyl;


R⁸, R⁹ and R¹⁰ are independently selected from H, alkyl, hydroxyalkyl, haloalkyl,
5 hydroxyhaloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy,
cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy,
alkoxyhaloalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, phenyl, substituted phenyl,
pyridinyl, substituted pyridinyl, pyrrolyl, substituted pyrrolyl, pyrrolydinyl, substituted
pyrrolydinyl, tetrahydrofuranyl, substituted tetrahydrofuranyl, halogen, hydroxy,
10 cyano, alkylsulfanyl, haloalkylsulfanyl, cycloalkylsulfanyl, alkylsulfinyl,
haloalkylsulfinyl, cycloalkylsulfinyl, alkylcarbonyl, haloalkylcarbonyl,
cycloalkylcarbonyl, alkylsulfonyl, haloalkylsulfonyl, cycloalkylsulfonyl, substituted
aminosulfonyl, substituted amino and substituted aminoalkyl, wherein substituted
aminosulfonyl, substituted amino and substituted aminoalkyl are substituted on the
15 nitrogen atom with one to two substituents independently selected from H, alkyl,
cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, alkylcarbonyl and
cycloalkylcarbonyl, and wherein substituted phenyl, substituted pyrrolyl, substituted
pyrrolydinyl, substituted tetrahydrofuranyl, and substituted pyridinyl are substituted
with one to three substituents independently selected from alkyl, halogen, haloalkyl,
20 alkoxy and haloalkoxy;

m, n, p and q are independently selected from 1 or 2;


-261-

**B**


,

**C**


,

**D**


,

**E**


,

**F**


,

**G**


,

**H**


,

**I**


,

**J**


,

**K**


,

**L**


,

**M**


,

**N**


,

**O**


,

**P**


,

**Q**

,


**R**

,


**S**

,


-262-

**T**


,

**U**


,

**V**


,

**X**


,

**Z**


,

**AA**


,

**AB**


,

**AC**


,

**AD**


,

**AE**


,

**AF**


,

**AG**


,

**AH**

,

**AI**

,

**AJ**

;



R^{11} is H, alkyl, haloalkyl or cycloalkyl;

R^{12} is alkyl, halogen, haloalkyl and alkoxy; R^2 is selected from the ring systems B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, X, Z, AA, AB, AC, AD, AE, AF, AG, AH, AI and AJ;

and pharmaceutically acceptable salts.

2. A compound according to claim 1, wherein

R^1 is alkyl, haloalkyl, substituted cycloalkyl, substituted cycloalkylalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted phenylalkynyl, substituted pyridinyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted pyridinylalkynyl, substituted thiophenyl, substituted thiophenylalkyl, substituted thiophenylalkenyl, substituted thiophenylalkynyl, substituted 2,3-dihydro-1H-isoin-
 5 2-yl, substituted 1H-indol-2-yl or substituted benzofuran-2-yl wherein substituted cycloalkyl, substituted cycloalkylalkyl, substituted phenyl, substituted phenylalkyl, substituted phenylalkynyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted pyridinylalkynyl, substituted thiophenyl, substituted thiophenylalkyl, substituted thiophenylalkenyl, substituted thiophenylalkynyl,
 10 substituted 2,3-dihydro-1H-isoin-2-yl, substituted 1H-indol-2-yl and substituted benzofuran-2-yl are substituted with R^8 , R^9 and R^{10} ;

Y is $-OC(O)-$, $-NR^7C(O)-$, $-C(O)-$, $-S(O)_2-$,  or  ;

A is $-N-$ or CR^5- ;

W is $-O-$, $-S-$, $-NR^6-$, $-C(O)-$, $-S(O)_2-$, $-C(O)-NR^6-$ or $-CR^3R^4-$;

20 R^3 and R^4 are independently selected from H, halogen, alkyl and cycloalkyl;


R^5 , R^6 and R^7 are independently selected from H, alkyl and cycloalkyl;

R^8 , R^9 and R^{10} are independently selected from H, alkyl, hydroxyalkyl, haloalkyl, hydroxyhaloalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkoxy, cycloalkoxy, cycloalkoxyalkyl, cycloalkylalkoxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, alkoxyhaloalkyl, alkoxyalkoxy, alkoxyalkoxyalkyl, phenyl, substituted phenyl, pyridinyl, substituted pyridinyl, halogen, hydroxy, cyano, alkylsulfanyl, haloalkylsulfanyl, cycloalkylsulfanyl, alkylsulfinyl, haloalkylsulfinyl,


5 cycloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, cycloalkylsulfonyl, substituted aminosulfonyl, substituted amino and substituted aminoalkyl, wherein substituted aminosulfonyl, substituted amino and substituted aminoalkyl are substituted on the nitrogen atom with one to two substituents independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, alkylcarbonyl and cycloalkylcarbonyl, and wherein substituted phenyl and substituted pyridinyl are optionally substituted with one to three substituents independently selected from alkyl, halogen, haloalkyl, alkoxy and haloalkoxy;

m, n, p and q are independently selected from 1 or 2;


-265-

**B**


,

**C**


,

**D**


,

**E**


,

**F**


,

**G**


,

**H**


,

**I**


,

**J**


,

**K**


,

**L**


,

**M**


,

**N**


,

**O**


,

**P**


,

**Q**

,


**R**

,


**S**

,


-266-

**T**


,

**U**


,

**V**


,

**X**


,

**Z**


,

**AA**


,

**AB**


,

**AC**

,

**AD**

,

**AE**

;

R^{11} is H, alkyl, haloalkyl or cycloalkyl;

R^2 is selected from the ring systems B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, X, Z, AA, AB, AC, AD and AE;

5 and pharmaceutically acceptable salts.

3. A compound according to claim 1, wherein R^1 is substituted cycloalkylalkyl, substituted piperazinyl, substituted piperidinyl, substituted indanyloxyalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, naphthyl, substituted naphthyl, substituted quinolinyl, substituted isoquinolinyl, or substituted 1H-
- 10

-267-


indol-2-yl, wherein substituted cycloalkylalkyl, substituted piperazinyl, substituted piperidinyl, substituted indanyloxyalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, substituted naphthyl, substituted quinolinyl, substituted isoquinolinyl and substituted 1H-indol-2-yl are substituted with R⁸, R⁹ and R¹⁰.

4. A compound according to any one of claims 1 to 3, wherein R¹ is substituted cycloalkylalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinylalkyl, substituted pyridinylalkenyl or substituted 1H-indol-2-yl, wherein substituted cycloalkylalkyl, substituted phenyl, substituted phenylalkyl, substituted phenoxyalkyl, substituted phenylcycloalkyl, substituted phenylalkenyl, substituted pyridinylalkyl, substituted pyridinylalkenyl, and substituted 1H-indol-2-yl are substituted with R⁸, R⁹ and R¹⁰.

5. A compound according to any one of claims 1 to 4, wherein R¹ is substituted phenylalkyl or substituted phenylalkenyl, wherein substituted phenylalkyl and substituted phenylalkenyl are substituted with R⁸, R⁹ and R¹⁰.

6. A compound according to any one of claims 1 to 5, wherein R¹ is phenylalkyl substituted with R⁸, R⁹ and R¹⁰.

7. A compound according to any one of claims 1 to 6, wherein Y is -OC(O)-, -C(O)-, -S(O)₂-



8. A compound according to any one of claims 1 to 7, wherein Y is -OC(O)- or -C(O)-.

9. A compound according to any one of claims 1 to 8, wherein Y is -OC(O)-.

10. A compound according to any one of claims 1 to 8, wherein Y is -C(O)-.

11. A compound according to any one of claims 1 to 10, wherein A is -N-.

12. A compound according to any one of claims 1 to 11, wherein W is -O-, -NR⁶-, -C(O)-, -S(O)₂-, -C(O)-NR⁶- or -CR³R⁴-.
13. A compound according to any one of claims 1 to 12, wherein W is -C(O)-, -C(O)-NR⁶- or -CR³R⁴-.
- 5 14. A compound according to any one of claims 1 to 13, wherein W is -C(O)-.
15. A compound according to any one of claims 1 or 3 to 14, wherein R² is selected from the ring systems B, C, D, E, G, H, M, O, P, R, S, T, U, V, X, Z, AA, AB, AC, AD, AE, AF, AG, AH and AI.
- 10 16. A compound according to any one of claims 1 to 15, wherein R² is selected from the ring systems B, C, D, E, G, H, M, O, P, R, S, T, U, V, X, Z, AA, AB, AC, AD and AE.
17. A compound according to any one of claims 1 to 16, wherein R² is selected from the ring systems B, D, H, M, O, R and AJ.
18. A compound according to any one of claims 1 to 17, wherein R² is selected from the ring systems B, D, H, O and R.
- 15 19. A compound according to any one of claims 1 to 18, wherein R² is selected from the ring systems B and D.
20. A compound according to any one of claims 1 to 19, wherein R² is the ring system D.
21. A compound according to any one of claims 1 to 20, wherein R³ and R⁴ are H.
22. A compound according to any one of claims 1 to 10 and 12 to 21, wherein R⁵ is H.
- 20 23. A compound according to any one of claims 1 to 13 and 14 to 22, wherein R⁶ is H or alkyl.
24. A compound according to any one of claims 1 to 23, wherein R⁸, R⁹ and R¹⁰ are independently selected from H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen, cyano, haloalkylsulfanyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, pyrrolyl substituted with one alkyl, pyrrolydiny, 25 tetrahydrofuranyl, alkylcarbonyl, and aminosulfonyl substituted on the nitrogen atom with

one to two substituents independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, alkylcarbonyl and cycloalkylcarbonyl.

25. A compound according to any one of claims 1 to 24, wherein R^8 , R^9 and R^{10} are independently selected from H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen, cyano, haloalkylsulfanyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl and aminosulfonyl substituted on the nitrogen atom with one to two substituents independently selected from H, alkyl, cycloalkyl, cycloalkylalkyl, hydroxyalkyl, alkoxyalkyl, alkylcarbonyl and cycloalkylcarbonyl.
- 5
26. A compound according to any one of claims 1 to 25, wherein R^8 , R^9 and R^{10} are independently selected from H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen, cyano, haloalkylsulfanyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl and aminosulfonyl substituted on the nitrogen atom with two alkyl.
- 10
27. A compound according to any one of claims 1 to 26, wherein R^8 , R^9 and R^{10} are independently selected from H, alkyl, haloalkyl, haloalkoxy, halogen and alkylsulfonyl.
- 15
28. A compound according to any one of claims 1 to 27, wherein R^8 is H, alkyl, haloalkyl, hydroxyhaloalkyl, alkoxy, haloalkoxy, alkoxyhaloalkyl, phenyl, pyridinyl, halogen, cyano, haloalkylsulfanyl, haloalkylsulfinyl, alkylsulfonyl, haloalkylsulfonyl, pyrrolyl substituted with one alkyl, pyrrolydinyl, tetrahydrofuranyl, alkylcarbonyl, or aminosulfonyl substituted on the nitrogen atom with two alkyl.
- 20
29. A compound according to any one of claims 1 to 28, wherein R^8 is haloalkyl, haloalkoxy, halogen or alkylsulfonyl.
30. A compound according to any one of claims 1 to 29, wherein R^8 is haloalkoxy or halogen.
31. A compound according to any one of claims 1 to 30, wherein R^8 is halogen.
- 25
32. A compound according to any one of claims 1 to 31, wherein R^9 is H, alkyl, haloalkyl, cycloalkyl, cycloalkoxy, alkoxy, haloalkoxy, alkoxyalkoxy, cyano or halogen.
33. A compound according to any one of claims 1 to 32, wherein R^9 is H, alkyl, haloalkyl, alkoxy or halogen.

34. A compound according to any one of claims 1 to 33, wherein R⁹ is H, alkyl or halogen.
35. A compound according to any one of claims 1 to 34, wherein R⁸ and R⁹ are halogen.
36. A compound according to any one of claims 1 to 35, wherein R¹⁰ is H, alkyl, alkoxy or halogen.
37. A compound according to any one of claims 1 to 36, wherein R¹⁰ is H or alkyl.
38. A compound according to any one of claims 1 to 37, wherein R¹⁰ is H.
39. A compound according to any one of claims 1 to 38, wherein R¹¹ is haloalkyl.
40. A compound according to any one of claims 1 to 39, wherein m is 1.
41. A compound according to any one of claims 1 to 40, wherein n is 1.
42. A compound according to any one of claims 1 to 41, wherein m and n are 1.
43. A compound according to any one of claims 1 to 42, wherein p and q are 1.
44. A compound according to any one of claims 1 to 43, wherein m, n, p and q are 1.
45. A compound according to any one of claims 1 to 44, selected from
 - (E)-1-[(3aS,8aR)-2-(4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;
 - 1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)propan-1-one;
 - 6-((3aR,6aS)-5-(3-(3,5-dichlorophenyl)propanoyl)octahydropyrrolo[3,4-c]pyrrole-2-carbonyl)benzo[d]oxazol-2(3H)-one;
 - (3aR,6aS)-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;
 - (3aR,6aS)-3,5-dichlorobenzyl 5-(9H-pyrido[3,4-b]indole-3-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

-271-

(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-indole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aR,6aS)-3,5-dichlorobenzyl 5-(9H-carbazole-3-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

5 (3aR,6aS)-3,5-dichlorobenzyl 5-(1H-indazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aR,6aS)-3,5-dichlorobenzyl 5-(1H-benzo[d]imidazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

10 trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

cis-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

(3aR,8aS)-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepine-6(7H)-carboxylate;

15 (1H-benzotriazol-5-yl)-{(3aS,6aR)-5-[2-(3-chloro-phenyl)-ethanesulfonyl]-hexahydropyrrolo[3,4-c]pyrrol-2-yl}-methanone;

(3aR,6aS)-3,5-dichlorobenzyl 5-(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

20 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chlorophenyl)-2,2-dimethylpropan-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(3aSR,6SR,7aSR)-6-{(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl}-hexahydro-benzooxazol-2-one;

25 (E)-1-[(3aS,8aR)-2-(benzo[c][1,2,5]oxadiazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

-272-

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-methyl-4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-methoxyphenyl)prop-2-en-1-one;

5 (E)-1-[(3aS,8aR)-2-((S)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one

(E)-1-[(3aS,8aR)-2-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one (E)-1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(2-isopropyl-phenyl)-prop-2-ene-1-one;

10 trans-3,5-dichlorobenzyl 2-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

(3aR,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

15 6-[(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl]-3H-benzooxazol-2-one;

(3aR,5s,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yloxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate;

20 (3aR,5r,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-yloxy)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate;

(3aS,6aS)-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-2-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester;

25 (3aR,6aR)-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

-273-

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

5 (3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester;

(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

10 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-methanesulfonyl-5-trifluoromethyl-benzyl ester;

cis-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate;

(3aS,7aR)-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

15 trans-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate;

(3aR,8aS)-3,5-dichlorobenzyl 6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxylate;

20 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 1-(3-chloro-phenyl)-cyclopropyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid bicyclo[4.1.0]hept-7-ylmethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid adamantan-2-ylmethyl ester;

25 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 1-fluoro-cyclohexylmethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-adamantan-2-yl-ethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-adamantan-1-yl-ethyl ester;

5 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid adamantan-1-ylmethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid cyclohexylmethyl ester;

10 cis-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2,2-trifluoro-1-methoxy-ethyl)-benzyl ester;

cis-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-(2,2,2-trifluoro-1-hydroxy-ethyl)-benzyl ester;

(3aR,6aS)-2-cyclohexylethyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

15 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-5-trifluoromethoxy-benzyl ester;

(3aR,6aS)-3-chloro-5-cyanobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

20 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-trifluoromethoxy-benzyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-5-trifluoromethyl-benzyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-trifluoromethoxy-benzyl ester;

25 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-fluoro-3-trifluoromethoxy-benzyl ester;

(3aR,6aS)-3-cyano-5-fluorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(3aR,6aS)-3-chloro-5-methoxybenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

5 (3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1S,4R)-3-methyl-bicyclo[2.2.1]hept-2-ylmethyl ester;

(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1R,4S)-1-bicyclo[2.2.1]hept-2-ylmethyl ester;

10 (3aR,5S,6aS)-5-[(3H-[1,2,3]triazol-4-ylmethyl)-carbamoyl]-hexahydro-cyclopenta[c]pyrrole-2-carboxylic acid 3,5-dichloro-benzyl ester;

(3aS,6aS)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

(3aS,6aR)-5-[(1H-[1,2,3]triazol-4-ylmethyl)-carbamoyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3,5-dichloro-benzyl ester;

15 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aR,5r,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-ylamino)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate;

20 (3aR,6aS)-3,5-dichlorobenzyl 5-((1H-benzo[d]imidazol-5-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

1-((3aR,6aS)-5-((1H-benzo[d][1,2,3]triazol-5-yl)methyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)propan-1-one;

(3aR,6aS)-3,5-dichlorobenzyl 5-((1H-indazol-5-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

25 (3aR,6aS)-3,5-dichlorobenzyl 5-((2-oxo-2,3-dihydrobenzo[d]oxazol-6-yl)methyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

-276-

6-(((3aR,6aS)-5-(3-(3,5-dichlorophenyl)propanoyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methyl)benzo[d]oxazol-2(3H)-one;

4-{(E)-3-[(3aS,8aR)-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-oxo-propenyl}-benzonitrile;

5 (E)-1-[(3aS,8aR)-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(3aR,6aS)-3,5-dichlorobenzyl 5-(2-oxo-2,3-dihydrobenzo[d]oxazol-6-ylsulfonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

10 (3aR,6aS)-3,5-dichlorobenzyl 5-(4,5,6,7-tetrahydro-1H-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate;

(E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-pyrazolo[4,3-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

15 cis-5-((3aR,8aS)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)decahydropyrrolo[3,4-d]azepine-2-carbonyl)hexahydrooxazolo[5,4-c]pyridin-2(1H)-one;

6-{(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl}-5,6,7,8-tetrahydro-1H-[1,6]naphthyridin-2-one;

20 (3aR,7aR)-5-{(3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carbonyl}-hexahydro-oxazolo[5,4-c]pyridin-2-one;

(E)-1-[(3aS,8aR)-2-(7,8-Dihydro-5H-[1,6]naphthyridine-6-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

25 (E)-3-(4-trifluoromethoxy-phenyl)-1-[(3aS,8aR)-2-(2-trifluoromethyl-5,6-dihydro-8H-[1,2,4]triazolo[1,5-a]pyrazine-7-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-prop-2-ene-1-one;

-277-

(3aS,8aR)-6-[(E)-3-(3-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid (1H-[1,2,3]triazol-4-ylmethyl)-amide;

(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

5 (3aS,6aR)-5-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid (1H-[1,2,3]triazol-4-ylmethyl)-amide;

(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

10 (3aS,8aR)-6-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid (4H-[1,2,4]triazol-3-ylmethyl)-amide;

(E)-1-[(3aS,8aR)-2-(6,7-dihydro-4H-[1,2,3]triazolo[1,5-a]pyrazine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aS,8aR)-2-(1,4,6,7-tetrahydro-imidazo[4,5-c]pyridine-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

15 (3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

(3aS,8aR)-6-[3-(3-chloro-phenyl)-2,2-dimethyl-propionyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid (3H-[1,2,3]triazol-4-ylmethyl)-amide;

20 (3aR,8aS)-N-(2-(1H-1,2,3-triazol-5-yl)ethyl)-6-((E)-3-(4-(trifluoromethoxy)phenyl)acryloyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

(3aR,7aS)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3,5-dichloro-benzyl ester;

(3aS,7aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3,5-dichloro-benzyl ester;

25 (+)-trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

-278-

(-)-trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

(-)-trans-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate;

5 (+)-trans-3,5-dichlorobenzyl 5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate;

(E)-1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(3,5-dichloro-phenyl)-prop-2-en-1-one;

10 (1H-benzotriazol-5-yl)-{trans-2-[5-(4-chloro-phenyl)-[1,3,4]oxadiazol-2-yl]-octahydro-pyrrolo[3,4-c]pyridin-5-yl}-methanone;

(E)-1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(1H-benzotriazol-5-yl)-[(3aR,6aS)-5-(5-chloro-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

15 (E)-1-[(3aR,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-5-trifluoromethyl-phenyl)-prop-2-ene-1-one;

1-[(3aR,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-5-trifluoromethyl-phenyl)-propan-1-one;

20 (1H-benzotriazol-5-yl)-[(3aR,6aS)-5-(6-chloro-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylsulfonyl)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-chlorophenyl)prop-2-en-1-one;

25 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-p-tolylprop-2-en-1-one;

4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)-N,N-dimethylbenzenesulfonamide;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxyphenyl)prop-2-en-1-one;

5 (E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)benzonitrile;

10 (E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluorophenyl)prop-2-en-1-one;

15 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-phenylprop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-2-yl)prop-2-en-1-one;

20 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-3-yl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chlorophenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(4-chlorophenyl)prop-2-en-1-one;

25 (E)-1-((3aR,8aS)-6-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-2(1H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(difluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one;

5 4-((E)-3-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-oxoprop-1-enyl)benzonitrile;

(E)-1-((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

10 (-)-(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(+)-(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3,5-dichlorophenyl)prop-2-en-1-one;

15 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(pyridin-4-yl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2,4-difluorophenyl)prop-2-en-1-one;

20 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2,4-dichlorophenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3,4-dichlorophenyl)prop-2-en-1-one;

(E)-1-[(3aS,7aS)-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-difluoromethoxy-phenyl)-prop-2-ene-1-one;

25 4-{(E)-3-[(3aS,7aS)-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-oxo-propenyl}-benzonitrile;

4-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)-3-fluorobenzonitrile;

4-((E)-3-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-oxoprop-1-enyl)-3-fluorobenzonitrile;

5 (E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(difluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-[cis-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

10 3-((E)-3-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-oxoprop-1-enyl)benzonitrile;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(2-fluoro-4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

15 (E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-chloro-2-fluorophenyl)prop-2-en-1-one;

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(3,5-dichlorophenyl)prop-2-en-1-one;

20 (E)-1-((3aR,6aS)-5-(4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

25 (E)-1-[(3aS,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-methanesulfonyl-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3,5-dimethoxy-phenyl)-prop-2-ene-1-one;

5 (E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-chloro-5-methoxy-phenyl)-prop-2-ene-1-one;

10 3-[(E)-3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-oxo-propenyl]-5-chloro-benzonitrile;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-methoxy-5-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-ene-1-one;

15 1-[(3aR,6aR)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

(3aR,7aR)-5-[(3aR,6aR)-5-[(E)-3-(4-trifluoromethoxy-phenyl)-acryloyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carbonyl]-hexahydro-oxazolo[5,4-c]pyridin-2-one;

20 (3aR,7aR)-5-[(3aR,6aR)-5-[3-(4-trifluoromethoxy-phenyl)-propionyl]-hexahydro-pyrrolo[3,4-c]pyrrole-2-carbonyl]-hexahydro-oxazolo[5,4-c]pyridin-2-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-phenyl-prop-2-ene-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-phenyl-propan-1-one;

25 (E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-prop-2-ene-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-propan-1-one;

(3aR,6aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-5-(3-(4-(trifluoromethoxy)phenyl)propanoyl)hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxamide;

5 (3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-((E)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)acryloyl)-N-methyloctahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-trifluoromethyl-pyridin-3-yl)-propan-1-one;

10 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-trifluoromethoxy-phenoxy)-ethanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

15 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-biphenyl-4-yl-propan-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluoro-2-(trifluoromethyl)phenyl)prop-2-en-1-one;

1-[(3aS,6aR)-5-(1H-Benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

20 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(methylsulfonyl)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylthio)phenyl)prop-2-en-1-one;

25 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-2-(3-(trifluoromethoxy)phenoxy)ethanone;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-(trifluoromethoxy)phenyl)prop-2-en-1-one;

5 (E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3-trifluoromethoxy-phenyl)-prop-2-en-1-one;

(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(4-trifluoromethoxy-phenyl)-prop-2-en-1-one;

10 (E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3-chloro-5-trifluoromethoxy-phenyl)-prop-2-en-1-one;

(E)-1-[trans-5-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-2-yl]-3-(3,5-dichloro-phenyl)-prop-2-en-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-phenyl-pyridin-3-yl)-prop-2-ene-1-one;

15 (E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-trifluoromethyl-pyridin-2-yl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-4-yl-phenyl)-prop-2-ene-1-one;

20 (E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-3-yl-phenyl)-prop-2-ene-1-one;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-2-yl-phenyl)-prop-2-ene-1-one;

1-[(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-3-methyl-phenoxy)-ethanone;

25 1-[(3aS,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-methyl-phenoxy)-ethanone;

(E)-1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-phenyl-pyridin-2-yl)-prop-2-ene-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-(trifluoromethylsulfinyl)phenyl)prop-2-en-1-one;

5 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-(trifluoromethoxy)phenyl)propan-1-one;

(3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-N-methyl-6-(3-(4-(trifluoromethoxy)phenyl)propanoyl)octahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

10 (3aR,8aS)-N-((1H-1,2,3-triazol-5-yl)methyl)-6-(3-(3-fluoro-4-(trifluoromethoxy)phenyl)propanoyl)-N-methyloctahydropyrrolo[3,4-d]azepine-2(1H)-carboxamide;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-difluoromethoxy-phenyl)-propan-1-one;

15 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-fluoro-4-trifluoromethoxy-phenyl)-propan-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-fluoro-2-(trifluoromethyl)phenyl)propan-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-methyl-4-(trifluoromethoxy)phenyl)propan-1-one;

20 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-fluoro-4-methoxyphenyl)propan-1-one;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-3-(2-isopropyl-phenyl)-propan-1-one;

25 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-trifluoromethyl-pyridin-2-yl)-propan-1-one;

-286-

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(5-phenyl-pyridin-2-yl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-4-yl-phenyl)-propan-1-one;

5 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-3-yl-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-pyridin-2-yl-phenyl)-propan-1-one;

10 (3aS,8aR)-6-[3-(4-trifluoromethoxy-phenyl)-propionyl]-octahydro-pyrrolo[3,4-d]azepine-2-carboxylic acid [2-(3H-[1,2,3]triazol-4-yl)-ethyl]-amide;

(E)-3-[4-(trifluoro-methoxy)-phenyl]-1-[(3aS,8aR)-2-((S)-4,5,6,7-tetrahydro-1H-imidazo[4,5-c]pyridine-6-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-prop-2-ene-1-one hydrochloride;

and pharmaceutically acceptable salts thereof.

15 46. A compound according to any one of claims 1 to 44, selected from

trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 4-trifluoromethoxy-benzyl ester;

20 1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-2-(4-trifluoromethoxy-phenoxy)-ethanone;

(E)-1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propenone;

25 (E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chloro-5-(trifluoromethyl)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxy-2-(trifluoromethyl)phenyl)prop-2-en-1-one;

(E)-1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-cyclopropylphenyl)prop-2-en-1-one;

5 trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 4-fluoro-2-trifluoromethyl-benzyl ester;

trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 2-cyclopropyl-4-trifluoromethyl-benzyl ester;

10 1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(2-trifluoromethoxy-phenoxy)-ethanone;

trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridine-5-carboxylic acid 2-methoxy-4-trifluoromethoxy-benzyl ester;

4-{2-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-oxo-ethoxy}-3-trifluoromethyl-benzonitrile;

15 1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-[4-methyl-2-(1-methyl-pyrrolidin-3-yl)-phenoxy]-ethanone;

20 1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepin-6-yl]-2-(2-chloro-4-fluoro-phenoxy)-ethanone;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(2-chloro-4-(trifluoromethyl)phenoxy)ethanone;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(6-isopropyl-3,3-dimethyl-2,3-dihydro-1H-inden-5-yloxy)ethanone;

25 (3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(5-chloro-2-(trifluoromethyl)phenoxy)ethanone;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-2-(2-tert-butyl-4-methoxyphenoxy)ethanone;

5 4-[2-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-2-oxoethoxy]-3-propan-2-ylbenzonitrile;

1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-3-[3-fluoro-4-(2,2,2-trifluoroethoxy)phenyl]propan-1-one;

10 1-[(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,5,7,8,8a-octahydropyrrolo[3,4-d]azepin-6-yl]-3-[2-fluoro-4-(2,2,2-trifluoroethoxy)phenyl]propan-1-one;

(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 3-fluoro-4-(2,2,2-trifluoro-ethoxy)-benzyl ester;

(3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 2-fluoro-4-(2,2,2-trifluoro-ethoxy)-benzyl ester;

15 (3aS,8aR)-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-d]azepine-6-carboxylic acid 4-(2,2,2-trifluoro-ethoxy)-benzyl ester;

(3aS,6aS)-5-(3H-[1,2,3]triazolo[4,5-b]pyridine-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

20 1-[(3aR,6aR)-5-(1H-triazolo[4,5-b]pyridine-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[4-(trifluoromethoxy)phenyl]propan-1-one;

(3aS,6aS)-5-(3H-[1,2,3]triazolo[4,5-c]pyridine-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(4-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

25 (3aS,6aS)-5-(7-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(6-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(4-chloro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

5 (3aS,6aS)-5-(6-trifluoromethyl-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(4-methyl-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

10 (3aS,6aS)-5-(6-methyl-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

1-[(3aR,6aR)-5-(4-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

(4-ethoxyquinolin-2-yl)((3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone;

15 (4-ethoxyquinolin-2-yl)((3aS,6aS)-5-(4-fluoro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)methanone;

6-[(3aR,6aR)-2-[3-[4-(trifluoromethoxy)phenyl]propanoyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrole-5-carbonyl]-3H-1,3-benzoxazol-2-one;

20 (3aS,6aS)-5-(2-oxo-2,3-dihydro-benzooxazole-6-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-(1,1,2,2-tetrafluoro-ethoxy)-benzyl ester;

25 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-difluoromethoxy-3-fluoro-benzyl ester;

(3aS,6aS)-5-(1H-[1,2,3]triazolo[4,5-b]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-difluoromethoxy-benzyl ester;

5 (3aS,6aS)-5-(1H-[1,2,3]triazolo[4,5-b]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester;

10 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-(2,2,2-trifluoro-ethoxy)-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 5-trifluoromethoxy-pyridin-2-ylmethyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-benzyl ester;

15 (3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-5-methyl-benzyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester;

20 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-ethoxy-benzyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-fluoro-4-trifluoromethoxy-benzyl ester;

25 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropyl-5-methyl-benzyl ester;

(3aS,6aS)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

5 (3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 2-fluoro-4-trifluoromethyl-benzyl ester;

10 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethyl-benzyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethyl-benzyl ester;

15 (3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-methanesulfonyl-benzyl ester;

20 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-ethoxy-5-fluoro-benzyl ester;

(3aS,6aS)-5-(4-methoxy-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-cyclobutoxy-benzyl ester;

25 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-isopropoxy-benzyl ester;

-292-

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-cyano-2-(2,2,2-trifluoro-ethoxy)-benzyl ester;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-chloro-2-ethoxy-5-fluoro-benzyl ester;

5 (E)-1-[trans-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propenone;

(3aS,6aS)-5-(4-sulfamoyl-piperidine-1-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

10 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(6-phenyl-pyridin-3-yl)-propan-1-one;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-isopropyl-phenoxy)-ethanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-trifluoromethyl-phenoxy)-ethanone;

15 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(biphenyl-2-yloxy)-ethanone;

(E)-1-[(3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propenone;

20 1-((3aR,6aR)-5-((R)-4,5,6,7-tetrahydro-1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-chloro-4-trifluoromethoxy-phenoxy)-ethanone;

25 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-pyrrol-1-yl-phenoxy)-ethanone;

-293-

4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-methoxy-benzonitrile;

4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-benzonitrile;

5 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-phenoxy-ethanone;

2-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-5-trifluoromethoxy-benzonitrile;

10 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-2-(2-isopropyl-5-methylphenoxy)ethanone;

(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(6-trifluoromethoxy-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(5-trifluoromethoxy-1H-indole-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

15 1-[trans-2-(1H-benzotriazole-5-carbonyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

1-[trans-2-(1H-benzotriazol-5-ylmethyl)-octahydro-pyrrolo[3,4-c]pyridin-5-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

20 1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(3-chloro-5-(trifluoromethyl)phenyl)propan-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(4-methoxy-2-(trifluoromethyl)phenyl)propan-1-one;

1-((3aR,8aS)-2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)octahydropyrrolo[3,4-d]azepin-6(7H)-yl)-3-(2-cyclopropylphenyl)propan-1-one;

25 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[3-methoxy-5-(trifluoromethoxy)phenyl]propan-1-one;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-isopropyl-5-methyl-phenoxy)-ethanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-bromo-4-trifluoromethoxy-phenoxy)-ethanone;

5 (1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(4'-chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-isopropyl-benzonitrile;

10 2-(2-Acetyl-phenoxy)-1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-ethanone;

4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-5-isopropyl-2-methyl-benzonitrile;

(1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

15 (1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(4-methoxy-naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

4-{2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-oxo-ethoxy}-3-ethoxy-benzonitrile;

20 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-4-trifluoromethoxy-phenyl)-propan-1-one;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-phenoxy)-ethanone;

[(3aS,6aS)-5-(4'-Chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-(R)-4,5,6,7-tetrahydro-1H-benzotriazol-5-yl-methanone;

25 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

(1H-benzotriazol-5-yl)-[(3aS,6aS)-5-(4'-chloro-biphenyl-4-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-[2-(tetrahydro-furan-2-yl)-phenoxy]-ethanone;

5 (1H-benzotriazol-5-yl)-[(3aR,6aR)-5-(4-methoxy-naphthalene-2-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-methanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(2-tert-butyl-phenoxy)-ethanone;

10 [(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-[trans-4-(4-chloro-phenyl)-cyclohexyl]-methanone;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(3-fluoro-4-trifluoromethyl-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(2-fluoro-4-trifluoromethyl-phenyl)-propan-1-one;

15 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(2-pyridin-3-ylphenoxy)ethanone;

4-[3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-oxopropyl]-2-methyl-5-propan-2-ylbenzonitrile;

20 4-[3-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-oxopropyl]-3-propan-2-ylbenzonitrile;

[(3aR,6aR)-5-[1-(4-chlorophenyl)piperidine-4-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(1H-benzotriazol-5-yl)methanone;

[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(4-propan-2-yloxynaphthalen-2-yl)methanone;

25 [(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(4-propan-2-yloxyquinolin-2-yl)methanone;

1-[(3aR,6aR)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-3-[2-fluoro-4-(trifluoromethoxy)phenyl]propan-1-one;

5 4-[2-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-2-methyl-5-propan-2-ylbenzonitrile;

[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-[1-(2,2,2-trifluoroethoxy)isoquinolin-3-yl]methanone;

1-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(4-bromo-2-tert-butylphenoxy)ethanone;

10 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(4-bromo-2-tert-butylphenoxy)ethanone;

4-[2-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-3-tert-butylbenzonitrile;

15 4-[2-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-oxoethoxy]-3-tert-butylbenzonitrile;

[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-[1-methyl-5-(trifluoromethoxy)indol-2-yl]methanone;

1-[(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-[4-(trifluoromethoxy)phenoxy]ethanone;

20 [(3aS,6aS)-5-[(5R)-4,5,6,7-tetrahydro-1H-benzotriazole-5-carbonyl]-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-(1-ethoxyisoquinolin-3-yl)methanone;

1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-2-yl]-2-(2-tert-butyl-4-methoxyphenoxy)ethanone;

25 ((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(4-ethoxyquinolin-2-yl)methanone;

-297-

((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(4-(2,2,2-trifluoroethoxy)quinolin-2-yl)methanone;

((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(6-cyclobutoxy-5-(trifluoromethyl)pyridin-3-yl)methanone;

5 ((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-bromo-6-(2-methoxyethoxy)pyridin-3-yl)methanone;

((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-bromo-6-(cyclopropylmethoxy)pyridin-3-yl)methanone;

10 ((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(5-cyclopropyl-6-(2,2,2-trifluoroethoxy)pyridin-3-yl)methanone;

((3aS,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)(6-(2,2,2-trifluoroethoxy)-5-(trifluoromethyl)pyridin-3-yl)methanone;

(1H-benzotriazol-5-yl)-{(3aS,6aS)-5-[4-(4-chloro-phenyl)-piperidine-1-carbonyl]-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl}-methanone;

15 (1H-benzotriazol-5-yl)-{(3aS,6aS)-5-[4-(4-chloro-phenyl)-piperazine-1-carbonyl]-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl}-methanone;

and pharmaceutically acceptable salts thereof.

47. A compound according to any one of claims 1 to 45, selected from

20 trans-3,5-dichlorobenzyl 2-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

trans-3,5-dichlorobenzyl 2-(2-oxo-2,3-dihydrobenzo[d]oxazole-6-carbonyl)hexahydro-1H-pyrrolo[3,4-c]pyridine-5(6H)-carboxylate;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 3-chloro-5-methanesulfonyl-benzyl ester;

25 (3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

-298-

(E)-1-((3aR,6aR)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)prop-2-en-1-one;

1-((3aR,6aS)-5-(1H-benzo[d][1,2,3]triazole-5-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(4-(trifluoromethoxy)phenyl)propan-1-one;

5 1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

1-[(3aR,6aR)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethyl-phenyl)-propan-1-one;

10 1-[(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-2-(4-chloro-2-isopropyl-5-methyl-phenoxy)-ethanone;

and pharmaceutically acceptable salts thereof.

48. A compound according to any one of claims 1 to 44 or 46, selected from

(3aS,6aS)-5-(4-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

15 1-[(3aR,6aR)-5-(4-fluoro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrol-2-yl]-3-(4-trifluoromethoxy-phenyl)-propan-1-one;

(3aS,6aS)-5-(1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-difluoromethoxy-3-fluoro-benzyl ester;


20 (3aS,6aS)-5-(1,4,6,7-tetrahydro-[1,2,3]triazolo[4,5-c]pyridine-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

(3aS,6aS)-5-((R)-4,5,6,7-Tetrahydro-1H-benzotriazole-5-carbonyl)-hexahydro-pyrrolo[3,4-c]pyrrole-2-carboxylic acid 4-trifluoromethoxy-benzyl ester;

and pharmaceutically acceptable salts thereof.

25 49. A process to prepare a compound according to any one of claims 1 to 48 comprising the reaction of a compound of formula (II) in the presence of a compound of formula (III), wherein R^1 , R^2 , A, W, m, n, p and q are as defined above, Y is -OC(O)-.

-299-



50. A compound according to any one of claims 1 to 48 for use as therapeutically active substance.
51. A pharmaceutical composition comprising a compound according to any one of claims 1 to 48 and a therapeutically inert carrier.
52. The use of a compound according to any one of claims 1 to 48 for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, fibrotic diseases and acute and chronic organ transplant rejection.
53. A compound according to any one of claims 1 to 48 for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, fibrotic diseases and acute and chronic organ transplant rejection.
54. The use of a compound according to any one of claims 1 to 48 for the preparation of a medicament for the treatment or prophylaxis of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, fibrotic diseases and acute and chronic organ transplant rejection.
55. A method for the treatment or prophylaxis a renal condition selected from the group consisting of renal conditions, liver conditions, inflammatory conditions, conditions of the nervous system, fibrotic diseases and acute and chronic organ transplant rejection, which method comprises administering an effective amount of a compound according to any one of claims 1 to 48.
56. A compound according to any one of claims 1 to 48, when manufactured according to a process of claim 49.
57. The invention as hereinbefore described.