



(11) **EP 2 402 465 A2**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**04.01.2012 Bulletin 2012/01**

(51) Int Cl.:  
**C12Q 1/70 (2006.01) C12Q 1/68 (2006.01)**

(21) Application number: **11176258.9**

(22) Date of filing: **13.07.2005**

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI  
SK TR**

(30) Priority: **13.07.2004 US 587734 P**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**05800781.6 / 1 771 585**

(71) Applicant: **GEN-PROBE INCORPORATED**  
**San Diego, CA 92121-4362 (US)**

(72) Inventors:  
• **Carlson, James D.**  
**San Diego, CA 92019 (US)**

• **Brentano, Steven T.**  
**Santee, CA 92071 (US)**

(74) Representative: **Viering, Jentschura & Partner**  
**Kennedydamm 55/Roßstr.**  
**40476 Düsseldorf (DE)**

Remarks:

- Claims filed after the date of the date of receipt of the divisional application (Rule 68(4) EPC).
- This application was filed on 02-08-2011 as a divisional application to the application mentioned under INID code 62.

(54) **Compositions and methods for detection of Hepatitis A virus nucleic acid**

(57) Nucleic acid oligomeric sequences and in vitro nucleic acid amplification and detection methods for detecting the presence of HAV RNA sequences in samples are disclosed. Kits comprising nucleic acid oligomers for

amplifying and detecting HAV nucleic acid sequences are disclosed.

**EP 2 402 465 A2**

**Description**FIELD OF THE INVENTION

5     **[0001]** This invention relates to diagnostic detection of a human virus, and specifically relates to assays to detect human hepatitis A virus sequences by using *in vitro* nucleic acid amplification and detection of amplified sequences.

BACKGROUND OF THE INVENTION

10    **[0002]** Hepatitis A virus (HAV) is the causative agent of one form of hepatitis that may produce symptoms that include fever, fatigue, nausea, abdominal pain, diarrhea, loss of appetite, and jaundice over less than two months. Of those infected with HAV, about 10% to 15% have a prolonged or relapsing symptoms over a six to nine months following infection. Immunity to HAV, based on the individual's production of anti-HAV immunoglobulin G (IgG), follows both symptomatic and asymptomatic infections.

15    **[0003]** Although the incidence of HAV infections has dramatically decreased in parts of the world in which vaccination for HAV (e.g., by using inactivated HAV) has been widely used since the late 1990's, epidemics of HAV infections (greater than 700 cases per 100,000 population) may occur in non-immune populations where poor sanitary conditions exist, even temporarily, e.g. following an earthquake. HAV is shed in feces of infected persons and is usually transmitted by the fecal-oral route. Community-wide outbreaks may result from foodborne transmission that occurs when an HAV-  
20    infected food handler contaminates food during preparation, or when food materials are contaminated during growing, harvesting, packing, or processing in the distribution system. Transmission may also result from contact with HAV-contaminated serum, blood products, or contaminated needles, e.g., by transfusion or injection drug use. Persons at risk of HAV infection include those who have household or sex contacts with HAV-infected persons, persons who have clotting-factor disorders (e.g., hemophilia) or chronic liver disease, persons who travel in countries where hepatitis A is  
25    common, men who have sex with men, illegal drug users, and children who live in areas with high rates of hepatitis A (e.g., > 20 cases per 100,000 population).

30    **[0004]** HAV is a 27-nm RNA virus (picornavirus) that contains a plus-sense single-stranded RNA genome of about 7.5 kb, for which a single serotype has been found worldwide. HAV replicates in the liver, is excreted in bile, and is shed in feces (up to  $10^8$  virus per ml) during the acute phase of an infection. The incubation period is usually two to six weeks before symptoms appear. Diagnosis of hepatitis A cannot be differentiated from other types of viral hepatitis by symptoms or other clinical features (e.g., elevated serum aminotransferase levels). Typically, hepatitis A diagnosis is confirmed by serological testing that provides positive results for the presence of anti-HAV immunoglobulins (Ig). Anti-HAV IgM is generally present five to ten days before the onset of symptoms and is undetectable in most patients by six months later, whereas anti-HAV IgG appears early during infection and remains detectable for the individual's lifetime. HAV RNA can  
35    be detected in the blood and stool of most persons during the acute phase of infection by using nucleic acid testing methods, e.g., amplification by the polymerase chain reaction (PCR), and nucleic acid sequencing, which has been used to identify the genetic relatedness of HAV following community-wide infections (Data et al., Morbidity Mortality Wkly. Rpt., 2003, 52(47): 1155-57; LaPorte et al., Morbidity Mortality Wkly. Rpt., 2003, 52(24): 565-67). These methods, however, are not generally used for diagnostic purposes.

40    **[0005]** In the USA, every year about 100 persons die from acute liver failure due to hepatitis A (death rate of about 0.015%). Even in nonfatal hepatitis A cases, substantial costs are associated with HAV infections, including the costs of patient hospitalization, outpatient visits, and lost work days. Public health costs associated with hepatitis A outbreaks include locating and administering immune globulin to people exposed to an infected individual or infectious source (e.g., contaminated water or food) within two weeks of exposure. Substantial psychological costs and economic losses may  
45    result from the perceived risk of infection, particularly for community-wide outbreaks. Because of the relative ease of HAV transmission in contaminated food and water, and the morbidity associated with hepatitis A, HAV is a potential agent for use in biological terrorism.

50    **[0006]** There exists a need to accurately detect the presence of HAV in biological and environmental samples. There exists a need to rapidly diagnose HAV-infected individuals. For example, because immune globulin must be administered to a person within two weeks of HAV exposure to be effective, there exists a need for a rapid and accurate assay to promptly evaluate food handlers with hepatitis symptoms and report HAV-positive sources to public health agencies. There is a need to detect HAV present in contaminated materials, such as water and food, to prevent community-wide outbreaks or epidemics resulting from use or consumption of these materials. There is also a need to detect HAV contamination in products that may be used in medical treatment, e.g., blood or serum used for transfusions or for the  
55    manufacture of factors derived from human fluids.

**[0007]** The present invention responds to these needs by disclosing oligonucleotide sequences used in nucleic acid testing methods to detect the presence of HAV nucleic acid in a sample.

SUMMARY OF THE INVENTION

**[0008]** The invention includes nucleic acid oligomers useful for purification, amplification and detection of HAV target sequences. Such oligomers or combinations of oligomers may be contained a kit configuration, embodiments of which may include additional oligomers and/or other reagents for amplifying and/or detecting a HAV sequence. The invention also includes methods of detecting HAV in a sample that use steps of purifying HAV nucleic acid from other components in the sample, amplifying a HAV RNA target sequence or cDNA made therefrom by using a nucleic acid polymerase *in vitro* and any combination of amplification oligomers as described herein to produce an amplified product, and detecting the amplified product by using a detection probe that hybridizes specifically with at least a portion of the amplified product. In one embodiment, HAV nucleic acid is purified by using at least one capture oligomer that includes a sequence that hybridizes specifically to a HAV RNA target region to form a hybridization complex that includes the HAV RNA which is separated from other sample components.

**[0009]** One aspect of the invention is a combination of at least two oligomers specific for amplifying a HAV target region that include: for a first HAV target region, oligomers of about 23 to 26 nt contained in the sequence of SEQ ID NO:138 that include at least the sequence of SEQ ID NO:139 or SEQ ID NO:140, or oligomers in a size range of about 19 to 25 nt contained in the sequence of SEQ ID NO:141 that contain at least one sequence of SEQ ID Nos. 142 to 146, or promoter primer oligomers in a size range of about 50 to 53 nt that include HAV target-specific portions of any one of SEQ ID Nos. 21 to 27; for a second HAV target region, oligomers of about 21 to 27 nt contained in the sequence of SEQ ID NO:60 or contained in the sequence of SEQ ID NO:86 that include at least the sequence of SEQ ID NO:156, or promoter primer oligomers in a size range of about 48 to 54 nt that include HAV target-specific portions of any one of SEQ ID Nos. 29 to 32; for a third HAV target region, oligomers of about 24 to 30 nt contained in the sequence of SEQ ID NO:147 that include at least the sequence of SEQ ID NO:148, or contained in the sequence of SEQ ID NO:157 that include at least the sequence of SEQ ID NO:158, or promoter primer oligomers that include HAV target-specific portions of SEQ ID NO:31 or SEQ ID NO:32; for a fourth HAV target region, oligomers of about 18 to 27 nt contained in the sequence of SEQ ID NO:93 or SEQ ID NO:95 that contain at least the sequence of SEQ ID NO:97, SEQ ID NO:159, or SEQ ID NO:160, or a promoter primer oligomer that includes a HAV target-specific portion of SEQ ID NO:33; for a fifth HAV target region, oligomers of about 19 to 31 nt contained in the sequence of SEQ ID NO: 149 that include at least the sequence of SEQ ID NO:150, or promoter primer oligomers in a size range of about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 34 to 40; for a sixth HAV target region, oligomers of about 24 to 28 nt contained in the sequence of SEQ ID NO:161 that include at least the sequence of SEQ ID NO:162, or promoter primer oligomers are embodiments of promoter primers that include HAV target-specific portions of SEQ ID NO:41 or SEQ ID NO:42; and for a seventh HAV target region, oligomers of about 20 to 30 nt contained in the sequence of SEQ ID NO: 151 that include at least any one of the sequences of SEQ ID NO:152 to SEQ ID NO:155, or contained in SEQ ID NO: 163 that include at least the sequence of SEQ ID NO:164, or contained in SEQ ID NO:165 that include at least any one of the sequences of SEQ ID Nos.166 to 168, or promoter primer oligomers in a size range of about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 43 to 49. Preferred embodiments of combinations of at least two oligomers specific for the first HAV target region are selected from SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO: 25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:143, SEQ ID NO:144, and SEQ ID NO:145. Preferred embodiments of combinations of at least two oligomers specific for the second HAV target region are selected from SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, and SEQ ID NO:156. Preferred embodiments of combinations of at least two oligomers specific for the third HAV target region are selected from SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:61, SEQ ID NO: 62, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, and SEQ ID NO:148. Preferred embodiments of combinations of at least two oligomers specific for the fourth HAV target region are selected from SEQ ID NO:33, SEQ ID NO:63, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, and SEQ ID NO:97. Preferred embodiments of combinations of at least two oligomers specific for the fifth HAV target region are selected from SEQ ID NO: 34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:64, SEQ ID NO: 65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:97, SEQ ID NO:149, and SEQ ID NO:150. Preferred combinations of at least two oligomers specific for the sixth HAV target region are selected from SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:161, and SEQ ID NO:162. Preferred combinations of at least two oligomers specific for the seventh HAV target region are selected from SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO: 78, SEQ ID NO:79, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO: 107, SEQ ID NO:108, SEQ ID NO:152, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:163, SEQ ID NO:164, SEQ ID

NO:165, SEQ ID NO:166, SEQ ID NO:167, and SEQ ID NO:168. Other preferred embodiments further include at least one capture probe oligomer selected from SEQ ID Nos. 1 to 14. Still other embodiments further include at least one detection probe oligomer selected from SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:113, SEQ ID NO:115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO:121 to SEQ ID NO:124, and SEQ ID NO:126 to SEQ ID NO:130. Preferred  
 5 embodiments of combinations of oligomers include at least two oligomers specific for amplifying a selected HAV target region and at least one detection probe oligomer that is specific for a sequence contained in HAV genomic sequence located between the selected two oligomers specific for amplifying the selected HAV target region. Preferred embodiments of such combinations of oligomers may be packaged together in a kit, which may further contain other reagents such as reagents used in purifying HAV RNA from a sample and/or reagents used in *in vitro* nucleic acid amplification,  
 10 and/or reagents used in producing a detectable signal from a detection probe oligomer.

**[0010]** Another aspect of the invention is a method of detecting the presence of HAV in a sample that includes the steps of purifying HAV nucleic acid from other components in a sample containing HAV; amplifying a HAV target sequence in the purified HAV nucleic acid, or a cDNA made therefrom, by using an *in vitro* amplification reaction that includes at least two amplification oligomers specific for a selected HAV target region, which include: for a first HAV target region,  
 15 oligomers of about 23 to 26 nt contained in the sequence of SEQ ID NO:138 that include at least the sequence of SEQ ID NO:139 or SEQ ID NO:140, or oligomers in a size range of about 19 to 25 nt contained in the sequence of SEQ ID NO:141 that contain at least one sequence of SEQ ID Nos. 142 to 146, or promoter primer oligomers in a size range of about 50 to 53 nt that include HAV target-specific portions of any one of SEQ ID Nos. 21 to 27; for a second HAV target region, oligomers of about 21 to 27 nt contained in the sequence of SEQ ID NO:60 or contained in the sequence of SEQ  
 20 ID NO:86 that include at least the sequence of SEQ ID NO:156, or promoter primer oligomers in a size range of about 48 to 54 nt that include HAV target-specific portions of any one of SEQ ID Nos. 29 to 32; for a third HAV target region, oligomers of about 24 to 30 nt contained in the sequence of SEQ ID NO:147 that include at least the sequence of SEQ ID NO:148, or contained in the sequence of SEQ ID NO:157 that include at least the sequence of SEQ ID NO:158, or promoter primer oligomers that include HAV target-specific portions of SEQ ID NO:31 or SEQ ID NO:32; for a fourth  
 25 HAV target region, oligomers of about 18 to 27 nt contained in the sequence of SEQ ID NO:93 or SEQ ID NO:95 that contain at least the sequence of SEQ ID NO:97, SEQ ID NO:159, or SEQ ID NO:160, or a promoter primer oligomer that includes a HAV target-specific portion of SEQ ID NO:33; for a fifth HAV target region, oligomers of about 19 to 31 nt contained in the sequence of SEQ ID NO:149 that include at least the sequence of SEQ ID NO:150, or promoter primer oligomers in a size range of about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID  
 30 Nos. 34 to 40; for a sixth HAV target region, oligomers of about 24 to 28 nt contained in the sequence of SEQ ID NO:161 that include at least the sequence of SEQ ID NO:162, or promoter primer oligomers are embodiments of promoter primers that include HAV target-specific portions of SEQ ID NO:41 or SEQ ID NO:42; and for a seventh HAV target region, oligomers of about 20 to 30 nt contained in the sequence of SEQ ID NO:151 that include at least any one of the sequences of SEQ ID NO:152 to SEQ ID NO:155, or contained in SEQ ID NO:163 that include at least the sequence  
 35 of SEQ ID NO:164, or contained in SEQ ID NO:165 that include at least any one of the sequences of SEQ ID Nos. 166 to 168, or promoter primer oligomers in a size range of about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 43 to 49, to produce an amplified product of the selected HAV target region; and detecting the amplified product by using a detection probe that hybridizes specifically with at least a portion of the amplified product. A preferred embodiment in the purifying step contacts the sample with at least one capture probe oligomer comprising  
 40 a sequence contained in any one of SEQ ID Nos. 1 to 14 that hybridizes specifically to a sequence in HAV RNA to form a hybridization complex with the HAV RNA, and separates the hybridization complex that contains the HAV RNA from other sample components. Preferred embodiments that amplify a sequence in the first HAV target region use at least two oligomers specific for the first HAV target region selected from SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:  
 45 26, SEQ ID NO:27, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:143, SEQ ID NO:144, and SEQ ID NO:145; and then use at least one detection probe that hybridizes specifically to the amplified product of the first HAV target region. Preferred embodiments that amplify a sequence in the second HAV target region use at least two oligomers specific for the second HAV target region selected  
 50 from SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, and SEQ ID NO:156; and then use at least one detection probe that hybridizes specifically to the amplified product of the second HAV target region. Preferred embodiments that amplify a sequence in the third HAV target region use at least two oligomers specific for the third HAV target region selected from SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:89, SEQ ID NO:90, SEQ ID  
 55 NO:91, and SEQ ID NO:148; and then use at least one detection probe that hybridizes specifically to the amplified product of the third HAV target region. Preferred embodiments that amplify a sequence in the fourth HAV target region use at least two oligomers specific for the fourth HAV target region selected from SEQ ID NO:33, SEQ ID NO:63, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, and SEQ ID NO:97; and then use at least

one detection probe that hybridizes specifically to the amplified product of the fourth HAV target region. Preferred embodiments that amplify a sequence in the fifth HAV target region use at least two oligomers specific for the fifth HAV target region selected from SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:97, SEQ ID NO:149, and SEQ ID NO:150; and then use at least one detection probe that hybridizes specifically to the amplified product of the fifth HAV target region. Preferred embodiments that amplify a sequence in the sixth HAV target region use at least two oligomers specific for the sixth HAV target region selected from SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:161, and SEQ ID NO:162; and then used at least one detection probe that hybridizes specifically to the amplified product of the sixth HAV target region. Preferred embodiments that amplify a sequence in the seventh HAV target region use at least two oligomers specific for the seventh HAV target region selected from SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:152, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:163, SEQ ID NO:164, SEQ ID NO:165, SEQ ID NO:166, SEQ ID NO:167, and SEQ ID NO:168; and then use at least one detection probe that hybridizes specifically to the amplified product of the seventh HAV target region.

## DETAILED DESCRIPTION OF THE INVENTION

**[0011]** The present invention includes methods of detecting HAV present in samples which may be biological samples derived from humans (e.g., feces, blood, serum, saliva or urine), environmental samples (e.g., water, soil) or other materials (e.g., foodstuffs) that are potentially contaminated with HAV. The methods are based on detecting the presence of HAV nucleic acid sequences by amplifying *in vitro* a region of the HAV genome and detecting the amplified nucleic acid by using a probe that binds specifically to a sequence in the amplified nucleic acid. One embodiment of the method includes a step of isolating or purifying HAV nucleic acid from a sample before the step of amplifying a region of the HAV genome. This embodiment isolates HAV genomic RNA by using a capture oligomer that binds specifically to a sequence in the HAV genome, preferably outside of the region of the HAV genome that is amplified, and separating the complex made up of the capture oligomer and the bound HAV RNA from other sample components by using a capture support, such as a particle to which the capture oligomer also binds. Amplifying a portion of the HAV genomic sequence uses one or more amplification oligomers that bind specifically to HAV RNA or a complementary sequence, and enzymatic synthesis *in vitro* to make additional copies of a portion of the HAV genomic sequence or a complementary sequence by using the amplification oligomers as primers for synthesis of the additional copies. A preferred embodiment uses an isothermal amplification reaction to make additional copies of a portion of the HAV genomic sequence. The amplified HAV sequence is then detected by specifically binding one or more probe oligomers to the amplified nucleic acid and detecting a signal that results from the probe oligomer bound to the amplified sequence. Detection of a signal resulting from the probe oligomer bound to the amplified HAV sequence indicates the presence of HAV in the sample. These methods are useful for detecting the presence of HAV in a variety of samples, such as biological samples used to diagnose a HAV infection in a human, or HAV-contaminated environmental samples to prevent the spread of HAV resulting from use or consumption of the contaminated source. These methods are also useful for testing human fluid samples for the presence of HAV, such as in serum or plasma, to prevent subsequent HAV infections resulting from use of the human fluid in transfusion or for preparation of therapeutic factors. The methods of the present invention are also useful for screening human tissue or organs for the presence of HAV to prevent their use in transplantation therapy. Thus, these methods are especially important for detecting HAV contamination in human samples or products derived from human tissue.

**[0012]** The present invention encompasses nucleic acid compositions, such as oligomers that hybridize specifically to HAV RNA or nucleic acids derived from HAV RNA, e.g., cDNA or amplified sequences made from HAV RNA. One such composition is a capture oligomer used to purify HAV RNA from a complex mixture such as a sample by hybridizing specifically to HAV RNA and attaching the hybridized HAV RNA to a capture support that permits separation of the captured HAV RNA from other sample components. The method of purification that uses such a capture oligomer is generally referred to as target capture, where HAV RNA is the specific target nucleic acid. Another oligomer of the invention is a nucleic acid amplification oligomer (sometimes referred to as a primer). Additional embodiments include probe oligomers that hybridize specifically to HAV RNA or amplified HAV nucleic acid sequences to provide a signal that detects the presence of an HAV specific sequence. These nucleic acid sequences are useful for capturing, amplifying and detecting HAV specific sequences and, thus, function together for detecting the presence of HAV in a sample.

**[0013]** A sample includes any liquid that may contain HAV or solid that may contain or have surface HAV. Samples include, for example, those from environmental sources such as water, biological sources such as human fluids or wastes, and food, packaging materials, or other components used in food processing. A biological sample includes any tissue

or material derived from a living or dead human which may contain HAV or HAV nucleic acid, including, for example, saliva, blood, plasma, serum, biopsy tissue, gastrointestinal tissue, urine, feces, or other body fluids, tissues or materials. A sample may be treated to physically or mechanically disrupt its physical state to release HAV particles or HAV RNA into an aqueous solution or solvent by using standard methods.

**[0014]** Nucleic acids include DNA or an analog thereof, RNA or an analog thereof, or mixed DNA-RNA polymers or oligomers, made up of at least two, and preferably ten or more bases linked by a backbone structure. DNA and RNA may be made up of the common bases (A, T, G and C for DNA, and A, G, C and U for RNA), although base analogs (e.g., inosine) and abasic positions (i.e., a phosphodiester backbone that lacks a nucleotide at one or more positions, see U.S. Pat. No. 5,585,481) are also included in these terms. Polymers may be many hundred or thousands of nucleotides long, whereas oligomers generally refer to nucleic acids of 1,000 or fewer linked nucleotides, and often comprise two to about 100 linked nucleotides. Oligomers generally fall in a size range having a lower limit of about 10 bases and an upper limit of about 150 bases, preferably in a size range of about 15 to about 70 bases. Oligomers may be purified from naturally occurring biological sources, but preferably are synthesized in vitro using any of a variety of well known enzymatic or chemical methods (e.g., Caruthers et al., 1987, *Methods in Enzymol.*, 154; 287).

**[0015]** A nucleic acid backbone refers to groups or linkages known in the art (Eschenmoser, 1999, *Science* 284: 2118-2124), e.g., sugar-phosphodiester linkages, 2'-O-methyl linkages, guanidine linkers in DNA ("DNG"), S-methylthiourea linkers, methylphosphonate linkages, phosphoramidate linkages, amide backbone modifications as in polyamide or peptide nucleic acids (PNA), phosphoromioate linkages, phosphonic ester nucleic acid linkages, pyranosyl oligonucleotide linkages, bicyclo- and tricyclo-nucleic acid linkages, formacetal and 3'-thioformacetal linkages, morpholino linkages, or other modifications of the natural phosphodiester internucleoside bond, or combinations thereof (Majlessi et al., 1998, *Nucl. Acids Res.* 28(9):2224-2229; Dempcy et al., 1995, *Proc. Natl. Acad. Sci. USA* 92:6497-6101; Browne et al., 1995, *Proc. Natl. Acad. Sci. USA* 92:7051-7055; Arya & Bruice, 1998, *J. Am. Chem. Soc.* 120:6619-6620; Reynolds et al., 1996, *Nucl. Acids Res.* 24(22):4584-4591; Gryaznov & Chen, 1994, *Am. Chem. Soc.* 116:3143-3144; Chaturvedi et al., 1996, *Nucl. Acids Res.* 24(12):2318-2323; Hyrup & Nielsen, 1996, *Bioorg. & Med. Chem.* 4:5-23; Hydig-Hielsen et al., PCT Pat. App. WO 95/32305; Mesmaeker et al., *Syn. Lett.*, Nov. 1997:1287-1290; Peyman et al., 1996, *Angew. Chem. Int. Ed. Engl.* 35(22):2636-2638; Aerschot et al., 1995, *Angew. Chem. Int. Ed. Engl.* 34(12):1338-1339; Koshkin et al., 1998, *J. Am. Chem. Soc.* 120:13252-13253; Steffens & Leumann, 1997, *J. Am. Chem. Soc.* 119:11548-11549; Jones et al., 1993, *J. Org. Chem.* 58:2983-2991; Summerton & Weller, 1997, *Antisense & Nucl. Acid Drug Dev.* 7: 187-195; Stirchak et al., 1989, *Nucl. Acids Res.* 17(15):6129-6141). A nucleic acid backbone may include a mixture of linkages in the same oligomer or polymer (e.g., one or more sugar-phosphodiester linkages and one or more 2'-O-methyl linkages in the strand) or may have the same linkages throughout the strand (e.g. all 2'-O-methyl or all amide modification linkages).

**[0016]** A target, target sequence, or target nucleic acid may refer to a large sequence (e.g., greater than 1000 nt) or a smaller sequence within a larger nucleic acid, to which another sequence binds, e.g., by using standard complementary base pairing. A target nucleic acid may be RNA or DNA, which is naturally occurring or made synthetically. For example, a target may be a relatively large nucleic acid such as the HAV genome, or a target may be a smaller subsequence contained in HAV RNA, its complement, or an amplification product made from it, which binds specifically another sequence in an oligomer. Those skilled in the art will appreciate that a target nucleic acid may exist in any form, e.g., a sense or antisense (+ or -) strand.

**[0017]** Complementary nucleic acids (or nucleic acid complementarity) refers to a base sequence in one strand of nucleic acid that, due to orientation of its functional groups, binds to a base sequence in an opposing strand, e.g., by hydrogen bonding between A and T or U bases, and between C and G bases. Substantially complementary means that a base sequence in one strand is not completely or perfectly complementary to a base sequence in an opposing strand, but that sufficient bonding occurs between bases of the two strands to form a stable hybridized complex in a set of conditions (e.g., salt concentration in an aqueous solution, or a temperature). Such conditions may be predicted by using the base sequences and standard mathematical calculations known to those skilled in the art for determining the melting temperature ( $T_m$ ) at which 50% of hybridized strands are denatured, or by empirical determination of  $T_m$  by using routine methods (e.g., see Sambrook et al., *Molecular Cloning, A Laboratory Manual*, 2nd Ed., (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989), at 9.50-51, 11.46-49, 11.55-57).

**[0018]** A hybridization condition refers to the cumulative environment in which one nucleic acid strand bonds to a second nucleic acid strand by complementary strand interactions to produce a hybridization complex. Such conditions include, e.g., temperature, chemical components and concentrations of compounds (e.g., salts, buffers, chelating agents, organic compounds) in aqueous and/or organic solutions that contain the nucleic acids. Other factors, such incubation time or reaction chamber dimensions may contribute to hybridization conditions, which are well known in the art (e.g., see Sambrook et al., *id.*, at 1.90-1.91, 9.47-9.51, 11.47-11.57).

**[0019]** A label refers to a molecular moiety that is detectable or produces a detectable response directly or indirectly, e.g., by catalyzing a reaction that produces a signal. Labels include luminescent moieties (e.g., fluorescent, bioluminescent, or chemiluminescent compounds), radioisotopes, members of binding pairs (e.g., biotin and avidin or streptavidin),

enzymes or enzyme substrates, reactive groups, or chromophores, e.g., a dye or particle that results in a detectable color. A detectable response or signal is any perceptible or measurable output that indicates the presence of a label, e.g., light, color, radioactive decay emission, electrical signal, magnetic field, or signal blockage, such as from quenching or turbidity.

**[0020]** An immobilized oligomer or probe refers to an oligomer that is connected or attached, covalently or noncovalently, to a capture support matrix, which provides a means for joining a capture hybrid containing a target nucleic acid to the capture support. A preferred immobilized probe is an oligomer that binds, directly or indirectly, to a target nucleic acid to facilitate separation of the bound target nucleic acid from unbound sample materials. In one embodiment, the target is indirectly bound to the immobilized probe via a capture probe that links the target and immobilized probe in a hybridization complex (see U.S. Pat. Nos. 6,110,678 and 6,280,952, Weisburg et al.). Any of a variety of supports may be used, such as matrices or particles made of, e.g., nitrocellulose, nylon, glass, polyacrylate, mixed polymers, polystyrene, silane polypropylene, and magnetic materials. Monodisperse magnetic particles of relatively uniform size that can be readily retrieved from solution by applying a magnetic force are a preferred embodiment of a support.

**[0021]** A capture oligomer or probe joins a target nucleic acid and an immobilized probe, i.e., by using a target-specific moiety that binds the target sequence and a moiety that attaches the capture probe to an immobilized probe. In one embodiment, both attachments result from hybridization of complementary base sequences, i.e., hybridization of a target sequence with a target-complementary sequence of the capture probe, and hybridization of another portion of the capture probe to a complementary sequence of the immobilized probe. In other embodiments, one or more attachments may occur by using members of a specific binding pair (e.g., biotin and avidin or streptavidin), which are well-known in the art. Compositions and methods that use capture probes are known (U.S. Pat. No. 6,910,878).

**[0022]** Separating or purifying refers to removing one or more components of a sample from other sample components. Sample components include nucleic acids in a generally aqueous solution phase which may also include materials such as proteins, carbohydrates, lipids, and other compounds. Preferably, separating or purifying a nucleic acid removes at least about 70%, more preferably at least about 90% and, even more preferably, at least about 95% of the nucleic acid from other sample components.

**[0023]** An amplification oligonucleotide or oligomer refers to an oligomer that hybridizes to a target nucleic acid, or its complementary sequence, and participates in a nucleic acid amplification reaction by serving as a primer for synthesis of nucleic acid *in vitro*. Amplification oligomers may contain additional functional sequences, such a promoter sequence that binds an RNA polymerase in an oligomer referred to as a promoter primer. An amplification oligonucleotide generally contains at least about 10 contiguous bases, preferably at least about 12 contiguous bases, that are complementary to a target sequence (or a complementary strand thereof). The contiguous bases are preferably at least about 80%, more preferably at least about 90%, and most preferably about 100% complementary to the sequence that binds to the amplification oligomer. An amplification oligomer may be RNA, DNA, or mixed Dhfa-RNA, and optionally may include modified nucleotides or backbone linkages.

**[0024]** A primer refers to an oligonucleotide that hybridizes to a template nucleic acid and which has an end (usually 3') that can be extended in a polymerization reaction catalyzed by an enzyme. The 5' region of the primer may be non-complementary to the target nucleic acid, e.g., as in a promoter primer that includes a 5' promoter sequence that is not present in the target sequence. Those skilled in the art will appreciate that a promoter primer may function as a primer independent of its promoter sequence (i.e., with or without the promoter sequence) and that any amplification oligomer may be modified to include a 5' promoter sequence, and thus function as a promoter primer.

**[0025]** Amplification refers to any known procedure for obtaining multiple copies of a target sequence, its complement, or fragments thereof. Amplification of fragments refers to production of an amplified nucleic acid that contains less than the complete target nucleic acid sequence or its complement, e.g., amplification of a portion of the complete HAV genome. Amplification of a fragment or portion of the complete target may result from using an amplification oligomer that which hybridizes to, and initiates polymerization from an internal position of the target nucleic acid. Known amplification methods include, e.g., transcription-mediated amplification (TMA), replicase-mediated amplification, the polymerase chain reaction (PCR), ligase chain reaction (LCR) and strand-displacement amplification (SDA). Replicase-mediated amplification uses self replicating RNA molecules, and a replicase such as QB-replicase (e.g., U.S. Pat. No. 4,786,600 Kramer et al.). PCR uses a DNA polymerase, multiple primers and thermal cycling to synthesize many copies of two complementary strands of DNA or cDNA (e.g., U.S. Pat. Nos. 4,683,195, 4,683,242, and 4,800,159, Mullis et al.). LCR uses at least four separate oligomers to amplify a target and its complementary strand by using multiple cycles of hybridization, ligation, and denaturation (e.g., U.S. Pat. Nos. 5,427,930 Biekenmeyer et al., and 5,494,810 Barany et al.), SDA uses a primer that contains a recognition site for a restriction endonuclease and the endonuclease nicks one strand of a hemimodified DNA duplex that includes the target sequence, followed by a series of primer extension and strand displacement steps (e.g., U.S. Pat. No. 5,422,252 Walker et al.) Transcription-mediated or transcription-associated amplification reactions use a polymerase to make a complementary strand to the target in a double-stranded form that contains a functional promoter for a specific RNA polymerase that makes transcripts that can cycle isothermally to produce additional copies of transcripts that are detectable amplification products.

**[0026]** Transcription-mediated or transcription-associated amplification uses an RNA polymerase to produce multiple RNA transcripts from a nucleic acid template in isothermal reactions that use an RNA polymerase, a DNA polymerase, deoxyribonucleoside triphosphates, ribonucleoside triphosphates, and a promoter-primer, and optionally may include one or more additional oligonucleotides. These methods of amplification and reaction conditions have been described in detail previously (e.g., see U.S. Pat. Nos. 5,399,491 and 5,554,516 Kacian et al., U.S. Pat. No. 5,437,990 Burg et al., PCT Nos. WO 88/09302 and WO 88/10315 Gingeras et al., U.S. Pat. No. 5,130,238 Malek et al., U.S. Pat. Nos. 4,868,105 and 5,124,246 Urdea et al.).

**[0027]** Preferred embodiments of the present invention use transcription-mediated amplification (TMA, described in U.S. Pat. Nos. 5,399,491 and 5,554,596). It will, however, be apparent to one skilled in the art that the methods and oligonucleotide primer sequences described herein are readily applicable to use with any nucleic acid amplification method that synthetically extends primers by using a polymerase.

**[0028]** A detection probe is a oligomer that binds to a specific target sequence and, by binding, produces, directly or indirectly, a detectable signal that indicates the presence of the target sequence. A detection probe need not be labeled to produce a detectable signal, such as an electrical impulse that results from the probe binding to the target. A labeled probe is made up of an oligomer that is linked, directly or indirectly, to a label. Methods of making and/or using labeled probes are well known (e.g., Sambrook et al., *id.*, Chapt. 10; U.S. Pat. Nos. 6,361,945 Becker et al., 5,658,737 Nelson et al., 5,656,207 Woodhead et al., 5,547,842 Hogan et al., 5,283,174 Arnold et al., 4,581,333 Kourilsky et al., and 5,731,148 Becker et al.). Detection probes may include a synthetic linker (U.S. Pat. Nos. 5,585,481 and 5,639,604 Arnold et al.), and a chemiluminescent label, such as an acridinium ester (AE) compound (U.S. Pat. Nos. 5,185,439, 5,656,207, and 5,658,737).

**[0029]** A homogeneous detectable label is a label that can be detected in a homogeneous manner depending on whether the label is bound or unbound to a target. That is, detection of a label in a homogeneous reaction does not require physical separation of unbound forms of the label from the mixture in which the signal is detected. It will be appreciated by skilled artisans that a homogeneous reaction may occur in solution or on a support, e.g. on an array, biochip or gene chip. Homogeneous detectable labels and conditions for their detection are well known (e.g., U.S. Pat. Nos. 5,283,174, 5,656,207, and 5,658,737).

**[0030]** By "consisting essentially of" is meant that additional component(s), composition(s) or method step(s) that do not materially change the basic and novel characteristics of the present invention may be included in the compositions, kits, or methods of the present invention. Such characteristics include the ability to detect specifically the presence of HAV nucleic acid in a sample with a sensitivity of at least 80% for samples containing 25 to 30 copies of HAV per ml by using a combination of capture probe, amplification primers, and detection probe oligomers as described herein. Any component(s), composition(s), or method step(s) that have a material effect on the specificity and/or sensitivity of detection of HAV present in a sample by using the nucleic acid oligomers and *in vitro* methods described herein, would fall outside of this term.

**[0031]** Unless defined otherwise, all scientific and technical terms used herein have the same meaning as commonly understood by those skilled in the relevant art. Definitions of many of the terms used herein are provided in, e.g., Dictionary of Microbiology and Molecular Biology, 2nd ed. (Singleton et al., 1994, John Wiley & Sons, New York, NY), The Encyclopedia of Molecular Biology (Kendrew, Ed., 1994, Blackwell Science Ltd., Cambridge, MA), or The Harper Collins Dictionary of Biology (Hale & Marham, 1991, Harper Perennial, New York, NY). Unless mentioned otherwise, the techniques employed or contemplated herein are standard methodologies well known to one of ordinary skill in the art. Examples are included to illustrate preferred embodiments.

**[0032]** The present invention includes compositions (nucleic acid amplification oligomers, detection probes, and optionally capture oligomers) and methods for detecting HAV nucleic acid in a sample. To select sequences appropriate for use as the oligomers disclosed herein, known HAV genomic sequences (Beneduce, et al., 1995, Virus Res. 36 (2-3): 299-309, Fujiwara, et al., 2001, J. Hepatol. 35 (1): 112-119, Hu, et al., 2002, Acta Virol. 46 (3): 153-157), including those of different isolates, partial sequences, and complementary sequences available on a public database (e.g., GenBank accession nos. AB020564 to AB020569) were aligned by matching regions of the same or similar sequences and the aligned sequences were compared using well known techniques. Although sequence comparisons may be facilitated by use of algorithms, those skilled in the art can readily perform such comparisons manually and visually. Portions of HAV sequences that contain relatively few sequence variants between the compared sequences were chosen as a basis for designing synthetic oligomers suitable for use in the capture, amplification and detection steps described herein. Other well known sequence characteristics, such as the GC content and the relative abundance of predicted secondary structures (e.g., hairpin turns or intramolecular pairing), were also considered in selecting the oligomer sequences.

**[0033]** Based on these analyses, regions of the HAV genome around nucleotides 200, 3700, 4700, 5700, 5800, 6000, and 7000 were chosen as potential target regions for detection of amplified HAV sequences. For each region, oligomers were designed for use in capturing the HAV RNA from a sample to purify it from other sample components, as amplification oligomers, and as probe sequences. Preferred embodiments of target regions are in portions of 0 to 305 nt, 4714 to 4765 nt, 5495 to 5788 nt, 5788 to 6069 nt, and 6952 to 7413 nt of the HAV genome.



**[0034]** Capture oligomer sequences generally include a sequence that binds specifically to a sequence near the target region to be amplified and a "tail" region used in attaching the hybridization complex that includes the target to a solid support, for example via hybridization to an immobilized oligomer (e.g., U.S. Pat. No. 6,110,678). Preferred capture oligomers include a target-specific sequence that binds specifically to a HAV RNA sequence and a covalently attached tail sequence (e.g., dT<sub>3</sub>dA<sub>30</sub>), as shown in SEQ ID Nos. 1 to 7. Those skilled in the art will understand that the target-specific portion of a capture oligomer (SEQ ID Nos. 8 to 14) or its RNA equivalent may be linked to any moiety that allows it to bind to an immobilized probe (e.g., a different tail sequence or a member of a binding pair, such as biotin or avidin). Any backbone may link the base sequence of a capture oligomer. Some embodiments use 2'-O-methyl linkages in the target-specific portion of a capture oligomer and standard DNA linkages in the tail portion. A polynucleotide tail sequence may be any sequence complementary to a sequence of an immobilized probe, and generally has a sequence length of about 5 to 50 residues, and is preferably a substantially homopolymeric sequence in a range of about 10 to about 40 residues (e.g., C<sub>10</sub> to C<sub>40</sub>) that is complementary to an immobilized homopolymeric sequence (e.g., G<sub>15</sub>).

**[0035]** Primer sequences bind specifically to an HAV RNA target sequence or a complementary strand and flank a target sequence that is amplified, although primer sequences may contain additional sequences that do not bind to the target or its complementary sequence. A primer may be a promoter primer and include a 5' promoter sequence, such as a T7 RNA polymerase promoter (SEQ ID NO:19). Embodiments of promoter primers include those of SEQ ID Nos. 20 to 49. Other embodiments of HAV-specific primers may include ancillary sequences, such as restriction endonuclease recognition sequences (SEQ ID Nos. 132 to 135). Those skilled in the art will appreciate that a target-specific sequence of a primer, with or without an attached promoter or ancillary sequence, may serve as a primer in a variety of *in vitro* amplification conditions. Amplification oligomers were designed for sequences in targeted regions of the HAV genome (e.g., around nucleotide positions 200, 3700, 4700, 5700, 5800, 6000, and 7000). Those skilled in the art will appreciate that these numbers refer to HAV target regions that are approximate only and that oligomers may function in an assay for more than one target region. That is, the oligomers are not restricted functionally by the identifying target region numbers which are provided as a shorthand reference for grouping preferred embodiments of the invention. Amplification oligomers may be synthesized as DNA, RNA, complementary DNA or RNA sequences, or mixed DNA and RNA sequences, and may include one or more non-standard nucleic acid backbone linkages. For example, an oligomer of SEQ ID NO:106 was synthesized with RNA bases and 2'-O-methyl linkages at residues 1 to 4 and standard DNA bases and linkages at the other residues.

**[0036]** For a first HAV target region (around position 200), amplification oligomers include those in a size range of about 23 to 26 nt that are contained in the sequence of SEQ ID NO:138, and include at least the sequence of SEQ ID NO:139 or SEQ ID NO:140. Embodiments of such oligomers includes those of SEQ ID NO:51 to SEQ ID NO:57. Embodiments of promoter primers for this region, in a size range of about 50 to 53 nt, are those that include target-specific portions of SEQ ID Nos. 21 to 27. Amplification oligomers for this target region also include those in a size range of about 19 to 25 nt contained in SEQ ID NO:141, and contain at least the sequence of any one of SEQ ID Nos. 142 to 146. Embodiments of amplification oligomers for this target region include those of SEQ ID Nos. 15 to 18, 20 to 27, 50 to 57, and 80 to 85.

**[0037]** For a second HAV target region (around position 3700), amplification oligomers include those in a size range of about 21 to 27 nt, contained in the sequence of SEQ ID NO:60, or in SEQ ID NO:86 and include at least SEQ ID NO: 156. Embodiments of promoter primers that include such target-specific portions for this region and are in a size range of about 48 to 54 nt include those of SEQ ID Nos. 29 to 32. Embodiments of amplification oligomers for this target region include those of SEQ ID Nos. 28 to 30, 58 to 60, and 86 to 88.

**[0038]** For a third HAV target region (around position 4700), amplification oligomers include those in a size range of about 24 to 30 nt that are contained in SEQ ID NO:147 and include at least the sequence of SEQ ID NO:148, or are contained in SEQ ID NO:157 and include at least the sequence of SEQ ID NO:158. Embodiments of amplification oligomers for this target region include those of SEQ ID Nos. 31, 32, 61, 62, 89, 90, and 91, of which SEQ ID NO:31 and SEQ ID NO:32 are promoter-primer embodiments which include a 5' promoter sequence attached to the target-specific sequence.

**[0039]** For a fourth HAV target region (around position 5700), amplification oligomers include those in a size range of about 18 to 27 nt that are contained in the sequence of SEQ ID NO:93 or SEQ ID NO:95. Embodiments of such oligomers include those that contain at least any one of SEQ ID NO:97, SEQ ID NO:159, and SEQ ID NO:160. Embodiments of amplification oligomers for this target region include those of SEQ ID Nos. 33, 63, and 92 to 97, of which SEQ ID NO: 33 is a promoter primer embodiment which includes a 5' promoter sequence attached to the target-specific sequence.

**[0040]** For a fifth HAV target region (around position 5800), amplification oligomers include those in a size range of about 19 to 31 nt that are contained in SEQ ID NO:149 and include at least the sequence of SEQ ID NO:150. Embodiments of promoter primers, in a size range of about 51 to 56 nt, that include such target-specific portions are those of SEQ ID Nos. 34 to 40. Other embodiments of amplification oligomers for this target region include those of SEQ ID Nos. 64 to 70, and 97.

**[0041]** For a sixth HAV target region (around position 6000), amplification oligomers include those of about 24 to 28

nt contained in the sequence of SEQ ID NO:161 and include the sequence of SEQ ID NO:162. Embodiments of amplification oligomers for this target region include those of SEQ ID Nos. 41, 42, 71, 72, 98, 99, and 101, of which SEQ ID Nos. 41 and 42 are embodiments of promoter primers which include a 5' promoter sequence attached to the target-specific sequence.

**[0042]** For a seventh HAV target region (around position 7000), amplification oligomers include those in a size range of about 20 to 30 nt contained in SEQ ID NO: 951 and that include at least any one of the sequences of SEQ ID NO: 152 to SEQ ID NO:155. Other embodiments of amplification oligomers for this target region are contained in SEQ ID NO:163 and include at least the sequence of SEQ ID NO:164. Additional embodiments are amplification oligomers that are contained in SEQ ID NO:965 and include at least any one of the sequences of SEQ ID Nos. 166 to 168. Embodiments of promoter primers, in a size range of about 51 to 56 nt, that include HAV target-specific portions for this region are SEQ ID Nos. 43 to 49. Other embodiments of amplification oligomers for this region include those of SEQ ID Nos. 73 to 79, and 102 to 108.

**[0043]** Oligomers were designed to hybridize to and detect amplified HAV sequences, which include the detection probes of SEQ ID Nos.109, 111, 113, 116, 117, 119, 121, 122, 123, 124, and 926 to 130. Those skilled in the art will appreciate that a detection probe will be chosen to hybridize to a sequence contained within an amplified sequence that is determined by the combination of amplification oligomers that are used. Detection probe oligomers may be synthesized as DNA, RNA, or mixed DNA and RNA polymers, and may include alternative backbone linkages such as 2'-O-methyl linkages. For example, oligomers of SEQ ID Nos.109, 111, 117, 119, 121,122, 928, and 130 were synthesized with 2'-O-methyl linkages, and oligomers of SEQ ID Nos. 124 and 127 were synthesized as mixed DNA and RNA nucleotides with 2'-O-methyl linkages from the second residue to the 3' terminal residue. Preferred embodiments of detection probes have an attached chemiluminescent label, preferably an acridinium ester (AE) compound (U.S. Pat. Nos. 5,185,439, 5,639,604, 5,585,481, and 5,656,744), which in preferred embodiments are attached to the probe by a non-nucleofide linker (see U.S. Pat. Nos. 5,585,481,5,656,744, and 5,639,604, particularly at column 10, line 6 to column 11, line 3, and Example 8). Embodiments of the probe oligomers were labeled by using known methods with an AE compound between residues 9 and 10 for SEQ ID Nos. 119, 121 and 124, between residues 10 and 11 for SEQ ID Nos. 115, 117, 126, 127 and 128, between residues 11 and 12 for SEQ ID Nos.109, 111, 123, 124 and 130, between residues 12 and 13 for SEQ ID Nos. 113, 122 and 129, and between residues 13 and 14 for SEQ ID NO:122. Probe oligomers were tested and characterized by hybridization with complementary oligomer sequences, by using standard methods for determining T<sub>m</sub> and/or differential hydrolysis of the acridinium ester in a hybridization complex (described in detail in U.S. Pat. No. 5,283,174). For example, hybridizations were performed by using complementary pairs of sequences of SEQ ID NO:109 and SEQ ID NO:110, SEQ ID NO:111 and SEQ ID NO:112, SEQ ID NO:113 and SEQ ID NO:114, SEQ ID NO:115 and SEQ ID NO:116, SEQ ID NO:117 and SEQ ID NO:118, SEQ ID NO:119 and SEQ ID NO:120, SEQ ID NO:124 and SEQ ID NO:125, SEQ ID NO:128 and SEQ ID NO:100, and SEQ ID NO:130 and SEQ ID NO:131. Hybridization testing may be performed by using other complementary sequences, such as SEQ ID NO:124 with SEQ ID NO: 137, and SEQ ID NO:129 with SEQ ID NO:136,

**[0044]** Compositions of the present invention include kits for detecting HAV nucleic acid sequences. Such kits include amplification oligomers as disclosed herein that function as primers for amplifying HAV nucleic acid sequences *in vitro*. Exemplary kits include a first amplification oligomer that hybridizes specifically to a sequence in a target region of the HAV RNA genome or its complementary sequence, and a second amplification oligomer that hybridizes specifically to another HAV sequence in the target region, preferably one complementary to the HAV RNA genomic sequence. Embodiments of kits include amplification oligomers that are combinations of primers and promoter primers as described herein. Kits may also contain one or more oligomers that serve as detection probes for detecting amplified HAV sequences of the target region of the primers selected for the kit, Embodiments kits that include of probe oligomers use one or more of the detection probe sequences as described herein, which may include a label attached directly or indirectly to the probe oligomer. Kits may also contain oligomers that serve as capture oligomers for purifying a target HAV RNA from a sample. Embodiments of such capture oligomers as described herein may contain a covalently attached tail sequence or other binding moiety used in target capture. Kits useful for practicing the methods described herein are also included in the invention, and preferred embodiments include at least two amplification oligomers as described herein, and may also include reagents for performing *in vitro* amplification, e.g., enzymes, shaft solutions, and nucleic acid synthesis substrate compounds. Oligomers described herein may be packaged in a variety of different embodiments, and thus, those skilled in the art will appreciate that the invention embraces many different kit configurations. For example, a kit may include amplification oligomers for only one target region of the HAV genome, or it may include amplification oligomers for multiple HAV target regions. Those skilled in the art will appreciate that a kit that includes a detection probe will include a probe that binds to a sequence amplified by the amplification oligomers of the kit. That is, selection of amplification oligomers and detection probe oligomers for a kit will be linked by their intended target region.

**[0045]** An embodiment of the assay to detect HAV nucleic acid in a sample includes the steps of capturing the HAV target nucleic acid from a sample by using a capture oligomer, amplifying a region of the captured HAV nucleic acid by using a combination of at least two primers, and detecting the amplified HAV sequence by specifically hybridizing it with

a detection probe oligomer and detecting a signal that results from the probe bound to the amplified HAV sequence. Preferred embodiments use a transcription-associated or transcription-mediated amplification reaction. Either the amplified nucleic acid or the probe may be labeled, or both may be unlabeled and a detectable signal results from an indirect label or response associated with the hybridization complex, such as an electrical impulse resulting from hybridization of the probe and the amplified nucleic acid.

**[0046]** The capturing step preferably uses a capture oligomer that includes a target-specific sequence (e.g., SEQ ID Nos. 8 to 14) that specifically hybridizes to a HAV target sequence and a moiety that permits the hybridized target nucleic acid to be separated from other sample components. The capturing step may use a capture oligomer that also includes a tail portion, e.g. as in SEQ ID Nos. 1 to 7, that serve as the moiety that allows the target nucleic acid to be separated from other sample components by hybridization of the tail portion to an immobilized probe, as previously described (U.S. Pat. No.6,110,678). Preferred embodiments use supports that are magnetic spheres that are monodisperse (i.e., uniform in size  $\pm$  about 5%) with covalently attached or immobilized poly-dT oligomers that hybridize to a complementary tail sequence of the capture oligomer. The hybridization complex that includes at least the target nucleic acid and the capture oligomer, and preferably also includes the immobilized probe, is separated from the other sample components by using standard physical separation methods (e.g., application of magnetic force, filtration, or centrifugation) and the captured target nucleic acid may be washed one or more times to further purify the target nucleic acid from other sample components. For example, particles with the attached target nucleic acid in a hybridization complex are suspended one or more times in a washing solution that maintains the complex and then the particles with the attached complex are retrieved from the washing solution as described above.

**[0047]** Amplifying the captured HAV target sequence uses an *in vitro* amplification reaction that uses at least two primers that flank the sequence to be amplified, e.g., an HAV sequence flanked by SEQ ID NO:66 and SEQ ID NO:95, or their complementary sequences. One embodiment uses a transcription-associated amplification, reaction that makes many RNA copies of a sequence in substantially isothermal conditions (as described previously in U.S. Pat. Nos. 5,399,491 and 5,554,516). Transcription-associated amplification uses two types of primers (one a promoter primer that contains a promoter sequence for an RNA polymerase), enzymes (a reverse transcriptase and an RNA polymerase), substrates (deoxyribonucleoside triphosphates, ribonucleoside triphosphates) and appropriate salts and buffers in solution to produce multiple RNA transcripts from a nucleic acid template. Briefly, a promoter primer hybridizes specifically to a target RNA sequence and reverse transcriptase creates a first strand cDNA by extension from the 3' end of the promoter primer and degrades the template strand in the resulting DNA:RNA duplex by using RNase H activity. A second primer binds to the cDNA and another DNA strand is synthesized by the reverse transcriptase from the end of the second primer, to create a double-stranded DNA with a functional promoter sequence to which the RNA polymerase binds. Multiple RNA transcripts ("amplicons") are transcribed and each can be a template in a new round of replication as described above, thus generating large amounts of single-stranded amplified sequence (e.g., about 100 to 3,000 transcripts from a single template). Embodiments of the invention that use a transcription-associated amplification reaction may use promoter primers (SEQ ID Nos. 20 to 49) with other primers (SEQ ID Nos. 15 to 18, 80 to 99, and 101 to 108) to amplify selected HAV sequences for detection.

**[0048]** The detecting step uses at least one probe that binds specifically to the amplified HAV sequences. Embodiments may use any known detection method (e.g., detection of a radioactive, fluorescent, enzymatic, colorimetric, electrical, or luminescent signal) to detect binding of the detection probe to the amplified HAV sequences, and the detected signal indicates the presence of HAV in the sample. Embodiments of probe oligomers (SEQ ID NOs. 109, 111, 113, 115, 117, 119, 121, 122 to 124, 126 to 130) may be unlabeled or labeled using any of a variety of known labels. In preferred embodiments, the detection step is performed in a homogeneous detection reaction without removing the unbound detection probe from the mixture. Embodiments of the probe oligomers for use in homogeneous detection reactions are preferably labeled with one of a variety of AE compounds which produce a chemiluminescent signal that is detected as described in detail previously (U.S. Pat. Nos. 5,283,174, 5,656,744, and 5,658,737).

**[0049]** A preferred assay embodiment generally includes the following steps. An HAV-containing sample is provided, which may be prepared by using standard laboratory methods to make a substantially aqueous solution or suspension that contains HAV. An aliquot (0.5 ml) of the sample solution or suspension is mixed with about an equal volume (0.4 to 0.5 ml) of a target capture reagent, i.e., a solution that contains one or more capture oligomers (4 pmol/reaction), magnetic particles with attached immobilized probes complementary to a portion of the capture oligomers, and salt compounds to provide a hybridization condition. The target capture reagent preferably includes a detergent or other chaotropic agent that disrupts the HAV particles and releases HAV RNA for hybridization with capture oligomers. The mixture is incubated 20-30 min at 60°C to allow hybridization of the target-specific portion of the capture oligomer to the HAV target sequence and then at room temperature for 20-30 min to allow binding of the capture oligomer and immobilized probe. A magnetic field is applied to the outside of the reaction container for about 10 min to separate the particles with the attached hybridization complexes that include HAV RNA, and the solution phase containing other sample components is aspirated away. To wash the particles with attached hybridization complexes, they are suspended in 1 ml of a wash buffer, separated from the solution substantially as described above, and the solution is removed. Particles with attached

hybridization complexes that include the purified HAV RNA are mixed with a solution that contains amplification reagents (buffers, salts, dXTP and XTP substrates), and a combination of amplification oligomers (a promoter primer and a primer combination, each at 3 to 30 pmol, generally 15 pmol each), and covered with oil (0.2 ml of filtered silicon oil) to prevent evaporation, and incubated for 10 min at 60°C, then for 10 min at 42°C, and then enzymes are added (reverse transcriptase and RNA polymerase), and the mixture is incubated for 60 min at 42°C. For detection, the amplification reaction mixture is incubated with at least one acridinium labeled detection probe oligomer to provide a maximum detectable signal (relative light units or RLU) of 2 million or less, as detected by using standard methods on a luminometer (e.g., Gen-Probe Leader®, Gen-Probe Incorporated, San Diego, CA). Detection probe is mixed with undiluted or a diluted aliquot of the amplification reaction mixture in a hybridization solution, incubated for 20 min at 60°C to allow hybridization of the probe oligomer to the amplified target sequence. Then, label on unbound probes is hydrolyzed by using a selection reagent (e.g., a base) and incubated for 10 min at 60°C, followed by adding a detection reagent (e.g., H<sub>2</sub>O<sub>2</sub>) to produce chemiluminescence, followed by pH neutralization (e.g., by adding acid), and detecting the chemiluminescent signal (RLU) on a luminometer (e.g., 1-5 sec).

**[0050]** For use in the methods described above, capture oligomers, amplification oligomers and detection probes may be synthesized using standard methods to produce DNA, RNA, or mixed DNA and RNA polymers. Such oligomers may include standard or modified linkages and/or naturally-occurring nucleosides (A, T or U, G, C), analogs (e.g., inosine), or synthetic purine and pyrimidine derivatives (e.g., P or K bases) (Lin & Brown, 1989, Nucl. Acids Res. 17:10373-83; Lin & Brown, 1992, Nucl. Acids Res. 20: 5149-52).

**[0051]** The general principles of the present invention may be more fully appreciated by reference to the following examples describe some embodiments of the present invention. In addition to the specific components described in the examples, generally the following reagents were used in the experiments described below.

**[0052]** Target capture reagent was made up of 790 mM HEPES, 680 mM LiOH, 10% (v/v) lithium lauryl sulfate (LLS), 230 mM succinic acid, 0.03% (v/v) anti-foaming agent, 100 µg/ml magnetic particles (1 micron SERA-MAG™ particles, Seradyn, Inc. Indianapolis, IN) with covalently attached poly-dT<sub>14</sub>, and one or more capture oligomers, each at 4 pmol per 400 µl. Wash buffer was made up of 150 mM NaCl, 10 mM HEPES, 6.5 mM NaOH, 1 mM EDTA, 0.3% (v/v) ethanol, 0.1% SDS, 0.02% (w/v) methyl paraben, 0.01% (w/v) propyl paraben, at pH 7.5. Amplification reagent was made up of 11.6 mM Tris base, 15 mM Tris-HCl, 22.7 mM MgCl<sub>2</sub>, 23.3 mM KCl, 3.33% glycerol, 0.05 mM Zn-acetate, 0.665 mM dATP, 0.665 mM dCTP, 0.665 mM dGTP, 0.665 mM dTTP, 5.32 mM ATP, 5.32 mM CTP, 5.32 mM GTP, and 5.32 mM UTP, at pH 7. Enzyme reagent was made up of 140 U/µl T7 RNA polymerase, 224 RTU/µl of Moloney Murine Leukemia Virus reverse transcriptase (MMLV-RT), 16 mM HEPES, 70 mM N-acetyl-L-cysteine, 3 mM EDTA, 0.05% (w/v) Na-azide, 20 mM Tris, 50 mM KCl, 20% (v/v) glycerol, 10% (v/v) TRITON® X-102, 150 mM trehalose, at pH 7. (Enzyme units typically are 1 U of T7 RNA polymerase incorporates 1 nmol of ATP into RNA in 1 hr at 37 ° C using a DNA template containing a T7 promoter, and 1 U of MMLV-RT incorporates 1 nmol of dTTP in 10 min at 37 ° C using 200-400 µM oligo-dT primer and poly-A template.) Probe reagent was made up of 100 mM succinic acid, 2% (w/v) LLS, 230 mM LiOH, 15 mM Aldrithiol-2, 1.2 M LiCl, 20 mM EDTA, 20 mM EGTA, 3% (v/v) ethanol, adjusted to pH 4.7 with LiOH. Selection reagent was made up of 600 mM boric acid, 182 mM NaOH, 1% (v/v) octoxynol (TRITON® X-100), at pH 8.5. Detection reagents were Detect Reagent I, which contained 1 mM nitric acid and 32 mM H<sub>2</sub>O<sub>2</sub>, and Detect Reagent II (to neutralize pH), which was 1.5 M NaOH (see U.S. Pat. No. 5,283,174 for details).

#### Example 1: Detection Probe Characterization

**[0053]** Oligomers of SEQ ID Nos. 109, 111, 113, 119, 123, 126, and 130 were synthesized using standard phosphoramidite chemistry (Caruthers et al., 1987, Methods in Enzymol., 154: 287) and an acridinium ester (AE) label was attached via a linker by using well known methods (U.S. Pat. Nos. 5,185,439 and 5,283,174), and probes were purified by using routine chromatographic methods (e.g., HPLC). Probes were AE labeled between residues 11 and 12 for SEQ ID NO:109, SEQ ID NO:111, SEQ ID NO:123, and SEQ ID NO:130, between residues 12 and 13 for SEQ ID NO:113, between residues 10 and 11 for SEQ ID NO:126, and between residues 9 and 10 for SEQ ID NO:119. To characterize the probe oligomers, each was hybridized with a complementary DNA and/or RNA oligomer (e.g., SEQ ID NO:109 with SEQ ID NO:110, SEQ ID NO:111 with SEQ ID NO:112, SEQ ID NO:113 with SEQ ID NO:114, SEQ ID NO:119 with SEQ ID NO:120, and SEQ ID NO:130 with SEQ ID NO:131), at temperatures below the predicted T<sub>m</sub> of the probe, and then the T<sub>m</sub> was experimentally determined by using standard methods. The differential hydrolysis of the AE label in probes hybridized to a complementary oligomer compared to AE in unbound probe was also experimentally determined by using standard methods (see U.S. Pat. No. 5,283,174). Briefly, the ratio of the time required for half of the signal to be lost due to AE hydrolysis in the hybrid compared to the time required for hydrolysis of half of the label in unbound probe was determined. The T<sub>m</sub>'s were in the range of 59° C to 66°C for oligomers of SEQ ID NOs.109,111,113,119 and 130 when hybridized to a complementary DNA, and T<sub>m</sub>'s were in the range of 76°C to 81 °C for oligomers of SEQ ID Nos. 109, 111, 123, 126 and 130 when hybridized to a complementary RNA. The differential hydrolysis ratios were in the range of 12 to 25 for probes of SEQ ID NOs.109, 111, 113, 119 and 130 when hybridized to complementary DNA,

and the differential hydrolysis ratios were in the range of 18 to 104 for probes of SEQ ID NOs.109,119,123, 126 and 130 when hybridized to complementary RNA. Separately, similar hybridization and differential hydrolysis tests were performed for probes of SEQ ID NO:121 labeled between residues 9 and 10, SEQ ID NO:122 labeled between residues 13 and 14, SEQ ID NO:124 labeled between residues 9 and 10, and SEQ ID NO:130 labeled between residues 11 and 12, and the differential hydrolysis ratios were in the range of 43 to 190 when the probes were hybridized to complementary RNA. These results showed that all of these synthetic probe oligomers hybridized specifically to their complementary target sequences and produced detectable signals useful for specifically detecting amplified HAV sequences.

### Example 2: Purification of HAV RNA from Samples

**[0054]** Capture oligomers of SEQ ID Nos. 1 to 7, synthesized by using standard phosphoramidite chemistry and purified using standard methods, were tested for their ability to capture HAV RNA released from virus in human plasma samples. Samples were made by adding HAV particles at known concentrations to normal human plasma (0.5 ml) and the samples containing HAV (e.g., 500 to 1000 per reaction) were mixed with an equal volume of the target capture reagent containing each capture oligomer individually (4 pmol/reaction) and polydT-magnetic particles. The mixtures were incubated for 30 min at 60°C, and then for 30 min at room temperature to form hybridization complexes that capture HAV RNA to the particles. The magnetic particles with attached captured HAV RNA were separated by applying a magnetic field for 10 min to the outside of the container, then the solution phase was aspirated away to remove other sample components, and the particles with attached hybridization complexes were washed twice sequentially, each using 1 ml of the wash buffer at room temperature and aspirating the washing solution away from the particles. Particles with attached hybridization complexes were then suspended in probe reagent (0.1 ml) containing a labeled detection probe, as described in Example 1, and incubated for 20 min at 60°C, followed by addition of selection reagent (0.2 ml), mixing and incubation for 10 min at 60°C. Production and detection of the chemiluminescent signal was performed by adding 200 µl of a detect reagent 1, incubation, and pH neutralization of the mixture by adding 200 µl of detect reagent II, and measuring RLU by using a luminometer, substantially as described above. For all of the capture oligomers tested, the presence of the HAV RNA in the sample was detected by detecting a positive signal significantly higher than background (RLU for a similar sample that contained no HAV). The assays showed little significant performance differences between the capture oligomers.

### Example 3: Amplification and Detection of HAV Sequences

**[0055]** HAV samples in normal human plasma were prepared substantially as described in Example 2 and the HAV RNA was captured by using various combinations of capture oligomers for assays to amplify and detect selected target regions of the HAV genome. For a target region of 0-305 residues of the genome, SEQ ID Nos. 2, 3, and 4 were used in the capture step. For a target region of 4714-4765 residues of the genome, SEQ ID NOS. 4, 5, 6, and 7 were used in the capture step. For a target region of 5495-5788 residues of the genome, SEQ ID NOS. 1 and 6 were used in the capture step. For a target region of 5788-6069 residues of the genome, SEQ ID NO:2 was used in the capture step. For a target region of 6952-7413 residues of the genome, SEQ ID NOS. 1,4, 5 and 7 were used in the capture step. The capture steps were performed substantially as described in Example 2.

**[0056]** The captured HAV RNA was amplified in reactions substantially as described above that contained different combinations of amplification oligomers to serve as primers for different target regions in the HAV genome. The primers used to amplify the target regions were as follows: SEQ ID NO:16 and SEQ ID NO:22 for the 0-305 residues region, SEQ ID NO:89 and SEQ ID NO:32 for the 4714-4765 residues region, SEQ ID NO:92 and SEQ ID NO:33 for the 5495-5788 residues region, SEQ ID NO:94 and SEQ ID NO:37 for the 5788-6069 residues region, and SEQ ID NO:108 and SEQ ID NO:46 for the 6952-7413 residues region. The amplification reactions were all performed substantially the same as described above. That is, particles with the attached HAV RNA from the target capture step were mixed with amplification reagent and the individual combination of amplification oligomers described above (generally 15 pmol each), and covered with silicon oil (0.2 ml) to prevent evaporation, and incubated for 10 min at 60°C and then for 10 min at 42°C. The enzyme reagent was added (reverse transcriptase and RNA polymerase), and the amplification reactions were incubated for 60 min at 42°C.

**[0057]** For detection, the amplification mixture was incubated with a labeled detection probe oligomer that hybridizes specifically to sequences contained in the amplified region. These included SEQ ID NO:109 or SEQ ID NO:111 for the 0-305 residues region, SEQ ID NO:115 for the 4714-4765 residues region, SEQ ID NO:117 for the 5495-5788 residues region, SEQ ID NO:121 and/or SEQ ID NO:122 for the 5788-6069 residues region, and SEQ ID NO:129 or SEQ ID NO:130 for the 6952-7413 residues region. The probes were provided in the probe reagent in an amount previously determined based on the specific activity of the labeled probe to produce a maximum detectable signal of 2 million RLU or less from the hybridized labeled probe. The probes and amplified sequences were incubated in the probe reagent at 55-60°C, and the chemiluminescent signal was produced from hybridized probes and detected substantially as describe in Ex-

amples 1 and 2. For all of the primer combinations tested with the captured HAV RNA, the sensitivity of the amplification assay detected between 400 and 1000 copies of HAV RNA present in the samples.

**[0058]** The combinations of capture oligomers, amplification oligomers and detection probes used for in these tests are summarized in Table 1.

**Table 1: Combinations of Oligomers for Testing for HAV in Samples**

Target Region (Residues)	Capture Oligomer(s)	Amplification Oligomers	Detection Probe(s)
0-305	SEQ ID Nos. 2, 3 and 4	SEQ ID Nos.16 and 22	SEQ ID No. 109 or 111
4714-4765	SEQ ID Nos. 4, 5, 6 and 7	SEQ ID Nos. 32 and 89	SEQ ID No.115
5495-5788	SEQ ID Nos. 1 and 6	SEQ ID Nos. 33 and 92	SEQ ID No.117
5788-6069	SEQ ID No. 2	SEQ ID Nos. 37 and 94	SEQ ID Nos. 121 and/or 122
6952-7413	SEQ ID Nos. 1, 4, 5 and 7	SEQ ID Nos. 46 and 108	SEQ ID No. 129 or 130

**[0059]** Similar experiments were performed using the different capture oligomers (SEQ ID Nos. 1, 2, 3, 4, 5, 6, and 7) separately with HAV-containing samples, in which the target capture step was performed substantially as described above on nine replicates for each assay condition. For all of these tests the target region was residues 5788 to 6069 of HAV, for which the same amplification oligomers of SEQ ID NO:36 and SEQ ID NO:96 were used in the amplification reactions with the captured HAV RNA performed as described above, and amplified products were detected by measuring chemiluminescence from hybridized detection probe (SEQ ID NO:123 or 124, labeled with AE between residues 11 and 12) as described above. The results of these assays are shown in Table 2 (average RLU for nine replicates).

**Table 2: Amplification and Detection of the 5788-6069 residue Target Region**

Purification by Capture Oligomer	Detected Signal (mean RLU)
SEQ ID NO:1	292,136
SEQ ID NO:2	275,732
SEQ ID NO:3	478,463
SEQ ID NO:4	522,837
SEQ ID NO:5	443,830
SEQ ID NO:6	416,905
SEQ ID NO:7	369,337

**[0060]** These results show that all of the capture oligomers sufficiently purified HAV RNA from samples to be amplified and detected to indicate the presence of HAV in the samples.

#### Example 4: Detection of HAV in Plasma Samples

**[0061]** This example uses an assay that detected HAV nucleic acid in HAV-positive plasma samples. To prepare samples, a commercially available stock of HAV in human plasma was diluted into HAV-negative plasma to obtain samples with 25, 30, 100, 300 and 500 HAV copies/ml; a negative control was plasma with no HAV. For each assay, performed on 20 replicate samples, 0.5 ml samples were mixed with 0.4 ml of target capture reagent containing capture oligomers of SEQ ID NO:4 (6.5 pmol/reaction) and SEQ ID NO:5 (1.3 pmol/reaction) and the target capture step was performed substantially as described in Example 3, except that a 20 min 60 °C incubation was used. For each assay, washed magnetic particles with the attached hybridization complexes that included capture oligomers of SEQ ID Nos. 4 and 5 bound to HAV RNA were used in amplification reactions that contained 75 µl of amplification reagent containing amplification oligomers (SEQ ID NO:36 at 13 pmol/reaction and SEQ ID NO:96 at 20 pmol/reaction). As described above, each mixture was covered with an oil layer, incubated 10 min at 60°C, then enzyme reagent (25 µl) was added, and the mixture was incubated for 60 min at 41.5°C to allow amplification of the HAV target sequence. Amplified sequences were detected by using 2-methyl-AE-labeled detection probes (SEQ ID Nos.121 and 122, each at 0.007-0.13 pmol/reaction in 25 µl of probe reagent) which were incubated for 15 min at 60 °C for hybridization of the probes to the amplified

HAV sequences. Then 250  $\mu$ l of selection reagent was added and the mixture was incubated 10 min at 60°C to hydrolyze label on unbound probes, and detection was performed as described above using Detect Reagents I and II to produce chemiluminescence (RLU) measured in a luminometer (LEADER™ HC Plus, Gen-Probe Inc.).

[0062] The results showed that the assay has a sensitivity of about 80 to 100% for samples containing 25 HAV per ml, about 90 to 100% for samples containing 30 HAV per ml, about 98 to 100% for samples containing 100 HAV per ml, and 100% for samples containing 300 and 500 HAV per ml. No positive results were detected for the negative controls (containing no HAV). These results show that the assay detects HAV in clinical samples with a sensitivity of about 25 copies of HAV per ml of sample.

#### Example 5: Detection of HAV RNA and Another Viral Target in the Same Sample

[0063] This assay includes the steps of target capture, amplification and detection substantially as described in Examples 1 to 4 to detect HAV and includes additional oligomers to capture, amplify and detect another target, human parvovirus B19, in the sample. To detect HAV, capture oligomers are SEQ ID Nos. 4 and 5, amplification oligomers are SEQ ID Nos. 36 and 96, and the detection probes are SEQ ID Nos. 121 and 122, used substantially as described in Example 4. To detect parvovirus B19 nucleic acid, a capture probe of SEQ ID NO:169, amplification oligomers of SEQ ID Nos. 170 and 171, and detection probes of SEQ ID NO:173 labeled between residues 5 and 6 and SEQ ID NO:174 labeled between residues 9 and 10 are used, similar to a previously described assay to detect parvovirus B19 (U.S. Pat. Pub. No. US-2003-0124578-A1). Samples of 0.5 ml normal human plasma are prepared to contain known amounts of HAV and parvovirus B19, and then mixed with 0.4 ml of target capture reagent containing the capture oligomers (SEQ ID Nos. 4, 5 and 169). Target capture steps are performed substantially as described in Example 2 and the purified viral targets are amplified in the same amplification reaction mixture, substantially as described in Examples 3 and 4, but using the HAV-specific and parvovirus-specific amplification oligomers described above. Following amplification of the viral target sequences, the detection step uses HAV-specific probes and parvovirus-specific probes as described above, but the probes for the different viral targets are labeled with different AE compounds to allow detection of different signals in the detection step by using differential kinetics (described in U.S. Pat. No. 5,658,737). In these assays, both HAV and parvovirus B19 nucleic acids are detected in the samples that contain both targets. The sensitivity of the assay for detection of HAV is about 25 copies/ml, i.e., positive signals are detected for 80% to 100% of the samples containing 25, 30, 100, 300 and 500 copies/ml of HAV. The sensitivity of the assay for detection of parvovirus B19 in the sample is as low as 150 international units/ml (IU/ml), i.e., positive signals are detected for 20% to 40% of samples containing 150 IU/ml. The assay reliably detects 400 or more parvovirus B19 IU/ml, i.e., 70% to 100% positive detection for samples containing 400, 600, 800, 1600 and 3000 IU/ml. These results show that HAV nucleic acid is specifically detected when the sample includes HAV and another virus, parvovirus B19, which is also specifically detected.

#### Further Embodiments

##### [0064]

1. A combination of at least two oligomers for amplifying a HAV target region comprising:

for a first HAV target region, oligomers of about 23 to 26 nt contained in the sequence of SEQ ID NO:138 that include at least the sequence of SEQ ID NO:139 or SEQ ID NO:140, or oligomers in a size range of about 19 to 25 nt contained in the sequence of SEQ ID NO: 141 that contain at least one sequence of SEQ ID Nos. 142 to 146, or promoter primer oligomers in a size range of about 50 to 53 nt that include HAV target-specific portions of any one of SEQ ID Nos. 21 to 27;

for a second HAV target region, oligomers of about 21 to 27 nt contained in the sequence of SEQ ID NO:60 or contained in the sequence of SEQ ID NO:86 that include at least the sequence of SEQ ID NO:156, or promoter primer oligomers in a size range of about 48 to 54 nt that include HAV target-specific portions of any one of SEQ ID Nos. 29 to 32;

for a third HAV target region, oligomers of about 24 to 30 nt contained in the sequence of SEQ ID NO:147 that include at least the sequence of SEQ ID NO:148, or contained in the sequence of SEQ ID NO:157 that include at least the sequence of SEQ ID NO:158, or promoter primer oligomers that include HAV target-specific portions of SEQ ID NO:31 or SEQ ID NO:32;

for a fourth HAV target region, oligomers of about 18 to 27 nt contained in the sequence of SEQ ID NO:93 or SEQ ID NO:95 that contain at least the sequence of SEQ ID NO:97, SEQ ID NO:159, or SEQ ID NO:160, or a promoter primer oligomer that includes a HAV target-specific portion of SEQ ID NO:33;

for a fifth HAV target region, oligomers of about 19 to 31 nt contained in the sequence of SEQ ID NO:149 that include at least the sequence of SEQ ID NO:150, or promoter primer oligomers in a size range of about 51 to

56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 34 to 40;  
for a sixth HAV target region, oligomers of about 24 to 28 nt contained in the sequence of SEQ ID NO:161 that  
include at least the sequence of SEQ ID NO:162, or promoter primer oligomers are embodiments of promoter  
primers that include HAV target-specific portions of SEQ ID NO:41 or SEQ ID NO:42; and  
for a seventh HAV target region, oligomers of about 20 to 30 nt contained in the sequence of SEQ ID NO: 151  
that include at least any one of the sequences of SEQ ID NO:152 to SEQ ID NO:155, or contained in SEQ ID  
NO:163 that include at least the sequence of SEQ ID NO:164, or contained in SEQ ID NO:166 that include at  
least any one of the sequences of SEQ ID Nos. 166 to 168, or promoter primer oligomers in a size range of  
about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 43 to 49.

2. The combination of at least two oligomers of embodiment 1 for the first HAV target region selected from the group  
consisting of: SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:21, SEQ  
ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:50, SEQ  
ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ  
ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:143, SEQ  
ID NO:144, and SEQ ID NO:145.

3. The combination of at least two oligomers of embodiment 1 for the second HAV target region selected from the  
group consisting of: SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:  
58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, and SEQ ID NO:156.

4. The combination of at least two oligomers of embodiment 1 for the third HAV target region selected from the  
group consisting of: SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:89, SEQ ID NO:  
90, SEQ ID NO:91, and SEQ ID NO:148.

5. The combination of at least two oligomers of embodiment 1 for the fourth HAV target region selected from the  
group consisting of: SEQ ID NO:33, SEQ ID NO:63, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:  
95, SEQ ID NO:96, and SEQ ID NO:97.

6. The combination of at least two oligomers of embodiment 1 for the fifth HAV target region selected from the group  
consisting of: SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ  
ID NO:40, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ  
ID NO:70, SEQ ID NO:97, SEQ ID NO:149, and SEQ ID NO:150.

7. The combination of at least two oligomers of embodiment 1 for the sixth HAV target region selected from the  
group consisting of: SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:98, SEQ ID NO:  
99, SEQ ID NO:101, SEQ ID NO:161, and SEQ ID NO:162.

8. The combination of at least two oligomers of embodiment 1 for the seventh HAV target region selected from the  
group consisting of: SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:  
48, SEQ ID NO:49, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:  
78, SEQ ID NO:79, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:104, SEQ ID NO:106, SEQ ID  
NO:107, SEQ ID NO:108, SEQ ID NO:152, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:163, SEQ ID NO:164,  
SEQ ID NO:165, SEQ ID NO:166, SEQ ID NO:167, and SEQ ID NO:168.

9. The combination of at least two oligomers of embodiment 1, further comprising at least one capture probe oligomer  
selected from the group consisting of SEQ ID Nos. 1 to 14.

10. The combination of at least two oligomers of embodiment 1, further comprising at least one detection probe  
oligomer selected from the group consisting of SEQ ID NO: 111, SEQ ID NO:113, SEQ ID NO:115, SEQ  
ID NO:117, SEQ ID NO:119, SEQ ID NO:121 to SEQ ID NO:124, and SEQ ID NO:126 to SEQ ID NO:130.

11. A kit comprising a combination of at least two oligomers according to embodiment 1.

12. A method of detecting the presence of HAV in a sample comprising the steps of:

purifying HAV nucleic acid from other components in a sample containing HAV;  
amplifying a HAV target sequence in the purified HAV nucleic acid, or a cDNA made therefrom, by using an *in*  
*vitro* amplification reaction that includes at least two amplification oligomers specific for a selected HAV target  
region, which include:

for a first HAV target region, oligomers of about 23 to 26 nt contained in the sequence of SEQ ID NO:138  
that include at least the sequence of SEQ ID NO:139 or SEQ ID NO:140, or oligomers in a size range of  
about 19 to 25 nt contained in the sequence of SEQ ID NO: 141 that contain at least one sequence of SEQ  
ID Nos. 142 to 146, or promoter primer oligomers in a size range of about 50 to 53 nt that include HAV  
target-specific portions of any one of SEQ ID Nos. 21 to 27;

for a second HAV target region, oligomers of about 21 to 27 nt contained in the sequence of SEQ ID NO:  
60 or contained in the sequence of SEQ ID NO:86 that include at least the sequence of SEQ ID NO:156,



or promoter primer oligomers in a size range of about 48 to 54 nt that include HAV target-specific portions of any one of SEQ ID Nos. 29 to 32;

for a third HAV target region, oligomers of about 24 to 30 nt contained in the sequence of SEQ ID NO:147 that include at least the sequence of SEQ ID NO:148, or contained in the sequence of SEQ ID NO:157 that include at least the sequence of SEQ ID NO:158, or promoter primer oligomers that include HAV target-specific portions of SEQ ID NO:31 or SEQ ID NO:32;

for a fourth HAV target region, oligomers of about 18 to 27 nt contained in the sequence of SEQ ID NO:93 or SEQ ID NO:95 that contain at least the sequence of SEQ ID NO:97, SEQ ID NO:159, or SEQ ID NO:160, or a promoter primer oligomer that includes a HAV target-specific portion of SEQ ID NO:33;

for a fifth HAV target region, oligomers of about 19 to 31 nt contained in the sequence of SEQ ID NO:149 that include at least the sequence of SEQ ID NO:150, or promoter primer oligomers in a size range of about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 34 to 40;

for a sixth HAV target region, oligomers of about 24 to 28 nt contained in the sequence of SEQ ID NO:161 that include at least the sequence of SEQ ID NO:162, or promoter primer oligomers are embodiments of promoter primers that include HAV target-specific portions of SEQ ID NO:41 or SEQ ID NO:42; and

for a seventh HAV target region, oligomers of about 20 to 30 nt contained in the sequence of SEQ ID NO:151 that include at least any one of the sequences of SEQ ID NO:152 to SEQ ID NO:155, or contained in SEQ ID NO:163 that include at least the sequence of SEQ ID NO:164, or contained in SEQ ID NO:165 that include at least any one of the sequences of SEQ ID Nos. 166 to 168, or promoter primer oligomers in a size range of about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 43 to 49

to produce an amplified product of the selected HAV target region; and

detecting the amplified product by using a detection probe that hybridizes specifically with at least a portion of the amplified product.

13. The method of embodiment 12, wherein the purifying step contacts the sample with at least one capture probe oligomer comprising a sequence contained in any one of SEQ ID Nos. 1 to 14 that hybridizes specifically to a sequence in HAV RNA to form a hybridization complex with the HAV RNA, and separates the hybridization complex that contains the HAV RNA from other sample components.

14. The method of embodiment 12, wherein the amplifying step amplifies a sequence in the first HAV target region by using at least two oligomers specific for the first HAV target region selected from the group consisting of: SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:63, SEQ ID NO:54, SEQ ID NO:65, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:143, SEQ ID NO:144, and SEQ ID NO:145; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the first HAV target region.

15. The method of embodiment 12, wherein the amplifying step amplifies a sequence in the second HAV target region by using at least two oligomers specific for the second HAV target region selected from the group consisting of: SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, and SEQ ID NO:156; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the second HAV target region.

16. The method of embodiment 12, wherein the amplifying step amplifies a sequence in the third HAV target region by using at least two oligomers specific for the third HAV target region selected from the group consisting of: SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, and SEQ ID NO:148; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the third HAV target region.

17. The method of embodiment 12, wherein the amplifying step amplifies a sequence in the fourth HAV target region by using at least two oligomers specific for the fourth HAV target region selected from the group consisting of: SEQ ID NO:33, SEQ ID NO:63, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, and SEQ ID NO:97; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the fourth HAV target region.

18. The method of embodiment 12, wherein the amplifying step amplifies a sequence in the fifth HAV target region by using at least two oligomers specific for the fifth HAV target region selected from the group consisting of: SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ

ID NO:97, SEQ ID NO:149, and SEQ ID NO:150; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the fifth HAV target region.

19. The method of embodiment 12, wherein the amplifying step amplifies a sequence in the sixth HAV target region by using at least two oligomers specific for the sixth HAV target region selected from the group consisting of: SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:161, and SEQ ID NO:162; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the sixth HAV target region.

20. The method of embodiment 12, wherein the amplifying step amplifies a sequence in the seventh HAV target region by using at least two oligomers specific for the seventh HAV target region selected from the group consisting of: SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:152, SEQ ID NO:163, SEQ ID NO:155, SEQ ID NO:163, SEQ ID NO:164, SEQ ID NO:165, SEQ ID NO:166, SEQ ID NO:167, and SEQ ID NO:168; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the seventh HAV target region.

## EP 2 402 465 A2

### SEQUENCE LISTING

5           <110> Gen-Probe Incorporated

            <120> Compositions and methods for detection of hepatitis A virus nucleic aci

10           <130> GP135-PCT

15           <140> To Be Assigned

            <141> 2005-07-13

20           <150> US 60/587,734

            <151> 2004-07-13

25           <160> 174

            <170> PatentIn version 3.1

30           <210> 1

            <211> 54

35           <212> DNA

            <213> Artificial Sequence

40           <220>

            <223> synthetic oligomer, capture probe

45           <400> 1  
ggacuuccaa gaggggcucc gtttaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 54

            <210> 2

50           <211> 60

            <212> DNA

            <213> Artificial Sequence

55           <220>

## EP 2 402 465 A2

<223> synthetic oligomer, capture probe  
 <400> 2  
 uuuagacucc uacagcucca ugcuaauttt aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60  
 5  
 <210> 3  
 <211> 61  
 10  
 <212> DNA  
 <213> Artificial Sequence  
 15  
 <220>  
 <223> synthetic oligomer, capture probe  
 <400> 3  
 20 uucauuucug uccauuucuc aucauucatt taaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60  
 a 61  
 25  
 <210> 4  
 <211> 58  
 <212> DNA  
 30  
 <213> Artificial Sequence  
 <220>  
 35  
 <223> synthetic oligomer, capture probe  
 <400> 4  
 gaaauugaau aguaaguucc accuctttaa aaaaaaaaaa aaaaaaaaaa aaaaaaaa 58  
 40  
 <210> 5  
 <211> 60  
 <212> DNA  
 45  
 <213> Artificial Sequence  
 <220>  
 50  
 <223> synthetic oligomer, capture probe  
 <400> 5  
 gcuaugcugc aggaaaaaua aucauggttt aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60  
 55  
 <210> 6

## EP 2 402 465 A2

<211> 56  
 <212> DNA  
 5 <213> Artificial Sequence  
  
 <220>  
 10 <223> synthetic oligomer, capture probe  
 <400> 6  
 gcacagcugc agggaaaaua aucttttaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 56  
 15  
 <210> 7  
 <211> 58  
 <212> DNA  
 20 <213> Artificial Sequence  
  
 <220>  
 25 <223> synthetic oligomer, capture probe  
 <400> 7  
 gacaaaagaa aacuggagac uuucctttaa aaaaaaaaaa aaaaaaaaaa aaaaaaa 58  
 30  
 <210> 8  
 <211> 21  
 35 <212> DNA  
 <213> Artificial Sequence  
  
 40 <220>  
 <223> synthetic oligomer  
 <400> 8  
 45 ggacttccaa gaggggctcc g 21  
  
 <210> 9  
 <211> 27  
 50 <212> DNA  
 <213> Artificial Sequence  
  
 55 <220>

## EP 2 402 465 A2

	<223> synthetic oligomer	
	<400> 9	
5	tttagactcc tacagctcca tgctaata	27
	<210> 10	
	<211> 28	
10	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> synthetic oligomer	
	<400> 10	
20	ttcattttctg tccattttctc atcattca	28
	<210> 11	
	<211> 25	
25	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> synthetic oligomer	
	<400> 11	
35	gaaattgaat agtaagttcc acctc	25
	<210> 12	
40	<211> 27	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> synthetic oligomer	
	<400> 12	
50	gcatagtctgc aggaaaatta atcatgg	27
	<210> 13	
55	<211> 23	

## EP 2 402 465 A2

<212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> synthetic oligomer  
 10  
 <400> 13  
 gcatagctgc aggaaaatta atc 23  
 <210> 14  
 15  
 <211> 25  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> synthetic oligomer  
 25  
 <400> 14  
 gacaaaagaa aactggagac ttcc 25  
 30  
 <210> 15  
 <211> 23  
 <212> DNA  
 35  
 <213> Artificial Sequence  
 <220>  
 40  
 <223> synthetic oligomer  
 <400> 15  
 ccatggtgag gggacttgat acc 23  
 45  
 <210> 16  
 <211> 22  
 <212> DNA  
 50  
 <213> Artificial Sequence  
 <220>  
 55  
 <223> synthetic oligomer

## EP 2 402 465 A2

	<400> 16 cLLgatacctL caccgccgLL Lg	22
5	<210> 17  <211> 23  <212> DNA	
10	<213> Artificial Sequence	
15	<220>  <223> synthetic oligomer  <400> 17 ttgatacctc accgccgttt gcc	23
20	<210> 18  <211> 25  <212> DNA	
25	<213> Artificial Sequence	
30	<220>  <223> synthetic oligomer  <400> 18 galacctcac cgccgLLLgc clagg	25
35	<210> 19  <211> 27  <212> DNA	
40	<213> Artificial Sequence	
45	<220>  <223> bacteriophage T7 promoter  <400> 19 aatttaatac gactcactat agggaga	27
50	<210> 20  <211> 49  <212> DNA	
55		



## EP 2 402 465 A2

<213> Artificial Sequence

5

<220>

<223> synthetic oligomer, promoter primer

<220>

10

<221> promoter

<222> (1)..(27)

<223>

15

<400> 20

aattttaatac gactcactat agggagaaga gaaacagatt aaagaaccc

49

20

<210> 21

<211> 53

<212> DNA

25

<213> Artificial Sequence

30

<220>

<223> synthetic oligomer, promoter primer

<220>

35

<221> promoter

<222> (1)..(27)

<223>

40

<400> 21

aattttaatac gactcactat agggagagga agaaagaaga cagaaagcgt gaa

53

45

<210> 22

<211> 51

<212> DNA

50

<213> Artificial Sequence

55

<220>

<223> synthetic oligomer, promoter primer

## EP 2 402 465 A2

<220>  
 <221> promoter  
 5 <222> (1)..(27)  
 <223>  
  
 10 <400> 22  
 aattttaatac gactcactat agggagagga agaaagaaga cagaaagcgt g 51  
  
 <210> 23  
 15 <211> 50  
 <212> DNA  
 <213> Artificial Sequence  
 20  
  
 <220>  
 25 <223> synthetic oligomer, promoter primer  
 <220>  
 <221> promoter  
 30 <222> (1)..(27)  
 <223>  
  
 35 <400> 23  
 aattttaatac gactcactat agggagagaa gaaagaagac agaaagcgtg 50  
  
 <210> 24  
 40 <211> 52  
 <212> DNA  
 <213> Artificial Sequence  
 45  
  
 <220>  
 50 <223> synthetic oligomer, promoter primer  
 <220>  
 <221> promoter  
 55 <222> (1)..(27)  
 <223>

## EP 2 402 465 A2

<400> 24  
 aatlllaatalac gactcactat agggagatgg aagaaagaag acagaaagcg lg 52

5 <210> 25  
 <211> 51  
 <212> DNA  
 10 <213> Artificial Sequence

15 <220>  
 <223> synthetic oligomer, promoter primer  
 <220>

20 <221> promoter  
 <222> (1)..(27)  
 <223>

25 <400> 25  
 aattttaatalac gactcactat agggagatgg aagaaagaag acagaaagcg t 51

30 <210> 26  
 <211> 51  
 <212> DNA  
 35 <213> Artificial Sequence

40 <220>  
 <223> synthetic oligomer, promoter primer  
 <220>

45 <221> promoter  
 <222> (1)..(27)  
 <223>

50 <400> 26  
 aattttaatalac gactcactat agggagactg gaagaaagaa gacagaaagc g 51

55 <210> 27

## EP 2 402 465 A2

<211> 52  
 <212> DNA  
 5 <213> Artificial Sequence  
  
 <220>  
 10 <223> synthetic oligomer, promoter primer  
 <220>  
 <221> promoter  
 15 <222> (1)..(27)  
 <223>  
  
 20 <400> 27  
 aatttaatac gactcactat agggagagca aggggagagc cctggaagaa ag 52  
  
 <210> 28  
 25 <211> 54  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> synthetic oligomer, promoter primer  
 35 <220>  
 <221> promoter  
 <222> (1)..(27)  
 40 <223>  
  
 45 <400> 28  
 aatllaatac gactcactat agggagacag tatllataat tccaacagtc acag 54  
  
 <210> 29  
 50 <211> 48  
 <212> DNA  
 <213> Artificial Sequence  
 55  
 <220>

## EP 2 402 465 A2

<223> synthetic oligomer, promoter primer  
 <220>  
 5 <221> promoter  
 <222> (1)..(27)  
 <223>  
 10  
 <400> 29  
 aatttaatac gactcactat agggagatct caacaaacca attatgtg 48  
 15 <210> 30  
 <211> 54  
 <212> DNA  
 20 <213> Artificial Sequence  
 <220>  
 25 <223> synthetic oligomer, promoter primer  
 <220>  
 30 <221> promotet  
 <222> (1)..(27)  
 <223>  
 35  
 <400> 30  
 aaLLLaalac gactcactat agggagacat gactcLcaac aaaccaalla LgLg 54  
 40 <210> 31  
 <211> 51  
 <212> DNA  
 45 <213> Artificial Sequence  
 <220>  
 50 <223> synthetic oligomer, promoter primer  
 <220>  
 55 <221> promoter  
 <222> (1)..(27)

## EP 2 402 465 A2

<223>

5           <400> 31  
aatttaatac gactcactat agggagacaa ttgcttcctt aacataaact g 51

<210> 32

10           <211> 53  
            <212> DNA  
            <213> Artificial Sequence

15

<220>

<223> synthetic oligomer, promoter primer

20           <220>

<221> promoter

<222> (1)..(27)

25           <223>

<400> 32

30           aatttaatac gactcactat agggagacga tcaattgctt ccttaacata aac 53

<210> 33

<211> 47

35           <212> DNA  
            <213> Artificial Sequence

40

<220>

<223> synthetic oligomer, promoter primer

<220>

45           <221> promoter

<222> (1)..(27)

<223>

50

<400> 33

55           aatttaatac gactcactat agggagacct ttctctctcc atgcttg 47

<210> 34

## EP 2 402 465 A2

<211> 52  
<212> DNA  
5 <213> Artificial Sequence  
  
<220>  
10 <223> synthetic oligomer, promoter primer  
<220>  
<221> promoter  
15 <222> (1)..(27)  
<223>  
  
20 <400> 34  
aatttaatac gactcactat agggagagaa ttgaatttcc tccagcaaca tg 52  
  
<210> 35  
25 <211> 56  
<212> DNA  
<213> Artificial Sequence  
30  
  
<220>  
<223> synthetic oligomer, promoter primer  
35 <220>  
<221> promoter  
<222> (1)..(27)  
40 <223>  
  
45 <400> 35  
aatllaatac gactcactat agggagaaca agaattgaat ttctccagc aacatg 56  
  
<210> 36  
50 <211> 54  
<212> DNA  
<213> Artificial Sequence  
55  
  
<220>

## EP 2 402 465 A2

<223> synthetic oligomer, promoter primer  
 <220>  
 5 <221> promoter  
 <222> (1)..(27)  
 <223>  
 10  
 <400> 36  
 aatttaatac gactcactat aggagaaca agaattgaat ttctccagc aaca 54  
 15  
 <210> 37  
 <211> 52  
 <212> DNA  
 20 <213> Artificial Sequence  
 <220>  
 25 <223> synthetic oligomer, promoter primer  
 <220>  
 30 <221> promoter  
 <222> (1)..(27)  
 <223>  
 35  
 <400> 37  
 aalllaalac gactcactat aggagacaa gaallgaall tctccagca ac 52  
 40  
 <210> 38  
 <211> 51  
 <212> DNA  
 45 <213> Artificial Sequence  
 <220>  
 50 <223> synthetic oligomer, promoter primer  
 <220>  
 55 <221> promoter  
 <222> (1)..(27)



# EP 2 402 465 A2

<223>

5                   <400> 38  
aatttaatac gactcactat agggagaaag aattgaattt cctccagcaa c 51

<210> 39

10                   <211> 53

<212> DNA

<213> Artificial Sequence

15

<220>

<223> synthetic oligomer, promoter primer

20

<220>

<221> promoter

<222> (1)..(27)

25

<223>

30                   <400> 39  
aatttaatac gactcactat agggagaaca agaattgaat ttctctcagc aac 53

<210> 40

<211> 52

35

<212> DNA

<213> Artificial Sequence

40

<220>

<223> synthetic oligomer, promoter primer

<220>

45

<221> promoter

<222> (1)..(27)

50

<223>

55                   <400> 40  
aatttaatac gactcactat agggagacca caagaattga atttctctca gc 52

<210> 41

## EP 2 402 465 A2

<211> 52  
 <212> DNA  
 5 <213> Artificial Sequence  
  
 <220>  
 10 <223> synthetic oligomer, promoter primer  
 <220>  
 <221> promoter  
 15 <222> (1)..(27)  
 <223>  
  
 20 <400> 41  
 aatttaatac gactcactat agggagactc tgagccaatc ttggatgaac tc 52  
  
 <210> 42  
 25 <211> 52  
 <212> DNA  
 <213> Artificial Sequence  
 30  
 <220>  
 <223> synthetic oligomer, promoter primer  
 35 <220>  
 <221> promoter  
 <222> (1)..(27)  
 40 <223>  
  
 <400> 42  
 45 aatllaatac gactcactat agggagacag aacaatlltc calcatgaca gl 52  
  
 <210> 43  
 50 <211> 55  
 <212> DNA  
 <213> Artificial Sequence  
 55  
 <220>

## EP 2 402 465 A2

<223> synthetic oligomer, promoter primer

<220>

5 <221> promoter

<222> (1)..(27)

<223>

10

<400> 43  
aatttaatac gactcactat aggagagagt cataaaatct cattctccac caatc 55

15 <210> 44

<211> 56

<212> DNA

20 <213> Artificial Sequence

<220>

25 <223> synthetic oligomer, promoter primer

<220>

30 <221> promoter

<222> (1)..(27)

<223>

35

<400> 44  
aaLLLaalac gactcactaL aggagagagaa acacLggLca Laaaalclca LLclcc 56

40 <210> 45

<211> 52

<212> DNA

45 <213> Artificial Sequence

<220>

50 <223> synthetic oligomer, promoter primer

<220>

55 <221> promoter

<222> (1)..(27)

## EP 2 402 465 A2

<223>

5           <400> 45  
aatttaatac gactcactat agggagaggt cacaaatgaa acactggcca ta 52

<210> 46

10           <211> 53  
            <212> DNA  
            <213> Artificial Sequence

15

<220>

<223> synthetic oligomer, promoter primer

20           <220>

<221> promoter

<222> (1)..(27)

25           <223>

30           <400> 46  
aatttaatac gactcactat agggagagaa aggtcacaaa tgaacactg gtc 53

<210> 47

<211> 52

35           <212> DNA  
            <213> Artificial Sequence

40

<220>

<223> synthetic oligomer, promoter primer

45           <220>

<221> promoter

<222> (1)..(27)

50           <223>

55           <400> 47  
aatttaatac gactcactat agggagaaaa ggtcacaaat gaaacactgg tc 52

<210> 48

## EP 2 402 465 A2

<211> 51  
<212> DNA  
5 <213> Artificial Sequence  
  
<220>  
10 <223> synthetic oligomer, promoter primer  
<220>  
<221> promoter  
15 <222> (1)..(27)  
<223>  
  
20 <400> 48  
aatttaatac gactcactat agggagagaa aggtcacaaa tgaacactg g 51  
  
<210> 49  
25 <211> 53  
<212> DNA  
<213> Artificial Sequence  
30  
  
<220>  
<223> synthetic oligomer, promoter primer  
35 <220>  
<221> promoter  
<222> (1)..(27)  
40 <223>  
  
<400> 49  
45 aatllaatac gactcactat agggagacaa atcatgaaag gtcacaaatg aaa 53  
  
<210> 50  
50 <211> 22  
<212> DNA  
<213> Artificial Sequence  
55  
  
<220>

## EP 2 402 465 A2

	<223> synthetic oligomer, primer	
	<400> 50	
5	agagaaacag attaaagaac cc	22
	<210> 51	
	<211> 26	
10	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> synthetic oligomer, primer	
	<400> 51	
20	ggaagaaaga agacagaaag cgtgaa	26
	<210> 52	
	<211> 24	
25	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> synthetic oligomer, primer	
	<400> 52	
35	ggaagaaaga agacagaaag cgtg	24
	<210> 53	
40	<211> 23	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> synthetic oligomer, primer	
	<400> 53	
50	gaagaaagaa gacagaaagc gtg	23
	<210> 54	
55	<211> 25	

## EP 2 402 465 A2

<212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> synthetic oligomer, primer  
 10  
 <400> 54  
 tggaagaaag aagacagaaa gcgtg 25  
 <210> 55  
 15  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> synthetic oligomer, primer  
 25  
 <400> 55  
 tggaagaaag aagacagaaa gcgt 24  
 30  
 <210> 56  
 <211> 24  
 <212> DNA  
 35  
 <213> Artificial Sequence  
 <220>  
 40  
 <223> synthetic oligomer, primer  
 <400> 56  
 ctggaagaaa gaagacagaa agcg 24  
 45  
 <210> 57  
 <211> 25  
 <212> DNA  
 50  
 <213> Artificial Sequence  
 <220>  
 55  
 <223> synthetic oligomer, primer

## EP 2 402 465 A2

<400> 57  
gcaaggggag agccclggaa gaaag 25

5 <210> 58  
<211> 27  
<212> DNA

10 <213> Artificial Sequence

<220>

15 <223> synthetic oligomer, primer  
<400> 58  
cagtatttat aatttcaaca gtcacag 27

20 <210> 59  
<211> 21  
<212> DNA

25 <213> Artificial Sequence

<220>

30 <223> synthetic oligomer, primer  
<400> 59  
Lclcaacaaa ccaallatgt g 21

35 <210> 60  
<211> 27  
<212> DNA

40 <213> Artificial Sequence

<220>

45 <223> synthetic oligomer, primer  
<400> 60  
catgactctc aacaaacca ttatgtg 27

50 <210> 61  
<211> 24  
<212> DNA

55



## EP 2 402 465 A2

<213> Artificial Sequence

5	<220>		
	<223>	synthetic oligomer, primer	
	<400>	61	
10		caattgcttc cttaacataa actg	24
	<210>	62	
	<211>	26	
15	<212>	DNA	
	<213>	Artificial Sequence	
20	<220>		
	<223>	synthetic oligomer, primer	
	<400>	62	
25		cgatcaattg cttccttaac ataaac	26
	<210>	63	
	<211>	20	
30	<212>	DNA	
	<213>	Artificial Sequence	
35	<220>		
	<223>	synthetic oligomer, primer	
	<400>	63	
40		ccLLLLccLc LccatgccLg	20
	<210>	64	
45	<211>	25	
	<212>	DNA	
	<213>	Artificial Sequence	
50	<220>		
	<223>	synthetic oligomer, primer	
55	<400>	64	

# EP 2 402 465 A2

gaallgaall lcccccagca acatg

25

<210> 65

5

<211> 29

<212> DNA

<213> Artificial Sequence

10

<220>

<223> synthetic oligomer, primer

15

<400> 65

acaagaattg aatttcctcc agcaacatg

29

<210> 66

20

<211> 27

<212> DNA

<213> Artificial Sequence

25

<220>

<223> synthetic oligomer, primer

30

<400> 66

acaagaattg aatttcctcc agcaaca

27

<210> 67

35

<211> 25

<212> DNA

<213> Artificial Sequence

40

<220>

<223> synthetic oligomer, primer

45

<400> 67

caagaattga atttctcca gcaac

25

<210> 68

50

<211> 24

<212> DNA

<213> Artificial Sequence

55

## EP 2 402 465 A2

	<220>	
	<223> synthetic oligomer, primer	
5	<400> 68 aagaattgaa ttctctccag caac	24
	<210> 69	
10	<211> 26	
	<212> DNA	
15	<213> Artificial Sequence	
	<220>	
20	<223> synthetic oligomer, primer	
	<400> 69 acaagaattg aatttccctcc agcaac	26
25	<210> 70	
	<211> 25	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
35	<223> synthetic oligomer, primer	
	<400> 70 ccacaagaat tgaatttctt ccagc	25
40	<210> 71	
	<211> 25	
	<212> DNA	
45	<213> Artificial Sequence	
	<220>	
50	<223> synthetic oligomer, primer	
	<400> 71 ctctgagcca atcttgatg aactc	25
55	<210> 72	

## EP 2 402 465 A2

<211> 25  
 <212> DNA  
 5 <213> Artificial Sequence  
  
 <220>  
 10 <223> synthetic oligomer, primer  
 <400> 72  
 cagaacaatt ttccatcatg acagt 25  
 15  
 <210> 73  
 <211> 28  
 <212> DNA  
 20 <213> Artificial Sequence  
  
 <220>  
 25 <223> synthetic oligomer, primer  
 <400> 73  
 ggtcataaaa tctcattctc caccaatc 28  
 30  
 <210> 74  
 <211> 29  
 35 <212> DNA  
 <213> Artificial Sequence  
  
 40 <220>  
 <223> synthetic oligomer, primer  
 <400> 74  
 45 gaaacactgg tcataaaatc tcattctcc 29  
  
 <210> 75  
 <211> 25  
 50 <212> DNA  
 <213> Artificial Sequence  
 55  
 <220>

## EP 2 402 465 A2

	<223> synthetic oligomer, primer	
5	<400> 75 ggtcacaaat gaaacactgg tcata	25
	<210> 76	
	<211> 26	
10	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> synthetic oligomer, primer	
20	<400> 76 gaaaggtcac aaatgaaaca ctggtc	26
	<210> 77	
25	<211> 25	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> synthetic oligomer, primer	
35	<400> 77 aaaggtcaca aatgaaacac tggtc	25
	<210> 78	
40	<211> 24	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> synthetic oligomer, primer	
50	<400> 78 gaaaggtcac aaatgaaaca ctgg	24
	<210> 79	
55	<211> 26	

	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> synthetic oligomer, primer	
10	<400> 79 caaatcatga aaggtcacaa atgaaa	26
	<210> 80	
15	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> synthetic oligomer, primer	
25	<400> 80 gacttgatac ctcaccgcc	19
	<210> 81	
30	<211> 23	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
40	<223> synthetic oligomer, primer	
	<400> 81 acttgatacc tcaccgccgt ttg	23
45	<210> 82	
	<211> 23	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
55	<223> synthetic oligomer, primer	

## EP 2 402 465 A2

	<400> 82 cLLgatacct caccgccgLL tgc	23
5	<210> 83  <211> 24  <212> DNA	
10	<213> Artificial Sequence	
	<220>	
15	<223> synthetic oligomer, primer  <400> 83 cttgatacct caccgccggtt tgcc	24
20	<210> 84  <211> 22  <212> DNA	
25	<213> Artificial Sequence	
	<220>	
30	<223> synthetic oligomer, primer  <400> 84 cuugatacct caccgccgLL tg	22
35	<210> 85  <211> 23  <212> DNA	
40	<213> Artificial Sequence	
	<220>	
45	<223> synthetic oligomer, primer  <400> 85 cuugatacct caccgccggtt tgc	23
50	<210> 86  <211> 26  <212> DNA	
55		

## EP 2 402 465 A2

<213> Artificial Sequence

5	<220>	
	<223> synthetic oligomer, primer	
	<400> 86	
10	gcagatagaa tgcttggatt gtctgg	26
	<210> 87	
	<211> 25	
15	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> synthetic oligomer, primer	
	<400> 87	
25	gcagatagaa tgcttggatt gtctg	25
	<210> 88	
	<211> 25	
30	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> synthetic oligomer, primer	
	<400> 88	
40	cagatagaal gcttggallg tctgg	25
	<210> 89	
45	<211> 30	
	<212> DNA	
	<213> Artificial Sequence	
50	<220>	
	<223> synthetic oligomer, primer	
55	<400> 89	



LcLccLLLLa LaaLagcaac LlcaaaLlgg

30

<210> 90

5

<211> 29

<212> DNA

<213> Artificial Sequence

10

<220>

<223> synthetic oligomer, primer

15

<400> 90

ctccttttat aatagcaact tcaaatgg

29

<210> 91

20

<211> 25

<212> DNA

<213> Artificial Sequence

25

<220>

<223> synthetic oligomer, primer

30

<400> 91

ttataatagc aacttcaaat tggtc

25

<210> 92

35

<211> 20

<212> DNA

<213> Artificial Sequence

40

<220>

<223> synthetic oligomer, primer

45

<400> 92

ggcaacatta gtgacaactg

20

<210> 93

50

<211> 27

<212> DNA

<213> Artificial Sequence

55

## EP 2 402 465 A2

	<220>	
	<223> synthetic oligomer, primer	
5	<400> 93 ggcaacatta gtgacaaactg ttaatgg	27
	<210> 94	
10	<211> 18	
	<212> DNA	
15	<213> Artificial Sequence	
	<220>	
20	<223> synthetic oligomer, primer	
	<400> 94 caggcatgga gaggaaaa	18
25	<210> 95	
	<211> 20	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
35	<223> synthetic oligomer, primer	
	<400> 95 caggcatgga gaggaaaagg	20
40	<210> 96	
	<211> 19	
	<212> DNA	
45	<213> Artificial Sequence	
	<220>	
50	<223> synthetic oligomer, primer	
	<400> 96 caggcatgga gaggaaaag	19
55	<210> 97	

## EP 2 402 465 A2

<211> 18

<212> DNA

5 <213> Artificial Sequence

<220>

10 <223> synthetic oligomer, primer

<400> 97  
ggcatggaga ggaaaagg 18

15 <210> 98

<211> 23

<212> DNA

20 <213> Artificial Sequence

<220>

25 <223> synthetic oligomer, primer

<400> 98  
taagaaaatt gaaatgcaga gaa 23

30 <210> 99

<211> 26

35 <212> DNA

<213> Artificial Sequence

40 <220>

<223> synthetic oligomer, primer

<400> 99  
ctccaaaacg ctttttagaa agagtc 26

45 <210> 100

<211> 22

50 <212> DNA

<213> Artificial Sequence

55 <220>

## EP 2 402 465 A2

	<223> synthetic oligomer	
	<400> 100	
5	ccatgattaa ttttcctgca gc	22
	<210> 101	
	<211> 26	
10	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> synthetic oligomer, primer	
	<400> 101	
20	ccaaaacgct ttttagaaag agtccc	26
	<210> 102	
	<211> 25	
25	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> synthetic oligomer, primer	
	<400> 102	
35	cgctgagttt gagcagaatt tagaa	25
	<210> 103	
40	<211> 28	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> synthetic oligomer, primer	
	<400> 103	
50	ctgaglllga gcagaallla gaaaatgc	28
	<210> 104	
55	<211> 24	

## EP 2 402 465 A2

	<212> DNA	
	<213> Artificial Sequence	
5	<220>	
	<223> synthetic oligomer, primer	
10	<400> 104 atgcatggct atgagtttta tcag	24
	<210> 105	
15	<211> 26	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> synthetic oligomer, primer	
25	<400> 105 gcatggctat gagttttatc agaaat	26
	<210> 106	
30	<211> 27	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
40	<223> synthetic oligomer, primer	
	<400> 106 gcauggctat gagttttatc agaaatt	27
45	<210> 107	
	<211> 26	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
55	<223> synthetic oligomer, primer	

## EP 2 402 465 A2

	<400> 107 caTggcTatg agTtTtTatCa gaaatt	26
5	<210> 108 <211> 27 <212> DNA	
10	<213> Artificial Sequence	
15	<220> <223> synthetic oligomer, primer <400> 108 gcTggcTat gagTtTtTatC agaaatt	27
20	<210> 109 <211> 20 <212> DNA	
25	<213> Artificial Sequence	
30	<220> <223> synthetic oligomer probe <400> 109 gccgTtTgcc TaggcTatag	20
35	<210> 110 <211> 20 <212> DNA	
40	<213> Artificial Sequence	
45	<220> <223> synthetic oligomer <400> 110 ctatagccta ggcaaacggc	20
50	<210> 111 <211> 25 <212> DNA	
55		

# EP 2 402 465 A2

<213> Artificial Sequence

5	<220>		
	<223>	synthetic oligomer probe	
	<400>	111	
10		caggggttctt taatctgttt ctcta	25
	<210>	112	
	<211>	25	
15	<212>	DNA	
	<213>	Artificial Sequence	
20	<220>		
	<223>	synthetic oligomer	
	<400>	112	
25		tagagaaaca gattaaagaa ccctg	25
	<210>	113	
	<211>	27	
30	<212>	DNA	
	<213>	Artificial Sequence	
35	<220>		
	<223>	synthetic oligomer, probe	
	<400>	113	
40		atgatgtttg gatllcatca ttctgtg	27
	<210>	114	
45	<211>	27	
	<212>	DNA	
	<213>	Artificial Sequence	
50	<220>		
	<223>	synthetic oligomer	
55	<400>	114	

# EP 2 402 465 A2

cacagaa1ga 1gaaalccaa acalcal

27

5

<210> 115

<211> 22

<212> DNA

<213> Artificial Sequence

10

<220>

<223> synthetic oligomer, probe

15

<400> 115

ggtcaaatacc aagtccaaaa ac

22

20

<210> 116

<211> 22

<212> DNA

<213> Artificial Sequence

25

<220>

<223> synthetic oligomer

30

<400> 116

gtttttggac ttggatttga cc

22

35

<210> 117

<211> 20

<212> RNA

<213> Artificial Sequence

40

<220>

<223> synthetic oligomer, probe

45

<400> 117

ccuauguuaa uuucugaggg

20

50

<210> 118

<211> 20

<212> DNA

<213> Artificial Sequence

55



# EP 2 402 465 A2

	<220>	
	<223> synthetic oligomer	
5	<400> 118 ccctcagaaa ttaacatagg	20
	<210> 119	
10	<211> 20	
	<212> RNA	
15	<213> Artificial Sequence	
	<220>	
20	<223> synthetic oligomer, probe	
	<400> 119 caggcaugga gaggaaaagg	20
25	<210> 120	
	<211> 20	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
35	<223> synthetic oligomer	
	<400> 120 ccttttcctc tccatgcctg	20
40	<210> 121	
	<211> 23	
	<212> RNA	
45	<213> Artificial Sequence	
	<220>	
50	<223> synthetic oligomer, probe	
	<400> 121 ggucuuccug gaaugugugg ugg	23
55	<210> 122	

# EP 2 402 465 A2

<211> 23  
 <212> RNA  
 5 <213> Artificial Sequence  
  
 <220>  
 10 <223> synthetic oligomer, probe  
 <400> 122  
 gucuuccugg aauguguggu ggg 23  
 15 <210> 123  
 <211> 21  
 <212> DNA  
 20 <213> Artificial Sequence  
  
 <220>  
 25 <223> synthetic oligomer, probe  
 <400> 123  
 tcttcctgga atgtgtggtg g 21  
 30 <210> 124  
 <211> 21  
 35 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 40 <223> synthetic oligomer, probe  
 <400> 124  
 45 tcuuccugga auguguggug g 21  
 <210> 125  
 <211> 20  
 50 <212> DNA  
 <213> Artificial Sequence  
 55 <220>

## EP 2 402 465 A2

	<223> synthetic oligomer	
	<400> 125	
5	caccacacat tccaggaaga	20
	<210> 126	
	<211> 20	
10	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> synthetic oligomer, probe	
	<400> 126	
20	cttcctggaa tgtgtggtgg	20
	<210> 127	
	<211> 20	
25	<212> RNA	
	<213> Artificial Sequence	
30	<220>	
	<223> synthetic oligomer, probe	
35	<400> 127	
	cuuccaggaa uguguggugg	20
	<210> 128	
40	<211> 22	
	<212> RNA	
	<213> Artificial Sequence	
45	<220>	
	<223> synthetic oligomer, probe	
50	<400> 128	
	gcugcaggaa aaauaaucuu gg	22
	<210> 129	
55	<211> 26	

## EP 2 402 465 A2

	<212> DNA	
	<213> Artificial Sequence	
5		
	<220>	
	<223> synthetic oligomer, probe	
10	<400> 129 tggagaaaga gaugauagaa uauagg	26
	<210> 130	
15	<211> 24	
	<212> RNA	
	<213> Artificial Sequence	
20		
	<220>	
	<223> synthetic oligomer, probe	
25	<400> 130 ggagaaugag auuuuauagac cagu	24
	<210> 131	
30	<211> 24	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
40	<223> synthetic oligomer	
	<400> 131 actggtcata aaatctcatt ctcc	24
45	<210> 132	
	<211> 28	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
55	<223> synthetic oligomer, primer	

	<220>	
	<221> Cla I site	
5	<222> (3)..(8)	
	<223>	
10	<220>	
	<221> Cla I recognition site	
	<222> (3)..(8)	
15	<223>	
20	<400> 132 ccatcgatgc gttttggaga ctacattc	28
	<210> 133	
25	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> synthetic oligomer, primer	
35	<220>	
	<221> Pst I recognition site	
	<222> (4)..(9)	
40	<223>	
45	<400> 133 aaactgcaga tgaaggttcc tacaattcc	29
	<210> 134	
50	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	

## EP 2 402 465 A2

<223> synthetic oligomer, primer  
 <220>  
 5 <221> Kpn I recognition site  
 <222> (4)..(9)  
 <223>  
 10  
 <400> 134  
 cggggtaccg cgttttggag actacattc 29  
 15 <210> 135  
 <211> 29  
 <212> DNA  
 20 <213> Artificial Sequence  
 <220>  
 25 <223> synthetic oligomer, primer  
 <220>  
 30 <221> Pst I recognition site  
 <222> (4)..(9)  
 <223>  
 35  
 <400> 135  
 aaactgcaga gagtggaac ttactattc 29  
 40 <210> 136  
 <211> 38  
 <212> RNA  
 45 <213> Artificial Sequence  
 <220>  
 50 <223> synthetic oligomer  
 <400> 136  
 uuaagccuau auucuaucan cucuuucucc aaacagga 38  
 55 <210> 137

## EP 2 402 465 A2

<211> 27  
 <212> DNA  
 5 <213> Artificial Sequence  
  
 <220>  
 10 <223> synthetic oligomer  
 <400> 137  
 gccccaccac acauuccagg aagacct 27  
 15  
 <210> 138  
 <211> 42  
 <212> DNA  
 20 <213> Artificial Sequence  
  
 <220>  
 25 <223> synthetic oligomer  
 <400> 138  
 gcaaggggag agccctggaa gaaagaagac agaaagcgtg aa 42  
 30  
 <210> 139  
 <211> 8  
 35 <212> DNA  
 <213> Artificial Sequence  
  
 40 <220>  
 <223> synthetic oligomer  
 <400> 139  
 45 gaagaaag 8  
  
 <210> 140  
 <211> 21  
 50 <212> DNA  
 <213> Artificial Sequence  
 55  
 <220>

## EP 2 402 465 A2

	<223> synthetic oligomer	
	<400> 140	
5	gaagaaagaa gacagaaagc g	21
	<210> 141	
	<211> 42	
10	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> synthetic oligomer	
	<400> 141	
20	ccatggtgag gggacttgat acctcaccgc cgtttgccta gg	42
	<210> 142	
	<211> 9	
25	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> synthetic oligomer	
35	<400> 142	
	cttgatacc	9
	<210> 143	
40	<211> 11	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> synthetic oligomer	
50	<400> 143	
	gacLLgalac c	11
	<210> 144	
55	<211> 22	



## EP 2 402 465 A2

<212> DNA

<213> Artificial Sequence

5

<220>

<223> synthetic oligomer

10

<400> 144  
cttgatacct caccgccgtt tg 22

<210> 145

15

<211> 19

<212> DNA

<213> Artificial Sequence

20

<220>

<223> synthetic oligomer

25

<400> 145  
gatacctcac cgcggtttg 19

<210> 146

30

<211> 17

<212> DNA

<213> Artificial Sequence

35

<220>

40

<223> synthetic oligomer

<400> 146  
cttgatacct caccgcc 17

<210> 147

45

<211> 28

<212> DNA

50

<213> Artificial Sequence

<220>

55

<223> synthetic oligomer

## EP 2 402 465 A2

	<400> 147 cgatcaattg ctccctaac ataaactg	28
5	<210> 148 <211> 22 <212> DNA	
10	<213> Artificial Sequence	
	<220>	
15	<223> synthetic oligomer	
	<400> 148 caattgcttc cttaacataa ac	22
20	<210> 149 <211> 31 <212> DNA	
25	<213> Artificial Sequence	
	<220>	
30	<223> synthetic oligomer	
	<400> 149 ccacaagaat tgaatttccct ccagcaacat g	31
35	<210> 150 <211> 19 <212> DNA	
40	<213> Artificial Sequence	
	<220>	
45	<223> synthetic oligomer	
	<400> 150 gaattgaatt tctccagc	19
50	<210> 151 <211> 58 <212> DNA	
55		

## EP 2 402 465 A2

<213> Artificial Sequence

5	<220>		
	<223>	synthetic oligomer	
	<400>	151	
10		caaatcatga aaggtcacaa atgaaacact ggtcataaaa tctcattctc caccaatc	58
	<210>	152	
	<211>	20	
15	<212>	DNA	
	<213>	Artificial Sequence	
20	<220>		
	<223>	synthetic oligomer	
	<400>	152	
25		ggtcacaaat gaaacactgg	20
	<210>	153	
	<211>	23	
30	<212>	DNA	
	<213>	Artificial Sequence	
35	<220>		
	<223>	synthetic oligomer	
	<400>	153	
40		aaaggtcaca aatgaaacac tgg	23
	<210>	154	
45	<211>	12	
	<212>	DNA	
	<213>	Artificial Sequence	
50	<220>		
	<223>	synthetic oligomer	
55	<400>	154	

	gaaacacLgg Lc	12
5	<210> 155 <211> 21 <212> DNA <213> Artificial Sequence	
10	<220> <223> synthetic oligomer	
15	<400> 155 ggtcataaaa tctcattctc c	21
20	<210> 156 <211> 24 <212> DNA <213> Artificial Sequence	
25	<220> <223> synthetic oligomer	
30	<400> 156 cagatagaat gcttggattg tctg	24
35	<210> 157 <211> 32 <212> DNA <213> Artificial Sequence	
40	<220> <223> synthetic oligomer	
45	<400> 157 tctcctttta taatagcaac ttcaaattgg tc	32
50	<210> 158 <211> 23 <212> DNA <213> Artificial Sequence	
55		

# EP 2 402 465 A2

	<220>	
	<223> synthetic oligomer	
5	<400> 158 ttataatagc aacttcaa at tgg	23
	<210> 159	
10	<211> 16	
	<212> DNA	
15	<213> Artificial Sequence	
	<220>	
20	<223> synthetic oligomer	
	<400> 159 ggcatggaga ggaaaa	16
25	<210> 160	
	<211> 17	
	<212> DNA	
30	<213> Artificial Sequence	
	<220>	
35	<223> synthetic oligomer	
	<400> 160 ggcatggaga ggaaaag	17
40	<210> 161	
	<211> 28	
	<212> DNA	
45	<213> Artificial Sequence	
	<220>	
50	<223> synthetic oligomer	
	<400> 161 ctccaaaacg ctttttagaa agagtccc	28
55	<210> 162	

## EP 2 402 465 A2

	<211> 24	
	<212> DNA	
5	<213> Artificial Sequence	
	<220>	
10	<223> synthetic oligomer	
	<400> 162	
	ccaaaacgct ttttagaaag agtc	24
15	<210> 163	
	<211> 30	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
25	<223> synthetic oligomer	
	<400> 163	
	cgctgagttt gagcagaatt tagaaaatgc	30
30	<210> 164	
	<211> 23	
35	<212> DNA	
	<213> Artificial Sequence	
40	<220>	
	<223> synthetic oligomer	
	<400> 164	
45	ctgagtttga gcagaattta gaa	23
	<210> 165	
50	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	

## EP 2 402 465 A2

	<223> synthetic oligomer	
5	<400> 165 atgcatggct atgagtttta tcagaaatt	29
	<210> 166	
	<211> 21	
10	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> synthetic oligomer	
20	<400> 166 catggctatg agttttatca g	21
	<210> 167	
25	<211> 22	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> synthetic oligomer	
35	<400> 167 gcatggctat gattttatc ag	22
	<210> 168	
40	<211> 25	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> synthetic oligomer	
50	<400> 168 catggctatg agttttatca gaaat	25
	<210> 169	
55	<211> 58	

## EP 2 402 465 A2

<212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 <223> synthetic oligomer, capture probe  
 10  
 <400> 169  
 gttggctata cctaaagtca tgaatcctaa aaaaaaaaaa aaaaaaaaaa aaaaaaaa 58  
 <210> 170  
 15  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> synthetic oligomer, primer  
 25  
 <400> 170  
 cccctagaaa acccatcctc t 21  
 30  
 <210> 171  
 <211> 52  
 <212> DNA  
 35  
 <213> Artificial Sequence  
 <220>  
 40  
 <223> synthetic oligomer, promoter primer  
 <220>  
 <221> promoter  
 45  
 <222> (1)..(27)  
 <223>  
 50  
 <400> 171  
 aatttaatac gactcactat agggagaagt accgggtagt tgtacgctaa ct 52  
 55  
 <210> 172  
 <211> 25



<212> DNA  
 <213> Artificial Sequence  
 5  
 <220>  
 10 <223> synthetic oligomer, primer  
 <400> 172  
 agtaccgggt agttgtacgc taact 25  
 15  
 <210> 173  
 <211> 20  
 20 <212> RNA  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> synthetic oligomer, probe  
 30 <400> 173  
 gucauggaca guuauucugac 20  
 35  
 <210> 174  
 <211> 22  
 <212> RNA  
 40 <213> Artificial Sequence  
 45  
 <220>  
 <223> synthetic oligomer, probe  
 <400> 174  
 50 guauuauca gugaagacuu ac 22

# Claims

- 55 1. A method of detecting the presence of HAV in a sample comprising the steps of:  
 (a) purifying HAV nucleic acid from other components in a sample containing HAV by contacting the sample with a capture oligomer that binds specifically to a sequence in the HAV genome and separating the complex

made up of the capture oligomer and the bound HAV RNA from other sample components by using a capture support, wherein the target capture oligomer is selected from the group consisting of SEQ ID NOS:1-7 and SEQ ID NOS:8-14 linked to a sequence or moiety that binds to an immobilized probe;

(b) amplifying a HAV target sequence in the purified HAV nucleic acid, or a cDNA made therefrom, by using an *in vitro* amplification reaction that includes one or more amplification oligomers specific for a selected HAV target region to produce an amplified product of the selected HAV target region; and

(c) detecting the amplified product by using a detection probe oligomer that hybridizes specifically with at least a portion of the amplified product and detecting a signal that results from the probe oligomer bound to the amplified sequence.

2. A method of detecting the presence of HAV in a sample comprising the steps of:

(a) purifying HAV nucleic acid from other components in a sample containing HAV;

(b) amplifying a HAV target sequence in the purified HAV nucleic acid, or a cDNA made therefrom, by using an *in vitro* amplification reaction that includes one or more amplification oligomers specific for a selected HAV target region to produce an amplified product of the selected HAV target region; and

(c) detecting the amplified product by using a detection probe oligomer that hybridizes specifically with at least a portion of the amplified product and detecting a signal that results from the probe oligomer bound to the amplified sequence, wherein the detection probe is selected from the group consisting of SEQ ID NO: 113, SEQ ID NO: 115, SEQ ID NO:117, SEQ ID NO:119, SEQ ID NO: 121 to SEQ ID NO: 124, and SEQ ID NO: 126 to SEQ ID NO:130.

3. The method of claim 1 or claim 2, wherein the at least two amplification oligomers are selected from:

(a) for a first HAV target region, oligomers of about 23 to 26 nt contained in the sequence of SEQ ID NO: 138 that include at least the sequence of SEQ ID NO: 139 or SEQ ID NO:140, or oligomers in a size range of about 19 to 25 nt contained in the sequence of SEQ ID NO: 141 that contain at least one sequence of SEQ ID Nos. 142 to 146, or promoter primer oligomers in a size range of about 50 to 53 nt that include HAV target-specific portions of any one of SEQ ID Nos. 21 to 27;

(b) for a second HAV target region, oligomers of about 21 to 27 nt contained in the sequence of SEQ ID NO: 60 or contained in the sequence of SEQ ID NO: 86 that include at least the sequence of SEQ ID NO: 156, or promoter primer oligomers in a size range of about 48 to 54 nt that include HAV target-specific portions of any one of SEQ ID Nos. 29 to 32;

(c) for a third HAV target region, oligomers of about 24 to 30 nt contained in the sequence of SEQ ID NO: 147 that include at least the sequence of SEQ ID NO: 148, or contained in the sequence of SEQ ID NO: 157 that include at least the sequence of SEQ ID NO: 158, or promoter primer oligomers that include HAV target-specific portions of SEQ ID NO:31 or SEQ ID NO:32;

(d) for a fourth HAV target region, oligomers of about 18 to 27 nt contained in the sequence of SEQ ID NO:93 or SEQ ID NO:95 that contain at least the sequence of SEQ ID NO:97, SEQ ID NO:159, or SEQ ID NO: 160, or a promoter primer oligomer that includes a HAV target-specific portion of SEQ ID NO:33;

(e) for a fifth HAV target region, oligomers of about 19 to 31 nt contained in the sequence of SEQ ID NO: 149 that include at least the sequence of SEQ ID NO: 150, or promoter primer oligomers in a size range of about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 34 to 40;

(f) for a sixth HAV target region, oligomers of about 24 to 28 nt contained in the sequence of SEQ ID NO: 161 that include at least the sequence of SEQ ID NO: 162, or promoter primer oligomers are embodiments of promoter primers that include HAV target-specific portions of SEQ ID NO:41 or SEQ ID NO:42; and

(g) for a seventh HAV target region, oligomers of about 20 to 30 nt contained in the sequence of SEQ ID NO: 151 that include at least any one of the sequences of SEQ ID NO:152 to SEQ ID NO:155, or contained in SEQ ID NO:163 that include at least the sequence of SEQ ID NO:164, or contained in SEQ ID NO:165 that include at least any one of the sequences of SEQ ID Nos. 166 to 168, or promoter primer oligomers in a size range of about 51 to 56 nt that include HAV target-specific portions of any one of SEQ ID Nos. 43 to 49.

4. The method of claim 1 or claim 2, wherein

(a) the amplifying step amplifies a sequence in the first HAV target region by using at least two oligomers specific for the first HAV target region selected from the group consisting of: SEQ NO:15, SEQ ID NO:16, SEQ ID NO: 17, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ

ID NO:54, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:80, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:83, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:143, SEQ ID NO:144, and SEQ ID NO:145; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the first HAV target region;

(b) the amplifying step amplifies a sequence in the second HAV target region by using at least two oligomers specific for the second HAV target region selected from the group consisting of: SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:86, SEQ ID NO:87, SEQ ID NO:88, and SEQ ID NO:156; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the second HAV target region;

(c) wherein the amplifying step amplifies a sequence in the third HAV target region by using at least two oligomers specific for the third HAV target region selected from the group consisting of: SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:89, SEQ ID NO:90, SEQ ID NO:91, and SEQ ID NO:148; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the third HAV target region;

(d) wherein the amplifying step amplifies a sequence in the fourth HAV target region by using at least two oligomers specific for the fourth HAV target region selected from the group consisting of: SEQ ID NO:33, SEQ ID NO:63, SEQ ID NO:92, SEQ ID NO:93, SEQ ID NO:94, SEQ ID NO:95, SEQ ID NO:96, and SEQ ID NO:97; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the fourth HAV target region;

(e) wherein the amplifying step amplifies a sequence in the fifth HAV target region by using at least two oligomers specific for the fifth HAV target region selected from the group consisting of: SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:97, SEQ ID NO:149, and SEQ ID NO:150; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the fifth HAV target region;

(f) wherein the amplifying step amplifies a sequence in the sixth HAV target region by using at least two oligomers specific for the sixth HAV target region selected from the group consisting of: SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:98, SEQ ID NO:99, SEQ ID NO:101, SEQ ID NO:161, and SEQ ID NO:162; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the sixth AV target region; and

(g) wherein the amplifying step amplifies a sequence in the seventh V target region by using at least two oligomers specific for the seventh HAV target region selected from the group consisting of: SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:102, SEQ ID NO:103, SEQ ID NO:104, SEQ ID NO:104, SEQ ID NO:106, SEQ ID NO:107, SEQ ID NO:108, SEQ ID NO:152, SEQ ID NO:153, SEQ ID NO:155, SEQ ID NO:163, SEQ ID NO:164, SEQ ID NO:165, SEQ ID NO:166, SEQ ID NO:167, and SEQ ID NO:168; and wherein the detecting step uses at least one detection probe that hybridizes specifically to the amplified product of the seventh HAV target region.

5. The method of claim 1 or claim 2 wherein the sample is a biological sample derived from humans, selected from the group consisting of blood samples or plasma samples.

6. The method of claim 1 or 2, wherein the purifying step (a) comprises mixing the sample or an aliquot of the sample with a target capture reagent containing one or more capture oligomers, magnetic particles with attached immobilized probes complementary to a portion of the capture oligomers, and salt compounds to provide hybridization conditions.

7. The method of claim 6, wherein the target capture reagent further comprises a detergent or other chaotropic agent that disrupts the HAV particles and releases HAV RNA for hybridization with capture oligomers.

8. The method of claim 6 or 7, wherein the mixture is incubated at 60°C to allow hybridization of the target-specific portion of the capture oligomer to the HAV target sequence and then at room temperature to allow binding of the capture oligomer and the immobilized probe.

9. The method of any one of claims 6-8, wherein the target capture reagent contains magnetic particles with covalently attached poly-dT<sub>14</sub> oligomer as an immobilized probe.

10. The method of any one of claims 6-9, wherein the hybridization complexes of HAV RNA with the target capture oligomer, which is in turn hybridized with the immobilized probe which are attached to the magnetic particles are

separated from other sample components by use of a magnetic field and are then optionally mixed with a wash buffer, and then separated from the wash buffer.

- 5      **11.** The method of claim 1 or claim 2, wherein the detecting step is an endpoint detecting step and the detection probe oligomers are labeled with a chemiluminescent compound.
12. The method of claim 11, wherein the labeled detection probes are contacted with a detection reaction mixture containing a detection reagent.
- 10      **13.** The method of any one of claims 1 to 12, wherein the presence of an additional viral target in the sample is detected.
- 14.** The method of claim 13, wherein the HAV RNA and the additional viral target are amplified in the same amplification reaction mixture and/or. wherein the detection step uses different probes for the different target viruses.
- 15      **15.** A kit containing at least one reagent useful for performing any of the method steps of claims 1 - 14.

20

25

30

35

40

45

50

55

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- US 5585481 A [0014] [0028]
- WO 9532305 A, Hydig-Hielsen [0015]
- US 6110678 A [0020] [0034] [0046]
- US 6280952 B, Weisburg [0020]
- US 6910878 B [0021]
- US 4786600 A, Kramer [0025]
- US 4683195 A [0025]
- US 4683242 A [0025]
- US 4800159 A, Mullis [0025]
- US 5427930 A, Biekenmeyer [0025]
- US 5494810 A, Barany [0025]
- US 5422252 A, Walker [0025]
- US 5399491 A [0026] [0027] [0047]
- US 5554516 A, Kacian [0026] [0047]
- US 5437990 A, Burg [0026]
- WO 8809302 A [0026]
- WO 8810315 A, Gingeras [0026]
- US 5130238 A, Malek [0026]
- US 4868105 A [0026]
- US 5124246 A, Urdea [0026]
- US 5554596 A [0027]
- US 6361945 B, Becker [0028]
- US 5658737 A, Nelson [0028] [0029] [0048] [0063]
- US 5656207 A, Woodhead [0028] [0029]
- US 5547842 A, Hogan [0028]
- US 5283174 A, Arnold [0028] [0029] [0043] [0048] [0052] [0053]
- US 4581333 A, Kourilsky [0028]
- US 5731148 A, Becker [0028]
- US 5639604 A, Arnold [0028]
- US 5185439 A [0028] [0053]
- US 5656744 A [0048]
- US 20030124578 A1 [0063]

## Non-patent literature cited in the description

- **DATA et al.** *Morbidity Mortality Wkly. Rpt.*, 2003, vol. 52 (47), 1155-57 [0004]
- **LAPORTE et al.** *Morbidity Mortality Wkly. Rpt.*, 2003, vol. 52 (24), 565-67 [0004]
- **CARUTHERS et al.** *Methods in Enzymol.*, 1987, vol. 154, 287 [0014]
- **ESCHENMOSER.** *Science*, 1999, vol. 284, 2118-2124 [0015]
- **MAJLESSI et al.** *Nucl. Acids Res.*, 1998, vol. 28 (9), 2224-2229 [0015]
- **DEMPCY et al.** *Proc. Natl. Acad. Sci. USA*, 1995, vol. 92, 6497-6101 [0015]
- **BROWNE et al.** *Proc. Natl. Acad. Sci. USA*, 1995, vol. 92, 7051-7055 [0015]
- **ARYA ; BRUICE.** *J. Am. Chem. Soc.*, 1998, vol. 120, 6619-6620 [0015]
- **REYNOLDS et al.** *Nucl. Acids Res.*, 1996, vol. 24 (22), 4584-4591 [0015]
- **GRYAZNOV ; CHEN.** *Am. Chem. See.*, 1994, vol. 116, 3143-3144 [0015]
- **CHATURVEDI et al.** *Nucl. Acids Res.*, 1996, vol. 24 (12), 2318-2323 [0015]
- **HYRUP ; NIELSEN.** *Bioorg. & Med. Chem.*, 1996, vol. 4, 5-23 [0015]
- **MESMAEKER et al.** *Syn. Lett.*, 1997, 1287-1290 [0015]
- **PEYMAN et al.** *Angew. Chem. Int Ed. Engl.*, 1996, vol. 35 (22), 2636-2638 [0015]
- **AERSCHOT et al.** *Angew. Chem. Int. Ed. Engl.*, 1995, vol. 34 (12), 1338-1339 [0015]
- **KOSHKIN et al.** *J. Am. Chem. Soc.*, 1998, vol. 120, 13252-13253 [0015]
- **STEFFENS ; LEUMANN.** *J. Am. Chem. Soc.*, 1997, vol. 119, 11548-11549 [0015]
- **JONES et al.** *J. Org. Chem.*, 1993, vol. 58, 2983-2991 [0015]
- **SUMMERTON ; WELLER.** *Antisense & Nucl. Acid Drug Dev.*, 1997, vol. 7, 187-195 [0015]
- **STIRCHAK et al.** *Nucl. Acids Res.*, 1989, vol. 17 (15), 6129-6141 [0015]
- **SAMBROOK et al.** *Molecular Cloning, A Laboratory Manual.* Cold Spring Harbor Laboratory, 1989, 9.50-5111.46-4911.55-57 [0017]
- *Dictionary of Microbiology and Molecular Biology.* John Wiley & Sons, 1994 [0031]
- *The Encyclopedia of Molecular Biology.* Blackwell Science Ltd, 1994 [0031]
- *The Harper Collins Dictionary of Biology.* Hale & Marham, 1991 [0031]
- **BENEDUCE et al.** *Virus Res.*, 1995, vol. 36 (2-3), 299-309 [0032]
- **FUJIWARA et al.** *J. Hepatol.*, 2001, vol. 35 (1), 112-119 [0032]
- **HU et al.** *Acta Virol.*, 2002, vol. 46 (3), 153-157 [0032]
- **LIN ; BROWN.** *Nucl. Acids Res.*, 1989, vol. 17, 10373-83 [0050]

## EP 2 402 465 A2

- **LIN ; BROWN.** *Nucl. Acids Res.*, 1992, vol. 20, 5149-52 [0050]
- **CARUTHERS et al.** *Methods in Enzymol.*, 1997, vol. 154, 287 [0053]