



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2944962 A1 2015/11/05

(21) **2 944 962**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) **Date de dépôt PCT/PCT Filing Date:** 2015/04/30
(87) **Date publication PCT/PCT Publication Date:** 2015/11/05
(85) **Entrée phase nationale/National Entry:** 2016/10/05
(86) **N° demande PCT/PCT Application No.:** EP 2015/059562
(87) **N° publication PCT/PCT Publication No.:** 2015/166073
(30) **Priorité/Priority:** 2014/04/30 (EP14166729.5)

(51) **Cl.Int./Int.Cl.** **C12N 15/13** (2006.01),
A61K 39/395 (2006.01), **A61P 35/00** (2006.01),
A61P 37/06 (2006.01), **C07K 16/28** (2006.01),
C07K 16/44 (2006.01), **C12N 5/10** (2006.01)
(71) **Demandeur/Applicant:**
MAX-DELBRUCK-CENTRUM FUR MOLEKULARE
MEDIZIN IN DER HELMHOLTZ-GEMEINSCHAFT,
DE
(72) **Inventeurs/Inventors:**
ODEN, FELIX, DE;
MARINO, STEPHEN, DE;
DAUMKE, OLIVER, DE
(74) **Agent:** CAMERON IP


(54) **Titre : ANTICORPS HUMANISES DIRIGES CONTRE CD269 (BCMA)**
(54) **Title: HUMANIZED ANTIBODIES AGAINST CD269 (BCMA)**

(57) **Abrégé/Abstract:**

The invention relates to humanized antibodies or antibody fragments that bind CD269 (BCMA), thereby disrupting the interaction between CD269 and its native ligands (BAFF and APRIL), and their use in the treatment of plasma cell-mediated diseases such as multiple myeloma and autoimmune diseases.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)
(19) World Intellectual Property
Organization
International Bureau


WIPO | PCT



(43) International Publication Date
5 November 2015 (05.11.2015)

WIPO | PCT

(10) International Publication Number


WO 2015/166073 A1

- (51) International Patent Classification:
C07K 16/28 (2006.01) A61K 39/00 (2006.01)
- (21) International Application Number:
PCT/EP2015/059562
- (22) International Filing Date:
30 April 2015 (30.04.2015)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
14166729.5 30 April 2014 (30.04.2014) EP
- (71) Applicant: MAX-DELBRÜCK-CENTRUM FÜR MOLEKULARE MEDIZIN [DE/DE]; Robert-Rössle-Str. 10, 13125 Berlin (DE).
- (72) Inventors: ODEN, Felix; Kärntener Straße 28, 10827 Berlin (DE). MARINO, Stephen; Katzbachstrasse 3, 10965 Berlin (DE). DAUMKE, Oliver; Sredzkistrasse 19, 10435 Berlin (DE).
- (74) Agent: BUCHANAN, Luke; HERTIN und Partner PartG mbB, Rechts- und Patentanwälte, Kurfürstendamm 54/55, 10707 Berlin (DE).

- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:
- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: HUMANIZED ANTIBODIES AGAINST CD269 (BCMA)

(57) Abstract: The invention relates to humanized antibodies or antibody fragments that bind CD269 (BCMA), thereby disrupting the interaction between CD269 and its native ligands (BAFF and APRIL), and their use in the treatment of plasma cell-mediated diseases such as multiple myeloma and autoimmune diseases.



WO 2015/166073 A1

HUMANIZED ANTIBODIES AGAINST CD269 (BCMA)

DESCRIPTION

The invention relates to humanized antibodies or antibody fragments that bind CD269 (BCMA), thereby disrupting the interaction between CD269 and its native ligands (BAFF and APRIL), and their use in the treatment of plasma cell-mediated diseases such as multiple myeloma and autoimmune diseases.

BACKGROUND OF THE INVENTION

The B cell maturation antigen (BCMA) is member 17 of the tumor necrosis factor receptor superfamily (TNFRSF). Its native ligands are the B cell activating factor (BAFF; also called BLyS or TALL-1, TNFSF13B) and a proliferation-inducing ligand (APRIL, TNFSF13, CD256) which are involved (through interaction with further ligands) in regulating various aspects of humoral immunity, B cell development, and homeostasis.

BCMA is highly expressed on malignant plasma cells, for example in multiple myeloma, (MM), which is a B cell non-Hodgkin lymphoma of the bone marrow, and plasma cell leukemia (PCL), which is more aggressive than MM and constitutes around 4% of all cases of plasma cell disorders. In addition to MM and PCL, BCMA has also been detected on Hodgkin and Reed-Sternberg cells in patients suffering from Hodgkin's lymphoma (Chiu et al. (2007) Blood 109:729-739). Similar to its function on plasma cells, ligand binding to BCMA has been shown to modulate the growth and survival of multiple myeloma cells expressing BCMA (Novak et al. (2004) Blood 103:689-694). Signalling of BAFF and APRIL via BCMA are considered as pro-survival factors for malignant plasma cells; hence, the depletion of BCMA-positive tumour cells and/or the disruption of ligand-receptor interaction should improve the therapeutic outcome for multiple myeloma and autoantibody-dependent autoimmune diseases.

There are presently various approaches available for the treatment of multiple myeloma (Raab et al. (2009) Lancet 374:324-339). Chemotherapy leads in most subjects only to partial control of multiple myeloma; only rarely does chemotherapy lead to complete remission. Combination approaches are therefore often applied, commonly involving an additional administration of corticosteroids, such as dexamethasone or prednisone. Corticosteroids are however plagued by side effects, such as reduced bone density. Stem cell transplantation has also been proposed, using one's own stem cells (autologous) or using cells from a close relative or matched unrelated donor (allogeneic). In multiple myeloma, most transplants performed are of the autologous kind. Such transplants, although not curative, have been shown to prolong life in selected patients (Suzuki (2013) Jpn J Clin Oncol 43:116-124). Alternatively thalidomide and derivatives thereof have recently been applied in treatment but are also associated with sub-optimal success rates and high costs. More recently, the proteasome inhibitor bortezomib (PS-341) has been approved for the treatment of relapsed and refractory MM and was used in numerous clinical trials alone or in combination with established drugs resulting in an encouraging clinical outcome (Richardson et al. (2003) New Engl J Med 348:2609-2617;

Kapoor *et al.* (2012) *Semin Hematol* 49:228-242). The costs for combined treatments are correspondingly high and success rates still leave significant room for improvement. The combination of treatment options is also not ideal due to an accumulation of side effects if multiple medicaments are used simultaneously. Novel approaches for the treatment of plasma cell diseases, in particular multiple myeloma, are required.

The ability to specifically target plasma cells is also of great benefit for the treatment of autoimmune diseases. Mild forms of autoimmune disease are usually initially treated with nonsteroidal anti-inflammatory drugs (NSAID) or disease-modifying anti-rheumatic drugs (DMARD). More severe forms of Systemic Lupus Erythematosus (SLE), involving organ dysfunction due to active disease, usually are treated with steroids in conjunction with strong immunosuppressive agents such as cyclophosphamide, a cytotoxic agent that targets cycling cells. Only recently Belimumab, an antibody targeting the cytokine BAFF, which is found at elevated levels in serum of patients with autoimmune diseases, received approval by the Food and Drug Administration (FDA) for its use in SLE. However, only newly formed B cells rely on BAFF for survival in humans, whereas memory B cells and plasma cells are less susceptible to selective BAFF inhibition (Jacobi *et al.* (2010) *Arthritis Rheum* 62:201-210). For rheumatoid arthritis (RA), TNF inhibitors were the first licensed biological agents, followed by abatacept, rituximab, and tocilizumab and others: they suppress key inflammatory pathways involved in joint inflammation and destruction, which, however, comes at the price of an elevated infection risk due to relative immunosuppression (Chan *et al.* (2010) *Nat Rev Immunol* 10:301-316, Keyser (2011) *Curr Rheumatol Rev* 7:77-87). Despite the approval of these biologicals, patients suffering from RA and SLE often show a persistence of autoimmune markers, which is most likely related to the presence of long-lived, sessile plasma cells in bone marrow that resist e.g. CD20-mediated ablation by rituximab and high dosage glucocorticoid and cyclophosphamid therapy.

Antibodies which bind CD269 (BCMA) and their use in the treatment of various B-cell related medical disorders have been described in the art. Ryan *et al* (*Molecular Cancer Therapeutics*, 2007 6(11), 3009) describe an anti-BCMA antibody obtained via vaccination in rats using a peptide of amino acids 5 to 54 of the BCMA protein. WO 2012/163805 describes BCMA binding proteins, such as chimeric and humanized antibodies, their use to block BAFF and/or APRIL interaction with BCMA and their potential use in treating plasma cell malignancies such as multiple myeloma. The antibody disclosed therein was obtained via vaccination in mouse using a recombinant peptide of amino acids 4 to 53 of the BCMA protein. WO 2010/104949 also discloses various antibodies that bind preferably the extracellular domain of BCMA and their use in treating B cell mediated medical conditions and disorders. WO 2002/066516 and WO 2012/066058 disclose bivalent antibodies that bind both BCMA and additional targets and their potential use in the treatment of B cell related medical disorders. Details regarding the binding properties and specific epitopes of the bivalent antibodies are not provided in either disclosure.

40 SUMMARY OF THE INVENTION

In light of the prior art the technical problem underlying the invention was the provision of an agent suitable for treating human diseases associated with pathogenic plasma cells, such as

multiple myeloma and autoimmune diseases. This problem is solved by the features of the independent claims. Preferred embodiments of the present invention are provided by the dependent claims.

Therefore, an object of the invention is to provide a humanized antibody or antibody fragment
5 that binds CD269 (BCMA), in particular an epitope of the extracellular domain of CD269 (BCMA).

The antibodies disclosed herein comprise humanized sequences, especially of the preferred VL and VH binding regions, which maintain the appropriate ligand affinities as described with respect to the chimeric antibody J22.9-xi.

10 In various embodiments of the invention the amino acid sequence modification to obtain said humanized sequences may occur in either the CDR regions of the original chimeric antibody J22.9-xi or in the framework regions, wherein the framework region is to be understood as a region in the variable domain of a protein which belongs to the immunoglobulin superfamily, and which is less "variable" than the CDRs.

15 It was entirely surprising that the particular humanized sequences provided herein, preferably the CDR regions of the VL and VH regions involved in binding, exhibit the specific and strong binding as demonstrated in the experimental examples, and maintain the binding characteristics of the original chimeric antibody J22.9-xi to such an extent as to maintain their desired therapeutic effect.

20 The invention therefore relates to an antibody or antibody fragment, comprising a VH domain that comprises CDR sequences of:

- RYWX₁S (H-CDR1; SEQ ID NO. 15), wherein X₁: I, F, L, V, Y, C, G, A, S, T);
- EINPX₂X₃STINYAPSLKDK (H-CDR2; SEQ ID No. 16), wherein X₂X₃: SS, NS, TS, GS, KS, RS, SD, SN, DE; and/or
- 25 – SLYX₄DYGDAX₅DYW (H-CDR3; SEQ ID NO. 17), wherein X₄: Y, L, A, V, F, I, W, and/or X₅: Y, L, F, I, V, A, C,

wherein said antibody or fragment thereof specifically binds an epitope of the extracellular domain of CD269 (BCMA).

30 It was particularly surprising that the humanized antibodies described herein, that exhibit sequence changes compared to the original chimeric antibody, in particular sequence changes in the CDRs of said chimeric antibody, maintain sufficient binding properties towards their target for therapeutic efficacy.

A skilled person would not have expected that the binding characteristics of the humanized variants would be similar to the original chimeric or mouse antibody. Considering the
35 sequence changes in the variable domains, in particular the CDRs, the beneficial binding characteristics of the humanized sequences demonstrated herein are considered a surprising technical effect. The comparison in binding characteristics between the partially and fully

humanized antibodies also shows improvements in the fully humanized sequences. This represents an entirely unexpected result. The initial modifications to the chimera (partially humanized) led to some loss in binding affinity. However, the introduction of further humanizations subsequently lead to enhanced binding, whereby the "fully humanized" sequences show similar binding properties compared to the original chimera, thereby showing a surprising technical effect after having made such significant sequence modifications without severe loss of binding affinity.

In a preferred embodiment the antibody or antibody fragment as described herein is characterised in that the VH domain comprises the CDR sequence RYWIS (SEQ ID NO. 18) or RYWFS (SEQ ID NO. 19).

In a preferred embodiment the antibody or antibody fragment as described herein is characterised in that said VH domain comprises the CDR sequence EINPNSSTINYAPSLKDK (SEQ ID No. 20) or EINPSSSTINYAPSLKDK (SEQ ID No. 21).

In further embodiments of the invention amino acid 54 of the VH domain may relate to any given amino acid or modified amino acid. As shown in the examples below, potential glycosylation of the N amino acid at this residue does not significantly disrupt specific and strong binding of the antibody to the target epitope. In light of this information, the invention relates to an antibody or antibody fragment comprising an CDR2 sequence as described herein, wherein any given amino acid or modified amino acid may be present at amino acid 54 of the VH domain in the CDR2 sequence.

In a preferred embodiment the antibody or antibody fragment as described herein is characterised in that said VH domain comprises the CDR sequence SLYYDYGDYDYW (SEQ ID NO. 22).

The invention further relates to an antibody or antibody fragment, comprising a VL domain that comprises CDR sequences of:

- KASQSVX₁X₂NVA (L-CDR1; SEQ ID NO. 23), wherein X₁X₂: ES, SS, TS, QS, HS, DH;
- SASLRFS (L-CDR2; SEQ ID NO 24); and/or
- QQYNNYPLTFG (L-CDR3; SEQ ID NO. 25),

wherein said antibody or fragment thereof specifically binds an epitope of the extracellular domain of CD269 (BCMA).

Also with respect to the LC sequence, it was surprising that the modified sequence of the CDR3 sequence had no significant detrimental effect on binding to the BCMA target.

Furthermore, antibodies of the present invention also show unexpected and beneficial stability characteristics when in solution, both when isolated or purified, or in vitro, and in vivo, post administration, that would not have been expected from the sequence changes carried out to the original chimeric antibody.

In a preferred embodiment the antibody or antibody fragment as described herein is characterised in that the VL domain comprises the CDR sequence KASQSVDSNVA (SEQ ID NO. 26).

- 5 In a preferred embodiment the antibody or antibody fragment as described herein comprises a VH domain that comprises the sequence
 EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWX₁SWVRQAPGKGLVWVGEINPX₂X₃STIN
 YAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASLYX₄DYGDAX₅DYWGQGTLVTVSS
 (SEQ ID NO. 4), wherein X₁: I, F, L, V, Y, C, G, A, S, T; X₂X₃: SS, NS, TS, GS, KS, RS, SD, SN, DE, preferably SS; X₄: Y, L, A, V, F, I, W; and X₅: Y, L, F, I, V, A, C.

- 10 In a preferred embodiment the antibody or antibody fragment as described herein is characterised in that the antibody or fragment comprises a VH domain that comprises the sequence according to SEQ ID NO. 6, SEQ ID NO. 7, SEQ ID NO. 8 or SEQ ID NO. 9.

In a preferred embodiment the antibody or antibody fragment as described herein comprises a VL domain that comprises the sequence

- 15 EIVMTQSPATLSVSPGERATLSCASQSVX₁X₂NVAWYQQKPGQAPRALIYSASLRFSGIPAR
 FSGSGSGTEFTLTISSLQSEDAVYYCQQYNNYPLTFGAGTKLELKR (SEQ ID NO. 12),
 wherein X₁X₂: ES, SS, TS, QS, HS, DH.

In a preferred embodiment the antibody or antibody fragment as described herein comprises a VL domain that comprises the sequence according to SEQ ID NO. 14.

- 20 In a preferred embodiment the antibody or antibody fragment as described herein comprises a VH domain that comprises the sequence according to SEQ ID NO. 6, SEQ ID NO. 7, SEQ ID NO. 8 or SEQ ID NO. 9 and a VL domain that comprises the sequence according to SEQ ID NO. 14.

- 25 The invention further relates to an antibody or antibody fragment as described herein comprising a VH domain, wherein said VH domain comprises a sequence according to
 X₁VQLX₂X₃SGGGLVQPGGSLX₄LSCAASGX₅X₆FX₇X₈YWZ₁SWVRX₉APGKGLEWX₁₀GEINPZ
₂SSTINYAPSLKX₁₁X₁₂FX₁₃ISRDNKNTLYLQMX₁₄X₁₅X₁₆RX₁₇EDTAX₁₈YYCASLYYDYGDAZ₃
 DYWGQGTGX₁₉VTVSS (SEQ ID No. 41), wherein X₁: Q, E; X₂: Q, V; X₃: Q, E; X₄: K, R; X₅: I, F; X₆: D, T; X₇: S, D; X₈: R, D; X₉: R, Q; X₁₀: I, V; X₁₁: D, G; X₁₂: K, R; X₁₃: I, T; X₁₄: S, N;
 30 X₁₅: K, S; X₁₆: V, L; X₁₇: S, A; X₁₈: L, V; X₁₉: S, L;
 and wherein at least one of Z₁: I, F, L, V, Y, C, G, A, S, T, preferably I or F; Z₂: S, N, T, G, K, R, D, preferably S and/or Z₃: Y, L, F, I, V, A, C, preferably Y;
 and wherein said antibody or fragment thereof specifically binds an epitope of the extracellular domain of CD269 (BCMA).

- 35 This embodiment encompasses various humanized antibodies, in particular the VH sequences thereof, all variants defined by the advantageous humanization carried out in the CDRs as described herein.

The invention further relates to an antibody or antibody fragment as described herein comprising a VL domain, wherein said VL domain comprises a sequence according to

DIVMTQSX₁X₂X₃X₄X₅X₆SVGD₇VX₈X₉TCKASQSVESNVAWYQQKPX₁₀QX₁₁PKX₁₂LIX₁₃SX₁₄
 X₁₅LRFSGVPARFX₁₆GSGSGTDFTLTISX₁₇LQSEDX₁₈AX₁₉YX₂₀CQQYNNYPLTFGAGTKLELK
 R (SEQ ID No. 42), wherein X1: Q, P; X2: R, A; X3: F, T; X4: M, L; X5: T, S; X6: T, V; X7: R,
 E; X8: S, T; X9: V, L; X10: R, G; X11: S, A; X12: A, L; X13: F, Y; X14: A, D; X15: S, D; X16: T,
 S; X17: N, S; X18: L, F; X19: E, V; X20: F, Y;

and wherein said antibody or fragment thereof specifically binds an epitope of the extracellular domain of CD269 (BCMA). This embodiment encompasses various humanized antibodies, in particular the VL sequences thereof, all variants defined by the advantageous humanization carried out in the CDRs as described herein.

10 Preferred embodiments regarding humanized antibody variants

As disclosed in detail herein, the sequence of the preferred embodiments of the invention according to J22.9-xi was humanized in order to provide a more compatible reagent for administration in human subjects. Various humanized sequence variants of J22.9-xi have been generated and tested for their binding affinity and specificity to both human and cynomolgus CD269 (BCMA). The results from binding assays demonstrate that the humanized sequences maintain the desired binding properties of the chimeric reagent J22.9-xi. In the below sequences the underlined regions represent the CDRs or putative CDRs.

Preferred embodiments regarding humanized VH variants

Additional information is provided below on the humanized antibody sequence of the present invention.

Chimeric sequence:

HC mouse (SEQ ID No. 1):

QVQLQQSGGGLVQPGGSLKLSCAASGIDFSRYWMSWVRRAPGKGLEWIGEINPDSSTINYAPSLKDKFI
 ISRDNAKNTLYLQMSKVRSEDTALYYCASLYYDYGDAMDYWGQGTSTVTVSS

The HC mouse sequence represents the variable region of the heavy chain (VH) originally developed for the chimeric antibody J22.9-xi, which comprises VL and VH domains obtained from a mouse antibody, capable of binding an epitope of the extracellular domain of CD269 (BCMA), and the VL and VH domains are fused to human CL and CH domains, respectively.

Partially humanized sequences:

30 HC partially humanized (SEQ ID No. 2):

EVQLVESGGGLVQPGGSLRLSCAASGFTFDYWMSWVRQAPGKGLEWVGEINPDSSTINYAPSLKGRFT
 ISRDNAKNTLYLQMNSLRAEDTAVYYCASLYYDYGDAMDYWGQGTSLVTVSS

The HC partially humanized sequence represents a modified amino acid sequence (via amino acid substitutions) in comparison to the chimeric antibody disclosed herein, whereby the VL and VH binding regions have been modified with respect to their sequence to make them more suitable for administration in humans.

Humanized VH sequence:

WO 2015/166073

PCT/EP2015/059562

- 7 -

hHC01 (SEQ ID No. 3)

EVQLV**ES**GGGLVQPGGSLRLSCAASG**FT**FSRYWMSWVR**Q**APGKGLVWVGEINPDSSTINYAPSLKDK**FT**
 ISRDNAKNTLYLQM**NSL**RAEDTAVYYCASLYYDYGDAMDYWGQGT**L**TVSS

Humanized VH sequence with removal of post translational modification motifs:

5 hHC02 (SEQ ID No. 4)

EVQLV**ES**GGGLVQPGGSLRLSCAASGFTFSRYW**X₁**SWVRQAPGKGLVWVGEINP**X₂X₃**STINYAPSLKDK
 FTISRDNAKNTLYLQM**NSL**RAEDTAVYYCASLY**X₄**DYGD**X₅**DYWGQGT**L**TVSS

Wherein:

X₁: I, F, L, V, Y, C, G, A, S, T, preferably I or F;

10 X₂X₃: SS, NS, TS, GS, KS, RS, SD, SN, DE, preferably SS;

X₄: Y, L, A, V, F, I, W, preferably Y; and/or

X₅: Y, L, F, I, V, A, C, preferably Y;

The "hHC01" and "hHC02" humanized sequences represent novel amino acid sequences that
 comprise sequence changes in comparison to both the original chimeric sequence and the
 15 partially humanized sequences described herein.

The PTM mutations are intended to remove potentially detrimental post translational
 modification motifs from said proteins, whilst maintaining the advantageous binding properties.
 The positions 1, 5, 6, 19, 27, 28, 34, 39, 46, 48, 54, 69, 84, 85, 86, 88, 93, 107 and/or 115 of
 hHC01 and hHC02 are preferably mutated (substituted) in comparison to the original chimeric
 20 sequence. The importance of the substitution relates primarily to the resulting amino acid, not
 the originating amino acid. The change may therefore also be carried out from the
 corresponding amino acid of the original chimeric amino acid or other variant, such as the
 partially humanized sequences.

The following substitutions are novel in comparison to the chimeric (SEQ ID No 1) sequence:

- 25
- the amino acid M34 of the HC (VH) sequence is substituted with any amino acid, preferably I, L, F, V, Y, C, G, A, S, T;
 - the amino acid E46 of the HC (VH) sequence is substituted with V;
 - the amino acids D54 and S55 of the HC (VH) sequence is substituted with any amino acid combination, preferably SS, TS, GS, KS, RS, SD, SN, DE;

30

 - the amino acid Y101 of the HC (VH) sequence is substituted with any amino acid, preferably L, A, V, F, I, W; and/or
 - the amino acid M107 of the HC (VH) sequence is substituted with any amino acid, preferably L, Y, F, I, V, A, C.

Sequences that may be modified at those residues required for direct interaction with BCMA:

35 hHC03 – modified amino acids involved in interaction with BCMA (SEQ ID No 5):

WO 2015/166073

PCT/EP2015/059562

- 8 -

EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYX₁MX₂WVRQAPGKGLVX₃VGX₄INPDSSTINYAPSLKDK
 FTISRDNAKNTLYLQMNSLRAEDTAVYYCASX₅X₆X₇DYGDX₈MDYWGQGTLVTVSS

Wherein preferred amino acids are:

- X₁: W, F, Y, preferred W;
- 5 X₂: S, T, N, Q, D, E, preferred S;
- X₃: W, F, Y, preferred W;
- X₄: E, Q, preferred E;
- X₅: L, I, V, G, A, preferred L;
- X₆: Y, X, preferred Y;
- 10 X₇: Y, F, L, I, V, M, preferred Y; and/or
- X₈: A, G, V, preferred A.

The "hHC03" humanized sequence represents novel amino acid sequences that comprise amino acid sequence changes in comparison to both the original chimeric sequence and the partially humanized sequence. These sequence changes are intended to reflect potential changes in the amino acids that bind the BCMA target, which may be substituted, whilst maintaining the advantageous binding properties. The importance of the substitution relates primarily to the resulting amino acid, not the originating amino acid. The change may therefore also be carried out from the corresponding amino acid of the original chimeric amino acid or other variant.

20 For example:

- the amino acid W33 of the HC (VH) sequence is W, F, Y;
- the amino acid S35 of the HC (VH) sequence is S, T, N, Q, D, E;
- the amino acid W47 of the HC (VH) sequence is W, F, Y;
- the amino acid E50 of the HC (VH) sequence is E, Q;
- 25 - the amino acid L99 of the HC (VH) sequence is L, I, V, G, A;
- the amino acid Y100 of the HC (VH) sequence is Y, X;
- the amino acid Y101 of the HC (VH) sequence is Y, F, L, I, V, M; and/or
- the amino acid A106 of the HC (VH) sequence is A, G, V.

In general, any change to a CDR region made during humanization may also be considered as a feature of a CDR sequence when considered independently of the framework sequence as a whole. Such modified CDR sequences may be considered defining features of the present invention, either within or independent of their context in the entire framework region described herein. For example, the CDR sequences identified by underline in the hHC01 to hHC03 may be considered a defining feature of the invention independently of the surrounding framework sequence.

Specific examples of humanized HC (VH) sequences:

hHC04 (SEQ ID NO 6):

EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWISWVRQAPGKGLVWVGEINPNSSTINYAPSLKD
KFTISRDNAKNTLYLQMNSLRAEDTAVYYCASLYYDYGDAYDYWGQGTLLTVSS

5 hHC05 (SEQ ID NO 7):

EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWFSWVRQAPGKGLVWVGEINPNSSTINYAPSLKD
KFTISRDNAKNTLYLQMNSLRAEDTAVYYCASLYYDYGDAYDYWGQGTLLTVSS

hHC06 (SEQ ID NO 8):

10 EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWISWVRQAPGKGLVWVGEINPSSSTINYAPSLKD
KFTISRDNAKNTLYLQMNSLRAEDTAVYYCASLYYDYGDAYDYWGQGTLLTVSS

hHC07 (SEQ ID NO 9):

EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWFSWVRQAPGKGLVWVGEINPSSSTINYAPSLKD
KFTISRDNAKNTLYLQMNSLRAEDTAVYYCASLYYDYGDAYDYWGQGTLLTVSS

Alignments:

- 15 A CLUSTAL W (1.83) multiple sequence alignment of the various substituted positions within the HC sequence provides appropriate sequence comparisons in Fig. 8. The "General sequence" represents an HC sequence, whereby each X represents a potential amino acid change to any given amino acid. Preferred amino acid substitutions are those described above for each of the potentially mutated positions.

20 **Preferred embodiments regarding humanized VL variants**

Chimeric sequence:

LC mouse (SEQ ID No. 43):

DIVMTQSQRFM TTSVGDRVSVTCKASQSVDSNVAWYQQKPRQSPKALIFASALRFSGV PARFTGSGSGT
DFTLTISNLQSEDLAEYFCQQYNNYPLTFGAGTKLELKR

- 25 The LC mouse sequence represents the variable region of the light chain (VL) originally developed for the chimeric antibody J22.9-xi, which comprises VL and VH domains obtained from a mouse antibody, capable of binding an epitope of the extracellular domain of CD269 (BCMA), and the VL and VH domains are fused to human CL and CH domains, respectively.

Partially humanized sequences:

30 **LC partially humanized (SEQ ID NO 10):**

DIVMTQSPATLSVSVGDEVTLTCKASQSVDSNVAWYQQKPGQAPKLLIYSDDLRFSGV PARFSGSGSGT
DFTLTISNLQSEDFAVYYCQQYNNYPLTFGAGTKLELKR

- 35 The LC partially humanized sequence represents a modified sequence (via amino acid substitutions) in comparison to the chimeric antibody disclosed in the examples of the present invention, whereby the VL and VH binding regions have been modified with respect to their sequence to make them more suitable for administration in humans.

WO 2015/166073

PCT/EP2015/059562

- 10 -

Humanized VL sequence:hLC01 (SEQ ID NO 11):

EIVMTQSPATLSVSPGERATLSCASQSVDSNVAWYQQKPGQAPRALIYSASLRFSGIPARFSGSGSGT
EFTLTISLQSEDFAVYYCQQYNNYPLTFGAGTKLELKR

5 Humanized VL sequence with removal of post translational modification motifs:

hLC02 (SEQ ID NO 12):

EIVMTQSPATLSVSPGERATLSCASQSVX₁X₂NVAWYQQKPGQAPRALIYSASLRFSGIPARFSGSGSG
TEFTLTISLQSEDFAVYYCQQYNNYPLTFGAGTKLELKR

Wherein:

10 X₁X₂: ES, SS, TS, QS, HS, DH, preferably ES.

The "hLC01" and "hLC02" humanized sequences represent novel amino acid sequences that comprise amino acid sequence changes in comparison to both the original chimeric sequence and the partially humanized sequences described herein.

15 The PTM mutations are intended to remove potentially detrimental post translational modification motifs from said proteins, whilst maintaining the advantageous binding properties. The positions 1, 8, 9, 10, 13, 15, 17, 19, 20, 21, 22, 30, 41, 43, 45, 49, 58, 63, 70, 77, 83, 85 and/or 87 of hLC01 and hLC02 are preferably mutated (substituted) in comparison to the original chimeric sequence.) The importance of the substitution relates primarily to the resulting amino acid, not the originating amino acid. The change may therefore also be carried
20 out from the corresponding amino acid of the original chimeric amino acid or other variant.

The following substitutions are novel over the chimeric and partially humanized sequences:

- the amino acid D1 of the LC (VL) sequence is substituted with E;
- the amino acid V15 of the LC (VL) sequence is substituted with P;
- the amino acid D17 of the LC (VL) sequence is substituted with E;
- 25 - the amino acid V19 of the LC (VL) sequence is substituted with A;
- the amino acid T22 of the LC (VL) sequence is substituted with S;
- the amino acids D30 and S31 of the LC (VL) sequence is substituted with any amino acid combination, preferably ES, SS, TS, QS, HS, DH;
- the amino acid V58 of the LC (VL) sequence is substituted with I; and/or
- 30 - the amino acid D70 of the LC (VL) sequence is substituted with E.

Sequences that may be modified in their CDR binding regions at those residues required for interaction with BCMA:

hLC03 – modified amino acids involved in interaction with BCMA (SEQ ID NO 13):

WO 2015/166073

PCT/EP2015/059562

- 11 -

EIVMTQSPATLSVSPGERATLSCKASQSVDX₁X₂VX₃WX₄QQKPGQAPRALIX₅X₆AX₇X₈RX₉SGIPARFS
 GSX₁₀X₁₁GTEFTLTISLQSEDFAVYYCX₁₂QX₁₃NNX₁₄PX₁₅TFGAGTKLELKR

Wherein preferred amino acids are:

X₁: S, H, T, N, D, Q;

5 X₂: N, E, Q;

X₃: A, G, V, S, T, L, I;

X₄: Y, F, L, I, V, A, G;

X₅: Y, F, L;

X₆: S, T;

10 X₇: S, T, D, N, H, E, Q;

X₈: L, V, I, M;

X₉: F, L, I, V, Y, M;

X₁₀: G, X;

X₁₁: S, X;

15 X₁₂: Q, V, L, I, M;

X₁₃: Y, F, L, I, Q;

X₁₄: Y, F, R, Q, K; and/or

X₁₅: L, I, V, F.

20 The "hLC03 humanized sequence" represents novel amino acid sequences that comprise amino acid sequence changes in comparison to both the original chimeric sequence and the partially humanized sequence. These sequence changes are intended to reflect potential changes in the amino acids that bind the BCMA target, which may be substituted, whilst maintaining the advantageous binding properties. The importance of the substitution relates primarily to the resulting amino acid, not the originating amino acid. The change may therefore
 25 also be carried out from the corresponding amino acid of the original chimeric amino acid or other variant.

For example:

- the amino acid S31 of the LC (VL) sequence is S, H, T, N, D, Q;
- the amino acid N32 of the LC (VL) sequence is N, E, Q;
- 30 - the amino acid A34 of the LC (VL) sequence is A, G, V, S, T, L, I;
- the amino acid Y36 of the LC (VL) sequence is Y, F, L, I, V, A, G;
- the amino acid Y49 of the LC (VL) sequence is Y, F, L;

- the amino acid S50 of the LC (VL) sequence is S, T;
- the amino acid S52 of the LC (VL) sequence is S, T, D, N, H, E, Q;
- the amino acid L53 of the LC (VL) sequence is L, V, I, M;
- the amino acid F55 of the LC (VL) sequence is F, L, I, V, Y, M;
- 5 - the amino acid G66 of the LC (VL) sequence is G, X;
- the amino acid S67 of the LC (VL) sequence is S, X;
- the amino acid Q89 of the LC (VL) sequence is Q, V, L, I, M;
- the amino acid Y91 of the LC (VL) sequence is Y, F, L, I, Q;
- the amino acid Y94 of the LC (VL) sequence is Y, F, R, Q, K; and/or
- 10 - the amino acid L96 of the LC (VL) sequence is L, I, V, F.

In general, any change to a CDR region may also be considered as a feature of a CDR sequence when considered independently of the framework sequence as a whole. Such modified CDR sequences may be considered defining features of the present invention, either within or independent of their context in the entire framework region described herein. For example, the CDR sequences identified by underline in the hLC01 to hLC03 may - in their unmodified or substituted form – be considered a defining feature of the invention independently of the surrounding framework sequence.

Example of humanized LC sequence:

hLC04 (SEQ ID NO 14):

20 EIVMTQSPATLSVSPGERATLSCKASQSVESNVAWYQQKPGQAPRALIYSASLRFSGIPARFSGSGSGT
EFTLTISLQSEDFAVYYCQQYNNYPLTFGAGTKLELKR

Alignments:

A CLUSTAL W (1.83) multiple sequence alignment of the various potentially amended sites within the LC sequence provides appropriate sequence comparisons in Fig. 9. The “General sequence” represents an LC sequence, whereby each X represents a potential amino acid change. Preferred amino acid substitutions are those described above for each of the potentially mutated positions.

The present invention therefore relates to the humanized sequences according to hHC01, hHC02, hHC03, hHC04, hHC05, hHC06, hHC07, hLC01, hLC02, hLC03 and/or hLC04, or any given combination thereof.

All possible combinations of potential modifications for any given potentially variant residue proposed herein (as identified by X in the “general” sequence) are encompassed by the present invention. By combining one or more of these various substitutions, humanized variants may be generated that exhibit the desired binding properties of the chimeric antibody originally developed and demonstrated herein. The antibodies or parts thereof described herein also encompass a sequence with at least 80%, preferably 90%, sequence identity to those humanized sequences disclosed explicitly or disclosed through a sequence formula.

Preferred embodiments regarding the antibody epitope

The invention therefore relates to an isolated antibody or antibody fragment that binds CD269 (BCMA), wherein the antibody binds an epitope comprising one or more amino acids of residues 13 to 32 of CD269 (BCMA).

5 The amino acid sequence of residues 13 to 32 of CD269 are shown in SEQ ID No. 40. The N-terminus sequence of CD269 is provided in SEQ ID No. 39. The extracellular domain of CD269 is provided as SEQ ID No. 38.

10 An antigen comprising the extracellular domain of CD269 according to SEQ ID No. 38 was used in vaccination in order to generate the binding specificity of the mouse and chimeric antibody described herein. Use of the entire CD269 protein, or fragments thereof comprising either a membrane-bound or intracellular domain, as an antigen during antibody generation could produce antibodies that bind concealed or intracellular domains of CD269, thereby rendering such agents unsuitable or disadvantageous for therapeutic application. The antibodies of the present invention are therefore defined by their binding to the extracellular portion of CD269. The specific epitope within the extracellular domain also represents a
15 preferred novel and unexpected characterising feature of the invention.

Fab fragments prepared from mouse or chimeric antibodies were crystallized in complex with the purified BCMA extracellular domain and the complex structure solved. The structural analysis has revealed detailed information of the epitope of the antibody of the present invention and its biological relevance. The binding of an epitope comprising one or more
20 amino acids of residues 13 to 32 of CD269 (BCMA) of the extracellular domain by the antibody of the present invention is an advantageous property, as this region shows a significant overlap with the binding sites of BAFF and APRIL, the two natural ligands of CD269. No anti-CD269 antibody described in art to date has shown such comprehensive overlap with the BAFF and APRIL binding sites.

25 In one embodiment the isolated antibody or antibody fragment of the present invention is characterised in that the antibody binds an epitope comprising one or more of amino acids 13, 15, 16, 17, 18, 19, 20, 22, 23, 26, 27 or 32 of CD269 (BCMA). In another embodiment the isolated antibody or antibody fragment of the present invention is characterised in that the antibody binds an epitope consisting of amino acids 13, 15, 16, 17, 18, 19, 20, 22, 23, 26, 27
30 and 32 of CD269 (BCMA). These residues represent the amino acids that interact directly with the antibody of the present invention, as identified by the crystal structure data provided herein. The numbering of these residues has been carried out with respect to SEQ ID No. 39, which provides the N-terminal sequence of CD269.

35 In one embodiment of the invention the isolated antibody or antibody fragment is characterised in that the antibody binding to CD269 (BCMA) disrupts the BAFF-CD269 and/or APRIL-CD269 interaction.

The binding of the antibodies of the present invention to the extracellular domain of CD269 disrupts the BAFF-CD269 interaction. Due to the fact that the binding sites of APRIL and BAFF are positioned at similar sites to the antibody epitope, the binding of the antibody to
40 CD269 will also block the APRIL-CD269 interaction.

Comparison of the specific epitope of the antibody of the present invention with the binding sites of APRIL and BAFF, for which crystal structures have been solved and their interaction sites mapped, reveals a comprehensive overlap in the binding sites of the natural ligands and the antibody as described herein. This represents a beneficial and unexpected aspect of the invention and enables a reliable and effective disruption of BAFF-CD269 and/or APRIL-CD269 interactions.

The invention therefore relates to an isolated antibody or antibody fragment as described herein, wherein the antibody disrupts the APRIL-CD269 interaction by binding an epitope comprising one or more amino acids of residues 13, 15, 17, 18, 19, 22, 26, 27, 30, 31, 32, 33, 34, 35 of CD269 (BCMA), in particular consisting of amino acids 13, 15, 17, 18, 19, 22, 26, 27, 32. These amino acids correspond to the binding site of APRIL on CD269, and the overlapping residues of CD269 that bind both the antibody as described herein and APRIL, respectively.

The invention therefore relates in another embodiment to an isolated antibody or antibody fragment as described herein, wherein the antibody disrupts the BAFF-CD269 interaction by binding an epitope comprising one or more amino acids of residues 13, 15, 16, 17, 18, 19, 22, 25, 26, 27, 29, 30, 31, 32, 34, 35 of CD269 (BCMA), in particular an epitope consisting of amino acids 13, 15, 16, 17, 18, 19, 22, 26, 27, 32. These amino acids correspond to the binding site of BAFF on CD269, and the overlapping residues of CD269 that bind both the antibody as described herein and BAFF, respectively.

Although antibodies that bind CD269 have been described in the art that also potentially disrupt APRIL- or BAFF-interactions with CD269, no relevant disclosure is provided relating to the specific epitope of such antibodies. It cannot be assumed that the previously described antibodies also bind an epitope with such a comprehensive overlap as the antibodies of the present invention. Even if APRIL- or BAFF-interactions with CD269 have been shown to be disrupted, this could potentially occur due to binding a considerably different epitope and subsequent steric hindrance of APRIL or BAFF docking. The degree of disruption of APRIL- or BAFF-interactions with CD269 caused by the antibodies of the prior has not been documented previously.

The antibodies of the present invention enable an effective and reliable disruption, which potentially represents an improved technical effect in comparison to those antibodies described in the art. An in vitro blocking assay can be performed for determination and comparison of BAFF and/or APRIL disruption, for example with the extracellular domain of human BCMA and recombinant BAFF or APRIL.

In a preferred embodiment the epitope specificity, in combination with the high affinity shown by the antibodies described herein, represents a novel and unexpected technical effect. In essence, the exceptionally high affinity of the J22.9 antibody and the humanized variants thereof, provides not only "disruption" or "blocking" of the binding of the natural ligands; but rather the ultra-high affinity of the antibodies of the invention ensures that the native ligands are essentially excluded completely or almost completely from binding their BCMA target when the antibody is present.

As disclosed in the examples below, the affinity of the humanized antibodies as described herein is surprisingly high and comparatively better than similar approaches attempted in the prior art. A Kd in the pM range (as shown below) is commonly accepted as an outstanding affinity not to be expected in common practice.

- 5 In another aspect the humanized antibody or antibody fragment of the invention binds CD269 with high affinity, for example when measured by surface plasmon resonance, such as Biacore, the antibody binds to human CD269 with an affinity of 100nM, 90, 80, 70, 60, 50, 40, 30nM or less, or 20nM or less, or an affinity of 15nM or less, or an affinity of 5nM or less, or an affinity of 1000 pM or less, or an affinity of 500pM or less, or an affinity of 100pM or less, or
10 80pM or less, or for example about 50pM.

In a further embodiment the antibody binds to human CD269 when measured by surface plasmon resonance, such as Biacore, of between about 1pM and about 100nM, or between about 100pM and about 50nM, or between about 200pM and about 20nM.

Further preferred embodiments of the invention

- 15 In one embodiment the invention relates to an antibody or antibody fragment comprising an amino acid sequence defined by one or more of the amino acids that directly interact with the CD269 target and/or one or more amino acids that interact via water interactions (see Tables 1 to 6). The large number of water interactions involved in the binding of the antibody as described herein to the epitope represents an unusual and surprising aspect of the binding. In
20 particular the high affinity of the antibody directed to the particular epitope described herein, in combination with the large number of water interactions involved in the binding surface between the antibody and epitope, represents a surprising and unexpected aspect of the invention.

- The invention therefore relates to an antibody or antibody fragment comprising an amino acid
25 sequence as described herein, wherein the sequence is characterised by the presence of the specific amino acid residues that are involved in the interaction surface with the target epitope via a water bridge according to table 5, selected from the group comprising Ser31, Asn32, Tyr36, Ser50, Ser52, Gly66, Gln89, Tyr91 and/or Tyr94 of the light chain, and/or Trp33, Ser35, Trp47, Glu50, Leu99 and/or Tyr101 of the heavy chain, with respect to the chimera disclosed
30 herein, or with respect to the corresponding residue of the humanized sequence variants disclosed herein.

- Although the examples with respect to water bridge formation of the antibody were carried out with the J22.9-xi chimera, the inventors assert that this technical effect is maintained in the humanized variants of the present invention due to the maintenance of binding characteristics
35 in the humanized variants compared to the original chimeric antibody tested. In the heavy chain, the only mutated water bridge residue is Y101, but its water interaction involves a main chain (i.e. backbone) atom, and therefore can be reasonably assumed not to change due to mutating the sidechain; in the light chain there are no mutations of residues involved in water bridges.

The antibody of the invention can be further characterised by the amino acid residues of the epitope involved in the interaction via water bridges with the antibody as described herein. The relevant features are provided in Table 5. The invention is therefore, for example in one embodiment, characterised in that residue Ser31 of the light chain interacts with Thr32 of CD269 via a water molecule. Such a description of the binding properties of the antibody of the present invention is intended for each interaction as provided in Table 5.

Furthermore, sequence variants of the antibodies described herein are encompassed in the present invention, in which one or more residues involved in a "water bridging" interaction is modified in order to "substitute" a direct side-chain interaction into the sequence at the expense of a water "bridge". For example, a mutation or change could be made in the amino acid sequence displacing the water from the interaction interface but not substantially affecting the affinity of the interaction. The invention therefore relates to an antibody or antibody fragment comprising an amino acid sequence as described herein, wherein the sequence is characterised by sequence variation of those amino acid residues that are involved in the interaction surface with the target epitope via a water bridge according to table 5, selected from the group comprising Ser31, Ser31, Asn32, Tyr36, Ser50, Ser52, Gly66, Gln89, Tyr91 and/or Tyr94 of the light chain, and/or Trp33, Ser35, Trp47, Glu50, Leu99 and/or Tyr101 of the heavy chain, with respect to the chimera disclosed herein, or with respect to the corresponding residue of the humanized sequence variants disclosed herein. Variation at the corresponding positions of the humanized antibodies described herein may relate to any given amino acid substitution, preferably an amino acid substitution that would effectively displace the water from the interaction but maintain similar binding properties with respect to epitope affinity and specificity.

In one embodiment of the invention the isolated antibody or antibody fragment is characterised in that the antibody is glycosylated, preferably comprising an N-linked oligosaccharide chain, preferably at Asn297 of the heavy chain.

Glycosylation of the antibody refers to the attachment of carbohydrates or glycans to the antibody. N-linked glycans are attached to the nitrogen of asparagine or arginine side-chains. The carbohydrate chains attached to the target proteins serve various functions. For instance, some proteins do not fold correctly unless they are glycosylated first. Also, polysaccharides linked at the amide nitrogen of asparagine in the protein can confer stability on some secreted glycoproteins. Glycosylation in this case is not a strict requirement for proper folding, but unglycosylated protein can be degraded more quickly.

As is demonstrated in the examples of the present invention, the deglycosylation of the antibody disclosed therein leads to a reduction in therapeutic effect in comparison to glycosylated forms of the antibody. It was surprising, that the glycosylation would play a significant role in maintaining activity of the antibody. The glycosylation therefore represents a preferred embodiment of the invention associated with unexpected technical advantages.

As demonstrated in the examples herein, although the overall tumor load of animals treated with J22.9-xi-N-glycan (deglycosylated) was not significantly different from animals receiving the isotype control antibody, the lifespan of these mice was substantially increased compared

to the isoAb-treated group. Since J22.9-xi-N-glycan was shown to be unable to induce ADCC or CDC, this result indicates that alone the binding of J22.9-xi to BCMA hinders tumor growth. It may be reasonably considered that this is due to blocking of the interaction between the receptor and its native ligands (APRIL and BAFF). This aspect of the invention and the antibodies described herein represents a surprising technical effect, which could not have been derived from the antibodies of the prior art. The J22.9-xi-N-glycan (deglycosylated) can be considered a control sample in the experiments described which enables the binding of the antibody to its target epitope, without the downstream effects of ADCC or CDC, to be assessed for potentially therapeutic effect. The antibodies of the invention, preferably with glycosylation, therefore demonstrate such an effective epitope binding that enables the prevention (or significant disruption) of binding by the natural ligands to lead to cell toxicity. This characteristic of the antibodies described herein has not been described for similar antibodies described in the art.

Although the examples with respect to glycosylation of the antibody were carried out with the J22.9-xi chimera, the inventors assert that this technical effect is maintained in the humanized variants of the present invention due to the maintenance of binding characteristics in the humanized variants compared to the original chimeric antibody tested. The preferred position of glycosylation (Asn297 of the heavy chain) has no direct connection to any of the mutated residues and lies in the human constant region of the full IgG. It is therefore reasonable to assume that no differences in the glycosylation pattern at this position exist in any of the J22.9 variants compared to the chimeric antibody.

Use and functional aspects of the invention

The antibodies of the present invention are capable of binding the epitopes described herein, blocking interaction of the natural ligands of this epitope, and inducing CDC and ADCC.

Antibody-Dependent Cell-Mediated Cytotoxicity (ADCC) is one factor induced by the antibody of the present invention that generates the desired therapeutic effect. ADCC is a mechanism of cell-mediated immune defense, whereby an effector cell of the immune system actively lyses a target cell, whose membrane-surface antigens have been bound by specific antibodies. After binding of CD269-expressing cells by the antibodies of the present invention ADCC may be induced. Classical ADCC is mediated by natural killer (NK) cells; macrophages, neutrophils and eosinophils can also mediate ADCC. ADCC is part of the adaptive immune response due to its dependence on a prior antibody response. Experiments in mice may indicate that ADCC is an important mechanism of action of therapeutic antibodies as described herein.

A preferred embodiment of the invention relates to the isolated antibody or antibody fragment as described herein for use as a medicament in the treatment of a medical disorder associated with the presence of pathogenic B cells.

In one embodiment of the invention the medical disorder is a CD269-associated disorder, preferably associated with pathogenic B cells, which is preferably a disease of plasma cells and/or memory B cells.

A disease of plasma cells may be a cancer of plasma cells, for example multiple myeloma, plasmacytoma, Waldenström macroglobulinemia or plasma cell leukemia. A disease of plasma cells may be a cancer of B lymphocytes, such as Hodgkin's disease.

5 In one embodiment of the invention the medical disorder is an autoimmune disease associated with autoreactive plasma cells and/or autoreactive memory B cells, such as an inflammatory autoimmune disease, for example systemic lupus erythematosus or rheumatic arthritis.

10 The invention therefore also encompasses a method of treatment for the medical disorders as disclosed herein, preferably comprising the administration of a therapeutically effective amount of antibody to a subject in need of such treatment.

A further aspect of the invention relates to an antibody-drug conjugate (ADC) comprising the antibody or antibody fragment as described herein. Anti-CD269 Antibody-Drug Conjugates "anti-CD269 ADC" can be described as an anti-CD269 antibody or fragment thereof conjugated to a therapeutic agent. In certain embodiments, the ADC comprises an anti-CD269
15 antibody (e.g., a humanized variant of J22.9-xi as described herein).

The ADCs or ADC derivatives as described herein produce clinically beneficial effects on CD269-expressing cells when administered to a subject with a CD269-expressing medical condition, such as cancer or autoimmune disorder. In one embodiment, the anti-CD269 antibody or derivative thereof is conjugated to a cytotoxic agent, such that the resulting ADC
20 or ADC derivative exerts a cytotoxic effect on a CD269-expressing cancer cell, preferably when taken up or internalized by the cell.

The anti-CD269 ADC or ADC derivative is preferably internalized and accumulates within a CD269-expressing cell, where the ADC or ADC derivative exerts a therapeutic effect (e.g., a cytotoxic effect). Particularly suitable moieties for conjugation to antibodies or antibody
25 derivatives are chemotherapeutic agents, prodrug converting enzymes, radioactive isotopes or compounds, or toxins. For example, an anti-CD269 antibody or derivative thereof can be conjugated to a cytotoxic agent such as a chemotherapeutic agent, or a toxin (e.g., a cytostatic or cytotoxic agent).

30 Another aspect of the invention relates to a preferably isolated nucleic acid molecule selected from the group consisting of:

a) a nucleic acid molecule comprising a nucleotide sequence

- which encodes an isolated antibody or antibody fragment according to any one of the preceding claims,
- which encodes an amino acid sequence selected from the group consisting of those sequences according to SEQ ID 1 to 31 and 41 to 42,
35
- comprising a sequence or sequence fragment of SEQ ID No. 32 to 36,

- b) a nucleic acid molecule which is complementary to a nucleotide sequence in accordance with a);
- c) a nucleic acid molecule comprising a nucleotide sequence having sufficient sequence identity to be functionally analogous/equivalent to a nucleotide sequence according to a) or b), comprising preferably a sequence identity to a nucleotide sequence according to a) or b) of at least 80%, preferably 90%, more preferably 95% ;
- d) a nucleic acid molecule which, as a consequence of the genetic code, is degenerated into a nucleotide sequence according to a) through c); and
- e) a nucleic acid molecule according to a nucleotide sequence of a) through d) which is modified by deletions, additions, substitutions, translocations, inversions and/or insertions and functionally analogous/equivalent to a nucleotide sequence according to a) through d).

A further aspect of the invention relates to a host cell, such as a bacterial cell or mammalian cell, preferably a hybridoma cell or cell line, capable of producing an antibody or antibody fragment as described herein, and/or comprising a nucleic acid molecule as described herein.

A further aspect of the invention relates to a pharmaceutical composition comprising the isolated antibody or antibody fragment as described herein, a nucleic acid molecule as described herein or a host cell as described herein, together with a pharmaceutically acceptable carrier.

An additional and surprising aspect of the invention is an improved stability of the antibody as disclosed herein. The antibody can readily be stored for extended periods under appropriate conditions without any loss of binding affinity. Appropriate tests have been carried out regarding maintenance of activity after storage at either -80 or 4 deg C, which demonstrate unexpectedly good stability of the antibody and maintenance of activity after storage at both aforementioned temperatures (Fig. 3c). This improved stability is evident for the chimeric antibody, and surprisingly also for the humanized variants thereof. Unexpectedly the stability of the humanized variants is improved over the chimera under long term storage.

A further advantage of the antibodies as described herein the effective systemic depletion of myeloma cells as demonstrated in the examples. Antibodies previously disclosed in the prior art have not been demonstrated to exhibit the desired anti-plasma cell effect in a systemic manner. Studies carried out on the antibodies of the prior art disclose only sub-cutaneous injection with myeloma cells and subsequent treatment of the isolated cell mass. The present invention provides an antibody capable of systemic depletion of cancerous multiple myeloma cells after their i.v. injection, as demonstrated in the examples. The effective depletion of targeted cells represents a technical effect that has not previously been demonstrated in the prior art in addition to a beneficial property of the antibodies of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

As used herein, an "antibody" generally refers to a protein consisting of one or more polypeptides substantially encoded by immunoglobulin genes or fragments of immunoglobulin genes. Where the term "antibody" is used, the term "antibody fragment" may also be considered to be referred to. The recognized immunoglobulin genes include the kappa, 5 lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD, and IgE, respectively. The basic immunoglobulin (antibody) structural unit is known to comprise a tetramer or dimer. Each tetramer is 10 composed of two identical pairs of polypeptide chains, each pair having one "light" (L) (about 25 kD) and one "heavy" (H) chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids, primarily responsible for antigen recognition. The terms "variable light chain" and "variable heavy chain" refer to these variable regions of the light and heavy chains respectively. Optionally, the antibody or the 15 immunological portion of the antibody, can be chemically conjugated to, or expressed as, a fusion protein with other proteins.

The antibodies of the invention are intended to bind against mammalian, in particular human, protein targets. The use of protein names may correspond to either mouse or human versions of a protein.

20 "Specific binding" is to be understood as via one skilled in the art, whereby the skilled person is clearly aware of various experimental procedures that can be used to test binding and binding specificity. Some cross-reaction or background binding may be inevitable in many protein-protein interactions; this is not to detract from the "specificity" of the binding between antibody and epitope. The term "directed against" is also applicable when considering the term 25 "specificity" in understanding the interaction between antibody and epitope.

Antibodies of the invention include, but are not limited to polyclonal, monoclonal, bispecific, human, humanized or chimeric antibodies, single variable fragments (ssFv), single domain antibodies (such as VHH fragments from nanobodies), single chain fragments (scFv), Fab fragments, F(ab')₂ fragments, fragments produced by a Fab expression library, anti-idiotypic 30 antibodies and epitope-binding fragments or combinations thereof of any of the above, provided that they retain the original binding properties. Also mini-antibodies and multivalent antibodies such as diabodies, triabodies, tetravalent antibodies and peptabodies can be used in a method of the invention. The immunoglobulin molecules of the invention can be of any class (i.e. IgG, IgE, IgM, IgD and IgA) or subclass of immunoglobulin molecules. Thus, the 35 term antibody, as used herein, also includes antibodies and antibody fragments either produced by the modification of whole antibodies or synthesized de novo using recombinant DNA methodologies.

Humanized antibody comprising one or more CDRs of antibodies of the invention or one or more CDRs derived from said antibodies can be made using any methods known in the art. 40 For example, four general steps may be used to humanize a monoclonal antibody. These are: (1) determining the nucleotide and predicted amino acid sequence of the starting antibody light and heavy variable domains (2) designing the humanized antibody, i.e., deciding which

antibody framework region to use during the humanizing process (3) the actual humanizing methodologies/techniques and (4) the transfection and expression of the humanized antibody. See, for example, U.S. Pat. Nos. 4,816,567; 5,807,715; 5,866,692; 6,331,415; 5,530,101; 5,693,761; 5,693,762; 5,585,089; 6,180,370; 5,225,539; 6,548,640.

- 5 The term humanized antibody means that at least a portion of the framework regions, and optionally a portion of CDR regions or other regions involved in binding, of an immunoglobulin is derived from or adjusted to human immunoglobulin sequences. The humanized, chimeric or partially humanized versions of the mouse monoclonal antibodies can, for example, be made by means of recombinant DNA technology, departing from the mouse and/or human genomic
- 10 DNA sequences coding for H and L chains or from cDNA clones coding for H and L chains. Humanized forms of mouse antibodies can be generated by linking the CDR regions of non-human antibodies to human constant regions by recombinant DNA techniques (Queen et al., 1989; WO 90/07861). Alternatively the monoclonal antibodies used in the method of the invention may be human monoclonal antibodies. Human antibodies can be obtained, for
- 15 example, using phage-display methods (WO 91/17271; WO 92/01047).

As used herein, humanized antibodies refer also to forms of non-human (e.g. murine, camel, llama, shark) antibodies that are specific chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin.

- 20 As used herein, human or humanized antibody means an antibody having an amino acid sequence corresponding to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies known in the art or disclosed herein. Human antibodies can be selected by competitive binding experiments, or otherwise, to have the same epitope specificity as a particular mouse antibody. The humanized antibodies of the
- 25 present invention surprisingly share the useful functional properties of the mouse antibodies to a large extent. Human polyclonal antibodies can also be provided in the form of serum from humans immunized with an immunogenic agent. Optionally, such polyclonal antibodies can be concentrated by affinity purification using amyloid fibrillar and/or non-fibrillar polypeptides or fragments thereof as an affinity reagent. Monoclonal antibodies can be obtained from serum
- 30 according to the technique described in WO 99/60846.

- The present invention further relates to the use of the antibodies, or fragments thereof, as described herein, for example the variable regions, in recognition molecules or affinity reagents that are suitable for selective binding to a target. The affinity reagent, antibody or fragment thereof according to the invention may be PEGylated, whereby PEGylation refers to
- 35 covalent attachment of polyethylene glycol (PEG) polymer chains to the inventive antibody. PEGylation may be routinely achieved by incubation of a reactive derivative of PEG with the target molecule. PEGylation to the antibody can potentially mask the agent from the host's immune system, leading to reduced immunogenicity and antigenicity or increase the hydrodynamic size of the agent which may prolong its circulatory time by reducing renal
- 40 clearance.

A variable region of an antibody refers to the variable region of the antibody light chain or the variable region of the antibody heavy chain, either alone or in combination. The variable regions of the heavy and light chain each consist of four framework regions (FR) connected by three complementarity determining regions (CDRs) also known as hypervariable regions. The CDRs in each chain are held together in close proximity by the FRs and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies. There are at least two techniques for determining CDRs: (1) an approach based on cross-species sequence variability (i.e., Kabat *et al.* Sequences of Proteins of Immunological Interest, (5th ed., 1991, National Institutes of Health, Bethesda Md.)); and (2) an approach based on crystallographic studies of antigen-antibody complexes (Al-lazikani *et al.* (1997) J. Molec. Biol. 273:927-948). As used herein, a CDR may refer to CDRs defined by either approach or by a combination of both approaches.

In some embodiments, the invention provides an antibody, which comprises at least one CDR, at least two, at least three, or more CDRs that are substantially identical to at least one CDR, at least two, at least three, or more CDRs of the antibody of the invention. Other embodiments include antibodies which have at least two, three, four, five, or six CDR(s) that are substantially identical to at least two, three, four, five or six CDRs of the antibodies of the invention or derived from the antibodies of the invention. In some embodiments, the at least one, two, three, four, five, or six CDR(s) are at least about 85%, 86%, 87%, 88%, 89%, 90%, 95%, 96%, 97%, 98%, or 99% identical to at least one, two or three CDRs of the antibody of the invention. It is understood that, for purposes of this invention, binding specificity and/or overall activity is generally retained, although the extent of activity may vary compared to said antibody (may be greater or lesser).

The half life and cytotoxic potential of an antibody are dependent primarily on the interaction of the Fc-domain with different Fc-gamma-receptors. In the case of the antibody half life, the neonatal Fc receptor (FcRn) plays a major role. This receptor is expressed on several cell types and tissues such as monocytes and vascular endothelia cells that are able to take up serum proteins into their recycling endosomes. In the endosomes, the pH is decreased to approximately 6 and under these conditions the antibodies are able to bind to FcRn. This interaction protects the antibodies from degradation until they are again released into the blood where the physiological pH disrupts the binding to the receptor (Roopenian and Akilesh (2007) Nat Rev Immunol 7:715-725). The higher the affinity of the antibody to the FcRn at pH 6, the greater the half life of that antibody. Fc-fragment mutations known to stabilize this interaction are summarised in Presta (2008, Curr Opin Immunol 20:460-470).

Therapeutic antibodies can act through several mechanisms upon binding to their target. The binding itself can trigger signal transduction, which can lead to programmed cell death (Chavez-Galan *et al.* (2009) Cell Mol Immunol 6:15-25). It can also block the interaction of a receptor with its ligand by either binding to the receptor or the ligand. This interruption can cause apoptosis if signals important for survival are affected (Chiu *et al.* (2007) Blood 109:729-739). With regard to cell-depletion there are two major effector mechanisms known. The first is the complement-dependent cytotoxicity (CDC) towards the target cell. There are three different pathways known. However, in the case of antibodies the important pathway for

CDC is the classical pathway which is initiated through the binding of C1q to the constant region of IgG or IgM (Wang and Weiner (2008) Expert Opin Biol Ther 8:759-768).

The second mechanism is called antibody-dependent cellular cytotoxicity (ADCC). This effector function is characterized by the recruitment of immune cells which express Fc-receptors for the respective isotype of the antibody. ADCC is largely mediated by activating Fc-gamma receptors (FcγR) which are able to bind to IgG molecules either alone or as immune complexes. Mice exhibit three (FcγRI, FcγRIII and FcγRIV) and humans five (FcγRI, FcγRIIA, FcγRIIC, FcγRIIIA and FcγRIIIB) activating Fcγ-receptors. These receptors are expressed on innate immune cells like granulocytes, monocytes, macrophages, dendritic cells and natural killer cells and therefore link the innate with the adaptive immune system. Depending on the cell type there are several modes of action of FcγR-bearing cells upon recognition of an antibody-marked target cell. Granulocytes generally release vasoactive and cytotoxic substances or chemoattractants but are also capable of phagocytosis. Monocytes and macrophages respond with phagocytosis, oxidative burst, cytotoxicity or the release of pro-inflammatory cytokines whereas Natural killer cells release granzymes and perforin and can also trigger cell death through the interaction with FAS on the target cell and their Fas ligand (Nimmerjahn and Ravetch (2008) Nat Rev Immunol 8:34-47; Wang and Weiner (2008) Expert Opin Biol Ther 8:759-768; Chavez-Galan *et al.* (2009) Cell Mol Immunol 6:15-25).

The antibody-dependent cellular cytotoxicity (ADCC) can also be improved by strengthening the binding of the Fc-domain to activating Fc-gamma receptors (FcγR). This can also be achieved through mutations in the Fc-gamma domain as summarized in Presta (2008, Curr Opin Immunol 20:460-470).

Another way to change the ADCC is manipulation of the sugar moiety present on each IgG at Asn297. Defucylation and removal of sialic acid from the end of the sugar molecules are known to increase the cytotoxic potential of an antibody (Anthony and Ravetch (2010) J Clin Immunol 30 Suppl 1:S9-14).

Sequence variants of the claimed nucleic acids, proteins and antibodies, for example defined by the claimed % sequence identity, that maintain the said properties of the invention are also included in the scope of the invention. Such variants, which show alternative sequences, but maintain essentially the same binding properties, such as target specificity, as the specific sequences provided are known as functional analogues, or as functionally analogous. Sequence identity relates to the percentage of identical nucleotides or amino acids when carrying out a sequence alignment.

It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology or sequence identity to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Deletions, substitutions and other changes in sequence that fall under the described sequence identity are also encompassed in the invention.

Protein sequence modifications, which may occur through substitutions, are also included within the scope of the invention. Substitutions as defined herein are modifications made to the amino acid sequence of the protein, whereby one or more amino acids are replaced with the same number of (different) amino acids, producing a protein which contains a different amino acid sequence than the primary protein, preferably without significantly altering the function of the protein. Like additions, substitutions may be natural or artificial. It is well known in the art that amino acid substitutions may be made without significantly altering the protein's function. This is particularly true when the modification relates to a "conservative" amino acid substitution, which is the substitution of one amino acid for another of similar properties. Such "conserved" amino acids can be natural or synthetic amino acids which because of size, charge, polarity and conformation can be substituted without significantly affecting the structure and function of the protein. Frequently, many amino acids may be substituted by conservative amino acids without deleteriously affecting the protein's function.

In general, the non-polar amino acids Gly, Ala, Val, Ile and Leu; the non-polar aromatic amino acids Phe, Trp and Tyr; the neutral polar amino acids Ser, Thr, Cys, Gln, Asn and Met; the positively charged amino acids Lys, Arg and His; the negatively charged amino acids Asp and Glu, represent groups of conservative amino acids. This list is not exhaustive. For example, it is well known that Ala, Gly, Ser and sometimes Cys can substitute for each other even though they belong to different groups.

Substitution variants have at least one amino acid residue in the antibody molecule removed and a different residue inserted in its place. The sites of greatest interest for substitutional mutagenesis include the hypervariable regions, but FR alterations are also contemplated. If such substitutions result in a change in biological activity, then more substantial changes, denominated "exemplary substitutions" in the table immediately below, or as further described below in reference to amino acid classes, may be introduced and the products screened.

Potential Amino Acid Substitutions:

Original residue	Preferred conservative substitutions	Examples of exemplary substitutions
Ala (A)	Val	Val; Leu; Ile
Arg (R)	Lys	Lys; Gln; Asn
Asn (N)	Gln	Gln; His; Asp, Lys; Arg
Asp (D)	Glu	Glu; Asn
Cys (C)	Ser	Ser; Ala
Gln (Q)	Asn	Asn, Glu

Glu (E)	Asp	Asp; Gln
Gly (G)	Ala	Ala
His (H)	Arg	Asn; Gln; Lys; Arg
Ile (I)	Leu	Leu; Val; Met; Ala; Phe; Norleucine
Leu (L)	Ile	Norleucine; Ile; Val; Met; Ala; Phe
Lys (K)	Arg	Arg; Gln; Asn
Met (M)	Leu	Leu; Phe; Ile
Phe (F)	Tyr	Leu; Val; Ile; Ala; Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr	Tyr; Phe
Tyr (Y)	Phe	Trp; Phe; Thr; Ser
Val (V)	Leu	Ile; Leu; Met; Phe; Ala; Norleucine

Substantial modifications in the biological properties of the antibody are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain.

Conservative amino acid substitutions are not limited to naturally occurring amino acids, but also include synthetic amino acids. Commonly used synthetic amino acids are omega amino acids of various chain lengths and cyclohexyl alanine which are neutral non-polar analogs; citrulline and methionine sulfoxide which are neutral non-polar analogs, phenylglycine which is an aromatic neutral analog; cysteic acid which is a negatively charged analog and ornithine which is a positively charged amino acid analog. Like the naturally occurring amino acids, this list is not exhaustive, but merely exemplary of the substitutions that are well known in the art.

The antibodies of the present invention may be produced by transfection of a host cell with an expression vector comprising the coding sequence for the antibody of the invention. An expression vector or recombinant plasmid is produced by placing these coding sequences for the antibody in operative association with conventional regulatory control sequences capable of controlling the replication and expression in, and/or secretion from, a host cell. Regulatory sequences include promoter sequences, e.g., CMV promoter, and signal sequences which can be derived from other known antibodies. Similarly, a second expression vector can be produced having a DNA sequence which encodes a complementary antibody light or heavy chain. In certain embodiments this second expression vector is identical to the first except insofar as the coding sequences and selectable markers are concerned, so to ensure as far as

possible that each polypeptide chain is functionally expressed. Alternatively, the heavy and light chain coding sequences for the antibody may reside on a single vector.

5 A selected host cell is co-transfected by conventional techniques with both the first and second vectors (or simply transfected by a single vector) to create the transfected host cell of the invention comprising both the recombinant or synthetic light and heavy chains. The transfected cell is then cultured by conventional techniques to produce the engineered antibody of the invention. The antibody which includes the association of both the recombinant heavy chain and/or light chain is screened from culture by appropriate assay, such as ELISA or RIA. Similar conventional techniques may be employed to construct other antibodies.

10 Suitable vectors for the cloning and subcloning steps employed in the methods and construction of the compositions of this invention may be selected by one of skill in the art. For example, the conventional pUC series of cloning vectors may be used. One vector, pUC19, is commercially available. The components of such vectors, e.g. replicons, selection genes, enhancers, promoters, signal sequences and the like, may be obtained from commercial or
15 natural sources or synthesized by known procedures for use in directing the expression and/or secretion of the product of the recombinant DNA in a selected host. Other appropriate expression vectors of which numerous types are known in the art for mammalian, bacterial, insect, yeast, and fungal expression may also be selected for this purpose.

20 The present invention also encompasses a cell line transfected with a recombinant plasmid containing the coding sequences of the antibodies of the present invention. Host cells useful for the cloning and other manipulations of these cloning vectors are also conventional.

Suitable host cells or cell lines for the expression of the antibodies of the invention include mammalian cells such as NS0, Sp2/0, CHO (e.g. DG44), COS, HEK, a fibroblast cell (e.g., 3T3), and myeloma cells, for example it may be expressed in a CHO or a myeloma cell.
25 Human cells may be used, thus enabling the molecule to be modified with human glycosylation patterns. Alternatively, other prokaryotic or eukaryotic cell lines may be employed. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art.

In accordance with the present invention there is provided a method of producing an anti-
30 CD269-antibody of the present invention which binds to and neutralises the activity of human CD269 which method comprises the steps of; providing a first vector encoding a heavy chain of the antibody; providing a second vector encoding a light chain of the antibody; transforming a mammalian host cell (e.g. CHO) with said first and second vectors; culturing the host cell of step (c) under conditions conducive to the secretion of the antibody from said host cell into
35 said culture media; recovering the secreted antibody of step (d). Once expressed, the antibody can be assessed for the desired binding properties using methods as described herein.

The invention encompasses immunoconjugates (interchangeably referred to as "antibody-drug conjugates" or "ADCs") comprising an antibody according to the invention as herein described including, but not limited to, an antibody conjugated to one or more cytotoxic agents, such as a
40 chemotherapeutic agent, a drug, a growth inhibitory agent, a toxin (e.g., a protein toxin, an

enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate). Techniques for conjugating therapeutic agents to proteins, and in particular to antibodies, such as for the Anti-CD269 Antibody-Drug Conjugates of the present invention, are well-known. (See, e.g., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy," in *Monoclonal Antibodies And Cancer Therapy* (Reisfeld *et al.* eds., Alan R. Liss, Inc., 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery," in *Controlled Drug Delivery* (Robinson *et al.* eds., Marcel Dekker, Inc., 2nd ed. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review," in *Monoclonal Antibodies '84: Biological And Clinical Applications* (Pinchera *et al.* eds., 1985); "Analysis, Results, and Future Prospective of the Therapeutic Use of Radiolabeled Antibody In Cancer Therapy," in *Monoclonal Antibodies For Cancer Detection And Therapy* (Baldwin *et al.* eds., Academic Press, 1995); and Thorpe *et al.*, 1982, *Immunol. Rev.* 62:119-58. See also, e.g., PCT publication WO 89/12624.)

Typically, the ADC or ADC derivative comprises a linker region between the therapeutic agent and the anti-CD269 antibody or derivative thereof. As noted supra, in typical embodiments, the linker is cleavable under intracellular conditions, such that cleavage of the linker releases the therapeutic agent from the antibody in the intracellular environment. For example, in some embodiments, the linker is cleavable by a cleaving agent that is present in the intracellular environment (e.g., within a lysosome or endosome or caveolae). The linker can be, e.g., a peptidyl linker that is cleaved by an intracellular peptidase or protease enzyme, including, but not limited to, a lysosomal or endosomal protease. In other embodiments, the cleavable linker is pH-sensitive, i.e., sensitive to hydrolysis at certain pH values. Typically, the pH-sensitive linker is hydrolyzable under acidic conditions. In yet other embodiments, the linker is cleavable under reducing conditions (e.g., a disulfide linker). A variety of disulfide linkers are known in the art (See for example Wawrzynczak *et al.*, In *Immunoconjugates: Antibody Conjugates in Radioimaging and Therapy of Cancer* (C. W. Vogel ed., Oxford U. Press, 1987. See also U.S. Patent No. 4,880,935.)

Typically, the linker is not substantially sensitive to the extracellular environment. In other, non-mutually exclusive embodiments, the linker promotes cellular internalization. In certain embodiments, the linker promotes cellular internalization when conjugated to the therapeutic agent (i.e., in the milieu of the linker-therapeutic agent moiety of the ADC or ADC derivative as described herein). In yet other embodiments, the linker promotes cellular internalization when conjugated to both the therapeutic agent and the anti-CD269 antibody or derivative thereof (i.e., in the milieu of the ADC or ADC derivative as described herein). A variety of linkers that can be used with the present compositions and methods are described in WO 2004010957 entitled "Drug Conjugates and Their Use for Treating Cancer, An Autoimmune Disease or an Infectious Disease" filed July 31, 2003, and U.S. Provisional Application No. 60/400,403, entitled "Drug Conjugates and their use for treating cancer, an autoimmune disease or an infectious disease", filed July 31, 2002 (the disclosure of which is incorporated by reference herein).

In certain embodiments, an immunoconjugate comprises an antibody as described herein, including but not limited to, an antibody and a chemotherapeutic agent or other toxin.

Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S),
 5 momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies.

Antibodies or fragments thereof of the present invention may also be conjugated to one or more toxins, including, but not limited to, a calicheamicin, maytansinoids, dolastatins,
 10 aurostatins, a tricothecene, and CC1065, and the derivatives of these toxins that have toxin activity. Suitable cytotoxic agents include, but are not limited to, an auristatin including dovaline-valine-dolaisoleuine-dolaproine-phenylalanine (MMAF) and monomethyl auristatin E (MMAE) as well as ester forms of MMAE, a DNA minor groove binding agent, a DNA minor groove alkylating agent, an enediyne, a lexitropsin, a duocarmycin, a taxane, including
 15 paclitaxel and docetaxel, a puromycin, a dolastatin, a maytansinoid, and a vinca alkaloid. Specific cytotoxic agents include topotecan, morpholino-doxorubicin, rhizoxin, cyanomorpholino-doxorubicin, dolastatin-10, echinomycin, combretastatin, calicheamicin, maytansine, DM-1, DM-4, netropsin. Other suitable cytotoxic agents include anti-tubulin agents, such as an auristatin, a vinca alkaloid, a podophyllotoxin, a taxane, a baccatin
 20 derivative, a cryptophysin, a maytansinoid, a combretastatin, or a dolastatin. Antitubulin agent include dimethylvaline-valine-dolaisoleuine-dolaproine-phenylalanine-p-phenylenediamine (AFP), MMAF, MMAE, auristatin E, vincristine, vinblastine, vindesine, vinorelbine, VP-16, camptothecin, paclitaxel, docetaxel, epothilone A, epothilone B, nocodazole, colchicines, colcimid, estramustine, cemadotin, discodermolide, maytansine, DM-1, DM-4 or eleutherobin.

25 In some embodiments, the immunoconjugate comprises an antibody conjugated to dolastatins or dolostatin peptidic analogs and derivatives, the auristatins (U.S. Pat. Nos. 5,635,483; 5,780,588). Dolastatins and auristatins have been shown to interfere with microtubule dynamics, GTP hydrolysis, and nuclear and cellular division (Woyke et al. (2001) Antimicrob. Agents and Chemother. 45(12):3580-3584) and have anticancer (U.S. Pat. No. 5,663,149)
 30 and antifungal activity (Pettit et al. (1998) Antimicrob. Agents Chemother. 42:2961-2965). The dolastatin or auristatin (which are pentapeptide derivatives of dolastatins) drug moiety may be attached to the antibody through the N (amino) terminus or the C (carboxyl) terminus of the peptidic drug moiety (WO 02/088172). Exemplary auristatin embodiments include the N-terminus linked monomethylauristatin drug moieties DE and DF, disclosed in
 35 "Monomethylvaline Compounds Capable of Conjugation to Ligands," U.S. Patent No. 7,498,298. As used herein, the abbreviation "MMAE" refers to monomethyl auristatin E. As used herein the abbreviation "MMAF" refers to dovaline-valine-dolaisoleuine-dolaproine-phenylalanine.

Typically, peptide-based drug moieties can be prepared by forming a peptide bond between
 40 two or more amino acids and/or peptide fragments. Such peptide bonds can be prepared, for example, according to the liquid phase synthesis method (see E. Schroder and K. Lubke, "The

Peptides," volume 1, pp 76-136, 1965, Academic Press) that is well known in the field of peptide chemistry.

Maytansinoids may be used as an active agent coupled to the antibody or fragment thereof according to the invention. Maytansinoids are mitototic inhibitors which act by inhibiting tubulin polymerization. Maytansine was first isolated from the east African shrub *Maytenus serrata* (U.S. Pat. No. 3,896, 1 1 1). Subsequently, it was discovered that certain microbes also produce maytansinoids, such as maytansinol and C-3 maytansinol esters (U.S. Pat. No. 4, 151 ,042). Highly cytotoxic maytansinoid drugs can be prepared from ansamitocin precursors produced by fermentation of microorganisms such as *Actinosynnema*. Antibody-maytansinoid conjugates are prepared by chemically linking an antibody to a maytansinoid molecule without significantly diminishing the biological activity of either the antibody or the maytansinoid molecule. See, e.g., U.S. Pat. No. 5,208,020. An average of 3-4 maytansinoid molecules conjugated per antibody molecule has shown efficacy in enhancing cytotoxicity of target cells without negatively affecting the function or solubility of the antibody, although even one molecule of toxin/antibody would be expected to enhance cytotoxicity over the use of naked antibody. Maytansinoids are well known in the art and can be synthesized by known techniques or isolated from natural sources.

Selected examples of the calicheamicin family of antibiotics may be used as an active agent coupled to the antibody or fragment thereof according to the invention. The calicheamicin family of antibiotics is capable of producing double-stranded DNA breaks at sub-picomolar concentrations. For the preparation of conjugates of the calicheamicin family, see U.S. Pat. Nos. 5,712,374, 5,714,586, 5,739,1 16, 5,767,285, 5,770,701 , 5,770,710, 5,773,001 , 5,877,296. Another anti-tumor drug that the antibody can be conjugated is QFA which is an antifolate. Both calicheamicin and QFA have intracellular sites of action and do not readily cross the plasma membrane. Therefore, cellular uptake of these agents through antibody mediated internalization greatly enhances their cytotoxic effects.

Other antitumor agents that can be conjugated to the antibodies include BCNU, streptozocin, vincristine and 5-fluorouracil, the family of agents known collectively LL-E33288 complex described in U.S. Pat. Nos. 5,053,394, 5,770,710, as well as esperamicins (U.S. Pat. No. 5,877,296). The present invention further contemplates an immunoconjugate formed between an antibody and a compound with nucleolytic activity (e.g., a ribonuclease or a DNA endonuclease such as a deoxyribonuclease; DNase). For selective destruction of the tumor, the antibody may comprise a highly radioactive atom.

A pharmaceutically acceptable carrier in the sense of the present invention may be any non-toxic material that does not significantly interfere in a detrimental sense with the effectiveness of the biological activity of the antibodies of the present invention. Evidently, the characteristics of the carrier will depend on the route of administration. Such a composition may contain, in addition to the active substance and carrier, diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. Formulation of pharmaceutically-acceptable excipients and carrier solutions is well-known to those of skill in the art, as is the development of suitable dosing and treatment regimens for using the particular compositions

described herein in a variety of treatment regimens, including e.g., oral, parenteral, intravenous, intranasal, and intra-muscular administration and formulation.

The medicament, otherwise known as a pharmaceutical composition, containing the active ingredient (antibody or antibody fragment) may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated. The present invention also refers to a pharmaceutical composition for topical application, oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. A skilled person is aware of the carriers and additives required for particular application forms.

When a therapeutically effective amount of the active substance (antibody or antibody fragment) of the invention is administered by intravenous, cutaneous or subcutaneous injection, the active substance may be in the form of a pyrogen-free, parenterally acceptable aqueous solution.

The invention also relates to administration of a therapeutically relevant amount of antibody as described herein in the treatment of a subject who has the medical disorders as disclosed herein. As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit. The amount of active substance in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Larger doses may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further.

The preparation of such parenterally acceptable solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to the active substance, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present

invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

The dose of the antibody administered evidently depends on numerous factors well-known in the art such as, e.g., the chemical nature and pharmaceutical formulation of the antibody, and of body weight, body surface, age and sex of the patient, as well as the time and route of administration. For an adult, the dose may exemplarily be between 0.001 µg and 1 g per day, preferably between 0.1 µg and 100 mg per day, more preferably between 1 µg and 100 mg per day, even more preferably between 5 µg and 10 mg per day. In a continuous infusion, the dose may exemplarily be between 0.01 µg and 100 mg, preferably between 1 µg and 10 mg per kilogram body mass per minute.

In another aspect of the present invention there is provided an antibody according to the invention as herein described for use in the treatment of a B-cell mediated or plasma cell mediated disease or antibody mediated disease or disorder selected from Multiple Myeloma (MM), chronic lymphocytic leukemia (CLL), Non-secretory multiple myeloma, Smoldering multiple myeloma, Monoclonal gammopathy of undetermined significance (MGUS), Solitary plasmacytoma (Bone, Extramedullar), Lymphoplasmacytic lymphoma (LPL), Waldenstrom's Macroglobulinemia, Plasma cell leukemia, Primary Amyloidosis (AL), Heavy chain disease, Systemic lupus erythematosus (SLE), POEMS syndrome / osteosclerotic myeloma, Type I and II cryoglobulinemia, Light chain deposition disease, Goodpasture's syndrome, Idiopathic thrombocytopenic purpura (ITP), Acute glomerulonephritis, Pemphigus and Pemphigoid disorders, and Epidermolysis bullosa acquisita; or any Non-Hodgkin's Lymphoma B-cell leukemia or Hodgkin's lymphoma (HL) with BCMA expression or any diseases in which patients develop neutralising antibodies to recombinant protein replacement therapy wherein said method comprises the step of administering to said patient a therapeutically effective amount of the antibody as described herein.

B-cell disorders can be divided into defects of B-cell development/immunoglobulin production (immunodeficiencies) and excessive/uncontrolled proliferation (lymphomas, leukemias). As used herein, B-cell disorder refers to both types of diseases, and methods are provided for treating B-cell disorders with an antibody.

In one aspect of the present invention the disease is Multiple Myeloma.

Use of the antibody as described herein in the manufacture of a medicament for the treatment of diseases and disorders as described herein is also provided.

For example in one aspect of the invention there is provided the use of the antibody as described herein for use in the treatment or prophylaxis of diseases and disorders responsive to modulation (such as inhibiting or blocking) of the interaction between BCMA and the ligands BAFF and APRIL.

In one embodiment of the invention the isolated antibody or antibody fragment is intended for use in the treatment of B lymphocyte cancers, such as Hodgkin's lymphoma.

In one embodiment of the invention the isolated antibody or antibody fragment is intended for use in the treatment of an autoimmune disease, such as a medical disorder associated with inflammation, preferably autoimmune disease with an inflammatory component, whereby the autoimmune disease is selected from Takayasu Arteritis, Giant-cell arteritis, familial

5 Mediterranean fever, Kawasaki disease, Polyarteritis nodosa, cutaneous Polyarteritis nodosa, Hepatitis-associated arteritis, Behcet's syndrome, Wegener's granulomatosis, ANCA-vasculitides, Churg-Strauss syndrome, microscopic polyangiitis, Vasculitis of connective tissue diseases, Hennoch-Schönlein purpura, Cryoglobulinemic vasculitis, Cutaneous leukocytoclastic angiitis, Tropical aortitis, Sarcoidosis, Cogan's syndrome, Wiskott-Aldrich

10 Syndrome, Lepromatous arteritis, Primary angiitis of the CNS, Thromboangiitis obliterans, Paraneoplastic arteritis, Urticaria, Dego's disease, Myelodysplastic syndrome, Erythema elevatum diutinum, Hyperimmunoglobulin D, Allergic Rhinitis, Asthma bronchiale, chronic obstructive pulmonary disease, periodontitis, Rheumatoid Arthritis, atherosclerosis, Amyloidosis, Morbus Chron, Colitis ulcerosa, Autoimmune Myositis, Diabetes mellitus, Multiple

15 sclerosis, Guillain-Barre Syndrome, histiocytosis, Osteoarthritis, atopic dermatitis, periodontitis, chronic rhinosinusitis, Psoriasis, psoriatic arthritis, Microscopic colitis, Pulmonary fibrosis, glomerulonephritis, Whipple's disease, Still's disease, erythema nodosum, otitis, cryoglobulinemia, Sjogren's syndrome, Lupus erythematosus, aplastic anemia, Osteomyelofibrosis, chronic inflammatory demyelinating polyneuropathy, Kimura's disease,

20 systemic sclerosis, chronic periaortitis, chronic prostatitis, idiopathic pulmonary fibrosis, chronic granulomatous disease, Idiopathic achalasia, bleomycin-induced lung inflammation, cytarabine-induced lung inflammation, Autoimmunthrombocytopenia, Autoimmunneutropenia, Autoimmunhemolytic anemia, Autoimmunlymphocytopenia, Chagas' disease, chronic autoimmune thyroiditis, autoimmune hepatitis, Hashimoto's Thyroiditis, atropic thyroiditis,

25 Graves disease, Autoimmune polyglandular syndrome, Autoimmune Addison Syndrome, Pemphigus vulgaris, Pemphigus foliaceus, Dermatitis herpetiformis, Autoimmune alopecia, Vitiligo, Antiphospholipid syndrome, Myasthenia gravis, Stiff-man syndrome, Goodpasture's syndrome, Sympathetic ophthalmia, Folliculitis, Sharp syndrome and/or Evans syndrome, in particular hay fever, periodontitis, atherosclerosis, rheumatoid arthritis, preferably rheumatoid

30 arthritis or multiple sclerosis.

SEQUENCES

Preferred antibody sequences of the invention:

SEQ ID No.	Sequence	Description
SEQ ID No. 1	QVQLQQSGGGLVQPGGSLKLSCAASGIDFSRYWMSWVR RAPGKGLEWIGEINPDSSTINYAPSLKDKFIIISRDNAK NTLYLQMSKVRSEDTALYYCASLYYDYGDAMDYWGQGT SVTVSS	HC (VH) mouse
SEQ ID No. 2	EVQLVESGGGLVQPGGSLRLSCAASGFTFDDYWMSWVR QAPGKGLEWVGEINPDSSTINYAPSLKGRFTISRDNAK NTLYLQMNSLRAEDTAVYYCASLYYDYGDAMDYWGQGT LVTVSS	HC partially humanized

SEQ ID No. 3	EVQL VE SGGGLVQPGGSLRLSCAASG FT FSRYWMSWVR QAPGKGL VWV GEINPDSSTINYAPSLKDK FT ISRDN AKNTLYLQ MNSLRA EDTAVYYCASLYDYGDAMDYWGQGT LVTVSS	hHC01
SEQ ID No. 4	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYW X₁ SWV RQAPGKGLVWVGEINP X₂X₃ STINYAPSLKDKFTISR DNAKNTLYLQ MNSLRA EDTAVYYCASLY X₄ DYGD X₅ DY WGQGT L TVSS Wherein X ₁ : I, F, L, V, Y, C, G, A, S, T, preferably I or F; X ₂ X ₃ : SS, NS, TS, GS, KS, RS, SD, SN, DE, preferably SS; X ₄ : Y, L, A, V, F, I, W, preferably Y; and/or X ₅ : Y, L, F, I, V, A, C, preferably Y	hHC02
SEQ ID No. 5	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRY X₁MX₂ W VRQAPGKGLV X₃ VG X₄ INPDSSTINYAPSLKDKFTISR DNAKNTLYLQ MNSLRA EDTAVYYCAS X₅X₆X₇ DYGD X₈ M DYWGQGT L TVSS Wherein X ₁ : W, F, Y, preferred W; X ₂ : S, T, N, Q, D, E, preferred S; X ₃ : W, F, Y, preferred W; X ₄ : E, Q, preferred E; X ₅ : L, I, V, G, A, preferred L; X ₆ : Y, X, preferred Y; X ₇ : Y, F, L, I, V, M, preferred Y; and/or X ₈ : A, G, V, preferred A	hHC03
SEQ ID No. 6	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYW ISW VRQAPGKGLVWVGEINP NS STINYAPSLKDKFTISR DNAKNTLYLQ MNSLRA EDTAVYYCASLYDYGD AYD YWGQGT L TVSS	hHC04
SEQ ID No. 7	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYW FSW VRQAPGKGLVWVGEINP NS STINYAPSLKDKFTISR DNAKNTLYLQ MNSLRA EDTAVYYCASLYDYGD AYD YWGQGT L TVSS	hHC05
SEQ ID No. 8	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYW ISW VRQAPGKGLVWVGEINP SS STINYAPSLKDKFTISR DNAKNTLYLQ MNSLRA EDTAVYYCASLYDYGD AYD YWGQGT L TVSS	hHC06

SEQ ID No. 9	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYW FSW VRQAPGKGLVWVGEINP SS STINYAPSLKDKFTISR DNAKNTLYLQMNSLRAEDTAVYYCASLYDYGD AYD <u>YWGQGT</u> LVTVSS	hHC07
SEQ ID No. 43	DIVMTQSQRFMTTSVGDRVSVTCKASQSVDSNVAWY QQKPRQSPKALIF SAS LRFSGV PAR FTGSGSGTDFT LTISNLQSEDLAEYFC QQYNNYPL TFGAGTKLELKR	LC (VL) mouse
SEQ ID No. 10	DIVMTQSPATLSVSVGDEVTLTCKASQSVDSNVAWYQQ KPGQAPKLLIY SDDL RFSGV PAR FSGSGSGTDFTLTIS SLQSEDFAVYYC QQYNNYPL TFGAGTKLELKR	LC partially humanized
SEQ ID No. 11	E IVMTQSP PATLSVSPGERATLS CKASQSVDSNVAWYQQ KPGQ APRALIY SASLRFS GIPARFSGSGSGTEFT LTIS SLQSEDFAVYYCQQYNNYPL TFGAGTKLELKR	hLC01
SEQ ID No. 12	EIVMTQSPATLSVSPGERATLSCKASQSV X₁X₂ NVAWY QQKPGQAPRALIY SAS LRFS GIPAR FSGSGSGTEFTLT ISSLQSEDFAVYYC QQYNNYPL TFGAGTKLELKR Wherein: X ₁ X ₂ : ES, SS, TS, QS, HS, DH, preferably ES.	hLC02
SEQ ID No. 13	EIVMTQSPATLSVSPGERATLSCKASQSV D X ₁ X ₂ VX ₃ W X₄ QQKPGQAPRALI X₅X₆AX₇X₈R X ₉ SGIPARFSGS X₁₀X₁₁ GTEFTLTISSLQSEDFAVYYC X₁₂QX₁₃NNX₁₄PX₁₅ TFG AGTKLELKR Wherein: X ₁ : S, H, T, N, D, Q; X ₂ : N, E, Q; X ₃ : A, G, V, S, T, L, I; X ₄ : Y, F, L, I, V, A, G; X ₅ : Y, F, L; X ₆ : S, T; X ₇ : S, T, D, N, H, E, Q; X ₈ : L, V, I, M; X ₉ : F, L, I, V, Y, M; X ₁₀ : G, X; X ₁₁ : S, X; X ₁₂ : Q, V, L, I, M; X ₁₃ : Y, F, L, I, Q; X ₁₄ : Y, F, R, Q, K; and/or	hLC03

	X ₁₅ : L, I, V, F	
SEQ ID No. 14	EIVMTQSPATLSVSPGERATLSCKASQSVESNVAWYQQ KPGQAPRALIYSASLRFSGIPARFSGSGSGTEFTLTIS SLQSEDFAVYYCQQYNNYPLTFGAGTKLELKR	hLC04
SEQ ID No. 15	RYWX ₁ S Wherein: X ₁ : I, F, L, V, Y, C, G, A, S, T, preferably I or F	H-CDR1 PTM
SEQ ID No. 16	EINPX ₂ X ₃ STINYAPSLKDK Wherein: X ₂ X ₃ : SS, NS, TS, GS, KS, RS, SD, SN, DE, preferably SS	H-CDR2 PTM
SEQ ID No. 17	SLYX ₄ DYGDAX ₅ DYW Wherein: X ₄ : Y, L, A, V, F, I, W, preferably Y; and/or X ₅ : Y, L, F, I, V, A, C, preferably Y	H-CDR3 PTM
SEQ ID No. 18	RYWIS	H-CDR1 PTM a
SEQ ID No. 19	RYWFS	H-CDR1 PTM b
SEQ ID No. 20	EINPNSSTINYAPSLKDK	H-CDR2 PTM a
SEQ ID No. 21	EINPSSSTINYAPSLKDK	H-CDR2 PTM b
SEQ ID No. 22	SLYYDYGDAYDYW	H-CDR3 PTM a
SEQ ID No. 23	KASQSVX ₁ X ₂ NVA Wherein: X ₁ X ₂ : ES, SS, TS, QS, HS, DH, preferably ES	L-CDR1 PTM
SEQ ID No. 24	SASLRFS	L-CDR2 PTM
SEQ ID No. 25	QQYNNYPLTFG	L-CDR3 PTM
SEQ ID No. 26	KASQSVDSNVA	L-CDR1 PTM a
SEQ ID No. 27	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWMSWVR QAPGKGLVWGEINPDSSTINYAPSLKDKFTISRDNK NTLYLQMNSLRAEDTAVYYCASLYYDYGDAMDYWGQGT LVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDY FPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT VPSSSLGTQTYICNVNHKPSNTKVDKRVKPKSCDKTHT	Full length humanized HC

	CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	
SEQ ID No. 28	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWFSWVR QAPGKGLVWGEINPSSSTINYAPSLKDKFTISRDNAL NTLYLQMNLSRAEDTAVYYCASLYDYGDYDYGQGT LVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDY FPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT VPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHT CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	Full length humanized HC with PTM mutations 1 (FSY)
SEQ ID No. 29	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWISWVR QAPGKGLVWGEINPSSSTINYAPSLKDKFTISRDNAL NTLYLQMNLSRAEDTAVYYCASLYDYGDYDYGQGT LVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDY FPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT VPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHT CPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVV VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	Full length humanized HC with PTM mutations 2 (ISY)
SEQ ID No. 30	EIVMTQSPATLSVSPGERATLSCKASQSVDSNVAWYQQ KPGQAPRALIYSASLRFSGI PARFSGSGSGTEFTLTIS SLQSEDFAVYYCQQYNNYPLTFGAGTKLELKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHK VYACEVTHQGLSPVTKSFNRGEC	Full length humanized LC
SEQ ID No. 31	EIVMTQSPATLSVSPGERATLSCKASQSVESNVAWYQQ KPGQAPRALIYSASLRFSGI PARFSGSGSGTEFTLTIS SLQSEDFAVYYCQQYNNYPLTFGAGTKLELKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHK VYACEVTHQGLSPVTKSFNRGEC	Full length humanized LC with PTM mutations

Preferred Nucleotide Sequences

SEQ ID No. 32	GAATTCCACCATGGGATGGTCATGTATCATCCTTTTCTA GTAGCAACTGCAACCGGTGCCACAGTGAAGTGCAGCTGG TCGAATCTGGAGGAGGCCTGGTTCAGCCTGGTGGCAGCCT TAGGCTCTCTTGTGCAGCCTCTGGCTTTACCTTCTCACGG TATTGGATGAGCTGGGTGAGACAGGCTCCAGGGAAAGGTC TGGTGTGGGTAGGGGAGATAAACCCGATAGCAGCACGAT CAACTATGCTCCGTCAGTGAAGACAAGTTCACCATTTCC	Full length humanized HC
---------------	--	-----------------------------

	CGCGATAATGCCAAGAACACTCTCTACTTGCAGATGAATT CCCTTCGAGCCGAGGATACAGCGGTGTACTACTGCGCCAG TCTGTACTacgactATGGGGACGCAATGGACTATTGGGGA CAAGGCACACTGGTGACTGTTAGCTCCGCGTCGACCAAGG GCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCAC CTCTGGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGAC TACTTCCCCGAACCTGTGACGGTCTCGTGGAACCTCAGGCG CCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCCTACA GTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTG CCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACG TGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGT TGAGCCCCAAATCTTGTGACAAAACCTCACACATGCCACCG TGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCC TCTTCCCCC AAAACCCAAGGACACCCTCATGATCTCCCG GACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCAC GAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCG TGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCCTC CTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCA AGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAAC CATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTG TACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACC AGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAG CGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACG GCTCCTTCTTCTCTATAGCAAGCTCACCCTGGACAAGAG CAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATG CATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCT CCCTGTCCCCGGGTAAATGAGTGCGACGGCCGGGCGGCGG CGGCGGATCC	
SEQ ID No. 33	GAATTCCACCATGGGATGGTTCATGTATCATCCTTTTTCTA GTAGCAACTGCAACCGGTGTCCACAGTGAAGTGCAGCTGG TCGAATCTGGAGGAGGCCTGGTTCAGCCTGGTGGCAGCCT TAGGCTCTCTTGTGCAGCCTCTGGCTTTACCTTCTCACGG TATTGGTTTCAGCTGGGTGAGACAGGCTCCAGGGAAAGGTC TGGTGTGGGTAGGGGAGATAAAACCCAGCAGCAGCACGAT CAACTATGCTCCGTCACTGAAAGACAAGTTCACCATTTCC CGCGATAATGCCAAGAACACTCTCTACTTGCAGATGAATT CCCTTCGAGCCGAGGATACAGCGGTGTACTACTGCGCCAG TCTGTACTACGACTATGGGGACGCATACGACTATTGGGGA CAAGGCACACTGGTGACTGTTAGCTCCGCGTCGACCAAGG GCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCAC CTCTGGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGAC TACTTCCCCGAACCTGTGACGGTCTCGTGGAACCTCAGGCG CCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCCTACA GTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTG CCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACG TGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGT TGAGCCCCAAATCTTGTGACAAAACCTCACACATGCCACCG TGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCC TCTTCCCCC AAAACCCAAGGACACCCTCATGATCTCCCG GACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCAC GAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCG TGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCCTC CTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCA AGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAAC CATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTG TACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACC AGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAG CGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACG	Full length humanized HC with PTM mutations 1

	GCTCCTTCTTCTCTATAGCAAGCTCACCGTGGACAAGAG CAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATG CATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCT CCCTGTCCCCGGGTAAATGAGTGCGACGGCCGGGCGGCGG CGGCGGATCC	
SEQ ID No. 34	GAATTCCACCATGGGATGGTCATGTATCATCCTTTTTCTA GTAGCAACTGCAACCGGTGTCCACAGTGAAGTGCAGCTGG TCGAATCTGGAGGAGGCCTGGTTCAGCCTGGTGGCAGCCT TAGGCTCTCTTGTGCAGCCTCTGGCTTTACCTTCTCACGG TATTGGaTCAGCTGGGTGAGACAGGCTCCAGGGAAAGGTC TGGTGTGGGTAGGGGAGATAAACCCAGCAGCAGCACGAT CAACTATGCTCCGTCACTGAAAGACAAGTTCACCATTTC CGCGATAATGCCAAGAACAACCTCTCTACTTGCAGATGAATT CCCTTCGAGCCGAGGATACAGCGGTGTACTACTGCGCCAG TCTGTACTACGACTATGGGGACGCATACGACTATTGGGGA CAAGGCACACTGGTGACTGTTAGCTCCGCGTCGACCAAGG GCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCAC CTCTGGGGGACAGCGGCCCTGGGCTGCCTGGTCAAGGAC TACTTCCCCGAACCTGTGACGGTCTCGTGGAACCTCAGGCG CCCTGACCAGCGGCGTGCACACCTTCCCGGCTGTCCTACA GTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTG CCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACG TGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGT TGAGCCCCAAATCTTGTGACAAAACCTCACACATGCCACCG TGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCC TCTTCCCCCCCCAAAACCCAAGGACACCCTCATGATCTCCCG GACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCAC GAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCG TGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTC CTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCA AGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAAC CATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTG TACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACC AGGTACAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAG CGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACCTACAAGACCACGCCTCCCGTGCTGGACTCCGACG GCTCCTTCTTCTCTATAGCAAGCTCACCGTGGACAAGAG CAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATG CATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCT CCCTGTCCCCGGGTAAATGAGTGCGACGGCCGGGCGGCGG CGGCGGATCC	Full length humanized HC with PTM mutations 2
SEQ ID No. 35	GAATTCCACCATGGGATGGTcATGTATCATCCTTTTTCTA GTAGCAACTGCAACCGGTGTACACTCCGAGATCGTGATGA CCCAGTCTCCTGCTACCCTGAGCGTTTCTCCCGGTGAAAG GGCCACACTCAGCTGCAAAGCCTCTCAAAGCGTGGACAGC AATGTGCGCTGGTATCAGCAGAAACCTGGCCAAGCTCCGA GAGCACTGATCTATTCCGCGTCATTGCGCTTTTCCGGCAT ACCAGCACGGTTTAGTGGCTCAGGGAGTGGGACTGAGTTC ACTCTGACGATTAGCTCCCTTCAGTCAGAGGATTTGCGCG TGTAATACTGTGAGCAGTACAACAACCTATCCCCTCACATT CGGAGCTGGAACCAAGCTGGAAGTGAAGCGTACGGTGGCT GCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGT TGAAATCTGGAAGTGCCTCTGTGTGTGCCTGCTGAATAA CTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGAT AACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAG AGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCAC CCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTC TACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCG TCACAAAGAGCTTCAACAGGGGAGAGTGTTAGGGATCC	Full length humanized LC

WO 2015/166073

PCT/EP2015/059562

- 39 -

SEQ ID No. 36	GAATTCCACCATGGGATGGTcATGTATCATCCTTTTTCTA GTAGCAACTGCAACCGGTGTACACTCCGAGATCGTGATGA CCCAGTCTCCTGCTACCCTGAGCGTTTCTCCCGGTGAAAG GGCCACACTCAGCTGCAAAGCCTCTCAAAGCGTGGAGAGC AATGTCGCCTGGTATCAGCAGAAACCTGGCCAAGCTCCGA GAGCACTGATCTATTCCGCGTCATTGCGCTTTTCCGGCAT ACCAGCACGGTTTAGTGGCTCAGGGAGTGGGACTGAGTTC ACTCTGACGATTAGCTCCCTTCAGTCAGAGGATTTGCGCG TGTACTACTGTCAGCAGTACAACAACCTATCCCCTCACATT CGGAGCTGGAACCAAGCTGGAAGTGAAGCGTACGGTGGCT GCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGT TGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAA CTTCTATCCCAGAGAGGGCCAAAGTACAGTGAAGGTGGAT AACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAG AGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCAC CCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTC TACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCG TCACAAAGAGCTTCAACAGGGGAGAGTGTTAGGGATCC	Full length humanized LC with PTM mutations
---------------	--	---

Preferred sequences of the invention pertaining to CD269 (BCMA):

SEQ ID No.	Sequence	Description
SEQ ID No. 37	MSPILGYWKIKGLVQPTRLLLEYLEEKYEELHYERDEGDK WRNKKFELGLEFPNLPYYIDGDVKLTQSMAIIRYIADKHN MLGGCPKERAIEISMLEGAVLDIRYGVSRIAYSKDFETLKV DFLSKLPEMLKMFEDRLCHKTYLNGDHVTHPDFMLYDALD VVLYMDPMCLDAFPKLVCFKKRIEAIPOIDKYLKSSKYIA WPLQGWQATFGGGDHPKSDLVPRGSMAGQCSQNEYFDSL LHACIPCQLRCSSNTPPLTCQRYCNASVTNSVKGTNALEH HHHHH	GST-BCMA-His
SEQ ID No. 38	MAGQCSQNEYFDSLHACIPCQLRCSSNTPPLTCQRYCNA SVTNSVKGTNALE	BCMA extracellular domain
SEQ ID No. 39	MLQMAGQCSQNEYFDSLHACIPCQLRCSSNTPPLTCQRY CNASVTNSVKGTNALE	BCMA N-terminus sequence
SEQ ID No. 40	YFDSLHACIPCQLRCSSNT	BCMA antibody epitope – amino acids 13 to 32 of BCMA

Preferred generalized amino acid sequences comprising the humanized sequence modifications:

5

SEQ ID No.	Sequence	Description
SEQ ID No. 41	X ₁ VQLX ₂ X ₃ SGGGLVQPGGSLX ₄ LSCAASGX ₅ X ₆ FX ₇ X ₈ YWZ ₁ SWVRX ₉ APGKGLEWX ₁₀ GEINPZ ₂ SSTINYAPSLKX ₁₁ X ₁₂ F X ₁₃ ISRDNKNTLYLQMX ₁₄ X ₁₅ X ₁₆ RX ₁₇ EDTAX ₁₈ YYCASLY	General sequence for humanized HC antibodies comprising

	YDYGDAZ ₃ DYWGQGT ₁₉ VTVSS wherein X1: Q, E; X2: Q, V; X3: Q, E; X4: K, R; X5: I, F; X6: D, T; X7: S, D; X8: R, D; X9: R, Q; X10: I, V; X11: D, G; X12: K, R; X13: I, T; X14: S, N; X15: K, S; X16: V, L; X17: S, A; X18: L, V; X19: S, L; and wherein at least one of Z ₁ : I or F; Z ₂ : S and/or Z ₃ : Y.	the PTM deletion modifications
SEQ ID No. 42	DIVMTQ ₁ SX ₂ X ₃ X ₄ X ₅ X ₆ SVGD ₇ VX ₈ X ₉ TCKASQSVESNVA WYQQK ₁₀ PK ₁₁ LIX ₁₂ SX ₁₃ X ₁₄ LRFS ₁₅ GV ₁₆ PARFX ₁₇ GS GSGTDF ₁₈ TLTISX ₁₉ LQSE ₂₀ DX ₂₁ AX ₂₂ YX ₂₃ CQQYNNYPLTFG AGTKLELKR wherein X1: Q, P; X2: R, A; X3: F, T; X4: M, L; X5: T, S; X6: T, V; X7: R, E; X8: S, T; X9: V, L; X10: R, G; X11: S, A; X12: A, L; X13: F, Y; X14: A, D; X15: S, D; X16: T, S; X17: N, S; X18: L, F; X19: E, V; X20: F, Y	General sequence for humanized LC antibodies comprising the PTM deletion modifications

FIGURES

The invention is demonstrated by way of the example by the examples and figures disclosed herein. The figures provided herein represent particular embodiments of the invention and are not intended to limit the scope of the invention. The figures are to be considered as providing a further description of possible and potentially preferred embodiments that enhance the technical support of one or more non-limiting embodiments.

Short description of the figures:

Figure 1: In vitro characterization of J22.9-xi.

Figure 2: The structure of CD269 (BCMA) and the J22.9-xi Fab:CD269 complex

Figure 3: In vitro cytotoxicity of J22.9-xi.

Figure 4: Efficacy of J22.9-xi in xenografted NSG mice.

Figure 5: Treatment of established tumors.

Figure 6: Tumor treatment in the early phase of disease.

Figure 7: Instability of hybridoma J22.9.

Figure 8: Sequences of the humanized sequences of the HC compared to J22.9-xi.

Figure 9: Sequences of the humanized sequences of the LC compared to J22.9-xi.

Figure 10: Sequence optimized variants of J22.9-xi show similar binding in ELISA.

Figure 11: Sequence optimized variants of J22.9-xi show similar binding in flow cytometry.

Figure 12: SPR raw data.

Figure 13: Gel electrophoresis of antibody variants

Detailed description of the figures:

Figure 1: In vitro characterization of J22.9-xi. Concentration dependent binding of J22.9-xi to BCMA in (a) ELISA and (b) by flow cytometry using CD269-positive MM.1S cells. (c) Binding affinity of J22.9-xi to BCMA was determined from surface plasmon resonance measurements with the indicated concentrations of BCMA. (d) J22.9-xi blocks the interaction between BAFF and BCMA adsorbed onto microtiter plates.

Figure 2: (a) CD269 (BCMA) recognition surface. Three views of the extracellular domain of CD269 (BCMA) showing the binding epitope residues for BAFF/APRIL and J22.9-xi. At top, a view directly on the binding face of CD269 (BCMA); the light grey shading indicates all residues comprising the binding epitope of BAFF and APRIL as identified from their crystal structures, black residues (shown as spheres) do not contact either BAFF, APRIL or J22.9-xi; the subset of epitope residues involved in J22.9-xi binding are shown in surface representation; remaining light grey residues shown as spheres are part of both the BAFF and APRIL epitopes but make no direct contacts to J22.9-xi. The middle and lower panels show the same representation as in the top panel but rotated 90° toward and away from the viewer, respectively. (b) Two views of the J22.9-xi Fab:CD269 complex. J22.9-xi is shown in surface representation with the heavy chain coloured light grey and the light chain in dark grey. CD269 (BCMA) is shown in ribbon representation bound to the J22.9-xi antigen pocket. At left, full view of the Fab:CD269 complex; at right, the complex tilted toward the viewer to show the binding pocket.

Figure 3: In vitro cytotoxicity of J22.9-xi. (a) CD269-positive MM.1S-Luc cells mixed with human PBMCs at an effector to target ratio of 20:1 were incubated with the indicated concentrations of J22.9-xi for 4 hours. Open symbols indicate cytotoxic activity of J22.9-xi without -N-glycans when incubated with PBMCs from donor 1 and 2. Error bars indicate SEM. (b) Deglycosylation does not affect J22.9-xi binding to MM.1S cells. (c) Storage of J22.9-xi for 3 weeks at 4 °C or -80 °C does not affect cytotoxicity.

Figure 4: Efficacy of J22.9-xi in xenografted NSG mice. (a) Tumor development over time with administration of 200 µg of J22.9-xi or the control antibody twice a week, and untreated control mice. (b) Total tumor burden between day 6 and 41 (Area under the curve (AUC) of (a)). Plotted are the mean values with SEM (**P < 0.01, ***P < 0.001, t-test). (c) Overall survival of J22.9-xi and isotype control mice. The P value was calculated using the Log-rank (Mantel-Cox) Test. (d-1) Detection of MM.1S-Luc cells in the indicated groups without administration of therapeutic antibody. Below the rightmost image the numbers (41, 41, 44, 40) indicate the days post-tumor cell injection on which the specific mouse died. (d-2) Detection of MM.1S-Luc cells in the indicated groups at day 21 and 28. Dorsal view. (e) Relationship between J22.9-xi concentration and tumor development. (f) Total tumor load between day 6 and 42 (AUC of (e)). Mean values with SEM (**P < 0.01, ***P < 0.001, t-test). (g) Overview of experimental time line.

Figure 5: Treatment of established tumors. (a) Tumor development over time with administration of 200 µg of J22.9-xi or control antibody twice weekly, and untreated control mice. (b) Total tumor load between day 8 and 48 (AUC of (a)). Plotted are the mean values with SEM (*P < 0.05, **P < 0.01, t-test). (c) Overall survival of J22.9-xi and isotype control mice. The P value was calculated using the Log-rank (Mantel-Cox) Test. An overview of the experimental time line is provided in Fig. 5d.

Figure 6: Tumor treatment in the early phase of disease. (a) Course of tumor growth when treated with 2 µg, 20 µg or 200 µg of J22.9-xi or 200 µg of either the isotype control antibody or J22.9-xi without -N-glycans, and without tumor. (b) Total tumor burden within day 9 to 44 (AUC of (a)). Shown are the mean values with SEM (*P < 0.05, **P < 0.01, t-test). (c) Survival of antibody-treated and control xenograft SCID-Beige mice. The P values were calculated using Log-rank (Mantel-Cox) Test. An overview of the experimental time line is provided in Fig. 6d.

Figure 7: Instability of hybridoma J22.9. The supernatant of hybridoma J22.9 tested positive for binding to BCMA in ELISA on BCMA-coated microtiter plates at day 1. Later analysis at indicated time points revealed a reduction of binding capacity of the supernatant. The medium was exchanged on days 7, 14 and 21.

Figure 8: Summary of the sequences of the humanized antibodies compared to J22.9-xi. Sequence comparisons were carried out using standard alignment software.

Figure 9: Summary of the sequences of the humanized antibodies compared to J22.9-xi. Sequence comparisons were carried out using standard alignment software.

Figure 10: Binding of the chimeric J22.9-xi and humanized variants was tested via ELISA using human BCMA (hBCMA) or cynomolgous BCMA (cyBCMA) coated microtiter plates (J22.9-H corresponds to humanized sequence SEQ ID No. 27; J22.9-FSY corresponds to humanized and PTM modified SEQ ID No. 28; J22.9-ISY corresponds to humanized and PTM modified SEQ ID No. 29).

Figure 11: Binding of the chimeric J22.9-xi and humanized variants was tested via flow cytometry using the human MM cell line RPMI-8226 (J22.9-FSY corresponds to humanized and PTM modified SEQ ID No. 28; J22.9-ISY corresponds to humanized and PTM modified SEQ ID No. 29).

Figure 12: SPR raw data: Binding affinities of the chimeric J22.9-xi and humanized variants to human (Fig. 12 A) and cynomolgus (Fig. 12 B) BCMA was measured by Surface Plasmon Resonance (SPR) Spectrometry. IgGs were immobilized via amine chemistry to a Proteon GLH sensor chip and binding measured with BCMA in the mobile phase. The order of raw data traces in the graph corresponds to the order to samples listed in the legend of the figure.

Figure 13: Gel electrophoresis of antibody variants. Antibody variants were run in non-reduced SDS-PAGE and stained to show protein migration.

EXAMPLES

The invention is demonstrated by way of the examples disclosed herein. The examples provided herein represent only particular embodiments of the invention and are not intended to limit the scope of the invention. The examples are to be considered as providing a further description of possible and potentially preferred embodiments that enhance the technical support of one or more non-limiting embodiments.

Although the examples with respect to crystallization of the antibody-epitope complex and the *in vitro* and *in vivo* anti-tumor effect was carried out using the original chimeric antibody J22.9-xi, the inventors assert that these technical effects are maintained in the humanized variants of the present invention, due to the maintenance of binding characteristics in the humanized variants compared to the original chimeric antibody tested. The data provided from the chimeric antibody is therefore provided as reference material and an indication of the industrial applicability and usefulness of the claimed human variants. Preliminary biological data indicates comparable effects between J22.9-xi and the humanized variants.

Binding and blocking characteristics of the J22.9-xi and BCMA interaction

The novel chimeric antibody (J22.9-xi) binds to the extracellular domain of human CD269 (BCMA, TNFRSF17). This was initially ascertained by ELISA and flow cytometry on the human multiple myeloma cell line MM.1S (Fig. 1a,b). The affinity of J22.9-xi to BCMA was determined using surface plasmon resonance (SPR). The mean K_d is 54 pM as shown in Figure 1c.

BCMA is known to trigger signals important for the survival of multiple myeloma and plasma cells *in vivo* through interaction with its ligands BAFF and/or APRIL (Mackay F *et al.* (2003) *Annu Rev Immunol* 21:231-264). An *in vitro* blocking assay was therefore performed with the extracellular domain of human BCMA and recombinant BAFF. The binding of J22.9-xi to BCMA clearly blocks the interaction between the receptor and its ligand BAFF. Using the isotype control antibody instead of J22.9-xi, recombinant BAFF binding to BCMA is unaffected (Fig. 1d).

The crystal structure of the J22.9-xi-Fab-BCMA-complex reveals an extensive binding interface with BCMA

Fab fragments prepared from J22.9-xi were crystallized in complex with the purified 46 amino acid residue BCMA extracellular domain and the complex structure solved to 1.9 angstroms resolution. High quality electron density is observable for residues 6 to 41 of BCMA and shows an extensive interaction with J22.9-xi, primarily with the light chain of the antibody (Fig. 2B). This interface, which buries 740.4 square angstroms and involves one third of the BCMA residues, covers 12 of 16 residues of the identical epitope observed in the crystal structures of BMCA complexes with APRIL and sTALL1 (also known as BAFF), including the conserved DxL motif (Gordon N C, *et al.* (2003), BAFF/BlysS receptor 3 comprises a minimal TNF receptor-like module that encodes a highly focused ligand-binding site. *Biochemistry* 42(20): 5977-83, and Patel D R, *et al.* (2004), Engineering an APRIL-specific B Cell Maturation Antigen. *JBC* 279(16): 16727-35), providing clear rationalization of the blocking effect seen in the *in vitro* assays with BAFF (Fig. 2A). The interaction with J22.9-xi additionally comprises a

direct side chain contact with Ala20 and Pro23 in BCMA, residues not part of the binding epitope covered by BAFF and APRIL, and several water-mediated hydrogen bonds. The overall conformation of BCMA in all three structures is very similar, with a C-alpha rmsd of 1.4 angstroms between the J22.9-xi and APRIL complexes and 1.5 angstroms between the J22.9-xi and sTALL1 complexes; the respective C-alpha rmsds for the J22.9-xi BCMA binding epitope (residues 13 – 30) are 0.98 and 0.88 angstroms. Although recognizing the same BCMA epitope having the DxL motif at its core, the binding site of J22.9-xi is very different from those of sTALL1 and APRIL, as is the collection of interactions comprising the interface.

As can be seen in Fig. 2B and Tables 1 and 2, 19 amino acids from J22.9-xi (6 from the heavy chain (Table 1), 13 from the light chain (Table 2)) form direct linkages to 12 residues from the extracellular domain of CD269.

Table 1: Amino acid interaction list between heavy chain of J22.9-xi and BCMA. These interaction lists were generated using the software PDBsum (Laskowski R A (2009)).

Heavy chain		CD269
Trp33	>	His19
Glu50	>	His19
Leu99	> >	Leu17 Leu18
Tyr100		Leu18
Tyr101	> > >	Ala20 Ile22 Pro23
Ala106	>	Leu18

Table 2: Amino acid interaction list between light chain of J22.9-xi and BCMA. These interaction lists were generated using the software PDBsum (Laskowski R A (2009)).

Light chain		CD269
Ser31	>	Arg27 Thr32
Ala34	>	Leu17
Tyr36	>	Leu17
Phe49	>	Leu18 Asp15
Ser50	>	Tyr13 Asp15 Arg27
Ser52	>	Arg27
Ser67	>	Thr32
Leu53	>	Tyr13 Leu26 Arg27
Phe55	>	Leu18

WO 2015/166073

PCT/EP2015/059562

- 45 -

Gln89	>	Leu17
Tyr91	>	Asp15 Ser16 Leu17
Tyr94	>	His19
Leu96	>	Leu17

Table 3: Interaction list of the residues involved in CD269:APRIL und CD269:BAFF binding (residues NOT directly contacted by J22.9 are underlined). These interaction lists were generated using the software PDBsum (Laskowski R A (2009)).

5

<u>APRIL</u>	<u>CD269</u>
Asp121	<u>Leu35</u>
Asp123	<u>Pro33</u>
	<u>Pro34</u>
	<u>Leu35</u>
Asp164	<u>Asn31</u>
Thr166	Arg27
	<u>Ser30</u>
Phe167	Tyr13
	Leu18
	Ile22
	Leu26
	Arg27
	<u>Asn31</u>
Thr168	Leu18
	Leu26
Met169	Leu17
Gly170	Leu17
	His19
Gln171	Leu17
Arg186	Leu17
	Leu18
	His19
Cys187	Leu17
Ile188	Asp15
	Leu17
	Leu18
Asp196	Leu26
Arg197	Leu26
Tyr199	Leu18
Pro221	Leu17
	His19
Arg222	Asp15
	Leu17
	Arg27
Asn224	Thr32
Lys226	<u>Asn31</u>
His232	His19
<u>BAFF</u>	<u>CD269</u>

WO 2015/166073

PCT/EP2015/059562

- 46 -

Tyr22	Ser16
Asp62	<u>Asn31</u>
Lys63	<u>Ser30</u>
	<u>Asn31</u>
Thr64	Arg27
	<u>Ser30</u>
	<u>Asn31</u>
Tyr65	Tyr13
	Asp15
	Leu18
	Ile22
Ala66	Leu17
Met67	Leu17
Gly68	Leu17
Arg90	Leu17
	His19
Cys91	Leu17
Ile92	Leu17
	Leu18
Glu97	<u>Ser29</u>
	<u>Ser30</u>
	<u>Asn31</u>
Leu99	Ile22
	Leu26
Asn101	Leu18
Pro123	Ser16
	Leu17
Arg124	Tyr13
	Asp15
	Leu17
	Arg27
Glu125	Arg27
	<u>Thr25</u>
	<u>Pro34</u>
	<u>Leu35</u>
Asn126	Thr32
Asp132	His19

Table 4: Residues of the CD269 target bound by direct contacts of J22.9, APRIL and/or BAFF. Residues of the CD269 target directly contacted only by J22.9 are underlined (20, 23). Residues of the CD269 target NOT directly contacted by J22.9 are in bold type (30, 31, 33, 34, 35 for APRIL; 25, 29, 30, 31, 34, 35 for BAFF).

J22.9: 13, 15, 16, 17, 18, 19, 20, 22, 23, 26, 27, 32

APRIL: 13, 15, 17, 18, 19, 22, 26, 27, **30, 31, 32, 33, 34, 35**

BAFF: 13, 15, 16, 17, 18, 19, 22, **25**, 26, 27, **29, 30, 31, 32, 34, 35**

Table 5: J22.9 Water interactions (J22.9-xi:H2O:CD269). The data in table 5 was generated using the software LigPlot (Wallace and Laskowski, European Bioinformatics Institute). (sc = side chain H-bond; mc = main chain H-bond)

Light Chain	H₂O#	CD269
Ser31 (sc)	285	Thr32 (sc, mc)
	285, 286	Arg27 (sc)
	285, 286	Ser30 (sc)
Ser31 (mc)	283, 284	Arg27 (sc)
Asn32 (sc)	105	Asp15 (sc)
	105, 284	Arg27 (sc)
	56	Ser16 (sc)
Tyr36	66, 93, 450	Leu17 (mc)
Ser50 (sc)	105	Asp15 (sc)
Ser52 (sc)	286	Ser30 (sc)
	286	Arg27 (sc)
	286, 285	Thr32 (sc, mc)
Gly66 (mc)	287	Thr32 (sc)
	285, 286	Arg27 (sc)
	285, 286	Ser30 (sc)
Gln89 (sc)	66, 93, 450	Leu17 (mc)
Tyr91 (mc)	282	Ser16 (sc)
	282, 281	Ser16 (mc)
Tyr94 (sc)	281	Ser16 (mc)
	281, 282	Ser16 (sc)
Heavy Chain	H₂O#	CD269
Trp33 (mc)	42, 280	Leu17 (mc)
	183, 279, 26	Leu18 (mc)
Ser35 (sc)	42, 66, 93, 280, 450	Leu17 (mc)
Trp47 (sc)	93, 450	Leu17 (mc)
Glu50 (sc)	281	Ser16 (mc)
	281, 282	Ser16 (sc)
	450	Leu17 (mc)
	450, 280	Leu17 (mc)
Leu99 (mc)	280	Leu17 (mc)
Tyr101 (mc)	26	Leu18 (mc)

5

Strong cytotoxic efficacy of J22.9-xi is strongly decreased after deglycosylation

A luciferase-based cytotoxicity assay was established using the luciferase transduced MM.1S-Luc cell line. In this assay, bioluminescence is only detected from living cells since luciferase released by dead cells is unable to function due to the lack of ATP in the medium. PBMCs from healthy donors were isolated and mixed with MM.1S-Luc cells in a ratio of 20 to 1. After 4 hours the bioluminescence was measured.

10

With a selection of 4 unstimulated donor PBMC preparations, the in vitro cytotoxicity of J22.9-xi was determined. The cytotoxic potential varies slightly between PBMCs from different donors. Within 4 hours of incubation, cell lysis reached 18 to 35 % at a concentration of 125

ng/ml J22.9-xi. Increasing the J22.9-xi concentration to 1 µg/ml increased cell lysis up to 56% (Fig. 3a).

After deglycosylation of J22.9-xi (J22.9-xi-N-glycan) with PNGase F, the cytotoxic activity dropped to below 8 %, whereas the binding of J22.9-xi-N-glycan to BCMA-positive MM.1S cells remained unaltered (Fig. 3a,b).

J22.9-xi reduces tumor burden in xenografted mice and prolongs survival

We used NOD scid common gamma chain knock out (NSG) mice lacking functional B, T and NK cell populations. These mice, injected with 1×10^7 MM.1S-Luc cells intravenously, develop hind limb paralysis within 6 weeks (Fig. 4d-1). The day on which the first symptom appears, defines the day of killing.

After injection of 1×10^7 MM.1S-Luc in the tail vein, the mice were divided randomly into 3 groups. The first group (n=2) received no treatment until the end of the experiment, whereas the second (n=5) and the third (n=6) group received twice weekly injections of 200 µg of an isotype control or the J22.9-xi antibody, respectively. The antibodies were administered for a period of 6 weeks intraperitoneally (i.p.) starting with the day of tumor cell injection. Tumor growth was monitored once a week using the IVIS Spectrum. Bioluminescence was measured 3 minutes after i.p. injection of luciferin.

A similar course of tumor development was seen in both the untreated group and the group receiving the control antibody, whereas the group treated with J22.9-xi showed significantly less tumor burden, already beginning at the first measurement point at day six (Fig. 4a). In addition, this group showed a smaller overall tumor load during the whole monitoring period (Fig. 4b). Isotype control treated animals had a median survival of 46 days after cell injection. Mice receiving J22.9-xi lived an average of 26 days longer. This corresponds to an extended survival of 55 % compared with mice receiving the control antibody (Fig. 4c). Massive infiltrations of tumor cells into the spine and inguinal lymph nodes were seen in non-treated mice and in mice receiving the isotype control antibody by day 28 after cell injection (Fig. 4d-2).

Administration of 200 µg of an antibody to a mouse corresponds to approximately 10 mg/kg bodyweight. To test the efficacy of J22.9-xi at lower doses we divided MM.1S-Luc-xenografted mice into four groups. The first group (n=7) received 200 µg of the control antibody twice weekly, and groups 2, 3 (each n=3) and 4 (n=9) were injected with 2 µg, 20 µg or 200 µg twice a week, respectively. Injection and monitoring were performed as described above.

Although tumors developed as expected in the control group mice, dramatically restricted tumor growth was observed in the groups receiving 20 µg or 200 µg of J22.9-xi (Fig. 4e,f). An overview of the experimental timeline is provided in Fig. 4g.

Growth of established tumors arrests for 5 weeks during J22.9-xi treatment

Therapeutic administration was mimicked by delaying the start of antibody treatment to 5 days after tumor cell injection. The xenografted mice were divided into 2 groups (n=6). The animals received 200 µg per injection of either the isotype control or J22.9-xi antibody twice a week.

The first measurement was done at day 8 post cell injection. While there is no tumor-derived bioluminescence measurable to day 35 in the group receiving J22.9-xi (n=5), a steady increase in tumor load was seen in animals receiving the isotype control antibody (n=6) (Fig. 5a,b). Mice from the isotype control group survived an average of 56 days after the cell injection, whereas all mice receiving J22.9-xi are still alive at day 77 (Fig. 5c). An overview of the experimental timeline is provided in Fig. 5d.

Intensive early phase treatment with J22.9-xi prevents tumor growth for 7 weeks

In order to further assess the effect of treatment timing on tumor growth, different antibodies were administered for five consecutive days starting from the day of tumor cell injection. Subsequent to i.v. cell injection, the animals were divided randomly into 5 groups. Group 1 (n=5) was treated with 200 µg of the isotype control antibody per injection (i.p.), whereas group 2 (n=6) received 200 µg/injection of the J22.9-xi-N-glycan antibody. The mice from groups 3 (n=4), group 4 (n=5) and group 5 (n=5) obtained 200 µg, 20 µg and 2 µg per injection of the J22.9-xi antibody, respectively. Bioluminescence measurements began at day 9 post cell injection. Up to day 44, no tumor-derived bioluminescence was seen in any of the groups receiving the intact J22.9-xi antibody. Although the tumor growth in the animals treated with J22.9-xi-N-glycan is decelerated, the overall tumor load is not significantly different from those animals receiving the isotype control antibody (Fig. 6a,b). Although the overall tumor load of animals treated with J22.9-xi-N-glycan (deglycosylated) was not significantly different (Fig 6b), the lifespan of these mice was substantially increased compared to the isoAb-treated group (Fig 6c). Since J22.9-xi-N-glycan was shown to be unable to induce ADCC or CDC, this result indicates that alone the binding of J22.9-xi to BCMA hinders tumor growth. It may be reasonably considered that this is due to blocking of the interaction between the receptor and its native ligands (APRIL and BAFF). An overview of the experimental timeline is provided in Fig. 6d.

Humanisation of J22.9-xi

The J22.9-xi antibody was humanized based on sequence alignment and the data obtained from the crystal structure. The sequences of the variable regions were aligned to their respective human homologs using IgBLAST (NCBI) or Clustal (EBI). Each proposed mutation was evaluated by visual inspection of the structure before alteration.

Binding of humanized variants to BCMA target

Binding of the mutants to BCMA was tested using flow cytometry, ELISA and SPR. The affinity of the humanized antibodies was measured using surface plasmon resonance (ProteOnTM XPR36; Bio-Rad). The binding data show surprising results with respect to the specificity and affinity of the humanized antibody variants to the same epitope as tested for J22.9-xi binding. As shown in the table below, it was entirely surprising that the humanized antibodies as described herein exhibited comparable binding characteristics as the original chimeric antibody. The SPR data reveals that the affinities of the humanized variants are similar to those of the chimera and are sufficient to assume their clinical relevance in light of the data provided herein for the original chimeric antibody. A skilled person would not have expected

that through the modification of the CDRs during humanization of the chimera that the binding characteristics would be maintained to such an extent.

ELISA was carried out as described herein using BCMA-coated microtiter plates (1microg/ml). As observed in Figure 10, binding was comparable for all humanized variants in comparison to J22.9-xi using both human and cynomolgous BCMA.

Flow cytometry was also carried out using the humanized variants described herein and equivalent binding to both human and cynomolgous BCMA for all humanized variants tested was shown (refer Figure 11).

SPR analysis was also conducted and affinities measured for humanized antibody variants. As can be observed in the table below (table 6), the affinities of the humanized variants (J22.9-H corresponds to humanized sequence SEQ ID No. 27; J22.9-FSY corresponds to humanized and PTM modified SEQ ID No. 28; J22.9-ISY corresponds to humanized and PTM modified SEQ ID No. 29).

Table 6. SPR Data

Name	ELISA (human)	ELISA (cynomolgous)	Flow cytometry	Melting temperatures	Affinity (SPR) (human) (n=3)	Affinity (SPR) (cynomolgous) (n=2)
J22.9-xi	+++	+++	+++	86 / 94 °C	$2.8 \pm 0.7 \times 10^{-10} \text{ M}$	$2.7 \times 10^{-9} \text{ M}$
J22.9-H	++	+	nd	86 / 94 °C	$1.5 \pm 0.3 \times 10^{-8} \text{ M}$	$2.0 \times 10^{-7} \text{ M}$
J22.9-FSY	+++	+++	+++	87 / 94 °C	$2.2 \pm 0.3 \times 10^{-9} \text{ M}$	$2.0 \times 10^{-8} \text{ M}$
J22.9-ISY	+++	+++	+++	86 / 94 °C	$2.0 \pm 0.2 \times 10^{-9} \text{ M}$	$1.7 \times 10^{-8} \text{ M}$

Amino acid 54 in CDR2 of the J22.9 heavy chain:

In order to remove a potential post-translational modification site in the humanized J22.9,

residue D54 of the heavy chain CDR2 was mutated to asparagine (N), inadvertently creating a new potential modification site for N-linked glycosylation. The mutated heavy chain containing N54 migrated slower on SDS gels (Fig. 13), indicating a larger size and that the CDR was glycosylated.

- 5 The corresponding IgG, J22.9-FNY, nevertheless bound BCMA in FACS and ELISA, and was crystallized in complex with BCMA. Although not completely refined, the 2.7 Angstrom resolution structure shows clear electron density extending from the N54 side chain - consistent with a sugar modification of the residue. It is surprising that such a large extension of the side chain would not disrupt binding to BCMA and it could be expected from these observations that multiple and various amino acid substitutions would be tolerated at this position, potentially also derivatizations other than sugars.

METHODS

Cell lines and culture

- 15 The human multiple myeloma cell line MM.1S (Greenstein *et al.* (2003) Exp Hematol 31:271-282) was obtained from Prof. B. Dörken (MDC, Berlin, Germany). For *in vivo* monitoring of tumor cell growth, Luciferase and GFP were cloned into the pFU vector of the lentiviral vector system ViraPower (Invitrogen). Via GFP-expression of transduced cells, monoclonal cell lines were isolated using fluorescence-activated single cell sorting. Cell lines were cultured in RPMI-1640 medium without phenol red, containing 10 % fetal calf serum, 100 units/ml of penicillin, and 100 µg/ml of streptomycin (all from PAA).

The HEK293-6E cells, purchased from the National Research Council of Canada, were maintained in Freestyle F17 medium (Invitrogen) supplemented with 7.5 mM L-Glutamine (PAA), 0.1 % Pluronic F-68 (Invitrogen), and 25 µg/ml G418 (Invitrogen). Cells were grown in Erlenmeyer flasks (Corning) at 110 rpm and 37 °C in a 5 % CO₂ atmosphere.

25 Antibody production and purification

- To obtain a BCMA-binding antibody, standard hybridoma technique was used. 4 BL/6 wild type mice were immunized 6 times with incomplete Freund's adjuvant and 30 µg of the extracellular domain of human BCMA C-terminally fused to Glutathione S-transferase (GST). After cell fusion followed by a screening period the J22.9 hybridoma was shown to secrete an anti-BCMA antibody.

- Due to the instability of the hybridomas the variable regions of the light and heavy chain of hybridoma J22.9 were amplified and cloned upstream of the human kappa or the IgG1 constant domain genes, respectively. The chimeric J22.9-xi antibody was produced by transient cotransfection of 293-6E cells with a 1:2 DNA plasmid mixture encoding the light and heavy chains, respectively. In brief: 293-6E cells were resuspended to 1.7×10^6 cells/ml in serum free Freestyle F17 medium and transfected using polyethyleneimine at a final concentration of 1 µg/ml culture. Two days after transfection, cells were fed with 100 % of the transfection volume Freestyle F17 medium containing 1 % tryptone N1 (Organo Technie). At day 7 cells were harvested by centrifugation and the filtered (0.45 µm) culture medium was

passed over a 3.5 ml Protein A Sepharose column (Bio-Rad). The column was washed with 10 ml phosphate buffered saline (PBS) and antibody eluted by addition of 20 mM sodium acetate, 150 mM NaCl, pH 3.5. Fractions of 2 ml were collected directly into tubes containing 100 µl 1 M HEPES, pH 7.5 for neutralization. The final yield of full length IgG was approximately 40 mg/l culture.

Since hybridoma J22.9 lost the capacity to produce/secrete the anti-BCMA antibody (Fig. 8), the variable regions of the heavy and light chains were amplified using PCR and subsequently cloned at the 5' end of the human constant IgG1 and K light chain genes, respectively. Through co-transfection of 293-6E cells with these two plasmids, the chimeric J22.9-xi antibody was produced. The production of the antibody of the invention was therefore inherently difficult and not achievable by straightforward routine methods.

The isotype control antibody composed of the J22.9-xi heavy chain and a random chimeric kappa light chain was produced in parallel with the J22.9-xi antibody. This antibody was shown by ELISA and flow cytometry to be unable to bind to BCMA.

The N-linked oligosaccharide chains at Asn297 of the heavy chain of J22.9-xi were removed enzymatically using N-Glycosidase F (PNGase F) (NEB). 10 mg of J22.9-xi were incubated with 15,000 units PNGase F in 500 µl PBS (pH 7.4) for 36 hours at 37 °C followed by buffer exchange into sterile PBS.

Determination of binding and blocking capacities of J22.9-xi by Enzyme-Linked Immunosorbent Assays (ELISA)

Microtiter plates were coated with 10 µg/ml of the extracellular domain of human BCMA. Coated BCMA was detected with serial dilution of J22.9-xi and the isotype control ranging from 1 to 1000 ng. Binding of J22.9-xi or isotype control antibody to the coated BCMA was detected with horseradish peroxidase (HRP)-conjugated goat anti-human secondary antibody (Jackson ImmunoResearch, 109-035-098, dilution 1:5,000).

Microtiter plates were coated with 1 µg/ml of the extracellular domain of human or cynomolgous BCMA (hBCMA or cyBCMA, respectively). Coated BCMA was detected with serial dilution of J22.9-xi, J22.9-H, J22.9-ISY and J22.9-FSY ranging from 0.26 pM to 500 nM. Binding of antibodies to the coated BCMA was detected with horseradish peroxidase (HRP)-conjugated goat anti-human secondary antibody (Jackson ImmunoResearch, 109-035-098, dilution 1:5,000).

For the blocking experiment, 1 mg/ml of human recombinant BAFF fused to a His-tag (Biomol) was applied after the antibodies and washing and detected using the mouse anti-His tag (AbD Serotec, AD1.1.10, dilution 1:5,000, HRP-conjugated) antibody. All ELISAs were developed using BD OptEIA reagents A and B (BD Bioscience) and measured with a microplate spectrophotometer (BioTek) at 450 nm and 570 nm.

Flow cytometry analysis

For cell surface antigen detection experiments, self-made antibodies (J22.9-xi, J22.9-H, J22.9-ISY, J22.9-FSY and the isotype control) and commercially available mouse anti-His tag (AbD

Serotec, AD1.1.10, dilution 1:100, Alexa Fluor 488-conjugated) and goat anti-human IgG1 (Jackson ImmunoResearch, 109-116-098, dilution 1:400, PE-conjugated) antibodies and human recombinant BAFF fused to a His-tag (Biomol) were used. Experiments were performed on a FACSCalibur or a FACSCanto II flow cytometer (BD Bioscience). The data were analysed with Flowjo software version 7.6 (TreeStar Inc.).

Generation of Fab and Fab:BCMA complexes

(Fab)₂ fragments were generated from full length J22.9-xi IgG by incubation with pepsin. J22.9-xi was passed over a PD-10 buffer exchange column into 50 mM sodium acetate, pH 3.5 and pepsin added at 30 µg per milligram J22.9-xi. Incubation at 37 °C for 2.5 hours was sufficient to completely digest the fragment crystallizable (Fc) region and pepsin was inactivated by exchange over a PD-10 column into PBS (pH 7.2). The reduction of the (Fab)₂ fragments to individual Fabs was accomplished in PBS by addition of 2-Mercaptoethylamine (50 mM) in the presence of 5 mM ethylenediaminetetraacetic acid (EDTA). After incubation for 90 minutes at 37°C, the reduced cysteines were blocked by alkylation with 500 µM iodoacetamide for 30 minutes followed by buffer exchange into fresh PBS. The Fab fragments were combined with 1.5 molar equivalents of purified BCMA and the complexes isolated by size exclusion chromatography on a Superdex 75 16/60 column. Fractions were analyzed on 4-12 % SDS polyacrylamide gels and fractions containing both Fab and BCMA were pooled and concentrated for crystallization trials.

Crystallization of Fab:BCMA complexes

Concentrated complexes were supplemented with 0.5 molar equivalents of pure BCMA to ensure saturation and were subjected to crystallization screening. Initial Fab:BCMA crystallization conditions were identified from commercial screens (Qiagen) in 96-well sitting drop format plates using a Gryphon pipetting robot (200nl drops) and optimized in 24 well plates in hanging drops (2-3µl). The complex was concentrated to 8 mg/ml and crystallized in 21 % PEG 3350, 0.1 M BisTris pH 6.5 and 5 mM CuCl₂ at 20 °C. Crystals appeared after three days as clusters of thin plates and attained their final size (0.2 – 0.3 mm) within approximately 7 days. Clusters were separated and individual plates were flash frozen in liquid nitrogen in mother liquor with 20 % glycerol as cryoprotectant. Complete diffraction data was collected from a single crystal at the BESSY synchrotron of the Helmholtz Zentrum Berlin. The structure was solved to a resolution of 1.9 angstroms by molecular replacement using the experimental phases from the structure of Efalizumab (3EO9) as the search model. Data processing was performed with the ccp4 suite of programs, structure refinement was performed using Phenix (Adams P D, *et al.* (2010), Acta Cryst. D66: 213-221) and model building and assessment using Coot. (Emsley et al, Acta Crystallographica Section D - Biological Crystallography, 2010, 66:486-501) Images were made using PyMOL (The PyMOL Molecular Graphics System, Version 1.5.0.4 Schrödinger, LLC).

In vitro cytotoxicity assay

In this assay the cytotoxic effect of J22.9-xi was determined by measuring the luminescence of the remaining living cells in a bioluminescence reader. In short: freshly obtained human filter

buffy coats (FBC) were back-flushed by gravity with 160 ml elution buffer (PBS (pH 7.4) containing 5 mM Na₂-EDTA and 2.5 [w/v] sucrose). Mononuclear cells were isolated from the eluted cells by Ficoll gradient centrifugation. Mononuclear cells from the interphases were taken and washed twice in elution buffer. After erythrocyte lysis, PBMCs were washed again, counted and adjusted by dilution in RPMI/10% FCS w/o phenol red to 1×10^7 cells/ml. 5×10^4 MM.1S-Luc cells in 50 μ l RPMI were plated in microtiter plates. Ten minutes prior to the addition of 100 μ l PBMCs, the MM.1S-Luc cells were incubated with J22.9-xi or the isotype control antibody serial dilutions in a sample volume of 200 μ l. After addition of target cells, antibodies and effector cells, microtiter plates were centrifuged (300 x g) for 2 minutes at room temperature (RT) and stored at 37 °C with 5 % CO₂. Control wells were treated with 1 % Triton X instead of antibody for complete lysis. After 4 hours of incubation, 25 μ l of PBS with luciferin (250 ng/ml) were applied to each well, and the bioluminescence of the living cells was measured in a bioluminescence reader (Tecan). The specific cytotoxicity was calculated according to the following formula:

$$100 - [\text{value (J22.9-xi)} - \text{value (total lysis)}] / [\text{value (isotype control)} - \text{value (total lysis)}] * 100.$$

In vivo studies

NOD.Cg-Prkdcscid Il2rgtm1Wjl Tg(CSF2)2Ygy Tg(IL3)1Ygy Tg(KITLG)3YgyJGckRolyJ mice (NSG) from The Jackson Laboratory and CB17.Cg-Prkdcscid Lystbg/Crl mice from Charles River Deutschland (Sulzfeld, Germany) were used. Experiments were performed with mice between 8-14 weeks old. All animal studies were performed according to institutional and state guidelines, under specific pathogen-free conditions. In the experimental examples relating to treatment of established tumours and tumour treatment in the early phase of disease the CB17.Cg-Prkdcscid Lystbg/Crl mice were used. The phenotype of the two mice strains mentioned herein is very similar. The animals have no functional B-, T- and NK-cells. A slightly slower tumour growth was observed in the CB17.Cg mice, indicating an even more promising effect of the therapeutic antibody of the present invention.

The xenograft model of multiple myeloma was induced by intravenous injection of 1×10^7 MM.1S-Luc cells in the tail vein at day zero. In this model, untreated animals develop hind limb paralysis within 6 weeks. Occurrence of this symptom indicates the end point of the experiment.

For the efficacy studies, the antibodies were administered intraperitoneally (i.p.) twice a week or on 5 consecutive days starting at day zero. The J22.9-xi antibody was given in doses of 2 μ g, 20 μ g or 200 μ g per injection; for the isotype control antibody, 200 μ g/injection was used. The bioluminescence of the MM.1S-Luc cells was measured after i.p. injection of 150 μ g luciferin using the IVIS Spectrum (Caliper Life Sciences). Measurements were done weekly. At each timepoint, 3 untreated control mice were also administered luciferin. Total flux values of these animals are either subtracted from each measurement or shown in the graphs.

To treat established tumors, antibody therapy was begun 5 days after injection of the MM.1S-Luc cells. 200 μ g of the J22.9-xi or isotype control antibody was administered twice a week for a period of 6 weeks.

Humanization of J22.9-xi

The heavy and light chain variable region sequences (mouse) were aligned with those from the corresponding heavy and light chain subtype human sequences to determine which residue alterations were required to produce a fully humanized sequence variant. Using the crystal structure of the J22.9-xi:hBCMA complex, each modification was first assessed *in silico* to identify mutations that could potentially disrupt binding of the antibody to BCMA. Two complete J22.9 variable region genes for each chain were synthesized, one with the original mouse sequence and one with a completely humanized sequence (i.e. containing all of the necessary humanizing mutations) with two added restriction enzyme sites to divide the genes into three cassettes each. After flagging potentially problematic mutations, various combinations of the original mouse and fully humanized gene cassettes were produced and their corresponding IgGs were expressed, purified and subjected to FACS analysis with BCMA positive cells to assess binding. Flagged problematic residues were mutated individually using PCR to verify their effect on affinity to BCMA and the final optimized constructs were subsequently quantitatively assessed for binding to both human and cynomolgus BCMA via SPR.

SPR

SPR was performed on a ProteonXPR36 using phosphate buffered saline supplemented with 0.005% Tween-20 (PBST). Whole IgG at a concentration of 15 ug/ml was immobilized to a Proteon GLH sensor chip using standard amine chemistry according to the manufacturer's instructions. For binding experiments, human or cynomolgus BCMA in PBST was used as the mobile phase. Binding affinities (K_d) were calculated from association (k_{on}) and dissociation (k_{off}) constants determined in parallel at multiple concentrations of BCMA (ranging from 0.4 to 800 nM for hBCMA and 2.7 nM to 1 μ M for cynoBCMA) assuming a single-site binding model. Additionally, further experimentation shows that the preferred embodiments of the invention provide surprising and unexpected effects, thereby solving the problem of the invention in a non-obvious fashion.

References

- Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., *et al.* (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* 66, 213-221.
- Al-Lazikani, B., Lesk, A.M., and Chothia, C. (1997). Standard conformations for the canonical structures of immunoglobulins. *J Mol Biol* 273, 927-948.
- Anthony, R.M., and Ravetch, J.V. (2010). A novel role for the IgG Fc glycan: the anti-inflammatory activity of sialylated IgG Fcs. *J Clin Immunol* 30 Suppl 1, S9-14.
- Chan *et al.* (2010) *Nat Rev Immunol* 10:301-316
- Chavez-Galan, L., Arenas-Del Angel, M.C., Zenteno, E., Chavez, R., and Lascurain, R. (2009). Cell death mechanisms induced by cytotoxic lymphocytes. *Cell Mol Immunol* 6, 15-25.
- Chiu, A., Xu, W., He, B., Dillon, S.R., Gross, J.A., Sievers, E., Qiao, X., Santini, P., Hyjek, E., Lee, J.W., *et al.* (2007). Hodgkin lymphoma cells express TACI and BCMA receptors and generate survival and proliferation signals in response to BAFF and APRIL. *Blood* 109, 729-739.
- Gordon, N.C., Pan, B., Hymowitz, S.G., Yin, J., Kelley, R.F., Cochran, A.G., Yan, M., Dixit, V.M., Fairbrother, W.J., and Starovasnik, M.A. (2003). BAFF/BLyS receptor 3 comprises a minimal TNF receptor-like module that encodes a highly focused ligand-binding site. *Biochemistry* 42, 5977-5983.
- Greenstein, S., Krett, N.L., Kurosawa, Y., Ma, C., Chauhan, D., Hideshima, T., Anderson, K.C., and Rosen, S.T. (2003). Characterization of the MM.1 human multiple myeloma (MM) cell lines: a model system to elucidate the characteristics, behavior, and signaling of steroid-sensitive and -resistant MM cells. *Exp Hematol* 31, 271-282.
- Jacobi, A.M., Huang, W., Wang, T., Freimuth, W., Sanz, I., Furie, R., Mackay, M., Aranow, C., Diamond, B., and Davidson, A. (2010). Effect of long-term belimumab treatment on B cells in systemic lupus erythematosus: extension of a phase II, double-blind, placebo-controlled, dose-ranging study. *Arthritis Rheum* 62, 201-210.
- Kapoor, P., Ramakrishnan, V., and Rajkumar, S.V. (2012). Bortezomib combination therapy in multiple myeloma. *Semin Hematol* 49, 228-242.
- Keyser, F.D. (2011). Choice of Biologic Therapy for Patients with Rheumatoid Arthritis: The Infection Perspective. *Curr Rheumatol Rev* 7, 77-87.
- Laskowski, R.A. (2009). PDBsum new things. *Nucleic Acids Res* 37, D355-359.
- Mackay, F., Schneider, P., Rennert, P., and Browning, J. (2003). BAFF AND APRIL: a tutorial on B cell survival. *Annu Rev Immunol* 21, 231-264.
- Nimmerjahn, F., and Ravetch, J.V. (2008). Fcgamma receptors as regulators of immune responses. *Nat Rev Immunol* 8, 34-47.
- Novak, A.J., Darce, J.R., Arendt, B.K., Harder, B., Henderson, K., Kindsvogel, W., Gross, J.A., Greipp, P.R., and Jelinek, D.F. (2004). Expression of BCMA, TACI, and BAFF-R in multiple myeloma: a mechanism for growth and survival. *Blood* 103, 689-694.
- Queen *et al.*, 1989; WO 90/07861

- Patel, D.R., Wallweber, H.J., Yin, J., Shriver, S.K., Marsters, S.A., Gordon, N.C., Starovasnik, M.A., and Kelley, R.F. (2004). Engineering an APRIL-specific B cell maturation antigen. *J Biol Chem* 279, 16727-16735.
- Presta, L.G. (2008). Molecular engineering and design of therapeutic antibodies. *Curr Opin Immunol* 20, 460-470.
- Raab, M.S., Podar, K., Breitkreutz, I., Richardson, P.G., and Anderson, K.C. (2009). Multiple myeloma. *Lancet* 374, 324-339.
- Richardson *et al.* (2003) *New Engl J Med* 348:2609-2617.
- Ryan et al (Molecular Cancer Therapeutics, 2007 6(11), 3009)
- Roopenian, D.C., and Akilesh, S. (2007). FcRn: the neonatal Fc receptor comes of age. *Nat Rev Immunol* 7, 715-725.
- Suzuki, K. (2013). Current therapeutic strategy for multiple myeloma. *Jpn J Clin Oncol* 43, 116-124.
- Thorpe *et al.*, 1982, *Immunol. Rev.* 62:119-58
- Wang, S.Y., and Weiner, G. (2008). Complement and cellular cytotoxicity in antibody therapy of cancer. *Expert Opin Biol Ther* 8, 759-768.
- Woyke et al. (2001) *Antimicrob. Agents and Chemother.* 45(12):3580-3584

CLAIMS

1. Antibody or antibody fragment, comprising a VH domain that comprises CDR sequences of:
 - RYWX₁S (H-CDR1; SEQ ID NO. 15), wherein X₁: I, F, L, V, Y, C, G, A, S, T);
 - EINPX₂X₃STINYAPSLKDK (H-CDR2; SEQ ID No. 16), wherein X₂X₃: SS, NS, TS, GS, KS, RS, SD, SN, DE; and
 - SLYX₄DYGDAX₅DYW (H-CDR3; SEQ ID NO. 17), wherein X₄: Y, L, A, V, F, I, W, and/or X₅: Y, L, F, I, V, A, C,

wherein said antibody or fragment thereof specifically binds an epitope of the extracellular domain of CD269 (BCMA).
2. Antibody or antibody fragment according to any one of the preceding claims, wherein the VH domain comprises the CDR sequence RYWIS (SEQ ID NO. 18) or RYWFS (SEQ ID NO. 19).
3. Antibody or antibody fragment according to any one of the preceding claims, wherein said VH domain comprises the CDR sequence EINPNSSTINYAPSLKDK (SEQ ID No. 20) or EINPSSSTINYAPSLKDK (SEQ ID No. 21).
4. Antibody or antibody fragment according to any one of the preceding claims, wherein said VH domain comprises the CDR sequence SLYYDYGDAYDYW (SEQ ID NO. 22).
5. Antibody or antibody fragment, comprising a VL domain that comprises CDR sequences of:
 - KASQSVX₁X₂NVA (L-CDR1; SEQ ID NO. 23), wherein X₁X₂: ES, SS, TS, QS, HS, DH;
 - SASLRFS (L-CDR2; SEQ ID NO 24); and
 - QQYNNYPLTFG (L-CDR3; SEQ ID NO. 25),

wherein said antibody or fragment thereof specifically binds an epitope of the extracellular domain of CD269 (BCMA).
6. Antibody or antibody fragment according to any one of the preceding claims, wherein the VL domain comprises the CDR sequence KASQSVDSNVA (SEQ ID NO. 26).
7. Antibody or antibody fragment according to any one of the preceding claims, comprising a VH domain that comprises the sequence
 EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWX₁SWVRQAPGKGLVWVGEINPX₂
 X₃STINYAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASLYX₄DYGDAX₅DYW
 GGGLTVTVSS (SEQ ID NO. 4), wherein X₁: I, F, L, V, Y, C, G, A, S, T; X₂X₃: SS, NS, TS, GS, KS, RS, SD, SN, DE, preferably SS; X₄: Y, L, A, V, F, I, W; and X₅: Y, L, F, I, V, A, C.
8. Antibody or antibody fragment according to any one of the preceding claims, comprising a VH domain that comprises the sequence according to SEQ ID NO. 6, SEQ ID NO. 7, SEQ ID NO. 8 or SEQ ID NO. 9.

9. Antibody or antibody fragment according to any one of the preceding claims, comprising a VL domain that comprises the sequence
EIVMTQSPATLSVSPGERATLSCASQSVX₁X₂NVAWYQQKPGQAPRALIYSASLRFS
GIPARFSGSGSGTEFTLTISLQSEDFAVYYCQQYNNYPLTFGAGTKLELKR (SEQ ID NO. 12), wherein X₁X₂: ES, SS, TS, QS, HS, DH.
10. Antibody or antibody fragment according to any one of the preceding claims, comprising a VL domain that comprises the sequence according to SEQ ID NO. 14.
11. Antibody or antibody fragment according to any one of the preceding claims, comprising a VH domain that comprises the sequence according to SEQ ID NO. 6, SEQ ID NO. 7, SEQ ID NO. 8 or SEQ ID NO. 9 and a VL domain that comprises the sequence according to SEQ ID NO. 14.
12. Antibody or antibody fragment according to any one of the preceding claims comprising a VH domain, wherein said VH domain comprises a sequence according to X₁VQLX₂X₃SGGGLVQPGGSLX₄LSCAASGX₅X₆FX₇X₈YWZ₁SWVRX₉APGKGLEWX₁₀GEINPZ₂SSTINYAPSLKX₁₁X₁₂FX₁₃ISRDNAKNTLYLQMX₁₄X₁₅X₁₆RX₁₇EDTAX₁₈YYCASLYYDYGDAZ₃DYWGQGTGX₁₉VTVSS (SEQ ID No. 41), wherein X₁: Q, E; X₂: Q, V; X₃: Q, E; X₄: K, R; X₅: I, F; X₆: D, T; X₇: S, D; X₈: R, D; X₉: R, Q; X₁₀: I, V; X₁₁: D, G; X₁₂: K, R; X₁₃: I, T; X₁₄: S, N; X₁₅: K, S; X₁₆: V, L; X₁₇: S, A; X₁₈: L, V; X₁₉: S, L; and wherein at least one of Z₁: I or F; Z₂: S and/or Z₃: Y; and wherein said antibody or fragment thereof specifically binds an epitope of the extracellular domain of CD269 (BCMA).
13. Isolated antibody or antibody fragment according to any one of the preceding claims comprising a VL domain, wherein said VL domain comprises a according to DIVMTQSX₁X₂X₃X₄X₅X₆SVGD₇VX₈X₉TCKASQSVE SNVAWYQQKPX₁₀QX₁₁PKX₁₂LIX₁₃SX₁₄X₁₅LRFSGVPARFX₁₆GSGSGTDFTLTISX₁₇LQSEDX₁₈AX₁₉YX₂₀CQQYNNYPLTFGAGTKLELKR (SEQ ID No. 42), wherein X₁: Q, P; X₂: R, A; X₃: F, T; X₄: M, L; X₅: T, S; X₆: T, V; X₇: R, E; X₈: S, T; X₉: V, L; X₁₀: R, G; X₁₁: S, A; X₁₂: A, L; X₁₃: F, Y; X₁₄: A, D; X₁₅: S, D; X₁₆: T, S; X₁₇: N, S; X₁₈: L, F; X₁₉: E, V; X₂₀: F, Y; and wherein said antibody or fragment thereof specifically binds an epitope of the extracellular domain of CD269 (BCMA).
14. Isolated antibody or antibody fragment according to any one of the preceding claims, wherein the antibody binds an epitope consisting of amino acids 13, 15, 16, 17, 18, 19, 20, 22, 23, 26, 27 or 32 of CD269.
15. Isolated antibody or antibody fragment according to any one of the preceding claims, wherein the antibody binding to CD269 (BCMA) disrupts BAFF-CD269 and/or APRIL-CD269 interaction.
16. Isolated antibody or antibody fragment according to any one of the preceding claims, wherein the antibody is glycosylated.
17. Isolated antibody or antibody fragment according to the preceding claim, wherein the glycan is an N-linked oligosaccharide chain at Asn297 of the heavy chain.
18. Isolated antibody or antibody fragment according to any one of the preceding claims for use as a medicament in the treatment of a medical disorder associated with the presence of pathogenic B cells.

19. Isolated antibody or antibody fragment according to any one of the preceding claims, wherein the medical disorder associated with pathogenic B cells is a disease of plasma cells and/or memory B cells.
20. Isolated antibody or antibody fragment according to any one of the preceding claims, wherein the medical disorder associated with pathogenic B cells is a cancer of plasma cells, for example multiple myeloma, plasmacytoma, Waldenström macroglobulinemia or plasma cell leukemia, or a cancer of B lymphocytes, such as Hodgkin's disease.
21. Isolated antibody or antibody fragment according to any one of the preceding claims, wherein the medical disorder associated with pathogenic B cells is an autoimmune disease associated with autoreactive plasma cells and/or auto reactive memory B cells, such as an inflammatory autoimmune disease, for example systemic lupus erythematosus (SLE) or rheumatic arthritis.
22. An antibody-drug conjugate (ADC) comprising the antibody or antibody fragment according to any one of the preceding claims conjugated to a therapeutic agent.
23. A nucleic acid molecule selected from the group comprising:
 - a) a nucleic acid molecule comprising a nucleotide sequence
 - which encodes an isolated antibody or antibody fragment according to any one of the preceding claims,
 - which encodes an amino acid sequence selected from the group consisting of those sequences according to SEQ ID 1 to 31 and 41 to 42,
 - comprising a sequence or sequence fragment of SEQ ID No. 32 to 36,
 - b) a nucleic acid molecule which is complementary to a nucleotide sequence in accordance with a);
 - c) a nucleic acid molecule comprising a nucleotide sequence having sufficient sequence identity to be functionally analogous/equivalent to a nucleotide sequence according to a) or b), comprising preferably a sequence identity to a nucleotide sequence according to a) or b) of at least 80%, preferably 90%, more preferably 95% ;
 - d) a nucleic acid molecule which, as a consequence of the genetic code, is degenerated into a nucleotide sequence according to a) through c); and
 - e) a nucleic acid molecule according to a nucleotide sequence of a) through d) which is modified by deletions, additions, substitutions, translocations, inversions and/or insertions and functionally analogous/equivalent to a nucleotide sequence according to a) through d).
24. A host cell, such as a bacterial cell or mammalian cell, preferably a hybridoma cell or cell line, capable of producing an antibody or antibody fragment and/or comprising a nucleic acid molecule according to any one of the preceding claims.
25. Pharmaceutical composition comprising the isolated antibody or antibody fragment, a nucleic acid molecule, an ADC or a host cell according to any one of the preceding claims, together with a pharmaceutically acceptable carrier.

Fig. 1




Fig. 2



Fig. 2 B




Fig. 3





Fig. 4



WO 2015/166073

PCT/EP2015/059562

6/20

Fig. 4 (cont.)

d-1






Fig. 4 (cont.)

d-2






Fig. 4 (cont.)



g

Experimental time line

tumor cell injection (i.v.)
1*10⁷ cells/mouse

weekly tumor monitoring (IVIS)





Fig. 5

**d****Experimental time line**

tumor cell injection (i.v.)
 1×10^7 cells/mouse

weekly tumor monitoring (IVIS)





Fig. 6



WO 2015/166073

PCT/EP2015/059562

11/20

Fig. 6 (cont.)

d

Experimental time linetumor cell injection (i.v.)
 1×10^7 cells/mouse

tumor monitoring (IVIS) every week





Fig. 7



WO 2015/166073

PCT/EP2015/059562

12/20

Fig. 8

HC

	1	10	20	30	40	50
HCg	XVQLXXSGGGLVQPGGSLXLSCAASGXXFXXYXXXWVRXAPGKGLXXXGX					
HCm	QVQLQQSGGGLVQPGGSLKLSCAASGIDFSRYWMSWVRRAPGKGLEWIGE					
HCpH	EVQLVESGGGLVQPGGSLRLSCAASGFTFDDYWMSWVRQAPGKGLEWVGE					
hHC01	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWMSWVRQAPGKGLVWVGE					
hHC02	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWXSWSVRQAPGKGLVWVGE					
hHC03	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYXMXWVRQAPGKGLVXVGX					
hHC04	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWISWVRQAPGKGLVWVGE					
hHC05	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWFSWVRQAPGKGLVWVGE					
hHC06	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWISWVRQAPGKGLVWVGE					
hHC07	EVQLVESGGGLVQPGGSLRLSCAASGFTFSRYWFSWVRQAPGKGLVWVGE					

	51	60	70	80	90	100
HCg	INPXXSTINYAPSLKXXFXISRDNAKNTLYLQMXXRSEDTAXYYCASXX					
HCm	INPDSSTINYAPSLKDKFTISRDNKNTLYLQMSKVRSEDTALYYCASLY					
HCpH	INPDSSTINYAPSLKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCASLY					
hHC01	INPDSSTINYAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASLY					
hHC02	INPXXSTINYAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASLY					
hHC03	INPDSSTINYAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASXX					
hHC04	INPNSSTINYAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASLY					
hHC05	INPNSSTINYAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASLY					
hHC06	INPSSSTINYAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASLY					
hHC07	INPSSSTINYAPSLKDKFTISRDNKNTLYLQMNSLRAEDTAVYYCASLY					

	101	110
HCg	XDYGDXXDYWGQGTXTVTVSS	
HCm	YDYGDAMDYWGQGTSTVTVSS	
HCpH	YDYGDAMDYWGQGTTLVTVSS	
hHC01	YDYGDAMDYWGQGTTLVTVSS	
hHC02	XDYGDAXDYWGQGTTLVTVSS	
hHC03	XDYGDXXMDYWGQGTTLVTVSS	
hHC04	YDYGDAYDYWGQGTTLVTVSS	
hHC05	YDYGDAYDYWGQGTTLVTVSS	
hHC06	YDYGDAYDYWGQGTTLVTVSS	
hHC07	YDYGDAYDYWGQGTTLVTVSS	

X: Variable amino acid according to humanized sequences

HCg: General heavy chain variable sequence

HCm: mouse heavy chain variable sequence

HCpH: partially humanized heavy chain variable sequence

CDRs are underlined in general and mouse sequences.

Fig. 9

LC

	1	10	20	30	40	50
LCg	<u>XIVMTQSXXXXXXXXSXGXXV</u> <u>XXXCKASQSVXXXVXWXQQK</u> <u>PXQXP</u> <u>KXLIXX</u>					
LCm	<u>DIVMTQSQR</u> <u>FMTTSVGDRVSV</u> <u>TCKASQSVDSN</u> <u>VAWYQQK</u> <u>PRQSP</u> <u>KALIFS</u>					
LCpH	<u>DIVMTQSPATLSV</u> <u>SVGDEV</u> <u>TLTCKASQSVDSN</u> <u>VAWYQQK</u> <u>PGQAP</u> <u>KLLIYS</u>					
hLC01	<u>EIVMTQSPATLSV</u> <u>SPGERAT</u> <u>LSCKASQSVDSN</u> <u>VAWYQQK</u> <u>PGQAP</u> <u>RALIYS</u>					
hLC02	<u>EIVMTQSPATLSV</u> <u>SPGERAT</u> <u>LSCKASQSVXXN</u> <u>VAWYQQK</u> <u>PGQAP</u> <u>RALIYS</u>					
hLC03	<u>EIVMTQSPATLSV</u> <u>SPGERAT</u> <u>LSCKASQSV</u> <u>DXVXWXQQK</u> <u>PGQAP</u> <u>RALIXX</u>					
hLC04	<u>EIVMTQSPATLSV</u> <u>SPGERAT</u> <u>LSCKASQSVESN</u> <u>VAWYQQK</u> <u>PGQAP</u> <u>RALIYS</u>					
	51	60	70	80	90	100
LCg	<u>XXXRXSGXPARFXGSXXG</u> <u>TXFTLTISXLQ</u> <u>SEDXAXYXCXQXNNXP</u> <u>XTFGA</u>					
LCm	<u>ASLRFS</u> <u>GVPARFTGSGSGT</u> <u>DFTLTISNLQ</u> <u>SEDLAEYFCQ</u> <u>QYNNYPL</u> <u>TFGA</u>					
LCpH	<u>DDLRFSGV</u> <u>PARFSGSGSGT</u> <u>DFTLTIS</u> <u>SLQSEDFAVYYC</u> <u>QYNNYPL</u> <u>TFGA</u>					
hLC01	<u>ASLRFS</u> <u>GIPARFSGSGSGT</u> <u>EFTLTIS</u> <u>SLQSEDFAVYYC</u> <u>QYNNYPL</u> <u>TFGA</u>					
hLC02	<u>ASLRFS</u> <u>GIPARFSGSGSGT</u> <u>EFTLTIS</u> <u>SLQSEDFAVYYC</u> <u>QYNNYPL</u> <u>TFGA</u>					
hLC03	<u>AXXR</u> <u>SGIPARFSGSXXG</u> <u>TFTLTIS</u> <u>SLQSEDFAVYYC</u> <u>XQXNNXP</u> <u>XTFGA</u>					
hLC04	<u>ASLRFS</u> <u>GIPARFSGSGSGT</u> <u>EFTLTIS</u> <u>SLQSEDFAVYYC</u> <u>QYNNYPL</u> <u>TFGA</u>					
	101					
LCg	GTKLELKR					
LCm	GTKLELKR					
LCpH	GTKLELKR					
hLC01	GTKLELKR					
hLC02	GTKLELKR					
hLC03	GTKLELKR					
hLC04	GTKLELKR					

X: Variable amino acid according to either mouse or humanized sequences

LCg: General heavy chain variable sequence


LCm: mouse heavy chain variable sequence

LCpH: partially humanized heavy chain variable sequence

CDRs are underlined in general and mouse sequences.

Fig. 10

Binding to hBCMA



Binding to cyBCMA




Fig. 11




Fig. 12 A




Fig. 12 A (cont.)




Fig. 12 B





Fig. 12 B (cont.)



WO 2015/166073

PCT/EP2015/059562

20/20

Fig. 13

