

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
14 December 2006 (14.12.2006)

PCT

(10) International Publication Number
WO 2006/131484 A1(51) International Patent Classification:
C07D 209/18 (2006.01) A61P 35/00 (2006.01)
A61K 31/405 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:
PCT/EP2006/062798

(22) International Filing Date: 31 May 2006 (31.05.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
05012562.4 10 June 2005 (10.06.2005) EP(71) Applicant (for all designated States except US):
SIGMA-TAU Industrie Farmaceutiche Riunite S.p.A. [IT/IT]; Viale Shakespeare, 47, I-00144 Rome (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PISANO, Claudio** [IT/IT]; Via Ceherubini, 60/B, I-04011 Aprilia (LT) (IT). **BATTISTUZZI, Gianfranco** [IT/IT]; Via Lucio Mario Perpetuo, 31, I-00174 Rome (IT). **DI MARZO, Maria** [IT/IT]; Via Nazionale, 214, I-81028 Santa Maria A Vico (CASERTA) (IT). **GIANNINI, Giuseppe** [IT/IT]; Via Boezio, 46/B, I-00040 Pomezia (RM) (IT). **MARZI, Mauro** [IT/IT]; Via Cina, 451, I-00144 Rome (IT). **VESCI, Loredana** [IT/IT]; Via Elio Petri, 3, I-00128 Rome (IT). **ZUNINO, Franco** [IT/IT]; Via dei Valtorta, 10, I-20127 Milan (IT).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:


- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:


- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: INDOLE DERIVATIVES HAVING ANTITUMOR ACTIVITY



(57) Abstract: Indole derivatives of Formula (I+A) having antitumour and chemosensitizing activity are described. Also described are pharmaceutical compositions containing the above-mentioned compounds, for the treatment of tumours.



INDOLE DERIVATIVES HAVING ANTITUMOUR ACTIVITY

FIELD OF THE INVENTION

The present invention relates to indole derivatives having antitumour activity as well as to the pharmaceutical compositions containing the above-mentioned compounds, for the treatment of tumours

BACKGROUND OF THE INVENTION

The therapy of tumours is being currently achieved by surgical intervention, radiation treatment and chemotherapy. The drawbacks of this latter are mainly due to the toxicity of the cytotoxic drugs, which is usually not limited to the cancer cells, and to the acquired resistance of the cancer cells to some of the most widely used drugs, which reduces the long-term efficacy of the therapy.

The elimination of the primary tumour by surgery is not always possible and in any case does not prevent the most metastasizing tumours, such as for example breast cancer or melanoma, to invade other target organs.

It has become evident that the therapy of the metastasizing tumours is unlikely to bring to the complete cure of the patient; therefore, the treatment with cytotoxic drugs is now seen as a palliative and life-prolonging method rather than a curative method. A chronic treatment with a drug having low toxicity would be preferable, while targeted to the control of the progression of the disease.

During the last years cancer drug development has moved from conventional cytotoxic chemotherapeutics to a more mechanism-based targeted

approach towards the common goal of tumour growth arrest. The rapid progress in chromatin research and understanding epigenetic control has supplied a plethora of potential targets for intervention in cancer.

Histone deacetylases (HDACs) have been widely implicated in growth 5 and transcriptional control, and inhibition of HDAC activity using small molecules causes apoptosis in tumour cells. Histone deacetylase inhibitors are now known to be potent inducers of growth arrest, differentiation, or apoptotic cell death in a variety of transformed cells in culture and in tumour bearing animals (Marks, P.A., Current Opinions in Oncology, 2001, Nov. 13 (6): 477-83; 10 Marks, P., Nat. Rev. Cancer 2001 Dec.1 (3):194-202).

On the other hand, as anticipated before, another very important and keenly perceived aspect of oncological therapy is the onset of resistance to the drug used by the tumour cells treated. The cells that develop resistance to a drug are often capable of resisting the effects of many other antitumour drugs, 15 even if these are unrelated chemically or act with different mechanisms of action. (Annu. Rev. Med 1991,42: 277-286; Drugs of the Future 1997,22: 653-660).

A number of tumours, such as, for instance, tumours of the adrenal cortex, colon, kidneys and jejunum and liver carcinoma manifest drug 20 resistance right from the very start of treatment with antitumour drugs (Barrows, L. R. Antineoplastic and Immunoactive Drugs, 1995 ; 75; 1236-1262).

In other cases, the tumour cells acquire resistance in a manner similar to that of bacterial resistance to antibiotics. This type of resistance has genetic or

epigenetic changes; these changes allow the daughter cells to proliferate in a *milieu* in which the antitumour agent is present.

Whatever the cause of the resistance, it leads to inefficacy of the antineoplastic treatment in the long term.

5 Patent application WO99/00381 describes bis-indole derivatives with antimetastatic activity.

Patent application WO02/36561, filed in the name of the applicant discloses compounds bis-heterocyclic compounds useful as antitumour agents.

In conclusion, in the oncology field there is a strongly perceived need for
10 new compounds endowed with antitumour and/or chemosensitising activity, i.e. compounds which are active against drug-resistant tumours and/or capable of making known antitumour drugs active against tumours against which they were ineffective owing to the onset of the above-mentioned conditions of drug resistance.


15 **DESCRIPTION OF THE INVENTION**

We have found that a class of indole derivatives possess the requisites essential for such antitumour, antimetastatic and chemosensitizing activity.

The compounds of the present invention showed to be more potent than the homologue compounds described in the patent application WO02/36561, to
20 inhibit the proliferation of NB4 promyelocytic leukaemia cells (see the results presented in Table 1 and Table 2 under the section Examples). These surprising anti-proliferative results, revealed for this new class of compounds an

anticancer potentiality for the treatment of cancer patients resistant to the therapies currently used.


Therefore the main object of the present invention are the indole compounds of Formula (I) below, which are useful agents as antitumour, 5 antimetastatic and chemosensitiser agents.



(I)

where


10



15

- A₁ is either H or A, wherein R, R' and R₁ are the same for both indole rings;
- X is a saturated or unsaturated (alkenylene or alkynylene), linear or branched (C₂-C₁₀) alkylene, optionally substituted with OH, or a (C₆-C₁₂) aryl, (C₃-C₁₀) heteroaryl containing at least one heteroatom selected from

N, O or S, wherein in the cyclic groups at least one of the -CH- is optionally substituted with C-halogen or C-(C₁-C₃) alkyl;



5 where,

- G is either H or a glycosyl;
- R₂, R₃ are the same or different and are either H or a (C₁-C₄) alkyl;
- R₁ is selected from the group consisting of H, (C₁-C₄) alkyl, (C₆-C₁₂) aryl, (C₆-C₁₂) aryl-(C₁-C₄) alkylene, (C₁-C₄) alkanoyl and (C₁-C₄)-alkyl-(C₆-C₁₀)-arylene;
- R and R', the same or different, are selected from the group consisting of:
 - H;
 - saturated or unsaturated, linear or branched (C₁-C₁₀) alkyl, optionally substituted with a (C₃-C₁₀) heteroaryl or (C₃-C₁₀) heterocyclyl-(C₁-C₄) alkylene, where the heterocycle contains at least one heteroatom selected from N, O or S, or with a group -NR₅R₆, where R₅, R₆ are the same or different and are H, linear or branched (C₁-C₄) alkyl, (C₁-C₄)-alkanoyl;
 - OR₄ where R₄ = H, (C₁-C₄) alkyl, mesyl, tosyl, (C₁-C₄) alkanoyl, glycosyl;

- halogen, azide, nitro, nitrile and NR_5R_6 .

The present invention also comprises tautomers, geometrical isomers, optically active forms as enantiomers, diastereomers and racemate forms, as well as pharmaceutically acceptable salts of the compounds of Formula (I).

5 Preferred pharmaceutically acceptable salts of the Formula (I) are acid addition salts formed with pharmaceutically acceptable acids like hydrochloride, hydrobromide, sulfate or bisulfate, phosphate or hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, citrate, tartrate, gluconate, methanesulfonate, benzenesulfonate, and para-toluenesulfonate salts.

10 Within the framework of the present invention, examples of the linear or branched ($\text{C}_1\text{-C}_4$) alkyl group, are understood to include methyl, ethyl, propyl and butyl and their possible isomers, such as, for example, isopropyl, isobutyl, and ter-butyl.

15 The term "alkylene" refers to a linear or branched chain divalent hydrocarbon radical. Examples of "alkylene" as used herein include, but are not limited to, methylene, ethylene, propane-1,3-diyl, propane-1,2-diyl and the like.

Examples of the ($\text{C}_6\text{-C}_{12}$) aryl or ($\text{C}_6\text{-C}_{12}$) aryl - ($\text{C}_1\text{-C}_4$) alkyl group are phenyl, 1- or 2-naphthyl, anthracenyl, benzyl, 2-phenylethyl 1-phenylethyl, 3-phenylpropyl, 2-anthracenylpropyl, 1-anthracenylpropyl, naphthylmethyl, 2-naphthylethyl, 1-naphthylethyl, 3-naphthylpropyl, 2-naphthylpropyl, 1-naphthylpropyl.

As used herein, the term " $(\text{C}_3\text{-C}_6)$ heterocyclo" or the term " $(\text{C}_3\text{-C}_6)$ heterocyclyl" refers to a monovalent three to six-membered non-aromatic ring


containing one or more heteroatomic substitutions independently selected from S, O, or N and having zero to five degrees of unsaturation. Examples of "heterocyclic" as used herein include, but are not limited to, tetrahydrofuryl, pyranyl, 1,4-dioxanyl, 1,3-dioxanyl, piperidinyl, pyrrolidinyl, 5 tetrahydrothiopyranyl, tetrahydrothiophenyl, and the like.

As used herein, the term "(C₃-C₆) heterocyclene" refers to a divalent three to six membered non-aromatic heterocyclic ring radical containing one or more heteroatoms independently selected from S, O, or N and having zero to five degrees of unsaturation. Examples of "heterocyclene" as used herein 10 include, but are not limited to, tetrahydrofuran-2,5-diyl, pyran-2,4-diyl, 1,4-dioxane-2,3-diyl, 1,3-dioxane-2,4-diyl, piperidine-2,4-diyl, piperidine-1,4-diyl, pyrrolidine-1,3-diyl, and the like.

What is meant by halogen is fluorine, chlorine, bromine and iodine.

Examples of the glycosyl residue are 6-D-galactosyl and 6-D-glucosyl.

15 According to independently preferred embodiments of the invention, X is a saturated or unsaturated (alkenylene or alkynylene), linear (C₂-C₁₀) alkylene; A₁ is A;



G, R₁, R₂ and R₃ are H;

and R and R' are the same or different and are H, linear (C₁-C₁₀) alkyl, optionally substituted with a (C₃-C₁₂) heteroaryl containing at least one heteroatom selected from N, O or S, or with a group -NR₅R₆ or OR₄ where R₄ is (C₁-C₄) alkyl.

5 The following are some of the most preferred compounds according to the invention:

3-{4-[Bis-(1H-indol-3-yl)-methyl]-phenyl}-N-hydroxy-acrylamide (ST2887),
5,5-Bis-(1H-indol-3-yl)-pentanoic acid hydroxyamide (ST2743),
6,6-Bis-(1H-indol-3-yl)-hexanoic acid hydroxyamide (ST2754),
10 7,7-Bis-(1H-indol-3-yl)-heptanoic acid hydroxyamide (ST2741),
7,7-Bis-(4-fluoro-1H-indol-3-yl)-heptanoic acid hydroxyamide (ST3052),
7,7-Bis-(5-methyl-1H-indol-3-yl)-heptanoic acid hydroxyamide (ST3044),
7,7-Bis-(7-ethyl-1H-indol-3-yl)-heptanoic acid hydroxyamide (ST3126),
7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid hydroxyamide (ST3043),
15 8,8-Bis-(1H-indol-3-yl)-octanoic acid hydroxyamide (ST2889),
N-Hydroxy-4,4-bis-(1H-indol-3-yl)-butyramide (ST2408),
N-Hydroxy-6-(1H-indol-3-yl)-hexanamide (ST2995),
7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid hydroxyamide
(ST3307),
20 7-(1H-Indol-3-yl)-7-(1H-indol-2-yl)-heptanoic acid hydroxyamide (ST3292), and
7,7-Bis-(1H-indol-3-yl)-heptanoic acid (ST3127).

The experimental results obtained (reported in the section entitled "Examples") show that the compounds of Formula (I), both alone and in

combination with other known antitumour drugs, are useful agents for the treatment of tumours.

A further object of the invention described herein are compounds with general Formula (I) and their use in the medical field.

5 A further object of the invention described herein is a pharmaceutical composition containing as active ingredient a compound of Formula (I) and at least a pharmaceutically acceptable excipient and/or diluent.

A further object of the invention described herein are compounds with general Formula (I) and a process for their preparation.

10 A further object of the invention described herein is a pharmaceutical composition containing as active ingredient a compound of Formula (I), for the treatment of a tumour pathology, in which the tumour is selected from the group consisting of sarcoma, carcinoma, carcinoid, bone tumour, neuroendocrine tumour, lymphoid leukaemia, acute promyelocytic leukaemia, myeloid 15 leukaemia, monocytic leukaemia, megakaryoblastic leukaemia and Hodgkin's disease.

A further object of the invention described herein is a pharmaceutical composition containing as active ingredient a compound Formula (I), for the treatment of a tumour pathology, in which the tumour has shown drug 20 resistance to the previous antibiotics used for its treatment, in which said compound of Formula (I) exerts a chemosensitising effect on said drug resistant tumour.

A further object of the invention described herein is a pharmaceutical composition containing as active ingredient a compound of Formula (I), in combination with one or more known antitumour agents, in which the antitumour compound is selected from the group consisting of alkylating agents, 5 topoisomerase inhibitors, antitubulin agents, intercalating compounds, anti metabolites, natural products such as vinca alkaloids, epipodophyllotoxins, antibiotics, enzymes, taxans, and cytodifferentiating compounds. Among the cytodifferentiating antitumour agents the one preferred is all-trans retinoic acid.

10 Another object of the present invention is a process for preparing the pharmaceutical compositions, as explained before, comprising mixing the compound(s) of Formula (I) with the suitable excipient(s) and/or diluent(s).

15 A further object of the invention described herein is the use of a compound of Formula (I) for the preparation of a medicine for the treatment of a tumour pathology.

20 A further object of the invention described herein is the use of a compound of Formula (I) for the preparation of a medicine for the treatment of a tumour pathology in which the tumour has shown drug resistance to the previous antitumour drugs used for its treatment, in which said compound of Formula (I) exerts a chemosensitising effect on said drug-resistant tumour.

A further object of the invention described herein is the use of a compound of Formula (I), in combination with one or more known antitumour

agents, for the preparation of a medicine for the treatment of tumour pathologies.

A further object of the invention described herein is the use of a compound of Formula (I) in combination with all-trans retinoic acid for the preparation of a medicine for the treatment of acute promyelocytic leukaemia.

Another object of the invention is a method of treating a mammal suffering from a tumour pathology, comprising administering a therapeutically effective amount of the compound(s) of Formula (I).

“Therapeutically effective amount” is an amount effective to achieve the medically desirable result in the treated subject. The pharmaceutical compositions may contain suitable pharmaceutical acceptable carriers, biologically compatible vehicles suitable for administration to an animal (for example, physiological saline) and eventually comprising auxiliaries (like excipients, stabilizers or diluents) which facilitate the processing of the active compounds into preparations which can be used pharmaceutical.

The pharmaceutical compositions may be formulated in any acceptable way to meet the needs of the mode of administration. The use of biomaterials and other polymers for drug delivery, as well the different techniques and models to validate a specific mode of administration, are disclosed in literature.

Modifications of the compounds of the invention to improve penetration of the blood-brain barrier would also be useful.

Any accepted mode of administration can be used and determined by those skilled in the art. For example, administration may be by various

parenteral routes such as subcutaneous, intravenous, intradermal, intramuscular, intraperitoneal, intranasal, transdermal, oral, or buccal routes.

Parenteral administration can be by bolus injection or by gradual perfusion over time. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions, which may contain auxiliary agents or excipients known in the art, and can be prepared according to routine methods. In addition, suspension of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, sesame oil, or synthetic fatty acid esters, for example, ethyloleate or triglycerides.

Aqueous injection suspensions that may contain substances increasing the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

Pharmaceutical compositions include suitable solutions for administration by injection, and contain from about 0.01 to 99 percent, preferably from about 20 to 75 percent of active compound together with the excipient. Compositions which can be administered rectally include suppositories.

It is understood that the dosage administered will be dependent upon the age, sex, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired. The dosage will be tailored to the individual subject, as is understood and determinable by

one of skill in the art. The total dose required for each treatment may be administered by multiple doses or in a single dose. The pharmaceutical composition of the present invention may be administered alone or in conjunction with other therapeutics directed to the condition, or directed to other 5 symptoms of the condition. Usually a daily dosage of active ingredient is comprised between 0.01 to 100 milligrams per kilogram of body weight.

The compounds of the present invention may be administered to the patient intravenously in a pharmaceutical acceptable carrier such as physiological saline.

10 Standard methods for intracellular delivery of peptides can be used, e. g. delivery via liposomes. Such methods are well known to those of ordinary skill in the art. The formulations of this invention are useful for parenteral administration, such as intravenous, subcutaneous, intramuscular, and intraperitoneal.

15 As well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently.

20 All references cited herein are entirely incorporated by reference herein, including all data, tables, figures, and text presented in the cited references.

Additionally, the entire contents of the references cited within the references cited herein are also entirely incorporated by reference. Reference to known method steps, conventional method steps, known methods or

conventional methods is not in any way an admission that any aspect, description or embodiment of the present invention is disclosed, taught or suggested in the relevant art.

Once understood the features of the methods and products disclosed in 5 present application, the necessity and kind of additional steps can be easily deduced by reviewing prior art, as well as the non-limiting following figures and examples describing the basic details and some applications of the invention


The compounds of Formula (I) may be prepared from readily available starting materials using the following general methods and procedures. It will be 10 appreciated that where typical or preferred experimental conditions (i.e. reaction temperatures, time, moles of reagents, solvents, etc.) are given, other experimental conditions can also be used, unless otherwise stated. Optimum reaction conditions may vary with the particular reactants or solvents used, but such conditions can be determined by one skilled in the art by routine 15 optimisation procedures.

A process for preparing the compounds of Formula (I), in which A is A1, comprises dimerizing the starting indole derivative in the presence of an aldheyde or acetal.

The compounds of the present invention can be prepared for example 20 according to the following general scheme.

GENERAL SCHEME I

[for Q = CONHOH; COOH; CH=CH-CONHOH; CONH(2-amino-phenyl)]



Y = CHO; CH(OR'')₂
 Z = OH; O-benzyl; 2-aminophenyl

5 and R'' is C₁-C₄ alkyl

General procedure for Step 1:

Step 1 is carried out in organic, polar or non-polar, protic or aprotic solvents, or mixtures of them, i.e. dichloromethane, dichloroethane, acetonitrile, methanol, ethanol, dimethylformamide, water, acetic acid. Reagent B is reacted 10 as free aldehyde or acetale and the reaction is catalysed by organic, inorganic, or Lewis acids, i.e. HCl (Herbert R *et al.*, *J Chem Soc*, 1969, 1505), trifluoroacetic acid [Tominaga Y *et al.*, *Heterocycles*, 2001, 55 (8), 1447-1450], acetic acid (Wang QM *et al.*, *Synlett*, 1995, 12, 1267-1268), dysprosium triflate [Chen D *et al.*, *Tetrahedron Lett*, 1996, 37 (26), 4467-4470], lithium perchlorate 15 [Yadav JS *et al.*, *Synthesis*, 2001, (5), 783-787]. Temperatures from -10 °C to 150 °C are used. Compounds C can be purified by flash silica gel chromatography.

General procedure for Step 2:

Step 2 is normally carried out in a mixture of water and organic solvents, such as methanol, ethanol, by using an inorganic base such as NaOH, LiOH. After stirring at room temperature, reaction mixture is acidified with HCl 1N and compounds D are purified by extraction with organic solvents, i.e. 5 dichloromethane, ethyl acetate.

General procedure for Step 3:

Step 3 is carried out in organic, polar or non-polar, aprotic solvents, or mixtures of them, i.e. dichloromethane, dichloroethane, dimethylformamide, tetrahydrofuran, dioxane. Different condensating and/or activating agents can 10 be used, such as DCC, EDC, HOBr, HATU, PyBOP [Lloyd-Williams P., Albericio F. and Giralt E., *Chemical Approaches to the synthesis of peptides and proteins*, (1997) and refs] and the reaction is catalysed by using an organic base, such as 15 N-methylmorpholine, N,N-diisopropylethylamine, triethylamine, DBU [Sandanayaka VP *et al.*, *Tetrahedron Lett*, 2001, 42 (28), 4605-4607. Bailen MA et al., *Tetrahedron Lett*, 2001, 42 (30), 5013-5016]. Temperatures from -10 °C to 100 °C are used. Compounds F are purified by silica gel chromatography or by reversed-phase high performance chromatography (RP-HPLC).

General procedure for Step 4:


If Z of intermediate F is a O-benzyl group, the reaction is carried out in 20 organic solvents, i.e. methanol, by using hydrogen pressures from 15 psi to 60 psi, and different catalysts such as Pd/C 10 %, Pd/BaSO₄ 5 or 10 % [Nikam SS *et al.*, *Tetrahedron Lett*, 1995, 36 (2), 197-200]. The final products G are

purified by solvent evaporation and eventually column chromatography on silica gel or high-performance reversed-phase chromatography.

When Q is $-\text{SH}$ the following Scheme II may be followed.


SCHEME II

5 [for Q = $-\text{SH}$]



According to the above Scheme II, the starting compound of step 1
10 (which corresponds to compound C of Scheme I) is treated with NaBH_4 .
According to step 2a the compound obtained through step 1 is reacted with
Lawesson's reagent [see Rajagopalan, S. et al., *Synth. Comm.* 1997, 27 (1),
187-194]. According to step 2b the compound obtained through step 1 is
reacted with CH_3COSH , DEAD, TPP [see Wisniewski, K. et al., *Bioorg. Med.*
15 *Chem. Lett.*, 1999 (9), 2801-2804]. In step 3 the compounds obtained through
step 2 a or step 2b are treated with NaOH to give the corresponding
compounds of Formula I.

When Q is $-\text{COCF}_3$ the following Scheme III may be followed.

SCHEME III

5


where W = A or Y and R'' = C₁-C₄ alkyl

According to the above Scheme III, the starting compound of step 1a is treated with (CH₃)₃SiCF₃ and NBu₄F. [see Wiedemann, J. *et al.*, *Angew. Chem. Int. Ed.*, 1998, 37 (6), 820-821]. If step 1b is performed, the starting compound 10 is treated with ClCOCOCl or SOCl₂; TFAA, Pyridine [see Boivin, J. *et al.*, *Tetrahedron* 1995, 51 (9), 2573-2584, or Frey, R.R. *et al.*, *Bioorg. Med. Chem. Lett.*, 2002 (12), 3443-3447].

Step 2 is carried out as step 1 of Scheme I [for Q = CONHOH; COOH; CH=CH-CONHOH; CONH(2-amino-phenyl)]

15 When Q is -COCONR₂R₃ the following Scheme IV may be followed.

SCHEME IV



where R" is C₁-C₄ alkyl

According to the above Scheme IV, the starting compound of step 1 is

5 treated with LiHMDS and Davis' oxaziridine.

According to step 2 the compound obtained through step 1 is reacted with R₂R₃NH.

According to step 3 the compound obtained through step 2 is reacted with Dess-Martin periodinane. [Wada, C.K. *et al.*, *Bioorg. Med. Chem. Lett.*,
10 2003 (13), 3331-3335].

Step 4 is carried out as step 1 in General Scheme I [for Q = CONHOH;
COOH; CH=CH-CONHOH; CONH(2-amino-phenyl)].

The following examples further illustrate the invention without limiting its scope.

EXAMPLES**EXAMPLE 1:****Preparation of 7,7-Bis-(1H-indol-3-yl)-heptanoic acid hydroxyamide (ST2741)**

Step 1. A solution of B (7-Oxo-heptanoic acid ethyl ester, 274.5 mg, 1.59 mmol), A (1H-Indole, 373.3 mg, 3.18 mmol) and Dysprosium triflate (1.45 g, 2.38 mmol) in MeOH/H₂O (7.5 mL/2.5 mL) was stirred for 36 h at room temperature. Then the reaction mixture was diluted with dichloromethane (DCM) and washed with H₂O (x 3 times). The organic layer was dried over sodium sulfate, filtered and evaporated under reduced pressure. The residue 10 was purified by column chromatography on silica gel using a gradient elution of hexane/ethyl acetate to give 0.4 g of C [7,7-Bis-(1H-indol-3-yl)-heptanoic acid ethyl ester, y = 65 %].

¹H-NMR (200 MHz, CDCl₃) δ (ppm): 1.30 (t, 3H, CH₃); 1.46 (br m, 4H, CH₂); 1.66 (br m, 2H, CH₂); 2.30 (m, 4H, CH₂); 4.18 (q, 2H, OCH₂); 4.51 (t, 1H, CH); 15 6.92 (s, 2H); 7.00-7.40 (m, 6H, CH arom); 7.65 (d, 2H, CH arom); 7.88 (s, 2H).

¹³C-NMR (50 MHz, CDCl₃) δ (ppm): 14.48; 25.18; 28.16; 29.47; 34.15; 34.60; 35.87; 60.43; 111.27; 119.07; 119.71; 120.37; 121.61; 121.78; 127.22; 136.67; 174.16.

ES-MS m/z: 411.6 [M+Na]⁺.

Step 2. To a solution of intermediate C [7,7-Bis-(1H-indol-3-yl)-heptanoic acid ethyl ester, 380 mg, 0.98 mmol] in THF: MeOH: H₂O 3:3:1 (3.8 mL) was added solid NaOH (235 mg, 5.88 mmol) and the reaction mixture was stirred at room temperature for 15 h. The solution was acidified (pH~1) with HCl 1N and then

extracted with DCM. The organic layer was washed with H₂O (x 3), then dried over sodium sulfate, filtered and evaporated under reduced pressure to give 0.33 g of D [7,7-Bis-(1H-indol-3-yl)-heptanoic acid, ST 3127, y = 93 %].

¹H-NMR (300 MHz, Acetone-d₆) δ (ppm): 1.45 (br m, 4H, CH₂); 1.59 (quint, 2H, CH₂); 2.26 (m, 4H, CH₂); 4.52 (t, 1H, CH); 6.87-6.95 (m, 2H, CH_{arom}); 6.99-7.07 (m, 2H, CH_{arom}); 7.21 (d, 2H, CH_{arom}); 7.34 (d, 2H, CH_{arom}); 7.59 (d, 2H, CH_{arom}); 9.89 (s, 2H, NH).

¹³C-NMR (75 MHz, Acetone-d₆) δ (ppm): 25.06; 28.14; 29.29; 33.57; 34.19; 35.81; 111.39; 118.42; 119.50; 119.95; 121.16; 121.99; 127.56; 137.39; 174.02.

10 ES-MS m/z: 359.2 [M-H]⁻ and 383.2 [M+Na]⁺.

Step 3. To a solution of intermediate D [7,7-Bis-(1H-indol-3-yl)-heptanoic acid, 140 mg, 0.388 mmol] and PyBOP (259.5 mg, 0.465 mmol) in DCM (1 mL) were added NMM (0.213 mL, 1.94 mmol) and E (O-benzylhydroxylamine hydrochloride, 74.3 mg, 0.465 mmol). The solution was stirred at room 15 temperature for 2.5 h, then diluted with ethyl acetate and washed with H₂O (x 3). The organic layer was dried over sodium sulfate, filtered and evaporated under reduced pressure. The residue was purified by a preparative TLC on silica gel using hexane/ethyl acetate 7:3 as eluent system, to give 0.12 g of F [7,7-Bis-(1H-indol-3-yl)-heptanoic acid benzyloxy-amide, y = 70 %].

20 ¹H-NMR (300 MHz, Acetone-d₆) δ (ppm): 1.41 (br m, 4H, CH₂); 1.57 (quint, 2H, CH₂); 2.02 (br m, 2H, CH₂); 2.26 (q, 2H, CH₂); 4.49 (t, 1H, CH); 4.84 (s, 2H, CH₂); 6.89 (m, 2H, CH_{arom}); 7.01 (m, 2H, CH_{arom}); 7.21 (s-d, [J = 1.59 Hz], 2H, CH); 7.28-7.42 (m, 7H, CH_{arom}); 7.57 (d, 2H, CH_{arom}); 9.92 (br s, 2H,

NH).

¹³C-NMR (75 MHz, Acetone-d₆) δ (ppm): 25.46; 28.13; 29.29; 32.91; 34.17; 35.81; 77.46; 111.32; 111.37; 118.34; 119.47; 119.94; 121.10; 121.81; 121.96; 127.52; 127.55; 128.35; 128.42; 129.12; 137.32; 161.09.

5 ES-MS m/z: 464.4 [M-H]⁻ and 488.5 [M+Na]⁺.

Step 4. Palladium on activated carbon (10%, cat.) was added to a solution of intermediate F [7,7-Bis-(1H-indol-3-yl)-heptanoic acid benzyloxy-amide, 79 mg, 0,170 mmol] in MeOH (4 mL). After 6 h of stirring under hydrogen atmosphere (15psi) the reaction mixture was filtered and the filtrate was evaporated under 10 reduced pressure. The residue was purified by preparative reversed-phase chromatography (column Lichrosorb RP-18, 7μm, eluents: CH₃CN/H₂O 1:1, flow = 10 mL/min) gave 50.0 mg of G [7,7-Bis-(1H-indol-3-yl)-heptanoic acid hydroxyamide, ST 2741, y = 78%].

15 ¹H-NMR (300 MHz, Acetone-d₆) δ (ppm): 1.42 (br, 4H, CH₂); 1.57 (quint, 2H, CH₂); 2.06 (br, 2H, CH₂); 4.48 (t, 1H, CH); 6.89 (m, 2H, CH arom); 6.99 (m, 2H, CH arom); 7.20 (s, 2H, CH arom); 7.33 (d, 2H, CH arom); 7.57 (d, 2H, CH arom); 9.92 (br, 2H, NH).

13C-NMR (75 MHz, Acetone-d₆) δ (ppm): 24.66; 31.65; 33.22; 34.84 (some signals under solvent signal); 110.37; 117.32; 118.45; 118.89; 120.09; 120.80; 20 120.94; 126.50; 136.28; 169.81.

ES-MS m/z: 374.4 [M-H]⁻.

EXAMPLE 2:

Preparation of 5,5-Bis-(1H-indol-3-yl)-pentanoic acid hydroxyamide (ST2743)

Step 1. A solution of A (1H-Indole, 1.12 g, 9.75 mmol), B [5,5-Dimethoxy-pentanoic acid methyl ester; 521.3 mg, 2.96 mmol] and Dysprosium triflate (2.64 g, 4.33 mmol) in MeOH/H₂O (6 mL/4 mL) reacted as in example 1 (at 60°C) to give 0.74 g of C [5,5-Bis-(1H-indol-3-yl)-pentanoic acid methyl ester, y 5 = 72 %].

¹H-NMR (200 MHz, CDCl₃) δ (ppm): 1.78 (m, 2H, CH₂); 2.28 (q, 2H, CH₂); 2.41 (t, 2H, CH₂); 3.68 (s, 3H, CH₃); 4.53 (t, 1H, CH); 6.95 (s, 2H); 7.00-7.32 (m, 6H, CH arom); 7.63 (d, 2H, CH arom); 7.89 (s, 2H, NH).

¹³C-NMR (50 MHz, CDCl₃) δ (ppm): 23.84; 34.02; 34.26; 35.28; 51.65; 111.25;

10 119.13; 119.64; 119.88; 121.69; 121.84; 127.11; 136.68; 174.40.

ES-MS m/z: 369.5 [M+Na]⁺.

Step 2. Intermediate C [5,5-Bis-(1H-indol-3-yl)-pentanoic acid methyl ester, 495 mg, 1.43 mmol] and solid NaOH (457 mg, 11.4 mmol) in MeOH/DCM (3 mL/1mL) reacted as in example 1 to give 0.45 g of D [5,5-Bis-(1H-indol-3-yl)-15 pentanoic acid, y = 95 %].

¹H-NMR (200 MHz, CDCl₃) δ (ppm): 1.78 (m, 2H, CH₂); 2.28 (q, 2H, CH₂); 2.41 (t, 2H, CH₂); 4.53 (t, 1H, CH); 6.95 (s, 2H); 7.00-7.36 (m, 6H, CH arom); 7.63 (d, 2H, CH arom); 7.89 (s, 2H, NH).

¹³C-NMR (50 MHz, CDCl₃) δ (ppm): 23.58; 33.96; 34.18; 35.17; 111.29; 119.14;

20 119.61; 119.76; 121.71; 121.85; 122.04; 127.07; 136.64; 179.85.

ES-MS m/z: 331.4 [M-H]⁻.

Step 3. A solution of intermediate D [5,5-Bis-(1H-indol-3-yl)-pentanoic acid, 202 mg, 0.608 mmol], PyBOP (406.2 mg, 0.73 mmol), NMM (0.33 mL, 3.04 mmol)

and E (O-benzylhydroxylamine hydrochloride, 116.5 mg, 0.73 mmol) in DCM (4 mL) reacted in the same conditions of example 1 to give 0.16 g of F [5,5-Bis-(1H-indol-3-yl)-pentanoic acid benzyloxy-amide, $\gamma = 60\%$].

$^1\text{H-NMR}$ (200 MHz, CDCl_3) δ (ppm): 1.78 (m, 2H, CH_2); 2.07 (br, 2H, CH_2); 2.20

5 (m, 2H, CH_2); 4.48 (t, 1H, CH); 4.86 (s, 2H, CH_2O); 6.97-7.22 (m, 6H, CH arom);

7.27-7.43 (br, 2H, CH arom); 7.56 (d, 2H, CH arom); 8.08 (s, 2H, NH).

ES-MS m/z : 436.4 [$\text{M}-\text{H}$]⁻ and 460.3 [$\text{M}+\text{Na}$]⁺.

Step 4. The reaction of intermediate F [5,5-Bis-(1H-indol-3-yl)-pentanoic acid benzyloxy-amide, 132 mg, 0.30 mmol] was carried out as in example 1 to give 10 0.10g of G [5,5-Bis-(1H-indol-3-yl)-pentanoic acid hydroxyamide, ST 2743, $\gamma = 97\%$].

$^1\text{H-NMR}$ (300 MHz, CD_3OH) δ (ppm): 1.69 (br quint, 2H, CH_2); 2.22 (br m, 4H, CH_2); 4.44 (t, 1H, CH); 6.86 (m, 2H, CH arom); 7.00 (t, 2H, CH arom); 7.20 (m, 4H, CH arom); 7.27 (d, 2H, CH arom); 7.47 (d, 2H, CH arom).

15 $^{13}\text{C-NMR}$ (75 MHz, CD_3OH) δ (ppm): 25.80; 35.22; 36.45; 36.62; 112.11; 119.18; 120.24; 120.42; 122.08; 122.84; 128.41; 138.40; 179.45.

ES-MS m/z : 346.4 [$\text{M}-\text{H}$]⁻.

EXAMPLE 3:

Preparation of 6,6-Bis-(1H-indol-3-yl)-hexanoic acid hydroxyamide (ST2754)

20 ST 2754 was synthesised as described in example 1.

Step 1. A solution of A (1H-Indole, 740 mg, 6.32 mmol), B [6-Oxo-hexanoic acid ethyl ester; 500 mg, 3.16 mmol] and Dysprosium triflate (2.89 g, 4.74 mmol) in $\text{MeOH}/\text{H}_2\text{O}$ (13 mL/8 mL) reacted as in example 1 (at 65°C) to give 0.60 g of C

[6,6-Bis-(1H-indol-3-yl)-hexanoic acid ethyl ester, y = 50 %].

¹H-NMR (200 MHz, CDCl₃) δ (ppm): 1.25 (t, 3H, CH₃); 1.48 (m, 2H, CH₂); 1.74 (q, 2H, CH₂); 2.28 (m, 4H, CH₂); 4.13 (q, 2H, CH₂); 4.50 (t, 1H, CH); 6.94 (s, 2H); 7.02-7.40 (m, 6H, CH arom); 7.63 (d, 2H, CH arom); 7.91 (s, 2H, NH).

5 ¹³C-NMR (50 MHz, CDCl₃) δ (ppm): 14.33; 25.26; 27.94; 34.02; 34.49; 35.58; 60.34; 111.24; 119.11; 119.71; 120.27; 121.60; 121.83; 127.20; 136.70; 174.12. ES-MS m/z: 375.6 [M+H]⁺.

Step 2. Intermediate C [6,6-Bis-(1H-indol-3-yl)-hexanoic acid ethyl ester], 300 mg, 0.80 mmol] and solid NaOH (400 mg, 9.6 mmol) in MeOH/DCM (4 mL/0.5 mL) reacted as in example 1 to give 0.27 g of D [6,6-Bis-(1H-indol-3-yl)-hexanoic acid, y= 97 %].

¹H-NMR (200 MHz, CDCl₃) δ (ppm): 1.48 (m, 2H, CH₂); 1.71 (m, 2H, CH₂); 2.29 (m, 4H, CH₂); 4.50 (t, 1H, CH); 6.90-7.40 (m, 8H, CH arom); 7.63 (d, 2H, CH arom); 7.90 (s, 2H, NH).

15 ¹³C-NMR (50 MHz, CDCl₃) δ (ppm): 15.35; 24.91; 27.82; 34.01; 35.44; 111.20; 119.15; 119.70; 120.24; 121.54; 121.87; 127.17; 136.69; 179.48.

ES-MS m/z: 345.5 [M-H]⁻.

Step 3. A solution of intermediate D [6,6-Bis-(1H-indol-3-yl)-hexanoic acid, 232 mg, 0.69 mmol], PyBOP (422.2 mg, 0.76 mmol), NMM (0.38 mL, 3.45 mmol) 20 and E (O-benzylhydroxylamine hydrochloride, 121.1 mg, 0.76 mmol) in DCM (3 mL) reacted in the same conditions of example 1 to give 0.19 g of F [6,6-Bis-(1H-indol-3-yl)-hexanoic acid benzyloxy-amide, y = 62 %].

¹H-NMR (200 MHz, CDCl₃) δ (ppm): 1.32 (br, 2H, CH₂); 1.63 (m, 2H, CH₂); 1.86

(br, 2H, CH_2); 2.17 (m, 2H, CH_2); 4.43 (t, 1H, CH); 4.78 (s, 2H, CH_2O); 6.80-7.20 (m, 6H, CH arom); 7.33 (s, 5H, CH arom); 7.58 (d, 2H, CH arom); 7.95 (s, 2H, NH).

^{13}C -NMR (50 MHz, $CDCl_3$) δ (ppm): 25.46; 27.71; 32.99; 33.95; 35.37; 60.54;

5 111.30; 119.07; 119.64; 119.87; 120.09; 120.80; 121.68; 121.81; 127.12; 128.72; 129.34; 136.66; 161.09.

ES-MS m/z: 452.7 $[M+H]^+$.

Step 4. The reaction of intermediate F [6,6-Bis-(1H-indol-3-yl)-hexanoic acid benzyloxy-amide, 162.7 mg, 0.36 mmol] was carried out as in example 1 to give 10 67 mg of G [6,6-Bis-(1H-indol-3-yl)-hexanoic acid hydroxyamide, ST 2754, y = 54%].

1H -NMR (200 MHz, CD_3OH) δ (ppm): 1.44 (m, 2H, CH_2); 1.69 (m, 2H, CH_2); 2.05 (t, 2H, CH); 2.25 (q, 2H, CH_2); 4.44 (t, 1H, CH); 6.89 (t, 2H, CH arom); 7.05 (m, 4H, CH arom); 7.31 (d, 2H, CH arom); 7.50 (d, 2H, CH arom).

15 ^{13}C -NMR (50 MHz, CD_3OH) δ (ppm): 25.03; 27.28; 32.05; 33.35; 34.75; 110.05; 117.65; 118.25; 119.77; 120.08; 120.69; 126.39; 136.20; 179.45.

ES-MS m/z: 360.3 $[M-H]^-$ and 384.3 $[M+Na]^+$.

EXAMPLE 4:

Preparation of N-Hydroxy-4,4-bis-(1H-indol-3-yl)-butyramide (ST2408)

20 **Step 1.** A solution of A (1H-Indole, 3.1 g, 26.5 mmol), B [4-Oxo-butyric acid; 4.8 mL, 7.69 mmol] and Dysprosium triflate (1.55 g, 2.5 mmol) in EtOH 95%/H₂O (32 mL/16 mL) reacted as in example 1 to give 0.36 g of C [5,5-Bis-(1H-indol-3-yl)-pentanoic acid methyl ester, y = 15 %].

¹H-NMR (200 MHz, DMSO-d₆) δ (ppm): 2.28 (m, 2H, CH₂); 2.48 (q, 2H, CH₂); 4.48 (t, 1H, CH); 6.80-7.16 (m, 6H, CH arom); 7.36 (d, 2H, CH arom); 7.56 (d, 2H, CH arom); 9.20 (s, 2H).

ES-MS m/z: 317.4 [M-H]⁻.

5 **Step 3.** A solution of intermediate D [4,4-Bis-(1H-indol-3-yl)-butyric acid, ST 1961, 220 mg, 0.69 mmol], PyBOP (392 mg, 0.70 mmol), TEA (0.40 mL, 2.77 mmol) and E (O-benzylhydroxylamine hydrochloride, 113 mg, 0.70 mmol) in DCM (40 mL) reacted in the same conditions of example 1 to give 0.20 g of F [N-Benzylxy-4,4-bis-(1H-indol-3-yl)-butyramide, y = 68 %].

10 ¹H-NMR (200 MHz, CD₃OD) δ (ppm): 2.18 (t, 2H, CH₂); 2.57 (q, 2H, CH₂); 4.43 (t, 1H, CH); 4.80 (s, 2H, CH₂O); 6.92 (t, 2H, CH arom); 6.98-7.17 (m, 4H, CH arom); 7.27 (m, 7H, CH arom); 7.56 (d, 2H, CH arom).

¹³C-NMR (50 MHz, CD₃OH) δ (ppm): 30.65; 30.74; 33.11; 77.00; 110.25; 117.38; 117.78; 118.34; 120.20; 121.09; 126.39; 127.50; 127.62; 128.34; 15 135.04; 136.48; 171.18.

ES-MS m/z: 422.3 [M-H]⁻ and 424.5 [M+H]⁺.

Step 4. The reaction of intermediate F, [N-Benzylxy-4,4-bis-(1H-indol-3-yl)-butyramide, 200 mg, 0.47 mmol] was carried out as in example 1 to give 0.11g of G [N-Hydroxy-4,4-bis-(1H-indol-3-yl)-butyramide, ST 2408, y= 71%].

20 ¹H-NMR (300 MHz, CD₃OH) δ (ppm): 2.20 (t, 2H, CH₂); 2.57 (q, 2H, CH₂); 4.46 (t, 1H, CH); 6.95 (t, 2H, CH arom); 6.98-7.19 (m, 4H, CH arom); 7.27 (d, 2H, CH arom); 7.47 (d, 2H, CH arom).

¹³C-NMR (50 MHz, CD₃OH) δ (ppm): 30.80; 30.93; 33.19; 110.23; 117.36;

117.87; 118.32; 120.18; 121.05; 126.41; 136.48; 171.31.

ES-MS m/z: 332.4 [M-H]⁻.

EXAMPLE 5:

Preparation of 8,8-Bis-(1H-indol-3-yl)-octanoic acid hydroxyamide (ST2889)

5 **Step 1.** A solution of B (8-Oxo-octanoic acid ethyl ester, 920.0 mg, 4.94 mmol), A (1H-Indole, 1.16 g mg, 9.88 mmol) and Dysprosium triflate (4.5 g, 7.41 mmol) in MeOH/H₂O (21 mL/7 mL) reacted in the same conditions of example 1 (16 h at 50 °C) to give 0.4 g of C [8,8-Bis-(1H-indol-3-yl)-octanoic acid ethyl ester, y = 23 %].

10 ¹H-NMR (200 MHz, CDCl₃) δ (ppm): 1.31 (t, 3H, CH₃); 1.42 (br, 4H, CH₂); 1.64 (m, 2H, CH₂); 2.30 (m, 4H, CH₂); 4.18 (q, 2H, OCH₂); 4.51 (t, 1H, CH); 6.95 (s, 2H); 7.10 (t, 2H, CH arom); 7.21 (t, 2H, CH arom); 7.33 (d, 2H, CH arom); 7.66 (d, 2H, CH arom); 7.90 (s, 2H, NH).

15 ¹³C-NMR (50 MHz, CDCl₃) δ (ppm): 14.39; 25.13; 28.26; 29.24; 29.53; 34.12; 34.52; 35.94; 60.35; 111.25; 119.08; 119.76; 120.51; 121.60; 121.80; 127.29; 136.73; 174.18.

ES-MS m/z: 401.3 [M-H]⁻ and 425.5 [M+Na]⁺.

20 **Step 2.** The reaction of intermediate C [8,8-Bis-(1H-indol-3-yl)-octanoic acid ethyl ester, 340 mg, 0.846 mmol] and solid LiOH (177.5 mg, 4.23 mmol) in MeOH/H₂O (2.9 mL/0.3 mL) was carried out as in example 1 to give 0.3 g of D [8,8-Bis-(1H-indol-3-yl)-octanoic acid, y = 95 %].

¹H-NMR (200 MHz, CDCl₃) δ (ppm): 1.39 (br, 6H, CH₂); 1.62 (quint, 2H, CH₂); 2.24 (m, 2H, CH₂); 2.33 (t, 2H, CH₂); 4.49 (t, 1H, CH); 6.98 (s, 2H, CH arom);

7.07 (t, 2H, *CH*arom); 7.18 (t, 2H, *CH*arom); 7.34 (d, 2H, *CH*arom); 7.63 (d, 2H, *CH*arom); 7.86 (s, 2H, *NH*).

¹³C-NMR (50 MHz, CDCl₃) δ (ppm): 24.75; 28.16; 29.08; 29.42; 34.02; 34.11; 35.82; 111.17; 119.11; 119.75; 120.54; 121.50; 121.83; 127.26; 136.70; 179.73.

5 ES-MS m/z: 373.0 [M-H]⁻ and 397.4 [M+Na]⁺.

Step 3. In a flask reagent E (hydroxylamine hydrochloride, 56.0 mg, 0.80 mmol) and DBU (122.6 mg, 0.80 mmol) were dissolved in DMF (0.8 mL) and the resulting solution was added to a solution of intermediate D [8,8-Bis-(1H-indol-3-yl)-octanoic acid, 200 mg, 0.537 mmol], HATU (224.6 mg, 0.591 mmol) and 10 DIEA (187.1 μL, 1.07 mmol). The reaction mixture was stirred at room temperature for 30 min and then diluted with ethyl acetate. The organic layer was washed with H₂O (x 3), dried over sodium sulfate, filtered and evaporated under reduced pressure. The residue was purified by a column chromatography on silica gel using a gradient elution of DCM/MeOH to give 0.12 g of F [8,8-Bis-(1H-indol-3-yl)-octanoic acid hydroxyamide, ST 2889, y = 57%].

¹H-NMR (300 MHz, DMSO-d₆) δ (ppm): 1.14-1.28 (br m, 6H, CH₂); 1.41 (quint, 2H, CH₂); 1.87 (t, 2H, CH₂); 2.13 (br, 2H, CH₂); 4.32 (t, 1H, CH); 6.83 (t, 2H, CH arom); 6.96 (t, 2H, CH arom); 7.18 (s, 2H, CH arom); 7.26 (d, 2H, CH arom); 7.46 (d, 2H, CH arom); 8.59 (s, 1H); 10.26 (s, 1H); 10.68 (s, 2H, NH).

20 ¹³C-NMR (75 MHz, DMSO-d₆) δ (ppm): 25.83; 28.48; 29.35; 29.48; 32.93; 34.09; 35.73; 110.95; 118.51; 119.57; 119.65; 121.21; 122.52; 127.37; 137.14; 169.80.

ES-MS m/z: 388.2 [M-H]⁻ and 412.7 [M+Na]⁺.

EXAMPLE 6:**Preparation of 7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid hydroxyamide**

(ST3043)

Step 1. A solution of B (7-Oxo-heptanoic acid ethyl ester, 58 mg, 0.34 mmol), A5 (7-Methoxy-1H-indole, 100 mg, 0.68 mmol) and trifluoroacetic acid (0.03 mmol mmol) in CH₃CN (1 mL) was stirred for 72 h at room temperature, then the solvent was evaporated under reduced pressure. Compound C [7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid ethyl ester] was not isolated and was characterised by ES-MS (m/z: 449.6 [M+H]⁺).10 **Step 2.** The crude of intermediate C [7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid ethyl ester], obtained from step 1 reacted with solid LiOH (43 mg, 1.02 mmol) in MeOH/THF/H₂O 3:3:1 (2 mL) as described in example 1 to give product D [7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid], that was not isolated and was characterised by ES-MS (m/z: 419.4 [M-H]⁻).15 **Step 3.** The crude of intermediate D [7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid] of step 2 reacted with PyBOP (265.2 mg, 0.51 mmol), DIET (0.30 mL, 1.7 mmol) and E (O-benzylhydroxylamine hydrochloride, 271.3 mg, 1.7 mmol) in DCM (2 mL) as in example 1 to give F [7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid benzyloxy-amide]; ES-MS (m/z: 526.8 [M+H]⁺).20 **Step 4.** The intermediate F [7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid benzyloxy-amide] obtained from step 3 was dissolved in MeOH and Pd/BaSO₄ 10% (cat) was added. The solution was mechanically stirred under hydrogen atmosphere (55 psi) overnight. The reaction mixture was filtered and the filtrate

was evaporated under reduced pressure. The solid residue was purified by preparative RP-HPLC (column Lichrosorb RP-18, 7 μ m, eluents: CH₃CN/H₂O + 0.1 %TFA 60: 40) gave 60.0 mg of G [7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid hydroxyamide, ST 3043, overall yield = 40%].

5 ¹H-NMR (200 MHz, DMSO-d₆) δ (ppm): 1.27 (br, 4H, CH₂); 1.42 (m, 2H, CH₂); 1.87 (t, 2H, CH₂); 2.09 (br, 2H, CH₂); 4.27 (t, 1H, CH); 6.56 (d, 2H, CH arom); 6.78 (t, 2H, CH arom); 7.05 (m, 4H, CH arom); 8.63 (s, 1H); 10.29 (s, 1H); 10.79 (s, 2H, NH).
10 ¹³C-NMR (50 MHz, DMSO-d₆) δ (ppm): 25.97; 28.37; 29.44; 33.11; 34.32; 35.79; 55.75; 101.96; 112.73; 119.14; 120.20; 122.24; 127.20; 128.96; 146.82; 169.89.
ES-MS m/z: 434.4 [M-H]⁻ and 458.3 [M+Na]⁺.

EXAMPLE 7:

15 Preparation of 7,7-Bis-(5-methyl-1H-indol-3-yl)-heptanoic acid hydroxyamide (ST3044)

ST 3044 was synthesised as in example 6.

20 **Step 1.** Reagent A (5-Methyl-1H-indole, 100 mg, 0.762 mmol) and B [7-Oxo-heptanoic acid ethyl ester, 65.6 mg, 0.381 mmol] reacted to give C [7,7-Bis-(5-methyl-1H-indol-3-yl)-heptanoic acid ethyl ester, that was not isolated and was characterised by ES-MS (m/z: 417.6 [M+H]⁺).

Step 2. The intermediate D [7,7-Bis-(5-methyl-1H-indol-3-yl)-heptanoic acid], not isolated, was characterised by ES-MS (m/z: 387.4 [M-H]⁻).

Step 3. The intermediate F [7,7-Bis-(5-methyl-1H-indol-3-yl)-heptanoic acid benzyloxy-amide] was purified by a flash chromatography on silica gel with a gradient elution of hexane/ethyl acetate and characterised by ES-MS (m/z: 494.7 [M+H]⁺).

5 **Step 4.** Product G [7,7-Bis-(5-methyl-1H-indol-3-yl)-heptanoic acid hydroxyamide, ST3044] was purified by filtration of reaction mixture and solvent evaporation under reduced pressure (55 mg, overall yield = 36%).

10 ¹H-NMR (300 MHz, DMSO-d₆) δ (ppm): 1.27 (br, 4H, CH₂); 1.42 (m, 2H, CH₂); 1.88 (t, 2H, CH₂); 2.09 (br, 2H, CH₂); 2.28 (s, 3H, CH₃); 4.25 (t, 1H, CH); 6.79 (d, 2H, CH arom); 7.09 (s, 2H, CH arom); 7.15 (d, 2H, CH arom); 7.25 (s, 2H, CH arom); 8.61 (s, 1H); 10.27 (s, 1H); 10.53 (s, 2H, NH).
15 ¹³C-NMR (75 MHz, DMSO-d₆) δ (ppm): 22.03; 25.93; 28.37; 29.39; 33.04; 34.39; 35.75; 111.69; 119.00; 119.23; 122.64; 122.85; 126.77; 127.58; 135.56; 169.82.

15 ES-MS m/z: 402.4 [M-H]⁻ and 426.4 [M+Na]⁺.

EXAMPLE 8:

Preparation of 7,7-Bis-(4-fluoro-1H-indol-3-yl)-heptanoic acid hydroxyamide (ST3052).

ST 3052 was synthesised as in example 6.

20 **Step 1.** Reagent A (4-Fluoro-1H-indole, 100 mg, 0.74 mmol) and B [7-Oxo-heptanoic acid ethyl ester, 63.7 mg, 0.37 mmol] reacted to give C [7,7-Bis-(4-fluoro-1H-indol-3-yl)-heptanoic acid ethyl ester], that was not isolated and was characterised by ES-MS (m/z: 425.6 [M+H]⁺).

Step 2. The intermediate D [7,7-Bis-(4-fluoro-1H-indol-3-yl)-heptanoic acid] was not isolated and was characterised by ES-MS (m/z: 395.4 [M-H]⁻).

Step 3. The intermediate F [7,7-Bis-(4-fluoro-1H-indol-3-yl)-heptanoic acid benzyloxy-amide] was roughly purified by a flash chromatography on silica gel

5 with a gradient elution of hexane/ethyl acetate and characterised by ES-MS (m/z: 502.7 [M+H]⁺).

Step 4. Product G [7,7-Bis-(4-fluoro-1H-indol-3-yl)-heptanoic acid hydroxyamide, ST 3052] was purified by filtration of reaction mixture, solvent evaporation and preparative RP-HPLC (column Lichrosorb RP-18, 7μm; 10 eluents: H₂O/CH₃CN 50:50; flow= 10 mL/min, 55 mg, overall yield = 35%).

¹H-NMR (300 MHz, DMSO-d₆) δ (ppm): 1.29 (br, 4H, CH₂); 1.42 (quint, 2H, CH₂); 1.87 (t, 2H, CH₂); 2.03 (q, 2H, CH₂); 4.67 (t, 1H, CH); 6.58-6.66 (m, 2H, CH arom); 6.98 (m, 4H, CH arom); 7.13 (d, 2H, CH arom); 8.60 (s, 1H); 10.27 (s, 1H); 11.01 (s, 2H, NH).

15 ¹³C-NMR (75 MHz, DMSO-d₆) δ (ppm): 25.87; 28.47; 29.39; 33.01; 35.15; 37.89; 103.98 (d, 1C arom, J= 19.6 Hz); 108.59; 115.75 (d, 1C arom J= 19.0 Hz); 118.89 (d, 1C arom, J= 3.97 Hz); 121.80 (d, 1C arom, J= 7.95 Hz); 123.12; 139.97 (d, 1C arom, J= 12.0 Hz); 157.11 (d, 1C, C-F, J= 243.7 Hz); 169.78.

ES-MS m/z: 410.5 [M-H]⁻; 434.4 [M+Na]⁺ and 450.4 [M+K]⁺.

20 **EXAMPLE 9:**

Preparation of 3-{4-[Bis-(1H-indol-3-yl)-methyl]-phenyl}-N-hydroxy-acrylamide (ST2887).

Step 1. The reaction of A (1H-Indole, 234 mg, 2 mmol) and B [3-(4-Formyl-

phenyl)-acrylic acid, 176 mg, 1 mmol] was carried out as in example 1 to obtain the acid C [3-{4-[Bis-(1H-indol-3-yl)-methyl]-phenyl}-acrylic acid] utilised as crude for step 2.

¹H-NMR (200 MHz, DMSO-d₆) δ (ppm): 5.86 (s, 1H, CH); 6.44 (d, 1H, CH, J = 16.1 Hz); 6.86 (m, 4H, CH arom); 7.04 (t, 2H, CH arom); 7.25-7.59 (m, 9H, 8CH arom + 1CH double bond); 10.85 (s, 2H, NH).

¹³C-NMR (50 MHz, DMSO-d₆) δ (ppm): 57.58; 112.24; 118.24; 118.98; 119.79; 121.68; 124.36; 127.28; 128.87; 129.56; 132.62; 137.32; 144.68; 148.30; 168.38.

ES-MS m/z: 391.15 [M-H]⁻.

Step 2. The reaction was carried out as in Step 3 of example 3. Purification by preparative RP-HPLC (column Lichrosorb RP-18, 7μm; eluents: H₂O/CH₃CN 50:50; flow= 10 mL/min) gave the product F [3-{4-[Bis-(1H-indol-3-yl)-methyl]-phenyl}-N-hydroxy-acrylamide, ST 2887].

¹H-NMR (300 MHz, DMSO-d₆) δ (ppm): 5.85 (s, 1H, CH); 6.39 (d, 1H, CH, J = 15.95 Hz); 6.86 (m, 4H, CH arom); 7.03 (t, 2H, CH arom); 7.25-7.48 (m, 9H, 8CH arom + 1CH double bond); 8.99 (s, 1H); 10.71 (s, 1H); 10.85 (s, 2H, NH).

¹³C-NMR (75 MHz, DMSO-d₆) δ (ppm): 57.58; 114.16; 120.28; 120.90; 121.72; 123.60; 126.11; 126.27; 129.23; 130.03; 131.52; 135.13; 139.26; 140.96;

149.35; 165.58.

ES-MS m/z: 405.71 [M-H]⁻.

EXAMPLE 10:

Preparation of 7,7-Bis-(1H-indol-3-yl)-heptanoic acid (2-amino-phenyl)-amide

(ST3071)

Step 3. The intermediate D [7,7-Bis-(1H-indol-3-yl)-heptanoic acid, 60 mg, 0.167 mmol] obtained from step 2 in example 1 was dissolved in DCM (1 mL), then PyBOP (95.3 mg, 0.184 mmol) and DIEA (87 μ L, 0.50 mmol) were added.

5 After 5 min of stirring the resulting solution was added to a solution of E (2-aminophenylamine; 86 mg, 0.835 mmol) in DCM (1 mL) and the reaction mixture was stirred at room temperature for 2h. Then the solvent was removed under reduced pressure and the residue dissolved in ethyl acetate. The solution was washed with HCl 0.5N, a saturated solution of NaHCO₃ and water (x2). The 10 organic layer was dried over sodium sulfate, filtered and evaporated under reduced pressure. The residue was roughly purified by a flash chromatography on silica gel by using hexane/ethyl acetate as eluent system, to give F [7,7-Bis-(1H-indol-3-yl)-heptanoic acid (2-amino-phenyl)-amide, ST 3071, 50 mg, y = 67%]

15 ¹H-NMR (200 MHz, DMSO-d₆) δ (ppm): 1.36 (br, 4H, CH₂); 1.54 (br m, 2H, CH₂); 2.23 (m, 4H, CH₂); 4.36 (t, 1H, CH); 4.79 (s, 2H, CH₂-O); 6.52 (t, 2H, CH arom); 6.70 (d, 2H, CH arom); 6.80-7.05 (m, 4H, CH arom); 7.06-7.32 (m, 4H, CH arom); 7.50 (d, 2H, CH arom); 9.06 (s, 1H, NH); 10.72 (s, 2H, NH).

13C-NMR (50 MHz, DMSO-d₆) δ (ppm): 26.16; 28.45; 29.53; 34.11; 35.70; 20 36.60; 112.02; 116.60; 116.89; 118.58; 119.57; 119.73; 121.29; 122.61; 124.30; 126.01; 126.40; 127.40; 137.19; 142.62; 171.89.

ES-MS m/z: 451.2 [M+H]⁺ and 449.4 [M-H]⁻.

EXAMPLE 11:

Preparation of N-Hydroxy-3-(1H-indol-3-yl)-acrylamide (ST2913)

Reagent E (hydroxylamine hydrochloride, 112.0 mg, 1.61 mmol) and DBU (254.0 mg, 1.61 mmol) were dissolved in DMF (0.5 mL) and the resulting solution was added to a suspension of D [3-(1H-Indol-3-yl)-acrylic acid, 151 mg, 5 0.806 mmol], HATU (337.1 mg, 0.887 mmol) and DIEA (281.0 μ L, 1.61 mmol) in DMF/DCM (1.5 mL/2 mL). The reaction was treated as described in step 3 of example 3 and the final purification by preparative RP-HPLC (column Lichrosorb RP-18, 7 μ m; eluents: H_2O/CH_3CN 70:30; flow= 10 mL/min) gave the product F [N-Hydroxy-3-(1H-indol-3-yl)-acrylamide, ST 2913, 70 mg, y = 10 43%].

1H -NMR (300 MHz, DMSO-d₆) δ (ppm): 6.40 (d, 1H, CH=C, J = 15.8 Hz); 7.08-7.22 (m, 2H, CH arom); 7.43 (m, 1H, CH arom); 7.54 (d, 1H, CH=C, J = 15.9 Hz); 7.78 (m, 2H, CH arom); 8.82 (s, 1H); 10.42 (s, 1H); 11.53 (s, 1H, NH).
 ^{13}C -NMR (75 MHz, DMSO-d₆) δ (ppm): 112.75; 112.97; 113.56; 120.39; 121.02; 15 122.87; 125.57; 130.83; 133.40; 138.02; 165.28.
ES-MS m/z: 201.1 [M-H]⁻ and 224.86 [M+Na]⁺.

EXAMPLE 12:Preparation of N-hydroxy-6-(1H-indol-3-yl)hexanamide (ST2995)

Step 3. In a flask reagent E (hydroxylamine hydrochloride, 120.8 mg, 1.74 mmol) and DIEA (302.6 μ L, 1.74 mmol) were dissolved in DMF (0.5 mL) and the resulting solution was added to a solution of D [6-(1H-indol-3-yl)hexanoic acid, 200 mg, 0.869 mmol], HATU (363 mg, 0.956 mmol) and DIEA (302.6 μ L, 1.74 mmol) in DMF (2 mL). The reaction mixture was stirred at room temperature for

2h and then diluted with ethyl acetate. The organic layer was washed with HCl 1N, NaHCO₃ 3%, brine (x 2), dried over sodium sulfate, filtered and evaporated under reduced pressure. Precipitation from DCM, filtration and final washings of the precipitate with DCM gave 0.17 g of F [8,8-Bis-(1H-indol-3-yl)-octanoic acid hydroxyamide, ST 2995, y = 80%].

¹H-NMR (200 MHz, DMSO-d₆) δ (ppm): 1.33 (m, 2H, CH₂); 1.65 (m, 4H, CH₂); 1.94 (t, 2H, CH₂); 2.66 (t, 2H, CH₂); 6.95 (t, 1H, CH arom); 7.05 (m, 2H, CH arom); 7.32 (d, 1H, CH arom); 7.48 (d, 1H, CH arom); 8.65 (s, 1H); 10.32 (s, 1H); 10.73 (s, 1H).

¹³C-NMR (50 MHz, DMSO-d₆) δ (ppm): 25.32; 25.79; 29.34; 30.39; 33.06; 112.03; 115.34; 118.78; 119.03; 121.49; 122.80; 127.96; 137.05; 169.86. ES-MS m/z: 245.2 [M-H]⁻ and 247.3 [M+H]⁺ and 269.2 [M+Na]⁺.

EXAMPLE 13:

Preparation of 7,7-Bis-(7-Ethyl-1H-indol-3-yl)-heptanoic acid hydroxyamide

(ST3126)

Step 1. A solution of B (7-Oxo-heptanoic acid ethyl ester, 90 mg, 0.516 mmol), A (7-ethyl-1H-Indole, 100.2 mg, 0.688 mmol) and Dysprosium triflate (420 mg, 0.688 mmol) in AcCN (1.2 mL) reacted as in example 1 (at room temperature for 2 days) to give C [7,7-Bis-(7-ethyl-1H-indol-3-yl)-heptanoic acid ethyl ester,

that was not isolated and was characterised by ES-MS (m/z: 445.8 [M+H]⁺).

Step 2. The crude of intermediate C [7,7-Bis-(7-ethyl-1H-indol-3-yl)-heptanoic acid ethyl ester], obtained from step 1 reacted as described in example 6 to give

product D [7,7-Bis-(7-ethyl-1H-indol-3-yl)-heptanoic acid], that was not isolated and was characterised by ES-MS (m/z: 415.4 [M-H]⁻).

Step 3. The crude of intermediate D [7,7-Bis-(7-ethyl-1H-indol-3-yl)-heptanoic acid] obtained from step 2 reacted as described in example 6 to give F [7,7-Bis-

5 (7-ethyl-1H-indol-3-yl)-heptanoic acid benzyloxy-amide], that was not isolated and was characterised by ES-MS (m/z: 522.8 [M+H]⁺).

Step 4. The crude of intermediate F [7,7-Bis-(7-ethyl-1H-indol-3-yl)-heptanoic acid benzyloxy-amide] obtained from step 3 reacted as described in example 6.

Purification by preparative RP-HPLC (column Lichrosorb RP-18, 7 \square m, eluents:

10 CH₃CN/H₂O 60: 40) gave 100.0 mg of G [7,7-Bis-(7-ethyl-1H-indol-3-yl)-heptanoic acid hydroxyamide, ST3126, overall yield = 67%].

¹H-NMR (300 MHz, DMSO-d₆) δ (ppm): 1.21 (t, 6H, CH₃); 1.28 (br, 4H, CH₂); 1.42 (m, 2H, CH₂); 1.88 (t, 2H, CH₂); 2.14 (br, 2H, CH₂); 2.77 (q, 4H, CH₂); 4.30 (t, 1H, CH); 6.76-6.81 (m, 4H, CH arom); 7.13 (s, 2H, CH arom); 7.34 (m, 2H, CH arom); 8.60 (s, 1H); 10.26 (s, 1H); 10.63 (s, 2H, NH).

¹³C-NMR (75 MHz, DMSO-d₆) δ (ppm): 14.99; 24.29; 25.91; 28.40; 29.41; 33.05; 34.24; 35.68; 111.37; 118.83; 119.91; 120.01; 122.01; 122.17; 127.23; 127.29; 135.66; 135.81; 169.80.

ES-MS m/z: 430.4 [M-H]⁻ and 454.0 [M+Na]⁺.

20 **EXAMPLE 14:**

Preparation 7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid hydroxyamide (ST3307)

Step 1. A solution of B (7-oxo-heptanoic acid ethyl ester, 2.7 g, 15.68 mmol) and A (5-morpholin-4-ylmethyl-3H-indole, 2.8 g, 12.95 mmol) in glacial CH₃COOH (20 mL) reacted at reflux overnight to give C 7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid ethyl ester, that was not isolated and was 5 characterised by ES-MS (m/z: 587.4 [M+H]⁺ and 609.4 [M+23]⁺.

Step 2. The crude of intermediate C 7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid ethyl ester, obtained from step 1 reacted as described in example 6 to give product D 7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid, that was not isolated and was characterised by ES-MS (m/z: 10 557.6 [M-H]⁻).

Step 3. The crude of intermediate D 7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid obtained from step 2 reacted as described in example 6 to give F 7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid benzyloxy-amide, that was not isolated and was characterised by ES-MS (m/z: 15 664.9 [M+H]⁺).

Step 4. The crude of intermediate F 7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid benzyloxy-amide obtained from step 3 reacted as described in example 6. Purification by preparative RP-HPLC (column Lichrosorb RP-18, 7 μ m, eluents: CH₃OH + 0.1 % TFA/H₂O + 0.1 % TFA = 45:55) gave 100.0 mg of 20 G 7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid hydroxyamide, ST3307, overall yield = 40%].

¹H-NMR (300 MHz, DMSO-d₆) δ (ppm): 1.28 (bm, 4H, 2xCH₂); 1.42 (bm, 2H, CH₂); 1.88 (m, 2H, CH₂); 2.12 (bm, 2H, CH₂); 2.32 (bm, 8H, 4xCH₂); 3.34 (bm,

4H, 2xCH₂); 3.53 (bm, 8H, 4xCH₂) 4.30 (t, J = 7.1 Hz, 1H, CH); 6.94 (d, J = 8.1 Hz, 2H, CH arom); 7.14 (s, 2H, CH arom); 7.22 (d, J = 8.1 Hz, 2H, CH arom); 7.33 (s, 2H, CH arom); 8.63 (bs, 2H); 10.30 (s, 2H); 10.71 (s, 2H, NH).

¹³C-NMR (75.5 MHz, DMSO-d₆) δ (ppm): 26.00; 27.90; 28.25; 33.99; 34.62;

5 35.49; 53.63; 62.29; 67.17; 109.78; 118.51; 121.57; 122.73; 123.94; 127.23; 130.13; 136.36; 170.05.

ES-MS m/z: 574.4 [M+H]⁺.

EXAMPLE 15:

Preparation 7-(1H-Indol-3-yl)-7-(1H-indol-2-yl)-heptanoic acid hydroxyamide

10 (ST3292)

During synthesis of ST2741, example 1, 7-(1H-Indol-3-yl)-7-(1H-indol-2-yl)-heptanoic acid hydroxyamide was obtained as impurity. The product was purified and characterized.

¹H-NMR (300 MHz, DMSO-d₆) δ (ppm): 1.28 (bm, 4H, 2XCH₂); 1.44 (bm, 2H,

15 CH₂); 1.43 (bm, 2H, CH₂); 1.88 (t, 2H, CH₂); 2.12 (bm, 2H, CH₂); 4.27 (t, J = 7.6 Hz, 1H, CH); 6.23 (s, 1H, CH arom); 6.86 (t, J = 7.0 Hz, 2H, CH arom); 6.92 (t, J = 7.0 Hz, 1H, CH arom); 7.00 (t, J = 7.1 Hz, 1H, CH arom); 7.18-7.24 (m, 2H, CH arom); 7.30 (d, J = 8.0 Hz, 1H, CH arom); 7.36 (d, J = 7.7 Hz, 1H, CH arom); 7.45 (d, J = 7.8 Hz, 1H, CH arom); 8.63 (s, 1H); 10.28 (s, 1H); 10.75 (s,

20 1H); 10.86 (s, 1H).

¹³C-NMR (75.5 MHz, DMSO-d₆) δ (ppm): 25.81; 28.04; 29.26; 33.00; 34.96;

36.51; 98.56; 111.39; 112.05; 117.55; 118.82; 119.07; 119.39; 119.82; 120.53; 121.49; 122.88; 127.14; 128.75; 136.66; 137.02; 144.31; 169.78..

ES-MS m/z: 376.2 [M+H]⁺, 398.1 [M+Na]⁺ and 374.2 [M-H]⁻.

EXAMPLE 16:

Preparation of 7,7-Bis-(1H-indol-3-yl)-heptanoic acid (ST3127)

7,7-Bis-(1H-indol-3-yl)-heptanoic acid (ST3127) was prepared as described in

5 example 1, step 1 and step 2.

¹H-NMR (300 MHz, Acetone-d₆) δ (ppm): 1.45 (br m, 4H, CH₂); 1.59 (quint, 2H, CH₂); 2.26 (m, 4H, CH₂); 4.52 (t, 1H, CH); 6.87-6.95 (m, 2H, CH_{arom}); 6.99-7.07 (m, 2H, CH_{arom}); 7.21 (d, 2H, CH_{arom}); 7.34 (d, 2H, CH_{arom}); 7.59 (d, 2H, CH_{arom}); 9.89 (s, 2H, NH).

10 ¹³C-NMR (75 MHz, Acetone-d₆) δ (ppm): 25.06; 28.14; 29.29; 33.57; 34.19; 35.81; 111.39; 118.42; 119.50; 119.95; 121.16; 121.99; 127.56; 137.39; 174.02.

ES-MS m/z: 359.2 [M-H]⁻ and 383.2 [M+Na]⁺.

List of abbreviations:

AcCN	Acetonitrile
15 DBU	1,8-Diazabicyclo[5.4.0]undec-7-ene
DCC	Dicyclohexylcarbodiimide
DCM	Dichloromethane
DIEA	N,N,N-diisopropylethylamine
DMAP	4-Dimethylaminopyridine
20 DMF	Dimethylformamide
DMSO	Dimethylsulfoxide
EDC	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride

	EtOH	Ethanol
	TEA	Triethylamine
	HATU	Hydroxyazabenzotriazol-N,N,N,N-tetramethyluronium hexafluorophosphate
5	HOEt	Hydroxybenzotriazole
	MeOH	Methanol
	NMM	N- Methylmorpholine
	PyBOP	(Benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate
10	RP-HPLC	Reversed Phase-HPLC
	TFA	trifluoroacetic acid
	THF	Tetrahydrofuran

Biological results

Cytotoxicity studies

15 To test the effects of the compounds on cell growth, NB4 human promyelocytic leukaemia, NCI-H460 non-small cell carcinoma cells and HCT-116 human colon carcinoma cells were used. NB4 and NCI-H460 tumour cells were grown RPMI 1640 containing 10% fetal bovine serum (GIBCO), whereas HCT-116 tumour cells were grown in McCoy's 5A containing 10% fetal bovine serum (GIBCO).

20 Tumour cells were seeded in 96-well tissue culture plates (Corning) at approximately 10% confluence and were allowed to attach and recover for at least 24 h. Varying concentrations of the drugs were then added to each well to calculate their IC₅₀ value (the concentration which inhibits the 50% of cell

survival). The plates were incubated for 24 h at 37 °C. At the end of the treatment, for NB4 tumour cells in suspension, the procedure was performed as follows: medium culture was removed by centrifugation of the plates at 1600 x g for 10 min and the surnatant was removed. 250 µl PBS were added, then the 5 plates were centrifuged at 1600 x g for 10 min, the surnatant was removed. 200 µl/well of medium culture RPMI 1640 containing 10% FCS were added and the plates were incubated at 37 °C for other 48 h. The plates were centrifuged again at 1600 x g for 10 min, the medium culture was removed and 200 µl PBS and 50 µl of cold 80%TCA were added. The plates were incubated on ice for at 10 least 1 h. TCA was removed, the plates were washed 3 times for immersion in distilled-water and dried on paper and at 40°C for 5 min. Then 200 µl of 0.4% sulphorodamine B in 1% acetic acid were added. The plates were incubated at room temperature for other 30 min. Sulphorodamine B was removed, the plates were washed for immersion in 1% acetic acid for 3 times, then they were dried 15 on paper and at 40 °C for 5 min. Then 200 µl Tris 10 mM were added, the plates were kept under stirring for 20 min. The survival cell was determined as optical density by a Multiskan spectrofluorimeter at 540 nm. For the tumour cells in adhesion (NCI-H460 and HCT-116), the procedure was as above mentioned, except that at the end of the treatment, the plates were washed by remotion of 20 the surnatant and addition of PBS 3 times without centrifugation. Also the last day of the assay, the surnatant was removed without centrifugation.

The amount of cells killed was calculated as the percentage decrease in sulphorodamine B binding compared with control cultures. The IC₅₀ values (the

concentration which inhibits the 50% of cell survival) were calculated with the "ALLFIT" program. In the table 1 the cytotoxicity evaluated on NB4 tumor cells showed that ST3044 was the most potent compound with an IC50 value of 0.15 μ M followed by other molecules (ST3052, ST3043, ST3292, ST3307, ST2741, 5 ST2887, ST2889, ST2754) with IC50 values of 0.4-2.3 μ M. On NCI-H460 non-small cell lung cancer and HCT-116 colon cancer, ST3044, ST3043 and ST3052 were the most potent molecules with IC50 values which ranged from 0.27 to 0.7 μ M, followed by ST3292, ST3307, ST2741, ST2887, ST2889, ST2754 with IC50 values of 0.6-5.8 μ M. Other molecules (ST2408, ST2743, 10 ST2995, ST3127) showed a minor cytotoxicity on NB4 or NCI-H460 and HCT-116 tumor cells, since IC50 values ranged from 8.8 μ M to 88.8 μ M.

Moreover, the molecules of the present invention resulted to be more potent than the homologues bis-heterocyclic compounds in the cytotoxicity test on NB4 tumor cells (see Table 2).

**Table 1.: Cytotoxicity of different compounds on NB4, NCI-H460 and HCT-116
tumour cells**


Compound	NB4	NCI-H460	HCT-116
IC50±SD, μ M			
ST2408	14.6±0.6	25.9±0.3	
ST2741	0.6±0.01	1.2±0.1	0.6±0.02
ST2743		88.8±2.8	
ST2754	1.6±0.06	5.2±0.3	5.8±0.2
ST2887	0.6±0.01	2.6±0.3	3.9±0.5
ST2889	2.3±0.1	2.7±0.2	3.5±0.2
ST2995	8.8±0.5	22±1.7	
ST3043	0.8±0.01	0.55±0.03	0.41±0.03
ST3044	0.15±0.02	0.51±0.1	0.27±0.01
ST3052	0.6±0.09	0.7±0.09	0.4±0.1
ST3127	>20	>20	>20
ST3292	0.67±0.01	0.82±0.07	
ST3307	0.42±0.02	1.1±0.1	

**Table 2.: Cytotoxicity of the homologues bis-heterocyclic compounds on NB4
tumour cells**

Compound	NB4
	IC50±SD, μ M
ST1346	44.3±0.6
ST1383	48±11
ST1393	16.2±0.3
ST1422	47.3±5.6
ST1730	5.7±0.2
ST1880	22±6
ST1896	33.6±6.5


CLAIMS

1. A compound of Formula (I),




(I)

5 where



- A₁ is either H or A, wherein R, R' and R₁ are the same for both indole rings;
- 10 ➤ X is a saturated or unsaturated (alkenylene or alkynylene), linear or branched (C₂-C₁₀) alkylene, optionally substituted with OH, or a (C₆-C₁₂) aryl, (C₃-C₁₀) heteroaryl containing at least one heteroatom selected from N, O or S, wherein in the cyclic groups at least one of the -CH- is optionally substituted with C-halogen or C-(C₁-C₃) alkyl;
- 15 ➤ Q =SH, COCONR₂R₃, COCF₃, or



where,

- G is either H or a glycosyl;
- R₂, R₃ are the same or different and are either H or a (C₁-C₄) alkyl;
- R₁ is selected from the group consisting of H, (C₁-C₄) alkyl, (C₆-C₁₂) aryl, (C₆-C₁₂) aryl-(C₁-C₄) alkylene, (C₁-C₄) alkanoyl and (C₁-C₄)-alkyl-(C₆-C₁₀)-arylene;
- R and R', the same or different, are selected from the group consisting of:
 - H
 - saturated or unsaturated, linear or branched (C₁-C₁₀) alkyl, optionally substituted with a (C₃-C₁₀) heteroaryl or (C₃-C₁₀) heterocycl-(C₁-C₄) alkylene, where the heterocycle contains at least one heteroatom selected from N, O or S, or with a group -NR₅R₆, where R₅, R₆ are the same or different and are H, linear or branched (C₁-C₄) alkyl, (C₁-C₄)-alkanoyl
 - OR₄ where R₄ = H, (C₁-C₄) alkyl, mesyl, tosyl, (C₁-C₄) alkanoyl, glycosyl
 - halogen, azide, nitro, nitrile and NR₅R₆.

20 2. The compound according to claim 1, wherein A is A₁.

3. The compound according to claim 1, wherein X is a saturated or unsaturated linear C₂-C₁₀ alkylene.

4. The compound according to claim 1, wherein Q is CONHOG.

5. The compound according to claims 1 or 4, wherein G, R₁, R₂ and R₃ are the same and are H.

6. The compound according to claim 1, wherein R and R' are the same and are selected from the group consisting of H, (C₁-C₃) alkylamine and (C₁-C₃) alkylmorpholine.

7. The compound of Formula (I) of claim 1, which is selected from the group 10 consisting of:

3-{4-[Bis-(1H-indol-3-yl)-methyl]-phenyl}-N-hydroxy-acrylamide,

5,5-Bis-(1H-indol-3-yl)-pentanoic acid hydroxyamide,

6,6-Bis-(1H-indol-3-yl)-hexanoic acid hydroxyamide,

7,7-Bis-(1H-indol-3-yl)-heptanoic acid hydroxyamide,

15 7,7-Bis-(4-fluoro-1H-indol-3-yl)-heptanoic acid hydroxyamide,

7,7-Bis-(5-methyl-1H-indol-3-yl)-heptanoic acid hydroxyamide,

7,7-Bis-(7-ethyl-1H-indol-3-yl)-heptanoic acid hydroxyamide,

7,7-Bis-(7-methoxy-1H-indol-3-yl)-heptanoic acid hydroxyamide,

7,7-Bis-(5-morpholin-4-ylmethyl-3H-indol-3-yl)-heptanoic acid hydroxyamide,

20 7-(1H-Indol-3-yl)-7-(1H-indol-2-yl)-heptanoic acid hydroxyamide,

7,7-Bis-(1H-indol-3-yl)-heptanoic acid,

8,8-Bis-(1H-indol-3-yl)-octanoic acid hydroxyamide,

N-Hydroxy-4,4-bis-(1H-indol-3-yl)-butyramide, and

N-Hydroxy-6-(1*H*-indol-3-yl)hexanamide,

8. A process for preparing the compounds according to any of the preceding claims, in which A is A1, comprising dimerizing the starting indole derivative in the presence of an aldehyde or acetal.

5 9. A compound of any of claims 1 to 7, as a medicament.

10. Use of a compound of any of claims 1 to 7, for the preparation of a medicine with antitumour activity.

11. Use of a compound of any of claims 1 to 7, for the preparation of a medicine for the treatment of a tumour pathology, in which the tumour has
10 shown drug resistance to previous antitumour drugs used for its treatment, in which said formula (I) compound exerts a chemosensitising effect on said drug-resistant tumour.

12. The use according to claims 10 or 11, in which the tumour is selected from the group consisting of sarcoma, carcinoma, carcinoid, bone tumour,
15 neuroendocrine tumour, lymphoid leukaemia, acute promyelocytic leukaemia, myeloid leukaemia, monocytic leukaemia, megakaryocytic leukaemia and Hodgkin's disease.

13. The use according to claim 10 or 11 in which the compound of Formula (I) is combined with one or more known antitumour agents.

20 14. The use according to claim 13, in which the known antitumour agent is all-trans retinoic acid.

15. A pharmaceutical composition containing as active ingredient a compound of formula (I) of any of claims 1 to 7, and at least one pharmaceutically acceptable excipient and/or diluent.

16. The composition according to claim 15, for the treatment of tumour

5 pathologies.

17. The composition according to claim 15, for the treatment of a tumour pathology, in which the tumour has shown drug resistance to previous antitumour drugs used for its treatment, in which said formula (I) compound exerts a chemosensitising action on said drug-resistant tumour.

10 18. The composition according to claims 16 or 17, in which the tumour pathology is selected from the group consisting of sarcoma, carcinoma, carcinoid, bone tumour, neuroendocrine tumour, lymphoid leukaemia, acute promyelocytic leukaemia, myeloid leukaemia, monocytic leukaemia, megakaryoblastic leukaemia and Hodgkin's disease.

15 19. The composition according to claims 16 or 17, in which the compound of Formula (I) is administered in combination with one or more known antitumour agents.


20 20. The composition according to claims 16 or 17, in which the known antitumour agent is selected from the group consisting of alkylating agents, topoisomerase inhibitors, antitubulin agents, intercalating compounds, antimetabolites, natural products such as vinca alkaloids, epipodophyllotoxins, antibiotics, enzymes, taxans, and cytodifferentiating compounds.

21. The composition according to claim 20, in which the cytodifferentiating antitumour compound is all-trans retinoic acid.

22. A process for preparing the composition of any claims from 15 to 21, comprising mixing the compound(s) of any claims from 1 to 7 with the suitable

5 excipient(s) and/or diluent(s).

23. Method of treating a mammal suffering from a disease according any of claims 10 to 12, comprising administering a therapeutically effective amount of the compound of any of claims 1 to 7.



Formula I

where

