




(51) International Patent Classification ⁶ :		A1	(11) International Publication Number:	WO 99/08653				
A61K 7/42			(43) International Publication Date:	25 February 1999 (25.02.99)				
(21) International Application Number:	PCT/EP98/05042							
(22) International Filing Date:	8 August 1998 (08.08.98)							
(30) Priority Data:	97810585.6	20 August 1997 (20.08.97)	EP					
(71) Applicant (for all designated States except US):	CIBA SPECIALTY CHEMICALS HOLDING INC. [CH/CH]; Klybeckstrasse 141, CH-4057 Basel (CH).							
(72) Inventors; and								
(75) Inventors/Applicants (for US only):	HÜGLIN, Dietmar [DE/DE]; Dorfstrasse 3, D-79591 Eimeldingen (DE). LUTHER, Helmut [DE/DE]; Tüllingerweg 3a, D-79639 Grenzach-Wyhlen (DE). REINEHR, Dieter [DE/DE]; Wolfsheule 10, D-79400 Kandern (DE).							
(74) Common Representative:	CIBA SPECIALTY CHEMICALS HOLDING INC.; Patentabteilung, Klybeckstrasse 141, CH-4057 Basel (CH).							
(81) Designated States:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).							
Published								
With international search report.								
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.								

(54) Title: COSMETIC USE OF BIS(RESORCINYL)TRIAZINE DERIVATIVES

(57) Abstract

A description is given of the use of bis(resorcinyl)triazines of formula (1) wherein R_1 and R_2 are each independently of the other hydrogen; C_1-C_{18} alkyl; C_2-C_{18} alkenyl; a radical of formula $-CH_2-CH(-OH)-CH_2-O-T_1$; a radical of formula (1a); or a radical of formula (1b); R_3 and R_4 are each independently of the other hydrogen or C_1-C_5 alkyl; R_5 is hydroxy; C_1-C_5 alkoxy which is unsubstituted or substituted by one or several OH groups; amino; mono- or di- C_1-C_5 alkylamino; M ; a radical of formula (1c); (1d); (1e); (1f); (1g); or (1h); R_6 is a direct bond; a straight-chain or branched C_1-C_4 alkylene radical or a radical of formula (a) or (b); R_7 , R_8 and R_9 are each independently of one another C_1-C_{18} alkyl; C_1-C_{18} alkoxy or a radical of formula (1i); R_{10} , R_{11} and R_{12} are each independently of one another C_1-C_{14} alkyl which is unsubstituted or substituted by one or several OH groups; R_{13} is hydrogen; M ; C_1-C_5 alkyl; or a radical of formula (c) R_{14} is C_1-C_5 alkyl; M is a metal cation; T_1 is hydrogen; or C_1-C_8 alkyl; m_1 is 1 to 3; m_2 is 2 to 14; and p_1 is 0; or a number from 1 to 5, for protecting human and animal skin and hair from the damaging effect of UV radiation. The compounds used according to this invention are very powerful UVA absorbers with fractions in the UVB range compositions.



to the invention are very powerful UV-A absorbers with fractions in the UVB range and are particularly suitable as sunscreens in cosmetic, pharmaceutical and veterinary compositions.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Cosmetic use of bis(resorcinyl)triazine derivatives

The present invention relates to the use of selected bis(resorcinyl)triazine derivatives for protecting human and animal skin and hair from the damaging effect of UV radiation and to a cosmetic formulation comprising these bis(resorcinyl)triazine derivatives.


The bis(resorcinyl)triazines used according to this invention correspond to formula



wherein


R₁ and R₂ are each independently of the other hydrogen; branched C₅-C₁₈alkyl; C₂-

C₁₈alkenyl; a radical of formula -CH₂-CH(-OH)-CH₂-O-T₁ ; a radical of formula






R₃ and R₄ are each independently of the other hydrogen or C₁-C₅alkyl;

R₅ is hydroxy; C₁-C₅alkoxy which is unsubstituted or substituted by one or several OH groups; amino; mono- or di-C₁-C₅alkylamino; M; a radical of formula




- 2 -



R_6 is a direct bond; a straight-chain or branched C₁-C₄alkylene radical or a radical of formula $-C_{m_1}H_{2m_1}$ or $-C_{m_1}H_{2m_1}O-$;

R_7 , R_8 and R_9 are each independently of one another C₁-C₁₈alkyl; C₁-C₁₈alkoxy or a radical of



R_{10} , R_{11} and R_{12} are each independently of one another C₁-C₁₄alkyl which is unsubstituted or substituted by one or several OH groups;

R_{13} is hydrogen; M; C₁-C₅alkyl; or a radical of formula $-(CH_2)_{m_2}-O-T_1$;

R_{14} is C₁-C₅alkyl;

M is a metal cation;

T_1 is hydrogen; or C₁-C₈alkyl;

m_1 is 1 to 3;

m_2 is 2 to 14; and

p_1 is 0; or a number from 1 to 5.

C₁-C₅Alkyl, C₁-C₈alkyl and C₁-C₁₈alkyl are straight-chain or branched alkyl radicals, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, amyl, isoamyl or tert-amyl,

heptyl, octyl, iso-octyl, nonyl, decyl, undecyl, dodecyl, tetradecyl, pentadecyl, hexadecyl, heptadecyl or octadecyl.


C_1-C_5 Alkoxy and C_1-C_{18} alkoxy are straight-chain or branched alkoxy radicals, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butyloxy, sec-butyloxy, tert-butyloxy, amyloxy, iso-amylloxy or tert-amylloxy, heptyloxy, octyloxy, iso-octyloxy, nonyloxy, decyloxy, undecyloxy, dodecyloxy, tetradecyloxy, pentadecyloxy, hexadecyloxy, heptadecyloxy or octadecyloxy.

C_2-C_{18} Alkenyl is, for example, allyl, methallyl, isopropenyl, 2-butenyl, 3-butenyl, isobut enyl, n-penta-2,4-dienyl, 3-methyl-but-2-enyl, n-oct-2-enyl, n-dodec-2-enyl, iso-dodecenyl, n-dodec-2-enyl or n-octadec-4-enyl.

Examples of mono- or di- C_1-C_5 alkylamino are methylamino, ethylamino, propylamino, n-butylamino, sec-butylamino, tert-butylamino, pentylamino, dimethylamino, diethylamino; dipropylamino; dibutylamino or methylethylamino.

Examples of metal cations are the lithium, potassium, sodium, calcium, magnesium, copper or zinc ion.

According to this invention, it is preferred to use bis(resorcinyl) compounds of formula



wherein


R_{15} and R_{16} are each independently of the other branched C_5-C_{18} alkyl; or
 $-CH_2-CH(-OH)-CH_2-O-T_1$;

R_{17} and R_{18} are each independently of the other hydrogen or C_1-C_5 alkyl; and
 T_1 is hydrogen; or C_1-C_5 alkyl.

Preeminently interesting compounds are those of formula (2), wherein R₁₅ and R₁₆ are each independently of the other branched C₅-C₁₈alkyl or -CH₂-CH(-OH)-CH₂-O-T₁;
 R₁₇ and R₁₈ are hydrogen; and
 T₁ is hydrogen; or C₁-C₅alkyl;
 and in particular those compounds of formula (2), wherein R₁₅ and R₁₆ are each independently of the other branched C₅-C₁₈alkyl; and R₁₇ and R₁₈ are hydrogen.

Very particularly preferred are those triazine compounds of formula (2), wherein R₁₅ and R₁₆ have the same meaning.

Other triazine derivatives which may be used according to this invention are those corresponding to formula



wherein

R₁₉ and R₂₀ are each independently of the other the radical of formula $-\text{CH}_2\text{C}(=\text{O})\text{O}-\text{R}_{23}$; and

R₂₁, R₂₂ and R₂₃ are each independently of one another hydrogen; or C₁-C₁₀alkyl.

Examples to be mentioned of compounds of formula (1) are:

2-(4'-methylphenyl)-4,6-bis(2'-hydroxy-4'-n-octyloxyphenyl)-1,3,5-triazine;

2-(2',4'-dimethylphenyl)-4,6-bis(2'-hydroxy-4'-n-octyloxyphenyl)-1,3,5-triazine;

2-phenyl-4,6-bis(2'-hydroxy-4'-n-octyloxyphenyl)-1,3,5-triazine;

2-phenyl-4,6-bis(2'-hydroxy-4'-n-octyloxyphenyl)-1,3,5-triazine;

2-phenyl-4-{{4-(3-(2-propyloxy)-2-hydroxypropyloxy)-2-hydroxy]phenyl}-6-(4-methoxyphenyl)-1,3,5-triazine;
 2-(4'-methylphenyl)-4,6-bis{{4-(2-ethylhexyloxy)-2-hydroxy]phenyl}-1,3,5-triazine;
 2-(2',4'-dimethylphenyl)-4,6-bis{{4-(2-ethylhexyloxy)-2-hydroxy]phenyl}-1,3,5-triazine;
 2-phenyl-4,6-bis{{4-(2-ethylhexyloxy)-2-hydroxy]phenyl}-1,3,5-triazine;
 2-phenyl-4,6-bis{{4-(tris(trimethylsiloxy)silylpropyloxy)-2-hydroxy]phenyl}-1,3,5-triazine;
 2-phenyl-4,6-bis{{4-(2"methylpropenyloxy)-2-hydroxy]phenyl}-1,3,5-triazine;
 2-phenyl-4,6-bis{{4-(3-(2-propyloxy)-2-hydroxypropyloxy)-2-hydroxy]phenyl}-1,3,5-triazine; or
 2-phenyl,4,6-bis{{4-(2-ethylhexyloxy)-2-hydroxy]phenyl}-1,3,5-triazine.

Other examples of triazine derivatives are listed in Table 1:

Table 1:

R _a	R _b	R _c
H	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{O}- \\ \\ \text{CH}_3 \end{array} $	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{O}- \\ \\ \text{CH}_3 \end{array} $
CH ₃	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{O}- \\ \\ \text{CH}_3 \end{array} $	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\text{C}-\text{O}- \\ \\ \text{CH}_3 \end{array} $
H	$ -\text{NH}-\text{C}_6\text{H}_4-\text{COOC}_4\text{H}_9 $	$ -\text{NH}-\text{C}_6\text{H}_4-\text{COOC}_4\text{H}_9 $
CH ₃	$ -\text{NH}-\text{C}_6\text{H}_4-\text{COOC}_4\text{H}_9 $	$ -\text{NH}-\text{C}_6\text{H}_4-\text{COOC}_4\text{H}_9 $
H	-OH	-OH

Table 1 (continuation)






<u>R_a</u>	<u>R_b</u>	<u>R_c</u>
CH ₃	-OH	-OH
H	-OH	-OH
H	-OM M= alkali metal, alkaline earth metal, Cu, Zn, Mg	-OM M= alkali metal, alkaline earth metal, Cu, Zn, Mg
CH ₃	-OM M= alkali metal, alkaline earth metal, Cu, Zn, Mg	-OM M= alkali metal, alkaline earth metal, Cu, Zn, Mg
H	O ⁻ + N(CH ₂ CH ₂ OH) ₃	O ⁻ + N(CH ₂ CH ₂ OH) ₃
CH ₃	O ⁻ + N(CH ₂ CH ₂ OH) ₃	O ⁻ + N(CH ₂ CH ₂ OH) ₃
H	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\overset{+}{\text{N}}\text{---}(\text{CH}_2)_n\text{---O---} \\ \\ \text{CH}_3 \end{array} \quad n=2-14 $	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\overset{+}{\text{N}}\text{---}(\text{CH}_2)_n\text{---O---} \\ \\ \text{CH}_3 \end{array} \quad n=2-14 $
CH ₃	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\overset{+}{\text{N}}\text{---}(\text{CH}_2)_n\text{---O---} \\ \\ \text{CH}_3 \end{array} \quad n=2-14 $	$ \begin{array}{c} \text{CH}_3 \\ \\ \text{H}_3\text{C}-\overset{+}{\text{N}}\text{---}(\text{CH}_2)_n\text{---O---} \\ \\ \text{CH}_3 \end{array} \quad n=2-14 $
H		
CH ₃		
H		

Table 1 (continuation)

<u>R_a</u>	<u>R_b</u>	<u>R_c</u>
CH ₃		
H		
CH ₃		
H		
CH ₃		

The bis(resorcinyl)triazines which may be used in accordance with this invention can be prepared in different manner, for example using aromatic nitrile compounds as starting compounds. The dichlorotriazine compound of formula



can be prepared, for example, by reacting benzonitrile with dicyandiamide to the diamino-triazine compound which is subsequently saponified to the dihydroxytriazine (tautomer = -dione) and then converted to the dichlorotriazine compound of formula (1k) with thionyl chloride. The two resorcinol groups are then introduced in commonly known manner by Friedel-Crafts acylation of resorcinol in the presence of a Lewis acid, preferably aluminium chloride. In the third step, the free hydroxyl groups in p-position are etherified, depending on the meaning of R₁ and R₂ in the compound of formula (1) by alkylation or acid-catalysed addition of glycidyl ethers.

Other processes for the preparation of the triazine derivatives which can be used in accordance with this invention are described in EP-A-0,775,698.

The bis(resorcinyl)triazine compounds of formula (1) used according to this invention are particularly suitable as UV filters, i.e. for protecting ultraviolet-sensitive organic materials, in particular human and animal skin and hair, from the harmful effect of UV radiation. These compounds are, in particular, very powerful UVA absorbers with fractions in the UVB range and can be produced at low cost. Accordingly, these compounds are suitable for use as light stabilisers in cosmetic, pharmaceutical and veterinary compositions. They can be used in dissolved as well as in micronised state.

Accordingly, this invention also relates to a cosmetic formulation, which comprises at least one compound of formula (1) as well as cosmetically compatible carriers or auxiliaries.



For cosmetic use, the light stabilisers of this invention usually have an average particle size in the range from 0.02 to 2, preferably from 0.05 to 1.5 and, most preferably, from 0.1 to 1.0 m. The insoluble UV absorbers used according to this invention can be brought to the desired particle size by customary methods, for example by grinding e.g. with a jet, ball, vibratory or hammer mill. Grinding is preferably carried out in the presence of 0.1 to 30 by

weight, preferably of 0.5 to 15 % by weight, based on the UV absorber, of a grinding aid such as an alkylated vinylpyrrolidone polymer, vinylpyrrolidone-vinylacetate copolymer, acylglutamate or, preferably, phospholipid.



In addition to the inventive UV absorbers, the cosmetic formulation can also contain one or more than one other UV protective, for example an organic UV absorber from the classes of the p-aminobenzoic acid derivatives, salicylic acid derivatives, benzophenone derivatives, dibenzoylmethane derivatives, diphenylacrylate derivatives, benzofuran derivatives, polymeric UV absorbers comprising one or more than one organosilicon radical, cinnamic acid derivatives, camphor derivatives, trianilino-s-triazine derivatives, phenylbenzimidazolesulfonic acid and the salts thereof, menthylanthranilates, benzotriazole derivatives, and/or an inorganic micropigment selected from the group consisting of aluminium oxide- or silicium dioxide-coated TiO_2 , zinc oxide or mica.

- Examples of p-aminobenzoic acid derivative compounds:

4-aminobenzoic acid (PABA); ethyldihydroxypropyl-PABA of formula








wherein m, n and x have the same meaning and are each at most 25;
octyldimethyl PABA of formula



- Examples of salicylic acid derivative compounds:



homomenthylsalicylate of formula




- Examples of benzophenone derivative compounds:
benzophenone-3-(2-hydroxy-4-methoxybenzophenone), benzophenone-4-(2-hydroxy-4-methoxybenzophenone-5-sulfonic acid) or benzophenone-8-(2,2'-dihydroxy-4-methoxybenzophenone).
- Examples of dibenzoylmethane derivative compounds:
butylmethoxydibenzoylmethane[1-(4-tert-butyl)-3-(4-methoxyphenyl)propane-1,3-dione].
- Examples of diphenylacrylate derivative compounds:
octocrylene(2-ethylhexyl-2-cyano-3,3'-diphenylacrylate) or etocrylene(ethyl-2-cyano-3,3'-diphenylacrylate).

- Examples of benzofuran derivative compounds:

3-(benzofuranyl)-2-cyanoacrylate, 2-(2-benzofuranyl)-5-tert-butylbenzoxazole or 2-(p-aminophenyl)benzofurane and, in particular, the compound of formula




benzylidenemalonate derivative, in particular the compound of formula




wherein

R₂₄ is hydrogen or O-Me and

r is approximately 7; the compound of formula





- Examples of cinnamate compounds:
octylmethoxycinnamate (4-methoxycinnamic acid-2-ethylhexyl), diethanolaminomethoxycinnamate (the diethanolamine salt of 4-methoxycinnamic acid), isoamyl-p-methoxycinnamate (4-ethoxycinnamic acid-2-isoamyl), 2,5-diisopropylmethylicinnamate or a cinnamic acid amido derivative.
- Examples of camphor derivative compounds:
4-methyl-benzylidenecamphor [3-(4'-methyl)benzylidenebornan-2-one], 3-benzylidene-camphor (3-benzylidenebornan-2-one), polyacrylamidomethylbenzylidenecamphor {N-[2(and 4)-2-oxyborn-3-ylidenemethyl]benzyl]acrylamide polymer}, trimoniumbenzylidenecamphorsulfate-[3-(4'-trimethylammonium)benzylidenebornan-2-onemethylsulfate], terephthalydenedicamphorsulfonic acid {3,3'-(1,4-phenylenedimethine)-bis(7,7-dimethyl-2-oxobicyclo-[2.2.1]heptane-1-methanesulfonic acid} or the salts thereof, or benzylidenecamphorsulfonic acid [3-(4'-sulfo)benzylidenebornan-2-one] or the salts thereof.
- Examples of trianilino-s-triazine derivative compounds:
octyltriazine-[2,4,6-trianilino-(p-carbo-2'-ethyl-1'-oxy)-1,3,5-triazine, and also the trianilino-s-triazine derivatives disclosed in US-A-5,332,568, US-A-5,252,323, WO 93/17002 and WO 97/03642 and EP-A-0,517,104.
- Examples of benzotriazole compounds:
2-(2-hydroxy-5-methylphenyl)benzotriazole.

The novel cosmetic formulation comprises 0.1 to 15 by weight, preferably 0.5 to 10 % by weight, based on the total weight of the formulation, of a UV absorber or of a mixture of UV absorbers as well as a cosmetically compatible auxiliary.

The cosmetic formulation can be prepared by physically mixing the UV absorber(s) with the auxiliary by customary methods, for example by simply stirring the individual components together.

The novel cosmetic formulation can be formulated as water-in-oil or oil-in-water emulsion, as oil-in-alcohol lotion, as vesicular dispersion of a ionic or nonionic amphiphilic lipid, as gel, solid stick or as aerosol formulation.

As water-in-oil or oil-in-water emulsion, the cosmetically compatible auxiliary preferably comprises 5 to 50% of an oil phase, 5 to 20% of an emulsifier and 30 to 90% of water. The oil phase can contain any oil suitable for cosmetic formulations, for example one or more than one hydrocarbon oil, wax, natural oil, silicone oil, fatty acid ester or fatty alcohol. Preferred mono- or polyols are ethanol, isopropanol, propylene glycol, hexylene glycol, glycerol and sorbitol.

It is possible to use any conventional emulsifier for the inventive cosmetic formulation, for example one or more than one ethoxylated ester of natural derivatives, such as polyethoxylated ester of hydrogenated castor oil; or a silicone oil emulsifier such as silicone polyol; an optionally ethoxylated fatty acid soap; an ethoxylated fatty alcohol; an optionally ethoxylated sorbitan ester; an ethoxylated fatty acid; or an ethoxylated glyceride.

The cosmetic formulation can also contain other components, for example emollients, emulsion stabilisers, skin moisturisers, suntan accelerators, thickeners such as xanthane, moisture retention agents such as glycerol, preservatives, fragrances and colourants.


The novel cosmetic formulation is distinguished by excellently protecting the human skin against the harmful effect of sunlight.

In the following Examples percentages are by weight. The amounts given for the bis(resorcinyl)triazine compounds refer to the pure substance.

Examples:Use Examples for cosmetic light protection

The light stabilisers are determined in accordance with the method of Diffey and Robson, J. Soc. Cosmet. Chem. 40, 127 - 133 (1989) using an SPF analyser (Optometrix, SPF 290).

To determine the photostabilities, the filter substances are dissolved in ethanol ($c = 1 \cdot 10^{-5} - 5 \cdot 10^{-5}$ M) and irradiated, with stirring, in a quartz cuvette using a metal halide lamp (Macam) ($I_{UVB} = 0.4 - 8.0$ mW/cm²). To convert to the solar spectrum (CIE D65-norm daylight, standardised to $I_{UVB} = 0.127$ mW/cm²), the integral over the products of the wavelength-resolved lamp intensity and the corresponding absorption values of the respective UV absorbers from 290 to 400 nm is calculated and is then divided by the integral over the products of the D65 light intensities and the corresponding absorption values of the respective UV absorber in the range from 290 to 400 nm. This factor is multiplied with the half life period for degradation under metal halide lamp irradiation to obtain the corresponding half life period under solar irradiation. The half life period for the photodegradation under lamp irradiation is determined via UV-spectroscopic measurement of the extinction at the wavelength of the maximum absorption and subsequent exponential fit. Using the process described, the half life periods for the photodegradation in D65 light are thus obtained.

Example 1: o/w emulsion with the compound of formula (101)

λ_{\max} : 356 nm (ethanol)

ϵ_{\max} : 41 400 M⁻¹cm⁻¹

(A):

triazine UV absorber of formula (101)	4 g
sesame oil	10 g
glyceryl stearate	4 g
stearic acid	1 g
cetyl alcohol	0.5 g
Polysorbate 20	0.2 g

(B):

propylene glycol	4 g
propylparaben	0.05 g
methylparaben	0.15
triethanolamine	0.1 g
Carbomer 934	0.1 g
water	ad 100 ml

Preparation of the emulsionPhase (A):

The UV absorber is first dissolved in sesame oil and the other components of (A) are then added and fused together.

Phase (B):

Propylparaben and methylparaben are dissolved in propylene glycol. 60ml water are then added, the mixture is heated to 70°C and Carbomer 934 is emulsified therein.

Emulsion:

(A) is slowly added to B with vigorous mechanical energy input. The volume is adjusted to 100 ml by addition of water.

The sunscreen factors and photostabilities obtained are compiled in Table 1.

<u>Table 1:</u>	<u>Concentration</u>	<u>Sunscreen factor *)</u>	<u>Photostability**) [h]</u>
compound of formula (101)	4%	7.1	6000

*) according to Diffey and Robson

**) as half life period of the photodegradation in D65 light in ethanolic solution

The sunscreen factor can be varied via the UV absorber concentration.

The results demonstrate that the active substance has high photostability and that a good sunscreen factor may be achieved even at low concentration.

Other physically defined measures for UVA characterisation:

Table 2: Other physically defined measures for UVA characterisation

	<u>compound of formula (101)</u> <u>4% emulsion</u>
T(360) ¹	0.6 %
UVA/UVB ratio ²	0.79
λ_{crit} [nm] ³	373

Example 2: Sunscreen Emulsion containing new oil soluble filter (101) and Octyl methoxycinnamate

<u>Phase</u>	<u>Component</u>	<u>% b.w.</u>
(A)	Water (aqua), deionized	54.90
	Xanthan gum, 2% solution	15.00
	Propylene glycol	1.50
	Tetrasodium EDTA	0.10
(B)	Isopropylmyristate	10.00
	Triazine UV-absorber with formula (101)	3.00
(C)	Octyl methoxycinnamate	7.50
	Glyceryl stearate (and) PEG-100 stearate	3.00
	Cyclomethicone	1.00
	Cetearyl alcohol (and) ceteareth-20 (Amerchol)	3.00
(D)	Propylene glycol (and) methylparaben (and) propylparaben (and) diazolidinyl urea (ISP)	1.00

¹ Transmission at 360 nm


² Ratio of the areas standardised to the wavelengths under the absorption curve from 290 to 320 and from 320 to 400 nm

³ Wavelength at which 90% of the area under the absorption curve is reached when starting the integration at 290 nm and defining the area between 290 and 400 nm as 100%.

Procedure: The components of (A) are mixed in order and heated to 75°C. Components of (B) are mixed, heated to 75°C and stirred until filter is dissolved. Then (C) is added to B and mixed for 20 minutes (= mixture (BC)). (D) is then added to (BC) (= mixture (BCD)). (A) is added to (BCD) while mixing and cooling to room temperature.

Resulting SPF (Sun Protection Factor): 16 (method according example 1)

Example 3: Sunscreen Formulation containing new water soluble filter of the formula (102) and Octyl methoxycinnamate



$\lambda_{\text{max}} = 352\text{nm}$ (ethanol)

$\epsilon_{\text{max}} = 39\,800\text{ M}^{-1}\text{ cm}^{-1}$


<u>Phase</u>	<u>Component</u>	<u>% b.w.</u>
(A)	Triazine UV-absorber with formula (102)	4.00
	Carbomer	0.30
(B)	water (aqua), deionized	57.93
(C)	Propylene glycol	5.00
	Methylparaben	0.20
	Propylparaben	0.10
	Triethanolamine	0.45
	Tetrasodium EDTA	0.02
(D)	Octyl methoxycinnamate	5.00
	Glyceryl stearate (and) PEG-100 stearate	1.00
	Cyclomethicone	5.00
	Glyceryl stearate	4.00
	Stearic acid	2.50
	Isostearyl isostearate	10.00
	Hydrogenated castor oil	2.00
	C ₁₂₋₁₅ alkyl benzoate	2.50

Procedure: (A) is dispersed in (B) and, when uniform, components of (C) (= mixture (ABC)) are added in order. While heating to 85°C, components of (D) are combined in a separate vessel and heated with mixing to 85°C. When (ABC) and (D) are both at 85°C and uniform, (D) is slowly added to ABC with strong mixing. 15 minutes after the last of D has been added cooling batch is started and cool to room temperature.




Resulting SPF (Sun Protection Factor): 12 (method according example 1)

What is claimed is


1. Use of a UV absorber of formula



wherein


 R_1 and R_2 are each independently of the other hydrogen; branched C_5 - C_{18} alkyl; C_2 - C_{18} alkenyl; a radical of formula $-\text{CH}_2\text{-CH}(\text{-OH})\text{-CH}_2\text{-O-T}_1$; a radical of formula R_3 and R_4 are each independently of the other hydrogen or C_1 - C_5 alkyl; R_5 is hydroxy; C_1 - C_5 alkoxy which is unsubstituted or substituted by one or several OH groups; amino; mono- or di- C_1 - C_5 alkylamino; M ; a radical of formula

- 21 -



R_6 is a direct bond; a straight-chain or branched C₁-C₄alkylene radical or a radical of formula $-C_{m_1}H_{2m_1}$ or $-C_{m_1}H_{2m_1}O-$;

R_7 , R_8 and R_9 are each independently of one another C₁-C₁₈alkyl; C₁-C₁₈alkoxy or a radical of



R_{10} , R_{11} and R_{12} are each independently of one another C₁-C₁₄alkyl which is unsubstituted or substituted by one or several OH groups;

R_{13} is hydrogen; M; C₁-C₅alkyl; or a radical of formula $-(CH_2)_{m_2}-O-T_1$;

R_{14} is C₁-C₅alkyl;

M is a metal cation;

T_1 is hydrogen; or C₁-C₈alkyl;

m_1 is 1 to 3;


m_2 is 2 to 14; and

p_1 is 0; or a number from 1 to 5,

for protecting human and animal skin and hair from the damaging effect of UV radiation.

2. Use according to claim 1, wherein the bis(resorcinyl)triazine used is the compound of formula

- 22 -



wherein

R_{15} and R_{16} are each independently of the other branched C_5 - C_{18} alkyl or $-CH_2-CH(-OH)-CH_2-$ $O-T_1$;

R_{17} and R_{18} are hydrogen; and

T_1 is hydrogen; or C_1 - C_5 alkyl.

3. Use according to claim 2, wherein, in formula (2),

R_{15} and R_{16} are each independently of the other C_1 - C_{18} alkyl or $-CH_2-CH(-OH)-CH_2-O-T_1$;

R_{17} and R_{18} are hydrogen; and

T_1 is hydrogen; or C_1 - C_5 alkyl.


4. Use according to claim 3, wherein, in formula (2),

R_{15} and R_{16} are each independently of the other branched C_5 - C_{18} alkyl; and

R_{17} and R_{18} are hydrogen.

5. Use according to any one of claims 2 to 4, wherein R_{15} and R_{16} in formula (2) have the same meaning.

6. Use according to claim 1, which comprises using a bis(resorcinyl)triazine of formula



wherein

R_{19} and R_{20} are each independently of the other the radical of formula $-\text{CH}_2-\text{C}(=\text{O})-\text{O}-R_{23}$; and

R_{21} , R_{22} and R_{23} are each independently of one another hydrogen; or C_1-C_{10} alkyl.

7. A cosmetic formulation, which comprises a UV absorber of formula (1) according to claim 1 together with cosmetically compatible carriers or auxiliaries.

8. A formulation according to claim 7, which comprises additional UV protectives.

9. A formulation according to either claim 7 or claim 8, wherein the additional UV protectives are an organic UV absorber from the classes of the p-aminobenzoic acid derivatives, salicylic acid derivatives, benzophenone derivatives, dibenzoylmethane derivatives, diphenylacrylate derivatives, benzofuran derivatives, polymeric UV absorbers comprising one or more than one organosilicon radical, cinnamic acid derivatives, camphor derivatives, trianilino-s-triazine derivatives, phenylbenzimidazolesulfonic acid and the salts thereof, menthylanthranilates, benzotriazole derivatives, and/or an inorganic micropigment selected from aluminium oxide- or silicium dioxide-coated TiO_2 , zinc oxide or mica.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/05042

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K7/42

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category ^o	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CH 480 090 A (CIBA) 31 October 1969 see the whole document ---	1-9
A	BE 639 329 A (CIBA) see the whole document ---	1-9
A	BE 650 932 A (CIBA) 25 January 1965 see the whole document ---	1-9
A	EP 0 775 698 A (CIBA SPECIALTY CHEMICALS HOLDING) 28 May 1997 cited in the application see the whole document ---	1-9
A	WO 95 22959 A (CIBA-GEIGY) 31 August 1995 see the whole document ---	1-9
		-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

^o Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority, claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

14 December 1998

18/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fischer, J.P.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/05042

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	EP 0 878 469 A (CIBA SPECIALTY CHEMICALS HOLDING) 18 November 1998 see the whole document -----	1-9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/05042

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
CH 480090	A 31-10-1969	NONE		
BE 639329	A	BE 639330 A		
		CH 467833 A		
		DE 1215712 B		
		DE 1216874 B		
		FR 1379138 A	01-03-1965	
		FR 1381452 A	19-03-1965	
		GB 1033387 A		
		GB 1033388 A		
		NL 123744 C		
		NL 299880 A		
		NL 299881 A		
		SE 315891 B	13-10-1969	
		SE 315892 B	13-10-1969	
		US 3242175 A	22-03-1966	
BE 650932	A 25-01-1965	CH 469053 A		
		DE 1291511 B		
		FR 1405559 A	22-11-1965	
		GB 1065425 A		
		NL 6408514 A	27-01-1965	
		SE 316470 B	27-10-1969	
		US 3249608 A	03-05-1966	
EP 775698	A 28-05-1997	DE 19543730 A	28-05-1997	
		AU 7194696 A	29-05-1997	
		JP 9188666 A	22-07-1997	
		NZ 299786 A	24-09-1998	
		SG 45516 A	16-01-1998	
WO 9522959	A 31-08-1995	AU 1665395 A	11-09-1995	
		BR 9506935 A	09-09-1997	
		EP 0746305 A	11-12-1996	
		GB 2286774 A, B	30-08-1995	
		JP 9509421 T	22-09-1997	
		ZA 9501459 A	23-08-1995	
EP 878469	A 18-11-1998	AU 6598798 A	19-11-1998	