

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

(10) **DK/EP 3470535 T3**

-
- (51) Int.Cl.: **C 12 Q 1/6886 (2018.01)**
- (45) Oversættelsen bekendtgjort den: **2020-06-29**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-04-01**
- (86) Europæisk ansøgning nr.: **18190058.0**
- (86) Europæisk indleveringsdag: **2004-06-17**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-04-17**
- (30) Prioritet: **2003-06-24 US 482339 P**
- (62) Stamansøgningsnr: **16196459.8**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LI LU MC NL PL PT RO SE SI SK TR**
- (73) Patenthaver: **Genomic Health, Inc., 301 Penobscot Drive, Redwood City, CA 94063, USA**
NSABP Foundation, Inc., Two Allegheny Center, Suite 1200, Pittsburgh PA 15212-5402, USA
- (72) Opfinder: **BRYANT, John L., -, deceased, USA**
PAIK, Soonmyung, Yanghwa-ro 45, Mapo-guMecenatpolis 102-2202, Seoul 04036, Sydkorea
BAKER, Joffre, 1400 Avery Street, Montara, CA 94037, USA
SHAK, Steve, 648 Fairway Circle, Hillsborough, CA 94010, USA
- (74) Fuldmægtig i Danmark: **AWA Denmark A/S, Strandgade 56, 1401 København K, Danmark**
- (54) Benævnelse: **FORUDSIGELSE AF SANDSYNLIGHED FOR TILBAGEVENDEN AF CANCER**
- (56) Fremdragne publikationer:
US-A1- 2002 156 263
REINHARD BOS ET AL: "Levels of hypoxia-inducible factor-1? independently predict prognosis in patients with lymph node negative breast carcinoma", CANCER, vol. 97, no. 6, 15 March 2003 (2003-03-15) , pages 1573-1581, XP055132674, ISSN: 0008-543X, DOI: 10.1002/cncr.11246
PINTO A E ET AL: "c-erbB-2 oncoprotein overexpression identifies a subgroup of estrogen receptor positive (ER+) breast cancer patients with poor prognosis", ANNALS OF ONCOLOGY, OXFORD UNIVERSITY PRESS, GB, vol. 12, 1 January 2001 (2001-01-01), pages 525-533, XP002406335, ISSN: 1569-8041, DOI: 10.1023/A:1011163211802
R. SILVESTRI ET AL: "p53 as an Independent Prognostic Marker in Lymph Node-Negative Breast Cancer Patients", JNCI JOURNAL OF THE NATIONAL CANCER INSTITUTE, vol. 85, no. 12, 16 June 1993 (1993-06-16) , pages 965-970, XP055132808, ISSN: 0027-8874, DOI: 10.1093/jnci/85.12.965
ADAM AMEUR ET AL: "Total RNA sequencing reveals nascent transcription and widespread co-transcriptional splicing in the human brain", NATURE STRUCTURAL & MOLECULAR BIOLOGY, vol. 18, no. 12, 6 November 2011 (2011-11-06), pages 1435-1440, XP055086600, ISSN: 1545-9993, DOI: 10.1038/nsmb.2143
TUTT ANDREW ET AL: "Risk estimation of distant metastasis in node-negative, estrogen receptor-positive breast cancer patients using an RT-PCR based prognostic expression signature", BMC CANCER, BIOMED CENTRAL, LONDON, GB, vol. 8, no. 1, 21 November 2008 (2008-11-21), page 339, XP021048927, ISSN: 1471-

Fortsættes ...

DescriptionBackground of the Invention5 Field of the Invention

[0001] The present disclosure provides gene sets the expression of which is important in the diagnosis and/or prognosis of cancer.

10 Description of the Related Art

[0002] Oncologists have a number of treatment options available to them, including different combinations of chemotherapeutic drugs that are characterized as "standard of care," and a number of drugs that do not carry a label claim for particular cancer, but for which there is evidence of efficacy in that cancer. Best likelihood of good treatment outcome requires that patients be assigned to optimal available cancer treatment, and that this assignment be made as quickly as possible following diagnosis.

[0003] Currently, diagnostic tests used in clinical practice are single analyte, and therefore do not capture the potential value of knowing relationships between dozens of different markers. Moreover, diagnostic tests are frequently not quantitative, relying on immunohistochemistry. This method often yields different results in different laboratories, in part because the reagents are not standardized, and in part because the interpretations are subjective and cannot be easily quantified. RNA-based tests have not often been used because of the problem of RNA degradation over time and the fact that it is difficult to obtain fresh tissue samples from patients for analysis. Fixed paraffin-embedded tissue is more readily available and methods have been established to detect RNA in fixed tissue. However, these methods typically do not allow for the study of large numbers of genes (DNA or RNA) from small amounts of material. Thus, traditionally fixed tissue has been rarely used other than for immunohistochemistry detection of proteins.

[0004] In the past few years, several groups have published studies concerning the classification of various cancer types by microarray gene expression analysis (see, e.g. Golub et al., Science 286:531-537 (1999); Bhattacharjee et al., Proc. Natl. Acad. Sci. USA 98:13790-13795 (2001); Chen-Hsiang et al., Bioinformatics 17 (Suppl. 1):S316-S322 (2001); Ramaswamy et al., Proc. Natl. Acad. Sci. USA 98:15149-15154 (2001)). Certain classifications of human breast cancers based on gene expression patterns have also been reported (Martin et al., Cancer Res. 60:2232-2238 (2000); West et al., Proc. Natl. Acad. Sci. USA 98:11462-11467 (2001); Sorlie et al., Proc. Natl. Acad. Sci. USA 98:10869-10874 (2001); Yan et al., Cancer Res. 61:8375-8380 (2001)). However, these studies mostly focus on improving and refining the already established classification of various types of cancer, including breast cancer, and generally do not provide new insights into the relationships of the differentially expressed genes, and do not link the findings to treatment strategies in order to improve the clinical outcome of cancer therapy.

[0005] Although modern molecular biology and biochemistry have revealed hundreds of genes whose activities influence the behavior of tumor cells, state of their differentiation, and their sensitivity or resistance to certain therapeutic drugs, with a few exceptions, the status of these genes has not been exploited for the purpose of routinely making clinical decisions about drug treatments. One notable exception is the use of estrogen receptor (ER) protein expression in breast carcinomas to select patients to treatment with anti-estrogen drugs, such as tamoxifen. Another exceptional example is the use of ErbB2 (Her2) protein expression in breast carcinomas to select patients with the Her2 antagonist drug Herceptin® (Genentech, Inc., South San Francisco, CA).

[0006] Despite recent advances, the challenge of cancer treatment remains to target specific treatment regimens to pathogenically distinct tumor types, and ultimately personalize tumor treatment in order to maximize outcome. Hence, a need exists for tests that simultaneously provide predictive information about patient responses to the variety of treatment options. This is particularly true for breast cancer, the biology of which is poorly understood. It is clear that the classification of breast cancer into a few subgroups, such as ErbB2⁺ subgroup, and subgroups characterized by low to absent gene expression of the estrogen receptor (ER) and a few additional transcriptional factors (Perou et. al., Nature 406:747-752 (2000)) does not reflect the cellular and molecular heterogeneity of breast cancer, and does not allow the design of treatment strategies maximizing patient response.

[0007] In particular, once a patient is diagnosed with cancer, such as breast or ovarian cancer, there is a strong need for methods that allow the physician to predict the expected course of disease, including the likelihood of cancer recurrence, long-term survival of the patient, and the like, and select the most appropriate treatment option accordingly.

[0008] Bos et al. (Cancer, Vol. 97, No 6, 1573-1518, 2003) discloses using HIF- α RNA to predict the likelihood of disease-free survival of a node-negative, invasive ductal breast cancer patient. US2002/156263 discloses MELK (KIAA0175) as one of a large number of genes found to be differentially expressed in breast cancer.

Summary of the Invention

[0009] The present invention is defined by the appended claims 1 to 8.

[0010] The present invention accommodates the use of archived paraffin-embedded biopsy material for assay of all markers in the set, and therefore is compatible with the most widely available type of biopsy material. It is also compatible with several different methods of tumor tissue harvest, for example, via core biopsy or fine needle aspiration. Further, for each member of the gene set, the invention specifies oligonucleotide sequences that can be used in the test.

[0011] In one aspect, the present invention concerns a method of predicting the likelihood of disease-free survival of a node-negative, ER positive, invasive ductal breast cancer patient, comprising determining the expression level of the RNA transcript of MELK in a breast cancer cell or tissue sample obtained from the patient, normalized against the expression level of a reference set of RNA transcripts comparing the normalized expression level of MELK from the patient to a normalized expression level of MELK in a breast cancer tissue reference set comprising patients who were (a) alive without local, regional, or distant breast cancer recurrence, (b) alive with contralateral breast cancer recurrence, (c) alive with non-breast second primary cancer, or (d) died prior to breast cancer recurrence, wherein an increased normalized expression level of MELK from the patient indicates a decreased likelihood of disease-free survival.

[0012] In various embodiments, the expression level of at least 2, or at least 5, or at least 10, or at least 15, or at least 20, or at least 25 prognostic RNA transcripts or their expression products is determined.

[0013] In another embodiment, the cancer is breast cancer or ovarian cancer.

[0014] The cancer is node negative, ER positive breast cancer. In a further embodiment, the RNA comprises intronic RNA. The expression level of the RNA transcript of MELK is determined, wherein expression of MELK indicates a decreased likelihood of long-term survival without cancer recurrence.

[0015] The invention concerns a method of predicting the likelihood of long-term survival of a cancer patient without the recurrence of cancer, comprising determining the expression level of MELK RNA transcripts in a cancer cell obtained from said patient, normalized against the expression level of all RNA transcripts or their products in the cancer cell, or of a reference set of RNA transcripts or their expression products, wherein increased expression of MELK indicates a decreased likelihood of long term survival without cancer recurrence.

[0016] In an embodiment of this method, the RNA is isolated from a fixed, wax-embedded breast cancer tissue specimen of the patient.

[0017] In another embodiment, the RNA is isolated from core biopsy tissue or fine needle aspirate cells.

[0018] In a different aspect, the disclosure concerns an array comprising polynucleotides hybridizing to two or more of the following genes: B_Catenin; BAG1; BIN1; BUB1; C20_orf1; CCNB1; CCNE2; CDC20; CDH1; CEGP1; CIAP1; cMYC; CTSL2; DKFZp586M07; DR5; EpCAM; EstR1; FOXM1; GRB7; GSTM1; GSTM3; HER2; HNRPA; ID1; IGF1R; ITGA7; Ki_67; KNSL2; LMNB1; MCM2; MELK; MMP12; MMP9; MYBL2; NEK2; NME1; NPD009; PCNA; PR; PREP; PTTG1; RPLPO; Src; STK15; STMY3; SURV; TFRC; TOP2A; and TS, immobilized on a solid surface.

[0019] In an aspect of the disclosure, the array comprises polynucleotides hybridizing to two or more of the following genes: MMP9, GSTM1, MELK, PR, DKFZp586M07, GSTM3, CDC20, CCNB1, STMY3, GRB7, MYBL2, CEGP1, SURV, LMNB1, CTSL2, PTTG1, BAG1, KNSL2, CIAP1, PREP, NEK2, EpCAM, PCNA, C20_orf1, ITGA7, ID1 B_Catenin, EstR1, CDH1, TS HER2, and cMYC.

[0020] In another aspect, the array comprises polynucleotides hybridizing to two or more of the following genes: GRB7, SURV, PR, LMNB1, MYBL2, HER2, GSTM1, MELK, S20_orf1, PTTG1, BUB1, CDC20, CCNB1, STMY3, KNSL2, CTSL2, MCM2, NEK2, DR5, Ki_67, CCNE2, TOP2A, PCNA, PREP, FOXM1, NME1, CEGP1, BAG1, STK15, HNRPA, EstR1, MMP9, DKFZp586M07, TS, Src, BIN1, NP009, RPLPO, GSTM3, MMP12, TFRC, and IGF1R.

[0021] In a further aspect, the arrays comprise polynucleotides hybridizing to at least 3, or at least 5, or at least 10, or at least 15, or at least 20, or at least 25 of the listed genes.

[0022] In a still further aspect, the arrays comprise polynucleotides hybridizing to all of the listed genes.

[0023] In yet another aspect, the arrays comprise more than one polynucleotide hybridizing to the same gene.

[0024] In an additional aspect, the arrays comprise intron-based sequences.

[0025] In another aspect, the polynucleotides are cDNAs, which can, for example, be about 500 to 5000 bases long.

[0026] In yet another aspect, the polynucleotides are oligonucleotides, which can, for example, be about 20 to 80 bases long.

[0027] The arrays can, for example, be immobilized on glass, and can contain hundreds of thousand, e.g. 330,000 oligonucleotides.

[0028] In a further aspect, the disclosure concerns a method of predicting the likelihood of long-term survival of a patient diagnosed with invasive breast cancer, without the recurrence of breast cancer, comprising the steps of

- (a) determining the expression levels of the RNA transcripts or the expression products of genes of a gene set selected from the group consisting of B_Catenin; BAG1; BIN1; BUB1; C20_orf1; CCNB1; CCNE2; CDC20; CDH1; CEGP1; CIAP1; cMYC; CTSL2; DKFZp586M07; DR5; EpCAM; EstR1; FOXM1; GRB7; GSTM1; GSTM3; HER2;

HNRPA7; ID1; IGF1R; ITGA7; Ki_67; KNSL2; LMNB1; MCM2; MELK; MMP12; MMP9; MYBL2; NEK2; NME1; N-
PD009; PCNA; PR; PREP; PTTG1; RPLPO; Src; STK15; STMY3; SURV; TFRC; TOP2A; and TS in a breast cancer
cell obtained from the patient, normalized against the expression levels of all RNA transcripts or their expression
products in said breast cancer cell, or of a reference set of RNA transcripts or their products;

(b) subjecting the data obtained in step (a) to statistical analysis; and;

(c) determining whether the likelihood of said long-term survival has increased or decreased.

[0029] In a still further aspect, the disclosure concerns a method of preparing a personalized genomics profile for a
patient, comprising the steps of

(a) subjecting RNA extracted from a breast tissue obtained from the patient to gene expression analysis;

(b) determining the expression level in the tissue of one or more genes selected from the breast cancer gene set
listed in any one of Tables 1 and 2, wherein the expression level is normalized against a control gene or genes and
optionally is compared to the amount found in a breast cancer reference tissue set; and

(c) creating a report summarizing the data obtained by said gene expression analysis. The breast tissue may
comprise breast cancer cells.

[0030] In another embodiment, the breast tissue is obtained from a fixed, paraffin-embedded biopsy sample, in which
the RNA may be fragmented.

[0031] The report may include prediction of the likelihood of long term survival of the patient and/or a recommendation
for a treatment modality of said patient.

[0032] In a further aspect, the disclosure concerns a method for measuring levels of mRNA products of genes listed
in Tables 1 and 2 by real time polymerase chain reaction (RT-PCR), by using an amplicon listed in Table and a primer-
probe set listed in Tables 4A - 4D.

[0033] In a still further aspect, the disclosure concerns a PCR primer-probe set listed in Tables 4A - 4D, and a PCR
amplicon listed in Table 3.

A. Definitions

[0034] Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly
understood by one of ordinary skill in the art to which this invention belongs. Singleton et al., Dictionary of Microbiology
and Molecular Biology 2nd ed., J. Wiley & Sons (New York, NY 1994), and March, Advanced Organic Chemistry Reac-
tions, Mechanisms and Structure 4th ed., John Wiley & Sons (New York, NY 1992), provide one skilled in the art with
a general guide to many of the terms used in the present application.

[0035] One skilled in the art will recognize many methods and materials similar or equivalent to those described herein,
which could be used in the practice of the present invention. Indeed, the present invention is in no way limited to the
methods and materials described. For purposes of the present invention, the following terms are defined below.

[0036] The term "microarray" refers to an ordered arrangement of hybridizable array elements, preferably polynucle-
otide probes, on a substrate.

[0037] The term "polynucleotide," when used in singular or plural, generally refers to any polyribonucleotide or polyde-
oxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. Thus, for instance, polynucleotides
as defined herein include, without limitation, single- and double-stranded DNA, DNA including single- and double-stranded
regions, single- and double-stranded RNA, and RNA including single- and double-stranded regions, hybrid molecules
comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or include single- and double-
stranded regions. In addition, the term "polynucleotide" as used herein refers to triple-stranded regions comprising RNA
or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules.
The regions may include all of one or more of the molecules, but more typically involve only a region of some of the
molecules. One of the molecules of a triple-helical region often is an oligonucleotide. The term "polynucleotide" specifically
includes cDNAs. The term includes DNAs (including cDNAs) and RNAs that contain one or more modified bases. Thus,
DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotides" as that term is intended
herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritiated bases,
are included within the term "polynucleotides" as defined herein. In general, the term "polynucleotide" embraces all
chemically, enzymatically and/or metabolically modified forms of unmodified polynucleotides, as well as the chemical
forms of DNA and RNA characteristic of viruses and cells, including simple and complex cells.

[0038] The term "oligonucleotide" refers to a relatively short polynucleotide, including, without limitation, single-stranded
deoxyribonucleotides, single- or double-stranded ribonucleotides, RNA:DNA hybrids and double-stranded DNAs. Oli-
gonucleotides, such as single-stranded DNA probe oligonucleotides, are often synthesized by chemical methods, for
example using automated oligonucleotide synthesizers that are commercially available. However, oligonucleotides can

be made by a variety of other methods, including *in vitro* recombinant DNA-mediated techniques and by expression of DNAs in cells and organisms.

[0039] The terms "differentially expressed gene," "differential gene expression" and their synonyms, which are used interchangeably, refer to a gene whose expression is activated to a higher or lower level in a subject suffering from a disease, specifically cancer, such as breast cancer, relative to its expression in a normal or control subject. The terms also include genes whose expression is activated to a higher or lower level at different stages of the same disease. It is also understood that a differentially expressed gene may be either activated or inhibited at the nucleic acid level or protein level, or may be subject to alternative splicing to result in a different polypeptide product. Such differences may be evidenced by a change in mRNA levels, surface expression, secretion or other partitioning of a polypeptide, for example. Differential gene expression may include a comparison of expression between two or more genes or their gene products, or a comparison of the ratios of the expression between two or more genes or their gene products, or even a comparison of two differently processed products of the same gene, which differ between normal subjects and subjects suffering from a disease, specifically cancer, or between various stages of the same disease. Differential expression includes both quantitative, as well as qualitative, differences in the temporal or cellular expression pattern in a gene or its expression products among, for example, normal and diseased cells, or among cells which have undergone different disease events or disease stages. For the purpose of this invention, "differential gene expression" is considered to be present when there is at least an about two-fold, preferably at least about four-fold, more preferably at least about six-fold, most preferably at least about ten-fold difference between the expression of a given gene in normal and diseased subjects, or in various stages of disease development in a diseased subject.

[0040] The term "over-expression" with regard to an RNA transcript is used to refer to the level of the transcript determined by normalization to the level of reference mRNAs, which might be all measured transcripts in the specimen or a particular reference set of mRNAs.

[0041] The phrase "gene amplification" refers to a process by which multiple copies of a gene or gene fragment are formed in a particular cell or cell line. The duplicated region (a stretch of amplified DNA) is often referred to as "amplicon." Usually, the amount of the messenger RNA (mRNA) produced, *i.e.*, the level of gene expression, also increases in the proportion of the number of copies made of the particular gene expressed.

[0042] The term "prognosis" is used herein to refer to the prediction of the likelihood of cancer-attributable death or progression, including recurrence, metastatic spread, and drug resistance, of a neoplastic disease, such as breast cancer. The term "prediction" is used herein to refer to the likelihood that a patient will respond either favorably or unfavorably to a drug or set of drugs, and also the extent of those responses, or that a patient will survive, following surgical removal or the primary tumor and/or chemotherapy for a certain period of time without cancer recurrence. The predictive methods of the present invention can be used clinically to make treatment decisions by choosing the most appropriate treatment modalities for any particular patient. The predictive methods of the present invention are valuable tools in predicting if a patient is likely to respond favorably to a treatment regimen, such as surgical intervention, chemotherapy with a given drug or drug combination, and/or radiation therapy, or whether long-term survival of the patient, following surgery and/or termination of chemotherapy or other treatment modalities is likely.

[0043] The term "long-term" survival is used herein to refer to survival for at least 3 years, more preferably for at least 8 years, most preferably for at least 10 years following surgery or other treatment.

[0044] The term "tumor," as used herein, refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues.

[0045] The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of cancer include, but are not limited to, breast cancer, ovarian cancer, colon cancer, lung cancer, prostate cancer, hepatocellular cancer, gastric cancer, pancreatic cancer, cervical cancer, liver cancer, bladder cancer, cancer of the urinary tract, thyroid cancer, renal cancer, carcinoma, melanoma, and brain cancer.

[0046] The "pathology" of cancer includes all phenomena that compromise the well-being of the patient. This includes, without limitation, abnormal or uncontrollable cell growth, metastasis, interference with the normal functioning of neighboring cells, release of cytokines or other secretory products at abnormal levels, suppression or aggravation of inflammatory or immunological response, neoplasia, premalignancy, malignancy, invasion of surrounding or distant tissues or organs, such as lymph nodes, etc.

[0047] "Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., *Current Protocols in Molecular Biology*, Wiley

Interscience Publishers, (1995).

[0048] "Stringent conditions" or "high stringency conditions", as defined herein, typically: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

[0049] "Moderately stringent conditions" may be identified as described by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc., as necessary to accommodate factors such as probe length and the like.

[0050] In the context of the present invention, reference to "at least one," "at least two," "at least five," etc. of the genes listed in any particular gene set means any one or any and all combinations of the genes listed.

[0051] The term "node negative" cancer, such as "node negative" breast cancer, is used herein to refer to cancer that has not spread to the lymph nodes.

[0052] The terms "splicing" and "RNA splicing" are used interchangeably and refer to RNA processing that removes introns and joins exons to produce mature mRNA with continuous coding sequence that moves into the cytoplasm of an eukaryotic cell.

[0053] In theory, the term "exon" refers to any segment of an interrupted gene that is represented in the mature RNA product (B. Lewin. *Genes IV* Cell Press, Cambridge Mass. 1990). In theory the term "intron" refers to any segment of DNA that is transcribed but removed from within the transcript by splicing together the exons on either side of it. Operationally, exon sequences occur in the mRNA sequence of a gene as defined by Ref. SEQ ID numbers. Operationally, intron sequences are the intervening sequences within the genomic DNA of a gene, bracketed by exon sequences and having GT and AG splice consensus sequences at their 5' and 3' boundaries.

B. Detailed Description

[0054] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, and biochemistry, which are within the skill of the art. Such techniques are explained fully in the literature, such as, "Molecular Cloning: A Laboratory Manual", 2nd edition (Sambrook et al., 1989); "Oligonucleotide Synthesis" (M.J. Gait, ed., 1984); "Animal Cell Culture" (R.I. Freshney, ed., 1987); "Methods in Enzymology" (Academic Press, Inc.); "Handbook of Experimental Immunology", 4th edition (D.M. Weir & C.C. Blackwell, eds., Blackwell Science Inc., 1987); "Gene Transfer Vectors for Mammalian Cells" (J.M. Miller & M.P. Calos, eds., 1987); "Current Protocols in Molecular Biology" (F.M. Ausubel et al., eds., 1987); and "PCR: The Polymerase Chain Reaction", (Mullis et al., eds., 1994).

1. *Gene Expression Profiling*

[0055] Methods of gene expression profiling include methods based on hybridization analysis of polynucleotides, methods based on sequencing of polynucleotides, and proteomics-based methods. The most commonly used methods known in the art for the quantification of mRNA expression in a sample include northern blotting and in situ hybridization (Parker & Barnes, *Methods in Molecular Biology* 106:247-283 (1999)); RNAse protection assays (Hod, *Biotechniques* 13:852-854 (1992)); and PCR-based methods, such as reverse transcription polymerase chain reaction (RT-PCR) (Weis et al., *Trends in Genetics* 8:263-264 (1992)). Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. Representative methods for sequencing-based gene expression analysis include Serial Analysis of Gene Expression (SAGE), and gene expression analysis by massively parallel signature sequencing (MPSS).

2. *PCR-based Gene Expression Profiling Methods*

a *Reverse Transcriptase PCR (RT-PCR)*

[0056] Of the techniques listed above, the most sensitive and most flexible quantitative method is RT-PCR, which can be used to compare mRNA levels in different sample populations, in normal and tumor tissues, with or without drug treatment, to characterize patterns of gene expression, to discriminate between closely related mRNAs, and to analyze RNA structure.

[0057] The first step is the isolation of mRNA from a target sample. The starting material is typically total RNA isolated from human tumors or tumor cell lines, and corresponding normal tissues or cell lines, respectively. Thus RNA can be isolated from a variety of primary tumors, including breast, lung, colon, prostate, brain, liver, kidney, pancreas, spleen, thymus, testis, ovary, uterus, etc., tumor, or tumor cell lines, with pooled DNA from healthy donors. If the source of mRNA is a primary tumor, mRNA can be extracted, for example, from frozen or archived paraffin-embedded and fixed (e.g. formalin-fixed) tissue samples.

[0058] General methods for mRNA extraction are well known in the art and are disclosed in standard textbooks of molecular biology, including Ausubel et al., *Current Protocols of Molecular Biology*, John Wiley and Sons (1997). Methods for RNA extraction from paraffin embedded tissues are disclosed, for example, in Rupp and Locker, *Lab Invest.* 56:A67 (1987), and De Andrés et al., *BioTechniques* 18:42044 (1995). In particular, RNA isolation can be performed using purification kit, buffer set and protease from commercial manufacturers, such as Qiagen, according to the manufacturer's instructions. For example, total RNA from cells in culture can be isolated using Qiagen RNeasy mini-columns. Other commercially available RNA isolation kits include MasterPure™ Complete DNA and RNA Purification Kit (EPICENTRE®, Madison, WI), and Paraffin Block RNA Isolation Kit (Ambion, Inc.). Total RNA from tissue samples can be isolated using RNA Stat-60 (Tel-Test). RNA prepared from tumor can be isolated, for example, by cesium chloride density gradient centrifugation.

[0059] As RNA cannot serve as a template for PCR, the first step in gene expression profiling by RT-PCR is the reverse transcription of the RNA template into cDNA, followed by its exponential amplification in a PCR reaction. The two most commonly used reverse transcriptases are avian myeloblastosis virus reverse transcriptase (AMV-RT) and Moloney murine leukemia virus reverse transcriptase (MMLV-RT). The reverse transcription step is typically primed using specific primers, random hexamers, or oligo-dT primers, depending on the circumstances and the goal of expression profiling. For example, extracted RNA can be reverse-transcribed using a GeneAmp RNA PCR kit (Perkin Elmer, CA, USA), following the manufacturer's instructions. The derived cDNA can then be used as a template in the subsequent PCR reaction.

[0060] Although the PCR step can use a variety of thermostable DNA-dependent DNA polymerases, it typically employs the Taq DNA polymerase, which has a 5'-3' nuclease activity but lacks a 3'-5' proofreading endonuclease activity. Thus, TaqMan® PCR typically utilizes the 5'-nuclease activity of Taq or Tth polymerase to hydrolyze a hybridization probe bound to its target amplicon, but any enzyme with equivalent 5' nuclease activity can be used. Two oligonucleotide primers are used to generate an amplicon typical of a PCR reaction. A third oligonucleotide, or probe, is designed to detect nucleotide sequence located between the two PCR primers. The probe is non-extendible by Taq DNA polymerase enzyme, and is labeled with a reporter fluorescent dye and a quencher fluorescent dye. Any laser-induced emission from the reporter dye is quenched by the quenching dye when the two dyes are located close together as they are on the probe. During the amplification reaction, the Taq DNA polymerase enzyme cleaves the probe in a template-dependent manner. The resultant probe fragments disassociate in solution, and signal from the released reporter dye is free from the quenching effect of the second fluorophore. One molecule of reporter dye is liberated for each new molecule synthesized, and detection of the unquenched reporter dye provides the basis for quantitative interpretation of the data.

[0061] TaqMan® RT-PCR can be performed using commercially available equipment, such as, for example, ABI PRISM 7700™ Sequence Detection System™ (Perkin-Elmer-Applied Biosystems, Foster City, CA, USA), or Lightcycler (Roche Molecular Biochemicals, Mannheim, Germany). In a preferred embodiment, the 5' nuclease procedure is run on a real-time quantitative PCR device such as the ABI PRISM 7700™ Sequence Detection System™. The system consists of a thermocycler, laser, charge-coupled device (CCD), camera and computer. The system amplifies samples in a 96-well format on a thermocycler. During amplification, laser-induced fluorescent signal is collected in real-time through fiber optics cables for all 96 wells, and detected at the CCD. The system includes software for running the instrument and for analyzing the data.

[0062] 5'-Nuclease assay data are initially expressed as Ct, or the threshold cycle. As discussed above, fluorescence values are recorded during every cycle and represent the amount of product amplified to that point in the amplification reaction. The point when the fluorescent signal is first recorded as statistically significant is the threshold cycle (C_t).

[0063] To minimize errors and the effect of sample-to-sample variation, RT-PCR is usually performed using an internal standard. The ideal internal standard is expressed at a constant level among different tissues, and is unaffected by the experimental treatment. RNAs most frequently used to normalize patterns of gene expression are mRNAs for the house-

keeping genes glyceraldehyde-3-phosphate-dehydrogenase (GAPDH) and β -actin.

[0064] A more recent variation of the RT-PCR technique is the real time quantitative PCR, which measures PCR product accumulation through a dual-labeled fluorogenic probe (i.e., TaqMan® probe). Real time PCR is compatible both with quantitative competitive PCR, where internal competitor for each target sequence is used for normalization, and with quantitative comparative PCR using a normalization gene contained within the sample, or a housekeeping gene for RT-PCR. For further details see, e.g. Held et al., *Genome Research* 6:986-994 (1996).

[0065] The steps of a representative protocol for profiling gene expression using fixed, paraffin-embedded tissues as the RNA source, including mRNA isolation, purification, primer extension and amplification are given in various published journal articles {for example: T.E. Godfrey et al. *J. Molec. Diagnostics* 2: 84-91 [2000]; K. Specht et al., *Am. J. Pathol.* 158: 419-29 [2001]}. Briefly, a representative process starts with cutting about 10 μ m thick sections of paraffin-embedded tumor tissue samples. The RNA is then extracted, and protein and DNA are removed. After analysis of the RNA concentration, RNA repair and/or amplification steps may be included, if necessary, and RNA is reverse transcribed using gene specific promoters followed by RT-PCR.

b. MassARRAY System

[0066] In the MassARRAY-based gene expression profiling method, developed by Sequenom, Inc. (San Diego, CA) following the isolation of RNA and reverse transcription, the obtained cDNA is spiked with a synthetic DNA molecule (competitor), which matches the targeted cDNA region in all positions, except a single base, and serves as an internal standard. The cDNA/competitor mixture is PCR amplified and is subjected to a post-PCR shrimp alkaline phosphatase (SAP) enzyme treatment, which results in the dephosphorylation of the remaining nucleotides. After inactivation of the alkaline phosphatase, the PCR products from the competitor and cDNA are subjected to primer extension, which generates distinct mass signals for the competitor- and cDNA-derivatives PCR products. After purification, these products are dispensed on a chip array, which is pre-loaded with components needed for analysis with matrix-assisted laser desorption ionization time-of-flight mass spectrometry (MALDI-TOF MS) analysis. The cDNA present in the reaction is then quantified by analyzing the ratios of the peak areas in the mass spectrum generated. For further details see, e.g. Ding and Cantor, *Proc. Natl. Acad. Sci. USA* 100:3059-3064 (2003).

c. Other PCR-based Methods

[0067] Further PCR-based techniques include, for example, differential display (Liang and Pardee, *Science* 257:967-971 (1992)); amplified fragment length polymorphism (AFLP) (Kawamoto et al., *Genome Res.* 12:1305-1312 (1999)); BeadArray™ technology (Illumina, San Diego, CA; Oliphant et al., *Discovery of Markers for Disease* (Supplement to *Biotechniques*), June 2002; Ferguson et al., *Analytical Chemistry* 72:5618 (2000)); BeadsArray for Detection of Gene Expression (BADGE), using the commercially available Luminex100 LabMAP system and multiple color-coded microspheres (Luminex Corp., Austin, TX) in a rapid assay for gene expression (Yang et al., *Genome Res.* 11:1888-1898 (2001)); and high coverage expression profiling (HiCEP) analysis (Fukumura et al., *Nucl. Acids. Res.* 31(16) e94 (2003)).

3. Microarrays

[0068] Differential gene expression can also be identified, or confirmed using the microarray technique. Thus, the expression profile of breast cancer-associated genes can be measured in either fresh or paraffin-embedded tumor tissue, using microarray technology. In this method, polynucleotide sequences of interest (including cDNAs and oligonucleotides) are plated, or arrayed, on a microchip substrate. The arrayed sequences are then hybridized with specific DNA probes from cells or tissues of interest. Just as in the RT-PCR method, the source of mRNA typically is total RNA isolated from human tumors or tumor cell lines, and corresponding normal tissues or cell lines. Thus RNA can be isolated from a variety of primary tumors or tumor cell lines. If the source of mRNA is a primary tumor, mRNA can be extracted, for example, from frozen or archived paraffin-embedded and fixed (e.g. formalin-fixed) tissue samples, which are routinely prepared and preserved in everyday clinical practice.

[0069] In a specific embodiment of the microarray technique, PCR amplified inserts of cDNA clones are applied to a substrate in a dense array. Preferably at least 10,000 nucleotide sequences are applied to the substrate. The microarrayed genes, immobilized on the microchip at 10,000 elements each, are suitable for hybridization under stringent conditions. Fluorescently labeled cDNA probes may be generated through incorporation of fluorescent nucleotides by reverse transcription of RNA extracted from tissues of interest. Labeled cDNA probes applied to the chip hybridize with specificity to each spot of DNA on the array. After stringent washing to remove non-specifically bound probes, the chip is scanned by confocal laser microscopy or by another detection method, such as a CCD camera. Quantitation of hybridization of each arrayed element allows for assessment of corresponding mRNA abundance. With dual color fluorescence, separately labeled cDNA probes generated from two sources of RNA are hybridized pairwise to the array. The relative

abundance of the transcripts from the two sources corresponding to each specified gene is thus determined simultaneously. The miniaturized scale of the hybridization affords a convenient and rapid evaluation of the expression pattern for large numbers of genes. Such methods have been shown to have the sensitivity required to detect rare transcripts, which are expressed at a few copies per cell, and to reproducibly detect at least approximately two-fold differences in the expression levels (Schena et al., Proc. Natl. Acad. Sci. USA 93(2): 106-149 (1996)). Microarray analysis can be performed by commercially available equipment, following manufacturer's protocols, such as by using the Affymetrix GenChip technology, or Incyte's microarray technology.

[0070] The development of microarray methods for large-scale analysis of gene expression makes it possible to search systematically for molecular markers of cancer classification and outcome prediction in a variety of tumor types.

4. Serial Analysis of Gene Expression (SAGE)

[0071] Serial analysis of gene expression (SAGE) is a method that allows the simultaneous and quantitative analysis of a large number of gene transcripts, without the need of providing an individual hybridization probe for each transcript. First, a short sequence tag (about 10-14 bp) is generated that contains sufficient information to uniquely identify a transcript, provided that the tag is obtained from a unique position within each transcript. Then, many transcripts are linked together to form long serial molecules, that can be sequenced, revealing the identity of the multiple tags simultaneously. The expression pattern of any population of transcripts can be quantitatively evaluated by determining the abundance of individual tags, and identifying the gene corresponding to each tag. For more details see, e.g. Velculescu et al., Science 270:484-487 (1995); and Velculescu et al., Cell 88:243-51 (1997).

5. Gene Expression Analysis by Massively Parallel Signature Sequencing (MPSS)

[0072] This method, described by Brenner et al., Nature Biotechnology 18:630-634 (2000), is a sequencing approach that combines non-gel-based signature sequencing with *in vitro* cloning of millions of templates on separate 5 μ m diameter microbeads. First, a microbead library of DNA templates is constructed by *in vitro* cloning. This is followed by the assembly of a planar array of the template-containing microbeads in a flow cell at a high density (typically greater than 3×10^6 microbeads/cm²). The free ends of the cloned templates on each microbead are analyzed simultaneously, using a fluorescence-based signature sequencing method that does not require DNA fragment separation. This method has been shown to simultaneously and accurately provide, in a single operation, hundreds of thousands of gene signature sequences from a yeast cDNA library.

6. Immunohistochemistry

[0073] Immunohistochemistry methods are also suitable for detecting the expression levels of the prognostic markers of the present invention. Thus, antibodies or antisera, preferably polyclonal antisera, and most preferably monoclonal antibodies specific for each marker are used to detect expression. The antibodies can be detected by direct labeling of the antibodies themselves, for example, with radioactive labels, fluorescent labels, hapten labels such as, biotin, or an enzyme such as horse radish peroxidase or alkaline phosphatase. Alternatively, unlabeled primary antibody is used in conjunction with a labeled secondary antibody, comprising antisera, polyclonal antisera or a monoclonal antibody specific for the primary antibody. Immunohistochemistry protocols and kits are well known in the art and are commercially available.

7. Proteomics

[0074] The term "proteome" is defined as the totality of the proteins present in a sample (e.g. tissue, organism, or cell culture) at a certain point of time. Proteomics includes, among other things, study of the global changes of protein expression in a sample (also referred to as "expression proteomics"). Proteomics typically includes the following steps: (1) separation of individual proteins in a sample by 2-D gel electrophoresis (2-D PAGE); (2) identification of the individual proteins recovered from the gel, e.g. by mass spectrometry or N-terminal sequencing, and (3) analysis of the data using bioinformatics. Proteomics methods are valuable supplements to other methods of gene expression profiling, and can be used, alone or in combination with other methods, to detect the products of the prognostic markers of the present invention.

8. General Description of the mRNA Isolation, Purification and Amplification

[0075] The steps of a representative protocol for profiling gene expression using fixed, paraffin-embedded tissues as the RNA source, including mRNA isolation, purification, primer extension and amplification are provided in various

published journal articles (for example: T.E. Godfrey et al., J. Molec. Diagnostics 2: 84-91 [2000]; K. Specht et al., Am. J. Pathol. 158: 419-29 [2001]). Briefly, a representative process starts with cutting about 10 μ m thick sections of paraffin-embedded tumor tissue samples. The RNA is then extracted, and protein and DNA are removed. After analysis of the RNA concentration, RNA repair and/or amplification steps may be included, if necessary, and RNA is reverse transcribed using gene specific promoters followed by RT-PCR. Finally, the data are analyzed to identify the best treatment option(s) available to the patient on the basis of the characteristic gene expression pattern identified in the tumor sample examined, dependent on the predicted likelihood of cancer recurrence.

9. Breast Cancer Gene Set, Assayed Gene Subsequences, and Clinical Application of Gene Expression Data

[0076] An important aspect of the present invention is to use the measured expression of certain genes by breast cancer tissue to provide prognostic information. For this purpose it is necessary to correct for (normalize away) both differences in the amount of RNA assayed and variability in the quality of the RNA used. Therefore, the assay typically measures and incorporates the expression of certain normalizing genes, including well known housekeeping genes, such as GAPDH and Cyp1. Alternatively, normalization can be based on the mean or median signal (Ct) of all of the assayed genes or a large subset thereof (global normalization approach). On a gene-by-gene basis, measured normalized amount of a patient tumor mRNA is compared to the amount found in a breast cancer tissue reference set. The number (N) of breast cancer tissues in this reference set should be sufficiently high to ensure that different reference sets (as a whole) behave essentially the same way. If this condition is met, the identity of the individual breast cancer tissues present in a particular set will have no significant impact on the relative amounts of the genes assayed. Usually, the breast cancer tissue reference set consists of at least about 30, preferably at least about 40 different FPE breast cancer tissue specimens. Unless noted otherwise, normalized expression levels for each mRNA/tested tumor/patient will be expressed as a percentage of the expression level measured in the reference set. More specifically, the reference set of a sufficiently high number (e.g. 40) of tumors yields a distribution of normalized levels of each mRNA species. The level measured in a particular tumor sample to be analyzed falls at some percentile within this range, which can be determined by methods well known in the art. Below, unless noted otherwise, reference to expression levels of a gene assume normalized expression relative to the reference set although this is not always explicitly stated.

10. Design of Intron-Based PCR Primers and Probes

[0077] According to one aspect of the present invention, PCR primers and probes are designed based upon intron sequences present in the gene to be amplified. Accordingly, the first step in the primer/probe design is the delineation of intron sequences within the genes. This can be done by publicly available software, such as the DNA BLAT software developed by Kent, W.J., Genome Res. 12(4):656-64 (2002), or by the BLAST software including its variations. Subsequent steps follow well established methods of PCR primer and probe design.

[0078] In order to avoid non-specific signals, it is important to mask repetitive sequences within the introns when designing the primers and probes. This can be easily accomplished by using the Repeat Masker program available on-line through the Baylor College of Medicine, which screens DNA sequences against a library of repetitive elements and returns a query sequence in which the repetitive elements are masked. The masked intron sequences can then be used to design primer and probe sequences using any commercially or otherwise publicly available primer/probe design packages, such as Primer Express (Applied Biosystems); MGB assay-by-design (Applied Biosystems); Primer3 (Steve Rozen and Helen J. Skaletsky (2000) Primer3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S (eds) Bioinformatics Methods and Protocols: Methods in Molecular Biology. Humana Press, Totowa, NJ, pp 365-386).

[0079] The most important factors considered in PCR primer design include primer length, melting temperature (T_m), and G/C content, specificity, complementary primer sequences, and 3'-end sequence. In general, optimal PCR primers are generally 17-30 bases in length, and contain about 20-80%, such as, for example, about 50-60% G+C bases. T_m's between 50 and 80 °C, e.g. about 50 to 70 °C are typically preferred.

[0080] For further guidelines for PCR primer and probe design see, e.g. Dieffenbach, C.W. et al., "General Concepts for PCR Primer Design" in: PCR Primer, A Laboratory Manual, Cold Spring Harbor Laboratory Press, New York, 1995, pp. 133-155; Innis and Gelfand, "Optimization of PCRs" in: PCR Protocols, A Guide to Methods and Applications, CRC Press, London, 1994, pp. 5-11; and Plasterer, T.N. Primerselect: Primer and probe design. Methods Mol. Biol. 70:520-527 (1997).

[0081] Further details of the invention will be described in the following non-limiting Example.

Example

A Phase II Study of Gene Expression in 242 Malignant Breast Tumors

[0082] A gene expression study was designed and conducted with the primary goal to molecularly characterize gene expression in paraffin-embedded, fixed tissue samples of invasive breast ductal carcinoma, and to explore the correlation between such molecular profiles and disease-free survival.

Study design

[0083] Molecular assays were performed on paraffin-embedded, formalin-fixed primary breast tumor tissues obtained from 252 individual patients diagnosed with invasive breast cancer. All patients were lymph node-negative, ER-positive, and treated with Tamoxifen. Mean age was 52 years, and mean clinical tumor size was 2 cm. Median follow-up was 10.9 years. As of January 1, 2003, 41 patients had local or distant disease recurrence or breast cancer death. Patients were included in the study only if histopathologic assessment, performed as described in the Materials and Methods section, indicated adequate amounts of tumor tissue and homogeneous pathology.

Materials and Methods

[0084] Each representative tumor block was characterized by standard histopathology for diagnosis, semi-quantitative assessment of amount of tumor, and tumor grade. When tumor area was less than 70% of the section, the tumor area was grossly dissected and tissue was taken from 6 (10 micron) sections. Otherwise, a total of 3 sections (also 10 microns in thickness each) were prepared. Sections were placed in two Costar Brand Microcentrifuge Tubes (Polypropylene, 1.7 mL tubes, clear). If more than one tumor block was obtained as part of the surgical procedure, the block most representative of the pathology was used for analysis.

Gene Expression Analysis

[0085] mRNA was extracted and purified from fixed, paraffin-embedded tissue samples, and prepared for gene expression analysis as described in chapter 6 above.

[0086] Molecular assays of quantitative gene expression were performed by RT-PCR, using the ABI PRISM 7900™ Sequence Detection System™ (Perkin-Elmer-Applied Biosystems, Foster City, CA, USA). ABI PRISM 7900™ consists of a thermocycler, laser, charge-coupled device (CCD), camera and computer. The system amplifies samples in a 384-well format on a thermocycler. During amplification, laser-induced fluorescent signal is collected in real-time through fiber optics cables for all 384 wells, and detected at the CCD. The system includes software for running the instrument and for analyzing the data.

Analysis and Results

[0087] Tumor tissue was analyzed for 187 cancer-related genes and 5 reference genes. Adequate RT-PCR profiles were obtained from 242 of the 252 patients. The threshold cycle (CT) values for each patient were normalized based on the median of the 7 reference genes for that particular patient. Clinical outcome data were available for all patients from a review of registry data and selected patient charts. Outcomes were classified as:

Event: Alive with local, regional or distant breast cancer recurrence or death due to breast cancer.

No Event: Alive without local, regional or distant breast cancer recurrence or alive with contralateral breast cancer recurrence or alive with non-breast second primary cancer or died prior to breast cancer recurrence.

[0088] Analysis was performed by:

- A. determination of the relationship between normalized gene expression and the binary outcomes of 0 or 1;
- B. Analysis of the relationship between normalized gene expression and the time to outcome (0 or 1 as defined above) where patients who were alive without breast cancer recurrence or who died due to a cause other than breast cancer were censored. This approach was used to evaluate the prognostic impact of individual genes and also sets of multiple genes.

Analysis of patients with invasive breast carcinoma by binary approach

[0089] In the first (binary) approach, analysis was performed on all 242 patients with invasive breast carcinoma. A t test was performed on the groups of patients classified as either no recurrence and no breast cancer related death at 10 years, versus recurrence, or breast cancer-related death at 10 years, and the p-values for the differences between the groups for each gene were calculated.

[0090] Table 1 lists the 33 genes for which the p-value for the differences between the groups was <0.05. The first column of mean expression values pertains to patients who had a metastatic recurrence of nor died from breast cancer. The second column of mean expression values pertains to patients who neither had a metastatic recurrence of nor died from breast cancer.

Table 1

Gene	Mean group A Event	Mean group B No event	T statistic	P value
MMP9	-3.15	-4.27	3.75	0.00
GSTM1	-5.02	-4.03	-3.56	0.00
MELK	-3.89	-4.66	3.34	0.00
PR	-4.56	-3.18	-3.27	0.00
DKFZp586M07	-3.83	-2.94	-3.09	0.00
GSTM3	-2.56	-1.69	-3.06	0.00
MCM2	-3.51	-4.08	3.03	0.00
CDC20	-3.01	-3.75	3.01	0.00
CCNB1	-4.48	-5.17	3.02	0.00
STMY3	-0.58	-1.20	2.95	0.00
GRB7	-1.93	-3.01	2.98	0.00
MYBL2	-3.91	-4.78	2.91	0.01
CEGP1	-3.00	-1.85	-2.89	0.01
SURV	-4.23	-5.06	2.88	0.01
LMNB1	-2.40	-2.91	2.81	0.01
CTSL2	-5.74	-6.39	2.83	0.01
PTTG1	-3.49	-4.14	2.72	0.01
BAG1	-1.76	-1.30	-2.58	0.01
KNSL2	-3.35	-4.06	2.60	0.01
CIAP1	-4.44	-4.02	-2.58	0.01
PREP	-3.34	-3.74	2.56	0.01
NEK2	-5.25	-5.80	2.53	0.01
EpCAM	-1.95	-2.31	2.50	0.01
PCNA	-2.79	-3.13	2.42	0.02
C20_orf1	-2.43	-3.09	2.39	0.02
ITGA7	-4.53	-3.87	-2.37	0.02
ID1	-2.58	-2.17	-2.30	0.02
B_Catenin	-1.32	-1.08	-2.28	0.03
EstR1	-0.78	-0.12	-2.28	0.03
CDH1	-2.76	-3.27	2.20	0.03
TS	-2.86	-3.29	2.18	0.03
HER2	0.53	-0.22	2.18	0.03
cMYC	-3.22	-2.85	-2.16	0.04

[0091] In the foregoing Table 1, negative t-values indicate higher expression, associated with better outcomes, and, inversely, higher (positive) t-values indicate higher expression associated with worse outcomes. Thus, for example, elevated expression of the CCNB1 gene (t-value = 3.02; CT mean alive < CT mean deceased) indicates a reduced likelihood of disease free survival. Similarly, elevated expression of the GSTM1 gene (t-value = -3.56; CT mean alive > CT mean deceased) indicates an increased likelihood of disease free survival.

[0092] Thus, based on the data set forth in Table 1, the expression of any of the following genes in breast cancer

indicates a reduced likelihood of survival without cancer recurrence: C20_orf1; CCNB1; CDC20; CDH1; CTSL2; EpCAM; GRB7; HER2; KNSL2; LMNB1; MCM2; MMP9; MYBL2; NEK2; PCNA; PREP; PTTG1; STMY3; SURV; TS; MELK.

[0093] Based on the data set forth in Table 1, the expression of any of the following genes in breast cancer indicates a better prognosis for survival without cancer recurrence: BAG1; BCatenin; CEGP1; CIAP1; cMYC; DKFZp586M07; EstR1; GSTM1; GSTM3; ID1; ITGA7; PR.

Analysis of multiple genes and indicators of outcome

[0094] Two approaches were taken in order to determine whether using multiple genes would provide better discrimination between outcomes. First, a discrimination analysis was performed using a forward stepwise approach. Models were generated that classified outcome with greater discrimination than was obtained with any single gene alone.; According to a second approach (time-to-event approach), for each gene a Cox Proportional Hazards model (see, e.g. Cox, D. R., and Oakes, D. (1984), Analysis of Survival Data, Chapman and Hall, London, New York) was defined with time to recurrence or death as the dependent variable, and the expression level of the gene as the independent variable. The genes that have a p-value < 0.05 in the Cox model were identified. For each gene, the Cox model provides the relative risk (RR) of recurrence or death for a unit change in the expression of the gene. One can choose to partition the patients into subgroups at any threshold value of the measured expression (on the CT scale), where all patients with expression values above the threshold have higher risk, and all patients with expression values below the threshold have lower risk, or vice versa, depending on whether the gene is an indicator of bad (RR>1.01) or good (RR<1.01) prognosis. Thus, any threshold value will define subgroups of patients with respectively increased or decreased risk. The results are summarized in Table 2, which lists the 42 genes for which the p-value for the differences between the groups was <0.05.

Table 2

Gene	Relative Risk	p-value
GRB7	1.52	0.000011
SURV	1.57	0.000090
PR	0.74	0.000129
LMNB1	1.92	0.000227
MYBL2	1.46	0.000264
HER2	1.46	0.000505
GSTM1	0.68	0.000543
MELK	1.59	0.000684
C20_orf1	1.59	0.000735
PTTG1	1.63	0.001135
BUB1	1.58	0.001425
CDC20	1.54	0.001443
CCNB1	1.60	0.001975
STMY3	1.47	0.002337
KNSL2	1.48	0.002910
CTSL2	1.43	0.003877
MCM2	1.59	0.005203
NEK2	1.48	0.006533
DR5	0.62	0.006660
Ki_67	1.46	0.008188
CCNE2	1.38	0.009505
TOP2A	1.38	0.009551
PCNA	1.67	0.010237
PREP	1.69	0.012308
FOXM1	1.52	0.012837
NME1	1.46	0.013622
CEGP1	0.84	0.013754
BAG1	0.68	0.015422
STK15	1.46	0.017013

(continued)

	Gene	Relative Risk	p-value
5	HNRPAB	1.96	0.017942
	EstR1	0.80	0.018877
	MMP9	1.19	0.019591
	DKFZp586M07	0.79	0.020073
	TS	1.44	0.025186
10	Src	1.70	0.037398
	BIN1	0.75	0.038979
	NPD009	0.80	0.039020
	RPLPO	0.52	0.041575
	GSTM3	0.84	0.041848
15	MMP12	1.27	0.042074
	TFRC	1.57	0.046145
	IGF1R	0.78	0.046745

20 **[0095]** Based on the data set forth in Table 2, the expression of any of the following genes in breast cancer indicates a reduced likelihood of survival without cancer recurrence: GRB7; SURV; LMNB1; MYBL2; HER2; MELK; C20_orf1; PTTG1; BUB1; CDC20; CCNB1; STMY3; KNSL2; CTSL2; MCM2; NEK2; Ki_67; CCNE2; TOP2A-4; PCNA; PREP; FOXM1; NME1; STK15; HNRPAB; MMP9; TS; Src; MMP12; TFRC.

25 **[0096]** Based on the data set forth in Table 2, the expression of any of the following genes in breast cancer indicates a better prognosis for survival without cancer recurrence: PR; GSTM1; DR5; CEGP1; BAG1; EstR1; DKFZp586M07; BIN1; NPD009; RPLPO; GSTM3; IGF1R.

30 **[0097]** The binary and time-to-event analyses, with few exceptions, identified the same genes as prognostic markers. For example, comparison of Tables 1 and 2 shows that 10 genes were represented in the top 15 genes in both lists. Furthermore, when both analyses identified the same gene at [p<0.10], which happened for 26 genes, they were always concordant with respect to the direction (positive or negative sign) of the correlation with survival/recurrence. Overall, these results strengthen the conclusion that the identified markers have significant prognostic value.

Multivariate gene analysis of 242 patients with invasive breast carcinoma

35 **[0098]** For Cox models comprising more than two genes (multivariate models), stepwise entry of each individual gene into the model is performed, where the first gene entered is pre-selected from among those genes having significant univariate p-values, and the gene selected for entry into the model at each subsequent step is the gene that best improves the fit of the model to the data. This analysis can be performed with any total number of genes. In the analysis the results of which are shown below, stepwise entry was performed for up to 10 genes.

40 **[0099]** Multivariate analysis was performed using the following equation::

Multivariate analysis was performed using the following equation::

$$R = \exp[\text{coef}(\text{geneA}) \times \text{Ct}(\text{geneA}) + \text{coef}(\text{geneB}) \times \text{Ct}(\text{geneB}) + \text{coef}(\text{geneC}) \times \text{Ct}(\text{geneC}) + \dots]$$

45 **[0100]** In this equation, coefficients for genes that are predictors of beneficial outcome are positive numbers and coefficients for genes that are predictors of unfavorable outcome are negative numbers. The "Ct" values in the equation are ΔCt s, i.e. reflect the difference between the average normalized Ct value for a population and the normalized Ct measured for the patient in question. The convention used in the present analysis has been that ΔCt s below and above the population average have positive signs and negative signs, respectively (reflecting greater or lesser mRNA abundance). The relative risk (RR) calculated by solving this equation will indicate if the patient has an enhanced or reduced chance of long-term survival without cancer recurrence.

50 **[0101]** A multivariate stepwise analysis, using the Cox Proportional Hazards Model, was performed on the gene expression data obtained for all 242 patients with invasive breast carcinoma. The following ten-gene set has been identified by this analysis as having particularly strong predictive value of patient survival: GRB7; LMNB1; ER; STMY3; KLK10; PR; KRT5; FGFR1; MCM6; SNRPF. In this gene set ER, PR, KRT5 and MCM6 contribute to good prognosis, while GRB7, LMNB1, STMY3, KLK10, FGFR1, and SNRPF contribute to poor prognosis.

[0102] While the present invention has been described with reference to what are considered to be the specific embodiments, it is to be understood that the invention is not limited to such embodiments. To the contrary, the invention is intended to cover various modifications included within the scope of the appended claims.

5

10

15

20

25

30

35

40

45

50

55

TABLE 3

Gene	Accession	Start	Stop	SEQ ID NO.	Sequence
B-Catenin	NM_001904	1549	1629	SEQ ID NO: 1	GGCTGTTGTGGTACTGCTTCGGGCTGTGACAGGGAAGACATCACTGAGGCTGCATCTGTGCTCTCGTCATCTGA
BAG1	NM_004323	673	754	SEQ ID NO: 2	CGTTGTGACACCTTGAATACAAAGATGGTTCCGGGTGTCATGTTAAATTTGGGAAAGAAACAGTCCACAGGAAGAGGTGAAC
BIN1	NM_004305	866	942	SEQ ID NO: 3	CCTGAAAGGGAACAAGAGCCCTTCGCCCTCCAGATGGCTCCCTGCCGCCACCCCGAGATCAAGTCAACCCAG
BUB1	NM_004336	1002	1070	SEQ ID NO: 4	CCGAGGTTAATCCAGCACGATATGGGCCAAGTGTAGGCTCCAGCAGAACTGAGAGGCCCATGTCTT
C20 orf1	NM_012112	2675	2740	SEQ ID NO: 5	TCAGCTGTGAGCTGCGGATACCCGCCGGCAAATGGGACCTGCTCTTAACCTCAAACCTAGGACCCGT
CCNB1	NM_031966	823	907	SEQ ID NO: 6	TTCAAGGTTTGGCAGGAGACCATGTACATGACTGTCTCCATTATTGATCGGTTTCATGCAGAAATAATTGTGTGCCCAAGAGATG
CCNE2	NM_057749	2026	2108	SEQ ID NO: 7	ATGCTGTGGCTCCTTCCTTAAGTGGGCTTTCTTGACATGATAGTTGCTTGGTAATAACCTTTTGTATATCACAAATTTGGGT
CDC20	NM_001255	679	747	SEQ ID NO: 8	TGGATTGGAGTTCTGGGAATGTAAGTGGCGGTGGCACTGGACAAACAGTGTGTACCTGTGAGTSCAAGC
CDH1	NM_004360	2499	2580	SEQ ID NO: 9	TGAGTGTCCCGCGGTATCTTCCCGCCCTGCCAATCCCGATGAAATTTGGAATTTTATTGATGAAAATCTGAAGGGGCTG
CEGP1	NM_020974	563	640	SEQ ID NO: 10	TGACAATCAGACACCTGCAATCACCGCTUGGAAGAGGGCTTGAGCTGCATGAATAAGGATCAAGCTGTAGTCACA
CIAP1	NM_001166	1822	1894	SEQ ID NO: 11	TGCTGTGTGGGAAGCTCAGTAACATGGGAACCAAGGATGATGCTATGTACAGAACACCGGAGGCATTTTC
cMYC	NM_002467	1494	1578	SEQ ID NO: 12	TCCCTCCACTCGGAAGGACTATCCTCTGCCAAGAGGTCAGTTGGACAGTGTACAGTCCCTGAACAGATCAGCAACAACCG
CTSL2	NM_001333	671	738	SEQ ID NO: 13	TGTCCTACTGACGAGCAGAATCTGGTGGACTGTTCCGGTCCCTCAAGGCAATCAGGGCTGCAATGGT
DKFZp586	AL050227	559	633	SEQ ID NO: 14	TCCATTTCTACCTGTTAACCTTCATCATTTTGTGCAGGCCCTGGAAGCAAGAGAGGAAGGGACCGACTGCAT
DR5	NM_003842	1127	1211	SEQ ID NO: 15	CTCTGAGACAGTGTTCGATGACTTTGCAGACTTTGGTGCCCTTTGACTCCTGGGAGCCGCTCATGAGGAAGTTGGCCCTCATGG
EpcAM	NM_002354	435	510	SEQ ID NO: 16	GGGCCCTCCAGAACAAATGATGGGCTTTATCATCCTGACTGGATGAGAGCGGGGCTCTTTAAGGCCAAGCAGTGCA
EstR1	NM_000125	1956	2024	SEQ ID NO: 17	CGTGGTGGCCCTCTATGACCTGCTGCTGGAGATGCTGGAGGCCACCCGCTACATGGGCCCACTAGCC
FGFR1	NM_023109	2685	2759	SEQ ID NO: 18	CACGGGACATTCACACATCGACTACTATAAAAGACAAACCAACGCGCGACTGCCTGTGAAGTGGATGGCACCC
FOXM1	NM_021953	1898	1980	SEQ ID NO: 19	CCACCCCGAGCAAAATCTGTCTCCCGAGAACCCCTGAATCTCTGGAGGCTCACGCCCCCGAGCCAAAATAGGGGACTGGATT
GRB7	NM_005310	1275	1342	SEQ ID NO: 20	CCATCTGCATCCATCTTGTTTGGGCTCCCGCCCTTGAGAAGTGCCTCAGATAATACCGTGGTGGCC
GSTM1	NM_000581	93	179	SEQ ID NO: 21	AAGCTATGAGGAAAGAGTACACGATGGGGACGCTCCTGATTATGACAGAAGCCAGTGGCTGAATGAATAATTCAGCTGGGCG
GSTM3	NM_000849	248	324	SEQ ID NO: 22	CAATGCCATCTTTGGGCTACATGCTCGCAAGCAACACATGTGTGTGGAGACTGAAGAAGAAAGATTCGAGTGGAC

(continued)

Gene	Accession	Start	Stop	SEQ ID NO.	Sequence
HER2	NM_004448	1138	1208	SEQ ID NO: 23	CGGTGTGAGAAAGTGACGAAGCCCTGTGCCGAGTGTGCTATGGTCTGGGCATGGAGCACTTGC GAGAGG
HNRPAB	NM_004499	1086	1170	SEQ ID NO: 24	CAAGGGACGACCAACTGATCGACACATGCTTTGTTGGATGGAGTGAACAATATGTACC AAATTTAACTTGGCAAC
ID1	NM_002165	286	356	SEQ ID NO: 25	AGAACCGCAAGGTGAGCAAGGTGGAGATTCTCCAGCAOGTCATCGACTACATCAGGGACCTTCAG TTGGA
IGF1R	NM_000875	3467	3550	SEQ ID NO: 26	GCATGGTAGCCGAAGATTTCACAGTCAAAAATCGGAGATTTTGGTATGACGGGAGATATCTATGAGA CAGACTATTACGGGAAA
ITGA7	NM_002206	633	712	SEQ ID NO: 27	GATATGATTGGTCGCTGCTTTGTGCTCAGCCAGGACCTGGCCATCCGGGATGATTTGGATGGTGG GGAATGGAAGTTCT
KI-67	NM_002417	42	122	SEQ ID NO: 28	CGGACITTTGGGTGCGACTTGACGAGCGGTGGTTGACAAAGTGGCCTTTCGGGGCCGATCGTCCC AGTGGAAAGATTGTAA
KLK10	NM_002776	966	1044	SEQ ID NO: 29	GCCCAGAGGCTCCATCGTCGCTCTTCTCCCAAGTCGGGTGAACCTCTCCCTTGTCTGCACT GTTCAACGCTCG
KNSL2	BC000712	1266	1343	SEQ ID NO: 30	CCACCTCGCCATGATTTTTCCTTTGACCGGGTATTCACAGGAAAGTGGACAGGATGAAGTGTT GAAGAGATTGC
KRT5	NM_000424	1605	1674	SEQ ID NO: 31	TCAGTGGAGAAGGAGTTGACACAGTCAACATCTCTGTTGTACAAAGCAGTGTTCCTCTGGATATG GCA
LMNB1	NM_005573	1500	1566	SEQ ID NO: 32	TGCAAAAGCTGGTGTACAGGCCAGCCGCCCAACTGACCTCATCTGGAAGAACCAGAACTCGTGGG G
MCM2	NM_004526	2442	2517	SEQ ID NO: 33	GACTTTTGGCCGCTACCTTTTCATTCGCGCGTGACAACAATGAGCTGTTGCTCTTCTCATCTGAAGCA GTTAGTGGC
MCM6	NM_005915	2669	2751	SEQ ID NO: 34	TGATGTCTCTATGTGTACATTCATCAGAGTTTCATACCAACACAGGCTTCAGCACTTCCTTTGGT GTGTTTCTGTCTCCCA
MELK	NM_014791	22	87	SEQ ID NO: 35	AACCCGGCGATCGAAAAGATTCTTAGGAACGCCGTACCCAGCCGCTCTCTCAGGACAGCAGGCCCC
MMP12	NM_002426	816	894	SEQ ID NO: 36	CCAACGTTGCCAAATCTGACAAATTCAGAACCAAGCTCTGTGACCCCAATTTGAGTTTTGATGCT GTCACTACCGT
MMP9	NM_004994	124	191	SEQ ID NO: 37	GAGAACCAATCTCACCGACAGGAGCTGGCAGAGGAATACCTGTACCGCTATGGTTACACTCGGG TG
MYBL2	NM_002466	599	673	SEQ ID NO: 38	GCCGAGATCGCCAAGATGTTGCCAGGGAGGACAGACAATGCTGTGAAGAATCACTGGAACCTCTAC CATCAAAAG
NEK2	NM_002497	102	181	SEQ ID NO: 39	GTGAGGACGCGGACTCTGCGGACTGCGCGCCATGCGTTCCCGGGCTGAGGACTATGAAGTGT TGACACCAATTGGCA
NME1	NM_000269	365	439	SEQ ID NO: 40	CCAACCTGACAGACTCCAAGCCCTGGGACCATCCGTGGAGACTTCTGCATACAAGTTGGCAGGAAC ATTATACAT
NPD009	NM_020686	589	662	SEQ ID NO: 41	GGCTGTGGCTGAGGCTGTAGCATCTCTGTGGAGGTGAGACACTCTGSGAACTGATTTGACCTCG AATGCTCC
PCNA	NM_002592	157	228	SEQ ID NO: 42	GAAGGTGTTGGAGGCACTCAAGGACCTCATCAACGAGGCCCTGCTGGGATATTAGCTCCAGCGGTG TAAACC
PR	NM_000926	1895	1980	SEQ ID NO: 43	GCATCAGGCTGTATTATGGTGTCTTACCTGTGGAGGTGTAAGGTCTTCTTTAAGAGGCGCAATG GAAGGGCAGCAACTACT

5
10
15
20
25
30
35
40
45
50
55

(continued)

Gene	Accession	Start	Stop	SEQ ID NO.	Sequence
PREP	NM_002726	859	965	SEQ ID NO: 44	GGGACGGGTTCACATTCAGACGAATCGCAGTCTCCCAACTATCGCGTGATCAACATTGACTTC TGGATCCTG
PTTG1	NM_004219	48	122	SEQ ID NO: 45	GGCTACTCTGATCTATGTTGATAGGAAATGGAGAACCAAGGACCCCGTGTGGTTGCTAAGGATG GGCTGAAGC
RPLPO	NM_001002	791	866	SEQ ID NO: 46	CCATTCTATCATCAACGGGTACAAACGAGTCCCTGGCCTTGCTGTGTGGAGACGGATTACACCTTCCG ACTTGTGA
SNRPF	NM_003095	71	150	SEQ ID NO: 47	GGCTGGTCGGCAGAGAGTAGGCTGCAACATTCCGGCCGTGGTTTACATGAGTTTACCCCTCAATCC CAAACCTTTCTCA
Src	NM_004383	979	1043	SEQ ID NO: 48	CCTGAACATGAAGGAGCTGAAGCTGCTGCAGACCATCGGGAAGGGGAGTTCCGGAGACGTGATG
STK15	NM_003600	1101	1170	SEQ ID NO: 49	CATCTTCCAGGAGGACCACCTCTCTGTGGCACCCCTGGACTACCTGCCCCCTGAAATGATTGAAGGT CGGA
STMY3	NM_005940	2090	2180	SEQ ID NO: 50	CCTGGAGGCTGCAACATACCTCAATCCTGTCCACAGCCGGATCCTCCTGAAGCCCTTTTTCGCAGC ACTGCTATCCTCCAAAGCCATTGTA
SURV	NM_001168	737	817	SEQ ID NO: 51	TGTTTTGATTCCCGGGCTTACCAGGTGAGAA GTGAGGAGGGAAGAGGCAGTGTCCCTTTTGCTA GAGCTGACAGCTTG
TFR3	NM_003234	2110	2178	SEQ ID NO: 52	GCCAACTGCTTTTCATTTGTGAGGGATCTGAACCAATACAGAGCAGACATAAAGGAAATGGGCTGA GT
TOP2A	NM_001067	4505	4577	SEQ ID NO: 53	AATCCAAGGGGAGAGTGATGACTTCCATATGGACTTTGACTCAGCTGTGGCTCCTCGGGCAAAA TCTGTAC
TS	NM_001071	764	829	SEQ ID NO: 54	GCCTGGGTGTGCCCTTTCAACATCGCCAGCTACGCCCTGCTCACGTACATGATTGGGCACATCAGC

TABLE 4

Gene	Accession	Name	SEQ ID NO	Sequence
B-Catenin	NM_001904	S2150/B-Cate.f3	SEQ ID NO:55	GGCTCTTGTCGCTACTGTCTT
B-Catenin	NM_001904	S2151/B-Cate.f3	SEQ ID NO:56	TCAGATGACGAAGACACAGATG
B-Catenin	NM_001904	S5046/B-Cate.p3	SEQ ID NO:57	AGGCTCAGTGATGTCTTCCCTGTCACCAG
BAG1	NM_004323	S1386/BAG1.f2	SEQ ID NO:58	CGTTGTGACGACCTTGGAAATACAA
BAG1	NM_004323	S1387/BAG1.f2	SEQ ID NO:59	GTTCACCTCTTCCCTGTGGACTGT
BAG1	NM_004323	S4731/BAG1.p2	SEQ ID NO:60	CCCAATTAAACATGACCCGGCAACCAT
BIN1	NM_004305	S2651/BIN1.f3	SEQ ID NO:61	CCTGCAAAAGGGAACAAGAG
BIN1	NM_004305	S2652/BIN1.f3	SEQ ID NO:62	CGTGGTTGACTCTGATCTCG
BIN1	NM_004305	S4954/BIN1.p3	SEQ ID NO:63	CTTCGCCTCCAGATGGCTCCC
BUB1	NM_004336	S4294/BUB1.f1	SEQ ID NO:64	CCGAGGTTAATCCAGCACGTA
BUB1	NM_004336	S4295/BUB1.f1	SEQ ID NO:65	AAGACATGGCGCTCTCAGTTC
BUB1	NM_004336	S4296/BUB1.p1	SEQ ID NO:66	TGCTGGGAGCCTACACTTGGCCC
C20 orf1	NM_012112	S3560/C20 or.f1	SEQ ID NO:67	TCAGCTGTGAGCTGCGGATA
C20 orf1	NM_012112	S3561/C20 or.f1	SEQ ID NO:68	ACGGTCCTAGGTTTGAGGTTAAGA
C20 orf1	NM_012112	S3562/C20 or.p1	SEQ ID NO:69	CAGGTCCCATTCGCGGGCG
CCNB1	NM_031966	S1720/CCNB1.f2	SEQ ID NO:70	TTCAGGTTGTTGCAGGAGAC
CCNB1	NM_031966	S1721/CCNB1.f2	SEQ ID NO:71	CATCTTCTTGGGCACACAAT
CCNB1	NM_031966	S4733/CCNB1.p2	SEQ ID NO:72	TGCTCCATTATTGATCGGTTTCATGCA
CCNE2	NM_057749	S1458/CCNE2.f2	SEQ ID NO:73	ATGCTGTGCTCCTTCTCTAACT
CCNE2	NM_057749	S1459/CCNE2.f2	SEQ ID NO:74	ACCCAAATTGTGATATACAAAAGGTT
CCNE2	NM_057749	S4945/CCNE2.p2	SEQ ID NO:75	TACCAAGCAACCTACATGTCAAGAAAGCCC
CDC20	NM_001255	S4447/CDC20.f1	SEQ ID NO:76	TGGATTGGAGTTCTGGGAATG
CDC20	NM_001255	S4448/CCC20.f1	SEQ ID NO:77	GCTTGCACTCCACAGGTACACA
CDC20	NM_001255	S4449/CDC20.p1	SEQ ID NO:78	ACTGGCCGTGGCACTGGACAACA
CDH1	NM_004360	S0073/CDH1.f3	SEQ ID NO:79	TGAGTGTCCCCCGGTATCTTC
CDH1	NM_004360	S0075/CDH1.f3	SEQ ID NO:80	CAGCCGCTTTCAGATTTTCAT
CDH1	NM_004360	S4990/CDH1.p3	SEQ ID NO:81	TGCCAATCCCGATGAAATTGGAAATTT
CEGP1	NM_020974	S1494/CEGP1.f2	SEQ ID NO:82	TGACAAATCAGCACACCTGCAT
CEGP1	NM_020974	S1495/CEGP1.f2	SEQ ID NO:83	TGTGACTACAGCCGTGATCCTTA
CEGP1	NM_020974	S4735/CEGP1.p2	SEQ ID NO:84	CAGGCCCTCTTCCGAGCGGT
CIAP1	NM_001166	S0764/CIAP1.f2	SEQ ID NO:85	TGCCTGTGTGGGGAAGCT
CIAP1	NM_001166	S0765/CIAP1.f2	SEQ ID NO:86	GGAAAATGCCTCCGGTGTT
CIAP1	NM_001166	S4802/CIAP1.p2	SEQ ID NO:87	TGACATAGCATCATCCTTTGGTTCCCAGTT

(continued)

Gene	Accession	Name	SEQ ID NO	Sequence
cMYC	NM_002467	S0085/cMYC.f3	SEQ ID NO:88	TCCCTCCACTCGGAAGGACTA
cMYC	NM_002467	S0087/cMYC.f3	SEQ ID NO:89	CGGTTGTTGCTGATCTGTCTCA
cMYC	NM_002467	S4994/cMYC.p3	SEQ ID NO:90	TCTGACACTGTCCAAC TTGACCCTCTT
CTSL2	NM_001333	S4354/CTSL2.f1	SEQ ID NO:91	TGTC TACTGAGCGAGCAGAA
CTSL2	NM_001333	S4355/CTSL2.r1	SEQ ID NO:92	ACCATTCAGCCCTGATTG
CTSL2	NM_001333	S4356/CTSL2.p1	SEQ ID NO:93	CTTGAGGACGCGAACAGTCCACCA
DKFZp586M0723	AL050227	S4396/DKFZp5.f1	SEQ ID NO:94	TCCATTTTCTACCTGTTAACCTTCATC
DKFZp586M0723	AL050227	S4397/DKFZp5.r1	SEQ ID NO:95	ATGCAGTCGGTCCC TTCCCT
DKFZp586M0723	AL050227	S4398/DKFZpS.p1	SEQ ID NO:96	TTGCTTCCAGGGCCTGCACAAAA
DR5	NM_003842	S2551/DR5.f2	SEQ ID NO:97	CTCTGAGACAGTGCTTCGATGACT
DR5	NM_003842	S2552/DR5.r2	SEQ ID NO:98	CCATGAGGCCCAACTTCCT
DR5	NM_003842	S4979/DR5.p2	SEQ ID NO:99	CAGACTTGGTGCCCTTTGACTCC
EpCAM	NM_002354	S1807/EpCAM.f1	SEQ ID NO:100	GGGCCCCCAGAACAAATGAT
EpCAM	NM_002354	S1808/EpCAM.r1	SEQ ID NO:101	TGCACTGCTTGGCCTTAAAGA
EpCAM	NM_002354	S4984/EpCAM.p1	SEQ ID NO:102	CCGCTCTCATCGCAGTCAGGATCAT
EstR1	NM_C00125	S0115/EstR1.f1	SEQ ID NO:103	CGTGGTGCCCTCTATGAC
EstR1	NM_000125	S0117/EstR1.r1	SEQ ID NO:104	GGCTAGTGGCGGCATGTAG
EstR1	NM_000125	S4737/EstR1.p1	SEQ ID NO:105	CTGGAGATGCTGGACGCCC
FGFR1	NM_023109	S0818/FGFR1.f3	SEQ ID NO:106	CACGGGACATTCAACACATC
FGFR1	NM_023109	S0819/FGFR1.r3	SEQ ID NO:107	GGGTGCCATCCACTTCACA
FGFR1	NM_023109	S4816/FGFR1.p3	SEQ ID NO:108	ATAAAAAGACAACCAACGGCGACTGC
FOX M1	NM_021953	S2006/FOX M1.f1	SEQ ID NO:109	CCACCCCGAGCAAATCTGT
FOX M1	NM_021953	S2007/FOX M1.r1	SEQ ID NO:110	AAATCCAGTCCCCCTACTTTGG
FOX M1	NM_021953	S4757/FOX M1.p1	SEQ ID NO:111	CCTGAATCCTGGAGGCTCACGCC
GRB7	NM_005310	S0130/GRB7.f2	SEQ ID NO:112	ccatctgcattccatctgft
GRB7	NM_005310	S0132/GRB7.r2	SEQ ID NO:113	ggccaccagggtattatctg
GRB7	NM_005310	S4726/GRB7.p2	SEQ ID NO:114	ctccccacccttgagaagtcct
GSTM1	NM_000561	S2026/GSTM1.r1	SEQ ID NO:115	GGCCACAGCTTGAATTTTCA
GSTM1	NM_000561	S2027/GSTM1.f1	SEQ ID NO:116	AAGCTATGAGGAAAAAGATACACGAT
GSTM1	NM_000561	S4739/GSTM1.p1	SEQ ID NO:117	TCAGCCACTGGCTTCTGTCAATAATCAGGAG
GSTM3	NM_000849	S2038/GSTM3.f2	SEQ ID NO:118	CAATGCCATCTTGGCTACAT
GSTM3	NM_000849	S2039/GSTM3.r2	SEQ ID NO:119	GTCCACTCGAATCTTTTCTCTTCA
GSTM3	NM_000849	S5064/GSTM3.p2	SEQ ID NO:120	CTCGCAAGCACAAACATGTGTGGTGAGA

(continued)

Gene	Accession	Name	SEQ ID NO	Sequence
HER2	NM_004448	S0142/HER2.f3	SEQ ID NO:121	CGGTGTGAGAAAGTGCAGCAA
HER2	NM_004448	S0144/HER2.f3	SEQ ID NO:122	CCTCTCGCAAGTGCTCCAT
HER2	NM_004448	S4729/HER2.p3	SEQ ID NO:123	CCAGACCATAGCACACTCGGGCAC
HNRPAB	NM_004499	S4510/HNRPAB.f3	SEQ ID NO:124	CAAGGAGCGACCAACTGA
HNRPAB	NM_004499	S4511/HNRPAB.f3	SEQ ID NO:125	GTTTGCCAAGTTAAATTTGGTACATAAT
HNRPAB	NM_004499	S4512/HNRPAB.p3	SEQ ID NO:126	CTCCATATCCAAACAAAGCATGTGTGCG
ID1	NM_002165	S0820/ID1.f1	SEQ ID NO:127	AGAACCGCAAGGTGAGCAA
ID1	NM_002165	S0821/ID1.r1	SEQ ID NO:128	TCCAACTGAAGGTCCCTGATG
ID1	NM_002165	S4832/ID1.p1	SEQ ID NO:129	TGGAGATTCTCCAGCACGTCATCGAC
IGF1R	NM_000875	S1249/IGF1R.f3	SEQ ID NO:130	GCATGGTAGCCGAAGATTTC
IGF1R	NM_000875	S1250/IGF1R.f3	SEQ ID NO:131	TTTCCGGTAATAGTCTGTCTCATAGATATC
IGF1R	NM_000875	S4895/IGF1R.p3	SEQ ID NO:132	CGCGTCATACCAAAATCTCCGATTTTGA
ITGA7	NM_002206	S0859/ITGA7.f1	SEQ ID NO:133	GATATGATTGGTCGCTGCTTG
ITGA7	NM_002206	S0920/ITGA7.r1	SEQ ID NO:134	AGAACTTCCATTCCCCACCAT
ITGA7	NM_002206	S4795/ITGA7.p1	SEQ ID NO:135	CAGCCAGGACCTGGCCATCCG
Ki-67	NM_002417	S0436/Ki-67.f2	SEQ ID NO:136	CGGACTTTGGGTGCGACTT
Ki-67	NM_002417	S0437/Ki-67.r2	SEQ ID NO:137	TTACAACTCTTCCACTGGGACGAT
Ki-67	NM_002417	S4741/Ki-67.p2	SEQ ID NO:138	CCACTTGTGCAACCAACCGCTCGT
KLK10	NM_002776	S2624/KLK1 0.f3	SEQ ID NO:139	GCCCAGAGGCTCCATCGT
KLK10	NM_002776	S2625/KLK10.r3	SEQ ID NO:140	CAGAGGTTTGAACAGTGCAGACA
KLK10	NM_002776	S4978/KLK10.p3	SEQ ID NO:141	CCTCTTCTCTCCCCAGTGGCTGA
KNSL2	BC000712	S4432/KNSL2.f2	SEQ ID NO:142	CCACCTGCGCCATGATTTTTC
KNSL2	BC000712	S4433/KNSL2.r2	SEQ ID NO:143	GCAATCTCTTCAAACACTTCATCCT
KNSL2	BC000712	S4434/KNSL2.p2	SEQ ID NO:144	TTTGACCGGGTATTTCCACCAGGAA
KRT5	NM_000424	S0175/KRT5.f3	SEQ ID NO:145	tcagtggagaaggagtggga
KRT5	NM_000424	S0177/KRT5.f3	SEQ ID NO:146	tgccatatccagaggaaaca
KRT5	NM_000424	S5015/KRT5.p3	SEQ ID NO:147	ccagtcacatctctgtgtcacaagca
LMNB1	NM_005573	S4477/LMNB1.f1	SEQ ID NO:148	TGCAAAACGCTGGTGTACACA
LMNB1	NM_006573	S4478/LMNB1.r1	SEQ ID NO:149	CCCCACGAGTTCTGGTTCTTC
LMNB1	NM_005573	S4479/LMNB1.p1	SEQ ID NO:150	CAGCCCCCAACTGACCTCATC
MCM2	NM_004526	S1602/MCM2.f2	SEQ ID NO:151	GACTTTTGCCCGCTACCTTTC
MCM2	NM_004526	S1603/MCM2.r2	SEQ ID NO:152	GCCACTAACTGCTTTCAGTATGAAGAG
MCM2	NM_004526	S4900/MCM2.p2	SEQ ID NO:153	ACAGCTCATTGTTGTACGCGCGGA

(continued)

Gene	Accession	Name	SEQ ID NO	Sequence
MCM6	NM_005915	S1704/MCM6.f3	SEQ ID NO:154	TGATGGTCTTATGTGTCACATTCA
MCM6	NM_005915	S1705/MCM6.r3	SEQ ID NO:155	TGGACAGGAAACACACCAA
MCM6	NM_005915	S4919/MCM6.p3	SEQ ID NO:156	CAGGTTTCATACCAACACAGGCTTCAGCAC
MELK	NM_014791	S4318/MELK.f1	SEQ ID NO:157	AACCCGGGATCGAAAAG
MELK	NM_014791	S4319/MELK.r1	SEQ ID NO:158	GGGCCTGCTGTCCTGAGA
MELK	NM_014791	S4320/MELK.p1	SEQ ID NO:159	TCTTAGGAACGCCGTACCAAGCCGC
MMP12	NM_002426	S4381/MMP12.f2	SEQ ID NO:160	CCAACGCTTGCCAAATCCT
MMP12	NM_002426	S4382/MMP12.r2	SEQ ID NO:161	ACGGTAGTGACAGCATCAAACTC
MMP12	NM_002426	S4383/MMP12.p2	SEQ ID NO:162	AACCAGCTCTCTGTGACCCCAATT
MMP9	NM_004994	S0656/MMP9.f1	SEQ ID NO:163	GAGAACCAATCTCACCGACA
MMP9	NM_004994	S0657/MMP9.r1	SEQ ID NO:164	CACCCGAGTGTAAACATAGC
MMP9	NM_004994	S4760/MMP9.p1	SEQ ID NO:165	ACAGGTATTCCTCTGCCAGCTGCC
MYBL2	NM_002466	S3270/MYBL2.f1	SEQ ID NO:166	GCCGAGATCGCCCAAGATG
MYBL2	NM_002466	S3271/MYBL2.r1	SEQ ID NO:167	CTTTTGATGGTAGAGTCCAGTGATTG
MYBL2	NM_002466	S4742/MYBL2.p1	SEQ ID NO:168	CAGCATTGTCTGTCTCCTCGGCA
NEK2	NM_002497	S4327/NEK2.f1	SEQ ID NO:169	GTGAGGCAGCGCGACTCT
NEK2	NM_002497	S4328/NEK2.r1	SEQ ID NO:170	TGCCAATGGTGTAACAACACTTCA
NEK2	NM_002497	S4329/NEK2.p1	SEQ ID NO:171	TGCCTTCCCGGGCTGAGGACT
NME1	NM_000269	S2526/NME1.f3	SEQ ID NO:172	CCAACCTGCAGACTCCAA
NME1	NM_000269	S2527/NME1.r3	SEQ ID NO:173	ATGTATAATGTTCTGCCAACTTGATG
NME1	NM_000269	S4949/NME1.p3	SEQ ID NO:174	CCTGGACCATCCGTGGAGACTTCT
NPD009	NM_020686	S4474/NPD009.f3	SEQ ID NO:175	GGCTGTGGCTGAGGCTGTAG
NPD009	NM_020686	S4475/NPD009.r3	SEQ ID NO:176	GGAGCATTCGAGGTCAAATCA
NPD009	NM_020686	S4476/NPD009.p3	SEQ ID NO:177	TTCCCAGAGTGTCTCACCTCCAGCAGAG
PCNA	NM_002592	S0447/PCNA.f2	SEQ ID NO:178	GAAGGTGTTGGAGGCACTCAAG
PCNA	NM_002592	S0448/PCNA.r2	SEQ ID NO:179	GGTTTACACCGCTGGAGCTAA
PCNA	NM_002592	S4784/PCNA.p2	SEQ ID NO:180	ATCCCAGCAGGCCCTCGTTGATGAG
PR	NM_000926	S1336/PR.f6	SEQ ID NO:181	GCATCAGGCTGTCATTATGG
PR	NM_000926	S1337/PR.r6	SEQ ID NO:182	AGTAGTTGTGCTGCCCCTTCC
PR	NM_000926	S4743/PR.p6	SEQ ID NO:183	TGTCCTTACCTGTGGAGCTGTAAGGTC
PREP	NM_002726	S1771/PREP.f1	SEQ ID NO:184	GGGACGGTGTTCACATTCAAG
PREP	NM_002726	S1772/PREP.r1	SEQ ID NO:185	CAGGATCCCAGAAAGTCAATGTTG
PREP	NM_002726	S4929/PREP.p1	SEQ ID NO:186	TCGCCAGTCTCCCAACTATCGCGT

(continued)

Gene	Accession	Name	SEQ ID NO	Sequence
PTTG1	NM_004219	S4525/PTTG1.f2	SEQ ID NO:187	GGCTACTCTGATCTATGTTGATAAGGAA
PTTG1	NM_004219	S4526/PTTG1.r2	SEQ ID NO:188	GCTTCAGCCCATCCTTAGCA
PTTG1	NM_004219	S4527/PTTG1.p2	SEQ ID NO:189	CACACGGGTGCCTGGTTCTCCA
RPLPO	NM_001002	S0256/RPLPO.f2	SEQ ID NO:190	CCATTCTATCATCAACGGGTACAA
RPLPO	NM_001002	S0258/RPLPO.r2	SEQ ID NO:191	TCAGCAAGTGGGAAGGTGTAATC
RPLPO	NM_001002	S4744/RPLPO.p2	SEQ ID NO:192	TCTCCACAGACAAAGGCCAGGACTCG
SNRPF	NM_003095	S4489/SNRPF.f2	SEQ ID NO:193	GGCTGGTCGGCAGAGAGTAG
SNRPF	NM_003095	S4490/SNRPF.r2	SEQ ID NO:194	TGAGGAAAGGTTTGGGATTGA
SNRPF	NM_003095	S4491/SNRPF.p2	SEQ ID NO:195	AAACTCATGTAACCCACGGCCGAATGTTG
Src	NM_004383	S1820/Src.f2	SEQ ID NO:196	CCTGAACATGAAGGAGCTGA
Src	NM_004383	S1821/Src.r2	SEQ ID NO:197	CATCACGTCTCCGAACTCC
Src	NM_004383	S5034/Src.p2	SEQ ID NO:198	TCCCGATGGTCTGCAGCAGCT
STK15	NM_003600	S0794/STK15.f2	SEQ ID NO:199	CATCTCCAGGAGGACCACT
STK15	NM_003600	S0795/STK15.r2	SEQ ID NO:200	TCCGACCTTCAATCATTTCA
STK15	NM_003600	S4745/STK15.p2	SEQ ID NO:201	CTCTGTGGCACCCCTGGACTACCTG
STMY3	NM_005940	S2067/STMY3.f3	SEQ ID NO:202	CCTGGAGGCTGCAACATACC
STMY3	NM_005940	S2068/STMY3.r3	SEQ ID NO:203	TACAATGGCTTTGGAGGATAGCA
STMY3	NM_005940	S4746/STMY3.p3	SEQ ID NO:204	ATCCTCCTGAAGCCCTTTTCGCAGC
SURV	NM_001168	S02591/SURV.f2	SEQ ID NO:205	TGTTTTGATTCCTCCGGGCTTA
SURV	NM_001168	S0261/SURV.r2	SEQ ID NO:206	CAAAGCTGTCAAGCTCTAGCAAAAG
SURV	NM_001168	S4747/SURV.p2	SEQ ID NO:207	TGCCCTTCTCCTCCCTCACTTCTCACCT
TFRC	NM_003234	S1352/TFRC.f3	SEQ ID NO:208	GCCAACTGCTTTCAATTTGTG
TFRC	NM_003234	S1353/TFRC.r3	SEQ ID NO:209	ACTCAGGCCCATTTCCCTTA
TFRC	NM_003234	S4748/TFRC.p3	SEQ ID NO:210	AGGGATCTGAACCAATACAGAGCAGACA
TOP2A	NM_001067	S0271/TOP2A.f4	SEQ ID NO:211	AATCCAAGGGGGAGAGTGAT
TOP2A	NM_001067	S0273/TOP2A.r4	SEQ ID NO:212	GTACAGATTTTGCCCGAGGA
TOP2A	NM_001067	S4777/TOP2A.p4	SEQ ID NO:213	CATATGGACTTTGACTCAGCTGTGGC
TS	NM_001071	S0280/TS.f1	SEQ ID NO:214	GCCTCGGTGTGCCCTTTCA
TS	NM_001071	S0282/TS.r1	SEQ ID NO:215	CGTGATGTGCGCAATCATG
TS	NM_001071	S4780/TS.p1	SEQ ID NO:216	CATGCCAGCTACGCCCTGCTC

SEQUENCE LISTING

[0103]

5 <110> Genomic Health

 <120> Prediction of Likelihood of Cancer Recurrence

 <130> GRF/FP6349666

10 <140> 04809450.2

 <141> 2004-06-17

 <150> PCT/US2004/019567

15 <151> 2004-06-17

 <150> US 60/482,339

 <151> 2003-06-24

20 <160> 216

 <170> FastSEQ for Windows Version 4.0

 <210> 1

25 <211> 80

 <212> DNA

 <213> Artificial Sequence

 <220>

30 <223> PCR Amplicon

 <400> 1

35 ggctcttgtg cgtactgtcc ttcgggctgg tgacagggaa gacatcactg agcctgccat 60

 ctgtgctctt cgtcactctga 80

 <210> 2

 <211> 81

40 <212> DNA

 <213> Artificial Sequence

 <220>

 <223> PCR Amplicon

45 <400> 2

 cgttgtcagc acttgggaata caagatgggt gccgggtcat gttaattggg aaaaagaaca 60

50 gtccacagga agaggttgaa c 81

 <210> 3

 <211> 76

 <212> DNA

 <213> Artificial Sequence

55 <220>

 <223> PCR Amplicon

<400> 3

5 cctgcaaaag ggaacaagag cccttcgcct ccagatggct cccctgccgc ccccccgag 60
atcagagtca accacg 76

<210> 4

<211> 68

<212> DNA

10 <213> Artificial Sequence

<220>

<223> PCR Amplicon

15 <400> 4

ccgaggttaa tccagcacgt atggggccaa gtgtaggctc ccagcaggaa ctgagagcgc 60
catgtctt 68

20

<210> 5

<211> 65

<212> DNA

<213> Artificial Sequence

25

$\langle 220 \rangle$

<223> PCR Amplicon

<400> 5

30

```
ttagctgtga gctgcggata ccgcccggca atgggaacctg ctcttaacct caaacctagc 60
accgt                                           65
```

35

<210> 6

<211> 84

<212> DNA

<213> Artificial Sequence

40

$\langle 220 \rangle$

<223> PCR Amplicon

<400> 6

45

```
ttcaggttgt  tgcaggagac  catgtacatg  actgtctcca  ttattgatcg  gttcatgcag  60
aataattgtg  tgcccaagaa  gatg                                     84
```

<210> 7

<211> 82

<212> DNA

<213> Artificial Sequence

<220>

55 <223> PCR Amplicon

<400> 7

atgctgtggc tccttcctaa ctggggcttt cttgacatgt aggttgcttg gtaataacct 60
 ttttgtatat cacaatttgg gt 82

5 <210> 8
 <211> 68
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> PCR Amplicon
 <400> 8

15 tggattggag ttctgggaat gtactggccg tggcactgga caacagtgtg tacctgtgga 60
 gtgcaagc 68

20 <210> 9
 <211> 81
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> PCR Amplicon
 <400> 9

30 tgagtgtccc ccggtatctt ccccgccctg ccaatcccga tgaaattgga aattttattg 60
 atgaaaaatct gaaagcggct g 81

35 <210> 10
 <211> 77
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> PCR Amplicon
 <400> 10

45 tgacaatcag cacacctgca ttcaccgctc ggaagagggc ctgagctgca tgaataagga 60
 tcacggctgt agtcaca 77

50 <210> 11
 <211> 72
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Amplicon

55 <400> 11

tgctgtggt gggaagctca gtaactggga accaaaggat gatgctatgt cagaacaccg 60
gaggcatattt cc 72

5 <210> 12
<211> 84
<212> DNA
<213> Artificial Sequence

10 <220>
<223> PCR Amplicon

<400> 12

15 tccctccact cggaaggact atcctgctgc caagaggggc aagttggaca gtgtcagagt 60
cctgagacag atcagcaaca accg 84

20 <210> 13
<211> 67
<212> DNA
<213> Artificial Sequence

25 <220>
<223> PCR Amplicon

<400> 13

30 tgtctcactg agcgagcaga atctggtgga ctgttcgcgt cctcaaggca atcagggctg 60
caatggt 67

35 <210> 14
<211> 74
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Amplicon

40 <400> 14

tccattttct acctgttaac cttcatcatt ttgtgcaggc cctggaagca aagagaggaa 60

45 gggaccgact gcat 74

50 <210> 15
<211> 84
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Amplicon

55 <400> 15

ctctgagaca gtgcttcgat gactttgcag acttggtgcc ctttgactcc tgggagccgc 60
tcatgaggaa gttgggcctc atgg 84

5 <210> 16
<211> 75
<212> DNA
<213> Artificial Sequence

10 <220>
<223> PCR Amplicon

<400> 16

15 gggccctcca gaacaatgat gggctttatg atcctgactg cgatgagagc gggctcttta 60
aggccaagca gtgca 75

20 <210> 17
<211> 68
<212> DNA
<213> Artificial Sequence

25 <220>
<223> PCR Amplicon

<400> 17

30 cgtggtgccc ctctatgacc tgctgctgga gatgctggac gccaccgcc tacatgcgcc 60
cactagcc 68

35 <210> 18
<211> 74
<212> DNA
<213> Artificial Sequence

40 <220>
<223> PCR Amplicon

<400> 18

45 cacgggacat tcaccacatc gactactata aaaagacaac caacggccga ctgcctgtga 60
agtggatggc accc 74

50 <210> 19
<211> 82
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Amplicon

55 <400> 19

ccaccccgag caaatctgtc ctccccagaa cccctgaatc ctggaggctc acgccccag 60
 ccaaagtagg gggactggat tt 82

5 <210> 20
 <211> 67
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> PCR Amplicon
 <400> 20

15 ccatctgcat ccatcttgtt tgggctcccc acccttgaga agtgcctcag ataataccct 60
 ggtggcc 67

20 <210> 21
 <211> 86
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> PCR Amplicon
 <400> 21

30 aagctatgag gaaaagaagt acacgatggg ggacgctcct gattatgaca gaagccagtg 60
 gctgaatgaa aaattcaagc tgggcc 86

35 <210> 22
 <211> 76
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> PCR Amplicon
 <400> 22

45 caatgccatc ttgcgctaca tcgctcgcaa gcacaacatg tgtggtgaga ctgaagaaga 60
 aaagattcga gtggac 76

50 <210> 23
 <211> 70
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Amplicon

55 <400> 23

cggtgtgaga agtgcagcaa gccctgtgcc cgagtgtgct atggtctggg catggagcac 60
 ttgcgagagg 70

5 <210> 24
 <211> 84
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> PCR Amplicon
 <400> 24

15 caagggagcg accaactgat cgcacacatg ctttgtttgg atatggagtg aacacaatta 60
 tgtaccaaat ttaacttggc aaac 84

20 <210> 25
 <211> 70
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> PCR Amplicon
 <400> 25

30 agaaccgcaa ggtgagcaag gtggagattc tccagcacgt catcgactac atcagggacc 60
 ttcagttgga 70

35 <210> 26
 <211> 83
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> PCR Amplicon
 <400> 26

45 gcatggtagc cgaagatttc acagtcaaaa tcggagattt tggtagacg cgagatatct 60
 atgagacaga ctattaccgg aaa 83

50 <210> 27
 <211> 79
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> PCR Amplicon
 <400> 27

gatatgattg gtcgctgctt tgtgctcagc caggacctgg ccatccggga tgagttggat 60
 ggtggggaat ggaagttct 79

5 <210> 28
 <211> 80
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> PCR Amplicon
 <400> 28

15 cggacttttg gtgcgacttg acgagcgggtg gttcgacaag tggccttgcg ggccggatcg 60
 tcccagtgga agagttgtaa 80

20 <210> 29
 <211> 78
 <212> DNA
 <213> Artificial Sequence

25 <220>
 <223> PCR Amplicon
 <400> 29

30 gccacagaggc tccatcgctc atcctcttcc tcccagtcg gctgaactct ccccttgtct 60
 gcactgttca aacctctg 78

35 <210> 30
 <211> 77
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> PCR Amplicon
 <400> 30

45 ccacctcgcc atgatttttc ctttgaccgg gtattccac caggaagtgg acaggatgaa 60
 gtgtttgaag agattgc 77

50 <210> 31
 <211> 69
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Amplicon

55 <400> 31

tcagtggaga aggagttgga ccagtcaaca tctctgttgt cacaagcagt gtttcctctg 60
gatatggca 69

5 <210> 32
<211> 66
<212> DNA
<213> Artificial Sequence

10 <220>
<223> PCR Amplicon

<400> 32

15 tgcaaacgct ggtgtcacag ccagcccccc aactgacctc atctggaaga accagaactc 60
gtgggg 66

20 <210> 33
<211> 75
<212> DNA
<213> Artificial Sequence

25 <220>
<223> PCR Amplicon

<400> 33

30 gacttttgcc cgctaccttt cattccggcg tgacaacaat gagctgttgc tcttcatact 60
gaagcagtta gtggc 75

35 <210> 34
<211> 82
<212> DNA
<213> Artificial Sequence

40 <220>
<223> PCR Amplicon

<400> 34

45 tgatggtcct atgtgtcaca ttcattcacag gtttcatacc aacacaggct tcagcacttc 60
ctttggtgtg tttcctgtcc ca 82

50 <210> 35
<211> 65
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Amplicon

55 <400> 35

aacccggcga tcgaaaagat tcttaggaac gccgtaccag ccgcgtctct caggacagca 60

ggccc

65

5
 <210> 36
 <211> 78
 <212> DNA
 <213> Artificial Sequence

10
 <220>
 <223> PCR Amplicon

<400> 36

15
 ccaacgcttg ccaaattcctg acaattcaga accagctctc tgtgacccca atttgagttt 60
 tgatgctgtc actaccgt 78

20
 <210> 37
 <211> 67
 <212> DNA
 <213> Artificial Sequence

25
 <220>
 <223> PCR Amplicon

<400> 37

30
 gagaaccaat ctcaccgaca ggcagctggc agaggaatac ctgtaccgct atgggttacac 60
 tcgggtg 67

35
 <210> 38
 <211> 74
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Amplicon

40
 <400> 38

45
 gccgagatcg ccaagatgtt gccagggagg acagacaatg ctgtgaagaa tcactggaac 60
 tctaccatca aaag 74

50
 <210> 39
 <211> 79
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Amplicon

<400> 39

55
 gtgaggcagc gcgactctgg cgactggccg gccatgcctt cccgggctga ggactatgaa 60
 gtgttgatca ccattggca 79

<210> 40
 <211> 74
 <212> DNA
 <213> Artificial Sequence

5

<220>
 <223> PCR Amplicon

10

<400> 40
 ccaaccctgc agactccaag cctgggacca tccgtggaga cttctgcata caagttggca 60
 ggaacattat acat 74

15

<210> 41
 <211> 73
 <212> DNA
 <213> Artificial Sequence

20

<220>
 <223> PCR Amplicon

<400> 41

25

ggctgtggct gaggctgtag catctctgct ggaggtgaga cactctggga actgatttga 60
 cctcgaatgc tcc 73

30

<210> 42
 <211> 71
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Amplicon

35

<400> 42

40

gaaggtgttg gaggcactca aggacctcat caacgaggcc tgctgggata ttagctccag 60
 cggtgtaaac c 71

45

<210> 43
 <211> 85
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Amplicon

50

<400> 43

55

gcatcaggct gtcattatgg tgccttacc tgtgggagct gtaaggcttt ctttaagagg 60
 gcaatggaag ggcagcacia ctact 85

<210> 44
 <211> 76

<212> DNA

<213> Artificial Sequence

$\langle 220 \rangle$

<223> PCR Amplicon

<400> 44

```

gggacggtgt  tcacattcaa  gacgaatcgc  cagtctccca  actatcgcgt  gatcaacatt  60
gacttctggg  atcctg                                           76

```

<210> 45

<211> 74

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Amplicon

<400> 45

ggctactctg atctatgttg ataaggaaaa tggagaacca ggcacccgtg tggttgctaa 60
 ggatgggctg aagc 74

<210> 46

<211> 75

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Amplicon

<400> 46

```
ccattctatc atcaacgggt acaaacgagt cctggccttg tctgtggaga cggattacac 60
cttcccactt gctga                                     75
```

<210> 47

<211> 79

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Amplicon

<400> 47

```
ggctgggtcgg cagagagtag cctgcaacat tcggccgtgg tttacatgag tttacccctc 60
aatcccaaac ctttcctca                                     79
```

<210> 48

<211> 64

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Amplicon

<400> 48

5

cctgaacatg aaggagctga agctgctgca gaccatcggg aagggggagt tcggagacgt 60
 gatg 64

<210> 49

10

<211> 69

<212> DNA

<213> Artificial Sequence

<220>

15

<223> PCR Amplicon

<400> 49

20

catctttccag gaggaccact ctctgtggca ccctggacta cctgccccct gaaatgattg 60
 aaggtcggg 69

<210> 50

25

<211> 90

<212> DNA

<213> Artificial Sequence

<220>

30

<223> PCR Amplicon

<400> 50

35

cctggagggt gcaacatacc tcaatcctgt ccagggccgg atcctcctga agcccttttc 60
 gcagcaactgc taccctccaa agccattgta 90

<210> 51

40

<211> 79

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Amplicon

45

<400> 51

tgtttttgatt cccgggctta ccaggtgaga agtgaggag gaagaaggca gtgtcccttt 60
 tgctagagct gacagcttg 79

50

<210> 52

<211> 68

<212> DNA

<213> Artificial Sequence

55

<220>

<223> PCR Amplicon

<400> 52

5 gccaaactgct ttcatattgtg agggatctga accaatacag agcagacata aaggaaatgg 60
gcctgagt 68

<210> 53

<211> 72

<212> DNA

10 <213> Artificial Sequence

$\langle 220 \rangle$

<223> PCR Amplicon

15 <400> 53

```
aatccaaggg ggagagtgat gacttccata tggactttga ctcagctgtg gctcctcggg 60
caaaatctgt ac 72
```

20

<210> 54

<211> 65

<212> DNA

<213> Artificial Sequence

25

$\langle 220 \rangle$

<223> PCR Amplicon

<400> 54

30

```
gcctcgggtgt gcctttcaac atcgccagct acgccttgc cactacatg attgcgcaca 60
tcacg                                             65
```

35

<210> 55

<211> 22

<212> DNA

<213> Artificial Sequence

40

$\langle 220 \rangle$

<223> PCR primer-probe

<400> 55

ggctcttg tg cgtactgtcc tt 22

45

<210> 56

<211> 23

<212> DNA

<213> Artificial Sequence

50

$\langle 220 \rangle$

<223> PCR primer-probe

<400> 56

55 tcagatgacg aagagcacag atg 23

<210> 57

<211> 29

<212> DNA
 <213> Artificial Sequence

5 <220>
 <223> PCR primer-probe

 <400> 57
 aggctcagtg atgtcttccc tgcaccag 29

10 <210> 58
 <211> 23
 <212> DNA
 <213> Artificial Sequence

15 <220>
 <223> PCR primer-probe

20 <400> 58
 cgtgtgcagc acttgaata caa 23

25 <210> 59
 <211> 24
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> PCR primer-probe

 <400> 59
 gttcaacctc ttctgtgga ctgt 24

35 <210> 60
 <211> 26
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> PCR primer-probe

40 <400> 60
 cccaattaac atgaccggc aaccat 26

45 <210> 61
 <211> 20
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> PCR primer-probe

50 <400> 61
 cctgcaaaag ggaacaagag 20

55 <210> 62
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

5 <400> 62
 cggtggtgac tctgatctcg 20

<210> 63
 <211> 21
 <212> DNA
 10 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

15 <400> 63
 cttcgcctcc agatggctcc c 21

<210> 64
 <211> 21
 20 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

25 <400> 64
 ccgaggtaa tccagcacgt a 21

<210> 65
 <211> 21
 30 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

35 <400> 65
 aagacatggc gctctcagtt c 21

40 <210> 66
 <211> 23
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <223> PCR primer-probe

<400> 66
 50 tgctgggagc ctacacttgg ccc 23

<210> 67
 <211> 20
 <212> DNA
 55 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

<400> 67
 tcagctgtga gctgcggata 20

5
 <210> 68
 <211> 24
 <212> DNA
 <213> Artificial Sequence

10
 <220>
 <223> PCR primer-probe

<400> 68
 acggtcctag gttgaggtt aaga 24

15
 <210> 69
 <211> 19
 <212> DNA
 <213> Artificial Sequence

20
 <220>
 <223> PCR primer-probe

<400> 69
 caggtcccat tgccgggcg 19

25
 <210> 70
 <211> 20
 <212> DNA
 <213> Artificial Sequence

30
 <220>
 <223> PCR primer-probe

<400> 70
 ttcaggttgt tgcaggagac 20

35
 <210> 71
 <211> 20
 <212> DNA
 <213> Artificial Sequence

40
 <220>
 <223> PCR primer-probe

45
 <400> 71
 catcttcttg ggcacacaat 20

<210> 72
 <211> 27
 <212> DNA
 <213> Artificial Sequence

50
 <220>
 <223> PCR primer-probe

55
 <400> 72
 tgtctccatt attgatcggt tcatgca 27

<210> 73
 <211> 22
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> PCR primer-probe
 <400> 73
 10 atgctgtggc tccttcctaa ct 22
 <210> 74
 <211> 27
 <212> DNA
 15 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 20 <400> 74
 acccaaattg tgatatacaa aaaggtt 27
 <210> 75
 <211> 30
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 30 <400> 75
 taccaagcaa cctacatgtc aagaaagccc 30
 <210> 76
 <211> 21
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 40 <400> 76
 tggattggag ttctgggaat g 21
 <210> 77
 <211> 22
 45 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 50 <400> 77
 gcttgactc cacaggtaca ca 22
 55 <210> 78
 <211> 23
 <212> DNA

<213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 5
 <400> 78
 actggccgtg gactggaca aca 23
 <210> 79
 10 <211> 21
 <212> DNA
 <213> Artificial Sequence
 <220>
 15 <223> PCR primer-probe
 <400> 79
 tgagtgtccc ccggtatctt c 21
 20 <210> 80
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> PCR primer-probe
 <400> 80
 30 cagccgcttt cagatttca t 21
 <210> 81
 <211> 27
 <212> DNA
 <213> Artificial Sequence
 35 <220>
 <223> PCR primer-probe
 <400> 81
 40 tgccaatccc gatgaaattg gaaattt 27
 <210> 82
 <211> 21
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 50 <400> 82
 tgacaatcag cacacctgca t 21
 <210> 83
 <211> 23
 55 <212> DNA
 <213> Artificial Sequence
 <220>

<223> PCR primer-probe
 <400> 83
 tgtgactaca gccgtgatcc tta 23
 5
 <210> 84
 <211> 20
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> PCR primer-probe
 <400> 84
 15 caggccctct tccgagcgg 20
 <210> 85
 <211> 18
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 25 <400> 85
 tgcctgtggt gggaagct 18
 <210> 86
 <211> 19
 30 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 35 <400> 86
 ggaaaatgcc tccggtgtt 19
 <210> 87
 40 <211> 30
 <212> DNA
 <213> Artificial Sequence
 <220>
 45 <223> PCR primer-probe
 <400> 87
 tgacatagca tcataccttg gttcccagtt 30
 50 <210> 88
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 55 <220>
 <223> PCR primer-probe
 <400> 88

tccctccact cggaaggact a 21
 <210> 89
 <211> 22
 5 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 10
 <400> 89
 cggtgtgtgc tgatctgtct ca 22
 <210> 90
 15 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 20 <223> PCR primer-probe
 <400> 90
 tctgacactg tccaactga ccctctt 27
 25 <210> 91
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 30 <220>
 <223> PCR primer-probe
 <400> 91
 35 tgtctcactg agcgagcaga a 21
 <210> 92
 <211> 19
 <212> DNA
 <213> Artificial Sequence
 40 <220>
 <223> PCR primer-probe
 <400> 92
 45 accattgcag ccctgattg 19
 <210> 93
 <211> 24
 <212> DNA
 50 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 55 <400> 93
 cttgaggacg cgaacagtcc acca 24
 <210> 94

<211> 27
 <212> DNA
 <213> Artificial Sequence

5 <220>
 <223> PCR primer-probe

<400> 94
 tccattttct acctgttaac cttcatc 27

10 <210> 95
 <211> 19
 <212> DNA
 <213> Artificial Sequence

15 <220>
 <223> PCR primer-probe

<400> 95
 20 atgcagtcgg tcccttcct 19

<210> 96
 <211> 23
 <212> DNA
 25 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

30 <400> 96
 ttgcttccag ggcctgcaca aaa 23

<210> 97
 <211> 24
 35 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

40 <400> 97
 ctctgagaca gtgcttcgat gact 24

<210> 98
 45 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 50 <223> PCR primer-probe

<400> 98
 ccatgaggcc caacttcct 19

55 <210> 99
 <211> 23
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

 <400> 99
 5 cagacttggt gcccttgac tcc 23

 <210> 100
 <211> 20
 <212> DNA
 10 <213> Artificial Sequence

 <220>
 <223> PCR primer-probe

 <400> 100
 15 gggccctcca gaacaatgat 20

 <210> 101
 <211> 21
 20 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> PCR primer-probe
 25
 <400> 101
 tgcactgctt ggccttaaag a 21

 <210> 102
 <211> 25
 30 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> PCR primer-probe
 35
 <400> 102
 ccgctctcat cgcagtcagg atcat 25

 <210> 103
 <211> 19
 40 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> PCR primer-probe
 45
 <400> 103
 50 cgtggtgccc ctctatgac 19

 <210> 104
 <211> 19
 <212> DNA
 <213> Artificial Sequence
 55
 <220>
 <223> PCR primer-probe

<400> 104
 ggctagtggg cgcatgtag 19

5 <210> 105
 <211> 19
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> PCR primer-probe

<400> 105
 ctggagatgc tggacgccc 19

15 <210> 106
 <211> 20
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> PCR primer-probe

<400> 106
 cacgggacat tcaccacatc 20

25 <210> 107
 <211> 19
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> PCR primer-probe

<400> 107
 ggggtgccatc cacttcaca 19

35 <210> 108
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

45 <400> 108
 ataaaaagac aaccaacggc cgactgc 27

<210> 109
 <211> 19
 <212> DNA
 <213> Artificial Sequence

50 <220>
 <223> PCR primer-probe

55 <400> 109
 ccaccccgag caaatctgt 19

<210> 110
 <211> 22
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> PCR primer-probe
 <400> 110
 10 aaatccagtc ccctacttt gg 22
 <210> 111
 <211> 23
 <212> DNA
 15 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 20 <400> 111
 cctgaatcct ggaggctcac gcc 23
 <210> 112
 <211> 20
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 30 <400> 112
 ccatctgcat ccatctgtt 20
 <210> 113
 <211> 20
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 40 <400> 113
 ggccaccagg gtattatctg 20
 <210> 114
 <211> 23
 45 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 50 <400> 114
 ctccccaccc ttgagaagtg cct 23
 55 <210> 115
 <211> 20
 <212> DNA

<213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 5
 <400> 115
 ggcccagctt gaattttca 20
 <210> 116
 10 <211> 27
 <212> DNA
 <213> Artificial Sequence
 <220>
 15 <223> PCR primer-probe
 <400> 116
 aagctatgag gaaaagaagt acacgat 27
 20 <210> 117
 <211> 30
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> PCR primer-probe
 <400> 117
 30 tcagccactg gcttctgtca taatcaggag 30
 <210> 118
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 35 <220>
 <223> PCR primer-probe
 <400> 118
 40 caatgccatc ttgcgctaca t 21
 <210> 119
 <211> 25
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 50 <400> 119
 gtccactcga atcttttctt ctca 25
 <210> 120
 <211> 27
 55 <212> DNA
 <213> Artificial Sequence
 <220>

<223> PCR primer-probe
 <400> 120
 ctcgcaagca caacatgtgt ggtgaga 27
 5
 <210> 121
 <211> 20
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> PCR primer-probe
 <400> 121
 cggtgtgaga agtgcagcaa 20
 15
 <210> 122
 <211> 19
 <212> DNA
 <213> Artificial Sequence
 20
 <220>
 <223> PCR primer-probe
 <400> 122
 cctctcgcaa gtgctccat 19
 25
 <210> 123
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> PCR primer-probe
 <400> 123
 ccagaccata gcacactcgg gcac 24
 35
 <210> 124
 <211> 19
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> PCR primer-probe
 <400> 124
 caagggagcg accaactga 19
 45
 <210> 125
 <211> 28
 <212> DNA
 <213> Artificial Sequence
 50
 <220>
 <223> PCR primer-probe
 <400> 125

gtttgccaag ttaaatttgg tacataat 28

5 <210> 126
 <211> 28
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> PCR primer-probe

<400> 126
 ctccatatcc aaacaaagca tgtgtgcg 28

15 <210> 127
 <211> 19
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> PCR primer-probe

<400> 127
 agaaccgcaa ggtgagcaa 19

25 <210> 128
 <211> 21
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> PCR primer-probe

<400> 128
 tccaactgaa ggtccctgat g 21

35 <210> 129
 <211> 26
 <212> DNA
 <213> Artificial Sequence

40 <220>
 <223> PCR primer-probe

45 <400> 129
 tggagattct ccagcacgtc atcgac 26

<210> 130
 <211> 21
 <212> DNA
 50 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

55 <400> 130
 gcatggtagc cgaagatttc a 21

<210> 131

<211> 30
 <212> DNA
 <213> Artificial Sequence

5 <220>
 <223> PCR primer-probe

<400> 131
 10 ttccggtaa tagtgtgtct catagatatc 30

<210> 132
 <211> 28
 <212> DNA
 <213> Artificial Sequence

15 <220>
 <223> PCR primer-probe

<400> 132
 20 cgcgtcatac caaaatctcc gattttga 28

<210> 133
 <211> 22
 <212> DNA
 25 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

30 <400> 133
 gatatgattg gtcgctgctt tg 22

<210> 134
 <211> 21
 35 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

40 <400> 134
 agaactcca ttccccacca t 21

<210> 135
 45 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 50 <223> PCR primer-probe

<400> 135
 cagccaggac ctggccatcc g 21

55 <210> 136
 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

 <400> 136
 5 cggactttgg gtgcgactt 19

 <210> 137
 <211> 24
 <212> DNA
 10 <213> Artificial Sequence

 <220>
 <223> PCR primer-probe

 15 <400> 137
 ttacaactct tcactggga cgat 24

 <210> 138
 <211> 23
 20 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> PCR primer-probe
 25
 <400> 138
 ccacttgctg aaccaccgct cgt 23

 <210> 139
 30 <211> 18
 <212> DNA
 <213> Artificial Sequence

 <220>
 35 <223> PCR primer-probe

 <400> 139
 gccagaggc tccatcgt 18
 40
 <210> 140
 <211> 23
 <212> DNA
 <213> Artificial Sequence

 45 <220>
 <223> PCR primer-probe

 <400> 140
 50 cagaggttg aacagtgcag aca 23

 <210> 141
 <211> 23
 <212> DNA
 <213> Artificial Sequence
 55
 <220>
 <223> PCR primer-probe

<400> 141
 cctcttcctc cccagtcggc tga 23

5
 <210> 142
 <211> 20
 <212> DNA
 <213> Artificial Sequence

10
 <220>
 <223> PCR primer-probe

<400> 142
 ccacctcgcc atgatttttc 20

15
 <210> 143
 <211> 25
 <212> DNA
 <213> Artificial Sequence

20
 <220>
 <223> PCR primer-probe

<400> 143
 gcaatctctt caaacacttc atcct 25

25
 <210> 144
 <211> 25
 <212> DNA
 <213> Artificial Sequence

30
 <220>
 <223> PCR primer-probe

<400> 144
 tttgaccggg tattccacc aggaa 25

35
 <210> 145
 <211> 20
 <212> DNA
 <213> Artificial Sequence

40
 <220>
 <223> PCR primer-probe

45
 <400> 145
 tcagtggaga aggagttgga 20

<210> 146
 <211> 20
 <212> DNA
 <213> Artificial Sequence

50
 <220>
 <223> PCR primer-probe

55
 <400> 146
 tgccatatcc agaggaaaca 20

<210> 147
 <211> 28
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> PCR primer-probe
 <400> 147
 10 ccagtcaaca tctctgttgt cacaagca 28
 <210> 148
 <211> 19
 <212> DNA
 15 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 20 <400> 148
 tgcaaacgct ggtgtcaca 19
 <210> 149
 <211> 21
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 30 <400> 149
 ccccacgagt tctggttct c 21
 <210> 150
 <211> 22
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 40 <400> 150
 cagccccca actgacctca tc 22
 <210> 151
 <211> 21
 45 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 50 <400> 151
 gacttttgcc cgctaccttt c 21
 55 <210> 152
 <211> 26
 <212> DNA

<213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 5
 <400> 152
 gccactaact gcttcagtat gaagag 26
 <210> 153
 10 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 15 <223> PCR primer-probe
 <400> 153
 acagctcatt gttgtcacgc cgga 24
 20 <210> 154
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> PCR primer-probe
 <400> 154
 30 tgatggctct atgtgtcaca ttca 24
 <210> 155
 <211> 20
 <212> DNA
 <213> Artificial Sequence
 35 <220>
 <223> PCR primer-probe
 <400> 155
 40 tgggacagga aacacaccaa 20
 <210> 156
 <211> 30
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 50 <400> 156
 caggtttcat accaacacag gcttcagcac 30
 <210> 157
 <211> 18
 55 <212> DNA
 <213> Artificial Sequence
 <220>

<223> PCR primer-probe
 <400> 157
 aacccggcga tcgaaaag 18
 5
 <210> 158
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> PCR primer-probe
 <400> 158
 15 ggcctgctg tcctgaga 18
 <210> 159
 <211> 24
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 25 <400> 159
 tcttaggaac gccgtaccag ccgc 24
 <210> 160
 <211> 19
 30 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 35 <400> 160
 ccaacgcttg ccaaactct 19
 <210> 161
 40 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 45 <223> PCR primer-probe
 <400> 161
 acggtagtga cagcatcaaa actc 24
 50 <210> 162
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 55 <220>
 <223> PCR primer-probe
 <400> 162

aaccagctct ctgtgacccc aatt 24

<210> 163
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

<400> 163
 gagaaccaat ctcaccgaca 20

<210> 164
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

<400> 164
 cacccgagtg taaccatagc 20

<210> 165
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

<400> 165
 acaggtattc ctctgccagc tgcc 24

<210> 166
 <211> 18
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

<400> 166
 gccgagatcg ccaagatg 18

<210> 167
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

<400> 167
 cttttgatgg tagagttcca gtgattc 27

<210> 168

<211> 24
 <212> DNA
 <213> Artificial Sequence

5 <220>
 <223> PCR primer-probe

<400> 168
 cagcattgtc tgcctccct ggca 24

10 <210> 169
 <211> 18
 <212> DNA
 <213> Artificial Sequence

15 <220>
 <223> PCR primer-probe

<400> 169
 20 gtgaggcagc gcgactct 18

<210> 170
 <211> 23
 <212> DNA
 25 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

30 <400> 170
 tgccaatggt gtacaacact tca 23

<210> 171
 <211> 21
 35 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

40 <400> 171
 tgccttcccg ggctgaggac t 21

<210> 172
 45 <211> 19
 <212> DNA
 <213> Artificial Sequence

<220>
 50 <223> PCR primer-probe

<400> 172
 ccaaccctgc agactccaa 19

55 <210> 173
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

5 <400> 173
 atgtataatg ttctgccaa ctgtatg 28

<210> 174
 <211> 25
 <212> DNA
 10 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

15 <400> 174
 cctgggacca tccgtggaga cttct 25

<210> 175
 <211> 20
 20 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

25 <400> 175
 ggctgtggct gaggctgtag 20

<210> 176
 <211> 21
 30 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

35 <400> 176
 ggagcattcg aggtcaaac a 21

<210> 177
 <211> 28
 40 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

45 <400> 177
 ttccagagt gtctcacctc cagcagag 28

50 <210> 178
 <211> 22
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> PCR primer-probe

<400> 178
 gaaggtgttg gaggcactca ag 22

5
 <210> 179
 <211> 21
 <212> DNA
 <213> Artificial Sequence

10
 <220>
 <223> PCR primer-probe

<400> 179
 ggtttacacc gctggagcta a 21

15
 <210> 180
 <211> 24
 <212> DNA
 <213> Artificial Sequence

20
 <220>
 <223> PCR primer-probe

<400> 180
 atcccagcag gcctcgttga tgag 24

25
 <210> 181
 <211> 20
 <212> DNA
 <213> Artificial Sequence

30
 <220>
 <223> PCR primer-probe

<400> 181
 gcatcaggct gtcattatgg 20

35
 <210> 182
 <211> 20
 <212> DNA
 <213> Artificial Sequence

40
 <220>
 <223> PCR primer-probe

45
 <400> 182
 agtagttgtg ctgcccttcc 20

50
 <210> 183
 <211> 28
 <212> DNA
 <213> Artificial Sequence

55
 <220>
 <223> PCR primer-probe

<400> 183
 tgtccttacc tgtgggagct gtaaggtc 28

<210> 184
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> PCR primer-probe
 <400> 184
 10 gggacggtgt tcacattcaa g 21
 <210> 185
 <211> 23
 <212> DNA
 15 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 20 <400> 185
 caggatccca gaagtcaatg ttg 23
 <210> 186
 <211> 24
 25 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 30 <400> 186
 tcgccagtct cccaactatc gcgt 24
 <210> 187
 <211> 28
 35 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 40 <400> 187
 ggctactctg atctatgttg ataaggaa 28
 <210> 188
 <211> 20
 45 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 50 <400> 188
 gcttcagccc atccttagca 20
 55 <210> 189
 <211> 22
 <212> DNA

<213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 5
 <400> 189
 cacacgggtg cctggtctc ca 22
 <210> 190
 10 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 15 <223> PCR primer-probe
 <400> 190
 ccattctatc atcaacgggt acaa 24
 20 <210> 191
 <211> 23
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> PCR primer-probe
 <400> 191
 30 tcagcaagtg ggaaggtga atc 23
 <210> 192
 <211> 25
 <212> DNA
 <213> Artificial Sequence
 35 <220>
 <223> PCR primer-probe
 <400> 192
 40 tctccacaga caaggccagg actcg 25
 <210> 193
 <211> 20
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 50 <400> 193
 ggctggtcgg cagagagtag 20
 <210> 194
 <211> 21
 55 <212> DNA
 <213> Artificial Sequence
 <220>

<223> PCR primer-probe
 <400> 194
 tgaggaaagg ttgggattg a 21
 5
 <210> 195
 <211> 29
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> PCR primer-probe
 <400> 195
 15 aaactcatgt aaaccacggc cgaatgttg 29
 <210> 196
 <211> 20
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 25 <400> 196
 cctgaacatg aaggagctga 20
 <210> 197
 <211> 19
 30 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> PCR primer-probe
 35 <400> 197
 catcacgtct ccgaactcc 19
 <210> 198
 40 <211> 21
 <212> DNA
 <213> Artificial Sequence
 <220>
 45 <223> PCR primer-probe
 <400> 198
 tccgatggg ctgcagcagc t 21
 50 <210> 199
 <211> 20
 <212> DNA
 <213> Artificial Sequence
 55 <220>
 <223> PCR primer-probe
 <400> 199

	catcttcag gaggaccact	20
	<210> 200	
	<211> 20	
5	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> PCR primer-probe	
10		
	<400> 200	
	tccgaccttc aatcatttca	20
	<210> 201	
15	<211> 24	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
20	<223> PCR primer-probe	
	<400> 201	
	ctctgtggca ccctggacta cctg	24
25	<210> 202	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> PCR primer-probe	
	<400> 202	
35	cctggaggct gcaacatacc	20
	<210> 203	
	<211> 23	
	<212> DNA	
40	<213> Artificial Sequence	
	<220>	
	<223> PCR primer-probe	
	<400> 203	
45	tacaatggct ttgaggata gca	23
	<210> 204	
	<211> 25	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
	<223> PCR primer-probe	
55	<400> 204	
	atcctcctga agcccttttc gcagc	25
	<210> 205	

<211> 20
 <212> DNA
 <213> Artificial Sequence

5 <220>
 <223> PCR primer-probe

<400> 205
 10 tgtttgatt cccgggctta 20

<210> 206
 <211> 24
 <212> DNA
 <213> Artificial Sequence

15 <220>
 <223> PCR primer-probe

<400> 206
 20 caaagctgtc agctctagca aaag 24

<210> 207
 <211> 28
 <212> DNA
 25 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

30 <400> 207
 tgccttcttc ctccctcact tctcacct 28

<210> 208
 <211> 20
 35 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

40 <400> 208
 gccaaactgct ttcattgtg 20

<210> 209
 45 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 50 <223> PCR primer-probe

<400> 209
 actcaggccc atttccttta 20

55 <210> 210
 <211> 28
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

5 <400> 210
 agggatctga accaatacag agcagaca 28

<210> 211
 <211> 20
 <212> DNA
 10 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

15 <400> 211
 aatccaaggg ggagagtgat 20

<210> 212
 <211> 20
 20 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

25 <400> 212
 gtacagattt tgcccgagga 20

<210> 213
 <211> 26
 30 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

35 <400> 213
 catatggact ttgactcagc tgtggc 26

40 <210> 214
 <211> 18
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <223> PCR primer-probe

<400> 214
 50 gcctcgggtg gccttca 18

<210> 215
 <211> 19
 <212> DNA
 55 <213> Artificial Sequence

<220>
 <223> PCR primer-probe

	<400> 215	
	cgatgatgtgc gcaatcatg	19
5	<210> 216	
	<211> 22	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> PCR primer-probe	
	<400> 216	
	catcgccagc tacgccctgc tc	22
15	1	

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 2002156263 A [0008]
- US 04809450 [0103]
- US 2004019567 W [0103]
- US 60482339 B [0103]

Non-patent literature cited in the description

- GOLUB et al. *Science*, 1999, vol. 286, 531-537 [0004]
- BHATTACHARJAE et al. *Proc. Natl. Acad. Sci. USA*, 2001, vol. 98, 13790-13795 [0004]
- CHEN-HSIANG et al. *Bioinformatics*, 2001, vol. 17 (1), S316-S322 [0004]
- RAMASWAMY et al. *Proc. Natl. Acad. Sci. USA*, 2001, vol. 98, 15149-15154 [0004]
- MARTIN et al. *Cancer Res.*, 2000, vol. 60, 2232-2238 [0004]
- WEST et al. *Proc. Natl. Acad. Sci. USA*, 2001, vol. 98, 11462-11467 [0004]
- SORLIE et al. *Proc. Natl. Acad. Sci. USA*, 2001, vol. 98, 10869-10874 [0004]
- YAN et al. *Cancer Res.*, 2001, vol. 61, 8375-8380 [0004]
- PEROU. *Nature*, 2000, vol. 406, 747-752 [0006]
- BOS et al. *Cancer*, 2003, vol. 97 (6), 1573-1518 [0008]
- SINGLETON et al. *Dictionary of Microbiology and Molecular Biology*. J. Wiley & Sons, 1994 [0034]
- Advanced Organic Chemistry Reactions. MARCH. Mechanisms and Structure. John Wiley & Sons, 1992 [0034]
- AUSUBEL et al. *Current Protocols in Molecular Biology*. Wiley Interscience Publishers, 1995 [0047]
- SAMBROOK et al. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Press, 1989 [0049]
- B. LEWIN. *Genes IV*. Cell Press, 1990 [0053]
- SAMBROOK et al. *Molecular Cloning: A Laboratory Manual*. 1989 [0054]
- Oligonucleotide Synthesis. 1984 [0054]
- Animal Cell Culture. 1987 [0054]
- Methods in Enzymology. Academic Press, Inc, [0054]
- Handbook of Experimental Immunology. Blackwell Science Inc, 1987 [0054]
- Gene Transfer Vectors for Mammalian Cells. 1987 [0054]
- Current Protocols in Molecular Biology. 1987 [0054]
- PCR: The Polymerase Chain Reaction. 1994 [0054]
- PARKER ; BARNES. *Methods in Molecular Biology*, 1999, vol. 106, 247-283 [0055]
- HOD. *Biotechniques*, 1992, vol. 13, 852-854 [0055]
- WEIS et al. *Trends in Genetics*, 1992, vol. 8, 263-264 [0055]
- AUSUBEL et al. *Current Protocols of Molecular Biology*. John Wiley and Sons, 1997 [0058]
- RUPP ; LOCKER. *Lab Invest.*, 1987, vol. 56, A67 [0058]
- DE ANDRÉS et al. *BioTechniques*, 1995, vol. 18, 42044 [0058]
- HELD et al. *Genome Research*, 1996, vol. 6, 986-994 [0064]
- T.E. GODFREY et al. *J. Molec. Diagnostics*, 2000, vol. 2, 84-91 [0065] [0075]
- K. SPECHT et al. *Am. J. Pathol.*, 2001, vol. 158, 419-29 [0065] [0075]
- DING ; CANTOR. *Proc. Natl. Acad. Sci. USA*, 2003, vol. 100, 3059-3064 [0066]
- LIANG ; PARDEE. *Science*, 1992, vol. 257, 967-971 [0067]
- KAWAMOTO et al. *Genome Res.*, 1999, vol. 12, 1305-1312 [0067]
- OLIPHANT et al. *Discovery of Markers for Disease (Supplement to Biotechniques)*, June 2002 [0067]
- FERGUSON et al. *Analytical Chemistry*, 2000, vol. 72, 5618 [0067]
- YANG et al. *Genome Res.*, 2001, vol. 11, 1888-1898 [0067]
- FUKUMURA et al. *Nucl. Acids. Res.*, 2003, vol. 31 (16), e94 [0067]
- SCHENA et al. *Proc. Natl. Acad. Sci. USA*, 1996, vol. 93 (2), 106-149 [0069]
- VELCULESCU et al. *Science*, 1995, vol. 270, 484-487 [0071]
- VELCULESCU et al. *Cell*, 1997, vol. 88, 243-51 [0071]
- BRENNER et al. *Nature Biotechnology*, 2000, vol. 18, 630-634 [0072]
- KENT, W.J. *Genome Res.*, 2002, vol. 12 (4), 656-64 [0077]

- Primer3 on the WWW for general users and for biologist programmers. **STEVE ROZEN ; HELEN J. SKALETSKY**. Bioinformatics Methods and Protocols: Methods in Molecular Biology. Humana Press, 2000, 365-386 **[0078]**
- General Concepts for PCR Primer Design. **DIEFFENBACH, C.W. et al**. PCR Primer, A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1995, 133-155 **[0080]**
- Optimization of PCRs. **INNIS ; GELFAND**. PCR Protocols, A Guide to Methods and Applications. CRC Press, 1994, 5-11 **[0080]**
- **PLASTERER, T.N**. Primerselect: Primer and probe design. *Methods Mol. Biol.*, 1997, vol. 70, 520-527 **[0080]**
- **COX, D. R. ; OAKES, D**. Analysis of Survival Data. Chapman and Hall, 1984 **[0094]**

P A T E N T K R A V

1. Fremgangsmåde til forudsigelse af sandsynligheden for sygdomsfri overlevelse af en patient med lymfeknude-negativ, ER-positiv, invasiv duktal brystcancer omfattende bestemmelse af ekspressionsniveauet af RNA-transskriptet af MELK i en brystcancercelle-
5 eller vævsprøve taget fra patienten, normaliseret mod ekspressionsniveauet af et referencesæt af RNA-transskripter, sammenligning af det normaliserede MELK-ekspressionsniveau fra patienten med et normaliseret MELK-ekspressionsniveau i et referencesæt af brystcancervæv omfattende patienter, der var (a) i live uden lokalt, regionalt eller fjernt brystcancerrecidiv, (b) i live med kontralateralt brystcancerrecidiv, (c) i live med en anden primær
10 ikkebryst-cancer, eller (d) døde før brystcancerrecidiv, hvor et øget normaliseret MELK-ekspressionsniveau fra patienten indikerer en nedsat sandsynlighed for sygdomsfri overlevelse.
2. Fremgangsmåde ifølge krav 1, hvor RNA'et omfatter intron-RNA.
3. Fremgangsmåde ifølge krav 1 eller 2, hvor RNA'et er isoleret fra en fikseret, voks-
15 indlejret brystcancervævsprøve fra patienten.
4. Fremgangsmåde ifølge krav 1, 2 eller 3, hvor RNA'et er isoleret fra kernebiopsivæv.
5. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor RNA'et er isoleret fra finnålsaspirerede celler.
- 20 6. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor ekspressionsniveauet af RNA-transskriptet bestemmes ved anvendelse af et array omfattende et polynukleotid, der hybridiserer med genet, som er immobiliseret på en fast overflade.
7. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor ekspressionsniveauet af RNA-transskriptet bestemmes ved revers transskription-
25 polymerasekædereaktion (RT-PCR).
8. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor RNA'et er fragmenteret.