



(43) International Publication Date
12 December 2013 (12.12.2013)

(10) International Publication Number
WO 2013/185021 A2

- (51) **International Patent Classification:**
C07F 7/08 (2006.01)
- (21) **International Application Number:**
PCT/US2013/044689
- (22) **International Filing Date:**
7 June 2013 (07.06.2013)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/657,309 8 June 2012 (08.06.2012) US
61/807,192 1 April 2013 (01.04.2013) US
- (71) **Applicant: THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA** [US/US]; 3160 Chestnut Street, Suite 200, Philadelphia, PA 19104 (US).
- (72) **Inventors; and**
- (71) **Applicants :** **HOYE, Adam, T.** [US/US]; 267 Rochelle Avenue, Apt. 1, Philadelphia, PA 19128 (US). **KIM, Won-suk** [KR/KR]; 168-31 Jangnan 1-dong, Dondaemungo, Seoul, 130-101 (KR). **MARTINEZ-SO-LORIO, Dionicio** [US/US]; 101 Conshohocken State Road, Apt. B4, Bala Cynwyd, PA 19004 (US). **SMITH, Amos, B.** [US/US]; 517 General Lafayette Road, Merion, PA 19066 (US). **SANCHEZ, Luis** [US/US]; 3160 Chestnut Street, Suite 200, Philadelphia, PA 19104 (US).
- (72) **Inventors: TONG, Rongbiao;** Tower 1,9b, Senior Staff Quarters, The Hong Kong University Of Science, And Technology, Clear Water Bay, Kowloon, Hong Kong
- (CN). **NGUYEN, Minh, Huu;** 4039 Chestnut Street, Apt. 9, Philadelphia, PA 19104 (US).
- (74) **Agents: CALDWELL, John, W** et al.; Woodcock Washburn LLP, Cira Centre, 12th Floor, 2929 Arch Street, Philadelphia, PA 19104 (US).
- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- Published:**
- *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*

(54) **Title:** SILICON-BASED CROSS COUPLING AGENTS AND METHODS OF THEIR USE

(57) **Abstract:** Compositions and methods using silicon-based cross-coupling agents in the formation of carbon-carbon and carbon-nitrogen bonds are described.



WO 2013/185021 A2

SILICON-BASED CROSS COUPLING AGENTS AND METHODS OF THEIR USE

REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 61/657,309, filed June 8, 2012 and U.S. Provisional Application No. 61/807,192, filed April 1, 2013, the entireties of which are incorporated by reference herein.

GOVERNMENT RIGHTS

[0002] At least a portion of this work was supported by a grant from the National Institutes of Health through Grant GM-29028. Pursuant to 35 U.S.C. § 202, the government may have certain rights in the invention.

TECHNICAL FIELD


[0003] The present invention is directed to silicon-based cross-coupling agents and methods of using them in cross-coupling reactions.

BACKGROUND

[0004] Cross-coupling reactions (“CCRs”) of organometallic/main group reagents with, for example, organic halides permit the construction of carbon-carbon and carbon-nitrogen bonds. Although CCRs are atom-efficient processes, the known CCRs have drawbacks, including the undesired formation of homo-coupled products and the use of toxic metals. As such, new methods for the cross coupling of organic compounds to form new carbon-carbon bonds and new carbon-nitrogen bonds are needed.

SUMMARY

[0005] The present invention is directed to compounds of formula I:



wherein Y is CH or N;

R₁ and R₂ are independently C₁₋₁₀ straight or branched-chain alkyl optionally substituted with one or more halogen, nitro, C₁₋₆alkoxy, or aryl;

R₃ is H;

aryl optionally substituted with one or more halogen, nitro, diC₁₋₆alkylamino;

C₁₋₆alkoxy, or C₁₋₆alkyl;

heteroaryl optionally substituted with one or more halogen, nitro, diC₁₋₆alkylamino;

C₁₋₆alkoxy, or C₁₋₆alkyl;

C₁₋₁₀ straight or branched-chain alkyl optionally substituted with one or more halogen, nitro, C₁₋₆alkoxy, or aryl;

a polymer; or

a resin support;


R_{3a} is H or C₁₋₆alkyl optionally substituted with one or more halogen; and

at least one R₄, wherein each R₄ is independently hydrogen, halogen, nitro, C₁₋₆alkoxy, C₁₋₆alkyl, aryl, or a resin support.

Use of the compounds of formula I as silicon-based cross-coupling agents in the formation of carbon-carbon and carbon-nitrogen bonds is also described.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0006] The present invention is directed to compounds of formula I for use as silicon-based cross-coupling agents in the formation of carbon-carbon and carbon-nitrogen bonds:



wherein

Y is CH or N;

R₁ and R₂ are independently C₁₋₁₀ straight or branched-chain alkyl optionally substituted with one or more halogen, nitro, C₁₋₆alkoxy, or aryl;

R₃ is H;

aryl optionally substituted with one or more halogen, nitro, diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl;

heteroaryl optionally substituted with one or more halogen, nitro, diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl;

C₁₋₁₀ straight or branched-chain alkyl optionally substituted with one or more halogen, nitro, C₁₋₆alkoxy, or aryl;


a polymer; or

a resin support;

R_{3a} is H or C₁₋₆alkyl optionally substituted with one or more halogen; and

at least one R₄, wherein each R₄ is independently hydrogen, halogen, nitro, C₁₋₆alkoxy, C₁₋₆alkyl, aryl, or a resin support.

In some embodiments of the invention, the compound of formula I is not



[0007] The reactivity of the compound of formula I can be attenuated by manipulation of R₁ and R₂. For example, in preferred embodiments, R₁ and R₂ are independently methyl, ethyl, *n*-propyl, or isopropyl. Preferably, R₁ and R₂ are independently methyl or isopropyl. In exemplary embodiments, R₁ and R₂ are each methyl. In other embodiments, R₁ and R₂ are each

isopropyl. In still other embodiments, R₁ and R₂ are each ethyl. In yet other embodiment, R₁ and R₂ can be optionally substituted with one or more halogen, nitro, C₁₋₆alkoxy, or aryl.

[0008] The reactivity of the compound of formula I can be attenuated by manipulation of R₃. For example, in some embodiments, R₃ is C₁₋₁₀, preferably C₁₋₆, straight or branched-chain alkyl optionally substituted with one or more halogen or C₁₋₆alkoxy. Preferably, R₃ is C₁₋₄ straight or branched-chain alkyl optionally substituted with one or more halogen or C₁₋₆alkoxy. More preferably, R₃ is C₁₋₄ straight or branched-chain alkyl. In exemplary embodiments, R₃ is *n*-butyl, isobutyl, *sec*-butyl, or *tert*-butyl. In most preferred embodiments, R₃ is *n*-butyl. In those embodiments wherein R₃ is C₁₋₆alkyl substituted with one or more halogen, R₃ is preferably CF₃.

[0009] In other embodiments, R₃ is aryl, preferably phenyl, optionally substituted with one or more halogen, nitro, diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl. Preferably, R₃ is phenyl optionally substituted with one or more diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl. In exemplary embodiments, R₃ is phenyl.

[0010] In still other embodiments, R₃ is heteroaryl, preferably pyridyl, optionally substituted with one or more halogen, nitro, diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl. Preferably, R₃ is pyridyl optionally substituted with one or more diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl. In exemplary embodiments, R₃ is pyridyl.

[0011] In other preferred embodiments, R₃ is a polymer. Preferably, this polymer acts as a solid support for the siloxane transfer agent. Such polymers are known in the art *per se*. Preferred polymers are described herein.

[0012] In still other embodiments, R₃ is a resin support.


[0013] In preferred embodiments of the invention, R_{3a} is H or C₁₋₆alkyl optionally substituted with one or more halogen. Preferably, R_{3a} is H. In those embodiments wherein R_{3a} is C₁₋₆alkyl, exemplary moieties include methyl and ethyl. In other embodiments wherein R_{3a} is C₁₋₆alkyl substituted with one or more halogen, R_{3a} is preferably CF₃.


[0014] The reactivity of the compound of formula I can be attenuated by manipulation of R₄. The compounds of formula I can include one, two, three, or four R₄ groups, which can each be the same or different. R₄ is preferably hydrogen, halogen, nitro, C₁₋₆alkoxy, C₁₋₆alkyl, or aryl. More preferably, R₄ is hydrogen, halogen, for example F, C₁₋₆alkoxy, or C₁₋₆alkyl. In exemplary embodiments, R₄ is hydrogen. In other embodiments, R₄ is halogen, for example, F, Cl, or Br, with F being particularly preferred. R₄ can also be a resin support.

[0015] In those embodiments wherein the compound of formula I include a resin support, the resin support is at either R₃ or R₄. That is, in certain embodiments wherein R₃ is a


resin support, each R_4 is independently hydrogen, halogen, nitro, C_{1-6} alkoxy, C_{1-6} alkyl, or aryl. In those embodiments wherein R_4 is a resin support, R_3 is H; aryl optionally substituted with one or more halogen, nitro, diC_{1-6} alkylamino, C_{1-6} alkoxy, or C_{1-6} alkyl; heteroaryl optionally substituted with one or more halogen, nitro, diC_{1-6} alkylamino, C_{1-6} alkoxy, or C_{1-6} alkyl; C_{1-10} straight or branched-chain alkyl optionally substituted with one or more halogen, nitro, C_{1-6} alkoxy, or aryl; or a polymer.

[0016] It is preferred that the aromatic moiety of the compounds of formula I is phenyl, that is, wherein Y is CH or CR_4 . In such embodiments, preferred compounds of formula I include, for example:






“polymer-supported siloxane transfer agent (saturated),”
for example, having 20 to 200 repeating units,




[0017] Exemplary embodiments of the invention wherein R_3 is a polymer include compounds of the formula:




wherein n is independently about 20 to 300, preferably 20 to 200, or 150 to about 300. Most preferably, n is about 200 or 250.


[0018] Exemplary embodiments of the invention wherein R_3 or R_4 is a resin support include:



[0019] In other embodiments, the aromatic moiety of the compounds of formula I is pyridyl, that is, wherein Y is N.



[0020] Exemplary embodiments of the invention wherein Y is N are:



[0021] Based on its physico-chemical properties, a compound of formula I including a pyridyl group may facilitate removal of the cross-coupling transfer agent from the products of the cross-coupling reaction mixture by treatment of the crude reaction mixture with Brønsted or Lewis acids upon workup.

[0022] In accordance to the present invention, and as described above and below, the siloxane motif maybe incorporated into polymers using monomers derived from the silicon transfer agent. This polymeric material may improve the ease of purification following the cross-coupling reaction, and may make the transfer agent easier to handle by altering its physical properties.

[0023] The invention is also directed to methods of cross-coupling compounds of formula NuLi with a compound of formula E-X to form a compound of formula Nu-E. These methods comprise contacting the compound of formula NuLi with the compound of formula E-X in the presence of a siloxane compound of the invention as described herein, a catalyst or a catalyst system, and an ethereal solvent, for a time and under conditions sufficient to produce the compound of formula Nu-E. In these methods, Nu is an aryl compound or an alkenyl compound; E is an aryl compound or an alkenyl compound; and X is iodo or bromo or X is iodo, chloro, or bromo.

[0024] The invention is also directed to methods of cross-coupling compounds of formula Nu-Li with a compound of formula E-X to form a compound of formula Nu-E. These methods comprise contacting the compound of formula NuLi with the compound of formula E-x in the presence of a siloxane compound of the invention as described herein, a catalyst or a catalyst system, and an ethereal solvent, for a time and under conditions sufficient to produce the compound of formula Nu-E. In these methods, Nu is an aryl compound or an alkenyl compound; E is a disubstituted amine, and X is -O-benzoyl.


[0025] As used herein, “aryl compound” refers to an organic compound comprising a phenyl or naphthyl group that is optionally substituted with one or more substituents. Examples of substituents include alkyl, aryl, halogen, nitro, cyano, ester, alkoxy, siloxy, and the like.

[0026] As used herein, “alkenyl compound” refers to an organic compound comprising a carbon-carbon double bond that is optionally substituted with one or more substituents. Examples of substituents include alkyl, aryl, halogen, nitro, cyano, ester, alkoxy, siloxy, and the like.

[0027] As used herein, “C₁₋₁₀ straight or branched-chain alkyl” refers to an aliphatic hydrocarbon including from 1 to 10 carbon atoms. Examples include methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, tert-butyl, n-pentyl, iso-pentyl, n-hexyl, and the like.

[0028] As used herein, “diC₁₋₆alkylamino” refers to an amino moiety substituted with two C₁₋₆alkyl group. The C₁₋₆alkyl groups can be the same or different. Preferred diC₁₋₆alkylamino groups for use in the invention include dimethylamino, diethylamino, methylethylamino, diisopropylamino, and the like.

[0029] As used herein, “disubstituted amine” refers to a nitrogen atom having two substituents, which are the same or different. Disubstituted amine also refers to compounds wherein the two substituents are joined to form a ring. Examples of disubstituted amines include



[0030] As used herein, “aryl” refers to an aromatic 6-13 membered mono- or bi-cyclic ring such as phenyl or naphthyl.

[0031] As used herein, “halogen” refers to iodo, bromo, chloro, or fluoro.


[0032] As used herein, “heteroaryl” refers to a mono-or bicyclic aromatic ring structure including carbon atoms as well as up to four heteroatoms selected from nitrogen, oxygen, and sulfur. Heteroaryl rings can include a total of 5, 6, 9, or 10 ring atoms. Preferred heteroaryl groups include pyridyl and pyrimidinyl.

[0033] As used herein, “resin support” refers to solid supports used in, for example, combinatorial chemistry. “Wang resins” and “Merrifield resin” are examples of such resins. Resin supports and their use and incorporation are known to those skilled in the art, *per se*.

[0034] The compounds of formula I can be used in organic synthesis, most preferably as silicon-based cross-coupling agents. In exemplary embodiments, the compound of formula I is used in the presence of, for example, an organo-lithium compound and an organo-halogen compound to form the resulting cross-coupled organic compound. *See, e.g.*, Scheme 1 for an example using one embodiment of the invention. Preferably, the organo-lithium compound is an aryl-lithium or alkenyl-lithium compound. The organo-halogen compound is preferably an aryl-halogen compound or an alkenyl-halogen compound. In such embodiments, the halogen is

preferably iodo, chloro, or bromo. The resulting cross-coupled organic compounds are thus aryl-aryl compounds, aryl-alkenyl, or alkenyl-alkenyl compounds.




Scheme 1



[0035] In such cross-coupling reactions, the compounds of formula I can be used in stoichiometric amounts, that is, one molar equivalent, as compared to either the organo-lithium or organo-halogen compound. In some embodiments, the compounds of formula I are used in greater than stoichiometric amounts, that is, greater than one molar equivalent as compared to either the organo-lithium or organo-halogen compound. In such embodiments, up to three molar equivalents of the compound of formula I, as compared to either the organo-lithium or organo-halogen compound, can be used. Within the scope of the invention, it is envisioned that 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.8, 2.0, 2.2, 2.5, or 2.8 molar equivalents of the compound of formula I, as compared to either the organo-lithium or organo-halogen compound, can be used. It is preferred to use 1.6, 1.8, or 2.0 molar equivalents of the compound of formula I. Most preferred is 1.8 molar equivalents of the compound of formula I. The amount of the compound of formula I required for cross-coupling a particular organo-lithium compound with a particular organo-halogen compound can be determined without undue experimentation by someone of skill in the art.

[0036] Alternatively, the compound of formula I can be used in catalytic amounts, that is, less than one molar equivalent, as compared to either the organo-lithium or organo-halogen compound.

[0037] In some embodiments, cross-coupling of an organo-lithium compound with an organo-halogen compound in the presence of the silicon-based cross-coupling agent compounds of formula I of the invention requires the use of a catalyst or catalyst system. Preferred catalysts or catalyst systems comprise palladium compounds, copper compounds, nickel compounds, or mixtures thereof, optionally in the presence of a compound such as cyclohexyl-(2-diphenylphosphanyl-benzylidene)-amine (dpca), XPhos (2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl), or Johnphos ((2-biphenyl)di-*tert*-butylphosphine):



[0038] Preferably, the palladium compound is PdCl_2 , $\text{Pd}(\text{PPh}_3)_4$, or $\text{Pd}(\text{OAc})_2$. The copper compound is preferably a copper halide, for example, CuI . A preferred catalyst system for use in the methods of the invention comprises PdCl_2 , CuI , and dpca. Another preferred catalyst system includes $\text{Pd}(\text{OAc})_2$ and XPhos. Yet another catalyst system includes CuI and dpca. Another catalyst system include CuI and Johnphos.

[0039] Those of skill in the art can readily determine the mole percentage of the catalyst or each component of the catalyst system for use in the methods of the invention. For example, the catalyst or component of the catalyst system can be present in an amount of from about 1 mol% to about 15 mol%. Alternatively, the catalyst or component of the catalyst system can be present in an amount of from about 3 mol% to about 10 mol%.

[0040] In embodiments incorporating the use of a palladium compound as the catalyst or component of the catalyst system, the palladium compound is present in an amount of from about 1 mol% to about 5 mol% or to about 10 mol%. Preferably, the palladium compound, for example, PdCl_2 , is present at about 3 mol%.

[0041] In embodiments incorporating the use of a copper compound as the catalyst or component of the catalyst system, the copper compound is present in an amount of from about 0.1 mol% to about 10 mol%. In other embodiments, the copper compound is present in an amount of from about 1 mol% to about 10 mol%. Alternatively, the copper compound is present in an amount of from about 5 mol% to about 12 mol%. More preferably, the copper compound is present in an amount of from about 9 mol% to about 11 mol%. Most preferably, for example when the copper compound is CuI , it is present at about 10 mol%.

[0042] In those embodiments incorporating dpca in the catalyst systems of the invention, the dpca is present from about 1 mol% to about 10 mol%. Preferably, the dpca is

present in an amount from about 2 mol% to about 8 mol%. Particularly preferred are embodiments wherein the dpca is present at about 4 mol%. In those embodiments employing dpca, the dpca is used in conjunction with PdCl₂. Dpca can also be used in conjunction with CuI. Preferably, the dpca is present in an amount greater than the amount of PdCl₂ or CuI. For example, the dpca is present at about 4 mol% and the PdCl₂ is present at about 3 mol%.

[0043] In those embodiments incorporating XPhos in the catalyst systems of the invention, the XPhos is present from about 1 mol% to about 50 mol%. Preferably, the XPhos is present in an amount from about 10 mol% to about 30 mol%. Particularly preferred are embodiments wherein the XPhos is present at about 20 mol%.


[0044] In embodiments incorporating Johnphos in the catalyst systems of the invention, the Johnphos is present from about 1 mol% to about 20 mol%. Preferably, the XPhos is present in an amount from about 5 mol% to about 15 mol%. Particularly preferred are embodiments wherein the XPhos is present at about 10 mol%.


[0045] Preferred solvents for use in the cross-coupling reactions of the invention include ethereal solvents, for example tetrahydrofuran (THF), tetrahydropyran (THP), diisopropyl ether and diethyl ether, with THF being particularly preferred.

[0046] Siloxanes of the invention can be prepared according to the methods described herein. Two exemplary methods are shown in Schemes 2A and 2B. Other siloxanes within the scope of the invention can be prepared similarly.

Scheme 2

A. Two-Pot Synthesis of Siloxane **1** from Benzaldehyde.



B. One-Pot Synthesis of Siloxane **2a** from Benzaldehyde.

[0047] Preferred compounds of formula I have been successfully employed to cross-couple a variety of organo-lithium and organo-halogen compounds, while limiting the amount of homo-coupled product. For example, Scheme 3 depicts the results of several experiments where the organo-halogen compound was 4-iodoanisole and the organo-lithium compound was phenyl lithium.

Scheme 3^a

Entry	Solvent	Equiv. 1	Equiv. PhLi	Time (step a)	Time (step b)	¹ H NMR Results (A:B:C) ^b
1	THF	2.0	2.1	0.5 h	2 h	13.4 : 1 : nd
2	THF	1.2	1.1	0.5 h	0.5 h	7.2 : 1.0 : 4.0
3	THF	1.6	1.5	1 h	2 h	17.8 : 1.0 : 2.2
4 ^c	THF	1.6	1.5	1 h	2 h	>20 : 1.0 : 2.8
5 ^c	THF	2.0	1.5	1 h	2 h	>20 : nd : nd
6 ^c	THF	2.0	1.2	1 h	2 h	3.4 : nd : 1.0
7 ^c	THF	--	1.5	1 h	2 h	2.5 : 0.6 : 1.0 ^d
8 ^c	THF	1.8	1.5	1 h	2 h	>20 : nd : nd

^a All reactions were performed on 0.45 mmol scale with 4-iodoanisole as the limiting reagent.

^b determined by ¹H NMR analysis of the crude mixture of reaction products following aqueous workup and extraction with Et₂O. nd = not detected


^c PdCl₂ (3 mol %), dpca (4 mol %), and CuI (10 mol %) were premixed for 30 minutes in THF prior to the addition of 4-iodoanisole, after which the PhLi and **1** reaction mixture was introduced via cannula.







^d No 1-oxa-2-silacyclopentene was used in the reaction; a solution of PhLi in THF was used as a substitute.

[0048] Referring to Scheme 3, compound **1** is a compound of formula I within the scope of the invention. Good conversion of the aryl iodide occurred when steps a and b were allowed to proceed for longer times (see entry 3). A significant increase in the efficiency of the process was observed when the catalyst system of PdCl₂, dpca, and CuI was premixed for about 30 minutes in THF at room temperature prior to introduction of the aryl halide, which was followed by immediate addition of a mixture of PhLi and **1** in THF (see entry 4). Using this protocol in conjunction with the use of 2.0 molar equivalents of compound **1** led to complete conversion of 4-iodoanisole with no detectable homocoupled product. Notably, significant homocoupling and catalyst decomposition was observed in the absence of silicon cross coupling agent **1** (see entry 7). The best results for these reagents, as shown in Scheme 3, resulted using 1.8 molar equivalents of compound **1** and 1.5 molar equivalents of PhLi.

[0049] As shown in Scheme 4, organo-halides other than 4-iodoanisole can be used in cross-coupling reactions of the invention using the silicon cross-coupling agents of the present invention, *i.e.*, compounds of formula I.

Scheme 4



2	 9b	92% (10b)	6	 9g	85% (10e)
3 ^b	 9c	92% (10b)	7 ^b	 9h	95% (10f)
4	 9e	75% (10c)	8	 9i	67% (10g)

^a Isolated yields.

^b The reaction was allowed to proceed for 12 h following addition of the PhLi.

[0050] As demonstrated in Scheme 4, electron-rich and electron-deficient substrates were well-tolerated in the reaction, as were a variety of common functional groups, for example, ester, nitriles, and azoheterocycles, to provide the biaryl compounds **10a-g** in good yields. In all cases, silicon cross-coupling agent **1**, which is a compound of formula I of the invention, emerged from the reaction intact, as observed by ¹H NMR analysis of the crude reaction mixtures. For products possessing polarities similar to that of compound **1**, an oxidative Tamao-Fleming workup (Simmons, E.M. et al., J. Am. Chem. Soc. 2010, 132, 17092; Tamao, K. et al. Organometallics 1983, 2, 1694; Fleming, I. et al., J. Chem. Soc. Chem. Commun. 1984, 29) of the initial reaction mixture was used to convert **1** to the corresponding diol, which could be easily separated by column chromatography.

Scheme 5

Nu-Li (1.5 equiv) $\text{THF, -78 } ^\circ\text{C to rt, 1 h}$ <i>via cannula</i> rt Nu-E $\text{PdCl}_2 \text{ (3 mol\%)} + \text{dpca (4 mol\%)} + \text{Cul (10 mol\%)}$ THF, rt, 30 min E-X (1.0 equiv) then rt Nu-E				
	Nu-Li	E-X	Product (Nu-E)	Yield ^a
1				81%
2			14a	82% (gram-scale)
3	13a			84%
4		9a	14b	82%
5	13a			79%
6		9a	14c	77%
7		9a		67%
8		9a		81%
9 ^b	13f			72%

	Nu-Li	E-X	Product (Nu-E)	Yield^a
10	 OTBS 13g	 DMBO (+)-9n	 OTBS (+)-14g	78%

^a isolated yields.

^b The crude product mixture was treated with TBAF in THF to remove the silyl group prior to purification.

[0051] The methods of the claimed invention can also be extended to alkenyl substrates. As shown in Scheme 5 above, vinyl halides **9j-l** were good coupling partners with **13a**, forming styrenes **14a-c** with retention of the alkene geometry (see entries 1, 3, and 5). Notably, the roles of the coupling partners could be reversed by using the corresponding vinylolithium and aryl halide to access identical coupling products in comparable yields (see entries 2, 4, and 6), demonstrating the flexibility of the methods of the invention with respect to the choice of nucleophilic and electrophilic components. Geminal and vicinal substitution patterns were tolerated in the cross coupling process from silicon, providing coupled products **14c** and **14d**. Also noteworthy was the successful vinyl-vinyl couplings of **13f** and **9m** and between **13g** and **(+)-9n** to provide dienes **14f** and **(+)-14g**.

[0052] Scheme 6 details examples demonstrating the scope and utility of the methods of the invention using preferred embodiments of compounds of formula I as silicon cross coupling agents.

Scheme 6


Entry	Siloxane	Nu-Li	E-X	Catalyst System (Pd, ligand, Cu)	Product	Isolated Yield
1		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		96%
2		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		92%
3		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		92%
4		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		75%
5		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		76%
6		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		85%
7		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		95%
8		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		67%
9				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		81%
10				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		82%
11				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		84%
12				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		82%
13				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		79%

Entry	Siloxane	Nu-Li	E-X	Catalyst System (Pd, ligand, Cu)	Product	Isolated Yield
14				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		77%
15				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		67%
16				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		81%
17				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		96%
18				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		72%
19				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		78%
20		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		--
21		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		--
22				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		93%
23				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		96%
24				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		84%
25				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		80%
26				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		81%

Entry	Siloxane	Nu-Li	E-X	Catalyst System (Pd, ligand, Cu)	Product	Isolated Yield
27				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		74%
28		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		84%
29				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		96%
30				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		83%
31				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		87%
32				PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		80%
33		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		69%
34		PhLi		PdCl ₂ (3 mol%) dpca (4 mol%) CuI (10 mol%)		79%

[0053] As can be seen in Scheme 7, the synthetic utility of the cross-coupling method may be expanded to include the use of sp³-hybridized organolithium species and alkyl halides. By employing Pd- or Ni-catalysis, it may be possible access sp²-sp³ coupled products. Additionally, sp³-sp³ cross-couplings involving secondary alkyl halides may also be possible. It is preferred, in some examples, that iso-propyl group or groups are present on the silicon in order to avoid competitive transfer of primary alkyl groups from the activated silicon species (*e.g.*, methyl).






Scheme 7











[0054] Additional examples of cross-coupling reactions performed using siloxanes of the invention are set forth in Scheme 8.

Scheme 8

Siloxane (1.8 equiv) $\xrightarrow[\text{solvent (THF or diethyl ether), rt, 2 h}]{\text{PhLi (1.5 equiv)}}$ $\xrightarrow[\text{via cannula, rt, 2 h}]{\text{THF, rt, 30 min}}$ PdCl_2 (3 mol%) + dpca (4 mol%) + Cul (10 mol%) $\xrightarrow[\text{THF, rt, 30 min}]{\text{then}}$ $\text{I-C}_6\text{H}_4\text{-OMe}$ (1.0 equiv) Products: A (1,4-dimethoxybiphenyl), B (1,3-dimethoxybiphenyl), and C (4-methoxyiodobenzene).				
Entry	Siloxane	$^1\text{H NMR}$ Ratio ^b A:B:C	Siloxane Recovery (%)	Yield (A) (%) ^d
1	 (1)	>20 : -- : --	<i>e</i>	96
2	 (1a)	>20 : -- : --	<i>e</i>	92

Siloxane (1.8 equiv) $\xrightarrow[\text{solvent (THF or diethyl ether), rt, 2 h}]{\text{PhLi (1.5 equiv)}}$ $\xrightarrow[\text{THF, rt, 30 min}]{\text{PdCl}_2 \text{ (3 mol\%)} + \text{dpca (4 mol\%)} + \text{CuI (10 mol\%)}}$ $\xrightarrow[\text{rt, 2 h}]{\text{then, via cannula}}$ + $\text{I}-\text{C}_6\text{H}_4-\text{OMe}$ (1.0 equiv) $\rightarrow$ A + B + C				
Entry	Siloxane	$^1\text{H NMR}$ Ratio ^b A:B:C	Siloxane Recovery (%)	Yield (A) (%) ^d
3	 (1e)	13 : 1.0 : --	<i>e</i>	63 ¹
4	 (1f)	1.7 : 1.2 : 1.0	<i>e</i>	<i>f</i>
5	 (2c)	2.0 : 1.0 : : --	92 ^c	42
6	 (2d)	2.5 : 1.0 : : --	94 ^c	44
7	 (2a)	>20 : -- : --	81 ^c	96

Siloxane (1.8 equiv) $\xrightarrow[\text{solvent (THF or diethyl ether), rt, 2 h}]{\text{PhLi (1.5 equiv)}}$ $\xrightarrow[\text{THF, rt, 30 min}]{\text{PdCl}_2 \text{ (3 mol\%)} + \text{dpca (4 mol\%)} + \text{CuI (10 mol\%)}}$ $\xrightarrow[\text{then, rt, 2 h}]{\text{I-C}_6\text{H}_4\text{-OMe (1.0 equiv)}}$ $\xrightarrow[\text{via cannula}]{}$ + $\text{C}_6\text{H}_5\text{-OMe}$ + $\text{C}_6\text{H}_4\text{(OMe)-I}$ + $\text{C}_6\text{H}_4\text{(OMe)-OMe}$ A B C				
Entry	Siloxane	¹ H NMR Ratio ^b A:B:C	Siloxane Recovery (%)	Yield (A) (%) ^d
8	 (2b)	>20 : -- : --	85 ^c	98
9	 (3a)	>20 : -- : --	<i>e</i>	98
10	 (3b)	>20 : -- : --	<i>e</i>	98
11	 (3c)	>20 : -- : --	67 ^x	92
12	 (3d)	>20 : -- : --	74 ^d	94

Siloxane (1.8 equiv) $\xrightarrow[\text{solvent (THF or diethyl ether), rt, 2 h}]{\text{PhLi (1.5 equiv)}}$ $\xrightarrow[\text{THF, rt, 30 min}]{\text{PdCl}_2 \text{ (3 mol\%)} + \text{dpca (4 mol\%)} + \text{CuI (10 mol\%)}}$ $\xrightarrow[\text{rt, 2 h}]{\text{then 4-iodoanisole (1.0 equiv)}}$ $\xrightarrow{\text{via cannula}}$ Products: A (4-methoxybiphenyl), B (4-methoxy-1-phenyl-4-methoxybenzene), C (4-iodoanisole)				
Entry	Siloxane	¹ H NMR Ratio ^b A:B:C	Siloxane Recovery (%)	Yield (A) (%) ^d
13	 (3e)	>20 : -- : --	96 ^d	98
14	 (3f)	>20 : -- : --	76 ^d	96
15	 (3g)	>20 : -- : --	96 ^d	98

^aAll reactions were performed on 0.45 mmol scale with 4-iodoanisole as the limiting reagent.


^bRequires heating to 50 °C to participate in the cross-coupling reaction.

^cDetermined by ¹H NMR analysis of the crude reaction mixture.

^dRecovered via column chromatography.


[0055] Methods of the invention can also be accomplished without copper in the catalyst system. In such embodiments, the catalyst system comprises a palladium-based catalysts, for example, PdCl₂ or Pd(OAc)₂. These embodiments can also employ the use of, for example, XPhos to form a palladium complex catalyst system. An example of such a method is set forth in Scheme 9.

Scheme 9



[0056] Polymeric siloxane compounds are also within the scope of the invention. As can be seen in Scheme 10 (wherein *n* is about 150 to about 300, preferably about 200 or 250), the siloxane motif may be incorporated into a polymer *via* ring-opening metathesis polymerization (ROMP). For example, according to Scheme 10, treatment of commercially available 5-norbornene-2-carboxaldehyde (a mixture of endo- and exo-isomers) with phenylmagnesium bromide generated the benzylic alcohol. Subsequent ortho-lithiation with *n*-BuLi, followed by trapping with Me₂SiHCl and treatment with H₂O provided a mixture of benzyl alcohol and siloxane (observed by ¹H-NMR), which was treated with catalytic K⁺OtBu to furnish complete conversion to the desired siloxane monomer. This sequence can be performed on multigram-scale. ROMP (ring opening metathesis polymerization) of this monomer was then achieved smoothly with the first generation Grubbs catalyst.

Scheme 10




[0057] The polymer can be obtained in near quantitative yield without using cross-linking units or co-polymerization agents. As such, the loading of the polymer with siloxane units should be nearly identical to the molarity of the monomer, *e.g.*, 3.9 mmol/g for the example set forth in Scheme 9, with each polymer chain having a relative length of 200-mers. The number of repeating siloxane units on each polymer chain can be adjusted by changing the amount of Grubbs catalyst used during the polymerization process.

[0058] These polymers of the invention are generally soluble in common organic solvents such as those used in the cross-coupling reaction.

[0059] The silicon transfer reagent may also be incorporated into a solid support, for example a resin, using "click chemistry," which is understood by those in the art to include, for example, the reaction of an alkyne and an azide to produce a 1,2,3-triazole. *See* Scheme 11. In preferred embodiments of the invention, the reaction is between an alkyne-capped resin, such as those known in the art, and an azido-siloxane, which can be prepared according to known methods. (Scheme 11). In these embodiments, the compounds of formula I can be tethered to a resin via a triazolyl linker. These embodiments of the inventions may facilitate removed of the


transfer reagent from the reaction mixture by mechanical means such as filtration. Additionally, the transfer agent may be regenerated and reused.

Scheme 11



[0060] To further exemplify the invention, STA-I₂₀₀, one embodiment of the invention, was used to mediate the cross-coupling of phenyl lithium and 1-iodoanisole. See Scheme 12.

Scheme 12



Entry	Equiv. PhLi	Equiv STA-I ₂₀₀	Conc. STA-I ₂₀₀	Time	¹ H-NMR Results ^b (B : A : C)
1	1.5	2.0	15 mg/mL	25 h	57 : 40 : 3
2 ^d	2.5	3.0	15 mg/mL	15 h	90 : 5 : 5
3 ^e	2.5	3.0	10 mg/mL	2 h	100 : nd : <1
4	1.5	3.0	10 mg/mL	15 h	84 : 11 : 5
5	2.0	3.0	10 mg/mL	2 h	72 : 18 : <1
6	2.0	2.5	10 mg/mL	2 h	88 : 9 : 3
7 ^f	2.5	--	--	2 h	14 : 77 : 9 ^g

^aAll reactions were performed on 0.3 mmol scale in THF with 4-iodoanisole as the limiting reagent.

^bDetermined by ¹H-NMR analysis of crude mixture of reaction products following aqueous workup, extraction with Et₂O and removal of recovered polymer via precipitation in acetonitrile.

^cPart of recovered polymer remained insoluble in THF.

^dSmall amount of recovered polymer remained insoluble in THF; 82% isolated yield of **B**.

^ePolymer was recovered quantitatively and re-useable; 98% isolated yield of **B**.


^fNo siloxane polymer was used in the reaction; a solution of PhLi in THF was used as a substitute.




















^gMixture of other products was also observed.

nd = not detected.

[0061] The polymeric siloxanes of the invention have utility in, for example, the cross-coupling between any organolithiums and aryl or alkenyl iodides. Examples of such cross-coupling reactions are set forth in Scheme 13.


Scheme 13



Entry	Nu-Li	E-X	Time	Product (Nu-E)	Yield
1	PhLi		2 h		98%
2	PhLi		2 h		91%
3	PhLi		2 h		93%
4	PhLi		24 h		7%
5	PhLi		12 h		89%
6			2 h		98%
7			2 h		71%
8			17 h		70%

Entry	Nu-Li	E-X	Time	Product (Nu-E)	Yield
9	TBSO-CH ₂ -CH=CH-Li	I-CH=CH-CH ₂ -CH ₂ -Ph	15 h	HO-CH ₂ -CH=CH-CH=CH-CH ₂ -CH ₂ -Ph	72%

[0062] Polymeric siloxanes of the invention are also recyclable and retain activity through multiple cross-coupling reaction cycles, using the same nucleophile for each transformation. *See, e.g.*, Scheme 14.



Cycle	Isolated yield of B	Polymer Recovered ^a	¹ H-NMR Results (B : A : C)
0		(Mn = 63500, PDI = 1.3)	
1 st	98%	98% (Mn = 73000, PDI = 1.6)	100 : nd : <1
2 nd	91%	99% (Mn = 104400, PDI = 2.3)	97 : nd : 3
3 rd	81%	99% (Mn = 94600, PDI = 2.6)	86 : 11 : 3


^aPoly(methyl methacrylate) standards were used to determine Mn and PDI values.











Mn = number average molecule weight.

PDI = polydispersity index.

[0063] Polymeric siloxanes of the invention can also transfer different nucleophiles in repeated cycles. *See, e.g.*, Scheme 15.

Scheme 15



Cycle	Nu-Li	¹ H-NMR Ratio of Products ^a		
1 st	<chem>C5H11CH=CHLi</chem>	 78	 20	 2
2 nd	PhLi	 91	 5	 4
3 rd	PhLi	 75	 22	 2  1

^aDetermined by ¹H-NMR analysis of crude mixture of reaction products following aqueous workup, extraction with Et₂O and removal of recovered polymer via precipitation in acetonitrile.

[0064] Also within the scope of the invention are methods of forming carbon-nitrogen bonds using the siloxanes of the invention. Examples of such methods are set forth in Scheme 16.

Scheme 16

Entry	R-Li	(R') ₂ N-OBz	R-N(R') ₂ Yield	Recovered siloxane
1	PhLi		98%	--
2	PhLi		91%	85%
3	PhLi		93%	87%
4	PhLi		65% (using 10% CuI)	75%
5	PhLi		83% (using 20% CuI)	86%
6			73%	80%
7			40%	--


[0065] The following examples are only illustrative and are not meant to limit the invention. Those skilled in the art will be able to practice the full breadth of the invention in view of the following examples, this specification, and the knowledge of the art.

EXAMPLES

[0066] General. All moisture-sensitive reactions were performed using syringe-septum cap techniques under an inert atmosphere of N₂. All glassware was flame dried or dried in an oven (140 °C) for at least 4 h prior to use. Reactions were magnetically stirred unless otherwise stated. Tetrahydrofuran (THF), dichloromethane (CH₂Cl₂) and diethyl ether (Et₂O) were dried by passage through alumina in a Pure Solve™ PS-400 solvent purification system. Unless otherwise stated, solvents and reagents were used as received. Analytical thin layer chromatography was performed on pre-coated silica gel 60 F-254 plates (particle size 40-55 micron, 230-400 mesh) and visualized by a uv lamp or by staining with PMA (2 g phosphomolybdic acid dissolved in 20 mL absolute ethanol), KMnO₄ (1.5 g of KMnO₄, 10 g of K₂CO₃ and 2.5 mL of 5% aq. NaOH in 150 mL H₂O), or CAM (4.8 g of (NH₄)₆Mo₇O₂₄·4H₂O and 0.2 g of Ce(SO₄)₂ in 100 mL of a 3.5 N H₂SO₄ solution) stain. Column chromatography was performed using silica gel (Silacyle Silaflash® P60, 40-63 micron particle size, 230-300 mesh) and compressed air pressure with commercial grade solvents. Yields refer to chromatographically and spectroscopically pure compounds, unless otherwise stated. NMR spectra were recorded at 500 MHz/125 MHz (¹H NMR/¹³C NMR) on a Bruker Avance III 500 MHz spectrometer at 300 K. Chemical shifts are reported in parts per million with the residual solvent peak as an internal standard. ¹H NMR spectra are tabulated as follows: chemical shift, multiplicity (s=singlet, d=doublet, t=triplet, q=quartet, qn=quintet, dd=doublet of doublets, ddd=doublet of doublet of doublets, dddd=doublet of doublet of doublet of doublets, dt=doublet of triplets, m=multiplet, b=broad), coupling constant and integration. ¹³C NMR spectra are tabulated by observed peak. Melting points were determined using a Thomas-Hoover capillary melting point apparatus and are uncorrected. Infrared spectra were measured on a Jasco FT/IR 480 plus spectrometer. High-resolution mass spectra (HRMS) were obtained at the University of Pennsylvania on a Waters GCT Premier spectrometer. Single crystal X-ray structures were determined at the University of Pennsylvania. X-ray intensity data were collected on a Rigaku Mercury CCD or Bruker APEXII CCD area detector employing graphite-monochromated Mo-Kα radiation (λ = 0.71073 Å) at a temperature of 143(1) K.

Preparation of water-washed silica gel for column chromatography (where specified):

[0067] Silica gel was suspended in H₂O and the slurry mixture was then packed into a prepared column. The obtained H₂O-washed silica gel packed column was then rinsed with 2 column volumes of acetone, 1 column volume of EtOAc and 2 column volumes of hexanes, successively. The obtained column was then ready for use.


[0068] Preparation of 3-butyl-1,1-dimethyl-1,3-dihydro-benzo[c][1,2]oxasilole:

[0069] *Synthesis of 1-(2-bromophenyl)pentan-1-ol*: Following the procedure described by Wagner (P. J. Wagner and E. J. Siebert, J. Am. Chem. Soc, 1981, 103, 7329), $n\text{-BuMgCl}$ (2.0 M in Et_2O , 14.3 mL, 38.1 mmol) was added slowly to a vigorously stirred solution of 2-bromobenzaldehyde (6.4 g, 34.6 mmol) in Et_2O (50 mL) at rt. The resulting suspension was heated to reflux for 2 h and cooled to rt. The reaction mixture was quenched, using extreme caution, by slow addition of aqueous 1 N HCl solution. The organic layer was collected and the aqueous layer was extracted using Et_2O (2 x 60 mL). The combined organic layers were washed with brine, dried over MgSO_4 and concentrated *in vacuo*. The crude product was purified by flash chromatography on silica gel (1:10, EtOAc /Hexane) to afford the desired alcohol (6.9 g, 83%) as light yellow oil: ^1H NMR (500 MHz, CDCl_3) δ 7.55 (dd, $J = 7.5$, 1.0 Hz, 1H), 7.52 (dd, $J = 8.0$, 1.0 Hz, 1H), 7.34 (td, $J = 7.5$, 1.0 Hz, 1H), 7.12 (td, $J = 8.0$, 1.5 Hz, 1H), 5.07 (dd, $J = 8.0$, 4.5 Hz, 1H), 2.08 (OH, 1H), 1.80 (m, 1H), 1.69 (m, 1H), 1.51 (m, 1H), 1.38 (m, 3H), 0.94 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 144.1, 132.8, 128.9, 127.5, 122.2, 73.1, 37.6, 28.2, 22.7, 14.2.

[0070] Following Hiyama's procedure (Y. Nakao, H. Imanaka, A. K. Sahoo, A. Yada, T. Hiyama, J. Am. Chem. Soc. 2005, 127, 6952), the alcohol (6.9 g, 28.5 mmol) obtained above was dissolved in 3,4-dihydro-2H-pyran (3.0 g, 31.4 mmol) at rt. A drop of concentrated HCl (37.5%, about 10 μL) was added and the resulting mixture was stirred for 24 h at rt. The reaction mixture was diluted with diethyl ether (20 mL) and quenched by addition of aqueous NaHCO_3 (10 mL). The organic layer was collected and the aqueous layer was extracted using Et_2O (2 x 20 mL). The combined organic layers were washed with brine, dried over MgSO_4 and concentrated *in vacuo*. The crude product was purified by short flash chromatography on silica gel (1:10, EtOAc /Hexane) to afford a colorless oil (7.9 g, 24.2 mmol, 85%), which was dissolved in dry THF (60 mL) and cooled to -78°C . $n\text{-Butyllithium}$ (2.0 M in hexane, 13.3 mL, 26.6 mmol) was added dropwise and the resulting solution was allowed to warm to -30°C over 1 hr and stirred for an additional 30 min at -30°C . The solution was then cooled to -78°C and chlorodimethylsilane (Me_2SiHCl , 4.6 g, 48.4 mmol) was added dropwise over 10 minutes. The

resulting reaction mixture was allowed to warm to rt and was stirred overnight. The reaction mixture was quenched by addition of saturated aqueous NaHCO_3 (30 mL) and extracted with Et_2O (3 x 50 mL). The combined organic layers were washed with brine, dried with MgSO_4 and concentrated *in vacuo*. The crude product was purified by short flash chromatography on silica gel (1:10, EtOAc/Hexane) to afford the desired silylated product (6.22 g, 20.3 mmol, 84%) as a colorless oil. The silylated product was then dissolved in MeOH (40 mL) and $p\text{-TsOH}\cdot\text{H}_2\text{O}$ (0.08 g, 0.4 mmol) was added. The resulting solution was stirred for 24 hrs at rt and concentrated *in vacuo*. Distillation of the crude product under vacuum afforded 1-oxa-2-silacyclopentane ($\pm$)-2 (3.89 g, 17.7 mmol, 87%) as a colorless oil: IR (neat, cm^{-1}) 3059, 2958, 2927, 2858, 1594, 1443, 1250, 1080, 919, 865, 830, 789, 744.; ^1H NMR (500 MHz, CDCl_3) δ 7.56 (d, $J = 7.0$ Hz, 1H), 7.41 (td, $J = 7.5, 1.0$ Hz, 1H), 7.29 (t, $J = 7.0$ Hz, 1H), 7.22 (d, $J = 7.5$ Hz, 1H), 5.26 (dd, $J = 7.5, 3.5$ Hz, 1H), 1.93 (m, 1H), 1.62 (m, 1H), 1.50-1.20 (m, 4H), 0.90 (t, $J = 7.5$ Hz, 3H), 0.41 (s, 3H), 0.38 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 153.5, 135.8, 131.0, 129.7, 127.0, 122.4, 81.9, 38.9, 27.5, 23.0, 14.2, 1.5, 0.8; HRMS (ES^+) m/z ($\text{M}+\text{H}$) $^+$: Calcd for $\text{C}_{13}\text{H}_{21}\text{OSi}$: 221.1362, found: 221.1367.


Preparation of Silicon-Based Cross Coupling Agents



[0071] To a solution of benzaldehyde (5.1 mL, 50 mmol) in a mixture of hexanes (250 mL) and Et_2O (200 mL) under an atmosphere of N_2 at room temperature was added a solution of n -butyllithium in hexanes (110 mmol) dropwise, maintaining the reaction mixture at room temperature using a water bath. The resulting solution was heated to reflux for 16 h, and the brown mixture was cooled to room temperature, then to -78°C using an acetone/ $\text{CO}_2(\text{s})$ bath. To this solution was added $\text{R}_1(\text{R}_2)\text{SiHCl}$ (110 mmol), and the resulting pale yellow slurry was allowed to warm to room temperature and stirred for 16 h. The reaction mixture was diluted with 0.5 M aq. KHCO_3 (100 mL) and extracted with Et_2O (3 x 50 mL). The combined organic layers were washed with brine, dried (MgSO_4), filtered and concentrated under reduced pressure to afford an oil.

[0072] The crude oil was taken up in THF (500 mL) in a 2 L flask, and $4 \text{ \AA}$ molecular sieves were added. To the stirred mixture was added KOt-Bu (280 mg, 2.5 mmol), and vigorous


gas evolution resulted. Once gas production had ceased, the turbid mixture was diluted with H₂O (100 mL), and the aqueous phase was extracted with Et₂O (3 x 50 mL). The organic extracts were dried (MgSO₄), filtered and concentrated. The resulting yellow oil was purified by chromatography on SiO₂ (100% Hex then 5% Et₂O/Hex). In some cases, the title compound was purified a second time by bulb-to-bulb distillation under vacuum to remove co-eluting impurities.



3-Butyl-1,1-dimethyl-1,3-dihydrobenzo[c][1,2]oxasilole (1). To a cooled solution of benzaldehyde (21.0 g, 198 mmol, 1.00 equiv) in hexanes (733 mL, pre-dried over MgSO₄) and Et₂O (587 mL) at 0 °C was added *n*-BuLi (224 mL, 1.94 M in hexanes, 435 mmol, 2.20 equiv) dropwise. The reaction mixture was allowed to warm to room temperature and was stirred for 30 minutes. The flask was then fitted with a reflux condenser and the reaction mixture was heated to reflux (75 °C) for 16 h. A dark solution resulted and the reaction mixture was allowed to cool to room temperature. The reflux condenser was removed and the reaction mixture was cooled to -78 °C and Me₂SiHCl (48.4 mL, 435 mmol, 2.20 equiv) was then added dropwise. The reaction mixture was allowed to slowly warm to room temperature and stirred for 8 h. The resulting pale yellow slurry was then quenched with H₂O (300 mL) and the aqueous phase extracted with Et₂O (3 × 100 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure.

[0073] The resulting crude orange oil was taken up in THF (200 mL) at room temperature and 4 Å molecular sieves (3.00 g) and KO^{*t*}-Bu (1.10 g, 9.90 mmol, 0.05 equiv) were added as a single portion and vigorous evolution of H₂ was observed. The reaction mixture was allowed to stir at room temperature for 5 h, quenched with H₂O (50 mL) and the aqueous phase extracted with Et₂O (3 × 100 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on water-washed silica (100 % hexanes then 1% EtOAc in hexanes) followed by Kugelrohr distillation (85-95 °C, 0.05 mmHg) provided **1** (22.7 g, 103 mmol, 52% yield) as a colorless oil. Analytical data matches that which has been previously reported:⁷ **R_f** 0.35 (1% EtOAc in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.56 (d, *J* = 7.0 Hz, 1H), 7.41 (td, *J* = 7.5, 1.0 Hz, 1H), 7.29 (t, *J* = 7.0 Hz, 1H), 7.22 (d, *J* = 7.5 Hz, 1H), 5.26 (dd, *J* = 7.5, 3.5 Hz, 1H), 1.93 (m, 1H), 1.62 (m, 1H), 1.50-1.20 (m, 4H), 0.90 (t, *J* = 7.5 Hz, 3H), 0.41 (s, 3H), 0.38 (s, 3H); ¹³C NMR (125 MHz,

CDCl₃) δ 153.5, 135.8, 131.0, 129.7, 127.0, 122.4, 81.9, 38.9, 27.5, 23.0. 14.2, 1.5, 0.8; **IR** (neat) 3059 (s), 2958 (m), 2927 (m), 2858 (m), 1594 (s), 1443 (s), 1250 (s), 1080 (s), 919 (m), 865 (s), 830 (s), 789 (s), 744 (s) cm⁻¹; **HRMS** (ES⁺) m/z calcd for C₁₃H₂₁OSi [M+H]⁺ 221.1362, found 221.1367.




1,1-Dimethyl-3-phenyl-1,3-

dihydrobenzo[c][1,2]oxasilole (1a): Phenylmagnesium bromide (10.4 mL, 3.00 M in Et₂O, 31.3 mmol, 1.20 equiv) was added dropwise to a vigorously stirred solution of 2-bromobenzaldehyde (4.82 g, 26.1 mmol, 1.00 equiv) in Et₂O (75 mL) at 0 °C. The reaction mixture was stirred at room temperature for 12 h, quenched with sat. aq. NH₄Cl (25 mL) and the aqueous phase extracted with Et₂O (2 x 25 mL). The combined organic layers were washed with brine, dried (MgSO₄), and concentrated under reduced pressure. Flash chromatography on silica (1% Et₂O in hexanes then 10% Et₂O in hexanes) provided the desired (2-bromophenyl)(phenyl)methanol (**S1**) (6.44 g, 24.5 mmol, 94% yield) as a white solid. Analytical data matches that which has been previously reported:⁸ **R_f** 0.3 (10% EtOAc in hexanes); **¹H NMR** (500 MHz, CDCl₃) δ 7.61-7.52 (m, 2 H), 7.41 (d, J = 6.9 Hz, 2 H), 7.37-7.31 (m, 3 H), 7.28 (t, J = 7.1 Hz, 1 H), 7.15 (dt, J = 1.6, 7.6 Hz, 1 H), 6.21 (d, J = 3.8 Hz, 1 H), 2.34 (d, J = 4.0 Hz, 1 H); **¹³C NMR** (125 MHz, CDCl₃) δ 142.6, 142.3, 133.0, 129.3, 128.6, 127.9, 127.8, 127.2, 126.7, 122.9, 74.9.

[0074] The resulting (2-bromophenyl)(phenyl)methanol (**S1**) (4.29 g, 16.3 mmol, 1.00 equiv) was dissolved in THF (50 mL) and *n*-BuLi (14.7 mL, 2.45 M in hexanes, 35.9 mmol, 2.20 equiv) was added dropwise at -78 °C. The reaction mixture was stirred for 1 h, followed by the addition of Me₂SiHCl (3.90 mL, 35.9 mmol, 2.20 equiv) at -78 °C. The resulting reaction mixture was allowed to warm to room temperature and stirred for 12 h. The reaction mixture was quenched by addition of *t*-BuOH (20 mL) and stirred for 5 h, followed by the addition of H₂O (20 mL) and stirred for another 2 h. The aqueous phase was then extracted with Et₂O (2 x 25 mL) and the combined organic layers were washed with brine, dried (MgSO₄), and concentrated under reduced pressure. Flash chromatography on water-washed silica gel (100% hexanes then 1% Et₂O/Hexanes) provided the desired siloxane **1a** (1.82 g, 26.8 mmol, 46% yield) as a white crystalline solid: **m.p.** 45.5 - 46.5 °C; **R_f** 0.3 (5% Et₂O in hexanes); **¹H NMR** (500 MHz, CDCl₃)

δ 7.63-7.59 (m, 1 H), 7.36-7.26 (m, 7 H), 7.05-7.01 (m, 1 H), 6.17 (s, 1 H), 0.53 (s, 3 H), 0.45 (s, 3 H); ^{13}C NMR (125 MHz, CDCl_3) δ 152.6, 143.9, 135.3, 130.8, 129.9, 128.7, 127.9, 127.3, 127.3, 123.9, 84.2, 1.4, 0.7; IR (CH_2Cl_2) 3057 (m), 2964 (m), 2874 (m), 1447 (m), 1265 (s), 1181 (m), 1136 (m), 1042 (s), 1016 (s), 862 (s), 822 (s), 793 (s), 740 (s), 702 (s), 656 (m) cm^{-1} ; HRMS (CI^+) m/z calculated for $\text{C}_{15}\text{H}_{17}\text{SiO}$ $[\text{M}+\text{H}]^+$ 241.1049, found 241.1034.




3-Isopropyl-1,1-dimethyl-1,3-dihydrobenzo[c][1,2]oxasilole (1b):

Phenylmagnesium bromide (11.7 mL, 3.00 M in Et_2O , 35.1 mmol, 1.20 equiv) was added dropwise to a vigorously stirred solution of isobutyraldehyde (2.11 g, 29.3 mmol, 1.00 equiv) in Et_2O (50 mL) at 0 °C. The reaction mixture was warmed to room temperature and stirred for 12 h, then quenched with sat. aq. NH_4Cl (25 mL). The aqueous phase was extracted with Et_2O (2 x 25 mL) and the combined organic layers were washed with brine, dried (MgSO_4) and concentrated under reduced pressure. Flash chromatography on silica, (10% Et_2O in hexanes) provided the desired 2-methyl-1-phenylpropan-1-ol (**S2**) (4.09 g, 27.2 mmol, 93% yield) as a colorless oil. Analytical data matches that which has been previously reported:⁹ R_f 0.15 (10% Et_2O in hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.36-7.26 (m, 5 H), 4.37 (dd, J = 3.3, 6.8 Hz, 1 H), 2.02-1.91 (m, 1 H), 1.82 (d, J = 3.4 Hz, 1 H), 1.01 (d, J = 6.7 Hz, 3 H), 0.81 (d, J = 6.7 Hz, 3 H); ^{13}C NMR (125 MHz, CDCl_3) δ 143.8, 128.3, 127.6, 126.7, 80.2, 35.4, 19.2, 18.4.

[0075] The resulting 2-methyl-1-phenylpropan-1-ol (**S2**) (2.76 g, 18.4 mmol, 1.00 equiv) was dissolved in hexanes (125 ml, pre-dried over MgSO_4) and Et_2O (100 ml) at 0 °C and $n\text{-BuLi}$ (22.9 ml, 1.77 M in hexanes, 40.5 mmol, 2.20 equiv) was added dropwise. The reaction mixture was allowed to warm to room temperature and was stirred for 30 minutes. The flask was then fitted with a reflux condenser and the system was heated to reflux (75 °C) for 16 h. A dark solution resulted and the reaction mixture was allowed to cool to room temperature. The reflux condenser was removed and the reaction mixture was cooled to -78 °C. Me_2SiHCl (4.40 ml, 40.5 mmol, 2.20 equiv) was added dropwise. The reaction mixture was allowed to slowly warm to room temperature and stirred for 12 h. The slurry was quenched by addition of $t\text{-BuOH}$ (20 mL) and stirred for 5 h, followed by the addition of H_2O (20 ml) and stirred for another 2 h. The aqueous phase was extracted with Et_2O (2 x 25 mL) and the combined organic layers were washed with brine, dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography on water-washed silica gel (100% hexanes then 1% Et_2O /Hexanes) followed by


Kugelrohr distillation (75-80 °C, 0.025 mmHg) provided **1b** (1.48 g, 7.17 mmol, 39% yield) as a colorless oil: R_f 0.5 (5% Et₂O in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.54 (d, *J* = 7.1 Hz, 1 H), 7.39 (dt, *J* = 1.0, 7.5 Hz, 1 H), 7.29 (t, *J* = 7.2 Hz, 1 H), 7.20 (d, *J* = 7.7 Hz, 1 H), 5.17 (d, *J* = 2.2 Hz, 1 H), 2.12 (dspt, *J* = 2.5, 6.8 Hz, 1 H), 1.17 (d, *J* = 6.9 Hz, 3 H), 0.60 (d, *J* = 6.7 Hz, 3 H), 0.41 (s, 3 H), 0.36 (s, 3 H); ¹³C NMR (125 MHz, CDCl₃) δ 152.5, 136.4, 130.8, 129.7, 127.0, 122.4, 86.5, 34.6, 20.4, 15.0, 1.0, 0.8; IR (neat) 3060 (m), 2963 (s), 2873 (m), 1594 (m), 1469 (m), 1442 (m), 1383 (m), 1365 (m), 1251 (s), 1198 (m), 1137 (m), 1120 (m), 1102 (m), 1067 (s), 1014 (s), 953 (s), 874 (s), 830 (s), 788 (s), 744 (s), 704 (m), 651 (m) cm⁻¹; HRMS (CI⁺) *m/z* calculated for C₁₁H₁₅SiO [M-Me]⁺ 191.0892, found 191.0893.



3-(sec-Butyl)-1,1-dimethyl-1,3-dihydrobenzo[c][1,2]oxasilole (1c):


To a cooled solution of benzaldehyde (3.25 g, 30.6 mmol, 1.00 equiv) in hexanes (125 ml, pre-dried over MgSO₄) and Et₂O (100 mL) at 0 °C was added *sec*-BuLi (26.2 mL, 1.4 M in cyclohexane, 36.7 mmol, 1.20 equiv) dropwise. The reaction mixture was allowed to warm to room temperature and was stirred for 5 h, at which time *n*-BuLi (20.7 mL, 1.77 M in hexanes, 36.7 mmol, 1.20 equiv) was added dropwise. The flask was then fitted with a reflux condenser and the system was heated to reflux (75 °C) for 16 h. A dark solution resulted and the reaction mixture was allowed to cool to room temperature. The reflux condenser was removed and the reaction mixture was cooled to -78 °C and Me₂SiHCl (7.97 ml, 67.3 mmol, 2.20 equiv) was added dropwise. The reaction mixture was allowed to slowly warm to room temperature and stirred for 12 h. The slurry was quenched by addition of *t*-BuOH (20 mL) and stirred for 5 h, followed by the addition of H₂O (20 ml) and stirred for another 2 h. The aqueous phase was extracted with Et₂O (2 x 25 mL) and the combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on water-washed silica gel (100% hexanes then 1% Et₂O/Hexanes) followed by Kugelrohr distillation (140-160 °C, 0.01 mmHg) provided **1c**, isolated as a 5:3 ratio of diastereomers by ¹H NMR spectroscopy (2.02 g, 9.17 mmol, 30% yield) as a colorless oil: R_f 0.55 (5% Et₂O in hexanes); Major diastereoisomer: ¹H NMR (500 MHz, CDCl₃) δ 7.54 (d, *J* = 7.1 Hz, 1 H), 7.39 (t, *J* = 7.5 Hz, 1 H), 7.29 (t, *J* = 7.1 Hz, 1 H), 7.21 (d, *J* = 7.7 Hz, 1 H), 5.20 (d, *J* = 2.2 Hz, 1 H), 1.88-1.79 (m, 1 H), 1.13 (d, *J* = 6.9 Hz, 3 H), 1.08-1.01 (m, 2 H), 0.79 (t, *J* = 7.4 Hz, 3 H), 0.40 (s, 3 H), 0.36 (s, 3 H); ¹³C NMR (125 MHz, CDCl₃) δ 152.3, 136.6, 130.9, 129.6, 126.9, 122.4, 86.7,

41.8, 22.5, 16.9, 12.5, 1.0, 0.8. *Minor diastereoisomer*: $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.54 (d, J = 7.1 Hz, 1 H), 7.39 (t, J = 7.5 Hz, 1 H), 7.29 (t, J = 7.1 Hz, 1 H), 7.19 (d, J = 7.7 Hz, 1 H), 5.20 (d, J = 1.2 Hz, 1 H), 1.88-1.79 (m, 1 H), 1.74-1.65 (m, 1 H), 1.53-1.43 (m, 1 H), 1.04 (t, J = 7.7 Hz, 3 H), 0.55 (d, J = 6.7 Hz, 3 H), 0.40 (s, 3 H), 0.36 (s, 3 H); $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ 152.7, 136.4, 130.8, 129.7, 126.9, 122.2, 84.62, 41.5, 31.9, 27.5, 12.5, 1.0, 0.7; **IR** (neat) 3060 (m), 2962 (s), 2875 (s), 1594 (m), 1443 (s), 1375 (m), 1323 (m), 1250 (s), 1197 (m), 1137 (m), 1107 (s), 1068 (s), 1040 (s), 1014 (s), 960 (s), 875 (s), 826 (s), 789 (s), 746 (s), 705 (m), 653 (m) cm^{-1} ; **HRMS** (Cl^+) m/z calculated for $\text{C}_{13}\text{H}_{19}\text{SiO}$ $[\text{M}-\text{H}]^+$ 221.1362, found 221.1368.



1,1,3-Trimethyl-1,3-dihydrobenzo[c][1,2]oxasilole (1d): To a cooled solution of benzaldehyde (2.63 g, 24.8 mmol, 1.00 equiv) in hexanes (125 ml, pre-dried over MgSO_4) and Et_2O (100 ml) at 0 °C was added MeLi (1.36 M in Et_2O , 29.7 mmol, 1.20 equiv) dropwise. The resulting solution was allowed to warm to room temperature and stirred for 5 h, at which time $n\text{-BuLi}$ (12.1 mL, 2.45 M in hexanes, 29.7 mmol, 1.20 equiv) was added dropwise. The flask was then fitted with a reflux condenser and the system was heated to reflux (75 °C) for 16 h. A dark solution resulted and the reaction mixture was allowed to cool to room temperature. The reflux condenser was removed and the reaction mixture was cooled to -78 °C and Me_2SiHCl (5.20 ml, 54.6 mmol, 2.20 equiv) was added dropwise. The reaction mixture was allowed to slowly warm to room temperature and stirred for 12 h. The reaction mixture was quenched by addition of $t\text{-BuOH}$ (20 mL) and stirred for 5 h, followed by the addition of H_2O (20 ml) and stirred for another 2 h. The aqueous phase was extracted with Et_2O (2 x 25 mL) and the combined organic layers were washed with brine, dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography on water-washed silica gel (100% hexanes then 1% Et_2O /Hexanes) followed by Kugelrohr distillation (55 °C, 0.01 mmHg) provided **1d** (1.55 g, 8.69 mmol, 35 % yield) as a colorless oil: R_f 0.45 (5% Et_2O in hexanes); $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.56 (d, J = 7.1 Hz, 1 H), 7.40 (dt, J = 0.8, 7.5 Hz, 1 H), 7.30 (t, J = 7.3 Hz, 1 H), 7.22 (d, J = 7.7 Hz, 1 H), 5.34 (q, J = 6.5 Hz, 1 H), 1.51 (d, J = 6.5 Hz, 3 H), 0.41 (s, 3 H), 0.37 (s, 3 H); $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ 154.5, 135.2, 131.0, 129.8, 127.0, 122.3, 78.0, 25.4, 1.7, 0.6; **IR** (neat) 3060 (m), 2968 (s), 2924 (m), 2867 (m), 1595 (m), 1444 (s), 1368 (m), 1318 (s), 1251 (s), 1199 (m), 1137 (m), 1086 (s), 1028 (s), 929 (s), 855 (s), 828 (s), 793 (s), 759 (s), 742 (s), 696 (m), 653 (m) cm^{-1} ; **HRMS** (Cl^+) m/z calculated for $\text{C}_9\text{H}_{11}\text{SiO}$ $[\text{M}-\text{Me}]^+$ 163.0579, found

163.0578.

**1,1,3,3-Tetramethyl-1,3-dihydrobenzo[c][1,2]oxasilole (1f):¹**


Following a previously reported procedure,¹ methyl 2-bromobenzoate (9.70 g, 45.0 mmol, 1.00 equiv) was dissolved in Et₂O (100 ml) and cooled to 0 °C. Methylmagnesium bromide (99 mL, 1.0 M in Bu₂O, 99.0 mmol, 2.20 equiv) was then added via cannula into the reaction mixture. The resulting solution was heated to reflux (40 °C) for 2 h then cooled to room temperature before being quenched with sat. aq. NH₄Cl (75 mL). The aqueous phase extracted with Et₂O (2 x 50 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure to afford 2-(2-bromophenyl)propan-2-ol (**S3**) (8.30 g, ca. 38.8 mmol, 1.00 equiv), which was used without further purification.

[0076] To the crude alcohol was added conc. HCl (3 drops) and 3,4-dihydro-2H-pyran (**S3**) (4.07 g, 48.5 mmol, 1.20 equiv) and stirred neat at room temperature for 24 h. The mixture was diluted with Et₂O (50 mL), and washed with a sat. aq. NaHCO₃ (25 mL). The aqueous layer was extracted with Et₂O (2 x 50 mL) and the combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure to afford 2-((2-(2-bromophenyl)propan-2-yl)oxy)tetrahydro-2H-pyran (**S4**) (8.30 g, ca. 38.8 mmol, 1.00 equiv) which was used without further purification.

[0077] The crude THP-protected product (**S4**) was dissolved in THF (50 mL) and *n*-BuLi (23.3 mL, 2.00 M in hexanes, 46.6 mmol, 1.20 equiv) was added dropwise at -78 °C. The reaction mixture was stirred for 1 h, followed by the addition of Me₂SiHCl (6.30 mL, 58.2 mmol, 1.50 equiv) at -78 °C. The resulting reaction mixture was allowed to warm to room temperature and stirred for 12 h. The reaction mixture was quenched by addition of H₂O (50 ml). The aqueous phase was then extracted with Et₂O (2 x 30 mL) and the combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure to afford dimethyl(2-(2-((tetrahydro-2H-pyran-2-yl)oxy)propan-2-yl)phenyl)silane (**S5**) (8.30 g, ca. 38.8 mmol, 1.00 equiv) which was used without further purification.


[0078] To this crude material (**S5**) was added MeOH (50 mL) and *p*-toluenesulfonic acid monohydrate (370 mg, 1.94 mmol, 0.05 equiv), and the mixture was stirred at room temperature for 12 h before concentration under reduced pressure. The residue was diluted with Et₂O (25 mL), and washed with a sat. aq. NaHCO₃ (25 mL). The aqueous layer was extracted

with Et₂O (2 x 25 mL), and the combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Kugelrohr distillation (60 °C, 0.01 mmHg) provided **1f** (3.27 g, 17.0 mmol, 38% yield from methyl 2-bromobenzoate) as a colorless oil. Analytical data matches that which has been previously reported:² **R_f** 0.3 (1% Et₂O in hexanes); ¹**H NMR** (500 MHz, CDCl₃) δ 7.52 (dt, *J* = 7.1, 0.9 Hz, 1 H), 7.41 (td, *J* = 7.5, 1.3 Hz, 1 H), 7.30 (dt, *J* = 7.2, 0.9 Hz, 1 H), 7.22 (d, *J* = 7.7 Hz, 1 H), 1.55 (s, 6 H), 0.39 (s, 6 H); ¹³**C NMR** (125 MHz, CDCl₃) δ 157.6, 134.4, 130.6, 129.7, 126.6, 122.1, 83.5, 32.1, 1.37.



3-Butyl-1,1-diisopropyl-1,3-dihydrobenzo[c][1,2]oxasilole (2c): To

a cooled solution of benzaldehyde (3.08 g, 29.0 mmol, 1.00 equiv) in hexanes (125 ml, pre-dried over MgSO₄) and Et₂O (100 ml) at 0 °C was added *n*-BuLi (27.8 mL, 2.3 M in hexanes, 63.9 mmol, 2.20 equiv) dropwise. The reaction mixture was allowed to warm to room temperature and was stirred for 30 minutes. The flask was then fitted with a reflux condenser and the system was heated to reflux (75 °C) for 16 h. A dark solution resulted and was allowed to cool to room temperature, the reflux condenser was removed and the reaction mixture was cooled to -78 °C and *i*-Pr₂SiHCl (10.9 ml, 63.8 mmol, 2.20 equiv) was added dropwise. The reaction mixture was allowed to slowly warm to room temperature and stirred for 12 h. The reaction mixture was quenched by addition of 3M HCl in MeOH (40 mL) at 0 °C and stirred for 12 h at room temperature. Water (100 mL) was added and stirred for another 2 h and the aqueous phase extracted with Et₂O (2 x 50 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on silica (1% Et₂O/Hexanes) provided **2c** (5.94 g, 21.5 mmol, 74% yield) as a colorless oil: **R_f** 0.70 (5% Et₂O in hexanes); ¹**H NMR** (500 MHz, CDCl₃) δ 7.53 (d, *J* = 7.1 Hz, 1 H), 7.38 (t, *J* = 7.5 Hz, 1 H), 7.27 (t, *J* = 7.1 Hz, 1 H), 7.20 (d, *J* = 7.7 Hz, 1 H), 5.18-5.14 (m, 1 H), 1.94-1.86 (m, 1 H), 1.60-1.48 (m, 3 H), 1.45-1.30 (m, 2 H), 1.25-1.16 (m, 2 H), 1.06 (d, *J* = 6.7 Hz, 3 H), 1.05 (d, *J* = 7.1 Hz, 3 H), 1.00 (d, *J* = 7.5 Hz, 3 H), 0.93 (d, *J* = 7.3 Hz, 3 H), 0.92 (t, *J* = 7.3 Hz, 3 H); ¹³**C NMR** (125 MHz, CDCl₃) δ 154.7, 132.4, 132.2, 129.5, 126.7, 122.2, 82.3, 39.1, 28.1, 23.0, 17.6, 17.5, 17.2, 17.1, 14.2, 13.4, 12.9; **IR** (neat) 3059 (m), 3000 (m), 2941 (s), 2864 (s), 1595 (m), 1464 (s), 1443 (m), 1381 (m), 1261 (m), 1111 (m), 1080 (s), 1054 (m), 1012 (m), 988 (m), 973 (m), 917 (s), 880 (s), 846 (m), 831 (m), 816 (m), 749 (s), 716 (s), 668 (s), 648 (m), 612 (m) cm⁻¹; **HRMS** (CI⁺) *m/z* calculated for C₁₄H₂₁SiO [M-C₃H₇]⁺ 233.1726, found 233.1726.




1,1-Diisopropyl-3-phenyl-1,3-dihydrobenzo[c][1,2]oxasilole (2d):


To the previously prepared (2-bromophenyl)(phenyl)methanol (**S1**) (5.18 g, 19.7 mmol, 1.00 equiv) dissolved in THF (50 mL) was added *n*-BuLi (19.0 mL, 2.28 M in hexanes, 43.3 mmol 2.20 equiv) dropwise at -78°C . The reaction mixture was stirred for 1 h, followed by the addition of *i*-Pr₂SiHCl (7.39 mL, 43.3 mmol, 2.20 equiv) at -78°C . The resulting reaction mixture was allowed to warm to room temperature and stirred for 12 h. The slurry was quenched by addition of 3M HCl in MeOH (40 mL) at 0°C and stirred for 4 h at room temperature. Water (100 mL) was added and stirred for another 2 h and the aqueous phase extracted with Et₂O (2 x 50 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on silica (1% Et₂O/Hexanes) provided **2d** (4.12 g, 13.9 mmol, 71% yield) as a colorless oil: *R*_f 0.45 (5% Et₂O in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.63-7.59 (m, 1 H), 7.37-7.27 (m, 7 H), 6.99 (d, *J* = 6.9 Hz, 1 H), 6.12 (s, 1 H), 1.39-1.24 (m, 2 H), 1.16 (d, *J* = 7.5 Hz, 3 H), 1.13 (d, *J* = 7.5 Hz, 3 H), 1.06 (d, *J* = 5.7 Hz, 3 H), 1.04 (d, *J* = 5.7 Hz, 3 H); ¹³C NMR (125 MHz, CDCl₃) δ 153.4, 143.8, 132.7, 131.9, 129.8, 128.6, 128.0, 127.8, 127.0, 123.9, 84.6, 17.7, 17.2, 13.6, 13.3; IR (neat) 3061 (m), 3031 (m), 3000 (m), 2942 (s), 2891 (m), 2864 (s), 1592 (m), 1494 (m), 1463 (s), 1442 (s), 1383 (m), 1263 (m), 1182 (m), 1133 (m), 1077 (m), 1065 (s), 1038 (s), 1014 (s), 988 (s), 919 (m), 881 (m), 815 (s), 747 (s), 732 (s), 713 (s), 698 (s), 669 (s), 633 (m) cm⁻¹; HRMS (CI⁺) *m/z* calculated for C₁₆H₁₇SiO [M-C₃H₇]⁺ 253.1049, found 253.1056.

[0079] Preparation of Et₂SiHCl from commercially available Et₂SiH₂:⁴ Anhydrous CuCl₂ (96.5 g, 714 mmol, 2.10 equiv) was dried under vacuum at 200°C (utilizing a sand bath) in a 2000 mL two-necked round-bottomed flask (RBF) for 12 hours. Upon cooling to room temperature, CuI (3.24 g, 17.0 mmol, 0.05 equiv), Et₂O (680 mL ~ 0.5 M) and Et₂SiH₂ (30.0 g, 340 mmol, 1.00 equiv) were successively added and the resulting slurry stirred at room temperature for 43 hours. After 8 hours, it was noted that the reaction went from brown to light grey in color with black precipitate forming. After 43 hours, the reaction mixture was filtered under N₂ atmosphere and the flask rinsed with Et₂O (2 x 50 mL). The Et₂O was distilled (35°C) and the reaction mixture transferred to a 100 mL flame-dried RBF via syringe, carefully leaving behind any remaining Cu salts. Kugelrohr distillation ($95-100^{\circ}\text{C}$) of this crude reaction mixture

under N₂ provided clean Et₂SiHCl (32.2 g, 263 mmol, 77% yield) whose analytical data matches that which has been previously reported.⁴




3-Butyl-1,1-diethyl-1,3-dihydrobenzo[c][1,2]oxasilole (2a): To a cooled solution of benzaldehyde (10.0 g, 94.2 mmol, 1.00 equiv) in hexanes (500 mL, pre-dried over MgSO₄) and Et₂O (400 mL) at 0 °C was added *n*-BuLi (86.4 mL, 2.40 M in hexanes, 2.20 equiv) dropwise. The reaction mixture was allowed to warm to room temperature and was stirred for 30 minutes. The flask was then fitted with a reflux condenser and the system was heated to reflux (75 °C) for 16 h. A dark solution resulted and the reaction mixture was allowed to cool to room temperature. The reflux condenser was removed and the reaction mixture was cooled to -78 °C and Et₂SiHCl (28.9 mL, 207 mmol, 2.20 equiv) was added dropwise. The reaction mixture was allowed to slowly warm to room temperature and stirred for 8 h. The resulting pale yellow slurry was then quenched with H₂O (300 mL) with vigorous evolution of H₂ gas observed. The reaction mixture was allowed to stir for 5 h and the aqueous layer extracted with Et₂O (3 × 100 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on silica (100 % hexanes to 1% EtOAc in hexanes) provided **2a** (6.2 g, 25.0 mmol, 53% yield) as a colorless oil: *R*_f 0.45 (1% EtOAc in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.54 (d, *J* = 7.1 Hz, 1H), 7.39 (td, *J* = 7.5, 1.0 Hz, 1H), 7.28 (t, *J* = 7.2 Hz, 1H), 7.22 (d, *J* = 7.5 Hz, 1H), 5.22 (dd, *J* = 8.0, 3.3 Hz, 1H), 1.92 (m, 1H), 1.59 (m, 1H), 1.52-1.30 (m, 4H), 0.99 (t, *J* = 7.7 Hz, 3H), 0.94-0.88 (m, 6H), 0.88-0.78 (m, 4H); ¹³C NMR (125 MHz, CDCl₃) δ 154.2, 133.7, 131.6, 129.6, 126.8, 122.3, 82.12, 39.1, 27.8, 22.9, 14.2, 7.38, 7.16, 6.88, 6.60; IR (neat) 3058 (m), 2956 (s), 2932 (m), 2874 (m), 1459 (s), 1080 (s), 1013 (m), 916 (s), 741 (s) cm⁻¹; HRMS (CI⁺) *m/z* calculated for C₁₅H₂₄OSi [M]⁺ 248.1596, found 248.1593.



1,1-Diethyl-3-phenyl-1,3-dihydrobenzo[c][1,2]oxasilole (2b): To the previously prepared (2-bromophenyl)(phenyl)methanol (**S1**) (4.16 g, 15.8 mmol, 1.00 equiv) dissolved in THF (120 mL) was added *n*-BuLi (14.6 mL, 2.39 M in hexanes, 34.8 mmol, 2.20

equiv) dropwise at $-78\text{ }^{\circ}\text{C}$. The reaction mixture was stirred for 1 h, followed by the addition of Et_2SiHCl (4.84 mL, 34.8 mmol, 2.20 equiv) at $-78\text{ }^{\circ}\text{C}$. The resulting pale yellow slurry was then quenched with H_2O (300 mL) with vigorous evolution of H_2 gas observed. The reaction mixture was allowed to stir for 5 h and the aqueous phase was extracted with Et_2O (2 x 50 mL), the combined organic layers were washed with brine, dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography on silica (100% hexanes to 5% Et_2O in hexanes) provided **2b** (1.74 g, 6.48 mmol, 41% yield) as a colorless oil: R_f 0.45 (5% Et_2O in hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.63-7.60 (m, 1 H), 7.36-7.27 (m, 7 H), 7.02 (d, $J = 7.1$ Hz, 1 H), 6.16 (s, 1 H), 1.08 (t, $J = 7.9$ Hz, 3 H), 1.02-0.95 (m, 5 H), 0.95-0.82 (m, 2 H); ^{13}C NMR (125 MHz, CDCl_3) δ 153.2, 143.9, 133.6, 131.4, 129.9, 128.6, 128.0, 127.5, 127.1, 123.9, 84.4, 7.4, 7.1, 7.0, 6.7; IR (neat) 3059 (m), 2999 (m), 2956 (s), 2878 (s), 1595 (m), 1486 (m), 1451 (s), 1412 (m), 1343 (w), 1265 (m), 1236 (m), 1182 (m), 1133 (m), 1012 (bs), 964 (m), 926 (m), 871 (m), 817 (s), 739 (bs), 661 (m), 629 (m) cm^{-1} ; HRMS (CI^+) m/z calculated for $\text{C}_{17}\text{H}_{21}\text{SiO}$ $[\text{M}+\text{H}]^+$ 269.1362, found 269.1366.




3-(1,1-Dimethyl-1,3-dihydrobenzo[c][1,2]oxasilol-3-yl)pyridine

(3d): To a cooled solution of Et_2O (420 mL) containing $n\text{-BuLi}$ (48.7 mL, 2.00 M in hexanes, 1.30 equiv) at $-78\text{ }^{\circ}\text{C}$ was added 3-bromopyridine (20.0 g, 127 mmol, 1.20 equiv) dropwise. The resulting yellow slurry was allowed to stir at $-78\text{ }^{\circ}\text{C}$ for 30 min. 2-Bromobenzaldehyde (19.5 g, 106 mmol, 1.00 equiv) was added dropwise and the reaction mixture was allowed to stir for 5 h at $-78\text{ }^{\circ}\text{C}$. The reaction was then warmed to $0\text{ }^{\circ}\text{C}$ and quenched with 3M aq. HCl (100 mL). The organic layer was washed with another portion of 3M aq. HCl (50 mL). The acidic aqueous layers were collected and neutralized to pH 8-9 with 1M aq. NaOH (350 mL), producing a white turbid mixture that was extracted with Et_2O (3 x 150 mL). The combined organic layers were dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography on silica (0.5% to 1% to 5% MeOH in CH_2Cl_2) provided (2-bromophenyl)(pyridin-3-yl)methanol (**S6**) (23.6 g, 89.4 mmol, 85% yield) as an off-white crystalline solid: Analytical data matches that which has been previously reported:¹⁰ R_f 0.35 (5% MeOH in CH_2Cl_2); ^1H NMR (500 MHz, CDCl_3) δ 8.57 (d, $J = 1.8$ Hz, 1H), 8.41 (dd, $J = 4.7$, 1.1 Hz, 1H), 7.69 (td, $J = 7.6$, 1.6 Hz, 1H), 7.62 (dd, $J = 7.8$, 1.2 Hz, 1H), 7.53 (dd, $J = 7.9$, 1.0 Hz, 1H), 7.36 (t, $J = 7.5$ Hz, 1 H), 7.23 (dd, $J = 7.8$, 4.8 Hz, 1H), 7.16 (td, $J = 7.5$, 1.5 Hz, 1H),

6.21 (s, 1 H), 3.93 (s, 1 H); ^{13}C NMR (125 MHz, CDCl_3) δ 148.9, 148.7, 142.1, 138.2, 134.9, 133.1, 129.6, 128.5, 128.1, 123.6, 122.7, 72.7.


[0080] To a cooled solution of (2-bromophenyl)(pyridin-3-yl)methanol (**S6**) (7.00 g, 26.5 mmol, 1.00 equiv) in a mixture (1:1) of THF (100 mL) and Et_2O (100 mL) at -78°C was added *n*-BuLi (29.2 mL, 2.00 M in hexanes, 2.20 equiv) dropwise. The resulting dark yellow slurry was allowed to stir at -78°C for 30 min and Me_2SiHCl (6.50 mL, 58.3 mmol, 2.20 equiv) was added dropwise. The reaction mixture was allowed to slowly warm to room temperature and stirred for 12 h. The resulting orange slurry was then quenched with H_2O (100 mL) with vigorous evolution of H_2 gas observed. The reaction mixture was allowed to stir for 5 h and extracted with Et_2O (3×50 mL). The combined organic layers were collected and washed with 1M aq. HCl (3×50 mL). The acidic aqueous layers were collected and neutralized to pH 8-9 with 1M aq. NaOH (150 mL) producing a white turbid mixture that was extracted with Et_2O (3×100 mL). The combined organic layers were dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography on silica (0.5% to 1% to 5% MeOH in CH_2Cl_2) followed by Kugelrohr distillation ($135 - 150^\circ\text{C}$, 0.01 mmHg) provided **3d** (3.14 g, 13.0 mmol, 49% yield) as an off-white amorphous solid: R_f 0.2 (5% MeOH in CH_2Cl_2); ^1H NMR (500 MHz, CDCl_3) δ 8.61 (d, $J = 1.7$ Hz, 1H), 8.53 (dd, $J = 4.7, 1.5$ Hz, 1H), 7.63 (m, 1 H), 7.53 (dt, $J = 7.8, 1.8$ Hz, 1H), 7.33 (m, 2 H), 7.25 (m, 1 H), 7.00 (m, 1 H), 6.20 (s, 1 H), 0.53 (s, 3 H), 0.46 (s, 3 H); ^{13}C NMR (125 MHz, CDCl_3) δ 151.4, 149.3, 148.8, 139.3, 135.2, 134.7, 131.0, 130.1, 127.6, 81.7, 1.23, 0.56; IR (neat) 3057 (m), 2959 (s), 1662 (m), 1579 (m), 1475 (s), 1427 (s), 1252 (s), 1181 (s), 1136 (s), 1051 (m), 860 (m), 822 (s), 791 (s), 750 (s), 713 (s) cm^{-1} ; HRMS (ES^+) m/z calculated for $\text{C}_{14}\text{H}_{16}\text{NOSi}$ $[\text{M}+\text{H}]^+$ 242.1001, found 242.0993.




3-(1,1-Diethyl-1,3-dihydrobenzo[c][1,2]oxasilol-3-yl)pyridine (3e):

To a cooled solution of (2-bromophenyl)(pyridin-3-yl)methanol (**S6**) (7.00 g, 26.5 mmol, 1.00 equiv) in a mixture (1 : 1) of dry THF (100 mL) and dry Et_2O (100 mL) at -78°C was added *n*-BuLi (26.5 mL, 2.20 M in hexanes, 2.20 equiv) dropwise. The resulting dark yellow slurry was allowed to stir at -78°C for 30 min and Et_2SiHCl (8.15 mL, 58.3 mmol, 2.20 equiv) was added dropwise. The reaction mixture was allowed to slowly warm to room temperature and stirred for 12 h. The resulting red-brown solution was then quenched with H_2O (100 mL) with vigorous evolution of H_2 gas observed. The reaction mixture was allowed to stir for 5 h and extracted with


Et₂O (3 × 50 mL). The combined organic layers were collected and washed with 1M aq. HCl (3 × 50 mL). The acidic aqueous layers were collected and neutralized to pH 8-9 with 1M aq. NaOH (150 mL) producing a white turbid mixture that was extracted with Et₂O (3 × 100 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on silica (0.5% to 1% to 3% MeOH in CH₂Cl₂) provided **3e** (3.6 g, 13.4 mmol, 51% yield) as pale yellow oil: *R_f* 0.45 (5% MeOH in CH₂Cl₂); ¹H NMR (500 MHz, CDCl₃) δ 8.62 (d, *J* = 1.0 Hz, 1H), 8.53 (dd, *J* = 4.6, 1.2 Hz, 1H), 7.62 (m, 1 H), 7.53 (dt, *J* = 7.7, 1.6 Hz, 1H), 7.33 (m, 2 H), 7.23 (m, 1 H), 6.99 (d, *J* = 6.6 Hz, 1H), 6.19 (s, 1 H), 1.05 (m, 3 H), 1.0-0.88 (m, 7 H); ¹³C NMR (125 MHz, CDCl₃) δ 152.0, 149.4, 149.2, 139.3, 134.8, 133.6, 131.5, 130.1, 127.5, 123.7, 123.6, 81.9, 7.25, 7.00, 6.87, 6.51; IR (neat) 3057 (m), 2957 (m), 2876 (s), 2942 (s), 1580 (s), 1425 (m), 1235 (s), 1183 (s), 1134 (s), 1050 (m), 1017 (m), 817 (m), 744 (s) cm⁻¹; HRMS (ES⁺) *m/z* calculated for C₁₆H₂₀NOSi [M+H]⁺ 270.1314, found 270.1314.



1-(3-Bromopyridin-4-yl)pentan-1-ol (S7): To a cooled THF solution (315 mL) of LDA, prepared from diisopropylamine (15.6 mL, 110 mmol, 1.10 equiv) and *n*-BuLi (42.4 mL, 2.50 M in hexanes, 106 mmol, 1.05 equiv), 3-bromopyridine (15.8 g, 100 mmol, 1.00 equiv) was added dropwise as a solution in THF (16.6 mL) at -78 °C. The resulting solution temperature was further lowered to -100 °C and stirred for 10 min at this temperature before a THF solution (47.6 mL) of pentanal (21.4 mL, 200 mmol, 2.00 equiv) was added dropwise over 10 min. The reaction mixture was stirred at -100 °C for 1 h then warmed to -20 °C over 20 min before it was quenched with sat. aq. NH₄Cl (150 mL). The aqueous phase was extracted with EtOAc (3 × 75 mL). The combined organic layers were washed with brine, dried (MgSO₄) and concentrated under reduced pressure. Flash chromatography on silica (10% to 20% EtOAc in hexanes) provided 1-(3-bromopyridin-4-yl)pentan-1-ol (**S7**) (17.5 g, 71.7 mmol, 72% yield) as a yellow oil: *R_f* 0.1 (30% EtOAc in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 8.56 (s, 1 H), 8.43 (d, *J* = 5.0 Hz, 1H), 7.50 (d, *J* = 5.0 Hz, 1H), 4.98 (dd, *J* = 8.0 Hz, 3.3 Hz, 1 H), 3.03 (bs, 1 H), 1.76 (m, 1 H), 1.60 (m, 1 H), 1.51-1.31 (m, 4 H), 0.90 (t, *J* = 7.2 Hz, 3 H); ¹³C NMR (125 MHz, CDCl₃) δ 153.4, 151.6, 148.5, 122.3, 120.1, 72.0, 37.0, 27.9, 22.6, 14.1; IR (neat) 3272 (bs), 2956 (m), 2929 (m), 2859 (s), 1585 (m), 1463 (s), 1401 (m), 1082 (m), 1021 (s), 845 (s) cm⁻¹; HRMS (CI⁺) *m/z* calculated for C₁₀H₁₅BrNO [M+H]⁺ 244.0259, found 244.0259.


**3-Butyl-1,1-dimethyl-1,3-dihydro-[1,2]oxasilolo[3,4-c]pyridine (3a):**

1-(3-bromopyridin-4-yl)pentan-1-ol (**S7**) (1.00 g, 4.10 mmol, 1.00 equiv) was dissolved in THF (27.4 mL) and *n*-BuLi (4.4 mL, 2.05 M in hexanes, 9.02 mmol, 2.20 equiv) was added dropwise at -78°C . The reaction mixture was stirred for 30 min, followed by the addition of Me_2SiHCl (0.98 mL, 9.02 mmol, 2.20 equiv) at -78°C . The resulting reaction mixture was allowed to warm to room temperature and was stirred for 12 h. The reaction mixture was then quenched with H_2O (20 mL) with evolution of H_2 gas observed. The reaction mixture was allowed to stir for 5 h and extracted with EtOAc (3×20 mL). The combined organic layers were collected and washed with 1M aq. HCl (3×40 mL). The acidic aqueous layers were collected and neutralized to pH 8-9 with 1M aq. NaOH (120 mL) producing a white turbid mixture that was extracted with Et_2O (3×75 mL). The combined organic layers were dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography on silica (50% EtOAc in hexanes) followed by Kugelrohr distillation (120°C , 0.01 mmHg) provided **3a** (281 mg, 1.27 mmol, 31% yield) as an off-white amorphous solid: R_f 0.3 (70% EtOAc in hexanes); $^1\text{H NMR}$ (500MHz, CDCl_3) δ 8.76 (bs, 1 H), 8.56 (d, $J = 5.0$ Hz, 1H), 7.13 (d, $J = 5.2$ Hz, 1H), 5.20 (dd, $J = 7.5$ Hz, $J = 3.5$ Hz, 1 H), 1.90 (m, 1 H), 1.59 (m, 1 H), 1.45-1.29 (m, 4 H), 0.89 (t, $J = 7.0$ Hz, 3 H), 0.44 (s, 3 H), 0.41 (s, 3 H); $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ 152.6, 143.9, 135.3, 130.8, 129.9, 128.7, 127.9, 127.3, 127.3, 123.9, 84.2, 1.4, 0.7; **IR** (CH_2Cl_2) 3057 (m), 2964 (m), 2874 (m), 1447 (m), 1265 (s), 1181 (m), 1136 (m), 1042 (s), 1016 (s), 862 (s), 822 (s), 793 (s), 740 (s), 702 (s), 656 (m) cm^{-1} ; **HRMS** (CI^+) m/z calculated for $\text{C}_{12}\text{H}_{20}\text{NOSi}$ [$\text{M}+\text{H}$] $^+$ 222.1236, found 222.1233.

**3-Butyl-1,1-diethyl-1,3-dihydro-[1,2]oxasilolo[3,4-c]pyridine (3b):**


1-(3-Bromopyridin-4-yl)pentan-1-ol (**S7**) (1.00 g, 4.10 mmol, 1.00 equiv) was dissolved in THF (27.4 mL) and *n*-BuLi (4.4 mL, 2.05 M in hexanes, 9.02 mmol, 2.20 equiv) was added dropwise at -78°C . The reaction mixture was stirred for 30 min, followed by the addition of Et_2SiHCl (1.12 mL, 9.02 mmol, 2.20 equiv) at -78°C . The resulting reaction mixture was allowed to warm to room temperature and stirred for 12 h. The slurry was quenched with H_2O (20 mL) with vigorous evolution of H_2 gas observed. The solution was allowed to stir for 5 h and extracted

with EtOAc (3 × 20 mL). The combined organic layers were collected and washed with 1M aq. HCl (3 × 40 mL). The acidic aqueous layers were collected and neutralized to pH 8-9 with 1M aq. NaOH (120 mL) producing a white turbid mixture that was extracted with Et₂O (3 × 75 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on silica (30% EtOAc in hexanes) followed by Kugelrohr distillation (130 °C, 0.01 mmHg) provided **3b** (540 mg, 2.17 mmol, 43% yield) as a yellow oil: **R_f** 0.2 (40% EtOAc in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 8.69 (bs, 1 H), 8.50 (d, *J* = 5.0 Hz, 1H), 7.09 (d, *J* = 5.0 Hz, 1H), 5.12 (dd, *J* = 7.8 Hz, *J* = 3.4 Hz, 1 H), 1.84 (m, 1 H), 1.51 (m, 1 H), 1.44-1.24 (m, 4 H), 0.93 (t, *J* = 7.5 Hz, 3 H), 0.88-0.77 (m, 10 H); ¹³C NMR (125 MHz, CDCl₃) δ 162.9, 152.6, 149.7, 129.0, 117.7, 81.3, 38.0, 27.4, 22.6, 13.9, 7.09, 6.93, 6.55, 6.26; IR (CH₂Cl₂) 3057 (m), 2964 (m), 2874 (m), 1447 (m), 1265 (s), 1181 (m), 1136 (m), 1042 (s), 1016 (s), 862 (s), 822 (s), 793 (s), 740 (s), 702 (s), 656 (m) cm⁻¹; HRMS (CI⁺) *m/z* calculated for C₁₄H₂₄NOSi [M+H]⁺ 250.1549, found 250.1547.




(2-Bromophenyl)(3-(dimethylamino)phenyl)methanol (S8): In a two-necked 200 mL round-bottom flask, Mg turnings (5.30 g, 216 mmol, 10.0 equiv) were flame-dried under vacuum. Upon cooling to room temperature, THF (55 mL), 3-bromo-*N,N*-dimethylaniline (6.50 g, 32.4 mmol, 1.50 equiv) as a solution in THF (55 mL) and a crystal of I₂ were added. The flask was fitted with a reflux condenser and the solution vigorously stirred. Once the resulting exotherm had subsided, the reaction mixture was heated to reflux (70 °C) for 1 h. This Grignard reagent was then allowed to cool to room temperature and added to a cooled solution of 2-bromobenzaldehyde (4.00 g, 21.6 mmol, 1.00 equiv) in THF (32 mL) at 0 °C via cannula over 10 min. The reaction mixture was then allowed to reach room temperature and stirred for 12 h and quenched with sat. aq. NH₄Cl (75 mL). The aqueous phase was then extracted with EtOAc (3 × 50 mL). The combined organic layers were washed with brine, dried (Na₂SO₄) and concentrated under reduced pressure. Flash chromatography on silica (1% to 5% EtOAc in hexanes) provided the desired (2-bromophenyl)(3-(dimethylamino)phenyl)methanol (**S8**) (6.36 g, 20.8 mmol, 96% yield) a yellow viscous oil: **R_f** 0.4 (20% EtOAc in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.59 (dd, *J* = 7.7, 1.6 Hz, 1 H), 7.53 (dd, *J* = 7.9, 1.1 Hz, 1 H), 7.33 (td, *J* = 7.5, 1.0 Hz, 1 H), 7.20 (t, *J* = 7.9, 1 H), 7.13 (td, *J* = 7.7, 1.6 Hz, 1 H), 6.84 (bs, 1 H), 6.72 (d, *J* = 7.5, 1 H), 6.66 (dd, *J* = 8.2, 2.2 Hz, 1 H), 6.16 (s, 1 H), 2.93 (s, 6 H), 2.38 (bs, 1 H);

^{13}C NMR (125 MHz, CDCl_3) δ 150.8, 143.3, 142.9, 132.9, 129.3, 129.1, 128.7, 127.8, 123.1, 115.3, 112.2, 111.4, 75.3, 40.8; **IR** (neat) 3392 (bs), 2917 (s), 2848 (s), 1604 (s), 1498 (s), 1437 (s), 1353 (s), 1018 (m), 996 (s), 744 (s) cm^{-1} ; **HRMS** (ES^+) m/z calculated for $\text{C}_{15}\text{H}_{17}\text{NOBr}$ $[\text{M}+\text{H}]^+$ 306.0494, found 306.0492.




3-(1,1-Dimethyl-1,3-dihydrobenzo[c][1,2]oxasilol-3-yl)-N,N-dimethylaniline (3c): (2-Bromophenyl)(3-(dimethylamino)phenyl)methanol (**S8**) (6.90 g, 22.5 mmol, 1.00 equiv) was dissolved in THF (90 mL) and *n*-BuLi (20.8 mL, 2.40 M in hexanes, 50.0 mmol, 2.20 equiv) was added dropwise at -78°C . The reaction mixture was stirred for 45 min, followed by the addition of Me_2SiHCl (5.60 mL, 50.0 mmol, 2.20 equiv) at -78°C . The resulting reaction mixture was allowed to warm to room temperature and stirred for 12 h. The slurry was quenched with H_2O (75 mL) with vigorous evolution of H_2 gas observed. The reaction mixture was allowed to stir for 5 h and extracted with EtOAc (3×50 mL). The combined organic layers were collected and washed with 3M aq. HCl (3×50 mL). The acidic aqueous layers were collected and neutralized to pH 8-9 with 1M aq. NaOH producing a white turbid mixture that was extracted with Et_2O (3×75 mL). The combined organic layers were dried (MgSO_4), filtered and concentrated under reduced pressure. Flash chromatography on silica (5% to 10% EtOAc in hexanes) followed by Kugelrohr distillation (150°C , 0.01 mmHg) provided **3c** (2.8 g, 9.88 mmol, 44% yield) as an orange viscous oil: R_f 0.1 (40% EtOAc in hexanes); ^1H NMR (500 MHz, CDCl_3) δ 7.63 (dd, $J = 6.2, 1.6$ Hz, 1 H), 7.33 (m, 2 H), 7.22 (t, $J = 7.6$ Hz, 1 H), 7.13 (d, $J = 7.4$ Hz, 1 H), 6.71-6.62 (m, 3 H), 6.15 (s, 1 H), 2.94 (s, 6 H), 0.56 (s, 3 H), 0.48 (s, 3 H); ^{13}C NMR (125 MHz, CDCl_3) δ 152.8, 150.88, 144.71, 135.04, 130.6, 129.9, 129.3, 127.3, 127.1, 123.8, 115.6, 112.2, 111.6, 84.7, 40.7, 1.51, 0.65; **IR** (neat) 2953 (bs), 2874 (bs), 2802 (s), 1605 (s), 1499 (s), 1440 (s), 1353 (s), 1251 (s), 1135 (s), 1066 (s), 1043 (s), 995 (s), 870 (s), 789 (s), 743 (s) cm^{-1} ; **HRMS** (ES^+) m/z calculated for $\text{C}_{17}\text{H}_{22}\text{NOSi}$ $[\text{M}+\text{H}]^+$ 284.1471, found 284.1476.




(2-Bromophenyl)(4-(dimethylamino)phenyl)methanol (S9): In a two-necked 500 mL round-bottom flask, Mg turnings (10.5 g, 432 mmol, 10.0 equiv) were

flame-dried under vacuum. Upon cooling to room temperature, THF (72 mL), 4-bromo-*N,N*-dimethylaniline (12.9 g, 64.9 mmol, 1.50 equiv) as a solution in THF (72 mL) and a crystal of I₂ were added. The flask was fitted with a reflux condenser and the solution was vigorously stirred. Once the resulting exotherm had subsided, the reaction mixture was heated to reflux (70 °C) for 1 h. This Grignard reagent was then allowed to cool to room temperature and added to a cooled solution of 2-bromobenzaldehyde (8.00 g, 43.2 mmol, 1.00 equiv) in THF (87 mL) at 0 °C via cannula over 10 min. The reaction was then allowed to reach room temperature and stirred for 12 h and quenched with sat. aq. NH₄Cl (100 mL). The aqueous phase was then extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine, dried (Na₂SO₄) and concentrated under reduced pressure. Flash chromatography on silica (3% to 10% EtOAc in hexanes) provided of the desired (2-bromophenyl)(4-(dimethylamino)phenyl)methanol (**S9**) (12.6 g, 41.2 mmol, 95% yield) as a blue crystalline solid: **R_f** 0.2 (10% EtOAc in hexanes); **¹H NMR** (500MHz, CDCl₃) δ 7.71 (dd, *J* = 7.7, 1.5 Hz, 1 H), 7.53 (dd, *J* = 7.9, 1.0 Hz, 1 H), 7.37 (td, *J* = 7.6, 1.0 Hz, 1 H), 7.24 (m, 2 H), 7.14 (td, *J* = 7.7, 1.5 Hz, 1 H), 6.69 (m, 2 H), 6.07 (s, 1 H), 2.94 (s, 6 H), 2.58 (bs, 1 H); **¹³C NMR** (125 MHz, CDCl₃) δ 150.3, 143.2, 132.9, 130.3, 128.8, 128.6, 128.4, 128.3, 127.7, 122.8, 112.8, 112.6, 112.3, 74.8, 40.7; **IR** (neat) 3387 (bs), 2887 (m), 2802 (s), 1613 (s), 1521 (s), 1352 (s), 1162 (s), 1017 (m), 810 (s), 758 (s), 741 (s) cm⁻¹; **HRMS** (ES⁺) *m/z* calculated for C₁₅H₁₇NOBr [M+H]⁺ 306.0494, found 306.0497.



4-(1,1-Dimethyl-1,3-dihydrobenzo[c][1,2]oxasilol-3-yl)-*N,N*-dimethylaniline (3f): (2-Bromophenyl)(4-(dimethylamino)phenyl)methanol (**S9**) (2.33 g, 7.61 mmol, 1.00 equiv) was dissolved in THF (25 mL) and *n*-BuLi (7.6 mL, 2.20 M in hexanes, 16.7 mmol, 2.20 equiv) was added dropwise at -78 °C. The reaction mixture was stirred for 45 min, followed by the addition of Me₂SiHCl (2.00 mL, 16.7 mmol, 2.20 equiv) at -78 °C. The resulting reaction mixture was allowed to warm to room temperature and stirred for 12 h. The slurry was quenched with H₂O (50 mL) with vigorous evolution of H₂ gas observed. The reaction mixture was allowed to stir for 5 h and extracted with EtOAc (3 × 25 mL). The combined organic layers were collected and washed with 3M aq. HCl (3 × 25 mL). The acidic aqueous layers were collected and neutralized to pH 8-9 with 1M aq. NaOH producing a white turbid mixture that was extracted with Et₂O (3 × 75 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on

silica (5% Et₂O in hexanes) provided **3f** (795 mg, 2.81 mmol, 37% yield) as an yellow crystalline solid: **R_f** 0.1 (25% EtOAc in hexanes); **¹H NMR** (500 MHz, CDCl₃) δ 7.63 (dd, *J* = 6.0, 1.8 Hz, 1 H), 7.33 (m, 2 H), 7.14 (m, 2 H), 7.06 (m, 1 H), 6.72 (m, 2 H), 6.14 (s, 1 H), 2.95 (s, 6 H), 0.52 (s, 3 H), 0.46 (s, 3 H); **¹³C NMR** (125 MHz, CDCl₃) δ 153.3, 150.5, 135.5, 132.1, 130.6, 129.8, 128.5, 128.3, 127.1, 124.1, 112.7, 84.2, 40.8, 1.57, 0.66; **IR** (neat) 3419 (bs), 2921 (m), 1615 (s), 1523 (s), 1443 (s), 1348 (s), 1251 (s), 1163 (s), 1134 (s), 861 (s), 822 (s), 789 (s), 743 (s) cm⁻¹; **HRMS** (CI⁺) *m/z* calculated for C₁₇H₂₂NOSi [M+H]⁺ 284.1471, found 284.1459.



4-(1,1-Diethyl-1,3-dihydrobenzo[c][1,2]oxasilol-3-yl)-N,N-dimethylaniline (3g): (2-Bromophenyl)(4-(dimethylamino)phenyl)methanol (**S9**) (7.00 g, 22.9 mmol, 1.00 equiv) was dissolved in THF (230 mL), *n*-BuLi (22.9 mL, 2.20 M in hexanes, 50.4 mmol, 2.20 equiv) was added dropwise at -78 °C. The reaction mixture was stirred for 45 min, followed by the addition of Et₂SiHCl (7.00 mL, 50.4 mmol, 2.20 equiv) at -78 °C. The resulting reaction mixture was allowed to warm to room temperature and stirred for 12 h. The slurry was quenched with H₂O (100 mL) with vigorous evolution of H₂ gas observed. The reaction mixture was allowed to stir for 5 h and extracted with EtOAc (3 × 50 mL). The combined organic layers were collected and washed with 3M aq. HCl (3 × 50 mL). The acidic aqueous layers were collected and neutralized to pH 8-9 with 1M aq. NaOH (300 mL) producing a white turbid mixture that was extracted with Et₂O (3 × 100 mL). The combined organic layers were dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography on silica (5% EtOAc in hexanes) provided **3g** (3.6 g, 11.6 mmol, 51% yield) as an yellow oil: **R_f** 0.3 (15% EtOAc in hexanes); **¹H NMR** (500 MHz, CDCl₃) δ 7.64 (dd, *J* = 6.0, 1.5 Hz, 1 H), 7.37-7.30 (m, 2 H), 7.17-7.15 (m, 2 H), 7.06 (m, 1 H), 6.74- 6.70 (m, 2 H), 6.15 (s, 1 H), 2.95 (s, 6 H), 1.08 (m, 3 H), 1.04-0.88 (m, 7 H); **¹³C NMR** (125 MHz, CDCl₃) δ 153.8, 150.4, 133.8, 132.0, 131.2, 129.7, 128.7, 126.8, 124.1, 112.5, 84.3, 40.7, 7.43, 7.11, 6.99, 6.65; **IR** (neat) 2955 (m), 2875 (m), 2800 (s), 1614 (s), 1523 (s), 1444 (m), 1348 (s), 1230 (s), 1162 (s), 1132 (s), 997 (m), 949 (s), 823 (m), 723 (m) cm⁻¹; **HRMS** (ES⁺) *m/z* calculated for C₁₉H₂₆NOSi [M+H]⁺ 312.1784, found 312.1783.


[0081] General Procedure A: To a cooled solution of siloxane (0.81 mmol, 1.8 equiv) in dry Et₂O (1.2 mL) at room temperature was added a solution of PhLi in Bu₂O (0.68 mmol, 1.5

equiv) and allowed to stir for 2 h. After 1.5 h had elapsed following PhLi addition, in a separate flask were combined PdCl₂ (2.50 mg, 0.014 mmol, 0.03 equiv), CuI (8.60 mg, 0.045 mmol, 0.1 equiv) and dpca (6.7 mg, 0.018 mmol, 0.04 equiv) in dry THF (1 mL) at room temperature and stirred for 30 min. The aryl halide (0.45 mmol, 1.0 equiv) was added to the orange slurry, followed by addition of the siloxane/PhLi reaction mixture by cannula (flask rinsed with 0.5 mL of THF). After 2 h at room temperature, the reaction mixture was diluted with Et₂O (2 mL) and quenched according to the siloxane used in the reaction; **1**, **1a-f**, **2a-d** quenched with sat. aq. NH₄Cl (5 mL); **3a-b** and **3d-e** quenched with 1M aq. HCl (5 mL); **3c** and **3f-g** quenched with 3M aq. HCl (5 mL).

[0082] For siloxanes **1**, **1a-f**, **2a-d** the aqueous layer was extracted with Et₂O (3 × 5 mL) and the combined organic layers were dried (MgSO₄), filtered, and concentrated under reduced pressure. Flash chromatography provided the cross-coupled product and the recovered siloxane (where possible as with siloxanes **2a-d**).


[0083] For siloxanes **3a-g**, the organic layer was washed with either 1M or 3M aq. HCl (3 × 5 mL) (according to siloxane, see above), and the acidic aqueous layers were collected. The organic layer was then washed with sat. aq. NaHCO₃ (5 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure. Flash chromatography provided the cross-coupled product. The acidic aqueous layer was then basified to pH 8-9 with 1M aq. NaOH, producing a white turbid mixture that was then extracted with Et₂O 3 × 50 mL). The combined organic layers were dried (MgSO₄), filtered, and concentrated under reduced pressure to provide the recovered siloxane.

Preparation of fluorinated siloxanes.




Procedure: To a cooled solution of (Hexafluoro-2-hydroxyisopropyl)benzene (3.00 g, 12.3 mmol, 1.00 equiv) in hexanes (75 mL, pre-dried over MgSO₄) and Et₂O (60 mL) at 0 °C was added *n*-BuLi (12.6 mL, 2.15 M in hexanes, 27.1 mmol, 2.20 equiv) dropwise under N₂. The reaction mixture was warmed to room temperature and stirred for 30 minutes. The flask was then fitted with a reflux condenser and the reaction mixture was heated to reflux (70 °C) for 16 h. A dark brown solution resulted and the reaction mixture was allowed to cool to room

temperature. The reflux condenser was removed and the reaction mixture was cooled to $-78\text{ }^{\circ}\text{C}$ before either *i*-Pr₂SiHCl (4.60 mL, 27.06 mmol, 2.20 equiv) or Et₂SiHCl was added dropwise under N₂. The reaction mixture was allowed to slowly warm to room temperature and stirred for 12 h. The resulting pale yellow slurry was then quenched with H₂O (75 mL) and the aqueous phase extracted with Et₂O (3 × 40 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography (100% hexanes) provided 2.76 g (63% yield) of the desired diisopropyl siloxane as a white crystalline solid (m.p. $39\text{--}41\text{ }^{\circ}\text{C}$).




Procedure: To a cooled solution of 1,2-Dibromotetrafluorobenzene (5.00 g, 16.2 mmol, 1.00 equiv) in Et₂O (162 mL) at $-78\text{ }^{\circ}\text{C}$ was added *n*-BuLi (7.75 mL, 2.10 M in hexanes, 16.2 mmol, 1.00 equiv) dropwise over 30 min under N₂. The reaction mixture was stirred for another 30 minutes before pentanal (1.90 mL, 17.9 mmol, 1.10 equiv) was added dropwise. The reaction mixture was stirred for 5 h at $-78\text{ }^{\circ}\text{C}$ before it was warmed to $0\text{ }^{\circ}\text{C}$ and quenched with H₂O (75 mL). The aqueous phase extracted with Et₂O (3 × 30 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Flash chromatography (2% to 5% ethyl acetate in hexanes) provided 4.34 g (85% yield) of the perfluorinated bromo benzylic alcohol.

The resulting perfluorinated bromo benzylic alcohol (332 mg, 1.05 mmol, 1.00 equiv) was dissolved in THF (5.5 mL) and *n*-BuLi (1.1 mL, 2.10 M in hexanes, 2.21 mmol, 2.10 equiv) was added dropwise at $-78\text{ }^{\circ}\text{C}$. The reaction mixture was stirred for 45 min followed by the addition of either *i*Pr₂SiHCl (0.40 mL, 2.32 mmol, 2.20 equiv) or Et₂SiHCl at $-78\text{ }^{\circ}\text{C}$. The resulting reaction mixture was allowed to warm to room temperature and stirred for 12 h. The reaction mixture was quenched by addition of H₂O (20 mL) and stirred for 1 h. The aqueous phase was then extracted with Et₂O (2 × 15 mL) and the combined organic layers were washed with brine, dried (MgSO₄), and concentrated under reduced pressure. Flash chromatography (100% hexanes) provided the desired siloxane (223 mg, 61% yield) as a light yellow oil.

**4-Methoxy-1,1'-biphenyl (S10):**

Following General Procedure A, the product was purified by chromatography on SiO₂ (1% EtOAc in hexanes) to afford **S7** (80.0 mg, 0.43 mmol, 96% with siloxane **2a**; 81.0 mg, 0.44 mmol, 98% with siloxane **2b**; 81.0 mg, 0.44 mmol, 98% with siloxane **3e**; 81.0 mg, 0.44 mmol, 98% with siloxane **3g**) as a colorless solid. Analytical data matches that which has been previously reported for **S10**:¹¹ *R_f* 0.5 (1% EtOAc in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.57-7.53 (m, 4 H), 7.42 (t, *J* = 7.6 Hz, 2 H), 7.31 (t, *J* = 7.4 Hz, 1 H), 6.99 (d, *J* = 8.7 Hz, 2 H), 3.86 (s, 3 H); ¹³C NMR (125 MHz, CDCl₃) δ 159.3, 141.0, 133.9, 128.9, 128.3, 126.9, 126.8, 114.3, 55.5.

**[1,1'-Biphenyl]-4-carbonitrile (S11):**


Following General Procedure A, the product was purified by chromatography on SiO₂ (1% EtOAc in hexanes) to afford **S8** (74.0 mg, 0.41 mmol, 92% with siloxane **2a**; 73.0 mg, 0.40 mmol, 91% with siloxane **2b**; 77.0 mg, 0.43 mmol, 96% with siloxane **3e**; 75.0 mg, 0.42 mmol, 94% with siloxane **3g**) as a colorless solid. Analytical data matches that which has been previously reported for **S11**:¹² *R_f* 0.3 (1% EtOAc in hexanes); ¹H NMR (500 MHz, CDCl₃) 7.73 (d, *J* = 8.3 Hz, 2 H), 7.69 (d, *J* = 8.4 Hz, 2 H), 7.59 (d, *J* = 7.4 Hz, 2 H), 7.49 (t, *J* = 7.2 Hz, 2 H), 7.43 (t, *J* = 7.3 Hz, 1 H); ¹³C NMR (125 MHz, CDCl₃) δ 145.8, 139.3, 132.8, 129.3, 128.8, 127.9, 127.4, 119.1, 111.1.

[0084] General Procedure B: To a solution of alkenyl iodide or aryl iodide (0.68 mmol, 1.50 equiv) in Et₂O (1 mL) at -78 °C was added *t*-BuLi in pentane (1.35 mmol, 3.00 equiv), and a white or yellow (depending on alkenyl iodide or aryl iodide) slurry developed. The reaction mixture was allowed to stir for 40 min at -78 °C and 20 min at room temperature, at which time a solution of siloxane (0.81 mmol, 1.80 equiv) in THF (0.5 mL + 0.2 mL rinse) was added and allowed to stir at room temperature for 2 h. After 1.5 h had elapsed following siloxane addition, in a separate flask were combined PdCl₂ (2.50 mg, 0.014 mmol, 0.03 equiv), CuI (8.60 mg, 0.045 mmol, 0.1 equiv) and dpca (6.7 mg, 0.018 mmol, 0.04 equiv) in dry THF (1 mL) at room temperature and stirred for 30 min. The alkenyl halide or aryl iodide (0.45 mmol, 1.00 equiv) was added as a solution in THF (0.3 mL) to the orange slurry, immediately followed by addition of the siloxane reaction mixture by cannula (flask rinsed with 0.5 mL of THF). After 2-12 h at room temperature, the reaction mixture was diluted with Et₂O (2 mL) and quenched according to the siloxane used in the reaction; **1**, **1a-f**, **2a-d** quenched with sat. aq. NH₄Cl (5

mL); **3a-b** and **3d-e** quenched with 1M aq. HCl (5 mL); **3c** and **3f-g** quenched with 3M aq. HCl (5 mL).

[0085] For siloxanes **1**, **1a-f**, **2a-d**, the aqueous layer was extracted with Et₂O (3 × 5 mL) and the combined organic layers were dried (MgSO₄), filtered, and concentrated under reduced pressure. Flash chromatography provided the cross-coupled product and the recovered siloxane (where possible as with siloxanes **2a-d**).

[0086] For siloxanes **3a-g**, the organic layer was washed with either 1M or 3M aq. HCl (3 × 5 mL) (according to siloxane, see above), and the acidic aqueous layers were collected. The organic layer was then washed with sat. aq. NaHCO₃ (5 mL), dried (MgSO₄), filtered, and concentrated under reduced pressure. Flash chromatography provided the cross-coupled product. The acidic aqueous layer was then basified to pH 8-9 with 1M aq. NaOH, producing a white turbid mixture which was then extracted with Et₂O 3 × 50 mL). The combined organic layers were dried (MgSO₄), filtered, and concentrated under reduced pressure to provide the recovered siloxane.




methoxyphenyl)allyl)oxy)dimethylsilane (S12): Following General Procedure B, the product was purified by chromatography on SiO₂ (1% EtOAc in hexanes) to afford **S9** in >20:1 E/Z ratio as a colorless oil (110 mg, 0.40 mmol, 88% with siloxane **2a**; 115 mg, 0.41 mmol, 92% with siloxane **2b**; 124 mg, 0.44 mmol, 99% with siloxane **3e**; 121 mg, 0.44 mmol, 97% with siloxane **3g**). Analytical data matches that which has been previously reported for **S12**:¹³ *R*_f 0.3 (1% EtOAc in hexanes); ¹H NMR (500 MHz, CDCl₃) δ 7.31 (d, *J* = 8.8 Hz, 2 H), 6.85 (d, *J* = 8.8 Hz, 2 H), 6.53 (d, *J* = 16.0 Hz, 1 H), 6.15 (dt, *J* = 5.2, 16.0 Hz, 1 H), 4.33 (d, *J* = 5.3 Hz, 2 H), 3.81 (s, 3 H), 0.94 (s, 9 H), 0.11 (s, 6 H); ¹³C NMR (125 MHz, CDCl₃) δ 159.2, 130.1, 129.3, 127.7, 127.1, 114.07, 64.2, 55.4, 26.1, 18.6, -4.97.




1-yl)oxy)silane (S13): Following General Procedure B, the product was purified by chromatography on SiO₂ (1% EtOAc in hexanes) to afford **S13** in >20:1 E/Z ratio as a colorless oil (125 mg, 0.41 mmol, 92% with siloxane **2a**; 124 mg, 0.41 mmol, 91% with siloxane **2b**; 131 mg, 0.43 mmol, 96% with siloxane **3e**; 135 mg, 0.44 mmol, 99% with siloxane **3g**): *R*_f 0.3 (1%

EtOAc in hexanes); $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.31-7.25 (m, 2 H), 7.21-7.17 (m, 3 H), 6.19 (dd, $J = 10.6, 15.0$ Hz, 1 H), 6.08 (dd, $J = 10.6, 15.0$ Hz, 1 H), 5.75-5.62 (m, 2H), 4.21 (d, $J = 5.5$ Hz, 2 H), 2.72 (t, $J = 7.5$ Hz, 2 H), 2.41 (q, $J = 7.5$ Hz, 2 H), 0.93 (s, 9 H), 0.83 (s, 6 H); $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ 141.9, 133.4, 132.4, 130.8, 130.4, 130.3, 129.5, 128.6, 128.5, 126.0, 63.8, 63.7, 35.9, 34.6, 29.9, 26.1, 18.6, -5.01. **IR** (neat) 3026 (m), 2954 (m), 2930 (s), 2856 (s) 1468 (m), 1254 (s), 1109 (m), 1066 (bs), 988 (s), 837 (s), 776 (s) cm^{-1} ; **HRMS** (CI^+) m/z calculated for $\text{C}_{15}\text{H}_{21}\text{OSi} [\text{M}-\text{C}_4\text{H}_9]^+$ 245.1362, found 245.1368.



(E)-1-(Hept-1-en-1-yl)-4-methoxybenzene (S14):


Following General Procedure B, the product was purified by chromatography on SiO_2 (1% EtOAc in hexanes) to afford **S14** in >20:1 E/Z ratio as a colorless oil (89.0 mg, 0.44 mmol, 97% with siloxane **2a**; 88.0 mg, 0.43 mmol, 96% with siloxane **2b**; 87.0 mg, 0.43 mmol, 95% with siloxane **3e**; 87.0 mg, 0.42 mmol, 94% with siloxane **3g**) as a colorless solid. Analytical data matches that which has been previously reported for **S14**:¹⁴ **R_f** 0.3 (1% EtOAc in hexanes); $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.28 (dd, $J = 2.0, 6.5$ Hz, 2 H), 6.84 (dd, $J = 2.1, 6.6$ Hz, 2 H), 6.32 (d, $J = 15.7$ Hz, 1 H), 6.09 (dt, $J = 7.2, 15.8$ Hz, 1 H), 3.80 (s, 3 H), 2.18 (qd, $J = 1.3, 7.3$ Hz, 2 H), 1.46 (qn, $J = 7.2$ Hz, 2 H), 1.40-1.37 (m, 4 H), 0.90 (t, $J = 7.1$ Hz, 3 H); $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ 158.6, 131.2, 129.6, 129.5, 127.1, 114.2, 55.7, 33.5, 31.4, 29.5, 22.7, 14.1.



(Z)-1-(Hept-1-en-1-yl)-4-methoxybenzene (S15):

Following General Procedure A, the product was purified by chromatography on SiO_2 (1% EtOAc in hexanes) to afford **S15** in >20:1 Z/E ratio as a colorless oil (80.0 mg, 0.39 mmol, 87% with siloxane **2a**; 82.0 mg, 0.40 mmol, 89 % with siloxane **2b**; 88.0 mg, 0.43 mmol, 96% with siloxane **3e**, 87.0 mg, 0.42 mmol, 94% with siloxane **3g**). Analytical data matches that which has been previously reported for **S15** (Denmark, S. E.; Sweis, R. F. J. Am. Chem. Soc. 2001, 123, 6439). **R_f** 0.3 (1% EtOAc in hexanes); $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.23 (d, $J = 8.6$ Hz, 2 H), 6.87 (d, $J = 8.8$ Hz, 2 H), 6.33 (d, $J = 11.7$ Hz, 1 H), 5.55 (dt, $J = 7.3, 11.6$ Hz, 1 H), 3.82 (s, 3 H), 2.29 (qd, $J = 1.3, 7.3$ Hz, 2 H), 1.45 (qn, $J = 7.6$ Hz, 2 H), 1.36-1.28 (m, 4 H), 0.90 (t, $J = 7.2$ Hz, 3 H); $^{13}\text{C NMR}$ (125 MHz, CDCl_3) δ 158.6, 131.7, 130.5, 130.1, 128.4, 113.5, 55.1, 31.4, 29.7, 28.9, 22.6, 14.2.

Preparation of Siloxane Polymers




[0087] To a stirred solution of aldehyde **20** (5.42 g, 44.4 mmol, mixture of *endo* and *exo* isomers) in THF (200 mL) at 0 °C was added solution of PhMgBr in Et₂O (17.8 mL, 3.00 M, 53.3 mmol) dropwise. The solution was stirred for 15 h at room temperature and then diluted with sat. aq. NH₄Cl (50 mL). The aqueous phase was extracted with Et₂O and the combined organic layers were washed with brine, dried with MgSO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography on SiO₂ (5% Et₂O/ hexanes) to afford **21** (7.72 g, 87%, mixture of 4 diastereomers) as a pale yellow oil.

[0088] To a stirred solution of alcohol **21** (6.83 g, 34.1 mmol) in hexanes (125 mL, dried over MgSO₄) and Et₂O (100 mL) at 0 °C was added *n*-BuLi in hexanes 35.8 mL, 2.10 M, 75.1 mmol) dropwise. The solution was stirred at room temperature for 1 h, at which time the solution was heated to reflux for 23 h. The solution was then cooled to -78 °C and Me₂SiHCl (8.16 mL, 75.1 mmol) was added dropwise. The solution was warmed to room temperature, and after 23 h, H₂O (100 mL) was added and the biphasic mixture was stirred for another 24 h. The aqueous phase was extracted with hexanes and the combined organic layers were washed with brine, dried with MgSO₄, filtered and concentrated under reduced pressure. The crude product was purified by column chromatography on SiO₂ (0-1% Et₂O/ hexanes; the column was slurry-packed using SiO₂ in H₂O, which was then washed with acetone, EtOAc, and hexanes, successively, prior to chromatographic separation), followed by Kugelrohr distillation at 120-130 °C (0.025 mmHg) to afford **22** as a colorless oil.


[0089] Grubbs' 1st generation catalyst (6.1 mg, 7.4 μmol) was dissolved in CH₂Cl₂ (0.5

mL) and the solution was stirred for 30 min. The catalyst solution was then introduced via cannula to another flask containing **22** (378 mg, 1.48 mmol) in CH₂Cl₂ (1 mL), and the resulting solution was then stirred at room temperature. After 17 h, the reaction mixture was heated to reflux. After 19 h, the reaction was quenched with ethylvinylether (0.5 mL) in CH₂Cl₂ (1.5 mL), and heating of the mixture at 50 °C was continued for another 2 h. The solution was then cooled to room temperature and diluted with CH₂Cl₂ (5 mL). Tris(hydroxymethyl)phosphine (53.9 mg, 0.3 eq.) in H₂O (10 mL) was added to the solution and the biphasic mixture was stirred vigorously for 15 min. The aqueous phase was extracted with CH₂Cl₂ and the combined organic layers were washed with brine, dried with MgSO₄, filtered and concentrated under reduced pressure. The obtained white solid was washed extensively with MeCN and dried under vacuum to afford **23** (363 mg, 96%, M_n = 73689, PDI = 1.272) as a white solid. n is about 250.

Polymer-Mediated Cross-Coupling



[0090] To a stirred solution of polymer **23** (196 mg, see above) in THF (4 mL) at -78 °C was added PhLi in dibutylether (479 μL, 1.23 M, 0.575 mmol) dropwise, followed by THF (6 mL). The reaction mixture was stirred for 3 h at room temperature and a colorless slurry resulted. A mixture of 4-iodobenzonitrile (87.7 mg, 0.383 mmol), PdCl₂ (2.0 mg, 0.011 mmol, 3 mol%), CuI (7.3 mg, 0.038 mmol), and dpca (N-[2-(Diphenylphosphino)benzylidene]cyclohexylamine, 5.7 mg, 0.015 mmol) was added to the flask in a single portion and the slurry was stirred for 2 d at room temperature. The reaction mixture was diluted with sat. aq. NH₄Cl (5 mL), and the aqueous phase was extracted with Et₂O. The combined organic layers were washed with brine, dried with MgSO₄, filtered and concentrated under reduced pressure. The resulting residue was washed extensively with acetonitrile, and the washings were combined and concentrated to obtain the crude product, which was purified by chromatography on SiO₂ (1% Et₂O/ hexanes) to afford the product (26.1 mg, 0.146 mmol, 38%) as a white solid.

Alternative procedure for polymer-mediated cross-coupling

[0091] To a cooled solution of siloxane polymer in THF (0.193 g, 0.752 mmol, 3.0 equiv, 10 mg/mL) at -78 °C was added PhLi in Bu₂O (348 µL, 1.8 M, 0.627 mmol, 2.5 equiv) dropwise. The reaction mixture was allowed to warm to rt and was stirred for 3 h, and a white, cloudy solution developed. A solid mixture of PdCl₂ (1.3 mg, 7.5 µmol, 0.03 equiv), CuI (4.8 mg, 0.025 mmol, 0.1 equiv), and dpca (3.7 mg, 0.01 mmol, 0.04 equiv) was combined and added to the reaction flask, followed by addition of 4-iodobenzonitrile (56.8 g, 0.248 mmol, 1.0 equiv). The obtained reaction mixture was stirred vigorously at rt. Care should be taken so that all reactants are submerged in THF and no solid is deposited on the side of the flask. After 2 h, the reaction mixture was quenched with sat. aq. NH₄Cl (5 mL), followed by addition of d.i. H₂O (5 mL). The organic layer was collected and the aqueous layer was extracted with Et₂O (2 x 10 mL). The combined organic layers were washed with brine, dried with MgSO₄, and concentrated *in vacuo* to 1-2 mL in volume. The obtained concentrated solution was added dropwise into a vigorously stirred solution of CH₃CN (250 mL). The precipitated polymer was filtered and the supernatant was concentrated *in vacuo* to provide the crude product. Following removal from the supernatant, the polymer was re-dissolved in DCM (20 mL) and filter through a glass fritted funnel to remove insoluble particles, if there is any. The obtained solution was then concentrated *in vacuo* to provide the recovered polymer (192 mg, 99 % recovery). The crude product was purified by chromatography on SiO₂ (2 % Et₂O/ hexanes) to afford 4-cyanobiphenyl (40.3 mg, 0.225 mmol, 91 %) as a colorless solid.

General Cross-Coupling Reaction Procedure:

[0092] To a cooled solution of siloxane (117 mg, 0.805 mmol) in dry THF (1.0 mL) at -78 °C was added a solution of PhLi in Bu₂O (0.67 mmol). The reaction mixture was allowed to warm to room temperature and was stirred for 1 h. After 30 min had elapsed following PhLi

addition, in a separate flask were combined PdCl_2 (2.4 mg, 0.013 mmol), CuI (8.6 mg, 0.045 mmol) and dpca (6.8 mg, 0.018 mmol) in dry THF (1.0 mL) at room temperature and stirred for 30 min. The aryl halide (0.45 mmol) was added to the orange slurry, followed by cannulation of the siloxane/ PhLi reaction mixture (flask rinsed with 0.5 mL THF). After 2 h at room temperature, the reaction mixture was diluted with Et_2O (2 mL), quenched with sat. aq. NH_4Cl (5 mL) and extracted with Et_2O (3 x 5 mL). The combined organic layers were dried (MgSO_4), filtered and concentrated under reduced pressure. The cross-coupled product was purified using column chromatography on SiO_2 .

Siloxane-mediated cross-coupling of 4-chloroanisole and phenyllithium using palladium


[0093] To a small pear-shaped flask containing a stirring solution of siloxane **2a** (223.6 mg, 0.5 mmol, 1.8 equiv) and THF (0.35 mL) at -78°C under inert atmosphere, was added PhLi (1.75 M in dibutyl ether, 0.43 mL, 0.75 mmol, 1.5 equiv) dropwise via syringe. The resulting yellow solution (viscous when cold) was allowed to reach room temperature and stirred for 2 h (nucleophile solution). Another flask containing $\text{Pd}(\text{OAc})_2$ (11.2 mg, 0.05 mmol, 10 mol %) and XPhos (47.7 mg, 0.1 mmol, 20 mol %) under nitrogen atmosphere was charged with THF (2 mL). This mixture was stirred at room temperature for 20 minutes, turning dark red (catalyst solution). Then, to a round-bottomed flask containing 4-chloroanisole (71.0 mg, 0.5 mmol, 1.0 equiv) under nitrogen atmosphere, were added sequentially catalyst solution and nucleophile solution via syringe and the resulting mixture was stirred at room temperature overnight. The solution was concentrated *in vacuo* and filtered through a short plug of silica gel eluting with ether. The filtrate was again concentrated *in vacuo* and subjected to column chromatography eluting with ether/hexanes (0–5%) to afford 4-methoxy-biphenyl (69 mg, 75% yield) with 87% purity as determined by $^1\text{H-NMR}$. Impurities include: biphenyl and residual siloxane.

Alternative procedure for siloxane-mediated cross-coupling of 4-chloroanisole and phenyllithium.


[0094] To a small pear-shaped flask containing a stirring solution of siloxane **2a** (447.2 mg, 1.8 mmol, 1.8 equiv) and THF (0.7 mL) at -78°C under inert atmosphere, was added PhLi (1.75 M in dibutyl ether, 0.86 mL, 1.5 mmol, 1.5 equiv) dropwise via syringe. The resulting yellow solution (viscous when cold) was allowed to reach room temperature and stirred for 2 h (nucleophile solution). Another flask containing $\text{Pd}(\text{OAc})_2$ (22.4 mg, 0.1 mmol, 10 mol %) and XPhos (95.4 mg, 0.2 mmol, 20 mol %) under nitrogen atmosphere was charged with THF (2 mL). This mixture was stirred at room temperature for 20 minutes, turning dark red (catalyst solution). Then, to a round-bottomed flask containing 4-chloroanisole (142.6 mg, 1.0 mmol, 1.0

equiv) under nitrogen atmosphere, were added sequentially catalyst solution via syringe and nucleophile solution via cannula and the resulting mixture was stirred at room temperature overnight. The solution was concentrated *in vacuo* and the residue was directly subjected to column chromatography eluting with dichloromethane/hexanes (1–2%) to afford the product (127.1 mg, 69% yield).

Carbon-nitrogen bond formation



[0095] To a cooled solution of siloxane **2a** (112.5 mg, 0.453 mmol, 1.8 equiv) in 4 mL THF at -78 °C was added PhLi in Bu₂O (210 μL, 0.377 mmol, 1.8 M, 1.5 equiv), dropwise. The reaction mixture was allowed to warm to room temperature (rt) and was stirred for 2 h. A solid mixture of CuI (4.8 mg, 0.025 mmol, 0.1 equiv) and dpca (9.3 mg, 0.025 mmol, 0.1 equiv) was combined and added to the reaction flask, followed by addition of piperidines-1-yl benzoate (51.7 mg, 0.252 mmol, 1.0 equiv). The obtained reaction mixture was stirred at rt. After 2 h, the reaction mixture was quenched with sat. aq. NH₄Cl (5 mL), followed by addition of d.i. H₂O (5 mL). The organic layer was collected and the aqueous layer was extracted with Et₂O (2 x 10 mL). The combined organic layers were washed with brine, dried with MgSO₄, and concentrated *in vacuo*. Flash chromatography on silica gel (2% Et₂O/hexanes) afforded 1-phenyl-piperidine (37.7 mg, 0.234 mmol, 93%) as a colorless oil, and recovered siloxane **2a** (97.9 mg, 0.394 mmol, 87%).

Polymer-mediated carbon-nitrogen cross-coupling

[0096] To a cooled solution of siloxane polymer in THF (0.131 g, 0.512 mmol, 3.0 equiv, 10 mg/mL) at -78°C was added PhLi in Bu₂O (237 μL , 1.8 M, 0.627 mmol, 2.5 equiv) dropwise. The reaction mixture was allowed to warm to rt and was stirred for 3 h, and a white, cloudy solution developed. A solid mixture of CuI (1.6 mg, 0.00855 mmol, 0.05 equiv), and Johnphos (5.1 mg, 0.0171 mmol, 0.1 equiv) was combined and added to the reaction flask, followed by addition of O-benzoyl-N,N-dibenzylhydroxylamine (54.3 mg, 0.171 mmol, 1.0 equiv). The obtained reaction mixture was stirred vigorously at rt. Care should be taken so that all reactants are submerged in THF and no solid is deposited on the side of the flask. After 2 h, the reaction mixture was quenched with sat. aq. NH₄Cl (5 mL), followed by addition of d.i. H₂O (5 mL). The organic layer was collected and the aqueous layer was extracted with Et₂O (2 x 10 mL). The combined organic layers were washed with brine, dried with MgSO₄, and concentrated *in vacuo* to 1-2 mL in volume. The obtained concentrated solution was added dropwise into a vigorously stirred solution of CH₃CN (250 mL). The precipitated polymer was filtered and the supernatant was concentrated *in vacuo* to provide the crude product. Following removal from the supernatant, the polymer was re-dissolved in DCM (20 mL) and filter through a glass fritted funnel to remove insoluble particles, if there is any. The obtained solution was then concentrated *in vacuo* to provide the recovered polymer (114.2 mg, 87 % recovery). The crude product was purified by chromatography on SiO₂ (1 % Et₂O/ hexanes) to afford N,N-dibenzylaniline (40.8 mg, 0.149 mmol, 87 %) as a colorless solid.


References:

- ¹ Nakao, Y.; Imanaka, H.; Chen, J.; Yada, A.; Hiyama, T. *J. Organomet. Chem.* **2007**, 692, 585.
- ² Nakao, Y.; Takeda, M.; Matsumoto, T.; Hiyama, T. *Angew. Chem. Int. Ed.* **2010**, 49, 4447.
- ³ Spino, C.; Gund, V. G.; Nadeau, C. *J. Comb. Chem.* **2005**, 7, 345.
- ⁴ (a) Kunai, A.; Kawakami, T.; Toyoda, K.; Ishikawa, M. *Organometallics*, **1992**, 11, 2708. (b) Kunai, A.; Ohshita, J. *J. Organomet. Chem.* **2003**, 686, 3.

- ⁵ Huang, Z.; Negishi, E. *Org. Lett.* **2006**, *8*, 3675.
- ⁶ Wang, Z.; Denmark, S. E. *Org. Synth.* **2005**, *81*, 42.
- ⁷ Smith, A. B., III; Tong, R.; Kim, W.-S.; Maio, W. M, *Angew. Chem. Int. Ed.* **2011**, *50*, 8904.
- ⁸ Lin, S., Lu, X. *J. Org. Chem.* **2007**, *72*, 9757.
- ⁹ Bastug, G., Dierick, S., Lebreux, F., Marko, I. E. *Org. Lett.* **2012**, *14*, 1306.
- ¹⁰ Harrowven, D.C.; Sutton, B.J.; Coulton, S. *Org. Biomol. Chem.* **2003**, *1*, 4047.
- ¹¹ Manolikakes, G.; Knochel, P. *Angew. Chem. Int. Ed.* **2009**, *48*, 205.
- ¹² Kobayashi, O.; Uraguchi, D.; Yamakawa, T. *Org. Lett.* **2009**, *11*, 2679.
- ¹³ Seki, M.; Mori, K. *Eur. J. Org. Chem.* **1999**, 2965.
- ¹⁴ Denmark, S. E.; Sweis, R. F. *J. Am. Chem. Soc.* **2001**, *123*, 6439.
- ¹⁵ Smith, A.B., III et al., *J. Am. Chem. Soc.* 2012, *134*, 4533-4536.
- ¹⁶ Son, E.-C., e tal., *Bull. Chem. Soc. Jpn.* 2006, *79*, 492.
- ¹⁷ Smith, A.B., III et al. *Angew. Chem. Int. Ed.* 2011, *50*, 8904-8907.

What is Claimed:

1. A compound of formula I:



I

wherein

Y is CH or N;

R₁ and R₂ are independently C₁₋₁₀ straight or branched-chain alkyl optionally substituted with one or more halogen, nitro, C₁₋₆alkoxy, or aryl;

R₃ is H;

aryl optionally substituted with one or more nitro, diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl;

heteroaryl optionally substituted with one or more nitro, diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl;


C₁₋₁₀ straight or branched-chain alkyl optionally substituted with one or more halogen, nitro, C₁₋₆alkoxy, or aryl;

a polymer; or

a resin support;

R_{3a} is H or C₁₋₆alkyl optionally substituted with one or more halogen; and


at least one R₄, wherein each R₄ is independently hydrogen, halogen, nitro, C₁₋₆alkoxy, C₁₋₆alkyl, aryl, or a resin support.






2. The compound of claim 1, wherein the compound is not

3. The compound of claim 1, wherein Y is CH.
4. The compound of claim 1, wherein Y is N.
5. The compound of any one of claims 1 to 4, wherein R₁ and R₂ are independently methyl, ethyl, propyl, or isopropyl.
6. The compound of claim 5, wherein R₁ and R₂ are independently methyl, ethyl, or isopropyl.
7. The compound of claim 5, wherein R₁ and R₂ are each methyl.
8. The compound of claim 5, wherein R₁ and R₂ are each isopropyl.
9. The compound of claim 5, wherein R₁ and R₂ are each ethyl.
10. The compound of any one of claims 1 to 9, wherein R₃ is C₁₋₆ straight or branched-chain alkyl optionally substituted with one or more C₁₋₆alkoxy.
11. The compound of claim 10, wherein R₃ is C₁₋₄ straight or branched-chain alkyl optionally substituted with one or more C₁₋₆alkoxy.
12. The compound of any one of claims 1 to 9, wherein R₃ is C₁₋₄ straight or branched-chain alkyl.
13. The compound of claim 12, wherein R₃ is n-butyl, isobutyl, or tert-butyl.
14. The compound of claim 12, wherein R₃ is n-butyl.
15. The compound of any one of claims 1 to 9, wherein R₃ is C₁₋₄ straight or branched-chain alkyl substituted with one or more halogen.
16. The compound of claim 15, wherein R₃ is -CF₃.


17. The compound of any one of claims 1 to 9, wherein R_3 is phenyl optionally substituted with one or more nitro, diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl.
18. The compound of claim 17, wherein R_3 is phenyl optionally substituted with one or more diC₁₋₆alkylamino, C₁₋₆alkoxy, or C₁₋₆alkyl.
19. The compound of claim 17, wherein R_3 is phenyl.
20. The compound of any one of claims 1 to 9, wherein R_3 is pyridyl.
21. The compound of any one of claims 1 to 9, wherein R_3 is a polymer.
22. The compound of any one of claims 1 to 21, wherein R_{3a} is H.
23. The compound of any one of claims 1 to 21, wherein R_{3a} is $-\text{CF}_3$.
24. The compound of any one of claims 1 to 23, wherein R_4 is hydrogen, C₁₋₆alkoxy, C₁₋₆alkyl, or aryl.
25. The compound of claim 24, wherein R_4 is hydrogen, C₁₋₆alkoxy, or C₁₋₆alkyl.
26. The compound of claim 24, wherein R_4 is hydrogen.
27. The compound of any one of the preceding claims that is










wherein n is 20 to 200,







28. A method of cross-coupling a compound of formula NuLi with a compound of formula E-X to form a compound of formula Nu-E comprising
- contacting the compound of formula NuLi with the compound of formula E-X in the presence of a compound of claim 1, catalyst or catalyst system, and an ethereal solvent;
- for a time and under conditions sufficient to produce the compound of formula Nu-E;
- wherein Nu is an aryl compound or an alkenyl compound; E is an aryl compound or an alkenyl compound; and X is iodo, chloro, or bromo.




29. The method of claim 28, wherein the compound of claim 1 is not .
30. The method of claim 28, wherein the solvent is tetrahydrofuran.
31. The method of claim 28, wherein the catalyst system comprises palladium, copper, or a mixture thereof
32. The method of claim 28, wherein the catalyst system comprises PdCl₂, dpca, and CuI.
33. The method of claim 28, wherein the catalyst system comprises dpca and CuI.
34. The method of claim 28, wherein the catalyst system comprises CuI and Johnphos
35. The method of claim 28, wherein the catalyst system comprises Pd(OAc)₂ and XPhos.
36. The method of any one of claims 28 to 35, wherein the compound of claim 1 is







37. A method of cross-coupling a compound of formula NuLi with a compound of formula E-X to form a compound of formula Nu-E comprising contacting the compound of formula NuLi with the compound of formula E-X in the presence of a compound of claim 1, catalyst or catalyst system, and an ethereal solvent;
- for a time and under conditions sufficient to produce the compound of formula Nu-E;
- wherein Nu is an aryl compound or an alkenyl compound; E is a disubstituted amine; and X is -O-benzoyl.



38. The method of claim 37, wherein the compound of claim 1 is not

39. The method of claim 37, wherein the solvent is tetrahydrofuran.

40. The method of claim 37, wherein the catalyst system comprises palladium, copper, or a mixture thereof.

41. The method of claim 37, wherein the catalyst system comprises dpca and CuI.
42. The method of claim 37, wherein the catalyst system comprises PdCl₂, dpca, and CuI.
43. The method of claim 37, wherein the catalyst system comprises Pd(OAc)₂ and XPhos.
44. The method of claim 37, wherein the catalyst system comprises CuI and Johnphos.
45. The method of any one of claims 37 to 44, wherein the compound of claim 1 is

