



US006476074B1

(12) **United States Patent**
Stjernschantz et al.

(10) **Patent No.:** **US 6,476,074 B1**
(45) **Date of Patent:** **Nov. 5, 2002**

(54) **METHOD AND COMPOSITION FOR TREATMENT OF ERECTILE DYSFUNCTION**

(58) **Field of Search** 604/517; 424/449; 560/121

(75) **Inventors:** **Johan Stjernschantz**, Uppsala (SE);
Bahram Resul, Uppsala (SE)

(56) **References Cited**

(73) **Assignee:** **Synphora AB**, Uppsala (SE)

U.S. PATENT DOCUMENTS

(*) **Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

4,190,670 A * 2/1980 Marx et al. 424/305
6,031,001 A * 2/2000 Stjernschantz et al. 514/573

FOREIGN PATENT DOCUMENTS

(21) **Appl. No.:** **09/462,450**

EP WO 39/00894 * 1/1993 A61K/13/195
WO WO 9300894 1/1993
WO WO 9632141 10/1996

(22) **PCT Filed:** **Jul. 10, 1998**

* cited by examiner

(86) **PCT No.:** **PCT/SE98/01367**

§ 371 (c)(1),
(2), (4) **Date:** **Apr. 10, 2000**

Primary Examiner—Thurman K. Page
Assistant Examiner—Isis A. D. Ghali
(74) *Attorney, Agent, or Firm*—Dinsmore & Shohl LLP

(87) **PCT Pub. No.:** **WO99/02164**

PCT Pub. Date: **Jan. 21, 1999**

(57) **ABSTRACT**

(30) **Foreign Application Priority Data**

Jul. 10, 1997 (SE) 9702681


Compositions and methods for treatment of impotence or erectile dysfunction employ prostaglandins that are selective EP₂ or EP₄ prostanoid receptor agonists. The compositions can be formulated for intracavernous injection, or for transurethral or transdermal application.

(51) **Int. Cl.**⁷ **A01N 37/08**; A61F 6/06;
A61F 13/00; A61P 15/08; A61P 15/10

(52) **U.S. Cl.** **514/573**; 424/430; 424/449

14 Claims, 2 Drawing Sheets

HUMAN CORPUS C.
n=8



HUMAN CORPUS C.
n=8




FIG. 1

RABBIT PENILE BLOOD VESSELS




FIG. 2

METHOD AND COMPOSITION FOR TREATMENT OF ERECTILE DYSFUNCTION

FIELD OF THE INVENTION

The present invention relates to the treatment of impotence or erectile dysfunction, and more particularly to a novel prostaglandin based composition therefore and the use thereof for treating impotence or erectile dysfunction.

BACKGROUND OF THE INVENTION

Erectile dysfunction is a disorder characterized by the inability of the male to develop and maintain erection for satisfactory sexual intercourse. Erectile dysfunction is a frequent disorder particularly amongst elderly men which may lead to reduced quality of life and psychological problems. It is estimated that there may be as many as 10–20 million people in the United States suffering from erectile dysfunction, and an estimated 30 million males with at least partial erectile dysfunction (NIH Consensus Conference 1993). The prevalence of erectile dysfunction has been reported to be about 5% at age 40, and up to 25% at age 65 or older. Thus erectile dysfunction is a major clinical challenge of increasing importance with the increased standard of living and demand of a better quality of life.

The etiology of erectile dysfunction may be psychogenic or organic. The latter seems to account for the majority of cases. Such organic causes include vascular, endocrinological, and neurological diseases as well as trauma. Patients suffering from diabetes are typically at risk. While in many cases erectile dysfunction caused by psychogenic factors may be reversible, impotence caused by organic factors needs adequate therapy. Such therapy comprises surgical intervention, devices and medical treatment. With more effective and better tolerated drugs there is a clear tendency towards medical therapy in the treatment of erectile dysfunction. The main modalities of medical therapy consist of systemic medication, usually peroral, and local medication in the genitourinary tract. Typical drugs given orally include e.g. yohimbine, an alpha-2 adrenergic antagonist, and testosterone, the male sex hormone. Furthermore bromocriptine has also been used, as well as antiserotonergic agents such as trazodone, ketanserin and mianserin. Recently, a selective type 5-phosphodiesterase inhibitor sildenafil (Viagra™) has been approved for clinical use. In addition there are many other drugs that have been tested and used for the treatment of erectile dysfunction. Drugs given locally include e.g. papaverine, a smooth muscle relaxing agent, and phentolamine, an alpha-adrenergic antagonist as well as prostaglandins, particularly prostaglandin E₁ (PGE₁; alprostadil). These drugs relax smooth muscle, thus promoting the development of erection, and are given by local injection into the cavernous tissue of the penis. Formulations for intraurethral (transurethral) administration of prostaglandins have also been developed (Wolfson et al., 1993; Bradley et al., 1996) and are described in several patents and patent applications, for instance in WO 93/00894, WO 91/16021 and EP-A-357581.

It is a clear advantage to administer the drug locally into the diseased or dysfunctional organ since a better effect is achieved, and usually less systemic side-effects are induced. However, the local side-effects, mainly pain and priapism, may constitute a problem. The latter meaning prolonged erection which potentially can lead to necrosis and irreversible damage to the reproductive organ has been minimized by the use of prostaglandins, above all PGE₁. However,

PGE₁, given by intracavernous injection or transurethrally causes local pain in as many as 10–30% of the patients. In a recently published study transurethral administration of PGE₁ resulted in pain sensation in about 35% of the patients, and 11% of the patients reported pain after every treatment with alprostadil (Padma-Nathan et al., 1997). It is thus obvious that local pain constitutes one of the most significant side-effects of PGE₁, and prostaglandin analogues without pain inducing effect would be desirable for clinical use.

SUMMARY OF THE INVENTION

We have now unexpectedly found that certain prostaglandin analogues of the E-type, more specifically agonists of the EP₂ prostanoid receptor, exert good relaxant effect in the penile cavernous tissue, and in blood vessels while exhibiting markedly reduced pain inducing effect. In particular we have found that PGE₂ analogues substituted in carbon 18, 19 or 20 with hydroxyl (OH) exert beneficial effects, and especially one analogue, namely 19R-OH-PGE₂, exhibited a surprisingly advantageous effect with respect to relaxation of penile cavernous tissue and blood vessels without inducing pain as studied in an animal model.

Thus, in a first aspect, the present invention provides a composition for the treatment of impotence or erectile dysfunction, comprising a prostaglandin which is a selective EP₂ or EP₄ receptor agonist, or an active derivative including the free acid, a salt or an ester thereof, optionally together with a physiologically acceptable carrier or vehicle.

In a second aspect, the invention provides a method of treating impotence or erectile dysfunction, which method comprises administering a therapeutically active and physiologically acceptable amount of the composition to an individual in need thereof.

In a third aspect, the invention provides the use of a selective EP₂ or EP₄ receptor agonist, or an active derivative including the free acid, a salt or an ester thereof, for the preparation of a medicament for the treatment of impotence or erectile dysfunction.

In still another aspect, the invention provides a method for eliminating or at least considerably reducing side effects, such as pain and irritation, observed in connection with the use of prostaglandin derivatives for treatment of impotence.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 is a diagram showing the concentration-response relation for PGE₁ and 19R-hydroxy-PGE₂ in isolated preparation of human corpus cavernosum precontracted with 10⁻⁶ M norepinephrine.

FIG. 2 is a diagram showing the vasodilatory effect of PGE₁ and 19R-hydroxy-PGE₂ on rabbit penile blood vessels precontracted with 10⁻⁶ M norepinephrine.

DETAILED DESCRIPTION OF THE INVENTION


Penile erection is based on three main physiological events: an increase in the arterial blood flow, a relaxation of the expansive tissue of the corpora cavernosa and the corpus spongiosum, and an obstruction of the venous return by mechanical compression of the veins caused by the expansive tissue. PGE₁ and the EP₂ receptor agonists cause vasodilatation and relax the expansive tissue to about the same extent, and thus promote the development of erection. However, PGE₁ being a natural prostaglandin with low selectivity for specific prostanoid receptors has a marked

3

irritant effect and causes pain as evident in clinical trials performed with alprostadil (e.g. Padha-Natham, 1997) and as will also be shown in the experimental part further below.


According to the invention, however, prostaglandin analogues with selectivity primarily for the EP₂ receptor which induce the same beneficial effect but with significantly less irritant effect are much more preferable as it can be predicted that such analogues will cause no or only minimal pain in clinical practice. To facilitate the understanding of the invention, a general description of prostaglandins is first given.

The prostaglandins are fatty acids usually derived from the precursors eicosatrienoic, eicosatetraenoic or eicosapentanoic acid through metabolic steps involving oxygenation. The prostaglandins typically carry a cyclopentane ring to which two carbon chains link, the upper chain usually being called the alpha chain and the lower chain the omega chain. The alpha chain is a 7 carbon chain with a terminal carboxylic acid moiety whereas the omega chain is an 8 carbon methyl-terminated aliphatic chain. Depending on the number of double bonds in these chains substituents of 1 to 3 are used. In prostaglandins with subscript 1, e.g. PGE₁, the double bond is situated between carbons 13 and 14 in the omega chain, and it exhibits trans configuration in naturally occurring prostaglandins. In prostaglandins with subscript 2, e.g. PGE₂, an additional double bond in the cis configuration exists between carbons 5 and 6 in the alpha chain, and finally in prostaglandins with subscript 3 a third double bond is situated between carbons 17 and 18 in the omega chain. This double bond also exhibits cis configuration in naturally occurring prostaglandins. All naturally occurring prostaglandins carry a hydroxyl group in carbon 15, which is essential for biological activity. The substituents of the cyclopentane ring determine whether the prostaglandin is of the A, B, C, D, E, F, G, H, I or J type as depicted in Scheme 1 below. The prostaglandins that have been used to exemplify the present invention are of the E-type, and the chemical structures of these prostanoids are depicted in Scheme 2. Except for 18RS-OH-PGE₂-methyl ester the prostaglandin analogues were also used as acids.




4

-continued



Scheme 2



The prostaglandins exert their pharmacologic effect through specific G protein coupled membrane receptors. There are at least 8 different receptors for the endogenous prostaglandins, namely the FP (PGF_{2α}), EP₁, EP₂, EP₃, EP₄ (PGE₂), DP (PGD₂), IP (PGI₂), and TP (TxA₂) receptors, the most common naturally occurring ligand for respective receptor being shown within parentheses. At least for the EP₃ receptor splice variants have been shown to exist. The EP₂ and EP₄ receptors are closely related, and probably mediate similar effects. We have therefore studied the effect only on the EP₂ receptor, and regard it to represent the EP₄ receptors as well in the case the compounds studied would have an effect on that receptor, too. The molecular biology and pharmacology of the prostanoid receptors have recently been reviewed by Coleman et al., 1994.

From a therapeutic point of view a problem with the endogenous prostaglandins is that they exert effects on many different prostanoid receptors. Each endogenous prostaglandin has a preference for one specific receptor type, but is not very selective and usually distinguishes poorly between the

receptor subtypes, i.e. the EP receptors. Thus PGE₁ and PGE₂ are good ligands for all subtypes of the EP receptor. Consequently selective effects on one of the subtypes of the EP receptor is impossible to achieve with the endogenous prostaglandins. However, certain synthetic prostaglandin analogues, e.g. butaprost, 11-deoxy-PGE₁ and AH13205 as well as a naturally occurring metabolite of PGE₂, namely 19R-OH-PGE₂, are selective EP₂ prostanoid receptor agonists.

18-OH-PGE₂, 19R-OH-PGE₂ and 20-OH-PGE₂ are effective EP₂ receptor agonists with selectivity for the EP₂ receptor over the EP₃ receptor. The endogenous PGE₁ is unselective and does not distinguish between the EP receptor subtypes sufficiently, and furthermore significantly spills over on the DP/IP receptors which e.g. the 18-, 19- and 20-OH substituted PGE₂ analogues do not. However, PGE₁ has been included as a reference substance as it is the only prostaglandin currently in clinical use for the treatment of erectile dysfunction.

Accordingly, high selectivity or specificity to the EP₂ receptor compared to other prostaglandin receptors, particularly the EP₃ receptor, characterizes the compounds to be used in the method or compositions according to the present invention. It need not be said that the more specific the compound is for the EP₂ receptor the better results are obtained, but a certain advantage is, of course, achieved also in cases of some interaction with other receptors. High selectivity in this connection means that the effect on the EP₂ receptor is at least more than S times, especially more than 10 times, and in particular more than 100 or 1000 times the effect on other prostaglandin receptors.


As stated above, according to the present invention, in particular the selective EP₂ receptor agonist 19R-OH-PGE₂ and its carboxylic acid esters appear to be unique and ideally suited for the treatment of erectile dysfunction. In this context it should be mentioned that 19R-OH-PGE₂ is a metabolite of PGE₂ in the genital tract, and can be found in large quantities normally in human semen (Taylor and Kelly, 1974). However, the physiological role of this unique metabolite is unknown. Thus, 19R-OH-PGE₂ and its carboxylic acid esters constitute a very attractive alternative to PGE₁ as medication for impotence since this analogue causes no pain, is as effective as PGE₁, and furthermore occurs normally in the body.

With respect to 19R-OH-PGE₂ which appears ideally suited for treatment of erectile dysfunction, it should be observed that various modifications or substitutions of the molecule are possible as long as the new derivatives exhibit selective agonism on the EP₂ receptor.

In Formula I, the wavy bonds represent the α (S) or β (R) configuration, and the dashed bonds represent a single, a triple or a double bond in the cis or trans configuration. R in the formula is hydrogen, a salt moiety, e.g. an alkali or ammonium salt, a straight or branched saturated or unsaturated alkyl group, preferably with 1–10 carbons, an alicyclic ring, preferably with 3–8 carbons, arylalkyl, preferably aryl-C₂₋₅ alkyl, or an aryl ring. X is a straight chain, saturated or unsaturated, preferably consisting of 2–5 carbons optionally interrupted by a heteroatom (O, S, N), and the chain may contain an alicyclic ring, preferably C₃–C₇ cycloalkyl, or an aryl or heteroaryl ring. R1 and R2, which are identical or different, are hydrogen, hydroxy, halogen, oxygen (=O or alkoxy) or an alkyl group with 1–3 carbons or an alkoxy group with 1–3 carbons or an ester OCOR₃, where R₃ is a straight or branched, saturated or unsaturated alkyl group preferably containing 1–10, especially 1–6

carbons, or a cycloalkyl, preferably containing 3–7 carbons, or an aryl or arylalkyl group, preferably aryl-C₂₋₅ alkyl (e.g. benzyl). Optionally C10 can be substituted e.g. to contain a mono- or dialkyl group.

Z is an alkyl chain of 1–8, e.g. 3–8 and 24 carbons, saturated or unsaturated, optionally interrupted by one or more heteroatoms (O, S, N), and containing one or more, preferably one substituent Y, attached to a carbon atom within the chain or a carbon atom in terminal position. The chain Z may be branched, containing alkyl (preferably methyl) substituents and/or contain an alicyclic ring, e.g. a cycloalkyl, or an aryl (esp. a benzene) or heteroaryl ring, in inter- or terminal position. Y, is hydroxy, sulfhydryl, amino, methylamino, dimethylamino, C₁₋₃ alkoxy or halogen (Cl, Br, F) or oxygen (keto). In a preferred embodiment of the invention Y is attached to carbon 18, 19 or 20. In the most preferred embodiment Y is OH and attached to carbon 19. At present it is believed that the inventive concept is based on the electronegative effect introduced by Y.



The prostaglandins may be epimeric mixtures as well as in the form of the individual epimers.

DESCRIPTION OF SUITABLE EMBODIMENTS

The EP₂ prostanoid receptor agonists according to the present invention can be used as the normal carboxylic acids, salts (e.g. cationic) or as ester prodrugs, preferentially carboxylic acid alkyl esters. The active compounds can either be administered by intracavernous injection, transurethral, or transdermal application (including on the glans of the penis) in a pharmaceutically acceptable delivery medium. For intracavernous injection sterile isotonic water based solutions are preferred. These should be buffered and have a pH of around 7.0–7.5 or at least in the interval of 6.0–8.0. For solubilisation of the prostaglandin different micellar systems can be used such as polysorbate. Cyclodextrins may also be employed for solubilisation. If the prostaglandin analogues to be used are unstable in water solution the compounds may be lyophilized and dissolved immediately before use or stabilized with different stabilizing agents such as cyclodextrins. Different slow-release formulations adapted to the requirements of injectable solutions may also be employed. If the new medicament is to be administered transurethral the active principle may be formulated in creams, gels or ointments, suppositories, or other solid state forms. Furthermore a device (applicator) for introducing the medicament into the urethra is also needed. It is understood that such a device can be designed in a variety of ways and consist of different materials. If the new medicament is to be administered transdermally various forms of creams, ointments, gels, and slow release systems such as patches may be employed. Also the inner surface of condoms or bandages may be lined by a suitable formulation containing the new medicament. Gels, creams, ointments, and different solid state formulations may or may not contain preservative such as benzalkonium chloride, chlorhexidine, thiomersal, parabenoic acid, and other compounds with satisfactory antimicrobial effect. For intracavernous injec-

tion the dose interval is 0.001–1 mg, typically 0.01–0.1 mg per injection. For transurethral and transdermal administration the dose interval is 0.01–10 mg, typically 0.1–1 mg for transurethral administration, and 0.1–10 mg for transdermal administration.

Accordingly the new medicament should be administered locally to the penis either by injection, or by applying it into the urethra with an applicator or syringe, or it should be applied topically on the skin or the mucous membrane of the penis. Such treatment should be initiated typically 5–60 min, depending on the mode of administration, before intercourse.

EXEMPLIFICATION OF THE INVENTION

PGE₁ was obtained from Chinoïn, Pharmaceutical and Chemical Works Co. Ltd., Budapest, Hungary.

DBU (644 mg, 4.2 mmol) in acetonitrile (1 ml) was added dropwise to a stirred solution of PGE₁ (300 mg, 0.84 mmol) in acetonitrile (3ml) at 0° C. The mixture was allowed to warm to room temperature whereupon isopropyl iodide (1142 mg, 6.72 mmol) in acetonitrile (2 ml) was added dropwise. After being stirred for 10 h (TLC monitoring), the reaction mixture was quenched with water (8 ml), the mixture was extracted with ethyl acetate (2×50 ml), and the extract was washed with brine (10 ml), citric acid 3% (10 ml), and finally sodium hydrogen carbonate 5% (2×10 ml). After drying with anhydrous sodium sulphate, the solvent was removed in vacuo and the residual oil was chromatographed on silica gel using ethyl acetate as eluent. This afforded 230 mg of the product (69%) of the title compound as a colorless oil: R_f=0.18 (ethyl acetate); ¹H NMR (CDCl₃) δ 0.89 (m, 3H), 1.2 (d, 6H), 1.21–1.4 (m, 10H), 1.42–1.62 (dm, 6H), 2.2–2.4 (dm, 4H) 2.7–2.75 (dd, 1H, 4.0–4.17 (dm, 2H), 5.0 (m, 1H, 5.5–5.7 (dm, 2H).

Synthesis of 18RS-Hydroxy-PGE₂-methyl Ester

1. Preparation of Dimethyl-(2-oxo-5-heptyne)-phosphonate. (2)

To a stirred suspension of dry sodium hydride (3.13 g, 124,07 mmol) in dry THF (100 ml) at room temperature was added dropwise a solution of dimethyl-(2-oxopropyl)-phosphonate (20,61 g, 124,07 mmol) in dry THF (50 ml) under N₂. The reaction mixture was stirred for 1 h, then cooled in an ice-bath and treated with a solution of n-butyllithium (7.95 g, 124, 07 mmol) in hexane, causing a dark brown solution to be formed. Stirring was continued for 1 h at 0° C., followed by dropwise addition of 1-bromo-2-butyne.1 (15 g, 112,79 mmol) in THF (30 ml). The reaction mixture was gradually warmed to room temperature and after 1 h (TLC monitoring) the reaction mixture was quenched with ice-water, HCl 1M to pH about 4 and extracted, twice, with ethyl acetate. The organic layer was washed with brine and chromatographed on silica gel using EtOAc as eluent. R_f=0.38 (silica gel, ethylacetate), yield 18 g (73%).

2. (1S,5R,6R,7R)-6-Formyl-7-[(4-phenylbenzoyl)oxy]-2-oxabicyclo{3.3.0}octan-3-one.(3)

A mixture of Coreys lactone (23,06 g, 65,44 mmol), DCC (40,50 g, 196,31 mmol), DMSO (27,8 ml, 392,6 mmol) and phosphoric acid 85% (2,2 ml) in DME (130 ml) was stirred at room temperature for 2h (TLC monitoring). The precipitate was removed on silica-gel pad washed with DME (2×50 ml). The filtrate was concentrated in vacuo and used for the next step without isolation.

4. (1S,5R,6R,7R)-6-(3-Oxo-6yn-1E-1-octenyl)-7-[(4-phenylbenzoyl)oxy]-2-oxabicyclo{3.3.0}octan-3-one.(4)

To a stirred suspension of sodium hydride (1,98 g, 78,53 mmol) in DME (140 ml) under N₂ was added dropwise the

above phosphonate 2 (18,56 g, 85,07 mmol) and stirred mechanically at room temperature for 1,5 h. The mixture was then cooled to –5° C., and a solution of the above crude Coreys aldehyde 3 was added dropwise. After 30 min at 0° C. and 2 h at room temperature (TLC monitoring), the reaction mixture was neutralised with citric acid 5% and extracted with ethyl acetate (2×100 ml). The organic layer was dried and evaporated. The residue (oil) was chromatographed on silica-gel using ethyl acetate and 10% methanol in ethyl acetate successively, giving a light yellowish oil. R_f=0.58(silica-gel, ethyl acetate), yield 52%.

5. (1S,5R,6R,7R)-6-(3,6-Dioxo-1E-1-octenyl)-7-[(4-phenylbenzoyl)oxy]-2-oxabicyclo{3.3.0}octan-3-one.(5)

To the acetylenic solution 4 in acetonitrile:water 2:1 (100 ml) was added mercuric oxide (13,8 g, 63,73 mmol) and sulfuric acid 1M (25,49 ml, 25,49 mmol). The reaction mixture was stirred magnetically. After about 1 h at room temperature (TLC monitoring) the reaction mixture was worked-up by addition of ethylacetate and HCl 1M. The organic layer was dried and evaporated. The crude oil was used for the next step without purification. R_f=0.44 (silicagel, EtOAc), yield=41%.

6. (1S,5R,6R,7R)-6-(3RS,6RS-Dihydroxy-1E-1-octenyl)-7-[(4-phenylbenzoyl)oxy]-2-oxabicyclo{3.3.0}octan-3-one. (6)

To a stirred solution of the above diketone 5 (12,0 g, 26.1 mmol) and cerium chloride hepta hydrate (5.83 g, 15,64 mmol) at –20° C. in methanol: methylene chlorider 1:1 was added sodium borohydride (1.48 g, 39,09 mmol) in small portions under N₂. After 30 min (TLC monitoring). The reaction mixture was quenched with HCl 1M to pH about 4–5, and diluted with water (50 ml) and ethyl acetate (100 ml). The organic layer was separated and the water layer was washed twice with EtOAc, dried and evaporated. The residue was purified on silica gel using EtOAc as eluent. The title compound 6 was obtained as colorless oil: yield 8.8 g (71%). R_f=0,15, 0,13 corresponding to the two isomers 15α and 15β (silicagel, EtOAc).

¹H NMR (CDCl₃) δ 0,9 (m, 3H), 1,4–1,8 (dm, 6H), 2,3 (d, 1H, 2,5–2,9 (dm, 5H), 3,5 (m, 1H), 4,2 (m, 1H, 5.1 (m, 1H, 5,3 (m, 1H), 5,7 (m, 2H), 7,4 (m, 1H, 7,5 (m, 2H), 7,6–7,7 (dd, 4H), 8,1 (d, 2H); ¹³C NMR (CDCl₃) d 176,40 (C6H4C=O), 16591 (lactone C=O), 146,07, 139,83, 136, 21, 130,15, 128,91, 128,31, 127,15, 83,27, 79,13, 73,11, 71,79, 71,48, 54,08, 42,84, 42,75, 37,55, 34,85, 34,01, 33,43, 32,95, 32,09, 30,17, 9,96.

7. (1S,5R,6R,7R)-6-(3RS,6RS-Di t Butyldimethylsilyloxy-1E-1-octenyl)-7-[(4-phenylbenzoyl)oxy]-2-oxabicyclo{3.3.0}octan-3-one.(7)

To a stirred solution of the above dihydroxy compound 6 (8,6 g, 18,51 mmol) in dichloromethane was added triethyl amine (12,83 ml, 92,56 mmol), t-butyldimethylsilyl chloride (13,95 g, 92,56 mmol) and 4-dimethylamino pyridine (1,13 g, 9,26 mmol). The mixture was stirred magnetically for 15 h at room temperature. The reaction mixture was diluted with ether, filtered and the precipitate was washed with ether. The organic layer was washed with brine, dried and concentrated in vacuo. The residue was chromatographed on silica gel using 5% ether in dichloromethane. R_f=0.58 (silica gel, 5% ether in CH₂Cl₂).Yield 12,2 g (92%)

8. (1S,5R,6R,7R)-6-(3RS,6RS-Di t Butyldimethylsilyloxy-1E-1-octenyl)-7-[(4-hydroxy)-2-oxabicyclo{3.3.0}octan-3-one.(8)

To a stirred solution of the above disilyl ether 7 (12 g, 17,31 mmol) in methanol was added potassium carbonate (1,2 g, 8,66 mmol). The reaction mixture was stirred at room temperature for 4 h (TLC monitoring). The mixture was

9

neutralised with citric acid 5%, extracted twice with ethyl acetate (100 ml), dried and concentrated in vacuo. The oil was chromatographed on silica gel using gradient elution with 5% ether in CH_2Cl_2 and EtOAc:acetone 1:1 successively. $R_f=0.43$ (silica gel, ethyl acetate), yield=8,86 g (74%)

9. (1S,5R,6R,7R)-6-(3RS,6RS-Di t Butyldimethylsilyloxy-1E-1-octenyl)-7-[(4-t-butylidimethylsilyloxy)-2-oxabicyclo{3.3.0}octan-3-one (9)

To a stirred solution of the above disilyloxy ether 8 (6,6 g, 12,86 mmol) in CH_2Cl_2 at room temperature was added triethyl amine (7,14 ml, 51,47 mmol) t-butylidimethylsilyl chloride (7,76 g, 51,47 mmol) and 4-dimethyl aminopyridine (0,47 g, 3,9 mmol). The reaction mixture was worked-up as in 7. The crude product was chromatographed on silica gel using 5% ether in dichloromethane to give a pure trisilyloxy ether product 9 as an oil. $R_f=0.62$ (silica gel, 5% ether in CH_2Cl_2), yield 7,8 g, (96%)

10. (1S,5R,6R,7R)-6-(3RS,6RS-Di t Butyldimethylsilyloxy-1E-1-octenyl)-7-[(4-t-butylidimethylsilyloxy)-2-oxabicyclo{3.3.0}octan-3-ol (10)

A solution of diisobutyl aluminium hydride (DIBAL-H) (2,9 g, 20,87 mmol) in dry THF was added dropwise to a stirred solution of the above trisilyl ether lactone 9 (7,7 g, 12,24 mmol) in THF (80 ml) at $-72/-80^\circ\text{C}$. After 1 h (TLC monitoring), the reaction mixture was quenched with ice (15 g) and ethyl acetate (150 ml), filtered and the filtrate was concentrated in vacuo. The residue was used directly without separation in the next step. $R_f=0.27$ (Silica gel, 5% ether in dichloromethane).

11. 11,15RS,18RS-Tri-t-butylidimethylsilyloxy-PGF_{2α} (11)

To a stirred suspension of 4-carboxybutyl-triphenylphosphonium bromide (21,70 g, 48,95 mmol) in dry THF under N_2 at -5°C was added potassium tert-butoxide (10,99 g, 97,91 mmol) and the mixture was stirred at room temperature for 30 min. To the resultant red-orange solution of ylide at $-15/-10^\circ\text{C}$ was added the lactol 10 (7,7 g, 12,24 mmol) in THF, and the mixture was stirred for 2-3 h (TLC monitoring) at room temperature. The reaction mixture was diluted with water and quenched with citric acid 15% to pH 6-7 and extracted with EtOAc, dried and concentrated in vacuo. The resultant slurry was used directly without isolation for the next step.

12. 11,15,18RS-Tri-t-butylidimethylsilyloxy-PGF_{2α} Methyl Ester (12)

Methyl iodide (8,6 g, 61,20 mmol), was added to a stirred solution of the crude product 10 (8,73 g, 12,24 mmol) and N,N-Diisopropyl ethyl amine (9,473 g, 73,44 mmol) in acetonitrile at room temperature. After 15h (TLC monitoring) the mixture was diluted with water (100 ml) and ethyl acetate (150 ml), washed with sodium hydrogen carbonate 5% (60 ml) and brine (70 ml). The organic layer was dried and evaporated in vacuo. The residue was chromatographed on silica gel using EtOAc as eluent. $R_f=0.18$ (silica gel, EtOAc:hexane 1:9).

13. 11,15,18RS-Tri-t-butylidimethylsilyloxy-PGE₂ Methyl Ester (13)

To a stirred solution of the compound 11 (3,3 g, 4,54 mmol) in dichloromethane was added pyridinium chlorochromate (PCC) (3,9 g, 18,15 mmol) treated with aluminium oxide (1 g PCC was stirred with 5 g aluminium oxide in acetone, the solvent was removed in vacuo giving a light yellow powder). The resulting suspension was stirred at room temperature for 4 h (TLC monitoring). The suspension was filtered on silica gel pad, washed with dichloromethane. The solvent was removed and the resulting oil was diluted with ether and washed with water (50 ml), sodium hydrogen carbonate 5% (50 ml). The solvent was removed

10

in vacuo. The residue was chromatographed on silica gel using 5% ether in CH_2Cl_2 . $R_f=0.32$ (silica gel, EtOAc:hexane 1:9), yield 3,1 g (86%)


14. 18RS-Hydroxy PGE₂ Methyl Ester (14)

The protecting groups 11,15,18-tri-t-butylidimethylsilyl chloride were removed by addition of HF 4% (108 ml) to a solution of 13 (3,0 g, 3,98 mmol) in acetonitrile (300 ml). The reaction mixture was stirred at room temperature for about 8 h (TLC monitoring). The reaction mixture was worked-up by addition of EtOAc (200 ml). The organic layer was separated and washed with sodium bicarbonate 5% and the pH was adjusted to about 6. The organic layer was washed with brine, dried and concentrated in vacuo. The residue was purified by column chromatography on silica gel using gradient elution with CH_2Cl_2 , EtOAc and 5-10% methanol in ether successively (The stationary phase, silica gel, in the column must be washed with the eluent containing triethylamine before purification, to avoid isomerization) 18RS-hydroxy-15S-PGE₂ Me ester $R_f=0.16$ (5% MeOH in ether); yield 310 mg.

18RS-hydroxy-15R-PGE₂ Me ester $R_f=0.20$ (5% MeOH in ether); yield 248 mg.






18RS-hydroxy-PGE₂ methyl ester 1H NMR (CDCl₃) δ 0,9 (t, 3H), 1,4-1,7 (dm, 8H), 2,1 (m, 2H), 2,2 (m, 2H), 2,3-2,4 (m, 5H), 2,7 (m, 1H), 3,6 (m, 1H, 18-CHOH), 3,7 (s, 3H), 4,05, (m, 1H, 15-CHOH), 4,2 (m, 1H, 11-CHOH), 5,3-5,5 (dm, 2H, db), 5,6-5,8 (dm, 2H, db); 13C (NMR) δ 214, 174,24, 136,7, 131,3, 130,8, 126,5, 72,36, 71,78, 71,70, 55,54, 54,54, 53,70, 51,60,46,06, 34,06, 33,42, 33,00, 32,10, 30,34, 30,29, 30,07.

Scheme 3












11

-continued



12

-continued



Synthesis of 19-R-Hydroxy-prostaglandin E₂-Methyl Ester

19R-hydroxy-prostaglandin E₂ was obtained from Cayman Chemicals, Ann Arbor, Mich., USA. Methyl iodide (9.2 mg, 0.065 mmol) in acetonitrile (1.0 ml) was added dropwise to a stirred solution of 19R-hydroxy-prostaglandin E₂ (4 mg, 0.011 mmol), and N,N-diisopropyl ethyl amine (7 mg, 0.054 mmol) in acetonitrile. More methyl iodide (4.5 mg, 0.032 mmol) in acetonitrile was added after 6 h and the stirring was continued for 12 h (TLC monitoring). The reaction mixture was quenched with water (5.0 ml) and extracted with ethyl acetate (2×10 ml), and the organic phase was washed with sodium hydrogen carbonate 5% (5 ml) and hydrochloric acid 0.5M (5 ml). After drying with anhydrous sodium sulphate, the solvent was removed in vacuo and the residue was chromatographed on silica-gel using ethyl

acetate:acetone 1:1 and acetone as eluent. This afforded 3.2 mg (72.7%) of the product as a colorless oil: $R_f=0.17$ (ethyl acetate:acetone:acetic acid 1:1:0.01) $^1\text{H NMR}$ (CDCl_3) δ 1.25 (d, 3H), 1.5–1.7 (m, 8H), 2.1–2.6 (m, 9H), 3.7, (s, 3H), 3.8 (m, 1H), 4.1 (m, 1H), 4.2 (m, 1H), 5.3–5 (dm, 2H), 5.6–5.8 (dm, 2H).

Synthesis of 20-OH-Prostaglandin E_2 -Methyl Ester

The commercially available 20-hydroxy PGE_2 (Cayman Chemicals, Ann Arbor, Mich.) (2.0 mg, 0.0054 mmol) was esterified in acetonitrile (2.0 ml) with methyl iodide (4.6 mg, 0.0327 mmol) in the presence of N,N -diisopropyl ethylamine (3.5 mg, 0.027 mmol). The reaction mixture was stirred at room temperature for 10 h (TLC monitoring, silica gel, ethyl acetate). The reaction mixture was quenched with water (3.0 ml) and extracted with ethyl acetate (2×10 ml). The organic layer was dried and concentrated in vacuo and the residual oil was chromatographed on silica gel using ethyl acetate:acetone 1:0.5 as eluent; $R_f=0.38$ (silica gel, ethyl acetate:acetone 1:1).

PGE_1 -isopropyl ester, 18RS-hydroxy- PGE_2 -methyl ester, 19R-hydroxy- PGE_2 -methyl ester and 20-hydroxy- PGE_2 -methyl ester were dissolved in 0.5% polysorbate-80 as a stock solution, and were diluted in 0.5% polysorbate-80 to the appropriate concentrations.

Pharmacological Experiments

Human penile cavernous tissue was obtained fresh from surgery, and representative tissue samples were mounted in smooth muscle tissue baths containing a modified Krebs's solution consisting of NaCl 119 mM, KCl 4.6 mM, MgCl_2 1.2 mM, NaH_2PO_4 1.2 mM, NaHCO_3 15 mM, CaCl_2 1.5 mM and glucose 11 mM. The solution also contained indomethacin at a final concentration of about 3 μM . The solution was continuously bubbled by 95% O_2 and 5% CO_2 , and the temperature was kept at 37° C.

The tissue preparations were stretched by a force corresponding to 500 mg, and were given a contractile tone by the addition of norepinephrine at a concentration of 10^{-6} M. Concentration-response curves were then constructed by adding cumulatively increasing concentrations of the prostaglandin analogues to the bath in a routine way. The relaxant effect was normalized by comparing to that of carbachol in the same preparation. PGE_1 and 19R-OH- PGE_2 , respectively, were used as acids.

To study the effect of the PGE_1 and 19R-OH- PGE_2 on the penile vasculature, penile blood vessels of the rabbit were isolated and mounted as ring segments in a small vessel myograph (J.P. Trading, Denmark) containing a solution consisting of NaCl 119 mM, KCl 4.7 mM, CaCl_2 1.5 mM, MgSO_4 1.17 mM, KH_2PO_4 1.18 mM, NaHCO_3 25 mM, EDTA 0.027 mM and glucose 11 mM. The solution also contained indomethacin at a final concentration of about 3 μM , and was continuously bubbled by 95% O_2 and 5% CO_2 . The temperature was kept constant at 37° C. The vessels were stretched and then precontracted using 10^{-6} M norepinephrine, and the vasorelaxant effect of the prostaglandin analogues was normalized by comparing it to that of papaverine in the same preparation. Cumulative concentration-response curves were constructed for the analogues.

The vasodilatory effect of the prostaglandin analogues was also studied in the rat by registering the blood pressure reducing effect. This is a relevant *in vitro* model for showing general vasodilatory effect of importance for inducing erection since vasodilation in penis is needed for achieving

erection. Rats were anaesthetized with a mixture of ketamine and xylazine and the prostaglandin analogues were infused intravenously. The blood pressure was continuously registered in a femoral artery. Each analogue was infused in 3 escalating doses in the same animal. None of the compounds was found to have any significant effect of the heart rate and the immediate reduction in arterial blood pressure therefore reflects an acute vasodilatory response to the prostaglandin analogues tested. In all experiments the blood pressure reduction was transient and the blood pressure increased immediately after termination of the infusion. The analogues were used both as esters and acids for the experiments.

The irritant effect of the prostaglandin analogues was tested using a behavioural model in cats. In this model the irritant effect on the eye is studied by registering the degree of eyelid closure and the behaviour of the animals. Compounds that cause discomfort and pain in the eye cause the animals to close their eyes. The analogues were applied as carboxylic acid esters, to enhance bioavailability, topically on the eye as a single dose at different dose levels. The animals were then followed for several hours at regular intervals. Each compound and dose was tested on a group of 3–6 cats. At least 3 days elapsed between two consecutive tests on the same animals. An arbitrary scale from 0 (absence of irritation) to 3 (marked irritation), with half steps was used.

The relaxant effect on the human penile cavernous tissue of PGE_1 and 19R-OH- PGE_2 are presented in Table I. It can be seen that PGE_1 and 19R-OH- PGE_2 were about equipotent and comparable to carbachol, however PGE_1 was slightly more potent than 19R-OH- PGE_2 . Since 19R-OH- PGE_2 is a selective EP_2 receptor agonist this finding indicates that the EP_2 receptor accounts for most of the relaxant effect of PGE_1 . Furthermore, in FIG. 1 it is demonstrated that the concentration-response curves of PGE_1 and 19R-OH- PGE_2 are parallel and differ only with a factor of about 2–3, i.e. the latter analogue is about half to one third as potent as the former.

TABLE I

Relaxation of human corpus cavernosum tissue induced by prostaglandins compared to carbachol (10^{-6} M).			
(Mean \pm SEM)			
Compound	n	Reduction in tension (%)	EC-50 (Moles/l)
Carbachol	8	100.0 \pm 0.0	—
PGE_1	8	90.9 \pm 3.7	2.5×10^{-7}
19R-OH- PGE_2	8	96.9 \pm 3.1	6.9×10^{-7}

The vasodilatory effect of PGE_1 and the EP_2 receptor agonist 19R-OH- PGE_2 on rabbit penile blood vessels is demonstrated in FIG. 2. It can be seen that the two prostaglandins had the same efficacy in inducing vasodilatation and were also about equipotent. This furthermore demonstrates that the EP_2 receptor mediates most of the vasodilatory effect of PGE_1 . The vasodilatory effect of all the prostaglandin analogues as studied in the rat by investigating the immediate reduction in blood pressure upon intravenous infusion is presented in Table II. As can be seen in the table PGE_1 and the three hydroxy-substituted PGE_2 analogues effectively reduced the blood pressure in the anaesthetized rat, demonstrating an acute vasodilatory response.

TABLE II

Prostaglandin analogue	2.5-3.5 $\mu\text{g}/\text{kg}$		7.5-10 $\mu\text{g}/\text{kg}$		25-35 $\mu\text{g}/\text{kg}$	
	Blood pressure o (mmHg)	Reduction (%)	Blood pressure o (mmHg)	Reduction (%)	Blood pressure o (mmHg)	Reduction (%)
PGE ₁	114 \pm 20	13 \pm 4	105 \pm 15	35 \pm 5	103 \pm 9	52 \pm 3
18RS-OH-PGE ₂	122 \pm 28	7 \pm 3	123 \pm 33	15 \pm 3	115 \pm 32	32 \pm 2
19R-OH-PGE ₂	115 \pm 14	13 \pm 10	103 \pm 12	31 \pm 3	91 \pm 9	43 \pm 6
20-OH-PGE ₂	91 \pm 11	26 \pm 5	85 \pm 10	38 \pm 3	91 \pm 6	44 \pm 0

Blood pressure o = blood pressure before infusion of prostaglandin analogue.

The irritant effects of the three prostaglandin analogues and PGE₁ as studied in the feline eye are presented in Table III. All three hydroxy-substituted PGE₂-analogues had significantly less irritant effect than PGE₂-isopropyl ester and PGE₁ acid. Most surprisingly we found that both 18RS-OH-PGE₂-methyl ester and 19R-OH-PGE₂-methyl ester had no or markedly reduced irritant effect, even at doses 1000 times higher than PGE₁-isopropyl ester. Previously it has been shown that 19R-OH-PGE₂ does not cause lid closure in rabbits but induces an irritant effect by inducing swelling of the ocular structures (Hall and Jaidly, 1977). We found no

PGE₁ acid, which is a much less lipophilic compound than the 18-, 19- and 20-hydroxy-substituted PGE₂-methyl ester analogues, caused significantly more irritation, and at much lower dose, than the hydroxy-substituted analogues. Thus the absence of pain and irritation after administration of 18RS-OH-PGE₂, 19R-OH-PGE₂ and 20OH-PGE₂ cannot be explained on the basis of increased hydrophilicity and poor bioavailability. Accordingly, selective EP₂ prostanoid receptor agonists seem to be very advantageous in that they exhibit no, or markedly reduced irritant effect.

TABLE III

Analogue	log p-value	Maximum irritative effect of prostaglandin analogues as studied in cat eyes. The log P values were estimated based on thin layer chromatography with PGF _{2α} isopropyl ester as reference (log P 4.5). ie = isopropyl ester, me = methyl ester. Maximum irritation = 3. Mean \pm SEM (n = 3-6).							
		Dose: 0.01 μg	Dose: 0.1 μg	Dose: 0.3 μg	Dose: 1 μg	Dose: 3 μg	Dose: 10 μg	Dose: 30 μg	Dose: 100 μg
PGE ₁ -ie	4.7	0.4 \pm 0.2	1.7 \pm 0.3	2.8 \pm 0.1	—	—	—	—	—
PGE ₁ acid	0.7	—	0.6 \pm 0.1	—	1.6 \pm 0.2	—	3.0 \pm 0.0	—	—
18RS-OH-PGE ₂ -me	2.7	—	—	—	0.0 \pm 0.0	0.0 \pm 0.0	0.0 \pm 0.0	0.3 \pm 0.2	—
19R-OH-PGE ₂ -me	2.5	—	0.0 \pm 0.0	—	0.1 \pm 0.1	—	0.1 \pm 0.1	—	0.7 \pm 0.4
20-OH-PGE ₂ -me	2.2	—	—	—	—	0.5 \pm 0.0	1.2 \pm 0.2	1.2 \pm 0.4	—

evidence of such irritation in the cat eye with 19R-OH-PGE₂ or the other hydroxy-substituted PGE₂ analogues.

To confirm that 19R-OH-PGE₂-methyl ester, which is more hydrophilic than PGE₁-isopropyl ester, indeed penetrates into the cornea and the intraocular parts of the eye we measured the intraocular pressure in 3 cats under local anaesthesia before and 1 hour after topical administration of the prostaglandin to the eye. The intraocular pressure was measured by pneumatonometry. One eye was treated with the test compound and the other eye received the vehicle only. In the eyes treated topically with 19R-OH-PGE₂-methyl ester the intraocular pressure decreased from 16.3 \pm 0.9 mmHg to 12.7 \pm 1.2 mmHg, whereas it was 16.3 \pm 0.9 mmHg and 15.7 \pm 0.7 mmHg in the control eyes at the same time points. It is well known that prostaglandins reduce the intraocular pressure in cats (Bito et al., 1989), and it can thus be taken as an evidence that the drug has penetrated into the eye. Furthermore, from Table III it can also be seen that the

References

- Anawalt, B. D., Bebb, R. A. and Matsumoto, A. M. 1996. Medical therapy for erectile dysfunction. *Current Opinion in Endocrinology and Diabetes*. 3; 472-477.
- Bito, L. Z., Carmras, C. B., Gum, G. G. et al. 1989. The ocular hypotensive effect and side-effects of prostaglandins on the eyes of experimental animals. In: *The ocular effects of prostaglandins and other eicosanoids* (Eds: L. Z. Bito and J. Stjemschantz). Alan R. Liss Inc. New York. pp:349-368.
- Coleman, R. A., Smith, W. L. and Narumiya, S. 1994. VIII. International Union of Pharmacology classification of prostanoid receptors: Properties, distribution, and structure of the receptors and their subtypes. *Pharmacological Reviews*. 46; 205-229.
- Hall, D. W. R. and Jaidly, K. D. 1977. Inflammatory responses of the rabbit eye to prostaglandins. *Agents-Actions-Suppl.* 2: 123-133.
- NIH Consensus Conference on Impotence. 1993. *JAMA*. 270;83-90.

Pada-Nathan, H., Hellstrom, W. G. J., Kaiser, F. E. et al. 1997. Treatment of men with erectile dysfunction with transurethral alprostadil. *New Eng. J. Med.* 336;1-7.

Taylor, P. L. and Kelly R. W. 1974. 19-hydroxylated E prostaglandins of human semen. *Nature.* 250; 665-667. 5

Wolfson, B., Pickett, S., Scott, N. E. et al. 1993. Intraurethral prostaglandin E-2 cream: A possible alternative treatment for erectile dysfunction. *Urology.* 42; 73-75.

What is claimed is:

1. A method of treating impotence or erectile dysfunction with reduced pain as compared with a method of treating impotence or erectile dysfunction using PGE₁ or PGE₂, the method comprising administering to a human a therapeutically effective and physiologically acceptable amount of 18, 19, or 20-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof. 10 15

2. The method according to claim 1, wherein 18-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof, is administered.

3. The method according to claim 1, wherein 19-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof, is administered. 20

4. The method according to claim 1, wherein 19-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof, is administered. 25

5. The method according to claim 1, wherein methyl or isopropyl ester of 19-OH-PGE₂, is administered.

6. The method according to claim 1, wherein 20-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof, is administered. 30

7. The method according to claim 1, wherein 18, 19, or 20-OH-PGE₂ is administered to the patient by intracavernous injection or by transurethral or transdermal application.

8. A method of treating impotence or erectile dysfunction with reduced pain as compared with a method of treating impotence or erectile dysfunction using PGE₁ or PGE₂, the method comprising administering to human a composition comprising a therapeutically effective and physiologically acceptable amount of an active agent adapted to treat erectile dysfunction or impotence with reduced pain as compared with administration of PGE₁ or PGE₂, wherein the active agent consists essentially of 18, 19, or 20-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof.

9. The method according to claim 8, wherein the active agent consists essentially of 18-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof.

10. The method according to claim 8, wherein the active agent consists essentially of 19-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof.

11. The method according to claim 8, wherein the active agent consists essentially of 19-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof.

12. The method according to claim 8, wherein the active agent consists essentially of methyl or isopropyl ester of 19-OH-PGE₂.

13. The method according to claim 8, wherein the active agent consists essentially of 20-OH-PGE₂, or a pharmaceutically acceptable salt or ester thereof.

14. The method according to claim 8, wherein the active agent is administered to the patient by intracavernous injection or by transurethral or transdermal application.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,476,074 B1
DATED : November 5, 2002
INVENTOR(S) : Johan Stjernschantz et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 17,

Lines 23-24 and 27, change "19-OH-PGE₂" to -- 19R-OH-PGE₂ --.

Column 18,

Line 20, change "19-OH-PGE₂" to -- 19R-OH-PGE₂ --.

Signed and Sealed this

Eighteenth Day of March, 2003

A handwritten signature in black ink, appearing to read "James E. Rogan", written over a horizontal line.

JAMES E. ROGAN
Director of the United States Patent and Trademark Office