



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/56, 9/24	A1	(11) International Publication Number: WO 92/17592 (43) International Publication Date: 15 October 1992 (15.10.92)
(21) International Application Number: PCT/NL92/00056 (22) International Filing Date: 20 March 1992 (20.03.92) (30) Priority data: 91200720.0 27 March 1991 (27.03.91) EP <i>(34) Countries for which the regional or international application was filed:</i> NL et al. (71) Applicant (for all designated States except US): GIST-BROCADES N.V. [NL/NL]; Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL). (72) Inventors; and (75) Inventors/Applicants (for US only) : VAN HEUVEL, Margaretha [NL/NL]; Cobetstraat 19, NL-2313 KA Leiden (NL). BAKHUIS, Janna, Gardina [NL/NL]; Van der Haertstraat 72, NL-2613 ZC Delft (NL). COUTEL, Yves [FR/FR]; 56, rue Culmette-Guirin, F-59710 Ennevelin (FR). HARDER, Abraham [NL/NL]; De Vaert 9, NL-2651 EP Berkel en Rodenrijs (NL). DE GRAAFF, Leendert, Hendrik [NL/NL]; Cornelis Koningstraat 8, NL-6862 CK Oosterbeek (NL). FLIPPPI, Michel, Johannes, Anthonie [NL/NL]; Thorbeckestraat 234, NL-6702 BZ		Wageningen (NL). VAN DER VEEN, Peter [NL/NL]; A. Kuiperstraat 10, NL-6703 BL Wageningen (NL). VISSER, Jacob [NL/NL]; Hinkeloordseweg 5, NL-6703 CK Wageningen (NL). ANDREOLI, Peter, Michael [BE/BE]; Bellegemsestraat 57, B-8510 Bellegem-Kortrijk (BE). (74) Agents: HUYGENS, Arthur, Victor et al.; Gist-Brocades N.V., Patents and Trademarks Department, Wateringseweg 1, P.O.Box 1, NL-2600 MA Delft (NL). (81) Designated States: AU, CA, FI, JP, NO, PL, US. Published <i>With international search report.</i>
(54) Title: CLONING AND EXPRESSION OF DNA MOLECULES ENCODING ARABINAN-DEGRADING ENZYMES OF FUNGAL ORIGIN (57) Abstract The present invention provides the cloning and expression of purified and isolated DNA molecules, obtainable from fungi, which encode enzymes having arabinan-degrading activity. The present invention also provides DNA constructs containing these DNA molecules and methods for their enhanced expression in selected microbial host cells. The present invention further provides methods for the use of the thus-produced arabinan-degrading enzymes.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
DE	Germany	MC	Monaco	TG	Togo
DK	Denmark			US	United States of America

Cloning and Expression of DNA Molecules Encoding
Arabinan-Degrading Enzymes of Fungal Origin

The present invention relates to the field of molecular biology. In particular, the present invention relates to the cloning and expression of genes encoding enzymes of fungal origin.

Background of the Invention

The composition of the plant cell wall is complex and variable. Polysaccharides are mainly found in the form of long chains of cellulose (the main structural component of the plant cell wall), hemicellulose (comprising various β -xylan chains) and pectic substances (consisting of galacturonans and rhamnogalacturonans; arabinans; and galactans and arabinogalactans). From the standpoint of the food industry, the pectic substances, arabinans in particular, have become one of the most important constituents of plant cell walls (Whitaker, J.R. (1984) Enzyme Microb. Technol., 6, 341).

Arabinans consist of a main chain of α -L-arabinose subunits linked α -(1 $\rightarrow$ 5) to one another. Side chains are linked α -(1 $\rightarrow$ 3) or sometimes α -(1 $\rightarrow$ 2) to the main α -(1 $\rightarrow$ 5)-L-arabinan backbone. In apple, for example, one third of the total arabinose is present in the side chains. The molecular weight of arabinan is normally about 15 kDa.

Enzymes capable of degrading arabinans are becoming increasingly important to the food industry. In juice production, for example, the demand to increase yields in order to reduce production costs has necessitated the modification of traditional processes. The utilization of enzymatic pre-treatments of the fruit pulp before pressing with specific enzymatic products drastically improves the juice yield by solubilizing the cell wall polysaccharides.

- 2 -

However, a persistent turbidity commonly referred to as "arabinan haze" has been a source of problems in the production of concentrated juices. The arabinan haze is more often present in concentrated juice than in non-concentrated juice. This may indicate that water activity has an influence on the solubility of arabinan. Furthermore, it has been found that this haze is soluble between 60 and 80°C.

It has also been found that while branched arabinan is soluble in concentrated chilled apple and pear juices, the linear, debranched α -(1 $\rightarrow$ 5)-L-arabinan is much less soluble. This debranched α -(1 $\rightarrow$ 5)-L-arabinan is formed from the L-arabinan by the action of an arabinan-degrading enzyme present in the commercial pectic enzyme preparations from Aspergillus niger, commonly used to increase juice yield after pulp treatment but before pressing.

On the other hand, debranched arabinans are considered desirable for certain other applications. WO 90/06343 discloses the debranching of sugar beet arabin by the action of an α -L-arabinofuranosidase, free of endo arabinanase activity, which is isolated from a culture filtrate of Aspergillus niger or from a commercial pectinase mixture using ion-exchange and gel filtration chromatography procedures. The debranched arabin may be used as a fat substitute in foods.

Arabinan-degrading enzymes are known to be produced by a variety of plants and microorganisms, among these, fungi such as those of the genera Aspergillus, Corticium, Rhodotorula (Kaji, A. (1984) Adv. Carbohydr. Chem. Biochem., 42, 383), Dichotomitus (Brillouet et al. (1985) Carbohydrate Research, 144, 113), Ascomycetes and Basidiomycetes (Sydow, G. (1977) DDR Patent Application No. 124,812).

In particular, the filamentous fungus Aspergillus niger is known to produce three different arabinan-degrading enzymes: an α -L-arabinanase having a molecular weight of approximately 35 kDa and two α -L-arabinofuranosidases having

molecular weights of approximately 118 and 60 kDa, respectively, (Rombouts et al. (1988) Carbohydrate Polymers, 9, 25). [N.B. van der Veen et al. ((1991) Arch. Microbial., 157, 23) reports molecular weights of 43, 83 and 67 kDa for these same three enzymes, respectively.]

The 35 kDa arabinanase (also known as ABN A) has endo activity and exclusively cleaves 1→5 linkages. The activity of this enzyme decreases as the 1,5- α -L-arabinan sequences become shorter and the concentration of arabinose dimers and trimers increased (Rombouts et al., supra). [N.B. van der Veen et al. (1991) reported a molecular weight of 43 kDa.]

The 118 kDa α -L-arabinofuranosidase (also known as arabinofuranosidase A, ABF A or EXO A) exclusively cleaves 1→5 linkages with exo-type activity as shown by the accumulation of arabinose monomers. This enzyme displays the highest activity on low molecular weight substrates (Rombouts et al., supra). [N.B. van der Veen et al. (1991) reported a molecular weight of 83 kDa.]

The 60 kDa α -L-arabinofuranosidase (also known as arabinofuranosidase B, ABF B or EXO B), also having exo-type activity, predominantly cleaves 1→3 and 1→2 α -L-arabinans, yet also demonstrates the ability to cleave 1→5 α -L-arabinans. Again, only arabinose monomers are detected after degradation of an α -L-arabinan-containing substrate (Rombouts et al., supra). [N.B. van der Veen et al. (1991) reported a molecular weight of 67 kDa.]

The A. niger ABF B enzyme has also demonstrated the ability to cleave the 1→6 linkage between a terminal arabinofuranosyl unit and the intermediate glucose of monoterpenyl α -L-arabinofuranosylglucosides (Gunata et al. (1989) European Patent Application No. 332.281; Gunata et al. (1990) J. Agric. Food Chem., 38, 772).

Enzymes having an activity similar to that of the A. niger ABF B enzyme have been identified in other fungi such as

Dichotomitus squalens (Brillouet et al., supra), Corticium rolfsii and Rhodotorula flava (Kaji, A., supra).

5 A possible solution to the problem of haze formation in juices may be to use an enzyme formulation having an improved balance between the endo arabinanase (ABN A) and the ABF A and ABF B enzymes, or alternatively, enzymes having similar activities, i.e. one or more arabinofuranosidases obtained from another microorganism which is capable of cleaving 1→3 and 1→2 α -L-arabinans, as well as 1→5 α -L-arabinans. However, 10 normal fermentation of A. niger fails to yield sufficient levels of the desired arabinofuranosidases. Moreover, it would be advantageous to be able to attenuate the amount of ABF A, ABF B and ABN A activities for optimal results in specific applications.

15 The ability of the ABF B enzyme to cleave the terminal 1→6 linkage of monoterpenyl α -L-arabinofuranosylglucosides may be used to assist in the release of aroma components from various fruit juices and thus, the enhancement of their flavors (Gunata, Z. et al. (1989 and 1990), supra). However, 20 it would be preferable to use a purified ABF B or ABF B-like enzyme for this purpose since the presence of other enzymatic activity may degrade other important components of the juice and, in so doing, have a detrimental effect on the ultimate quality of the juice.

25 In nature, microbial arabinan-degrading enzymes are always produced together with other enzymes having polysaccharide-degrading activities, such as pectinases, xylanases, acetyl xylan esterases, acetyl esterases, galactanases and cellulases. As mentioned above, for some 30 applications, these enzyme activities are not needed or are unwanted. Moreover, due to relatively low expression levels in the wild-type strains, the arabinan-degrading enzymes have proven to be somewhat difficult to isolate both from each other and from other enzymes produced by A. niger.

It is known that fermentation conditions may be varied to favor the production of an enzyme of interest. It is also known that the cloning of the gene encoding the desired enzyme and overexpressing it in its natural host, or in another compatible expression host will specifically enhance the production of the enzyme of interest. This latter method is particularly useful if the enzyme of interest is to be obtained in increased amounts, or in a form which is free of other, undesired enzymatic activity.

Clearly, it would be useful to increase arabinan-degrading activity via recombinant DNA techniques. However, until now, the genes encoding the A. niger arabinan-degrading enzymes have not been available. Accordingly, it would be of great importance to obtain genes encoding arabinan-degrading enzymes of fungal origin which may be brought to expression in its native host or, alternatively, in other microbial hosts wherein high expression levels of one or more arabinan-degrading enzymes may be achieved.

The expression of recombinant bacterial α -L-arabinofuranosidase has been previously described by Schwarz et al. ((1990) Biochem. Biophys. Res. Commun., 170, 368). The gene encoding a bacterial arabinofuranosidase was isolated from Clostridium stercoarium and brought to expression in an E. coli host.

A gene encoding an arabinosidase was cloned from the anaerobic bacterium Bacteriodes ovatus and brought to expression in E. coli, as disclosed by Whitehead & Hespell ((1990) J. Bacteriol., 172, 2408).

However, E. coli expression hosts are, in some instances, considered to be unsafe for the production of proteins by recombinant DNA methods due to their production of unacceptable by-products such as toxins. Furthermore, in E. coli expression, large amounts of heterologous protein tend to accumulate inside the cell. Subsequent purification of the

desired protein from the bulk of E. coli intracellular proteins can sometimes be difficult.

Moreover, since bacterial genes contain no introns, one is confronted with few problems in cloning and expressing such genes in prokaryotic hosts. On the other hand, the expression of eukaryotic genes is not always so straightforward. It is well known that genes isolated from eukaryotic strains may contain introns. This inherently introduces complications in the cloning and expression of these genes, should a prokaryotic host be preferred.

Furthermore, certain differences exist between the physical characteristics of arabinan-degrading enzymes of fungal origin and those from bacteria. In general, fungal enzymes have a pH optimum in the range from $\leq 3.0 - 6.0$. A few fungal species produce arabinan-degrading enzymes having pH optima as low as pH 2.0. These pH optima are generally significantly lower as compared to similar enzymes from bacterial strains which have a pH optimum in the range of pH 5.0 - 7.0 (Karimi and Ward (1989) J. Indust. Microbiol., 4, 173; Lee and Forsberg (1987) Can. J. Microbiol., 33, 1011). Thus, it is clear that bacterial arabinan-degrading enzymes are less suitable for use in, for example, processes requiring lower pH conditions.

Summary of the Invention

The present invention provides purified and isolated DNA molecules of fungal origin, each encoding an enzyme having arabinan-degrading activity.

It is also an object of the present invention to provide DNA constructs for the enhanced microbial expression of the DNA molecules of the present invention. The DNA (or expression) constructs provided contain the DNA molecule encoding the desired arabinan-degrading enzyme with either its native 5' and 3' regulatory regions or, in an alternative

- 7 -

embodiment, hybrid DNA constructs are provided in which the DNA molecule is operably linked to regulatory regions such as promoter, secretion leader and terminator signals which may be selected to provide for the optimal expression of the enzyme in the desired expression host and, if desired, the secretion therefrom.

It is a further object of the present invention to provide microbial expression hosts, transformed with one or more DNA constructs of the present invention, which are capable of the enhanced expression and, if desired, the secretion of arabinan-degrading enzymes of fungal origin.

It is yet a further object of the present invention to provide methods for the production of large quantities of arabinan-degrading enzymes of interest which may be advantageously used in industrial processes. Typically, industrial processes which require the activity of arabinan-degrading enzymes at a lower pH than that at which similar enzymes of bacterial origin optimally function.

Brief Description of the Figures

- Figure 1 Schematic diagram of the purification procedure for obtaining the α -L-arabinofuranosidase (ABF B) enzyme from A. niger.
- Figure 2 Amino acid sequence of the N-terminus of A. niger ABF B protein (Formula 1) and the nucleotide sequence of the oligonucleotide mixture AB1719 (Formula 1a).
- Figure 3 Amino acid sequence of the N-terminus of the 15 kDa CNBr fragment of the A. niger ABF B protein (Formula 3) and the nucleotide sequence of the oligonucleotide mixture AB2306 (Formula 3a).
- Figure 4 Partial restriction enzyme map of the hybridizing phage clone 7.

- Figure 5 Nucleotide sequence of the A. niger abfB gene (Formula 4).
- Figure 6 Amino acid sequence (Formula 5) of the A. niger ABF B protein, derived from the abfB gene sequence and confirmed by the abfB cDNA sequence.
- Figure 7 Physical map of pAB 6-1. The 14.5 kb HindIII DNA insert in pUC19 contains the entire amyloglucosidase (AG) locus from A. niger.
- Figure 8 A schematic view of the generation of AG/abfB gene fusions by the polymerase chain reaction (PCR). The sequences of all oligonucleotide primers used are indicated in the text.
- Figure 9 A schematic representation of the construction of pAGabfB2 and pAGabfB3. Key to the symbols used:
- ▨ : sequences encoding the 18 amino acid signal peptide of the A. niger amyloglucosidase (AG) gene (glaA).
- : sequences encoding the mature ABF B protein
- Amp : the ampicillin resistance gene
- ori : E. coli origin of replication
- AG : the A. niger amyloglucosidase (AG) promoter
- Figure 10 The construction of the expression cassette pAGabfB4. Key to the symbols used:
- pyrA : the A. niger gene coding for orotidine-5'-phosphate-decarboxylase
- The remaining symbols are the same as those presented in the legends of Figure 9.
- Figure 11 The construction of the expression cassette pAGabfB5. Key to the symbols used:
- amdS : the A. nidulans gene coding for an acetamidase

The remaining symbols are the same as those presented in the legends of Figure 9.

- Figure 12 Restriction map of cDNA clone pClX1. Only the KpnI,
5 PstI and XhoI sites are indicated. Restriction
sites from the polylinker of the pBluescript SK⁻
vector part of pClX1 are denoted between brackets
and not drawn on scale.
- Figure 13 Restriction map of pIM900. The bar represents the
10 8.5 kb XbaI/NsiI fragment cloned in the vector
pGEM-7Zf(+). The hatched box represents the
approximate position of the abfA gene. Only the
KpnI, PstI and XhoI sites are indicated.
- Figure 14 Nucleotide sequence of the A. niger abfA gene
(Formula 20).
- 15 Figure 15 Amino acid sequence (Formula 21) of the A. niger
ABF A protein derived from the abfA cDNA sequence.
- Figure 16 A: Analysis of culture filtrate of A. nidulans abfA
transformants and wild type WG096 grown for 24
hours on sugar beet pulp by SDS-PAGE followed by
20 Coomassie Brilliant Blue R250 staining.
lane 1: 1 μ g purified α -L-arabinofuranosidase A
lane 2: WG096
lanes 3 to 6: randomly chosen abfA transformants
B: Analysis of culture filtrate of A. niger abfA
25 transformants and wild type N402 grown for 24 hours
on sugar beet pulp by Western blot analysis.
lane 1: 1 μ g purified α -L-arabinofuranosidase A
lane 2: N402
lanes 3 to 6: randomly chosen abfA transformants
- 30 Figure 17 Analysis of the induction pattern of ABF B and ABN
A in A. niger N572 by Western blot analysis of
culture filtrates using anti ABF B and anti ABN A
antibodies.
lane 1: 24 hours growth on 1% D-glucose

- lane 2: 1 hour induction on 1% L-arabitol
lane 3: 2 hours induction on 1 % L-arabitol
lane 4: 3 hours induction on 1 % L-arabitol
lane 5: 4 hours induction on 1 % L-arabitol
5 lane 6: 5 hours induction on 1 % L-arabitol
lane 7: 6 hours induction on 1 % L-arabitol
lane 8: 7 hours induction on 1 % L-arabitol
lane 9: 1 μ g purified α -L-arabinofuranosidase B
- 10 Figure 18 Restriction map of pIM950. The bar represents the 3.1 HindIII fragment cloned in the vector pEMBL19. The hatched box represents the approximate position of the abnA gene.
- Figure 19 Nucleotide sequence of the A. niger abnA gene (Formula 22).
- 15 Figure 20 Amino acid sequence (Formula 23) of the A. niger ABN A protein derived from the abnA cDNA sequence.
- Figure 21 Analysis of culture filtrate of A. nidulans abnA transformants and wild type WG096 grown for 24 hours on sugar beet pulp by Western blot analysis.
- 20 lane 1: 1 μ g purified endo 1,5- α -L arabinanase from A. niger
lane 2: 1 μ g purified endo 1,5- α -L arabinanase from A. nidulans
lanes 3 and 4: A. nidulans WG096
- 25 lane 5: Transformant #20
lane 6: Transformant #18
lane 7: Transformant #17
lane 8: Transformant #16
lane 9: Transformant #12
- 30 lane 10: Transformant #11
lane 11: Transformant #7
lane 12: Transformant #6
lane 13: Transformant #2
lane 14: Transformant #1

- 11 -

Figure 22 Western blot analysis of culture filtrates of A. niger abnA transformants and wild type N402 grown for 24 hours on sugar beet pulp by Western blot analysis.

5

A:

lane 1: 1 μ g purified endo 1,5- α -L arabinanase
from A. niger

lane 2: N402

lanes 3 to 9: Transformants 1 to 7

10

B:

lane 1: 1 μ g purified endo 1,5- α -L arabinanase
from A. niger

lane 2: N402

lanes 3 to 9: Transformants 8 to 14

15

Detailed Description of the Invention

The present invention provides purified and isolated DNA molecules obtainable from fungi, and genetic variants thereof, which encode enzymes having arabinan-degrading activity. Genetic variants include DNA molecules encoding mutant arabinan-degrading proteins and degenerate DNA molecules wherein the desired activity of the enzyme expressed therefrom is retained.

The present invention also provides DNA constructs (also referred to herein as expression constructs) for the expression of one or more arabinan-degrading enzymes in a desired expression host. The DNA constructs provided may comprise the DNA molecule encoding the desired arabinan-degrading enzyme with its native 5' and 3' regulatory regions. Alternatively, hybrid DNA constructs are provided which comprise a DNA molecule encoding the arabinan-degrading enzyme operably linked to regulatory regions, such as promoter, secretion and terminator signals originating from homologous or heterologous organisms, these regulatory regions being

- 12 -

capable of directing the enhanced expression of the structural gene in an appropriate host. Preferably, the expression construct will be integrated into the genome of the selected expression host.

5 The present invention further provides vectors, preferably plasmids, for the cloning and/or transformation of microbial hosts via the introduction into the microbial host of the DNA constructs for the expression of the desired arabinan-degrading enzyme.

10 In addition, the present invention provides microbial hosts transformed with one or more vectors, each containing at least one of the DNA constructs described above. Microbial expression hosts may be selected from bacteria, yeasts or fungi.

15 Within the context of the present invention, the term "enhanced expression" is defined as the expression of the DNA construct whereby the arabinan-degrading enzyme of interest is produced at levels higher than that which are ordinarily encountered in the homologous wild-type organism. In the same
20 context, enhanced expression also intends expression in a heterologous organism which does not normally produce such arabinan-degrading enzyme except for the introduction of the DNA molecule encoding the arabinan-degrading enzyme of interest, together with appropriate regulatory regions, into
25 the heterologous expression host. Progeny of these expression hosts are, of course, also to be understood to be embraced by the present invention.

 Within the context of the present invention, the term "homologous" is understood to intend all that which is native
30 to the DNA molecule encoding the arabinan-degrading enzyme of interest, including its regulatory regions. A homologous host is defined as the species from which such DNA molecule may be isolated.

The term "heterologous" is thus defined as all that which is not native to the DNA molecule encoding the arabinan-degrading enzyme of interest itself, including regulatory regions. A "heterologous" host is defined as any microbial species other than that from which the arabinan-degrading enzyme-encoding gene has been isolated.

Within the scope of the present invention, the term "arabinan-degrading enzyme of interest" is understood to include arabinan-degrading enzymes of fungal origin.

Arabinan-degrading activity, as defined in the context of the present invention, is the ability of an enzyme to release arabinose residues, either monomers or oligomers, from arabinan backbones or from arabinan-containing sidechains of other hemicellulose backbone structures such as arabinoxylans or arabinogalactans, or even the release of arabinose monomers via the cleavage of the 1→6 linkage between the terminal arabinofuranosyl unit and the intermediate glucosyl unit of monoterpenyl α -L-arabinofuranosyl glucosides.

Examples of preferred "arabinan-degrading activity" may be selected from:

- a) the ability to cleave (1→2)- α -L-arabinosidic linkages;
- b) the ability to cleave (1→3)- α -L-arabinosidic linkages;
- c) the ability to cleave (1→5)- α -L-arabinosidic linkages;
- d) the ability to cleave the 1→6 linkage between the terminal arabinofuranosyl unit and the intermediate glucosyl unit of monoterpenyl α -L-arabinofuranosyl glucosides.

Most preferred arabinan-degrading enzymes are α -L-arabinofuranosidases and α -L-arabinanases which have 1) exo-type arabinan-degrading activity on (1→5)- α -L-arabinosidic linkages; or 2) exo-type arabinan-degrading activity on (1→3)-

- 14 -

α -L-arabinosidic linkages and (1 $\rightarrow$ 2)- α -L-arabinosidic linkages; 3) endo-type arabinan-degrading activity on (1 $\rightarrow$ 5)- α -L-arabinosidic linkages; or 4) the ability to cleave the 1 $\rightarrow$ 6 linkage between the terminal arabinofuranosyl unit and the intermediate glucosyl unit of monoterpenyl α -L-arabinofuranosyl glucosides.

Preferred DNA molecules encoding arabinan-degrading enzyme of the present invention are those which are obtainable from filamentous fungi of the genera Aspergillus (especially A. niger, A. niger var. tubigensis (see also Kusters-van Someren (1991) Curr. Genet., 19, 21), A. niger var. awamori, A. nidulans (with the exception of the abfA gene) and A. aculeatis), Dichotomitus (especially D. squalens), Corticium (especially C. rolfsii), Penicillium (especially P. chrysogenum) and Rhodotorula (especially R. flava). Especially preferred are the α -L-arabinanase (ABN A), and the α -L-arabinofuranosidases (ABF A and ABF B) which are obtainable from Aspergillus niger, Aspergillus niger var. tubigensis or Aspergillus niger var. awamori.

The present invention also includes DNA sequences which hybridize under low stringency conditions to the DNA sequences obtainable from the fungi described above, but which may differ in codon sequence due to the degeneracy of the genetic code or cross-species variation.

Arabinofuranosidases having the desired activity may be identified by various methods not critical to the present invention. For example, assays for the presence of para-nitrophenol or arabinose monomers are indicative of the ability of the enzyme to cleave 1 $\rightarrow$ 3 or 1 $\rightarrow$ 2- α -L-arabinosidic linkages using para-nitrophenyl- α -L-arabinoside or other arabinan-containing substrates, respectively, using methods such as those described by Rombouts et al. ((1988) supra) or van der Veen et al. ((1991) supra). The ability of an enzyme to cleave 1 $\rightarrow$ 5- α -L-arabinosidic linkages may be determined

using an apple juice ultrafiltration retentate (UFR) arabinan substrate and assaying for the presence of small arabinoside oligomers as described by Rombouts et al., supra. Alternatively, endo 1,5- α -L-arabinanase activity of a culture filtrate may be measured using a test kit such as Arabina-ZymeTM tablets as produced by Megazyme Pty. Ltd. (North Rocks, New South Wales, Australia).

Once an arabinan-degrading enzyme of interest has been identified, the DNA molecule encoding such arabinan-degrading enzyme may be obtained from the filamentous fungus which naturally produces it by culturing the fungus in a suitable medium, isolating the desired arabinan-degrading enzyme using known methods such as those outlined in Figure 1 (ABF B enzyme) or those as described by Rombouts et al. ((1988) supra) or van der Veen et al. ((1991) supra) and determining at least a portion of the amino acid sequence of the purified protein.

DNA probes may thereafter be obtained by designing oligonucleotides based on the deduced partial amino acid sequence. Amino acid sequences may be determined from the N-terminus of the complete protein and/or from the N-termini of internal peptide fragments obtained via proteolytic or chemical digestion of the complete protein. Once obtained, the DNA probe(s) are then used to screen a genomic or cDNA library.

A genomic library may be prepared by partially digesting the fungal chromosomal DNA with a restriction enzyme which recognizes a DNA sequence of four successive nucleotides, e.g. Sau3A, and cloning the resulting fragments in a suitable plasmid or lambda phage vector, e.g. lambda GEM-11.

Alternatively, a cDNA library may be prepared by cloning cDNA, synthesized from mRNA isolated from fungal cells induced for the synthesis of arabinan-degrading enzyme, into an

appropriate phage vector, e.g. lambda gt10, such as described by Harmsen et al. (1990) Curr. Genet., 18, 161.

Subsequently, after plating of a sufficient amount of colonies or plaques, the genomic or cDNA library may be
5 screened with a suitable DNA probe.

If this method is unsuccessful, the genomic or cDNA library may be differentially screened with cDNA probes obtained from mRNA from non-induced and induced cells. Induced mRNA is prepared from cells grown on media containing arabinan
10 as a carbon source, while non-induced mRNA must be isolated from cells grown on a carbon source other than arabinan, e.g. glucose. Among the clones which only hybridize with the induced cDNA probe, a clone containing the gene encoding the desired arabinan-degrading enzyme may be recovered.
15 Alternatively, a gene encoding an arabinan-degrading enzyme of interest may be identified by cross-hybridization with a gene from a related arabinan-degrading enzyme.

In the case of the ABF B enzyme, oligonucleotide probes are obtained from the N-terminal amino acid sequence (see
20 Figure 2, Formula 1) of an arabinan-degrading enzyme having an apparent molecular weight of 60 kDa (in glycosylated form) purified from an Aspergillus niger culture filtrate and/or from the amino acid sequence of an internal peptide fragment (see Figure 3, Formula 3) obtained by digestion of the enzyme
25 with CNBr. The oligonucleotide mixtures AB1719 (Figure 2, Formula 1a) and AB2306 (Figure 3, Formula 3a) are complementary to the corresponding deduced arabinofuranosidase mRNA. Four positive phage clones were obtained from the screening of a lambda GEM-11 library, prepared from partially
30 Sau3A digested DNA isolated from Aspergillus niger, with the N-terminal oligo mixture AB 1719.

DNA isolated from the four phage clones hybridized with the N-terminal oligonucleotide mixture AB1719 (see Figure 2, Formula 1a). A 2.8 kb SacI fragment was obtained (see Figure

4) and has been sequenced. The nucleotide sequence and encoded amino acid sequence are depicted in Figures 5 and 6 (Formulas 4 and 5, respectively). The SacI fragment, in plasmid pGBabfB1, was deposited in E. coli strain DH5 α at the Centraal Bureau voor Schimmelcultures (Baarn, the Netherlands on March 11, 1991, under accession number CBS 156.91).

The abfB gene encodes a protein 499 amino acids in length having a deduced molecular weight of 52523 Da (Figure 6, Formula 5), as derived from the abfB gene sequence (Figure 5) and confirmed by the abfB cDNA sequence. The N-terminal amino acid sequence, as determined in Example 12.2 (Formula 14) is preceded by a 18 amino acid-long hydrophobic sequence. The amino acid sequences determined from the CNBr peptides (Formulas 15, 16 and 17) are found in the sequence from amino acid position 203 until position 219, 267 until 286 and 294 until 312, respectively. The mature ABF B protein is 481 amino acids in length, and the unglycosylated protein has a deduced molecular weight of 50663 Da and a theoretical IEP of 3.8.

In the case of the ABF A enzyme, a cDNA library was made after the induction of mRNA in a strain of Aspergillus niger, using L-arabitol as inducer. Poly A⁺ RNA was isolated and used to synthesize cDNA which was in turn ligated into a bacteriophage lambda vector. The cDNA expression library was screened and phages expressing a fusion protein containing a part of the α -L-arabinofuranosidase A (ABF A) protein were identified by probing the filters with anti α -L-arabinofuranosidase A antiserum and subsequent detection using an alkaline phosphatase conjugate. The clone containing the largest insert (1.3 kb), designated pC1X1, was subjected to a limited restriction analysis (Figure 12).

An Aspergillus niger genomic library was also constructed and screened for the abfA gene. Plaque hybridization, using nitrocellulose replicas, was performed and 18 hybridizing plaques, appearing in duplicate on the replica

filters, were identified: λ_{abfA1} to λ_{abfA18} . The DNA isolated from phages λ_{abfA1} , λ_{abfA5} , λ_{abfA6} and λ_{abfA12} was analyzed by restriction analysis. A partial restriction map of the genomic region of the abfA gene was
5 obtained. An 8.5 kb NsiI/XbaI fragment from phage λ_{abfA5} was ligated in the vector pGEM-7Zf(+), resulting in the plasmid pIM900. The plasmid pIM900 was further analyzed using restriction enzymes resulting in a restriction map as shown in Figure 13. The plasmid pIM900 containing the abfA gene was
10 recloned into E. coli JM109 and was deposited at the Centraal Bureau voor Schimmelcultures (Baarn, the Netherlands on March 17, 1992, under accession number CBS 187.92). The sequence of the abfA gene is provided in Figure 14 (Formula 20).

The abfA gene encodes a protein 628 amino acids in
15 length (Figure 15, Formula 21) as derived from the abfA cDNA sequence. The N-terminal amino acid sequence, as determined in Example 12.1 (Formula 12) is preceded by a 25 amino acids long hydrophobic sequence. The amino acid sequence determined from the CNBr peptide (Formula 13) is found in the sequence from
20 position 38 until position 52. The mature ABF A protein is 603 amino acids in length, and the unglycosylated protein has a deduced molecular weight of 65378 Da and a theoretical IEP of 3.7.

In the case of the ABN A enzyme, a cDNA library was made
25 after the induction of mRNA in a strain of Aspergillus niger, using L-arabitol as inducer. The strain was cultured and mycelium was harvested at regular time intervals. RNA was isolated and used to synthesize cDNA and construct an expression library. The library was amplified and screened
30 using antibodies raised against ABN A. A single positive clone was isolated. The length of the cDNA insert was determined by digestion with EcoRI/XhoI and subsequent agarose electrophoresis. The cDNA fragment was approximately 700 bp in length.

An Aspergillus niger genomic library was also constructed and screened for the abnA gene. Plaque hybridization was performed as for the ABN A enzyme and 6 hybridizing plaques were isolated. DNA was isolated from each of the phages and was used for restriction analysis which resulted in a partial restriction map. Using this restriction map, a 3.1 kb HindIII was selected for subcloning of the abnA gene into the vector pEMBL19, which resulted in the plasmid pIM950. This plasmid was further analyzed using restriction enzymes, providing the restriction map shown in Figure 18. The plasmid pIM950 containing the abnA gene was recloned into E. coli JM109 and was deposited at the Centraal Bureau voor Schimmelcultures (Baarn, the Netherlands on March 17, 1992, under accession number CBS 188.92). The sequence of the abnA gene is provided in Figure 19 (Formula 22).

The abnA gene encodes a protein 346 amino acids in length, having a molecular weight of 37184 Da (Figure 20, Formula 23) as derived from the abnA cDNA sequence. The N-terminal sequence, as determined in Example 12.3 (Formula 18) is preceded by a 19 amino acids long hydrophobic sequence. The amino acid sequence as determined from a CNBr peptide (Example 12.3, Formula 19) is found from position 106 until position 125 of the amino acid sequence. The mature protein is 327 amino acids in length and the unglycosylated protein has deduced a molecular weight of 35204 Da and a theoretical IEP of 3.6.

The availability of a DNA molecule encoding an arabinan-degrading enzyme of interest enables the construction of mutant arabinan-degrading enzymes by site-directed mutagenesis. If the tertiary structure of the arabinan-degrading enzyme is known, and its catalytic and substrate binding domains are localized, amino acids may be selected for mutagenesis (for example with the aid of computer modelling) which most likely affect catalytic and/or substrate binding

- 20 -

functions. If the tertiary structure of the protein is not available, random mutants may be either generated along with the entire coding sequence, or the tertiary structure of the protein may be predicted by comparison with similar known arabinan-degrading enzymes isolated from another microorganism.

To facilitate the insertion of the DNA fragment containing the same DNA molecule encoding the arabinan-degrading enzyme into expression constructs comprising one or more heterologous regulatory regions, the polymerase chain reaction (PCR) (PCR Technology: Principles and Applications for DNA Amplification, (1989) H.A. Ehrlich, ed., Stockton Press, New York) may be used for introduction of appropriate restriction enzyme sites in the 5' and 3' ends of the coding sequence. The choice of restriction sites depends on the nucleotide sequence of the expression vector, i.e. the presence of other restriction sites within the DNA molecule.

To obtain the enhanced expression of the DNA construct for the production of the arabinan-degrading enzyme of interest in the original (homologous) production species, or alternatively in a heterologous fungal strain, the DNA molecule encoding the enzyme of interest, including its native regulatory regions, may be introduced into the selected expression host to increase the copy number of the construct and, consequently, protein expression.

If a heterologous expression host is preferred, and a yeast or a bacterial strain is selected, an uninterrupted (intronless) DNA molecule is used for the construction of a heterologous expression construct in order to avoid the possibility that splice signals residing on the genomic fragment are not recognized by the heterologous host. This uninterrupted DNA molecule may be obtained from a cDNA library constructed from mRNA isolated from cells, induced for the synthesis of arabinan-degrading enzymes. This library may be

screened with an oligonucleotide or cDNA probe obtained as described before. Alternatively, an uninterrupted DNA molecule may be obtained by applying a polymerase chain reaction using appropriate 5' and 3' oligonucleotides on the first strand cDNA synthesized from the RNA of arabinan-induced cells.

Enhanced expression of the DNA molecule encoding the arabinan-degrading enzyme of interest may also be achieved by the selection of heterologous regulatory regions, e.g. promoter, secretion leader and terminator regions, which serve to increase expression and, if desired, secretion levels of the protein of interest from the chosen expression host and/or to provide for the inducible control of the expression of the arabinan-degrading enzyme of interest.

Aside from the promoter native to the gene encoding the arabinan-degrading enzyme of interest, other promoters may be used to direct its expression. The promoter may be selected for its efficiency in directing the expression of the arabinan-degrading enzyme of interest in the desired expression host.

In another embodiment, a constitutive promoter may be selected to direct the expression of the desired arabinan-degrading enzyme. Such an expression construct may provide additional advantages since it circumvents the need to culture the expression hosts on a medium containing arabinans as an inducing substrate.

Examples of strong constitutive and/or inducible promoters which are preferred for use in fungal expression hosts are those which are obtainable from the fungal genes for xylanase (xlnA), phytase, ATP-synthetase, subunit 9 (oliC), triose phosphate isomerase (tpi), alcohol dehydrogenase (adhA), α -amylase (amy), amyloglucosidase (AG - from the glaA gene), acetamidase (amdS) and glyceraldehyde-3-phosphate dehydrogenase (gpd) promoters.

- 22 -

Examples of strong yeast promoters are those obtainable from the genes for alcohol dehydrogenase, lactase, 3-phosphoglycerate kinase and triosephosphate isomerase promoters.

5 Examples of strong bacterial promoters are the α -amylase and Spo2 promoters as well as promoters from extracellular protease genes.

Hybrid promoters may also advantageously be used to improve inducible regulation of the expression construct.

10 Often, it is desirable for the arabinan-degrading enzyme of interest to be secreted from the expression host into the culture medium from where the arabinan-degrading enzyme may be more easily recovered.

15 According to the present invention, the arabinan-degrading enzyme of interest's native secretion leader sequence may be used to effect the secretion of the expressed arabinan-degrading enzyme.

20 However, an increase in the expression of the arabinan-degrading enzyme sometimes results in the production of the protein in levels beyond that which the expression host is capable of processing and secreting, creating a bottleneck such that the protein product accumulates within the cell. Accordingly, the present invention also provides heterologous leader sequences to provide for the most efficient secretion
25 of the arabinan-degrading enzyme from the chosen expression host.

30 According to the present invention, the secretion leader may be selected on the basis of the desired expression host. A heterologous secretion leader may be chosen which is homologous to the other regulatory regions of the expression construct. For example, the leader of the highly secreted amyloglucosidase (AG) protein may be used in combination with the amyloglucosidase (AG) promoter itself, as well as in combination with other promoters. Hybrid signal sequences may

also advantageously be used within the context of the present invention.

5 Examples of preferred heterologous secretion leader sequences are those originating from the fungal amyloglucosidase (AG) gene (glaA - both 18 and 24 amino acid versions e.g. from Aspergillus), the α -factor gene (yeasts e.g. Saccharomyces and Kluyveromyces) or the α -amylase gene (Bacillus).

10 In general, terminators are not considered to be critical elements for the enhanced expression of genes. If desired, a terminator may be selected from the same genes as the promoters, or alternatively, the homologous terminator may be employed.

15 In addition to the genomic fragment mentioned above, the transforming DNA may contain a selection marker to discriminate cells which have incorporated the desired gene from the bulk of untransformed cells. This selection marker, provided with the appropriate 5' and 3' regulatory sequences, may reside on the same DNA molecule containing the desired
20 gene or be present on a separate molecule. In the latter case, a co-transformation must be performed. The ratio of the expression vector/selection vector must be adjusted in such a manner that a high percentage of the selected transformants also have incorporated the vector containing the expression
25 construct of the arabinan-degrading enzyme of interest.

The most suitable selection systems for industrial micro-organisms are those formed by the group of selection markers which do not require a mutation in the host organism. Examples of fungal selection markers are the genes for
30 acetamidase (amdS), ATP synthetase, subunit 9 (oliC), orotidine-5'-phosphate-decarboxylase (pyrA), phleomycin (Durand et al. (1991) Curr. Genet., 19, 149) and benomyl resistance (benA). Exemplary of non-fungal selection markers are the bacterial G418 resistance gene (this may also be used

- 24 -

in yeast, but not in fungi), the ampicillin resistance gene (E. coli), the neomycin resistance gene (Bacillus) and the E. coli uidA gene, coding for β -glucuronidase (GUS).

5 Once the desired expression construct has been assembled, it is transformed into a suitable cloning host such as E. coli to propagate the construct. Afterwards, the expression construct is introduced into a suitable expression host wherein the expression construct is preferably integrated into the genome. Certain hosts such as Bacillus species may be
10 used as both cloning and expression hosts, thus avoiding an extra transformation step.

According to the present invention, a variety of organisms may be used as hosts for the production of the arabinan-degrading enzyme of interest. If more than one
15 arabinan-degrading enzyme of interest is to be produced, multiple vectors, each containing an expression construct for an arabinan-degrading enzyme of interest (e.g. ABF A, ABF B and/or ABN A) may be introduced into the same expression host. Alternatively, each desired arabinan-degrading enzyme may be
20 produced independently in separate expression hosts.

In one embodiment of the present invention, a homologous expression host may be used. This involves the introduction of the expression construct back into the strain from which the arabinan-degrading enzyme encoding DNA molecule was isolated
25 either in increased gene copy numbers, or under the control of heterologous regulatory regions as described above, or both.

In another embodiment, one or more arabinan-degrading enzymes of interest may be produced by introducing and expressing one or more DNA molecule(s) encoding the arabinan-degrading enzyme(s) of interest, each under the control of
30 appropriate regulatory regions, into heterologous hosts such as bacteria, yeasts or fungi. For that purpose, a DNA molecule encoding an arabinan-degrading enzyme of interest is preferably expressed under the control of promoter and

terminator sequences originating from the heterologous host. In addition, it may be necessary to replace the native secretion leader sequence of the gene of the arabinan-degrading enzyme of interest with a leader sequence homologous to the expression host in order to achieve the most efficient expression and secretion of the product.

Factors such as the size (molecular weight), the need for proper glycosylation or the desirability of the extracellular secretion of an arabinan-degrading enzyme of interest play an important role in the selection of the expression host.

The gram-negative bacterium E. coli is widely used as a host for heterologous gene expression. However, large amounts of heterologous protein tend to accumulate inside the cell. Subsequent purification of the desired protein from the bulk of E. coli intracellular proteins can sometimes be difficult.

In contrast to E. coli, bacteria from the genus Bacillus are very suitable as heterologous hosts because of their capability to secrete proteins into the culture medium. Other bacteria suitable as hosts are those from the genera Streptomyces and Pseudomonas.

Depending on the nature of the DNA molecule encoding the arabinan-degrading enzyme of interest, and/or the desirability for further processing of the expressed protein, eukaryotic hosts such as yeasts or fungi may be preferred. In general, yeast cells are preferred over fungal cells because they are easier to manipulate. However, some proteins are either poorly secreted from the yeast cell, or in some cases are not processed properly (e.g. hyperglycosylation in yeast). In these instances, a fungal host organism should be selected.

A heterologous host may also be chosen wherein the arabinan-degrading enzyme of interest is produced in a form which is substantially free from other polysaccharide-degrading enzymes. This may be achieved by choosing a host

- 26 -

which does not normally produce such enzymes such as Kluyveromyces lactis.

Examples of preferred expression hosts within the scope of the present invention are fungi such as Aspergillus species (described in European Patent Applications 184.438 and 284.603) and Trichoderma species; bacteria such as Bacillus species (described in European Patent Applications 134.048 and 253.455), Streptomyces species and Pseudomonas species; and yeasts such as Kluyveromyces species (described in European Patent Applications 96.430 and 301.670) and Saccharomyces species.

Particularly preferred expression hosts may be selected from Aspergillus niger, Aspergillus niger var. tubigenis, Aspergillus niger var. awamori, Aspergillus aculeatis, Aspergillus nidulans, Aspergillus oryzae, Trichoderma reesei, Bacillus subtilis, Bacillus licheniformis, Bacillus amyloliquefaciens, Kluyveromyces lactis and Saccharomyces cerevisiae.

According to the present invention, the production of arabinan-degrading enzyme(s) of interest is effected by the culturing of microbial expression hosts which have been transformed with one or more DNA constructs of the present invention, in a conventional nutrient fermentation medium.

The fermentation medium consists of an ordinary culture medium containing a carbon source (e.g. glucose, maltose, molasses, etc.), a nitrogen source (e.g. ammonium sulphate, ammonium nitrate, ammonium chloride, etc.), an organic nitrogen source (e.g. yeast extract, malt extract, peptone, etc.) and inorganic nutrient sources (e.g. phosphate, magnesium, potassium, zinc, iron, etc.). Optionally, an inducer (e.g. sugar beet arabinan) may be included.

The selection of the appropriate medium may be based on the choice of expression hosts and/or based on the regulatory requirements of the expression construct. Such media are well-

known to those skilled in the art. The medium may, if desired, contain additional components favoring the transformed expression hosts over other potentially contaminating microorganisms.

5 The fermentation is performed over a period of 0.5-20 days in a batch or fed-batch process at a temperature in the range of between 0 and 45°C and a pH between 2 and 10. Preferred fermentation conditions are a temperature in the range of between 20 and 37°C and a pH between 3 and 9. The
10 appropriate conditions are selected based on the choice of the expression host.

 After fermentation, the cells are removed from the fermentation broth by means of centrifugation or filtration. After removal of the cells, The arabinan-degrading enzyme of
15 interest may then be recovered and, if desired, purified and isolated by conventional means.

 The product is stably formulated either in liquid or dry form. For certain applications, immobilization of the enzyme on a solid matrix may be preferred.

20 Arabinan-degrading enzymes of interest, produced by means of the present invention, may be applied either singly, or as a mixture of arabinan-degrading enzyme activities (i.e. combinations of ABF A and/or ABF B and/or ABN A activities), and optionally together with other selected enzymes in a
25 variety of processes requiring the action of arabinan-degrading enzymes. Moreover, the fungal arabinan-degrading enzymes of the present invention, which generally have lower pH optima than arabinan-degrading enzymes of bacterial origin, are particularly well suited for use in industrial processes
30 which are performed at low pH.

 It has been found that, owing to the fact that the ABF A enzyme acts only on low oligomeric arabinosides, the ABF B enzyme preferably hydrolyzes the side chain L-arabinose residues (with some endo activity) and the ABN A acts

- 28 -

optimally on linear arabinan, mixtures of the ABF A and/or ABF B enzyme act synergistically when combined with endo arabinanase (ABN A) activity. The present invention allows the skilled artisan to produce mixtures of the desired arabinan-degrading enzyme activities having optimal proportions of the
5 desired activities for specific industrial applications.

In accordance with the present invention, it has been found that the arabinan-degrading enzymes produced via the present invention, may be used in the production of
10 concentrated juices, especially fruit (e.g. apple, pear and the like) or vegetable juices to eliminate "arabinan haze".

The incorporation of an amount of ABF B, for example, to an enzyme preparation (such as that obtained from an Aspergillus species) which in turn is added to the pulp prior
15 to pressing provides an increased yield of the resultant juice without the presence of the undesirable arabinan haze in either concentrated or non-concentrated forms of the juice. A mixture of all three arabinan-degrading enzymes produced according to the present invention, may be supplemented in the
20 processing of concentrated juices from fruits having higher amounts arabinans (e.g. pear juice). Additional enzymatic activity such as pectinase activity may also be added for optimal results as may be determined by one skilled in the art.

Moreover, the incorporation of arabinan-degrading
25 enzymes produced according to the present invention in fruit and vegetable juices improve the filterability of the juices such as grape juice. In order to breakdown the arabinogalactans which are known to cause viscosity, the juice
30 is first treated with an enzyme mixture having high amounts of arabinanases and arabinofuranosidases produced according to the present invention, followed by treatment with β -1-3- and β -1-6- galactanases.

Alternatively, α -L-arabinofuranosidases, particularly the ABF B enzyme, may be used to assist in the liberation of aroma compounds from substrates such as juices or wines, as described by Gunata et al. ((1989) and (1990), *supra*). This is achieved in a two step process wherein the first step comprises the use of an α -L-arabinofuranosidase, preferably having ABF B-like activity, to catalyze the release of arabinose residues from monoterpenyl α -L-arabinofuranosyl glucosides contained in the fruit or vegetable juice via the cleavage of the (1 $\rightarrow$ 6) linkage between a terminal arabinofuranosyl unit and the intermediate glucose of a monoterpenyl α -L-arabinofuranosylglucoside. The α -L-arabinofuranosidase preferably being in purified form to avoid the undesirable degradation of other components of the juice which may be detrimental to the ultimate quality of the juice. The resulting desarabinosylated monoterpenyl glucoside is then treated with a β -glucosidase to yield the free terpenol. If desired, both reaction steps may be performed in the same reaction vessel without the need to isolate the intermediate product (Gunata et al. (1989), *supra*).

The liberation of these aroma compounds improves the flavor of the juice or wine. Moreover, in the case of wine, the control of the liberation of aroma compounds provides wines with a more consistent flavor, thus reducing or eliminating the undesirable effect of "poor vintage years".

Arabinan-degrading enzymes may also be added to animal feeds which are rich in arabinans. When added to feeds (including silage) for monogastric animals (e.g. poultry or swine) which contain cereals such as barley, wheat, maize, rye or oats or cereal by-products such as wheat bran or maize bran, the enzyme significantly improves the break-down of plant cell walls which leads to better utilization of the plant nutrients by the animal. As a consequence, growth rate and/or feed conversion are improved. Moreover, arabinan-

- 30 -

degrading enzymes may be used to the reduce the viscosity of feeds containing arabinans.

An arabinan-degrading enzyme may be added beforehand to the feed or silage if pre-soaking or wet diets are preferred. More advantageously, the arabinan-degrading enzymes produced via the present invention continue to hydrolyze arabinans in the feed in vivo. Fungal arabinan-degrading enzymes, which generally have lower pH optima, are capable of releasing important nutrients in such acidic environments as the stomach of the animal ingesting such arabinan-degrading enzyme-supplemented feed.

Another application for the arabinan-degrading enzymes produced according to the present invention is in the pulp and paper industry. The application of xylanases is often reported to be beneficial in the removal of lignins and terpenoids from the cellulose and hemicellulose residues of a hemicellulose backbone, an essential step in the processing of wood, wood pulp or wood derivative product for the production of paper. The addition of arabinan-degrading enzymes, produced according to the present invention, to the xylanase treatment step assists in the degradation of an arabinan-containing hemicellulose backbone and thus facilitates an improved, more efficient removal of both lignins and terpenoids. The application of arabinan-degrading enzymes is particularly advantageous in the processing of soft woods in which the hemicellulose backbone contains arabinoglucuronic xylans.

In addition, the arabinan-degrading enzymes produced via the present invention may be used in other processes such as to increase yield in the enzymatic hydrolysis of sugar beet pulp, the resulting hydrolyzed fraction being capable of use in microorganism culture medium; and the hydrolysis of sycamore or gum arabic or agricultural residues such wheat-straw.

The following examples are provided so as to give those of ordinary skill in the art a complete disclosure and description of how to make and use the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, pH, etc.) but some experimental errors and deviation should be accounted for. Unless indicated otherwise, temperature is in degrees Celsius and pressure is at or near atmospheric.

Example 1

Enzyme Assays

Protein content, used for the determination of specific enzymatic activity, was measured according to the method of Lowry et al. ((1951) J. Biol. Chem 193, 265), except where otherwise denoted.

Arabinofuranosidase activity may be determined by the addition of 0.250 ml enzyme solution to 8 mM p-nitrophenyl α -L-arabinofuranoside (PNA - in 0.1 M sodium acetate buffer; pH 4.4). After incubation for 10 min at 40°C, the reaction was stopped by the addition of 2 ml of 0.5 M glycine-sodium buffer, pH 9.0. The concentration of p-nitrophenol (pNP) was determined by reading the absorbance at 400 nm. 1 nkat of activity corresponds to the liberation of 1 nmol of pNP per second under the test conditions. Alternatively, arabinofuranosidase activity may be expressed in Units per milliliter (U/ml) wherein a unit of arabinofuranosidase activity is defined as the amount of enzyme which releases 1 μ mol pNP per minute from PNA (see van der Veen et al. (1991) *supra*).

β -D-glucopyranosidase and α -L-rhamnopyranosidase activities were measured under the same conditions, according to their action on p-nitrophenyl β -D-glucopyranoside and p-nitrophenyl α -L-rhamnopyranoside substrates, respectively.

- 32 -

Endo 1,5- α -L-arabinanase activity of a culture filtrate was measured and calculated using the ArabinaZymeTM test kit as produced by Megazyme Pty. Ltd. (North Rocks, New South Wales, Australia), according to the manufacturer's instructions. One unit of activity is defined as the amount of enzyme required to release 1 μ mol of arabinose reducing sugar equivalents from carboxymethyl linear (1 $\rightarrow$ 5)- α -L-arabinan per minute under the assay conditions.

10

Example 2

Purification of *Aspergillus niger* α -L-arabinofuranosidase B (ABF B)

15

A schematic diagram of the procedure which was applied for the purification of the α -L-arabinofuranosidase (ABF B) enzyme from a culture filtrate obtained from the *A. niger* HEM strain is shown in Figure 1. Five steps were used for purification: 1) fractionated precipitation with ammonium sulphate; 2) size exclusion; 3) ion exchange chromatography; 4) gel permeation; and 5) adsorption chromatography.

20

Eluted fractions were analyzed for arabinan-degrading, rhamnosidase, glucopyranosidase activities via the colorimetric method as described in Example 1. Protein content was measured with a Bradford microtest on a titration microplate (Margaret *et al.* (1985) Anal. Biochem. 147, 144). Chromatography experiments were carried out at 5°C.

25

Example 2.1: Fractionated precipitation with (NH₄)₂SO₄

30

The sample of a culture filtrate obtained from *A. niger* strain HEM containing the ABF B enzyme, inter alia, was adjusted to pH 5.5 (the isoelectric point of α -L-arabinofuranosidase), with dilute acetic acid and recovered by fractionated precipitation with (NH₄)₂SO₄ at concentrations of 50, 60, 70, 80 and 90% of saturation at a temperature of 5°C. The precipitate was solubilized in sodium acetate buffer (pH

- 33 -

4.5; 0.05 M) and diafiltered in the same buffer in order to remove salts. Characteristics of the different fractions are summarized in Table 1.

Table 1
Enzymatic activities of the precipitates

5

Enzymatic Activity (<u>nkat/ml</u>)	% (NH ₄) ₂ SO ₄		
	<u>50</u>	<u>60</u>	<u>70</u>
10 α-L-arabinofuranosidase	1160	1100	260
15 β-D-glucopyranosidase	1300	1700	230
α-L-rhamnopyranosidase	131	140	63
		80	90
		43	12
		37	12
		62	2.7

Example 2.2: Size exclusion on BIO-GEL P10

A BIO-GEL P 10 column (BIO-RAD; Richmond, Va., USA) was pre-equilibrated with sodium acetate buffer (pH 4.5; 0.05 M) at a flow rate of 60 ml/hour. A sample of 100 ml of the 50% $\text{NH}_4)_2\text{SO}_4$ precipitate was applied to the column. Proteins were eluted under the same conditions. Fractions of 20 ml were collected and assayed for their α -L-arabinofuranosidase activity. Fractions 14 to 22 were pooled and concentrated by ultrafiltration (cut-off = 10,000 x D) at the start volume.

Example 2.3: Anion exchange chromatography on DEAE ART LS

A DEAE ART LS column (5,2 x 36 cm; IBF Biotechnics, France) was pre-equilibrated in 0.05 M; pH 4.5 sodium acetate buffer at a flow rate of 100 ml/hour. A 100 ml volume of the pool of α -L-arabinofuranosidase obtained from Example 2.2 (above) was applied to the column. The protein was eluted using a gradient of sodium chloride from 0 to 0.5 M in the start buffer. Fractions of 15 ml were collected and assayed for enzyme activities.

An α -L-rhamnosidase peak appeared near the void volume of the column, showing that this enzyme was in the neutral or cationic form at this pH (pool CY183 III-1). β -D-glucopyranosidase and α -L-arabinofuranosidase each eluted separately with good resolution using a sodium chloride gradient.

Fractions 164 to 166 were pooled (pool CY183 III-3); ultrafiltered and stabilized on glycerol 50/50 w/w. The characteristics of the different pools are summarized in Table 2.

Table 2

Characteristics of the glycosidases purified
by DEAE TRISACRYL ART LS

	Product	Enzymatic activity (nkat/ml)		
		α -L-A*	α -L-R**	β -D-G***
5				
10	Rhamnopyranosidase (<u>A. niger</u>)	34.5	231	15.0
	Glucopyranosidase (<u>A. niger</u>)	0	0	243
	Arabinofuranosidase (<u>A. niger</u>)	4240	0	14.8
15				
	* α -L-arabinofuranosidase			
	** α -L-rhamnopyranosidase			
	*** β -D-glucopyranosidase			

20 Example 2.4: Gel permeation

A BIO-GEL P60 column (2.8 x 30 cm; BIO-RAD; Richmond, Va., USA) was pre-equilibrated with 0.01 M sodium phosphate buffer (pH 8.8), at a flow rate of 30 ml/hour. A sample of 5 ml α -L-arabinofuranosidase stabilized on glycerol 1/1 w/w obtained from DEAE ART LS chromatography (Example 2.3, above) was applied to the column. The eluted protein was detected with a UV detector at an absorbance of 280 nm.

Fractions of 5 ml were collected and analyzed for their activity on p-nitrophenyl α -L-arabinofuranoside. Proteins were eluted in a single sharp peak which contained the α -L-arabinofuranosidase activity.

30 Example 2.5: Adsorption chromatography on BIO-GEL HTP

A BIO-GEL HTP column (2.8 x 20 cm; BIO-RAD; Richmond, Va., USA) was pre-equilibrated with 0.01 M sodium phosphate buffer (pH 8.8), at a flow rate of 30 ml/hour. The pool of α -L-arabinofuranosidase from BIO-GEL P60 (35 ml) (obtained from Example 2.4, above) was applied to the pre-equilibrated BIO-GEL HTP column and chromatographed at a flow rate of 30 ml/hour. The proteins were eluted with an appropriate sodium phosphate buffer gradient (pH 8.8) from 0.01 to 0.2 M.

- 37 -

Fractions of 5 ml were collected and analyzed for their α -L-arabinofuranosidase, β -D-glucopyranosidase and α -L-rhamnopyranosidase activities. The two latter enzymes were not detected in the standard conditions. The ABF B enzyme was eluted with the first major peak.

Example 3

Characterization of the purified *Aspergillus niger*

α -L-arabinofuranosidase B (ABF B) enzyme

10 Example 3.1: High performance size exclusion chromatography (HPSEC) on TSK G3000 SW

To determine the homogeneity of the enzyme and to estimate the molecular weight, fractions 63, 64, 65, 66, 67 (as obtained from Example 2, above) were chromatographed on a TSK G3000 SW column (LKB, Produkter AB; Bromma, Sweden) under the following conditions. 100 μ l aliquots of each fraction were injected on the TSK G3000 SW column, which was pre-equilibrated with sodium-potassium phosphate buffer (pH 7.00; 0.1 M) at a flow rate of 0.5 ml/min. The eluted proteins were detected spectro-photometrically at an absorbance of 280 nm. Fractions of 0.5 ml were collected and the α -L-arabinofuranosidase activity was tested qualitatively with the colorimetric test on a microtitre plate.

25 The column was calibrated with standard molecular weight proteins (Pharmacia Gel filtration calibration kit; Pharmacia AB, Uppsala, Sweden). The chromatographic profile reveals the high homogeneity of the protein eluted in fraction 63 to 66. The molecular weight of the α -L-arabinofuranosidase B enzyme was determined to be approximately 70 kDa which agrees fairly closely to the molecular weight for the same enzyme as reported by van der Veen et al. ((1991) *supra*).

Example 3.2: SDS-PAGE Electrophoresis

Fractions 62 to 66 (obtained from Example 2, above) were pooled and concentrated by ultrafiltration (cut-off = 10,000 x D). The characteristics of this sample were determined and are summarized below:

- Protein concentration : 0.25 mg/ml
- Activity on pNP-arabinofuranoside: 38 nkat/ml
- Specific activity : 153 nkat/mg

SDS PAGE electrophoresis confirms the high purity of the enzyme and the molecular weight as determined by HPSEC.

Example 4

Amino acid sequence determination of the N-termini of the
mature ABF B protein and two CNBr-generated
peptide fragments

Approximately 10 µg of the purified ABF B was electrophoresed on a 7.5% SDS-polyacrylamide gel and electroblotted onto Immobilon membrane (Millipore), according to the method as described by Matsudaira ((1987) J. Biol. Chem., 262, 10035). The appropriate band was cut out and sent in for sequence determination (Eurosequence, Groningen). The following N-terminal sequence was obtained:
Gly-Pro-Xaa-Asp-Ile-Tyr-Glu-Ala-Gly-Asp-Thr-Pro-Xaa-

Val-Ala-Ala (Formula 1)

Approximately 100 µg of pure ABF B was digested overnight at room temperature in a volume of 150 µl 0.15 CNBr in 70% formic acid. Digestion products were separated on a 15% SDS-polyacrylamide gel and electroblotted onto Immobilon (Millipore; Matsudaira, supra). Two bands, of 29 and 15 kDa, were cut out and sequenced (Eurosequence, Groningen, the Netherlands). The following sequences were obtained from respectively the 29 (Formula 2) and 15 kDa fragment (Formula 3):

Gly-Pro-Xaa-Asp-Ile-Tyr-Glu-Ala-Gly-Asp-Thr-Pro-Xaa-

Val-Ala-Ala

(Formula 2)

- 39 -

Xaa-Lys-Glu-Xaa-Ala-Ile-Ile-Leu-Gly-Ile-Gly-Gly-Asp-Xaa-
Xaa-Asn-Gly-Ala

(Formula 3)

5 The amino acid sequence as depicted in Formula 2 is identical to the N-terminal sequence of the ABF B protein (see Formula 1), the 29 kDa band represents the N-terminal fragment of the protein, while the 15 kDa fragment represents an internal fragment.

10

Example 5

Construction of an

Aspergillus niger genomic library

15 5 g of A. niger HEM mycelium was ground in a mortar in liquid nitrogen until a fine white powder was obtained. The powder was lysed in 35 ml TES (10 mM Tris·HCl, pH 7.5; 50 mM EDTA; 150 mM NaCl) + 1% SDS at 55°C for a period of 2-4 hours. The lysate was extracted once with one volume phenol and several times with one volume phenol-chloroform, until
20 the interphase was no longer visible. After a final extraction with one volume chloroform, the DNA was ethanol-precipitated. The pellet was dissolved in TE (10 mM Tris; 1 mM EDTA; pH 8.0) and RNase was added to a concentration of 1 µg/ml.

25 10 µg of this DNA was partially digested with Sau3AI in such a way that fragments were obtained with an average length of 10-20 kb. The resulting Sau3AI ends were partially filled in with Klenow polymerase using the nucleotides dGTP and dATP and ligated to LambdaGEMTM-11 XhoI half-site arms (Promega). These arms contain partially filled-in XhoI sites
30 (with dTTP and dCTP) and are purchased ready-for-use from the manufacturer. The ligated DNA was packaged using the Packagene in vitro packaging system (Promega). Titration of this primary genomic library on E. coli MB406 resulted in
35 8000 plaque-forming units (pfu).

Example 6Screening of the *A. niger* genomic library for the α -L-arabinofuranosidase B (abfB) gene and partial characterization of hybridizing phage clones

5 Details of molecular cloning techniques are described by Sambrook et al. in Molecular Cloning: A Laboratory Manual, 2nd edition (1989; Cold Spring Harbor Laboratory Press). Enzyme incubations are performed following instructions described by the manufacturer.

10 The amino acid sequences as depicted in Formulas 1 and 3 were used for the construction of oligonucleotide mixtures AB1719 and AB2306 (see Figures 2 and 3). Inosine nucleotides were used in case of a degeneracy of the genetic code of four or more nucleotides.

15 Southern blots of 10 μ g chromosomal *A. niger* HEM DNA digested with either BamHI, BglII, EcoRI or HindIII were hybridized overnight at 35°C in 6 x SSC, 5 x Denhardt's solution, 0.5% SDS and 100 μ g/ml denatured calf thymus DNA with 15 pmol of these oligo mixtures, after which the blots
20 were washed two times during 10 min at 35°C with 6 x SSC, 0.1% SDS. Oligo AB1719 (Figure 2, Formula 1a) gave a pattern of several distinct hybridizing bands per enzyme digest.

 For the screening of the *A. niger* HEM genomic library, 2 x 10³ pfu per plate were plated on three plates of 15 cm
25 diameter using *E. coli* MB406 as plating bacterium. TB (10 g/l Bacto-tryptone; 5 g/l NaCl) medium plus 0.9% and 0.75% agar were used for the bottom and top, respectively. Duplicate filters (nitrocellulose, Millipore) were prepared from each plate and screened with oligo AB1719 using
30 hybridization and washing conditions similar to those used for Southern Blot hybridizations. Five in duplo hybridizing plaques could be detected. Each plaque was removed with a Pasteur pipette and the phages were eluted from the agar plug in 0.5 ml Phage Buffer (20 mM Tris·HCl, pH 7.5; 100 mM
35 NaCl; 10 mM MgSO₄). Plaques were purified by repeated plating of phages eluted from isolated agar plugs and

subsequent screening with oligo AB1719. The five plaques were found to hybridize to a different extent with oligo AB1719:

	<u>Plaque</u>	<u>Hybridization signal</u>
5	7	+++++
	8	+++++
	12	±
	45	++
	53	+++++
10	After plaque purification, DNA was isolated from phages originating from a single hybridizing plaque. Number 12 was omitted because of its low hybridization signal. Between 30,000-50,000 plaques were plated on 15 cm plates using TB without NaCl and agarose instead of agar in bottom	
15	and top. After overnight incubation, phages were eluted from the top agar by continuous shaking of the plate on a rocker platform during 90 minutes with 15 ml Phage Buffer. Halfway, the plates were given a quarter turn. The phage suspension was collected from the plates and centrifuged to remove cell	
20	debris. DNase and RNase were added to a final concentration of 1 µg/ml and the phage suspension was incubated 45 min at 37°C. Phages were PEG-precipitated for one hour on ice in 1 M NaCl and 10% PEG 6000. After centrifugation, the phage pellet was resuspended in 1 ml Phage Buffer per plate and	
25	insoluble debris removed by centrifugation. SDS and EDTA were added to a final concentration of 0.2% and 15 mM, after which proteinase K was added to 50 µg/ml and the mixture incubated for 30 min at 65°C. After successive extractions with equal volumes of phenol, phenol/chloroform and	
30	chloroform, the DNA was precipitated from the aqueous phase with isopropanol. The DNA was dissolved in TE, containing 0.1 µg/ml RNase.	

Of each phage DNA, a partial restriction map was constructed (see Figure 4 - phage clone 7). Two of the phages clones (8 and 53) contained a fragment from the same

genomic region of A. niger, while the others (7 and 45) contained different genomic fragments.

Example 7

5 Isolation and characterization of the abfB gene

Phage DNA was digested until completion with enzymes recognizing a sequence of four successive nucleotides and the resulting fragments were ligated in pTZ18R or pUC18. The resulting colonies were transferred to nitrocellulose
10 membranes and screened with oligo AB1719 to select plasmids with a hybridizing insert. Sufficiently small inserts were sequenced to identify the insert. Sequence analysis of a 2.8 kb SacI fragment of phage clone 7 was performed using a Sequenase^R version 2.0 reagent kit (United States
15 Biochemical) according to the manufacturer's instructions. The results of the sequence analysis (Figure 5, Formula 4) demonstrate a sequence in which both the nucleotide sequence of oligo AB1719 and the amino sequence of the N-terminus of the ABF B protein could be detected (compare Formula 1 and
20 Figure 6, Formula 5).

The 2.8 kb SacI fragment was cloned into the E. coli plasmid pUC18. The resulting plasmid was given the designation pGBabfB and was deposited in E. coli strain DH5 α at the Centraal Bureau voor Schimmelcultures (Baarn, the
25 Netherlands on March 11, 1991, under accession number CBS 156.91).

Example 8

30 Introduction of the abfB gene in Aspergillus niger N 593 by co-transformation

The phage clone 7 DNA, obtained in Example 6 is introduced into Aspergillus niger by co-transformation of
35 Aspergillus niger strain N593 using the Aspergillus niger pyrA gene as a selective marker on the plasmid pGW635 (Goosen et al. (1989) Mol. Gen. Genet. 219, 282) and the phage clone 7 DNA as the co-transforming DNA. For

co-transformation, 1:20 molar ratios of transforming DNAs were added in order to favor the frequency of co-transformation.

To assay the co-transformation frequencies, the plasmid pNOM102 (Roberts *et al.* (1989) *Curr. Genet.*, 15, 177) harboring the *E. coli uidA* gene, coding for β -glucuronidase (GUS) was used as co-transforming DNA.

PYRA⁺ transformants were subsequently transferred to X-glucuronide medium (Clontech, Palo Alto, CA, U.S.A.; Roberts *et al.* (1989) *supra*) and scored for GUS activity. Blue colonies were scored as GUS⁺ phenotype, colourless as GUS⁻ phenotype. *A. niger*, co-transformed with pNOM102 and pGW635 plasmids for transformation/selection respectively, gave co-transformation frequencies up to 85 %.

Protoplasts were prepared from mycelium by growing *Aspergillus niger* N593 on minimal medium supplemented with 50 mM glucose, 0.5 % yeast extract, 0.2 % casamino acids and 10 mM uridine for 20 hours at 30°C. The minimal medium had the following composition (per 1000 ml): 6.0 g NaNO₃, 1.5 g KH₂PO₄, 0.5 g MgSO₄·7H₂O, 0.5 g KCl, 1 ml Visniac solution (Visniac, W. & Santer, M. (1957) *Bact. Rev.*, 21, 195), carbon source as indicated, pH 6.0. The preparation of protoplasts of *A. niger* N593 and the transformation procedure was performed as described by Goosen *et al.* (1987) *Current Genet.*, 11, 499.

The resulting PYR⁺ transformants were isolated, purified and tested for arabinofuranosidase production in shake flasks (see Example 9). Since no plate assay for the co-transformed gene was available, a Polymerase Chain Reaction (PCR) amplification test developed by D. Seth (Biotech International Ltd., Australia) as a quick screen to identify those transformants harbouring the non-selected gene was utilized. This PCR amplification test was performed as follows :

- A few young strands of mycelium from PYR⁺ transformants were put in a 1.5 ml tube;

- 44 -

- 50 μ l of a 2 % SDS solution was added to this tube and heated at 95°C for 10 minutes;
- The heated sample was diluted 100 times with sterilized ultrapure water;
- 5 - 5 μ l of the diluted sample was used for the PCR amplification;
- The polymerase chain reactions were performed according to the supplier of AmpliTaq™ polymerase (Perkin Elmer Cetus). After denaturation (8 minutes at 100°C) and addition of 1 μ l AmpliTaq™ polymerase, the
10 reaction mixtures were subjected to 25 amplification cycles (each : 2' at 94°C, 2' at 55°C, 3' at 72°C) in a PHC-2 DNA-amplifier (Techne). After the last cycle, the polymerization step at 72°C was extended to 7
15 minutes to complete all strands;
- The PCR products were analyzed by agarose gel electrophoresis. The co-transformation frequencies ranged from 5-45 % for this pyrA/PCR screening.

20

Example 9

Screening of co-transformants for the expression of the abfB gene

The PYR^+ transformants obtained in Example 8 were analyzed for the expression of the abfB gene. The PYR^+
25 transformants harboring bacteriophage lambda 7 DNA are selected by the PCR amplification test.

Spores of these transformants were collected from cells grown for 5 days at 34°C on potato-dextrose agar (Difco) plates.

30 Arabinofuranosidase production was tested in shake flasks under the following conditions :

Approximately 1×10^6 spores were inoculated in 100 ml pre-culture medium containing (per liter): 30 g sucrose, 2 g NaNO_3 , 1 g K_2HPO_4 , 0.5 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.5 g KCl, 0.01 g
35 $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, .5 g yeast-extract, 10 g malt-extract and 3 g

- 45 -

carboxypolymethylene (B.F. Goodrich Company). The pH was adjusted to 6.1 with NaOH.

After growing for 48 hours at 34°C in a rotary shaker (250 rpm), 5 ml of the grown culture was inoculated in a 100 ml main-culture P medium containing (per liter) : 5 g soya meal, 4.4 g beet pulp, 3 g (NH₄)₂SO₄, 3 g glycine, 12 g (NH₄)₂HPO₄ and 0.24 g MnSO₄·H₂O. The pH was adjusted to 6.2 with H₃PO₄.

The mycelium was grown for at least a further 100 hours at 34°C and at 250 rpm. Samples were taken at selected intervals. The mycelium was removed by filtration and the culture filtrate analyzed by SDS-polyacrylamide gel electrophoresis and by the arabinofuranosidase assay as described in Example 1.

The results of the arabinofuranosidase assays are presented in Table 3.

Table 3

Arabinofuranosidase production by co-transformation of <u>A. niger</u> N593 with Lambda clone 7						
PYR ⁺ transformant* of strain N593	PCR assay ^a	α -L-arabinofuranosidase activity (nkat/ml)				
		hours after inoculation				
		24	48	72	96	
* 2	-	8	13	22	6	
* 6	-	12	30	34	22	
* 7	-	13	31	36	25	
* 9	-	17	33	30	19	
* 1	+	37	50	43	35	
* 8	+	33	52	49	37	
* 21	+	44	57	62	48	
* 34	+	46	53	59	41	

^a - indicates the absence of a PCR DNA fragment
+ indicates the presence of a PCR DNA fragment
after agarose gel electrophoresis.

From the results presented in Table 3, it may be concluded that PCR positive transformants produce only 1.5-2 times more arabinofuranosidase than the PCR negative transformants, i.e. those who received only the selective pyrA marker from the plasmid pGW635.

Example 10

Arabinofuranosidase expression in Aspergillus transformed with expression vectors containing the abfB gene fused to the promoter and signal sequences of the A. niger amyloglucosidase (glaA) gene

Example 10.1: Construction of the expression vector

All constructs were made using standard molecular biological procedures as described e.g. in Sambrook et al. (1989) Molecular Cloning: A laboratory manual, 2nd edition, Cold Spring Harbor Laboratory Press, N.Y..

To obtain overexpression of arabinofuranosidase in Aspergillus niger, an additional expression cassette has been made in which the abfB gene is under control of the A. niger amyloglucosidase (AG) promoter in combination with the 18 amino acid signal sequence of the glaA gene.

Example 10.2: Construction of pAB6-1, pAB-6-3, pAB6-4, pAB6-31 and pAGEK1

The amyloglucosidase (AG) gene (glaA) of A. niger was isolated from plasmid libraries containing 3-4 kb EcoRI fragments or 13-15 kb HindIII fragments in the E. coli vector pUC19 (Yanisch-Perron et al. (1985) Gene, 33, 103; obtainable from e.g. Pharmacia LKB Biotechnology, Sweden) using the following AG-specific oligos:

AG-1 : 5'-GACAATGGCTACACCAGCACCGCAACGGACATTGTTTGGCCC-3'

(Formula 6)

AG-2 : 5'-AAGCAGCCATTGCCCCGAAGCCGAT-3'

(Formula 7)

both based on the nucleotide sequence published for A. niger (Boel et al. (1984) EMBO J., 3, 1097-1102 ; Boel et al. (1984) Mol. Cell. Biol., 4, 2306). The oligonucleotide probes

- 47 -

were derived from the sequence surrounding intron 2: oligo AG-1 is located 3' of the intron and has a polarity identical to the AG mRNA and oligo AG-2 is found upstream of intron 2 and is chosen antiparallel to the AG mRNA.

5 From this screening, plasmid pAB6-1 was obtained which contains the glaA gene on a 14.5 kb HindIII fragment (see Figure 7).

10 From plasmid pAB6-1 several subclones were made in pUC19: pAB6-3 which contains a 1.8 kb EcoRI fragment just upstream of the glaA gene and probably harbors regulatory sequences; pAB6-4 which contains a 4.6 kb HindIII - BglII fragment comprising the promoter of the glaA gene and a portion of the 5'-end of this gene.

15 Next, plasmid pAB6-3 was partially digested with EcoRI and treated with T4 polymerase. Into this plasmid, the HindIII plus EcoRI fragment of plasmid pAB6-4 was ligated, again after treatment with T4 polymerase. The resulting construct was designated pAB6-31; this construct contains a 3.6 kb upstream fragment of the glaA gene with a destroyed EcoRI site in the middle and an unique EcoRI site close to the glaA gene.

20 From plasmid pAB6-31, a 3.4 kb KpnI/EcoRI fragment harboring the same upstream sequences of the glaA gene was subcloned into the vector pTZ18R (Pharmacia, Sweden). The resulting construct was designated pAGEK1.

Example 10.3: Construction of pAGabfB1

30 Fusion of the AG-promoter and the 18 amino acid leader sequence of the glaA gene to the abfB structural gene encoding the mature protein were performed by the Polymerase Chain Reaction (PCR) method as presented schematically in Figure 8.

As primers for the PCR-amplifications, four synthetic oligonucleotides were designed with the following sequences:

primer 2796: 5'-CTCTGCAGGAATTCAAGCTAG-3'

an AG-specific sequence around the EcoRI site
approximately 250 bp upstream from the ATG initiation
codon. (Formula 8)

5 primer 2628: 5'-GTAGATGTCACAGGGGCTGCCAACCCTGTGCAGAC-3'
reversed mature abfB $\longleftrightarrow$ reversed 18 amino
(Formula 9) acid AG-leader

primer 2629: 5'-GTCTGCACAGGGTTGGCAGGCCCTGTGACATCTAC-3'
18 amino acid AG-leader $\longleftrightarrow$ mature abfB
10 (Formula 10)

10 primer 2630: 5'-GCTTAGCCCGGGGTGCTTGGGTCAGG-3'

an abfB-specific sequence located at the SmaI site on
position 483. (Formula 11)

The PCR was performed as described by Pomp & Medrano
15 (1991) Biotechniques, 10, 58, with minor modifications (see
Example 8).

To fuse the AG sequences to the abfB coding sequences,
two separate PCR's were carried out:

- 20 - the first reaction (PCR(1) in Figure 8) using pAB6-1 as
template and oligos 2796 and 2628 as primers to amplify
a 230 bp DNA fragment containing the 3'-fragment of the
AG promoter and the 18 amino acid AG-leader sequence
flanked at the 3'-border by the nucleotides of the abfB
gene;
- 25 - the second reaction (PCR(2) in Figure 8) using pGBabfB1
as template and oligos 2629 and 2630 as primers to
amplify a 300 bp DNA fragment containing the 5'- portion
of the mature arabinofuranosidase B (abfB) gene flanked
at the 5'-border by 18 nucleotides of the
30 amyloglucosidase (AG) signal peptide.

The two DNA fragments generated by PCR were purified by
gel electrophoresis and SephaglasTM Band prep kit (Pharmacia,
Sweden).

35 These two purified DNA fragments were used as templates
in the third reaction (PCR (3) in Figure 8) using oligos 2796
and 2630 as primers to generate the AG-arabinofuranosidase B
fusion. The 530 bp DNA fragment was digested with SmaI,

- 49 -

purified by agarose gel electrophoresis and SephaglasTM Band prep kit, and ligated into the SmaI-cut pTZ18R vector.

The ligated DNA was transformed into E. coli electrocompetent Top10 cells as described by the supplier (Invitrogen corporation; San Diego, CA, U.S.A.). Selection was made on Luria Broth (LB) plates containing Xgal (5-bromo-4-chloro-3-indolyl- β -D-galactoside), IPTG (isopropylthiogalactoside) and ampicillin. Plasmid DNA from white transformants was prepared as described by Andreoli, P. (1985) Mol. Gen. Genet., 199, 372 and analyzed with the restriction enzymes EcoRI and SmaI. Those transformants containing the 530 bp fusion DNA fragment were sequenced and designated pAGabfB1.

A schematic view of the construction of plasmid pAGabfB1 is presented in Figure 8.

Example 10.4: Construction of pAGabfB2 and pAGabfB3

A schematic view of these two plasmid constructions is presented in Figure 9.

The E. coli vector pBluescript II SK(+), obtainable from Stratagene cloning systems (La Jolla, CA, U.S.A.), was digested with EcoRI and XhoI and purified by agarose gelelectrophoresis.

The 530 bp EcoRI-SmaI fragment from plasmid pAGabfB1 was isolated and the 1.8 kb SmaI-XhoI fragment from pGBabfB1 (containing the mature abfB gene) was isolated (see Example 7). These two fragments were ligated into the EcoRI-XhoI cut-pBluescript II SK(+) vector and transformed to E. coli DH5 α competent cells prepared according to a modified Hanahan procedure ((1983) J. Mol. Biol., 166, 557). Selection was made on LB plates containing Xgal, IPTG and ampicillin. Plasmid DNA from white transformants was prepared and analyzed with the restriction enzymes EcoRI and XhoI. The plasmid that harbored the desired 2.3 kb EcoRI-XhoI fragment was designated pAGabfB2.

For the construction of pAGabfB3, the remaining 3.4 kb upstream region of the AG-promoter was obtained by digestion

- 50 -

of plasmid pAGEK1 with KpnI and EcoRI and purified by agarose gel electrophoresis.

The 2.3. kb EcoRI-XhoI fragment containing the AG-arabanase gene fusion was purified from plasmid pAGabfB2.

5 These two fragments were ligated into the pBluescript II SK(+) vector (which was cut with KpnI-XhoI) and used to transform E. coli DH5 α competent cells as described above.

10 After plasmid DNA isolations and restriction enzyme analyses, the desired expression cassette pAGabfB3 was obtained (see Figure 9).

Example 10.5: Construction of pAGabfB4

15 To introduce a homologous selection marker for the transformation of Aspergillus niger strain N593 (Goosen et al. (1987) supra), the 3.8 kb XbaI fragment of plasmid pGW635 containing the pyrA gene (Goosen et al. (1989) supra), was inserted into the unique XbaI site of the pAGabfB3 expression cassette. The resulting expression/ selection plasmid was designated pAGabfB4 and is shown in Figure 10.

20

Example 10.6: Construction of pAGabfB5

25 A 3.8 kb HindIII-XbaI fragment from plasmid pFYT3 (van Gorcom, R.F. et al. (1991) European Patent Application 0 420 358 A1) comprising the amdS selection gene from Aspergillus nidulans (Corrick et al. (1987) Gene, 53, 63) was isolated and inserted into the HindIII-XbaI cut-pAGabfB3 plasmid. The resulting expression/selection plasmid was designated pAGabfB5 and is depicted in Figure 11.

30 Example 10.7: Introduction of the pAGabfB3 and pAGabfB4 expression cassettes into Aspergillus niger N593

35 The plasmid pAGabfB3 was introduced into A. niger by co-transformation of strain N593 using the pyrA gene as a selective marker on the plasmid pGW635 (Goosen et al. (1989) supra) and the expression cassette pAGabfB3 as the co-transforming DNA.

- 51 -

Transformation of strain N593 with the expression cassette pAGabfB4 was performed as described by Kusters-van Someren et al. (1991) Curr. Genet., 20, 293.

5 The resulting PYR⁺ transformants were isolated, purified and tested for arabinofuranosidase production in shake flasks using the process as described in Example 11.

As a control, transformants possessing only the vector pGW635 were tested.

10 Example 10.8: Introduction of the pAGabfB5 expression cassette in Aspergillus niger strain CBS 513.88

The plasmid pAGabfB5 was introduced in Aspergillus niger strain CBS 513.88 (deposited October 10, 1988) using
15 transformation procedures as described by Tilburn et al. (1983) Gene, 26, 205 and Kelly & Hynes (1985) EMBO J., 4, 475 with the following modifications:

- mycelium was grown on Aspergillus minimal medium (Cove (1966) Biochem. Biophys. Acta, 113, 51) supplemented
20 with 10 mM arginine and 10 mM proline for 16 hours at 30°C in a rotary shaker at 300 rpm;
- only Novozym 234 (NOVO Industri), and no helicase, was used for formation of protoplasts;
- after 90 minutes of protoplast formation, 1 volume of
25 STC buffer (1.2 M sorbitol, 10 mM Tris-HCl pH 7.5, 50 mM CaCl₂) was added to the protoplast suspension and centrifuged at 2500 g at 4° for 10 minutes in a swinging-bucket rotor. The protoplasts were washed twice and resuspended in STC-buffer at a concentration of 108
30 cells/ml;
- plasmid DNA was added in a volume of 10 µl in TE buffer (10 mM Tris-HCl pH 7.5, 0.1 mM EDTA) to 100 µl of the protoplast suspension;
- after incubation of the DNA-protoplast suspension at 0°C
35 for 15 minutes, 200 µl of PEG solution was added dropwise (25 % PEG 4000 (Merck), 10 mM Tris-HCl pH 7.5, 50 mM CaCl₂) and the incubation continued at room

temperature for 10 minutes. Subsequently, 1 ml of PEG solution (60 % PEG 4000 in 10 mM Tris-HCl pH 7.5, 50 mM CaCl_2) was added slowly, with repeated mixing of the tubes. After incubation at room temperature for 20 minutes, the suspensions were diluted with STC-buffer, mixed by inversion and centrifuged at 2000 x g at 4°C for 10 minutes. The protoplasts were resuspended gently in 200 μl STC-buffer and plated on *Aspergillus* selective minimal medium with 10 mM acetamide as the sole nitrogen source, 1 M sucrose, solidified with 0.75 % bacteriological agar n°1 (Oxoid). Growth was performed at 30°C for 6-10 days.

The resulting transformants were replica-plated to selective acetamide minimal medium 1.2 % agarose plates and incubated at 30°C for 4-7 days.

After 4-7 days, fast growing and good sporulating transformants appear, against a very low background.

The AMDS⁺ transformants were purified by streaking for single colonies on selective acetamide minimal medium 1.2 % agarose plates and tested for arabinofuranosidase production in shake flasks using the process as described in Example 11.

Example 11

Expression of the arabinofuranosidase B gene under the control of the AG promoter and 18 amino acid AG leader sequence in *Aspergillus niger*

Spores of purified transformants were collected from cells grown for 3-5 days at 34°C on potato-dextrose agar (Difco) plates.

Approximatively 1×10^8 spores were inoculated in 100 ml pre-culture medium containing (per liter): 1 g KH_2PO_3 ; 30 g saccharose (maltose); 5 g yeast-extract; 10 g casein-hydrolysate; 0.5 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ and 3 g Tween 80. The pH was adjusted to 5.5.

After growing overnight at 34°C in a rotary shaker (250 rpm), 1 ml of the growing culture was inoculated in a 100 ml main-culture containing (per liter): 1 g KH_2PO_3 ; 70 g

maltodextrin (Maldex MDO3, Amylum); 12.5 g yeast-extract; 25 g casein-hydrolysate; 2 g K_2SO_4 ; 0.5 g $MgSO_4 \cdot 7H_2O$; 0.03 g $ZnCl_2$; 0.02 g $CaCl_2$; 0.05 g $MnSO_4 \cdot 4H_2O$ and 0.3 g $FeSO_4$. The pH was adjusted to 5.6.

5 The mycelium was grown for at least a further 140 hours
at 34°C and at 250 rpm. Samples were taken at selected
intervals. The mycelium was removed by filtration and the
culture filtrate analyzed by SDS-polyacrylamide gel
electrophoresis and by the arabinofuranosidase assay as
10 described in Example 1.

The N593 transformants harboring the pAGabfB3 or the pAGabfB4 expression cassette exhibit enhanced production of the ABF B enzyme as compared to the N593 transformants harboring only the pyrA selection marker.

15 The CBS 513.88 transformants harboring the pAGabfB5
expression cassette exhibit enhanced production of the ABF B
enzyme as compared to the CBS 513.88 transformants harboring
the p18FYT3 expression cassette.

20 Example 12

Determination of amino acid sequences

Example 12.1: Amino acid sequence determination of α -L-arabinofuranosidase A (ABF A)

Approximately 1-2 nmol α -L-arabinofuranosidase A (ABF A) obtained from a culture filtrate of A. niger N400, purified as described by van der Veen et al. ((1991) *supra*), was used in gas-phase sequencing (SON facility, Leiden, NL). The following sequence was determined:

Xaa-Xaa-(Leu)-Lys-Val-Xaa-Thr-Gln-(Gly)-(Gly)

30 (Formula 12)

In addition, 2-4 nmol of the protein was cleaved by CNBr using the following methods: the protein was dialyzed against bidest for 18 hours after which the protein was freeze-dried. The powder was resuspended in 70 % formic acid at a concentration of 1 mg/ml. To this solution a 200 fold excess of CNBr, with respect to the number of expected methionine residues, was added and the protein solution was incubated in

- 54 -

the dark at room temperature for 24 hours. The reaction mixture was freeze-dried and washed twice with bidest and resuspended in sample buffer (50 Mm Tris pH 6.8; 100 Mm dithiothreitol; 2 % SDS 0.1 % bromphenol blue and 10 % glycerol). This solution was heated for 3 minutes at 100°C, after which the peptides were separated on a 15 % SDS-polyacrylamide gel, followed by blotting onto Immobilon-P membrane (Millipore) according to the method described by Matsudaira, P. ((1987) J. Biol. Chem., 262, 10035). Membrane fragments containing 2-3 nmol of the particular peptide were washed with bidest and subjected to sequence analysis, according to the method described by Amons, R. ((1987) FEBS Lett., 212, 68). The following amino acid sequence was determined:

Leu-Gln-Asn-Pro-Gly-Leu-Gln-Gly-Thr-Ala-Pro-Xaa-Leu-Thr-Ala
(Gly)
(Formula 13)

Example 12.2: Amino acid sequence determination of α -L-arabinofuranosidase B (ABF B)

A sample of the α -L-arabinofuranosidase B (ABF B) protein was obtained from a culture filtrate of A. niger N400 and purified as described by van der Veen et al. ((1991) supra). The following N-terminal amino acid sequence was determined as described in Example 12.1:

Xaa-Pro-Xaa-Asp-Ile-Tyr-Glu-Ala-Gly-Asp-Thr-Pro
(Formula 14)

In addition to this N-terminal sequence, additional sequences were obtained by sequencing CNBr peptides as described in Example 12.1:

Glu-Asn-Asn-Leu-Phe-Ser-(Gly)-Ala-Asp-Glu-(Gly)-Tyr-Asn-Ser-(Thr)-Asp-Pro-Thr
(Formula 15)

Ser-Lys-Glu-Gly-Ala-Ile-Ile-Leu-Gly-Ile-Gly-Gly-Asp-Asn-Ser-Asn-Gly-Ala-Gln-Gly
(Formula 16)

Thr-Ser-Gly-Tyr-Pro-Ser-Asp-Asp-Val-Glu-Asn-(Ser)-Val-Xaa-Gln-Ile-Val-Ala

(Formula 17)

5 The amino acid sequences of Formulas 14 - 17 are also found in the derived amino acid sequence as provided in Figure 6. This confirms that the ABF B enzyme is conserved in varying strains of Aspergillus niger.

10 Example 12.3: Amino acid sequence determination of endo 1,5- α -L-arabinanase (ABN A)

A sample of the endo 1,5- α -L-arabinanase (ABN A) protein was obtained from a culture filtrate of A. niger N400 and purified as described by van der Veen et al. ((1991) supra). The following N-terminal amino acid sequence was determined as described in Example 12.1:

15 Tyr-Ala-Asp-Pro-Gly-Ala-Xaa-Ser-Gly-Val-Xaa-Thr-Thr

(Formula 18)

20 In addition to this N-terminal sequence, additional sequences were obtained by sequencing CNBr peptides as described in Example 12.1:

Glu-Tyr-Gly-Ser-Trp-Thr-Asp-His-Gly-Ser-Thr-Gly-Ile-Ala-Ser-(Arg)-Xaa-Ala-Lys-Ile

(Formula 19)

25 Example 13

Molecular cloning and analysis of the *Aspergillus niger* α -L-arabinofuranosidase A gene (abfA)

Example 13.1: Construction of the cDNA expression library

Example 13.1.1: Induction and isolation of mRNA

30 A. niger N572 (Witteveen et al. (1989) J. Gen. Microbiol., 135, 2163) was precultivated for 24 hours on minimal medium containing 1% D-glucose as a carbon source, after which mycelium was harvested by filtration and washed with sterile saline. The mycelium was then transferred to fresh
35 medium containing 1 % (w/v) L-arabitol. After 16 hours induction, the mycelium was recollected by filtration and washed thoroughly with sterile saline. The mycelium was

- 56 -

subsequently frozen in liquid nitrogen after which it was powdered using a Microdismembrator (Braun). Total RNA was isolated from mycelial powder according to the guanidine monothiocyanate / LiCl protocol of Cathala et al. ((1983) DNA 2, 329), except that SDS was omitted from the solubilization buffer. Poly A⁺ RNA was isolated from 1 mg of total RNA by oligo(dT)-cellulose chromatography (Aviv & Leder (1972) Proc. Nat. Acad. Sci. U.S.A., 69, 1408; Sambrook et al. (1989) Molecular Cloning: a Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, N.Y.) with the following modifications: 10 mM HEPES pH 7.6 was used as a buffer and SDS was omitted from all solutions, the loading buffer was supplemented with 9 % (v/v) dimethylsulfoxide.

15 Example 13.1.2: Construction and immunochemical screening of the cDNA expression library

cDNA was synthesized from 5 µg poly A⁺RNA and ligated into bacteriophage lambda Uni-ZAP XR using the ZAPTM-cDNA synthesis kit (Stratagene) according to the manufacturers instructions. After ligation of the cDNA into Uni-ZAP XR vector-arms, the phage DNA was packaged using PackageneTM extracts (Promega). Ligation of 200 ng cDNA in 1 µg vector arms and subsequent packaging of one-fifth of the reaction mixture results in a primary library consisting of 2 x 10⁵ recombinant phages. The primary library was amplified using E. coli PLK-F', titrated and stored at 4°C.

Screening of the cDNA expression library obtained was performed essentially as described by Young and Davies ((1983) Science 222, 778). In short, 5000 plate forming units (pfu) of the amplified stock were plated on NZYCM medium (per 1000 ml: 10 g NZ amine, 5 g NaCl, 5 g yeast extract, 1 g casamino acids, 2 g MgSO₄·7H₂O, pH 7.5 - for plates, 12 g agar was added, for topagarose 7 g agarose) using E. coli BB4 (Stratagene) cells as a host in 0.6 % topagarose. Plates were incubated for 5 hours at 37°C, after which they were covered with nitrocellulose filters which were previously soaked in 10 mM IPTG and air-dried. Plates were then further incubated

for 6 hours at 37°C. Plates were cooled to 4°C and the position of the filters on the plates was marked before they were removed. The filters were incubated for 15 minutes in 0.5 M NaCl, 0.05 % Tween 20 (Biorad), 20 mM Tris-HCl pH 7.5 with gentle shaking, this was repeated once. The bacterial debris was removed by gentle scrubbing with gloved hands. Phages expressing a fusion protein containing a part of the α -L-arabinofuranosidase A (ABF A) protein were identified by probing the filters with anti α -L-arabinofuranosidase A antiserum and subsequent detection using an alkaline phosphatase conjugate, according to the procedure described for Western blots in the manufacturer's instructions (Biorad). In two experiments, 5×10^3 and 5×10^4 pfu of the amplified library were screened for expression of α -L-arabinofuranosidase A cDNA: two and nine positives were found, respectively. Upon purification and excision, the plasmids were isolated by growing the resulting colonies overnight in LB medium (per 1000 ml: 10 g trypticase peptone (BBL), 5 g yeast extract (BBL), 10 g NaCl, 0.5 mM Tris-HCl pH 7,5) containing 100 μ g/ml ampicillin. From the cultures, plasmid DNA was isolated by the alkaline lysis method as described by Maniatis *et al.* ((1982) Molecular Cloning, a Laboratory Manual, Cold Spring Harbor Laboratory, N.Y., 368-369). The lengths of the cDNA insert were determined by digestion with EcoRI and XhoI and subsequent agarose electrophoresis. The clone containing the largest insert (1.3 kb), designated pClX1, was subjected to a limited restriction analysis (Figure 12).

Example 13.2: Screening of the A. niger genomic library for the α -L-arabinofuranosidase A gene (abfA) and isolation of the gene

Example 13.2.1.: Screening of the A. niger genomic library for abfA gene

An A. niger N400 genomic library was constructed as described by Harmsen *et al.* ((1990) *supra*). To screen the library for the abfA gene, 10^4 pfu per plate were plated in

- 58 -

NZYCM topagarose containing 0.7% agarose on 85-mm-diameter NZYCM (1.2% agar) plates as described (Maniatis et al. (1982) *supra*, 64).

Plaque hybridization, using nitrocellulose replicas, was performed as follows: 6 x 10⁴ pfu were plated with E. coli BB4 cells in 0.6 % topagarose. After overnight incubation of the plates at 37°C, two replicas of each plate were made on nitrocellulose filters as described by Maniatis et al. ((1982) *supra*, 320). The filters were wetted and washed for 60 minutes at room temperature in 3 x SSC, after which they were prehybridized at 68°C for two hours in prehybridization buffer containing: 6 x SSC, 0.5 % SDS, 5 x Denhardt's solution and 100 µg/ml heat denatured herring sperm DNA (Boehringer Mannheim). The SSC solution was made from a 20 x SSC stock (per 1000 ml: 175.3 g NaCl, 107.1 g sodium citrate·5.5H₂O, pH 7.0). The 5 x Denhardt's solution was made from a 100 x stock solution (per 500 ml: 10 g Ficoll-400, 10 g polyvinyl pyrrolidone, 10 g Bovine Serum Albumin (Pentax Fraction V)). After two hours prehybridization, the prehybridization buffer was replaced by hybridization buffer which was identical to the prehybridization buffer, except that this buffer contained the ³²P-labelled 1.0 kb PstI fragment from cDNA clone pClX1 (Figure 12), isolated and labelled as described by van den Broeck et al. ((1992) European Patent Application 0 463 706 A1 - see Examples 2.1 and 7.1). The filters were hybridized for 18 hours at 68°C. After hybridization, the filters were washed twice at 68°C for 30 minutes in 2 x SSC / 0.1 % SDS, followed by two washes for a period of 30 minutes at 68°C in 0.2 x SSC / 0.1 % SDS. The air-dried filters were taped on a sheet of Whatman 3MM paper, keying marks were made with radioactive ink and the Whatman paper and filters were covered with Saran WrapTM. Hybridizing plaques were identified by exposure of Kodak XAR X-ray film for 18 hours at -70°C using an intensifying screen.

18 hybridizing plaques, appearing in duplicate on the replica filters, were identified: lambda_{abf}A1 to lambda_{abf}A18.

Each positive plaque was picked from the plate using a Pasteur pipette and the phages were eluted from the agar plug in 1 ml of SM buffer (per 1000 ml: 5.8 g NaCl, 2.0 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 50 ml 1 M Tris-HCl pH 7.5, 5 ml 20% gelatin) containing 20 μl chloroform, as described by Maniatis *et al.* ((1982) *supra*, 64). The phages obtained were purified by repeating the procedure described above using filter replicas from plates containing 50-100 plaques of the isolated phages.

After purification, the phages were propagated by plating 5×10^3 phages on NZYCM medium. After overnight incubation at 37°C, confluent plates were obtained, from which the phages were eluted by adding 5 ml SM buffer and storing the plates for 2 hours at 4°C with intermittent shaking. After collection of the supernatant using a pipette, the bacteria were removed from the solution by centrifugation at 4,000 x g for 10 minutes at 4°C. To the supernatant, 0.3 % chloroform was added and the number of pfu was determined by titration as described van den Broeck *et al.* ((1992), *supra*, see Example 2.4). These phage stocks contained approximately 10^{10} pfu/ml.

Example 13.2.2: Isolation of DNA from bacteriophage lambda

Four of the isolated phages ($\lambda_{\text{abf}}\text{A1}$, $\lambda_{\text{abf}}\text{A5}$, $\lambda_{\text{abf}}\text{A6}$ and $\lambda_{\text{abf}}\text{A12}$) were propagated by combining 5×10^9 *E. coli* LE392 bacteria (Murray, N. (1977) *Mol. Gen. Genet.*, 150, 53) in 300 μl SM buffer with 2×10^6 phages and incubating at 37°C for 15 minutes. After the incubation period, the infected bacteria were used to inoculate 100 ml prewarmed (37°C) NZYCM medium and were subsequently incubated for 9-12 hours at 37°C in a New Brunswick rotation shaker at 250 rpm, after which period the bacteria were lysed. The bacterial debris was removed by centrifugation for 10 minutes at 10 krpm at 4°C in a Sorvall High speed centrifuge. The phages were precipitated from the supernatant obtained (100 ml) by the addition of 10 g polyethylene glycol-6000 and 11.7 g NaCl and storing the solution overnight at 4°C. The

precipitated phages were collected by centrifugation at 14,000 x g at 4°C for 20 minutes. The supernatant was removed by aspiration, while the last traces of liquid were removed using a paper towel. The phages were carefully resuspended in
5 4 ml SM buffer and extracted once with an equal volume of chloroform.

Before the DNA was extracted from the phage particles, DNA and RNA originating from the lysed bacteria was removed by incubation of the phage suspension with DNase I and RNase
10 A (both 100 µg/ml) for 30 minutes at 37°C. The phage DNA was subsequently released from the phages by the addition of EDTA to a final concentration of 20 mM, while the protein was removed from the solution by extracting twice with an equal volume of a solution of phenol / chloroform / isoamyl alcohol
15 (25:24:1). After separation of the phases by centrifugation using a Sorvall centrifuge (14,000 x g, 10 minutes), the aqueous phase was extracted once with an equal volume of a solution of chloroform / isoamyl alcohol (24:1). The phases were separated by centrifugation, after which the DNA was
20 precipitated from the aqueous phase by the addition of 0.1 vol. 5 M sodium perchlorate and 0.1 vol. isopropanol, followed by incubation on ice for 30 minutes. The DNA was recovered by centrifugation for 10 minutes at 4°C (14,000 x g). The supernatant was removed by aspiration, after which
25 the DNA was resuspended in 400 µl TE buffer. The DNA was precipitated once again from this solution by the addition of 0.1 vol. 3 M sodium acetate and 2 vol. ethanol. The DNA was collected by centrifugation for 10 minutes at 4°C (14,000 x g). The supernatant was removed by aspiration, the remaining
30 pellet was briefly dried under vacuum, after which the DNA was resuspended in 125 µl TE buffer containing 0.1 µg/ml RNase A. This purification procedure results in the isolation of approximately 50-100 µg DNA from each phage.

Example 13.2.3: Restriction analysis of abfA containing phages

The DNA isolated from phages lambda_{abf}A1, lambda_{abf}A5, lambda_{abf}A6 and lambda_{abf}A12 was analyzed by Southern analysis.

5 The DNA was digested for 3 hours at 37°C in a reaction mixture composed of the following solutions: 5 µl (approximately 1 µg) DNA solution; 2 µl of the appropriate 10 x React buffer (BRL); 10 Units restriction enzyme (BRL) and sterile distilled water to give a final volume of 50 µl.
10 After digestion, the DNA was precipitated by the addition of 0.1 vol. 3 M sodium acetate and 2 vol. ethanol. The DNA was collected by centrifugation for 10 minutes at room temperature (14,000 x g). The supernatant was removed by aspiration and the remaining pellet was briefly dried under vacuum and resuspended in 20 µl sterile distilled water.
15 After the addition of 4 µl DNA loading buffer (0.25 % (w/v) bromophenol blue, 0.25 % (w/v) xylene cyanol, 15 % (w/v) Ficoll type 400 in H₂O), the samples were incubated for 10 minutes at 65°C and rapidly cooled on ice, before loading the
20 samples on a 0.6 % agarose gel in TAE buffer (50 x TAE buffer per 1000 ml: 242.0 g Trizma base (Sigma), 57.1 ml glacial acetic acid, 100 ml 0.5 M EDTA pH 8.0). The DNA fragments were separated by electrophoresis at 25 V for 15-18 hours.

After electrophoresis, the DNA was transferred and
25 denatured by alkaline vacuum blotting (VacuGene XL, Pharmacia LKB) to nylon membrane (Gene Bind 45, Pharmacia LKB) as described in the instruction manual (pp. 25-26) and subsequently prehybridized and hybridized using a ³²P-labelled 1 kb PstI fragment and hybridization conditions as
30 described in Example 12.2.1. The hybridization pattern was obtained by exposure of Kodak XAR-5 X-ray film for 18 hours at -70°C using an intensifying screen.

The restriction patterns obtained were used to derive a partial restriction map of the genomic region of the abfA gene. A 8.5 kb NsiI/XbaI fragment was selected for
35 subcloning.

- 62 -

Example 13.2.4: Subcloning of the A. niger abfA gene

From phage lamda_{abf}A5, the 8.5 kb NsiI/XbaI fragment was isolated by digesting the phage DNA and subsequent agarose electrophoresis. The fragment was cut from the agarose gel, after which it was recovered from the piece of agarose by electro-elution using ISCO cups. A dialysis membrane was mounted on both the large and the small containers of this cup, the cup was filled with 0.005 x TAE and the piece of agarose was placed in the large container of the cup. The cup was subsequently placed in the electro-elution apparatus, the large container in the cathode chamber contained TAE and the small container at the anode chamber contained TAE / 3 M sodium acetate. The fragments were electro-eluted at 100 V for a period of 2 hours. After this period, the cup was taken from the electro-elution apparatus and the buffer was removed from the large container, while the buffer was only removed from the upper part of the small container. The remaining buffer (200 µl) containing the DNA fragments was dialyzed in the cup against distilled water for a period of 30 minutes. Finally, the DNA was precipitated by the addition of 0.1 vol. 3 M sodium acetate (pH 5.6) and 2 vol. cold (-20°C) ethanol. The DNA was collected by centrifugation (Eppendorf centrifuge) for 30 minutes at 14,000 x g. and 4°C. After removal of the supernatant, the DNA pellet was dried using a Savant Speedvac vacuum centrifuge. The DNA was dissolved in 10 µl TE buffer and the concentration was determined by agarose electrophoresis, using lambda DNA with a known concentration as a reference and ethidium bromide staining to detect the DNA.

The fragment obtained was ligated in the vector pGEM-7Zf(+), which was digested with NsiI and XbaI prepared as follows: 1 µl (1 µg/µl) pGEM-7Zf(+) was mixed with 2 µl 10 x React 1 (BRL), 1 µl (10 U/µl) NsiI, 1 µl (10 U/µl) XbaI and 16 µl sterile distilled water. The DNA was digested for 1 hour at 37°C. The vector was isolated from a 0.6% agarose gel as described above.

The 8.5 kb NsiI/XbaI fragment was ligated in the vector, resulting in the plasmid pIM900, by the following procedure: 100 ng pGEM-7Zf(+) fragment was mixed with 100 ng 8.5 kb NsiI/XbaI fragment and 4 μ l 5 x ligation buffer (composition: 500 mM Tris-HCl (pH 7.6); 100 mM MgCl₂; 10 mM ATP; 10 mM dithiotreitol; 25 % PEG-6000) and 1 μ l (1.2 U/ μ l). T₄ DNA ligase (BRL) was added to this mixture to a final volume of 20 μ l. After incubation for 16 hours at 14°C, the mixture was diluted to 100 μ l with sterile water. 10 μ l of the diluted mixture was used to transform E. coli JM101 competent cells (Yanish-Perron *et al.* (1985) Gene, 33, 103), prepared by the CM1, CM2 method as described in the Pharmacia Manual for the M13 cloning/sequencing system. A selection of six of the resulting colonies were grown overnight in LB medium containing 100 μ g/ml ampicillin. Plasmid DNA was isolated from the cultures by the alkaline lysis method as described by Maniatis *et al.* ((1982) *supra*, 368), which was used in restriction analysis to select a clone harboring the desired plasmid, pIM900. Plasmid DNA was isolated on a large scale from 500 ml cultures of E. coli JM101 which contained pIM900. The cultures were grown in LB medium containing 100 μ g/ml ampicillin (Maniatis *et al.* (1982) *supra*, 86). The plasmid was purified by CsCl centrifugation, phenolyzed, ethanol precipitated and dissolved in 400 μ l TE. Approximately 500 μ g was obtained.

The plasmid pIM900 was further analyzed using restriction enzymes resulting in a restriction map as shown in Figure 13.

The plasmid pIM900 containing the abfA gene was recloned into E. coli JM109 and was deposited at the Centraal Bureau voor Schimmelcultures (Baarn, the Netherlands on March 17, 1992, under accession number CBS 187.92).

Example 13.3: The primary structure of the abfA gene

Example 13.3.1: Sequence analysis of the *A. niger* *abfA* gene

The sequence of the *A. niger* *abfA* gene, its promoter-regulation region, the structural gene and the termination region, was determined by subcloning fragments from pIM900 in M13mp18/mp19, in combination with the use of specific oligonucleotides as primers in the sequencing reactions.

For nucleotide sequence analysis, restriction fragments were isolated as described in Example 13.2.4 and were then cloned in bacteriophage M13 mp18/19 RF DNA vectors (Messing, J. (1983) *Methods in Enzymology*, 101C, 20; Norrander *et al.* (1983) *Gene*, 26, 101), which were digested with the appropriate restriction enzymes. The nucleotide sequences were determined by the dideoxynucleotide chain-termination procedure (Sanger *et al.* (1977) *Proc. Nat. Acad. Sci. U.S.A.*, 74, 5463) using the Pharmacia T7 DNA polymerase sequencing kit. Computer analysis was performed using the PC/GENE program. The nucleotide sequence of the *abfA* gene as determined is provided in Figure 14 (Formula 20). The amino acid sequence of the ABF A protein, as derived from the *abfA* cDNA sequence, is provided in Figure 15 (Formula 21).

Example 13.3.2: Characterization of the *abfA* gene

The sequence comprising the *abfA* structural gene (Figure 14, Formula 20) is preceded by a 1246 nucleotide long upstream region. No putative TATA box is found, the sequence TTAATTT at position 1156-1162, resembles most closely such a sequence. However regions which are extreme rich in CT are found in front of the translation initiation site (Gurr *et al.* (1987) In: *Gene Structure in Eukaryotic Microbes*, Vol. 22, Kinghorn, J.R. (editor), IRL Press, Oxford, 93).

The structural portion of the *abfA* gene ranges from position 1247 until position 3483 and contains seven introns. Intron A is a putative intron, based on the consensus sequences for fungal introns (Gurr *et al.* (1987) *supra*), the position of the six other introns is established by sequencing the cDNA fragment in pC1X1 and pC2X1.

The abfA gene encodes a protein 628 amino acids in length (Figure 15, Formula 21), as derived from the abfA cDNA sequence. The N-terminal amino acid sequence, as determined in Example 12.1 (Formula 12) is preceded by a 25 amino acids long hydrophobic sequence. The amino acid sequence determined from the CNBr peptide (Formula 13) is found in the sequence from position 38 until position 52. The mature ABF A protein is 603 amino acids in length, and has a deduced molecular weight of 65378 Da and a theoretical IEP of 3.7. The deduced molecular weight values for this enzyme differs from those reported in the literature as mentioned above (see Rombouts et al., supra and van der Veen et al., supra). This is most likely to be attributable to the literature evaluations being made on the glycosylated protein whereas the deduced molecular weight of the ABF A protein found in the present example is based strictly on the unglycosylated protein.

Example 14

Expression of the cloned abfA gene in

A. niger and A. nidulans

Example 14.1: Introduction of the abfA gene in A. niger N593 and A. nidulans G191 by cotransformation

The plasmid pIM900 was introduced into A. niger by cotransformation of A. niger N593 using the A. niger pyrA gene, located on the plasmid pGW613 (Goosen et al. (1987) supra), as a selective marker and the plasmid pIM900 as the cotransforming plasmid. Analogously the gene was introduced into A. nidulans G191 (Balance & Turner (1985) Gene, 36, 321) using the above-mentioned plasmids.

Protoplasts from both strains were prepared from mycelium obtained after growth on minimal medium supplemented with 0.5 % yeast extract, 0.2 % casamino acids, 50 mM glucose and 10 mM uridine for 20 hours at 30°C. The preparation of protoplasts and the transformation procedure was performed as described by Goosen et al. ((1987) supra), using 3 µg pGW613 and 50 µg pIM900.

5 The PYR⁺ transformants obtained were then analyzed for the expression of the abfA gene by SDS-PAGE followed by Coomassie Brilliant Blue R250 staining or Western blot analysis and by measuring the activity in the culture filtrate.

Example 14.2: Screening of transformants for the expression of the abfA gene

10 The transformants obtained in Example 14.1 were analyzed for the expression of the abfA gene product, the ABF A protein. Four transformants of each strain were selected and grown on minimal medium supplemented with 0.1 % yeast extract containing 1 % sugar beet pulp as a sole carbon source. After 24 hours of growth at 30°C, the mycelium was removed by
15 filtration and the arabinofuranosidase activity of culture filtrate was measured using para-nitrophenyl- α -L-arabinofuranoside (PNA) as described by van der Veen et al. ((1991) supra). In addition, the culture filtrate was analyzed by SDS-polyacrylamide gel electrophoresis (Laemmli, U.K. (1970)
20 Nature, 220, 680) using a gel containing 10 % acrylamide followed by Coomassie Brilliant Blue R250 staining (in the case of the A. nidulans transformants) and followed by Western blotting (in the case of the A. niger transformants). The ABF A protein was detected on nitrocellulose after
25 electroblotting and incubation with polyclonal antibodies raised against the ABF A protein which was purified as described in Example 12.1. The bound antibody was detected after incubation with goat-anti-mouse antibody conjugated to alkaline phosphatase, according to the Biorad instruction
30 manual.

All four transformants of both strains analyzed overproduced the ABF A protein as detected by Coomassie Brilliant Blue staining (Figure 16A) or by Western blotting (Figure 16B). Furthermore, an increased level of activity was
35 found in the culture filtrate of both strains (Table 4 and Table 5).

- 67 -

Table 4

PNA hydrolyzing activity in culture medium of Aspergillus nidulans wild type WG096 (Uitzetter, J.H.A.A. (1982) "Studies on carbon metabolism in wild type and mutants of Aspergillus nidulans", PhD Thesis, Wageningen Agricultural University, Wageningen, The Netherlands) and abfA transformants at 24 hours of growth on sugar beet pulp as a carbon source.

Strain	Activity (U/ml culture medium)
WG096	0.16
G191 :: pIM900-1	1.34
G191 :: pIM900-2	2.64
G191 :: pIM900-3	2.13
G191 :: pIM900-4	2.63

Table 5

PNA hydrolyzing activity in culture medium of Aspergillus niger wild type N402 and abfA transformants at 24 hours of growth on sugar beet pulp as a carbon source.

Strain	Activity (U/ml culture medium)
N402	0.35
N593 :: pIM900-1	2.48
N593 :: pIM900-2	1.25
N593 :: pIM900-3	1.99
N593 :: pIM900-4	1.79

Example 15

Molecular cloning of the *Aspergillus niger*
endo 1,5- α -L-arabinanase (abnA) gene

Example 15.1: Construction of the cDNA expression libraryExample 15.1.1: Induction and isolation of mRNA

A. niger N572 was precultivated for 24 hours on minimal medium containing 1 % glucose as a carbon source and supplemented with 0.1 % yeast extract and 1 mg/L nicotinamide, after which the mycelium was harvested by filtration and washed with sterile saline. The mycelium was then divided into eight equal portions and each portion was

then transferred to a 250 ml erlenmeyer flask containing 50 ml minimal medium with 1 % arabitol. These cultures were then incubated at 30°C for time intervals of 0, 1, 2, 3, 4, 5, 6 and 7 hours. For each time-point, the mycelium was harvested, washed with cold saline, frozen using liquid nitrogen and stored at -70°C. The induction patterns of endo 1,5- α -L-arabinanase A (ABN A) and ABF B were studied by Western blot analysis of 50 times concentrated culture samples of each of the time-points taken. ABF B was present in the culture medium 1 hour after transfer, while ABN A was found after 2 hours (Figure 17).

From the mycelium harvested at time-points of 1, 2, 3, 4, 5 and 6 hours of incubation, RNA was isolated as described in Example 13.1.1.

Example 15.1.2: Construction and immunochemical screening of the cDNA expression library

Half of the RNA of each time-point was pooled and the combined RNA sample was used to synthesize cDNA and construct an expression library as described in Example 13.1.2. After ligation of the cDNA into the vector and subsequent packaging of the DNA, a primary library of 3×10^4 phages was obtained. This primary library was amplified and screened using antibodies raised against ABN A (van der Veen *et al.* (1991) *supra*) as described in Example 13.1.2. Screening of 5×10^4 phages resulted in the isolation of a single positive clone. The length of the cDNA insert was determined by digestion with EcoRI/XhoI and subsequent agarose electrophoresis. The cDNA fragment was approximately 700 bp in length.

Example 15.2: Screening of the *A. niger* genomic library for the endo 1,5- α -L-arabinanase (abnA) gene and isolation of the gene

Using a 550 bp KpnI fragment (which was derived from the cDNA described in Example 15.1.2. and isolated and labelled as described in Example 13.2.1), the *A. niger* N400 genomic library was screened for the abnA gene, as described in

Example 13.2.1. Screening of 6×10^4 phages resulted in the isolation of six hybridizing phages. These phages were purified as described in Example 13.2.1, following which DNA was isolated from each of the phages as described in Example 13.2.2. The DNA obtained was used for restriction analysis as described in Example 13.2.3, which resulted in a partial restriction map. Using this restriction map, a 3.1 kb HindIII was selected for subcloning of the abnA gene into the vector pEMBL19 as described in Example 13.2.4, which resulted in the plasmid pIM950. This plasmid was further analyzed using restriction enzymes, providing the restriction map shown in Figure 18.

The plasmid pIM950 containing the abnA gene was recloned into E. coli JM109 and was deposited at the Centraal Bureau voor Schimmelcultures (Baarn, the Netherlands on March 17, 1992, under accession number CBS 188.92).

Example 15.3: Screening for cDNA clones hybridizing to 5' end of the abfB gene and the abnA gene

The cDNA library described in Example 15.1.1 and 15.1.2 was screened using ^{32}P -labelled fragments. From the abfB gene, an approximately 550 bp EcoRI/SalI fragment from plasmid pGBabfB1, was used as a probe. From the abnA gene, an approximately 300 bp SphI/BamHI fragment from plasmid pIM950, was used as a probe. The cDNA library was plated and screened with the labelled fragments as described in Example 13.2.1. Screening of 10^4 phages resulted in the isolation of 36 phages hybridizing to the abnA fragment and 8 phages hybridizing to the abfB fragment. For both genes, three phages were purified, the plasmid excised and isolated as described in Example 13.1.2. Sequencing of the 5'-termini of these cDNA fragments gave identical 5' ends for all three clones of both the abfB as well as the abnA gene. The cDNA of the abfB gene starts at position 120 (Figure 5, Formula 4) and the cDNA of the abnA gene starts at position 1163 (Figure 19, Formula 22).

Example 15.4: The primary structure of the abnA gene

Example 15.4.1: Sequence analysis of the abnA gene

5 The sequence of the A. niger abnA gene, its promoter-regulation region, the structural gene and the termination region, was determined by subcloning fragments from pIM950 into M13mp18/mp19, pGEM-7Zf(+) and pBluescript vectors, in combination with the use of specific oligonucleotides as primers in the sequencing reactions.

10 For nucleotide sequence analysis, restriction fragments were isolated as described in Example 13.2.4 and were then cloned in bacteriophage M13 mp18/19 RF DNA vectors (Messing (1983) supra; Norrander et al. (1983) supra), digested with the appropriate restriction enzymes. The nucleotide sequences were determined by the dideoxynucleotide chain-termination
15 procedure (Sanger et al. (1977) supra) using the Pharmacia T7 DNA polymerase sequencing kit. Computer analysis was performed using the PC/GENE program. The nucleotide sequence determined is provided in Figure 19 (Formula 22).

20 Example 15.4.2: Characterization of the abnA gene

The sequence comprising the abnA structural gene (Figure 19, Formula 22) is preceded by a 1177 nucleotides long upstream region, having a putative TATA box at position 1106-1113.

25 The structural portion of the abnA gene ranges from position 1178 until position 2378 and contains three introns. The position of the introns is established by sequencing the cDNA fragment obtained in Example 4.3. These introns are found at positions 1445 until 1494, 1972 until 2030 and 2117
30 until 2167.

The abnA gene encodes a protein 346 amino acids in length, having a molecular weight of 37184 Da (Figure 20, Formula 23) as derived from the abnA cDNA sequence. The N-terminal sequence, as determined in Example 12.3 (Formula 18)
35 is preceded by a 19 amino acids long hydrophobic sequence. The amino acid sequence as determined from a CNBr peptide (Example 12.3, Formula 19) is found from position 106 until

position 125 of the amino acid sequence. The mature protein is 327 amino acids in length and has a deduced molecular weight of 35204 Da and a theoretical IEP of 3.6.

5

Example 16

Expression of the cloned abnA gene in *A. niger* and *A. nidulans*

Example 16.1: Introduction of the abnA gene in *A. niger* N593 and *A. nidulans* WG191 by cotransformation

10 The plasmid pIM950 was introduced into *A. niger* by
cotransformation of *A. niger* N593 using the *A. niger* *pyrA*
gene, located on the plasmid pGW613 (Goosen *et al.* (1989)
supra), as a selective marker and the plasmid pIM950 as the
cotransforming plasmid. Analogously, the gene was introduced
15 in *A. nidulans* G191 using the above-mentioned plasmids.
Transformation of both strains was performed as described in
Example 14.1.

20

Example 16.2: Screening of transformants for the expression of the abnA gene

 The transformants obtained in Example 16.1 were analyzed
for the expression of the *abnA* gene product, the ABN A
protein. Ten *A. nidulans* transformants and fourteen *A. niger*
transformants were selected and grown on minimal medium
25 containing 1 % sugar beet pulp as a sole carbon source. After
24 hours of growth at 37°C for *A. nidulans* and at 30°C for *A.*
niger, the mycelium was removed by filtration and the endo
1,5- α -L-arabinanase activity of culture filtrate was measured
using ArabinaZyme tablets (Megazyme Pty. Ltd., North Rocks,
30 New South Wales, Australia) according to the manufacturers
instructions. The results for both strains are summarized in
Table 6 and Table 7.

Table 6

5 Endo 1,5- α -L-arabinanase activity in culture medium of Aspergillus nidulans wild type and abnA transformants at 24 hours of growth on sugar beet pulp as a carbon source.

	Strain	Activity (U/ml culture medium)
10	G191 :: pGW635	0.31
	G191 :: pIM950-1	0.83
	G191 :: pIM950-2	0.83
	G191 :: pIM950-6	0.79
	G191 :: pIM950-7	1.88
15	G191 :: pIM950-11	1.17
	G191 :: pIM950-12	0.70
	G191 :: pIM950-16	0.70
	G191 :: pIM950-17	2.34
	G191 :: pIM950-18	1.18
20	G191 :: pIM950-20	2.26

Table 7

25 Endo 1,5- α -L-arabinanase activity in culture medium of Aspergillus niger wild type and abnA transformants at 24 hours of growth on sugar beet pulp as a carbon source.

	Strain	Activity (U/ml culture medium)
	N402	0.17
	N593 :: pIM950-1	0.91
35	N593 :: pIM950-2	0.35
	N593 :: pIM950-3	0.25
	N593 :: pIM950-4	1.13
	N593 :: pIM950-5	1.23
	N593 :: pIM950-6	0.28
40	N593 :: pIM950-7	1.13
	N593 :: pIM950-8	1.31
	N593 :: pIM950-9	1.09
	N593 :: pIM950-10	0.22
	N593 :: pIM950-11	1.23
45	N593 :: pIM950-12	0.28
	N593 :: pIM950-13	0.32
	N593 :: pIM950-14	0.91

50 In addition, the culture filtrates were analyzed by Western blotting as described in Example 14.2. (Figure 21 A and B - A. nidulans transformants; Figure 22 A and B - A. niger transformants).

Example 17Characterization of the abfB gene

The sequence comprising the abfB structural gene (Figure 5, Formula 4), as found in the 2.8 kb SacI fragment (see Example 7), is preceded by a 166 nucleotide long upstream region. A putative TATA box extends from position 54 until position 60.

The structural portion of the abfB gene ranges from position 167 until position 1666 and contains no introns as is determined by sequencing the cDNA fragment obtained in Example 15.3.

The abfB gene encodes a protein 499 amino acids in length having a deduced molecular weight of 52523 Da (Figure 6, Formula 5) as derived from the abfB gene sequence and confirmed by the abfB cDNA sequence. The N-terminal amino acid sequence, as determined in Example 12.2 (Formula 14) is preceded by a 18 amino acid-long hydrophobic sequence. The amino acid sequences determined from the CNBr peptides (Formulas 15, 16 and 17) are found in the sequence from amino acid position 203 until position 219, 267 until 286 and 294 until 312, respectively. The mature ABF B protein is 481 amino acids in length, and has a deduced molecular weight of 50663 Da and a theoretical IEP of 3.8. The deduced molecular weight value for the ABF B enzyme differs from those reported in the literature as mentioned above (see Rombouts et al., supra and van der Veen et al., supra). This is most likely to be attributable to the literature evaluations being made on the glycosylated protein whereas the deduced molecular weight found in the present example is based strictly on the unglycosylated protein.

Example 18ABF B enzymatic activity on arabinoside-containing glucosidic extracts obtained from grape

A glycosidic extract was obtained by extraction of a must of Muscat de Frontignan variety grapes according to the method as described by Gunata et al. ((1989) supra).

- 74 -

The experimental protocol for the enzymatic hydrolysis on the glycosidic extract using the ABF B enzyme was as described by Gunata et al. ((1989) supra).

5 To 200 μ l of the glycosidic extract, 50 μ l (0.15 nkat, as determined according to the method described by Gunata et al. ((1989) supra) of a solution of α -arabinosidase (ABF B) was added and incubated for 16 hours at 40°C, pH 4.4. The solution was then extracted with 5 x 250 μ l pentane. The hydrolysis was monitored via thin-layer and gas
10 chromatography.

Afterwards, 50 μ l (0.15 nkat, as determined according to the method described by Gunata (1989), supra) of a β -glucosidase solution was added to the solution. The thus-obtained solution was reincubated for 16 hours at 40 °C, pH
15 4.4 and extracted with 5 x 250 μ l pentane. The hydrolysis was monitored via thin-layer and gas chromatography.

The results obtained demonstrated that the ABF B enzyme was able to hydrolyze 70% of the arabinosyl glucoside from the glycosidic extract.

20

Example 19

In-vitro digestion of sugar beet pulp with β -L-galactanase and/or α -L-arabinofuranosidase

In a model system simulating the conditions found in the
25 porcine stomach and small intestine, sugar beet pulp was incubated with respectively α -L-arabinofuranosidase (ABF B), β -L-galactanase and a mixture of both enzymes.

The sugar beet pulp was first incubated for three hours at pH=3.0 and T=39°C (porcine stomach conditions).
30 Afterwards, the pulp was incubated for an additional three hours at pH=6.5 and T=39°C (porcine small intestine conditions).

After the enzymatic incubation, the difference between the initial dry mass and the insolubles remaining after the
35 enzymatic hydrolysis (the matter was dried for 24 hours at T=103°C) was measured. This difference in dry matter was considered to be a relative measure of the in-vitro digestion

- 75 -

of the sugar beet pulp. The results of the experiments are summarized in Table 8.

Table 8

5 Percentage of digestibility in terms of dry mass after
 enzymatic incubation of sugar beet pulp at pig stomach and
 small intestine conditions

	<u>% of digestion</u>
10 Blank	15,9
α -L-arabinofuranosidase (ABF B)	15,8
β -L-galactanase	16,0
α -L-arabinofuranosidase (ABF B) plus β -L-galactanase	24,8

15 Similar results were obtained from wheat bran.

20 While the present invention has been described with
reference to specific embodiments thereof, it should be
understood by those skilled in the art that various changes
may be made and equivalents may be substituted without
parting from the true spirit and scope of the invention. In
addition, many modifications may be made to adapt a
25 particular situation, material, process, process step or
steps to the object, spirit and scope of the invention. All
such modifications are intended to be within the scope of the
claims appended hereto.

- 76 -

Claims

1. A purified and isolated DNA molecule encoding an enzyme having arabinan-degrading activity characterized in that the DNA molecule is selected from the group consisting of:
- a) a DNA molecule obtainable from a fungal strain which encodes an enzyme having arabinan-degrading activity;
 - b) genetic variants of the DNA molecule of part a);
 - c) DNA molecules capable of hybridizing to the DNA molecules of either part a) or part b), above.
2. The purified and isolated DNA molecule of claim 1, wherein the arabinan-degrading activity is selected from:
- a) the ability to cleave (1→2)- α -L-arabinosidic linkages;
 - b) the ability to cleave (1→3)- α -L-arabinosidic linkages;
 - c) the ability to cleave (1→5)- α -L-arabinosidic linkages;
 - d) the ability to cleave the 1→6 linkage between the terminal arabinofuranosyl unit and the intermediate glucosyl unit of monoterpenyl α -L-arabinofuranosyl glucosides.
3. The purified and isolated DNA molecule of claim 2, wherein the enzyme encoded by the DNA molecule has exo-type arabinan-degrading activity on (1→5)- α -L-arabinosidic linkages.
4. The purified and isolated DNA molecule of claim 3, wherein the DNA molecule is selected from the group consisting of:
- a) a DNA molecule having a nucleotide sequence as shown in Figure 14;
 - b) genetic variants of the DNA molecule of part a);

- c) DNA molecules capable of hybridizing to the DNA molecules of either part a) or part b), above;
- d) encodes an enzyme having an amino acid sequence as shown in Figure 15.

5

5. The purified and isolated DNA molecule of claim 3, wherein the DNA molecule is contained in the plasmid pIM900 (CBS 187.92).

10

6. The purified and isolated DNA molecule of claim 2, wherein the enzyme encoded by the DNA molecule has exo-type arabinan-degrading activity on (1→3)- α -L-arabinosidic linkages and (1→2)- α -L-arabinosidic linkages.

15

7. The purified and isolated DNA molecule of claim 6, wherein the DNA molecule is selected from the group consisting of:

- a) a DNA molecule having a nucleotide sequence as shown in Figure 5;
- b) genetic variants of the DNA molecule of part a);
- c) DNA molecules capable of hybridizing to the DNA molecules of either part a) or part b), above;
- d) encodes an enzyme having an amino acid sequence as shown in Figure 6.

25

8. The purified and isolated DNA molecule of claim 6, wherein the DNA molecule is contained in the plasmid pGBabfB1 (CBS 156.91).

30

9. The purified and isolated DNA molecule of claim 2, wherein the enzyme encoded by the DNA molecule has endo-type arabinan-degrading activity on (1→5)- α -L-arabinosidic linkages.

35

10. The purified and isolated DNA molecule of claim 9, wherein the DNA molecule is selected from the group consisting of:

- 78 -

- 5 a) a DNA molecule having a nucleotide sequence as shown in Figure 19;
- b) genetic variants of the DNA molecule of part a);
- c) DNA molecules capable of hybridizing to the DNA molecules of either part a) or part b), above;
- d) encodes an enzyme having an amino acid sequence as shown in Figure 20.

10 11. The purified and isolated DNA molecule of claim 9, wherein the DNA molecule is contained in the plasmid pIM950 (CBS 188.92).

15 12. A purified and isolated DNA molecule according to claim 2, wherein the DNA molecule is obtainable from a fungus selected from the genera Aspergillus, Dichotomitus, Corticium, Phytophthora and Rhodotorula.

20 13. A purified and isolated DNA molecule according to claim 12, wherein the DNA molecule is obtainable from a fungus from the genus Aspergillus.

25 14. The purified and isolated DNA molecule of claim 12, wherein the DNA molecule is obtainable from a fungal species selected from Aspergillus niger, Aspergillus niger var. tubigenis, Aspergillus niger var. awamori, Aspergillus aculeatis, Dichotomitus squalens, Corticium rolfsii, Penicillium chrysogenum and Rhodotorula flava.

30 15. The purified and isolated DNA molecule of claim 12, wherein the DNA molecule is obtainable from Aspergillus niger, Aspergillus niger var. tubigenis or Aspergillus niger var. awamori.

35 16. A DNA construct capable of directing the enhanced expression of a DNA molecule encoding an enzyme having arabinan-degrading activity, characterized in that a DNA molecule according to any one of claims 1-15, is operably

- 79 -

linked to regulatory regions capable of directing the enhanced expression of the arabinan-degrading enzyme in an appropriate host.

5 17. The DNA construct of claim 16, further characterized in that the regulatory regions include a promoter, operably linked to a DNA molecule according to claim 1, wherein the promoter is obtainable from the following genes: fungal endo α -L-arabinanase (abnA) gene; fungal α -L-arabinofuranosidase A (10 (abfA) gene; fungal α -L-arabinofuranosidase B (abfB) gene; fungal xylanase (xlnA) gene; fungal phytase gene; fungal ATP-synthetase synthetase gene; fungal subunit 9 (oliC) gene; fungal triose phosphate isomerase (tpi) gene; fungal alcohol dehydrogenase (adhA) gene; fungal α -amylase (amy) gene; 15 fungal amyloglucosidase (glaA) gene; fungal acetamidase (amdS) gene; fungal glyceraldehyde-3-phosphate dehydrogenase (gpd) gene; yeast alcohol dehydrogenase gene; yeast lactase gene; yeast 3-phosphoglycerate kinase gene; yeast triosephosphate isomerase gene; bacterial α -amylase gene; 20 bacterial Spo2 gene; and bacterial extracellular protease genes.

18. The DNA construct of claim 16, further characterized in that the regulatory regions include a secretion leader 25 sequence, wherein the secretion leader is obtainable from the following genes: fungal amyloglucosidase (glaA) gene 18 amino acid AG leader; fungal amyloglucosidase (glaA) gene 24 amino acid AG leader; yeast α -factor gene; and bacterial α -amylase gene.

30 19. A vector, characterized in that the vector contains a DNA construct according to claim 16.

20. The vector of claim 19, further characterized in that 35 the vector is a plasmid.

- 80 -

21. The vector of claim 20, wherein the plasmid is selected from pAGabfB3, pAGabfB4, pAGabfB5, pIM900-1, pIM900-2, pIM900-3, pIM900-4, pIM950-1, pIM950-2, pIM950-3, pIM950-4, pIM950-5, pIM950-6, pIM950-7, pIM950-8, pIM950-9, pIM950-10, 5 pIM950-11, pIM950-12, pIM950-13, pIM950-14, pIM950-16, pIM950-17, pIM950-18 and pIM950-20.

22. A transformed microbial host capable of the enhanced expression of an enzyme having arabinan-degrading activity, 10 characterized in that the microbial host contains a DNA construct according to claim 16.

23. A transformed microbial host according to claim 22, further characterized in that the microbial host is selected 15 from the genera consisting of Aspergillus, Kluyveromyces, Trichoderma, Saccharomyces and Bacillus.

24. A transformed microbial host according to claim 23, further characterized in that the microbial host is selected 20 from the species consisting of Aspergillus niger, Aspergillus niger var. awamori, Aspergillus niger var. tubigenensis, Aspergillus aculeatus, Aspergillus oryzae, Trichoderma reesei, Bacillus subtilis, Bacillus licheniformis, Kluyveromyces lactis and Saccharomyces cerevisiae.

25. A method for the enhanced expression of an enzyme having arabinan-degrading activity, characterized by the steps of:
a) culturing a microbial host according to claim 22 under conditions conducive to the expression of the 30 enzyme having arabinan-degrading activity; and
b) recovering the enzyme having arabinan-degrading activity.

26. A method for the enhanced expression of an enzyme having arabinan-degrading activity according to claim 25, wherein 35 the microbial host contains a DNA construct which in turn contains a DNA molecule according to claim 3.

27. A method for the enhanced expression of an enzyme having arabinan-degrading activity according to claim 25, wherein the microbial host contains a DNA construct which in turn
5 contains a DNA molecule according to claim 6.

28. A method for the enhanced expression of an enzyme having arabinan-degrading activity according to claim 25, wherein the microbial host contains a DNA construct which in turn
10 contains a DNA molecule according to claim 9.

29. An enzyme having arabinan-degrading activity, characterized in that the enzyme is produced by the method of claim 26.
15

30. An enzyme having arabinan-degrading activity, characterized in that the enzyme is produced by the method of claim 27.

20 31. An enzyme having arabinan-degrading activity, characterized in that the enzyme is produced by the method of claim 28.

25 32. Use of an enzyme having arabinan-degrading activity according to any one of claims 29 - 31 in the degradation of an arabinan-containing substrate.

30 33. A method for eliminating "arabinan haze" in a fruit or vegetable juice, characterized in that the fruit or vegetable juice is treated with an enzyme having arabinan-degrading activity according to any one of claims 29 - 31 in an amount effective in eliminating the arabinan haze.

35 34. A method for reducing the viscosity of a fruit or vegetable juice, characterized by the steps of:

- a) treating the fruit or vegetable juice with an enzyme having arabinan-degrading activity

- 82 -

according to any one of claims 29 - 31 in an amount effective in degrading arabinogalactans contained in the fruit or vegetable juice;

- 5 b) treating the desarabinosylated galactans with a galactanase enzyme in an amount sufficient to degrade the galactan polymers.

35. A method for releasing terpenyl aroma components from a fruit or vegetable juice, characterized by the steps of:

- 10 a) treating the fruit or vegetable juice with an enzyme having arabinan-degrading activity according to any one of claims 29 - 31 in an amount effective in catalyzing the release of
15 arabinose residues from monoterpenyl α -L-arabinofuranosyl glucosides contained in the fruit or vegetable juice;
- b) treating the resulting desarabinosylated monoterpenyl glucoside with a β -glucosidase enzyme in an amount sufficient to release the terpenyl
20 aroma components.

36. A method for improving the in vivo uptake and utilization of plant nutrients by an animal characterized by treating an arabinan-containing feed composition with an
25 enzyme having arabinan-degrading activity according to any one of claims 29 - 31.

37. A method for improving the removal of lignins and terpenoids from the cellulose and hemicellulose residues of
30 an arabinan-containing hemicellulose backbone of a wood, wood pulp or wood derivative product, characterized by treating the wood, wood pulp, or wood derivative product with an enzyme having arabinan-degrading activity according to any one of claims 29 - 31 in an amount effective in catalyzing
35 the degradation of the hemicellulose backbone.

Schematic diagram of the purification procedure for obtaining the α -L-arabinofuranosidase (exo B) enzyme from A. niger.




Figure 1

Amino acid sequence of the N-terminus of A. niger exo B enzyme and nucleotide sequence of the oligonucleotide mixture AB1719 derived from amino acid 4 (Asp) to 11 (Thr) of this sequence.

1 5 10 15
Gly-Pro-Xaa-Asp-Ile-Tyr-Glu-Ala-Gly-Asp-Thr-Pro-Xaa-Val-Ala-Ala

(Formula 1)

AB1719: 5' IGTATCICCTTCATAAATATC 3'

Figure 2

Amino acid sequence of the N-terminus of a 15 kDa CNBr fragment of the exo B protein and nucleotide sequence of the oligonucleotide mixture AB2306 derived from amino acid 6 (Ile) to 13 (Asp) of this sequence.

1	5	10	15
Xaa-Lys-Glu-Xaa-Ala-Ile-Ile-Leu-Gly-Ile-Gly-Gly-Asp-Xaa-Xaa-Asn-Gly-Ala			

(Formula 3)

AB2306: 5' TCICCCCAATICCIAGAATAAT 3' .
 G G G

Figure 3

4/35

Partial restriction enzyme map of the hybridizing phage clone 7. The fragment hybridizing with oligonucleotide mixture AB1719 is indicated with a bold line.




Figure 4

FIGURE 5 (page 1 of 3)

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 2753 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: *Aspergillus niger*
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 161..1660
 - (C) IDENTIFICATION METHOD: experimental
 - (D) OTHER INFORMATION: /codon_start= 161
/product= "alpha-L-arabinofuranosidase B"
/evidence= EXPERIMENTAL
/gene= "exoB"
- (ix) FEATURE:
 - (A) NAME/KEY: sig_peptide
 - (B) LOCATION: 161..214
- (ix) FEATURE:
 - (A) NAME/KEY: mat_peptide
 - (B) LOCATION: 215..1660

(Formula 4)

```

GAGCTCGATC TTGCATTTC TGCATAGGCA CTGACTGCAG TAAAGAATAT AAATAAGCCT      60
CCCTTCGCAC CCTAGTAGCA TGATTTCTTT CTCCAAGCGG CCTATCTCAC TGTCTCCTCT      120
CCTAGCCCAG AACTTACTGA GCAGGCAGTA ATCCTCCACC ATGTTCTCCC GCCGAAACCT      180
CGTCGCCCTA GGGCTGGCAG CCACCGTCAG CGCCGGCCCC TGTGACATCT ACGAAGCCGG      240
CGACACGCCC TCGTAGCCG CGCACAGCAC CACCCGCGCC CTATACAGCT CCTTCAGCGG      300
CGCCCTCTAC CAGCTCCAAC GTGGCTCCGA CGACACCACC ACCACCATCT CCCCCTCAC      360
AGCCGGCGGC GTCGCCGACG CCTCCGCCCA AGACACCTTC TGTGCCAACA CCACCTGCTT      420
GATCACCATC ATCTACGACC AATCCGGCAA CGGCAACCAC CTGACCCAAG CACCCCCGGG      480
AGGCTTCGAC GGCCCCGACG TCGACGGCTA CGACAACCTC GCCAGCGCCA TCGGCGCCCC      540
CGTCACCCTC AATGGCCAAA AGGCCTACGG CGTGTTTCATG TCCCCGGGCA CGGGCTACCG      600

```

FIGURE 5 (page 2 of 3)

TAACAACGAA GCCACCGGCA CCGCCACCGG CGACGAACCC GAAGGCATGT ACGCCGTCCT	660
GGACGGCAGC CACTACAACG ACGCCTGCTG CTTGACTAC GGCAACGCCG AGACCAGCAG	720
CACCGACACT GGC GCCGCC ACATGGAGGC CATCTACCTC GGAACAGCA CCACCTGGGG	780
CTACGGCGCC GGTGACGGGC CCTGGATCAT GGTGATATG GAGAACAACC TCTTCTCCGG	840
TGCTGATGAG GGATATAACT CCGGAGATCC CTCGATCAGC TACAGCTTTG TCACTGCCGC	900
GGTCAAGGGC GGGGCTGATA AGTGGGCTAT TCGCGGTGGT AATGCTGCCT CTGGGTCCCT	960
CTCTACTTAC TACAGCGGCG CTCGCCCGGA TTA CTCCGGC TATAACCCCA TGAGCAAGGA	1020
GGGCGCTATC ATCCTGGGTA TCGGCGGTGA CAACAGCAAC GGGCGCCAGG GTACCTTCTA	1080
CGAGGGTGTC ATGACCTCCG GCTACCCGTC GGACGATGTC GAGAACTCCG TCCAGGAGAA	1140
CATCGTGGCT GCGAAATACG TCTCCGGCTC GCTGGTCAGC GGCCCGTCGT TCACCTCCGG	1200
AGAGGTGGTC TCGCTGCGTG TCACTACCCC CGGTTACAGC ACGCGGTATA TTGCGCACAC	1260
TGACACCACT GTGAACACGC AGGTGCTGGA CGACGATAGT TCCACCACGC TGAAGGAGGA	1320
GGCTAGCTGG ACCGTGGTGA CAGGTCTGGC TAATAGTCAG TGCTTCTCGT TCGAGTCGGT	1380
TGATACCCCT GGTAGCTATA TCCGGCATT AACTTTGAG TTGCTGCTTA ATGCCAACGA	1440
TGGCACGAAG CAGTTCCATG AGGATGCTAC TTTCTGTCCT CAGGCGCCGT TGAATGGAGA	1500
AGGTA CTTCG TTGAGATCGT GGAGTTATCC GACCAGGTAT TTCAGGCATT ATGAGAATGT	1560
CCTGTATGCT GCTAGTAATG GTGGTGTGCA GACGTTTGAT TCCAAGACGT CGTTTAATAA	1620
TGATGTTAGC TTTGAGATTG AGACGGCGTT TGCTTCGTAA GGGGGGAATT GGGGGTTGTT	1680

FIGURE 5 (page 3 of 3)

TGGTGGTTTG GGTGTGGGT TGTATCTGCC TGTGTTGGCG GGGAGAGTGG TTTGTAAATA	1740
GGTCATTTTT GGTATATAGC ATACGAATAC TATACGATAG TATACATGAT AGTGGTTGTT	1800
TCATGTAATC GAGAAATTAT TCAAGGCTTG CAATACAATA TCTTATTACT GTATCTCGTG	1860
GAGCATTCAC AGACTGAACG GCTGCACAAT GATTCTTATG CGGTGATATT GACTGTGATT	1920
AGATGATATA TTCAGGTGTG CTATTTTCT TGTATTTATG CTTCATGCGA ATATATGTAT	1980
TTCACGACAG ACAGAGAGGT TAAACAAAGC ATACTAACGA GCGATTCAGA TATGGCATGT	2040
ATGAATAAGA CAAAAGTACT GCGACACTCA AGACCAACAC AAACCAGTAT ACTTCATAAC	2100
AATATATGCT AGGCAGTACC TCCGATTCTA CGATGAGCTT TACCAGAAAA TGGATTCTT	2160
TGTCCCTCGT CATTCTCAAT TTTTGGGCCT TCCACGGCTG CGGCAACCTG TTGAGCAAGA	2220
CGTGTCTCT CGTACGGATG GACGTTGATC TTCTTCTCGA GCAGGTCCAA TTGCTCTTTC	2280
TCCAAATGGC GGACTCTGAC CTCACCGGAG GAATCACTCG GCAAGCATAT CTGGTAAGTC	2340
GCTGAAAATT TCTTCTCGAA ACTGGCGTAG CTTGGTGGCC ACTGCGGGAA GCGTCGGAAG	2400
TATTTGCCTG TCCAGACGGT CAATATGAAT GAGTCGGATA CTTCAATGCC CGCCTGCCTT	2460
CTACGGTCGA AGAAATGCTG GCCGATGCTT CTCACGGAGG CATAGTTTAT AACAGGCTTG	2520
ATGGGGTTTT GTATGTCGAG GAGACCTAAT TCAAAGTGCA TAGATAGGCT CCTGAGATTG	2580
TGAAAACCGG TTGCTAGTTT GTCGAAGACG TCGITGGGCT GTTGTGTGGT TATAGTTCAG	2640
TTCATGGGAT GTACTAAGAG GATTCTTTGG CTTACCCATT GATCGTTCTC TCTCTTGACT	2700
CCCACCTGCA GTGACTTCAG TTTGGGACAG CTAACATCGA GGIGGCTGAG CTC	2753

FIGURE 6 (page 1 of 2)

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 499 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(Formula 5)

```

Met Phe Ser Arg Arg Asn Leu Val Ala Leu Gly Leu Ala Ala Thr Val
-18          -15          -10          -5

Ser Ala Gly Pro Cys Asp Ile Tyr Glu Ala Gly Asp Thr Pro Cys Val
-1  +1          5          10

Ala Ala His Ser Thr Thr Arg Ala Leu Tyr Ser Ser Phe Ser Gly Ala
15          20          25          30

Leu Tyr Gln Leu Gln Arg Gly Ser Asp Asp Thr Thr Thr Thr Ile Ser
          35          40          45

Pro Leu Thr Ala Gly Gly Val Ala Asp Ala Ser Ala Gln Asp Thr Phe
          50          55          60

Cys Ala Asn Thr Thr Cys Leu Ile Thr Ile Ile Tyr Asp Gln Ser Gly
          65          70          75

Asn Gly Asn His Leu Thr Gln Ala Pro Pro Gly Gly Phe Asp Gly Pro
      80          85          90

Asp Val Asp Gly Tyr Asp Asn Leu Ala Ser Ala Ile Gly Ala Pro Val
95          100          105          110

Thr Leu Asn Gly Gln Lys Ala Tyr Gly Val Phe Met Ser Pro Gly Thr
          115          120          125

Gly Tyr Arg Asn Asn Glu Ala Thr Gly Thr Ala Thr Gly Asp Glu Pro
          130          135          140

Glu Gly Met Tyr Ala Val Leu Asp Gly Thr His Tyr Asn Asp Ala Cys
          145          150          155

Cys Phe Asp Tyr Gly Asn Ala Glu Thr Ser Ser Thr Asp Thr Gly Ala
          160          165          170

Gly His Met Glu Ala Ile Tyr Leu Gly Asn Ser Thr Thr Trp Gly Tyr
175          180          185          190

Gly Ala Gly Asp Gly Pro Trp Ile Met Val Asp Met Glu Asn Asn Leu
          195          200          205

Phe Ser Gly Ala Asp Glu Gly Tyr Asn Ser Gly Asp Pro Ser Ile Ser
          210          215          220

```

FIGURE 6 (page 2 of 2)

Tyr	Ser	Phe	Val	Thr	Ala	Ala	Val	Lys	Gly	Gly	Ala	Asp	Lys	Trp	Ala				
		225					230					235							
Ile	Arg	Gly	Gly	Asn	Ala	Ala	Ser	Gly	Ser	Leu	Ser	Thr	Tyr	Tyr	Ser				
	240					245					250								
Gly	Ala	Arg	Pro	Asp	Tyr	Ser	Gly	Tyr	Asn	Pro	Met	Ser	Lys	Glu	Gly				
255					260					265					270				
Ala	Ile	Ile	Leu	Gly	Ile	Gly	Gly	Asp	Asn	Ser	Asn	Gly	Ala	Gln	Gly				
				275					280					285					
Thr	Phe	Tyr	Glu	Gly	Val	Met	Thr	Ser	Gly	Tyr	Pro	Ser	Asp	Asp	Val				
			290					295					300						
Glu	Asn	Ser	Val	Gln	Glu	Asn	Ile	Val	Ala	Ala	Lys	Tyr	Val	Ser	Gly				
		305					310					315							
Ser	Leu	Val	Ser	Gly	Pro	Ser	Phe	Thr	Ser	Gly	Glu	Val	Val	Ser	Leu				
	320					325					330								
Arg	Val	Thr	Thr	Pro	Gly	Tyr	Thr	Thr	Arg	Tyr	Ile	Ala	His	Thr	Asp				
335					340					345					350				
Thr	Thr	Val	Asn	Thr	Gln	Val	Val	Asp	Asp	Asp	Ser	Ser	Thr	Thr	Leu				
				355					360					365					
Lys	Glu	Glu	Ala	Ser	Trp	Thr	Val	Val	Thr	Gly	Leu	Ala	Asn	Ser	Gln				
			370					375					380						
Cys	Phe	Ser	Phe	Glu	Ser	Val	Asp	Thr	Pro	Gly	Ser	Tyr	Ile	Arg	His				
		385					390					395							
Tyr	Asn	Phe	Glu	Leu	Leu	Leu	Asn	Ala	Asn	Asp	Gly	Thr	Lys	Gln	Phe				
	400					405					410								
His	Glu	Asp	Ala	Thr	Phe	Cys	Pro	Gln	Ala	Pro	Leu	Asn	Gly	Glu	Gly				
415					420					425				430					
Thr	Ser	Leu	Arg	Ser	Trp	Ser	Tyr	Pro	Thr	Arg	Tyr	Phe	Arg	His	Tyr				
				435					440					445					
Glu	Asn	Val	Leu	Tyr	Ala	Ala	Ser	Asn	Gly	Gly	Val	Gln	Thr	Phe	Asp				
			450					455					460						
Ser	Lys	Thr	Ser	Phe	Asn	Asn	Asp	Val	Ser	Phe	Glu	Ile	Glu	Thr	Ala				
		465					470					475							
Phe	Ala	Ser																	
	480																		




Figure 7

AG/abfB GENE FUSION BY PCR




Figure 8

12/35




Figure 9




Figure 10




Figure 11

pC1X1




Figure 12




Figure 13

17/35
FIGURE 14 (page 1 of 5)

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 3958 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: *Aspergillus niger*
- (ix) FEATURE:
 - (A) NAME/KEY: exon
 - (B) LOCATION: 1247..1390
- (ix) FEATURE:
 - (A) NAME/KEY: intron
 - (B) LOCATION: 1391..1442
- (ix) FEATURE:
 - (A) NAME/KEY: exon
 - (B) LOCATION: 1443..1957
- (ix) FEATURE:
 - (A) NAME/KEY: intron
 - (B) LOCATION: 1958..2005
- (ix) FEATURE:
 - (A) NAME/KEY: exon
 - (B) LOCATION: 2006..2089
- (ix) FEATURE:
 - (A) NAME/KEY: intron
 - (B) LOCATION: 2090..2137
- (ix) FEATURE:
 - (A) NAME/KEY: exon
 - (B) LOCATION: 2138..2214
- (ix) FEATURE:
 - (A) NAME/KEY: intron
 - (B) LOCATION: 2215..2262
- (ix) FEATURE:
 - (A) NAME/KEY: exon
 - (B) LOCATION: 2263..2295
- (ix) FEATURE:
 - (A) NAME/KEY: intron
 - (B) LOCATION: 2296..2346
- (ix) FEATURE:
 - (A) NAME/KEY: exon
 - (B) LOCATION: 2347..2498
- (ix) FEATURE:
 - (A) NAME/KEY: intron
 - (B) LOCATION: 2499..2548
- (ix) FEATURE:
 - (A) NAME/KEY: exon
 - (B) LOCATION: 2549..3037
- (ix) FEATURE:
 - (A) NAME/KEY: intron
 - (B) LOCATION: 3038..3092
- (ix) FEATURE:
 - (A) NAME/KEY: exon
 - (B) LOCATION: 3093..3485

FIGURE 14 (page 2 of 5)

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: join(1247..1390, 1443..1957, 2006..2089, 2138..2214, 2263..2295, 2347..2498, 2549..3037, 3093..3485)
 (C) IDENTIFICATION METHOD: experimental
 (D) OTHER INFORMATION: /codon_start= 1247
 /product= "alpha-L-arabinofuranosidase A"
 /evidence= EXPERIMENTAL
 /gene= "exoA"

(ix) FEATURE:

(A) NAME/KEY: sig_peptide
 (B) LOCATION: 1247..1321

(ix) FEATURE:

(A) NAME/KEY: mat_peptide
 (B) LOCATION: 1322..3485

(Formula 20)

```

AAGGGACTGA CTGAACCTCAG TGAGATGTGC CGCGAAGTAA CTAAGCGCTG TGCCTGCAGT      60
GAGGCGTGAT GGAACACTGC ATCCGATCTC CAAGACCCCG ATATGAACCT TAATGACAAT      120
GGTCTAAGTT TTTTGTTCAT CTCGCATAGC ACGGTAGGCG ACTAAGTTGA AGGATAAAGA      180
GAGCAGGATT AGGATGGCTC GAGATCAAAT TGAAAGTAAT AGAGTAACAG GGGAAAACTC      240
TAACAGTTGG CTATGACGAA GGTGGCAGCG ACTTCCCGGA GAGAGATTAA GCCGGGTAGA      300
ACCAAGGGAC GGCTGTGGCT GGATCAGGAA CCCAAATCTA TTCCAGTATC GGAGGATACT      360
CAGGGGTCCC GCCGGGGTTT TCCCTTGATC GGAGAACATC AACCACATCA AAATGCGGAG      420
TAACGATGAG GGGGAAGTGC ACGGCCGACG TCAGAGTTTC GCTTGGGTGG ATCTGCAATG      480
TTTAGGCGGC CGACTAGACC GACAGTCCAG CACCTGCCCT GGGAGTTCAA GACAGTGGTT      540
TGACCATCTT TTGAGAAGAC TGAAATTCCC GTAGATTCCA TCAAGATTAT TCATGTATCA      600
CTAGCGACTC AGCACTGGGT AAGCTGCACC CACGGATTTT CTCAGCTATC TGAATGTATC      660
AGAGACAAAG ATCTTCGTGG ATGGAGGCCA TCAACCGAGC TTTAATGCGC CAAACCTACT      720
GTATGACTGA ATAATGGATG GATCGGCATG TCGTACGATC GGGTAAGGGT CCTAAAAATG      780

```

FIGURE 14 (page 3 of 5)

GAGGGCCGGG GAATTCTGGG GTTAGCCGTC GGGATTATTT CGGTACGACA AACTCACTGT 840
TCACTTTGAT CTCGGCGTGG TTGCGGCTAA AAAGATGACC TTGAAGGTGG AGCGAACAAC 900
TGGAGGTGCT CGGAGGGACG CGGAGAAGAG AGGATCCAAG ACATCGGGGC AGGGAAGTCC 960
ATGCCGCGTG CAGTCACGCT TTTCCTCCCT CTGCTCTCTT CTCCTCCACA TTCCACATTC 1020
CACCTCCACA CTCCTCCACT CTTCTCTGG CACCTTCCTC TTCCTCGATT TCTTGGCTTC 1080
TCGGTTGATT GTCTTCGACC TCGTTGCGAT GACCCACCG CATTTTCTGC TGCCTTGGCC 1140
TGTCTGATGC CCCGGTTAAT TTCACGAAAG TCCGGTCATA AAAGGCGTTG TATCCTCCCT 1200
TTGCAACAAT GACACCAGGC TCAGCTTCCT CCAGACAGCC GGCAACATGG TGGCCTTTTC 1260
AGCTCTTTCG GGCCTCAGCG CCGTTTCTTT ACTACTATCC CTCGTTCAAA ATGCACACGG 1320
AATCTCCTTG AAGGTCTCCA CCCAGGGTGG CAACTCATCC AGCCCCATCC TATATGGCTT 1380
CATGTTTGAG GTAGGCCGCA GACTAAGCAG TAAAAGACAA TTGGTTCTCT GACGGTGATT 1440
AGGATATCAA TCACTCAGGA GATGGAGGAA TTTATGGGCA AATGCTGCAG AACCCCTGGCC 1500
TTCAGGGAAC GGCCTTAAT CTGACTGCCT GGGCGGCTGT CGGTGACGCT ACCATCGCCA 1560
TTGATGGCGA CAGTCCATTG ACCTCTGCCA TCCCAGCAC CATCAAGCTT AATATTGCGG 1620
ACGATGCTAC CGGTGCGGTG GGTCTACCA ATGAGGGATA TTGGGGCATC CCAGTCGATG 1680
GCAGCGAATT CCATAGTTCC TTCTGGATAA AGGGAGA!TA CTCGGCGAC ATCACCCTCC 1740
GACTGGTTGG AACTACACC GGCACGGAGT ACGGCTCTAC CACTATCACC CATACGTCCA 1800
CAGCAGACAA CTTACCCAA GCCTCCGTCA AGTTCCCCAC CACCAAGGCT CCAGATGGCA 1860

FIGURE 14 (page 4 of 5)

ACGTTTTGTA CGAGCTCACA GTTGATGGAA GCGTGGCTGC TGGCTCGTCT TTGAACTTCG	1920
GCTACTTGAC GCTTTTTGGC GAAACCTATA AGTCAAGGTT TGCTTCCCTA TACTCCGAAG	1980
AGTGAATTAG GGGCTAATTG TGTAGGGAAA ATGGCCTGAA GCCCCAGCTT GCCAATGTGT	2040
TGGATGATAT GAAAGGATCC TTCCTGAGAT TTCCCGGCGG TAACAACCTG TAAGTCTCAG	2100
CTCGCCCGGT AAGTGATAG AAGCTCATCA GAGGTAGTGA GGGAAACAGC GCAGAAAACC	2160
GCTGGAAGTG GAACGAGACA ATCGGCGATC TTTGTGATCG TCCCGGACGT GAAGGTATGT	2220
CTCATTATTA GGATTGAAGC ATTCATCCCT GACGGTATAT AGGCACTTGG ACTTACTATA	2280
ACACCGACGG ACTGGGTAAG TAAAGGGCTA TATCAAGTA CCTAGATACT GTACTAACGC	2340
TTGTAGGCCT TCACGAATAC TTTTACTGGT GTGAGGATTT GGGGCTCGTA CCGGTGCTCG	2400
GTGTCTGGGA TGGGTTGCT CTGGAGTCGG GTGGCAACAC CCCCCTCAGG GCGGACGCAC	2460
TGACCCCTTA TATCGACGAT GTCTTGAACG AGCTCGAGGT ATGTTGAGCG GCATATCAAA	2520
TTGATAGCTG AAGCTAACCC ATTGGCAGTA CATCTTGGGT GATACGAGCA CGACCTATGG	2580
AGCGTGGCGC GCGGCAAACG GACAGGAGGA GCCGTGGAAC CTTACCATGG TCGAGATTGG	2640
CAATGAGGAC ATGCTGGGAG GCGGATGCGA GTCCTACGCG GAACGTTTCA CTGCCTTCTA	2700
TGATGCGATT CATGCTGCTT ATCCGGACCT TATCCTGATT GCCAGCACCA GCGAGGCGGA	2760
TTGCTTGCCC GAGTCAATGC CCGAGGGTAG CTGGGTCGAC TACCAGACT ACAGCACGCC	2820
TGATGGACTG GTGGGCCAGT TCACTACTT CGACAATTA AACCGCTCGG TACCATACTT	2880
CATCGGCGAG TACTCGCGCT GGGAGATTGA CTGGCCCAAC ATGAAGGGAT CGGTCGCTGA	2940

FIGURE 14 (page 5 of 5)

AGCCGTCTTC ATGATCGGGT TCGAGAGGAA CAGCGACGTG GTGAAGATGG CGGCGTATGC 3000
GCCTTTGCTC CAGCTAATCA ACTCGACTCA GTGGACGGTA AGTCACGACT GAGCAGCGGG 3060
GTTTCAGGTA TGAATGAGCT AACGAGTGGT AGCCGGACCT GATCGGATAC ACGCAGTCAC 3120
CCGGTGACAT TTTCCTGTCG ACCAGCTACT ACGTGCAGGA GATGTTCTCG CGCAACCGGG 3180
GTGATACAAT TAAGGAGGTG ACGTCGGACA GCGACTTCGG ACCGTTGTAC TGGGTTGCGT 3240
CGAGCGCCGG GGA CTCTGAC TACGTGAAGC TGGCCAATA TGGCTCCGAG ACGCAAGACC 3300
TCACGGTGAG CATCCCAGGA ACGAGCACAG GCAAGTTGAC GGTGCTGGCG GACAGTGATC 3360
CGGATGCGTA TAACTCGGAC ACCCAGACGC TGGTCACGCC GAGTGAATCG ACGGTGCAGG 3420
CGAGCAATGG CACTTTTACC TTTAGTTTGC CGGCATGGGC GGTGGCTGTC CTGGCGGCGA 3480
ACTAGCGTTG ATTGGGGCGA GCTCGTATGG GCGGCAAGTC GAATATTATC TGCAGGTGTG 3540
GTGTCATGTA CTGCAGTATT TCTGATTAAC GATAGAGAGA TAGATCCATG CTATATACCT 3600
GCTTGATAT CCAAGGCATT TATCCAATGA TAGGCAACGC CACCACACGT GGCTATGCAC 3660
ATACTAGTCA GGAGGTAGTA GAGTGACTGA CTGCTATGTA GCGGTCGGCC GGGGATAGCA 3720
GTTGAATGAC CCCAATTAAC AGAACCGCCT GACTCTCCAA CGTGACCTC TAGGTTATTT 3780
TATATTTCTT GTTAAAGCCC CCCAGAAACA CAACCGAAGA ATCAATGGAT GCGATGAAAA 3840
CAAGGCACTG GTAAGCGAAG GAGGGAAAAA CAAGCGGATT GGTGGTTGT GGCCCGATGC 3900
AAACGGGCGG CCTCTCCACT CGAATTGGAC AGAAGCGAAG GCCGTGCGCC GAGACGGG 3958

FIGURE 15 (page 1 of 3)

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 628 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein

(Formula 21)

Met	Val	Ala	Phe	Ser	Ala	Leu	Ser	Gly	Val	Ser	Ala	Val	Ser	Leu	Leu	-24	-20	-15	-10
Leu	Ser	Leu	Val	Gln	Asn	Ala	His	Gly	Ile	Ser	Leu	Lys	Val	Ser	Thr	-5	-1	+1	5
Gln	Gly	Gly	Asn	Ser	Ser	Ser	Pro	Ile	Leu	Tyr	Gly	Phe	Met	Phe	Glu	10	15	20	
Asp	Ile	Asn	His	Ser	Gly	Asp	Gly	Gly	Ile	Tyr	Gly	Gln	Met	Leu	Gln	25	30	35	40
Asn	Pro	Gly	Leu	Gln	Gly	Thr	Ala	Pro	Asn	Leu	Thr	Ala	Trp	Ala	Ala	45	50	55	
Val	Gly	Asp	Ala	Thr	Ile	Ala	Ile	Asp	Gly	Asp	Ser	Pro	Leu	Thr	Ser	60	65	70	
Ala	Ile	Pro	Ser	Thr	Ile	Lys	Leu	Asn	Ile	Ala	Asp	Asp	Ala	Thr	Gly	75	80	85	
Ala	Val	Gly	Leu	Thr	Asn	Glu	Gly	Tyr	Trp	Gly	Ile	Pro	Val	Asp	Gly	90	95	100	
Ser	Glu	Phe	His	Ser	Ser	Phe	Trp	Ile	Lys	Gly	Asp	Tyr	Ser	Gly	Asp	105	110	115	120
Ile	Thr	Val	Arg	Leu	Val	Gly	Asn	Tyr	Thr	Gly	Thr	Glu	Tyr	Gly	Ser	125	130	135	
Thr	Thr	Ile	Thr	His	Thr	Ser	Thr	Ala	Asp	Asn	Phe	Thr	Gln	Ala	Ser	140	145	150	
Val	Lys	Phe	Pro	Thr	Thr	Lys	Ala	Pro	Asp	Gly	Asn	Val	Leu	Tyr	Glu	155	160	165	
Leu	Thr	Val	Asp	Gly	Ser	Val	Ala	Ala	Gly	Ser	Ser	Leu	Asn	Phe	Gly	170	175	180	
Tyr	Leu	Thr	Leu	Phe	Gly	Glu	Thr	Tyr	Lys	Ser	Arg	Glu	Asn	Gly	Leu	185	190	195	200
Lys	Pro	Gln	Leu	Ala	Asn	Val	Leu	Asp	Asp	Met	Lys	Gly	Ser	Phe	Leu	205	210	215	
Arg	Phe	Pro	Gly	Gly	Asn	Asn	Leu	Glu	Gly	Asn	Ser	Ala	Glu	Asn	Arg	220	225	230	

SUBSTITUTE SHEET

FIGURE 15 (page 2 of 3)

Trp	Lys	Trp	Asn	Glu	Thr	Ile	Gly	Asp	Leu	Cys	Asp	Arg	Pro	Gly	Arg		
		235					240					245					
Glu	Gly	Thr	Trp	Thr	Tyr	Tyr	Asn	Thr	Asp	Gly	Leu	Gly	Leu	His	Glu		
	250					255					260						
Tyr	Phe	Tyr	Trp	Cys	Glu	Asp	Leu	Gly	Leu	Val	Pro	Val	Leu	Gly	Val		
265					270					275					280		
Trp	Asp	Gly	Phe	Ala	Leu	Glu	Ser	Gly	Gly	Asn	Thr	Pro	Leu	Thr	Gly		
				285					290					295			
Asp	Ala	Leu	Thr	Pro	Tyr	Ile	Asp	Asp	Val	Leu	Asn	Glu	Leu	Glu	Tyr		
			300					305					310				
Ile	Leu	Gly	Asp	Thr	Ser	Thr	Thr	Tyr	Gly	Ala	Trp	Arg	Ala	Ala	Asn		
		315					320					325					
Gly	Gln	Glu	Glu	Pro	Trp	Asn	Leu	Thr	Met	Val	Glu	Ile	Gly	Asn	Glu		
	330					335					340						
Asp	Met	Leu	Gly	Gly	Gly	Cys	Glu	Ser	Tyr	Ala	Glu	Arg	Phe	Thr	Ala		
345					350					355					360		
Phe	Tyr	Asp	Ala	Ile	His	Ala	Ala	Tyr	Pro	Asp	Leu	Ile	Leu	Ile	Ala		
				365					370					375			
Ser	Thr	Ser	Glu	Ala	Asp	Cys	Leu	Pro	Glu	Ser	Met	Pro	Glu	Gly	Ser		
			380					385					390				
Trp	Val	Asp	Tyr	His	Asp	Tyr	Ser	Thr	Pro	Asp	Gly	Leu	Val	Gly	Gln		
		395					400					405					
Phe	Asn	Tyr	Phe	Asp	Asn	Leu	Asn	Arg	Ser	Val	Pro	Tyr	Phe	Ile	Gly		
	410					415					420						
Glu	Tyr	Ser	Arg	Trp	Glu	Ile	Asp	Trp	Pro	Asn	Met	Lys	Gly	Ser	Val		
425					430					435				440			
Ala	Glu	Ala	Val	Phe	Met	Ile	Gly	Phe	Glu	Arg	Asn	Ser	Asp	Val	Val		
				445					450					455			
Lys	Met	Ala	Ala	Tyr	Ala	Pro	Leu	Leu	Gln	Leu	Ile	Asn	Ser	Thr	Gln		
			460					465					470				
Trp	Thr	Pro	Asp	Leu	Ile	Gly	Tyr	Thr	Gln	Ser	Pro	Gly	Asp	Ile	Phe		
		475					480					485					
Leu	Ser	Thr	Ser	Tyr	Tyr	Val	Gln	Glu	Met	Phe	Ser	Arg	Asn	Arg	Gly		
	490					495					500						
Asp	Thr	Ile	Lys	Glu	Val	Thr	Ser	Asp	Ser	Asp	Phe	Gly	Pro	Leu	Tyr		
505					510					515					520		

FIGURE 15 (page 3 of 3)

Trp	Val	Ala	Ser	Ser	Ala	Gly	Asp	Ser	Tyr	Tyr	Val	Lys	Leu	Ala	Asn
				525					530					535	
Tyr	Gly	Ser	Glu	Thr	Gln	Asp	Leu	Thr	Val	Ser	Ile	Pro	Gly	Thr	Ser
			540					545					550		
Thr	Gly	Lys	Leu	Thr	Val	Leu	Ala	Asp	Ser	Asp	Pro	Asp	Ala	Tyr	Asn
		555					560					565			
Ser	Asp	Thr	Gln	Thr	Leu	Val	Thr	Pro	Ser	Glu	Ser	Thr	Val	Gln	Ala
	570					575					580				
Ser	Asn	Gly	Thr	Phe	Thr	Phe	Ser	Leu	Pro	Ala	Trp	Ala	Val	Ala	Val
585					590					595					600
Leu	Ala	Ala	Asn												
			604												




Figure 16



Figure 17




Figure 18

FIGURE 19 (page 1 of 4)

```

(i) SEQUENCE CHARACTERISTICS:
    (A) LENGTH: 3147 base pairs
    (B) TYPE: nucleic acid
    (C) STRANDEDNESS: double
    (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: DNA (genomic)
(iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO
(vi) ORIGINAL SOURCE:
    (A) ORGANISM: Aspergillus niger
(ix) FEATURE:
    (A) NAME/KEY: exon
    (B) LOCATION: 1178..1444
(ix) FEATURE:
    (A) NAME/KEY: intron
    (B) LOCATION: 1445..1494
(ix) FEATURE:
    (A) NAME/KEY: exon
    (B) LOCATION: 1495..1971
(ix) FEATURE:
    (A) NAME/KEY: intron
    (B) LOCATION: 1972..2030
(ix) FEATURE:
    (A) NAME/KEY: exon
    (B) LOCATION: 2031..2116
(ix) FEATURE:
    (A) NAME/KEY: intron
    (B) LOCATION: 2117..2167
(ix) FEATURE:
    (A) NAME/KEY: exon
    (B) LOCATION: 2168..2378
(ix) FEATURE:
    (A) NAME/KEY: CDS
    (B) LOCATION: join(1178..1444, 1495..1971, 2031..2116,
2168..2378)
    (C) IDENTIFICATION METHOD: experimental
    (D) OTHER INFORMATION: /codon_start= 1178
                          /product= "endo 1,5-alpha-arabinanase"
                          /evidence= EXPERIMENTAL
                          /gene= "abnA"
(ix) FEATURE:
    (A) NAME/KEY: sig_peptide
    (B) LOCATION: 1178..1234
(ix) FEATURE:
    (A) NAME/KEY: mat_peptide
    (B) LOCATION: 1235..2378

```

(Formula 22)

```

AAGCTTGGTC GTGTCCGGG TAGTACTACC TCCGTAGAGA CTCCTCTGG ATCTTCTCAA      60
AAGGGGGTTC CGTGTCTCTG TACAGACTCT TCTAGTTCTC GCTTACGGCT CCACCCATTG    120

```

FIGURE 19 (page 2 of 4)

CTGAAGAACC CCACTCCTGC TGTTTCCCAT TTATGTTCCG GTCTAACGAT GGCTTCTCCA 180
CCAAGAAATG GAAGTGTGCT CTCCTCTCAA AACTCTCCAA TTCTGGTTTG ATCGATGTGA 240
GCTTACCAAG AAAGAGCGAA GACTTCCATC TTGTACTTAC GGACCCGATT AGCTATGCTT 300
CCTACACGTT NAANGACGAT NGACATATNN TGCTACCTGT TACTTGTGAG ATGGTCTCTC 360
TAACGCTGGC GGNATGTGGA CCAGATTCTC TTCAAAGTTA TGCCAAGATT CGCATGGGCC 420
AATCCTGAAA AGCGAAAAGC GCAAGGAGGC GCAGCTGAGA CAGAATCCCC TTTCTGCGAG 480
TCACATCTCG AGTCCCTTAA GGCCACCGTC GAGTGGTGGG CGATAAACTC CGCATATCTG 540
CCACCTGAAA AATATCGGTC AATACATCTA TGGCCTAATG AGACCAGTGG TCAGAATACG 600
CCTAAGGCCG GAAACTACTC CCGCAAATTG CTGCGACGAA GACGAGGGGG GCGTAGTCAT 660
AAATTTGCGT GTCGTGTGCC TATGATAATT TCTTGGGGTC CGAAGAGCAC GCTAGGGATG 720
TCAGAACAAC TAATGTTGCA GGATTGGTCT CCATCTCATC CCTGTCACTG ACATCATATT 780
ACTTCCTAGT ATATCATGTT CACCATTGTA TAGAGAGACG CGCCCATCGC AACGCACGTG 840
TTGGTGAGCA CCTCCACTAA AAGTGTCCTT GTTGTCTACA CACCACTCGT TTCACGCTGA 900
TGTGCCGAAT CATACTACAT GCATCGGCCA TCGGCTTATA CCACCAGGTG CCTCCGGAAT 960
TTTGCTGTTT CAAATGTTT TTTGTAATC CCTGAAACGA CCAAAGTATC ACGGAGGTAG 1020
CTCCGGACAG GCGGAAAACA CAACCACTTT GTCCACGAAT CGACACAGTA GTCTTCCGGT 1080
GGGTTTAGTT GTCGCTTCAT GATAGTATAT ATAGTAATAG TAGATCTGGA TGCTTGGGAT 1140
AAGTCATCCA GTGGTGACAG CGCCTACCTA AATCGTCATG TATCAACTCC TATCAGTTGC 1200

FIGURE 19 (page 3 of 4)

CTCGGTTCTT	CTGCTGGCCA	GCCTCGTGCA	TGGCTATGCT	GACCCCGGAG	CATGCTCGGG	1260
TGTTTGTACC	ACCCATGACC	CCGGTTTGAT	CCGGAGAGAG	TCGGACGGTA	CATACTTCCT	1320
CTTCTCTACA	GGAAACAAAA	TCTCCTACGT	CTCTGCATCA	TCCATCGAAG	GACCATGGAC	1380
CAGCGTTGGG	TCTATGCTGC	CGGATGGATC	GTCCATCGAC	CTCGATGGCA	ACGATGATCT	1440
TTGGGTAACT	ATGTCCCACT	GCAAACATGA	TTGGCCTGTC	TGACACTAGC	CCAGGCCCCG	1500
GATGTCTCCT	ATGTAGATGG	TCTCTATTAT	GTATACTACG	CTGTGTCGAC	CTTTGGATCC	1560
CAAGATTCCG	CCATCGGACT	TGCAACGTCC	GAGACGATGG	AATATGGCTC	TTGGACGGAT	1620
CATGGCTCCA	CTGGCATTGC	GTCTTCCTCA	GCCAAGATTT	ATAATGCGAT	CGACCCCAAC	1680
CTGATCTACG	CCGATGGCAC	CTACTACATC	AACTTTGGGT	CGTTCTGGGA	TGACATTTAC	1740
CAAGTCCCGA	TGAAGTCGAC	CCCAACGGCA	GCTGCCTCTT	CCTCCTACAA	TCTTGCGTAT	1800
GACCCGTCCG	GTACCCATGC	GGAGGAGGGT	TCCTATATGT	TTCAGTACGG	TGACTACTAC	1860
TACCTCTTTT	ACTCGGCAGG	TATCTGCTGT	GGATACGATA	CATCCATGCC	CGCTTCCGGA	1920
GAGGAGTATC	ATATCAAGGT	CTGCCGTTTG	ACTTCGCCCCA	CGGGTGATTT	CGTAAGTATA	1980
TCCGACATTT	AGAGAAAATA	CTAAGTAGGG	AAAGCTAACC	TGGTGCTAAG	GTTGACTCCG	2040
ACGGTACGGC	GTGCACGGAT	GGCGGGGGCA	CGATGGTGCT	CGAAAGCCAT	GGAGAAGTCT	2100
ATGGCCCTGG	CGGACAGTAA	GTGATTCTGC	AGCTGCCCCA	CCGATATTTA	TGCTTATCAT	2160
GTTTTAGGGG	TGTGTATGAC	GATCCCAACC	TTGGTCCGGT	TCTTTACTAC	CACTACATGA	2220
ACACCACGAT	TGGATACGCG	GATTCTGACG	CGCAGTTTGG	GTGGAACACG	ATCGACTTCT	2280

FIGURE 19 (page 4 of 4)

CTGACGGATG GACCGGTTGT ATAAGCATCT TGGTTGGCGA TGGAAATGGT GCTGCCTACG	2340
AGAGGGTCTA CAAGTTGTAT ATAGGATTGC GGCTATGAAC TCGGCCGAGA TATAGTTAGT	2400
TAGTAACTAC ATAGTGAAT CGTTAGTCAC TAATGCAGCC CGCATGAGTG CTCTGCCTGA	2460
GGCTTCAGTA CTGTCCCTCA ATTTCTGTTT CTGTTTTTAG TTTAAGCTCT GCAGCTGCTC	2520
CCCACCCATT CATCCACTCA TTTGGTGAAG ACATTCATTC TTCCCTTCAA CCACTCTATT	2580
AGAGCATTTT CATCTTCTCA CACTGATCAT CGTAATGCAG ATTCAAGATC AGTTTGGATT	2640
GAGATCCCTA CTGCATGCAC TGTGTGCGTG ACAGAAGCTG GAACTGGTAC TCGCTCGATC	2700
CGTGCAAGCG GAGTCAGTCA CATCCGCCCT GCGCTTCAGA CTAGCCTGCC AATAGTGACT	2760
ATCACAAGGG CCGTACTAAG ATCGAATAAC AGCTTTCCTA ACAACATCGC GTTAACGTGT	2820
TACTTTCCTC CAGCTTACAT TTTGCTATTG AAGCCATATA CGATCCCTTC GATGCGCTCG	2880
GTTTAACTTC TGCCACGATC ACCCGTTCAT CGAAGATTCT GGGAGTCTGT CGGTCTTCAT	2940
GCCAGCAAAG ATCGACTCAA ATACACACAG TCAGCAACAT CTGTTGCTTG TTACGGATGG	3000
TGCCTATTAT AATGCATGAA TAGGGGACCA AGCAATTCTC CGCTGTCGGC GATGTTGAAT	3060
TAAGGAGGGA TAATATTTGA TCGAATGACA AAGGGAAGAA TTGTGCGACA CGAGGGAACC	3120
TTATACTTGT CTACGGCCTA GCAGCAA	3147

FIGURE 20 (page 1 of 2)

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 346 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein

(Formula 23)

```

Met Tyr Gln Leu Leu Ser Val Ala Ser Val Pro Leu Leu Ala Ser Leu
-19              -15              -10              -5

Val His Gly Tyr Ala Asp Pro Gly Ala Cys Ser Gly Val Cys Thr Thr
      -1  +1              5              10

His Asp Pro Gly Leu Ile Arg Arg Glu Ser Asp Gly Thr Tyr Phe Leu
  15              20              25

Phe Ser Thr Gly Asn Lys Ile Ser Tyr Val Ser Ala Ser Ser Ile Glu
  30              35              40              45

Gly Pro Trp Thr Ser Val Gly Ser Met Leu Pro Asp Gly Ser Ser Ile
              50              55              60

Asp Leu Asp Gly Asn Asp Asp Leu Trp Ala Pro Asp Val Ser Tyr Val
              65              70              75

Asp Gly Leu Tyr Tyr Val Tyr Tyr Ala Val Ser Thr Phe Gly Ser Gln
      80              85              90

Asp Ser Ala Ile Gly Leu Ala Thr Ser Glu Thr Met Glu Tyr Gly Ser
  95              100              105

Trp Thr Asp His Gly Ser Thr Gly Ile Ala Ser Ser Ser Ala Lys Ile
 110              115              120              125

Tyr Asn Ala Ile Asp Pro Asn Leu Ile Tyr Ala Asp Gly Thr Tyr Tyr
              130              135              140

Ile Asn Phe Gly Ser Phe Trp Asp Asp Ile Tyr Gln Val Pro Met Lys
      145              150              155

Ser Thr Pro Thr Ala Ala Ala Ser Ser Ser Tyr Asn Leu Ala Tyr Asp
      160              165              170

Pro Ser Gly Thr His Ala Glu Glu Gly Ser Tyr Met Phe Gln Tyr Gly
 175              180              185

Asp Tyr Tyr Tyr Leu Phe Tyr Ser Ala Gly Ile Cys Cys Gly Tyr Asp
 190              195              200              205

Thr Ser Met Pro Ala Ser Gly Glu Glu Tyr His Ile Lys Val Cys Arg
      210              215              220

```

FIGURE 20 (page 2 of 2)

Ser	Thr	Ser	Pro	Thr	Gly	Asp	Phe	Val	Asp	Ser	Asp	Gly	Thr	Ala	Cys
			225					230					235		
Thr	Asp	Gly	Gly	Gly	Thr	Met	Val	Leu	Glu	Ser	His	Gly	Glu	Val	Tyr
		240					245					250			
Gly	Pro	Gly	Gly	Gln	Gly	Val	Tyr	Asp	Asp	Pro	Asn	Leu	Gly	Pro	Val
	255					260					265				
Leu	Tyr	Tyr	His	Tyr	Met	Asn	Thr	Thr	Ile	Gly	Tyr	Ala	Asp	Ser	Asp
270					275					280					285
Ala	Gln	Phe	Gly	Trp	Asn	Thr	Ile	Asp	Phe	Ser	Asp	Gly	Trp	Thr	Gly
				290					295					300	
Cys	Ile	Ser	Ile	Leu	Val	Gly	Asp	Gly	Asn	Gly	Ala	Ala	Tyr	Glu	Arg
			305					310					315		
Val	Tyr	Lys	Leu	Tyr	Ile	Gly	Leu	Arg	Leu						
		320					325								

1 2 3 4 5 6 7 8 9 10 11 12 13 14




Figure 21




Figure 22

INTERNATIONAL SEARCH REPORT

PCT/NL 92/00056

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all)⁶

According to International Patent Classification (IPC) or to both National Classification and IPC
 Int.Cl. 5 C12N15/56; C12N9/24

II. FIELDS SEARCHED

Minimum Documentation Searched⁷

Classification System

Classification Symbols

Int.Cl. 5

C12N ;

A23L ;

A23K

Documentation Searched other than Minimum Documentation
 to the Extent that such Documents are Included in the Fields Searched⁸

III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹

Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS vol. 170, no. 1, 16 July 1990, DULUTH MINNESOTA US pages 368 - 374; W.H.SCHWARZ ET AL.: 'Xylan degradation by the thermophile Clostridium stercorarium:cloning and expression of xylanase,beta-D-xylosidase,and alfa-L-arabinofuranosidase genes in E.coli' cited in the application see the whole document --- -/--	1-3,9, 16,19, 22,25, 26,28, 29,31

¹⁰ Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search

Date of Mailing of this International Search Report

25 JUNE 1992

14. 07. 92

International Searching Authority

Signature of Authorized Officer

EUROPEAN PATENT OFFICE

GURDJIAN D.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y	CARBOHYDRATE POLYMERS vol. 9, 1988, BARKING GB pages 25 - 47; F.M.ROMBOUTS ET AL.: 'The arabinanases of Aspergillus niger-Purification and characterisation of two alfa-L-arabinofuranosidases and an endo-1,5-alfa-l-arabinanase' see the whole document	1,2,9, 16,19, 22,25, 28,31
A	---	12,15
Y	CARBOHYDRATE RESEARCH vol. 144, 1985, AMSTERDAM NL pages 113 - 126; J.M.BRILLOUET ET AL.: 'Production, purification , and properties of an alfa-L-arabinofuranosidase from Dichomitus squalens' see abstract	3,26,29
A	---	12,14
A	ADVANCES IN CARBOHYDRATE CHEMISTRY AND BIOCHEMISTRY vol. 42, 1984, NEW YORK US pages 383 - 394; AKIRA KAJI: 'l-arabinosidases' cited in the application see the whole document	9,28,31
A	---	16,19, 22,25
A	MEDLINE DATABASE Ab.No.90236896 Whitehead Tr. et.al.: "The genes for three xylan- degrading activities from Bacteroides ovatus are clustered in a 3.8kB region" & J.Bacteriol., 1990, vol.172, n.5, pages 2408-12 cited in the application	16,19, 22,25
A	EP,A,0 311 469 (PLANT GENETIC SYTEMS N.V. ET AL.) 12 April 1989 see abstract; claims 1,2	16,19, 22,25
A	---	33,36
A	WORLD PATENTS INDEX LATEST Week 8451, Derwent Publications Ltd., London, GB; AN 84-316055 & JP,A,59 198 978 (OKADA K) 10 November 1984 see abstract	33,34
A	---	33,34
A	WORLD PATENTS INDEX LATEST Week 9117, Derwent Publications Ltd., London, GB; AN 91-104119 & AU,D,6 205 490 (CIBA GEIGY AG) 7 March 1991 see abstract	33,34

-/--

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	WO,A,8 504 672 (VALTION TEKNILLINEN TUTKIMUSKESUS) 24 October 1985 see abstract see page 2, line 4 - line 25 ---	33,37
A	JOURNAL OF AGRICULTURAL AND FOOD CHEMISTRY vol. 38, no. 3, 1990, TOKYO pages 772 - 776; Gunata, Ziya; Brillouet, Jean Marc; Voirin, Stephane; Baumes, Raymond; Cordonnier, Robert: 'Purification and some properties of an .alpha.-L-arabinofuranosidase from Aspergillus niger. Action on grape monoterpenyl arabinofuranosylglucosides' see the whole document ---	35

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. NL 9200056
SA 58342**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 25/06/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0311469	12-04-89	WO-A- 8901970	09-03-89
		JP-T- 2501266	10-05-90
WO-A-8504672	24-10-85	EP-A- 0214971	25-03-87
		EP-A- 0312121	19-04-89
		US-A- 4894338	16-01-90