



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION

(13) **A1**

(86) **Date de dépôt PCT/PCT Filing Date:** 2022/10/27
(87) **Date publication PCT/PCT Publication Date:** 2023/05/04
(85) **Entrée phase nationale/National Entry:** 2024/04/15
(86) **N° demande PCT/PCT Application No.:** US 2022/078771
(87) **N° publication PCT/PCT Publication No.:** 2023/076991
(30) **Priorité/Priority:** 2021/10/28 (US63/272,899)

(51) **Cl.Int./Int.Cl.** **A61K 31/352** (2006.01),
A61P 35/00 (2006.01)
(71) **Demandeur/Applicant:**
VERASTEM, INC., US
(72) **Inventeurs/Inventors:**
PACHTER, JONATHAN A., US;
CHOWDHURY, SANJIB, US;
COMA, SILVIA, US
(74) **Agent:** SMART & BIGGAR LP

(54) **Titre : POLYTHÉRAPIE POUR LE TRAITEMENT D'UNE CROISSANCE CELLULAIRE ANORMALE**
(54) **Title: COMBINATION THERAPY FOR TREATING ABNORMAL CELL GROWTH**

			Synergy Score					
			VS-6766 + BRAFi			MEKi + BRAFi		
Cell line	MAPK alteration	PAM alterations	VS-6766 + vemurafenib	VS-6766 + dabrafenib	VS-6766 + encorafenib	cobimetinib + vemurafenib	trametinib + dabrafenib	binimetinib + encorafenib
A2058	BRAF V600E	PIK3R6	17.615	15.797	17.747	6.682	-5.699	13.419
IGR-1	BRAF V600E	PIK3C2G, PIK3R5, PIK3CG	16.434	9.543	13.119	14.073	-4.811	9.585
SK-MEL-5	BRAF V600E	PIK3CB, PIK3C2A, RICTOR	12.431	3.829	9.094	0.564	-7.886	8.691
WM-266-4	BRAF V600E		6.114	4.522	6.603	1.568	1.025	3.527
C32	BRAF V600E	PIK2CB, mTOR	11.793	9.287	9.268	3.14	0.486	7.802




FIG. 1

(57) **Abrégé/Abstract:**

Combinations of compounds as described herein, a BRAF V600 inhibitor, a pan-RAF inhibitor, a CRAF inhibitor, or a RAF inhibitor, and a dual RAF/MEK inhibitor, can be used, for example, in methods of treating abnormal cell growth/cancers in a subject in need thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
04 May 2023 (04.05.2023)



(10) International Publication Number
WO 2023/076991 A1

(51) International Patent Classification:

A61K 31/352 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2022/078771

(22) International Filing Date:

27 October 2022 (27.10.2022)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

63/272,899 28 October 2021 (28.10.2021) US

(71) Applicant: **VERASTEM, INC.** [US/US]; 117 Kendrick Street, Suite 500, Needham, Massachusetts 02494 (US).

(72) Inventors: **PACHTER, Jonathan A.**; 117 Kendrick Street, Suite 500, Needham, Massachusetts 02494 (US). **CHOWDHURY, Sanjib**; 117 Kendrick Street, Suite 500, Needham, Massachusetts 02494 (US). **COMA, Silvia**; 117 Kendrick Street, Suite 500, Needham, Massachusetts 02494 (US).

(74) Agent: **CHOI, Boyeon** et al.; Goodwin Procter LLP, 100 Northern Avenue, IP DOCKETING DEPT./7TH FL, Boston, Massachusetts 02210 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CV, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, CV, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, ME, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: COMBINATION THERAPY FOR TREATING ABNORMAL CELL GROWTH

			Synergy Score					
Cell line	MAPK alteration	PAM alterations	VS-6766 + BRAFi			MEKi + BRAFi		
			VS-6766 + vemurafenib	VS-6766 + dabrafenib	VS-6766 + encorafenib	cobimetinib + vemurafenib	trametinib + dabrafenib	binimetinib + encorafenib
A2058	BRAF V600E	PIK3R6	17.615	15.797	17.747	6.682	-5.699	13.419
IGR-1	BRAF V600E	PIK3C2G, PIK3R5, PIK3CG	16.434	9.543	13.119	14.073	4.811	9.565
SK-MEL-5	BRAF V600E	PIK3CB, PIK3C2A, RICTOR	12.431	3.829	9.094	0.564	-7.886	8.691
WM-266-4	BRAF V600E		6.114	4.522	6.603	1.568	1.025	3.527
C32	BRAF V600E	PIK2CB, mTOR	11.793	9.287	9.268	3.14	0.466	7.802




FIG. 1

(57) Abstract: Combinations of compounds as described herein, a BRAF V600 inhibitor, a pan-RAF inhibitor, a CRAF inhibitor, or a RAF inhibitor, and a dual RAF/MEK inhibitor, can be used, for example, in methods of treating abnormal cell growth/cancers in a subject in need thereof.

[Continued on next page]



WO 2023/076991 A1

WO 2023/076991 A1 **Declarations under Rule 4.17:**

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *of inventorship (Rule 4.17(iv))*

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

COMBINATION THERAPY FOR TREATING ABNORMAL CELL GROWTH**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims the benefit of and priority to U.S. Provisional Patent Application No. 63/272,899 filed October 28, 2021, the entire contents of which are incorporated herein by reference.

BACKGROUND

[0002] Components of the RAS/RAF/MEK/ERK (MAPK) signal transduction pathway represent opportunities for the treatment of abnormal cell growth, e.g., cancer. Mutations in RAS/RAF/MEK/ERK are frequently found in human cancers. These mutants result in a constitutively active MAPK kinase cascade, leading to tumor cell proliferation, differentiation, survival, and migration. Selective inhibitors of certain components of the RAS/RAF/MEK/ERK signal transduction pathway, such as RAS, RAF, MEK and ERK, are useful in the treatment of abnormal cell growth, in particular cancer, in mammals.

[0003] Due to the severity and breadth of diseases and disorders associated with abnormal cell growth, e.g., cancer, there is a need for effective therapeutic means and methods for treatment. The compounds, compositions, and methods described herein are directed toward this end.

SUMMARY

[0004] The present disclosure provides, in part, combinations (e.g., combinations of compounds as described herein, e.g., a BRAF V600 inhibitor, a pan-RAF inhibitor, a CRAF inhibitor, or a RAF inhibitor, and a dual RAF/MEK inhibitor), which can be used, for example, in methods of treating abnormal cell growth (e.g., cancer) in a subject in need thereof.

[0005] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a BRAF V600 inhibitor.

[0006] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a BRAF V600 inhibitor, wherein the

cancer is identified as having a BRAF V600 mutation. In some embodiments, the BRAF V600 mutation is BRAF V600E, BRAF V600K, BRAF V600D, BRAF V600R, and/or BRAF V600M mutation.

[0007] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a pan-RAF inhibitor.

[0008] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a pan-RAF inhibitor, wherein the cancer is identified as having ARAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation.


[0009] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a CRAF inhibitor.

[00010] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a CRAF inhibitor, wherein the cancer is identified as having ARAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation.

[00011] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a RAF inhibitor.


[00012] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a RAF inhibitor, wherein the cancer is identified as having ARAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation.

[00013] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (I):



(I), or a pharmaceutically acceptable salt thereof.

[00014] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (II):



including pharmaceutically acceptable salts thereof, wherein the variables are as defined herein.

[00015] In some embodiments, the dual RAF/MEK inhibitor is IMM-1-104, or a pharmaceutically acceptable salt thereof.

[00016] Other objects and advantages will become apparent to those skilled in the art from a consideration of the ensuing **Detailed Description, Examples, and Claims**.

BRIEF DESCRIPTION OF THE DRAWINGS

[00017] **FIG. 1** illustrates the synergy of VS-6766 (dual RAF/MEK inhibitor) + BRAF V600 inhibitor (BRAFi) is better than synergy of MEK inhibitor (MEKi) + BRAF V600 inhibitor (BRAFi) in BRAF V600E melanoma cell lines. 3D proliferation studies with VS-6766 + BRAFi or MEKi + BRAFi in a panel of BRAF V600E melanoma cell lines were performed to calculate synergy of VS-6766 + BRAFi and MEKi + BRAFi. Bliss, Loewe, HSA and ZIP synergy analyses were performed to generate a composite synergy score.

[00018] **FIG. 2** illustrates the synergy of VS-6766 (dual RAF/MEK inhibitor) + vemurafenib (BRAFi) is better than synergy of cobimetinib (MEKi) + vemurafenib (BRAFi) in BRAF V600E melanoma cell lines. Dose-response matrices were used to assess anti-proliferative effects of the combination of VS-6766 (1:5 dilutions starting at 5 μ M) + vemurafenib (1:3 dilutions starting at 5 μ M) and cobimetinib (1:5 dilutions starting at 5 μ M) + vemurafenib (1:3 dilutions starting at 5 μ M) in a panel of BRAF V600E melanoma cancer cell lines. Synergy scores were calculated using a combination of 4 different methods (Bliss, Loewe, HSA and ZIP). Bliss synergy analysis are shown. Red = synergy; Green = antagonism.

[00019] **FIG. 3** illustrates the synergy of VS-6766 (dual RAF/MEK inhibitor) + dabrafenib (BRAFi) is better than synergy of trametinib (MEKi) + dabrafenib (BRAFi) in BRAF V600E melanoma cell lines. Dose-response matrices were used to assess anti-proliferative effects of the combination of VS-6766 (1:5 dilutions starting at 5 μ M) + dabrafenib (1:3 dilutions starting at 5 μ M) and trametinib (1:5 dilutions starting at 5 μ M) +

dabrafenib (1:3 dilutions starting at 5 μ M) in a panel of BRAF V600E melanoma cancer cell lines. Synergy scores were calculated using a combination of 4 different methods (Bliss, Loewe, HSA and ZIP). Bliss synergy analysis are shown. Red = synergy; Green = antagonism.

[00020] **FIG. 4** illustrates the synergy of VS-6766 + BRAFi is better than synergy of MEKi + BRAFi in BRAF V600E colorectal carcinoma cell lines. 3D proliferation studies with VS-6766 + BRAFi or MEKi + BRAFi in a panel of BRAF V600E colorectal carcinoma cell lines were performed to calculate synergy of VS-6766 + BRAFi and MEKi + BRAFi. Bliss, Loewe, HSA and ZIP synergy analyses were performed to generate a composite synergy score.

[00021] **FIG. 5** illustrates the synergy of VS-6766 (dual RAF/MEK inhibitor) + vemurafenib (BRAFi) is better than synergy of cobimetinib (MEKi) + vemurafenib (BRAFi) in BRAF V600E colorectal carcinoma cell lines. Dose-response matrices were used to assess anti-proliferative effects of the combination of VS-6766 (1:5 dilutions starting at 5 μ M) + vemurafenib (1:3 dilutions starting at 5 μ M) and cobimetinib (1:5 dilutions starting at 5 μ M) + vemurafenib (1:3 dilutions starting at 5 μ M) in a panel of BRAF V600E CRC cell lines. Synergy scores were calculated using a combination of 4 different methods (Bliss, Loewe, HSA and ZIP). Bliss synergy analysis are shown. Red = synergy; Green = antagonism.

[00022] **FIG. 6** illustrates the synergy of VS-6766 (dual RAF/MEK inhibitor) + dabrafenib (BRAFi) is better than synergy of trametinib (MEKi) + dabrafenib (BRAFi) in BRAF V600E colorectal carcinoma cell lines. Dose-response matrices were used to assess anti-proliferative effects of the combination of VS-6766 (1:5 dilutions starting at 5 μ M) + dabrafenib (1:3 dilutions starting at 5 μ M) and trametinib (1:5 dilutions starting at 5 μ M) + dabrafenib (1:3 dilutions starting at 5 μ M) in a panel of BRAF V600E CRC cell lines. Synergy scores were calculated using a combination of 4 different methods (Bliss, Loewe, HSA and ZIP). Bliss synergy analysis are shown. Red = synergy; Green = antagonism.

[00023] **FIG. 7** illustrates strong synergy observed with VS-6766 (RAF/MEK clamp) + pan-RAFi in NRAS mt melanoma cell lines. 3D proliferation studies with VS-6766 + pan-RAFi in a panel of NRAS mutant melanoma cell lines were performed to calculate synergy of VS-6766 + pan-RAFi. Bliss, Loewe, HSA and ZIP synergy analyses were performed to generate a composite synergy score.

[00024] **FIG. 8** illustrates strong synergy observed with VS-6766 (dual RAF/MEK inhibitor) + pan-RAFi in NRAS mt melanoma cell lines. Dose-response matrices were used to assess anti-proliferative effects of the combination of VS-6766 (1:5 dilutions starting at 5

μM) + pan-RAFi (1:3 dilutions starting at 5 μM) in a panel of NRAS mutant melanoma cell lines. Synergy scores were calculated using a combination of 4 different methods (Bliss, Loewe, HSA and ZIP). Bliss synergy analysis are shown. Red = synergy; Green = antagonism.

DETAILED DESCRIPTION

[00025] As generally described herein, the present disclosure provides, in part, methods useful for treating abnormal cell growth (e.g., cancer) in a subject in need thereof, comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a BRAF V600 inhibitor, a pan-RAF inhibitor, a CRAF inhibitor, or a RAF inhibitor.

Definitions

Chemical definitions

[00026] Definitions of specific functional groups and chemical terms are described in more detail below. The chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, *Handbook of Chemistry and Physics*, 75th Ed., inside cover, and specific functional groups are generally defined as described therein.

Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in Thomas Sorrell, *Organic Chemistry*, University Science Books, Sausalito, 1999; Smith and March, *March's Advanced Organic Chemistry*, 5th Edition, John Wiley & Sons, Inc., New York, 2001; Larock, *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, 1989; and Carruthers, *Some Modern Methods of Organic Synthesis*, 3rd Edition, Cambridge University Press, Cambridge, 1987.

[00027] Compounds described herein can comprise one or more asymmetric centers, and thus can exist in various isomeric forms, e.g., enantiomers and/or diastereomers. For example, the compounds described herein can be in the form of an individual enantiomer, diastereomer or geometric isomer, or can be in the form of a mixture of stereoisomers, including racemic mixtures and mixtures enriched in one or more stereoisomer. Isomers can be isolated from mixtures by methods known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts; or preferred isomers can be prepared by asymmetric syntheses. See, for example, Jacques *et al.*, *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen *et al.*, *Tetrahedron* 33:2725 (1977); Eliel, *Stereochemistry of Carbon Compounds* (McGraw-

Hill, NY, 1962); and Wilen, *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972). The invention additionally encompasses compounds described herein as individual isomers substantially free of other isomers, and alternatively, as mixtures of various isomers.

[00028] As used herein a pure enantiomeric compound is substantially free from other enantiomers or stereoisomers of the compound (*i.e.*, in enantiomeric excess). In other words, an “S” form of the compound is substantially free from the “R” form of the compound and is, thus, in enantiomeric excess of the “R” form. The term “enantiomerically pure” or “pure enantiomer” denotes that the compound comprises more than 75% by weight, more than 80% by weight, more than 85% by weight, more than 90% by weight, more than 91% by weight, more than 92% by weight, more than 93% by weight, more than 94% by weight, more than 95% by weight, more than 96% by weight, more than 97% by weight, more than 98% by weight, more than 98.5% by weight, more than 99% by weight, more than 99.2% by weight, more than 99.5% by weight, more than 99.6% by weight, more than 99.7% by weight, more than 99.8% by weight or more than 99.9% by weight, of the enantiomer. In some embodiments, the weights are based upon total weight of all enantiomers or stereoisomers of the compound.

[00029] In the compositions provided herein, an enantiomerically pure compound can be present with other active or inactive ingredients. For example, a pharmaceutical composition comprising enantiomerically pure R-compound can comprise, for example, about 90% excipient and about 10% enantiomerically pure R-compound. In some embodiments, the enantiomerically pure R-compound in such compositions can, for example, comprise, at least about 95% by weight R-compound and at most about 5% by weight S-compound, by total weight of the compound. For example, a pharmaceutical composition comprising enantiomerically pure S-compound can comprise, for example, about 90% excipient and about 10% enantiomerically pure S-compound. In some embodiments, the enantiomerically pure S-compound in such compositions can, for example, comprise, at least about 95% by weight S-compound and at most about 5% by weight R-compound, by total weight of the compound. In some embodiments, the active ingredient can be formulated with little or no excipient or carrier.

[00030] Compound described herein may also comprise one or more isotopic substitutions. For example, H may be in any isotopic form, including ^1H , ^2H (D or deuterium), and ^3H (T or tritium); C may be in any isotopic form, including ^{12}C , ^{13}C , and ^{14}C ;

O may be in any isotopic form, including ^{16}O and ^{18}O ; F may be in any isotopic form, including ^{18}F and ^{19}F ; and the like.

[00031] The following terms are intended to have the meanings presented therewith below and are useful in understanding the description and intended scope of the present invention. When describing the invention, which may include compounds and pharmaceutically acceptable salts thereof, pharmaceutical compositions containing such compounds and methods of using such compounds and compositions, the following terms, if present, have the following meanings unless otherwise indicated. It should also be understood that when described herein any of the moieties defined forth below may be substituted with a variety of substituents, and that the respective definitions are intended to include such substituted moieties within their scope as set out below.

[00032] The term “halogen atom,” as used herein, means any one of the radio stable atoms of column 7 of the Periodic Table of the Elements, e.g., fluorine, chlorine, bromine, or iodine, with fluorine and chlorine being preferred.

[00033] The term “ester,” as used herein, refers to a chemical moiety with formula $-(\text{R})_n\text{-COOR}'$, where R and R' are independently selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon), and where n is 0 or 1.

[00034] The term “amide,” as used herein, refers to a chemical moiety with formula $-(\text{R})_n\text{-C(O)NHR}'$ or $-(\text{R})_n\text{-NHC(O)R}'$, where R and R' are independently selected from the group consisting of alkyl, cycloalkyl, aryl, heteroaryl (bonded through a ring carbon) and heteroalicyclic (bonded through a ring carbon), and where n is 0 or 1. An amide may be an amino acid or a peptide molecule attached to a molecule of the present invention, thereby forming a prodrug.

[00035] Any amine, hydroxyl, or carboxyl side chain on the compounds disclosed herein can be esterified or amidified. The procedures and specific groups to be used to achieve this end are known to those of skill in the art and can readily be found in reference sources such as Greene and Wuts, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley & Sons, New York, NY, 1999, which is incorporated herein in its entirety.

[00036] The term “aromatic,” as used herein, refers to an aromatic group which has at least one ring having a conjugated pi electron system and includes both carbocyclic aryl (e.g., phenyl) and heterocyclic aryl groups (e.g., pyridine). The term includes monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of carbon atoms) groups. The term “carbocyclic” refers to a compound which contains one or more covalently closed ring

structures, and that the atoms forming the backbone of the ring are all carbon atoms. The term thus distinguishes carbocyclic from heterocyclic rings in which the ring backbone contains at least one atom which is different from carbon. The term “hetero aromatic” refers to an aromatic group which contains at least one heterocyclic ring.

[00037] As used herein, “Ca to Cb” in which “a” and “b” are integers refer to the number of carbon atoms in an alkyl, alkenyl or alkynyl group, or the number of carbon atoms in the ring of a cycloalkyl, aryl, heteroaryl or heterocyclyl group. That is, the alkyl, alkenyl, alkynyl, ring of the cycloalkyl, ring of the aryl, ring of the heteroaryl or ring of the heterocyclyl can contain from “a” to “b”, inclusive, carbon atoms. Thus, for example, a “C1 to C4 alkyl” group or a “C1-C4 alkyl” group refers to all alkyl groups having from 1 to 4 carbons, that is, CH_3 -, CH_3CH_2 -, $\text{CH}_3\text{CH}_2\text{CH}_2$ -, $(\text{CH}_3)_2\text{CH}$ -, $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2$ -, $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)$ - and $(\text{CH}_3)_3\text{C}$ -. Likewise, for example, cycloalkyl group may contain from “a” to “b”, inclusive, total atoms, such as a C3-C8 cycloalkyl group, 3 to 8 carbon atoms in the ring(s). If no “a” and “b” are designated with regard to an alkyl, cycloalkyl, or cycloalkenyl, the broadest range described in these definitions is to be assumed. Similarly, a “4 to 7 membered heterocyclyl” group refers to all heterocyclyl groups with 4 to 7 total ring atoms, for example, azetidine, oxetane, oxazoline, pyrrolidine, piperidine, piperazine, morpholine, and the like. As used herein, the term “C1-C6” includes C1, C2, C3, C4, C5 and C6, and a range defined by any of the two preceding numbers. For example, C1-C6 alkyl includes C1, C2, C3, C4, C5 and C6 alkyl, C2-C6 alkyl, C1-C3 alkyl, etc. Similarly, C3-C8 carbocyclyl or cycloalkyl each includes hydrocarbon ring containing 3, 4, 5, 6, 7 and 8 carbon atoms, or a range defined by any of the two numbers, such as C3-C7 cycloalkyl or C5-C6 cycloalkyl. As another example, 3 to 10 membered heterocyclyl includes 3, 4, 5, 6, 7, 8, 9, or 10 ring atoms, or a range defined by any of the two preceding numbers, such as 4 to 6 membered or 5 to 7 membered heterocyclyl.

[00038] As used herein, “alkyl” refers to a straight or branched hydrocarbon chain fully saturated (no double or triple bonds) hydrocarbon group. The alkyl group may have 1 to 20 carbon atoms (whenever it appears herein, a numerical range such as “1 to 20” refers to each integer in the given range; e.g., “1 to 20 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc., up to and including 20 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). The alkyl group may also be a medium size alkyl having 1 to 10 carbon atoms. The alkyl group could also be a lower alkyl having 1 to 5 carbon atoms. The alkyl group of the compounds may be designated as “C1-C4 alkyl” or similar

designations. By way of example only, “C1-C4 alkyl” indicates that there are one to four carbon atoms in the alkyl chain, i.e., the alkyl chain is selected from the group consisting of methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and t-butyl. Exemplary alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl, hexyl, ethenyl, propenyl, butenyl, and the like.

[00039] The alkyl group may be substituted or unsubstituted. When substituted, the substituent group(s) is(are) one or more group(s) individually and independently selected from alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl, heteroalicycyl, aralkyl, heteroaralkyl, (heteroalicycyl)alkyl, hydroxy, protected hydroxyl, alkoxy, aryloxy, acyl, ester, mercapto, alkylthio, arylthio, cyano, halogen, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, protected C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, silyl, sulfenyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, and amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. Wherever a substituent is described as being “optionally substituted” that substituent may be substituted with one of the above substituents.

[00040] As used herein, “alkenyl” refers to an alkyl group that contains in the straight or branched hydrocarbon chain one or more double bonds. An alkenyl group may be unsubstituted or substituted. When substituted, the substituent(s) may be selected from the same groups disclosed above with regard to alkyl group substitution. The alkenyl group may have 2 to 20 carbon atoms, although the present definition also covers the occurrence of the term “alkenyl” where no numerical range is designated. The alkenyl group may also be a medium size alkenyl having 2 to 9 carbon atoms. The alkenyl group could also be a lower alkenyl having 2 to 4 carbon atoms. The alkenyl group of the compounds may be designated as “C2-C4 alkenyl” or similar designations. By way of example only, “C2-C4 alkenyl” indicates that there are two to four carbon atoms in the alkenyl chain, i.e., the alkenyl chain is selected from the group consisting of ethenyl, propen-1-yl, propen-2-yl, propen-3-yl, buten-1-yl, buten-2-yl, buten-3-yl, buten-4-yl, 1-methyl-propen-1-yl, 2-methyl-propen-1-yl, 1-ethyl-ethen-1-yl, 2-methyl-propen-3-yl, buta-1,3-dienyl, buta-1,2,-dienyl, and buta-1,2-dien-4-yl. Exemplary alkenyl groups include, but are in no way limited to, ethenyl, propenyl, butenyl, pentenyl, and hexenyl, and the like.

[00041] As used herein, “alkynyl” refers to an alkyl group that contains in the straight or branched hydrocarbon chain one or more triple bonds. An alkynyl group may be

unsubstituted or substituted. When substituted, the substituent(s) may be selected from the same groups disclosed above with regard to alkyl group substitution. The alkynyl group may have 2 to 20 carbon atoms, although the present definition also covers the occurrence of the term “alkynyl” where no numerical range is designated. The alkynyl group may also be a medium size alkynyl having 2 to 9 carbon atoms. The alkynyl group could also be a lower alkynyl having 2 to 4 carbon atoms. The alkynyl group of the compounds may be designated as “C2-C4 alkynyl” or similar designations. By way of example only, “C2-C4 alkynyl” indicates that there are two to four carbon atoms in the alkynyl chain, i.e., the alkynyl chain is selected from the group consisting of ethynyl, propyn-1-yl, propyn-2-yl, butyn-1-yl, butyn-3-yl, butyn-4-yl, and 2-butynyl. Exemplary alkynyl groups include, but are in no way limited to, ethynyl, propynyl, butynyl, pentynyl, and hexynyl, and the like.

[00042] As used herein, “heteroalkyl” refers to a straight or branched hydrocarbon chain containing one or more heteroatoms, that is, an element other than carbon, including but not limited to, nitrogen, oxygen and sulfur, in the chain backbone. The heteroalkyl group may have 1 to 20 carbon atoms although the present definition also covers the occurrence of the term “heteroalkyl” where no numerical range is designated. The heteroalkyl group may also be a medium size heteroalkyl having 1 to 9 carbon atoms. The heteroalkyl group could also be a lower heteroalkyl having 1 to 4 carbon atoms. The heteroalkyl group of the compounds may be designated as “C1-C4 heteroalkyl” or similar designations. The heteroalkyl group may contain one or more heteroatoms. By way of example only, “C1-C4 heteroalkyl” indicates that there are one to four carbon atoms in the heteroalkyl chain and additionally one or more heteroatoms in the backbone of the chain.

[00043] As used herein, “aryl” refers to a carbocyclic (all carbon) ring or two or more fused rings (rings that share two adjacent carbon atoms) that have a fully delocalized pi-electron system. Examples of aryl groups include, but are not limited to, benzene, naphthalene and azulene. An aryl group may be substituted or unsubstituted. When substituted, hydrogen atoms are replaced by substituent group(s) that is(are) one or more group(s) independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl, heteroalicycyl, aralkyl, heteroaralkyl, (heteroalicycyl)alkyl, hydroxy, protected hydroxyl, alkoxy, aryloxy, acyl, ester, mercapto, alkylthio, arylthio, cyano, halogen, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, protected C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, silyl, sulfenyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, and

amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. When substituted, substituents on an aryl group may form a non-aromatic ring fused to the aryl group, including a cycloalkyl, cycloalkenyl, cycloalkynyl, and heterocyclyl.

[00044] As used herein, “heteroaryl” refers to a monocyclic or multicyclic aromatic ring system (a ring system with fully delocalized pi-electron system), one or two or more fused rings that contain(s) one or more heteroatoms, that is, an element other than carbon, including but not limited to, nitrogen, oxygen and sulfur. Examples of heteroaryl rings include, but are not limited to, furan, thiophene, phthalazine, pyrrole, oxazole, thiazole, imidazole, pyrazole, isoxazole, isothiazole, triazole, thiadiazole, pyridine, pyridazine, pyrimidine, pyrazine and triazine. A heteroaryl group may be substituted or unsubstituted. When substituted, hydrogen atoms are replaced by substituent group(s) that is(are) one or more group(s) independently selected from alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl, heteroalicyclyl, aralkyl, heteroaralkyl, (heteroalicyclyl)alkyl, hydroxy, protected hydroxyl, alkoxy, aryloxy, acyl, ester, mercapto, alkylthio, arylthio, cyano, halogen, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, protected C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, silyl, sulfenyl, sulfinyl, sulfonyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, and amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. When substituted, substituents on a heteroaryl group may form a non-aromatic ring fused to the aryl group, including a cycloalkyl, cycloalkenyl, cycloalkynyl, and heterocyclyl.

[00045] As used herein, an “aralkyl” or “arylalkyl” refers to an aryl group connected, as a substituent, via an alkylene group. The alkylene and aryl group of an aralkyl may be substituted or unsubstituted. Examples include but are not limited to benzyl, substituted benzyl, 2-phenylethyl, 3-phenylpropyl, and naphthylalkyl. In some cases, the alkylene group is a lower alkylene group.

[00046] As used herein, a “heteroaralkyl” or “heteroarylalkyl” is heteroaryl group connected, as a substituent, via an alkylene group. The alkylene and heteroaryl group of heteroaralkyl may be substituted or unsubstituted. Examples include but are not limited to 2-thienylmethyl, 3-thienylmethyl, furylmethyl, thienylethyl, pyrrolylalkyl, pyridylalkyl, isoxazolylalkyl, and imidazolylalkyl, and their substituted as well as benzo-fused analogs. In some cases, the alkylene group is a lower alkylene group.

[00047] As used herein, a “alkylene” refers to a branched, or straight chain fully saturated di-radical chemical group containing only carbon and hydrogen that is attached to

the rest of the molecule via two points of attachment (i.e., an alkanediyl). The alkylene group may have 1 to 20 carbon atoms, although the present definition also covers the occurrence of the term alkylene where no numerical range is designated. The alkylene group may also be a medium size alkylene having 1 to 9 carbon atoms. The alkylene group could also be a lower alkylene having 1 to 4 carbon atoms. The alkylene group may be designated as “C1-C4 alkylene” or similar designations. By way of example only, “C1-C4 alkylene” indicates that there are one to four carbon atoms in the alkylene chain, i.e., the alkylene chain is selected from the group consisting of methylene, ethylene, ethan-1,1-diyl, propylene, propan- 1,1-diyl, propan-2, 2-diyl, 1 -methyl-ethylene, butylene, butan- 1,1-diyl, butan-2,2-diyl, 2-methyl-propan- 1,1-diyl, 1 -methyl-propylene, 2-methyl-propylene, 1,1 -dimethyl-ethylene, 1,2-dimethyl-ethylene, and 1-ethyl-ethylene.

[00048] As used herein, “alkenylene” refers to a straight or branched chain di radical chemical group containing only carbon and hydrogen and containing at least one carbon-carbon double bond that is attached to the rest of the molecule via two points of attachment. The alkenylene group may have 2 to 20 carbon atoms, although the present definition also covers the occurrence of the term alkenylene where no numerical range is designated. The alkenylene group may also be a medium size alkenylene having 2 to 9 carbon atoms. The alkenylene group could also be a lower alkenylene having 2 to 4 carbon atoms. The alkenylene group may be designated as “C2-C4 alkenylene” or similar designations. By way of example only, “C2 alkenylene” indicates that there are two to four carbon atoms in the alkenylene chain, i.e., the alkenylene chain is selected from the group consisting of ethenylene, ethen-1,1-diyl, propenylene, propen- 1,1-diyl, prop-2-en-1,1-diyl, 1-methyl-ethenylene, but-1-enylene, but-2-enylene, but-1,3-dienylene, buten- 1,1-diyl, but-1,3-dien-1,1-diyl, but-2-en- 1,1-diyl, but-3-en- 1,1-diyl, 1-methyl-prop-2-en- 1,1-diyl, 2-methyl-prop-2-en- 1,1-diyl, 1-ethyl-ethenylene, 1,2-dimethyl-ethenylene, 1-methyl-propenylene, 2-methyl-propenylene, 3-methyl-propenylene, 2-methyl-propen- 1,1-diyl, and 2, 2-dimethyl-ethen- 1,1-diyl.

[00049] As used herein, “alkylidene” refers to a divalent group, such as =CR'R”, which is attached to one carbon of another group, forming a double bond, alkylidene groups include, but are not limited to, methylidene (=CH₂) and ethylidene (=CHCH₃). As used herein, “arylalkylidene” refers to an alkylidene group in which either R' and R' ' is an aryl group. An alkylidene group may be substituted or unsubstituted.

[00050] As used herein, “alkoxy” refers to the formula -OR wherein R is an alkyl is defined as above, e.g. methoxy, ethoxy, n-propoxy, 1-methylethoxy (isopropoxy), n- butoxy,

iso-butoxy, sec-butoxy, tert-butoxy, amoxy, tert-amoxy and the like. An alkoxy may be substituted or unsubstituted.

[00051] As used herein, “alkylthio” refers to the formula -SR wherein R is an alkyl is defined as above, e.g. methylmercapto, ethylmercapto, n-propylmercapto, 1-methylethylmercapto (isopropylmercapto), n-butylmercapto, iso-butylmercapto, sec-butylmercapto, tert-butylmercapto, and the like. An alkylthio may be substituted or unsubstituted.

[00052] As used herein, “aryloxy” and “arylthio” refers to RO- and RS-, respectively, in which R is an aryl, such as but not limited to phenyl. Both an aryloxy and arylthio may be substituted or unsubstituted.

[00053] As used herein, “acyl” refers to -C(=O)R, wherein R is hydrogen, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, C3-C8 carbocyclyl, aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein. Non-limiting examples include formyl, acetyl, propanoyl, benzoyl, and acryl.

[00054] As used herein, “cycloalkyl” refers to a completely saturated (no double bonds) mono- or multi- cyclic hydrocarbon ring system. When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro-connected fashion. Cycloalkyl groups may range from C3 to C10, in other embodiments it may range from C3 to C6. A cycloalkyl group may be unsubstituted or substituted. Exemplary cycloalkyl groups include, but are in no way limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like. If substituted, the substituent(s) may be an alkyl or selected from those indicated above with regard to substitution of an alkyl group unless otherwise indicated. When substituted, substituents on a cycloalkyl group may form an aromatic ring fused to the cycloalkyl group, including an aryl and a heteroaryl.

[00055] As used herein, “cycloalkenyl” refers to a cycloalkyl group that contains one or more double bonds in the ring although, if there is more than one, they cannot form a fully delocalized pi-electron system in the ring (otherwise the group would be “aryl,” as defined herein). When composed of two or more rings, the rings may be connected together in a fused, bridged or spiro-connected fashion. A cycloalkenyl group may be unsubstituted or substituted. When substituted, the substituent(s) may be an alkyl or selected from the groups disclosed above with regard to alkyl group substitution unless otherwise indicated. When substituted, substituents on a cycloalkenyl group may form an aromatic ring fused to the cycloalkenyl group, including an aryl and a heteroaryl.

[00056] As used herein, “cycloalkynyl” refers to a cycloalkyl group that contains one or more triple bonds in the ring. When composed of two or more rings, the rings may be joined together in a fused, bridged or spiro-connected fashion. A cycloalkynyl group may be unsubstituted or substituted. When substituted, the substituent(s) may be an alkyl or selected from the groups disclosed above with regard to alkyl group substitution unless otherwise indicated. When substituted, substituents on a cycloalkynyl group may form an aromatic ring fused to the cycloalkynyl group, including an aryl and a heteroaryl.

[00057] As used herein, “heteroalicyclic” or “heteroalicyclyl” refers to a stable 3- to 18 membered ring which consists of carbon atoms and from one to five heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. The “heteroalicyclic” or “heteroalicyclyl” may be monocyclic, bicyclic, tricyclic, or tetracyclic ring system, which may be joined together in a fused, bridged or spiro-connected fashion; and the nitrogen, carbon and sulfur atoms in the “heteroalicyclic” or “heteroalicyclyl” may be optionally oxidized; the nitrogen may be optionally quaternized; and the rings may also contain one or more double bonds provided that they do not form a fully delocalized pi-electron system throughout all the rings. Heteroalicyclyl groups may be unsubstituted or substituted. When substituted, the substituent(s) may be one or more groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl, heteroalicyclyl, aralkyl, heteroaralkyl, (heteroalicyclyl)alkyl, hydroxy, protected hydroxyl, alkoxy, aryloxy, acyl, ester, mercapto, alkylthio, arylthio, cyano, halogen, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, protected C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, silyl, haloalkyl, haloalkoxy, trihalomethanesulfonyl, trihalomethanesulfonamido, and amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. Examples of such “heteroalicyclic” or “heteroalicyclyl” include but are not limited to, azepinyl, acridinyl, carbazolyl, cinnolinyl, dioxolanyl, imidazolyl, morpholinyl, oxiranyl, piperidinyl A-oxide, piperidinyl, piperazinyl, pyrrolidinyl, 4-piperidonyl, pyrazolidinyl, 2-oxopyrrolidinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, and thiamorpholinyl sulfone. When substituted, substituents on a heteroalicyclyl group may form an aromatic ring fused to the heteroalicyclyl group, including an aryl and a heteroaryl.

[00058] As used herein, the term “(cycloalkenyl)alkyl” refers to a cycloalkenyl group connected, as a substituent, via an alkylene group. The alkylene and cycloalkenyl of a

(cycloalkenyl)alkyl may be substituted or unsubstituted. In some cases, the alkylene group is a lower alkylene group.

[00059] As used herein, the term “(cycloalkynyl)alkyl” to a cycloalkynyl group connected, as a substituent, via an alkylene group. The alkylene and cycloalkynyl of a (cycloalkynyl)alkyl may be substituted or unsubstituted. In some cases, the alkylene group is a lower alkylene group.

[00060] As used herein, the term “O-carboxy” refers to a “RC(=O)O-” group in which R can be hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl, heteroalicyclic, aralkyl, or (heteroalicyclic)alkyl, as defined herein. An O-carboxy may be substituted or unsubstituted.

[00061] As used herein, the term “C-carboxy” refers to a “-C(=O)R” group in which R can be the same as defined with respect to O-carboxy. A C-carboxy may be substituted or unsubstituted.

[00062] As used herein, the term “trihalomethanesulfonyl” refers to an “X₃CSO₂-” group wherein X is a halogen.

[00063] As used herein, the term “cyano” refers to a “-CN” group.

[00064] As used herein, the term “cyanato” refers to an “-OCN” group.

[00065] As used herein, the term “isocyanato” refers to a “-NCO” group.

[00066] As used herein, the term “thiocyanato” refers to a “-SCN” group.

[00067] As used herein, the term “isothiocyanato” refers to an “-NCS” group.

[00068] As used herein, the term “sulfinyl” refers to a “-S(=O)-R” group in which R can be the same as defined with respect to O-carboxy. A sulfinyl may be substituted or unsubstituted.

[00069] As used herein, the term “sulfonyl” refers to an “-SO₂R” group in which R can be the same as defined with respect to O-carboxy. A sulfonyl may be substituted or unsubstituted.

[00070] As used herein, the term “S-sulfonamido” refers to a “-SO₂NRARB” group in which RA and RB can be the same as defined with respect to O-carboxy. An S-sulfonamido may be substituted or unsubstituted.

[00071] As used herein, the term “N-sulfonamido” refers to a “-SO₂N(RA)(RB)” group in which RA and RB can be the same as defined with respect to O-carboxy. A sulfonyl may be substituted or unsubstituted.

[00072] As used herein, the term “trihalomethanesulfonamido” refers to an “X₃CSO₂N(R)-“ group with X as halogen and R can be the same as defined with respect to O-carboxy. A trihalomethanesulfonamido may be substituted or unsubstituted.

[00073] As used herein, the term “O-carbamyl” refers to a “-OC(=O)NRARB” group in which RA and RB can be the same as defined with respect to O-carboxy. An O-carbamyl may be substituted or unsubstituted.

[00074] As used herein, the term “N-carbamyl” refers to an “ROC(=O)NRA group in which R and RA can be the same as defined with respect to O-carboxy. An N-carbamyl may be substituted or unsubstituted.

[00075] As used herein, the term “O-thiocarbamyl” refers to a “-OC(=S)-NRARB” group in which RA and RB can be the same as defined with respect to O-carboxy. An O-thiocarbamyl may be substituted or unsubstituted.

[00076] As used herein, the term “N-thiocarbamyl” refers to an “ROC(=S)NRA-” group in which R and RA can be the same as defined with respect to O-carboxy. An N-thiocarbamyl may be substituted or unsubstituted.

[00077] As used herein, the term “C-amido” refers to a “-C(=O)NRARB” group in which RA and RB can be the same as defined with respect to O-carboxy. A C-amido may be substituted or unsubstituted.

[00078] As used herein, the term “N-amido” refers to a “RC(=O)NRA-“ group in which R and RA can be the same as defined with respect to O-carboxy. An N-amido may be substituted or unsubstituted.

[00079] As used herein, the term “amino” refers to a “-NRARB” group in which RA and RB are each independently selected from hydrogen, C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, C3-C8 carbocyclyl, C6-C10 aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

[00080] As used herein, the term “aminoalkyl” refers to an amino group connected via an alkylene group.

[00081] As used herein, the term “ester” refers to a “-C(=O)OR” group in which R can be the same as defined with respect to O-carboxy. An ester may be substituted or unsubstituted.

[00082] As used herein, the term “lower aminoalkyl” refers to an amino group connected via a lower alkylene group. A lower aminoalkyl may be substituted or unsubstituted.

[00083] As used herein, the term “lower alkoxyalkyl” refers to an alkoxy group connected via a lower alkylene group. A lower alkoxyalkyl may be substituted or unsubstituted.

[00084] As used herein, the term “acetyl” refers to a $-C(=O)CH_3$ group.

[00085] As used herein, the term “perhaloalkyl” refers to an alkyl group where all of the hydrogen atoms are replaced by halogen atoms.

[00086] As used herein, the term “carbocyclyl” refers to a non-aromatic cyclic ring or ring system containing only carbon atoms in the ring system backbone. When the carbocyclyl is a ring system, two or more rings may be joined together in a fused, bridged or spiro-connected fashion. Carbocyclyls may have any degree of saturation provided that at least one ring in a ring system is not aromatic. Thus, carbocyclyls include cycloalkyls, cycloalkenyls, and cycloalkynyls. The carbocyclyl group may have 3 to 20 carbon atoms, although the present definition also covers the occurrence of the term “carbocyclyl” where no numerical range is designated. The carbocyclyl group may also be a medium size carbocyclyl having 3 to 10 carbon atoms. The carbocyclyl group could also be a carbocyclyl having 3 to 6 carbon atoms. The carbocyclyl group may be designated as “C3-C6 carbocyclyl” or similar designations. Examples of carbocyclyl rings include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, 2,3-dihydro-indene, bicycle[2.2.2]octanyl, adamantyl, and spiro[4.4]nonanyl.

[00087] As used herein, the term “(cycloalkyl)alkyl” refers to a cycloalkyl group connected, as a substituent, via an alkylene group. The alkylene and cycloalkyl of a (cycloalkyl)alkyl may be substituted or unsubstituted. Examples include but are not limited cyclopropylmethyl, cyclobutylmethyl, cyclopropylethyl, cyclopropylbutyl, cyclobutylethyl, cyclopropylisopropyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cycloheptylmethyl, and the like. In some cases, the alkylene group is a lower alkylene group.

[00088] As used herein, the term “cycloalkyl” refers to a fully saturated carbocyclyl ring or ring system. Examples include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

[00089] As used herein, the term “cycloalkenyl” means a carbocyclyl ring or ring system having at least one double bond, wherein no ring in the ring system is aromatic. An example is cyclohexenyl.

[00090] As used herein, the term “heterocyclyl” refers to three-, four-, five-, six-, seven-, and eight- or more membered rings wherein carbon atoms together with from 1 to 3 heteroatoms constitute said ring. A heterocyclyl can optionally contain one or more

unsaturated bonds situated in such a way, however, that an aromatic pi-electron system does not arise. The heteroatoms are independently selected from oxygen, sulfur, and nitrogen. A heterocyclcyl can further contain one or more carbonyl or thiocarbonyl functionalities, so as to make the definition include oxo- systems and thio- systems such as lactams, lactones, cyclic imides, cyclic thioimides, cyclic carbamates, and the like. A “heterocyclcyl” can refer to a non-aromatic cyclic ring or ring system containing at least one heteroatom in the ring backbone. Heterocyclcyls may be joined together in a fused, bridged or spiro-connected fashion. Heterocyclcyls may have any degree of saturation provided that at least one ring in the ring system is not aromatic. The heteroatom(s) may be present in either a non-aromatic or aromatic ring in the ring system. The heterocyclcyl group may have 3 to 20 ring members (i.e., the number of atoms making up the ring backbone, including carbon atoms and heteroatoms), although the present definition also covers the occurrence of the term “heterocyclcyl” where no numerical range is designated. The heterocyclcyl group may also be a medium size heterocyclcyl having 3 to 10 ring members. The heterocyclcyl group could also be a heterocyclcyl having 3 to 6 ring members. The heterocyclcyl group may be designated as “3-6 membered heterocyclcyl” or similar designations. In preferred six membered monocyclic heterocyclcyls, the heteroatom(s) are selected from one up to three of O, N or S, and in preferred five membered monocyclic heterocyclcyls, the heteroatom(s) are selected from one or two heteroatoms selected from O, N, or S. Examples of heterocyclcyl rings include, but are not limited to, azepinyl, acridinyl, carbazolyl, cinnolinyl, dioxolanyl, imidazolinyl, imidazolidinyl, morpholinyl, oxiranyl, oxepanyl, thiepanyl, piperidinyl, piperazinyl, dioxopiperazinyl, pyrrolidinyl, pyrrolidonyl, pyrrolidionyl, 4-piperidonyl, pyrazolinyl, pyrazolidinyl, 1,3-dioxinyl, 1,3-dioxanyl, 1,4- dioxinyl, 1,4-dioxanyl, 1,3-oxathianyl, 1,4-oxathiinyl, 1,4-oxathianyl, 2//-1,2-oxazinyl, trioxanyl, hexahydro-1,3,5-triazinyl, 1,3-dioxolyl, 1,3-dioxolanyl, 1,3-dithiolyl, 1,3- dithiolanyl, isoxazolinyl, isoxazolidinyl, oxazolinyl, oxazolidinyl, oxazolidinonyl, thiazolinyl, thiazolidinyl, 1,3-oxathiolanyl, indolinyl, isoindolinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, tetrahydro-1,4-thiazinyl, thiamorpholinyl, dihydrobenzofuranyl, benzimidazolidinyl, and tetrahydroquinoline.

[00091] As used herein, the term “(heterocyclcyl)alkyl” refers to a heterocyclcyl group connected, as a substituent, via an alkylene group. Examples include, but are not limited to, imidazolinylmethyl and indolinylethyl.

[00092] Substituted groups are based upon or derived from the unsubstituted parent group in which there has been an exchange of one or more hydrogen atoms for another atom

or group. Unless otherwise indicated, when a group is deemed to be “substituted,” the group is substituted with one or more substituents independently selected from C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, C1-C6 heteroalkyl, C3-C7 carbocyclyl (optionally substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 haloalkyl, and C1-C6 haloalkoxy), C3-C7-carbocyclyl-C1-C6-alkyl (optionally substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 haloalkyl, and C1-C6 haloalkoxy), 5-10 membered heterocyclyl (optionally substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 haloalkyl, and C1-C6 haloalkoxy), 5-10 membered heterocyclyl-C1-C6-alkyl (optionally substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 haloalkyl, and C1-C6 haloalkoxy), aryl (optionally substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 haloalkyl, and C1-C6 haloalkoxy), aryl(C1-C6)alkyl (optionally substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 haloalkyl, and C1-C6 haloalkoxy), 5-10 membered heteroaryl (optionally substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 haloalkyl, and C1-C6 haloalkoxy), 5-10 membered heteroaryl(C1-C6)alkyl (optionally substituted with halo, C1-C6 alkyl, C1-C6 alkoxy, C1-C6 haloalkyl, and C1-C6 haloalkoxy), halo, cyano, hydroxy, C1-C6 alkoxy, C1-C6 alkoxy(C1-C6)alkyl (i.e., ether), aryloxy, sulfhydryl (mercapto), halo(C1-C6)alkyl (e.g., -CF₃), halo(C1-C6)alkoxy (e.g., -OCF₃), C1-C6 alkylthio, arylthio, amino, amino(C1-C6)alkyl, nitro, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, acyl, cyanato, isocyanato, thiocyanato, isothiocyanato, sulfinyl, sulfonyl, and oxo (=O). Wherever a group is described as “optionally substituted” that group can be substituted with the above substituents.

[00093] In some embodiments, a substituted group is substituted with one or more substituent(s) individually and independently selected from C1-C4 alkyl, amino, hydroxy, and halogen.

[00094] It is to be understood that certain radical naming conventions can include either a mono-radical or a di-radical, depending on the context. For example, where a substituent requires two points of attachment to the rest of the molecule, it is understood that the substituent is a di-radical. For example, a substituent identified as alkyl that requires two points of attachment includes di-radicals such as -CH₂-, -CH₂CH₂-, -CH₂CH(CH₃)CH₂-, and the like. Other radical naming conventions clearly indicate that the radical is a di-radical such as “alkylene” or “alkenylene.”

[00095] Unless otherwise indicated, when a substituent is deemed to be “optionally substituted,” it is meant that the substituent is a group that may be substituted with one or more group(s) individually and independently selected from alkyl, alkenyl, alkynyl,

cycloalkyl, aryl, heteroaryl, heteroalicyclic, hydroxyl, alkoxy, aryloxy, mercapto, alkylthio, arylthio, cyano, halo, carbonyl, thiocarbonyl, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, isocyanato, thiocyanato, isothiocyanato, nitro, silyl, trihalomethanesulfonyl, and amino, including mono- and di-substituted amino groups, and the protected derivatives thereof. The protecting groups that may form the protective derivatives of the above substituents are known to those of skill in the art and may be found in references such as Greene and Wuts, above.

Other definitions

[00096] "About" and "approximately" shall generally mean an acceptable degree of error for the quantity measured given the nature or precision of the measurements. Exemplary degrees of error are within 20 percent (%), typically, within 10%, and more typically, within 5% of a given value or range of values.

[00097] As used herein, "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, Berge *et al.*, describes pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences* (1977) 66:1–19. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate,

stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Pharmaceutically acceptable salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, lower alkyl sulfonate, and aryl sulfonate.

[00098] As used herein, “pharmaceutically acceptable carrier” refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions described herein include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[00099] As used herein, a “subject” to which administration is contemplated includes, but is not limited to, humans (i.e., a male or female of any age group, e.g., a pediatric subject (e.g., infant, child, adolescent) or adult subject (e.g., young adult, middle-aged adult or senior adult)) and/or a non-human animal, e.g., a mammal such as primates (e.g., cynomolgus monkeys, rhesus monkeys), cattle, pigs, horses, sheep, goats, rodents, cats, and/or dogs. In some embodiments, the subject is a human. In some embodiments, the subject is a non-human animal. The terms “human,” “patient,” and “subject” are used interchangeably herein.

[000100] Disease, disorder, and condition are used interchangeably herein.

[000101] As used herein, and unless otherwise specified, the terms “treat,” “treating” and “treatment” contemplate an action that occurs while a subject is suffering from the specified disease, disorder or condition, which reduces the severity of the disease, disorder or condition, or retards or slows the progression of the disease, disorder or condition (also “therapeutic treatment”).

[000102] In general, the “effective amount” of a compound refers to an amount sufficient to elicit the desired biological response. As will be appreciated by those of

ordinary skill in this art, the effective amount of a compound of the invention may vary depending on such factors as the desired biological endpoint, the pharmacokinetics of the compound, the disease being treated, the mode of administration, and the age, weight, health, and condition of the subject. As used herein, and unless otherwise specified, a “therapeutically effective amount” of a compound is an amount sufficient to provide a therapeutic benefit in the treatment of a disease, disorder or condition, or to delay or minimize one or more symptoms associated with the disease, disorder or condition. A therapeutically effective amount of a compound means an amount of therapeutic agent, alone or in combination with other therapies, which provides a therapeutic benefit in the treatment of the disease, disorder or condition. The term “therapeutically effective amount” can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of disease or condition, or enhances the therapeutic efficacy of another therapeutic agent. As used herein, and unless otherwise specified, a “prophylactically effective amount” of a compound is an amount sufficient to prevent a disease, disorder or condition, or one or more symptoms associated with the disease, disorder or condition, or prevent its recurrence. A prophylactically effective amount of a compound means an amount of a therapeutic agent, alone or in combination with other agents, which provides a prophylactic benefit in the prevention of the disease, disorder or condition. The term “prophylactically effective amount” can encompass an amount that improves overall prophylaxis or enhances the prophylactic efficacy of another prophylactic agent.

[000103] As used herein, “prophylactic treatment” contemplates an action that occurs before a subject begins to suffer from the specified disease, disorder or condition.

[000104] The term, “oral dosage form,” as used herein, refers to a composition or medium used to administer an agent to a subject. Typically, an oral dosage form is administered via the mouth, however, “oral dosage form” is intended to cover any substance which is administered to a subject and is absorbed across a membrane, e.g., a mucosal membrane, of the gastrointestinal tract, including, e.g., the mouth, esophagus, stomach, small intestine, large intestine, and colon. For example, “oral dosage form” covers a solution which is administered through a feeding tube into the stomach.

[000105] A “cycle”, as used herein in the context of a cycle of administration of a drug, refers to a period of time for which a drug is administered and may further include a rest period of not administering the drug to a subject. In some embodiments, one cycle is four weeks.

[000106] A “RAS mutation” is a mutation in the RAS gene. For example, a “KRAS mutation” is a mutation of the KRAS gene (i.e., a nucleic acid mutation) or Kras protein (i.e., an amino acid mutation) that results in aberrant Kras protein function associated with increased and/or constitutive activity by favoring the active GTP-bound state of the Kras protein. The mutation may be at conserved sites that favor GTP binding and constitutively active Kras protein. In some instances, the mutation is at one or more of codons 12, 13, and 16 of the KRAS gene. For example, a KRAS mutation may be at codon 12 of the KRAS gene, for instance, as a single point substitution mutation at codon 12 (i.e., KRAS G12X mutation) (e.g., a KRAS G12V mutation arises from a single nucleotide change (c.35G>T) and results in an amino acid substitution of the glycine (G) at position 12 by a valine (V)). Exemplary KRAS G12X mutations include, but are not limited to, KRAS G12V, KRAS G12D, KRAS G12A, KRAS G12R, KRAS G12S, and KRAS G12C. As another example, an “NRAS mutation” is a mutation of the NRAS gene (e.g., a nucleic acid mutation) or Nras protein (e.g., an amino acid mutation) that results in aberrant Nras protein function.

[000107] A “RAF mutation” is a mutation in the RAF gene. For example, a “BRAF mutation” is a mutation in the BRAF gene. In some instances, the mutation in the BRAF gene is a BRAF V600 mutation including BRAF V600E, BRAF V600K, BRAF V600D, BRAF V600R, and BRAF V600M mutation. A “CRAF” mutation is a mutation in the CRAF gene, and an “ARAF” mutation is a mutation in the ARAF gene.

Methods of Treatment


[000108] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a BRAF V600 inhibitor.

[000109] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a BRAF V600 inhibitor, wherein the cancer is identified as having a BRAF V600 mutation.

[000110] In some embodiments, the BRAF V600 mutation is BRAF V600E, BRAF V600K, BRAF V600D, BRAF V600R, and/or BRAF V600M mutation. In some embodiments, the BRAF V600 mutation is BRAF V600E mutation. In some embodiments, the BRAF V600 mutation is BRAF V600K mutation. In some embodiments, the BRAF V600 mutation is BRAF V600D mutation. In some embodiments, the BRAF V600 mutation is


BRAF V600R mutation. In some embodiments, the BRAF V600 mutation is BRAF V600M mutation.

[000111] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (I):



(I), or a pharmaceutically acceptable salt thereof.

[000112] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula




(I):

(I).

[000113] In some embodiments, the dual RAF/MEK inhibitor is a potassium salt of the compound of formula (I).

[000114] In some embodiments, the dual RAF/MEK inhibitor is IMM-1-104, or a pharmaceutically acceptable salt thereof.

[000115] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (II):



(II)

including pharmaceutically acceptable salts thereof, wherein the variables are as defined herein.

[000116] In some embodiments, the dual RAF/MEK inhibitor is a compound selected from the compound of Table I, or a pharmaceutically acceptable salt thereof.

[000117] In some embodiments, the BRAF V600 inhibitor is dabrafenib, encorafenib, vemurafenib, FORE-8394 (PLX-8394), tinloragenib, AZ-304, agerafenib, KIN-2787, BGB-3245, ABM-1310, TQB-3233, UB-941, AFX-1251, ARQ 736, ASN003, AVB-BRAF, BDTX-4933, CFT1946, HLX208, RO5212054, RO7276389, or TQ-B3233, or pharmaceutically acceptable salts thereof. In some embodiments, the BRAF V600 inhibitor

is dabrafenib, encorafenib, vemurafenib, FORE-8394, tinloragenib, AZ-304, agerafenib, or KIN-2787, or pharmaceutically acceptable salts thereof. In some embodiments, the BRAF V600 inhibitor is dabrafenib, encorafenib, or vemurafenib, or pharmaceutically acceptable salts thereof. In some embodiments, the BRAF V600 inhibitor is dabrafenib, or a pharmaceutically acceptable salt thereof. In some embodiments, the BRAF V600 inhibitor is encorafenib, or a pharmaceutically acceptable salt thereof. In some embodiments, the BRAF V600 inhibitor is vemurafenib, or a pharmaceutically acceptable salt thereof.

[000118] In some embodiments, the BRAF V600 inhibitor is orally administered to the subject. In some embodiments, the BRAF V600 inhibitor is administered once a week. In some embodiments, the BRAF V600 inhibitor is administered twice a week. In some embodiments, the BRAF V600 inhibitor is administered thrice a week. In some embodiments, the BRAF V600 inhibitor is administered four times a week. In some embodiments, the BRAF V600 inhibitor is administered five times a week. In some embodiments, the BRAF V600 inhibitor is administered six times a week. In some embodiments, the BRAF V600 inhibitor is administered once daily. In some embodiments, the BRAF V600 inhibitor is administered twice daily. In some embodiments, the BRAF V600 inhibitor is dosed at 10 mg to 2000 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at 100 mg to 1000 mg per administration.

[000119] In some embodiments, the BRAF V600 inhibitor is dosed as a cycle comprising administering the BRAF V600 inhibitor for three weeks and then not administering the BRAF V600 inhibitor for one week.

[000120] In some embodiments, the dual RAF/MEK inhibitor and the BRAF V600 inhibitor are independently dosed cyclically.

[000121] In some embodiments, the dual RAF/MEK inhibitor and the BRAF V600 inhibitor are simultaneously dosed cyclically.

[000122] In some embodiments, the methods described herein further comprises administering to the subject an effective amount of a FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof). In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at about 100 mg to about 1000 mg. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at about 100 mg to about 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at 200 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at 400 mg per

administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is administered twice daily. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed as a cycle, comprising administering the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) for three weeks and then not administering the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) for one week. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is administered orally to the subject.

[000123] In some embodiments, the dual RAF/MEK inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000124] In some embodiments, the dual RAF/MEK inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000125] In some embodiments, the BRAF V600 inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000126] In some embodiments, the BRAF V600 inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000127] In some embodiments, the cancer is pancreatic cancer, gynecologic cancer (e.g., cervical cancer, ovarian cancer, uterine cancer, vaginal cancer, endometrial cancer, or vulvar cancer), liver cancer, prostate cancer, mesothelioma, breast cancer, bladder cancer, melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the cancer is melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the lung cancer is non-small cell lung cancer. In some embodiments, the lung cancer is metastatic non-small cell lung cancer. In some embodiments, the melanoma is unresectable melanoma. In some embodiments, the melanoma is metastatic melanoma. In some embodiments, the cancer is colorectal cancer. In some embodiments, the thyroid cancer is papillary thyroid cancer. In some embodiments, the thyroid cancer is follicular thyroid cancer. In some embodiments, the thyroid cancer is anaplastic thyroid cancer.

[000128] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of defactinib or a pharmaceutically acceptable salt thereof and an effective amount of a BRAF V600 inhibitor, wherein the cancer is identified as having a BRAF V600 mutation.


[000129] In some embodiments, the BRAF V600 inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000130] In some embodiments, the BRAF V600 inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000131] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a pan-RAF inhibitor.


[000132] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a pan-RAF inhibitor, wherein the cancer is identified as having ARAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation. In some embodiments, the cancer is identified as having ARAF, BRAF, and/or CRAF mutation. In some embodiments, the cancer is identified as having KRAS, NRAS, and/or HRAS mutation. In some embodiments, the cancer is identified as having EGFR, ALKR, FGFR, PDGFR, and/or NF1 mutation. In some embodiments, the cancer is identified as having EGFR and/or FGFR, mutation. In some embodiments, the cancer is identified as having ALKR mutation. In some embodiments, the cancer is identified as having NF1 mutation. In some embodiments, the cancer is identified as having SOS1 and/or SOS2 mutation. In some embodiments, the cancer is identified as having MEK1, and/or MEK2 mutation.

[000133] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (I):



(I), or a pharmaceutically acceptable salt thereof.

[000134] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula




(I):

(I).

[000135] In some embodiments, the dual RAF/MEK inhibitor is a potassium salt of the compound of formula (I).

[000136] In some embodiments, the dual RAF/MEK inhibitor is IMM-1-104, or a pharmaceutically acceptable salt thereof.

[000137] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (II):



including pharmaceutically acceptable salts thereof, wherein the variables are as defined herein.

[000138] In some embodiments, the dual RAF/MEK inhibitor is a compound selected from the compound of Table I, or a pharmaceutically acceptable salt thereof.

[000139] In some embodiments, the pan-RAF inhibitor is belvarafenib, naporafenib, lifirafenib, tovorafenib, BAL3833, LY3009120, REDX05358, IRICoR-Ipsen, JZP815, METiS-01, QLH11906, or SJ-C1044, or pharmaceutically acceptable salts thereof. In some embodiments, the pan-RAF inhibitor is belvarafenib, naporafenib, lifirafenib, tovorafenib, BAL3833, LY3009120, or REDX05358 or pharmaceutically acceptable salts thereof.

[000140] In some embodiments, the pan-RAF inhibitor is orally administered to the subject. In some embodiments, the pan-RAF inhibitor is administered once a week. In some embodiments, the pan-RAF inhibitor is administered twice a week. In some embodiments, the pan-RAF inhibitor is administered thrice a week. In some embodiments, the pan-RAF inhibitor is administered four times a week. In some embodiments, the pan-RAF inhibitor is administered five times a week. In some embodiments, the pan-RAF inhibitor is administered six times a week. In some embodiments, the pan-RAF inhibitor is administered once daily. In some embodiments, the pan-RAF inhibitor is administered twice daily. In some embodiments, the pan-RAF inhibitor is dosed at 10 mg to 2000 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at 100 mg to 1000 mg per administration.

[000141] In some embodiments, the pan-RAF inhibitor is dosed as a cycle comprising administering the pan-RAF inhibitor for three weeks and then not administering the pan-RAF inhibitor for one week.

[000142] In some embodiments, the dual RAF/MEK inhibitor and the pan-RAF inhibitor are independently dosed cyclically.

[000143] In some embodiments, the dual RAF/MEK inhibitor and the pan-RAF inhibitor are simultaneously dosed cyclically.

[000144] In some embodiments, the methods described herein further comprises administering to the subject an effective amount of a FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof). In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at about 100 mg to about 1000 mg. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at about 100 mg to about 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at 200 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is administered twice daily. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed as a cycle, comprising administering the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) for three weeks and then not administering the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) for one week. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is administered orally to the subject.

[000145] In some embodiments, the dual RAF/MEK inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000146] In some embodiments, the dual RAF/MEK inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000147] In some embodiments, the pan-RAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000148] In some embodiments, the pan-RAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000149] In some embodiments, the cancer is pancreatic cancer, gynecologic cancer (e.g., cervical cancer, ovarian cancer, uterine cancer, vaginal cancer, endometrial cancer, or vulvar cancer), liver cancer, prostate cancer, mesothelioma, breast cancer, bladder cancer, melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the cancer is melanoma, lung cancer, colorectal cancer, thyroid cancer,

glioblastoma, or renal cancer. In some embodiments, the lung cancer is non-small cell lung cancer. In some embodiments, the lung cancer is metastatic non-small cell lung cancer. In some embodiments, the melanoma is unresectable melanoma. In some embodiments, the melanoma is metastatic melanoma. In some embodiments, the cancer is colorectal cancer. In some embodiments, the thyroid cancer is papillary thyroid cancer. In some embodiments, the thyroid cancer is follicular thyroid cancer. In some embodiments, the thyroid cancer is anaplastic thyroid cancer.

[000150] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of defactinib or a pharmaceutically acceptable salt thereof and an effective amount of a pan-RAF inhibitor.


[000151] In some embodiments, the pan-RAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000152] In some embodiments, the pan-RAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000153] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a CRAF inhibitor.


[000154] embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a CRAF inhibitor, wherein the cancer is identified as having ARAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation. In some embodiments, the cancer is identified as having ARAF, BRAF, and/or CRAF mutation. In some embodiments, the cancer is identified as having KRAS, NRAS, and/or HRAS mutation. In some embodiments, the cancer is identified as having EGFR, ALKR, FGFR, PDGFR, and/or NF1 mutation. In some embodiments, the cancer is identified as having EGFR and/or FGFR, mutation. In some embodiments, the cancer is identified as having ALKR mutation. In some embodiments, the cancer is identified as having NF1 mutation. In some embodiments, the cancer is identified as having SOS1 and/or SOS2 mutation. In some embodiments, the cancer is identified as having MEK1, and/or MEK2 mutation.

[000155] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (I):



(I), or a pharmaceutically acceptable salt thereof.

[000156] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula




(I):

(I).

[000157] In some embodiments, the dual RAF/MEK inhibitor is a potassium salt of the compound of formula (I).

[000158] In some embodiments, the dual RAF/MEK inhibitor is IMM-1-104, or a pharmaceutically acceptable salt thereof.

[000159] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (II):



(II)

including pharmaceutically acceptable salts thereof, wherein the variables are as defined herein.

[000160] In some embodiments, the dual RAF/MEK inhibitor is a compound selected from the compound of Table I, or a pharmaceutically acceptable salt thereof.

[000161] In some embodiments, the CRAF inhibitor is MG005, Quanta-RAF1, or STX200, or pharmaceutically acceptable salts thereof.

[000162] In some embodiments, the CRAF inhibitor is orally administered to the subject. In some embodiments, the CRAF inhibitor is administered once a week. In some embodiments, the CRAF inhibitor is administered twice a week. In some embodiments, the CRAF inhibitor is administered thrice a week. In some embodiments, the CRAF inhibitor is administered four times a week. In some embodiments, the CRAF inhibitor is administered five times a week. In some embodiments, the CRAF inhibitor is administered six times a week. In some embodiments, the CRAF inhibitor is administered once daily. In some

embodiments, the CRAF inhibitor is administered twice daily. In some embodiments, the CRAF inhibitor is dosed at 10 mg to 2000 mg per administration. In some embodiments, the CRAF inhibitor is dosed at 100 mg to 1000 mg per administration.

[000163] In some embodiments, the CRAF inhibitor is dosed as a cycle comprising administering the CRAF inhibitor for three weeks and then not administering the CRAF inhibitor for one week.

[000164] In some embodiments, the dual RAF/MEK inhibitor and the CRAF inhibitor are independently dosed cyclically.

[000165] In some embodiments, the dual RAF/MEK inhibitor and the CRAF inhibitor are simultaneously dosed cyclically.

[000166] In some embodiments, the methods described herein further comprises administering to the subject an effective amount of a FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof). In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at about 100 mg to about 1000 mg. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at about 100 mg to about 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at 200 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is administered twice daily. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed as a cycle, comprising administering the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) for three weeks and then not administering the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) for one week. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is administered orally to the subject.

[000167] In some embodiments, the dual RAF/MEK inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000168] In some embodiments, the dual RAF/MEK inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000169] In some embodiments, the CRAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000170] In some embodiments, the CRAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000171] In some embodiments, the cancer is pancreatic cancer, gynecologic cancer (e.g., cervical cancer, ovarian cancer, uterine cancer, vaginal cancer, endometrial cancer, or vulvar cancer), liver cancer, prostate cancer, mesothelioma, breast cancer, bladder cancer, melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the cancer is melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the lung cancer is non-small cell lung cancer. In some embodiments, the lung cancer is metastatic non-small cell lung cancer. In some embodiments, the melanoma is unresectable melanoma. In some embodiments, the melanoma is metastatic melanoma. In some embodiments, the cancer is colorectal cancer. In some embodiments, the thyroid cancer is papillary thyroid cancer. In some embodiments, the thyroid cancer is follicular thyroid cancer. In some embodiments, the thyroid cancer is anaplastic thyroid cancer.

[000172] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of defactinib or a pharmaceutically acceptable salt thereof and an effective amount of a CRAF inhibitor.

[000173] In some embodiments, the CRAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.


[000174] In some embodiments, the CRAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000175] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a RAF inhibitor.

[000176] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a RAF inhibitor, wherein the cancer is identified as having ARAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation. In some embodiments, the cancer is identified as having ARAF, BRAF, and/or CRAF mutation. In some embodiments, the cancer is identified as having KRAS, NRAS, and/or HRAS mutation.


In some embodiments, the cancer is identified as having EGFR, ALKR, FGFR, PDGFR, and/or NF1 mutation. In some embodiments, the cancer is identified as having EGFR and/or FGFR, mutation. In some embodiments, the cancer is identified as having ALKR mutation. In some embodiments, the cancer is identified as having NF1 mutation. In some embodiments, the cancer is identified as having SOS1 and/or SOS2 mutation. In some embodiments, the cancer is identified as having MEK1, and/or MEK2 mutation.

[000177] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (I):



(I), or a pharmaceutically acceptable salt thereof.

[000178] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula




(I):

(I).

[000179] In some embodiments, the dual RAF/MEK inhibitor is a potassium salt of the compound of formula (I).

[000180] In some embodiments, the dual RAF/MEK inhibitor is IMM-1-104, or a pharmaceutically acceptable salt thereof.

[000181] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (II):



including pharmaceutically acceptable salts thereof, wherein the variables are as defined herein.

[000182] In some embodiments, the dual RAF/MEK inhibitor is a compound selected from the compound of Table I, or a pharmaceutically acceptable salt thereof.

[000183] In some embodiments, the RAF inhibitor is donafenib, lifirafenib, rigosertib, BMS-908662, XP-102, ABM-2526, DDC-PanRAF, FNX006, or VRN-XX, or pharmaceutically acceptable salts thereof.

[000184] In some embodiments, the BRAF V600 inhibitor is orally administered to the subject. In some embodiments, the BRAF V600 inhibitor is administered once a week. In some embodiments, the BRAF V600 inhibitor is administered twice a week. In some embodiments, the BRAF V600 inhibitor is administered thrice a week. In some embodiments, the BRAF V600 inhibitor is administered four times a week. In some embodiments, the BRAF V600 inhibitor is administered five times a week. In some embodiments, the BRAF V600 inhibitor is administered six times a week. In some embodiments, the BRAF V600 inhibitor is administered once daily. In some embodiments, the BRAF V600 inhibitor is administered twice daily. In some embodiments, the BRAF V600 inhibitor is dosed at 10 mg to 2000 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at 100 mg to 1000 mg per administration.

[000185] In some embodiments, the RAF inhibitor is dosed as a cycle comprising administering the RAF inhibitor for three weeks and then not administering the RAF inhibitor for one week

[000186] In some embodiments, the dual RAF/MEK inhibitor and the BRAF V600 inhibitor are independently dosed cyclically.

[000187] In some embodiments, the dual RAF/MEK inhibitor and the BRAF V600 inhibitor are simultaneously dosed cyclically.

[000188] In some embodiments, the methods described herein further comprises administering to the subject an effective amount of a FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof). In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at about 100 mg to about 1000 mg. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at about 100 mg to about 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at 200 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed at 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is administered twice daily. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is dosed as a cycle, comprising administering the FAK inhibitor (e.g., defactinib, or a pharmaceutically

acceptable salt thereof) for three weeks and then not administering the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) for one week. In some embodiments, the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) is administered orally to the subject.

[000189] In some embodiments, the dual RAF/MEK inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000190] In some embodiments, the dual RAF/MEK inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000191] In some embodiments, the RAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically.

[000192] In some embodiments, the RAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

[000193] In some embodiments, the cancer is pancreatic cancer, gynecologic cancer (e.g., cervical cancer, ovarian cancer, uterine cancer, vaginal cancer, endometrial cancer, or vulvar cancer), liver cancer, prostate cancer, mesothelioma, breast cancer, bladder cancer, melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the cancer is melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the lung cancer is non-small cell lung cancer. In some embodiments, the lung cancer is metastatic non-small cell lung cancer. In some embodiments, the melanoma is unresectable melanoma. In some embodiments, the melanoma is metastatic melanoma. In some embodiments, the cancer is colorectal cancer. In some embodiments, the thyroid cancer is papillary thyroid cancer. In some embodiments, the thyroid cancer is follicular thyroid cancer. In some embodiments, the thyroid cancer is anaplastic thyroid cancer.


[000194] In some embodiments, provided herein is a method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of defactinib or a pharmaceutically acceptable salt thereof and an effective amount of a RAF inhibitor.

[000195] In some embodiments, the RAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are independently dosed cyclically. In some embodiments, the RAF inhibitor and the FAK inhibitor (e.g., defactinib, or a pharmaceutically acceptable salt thereof) are simultaneously dosed cyclically.

Dual RAF/MEK Inhibitors


[000196] An exemplary dual RAF/MEK inhibitor described herein is VS-6766 (also referred to as CKI27, CH5126766, or RO5126766).

[000197] In some embodiments, the dual RAF/MEK inhibitor is a compound of formula (I):



or a pharmaceutically acceptable salt thereof.


[000198] In some embodiments, the compound of formula (I) is:




which is also referred to herein as Compound 1 or VS-6766 free form.

[000199] In some embodiments, the dual RAF/MEK inhibitor is a pharmaceutically acceptable salt of the compound of formula (I). In some embodiments, the dual RAF/MEK inhibitor is a potassium salt of the compound of formula (I), which is also referred to as VS-6766. Other pharmaceutically acceptable salts of the compound of formula (I) are contemplated herein.

[000200] In some embodiments, the dual RAF/MEK inhibitor is a compound having the structure of Formula (II):



including pharmaceutically acceptable salts thereof, wherein:



R^1 , R^2 , R^3 , and R^4 are each independently selected from the group consisting of H, deuterium, hydroxyl, halogen, cyano, nitro, optionally substituted amino, optionally substituted C-amido, optionally substituted N-amido, optionally substituted ester, optionally substituted sulfonyl, optionally substituted S-sulfonamido, optionally substituted N-sulfonamido, optionally substituted sulfonate, optionally substituted O-thiocarbamyl, optionally substituted N-thiocarbamyl, optionally substituted N-carbamyl, optionally substituted O-carbamyl, optionally substituted urea, optionally substituted C1 to C6 alkoxy, optionally substituted C1 to C6 alkyl, optionally substituted C2 to C6 alkenyl, optionally substituted C2 to C6 alkynyl, optionally substituted C3 to C8 cycloalkyl, optionally substituted C6 to C10 aryl, optionally substituted C3 to C8 heterocyclyl, optionally substituted C3 to C10 heteroaryl, and L; R^6 is selected from the group consisting of H, deuterium, hydroxyl, halogen, cyano, nitro, optionally substituted amino, optionally substituted C1 to C6 alkoxy, optionally substituted C1 to C6 alkyl, optionally substituted C2 to C6 alkenyl, and optionally substituted C2 to C6 alkynyl;

X is $C(R^5)_2$, $CH(R^5)$, CH_2 , $-O-$, $>C=O$, $>C=S$, or $>N-H$;

L is $-Z_1-Z_2$ or $-Z_1-Z_2-Z_3$;











Z_1 , Z_2 , and Z_3 are independently selected from the group consisting of $-CH_2-$, $-O-$, $-S-$, $S=O$, $-SO_2-$, $C=O$, $-CO_2-$, $-NO_2$, $-NH-$, $-CH_2CCH$, $-CH_2CN$, $-NR^5R^{5'}$, $-NH(CO)-$, $-(CO)NH-$, $-(CO)NR^5R^{5'}$, $-NH-SO_2-$, $-SO_2-NH-$, $-R^5CH_2-$, $-R^5O-$, $-R^5S-$, $R^5-S=O$, $-R^5SO_2-$, $R^5-C=O$, $-R^5CO_2-$, $-R^5NH-$, $-R^5NH(CO)-$, $-R^5(CO)NH-$, $-R^5NH-SO_2-$, $-R^5SO_2-NH-$, $-NHCH_2CO-$, $-CH_2R^5-$, $-OR^5-$, $-SR^5-$, $S=O-R^5$, $-SO_2R^5-$, $C=O-R^5$, $-CO_2R^5-$, $-NHR^5-$, $-NH(CO)R^5-$, $-(CO)NHR^5-$, $-NH-SO_2R^5-$, $-SO_2-NHR^5-$, optionally substituted C1 to C6 alkyl, optionally substituted C3 to C8 cycloalkyl, optionally substituted C6 to C10 aryl, optionally substituted C3 to C8 heterocyclyl, optionally substituted C3 to C10 heteroaryl, $-CH_2-(\text{optionally})$











substituted aryl), -CH₂-(optionally substituted C3 to C8 cycloalkyl), and - CH₂-(optionally substituted C3 to C10 heteroaryl); each R⁵ and R⁵ are independently selected from H, deuterium, optionally substituted C1 to C6 alkyl, optionally substituted C2 to C6 alkenyl, optionally substituted C2 to C6 alkynyl, optionally substituted C3 to C8 carbocyclyl, optionally substituted C6 to C10 aryl, optionally substituted C3 to C8 heterocyclyl, and optionally substituted C3 to C10 heteroaryl; and









Y is CH₂, NH, or O, with the proviso that R¹ is not -O-pyrimidyl.

[000201] In some embodiments, the dual RAF/MEK inhibitor is a compound selected from a compound in Table I:









Table I.









No.	Structure	No.	Structure
7		9	
10		11	
12		13	
14		15	
16		17	









No.	Structure	No.	Structure
18		19	
20		21	
22		23	
24		25	
26		27	









No.	Structure	No.	Structure
28		29	
30		31	
32		33	
34		35	









No.	Structure	No.	Structure
36		37	
38		39	
40		41	
42		43	









No.	Structure	No.	Structure
44		45	
46		47	
48		49	
50		51	









No.	Structure	No.	Structure
52		53	
54		55	
56		57	
58		59	











No.	Structure	No.	Structure
60		61	
62		63	
64		65	
66		67	

No.	Structure	No.	Structure
68		69	
70		71	
72		73	
74		75	











No.	Structure	No.	Structure
76		77	
78		79	
80		81	
82		83	

No.	Structure	No.	Structure
84		85	
86		87	
88		89	
90		91	

No.	Structure	No.	Structure
92		93	
94		95	
96		97	
98		99	

No.	Structure	No.	Structure
100		101	
102		103	
104		105	
106		107	
108		109	











No.	Structure	No.	Structure
110		111	
112		113	
114		115	
116		117	
118		119	

No.	Structure	No.	Structure
120		121	
122		123	
124		125	
126		127	
128		129	

No.	Structure	No.	Structure
130		131	
132		133	
134		135	
136		137	
138		139	











No.	Structure	No.	Structure
140		141	
142		143	
144		145	
146		147	
148		149	











No.	Structure	No.	Structure
150		151	
152		153	
154		155	
156		157	
158		159	

No.	Structure	No.	Structure
160		161	
162		163	
164		165	
166		167	
168		169	

No.	Structure	No.	Structure
170		171	
172		173	
174		175	
176		177	
178		179	

No.	Structure	No.	Structure
180		181	
182		183	
184		185	
186		187	
188		189	

No.	Structure	No.	Structure
190		191	
192		193	
194		195	
196		197	
198		199	

No.	Structure	No.	Structure
200		201	
202		203	
204		205	
206		207	
208		209	

No.	Structure	No.	Structure
210		211	
212		213	
214		215	
216		217	
218		219	

No.	Structure	No.	Structure
220		221	
222		223	
224		225	
226		227	
228		229	

No.	Structure	No.	Structure
230		231	
232		233	
234		235	

[000202] In some embodiments, the dual RAF/MEK inhibitor is IMM-1-104 (Immuneering) or a pharmaceutically acceptable salt thereof.

[000203] In some embodiments, the dual RAF/MEK inhibitor is dosed at least once a week (e.g., once a week, twice a week, three times a week, four times a week, five times a week, or six times a week). In some embodiments, the dual RAF/MEK inhibitor is dosed once a week. In some embodiments, the dual RAF/MEK inhibitor is dosed twice a week. In some embodiments, the dual RAF/MEK inhibitor is dosed three times a week.

[000204] In some embodiments, the dual RAF/MEK inhibitor is dosed at about 0.1 mg to about 100 mg, e.g., about 0.1 mg to about 50 mg, about 0.1 mg to about 10 mg, about 0.1 mg to about 5 mg, about 0.1 mg to about 4 mg, about 0.1 mg to about 3 mg, about 0.1 mg to about 2 mg, about 0.1 mg to about 1 mg, about 1 mg to about 5 mg, about 1 mg to about 10 mg, about 1 mg to about 20 mg, about 1 mg to about 40 mg, about 1 mg to about 60 mg, about 1 mg to about 80 mg, about 1 mg to about 100 mg, about 10 mg to about 100 mg, about

20 mg to about 100 mg, about 40 mg to about 100 mg, about 60 mg to about 100 mg, or about 80 mg to about 100 mg. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 0.5 mg to about 10 mg per administration. In some embodiments, dual RAF/MEK inhibitor is dosed at about 0.8 mg to about 10 mg per administration. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 1 mg to about 5 mg per administration. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 2 mg to about 4 mg per administration. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 0.1 mg, 0.2 mg, 0.5 mg, 1 mg, 1.5 mg, 3 mg, 4 mg, 5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg, 35 mg, 40 mg, 45 mg, 50 mg, 65 mg, 70 mg, 75 mg, 80 mg, 85 mg, 90 mg, 95 mg, or 100 mg per administration. In some embodiments, dual RAF/MEK inhibitor is dosed at about 4 mg per administration. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 3.2 mg per administration. In some embodiments, the dual RAF/MEK inhibitor is administered orally.

[000205] In some embodiments, the dual RAF/MEK inhibitor is dosed as a cycle. In some embodiments, the cycle comprises administering the dual RAF/MEK inhibitor for three weeks and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed once a week. In some embodiments, the dual RAF/MEK inhibitor is dosed twice a week. In some embodiments, the dual RAF/MEK inhibitor is dosed three times a week. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 0.8 mg to about 10 mg per administration. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 1 mg to about 5 mg per administration. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 2 mg to about 4 mg per administration. In some embodiments, dual RAF/MEK inhibitor is dosed at about 4 mg per administration. In some embodiments, the dual RAF/MEK inhibitor is dosed at about 3.2 mg per administration.

[000206] In some embodiments, the dual RAF/MEK inhibitor is dosed twice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of about 0.8 mg to about 10 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed twice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of about 1 mg to about 5 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed twice a week as a cycle, wherein the

cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of about 2 mg to about 4 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed twice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of 3.2 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed twice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of 4 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the cycle is repeated at least once.


[000207] In some embodiments, the dual RAF/MEK inhibitor is dosed thrice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of about 0.8 mg to about 10 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed thrice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of about 1 mg to about 5 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed thrice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of about 2 mg to about 4 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed thrice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of 3.2 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the dual RAF/MEK inhibitor is dosed thrice a week as a cycle, wherein the cycle comprises administering the dual RAF/MEK inhibitor for three weeks at a dose of 4 mg per administration and then not administering the dual RAF/MEK inhibitor for one week. In some embodiments, the cycle is repeated at least once.

[000208] In alternative embodiments, the dual RAF/MEK inhibitor is dosed continuously (i.e., without a period of time, e.g., one week, wherein the dual RAF/MEK inhibitor is not administered). In some embodiments, the dual RAF/MEK inhibitor is dosed once a week. In some embodiments, the dual RAF/MEK inhibitor is dosed twice a week. In some embodiments, the dual RAF/MEK inhibitor is dosed three times a week.

RAF inhibitors


[000209] Exemplary RAF inhibitors include, but are not limited to,

donafenib having the following structure:




, or a pharmaceutically acceptable salt thereof,

lifirafenib having the following structure:




, or a pharmaceutically acceptable salt thereof,

rigosertib having the following structure:




, or a pharmaceutically acceptable salt thereof,

BMS-908662 having the following structure:



, or a pharmaceutically acceptable salt thereof,

XP-102 having the following structure:



, or a pharmaceutically acceptable salt thereof,

ABM-2526 (ABM Therapeutics), DDC-PanRAF (Deciphera Pharmaceuticals), FNX006 (Chengdu Fanuoxi Biomedical Technology), and VRN-XX (Voronoi), and pharmaceutically acceptable salts thereof.

[000210] In some embodiments, the RAF inhibitor is dosed at about 0.1 mg to about 5000 mg, e.g., about 1 mg to about 3000 mg, about 10 mg to about 2000 mg, e.g., about 100 mg to about 2000 mg, about 100 mg to about 1500 mg, about 100 mg to about 1000 mg, about 100 mg to about 800 mg, about 100 mg to about 600 mg, about 100 mg to about 400 mg, about 100 mg to about 200 mg, about 200 mg to about 2000 mg, about 200 mg to about 1500 mg, about 200 mg to about 1000 mg, about 200 mg to about 800 mg, about 200 mg to about 600 mg, about 200 mg to about 400 mg, about 400 mg to about 2000 mg, about 400 mg to about 1500 mg, about 400 mg to about 1000 mg, about 400 mg to about 800 mg, about 400 mg to about 600 mg, about 600 mg to about 2000 mg, about 600 mg to about 1500 mg, about 600 mg to about 1000 mg, about 600 mg to about 800 mg, about 800 mg to about 2000 mg, about 800 mg to about 1500 mg, about 800 mg to about 1000 mg, about 600 mg to about 2000 mg, about 600 mg to about 1500 mg, about 600 mg to about 1000 mg, about 600 mg to about 800 mg per administration. In some embodiments, the RAF inhibitor is dosed at 10 mg to 2000 mg per administration. In some embodiments, the RAF inhibitor is dosed at 100 mg to 1000 mg per administration. In some embodiments, the RAF inhibitor is dosed at 100 mg to 200 mg per administration. In some embodiments, the RAF inhibitor is dosed at 300 mg to 500 mg per administration. In some embodiments, the RAF inhibitor is dosed at 200 mg to 600 mg per administration. In some embodiments, the RAF inhibitor is dosed at 800 mg to 1000 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 1 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 5 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 10 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 50 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 100 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 150 mg per

administration. In some embodiments, the RAF inhibitor is dosed at about 200 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 250 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 300 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 350 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 400 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 450 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 500 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 550 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 600 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 650 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 700 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 750 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 800 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 850 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 900 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 950 mg per administration. In some embodiments, the RAF inhibitor is dosed at about 1000 mg per administration.

[000211] In some embodiments, the RAF inhibitor is administered at least once a week. In some embodiments, the RAF inhibitor is administered once a week. In some embodiments, the RAF inhibitor is administered twice a week. In some embodiments, the RAF inhibitor is administered thrice a week. In some embodiments, the RAF inhibitor is administered four times a week. In some embodiments, the RAF inhibitor is administered five times a week. In some embodiments, the RAF inhibitor is administered six times a week. In some embodiments, the RAF inhibitor is administered at least once daily. In some embodiments, the RAF inhibitor is administered once daily. In some embodiments, the RAF inhibitor is administered twice daily. In some embodiments, the RAF inhibitor is administered orally.

[000212] In some embodiments, the RAF inhibitor is administered as a cycle. For example, in some embodiments, the RAF inhibitor is dosed as a cycle comprising administering the RAF inhibitor for three weeks and then not administering the RAF inhibitor for one week.

[000213] In some embodiments, the dual RAF/MEK inhibitor is administered before the RAF inhibitor is administered. In some embodiments, the dual RAF/MEK inhibitor is


administered after the RAF inhibitor is administered. In some embodiments, the dual RAF/MEK inhibitor is administered concurrently with the RAF inhibitor.

[000214] In some embodiments, the dual RAF/MEK inhibitor and the RAF inhibitor are independently dosed cyclically.


[000215] In some embodiments, the dual RAF/MEK inhibitor and RAF inhibitor are simultaneously dosed cyclically.

BRAF V600 inhibitors


[000216] Exemplary BRAF V600 inhibitors include, but are not limited to, dabrafenib having the following structure:




encorafenib having the following structure:



vemurafenib having the following structure:




FORE-8394 (PLX-8394) having the following structure:




, or a pharmaceutically acceptable salt thereof,

tinloragenib (PF-07284890 or ARRY-461) (Pfizer), having the following structure:




, or a pharmaceutically acceptable salt thereof,

AZ-304 having the following structure:




, or a pharmaceutically acceptable salt thereof,

agerafenib having the following structure:



, or a pharmaceutically acceptable salt thereof,

KIN-2787 having the following structure:



, or a pharmaceutically acceptable salt thereof,

BGB-3245 (MapKure), ABM-1310 (ABM Therapeutics), TQB-3233 (Jiangsu Chia-tai Tianqing Pharmaceutical), UB-941 (UBI Pharma), AFX-1251 (Afecta Pharmaceuticals),

ARQ 736 (Merck), ASN003 (Asana BioSciences), AVB-BRAF (Aravive), BDTX-4933 (Black Diamond Therapeutics), CFT1946 (C4 Therapeutics), HLX208 (NeuPharma), RO5212054 (Daiichi Sankyo), RO7276389 (Genentech), and TQ-B3233 (Chia Tai Tianqing Pharmaceutical), and pharmaceutically acceptable salts thereof. In some embodiments, the BRAF V600 inhibitor is dabrafenib, encorafenib, vemurafenib, FORE-8394, tinloragenib, AZ-304, agerafenib, or KIN-2787, or pharmaceutically acceptable salts thereof. In some embodiments, the BRAF V600 inhibitor is dabrafenib, encorafenib, or vemurafenib, or pharmaceutically acceptable salts thereof. In some embodiments, the BRAF V600 inhibitor is dabrafenib, or a pharmaceutically acceptable salt thereof. In some embodiments, the BRAF V600 inhibitor is encorafenib, or a pharmaceutically acceptable salt thereof. In some embodiments, the BRAF V600 inhibitor is vemurafenib, or a pharmaceutically acceptable salt thereof.

[000217] In some embodiments, the BRAF V600 inhibitor is dosed at about 0.1 mg to about 5000 mg, e.g., about 1 mg to about 3000 mg, about 10 mg to about 2000 mg, e.g., about 100 mg to about 2000 mg, about 100 mg to about 1500 mg, about 100 mg to about 1000 mg, about 100 mg to about 800 mg, about 100 mg to about 600 mg, about 100 mg to about 400 mg, about 100 mg to about 200 mg, about 200 mg to about 2000 mg, about 200 mg to about 1500 mg, about 200 mg to about 1000 mg, about 200 mg to about 800 mg, about 200 mg to about 600 mg, about 200 mg to about 400 mg, about 400 mg to about 2000 mg, about 400 mg to about 1500 mg, about 400 mg to about 1000 mg, about 400 mg to about 800 mg, about 400 mg to about 600 mg, about 600 mg to about 2000 mg, about 600 mg to about 1500 mg, about 600 mg to about 1000 mg, about 600 mg to about 800 mg, about 800 mg to about 2000 mg, 800 mg to about 1500 mg, about 800 mg to about 1000 mg, about 600 mg to about 2000 mg, about 600 mg to about 1500 mg, about 600 mg to about 1000 mg, about 600 mg to about 800 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at 10 mg to 2000 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at 100 mg to 1000 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at 100 mg to 200 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at 300 mg to 500 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at 200 mg to 600 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at 800 mg to 1000 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 1 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 5 mg per

administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 10 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 50 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 100 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 150 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 200 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 250 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 300 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 350 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 400 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 450 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 500 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 550 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 600 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 650 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 700 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 750 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 800 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 850 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 900 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 950 mg per administration. In some embodiments, the BRAF V600 inhibitor is dosed at about 1000 mg per administration.

[000218] In some embodiments, the BRAF V600 inhibitor is administered at least once a week. In some embodiments, the BRAF V600 inhibitor is administered once a week. In some embodiments, the BRAF V600 inhibitor is administered twice a week. In some embodiments, the BRAF V600 inhibitor is administered thrice a week. In some embodiments, the BRAF V600 inhibitor is administered four times a week. In some embodiments, the BRAF V600 inhibitor is administered five times a week. In some embodiments, the BRAF V600 inhibitor is administered six times a week. In some embodiments, the BRAF V600 inhibitor is administered at least once daily. In some embodiments, the BRAF V600 inhibitor is administered once daily. In some embodiments, the BRAF V600 inhibitor is administered twice daily. In some embodiments, the BRAF V600 inhibitor is administered orally.

[000219] In some embodiments, the BRAF V600 inhibitor is administered as a cycle. For example, in some embodiments, the BRAF V600 inhibitor is dosed as a cycle comprising administering the BRAF V600 inhibitor for three weeks and then not administering the BRAF V600 inhibitor for one week.


[000220] In some embodiments, the dual RAF/MEK inhibitor is administered before the BRAF V600 inhibitor is administered. In some embodiments, the dual RAF/MEK inhibitor is administered after the BRAF V600 inhibitor is administered. In some embodiments, the dual RAF/MEK inhibitor is administered concurrently with the BRAF V600 inhibitor.

[000221] In some embodiments, the dual RAF/MEK inhibitor and the BRAF V600 inhibitor are independently dosed cyclically.

[000222] In some embodiments, the dual RAF/MEK inhibitor and BRAF V600 inhibitor are simultaneously dosed cyclically.

pan-RAF inhibitors


[000223] Exemplary pan-RAF inhibitors include, but are not limited to, belvarafenib having the following structure:



, or a pharmaceutically acceptable salt


thereof,

naporafenib having the following structure:




, or a pharmaceutically acceptable salt thereof,

lifirafenib having the following structure:




, or a pharmaceutically acceptable salt thereof,

tovorafenib having the following structure:




, or a pharmaceutically acceptable salt thereof,

BAL3833 having the following structure:




, or a pharmaceutically acceptable salt thereof,

LY3009120, having the following structure:



, or a pharmaceutically acceptable salt thereof,

REDX05358 having the following structure:



, or a pharmaceutically acceptable salt thereof,

IRICoR-Ipsen (IRICoR), JZP815 (Jazz Pharmaceuticals), METiS-01 (METiS Therapeutics), QLH11906 (Qilu Pharmaceutical), and SJ-C1044 (Samjin Pharmaceutical), and pharmaceutically acceptable salts thereof.

[000224] In some embodiments, the pan-RAF inhibitor is dosed at about 0.1 mg to about 5000 mg, e.g., about 1 mg to about 3000 mg, about 10 mg to about 2000 mg, e.g., about 100 mg to about 2000 mg, about 100 mg to about 1500 mg, about 100 mg to about 1000 mg, about 100 mg to about 800 mg, about 100 mg to about 600 mg, about 100 mg to about 400 mg, about 100 mg to about 200 mg, about 200 mg to about 2000 mg, about 200 mg to about 1500 mg, about 200 mg to about 1000 mg, about 200 mg to about 800 mg, about 200 mg to about 600 mg, about 200 mg to about 400 mg, about 400 mg to about 2000 mg, about 400 mg to about 1500 mg, about 400 mg to about 1000 mg, about 400 mg to about 800 mg, about 400 mg to about 600 mg, about 600 mg to about 2000 mg, about 600 mg to about 1500 mg, about 600 mg to about 1000 mg, about 600 mg to about 800 mg, about 800 mg to about 2000 mg, 800 mg to about 1500 mg, about 800 mg to about 1000 mg, about 600 mg to about 2000 mg, about 600 mg to about 1500 mg, about 600 mg to about 1000 mg, about 600 mg to about 800 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at 10 mg to 2000 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at 100 mg to 1000 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at 100 mg to 200 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at 300 mg to 500 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at 200 mg to 600 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at 800 mg to 1000 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 1 mg per administration. In some

embodiments, the pan-RAF inhibitor is dosed at about 5 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 10 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 50 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 100 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 150 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 200 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 250 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 300 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 350 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 400 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 450 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 500 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 550 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 600 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 650 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 700 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 750 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 800 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 850 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 900 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 950 mg per administration. In some embodiments, the pan-RAF inhibitor is dosed at about 1000 mg per administration.

[000225] In some embodiments, the pan-RAF inhibitor is administered at least once a week. In some embodiments, the pan-RAF inhibitor is administered once a week. In some embodiments, the pan-RAF inhibitor is administered twice a week. In some embodiments, the pan-RAF inhibitor is administered thrice a week. In some embodiments, the pan-RAF inhibitor is administered four times a week. In some embodiments, the pan-RAF inhibitor is administered five times a week. In some embodiments, the pan-RAF inhibitor is administered six times a week. In some embodiments, the pan-RAF inhibitor is administered at least once daily. In some embodiments, the pan-RAF inhibitor is administered once daily. In some embodiments, the pan-RAF inhibitor is administered twice daily. In some embodiments, the pan-RAF inhibitor is administered orally.

[000226] In some embodiments, the pan-RAF inhibitor is administered as a cycle. For example, in some embodiments, the pan-RAF inhibitor is dosed as a cycle comprising

administering the pan-RAF inhibitor for three weeks and then not administering the pan-RAF inhibitor for one week.

[000227] In some embodiments, the dual RAF/MEK inhibitor is administered before the pan-RAF inhibitor is administered. In some embodiments, the dual RAF/MEK inhibitor is administered after the pan-RAF inhibitor is administered. In some embodiments, the dual RAF/MEK inhibitor is administered concurrently with the pan-RAF inhibitor.

[000228] In some embodiments, the dual RAF/MEK inhibitor and the pan-RAF inhibitor are independently dosed cyclically.

[000229] In some embodiments, the dual RAF/MEK inhibitor and pan-RAF inhibitor are simultaneously dosed cyclically.

CRAF inhibitors

[000230] Exemplary CRAF inhibitors include, but are not limited to, MG005 (Metagene Biotech), Quanta-RAF1 (Quanta Therapeutics), and STX200 (SyntheX), and pharmaceutically acceptable salts thereof.

[000231] In some embodiments, the CRAF inhibitor is dosed at about 0.1 mg to about 5000 mg, e.g., about 1 mg to about 3000 mg, about 10 mg to about 2000 mg, e.g., about 100 mg to about 2000 mg, about 100 mg to about 1500 mg, about 100 mg to about 1000 mg, about 100 mg to about 800 mg, about 100 mg to about 600 mg, about 100 mg to about 400 mg, about 100 mg to about 200 mg, about 200 mg to about 2000 mg, about 200 mg to about 1500 mg, about 200 mg to about 1000 mg, about 200 mg to about 800 mg, about 200 mg to about 600 mg, about 200 mg to about 400 mg, about 400 mg to about 2000 mg, about 400 mg to about 1500 mg, about 400 mg to about 1000 mg, about 400 mg to about 800 mg, about 400 mg to about 600 mg, about 600 mg to about 2000 mg, about 600 mg to about 1500 mg, about 600 mg to about 1000 mg, about 600 mg to about 800 mg, about 800 mg to about 2000 mg, about 800 mg to about 1500 mg, about 800 mg to about 1000 mg, about 600 mg to about 2000 mg, about 600 mg to about 1500 mg, about 600 mg to about 1000 mg, about 600 mg to about 800 mg per administration. In some embodiments, the CRAF inhibitor is dosed at 10 mg to 2000 mg per administration. In some embodiments, the CRAF inhibitor is dosed at 100 mg to 1000 mg per administration. In some embodiments, the CRAF inhibitor is dosed at 100 mg to 200 mg per administration. In some embodiments, the CRAF inhibitor is dosed at 300 mg to 500 mg per administration. In some embodiments, the CRAF inhibitor is dosed at 200 mg to 600 mg per administration. In some embodiments, the CRAF inhibitor is dosed at 800 mg to 1000 mg per administration. In some embodiments, the CRAF inhibitor is dosed at

about 1 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 5 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 10 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 50 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 100 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 150 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 200 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 250 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 300 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 350 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 400 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 450 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 500 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 550 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 600 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 650 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 700 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 750 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 800 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 850 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 900 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 950 mg per administration. In some embodiments, the CRAF inhibitor is dosed at about 1000 mg per administration.

[000232] In some embodiments, the CRAF inhibitor is administered at least once a week. In some embodiments, the CRAF inhibitor is administered once a week. In some embodiments, the CRAF inhibitor is administered twice a week. In some embodiments, the CRAF inhibitor is administered thrice a week. In some embodiments, the CRAF inhibitor is administered four times a week. In some embodiments, the CRAF inhibitor is administered five times a week. In some embodiments, the CRAF inhibitor is administered six times a week. In some embodiments, the CRAF inhibitor is administered at least once daily. In some embodiments, the CRAF inhibitor is administered once daily. In some embodiments, the CRAF inhibitor is administered twice daily. In some embodiments, the CRAF inhibitor is administered orally.

[000233] In some embodiments, the CRAF inhibitor is administered as a cycle. For example, in some embodiments, the CRAF inhibitor is dosed as a cycle comprising administering the RAF inhibitor for three weeks and then not administering the CRAF inhibitor for one week.

[000234] In some embodiments, the dual RAF/MEK inhibitor is administered before the CRAF inhibitor is administered. In some embodiments, the dual RAF/MEK inhibitor is administered after the CRAF inhibitor is administered. In some embodiments, the dual RAF/MEK inhibitor is administered concurrently with the CRAF inhibitor.

[000235] In some embodiments, the dual RAF/MEK inhibitor and the CRAF inhibitor are independently dosed cyclically.


[000236] In some embodiments, the dual RAF/MEK inhibitor and CRAF inhibitor are simultaneously dosed cyclically.

FAK Inhibitors

[000237] Potent inhibitors of the FAK protein tyrosine kinases may be adapted to therapeutic use as antiproliferative agents (e.g., anticancer), antitumor (e.g., effective against solid tumors), antiangiogenesis (e.g., stop or prevent proliferation of blood vessels) in mammals, particularly in humans. In some embodiments, the methods described herein further contemplate administering to the subject a FAK inhibitor described herein. The FAK inhibitors may be useful in the prevention and treatment of non-hematologic malignancies, a variety of human hyperproliferative disorders such as malignant and benign tumors of the liver, kidney, bladder, breast, gastric, ovarian, colorectal, prostate, pancreatic, lung, vulval, thyroid, hepatic carcinomas, sarcomas, glioblastomas, head and neck, and other hyperplastic conditions such as benign hyperplasia of the skin (e.g., psoriasis) and benign hyperplasia of the prostate (e.g., BPH), and in the prevention and treatment of disorders such as mesothelioma. In some embodiments, the compounds described herein, e.g., FAK inhibitors, inhibit protein tyrosine kinase 2 (PYK2).


[000238] In some embodiments, the methods described herein further contemplate administering to the subject an effective amount of a FAK inhibitor.

[000239] An exemplary FAK inhibitor includes, but is not limited to, defactinib having the following structure:




or a pharmaceutically acceptable salt thereof. Defactinib is also known as VS-6063 (e.g., VS-6063 free base) or PF-04554878. VS-6063 and related compounds are also disclosed in, for example, U.S. Patent No. 7,928,109, the content of which is incorporated herein by reference. In some embodiments, VS-6063 can form a pharmaceutically acceptable salt (e.g., VS-6063 hydrochloride).

[000240] In some embodiments, the FAK inhibitor is VS-4718, having the following structure:




or a pharmaceutically acceptable salt thereof.

[000241] In some embodiments, the FAK inhibitor is TAE226, having the following structure:




or a pharmaceutically acceptable salt thereof.

[000242] In some embodiments, the FAK inhibitor is GSK2256098, having the following structure:




or a pharmaceutically acceptable salt thereof.

[000243] In some embodiments, the FAK inhibitor is PF-03814735, having the following structure:



or a pharmaceutically acceptable salt thereof.

[000244] In some embodiments, the FAK inhibitor is BI-4464, having the following structure:



or a pharmaceutically acceptable salt thereof.

[000245] In some embodiments, the FAK inhibitor is BI-853520 (IN10018; Boehringer Ingelheim). In some other embodiments, the FAK inhibitor is APG-2449 (Ascentage Pharma Group).

[000246] In some embodiments, the FAK inhibitor is selected from the group consisting of defactinib, TAE226, BI-853520, GSK2256098, PF-03814735, BI-4464, VS-4718, and APG-2449, or a pharmaceutically acceptable salt thereof. For example, the FAK inhibitor is defactinib or a pharmaceutically acceptable salt thereof.

[000247] In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed at least once daily. For example, in some embodiments, the FAK inhibitor (e.g., defactinib) is dosed once daily. In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed twice daily.

[000248] In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed at about 100 mg to about 1000 mg, e.g., about 100 mg to about 800 mg, about 100 mg to about 600 mg, about 100 mg to about 400 mg, about 100 mg to about 200 mg, about 200 mg to about 1000 mg, about 400 mg to about 1000 mg, about 600 mg to about 1000 mg, about 800 mg to about 1000 mg, about 200 mg to about 800 mg, about 200 mg to about 600 mg, about 200 mg to about 400 mg, about 400 mg to about 800 mg, or about 400 mg to about 600 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed at about 200 mg to about 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed at about 100 mg per administration. In some embodiments, the FAK

inhibitor (e.g., defactinib) is dosed at about 200 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed at about 300 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed at about 400 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed at about 500 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib) is dosed at about 600 mg per administration. In some embodiments, the FAK inhibitor (e.g., defactinib) is administered orally.

[000249] In some embodiments, the FAK inhibitor is dosed as a cycle, wherein the cycle comprises administering the FAK inhibitor for three weeks and then not administering the FAK inhibitor for one week. In some embodiments, the cycle is repeated at least once.

Diseases and Disorders

Abnormal Cell Growth

[000250] Abnormal cell growth, as used herein and unless otherwise indicated, refers to cell growth that is independent of normal regulatory mechanisms (e.g., loss of contact inhibition). This includes the abnormal growth of: (1) tumor cells (tumors) that proliferate, for example, by expressing a mutated tyrosine kinase or overexpression of a receptor tyrosine kinase; (2) benign and malignant cells of other proliferative diseases, for example, in which aberrant tyrosine kinase activation occurs; (3) any tumors that proliferate, for example, by receptor tyrosine kinases; (4) any tumors that proliferate, for example, by aberrant serine/threonine kinase activation; and (5) benign and malignant cells of other proliferative diseases, for example, in which aberrant serine/threonine kinase activation occurs. Abnormal cell growth can refer to cell growth in epithelial (e.g., carcinomas, adenocarcinomas); mesenchymal (e.g., sarcomas (e.g. leiomyosarcoma, Ewing's sarcoma)); hematopoietic (e.g., lymphomas, leukemias, myelodysplasias (e.g., pre-malignant)); or other (e.g., melanoma, mesothelioma, and other tumors of unknown origin) cell.

Neoplastic Disorders

[000251] Abnormal cell growth can refer to a neoplastic disorder. A "neoplastic disorder" is a disease or disorder characterized by cells that have the capacity for autonomous growth or replication, e.g., an abnormal state or condition characterized by proliferative cell growth. An abnormal mass of tissue as a result of abnormal cell growth or division, or a "neoplasm," can be benign, pre-malignant (carcinoma in situ) or malignant (cancer).

[000252] Exemplary neoplastic disorders include: carcinoma, sarcoma, metastatic disorders (e.g., tumors arising from prostate, colon, lung, breast and liver origin),

hematopoietic neoplastic disorders, e.g., leukemias, metastatic tumors. Treatment with the compound may be in an amount effective to ameliorate at least one symptom of the neoplastic disorder, e.g., reduced cell proliferation, reduced tumor mass, etc.

Cancer

[000253] The inventive methods of the present invention may be useful in the prevention and treatment of cancer, including for example, solid tumors, soft tissue tumors, and metastases thereof. The disclosed methods are also useful in treating non-solid cancers. Exemplary solid tumors include malignancies (e.g., sarcomas, adenocarcinomas, and carcinomas) of the various organ systems, such as those of lung, breast, lymphoid, gastrointestinal (e.g., colon), and genitourinary (e.g., renal, urothelial, or testicular tumors) tracts, pharynx, prostate, and ovary. Exemplary adenocarcinomas include colorectal cancers, renal-cell carcinoma, liver cancer (e.g., Hepatocellular carcinoma), non-small cell carcinoma of the lung, pancreatic (e.g., metastatic pancreatic adenocarcinoma) and cancer of the small intestine.

[000254] The cancer can include mesothelioma; neurofibromatosis; e.g., neurofibromatosis type 2, neurofibromatosis type 1; renal cancer; lung cancer, non small cell lung cancer; liver cancer; thyroid cancer; ovarian; breast cancer; a nervous system tumor; schwannoma; meningioma; schwannomatosis; neuroma acoustic; adenoid cystic carcinoma; ependymoma; ependymal tumors, or any other tumor which exhibits decreased merlin expression and/or mutation, and/or deletion and/or promotor hypermethylation of the NF-2 gene. In some embodiments, the cancer is renal cancer.

[000255] For example, the cancer may include, but is not limited to, ovarian cancer, non-small cell lung cancer (e.g., NSCLC adenocarcinoma)), uterine endometrioid carcinoma, pancreatic adenocarcinoma, colorectal adenocarcinoma, colorectal cancer, pancreatic cancer, or lung adenocarcinoma. In some embodiments, the NSCLC is characterized as having a KRAS mutation. In some embodiments, the ovarian cancer is low grade serous ovarian cancer.

[000256] The cancer can include cancers characterized as comprising cancer stem cells, cancer associated mesenchymal cells, or tumor initiating cancer cells. The cancer can include cancers that have been characterized as being enriched with cancer stem cells, cancer associated mesenchymal cells, or tumor initiating cancer cells (e.g., a tumor enriched with cells that have undergone an epithelial-to-mesenchymal transition or a metastatic tumor).

[000257] The cancer can include pancreatic cancer, gynecologic cancer (e.g., cervical cancer, ovarian cancer, uterine cancer, vaginal cancer, endometrial cancer, or vulvar cancer),

liver cancer, prostate cancer, mesothelioma, breast cancer, bladder cancer, melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the cancer is melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer. In some embodiments, the lung cancer is non-small cell lung cancer. In some embodiments, the lung cancer is metastatic non-small cell lung cancer. In some embodiments, the melanoma is unresectable melanoma. In some embodiments, the melanoma is metastatic melanoma. In some embodiments, the cancer is colorectal cancer. In some embodiments, the thyroid cancer is papillary thyroid cancer. In some embodiments, the thyroid cancer is follicular thyroid cancer. In some embodiments, the thyroid cancer is anaplastic thyroid cancer.

[000258] The cancer can be a primary tumor, i.e., located at the anatomical site of tumor growth initiation. The cancer can also be metastatic, i.e., appearing at least a second anatomical site other than the anatomical site of tumor growth initiation. The cancer can be a recurrent cancer, i.e., cancer that returns following treatment, and after a period of time in which the cancer was undetectable. The recurrent cancer can be anatomically located locally to the original tumor, e.g., anatomically near the original tumor; regionally to the original tumor, e.g., in a lymph node located near the original tumor; or distantly to the original tumor, e.g., anatomically in a region remote from the original tumor.

[000259] The cancer can also include for example, but is not limited to, epithelial cancers, breast, lung, pancreatic, colorectal (e.g., metastatic colorectal, e.g., metastatic KRAS mutated), prostate, head and neck, melanoma (e.g., NRAS mutated locally advanced or metastatic malignant cutaneous melanoma), acute myelogenous leukemia, and glioblastoma. Exemplary breast cancers include triple negative breast cancer, basal-like breast cancer, claudin-low breast cancer, invasive, inflammatory, metaplastic, and advanced HER-2 positive or ER-positive cancers resistant to therapy.

[000260] In some embodiments, the cancer is characterized as having a RAS mutation. In some embodiments, the cancer is a cancer characterized as having a KRAS mutation. In some embodiments, the cancer is a cancer characterized as having a NRAS mutation. In some embodiments, the cancer is a cancer characterized as having a HRAS mutation.

[000261] In some embodiments, the cancer is a cancer characterized as having a RAF mutation. In some embodiments, the cancer is a cancer characterized as having a BRAF mutation.

[000262] The cancer can also include lung adenocarcinoma, colorectal cancer (CRC), uveal melanoma, ovarian cancer, uterine endometrioid carcinoma, bladder urothelial carcinoma, breast invasive lobular carcinoma, cervical squamous cell carcinoma, cutaneous melanoma, endocervical adenocarcinoma, hepatocellular carcinoma, pancreatic adenocarcinoma, biphasic type pleural mesothelioma, renal clear cell carcinoma, renal clear cell carcinoma, stomach adenocarcinoma, tubular stomach adenocarcinoma, uterine carcinosarcoma, or uterine malignant mixed Mullerian tumor.

[000263] In some embodiments, the cancer is unresectable or metastatic melanoma, melanoma with lymph node involvement or metastatic disease who have undergone complete resection, metastatic non-small cell lung cancer and progression on or after platinum-based chemotherapy, metastatic small cell lung cancer with progression after platinum-based chemotherapy and at least one other line of therapy, advanced renal cell carcinoma who have received prior antiangiogenic therapy, advanced renal cell carcinoma, classical Hodgkin lymphoma, recurrent or metastatic squamous cell carcinoma of the head and neck with disease progression on or after a platinum-based therapy, locally advanced or metastatic urothelial carcinoma, microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) metastatic colorectal cancer, or hepatocellular carcinoma.

[000264] In some embodiments, the cancer is melanoma, non-small cell lung cancer, small cell lung cancer, head and neck squamous cell cancer, classical Hodgkin lymphoma, primary mediastinal large B-cell lymphoma, urothelial carcinoma, microsatellite instability-high cancer, gastric cancer, esophageal cancer, cervical cancer, hepatocellular carcinoma, merkel cell carcinoma, renal cell carcinoma, or endometrial carcinoma.

[000265] Other cancers include but are not limited to, uveal melanoma, brain, abdominal, esophagus, gastrointestinal, glioma, liver, tongue, neuroblastoma, osteosarcoma, ovarian, retinoblastoma, Wilm's tumor, multiple myeloma, skin, lymphoma, blood and bone marrow cancers (e.g., advanced hematological malignancies, leukemia, e.g., acute myeloid leukemia (e.g., primary or secondary), acute lymphoblastic leukemia, acute lymphocytic leukemia, T cell leukemia, hematological malignancies, advanced myeloproliferative disorders, myelodysplastic syndrome, relapsed or refractory multiple myeloma, advanced myeloproliferative disorders), retinal, bladder, cervical, kidney, endometrial, meningioma, lymphoma, skin, uterine, lung, non small cell lung, nasopharyngeal carcinoma, neuroblastoma, solid tumor, hematologic malignancy, squamous cell carcinoma, testicular, thyroid, mesothelioma, brain vulval, sarcoma, intestine, oral, endocrine, salivary, spermatocyte seminoma, sporadic medullary thyroid carcinoma, non-proliferating testes cells,

cancers related to malignant mast cells, non-Hodgkin's lymphoma, and diffuse large B cell lymphoma.

[000266] In some embodiments, the tumor is a solid tumor. In some embodiments, the solid tumor is locally advanced or metastatic, hi some embodiments, the solid tumor is refractory (e.g., resistant) after standard therapy.

[000267] Methods described herein can reduce, ameliorate or altogether eliminate the disorder, and/or its associated symptoms, to keep it from becoming worse, to slow the rate of progression, or to minimize the rate of recurrence of the disorder once it has been initially eliminated (i.e., to avoid a relapse). A suitable dose and therapeutic regimen may vary depending upon the specific compounds, combinations, and/or pharmaceutical compositions used and the mode of delivery of the compounds, combinations, and/or pharmaceutical compositions. In some embodiments, the method increases the average length of survival, increases the average length of progression-free survival, and/or reduces the rate of recurrence, of subjects treated with the combinations described herein in a statistically significant manner.

[000268] In some embodiments, the cancer is lung cancer (e.g., non-small cell lung cancer CNSCLC), e.g., KRAS mutant NSCLC; metastatic cancer), bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer (e.g., unresectable low-grade ovarian, advanced or metastatic ovarian cancer), rectal cancer, cancer of the anal region, stomach cancer, colon cancer, breast cancer (e.g., triple-negative breast cancer (e.g., breast cancer which does not express the genes for the estrogen receptor, progesterone receptor, and Her2/neu)), uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, chronic or acute leukemia, lymphocytic lymphomas, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, neoplasms of the central nervous system (CNS), primary CNS lymphoma, spinal axis tumors, brain stem glioma, pituitary adenoma, mesothelioma (e.g., malignant pleural mesothelioma, e.g., surgical resectable malignant pleural mesothelioma) or a combination of one or more of the foregoing cancers. In some embodiments, the cancer is metastatic. In some embodiments, the abnormal cell growth is locally recurring (e.g., the subject has a locally recurrent disease, e.g., cancer).

Additional Therapies

[000269] In some embodiments, the methods and compositions described herein is administered together with an additional therapy or additional agent. In one embodiment, a mixture of one or more compounds or pharmaceutical compositions may be administered with the combination described herein to a subject in need thereof. In yet another embodiment, one or more compounds or compositions (e.g., pharmaceutical compositions) may be administered with the combination described herein for the treatment or avoidance of various diseases, including, for example, cancer, diabetes, neurodegenerative diseases, cardiovascular disease, blood clotting, inflammation, flushing, obesity, aging, stress, etc. In various embodiments, combination therapies comprising a compound or pharmaceutical composition described herein may refer to (1) pharmaceutical compositions that comprise one or more compounds in combination with the combination described herein; and (2) co-administration of one or more compounds or pharmaceutical compositions described herein with the combination described herein, wherein the compound or pharmaceutical composition described herein have not been formulated in the same compositions. In some embodiments, the combinations described herein is administered with an additional treatment (e.g., an additional cancer treatment). In some embodiments, the additional treatment (e.g., an additional cancer treatment) can be administered simultaneously (e.g., at the same time), in the same or in separate compositions, or sequentially. Sequential administration refers to administration of one treatment before (e.g., immediately before, less than 5, 10, 15, 30, 45, 60 minutes; 1, 2, 3, 4, 6, 8, 10, 12, 16, 20, 24, 48, 72, 96 or more hours; 4, 5, 6, 7, 8, 9 or more days; 1, 2, 3, 4, 5, 6, 7, 8 or more weeks before) administration of an additional, e.g., secondary, treatment (e.g., a compound or therapy). The order of administration of the first and secondary compound or therapy can also be reversed.

[000270] Exemplary cancer treatments include, for example: chemotherapy, targeted therapies such as antibody therapies, immunotherapy, and hormonal therapy. Examples of each of these treatments are provided below.

Chemotherapy

[000271] In some embodiments, a combination described herein is administered with a chemotherapy. Chemotherapy is the treatment of cancer with drugs that can destroy cancer cells. "Chemotherapy" usually refers to cytotoxic drugs which affect rapidly dividing cells in general, in contrast with targeted therapy. Chemotherapy drugs interfere with cell division in various possible ways, e.g., with the duplication of DNA or the separation of newly formed chromosomes. Most forms of chemotherapy target all rapidly dividing cells and are not

specific for cancer cells, although some degree of specificity may come from the inability of many cancer cells to repair DNA damage, while normal cells generally can.

[000272] Examples of chemotherapeutic agents used in cancer therapy include, for example, antimetabolites (e.g., folic acid, purine, and pyrimidine derivatives) and alkylating agents (e.g., nitrogen mustards, nitrosoureas, platinum, alkyl sulfonates, hydrazines, triazenes, aziridines, spindle poison, cytotoxic agents, topoisomerase inhibitors and others). Exemplary agents include Aclarubicin, Actinomycin, Alitretinon, Altretamine, Aminopterin, Aminolevulinic acid, Amrubicin, Amsacrine, Anagrelide, Arsenic trioxide, Asparaginase, Atrasentan, Belotecan, Bexarotene, endamustine, Bleomycin, Bortezomib, Busulfan, Camptotnecin, Capecitabine, Carboplatin, Carboquone, Carmofur, Carmustine, Celecoxib, Chlorambucil, Chlormethine, Cisplatin, Cladribine, Clofarabine, Crisantaspase, Cyclophosphamide, Cytarabine, Dacarbazine, Dactinomycin, Daunorubicin, Decitabine, Demecolcine, Docetaxel, Doxorubicin, Etoposide, Etoposide, Floxuridine, Fludarabine, Fluorouracil (5FU), Fotemustine, Gemcitabine, Gliadel implants, Hydroxycarbamide, Hydroxyurea, idarubicin, Ifosfamide, Irinotecan, Irofulven, Ixabepilone, Larotaxel, Leucovorin, Liposomal doxorubicin, Liposomal daunorubicin, Lonidamine, Lomustine, Lucanthone, Mannosulfan, Masoprocil, Melphalan, Mercaptopurine, Mesna, Methotrexate, Methyl aminolevulinate, Mitobronitol, Mitoguanine, Mitotane, Mitomycin, Mitoxantrone, Nedaplatin, Nimustine, Oblimersen, Omacetaxine, Ortoposide, Oxaliplatin, Paclitaxel, Pegaspargase, Pemetrexed, Pentostatin, Pirarubicin, Pixantrone, Plicamycin, Porfimer sodium, Prednimustine, Procarbazine, Raltitrexed, Ranimustine, Rubitecan, Sapacitabine, Semustine, Sitimagene ceradenovec, Strataplatin, Streptozocin, Talaporfm, Tegafur-uracil, Temoporfin, Temozolomide, Teniposide, Tesetaxel, Testolactone, Tetranitrate, Thiotepa, Tiazofurine, Tioguanine, Tipifarnib, Topotecan, Trabectedin, Triaziquone, Triethylenemelamine, Triplatin, Tretinoin, Treosulfan, Trofosfamide, Uramustine, Valrubicin, Verteporfin, Vinblastine, Vincristine, Vindesine, Vinflunine, Vinorelbine, Vorinostat, Zorubicin, and other cytostatic or cytotoxic agents described herein.

[000273] Because some drugs work better together than alone, two or more drugs are often given at the same time or sequentially. Often, two or more chemotherapy agents are used as combination chemotherapy. In some embodiments, the chemotherapy agents (including combination chemotherapy) can be used in combination with a combination described herein.

Targeted Therapy

[000274] In some embodiments, a combination described herein is administered with a targeted therapy. Targeted therapy constitutes the use of agents specific for the deregulated proteins of cancer cells. Small molecule targeted therapy drugs are generally inhibitors of enzymatic domains on mutated, overexpressed, or otherwise critical proteins within the cancer cell. Prominent examples are the tyrosine kinase inhibitors such as Axitinib, Bosutinib, Cediranib, dasatinib, erlotinib, imatinib, gefitinib, lapatinib, Lestaurtinib, Nilotinib, Semaxanib, Sorafenib, Sunitinib, and Vandetanib, and also cyclin-dependent kinase inhibitors such as Alvocidib and Seliciclib. Monoclonal antibody therapy is another strategy in which the therapeutic agent is an antibody which specifically binds to a protein on the surface of the cancer cells. Examples include the anti-HER2/neu antibody trastuzumab (HERCEPTIN®) typically used in breast cancer, and the anti-CD20 antibody rituximab and Tositumomab typically used in a variety of B-cell malignancies. Other exemplary antibodies include Ctuximab, Panitumumab, Trastuzumab, Alemtuzumab, Bevacizumab, Edrecolomab, and Gemtuzumab. Exemplary fusion proteins include Aflibercept and Denileukin diftitox. In some embodiments, the targeted therapy can be used in combination with a combination described herein.

[000275] Targeted therapy can also involve small peptides as "homing devices" which can bind to cell surface receptors or affected extracellular matrix surrounding the tumor. Radionuclides which are attached to these peptides (e.g., RGDs) eventually kill the cancer cell if the nuclide decays in the vicinity of the cell. An example of such therapy includes BEXXAR®.

Immunotherapy

[000276] In some embodiments, a combination described herein is administered with an immunotherapy. Cancer immunotherapy refers to a diverse set of therapeutic strategies designed to induce the subject's own immune system to fight the tumor.

[000277] Contemporary methods for generating an immune response against tumors include intravesicular BCG immunotherapy for superficial bladder cancer, and use of interferons and other cytokines to induce an immune response in subjects with renal cell carcinoma and melanoma. Allogeneic hematopoietic stem cell transplantation can be considered a form of immunotherapy, since the donor's immune cells will often attack the tumor in a graft- versus-tumor effect. In some embodiments, the immunotherapy agents can be used in combination with a combination as described herein.

Hormonal Therapy

[000278] In some embodiments, a combination described is administered with a hormonal therapy. The growth of some cancers can be inhibited by providing or blocking certain hormones. Common examples of hormone-sensitive tumors include certain types of breast and prostate cancers. Removing or blocking estrogen or testosterone is often an important additional treatment. In certain cancers, administration of hormone agonists, such as progestogens may be therapeutically beneficial. In some embodiments, the hormonal therapy agents can be used in combination with a combination described herein.

[000279] In some embodiments, the additional agent is an agent that modifies ER, PR, and/or AR. For example, the additional agent is an AR antagonist, which includes, but is not limited to, flutamide, bicalutamide and nilutamide. In some embodiments, the additional agent is an agent that blocks estrogen or progesterone, which includes, aromatase inhibitors including but is not limited to, anastrozole, letrozole, and exemestane. In some embodiments, the additional agent is an estrogen receptor modulator including, but not limited to, fulvestrant, tamoxifen and raloxifene.

Radiation Therapy

[000280] The combinations described herein can be used in combination with directed energy or particle, or radioisotope treatments, e.g., radiation therapies, e.g., radiation oncology, for the treatment of proliferative disease, e.g., cancer, e.g., cancer associated with cancer stem cells. The combinations described herein may be administered to a subject simultaneously or sequentially along with the directed energy or particle, or radioisotope treatments. For example, the combinations described herein may be administered before, during, or after the directed energy or particle, or radioisotope treatment, or a combination thereof. The directed energy or particle therapy may comprise total body irradiation, local body irradiation, or point irradiation. The directed energy or particle may originate from an accelerator, synchrotron, nuclear reaction, vacuum tube, laser, or from a radioisotope. The therapy may comprise external beam radiation therapy, teletherapy, brachy therapy, sealed source radiation therapy, systemic radioisotope therapy, or unsealed source radiotherapy. The therapy may comprise ingestion of, or placement in proximity to, a radioisotope, e.g., radioactive iodine, cobalt, cesium, potassium, bromine, fluorine, carbon. External beam radiation may comprise exposure to directed alpha particles, electrons (e.g., beta particles), protons, neutrons, positrons, or photons (e.g., radiowave, millimeter wave, microwave, infrared, visible, ultraviolet, X-ray, or gamma-ray photons). The radiation may be directed at any portion of the subject in need of treatment.

Surgery

[000281] The combinations described herein can be used in combination with surgery, e.g., surgical exploration, intervention, biopsy, for the treatment of proliferative disease, e.g., cancer, e.g., cancer associated with cancer stem cells. The combinations described herein may be administered to a subject simultaneously or sequentially along with the surgery. For example, the combinations described herein may be administered before (preoperative), during, or after (post-operative) the surgery, or a combination thereof. The surgery may be a biopsy during which one or more cells are collected for further analysis. The biopsy may be accomplished, for example, with a scalpel, a needle, a catheter, an endoscope, a spatula, or scissors. The biopsy may be an excisional biopsy, an incisional biopsy, a core biopsy, or a needle biopsy, e.g., a needle aspiration biopsy. The surgery may involve the removal of localized tissues suspected to be or identified as being cancerous. For example, the procedure may involve the removal of a cancerous lesion, lump, polyp, or mole. The procedure may involve the removal of larger amounts of tissue, such as breast, bone, skin, fat, or muscle. The procedure may involve removal of part of, or the entirety of, an organ or node, for example, lung, throat, tongue, bladder, cervix, ovary, testicle, lymph node, liver, pancreas, brain, eye, kidney, gallbladder, stomach, colon, rectum, or intestine. In one embodiment, the cancer is breast cancer, e.g., triple negative breast cancer, and the surgery is a mastectomy or lumpectomy.

Anti-Inflammatory Agents

[000282] A combination described herein can be administered with an anti-inflammatory agent. Anti-inflammatory agents can include, but are not limited to, non-steroidal anti-inflammatory agents (e.g., Salicylates (Aspirin (acetylsalicylic acid), Diflunisal, Salsalate), Propionic acid derivatives (Ibuprofen, Naproxen, Fenoprofen, Ketoprofen, Flurbiprofen, Oxaprozin, Loxoprofen), Acetic acid derivatives (Indomethacin, Sulindac, Etodolac, Ketorolac, Diclofenac, Nabumetone), Enolic acid (Oxicam) derivatives (Piroxicam, Meloxicam, Tenoxicam, Droxicam, Lomoxicam, Isoxicam), Fenamic acid derivatives (Fenamates) (Mefenamic acid, Meclofenamic acid, Flufenamic acid, Tolfenamic acid). Selective COX -2 inhibitors (Coxibs) (Celecoxib), Sulphonanilides (Nimesulide). Steroids (e.g. Hydrocortisone (Cortisol), Cortisone acetate, Prednisone, Prednisolone, Methylprednisolone, Dexamethasone, Betamethasone, Triamcinolone, Beclometasone, Fludrocortisone acetate, Deoxycorticosterone acetate, Aldosterone).

Analgesic Agents

[000283] Analgesics can include but are not limited to, opiates (e.g. morphine, codeine, oxycodone, hydrocodone, dihydromorphine, pethidine, buprenorphine, tramadol, venlafaxine), paracetomal and Nonsteroidal anti-inflammatory agents (e.g., Salicylates (Aspirin (acetylsalicylic acid), Diflunisal, Salsalate), Propionic acid derivatives (Ibuprofen, Naproxen, Fenoprofen, Ketoprofen, Flurbiprofen, Oxaprozin, Loxoprofen), Acetic acid derivatives (Indomethacin, Sulindac, Etodolac, Ketorolac, Diclofenac, Nabumetone), Enolic acid (Oxicam) derivatives (Piroxicam, Meloxicam, Tenoxicam, Droxicam, Lomoxicam, Isoxicam), Fenamic acid derivatives (Fenamates)(Mefenamic acid, Meclofenamic acid, Flufenamic acid. Tolfenamic acid). Selective COX-2 inhibitors (Coxibs) (Ceiecoxib), Sulphonanilides (Nimesulide).

Antiemetic Agents

[000284] A combination described herein can be administered with an antiemetic agent. Antiemetic agents can include, but are not limited to, 5-HT₃ receptor antagonists (Dolasetron (Anzemet), Granisetron (Kytril, Sancuso), Ondansetron (Zofran), Tropisetron (Navoban), Palonosetron (Aloxi), Mirtazapine (Remeron)), Dopamine antagonists (Domperidone, Olanzapine, Droperidol, Haloperidol, Chlorpromazine, Promethazine, Prochlorperazine, Metoclopramide (Reglan), Alizapride, Prochlorperazine (Compazine, Stemzine, Buccastem, Stemetil, Phenotil), NK₁ receptor antagonist (Aprepitant (Emend), Antihistamines (Cyclizine, Diphenhydramine (Benadryl), Dimenhydrinate (Gravol, Dramamine), Meclozine (Bonine, Antivert), Promethazine (Pentazine, Phenergan, Promacot), Hydroxyzine), benzodiazapines (Lorazepam, Midazolam), Anticholinergics (hyoscine), steroids (Dexamethasone).

Combinations

[000285] The phrase, "in combination with," and the terms "co-administration," "co-administering," or "co-providing", as used herein in the context of the administration of a compound described herein or a therapy described herein, means that two (or more) different compounds or therapies are delivered to the subject during the course of the subject's affliction with the disease or disorder (e.g., a disease or disorder as described herein, e.g., cancer), e.g., two (or more) different compounds or therapies are delivered to the subject after the subject has been diagnosed with the disease or disorder (e.g., a disease or disorder as described herein, e.g., cancer) and before the disease or disorder has been cured or eliminated or treatment has ceased for other reasons.

[000286] In some embodiments, the delivery of one compound or therapy is still occurring when the delivery of the second begins, so that there is overlap in terms of administration. This is sometimes referred to herein as "simultaneous" or "concurrent delivery." In other embodiments, the delivery of one compound or therapy ends before the delivery of the other compound or therapy begins. In some embodiments of either case, the treatment (e.g., administration of compound, composition, or therapy) is more effective because of combined administration. For example, the second compound or therapy is more effective, e.g., an equivalent effect is seen with less of the second compound or therapy, or the second compound or therapy reduces symptoms to a greater extent, than would be seen if the second compound or therapy were administered in the absence of the first compound or therapy, or the analogous situation is seen with the first compound or therapy. In some embodiments, delivery is such that the reduction in a symptom, or other parameter related to the disorder is greater than what would be observed with one compound or therapy delivered in the absence of the other. The effect of the two compounds or therapies can be partially additive, wholly additive, or great than additive (e.g., synergistic). The delivery can be such that the first compound or therapy delivered is still detectable when the second is delivered.

[000287] In some embodiments, the first compound or therapy and second compound or therapy can be administered simultaneously (e.g., at the same time), in the same or in separate compositions, or sequentially. Sequential administration refers to administration of one compound or therapy before (e.g., immediately before, less than 5, 10, 15, 30, 45, 60 minutes; 1, 2, 3, 4, 6, 8, 10, 12, 16, 20, 24, 48, 72, 96 or more hours; 4, 5, 6, 7, 8, 9 or more days; 1, 2, 3, 4, 5, 6, 7, 8 or more weeks before) administration of an additional, e.g., secondary, compound or therapy. The order of administration of the first and secondary compound or therapy can also be reversed.

[000288] The combinations described herein can be a first line treatment for abnormal cell growth, e.g., cancer, i.e., it is used in a subject who has not been previously administered another drug intended to treat the cancer; a second line treatment for the cancer, i.e., it is used in a subject in need thereof who has been previously administered another drug intended to treat the cancer; a third or fourth treatment for the cancer, i.e., it is used in a subject who has been previously administered two or three other drugs intended to treat the cancer.

Administration and Dosage

[000289] The combinations of this invention may be administered orally, parenterally, topically, rectally, or via an implanted reservoir, preferably by oral administration or administration by injection. In some cases, the pH of the composition (e.g., pharmaceutical composition) may be adjusted with pharmaceutically acceptable acids, bases or buffers to enhance the stability or efficacy of the composition.

[000290] In some embodiments, the subject is administered the composition (e.g., pharmaceutical composition) orally. In some embodiments the composition (e.g., pharmaceutical composition) is orally administered in any orally acceptable dosage form including, but not limited to, liqui-gel tablets or capsules, syrups, emulsions and aqueous suspensions. Liqui-gels may include gelatins, plasticisers, and/or opacifiers, as needed to achieve a suitable consistency and may be coated with enteric coatings that are approved for use, e.g., shellacs. Additional thickening agents, for example gums, e.g., xanthum gum, starches, e.g., corn starch, or glutens may be added to achieve a desired consistency of the composition (e.g., pharmaceutical composition) when used as an oral dosage. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

[000291] In some embodiments, the subject is administered the composition (e.g., pharmaceutical composition) in a form suitable for oral administration such as a tablet, capsule, pill, powder, sustained release formulations, solution, and suspension. The composition (e.g., pharmaceutical composition) may be in unit dosage forms suitable for single administration of precise dosages. Pharmaceutical compositions may comprise, in addition to a compound as described herein a pharmaceutically acceptable carrier, and may optionally further comprise one or more pharmaceutically acceptable excipients, such as, for example, stabilizers, diluents, binders, and lubricants. In addition, the tablet may include other medicinal or pharmaceutical agents, carriers, and or adjuvants. Exemplary pharmaceutical compositions include compressed tablets (e.g., directly compressed tablets).

[000292] Tablets are also provided comprising the active or therapeutic ingredient (e.g., compound as described herein). In addition to the active or therapeutic ingredients, tablets may contain a number of inert materials such as carriers. Pharmaceutically acceptable carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, sesame oil and the like. Saline solutions and aqueous dextrose can also be employed as liquid carriers. Oral dosage forms for use in accordance with the present invention thus may be formulated in conventional manner using one or more

pharmaceutically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active ingredients into preparations which, can be used pharmaceutically.

[000293] Excipients can impart good powder flow and compression characteristics to the material being compressed. Examples of excipients are described, for example, in the Handbook of Pharmaceutical Excipients (5th edition), Edited by Raymond C Rowe, Paul J. Sheskey, and Sian C. Owen; Publisher: Pharmaceutical Press.

[000294] For oral administration, the active ingredients, e.g., the compound as described herein can be formulated readily by combining the active ingredients with pharmaceutically acceptable carriers well known in the art. Such carriers enable the active ingredients of the invention to be formulated as tablets, pills, capsules, liquids, gels, syrups, slurries, powders or granules, suspensions or solutions in water or non-aqueous media, and the like, for oral ingestion by a subject. Pharmacological preparations for oral use can be made using a solid excipient, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries if desired, to obtain, for example, tablets. Suitable excipients such as diluents, binders or disintegrants may be desirable.

[000295] The dosage may vary depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the subject's condition. (See e.g., Fingl, et al., 1975, in 'The Pharmacological Basis of Therapeutics'). Lower or higher doses than those recited above may be required. Specific dosage and treatment regimens for any particular subject will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health status, sex, diet, time of administration, rate of excretion, drug combination, the severity and course of the disease, condition or symptoms, the subject's disposition to the disease, condition or symptoms, and the judgment of the treating physician. A course of therapy can comprise one or more separate administrations of a compound as described herein. A course of therapy can comprise one or more cycles of a compound as described herein.

[000296] In some embodiments, a cycle, as used herein in the context of a cycle of administration of a drug, refers to a period of time for which a drug is administered to a subject. For example, if a drug is administered for a cycle of 21 days, the periodic administration, e.g., daily or twice daily, is given for 21 days. A drug can be administered for more than one cycle. Rest periods may be interposed between cycles. A rest cycle may be 1, 2, 4, 6, 8, 10, 12, 16, 20, 24 hours, 1, 2, 3, 4, 5, 6, 7 days, or 1, 2, 3, 4 or more weeks in length.

[000297] Oral dosage forms may, if desired, be presented in a pack or dispenser device, such as an FDA approved kit, which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser may also be accompanied by a notice associated with the container in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions or human or veterinary administration. Such notice, for example, may be of labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert.

EXAMPLES

[000298] In order that the invention described herein may be more fully understood, the following examples are set forth. The examples described in this application are offered to illustrate the pharmaceutical compositions and methods provided herein and are not to be construed in any way as limiting their scope.

Example 1. Synergy of Dual RAF/MEK Inhibitor and BRAF V600 inhibitors or pan-RAF inhibitors in Cancer Cell Lines

Materials and Methods

3D proliferation assays in vitro

[000299] As examples, BRAF V600E human melanoma, BRAF V600E human colorectal carcinoma (CRC), and NRAS mutant human melanoma cell lines were grown in 3D conditions. Briefly, 96-well plates were coated with 50 μ L of Matrigel (100%) and incubated at 37°C and 5% CO₂ for 30 min in order for the Matrigel to solidify. Cells were seeded in 100 μ L of 2% Matrigel containing medium. After an overnight incubation (17-22 hours), cells were treated with VS-6766 +/- BRAF V600 inhibitor (e.g., vemurafenib, dabrafenib, encorafenib), VS-6766 +/- pan-RAF inhibitor (e.g., belvarafenib, naporafenib, lifirafenib, tovorafenib), or MEK inhibitor (e.g., cobimetinib, trametinib, binimetinib) +/- BRAF (e.g., vemurafenib, dabrafenib, encorafenib) inhibitor for 7 days. VS-6766 and MEK inhibitors were used at 1:5 dilutions starting at 5 μ M. BRAF V600 inhibitors and pan-RAF inhibitors were used at 1:3 dilutions starting at 5 μ M. Cell viability was measured using the cell viability CellTiter-Glo assay.

Synergy analysis

[000300] Raw data and metadata files were processed with a custom R-script for single agent and combination activity. Bliss, Loewe, Highest Single Agent (HSA) and ZIP synergy analysis were performed to generate a composite synergy score. Summary graphics and reports were saved for visualization and further analysis.

Results

[000301] 3D proliferation assays were performed to determine whether dual RAF/MEK inhibitors (e.g., VS-6766) augment the anti-proliferative activity of a BRAF V600 inhibitor (e.g., vemurafenib, dabrafenib, encorafenib) in a panel of BRAF V600 cancer cell lines (such as 5 BRAF V600E melanoma (**FIGS. 1-3**) and 3 BRAF V600E colorectal carcinoma (CRC) cell lines (**FIGS. 4-6**)). Synergy scores were calculated using a combination of 4 different methods (Bliss, Loewe, HSA and ZIP). As shown in **FIGS. 1-6**, the dual RAF/MEK inhibitor (e.g., VS-6766) was synergistic with BRAF V600 inhibitors in reducing cell viability of a panel of BRAF V600E melanoma and CRC cell lines. Furthermore, the synergy between the dual RAF/MEK inhibitor (e.g., VS-6766) + BRAF V600 inhibitors was compared to the synergy between a MEK inhibitor (e.g., cobimetinib, trametinib, binimetinib) + BRAF V600 inhibitor. It was found that synergy of the dual RAF/MEK inhibitor (e.g., VS-6766) + BRAF V600 inhibitor was better than synergy of MEK inhibitors + BRAF V600 inhibitor. These results support the clinical evaluation of the dual RAF/MEK inhibitor (e.g., VS-6766) in combination with BRAF V600 inhibitor for cancers harboring BRAF V600 mutations such as, but not limited to, BRAF V600E.

[000302] 3D proliferation assays were performed to determine whether a dual RAF/MEK inhibitor (e.g., VS-6766) augments the anti-proliferative activity of a pan-RAF inhibitor (e.g., belvarafenib, naporafenib, lifirafenib, tovorafenib) in a panel of mutant cell lines (such as 3 NRAS mutant melanoma cell lines (**FIGS. 7, 8**)). Synergy scores were calculated using a combination of 4 different methods (Bliss, Loewe, HSA and ZIP). As shown in **FIGS. 7, 8**, dual RAF/MEK inhibitor (e.g., VS-6766) was synergistic with pan-RAF inhibitors in reducing cell viability of a panel of NRAS mutant melanoma cell lines. These results support the clinical evaluation of VS-6766 in combination with pan-RAF inhibitors for cancers harboring RAS mutations, such as but not limited to NRAS mutations.

Equivalents and Scope

[000303] In the claims articles such as “a,” “an,” and “the” may mean one or more than one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention includes embodiments in which more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process.

[000304] Furthermore, the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, and descriptive terms from one or more of the listed claims is introduced into another claim. For example, any claim that is dependent on another claim can be modified to include one or more limitations found in any other claim that is dependent on the same base claim. Where elements are presented as lists, e.g., in Markush group format, each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements and/or features, some embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements and/or features. For purposes of simplicity, those embodiments have not been specifically set forth *in haec verba* herein. It is also noted that the terms “comprising” and “containing” are intended to be open and permits the inclusion of additional elements or steps. Where ranges are given, endpoints are included. Furthermore, unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or sub-range within the stated ranges in different embodiments of the invention, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.


[000305] This application refers to various issued patents, published patent applications, journal articles, and other publications, all of which are incorporated herein by reference. If there is a conflict between any of the incorporated references and the instant specification, the specification shall control. In addition, any particular embodiment of the present invention that falls within the prior art may be explicitly excluded from any one or more of the claims. Because such embodiments are deemed to be known to one of ordinary skill in the art, they

may be excluded even if the exclusion is not set forth explicitly herein. Any particular embodiment of the invention can be excluded from any claim, for any reason, whether or not related to the existence of prior art.

[000306] Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation many equivalents to the specific embodiments described herein. The scope of the present embodiments described herein is not intended to be limited to the above Description, but rather is as set forth in the appended claims. Those of ordinary skill in the art will appreciate that various changes and modifications to this description may be made without departing from the spirit or scope of the present invention, as defined in the following claims.


CLAIMS

1. A method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a BRAF V600 inhibitor, wherein the cancer is identified as having a BRAF V600 mutation.
2. The method of claim 1, wherein the BRAF V600 mutation is BRAF V600E, BRAF V600K, BRAF V600D, BRAF V600R, and/or BRAF V600M mutation.
3. The method of claim 2, wherein the BRAF V600 mutation is BRAF V600E mutation.
4. The method of claim 2, wherein the BRAF V600 mutation is BRAF V600K mutation.
5. The method of claim 2, wherein the BRAF V600 mutation is BRAF V600D mutation.
6. The method of claim 2, wherein the BRAF V600mutation is BRAF V600R mutation.
7. The method of claim 2, wherein the BRAF V600mutation is BRAF V600M mutation.
8. The method of any one of claims 1-7, wherein the dual RAF/MEK inhibitor is a compound of formula (I):



(I), or a pharmaceutically acceptable salt thereof.

9. The method of claim 8, wherein the dual RAF/MEK inhibitor is a compound of formula (I):



(I).


10. The method of claim 8, wherein the dual RAF/MEK inhibitor is a potassium salt of the compound of formula (I).

11. The method of any one of claims 1-10, wherein the dual RAF/MEK inhibitor is orally administered to the subject.
12. The method of any one of claims 1-11, wherein the dual RAF/MEK inhibitor is administered twice a week.
13. The method of any one of claims 1-12, wherein the dual RAF/MEK inhibitor is administered at a dose of 0.5 mg to about 10 mg per administration.
14. The method of claim 13, wherein the dual RAF/MEK inhibitor is dosed at 3.2 mg per administration.
15. The method of claim 13, wherein the dual RAF/MEK inhibitor is dosed at 4 mg per administration.
16. The method of any one of claims 1-15, wherein the dual RAF/MEK inhibitor is dosed as a cycle comprising administering the dual RAF/MEK inhibitor for three weeks and then not administering the dual RAF/MEK inhibitor for one week.
17. The method of any one of claims 1-16, wherein the BRAF V600 inhibitor is dabrafenib, encorafenib, vemurafenib, FORE-8394 (PLX-8394), tinloragenib, AZ-304, agerafenib, KIN-2787, BGB-3245, ABM-1310, TQB-3233, UB-941, AFX-1251, ARQ 736, ASN003, AVB-BRAF, BDTX-4933, CFT1946, HLX208, RO5212054, RO7276389, or TQ-B3233, or pharmaceutically acceptable salts thereof.
18. The method of any one of claims 1-17, wherein the BRAF V600 inhibitor is dabrafenib, encorafenib, or vemurafenib, or pharmaceutically acceptable salts thereof.
19. The method of any one of claims 1-18, wherein the BRAF V600 inhibitor is orally administered to the subject.
20. The method of any one of claims 1-19, wherein the BRAF V600 inhibitor is administered once daily.
21. The method of any one of claims 1-19, wherein the BRAF V600 inhibitor is administered twice daily.

22. The method of any one of claims 1-19, wherein the BRAF V600 inhibitor is dosed as a cycle comprising administering the BRAF V600 inhibitor for three weeks and then not administering the BRAF V600 inhibitor for one week.
23. The method of any one of claims 1-22, wherein the BRAF V600 inhibitor is dosed at 10 mg to 2000 mg per administration.
24. The method of any one of claims 1-23, wherein the BRAF V600 inhibitor is dosed at 100 mg to 1000 mg per administration.
25. The method of any one of claims 1-24, wherein the cancer is pancreatic cancer, gynecologic cancer (e.g., cervical cancer, ovarian cancer, uterine cancer, vaginal cancer, endometrial cancer, or vulvar cancer), liver cancer, prostate cancer, mesothelioma, breast cancer, bladder cancer, melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer.
26. The method of any one of claims 1-25, wherein the cancer is melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer.
27. The method of claim 25 or 26, wherein the lung cancer is non-small cell lung cancer (e.g., metastatic non-small cell lung cancer).
28. The method of claim 25 or 26, wherein the melanoma is unresectable melanoma or metastatic melanoma.
29. The method of claim 25 or 26, wherein the cancer is colorectal cancer.
30. The method of claim 25 or 26, wherein the thyroid cancer is papillary thyroid cancer, follicular thyroid cancer, or anaplastic thyroid cancer.
31. A method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a pan-RAF inhibitor, wherein the cancer is identified as having ARAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation.


32. The method of claim 31, wherein the pan-RAF inhibitor is BAL3833, LY3009120, REDX05358, IRICoR-Ipsen, JZP815, METiS-01, QLH11906, or SJ-C1044, or pharmaceutically acceptable salts thereof.
33. The method of claim 31 or 32, wherein the pan-RAF inhibitor is orally administered to the subject.
34. The method of any one of claims 31-33, wherein the pan-RAF inhibitor is administered once daily.
35. The method of any one of claims 31-34, wherein the pan-RAF inhibitor is administered twice daily.
36. The method of any one of claims 31-35, wherein the pan-RAF inhibitor is dosed at 10 mg to 2000 mg per administration.
37. The method of any one of claims 31-36, wherein the pan-RAF inhibitor is dosed at 100 mg to 1000 mg per administration.
38. A method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a CRAF inhibitor, wherein the cancer is identified as having ARAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation.
39. The method of claim 38 wherein the CRAF inhibitor is MG005, Quanta-RAF1, or STX200, or pharmaceutically acceptable salts thereof.
40. The method of claim 38 or 39, wherein the CRAF inhibitor is orally administered to the subject.
41. The method of any one of claims 38-40, wherein the CRAF inhibitor is administered once daily.
42. The method of any one of claims 38-41, wherein the CRAF inhibitor is administered twice daily.

43. The method of any one of claims 38-42, wherein the CRAF inhibitor is dosed at 10 mg to 2000 mg per administration.
44. The method of any one of claims 38-43, wherein the CRAF inhibitor is dosed at 100 mg to 1000 mg per administration.
45. A method of treating a cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of a dual RAF/MEK inhibitor and an effective amount of a RAF inhibitor, wherein the cancer is identified as having RAF, BRAF, CRAF, KRAS, NRAS, HRAS, EGFR, ALKR, FGFR, PDGFR, NF1, SOS1, SOS2, MEK1, and/or MEK2 mutation.
46. The method of claim 45 wherein the RAF inhibitor is donafenib, lifirafenib, rigosertib, BMS-908662, XP-102, ABM-2526, DDC-PanRAF, FNX006, or VRN-XX, or pharmaceutically acceptable salts thereof.
47. The method of claim 45 or 46, wherein the RAF inhibitor is orally administered to the subject.
48. The method of any one of claims 45-47, wherein the RAF inhibitor is administered once daily.
49. The method of any one of claims 45-48, wherein the RAF inhibitor is administered twice daily.
50. The method of any one of claims 45-49, wherein the RAF inhibitor is dosed at 10 mg to 2000 mg per administration.
51. The method of any one of claims 45-50, wherein the RAF inhibitor is dosed at 100 mg to 1000 mg per administration.
52. The method of any one of claims 31-51, wherein the dual RAF/MEK inhibitor is a compound of formula (I):



(I), or a pharmaceutically acceptable salt thereof.

53. The method of claim 52, wherein the dual RAF/MEK inhibitor is a compound of formula (I):



(I).

54. The method of claim 52, wherein the dual RAF/MEK inhibitor is a potassium salt of the compound of formula (I).

55. The method of any one of claims 31-54, wherein the dual RAF/MEK inhibitor is orally administered to the subject.

56. The method of any one of claims 31-55, wherein the dual RAF/MEK inhibitor is administered twice a week.

57. The method of any one of claims 31-56, wherein the dual RAF/MEK inhibitor is administered at a dose of 0.5 mg to about 10 mg per administration.

58. The method of claim 57, wherein the dual RAF/MEK inhibitor is dosed at 3.2 mg per administration.

59. The method of claim 57, wherein the dual RAF/MEK inhibitor is dosed at 4 mg per administration.

60. The method of any one of claims 31-59, wherein the dual RAF/MEK inhibitor is dosed as a cycle comprising administering the dual RAF/MEK inhibitor for three weeks and then not administering the dual RAF/MEK inhibitor for one week.

61. The method of any one of claims 31-60, wherein the cancer is pancreatic cancer, gynecologic cancer (e.g., cervical cancer, ovarian cancer, uterine cancer, vaginal cancer,

endometrial cancer, or vulvar cancer), liver cancer, prostate cancer, mesothelioma, breast cancer, bladder cancer, melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer.

62. The method of any one of claims 31-61, wherein the cancer is melanoma, lung cancer, colorectal cancer, thyroid cancer, glioblastoma, or renal cancer.

63. The method of claim 61 or 62, wherein the lung cancer is non-small cell lung cancer (e.g., metastatic non-small cell lung cancer).

64. The method of claim 61 or 62, wherein the melanoma is unresectable melanoma or metastatic melanoma.

65. The method of claim 61 or 62, wherein the cancer is colorectal cancer.

66. The method of claim 61 or 62, wherein the thyroid cancer is papillary thyroid cancer, follicular thyroid cancer, or anaplastic thyroid cancer.

67. The method of any one of claims 1-66, further comprising administering to the subject an effective amount of a FAK inhibitor.

68. The method of claim 67, wherein the FAK inhibitor is defactinib, or a pharmaceutically acceptable salt thereof.

69. The method of claim 67 or 68, wherein the FAK inhibitor is dosed at about 100 mg to about 1000 mg.

70. The method of claim 69, wherein the FAK inhibitor is dosed at about 100 mg to about 400 mg per administration.

71. The method of claim 70, wherein the FAK inhibitor is dosed at 200 mg per administration.

72. The method of claim 70, wherein the FAK inhibitor is dosed at 400 mg per administration.

73. The method of any one of claims 67-72, wherein the FAK inhibitor is administered once daily.

74. The method of any one of claims 67-72, wherein the FAK inhibitor is administered twice daily.
75. The method of any one of claims 67-74, wherein the FAK inhibitor is dosed as a cycle, comprising administering the FAK inhibitor for three weeks and then not administering the FAK inhibitor for one week.
76. The method of any one of claims 67-75, wherein the FAK inhibitor is orally administered to the subject.




FIG. 1




FIG. 2




FIG. 3






FIG. 4








FIG. 7




FIG. 8

			Synergy Score					
			VS-6766 + BRAFi			MEKi + BRAFi		
Cell line	MAPK alteration	PAM alterations	VS-6766 + vemurafenib	VS-6766 + dabrafenib	VS-6766 + encorafenib	cobimetinib + vemurafenib	trametinib + dabrafenib	binimetinib + encorafenib
A2058	BRAF V600E	PIK3R6	17.615	15.797	17.747	6.682	-5.699	13.419
IGR-1	BRAF V600E	PIK3C2G, PIK3R5, PIK3CG	16.434	9.543	13.119	14.073	-4.811	9.565
SK-MEL-5	BRAF V600E	PIK3CB, PIK3C2A, RICTOR	12.431	3.829	9.094	0.564	-7.886	8.691
WM-266-4	BRAF V600E		6.114	4.522	6.603	1.568	1.025	3.527
C32	BRAF V600E	PIK2CB, mTOR	11.793	9.287	9.268	3.14	0.466	7.802



FIG. 1