

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

26 March 2020 (26.03.2020)



(10) International Publication Number

WO 2020/059894 A1

(51) International Patent Classification:

A61K 31/4545 (2006.01) A61P 25/02 (2006.01)

A61K 45/06 (2006.01) A61P 25/06 (2006.01)

A61K 9/00 (2006.01)

Published:

— with international search report (Art. 21(3))

(21) International Application Number:

PCT/JP2019/038061

(22) International Filing Date:

19 September 2019 (19.09.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/733,927 20 September 2018 (20.09.2018) US

(71) Applicant: TAKEDA PHARMACEUTICAL COMPAN-

NY LIMITED [JP/JP]; 1-1, Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka, 5410045 (JP).

(72) Inventors: ASGHARNEJAD, Mahnaz; c/o Millennium

Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Massachusetts, 02139 (US). ARKILO, Dimitrios; c/o Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Massachusetts, 02139 (US). NISHI, Toshiya; c/o Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Massachusetts, 02139 (US).

(74) Agent: TAKASHIMA, Hajime; Meiji Yasuda Seimei Os-

aka Midotsuji Bldg., 1-1, Fushimimachi 4-chome, Chuo-ku, Osaka-shi, Osaka, 5410044 (JP).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: CH24H Inhibitors For Pain Use

(57) Abstract: The present disclosure relates to methods of treating pain by administration of (4-benzyl-4-hydroxypiperidin-1-yl) (2, 4'-bipyridin-3-yl) methanone or a pharmaceutically acceptable salt thereof.



WO 2020/059894 A1

DESCRIPTION

Title of the Invention: CH24H Inhibitors For Pain Use

Technical Field

5 [0001]

The present invention relates to (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof for use in treatment of pain.

10 [0002]

(Background of the Invention)

Complex regional pain syndrome (CRPS) was initially identified by Claude Bernard (1813-1878) when he noted that extreme pain could be associated with abnormalities in the
15 autonomic nervous system. Mitchell (1829-1914) coined the term 'causalgia' to describe this constellation occurring in veterans of the American Civil War. The term 'reflex sympathetic dystrophy' was used until recently when CRPS was agreed upon. CRPS can be subdivided into 3 types: (1) CRPS-I: previously known as reflex sympathetic dystrophy; (2) CRPS-II, previously known as causalgia and defined as CRPS with clinical and/or electrodiagnostic evidence of nerve damage; and 3) CRPS (not otherwise specified) which only partially meets diagnostic
20 criteria, but no better diagnosis can be discerned. CRPS-I and CRPS-II have similar outcomes and response to pain medication. Autonomic changes are often required for the diagnosis and may distinguish between acute CRPS ('hot' limb with edema and red coloration) and chronic ('cold' limb with atrophy and blue coloration). Although there is significant involvement of the
30 peripheral nervous system, chronic CRPS is thought to be a 'brain disease' with demonstrated alterations in both central nervous system (CNS) function as well as structural changes including alterations in cortical representations in both sensory and motor cortex. CRPS is known as one of the most

painful disorders and the risk of suicide is significantly higher in patients with CRPS with one study demonstrating that 75% of patients had a high risk for suicide. Amputation has rarely been considered as an option for treatment. Although the pathophysiology has not been established, overactivity of the *N*-methyl-D-aspartate (NMDA) receptors are thought to play a role and while no drugs are approved for CRPS, ketamine, a NMDA receptor antagonist, has established efficacy in a randomized control trial. Use of an NMDA antagonist along with morphine, decreased pain and cerebral pain representation consistent with brain involvement in CRPS.

Summary of the Invention

[0003]

Aspects of this disclosure relate to a method of treating pain in a mammal comprising administering an effective amount of a composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof to the mammal in need thereof. The pain optionally includes inflammatory pain, neuropathic pain, cancer inflammatory pain, post/perioperative pain, and idiopathic pain which is pain of unknown origin, for example, phantom limb pain.

In some embodiments, the pain is Complex Regional Pain Syndrome (CRPS).

In some embodiments, the pain is migraine.

[0004]

Neuropathic pain is caused by injury or infection of peripheral sensory nerves. It includes, but is not limited to pain from peripheral nerve trauma, herpes virus infection, diabetes mellitus, causalgia, plexus avulsion, neuroma, limb amputation, and vasculitis.

Neuropathic pain is also caused by nerve damage from chronic alcoholism, human immunodeficiency virus infection, hypothyroidism, uremia, or vitamin deficiencies.

Neuropathic pain includes, but is not limited to pain caused by nerve injury such as, for example, "the pain diabetics suffer from", burn pain, gout pain, osteoarthritic pain, trigeminal neuralgia pain, acute herpetic and
5 postherpetic pain, causalgia pain, chronic pain, diabetic neuropathic pain, fibromyalgia pain, neuropathic pain associated with diabetic peripheral neuropathy, post herpetic neuralgia, fibromyalgia, neuropathic pain associated with
10 spinal cord injury, Complex Regional Pain Syndrome (CRPS), migraine, cluster headache, labor pain, pruritus, and bone cancer pain.

[0005]

In addition, aspects of this disclosure relate to a method of treating autoimmune disease, cardiovascular disease,
15 diabetic disease, digestive organ disease, ophthalmologic disease or cancer disease in a mammal comprising administering an effective amount of a composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof to the mammal in need
20 thereof.

[0006]

The autoimmune disease includes rheumatoid arthritis, psoriasis, inflammatory bowel disease (e.g., Crohn's disease, ulcerative colitis etc.), Sjogren's syndrome, Behcet's disease,
25 multiple sclerosis, systemic lupus erythematosus, lupus nephritis, discoid lupus erythematosus, Castleman's disease, ankylopoietic spondylarthritis, polymyositis, dermatomyositis (DM), polyarteritis nodosa (PN), mixed connective tissue disease (MCTD), scleroderma, profundus lupus erythematosus,
30 chronic thyroiditis, Graves' disease, autoimmune gastritis, type I diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, graft versus host disease, Addison's disease, abnormal immunoresponse, arthritis,

dermatitis, radiodermatitis, primary biliary cirrhosis, urinary incontinence, chronic cough, asthma, cystitis, and the like.

[0007]

The cardiovascular disease includes hypertension, blood pressure circadian rhythm abnormality (e.g., early-morning hypertension, nocturnal hypertension etc.), heart diseases (e.g., cardiac hypertrophy, acute heart failure, chronic heart failure including cardiac failure, impaired vasodilation, cardiac myopathy, angina pectoris, myocarditis, atrial fibrillation, arrhythmia, tachycardia, myocardial infarction etc.), cerebrovascular disorders (e.g., asymptomatic cerebrovascular disorder, transient cerebral ischemia, cerebral apoplexy, cerebrovascular dementia, hypertensive encephalopathy, cerebral infarction etc.), cerebral edema, cerebral circulatory disorder, recurrence and sequela of cerebrovascular disorders (e.g., neurotic symptom, psychic symptom, subjective symptom, disorder in daily living activities etc.), ischemic peripheral circulation disorder, myocardial ischemia, venous insufficiency, progression of cardiac insufficiency after myocardial infarction, renal diseases (e.g., nephritis, glomerulonephritis, glomerulosclerosis, renal failure, thrombotic vasculopathy, complication of dialysis, organ dysfunction including nephropathy by radiation damage etc.), arteriosclerosis including atherosclerosis (e.g., aneurysm, coronary sclerosis, cerebral arteriosclerosis, peripheral arterial sclerosis etc.), vascular hypertrophy, vascular hypertrophy or obliteration and organ disorders after intervention (e.g., percutaneous transluminal coronary angioplasty, stenting, coronary angiography, intravascular ultrasound, intracoronary thrombolysis etc.), vascular re-obliteration and restenosis after bypass, polycythemia, hypertension, organ disorder and vascular hypertrophy after transplantation, rejection after transplantation, thrombosis, multiple organ disorder, endothelial dysfunction, hypertensive tinnitus, other

cardiovascular diseases (e.g., deep vein thrombosis, obstructive peripheral circulatory disorder, arteriosclerosis obliterans, thromboangiitis obliterans, ischemic cerebral circulatory disorder, Raynaud's disease, Buerger's disease
5 etc.), metabolic and/or nutritional disorders (e.g., obesity, hyperlipemia, hypercholesterolemia, hyperuricacidemia, hyperkalemia, hypernatremia etc.), and the like.

[0008]

The diabetic disease includes diabetes mellitus (e.g.,
10 type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes mellitus, obese diabetes mellitus), obesity (e.g., malignant mastocytosis, exogenous obesity, hyperinsular obesity, hyperplasmic obesity, hypophyseal adiposity, hypoplasmic obesity, hypothyroid obesity, hypothalamic obesity,
15 symptomatic obesity, infantile obesity, upper body obesity, alimentary obesity, hypogonadal obesity, systemic mastocytosis, simple obesity, central obesity and the like), hyperphagia, hyperlipidemia/dyslipidemia (e.g., hypertriglyceridemia, hypercholesterolemia, high LDL-cholesterolemia, low HDL-
20 cholesterolemia, postprandial hyperlipemia), diabetic complications [e.g., neuropathy, nephropathy, retinopathy, diabetic cardiomyopathy, cataract, macroangiopathy, osteopenia, hyperosmolar diabetic coma, infectious disease (e.g., respiratory infection, urinary tract infection,
25 gastrointestinal infection, dermal soft tissue infections, inferior limb infection), diabetic gangrene, xerostomia, hypacusis, cerebrovascular disorder, peripheral blood circulation disorder], metabolic syndrome (disease states having 3 or more selected from hypertriglycerid(TG)emia, low
30 HDL cholesterol(HDL-C)emia, hypertension, abdominal obesity and impaired glucose tolerance), and the like.

[0009]

The digestive organ disease includes an irritable bowel syndrome, inflammatory intestine disease, inflammatory bowel

disease, ulcerative colitis, Crohn's disease, diseases caused by a spiral urease-positive gram-negative bacterium (Helicobacter pylori, etc.) (gastritis, gastric ulcer, etc.), gastric cancer, postgastrostomy disorder, indigestion, 5 esophageal ulcer, pancreatitis, polyp of the colon, cholelithiasis, hemorrhoids, peptic ulcer, situational ileitis, gluttony, constipation, diarrhea, borborygmus, non-alcoholic fatty liver disease, visceral pain, gastrointestinal disorder, esophagitis, etc.

10 [0010]

The ophthalmologic disease includes terygium, spring catarrh, dry eye, superficial punctate keratopathy and the like.

[0011]

The cancer disease includes colorectal cancer (e.g. colon 15 cancer, rectal cancer, anal cancer, familial colorectal cancer, hereditary nonpolyposis colorectal cancer and gastrointestinal stromal tumor), lung cancer (e.g. non-small-cell lung cancer, small cell lung cancer and malignant mesothelioma), mesothelioma, pancreatic cancer (e.g. pancreatic ductal 20 carcinoma and pancreatic endocrine tumor), pharyngeal cancer, laryngeal cancer, esophageal cancer, stomach cancer (e.g. papillary adenocarcinoma, mucous adenocarcinoma and adenosquamous carcinoma), duodenal carcinoma, small intestinal cancer, breast cancer (e.g. infiltrating duct carcinoma, 25 noninfiltrating intraductal carcinoma and inflammatory breast cancer), ovarian cancer (e.g. epithelial ovarian cancer, extragonadal germ cell tumor, ovarian germ cell tumor and ovarian low malignant potential tumor), testicular tumor, prostate cancer (e.g. hormone-dependent prostate cancer, 30 hormone-independent prostate cancer and castration-resistant prostate cancer), liver cancer (e.g. hepatic cell carcinoma, primary hepatic cancer and cancer of extrahepatic bile duct), thyroid cancer (e.g. thyroid medullary carcinoma), kidney cancer (e.g. renal cell carcinoma (e.g. clear cell type renal

cell carcinoma) and transitional cell carcinoma of renal pelvis and ureter), uterine cancer (e.g. cervical cancer, corpus uteri cancer and uterine sarcoma), gestational choriocarcinoma, brain tumor (e.g. medulloblastoma, glioma, pineal astrocytoma, 5 pilocytic astrocytoma, diffuse astrocytoma, anaplastic astrocytoma and pituitary adenoma), retinoblastoma, skin cancer (e.g. basal cell carcinoma and malignant melanoma), sarcoma (e.g. rhabdomyosarcoma, leiomyosarcoma, soft tissue sarcoma and spindle cell sarcoma), malignant bone tumor, bladder cancer, 10 blood cancer (e.g. multiple myeloma, leukemia (e.g. acute myeloid leukemia), malignant lymphoma, Hodgkin's disease and chronic myeloproliferative disease), and cancer of unknown primary, and the like.

[0012]

15 In some embodiments, administering the effective amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof results in (i) a reduction in 24(S)-hydroxycholesterol (24HC) levels and/or (ii) a reduction in 20 NMDA receptor function.

[0013]

In some embodiments, the effective amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt 25 thereof is administered orally or parenterally (e.g., topically, rectally, intravenously, etc.).

[0014]

In some embodiments, the effective amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'- 30 bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof is administered as a single unit dose. The single unit dose is preferably about 0.01 mg/kg to 100 mg/kg, more preferably about 0.05 mg/kg to 100 mg/kg, still more preferably about 0.1 mg/kg to 10 mg/kg.

[0015]

In some embodiments, the effective amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof is administered according to a dose regimen.

[0016]

In some embodiments, the mammal is a human. In further embodiments, the human is an adolescent or an adult.

[0017]

In some embodiments, the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof further comprises a pharmaceutically acceptable carrier.

[0018]

In some embodiments, the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof further comprises an additional active agent. Non-limiting examples of additional active agents include: sumatriptan, gabapentin, naratriptan, rizatriptan, ibuprofen, naproxen, dihydroergotamine, amitriptyline, venlafaxine, duloxetine, atenolol, metoprolol, nadolol, propranolol, timolol, propranolol, topiramate, valproate, sodium valproate, erenumab, one-botulinum toxin A(buspirone hydrochloride, tandospirone citrate, osemototan hydrochloride etc.). Further examples of additional active agents include ketamine, biphosphonates (e.g., risedronate, zoledronate), calcitonin, opioids (e.g., oxycodone, morphine, hydrocodone, fentanyl).

[0019]

Aspects of this disclosure also relate to a pharmaceutical composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, which is for treating pain.

[0020]

Aspects of this disclosure also relate to (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof for use in treatment
5 of pain.

[0021]

(Detailed Description of the Invention)

Aspects of the disclosure relate to a method of treating pain in a mammal (e.g., human, bovine, horse, dog, cat, monkey,
10 mouse, rat; preferably human) comprising administering an effective amount of a composition comprising (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof to the mammal in need thereof. The pain optionally includes inflammatory pain,
15 neuropathic pain, cancer inflammatory pain, post/perioperative pain, and idiopathic pain which is pain of unknown origin, for example, phantom limb pain.

The pain includes preferably Complex Regional Pain Syndrome (CRPS) and migraine, more preferably Complex Regional
20 Pain Syndrome (CRPS).

[0022]

In addition, aspects of this disclosure relate to a method of treating autoimmune disease, cardiovascular disease, diabetic disease, digestive organ disease, ophthalmologic
25 disease or cancer disease in a mammal comprising administering an effective amount of a composition comprising (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof to the mammal in need thereof.

30 [0023]

The autoimmune disease includes rheumatoid arthritis, psoriasis, inflammatory bowel disease (e.g., Crohn's disease, ulcerative colitis etc.), Sjogren's syndrome, Behcet's disease, multiple sclerosis, systemic lupus erythematosus, lupus

nephritis, discoid lupus erythematosus, Castleman's disease, ankylopoietic spondylarthritis, polymyositis, dermatomyositis (DM), polyarteritis nodosa (PN), mixed connective tissue disease (MCTD), scleroderma, profundus lupus erythematosus, 5 chronic thyroiditis, Graves' disease, autoimmune gastritis, type I diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, graft versus host disease, Addison's disease, abnormal immunoreponse, arthritis, 10 dermatitis, radiodermatitis, primary biliary cirrhosis, urinary incontinence, chronic cough, asthma, cystitis, and the like.

[0024]

The cardiovascular disease includes hypertension, blood pressure circadian rhythm abnormality (e.g., early-morning 15 hypertension, nocturnal hypertension etc.), heart diseases (e.g., cardiac hypertrophy, acute heart failure, chronic heart failure including cardiac failure, impaired vasodilation, cardiac myopathy, angina pectoris, myocarditis, atrial fibrillation, arrhythmia, tachycardia, myocardial infarction 20 etc.), cerebrovascular disorders (e.g., asymptomatic cerebrovascular disorder, transient cerebral ischemia, cerebral apoplexy, cerebrovascular dementia, hypertensive encephalopathy, cerebral infarction etc.), cerebral edema, cerebral circulatory disorder, recurrence and sequela of cerebrovascular disorders 25 (e.g., neurotic symptom, psychic symptom, subjective symptom, disorder in daily living activities etc.), ischemic peripheral circulation disorder, myocardial ischemia, venous insufficiency, progression of cardiac insufficiency after myocardial infarction, renal diseases (e.g., nephritis, glomerulonephritis, 30 glomerulosclerosis, renal failure, thrombotic vasculopathy, complication of dialysis, organ dysfunction including nephropathy by radiation damage etc.), arteriosclerosis including atherosclerosis (e.g., aneurysm, coronary sclerosis, cerebral arteriosclerosis, peripheral arterial sclerosis etc.),

vascular hypertrophy, vascular hypertrophy or obliteration and organ disorders after intervention (e.g., percutaneous transluminal coronary angioplasty, stenting, coronary angioscopy, intravascular ultrasound, intracoronary
5 thrombolysis etc.), vascular re-obliteration and restenosis after bypass, polycythemia, hypertension, organ disorder and vascular hypertrophy after transplantation, rejection after transplantation, thrombosis, multiple organ disorder, endothelial dysfunction, hypertensive tinnitus, other
10 cardiovascular diseases (e.g., deep vein thrombosis, obstructive peripheral circulatory disorder, arteriosclerosis obliterans, thromboangiitis obliterans, ischemic cerebral circulatory disorder, Raynaud's disease, Buerger's disease etc.), metabolic and/or nutritional disorders (e.g., obesity,
15 hyperlipemia, hypercholesterolemia, hyperuricacidemia, hyperkalemia, hypernatremia etc.), and the like.

[0025]

The diabetic disease includes diabetes mellitus (e.g., type 1 diabetes mellitus, type 2 diabetes mellitus, gestational
20 diabetes mellitus, obese diabetes mellitus), obesity (e.g., malignant mastocytosis, exogenous obesity, hyperinsular obesity, hyperplasmic obesity, hypophyseal adiposity, hypoplasmic obesity, hypothyroid obesity, hypothalamic obesity, symptomatic obesity, infantile obesity, upper body obesity,
25 alimentary obesity, hypogonadal obesity, systemic mastocytosis, simple obesity, central obesity and the like), hyperphagia, hyperlipidemia/dyslipidemia (e.g., hypertriglyceridemia, hypercholesterolemia, high LDL-cholesterolemia, low HDL-cholesterolemia, postprandial hyperlipemia), diabetic
30 complications [e.g., neuropathy, nephropathy, retinopathy, diabetic cardiomyopathy, cataract, macroangiopathy, osteopenia, hyperosmolar diabetic coma, infectious disease (e.g., respiratory infection, urinary tract infection, gastrointestinal infection, dermal soft tissue infections,

inferior limb infection), diabetic gangrene, xerostomia, hypacusis, cerebrovascular disorder, peripheral blood circulation disorder], metabolic syndrome (disease states having 3 or more selected from hypertriglycerid(TG)emia, low HDL cholesterol(HDL-C)emia, hypertension, abdominal obesity and impaired glucose tolerance), and the like.

[0026]

The digestive organ disease includes an irritable bowel syndrome, inflammatory intestine disease, inflammatory bowel disease, ulcerative colitis, Crohn's disease, diseases caused by a spiral urease-positive gram-negative bacterium (*Helicobacter pylori*, etc.) (gastritis, gastric ulcer, etc.), gastric cancer, postgastrostomy disorder, indigestion, esophageal ulcer, pancreatitis, polyp of the colon, cholelithiasis, hemorrhoids, peptic ulcer, situational ileitis, gluttony, constipation, diarrhea, borborygmus, non-alcoholic fatty liver disease, visceral pain, gastrointestinal disorder, esophagitis, etc.

[0027]

The ophthalmologic disease includes terygium, spring catarrh, dry eye, superficial punctate keratopathy and the like.

[0028]

The cancer disease includes colorectal cancer (e.g. colon cancer, rectal cancer, anal cancer, familial colorectal cancer, hereditary nonpolyposis colorectal cancer and gastrointestinal stromal tumor), lung cancer (e.g. non-small-cell lung cancer, small cell lung cancer and malignant mesothelioma), mesothelioma, pancreatic cancer (e.g. pancreatic ductal carcinoma and pancreatic endocrine tumor), pharyngeal cancer, laryngeal cancer, esophageal cancer, stomach cancer (e.g. papillary adenocarcinoma, mucous adenocarcinoma and adenosquamous carcinoma), duodenal carcinoma, small intestinal cancer, breast cancer (e.g. infiltrating duct carcinoma, noninfiltrating intraductal carcinoma and inflammatory breast

cancer), ovarian cancer (e.g. epithelial ovarian cancer, extragonadal germ cell tumor, ovarian germ cell tumor and ovarian low malignant potential tumor), testicular tumor, prostate cancer (e.g. hormone-dependent prostate cancer, 5 hormone-independent prostate cancer and castration-resistant prostate cancer), liver cancer (e.g. hepatic cell carcinoma, primary hepatic cancer and cancer of extrahepatic bile duct), thyroid cancer (e.g. thyroid medullary carcinoma), kidney cancer (e.g. renal cell carcinoma (e.g. clear cell type renal 10 cell carcinoma) and transitional cell carcinoma of renal pelvis and ureter), uterine cancer (e.g. cervical cancer, corpus uteri cancer and uterine sarcoma), gestational choriocarcinoma, brain tumor (e.g. medulloblastoma, glioma, pineal astrocytoma, pilocytic astrocytoma, diffuse astrocytoma, anaplastic 15 astrocytoma and pituitary adenoma), retinoblastoma, skin cancer (e.g. basal cell carcinoma and malignant melanoma), sarcoma (e.g. rhabdomyosarcoma, leiomyosarcoma, soft tissue sarcoma and spindle cell sarcoma), malignant bone tumor, bladder cancer, blood cancer (e.g. multiple myeloma, leukemia (e.g. acute 20 myeloid leukemia), malignant lymphoma, Hodgkin's disease and chronic myeloproliferative disease), and cancer of unknown primary, and the like.

[0029]

Not to be bound by theory, it is believed that (4-benzyl- 25 4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof can reduce NMDA receptor function by blocking the production of 24(S)-hydroxycholesterol (24HC or cerebrosterol), which is the product of CH24H enzymatic activity and a potent (EC50 = 1.2 30 μM) and specific positive allosteric modulator of NDMA receptors. By blocking the production of 24(S)-hydroxycholesterol, specific inhibitors of CH24H have the potential to reduce excitatory neurotransmission through NMDA receptors. Accordingly, the CH24H inhibitors disclosed herein

are hypothesized to have the same or similar mechanism as ketamine, an NMDA receptor open channel blocker, and, accordingly, confer effects, specifically in a mammal suffering from any one of inflammatory pain, neuropathic pain, cancer inflammatory pain, post/perioperative pain, or idiopathic pain which is pain of unknown origin, for example, phantom limb pain.

Not to be bound by theory, these effects are believed to be dose dependent.

[0030]

Accordingly, in some embodiments, administering the effective amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof results in (i) a reduction in 24(S)-hydroxycholesterol (24HC) levels and/or (ii) a reduction in NMDA receptor function.


[0031]

For instance, a reduction in 24(S)-hydroxycholesterol (24HC) levels, recited in (i) above, may be measured by an LC/MS assay. Similarly, a reduction in NMDA receptor function, recited in (ii) above, may be measured electrophysiologically ex-vivo after administration or likely in dissociated neuronal cultures.

[0032]

In addition to (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone, the CH24H inhibitor is a compound represented by the formula (I):

[0033]




[0034]

wherein

X¹ is a carbon atom or a nitrogen atom;

Ring A is

[0035]



[0036]

each of which is optionally further substituted and optionally bridged;

R¹ is an optionally substituted C₁₋₆ alkyl group, an optionally substituted hydroxy group, an optionally substituted amino group, an optionally substituted carbocyclic group, or an optionally substituted heterocyclic group, or R¹ is optionally bonded to the atom on Ring A to form, together with Ring A, a spiro ring or a fused ring, each of which is substituted by an oxo group and optionally further substituted;

10 15

R² is an optionally substituted C₆₋₁₄ aryl group, or an optionally substituted aromatic heterocyclic group; and

R³ is a hydrogen atom or a substituent when X¹ is a carbon atom, or absent when X¹ is a nitrogen atom, (tert-butyl 4-(4-phenylpyrimidin-5-yl)piperazine-1-carboxylate is excluded) or a salt thereof has the same efficacy of (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone.

20

[0037]

A variety of compounds represented by formula (I) are provided in US 2015/0315209, the entirety of which is incorporated by reference herein. Further non-limiting examples of a CH24H inhibitor are (2R)-1-((1-(4-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)piperidin-4-yl)carbonyl)pyrrolidine-2-carbonitrile and a pharmaceutically acceptable salts thereof.

25 30

A variety of compounds represented by formula (I) may be a hydrate or a non-hydrate.

[0038]

Definitions

Technical and scientific terms used herein have the meanings commonly understood by one of ordinary skill in the art to which the present invention pertains, unless otherwise defined. Reference is made herein to various methodologies known to those of ordinary skill in the art. Publications and other materials setting forth such known methodologies to which reference is made are incorporated herein by reference in their entireties as though set forth in full. Any suitable materials and/or methods known to those of ordinary skill in the art can be utilized in carrying out the present invention. However, specific materials and methods are described. Materials, reagents, and the like to which reference are made in the following description and examples are obtainable from commercial sources, unless otherwise noted.

[0039]

The following terms are used throughout as defined below.

[0040]

As used herein, singular articles such as "a" and "an" and "the" and similar referents denote both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. Disclosure of ranges of values should be understood as a shorthand reference to each separate value falling within the range and at the endpoints, again unless otherwise clearly indicated, and each separate value is to be understood as being described herein as if it were individually set forth.

[0041]

All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context.

[0042]

The use of any and all examples, or exemplary language

("e.g." or "such as") herein, is intended merely to better illuminate the embodiments and does not pose a limitation on the scope of the claims unless otherwise stated. No language in the specification should be construed as indicating any non-
5 claimed feature as essential.

[0043]

As used herein, "about" will be understood by persons of ordinary skill in the art and will vary to some extent depending upon the context in which it is used. If there are
10 uses of the term which are not clear to persons of ordinary skill in the art, given the context in which it is used, "about" will mean up to plus or minus 10% of the particular term.

[0044]


15 As used herein, the term "active agent" means a biologically active component of a pharmaceutical composition, e.g. an agent that has a pharmacological effect on the subject or patient to which it is administered. Contemplated herein are (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-
20 yl)methanone or a pharmaceutically acceptable salt thereof and additional active agents for use in combination with (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof. Non-limiting examples of suitable additional active agents for use in
25 combination with the CH24H inhibitors include those mentioned above.

[0045]

The compound (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone, shown below as Formula (II) is
30 described in US Patent No. 8,648,079:

[0046]

Formula (II)



[0047]

5 The compound of Formula (II) is a CH24H inhibitor. (4-Benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof is safely administered to a mammal, preferably human.

[0048]

10 As used herein, the term "effective amount" means an amount effective to successfully achieve a particular biological effect. In the present case, the effective amount is an amount to effective to treat pain which includes inflammatory pain, neuropathic pain, cancer inflammatory pain, 15 post/perioperative pain, and idiopathic pain which is pain of unknown origin, for example, phantom limb pain are included especially.

Neuropathic pain is caused by injury or infection of peripheral sensory nerves. It includes, but is not limited to 20 pain from peripheral nerve trauma, herpes virus infection, diabetes mellitus, causalgia, plexus avulsion, neuroma, limb amputation, and vasculitis.

Neuropathic pain is also caused by nerve damage from chronic alcoholism, human immunodeficiency virus infection, 25 hypothyroidism, uremia, or vitamin deficiencies.

Neuropathic pain includes, but is not limited to pain caused by nerve injury such as, for example, "the pain diabetics suffer from", burn pain, gout pain, osteoarthritic pain, trigeminal neuralgia pain, acute herpetic and

postherpetic pain, causalgia pain, chronic pain, diabetic neuropathic pain, fibromyalgia pain, neuropathic pain associated with diabetic peripheral neuropathy, post herpetic neuralgia, fibromyalgia, neuropathic pain associated with spinal cord injury, Complex Regional Pain Syndrome (CRPS) and migraine. Suitable effective amounts may be determined according to methods well known in the art to determine single unit dosage and/or dose regimens.

[0049]

10 In the present another case, the effective amount is an amount to effective to treat autoimmune disease, cardiovascular disease, diabetic disease, digestive organ disease, ophthalmologic disease or cancer disease.

[0050]

15 The autoimmune disease includes rheumatoid arthritis, psoriasis, inflammatory bowel disease (e.g., Crohn's disease, ulcerative colitis etc.), Sjogren's syndrome, Behcet's disease, multiple sclerosis, systemic lupus erythematosus, lupus nephritis, discoid lupus erythematosus, Castleman's disease, ankylopoietic spondylarthritis, polymyositis, dermatomyositis (DM), polyarteritis nodosa (PN), mixed connective tissue disease (MCTD), scleroderma, profundus lupus erythematosus, chronic thyroiditis, Graves' disease, autoimmune gastritis, type I diabetes, autoimmune hemolytic anemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, graft versus host disease, Addison's disease, abnormal immunoresponse, arthritis, dermatitis, radiodermatitis, primary biliary cirrhosis, urinary incontinence, chronic cough, asthma, cystitis, and the like.

30 [0051]

The cardiovascular disease includes hypertension, blood pressure circadian rhythm abnormality (e.g., early-morning hypertension, nocturnal hypertension etc.), heart diseases (e.g., cardiac hypertrophy, acute heart failure, chronic heart

failure including cardiac failure, impaired vasodilation, cardiac myopathy, angina pectoris, myocarditis, atrial fibrillation, arrhythmia, tachycardia, myocardial infarction etc.), cerebrovascular disorders (e.g., asymptomatic
5 cerebrovascular disorder, transient cerebral ischemia, cerebral apoplexy, cerebrovascular dementia, hypertensive encephalopathy, cerebral infarction etc.), cerebral edema, cerebral circulatory disorder, recurrence and sequela of cerebrovascular disorders (e.g., neurotic symptom, psychic symptom, subjective symptom,
10 disorder in daily living activities etc.), ischemic peripheral circulation disorder, myocardial ischemia, venous insufficiency, progression of cardiac insufficiency after myocardial infarction, renal diseases (e.g., nephritis, glomerulonephritis, glomerulosclerosis, renal failure, thrombotic vasculopathy,
15 complication of dialysis, organ dysfunction including nephropathy by radiation damage etc.), arteriosclerosis including atherosclerosis (e.g., aneurysm, coronary sclerosis, cerebral arteriosclerosis, peripheral arterial sclerosis etc.), vascular hypertrophy, vascular hypertrophy or obliteration and
20 organ disorders after intervention (e.g., percutaneous transluminal coronary angioplasty, stenting, coronary angiography, intravascular ultrasound, intracoronary thrombolysis etc.), vascular re-obliteration and restenosis after bypass, polycythemia, hypertension, organ disorder and
25 vascular hypertrophy after transplantation, rejection after transplantation, thrombosis, multiple organ disorder, endothelial dysfunction, hypertensive tinnitus, other cardiovascular diseases (e.g., deep vein thrombosis, obstructive peripheral circulatory disorder, arteriosclerosis
30 obliterans, thromboangiitis obliterans, ischemic cerebral circulatory disorder, Raynaud's disease, Buerger's disease etc.), metabolic and/or nutritional disorders (e.g., obesity, hyperlipemia, hypercholesterolemia, hyperuricacidemia, hyperkalemia, hypernatremia etc.), and the like.

[0052]

The diabetic disease includes diabetes mellitus (e.g., type 1 diabetes mellitus, type 2 diabetes mellitus, gestational diabetes mellitus, obese diabetes mellitus), obesity (e.g., malignant mastocytosis, exogenous obesity, hyperinsular obesity, hyperplasmic obesity, hypophyseal adiposity, hypoplasmic obesity, hypothyroid obesity, hypothalamic obesity, symptomatic obesity, infantile obesity, upper body obesity, alimentary obesity, hypogonadal obesity, systemic mastocytosis, simple obesity, central obesity and the like), hyperphagia, hyperlipidemia/dyslipidemia (e.g., hypertriglyceridemia, hypercholesterolemia, high LDL-cholesterolemia, low HDL-cholesterolemia, postprandial hyperlipemia), diabetic complications [e.g., neuropathy, nephropathy, retinopathy, diabetic cardiomyopathy, cataract, macroangiopathy, osteopenia, hyperosmolar diabetic coma, infectious disease (e.g., respiratory infection, urinary tract infection, gastrointestinal infection, dermal soft tissue infections, inferior limb infection), diabetic gangrene, xerostomia, hypacusis, cerebrovascular disorder, peripheral blood circulation disorder], metabolic syndrome (disease states having 3 or more selected from hypertriglycerid(TG)emia, low HDL cholesterol(HDL-C)emia, hypertension, abdominal obesity and impaired glucose tolerance), and the like.

[0053]

The digestive organ disease includes an irritable bowel syndrome, inflammatory intestine disease, inflammatory bowel disease, ulcerative colitis, Crohn's disease, diseases caused by a spiral urease-positive gram-negative bacterium (Helicobacter pylori, etc.) (gastritis, gastric ulcer, etc.), gastric cancer, postgastrostomy disorder, indigestion, esophageal ulcer, pancreatitis, polyp of the colon, cholelithiasis, hemorrhoids, peptic ulcer, situational ileitis, gluttony, constipation, diarrhea, borborygmus, non-alcoholic

fatty liver disease, visceral pain, gastrointestinal disorder, esophagitis, etc.

[0054]

The ophthalmologic disease includes terygium, spring
5 catarrh, dry eye, superficial punctate keratopathy and the like.

[0055]

The cancer disease includes colorectal cancer (e.g. colon cancer, rectal cancer, anal cancer, familial colorectal cancer, hereditary nonpolyposis colorectal cancer and gastrointestinal
10 stromal tumor), lung cancer (e.g. non-small-cell lung cancer, small cell lung cancer and malignant mesothelioma), mesothelioma, pancreatic cancer (e.g. pancreatic ductal carcinoma and pancreatic endocrine tumor), pharyngeal cancer, laryngeal cancer, esophageal cancer, stomach cancer (e.g.
15 papillary adenocarcinoma, mucous adenocarcinoma and adenosquamous carcinoma), duodenal carcinoma, small intestinal cancer, breast cancer (e.g. infiltrating duct carcinoma, noninfiltrating intraductal carcinoma and inflammatory breast cancer), ovarian cancer (e.g. epithelial ovarian cancer,
20 extragonadal germ cell tumor, ovarian germ cell tumor and ovarian low malignant potential tumor), testicular tumor, prostate cancer (e.g. hormone-dependent prostate cancer, hormone-independent prostate cancer and castration-resistant prostate cancer), liver cancer (e.g. hepatic cell carcinoma, primary hepatic cancer and cancer of extrahepatic bile duct),
25 thyroid cancer (e.g. thyroid medullary carcinoma), kidney cancer (e.g. renal cell carcinoma (e.g. clear cell type renal cell carcinoma) and transitional cell carcinoma of renal pelvis and ureter), uterine cancer (e.g. cervical cancer, corpus uteri
30 cancer and uterine sarcoma), gestational choriocarcinoma, brain tumor (e.g. medulloblastoma, glioma, pineal astrocytoma, pilocytic astrocytoma, diffuse astrocytoma, anaplastic astrocytoma and pituitary adenoma), retinoblastoma, skin cancer (e.g. basal cell carcinoma and malignant melanoma), sarcoma

(e.g. rhabdomyosarcoma, leiomyosarcoma, soft tissue sarcoma and spindle cell sarcoma), malignant bone tumor, bladder cancer, blood cancer (e.g. multiple myeloma, leukemia (e.g. acute myeloid leukemia), malignant lymphoma, Hodgkin's disease and
5 chronic myeloproliferative disease), and cancer of unknown primary, and the like. Suitable effective amounts may be determined according to methods well known in the art to determine single unit dosage and/or dose regimens.

[0056]

10 The term "single unit dose" in this context refers to an effective amount provided in a single administration. Non-limiting examples of suitable single unit doses for use in the claimed methods include about 0.01 to 100 mg/kg body weight, preferably 0.05 to 30 mg/kg body weight, more preferably 0.1 to
15 10 mg/kg body weight for oral administration to an adult patient (body weight 60 kg).

[0057]

The term "dose regimen" in this context refers to an effective amount provided over a fixed number of
20 administrations over a specified duration of time.

In some embodiments, the effective amount is administered according to a dose regimen of either twice a day (BID) or once a day (QD) dosing.

[0058]

25 It is appreciated that single unit doses may be tailored to the mammal being treated. For example, for humans, non-limiting exemplary single unit doses include less than about 1350 mg, between about 50 mg and about 800 mg (preferably between about 100 mg and about 800 mg), about 100 mg, about 150
30 mg, about 200 mg, about 300 mg, about 400 mg, about 600 mg or about 800 mg.

[0059]

Suitable routes of administration and doses and formulations suited thereto are known in the art. Non-limiting

examples of routes of administration relevant to the claimed methods include oral and parenteral (e.g., topical, rectal, or intravenous) routes. Examples of the dosage form suited for a particular route of administration include oral preparations
5 such as tablet (including sugar-coated tablet, film-coated tablet, sublingual tablet, orally disintegrating tablet), capsules (including soft capsule, microcapsule), granule, powder, troche, syrup, emulsion, suspension, films (e.g., orally disintegrable films) and the like; and parenteral
10 preparations such as injection (e.g., subcutaneous injection, intravenous injection, intramuscular injection, intraperitoneal injection, drip infusion), external preparations (e.g., dermal preparation, ointment), suppository (e.g., rectal suppository, vaginal suppository), pellet, nasal preparation, pulmonary
15 preparation (inhalant), eye drop and the like. Optionally, these preparations may be a release control preparation (e.g., sustained-release microcapsule) such as an immediate-release preparation or a sustained-release preparation.

[0060]

20 The term "pharmaceutically acceptable salt," as used herein, represents salts or zwitterionic forms of the compounds of the present technology which are water or oil-soluble or dispersible; which are suitable for treatment of diseases without undue toxicity, irritation, and allergic-response;
25 which are commensurate with a reasonable benefit/risk ratio; and which are effective for their intended use. The salts can be prepared during the final isolation and purification of the compounds or separately by reacting the appropriate compound in the form of the free base with a suitable acid. Representative
30 acid addition salts include acetate, adipate, alginate, L-ascorbate, aspartate, benzoate, benzenesulfonate (besylate), bisulfate, butyrate, camphorate, camphorsulfonate, citrate, digluconate, formate, fumarate, gentisate, glutarate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate,

hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate (isethionate), lactate, maleate, malonate, DL-mandelate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphonate, picrate, pivalate, propionate, pyroglutamate, succinate, sulfonate, tartrate, L-tartrate, trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, para-toluenesulfonate (p-tosylate), and undecanoate. Also, basic groups in the compounds of the present technology can be quaternized with methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dimethyl, diethyl, dibutyl, and diamyl sulfates; decyl, lauryl, myristyl, and steryl chlorides, bromides, and iodides; and benzyl and phenethyl bromides.

Examples of acids which can be employed to form pharmaceutically acceptable addition salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, and citric. Salts can also be formed by coordination of the compounds with an alkali metal or alkaline earth ion. Hence, the present technology contemplates sodium, potassium, magnesium, and calcium salts of the compounds of the present technology and the like.

[0061]

As used herein, the term "pharmaceutically acceptable carrier" means one or more organic or inorganic carrier substances conventionally used in the formulation of pharmaceutical compositions. Suitable pharmaceutically acceptable carriers can be determined by methods well known in the art e.g. excipients, lubricants, binders and disintegrants for solid preparations; solvents, solubilizing agents, suspending agents, isotonicity agents, buffers, and soothing agents for liquid preparations; and/or preparation additives such as preservatives, antioxidants, colorants, and sweetening

agents. Non-limiting examples of such suitable pharmaceutically acceptable carriers include:

for an excipient: lactose, sucrose, D-mannitol, D-sorbitol, starch, gelatinated starch, dextrin, crystalline
5 cellulose, low-substituted hydroxypropylcellulose, sodium carboxymethylcellulose, gum arabic, pullulan, light anhydrous silicic acid, synthesis aluminum silicate and magnesium alumino metasilicate;

for a lubricant: magnesium stearate, calcium stearate,
10 talc and colloidal silica;

for a binder: gelatinated starch, sucrose, gelatin, gum arabic, methylcellulose, carboxymethylcellulose, sodium carboxymethylcellulose, crystalline cellulose, sucrose, D-mannitol, trehalose, dextrin, pullulan, hydroxypropylcellulose,
15 hydroxypropylmethylcellulose and polyvinylpyrrolidone;

for a disintegrant: lactose, sucrose, starch, carboxymethylcellulose, calcium carboxymethylcellulose, croscarmellose sodium, sodium carboxymethyl starch, light anhydrous silicic acid and low-substituted
20 hydroxypropylcellulose;

for a solvent: water, physiological brine, Ringer's solution, alcohol, propylene glycol, polyethylene glycol, sesame oil, corn oil, olive oil and cottonseed oil;

for a solubilizing agent: polyethylene glycol, propylene
25 glycol, D-mannitol, trehalose, benzyl benzoate, ethanol, trisaminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate, sodium salicylate and sodium acetate;

for a suspending agent: surfactants such as
30 stearyltriethanolamine, sodium lauryl sulfate, lauryl aminopropionate, lecithin, benzalkonium chloride, benzethonium chloride, and glycerol monostearate; hydrophilic polymers such as polyvinyl alcohol, polyvinylpyrrolidone, sodium carboxymethylcellulose, methylcellulose, hydroxymethylcellulose,

hydroxyethylcellulose, hydroxypropylcellulose and the like;
polysorbates, and polyoxyethylene hydrogenated castor oil;

for an isotonicity agent: sodium chloride, glycerol, D-mannitol, D-sorbitol and glucose;

5 for a buffer: phosphate, acetate, carbonate, and citrate;

for a soothing agent: benzyl alcohol;

for a preservative: p-oxybenzoates, chlorobutanol, benzyl alcohol, phenethyl alcohol, dehydroacetic acid, and sorbic acid;

10 for an antioxidant: sulfite and ascorbate;

for a colorant: aqueous water-soluble food tar colors (e.g., food colors such as Food Color Red Nos. 2 and 3, Food Color Yellow Nos. 4 and 5, Food Color Blue Nos. 1 and 2 and the like), water insoluble lake dyes (e.g., aluminum salt of the
15 above-mentioned water-soluble food tar color), and natural dyes (e.g., β -carotene, chlorophyll, ferric oxide red); and

for a sweetening agent: saccharin sodium, dipotassium glycyrrhizinate, aspartame, and stevia.

[0062]

20 As used herein, the term "treating" includes the prevention, reduction, and/or complete resolution of the symptoms associated with or the cause of the target indication and/or a lessening of severity of the condition.

In some embodiments, (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically
25 acceptable salt thereof may be used in combination with other therapies including spinal code stimulators, sympathectomy and amputation.

Examples

30 [0063]

The present invention is explained in detail in the following by referring to Examples, which are not to be construed as limitative, and the invention may be changed within the scope of the present invention.

[0064]

Example 1 - Efficacy data in Preclinical CRPS model

The following Example provides data from a mouse model of chronic post-ischemic pain, which recapitulates symptoms seen in CRPS patients with peripheral tissue injury and subsequent progression of allodynia. (4-Benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone was tested in the CRPS model for its potential effects on pain. Following 3-hour ischemia and 0.5-hour reperfusion given at a hind limb, mice (C57) were orally treated with (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone at the dose of 30 mg/kg once daily for 7 days (n = 8). Paw withdraw threshold was evaluated with von Frey filaments as a behavioral index of mechanical allodynia at 0 h for the base line, 6 h, Days 1, 2, 4 and 7. The cumulative score of paw withdraw threshold over the study period was 17.7 in the sham-treated control group without ischemia/reperfusion injury. In the groups given ischemic injury, the cumulative score was 9.0 and 16.2 for the vehicle control and (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone group, respectively. It was hence indicated that allodynia threshold was markedly increased by treatment with (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone. These results support the use of (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone to treat the pain sensitization in CRPS.

[0065]

Example 2 - Efficacy data in Preclinical Migraine model

The following Example provides data from a rat cortical spreading depression (CSD) model, which recapitulates symptoms seen in migraine patients with aura. (2R)-1-((1-(4-(4-Methyl-1H-pyrazol-1-yl)pyridin-3-yl)piperidin-4-yl)carbonyl)pyrrolidine-2-carbonitrile was tested in the CSD model for its potential effects on hyperperfusion events and cortical direct current (DC) -potentials. Adult female Wistar

rats from Charles River Germany were used for the experiment. Animals were housed at a standard temperature ($22 \pm 1^\circ\text{C}$) and in a light-controlled environment (lights on from 7 am to 8 pm) with ad libitum access to food and water. Rats were orally
5 treated with vehicle (0.5% methylcellulose in water) or (2R)-1-
((1-(4-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)piperidin-4-
yl)carbonyl)pyrrolidine-2-carbonitrile at the dose of 10 mg/kg
once daily for 6 days ($n = 8$). One hour after the last dosing,
rats were anesthetized with 5% isoflurane (in 70% N_2O and 30%
10 O_2 ; flow 300 ml/min), and placed in a stereotactic frame.
During the operation and CSD the concentration of anesthetic
was reduced to 1-1.5%. The rectal temperature was maintained at
 $37.0 \pm 1.0^\circ\text{C}$ with a homeothermic blanket system. The skin was
opened by a medial incision and retracted laterally. Three burr
15 holes were drilled under saline cooling over the right
hemisphere at the following coordinates (mm from bregma): (1)
posterior 4.5, lateral, 2.0 (occipital cortex): KCl application
site; (2) posterior 0.5, lateral 2.0 (parietal cortex) (3)
anterior 2, lateral 2 (frontal cortex). A laser-Doppler flow
20 probe (Oxyflow, Oxford Optronics, UK) to monitor cerebral blood
flow (CBF) was placed in the parietal cortex burr holes on the
intact dura. An invasive Ag/AgCl electrode for measuring DC
potential shifts was placed in the frontal cortex burr holes on
the intact dura. For the DC-potential measurement, a reference
25 electrode was fixed in the neck. After surgical preparation,
the cortex was allowed to recover for 15 minutes under saline
irrigation. One molar KCl was placed on the pial surface and
kept moist by placing 5 μl of KCl solution every 15 minutes.
The number of KCl-induced CSDs was checked over the period of 2
30 hours and the number of CBF changes was also measured. The
average number of KCl-induced CSDs was 10.00 and 4.75 for the
vehicle-treated group and the (2R)-1-((1-(4-(4-methyl-1H-
pyrazol-1-yl)pyridin-3-yl)piperidin-4-yl)carbonyl)pyrrolidine-
2-carbonitrile -treated group, respectively. The average number

of CBF changes was 22.75 and 15.75 for the vehicle-treated group and the (2R)-1-((1-(4-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)piperidin-4-yl)carbonyl)pyrrolidine-2-carbonitrile - treated group, respectively. It was hence indicated that CSD events were markedly reduced by treatment with (2R)-1-((1-(4-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)piperidin-4-yl)carbonyl)pyrrolidine-2-carbonitrile. These results support the use of (2R)-1-((1-(4-(4-methyl-1H-pyrazol-1-yl)pyridin-3-yl)piperidin-4-yl)carbonyl)pyrrolidine-2-carbonitrile to treat migraine.

The efficacy of (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone in the treatment of migraine may be confirmed in the same manner as in Example 2.

[0066]

Example 3 - Efficacy, Safety and Tolerability of (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone as an Adjunctive Therapy in Adult Subjects With Chronic Complex Regional Pain Syndrome

This is a randomized, double-blind, placebo-controlled, parallel-group study in adult subjects (≥ 18 years) with chronic (symptoms ≥ 6 months) CRPS. The objective will be to look for a signal to determine if (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone as adjunctive therapy can reduce pain as measured by the NPS (an 11-point scale by electronic pain diary). This study will also evaluate efficacy of (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone as measured by PROMIS-29 version 2, PGIC, and CSS.

Approximately 24 subjects will be randomized to ensure 20 completers in the double-blind phase of the study. Randomization will be 2:1 (treatment: placebo).

This study consists of 2 parts:

Part A: Double-Blind Treatment

- 2- to 4-week screening/baseline period.
- 3-week titration period.

- 4-week maintenance period.
 - Follow-up/taper period if not rolling over into Part B
- Part B: Open-Label Extension
- 12-week open label.
- 5 - Follow-up/taper period.

[0067]

Part A: Double-Blind Treatment (9-13 weeks)

At the screening visit (Visit 1), after obtaining informed consent, subjects will undergo screening procedures to
10 assess subject eligibility in accordance with study entry
criteria. Subjects who fulfill the CRPS Budapest Criteria, and
have symptoms for ≥ 6 months and meet inclusion/exclusion
criteria at the screening/baseline visit will be eligible for
entry into the study. For a minimum of 6 of the last 7
15 screening days prior to enrollment into the study, baseline
current pain intensity will be collected 3 times a day to
provide an average daily 24-hour pain intensity (NPS; an 11-
point scale by electronic pain diary). The baseline will be
defined as the mean of the average screening 24-hour pain
20 intensity score for the last 7 days prior to the first dose.
During the Part A, average 24-hour pain intensity will be
collected 3 times a day. Pain intensity score collected during
the last 7 days prior to Day 14 and Day 49 (or the last dose in
Part A) will be used as to calculate the average 24-hour pain
25 intensity score for the primary endpoint analysis.

At the end of the prospective screening/baseline period,
subjects will return to the clinic (Visit 2, Day 1) and if a
subject does not meet the eligibility criteria the subject will
be discontinued from the study and considered a screen failure.

30 On Day 1, the subjects who meet the entry criteria will
be randomized in a 2:1 ratio to double-blind treatment with
investigational product (IP), either (4-benzyl-4-
hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone 100 mg
tablets or matching placebo, for 7 weeks (3-week titration

period and 4-week maintenance period).

All screening/baseline assessments will be collected prior to initiating treatment.

At Visit 2 (Day 1), after randomization and all predose
5 procedures have been performed, subjects will be started on 100
mg BID IP (either (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-
bipyridin-3-yl)methanone 100 mg tablets or matching placebo)
for approximately 1 week. The site will contact the subject to
determine safety and tolerability and the dose will be
10 increased 200 mg BID IP for approximately 1 week.

At Visit 3, safety and tolerability will be assessed and
if the drug is well tolerated and the subject continues to need
a higher dose, the dose will be increased to 300 mg BID. After
a week at 300 mg BID, the site will contact the subject to
15 determine safety and tolerability of this dose. After
completing 1 week at 300 mg BID, the 3-week titration period
will be completed and the 4-week maintenance period will begin.

If at any time during the 3-week titration period the
subject cannot tolerate the dose, the dose may be decreased to
20 the previous tolerated dose. If the subject cannot tolerate the
minimum daily dose of 100 mg BID, the subject will be withdrawn
from the study. If the dose is decreased due to tolerance,
based on the investigator's review, the dose may be increased
to the next highest dose 1 time during the titration period.
25 Dose modifications/up-titrations outside of this period should
be discussed with the medical monitor.

The dose at end of the titration period will be continued
through the maintenance period. During the 4-week maintenance
period, nonemergency IP dose changes should be discussed with
30 the sponsor/medical monitor prior to initiation. The subject
should be instructed not to alter their dose without prior
approval from the investigator. Any change in dose will be
documented in the subject's clinic chart and dosing card.

Pain medications and nondrug treatments must be stable

(regimented per prescription) for 1 month prior to screening and should remain stable throughout Part A. Pain medication use may be adjusted under supervision during Part B. Concurrent treatment regimen data will be collected throughout the study.

5 A single effective rescue medication must be identified for each subject for use during the study.

The prescribed maximum dose must remain stable during Part A. The use of rescue pain medications will be assessed at each visit; subjects requiring significant increase of rescue
10 medication (frequency or dose 50% over pre-enrollment levels or over the prescribed maximum) during Part A will be considered for withdrawal from the study at the investigator's discretion.

During Part A, all subjects will continue to enter the current pain intensity score as described 3 times a day using
15 the NPS in the electronic pain diary. This data will be captured daily during Part A.

An unblinded interim analysis will be conducted when all subjects have completed the double-blind Part A.

Following completion of Part A, subjects will have the
20 option to continue into Part B, a 12-week open-label extension study part, or to enter a double-blind taper period (maximum 6 days).

In all subjects choosing the double-blind taper period, the IP dose will be reduced to the next lower dose every 3 days
25 (or less frequently based on the investigator's discretion) until IP dose is discontinued. After tapering, the subjects will complete a safety follow-up phone call approximately 15 days after the last dose of IP.

[0068]

30 Part B Open-Label Extension (12 weeks):

Since the therapy assignment will remain blinded, all subjects who choose to continue into Part B will start at 200 mg BID (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone (100 mg tablets), regardless of the treatment they

were on in Part A. Subjects should remain at this dose for 48 hours after which the dose may be increased or decreased based on tolerability. The subject should be instructed not to alter their dose without prior approval from the investigator. Any
5 change in dose will be documented in the subject's clinic chart and dosing card. Subjects will visit the clinic approximately every 6 weeks. The current pain intensity will be collected 3 times a day using the electronic pain diary. From this data an average 24-hour pain intensity score can be calculated. In
10 addition, using the data collected in the last 7 days prior to Day 91 and Day 133 (or the last dose in Part B), the average pain score prior to these visits will be derived.

During the taper period, the IP dose will be reduced to the next lower dose every 3 days (or less frequently based on
15 the investigator's discretion) until IP dose is discontinued. After tapering, the subjects will complete a safety follow-up phone call approximately 15 days after the last dose of IP and exit the study.

The total study duration from screening to the last visit
20 in Part B will be approximately 6 months.

[0069]

Example 4 - A Randomized, Double-blind, Placebo-controlled, Parallel-group Study to Evaluate the Efficacy, Tolerability and Safety of (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone as Monotherapy for Prophylaxis in Adult Patients with Episodic Migraine with and/or without Aura.
25

This is a randomized, double-blind, placebo-controlled, parallel-group study in adult patients (18-65 years, inclusive) with episodic migraine with and/or without aura for at least 1
30 year prior to screening (as per ICHD-3 criteria) and with onset prior to age 50. The primary objective will be to determine if (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone, as monotherapy, can reduce the frequency of migraine headache days per 28 days. The study will evaluate

efficacy as the change from baseline in migraine headache days as recorded in headache diary. In addition, the study will evaluate the potential improvement after treatment with (4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone
5 in the frequency of the headache days, in the intensity of migraine headaches and headaches overall, and in the Patient Global Impression of Severity (PGI-S).

Approximately 40 patients will be randomized.

Randomization will be 1: 1 (20 treatment:20 placebo).

10 This study consists of 2 parts:

Part A: Double-blind

- o 4-week Screening/Baseline Period
- o 3-week Titration Period
- o 12-week Maintenance Period
- 15 o 1-week Down-Titration Period if not rolling over into part B

Part B: Open-label extension

- o 12-week open label
- o 1-week Down-Titration Period

20 [0070]

Part A: Double-blind Part (19-20 weeks)

At the Screening Visit (Visit 1), after obtaining informed consent, patients will undergo screening procedures to assess patient eligibility in accordance with study entry
25 criteria. Patients who have symptoms for at least 1 year and meet all inclusion/exclusion criteria at the Screening/Baseline Visit will be eligible for entry into the study.

At the end of the prospective Screening/Baseline Period, patients will return to the clinic (Visit 2, Day 1). Patients
30 who do not meet the eligibility criteria or did not comply with the data collection will be discontinued from the study and considered screen failure.

On Day 1, the patients who meet the entry criteria will be randomized in a 1:1 ratio to double-blind treatment with

Investigational Product (IP), either (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or matching placebo for 15 weeks (3-week Titration Period, 12-week Maintenance Period). All Screening/Baseline assessments will be
5 collected prior to initiating treatment.

At Visit 2 (Day 1), after randomization and all pre-dose procedures have been performed, patients will be started on 100 mg BID Investigational Product (IP) (either (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or
10 identical placebo) for 1 week. The site will contact the patient via phone to determine safety and tolerability (Visit 3, Day 7) and the dose will be increased to 200 mg BID IP for one week. At Visit 4 (Day 14), risk/benefit will be assessed and if the drug is well tolerated, the dose will be increased to 300
15 mg BID. After a week at 300 mg BID, the site will contact the patient via phone to determine safety and tolerability of this dose (Visit 5, Day 21). After one-week dosing at 300 mg BID, the 3-week Titration Period will be completed, and the 12-week Maintenance Period will begin. If at any time during the 3-week
20 Titration Period the patient cannot tolerate the dose, the dose may be decreased to the previous tolerated dose. If the patient cannot tolerate the minimum daily dose of 100 mg BID, the patient will be withdrawn from the study. If the dose has been decreased due to tolerance, based on the Investigator's review,
25 the dose may be increased to the next highest dose one time during the Titration Period. The IP dose selected at end of titration period will be continued through maintenance period. During the 12-week Maintenance Period, IP non-emergency dose changes should be discussed with the Sponsor/Medical Monitor
30 prior to initiation. The subject should be instructed not to alter their dose without prior approval from the investigator.

At Weeks 7 and 14, sites will contact patients by phone to monitor study drug compliance, concomitant medication use changes, AEs, and provide instructions regarding continued

dosing. Any change in dose will be documented in the patient's source doc and dosing card.

Migraine Specific Medications (MSM, rescue medication only), and non-drug treatments must be stable for 3 months prior to enrollment (screening) and not be altered during the treatment period. Concurrent treatment regimen data will be collected throughout the study.

An effective MSM must be identified for each patient for use during the trial. The prescribed maximum dose for the rescue medication must remain stable during part A. The use of MSM will be assessed at visits; patients requiring significant increase of MSM (frequency or dose 50% over pre-screening levels or over the prescribed maximum) will be considered for withdrawal from the study at the Investigator discretion.

All patients will continue to log each occurring headache, including intensity, and aura characteristics (if any) daily and will follow-up with in-person visits on 6-week intervals (Visit 6, Day 63 and Visit 7, Day 105).

Following completion of part A, subjects will have the option to continue into part B, a 12-week, open-label drug extension or to enter a 1-week double-blind taper period.

In all subjects choosing the taper period, the IP dose will be reduced to the next lower dose every 3 days (or less frequently based on the investigator's discretion) until IP dose is discontinued. After tapering, the subjects will complete a safety follow-up phone call approximately 14 days after the last dose of IP.

[0071]

Part B Open-label Extension (12 weeks):

Since the treatment assignments will remain blinded, all subjects who choose to continue into part B will start at 200 mg BID (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone regardless of the treatment they were on in part A. Subjects should remain at this dose for 48 hours after which

the dose may be increased to 300 mg BID or decreased to 100 mg BID based on tolerability via phone based on investigator discretion. Subjects will follow-up with in-person visits on 6-week intervals (Visit 8, Day 147 and Visit 9, Day 189). The
5 subject should be instructed not to alter their dose without prior approval from the investigator. Any change in dose will be documented in the subject's clinic chart and dosing card. All patients will continue to log each occurring headache, including intensity and aura characteristics (if any), daily.

10 During the taper period, the IP dose will be reduced to the next lower dose every 3 days (or less frequently based on the investigator's discretion) until IP dose is discontinued. After tapering, the subjects will complete a safety follow-up phone call approximately 14 days after the last dose of IP and
15 exit the study.

The total study duration to the last visit will be approximately 8 months.

[0072]

Procedures:

20 **Scales:** The migraine headache and headache frequency, intensity and aura characteristics (if any) will be measured with a headache diary daily from Visit 1 up to Visit 9. The PGI-S will be administered at Visits 2, 6, 7 and 9.

Laboratory Sampling: Blood samples for clinical safety
25 laboratory tests (5 mL at each sample time point) including hematology and chemistry, will be collected at Visit 1 (Screening/Baseline), Visit 2, and Visit 6. Blood Samples for population PK analysis of (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone will be drawn at Visit 2,
30 Visit 6 and Visit 7. Blood draws for the biomarker 24HC will be drawn at Visit 1 (Baseline), Visit 2 and Visit 6. Blood for pregnancy testing will be collected for all females of childbearing potential at Visit 1 (Screening/Baseline) and Visit 6 and urine pregnancy testing will be collected for all

females of childbearing potential at Visit 2 and Visit 9. In addition, urinalysis for urine for drug (illicit) screening will be performed at Visit 1 (Screening//Baseline). Cannabis use will be allowed in the study.

5 **Electrocardiograms:** Echocardiograms will be administered at Visit 1, Visit 6, and Visit 8.

[0073]

Formulation Example

For (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-
10 yl)methanone, the formulation (film coating tablet) was produced in line with the following specification (Table 1).

[0074]

Table 1

	Components	Quantity per Tablet (mg)
		20 mg
Core Tablet (Internal granules)	(4-benzyl-4-hydroxypiperidin-1-yl) (2,4'-bipyridin-3-yl)methanone	20
	Microcrystalline cellulose (PH101)	1.5
	Low-Substituted Hydroxypropyl Cellulose (L-HPC 21)	1.25
	Hydroxypropyl Cellulose	0.75
(External granules)	Low-Substituted Hydroxypropyl Cellulose (L-HPC 21)	1.25
	Magnesium Stearate	0.25
Coating Solution	(OPADRY Red 03F45081)	0.508
	(OPADRY Yellow 03F42240)	0.508
	Hypromellose 2910 ¹⁾	(0.75)
	Polyethylene Glycol 8000 ¹⁾	(0.167)
	Titanium Dioxide ¹⁾	(0.083)
	Ferric Oxide, Red ¹⁾	(0.008)
	Ferric Oxide, Yellow ¹⁾	(0.008)
	Total	26.016

[0075]

1) These ingredients are components of OPADRY® Red 03F45081 and
 5 OPADRY (registered trademark) Yellow 03F42240 (premixed coating materials).

[0076]

The following references are incorporated by reference herein to the same extent as if individually incorporated.

10 1. Johanna C. M. Schilder. et al. Pain Relief Is Associated With Improvement in Motor Function in Complex Regional Pain Syndrome Type 1: Secondary Analysis of a Placedo-Controlled

- Study on the Effects of Ketamine. *The Journal of Pain* Vol 14, No. 11 (November) 2013: 1514-1521
2. Robert J Schwartzman. et al. The use of ketamine in complex regional pain syndrome: possible mechanisms. *Expert Rev. Neurother.* 11(5), 719-734 (2011)
3. En Lin Goh. et al. Complex regional pain syndrome: a recent update. *Burns & Trauma* (2017) 5: 2
4. O'Connell NE. et al. Interventions for treating pain and disability in adults with complex regional pain syndrome - an overview of systematic reviews (Review). *Cochrane Database of Systematic Reviews* 2013, Issue 4. Art. No.: CD9416
5. David W. Dodick. et al. Effect of Fremanezumab Compared With Placebo for Prevention of Episodic Migraine. *JAMA*, 2018; 319(19): 1999-2008
- 15 6. Virginia L. Stauffer. et al. Evaluation of Galcanezumab for the Prevention of Episodic Migraine. *JAMA Neurol.* doi: 10.1001/jamaneurol.2018.1212 [0077]

This application is based on patent applications No. US 20 62/733,927 filed on September 20, 2018 in USA, the contents of which are encompassed in full herein.

CLAIMS

1. A method of treating pain in a mammal comprising administering an effective amount of a composition comprising
5 (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof to the mammal in need thereof.
2. The method of claim 1, wherein the pain is Complex Regional
10 Pain Syndrome (CRPS).
3. The method of claim 1, wherein the pain is migraine.
4. The method of claim 1, wherein administering the effective
15 amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof results in (i) a reduction in 24(S)-hydroxycholesterol (24HC) levels and/or (ii) a reduction in NMDA receptor function.
20
5. The method of claim 1, wherein the effective amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof is administered orally or parenterally.
25
6. The method of claim 1, wherein the effective amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof is administered as a single unit dose.
30
7. The method of claim 6, wherein the single unit dose is about 0.01 mg/kg to 100 mg/kg.
8. The method of claim 6, wherein the single unit dose is about

0.05 mg/kg to 100 mg/kg.

9. The method of claim 6, wherein the single unit dose is about 0.1 mg/kg to 10 mg/kg.

5

10. The method of claim 1, wherein the effective amount of the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof is administered according to a dose regimen.

10

11. The method of claim 1, wherein the mammal is a human.

12. The method of claim 1, wherein the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof
15 further comprises a pharmaceutically acceptable carrier.

13. The method of claim 1, wherein the composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof
20 further comprises an additional active agent.

14. A pharmaceutical composition comprising (4-benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a
25 pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, which is for treating pain.

15. (4-Benzyl-4-hydroxypiperidin-1-yl)(2,4'-bipyridin-3-yl)methanone or a pharmaceutically acceptable salt thereof for
30 use in treatment of pain.

INTERNATIONAL SEARCH REPORT

International application No
PCT/JP2019/038061

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K31/4545 A61K45/06 A61K9/00 A61P25/02 A61P25/06
ADD.
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
Minimum documentation searched (classification system followed by classification symbols)
A61K A61P
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, WPI Data, BIOSIS, EMBASE, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2013/090341 A1 (KOIKE TATSUKI [JP] ET AL) 11 April 2013 (2013-04-11) cited in the application	14
Y	the whole document paragraphs [0045], [0504], [0521] - [0523] table 6; compound 44 examples 3-4	1-15
Y	----- US 2016/024049 A1 (KOIKE TATSUKI [JP] ET AL) 28 January 2016 (2016-01-28) the whole document paragraphs [0694] - [0695], [0698] ----- -/--	1-15

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search 2 December 2019	Date of mailing of the international search report 11/12/2019
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Houyvet-Landriscina

INTERNATIONAL SEARCH REPORT

International application No
PCT/JP2019/038061

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2017/114042 A1 (KOIKE TATSUKI [JP] ET AL) 27 April 2017 (2017-04-27) the whole document paragraph [0010] paragraphs [0443], [0471] -----	1,3-15
Y	US 2016/022701 A1 (REDDY KIRAN [US] ET AL) 28 January 2016 (2016-01-28) the whole document paragraph [0438] -----	1-15
A	S. M. PAUL ET AL: "The Major Brain Cholesterol Metabolite 24(S)-Hydroxycholesterol Is a Potent Allosteric Modulator of N-Methyl-D-Aspartate Receptors", THE JOURNAL OF NEUROSCIENCE, vol. 33, no. 44, 30 October 2013 (2013-10-30), pages 17290-17300, XP55543316, US ISSN: 0270-6474, DOI: 10.1523/JNEUROSCI.2619-13.2013 abstract -----	1-15
A	US 9 643 957 B2 (TAKEDA PHARMACEUTICALS CO [JP]) 9 May 2017 (2017-05-09) cited in the application tables 1-7; compound 70 -----	1-15
Y	WO 2013/056229 A1 (US HEALTH [US]; COOPER HEALTH SYSTEM [US] ET AL.) 18 April 2013 (2013-04-18) the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/JP2019/038061

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2013090341	A1	11-04-2013	
		AR 088226 A1	21-05-2014
		AU 2012321815 A1	24-04-2014
		BR 112014007257 A2	28-03-2017
		CA 2851087 A1	18-04-2013
		CL 2014000806 A1	05-09-2014
		CN 103842356 A	04-06-2014
		CO 6940432 A2	09-05-2014
		CR 20140151 A	26-05-2014
		DK 2763979 T3	01-04-2019
		DO P2014000064 A	15-07-2014
		EA 201490747 A1	29-08-2014
		EC SP14013329 A	31-05-2014
		EP 2763979 A1	13-08-2014
		ES 2716531 T3	13-06-2019
		GE P201706608 B	25-01-2017
		HR P20190491 T1	03-05-2019
		HU E043171 T2	28-08-2019
		JP 6117191 B2	19-04-2017
		JP 2014528904 A	30-10-2014
		KR 20140072902 A	13-06-2014
		LT 2763979 T	25-03-2019
		MA 35623 B1	01-11-2014
		MX 360706 B	14-11-2018
		NZ 623185 A	29-01-2016
		PE 20141120 A1	15-09-2014
		PL 2763979 T3	28-06-2019
		PT 2763979 T	12-04-2019
		SG 2014010417 A	27-06-2014
		SI 2763979 T1	30-04-2019
		TN 2014000122 A1	01-07-2015
		TR 201902788 T4	21-03-2019
		TW 201329068 A	16-07-2013
		UA 115133 C2	25-09-2017
		US 2013090341 A1	11-04-2013
		US 2014088118 A1	27-03-2014
		US 2014088146 A1	27-03-2014
		US 2014228373 A1	14-08-2014
		US 2015376205 A1	31-12-2015
		US 2016326136 A1	10-11-2016
		US 2017145031 A1	25-05-2017
		US 2019071455 A1	07-03-2019
		US 2019233436 A1	01-08-2019
		UY 34365 A	31-05-2013
		WO 2013054822 A1	18-04-2013
		ZA 201402424 B	29-07-2015
US 2016024049	A1	28-01-2016	
		EP 2982666 A1	10-02-2016
		JP 6272832 B2	31-01-2018
		JP WO2014163161 A1	16-02-2017
		US 2016024049 A1	28-01-2016
		WO 2014163161 A1	09-10-2014
US 2017114042	A1	27-04-2017	
		CA 2951521 A1	17-12-2015
		CN 106573916 A	19-04-2017
		EP 3152198 A1	12-04-2017
		JP 6533799 B2	19-06-2019
		JP 2017520540 A	27-07-2017
		US 2017114042 A1	27-04-2017

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/JP2019/038061

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		WO 2015190613 A1	17-12-2015
US 2016022701	A1 28-01-2016	AU 2014243819 A1	01-10-2015
		AU 2018278924 A1	03-01-2019
		BR 112015022934 A2	18-07-2017
		CA 2905346 A1	02-10-2014
		CN 105164161 A	16-12-2015
		EP 2970504 A1	20-01-2016
		HK 1220472 A1	05-05-2017
		IL 241576 A	31-10-2019
		JP 2016514967 A	26-05-2016
		RU 2015143463 A	19-04-2017
		US 2016022701 A1	28-01-2016
		WO 2014160441 A1	02-10-2014
		ZA 201506807 B	25-01-2017
US 9643957	B2 09-05-2017	AU 2013358112 A1	30-07-2015
		CA 2894492 A1	19-06-2014
		CL 2015001599 A1	09-10-2015
		CN 104981457 A	14-10-2015
		CR 20150364 A	20-09-2016
		DO P2015000140 A	15-07-2015
		EA 201591122 A1	30-11-2015
		EP 2933247 A1	21-10-2015
		ES 2695350 T3	03-01-2019
		GE P201706756 B	25-10-2017
		HK 1210769 A1	06-05-2016
		HU E039922 T2	28-02-2019
		IL 239278 A	31-10-2018
		JP 6224004 B2	01-11-2017
		JP WO2014092100 A1	12-01-2017
		KR 20150093240 A	17-08-2015
		PE 20151162 A1	28-08-2015
		PH 12015501316 A1	24-08-2015
		SG 11201504516P A	30-07-2015
		TN 2015000245 A1	03-10-2016
		TR 201814885 T4	22-10-2018
		UA 115156 C2	25-09-2017
		US 2015315209 A1	05-11-2015
		US 2017197983 A1	13-07-2017
		WO 2014092100 A1	19-06-2014
		ZA 201504980 B	28-09-2016
WO 2013056229	A1 18-04-2013	AU 2012323845 A1	24-04-2014
		CA 2851619 A1	18-04-2013
		CN 104395283 A	04-03-2015
		CN 109793725 A	24-05-2019
		EP 2766339 A1	20-08-2014
		JP 6296985 B2	20-03-2018
		JP 2015501302 A	15-01-2015
		US 2014296241 A1	02-10-2014
		US 2017049780 A1	23-02-2017
		US 2018098993 A1	12-04-2018
		WO 2013056229 A1	18-04-2013