




(86) Date de dépôt PCT/PCT Filing Date: 1999/11/03
 (87) Date publication PCT/PCT Publication Date: 2000/05/18
 (45) Date de délivrance/Issue Date: 2009/02/10
 (85) Entrée phase nationale/National Entry: 2001/04/11
 (86) N° demande PCT/PCT Application No.: SE 1999/001989
 (87) N° publication PCT/PCT Publication No.: 2000/027366
 (30) Priorité/Priority: 1998/11/05 (SE9803772-4)

(51) Cl.Int./Int.Cl. *A61K 31/4439* (2006.01),
A61K 47/38 (2006.01), *A61K 9/24* (2006.01),
A61P 1/00 (2006.01)
 (72) Inventeurs/Inventors:
 BERGSTRAND, PONTUS, SE;
 WANG, PETER, SE
 (73) Propriétaire/Owner:
 ASTRAZENECA AB, SE
 (74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : FORMULATION PHARMACEUTIQUE CONTENANT DE L'OMEPRAZOLE
 (54) Title: PHARMACEUTICAL FORMULATION COMPRISING OMEPRAZOLE



(57) Abrégé/Abstract:

An enteric coated oral pharmaceutical formulation comprising as active ingredient a compound selected from the group of omeprazole, an alkaline salt of omeprazole, one of the single enantiomers of omeprazole and an alkaline salt of one of the single

(57) **Abrégé(suite)/Abstract(continued):**

enantiomers of omeprazole, wherein the formulation comprises a core material that comprises the active ingredient and optionally an alkaline reacting compound, the active ingredient is in admixture with a pharmaceutically acceptable excipient, such as for instance a binding agent, and on said core material a separating layer and an enteric coating layer. A hydroxypropyl cellulose (HPC) with a specific cloud point is used in the manufacture of the claimed pharmaceutical formulations. Furthermore, the application describes the processes for their preparation and the use of the claimed formulations in medicine.




PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 9/24, 31/44, 47/38	A1	(11) International Publication Number: WO 00/27366 (43) International Publication Date: 18 May 2000 (18.05.00)
<p>(21) International Application Number: PCT/SE99/01989</p> <p>(22) International Filing Date: 3 November 1999 (03.11.99)</p> <p>(30) Priority Data: 9803772-4 5 November 1998 (05.11.98) SE</p> <p>(71) Applicant (for all designated States except US): AS-TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): BERGSTRAND, Pontus [SE/SE]; Astra Hässle AB, S-431 83 Mölndal (SE). WANG, Peter [SE/SE]; Astra Hässle AB, S-431 83 Mölndal (SE).</p> <p>(74) Agent: ASTRAZENECA AB; Intellectual Property, Patents, S-151 85 Södertälje (SE).</p>	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: PHARMACEUTICAL FORMULATION COMPRISING OMEPRAZOLE



(57) Abstract

An enteric coated oral pharmaceutical formulation comprising as active ingredient a compound selected from the group of omeprazole, an alkaline salt of omeprazole, one of the single enantiomers of omeprazole and an alkaline salt of one of the single enantiomers of omeprazole, wherein the formulation comprises a core material that comprises the active ingredient and optionally an alkaline reacting compound, the active ingredient is in admixture with a pharmaceutically acceptable excipient, such as for instance a binding agent, and on said core material a separating layer and an enteric coating layer. A hydroxypropyl cellulose (HPC) with a specific cloud point is used in the manufacture of the claimed pharmaceutical formulations. Furthermore, the application describes the processes for their preparation and the use of the claimed formulations in medicine.

PHARMACEUTICAL FORMULATION COMPRISING OMEPRAZOLE

Field of the invention.

The present invention relates to an oral pharmaceutical formulation comprising the acid labile H^+ , K^+ -ATPase inhibitor omeprazole, an alkaline salt of omeprazole, one of the single enantiomers thereof or an alkaline salt of one of the single enantiomers of omeprazole. In the following these compounds are referred to as omeprazole. The formulation is in the form of a multiple unit dosage form that comprises enteric coating layered units of omeprazole. More specifically, the units comprise a core material that comprises omeprazole optionally in admixture with an alkaline reacting substance, and in admixture with one or more pharmaceutically acceptable excipients such as a binding agent, a filling agent and/or a disintegrating agent. Furthermore, each unit comprises a separating layer to separate the enteric coating layer from the core material. The separating layer comprises a specific quality of hydroxypropyl cellulose (HPC), and optionally pharmaceutical excipients. More specifically, the HPC quality is defined by having a specific cloud point.

Furthermore, the present invention refers to the use of the specific quality of HPC in the manufacture of a pharmaceutical formulation comprising omeprazole, and the use of such a pharmaceutical formulation in medicine.

Background of the invention.

Omeprazole, an alkaline salt thereof, the single enantiomers of omeprazole and an alkaline salt of the single enantiomers of omeprazole, all compounds hereinafter referred to as omeprazole, are used in the treatment of gastric acid related diseases. Omeprazole and pharmaceutically acceptable salts thereof are described in EP 5129, and some specific alkaline salts of omeprazole are described in EP 124 495 and WO95/01977. Certain salts of the single enantiomers of omeprazole and their preparations are described in WO94/27988.

Omeprazole is generally known to be useful for inhibiting gastric acid secretion in mammals and man by controlling gastric acid secretion at the final step of the acid secretory pathway. Thus, in a more general sense, it may be used for prevention and treatment of gastric-acid related diseases in mammals and man, including e.g. reflux oesophagitis, gastritis, duodenitis, gastric ulcers and duodenal ulcers. Furthermore, it may be used for treatment of other gastrointestinal disorders where gastric acid inhibitory effect is desirable e.g. in patients on NSAID therapy, in patients with non ulcer dyspepsia, in patients with symptomatic gastro-oesophageal reflux disease, and in patients with gastrinomas. It may also be used in a patient in intensive care situations, in a patient with acute upper gastrointestinal bleeding, pre-and post-operatively to prevent aspiration of gastric acid and to prevent and treat stress ulceration. Further, it may be useful in the treatment of psoriasis as well as in the treatment of *Helicobacter* infections and diseases related to these, as well as in the treatment or prophylaxis of inflammatory conditions in mammals, including man.

15

Omeprazole is, however, susceptible to degradation or transformation in acidic and neutral media. The degradation is catalyzed by acidic compounds and is stabilized in mixtures with alkaline compounds. The chemical stability of omeprazole is also affected by moisture, heat, and organic solvents and to some degree by light.

20

Due to the chemical stability properties of omeprazole, it is obvious that an oral solid dosage form comprising omeprazole must be protected from contact with the acidic gastric juice. Omeprazole must also be transferred in intact form to that part of the gastrointestinal tract where pH is near neutral and where rapid absorption can occur.

25

A pharmaceutical oral dosage form of omeprazole is best protected from contact with acidic gastric juice by an enteric coating layer. For instance, EP 247 983 describes enteric coated formulations of omeprazole. Such as formulation contains omeprazole in the form of a core unit containing omeprazole together with an alkaline salt or containing an alkaline salt of omeprazole optionally together with an alkaline salt, the core unit is layered with a

30

separating layer and an enteric coating layer. In WO 96/01623 a multiple unit tableted dosage formulation comprising omeprazole is described.

The oral formulations described in EP 247 983 and the tablet formulations described in
5 WO 96/01623 are examples of enteric coating layered formulations that comprise or optionally comprise a separating layer to separate the acidic enteric coating material from omeprazole being an acid susceptible substance. HPC may be used in a layer that separates the core material from the enteric coating layer in the described formulations. All ingredients, including HPC qualities, used in a pharmaceutical preparation must fulfil strict
10 criteria, such as for instance requirements defined in pharmacopoeial monographs.

The rate of release of omeprazole from a pharmaceutical dosage form can influence the total extent of absorption of omeprazole into the general circulation (Pilbrant and Cederberg, Scand. J. Gastroenterology 1985; 20 (suppl. 108) p. 113-120). Therefore the
15 limits for rate of release of the omeprazole from the pharmaceutical formulation are stated in the marketing approval for the products. The release of omeprazole is affected both by the chemical stability of the active substance and the release stability of the pharmaceutical formulation. If the formulation is unstable with respect to the release rate, the drug will have a non-accepted storage time, i.e. the expiration period for the product will be too
20 short.

It has now surprisingly been found that different batches of HPC, which fulfil all pharmacopoeial requirements, used as material for the separating layer in a pharmaceutical formulation comprising omeprazole, may result in different release rate over time. Thus,
25 the storage period for the pharmaceutical formulation may not be acceptable. One parameter of interest for the HPC's influence on the release stability is its water solubility.

The aqueous solubility of HPC decreases with increasing temperature due to polymer phase separation. This is observed as a clouding of the polymer solution when the
30 temperature is increased. Cloud point is the temperature at which this polymer phase

23940-1250

4

separation occurs. Cloud point is determined by measuring the light transmission through the polymer solution. The light transmission of a specific system where the polymer is dissolved, that is a transparent polymer solution without clouding, is defined as light transmission 100 %. In this patent application cloud point is defined as the temperature
5 where the light transmission of a specific system is 96% when a commercial instrument from Mettler is used. For other cloud point systems and instruments another light transmisson may be specified for each system.

One problem that can be avoided by the new formulation and use of a specific quality of
10 HPC, is that the storage period for the dosage form can be extended and guaranteed. From an economical aspect it is advantageous to specify and check the HPC quality thereby keeping a long expire date of the dosage form.

Outline of the invention.

15 It has now been found that a quality of HPC with a cloud point of not less than 38°C determined as the temperature where the light transmission of a specified system is 96% measured by a Mettler FP90/FP 81C instrument is desirable in an enteric coating layered pharmaceutical formulation comprising omeprazole. Preferably, the HPC should have a cloud point of not less than 40°C, and more preferably not less than 41°C. When another
20 instrument is used for determination, the cloud point may be specified in other terms. An upper limit for the cloud point is not critical and therefore there is no need to specify that.

The HPC is used as a constituent of a separating layer separating the core material comprising omeprazole from the enteric coating layer. The HPC quality defined in the
25 present patent application is desirable in fulfilling the criteria on release rate stability and to be suitable for oral administration forms comprising omeprazole.

23940-1250

4a

In one formulation aspect, the invention provides an enteric coated oral pharmaceutical formulation comprising a core material that comprises, as active ingredient, a compound selected from the group consisting of omeprazole, an alkaline salt of omeprazole, one of the single enantiomers of omeprazole and an alkaline salt of one of the single enantiomers of omeprazole, optionally in admixture with an alkaline reacting compound, wherein the active ingredient is in admixture with one or more pharmaceutically acceptable excipients, wherein on said core material there is a separating layer and an enteric coating layer, wherein a hydroxypropyl cellulose (HPC) with a cloud point of at least 38°C determined as the temperature where the light transmission of the system is 96%, is used as a constituent of the separating layer, and wherein the cloud point is determined in the following way: the HPC is dissolved in a concentration of 1.0% (w/w) in a mixed solution of disodium hydrogenphosphate buffer 0.086 M and hydrochloric acid 0.1 M in the proportions 7:3 at a pH of 6.75-6.85.

In a use aspect, the invention provides use of the HPC as defined above in the manufacture of the formulation defined above.

In a process aspect, the invention provides a process for the manufacture of the formulation as defined above, wherein the active ingredient is mixed with the binding agent and, optionally mixed with the alkaline reacting compound, is layered on a seed and formulated into the core material and the separating layer is coating layered on said core material, and thereafter the enteric coating layer is applied, wherein the separating layer comprises HPC as defined above.

23940-1250

4b

In a further use aspect, the invention provides use of the formulation as defined above, for the manufacture of a medicament for the treatment of a gastrointestinal disease.

5 In a still further use aspect, the invention provides use of the formulation as defined above, for the treatment of a gastrointestinal disease

In a commercial package aspect, the invention provides a commercial package comprising the formulation as
10 defined above, and associated therewith instructions for the use thereof in the treatment of a gastrointestinal disease.

Detailed description of the drawings.

Figure 1 shows two graphs representing two different dosage forms based on two qualities of HPC named Type A and Type B. The graphs show released omeprazole from the dosage forms after 3 months and 6 months storage at accelerated conditions at 40°C and 75% relative humidity. The two HPC qualities are used as a constituent of the separating layer described in Example 2 below. With a separating layer comprising HPC Type A the release rate of omeprazole over time has decreased. With the HPC Type B the release rate of omeprazole over time is almost the same as for a freshly produced product.

Figure 2 shows two graphs representing two different qualities of HPC named Type A and Type B. The graphs show cloud point determinations for the two HPC qualities used as a constituent of the separating layer described in Examples 1 - 3 below.

Figure 3a) and Figure 3b) show graphs representing two different dosage forms based on two qualities of HPC named Type A and Type B. Figure 3a) shows released omeprazole from dosage forms comprising HPC type A, i.e. a reference. Figure 3b) shows released omeprazole from dosage forms comprising HPC type B, i.e. according to the invention. The two HPC qualities are used as a constituent of the separating layer described in Example 1 below.


Detailed description of the invention.

Core materials.

Omeprazole with formula Ia, is preferably formulated into an oral composition in the form of a pharmaceutically acceptable salt, such as an alkaline salt selected from the group of the Mg²⁺, Ca²⁺, Na⁺ and K⁺ salts, more preferably the Mg²⁺ salt. Omeprazole may also be used in the form of one of the single enantiomers of omeprazole or an alkaline salt of one of the single enantiomers of omeprazole, especially an alkaline salt of the (-)-enantiomer of omeprazole, and more preferably the Mg²⁺ salt of the (-)-enantiomer of omeprazole.

23940-1250

6



The core material for the individually enteric coating layered pellets can be composed and formulated according to different principles, such as described in EP 247 983 and WO 96/01623. For instance, omeprazole is mixed with one or more pharmaceutical constituents to obtain preferred handling and processing properties and also to obtain a suitable concentration of omeprazole in the final mixture. Pharmaceutical constituents such as fillers, binders, lubricants, disintegrating agents, surfactants and other pharmaceutically acceptable additives, can be used.

Preferably, omeprazole, optionally after mixing with an alkaline compound, is mixed with suitable constituents including a binding agent and formulated into a core material. Said core materials may be produced by extrusion/spheronization, balling or compression and by utilizing different process equipment. The formulated core materials may have a size of less than approximately 2 mm. The manufactured core materials can be layered further with additional ingredients, optionally comprising active substance, and/or be used for further processing.

Alternatively, inert seeds layered with active substance (the active substance is optionally mixed with alkaline compounds) can be used as the core material for the further processing. The seeds, which are to be layered with the active substance, can be water insoluble seeds comprising different oxides, celluloses, organic polymers and other materials, alone or in mixtures or water soluble seeds comprising different inorganic salts, sugars, non-pareils and other materials, alone or in mixtures.

Before the seeds are layered, for instance by using granulating or spray coating/layering equipment, omeprazole is mixed with a binding agent and optionally further components. Such further components can be binders, surfactants, fillers, disintegrating agents, alkaline additives or other pharmaceutically acceptable ingredients, alone or in mixtures.

5

The binders are for example celluloses such as hydroxypropyl methylcellulose, hydroxypropyl cellulose, microcrystalline cellulose and carboxymethyl-cellulose sodium, polyvinyl pyrrolidone, sugars, starches and other pharmaceutically acceptable substances with cohesive properties. Suitable surfactants are found in the groups of pharmaceutically acceptable non-ionic or ionic surfactants, such as for instance sodium lauryl sulphate.

10

The active substance may also be mixed with an alkaline pharmaceutically acceptable substance (or substances). Such substances can be chosen among, but are not restricted to, substances such as the sodium, potassium, calcium, magnesium and aluminium salts of phosphoric acid, carbonic acid, citric acid or other suitable weak inorganic or organic acids; aluminium hydroxide/sodium bicarbonate co-precipitate; substances normally used in antacid preparations such as aluminium, calcium and magnesium hydroxides; magnesium oxide or composite substances, such as $\text{Al}_2\text{O}_3 \cdot 6\text{MgO} \cdot \text{CO}_2 \cdot 12\text{H}_2\text{O}$, $\text{Mg}_6\text{Al}_2(\text{OH})_{16}\text{CO}_3 \cdot 4\text{H}_2\text{O}$, $\text{MgO} \cdot \text{Al}_2\text{O}_3 \cdot 2\text{SiO}_2 \cdot n\text{H}_2\text{O}$ or similar compounds; organic pH-buffering substances such as trihydroxy methyl amino methane, basic amino acids and their salts or other similar, pharmaceutically acceptable pH-buffering substances.

15

20

Alternatively, the aforementioned core material can be prepared by using spray drying or spray congealing technique.

25

Separating layer(s)

The core material containing omeprazole must, according to EP 247 983, be separated from the enteric coating polymer(s) containing free carboxyl groups, which may otherwise cause degradation/dicolouration of omeprazole during the coating process or during storage.

30

According to the present invention, the separating layer comprises a specific quality of HPC. This specific quality of HPC should preferably have a cloud point of at least 38°C determined by a Mettler instrument. The cloud point is determined in a mixed disodium hydrogenphosphate buffer 0.086 M and hydrochloric acid 0.1 M in the proportions 7:3. The mixed solution used for the cloud point determination has a pH of 6.75 - 6.85. The concentration of HPC in the mixed solution is 1.0% (w/w) for the Mettler instrument. For more detailed information on the composition of the mixed solution, see below in the experimental section. Preferably, the HPC has a low viscosity, such as for instance below 400 mPas in a 5% (w/w) water solution at 25°C.

10

Alternatively, the quality of HPC may be determined by a method that correlates with the method described above, e.g. NIR spectrophotometry.

Additives such as plasticizers, colorants, pigments, fillers, anti-tacking, buffering agents, and anti-static agents, such as for instance magnesium stearate, titanium dioxide, talc, and other additives may also be included in the separating layer(s).

Enteric coating layer(s)

One or more enteric coating layers are applied onto the core material covered with separating layer(s) by using a suitable coating technique. The enteric coating layer material may be dispersed or dissolved in either water or in a suitable organic solvent. As enteric coating layer polymers one or more, separately or in combination, of the following polymers can be used; e.g. solutions or dispersions of methacrylic acid copolymers, cellulose acetate phthalate, cellulose acetate butyrate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, polyvinyl acetate phthalate, cellulose acetate trimellitate, carboxymethylethylcellulose, shellac or other suitable enteric coating layer polymer(s). For environmental reasons, an aqueous coating process may be preferred. In such aqueous processes methacrylic acid copolymers are most preferred.

The enteric coating layers may contain pharmaceutically acceptable plasticizers to obtain desirable mechanical properties, such as flexibility and hardness of the enteric coating layers. Such plasticizers are for instance, but not restricted to, triacetin, citric acid esters, phthalic acid esters, dibutyl sebacate, cetyl alcohol, polyethylene glycols, polysorbates or other plasticizers. The amount of plasticizer is optimized for each enteric coating layer formula, in relation to selected enteric coating layer polymer(s), selected plasticizer(s) and the applied amount of said polymer(s). Additives such as dispersants, colorants, pigments, polymers e.g. poly(ethylacrylate, methylmethacrylate), anti-tacking and anti-foaming agents may also be included in the enteric coating layer(s). Other compounds may be added to increase film thickness and to decrease diffusion of acidic gastric juices into the acidic susceptible active substance.

To protect the acidic susceptible active substance, the enteric coating layer(s) preferably constitute(s) a thickness of at least approximately 10 μm . The maximum thickness of the applied enteric coating layer(s) is normally only limited by processing conditions.

The pellets or units covered with enteric coating layer(s) may further be covered with one or more over-coating layer(s). The over-coating layer(s) can be applied to the enteric coating layered pellets by coating or layering procedures in suitable equipment such as coating pan, coating granulator or in a fluidized bed apparatus using water and/or organic solvents for the layering process.

Final dosage form.

The prepared pellets may be filled in hard gelatine capsules or compressed with suitable tablet excipients into a tableted multiple unit formulation, and the latter is preferred. Final dosage forms may also include but is not restricted to effervescent tablets, and combinations of omeprazole with other active ingredients, such as for instance antibacterial substances, NSAID(s), motility stimulating agents or antacids.

Experimental section.

Example 1: Test of omeprazole multiple unit tablets, in which the pellets are layered with different types of HPC used as a constituent of the separation layer (laboratory scale).

5 Omeprazole tablets with the following composition were prepared according to the description in WO 96/01623. Sugar spheres were spray layered in a fluidized bed with an aqueous suspension of omeprazole magnesium salt and HPMC. The prepared pellets were layered with a separating layer and thereafter enteric coated. Enteric coated pellets were mixed with tablets excipients and compressed into a multiple unit tablet.

10

The composition of the tested omeprazole tablets (20 mg strength) was as follows.

	NAME OF INGREDIENT	FORMULA (mg/tablet)
	Omeprazole magnesium	20.6
15	Glyceryl monostearate	1.4
	Hydroxypropylcellulose	4.8
	Hydroxypropyl methylcellulose	4.6
	Magnesium stearate	0.7
	Methacrylic acid copolymer type C	27
20	Microcrystalline cellulose	220
	Polysorbate 80	0.1
	Polyvinylpyrrolidone crosslinked	4.6
	Sodium stearyl fumarate	0.5
	Sugar spheres	22
25	Talc	8.3
	Triethyl citrate	8.2

Omeprazole multiple unit tablets prepared with a separating layer on the pellets which separating layer comprises HPC, of either quality i.e type A or type B. HPC of the two

types fulfill all requirements in the PhEur as well as the USP. However, the HPC of the two types differ with respect to some physical/chemical characteristics, e.g. cloud point.

The prepared tablets were tested according to the description below. The tablets, i.e. the pellets, were prepared from the same batch of omeprazole magnesium, and with the same enteric coating material. The release of omeprazole was tested on stored tablets after 0 month, and 6 months storage. The amount of released omeprazole within 30 minutes in a buffer solution was determined.

The tablets were pre-exposed to hydrochloric acid at 37°C for 2 hours. Thereafter the drug release in buffer solution pH 6.8 at 30 minutes was determined by liquid chromatography. The buffer solution pH 6.8 was a mixture of disodium hydrogenphosphate buffer 0.086 M and hydrochloric acid 0.1 M in the proportions 7:3, pH should be between 6.75 and 6.85. The hydrochloric acid 0.1 M was prepared by dissolving 213 ml of conc. HCl in water and added with water to 25 000 ml. The disodium hydrogen phosphate solution 0.086 M was prepared by dissolving 382 g Na₂HPO₄·2H₂O in water and dilute to 25 000 ml with water.

The stability testing was performed on (20 mg strength) tablets packed in plastic bottles with desiccant (the tablets were not covered by a tablet coat).

Results are shown in Figure 3a) and Figure 3b). Figure 3a) shows results with the HPC quality type A, i.e. a reference, and Figure 3b) shows results with HPC type B, i.e. according to the instant invention.

Example 2. Release of omeprazole from tablets comprising different types of HPC as a constituent of the separating layer.

Pilot scale batches (using HPC of type A: 6 batches, and type B: 2 batches) were manufactured in order to confirm the improvement found during the laboratory testing in

Example 1. Results from stability studies are shown in Figure 1.

The comparison clearly indicates improved release rate stability for tablets containing HPC of type B relative to that of type A.

General compositions for omeprazole tablets (20-mg strength):

5	NAME OF INGREDIENT	FORMULA (mg/tablet)
	Omeprazole magnesium	20.6
	Colour iron oxide reddish-brown	0.3
	Glyceryl monostearate	1.4
	Hydroxypropylcellulose	4.8
10	Hydroxypropyl methylcellulose	15
	Magnesium stearate	0.7
	Methacrylic acid copolymer type C	27
	Microcrystalline cellulose	220
	Paraffin	0.2
15	Polyethylene glycol 6000	2.5
	Polysorbate 80	0.1
	Polyvinylpyrrolidone crosslinked	4.6
	Sodium stearyl fumarate	0.5
	Sugar spheres	22
20	Talc	8.3
	Titanium dioxide	2.2
	Triethyl citrate	8.2

The tablets were manufactured as described in example 1, with the additional step of a
25 tablet coat comprising HPMC, PEG 6 000, and pigment.

Example 3. Cloud point determinations.

Omeprazole tablets were manufactured in laboratory scale as described in example 1.

Cloud point determinations of the HPC types in the Mettler instrument was conducted in the following way. The cloud point of the HPC types was determined in a mixed phosphate buffer 0.086 M and hydrochloric acid 0.1 M in the proportions 7:3. The mixed solution used for the cloud point determination had a pH of 6.75 - 6.85. The concentration of HPC in the mixed solution was 1.0% (w/w). It is essential for the specificity of the cloud point determination that this system is used in the chosen instrument. The Mettler instrument comprises the following parts: Mettler FP90 Central processor, FP81C Measuring unit and ME-18572 boiling point tubes. A temperature range of 30.0 to 50.0°C was used and a heating rate of 1.0°C/min. The cloud point is defined as the temperature where the light transmission is 96%.

The results are shown in Figure 2.

23940-1250

14

CLAIMS:

1. An enteric coated oral pharmaceutical formulation comprising a core material that comprises, as active ingredient, a compound selected from the group consisting of omeprazole, an alkaline salt of omeprazole, one of the single enantiomers of omeprazole and an alkaline salt of one of the single enantiomers of omeprazole, optionally in admixture with an alkaline reacting compound, wherein the active ingredient is in admixture with one or more pharmaceutically acceptable excipients, wherein on said core material there is a separating layer and an enteric coating layer, wherein a hydroxypropyl cellulose (HPC) with a cloud point of at least 38°C determined as the temperature where the light transmission of the system is 96%, is used as a constituent of the separating layer, and wherein the cloud point is determined in the following way: the HPC is dissolved in a concentration of 1.0% (w/w) in a mixed solution of disodium hydrogenphosphate buffer 0.086 M and hydrochloric acid 0.1 M in the proportions 7:3 at a pH of 6.75-6.85.
2. A formulation according to claim 1, wherein the one or more excipients are a binding agent, a filling agent, a disintegrating agent or any mixture thereof.
3. A formulation according to claim 1 or 2, wherein the HPC has a cloud point of at least 40°C.
4. A formulation according to claim 3, wherein the HPC has a cloud point of at least 41°C.
5. A formulation according to any one of claims 1 to 4, wherein the enteric coating layer comprises a methacrylic acid copolymer.

23940-1250

15

6. A formulation according to any one of claims 1 to 5, wherein the HPC has a low viscosity.

7. A formulation according to any one of claims 1 to 6, wherein the active ingredient is omeprazole.

5 8. A formulation according to any one of claims 1 to 6, wherein the active ingredient is a magnesium salt of omeprazole.

9. A formulation according to any one of claims 1 to 6, wherein the active ingredient is a magnesium salt of
10 the (-)-enantiomer of omeprazole.

10. Use of the HPC as defined in any one of claims 1, 3, 4 and 6, in the manufacture of the formulation as defined in any one of claims 1, 2, 5 and 7 to 9.

11. A process for the manufacture of the formulation
15 as defined in any one of claims 2, 5 and 7 to 9, wherein the active ingredient is mixed with the binding agent and, optionally mixed with the alkaline reacting compound, is layered on a seed and formulated into the core material and the separating layer is coating layered on said core
20 material, and thereafter the enteric coating layer is applied, wherein the separating layer comprises HPC as defined in any one of claims 1, 3, 4 and 6.

12. Use of the formulation as defined in any one of claims 1 to 9, for the manufacture of a medicament for the
25 treatment of a gastrointestinal disease.

13. Use of the formulation as defined in any one of claims 1 to 9, for the treatment of a gastrointestinal disease.

23940-1250


16


14. The formulation as defined in any one of claims 1 to 9, for use in the manufacture of a medicament for the treatment of a gastrointestinal disease.

15. The formulation as defined in any one of claims 1 to 9, for use in the treatment of a gastrointestinal disease.

16. A commercial package comprising the formulation as defined in any one of claims 1 to 9, and associated therewith instructions for the use thereof in the treatment of a gastrointestinal disease.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA
PATENT AGENTS








Fig. 3b

