



Europäisches  
Patentamt  
European  
Patent Office  
Office européen  
des brevets



(11) EP 2 724 727 A1

(12)

## EUROPEAN PATENT APPLICATION

(43) Date of publication:  
30.04.2014 Bulletin 2014/18

(51) Int Cl.:  
A61K 39/395 (2006.01)

(21) Application number: 13195495.0

(22) Date of filing: 27.06.2008

(84) Designated Contracting States:  
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT  
RO SE SI SK TR  
Designated Extension States:  
AL BA MK RS

- Vournakis, John, N.  
Charleston, SC 29403 (US)
- Felber, Barbara, K.  
Rockville, MD 20850 (US)
- Finkielstein, Sergio  
Newton, MA 02459 (US)

(30) Priority: 27.06.2007 US 937471 P

(74) Representative: Jones Day  
Rechtsanwälte, Attorneys-at-Law  
Patentanwälte  
Intellectual Property  
Prinzregentenstraße 11  
80538 München (DE)

(71) Applicants:  
• The United States of America, as Represented by  
The Secretary, Department of Health and Human  
Services  
Bethesda, Maryland 20892-7660 (US)  
• Admune Therapeutics LLC  
Danvers, MA 01923 (US)

Remarks:

- This application was filed on 03-12-2013 as a divisional application to the application mentioned under INID code 62.
- Claims filed after the date of receipt of the divisional application (Rule 68(4) EPC.).

(72) Inventors:  
• Pavlakis, George, N.  
Rockville, MD 20850 (US)

### (54) Complexes of IL-15 and IL-15R alpha and uses thereof

(57) The invention provides a complex for use in a method of treating cancer in a subject, wherein  
(A) the complex comprises: (i) interleukin 15 (IL-15); and  
(ii) a soluble form of interleukin 15 receptor alpha (IL-15Ra), wherein the complex is administered with an antibody that specifically binds to a T cell receptor or a tumor-associated antigen;  
(B) the complex comprises: (i) a fusion protein comprising

IL-15 and a soluble form of IL-15Ra; and (ii) an antibody that specifically binds to a cancer antigen; or  
(C) the complex comprises: (i) IL-15; (ii) a soluble form of IL-15Ra, or a fusion protein comprising a soluble form of IL-15Ra and the Fc domain of an immunoglobulin or a fragment thereof; and (iii) an antibody that specifically binds to a cancer antigen.

**Description****RELATED APPLICATIONS**

5 [0001] This application claims the benefit of U.S. Provisional Application No. 60/937,471 filed June 27, 2007, which is incorporated by reference herein in its entirety.

**GOVERNMENTAL INTERESTS**

10 [0002] This invention was created in the performance of a Cooperative Research and Development Agreement with the National Institutes of Health, an Agency of the Department of Health and Human Services. The Government of the United States has certain rights in this invention.

**1. FIELD OF THE INVENTION**

15 [0003] The present invention relates to Therapeutic Agents that modulate IL-15-mediated function for the prevention, treatment and/or management of diseases involving IL-15-mediated signaling, including, but not limited to cancer, infectious disease, autoimmune disease, and transplantation rejection.

**2. BACKGROUND OF THE INVENTION**

20 [0004] The cytokine, interleukin-15 (IL-15), is a member of the four alpha-helix bundle family of lymphokines produced by many cells in the body. IL-15 plays a pivotal role in modulating the activity of both the innate and adaptive immune system, e.g., maintenance of the memory T-cell response to invading pathogens, inhibition of apoptosis, activation of dendritic cells, and induction of Natural Killer (NK) cell proliferation and cytotoxic activity.

25 [0005] The IL-15 receptor consists of three polypeptides, the type-specific IL-15 receptor alpha ("IL-15Ra"), the IL-2/IL-15 receptor beta (or CD122) (" $\beta$ "), and the common gamma chain (or CD132) (" $\gamma$ ") that is shared by multiple cytokine receptors. The IL-15Ra is thought to be expressed by a wide variety of cell types, but not necessarily in conjunction with  $\beta$  and  $\gamma$ . IL-15 signaling has been shown to occur through the heterodimeric complex of IL-15Ra,  $\beta$ , and  $\gamma$ ; through the heterodimeric complex of  $\beta$  and  $\gamma$ , or through a subunit, IL-15RX, found on mast cells.

30 [0006] IL-15 is a soluble protein, but endogenous IL-15 is not readily detectable in serum or body fluids - instead, it occurs predominantly as a membrane-bound form that is expressed or acquired by several types of accessory cells. For instance, although IL-15 mRNA is detected in cells of both hematopoietic and non-hematopoietic lineage, T cells do not produce IL-15. Instead, IL-15 binds to the IL-15Ra, forming cell-surface complexes on T cells. IL-15 specifically binds to the IL-15Ra with high affinity via the "sushi domain" in exon 2 of the extracellular domain of the receptor. After trans-endosomal recycling and migration back to the cell surface, these IL-15 complexes acquire the property to activate bystander cells expressing the IL-15R  $\beta\gamma$  low-affinity receptor complex, inducing IL-15-mediated signaling via the Jak/Stat pathway. A naturally occurring soluble form of IL-15Ra ("sIL-15Ra"), which is cleaved at a cleavage site in the extracellular domain immediately distal to the transmembrane domain of the receptor has been observed. Tumor necrosis factor-alpha-converting enzyme (TACE/ADAM17) has been implicated as a protease involved in this process.

35 [0007] Based on its multifaceted role in the immune system, various therapies designed to modulate IL-15-mediated function have been explored. For example, the administration of exogenous IL-15 can enhance the immune function of patients infected with human immunodeficiency virus (HIV). In keeping with its immune enhancing activity, increased expression of endogenous IL-15 is observed in patients with autoimmune diseases, e.g., rheumatoid arthritis, multiple sclerosis, ulcerative colitis, and psoriasis. Because some studies reported that the soluble form of the IL-15Ra (sIL-15Ra) is an antagonist of IL-15-mediated signaling, the sIL-15Ra has been explored for treating autoimmune inflammatory diseases. Nevertheless, recent reports suggest that IL-15, when complexed with the sIL-15Ra, or the sushi domain, maintains its immune enhancing function.

40 [0008] Despite the amount of progress made in understanding the function of IL-15, it is unclear how various forms of the IL-15Ra, alone or when complexed to IL-15, can be used to modulate IL-15 function as part of a therapeutic regimen.

**3. SUMMARY OF THE INVENTION**

45 [0009] The present invention relates to agents that modulate interleukin-15 ("IL-15") signal transduction or function ("Therapeutic Agents") and the use of those agents to modulate immune function. The Therapeutic Agents target the interaction between IL-15 and its receptor and modulate IL-15-induced signal transduction. The Therapeutic Agents may be formulated with polymers, such as poly- $\beta$ -1→4-N- acetylglucosamine, for administration to a human subject to modulate IL-15-mediated immune function.

[0010] The present invention provides Therapeutic Agents that induce IL-15 signal transduction and enhance IL-15-mediated immune function (*i.e.*, Agonistic Therapeutic Agents). The Agonistic Therapeutic Agents are useful for enhancing IL-15-mediated immune function in a subject in need of such therapy. In particular, the Agonistic Therapeutic Agents are useful for the prevention, treatment and/or management of disorders in which it is beneficial to enhance IL-

5 15-mediated immune function. Non-limiting examples of such disorders include cancer and infectious diseases. In a specific embodiment, the invention provides a method for treating a cancer or an infectious disease in a human subject, comprising administering to a human subject in need thereof an effective amount of an Agonistic Therapeutic Agent.

[0011] The Agonistic Therapeutic Agents include complexes that bind to the  $\beta\gamma$  subunits of the IL-15 receptor and comprise IL-15 covalently or noncovalently bound to interleukin-15 receptor alpha ("IL-15Ra") ("IL-15/IL-15Ra complexes"). The IL-15/IL-15Ra complex may comprise native IL-15 or an IL-15 derivative covalently or noncovalently bound to native IL-15Ra or an IL-15Ra derivative. In one embodiment, the IL-15/IL-15Ra complex comprises an IL-15Ra derivative and the IL-15Ra derivative is a soluble form of the native IL-15Ra. In another embodiment, the IL-15/IL-15Ra complex comprises an IL-15Ra derivative and the IL-15Ra derivative comprises mutations that inhibits cleavage by an endogenous protease. In a specific embodiment, the extracellular domain cleavage site of IL-15Ra is replaced with a cleavage site

10 15 that is specifically recognized by a heterologous protease. In a specific embodiment, the extracellular domain cleavage site of IL-15Ra that is cleaved by an endogenous processing enzyme is replaced with a heterologous domain (*e.g.*, heterologous transmembrane domain) or a synthetic amino acid sequence that does not allow cleavage and generation of soluble IL-15Ra. In certain embodiments, the extracellular domain cleavage site of IL-15Ra that is cleaved by an endogenous processing enzyme is mutated to inhibit cleavage and generation of soluble IL-15Ra. In one embodiment, the extracellular domain cleavage site of IL-15Ra is replaced with a heterologous extracellular domain cleavage site (*e.g.*, heterologous transmembrane domain that is recognized and cleaved by another enzyme unrelated to the endogenous processing enzyme that cleaves the IL-15Ra).

[0012] In addition to IL-15 and IL-15Ra, the IL-15/IL-15Ra complexes may comprise a heterologous molecule. The heterologous molecule may be conjugated to IL-15 and/or IL-15Ra. The heterologous molecule is conjugated to IL-15 or IL-15Ra in a manner that does not interfere or prevent IL-15 and IL-15Ra from binding to one another and does not interfere or prevent the interaction between the IL-15/IL-15Ra complex and the  $\beta\gamma$  subunits of the IL-15 receptor. In some embodiments, the heterologous molecule is an antigen associated with a disease that one intends to prevent, treat and/or manage. Non-limiting examples of such antigens include viral antigens, bacterial antigens, parasitic antigens, and tumor antigens. In other embodiments, the heterologous molecule is an antibody that specifically binds to an antigen associated with a disease that one intends to prevent, treat and/or manage. In some embodiments, the antibody specifically binds to a cellular antigen (*e.g.*, a receptor) expressed by a cell that one desires to target. In some embodiments, the heterologous molecule increases protein stability. In certain embodiments, the heterologous molecule is an Fc domain of an immunoglobulin or a fragment thereof. In other embodiments, the heterologous molecules is not an Fc domain of an immunoglobulin molecule or a fragment thereof.

[0013] The IL-15/IL-15Ra complexes may be formulated for administration to a human subject to enhance an IL-15-mediated immune function. In a specific embodiment, the IL-15/IL-15Ra complexes are formulated with a polymer, such as poly- $\beta$ -1 $\rightarrow$ 4-N-acetylglucosamine, for administration to a subject (preferably, a human subject). The IL-15/IL-15Ra complexes may also be administered to a non-human subject for veterinary uses and/or to produce antibodies that specifically bind to IL-15/IL-15Ra complexes.

[0014] In specific embodiments, the present invention encompasses a method for enhancing an IL-15-mediated immune function in a human subject, comprising administering to a human subject in need thereof a composition comprising an effective amount of an IL-15/IL-15Ra complex formulated with a poly- $\beta$ -1 $\rightarrow$  $\beta$ -N-acetylglucosamine polymer, wherein the IL-15/IL-15Ra complex comprises human IL-15 or a derivative thereof covalently or noncovalently linked to human IL-15Ra or a derivative thereof. In a further embodiment, the IL-15/IL-15Ra complex comprises human IL-15 and human IL-15Ra. In another embodiment, the IL-15/IL-15Ra complex comprises human IL-15 and a human IL-15Ra derivative. In yet another embodiment, the human IL-15Ra or human IL-15Ra derivative is soluble. In a particular embodiment, the method further comprises administering to the human subject one or more other therapeutic polypeptides (*e.g.*, cytokines or growthfactors) or therapy.

[0015] The Agonistic Therapeutic Agents also include nucleic acids encoding IL-15 and IL-15Ra that when expressed produce IL-15/IL-15Ra complexes, and cells engineered to recombinantly express IL-15/IL-15Ra complexes by introducing nucleic acids encoding IL-15 and IL-15Ra into the cells. The nucleic acids may be administered to a subject (preferably, a human subject) as part of a gene therapy protocol. In a specific embodiment, the nucleic acids are formulated with a polymer, such as poly- $\beta$ -1 $\rightarrow$ 4-N-acetylglucosamine, for administration to a subject (preferably, a human subject). Alternatively, the nucleic acids may be transfected (in a specific embodiment, stably transfected) into cells to produce large quantities of IL-15/IL-15Ra complex suitable for *in vitro* and/or *in vivo* uses. In one embodiment, the cells engineered to express the nucleic acids are cell lines. In another embodiment, the cells engineered to express the nucleic acids are primary cells from a subject (preferably, a human subject). In a specific embodiment, the cells engineered to express the nucleic acids are cancer cells or cells infected with a pathogen.

**[0016]** Cells engineered to express IL-15 and IL-15Ra may be used to generate large quantities of IL-15/IL-15Ra complex suitable for *in vitro* and *in vivo* uses. Cells engineered to express IL-15 and IL-15Ra may also be administered to a subject (preferably, a human subject) as part of a gene therapy protocol. In a specific embodiment, irradiated cancer cells engineered to express IL-15 and IL-15Ra are administered to a cancer patient. In a particular embodiment, cells engineered to express IL-15/IL-15Ra complexes are formulated for administration to a human subject to enhance an IL-15-mediated immune function. In another embodiment, the cells engineered to express IL-15/IL-15Ra complexes are formulated with a polymer, such as poly- $\beta$ -1 $\rightarrow$ 4-N- acetylglucosamine, for administration to a subject (preferably, a human subject). The present invention also relates to an irradiated cancer cell recombinantly expressing human IL-15 and human IL-15Ra produced by a method described herein, and a pharmaceutical composition comprising an irradiated cancer cell described herein. In a particular embodiment, the pharmaceutical composition comprises an irradiated cancer cell formulated with a poly- $\beta$ -1 $\rightarrow$ 4-N- acetylglucosamine polymer.

**[0017]** One aspect of the present invention relates to a method for making irradiated cancer cells recombinantly expressing human IL-15 and human IL-15Ra, said method comprises the steps of: (i) isolating cancer cells from a subject diagnosed with cancer; (ii) introducing a nucleic acid construct(s) encoding recombinant human IL-15 or a derivative thereof and human IL-15Ra or a derivative thereof; and (iii) irradiating said cancer cells. In a particular embodiment, the human IL-15Ra or human IL-15Ra derivative is soluble.

**[0018]** In specific embodiments, the present invention provides for a cell that recombinantly expresses a mammalian IL-15 or a derivative thereof and a mammalian IL-15Ra or a derivative thereof, wherein the cell expresses at least 0.6 pg of mammalian IL-15 or a derivative thereof. In particular embodiments, the cell expressing at least 0.6 pg of mammalian IL-15 or a derivative thereof grows in serum-free media.

**[0019]** The present invention relates to a method for treating cancer in a human subject, comprising administering to a human subject in need thereof a composition comprising irradiated cancer cells engineered to recombinantly co-express (i) human IL-15 or a derivative thereof, and (ii) human IL-15Ra or a derivative thereof. In a particular embodiment, the cancer cells are obtained by isolating cancer cells from the subject. In a further embodiment, the irradiated cancer cells are obtained by engineering cancer cells to recombinantly co-express human IL-15 or a derivative thereof and human IL-15Ra or a derivative thereof (prior to irradiation). In another embodiment, the irradiated cancer cells are obtained by engineering cancer cells to recombinantly co-express human IL-15 and a human IL-15Ra derivative (prior to irradiation). In yet another embodiment, the human IL-15Ra or human IL-15Ra derivative is soluble. In a certain embodiment, the irradiated cancer cells further recombinantly express one or more other therapeutic polypeptides (e.g., cytokines or growthfactors).

**[0020]** The present invention also provides Therapeutic Agents that reduce or inhibit IL-15 signal transduction and suppress IL-15-mediated immune function (*i.e.*, Antagonistic Therapeutic Agents"). The Antagonistic Therapeutic Agents include antibodies that specifically bind to an IL-15/IL-15Ra complex and prevent endogenous IL-15/IL-15Ra complexes from binding to the  $\beta\gamma$  subunits of the IL-15 receptor, and cells engineered to express such antibodies. The Antagonistic Therapeutic Agents are useful for suppressing IL-15-mediated immune function in a subject in need of such therapy. In particular, the Antagonistic Therapeutic Agents are useful for the prevention, treatment and/or management of disorders in which it is beneficial to suppress IL-15-mediated immune function. Non-limiting examples of such disorders include autoimmune disorders, graft versus host disease, transplantation rejection, and inflammatory disorders. In a specific embodiment, the invention provides a method for treating an autoimmune disorder or inflammatory disorder in a human subject, comprising administering to a human subject in need thereof an effective amount of an antibody that specifically binds to endogenous IL-15/IL-15Ra complexes. In a further embodiment, the antibody specifically binds to endogenous IL-15/IL-15Ra complexes and does not specifically bind to IL-15 alone or IL-15Ra alone when not complexed to each other. In specific embodiments, the present invention relates to a method for treating an autoimmune disorder or inflammatory disorder in a human subject, comprising administering to a human subject in need thereof an effective amount of an antibody that specifically binds to IL-15/IL-15Ra complexes and reduces the binding of the IL-15/IL-15Ra complexes to the beta-gamma receptor complex as determined in cell culture or *in vitro*. In a further embodiment of the method, the antibody is a monoclonal humanized antibody.

### 3.1. TERMINOLOGY

**[0021]** As used herein, the terms "about" and "approximately," when used to modify numeric value or numeric range, indicate that reasonable deviations from the value or range, typically 5% or 10% above and 5% or 10% below the value or range, remain within the intended meaning of the recited value or range.

**[0022]** As used herein, the terms "antibody" and "antibodies" refer to molecules that contain an antigen binding site, *e.g.*, immunoglobulins. Antibodies include, but are not limited to, monoclonal antibodies, multispecific antibodies, human antibodies, humanized antibodies, synthetic antibodies, chimeric antibodies, polyclonal antibodies, single domain antibodies, camelized antibodies, single-chain Fvs (scFv), single chain antibodies, Fab fragments, F(ab') fragments, disulfide-linked bispecific Fvs (sdFv), intrabodies, and anti-idiotypic (anti-Id) antibodies (including, *e.g.*, anti-Id and anti-anti-Id

antibodies to antibodies), and epitope-binding fragments of any of the above. In particular, antibodies include immunoglobulin molecules and immunologically active fragments of immunoglobulin molecules. Immunoglobulin molecules can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass.

**[0023]** As used herein, the terms "disease" and "disorder" are used interchangeably to refer to a condition, in particular, a pathological condition, and more particularly a disease affected by IL-15 signal transduction.

**[0024]** As used herein, the terms "immunospecifically binds," "immunospecifically recognizes," "specifically binds," "specifically recognizes" and analogous terms in the context of antibodies refer to molecules that specifically bind to an antigen (e.g., epitope or immune complex) and do not specifically bind to another molecule. A molecule that specifically binds to an antigen may bind to other peptides or polypeptides with lower affinity as determined by, e.g., immunoassays, BIACore, or other assays known in the art. Preferably, molecules that specifically bind an antigen do not cross react with other proteins. Molecules that specifically bind an antigen can be identified, for example, by immunoassays, BIACore, or other techniques known to those of skill in the art.

**[0025]** As used herein, the terms "specifically binds," "specifically recognizes" and analogous terms in the context of receptor (e.g., native IL-15Ra) and ligand (e.g., native IL-15) interaction refer to the specific binding or association between the ligand and receptor. Preferably, the ligand has higher affinity for the receptor than for other molecules. In a specific embodiment, the ligand is native IL-15 and the native receptor is IL-15Ra. In another specific embodiment, the ligand is the native IL-15/IL-15Ra complex and the native receptor is the  $\beta\gamma$  receptor complex. In a further embodiment, the IL-15/IL-15Ra complex binds to the  $\beta\gamma$  receptor complex and activates IL-15 mediated signal transduction. Ligands that specifically bind a receptor can be identified, for example, by immunoassays, BIAcore, or other techniques known to those of skill in the art.

[0026] As used herein, the terms "native IL-15" and "native interleukin-15" in the context of proteins or polypeptides refer to any naturally occurring mammalian interleukin-15 amino acid sequences, including immature or precursor and mature forms. Non-limiting examples of GeneBank Accession Nos. for the amino acid sequence of various species of native mammalian interleukin-15 include NP\_000576 (human, immature form), CAA62616 (human, immature form), NP\_001009207 (Felis catus, immature form), AAB94536 (rattus, immature form), AAB41697 (rattus, immature form), NP\_032383 (Mus musculus, immature form), AAR19080 (canine), AAB60398 (macaca mulatta, immature form), AAI00964 (human, immature form), AAH23698 (mus musculus, immature form), and AAH18149 (human). The amino acid sequence of the immature/precursor form of native human IL-15, which comprises the long signal peptide (underlined) and the mature human native IL-15 (*italicized*), is provided:

MRISKPHLRSISIQCYLCLLLNSHFLTEAGIHVFILGCFSAGLPKTEANWVNVISDLKKIE  
DLIQSMHIDATLYTESDVHPSCKVAMKCFLLELQVISLESGDASIHDTVENLIIANNSLSS NGNVTESGCK-  
ECEELEEKNIKEFLQSFVHIVQMFINTS (SEQ ID NO: 1). In some embodiments, native IL-15 is the immature or precursor form of a naturally occurring mammalian IL-15. In other embodiments, native IL-15 is the mature form of a naturally occurring mammalian IL-15. In a specific embodiment, native IL-15 is the precursor form of naturally occurring human IL-15. In another embodiment, native IL-15 is the mature form of naturally occurring human IL-15. In one embodiment, the native IL-15 protein/polypeptide is isolated or purified.

**[0028]** As used herein, the terms "IL-15 derivative" and "interleukin-15 derivative" in the context of proteins or polypeptides refer to: (a) a polypeptide that is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical to a native mammalian IL-15 polypeptide; (b) a polypeptide encoded by a nucleic acid sequence that is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical a nucleic acid sequence encoding a native mammalian IL-15 polypeptide; (c) a polypeptide that contains 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more amino acid mutations (*i.e.*, additions, deletions and/or substitutions) relative

to a native mammalian IL-15 polypeptide; (d) a polypeptide encoded by nucleic acids can hybridize under high, moderate or typical stringency hybridization conditions to nucleic acids encoding a native mammalian IL-15 polypeptide; (e) a polypeptide encoded by a nucleic acid sequence that can hybridize under high, moderate or typical stringency hybridization conditions to a nucleic acid sequence encoding a fragment of a native mammalian IL-15 polypeptide of at least 5

20 contiguous amino acids, at least 30 contiguous amino acids, at least 40 contiguous amino acids, at least 50 contiguous amino acids, at least 100 contiguous amino acids, or at least 150 contiguous amino acids; or (f) a fragment of a native mammalian IL-15 polypeptide. IL-15 derivatives also include a polypeptide that comprises the amino acid sequence of a naturally occurring mature form of a mammalian IL-15 polypeptide and a heterologous signal peptide amino acid sequence. In a specific embodiment, an IL-15 derivative is a derivative of a native human IL-15 polypeptide. In another embodiment, an IL-15 derivative is a derivative of an immature or precursor form of naturally occurring human IL-15 10 polypeptide. In another embodiment, an IL-15 derivative is a derivative of a mature form of naturally occurring human IL-15 polypeptide. In one embodiment, an IL-15 derivative is isolated or purified.

**[0029]** In a preferred embodiment, IL-15 derivatives retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% of the function of native mammalian IL-15 polypeptide to bind IL-15Ra polypeptide, as measured by assays well known in the art, e.g., ELISA, Biacore, co-immunoprecipitation. In another preferred embodiment, IL-15 derivatives retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% of the function of native mammalian IL-15 polypeptide to induce IL-15-mediated signal transduction, as measured by assays well-known in the art, e.g., electromobility shift assays, ELISAs and other immunoassays.

**[0030]** Percent identity can be determined using any method known to one of skill in the art. In a specific embodiment, the percent identity is determined using the "Best Fit" or "Gap" program of the Sequence Analysis Software Package (Version 10; Genetics Computer Group, Inc., University of Wisconsin Biotechnology Center, Madison, Wisconsin). Information regarding hybridization conditions (e.g., high, moderate, and typical stringency conditions) have been described, see, e.g., U.S. Patent Application Publication No. US 2005/0048549 (e.g., paragraphs 72-73).

**[0031]** As used herein, the terms "IL-15 derivative" and "interleukin-15 derivative" in the context of nucleic acids refer to: (a) a nucleic acid sequence that is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical to the naturally occurring nucleic acid sequence encoding a mammalian IL-15 polypeptide; (b) a nucleic acid sequence encoding a polypeptide that is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical the amino acid sequence of a native mammalian IL-15 polypeptide; (c) a nucleic acid sequence that contains 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more nucleic acid base mutations (i.e., additions, deletions and/or substitutions) relative to the naturally occurring nucleic acid sequence encoding a mammalian IL-15 polypeptide; (d) a nucleic acid sequence that hybridizes under high, moderate or typical stringency hybridization conditions to a naturally occurring nucleic acid sequence encoding a mammalian IL-15 polypeptide; (e) a nucleic acid sequence that hybridizes under high, moderate or typical stringency hybridization conditions to a fragment of a naturally occurring nucleic acid sequence encoding a mammalian IL-15 polypeptide; and (f) a nucleic acid sequence encoding a fragment of a naturally occurring nucleic acid sequence encoding a mammalian IL-15 polypeptide. In a specific embodiment, an IL-15 derivative in the context of nucleic acids is a derivative of a naturally occurring nucleic acid sequence encoding a human IL-15 polypeptide. In another embodiment, an IL-15 derivative in the context of nucleic acids is a derivative of a naturally occurring nucleic acid sequence encoding an immature or precursor form of a human IL-15 polypeptide. In another embodiment, an IL-15 derivative in the context of nucleic acids is a derivative of a naturally 30 occurring nucleic acid sequence encoding a mature form of a human IL-15 polypeptide.

**[0032]** IL-15 derivative nucleic acid sequences include codon-optimized nucleic acid sequences that encode native mammalian IL-15 polypeptide, including mature and immature forms of IL-15 polypeptide. In other embodiments, IL-15 derivative nucleic acids include nucleic acids that encode mammalian IL-15 RNA transcripts containing mutations that eliminate potential splice sites and instability elements (e.g., A/T or A/U rich elements) without affecting the amino acid 40 sequence to increase the stability of the mammalian IL-15 RNA transcripts.

**[0033]** In a preferred embodiment, IL-15 derivative nucleic acid sequences encode proteins or polypeptides that retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% of the function of a native mammalian IL-15 polypeptide to bind IL-15Ra, as measured by assays well known in the art, e.g., ELISA, Biacore, co-immunoprecipitation. In another preferred embodiment, IL-15 derivative nucleic acid sequences encode proteins or polypeptides that retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% of the function of a native mammalian IL-15 polypeptide to induce IL-15-mediated signal transduction, as measured by assays well-known in the art, e.g., electromobility shift assays, ELISAs and other immunoassays.

**[0034]** As used herein, the terms "IL-15" and "interleukin-15" refer to a native IL-15, an IL-15 derivative, or a native IL-15 and an IL-15 derivative.

**[0035]** As used herein, the terms "native IL-15Ra" and "native interleukin-15 receptor alpha" in the context of proteins or polypeptides refer to any naturally occurring mammalian interleukin-15 receptor alpha ("IL-15Ra") amino acid sequence, including immature or precursor and mature forms and naturally occurring isoforms. Non-limiting examples of GeneBank Accession Nos. for the amino acid sequence of various native mammalian IL-15Ra include NP\_002180

(human), ABK41438 (Macaca mulatta), NP\_032384 (Mus musculus), Q60819 (Mus musculus), CAI41082 (human). The amino acid sequence of the immature form of the native full length human IL-15Ra, which comprises the signal peptide (underlined) and the mature human native IL-15Ra (italicized), is provided: MAPRRARGCR TLGPALLL LLLRPPATRG ITCPPPMSE HADIWVKSYS LYSRERYICN SGFKRKAGTS SLTECVLNKA TNVAHWTPS LKCIRDPA LV HQR-PAPPSTV TTAGVTPQPE SLSPSGKEPA ASSPSSNNTA ATTAAIVPGS QLMPSKSPST GTTEISSHES SHGTPSQTTA KNWELTASAS HQPPGVYPQG HSDTTVAIST STVLLCGLSA VSLLACYLKS RQTPLASVE MEAMEALPVT WGTSS- RDEDL ENCSHHL (SEQ ID NO: 3). The amino acid sequence of the immature form of the native soluble human IL-15Ra, which comprises the signal peptide (underlined) and the mature human native IL-15Ra (italicized), is provided: MAPRRARGCR TLGPALLL LLLRPPATRG ITCPPPMSE HADIWVKSYS LYSRERYICN SGFKRKAGTS SLTECV- LNKA TNVAHWTPS LKCIRDPA LV HQR-PAPPSTV TTAGVTPQPE SLSPSGKEPA ASSPSSNNTA ATTAAIVPGS QLMPSKSPST GTTEISSHES SHGTPSQTTA KNWELTASAS HQPPGVYPQG HSDTT (SEQ ID NO: 4). In some embodiments, native IL-15Ra is the immature form of a naturally occurring mammalian IL-15Ra polypeptide. In other embodiments, native IL-15Ra is the mature form of a naturally occurring mammalian IL-15Ra polypeptide. In certain embodiments, native IL-15Ra is a soluble form of a naturally occurring mammalian IL-15Ra polypeptide. In other embodiments, native IL-15Ra is the full-length form of a naturally occurring mammalian IL-15Ra polypeptide. In one embodiment, a native IL-15Ra protein or polypeptide is isolated or purified.

**[0036]** As used herein, the terms "native IL-15Ra" and "native interleukin-15 receptor alpha" in the context of nucleic acids refer to any naturally occurring nucleic acid sequences encoding mammalian interleukin-15 receptor alpha, including the immature or precursor and mature forms. Non-limiting examples of GeneBank Accession Nos. for the nucleotide sequence of various species of native mammalian IL-15Ra include NM\_002189 (human), EF033114 (Macaca mulatta), and NM\_008358 (Mus musculus). The nucleotide sequence encoding the immature form of native human IL-15Ra, which comprises the nucleotide sequence encoding the signal peptide (underlined) and the nucleotide sequence encoding the mature human native IL-15Ra (italicized), is provided: atggccccc gcgccggcg cgccggctgcc ggaccctcg tctcccgcc ctgctactgc tgctgctgc ccggccgcgg gcacacgctgg ccatccatgc aatccatgc aacacgcaga catctgggtc aagagctaca gcttggactc cagggagcgg tacatgttgc actctggttt caagcgtaaa gcccggcacgt ccagctgtac ggatgtcg ttgaacaagg ccacaaatgt cgcccaactgg acaaccccca gtctcaaatg cattagagac cctggccctgg ttcaccaaaag gccagcccca ccctccacag taacgcacggc agggggtgacc ccacagccat cccttcgtgg aaagagcccg cagcttcatc tcccaatgc aacaacacag cgcccaacaac agcagctatt gtccgggc cccaatgt gcctcaaaa tcacccatca caggaaccac agagataaagc agtcatgatg cctccacgg cacccctct cagacaacag ccaagaactg ggaactcaca gcatccgc cccaccagcc gccagggtgt tatccacagg gccacagcg caccactgt gtctatccca cgttccactgt cctgtgtgt gggtctgatc ctgtgtct cctggcatgc tacatgttgc caagcgtaaa tcacccatgg ttcaccaaaag gccagcccca ccctccacag taacgcacggc agggggtgacc ccacagccat aggggtgacc ccacagccat aaagagcccg cagcttcatc tcccaatgc aacaacacag cgcccaacaac agcagctatt gtccgggc cccaatgt gcctcaaaa tcacccatca caggaaccac agagataaagc agtcatgatg cctccacgg cacccctct cagacaacag ccaagaactg ggaactcaca gcatccgc cccaccagcc gccagggtgt tatccacagg gccacagcg caccactgt (SEQ ID NO: 5). The nucleotide sequence encoding the immature form of native soluble human IL-15Ra protein or polypeptide, which comprises the nucleotide sequence encoding the signal peptide (underlined) and the nucleotide sequence encoding the mature human soluble native IL-15Ra (italicized), is provided: atggccccc gcgccggcg cgccggctgcc ggaccctcg tctcccgcc ctgctactgc tgctgctgc ccggccgcgg gcacacgctgg ccatccatgc aatccatgc aacacgcaga catctgggtc aagagctaca gcttggactc cagggagcgg tacatgttgc actctggttt caagcgtaaa gcccggcacgt ccagctgtac ggatgtcg ttgaacaagg ccacaaatgt cgcccaactgg acaaccccca gtctcaaatg cattagagac cctggccctgg ttcaccaaaag gccagcccca ccctccacag taacgcacggc agggggtgacc ccacagccat cccttcgtgg aaagagcccg cagcttcatc tcccaatgc aacaacacag cgcccaacaac agcagctatt gtccgggc cccaatgt gcctcaaaa tcacccatca caggaaccac agagataaagc agtcatgatg cctccacgg cacccctct cagacaacag ccaagaactg ggaactcaca gcatccgc cccaccagcc gccagggtgt tatccacagg gccacagcg caccactgt (SEQ ID NO: 6). In a specific embodiment, the nucleic acid is an isolated or purified nucleic acid. In some embodiments, naturally occurring nucleic acids encode the immature form of a naturally occurring mammalian IL-15Ra polypeptide. In other embodiments, naturally occurring nucleic acids encode the mature form of a naturally occurring mammalian IL-15Ra polypeptide. In certain embodiments, naturally occurring nucleic acids encode the soluble form of a naturally occurring mammalian IL-15Ra polypeptide. In other embodiments, naturally occurring nucleic acids encode the full-length form of a naturally occurring mammalian IL-15Ra polypeptide. In a specific embodiment, naturally occurring nucleic acids encode the precursor form of naturally occurring human IL-15 polypeptide. In another embodiment, naturally occurring nucleic acids encode the mature of naturally occurring human IL-15 polypeptide. In certain embodiments, naturally occurring nucleic acids encode the soluble form of a naturally occurring human IL-15Ra polypeptide. In other embodiments, naturally occurring nucleic acids encode the full-length form of a naturally occurring human IL-15Ra polypeptide.

**[0037]** As used herein, the terms "IL-15Ra derivative" and "interleukin-15 receptor alpha derivative" in the context of a protein or polypeptide refer to: (a) a polypeptide that is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical to a native mammalian IL-15 polypeptide; (b) a polypeptide encoded by a nucleic acid sequence that is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical

a nucleic acid sequence encoding a native mammalian IL-15Ra polypeptide; (c) a polypeptide that contains 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more amino acid mutations (*i.e.*, additions, deletions and/or substitutions) relative to a native mammalian IL-15Ra polypeptide; (d) a polypeptide encoded by a nucleic acid sequence that can hybridize under high, moderate or typical stringency hybridization conditions to a nucleic acid sequence encoding a native mammalian IL-15Ra polypeptide; (e) a polypeptide encoded by a nucleic acid sequence that can hybridize under high, moderate or typical stringency hybridization conditions to nucleic acid sequences encoding a fragment of a native mammalian IL-15 polypeptide of at least 20 contiguous amino acids, at least 30 contiguous amino acids, at least 40 contiguous amino acids, at least 50 contiguous amino acids, at least 100 contiguous amino acids, or at least 150 contiguous amino acids; or (f) a fragment of a native mammalian IL-15Ra polypeptide. IL-15Ra derivatives also include a polypeptide that comprises the amino acid sequence of a naturally occurring mature form of mammalian IL-15Ra polypeptide and a heterologous signal peptide amino acid sequence. In a specific embodiment, an IL-15Ra derivative is a derivative of a native human IL-15Ra polypeptide. In another embodiment, an IL-15Ra derivative is a derivative of an immature form of naturally occurring human IL-15 polypeptide. In another embodiment, an IL-15Ra derivative is a derivative of a mature form of naturally occurring human IL-15 polypeptide. In one embodiment, an IL-15Ra derivative is the soluble form of a native mammalian IL-15Ra polypeptide. In a specific embodiment, an IL-15Ra derivative is purified or isolated.

**[0038]** In a preferred embodiment, IL-15Ra derivatives retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% of the function of a native mammalian IL-15Ra polypeptide to bind an IL-15 polypeptide, as measured by assays well known in the art, *e.g.*, ELISA, Biacore, co-immunoprecipitation. In another preferred embodiment, IL-15Ra derivatives retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% of the function of a native mammalian IL-15Ra polypeptide to induce IL-15-mediated signal transduction, as measured by assays well-known in the art, *e.g.*, electromobility shift assays, ELISAs and other immunoassays.

**[0039]** As used herein, the terms "IL-15Ra derivative" and "interleukin-15 receptor alpha derivative" in the context of nucleic acids refer to: (a) a nucleic acid sequence that is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical to the naturally occurring nucleic acid sequence encoding a mammalian IL-15Ra polypeptide; (b) a nucleic acid sequence encoding a polypeptide that is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical to the amino acid sequence of a native mammalian IL-15Ra polypeptide; (c) a nucleic acid sequence that contains 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more nucleic acid mutations (*i.e.*, additions, deletions and/or substitutions) relative to the naturally occurring nucleic acid sequence encoding a mammalian IL-15Ra polypeptide; (d) a nucleic acid sequence that hybridizes under high, moderate or typical stringency hybridization conditions to a naturally occurring nucleic acid sequence encoding a mammalian IL-15Ra polypeptide; (e) a nucleic acid sequence that hybridizes under high, moderate or typical stringency hybridization conditions to a fragment of a naturally occurring nucleic acid sequence encoding a mammalian IL-15Ra polypeptide; and (f) a nucleic acid sequence encoding a fragment of a naturally occurring nucleic acid sequence encoding a mammalian IL-15Ra polypeptide. In a specific embodiment, an IL-15Ra derivative in the context of nucleic acids is a derivative of a naturally occurring nucleic acid sequence encoding a human IL-15Ra polypeptide. In another embodiment, an IL-15Ra derivative in the context of nucleic acids is a derivative of a naturally occurring nucleic acid sequence encoding an immature form of a human IL-15Ra polypeptide. In another embodiment, an IL-15Ra derivative in the context of nucleic acids is a derivative of a naturally occurring nucleic acid sequence encoding a mature form of a human IL-15Ra polypeptide. In one embodiment, an IL-15Ra derivative refers to a nucleic acid sequence encoding a mammalian IL-15Ra polypeptide that is soluble.

**[0040]** IL-15Ra derivative nucleic acid sequences include codon-optimized nucleic acid sequences that encode native IL-15Ra polypeptide, including mature and immature forms of IL-15Ra polypeptide. In other embodiments, IL-15Ra derivative nucleic acids include nucleic acids that encode IL-15Ra RNA transcripts containing mutations that eliminate potential splice sites and instability elements (*e.g.*, A/T or A/M rich elements) without affecting the amino acid sequence to increase the stability of the IL-15Ra RNA transcripts.

**[0041]** In a preferred embodiment, IL-15Ra derivative nucleic acid sequences encode proteins or polypeptides that retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% of the function of a native mammalian IL-15Ra polypeptide to bind IL-15, as measured by assays well known in the art, *e.g.*, ELISA, Biacore, co-immunoprecipitation. In another preferred embodiment, IL-15Ra derivative nucleic acid sequences encode proteins or polypeptides that retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% of the function of a native mammalian IL-15Ra to induce IL-15-mediated signal transduction, as measured by assays well-known in the art, *e.g.*, electromobility shift assays, ELISAs and other immunoassays.

**[0042]** As used herein, the terms "IL-15Ra" and "interleukin-15 receptor alpha" refer to a native IL-15Ra, an IL-15Ra derivative, or a native IL-15Ra and an IL-15Ra derivative.

**[0043]** As used herein, the term "IL-15/IL-15Ra complex" refers to a complex comprising IL-15 and IL-15Ra covalently or noncovalently bound to each other. In a preferred embodiment, the IL-15Ra has a relatively high affinity for IL-15, *e.g.*,  $K_d$  of 10 to 50 pM as measured by a technique known in the art, *e.g.*, KinEx A assay, plasma surface resonance

(e.g., BIACore assay). In another preferred embodiment, the IL-15/IL-15Ra complex induces IL-15-mediated signal transduction, as measured by assays well-known in the art, e.g., electromobility shift assays, ELISAs and other immunoassays. In some embodiments, the IL-15/IL-15Ra complex retains the ability to specifically bind to the  $\beta\gamma$  chain.

**[0044]** As used herein, the terms "subject" and "patient" are used interchangeably and refer to a mammal such as a non-primate (e.g., cows, pigs, horses, cats, dogs, rats etc.) and a primate (e.g., monkey and human), most preferably a human.

**[0045]** As used herein, the terms "purified" and "isolated" in the context of a compound or agent (including, e.g., proteinaceous agents such as antibodies) that is chemically synthesized refers to a compound or agent that is substantially free of chemical precursors or other chemicals when chemically synthesized. In a specific embodiment, the compound or agent is 60%, 65%, 75%, 80%, 85%, 90%, 95%, or 99% free (by dry weight) of other, different compounds or agents.

**[0046]** As used herein, the terms "purified" and "isolated" when used in the context of a compound or agent (including proteinaceous agents such as antibodies and polypeptides) that can be obtained from a natural source, e.g., cells, refers to a compound or agent which is substantially free of contaminating materials from the natural source, e.g., soil particles, minerals, chemicals from the environment, and/or cellular materials from the natural source, such as but not limited to cell debris, cell wall materials, membranes, organelles, the bulk of the nucleic acids, carbohydrates, proteins, and/or lipids present in cells. The phrase "substantially free of natural source materials" refers to preparations of a compound or agent that has been separated from the material (e.g., cellular components of the cells) from which it is isolated. Thus, a compound or agent that is isolated includes preparations of a compound or agent having less than about 30%, 20%, 10%, 5%, 2%, or 1% (by dry weight) of cellular materials and/or contaminating materials.

**[0047]** An "isolated" nucleic acid sequence or nucleotide sequence is one which is separated from other nucleic acid molecules which are present in a natural source of the nucleic acid sequence or nucleotides sequence. Moreover, an "isolated", nucleic acid sequence or nucleotide sequence, such as a cDNA molecule, can be substantially free of other cellular material or culture medium when produced by recombinant techniques, or substantially free of chemical precursors when chemically synthesized. In certain embodiments, an "isolated" nucleic acid sequence or nucleotide sequence is a nucleic acid sequence or nucleotide sequence that is recombinantly expressed in a heterologous cell.

**[0048]** In some embodiments, the terms "nucleic acid", "nucleotide" and "polynucleotide" refer to deoxyribonucleotides, deoxyribonucleic acids, ribonucleotides, and ribonucleic acids, and polymeric forms thereof, and include either single- or double-stranded forms. In certain embodiments, such terms include known analogues of natural nucleotides, for example, peptide nucleic acids ("PNA"s), that have similar binding properties as the reference nucleic acid. In some embodiments, such terms refer to deoxyribonucleic acids (e.g., cDNA or DNA). In other embodiments, such terms refer to ribonucleic acid (e.g., mRNA or RNA).

**[0049]** As used herein, the terms "therapies" and "therapy" can refer to any protocol(s), method(s), compositions, formulations, and/or agent(s) that can be used in the prevention, treatment, management, or amelioration of a disease, e.g., cancer, infectious disease, autoimmune disease, graft versus host disease, and transplantation rejection, or a symptom associated therewith. In certain embodiments, the terms "therapies" and "therapy" refer to biological therapy, supportive therapy, and/or other therapies useful in treatment, management, prevention, or amelioration of a disease or a symptom associated therewith known to one of skill in the art. In one embodiment, a therapy includes an Agonistic Therapeutic Agent. In one embodiment, a therapy includes an Antagonistic Therapeutic Agent. In one embodiment, a therapy is not an Agonistic Therapeutic Agent. In one embodiment, a therapy is not an Antagonistic Therapeutic Agent.

**[0050]** As used herein, the terms "protein(s)" and "polypeptide(s)" interchangeably to refer to a chain of amino acids linked together by peptide bonds. In some embodiments, the terms "protein(s)" and "polypeptide(s)" refer to a macromolecule which comprises amino acids that are linked together by peptide bonds.

**[0051]** As used herein, the term "fragment" in the context of a nucleotide sequence refers to a nucleotide sequence comprising an nucleic acid sequence of at least 5 contiguous nucleic acid bases, at least 10 contiguous nucleic acid bases, at least 15 contiguous nucleic acid bases, at least 20 contiguous nucleic acid bases, at least 25 contiguous nucleic acid bases, at least 40 contiguous nucleic acid bases, at least 50 contiguous nucleic acid bases, at least 60 contiguous nucleic acid bases, at least 70 contiguous nucleic acid bases, at least 80 contiguous nucleic acid bases, at least 90 contiguous nucleic acid bases, at least 100 contiguous nucleic acid bases, at least 125 contiguous nucleic acid bases, at least 150 contiguous nucleic acid bases, at least 175 contiguous nucleic acid bases, at least 200 contiguous nucleic acid bases, or at least 250 contiguous nucleic acid bases of the nucleotide sequence of the gene of interest, e.g., *IL-15*, *IL-15Ra*. The nucleic acid may be RNA, DNA, or a chemically modified variant thereof. In a specific embodiment, the fragment is a fragment of *IL-15* or *IL-15Ra*.

**[0052]** As used herein, the term "fragment" in the context of a fragment of a proteinaceous agent (e.g., a protein or polypeptide) refers to a fragment that is composed of 8 or more contiguous amino acids, 10 or more contiguous amino acids, 15 or more contiguous amino acids, 20 or more contiguous amino acids, 25 or more contiguous amino acids, 50 or more contiguous amino acids, 75 or more contiguous amino acids, 100 or more contiguous amino acids, 150 or more contiguous amino acids, 200 or more contiguous amino acids, 10 to 150 contiguous amino acids, 10 to 200 contiguous amino acids, 10 to 250 contiguous amino acids, 10 to 300 contiguous amino acids, 50 to 100 contiguous amino acids,

50 to 150 contiguous amino acids, 50 to 200 contiguous amino acids, 50 to 250 contiguous amino acids or 50 to 300 contiguous amino acids of a proteinaceous agent, e.g., IL-15 and IL-15Ra polypeptides.

**[0053]** As used herein, the term "in combination" refers to the use of more than one therapies (e.g., one or more prophylactic and/or therapeutic agents). The use of the term "in combination" does not restrict the order in which therapies are administered to a subject with a disease or disorder. A first therapy (e.g., a prophylactic or therapeutic agent) can be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concomitantly with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a second therapy (e.g., a prophylactic or therapeutic agent) to a subject with a disease or disorder or a symptom thereof.

**[0054]** As used herein, the term "premature human infant" refers to a human infant born at less than 37 weeks of gestational age.

**[0055]** As used herein, the term "human infant" refers to a newborn to 1 year old year human.

**[0056]** As used herein, the term "human child" refers to a human that is 1 year to 18 years old.

**[0057]** As used herein, the term "human adult" refers to a human that is 18 years or older.

**[0058]** As used herein, the term "elderly human" refers to a human 65 years or older.

**[0059]** As used herein, the terms "treat", "treating" and "treatment" in the context of the administration of a therapy to a subject refer to the beneficial effects that a subject derives from a therapy, such as, but not limited to, the reduction or inhibition of the progression, spread and/or duration of a disease or disorder, the reduction or amelioration of the severity of a disease or disorder, amelioration of one or more symptoms of a disease or disorder, and/or the reduction in the duration of one or more symptom of a disease or disorder resulting from the administration of one or more therapies. In specific embodiments, such terms in the context of cancer include, but are not limited to, one, two, or three or more results following the administration of a therapy to a subject: (1) a reduction in the growth of a tumor or neoplasm; (2) a reduction in the formation of a tumor; (3) an eradication, removal, or control of primary, regional and/or metastatic cancer; (4) a reduction in metastatic spread; (5) a reduction in mortality; (6) an increase in survival rate; (7) an increase in length of survival; (8) an increase in the number of patients in remission; (9) a decrease in hospitalization rate; (10) a decrease in hospitalization lengths; and (11) the maintenance in the size of the tumor so that it does not increase by more than 10%, or by more than 8%, or by more than 6%, or by more than 4%; preferably the size of the tumor does not increase by more than 2%.

**[0060]** As used herein, the terms "prevent," " preventing" and "prevention" in the context of the administration of a therapy to a subject refer to the inhibition of the onset or recurrence of a disease or disorder in a subject.

**[0061]** As used herein, the terms "manage," "managing," and "management," in the context of the administration of a therapy to a subject, refer to the beneficial effects that a subject derives from a therapy, which does not result in a cure of a disease or disorder. In certain embodiments, a subject is administered one or more therapies to "manage" a disease or disorder so as to prevent the progression or worsening of symptoms associated with a disease or disorder.

#### 4. BRIEF DESCRIPTION OF THE DRAWINGS

**[0062]** **FIGs. 1A-B:** Nucleic acid and amino acid sequences of native human IL-15. The nucleic acid sequence (FIG. 1A) (SEQ ID NO: 2) and amino acid sequence (FIG. 1B) (SEQ ID NO: 1) are shown. The amino acid sequence and nucleic acid sequence of the long signal peptide (underlined) and mature form (italicized) are indicated.

**[0063]** **FIGs. 2A-B:** Nucleic acid and amino acid sequences of full length native human IL-15Ra. The nucleic acid sequence (FIG. 2A) (SEQ ID NO: 5) and amino acid sequence (FIG. 2B) (SEQ ID NO: 3) are shown. The amino acid sequence and nucleic acid sequence of the signal peptide (underlined) and mature form (italicized) are indicated.

**[0064]** **FIGs. 3A-B:** Nucleic acid and amino acid sequences of soluble native human IL-15. The nucleic acid sequence (FIG. 3A) (SEQ ID NO: 6) and amino acid sequence (FIG. 3B) (SEQ ID NO: 4) are shown. The amino acid sequence and nucleic acid sequence of the signal peptide (underlined) and mature form (italicized) are indicated.

**[0065]** **FIGs. 4A-D:** AG32 nucleic acid construct encoding optimized human IL-15. The nucleic acid sequence (FIG. 4A) (SEQ ID NO: 9) and the amino acid sequence (FIG. 4B) (SEQ ID NO: 10) are shown. FIG. 4C shows a schematic of the nucleic acid construct comprising the CMV promoter. FIG. 4D shows an alignment of the amino acid sequence translated from the open reading frame of the nucleic acid sequence encoding optimized human IL-15 (AG32 hull15opt).

**[0066]** **FIGs. 5A-D:** AG59 nucleic acid construct encoding optimized human IL-15 with a signal peptide and pro-peptide of tPA. The nucleic acid sequence (FIG. 5A) (SEQ ID NO: 11) and the amino acid sequence (FIG. 5B) (SEQ ID NO: 12) are shown. FIG. 5C shows a schematic of the nucleic acid expression construct comprising the CMV promoter. FIG. 5D shows an alignment of the amino acid sequence translated from the open reading frame of the nucleic acid sequence encoding optimized human IL-15 (AG59 CMV hull15tPA6).

**[0067]** **FIGs. 6A-D:** AG79 nucleic acid construct encoding optimized human IL-15Ra. The nucleic acid sequence (FIG.

6A) (SEQ ID NO: 13) and the amino acid sequence (FIG. 6B) (SEQ ID NO: 14) are shown. FIG. 6C shows a schematic of the nucleic acid expression construct comprising the CMV promoter. FIG. 6D shows an alignment of the amino acid sequence translated from the open reading frame of the nucleic acid sequence encoding human IL-15Ra (AG79 hull15Ra).

5 [0068] **FIGs. 7A-D:** AG98 nucleic acid construct encoding optimized soluble human IL-15Ra. The nucleic acid sequence (FIG. 7A) (SEQ ID NO: 15) and the amino acid sequence

[0069] (FIG. 7B) (SEQ ID NO: 16) are shown. FIG. 7C shows a schematic of the nucleic acid expression construct comprising the CMV promoter. FIG. 7D shows an alignment of the amino acid sequence translated from the open reading frame of the nucleic acid sequence encoding soluble human IL-15Ra (AG98 CMV hu sIL15Ra).

10 [0070] **FIGs. 8A-D:** AG151 nucleic acid construct encoding optimized human IL-15 with a signal peptide of human GM-CSF. The nucleic acid sequence (FIG. 8A) (SEQ ID NO: 17) and the amino acid sequence (FIG. 8B) (SEQ ID NO: 18) are shown. FIG. 8C shows a schematic of the nucleic acid expression construct comprising the CMV promoter. FIG. 8D shows an alignment of the amino acid sequence translated from the open reading frame of the nucleic acid sequence encoding human IL-15 with a signal peptide of human GM-CSF (AG151 CMVhull15huGMCSF).

15

## 5. DETAILED DESCRIPTION OF THE INVENTION

[0071] The present invention relates to agents that modulate IL-15 signal transduction or function ("Therapeutic Agents") and the use of those agents to modulate immune function. The Therapeutic Agents target the interaction between IL-15 and its receptor and modulate IL-15-induced signal transduction. In a specific embodiment, the Therapeutic Agents modulate the interaction between the IL-15 receptor beta and gamma complex and complexes composed of IL-15 and IL-15Ra.

20 [0072] The present invention provides Therapeutic Agents that induce IL-15 signal transduction and enhance the immune function (*i.e.*, Agonistic Therapeutic Agents). Such Agonistic Therapeutic Agents include (i) IL-15/IL-15Ra complexes that bind to the beta/gamma receptor complex and induce IL-15-mediated signal transduction, (ii) nucleic acid sequences encoding IL-15 and IL-15Ra to form such complexes, and (iii) cells expressing such complexes in high amounts. In one embodiment, high amounts of IL-15/IL-15Ra complexes refer to amounts of IL-15/IL-15Ra complexes expressed by cells that are at least 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, or more than 20 fold higher than amounts of IL-15/IL-15Ra complexes expressed endogenously by control cells (*e.g.*, cells that have not been genetically engineered to recombinantly express IL-15, IL-15Ra, or both IL-15 and IL-15Ra, or cells comprising an empty vector). The Agonistic Therapeutic Agents are useful for the prevention, treatment and/or management of disorders in which it is beneficial to enhance certain aspects of the immune system function, in particular, immune system functions that are mediated by IL-15 signaling. Non-limiting examples of such disorders include cancer and infectious diseases.

25 [0073] The present invention also provides Therapeutic Agents that reduce or inhibit IL-15 signal transduction and suppress the immune system function (*i.e.*, Antagonistic Therapeutic Agents"). Such Antagonistic Therapeutic Agents include antibodies that immunospecifically bind to an IL-15/IL-15Ra complex and prevent endogenous IL-15/IL-15Ra from binding to the beta/gamma receptor complex and from inducing IL-15-mediated signal transduction. The Antagonistic Therapeutic Agents are useful for the prevention, treatment and/or management of disorders in which it is beneficial to 30 suppress certain aspects of the immune function, in particular, immune system functions that are mediated by IL-15 signaling. Non-limiting examples of such disorders include autoimmune diseases, inflammatory conditions, graft versus host disease, and transplant rejection.

35 [0074] In other aspects, the invention provides an Agonistic Therapeutic Agent or an Antagonistic Therapeutic Agent formulated with any natural polymer fiber suitable for biomedical use, including, but not limited to, chitin and chitosan, derived from shellfish, fungal, or microalgal sources. In a preferred embodiment, the polymer fiber is poly- $\beta$ -1 $\rightarrow$ 4-N-acetylglucosamine (p-GlcNAc), including deacetylated forms of pGlcNAc. In a specific embodiment, the Agonistic Therapeutic Agent or Antagonistic Therapeutic Agent formulated with a natural polymer is administered to a subject. In one embodiment, the Agonistic Therapeutic Agent or Antagonistic Therapeutic Agent is purified.

40 [0075] The Therapeutic Agents may be advantageously used in combination therapy. Combination therapy includes concurrent and successive administration of a Therapeutic Agent and another therapy. As used herein, the Therapeutic Agent and another therapy are said to be administered concurrently if they are administered to the patient on the same day, for example, simultaneously, or 1, 2, 3, 4, 5, 6, 7 or 8 hours apart. In contrast, the Therapeutic Agent and the therapy are said to be administered successively if they are administered to the patient on the different days, for example, the Therapeutic Agent and the therapy can be administered at a 1-day, 2-day or 3-day intervals. In the methods of the 45 present invention, administration of the Therapeutic Agent can precede or follow administration of the second therapy.

50 [0076] As a non-limiting example, the Therapeutic Agent and another therapy can be administered concurrently for a first period of time, followed by a second period of time in which the administration of the Therapeutic Agent and the other therapy is alternated.

[0077] When administered simultaneously, the Therapeutic Agent and the other therapy can be in the same pharmaceutical composition or in different pharmaceutical compositions.

[0078] The subsections below describe in more detail Therapeutic Agents, screening assays for identifying or validating Therapeutic Agents, methods for characterizing Therapeutic Agents, formulations comprising the Therapeutic Agents, and methods of using the Therapeutic Agents to modulate immune system function.

### 5.1. Therapeutic Agents

[0079] The invention provides for Therapeutic Agents that modulate IL-15-mediated function or signaling. In particular, the present invention provides Therapeutic Agents that enhance IL-15-mediated function or signaling (i.e., Agonistic Therapeutic Agents). The administration of an Agonistic Therapeutic Agent to a subject in need thereof enhances IL-15-mediated function or signaling, which in turn results in the enhancement of certain aspects of the immune function in the subject. The enhancement of the immune function can be in the form of, e.g., an antibody response (humoral response) or a cellular immune response, e.g., cytokine secretion (e.g., interferon-gamma), helper activity or cellular cytotoxicity. In one embodiment, the enhancement of the immune function is increased cytokine secretion, antibody production, effector function, T cell proliferation, and/or NK cell proliferation.

[0080] The present invention also provides Therapeutic Agents that suppress or reduce IL-15-mediated function or signaling (i.e., Antagonistic Therapeutic Agents). The administration of an Antagonistic Therapeutic Agent to a subject in need thereof suppresses or decreases IL-15-mediated function or signaling, which in turn results in the suppression of certain aspects of the immune function in the subject. The suppression of the immune function can be in the form of, e.g., a lower antibody response (humoral response) or a lower cellular immune response, e.g., cytokine secretion (e.g., interferon-gamma), helper activity or cellular cytotoxicity. In one embodiment, the suppression of the immune function is decreased cytokine secretion, antibody production, effector function, T cell proliferation, and/or NK cell proliferation.

#### 25 5.1.1. Protein Complexes

[0081] The present invention provides Therapeutic Agents that are complexes comprising IL-15 covalently or noncovalently bound to IL-15Ra ("IL-15/IL-15Ra complexes"). The IL-15/IL-15Ra complex is able to bind to the  $\beta\gamma$  receptor complex. In a specific embodiment, the Agonistic Therapeutic Agents are IL-15/IL-15Ra complexes that can induce IL-15-mediated signal transduction, e.g., IL-15-mediated Jak/Stat pathway signal transduction. Such induction in IL-15-mediated signaling results in enhancement of an immune function in a subject.

[0082] The IL-15/IL-15Ra complexes may be composed of native IL-15 or an IL-15 derivative and native IL-15Ra or an IL-15Ra derivative. In a specific embodiment, the IL-15/IL-15Ra complex is composed of an IL-15 derivative and an IL-15Ra derivative. In one embodiment, the IL-15Ra derivative is a soluble form of IL-15Ra. In a specific embodiment, the soluble form of IL-15Ra lacks the transmembrane domain of native IL-15Ra, and optionally, the intracellular domain of native IL-15Ra. In another embodiment, the IL-15Ra derivative is the extracellular domain of native IL-15Ra or a fragment thereof. In certain embodiments, the IL-15Ra derivative is a fragment of the extracellular domain comprising the sushi domain or exon 2 of native IL-15Ra. In some embodiments, the IL-15Ra derivative comprises a fragment of the extracellular domain comprising the sushi domain or exon 2 of native IL-15Ra and at least one amino acid that is encoded by exon 3. In certain embodiments, the IL-15Ra derivative comprises a fragment of the extracellular domain comprising the sushi domain or exon 2 of native IL-15Ra and an IL-15Ra hinge region or a fragment thereof. In a specific embodiment, the IL-15Ra derivative is encoded by a nucleic acid sequence optimized to enhance expression of IL-15Ra, e.g., using methods as described in U.S. Provisional Application No. 60/812,566, filed on June 9, 2006; and U.S. Patent Nos. 5,965,726; 6,174,666; 6,291,664; 6,414,132; and 6,794,498, which are incorporated by reference herein in their entireties. In another embodiment, the IL-15 derivative is encoded by a nucleic acid sequence optimized to enhance expression of IL-15, e.g., using methods as described in U.S. Provisional Application Nos. 60/812,566, filed on June 9, 2006 and 60/758,819, filed on January 13, 2006, and International Patent Application Publication No. WO 2007/084342; and U.S. Patent Nos. 5,965,726; 6,174,666; 6,291,664; 6,414,132; and 6,794,498, which are incorporated by reference herein in their entireties.

[0083] In another embodiment, the IL-15Ra derivative comprises a mutation in the extracellular domain cleavage site that inhibits cleavage by an endogenous protease that cleaves native IL-15Ra. In a specific embodiment, the extracellular domain cleavage site of IL-15Ra is replaced with a cleavage site that is recognized and cleaved by a heterologous known protease. Non-limiting examples of such heterologous protease cleavage sites include Arg-X-X-Arg (SEQ ID NO: 7), which is recognized and cleaved by furin protease; and A-B-Pro-Arg-X-Y (SEQ ID NO: 8) (A and B are hydrophobic amino acids and X and Y are nonacidic amino acids) and Gly-Arg-Gly, which are recognized and cleaved by thrombin protease.

[0084] In addition to IL-15 and IL-15Ra, the IL-15/IL-15Ra complexes may comprise a heterologous molecule. In some embodiments, the heterologous molecule is an antigen associated with a disease that one intends to prevent, treat

and/or manage (e.g., a viral antigen, bacterial antigen, parasitic antigen, or cancer antigen). Non-limiting examples of such antigens include antigens of the flavivirus, West Nile Virus (WNV) including structural proteins, e.g., C, M, and E, and non-structural proteins, e.g., NS1, NS2A, NS2B, NS3, NS4A, NS4B and NS5; human immunodeficiency virus (HIV) antigens gp41, gp120, gp160, Nef, Gag, and Rev, Tat, Vif, Vpu, Vpr, or vpx; influenza virus hemagglutinin; human respiratory syncytial virus G glycoprotein; core protein, matrix protein or other protein of Dengue virus; measles virus hemagglutinin; herpes simplex virus type 2 glycoprotein gB; poliovirus I VP1 (Emini et al., 1983, *Nature* 304:699); an envelope glycoprotein of HIV I; hepatitis B surface antigen; diphtheria toxin; streptococcus 24M epitope; gonococcal pilin; pseudorabies virus g50 (gpD); pseudorabies virus II (gpB); pseudorabies virus gIII (gpC); pseudorabies virus glycoprotein H; pseudorabies virus glycoprotein E; transmissible gastroenteritis glycoprotein 195; transmissible gastroenteritis matrix protein; swine rotavirus glycoprotein 38; swine parvovirus capsid protein; Serpulina hydodysenteriae protective antigen; bovine viral diarrhea glycoprotein 55; Newcastle disease virus hemagglutinin-neuraminidase; swine flu hemagglutinin; swine flu neuraminidase; antigens of foot and mouth disease virus; antigens of hog cholera virus; antigens of swine influenza virus; antigens of African swine fever virus; *Mycoplasma hyopneumoniae*; antigens of infectious bovine rhinotracheitis virus (e.g., infectious bovine rhinotracheitis virus glycoprotein E or glycoprotein G); antigens of infectious laryngotracheitis virus (e.g., infectious laryngotracheitis virus glycoprotein G or glycoprotein I); a glycoprotein of La Crosse virus; antigens of neonatal calf diarrhea virus; Venezuelan equine encephalomyelitis virus; punta toro virus; murine leukemia virus; mouse mammary tumor virus; hepatitis B virus core protein and/or hepatitis B virus surface antigen or a fragment or derivative thereof (see, e.g., U.K. Patent Publication No. GB 2034323A published June 4, 1980; Ganem and Varmus, 1987, *Ann. Rev. Biochem.* 56:651-693; Tiollais et al., 1985, *Nature* 317:489-495); antigen of equine influenza virus or equine herpesvirus (e.g., equine influenza virus type A/Alaska 91 neuraminidase, equine influenza virus type A/Miami 63 neuraminidase; equine influenza virus type A/Kentucky 81 neuraminidase; equine herpes virus type 1 glycoprotein B; equine herpes virus type 1 glycoprotein D); antigen of bovine respiratory syncytial virus or bovine parainfluenza virus (e.g., bovine respiratory syncytial virus attachment protein (BRSV G); bovine respiratory syncytial virus fusion protein (BRSV F); bovine respiratory syncytial virus nucleocapsid protein (BRSV N); bovine parainfluenza virus type 3 fusion protein; the bovine parainfluenza virus type 3 hemagglutinin neuraminidase); bovine viral diarrhea virus glycoprotein 48 or glycoprotein 53.

**[0085]** Other non-limiting examples of antigens include KS 1/4 pan-carcinoma antigen, ovarian carcinoma antigen (CA125), prostatic acid phosphate, prostate specific antigen, melanoma-associated antigen p97, melanoma antigen gp75, high molecular weight melanoma antigen (HMW-MAA), prostate specific membrane antigen, carcinoembryonic antigen (CEA), polymorphic epithelial mucin antigen, human milk fat globule antigen, Colorectal tumor-associated antigens such as: CEA, TAG-72, C017-1A; GICA 19-9, CTA-1 and LEA, Burkitt's lymphoma antigen-38.13, CD19, human B-lymphoma antigen-CD20, CD33, melanoma specific antigens such as ganglioside GD2, ganglioside GD3, ganglioside GM2, ganglioside GM3, tumor-specific transplantation type of cell-surface antigen (TSTA) such as virally-induced tumor antigens including T-antigen DNA tumor viruses and envelope antigens of RNA tumor viruses, oncofetal antigen-alpha-fetoprotein such as CEA of colon, bladder tumor oncofetal antigen, differentiation antigen such as human lung carcinoma antigen L6, L20, antigens of fibrosarcoma, human leukemia T cell antigen-Gp37, neoglycoprotein, sphingolipids, breast cancer antigen such as EGFR (Epidermal growth factor receptor), HER2 antigen (p185HER2), EphA2 receptor, polymorphic epithelial mucin (PEM), malignant human lymphocyte antigen-APO-1, differentiation antigen such as I antigen found in fetal erythrocytes and primary endoderm, I(Ma) found in gastric adenocarcinomas, M18 and M39 found in breast epithelium, SSEA-1 found in myeloid cells, VEP8, VEP9, Myl, VIM-D5, and D156-22 found in colorectal cancer, TRA-1-85 (blood group H), C 14 found in colonic adenocarcinoma, F3 found in lung adenocarcinoma, AH6 found in gastric cancer, Y hapten, Ley found in embryonal carcinoma cells, TL5 (blood group A), EGF receptor, E1 series (blood group B) found in pancreatic cancer, FC 10.2 found in embryonal carcinoma cells, gastric adenocarcinoma, CO-514 (blood group Lea) found in adenocarcinoma, NS-10 found in adenocarcinomas, CO-43 (blood group Leb), G49, 19.9 found in colon cancer, gastric cancer mucins, TSA7 found in myeloid cells, R24 found in melanoma, 4.2, GD3, D1.1, OFA-1, GM2, OFA-2, GD2, M1:22:25:8 found in embryonal carcinoma cells and SSEA-3, SSEA-4 found in 4-8-cell stage embryos.

**[0086]** In other embodiments, the heterologous molecule is an antibody that specifically binds to an antigen associated with a disease that one intends to prevent, treat and/or manage (e.g., an antibody that specifically binds to a viral antigen, bacterial antigen, parasitic antigen, or cancer antigen). Non-limiting examples of such antibodies include anti-CD34 antibody, anti-CD56 antibody, anti-CD8 antibody, anti-CD22 antibody, anti-CD20 antibody, anti-CD19 antibody, anti-CD3 antibody, anti-EGFR antibody, anti-HER2 antibody, anti-CD34 antibody, anti-ckit antibody, anti-flt3 antibody, anti-hemagglutinin antibody, anti-gp41 antibody, anti-gp120 antibody, and anti-HSV-II glycoprotein gB antibody. In other embodiments, the antibody immunospecifically binds to one of the antigens listed above. In some embodiments, the antibody specifically binds to a cellular antigen (e.g., a receptor or cell surface antigen) expressed by a cell that one desires to target. For example, the IL-15/IL-15Ra complex can be targeted to CD34+ progenitor cells with an anti-CD34 antibody to induce development of such cells into CD56+ NK cells. The IL-15/IL-15Ra complex can be targeted to CD56+ NK cells with an anti-CD56 antibody to induce proliferation of such cells.

**[0087]** In some embodiments, the heterologous molecule increases protein stability. Non-limiting examples of such

molecules include polyethylene glycol (PEG), Fc domain of an IgG immunoglobulin or a fragment thereof, or albumin that increase the half-life of IL-15 or IL-15Ra *in vivo*. In certain embodiments, the heterologous molecules is not an Fc domain of an immunoglobulin molecule or a fragment thereof.

[0088] In those IL-15/IL-15Ra complexes comprising a heterologous molecule, the heterologous molecule may be conjugated to IL-15 and/or IL-15Ra. In one embodiment, the heterologous molecule is conjugated to IL-15Ra. In another embodiment, the heterologous molecule is conjugated to IL-15.

[0089] The components of an IL-15/IL-15Ra complex may be directly fused, using either non-covalent bonds or covalent bonds (e.g., by combining amino acid sequences via peptide bonds), and/or may be combined using one or more linkers. In a specific embodiment, IL-15 and IL-15Ra are directly fused to each other using either non-covalent bonds or covalent bonds (e.g., by combining amino acid sequences via peptide bonds), and/or may be combined using one or more linkers. In specific embodiments, a polypeptide comprising IL-15 and IL-15Ra directly fused to each other using either non-covalent bonds or covalent bonds is functional (e.g., capable of specifically binding to the IL-15R  $\beta\gamma$  complex and inducing IL-15-mediated signal transduction and/or IL-15-mediated immune function). Linkers suitable for preparing the IL-15/IL-15Ra complexes comprise peptides, alkyl groups, chemically substituted alkyl groups, polymers, or any other covalently-bonded or non-covalently bonded chemical substance capable of binding together two or more components. Polymer linkers comprise any polymers known in the art, including polyethylene glycol ("PEG"). In some embodiments, the linker is a peptide that is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more amino acids long. In a specific embodiment, the linker is long enough to preserve the ability of IL-15 to bind to the IL-15Ra. In other embodiments, the linker is long enough to preserve the ability of the IL-15/IL-15Ra complex to bind to the  $\beta\gamma$  receptor complex and to act as an agonist to mediate IL-15 signal transduction.

[0090] The present invention relates to Therapeutic Agents comprising IL-15/IL-15Ra complexes for use in the methods described herein. In particular embodiments, IL-15/IL-15Ra complexes are pre-coupled prior to use in the methods described herein (e.g., prior to contacting cells with the IL-15/IL-15Ra complexes or prior to administering the IL-15/IL-15Ra complexes to a subject). In other embodiments, the IL-15/IL-15Ra complexes are not pre-coupled prior to use in the methods described herein. In specific embodiments, the IL-15/IL-15Ra complex is administered in combination with a vaccine composition to enhance the immune response elicited by the administration of the vaccine composition to a subject. In a specific embodiment, an Agonistic Therapeutic Agent comprising IL-15 and IL-15Ra directly fused to each other is administered in combination with a vaccine composition to enhance an immune response elicited by administration of the vaccine composition to a subject.

### 5.1.2. Nucleic Acids

[0091] The invention provides Therapeutic Agents that are nucleic acids that encode IL-15 and IL-15Ra. The nucleic acids encode IL-15 and IL-15Ra that are capable of covalently or noncovalently binding to each other to form the IL-15/IL-15Ra complexes described in Section 5.1.1, *supra*. Such IL-15/IL-15Ra complexes can bind to the  $\beta\gamma$  receptor complex, and induce IL-15-mediated signal transduction.

[0092] Nucleic acid sequences encoding native IL-15 are well known in the art and have been described, for a review, see, Fehniger and Caligiuri, Blood, 2001, 97:14-32, which is incorporated by reference herein in its entirety. For example, the nucleic acid sequences encoding native IL-15 can be readily found in publicly available publications and databases, e.g., National Center for Biotechnology Information website at ncbi.nlm.nih.gov. Nucleic acid sequences encoding native IL-15Ra have been described, e.g., see International Publication No. WO 95/30695, and can also be readily found in publicly available publications and databases, e.g., National Center for Biotechnology Information website at ncbi.nlm.nih.gov. Cloning techniques well known in the art can be used to generate nucleic acids encoding IL-15 and IL-15Ra. See, e.g., Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc. (1995); Sambrook et al., *Molecular Cloning, A Laboratory Manual* (2d ed.), Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1989); Birren et al., *Genome Analysis: A Laboratory Manual*, volumes 1 through 4, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1997-1999).

[0093] In one embodiment, the Therapeutic Agent comprises nucleic acids that encode native IL-15Ra. In another embodiment, the Therapeutic Agent comprises nucleic acids that encode an IL-15Ra derivative that is a soluble form of native IL-15Ra. In a specific embodiment, the Therapeutic Agent comprises nucleic acids that encode an IL-15Ra derivative that is a soluble form of IL-15Ra that lacks the transmembrane domain of native IL-15Ra, and optionally the intracellular domain of native IL-15Ra. In other embodiments, the Therapeutic Agents comprise nucleic acids that encode an IL-15Ra derivative that is the extracellular domain of native IL-15Ra or a fragment thereof. In certain embodiments, the Therapeutic Agents comprise nucleic acids that encode an IL-15Ra derivative that is a fragment of the extracellular domain comprising the sushi domain or exon 2 of native IL-15Ra. In some embodiments, Therapeutic Agents comprise nucleic acids that encode an IL-15Ra derivative comprising a fragment of the extracellular domain comprising the sushi domain or exon 2 of native IL-15Ra and at least one amino acid that is encoded by exon 3. In certain embodiments, the Therapeutic Agents comprise nucleic acids that encode an IL-15Ra derivative comprising a fragment of the extracellular

domain comprising the sushi domain or exon 2 of native IL-15Ra and an IL-15Ra hinge region or a fragment thereof. In some embodiments, the Therapeutic Agents comprise nucleic acids that encode an IL-15Ra derivative that consists essentially of the sushi domain or exon 2 of the receptor. In a specific embodiment, Therapeutic Agents comprise nucleic acids comprising sequences encoding a chimeric polypeptide comprising IL-15 and IL-15Ra directly fused to each other.

5 In another embodiment, the Therapeutic Agents comprise nucleic acids that encode an IL-15Ra derivative comprising a mutation in the extracellular domain cleavage site that inhibits cleavage by an endogenous protease that cleaves IL-15Ra. In a specific embodiment, Therapeutic Agents comprise nucleic acids that encode an IL-15Ra derivative comprising mutations (in the extracellular domain cleavage site) that inhibit cleavage by an endogenous protease. In a specific embodiment, Therapeutic Agents comprise nucleic acids comprising sequences that encode an IL-15Ra derivative, 10 wherein an extracellular domain cleavage site of IL-15Ra is replaced with a heterologous domain (e.g., heterologous transmembrane domain) or a synthetic amino acid sequence that does not allow cleavage and generation of soluble IL-15Ra. In certain embodiments, the extracellular domain cleavage site of IL-15Ra that is cleaved by an endogenous processing enzyme is mutated to inhibit cleavage and generation of soluble IL-15Ra. In one embodiment, the extracellular domain cleavage site of IL-15Ra is replaced with a heterologous extracellular domain cleavage site (e.g., heterologous 15 transmembrane domain that is recognized and cleaved by another enzyme unrelated to the endogenous processing enzyme that cleaves IL-15Ra).

20 [0094] In a specific embodiment, the Therapeutic Agents comprise nucleic acids that encode IL-15 and/or IL-15Ra that are optimized, e.g., by codon/RNA optimization, replacement with heterologous signal sequences, and elimination of mRNA instability elements. Methods to generate optimized nucleic acids encoding IL-15 and IL-15Ra for expression by introducing codon changes and/or eliminating inhibitory regions in the mRNA can be carried out by adapting the optimization methods described in, e.g., U.S. Patent Nos. 5,965,726; 6,174,666; 6,291,664; 6,414,132; and 6,794,498, for IL-15 and IL-15Ra. The contents of each of these references are incorporated by reference herein in its entirety. See also U.S. Provisional Application Nos. 60/812,566, filed on June 9, 2006, and 60/758,819, filed on January 13, 2007, and International Patent Application Publication No. WO 2007/084342, which are also incorporated by reference herein in their entireties. For example, potential splice sites and instability elements (e.g., A/T or A/U rich elements) within the RNA of IL-15 and IL-15Ra can be mutated without altering the amino acids encoded by the nucleic acid sequences to increase stability of the RNA for expression. The alterations utilize the degeneracy of the genetic code, e.g., using an alternative codon for an identical amino acid. In some embodiments, it may be desirable to alter one or more codons to encode a conservative mutation, e.g., a similar amino acid with similar chemical structure and properties and/or function 30 as the original amino acid. Such methods can increase expression of IL-15 and/or IL-15Ra proteins by at least 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 10 fold, 20 fold, 30 fold, 40 fold, 50 fold, 60 fold, 70 fold, 80 fold, 90 fold, or 100 fold or more relative to the expression of IL-15 and/or IL-15Ra proteins encoded by native nucleic acid sequences.

35 [0095] Further, the native signal peptide sequence of IL-15 and/or IL-15Ra can be replaced with a heterologous signal peptide, e.g., a signal peptide of human GM-CSF (see FIGs. 8A-D), tissue plasminogen activator (tPA) (see FIGs. 5A-D), prolactin, growth hormone or an immunoglobulin protein (e.g., IgE). In a specific embodiment, the signal peptide of IL-15 is replaced with the signal sequence of tPA. In other specific embodiments, the signal peptide of IL-15 is replaced with the signal peptide of human GM-CSF. Such alterations can increase expression of IL-15 and/or IL-15Ra proteins/polypeptides by at least 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 10 fold, 20 fold, 30 fold, 40 fold, 50 fold, 60 fold, 70 fold, 80 fold; 90 fold, or 100 fold or more relative to the expression of IL-15 and/or IL-15Ra proteins with the respective native signal peptide, as measured/detected by a technique known to one of skill in the art, e.g., ELISA.

40 [0096] In some embodiments, an optimized nucleotide sequence encoding IL-15 or IL-15Ra hybridizes to the nucleotide sequence encoding native IL-15 or IL-15Ra, respectively. In specific embodiments, an optimized nucleotide sequence encoding IL-15 or IL-15Ra hybridizes under high stringency conditions to a nucleotide sequence encoding native IL-15 or IL-15Ra, respectively, or a fragment thereof. In a specific embodiment, an optimized nucleotide sequence encoding IL-15 or IL-15Ra hybridizes under high stringency, intermediate or lower stringency hybridization conditions to a nucleotide sequence encoding native IL-15 or IL-15Ra, respectively, or a fragment thereof. Information regarding hybridization conditions have been described, see, e.g., U.S. Patent Application Publication No. US 2005/0048549 (e.g., paragraphs 72-73).

45 [0097] The present invention provides nucleic acids encoding IL-15, IL-15Ra, and a heterologous molecule in a form that allows IL-15 to covalently or noncovalently bind to the IL-15Ra to form IL-15/IL-15Ra complexes. In some embodiments, the heterologous molecule is an antigen associated with a disease that one intends to prevent, treat and/or manage. Non-limiting examples of such antigens include those listed above in Section 5.1.1. In other embodiments, the heterologous molecule is an antibody that specifically binds to an antigen associated with a disease that one intends to prevent, treat and/or manage. Non-limiting examples of such antibodies include those listed above in Section 5.1.1 and 50 those known in the art. In some embodiments, the antibody specifically binds to a cellular surface antigen (e.g., a receptor) expressed by a cell that one desires to target. In some embodiments, the heterologous molecule increases protein stability. Non-limiting examples of such molecules include polyethylene glycol (PEG), Fc domain of an IgG immunoglobulin or a fragment thereof, or albumin that increase the half-life of IL-15 or IL-15Ra *in vivo*. In certain embodiments, the

heterologous molecules is not an Fc domain of an immunoglobulin molecule or a fragment thereof.

**[0098]** In those IL-15/IL-15Ra complexes comprising a heterologous molecule, the heterologous molecule may be conjugated to IL-15 and/or IL-15Ra. In one embodiment, the heterologous molecule is conjugated to IL-15Ra. In another embodiment, the heterologous molecule is conjugated to IL-15.

**[0099]** In specific embodiments, IL-15 and IL-15Ra are encoded by one nucleic acid construct (e.g., bicistronic construct). In some embodiments, IL-15 and IL-15Ra are encoded by one nucleic acid construct comprising a single open reading frame (ORF) of IL-15 and IL-15Ra. In some embodiments, IL-15 or IL-15Ra encoded by a nucleic acid construct may be conjugated to a nucleic acid encoding a heterologous molecule, such as an antigen or an antibody of interest. In other embodiments, IL-15 and IL-15Ra are encoded by two nucleic acid constructs, wherein a first nucleic acid construct encodes IL-15 and a second nucleic acid construct encodes IL-15Ra. The IL-15 encoded by the first nucleic acid construct may be conjugated to a nucleic acid encoding a heterologous molecule, such as an antigen or an antibody of interest. Alternatively, or in addition, the IL-15Ra encoded by the second nucleic acid construct may be conjugated to a nucleic acid encoding a heterologous molecule, such as an antigen or an antibody of interest.

#### 15 5.1.2.1. Constructs

**[0100]** The nucleic acids encoding IL-15 and/or IL-15Ra can be inserted into nucleic acid constructs for expression in mammalian cells, bacteria, yeast, and viruses. IL-15 and IL-15Ra can be recombinantly expressed from the same nucleic acid construct (e.g., using a bicistronic nucleic acid construct) or from different nucleic acid constructs (e.g., using monocistronic nucleic acid constructs). In one embodiment, IL-15 and IL-15Ra can be recombinantly expressed from a single nucleic acid construct comprising a single open reading frame (ORF) of IL-15 and IL-15Ra.

**[0101]** The nucleic acid constructs may comprise one or more transcriptional regulatory element(s) operably linked to the coding sequence of IL-15 and/or IL-15Ra. The transcriptional regulatory elements are typically 5' to the coding sequence and direct the transcription of the nucleic acids encoding IL-15 and/or IL-15Ra. In some embodiments, one or more of the transcriptional regulatory elements that are found in nature to regulate the transcription of the native IL-15 and/or native IL-15Ra gene are used to control transcription. In other embodiments, one or more transcriptional regulatory elements that are heterologous to the native IL-15 and/or native IL-15Ra gene are used to control transcription. Any transcriptional regulatory element(s) known to one of skill in the art may be used. Non-limiting examples of the types of transcriptional regulatory element(s) include a constitutive promoter, a tissue-specific promoter, and an inducible promoter. In a specific embodiment, transcription is controlled, at least in part, by a mammalian (in some embodiments, human) transcriptional regulatory element(s). In a specific embodiment, transcription is controlled, at least in part, by a strong promoter, e.g., CMV.

**[0102]** Specific examples of promoters which may be used to control transcription include, but are not limited to, the SV40 early promoter region (Benoist & Chambon, 1981, *Nature* 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, *Cell* 22:787-7797), the herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, *Nature* 296:39-42); adenovirus (ADV), cytomegalovirus (CMV), bovine papilloma virus (BPV), parovirus B19p6 promoter, prokaryotic expression vectors such as the .beta.-lactamase promoter (Villa-Kamaroff et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.* 75:3727-3731), or the tac promoter (DeBoer et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:21-25); see also "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242:74-94; plant expression vectors comprising the nopaline synthetase promoter region (Herrera-Estrella et al., *Nature* 303:209-213) or the cauliflower mosaic virus 35S RNA promoter (Gardner, et al., 1981, *Nucl. Acids Res.* 9:2871), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., 1984, *Nature* 310:115-120); promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45:485-495), albumin gene control region which is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.* 5:1639-1648; Hammer et al., 1987, *Science* 235:53-58; alpha 1-antitrypsin gene control region which is active in the liver (Kelsey et al., 1987, *Genes and Devel.* 1:161-171), beta-globin gene control region which is active in myeloid cells (Mogram et al., 1985, *Nature* 315:338-340; Kollias et al., 1986, *Cell* 46:89-94; myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, *Cell* 48:703-712); myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, *Nature* 314:283-286), and gonadotropic

releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, *Science* 234:1372-1378). In other aspects, an inducible promoter can be used.

**[0103]** The nucleic acid constructs also may comprise one or more post-transcriptional regulatory element(s) operably linked to the coding sequence of IL-15 and/or IL-15Ra. The post-transcriptional regulatory elements can be 5' and/or 3' to the coding sequence and direct the post-transcriptional regulation of the translation of RNA transcripts encoding IL-15 and/or IL-15Ra.

**[0104]** In another aspect, the nucleic acid construct can be a gene targeting vector that replaces a gene's existing regulatory region with a regulatory sequence isolated from a different gene or a novel regulatory sequence as described, e.g., in International Publication Nos. WO 94/12650 and WO 01/68882, which are incorporated by reference herein in their entireties.

**[0105]** The nucleic acid construct chosen will depend upon a variety of factors, including, without limitation, the strength of the transcriptional regulatory elements and the host cell to be used to express IL-15 and/or IL-15Ra. The nucleic acid constructs can be a plasmid, phagemid, cosmid, viral vector, phage, artificial chromosome, and the like. In one aspect, the vectors can be episomal or non-/homologously integrating vectors, which can be introduced into the appropriate host cells by any suitable means (transformation, transfection, conjugation, protoplast fusion, electroporation, calcium phosphate-precipitation, direct microinjection, etc.) to transform them.

**[0106]** The nucleic acid constructs can be a plasmid or a stable integration vector for transient or stable expression of IL-15 and/or IL-15Ra in host cells. For stable expression, the vector can mediate chromosomal integration at a target site or a random chromosomal site. Non-limiting examples of host cell-vector systems that may be used to express IL-15 and/or IL-15Ra include mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, retroviruses, lentiviruses, etc.); insect cell systems infected with virus (e.g., baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, DNA, plasmid DNA, or cosmid DNA; and stable cell lines generated by transformation using a selectable marker. In some embodiments, the nucleic acid constructs include a selectable marker gene including, but not limited to, neo, gpt, dhfr, ada, pac, hyg, CAD and hisD.

**[0107]** The nucleic acid constructs can be monocistronic or multicistronic. A multicistronic nucleic acid construct may encode 2, 3, 4, 5, 6, 7, 8, 9, 10 or more, or in the range of 2-5, 5-10 or 10-20 genes/nucleotide sequences. For example, a bicistronic nucleic acid construct may comprise in the following order a promoter, a first gene (e.g., IL-15), and a second gene and (e.g., IL-15Ra). In such a nucleic acid construct, the transcription of both genes is driven by the promoter, whereas the translation of the mRNA from the first gene is by a cap-dependent scanning mechanism and the translation of the mRNA from the second gene is by a cap-independent mechanism, e.g., by an IRES.

**[0108]** Techniques for practicing aspects of this invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, and recombinant DNA manipulation and production, which are routinely practiced by one of skill in the art. See, e.g., Sambrook, 1989, *Molecular Cloning, A Laboratory Manual*, Second Edition; *DNA Cloning, Volumes I and II* (Glover, Ed. 1985); *Oligonucleotide Synthesis* (Gait, Ed. 1984); *Nucleic Acid Hybridization* (Hames & Higgins, Eds. 1984); *Transcription and Translation* (Hames & Higgins, Eds. 1984); *Animal Cell Culture* (Freshney, Ed. 1986); *Immobilized Cells and Enzymes* (IRL Press, 1986); *Perbal, A Practical Guide to Molecular Cloning* (1984); *Gene Transfer Vectors for Mammalian Cells* (Miller & Calos, Eds. 1987, Cold Spring Harbor Laboratory); *Methods in Enzymology, Volumes 154 and 155* (Wu & Grossman, and Wu, Eds., respectively), (Mayer & Walker, Eds., 1987); *Immunochemical Methods in Cell and Molecular Biology* (Academic Press, London, Scopes, 1987), *Expression of Proteins in Mammalian Cells Using Vaccinia Viral Vectors in Current Protocols in Molecular Biology, Volume 2* (Ausubel et al., Eds., 1991).

**[0109]** The nucleic acid construct comprising nucleic acids encoding IL-15 and/or IL-15Ra can be *administered in vivo* to a mammal or transfected into primary or immortalized cells in culture. In certain aspects, the nucleic acid constructs comprising nucleic acids encoding IL-15 and/or IL-15Ra are administered to a mammal for recombinant expression of IL-15 and IL-15Ra *in vivo* to enhance IL-15 mediated signal transduction and to enhance an immune function associated with IL-15 signaling *in vivo*. In other aspects, the nucleic acids encoding IL-15 and/or IL-15Ra are administered in combination with a vaccine composition to enhance the immune response elicited by the administration of the vaccine composition to a subject.

**[0110]** The nucleic acid constructs comprising nucleic acids encoding IL-15 and/or IL-15Ra can be used to generate cells that express IL-15 and IL-15Ra for enhancement of an immune function in a subject. In particular, such cells can transpresent the IL-15/IL-15Ra complex on the cell surface to adjacent cells that express the  $\beta\gamma$  receptor complex, thus inducing IL-15 signal transduction. In some embodiments, the cells are primary cells (e.g., tumor cells isolated from a patient). In other embodiments, the cells are mammalian cell lines.

**[0111]** In another aspect, the nucleic acids encoding IL-15 and/or IL-15Ra can be used to generate mammalian cells that recombinantly express IL-15 and IL-15Ra in high amounts for the isolation and purification of IL-15 and IL-15Ra, preferably IL-15 and the IL-15Ra are associated as complexes. In one embodiment, high amounts of IL-15/IL-15Ra complexes refer to amounts of IL-15/IL-15Ra complexes expressed by cells that are at least 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, or more than 20 fold higher than amounts of IL-15/IL-15Ra complexes

expressed endogenously by control cells (e.g., cells that have not been genetically engineered to recombinantly express IL-15, IL-15Ra, or both IL-15 and IL-15Ra, or cells comprising an empty vector). In a specific embodiment, the IL-15Ra is the soluble form of IL-15Ra. In a specific embodiment, the IL-15Ra is the soluble form of IL-15Ra associated with IL-15 in a stable heterodimer, which increases yields and simplifies production and purification of bioactive heterodimer IL-15/soluble IL-15Ra cytokine. Recombinant IL-15 and IL-15Ra can be purified using methods of recombinant protein production and purification are well known in the art, e.g., see International Publication No. WO 07/070488, which is incorporated by reference herein in its entirety. Briefly, the polypeptide can be produced intracellularly, in the periplasmic space, or directly secreted into the medium. Cell lysate or supernatant comprising the polypeptide can be purified using, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™ (gel filtration substance; Pharmacia Inc., Piscataway, New Jersey) chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available.

**[0112]** In some embodiments, IL-15 and IL-15Ra are synthesized or recombinantly expressed by different cells and subsequently isolated and combined to form an IL-15/IL-15Ra complex, *in vitro*, prior to administration to a subject. In other embodiments, IL-15 and IL-15Ra are synthesized or recombinantly expressed by different cells and subsequently isolated and simultaneously administered to a subject an IL-15/IL-15Ra complex *in situ* or *in vivo*. In yet other embodiments, IL-15 and IL-15Ra are synthesized or expressed together by the same cell, and the IL-15/IL-15Ra complex formed is isolated.

### 20 5.1.3. Antibodies

**[0113]** The present invention provides antibodies that specifically bind to IL-15/IL-15Ra complexes. In a specific embodiment, the present invention provides antibodies that specifically bind to the complex formed when IL-15 and IL-15Ra bind to each other. In a more specific embodiment, the present invention provide antibodies that specifically bind to the complex formed between IL-15 and IL-15Ra and prevent or reduce IL-15-mediated signal transduction. Such antibodies, in accordance with this embodiment, may block (sterically or non-sterically) the binding of the complex of IL-15 and IL-15Ra with the  $\beta\gamma$  (or CD122/CD132) receptor complex of the IL-15 receptor. In a specific embodiment, the antibody reduces the binding of the IL-15/IL-15Ra complex to the  $\beta\gamma$  complex by at least 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 11 fold, 12 fold, 13 fold, 14 fold, 15 fold, 16 fold, 17 fold, 18 fold, 19 fold, 20 fold, 30 fold, 40 fold, 50 fold, 60 fold, 70 fold, 80 fold, 90 fold, 100 fold, 500 fold, or 1000 fold relative to a negative control (e.g., binding affinity of the IL-15/IL-15Ra complex to the  $\beta\gamma$  receptor complex in the absence of the antibody) as determined by methods well known in the art, e.g., ELISA, cell-based assays including flow cytometry, KinEx A assay, and plasmon surface resonance assay (e.g., BiAcore assay).

**[0114]** Antibodies that specifically bind to the IL-15/IL-15Ra complex can be produced by any method well known in the art, e.g., as described in U.S. Patent Nos. 5,807,715, 6,331,415, and 6,818,216; U.S. Patent Application Publication Nos. US 2002/0098189, US 2004/0028685, US 2005/0019330, and US 2007/0086943; International Publication No. WO 02/46237; and Harlow et al., *Antibodies. A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references are incorporated by reference herein in their entireties).

**[0115]** The antibodies that specifically bind to IL-15/IL-15Ra complexes can be administered to a subject to prevent IL-15/IL-15Ra complexes from binding to the  $\beta\gamma$  receptor complex and from inducing IL-15-mediated signal transduction. Thus, such antibodies can suppress an immune function that is associated with IL-15 signal transduction. In some embodiments, antibodies described herein are useful for detecting the presence of IL-15/IL-15Ra complexes.

### 45 5.1.4. Cells

**[0116]** Cells can be engineered to express the protein(s) encoded by the nucleic acid constructs described in Section 5.1.2, *supra*, in high amounts. In one embodiment, high amounts of IL-15/IL-15Ra complexes refer to amounts of IL-15/IL-15Ra complexes expressed by cells that are at least 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, or more than 20 fold higher than amounts of IL-15/IL-15Ra complexes expressed endogenously by control cells (e.g., cells that have not been genetically engineered to recombinantly express IL-15, IL-15Ra, or both IL-15 and IL-15Ra, or cells comprising an empty vector). In addition, cells can be engineered to express the antibodies described in Section 5.1.3, *supra*, using techniques well-known to one of skill in the art, see, e.g., U.S. Patent Nos. 5,807,715, 6,331,415, and 6,818,216; U.S. Patent Application Publication Nos. US 2002/0098189, US 2004/0028685, US 2005/0019330, and US 2007/0086943; International Publication No. WO 02/46237; and Harlow et al., *Antibodies. A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references are incorporated by reference herein in their

entireties). The host cells chosen for expression of nucleic acids will depend upon the intended use of the cells. Factors such as whether a cell glycosylates similar to cells that endogenously express, e.g., IL-15 and/or IL-15Ra, may be considered in selecting the host cells.

**[0117]** In one embodiment, the invention also provides for a method of increasing the yield and bioactivity of IL-15 by 5 constructing cell lines expressing both IL-15 and soluble IL-15Ra, and purifying the stable heterodimer, which can be used *in vitro* or *in vivo*, e.g., can be administered to a human. In one embodiment, the stability of IL-15 is increased when produced from cell lines recombinantly expressing both IL-15 and IL-15Ra.

**[0118]** Non-limiting examples of hosts cells that can be used to express the protein(s) encoded by the nucleic acid constructs described in Section 5.1.2, *supra*, or the antibodies described in Section 5.1.3, *supra*, include mammalian 10 cells, bacterial cells, yeast cells, primary cells, immortalized cells, and insect cells. In a specific embodiment, the host cells are a mammalian cell line. Examples of mammalian cell lines include, but are not limited to, COS, CHO, HeLa, NIH3T3, HepG2, MCF7, HEK, 293T, RD, PC12, hybridomas, pre-B cells, 293, 293H, K562, SkBr3, BT474, A204, M07Sb, TF $\beta$ 1, Raji, Jurkat, MOLT-4, CTLL-2, MC-IXC, SK-N-MC, SK-N-MC, SK-N-DZ, SH-SY5Y, C127, N0, and BE(2)-C cells. Other mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell 15 lines available from the American Type Culture Collection (ATCC). In another embodiment, the host cells are immortalized cell lines derived from a subject. In another embodiment, the host cells are primary or secondary cells from a subject. In a particular embodiment, the host cells are cancer cells. In another embodiment, the host cells are epithelial cells or endothelial cells. In another embodiment, the host cells are fetal/embryonic cells. In some embodiments, the host cells are progenitor cells. In some embodiments, the host cells are lymphocytes (e.g., T cells and B cells). In another embodiment, the host cells are stem cells. In yet another embodiment, the host cells engineered to express the nucleic acid 20 constructs of Section 5.1.2, *supra*, are from an adult.

**[0119]** In some embodiments, isolated cells are utilized herein. In a specific embodiment, the isolated cells are at least 80%, 90%, 95% or 98% free of a different cell type as measured by a technique known to one of skill in the art, such as flow cytometry. In other words, at least 80%, 90%, 95% or 98% of the isolated cells are of the same cell type.

**[0120]** In a specific embodiment, the nucleic acid constructs encoding IL-15 or IL-15Ra can be co-transfected or transfected into the same host cells or different host cells. Optionally, a nucleic acid construct comprising nucleic acids encoding a selectable marker gene can also be transfected into the same cells to select for transfected cells. If the nucleic acid constructs comprising nucleic acids encoding IL-15 and IL-15Ra are transfected into different cells, IL-15 and IL-15Ra expressed by the different cells can be isolated and contacted with each other under conditions suitable 30 to form IL-15/IL-15Ra complexes described in Section 5.1.1, *supra*. Any techniques known to one of skill in the art can be used to transfet or transducer host cells with nucleic acids including, e.g., transformation, transfection, conjugation, protoplast fusion, electroporation, calcium phosphate-precipitation, direct microinjection, and infection with viruses, including but not limited to adenoviruses, lentiviruses, and retroviruses.

**[0121]** For long-term, high-yield production of a recombinant of IL-15 and IL-15Ra polypeptides, stable cell lines can 35 be generated. For example, cell lines can be transformed using the nucleic acid constructs of Section 5.1.2 which may contain a selectable marker gene on the same or on a separate nucleic acid construct. The selectable marker gene can be introduced into the same cell by co-transfection. Following the introduction of the vector, cells are allowed to grow for 1-2 days in an enriched media before they are switched to selective media to allow growth and recovery of cells that successfully express the introduced nucleic acids. Resistant clones of stably transformed cells may be proliferated using 40 tissue culture techniques well known in the art that are appropriate to the cell type. In a particular embodiment, the cell line has been adapted to grow in serum-free medium. In one embodiment, the cell line has been adapted to grow in serum-free medium in shaker flasks. In one embodiment, the cell line has been adapted to grow in stir or rotating flasks. In certain embodiments, the cell line is cultured in suspension. In particular embodiments, the cell line is not adherent or has been adapted to grow as nonadherent cells. In certain embodiments, the cell line has been adapted to grow in 45 low calcium conditions. In some embodiments, the cell line is cultured or adapted to grow in low serum medium.

**[0122]** In a specific embodiment, the host cell recombinantly expresses IL-15 and the full length IL-15Ra. In another specific embodiment, the host cell recombinantly expresses IL-15 and the soluble form of IL-15Ra. In another specific embodiment, the host cell recombinantly expresses IL-15 and a membrane-bound form of IL-15Ra which is not cleaved 50 from the surface of the cell and remains cell associated. In some embodiments, the host cell recombinantly expressing IL-15 and/or IL-15Ra (full-length or soluble form) also recombinantly expresses another polypeptide (e.g., a cytokine or fragment thereof).

**[0123]** In a specific embodiment, a particularly preferred method of high-yield production of a recombinant polypeptide of the present invention is through the use of dihydro folate reductase (DHFR) amplification in DHFR-deficient CHO cells, by the use of successively increasing levels of methotrexate as described in U.S. Patent No. 4,889,803, which is incorporated by reference herein in its entirety. The polypeptide obtained from such cells may be in a glycosylated form.

**[0124]** In a specific embodiment, the nucleic acid constructs are suitable for introduction and expression in primary cells isolated from a subject. The primary cells are engineered to express IL-15 and IL-15Ra. In a specific embodiment, the primary cells express IL-15 and the full length IL-15Ra. In a particular embodiment the IL-15Ra contains a mutation

that eliminated the extracellular domain cleavage site or that replaces the native extracellular domain cleavage site with a heterologous extracellular domain cleavage site. When administered to a subject, such cells can transpresent the IL-15/IL-15Ra complex to adjacent cells *in vivo*, thus mediating IL-15 signal transduction and enhancing an immune system function mediated by IL-15 signaling. In another specific embodiment, the primary cells express IL-15 and the soluble form of IL-15Ra.

**[0125]** In a particular embodiment, the primary cells are cancer cells: (i) isolated from a subject (diagnosed with cancer); (ii) engineered to recombinantly express either IL-15 or IL-15Ra (full-length or soluble form) or both; and (iii) irradiated prior to administration to a cancer patient for enhancing an immune response in the subject to antigens of the cancer cells. In a specific embodiment, the primary cells isolated from a subject are further engineered to recombinantly express another therapeutic polypeptide, e.g., a cytokine (e.g., IL-1, IL-2, IL-6, IL-11, IL-12, IL-13, TNF-alpha, GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , or interferon- $\gamma$ ), a growth factor or a fragment or derivative thereof. In a specific embodiment, the primary cells isolated from a subject are further engineered to recombinantly express an antigen of a cancer. In one embodiment, irradiated cells can be administered to the same subject or to a different subject from which the cells were obtained. In a particular embodiment, the primary cells are isolated from a tumor of the subject. In one embodiment, cells of a cancer cell line can be engineered to recombinantly express either IL-15 or IL-15Ra or both, wherein the cells are irradiated before administration to a subject.

**[0126]** In some embodiments, the present invention provides for methods for making cells described herein. In one embodiment, the present invention relates to a method for making irradiated cancer cells co-expressing IL-15 and IL-15Ra comprising the steps of: (i) isolating cancer cells from a subject (diagnosed with cancer); (ii) engineering said cancer cells to recombinantly express either IL-15, or IL-15Ra (full length or soluble form), or both; and (iii) irradiating said cancer cells. In some embodiment, the present invention relates to a method for making irradiated cancer cells co-expressing IL-15 and IL-15Ra comprising the steps of: (i) isolating cancer cells from a subject (diagnosed with cancer); (ii) introducing a nucleic acid construct(s) encoding recombinant IL-15 or a derivative thereof and human IL-15Ra or a derivative thereof; and (iii) irradiating said cancer cells. In particular embodiments, the irradiated cancer cells are administered to the subject from which the cancer cells were isolated (or obtained).

**[0127]** In one embodiment, the present invention relates to a method for making host cells capable of growing in serum-free medium comprising the steps of: (i) engineering host cells to recombinantly express IL-15 and/or IL-15Ra (full length or soluble form), (ii) culturing the host cells in medium comprising a 1:1 ratio of old medium (comprising 10% serum) to new medium; and (iii) repeating step (ii) until the host cells grow at a desired growth rate, wherein the new medium is serum-free, and wherein the desired growth rate is the desired growth rate is the growth rate the host cells grows when cultured in medium comprising 10% serum.

**[0128]** In some embodiments, the present invention encompasses a cell that recombinantly expresses a mammalian IL-15 or a derivative thereof and a mammalian IL-15Ra or a derivative thereof, wherein the cell expresses at least 0.6 pg of mammalian IL-15 or a derivative thereof. In certain embodiments, the cell expresses at least 0.1 pg, 0.5 pg, 1 pg, or 2 pg of mammalian IL-15 or a derivative thereof. In some embodiments, the cell expresses approximately 0.1 pg to 0.6 pg, 0.5 pg to 1 pg, 0.5 pg to 2 pg, or 0.1 pg to 2 pg of mammalian IL-15 or a derivative thereof. In particular embodiments, the cell recombinantly expresses a mammalian IL-15 or a derivative thereof that is more stable than endogenous IL-15 produced by a cell not recombinantly expressing both mammalian IL-15 and IL-15Ra. In specific embodiments, the protein stability of recombinant mammalian IL-15 produced by such a cell is at least 1 fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, 50 fold, or 100 fold more stable than endogenous IL-15 produced by a cell not recombinantly expressing both mammalian IL-15 and IL-15Ra as measured by a technique known to one of skill in the art, e.g., high performance size exclusion chromatography (HPSEC). In some embodiments, the mammalian IL-15 or derivative thereof is stable at 32°C or 37°C for 6 hours, 12 hours, 1 day, 2 days, 5 days, 7 days, 14 days, 1 month, 2 months or more. In particular embodiments, the cell recombinantly expresses a mammalian IL-15 or a derivative thereof that is degraded at a slower rate than endogenous IL-15 produced by a cell not recombinantly expressing both mammalian IL-15 and IL-15Ra. In specific embodiments, the protein degradation rate of recombinant mammalian IL-15 (*in vitro* or *in vivo*) produced by such a cell is at least 1 fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, 50 fold, or 100 fold smaller than the protein degradation rate of endogenous IL-15 produced by a cell not recombinantly expressing both mammalian IL-15 and IL-15Ra as measured by a technique known to one of skill in the art, e.g., ELISA, western blot or HPSEC.

**[0129]** In some embodiments, the present invention encompasses a cell that recombinantly expresses a mammalian IL-15 or a derivative thereof and a mammalian IL-15Ra or a derivative thereof, wherein the cell grows in serum-free media. In certain embodiments, the cell expresses at least 0.6 pg of mammalian IL-15 or a derivative thereof. In certain embodiments, the cell expresses at least 0.1 pg, 0.5 pg, 1 pg, or 2 pg of mammalian IL-15 or a derivative thereof. In some embodiments, the cell expresses approximately 0.1 pg to 0.6 pg, 0.5 pg to 1 pg, 0.5 pg to 2 pg, or 0.1 pg to 2 pg of mammalian IL-15 or a derivative thereof.

**[0130]** In some embodiments, the present invention relates to a population of cells that recombinantly expresses a mammalian IL-15 or a derivative thereof and a mammalian IL-15Ra or a derivative thereof, wherein the population of

5 cells express at least 600 ng/million cells of mammalian IL-15 or a derivative thereof. In specific embodiments, the population of cells express at least 150 ng/million cells per day cells of mammalian IL-15 or a derivative thereof. In some embodiments, the population of cells express at least 50 ng/million cells per day, 100 ng/million cells per day, 200 ng/million cells per day, 250 ng/million cells per day, or 300 ng/million cells per day cells of mammalian IL-15 or a derivative thereof. In particular embodiments, the population of cells express approximately 50 ng/million cells per day to 200 ng/million cells per day, 100 ng/million cells per day to 250 ng/million cells per day, or 50 ng/million cells per day to 300 ng/million cells per day cells of mammalian IL-15 or a derivative thereof. In certain embodiments, the population of cells express approximately 100 ng/million cells to 1000 ng/million cells, 500 ng/million cells to 1000 ng/million cells, 500 ng/million cells to 2000 ng/million cells, or 100 ng/million cells to 2000 ng/million cells of mammalian IL-15 or a derivative thereof.

10 **[0131]** In particular embodiments, the population of cells recombinantly expresses a mammalian IL-15 or a derivative thereof that is more stable than endogenous IL-15 produced by a cell not recombinantly expressing both mammalian IL-15 and IL-15Ra. In specific embodiments, the protein stability of recombinant mammalian IL-15 produced by such a population of cells is at least 1 fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, 50 fold, or 100 fold more stable than endogenous IL-15 produced by a population of cells not recombinantly expressing both mammalian IL-15 and IL-15Ra as measured by a technique known to one of skill in the art, e.g., high performance size exclusion chromatography (HPSEC). In some embodiments, the mammalian IL-15 or derivative thereof is stable at 32°C or 37°C for 6 hours, 12 hours, 1 day, 2 days, 5 days, 7 days, 14 days, 1 month, 2 months or more. In particular embodiments, the population of cells recombinantly expresses a mammalian IL-15 or a derivative thereof that is degraded at a slower rate than endogenous IL-15 produced by a cell not recombinantly expressing both mammalian IL-15 and IL-15Ra. In specific embodiments, the protein degradation rate of recombinant mammalian IL-15 (*in vitro* or *in vivo*) produced by such a population of cells is at least 1 fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, 50 fold, or 100 fold smaller than the protein degradation rate of endogenous IL-15 produced by a population of cells not recombinantly expressing both mammalian IL-15 and IL-15Ra as measured by a technique known to one of skill in the art, e.g., ELISA, western blot or HPSEC.

15 **[0132]** In some embodiments, the present invention relates to a population of cells that recombinantly expresses a mammalian IL-15 or a derivative thereof and a mammalian IL-15Ra or a derivative thereof, wherein the population of cells grows in serum-free media. In particular embodiments, the population of cells express at least 600 ng/million cells of mammalian IL-15 or a derivative thereof. In specific embodiments, the population of cells express at least 150 ng/million cells per day cells of mammalian IL-15 or a derivative thereof. In some embodiments, the population of cells express at least 50 ng/million cells per day, 100 ng/million cells per day, 200 ng/million cells per day, 250 ng/million cells per day, or 300 ng/million cells per day cells of mammalian IL-15 or a derivative thereof. In particular embodiments, the population of cells express approximately 50 ng/million cells per day to 200 ng/million cells per day, 100 ng/million cells per day to 250 ng/million cells per day, or 50 ng/million cells per day to 300 ng/million cells per day cells of mammalian IL-15 or a derivative thereof. In certain embodiments, the population of cells express approximately 100 ng/million cells to 1000 ng/million cells, 500 ng/million cells to 1000 ng/million cells, 500 ng/million cells to 2000 ng/million cells, or 100 ng/million cells to 2000 ng/million cells of mammalian IL-15 or a derivative thereof.

## 5.2. Polymers

40 **[0133]** The invention provides for any of the Therapeutic Agents described in Section 5.1 to be formulated with any natural polymer suitable for biomedical use, including poly- $\beta$ -1 $\rightarrow$ 4-N- acetylglucosamine ("p-GlcNAc"). p-GlcNAc can be found in chitin and chitosan derived from shellfish, fungi, or microalgal sources. In a preferred embodiment, the p-GlcNAc is purified as described in U.S. Patent Application Publication Nos. US 2004/0220140 and US 2001/0055807 and U.S. Patent Nos. 5,622,834; 5,623,064; 5,624,679; 5,686,115; 5,858,350; 6,599,720; 6,686,342; and 7,115,588, which are incorporated by reference herein in their entireties. In a specific embodiment, the polymer is in the form of fibers. However, other forms, such as powder, may be used.

45 **[0134]** Examples of suitable polymers for practicing the invention include, but are not limited to, cellulose-based polymers, xanthan, polyaramides, polyamides, polyimides, polyarnide/imides, polyamidehydrazides, polyhydrazides, poly-imidazoles, polybenzoxazoles, polyester/amide, polyester/imide, polycarbonate/amides, polycarbonate/imides, polysulfone/amides, polysulfone imides, and the like, copolymers and blends thereof. Other suitable classes of polymers that may be used include polyvinylidene fluorides and polyacrylonitriles. Examples of these polymers include those described in U.S. Pat. Nos. 4,705,540; 4,717,393; 4,717,394; 4,912,197; 4,838,900; 4,935,490; 4,851,505; 4,880,442; 4,863,496; and 4,961,539; and European Patent Application No. 0 219 878, all of which are incorporated by reference herein in their entireties. The polymers can include at least one of either of cellulose-based polymers, polyamides, polyaramides, polyamide/imides or polyimides. In certain embodiments, the polymers include polyaramides, polyester, urethan and polytetrafluoroethylene.

55 **[0135]** In some embodiments, polymerized N-acetylglucosamine or derivatives thereof are used. In a preferred em-

bodiment, the polymer is poly-N-acetylglucosamine or a derivative thereof. In certain embodiments, the poly-N-acetylglucosamine has a  $\beta$ -1 $\rightarrow$ 4 configuration. In other embodiments, the poly-N-acetylglucosamine has an  $\alpha$ -1 $\rightarrow$ 4 configuration.

[0136] In specific embodiments, the polymer is chitin, chitosan, cellulose, nylon or PET (polyethylene terephthalate).

[0137] In other specific embodiments, the polymer is biocompatible and/or biodegradable. Biocompatibility may be determined by a variety of techniques, including, but not limited to such procedures as the elution test, intramuscular implantation, or intracutaneous or systemic injection into animal subjects. Such tests are described in U.S. Patent No. 6,686,342, which is incorporated by reference herein in its entirety, or equivalent tests as set forth in ISO-10993 guidances. Biodegradable polymers preferably degrade within about 1 day, 2 day, 5 day, 8 day, 12 day, 17 day, 25 day, 30 day, 35 day, 40 day, 45 day, 50 day, 55 day, 60 day, 65 day, 70 day, 75 day, 80 day, 85 day, 90 day, 95 day, or 100 days after administration or implantation into a patient.

[0138] In certain aspects of the invention, the polymer is immunoneutral.

[0139] In one embodiment, Therapeutic Agents or compositions thereof are formulated with purified polymers, which may be about 100%, 99.9%, 99.8%, 99.5%, 99%, 98%, 95%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, or 20% pure. In a specific embodiment, the polymers used to formulate the Therapeutic Agents or compositions thereof are 90-100% pure.

[0140] In certain embodiments, the polymer that is used to formulate the Therapeutic Agent is not one or more of the following: an ionic synthetic hydrogel such as, but not limited to, crosslinked poly(AAn-acrylic acid) and poly(AAm-dimethylaminoethyl methacrylate), poly(vinyl alcohol) (PVA), poly(ethylene glycol) (PEG), poly(N-vinyl pyrrolidone), poly(methoxy-PEG methacrylate). In certain embodiments, the polymer is not one or more of: a poly-L-amino acid, such as poly L lysine, poly L arginine, poly L glutamic acid, poly L histidine, poly D glutamic acid or a mixture thereof. In certain embodiments, the polymer is not one or more of: an alginate polymer, such as sodium alginate, calcium alginate, strontium alginate, barium alginate, magnesium alginate or any other alginate or a mixture thereof. In certain embodiments, the polymer is not derived from one or more of: a shell fish, a crustacean, insect, fungi and/or yeasts. In certain embodiments, the polymers do not comprise collagen fibers. In certain embodiments, the polymers do not comprise elastin fibers. In certain embodiments, the polymers do not comprise block polymer polaxamer 407. In certain embodiments, the polymers do not comprise nondegradable polymeric matrices. In certain embodiments, the polymers do not comprise degradable polymeric matrices. In other embodiments, any of the above-referenced polymers are included in the Therapeutic Agents or compositions thereof. For example, in certain embodiments, the polymer that is used to formulate a Therapeutic Agent is one or more of the following: an ionic synthetic hydrogel such as, but not limited to, crosslinked poly(AAn-acrylic acid) and poly(AAm-dimethylaminoethyl methacrylate), poly(vinyl alcohol) (PVA), poly(ethylene glycol) (PEG), poly(N-vinyl pyrrolidone), poly(methoxy-PEG methacrylate). In certain embodiments, the polymer is not one or more of: a poly-L-amino acid, such as poly L lysine, poly L arginine, poly L glutamic acid, poly L histidine, poly D glutamic acid or a mixture thereof. In certain embodiments, the polymer is one or more of: an alginate polymer, such as sodium alginate, calcium alginate, strontium alginate, barium alginate, magnesium alginate or any other alginate or a mixture thereof. In certain embodiments, the polymer is derived from one or more of: a shell fish, a crustacean, insect, fungi and/or yeasts. In certain embodiments, the polymers does comprise collagen fibers. In certain embodiments, the polymers do comprise elastin fibers. In certain embodiments, the polymers do comprise block polymer polaxamer 407. In certain embodiments, the polymers do comprise nondegradable polymeric matrices. In certain embodiments, the polymers do comprise degradable polymeric matrices.

[0141] The polymers can be in the form of fibers. Fibers may be about 0.20, 0.25, 0.30, 0.35, 0.40, 0.54, 0.50, 0.55, 0.60, or 0.65 microns in thickness and/or diameter as determined by electron microscopy. In preferred embodiments the fibers are about 0.50 microns in width and range in length from about 20 to 100 microns as determined by electron microscopy, particularly, scanning electron microscopy. In another preferred embodiments the fibers are about 0.50 microns in width and range in length from about 50 to 100 microns as determined by electron microscopy, particularly, scanning electron microscopy. In yet another preferred embodiments the fibers are about 0.50 microns in width and range in length from about 75 to 100 microns as determined by electron microscopy, particularly, scanning electron microscopy.

[0142] In certain embodiments, at least 75%, at least 85%, at least 90%, or at least 95% of a Therapeutic Agent is formulated with one or more of the polymers listed above.

[0143] In one embodiment, the Therapeutic Agents comprise more than one type of polymer (e.g., poly- $\beta$ -1 $\rightarrow$ 4-N-acetylglucosamine and cellulose).

### 5.2.1. Poly- $\beta$ -1 $\rightarrow$ 4-N-Acetylglucosamine.

[0144] A preferred polymer for use in combination with a Therapeutic Agent is poly-N-acetylglucosamine or a derivative thereof.

[0145] As used herein, poly-N-acetylglucosamine includes any polymer of N-acetylglucosamine and/or glucosamine covalently linked in a  $\beta$ -1 $\rightarrow$ 4 or an  $\alpha$ -1 $\rightarrow$ 4 conformation and in any degree or form of crystallinity. In specific embodiments, poly-N-acetylglucosamine is (1) a semi-crystalline form of poly-N-acetylglucosamine; (2) a poly-N-acetylglucosamine

comprising about 50 to about 150,000 N-acetylglucosamine monosaccharides covalently attached in a  $\beta$ -1 $\rightarrow$ 4 conformation and having a molecular weight of about 10,000 daltons to about 30 million daltons; (3) a poly- $\beta$ -1 $\rightarrow$ 4-acetylglucosamine comprising about 50 to about 50,000 N-acetylglucosamine monosaccharides covalently attached in a  $\beta$ -1 $\rightarrow$ 4 conformation and having a molecular weight of about 10,000 daltons to about 10 million daltons; (4) a poly- $\beta$ -1 $\rightarrow$ 4-acetylglucosamine comprising about 50 to about 10,000 N-acetylglucosamine monosaccharides covalently attached in a  $\beta$ -1 $\rightarrow$ 4 conformation and having a molecular weight of about 10,000 daltons to about 2 million daltons; (5) a poly- $\beta$ -1 $\rightarrow$ 4-N-acetylglucosamine comprising about 50 to about 4,000 N-acetylglucosamine monosaccharides covalently attached in a  $\beta$ -1 $\rightarrow$ 4 conformation and having a molecular weight of about 10,000 daltons to about 800,000 daltons; and (6) deacetylated counterparts of (1)-(5) above in which the degree of deacetylation ranges from 1%-99%. In specific embodiments, the degree of deacetylation is at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90%.

**[0146]** Derivatives, such as chemical derivatives, of poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may also be used to formulate the Therapeutic Agents. For example, sulfated poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine derivatives, phosphorylated poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine derivatives, or nitrated poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine derivatives may be used. Additionally, one or more of the monosaccharide units of the poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may contain one or more sulfonyl groups one or more O-acyl groups. In addition, one or more of the monosaccharides of the deacetylated poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may contain an N-acyl group. One or more of the monosaccharides of the poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine or of its deacetylated derivative, may contain an O-alkyl group. One or more of the monosaccharide units of the poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may be an alkali derivative. One or more of the monosaccharide units of the deacetylated derivative of poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may contain an N-alkyl group. One or more of the monosaccharide units of the deacetylated derivative of poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may contain at least one deoxyhalogen derivative. One or more of the monosaccharide units of the deacetylated derivative of poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may form a salt. One or more of the monosaccharide units of the deacetylated derivative of poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may form a metal chelate. Preferably, the metal is zinc. One or more of the monosaccharide units of the deacetylated derivative of poly- $\beta$ -1 $\rightarrow$ 4 N-acetylglucosamine may contain an N-alkylidene or an N-arylidene group. Methods of making such derivatives are described in U.S. Patent No. 5,623,064, which is incorporated by reference herein in its entirety.

**[0147]** In specific embodiments, poly-N-acetylglucosamine is obtained by a process comprising a) treating a microalga comprising a cell body and a poly-N-acetylglucosamine fiber with a biological agent (such as hydrofluoric acid) capable of separating the N-acetylglucosamine fiber from the cell body for a sufficient time so that the poly-N-acetylglucosamine fiber is released from the cell body; b) segregating the poly-N-acetylglucosamine fiber from the cell body; and c) removing contaminants from the segregated poly-N-acetylglucosamine fiber. Detailed descriptions of making poly-N-acetylglucosamine according to such methods are described in U.S. Patent Application Publication Nos. US 2004/0220140 and US 2001/055807, and U.S. Patent Nos. 5,622,834; 5,623,064; 5,624,679; 5,686,115; 5,858,350; 6,599,720; 6,686,342; and 7,115,588, the contents of which are incorporated herein in their entireties.

**[0148]** As an alternative source to microalgal p-GlcNAc, p-GlcNAc purified from crustacean or fungal chitin or chitosan may be used to formulate a Therapeutic Agent according to the present invention.

**[0149]** Optionally, the p-GlcNAc is used in fiber form, for example as described above.

#### 40 5.3. Compositions

**[0150]** The invention provides compositions comprising the Therapeutic Agents, including Agonistic Therapeutic Agents and Antagonist Therapeutic Agents. The compositions include bulk drug compositions useful in the manufacture of pharmaceutical compositions (e.g., impure or non-sterile compositions) and pharmaceutical compositions (i.e., compositions that are suitable for administration to a subject or patient) which can be used in the preparation of unit dosage forms. The compositions comprise an effective amount of a Therapeutic Agent or a combination of Therapeutic Agents and a pharmaceutically acceptable carrier. In specific embodiments, the compositions comprise an effective amount of one or more Therapeutic Agents and a pharmaceutically acceptable carrier. In some embodiments, the composition further comprises an additional therapeutic, e.g., anti-cancer agent, anti-viral agent, anti-inflammatory agent, adjuvant. Non-limiting examples of such therapeutics are provided in Section 5.4.5, *infra*.

**[0151]** In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant (e.g., Freund's adjuvant (complete and incomplete) or, more preferably, MF59C.1 adjuvant available from Chiron, Emeryville, CA), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. In one embodiment, water is a carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers,

particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like.

5

### 5.3.1. Formulations with Polymers

[0152] The present invention provides for compositions (and pharmaceutical compositions) comprising a Therapeutic Agent that are formulated with a polymer, such as described in Section 5.2, *supra*. Various formulations comprising a Therapeutic Agent and a polymer are described herein.

10

[0153] The polymers can be formulated as a gel, solid, liquid, sponge, foam, spray, emulsion, suspension, solution, mat, string, gauze, suture, bead, microsphere, or microfibril.

15

[0154] In specific embodiments, the polymers are formulated as barriers, membranes, or films. Alternatively, the polymers are added to barriers, membranes, or films. A barrier, membrane, or film can be supplied in a variety of standard sizes, which can be further cut and sized to the area being treated. Alternatively, the polymer can be formulated as a barrier, membrane, or film made out of strings, microbeads, microspheres, or microfibrils, or the composition can be formulated as a barrier-forming mat. The pharmaceutical compositions comprising Therapeutic Agents and polymers can include a pharmaceutically acceptable carrier, a neutral liquid, neutral gel or neutral solid.

20

[0155] In addition, by varying the ratio of the components in said biodegradable matrices, the surgical handling properties of the cell matrix can be adjusted in a range from a dimensionally stable matrix, to a moldable putty-like consistency to a pliable gel or slurry, to a powder or to an injectable fluid.

25

[0156] In specific embodiments, the polymer is formulated as a gel. The gel can be of varying viscosity. For various embodiments, a gel with a low viscosity is desired. For injectable gels, higher viscosity may be desired if the Therapeutic Agents or compositions thereof are intended to remain in a location of the body rather than dissipate rapidly. Viscosity is the quantity that describes a fluid's resistance to flow measured in centipoise (cP). While the range of viscosity is a continuum. For example, as a frame of reference, not as a limitation of the meaning of viscosity, the viscosity values of about 1-4 cP generally are typified by fluid compositions. Viscosity values of about 5-14 cP generally are typified by gel-like compositions, while viscosity values of 15-20 cP are relatively hard compositions such as plastics. The viscosity of cell cytoplasm is about 11 cP. Viscosity can be measured with, for example, a Saybolt International B.V. (Vlaardingen, The Netherlands). One skilled in the art can also use other measurement techniques and devices common in the art.

30

[0157] In other embodiments, the polymer is formulated as a sponge. When the polymer is a sponge, a predetermined amount of a cell suspension can be transferred on top of a sponge matrix, and the cell suspension can be absorbed. Therapeutic Agents comprising cells expressing IL-15 and IL-15Ra in high amounts can be formulated with polymers which typically comprise a polymer fiber matrix with cells associated throughout the matrix. This allows for greater cell numbers to interact with fibers and the matrix is able to absorb large numbers of cells. In one embodiment, high amounts of IL-15/IL-15Ra complexes refer to amounts of IL-15/IL-15Ra complexes expressed by cells that are at least 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, or more than 20 fold higher than amounts of IL-15/IL-15Ra complexes expressed endogenously by control cells (e.g., cells that have not been genetically engineered to recombinantly express IL-15, IL-15Ra, or both IL-15 and IL-15Ra, or cells comprising an empty vector).

35

[0158] In certain embodiments, the polymer is formulated as a membrane. The membranes may be porous or relatively continuous. In some embodiments the membranes are made of woven polymer fibers that have been combined with Therapeutic Agents comprising cells expressing IL-15 and IL-15Ra. In specific embodiments, Such membranes are particularly useful for delivery on the surface of the skin, surface of internal organs, or surface of lining of body cavities.

40

[0159] In some embodiments, Therapeutic Agents comprising cells recombinantly expressing IL-15, and another therapeutic agent (e.g., a cytokine, e.g., IL-12 or IL-15, or a soluble receptor, e.g., soluble IL-15Ra) are formulated with a polymer, e.g., a polymer described in Section 5.2, *supra*. In some embodiments, Therapeutic Agents comprising cells recombinantly expressing IL-15Ra, and another therapeutic agent (e.g., a cytokine, e.g., IL-12 or IL-15) are formulated with a polymer, e.g., a polymer described in Section 5.2, *supra*. In some embodiments, Therapeutic Agents comprising cells recombinantly expressing (i) IL-15, or IL-15Ra, or IL-15 and IL-15Ra, and (ii) another therapeutic polypeptide (e.g., a cytokine, e.g., IL-12, IL-6, or GM-CSF), are formulated with a polymer, e.g., a polymer described in Section 5.2, *supra*. In a particular embodiment, the cells are irradiated. In a specific embodiment, the cells are irradiated cancer cells engineered to recombinantly express IL-15, or IL-15Ra, or IL-15 and IL-15Ra.

45

[0160] In certain embodiments, the number of cells is at least about 100, 200, 300, 400, 500, 700, 1,000, 5,000, 10,000, 25,000, 50,000, or 100,000 cells. In specific embodiments, the number of cells is at least about 100, 200, 300, 400, or 500 cells. In other embodiments, the number of cells is at least about 300, 400, 500, 600, 700, 800, 900, 1,000, 2,000, 3,000, 4,000 or 5,000 cells. In yet other specific embodiments, the number of cells is at least about 700, 1,000, 5,000, 10,000, 15,000, or 20,000 cells. In some embodiments, the number of cells is at least about 5,000, 10,000, 25,000,

50,000, 75,000 or 100,000 cells. In yet another embodiment, the number of cells is at least about 50,000, 100,000, 200,000, 300,000, 400,000 or 500,000 cells. In other embodiments, the number of cells is at least about  $1 \times 10^6$ ,  $5 \times 10^6$ ,  $1 \times 10^7$ ,  $5 \times 10^7$ ,  $1 \times 10^8$ ,  $5 \times 10^8$  or more cells per mg of polymer fiber. In other embodiments, the number of cells is at least about  $1 \times 10^6$ ,  $5 \times 10^6$ , or  $1 \times 10^7$  or more cells per mg of polymer fiber. In some embodiments, the number of cells is at least 1  $\times 10^7$ ,  $5 \times 10^7$ ,  $1 \times 10^8$  or more cells per mg of polymer fiber. In other embodiments, the number of cells is at least  $5 \times 10^7$ ,  $1 \times 10^8$ ,  $5 \times 10^8$  or more cells per mg of polymer fiber. In some embodiments, the number of cells is less than  $1 \times 10^4$ ,  $5 \times 10^4$ ,  $1 \times 10^5$ ,  $5 \times 10^5$ ,  $1 \times 10^6$ ,  $5 \times 10^6$ ,  $1 \times 10^7$ ,  $5 \times 10^7$ ,  $1 \times 10^8$ , or  $5 \times 10^8$  cells.

### 10 5.3.2. Formulations without Polymers

[0161] Pharmaceutical compositions for use in accordance with the present invention may be formulated in any conventional manner using one or more pharmaceutically acceptable carriers or excipients.

[0162] Generally, the components of the pharmaceutical compositions comprising Therapeutic Agents are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the Therapeutic Agent is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline (e.g., PBS). Where the Therapeutic Agent is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[0163] In some embodiments, Therapeutic Agents may be formulated for administration by any method known to one of skill in the art, including but not limited to, inhalation, insufflation (either through the mouth or the nose), oral, intradermal, transdermal, intraparenteral, intratumoral, and mucosal (such as buccal, vaginal, rectal, sublingual) administration.

[0164] In a specific embodiment, the Therapeutic Agents are formulated for local or systemic parenteral administration. In one embodiment, the Therapeutic Agents are formulated in a pharmaceutically compatible solution.

[0165] For oral administration, the pharmaceutical compositions comprising Therapeutic Agents that are polypeptides or nucleic acids may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring and sweetening agents as appropriate. Preparations for oral administration may be suitably formulated to give controlled release of the Therapeutic Agent or compositions thereof. For buccal administration the compositions may take the form of tablets or lozenges formulated in conventional manner.

[0166] For administration by inhalation, the Therapeutic Agents are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0167] The Therapeutic Agents can be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use.

[0168] The Therapeutic Agents may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

[0169] In addition to the formulations described previously, the Therapeutic Agents may also be formulated for implantation (for example subcutaneously or intramuscularly) or by intramuscular injection.

[0170] In a specific embodiment, the formulation and administration of various chemotherapeutic, biological/immuno-therapeutic and hormonal therapeutic agents for use in combination with Therapeutic Agents are known in the art and described in the Physician's Desk Reference, 61st ed. (2007). In some embodiments, the Therapeutic Agents are formulated with other therapies, such as those described in Section 5.4 below. In some embodiments, Therapeutic

Agents comprising cells recombinantly expressing IL-15, or IL-15Ra, or IL-15 and IL-15Ra are formulated as pharmaceutical compositions. In some embodiments, Therapeutic Agents comprising irradiated cancer cells recombinantly expressing IL-15, or IL-15Ra, or IL-15 and IL-15Ra are formulated as pharmaceutical compositions. In some embodiments, Therapeutic Agents comprising cells recombinantly expressing (i) IL-15, or IL-15Ra, or IL-15 and IL-15Ra; and (ii) another therapeutic polypeptide (e.g., a cytokine, e.g., IL-12, IL-6, or GM-CSF), are formulated as pharmaceutical compositions. In some embodiments, Therapeutic Agents comprising irradiated cancer cells recombinantly expressing (i) IL-15, or IL-15Ra, or IL-15 and IL-15Ra; and (ii) another therapeutic polypeptide (e.g., a cytokine, e.g., IL-12, IL-6, or GM-CSF), are formulated as pharmaceutical compositions. In some embodiments, a combination of (i) Therapeutic Agents that are cells recombinantly expressing IL-15, or IL-15Ra, or IL-15 and IL-15Ra; and (ii) one or more other therapies, e.g., a cytokine (e.g., IL-12, IL-6, GM-CSF); are formulated as pharmaceutical compositions. In some embodiments, a combination of (i) Therapeutic Agents that are irradiated cancer cells recombinantly expressing IL-15, or IL-15Ra, or IL-15 and IL-15Ra; and (ii) one or more other therapies, e.g., a cytokine (e.g., IL-12, IL-6, GM-CSF); are formulated as pharmaceutical compositions.

15 **5.4. Prophylactic/Therapeutic Methods**

**5.4.1. Enhancing Immune Function**

20 [0171] The invention provides methods for enhancing IL-15 mediated immune function in a subject, comprising administering an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject in need thereof. In a specific embodiment, the invention provides methods for preventing, treating, and/or managing diseases in which it is desirable to enhance immune function, comprising administering an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject in need thereof. In specific embodiments, the Agonistic Therapeutic Agents are formulated with polymers described in Section 5.2, *supra*. In other specific embodiments, the method comprises combination therapy, wherein the Agonistic Therapeutic Agent is administered to a subject in combination with another therapy, such as those described below, to enhance IL-15 mediated immune function. In a particular embodiment, the Agonistic Therapeutic Agent is administered in combination with a vaccine composition to induce or enhance the immune response elicited by the vaccine composition. Non-limiting examples of diseases that can be prevented, treated, or managed by an enhancement of immune function include, but are not limited to, cancer and infectious diseases. Various cancers and infectious diseases are described below. In a specific embodiment, an Agonistic Therapeutic Agent described herein can be used to treat or manage a condition associated with cancer or a condition resulting from the administration of an anti-cancer therapy (such as, e.g., chemotherapy or radiation). In another embodiment, an Agonistic Therapeutic Agent is administered to a patient diagnosed with cancer to increase the proliferation and/or effector function of one or more immune cell populations in the patient.

25 [0172] In a specific embodiment, an Agonistic Therapeutic Agent enhances or induces immune function in a subject by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the immune function in a subject not administered the Agonistic Therapeutic Agent using assays well known in the art, e.g., ELISPOT, ELISA, and cell proliferation assays. In a specific embodiment, the immune function is cytokine release (e.g., interferon-gamma, IL-2, IL-5, IL-10, IL-12, or transforming growth factor (TGF) -beta). In one embodiment, the IL-15 mediated immune function is NK cell proliferation, which can be assayed, e.g., by flow cytometry to detect the number of cells expressing markers of NK cells (e.g., CD56). In another embodiment, the IL-15 mediated immune function is antibody production, which can be assayed, e.g., by ELISA. In some embodiments, the IL-15 mediated immune function is effector function, which can be assayed, e.g., by a cytotoxicity assay or other assays well known in the art.

30 [0173] In specific embodiments, non-limiting examples of immune function enhanced by the Agonistic Therapeutic Agent are proliferation/expansion of lymphocytes (e.g., increase in the number of lymphocytes), inhibition of apoptosis of lymphocytes, activation of dendritic cells (or antigen presenting cells), and antigen presentation. In particular embodiments, an immune function enhanced by the Agonistic Therapeutic Agent is proliferation/expansion in the number of or activation of CD4<sup>+</sup> T cells (e.g., Th1 and Th2 helper T cells), CD8<sup>+</sup> T cells (e.g., cytotoxic T lymphocytes, alpha/beta T cells, and gamma/delta T cells), B cells (e.g., plasma cells), memory T cells, memory B cells, dendritic cells (immature or mature), antigen presenting cells, macrophages, mast cells, natural killer T cells (NKT cells), tumor-resident T cells, CD122<sup>+</sup> T cells, or natural killer cells (NK cells). In one embodiment, the Agonistic Therapeutic Agent enhances the proliferation/expansion or number of lymphocyte progenitors. In some embodiments, an Agonistic Therapeutic Agent increases the number of CD4<sup>+</sup> T cells (e.g., Th1 and Th2 helper T cells), CD8<sup>+</sup> T cells (e.g., cytotoxic T lymphocytes, alpha/beta T cells, and gamma/delta T cells), B cells (e.g., plasma cells), memory T cells, memory B cells, dendritic cells (immature or mature), antigen presenting cells, macrophages, mast cells, natural killer T cells (NKT cells), tumor-resident T cells, CD122<sup>+</sup> T cells, or natural killer cells (NK cells) by approximately 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, or more relative a negative control (e.g., number of the respective cells not treated, cultured,

or contacted with an Agonistic Therapeutic Agent).

#### 5.4.1.1. Cancer

5 [0174] The invention provides a method for preventing, treating, and/or managing cancer, comprising administering an effective amount of an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject in need thereof. In specific embodiments, the Agonistic Therapeutic Agents are formulated with polymers described in Section 5.2, *supra*.

10 [0175] The effect of an Agonistic Therapeutic Agent on proliferation of cancer cells can be detected by routine assays, such as by assays that measure the uptake of radiolabeled thymidine. Alternatively, cell viability can be measured by assays that measure lactate dehydrogenase (LDH), a stable cytosolic enzyme that is released upon cell lysis, or by the release of [<sup>51</sup>Cr] upon cell lysis. In one embodiment, necrosis measured by the ability or inability of a cell to take up a dye such as neutral red, trypan blue, or ALAMARTM blue (Page et al., 1993, *Intl. J. of Oncology* 3:473 476). In such an assay, the cells are incubated in media containing the dye, the cells are washed, and the remaining dye, reflecting cellular uptake of the dye, is measured spectrophotometrically.

15 [0176] In another embodiment, the dye is sulforhodamine B (SRB), whose binding to proteins can be used as a measure of cytotoxicity (Skehan et al., 1990, *J. Nat'l Cancer Inst.* 82:1107 12). In yet another embodiment, a tetrazolium salt, such as MTT, is used in a quantitative colorimetric assay for mammalian cell survival and proliferation by detecting living, but not dead, cells (see, e.g., Mosmann, 1983, *J. Immunol. Methods* 65:55 63).

20 [0177] In other embodiments, apoptotic cells are measured in both the attached and "floating" compartments of the cultures. Both compartments are collected by removing the supernatant, trypsinizing the attached cells, and combining both preparations following a centrifugation wash step (10 minutes, 2000 rpm). The protocol for treating tumor cell cultures with sulindac and related compounds to obtain a significant amount of apoptosis has been described in the literature (see, e.g., Piazza et al., 1995, *Cancer Research* 55:3110 16). Features of this method include collecting both 25 floating and attached cells, identification of the optimal treatment times and dose range for observing apoptosis, and identification of optimal cell culture conditions.

25 [0178] In another embodiment, apoptosis is quantitated by measuring DNA fragmentation. Commercial photometric methods for the quantitative *in vitro* determination of DNA fragmentation are available. Examples of such assays, including TUNEL (which detects incorporation of labeled nucleotides in fragmented DNA) and ELISA-based assays, are described in *Biochemica*, 1999, no. 2, pp. 34 37 (Roche Molecular Biochemicals).

30 [0179] In yet another embodiment, apoptosis can be observed morphologically.

[0180] Cancer cell lines on which such assays can be performed are well known to those of skill in the art. Apoptosis, necrosis and proliferation assays can also be performed on primary cells, e.g., a tissue explant.

35 [0181] In a specific embodiment, the proliferation or viability of cancer cells contacted with an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent is inhibited or reduced by at least 2 fold, preferably at least 2.5 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 7 fold, or at least 10 fold relative to the proliferation of the cancer cells when contacted with a negative control as measured using assays well known in the art, e.g., cell proliferation assays using CSFE, BrdU, and <sup>3</sup>H-Thymidine incorporation. In another embodiment, the proliferation of 40 cancer cells contacted with an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent is inhibited or reduced by at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% relative to cancer cells contacted with a negative control as measured using assays well known in the art, e.g., cell proliferation assays using CSFE, BrdU, and <sup>3</sup>H-Thymidine incorporation, or those assays described above. In one aspect, the composition comprising an Agonistic Therapeutic Agent further comprises cells (e.g., NK cells or cytotoxic T cells) 45 that are responsive to IL-15 signaling and that can target and exert cytotoxic effects on the cancer cells.

50 [0182] In a specific embodiment, the administration of an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject with cancer (in some embodiments, an animal model for cancer) inhibits or reduces the growth of a tumor by at least 2 fold, preferably at least 2.5 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 7 fold, or at least 10 fold relative to the growth of a tumor in a subject with cancer (in some embodiments, in the same animal model for cancer) administered a negative control as measured using assays well known in the art. In another embodiment, the administration of an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject with cancer (in some embodiments, an animal model for cancer) inhibits or reduces the growth of a tumor by at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% relative 55 to the growth of a tumor in a subject with cancer (in some embodiments, in the same animal model for cancer) administered a negative control as measured using assays well known in the art.

[0183] In a specific embodiment, the administration of an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject with cancer (in some embodiments, an animal model for cancer) reduces

the size of a tumor by at least 2 fold, preferably at least 2.5 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 7 fold, or at least 10 fold relative to the growth of a tumor in a subject with cancer (in some embodiments, the same animal model for cancer) administered a negative control as measured using assays well known in the art. In another embodiment, the administration of an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject with (in some embodiments, an animal model for cancer) reduces the size of a tumor by at least 25%, at least 5

30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% relative to the growth of a tumor in a subject with cancer (in some embodiments, the same animal model for cancer) administered a negative control as measured using assays well known in the art. In a specific embodiment, the cancer is melanoma, renal cancer, colon cancer, or prostate 10 cancer.

**[0184]** In addition, an Agonistic Therapeutic Agent that is useful in the treatment or management of cancer is an irradiated tumor (or cancer) cell or irradiated tumor (or cancer) cell line that recombinantly expresses IL-15/IL-15Ra complexes. Such Agonistic Therapeutic Agents can be prepared by a method that comprises transducing or transfecting 15 primary tumor cells or tumor cell lines with expression constructs for IL-15 and IL-15Ra, and following expression of the IL-15 and IL-15Ra proteins, irradiating the tumor cells. In one embodiment, the form of irradiation is  $\gamma$ -radiation. IL-15 and IL-15Ra can be expressed from the same construct or different constructs, and under the control of the same promoter or different promoters. The IL-15 and IL-15Ra expression construct(s) can be introduced into tumor (or cancer) 20 cells by any means known to those of skill in the art, such as using a viral vector or mammalian expression vector. Alternatively, the native IL-15 and native IL-15Ra of the tumor (or cancer) cells may be activated using gene activation 25 methods. Subsequently, the irradiated tumor (or cancer) cells recombinantly expressing IL-15/IL-15Ra complexes are administered into a cancer patient to induce and/or enhance an immune response to the cancer cells or to cancer antigens of the cancer cells. In a specific embodiment, the induced or enhanced immune response is increased production of antibodies to the cancer cells or to the cancer antigens of the cancer cells in the cancer patient. In another embodiment, the induced or enhanced immune response is an increase in effector cell function, e.g., antibody-dependent cellular cytotoxicity (ADCC) against cancer cells in the patient. In some embodiments, the induced or enhanced immune response is increase in lymphocyte number, lymphocyte proliferation, and/or lymphocyte activity.

**[0185]** An Agonistic Therapeutic Agent can be administered in combination with one or more other therapies, e.g., anti-cancer agents, cytokines or anti-hormonal agents, to treat and/or manage cancer. Non-limiting examples anti-cancer agents are described in Section 5.4.5.1 below. In one embodiment, the combination of an Agonistic Therapeutic Agent 30 and one or more other therapies provides an additive therapeutic effect relative to the therapeutic effects of the Agonistic Therapeutic Agent alone or the one or more other therapies alone. In one embodiment, the combination of an Agonistic Therapeutic Agent and one or more other therapies provides more than an additive therapeutic effect relative to the therapeutic effects of the Agonistic Therapeutic Agent alone or the one or more other therapies alone. In one embodiment, the combination of an Agonistic Therapeutic Agent and one or more other therapies provides a synergistic therapeutic 35 effect relative to the therapeutic effects of the Agonistic Therapeutic Agent alone or the one or more other therapies alone.

**[0186]** In one embodiment, the one or more therapies include, but are not limited to cytokines/growth factors, e.g., interleukin (IL) 1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, IL-11, IL-12, IL-15, TNF- $\alpha$ , TNF- $\beta$ , TGF- $\beta$ , GM-CSF, and interferon- $\gamma$ . In one embodiment, the one or more therapies include, but are not limited to receptors, antibodies, or other binding agents that bind to cytokines/growth factors, e.g., interleukin (IL) 1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, 40 IL-11, IL-12, TNF- $\alpha$ , TNF- $\beta$ , TGF- $\beta$ , GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , and interferon- $\gamma$ . In some embodiments, the one or more therapies include, but are not limited to, cells recombinantly expressing a therapeutic protein (or polypeptides), e.g., a cytokine, a growth factor, a chemokine, or a fragment or derivative thereof. In a particular embodiment, the one or more therapies include, but are not limited to, cells recombinantly expressing IL-12, IL-6, GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$  or TNF- $\alpha$ . In one embodiment, the one or more therapies are not cells recombinantly expressing IL-12. In one embodiment, the one or more therapies are not cells recombinantly expressing IL-6. In one embodiment, the one or more therapies are not cells recombinantly expressing GM-CSF. In one embodiment, the one or more therapies are not cells recombinantly expressing TNF- $\alpha$ . In a specific embodiment, the cells recombinantly expressing a therapeutic protein (polypeptide) are cancer cells. In a specific embodiment, the cells recombinantly expressing a therapeutic protein (polypeptide) are irradiated cancer cells. In a specific embodiment, the cells recombinantly expressing a therapeutic protein (polypeptide) are irradiated cancer cells obtained from a patient. In one embodiment, the cells recombinantly express one or more therapeutic proteins (polypeptides). In one embodiment, the present invention provides for a combination of an Agonistic Therapeutic Agent and one or more other therapies, wherein the Agonistic Therapeutic Agent comprises cells recombinantly express IL-15 and IL-15Ra, and wherein the one or more other therapies comprise one or more exogenous cytokines (e.g., IL-12, IL-6, GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$  or TNF- $\alpha$ ). In one embodiment, the present invention provides for a combination of an Agonistic Therapeutic Agent and one or more other therapies, wherein the Agonistic Therapeutic Agent comprises cells recombinantly express 50 IL-15 and IL-15Ra, and wherein the one or more other therapies comprise (i) exogenous IL-15 polypeptide or exogenous IL-15Ra polypeptide, and (ii) one or more other exogenous cytokines (e.g., IL-12, IL-6, GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$  or TNF- $\alpha$ ).

$\beta$ , interferon- $\gamma$  or TNF- $\alpha$ ). In one embodiment, an Agonistic Therapeutic Agent comprises cells engineered to recombinantly express (i) IL-15, IL-15Ra, or IL-15 and IL-15Ra; and (ii) one or more cytokines/growth factors.

[0187] An Agonistic Therapeutic Agent can also be administered in combination with radiation therapy comprising, e.g., the use of x-rays, gamma rays and other sources of radiation to destroy the cancer cells. In specific embodiments, the radiation treatment is administered as external beam radiation or teletherapy wherein the radiation is directed from a remote source. In other embodiments, the radiation treatment is administered as internal therapy or brachytherapy wherein a radioactive source is placed inside the body close to cancer cells or a tumor mass. In one aspect, the Agonistic Therapeutic Agent can enhance the immune function of cancer patient with a compromised immune system due to anti-cancer therapy. An Agonistic Therapeutic Agent can also be administered in combination with chemotherapy. In one embodiment, an Agonistic Therapeutic Agent can be used before, during or after radiation therapy or chemotherapy. In one embodiment, an Agonistic Therapeutic Agent can be used before, during or after surgery. In one embodiment, the present invention provides for a combination of transplant and an Agonistic Therapeutic Agent.

[0188] Non-limiting examples of anti-hormonal agents are anti-hormonal agents that act to regulate or inhibit hormone action on tumors, such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX® tamoxifen), raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON toremifene; aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE® megestrol acetate, AROMASIN®V exemestane, formestan, fadrozole, RIVISOR® vorozole, FEMARA® letrozole, and ARIMIDEX®D anastrozole; and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; as well as troxacitabine (a 1,3-dioxolane nucleoside cytosine analog); antisense oligonucleotides, particularly those which inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC-alpha, Raf and H-Ras; ribozymes such as a VEGF expression inhibitor (e.g., ANGIOZYME® ribozyme) and a HER2 expression inhibitor; vaccines such as gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEU-VECTIN® vaccine, and VAXID® vaccine; PROLEUKIN® rIL-2; LURTOTECAN® topoisomerase 1 inhibitor; ABARELX® rmRH; Vinorelbine and Esperamicins (see U.S. Pat. No. 4,675,187), and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0189] Cancers and related disorders that can be prevented, treated, or managed in accordance with the methods described herein include, but are not limited to, the following: Leukemias including, but not limited to, acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemias such as myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia leukemias and myelodysplastic syndrome, chronic leukemias such as but not limited to, chronic myelocytic (granulocytic) leukemia, and chronic lymphocytic leukemia, hairy cell leukemia; polycythemia vera; lymphomas such as but not limited to Hodgkin's disease, and non-Hodgkin's disease; multiple myelomas such as but not limited to smoldering multiple myeloma, nonsecretory myeloma, osteosclerotic myeloma, plasma cell leukemia, solitary plasmacytoma and extramedullary plasmacytoma; Waldenstrom's macroglobulinemia; monoclonal gammopathy of undetermined significance; benign monoclonal gammopathy; heavy chain disease; bone and connective tissue sarcomas such as but not limited to bone sarcoma, osteosarcoma, chondrosarcoma, Ewing's sarcoma, malignant giant cell tumor, fibrosarcoma of bone, chordoma, periosteal sarcoma, soft-tissue sarcomas, angiosarcoma (hemangiosarcoma), fibrosarcoma, Kaposi's sarcoma, leiomyosarcoma, liposarcoma, lymphangiosarcoma, neurilemmoma, rhabdomyosarcoma, and synovial sarcoma; brain tumors including but not limited to, glioma, astrocytoma, brain stem glioma, ependymoma, oligodendrogioma, nonglial tumor, acoustic neurinoma, craniopharyngioma, medulloblastoma, meningioma, pineocytoma, pineoblastoma, and primary brain lymphoma; breast cancer including, but not limited to, adenocarcinoma, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, Paget's disease, and inflammatory breast cancer; adrenal cancer, including but not limited to, pheochromocytom and adrenocortical carcinoma; thyroid cancer such as but not limited to papillary or follicular thyroid cancer, medullary thyroid cancer and anaplastic thyroid cancer; pancreatic cancer, including but not limited to, insulinoma, gastrinoma, glucagonoma, vipoma, somatostatinsecreting tumor, and carcinoid or islet cell tumor; pituitary cancers including but not limited to, Cushing's disease, prolactin-secreting tumor, acromegaly, and diabetes insipidus; eye cancers including but not limited to, ocular melanoma such as iris melanoma, choroidal melanoma, and ciliary body melanoma, and retinoblastoma; vaginal cancers, including but not limited to, squamous cell carcinoma, adenocarcinoma, and melanoma; vulvar cancer, including but not limited to, squamous cell carcinoma, melanoma, adenocarcinoma, basal cell carcinoma, sarcoma, and Paget's disease; cervical cancers including but not limited to, squamous cell carcinoma, and adenocarcinoma; uterine cancers including but not limited to, endometrial carcinoma and uterine sarcoma; ovarian cancers including but not limited to, ovarian epithelial carcinoma, borderline tumor, germ cell tumor, and stromal tumor; esophageal cancers including but not limited to, squamous cancer, adenocarcinoma, adenoid cystic carcinoma, mucoepidermoid carcinoma, adenosquamous carcinoma, sarcoma, melanoma, plasmacytoma, verrucous carcinoma, and oat cell (small cell) carcinoma; stomach cancers including but not limited to, adenocarcinoma, fungating (polypoid), ulcerating, superficial spreading, diffusely spreading, malignant lymphoma, liposarcoma, fibrosarcoma, and carcinosarcoma; colon cancers; rectal cancers; liver cancers including but not limited to hepatocellular carcinoma and hepatob-

lastoma; gallbladder cancers including but not limited to, adenocarcinoma; cholangiocarcinomas including but not limited to, papillary, nodular, and diffuse; lung cancers including but not limited to, non-small cell lung cancer, squamous cell carcinoma (epidermoid carcinoma), adenocarcinoma, large-cell carcinoma and small-cell lung cancer; testicular cancers including but not limited to, germinal tumor, seminoma, anaplastic, spermatocytic, nonseminoma, embryonal carcinoma,

5 teratoma carcinoma, choriocarcinoma (yolk-sac tumor); prostate cancers including but not limited to, adenocarcinoma, leiomyosarcoma, and rhabdomyosarcoma; penile cancers; oral cancers including but not limited to, squamous cell carcinoma; basal cancers; salivary gland cancers including but not limited to, adenocarcinoma, mucoepidermoid carcinoma,

10 and adenoidcystic carcinoma; pharynx cancers including but not limited to, squamous cell cancer, and verrucous; skin cancers including but not limited to, basal cell carcinoma, squamous cell carcinoma and melanoma, and superficial spreading melanoma, nodular melanoma, lentigo malignant melanoma, acral lentiginous melanoma; kidney cancers including but not limited to, renal cell cancer, renal cancer, adenocarcinoma, hypernephroma, fibrosarcoma, and transitional cell cancer (renal pelvis and/or uterus); Wilms' tumor; bladder cancers including but not limited to, transitional

15 cell carcinoma, squamous cell cancer, adenocarcinoma, and carcinosarcoma. In addition, cancers include myxosarcoma, osteogenic sarcoma, endothelioma, lymphangioendothelioma, mesothelioma, synovioma, hemangioblastoma, epithelial carcinoma, cystadenocarcinoma, bronchogenic carcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma and papillary adenocarcinomas (for a review of such disorders, see Fishman et al., 1985,

Medicine, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy et al., 1997, Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery, Viking Penguin, Penguin Books U.S.A., Inc., United States of America).

**[0190]** In one embodiment, the cancer is benign, e.g., polyps and benign lesions. In other embodiments, the cancer is metastatic. The Agonistic Therapeutic Agents can be used in the treatment of pre-malignant as well as malignant conditions. Pre-malignant conditions include hyperplasia, metaplasia, and dysplasia. Treatment of malignant conditions includes the treatment of primary as well as metastatic tumors. In a specific embodiment the cancer is melanoma, colon cancer, and lung cancer.

**[0191]** In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a subject suffering from or diagnosed with cancer. In other embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a subject predisposed or susceptible to developing cancer. In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a subject that lives in a region where there is a high occurrence rate of cancer. In a specific embodiment, the cancer is characterized by a pre-malignant tumor or a malignant tumor.

**[0192]** In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a mammal which is 0 to 6 months old, 6 to 12 months old, 1 to 5 years old, 5 to 10 years old, 10 to 15 years old, 15 to 20 years old, 20 to 25 years old, 25 to 30 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old, 50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human at risk developing cancer. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human with cancer. In certain embodiments, the patient is a human 0 to 6 months old, 6 to 12 months old, 1 to 5 years old, 5 to 10 years old, 10 to 15 years old, 15 to 20 years old, 13 to 19 years old, 20 to 25 years old, 25 to 30 years old, 20 to 65 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old, 50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In some embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human infant or a premature human infant. In other embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human child. In other embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human adult. In yet other embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to an elderly human.

**[0193]** In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a pet, e.g., a dog or cat. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a farm animal or livestock, e.g., pig, cows, horses, chickens, etc.

**[0194]** In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a primate, preferably a human, or another mammal, such as a pig, cow, horse, sheep, goat, dog, cat and rodent, in an immunocompromised state or immunosuppressed state or at risk for becoming immunocompromised or immunosuppressed. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject receiving

or recovering from immunosuppressive therapy. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that has or is at risk of getting AIDS, a viral infection, or a bacterial infection. In certain embodiments, a subject that is, will or has undergone surgery, chemotherapy and/or radiation therapy. In some embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that lives in a nursing home, a group home (*i.e.*, a home for 10 or more subjects), or a prison.

**[0195]** In some embodiments, a patient is administered an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is before any adverse effects or intolerance to therapies other than Agonistic Therapeutic Agents develops. In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to refractory patients. In a certain embodiment, refractory patient is a patient refractory to a standard anti-cancer therapy. In certain embodiments, a patient with cancer, is refractory to a therapy when the cancer has not significantly been eradicated and/or the symptoms have not been significantly alleviated. The determination of whether a patient is refractory can be made either *in vivo* or *in vitro* by any method known in the art for assaying the effectiveness of a treatment, using art-accepted meanings of "refractory" in such a context. In various embodiments, a patient with cancer is refractory when a cancerous tumor has not decreased or has increased.

**[0196]** In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a patient to prevent the onset or reoccurrence of cancer in a patient at risk of developing such cancer. In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a patient who are susceptible to adverse reactions to conventional therapies.

**[0197]** In some embodiments, one or more Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a patient who has proven refractory to therapies other than Agonistic Therapeutic Agents, but are no longer on these therapies. In certain embodiments, the patients being managed or treated in accordance with the methods described herein are patients already being treated with antibiotics, anti-cancer agents, or other biological therapy/immunotherapy. Among these patients are refractory patients, patients who are too young for conventional therapies, and patients with reoccurring viral infections despite management or treatment with existing therapies.

**[0198]** In some embodiments, the subject being administered one or more Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies has not received a therapy prior to the administration of the Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies. In other embodiments, one or more Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a subject who has received a therapy prior to administration of one or more Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies. In some embodiments, the subject administered an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent was refractory to a prior therapy or experienced adverse side effects to the prior therapy or the prior therapy was discontinued due to unacceptable levels of toxicity to the subject.

#### 5.4.1.2. Infectious Diseases

**[0199]** The invention provides a method for treating, preventing and/or managing an infectious disease in a subject, comprising administering an effective amount of an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject in need thereof.

**[0200]** In a particular embodiment, the Agonistic Therapeutic Agents are cells infected with a pathogen, wherein the cells are engineered to express IL-15 and IL-15Ra as described in Section 5.1.1, are irradiated, and are administered to a patient to induce and/or enhance an immune response to the infected cells or to antigens of the pathogen. In one embodiment, the pathogen is an intracellular pathogen, *e.g.*, intracellular bacteria or virus. In a specific embodiment, the induced or enhanced immune response is increased production in the patient of antibodies to the infected cells or to the antigens of the pathogen. In one embodiment, the form of irradiation is  $\gamma$ -radiation. IL-15 and IL-15Ra can be expressed from the same construct or different constructs, and under the control of the same promoter or different promoters. The IL-15 and IL-15Ra expression construct(s) can be introduced into cells infected with a pathogen by any means known to those of skill in the art, such as using a viral vector or mammalian expression vector. Alternatively, the native IL-15 and native IL-15Ra of the cells infected with a pathogen may be activated using gene activation methods. Subsequently, the irradiated cells infected with a pathogen recombinantly expressing IL-15/IL-15Ra complexes are administered into a subject to induce and/or enhance an immune response to the pathogen or to antigens of the pathogen. In a specific embodiment, the induced or enhanced immune response is increased production of antibodies to the pathogen. In another embodiment, the induced or enhanced immune response is an increase in effector cell function, *e.g.*, antibody-dependent cellular cytotoxicity (ADCC) against the pathogen and/or cells infected with a pathogen in the

patient. In some embodiments, the induced or enhanced immune response is increase in lymphocyte number, lymphocyte proliferation, and/or lymphocyte activity. In another embodiment, the induced or enhanced immune response is an increase in effector cell function, e.g., cytotoxic cells or antibody-dependent cellular cytotoxicity (ADCC) against the infected cells in the patient.

5 [0201] In other embodiments, an Agonistic Therapeutic Agent can be administered in combination with one or more other therapies. Non-limiting examples of other therapies that can be used in combination with Agonistic Therapeutic Agents are described in Sections 5.4.5.2 and 5.4.5.3. In one embodiment, the combination of an Agonistic Therapeutic Agent and one or more other therapies provides an additive therapeutic effect relative to the therapeutic effects of the Agonistic Therapeutic Agent alone or the one or more other therapies alone. In one embodiment, the combination of an  
10 Agonistic Therapeutic Agent and one or more other therapies provides more than an additive therapeutic effect relative to the therapeutic effects of the Agonistic Therapeutic Agent alone or the one or more other therapies alone. In one embodiment, the combination of an Agonistic Therapeutic Agent and one or more other therapies provides a synergistic therapeutic effect relative to the therapeutic effects of the Agonistic Therapeutic Agent alone or the one or more other therapies alone.

15 [0202] In one embodiment, the one or more therapies include, but are not limited to cytokines/growth factors, e.g., interleukin (IL) 1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, IL-11, IL-12, IL-15, TNF- $\alpha$ , TNF- $\beta$ , TGF- $\beta$ , GM-CSF, and interferon- $\gamma$ . In one embodiment, the one or more therapies include, but are not limited to receptors, antibodies, or other binding agents that bind to cytokines/growth factors, e.g., interleukin (IL) 1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-10, IL-11, IL-12, TNF- $\alpha$ , TNF- $\beta$ , TGF- $\beta$ , GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , and interferon- $\gamma$ . In some embodiments, the  
20 one or more therapies include, but are not limited to, cells recombinantly expressing a therapeutic protein (or polypeptides), e.g., a cytokine, a growth factor, a chemokine, or a fragment or derivative thereof. In a particular embodiment, the one or more therapies include, but are not limited to, cells recombinantly expressing IL-12, IL-6, GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$  or TNF- $\alpha$ . In one embodiment, the one or more therapies are not cells recombinantly expressing IL-12. In one embodiment, the one or more therapies are not cells recombinantly expressing IL-6. In one  
25 embodiment, the one or more therapies are not cells recombinantly expressing GM-CSF. In one embodiment, the one or more therapies are not cells recombinantly expressing TNF- $\alpha$ . In a specific embodiment, the cells recombinantly expressing a therapeutic protein (polypeptide) are cells infected with a pathogen or infectious agent. In a specific embodiment, the cells recombinantly expressing a therapeutic protein (polypeptide) are irradiated cells infected with a pathogen or infectious agent. In a specific embodiment, the cells recombinantly expressing a therapeutic protein (polypeptide) are irradiated cells infected with a pathogen or infectious agent obtained from a patient. In one embodiment, the cells recombinantly express one or more therapeutic proteins (polypeptides). In one embodiment, the present invention provides for a combination of an Agonistic Therapeutic Agent and one or more other therapies, wherein the Agonistic Therapeutic Agent comprises cells recombinantly express IL-15 and IL-15Ra, and wherein the one or more other therapies comprise one or more exogenous cytokines (e.g., IL-12, IL-6, GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$  or TNF- $\alpha$ ). In one embodiment, the present invention provides for a combination of an Agonistic Therapeutic Agent and one or  
30 more other therapies, wherein the Agonistic Therapeutic Agent comprises cells recombinantly express IL-15 and IL-15Ra, and wherein the one or more other therapies comprise (i) exogenous IL-15 polypeptide or exogenous IL-15R polypeptide, and (ii) one or more other exogenous cytokines (e.g., IL-12, IL-6, GM-CSF, interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$  or TNF- $\alpha$ ). In one embodiment, an Agonistic Therapeutic Agent comprises cells engineered to recombinantly express (i) IL-15, IL-15Ra, or IL-15 and IL-15Ra; and (ii) one or more cytokines/growth factors.

35 [0203] Infectious diseases that can be treated, prevented, and/or managed by Agonistic Therapeutic Agents are caused by infectious agents including but not limited to bacteria, fungi, protozoae, and viruses.

40 [0204] Viral diseases that can be prevented, treated and/or managed in accordance with the methods described herein include, but are not limited to, those caused by hepatitis type A, hepatitis type B, hepatitis type C, influenza, varicella, adenovirus, herpes simplex type I (HSV-I), herpes simplex type II (HSV-II), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, hantavirus, coxsackie virus, mumps virus, measles virus, rubella virus, polio virus, small pox, Epstein Barr virus, human immunodeficiency virus type I (HIV-I), human immunodeficiency virus type II (HIV-II), and agents of viral diseases such as viral meningitis, encephalitis, dengue or small pox.

45 [0205] Bacterial diseases caused by bacteria (e.g., *Escherichia coli*, *Klebsiella pneumoniae*, *Staphylococcus aureus*, *Enterococcus faecalis*, *Candida albicans*, *Proteus vulgaris*, *Staphylococcus viridans*, and *Pseudomonas aeruginosa*) that can be prevented, treated and/or managed in accordance with the methods described herein include, but are not limited to, *mycobacteria rickettsia*, *mycoplasma*, *neisseria*, *S. pneumonia*, *Borrelia burgdorferi* (Lyme disease), *Bacillus antracis* (anthrax), *tetanus*, *streptococcus*, *staphylococcus*, *mycobacterium*, *pertussis*, *cholera*, *plague*, *diphtheria*, *chlamydia*, *S. aureus* and *legionella*.

50 [0206] Protozoal diseases caused by protozoa that can be prevented, treated and/or managed in accordance with the methods described herein include, but are not limited to, leishmania, kokzidioa, trypanosoma or malaria.

55 [0207] Parasitic diseases caused by parasites that can be prevented, treated and/or managed in accordance with the

methods described herein include, but are not limited to, chlamydia and rickettsia.

**[0208]** In certain embodiments, administering an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject (in some embodiments, an animal model) infected with an infectious agent inhibits or reduces replication of the infectious agent by at least 20% to 25%, preferably at least 25% to 30%, at least 30% to 35%, at least 35% to 40%, at least 40% to 45%, at least 45% to 50%, at least 50% to 55%, at least 55% to 60%, at least 60% to 65%, at least 65% to 70%, at least 70% to 75%, at least 75% to 80%, or up to at least 85% relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In some embodiments, administering an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject (in some embodiments, an animal model) infected with an infectious agent inhibits or reduces replication of the infectious agent by at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 8 fold, 10 fold, 15 fold, 20 fold, or 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In other embodiments, administering an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject (in some embodiments, an animal model) infected with an infectious agent inhibits or reduces replication of the infectious agent by 1 log, 1.5 logs, 2 logs, 2.5 logs, 3 logs, 3.5 logs, 4 logs, 5 logs or more relative to a negative control as determined using an assay described herein or others known to one of skill in the art.

**[0209]** In certain embodiments, administering an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject (in some embodiments, an animal model) infected with an infectious agent reduces the titer of the infectious agent by at least 20% to 25%, preferably at least 25% to 30%, at least 30% to 35%, at least 35% to 40%, at least 40% to 45%, at least 45% to 50%, at least 50% to 55%, at least 55% to 60%, at least 60% to 65%, at least 65% to 70%, at least 70% to 75%, at least 75% to 80%, or up to at least 85% relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In some embodiments, administering an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject (in some embodiments, an animal model) infected with an infectious agent reduces the titer of the infectious agent by at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 8 fold, 10 fold, 15 fold, 20 fold, or 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In other embodiments, administering an Agonistic Therapeutic Agent or a composition comprising an Agonistic Therapeutic Agent to a subject (in some embodiments, an animal model) infected with an infectious agent reduces the titer of the infectious agent by 1 log, 1.5 logs, 2 logs, 2.5 logs, 3 logs, 3.5 logs, 4 logs, 5 logs or more relative to a negative control as determined using an assay described herein or others known to one of skill in the art.

**[0210]** In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a subject suffering from an infectious disease caused by infectious agents including, but not limited to bacteria, fungi, protozoae, and viruses. In other embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a subject predisposed or susceptible to an infectious disease. In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a subject that lives in a region where there has been or might be an outbreak with infections by infectious agents. In some embodiments, the infection is a latent infection. In other embodiments, the infection by the infectious agent is an active infection. In yet other embodiments, the infection by the infectious agent is a chronic viral infection. In a specific embodiment, the infection is a viral infection. In a specific embodiment, the virus infects humans.

**[0211]** In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a mammal which is 0 to 6 months old, 6 to 12 months old, 1 to 5 years old, 5 to 10 years old, 10 to 15 years old, 15 to 20 years old, 20 to 25 years old, 25 to 30 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old, 50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human at risk for a virus infection. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human with a virus infection. In certain embodiments, the patient is a human 0 to 6 months old, 6 to 12 months old, 1 to 5 years old, 5 to 10 years old, 5 to 12 years old, 10 to 15 years old, 15 to 20 years old, 13 to 19 years old, 20 to 25 years old, 25 to 30 years old, 20 to 65 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old, 50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In some embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human infant or premature human infant. In other embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human child. In other embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a human adult. In yet other embodiments, an Agonistic Therapeutic Agent,

composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to an elderly human.

**[0212]** In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a pet, e.g., a dog or cat. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a farm animal or livestock, e.g., pig, cows, horses, chickens, etc. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a bird, e.g., ducks or chicken.

**[0213]** In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a primate, preferably a human, or another mammal, such as a pig, cow, horse, sheep, goat, dog, cat and rodent, in an immunocompromised state or immunosuppressed state or at risk for becoming immunocompromised or immunosuppressed. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject receiving or recovering from immunosuppressive therapy. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that has or is at risk of getting cancer, AIDS, another infection, or a bacterial infection. In certain embodiments, a subject that is, will or has undergone surgery, chemotherapy and/or radiation therapy. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that has cystic fibrosis, pulmonary fibrosis, or another disease which makes the subject susceptible to an infection. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that has, will have or had a tissue transplant. In some embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that lives in a nursing home, a group home (*i.e.*, a home for 10 or more subjects), or a prison. In some embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that attends school (*e.g.*, elementary school, middle school, junior high school, high school or university) or daycare. In some embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that works in the healthcare area, such as a doctor or a nurse, or in a hospital. In certain embodiments, an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy is administered to a subject that is pregnant or will become pregnant.

**[0214]** In some embodiments, a patient is administered an Agonistic Therapeutic Agent, composition comprising an Agonistic Therapeutic Agent, or a combination therapy before any adverse effects or intolerance to therapies other than Agonistic Therapeutic Agents develops. In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to refractory patients. In a certain embodiment, refractory patient is a patient refractory to a standard therapy. In certain embodiments, a patient with an infection, is refractory to a therapy when the infection has not significantly been eradicated and/or the symptoms have not been significantly alleviated. The determination of whether a patient is refractory can be made either *in vivo* or *in vitro* by any method known in the art for assaying the effectiveness of a treatment of infections, using art-accepted meanings of "refractory" in such a context. In various embodiments, a patient with an infection is refractory when replication of the infectious agent has not decreased or has increased.

**[0215]** In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a patient to prevent the onset or reoccurrence of infections (*e.g.*, viral infections) in a patient at risk of developing such infections. In some embodiments, Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a patient who are susceptible to adverse reactions to conventional therapies.

**[0216]** In some embodiments, one or more Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a patient who has proven refractory to therapies other than Agonistic Therapeutic Agents, but are no longer on these therapies. In certain embodiments, the patients being managed or treated in accordance with the methods of this invention are patients already being treated with antibiotics, anti-virals, anti-fungals, or other biological therapy/immunotherapy. Among these patients are refractory patients, patients who are too young for conventional therapies, and patients with reoccurring viral infections despite management or treatment with existing therapies.

**[0217]** In some embodiments, the subject being administered one or more Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies has not received a therapy prior to the administration of the Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies.

**55** In other embodiments, one or more Agonistic Therapeutic Agents, compositions comprising Agonistic Therapeutic Agents, or combination therapies are administered to a subject who has received a therapy prior to administration of one or more Agonistic Therapeutic Agents or compositions comprising one or more Agonistic Therapeutic Agents, or combination therapies. In some embodiments, the subject administered an Agonistic Therapeutic Agent or a composition

comprising an Agonistic Therapeutic Agent was refractory to a prior therapy or experienced adverse side effects to the prior therapy or the prior therapy was discontinued due to unacceptable levels of toxicity to the subject.

#### 5.4.2. Suppressing Immune Function

5

**[0218]** The invention provides methods for suppressing the IL-15 mediated immune function in a subject, comprising administering an Antagonistic Therapeutic Agent or a composition comprising an Antagonistic Therapeutic Agent to a subject in need thereof. In a specific embodiment, the invention provides methods for preventing, treating, and/or managing diseases in which it is desirable to suppress immune function, comprising administering an Antagonistic Therapeutic Agent or a composition comprising an Antagonistic Therapeutic Agent to a subject in need thereof. In specific embodiments, the Antagonistic Therapeutic Agents are formulated with polymers described in Section 5.2, *supra*. In other embodiments, an Antagonistic Therapeutic Agent can be administered in combination with one or more other therapies to suppress an IL-15 mediated immune function. Non-limiting examples of diseases that can be prevented, treated, or managed by suppressing immune function include, but are not limited to, autoimmune disease, inflammatory disorders, graft versus host disease, and transplant rejection.

10

**[0219]** In a specific embodiment, an Antagonistic Therapeutic Agent suppresses or reduces IL-15 mediated immune function, e.g., NK cell proliferation and cytokine production.

15

**[0220]** In a specific embodiment, an Antagonistic Therapeutic Agent suppresses IL-15 mediated immune function in a subject (in some embodiments, an animal model) by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the immune function in a subject not administered the Therapeutic Agent using assays well known in the art, e.g., ELISPOT, ELISA, and cell proliferation assays. In a specific embodiment, the immune function is cytokine release or production (e.g., interferon-gamma). In one embodiment, the IL-15 mediated immune function is NK cell proliferation, which can be assayed, e.g., by flow cytometry to detect the number of cells expressing markers of NK cells (e.g., CD56). In one embodiment, the IL-15 mediated immune function is T cell (CD4<sup>+</sup> cell and/or CD8<sup>+</sup> cell) proliferation and/or activation.

20

**[0221]** In specific embodiments, an Antagonistic Therapeutic Agent can inhibit or reduce proliferation/expansion of lymphocytes (e.g., number of lymphocytes), induce or increase of apoptosis of lymphocytes, inhibit activation of dendritic cells (or antigen presenting cells), and antigen presentation. In particular embodiments, an immune function suppressed by an Antagonistic Therapeutic Agent is proliferation/expansion/activation of CD4<sup>+</sup> T cells (e.g., Th1 and Th2 helper T cells), CD8<sup>+</sup> T cells (e.g., cytotoxic T lymphocytes, alpha/beta T cells, and gamma/delta T cells), B cells (e.g., plasma cells), memory T cells, memory B cells, dendritic cells (immature or mature), antigen presenting cells, macrophages, mast cells, or natural killer cells. In one embodiment, an Antagonistic Therapeutic Agent decreases/reduces the proliferation/expansion or number of lymphocyte progenitors.

25

**[0222]** In a specific embodiment, an Antagonistic Therapeutic Agent suppresses IL-15 mediated immune function in a cell-based assay (e.g., assaying proliferation of an IL-15-responsive cell line such as CTLL-2 or TF $\beta$ 1) by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to a negative control as determined using assays well known in the art, e.g., cell proliferation assays with CSFE, BrdU, and <sup>3</sup>H-Thymidine incorporation. In a specific embodiment, the immune function is cytokine release (e.g., interferon-gamma). In particular embodiments, the activity of an Antagonistic Therapeutic Agent in cell-based assays and in animal model experiments correlate with the *in vivo* function of the Antagonistic Therapeutic Agent to suppress IL-15 mediated immune function.

30

**[0223]** Various autoimmune and inflammatory diseases that can be prevented, treated and/or managed are listed below.

35

##### 5.4.2.1. Autoimmune and Inflammatory Disorders

40

**[0224]** The invention provides a method for treating, preventing and/or managing an autoimmune disorder or inflammatory disorder in a subject, comprising administering an effective amount of an Antagonistic Therapeutic Agent or a composition comprising an Antagonistic Therapeutic Agent to a subject in need thereof. The invention also provides a method for reducing inflammation in a subject, comprising administering an effective amount of an Antagonistic Therapeutic Agent or a composition comprising an Antagonistic Therapeutic Agent to a subject in need thereof. Non-limiting examples of autoimmune disorders and inflammatory disorders include transplant rejection and graft versus host disease (GVHD). GVHD occurs when a donor's immune cells (e.g., donor's T cells) attack cells in the recipient subject's body. Transplant rejection occurs when a transplanted organ or tissue fails to be accepted by the body of the transplant recipient. In general, the transplant rejection is due to the immune system of the recipient (e.g., recipient's T cells) attacking the transplanted organ or tissue.

[0225] In a specific embodiment, administering an Antagonistic Therapeutic Agent or composition comprising an Antagonistic Therapeutic Agent to a subject (in some embodiments, an animal model) reduces the inflammation in an subject by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the inflammation in an subject not administered the Antagonistic Therapeutic Agent using methods known in the art. For example, reduction in inflammation can be measured by the reduction in cytokine secretion (e.g., tumor necrosis factor alpha, interferon gamma). In a specific embodiment, administering an Antagonistic Therapeutic Agent or composition comprising an Antagonistic Therapeutic Agent to a subject (in some embodiments, an animal model) reduces the inflammation in an subject by at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 8 fold, 10 fold, 15 fold, 20 fold, or 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to the inflammation in an subject not administered the Antagonistic Therapeutic Agent using methods known in the art. For example, reduction in inflammation can be measured by the reduction cytokine secretion (e.g., tumor necrosis factor alpha, interferon gamma).

[0226] In other embodiments, an Antagonistic Therapeutic Agent can be administered in combination with one or more other therapies to suppress IL-15 mediated immune function in a subject. Various anti-inflammatory agents known in the art can be used in combination with Antagonistic Therapeutic Agents.

[0227] Examples of autoimmune disorders that can be prevented, treated and/or managed in accordance with the methods described herein include, but are not limited to, celiac (coeliac disease), alopecia areata, ankylosing spondylitis, antiphospholipid syndrome, autoimmune Addison's disease, autoimmune diseases of the adrenal gland, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune oophoritis and orchitis, autoimmune thrombocytopenia, Behcet's disease, bullous pemphigoid, cardiomyopathy, celiac spruadermatitis, chronic fatigue immune dysfunction syndrome (CFIDS), chronic inflammatory demyelinating polyneuropathy, Churg-Strauss syndrome, cicatrical pemphigoid, CREST syndrome, cold agglutinin disease, Crohn's disease, discoid lupus, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, glomerulonephritis, Graves' disease, Guillain-Barre, Hashimoto's thyroiditis, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), IgA neuropathy, juvenile arthritis, lichen planus, lupus erythematosus, Ménière's disease, mixed connective tissue disease, multiple sclerosis, type 1 or immune-mediated diabetes mellitus, myasthenia gravis, pemphigus vulgaris, pernicious anemia, polyarteritis nodosa, polychondritis, polyglandular syndromes, polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, psoriatic arthritis, Raynauld's phenomenon, Reiter's syndrome, Rheumatoid arthritis, sarcoidosis, scleroderma, Sjögren's syndrome, stiff-man syndrome, systemic lupus erythematosus, lupus erythematosus, takayasu arteritis, temporal arteritis/ giant cell arteritis, ulcerative colitis, uveitis, vasculitides such as dermatitis herpetiformis vasculitis, vitiligo, and Wegener's granulomatosis. Examples of inflammatory disorders include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthropathy, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections. Some autoimmune disorders are associated with an inflammatory condition. Thus, there is overlap between what is considered an autoimmune disorder and an inflammatory disorder. Therefore, some autoimmune disorders may also be characterized as inflammatory disorders. Examples of inflammatory disorders which can be prevented, treated or managed in accordance with the methods described herein include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthropathy, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections.

[0228] In some embodiments, Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies are administered to a subject suffering from an autoimmune disease or inflammatory disorder. In other embodiments, Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies are administered to a subject predisposed or susceptible to developing an autoimmune disease or inflammatory disorder.

[0229] In certain embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a mammal which is 0 to 6 months old, 6 to 12 months old, 1 to 5 years old, 5 to 10 years old, 10 to 15 years old, 15 to 20 years old, 20 to 25 years old, 25 to 30 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old, 50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In certain embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a human at risk developing an autoimmune disease or inflammatory disorder. In certain embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a human with an autoimmune disease or inflammatory disorder. In certain embodiments, the patient is a human 0 to 6 months old, 6 to 12 months old, 1 to 5 years old, 5 to 10 years old, 5 to 12 years old, 10 to 15 years old, 15 to 20 years old, 13 to 19 years old, 20 to 25 years old, 25 to 30 years old, 20 to 65 years old, 30 to 35 years old, 35 to 40 years old, 40 to 45 years old, 45 to 50 years old,

50 to 55 years old, 55 to 60 years old, 60 to 65 years old, 65 to 70 years old, 70 to 75 years old, 75 to 80 years old, 80 to 85 years old, 85 to 90 years old, 90 to 95 years old or 95 to 100 years old. In some embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a human infant or premature human infant. In other embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a human child. In other embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a human adult. In yet other embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to an elderly human.

5 [0230] In certain embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a pet, e.g., a dog or cat. In certain embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a farm animal or livestock, e.g., pig, cows, horses, chickens, etc.

10 [0231] In certain embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a primate, preferably a human, or another mammal, such as a pig, cow, horse, sheep, goat, dog, cat and rodent, in an immunocompromised state or immunosuppressed state or at 15 risk for becoming immunocompromised or immunosuppressed. In certain embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a subject receiving or recovering from immunosuppressive therapy. In certain embodiments, an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy is administered to a subject that 20 has or is at risk of getting AIDS, a viral infection, or a bacterial infection. In certain embodiments, a subject that is, will or has undergone surgery, chemotherapy and/or radiation therapy.

25 [0232] In some embodiments, a patient is administered an Antagonistic Therapeutic Agent, composition comprising an Antagonistic Therapeutic Agent, or a combination therapy before any adverse effects or intolerance to therapies other than Antagonistic Therapeutic Agents develops. In some embodiments, Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies are administered to refractory patients. In a certain embodiment, refractory patient is a patient refractory to a standard therapy. In certain embodiments, a patient with an 30 autoimmune disease or inflammatory disorder, is refractory to a therapy when the autoimmune disease or inflammatory disorder, respectively, has not significantly been eradicated and/or the symptoms have not been significantly alleviated. The determination of whether a patient is refractory can be made either *in vivo* or *in vitro* by any method known in the art for assaying the effectiveness of a treatment, using art-accepted meanings of "refractory" in such a context. In various embodiments, a patient with a inflammatory disorder is refractory when inflammation has not decreased or has increased.

35 [0233] In some embodiments, Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies are administered to a patient who are susceptible to adverse reactions to conventional therapies.

40 [0234] In some embodiments, one or more Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies are administered to a patient who has proven refractory to therapies other than Antagonistic Therapeutic Agents, but are no longer on these therapies. In certain embodiments, the patients being managed or treated in accordance with the methods described herein are patients already being treated with antibiotics, anti-cancer agents, anti-inflammatory agents, or other biological therapy/immunotherapy. Among these patients are refractory patients, patients who are too young for conventional therapies.

45 [0235] In some embodiments, the subject being administered one or more Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies has not received a therapy prior to the administration of the Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies. In other embodiments, one or more Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies are administered to a subject who has received a therapy prior to administration of one or more Antagonistic Therapeutic Agents, compositions comprising Antagonistic Therapeutic Agents, or combination therapies. In some embodiments, the subject administered an Antagonistic Therapeutic Agent or a composition comprising an Antagonistic Therapeutic Agent was refractory to a prior therapy or experienced 50 adverse side effects to the prior therapy or the prior therapy was discontinued due to unacceptable levels of toxicity to the subject.

#### 5.4.3. Mode of Administration

55 [0236] Therapeutic Agents can be administered via any route known in the art. In a specific embodiment, Therapeutic Agents formulated with polymers are especially suited for local delivery, but such formulations can also be for systemic administration.

[0237] Therapeutic Agents or compositions thereof can be administered orally, or by any other convenient route, for example, by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa,

rectal, and intestinal mucosa) and may be administered together with another biologically active agent. Administration can be systemic or local. Various delivery systems are known, e.g., encapsulation in liposomes, microparticles, micro-capsules, capsules, and can be used to deliver the Therapeutic Agents or compositions thereof and pharmaceutically acceptable salts thereof.

5      **[0238]** Methods of administration include but are not limited to parenteral, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, oral, sublingual, intranasal, intracerebral, intravaginal, transdermal, rectally, by inhalation, intratumoral, or topically, particularly to the ears, nose, eyes, or skin. The mode of administration is left to the discretion of the practitioner. In some embodiments, administration will result in the release of a Therapeutic Agent into the bloodstream.

10     **[0239]** In specific embodiments, it may be desirable to administer a Therapeutic Agent locally. This may be achieved, for example, and not by way of limitation, by local infusion, topical application, e.g., in conjunction with a wound dressing, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers.

15     **[0240]** In certain embodiments, it may be desirable to introduce a Therapeutic Agent into the central nervous system by any suitable route, including intraventricular, intrathecal and epidural injection. Intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir.

20     **[0241]** Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent, or via perfusion in a fluorocarbon or synthetic pulmonary surfactant.

25     **[0242]** In certain embodiments, a Therapeutic Agent is formulated as a suppository, with traditional binders and vehicles such as triglycerides.

30     **[0243]** For viral infections or melanoma with cutaneous manifestations, the Therapeutic Agent can be administered topically.

35     **[0244]** In another embodiment, a Therapeutic Agent is delivered in a vesicle, in particular a liposome (See Langer, 1990, *Science* 249:1527 1533; Treat et al., in *Liposomes in the Therapy of Infectious Disease and Bacterial infection*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353 365 (1989); Lopez Berestein, *ibid.*, pp. 317 327).

40     **[0245]** In another embodiment, a Therapeutic Agent is delivered in a controlled release system (See, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115 138 (1984)). Examples of controlled-release systems are discussed in the review by Langer, 1990, *Science* 249:1527 1533 may be used. In one embodiment, a pump may be used (See Langer, *supra*; Sefton, 1987, *CRC Crit. Ref. Biomed. Eng.* 14:201; Buchwald et al., 1980, *Surgery* 88:507; Saudek et al., 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used (See *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, *J. Macromol. Sci. Rev. Macromol. Chem.* 23:61; See also Levy et al., 1985, *Science* 228:190; During et al., 1989, *Ann. Neurol.* 25:351; Howard et al., 1989, *J. Neurosurg.* 71:105). In a specific embodiment, a controlled-release system comprising a Therapeutic Agent is placed in close proximity to the tissue affected by the disease to be prevented, treated and/or managed. In accordance with this embodiment, the close proximity of the controlled-release system to the affected tissue may result in only a fraction of the dose of the Therapeutic Agent required if it is systemically administered.

45     **[0246]** In a specific embodiment, the Therapeutic Agent formulated with a polymer is administered locally to the area or tissue of a patient for enhancement or reduction of IL-15 function. In some embodiments, an Agonistic Therapeutic Agent formulated with polymers can be administered locally to a tumor in a cancer patient to enhance or induce IL-15 function and immune response to the tumor. In other embodiments, an Agonistic Therapeutic Agent formulated with polymers can be administered locally to a tissue infected with a pathogen in a subject to enhance or induce IL-15 function and an immune response to the pathogen. In certain embodiments, the Agonistic Therapeutic Agent includes the IL-15/IL-15Ra complex or nucleic acids encoding IL-15 and IL-15Ra. In other embodiments, the Agonistic Therapeutic Agent includes cells expressing IL-15 and IL-15Ra in high amounts and polymers. In one embodiment, high amounts of IL-15/IL-15Ra complexes refer to amounts of IL-15/IL-15Ra complexes expressed by cells that are at least 1 fold, 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 20 fold, or more than 20 fold higher than amounts of IL-15/IL-15Ra complexes expressed endogenously by control cells (e.g., cells that have not been genetically engineered to recombinantly express IL-15, IL-15Ra, or both IL-15 and IL-15Ra, or cells comprising an empty vector).

50     **[0247]** In some embodiments, an Antagonistic Therapeutic Agent formulated with polymers is administered locally to a site of inflammation in a subject suffering from an autoimmune or inflammatory disorder to suppress or reduce IL-15 function and the immune response elicited at the site of inflammation. In other embodiments, an Antagonistic Therapeutic Agent formulated with polymers is administered locally to a site of a transplanted tissue/organ in a subject to suppress or reduce IL-15 function and immune response to the transplanted tissue.

#### 5.4.4. Dosages & Frequency of Administration

**[0248]** The amount of a Therapeutic Agent, or the amount of a composition comprising a Therapeutic Agent, that will be effective in the prevention, treatment and/or management of a disease that is affected by IL-15 function can be determined by standard clinical techniques. *In vitro* or *in vivo* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed will also depend, e.g., on the route of administration, the type of symptoms, and the seriousness of the symptoms, and should be decided according to the judgment of the practitioner and each patient's or subject's circumstances.

**[0249]** In some embodiments, the dosage of a Therapeutic Agent or compositions thereof is determined by extrapolating from the no observed adverse effective level (NOAEL), as determined in animal studies. This extrapolated dosage is useful in determining the maximum recommended starting dose for human clinical trials. For instance, the NOAELs can be extrapolated to determine human equivalent dosages (HED). Typically, HED is extrapolated from a non-human animal dosage based on the doses that are normalized to body surface area (*i.e.*, mg/m<sup>2</sup>). In specific embodiments, the NOAELs are determined in mice, hamsters, rats, ferrets, guinea pigs, rabbits, dogs, primates, primates (monkeys, marmosets, squirrel monkeys, baboons), micropigs or minipigs. For a discussion on the use of NOAELs and their extrapolation to determine human equivalent doses, See Guidance for Industry Estimating the Maximum Safe Starting Dose in Initial Clinical Trials for Therapeutics in Adult Healthy Volunteers, U.S. Department of Health and Human Services Food and Drug Administration Center for Drug Evaluation and Research (CDER), Pharmacology and Toxicology, July 2005. In one embodiment, a Therapeutic Agent or composition thereof is administered at a dose that is lower than the human equivalent dosage (HED) of the NOAEL over a period of 1 week, 2 weeks, 3 weeks, 1 month, 2 months, three months, four months, six months, nine months, 1 year, 2 years, 3 years, 4 years or more.

**[0250]** In certain embodiments, a dosage regime for a human subject can be extrapolated from animal model studies using the dose at which 10% of the animals die (LD10). In general the starting dose of a Phase I clinical trial is based on preclinical testing. A standard measure of toxicity of a drug in preclinical testing is the percentage of animals that die because of treatment. It is well within the skill of the art to correlate the LD 10 in an animal study with the maximal-tolerated dose (MTD) in humans, adjusted for body surface area, as a basis to extrapolate a starting human dose. In some embodiments, the interrelationship of dosages for one animal model can be converted for use in another animal, including humans, using conversion factors (based on milligrams per meter squared of body surface) as described, e.g., in Freireich et al., Cancer Chemother. Rep., 1966, 50:219-244. Body surface area may be approximately determined from height and weight of the patient. See, e.g., Scientific Tables, Geigy Pharmaceuticals, Ardley, N. Y., 1970, 537. In certain embodiments, the adjustment for body surface area includes host factors such as, for example, surface area, weight, metabolism, tissue distribution, absorption rate, and excretion rate. In addition, the route of administration, excipient usage, and the specific disease or virus to target are also factors to consider. In one embodiment, the standard conservative starting dose is about 1/10 the murine LD10, although it may be even lower if other species (*i.e.*, dogs) were more sensitive to the Therapeutic Agent. In other embodiments, the standard conservative starting dose is about 1/100, 1/95, 1/90, 1/85, 1/80, 1/75, 1/70, 1/65, 1/60, 1/55, 1/50, 1/45, 1/40, 1/35, 1/30, 1/25, 1/20, 1/15, 2/10, 3/10, 4/10, or 5/10 of the murine LD10. In other embodiments, an starting dose amount of a Therapeutic Agent in a human is lower than the dose extrapolated from animal model studies. In another embodiment, an starting dose amount of a Therapeutic Agent in a human is higher than the dose extrapolated from animal model studies. It is well within the skill of the art to start doses of the active composition at relatively low levels, and increase or decrease the dosage as necessary to achieve the desired effect with minimal toxicity.

**[0251]** Exemplary doses of Therapeutic Agents comprising polypeptides or antibodies or compositions thereof include milligram or microgram amounts per kilogram of subject or sample weight (e.g., about 1 microgram per kilogram to about 500 milligrams per kilogram, about 5 micrograms per kilogram to about 100 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram). In specific embodiments, a daily dose is at least 50 mg, 75 mg, 100 mg, 150 mg, 250 mg, 500 mg, 750 mg, or at least 1 g.

**[0252]** In one embodiment, the dosage is a concentration of 0.01 to 5000 mM, 1 to 300 mM, 10 to 100 mM and 10 mM to 1 M. In another embodiment, the dosage is a concentration of at least 5  $\mu$ M, at least 10  $\mu$ M, at least 50  $\mu$ M, at least 100  $\mu$ M, at least 500  $\mu$ M, at least 1 mM, at least 5 mM, at least 10 mM, at least 50 mM, at least 100 mM, or at least 500 mM.

**[0253]** In one embodiment, the dosage is a concentration of 0.01 to 5000 mM, 1 to 300 mM, 10 to 100 mM and 10 mM to 1 M. In another embodiment, the dosage is a concentration of at least 5  $\mu$ M, at least 10  $\mu$ M, at least 50  $\mu$ M, at least 100  $\mu$ M, at least 500  $\mu$ M, at least 1 mM, at least 5 mM, at least 10 mM, at least 50 mM, at least 100 mM, or at least 500 mM. In a specific embodiment, the dosage is 0.25  $\mu$ g/kg or more, preferably 0.5  $\mu$ g/kg or more, 1  $\mu$ g/kg or more, 2  $\mu$ g/kg or more, 3  $\mu$ g/kg or more, 4  $\mu$ g/kg or more, 5  $\mu$ g/kg or more, 6  $\mu$ g/kg or more, 7  $\mu$ g/kg or more, 8  $\mu$ g/kg or more, 9  $\mu$ g/kg or more, or 10  $\mu$ g/kg or more, 25  $\mu$ g/kg or more, preferably 50  $\mu$ g/kg or more, 100  $\mu$ g/kg or more, 250  $\mu$ g/kg or more, 500  $\mu$ g/kg or more, 1 mg/kg or more, 5 mg/kg or more, 6 mg/kg or more, 7 mg/kg or more, 8 mg/kg or more, 9 mg/kg or more, or 10 mg/kg or more of a patient's body weight.

[0254] In another embodiment, the dosage is a unit dose of 5 mg, preferably 10 mg, 50 mg, 100 mg, 150 mg, 200 mg, 250 mg, 300 mg, 350 mg, 400 mg, 500 mg, 550 mg, 600 mg, 650 mg, 700 mg, 750 mg, 800 mg or more. In another embodiment, the dosage is a unit dose that ranges from about 5 mg to about 100 mg, about 100 mg to about 200  $\mu$ g, about 150 mg to about 300 mg, about 150 mg to about 400 mg, 250  $\mu$ g to about 500 mg, about 500 mg to about 800 mg, about 500 mg to about 1000 mg, or about 5 mg to about 1000 mg.

[0255] Exemplary doses of Therapeutic Agents comprising nucleic acids or compositions thereof include 0.1  $\mu$ g, 0.5  $\mu$ g, 1  $\mu$ g, 1.5  $\mu$ g, 2  $\mu$ g, 3  $\mu$ g, 4  $\mu$ g, 5  $\mu$ g, 6  $\mu$ g, 7  $\mu$ g, 8  $\mu$ g, 9  $\mu$ g, 10  $\mu$ g, 15  $\mu$ g, 20  $\mu$ g, 25  $\mu$ g, 30  $\mu$ g, 35  $\mu$ g, 40  $\mu$ g, 45  $\mu$ g, 50  $\mu$ g, or 60  $\mu$ g of nucleic acids per dose. In a specific embodiment, the dose is in the range of 10 ng to 100 mg, or 50 ng to 100 mg, or 100 ng to 100 mg of nucleic acids per dose. In some specific embodiments, the dose is in the range of 10 pg to 100 mg, or 50 pg to 100 mg, or 100 pg to 100 mg, or 100 pg to 100 ng of nucleic acids per dose.

[0256] In certain embodiments, suitable dosage ranges for oral administration are about 0.001 milligram to about 500 milligrams of a Therapeutic Agent, per kilogram body weight per day. In specific embodiments, the oral dose is about 0.01 milligram to about 100 milligrams per kilogram body weight per day, about 0.1 milligram to about 75 milligrams per kilogram body weight per day or about 0.5 milligram to 5 milligrams per kilogram body weight per day. The dosage amounts described herein refer to total amounts administered; that is, if more than one Therapeutic Agent is administered, then, in some embodiments, the dosages correspond to the total amount administered. In a specific embodiment, oral compositions contain about 10% to about 95% a Therapeutic Agent by weight.

[0257] Suitable dosage ranges for intravenous (i.v.) administration are about 0.01 milligram to about 100 milligrams per kilogram body weight per day, about 0.1 milligram to about 35 milligrams per kilogram body weight per day, and about 1 milligram to about 10 milligrams per kilogram body weight per day. In some embodiments, suitable dosage ranges for intranasal administration are about 0.01 pg/kg body weight per day to about 1 mg/kg body weight per day. Suppositories generally contain about 0.01 milligram to about 50 milligrams of a Therapeutic Agent per kilogram body weight per day and comprise active ingredient in the range of about 0.5% to about 10% by weight.

[0258] Recommended dosages for intradermal, intramuscular, intraperitoneal, subcutaneous, epidural, sublingual, intracerebral, intravaginal, transdermal administration or administration by inhalation are in the range of about 0.001 milligram to about 500 milligrams per kilogram of body weight per day. Suitable doses for topical administration include doses that are in the range of about 0.001 milligram to about 50 milligrams, depending on the area of administration.

[0259] In another embodiment, a subject is administered one or more doses of a prophylactically or therapeutically effective amount of a Therapeutic Agent or a composition thereof, wherein the prophylactically or therapeutically effective amount is not the same for each dose. In another embodiment, a subject is administered one or more doses of a prophylactically or therapeutically effective amount of a Therapeutic Agent or a composition thereof, wherein the dose of a prophylactically or therapeutically effective amount administered to said subject is increased by, e.g., 0.01  $\mu$ g/kg, 0.02  $\mu$ g/kg, 0.04  $\mu$ g/kg, 0.05  $\mu$ g/kg, 0.06  $\mu$ g/kg, 0.08  $\mu$ g/kg, 0.1  $\mu$ g/kg, 0.2  $\mu$ g/kg, 0.25  $\mu$ g/kg, 0.5  $\mu$ g/kg, 0.75  $\mu$ g/kg, 1  $\mu$ g/kg, 1.5  $\mu$ g/kg, 2  $\mu$ g/kg, 4  $\mu$ g/kg, 5  $\mu$ g/kg, 10  $\mu$ g/kg, 15  $\mu$ g/kg, 20  $\mu$ g/kg, 25  $\mu$ g/kg, 30  $\mu$ g/kg, 35  $\mu$ g/kg, 40  $\mu$ g/kg, 45  $\mu$ g/kg, or 50  $\mu$ g/kg, as treatment progresses. In another embodiment, a subject is administered one or more doses of a prophylactically or therapeutically effective amount of a Therapeutic Agent or composition thereof, wherein the dose is decreased by, e.g., 0.01  $\mu$ g/kg, 0.02  $\mu$ g/kg, 0.04  $\mu$ g/kg, 0.05  $\mu$ g/kg, 0.06  $\mu$ g/kg, 0.08  $\mu$ g/kg, 0.1  $\mu$ g/kg, 0.2  $\mu$ g/kg, 0.25  $\mu$ g/kg, 0.5  $\mu$ g/kg, 0.75  $\mu$ g/kg, 1  $\mu$ g/kg, 1.5  $\mu$ g/kg, 2  $\mu$ g/kg, 4  $\mu$ g/kg, 5  $\mu$ g/kg, 10  $\mu$ g/kg, 15  $\mu$ g/kg, 20  $\mu$ g/kg, 25  $\mu$ g/kg, 30  $\mu$ g/kg, 35  $\mu$ g/kg, 40  $\mu$ g/kg, 45  $\mu$ g/kg, or 50  $\mu$ g/kg, as treatment progresses.

[0260] For Therapeutic Agents comprising cells expressing IL-15 and IL-15Ra in high amounts, the suitable dosage range for administration by any route of administration can be at least 100, 200, 300, 400, 500, 700, 1,000, 5,000, 10,000, 25,000, 50,000, or 100,000 cells. In specific embodiments, the number of cells is at least 100, 200, 300, 400, 500 cells. In other embodiments, the number of cells is at least 300, 400, 500, 700, 1,000 cells. In yet other specific embodiments, the number of cells is at least 700, 1,000, 5,000, 10,000 cells. In some embodiments, the number of cells is at least 5,000, 10,000, 25,000, 50,000, or 100,000 cells. In yet another embodiment, the number of cells is at least 50,000, or 100,000 cells. In other embodiments, the number of cells is at least  $1 \times 10^6$ ,  $5 \times 10^6$ ,  $1 \times 10^7$ ,  $5 \times 10^7$ ,  $1 \times 10^8$ ,  $5 \times 10^8$  or more cells. In other embodiments, the number of cells is at least  $1 \times 10^6$ ,  $5 \times 10^6$ , or  $1 \times 10^7$  or more cells. In some embodiments, the number of cells is at least  $1 \times 10^7$ ,  $5 \times 10^7$ ,  $1 \times 10^8$  or more cells. In other embodiments, the number of cells is at least  $5 \times 10^7$ ,  $1 \times 10^8$ ,  $5 \times 10^8$  or more cells. In specific embodiments, the number of cells is between  $1 \times 10^4$  to  $1 \times 10^6$ ,  $5 \times 10^4$  to  $5 \times 10^6$ ,  $1 \times 10^5$  to  $1 \times 10^7$ ,  $1 \times 10^5$  to  $5 \times 10^8$ ,  $1 \times 10^6$  to  $1 \times 10^8$ , or  $1 \times 10^6$  to  $1 \times 10^7$ , or  $1 \times 10^4$  to  $1 \times 10^5$  cells.

[0261] In certain embodiments, a subject is administered a Therapeutic Agent or composition thereof in an amount effective to inhibit or reduce symptoms associated with a disease or disorder by at least 20% to 25%, preferably at least 25% to 30%, at least 30% to 35%, at least 35% to 40%, at least 40% to 45%, at least 45% to 50%, at least 50% to 55%, at least 55% to 60%, at least 60% to 65%, at least 65% to 70%, at least 70% to 75%, at least 75% to 80%, or up to at least 85% relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In certain embodiments to treat, a subject is administered a Therapeutic Agent or a composition thereof in an amount effective to inhibit or reduce symptoms associated with a disease or disorder by at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 8 fold, 10 fold, 15 fold, 20 fold, or 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to a

negative control as determined using an assay described herein or other known to one of skill in the art.

**[0262]** In certain embodiments to treat or manage an infectious disease, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to inhibit or reduce replication of an infectious agent by at least 20% to 25%, preferably at least 25% to 30%, at least 30% to 35%, at least 35% to 40%, at least 40% to 45%, at least 45% to 50%, at least 50% to 55%, at least 55% to 60%, at least 60% to 65%, at least 65% to 70%, at least 70% to 75%, at least 75% to 80%, or up to at least 85% relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In certain embodiments, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to inhibit or reduce replication of an infectious agent by at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 8 fold, 10 fold, 15 fold, 20 fold, or 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In other embodiments, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to inhibit or reduce replication of an infectious agent by at least 1 log, 1.5 logs, 2 logs, 2.5 logs, 3 logs, 3.5 logs, 4 logs, 5 logs or more relative to a negative control as determined using an assay described herein or others known to one of skill in the art.

**[0263]** In certain embodiments to prevent, treat, and/or manage cancer, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to inhibit or reduce tumor growth or cancer cell proliferation by at least 20% to 25%, preferably at least 25% to 30%, at least 30% to 35%, at least 35% to 40%, at least 40% to 45%, at least 45% to 50%, at least 50% to 55%, at least 55% to 60%, at least 60% to 65%, at least 65% to 70%, at least 70% to 75%, at least 75% to 80%, or up to at least 85% relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In some embodiments, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to inhibit or reduce tumor growth or cancer cell proliferation by at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 8 fold, 10 fold, 15 fold, 20 fold, or 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to a negative control as determined using an assay described herein or others known to one of skill in the art.

**[0264]** In certain embodiments to treat or manage autoimmune or inflammatory diseases, a subject is administered an Antagonistic Therapeutic Agent or composition thereof in an amount effective to suppress or reduce certain aspects of the immune function by at least 20% to 25%, preferably at least 25% to 30%, at least 30% to 35%, at least 35% to 40%, at least 40% to 45%, at least 45% to 50%, at least 50% to 55%, at least 55% to 60%, at least 60% to 65%, at least 65% to 70%, at least 70% to 75%, at least 75% to 80%, or up to at least 85% relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In some embodiments, a subject is administered an Antagonistic Therapeutic Agent or composition thereof in an amount effective to suppress or reduce certain aspects of the immune function by at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 8 fold, 10 fold, 15 fold, 20 fold, or 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to a negative control as determined using an assay described herein or others known to one of skill in the art.

**[0265]** In certain embodiments to, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to induce or enhance an immune response by at least 20% to 25%, preferably at least 25% to 30%, at least 30% to 35%, at least 35% to 40%, at least 40% to 45%, at least 45% to 50%, at least 50% to 55%, at least 55% to 60%, at least 60% to 65%, at least 65% to 70%, at least 70% to 75%, at least 75% to 80%, or up to at least 85% relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In some embodiments, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to induce or enhance an immune response by at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 8 fold, 10 fold, 15 fold, 20 fold, or 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to a negative control as determined using an assay described herein or others known to one of skill in the art.

**[0266]** In certain embodiments to, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to increase or enhance the number of lymphocytes (in some embodiments, in a specific target body compartment) by at least 20% to 25%, preferably at least 25% to 30%, at least 30% to 35%, at least 35% to 40%, at least 40% to 45%, at least 45% to 50%, at least 50% to 55%, at least 55% to 60%, at least 60% to 65%, at least 65% to 70%, at least 70% to 75%, at least 75% to 80%, or up to at least 85% relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In some embodiments, a subject is administered an Agonistic Therapeutic Agent or composition thereof in an amount effective to increase or enhance the number of lymphocytes (in some embodiments, in a specific target body compartment) by at least 1.5 fold, at least 2 fold, at least 2.5 fold, at least 3 fold, at least 4 fold, at least 5 fold, at least 8 fold, at least 10 fold, at least 15 fold; or at least 20 fold; or by approximately 2 to 5 fold, 2 to 10 fold, 5 to 10 fold, or 5 to 20 fold relative to a negative control as determined using an assay described herein or others known to one of skill in the art. In some embodiments, the specific target body compartment where the number of lymphocytes is increased or enhanced by an Agonistic Therapeutic Agent is the lung, stomach, heart, kidney, liver, small intestines, large intestines, breast, prostate, or bladder. In particular embodiments, the specific target body compartment where the number of lymphocytes is increased or enhanced is the body compartment affected by a disease or disorder (e.g., cancer or infectious disease). In some embodiments, the specific target body

compartment where the number of lymphocytes is increased or enhanced is the lymph node, spleen, or peripheral blood.

**[0267]** In certain embodiments, a dose of a Therapeutic Agent or composition thereof is administered to a subject every day, every other day, every couple of days, every third day, once a week, twice a week, three times a week, or once every two weeks. In other embodiments, two, three or four doses of a Therapeutic Agent or composition thereof is administered to a subject every day, every couple of days, every third day, once a week or once every two weeks. In some embodiments, a dose(s) of a Therapeutic Agent or composition thereof is administered for 2 days, 3 days, 5 days, 7 days, 14 days, or 21 days. In certain embodiments, a dose of a Therapeutic Agent or composition thereof is administered for 1 month, 1.5 months, 2 months, 2.5 months, 3 months, 4 months, 5 months, 6 months or more.

**[0268]** The dosages of prophylactic or therapeutic agents which have been or are currently used for the prevention, treatment and/or management of a disease or disorder that is affected by IL-15 function/signaling, e.g., cancer, infectious disease, autoimmune and inflammatory disease, and transplant rejection, can be determined using references available to a clinician such as, e.g., the Physicians' Desk Reference (61st ed. 2007). Preferably, dosages lower than those which have been or are currently being used to prevent, treat and/or manage the disease or disorder are utilized in combination with one or more Therapeutic Agents or compositions thereof.

**[0269]** For agents which have been approved for uses other than prevention, treatment or management of disease or disorder that is affected by IL-15 function/signaling, e.g., cancer, infectious disease, autoimmune and inflammatory disease, and transplant rejection, safe ranges of doses can be readily determined using references available to clinicians, such as e.g., the Physician's Desk Reference (61st ed. 2007).

**[0270]** The above-described administration schedules are provided for illustrative purposes only and should not be considered limiting. A person of ordinary skill in the art will readily understand that all doses are within the scope of the invention.

#### 5.4.5. Additional/Combination Therapy

**[0271]** Other therapies that can be used in combination with Therapeutic Agents (i.e., Agonistic Therapeutic Agents and Antagonistic Therapeutic Agents) for the prevention, treatment and/or management of a disease that is affected by IL-15 function/signaling, e.g., cancer, infectious disease, autoimmune and inflammatory disease, and transplant rejection, include, but are not limited to, small molecules, synthetic drugs, peptides (including cyclic peptides), polypeptides, proteins, nucleic acids (e.g., DNA and RNA nucleotides including, but not limited to, antisense nucleotide sequences, triple helices, RNAi, and nucleotide sequences encoding biologically active proteins, polypeptides or peptides), antibodies, synthetic or natural inorganic molecules, mimetic agents, and synthetic or natural organic molecules. Specific examples of such therapies include, but are not limited to, immunomodulatory agents (e.g., interferon), anti-inflammatory agents (e.g., adrenocorticoids, corticosteroids (e.g., beclomethasone, budesonide, flunisolide, fluticasone, triamcinolone, methylprednisolone, prednisolone, prednisone, hydrocortisone), glucocorticoids, steriods, and non-steriodal anti- inflammatory drugs (e.g., aspirin, ibuprofen, diclofenac, and COX-2 inhibitors), pain relievers, leukotriene antagonists (e.g., montelukast, methyl xanthines, zafirlukast, and zileuton), beta2-agonists (e.g., albuterol, biterol, fenoterol, isoetharie, metaproterenol, pirbuterol, salbutamol, terbutalin formoterol, salmeterol, and salbutamol terbutaline), anticholinergic agents (e.g., ipratropium bromide and oxitropium bromide), sulphosalazine, penicillamine, dapsone, antihistamines, anti-malarial agents (e.g., hydroxychloroquine), anti-viral agents (e.g., nucleoside analogs (e.g., zidovudine, acyclovir, gancyclovir, vidarabine, idoxuridine, trifluridine, and ribavirin), foscarnet, amantadine, rimantadine, saquinavir, indinavir, ritonavir, and AZT) and antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, erythomycin, penicillin, mithramycin, and anthramycin (AMC)).

**[0272]** Any therapy which is known to be useful, or which has been used or is currently being used for the prevention, management, and/or treatment of a disease that is affected by IL-15 function/signaling can be used in combination with Therapeutic Agents. See, e.g., Gilman et al., Goodman and Gilman's: The Pharmacological Basis of Therapeutics, 10th ed., McGraw-Hill, New York, 2001; The Merck Manual of Diagnosis and Therapy, Berkow, M.D. et al. (eds.), 17th Ed., Merck Sharp & Dohme Research Laboratories, Rahway, NJ, 1999; Cecil Textbook of Medicine, 20th Ed., Bennett and Plum (eds.), W.B. Saunders, Philadelphia, 1996, and Physicians' Desk Reference (61 st ed. 2007) for information regarding therapies (e.g., prophylactic or therapeutic agents) which have been or are currently being used for preventing, treating and/or managing disease or disorder that is affected by IL-15 function/signaling, e.g., cancer, infectious disease, autoimmune and inflammatory disease, graft versus host disease, and transplant rejection.

##### 5.4.5.1. Anti-Cancer Agents

**[0273]** Non-limiting examples of one or more other therapies that can be used in combination with a Therapeutic Agent include immunomodulatory agents, such as but not limited to, chemotherapeutic agents and non-chemotherapeutic immunomodulatory agents. Non-limiting examples of chemotherapeutic agents include methotrexate, cyclosporin A, leflunomide, cisplatin, ifosfamide, taxanes such as taxol and paclitaxol, topoisomerase I inhibitors (e.g., CPT-11, topo-

5 tecan, 9-AC, and GG-211), gemcitabine, vinorelbine, oxaliplatin, 5-fluorouracil (5-FU), leucovorin, vinorelbine, temodal, cytochalasin B, gramicidin D, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin homologs, and cytoxan. Examples of non-chemo-therapeutic immunomodulatory agents include, but are not limited to, anti-T cell receptor antibodies (e.g., anti-CD4 antibodies (e.g., cM-T412 (Boeringer), IDEC-CE9.1® (IDEC and SKB), mAB 4162W94, Orthoclone and OKTcdr4a (Janssen-Cilag)), anti-CD3 antibodies (e.g., Nuvion (Product Design Labs), OKT3 (Johnson & Johnson), or Rituxan (IDEC)), anti-CD5 antibodies (e.g., an anti-CD5 ricin-linked immunoconjugate), anti-CD7 antibodies (e.g., CHH-380 (Novartis)), anti-CD8 antibodies, anti-CD40 ligand monoclonal antibodies (e.g., IDEC-131 (IDEC)), anti-CD52 antibodies (e.g., CAMPATH 1H (Ilex)), anti-CD2 antibodies (e.g., MEDI-507 (MedImmune, Inc., International Publication Nos. WO 02/098370 and WO 02/069904), anti-CD11a antibodies (e.g., Xanelim (Genentech)), and anti-B7 antibodies (e.g., IDEC-114) (IDEC)); anti-cytokine receptor antibodies (e.g., anti-IFN receptor antibodies, anti-IL-2 receptor antibodies (e.g., Zenapax (Protein Design Labs)), anti-IL-4 receptor antibodies, anti-IL-6 receptor antibodies, anti-IL-10 receptor antibodies, and anti-IL-12 receptor antibodies), anti-cytokine antibodies (e.g., anti-IFN antibodies, anti-TNF- $\alpha$  antibodies, anti-IL-1 $\beta$  antibodies, anti-IL-6 antibodies, anti-IL-8 antibodies (e.g., ABX-IL-8 (Abgenix)), anti-IL-12 antibodies and anti-IL-23 antibodies); CTLA4-immunoglobulin; LFA-3TIP (Biogen, International Publication No. WO 93/08656 and U.S. Patent No. 6,162,432); soluble cytokine receptors (e.g., the extracellular domain of a TNF- $\alpha$  receptor or a fragment thereof, the extracellular domain of an IL-1 $\beta$  receptor or a fragment thereof, and the extracellular domain of an IL-6 receptor or a fragment thereof); cytokines or fragments thereof (e.g., interleukin (IL)-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-15, IL-23, TNF- $\alpha$ , TNF- $\beta$ , interferon (IFN)- $\alpha$ , IFN- $\beta$ , IFN- $\gamma$ , and GM-CSF); and anti-cytokine antibodies (e.g., anti-IL-2 antibodies, anti-IL-4 antibodies, anti-IL-6 antibodies, anti-IL-10 antibodies, anti-IL-12 antibodies, anti-IL-15 antibodies, anti-TNF- $\alpha$  antibodies, and anti-IFN- $\gamma$  antibodies), and antibodies that immunospecifically bind to tumor-associated antigens (e.g., Herceptin®). In certain embodiments, an immunomodulatory agent is an immunomodulatory agent other than a chemotherapeutic agent. In other embodiments an immunomodulatory agent is an immunomodulatory agent other than a cytokine or hemopoietic such as IL-1, IL-2, IL-4, IL-12, IL-15, TNF, IFN- $\alpha$ , IFN- $\beta$ , IFN- $\gamma$ , M-CSF, G-CSF, IL-3 or erythropoietin. In yet other embodiments, an immunomodulatory agent is an agent other than a chemotherapeutic agent and a cytokine or hemopoietic factor.

30 **[0274]** Non-limiting examples of anti-cancer agents that can be used as therapies in combination with Therapeutic Agents, include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxitene; droloxitene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitracin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulazole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofo-sine; interleukin II (including recombinant interleukin II, or rIL2), interferon alpha-2a; interferon alpha-2b; interferon alpha-1; interferon alpha-3; interferon beta-1a; interferon gamma-1b; iproplatin; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; maso-procol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedepa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogetimide; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiomustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprime; thioguanine; thiotepla; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; triciribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vineristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride. Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecyepenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; am-

5 rubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetrorelix; chlorIns; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentanthraquinones; cycloplatan; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; daclizimab; decitabine; dehydrodideamin B; deslorelin; dexamethasone; dexamofamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnor spermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine; droloxfene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflorenthine; el-temene; emitefur; epirubicin; epiristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium 20 texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; iso-homohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; 25 lentinan sulfate; leptoLstatin; letrozole; leukemia alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; HMG-CoA reductase inhibitor (such as but not limited to, Lovastatin, Pravastatin, Fluvastatin, Statin, Simvastatin, and Atorvastatin); loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; manostatin A; marimastat; masoprolol; maspin; 30 matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; molidamol; multiple drug resistance gene inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaper- 35 oxide B; mycobacterial cell wall extract; myriaporone; N-acetylinaline; N-substituted benzamides; nafarelin; nategestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoyl rhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; 40 protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived 45 inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain antigen binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotriptan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; triciribine; 50 55

trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrophostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; Vitaxin®; vorozole; zanoteron; zeniplatin; zilascorb; and zinostatin stimalamer. Additional anti-cancer drugs are 5-fluorouracil and leucovorin.

5 These two agents are particularly useful when used in methods employing thalidomide and a topoisomerase inhibitor. In specific embodiments, a anti-cancer agent is not a chemotherapeutic agent.

10 [0275] In other specific embodiments, a Therapeutic Agent can be administered in combination with the administration of one or more therapies such as, but not limited to anti-cancer agents such as those disclosed in Table 1 with standard doses. When used in a combination therapy, the dosages and/or the frequency of administration listed in Table 1 may be decreased.

TABLE 1

Therapeutic Agent	Dose/Administration/Formulation		
doxorubicin hydrochloride (Adriamycin RDF® and Adriamycin PFS®)	Intravenous	60-75 mg/m <sup>2</sup> on Day 1	21 day intervals
epirubicin hydrochloride (Ellence™)	Intravenous	100-120 mg/m <sup>2</sup> on Day 1 of each cycle or divided equally and given on Days 1-8 of the cycle	3-4 week cycles
fluorousacil	Intravenous	How supplied: 5 mL and 10 mL vials (containing 250 and 500 mg flourouracil respectively)	
docetaxel (Taxotere®)	Intravenous	60- 100 mg/m <sup>2</sup> over 1 hour	Once every 3 weeks
paclitaxel (Taxol®)	Intravenous	175 mg/m <sup>2</sup> over 3 hours	Every 3 weeks for 4 courses (administered sequentially to doxorubicin-containing combination chemotherapy)
tamoxifen citrate (Nolvadex®)	Oral (tablet)	20-40 mg Dosages greater than 20 mg should be given in divided doses (morning and evening)	Daily
leucovorin calcium for injection	intravenous or intramuscular injection	How supplied: 350 mg vial	Dosage is unclear from text. PDR 3610
luprolide acetate (Lupron®)	single subcutaneous injection	1 mg (0.2 mL or 20 unit mark)	Once a day
flutamide (Eulexin®)	Oral (capsule)	50 mg (capsules contain 125 mg flutamide each)	3 times a day at 8 hour intervals (total daily dosage 750 mg)
nilutamide (Nilandron®)	Oral (tablet)	300 mg or 150 mg (tablets contain 50 or 150 mg nilutamide each)	300 mg once a day for 30 days followed by 150 mg once a day
bicalutamide (Casodex®)	Oral (tablet)	50 mg (tablets contain 50 mg bicalutamide each)	Once a day
progesterone	Injection	USP in sesame oil 50 mg/mL	
ketoconazole (Nizoral®)	Cream	2% cream applied once or twice daily depending on symptoms	

(continued)

Therapeutic Agent		Dose/Administration/Formulation		
5	prednisone	Oral (tablet)	Initial dosage may vary from 5 mg to 60 mg per day depending on the specific disease entity being treated.	
10	estramustine phosphate sodium (Emcyt®)	Oral (capsule)	14 mg/ kg of body weight (i.e. one 140 mg capsule for each 10 kg or 22 1b of body weight)	Daily given in 3 or 4 divided doses
15	etoposide or VP-16	Intravenous	5 mL of 20 mg/ mL solution (100 mg)	
20	dacarbazine (DTIC-Dome®)	Intravenous	2-4.5 mg/kg	Once a day for 10 days. May be repeated at 4 week intervals
25	polifeprosan 20 with carmustine implant (BCNU) (nitrosourea) (Gliadel®)	wafer placed in resection cavity	8 wafers, each containing 7.7 mg of carmustine, for a total of 61.6 mg, if size and shape of resection cavity allows	
30	cisplatin	Injection	[n/a in PDR 861] How supplied: solution of 1 mg/mL in multi-dose vials of 50mL and 100mL	
35	mitomycin	Injection	supplied in 5 mg and 20 mg vials (containing 5 mg and 20 mg mitomycin)	
40	gemcitabine HCl (Gemzar®)	Intravenous	For NSCLC- 2 schedules have been investigated and the optimum schedule has not been determined 4 week schedule-administration intravenously at 1000 mg/m <sup>2</sup> over 30 minutes on 3 week schedule-Gemzar administered intravenously at 1250 mg/m <sup>2</sup> over 30 minutes	4 week schedule-Days 1,8 and 15 of each 28-day cycle. Cisplatin intravenously at 100 mg/m <sup>2</sup> on day 1 after the infusion of Gemzar. 3 week schedule-Days 1 and 8 of each 21 day cycle. Cisplatin at dosage of 100 mg/m <sup>2</sup> administered intravenously after administration of Gemzar on day 1.
45	carboplatin (Paraplatin®)	Intravenous	Single agent therapy: 360 mg/m <sup>2</sup> I.V. on day 1 (infusion lasting 15 minutes or longer) Other dosage calculations: Combination therapy with cyclophosphamide, Dose adjustment recommendations, Formula dosing, etc.	Every 4 weeks
50	ifosamide (Ifex®)	Intravenous	1.2 g/m <sup>2</sup> daily	5 consecutive days Repeat every 3 weeks or after recovery from hematologic toxicity
55	topotecan hydrochloride (Hycamtin®)	Intravenous	1.5 mg/m <sup>2</sup> by intravenous infusion over 30 minutes daily	5 consecutive days, starting on day 1 of 21 day course

(continued)

Therapeutic Agent	Dose/Administration/Formulation		
5 Bisphosphonates Pamidronate Alendronate Risedronate	Intravenous or Oral take with 6-8 oz water.	60 mg or 90 mg single infusion over 4-24 hours to correct hypercalcemia in cancer patients 5 mg/d daily for 2 years and then 10 mg/d for 9 month to prevent or control bone resorption. 5.0 mg to prevent or control bone resorption.	
10 Lovastatin (Mevacor™)	Oral	10 - 80 mg/day in single or two divided dose.	

#### 5.4.5.2. Antiviral Agents

15 [0276] Antiviral agents that can be used in combination with Therapeutic Agents include, but are not limited to, non-nucleoside reverse transcriptase inhibitors, nucleoside reverse transcriptase inhibitors, protease inhibitors, and fusion inhibitors. In one embodiment, the antiviral agent is selected from the group consisting of amantadine, oseltamivir phosphate, rimantadine, and zanamivir. In another embodiment, the antiviral agent is a non-nucleoside reverse transcriptase inhibitor selected from the group consisting of delavirdine, efavirenz, and nevirapine. In another embodiment, the antiviral agent is a nucleoside reverse transcriptase inhibitor selected from the group consisting of abacavir, didanosine, emtricitabine, emtricitabine, lamivudine, stavudine, tenofovir DF, zalcitabine, and zidovudine. In another embodiment, the antiviral agent is a protease inhibitor selected from the group consisting of amprenavir, atazanavir, fosamprenav, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir. In another embodiment, the antiviral agent is a fusion inhibitor such as enfuvirtide.

20 [0277] Additional, non-limiting examples of antiviral agents for use in combination with Therapeutic Agents include the following: rifampicin, nucleoside reverse transcriptase inhibitors (e.g., AZT, ddI, ddC, 3TC, d4T), non-nucleoside reverse transcriptase inhibitors (e.g., delavirdine efavirenz, nevirapine), protease inhibitors (e.g., aprenavir, indinavir, ritonavir, and saquinavir), idoxuridine, cidofovir, acyclovir, ganciclovir, zanamivir, amantadine, and palivizumab. Other 25 examples of anti-viral agents include but are not limited to acemannan; acyclovir; acyclovir sodium; adefovir; alovudine; alvircept sudotox; amantadine hydrochloride (SYMMETREL™); aranotin; arildone; atevirdine mesylate; avridine; cidofovir; cipamfylline; cytarabine hydrochloride; delavirdine mesylate; desciclovir; didanosine; disoxaril; edoxudine; envirodene; enviroxime; famciclovir; famotine hydrochloride; fiaicitabine; fialuridine; fosarilate; foscamet sodium; fosfonet sodium; ganciclovir; ganciclovir sodium; idoxuridine; kethoxal; lamivudine; lobucavir; memantine hydrochloride; methisazone; nevirapine; oseltamivir phosphate (TAMIFLUTM); penciclovir; pirodavir; ribavirin; rimantadine hydrochloride (FLUMADINETM); saquinavir mesylate; somantadine hydrochloride; sorivudine; statolon; stavudine; tilorone hydrochloride; trifluridine; valacyclovir hydrochloride; vidarabine; vidarabine phosphate; vidarabine sodium phosphate; viroxime; zalcitabine; zanamivir (RELENZATM); zidovudine; and zinviroxime.

#### 40 5.4.5.3. Antibacterial Agents

45 [0278] Antibacterial agents, including antibiotics, that can be used in combination with Therapeutic Agents include, but are not limited to, aminoglycoside antibiotics, glycopeptides, amphenicol antibiotics, ansamycin antibiotics, cephalosporins, cephamycins oxazolidinones, penicillins, quinolones, streptogramins, tetracyclines, and analogs thereof. In some embodiments, antibiotics are administered in combination with a Therapeutic Agent to prevent and/or treat a bacterial infection.

50 [0279] In a specific embodiment, Therapeutic Agents are used in combination with other protein synthesis inhibitors, including but not limited to, streptomycin, neomycin, erythromycin, carbomycin, and spiramycin.

55 [0280] In one embodiment, the antibacterial agent is selected from the group consisting of ampicillin, amoxicillin, ciprofloxacin, gentamycin, kanamycin, neomycin, penicillin G, streptomycin, sulfanilamide, and vancomycin. In another embodiment, the antibacterial agent is selected from the group consisting of azithromycin, cefonicid, cefotetan, cephalexin, cephamycin, chlortetracycline, clarithromycin, clindamycin, cycloserine, dalfopristin, doxycycline, erythromycin, linezolid, mupirocin, oxytetracycline, quinupristin, rifampin, spectinomycin, and trimethoprim.

55 [0281] Additional, non-limiting examples of antibacterial agents for use in combination with Therapeutic Agents include the following: aminoglycoside antibiotics (e.g., apramycin, arbekacin, bambamycins, butirosin, dibekacin, neomycin, neomycin, undecylenate, netilmicin, paromomycin, ribostamycin, sisomicin, and spectinomycin), amphenicol antibiotics (e.g., azidamfenicol, chloramphenicol, florfenicol, and thiamphenicol), ansamycin antibiotics (e.g., rifamide and rifampin),

carbacephems (e.g., loracarbef), carbapenems (e.g., biapenem and imipenem), cephalosporins (e.g., cefaclor, cefadroxil, cefamandole, cefatrizine, cefazidone, cefozopran, cefpimizole, cefpiramide, and cefpirome), cephams (e.g., cefbuperazone, cefmetazole, and cefminox), folic acid analogs (e.g., trimethoprim), glycopeptides (e.g., vancomycin), lincosamides (e.g., clindamycin, and lincomycin), macrolides (e.g., azithromycin, carbomycin, clarithromycin, dirithromycin, erythromycin, and erythromycin acistrate), monobactams (e.g., aztreonam, carumonam, and tigemonam), nitrofurans (e.g., furaltadone, and furazolium chloride), oxacephems (e.g., flomoxef, and moxalactam), oxazolidinones (e.g., linezolid), penicillins (e.g., amdinocillin, amdinocillin pivoxil, amoxicillin, bacampicillin, benzylpenicillinic acid, benzylpenicillin sodium, epicillin, fenbenicillin, floxacillin, penamccillin, penethamate hydriodide, penicillin o benethamine, penicillin O, penicillin V, penicillin V benzathine, penicillin V hydrabamine, penimepencycline, and phencihicillin potassium), quinolones and analogs thereof (e.g., cinoxacin, ciprofloxacin, clinafloxacin, flumequine, grepagloxacin, levofloxacin, and moxifloxacin), streptogramins (e.g., quinupristin and dalfopristin), sulfonamides (e.g., acetyl sulfamethoxypyrazine, benzylsulfamide, nopyrlsulfamide, phthalylsulfacetamide, sulfachrysoidine, and sulfacytine), sulfones (e.g., diathymosulfone, glucosulfone sodium, and solasulfone), and tetracyclines (e.g., apicycline, chlortetracycline, clomocycline, and demeclocycline). Additional examples include cycloserine, mupirocin, tuberin amphomycin, bacitracin, capreomycin, colistin, enduracidin, enviomycin, and 2,4 diaminopyrimidines (e.g., brodimoprim).

## 5.5. Biological Activity

### 5.5.1. Assays for Testing the Function of the Therapeutic Agent

#### 5.5.1.1. Cell-based Assays

**[0282]** The invention provides for methods to identify agents that modulate the activity of IL-15/IL-15Ra complexes. The activity of an agent can be assayed with an IL-15 sensitive cell line, e.g., CTLL-2 cells, a mouse cytotoxic T lymphoma cell line (ATCC Accession No. TIB-214) or TF1- $\beta$  cells. See, e.g., International Publication No. WO 05/085282. For example, to identify antagonists of IL-15 mediated function, proliferation of CTLL-2 or TF1- $\beta$  cells cultured with an IL-15/IL-15Ra complex in the presence or absence of one or more Antagonist Therapeutic Agents (e.g., antibody) can be assessed by  $^3$ H-thymidine incorporation assays well known in the art and described in International Publication No. WO 05/085282, which is incorporated by reference herein in its entirety.

**[0283]** To assess the activity of Therapeutic Agents (e.g., polypeptides or nucleic acids encoding IL-15 and/or IL-15Ra and cells expressing those polypeptides) as agonists, proliferation of CTLL-2 or TF1- $\beta$  cells cultured in the presence or absence of one or more Therapeutic Agents (e.g., IL-15/IL-15Ra complex) can be assessed by  $^3$ H-thymidine incorporation assays well known in the art and described in International Publication No. WO 05/085282.

**[0284]** Various assays known in the art can be used to assess whether a Therapeutic Agent enhances or suppresses immune function. In one aspect, the Therapeutic Agent increases an immune response that can be, e.g., an antibody response (humoral response) or a cellular immune response, e.g., cytokine secretion (e.g., interferon-gamma), helper activity or cellular cytotoxicity. In one embodiment, the increased immune response is increased cytokine secretion, antibody production, effector function, T cell proliferation, and/or NK cell proliferation. Various assays to measure such activities are well known in the art, and exemplary descriptions of such assays are provided below.

**[0285]** For example, enzyme-linked immunosorbent assays (ELISA) are well known in the art and are described, e.g., in Section 2.1 of Current Protocols in Immunology, Coligan et al. (eds.), John Wiley and Sons, Inc. 1997. ELISA can be used, e.g., to assay the amount or concentration of IL-15 or IL-15Ra polypeptide.

**[0286]** In another method, the "tetramer staining" assay (Altman et al., 1996, Science 274: 94-96) may be used to identify antigen-specific T-cells and to assess how Therapeutic Agents modulate (e.g., enhance or suppress) antigen-specific T cell responses. For example, an MHC molecule containing a specific peptide antigen, such as a tumor-specific antigen, is multimerized to make soluble peptide tetramers and labeled, for example, by complexing to streptavidin. The MHC-peptide antigen complex is then mixed with a population of T cells obtained from a subject administered with an immunogenic composition alone or in combination with a Therapeutic Agent. Biotin is then used to stain T cells which express the tumor-specific antigen of interest.

**[0287]** Furthermore, using the mixed lymphocyte target culture assay, the cytotoxicity of T cells can be tested in a  $^{51}$ Cr-release assay as described, e.g., in Palladino et al., 1987, Cancer Res. 47:5074-5079. Briefly, the mixed lymphocyte culture is added to a target cell suspension to give different effector:target (E:T) ratios (usually 1:1 to 40:1). The target cells are pre-labeled by incubating  $1 \times 10^6$  target cells in culture medium containing 500  $\mu$ Ci of  $^{51}$ Cr per ml for one hour at 37°C. The cells are washed three times following labeling. Each assay point (E:T ratio) is performed in triplicate and the appropriate controls incorporated to measure spontaneous  $^{51}$ Cr release (no lymphocytes added to assay) and 100% release (cells lysed with detergent). After incubating the cell mixtures for 4 hours, the cells are pelleted by centrifugation at 200g for 5 minutes. The amount of  $^{51}$ Cr released into the supernatant is measured by a gamma counter. The percent cytotoxicity is measured as cpm in the test sample minus spontaneously released cpm divided by the total detergent

released cpm minus spontaneously released cpm.

**[0288]** In another embodiment, an ELISPOT assay can be used to measure cytokine release *in vitro* by T cells after administration of an effective amount of a Therapeutic Agent to a subject. Cytokine release is detected by antibodies which are specific for a particular cytokine, e.g., interleukin-2, tumor necrosis factor  $\gamma$  or interferon- $\gamma$  (see, e.g., Scheibenbogen et al., 1997, Int. J. Cancer 71:932-936). The assay is carried out in a microtitre plate which has been pre-coated with an antibody specific for a cytokine of interest which captures the cytokine secreted by T cells. After incubation of T cells for 24-48 hours in the coated wells, the T cells are removed and replaced with a second labeled antibody that recognizes a different epitope on the cytokine. After extensive washing to remove unbound antibody, an enzyme substrate which produces a colored reaction product is added to the plate. The number of cytokine-producing cells is counted under a microscope. This method has the advantages of short assay time, and sensitivity without the need of a large number of cytotoxic T cells.

**[0289]** In some aspects, the immune response induced or enhanced by an Agonistic Therapeutic Agent is enhanced or increased by at least 2 fold, 3 fold, 4 fold, 5 fold, 6 fold, 7 fold, 8 fold, 9 fold, 10 fold, 11 fold, or 12 fold relative to an immune response elicited by a negative control as determined by any known assay in the art. In certain embodiments, the immune response induced by the Agonistic Therapeutic Agent is enhanced by at least 0.5-2 times, at least 2-5 times, at least 5-10 times, at least 10-50 times, at least 50-100 times, at least 100-200 times, at least 200-300 times, at least 300-400 times or at least 400-500 times relative to the immune response induced by a negative control as assayed by any known method in the art. In specific embodiments, the assay used to assess immune response measures the level of antibody production, cytokine production, or cellular cytotoxicity, and such assays are well known in the art. In some embodiments, the assay used to measure the immune response is an enzyme-linked immunosorbent assay (ELISA) that determines antibody or cytokine levels, an ELISPOT assay that determines cytokine release, or a  $^{51}\text{Cr}$  release assay that determines cellular cytotoxicity.

**[0290]** In specific embodiments, the Agonistic Therapeutic Agent induces or enhances an immune response in a subject that is measured by antibody titer in the serum of the subject, and the antibody titer is at least 0.2 to 5 times, 5 to 20 times, 10 to 30 times, 20 to 50 times, 50 to 200 times, 100 to 500, 200 to 1000 times, or 500 to 2,000 times higher as compared to the antibody titer in the serum of a subject administered a negative control. In specific embodiments, the mean serum antibody titer against the antigen in the subject administered the Agonistic Therapeutic Agent is increased by at least 0.5-2 times, at least 2-5 times, at least 5-10 times, at least 10-50 times, at least 50-100 times, at least 100-200 times, at least 200-300 times, at least 300-400 times or at least 400-500 times relative to the mean serum antibody titer in the subject administered a negative control as determined by methods well known in the art.

**[0291]** In another specific embodiment, the invention provides methods of administering Agonistic Therapeutic Agents to induce or enhance the level of cytokine production or secretion, e.g., interferon- $\gamma$ , (that may be 0.5 to 500 times higher) as compared to the level of cytokine production or secretion in a negative control sample. In specific embodiments, the Agonistic Therapeutic Agent induces or enhances an immune response that is measured by increased cytokine release, and the cytokine concentration is at least 0.2 to 5 times, 5 to 20 times, 10 to 30 times, 20 to 50 times, 50 to 200 times, 100 to 500, 200 to 1000 times, or 500 to 2,000 times higher as compared to the cytokine concentration of a negative control. In specific embodiments, the mean serum cytokine concentration of samples obtained from a subject administered the Agonistic Therapeutic Agent is increased by at least 0.5-2 times, at least 2-5 times, at least 5-10 times, at least 10-50 times, at least 50-100 times, at least 100-200 times, at least 200-300 times, at least 300-400 times or at least 400-500 times relative to the mean serum cytokine concentration of samples obtained from a subject administered a negative control as determined by methods well known in the art. In some embodiments, the negative control can be samples from the subject prior to administration of the Agonistic Therapeutic Agent.

**[0292]** In specific embodiments, the Agonistic Therapeutic Agent induces or enhances NK cell proliferation in a subject that by at least 0.2 to 5 times, 5 to 20 times, 10 to 30 times, 20 to 50 times, 50 to 200 times, 100 to 500, 200 to 1000 times, or 500 to 2,000 times higher relative to NK cell proliferation in a negative control. In specific embodiments, the Therapeutic Agent induces or enhances T cell proliferation in a subject that by at least 0.2 to 5 times, 5 to 20 times, 10 to 30 times, 20 to 50 times, 50 to 200 times, 100 to 500, 200 to 1000 times, or 500 to 2,000 times higher relative to T cell proliferation in a negative control as determined by methods well known in the art, e.g., flow cytometry, CSFE staining,  $^{3}\text{H}$ -thymidine incorporation.

**[0293]** The increase in antibody (humoral) or cellular immune response induced by an effective amount of the Therapeutic Agent can be assessed using various methods well known in the art.

**[0294]** For assessing the activity of an Antagonistic Therapeutic Agent that is an antibody that immunospecifically binds to the IL-15/IL-15Ra complex, a cell culture assay can be carried out to determine the ability of the antibody to reduce the binding affinity of the IL-15/IL-15Ra complex to the  $\beta\gamma$  receptor complex expressed on the surface of cells. Cells that endogenously or recombinantly express the  $\beta\gamma$  receptor complex can be used in this assay. The cells are contacted with the IL-15/IL-15Ra complex in the presence or absence of the Antagonistic Therapeutic Agent antibody. The IL-15/IL-15Ra complex is labeled with a fluorophore, radioisotope, or other detection markers, and the level of binding of the labeled IL-15/IL-15Ra complex to the  $\beta\gamma$  receptor complex expressed on the cell surface of cells are assayed

in the presence and absence of the antibody using methods known in the art, e.g., flow cytometry fluorescent markers, or other compatible machines to detect the detectin marker. In a specific embodiment, the Antagonistic Therapeutic Agent antibody reduces the amount of labeled IL-15/IL-15Ra complexes that bind to the  $\beta\gamma$  receptor complex on the cell surface.

5

### 5.5.1.2. *In vitro* Assays

[0295] The identification of antibodies that immunospecifically bind to the IL-15/IL-15Ra complex can be assessed using any method well known in the art, e.g., ELISA, coimmunoprecipitation, Biacore assays, and KinEx A assays.

10

[0296] Binding assays can be used to determine the binding affinity of an antibody to IL-15/IL-15Ra complexes. Binding assays may be performed either as direct binding assays or as competition-binding assays. Binding can be detected using standard ELISA or standard Flow Cytometry assays. In a direct binding assay, a candidate antibody is tested for binding to the IL-15/IL-15Ra complex.

15

[0297] Competition-binding assays, on the other hand, assess the ability of a candidate antibody to compete with a known antibodies or other compound that binds IL-15/IL-15Ra complexes.

20

[0298] In a direct binding assay, the IL-15/IL-15Ra complex is contacted with a candidate antibody under conditions that allow binding of the candidate antibody to the IL-15/IL-15Ra complexes. The binding may take place in solution or on a solid surface. Preferably, the candidate antibody is previously labeled for detection. Any detectable compound may be used for labeling, such as but not limited to, a luminescent, fluorescent, or radioactive isotope or group containing same, or a nonisotopic label, such as an enzyme or dye. After a period of incubation sufficient for binding to take place, the reaction is exposed to conditions and manipulations that remove excess or non-specifically bound antibody. Typically, it involves washing with an appropriate buffer. Finally, the presence of a IL-15/IL-15Ra-antibody complex is detected.

25

[0299] In a competition-binding assay, a candidate antibody is evaluated for its ability to inhibit or displace the binding of a known anti-IL-15/IL-15Ra complex antibody (or other compound) to the IL-15/IL-15Ra complex. A labeled known binder of IL-15/IL-15Ra complex may be mixed with the candidate antibody, and placed under conditions in which the interaction between them would normally occur, with and without the addition of the candidate antibody. The amount of labeled known binder of IL-15/IL-15Ra complex that binds the IL-15/IL-15Ra complex may be compared to the amount bound in the presence or absence of the candidate antibody.

30

[0300] In one embodiment, the binding assay is carried out with one or more components immobilized on a solid surface to facilitate antibody antigen complex formation and detection. In various embodiments, the solid support could be, but is not restricted to, polycarbonate, polystyrene, polypropylene, polyethylene, glass, nitrocellulose, dextran, nylon, polyacrylamide and agarose. The support configuration can include beads, membranes, microparticles, the interior surface of a reaction vessel such as a microtiter plate, test tube or other reaction vessel. The immobilization of IL-15/IL-15Ra complex, or other component, can be achieved through covalent or non-covalent attachments. In one embodiment, the attachment may be indirect, i.e., through an attached antibody. In another embodiment, the IL-15/IL-15Ra complex and negative controls are tagged with an epitope, such as glutathione S-transferase (GST) so that the attachment to the solid surface can be mediated by a commercially available antibody such as anti-GST (Santa Cruz Biotechnology).

35

[0301] For example, such an affinity binding assay may be performed using the IL-15/IL-15Ra complex which is immobilized to a solid support. Typically, the non-mobilized component of the binding reaction, in this case the candidate anti-IL-15/IL-15Ra complex antibody, is labeled to enable detection. A variety of labeling methods are available and may be used, such as luminescent, chromophore, fluorescent, or radioactive isotope or group containing same, and nonisotopic labels, such as enzymes or dyes. In one embodiment, the candidate antibody is labeled with a fluorophore such as fluorescein isothiocyanate (FITC, available from Sigma Chemicals, St. Louis). Such an affinity binding assay may be performed using the IL-15/IL-15Ra complex antigen immobilized on a solid surface. Antibodies are then incubated with the antigen and the specific binding of antibodies is detected by methods known in the art including, but not limited to, BiaCore Analyses, ELISA, FMET and RIA methods.

40

[0302] Finally, the label remaining on the solid surface may be detected by any detection method known in the art. For example, if the candidate antibody is labeled with a fluorophore, a fluorimeter may be used to detect complexes.

45

[0303] In one embodiment, the antibody is added to binding assays in the form of intact cells that express IL-15/IL-15Ra complex antigen, or isolated membranes containing IL-15/IL-15Ra complex. Thus, direct binding to IL-15/IL-15Ra complex antigen may be assayed in intact cells in culture or in animal models in the presence and absence of the candidate antibody. A labeled candidate antibody may be mixed with cells that express human IL-15/IL-15Ra complex, or with crude extracts obtained from such cells, and the candidate antibody may be added. Isolated membranes may be used to identify candidate antibodies that interact with IL-15/IL-15Ra complex. For example, in a typical experiment using isolated membranes, cells may be genetically engineered to express IL-15/IL-15Ra complex antigen. Membranes can be harvested by standard techniques and used in an *in vitro* binding assay. Labeled candidate antibody (e.g., fluorescent labeled antibody) is bound to the membranes and assayed for specific activity; specific binding is determined by comparison with binding assays performed in the presence of excess unlabeled (cold) candidate antibody. Alterna-

tively, soluble IL-15/IL-15Ra complex may be recombinantly expressed and utilized in non-cell based assays to identify antibodies that bind to the IL-15/IL-15Ra complex. The recombinantly expressed IL-15/IL-15Ra polypeptides can be used in the non-cell based screening assays.

**[0304]** Alternatively, the binding reaction may be carried out in solution. In this assay, the labeled component is allowed to interact with its binding partner(s) in solution. If the size differences between the labeled component and its binding partner(s) permit such a separation, the separation can be achieved by passing the products of the binding reaction through an ultrafilter whose pores allow passage of unbound labeled component but not of its binding partner(s) or of labeled component bound to its partner(s). Separation can also be achieved using any reagent capable of capturing a binding partner of the labeled component from solution, such as an antibody against the binding partner and so on.

**[0305]** In another specific embodiment, the solid support is membrane containing IL-15/IL-15Ra complex attached to a microtiter dish. Candidate antibodies, for example, can bind cells that express library antibodies cultivated under conditions that allow expression of the library members in the microtiter dish. Library members that bind to the IL-15/IL-15Ra complex are harvested. Such methods, are generally described by way of example in Parmley and Smith, 1988, Gene, 73:305-318; Fowlkes et al., 1992, BioTechniques, 13:422-427; PCT Publication No. WO94/18318; and in references cited hereinabove.

**[0306]** Various methods described above or known in the art can be adapted to assay the binding affinity of IL-15 derivatives to native IL-15Ra, IL-15Ra derivatives to native IL-15, IL-15 derivatives to IL-15Ra derivatives, and IL-15/IL-15Ra complexes to the  $\beta\gamma$  receptor complex.

#### 5.5.2. Animal Models

**[0307]** Therapeutic Agents are preferably assayed *in vivo* for the desired therapeutic or prophylactic activity prior to use in humans. For example, in one embodiment, a Therapeutic Agent can be administered to the animal at the same time as the onset of a disease or disorder in the animal. In another embodiment, a Therapeutic Agent can be administered to the animal prior to the onset of a disease or disorder in the animal. In another embodiment, a Therapeutic Agent can be administered to the animal subsequent to the onset of a disease or disorder in the animal. In a specific embodiment, the Therapeutic Agent is administered to the animal more than one time. In another specific embodiment, the Therapeutic Agent is administered in combination with another therapy.

**[0308]** Therapeutic Agent can be tested in animal models systems including, but are not limited to, rats, mice, chicken, cows, monkeys, pigs, goats, sheep, dogs, rabbits, guinea pigs, etc. In a specific embodiment, Therapeutic Agents are tested in a mouse model system. Such model systems are widely used and well-known to the skilled artisan.

**[0309]** In certain embodiments, an animal, such as six weeks old female Balb/c mice, are administered nucleic acids encoding IL-15 and IL-15Ra by hydrodynamic injection and the plasma levels of IL-15 and/or the bioactivity of IL-15 is assessed. Briefly, animals (e.g., mice) are injected with the IL-15 plasmid alone or in combination with IL-15Ra plasmid in sterile 0.9% NaCl through their tail vein within 7 seconds using a 27.5 gauge needle. Mice are bled after a certain number of days (e.g., at day 1 and day 3) after injection and the plasma levels of IL-15 are measured using, e.g., an IL-15 chemiluminescent immunoassay (QuantiGlo, R&D systems). After a certain number of days (e.g., 3 days) after injection, mice are sacrificed and liver, lungs, spleen, and mesenteric lymph nodes are collected and analyzed to assess IL-15 bioactivity. To make single cell suspensions, spleens are be gently squeezed through a 100  $\mu\text{m}$  Cell Strainer (Thomas) and washed in RPMI (Gibco) to remove any remaining organ stroma. The cells are resuspended in media (e.g., RPMI containing 10% fetal calf serum (FCS)) and counted using, e.g., Acridine Orange (Molecular Probes)/Ethidium Bromide (Fisher) dye. Lung and liver are minced and incubated with collagenase (Sigma) and DNase (Roche) for a period of time (e.g., 1 hour) at 37°C to make single cell suspensions. Single cells are collected and resuspended in media (e.g., complete RPMI with 10% FCS). The bioactivity of IL-15 *in vivo* may be measured in liver, lung and spleen using multicolor Flow Cytometry. Briefly, the cells are washed in FACS buffer containing 0.2% FCS and stained with the following panel of conjugated rat anti-mouse antibodies: CD3-APCCy7, CD4-PerCP, CD8-PECy7, CD44-APC, CD49b-FITC and CD62L-PE (BD Pharmingen). Samples are acquired using FACSaria (BD) and the data is analyzed by FlowJo software (Tree Star, San Carlos, CA).

**[0310]** The anti-cancer activity of the Therapeutic Agent can be determined by using various experimental animal models for the study of cancer well known in the art as described in, e.g., Relevance of Tumor Models for Anticancer Drug Development (1999, eds. Fiebig and Burger); Contributions to Oncology (1999, Karger); The Nude Mouse in Oncology Research (1991, eds. Boven and Winograd); and Anticancer Drug Development Guide (1997 ed. Teicher), incorporated herein by reference in their entireties.

**[0311]** Animal models for cancer can be used to assess the efficacy of a Therapeutic Agent, a composition thereof, or a combination therapy comprising a Therapeutic Agent. Non-limiting examples of animal models for lung cancer include, but are not limited to, lung cancer animal models described by Zhang & Roth (1994, *In vivo* 8(5):755-69) and a transgenic mouse model with disrupted p53 function (see, e.g., Morris et al., 1998, *J La State Med Soc* 150(4):179-85). An example of an animal model for breast cancer includes, but is not limited to, a transgenic mouse that overexpresses

5 cyclin D1 (see, e.g., Hosokawa et al., 2001, *Transgenic Res* 10(5):471-8). An example of an animal model for colon cancer includes, but is not limited to, a TCR- $\beta$  and p53 double knockout mouse (see, e.g., Kado et al., 2001, *Cancer Res* 61(6):2395-8). Examples of animal models for pancreatic cancer include, but are not limited to, a metastatic model of Panc02 murine pancreatic adenocarcinoma (see, e.g., Wang et al., 2001, *Int J Pancreatol* 29(1):37-46) and nu-nu mice generated in subcutaneous pancreatic tumours (see, e.g., Ghaneh et al., 2001, *Gene Ther* 8(3):199-208). Examples of animal models for non-Hodgkin's lymphoma include, but are not limited to, a severe combined immunodeficiency ("SCID") mouse (see, e.g., Bryant et al., 2000, *Lab Invest* 80(4):553-73) and an IgHmu-HOX11 transgenic mouse (see, e.g., Hough et al., 1998, *Proc Natl Acad Sci USA* 95(23):13853-8). An example of an animal model for esophageal cancer includes, but is not limited to, a mouse transgenic for the human papillomavirus type 16 E7 oncogene (see, e.g., Herber et al., 1996, *J Virol* 70(3):1873-81). Examples of animal models for colorectal carcinomas include, but are not limited to, Apc mouse models (see, e.g., Fodde & Smits, 2001, *Trends Mol Med* 7(8):369-73 and Kuraguchi et al., 2000, *Oncogene* 19(50):5755-63).

10 15 20 25 **[0312]** For animal models of infectious diseases, the effectiveness of a Therapeutic Agent relative to a negative control can be assessed in animals infected with virus. Samples obtained from these animals (e.g., serum, urine, sputum, semen, saliva, plasma, or tissue sample) can be tested for enhancement of immune function, e.g., enhancement in cytokine release, enhancement in antibody production, T cell proliferation, NK cell proliferation, with methods well known in the art and described herein. Samples obtained from these animals (e.g., serum, urine, sputum, semen, saliva, plasma, or tissue sample) can also be tested for reduction in viral replication via well known methods in the art, e.g., those that measure altered viral replication (as determined, e.g., by plaque formation) or the production of viral proteins (as determined, e.g., by Western blot, ELISA, or flow cytometry analysis) or viral nucleic acids (as determined, e.g., by RT-PCR, northern blot analysis or southern blot). For quantitation of virus in tissue samples, tissue samples are homogenized in phosphate-buffered saline (PBS), and dilutions of clarified homogenates are adsorbed for 1 hour at 37°C onto monolayers of cells (e.g., Vero, CEF or MDCK cells). In other assays, histopathologic evaluations are performed after infection, preferably evaluations of the organ(s) the virus is known to target for infection. Virus immunohistochemistry can be performed using a viral-specific monoclonal antibody. Non-limiting exemplary animal models described below can be adapted for other viral systems.

30 35 40 45 **[0313]** Various animal models for infectious diseases that are well known in the art can be employed to assess the efficacy of Therapeutic Agents in preventing, treating, and/or managing infectious diseases, e.g.: mouse models of herpes simplex virus (HSV) are described in Crute et al., *Nature Medicine*, 2002, 8:386-391 and Bolger et al., *Antiviral Res.*, 1997, 35:157-165; guinea pig models of HSV are described in Chen et al., *Virol. J.* 2004 Nov 23, 1:11; animal models of mouse cytomegalovirus (MCMV) and human cytomegalovirus (HCMV) are described in Kern et al., *Antimicrob. Agents Chemother.*, 2004, 48:4745-4753; Guinea pig models of CMV is described in Bourne et al., *Antiviral Res.*, 2000, 47:103-109, Bravo et al., *Antiviral Res.*, 2003, 60:41-49 and Bravo et al., *J. Infectious Diseases*, 2006, 193:591-597; animal models of influenza virus are described in Sidwell et al., *Antiviral Res.*, 2000, 48:1-16; and McCauley et al., *Antiviral Res.*, 1995, 27:179-186; mouse models of hepatitis B virus (HBV) are described in Cavanaugh et al., *J. Virol.*, 1997, 71:3236-3243 and Guidotti et al., *J. Virol.*, 1995, 69:6158-6169; mouse models of hepatitis C virus (HCV) are described in Zhu et al., *Antimicrobial Agents and Chemother.*, 2006, 50:3260-3268, Bright et al., *Nature*, 2005, 436:973-978, Hsu et al., *Nat. Biotechnol.*, 2003, 21:519-525, Ilan et al., *J. Infect. Dis.* 2002, 185:153-161, Kneteman et al., *Hepatology*, 2006, 43:1346-1353, Mercer et al., *Nat. Med.*, 2001, 7:927-933, and Wu et al., *Gastroenterology*, 2005, 128:1416-1423; animal models of HIV are described in Ayash-Rashkovsky et al., *FASEB J.*, 2005, 19:1149-1151, Mosier et al., *Semin. Immunol.*, 1996, 8:255-262, Mosier et al., *Hosp. Pract. (Off Ed.)*, 1996, 31:41-48, 53-55, 59-60, Bonyhadi et al., *Mol. Med. Today*, 1997, 3:246-253, Jolicoeur et al., *Leukemia*, 1999, 13:S78-S80, Browning et al., *Proc. Natl. Acad. Sci. USA*, 1997, 94:14637-14641, and Sawada et al., *J. Exp. Med.*, 1998, 187:1439-1449, and Schito et al., *Curr. HIV Res.*, 2006, 4:379-386.

50 55 **[0314]** Other animal models for viral infections can also be used to assess the efficacy of a Therapeutic Agent, a composition thereof, or a combination therapy comprising a Therapeutic Agent, e.g., animal models for viral infections such as EBV-associated diseases, gammaherpesviruses, infectious mononucleosis, simian immunodeficiency virus ("SIV"), Borna disease virus infection; hepatitis, varicella virus infection, viral pneumonitis, Epstein-Barr virus pathogenesis, feline immunodeficiency virus ("FIV"), HTLV type 1 infection, human rotaviruses, and genital herpes have been developed (see, e.g., Hayashi et al., 2002, *Histol Histopathol* 17(4):1293-310; Arico et al., 2002, *J Interferon Cytokine Res* 22(11):1081-8; Flano et al., 2002, *Immunol Res* 25(3):201-17; Sauermann, 2001, *Curr Mol Med* 1(4):515-22; Pletnikov et al., 2002, *Front Biosci* 7:d593-607; Engler et al., 2001, *Mol Immunol* 38(6):457-65; White et al., 2001, *Brain Pathol* 11(4):475-9; Davis & Matalon, 2001, *News Physiol Sci* 16:185-90; Wang, 2001, *Curr Top Microbiol Immunol* 258:201-19; Phillips et al., 2000, *J Psychopharmacol*. 14(3):244-50; Kazanji, 2000, *AIDS Res Hum Retroviruses*. 16(16):1741-6; Saif et al., 1996, *Arch Virol Suppl*. 12:153-61; and Hsiung et al., 1984, *Rev Infect Dis*. 6(1):33-50).

52 **[0315]** Other animal models for viral respiratory infections include, but not limited to, PIV (see, e.g., Shephard et al., 2003 *Res Vet Sci* 74(2): 187-190; Ottolini et al., 2002 *J Infect Dis* 186(12): 1713-1717), and RSV (see, e.g., Culley et al., 2002 *J Exp Med* 196(10): 1381-1386; and Curtis et al., 2002 *Exp Biol Med* 227(9): 799-802).

[0316] The Therapeutic Agent, composition thereof, or combination therapy comprising the Therapeutic Agent can be tested for their ability to decrease the time course of viral infection.

[0317] Animal models for bacterial infections can also be used to assess the efficacy of a Therapeutic Agent, a composition thereof, or a combination therapy comprising a Therapeutic Agent. Animal models for bacterial infections such as *H. pylori*-infection, *genital mycoplasmosis*, *primary sclerosing cholangitis*, *cholera*, chronic lung infection with *Pseudomonas aeruginosa*, Legionnaires' disease, gastroduodenal ulcer disease, bacterial meningitis, gastric *Helicobacter* infection, *pneumococcal otitis media*, experimental allergic neuritis, leprosy neuropathy, mycobacterial infection, endocarditis, Aeromonas-associated enteritis, *Bacteroides fragilis* infection, syphilis, *streptococcal* endocarditis, acute hematogenous osteomyelitis, human scrub typhus, toxic shock syndrome, anaerobic infections, *Escherichia coli* infections, and *Mycoplasma pneumoniae* infections have been developed (see, e.g., Sugiyama et al., 2002, *J Gastroenterol.* 37 Suppl 13:6-9; Brown et al., 2001, *Am J Reprod Immunol.* 46(3):232-41; Vierling, 2001, *Best Pract Res Clin Gastroenterol.* 15(4):591-610; Klose, 2000, *Trends Microbiol.* 8(4):189-91; Stotland et al., 2000, *Pediatr Pulmonol.* 30(5):413-24; Brieland et al., 2000, *Immunopharmacology* 48(3):249-52; Lee, 2000, *Baillieres Best Pract Res Clin Gastroenterol.* 14(1):75-96; Koedel & Pfister, 1999, *Infect Dis Clin North Am.* 13(3):549-77; Nedrud, 1999, *FEMS Immunol Med Microbiol.* 24(2):243-50; Prellner et al., 1999, *Microb Drug Resist.* 5(1):73-82; Vriesendorp, 1997, *J Infect Dis.* 176 Suppl 2:S164-8; Shetty & Antia, 1996, *Indian J Lepr.* 68(1):95-104; Balasubramanian et al., 1994, *Immunobiology* 191(4-5):395-401; Carbon et al., 1994, *Int J Biomed Comput.* 36(1-2):59-67; Haberberger et al., 1991, *Experientia.* 47(5):426-9; Onderdonk et al., 1990, *Rev Infect Dis.* 12 Suppl 2:S169-77; Wicher & Wicher, 1989, *Crit Rev Microbiol.* 16(3):181-234; Scheld, 1987, *J Antimicrob Chemother.* 20 Suppl A:71-85; Emslie & Nade, 1986, *Rev Infect Dis.* 8(6):841-9; Ridgway et al., 1986, *Lab Anim Sci.* 36(5):481-5; Quimby & Nguyen, 1985, *Crit Rev Microbiol.* 12(1):1-44; Onderdonk et al., 1979, *Rev Infect Dis.* 1(2):291-301; Smith, 1976, *Ciba Found Symp.* (42):45-72, and Taylor-Robinson, 1976, *Infection.* 4(1 Suppl):4-8).

[0318] The Therapeutic Agent, composition thereof, or combination therapy comprising the Therapeutic Agent can be tested for their ability to decrease the time course of bacterial infection, e.g., a bacterial respiratory infection by at least 25%, at least 50%, at least 60%, at least 75%, at least 85%, at least 95%, or at least 99% relative to a negative control using methods well known in the art.

[0319] The efficacy of Therapeutic Agents, compositions thereof, or combination therapies comprising Therapeutic Agents for the prevention, treatment and/or management of a fungal infection can be assessed in animal models for such infections. Animal models for fungal infections such as *Candida* infections, zygomycosis, *Candida mastitis*, progressive disseminated trichosporonosis with latent trichosporonemia, disseminated candidiasis, pulmonary paracoccidioidomycosis, pulmonary aspergillosis, *Pneumocystis carinii* pneumonia, cryptococcal meningitis, coccidioidal meningoencephalitis and cerebrospinal vasculitis, *Aspergillus niger* infection, *Fusarium* keratitis, paranasal sinus mycoses, *Aspergillus fumigatus* endocarditis, tibial dyschondroplasia, *Candida glabrata* vaginitis, oropharyngeal candidiasis, X-linked chronic granulomatous disease, tinea pedis, cutaneous candidiasis, mycotic placentitis, disseminated trichosporonosis, allergic bronchopulmonary aspergillosis, mycotic keratitis, *Cryptococcus neoformans* infection, fungal peritonitis, *Curvularia geniculata* infection, *staphylococcal* endophthalmitis, sporotrichosis, and dermatophytosis have been developed (see, e.g., Arendrup et al., 2002, *Infection* 30(5):286-91; Kamei, 2001, *Mycopathologia* 152(1):5-13; Guhad et al., 2000, *FEMS Microbiol Lett.* 192(1):27-31; Yamagata et al., 2000, *J Clin Microbiol.* 38(9):32606; Andrusis et al., 2000, *J Clin Microbiol.* 38(6):2317-23; Cock et al., 2000, *Rev Inst Med Trop Sao Paulo* 42(2):59-66; Shibuya et al., 1999, *Microb Pathog.* 27(3):123-31; Beers et al., 1999, *J Lab Clin Med.* 133(5):423-33; Najvar et al., 1999, *Antimicrob Agents Chemother.* 43(2):413-4; Williams et al., 1988, *J Infect Dis.* 158(4):1217-21; Yoshida, 1988, *Kansenshogaku Zasshi.* 1998 Jun;72(6):621-30; Alexandrakis et al., 1998, *Br J Ophthalmol.* 82(3):306-11; Chakrabarti et al., 1997, *J Med Vet Mycol.* 35(4):295-7; Martin et al., 1997, *Antimicrob Agents Chemother.* 41(1):13-6; Chu et al., 1996, *Avian Dis.* 40(3):715-9; Fidel et al., 1996, *J Infect Dis.* 173(2):425-31; Cole et al., 1995, *FEMS Microbiol Lett.* 15;126(2):177-80; Pollock et al., 1995, *Nat Genet.* 9(2):202-9; Uchida et al., 1994, *Jpn J Antibiot.* 47(10):1407-12; Maebashi et al., 1994, *J Med Vet Mycol.* 32(5):349-59; Jensen & Schonheyder, 1993, *J Exp Anim Sci.* 35(4):155-60; Gokaslan & Anaissie, 1992, *Infect Immun.* 60(8):3339-44; Kurup et al., 1992, *J Immunol.* 148(12):3783-8; Singh et al., 1990, *Mycopathologia.* 112(3):127-37; Salkowski & Balish, 1990, *Infect Immun.* 58(10):3300-6; Ahmad et al., 1986, *Am J Kidney Dis.* 7(2):153-6; Altur-Werber E, Edberg SC, 1985, *Mycopathologia.* 89(2):69-73; Kane et al., 1981, *Antimicrob Agents Chemother.* 20(5):595-9; Barbee et al., 1977, *Am J Pathol.* 86(1):281-4; and Maestrone et al., 1973, *Am J Vet Res.* 34(6):833-6). Animal models for fungal respiratory infections such as *Candida albicans*, *Aspergillus fumigatus*, invasive pulmonary aspergillosis, *Pneumocystis carinii*, pulmonary cryptococcosis, *Pseudomonas aeruginosa*, *Cunninghamella bertholletiae* (see, e.g., Aratani et al., 2002 *Med Mycol.* 40(6):557-563; Bozza et al., 2002 *Microbes Infect.* 4(13): 1281-1290; Kurup et al., 2002 *Int Arch Allergy Immunol.* 129(2):129-137; Hori et al., 2002 *Eur J Immunol.* 32(5): 1282-1291; Rivera et al., 2002 *J Immunol.* 168(7): 3419-3427; Vassallo et al., 2001, *Am J Respir Cell Mol Biol.* 25(2): 203-211; Wilder et al., 2002 *Am J Respir Cell Mol Biol.* 26(3): 304-314; Yonezawa et al., 2000 *J Infect Chemother.* 6(3): 155-161; Cacciapuoti et al., 2000 *Antimicrob Agents Chemother.* 44(8): 2017-2022; and Honda et al., 1998 *Mycopathologia.* 144(3):141-146).

[0320] The Therapeutic Agents, compositions thereof, or combination therapies comprising Therapeutic Agents can

be tested for their ability to decrease the time course of fungal respiratory infection by at least 25%, at least 50%, at least 60%, at least 75%, at least 85%, at least 95%, or at least 99%. Techniques known to those of skill in the art can be used to analyze the function of the Therapeutic Agents, compositions thereof, or combination therapies comprising Therapeutic Agents *in vivo*.

5      [0321] Animal models for autoimmune disorders can also be used to assess the efficacy of a Therapeutic Agent, composition thereof, or combination therapy comprising a Therapeutic Agent. Animal models for autoimmune disorders such as type 1 diabetes, thyroid autoimmunity, systemic lupus erythematosus, and glomerulonephritis have been developed (Flanders et al., 1999, Autoimmunity 29:235-246; Krogh et al., 1999, Biochimie 81:511-515; Foster, 1999, Semin. Nephrol. 19:12-24).

10     [0322] Efficacy in preventing, treating and/or managing an autoimmune disorder may be demonstrated, e.g., by detecting the ability of an antibody, a composition, or a combination therapy described herein to reduce one or more symptoms of the autoimmune disorder, to reduce mean absolute lymphocyte counts, to decrease T cell activation, to decrease T cell proliferation, to reduce cytokine production, or to modulate one or more particular cytokine profiles. Efficacy in preventing or treating psoriasis may be demonstrated, e.g., by detecting the ability of a Therapeutic Agent 15 or composition thereof to reduce one or more symptoms of psoriasis, to reduce mean absolute lymphocyte counts, to reduce cytokine production, to modulate one or more particular cytokine profiles, to decrease scaling, to decrease erythema, to decrease plaque elevation, to decrease T cell activation in the dermis or epidermis of an affected area, to decrease T cell infiltration to the dermis or epidermis of an affected area, to reduce PASI, to improve the physician's global assessment score, or to improve quality of life.

20     [0323] The anti-inflammatory activity of a Therapeutic Agent, a composition thereof, or a combination therapy comprising a Therapeutic Agent can be determined by using various experimental animal models of inflammatory arthritis known in the art and described in Crofford L.J. and Wilder R.L., "Arthritis and Autoimmunity in Animals," in Arthritis and Allied Conditions: A Textbook of Rheumatology, McCarty (eds.), Chapter 30 (Lee and Febiger, 1993). Experimental and spontaneous animal models of inflammatory arthritis and autoimmune rheumatic diseases can also be used to assess 25 the anti-inflammatory activity of the Therapeutic Agents, compositions thereof, or combination therapies comprising Therapeutic Agents.

25     [0324] The anti-inflammatory activity of a Therapeutic Agent, a composition thereof, or a combination therapy comprising a Therapeutic Agent can also be assessed by measuring the inhibition of carrageenan-induced paw edema in the rat, using a modification of the method described in Winter C. A. et al., "Carrageenan Induced Edema in Hind Paw 30 of the Rat as an Assay for Anti-inflammatory Drugs" Proc. Soc. Exp. Biol Med. 111, 544-547, (1962). This assay has been used as a primary *in vivo* screen for the anti-inflammatory activity of most NSAIDs, and is considered predictive 35 of human efficacy. The anti-inflammatory activity of the test therapies (e.g., The Therapeutic Agents, compositions thereof, or combination therapies comprising Therapeutic Agents) is expressed as the percent inhibition of the increase in hind paw weight of the test group relative to the vehicle dosed control group.

35     [0325] In a specific embodiment where the experimental animal model used is adjuvant-induced arthritis rat model, body weight can be measured relative to a control group to determine the anti-inflammatory activity of a Therapeutic Agent, a composition thereof, or a combination therapy.

40     [0326] Animal models for allergies and asthma are known in the art, such as constant-flow inflation with end-inspiratory occlusion described in Ewart et al., 1995 J Appl Physiol 79(2):560-566 and other assays described in, e.g., Komai et 45 al., 2003 Br J Pharmacol 138(5): 912-920; Kenyon et al., 2003 Toxicol Appl Pharmacol 186(2): 90-100; Path et al., 2002 Am J Resp & Critical Care Med 166(6): 818-826; Martins et al., 1990 Crit Care Med 19:515-519; Nicolaides et al., 1997 Proc Natl Acad Sci USA 94:13175-13180; McLane et al., 1998 19:713-720; and Temann et al., 1998 J Exp Med 188(7): 1307-1320. For example, the murine adoptive transfer model is an animal model used to assess the efficacy a Therapeutic Agent, a composition thereof, or a combination therapy for the prevention, treatment, management, and/or asthma 50 include. In the murine adoptive transfer model, aeroallergen provocation of TH1 or TH2 recipient mice results in TH effector cell migration to the airways and is associated with an intense neutrophilic (TH1) and eosinophilic (TH2) lung mucosal inflammatory response (Cohn et al., 1997, J. Exp. Med. 186:1737-1747). Airway hypersensitivity can be induced in mice by ovalbumin (Tomkinson et al., 2001, J. Immunol. 166:5792-5800) or Schistosoma mansoni egg antigen (Tesciuba et al., 2001, J. Immunol. 167:1996-2003).

55     [0327] Efficacy in preventing or treating an inflammatory disorder may be demonstrated, e.g., by detecting the ability of a Therapeutic Agent, a composition thereof, or a combination therapy comprising a Therapeutic Agent to reduce one or more symptoms of the inflammatory disorder, to decrease T cell activation, to decrease T cell proliferation, to modulate one or more cytokine profiles, to reduce cytokine production, to reduce inflammation of a joint, organ or tissue or to improve quality of life.

55     [0328] Changes in inflammatory disease activity may also be assessed through tender and swollen joint counts, patient and physician global scores for pain and disease activity, and the ESR/CRP. Progression of structural joint damage may be assessed by quantitative scoring of X-rays of hands, wrists, and feet (Sharp method). Changes in functional status in humans with inflammatory disorders may be evaluated using the Health Assessment Questionnaire (HAQ), and quality

of life changes are assessed with the SF.

[0329] The efficacy of a Therapeutic Agent, a composition thereof, or a combination therapy comprising a Therapeutic Agent in preventing, treating and/or managing Type I allergic reaction may be assessed by its ability to induce anti-IgE antibodies that inhibit IgE from binding to its receptor on mast cells or basophils *in vitro*. IgE levels can be assayed by immunoassays, gel electrophoresis followed by visualization, radioimmunosorbent test (RIST), radioallergosorbent test (RAST), or any other method known to those skilled in the art.

### 5.5.3. Toxicity

[0330] The toxicity and/or efficacy of the prophylactic and/or therapeutic protocols described herein can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Therapies that exhibit large therapeutic indices are preferred. While therapies that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such agents to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[0331] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage of the prophylactic and/or therapeutic agents for use in humans. The dosage of such agents lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any therapy used in the method, e.g., as described herein, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the Therapeutic Agent that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0332] In yet another embodiment, apoptotic cells are measured in both the attached and "floating" compartments of the cultures. Both compartments are collected by removing the supernatant, trypsinizing the attached cells, and combining both preparations following a centrifugation wash step (10 minutes, 2000 rpm).

[0333] In yet another embodiment, apoptosis is quantitated by measuring DNA fragmentation. Commercial photometric methods for the quantitative *in vitro* determination of DNA fragmentation are available. Examples of such assays, including TUNEL (which detects incorporation of labeled nucleotides in fragmented DNA) and ELISA-based assays, are described in Biochemica, 1999, no. 2, pp. 34-37 (Roche Molecular Biochemicals).

[0334] In yet another embodiment, apoptosis can be observed morphologically.

[0335] Cell lines on which such assays can be performed are well known to those of skill in the art. Apoptosis, necrosis and proliferation assays can also be performed on primary cells, e.g., a tissue explant.

## 6. EXAMPLES

[0336] Human 293 cells were transfected with nucleic acid expression construct for IL-15 (see, e.g., FIG. 5A-D) alone or in combination with a nucleic acid expression construct for IL-15sRa (see, e.g., FIG. 7A-D) or IL-15Ra (see, e.g., FIG. 6A-D) together with a plasmid conferring hygromycin resistance. IL15tPA indicates the optimized nucleic acid expression construct having the tPA prepro peptide (i.e., signal peptide) replacing the natural IL-15 secretory signal. "Ra" and "sRa" indicate the optimized expression vectors used for expression of the full-length IL-15Ra and the extracellular portion of IL-15Ra (soluble form), respectively. Transfected cells were treated with hygromycin (250 µg/ml), and rapidly growing resistant cell foci were isolated and expanded. Supernatants of the different clones were assayed for IL-15 expression after 2 days in culture by ELISA (R&D Systems Quantikine human IL-15 Elisa kit). IL-15 production is higher in cells receiving both genes (p=0.0166). Two measurements of IL-15 were determined for most clones.

Table 2:

IL-15/million cells (ng/ml)	clone#	genes
9.2	3.1	IL15tPA
14.0	3.1	IL15tPA
37.7	3.3	IL15tPA
74.0	3.3	IL15tPA

(continued)

	IL-15/million cells (ng/ml)	clone#	genes
5	147.2	7.5	IL15tPA+sRA
	136.8	6.4	IL15tPA+sRA
	266.2	6.4	IL15tPA+sRA
	135.9	6.5	IL15tPA+sRA
10	133.8	6.5	IL15tPA+sRA
	184.1	6.8	IL15tPA+sRA
	192.7	6.8	IL15tPA+sRA
	37.5	6.9	IL15tPA+sRA
15	37.8	6.9	IL15tPA+sRA
	101.3	7.2	IL15tPA+sRA
	176.0	7.2	IL15tPA+sRA
	294.5	7.21	IL15tPA+sRA
20	390.0	7.21	IL15tPA+sRA
	81.2	4.2	IL15tPA+RA
	194.2	4.2	IL15tPA+RA
25	54.0	4.4	IL15tPA+RA
	100.8	4.4	IL15tPA+RA
	18.3	5.6	IL15tPA+RA
	46.2	5.6	IL15tPA+RA
30	31.2	5.2	IL15tPA+RA
	22.5	5.2	IL15tPA+RA

[0337] Clone 7.21 was further cultured and selected for the ability to grow in serum-free medium in shake flasks and selected for cells with the ability to grow under such conditions. Briefly, the clones were cultured at a 1:1 medium ratio of old (10% fetal calf serum) to new media (serum-free) until two population doublings were observed. This procedure was repeated until cells that could be grown well in serum free medium were obtained. The serum free medium used was a 1:1 mix of 2 commercial media: (i) HyClone HyQ SFM4HEK293 (cat. # SH30521.02) and (ii) Invitrogen FreeStyle 293 (cat. # 12338-026). The typical yield of recombinant IL-15 produced by cells adapted to grow in serum-free medium in shake flasks from clone 7.21 is approximately 3 to 4 mg/L of medium and 0.6  $\mu$ g/10<sup>6</sup> cells as measured by ELISA (R&D Systems, Quantikine human IL-15 ELISA kit).

45

## 7. SPECIFIC EMBODIMENTS, CITATION AND REFERENCES

50

[0338] The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description and accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

[0339] Various references, including patent applications, patents, and scientific publications, are cited herein; the disclosure of each such reference is hereby incorporated herein by reference in its entirety.

55

### Certain Items

[0340]

1. A method for treating cancer in a human subject, comprising administering to a human subject in need thereof a composition comprising irradiated cancer cells engineered to recombinantly co-express (i) human IL-15 or a derivative thereof, and (ii) human IL-15 receptor alpha ("IL-15Ra") or a derivative thereof.

5 2. The method of item 1, wherein the irradiated cancer cells are generated using cancer cells obtained from the subject.

10 3. The method of item 1 or 2, wherein the irradiated cancer cells recombinantly express human IL-15 and human IL-15Ra or a derivative thereof.

15 4. The method of item 1 or 2, wherein the irradiated cancer cells recombinantly express a human IL-15 derivative thereof and human IL-15Ra.

15 5. The method of item 1 or 2, wherein the irradiated cancer cells recombinantly express a human IL-15 derivative and a human IL-15Ra derivative.

6. The method of any one of items 1 to 5, wherein the human IL-15Ra or human IL-15Ra derivative is soluble.

20 7. The method of any one of items 1 to 6, wherein the irradiated cancer cells further recombinantly express one or more other therapeutic polypeptides.

25 8. A method for treating an autoimmune disorder or inflammatory disorder in a human subject, comprising administering to a human subject in need thereof an effective amount of an antibody that specifically binds to IL-15/IL-15Ra complexes and reduces the binding of the IL-15/IL-15Ra complexes to the beta-gamma receptor complex as determined in cell culture or *in vitro*.

9. The method of item 8, wherein the antibody is a monoclonal humanized antibody.

30 10. A method for enhancing an IL-15-mediated immune function in a human subject, comprising administering to a human subject in need thereof a composition comprising an effective amount of an IL-15/IL-15Ra complex formulated with a poly- $\beta$ - $\rightarrow$ 4-N- acetylglucosamine polymer, wherein the IL-15/IL-15Ra complex comprises human IL-15 or a derivative thereof covalently or noncovalently linked to human IL-15Ra or a derivative thereof.

35 11. The method of item 10, wherein the IL-15/IL-15Ra complex comprises human IL-15 and human IL-15Ra or a derivative thereof.

12. The method of item 10, wherein the IL-15/IL-15Ra complex comprises a human IL-15 derivative thereof and human IL-15Ra.

40 13. The method of item 10, wherein the IL-15/IL-15Ra complex comprises a human IL-15 derivative and a human IL-15Ra derivative.

14. The method of any one of items 10 to 13, wherein the human IL-15Ra or human IL-15Ra derivative is soluble.

45 15. The method of any one of items 10 to 14, wherein said method further comprises administering to the human subject one or more other therapeutic polypeptides.

50 16. A method for making irradiated cancer cells recombinantly expressing human IL-15 and human IL-15Ra, said method comprises the steps of: (i) isolating cancer cells from a subject diagnosed with cancer; (ii) introducing a nucleic acid construct(s) encoding recombinant human IL-15 or a derivative thereof and human IL-15Ra or a derivative thereof; and (iii) irradiating said cancer cells.

17. The method of item 16, wherein the human IL-15Ra or human IL-15Ra derivative is soluble.

55 18. An irradiated cancer cell recombinantly expressing human IL-15 and human IL-15Ra produced by the method of item 16 or 17.

19. A pharmaceutical composition comprising the irradiated cancer cell of item 18.

**EP 2 724 727 A1**

20. The pharmaceutical composition of item 19, wherein the irradiated cancer cell is formulated with a poly- $\beta$ -1 $\rightarrow$ 4-N- acetylglucosamine polymer.

5 21. A cell that recombinantly expresses a mammalian IL-15 or a derivative thereof and a mammalian IL-15Ra or a derivative thereof, wherein the cell expresses at least 0.6 pg of mammalian IL-15 or a derivative thereof.

22. The cell of item 21, wherein the cell grows in serum-free media.

10

15

20

25

30

35

40

45

50

55

## SEQUENCE LISTING

5 <110> Admune Therapeutics LLC  
The United States Of America, As Represented By The Secretary,  
Department Of Health And Human Services, Office Of Technology  
Transfer National Institute Of Health

<120> COMPLEXES OF IL-15 AND IL-15RALPHA AND USES THEREOF

10 <130> 102385PCEPT1

<150> US20070937471  
<151> 2007-06-27

15 <160> 18

<170> FastSEO for Windows Version 4.0

20 <210> 1  
<211> 1  
<212> P  
<213> H

```

<220>
<223> immature/precursor form of native human IL-15

30 <400> 1
      Met Arg Ile Ser Lys Pro His Leu Arg Ser Ile Ser Ile Gln Cys Tyr
      1           5           10          15
      Leu Cys Leu Leu Leu Asn Ser His Phe Leu Thr Glu Ala Gly Ile His
      20          25          30
      Val Phe Ile Leu Gly Cys Phe Ser Ala Gly Leu Pro Lys Thr Glu Ala
      35          40          45
      Asn Trp Val Asn Val Ile Ser Asp Leu Lys Lys Ile Glu Asp Leu Ile
      50          55          60
      Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu Ser Asp Val His
      65          70          75          80
40   Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu Leu Glu Leu Gln
      85          90          95
      Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His Asp Thr Val Glu
      100         105         110
      Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser Asn Gly Asn Val
      115         120         125
45   Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu Glu Lys Asn Ile
      130         135         140
      Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln Met Phe Ile Asn
      145         150         155          160
      Thr Ser

```

50

<210> 2

<211> 489

<212> DNA

55 <213> *Homo sapiens*

<220>

<221> sig\_peptide  
 <222> (1)...(145)

5 <220>  
 <223> coding sequence of immature/precursor form of native human IL-15

10 <400> 2  
 atgagaattt ccaaaccaca tttgagaagt atttccatcc agtgcactt gtgtttactt 60  
 ctaaacagtc atttctaac tgaagctggc attcatgtct tcattttggg ctgtttcagt 120  
 gcagggcttc ctaaaacaga agccaactgg gtgaatgtaa taagtgattt gaaaaaaaaatt 180  
 gaagatctta ttcaatctat gcatattgtat gctactttat atacggaaag tgatgtcac 240  
 cccagttgca aagtaacagc aatgaagtgc tttctcttgg agttacaagt tatttcactt 300  
 gagtccggag atgcaagtat tcatgataca gtagaaaaatc tgatccatcct agcaaacaac 360  
 agtttgtctt ctaatggaa tgtaacagaa tctggatgca aagaatgtga ggaactggag 420  
 gaaaaaaaaata tttaagaatt tttgcagagt tttgtacata ttgtccaaat gttcatcaac 480  
 15 acttcttga 489

20 <210> 3  
 <211> 267  
 <212> PRT  
 <213> Homo sapiens

25 <220>  
 <221> SIGNAL  
 <222> (1)...(30)

30 <220>  
 <223> immature form of the native full length human IL-15  
 receptor alpha

<400> 3  
 Met Ala Pro Arg Arg Ala Arg Gly Cys Arg Thr Leu Gly Leu Pro Ala  
 1 5 10 15  
 Leu Leu Leu Leu Leu Leu Arg Pro Pro Ala Thr Arg Gly Ile Thr  
 20 25 30  
 Cys Pro Pro Pro Met Ser Val Glu His Ala Asp Ile Trp Val Lys Ser  
 35 40 45  
 Tyr Ser Leu Tyr Ser Arg Glu Arg Tyr Ile Cys Asn Ser Gly Phe Lys  
 50 55 60  
 Arg Lys Ala Gly Thr Ser Ser Leu Thr Glu Cys Val Leu Asn Lys Ala  
 65 70 75 80  
 Thr Asn Val Ala His Trp Thr Thr Pro Ser Leu Lys Cys Ile Arg Asp  
 85 90 95  
 40 Pro Ala Leu Val His Gln Arg Pro Ala Pro Pro Ser Thr Val Thr Thr  
 100 105 110  
 Ala Gly Val Thr Pro Gln Pro Glu Ser Leu Ser Pro Ser Gly Lys Glu  
 115 120 125  
 Pro Ala Ala Ser Ser Pro Ser Ser Asn Asn Thr Ala Ala Thr Thr Ala  
 130 135 140  
 45 Ala Ile Val Pro Gly Ser Gln Leu Met Pro Ser Lys Ser Pro Ser Thr  
 145 150 155 160  
 Gly Thr Thr Glu Ile Ser Ser His Glu Ser Ser His Gly Thr Pro Ser  
 165 170 175  
 Gln Thr Thr Ala Lys Asn Trp Glu Leu Thr Ala Ser Ala Ser His Gln  
 180 185 190  
 50 Pro Pro Gly Val Tyr Pro Gln Gly His Ser Asp Thr Thr Val Ala Ile  
 195 200 205  
 Ser Thr Ser Thr Val Leu Leu Cys Gly Leu Ser Ala Val Ser Leu Leu  
 210 215 220  
 Ala Cys Tyr Leu Lys Ser Arg Gln Thr Pro Pro Leu Ala Ser Val Glu  
 225 230 235 240  
 55 Met Glu Ala Met Glu Ala Leu Pro Val Thr Trp Gly Thr Ser Ser Arg  
 245 250 255

Asp Glu Asp Leu Glu Asn Cys Ser His His Leu  
 260 265

5           <210> 4  
 <211> 205  
 <212> PRT  
 <213> Homo sapines

10           <220>  
 <221> SIGNAL  
 <222> (1)...(30)

15           <220>  
 <223> immature form of the native soluble human IL-15 receptor alpha

20           <400> 4  
 Met Ala Pro Arg Arg Ala Arg Gly Cys Arg Thr Leu Gly Leu Pro Ala  
 1           5           10           15  
 Leu Leu Leu Leu Leu Leu Arg Pro Pro Ala Thr Arg Gly Ile Thr  
 20           25           30  
 Cys Pro Pro Pro Met Ser Val Glu His Ala Asp Ile Trp Val Lys Ser  
 35           40           45  
 Tyr Ser Leu Tyr Ser Arg Glu Arg Tyr Ile Cys Asn Ser Gly Phe Lys  
 50           55           60  
 Arg Lys Ala Gly Thr Ser Ser Leu Thr Glu Cys Val Leu Asn Lys Ala  
 65           70           75           80  
 25           Thr Asn Val Ala His Trp Thr Thr Pro Ser Leu Lys Cys Ile Arg Asp  
 85           90           95  
 Pro Ala Leu Val His Gln Arg Pro Ala Pro Pro Ser Thr Val Thr Thr  
 100           105           110  
 Ala Gly Val Thr Pro Gln Pro Glu Ser Leu Ser Pro Ser Gly Lys Glu  
 30           115           120           125  
 Pro Ala Ala Ser Ser Pro Ser Ser Asn Asn Thr Ala Ala Thr Thr Ala  
 130           135           140  
 Ala Ile Val Pro Gly Ser Gln Leu Met Pro Ser Lys Ser Pro Ser Thr  
 145           150           155           160  
 Gly Thr Thr Glu Ile Ser Ser His Glu Ser Ser His Gly Thr Pro Ser  
 35           165           170           175  
 Gln Thr Thr Ala Lys Asn Trp Glu Leu Thr Ala Ser Ala Ser His Gln  
 180           185           190  
 Pro Pro Gly Val Tyr Pro Gln Gly His Ser Asp Thr Thr  
 195           200           205

40

45           <210> 5  
 <211> 804  
 <212> DNA  
 <213> Homo sapiens

50           <220>  
 <221> sig\_peptide  
 <222> (1)...(90)

55           <220>  
 <223> coding sequence of immature form of the native full length  
 human IL-15 receptor alpha

            <400> 5  
 atggccccgc ggcggggcgcg cggctgccc accctcggtc tcccgccgtc gctactgctg 60  
 ctgctgctcc ggccggccggc gacgcggggc atcacgtgcc ctccccccat gtccgtggaa 120  
 cacgcagaca tctgggtcaa gagctacagc ttgtactcca gggagcggta catttgtaac 180  
 tctggtttca agcgtaaagc cggcacgtcc agcctgacgg agtgagtggtt gaacaaggcc 240

acgaatgtcg cccactggac aaccccccagt ctcaaatgca ttagagaccc tgccctgggt 300  
 caccaaaggc cagcggcacc ctccacagta acgacggcag gggtagccccc acagccagag 360  
 agcctctccc cttctggaaa agagcccgca gcttcatctc ccagctaaaa caacacagcg 420  
 gcccacaacag cagctattgt cccgggctcc cagctgatgc cttaaaaatc accttccaca 480  
 5 ggaaccacacag agataaagcag tcatgagtc tcccacggca cccctctca gacaacagcc 540  
 aagaactggg aactcacaacgc atccgcctcc caccagccgc caggtgtgta tccacagggc 600  
 cacagcgaca ccactgtggc tatctccacg tccactgtcc tgctgtgtgg gctgagcgt 660  
 gtgtctctcc tggcatgta cctcaagtca aggcaaaatc ccccgctggc cagcgttcaa 720  
 atggaagcca tggaggctct gccggtaact tgggggacca gcagcagaga tgaagacttg 780  
 gaaaactgct ctcaccaccc atga 804  
 10  
 <210> 6  
 <211> 615  
 <212> DNA  
 <213> Homo sapiens  
 15  
 <220>  
 <221> sig\_peptide  
 <222> (1)...(90)  
 <220>  
 20 <223> coding sequence of immature form of the native soluble human  
     IL-15 receptor alpha  
 <400> 6  
 atggcccccgc ggcggggcgcg cggctgccc accctcggtc tcccgccgt gctactgctg 60  
 ctgctgctcc ggccggccgc gacgcggggc atcacgtcc ctccccccat gtcggggaa 120  
 cacgcagaca tctgggtcaa gagctacagc ttgtactcca gggagccgt aattttaac 180  
 tctggttca agcgtaaagc cggcacgtcc agcctgacgg agtgcgtgtt gaacaaggcc 240  
 acgaatgtcg cccactggac aaccccccagt ctcaaatgca ttagagaccc tgccctgggt 300  
 caccaaaggc cagcggcacc ctccacagta acgacggcag gggtagccccc acagccagag 360  
 25 agcctctccc cttctggaaa agagcccgca gcttcatctc ccagctaaaa caacacagcg 420  
 gcccacaacag cagctattgt cccgggctcc cagctgatgc cttaaaaatc accttccaca 480  
 ggaaccacacag agataaagcag tcatgagtc tcccacggca cccctctca gacaacagcc 540  
 aagaactggg aactcacaacgc atccgcctcc caccagccgc caggtgtgta tccacaggc 600  
 cacagcgaca ccact 615  
 30  
 <210> 7  
 <211> 4  
 <212> PRT  
 <213> Artificial Sequence  
 <220>  
 35 <223> heterologous protease cleavage sites recognized by  
     furin protease  
 <220>  
 <221> VARIANT  
 <222> 2,3  
 <223> Xaa = any amino acid  
 <400> 7  
 Arg Xaa Xaa Arg  
 1  
 45  
 <210> 8  
 <211> 6  
 <212> PRT  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> heterologous protease cleavage sites recognized by

### thrombin protease

```
<220>
<221> VARIANT
<222> 1,2
<223> xaa = hydrophobic amino acids
```

<220>  
<221> VARIANT  
<222> 5,6  
<223> Xaa = nonacidic amino acids

<400> 8  
Xaa Xaa Pro Arg Xaa Xaa  
1 5

<210> 9  
<211> 1847  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> AG32 huIL15opt - nucleic acid construct encoding  
optimized human IL-15

<400> 9	cctggcatt	gcatacgttgcataccatatc	ataatatgtatccat	catttatatttatccat	ggctcatgtc	60
caacattacc	gccatgttga	cattgattat	tgactagttat	ttaatagtaat	tcaattacgg	120
ggtcattagt	tcataggccca	tatatggagt	tcggcggtac	ataacttacg	gtaaatggcc	180
cgcctggctg	accggccaaac	gaccccccgc	cattgacgtc	aataatgacg	tatgttccca	240
tagtaacgcc	aatagggact	ttccattgac	gtcaatgggt	ggagtattta	cggtaaactg	300
cccacttggc	agtacatcaa	gtgtatcata	tgccaagttac	gccccctatt	gacgtcaatg	360
atggtaaatg	gccccctgg	cattatgccc	agtacatgac	cttataggac	tttcctactt	420
ggcagtagat	ctacgttata	gtcatcgcta	ttaccatgg	gatgcgggtt	tggcagtagaca	480
tcaatggcg	tggatagcg	tttgcactac	ggggatttcc	aagtctccac	ccattgcacg	540
tcaatggag	tttgggg	caccaaaaatc	aacgggactt	tccaaaatgt	cgtacaact	600
ccgccccatt	gacgcattatg	ggcggttaggc	gtgtacgggt	ggaggcttat	ataagcagag	660
ctcggttagt	gaaccgtcag	atcgccctgg	gacgcattcc	acgctgtttt	gacccatata	720
gaagacacccg	ggaccgtatcc	agccctccgc	ggcgccgtc	gacaagaaaat	gcggatctcg	780
aagccgcacc	tgcgttcgtat	atcgatccag	tgctacctgt	gcctgctcct	gaactcgac	840
ttccctacgg	aggccggat	acacgttcc	atcctggct	gcttctcggt	ggggctggcc	900
aagacggagg	cgaactgggt	gaacgtgatc	tcggacctga	agaagatcg	ggacccatc	960
cagtgcgtgc	acatcgacgc	gacgcgtgtac	acggagtcgg	acgtccaccc	gtcggtcaag	1020
gtcacggca	tgaagtgtt	cctcctggag	ctccaagtca	tctcgctcg	gtcgggggac	1080
gcgtcgatcc	acgacacgg	ggagaacctg	atcatcctgg	cgaacaactc	gctgtcg	1140
aacgggaacg	tcacggagtc	gggctgcaag	gagtgcgagg	agctggagga	gaagaacatc	1200
aaggagttcc	tgcgttcgtt	cgtgcacatc	gtccagatgt	tcatcaacac	gtcggtaggg	1260
ccggcgcgc	cgaattcgcg	gatatcggtt	aacggatcca	gatctgtgt	gccttctagt	1320
tgccagccat	ctgtgttttgc	cccccccccc	gtgccttcct	tgaccctgg	aggtggcact	1380
cccactgtcc	tttccataata	aaatgaggaa	attgcacatgc	attgtctgag	taggtgtcat	1440
tctattctgg	gggggggggt	ggggcaggac	agcaaggggg	aggattggga	agacaatagc	1500
aggcatgtcg	gggatgcgg	gggctctatg	ggtacccagg	tgctgaagaa	ttgaccgg	1560
tcctcctgg	ccagaaagaa	gcaggccat	cccccttcct	gtgacacacc	ctgtccacgc	1620
ccctggttct	tagtccagc	cccactcata	ggacactcat	agtcaggag	ggctccgcct	1680
tcaatccac	ccgcattaaatg	acttggagcg	gtctctccct	ccctcatcg	cccaccaaac	1740
caaaccatgc	ctccaagatg	ggaaagaaat	taaagcaaga	taggtattt	agtgcagagg	1800
gagagaaaaat	qcctccaaca	tgtqagggaaq	taatqagaga	aatcata		1847

<210> 10  
<211> 162  
<212> PRT  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; AG32 huIL15opt - amino acid sequence of optimized human IL-15

&lt;400&gt; 10

5 Met Arg Ile Ser Lys Pro His Leu Arg Ser Ile Ser Ile Gln Cys Tyr

1 5 10 15

Leu Cys Leu Leu Leu Asn Ser His Phe Leu Thr Glu Ala Gly Ile His

20 25 30

Val Phe Ile Leu Gly Cys Phe Ser Ala Gly Leu Pro Lys Thr Glu Ala

35 40 45

10 Asn Trp Val Asn Val Ile Ser Asp Leu Lys Ile Glu Asp Leu Ile

50 55 60

Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu Ser Asp Val His

65 70 75 80

Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu Leu Glu Leu Gln

85 90 95

15 Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His Asp Thr Val Glu

100 105 110

Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser Asn Gly Asn Val

115 120 125

20 Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu Glu Lys Asn Ile

130 135 140

Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln Met Phe Ile Asn

145 150 155 160

Thr Ser

25

&lt;210&gt; 11

&lt;211&gt; 1808

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

30

&lt;220&gt;

<223> AG59 CMV huIL15tPA6 - nucleic acid construct encoding  
optimized human IL-15

&lt;400&gt; 11

35 cctggccatt gcatacggtt tatccatatac ataataatgtt catttatattt ggctcatgtc 60

caacattacc gccatgttga cattgattat tgacttagttt ttaatagttt tcaattacgg 120

ggtcattagt tcatacgccca tataatggagt tccgcgttac ataacttacg gttaatggcc 180

cgccctggctg accgccccaaac gaccccccggc cattgacgtc aataatgacg tatgttccca 240

tagtaacgccc aataggggact ttccattgtac gtcaatgggtt ggagtattta cggtaaaactg 300

cccacttggc agtacatcaa gtgtatcata tgccaatgtc gccccctattt gacgtcaatg 360

atggtaaatg gcccccttgg cattatgtccc agtacatgtac cttatgggac tttcctactt 420

ggcagttacat ctacgttataa gtcatcgcta ttaccatgggtt gatgcgggtt tggcagttaca 480

tcaatggggcg tggatagcgg tttgactcac ggggattttcc aagtctccac cccattgtacg 540

tcaatggggag tttgttttgg caccaaaaatc aacggggactt tccaaaatgtt cgtaacaact 600

ccgccccattt gacgtcaatg ggcggtaggc gtgtacgggtt ggaggcttatataaaggcag 660

ctcggttagt gaaccgtcag atgccttggc gacgtccatcc acgtgtttt gaccccttata 720

gaagacacccg ggaccgtatcc agcctccgcg ggccgcgttc gacaagaaat ggatgtcaatg 780

aagagagggc tctgtctgt gctgtctgt tttggagcag tcttcgtttt gcccagccag 840

gaaatccatg cccgatttgc aagaggagcc agaaaactggg tgaacgttat ctcggaccc 900

aagaagatcg aggacccatc ccagtcgtatc cacatcgacg cgacgtgtt cacggagtcg 960

50 gacgtccacc cgctgtccaa ggtcacggcg atgaagtgtt tccttcgtt gctccaagtc 1020

atctcgctcg agtcggggcg cgcgtcgatc cacgacacgg tggagaacctt gatcatccgt 1080

gccaacaactt cgctgtcgatc gaaacgggaaac gtcacgggtt cgggtgtt ggtgtgcgag 1140

gagctggagg agaagaacat caaggagttt ctgcgtgtt tggatgttgcacat cgtccagatg 1200

ttccatcaaca cgtcggtt gcccggcgcc cccatttgcg ggtatgttgcgtt taacggatcc 1260

55 agatctgtgt tgccttctat ttggccagccaa tctgttgcgtt gcccctcccc cgtgccttc 1320

ttgacccttgg aaggtgccac tccactgttccataaaatgttgcgtt aattgtcatcg 1380

cattgtctga gtaggtgtca ttctatttgcgtt ggggggtgggg tggggcagggc aagcaagggg 1440

5 gaggattggg aagacaatag caggcatgct gggatgcgg tggcttat ggttacccag 1500  
 gtgctgaaga attgaccgg ttcctcctgg gccagaaaga agcaggcaca tccccttc 1560  
 tgtgacacac cctgtccacg cccctggttc ttagttccag cccactcat agacactca 1620  
 tagctcagga gggctccgc ttcaatccca cccgctaaag tacttgagc ggtctctccc 1680  
 tccctcatca gcccacccaaa ccaaaccctag cctccaagag tggagaaa taaaagcaag 1740  
 ataggctatt aagtgcagag ggagagaaa tgcctccaac atgtgaggaa gtaatgagag 1800  
 aaatcata 1808

10 <210> 12  
 <211> 149  
 <212> PRT  
 <213> Artificial Sequence

15 <220>  
 <223> AG59 CMV huIL15tPA6 - amino acid sequence of optimized human IL-15

20 <400> 12  
 Met Asp Ala Met Lys Arg Gly Leu Cys Cys Val Leu Leu Cys Gly  
 1 5 10 15  
 Ala Val Phe Val Ser Pro Ser Gln Glu Ile His Ala Arg Phe Arg Arg  
 20 25 30  
 Gly Ala Arg Asn Trp Val Asn Val Ile Ser Asp Leu Lys Lys Ile Glu  
 35 40 45  
 Asp Leu Ile Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu Ser  
 50 55 60  
 Asp Val His Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu Leu  
 65 70 75 80  
 25 Glu Leu Gln Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His Asp  
 85 90 95  
 Thr Val Glu Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser Asn  
 100 105 110  
 Gly Asn Val Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu Glu  
 30 115 120 125  
 Lys Asn Ile Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln Met  
 130 135 140  
 Phe Ile Asn Thr Ser  
 145

35 <210> 13  
 <211> 2140  
 <212> DNA  
 <213> Artificial Sequence

40 <220>  
 <223> AG79 huIL15Ra - nucleic acid construct encoding  
 optimized human IL-15Ra

45 <400> 13  
 cctggccatt gcatacggtt tatccatatac ataatatgtt catttatatt ggctcatgtc 60  
 caacattacc gccatgttga cattgattat tgactagttt ttaatagtaa tcaattacgg 120  
 ggtcattagt tcatacgcca tatagggtt tccgcgttac ataacttacg gtaaatggcc 180  
 cgcctggctg accgcccacac gaccccccgc cattgacgtc aataatgacg tatgttccca 240  
 tagtaacgcc aataggact ttccattgtac gtcaatgggt ggagtattta cgtaaaactg 300  
 cccacttggc agtacatcaa gtgtatcata tgccaaagtac gccccctatt gacgtcaatg 360  
 50 atggtaaatg gccccctgg cattatgccc agtacatgtc cttatggac tttcctactt 420  
 ggcagtacat ctacgttata gtcatcgcta ttaccatggt gatgcgggtt tggcagtaca 480  
 tcaatggcg tggatagccg tttgactcac ggggatttcc aagtctccac cccattgacg 540  
 tcaatggag ttgtttgg caccaaaatc aacggggactt tccaaaatgt cgttacaact 600  
 55 ccgccttgcatt gacgcataatg ggcggtaggc gtgtacgggt ggaggcttat ataagcagag 660  
 ctcgttttagt gaaccgtcag atcgcttggaa gacgccttcc acgctgtttt gacctccata 720  
 gaagacaccg ggaccgatcc agcctccgcg ggcgcgcgtc gacgcttagca agaaatggcc 780  
 ccgaggcggg cgcgaggctg ccggaccctc ggtatcccg gctgtact gctcctgctg 840

ctccggccgc cggcgacgcg gggcatcacg tgccgcgc ccatgtccgt ggagcacgca 900  
 gacatctgg tcaagagcta cagcttgcac tcccgccgac ggtacatctg caactcggt 960  
 ttcaagcgga aggccgcac gtccagcctg acggagtcg tggtaacaa gcccacgaa 1020  
 gtcgcccact ggacgacccc ctcgctcaag tgcacccgcg accccgcct gttcaccag 1080  
 5 cggccgcgc caccctccac cgtAACGACG gcgggggtga ccccgccagcc ggagagcctc 1140  
 tccccgtcg gaaaggagcc cgccgcgtcg tcgcccagct cgaacaacac ggcggccaca 1200  
 actcgacga tcgtccggg ctcccagctg atgcgtcga agtcgcgtc cacggaaacc 1260  
 acggagatca gcagtcatga gtctccac ggcacccccc cgaacacgac gccaagaac 1320  
 tgggaactca cggcgtccgc ctcccaccc cggccggggg tggatccgca aggccacagc 1380  
 gacaccacgg tggcgtctc cagtcacg gtccgtcg tgggctgag cgccgtgtcg 1440  
 10 ctctggcg tctaccaa gtcaggcgactccccc tggcagcgt tgagatggag 1500  
 gccatggagg ctctggcg gacgtggggg accagcagca gggatgagga cttggagaac 1560  
 tgctcgacc acctataatg agaattcgat ccagatctc tggccctct agttgcacgc 1620  
 catctgttgc ttgcccctcc cccgtgcctt ccttgcaccc ggaaggtgcc actccactg 1680  
 tccttcata ataaaatgag gaaattgcat cgcattgtct gaggatgtt cattctattc 1740  
 15 tgggggggtgg ggtggggcag gacagcaagg gggaggattt ggaagacaat agcaggcatg 1800  
 ctggggatgc ggtgggtct atgggtaccc aggtgtcga gaattgaccc gttccctcc 1860  
 gggccagaaa gaagcaggca catccccctc tctgtgacac accctgtcca cgcggctgg 1920  
 tcttagttcc agcccaactc ataggacact catagctcag gaggctccg cttcaatcc 1980  
 caccgcataa agtacttggc gcgtctctc cttccat cagcccacca accaaacct 2040  
 agcctccaag agtggaaaga aatcaaagca agataggta ttaagtgcag agggagagaa 2100  
 20 aatgcctcca acatgtgagg aagtaatgag agaaatcata 2140

<210> 14  
 <211> 267  
 <212> PRT  
 <213> Artificial Sequence  
 25  
 <220>  
 <223> AG79 huIL15Ra - amino acid sequence of optimized human IL-15Ra

<400> 14  
 30 Met Ala Pro Arg Arg Ala Arg Gly Cys Arg Thr Leu Gly Leu Pro Ala  
 1 5 10 15  
 Leu Leu Leu Leu Leu Leu Arg Pro Pro Ala Thr Arg Gly Ile Thr  
 20 25 30  
 Cys Pro Pro Pro Met Ser Val Glu His Ala Asp Ile Trp Val Lys Ser  
 35 40 45  
 Tyr Ser Leu Tyr Ser Arg Glu Arg Tyr Ile Cys Asn Ser Gly Phe Lys  
 50 55 60  
 Arg Lys Ala Gly Thr Ser Ser Leu Thr Glu Cys Val Leu Asn Lys Ala  
 65 70 75 80  
 Thr Asn Val Ala His Trp Thr Thr Pro Ser Leu Lys Cys Ile Arg Asp  
 85 90 95  
 40 Pro Ala Leu Val His Gln Arg Pro Ala Pro Pro Ser Thr Val Thr Thr  
 100 105 110  
 Ala Gly Val Thr Pro Gln Pro Glu Ser Leu Ser Pro Ser Gly Lys Glu  
 115 120 125  
 Pro Ala Ala Ser Ser Pro Ser Ser Asn Asn Thr Ala Ala Thr Thr Ala  
 130 135 140  
 45 Ala Ile Val Pro Gly Ser Gln Leu Met Pro Ser Lys Ser Pro Ser Thr  
 145 150 155 160  
 Gly Thr Thr Glu Ile Ser Ser His Glu Ser Ser His Gly Thr Pro Ser  
 165 170 175  
 Gln Thr Thr Ala Lys Asn Trp Glu Leu Thr Ala Ser Ala Ser His Gln  
 180 185 190  
 Pro Pro Gly Val Tyr Pro Gln Gly His Ser Asp Thr Thr Val Ala Ile  
 195 200 205  
 Ser Thr Ser Thr Val Leu Leu Cys Gly Leu Ser Ala Val Ser Leu Leu  
 210 215 220  
 50 Ala Cys Tyr Leu Lys Ser Arg Gln Thr Pro Pro Leu Ala Ser Val Glu  
 225 230 235 240  
 Met Glu Ala Met Glu Ala Leu Pro Val Thr Trp Gly Thr Ser Ser Arg

245	250	255
Asp Glu Asp Leu Glu Asn Cys Ser His His Leu		
260	265	

	Cys Pro Pro Pro Met Ser Val Glu His Ala Asp Ile Trp Val Lys Ser
	35 40 45
	Tyr Ser Leu Tyr Ser Arg Glu Arg Tyr Ile Cys Asn Ser Gly Phe Lys
	50 55 60
5	Arg Lys Ala Gly Thr Ser Ser Leu Thr Glu Cys Val Leu Asn Lys Ala
	65 70 75 80
	Thr Asn Val Ala His Trp Thr Thr Pro Ser Leu Lys Cys Ile Arg Asp
	85 90 95
	Pro Ala Leu Val His Gln Arg Pro Ala Pro Pro Ser Thr Val Thr Thr
10	100 105 110
	Ala Gly Val Thr Pro Gln Pro Glu Ser Leu Ser Pro Ser Gly Lys Glu
	115 120 125
	Pro Ala Ala Ser Ser Pro Ser Ser Asn Asn Thr Ala Ala Thr Thr Ala
	130 135 140
15	Ala Ile Val Pro Gly Ser Gln Leu Met Pro Ser Lys Ser Pro Ser Thr
	145 150 155 160
	Gly Thr Thr Glu Ile Ser Ser His Glu Ser Ser His Gly Thr Pro Ser
	165 170 175
	Gln Thr Thr Ala Lys Asn Trp Glu Leu Thr Ala Ser Ala Ser His Gln
	180 185 190
20	Pro Pro Gly Val Tyr Pro Gln Gly His Ser Asp Thr Thr
	195 200 205

	<210> 17
	<211> 1754
	<212> DNA
25	<213> Artificial Sequence
	<220>
	<223> AG151 huIL-15 huGM-CSF - nucleic acid construct encoding optimized human IL-15 with a signal peptide of human GM-CSF
30	<400> 17
	cctggccatt gcatacgttg tattccatatac ataatatgtt catttatatt ggctcatgtc 60 caacattacc gccatgttga cattgattat tgactagttt ttaatagttt tcaattacgg 120 ggtcattatgt tcataggccca tataatggatg tccgcgttac ataaatctacg gtaaatggcc 180 cgccctggctg accgccccac gaccccccgc cattgacgtc aataatgacg tatgttccca 240 tagtaacgc aataggggact ttccattgtac gtcaatgggt ggagtttta cggtaaaactg 300 cccaacttggc agtacatcaa gtgtatcata tgccaaagtac gccccctatt gacgtcaatg 360 atggtaaatgg cccgcctgg cattatggcc agtacatgtac cttatgggac tttcctactt 420 ggcagttacat ctacgttata gtcatcgcc ttacatggat gatgcgggtt tggcagtaca 480 tcaatggcg tggatagcgg tttgactcac ggggattttcc aagtctccac cccattgacg 540 tcaatggcg tttgtttgg cacaaaaatc aacggggactt tccaaaatgt ctgtacaact 600 ccgccccatt gacgcaatg ggccgttaggc gtgtacgggt ggaggcttat ataaggcagag 660 ctcgtttagt gaaccgtcag atccctggc gacgcatttc acgctgtttt gacctccata 720 gaagacacccg ggaccgatcc agctccgcg ggcgcgcgtc gacaagaaat gtggctccag 780 agcctgtac tcctggggac ggtggcctgc agcatctcg actgggtgaa cgtgatctcg 840 gacctgaaga agatcgagga cctcatccag tcgtatcaca tcgacgcgac gctgtacacg 900 gagtccggacg tccaccggcgtc gtgcacaggc acggcgatgtc agtgccttgcct cctggagcgtc 960 caagtcatct cgctcgagtc gggggacgcg tcgatccacg acacgggttga gaacctgatc 1020 atccctggcga acaactcgct gtctcgaaac gggAACGTC cggagtcggg ctgcaaggag 1080 tgcgaggagc tggaggagaa gaacatcaag gagttccctgc agtcgttcgt gcacatcggtc 1140 cagatgttca tcaacacgtc gtgaggggccc ggcgcgcgtc attcgccat atcggttaac 1200 ggatcccgat ctgctgtgcc ttctagttgc cagccatctg ttgtttggccc ctcccccgtg 1260 ccttccttgc ccctggaaagg tgcactccctt actgtccctt cctaataaaaa tgagggaaatt 1320 gcatcgcatgt ctgtcgatgt gtgtcattctt attctggggg gtgggggtggg gcaggacacgc 1380 aaggggggagg attgggaaga caatagcagg catgctgggg atgcgggtggg ctctatgggt 1440 acccagggtgc tgaagaatttgc acccggttcc tccctggggca gaaagaagca ggcacatccc 1500 cttctctgttgc acacaccctg tccacgcggcc tgggttcttag ttccagcccc actcataggg 1560 cactcatagtc tcaggaggggc tccgccttca atccccccctt ctaaagtact tggagcgggtc 1620 tctccctccc tcatcgcccc accaaaaccaa acctagccctc caagagtggg aagaaaattaa 1680 agcaagatag gctttaatgtt gcagaggag agaaaatgcc tccaaatgtt gaggaagttaa 1740

tgagagaaat cata

1754

&lt;210&gt; 18

&lt;211&gt; 131

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> AG151 huIL-15 huGM-CSF - amino acid sequence of optimized  
10 human IL-15 with a signal peptide of human GM-CSF

&lt;400&gt; 18

Met Trp Leu Gln Ser Leu Leu Leu Gly Thr Val Ala Cys Ser Ile  
1 5 10 15Ser Asn Trp Val Asn Val Ile Ser Asp Leu Lys Lys Ile Glu Asp Leu  
15 20 25 30Ile Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu Ser Asp Val  
35 40 45His Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu Leu Glu Leu  
20 50 55 60Gln Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His Asp Thr Val  
65 70 75 80Glu Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser Asn Gly Asn  
85 90 95Val Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu Glu Lys Asn  
25 100 105 110Ile Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln Met Phe Ile  
115 120 125Asn Thr Ser  
130

30

**Claims**

1. A complex for use in a method of treating cancer in a subject, wherein

35

(A) the complex comprises: (i) interleukin 15 (IL-15); and (ii) a soluble form of interleukin 15 receptor alpha (IL-15Ra), wherein the complex is administered with an antibody that specifically binds to a T cell receptor or a tumor-associated antigen;

40

(B) the complex comprises: (i) a fusion protein comprising IL-15 and a soluble form of IL-15Ra; and (ii) an antibody that specifically binds to a cancer antigen; or

(C) the complex comprises: (i) IL-15; (ii) a soluble form of IL-15Ra, or a fusion protein comprising a soluble form of IL-15Ra and the Fc domain of an immunoglobulin or a fragment thereof; and (iii) an antibody that specifically binds to a cancer antigen.

45

2. The complex for use of claim 1(C), wherein the complex comprises (i) IL-15; (ii) a soluble form of IL-15 receptor alpha, and (iii) an antibody that specifically binds to a cancer antigen.

50

3. The complex for use of claim 1(C), wherein the complex comprises (i) IL-15, (ii) a fusion protein comprising a soluble form of IL-15Ra and the Fc domain of an immunoglobulin or a fragment thereof; and (iii) an antibody that specifically binds to a cancer antigen.

4. The complex for use of claim 1(B), wherein the fusion protein comprises a soluble form of IL-15Ra, a linker, and IL-15.

55

5. The complex for use of claim 4, wherein the linker is 5, 10, 15, 16, 17, 18, 19 or 20 amino acids long.

6. The complex for use of claim 1(B), 1(C), 2, 3, 4, or 5, wherein the antibody that specifically binds to a cancer antigen is fused or linked to the IL-15 or the soluble form of IL-15Ra.

7. The complex for use of claim 1(B), 1(C), 2, 3, 4, 5, or 6, wherein the cancer antigen is prostate specific membrane antigen, carcinoembryonic antigen (CEA), CD19, ganglioside GD2, ganglioside GD3, epidermal growth factor receptor (EGFR), or HER2.
- 5 8. The complex for use of claim 1(B), 1(C), 2, 3, 4, 5, or 6, wherein the antibody that specifically binds to a cancer antigen is an anti-CD20 antibody, anti-CD19 antibody, anti-EGFR antibody, or anti-HER2 antibody.
9. The complex for use of claim 1(A), wherein the antibody that specifically binds to a T cell receptor or a tumor-associated antigen is trastuzumab (Herceptin®) or rituximab (Rituxan®).
- 10 10. The complex for use of any one of the preceding claims, wherein the cancer is a leukemia, lymphoma, breast cancer, brain tumor, thyroid cancer, pancreatic cancer, skin cancer, lung cancer, uterine cancer, ovarian cancer, stomach cancer, colon cancer, testicular cancer, prostate cancer, renal cancer, bladder cancer, melanoma, non-small cell lung cancer, multiple myeloma, acute myelocytic leukemia, or non-Hodgkin lymphoma.
- 15 11. The complex for use of any one of the preceding claims, wherein the soluble form of IL-15Ra comprises exon 2 of IL-15Ra, exon 2 and the hinge region of IL-15Ra, or the extracellular domain of IL-15Ra.
12. The complex for use of any one of the preceding claims, wherein the IL-15 is human IL-15.
- 20 13. The complex for use of any one of the preceding claims, wherein the IL-15:
  - (a) has an amino acid sequence that is at least 95% identical to the amino acid sequence consisting of amino acid residues 49 to 162 of SEQ ID NO:1;
  - (b) is encoded by a nucleic acid sequence that is at least 80% identical to SEQ ID NO:2;
  - (c) is encoded by a nucleic acid sequence that is at least 80% identical to the nucleic acid sequence consisting of nucleic acid residues 145 to 489 of SEQ ID NO:2;
  - (d) has the amino acid sequence consisting of amino acid residues 49 to 162 of SEQ ID NO:1;
  - (e) is encoded by SEQ ID NO:2;
  - 30 (f) is encoded by the nucleic acid sequence consisting of nucleic acid residues 145 to 489 of SEQ ID NO:2;
  - (g) is encoded by the nucleic acid sequence consisting of nucleic acid residues 769 to 1257 of SEQ ID NO:9;
  - (h) is encoded by the nucleic acid sequence consisting of nucleic acid residues 769 to 1218 of SEQ ID NO:11; or
  - (i) is encoded by the nucleic acid sequence consisting of nucleic acid residues 769 to 1164 of SEQ ID NO:17.
- 35 14. The complex for use of any one of the preceding claims, wherein the soluble form of IL-15Ra is a soluble form of human IL-15Ra.
15. The complex for use of any one of the preceding claims, wherein the soluble form of IL-15Ra:
  - (a) has an amino acid sequence that is at least 95% identical to the amino acid sequence consisting of amino acid residues 31 to 205 of SEQ ID NO: 4;
  - (b) is encoded by a nucleic acid sequence that is at least 80% identical to SEQ ID NO:6;
  - (c) is encoded by a nucleic acid sequence that is at least 80% identical to the nucleic acid sequence consisting of nucleic acid residues 91 to 615 of SEQ ID NO:6;
  - 45 (d) has the amino acid sequence consisting of amino acid residues 31 to 205 of SEQ ID NO: 4;
  - (e) is encoded by SEQ ID NO:6;
  - (f) is encoded by the nucleic acid sequence consisting of nucleic acid residues 91 to 615 of SEQ ID NO:6; or
  - (g) is encoded by the nucleic acid sequence consisting of nucleic acid residues 775 to 1395 of SEQ ID NO:15.
- 50 16. The complex for use of any one of the preceding claims, wherein the subject is human.

native human IL-15

ATGAGAAT TTCGAAACCA CATTGAGAA GTATTCAT CCAGTGCTAC  
TTGTGTTAC TTCTAACAG TCATTTCTA ACTGAAGCTG GCATTCATGT  
CTTCATTTG GGCTGTTCA GTGCAGGGCT TCCTAAAACA GAAGCCAACT  
GGGTGAATGT AATAAGTGAT TTGAAAAAAA TTGAAGATCT TATTCAATCT  
ATGCATATTG ATGCTACTTT ATATACGGAA AGTGATGTTG ACCCCAGTTG  
CAAAGTAACA GCAATGAAGT GCTTCTCTT GGAGTTACAA GTTATTCAC  
TTGAGTCCGG AGATGCAAGT ATTCATGATA CAGTAGAAAA TCTGATCATC  
CTAGCAAACA ACAGTTGTC TTCTAATGGG AATGTAACAG AATCTGGATG  
CAAAGAATGT GAGGAAGTGG AGGAAAAAAA TATTAAGAA TTTTGAGA  
GTTTGTACA TATTGTCCAA ATGTTCATCA ACACCTCTTG A

**Fig. 1A**

MRISKPHLRSISIQCYLCLLNSHFLTEAGIHVFILGCFSA~~GLPKTEANWVNV~~  
ISDLKKIEDLIQSMHIDATLYTESDVHPSCKVTAMKCFLLELQVISLESGDASIHD  
VENLILANNSLSSNGNVTESGCKECEELEEKNIKEFLQSFVHIVQMFINTS

**Fig. 1B**

native human IL-15Ra

ATGGCCCC GCGGGGGCG CGGGCTGCC GGACCTCGG TCTCCGGCG  
CTGCTACTGC TGCTGCTGCT CGGGCGCCG GCGACGCGGG GCATCACGTG  
CCCTCCCCC ATGTCCGTGG AACACGCAGA CATCTGGTC AAGAGCTACA  
GCTTGTACTC CAGGGAGCGG TACATTTGTA ACTCTGGTT CAAGCGTAAA  
GCCGGCACGT CCAGCCTGAC GGAGTGCCTG TTGAACAAAGG CCACGAATGT  
CGCCCACGG ACAACCCCCA GTCTCAAATG CATTAGAGAC CCTGCCCTGG  
TTCACCAAAG GCCAGCGCCA CCCTCCACAG TAACGACGGC AGGGGTGACC  
CCACAGCCAG AGAGCCTCTC CCCTTCTGGA AAAGAGCCCG CAGCTTCATC  
TCCCAGCTCA AACAACACAG CGGCCACAAC AGCAGCTATT GTCCCGGGCT  
CCCAGCTGAT GCCTTCAAAA TCACCTTCCA CAGGAACCAC AGAGATAAGC  
AGTCATGAGT CCTCCCACGG CACCCCTCT CAGACAACAG CCAAGAACTG  
GGAACTCACA GCATCCGCCT CCCACCAGCC GCCAGGTGTG TATCCACAGG  
GCCACAGCGA CACCACTGTG GCTATCTCCA CGTCCACTGT CCTGCTGTGT  
GGGCTGAGCG CTGTGCTCT CTTGGCATGC TACCTCAAGT CAAGGCAAAC  
TCCCCCGCTG GCCAGCGTTG AAATGGAAGC CATGGAGGCT CTGCCGGTGA  
CTTGGGGGAC CAGCAGCAGA GATGAAGACT TGGAAAATG CTCTCACCAC  
CTATGA

**Fig. 2A**

MAPRRARGCRTLGLPALLLLLLRPPATRGITCPPPMSVEHADIWVKSYSLYSRERYICN  
SGFKRKAGTSSLTECVLNKATNVAHWTPSLCIRDPALVHQRPAAPPSTVTTAGVTPQES  
LSPSGKEPAASSPSSNNTAATTAAIVPGSQLMPSKSPSTGTTEISSHESSHGTPSQTAKNWE  
LTASASHQPPGVYPQGHSDTTVAISTSTVLLCGLSAVSLACYLKSRQTPPLASVEMEAMEA  
LPVTWGTSSRDEDLENCSHHL

**Fig. 2B**

native soluble human IL-15Ra

ATGGCCCC GCGGCGGGCG CGCGGCTGCC GGACCCTCGG TCTCCCGCGC  
CTGCTACTGC TGCTGCTGCT CCGGCCGCG GCGACGCGGG GCATCACGTG  
CCCTCCCCCC ATGTCCGTGG AACACGCAGA CATCTGGGTC AAGAGCTACA  
GCTTGTACTC CAGGGAGCGG TACATTTGTA ACTCTGGTTT CAAGCGTAAA  
GCCGGCACGT CCAGCCTGAC GGAGTGCGTG TTGAACAAAGG CCACGAATGT  
CGCCCACGTG ACAACCCCCA GTCTCAAATG CATTAGAGAC CCTGCCCTGG  
TTCACCAAAG GCCAGCGCCA CCCTCCACAG TAACGACGGC AGGGGTGACC  
CCACAGCCAG AGAGCCTCTC CCCTTCTGGA AAAGAGCCCG CAGCTTCATC  
TCCCAGCTCA ACAACACAG CGGCCACAAC AGCAGCTATT GTCCCGGGCT  
CCCAGCTGAT GCCTTCAAAA TCACCTTCCA CAGGAACCAC AGAGATAAGC  
AGTCATGAGT CCTCCCACGG CACCCCCCTCT CAGACAACAG CCAAGAACTG  
GGAACTCACA GCATCCGCCT CCCACCAGCC GCCAGGTGTG TATCCACAGG  
GCCACAGCGA CACCACT

**Fig. 3A**

MAPRRARGCRTLGLPALLLLLLRPPATRGITCPPPMSVEHADIWVKSYSLYSRERYICN  
SGFKRKAGTSSLTECVLNKATNVAHWTPSLKCIRDPAVLHQRPAPPSTVTTAGVTPQPE  
LSPSGKEPAASSPSSNNATAATTAAIVPGSQLMPSKSPSTGTTEISSHESSHGTPSQTAKNWE  
LTASASHQPPGVYPQGHSDTT

**Fig. 3B**

## AG32 huIL15opt

CCTGGCCATTGCATACGTTGATCCATATCATAATATGTACATTATGGCTCATGTCCA  
 ACATTACCGCCATGTTGACATTGATTATTGACTAGTTATTAAATAGTAATCAATTACGGGGTC  
 ATTAGTTCATAGCCCATAATGGAGTTCCCGCTTACATAACTACGGTAATGGCCCGCCTG  
 GCTGACCGCCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACG  
 CCAATAGGGACTTCCATTGACGTCAATGGGTGGAGTATTACGGTAAACTGCCACTTGGC  
 AGTACATCAAGTGTATCATATGCCAAGTACGCCCTATTGACGTCAATGATGGTAAATGGC  
 CCGCCTGGCATTATGCCAGTACATGACCTTATGGGACTTCCACTTGGCAGTACATCTAC  
 GTATTAGTCATCGCTATTACCATGGTATGCGGTTTGGCAGTACATCAATGGCGTGGATA  
 GCGGTTTGACTCACGGGATTCCAAGTCTCCACCCATTGACGTCAATGGGAGTTGTTT  
 GGCACCAAAATCAACGGACTTCCAAAATGTCGTAACAACCTCCACGGGCGGTATACACGTCT  
 GGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTAGTGAACCGTCAGAT  
 CGCCTGGAGACGCCATCCACGCTGTTTGACCTCCATAGAACAGACCCGGGACCGATCCAGCC  
 TCCCGGGGcgcgcgacaaga **ATGC GGATCTCGAAGCCGACCTGGCGTCGATATCGAT**  
**CCAGTGCTACCTGTGCCTGCTCTGA**ACTCGCACTTCCACGGAGGCGGTATACACGTCT  
**TCATCCTGGGCTGCTCTCGC**GGGGCTGCCAAGACGGAGGCAACTGGGTGAACGTGATC  
 TCGGACCTGAAGAAGATCGAGGACCTCATCCAGTCGATGCACATCGACGCGACGCTGTACAC  
 GGAGTCGGACGTCCACCCGTCGTGCAAGGTACGGCGATGAAGTGCCTCCTGGAGCTCC  
 AAGTCATCTCGCTCGAGTCGGGGACGCGTCGATCCACGACACGGTGGAGAACCTGATCATC  
 CTGGCGAACAACTCGCTGTCGAACGGGAACGTACGGAGTCGGGCTGCAAGGAGTGCAG  
 GGAGCTGGAGGAGAACATCAAGGAGTTCTGCAGTCGTTCTGCACATCGTCCAGATGT  
**TCATCAACACGTG**TGAgggccccggcgccgaattcgcgatatcggttaacggatccaGA  
 TCTGCTGTGCCTCTAGTTGCCAGCCATCTGTTGCCCCCTCCCCCGTGCCTCCTTGAC  
 CCTGGAAGGTGCCACTCCACTGTCCTTCTTAATAAAATGAGGAATTGCATCGCATTGTC  
 TGAGTAGGTGTCATTCTATTCTGGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGG  
 GAAGACAATAGCAGGCATGCTGGGATGCGGTGGCTCTATGGGTACCCAGGTGCTGAAGAA  
 TTGACCCGGTCCCTCTGGGCCAGAAAGAACAGGACATCCCCCTCTGTGACACACCCCT  
 GTCCACGCCCTGGTTCTAGTCCAGCCCCACTCATAGGACACTCATAGCTCAGGAGGGCT  
 CCGCCTCAATCCCACCCGCTAAAGTACTTGGAGCGGTCTCTCCCTCCCTCATCAGCCCACC  
 AAACCAAACCTAGCCTCCAAGAGTGGGAAGAAATTAAAGCAAGATAGGCTATTAAGTCAGA  
 GGGAGAGAAAATGCCTCCAACATGTGAGGAAGTAATGAGAGAAATCATA

Fig. 4A

M R I S K P H L R S I S I Q C Y L C L L L N S H F L T E A G I H V F I L G C  
**F S A G L P K T E A N W V N V I S D L K K I E D L I Q S M H I D A T L Y T E**  
 S D V H P S C K V T A M K C F L L E L Q V I S L E S G D A S I H D T V E N  
 L I I L A N N S L S S N G N V T E S G C K E C E E L E E K N I K E F L Q S F  
 V H I V Q M F I N T S .

Fig. 4B




Fig. 4C

1 CCTGGCCATTGCATACTTGTATCCATATCATAATATGTACATTATGGCTATGTCAAACATTACGCCAT  
 76 GTTGACATTGATTATTGACTAGTTATTAAATAGTAATCAATTACGGGGTCTAGTTCATAGCCATATATGGAGT  
 151 TCCGCCTTACATACTTACGGTAAATGGCCGCTGGCTGACGCCAACGACCCCCGCCATTGACGTCAATAA  
 226 TGACGTATGTTCCCATAGTAACGCCAATAGGGACTTCCATTGACGTCAATGGTGGAGTATTACGGTAACTG  
 301 CCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCTATTGACGTCAATGATGGTAAATGGCCG  
 376 CCTGGCATTATGCCAGTACATGACCTTATGGACTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTA  
 451 TTACCATGGTGTGATGGGTTTGGCAGTACATCAATGGCGTGGATAGCGGTTGACTCACGGGATTCCAAGTC  
 526 TCCACCCCATTGACGTCAATGGAGTTGTTGGCACCAAAATCAACGGGACTTCCAAAATGTCGTAACAACT  
 601 CCGCCCCATTGACGCAAATGGCGTAGGCGTGTACGGTGGAGGTCTATATAAGCAGAGCTGTTAGTGAACC  
 676 GTCAGATCGCTGGAGACGCCATCCACGCTGTTTGACCTCCATAGAAGACACCGGACCGATCCAGCCTCCGCG  
 751 GGcgcgctcgacaagaa ATG CGG ATC TCG AAG CCG CAC CTG CGG TCG ATA TCG ATC CAG  
 15▶ M R I S K P H L R S I S I Q  
 811 TGC TAC CTG TGC CTG CTC CTG AAC TCG CAC TTC CTC ACG GAG GCC GGT ATA CAC GTC  
 15▶ C Y L C L L N S H F L T E A G I H V  
 868 TTC ATC CTG GGC TGC TTC TCG GCG GGG CTG CCG AAG ACG GAG GCG AAC TGG GTG AAC  
 34▶ F I L G C F S A G L P K T E A N W V N  
 925 GTG ATC TCG GAC CTG AAG AAG ATC GAG GAC CTC ATC CAG TCG ATG CAC ATC GAC GCG  
 53▶ V I S D L K K I E D L I Q S M H I D A  
 982 ACG CTG TAC ACG GAG TCG GAC GTC CAC CCG TCG TGC AAG GTC ACG GCG ATG AAG TGC  
 72▶ T L Y T E S D V H P S C K V T A M K C  
 1039 TTC CTC CTG GAG CTC CAA GTC ATC TCG CTC GAG TCG GGG GAC GCG TCG ATC CAC GAC  
 91▶ F L L E L Q V I S L E S G D A S I H D  
 1096 ACG GTG GAG AAC CTG ATC ATC CTG GCG AAC AAC TCG CTG TCG TCG AAC AAC GGG AAC GTC  
 110▶ T V E N L I I L A N N S L S S N G N V  
 1153 ACG GAG TCG GGC TGC AAG GAG TGC GAG GAG CTG GAG GAG AAG AAC ATC AAG GAG TTC  
 129▶ T E S G C K E C E E L E E K N I K E F  
 1210 CTG CAG TCG TTC GTG CAC ATC GTC CAG ATG TTC ATC AAC ACG TCG TGA gggccggcgc  
 148▶ L Q S F V H I V Q M F I N T S •  
 1269 gccgaattcgccgatatcggttaacggatccaGATCTGCTGTGCCCTCTAGTTGCCAGCCATCTGTTGTTGCC  
 1344 CTCGGGGTGCCTCTTGACCTGGAAAGGTGCCACTCCACTGTCCTTCTTAATAAAATGAGGAAATTGCACTC  
 1419 GCATTGCTGAGTAGGTGTCATTCTATTCTGGGGGTGGGGTGGGGCAGGACAGCAAGGGGAGGATTGGGAAGA  
 1494 CAATAGCAGGCATGCTGGGATGCCGGTGGCTATGGTACCCAGGTGCTGAAGAATTGACCCGGTCTCTG  
 1569 GGCCAGAAAAGCAGGCACATCCCTCTCTGTGACACACCCCTGCCAGGCCCTGGTTCTAGTTCCAGCCCC  
 1644 ACTCATAGGACACTCATAGCTCAGGAGGGCTCCGCCCTCAATCCCACCCGCTAAAGTACTTGGAGCGGTCTCTC  
 1719 CTCCCTCATGCCAACCAAACCTAGCCTCAAGAGTGGGAAGAAATTAAAGCAAGATAGGCTATTAAGT  
 1794 GCAAGGGAGAGAAAATGCCCTCAACATGAGGAAGTAATGAGGAGAAATCATA  
 ▶

Fig. 4D

## AG59 CMV huIL15tPA6

CCTGGCCATTGCATACTGTTATCCATATCATAATATGTACATTATGGCTCATGTCCA  
 ACATTACCGCCATGTTGACATTGATTATTGACTAGTTATTAAATAGTAATCAATTACGGGTC  
 ATTGTTCATAGCCCATAATGGAGTCCCGCTTACATAACTACGGTAAATGGCCCGCTG  
 GCTGACCGCCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAAGTAACG  
 CCAATAGGGACTTCCATTGACGTCAATGGGTGGAGTATTACGGTAAACTGCCACTGGC  
 AGTACATCAAGTGTATCATATGCCAAGTACGCCCTATTGACGTCAATGATGGTAAATGGC  
 CCGCCTGGCATTATGCCAGTACATGACCTATGGACTTCCACTTGGCAGTACATCTAC  
 GTATTAGTCATCGCTATTACCATGGTATGCGGTTTGGCAGTACATCAATGGCGTGGATA  
 GCGGTTGACTCACGGGATTCCAAGTCTCCACCCATTGACGTCAATGGGAGTTGTTT  
 GGCACCAAAATCAACGGGACTTCCAAAATGCGTAACAACCTCCGCCCCATTGACGCAAATG  
 GGCAGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTAGTGAACCGTCAGAT  
 CGCCTGGAGACGCCATCCACGCTTTGACCTCATAGAACGACACGGGACCGATCCAGCC  
 TCCCGGGcgcgcgacaaagaa**ATGGATGCAATGAAGAGAGGGCTCTGCTGTGCTGCT**  
**GCTGTGTGGAGCAGTCTCGTTGCCAGCCAGGAAATCCATGCCGATTCAAGAGAGGAG**  
**CCAGAAACTGGGTGAACGTGATCTGGACCTGAAGAAGATCGAGGACCTCATCCAGTCAGT**  
**CACATCGACGCGACGCTGTACACGGAGTCGGACGTCCACCCCTCGTCAAGGTACGGCGAT**  
**GAAGTGCTTCCCTGGAGCTCCAAGTCATCTCGCTCGAGTCGGGGACCGCTGATCCACG**  
**ACACGGTGGAGAACCTGATCATCCTGGCGAACAACTCGCTGTCGTAACGGGAACGTCACG**  
**GAGTCGGGCTGCAAGGAGTGCAGGGAGCTGGAGGGAGAACATCAAGGAGTTCTGCAGTC**  
**GTTCGTGCACATGTCAGATGTTCATCAACACGTCGT**GA**ggggccggcgccgaattcgc**  
**ggatatcggttaacggatccaGATCTGCTGCTCTAGTTGCCAGCCATCTGTTGTTGC**  
**CCCTCCCCCGTGCCTCCTTGACCCCTGGAAAGGTGCCACTCCCACTGCTCTTCTAATAAAA**  
**TGAGGAAATTGCATCGCATTGTCAGTAGGTGTCATTCTATTCTGGGGGTGGGTGGGCT**  
**AGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCATGCTGGGATGCGGTGGCT**  
**ATGGGTACCCAGGTGCTGAAGAATTGACCCGGTTCCCTGGGCCAGAAAGAACAGGCACA**  
**TCCCCTCTGTGACACACCCCTGTCACGCCCTGGTTCTAGTTCCAGCCCCACTCATAG**  
**GACACTCATAGCTCAGGAGGGCTCCGCTTCAATCCCACCCGCTAAAGTACTTGGAGCGGTC**  
**TCTCCCTCCCTCATCAGCCCACAAACCTAGCCTCCAAGAGTGGGAAGAAATTAAAG**  
**CAAGATAGGCTATTAAGTGCAGAGGGAGAGAAAATGCCTCCAACATGTGAGGAAGTAATGAG**  
**AGAAATCATA**

Fig. 5A

M D A M K R G L C C V L L L C G A V F V S P S Q E I H A R F R R G A R N  
 W V N V I S D L K K I E D L I Q S M H I D A T L Y T E S D V H P S C K V T A  
 M K C F L L E L Q V I S L E S G D A S I H D T V E N L I I L A N N S L S S N  
 G N V T E S G C K E C E E L E E K N I K E F L Q S F V H I V Q M F I N T S •

Fig. 5B


AG59 CMV huIL15tPA6  
(1808 bp)

Fig. 5C

1 CCTGGCCATTGCATACTGTTATCCATATCATAATATGTACATTATGGCTCATGTCACATTACCGCCATGTTGACATTGATTGACTAGTTA  
 101 TTAATAGTAATCAATTACGGGTCTAGTCATAGCCATATGGAGTCCCGCTTACATACTTACGGTAATGGCCGCTGGCTGACCGCCCAAC  
 201 GACCCCCCCCATTGACGTCAATAATGACGTATGTCATAGTAACGCCAATAGGGACTTCCATTGACGTCAATGGTGGAGTATTACGGTAACTG  
 301 CCCACTTGGCAAGTACATCAAGTGTATCATGCCAAGTACGCCCTATTGACGTCAATGTTGAAATGGCCGCTGGCATTATGCCAGTACATGAC  
 401 CTTATGGGACTTCTACTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTATGCGCTGGAGTACATCAATGGGCTGGATAGCGG  
 501 TTTGACTCACGGGATTCCAAGTCTCCACCCATTGACGTCAATGGAGTTGTTGGCACCAAAATCAACGGACTTCCAAATGTCGAACAAC  
 601 CGCGCCATTGACGCAAATGGCGGTAGGGTGTACGGTGGGAGGTATATAAGCAGAGCTCGTTAGTGAACCGTCAGATGCCCTGGAGACGCCATCC  
 701 ACGCTGTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCGCGGgcgcgcgtcgacaagaa ATG GAT GCA ATG AAG AGA GGG CTC  
 793 TGC TGT GTG CTG CTG CTG TGT GGA GCA GTC TTC GTT TCG CCC AGC CAG GAA ATC CAT GCC CGA TTC AGA AGA GGA  
 868 99 C C V L L L C G A V F V S P S Q E I H A R F R R G  
 868 GCC AGA AAC TGG GTG AAC GTG ATC TCG GAC CTG AAG AAG ATC GAG GAC CTC ATC CAG TCG ATG CAC ATC GAC GCG  
 943 34 A R  
 943 ACG CTG TAC ACG GAG TCG GAC GTC CAC CCG TCG TGC AAG GTC ACG GCG ATG AAG TGC TTC CTC CTG GAG CTC CAA  
 1018 24 T L Y T E S D V H P S C K V T A M K C F L L E L Q  
 1018 GTC ATC TCG CTC GAG TCG GGG GAC GCG TCG ATC CAC GAC ACG GTG GAG AAC CTG ATC ATC CTG GCG AAC AAC TCG  
 1093 49 V I S L E S G D A S I H D T V E N L I I L A N N S  
 1093 CTG TCG TCG AAC GGG AAC GTC ACG GAG TCG GGC TGC AAG GAG TGC GAG GAG CTG GAG GAG AAC AAC ATC AAG GAG  
 1168 74 L S S N G N V T E S G C K E C E E L E E K N I K E  
 1168 TTC CTG CAG TCG TTC GTG CAC ATC GTC CAG ATG TTC ATC AAC ACG TCG TGA gggccggccggccgaaattcgccatcggt  
 1251 99 F L Q S F V H I V Q M F I N T S \*  
 1251 taacggatccaGATCTGCTGTGCTCTAGTTGCGCCATCTGTTGTTGCCCCCTCCCCGTGCTTCTGACCCCTGGAGGTGCGACTCCACTGTC  
 1351 1351 CTTTCTAATAAAATGAGGAATTGCGATGCGCTGGCTCTATGGTACCCAGGTGCTGAAGAATTGACCCGGTCTCTGGCCAGAAAGCAGGCACA  
 1451 1451 AAGACAATAGCAGGCATGCTGGGATGCGGTGGCTCTATGGTACCCAGGTGCTGAAGAATTGACCCGGTCTCTGGCCAGAAAGCAGGCACA  
 1551 1551 TCCCCCTCTGTGACACACCTGTCCACGCCCTGGTCTTAAGTCCAGCCCCACTCATAGGACACTCATAGCTCAGGAGGGCTCCGCTTCAATCCA  
 1651 1651 CCCGCTAAAGTACTTGGAGCGGTCTCCCTCCCTCATCAGCCACCAACCAACCTAGCTCCAGAGTGGGAAGAAATTAAAGCAAGATAGGCTATT  
 1751 1751 AAGTGCAGAGGGAGAGAAAATGCTCCACATGAGGAAGTAATGAGAGAAATCATA

Fig. 5D

AG79 huIL15Ra

CCTGGCCATTGCATACTGTATCCATATCATAATATGTACATTATTTGGCTCATGTCCA  
 ACATTACGCCATGTTGACATTGATTATTGACTAGTTATTAAATAGTAATCAATTACGGGTC  
 ATTAGTTCATAGCCATATGGAGTTCCCGTTACATAACTACGGTAATGGCCCGCTG  
 GCTGACCGCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACG  
 CCAATAGGGACTTCCATTGACGTCAATGGTGGAGTATTGACGTCAATGATGGTAAATGGC  
 AGTACATCAAGTGTATCATATGCCAAGTACGCCCTATTGACGTCAATGATGGTAAATGGC  
 CCGCCTGGCATTATGCCAGTACATGACCTATTGGACTTCCACTTGGCAGTACATCTAC  
 GTATTAGTCATCGCTATTACCATGGTATGCGGTTTGGCAGTACATCAATGGCGTGGATA  
 GCGGTTGACTCACGGGATTCCAAGTCTCCACCCATTGACGTCAATGGAGTTGTTT  
 GGCACCAAAATCAACGGACTTCAAATGTCGTACAACACTCCGCCATTGACGCAAATG  
 GCGGGTAGGCGTGTACGGTGGAGGTCTATATAAGCAGAGCTCGTTAGTGAACCGTCAGAT  
 CGCCTGGAGACGCCATCCACGCTGTTGACCTCCATAGAACGACACCGGACCGATCCAGCC  
 TCCGCGGcgcgctcgacgctagcaagaa**ATGGCCCCGAGGCGGGCGCAGGCTGCCGGAC**  
**CCTCGGTCTCCCGCGCTGCTACTGCTCCTGCTGCTCCGGCCGGCGACGCCGGCATCA**  
**CGTCCCCCCCCCATGTCGTGGAGCACGACATCTGGTCAAGAGCTACAGCTTGTAC**  
**TCCCGGGAGCGGTACATCTGCAACTCGGTTCAAGCGGAAGGCCGGCACGTCCAGCCTGAC**  
**GGAGTGCCTGTTGAACAAGGCCACGAATGTCGCCACTGGACGACCCCTCGCTCAAGTGA**  
**TCCCGACCCGGCCCTGGTACCCAGCGGGCGGCCACCCCTCACCGTAACGACGGCGGGG**  
**GTGACCCCGCAGCGGAGAGCCTCTCCCGTCGGAAAGGAGGCCGCCGTCGTCGCCAG**  
**CTCGAACAAACACGGCGCCACAACGTCAGCGATCGTCCCAGCTCCAGCTGATGCCGTCGA**  
**AGTCGCCGTCCACGGAAACCACGGAGATCAGCAGTCATGAGTCCTCCACGGCACCCCTCG**  
**CAAACGACGCCAAGAACTGGAAACTCACGGCGTCCGCTCCACCAGCCGGGGGTGTA**  
**TCCGCAAGGCCACAGCGACACCACGGTGGCGATCTCCACGTCCACGGTCTGCTGTGGC**  
**TGAGCGCGGTGTCGCTCTGGCGTGTACCTCAAGTCGAGGCAACTCCCCGCTGCCAGC**  
**GTTGAGATGGAGGCCATGGAGGCTCTGCCGTGACGTGGGGACCGACGAGGGATGAGGA**  
**CTTGGAGAACTGCTCGCACCCACCTATAATGA**gaattcgatccaGATCTGCTGTGCCTTCTAG  
 TTGCCAGCCATCTGTTGCCCCCTCCCCGTGCCTTCCTGACCTGGAAGGTGCCACTC  
 CCACGTCTCTTCTAATAAAATGAGGAATTGCACTCGCATTGTCAGTAGGTGTCATTCT  
 ATTCTGGGGGGTGGGGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCA  
 TGCTGGGGATGCGGTGGCTCTATGGTACCCAGGTGCTGAAGAATTGACCCGGTCTCCT  
 GGGCCAGAAAGCAGGCACATCCCCTCTGTGACACACCCCTGTCCACGCCCTGGTTC  
 TTAGTTCCAGCCCCACTCATAGGACACTCATAGCTCAGGAGGCTCCGCTTCAATCCCACC  
 CGCTAAAGTACTTGGAGCGGTCTCCCTCCCTCATCAGCCACCAACAAACCTAGCCTC  
 CAAGAGTGGGAAGAAATTAAAGCAAGATAGGCTATTAGTGCAGAGGGAGAGAAAATGCCTC  
 CAACATGTGAGGAAGTAATGAGAGAAATCATA

Fig. 6A

MAPRRARGCRTLGLPALLLRLPPATRGITCP  
 SVEHADIWVKSYSLYSRERYICNSGFKRKAGTSSLTE  
 CVLNKATNVAHWTPSLKCIRDPALVHQRPAPPSTVT  
 TAGVTPQPESLSPSGKEPAASSPSSNNTAATTAAIVP  
 GSQLMPSKSPSTGTTTEISSHESSHGTPSQTTAKNWEL  
 TASASHQPPGVYPQGHSDTTVAISTSTVLLCGLSAVS  
 LLACYLRQTPPLASVEMEAMEALPVTWGTSSRDED  
 LENCSHHL..

Fig. 6B




Fig. 6C

1 CCTGGCCATTGCATACTTGTATCCATATCATAATATGTACATTATGGCTATGTCACATTACGGCCATGT  
 78 TGACATTGATTGACTAGTTATTAACTAGTAATCAATTACGGGTCTTACGGCCATTGACGTCATAATGACGT  
 155 CGTTACATAACTTACGGTAATGGCCGCCCTGGCTGACCGCCAAACGACCCCCGCCATTGACGTCATAATGACGT  
 232 ATGTTCCCATAGTAACGCCAATAGGGACTTCCATTGACGTCATGGTGGAGTATTACGGTAAACTGCCACTTG  
 309 GCAGTACATCAAGTGTATCATATGCCAAGTACGCCCTTATTGACGTCATGATGGTAAATGGCCGCCCTGGCTTAA  
 386 TGCCCAGTACATGCCATTGGACTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGA  
 463 TCGGGTTTGGCAGTACATCAATGGCGTGGATACGGTTGACTCACGGGATTCAAGTCTCCACCCATTGAC  
 540 GTCAATGGGAGTTGTTGGCACAAAATCAACGGGACTTCCAAAATGCGTAACAACCTCCGCCATTGACGCA  
 617 AATGGGCGGTAGGCGTGTACGGTGGAGGTCTATAAACGAGCTCGTTAGTGAACCGTCAGATGCCCTGGAGAC  
 694 GCCATCCACGCTGTTGACCTCCATAGAACACCGGACCGATCCAGCCTCCGGggcgcgctcgacgctagca →  
 771 agaa ATG GCC CCG AGG CGG GCG CGA GGC TGC CGG ACC CTC GGT CTC CCG GCG CTG CTA  
 1 M A P R R A R G C R T L G L P A L L  
 829 CTG CTC CTG CTG CTC CGG CGG CGG GCG ACG CGG GGC ATC ACG TGC CCG CCC CCC ATG  
 19 L L L L R P P A T R G I T C P P P M  
 886 TCC GTG GAG CAC GCA GAC ATC TGG GTC AAG AGC TAC AGC TTG TAC TCC CGG GAG CGG  
 38 S V E H A D I W V K S Y S L Y S R E R  
 943 TAC ATC TGC AAC TCG GGT TTC AAG CGG AAG GGC ACG TCC AGC CTG AGC CGG TGC  
 57 Y I C N S G F K R K A G T S S L T E C  
 1000 GTG TTG AAC AAG GCC ACG AAT GTC GCC CAC TGG ACG ACC CCC TCG CTC AAG TGC ATC  
 76 V L N K A T N V A H W T T P S L K C I  
 1057 CGC GAC CCG GCC CTG GTT CAC CAG CGG CCC GCG CCA CCC TCC ACC GTA ACG ACG GCG  
 95 R D P A L V H Q R P A P P S T V T T A  
 1114 GGG GTG ACC CCG CAG CGG GAG AGC CTC TCC CCG TCG GGA AAG GAG CCC GCG GCG TCG  
 114 P G V T P Q P E S L S P S G K E P A A S  
 1171 TCG CCC AGC TCG AAC AAC ACG GCG ACA ACT GCA GCG ATC GTC CCG GGC TCC CAG  
 133 S P S S N N T A A T T A A I V P G S Q  
 1228 CTG ATG CCG TCG AAG TCG CGG TCC ACG GGA ACC ACG GAG ATC AGC AGT CAT GAG TCC  
 152 L M P S K S P S T G T T E I S S H E S  
 1285 TCC CAC GGC ACC CCC TCG CAA ACG ACG GGC AAG AAC TGG GAA CTC ACG GCG TCC GCC  
 171 S H G T P S Q T T A K N W E L T A S A  
 1342 TCC CAC CAG CCG CGG GGG GTG TAT CGG CAA GGC CAC AGC GAC ACC ACG GTG GCG ATC  
 190 P S H Q P P G V Y P Q G H S D T T V A I  
 1399 TCC TCC ACG TCG GTC CTG TGT GGG CTG AGC GCG GTG TCG CTC CTG GCG TGC TAC  
 209 S T S T V L L C G L S A V S L L A C Y  
 1456 CTC AAG TCG AGG CAG ACT CCC CGG CTG GCC AGC GTT GAG ATG GAG GGC ATG GAG GCT  
 228 L K S R Q T P P L A S V E M E A M E A  
 1513 CTG CGG GTG ACG TGG GGG ACC AGC AGC AGG GAT GAG GAC TTG GAG AAC TGC TCG CAC  
 247 L P V T W G T S S R D E D L E N C S H  
 1570 CAC CTA TAA TGA *gaattc* gat CTG CTG TGC CTT CTA GTT GCC AGC CAT CTG TTG  
 266 H L \* \*  
 1630 TTT GCC CCT CCC CGG TGC CTT CCT TGA CCC TGG AAG GTG CCA CTC CCA CTG TCC TTT  
 1687 CCT AAT AAA ATG AGG AAA TTG C ATCGCATTGTCGTGAGTAGGTGTCATTCTATTCTGGGGGGTGGGGTGG  
 1756 GCGAGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCATGCTGGGGATGCGGTGGGTCTATGGGTACCCAG  
 1833 GTGCTGAAGAATTGACCCGGTCTCTGGCCAGAAAGAAGCAGGCACATCCCCCTCTGTGACACACCCCTGTC  
 1910 ACGCCCTGGTTCTTAGTCCAGGCCCACTCATAGGACACTCATAGCTCAGGAGGGCTCCCTCAATCCACCCCG  
 1987 CTAAGTACTTGGAGCGGTCTCTCCCTCCCTCATAGCCACCAACCAACCTAGCCTCCAAGACTGGAGAAAT  
 2064 TAAAGCAAGATAGGCTATTAAGTGCAGAGGGAGAGAAAATGCCTCAACATGTGAGGAAGTAATGAGAGAAATCATA →

Fig. 6D

AG98 CMV hu sIL15Ra

CCTGGCCATTGCATACTTGTATCCATATCATAATATGTACATTATGGCTCATGTCCA  
 ACATTACGCCATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTC  
 ATTAGTCATAGCCCATAATGGAGTCCCGTTACATAACTACGGTAAATGGCCCGCTG  
 GCTGACCGCCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATACTAACG  
 CCAATAGGGACTTCCATTGACGTCAATGGGTGGAGTATTACGGTAAACTGCCACTTGGC  
 AGTACATCAAGTGTATCATATGCCAAGTACGCCCTATTGACGTCAATGATGGTAAATGGC  
 CCGCCTGGCATTATGCCAGTACATGACCTTATGGACTTCTACTTGGCAGTACATCTAC  
 GTATTAGTCATCGCTATTACCATGGTATGCGGTTTGGCAGTACATCAATGGCGTGGATA  
 GCGGTTGACTCACGGGATTCCAAGTCTCCACCCATTGACGTCAATGGAGTTGTTT  
 GGCACCAAAATCAACGGACTTCCAAAATGTCGTAACAACCTCCGCCATTGACGCAAATG  
 GGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTAGTGAACCGTCAGAT  
 CGCCTGGAGACGCCATCCACGCTGTTGACCTCCATAGAAGACACCGGACGATCCAGCC  
 TCCCGGGgcgcgcgtcgacgttagcaagaa**ATGGCCCCGAGGCCGGCGAGGCTGCCGGAC**  
 CCTCGGTCTCCGGCGTGTACTGCTCTGCTGCTCCGGCCCGCGACGCCGGCATCA  
 CGTCCCCGCCCATGTCGTTGGAGCACGCAGACATCTGGGTCAAGAGCTACAGCTTGTAC  
 TCCCGGGAGCGGTACATCTGCAACTCGGGTTCAAGCGGAAGGCCGGCACGTCCAGCCTGAC  
 GGAGTGGTGTGAACAAGGCCACGAATGTCGCCACTGGACGACCCCTCGCTCAAGTGCA  
 TCCCGGACCCGGCCCTGGTCACCAGCGGGCCCGCCACCCCTCACCGTAACGACGGCGGGG  
 GTGACCCCGCAGCCGGAGAGCCTCTCCCGTCGGAAAGGAGCCCGCCGTCGTCGCCAG  
 CTCGAACAACACGGCGGCCACAACGTCAGCGATCGTCCCAGCTGATGCCGTCGA  
 AGTCGCCGTCCACGGGAACCACGGAGATCAGCAGTCATGAGTCTCCACGGCACCCCTCG  
 CAAACGACGGCAAGAACTGGGAACTCACGGCGTCCGCCTCCACAGCCGCCGGGTGTA  
 TCCGCAAGGCCACAGCGACACCACGTAAT**GA**gaattcgcgatatcggttaacggatccaGA  
 TCTGCTGTGCCCTCTAGTTGCCAGCCATCTGTTGTTGCCCTCCCCGTGCCTCCTTGAC  
 CCTGGAAGGTGCCACTCCACTGTCCTTCTAATAAAATGAGGAAATTGCATCGCATTGTC  
 TGAGTAGGTGTCATTCTATTCTGGGGGTGGGGTGGGGCAGGACAGCAAGGGGAGGATTGG  
 GAAGACAATAGCAGGCATGCTGGGATGCCGTGGCTATGGGTACCCAGGTGCTGAAGAA  
 TTGACCCGGTTCCCTCTGGGCCAGAAAGAACAGCAGGCACATCCCTCTGTGACACACCCT  
 GTCCACGCCCTGGTTCTAGTCCAGCCCCACTCATAGGACACTCATAGCTCAGGAGGGCT  
 CCGCCTCAATCCCACCGCTAAAGTACTGGAGCGGTCTCCCTCCCTCATCAGCCCACC  
 AAACCAAACCTAGCCTCCAAGAGTGGGAAGAAATTAAAGCAAGATAGGCTATTAAGTGCAGA  
 GGGAGAGAAAATGCCTCCAACATGTGAGGAAGTAATGAGAGAAATCATA

Fig. 7A

M A P R R A R G C R T L G L P A L L L L L R P P A T R G I T C P P P M  
 S V E H A D I W V K S Y S L Y S R E R Y I C N S G F K R K A G T S S L T E  
 C V L N K A T N V A H W T T P S L K C I R D P A L V H Q R P A P P S T V T  
 T A G V T P Q P E S L S P S G K E P A A S S P S S N N T A A T T A A I V P  
 G S Q L M P S K S P S T G T T E I S S H E S S H G T P S Q T T A K N W E L  
 T A S A S H Q P P G V Y P Q G H S D T T ..

Fig. 7B

Construct: AG98 CMVhu-sIL15Ra  
(1971 bp)




Fig. 7C

1 CCTGGCCATTGCATACTGTTATCCATATCATAATATGTACATTATATGGCTATGCCAACATTACGCCATGTTGACATTGATTGACTAGTTA  
101 TTAATAGTAATCAATTACGGGGTCAATTAGTTCATAGCCATATATGGAGTCCGGTTACATAACTACGGTAATGGCCGCTGGCTGACCGCCAAAC  
201 GACCCCCGCCATTGACGTCATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTCCATTGACGTCATGGTGGAGTATTACGGTAAACTG  
301 CCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCTATTGACGTCATGATGGTAAATGGCCGCTGGCATTATGCCAGTACATGAC  
401 CTTATGGGACTTCTACTTGGCAGTACATCTACGTATTAGTCATCGTATTACCATGGTATGCGGTTTGGCAGTACATCAATGGCGTGGATAGCGG  
501 TTTGACTCACGGGGATTCCAACTCTCCACCCATTGACGTCATGGAGTTGGTGGACCAAAATCAACGGGACTTCCAAAATGTCGAACAACCT  
601 CGGCCCCATTGACGCAAATGGCGGTAGGCGTACGGTGGAGGCTATATAAGCAGAGCTGTTAGTGAACCGTCAGATGCCCTGGAGACGCCATCC  
701 ACGCTGTTTGACTCCATAGAAGACACGGGACCGATCCAGCCTCCGGggcgcgctcgacgctcgaa ATG GCC CCG AGG CGG GCG  
793 CGA GGC TGC CGG ACC CTC GGT CTC CGG GCG CTG CTA CTG CTC CTG CTG CTC CGG CGG CGG CGG CGG ACG CGG GGC ATC  
7 R G C R T L G L P A L L L L L R P P A T R G I  
868 ACG TGC CCG CCC CCC ATG TCC GTG GAG CAC GCA GAC ATC TGG GTC AAG AGC TAC AGC TTG TAC TCC CGG GAG CGG  
32 T C P P M S V E H A D I W V K S Y S L Y S R E R  
943 TAC ATC TGC AAC TCG GGT TTC AAG CGG AAG GCC GGC ACG TCC AGC CTG AGC GAG TGC GTG TTG AAC AAG GGC ACG  
57 Y I C N S G F K R K A G T S S L T E C V L N K A T  
1018 AAT GTC GCC CAC TGG ACG ACC CCC TCG CTC AAG TGC ATC CGC GAC CCG CGG GCC CTG GTT CAC CAG CGG CCC GCG CCA  
82 N V A H W T T P S L K C I R D P A L V H Q R P A P  
1093 CCC TCC ACC GTA ACG ACG GCG GGG GTG ACC CGG CAG CGG GAG AGC CTC TCC CGG TCG GGA AAG GAG CCC GCG  
107 P S T V T T A G V T P Q P E S L S P S G K E P A A  
1168 TCG TCG CCC AGC TCG AAC AAC ACG GCG GCC ACA ACT GCA GCG ATC GTC CGG GGC TCC CAG CTG ATG CGG TCG AAG  
132 S S P S S N N T A A T T A A I V P G S Q L M P S K  
1243 TCG CCG TCC ACG GGA ACC ACG GAG ATC ACG AGT CAT GAG TCC TCC CAC GGC ACC CCC TCG CAA ACG ACG GGC AAG  
157 S P S T G T T E I S S H E S S H G T P S Q T T A K  
1318 AAC TGG GAA CTC ACG GCG TCC GCC CAC CAG CGG CCG GGG GTG TAT CCG CAA GGC CAC AGC GAC ACC ACG TAA  
182 N W E L T A S A S H Q P P G V Y P Q G H S D T T •  
1393 TGA gaattcggatatcggttaacggatccatcgctgtgccttctatgtggccatctgtgtgtgcctcccccgtgccttcgtaccct  
207 \*  
1492 GGAAGGTGCCACTCCACTGTGCTTCTAATAAAATGAGGAATTGCATCGATTGTCAGTGGTGTATTCTGTTGGGGTGGGTGGGAG  
1592 GACAGCAAGGGGAGGATTGGGAGACAATAGCAGGCACTGCTGGGATCCGGTGGCTATGGTACCCAGGTCTGAAGAATTGACCCGGTTCCCTCCT  
1692 GGGCCAGAAAGAAGCAGGCACATCCCTCTCTGTGACACACCCCTGCCACGCCCTGGTTCTAGTCCAGCCCCACTCATGGACACTCATAGCTCAG  
1792 GAGGGCTCCGCTTCATCCACCCGCTAAAGTACTTGGAGGGTCTCCCTCCCTCATGCCACCAACCAACCTAGCTCCAGAGTGGAGAAG  
1892 AATTAAGCAAGATAGGCTATTAGTCAGAGGGAGAGAAAATGCCCTCAACATGTGAGGAAGTAATGAGAGAAAATCATA

Fig. 7D

AG151 huIL-15 huGM-CSF

CCTGGCCATTGCATACTTGTATCCATATCATAATATGTACATTATGGCTCATGTCCA  
ACATTACCGCCATGTTGACATTGATTATTGACTAGTTATTAAATAGTAATCAATTACGGGTC  
ATTAGTTCATAGCCCATATGGAGTTCCCGGTACATAACTACGGTAAATGGCCCGCTG  
GCTGACCGCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCATAGAACG  
CCAATAGGGACTTCCATTGACGTCAATGGGTGGAGTATTACGGTAAACTGCCACTTGGC  
AGTACATCAAGTGTATCATATGCCAAGTACGCCCTATTGACGTCAATGATGGTAAATGGC  
CCGCCTGGCATTATGCCAGTACATGACCTTATGGGACTTCCATTGGCAGTACATCTAC  
GTATTAGTCATCGCTATTACCATGGTATGCCGTTTGGCAGTACATCAATGGCGTGGATA  
GCGGTTTACTCACGGGATTCCAAGTCTCCACCCATTGACGTCAATGGGAGTTGTTT  
GGCACAAAATCAACGGGACTTCCAAAATGTCGTAACAACCTCCGCCATTGACGCAAATG  
GGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTAGTGAACCGTCAGAT  
CGCCTGGAGACGCCATCCACGCTGTTGACCTCCATAGAAGACACCGGACCGATCAGCC  
TCCGGGGggcgcgctcgacaagaatGTGGCTCCAGAGCCTGCTACTCCTGGGACGGTGGC  
CTGCAGCATCTGA~~ACTGGGTGAA~~CGTAC~~T~~CGGAC~~T~~GAAGAAGATCGAGGAC~~T~~CATCC  
AGTCGATGCACATCGACCGCAGCCTGTACACGGAGTCGGACGTCACCCGTGCAAGGTC  
ACGGCGATGAAGTGCTTCCCTGGAGCTCCAAGTCATCTCGCTCGAGTCGGGGACCGTC  
GATCCACGACACGGTGGAGAACCTGATCATCCTGGCAACAACTCGCTGTCGTAACGGGA  
ACGTACGGAGTCGGCTGCAAGGAGTGCAGGGAGCTGGAGGAGAAGAACATCAAGGAGTTC  
CTGCAGTCGTTCGTGCACATCGTCCAGATGTTCATCAACACGT~~CGT~~GA~~ggcccgccgcgc~~  
gaattcgcgatatcggttaacggatccaGATCTGCTGTGCCTCTAGTTGCCAGCCATCTG  
TTGTTGCCCTCCCCGTGCCTTCCTGACCTGGAAAGGTGCCACTCCACTGTCCCTTCC  
TAATAAAATGAGGAAATGCATCGCATTGCTGAGTAGGTGTCATTCTATTCTGGGGGTGG  
GGTGGGGCAGGACAGCAAGGGGGAGGATTGGAAGACAATAGCAGGCATGCTGGGATGCGG  
TGGGCTCTATGGGTACCCAGGTGCTGAAGAATTGACCCGGTTCCCTGGCCAGAAAGAAG  
CAGGCACATCCCTCTGTGACACACCCGTCCACGCCCTGGTCTTAGTTCCAGCCCC  
ACTCATAGGACACTCATAGCTCAGGAGGGCTCCGCCTCAATCCCACCCGCTAAAGTACTTG  
GAGCGGTCTCCCTCCCTCATCAGCCCACAAACCAACCTAGCCTCCAAGAGTGGGAAGA  
AATTAAAGCAAGATAGGTATTAAGTCAGAGGGAGAGAAAATGCCTCCAACATGTGAGGAA  
GTAATGAGAGAAAATCATA

Fig. 8A

M W L Q S L L L G T V A C S I S N W V N V I S D L K K I E D L I Q S M H I  
D A T L Y T E S D V H P S C K V T A M K C F L L E L Q V I S L E S G D A S I  
H D T V E N L I I L A N N S L S S N G N V T E S G C K E C E E L E E K N I K  
E F L Q S F V H I V Q M F I N T S .

Fig. 8B




Fig. 8C

1 CCTGGCCATTGCATACTTGTATCCATATCATATACTACATTATGGCTCATGTCACACATTACGCCATGTTGACATTGATTATTGACTAGTTA  
 101 TTAATAGTAATCAATTACGGGGTCAATTAGTTCATAGCCCATATATGGACTTCCGCGTACATAACTTACGGTAATGGCCGCTGGCTGACCCCAAC  
 201 GACCCCCGCCATTGACGTCAATAATGACGTATGTTCCATAGTAACGCCAATAGGGACTTTCATTGACGTCAATGGGGAGTATTACGGTAACTG  
 301 CCCACTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCTATTGACGTCAATGGTAAATGGCCGCTGGCATTGCCCAGTACATGAC  
 401 CTTATGGGACTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGTATTACCATGGTGTGGGGTTGGCAGTACATCAATGGGCTGGATAGGG  
 501 TTTGACTCAGGGGATTCCAAGTCTCACCCATTGACGTCAATGGGAGTTGGTGGCAGCAAAATCAACGGGACTTCCAAAATGTGTAACAACT  
 601 CGGCCCCATTGACGCAAATGOGCGTAGGCCTGACGTTGGAGGTCTATATAAGCAGAGCTGTTAGTGAACCCTGAGATGCCCTGGAGACGCCATCC  
 701 ACGCTGTTTGACCTCCATAGAACAGACCGGGACCGATCCAGCCTCCGGGGcgcgcgtcgacaaagaa ATG TGG CTC CAG AGC CTG CTA  
 790 CTC CTG GGG ACG GTG GCC TGC AGC ATC TCG AAC TGG GTG AAC GTG ATC TCG GAC CTG AAG AAG ATC GAG GAC CTC  
 8 L L G T V A C S I S N W V N V I S D L K K I E D L  
 865 ATC CAG TCG ATG CAC ATC GAC GCG ACG CTG TAC ACG GAG TCG GAC GTC CAC CCG TCG TGC AAG GTC ACG GCG ATG  
 33 I Q S M H I D A T L Y T E S D V H P S C K V T A M  
 940 AAG TGC TTC CTC CTG GAG CTC CAA GTC ATC TCG CTC GAG TCG GGG GAC GCG TCG ATC CAC GAC ACG GTG GAG AAC  
 58 K C F L L E L Q V I S L E S G D A S I H D T V E N  
 1015 CTG ATC ATC CTG GCG AAC AAC TCG CTG TCG TCG AAC GGG AAC GTC ACG GAG TCG GGC TGC AAG GAG TGC GAG GAG  
 83 L I I L A N N S L S S N G N V T E S G C K E C E E  
 1090 CTG GAG GAG AAC AAC ATC AAG GAG TTC CTG CAG TCG TTC GTG CAC ATC GTC CAG ATG TTC ATC AAC ACG TCG TGA  
 108 L E E K N I K E F L Q S F V H I V Q M F I N T S \*  
 1165 gggcccgccgcgcgcgatattcgccgatatcggttaacggatccaGATCTGCTGTGCTTCTAGTTGCGACCCATCTGTTGCCCCCTCCCCCGTGCTGCTT  
 1265 CCTTGACCCCTGGAAAGGTGCCACTCCACTGTCTTCTCTAAATAAAATGAGGAAATTGATCCTGCGTCTGACTAGGTGTCATTCTATCTGGGGGGTGG  
 1365 GGTGGGGCAGGACAGCAAGGGGGAGGATTGGGAAGACAATAGCAGGCATGCTGGGATCGGTGGGCTATGGTACCCAGGTGTAAGAATTGACCC  
 1465 GGTCTCTCTGGCCAGAAADAAGCAGGCACATCCCCCTCTGTGACACACCCCTGTCCACCCCCCTGGTTCTTAACTTCAGCCCCACTCATAGGACCT  
 1565 CATAAGCTCAAGGAGGGCTCCGCCTCAATCCCACCCGCTAAAGTACTTGGAGCGCTCTCCCTCCCTCATCACCCACCAAAACCTAGCTCCAAAG  
 1665 AGTGGGAAGAAATTAAAGCAAGATAGGTATTAAGTGCAGAGGGAGAGAAAATGCTCCAACTGTGAGGAAGTAATGAGAGAAATCATA

Fig. 8D



## EUROPEAN SEARCH REPORT

Application Number  
EP 13 19 5495

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (IPC)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	WO 2007/046006 A2 (INST NAT SANTE RECH MED [FR]; JACQUES YANNICK [FR]; PLET ARIANE [FR]);) 26 April 2007 (2007-04-26) * page 13, line 1 - line 17 * * page 14, line 16 - line 30 * * page 39, line 11 - page 41, line 34 * * page 54, line 10 - line 14 * * example 1 * ----- X WO 2007/001677 A2 (UNIV CONNECTICUT [US]; LEFRANCOIS LEO [US]; STOKLASEK THOMAS A [US]) 4 January 2007 (2007-01-04) * examples 1-2 * -----	1-16	INV. A61K39/395
The present search report has been drawn up for all claims			
2	Place of search Munich	Date of completion of the search 19 March 2014	Examiner Mauhin, Viviane
<b>CATEGORY OF CITED DOCUMENTS</b> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.

EP 13 19 5495

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

19-03-2014

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 2007046006	A2	26-04-2007	AT	527365 T	15-10-2011	
			BR	PI0619300 A2	27-09-2011	
			CA	2625694 A1	26-04-2007	
			CN	101360827 A	04-02-2009	
			DK	1934353 T3	09-01-2012	
			EP	1777294 A1	25-04-2007	
			EP	1934353 A2	25-06-2008	
			ES	2374166 T3	14-02-2012	
			JP	5394742 B2	22-01-2014	
			JP	2009512433 A	26-03-2009	
			JP	2013226155 A	07-11-2013	
			PT	1934353 E	10-01-2012	
			US	2009238791 A1	24-09-2009	
			WO	2007046006 A2	26-04-2007	
<hr/>						
WO 2007001677	A2	04-01-2007	AU	2006262711 A1	04-01-2007	
			CA	2608474 A1	04-01-2007	
			EP	1899364 A2	19-03-2008	
			JP	5236461 B2	17-07-2013	
			JP	2008545397 A	18-12-2008	
			JP	2013067632 A	18-04-2013	
			NZ	563263 A	30-04-2010	
			US	2006263857 A1	23-11-2006	
			US	2012177598 A1	12-07-2012	
			WO	2007001677 A2	04-01-2007	
<hr/>						

EPO FORM 50459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- US 93747107 P [0001]
- US 20050048549 A [0030] [0096]
- US 81256606 P [0082] [0094]
- US 5965726 A [0082] [0094]
- US 6174666 A [0082] [0094]
- US 6291664 A [0082] [0094]
- US 6414132 A [0082] [0094]
- US 6794498 A [0082] [0094]
- US 60758819 P [0082] [0094]
- WO 2007084342 A [0082] [0094]
- GB 2034323 A [0084]
- WO 9530695 A [0092]
- WO 9412650 A [0104]
- WO 0168882 A [0104]
- WO 07070488 A [0111]
- US 5807715 A [0114] [0116]
- US 6331415 B [0114] [0116]
- US 6818216 B [0114] [0116]
- US 20020098189 A [0114] [0116]
- US 20040028685 A [0114] [0116]
- US 20050019330 A [0114] [0116]
- US 20070086943 A [0114] [0116]
- WO 0246237 A [0114] [0116]
- US 4889803 A [0123]
- US 20040220140 A [0133] [0147]
- US 20010055807 A [0133]
- US 5622834 A [0133] [0147]
- US 5623064 A [0133] [0146] [0147]
- US 5624679 A [0133] [0147]
- US 5686115 A [0133] [0147]
- US 5858350 A [0133] [0147]
- US 6599720 A [0133] [0147]
- US 6686342 A [0133] [0147]
- US 7115588 A [0133] [0147]
- US 4705540 A [0134]
- US 4717393 A [0134]
- US 4717394 A [0134]
- US 4912197 A [0134]
- US 4838900 A [0134]
- US 4935490 A [0134]
- US 4851505 A [0134]
- US 4880442 A [0134]
- US 4863496 A [0134]
- US 4961539 A [0134]
- EP 0219878 A [0134]
- US 6686342 B [0137]
- US 2001055807 A [0147]
- US 4675187 A [0188]
- WO 02098370 A [0273]
- WO 02069904 A [0273]
- WO 9308656 A [0273]
- US 6162432 A [0273]
- WO 05085282 A [0282] [0283]
- WO 9418318 A [0305]

## Non-patent literature cited in the description

- **EMINI et al.** *Nature*, 1983, vol. 304, 699 [0084]
- **GANEM ; VARMUS.** *Ann. Rev. Biochem.*, 1987, vol. 56, 651-693 [0084]
- **TIOLLAIS et al.** *Nature*, 1985, vol. 317, 489-495 [0084]
- **FEHNIGER ; CALIGIURI.** *Blood*, 2001, vol. 97, 14-32 [0092]
- **AUSUBEL et al.** *Current Protocols in Molecular Biology*. John Wiley and Sons, Inc, 1995 [0092]
- **SAMBROOK et al.** *Molecular Cloning, A Laboratory Manual*. Cold Spring Harbor Press, 1989 [0092]
- **BIRREN et al.** *Genome Analysis: A Laboratory Manual*. Cold Spring Harbor Press, 1997, vol. 1-4 [0092]
- **BERNOIST ; CHAMBON.** *Nature*, 1981, vol. 290, 304-310 [0102]
- **YAMAMOTO et al.** *Cell*, 1980, vol. 22, 787-7797 [0102]
- **WAGNER et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1981, vol. 78, 1441-1445 [0102]
- **BRINSTER et al.** *Nature*, 1982, vol. 296, 39-42 [0102]
- **VILLA-KAMAROFF et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1978, vol. 75, 3727-3731 [0102]
- **DEBOER et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 1983, vol. 80, 21-25 [0102]
- Useful proteins from recombinant bacteria. *Scientific American*, 1980, vol. 242, 74-94 [0102]
- **HERRERA-ESTRELLA et al.** *Nature*, vol. 303, 209-213 [0102]
- **GARDNER et al.** *Nucl. Acids Res.*, 1981, vol. 9, 2871 [0102]
- **HERRERA-ESTRELLA et al.** *Nature*, 1984, vol. 310, 115-120 [0102]
- **SWIFT et al.** *Cell*, 1984, vol. 38, 639-646 [0102]

- **ORNITZ et al.** *Cold Spring Harbor Symp. Quant. Biol.*, 1986, vol. 50, 399-409 **[0102]**
- **MACDONALD.** *Hepatology*, 1987, vol. 7, 425-515 **[0102]**
- **HANAHAN.** *Nature*, 1985, vol. 315, 115-122 **[0102]**
- **GROSSCHEDL et al.** *Cell*, 1984, vol. 38, 647-658 **[0102]**
- **ADAMES et al.** *Nature*, 1985, vol. 318, 533-538 **[0102]**
- **ALEXANDER et al.** *Mol. Cell. Biol.*, 1987, vol. 7, 1436-1444 **[0102]**
- **LEDER et al.** *Cell*, 1986, vol. 45, 485-495 **[0102]**
- **PINKERT et al.** *Genes and Devel.*, 1987, vol. 1, 268-276 **[0102]**
- **KRUMLAUF et al.** *Mol. Cell. Biol.*, 1985, vol. 5, 1639-1648 **[0102]**
- **HAMMER et al.** *Science*, 1987, vol. 235, 53-58 **[0102]**
- **KELSEY et al.** *Genes and Devel.*, 1987, vol. 1, 161-171 **[0102]**
- **MOGRAM et al.** *Nature*, 1985, vol. 315, 338-340 **[0102]**
- **KOLLIAS et al.** *Cell*, 1986, vol. 46, 89-94 **[0102]**
- **READHEAD et al.** *Cell*, 1987, vol. 48, 703-712 **[0102]**
- **SANI.** *Nature*, 1985, vol. 314, 283-286 **[0102]**
- **MASON et al.** *Science*, 1986, vol. 234, 1372-1378 **[0102]**
- **SAMBROOK.** Molecular Cloning, A Laboratory Manual. 1989 **[0108]**
- DNA Cloning. 1985, vol. I, II **[0108]**
- Oligonucleotide Synthesis. 1984 **[0108]**
- Nucleic Acid Hybridization **[0108]**
- Transcription and Translation. 1984 **[0108]**
- Animal Cell Culture. 1986 **[0108]**
- Immobilized Cells and Enzymes. IRL Press, 1986 **[0108]**
- **PERBAL.** A Practical Guide to Molecular Cloning. 1984 **[0108]**
- Gene Transfer Vectors for Mammalian Cells. Cold Spring Harbor Laboratory, 1987 **[0108]**
- Methods in Enzymology. 1987, vol. 154, 155 **[0108]**
- Immunochemical Methods in Cell and Molecular Biology. Academic Press, 1987 **[0108]**
- Expression of Proteins in Mammalian Cells Using Vaccinia Viral Vectors in Current Protocols in Molecular Biology. 1991, vol. 2 **[0108]**
- **HARLOW et al.** Antibodies. A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1988 **[0114]** **[0116]**
- **HAMMERLING et al.** Monoclonal Antibodies and T-Cell Hybridomas. Elsevier, 1981, 563-681 **[0114]** **[0116]**
- Physician's Desk Reference. 2007 **[0170]** **[0269]**
- **PAGE et al.** *Intl. J. of Oncology*, 1993, vol. 3, 473-476 **[0175]**
- **SKEHAN et al.** *J. Nat'l Cancer Inst.*, 1990, vol. 82, 1107-12 **[0176]**
- **MOSMANN.** *J. Immunol. Methods*, 1983, vol. 65, 55-63 **[0176]**
- **PIAZZA et al.** *Cancer Research*, 1995, vol. 55, 3110-16 **[0177]**
- Biochemica. Roche Molecular Biochemicals, 1999, 34-37 **[0178]** **[0333]**
- **FISHMAN et al.** Medicine. J.B. Lippincott Co, 1985 **[0189]**
- **MURPHY et al.** Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery. Penguin Books U.S.A., Inc, 1997 **[0189]**
- **LANGER.** *Science*, 1990, vol. 249, 1527-1533 **[0244]** **[0245]**
- **TREAT et al.** Liposomes in the Therapy of Infectious Disease and Bacterial infection. Liss, 1989, 353-365 **[0244]**
- LIPOSOMES IN THE THERAPY OF INFECTIOUS DISEASE AND BACTERIAL INFECTION. 317-327 **[0244]**
- **GOODSON.** *Medical Applications of Controlled Release*, 1984, vol. 2, 115-138 **[0245]**
- **SEFTON.** *CRC Crit. Ref. Biomed. Eng.*, 1987, vol. 14, 201 **[0245]**
- **BUCHWALD et al.** *Surgery*, 1980, vol. 88, 507 **[0245]**
- **SAUDEK et al.** *N. Engl. J. Med.*, 1989, vol. 321, 574 **[0245]**
- Medical Applications of Controlled Release. CRC Pres, 1974 **[0245]**
- Controlled Drug Bioavailability, Drug Product Design and Performance. Wiley **[0245]**
- **RANGER ; PEPPAS.** *J. Macromol. Sci. Rev. Macromol. Chem.*, 1983, vol. 23, 61 **[0245]**
- **LEVY et al.** *Science*, 1985, vol. 228, 190 **[0245]**
- **DURING et al.** *Ann. Neurol.*, 1989, vol. 25, 351 **[0245]**
- **HOWARD et al.** *J. Neurosurg.*, 1989, vol. 71, 105 **[0245]**
- Guidance for Industry Estimating the Maximum Safe Starting Dose in Initial Clinical Trials for Therapeutics in Adult Healthy Volunteers. Pharmacology and Toxicology. U.S. Department of Health and Human Services Food and Drug Administration Center for Drug Evaluation and Research (CDER, July 2005 **[0249]**
- **FREIREICH et al.** *Cancer Chemother. Rep.*, 1966, vol. 50, 219-244 **[0250]**
- **GILMAN et al.** Goodman and Gilman's: The Pharmacological Basis of Therapeutics. McGraw-Hill, 2001 **[0272]**
- The Merck Manual of Diagnosis and Therapy. Merck Sharp & Dohme Research Laboratories, 1999 **[0272]**
- Cecil Textbook of Medicine. W.B. Saunders, 1996 **[0272]**
- Physicians' Desk Reference. 2007 **[0272]**
- Current Protocols in Immunology. John Wiley and Sons, Inc, 1997 **[0285]**
- **ALTMAN et al.** *Science*, 1996, vol. 274, 94-96 **[0286]**
- **PALLADINO et al.** *Cancer Res.*, 1987, vol. 47, 5074-5079 **[0287]**

- **SCHEIBENBOGEN et al.** *Int. J. Cancer*, 1997, vol. 71, 932-936 [0288]
- **PARMLEY ; SMITH.** *Gene*, 1988, vol. 73, 305-318 [0305]
- **FOWLKES et al.** *BioTechniques*, 1992, vol. 13, 422-427 [0305]
- Relevance of Tumor Models for Anticancer Drug Development. 1999 [0310]
- **KARGER.** *Contributions to Oncology*, 1999 [0310]
- The Nude Mouse in Oncology Research. 1991 [0310]
- Anticancer Drug Development Guide. 1997 [0310]
- **ZHANG ; ROTH.** *In vivo*, 1994, vol. 8 (5), 755-69 [0311]
- **MORRIS et al.** *J La State Med Soc*, 1998, vol. 150 (4), 179-85 [0311]
- **HOSOKAWA et al.** *Transgenic Res*, 2001, vol. 10 (5), 471-8 [0311]
- **KADO et al.** *Cancer Res*, 2001, vol. 61 (6), 2395-8 [0311]
- **WANG et al.** *Int J Pancreatol*, 2001, vol. 29 (1), 37-46 [0311]
- **GHANEH et al.** *Gene Ther*, 2001, vol. 8 (3), 199-208 [0311]
- **BRYANT et al.** *Lab Invest*, 2000, vol. 80 (4), 553-73 [0311]
- **HOUGH et al.** *Proc Natl Acad Sci USA*, 1998, vol. 95 (23), 13853-8 [0311]
- **HERBER et al.** *J Virol*, 1996, vol. 70 (3), 1873-81 [0311]
- **FODDE ; SMITS.** *Trends Mol Med*, 2001, vol. 7 (8), 369-73 [0311]
- **KURAGUCHI et al.** *Oncogene*, 2000, vol. 19 (50), 5755-63 [0311]
- **CRUTE et al.** *Nature Medicine*, 2002, vol. 8, 386-391 [0313]
- **BOLGER et al.** *Antiviral Res.*, 1997, vol. 35, 157-165 [0313]
- **CHEN et al.** *Virol. J*, 23 November 2004, vol. 1, 11 [0313]
- **KERN et al.** *Antimicrob. Agents Chemother.*, 2004, vol. 48, 4745-4753 [0313]
- **BOURNE et al.** *Antiviral Res.*, 2000, vol. 47, 103-109 [0313]
- **BRAVO et al.** *Antiviral Res.*, 2003, vol. 60, 41-49 [0313]
- **BRAVO et al.** *J. Infectious Diseases*, 2006, vol. 193, 591-597 [0313]
- **SIDWELL et al.** *Antiviral Res.*, 2000, vol. 48, 1-16 [0313]
- **MCCAULEY et al.** *Antiviral Res.*, 1995, vol. 27, 179-186 [0313]
- **CAVANAUGH et al.** *J. Virol.*, 1997, vol. 71, 3236-3243 [0313]
- **GUIDOTTI et al.** *J. Virol.*, vol. 69, 6158-6169 [0313]
- **ZHU et al.** *Antimicrobial Agents and Chemother.*, 2006, vol. 50, 3260-3268 [0313]
- **BRIGHT et al.** *Nature*, 2005, vol. 436, 973-978 [0313]
- **HSU et al.** *Nat. Biotechnol.*, 2003, vol. 21, 519-525 [0313]
- **ILAN et al.** *J. Infect. Dis.*, 2002, vol. 185, 153-161 [0313]
- **KNETEMAN et al.** *Hepatology*, 2006, vol. 43, 1346-1353 [0313]
- **MERCER et al.** *Nat. Med.*, 2001, vol. 7, 927-933 [0313]
- **WU et al.** *Gastroenterology*, 2005, vol. 128, 1416-1423 [0313]
- **AYASH-RASHKOVSKY et al.** *FASEB J.*, 2005, vol. 19, 1149-1151 [0313]
- **MOSIER et al.** *Semin. Immunol.*, 1996, vol. 8, 255-262 [0313]
- **MOSIER et al.** *Hosp. Pract.* 1996, vol. 31, 41-4853-5559-60 [0313]
- **BONYHADI et al.** *Mol. Med. Today*, 1997, vol. 3, 246-253 [0313]
- **JOLICOEUR et al.** *Leukemia*, 1999, vol. 13, S78-S80 [0313]
- **BROWNING et al.** *Proc. Natl. Acad. Sci. USA*, 1997, vol. 94, 14637-14641 [0313]
- **SAWADA et al.** *J. Exp. Med.*, 1998, vol. 187, 1439-1449 [0313]
- **SCHITO et al.** *Curr. HIV Res.*, 2006, vol. 4, 379-386 [0313]
- **HAYASHI et al.** *Histol Histopathol*, 2002, vol. 17 (4), 1293-310 [0314]
- **ARICO et al.** *J Interferon Cytokine Res*, 2002, vol. 22 (11), 1081-8 [0314]
- **FLANO et al.** *Immunol Res*, 2002, vol. 25 (3), 201-17 [0314]
- **SAUERMANN.** *Curr Mol Med*, 2001, vol. 1 (4), 515-22 [0314]
- **PLETNIKOV et al.** *Front Biosci*, 2002, vol. 7, d593-607 [0314]
- **ENGLER et al.** *Mol Immunol*, 2001, vol. 38 (6), 457-65 [0314]
- **WHITE et al.** *Brain Pathol*, 2001, vol. 11 (4), 475-9 [0314]
- **DAVIS ; MATALON.** *News Physiol Sci*, 2001, vol. 16, 185-90 [0314]
- **WANG.** *Curr Top Microbiol Immunol.*, 2001, vol. 258, 201-19 [0314]
- **PHILLIPS et al.** *J Psychopharmacol.*, 2000, vol. 14 (3), 244-50 [0314]
- **KAZANJI.** *AIDS Res Hum Retroviruses*, 2000, vol. 16 (16), 1741-6 [0314]
- **SAIF et al.** *Arch Virol*, 1996, vol. 12, 153-61 [0314]
- **HSIUNG et al.** *Rev Infect Dis.*, 1984, vol. 6 (1), 33-50 [0314]
- **SHEPHARD et al.** *Res Vet Sci*, 2003, vol. 74 (2), 187-190 [0315]
- **OTTOLINI et al.** *J Infect Dis*, 2002, vol. 186 (12), 1713-1717 [0315]
- **CULLEY et al.** *J Exp Med*, 2002, vol. 196 (10), 1381-1386 [0315]

- CURTIS et al. *Exp Biol Med*, 2002, vol. 227 (9), 799-802 [0315]
- SUGIYAMA et al. *J Gastroenterol*, 2002, vol. 37 (13), 6-9 [0317]
- BROWN et al. *Am J Reprod Immunol*, 2001, vol. 46 (3), 232-41 [0317]
- VIERLING. *Best Pract Res Clin Gastroenterol*, 2001, vol. 15 (4), 591-610 [0317]
- KLOSE. *Trends Microbiol*, 2000, vol. 8 (4), 189-91 [0317]
- STOTLAND et al. *Pediatr Pulmonol*, 2000, vol. 30 (5), 413-24 [0317]
- BRIELAND et al. *Immunopharmacology*, 2000, vol. 48 (3), 249-52 [0317]
- LEE. *Baillieres Best Pract Res Clin Gastroenterol*, 2000, vol. 14 (1), 75-96 [0317]
- KOEDEL ; PFISTER. *Infect Dis Clin North Am*, 1999, vol. 13 (3), 549-77 [0317]
- NEDRUD. *FEMS Immunol Med Microbiol*, 1999, vol. 24 (2), 243-50 [0317]
- PRELLNER et al. *Microb Drug Resist*, 1999, vol. 5 (1), 73-82 [0317]
- VRIESENDORP. *J Infect Dis*, 1997, vol. 176 (2), 164-8 [0317]
- SHETTY ; ANTIA. *Indian J Lepr*, 1996, vol. 68 (1), 95-104 [0317]
- BALASUBRAMANIAN et al. *Immunobiology*, 1994, vol. 191 (4-5), 395-401 [0317]
- CARBON et al. *Int J Biomed Comput*, 1994, vol. 36 (1-2), 59-67 [0317]
- HABERBERGER et al. *Experientia*, 1991, vol. 47 (5), 426-9 [0317]
- ONDERDONK et al. *Rev Infect Dis*, 1990, vol. 12 (2), 169-77 [0317]
- WICHER ; WICHER. *Crit Rev Microbiol*, 1989, vol. 16 (3), 181-234 [0317]
- SCHELD. *J Antimicrob Chemother*, 1987, vol. 20 (A), 71-85 [0317]
- EMSLIE ; NADE. *Rev Infect Dis*, 1986, vol. 8 (6), 841-9 [0317]
- RIDGWAY et al. *Lab Anim Sci*, 1986, vol. 36 (5), 481-5 [0317]
- QUIMBY ; NGUYEN. *Crit Rev Microbiol*, 1985, vol. 12 (1), 1-44 [0317]
- ONDERDONK et al. *Rev Infect Dis*, 1979, vol. 1 (2), 291-301 [0317]
- SMITH. *Ciba Found Symp*, 1976, 45-72 [0317]
- TAYLOR-ROBINSON. *Infection*, 1976, vol. 4 (1), 4-8 [0317]
- ARENDRUP et al. *Infection*, 2002, vol. 30 (5), 286-91 [0319]
- KAMEI. *Mycopathologia*, 2001, vol. 152 (1), 5-13 [0319]
- GUHAD et al. *FEMS Microbiol Lett*, 2000, vol. 192 (1), 27-31 [0319]
- YAMAGATA et al. *J Clin Microbiol*, 2000, vol. 38 (9), 32606 [0319]
- ANDRUTIS et al. *J Clin Microbiol*, 2000, vol. 38 (6), 2317-23 [0319]
- COCK et al. *Rev Inst Med Trop Sao Paulo*, 2000, vol. 42 (2), 59-66 [0319]
- SHIBUYA et al. *Microb Pathog*, 1999, vol. 27 (3), 123-31 [0319]
- BEERS et al. *J Lab Clin Med*, 1999, vol. 133 (5), 423-33 [0319]
- NAJVAR et al. *Antimicrob Agents Chemother*, 1999, vol. 43 (2), 413-4 [0319]
- WILLIAMS et al. *J Infect Dis*, 1988, vol. 158 (4), 1217-21 [0319]
- YOSHIDA. *Kansenshogaku Zasshi*, 1988, vol. 72 (6), 621-30 [0319]
- ALEXANDRAKIS et al. *Br J Ophthalmol*, 1998, vol. 82 (3), 306-11 [0319]
- CHAKRABARTI et al. *J Med Vet Mycol*, 1997, vol. 35 (4), 295-7 [0319]
- MARTIN et al. *Antimicrob Agents Chemother*, 1997, vol. 41 (1), 13-6 [0319]
- CHU et al. *Avian Dis*, 1996, vol. 40 (3), 715-9 [0319]
- FIDEL et al. *J Infect Dis*, 1996, vol. 173 (2), 425-31 [0319]
- COLE et al. *FEMS Microbiol Lett*, 1995, vol. 126 (2), 177-80 [0319]
- POLLOCK et al. *Nat Genet*, 1995, vol. 9 (2), 202-9 [0319]
- UCHIDA et al. *Jpn J Antibiot*, 1994, vol. 47 (10), 1407-12 [0319]
- MAEBASHI et al. *J Med Vet Mycol*, 1994, vol. 32 (5), 349-59 [0319]
- JENSEN ; SCHONHEYDER. *J Exp Anim Sci*, 1993, vol. 35 (4), 155-60 [0319]
- GOKASLAN ; ANAISIE. *Infect Immun*, 1992, vol. 60 (8), 3339-44 [0319]
- KURUP et al. *J Immunol*, 1992, vol. 148 (12), 3783-8 [0319]
- SINGH et al. *Mycopathologia*, 1990, vol. 112 (3), 127-37 [0319]
- SALKOWSKI ; BALISH. *Infect Immun*, 1990, vol. 58 (10), 3300-6 [0319]
- AHMAD et al. *Am J Kidney Dis*, 1986, vol. 7 (2), 153-6 [0319]
- ALTURE-WERBER E ; EDBERG SC. *Mycopathologia*, 1985, vol. 89 (2), 69-73 [0319]
- KANE et al. *Antimicrob Agents Chemother*, 1981, vol. 20 (5), 595-9 [0319]
- BARBEE et al. *Am J Pathol*, 1977, vol. 86 (1), 281-4 [0319]
- MAESTRONE et al. *Am J Vet Res*, 1973, vol. 34 (6), 833-6 [0319]
- ARATANI et al. *Med Mycol*, 2002, vol. 40 (6), 557-563 [0319]
- BOZZA et al. *Microbes Infect*, 2002, vol. 4 (13), 1281-1290 [0319]
- KURUP et al. *Int Arch Allergy Immunol*, 2002, vol. 129 (2), 129-137 [0319]

- **HORI** et al. *Eur J Immuno*, 2002, vol. 32 (5), 1282-1291 [0319]
- **RIVERA** et al. *J Immuno*, 2001, vol. 168 (7), 3419-3427 [0319]
- **VASSALLO** et al. *Am J Respir Cell Mol Biol*, 2001, vol. 25 (2), 203-211 [0319]
- **WILDER** et al. *Am J Respir Cell Mol Biol*, 2002, vol. 26 (3), 304-314 [0319]
- **YONEZAWA** et al. *J Infect Chemother*, 2000, vol. 6 (3), 155-161 [0319]
- **CACCIAPUOTI** et al. *Antimicrob Agents Chemother*, 2000, vol. 44 (8), 2017-2022 [0319]
- **HONDA** et al. *Mycopathologia*, 1998, vol. 144 (3), 141-146 [0319]
- **FLANDERS** et al. *Autoimmunity*, 1999, vol. 29, 235-246 [0321]
- **KROGH** et al. *Biochimie*, 1999, vol. 81, 511-515 [0321]
- **FOSTER**. *Semin. Nephrol.*, 1999, vol. 19, 12-24 [0321]
- Arthritis and Autoimmunity in Animals. **CROFFORD L.J.** ; **WILDER R.L.** Arthritis and Allied Conditions: A Textbook of Rheumatology. Lee and Febiger, 1993 [0323]
- **WINTER C. A.** et al. Carrageenan Induced Edema in Hind Paw of the Rat as an Assay for Anti-inflammatory Drugs. *Proc. Soc. Exp. Biol Med.*, 1962, vol. 111, 544-547 [0324]
- **EWART** et al. *J Appl Physiol*, 1995, vol. 79 (2), 560-566 [0326]
- **KOMAI** et al. *Br J Pharmacol*, 2003, vol. 138 (5), 912-920 [0326]
- **KENYON** et al. *Toxicol Appl Pharmacol*, 2003, vol. 186 (2), 90-100 [0326]
- **PATH** et al. *Am J Resp & Critical Care Med*, 2002, vol. 166 (6), 818-826 [0326]
- **MARTINS** et al. *Crit Care Med*, 1990, vol. 19, 515-519 [0326]
- **NICOLAIDES** et al. *Proc Natl Acad Sci USA*, 1997, vol. 94, 13175-13180 [0326]
- **TEMANN** et al. *J Exp Med*, 1998, vol. 188 (7), 1307-1320 [0326]
- **COHN** et al. *J. Exp. Med.*, 1997, vol. 186, 1737-1747 [0326]
- **TOMKINSON** et al. *J. Immunol.*, 2001, vol. 166, 5792-5800 [0326]
- **TESCIUBA** et al. *J. Immunol.*, 2001, vol. 167, 1996-2003 [0326]

## 摘要

本發明提供了一種在治療受試者的癌症的方法中使用的複合物，其中，(A)所述複合物包括：(i)白細胞介素 15 (IL-15)；和(ii)白細胞介素 15 受體  $\alpha$  (IL-15Ra) 的可溶形式，其中，所述複合物被施用一抗體，該抗體特異性結合 T 細胞受體或腫瘤相關抗原；(B)所述複合物包括：(i)一種融合蛋白，其包含 IL-15 和 IL-15Ra 的可溶形式；以及(ii)特異性結合癌抗原的抗體；或者(C) 所述複合物包括：(i)IL-15；(ii)IL-15Ra 的可溶形式，或者一種融合蛋白，其包含 IL-15Ra 的可溶形式和免疫球蛋白的 Fc 結構域或其片段；以及(iii)特異性結合癌抗原的抗體。