(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
3 June 2010 (03.06.2010)

PCT

(10) International Publication Number
WO 2010/061359 A1

(51) International Patent Classification:
A61K 39/00 (2006.01) *A61P 35/00* (2006.01)
A61K 31/00 (2006.01)

(21) International Application Number:
PCT/IB2009/055391

(22) International Filing Date:
27 November 2009 (27.11.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
08291118.1 28 November 2008 (28.11.2008) EP

(71) Applicant *(for all designated States except US)*: **SANOFI-AVENTIS** [FR/FR]; 174 avenue de France, F-75013 Paris (FR).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **LEJEUNE, Pascale** [FR/FR]; C/o Sanofi-Aventis Patent Department, 174 avenue de France, F-75013 Paris (FR). **VRIGNAUD, Patricia** [FR/FR]; C/o Sanofi-Aventis Patent Department, 174 avenue de France, F-75013 Paris (FR).

(74) Agent: **BOUVET, Philippe**; c/o Sanofi-Aventis, Patent Department, 174 avenue de France, F-75013 Paris (FR).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

— *with sequence listing part of description (Rule 5.2(a))*

WO 2010/061359 A1

(54) Title: ANTITUMOR COMBINATIONS CONTAINING ANTIBODIES RECOGNIZING SPECIFICALLY CD38 AND CYTARABINE

(57) Abstract: Pharmaceutical composition comprising an antibody specifically recognizing CD38 and cytarabine.

Antitumor combinations containing antibodies recognizing specifically CD38 and cytarabine.

The present invention relates to combinations of monoclonal antibodies directed against CD38 and cytarabine which are therapeutically useful in the treatment of neoplastic diseases.

CD38 is a 45 kD type II transmembrane glycoprotein with a long C-terminal extracellular domain and a short N-terminal cytoplasmic domain. The CD38 protein is a bifunctional ectoenzyme that can catalyze the conversion of NAD^+ into cyclic ADP-ribose (cADPR) and also hydrolyze cADPR into ADP-ribose. CD38 is upregulated and has been implicated in many hematopoietic malignancies.

Monoclonal antibodies 38SB13, 38SB18, 38SB19, 38SB30, 38SB31, and 38SB39, which specifically recognize CD38, are described in PCT application WO2008/047242. Said anti-CD38 antibodies are capable of killing $CD38^+$ cells by three different cytotoxic mechanisms, induction of apoptosis, antibody-dependent cell-mediated cytotoxicity (ADCC), and complement-dependent cytotoxicity (CDC). In addition, these antibodies are able to directly induce apoptosis of $CD38^+$ cells, even without the presence of stroma cells or stroma-derived cytokines. Cytarabine is an anti-metabolic agent used in chemotherapy. There is still a need for novel and efficacious medicaments for treating cancer.

It has now been found, and for this invention, that the efficacy of the humanized anti-CD38 antibodies may be considerably improved when it is administered in combination with at least one substance which is therapeutically useful in anticancer treatments and has a mechanism identical to or different from the one of the humanized anti-CD38 antibodies and which is limited in the present invention to cytarabine.

The term “antibody” is used herein in the broadest sense and specifically covers monoclonal antibodies (including full length monoclonal antibodies) of any isotype such as IgG, IgM, IgA, IgD and IgE, polyclonal antibodies, multispecific antibodies, chimeric antibodies, and antibody fragments. A typical IgG antibody is comprised of two identical heavy chains and two identical light chains that are joined by disulfide bonds. Each heavy and light chain contains a constant region and a variable region. Each variable region contains three segments called “complementarity-determining regions” (“CDRs”)

or “hypervariable regions”, which are primarily responsible for binding an epitope of an antigen. They are usually referred to as CDR1, CDR2, and CDR3, numbered sequentially from the N-terminus. The more highly conserved portions of the variable regions outside of the CDRs are called the “framework regions”.

As used herein, “V_H” or “VH” refers to the variable region of an immunoglobulin heavy chain of an antibody, including the heavy chain of an Fv, scFv, dsFv, Fab, Fab’ or F(ab’)2 fragment. Reference to “V_L” or “VL” refers to the variable region of the immunoglobulin light chain of an antibody, including the light chain of an Fv, scFv, dsFv, Fab, Fab’ or F(ab’)2 fragment.

The 38SB13 antibody comprises at least one heavy chain having an amino acid sequence consisting of SEQ ID NO: 50 and at least one light chain having an amino acid sequence consisting of SEQ ID NO: 38, said heavy chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 1, 2, and 3, and said light chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 4, 5, and 6.

The 38SB18 antibody comprises at least one heavy chain having an amino acid sequence consisting of SEQ ID NO: 52 and at least one light chain having an amino acid sequence consisting of SEQ ID NO: 40, said heavy chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 7, 8, and 9, and said light chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 10, 11, and 12.

The 38SB19 antibody comprises at least one heavy chain having an amino acid sequence consisting of SEQ ID NO: 54 and at least one light chain having an amino acid sequence consisting of SEQ ID NO: 42, said heavy chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 13, 14, and 15, and said light chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 16, 17, and 18.

The 38SB30 antibody comprises at least one heavy chain having an amino acid sequence consisting of SEQ ID NO: 56 and at least one light chain having an amino acid sequence consisting of SEQ ID NO: 44, said heavy chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 19, 20, and

21, and said light chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 22, 23, and 24.

The 38SB31 antibody comprises at least one heavy chain having an amino acid sequence consisting of SEQ ID NO: 58 and at least one light chain having an amino acid sequence consisting of SEQ ID NO: 46, said heavy chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 25, 26, and 27, and said light chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 28, 29, and 30.

The 38SB39 antibody comprises at least one heavy chain having an amino acid sequence consisting of SEQ ID NO: 60 and at least one light chain having an amino acid sequence consisting of SEQ ID NO: 48, said heavy chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 31, 32, and 33, and said light chain comprising three sequential CDRs having amino acid sequences consisting of SEQ ID NOS: 34, 35, and 36.

The hybridoma cell lines producing the 38SB13, 38SB18, 38SB19, 38SB30, 38SB31, and 38SB39 murine anti-CD38 antibodies have been deposited at the American Type Culture Collection (10801 University Bld, Manassas, VA, 20110-2209, USA), on June 21, 2006, under the deposit numbers PTA-7667, PTA-7669, PTA-7670, PTA-7666, PTA-7668, and PTA-7671, respectively (as described in WO 2008/047242).

The term “humanized antibody”, as used herein, refers to a chimeric antibody which contain minimal sequence derived from non-human immunoglobulin. The goal of humanization is a reduction in the immunogenicity of a xenogenic antibody, such as a murine antibody, for introduction into a human, while maintaining the full antigen binding affinity and specificity of the antibody. Humanized antibodies, or antibodies adapted for non-rejection by other mammals, may be produced using several technologies such as resurfacing and CDR grafting. As used herein, the resurfacing technology uses a combination of molecular modelling, statistical analysis and mutagenesis to alter the non-CDR surfaces of antibody variable regions to resemble the surfaces of known antibodies of the target host. The CDR grafting technology involves substituting the complementarity determining regions of, for example, a mouse antibody, into a human framework domain, e.g., see W0 92/22653. Humanized chimeric antibodies preferably have constant regions and variable regions other than

the complementarity determining regions (CDRs) derived substantially or exclusively from the corresponding human antibody regions and CDRs derived substantially or exclusively from a mammal other than a human.

Strategies and methods for the resurfacing of antibodies, and other methods for reducing immunogenicity of antibodies within a different host, are disclosed in US Patent 5,639,641, which is hereby incorporated in its entirety by reference. Antibodies can be humanized using a variety of other techniques including CDR-grafting (EP 0 239 400; WO 91/09967; U.S. Pat. Nos. 5,530,101; and 5,585,089), veneering or resurfacing (EP 0 592 106; EP 0 519 596; Padlan E. A., 1991, *Molecular Immunology* **28(4/5)**: 489-498; Studnicka G. M. *et al.*, 1994, *Protein Engineering*, **7(6)**: 805-814; Roguska M.A. *et al.*, 1994, *PNAS*, **91**: 969-973), chain shuffling (U.S. Pat. No. 5,565,332), and identification of flexible residues (PCT/US2008/074381). Human antibodies can be made by a variety of methods known in the art including phage display methods. See also U.S. Pat. Nos. 4,444,887, 4,716,111, 5,545,806, and 5,814,318; and international patent application publication numbers WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741 (said references incorporated by reference in their entireties).

The anti-CD38 antibodies of the pharmaceutical combination of the present invention are humanized antibodies which recognize CD38 and kill $CD38^+$ cells by apoptosis, ADCC, and CDC. In a further embodiment, the humanized antibodies of the invention are capable of killing said $CD38^+$ cells by apoptosis even in the absence of stroma cells or stroma-derived cytokines.

A preferred embodiment of such a humanized antibody is a humanized 38SB13, 38SB18, 38SB19, 38SB30, 38SB31, or 38SB39 antibody, or an epitope-binding fragment thereof.

The CDRs of the 38SB13, 38SB18, 38SB19, 38SB30, 38SB31, and 38SB39 antibodies are identified by modelling and their molecular structures have been predicted. Thus, in one embodiment, this invention provides humanized antibodies or epitope-binding fragment thereof comprising one or more CDRs having an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, and 36. In a preferred embodiment, a humanized version of 38SB13 is provided, which

comprises at least one heavy chain and at least one light chain, wherein said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 1, 2, and 3, and wherein said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 4, 5, and 6. In another preferred embodiment, a humanized version of 38SB18 is provided, which comprises at least one heavy chain and at least one light chain, wherein said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 7, 8, and 9, and wherein said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 10, 11, and 12. In another preferred embodiment, a humanized version of 38SB19 is provided, which comprises at least one heavy chain and at least one light chain, wherein said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 13, 14, and 15, and wherein said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 16, 17, and 18. In another preferred embodiment, a humanized version of 38SB30 is provided, which comprises at least one heavy chain and at least one light chain, wherein said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 19, 20, and 21, and wherein said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 22, 23, and 24. In another preferred embodiment, a humanized version of 38SB31 is provided, which comprises at least one heavy chain and at least one light chain, wherein said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 25, 26, and 27, and wherein said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 28, 29, and 30. In another preferred embodiment, a humanized version of 38SB39 is provided, which comprises at least one heavy chain and at least one light chain, wherein said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 31, 32, and 33, and wherein said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 34, 35, and 36.

In one embodiment, this invention provides humanized antibodies or fragments thereof which comprise a V_H having an amino acid sequence selected from the group of SEQ ID NOS: 66 and 72. In a preferred embodiment, a humanized 38SB19 antibody is provided which comprises a V_H having an amino acid sequence represented by SEQ ID NO: 66. In another preferred embodiment, a humanized 38SB31 antibody is provided which comprises a V_H having an amino acid sequence represented by SEQ ID NO: 72.

In another embodiment, this invention provides humanized antibodies or fragments thereof which comprise a V_L having an amino acid sequence selected from the group of SEQ ID NOS: 62, 64, 68, and 70. In a preferred embodiment, a humanized 38SB19 antibody is provided which comprises a V_L having an amino acid sequence chosen from the group of SEQ ID NOS: 62 and 64. In another preferred embodiment, a humanized 38SB31 antibody is provided which comprises a V_L having an amino acid sequence chosen from the group of SEQ ID NOS: 68 and 70.

Each of the humanized versions of the 38SB13, 38SB18, 38SB19, 38SB30, 38SB31, and 38SB39 antibodies has been shown to be particularly advantageous as an anticancer agent. The preparation, physical properties and beneficial pharmacological properties thereof are described in WO 2008/047242, which is incorporated by reference herein in its entirety. Generally, the doses used for treating human beings, which depend on factors distinctive to the subject to be treated, are between 1 and 150 mg/kg administered orally or between 1 and 150 mg/kg administered intravenously.

Cytosine arabinoside or cytarabine or araC (brand name: Aracytin™) is an antimetabolic agent (1β-arabinofuranosylcytosine). Cytarabine is rapidly converted into cytosine arabinoside triphosphosphate, which damages DNA when the cell cycle is in the S phase (synthesis of DNA). Rapidly dividing cells, which require DNA replication for mitosis, are therefore most affected. Cytarabine also inhibits both DNA and RNA polymerases and nucleotide reductase enzymes needed for DNA synthesis. Cytarabine is rapidly deaminated in the body into the inactive uracil derivative and therefore is often given by continuous intravenous infusion.

One aspect of the invention is a pharmaceutical composition comprising an anti-CD38 antibody in combination with at least cytarabine. Since the activity of the products depends on the doses used, it is thus possible to use lower doses and to increase the

activity while decreasing the toxicity phenomena. The improved efficacy of a combination according to the invention may be demonstrated by determination of the therapeutic synergy. A combination manifests therapeutic synergy if it is therapeutically superior to the best agent of the study used alone at its maximum tolerated dose or at its highest dose tested when toxicity cannot be reached in the animal species.

.

This efficacy may be quantified, for example, by the $\log_{10}$ cell kill, which is determined according to the following formula:

$$\log_{10} \text{ cell kill} = \text{T-C (days)}/3.32\times T_d$$

in which T - C represents the tumor growth delay, which is the median time in days for the tumors of the treated group (T) and the tumors of the control group (C) to reach a predetermined value (1 g for example), and T_d represents the time in days needed for the volume of the tumor to double in the control animals [T.H. Corbett et al., *Cancer*, **40**: 2660-2680 (1977); F.M. Schabel et al., *Cancer Drug Development*, Part B, Methods in Cancer Research, **17**: 3-51, New York, Academic Press Inc. (1979)]. A product is considered to be active if $\log_{10}$ cell kill is greater than or equal to 0.7. A product is considered to be very active if $\log_{10}$ cell kill is greater than 2.8.

The combination, in which each of the constituents will be present at a dose generally not exceeding its maximum tolerated dose, will manifest therapeutic synergy when the $\log_{10}$ cell kill is greater than the value of the $\log_{10}$ cell kill of the best constituent when it is administered alone and used at its maximum tolerated dose or at its highest dose tested.

The efficacy of the combinations on solid tumors may be determined experimentally in the following manner:

The animals subjected to the experiment, generally mice, are subcutaneously grafted bilaterally with 30 to 60 mg of a tumor fragment on day 0. The animals bearing tumors are randomized based on their tumor size before being subjected to the various treatments and controls. Chemotherapy begins when tumors have reached a predetermined size after grafting, depending on the type of tumor, and the animals are observed every day. The different animal groups are weighed daily during treatment

until the maximum weight loss is reached and subsequent full weight recovery has occurred. The groups are then weighed once or twice a week until the end of the trial.

The tumors are measured 1 to 5 times a week, depending on the tumor doubling time, until the tumor reaches approximately 2 g, or until the animal dies (if this occurs before the tumor reaches 2 g). The animals are necropsied immediately after euthanasia or death.

The antitumor activity is determined in accordance with the different parameters recorded.

Results obtained with combinations of hu38SB19 and cytarabine used at their optimal doses are indicated hereunder as examples.

The present invention also relates, therefore, to pharmaceutical compositions containing the combinations according to the invention.

The constituents of which the combination are composed may be administered simultaneously, semi-simultaneously, separately, or spaced out over a period of time so as to obtain the maximum efficacy of the combination; it being possible for each administration to vary in its duration from a rapid administration to a continuous perfusion.

As a result, for the purposes of the present invention, the combinations are not exclusively limited to those which are obtained by physical association of the constituents, but also to those which permit a separate administration, which can be simultaneous or spaced out over a period of time.

The compositions according to the invention are preferably compositions which can be administered parentally. However, these compositions may be administered orally, subcutaneously or intraperitoneally in the case of localized regional therapies.

The compositions for parental administration are generally pharmaceutically acceptable, sterile solutions or suspensions which may optionally be prepared as required at the time of use. For the preparation of non-aqueous solutions or suspensions, natural vegetable oils such as olive oil, sesame oil or liquid petroleum or injectable organic

esters such as ethyl oleate may be used. The sterile aqueous solutions can consist of a solution of the product in water. The aqueous solutions are suitable for intravenous administration provided the pH is appropriately adjusted and the solution is made isotonic, for example with a sufficient amount of sodium chloride or glucose. The sterilization may be carried out by heating or by any other means which does not adversely affect the composition. The combinations may also take the form of liposomes or the form of an association with carriers as cyclodextrins or polyethylene glycols.

The compositions for oral, subcutaneous or intraperitoneal administration are preferably aqueous suspensions or solutions.

In the combinations according to the invention, the application of the constituents of which may be simultaneous, separate or spaced out over a period of time, it is especially advantageous for the amount of humanized anti-CD38 antibody to represent from 10 to 90 % by weight of the combination, it being possible for this content to vary in accordance with the nature of the associated substance, the efficacy sought and the nature of the cancer to be treated.

The combinations according to the invention are especially useful in the treatment of several types of cancers including (but not limited to) the following: carcinomas and adenocarcinomas, including that of the bladder, breast, colon, head-and-neck, prostate, kidney, liver, lung, ovary, pancreas, stomach, cervix, thyroid and skin, and including squamous cell carcinoma ; hematopoietic tumors of lymphoid lineage, including multiple myeloma, leukemia, acute and chronic lymphocytic (or lymphoid) leukemia, acute and chronic lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, non-Hodgkin lymphoma (e.g. Burkitt's lymphoma) ; hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous (myeloid or myelocytic) leukemias, and promyelocytic leukemia; tumors of mesenchymal origin, including fibrosarcoma, osteosarcoma and rhabdomyosarcoma; tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma, and schwannomas; and other tumors, including melanoma, teratocarcinoma, xeroderma pigmentosum, keratoacanthoma, and seminoma, and other cancers yet to be determined in which CD38 is expressed. They are mainly useful for treating leukemia, lymphoma and cancers resistant to the commonly used anticancer agents as anti-CD38 antibodies have a unique mechanism of action.

Thus, the invention also encompasses the use of the above combinations for the manufacture of a medicament for the treatment of cancer.

Example:

In this example, the effectiveness of an anti-CD38 antibody/cytarabine combination of the invention for tumor growth inhibition was demonstrated in vivo.

The selected tumor model was a transplantable human T-cell acute lymphoblastic leukemia (T-ALL) cell line, DND-41, implanted in SCID mice.

Hu38SB19 was formulated in phosphate buffer saline, without Ca^{2+} and Mg^{2+}, pH7.4. Hu38SB19 was administered intravenously on days 18, 21, 24, 27 after tumor implantation.

Palmo Ara-C, a cytarabine derivative suitable for slow-release administration in mice, was formulated in 3 % ethanol, 1 % polysorbate 80, 96 % water, and was administered subcutaneously on days 18, 21, 24, 27 after tumor implantation.

The results of the experiment are reported in Table 1.

Tumor doubling time = 3.4 days.

The following end points have been used:

- Toxicity was declared at dosages inducing ≥ 20 % body weight loss or ≥ 10 % drug death,
- Antitumor efficacy was determined by evaluating log10 cell kill = (T-C) / [3.32 x (tumor doubling time in days)]

  (T meaning the median time of the treated mice to reach 750 mg and C the median time (26.9 days) of the control mice to reach the same size; tumor-free survivors are excluded from these calculations and are tabulated separately). No antitumor activity was declared for log cell kill < 0.7, and the treatment was declared highly active for log cell kill ≥ 2.8

- Tumor Free Survivors (TFS): correspond to complete regression below the limit of palpation (63 mg) for the entire duration of the study (>100 days post last treatment).
- Therapeutic Synergism: a combination has therapeutic synergism if it is more active than the best single agent of the study (by at least 1 log cell kill).

Toxicity for Palmo Ara-C alone was observed at a dose of 96.3 mg/kg/injection, with 6 drug-related deaths out of 6 mice treated. The highest nontoxic dose (HNTD) for Palmo Ara-C was 58.0 mg/kg/inj (total injected dose = 232.0 mg/kg). The 58.0 mg/kg/inj dose was found to be highly active with a $\log_{10}$ cell kill of 5.4 and 3/6 TFS on day 160. The dose below of 36.0 mg/kg/inj (total injected dose = 144.0 mg/kg) was also found to be highly active with a $\log_{10}$ cell kill of 4.3.The lowest dose of 22.3 mg/kg/inj was active with a 2.0 $\log_{10}$ cell kill.

Regarding hu38SB19, the product was well tolerated at a dose of 40 mg/kg/inj. No toxicity was observed, which can be explained by the lack of cross-reactivity of the antibody with murine CD38. The $\log_{10}$ cell kill was 0.5, indicating that hu38DB19 was not active under these conditions.

The combination of Palmo Ara-C at 96.3 mg/kg/inj and hu38SB19 at 40 mg/kg/inj was toxic, with 5 drug-related deaths out of 6 mice treated, i.e. very similar to what was observed with Palmo Ara-C alone at the same dose. The dose of 58.0 mg/kg/inj of Palmo Ara-C with 40 mg/kg/inj of hu38SB19 was considered to be the HNTD. Remarkably, at this dose, 5 out of 6 mice stayed TFS until the end of the study (day 160). The lower dose of 36.0 mg/kg/inj of Palmo Ara-C with 40 mg/kg/inj of hu38SB19 displayed a $\log_{10}$ cell kill of 8.1 and 3/6 TFS in comparison to a 4.3 $\log_{10}$ cell kill and no TFS for the equitoxic dose of Palmo Ara-C alone and was thus considered highly active. An antitumor activity of 2.7 $\log_{10}$ cell kill was recorded for the lowest dose of the combination (in comparison to 2.0 $\log_{10}$ cell kill for the equitoxic dose of Palmo Ara-C alone. We conclude that the combination shows a therapeutic synergism in comparison to the best single agent of the study, Palmo Ara-C.

Table I: Combination of hu38SB19 and Palmo Ara-C against advanced human T-cell acute lymphoblastic leukemia DND-41 implanted in SCID female mice.

hu38SB19 IV	Palmo Ara-C SC							
40.0 (160.0)	-	18, 21, 24, 27	0/6	-2.1 (19)	6.1	0.5	0/6	HDT - Inactive
-	96.3 (385.2)	18, 21, 24, 27	6/6	-26.6 (31)	-	-	-	Toxic
-	58.0 (232.0)		0/6	-6.1 (31)	60.5	5.4	3/6	HNTD – highly active
-	36.0 (144.0)		0/6	-4.6 (30)	48.6	4.3	0/6	Highly active
-	22.3 (89.2)		0/6	-0.5 (32)	22.1	2.0	0/6	Active
40.0 (160.0)	96.3 (385.2)	18, 21, 24, 27	5/6	-21.6 (34)	-	-	-	Toxic
40.0 (160.0)	58.0 (232.0)		0/6	-6.5 (30)			5/6	HNTD - Highly active
40.0 (160.0)	36.0 (144.0)		0/6	-3.2 (31)	90.9	8.1	3/6	Highly active
40.0 (160.0)	22.3 (89.2)		0/6	-1.7 (19)	30.8	2.7	0/6	active

Tumor doubling time = 3.4 days. Median tumor size at start of therapy = 124-136 mg. Time for median control tumor to reach 750 mg = 26.9 days. BWC = body weight change, T-C = tumor growth delay, HDT = highest dose tested, HNTD = highest nontoxic dose, TFS = tumor free survivors, IV = intravenous, SC = subcutaneous,. Formulations: hu38SB19 = phosphate buffer saline without Ca^{2+} and Mg^{2+}, pH 7.4, Palmo Ara-C = 3 % ethanol, 1 % polysorbate 80, 96 % water.

Claims

1. A pharmaceutical combination comprising an antibody specifically recognizing CD38 and at least cytarabine, wherein said antibody is capable of killing a $CD38^+$ cell by apoptosis, antibody-dependent cell-mediated cytotoxicity (ADCC), and complement-dependent cytotoxicity (CDC).

2. The combination of claim 1, wherein said antibody is a humanized antibody.

3. The combination of claim 2 wherein said antibody comprises one or more complementarity-determining region having an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, and 36.

4. The combination of claim 3 wherein said antibody comprises at least one heavy chain and at least one light chain, wherein said heavy chain has an amino acid sequence represented by SEQ ID NO: 66 and said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 13, 14, and 15, and wherein said light chain has an amino acid sequence selected from the group of SEQ ID NOS: 62 and 64, and said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 16, 17, and 18.

5. The combination of claim 3 wherein said antibody comprises at least one heavy chain and at least one light chain, wherein said heavy chain has an amino acid sequence represented by SEQ ID NO: 72 and said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 25, 26, and 27, and wherein said light chain has an amino acid sequence selected from the group of SEQ ID NOS: 68 and 70, and said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 28, 29, and 30.

6. Use of an antibody specifically recognizing CD38 for the preparation of a pharmaceutical combination according to claim 1 for the manufacture of a medicament for the treatment of cancer.

7. The use of claim 6, wherein said antibody is a humanized antibody.

8. The use of claim 7 wherein said antibody comprises one or more complementarity-determining region having an amino acid sequence selected from the group consisting of SEQ ID NOS: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, and 36.

9. The use of claim 8 wherein said antibody comprises at least one heavy chain and at least one light chain, wherein said heavy chain has an amino acid sequence represented by SEQ ID NO: 66 and said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 13, 14, and 15, and wherein said light chain has an amino acid sequence selected from the group of SEQ ID NOS: 62 and 64, and said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 16, 17, and 18.

10. The use of claim 8 wherein said antibody comprises at least one heavy chain and at least one light chain, wherein said heavy chain has an amino acid sequence represented by SEQ ID NO: 72 and said heavy chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 25, 26, and 27, and wherein said light chain has an amino acid sequence selected from the group of SEQ ID NOS: 68 and 70, and said light chain comprises three sequential complementarity-determining regions having amino acid sequences represented by SEQ ID NOS: 28, 29, and 30.

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2009/055391

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K39/00 A61K31/00 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, WPI Data, Sequence Search, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 1 914 242 A (SANOFI AVENTIS [FR]) 23 April 2008 (2008-04-23) page 3, line 47 - line 51 page 26, line 19 page 26, line 25 - line 33	1-10
X	WO 2006/099875 A (GENMAB AS [DK]; WEERS MICHEL DE [NL]; GRAUS YVO [NL]; OPRINS JUDITH [N) 28 September 2006 (2006-09-28) page 176, line 15 - line 29 page 168, line 10 - page 172, line 8	1-10
X	WO 99/62526 A (MAYO FOUNDATION [US]; LUST JOHN A [US]; DONOVAN KATHLEEN A [US]) 9 December 1999 (1999-12-09) claims 1,2 page 9, line 24 - page 10, line 2	1-10

-/--

[X] Further documents are listed in the continuation of Box C.

[X] See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
23 March 2010	06/04/2010

Name and mailing address of the ISA/	Authorized officer
European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Rodrigo-Simón, Ana

INTERNATIONAL SEARCH REPORT

International application No
PCT/IB2009/055391

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ELLIS JONATHAN H ET AL: "Engineered anti-CD38 monoclonal antibodies for immunotherapy of multiple myeloma" JOURNAL OF IMMUNOLOGY, AMERICAN ASSOCIATION OF IMMUNOLOGISTS, US, vol. 155, no. 2, 1 January 1995 (1995-01-01), pages 925-937, XP002146232 ISSN: 0022-1767 the whole document	1-10
A	PEIPP M ET AL: "FULLY HUMAN CD38 ANTIBODIES EFFICIENTLY TRIGGER ADCC AND CDC OF MULTIPLE MYELOMA AND PLASMA CELL LEUKEMIA CELLS" INTERNET CITATION, [Online] XP002473215 Retrieved from the Internet: URL:http://www.genmab.com/upload/poster_ash_2005_monday_05-12-2005.pdf> [retrieved on 2008-03-18] the whole document	1-10
A	PEIPP M ET AL: "FULLY HUMAN CD38 ANTIBODIES EFFICIENTLY TRIGGER ADCC OF MULTIPLE MYELOMA CELL LINES AND PRIMARY TUMOR CELLS" BLOOD, AMERICAN SOCIETY OF HEMATOLOGY, US, vol. 106, no. 11, PART 01, 1 November 2005 (2005-11-01), page 944A,ABSTRACTNO.3377, XP009069301 ISSN: 0006-4971 the whole document & PEIPP M ET AL: "Fully human CD38 antibodies efficiently trigger ADCC of multiple myeloma cell lines and primary tumor cells" BIOSIS,, 1 January 1900 (1900-01-01), XP002389878	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/IB2009/055391

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 1914242 A	23-04-2008	AR 063488 A1	28-01-2009
		AU 2007311511 A1	24-04-2008
		CA 2663209 A1	24-04-2008
		CL 29952007 A1	16-05-2008
		CN 101616933 A	30-12-2009
		CR 10679 A	24-06-2009
		EA 200970393 A1	30-12-2009
		EP 2076540 A2	08-07-2009
		WO 2008047242 A2	24-04-2008
		KR 20090078330 A	17-07-2009
		SV 2009003203 A	27-10-2009
		US 2009304710 A1	10-12-2009
		UY 30655 A1	31-05-2008
WO 2006099875 A	28-09-2006	AU 2006226733 A1	28-09-2006
		CA 2602375 A1	28-09-2006
		EA 200702053 A1	28-02-2008
		EP 1866338 A1	19-12-2007
		JP 2008533977 T	28-08-2008
		KR 20070116137 A	06-12-2007
		US 2009148449 A1	11-06-2009
WO 9962526 A	09-12-1999	AU 770718 B2	26-02-2004
		AU 4418599 A	20-12-1999
		CA 2329940 A1	09-12-1999
		EP 1085882 A2	28-03-2001
		JP 2003524587 T	19-08-2003