



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2518372 C 2013/07/09

(11)(21) **2 518 372**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(22) Date de dépôt/Filing Date: 2005/09/30

(41) Mise à la disp. pub./Open to Public Insp.: 2006/03/30

(45) Date de délivrance/Issue Date: 2013/07/09

(30) Priorité/Priority: 2004/09/30 (US60/614533)

(51) Cl.Int./Int.Cl. *G01N 33/569* (2006.01),
G01N 33/58 (2006.01)

(72) Inventeur/Inventor:
ZHENG, JIAN, US

(73) Propriétaire/Owner:
ORTHO-CLINICAL DIAGNOSTICS, INC., US

(74) Agent: NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) Titre : PEPTIDES POUR DISTINGUER DES PRIONS

(54) Title: PEPTIDES FOR DISCRIMINATION OF PRIONS

(57) **Abrégé/Abstract:**

The aim of the present invention is to provide a non-intrusive way to isolate, concentrate and monitor the TSE disease-related pathogenic prion protein. The invention described several peptides and their ability to capture PrP^{Sc} from brain homogenate of prion disease infected animal and human. These peptides do not capture cellular prion protein from individuals with no prion disease.



Abstract:

The aim of the present invention is to provide a non-intrusive way to isolate, concentrate and monitor the TSE disease-related pathogenic prion protein.

- 5 The invention described several peptides and their ability to capture PrP^{Sc} from brain homogenate of prion disease infected animal and human. These peptides do not capture cellular prion protein from individuals with no prion disease.

PEPTIDES FOR DISCRIMINATION OF PRIONS

5 Background

Prion diseases or transmissible spongiform encephalopathies (TSE) have captured the interest of the scientific community, not only because they challenged conventional thought about the nature of infectious agents, but also because of the potentially serious public health threat they pose to food and blood safety. Furthermore, the possibility of disease transmission between animals and humans has been dramatically underscored in recent years by the epidemic occurrence of a new variant disease form, vCJD. The present invention described the use of synthetic peptides for the diagnosis of Transmissible Spongiform Encephalopathies (TSE) diseases in animals and humans.

Prions – A Pathogenic Agent Causing TSE

Transmissible spongiform encephalopathies (TSEs) comprise a group of rapidly progressing, neurodegenerative fatal diseases that affect both humans and animals. TSEs have clinical and neuropathological characteristics which include devastating dementia, pyramidal and extrapyramidal signs with myoclonus, multifocal spongiform changes, astrogliosis, amyloid plaques, neuronal loss, absence of inflammatory reaction and are usually characterized by a long incubation period.

It was once suggested that TSE diseases might be caused by "slow viruses" or viroids (Gajdusek 1977). However, the extreme resistance of scrapie infectivity to radiation, nucleases, and other reagents damaging to genetic materials are inconsistent with the "virus" theory. All these "unusual" characteristics of the TSE infectious agent led Dr. Stanley Prusiner to propose the concept of "prions" in 1982 (Prusiner 1982). Prion (PrP), which stands for nucleic acid-free *proteinaceous infectious* particle, is a glycoprotein present in humans and animals. The cellular form of this protein (PrP^C) has

two N-link glycosylation sites and a GPI anchor at the C-terminus. It has been most commonly found in neurons, and, to a much lower extent, it has also been found in other cells such as leucocytes, monocytes and platelets (Holada 2000). The transmissible scrapie disease form of the prion protein (PrP^{Sc}) is a protease resistant isoform of its cellular precursor and is predominantly found in brain. At much lower level, it has also been found in tonsil, spleen, and lymph nodes in vCJD patients (Parizek 2001). As a result of Prusiner's concept of the "prion" as an infectious agent responsible for scrapie disease, and by extension, that of all TSE diseases gave rise to the notion of what are commonly referred to as Prion diseases to describe a class of pathologies believed to be linked to this protein.

Human prion disease is known as Creutzfeldt-Jakob disease (CJD) and usually affects elderly persons. Depending on disease phenotype, sporadic (sCJD) accounts for most of CJD, while about 10% are familial CJD (fCJD). Only recently, a new variant of CJD (vCJD) has emerged which occurs in young adults. Since 1995, more than 100 cases have been reported. Prion diseases affect many species of domestic and wild animals, manifestations of which are often given unique names, for example, scrapie in sheep and goats (McGowan 1922), chronic wasting disease (CWD, Williams 1980) in cervids and bovine spongiform encephalopathy (BSE, or "mad cow" disease, Wells 1987) in cattle.

Characteristics of PrP^C and PrP^{Sc}

According to the Prusiner's "protein only" hypothesis, the transmissible pathogen causing prion diseases is a protein. The major component of the prion appears to be abnormally folded PrP, designated as PrP^{Sc} (disease-specific, "scrapie" isoform). It is believed that PrP^{Sc} is generated from its normal counterpart, PrP^C, through a change in conformation (Cohen, 1998). Although there is still ambiguity concerning the mechanism of the conversion, it is still broadly accepted that in the presence of PrP^{Sc}, normal PrP^C, acting

as a substrate, undergoes a conformational structure change, and becomes PrP^{Sc} through an autocatalytic process and results in PrP^{Sc} aggregation and amyloid rod formation, hence causing cell death (Hope 1986, Horwich 1997). The structural change from PrP^{C} to PrP^{Sc} is most supported by a crucial

5 conformational change, involving a substantial increase in the amount of Beta-sheet structure of the protein, with possibly a small decrease in the amount of alpha-helix, indicated by circular dichroism and infrared spectroscopy (Pan 1993, Caughey 1991). It appears that the efficiency of conversion from PrP^{C} into PrP^{Sc} depends on the degree of sequence

10 homology between the two PrP conformers. Because this unique mechanism, prion diseases can be readily transmitted within the same species, and when the PrP sequence homology is sufficient, from species to species (Raymond, 2000). Therefore, if correct, the proposed mechanism of transmission of prion diseases provides the potential to trigger epidemics by transmission of these

15 brain disorders between animals and humans.

Protease resistance is another characteristic that distinguishes PrP^{Sc} from PrP^{C} . In cultured cells and brain or in samples from many patients with GSS, PrP^{Sc} is smaller than its cellular precursor PrP^{C} . Even though cellular prion

20 and scrapie prion are two isoform of same *PRNP* genomic product, PrP^{C} is completely degraded by Proteinase K treatment while PrP^{Sc} undergoes only limited digestion. The digestion yields a form of protein referred to as PrP 27-30 in which the N-terminus has been removed. PrP 27-30 has been postulated to be the PrP^{Sc} core required for PrP^{C} hosted PrP^{Sc} replication.

25 The protease treated prion molecule, PrP 27-30 or PrP^{res} , is tightly linked to scrapie infectivity (Gabizon 1988), and provides additional evidence that PrP^{Sc} is an infectious protein.

An additional attribute, perhaps linked to the significant increase in Beta-sheet

30 structure and concomitant protease-resistance, is the observed difference in solubility between PrP^{Sc} and PrP^{C} . While PrP^{C} is a soluble protein, the PrP^{Sc}

isoform is highly insoluble. Furthermore, PrP^C is found attached to the surface of neurons through a GPI tail anchored into membrane (Shyng 1994) while PrP^{res} is found in the cytoplasm of affected cells (Taraboulos 1990), most likely associated with late endosomal and lysosomal compartments (Arnold
 5 1995), and PrP^{Sc} is also localized in amorphous aggregates in enriched fractions from infected brain (Meyer 1986).

There is mounting evidence indicating a tight linkage between scrapie infectivity and PrP 27-30. Even in the purest samples, the estimated ratio of
 10 PrP molecules to infectious units is $\sim 10^4$ to 10^5 (Horwich 1997, Bolton 2001). At such low levels of infectivity, it is possible that other components, co-factors, or covalent modifications, are required for infectivity. The transgenic studies on the susceptibility of mice expressing chimeric human-mouse PrP^C suggest the presence of at least one host factor other than PrP^C, tentatively
 15 termed factor X, which might function as a molecular chaperone in the formation of PrP^{Sc} (Telling 1995).

Infectivity and Transmissibility of Prion Diseases

Prion diseases are transmissible. The transmissibility of animal prion
 20 diseases has long been established experimentally by inoculation of brain homogenates from affected animals into healthy ones, such scrapie from sheep to sheep (Cuillé 1936) and across species to goat (Pattison 1957), kuru and CJD from humans to chimpanzees (Gajdusek 1966, Gibbs 1968). The significant breakthrough was the successful transmission of scrapie to
 25 mice (Chandler 1961) which greatly facilitated TSE research by providing an experimental model. The cause of recent BSE in cattle and new variant CJD in human (vCJD) was considered a consequence of dietary exposure to the mix of scrapie sheep carcasses rendered for animal feed in the case of BSE
 30 (Brown 1997) and to beef from cattle affected with BSE in the case of vCJD (Bruce 1997). The link between vCJD and BSE is further supported by the neuropathologic evidence obtained from BSE-adapted macaques, the

nearest model to humans, and from the study on inbred mice inoculated with the agent causing BSE and vCJD (Lasmézas 1996). Of particular concern to an epidemic expanding of CWD in mule deer and elk in North America is whether CWD, like BSE, could be transmitted to humans who may be
5 exposed to the disease through hunting, or handling and eating infected meat. Tragic, unintended transmission of prion disease in humans has been documented, such as the kuru epidemic caused by cannibalistic ingestion of brain tissue from the deceased, and the iatrogenic transmission of CJD through the use of hormones, tissue transplants, and contaminated medical
10 devices.

There is no hard evidence indicating any of CJD diseases is related to animal TSEs that may have crossed species barriers. The epidemic of kuru has provided the largest body of evidence of acquired human prion disease.
15 Although no vCJD patient has been documented as a victim of human-to-human transmission, the close link between BSE and vCJD attracted considerable attention. Concerns about human infection have been based on the observation that PrP^{Sc} is readily detectable in BSE and vCJD lymphoreticular tissues but not in classic CJD (Hill 1997), followed by the
20 presumption that scrapie pathogen from sheep passage to cattle may have altered host range and become more adaptable to human. Experimental precedents for such behavior are well known: passage of mouse-adapted strains of scrapie through hamsters altered their transmissibility on back passage to mice (Kimberlin 1987, Kimberlin 1989); human strains of kuru or
25 CJD did not transmit to ferrets or goats until passaged through primates or cats (Gibbs 1979); and a bovine strain of BSE did not transmit to hamsters until passaged through mice (Foster 1994). Alternatively, if BSE originated from a spontaneous mutation in cattle, experimental studies of species susceptibility to this new strain of transmissible spongiform encephalopathy
30 (TSE) had not sufficiently advanced to predict that humans would not be susceptible.

Study on human CJD and vCJD disease indicated that genomic susceptibility might yet be another factor that may influence the spread of TSE in humans. The majority of sporadic CJD patients were found to be homozygous for
 5 Met/Met or for Val/Val at codon 129 (Belay 1999). Nevertheless, all reported vCJD cases have been found to be homozygous for Met/Met.

The size and duration of vCJD epidemic still remains uncertain. Depending on the assumptions made and the modeling calculations employed, different
 10 predictions were proposed. One estimation of total vCJD predicts as few as 205 cases (Valleron 2001). On the other hand, another prediction for vCJD mortality for the next 80 years ranges from 50 to 50,000 if infection comes only from BSE. It could reach up to 150,000 if BSE is proven to infect sheep and if subsequently it is allowed to enter human food chain (Ferguson 2002).
 15 Although it is impossible to make accurate predictions if the necessary parameters are either mistaken or not available, one thing is certain that if vCJD infectivity is present in blood, any prediction will be an underestimate. In addition, vCJD has been proven to be a new disease entity and not simply the result of increased surveillance of CJD in humans (Hillier 2002).

20

Countermeasures have been taken by government to eliminate the spread of BSE incidence. Ruminant protein feed was banned in US and UK (1988). A series of measures have also been taken to prevent potentially infected meat from entering human food chain. To further reduce the human risk, FDA and
 25 CBER has issued a new policy in Aug. 2001, which indefinitely defers any donor who stayed cumulative ≥ 6 month during 1980-1996 in the United Kingdom (FDA 2001).

Diagnostic Assay for Prion Disease

30 The development of sensitive and reliable assays for prion detection is absolutely essential for disease surveillance, risk assessment, and when

combined with future therapeutics, for disease prevention and eradication. Currently, there are basically three assay formats for diagnosis of prion diseases. **(1) Animal infectivity bioassays** are by far the most sensitive method for the measurement of infectious prion in experimental scrapie in rodents, usually accomplished by intracerebral injection of brain homogenates from sick animals into recipient animals. However, the quantitative measurement of prion disease infectivity in different animal species is limited due to the "species barrier" and distinct prion "strains" exist that differ in terms of pathology, incubation time, and molecular characteristics of PrP^{Sc}. Therefore, this time-consuming, expensive postmortem diagnosis is mostly used as a research tool for distinguishing different prion strains in rodents, and serves as a reference for calibration of infectious brain material. **(2) Current PrP Immunoassays** are based on the detection of protease-resistant PrP^{Sc}, the only known molecular hallmark of all prion diseases. For many years, the detection of PrP^{Sc} by immunochemical methods (immunohistochemistry and Western blotting) has provided the most accurate diagnosis for prion diseases in animals and humans (Schaller 1999, Biffiger 2002). They are widely used for postmortem diagnosis. Many monoclonal and polyclonal antibodies have been raised against various regions of PrP for this purpose such as widely used 3F4, 6H4 described in US4806627 and EP0861900. However, only few were claimed to be able to discriminate between PrP^C (often present in much larger quantities) and PrP^{Sc}. Monoclonal antibodies reported by Korth in 1997 and by Paramithiotis in 2003 were both IgMs and no diagnosis assay was developed by these antibodies. Consequently, almost all current immunochemical methods require a step to reduce or eliminate a PrP^C, usually by nonspecific proteolysis such as proteinase K (PK) digestion, prior to the detection of PrP^{Sc}. Such pretreatment cleaves the first 60 – 70 residues from PrP^{Sc} to yield a PK-resistant PrP27kDa-30 kDa core called PrP^{res}. Anti-PrP antibodies that recognize the remaining C-terminal region of the protein can then be used to detect the N-terminally truncated PrP^{Sc}, or PrP^{res}, present only in pathological

samples. For immunohistochemical staining, tissue sections are also pretreated, usually by acid hydrolysis to reduce the PrP^{C} related background. Therefore, PrP^{res} is a surrogate for the precursor PrP^{Sc} in these immunoassays. Among the various immunoassay formats, Western blotting
 5 has the advantage of revealing detailed molecular patterns of PrP based on the migration of di-, mono- and unglycosylated PrP bands. This method also has been widely used for distinguishing distinct brain PrP^{res} subtypes in human and animal prion diseases. Besides Western blot, other assay formats have now been developed for higher sample throughput, increased sensitivity,
 10 and better quantification, including traditional ELISA, dissociation-enhanced lanthanide-fluorescence-immunoassay (DELFI, Barnard 2000 and a method described in US20020137114A1) and conformation-dependent immunoassay (CDI) combined with ELISA and fluorescence detection (Safar 1998, US 20010001061A1, US20020001817A1). However, regardless of the format,
 15 PrP^{Sc} can be differentiated from PrP^{C} only after the mandatory PK digestion, which may be difficult to optimize for disparate biological samples. (3) Other methods have also been described for sample treatment including immunohistochemistry of third eyelid lymphoid tissue for preclinical diagnosis of ovine scrapie (O'Rourke 2000, US6165784, US6261790), chemical
 20 treatment with sodium phosphotungstate to enrich PrP^{Sc} from brain and from other peripheral tissue homogenates (Wadsworth 2001), and detection of a new isoform of the prion protein in the urine of infected animals and humans (Shaked 2001b, WO0233420A2). Other detection systems were also documented including capillary electrophoresis, and Fourier transform
 25 infrared spectroscopy. These methods are still in their initial stages of development and are technically complex. In addition to the traditional identification of pathogenic prion by eliminating cellular prion followed by non-discriminatory anti-prion antibody recognition, other reagents were found to be able to differentiate PrP^{Sc} from PrP^{C} , such as plasminogen and fibrinogen.
 30 The evidence provided suggested that a property common to PrP^{Sc} of various species, rather than prion primary sequence or the specific tertiary structure

of individual PrP^{Sc} molecules, could be responsible for binding to plasminogen (Fischer 2000, Maissen 2001). The application for the use of plasminogen and other serum/plasma proteins for the capture and detection of pathogenic prion protein has been described in WO0200713 and in
 5 US20010053533A1 (Aguzzi 2001).

All current manufactured prion diagnosis assays use brain tissue as a sample source. The European Commission in 1999 evaluated 4 BSE test kits from different manufacturers (Moynagh 1999). They all required a separate sample
 10 preparation procedure. Depending on the kit instructions, the brain tissue homogenate needed to be processed, including denaturation, PK digestion or PrP^{Sc} enrichment. The assay detection systems employed in DELFA, immunoblot, or in plate ELISA formats used either chemiluminescent or a colorimetric substrate.

15

Challenges to antemortem diagnosis of prion disease

An issue common to PrP^{Sc}-based antemortem assays is whether PrP^{Sc} is present in peripheral tissues or body fluids. Because of technical difficulties, little experimental data on the presence of PrP^{Sc} or its associated infectivity in
 20 body fluids are available, and this subject remains controversial. In the hamster model of scrapie, a low level of infectivity can be detected in blood. Although the infectivity in lymphocyte-rich buffy-coat derived from diseased hamster blood is greater than in plasma, it only accounts for relatively a small portion when compared to whole blood inoculums. The molecular definition of
 25 this infectious agent present in the blood is not clear. Searching for risk factors and possible sources of infection in sporadic CJD patients revealed no significant correlation of disease to diet, blood transfusion or receiving other blood product. Although early reports indicated the possible presence of infectivity in blood obtained from CJD patients after intracerebral inoculation
 30 to mice (Manuelidis 1985, Tateishi 1985), The highest amount of infectivity or PrP^{res} is invariably found in the central nervous system (CNS), but not

consistently found in peripheral tissues in classical human prion diseases, except in the case of vCJD. The presence of readily detectable PrP^{Sc} in the peripheral lymphoreticular tissues such as tonsils, spleen, and lymph nodes of vCJD patients has raised a serious concern that abundant amounts of PrP^{res} present in lymphoreticular tissues could interact with the circulatory system, and as a consequence, trace amounts of PrP^{Sc} may be present in blood of vCJD patients for possible blood transmission. Other TSE infectivity in blood has also been demonstrated in various experimental animals. Most blood for infectivity studies was obtained from TSE –adapted rodents such as mice and hamsters. Mice-adapted BSE, mice-adapted vCJD has been established through intracerebral and intravenous transmission. The only exception model was a study conducted in the sheep. In this experiment, a sheep transfused with whole blood, taken from another sheep inoculated with BSE brain lysate, developed symptoms of BSE (Houston 2000, Hunter 2002). However, these experimental results yet need to be fully evaluated. It is anticipated that finding of such infectious agent in blood would help us to better understand the relationship between PrP^{Sc} and TSE disease.

It is important to note that the harsh sample treatment to eliminate PrP^C background may not be suitable for other peripheral tissue specimens or body fluids due to differing protein content, the difficulty of applying this assay to large numbers of samples, the inevitable elimination of protease-sensitive PrP^{Sc} folding intermediates or even a fraction of authentic PrP^{Sc}, thus reducing the sensitivity of detection. This may be especially relevant for assays using peripheral tissues and body fluids, as only low levels of PrP^{Sc} may be present (Horiuchi 1999, Jackson 1999, Swietnicki 2000). These concerns necessitate the development of immunological reagents with a high affinity for PrP^{Sc}, allowing for specific detection without the need for proteolytic treatment.

30

Discovery of novel capture reagent for PrP^{Sc} detection

Whether or not one accepts the “protein only” or “prion only” hypothesis, there are continuous efforts underway to search for agents or molecules other than prion that may contribute to the pathogenesis of prion disease. This search is driven by many unanswered questions. For example, synthesized prion protein, free of any contamination, does not cause disease; the mechanism that triggers the conversion of normal PrP^{C} to the pathogenic PrP^{Sc} isoform is unknown. Another unresolved question involves the various prion disease phenotypes observed in animals and humans, defined by disease incubation period, glycosylation level and lesion patterns. After serial passage in inbred mice homozygous for a single *PRNP* genotype, all the scrapie strains retained their original disease profile. These observations led investigators to question whether varied phenotypic strains were dominated by different conformational isoforms of same cellular prion precursor, or whether there is another factor that determines the phenotype of the inheritable strain. In fact, in *vitro* conversion models PrP^{res} formed in cell-free reactions has never been shown to constitute new TSE infectivity in animals (Caughey 2003). These questions led many to believe that there is a missing element, dubbed “protein X” as Prusiner suggested, yet to be discovered.

The presence of a tightly bound RNA or DNA molecule in the prion particle was proposed to explain propagation of different strains of scrapie agent with distinct phenotypes in animals homozygous for the *PRNP* gene (Weissmann 1991). Analysis of highly purified scrapie prions by return refocusing gel electrophoresis revealed the small size of remaining nucleic acids (Kellings 1992). In a recent report, however, Narang indicated that animals inoculated with ssDNA purified from scrapie-hamster brains mixed with non-pathogenic prion developed clinical disease (Narang 2002). Based on his findings, he postulated that the “accessory protein” coded by the ssDNA may be involved in PrP^{C} to PrP^{Sc} conversion. Based on those *in vitro* conformation and conversion studies, it was hypothesized that DNA would act as a guardian of the PrP^{Sc} conformation as well as a catalyst to facilitate PrP^{Sc} conversion and

aggregation (Cordeiro 2001). Most recently, it was reported that stoichiometric transformation of PrP^{C} to PrP^{res} in vitro requires specific RNA molecules (Deleault 2003). The anti-nucleic acid monoclonal antibody developed by Ortho-Clinical Diagnostics that can discriminatively capture PrP^{Sc} but not PrP^{C}
 5 (US60/434,627, US60/446,217) is another evidence to demonstrate the association of PrP^{Sc} to nucleic acids.

It is known that PrP^{Sc} isolated from diseased brain is also associated with a variety of glycans. Those include 1,4-linked glucose units in prion rods,
 10 sphingolipids, polysaccharides and other membrane components in PrP^{Sc} aggregates (Appel 1999, Klein 1998), and sulfated proteoglycan in prion amyloid plaques (Snow 1990), a property that has been exploited in immunohistochemistry, where binding by heparan sulfate antibodies (anti-HS) and heparan sulfate proteoglycan antibodies (anti-HSPG), has been shown to
 15 correlate with abnormal PrP as early as 70 days post-infection and throughout the course of the disease (McBride 1998). Through a mechanism that is perhaps different from that by which nucleic acids participate in the conversion of PrP^{C} to PrP^{Sc} , glycan also convert cellular prion protein into Beta-sheet conformation. In *vitro* conversion from PrP^{C} to PrP^{Sc} and in prion
 20 infectivity reconstitution experiments, sulfate glycans have been shown either to facilitate the conversion or to escalate infectivity (Wong 2001, Shaked 2001a, Diaz-Nido 2002). With recombinant GST::full-length prion and GST::prion fragment, Warner recently demonstrated direct binding of recombinant prion to heparin and heparan sulfate (Warner 2002). The
 25 peptide region 23-52 in prion sequence was positive in all HS and HSPG binding tests. Since the peptide failed to compete with full-length prion for binding to heparin, the author suggested that there might be another major GAG-binding site in intact PrP^{C} . Another interesting observation is that plasminogen has been reported to bind brain-derived PrP^{Sc} , but not to PrP^{C} .
 30 Although it has not been demonstrated that plasminogen has a direct interaction with PrP^{Sc} , a binding site is suggested within the Kringle region of

plasminogen, a region that has a known affinity for heparin. Another noteworthy observation is that GAGs from different species (bovine and porcine) or from different organs (lung, kidney and intestine) have shown different affinities for prion binding. The difference in affinity may be due to prion sequence itself, or may depend on the presence of particular sugar unit in the tested GAGs.

In light of these observations, a number of peptides were proposed that are designed to selectively capture PrP^{Sc} but not PrP^{C} through peptide affinity to unique PrP^{Sc} conformation or to PrP^{Sc} associated molecules. They were screened by the ability to identify and to capture PrP^{Sc} from homogenates of diseased brains by immunoprecipitation without protease pretreatment.

(1) Heparin/heparin sulfate binding domain peptides:

Glycans (GAG) such as heparin and heparin sulfate were associated with PrP^{Sc} amyloid aggregates. Because the association affinity was much higher in $\text{GAG}::\text{PrP}^{\text{Sc}}$ than in $\text{GAG}::\text{PrP}^{\text{C}}$ it is possible to use peptides characterized as glycan binding domain to selectively capture PrP^{Sc} but not PrP^{C} . For this reason, peptides described as heparin or heparin sulfate binding domain were synthesized: (1) WQPPRARI of carboxy-terminal fibronectin (Woods 1993, Hines 1994) (2) NWCKRGRKQCKTH of amyloid protein precursor (Small 1994), (3) NYKKPKL of N-terminal fibroblast growth factor (FGF)-1 (Lou 1996) and (4) KDFLSIELVRGRVK of the C-terminal G-domain of the laminin alpha1 chain (Yoshida 1999).

(2) “Condensed” Kringle peptides:

Kringle region was involved in the selective binding of plasminogen to PrP^{Sc} . It was also known that Kringle region had heparin/heparin sulfate binding activity (Mizuno 1994) for which positively charged amino acids (such as Arg and Lys) were involved (Soeda 1989). In a separate publication, two conserved tripeptides “YYR”, or rather three discontinued “YYX” in prion

sequence as suggested, were found to interact with PrP^{Sc} but not interact with PrP^C (Paramithiotis 2003, WO0078344A1). Interestingly found in human plasminogen sequence, there were four Tyr-Arg-Gly sequences in four out of five Kringle regions (Kringle region 1, 3, 4 and 5); there were six Tyr-Arg or Arg-Tyr or Arg-Lys sequences in plasminogen. Furthermore, several distant Tyr, Arg or Lys residues were brought closely together by disulphide bridges formed in Kringle loops (Figure 1). Based on these observations, it was postulated that amino acids Tyr, Arg and Lys in plasminogen Kringle region and in prion sequence, possibly through an interaction to glycan that was associated with PrP^{Sc} complex, could be accounted for the selective binding to PrP^{Sc}. For this reason, two "condensed Kringle" peptides YRGYRGYRGYRG and YRGYRGYKGYGYRG were synthesized.

(3) Nucleic acid binding peptides:

Nucleic acids are another category of molecules that associated with PrP^{Sc} aggregates. Anti-DNA antibodies have been used effectively to capture PrP^{Sc}. Histones are a group of proteins known to bind nuclear DNA. Therefore, three peptides (1) AQKKDGKKRKRSRKESYSIYV of H2B(21-41), (2) ARTKQTARKSTGGKAPRKQLA of H3(1-21), and (3) SGRGKGGKGLGKGGAKRHRKVLR of H4(2-24) were synthesized to evaluate their ability of the capture of PrP^{Sc}.

Summary of the Invention

The aim of the present invention is to provide a non-intrusive way to isolate, concentrate and monitor the TSE disease-related pathogenic prion protein. The invention describes several peptides and their ability to capture PrP^{Sc} from brain homogenate of prion disease infected animal and human. These peptides do not capture cellular prion protein from individuals with no prion disease. The evidence provided in support of this invention demonstrated that PrP^{Sc} is associated with high affinity to many other molecules such as nucleic acid and glycans as investigated. The evidence also demonstrated that the

association was strong, resistant to PK digestion treatment, and that PrP^{Sc} could be readily isolated by peptides through recognition of such associated binding partner.

5 This invention relates to the use of described peptides to capture PrP^{Sc} through glycans or nucleic acids associated with high affinity to PrP^{Sc}, in combination with any prion sequence-specific antibody for the detection of PrP^{Sc}.

10 In another aspect, this invention relates to the peptides, as described above, that preferably binds to pathogenic prion protein but not to the normal cellular form of prion protein.

In another aspect, this invention relates to the peptides, as described above,
15 for the detection of PrP^{Sc} through high affinity recognition of associated glycans or nucleic acids in combination of prion sequence specific antibodies.

In another aspect, this invention relates to the peptides, as described above,
for the isolation, purification, capture, elimination and monitoring PrP^{Sc} in
20 biological reagent production.

In another aspect, this invention relates to compositions and kits for
determining the presence of PrP^{Sc}, comprising peptides, as described above,
for either capture or for detection step in the assay procedure.

25

In another aspect, this invention relates to compositions and kits for
determining the presence of PrP^{Sc} antibody produced in response to high
affinity associated glycans or nucleic acids as a binding partner to pathogenic
prion protein.

30

In yet another aspect, this invention relates to anti-PrP^{Sc} antibodies and their production using the said glycans or nucleic acids that can interact with and/or associate with PrP^{Sc}, and their use in detecting glycan::PrP^{Sc} or nucleic acid::PrP^{Sc} complex and prion disease infection.

- 5 In another aspect, this invention relates to a non-harsh sample treatment procedure involving glycanase or nuclease digestion for the benefit of the use of described peptides as described above.

In one aspect, the invention relates to a method for detection of PrP^{Sc} comprising: first contacting a sample with a peptide consisting of an amino acid sequence selected from
 10 the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRSRKESYSIYV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9) SGRGKGGKGLGKGGAKRHRKVLR; and then detecting PrP^{Sc} with a
 15 labeled prion specific antibody, wherein the peptide specifically captures PrP^{Sc} and the labeled prion specific antibody detects the captured PrP^{Sc}.

In another aspect, the invention relates to an immunoassay for detecting PrP^{Sc} comprising: providing a solid support having bound thereto a peptide, consisting of an amino acid sequence selected from the group consisting of: (SEQ ID NO:1)
 20 WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRSRKESYSIYV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9) SGRGKGGKGLGKGGAKRHRKVLR; contacting the solid support with a sample; washing the support to remove any unbound sample; contacting the solid
 25 support with a labeled prion specific antibody; and carrying out a detection step to determine if PrP^{Sc} are bound to the solid support, wherein the peptide specifically captures PrP^{Sc} and the labeled prion specific antibody detects the captured PrP^{Sc}.

In another aspect, the invention relates to a kit for the detection of PrP^{Sc} comprising: a solid support having bound thereto a peptide, consisting of an amino acid sequence

selected from the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRKRSRKESYSIYV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and
 5 (SEQ ID NO:9) SGRGKGGKGLGKGGAKRHRKVLR; and a labeled prion specific antibody, wherein the peptide specifically captures PrP^{Sc} and the labeled prion specific antibody detects the captured PrP^{Sc}.

In another aspect, the invention relates to a method for detection of PrP^{Sc} comprising: first contacting a sample with a prion specific antibody; and then detecting PrP^{Sc} with a
 10 labeled peptide consisting of an amino acid sequence selected from the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRKRSRKESYSIYV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9)
 15 SGRGKGGKGLGKGGAKRHRKVLR, wherein the prion specific antibody captures PrP^C and PrP^{Sc} and the labeled peptide specifically detects the captured PrP^{Sc}.

In another aspect, the invention relates to an immunoassay for detecting PrP^{Sc} comprising: providing a solid support having bound thereto a prion specific antibody; contacting the solid support with a sample; washing the support to remove any unbound
 20 sample; contacting the solid support with a labeled peptide consisting of an amino acid sequence selected from the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRKRSRKESYSIYV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and
 25 (SEQ ID NO:9) SGRGKGGKGLGKGGAKRHRKVLR; and carrying out a detection step to determine if PrP^{Sc} are bound to the solid support, wherein the prion specific antibody captures PrP^C and PrP^{Sc} and the labeled peptide specifically detects the captured PrP^{Sc}.

In another aspect, the invention relates to a kit for the detection of PrP^{Sc} comprising: a solid support having bound thereto a prion specific antibody; and a labeled peptide consisting of an amino acid sequence selected from the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQQKDGKKRKRSRKESYSIV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9) SGRGKGGKGLGKGGAKRHRKVLR, wherein the prion specific antibody captures PrP^C and PrP^{Sc} and the labeled peptide specifically detects the captured PrP^{Sc}.

10 Brief Description of the Drawings

Figure 1 is an illustration of human plasminogen sequence and Kringle regions.

Figure 2 is an illustration of an IP capture and immunoblot of hamster PrP^{Sc}.

15 Figure 3 is an illustration of an IP capture and immunoblot of BSE PRP^{Sc}.

Figure 4 is an illustration of immunocapture of PRP^{Sc} from brains of vCJD by peptides.

Brain homogenate preparation:

20 Normal and scrapie hamster brain lysate were obtained from Baltimore Research and Education Foundation as 10% whole brain tissue homogenate in PBS (w/v). The lysate was further treated by adding 1/10 volume of 10X detergent homogenate buffer, composed of 5% sodium deoxycolate and 5% Igepal CA-630 (equivalent to NP-40) in PBS, incubating at 4C for 1hr., followed by centrifugation at 1000g for 10 minutes. The resulting supernatant was collected and used in the assay.

Normal and BSE bovine brain tissue were provided by Veterinary Laboratories Agency (VLA), UK. Human vCJD and Lewy body dementia brain tissue were provided by
30 National CJD Surveillance Unit (NCJDSU), UK. Brain

tissue was processed the same way (or similar) as hamster brain homogenate preparation.

Synthetic peptides:

- 5 Ten peptides were either synthesized by ResGen (currently a division of Invitrogen Corporation), or purchased from Upstate Group, Inc. Each peptide was labeled with biotin either on the C-terminal Lysine bearing an aminohexanoyl spacer (K(Lc)), or at the N-terminal through an aminohexanoyl spacer (AMCAP).

10

Peptide ID	Peptide Sequence	Sequence Origin
	Made by	
SEQ ID No.: 1:	WQPPRARIG-K(Lc)-Biotin	Fibronectin
	ResGen	
15	SEQ ID No.: 2	Biotin-AMCAP-NWCKRGRKNCKTH Amyloid protein
	precursor ResGen	
	SEQ ID No.: 3	Biotin-AMCAP-NYKKPKLG Fibroblast growth factor (FGF)-1
	ResGen	
	SEQ ID No.: 4	Biotin-AMCAP-KDFLSIELVRGRVK Laminin A chain
20	ResGen	
	SEQ ID No.: 5	YRGYRGYRGYRG-K(Lc)-Biotin "condensed"
	Kringle-A	
	ResGen	
	SEQ ID No.: 6	YRGRYGYKGKYGYRG-K(Lc)-Biotin "condensed"
25	Kringle-B	
	ResGen	
	SEQ ID No.: 7	AQKKDGKKRKRSRKESYSIYV-GGK(Lc)-Biotin Human histon 2B
	Upstate	
	SEQ ID No.: 8	ARTKQTARKSTGGKAPRKQLA-GGK(Lc)-Biotin Human
30	histon 3	
	Upstate	
	SEQ ID No.: 9	SGRGKGGKGLGKGGAKRHRKVLRLR-GSGSK(Lc)-Biotin
	Human histon 4	

Upstate

SEQ ID No.: 10: **Biotin-AMCAP-TIADRYYRETAR** VP16 protein,
 HSV-2
 ResGen

5

Biotinylated peptide was dissolved in PBS at 1mg/mL and stored at -20°C until use.

Conjugation of biotinylated peptide to streptavidin magnetic beads:

- 10 0.5 mL Dynabeads® M-280 Streptavidin (DynaL Biotech, NY, USA, Cat.# 112.06) were washed twice with PBS and the beads isolated from buffer with the magnet (DynaL Magnetic Particle Concentrator, MPC). 100ug peptide and 1mL PBS was added to the washed beads. Incubation with rotation was performed at 37C for 1-2 hours. The beads were isolated from the buffer with
- 15 the MPC, washed twice with 1 ml PBS (0.1% BSA), and rotated for 5 minutes at room temperature while washing. The peptide -conjugated beads were then blocked for 2 hours, 37°C with 1 mM Biotin in 0.2 M Tris-HCl, pH 8.0, containing 0.1% BSA. The beads were subsequently washed 2 times with 1 ml PBS (0.1% BSA) and once with 1 ml PBS (0.1% BSA, 1% Tween 20)
- 20 incubating each time for 10 minutes at room temp. The beads were then washed once with 1 ml PBS (0.1%BSA) and then stored in 1ml PBS (0.05% sodium azide) at 4°C.

Proteinase K digestion:

- 25 Conditions for the PK digestion of brain lysate: Brain homogenate was suspended in PBS buffer with or without non-ionic detergent. The total homogenate protein concentration was no more than 2.5 mg/mL. PK (Roche Diagnostics, IN, USA, Cat.# 1373196) was added to a final concentration of 50ug/mL. Incubation was at 37C for 0.5 to 1 hour. Digestion was stopped by
- 30 adding Pefabloc SC (Roche Diagnostics, IN, USA, Cat.# 1585916) to a final concentration of 4mM.

Immunoprecipitation (IP), non-reducing electrophoresis and immunoblot detection of PrP^{Sc}:

Peptide conjugated magnetic beads were used to capture PrP^{Sc} from brain homogenate by immunoprecipitation. The IP procedure consists of the following protocol: mix 50uL peptide conjugated beads with PK treated or non-PK treated brain homogenate in a total of 1 mL IP buffer (3% Tween20TM and 3% Igpal CA-630 in PBS) and incubate at 25C for 2.5 hours with rotation → Separate beads using MPC device and wash beads 3 times of 30 second vortexing with IP wash buffer (2% Tween20TM and 2% Igpal CA-630 in PBS) →

10 Elute captured PrP^{Sc} by heating beads with NuPAGE sample buffer for 10-15 minutes. The eluted sample from IP capture were loaded onto a 4-12% NuPAGE® Bis-Tris Gel (Invitrogen, CA, USA, Cat.# NP0302) and subjected to non-reducing electrophoresis at 200V for 45 minutes. The immunoblot procedure was performed as follows: transfer separated proteins in the gel to

15 a 0.2um PVDF membrane (Invitrogen, Cat# LC2002) at 30V for 60 minutes → Block the membrane with BlockerTM Casein in TBS (0.05% Tween20TM) (Pierce Chemical Corp., IL, USA, Cat.# 37532) either at 25C for 1 hour with shaking or at 4C overnight. → As primary antibody, use 3F4 (Signet, MA, USA, Cat.# 9620-02) at 1:3000 dilution or 6H4 (Prionics AG, Switzerland, Cat.# 01-011)

20 at 1:5000 dilution to detect PrP^{Sc}. Incubate the membrane with diluted primary antibody in 10% BlockerTM Casein in TBST buffer (25mM Tri-Cl, 0.2M NaCl, 0.2% Tween20TM, pH 8.0) at 25C for 1 hour with shaking. → Wash 3X 5 minutes with TBST buffer with shaking. → Incubate membrane with horseradish peroxidase conjugated goat polyclonal anti-mouse IgG (H+L)

25 (Jackson ImmunoResearch Laboratories, PA, USA, Cat.# 115-035-003) at 1:10,000 to 1:30,000 dilution in 50% BlockerTM Casein in TBST buffer at 25C for 1 hour with shaking. → Wash 6X 5 minutes with TBST buffer with shaking. → Add ECL chemiluminescence substrate (Amersham Biosciences, NJ, USA, Cat.# RPN2109) or SuperSignal West Dura chemiluminescence substrate

30 (Pierce) on membrane to develop for 5 minutes. → Take image by exposure

either to Bio Max MR-2 film (Kodak, NY, USA) or to the ChemiDocTM Gel Documentation System (Bio-Rad, CA, USA).

Advantages.

5 The present invention uses peptide to capture PrP^{Sc} by recognition of high affinity associated molecules in PrP^{Sc} complex such as glycans, nucleic acid. Because the tight association of these molecules only to PrP^{Sc} but not to PrP^C, the present invention provided a non-intrusive means for the detection of PrP^{Sc} while no PK digestion or other protein modification procedure
10 required. It is anticipated that the mild conditions will preserve the original structure and conformation of the pathogenic prion protein, thereby offering opportunity to determine the presence of true PrP^{Sc} while minimizing the generation of PrP^{Sc} due to harsh sample treatment.

15 The use of synthetic peptides offer advantages in that they display the binding specificity but can also be easily handled in direct coating to a solid phase as well as be conjugated to link to signal given reagents such as horseradish peroxidase (HRP), or to be adopted into other desired diagnosis assay format.

20

Literature cited.

Aguzzi A, Fischer MB, (2001) Prion-Binding Activity in Serum and Plasma., US 20010053533A1

25 Appel TR, Dumpitak C, Matthiesen U, Riesner D. (1999) Prion rods contain an inert polysaccharide scaffold. **Biol Chem** 380(11):1295-306

Arnold, JE, Tipler C, Laszlo L, Hope J, Landon M, Mayer RJ (1995) The abnormal isoform of the prion protein accumulates in late-endosome-like
30 organelles in scrapie-infected mouse brain. **J. Pathol.** 176:403-411

Barnard G, Helmick B, Madden S, Gilbourne C and Patel R (2000) The measurement of prion protein in bovine brain tissue using differential extraction and DELFIA as a diagnostic test for BSE **Luminescence** 2000;15:357-362

5

Belay ED (1999) Transmissible spongiform encephalopathies in humans. **Annu. Rev. Microbiol.** 53:283-314.

Biffiger K, Zwald D, Kaufmann L, Briner A, Nayki I, Purro M, Bottcher S, Struckmeyer T, Schaller O, Meyer R, Fatzer R, Zurbriggen A, Stack M, Moser M, Oesch B, Kubler E. (2002) Validation of a luminescence immunoassay for the detection of PrP(Sc) in brain homogenate. **J Virol Methods** 101(1-2):79-84.

Bolton DC (2001) Prions and proteins: distinguishing between conformations. **The Lancet** 358(9277):164-165

Brown P, Cathala F, Raubertas RF, Gajdusek DC, Castaigne P. (1987) The epidemiology of Creutzfeldt-Jakob disease: conclusion of a 15 year investigation in France and review of the world literature. **Neurology** 37:895-904.

Brown P. The risk of bovine spongiform encephalopathy ("mad cow disease") to human health. **JAMA** 1997; 278: 1008-1011

25

Bruce ME, Will RG, Ironside JW, McConnell I, Drummond D, Suttie A, McCardle L, Chree A, Hope J, Birkett C, Cousens S, Fraser H and Bostock CJ. (1997) Transmissions to mice indicate that 'new variant' CJD is caused by the BSE agent. **Nature** 389:498-501

30

Caughey BW, Dong A, Bhat KS, Ernst D, Hayes SF, Caughey, W.S. (1991) Secondary structure analysis of the scrapie-associated protein PrP27-30 in water by infrared spectroscopy. **Biochemistry** 30:7672-7680.

- 5 Caughey B and Kocisko DA. (2003) A nucleic-acid accomplice? **Nature** Oct 16;425(6959):673-4

Chandler R, (1961) Encephalopathy in mice produced with scrapie brain material. **Lancet** 1:1378-1379

10

Cohen FE, Prusiner SB. (1998) Pathologic conformations of prion proteins. **Annu Rev Biochem** 67:793-819

- 15 Cordeiro Y, Machado F, Juliano L, Juliano MA, Brentani RR, Foguel D, Silva JL. (2001) DNA converts cellular prion protein into the beta-sheet conformation and inhibits prion peptide aggregation. **J Biol Chem.** 276(52):49400-9.

- 20 Cuillé J, Chelle PL (1936) La maladie dite tremblante du mouton est-elle inocuable? **C. R. Acad. Sci.** 203, 1552-1554

Deleault ND, Lucassen RW and Supattapone S. (2003) RNA molecules stimulate prion protein conversion. **Nature** Oct 16;425(6959):717-20

- 25 D'iaz-Nido J., Wandosell F. and Avila J (2002) Glycosaminoglycans and beta-amyloid, prion and tau peptides in neurodegenerative diseases. **J Biol Chem** Jul;23(7):1323-32

- 30 FDA and CBER, U.S. Department of Health and Human Services (2001) IV. RECOMMENDATIONS FOR DONOR DEFERRAL in Guidance for Industry: Revised Preventive Measures to Reduce the Possible Risk of

Transmission of Creutzfeldt-Jakob Disease (CJD) and Variant Creutzfeldt-Jakob Disease (vCJD) by Blood and Blood Products

Ferguson NM, Ghani AC, Donnelly CA, Hagenshaars TJ, Anderson RM.

- 5 (2002) Estimating the human health risk from possible BSE infection of the British sheep flock. **Nature** 415(6870):420-4

Fischer MB, Roeckl C, Parizek P, Schwarz HP, Aguzzi A, (2000) Binding of disease-associated prion protein to plasminogen. **Nature** 408:479-83.

10

Foster JD, Hope J, McConnell I, Bruce M, Fraser H. (1994) Transmission of bovine spongiform encephalopathy to sheep, goats, and mice. **Ann N Y Acad Sci** 724:300-3.

- 15 Gabizon R, McKinley MP, Groth D, Prusiner SB (1988). Immunoaffinity purification and neutralization of scrapie prion infectivity. **Proc. Natl. Acad. Sci. USA** 85, 6617-6621.

- 20 Gajdusek DC, Gibbs CJJ, Alpers MP (1966) Experimental transmission of a kuru-like syndrome to chimpanzees. **Nature** 209:794-796

Gajdusek DC, (1977) Unconventional viruses and the origin and disappearance of kuru. **Science** 197:943-60

- 25 Gibbs CJJ, Gajdusek DC, Asher DM, Alpers MP, Beck E, Daniel PM, Matthews WB (1968) Creutzfeldt-Jakob disease (spongiform encephalopathy): transmission to the chimpanzee. **Science** 161, 388-389

- 30 Gibbs CJ Jr, Gajdusek DC, Amyx H. (1979) Strain variation in the viruses of Creutzfeldt-Jakob disease and kuru. In: Prusiner SB, Hadlow WJ,

editors. Slow transmissible diseases of the nervous system. Volume 2.
New York: Academic Press; p. 87-110.

- 5 Hill AF, Zeidler M, Ironside J, Collinge J (1997) Diagnosis of new variant
Creutzfeldt-Jakob disease by tonsil biopsy. **Lancet** 349(9045):99-100

Hillier CE, Salmon RL, Neal JW, Hilton DA. (2002) Possible
underascertainment of variant Creutzfeldt-Jakob disease: a systematic
study. **J Neurol Neurosurg Psychiatry** 72(3):304-9

10

- Hines KL, Kulkarni AB, McCarthy JB, Tian H, Ward JM, Christ M,
McCartney-Francis NL, Furcht LT, Karlsson S, Wahl SM. (1994) Synthetic
fibronectin peptides interrupt inflammatory cell infiltration in transforming
growth factor beta 1 knockout mice. **Proc Natl Acad Sci U S A** May
 15 24;91(11):5187-91.

Holada K, Simak J, Vostal JG (2000) Transmission of BSE by blood
transfusion. **Lancet** 356(9243):1772

- 20 Hope J, Morton LJD, Farquhar CF, Multhaup G, Beyreuther K, Kimberlin
RH (1986). The major polypeptide of scrapie-associated fibrils (SAF) has
the same size, charge distribution and N-terminal protein sequence as
predicted for the normal brain protein (PrP). **EMBO J.** 5, 2591-2597.

- 25 Horiuchi M, Caughey B, (1999) Specific binding of normal prion protein to
the scrapie form via a localized domain initiates its conversion to the
protease-resistant state. **EMBO J** 18(12):3193-203.

- Horwich AL, Weissman JS (1997) Deadly Conformations—Protein
 30 Misfolding in Prion Disease. **Cell**, 89:499-510

Houston F (2000) Transmission of BSE by blood transfusion in sheep.

Lancet 356(9234):999-1000

Hunter N, Foster J, Chong A, McCutcheon S, Parnham D, Eaton S,

5 MacKenzie C, Houston F. (2002) Transmission of prion diseases by blood transfusion. **J Gen Virol** Nov;83(Pt 11):2897-905

Jackson GS, Hosszu LL, Power A, Hill AF, Kenney J, Saibil H, Craven CJ,

10 Waltho JP, Clarke AR, Collinge J. (1999) Reversible conversion of monomeric human prion protein between native and fibrillogenic conformations. **Science** 283(5409):1935-7.

Kellings K, Meyer N, Mirenda C, Prusiner SB, Riesner D. (1992). Further

15 analysis of nucleic acids in purified scrapie prion preparations by improved return refocusing gel electrophoresis. **J. Gen. Virol.** 73, 1025-1029.

Kimberlin RH, Cole S, Walker CA. (1987) Temporary and permanent

modifications to a single strain of mouse scrapie on transmission to rats and hamsters. **J Gen Virol** 68:1875-81.

20

Kimberlin RH, Walker CA, Fraser H. (1989) The genomic identity of

different strains of mouse scrapie is expressed in hamsters and preserved on reisolation in mice. **J Gen Virol** 70:2017-25.

25 Klein TR, Kirsch D, Kaufmann R. and Riesner D (1998) **Biol. Chem.** 379:655-666

Korth C, Stierli B, Streit P, Moser M, Schaller O, Fischer R, Schulz-

Schaeffer W, Kretzschmar H, Raeber A, Braun U, Ehrensperger F,

30 Hornemann S, Glockshuber R, Riek R, Billeter M, Wuthrich K, Oesch B.

(1997) Prion (PrP^{Sc})-specific epitope defined by a monoclonal antibody.
Nature 390(6655):74-7

Lasmézas CI, Deslys JP, Demaimay R, Adjou KT, Lamoury F, et al. (1996)
 5 BSE transmission to macaques. **Nature** 381:743-44

Luo Y, Gabriel JL, Wang F, Zhan X, Maciag T, Kan M, McKeehan WL.
(1996) Molecular modeling and deletion mutagenesis implicate the nuclear
translocation sequence in structural integrity of fibroblast growth factor-1. **J**
 10 **Biol Chem.** Oct 25;271(43):26876-83.

Maissen M, Roeckl C, Markus G, Goldman W, Aguzzi A. (2001)
Plasminogen binds to disease-associated prion protein of multiple species.
Lancet 357:2026-8.

15 Manuelidis EE, Kim JH, Mericangas JR, Manuelidis L. (1985) Transmission
to animals of Creutzfeldt-Jakob disease from human blood. **Lancet** 2:896-
97

20 McBride PA, Wilson MI, eikelenboom P, Tunstall A, Bruce ME (1998)
Heparan Sulfate Proteoglycan is Associated with Amyloid Plaques and
Neuroanatomically Targeted PrP Pathology throughout the Incubation
Period of Scrapie-Infected Mice. **Experimental Neurol.** 149:447-454

25 McGowan JP. (1922) Scrapie in sheep. **Scott. J. Agric.** 5:365-75

Meyer RK, McKinley MP, Bowman KA, Braunfeld MB, Barry RA, Prusiner
SB (1986). Separation and properties of cellular and scrapie prion proteins.
Proc. Natl. Acad. Sci. USA 83:2310-2314.

30

- Mizuno K, Inoue H, Hagiya M, Shimizu S, Nose T, Shimohigashi Y, Nakamura T. (1994) Hairpin loop and second kringle domain are essential sites for heparin binding and biological activity of hepatocyte growth factor. **J Biol Chem.** Jan 14;269(2):1131-6. **Plasminogen**
- 5
- Moynagh J, Schimmel H, (1999) The evaluation of tests for the diagnosis of Transmissible Spongiform Encephalopathy in Bovines (8 July 1999) <http://europa.eu.int/comm/food/fs/bse/bse12-en.html>.
- 10 Narang HK (2002) A critical review of the nature of the spongiform encephalopathy agent: prion theory versus virus theory. **Exp Biol. Med.** (Maywood) 227(1):4-19
- O'Rourke KI, Baszler TV, Besser TE, Miller JM, Cutlip RC, Wells GA, et al. (2000) Preclinical diagnosis of scrapie by immunohistochemistry of third eyelid lymphoid tissue. **J Clin Microbiol** 38:3254-9.
- 15
- Pan KM, Baldwin M, Nguyen J, Gasset M, Serban A, Groth D, Mehlhorn I, Huang Z, Fletterick RJ, Cohen FE, Prusiner SB. (1993) Conversion of -helices into -sheets features in the formation of the scrapie prion proteins. **Proc. Natl. Acad. Sci. USA** 90:10962-10966
- 20
- Paramithiotis E et al (2003) A prion protein epitope selective for the pathologically misfolded conformation **Nat Med** Jul;9(7):893-9
- 25
- Parizek P, Roeckl C, Weber J, Flechsig E, Aguzzi A, Raeber AJ. (2001) Similar turnover and shedding of the cellular prion protein in primary lymphoid and neuronal cells.
- 30 Pattison I (1957) Transmission of scrapie to the goat. **Lancet** 272:104-105.

Prusiner SB, Novel proteinaceous infectious particles cause scrapie.
Science (1982) 216:136-44

5 Raymond GJ et al. (2000) Evidence of a molecular barrier limiting
susceptibility of humans, cattle and sheep to chronic wasting disease.
EMBO J Sept 1; 19(17): 4425-30

10 Safar J, Wille H, Itri V, Groth D, Serban H, Torchia M, Cohen FE, Prusiner
SB. (1998) Eight prion strains have PrP(Sc) molecules with different
conformations. Nat. Med. 4(10):1157-65

15 Schaller O, Fatzer R, Stack M, Clark J, Cooley W, Biffiger K, Egli S, Doherr
M, Vandeveld M, Heim D, Oesch B, Moser M. (1999) Validation of a
western immunoblotting procedure for bovine PrP(Sc) detection and its use
as a rapid surveillance method for the diagnosis of bovine spongiform
encephalopathy (BSE). Acta Neuropathol (Berl) 98(5):437-43.

20 Shaked GM, Meiner Z, Avraham I, Taraboulos A, Gabizon R (2001a)
Reconstitution of Prion Infectivity from Solubilized Protease-resistant PrP
and Nonprotein Components of Prion Rods. J. Biol. Chem.
276(17):14324-14328

25 Shaked GM, Shaked Y, Kariv-Inbal Z, Halimi M, Avraham I, Gabizon R,
(2001b) A Protease-resistant Prion Protein Isoform Is Present in Urine of
Animals and Humans Affected with Prion Diseases J. Biol. Chem.,
276(34):31479-82

30 Shyng SL, Heuser JE, Harris DA. (1994) Aglycolipid-anchored prion protein
is endocytosed via clathrin-coated pits. J. Cell Biol. 125:1239-50

Small DH, Nurcombe V, Reed G, Clarris H, Moir R, Beyreuther K, Masters CL. (1994) A heparin-binding domain in the amyloid protein precursor of Alzheimer's disease is involved in the regulation of neurite outgrowth. **J Neurosci.** Apr;14(4):2117-27

5

Snow AD (1990) Immunolocalization of heparan sulfate proteoglycans to the prion protein amyloid plaques of Gerstmann-Straussler syndrome, Creutzfeldt-Jakob disease and scrapie. **Lab Invest.** 63(5):601-1

- 10 Soeda S, Ohki H, Shimeno H, Nagamatsu A (1989) Further characterization of the binding of plasminogen to heparin: evidence for the involvement of lysine residues. **Biochim Biophys Acta.** Nov 9;999(1):29-35. **Plasminogen**

- 15 Swietnicki W, Morillas M, Chen SG, Gambetti P, Surewicz WK. (2000) Aggregation and fibrillization of the recombinant human prion protein huPrP90-231. **Biochemistry** 39(2):424-31.

- 20 Taraboulos A, Serban D, Prusiner SB. (1990) Scrapie prion proteins accumulate in the cytoplasm of persistently infected cultured cells. **J. Cell. Biol.** 110:2117-32

Tateishi J. (1985) Transmission of Creutzfeldt-Jakob disease from human blood and urine into mice. **Lancet** 2:1074

25

Telling GC, Scott M, Mastrianni J, Gabizon R., Torchia M, Cohen FE, DeArmond SJ, Prusiner SB (1995). Prion propagation in mice expressing human and chimeric PrP transgenes implicates the interaction of cellular PrP with another protein. **Cell** 83:79-90.

30

- Valleron AJ, Boelle PY, Will R, Cesbron JY. (2001) Estimation of epidemic size and incubation time based on age characteristics of vCJD in the United Kingdom. **Science** 294(5547):1726-8
- 5 Wadsworth JD, Joiner S, et al. (2001) Tissue distribution of protease resistant prion protein in variant Creutzfeldt-Jakob disease using a highly sensitive immunoblotting assay. **Lancet** 358(9277):171-80
- Warner RG, Hundt C, Weiss S, Turnbull JE (2002) Identification of the
 10 heparan sulfate binding sites in the cellular prion protein. **J Biol Chem.** May 24;277(21):18421-30
- Wells GAH, Scott AC, Johnson CT, Gunning RF, Hancock RD, et al. (1987) A novel progressive spongiform encephalopathy in cattle. **Vet. Rec.**
 15 31:419-20
- Williams ES, Young S. (1980) Chronic wasting disease of captive mule deer: a spongiform encephalopathy. **J. Wildl. Dis.** 16:89-98
- 20 Wong C, Xiong LW, Horiuchi M, Raymond L, Wehrly K, Chesebro B, Caughey B (2001) Sulfated glycans and elevated temperature stimulate PrPSc-dependent cell-free formation of protease-resistant prion protein **EMBO J** 20(3):377-386
- 25 Woods A, McCarthy JB, Furcht LT, Couchman JR. (1993) A synthetic peptide from the COOH-terminal heparin-binding domain of fibronectin promotes focal adhesion formation. **Mol Biol Cell.** Jun;4(6):605-13
- Yoshida I, Tashiro K, Monji A, Nagata I, Hayashi Y, Mitsuyama Y, Tashiro N.
 30 (1999) Identification of a heparin binding site and the biological activities of

the laminin alpha1 chain carboxy-terminal globular domain. **J Cell Physiol.**
Apr;179(1):18-28.

SEQUENCE LISTING

<110> Ortho-Clinical Diagnostics, Inc.

5 <120> Peptides for Discrimination of Prions

<130> 1011-4943CA

<141> 2005-09-30

10

<150> U.S. 60/614,533

<151> 2004-09-30

<160> 10

15

<170> PatentIn version 3.1

<210> 1

<211> 9

20 <212> PRT

<213> peptide

<400> 1

25 Trp Gln Pro Pro Arg Ala Arg Ile Gly

1 5

<210> 2

30 <211> 13

<212> PRT

<213> peptide

<400> 2

5 Asn Trp Cys Lys Arg Gly Arg Lys Asn Cys Lys Thr His

1 5 10

<210> 3

<211> 8

10 <212> PRT

<213> peptide

<400> 3

15 Asn Tyr Lys Lys Pro Lys Leu Gly

1 5

<210> 4

<211> 14

20 <212> PRT

<213> peptide

<400> 4

25 Lys Asp Phe Leu Ser Ile Glu Leu Val Arg Gly Arg Val Lys

1 5 10

<210> 5

<211> 12

30 <212> PRT

<213> peptide

<400> 5

Tyr Arg Gly Tyr Arg Gly Tyr Arg Gly Tyr Arg Gly
5 1 5 10

<210> 6

<211> 15

<212> PRT

10 <213> peptide

<400> 6

Tyr Arg Gly Arg Tyr Gly Tyr Lys Gly Lys Tyr Gly Tyr Arg Gly
15 1 5 10 15

<210> 7

<211> 21

<212> PRT

20 <213> peptide

<400> 7

Ala Gln Lys Lys Asp Gly Lys Lys Arg Lys Arg Ser Arg Lys Glu Ser
25 1 5 10 15

Tyr Ser Ile Tyr Val
20

30 <210> 8
<211> 21

<212> PRT

<213> peptide

<400> 8

5

Ala Arg Thr Lys Gln Thr Ala Arg Lys Ser Thr Gly Gly Lys Ala Pro

1 5 10 15

Arg Lys Gln Leu Ala

10 20

<210> 9

<211> 23

<212> PRT

15 <213> peptide

<400> 9

Ser Gly Arg Gly Lys Gly Gly Lys Gly Leu Gly Lys Gly Gly Ala Lys

20 1 5 10 15

Arg His Arg Lys Val Leu Arg

20

25 <210> 10

<211> 12

<212> PRT

<213> peptide

30 <400> 10

Thr Ile Ala Asp Arg Tyr Tyr Arg Glu Thr Ala Arg
1 5 10

We claim:

1. A method for detection of PrP^{Sc} comprising:
 first contacting a sample with a peptide consisting of an amino acid
 5 sequence selected from the group consisting of: (SEQ ID NO:1)
 WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3)
 NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6)
 YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRSRKESYSIYV,
 (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9)
 10 SGRGKGGKGLGKGGAKRHRKVLR; and
 then detecting PrP^{Sc} with a labeled prion specific antibody, wherein the
 peptide specifically captures PrP^{Sc} and the labeled prion specific antibody
 detects the captured PrP^{Sc}.

- 15 2. An immunoassay for detecting PrP^{Sc} comprising:
 providing a solid support having bound thereto a peptide, consisting of an
 amino acid sequence selected from the group consisting of: (SEQ ID
 NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3)
 NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6)
 20 YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRSRKESYSIYV,
 (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9)
 SGRGKGGKGLGKGGAKRHRKVLR;
 contacting the solid support with a sample;
 washing the support to remove any unbound sample;
 25 contacting the solid support with a labeled prion specific antibody; and
 carrying out a detection step to determine if PrP^{Sc} are bound to the solid
 support, wherein the peptide specifically captures PrP^{Sc} and the labeled
 prion specific antibody detects the captured PrP^{Sc}.

- 30 3. A kit for the detection of PrP^{Sc} comprising:

a solid support having bound thereto a peptide, consisting of an amino acid sequence selected from the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRSRKESYSIYV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9) SGRGKGGKGLGKGGAKRHRKVLR; and
a labeled prion specific antibody, wherein the peptide specifically captures PrP^{Sc} and the labeled prion specific antibody detects the captured PrP^{Sc}.

4. Method for detection of PrP^{Sc} comprising:

first contacting a sample with a prion specific antibody; and
then detecting PrP^{Sc} with a labeled peptide consisting of an amino acid sequence selected from the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRSRKESYSIYV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9) SGRGKGGKGLGKGGAKRHRKVLR, wherein the prion specific antibody captures PrP^C and PrP^{Sc} and the labeled peptide specifically detects the captured PrP^{Sc}.

5. An immunoassay for detecting PrP^{Sc} comprising:

providing a solid support having bound thereto a prion specific antibody;
contacting the solid support with a sample;
washing the support to remove any unbound sample;
contacting the solid support with a labeled peptide consisting of an amino acid sequence selected from the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRSRKESYSIYV,

(SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9)
SGRGKGGKGLGKGGAKRHRKVLRL; and

carrying out a detection step to determine if PrP^{Sc} are bound to the solid support, wherein the prion specific antibody captures PrP^{C} and PrP^{Sc} and the labeled peptide specifically detects the captured PrP^{Sc} .

6. A kit for the detection of PrP^{Sc} comprising:
a solid support having bound thereto a prion specific antibody; and
a labeled peptide consisting of an amino acid sequence selected from the group consisting of: (SEQ ID NO:1) WQPPRARI; (SEQ ID NO:2) NWCKRGRKQCKTH; (SEQ ID NO:3) NYKKPKL; (SEQ ID NO:5) YRGYRGYRGYRG; (SEQ ID NO:6) YRGRYGYKGKYGYRG; (SEQ ID NO:7) AQKKDGKKRKRSRKESYSIYV, (SEQ ID NO:8) ARTKQTARKSTGGKAPRKQLA; and (SEQ ID NO:9) SGRGKGGKGLGKGGAKRHRKVLRL, wherein the prion specific antibody captures PrP^{C} and PrP^{Sc} and the labeled peptide specifically detects the captured PrP^{Sc} .

Application number / Numéro de demande: 2518372

Figures: _____

Pages: 2/4, 3/4, 4/4

Unscannable items received with this application
(Request original documents in File Prep. Section on the 10th floor)

Documents reçu avec cette demande ne pouvant être balayés
(Commander les documents originaux dans la section de la préparation
des dossiers au 10ième étage)

FIGURE-1 Human Plasminogen Sequence and Kringle regions

- ▲ There are 4 YRG sequence in 4 out of 5 Kringle regions. ■ There are 6 YK, RY, RK sequences
- There are lots of Y, R, or K around several Kringle loop disulphide bridges.

