

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2019391097 B2**

(54) Title
CDK9 inhibitors and polymorphs thereof for use as agents for treatment of cancer

(51) International Patent Classification(s)
A61K 31/675 (2006.01) **A61K 47/12** (2006.01)
A61K 47/02 (2006.01) **A61K 47/26** (2006.01)

(21) Application No: **2019391097** (22) Date of Filing: **2019.12.04**

(87) WIPO No: **WO20/117988**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/775,303	2018.12.04	US

(43) Publication Date: **2020.06.11**

(44) Accepted Journal Date: **2025.07.03**

(71) Applicant(s)
Sumitomo Pharma Oncology, Inc.

(72) Inventor(s)
SIDDIQUI-JAIN, Adam;FLYNN, Paul;FUJIWARA, Yuji;MASUMOTO, Shuji;TANAKA, Hiroaki;KUREBAYASHI, Hirotaka;HASHIZUKA, Takahiko;ARIKAWA, Yuka

(74) Agent / Attorney
FPA Patent Attorneys Pty Ltd, Level 19, South Tower 80 Collins Street, Melbourne, VIC, 3000, AU

(56) Related Art
WO 2016/187316 A1
Balbach, S. et al., "Pharmaceutical evaluation of early development candidates "the 100 mg-approach", International Journal of Pharmaceutics. 2004, vol. 275, pages 1-12
Singhal, D. et al., "Drug polymorphism and dosage form design: a practical perspective", Advanced Drug Delivery Reviews. 2004, vol. 56, pages 335-347
Caira, M. R., "Crystalline Polymorphism of Organic Compounds", Design of organic solids, Weber, E. et al "ED", Springer, 1998, pages 164-208
Bastin, R. J. et al., "Salt Selection and Optimisation Procedures for Pharmaceutical New Chemical Entities", Organic Process Research & Development. 2000, vol 4, 427-435

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
11 June 2020 (11.06.2020)



(10) International Publication Number
WO 2020/117988 A1

(51) International Patent Classification:

A61K 31/675 (2006.01) A61K 47/12 (2006.01)
A61K 47/02 (2006.01) A61K 47/26 (2006.01)

(21) International Application Number:

PCT/US2019/064549

(22) International Filing Date:

04 December 2019 (04.12.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/775,303 04 December 2018 (04.12.2018) US

(71) Applicant: **TOLERO PHARMACEUTICALS, INC.**
[US/US]; 3900 N Traverse Mountain Blvd, Lehi, UT 84043
(US).

(72) Inventors: **SIDDIQUI-JAIN, Adam**; 3900 N Traverse
Mountain Blvd, Lehi, UT 84043 (US). **FLYNN, Paul**; 3900
N Traverse Mountain Blvd, Lehi, UT 84043 (US). **FU-
JIWARA, Yuji**; 3900 N Traverse Mountain Blvd, Lehi, UT
84043 (US). **MASUMOTO, Shuji**; c/o Sumitomo Dainip-

pon Pharma Company, Ltd., 6-8, Dosho-Machi 2-Chome,
Chuo-ku, Osaka-shi, Osaka 541-0045 (JP). **TANAKA, Hi-
roaki**; c/o Sumitomo Dainippon Pharma Company, Ltd.,
6-8, Dosho-Machi 2-Chome, Chuo-ku, Osaka-shi, Osa-
ka 541-0045 (JP). **KUREBAYASHI, Hirotaka**; c/o Su-
mitomo Dainippon Pharma Company, Ltd., 6-8, Dosho-
Machi 2-Chome, Chuo-ku, Osaka-shi, Osaka 541-0045
(JP). **HASHIZUKA, Takahiko**; c/o Sumitomo Dainip-
pon Pharma Company, Ltd., 6-8, Dosho-Machi 2-Chome,
Chuo-ku, Osaka-shi, Osaka 541-0045 (JP). **ARIKAWA,
Yuka**; c/o Sumitomo Dainippon Pharma Company, Ltd.,
6-8, Dosho-Machi 2-Chome, Chuo-ku, Osaka-shi, Osaka
541-0045 (JP).

(74) Agent: **HALLER, Rachel, A.** et al.; Seed Intellectual Prop-
erty Law Group LLP, Suite 5400, 701 Fifth Avenue, Seat-
tle, WA 98104-7064 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,

(54) Title: CDK9 INHIBITORS AND POLYMORPHS THEREOF FOR USE AS AGENTS FOR TREATMENT OF CANCER





FIG. 1

(57) Abstract: A crystalline form and/or polymorph of a compound having the following structure (I), including tautomeric and zwitterionic forms thereof, are provided. Methods associated with preparation and use of the polymorphs, and pharmaceutical compositions comprising the same are also provided. Also provided are methods for preparing a compound having formula (I), or a salt, tautomer or zwitterionic form thereof.

[Continued on next page]



WO 2020/117988 A1

KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*
- *of inventorship (Rule 4.17(iv))*

Published:

- *with international search report (Art. 21(3))*

CDK9 INHIBITORS AND POLYMORPHS THEREOF FOR USE AS AGENTS FOR
TREATMENT OF CANCER

RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 62/775,303, filed on December 4, 2018. The entire teachings of the above application is incorporated herein by reference.

BACKGROUND


[0002] Alvocidib (flavopiridol) is a potent cyclin-dependent kinase (CDK) inhibitor with selectivity for CDKs 9, 1, 2, 4 and 7. 2-(2-Chlorophenyl)-5-hydroxy-8-((3S,4R)-3-hydroxy-1-methylpiperidin-4-yl)-4-oxo-4H-chromen-7-yl dihydrogen phosphate, disclosed in International Publication No. WO 2016/187316, is an orally bioavailable prodrug of alvocidib.

[0003] There is a need for crystalline forms and/or polymorphs of 2-(2-chlorophenyl)-5-hydroxy-8-((3S,4R)-3-hydroxy-1-methylpiperidin-4-yl)-4-oxo-4H-chromen-7-yl dihydrogen phosphate that are suitable for use in pharmaceutical compositions and methods for treating diseases associated with overexpression of a CDK. There is a further need for methods for preparing 2-(2-chlorophenyl)-5-hydroxy-8-((3S,4R)-3-hydroxy-1-methylpiperidin-4-yl)-4-oxo-4H-chromen-7-yl dihydrogen phosphate that enable the manufacture of 2-(2-chlorophenyl)-5-hydroxy-8-((3S,4R)-3-hydroxy-1-methylpiperidin-4-yl)-4-oxo-4H-chromen-7-yl dihydrogen phosphate for use in pharmaceutical compositions and methods for treating diseases associated with overexpression of a CDK.

BRIEF SUMMARY

[0004] Embodiments of the present invention are generally directed to CDK (*e.g.*, CDK9) inhibitors and polymorphs thereof as well as pharmaceutical compositions comprising the same, for use as therapeutic or prophylactic agents, for example, for treatment of cancer (*e.g.*, hematological cancer) and other conditions.

[0005] Some embodiments provide a crystalline form of a compound having the structure (I):



or a tautomer or zwitterionic form thereof.

[0006] More specifically, aspects of the present disclosure are directed to a polymorph of a compound having the structure (I), or a tautomer or zwitterionic form thereof. For example, in some embodiments the polymorph has an X-ray powder diffraction pattern comprising the following: D space (Å): 18.3 ± 0.09 , 8.1 ± 0.06 , 6.4 ± 0.08 , 5.9 ± 0.06 , 4.4 ± 0.05 expressed in terms of “D” spacing.

[0007] Another embodiment affords a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof, the polymorph being a crystallized form having a monoclinic space group $P2_1$ with lattice parameters of $a = 6.46(1) \text{ \AA}$, $b = 9.07(2) \text{ \AA}$, $c = 18.25(4) \text{ \AA}$, $\beta = 95.457(2)^\circ$, and a volume of $1066.11(4) \text{ \AA}^3$.

[0008] One embodiment provides a crystalline form of a compound of structure (I), or a tautomer or zwitterionic form thereof, having an x-ray powder diffraction pattern substantially in accordance with that depicted in FIG. 1.

[0009] Yet another embodiment provides a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof, wherein the polymorph is Form B as described herein.

[0010] Still another embodiment affords a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof, wherein the polymorph is formed by a method comprising (a) contacting an amorphous compound of structure (I) with a lattice forming reagent; and (b) treating the product of step (a) with solvent having water content less than about 0.05% v/v and removing the solvent, thereby forming the polymorph.

[0011] One embodiment provides a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof, wherein the polymorph has an initial purity of at


least 99.5% and a subsequent purity of at least 99.5% after being stored from about 12 hours up to about 7 days at about $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at a relative humidity of 60%.

[0012] One embodiment provides a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof having an endotherm peak value at about $256^{\circ}\text{C} - 268^{\circ}\text{C}$ as determined by differential scanning calorimetry.

[0013] Pharmaceutical compositions comprising a compound having structure (I) (*e.g.*, a polymorph or crystalline form of a compound having structure (I)), or a tautomer or zwitterionic form thereof, and a pharmaceutically acceptable carrier or excipient are also provided in various other embodiments. One embodiment provides a pharmaceutical composition comprising from about 0.5 weight percent to about 11.25 weight percent of a compound having the structure (I), or a tautomer or zwitterionic form thereof, and from about 85 weight percent to about 99.5 weight percent of a pharmaceutically acceptable carrier or excipient. One embodiment provides a pharmaceutical composition comprising about 0.6 weight percent of a compound having structure (I), or a tautomer or zwitterionic form thereof; about 97.4 weight percent of anhydrous lactose; about 1 weight percent colloidal silicon dioxide; and about 1 weight percent magnesium stearate. Another embodiment provides a pharmaceutical composition comprising about 11 weight percent of a compound having the structure (I), or a tautomer or zwitterionic form thereof; about 87 weight percent of anhydrous lactose; about 1 weight percent colloidal silicon dioxide; and about 1 weight percent magnesium stearate. Yet another embodiment provides a pharmaceutical composition comprising about 0.6 weight percent of a compound having structure (I), or a tautomer or zwitterionic form thereof; and from about 98 weight percent to about 99.5 weight percent of cornstarch. Another embodiment provides a pharmaceutical composition comprising about 11 weight percent of a compound of the structure (I), or a tautomer or zwitterionic form thereof; and about 88 weight percent of cornstarch.


[0014] Other embodiments provide methods including a method for treatment of a disease associated with overexpression of a cyclin-dependent kinase (CDK), such as cancer (*e.g.*, a hematologic cancer), in a mammal in need thereof, for example, by administering to the mammal a therapeutically effective amount of a crystalline form or a polymorph of a compound having structure (I), or a tautomer or zwitterionic form thereof.

[0015] In other embodiments, the present disclosure provides a method for preparing a polymorph or crystalline form of the compound of structure (I). One embodiment provides a method for preparing crystalline Form B of a compound having structure (II):




(II)

the method comprising contacting amorphous compound having the structure (I), or a tautomer or zwitterionic form thereof, with an acid in a solvent, thereby preparing the crystalline Form B of the compound having structure (II). Another embodiment provides a method for preparing crystalline Form B of the compound having structure (II), the method comprising contacting a compound having structure (V):



(V)

or a tautomer, salt or zwitterionic form thereof, with an acid in a solvent, thereby preparing the crystalline Form B of the compound having structure (II). Yet another embodiment provides a method for preparing crystalline Form B of a compound having structure (II), the method comprising contacting a compound having the following structure (IV):



(IV)

or a tautomer or salt thereof, with di-*tert*-butylhalophosphonate (*e.g.*, di-*tert*-butylchlorophosphonate, di-*tert*-butylbromophosphonate) in the presence of an amine base, thereby forming a compound having structure (V), or a tautomer or salt thereof; and contacting the compound having structure (V), or a tautomer or salt thereof, in an organic solvent, with an acid having a pK_a value of greater than about 1 (*e.g.*, an acid having at least one pK_a value that is less than about 5 and a pK_a value of greater than about 1), thereby preparing crystalline Form B of a compound having structure (II).

Another embodiment is a method for preparing crystalline Form B of a compound having structure (II), the method comprising contacting a compound having structure (IV), or a tautomer or salt thereof, with di-*tert*-butylhalophosphonate (*e.g.*, di-*tert*-butylchlorophosphonate, di-*tert*-butylbromophosphonate) in the presence of an amine base, thereby forming a compound having structure (V), or a tautomer or salt thereof; contacting the compound having structure (V), or a tautomer or salt thereof, with an acid having a pK_a value of less than about 1, thereby forming a salt of a compound having structure (I); contacting the salt of a compound having structure (I) with a base, thereby forming amorphous compound having structure (I); and contacting the amorphous form of the compound having structure (I), in an organic solvent, with an acid having a pK_a value of greater than about 1 (*e.g.*, an acid having at least one pK_a value that is less than about 5 and a pK_a value of greater than about 1), thereby preparing crystalline Form B of the compound having structure (II).

[0016] The present disclosure also provides methods for preparing a compound having structure (I), or a salt, tautomer or zwitterionic form thereof. The method comprises contacting a compound having structure (IV), or a tautomer or salt thereof, with di-*tert*-butylhalophosphonate (*e.g.*, di-*tert*-butylchlorophosphonate, di-*tert*-butylbromophosphonate) in the presence of an amine base, thereby forming a compound having structure (V), or a tautomer or salt thereof; and contacting the

compound having structure (V), or a tautomer or salt thereof, with an acid, thereby preparing a compound having structure (I), or a salt, tautomer or zwitterionic form thereof.

[0017] These and other aspects of the disclosure will be apparent upon reference to the following detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018] In the figures, identical reference numbers identify similar elements. The sizes and relative positions of elements in the figures are not necessarily drawn to scale and some of these elements are enlarged and positioned to improve figure legibility. Further, the particular shapes of the elements as drawn are not intended to convey any information regarding the actual shape of the particular elements, and have been solely selected for ease of recognition in the figures.

[0019] FIG. 1 illustrates an x-ray diffractogram obtained from XRPD analysis for polymorph Form B.

[0020] FIG. 2 shows a stability profile in the form of a TG-FTIR thermogram and a DSC spectrum for the amorphous form of structure (I) (*i.e.*, Form A).

[0021] FIG. 3 illustrates a comparison of lattice forming reagents and the similarity of diffractogram patterns as determined by XRPD for polymorph Form B.

[0022] FIG. 4 depicts the Oak Ridge Thermal Ellipsoid Plot (ORTEP) diagrams of single crystal structure obtained for polymorph Form B.

[0023] FIG. 5 shows diffractograms from XRPD analysis of three samples of polymorph Form C.

[0024] FIG. 6A-6B show TG-FTIR thermograms of polymorph Form C before (FIG. 6A) and after drying (FIG. 6B).

[0025] FIG. 7 shows a comparison of ¹H NMR spectra acquired for forms A (amorphous), form B (polymorph) and form C.

[0026] FIG. 8 shows the differential scanning calorimetry output of heat flow plotted as a function of temperature for polymorph form B.

[0027] FIG. 9A is a graph of percentage of drug released versus time, and shows the amount of API released from various formulations of API and filler in the excipient compatibility study described in Example 13.

[0028] FIG. 9B is a graph of percentage of drug released versus time, and shows the amount of API released from various formulations of API and lubricant in the excipient compatibility study described in Example 13.

[0029] FIG. 9C is a graph of percentage of drug released versus time, and shows the amount of API released from various formulations of API and disintegrant in the excipient compatibility study described in Example 13.

[0030] FIG. 9D is a graph of percentage of drug released versus time, and shows the amount of API released from various formulations of API and fluidizer/other excipient in the excipient compatibility study described in Example 13.

[0031] FIG. 10A is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-01 of Example 14 stored at 5 °C.

[0032] FIG. 10B is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-01 of Example 14 stored at 25 °C and 60% relative humidity (RH).

[0033] FIG. 10C is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-01 of Example 14 stored at 40 °C and 75% RH.

[0034] FIG. 10D is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-02 of Example 14 stored at 5 °C.

[0035] FIG. 10E is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-02 of Example 14 stored at 25 °C and 60% RH.

[0036] FIG. 10F is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-02 of Example 14 stored at 40 °C and 75% RH.

[0037] FIG. 10G is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-03 of Example 14 stored at 5 °C.

[0038] FIG. 10H is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-03 of Example 14 stored at 25 °C and 60% RH.

[0039] FIG. 10I is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-03 of Example 14 stored at 40 °C and 75% RH.

[0040] FIG. 10J is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-04 of Example 14 stored at 5 °C.

[0041] FIG. 10K is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-04 of Example 14 stored at 25 °C and 60% RH.

[0042] FIG. 10L is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-04 of Example 14 stored at 40 °C and 75% RH.

[0043] FIG. 10M is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-05 of Example 14 stored at 5 °C.

[0044] FIG. 10N is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-05 of Example 14 stored at 25 °C and 60% RH.

[0045] FIG. 10O is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-05 of Example 14 stored at 40 °C and 75% RH.

[0046] FIG. 10P is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-06 of Example 14 stored at 5 °C.

[0047] FIG. 10Q is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-06 of Example 14 stored at 25 °C and 60% RH.

[0048] FIG. 10R is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-06 of Example 14 stored at 40 °C and 75% RH.

[0049] FIG. 10S is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-08 of Example 14 stored at 5 °C.

[0050] FIG. 10T is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-08 of Example 14 stored at 25 °C and 60% RH.

[0051] FIG. 10U is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-08 of Example 14 stored at 40 °C and 75% RH.

[0052] FIG. 10V is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-09 of Example 14 stored at 5 °C.

[0053] FIG. 10W is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-09 of Example 14 stored at 25 °C and 60% RH.

[0054] FIG. 10X is a graph of dissolution (%) versus time, and shows the amount of compound of structure (I) released from Formulation No. 401-09 of Example 14 stored at 40 °C and 75% RH.

[0055] FIG. 11A is a graph of plasma alvocidib concentration (ng/mL) versus time, and shows the concentration of alvocidib in the plasma of patients in cohort 1 on day 1 following daily oral QD dosing with a 1-mg strength capsule containing Formulation No. 401-01.

[0056] FIG. 11B is a graph of plasma alvocidib concentration (ng/mL) versus time, and shows the concentration of alvocidib in the plasma of patients in cohort 1 on day 14 following daily oral QD dosing with a 1-mg strength capsule containing Formulation No. 401-01.

[0057] FIG. 11C is a graph of plasma alvocidib concentration (ng/mL) versus time, and shows the concentration of alvocidib in the plasma of patients in cohort 2 on day 1

following daily oral BID dosing with a 1-mg strength capsule containing Formulation No. 401-01.

[0058] FIG. 11D is a graph of plasma alvocidib concentration (ng/mL) versus time, and shows the concentration of alvocidib in the plasma of patients in cohort 2 on day 14 following daily oral BID dosing with a 1-mg strength capsule containing Formulation No. 401-01.

[0059] FIG. 11E is a graph of plasma alvocidib concentration (ng/mL) versus time, and shows the concentration of alvocidib in the plasma of patients in Cohort 5 on day 1 following daily oral BID dosing with 6 mg of Formulation No. 401-01.

[0060] FIG. 11F is a graph of plasma alvocidib concentration (ng/mL) versus time, and shows the concentration of alvocidib in the plasma of patients in Cohort 5 on day 14 following daily oral BID dosing with 6 mg of Formulation No. 401-01.

[0061] FIG. 11G is a graph of alvocidib (ng/mL) versus cohort, and shows the average C_{max} of alvocidib on day 1 and day 14 following daily oral QD dosing with a 1-mg strength capsule containing Formulation No. 401-01.

[0062] FIG. 11H is a graph of alvocidib (ng*hr/mL) versus cohort, and shows the area under the curve (AUC) of alvocidib on day 1 (AUC₀₋₈) and day 14 (AUC₀₋₈ and AUC₀₋₂₄) following daily oral BID dosing with a 1-mg strength capsule containing Formulation No. 401-01.

[0063] FIG. 11I is a graph of mean concentration of alvocidib (nM) versus time, and shows the mean concentration of alvocidib in plasma of Cohort 5 patients over a 24-hour period.

[0064] FIG. 12 is a graph, and shows the completed cycles on the study described in Example 15 through Cohort 5.

DETAILED DESCRIPTION

[0065] In the following description, certain specific details are set forth in order to provide a thorough understanding of various embodiments of the disclosure. However, one skilled in the art will understand that the disclosure may be practiced without these details.

[0066] Unless the context requires otherwise, throughout the present specification and claims, the word "comprise" and variations thereof, such as, "comprises" and

“comprising” are to be construed in an open, inclusive sense (*i.e.*, as “including, but not limited to”).

[0067] Reference throughout this specification to “one embodiment” or “an embodiment” means that a particular feature, structure or characteristic described in connection with the embodiment is included in at least one embodiment of the present invention. Thus, the appearances of the phrases “in one embodiment” or “in an embodiment” in various places throughout this specification are not necessarily all referring to the same embodiment. Furthermore, the particular features, structures, or characteristics may be combined in any suitable manner in one or more embodiments.

[0068] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. As used in the specification and claims, the singular form “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise.

[0069] “Crystalline,” as used herein, refers to a homogeneous solid formed by a repeating, three-dimensional pattern of atoms, ions or molecules having fixed distances between constituent parts. The unit cell is the simplest repeating unit in this pattern. Notwithstanding the homogenous nature of an ideal crystal, a perfect crystal rarely, if ever, exists. “Crystalline,” as used herein, encompasses crystalline forms that include crystalline defects, for example, crystalline defects commonly formed by manipulating (*e.g.*, preparing, purifying) the crystalline forms described herein. A person skilled in the art is capable of determining whether a sample of a compound is crystalline notwithstanding the presence of such defects.

[0070] “Polymorph,” as used herein, refers to a crystalline form of a compound characterized by a distinct arrangement of its molecules in a crystal lattice. Polymorphs can be characterized by analytical methods such as x-ray powder diffraction (XRPD), differential scanning calorimetry (DSC) and thermogravimetric analysis.

[0071] The crystalline forms and/or polymorphs described herein can be substantially pure. As used herein, “substantially pure,” used without further qualification, means the indicated compound has a purity greater than 90 weight percent, for example, greater than 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99 weight percent, and also including a purity equal to about 100 weight percent, based on the weight of the compound. The remaining material comprises other form(s) of the

compound, and/or reaction impurities and/or processing impurities arising from its preparation (*e.g.*, alvocidib). Purity can be assessed using techniques known in the art, for example, using an HPLC assay described herein. “Substantially pure” can also be qualified as in “substantially pure of other physical forms of the compound having structure (I), or a tautomer or zwitterionic form thereof” or “substantially pure of alvocidib.” When qualified thus, “substantially pure” means that the indicated compound contains less than 10%, preferably less than 5%, more preferably less than 3%, most preferably, less than 1% by weight of the indicated impurity (*e.g.*, any other physical forms of an indicated crystalline form of a compound; alvocidib).

[0072] As used herein, the term “alvocidib” means 2-(2-chlorophenyl)-5,7-dihydroxy-8-[(3S,4R)-3-hydroxy-1-methylpiperidin-4-yl]chromen-4-one, or a salt (*e.g.*, a pharmaceutically acceptable salt) thereof (*e.g.*, 2-(2-chlorophenyl)-5,7-dihydroxy-8-[(3S,4R)-3-hydroxy-1-methylpiperidin-4-yl]chromen-4-one hydrochloride).

[0073] An XRPD pattern or DSC thermogram that is “substantially in accordance” with one or more figures herein showing an XRPD pattern or diffractogram or DSC thermogram, respectively, is one that would be considered by one skilled in the art to represent the same single crystalline form of the compound having structure (I), or a tautomer or zwitterionic form thereof, as the sample of the compound having structure (I), or a tautomer or zwitterionic form thereof, that provided the pattern or diffractogram or thermogram of one or more figures provided herein. Thus, an XRPD pattern or DSC thermogram that is substantially in accordance may be identical to that of one of the figures or, more likely, may be somewhat different from one or more of the figures. For example, an XRPD pattern that is somewhat different from one or more of the figures may not necessarily show each of the lines of the diffraction pattern presented herein and/or may show a slight change in appearance or intensity of the lines or a shift in the position of the lines. These differences typically result from differences in the conditions involved in obtaining the data or differences in the purity of the sample used to obtain the data. A person skilled in the art is capable of determining if a sample of a crystalline compound is of the same form as or a different form from a form disclosed herein by comparison of the XRPD pattern or DSC thermogram of the sample and the corresponding XRPD pattern or DSC thermogram disclosed herein.

[0074] The crystalline forms provided herein can also be identified on the basis of differential scanning calorimetry (DSC) and/or thermogravimetric analysis (TGA). DSC is a thermoanalytical technique in which the difference in the amount of heat required to increase the temperature of a sample is measured as a function of temperature. DSC can be used to detect physical transformations, such as phase transitions, of a sample. For example, DSC can be used to detect the temperature(s) at which a sample undergoes crystallization, melting or glass transition. It is to be understood that any temperature associated with DSC specified herein, with the exception of the DSC temperatures in the Figures or Examples, means the specified value ± 5 °C or less. For example, when an embodiment or a claim specifies an endothermic peak at 264 °C, this is to be understood to mean 264 °C ± 5 °C or less, that is a temperature of from 259 °C to 269 °C. In preferred embodiments, a DSC is the specified value ± 3 °C or less, in more preferred embodiments, ± 2 °C or less.

[0075] “Lattice forming reagent” or “salt former” refers to a chemical substance that is used in combination with another compound to facilitate the formation of a crystalline or polymorphic solid. For example, in some embodiments, a lattice forming reagent facilitates the formation of polymorph Form B of the compound of structure (I). Lattice forming reagents include acids, bases, sugars, peptides, and the like. In some embodiments, the lattice forming reagent is an acid (*e.g.*, an acid having at least one pK_a value that is less than about 5, such as about 4). For example, a lattice forming reagent may be an organic acid (*e.g.*, maleic acid, acetic acid, fumaric acid, tartaric acid). In some embodiments, the acid is a diprotic acid.

[0076] The term “effective amount” or “therapeutically effective amount” refers to that amount of a compound (*e.g.*, polymorph Form B of a compound of structure (I)) described herein that is sufficient to effect the intended application including disease treatment, as defined below. The therapeutically effective amount may vary depending upon the intended treatment application (*in vivo*), or the subject and disease condition being treated, *e.g.*, the weight and age of the subject, the severity of the disease condition, the manner of administration and the like, which can readily be determined by one of ordinary skill in the art. The term also applies to a dose that will induce a particular response in target cells, *e.g.*, reduction of platelet adhesion and/or cell migration. The specific dose will vary depending on the particular compound chosen,

the dosing regimen to be followed, whether it is administered in combination with other compounds, timing of administration, the tissue to which it is administered, and the physical delivery system in which it is carried.

[0077] As used herein, “treatment” or “treating” refers to an approach for obtaining beneficial or desired results with respect to a disease, disorder or medical condition including a therapeutic benefit and/or a prophylactic benefit. By therapeutic benefit is meant eradication or amelioration of the underlying disorder being treated. Also, a therapeutic benefit is achieved with the eradication or amelioration of one or more of the physiological symptoms associated with the underlying disorder such that an improvement is observed in the subject, notwithstanding that the subject may still be afflicted with the underlying disorder. In certain embodiments, for prophylactic benefit, the compositions are administered to a subject at risk of developing a particular disease, or to a subject reporting one or more of the physiological symptoms of a disease, even though a diagnosis of this disease may not have been made.

[0078] A “therapeutic effect,” as that term is used herein, encompasses a therapeutic benefit and/or a prophylactic benefit as described above. A prophylactic effect includes delaying or eliminating the appearance of a disease or condition, delaying or eliminating the onset of symptoms of a disease or condition, slowing, halting, or reversing the progression of a disease or condition, or any combination thereof.

[0079] The terms “co-administration,” “administered in combination with,” and their grammatical equivalents, as used herein, encompass administration of two or more agents to a subject, such as an animal, including humans, to treat a disease, disorder or condition described herein. In some embodiments, administration of the two or more agents is such that both agents and/or their metabolites are present in the subject at the same time. Co-administration includes simultaneous administration in separate compositions, administration at different times in separate compositions, or administration in a composition in which the two or more agents are present.

[0080] An “anti-cancer agent,” “anti-tumor agent” or “chemotherapeutic agent” refers to any agent useful in the treatment of a neoplastic condition. One class of anti-cancer agents comprises chemotherapeutic agents. “Chemotherapy” means the administration of one or more chemotherapeutic drugs and/or other agents to a cancer

patient by various methods, including intravenous, oral, intramuscular, intraperitoneal, intravesical, subcutaneous, transdermal, buccal, or inhalation or in the form of a suppository.

[0081] “Subject” refers to an animal, such as a mammal, for example, a human. The methods described herein can be useful in both human therapeutics and veterinary applications. In some embodiments, the subject is a mammal, and in some embodiments, the subject is human.

[0082] “Mammal” includes humans and both domestic animals such as laboratory animals and household pets (*e.g.*, cats, dogs, swine, cattle, sheep, goats, horses, rabbits), and non-domestic animals, such as wildlife and the like. In some embodiments, the mammal is a human.

[0083] “Radiation therapy” means exposing a subject, using routine methods and compositions known to the practitioner, to radiation emitters such as alpha-particle emitting radionuclides (*e.g.*, actinium and thorium radionuclides), low linear energy transfer (LET) radiation emitters (*i.e.*, beta emitters), conversion electron emitters (*e.g.*, strontium-89 and samarium-153-EDTMP, or high-energy radiation, including without limitation x-rays, gamma rays, and neutrons.

[0084] The term “*in vivo*” refers to an event that takes place in a subject’s body.

[0085] Embodiments of the invention disclosed herein are also meant to encompass crystalline forms and/or polymorphs of a compound of structure (I), or a tautomer or zwitterionic form thereof, being isotopically-labelled by having one or more atoms replaced by an atom having a different atomic mass or mass number (*i.e.*, an “isotopic form” of the crystalline forms and/or polymorphs of a compound of structure (I), or a tautomer or zwitterionic form thereof). Examples of isotopes that can be incorporated into the disclosed crystalline forms and/or polymorphs, or a tautomer or zwitterionic form thereof, include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, chlorine, and iodine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I , and ^{125}I , respectively. These radiolabeled compounds could be useful to help determine or measure the effectiveness of the compounds, by characterizing, for example, the site or mode of action, or binding affinity to pharmacologically important site of action. Certain isotopically-labeled crystalline forms and/or polymorphs of compounds of structure (I), or a tautomer or zwitterionic

form thereof, for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium (*i.e.*, ^3H), and carbon-14 (*i.e.*, ^{14}C) are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

[0086] Substitution with heavier isotopes such as deuterium (*i.e.*, ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence are preferred in some circumstances.

[0087] Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O and ^{13}N , can be useful in positron emission topography (PET) studies for examining substrate receptor occupancy. Isotopically-labeled crystalline forms and/or polymorphs of compounds of structure (I), or a tautomer or zwitterionic form thereof, can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the Examples as set out below using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

[0088] It is understood that in the present description, combinations of substituents and/or variables of the depicted formulae are permissible only if such contributions result in stable compounds. “Stable compound” and “stable structure” are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

[0089] A “pharmaceutical composition” refers to a formulation of a compound of the disclosure and a medium generally accepted in the art for the delivery of the biologically active compound to mammals, *e.g.*, humans. Such a medium includes all pharmaceutically acceptable carriers, diluents or excipients therefor.

[0090] “Pharmaceutically acceptable carrier or excipient” includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals.

[0091] “Pharmaceutically acceptable salt” includes both acid and base addition salts.

[0092] “Pharmaceutically acceptable acid addition salt” refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, *p*-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

[0093] “Pharmaceutically acceptable base addition salt” refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine,

procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, piperazine, piperidine, *N*-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

[0094] A “tautomer” refers to a proton shift from one atom of a molecule to another atom of the same molecule. Embodiments thus include tautomers of the disclosed compounds, crystalline forms and polymorphs.


[0095] “Zwitterionic form” refers to a form of a compound, wherein at least one functional group has a positive electrical charge, at least one functional group has a negative electrical charge, and the net charge of the entire molecule is zero. For example, the phosphate group ($-\text{PO}_3\text{H}_2$) of a compound having structure (I) may exist in an anionic form (*e.g.*, $-\text{PO}_3\text{H}^-$), and the nitrogen atom of a compound having structure (I) may exist in the protonated (cationic) form. The compound having structure (II) is a zwitterionic form of the compound having structure (I), for example. Embodiments include zwitterions of the disclosed compounds, crystalline forms and polymorphs.

[0096] The chemical naming protocol and structure diagrams used herein are a modified form of the I.U.P.A.C. nomenclature system, using the ACD/Name Version 9.07 software program and/or ChemDraw Ultra Version 11.0.1 software naming program (CambridgeSoft). For complex chemical names employed herein, a substituent group is typically named before the group to which it attaches. For example, cyclopropylethyl comprises an ethyl backbone with a cyclopropyl substituent. Except as described below, all bonds are identified in the chemical structure diagrams herein, except for all bonds on some carbon atoms, which are assumed to be bonded to sufficient hydrogen atoms to complete the valency.

Crystalline and Polymorph Forms of Compounds of Structure (I)

[0097] It has been found that compounds having structure (I), or a tautomer or zwitterionic form thereof, can exist in various crystalline and/or polymorphic forms.


[0098] Accordingly, one embodiment provides a crystalline form of a compound having the following structure (I):



(I)

or a tautomer or zwitterionic form thereof. In some embodiments, the crystalline form comprises Form B. In some embodiments, the crystalline form consists essentially of Form B. In some embodiments, the crystalline form consists of Form B. In some embodiments, the crystalline form (*e.g.*, Form B) is substantially pure (*e.g.*, of other physical forms of the compound having structure (I), or a tautomer or zwitterionic form thereof; of impurities; of alvocidib). In some embodiments, the crystalline form is of a compound having structure (II).

[0099] Form B has structure (II):




(II)

and is characterized, in some embodiments, by an x-ray powder diffraction (XRPD) pattern comprising at least three peaks (*e.g.*, three peaks, at least four peaks, four peaks, at least five peaks, five peaks, six peaks) at 2-theta angles selected from the group consisting of $4.8 \pm 0.2^\circ$, $10.8 \pm 0.2^\circ$, $13.7 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$, $20.0 \pm 0.2^\circ$ and $24.6 \pm 0.2^\circ$. In some embodiments, Form B is characterized by an XRPD pattern comprising peaks at the following 2-theta angles: $10.8 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$ and $20.0 \pm 0.2^\circ$. In some embodiments, Form B is characterized by an XRPD pattern comprising peaks at the following 2-theta angles: $4.8 \pm 0.2^\circ$, $10.8 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$ and $20.0 \pm 0.2^\circ$. In some embodiments, Form B is characterized by an XRPD pattern comprising peaks at the following 2-theta angles: $4.8 \pm 0.2^\circ$, $10.8 \pm 0.2^\circ$, $13.7 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$ and $20.0 \pm 0.2^\circ$. In some embodiments, Form B has an XRPD pattern substantially in accordance

with that depicted in FIG. 1. In some embodiments, Form B is characterized by a DSC thermogram comprising an endothermic peak at about 264 °C. In some embodiments, Form B is characterized by a DSC thermogram substantially in accordance with that depicted in FIG. 8.

[00100] One embodiment provides a polymorph of a compound having the following structure (I):



(I)

or a tautomer or zwitterionic form thereof, the polymorph having an X-ray powder diffraction pattern comprising the following:

D space (Å):

18.3 ± 0.09

8.1 ± 0.06

6.4 ± 0.08

5.9 ± 0.06

4.4 ± 0.05

expressed in terms of "D" spacing.

[00101] In a related embodiment, the polymorph has an X-ray powder diffraction pattern comprising the following:

D space (Å):

18.38 ± 0.003

8.15 ± 0.008

6.47 ± 0.002

5.95 ± 0.007

4.44 ± 0.006

expressed in terms of "D" spacing.

[00102] In yet another related embodiment, the polymorph has an X-ray powder diffraction pattern comprising the following:

D space (Å):

18.382

8.157

6.472

5.956

4.445

expressed in terms of "D" spacing.

[00103] One embodiment provides a polymorph of a compound of structure (I), or a tautomer or zwitterion thereof, wherein the polymorph is a crystalized form having a monoclinic space group $P2_1$ with lattice parameters of:

$a = 6.46(1) \text{ \AA}$;

$b = 9.07(2) \text{ \AA}$;

$c = 18.25(4) \text{ \AA}$; and

$\beta = 95.457(2)^\circ$;

and a volume of $1066.11(4) \text{ \AA}^3$.

[00104] Another embodiment provides a polymorph of a compound of structure (I), or a tautomer or zwitterion thereof, wherein the polymorph is a crystalized form having a monoclinic space group $P2_1$ with lattice parameters of:

$a = 6.4695(1) \text{ \AA}$;

$b = 9.0692(2) \text{ \AA}$;

$c = 18.2530(4) \text{ \AA}$; and

$\beta = 95.457(2)^\circ$,

and a volume of $1066.11(4) \text{ \AA}^3$.

[00105] Yet another embodiment provides a polymorph of structure (I) or a tautomer or zwitterionic form thereof, wherein the polymorph has the X-ray diffraction pattern set forth in FIG. 1 (or substantially similar). In some embodiments, the polymorph has the X-ray diffraction pattern substantially similar to Table 6.

[00106] Another related embodiment provides a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof, wherein the polymorph is Form B.

[00107] One embodiment provides a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof wherein the polymorph is formed by a method comprising:

a) contacting an amorphous compound of structure (I) with a lattice forming reagent; and

b) treating the product of step a) with solvent having water content less than about 0.05 % v/v and removing the solvent, thereby forming the polymorph.

[00108] In some more specific embodiments, the lattice forming reagent is maleic acid. In some embodiments, the solvent is ethanol or tetrahydrofuran. In some other embodiments, the contacting further comprises suspending the amorphous compound of structure (I) in aqueous tetrahydrofuran.

[00109] Another embodiment affords a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof, wherein the polymorph has an initial purity of at least 99.5% and a subsequent purity of at least 99.5% after being stored from about 12 hours up to about 7 days above a temperature of about 22°C.


[00110] In certain related embodiments, the subsequent purity is at least 99.5% after being stored greater than about 7 days above a temperature of about 22°C. In some more specific embodiments, the subsequent purity is at least 99.5% after being stored greater than about 30 days above a temperature of about 20°C. In some embodiments, the initial purity and subsequent purity are as determined by HPLC.

[00111] Without wishing to be bound by theory, it is thought that the water content of the polymorph can have a significant effect on the purity and storage stability of the polymorph. That is, the polymorph can undergo a hydrolysis reaction that converts that phosphate moiety to a hydroxyl group. As such, an impurity may be present in the form of hydrolyzed structure (I) (*i.e.*, alvocidib). However, it was unexpectedly discovered that keeping the water content of the polymorph and any subsequently formed compositions provided an active substance (*i.e.*, the polymorph of structure (I)) with much more robust stability.

[00112] Accordingly, some embodiments provide a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof, wherein the polymorph has an initial purity of at least 99.5% and a subsequent purity of at least 99.5% after being stored from about 12 hours up to about 7 days above a temperature at about $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at a relative humidity of 60%. In some embodiments, the subsequent purity is at least 99.5% after being stored for greater than about 7 days at about $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at a relative humidity of 60%. In other embodiments, the subsequent purity is at least 99.5% after being stored for greater than about 30 days at about $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at a relative humidity of 60%. In some of the foregoing embodiments, the initial purity and subsequent purity are as determined by HPLC.

[00113] Still other embodiments provide a polymorph of a compound of structure (I) or a tautomer or zwitterionic form thereof, wherein the polymorph has an endotherm peak value at about $256.0^{\circ}\text{C} - 268.0$ (*i.e.*, $262.0^{\circ}\text{C} \pm 6.0$) as determined by differential scanning calorimetry. In some embodiments, the endotherm peak value is at a temperature ranging from 256°C to 268°C , from 257.0°C to 266.0°C or from 258.0°C to 265.0°C . In some more specific embodiments, the endotherm peak value is at about $259.0^{\circ}\text{C} - 265.0^{\circ}\text{C}$.

[00114] In some embodiments of the foregoing, the polymorph comprises a zwitterionic form having the following structure (II):



(II)

[00115] In some of the foregoing embodiments, the polymorph has water content less than 0.50 % as determined by Karl Fischer titration. For example in some embodiments, the polymorph has water content less than 0.45 %, less than 0.40 %, less than 0.35 %, less than 0.30 %, less than 0.25 %, less than 0.20 %, less than 0.15 %, or less than 0.10 % as determined by Karl Fischer titration.


Preparation of Compounds Having Structure (I)

[00116] It will be appreciated by those skilled in the art that the processes and reactions for preparing the compounds described herein may be modified in accordance with standard techniques to include alternative reagents and/or reaction conditions. For example, a reaction including an acid as lattice forming reagent may alternatively employ another suitable reagent such as a base, a sugar derivative, a peptide or other reagent.

[00117] One embodiment provides a method for preparing a polymorph, the method comprising:

a) contacting an amorphous compound having the following structure

(I):



(I)

with a lattice forming reagent; and

b) treating the product of step a) with solvent having water content less than about 0.05 % v/v; and removing the solvent, thereby forming the polymorph.

[00118] In some embodiments, the lattice forming reagent is an acid. For example, in some embodiments, the acid is an organic acid. In some embodiments the acid is an acid having a pK_a greater than 1.0. In some embodiments the acid is a diprotic acid. In some specific embodiments, the acid is selected from the group consisting of maleic acid, fumaric acid, L-tartaric acid, hippuric acid, nicotinic acid, acetic acid, and combinations thereof. In some more specific embodiments, the lattice forming reagent is maleic acid.

[00119] In some specific embodiments, the contacting comprises adding the lattice forming reagent (*e.g.*, maleic acid) in a 1:1 ± 0.5 ratio (*e.g.*, molar ratio) of the amorphous compound of structure (I) to maleic acid. In more specific embodiments the ratio (*e.g.*, molar ratio) is 1:1 ± 0.3, 1:1 ± 0.2, or 1:1 ± 0.1 compound of structure (I) to


maleic acid. In other embodiments, the ratio (*e.g.*, molar ratio) is from about 1:0.1 to about 1:10 (*e.g.*, from about 1:0.1 to about 1:5) compound of structure (I) to maleic acid.

[00120] The lattice forming reagent is not limited to acids and can be any other compound that can be mixed with a compound of structure (I) and forms polymorph Form B. In particular, in some embodiments, the lattice forming reagent is a sugar or a peptide. In more specific embodiments, the lattice forming reagent is D-xylose. In other specific embodiments, the lattice forming reagent is aspartame.

[00121] The removing step is performed to remove residual solvent and lattice forming reagent and afford a solid product as a free base. Accordingly, in some embodiments the removing may involve reduced pressure (*e.g.*, below 1 atm), a flow of air or inter gas (*e.g.*, N₂), increased temperature (*e.g.*, above 25°C) or combinations thereof. In specific embodiments, the removing is at a pressure less than about 1 atm and a temperature greater than about 35°C. In some more specific embodiments, the pressure is less than about 0.9 atm, about 0.8 atm, about 0.7 atm, about 0.6 atm, or about 0.5 atm. In some embodiments, the temperature is above about 36 °C, about 37 °C, about 38 °C, about 39 °C, about 40 °C, about 42 °C, or about 45 °C. In some embodiments, the temperature is above the boiling point of residual solvent, *e.g.*, about 100 °C. Each of the temperature and pressure values recited above include a range of ± 5% of the recited value (*i.e.*, about 45°C includes 45°C ± 2.25°C).


[00122] In some embodiments, the removing comprises filtering the solid product away from residual solvent and/or lattice forming reagent.

[00123] Additionally, some embodiments of the methods recited above further comprise preparing a compound of structure (III):




(III)

by reacting a compound having the following structure:




or a salt thereof, with a base and a compound having the following structure:



[00124] In some of the foregoing embodiments, the base is an amine base, (*e.g.*, triethylamine, diisopropylethyl amine). In some of the foregoing embodiments, amine base is added in an organic solvent.

[00125] In some embodiments, the method further comprises preparing the amorphous compound of structure (I):



(I)

by reacting a compound of structure (III), or a salt thereof, with a phosphoester cleaving compound and treating with a basic solution.


[00126] In some embodiments, the phosphoester cleaving compound is an organosilicon (*e.g.*, trimethyl silyl bromide, "TMSBr"). In some embodiments, the basic solution comprises ammonium (*e.g.*, ammonium bicarbonate). In some embodiments, the basic solution comprises acetonitrile. In some embodiments, the basic solution comprises aqueous acetonitrile.

[00127] In some embodiments, the polymorph is a polymorph as described in the foregoing embodiments. Additionally, one embodiment provides a polymorph that is prepared according to any of the methods described herein.

[00128] Thus, in any of the embodiments described herein involving polymorph conversion/formation (*e.g.*, preparation of crystalline Form B of a compound having structure (II)), the polymorph conversion/formation step(s) comprise crystallizing compound having structure (I) (*e.g.*, amorphous compound having structure (I)), or a tautomer or zwitterionic form thereof, from an acidic solution (*e.g.*, having a pH of greater than about 1; having a pH of greater than about 1 to less than about 5, for example, from about 1 to about 4, from about 2 to about 5). In some embodiments, the acidic solution comprises a lattice forming reagent (*e.g.*, an acid, such as maleic acid; an acid having a pK_a of greater than about 1; an acid having at least one pK_a of less than about 5 and a pK_a of greater than about 1, for example, at least one pK_a of less than about 4 and a pK_a of greater than about 1, at least one pK_a of less than about 5 and a pK_a of greater than about 2) and a solvent.

[00129] One embodiment is a method for preparing crystalline Form B of a compound having structure (II), the method comprising contacting amorphous compound having structure (I), or a tautomer or zwitterionic form thereof, with a lattice forming reagent (*e.g.*, an acid, such as maleic acid; an acid having a pK_a of greater than about 1; an acid having at least one pK_a of less than about 5 and a pK_a of greater than about 1, for example, an acid having at least one pK_a of less than about 4 and a pK_a of greater than about 1, an acid having at least one pK_a of less than about 5 and a pK_a of greater than about 2) in a solvent, thereby preparing the crystalline Form B of the compound having structure (II).

[00130] One embodiment is a method for preparing crystalline Form B of a compound having structure (II), the method comprising contacting a compound having structure (V):



or a tautomer, salt or zwitterionic form thereof, with an acid in a solvent, thereby preparing the crystalline Form B of the compound having structure (II). In some


embodiments, the molar ratio of acid to compound having structure (V), or a tautomer, salt or zwitterionic form thereof, is from about 0.1:1 to about 10:1 (*e.g.*, from about 0.1:1 to about 5:1; from about 0.4:1 to about 1.1:1; about 0.5:1).

[00131] In one embodiment, the lattice forming reagent is an acid. In some embodiments, the acid for preparing crystalline Form B has at least one pK_a value that is less than about 5 (*e.g.*, less than about 4) and/or a pK_a value of greater than about 1 (*e.g.*, greater than about 2).

[00132] In some embodiments for preparing crystalline Form B, the acid is an organic acid. As used herein, “organic acid” is an organic compound with acidic properties. Typically, an organic acid contains at least one carboxyl group (-COOH). Organic acids include, but are not limited to, maleic acid, acetic acid, citric acid and propionic acid. In some embodiments, the organic acid is maleic acid.

[00133] As used herein, “solvent” refers to a liquid that serves as a medium for a chemical reaction or other procedure in which compounds are being manipulated (*e.g.*, crystallization). Typically, the solvent in the methods disclosed herein is an organic solvent or water, or a combination thereof. Examples of organic solvents include polar, protic solvents (*e.g.*, an alcohol such as methanol, ethanol, butanol), polar aprotic solvents (*e.g.*, dimethylformamide, tetrahydrofuran, ethyl acetate, acetone, methyl ethyl ketone) or nonpolar solvents (*e.g.*, diethyl ether). In some embodiments, the solvent comprises tetrahydrofuran. In some embodiments, the solvent is a mixture of methanol and acetone.

[00134] Also provided are methods for preparing a compound having structure (I), or a salt, tautomer or zwitterionic form thereof (*e.g.*, a compound having structure (II)). The method comprises contacting a compound having the following structure (IV):



(IV)

or a tautomer or salt thereof, with di-*tert*-butylhalophosphonate (*e.g.*, di-*tert*-butylchlorophosphonate, di-*tert*-butylbromophosphonate) in the presence of an amine

base, thereby forming a compound having structure (V), or a tautomer or salt thereof, and contacting the compound having structure (V), or a tautomer or salt thereof, with an acid (*e.g.*, maleic acid), thereby preparing a compound having structure (I), or a salt, tautomer or zwitterionic form thereof. In some embodiments, the method further comprises contacting di-*tert*-butylphosphonate with carbon tetrahalide (*e.g.*, carbon tetrachloride, carbon tetrabromide) to prepare the di-*tert*-butylhalophosphonate (*e.g.*, di-*tert*-butylchlorophosphonate, di-*tert*-butylbromophosphonate, respectively). In some embodiments, the di-*tert*-butylhalophosphonate is prepared *in situ* with the compound having structure (IV), or a salt thereof, by contacting di-*tert*-butylphosphonate with carbon tetrahalide (*e.g.*, carbon tetrachloride, carbon tetrabromide). In some embodiments, the acid is hydrochloric acid or acetic acid, or a combination thereof.

[00135] “Amine base,” as used herein, refers to a compound that contains a nitrogen atom with a lone pair. “Amine base” includes primary, secondary and tertiary amine bases, as well as ammonia. Typically, the amine base is an organic amine base. Examples of organic amine bases include, but are not limited to, diisopropylethylamine, *tert*-butylamine and triethylamine.

[00136] It will be appreciated that when an acid having a pK_a of less than about the pK_a of the phosphate group (-PO₂OH) of a compound having structure (I) (*e.g.*, hydrochloric acid) is used in the methods described herein, a salt (*e.g.*, a hydrochloride salt) of the compound having structure (I) will result. Thus, in some embodiments, the compound formed by contacting a compound having structure (V), or a tautomer or salt thereof, with an acid is a salt of a compound having structure (I). In such instances, in some embodiments, the method further comprises contacting the salt of the compound having structure (I) with a base (*e.g.*, ammonium bicarbonate), thereby forming the compound having structure (I), or a tautomer or zwitterionic form thereof.

[00137] One embodiment is a method for preparing crystalline Form B of a compound having structure (II). The method comprises contacting a compound having structure (IV), or a tautomer or salt thereof, with di-*tert*-butylhalophosphonate (*e.g.*, di-*tert*-butylchlorophosphonate, di-*tert*-butylbromophosphonate) in the presence of an amine base, thereby forming a compound having structure (V), or a tautomer or salt thereof; and contacting the compound having structure (V), or a tautomer or salt thereof, in an organic solvent, with an acid having a pK_a value of greater than about 1


(*e.g.*, an acid having at least one pK_a value that is less than about 5 and a pK_a value of greater than about 1, for example, at least one pK_a value that is less than about 4 and a pK_a value of greater than about 1, at least one pK_a value that is less than about 5 and a pK_a value of greater than about 2), thereby preparing crystalline Form B of a compound having structure (II).

[00138] Another embodiment is a method for preparing crystalline Form B of a compound having structure (II), comprising contacting a compound having structure (IV), or a tautomer or salt thereof, with di-*tert*-butylhalophosphonate (*e.g.*, di-*tert*-butylchlorophosphonate, di-*tert*-butylbromophosphonate) in the presence of an amine base, thereby forming a compound having structure (V), or a tautomer or salt thereof; and contacting the compound having structure (V), or a tautomer or salt thereof (*e.g.*, in an organic solvent) with an acid having a pK_a value of less than about 1, thereby forming a salt of a compound having structure (I); contacting the salt of a compound having structure (I) with a base, thereby forming amorphous compound having structure (I); and contacting the amorphous compound having structure (I) with an acid having a pK_a value of greater than about 1 (*e.g.*, an acid having at least one pK_a value that is less than about 5 and a pK_a value of greater than about 1, for example, at least one pK_a value that is less than about 4 and a pK_a value of greater than about 1, at least one pK_a value that is less than about 5 and a pK_a value of greater than about 2), thereby preparing crystalline Form B of a compound having structure (II).

[00139] It will also be appreciated by those skilled in the art that in the processes for preparing the compounds described herein, the functional groups of intermediate compounds may need to be protected by suitable protecting groups. Such functional groups include hydroxy, amino, mercapto and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl or diarylalkylsilyl (for example, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl, and the like. Suitable protecting groups for amino, amidino and guanidino include *t*-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for mercapto include -C(O)-R" (where R" is alkyl, aryl or arylalkyl), *p*-methoxybenzyl, trityl and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters. Protecting groups are optionally added or removed in accordance with standard techniques, which are known to one skilled in the art and described herein.

The use of protecting groups is described in detail in Green, T.W. and P.G.M. Wutz, *Protective Groups in Organic Synthesis* (1999), 3rd Ed., Wiley. As one of skill in the art would appreciate, the protecting group may also be a polymer resin such as a Wang resin, Rink resin or a 2-chlorotrityl-chloride resin.

[00140] The following General Reaction Scheme 1 illustrates an exemplary method of forming a polymorph of a compound of structure (I):




(I)

[00141] It is understood that one skilled in the art may be able to make these compounds by similar methods or by combining other methods known to one skilled in the art. It is also understood that one skilled in the art would be able to make, in a similar manner as described below, a compound of structure (I), or a tautomer, salt or zwitterionic form thereof, not specifically illustrated below by using the appropriate starting components and modifying the parameters of the synthesis as needed (*e.g.*, reagents, solvents, reaction times and temperatures, etc.). In general, starting components may be obtained from sources such as Sigma Aldrich, Lancaster Synthesis, Inc., Maybridge, Matrix Scientific, TCI, and Fluorochem USA, etc. or synthesized according to sources known to those skilled in the art (*see*, for example, *Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 5th edition (Wiley, December 2000)) or prepared as described in this invention.

[00142] Polymorphs and compounds were analyzed using techniques known in the art, for example, by x-ray powder diffraction (XRPD), dynamic vapor sorption (DVS), thermal gravimetric analysis (TGA), differential scanning calorimetry (DSC), mass spectrometry, high performance (or pressure) liquid chromatography (HPLC), ¹³C NMR, ³¹P NMR and/or ¹H NMR. Synthetic procedures are described in more detail below.

General Reaction Scheme 1




[00143] Embodiments of the polymorph of a compound of structure (I) as shown can be prepared according to General Reaction Scheme 1. Certain intermediates useful for preparation of the compound having structure (I), or a tautomer, salt or zwitterionic form thereof (*e.g.*, A-1, A-2, A-3, A-4, A-10, A-11) can be prepared according to methods described in U.S. Pat. No. 9,758,539 and U.S. Pub. No. 2007/0015802, which are incorporated herein by reference in their entireties. As shown in the General Reaction Scheme 1, A-1 can be purchased from commercial sources or prepared according to methods familiar to one of ordinary skill in the art. Reaction of A-1 with a strong Lewis acid (*e.g.*, boron tribromide) affords A-2, which can then be phosphorylated with an activated protected phosphate reagent (*e.g.*, diethyl chlorophosphate) under basic conditions (*e.g.*, using diisopropyl ethyl amine as a base) in an appropriate solvent (*e.g.*, NMP). The phosphorylated product (A-3) can then be deprotected under standard conditions (*e.g.*, TMS-Br) and neutralized using suitable conditions (*e.g.*, ammonium carbonate in aqueous acetonitrile). The resultant compound A-4 (*i.e.*, an amorphous solid or "Form A") which is then converted to the polymorph A-5 using an appropriate lattice forming reagent (*e.g.*, maleic acid) and a suitable solvent system (*e.g.*, THF and water at 19:1 v/v). Finally, the lattice forming

reagent is removed using a suitable, substantially anhydrous solvent (*e.g.*, an alcohol such as ethanol) and dried to afford polymorph Form B.

[00144] It should be noted that the General Reaction Scheme 1 only depicts an exemplary method for preparation of the polymorph A-5 and other methods are available, using different reagents, and/or different intermediates, *etc.*


[00145] For example, General Reaction Scheme 2 describes another method for preparing Form A and Form B.



[00146] Reaction of A-1 with a strong Lewis acid (*e.g.*, boron tribromide) affords A-2, which can then be phosphorylated with an activated protected phosphate reagent (*e.g.*, di-*tert*-butylhalophosphonate, such as di-*tert*-butylchlorophosphonate or di-*tert*-butylbromophosphonate) under basic conditions (*e.g.*, using diisopropylethylamine as a base) in an appropriate solvent (*e.g.*, dimethylformamide; DMF). In some embodiments employing di-*tert*-butylhalophosphonate, the di-*tert*-butylhalophosphonate can be prepared (*e.g.*, *in situ* with A-2) by contacting di-*tert*-butylphosphonate with carbon tetrahalide (*e.g.*, carbon tetrachloride, carbon tetrabromide). The phosphorylated product (A-10) can then be deprotected under acidic conditions (*e.g.*, using hydrochloric and acetic acids). The resulting hydrochloride salt can then be converted to its free base and/or zwitterionic form, Form A (A-4), under neutralizing conditions (*e.g.*, ammonium carbonate in aqueous acetonitrile). The resultant compound A-4 can then be converted to Form B (A-5) using an appropriate

lattice forming reagent (*e.g.*, an acid, such as maleic acid, acetic acid, citric acid, propionic acid) and a suitable solvent (*e.g.*, THF and water at 19:1 v/v, methanol, methanol and acetone at 1:1 v/v). In some embodiments, the lattice forming reagent is removed from A-5 using a suitable, substantially anhydrous solvent (*e.g.*, an alcohol such as ethanol), and resulting A-5 is dried to afford Form B.

[00147] General Reaction Scheme 3 provides yet another method for preparing Form B.



[00148] As an alternative to performing the deprotection, free base/zwitterion formation and polymorph conversion steps in three, separate steps, some embodiments provide for deprotection and polymorph formation in a single step. Briefly, A-10 can be deprotected and converted to Form B (A-5) under acidic conditions (*e.g.*, maleic acid in methanol and acetone at 1:1 v/v) to afford Form B.

Pharmaceutical Compositions

[00149] In some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is formulated into a pharmaceutical composition. In any of the pharmaceutical compositions described herein, the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof can be the compound having structure (II), or any crystalline form or polymorph of a compound of structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof described herein, *e.g.*, Form B.

[00150] A pharmaceutical composition, as used herein, refers to a mixture of the compound having structure (I) (*e.g.* a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, with other chemical components, such as carriers, stabilizers, diluents, dispersing agents, suspending agents, thickening agents, and/or excipients.

[00151] In particular, one embodiment provides a pharmaceutical composition comprising a crystalline form and/or polymorph of a compound of structure (I), or a tautomer or zwitterionic form thereof, and a pharmaceutically acceptable carrier or excipient.

[00152] In some embodiments, the pharmaceutical composition is formulated for oral administration. Pharmaceutical compositions described herein are formulated by combining the active (*e.g.*, crystalline form and/or polymorph of a compound having structure (I), or a tautomer or zwitterionic form thereof) with, *e.g.*, pharmaceutically acceptable carriers or excipients. In various embodiments, the crystalline form and/or polymorph of a compound of structure (I), or a tautomer or zwitterionic form thereof is formulated in an oral dosage form that includes, by way of example only, tablets, powders, pills, dragees, capsules, liquids, gels, syrups, elixirs, slurries, suspensions and the like. In some embodiments, the pharmaceutical composition comprises an oral capsule.

[00153] In some embodiments, the pharmaceutical composition comprises the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, in crystalline or polymorph form), in a concentration ranging from about 0.5 weight percent to about 11.25 weight percent. In other specific embodiments, the pharmaceutical composition comprises the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, in crystalline or polymorph form) in a concentration ranging from about 0.6 weight percent to about 11.05 weight percent.

[00154] In more specific embodiments, the pharmaceutical composition comprises about 0.6 weight percent of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, in crystalline or polymorph form). In other specific embodiments, the pharmaceutical composition comprises about 2.5 weight percent of the compound having structure (I), or a tautomer,

pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, in crystalline or polymorph form). In still other specific embodiments, the pharmaceutical composition comprises about 11.0 weight percent of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, in crystalline or polymorph form).

[00155] In some more specific embodiments, the pharmaceutical composition comprises about 1 milligram (mg) of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, in crystalline or polymorph form). In other specific embodiments, the pharmaceutical composition comprises about 4 mg of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, in crystalline or polymorph form). In some embodiments, the pharmaceutical composition comprises about 16 mg of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, in crystalline or polymorph form).

[00156] In embodiments, the excipient is anhydrous. In some embodiments, the excipient is anhydrous lactose or microcrystalline cellulose. In more specific embodiments, the excipient is anhydrous lactose. In some embodiments, the excipient is microcrystalline cellulose. In specific embodiments, the excipient is anhydrous microcrystalline cellulose.

[00157] In other embodiments, the excipient is anhydrous lactose or cornstarch (*e.g.*, cornstarch).

[00158] In some of the foregoing embodiments, the pharmaceutical composition comprises the excipient in a concentration ranging from about 85 weight percent to about 98 weight percent. In more specific embodiments, the pharmaceutical composition comprises the excipient in a concentration ranging from about 87 weight percent to about 97.5 weight percent. In some embodiments, the pharmaceutical composition comprises about 87 weight percent of the excipient. In other embodiments, the pharmaceutical composition comprises about 95.5 weight percent of the excipient.

[00159] In related embodiments, the pharmaceutical composition further comprises a glidant. In some embodiments, the glidant is colloidal silicon dioxide. In some of these

embodiments, the pharmaceutical composition comprises about 1 weight percent of the glidant (*e.g.*, colloidal silicon dioxide).

[00160] In some related embodiments, the pharmaceutical composition further comprises a lubricant. In some embodiments, the lubricant is magnesium stearate. In some of these embodiments, the pharmaceutical composition comprises about 1 weight percent of the lubricant (*e.g.*, magnesium stearate). Other examples of lubricants are described in Example 13 herein.

[00161] In some related embodiments, the pharmaceutical composition further comprises a disintegrant. In some embodiments, the disintegrant is partly pregelatinized starch, low-substituted hydroxyl propyl cellulose or carmellose calcium. Other examples of disintegrants are described in Example 13 herein.

[00162] One embodiment is a pharmaceutical composition comprising from about 0.5 weight percent to about 11.25 weight percent (*e.g.*, from about 0.5 weight percent to about 1 weight percent, from about 10.5 weight percent to about 11.25 weight percent) of a compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, a compound having structure (II)), and from about 85 weight percent to about 99.5 weight percent (*e.g.*, from about 95 weight percent to about 99.5 weight percent, from about 85 weight percent to about 90 weight percent, respectively) of a pharmaceutically acceptable carrier or excipient.

[00163] One embodiment is a pharmaceutical composition comprising about 0.6 weight percent of a compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, a compound having structure (II)), about 97.4 weight percent of anhydrous lactose, about 1 weight percent colloidal silicon dioxide, and about 1 weight percent magnesium stearate. In a specific aspect of this embodiment, the pharmaceutical composition comprises about 1 mg of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, a compound having structure (II)).

[00164] One embodiment is a pharmaceutical composition comprising about 11 weight percent of a compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, a compound having structure (II)), about 87 weight percent of anhydrous lactose, about 1 weight percent colloidal silicon dioxide, and about 1 weight percent magnesium stearate. In a specific aspect of this

embodiment, the pharmaceutical composition comprises about 16 mg of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, a compound having structure (II)).

[00165] One embodiment is a pharmaceutical composition comprising about 0.6 weight percent of a compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, a compound having structure (II)), and from about 98 weight percent to about 99.5 weight percent of cornstarch. In a specific aspect of this embodiment, the pharmaceutical composition further comprises about 1 weight percent magnesium stearate.

[00166] One embodiment is a pharmaceutical composition comprising about 11 weight percent of a compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof (*e.g.*, a compound having structure (II)), and about 88 weight percent of cornstarch. In a specific aspect of this embodiment, the pharmaceutical composition further comprises about 1 weight percent magnesium stearate.

[00167] In some of the foregoing embodiments, the pharmaceutical compositions are in the form of a capsule for oral administration.

[00168] In embodiments, a pharmaceutical composition described herein (*e.g.*, comprising a crystalline form and/or polymorph of a compound of structure (I), or a tautomer or zwitterionic form thereof) is provided in the form of a rapid release formulation. As used herein, “rapid release formulation” refers to a formulation that does not delay or prolong release of an active contained therein. Typically, rapid release formulations release at least 70% of the active contained therein within 1 hour. In some embodiments, a rapid release formulation releases at least 75% of an active contained therein (*e.g.*, a compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof) within about 45 minutes. “Rapid release formulation” is also referred to as “immediate release formulation.”

[00169] In embodiments, a pharmaceutical composition described herein (*e.g.*, comprising a crystalline form and/or polymorph of a compound of structure (I), or a tautomer or zwitterionic form thereof) is provided in the form of an extended release formulation. As used herein, “extended release formulation” refers to a formulation that

releases an active contained therein in a controlled manner during an extended period of time, at a predetermined rate, duration, and location following administration.

[00170] In embodiments, a pharmaceutical composition described herein (*e.g.*, comprising a crystalline form and/or polymorph of a compound of structure (I), or a tautomer or zwitterionic form thereof) is provided in the form of an intermediate release formulation. As used herein, “intermediate release formulation” refers to a formulation that delays or extends release of an active contained therein, or alters its site of release, but is not an extended release formulation, as that term is described herein.

[00171] The pharmaceutical composition may also be formulated for a route of administration other than oral. Other suitable routes of administration include intravenous, rectal, aerosol, parenteral, ophthalmic, pulmonary, transmucosal, transdermal, vaginal, otic, nasal, and topical administration. In addition, by way of example only, parenteral delivery includes intramuscular, subcutaneous, intravenous, intramedullary injections, as well as intrathecal, direct intraventricular, intraperitoneal, intralymphatic, and intranasal injections.

[00172] In certain embodiments, a pharmaceutical composition described herein is administered in a local rather than systemic manner, for example, via injection of the pharmaceutical composition directly into an organ, often in a depot preparation or sustained release formulation. In specific embodiments, long-acting formulations are administered by implantation (for example, subcutaneously or intramuscularly) or by intramuscular injection. Furthermore, in other embodiments, the pharmaceutical composition is delivered in the form of a targeted drug delivery system, for example, a liposome coated with organ-specific antibody. In such embodiments, the liposomes are targeted to and taken up selectively by the organ. In yet other embodiments, the pharmaceutical composition is provided in the form of a rapid release formulation, in the form of an extended release formulation, or in the form of an intermediate release formulation. In yet other embodiments, the pharmaceutical composition is administered topically.

[00173] In other embodiments, the pharmaceutical composition is formulated for transmucosal administration. In specific embodiments, transmucosal formulations include penetrants that are appropriate to the barrier to be permeated. In still other embodiments, wherein the pharmaceutical composition is formulated for other

parenteral injections; appropriate formulations include aqueous or non-aqueous solutions. In specific embodiments, such solutions include physiologically compatible buffers and/or excipients.

[00174] In other embodiments, pharmaceutical compositions described herein are formulated for buccal or sublingual administration. Formulations suitable for buccal or sublingual administration include, by way of example only, tablets, lozenges, or gels.

[00175] In still other embodiments, the pharmaceutical composition described herein is formulated for parental injection, including bolus injection or continuous infusion. In specific embodiments, formulations for injection are presented in unit dosage form (*e.g.*, in ampoules) or in multi-dose containers. Preservatives are, optionally, added to the injection formulations. In still other embodiments, the pharmaceutical compositions are formulated in a form suitable for parenteral injection as sterile suspensions, solutions or emulsions in oily or aqueous vehicles. Parenteral injection formulations optionally contain formulatory agents such as suspending, stabilizing and/or dispersing agents. In additional embodiments, suspensions are prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles for use in the pharmaceutical compositions described herein include, by way of example only, fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. In certain specific embodiments, aqueous injection suspensions contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension contains suitable stabilizers or agents which increase the solubility of the active to allow for the preparation of highly concentrated solutions. Alternatively, in other embodiments, the active ingredient is in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

[00176] In some embodiments, the pharmaceutical composition is administered in dosages. It is known in the art that due to intersubject variability in compound pharmacokinetics, individualization of dosing regimen is necessary for optimal therapy. Dosing for the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, may be found by routine experimentation in light of the instant disclosure.

[00177] The compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, according to certain embodiments, is effective over a wide dosage range. For example, in the treatment of adult humans, dosages from 0.01 to 1000 mg, from 0.5 to 100 mg, from 1 to 50 mg per day, and from 5 to 40 mg per day are examples of dosages that are used in some embodiments. An exemplary dosage is 10 to 30 mg per day. In various embodiments, the dosage is 3, 6, 9, 12, 16, 21, 28, 32, 42, or 50 mg per day. In some embodiments, the dosage ranges from about 1 mg to about 30 mg per day, *e.g.*, about 1 mg, about 2 mg, about 4 mg, about 8 mg, about 12 mg, about 16 mg or about 22 mg per day. The exact dosage will depend upon the route of administration, the form in which the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is administered, the subject to be treated, the body weight of the subject to be treated, and the preference and experience of the attending physician.

[00178] In some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is administered in a single dose. Routes of administration are selected and used as appropriate. A single dose of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, may also be used for treatment of an acute condition.

[00179] In some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is administered in multiple doses. In some embodiments, dosing is about once, twice, three times, four times, five times, six times, or more than six times per day (*e.g.*, twice per day). In other embodiments, dosing is about once a month, once every two weeks, once a week, or once every other day.

[00180] In another embodiment, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is administered about once per day to about 6 times per day. In another embodiment, the administration of the polymorph of a compound of structure (I) continues for less than about 7 days. In

yet another embodiment, the administration continues for more than about 6, 10, 14, 28 days, two months, six months, or one year. In some cases, continuous dosing is achieved and maintained as long as necessary.

[00181] Administration of the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, may continue as long as necessary. In some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is administered for more than 1, 2, 3, 4, 5, 6, 7, 14, or 28 days. In some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is administered for less than 28, 14, 7, 6, 5, 4, 3, 2, or 1 day. In some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is administered chronically on an ongoing basis, *e.g.*, for the treatment of chronic effects. In some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof is administered once daily for the first 21 days out of a 28-day cycle. In some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof is administered twice daily for the first 21 days out of a 28-day cycle. A cycle may be repeated at least once, at least twice, at least three times, or at least four times.

[00182] In certain embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is administered as a pharmaceutical composition in which the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form, is mixed with one or more other active ingredients, as in combination therapy. Accordingly, in embodiments, provided is a pharmaceutical combination comprising a compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, and one or

more additional therapeutic agents (*e.g.*, chemotherapeutic agents). Encompassed herein are all combinations of actives set forth in the combination therapies section below and throughout this disclosure.

[00183] In some of the foregoing embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, has a water content less than 0.50%, as determined by Karl Fischer titration. For example, in some embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, has a water content less than 0.45%, less than 0.40%, less than 0.35%, less than 0.30%, less than 0.25%, less than 0.20%, less than 0.15%, or less than 0.10%, as determined by Karl Fischer titration.

[00184] Some particular embodiments provide a unit dose form comprising a pharmaceutical composition as described herein. In various embodiments, the unit dose form is formulated for oral administration. In some embodiments, the unit dose form is a capsule. In some embodiments, the unit dose form is a tablet.

[00185] In some embodiments, dosage forms, such as dragee cores and tablets, are provided with one or more suitable coatings. In specific embodiments, concentrated sugar solutions are used for coating the dosage form. The sugar solutions, optionally contain additional components, such as, by way of example only, gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs and/or pigments are also optionally added to the coatings for identification purposes. Additionally, the dyestuffs and/or pigments are optionally utilized to characterize different combinations of active compound doses.

[00186] In some embodiments, the unit dose form comprises the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, in a concentration ranging from about 0.5 weight percent to about 11.25 weight percent. In other specific embodiments, the unit dose form comprises the polymorph in a concentration ranging from about 0.6 weight percent to about 11.05 weight percent.

[00187] In more specific embodiments, the unit dose form comprises about 0.6 weight percent of the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof. In other specific embodiments, the unit dose form comprises about 2.5 weight percent of the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof. In still other specific embodiments, the unit dose form comprises about 11.0 weight percent of the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof.

[00188] In some more specific embodiments, the unit dose form comprises about 1 milligram (mg) of the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof. In other specific embodiments, the unit dose form comprises about 4 mg of the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof. In some embodiments, the unit dose form comprises about 16 mg of the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof.

[00189] In some of the foregoing embodiments, the unit dose form comprises the excipient in a concentration ranging from about 85 weight percent to about 98 weight percent. In more specific embodiments, the unit dose form comprises the excipient in a concentration ranging from about 87 weight percent to about 97.5 weight percent. In some embodiments, the unit dose form comprises about 87 weight percent of the excipient. In other embodiments, the unit dose form comprises about 95.5 weight percent of the excipient.

[00190] In embodiments, the excipient is anhydrous. In some embodiments, the excipient is anhydrous lactose or microcrystalline cellulose. In more specific embodiments, the excipient is anhydrous lactose. In some embodiments, the excipient

is microcrystalline cellulose. In specific embodiments, the excipient is anhydrous microcrystalline cellulose.

[00191] In some embodiments, the excipient is anhydrous lactose or cornstarch.

[00192] In related embodiments, the unit dose form further comprises a glidant. In some embodiments, the glidant is colloidal silicon dioxide. In some of these embodiments, the unit dose form comprises about 1 weight percent of the glidant (*e.g.*, colloidal silicon dioxide).

[00193] In some related embodiments, the unit dose form further comprises a lubricant. In some embodiments, the lubricant is magnesium stearate. In some of these embodiments, the unit dose form comprises about 1 weight percent of the lubricant (*e.g.*, magnesium stearate). Other examples of lubricants are described in Example 13 herein.

[00194] In some related embodiments, the pharmaceutical composition further comprises a disintegrant. In some embodiments, the disintegrant is partly pregelatinized starch, low-substituted hydroxyl propyl cellulose or carmellose calcium. Other examples of disintegrants are described in Example 13 herein.

[00195] In particular embodiments, the unit dose form is in the form of a capsule for oral administration, *e.g.*, a hydroxypropylmethylcellulose capsule. In certain embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, are formulated into other oral dosage forms. Oral dosage forms include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. In specific embodiments, push-fit capsules contain the active ingredients in admixture with one or more fillers. Fillers include, by way of example only, lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In other embodiments, soft capsules contain the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, dissolved or suspended in a suitable liquid. Suitable liquids include, by way of example only, one or more fatty oil, liquid paraffin, or liquid polyethylene glycol. In addition, stabilizers are optionally added.

[00196] The pharmaceutical compositions described herein are formulated in any conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compound(s) into preparations which can be used pharmaceutically. In certain embodiments, the formulation of the pharmaceutical composition facilitates administration to a subject.

[00197] Proper formulation is dependent upon the route of administration chosen. Any pharmaceutically acceptable techniques, carriers, and excipients are optionally used as suitable to formulate the pharmaceutical compositions described herein:

Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999), the relevant teachings of which are incorporated herein by reference in their entireties.

[00198] In embodiments, pharmaceutical compositions described herein are manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

[00199] In particular embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is mixed with an excipient and, optionally, a lubricant and a glidant using direct blending techniques, and capsules are filled with the resulting mixture on a manual capsule filling machine. In some particular embodiments, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, the excipient and the optional glidant are put through a 500 μm screen, then mixed together, and blended for 16 minutes at 30 rpm. The optional lubricant is then sieved through a 250 μm screen and added to the mixture. The resulting mixture is then blended for 5 minutes at 30 rpm. Capsules are then filled using an overage of from 0% to about 5% (*e.g.*, about 1%), using tamping and/or tapping, as needed.

[00200] In certain embodiments, pharmaceutical preparations for oral use are obtained by mixing one or more solid excipients with the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, optionally grinding the resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as: for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methylcellulose, microcrystalline cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose; or others, such as polyvinylpyrrolidone (PVP or povidone) or calcium phosphate. In specific embodiments, disintegrating agents are optionally added. Disintegrating agents include, by way of example only, cross-linked croscarmellose sodium, polyvinylpyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

[00201] Additionally, the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, can encompass unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as, for example, ethanol, and the like. The solvated forms of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, are also considered to be disclosed herein. In addition, the pharmaceutical compositions optionally include other medicinal or pharmaceutical agents, carriers, adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure, buffers, and/or other therapeutically valuable substances.

[00202] Methods for the preparation of compositions comprising the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, include formulating with one or more inert, pharmaceutically acceptable excipients or carriers to form a solid, semi-solid or liquid. Solid compositions include powders, tablets, dispersible granules, capsules, cachets, and suppositories. Liquid compositions include solutions in which the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is dissolved,

emulsions comprising the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, or a solution containing liposomes, micelles, or nanoparticles comprising the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof. Semi-solid compositions include gels, suspensions and creams. The form of the pharmaceutical compositions described herein include liquid solutions or suspensions, solid forms suitable for solution or suspension in a liquid prior to use, or as emulsions. These compositions also optionally contain minor amounts of nontoxic, auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, and so forth.

[00203] In some embodiments, the pharmaceutical composition comprising the compound having structure (I) (*e.g.*, a crystalline form and/or polymorph of a compound having structure (I)), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, takes the form of a liquid where the agents are present in solution, in suspension or both. Typically, when the composition is administered as a solution or suspension, a first portion of the agent is present in solution and a second portion of the agent is present in particulate form, in suspension in a liquid matrix. In some embodiments, a liquid composition includes a gel formulation. In other embodiments, the liquid composition is aqueous.

[00204] In certain embodiments, useful aqueous suspensions contain one or more polymers as suspending agents. Useful polymers include water-soluble polymers, such as cellulosic polymers, *e.g.*, hydroxypropylmethylcellulose, and water-insoluble polymers, such as cross-linked carboxyl-containing polymers. Certain pharmaceutical compositions described herein comprise a mucoadhesive polymer, selected, for example, from carboxymethylcellulose, carbomer (acrylic acid polymer), poly(methylmethacrylate), polyacrylamide, polycarbophil, acrylic acid/butyl acrylate copolymer, sodium alginate or dextran.

[00205] Useful pharmaceutical compositions also, optionally, include solubilizing agents to aid in the solubility of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof. The term “solubilizing agent” generally includes agents that result in formation of a micellar solution or a true solution of the agent. Certain acceptable nonionic surfactants, for example, polysorbate

80, are useful as solubilizing agents, as are ophthalmically acceptable glycols, polyglycols, *e.g.*, polyethylene glycol 400, and glycol ethers.

[00206] Furthermore, pharmaceutical compositions optionally include one or more pH adjusting agents or buffering agents, including acids, such as acetic, boric, citric, lactic, phosphoric and hydrochloric acids; bases such as sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, sodium lactate and tris-hydroxymethylaminomethane; and buffers such as citrate/dextrose, sodium bicarbonate and ammonium chloride. Such acids, bases and buffers are included in an amount required to maintain pH of the composition in an acceptable range.

[00207] Pharmaceutical compositions also, optionally, include one or more salts in an amount required to bring osmolality of the composition into an acceptable range. Such salts include those having, for example, sodium, potassium or ammonium cations and chloride, citrate, ascorbate, borate, phosphate, bicarbonate, sulfate, thiosulfate or bisulfite anions; suitable salts include sodium chloride, potassium chloride, sodium thiosulfate, sodium bisulfite and ammonium sulfate.

[00208] Pharmaceutical compositions optionally include one or more preservatives to inhibit microbial activity. Suitable preservatives include mercury-containing substances such as merfen and thiomersal; stabilized chlorine dioxide; and quaternary ammonium compounds such as benzalkonium chloride, cetyltrimethylammonium bromide and cetylpyridinium chloride.

[00209] Still other pharmaceutical compositions include one or more surfactants to enhance physical stability or for other purposes. Suitable nonionic surfactants include polyoxyethylene fatty acid glycerides and vegetable oils, *e.g.*, polyoxyethylene (60) hydrogenated castor oil; and polyoxyethylene alkylethers and alkylphenyl ethers, *e.g.*, octoxynol 10, octoxynol 40.

[00210] Still other pharmaceutical compositions include one or more antioxidants to enhance chemical stability where required. Suitable antioxidants include, by way of example only, ascorbic acid and sodium metabisulfite.

[00211] In certain embodiments, aqueous suspension compositions are packaged in single-dose non-reclosable containers. Alternatively, multiple-dose, reclosable containers are used, in which case it is typical to include a preservative in the composition.

[00212] In alternative embodiments, other delivery systems for hydrophobic pharmaceutical compounds are employed. Liposomes and emulsions are examples of delivery vehicles or carriers useful herein. In certain embodiments, organic solvents such as N-methylpyrrolidone are also employed. In additional embodiments, the compounds described herein are delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the active. Various sustained-release materials are useful herein. In some embodiments, sustained-release capsules release the active for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the active, additional strategies for protein stabilization are employed.

[00213] In certain embodiments, the formulations described herein comprise one or more antioxidants, metal chelating agents, thiol-containing compounds and/or other general stabilizing agents. Examples of such stabilizing agents, include: (a) about 0.5% to about 2% w/v glycerol, (b) about 0.1% to about 1% w/v methionine, (c) about 0.1% to about 2% w/v monothioglycerol, (d) about 1 mM to about 10 mM EDTA, (e) about 0.01% to about 2% w/v ascorbic acid, (f) 0.003% to about 0.02% w/v polysorbate 80, (g) 0.001% to about 0.05% w/v. polysorbate 20, (h) arginine, (i) heparin, (j) dextran sulfate, (k) cyclodextrins, (l) pentosan polysulfate and other heparinoids, (m) divalent cations such as magnesium and zinc; or (n) combinations thereof.

[00214] In some embodiments, the concentration of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, provided in a pharmaceutical composition or unit dose form is no more than 100%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1%, 0.09%, 0.08%, 0.07%, 0.06%, 0.05%, 0.04%, 0.03%, 0.02%, 0.01%, 0.009%, 0.008%, 0.007%, 0.006%, 0.005%, 0.004%, 0.003%, 0.002%, 0.001%, 0.0009%, 0.0008%, 0.0007%, 0.0006%, 0.0005%, 0.0004%, 0.0003%, 0.0002%, or 0.0001% w/w, w/v or v/v.

[00215] In some embodiments, the concentration of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is in the range of from approximately 0.0001% to approximately 50%, approximately 0.001% to approximately 40 %, approximately 0.01% to approximately 30%,

approximately 0.02% to approximately 29%, approximately 0.03% to approximately 28%, approximately 0.04% to approximately 27%, approximately 0.05% to approximately 26%, approximately 0.06% to approximately 25%, approximately 0.07% to approximately 24%, approximately 0.08% to approximately 23%, approximately 0.09% to approximately 22%, approximately 0.1% to approximately 21%, approximately 0.2% to approximately 20%, approximately 0.3% to approximately 19%, approximately 0.4% to approximately 18%, approximately 0.5% to approximately 17%, approximately 0.6% to approximately 16%, approximately 0.7% to approximately 15%, approximately 0.8% to approximately 14%, approximately 0.9% to approximately 12%, or approximately 1% to approximately 10% w/w, w/v or v/v.

[00216] In some embodiments, the concentration of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, is in the range of from approximately 0.001% to approximately 10%, approximately 0.01% to approximately 5%, approximately 0.02% to approximately 4.5%, approximately 0.03% to approximately 4%, approximately 0.04% to approximately 3.5%, approximately 0.05% to approximately 3%, approximately 0.06% to approximately 2.5%, approximately 0.07% to approximately 2%, approximately 0.08% to approximately 1.5%, approximately 0.09% to approximately 1%, or approximately 0.1% to approximately 0.9% w/w, w/v or v/v.

[00217] In some embodiments, the amount of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, provided in a pharmaceutical composition or unit dose form is no more than 10 g, 9.5 g, 9.0 g, 8.5 g, 8.0 g, 7.5 g, 7.0 g, 6.5 g, 6.0 g, 5.5 g, 5.0 g, 4.5 g, 4.0 g, 3.5 g, 3.0 g, 2.5 g, 2.0 g, 1.5 g, 1.0 g, 0.95 g, 0.9 g, 0.85 g, 0.8 g, 0.75 g, 0.7 g, 0.65 g, 0.6 g, 0.55 g, 0.5 g, 0.45 g, 0.4 g, 0.35 g, 0.3 g, 0.25 g, 0.2 g, 0.15 g, 0.1 g, 0.09 g, 0.08 g, 0.07 g, 0.06 g, 0.05 g, 0.04 g, 0.03 g, 0.02 g, 0.01 g, 0.009 g, 0.008 g, 0.007 g, 0.006 g, 0.005 g, 0.004 g, 0.003 g, 0.002 g, 0.001 g, 0.0009 g, 0.0008 g, 0.0007 g, 0.0006 g, 0.0005 g, 0.0004 g, 0.0003 g, 0.0002 g, or 0.0001 g.

Methods of Treatment

[00218] Embodiments of the present disclosure include methods for treating a disease associated with overexpression of a cyclin-dependent kinase (CDK), such as

CDK 9, 1, 2, 4 or 7 (*e.g.*, CDK9), in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound having structure (I) (*e.g.*, a crystalline form and/or polymorph thereof), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof; pharmaceutical composition; and/or unit dose form described herein. In some embodiments, the disease associated with overexpression of a CDK is cancer.

[00219] Some embodiments include methods for treatment of cancer comprising administering an effective amount of a compound having structure (I) (*e.g.*, a crystalline form and/or polymorph thereof), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, to a subject in need thereof. Certain embodiments provide a method for treatment of cancer, the method comprising administering an effective amount of a pharmaceutical composition comprising a compound having structure (I) (*e.g.*, a crystalline form and/or polymorph thereof), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, to a subject in need thereof. The compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, may be used singly or in combination with one or more therapeutic agents, for example, as components of one or more mixtures.

[00220] In some embodiments, the cancer treated is a hematologic cancer. Hematologic malignancies that can be treated with a compound having structure (I), or a tautomer or zwitterionic form thereof, include leukemias and lymphomas. In some embodiments, the hematologic cancer is selected from acute myelogenous leukemia (AML), follicular lymphoma, acute lymphoblastic leukemia (ALL), chronic lymphocytic leukemia (CLL), multiple myeloma (MM) and non-Hodgkin's lymphoma (*e.g.*, AML, follicular lymphoma, ALL, CLL and non-Hodgkin's lymphoma). In more specific embodiments, the hematological cancer is AML. In other more specific embodiments, the hematologic cancer is CLL. In more specific embodiments, the hematologic cancer is MM. In still other specific embodiments, the hematologic cancer is myelodysplastic syndrome (MDS).

[00221] In other embodiments, the cancer comprises a solid tumor. Accordingly, in some embodiments, the cancer is a solid tumor cancer. In various embodiments, the solid tumor cancer is breast cancer, bladder cancer, liver cancer, pancreatic cancer, lung cancer, colorectal cancer, ovarian cancer, prostate cancer, or melanoma. In some

embodiments, the cancer is bladder cancer. In some embodiments, the cancer is lung cancer. In other embodiments, the cancer is liver cancer. In some embodiments, the cancer is a sarcoma, bladder cancer or renal cancer. In some embodiments, the cancer is prostate cancer (*e.g.*, castration-resistant prostate cancer, castration-sensitive prostate cancer). In other embodiments, the cancer is bladder cancer, pancreatic cancer, colorectal cancer, kidney cancer, non-small cell lung carcinoma, prostate cancer, sarcoma, skin cancer, thyroid cancer, testicular cancer or vulvar cancer. In some embodiments, the cancer is endometrial cancer, pancreatic cancer, testicular cancer, renal cancer, melanoma, colorectal cancer, thyroid cancer, bladder cancer, pancreatic cancer, vulvar cancer, sarcoma, prostate cancer, lung cancer or anal cancer. In some embodiments, the cancer is a sarcoma. In some embodiments, the cancer is a renal cell carcinoma.

[00222] “Castration-resistant prostate cancer” refers to prostate cancer that progresses in a subject following administration of one or more androgen depletion therapies (ADTs). Progression of prostate cancer can be evidenced by, for example, a prostate-specific antigen doubling time (PSADT) of less than or equal to 10 months, the progression of pre-existing disease (*e.g.*, radiographic progression, clinical progression, a skeletal-related event, prostate-specific antigen (PSA) progression), and/or the appearance of new metastases in a subject, and is typically driven by androgens, which are a class of hormones including testosterone and dihydrotestosterone (DHT). These androgens bind to the androgen receptor (AR), which is a transcription activator that promotes growth and survival of prostate cells, including prostate cancer cells. ADT refers to a therapy to suppress androgen levels (*e.g.*, surgical castration or chemical castration) or androgen signaling (*e.g.*, by reducing androgen binding to androgen receptor), which may be used to slow the progression of prostate cancer. Androgen deprivation therapy typically causes a temporary reduction in tumor burden concomitant with a decrease in serum PSA. Mechanisms of castration resistance include the emergence of AR variants that are active in the absence of androgen, including splice variants, point mutations to AR, and AR gene amplifications. Castration resistance can be biochemically characterized before the onset of symptoms by a rising titer of serum PSA (Miller, *et al.*, 1992 J. Urol. 147, 956 961). Radiographic progression can be assessed with the use of sequential imaging, and is evidenced by, for

example, bone scan identification of two or more new bone lesions with confirmation (according to the Prostate Cancer Clinical Trials Working Group 2 criteria). Response Evaluation Criteria in Solid Tumors (RECIST v 1.1) criteria can also be used to assess radiographic progression of soft tissue lesions. Guidelines for monitoring prostate cancer, including progression of prostate cancer, are described in NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer, version 4.2019, August 19, 2019, the relevant contents of which are incorporated herein by reference in their entirety.

“Castration-resistant prostate cancer” is used interchangeably herein with “androgen-resistant prostate cancer”, “androgen-independent prostate cancer” and “hormone-resistant prostate cancer”.

[00223] “Castration-sensitive prostate cancer” refers to prostate cancer that does not progress (*e.g.*, responds) following administration of one or more ADTs. Progression of prostate cancer can be assessed according to criteria described herein, for example, with respect to “castration-resistant prostate cancer,” and guidelines for monitoring prostate cancer, including progression of prostate cancer, are described in NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer, version 4.2019, August 19, 2019, the relevant contents of which are incorporated herein by reference in their entirety. “Castration-sensitive prostate cancer” is used interchangeably herein with “androgen-sensitive prostate cancer”, “androgen-dependent prostate cancer” and “hormone-sensitive prostate cancer”.

[00224] Further examples of cancers treatable according to the methods described herein include, but are not limited to, Acute Lymphoblastic Leukemia (ALL); Acute Myeloid Leukemia (AML); Adrenocortical Carcinoma; Adrenocortical Carcinoma, Childhood; AIDS-Related Cancer (*e.g.*, Kaposi Sarcoma, AIDS-Related Lymphoma, Primary CNS Lymphoma); Anal Cancer; Appendix Cancer; Astrocytomas, Childhood; Atypical Teratoid/Rhabdoid Tumor, Childhood, Central Nervous System; Basal Cell Carcinoma of the Skin; Bile Duct Cancer; Bladder Cancer; Bladder Cancer, Childhood; Bone Cancer (including Ewing Sarcoma, Osteosarcoma and Malignant Fibrous Histiocytoma); Brain Tumors/Cancer; Breast Cancer; Burkitt Lymphoma; Carcinoid Tumor (Gastrointestinal); Carcinoid Tumor, Childhood; Cardiac (Heart) Tumors, Childhood; Embryonal Tumors, Childhood; Germ Cell Tumor, Childhood; Primary CNS Lymphoma; Cervical Cancer; Childhood Cervical Cancer; Cholangiocarcinoma;

Chordoma, Childhood; Chronic Lymphocytic Leukemia (CLL); Chronic Myelogenous Leukemia (CML); Chronic Myeloproliferative Neoplasms; Colorectal Cancer; Childhood Colorectal Cancer; Craniopharyngioma, Childhood; Cutaneous T-Cell Lymphoma (*e.g.*, Mycosis Fungoides and Sézary Syndrome); Ductal Carcinoma In Situ (DCIS); Embryonal Tumors, Central Nervous System, Childhood; Endometrial Cancer (Uterine Cancer); Ependymoma, Childhood; Esophageal Cancer; Childhood Esophageal Cancer; Esthesioneuroblastoma; Ewing Sarcoma; Extracranial Germ Cell Tumor, Childhood; Extragonadal Germ Cell Tumor; Eye Cancer; Childhood Intraocular Melanoma; Intraocular Melanoma; Retinoblastoma; Fallopian Tube Cancer; Fibrous Histiocytoma of Bone, Malignant, and Osteosarcoma; Gallbladder Cancer; Gastric (Stomach) Cancer; Childhood Gastric (Stomach) Cancer; Gastrointestinal Carcinoid Tumor; Gastrointestinal Stromal Tumors (GIST); Childhood Gastrointestinal Stromal Tumors; Germ Cell Tumors; Childhood Central Nervous System Germ Cell Tumors (*e.g.*, Childhood Extracranial Germ Cell Tumors, Extragonadal Germ Cell Tumors, Ovarian Germ Cell Tumors, Testicular Cancer); Gestational Trophoblastic Disease; Hairy Cell Leukemia; Head and Neck Cancer; Heart Tumors, Childhood; Hepatocellular (Liver) Cancer; Histiocytosis, Langerhans Cell; Hodgkin Lymphoma; Hypopharyngeal Cancer; Intraocular Melanoma; Childhood Intraocular Melanoma; Islet Cell Tumors, Pancreatic Neuroendocrine Tumors; Kaposi Sarcoma; Kidney (Renal Cell) Cancer; Langerhans Cell Histiocytosis; Laryngeal Cancer; Leukemia; Lip and Oral Cavity Cancer; Liver Cancer; Lung Cancer (Non-Small Cell and Small Cell); Childhood Lung Cancer; Lymphoma; Male Breast Cancer; Malignant Fibrous Histiocytoma of Bone and Osteosarcoma; Melanoma; Childhood Melanoma; Melanoma, Intraocular (Eye); Childhood Intraocular Melanoma; Merkel Cell Carcinoma; Mesothelioma, Malignant; Childhood Mesothelioma; Metastatic Cancer; Metastatic Squamous Neck Cancer with Occult Primary; Midline Tract Carcinoma With NUT Gene Changes; Mouth Cancer; Multiple Endocrine Neoplasia Syndromes; Multiple Myeloma/Plasma Cell Neoplasms; Mycosis Fungoides; Myelodysplastic Syndromes, Myelodysplastic/Myeloproliferative Neoplasms; Myelogenous Leukemia, Chronic (CML); Myeloid Leukemia, Acute (AML); Myeloproliferative Neoplasms, Chronic; Nasal Cavity and Paranasal Sinus Cancer; Nasopharyngeal Cancer; Neuroblastoma; Non-Hodgkin Lymphoma; Non-Small Cell Lung Cancer; Oral Cancer,

Lip and Oral Cavity Cancer and Oropharyngeal Cancer; Osteosarcoma and Malignant Fibrous Histiocytoma of Bone; Ovarian Cancer; Childhood Ovarian Cancer; Pancreatic Cancer; Childhood Pancreatic Cancer; Pancreatic Neuroendocrine Tumors; Papillomatosis (Childhood Laryngeal); Paraganglioma; Childhood Paraganglioma; Paranasal Sinus and Nasal Cavity Cancer; Parathyroid Cancer; Penile Cancer; Pharyngeal Cancer; Pheochromocytoma; Childhood Pheochromocytoma; Pituitary Tumor; Plasma Cell Neoplasm/Multiple Myeloma; Pleuropulmonary Blastoma; Pregnancy and Breast Cancer; Primary Central Nervous System (CNS) Lymphoma; Primary Peritoneal Cancer; Prostate Cancer; Rectal Cancer; Recurrent Cancer; Renal Cell (Kidney) Cancer; Retinoblastoma; Rhabdomyosarcoma, Childhood; Salivary Gland Cancer; Sarcoma (*e.g.*, Childhood Rhabdomyosarcoma, Childhood Vascular Tumors, Ewing Sarcoma, Kaposi Sarcoma, Osteosarcoma (Bone Cancer), Soft Tissue Sarcoma, Uterine Sarcoma); Sézary Syndrome; Skin Cancer; Childhood Skin Cancer; Small Cell Lung Cancer; Small Intestine Cancer; Soft Tissue Sarcoma; Squamous Cell Carcinoma of the Skin; Squamous Neck Cancer with Occult Primary, Metastatic; Stomach (Gastric) Cancer; Childhood Stomach (Gastric) Cancer; T-Cell Lymphoma, Cutaneous (*e.g.*, Mycosis Fungoides and Sézary Syndrome); Testicular Cancer; Childhood Testicular Cancer; Throat Cancer (*e.g.*, Nasopharyngeal Cancer, Oropharyngeal Cancer, Hypopharyngeal Cancer); Thymoma and Thymic Carcinoma; Thyroid Cancer; Transitional Cell Cancer of the Renal Pelvis and Ureter; Ureter and Renal Pelvis, Transitional Cell Cancer; Urethral Cancer; Uterine Cancer, Endometrial; Uterine Sarcoma; Vaginal Cancer; Childhood Vaginal Cancer; Vascular Tumors; Vulvar Cancer; and Wilms Tumor and Other Childhood Kidney Tumors.

[00225] Metastases of the aforementioned cancers can also be treated in accordance with the methods described herein. Thus, in some embodiments, the cancer is a metastatic cancer. In other embodiments, the cancer is a primary cancer.

[00226] In some embodiments, the cancer is MCL-1 dependent. As used herein, “MCL-1-dependent” refers to the subset of cancers (*e.g.*, hematologic cancers) wherein myeloid cell leukemia 1 (MCL-1) is the primary driver of suppressing apoptosis. Typically, MCL-1 dependency promotes blast survival, and is associated with treatment resistance and relapse. MCL-1 dependence can be assessed, for example, by contacting a subject’s cancer cell with a profiling peptide, as described in International Publication

Nos. WO 2016/172214 and WO 2018/119000, the relevant contents of which are incorporated herein by reference in their entireties.

[00227] In some embodiments, the cancer is c-Myc-altered. As used herein, “c-Myc-altered” refers to the subset of cancers wherein c-Myc is altered compared to its native sequence, where its expression is amplified compared to an appropriate control (*e.g.*, corresponding normal cells), and where protein levels suggest overexpression of c-Myc. For example, it has been found that c-Myc drives androgen independence in prostate cancer, and overexpression attenuates the anti-tumor activity of androgen receptor suppression. In addition, c-Myc is significantly upregulated in androgen receptor-sensitive prostate cancer. Examples of cancers that can be c-Myc-altered include, but are not limited to, lymphoma (*e.g.*, Burkitt lymphoma, B-cell lymphoma, T-cell lymphoma), cervical cancer, colon cancer, ovarian cancer, breast cancer, lung cancer, prostate cancer, colorectal cancer, pancreatic cancer, gastric cancer and uterine cancer.

[00228] In some embodiments of the methods of treatment, the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, the pharmaceutical composition, or the unit dose form is administered orally.

[00229] Some embodiments provide methods for combination therapies in which an agent known to modulate other pathways, or other components of the same pathway, or even overlapping sets of target enzymes are used in combination with a compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof. In one aspect, such therapy includes the combination of the compound having structure (I), or a tautomer, pharmaceutically acceptable salt or zwitterionic form thereof, with a chemotherapeutic agent, therapeutic antibody, and/or radiation treatment, *e.g.*, to provide a synergistic or additive therapeutic effect.

[00230] Many chemotherapeutics are presently known in the art and can be used in combination with a compound having structure (I), or a pharmaceutically acceptable salt or zwitterionic form thereof. In some embodiments, the chemotherapeutic is selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites, intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, biological response modifiers, anti-hormones, angiogenesis inhibitors, and anti-androgens.

[00231] For example, the further therapeutic agent may comprise an alkylating agent, such as chlorambucil, cyclophosphamide, cisplatin; a mitotic inhibitor such as docetaxel (Taxotere; 1,7 β ,10 β -trihydroxy-9-oxo-5 β ,20-epoxytax-11-ene-2 α ,4,13 α -triyl 4-acetate 2-benzoate 13-{(2R,3S)-3-[(tert-butoxycarbonyl)amino]-2-hydroxy-3-phenylpropanoate}) or paclitaxel; an antimetabolite such as 5-fluorouracil, cytarabine, methotrexate, or pemetrexed; an anti-tumor antibiotic such as daunorubicin or doxorubicin; a corticosteroid such as prednisone or methylprednisone; or a Bcl-2 inhibitor such as venetoclax.

[00232] In certain embodiments, the further therapeutic agent is docetaxel. Docetaxel (trade name TAXOTERE®) is a type of chemotherapeutic agent known as an antimicrotubule agent. Docetaxel is used for treating a variety of cancers, such as metastatic prostate cancer. Docetaxel treatment is often administered intravenously, and often includes premedication with a corticosteroid such as prednisone.

[00233] In certain aspects of all embodiments, the further therapeutic agent is venetoclax (GDC-0199, ABT199, RG7601, trade name VENCLEXTA® or VENCLYXTO®), which is a Bcl-2 inhibitor that can induce apoptosis in cancer cells. Venetoclax is typically administered orally.

[00234] The further therapeutic agent may be a pharmacological agent that is currently approved by the Food and Drug Administration (FDA) in the U.S. (or elsewhere by any other regulatory body) for use as pharmacological treatment of prostate cancer, or is currently being used experimentally as part of a clinical trial program that relates to prostate cancer. For example, the further therapeutic agents may comprise, without limitation, the chemical entity known as enzalutamide or MDV3100 (4-(3-(4-cyano-3-(trifluoromethyl)phenyl)-5,5-dimethyl-4-oxo-2-thioxoimidazolidin-1-yl)-2-fluoro-N-methylbenzamide) and related compounds; the chemical entity known as TOK 001 and related compounds; the chemical entity known as ARN-509; the chemical entity known as abiraterone (or CB-7630; (3S,8R,9S,10R,13S,14S)-10,13-dimethyl-17-(pyridin-3-yl) 2,3,4,7,8,9,10,11,12,13,14,15-dodecahydro-1H-cyclopenta[a]phenanthren-3-ol), and related molecules; the chemical entity known as bicalutamide (N-[4-cyano-3-(trifluoromethyl)phenyl]-3-[(4-fluorophenyl)sulfonyl]-2-hydroxy-2-methylpropanamide) and related compounds; the chemical entity known as nilutamide (5,5-dimethyl-3-[4-nitro-3-(trifluoromethyl)phenyl] imidazolidine-2,4-

dione) and related compounds; the chemical entity known as flutamide (2-methyl-N-[4-nitro-3-(trifluoromethyl)phenyl]-propanamide) and related compounds; the chemical entity known as cyproterone acetate (6-chloro-1 β ,2 β -dihydro-17-hydroxy-3'H-cyclopropa[1,2]pregna-4,6-diene-3,20-dione) and related compounds, which is currently used to treat prostate cancer, the chemical entity known as docetaxel and related compounds, which is currently used alone or in combination with prednisone to treat prostate cancer, the chemical entity known as bevacizumab (Avastin), a monoclonal antibody that may be used to treat prostate cancer, the chemical entity known as OSU-HDAC42 ((S)-(+)-N-hydroxy-4-(3-methyl-2-phenylbutyrylamino)-benzamide), and related compounds; the chemical entity known as VITAXIN, which may be used to treat prostate cancer, the chemical entity known as sunitumib (N-(2-diethylaminoethyl)-5-[(Z)-(5-fluoro-2-oxo-1H-indol-3-ylidene)methyl]-2,4-dimethyl-1H-pyrrole-3-carboxamide) and related compounds, which may be used for treatment of prostate cancer, the chemical entity known as ZD-4054 (N-(3-Methoxy-5-methylpyrazin-2-yl)-2-[4-(1,3,4-oxadiazol-2-yl)phenyl]pyridin-3-sulfonamid) and related compounds; the chemical entity known as VN/124-1 (3 β -Hydroxy-17-(1H-benzimidazol-1-yl)androsta-5,16-diene), and related compounds; the chemical entity known as cabazitaxel (XRP-6258), and related compounds; the chemical entity known as MDX-010 (Ipilimumab); the chemical entity known as OGX 427; the chemical entity known as OGX 011; the chemical entity known as finasteride (Proscar, Propecia; N-(1,1-dimethylethyl)-3-oxo-(5 α ,17 β)-4-azaandrost-1-ene-17-carboxamide), and related compounds; the chemical entity known as dutasteride (Avodart; 5 α ,17 β)-N-{2,5-bis(trifluoromethyl)phenyl}-3-oxo-4-azaandrost-1-ene-17-carboxamide) and related compounds; the chemical entity known as turosteride ((4aR,4bS,6aS,7S,9aS,9bS,11aR)-1,4a,6a-trimethyl-2-oxo-N-(propan-2-yl)-N-(propan-2-ylcarbonyl)hexadecahydro-1H-indeno[5,4-f]quinoline-7-carboxamide), and related compounds; the chemical entity known as bexlosteride (LY-191,704; (4aS,10bR)-8-chloro-4-methyl-1,2,4a,5,6,10b-hexahydrobenzo[f]quinolin-3-one), and related compounds; the chemical entity known as izonsteride (LY-320,236; (4aR,10bR)-8-[(4-ethyl-1,3-benzothiazol-2-yl)sulfanyl]-4,10b-dimethyl-1,4,4a,5,6,10b-hexahydrobenzo[f]quinolin-3(2H)-one) and related compounds; the chemical entity known as FCE 28260 and related compounds; the chemical entity known as SKF105,111, and related compounds; the chemical entity

known as AZD3514; the chemical entity known as EZN-4176; the chemical entity known as ODM-201, sipuleucel-T, cabazitaxel; a combination of bevacizumab, docetaxel, thalidomide and prednisone; and/or abiraterone. In certain aspects of all embodiments, the further therapeutic agent is an androgen receptor antagonist that blocks androgen binding to androgen receptor. Examples of therapies that block androgen binding to androgen receptor include enzalutamide and apalutamide. In particular embodiments, the further therapeutic agent is enzalutamide. Enzalutamide (trade name XTANDI®) is an androgen receptor (AR) antagonist that is used for treating non-metastatic castration-resistant prostate cancer and metastatic castration-resistant prostate cancer. Enzalutamide treatment may be combined with castration (surgical or chemical).

[00235] In certain aspects of all embodiments, the further therapeutic agent is abiraterone. Abiraterone (trade name ZYTIGA®) is a CYP17A1 inhibitor, which significantly decreases testosterone production. Abiraterone treatment may be combined with other further therapies, such as a corticosteroid (*e.g.*, prednisone) and/or castration (surgical or chemical).

[00236] In certain aspects of all embodiments, the further therapeutic agent is selected from at least one of: a bromodomain inhibitor, a histone methyltransferase inhibitor, a histone deacetylase inhibitor, or a histone demethylases inhibitor.

[00237] In certain aspects of all embodiments, the further therapeutic agent is a bromodomain inhibitor, for example, an inhibitor of a bromodomain protein such as Brd2, Brd3, Brd4 and/or BrdT. In particular embodiments, the further therapeutic agent comprises a BRD4 inhibitor. In some of these embodiments, the further therapeutic agent is JQ-1 (Nature 2010 Dec 23;468(7327):1067-73), BI2536 (ACS Chem. Biol. 2014 May 16;9(5):1160-71; Boehringer Ingelheim), TG101209 (ACS Chem. Biol. 2014 May 16;9(5):1160-71), OTX015 (Mol. Cancer Ther. November 2013 12; C244; Oncoethix), IBET762 (J Med Chem. 2013 Oct 10;56(19):7498-500; GlaxoSmithKline), IBET151 (Bioorg. Med. Chem. Lett. 2012 Apr 15;22(8):2968-72; GlaxoSmithKline), PFI-1 (J. Med. Chem. 2012 Nov 26;55(22):9831-7; Cancer Res. 2013 Jun 1;73(11):3336-46; Structural Genomics Consortium) or CPI-0610 (Constellation Pharmaceuticals). In other embodiments, the BRD inhibitor is IBET 762 (GSK525762), TEN-010 (Tensha Therapeutics), CPI-203 (*Leukemia*. 28 (10): 2049–59, 2014), RVX-

208 (*Proceedings of the National Academy of Sciences of the United States of America*. 110 (49): 19754–9, 2013), LY294002 (*ACS Chemical Biology*. 9 (2): 495–502, 2014), AZD5153 (*Journal of Medicinal Chemistry*. 59 (17): 7801–17, 2016), MT-1 (*Nature Chemical Biology*. 12 (12): 1089–1096 2016) or MS645 (*Proceedings of the National Academy of Sciences of the United States of America*. 115 (31): 7949–7954, 2018).

[00238] In certain aspects of all embodiments, the further therapeutic agent is a histone methyltransferase inhibitor. In some of these embodiments, the further therapeutic agent comprises a DOT1-like histone methyltransferase (DOT1L) inhibitor. DOT1L is a histone methyltransferase enzyme that targets lysine 79 in the globular domain of histone H3 for mono-, di-, or trimethylation. In some of these embodiments, the further therapeutic agent is EPZ004777, EPZ-5676 (*Blood*. 2013 Aug 8;122(6):1017-25) or SGC0946 (*Nat. Commun*. 2012;3:1288), for example, EPZ-5676.

[00239] In certain aspects of all embodiments, the further therapeutic agent is a histone deacetylase (HDAC) inhibitor. HDAC proteins may be grouped into classes based on homology to yeast HDAC proteins with Class I made up of HDAC1, HDAC2, HDAC3 and HDAC 8; Class IIa made up of HDAC4, HDAC5, HDAC7 and HDAC 9; Class IIb made up of HDAC6 and HDAC10; and Class IV made up of HDAC11. In some of these embodiments, the further therapeutic agent is trichostatin A, vorinostat (*Proc. Natl. Acad. Sci. U.S.A.* 1998 Mar 17;95(6):3003-7), givinostat, abexinostat (*Mol. Cancer Ther.* 2006 May;5(5):1309-17), belinostat (*Mol. Cancer Ther.* 2003 Aug;2(8):721-8), panobinostat (*Clin. Cancer Res.* 2006 Aug 1;12(15):4628-35), resminostat (*Clin. Cancer Res.* 2013 Oct 1;19(19):5494-504), quisinostat (*Clin. Cancer Res.* 2013 Aug 1;19(15):4262-72), depsipeptide (*Blood*. 2001 Nov 1;98(9):2865-8), entinostat (*Proc. Natl. Acad. Sci. U.S.A.* 1999 Apr 13;96(8):4592-7), mocetinostat (*Bioorg. Med. Chem. Lett.* 2008 Feb 1;18(3):1067-71) or valproic acid (*EMBO J.* 2001 Dec 17;20(24):6969-78). For example, in some embodiments, the further therapeutic agent is panobinostat. In other embodiments, the further therapeutic agent is panobinostat or SAHA.

[00240] In certain aspects of all embodiments, the further therapeutic agent is a histone demethylase inhibitor. In particular embodiments, the histone demethylase inhibitor is a lysine-specific demethylase 1A (Lsd1) inhibitor. In some of these embodiments, the further therapeutic agent is HCI-2509 (*BMC Cancer*. 2014 Oct

9;14:752), tranilcypromine or ORY-1001 (J. Clin. Oncol 31, 2013 (suppl; abstr e13543). In other embodiments, the further therapeutic agent is HCI-2509.

[00241] In certain aspects of all embodiments, the further therapeutic agent is a MLL-menin inhibitor. Menin is a co-factor of the oncogenic MLL fusion protein, and an MLL-menin inhibitor blocks the interaction of the two proteins. Examples of MLL-menin inhibitors include MI-453, M-525, and MI-503.

[00242] In certain aspects of all embodiments, the further therapeutic agent is a B-cell receptor signaling antagonist (*e.g.*, a Bruton's tyrosine kinase (BTK) inhibitor, such as ibrutinib).

[00243] In certain aspects of all embodiments, the further therapeutic agents is an immunomodulator. Immunomodulators of particular interest for use in combination with compounds of the present disclosure include: afutuzumab (available from ROCHE®); pegfilgrastim (NEULASTA®); lenalidomide (CC-5013, REVLIMID®); thalidomide (THALOMID®); actimid (CC4047); and IRX-2 (mixture of human cytokines including interleukin 1, interleukin 2, and interferon γ , CAS 951209-71-5, available from IRX Therapeutics).

[00244] In certain aspects of all embodiments, the further therapeutic agent comprises a chimeric antigen receptor T-cell (CAR-T) therapy. CAR-T therapies of particular interest for use in combination with compounds of the present disclosure include: tisagenlecleucel (Novartis), axicabtagene ciloleucel (Kite), and tocilizumab and atlizumab (Roche).

[00245] In certain aspects of all embodiments, the further therapeutic agent is an immune checkpoint inhibitor (*e.g.*, a PD-1 inhibitor, such as pembrolizumab or nivolumab; a PD-L1 inhibitor, such as atezolizumab, avelumab, or durvalumab; a CTLA-4 inhibitor; a LAG-3 inhibitor; or a Tim-3 inhibitor). Other immune checkpoint inhibitors of interest for use in combination with compounds of the present disclosure include: PD-1 inhibitors, such as pembrolizumab (KEYTRUDA®), nivolumab (OPDIVO®), cemiplimab (LIBTAYO®), spartalizumab (PDR001), pidilizumab (CureTech), MEDI0680 (Medimmune), cemiplimab (REGN2810), dostarlimab (TSR-042), PF-06801591 (Pfizer), tislelizumab (BGB-A317), camrelizumab (INCSHR1210, SHR-1210), and AMP-224 (Amplimmune); PD-L1 inhibitors, such as atezolizumab (TECENTRIQ®), avelumab (BAVENCIO®), durvalumab (IMFINZI®), FAZ053

(Novartis), and BMS-936559 (Bristol-Myers Squibb); and drugs that target CTLA-4, such as ipilimumab (YERVOY®).

[00246] In various embodiments, the immune checkpoint inhibitor is a PD-1 inhibitor. In specific embodiments, the PD-1 inhibitor is pembrolizumab, nivolumab, or a combination thereof. In particular embodiments, the PD-1 inhibitor is pembrolizumab (also known as lambrolizumab, MK-3475, MK03475, SCH-900475, or KEYTRUDA®). Pembrolizumab and other anti-PD-1 antibodies are disclosed in Hamid, O., *et al.* (2013) *New England Journal of Medicine* 369 (2): 134–44, US 8,354,509, and WO 2009/114335, incorporated by reference in their entireties. In particular embodiments, the PD-1 inhibitor is nivolumab (also known as MDX-1106, MDX-1106-04, ONO-4538, BMS-936558, or OPDIVO®). Nivolumab (clone 5C4) and other anti-PD-1 antibodies are disclosed in US 8,008,449 and WO 2006/121168, incorporated by reference in their entireties. In some other embodiments, the PD-1 inhibitor is AMP-224 (Amplimmune), CBT-501 (CBT Pharmaceuticals), CBT-502 (CBT Pharmaceuticals), JS001 (Junshi Biosciences), IBI308 (Innovent Biologics), INCSHR1210 (Incyte), also known as SHR-1210 (Hengrui Medicine), BGBA317 (Beigene), BGB-108 (Beigene), BAT-I306 (Bio-Thera Solutions), GLS-010 (Gloria Pharmaceuticals; WuXi Biologics), AK103, AK104, AK105 (Akesio Biopharma; Hangzhou Hansi Biologics; Hanzhong Biologics), LZM009 (Livzon), HLX-10 (Henlius Biotech), MEDI0680 (Medimmune), PDF001 (Novartis), PF-06801591 (Pfizer), pidilizumab (CureTech), REGN2810 (Regeneron), TSR-042 (Tesar), also known as ANB011, or CS1003 (CStone Pharmaceuticals). MEDI0680 (Medimmune) is also known as AMP-514. MEDI0680 and other anti-PD-1 antibodies are disclosed in US 9,205,148 and WO 2012/145493, incorporated by reference in their entireties. Pidilizumab is also known as CT-011. Pidilizumab and other anti-PD-1 antibodies are disclosed in Rosenblatt, J., *et al.* (2011) *J Immunotherapy* 34(5): 409-18, US 7,695,715, US 7,332,582, and US 8,686,119, incorporated by reference in their entireties.

[00247] In one embodiment, the anti-PD-1 antibody molecule is cemiplimab. In one embodiment, the anti-PD-1 antibody molecule is sintilimab. In one embodiment, the anti-PD-1 antibody molecule is toripalimab. In one embodiment, the anti-PD-1 antibody molecule is camrelizumab.

[00248] Further known anti-PD-1 antibody molecules include those described, *e.g.*, in WO 2015/112800, WO 2016/092419, WO 2015/085847, WO 2014/179664, WO 2014/194302, WO 2014/209804, WO 2015/200119, US 8,735,553, US 7,488,802, US 8,927,697, US 8,993,731, and US 9,102,727, incorporated by reference in their entireties.

[00249] In one embodiment, the PD-1 inhibitor is an anti-PD-1 antibody molecule as described in US 2015/0210769. In one embodiment, the anti-PD-1 antibody molecule comprises the CDRs, variable regions, heavy chains and/or light chains of BAP049-Clone-E or BAP049-Clone-B disclosed in US 2015/0210769. The antibody molecules described herein can be made by vectors, host cells, and methods described in US 2015/0210769, incorporated by reference in its entirety.

[00250] In one embodiment, the PD-1 inhibitor is a peptide that inhibits the PD-1 signaling pathway, *e.g.*, as described in US 8,907,053, incorporated by reference in its entirety. In one embodiment, the PD-1 inhibitor is an immunoadhesin (*e.g.*, an immunoadhesin comprising an extracellular or PD-1 binding portion of PD-L1 or PD-L2 fused to a constant region (*e.g.*, an Fc region of an immunoglobulin sequence). In one embodiment, the PD-1 inhibitor is AMP-224 (B7-DCIg (Amplimmune), *e.g.*, disclosed in WO 2010/027827 and WO 2011/066342, incorporated by reference in their entireties).

[00251] In some embodiments, the immune checkpoint inhibitor is a PD-L1 inhibitor. In some such embodiments, the PD-L1 inhibitor is atezolizumab, avelumab, durvalumab, or a combination thereof. In particular embodiments, the PD-L1 inhibitor is atezolizumab, also known as MPDL3280A, RG7446, RO5541267, YW243.55.S70, or TECENTRIQ™. Atezolizumab and other anti-PD-L1 antibodies are disclosed in US 8,217,149, incorporated by reference in its entirety. In particular embodiments, the PD-L1 inhibitor is avelumab, also known as MSB0010718C. Avelumab and other anti-PD-L1 antibodies are disclosed in WO 2013/079174, incorporated by reference in its entirety. In particular embodiments, the PD-L1 inhibitor is durvalumab, also known as MEDI4736. Durvalumab and other anti-PD-L1 antibodies are disclosed in US 8,779,108, incorporated by reference in its entirety. In certain embodiments, the PD-L1 inhibitor is KN035 (Alphamab; 3DMed), BMS 936559 (Bristol-Myers Squibb), CS1001 (CStone Pharmaceuticals), FAZ053 (Novartis), SHR-1316 (Hengrui

Medicine), TQB2450 (Chiatai Tianqing), STI-A1014 (Zhaoke Pharm; Lee's Pharm), BGB-A333 (Beigene), MSB2311 (Mabspace Biosciences), or HLX-20 (Henlius Biotech). In one embodiment, the anti-PD-L1 antibody molecule is BMS-936559 (Bristol-Myers Squibb), also known as MDX-1105 or 12A4. BMS-936559 and other anti-PD-L1 antibodies are disclosed in US 7,943,743 and WO 2015/081158, incorporated by reference in their entireties. In some embodiments, the PD-L1 inhibitor is a monoclonal antibody (*e.g.*, as made by Hisun Pharm and applying for clinical trials).

[00252] In one embodiment, the PD-L1 inhibitor is an anti-PD-L1 antibody molecule. In one embodiment, the PD-L1 inhibitor is an anti-PD-L1 antibody molecule as disclosed in US 2016/0108123, incorporated by reference in its entirety. In one embodiment, the anti-PD-L1 antibody molecule comprises the CDRs, variable regions, heavy chains and/or light chains of BAP058-Clone O or BAP058-Clone N disclosed in US 2016/0108123.

[00253] Further known anti-PD-L1 antibodies include those described, *e.g.*, in WO 2015/181342, WO 2014/100079, WO 2016/000619, WO 2014/022758, WO 2014/055897, WO 2015/061668, WO 2013/079174, WO 2012/145493, WO 2015/112805, WO 2015/109124, WO 2015/195163, US 8,168,179, US 8,552,154, US 8,460,927, and US 9,175,082, incorporated by reference in their entireties.

[00254] In some embodiments, the immune checkpoint inhibitor is a CTLA-4 inhibitor. In certain embodiments, the CTLA-4 inhibitor is ipilimumab. In other embodiments, the CTLA4 inhibitor is tremelimumab.

[00255] In some embodiments, the immune checkpoint inhibitor is a LAG-3 inhibitor. In some embodiments, the LAG-3 inhibitor is chosen from LAG525 (Novartis), BMS-986016 (Bristol-Myers Squibb), or TSR-033 (Tesaro). In one embodiment, the LAG-3 inhibitor is an anti-LAG-3 antibody molecule. In one embodiment, the LAG-3 inhibitor is an anti-LAG-3 antibody molecule as disclosed in US 2015/0259420, incorporated by reference in its entirety. In one embodiment, the anti-LAG-3 antibody molecule comprises the CDRs, variable regions, heavy chains and/or light chains of BAP050-Clone I or BAP050-Clone J disclosed in US 2015/0259420.

[00256] In one embodiment, the anti-LAG-3 antibody molecule is BMS-986016 (Bristol-Myers Squibb), also known as BMS986016. BMS-986016 and other anti-LAG-3 antibodies are disclosed in WO 2015/116539 and US 9,505,839, incorporated by reference in their entireties. In one embodiment, the anti-LAG-3 antibody molecule is TSR-033 (Tesaro). In one embodiment, the anti-LAG-3 antibody molecule is IMP731 or GSK2831781 (GSK and Prima BioMed). IMP731 and other anti-LAG-3 antibodies are disclosed in WO 2008/132601 and US 9,244,059, incorporated by reference in their entireties. In one embodiment, the anti-LAG-3 antibody molecule is IMP761 (Prima BioMed).

[00257] Further known anti-LAG-3 antibodies include those described, *e.g.*, in WO 2008/132601, WO 2010/019570, WO 2014/140180, WO 2015/116539, WO 2015/200119, WO 2016/028672, US 9,244,059, US 9,505,839, incorporated by reference in their entireties.

[00258] In one embodiment, the anti-LAG-3 inhibitor is a soluble LAG-3 protein, *e.g.*, IMP321 (Prima BioMed), *e.g.*, as disclosed in WO 2009/044273, incorporated by reference in its entirety.

[00259] In some embodiments, the immune checkpoint inhibitor is a TIM-3 inhibitor. In some embodiments, the TIM-3 inhibitor is MGB453 (Novartis) or TSR-022 (Tesaro).

[00260] In one embodiment, the TIM-3 inhibitor is an anti-TIM-3 antibody molecule. In one embodiment, the TIM-3 inhibitor is an anti-TIM-3 antibody molecule as disclosed in US 2015/0218274, incorporated by reference in its entirety. In one embodiment, the anti-TIM-3 antibody molecule comprises the CDRs, variable regions, heavy chains and/or light chains of ABTIM3-hum11 or ABTIM3-hum03 disclosed in US 2015/0218274.

[00261] In one embodiment, the anti-TIM-3 antibody molecule is TSR-022 (AnaptysBio/Tesaro). In one embodiment, the anti-TIM-3 antibody molecule comprises one or more of the CDR sequences (or collectively all of the CDR sequences), the heavy chain or light chain variable region sequence, or the heavy chain or light chain sequence of APE5137 or APE5121. APE5137, APE5121, and other anti-TIM-3 antibodies are disclosed in WO 2016/161270, incorporated by reference in its entirety. In one embodiment, the anti-TIM-3 antibody molecule is the antibody clone F38-2E2.

[00262] Further known anti-TIM-3 antibodies include those described, *e.g.*, in WO 2016/111947, WO 2016/071448, WO 2016/144803, US 8,552,156, US 8,841,418, and US 9,163,087, incorporated by reference in their entireties.

[00263] In an effort to protect normal cells from treatment toxicity and to limit organ toxicities, cytoprotective agents (such as neuroprotectants, free-radical scavengers, cardioprotectors, anthracycline extravasation neutralizers, nutrients and the like) may be used as an adjunct therapy in combination with compounds of the present disclosure. Suitable cytoprotective agents include amifostine (ETHYOL®), glutamine, dimesna (TAVOCEPT®), mesna (MESNEX®), dexrazoxane (ZINECARD® or TOTECT®), xaliproden (XAPRILA®), and leucovorin (also known as calcium leucovorin, citrovorum factor and folinic acid).


[00264] Some patients may experience allergic reactions to compounds of the present disclosure and/or other therapeutic agent(s) (*e.g.*, anti-cancer agent(s)) during or after administration. Therefore, anti-allergic agents can be administered in combination with compounds of the present disclosure and/or other therapeutic agent(s) (*e.g.*, anti-cancer agent(s)) to minimize the risk of an allergic reaction. Suitable anti-allergic agents include corticosteroids (Knutson, S., *et al.*, *PLoS One*, DOI:10.1371/journal.pone.0111840 (2014)), such as dexamethasone (*e.g.*, DECADRON®), beclomethasone (*e.g.*, BECLOVENT®), hydrocortisone (also known as cortisone, hydrocortisone sodium succinate, hydrocortisone sodium phosphate, sold under the tradenames ALA-CORT®, hydrocortisone phosphate, SOLU-CORTEF®, HYDROCORT ACETATE® and LANACORT®), prednisolone (sold under the tradenames DELTA-CORTEL®, ORAPRED®, PEDIAPRED® and PRELONE®), prednisone (sold under the tradenames DELTASONE®, LIQUID RED®, METICORTEN® and ORASONE®), methylprednisolone (also known as 6-methylprednisolone, methylprednisolone acetate, methylprednisolone sodium succinate, sold under the tradenames DURALONE®, MEDRALONE®, MEDROL®, M-PREDNISOL® and SOLU-MEDROL®); antihistamines, such as diphenhydramine (*e.g.*, BENADRYL®), hydroxyzine, and cyproheptadine; and bronchodilators, such as the beta-adrenergic receptor agonists, albuterol (*e.g.*, PROVENTIL®), and terbutaline (BRETHINE®).

[00265] Some patients may experience nausea during and after administration of the compounds described herein and/or other therapeutic agent(s) (*e.g.*, anti-cancer agent(s)). Therefore, anti-emetics can be used in combination with compounds of the present disclosure and/or other therapeutic agent(s) (*e.g.*, anti-cancer agent(s)) to prevent nausea (upper stomach) and vomiting. Suitable anti-emetics include aprepitant (EMEND®), ondansetron (ZOFTRAN®), granisetron HCl (KYTRIL®), lorazepam (ATIVAN®), dexamethasone (DECADRON®), prochlorperazine (COMPazine®), casopitant (REZONIC® and ZUNRISA®), and combinations thereof.

[00266] Medication to alleviate the pain experienced during treatment is often prescribed to make the patient more comfortable. Common over-the-counter analgesics, such as TYLENOL®, can also be used in combination with compounds of the present disclosure and/or other therapeutic agent(s) (*e.g.*, anti-cancer agent(s)). Opioid analgesic drugs such as hydrocodone/paracetamol or hydrocodone/acetaminophen (*e.g.*, VICODIN®), morphine (*e.g.*, ASTRAMORPH® or AVINZA®), oxycodone (*e.g.*, OXYCONTIN® or PERCOCET®), oxymorphone hydrochloride (OPANA®), and fentanyl (*e.g.*, DURAGESIC®) can be useful for moderate or severe pain, and can be used in combination with compounds of the present disclosure and/or other therapeutic agent(s) (*e.g.*, anti-cancer agent(s)).

EMBODIMENTS

[00267] Embodiment 1. A polymorph of a compound having the following structure (I):



(I)

or a tautomer or zwitterionic form thereof, the polymorph having an X-ray powder diffraction pattern comprising the following:

D space (Å):

18.3 ± 0.09

8.1 ± 0.06

6.4 ± 0.08

5.9 ± 0.06

4.4 ± 0.05

expressed in terms of "D" spacing.

[00268] Embodiment 2. The polymorph of Embodiment 1, wherein the X-ray powder diffraction pattern comprises the following:

D space (Å):

18.38 ± 0.003

8.15 ± 0.008

6.47 ± 0.002

5.95 ± 0.007

4.44 ± 0.006

expressed in terms of "D" spacing.

[00269] Embodiment 3. The polymorph of Embodiment 1, wherein the X-ray powder diffraction pattern comprises the following:

D space (Å):

18.382

8.157


6.472

5.956

4.445

expressed in terms of "D" spacing.

[00270] Embodiment 4. A polymorph of a compound having the following structure (I):



(I)

or a tautomer or zwitterion thereof, the polymorph being a crystalized form having a monoclinic space group $P2_1$ with lattice parameters of:

$$a = 6.46(1) \text{ \AA};$$


$$b = 9.07(2) \text{ \AA};$$

$$c = 18.25(4) \text{ \AA}; \text{ and}$$

$$\beta = 95.457(2)^\circ;$$

and a volume of $1066.11(4) \text{ \AA}^3$.


[00271] Embodiment 5. A polymorph of a compound having the following structure (I):



(I)

or a tautomer or zwitterionic form thereof, wherein the polymorph has the X-ray diffraction pattern set forth in Figure 1.


[00272] Embodiment 6. A polymorph of a compound having the following structure (I):



(I)

or a tautomer or zwitterionic form thereof, wherein the polymorph is Form B.

[00273] Embodiment 7. A polymorph of a compound having the following



structure (I):

(I)

or a tautomer or zwitterionic form thereof wherein the polymorph is formed by a method comprising:

a) contacting an amorphous compound of structure (I) with a lattice forming reagent; and

b) treating the product of step a) with a solvent having water content less than about 0.05 % v/v and removing the solvent, thereby forming the polymorph.


[00274] Embodiment 8. The polymorph of Embodiment 7, wherein the lattice forming reagent is maleic acid.

[00275] Embodiment 9. The polymorph of Embodiment 8, wherein the maleic acid is added at a ratio of 1:1 ± 0.2 of amorphous compound to maleic acid.

[00276] Embodiment 10. The polymorph of any one of Embodiments 7-9, wherein the solvent is ethanol or tetrahydrofuran.

[00277] Embodiment 11. The polymorph of any one of Embodiments 7-10, wherein the contacting further comprises suspending the amorphous compound of structure (I) in aqueous tetrahydrofuran.

[00278] Embodiment 12. A polymorph of a compound having the following structure (I):



(I)


or a tautomer or zwitterionic form thereof, wherein the polymorph has an initial purity of at least 99.5% and a subsequent purity of at least 99.5% after being stored from about 12 hours up to about 7 days at about $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at a relative humidity of 60%.

[00279] Embodiment 13. The polymorph of Embodiment 12, wherein the subsequent purity is at least 99.5% after being stored for greater than about 7 days at about $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at a relative humidity of 60%.

[00280] Embodiment 14. The polymorph of Embodiment 12 or 13, wherein the subsequent purity is at least 99.5% after being stored for greater than about 30 days at about $25^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at a relative humidity of 60%.

[00281] Embodiment 15. The polymorph of any one of Embodiments 12-14, wherein the initial purity and subsequent purity are as determined by HPLC.

[00282] Embodiment 16. A polymorph of a compound having the following structure (I):




(I)

or a tautomer or zwitterionic form thereof, wherein the polymorph has an endotherm peak value at about $256^{\circ}\text{C} - 268^{\circ}\text{C}$ as determined by differential scanning calorimetry.

[00283] Embodiment 17. The polymorph of Embodiment 16, wherein the endotherm peak value is at about $257.0^{\circ}\text{C} - 266.0^{\circ}\text{C}$.

[00284] Embodiment 18. The polymorph of Embodiment 16 or 17, wherein the endotherm peak value is at about $258.0^{\circ}\text{C} - 265.0$.

[00285] Embodiment 19. The polymorph of any one of claims 1-18, wherein the polymorph comprises a zwitterionic form having the following structure (II):



(II)

[00286] Embodiment 20. A pharmaceutical composition comprising the polymorph of any one of Embodiments 1-19 and a pharmaceutically acceptable carrier or excipient.

[00287] Embodiment 21. The pharmaceutical composition of Embodiment 20, wherein the pharmaceutical composition is formulated for oral administration.

[00288] Embodiment 22. The pharmaceutical composition of Embodiment 20 or 21, wherein the pharmaceutical composition comprises the polymorph in a concentration ranging from about 0.5 wt.% to about 11.25 wt.%.

[00289] Embodiment 23. The pharmaceutical composition of any one of Embodiments 20-22, wherein the pharmaceutical composition comprises the polymorph in a concentration ranging from about 0.6 wt.% to about 11.05 wt.%.

[00290] Embodiment 24. The pharmaceutical composition of any one of Embodiments 20-23, wherein the pharmaceutical composition comprises about 0.6 wt.% of the polymorph.

[00291] Embodiment 25. The pharmaceutical composition of any one of Embodiments 20-23, wherein the pharmaceutical composition comprises about 2.5 wt.% of the polymorph.

[00292] Embodiment 26. The pharmaceutical composition of any one of Embodiments 20-23, wherein the pharmaceutical composition comprises about 11.0 wt.% of the polymorph.

[00293] Embodiment 27. The pharmaceutical composition of any one of Embodiments 20-26, wherein the pharmaceutical composition comprises about 1 milligram (mg) of the polymorph.

[00294] Embodiment 28. The pharmaceutical composition of any one of Embodiments 20-26, wherein the pharmaceutical composition comprises about 4 mg of the polymorph.

[00295] Embodiment 29. The pharmaceutical composition of any one of Embodiments 20-26, wherein the pharmaceutical composition comprises about 16 mg of the polymorph.

[00296] Embodiment 30. The pharmaceutical composition of any one of Embodiments 20-29, wherein the excipient is anhydrous lactose or microcrystalline cellulose.

[00297] Embodiment 31. The pharmaceutical composition of any one of Embodiments 20-30, wherein the excipient is anhydrous lactose.

[00298] Embodiment 32. The pharmaceutical composition of any one of Embodiments 20-30, wherein the excipient is microcrystalline cellulose.

[00299] Embodiment 33. The pharmaceutical composition of any one of Embodiments 20-32, wherein the pharmaceutical composition comprises the excipient in a concentration ranging from about 85 wt.% to about 98 wt.%.

[00300] Embodiment 34. The pharmaceutical composition of any one of Embodiments 20-33, wherein the pharmaceutical composition comprises the excipient in a concentration ranging from about 87 wt.% to about 97.5 wt.%.

[00301] Embodiment 35. The pharmaceutical composition of any one of Embodiments 20-34, wherein the pharmaceutical composition comprises about 87 wt.% of the excipient.

[00302] Embodiment 36. The pharmaceutical composition of any one of Embodiments 20-34, wherein the pharmaceutical composition comprises about 95.5 wt.% of the excipient.

[00303] Embodiment 37. The pharmaceutical composition of any one of Embodiments 20-34, wherein the pharmaceutical composition comprises about 97.5 wt.% of the excipient.

[00304] Embodiment 38. The pharmaceutical composition of any one of Embodiments 20-37, further comprising colloidal silicon dioxide.

[00305] Embodiment 39. The pharmaceutical composition of Embodiment 38, wherein the pharmaceutical composition comprises about 1 wt.% of the colloidal silicon dioxide.

[00306] Embodiment 40. The pharmaceutical composition of any one of Embodiments 20-39 further comprising magnesium stearate.

[00307] Embodiment 41. The pharmaceutical composition of Embodiment 40, wherein the pharmaceutical composition comprises about 1 wt.% of the magnesium stearate.

[00308] Embodiment 42. The pharmaceutical composition of any one of Embodiments 20-41, in the form of a capsule for oral administration.


[00309] Embodiment 43. A pharmaceutical composition comprising a polymorph of any one of Embodiments 1-19 having water content less than 0.50 % as determined by Karl Fischer titration.

[00310] Embodiment 44. A unit dose form comprising a pharmaceutical composition of any one of Embodiments 20-43.

[00311] Embodiment 45. A method for preparing a polymorph, the method comprising:

a) contacting an amorphous compound having the following structure

(I):



(I)

with a lattice forming reagent; and

b) treating the product of step a) with solvent having water content less than about 0.05 % v/v; and removing the solvent, thereby forming the polymorph.

[00312] Embodiment 46. The method of Embodiment 45, wherein the lattice forming reagent is an acid.

[00313] Embodiment 47. The method of Embodiment 46, wherein the acid is selected from the group consisting of maleic acid, fumaric acid, L-tartaric acid, hippuric acid, nicotinic acid, acetic acid, and combinations thereof.


[00314] Embodiment 48. The method of any one of Embodiments 45-47, wherein the lattice forming reagent is maleic acid.

[00315] Embodiment 49. The method of Embodiment 45, wherein the lattice forming reagent is aspartame.

[00316] Embodiment 50. The method of Embodiment 45, wherein the lattice forming reagent is D-xylose.


[00317] Embodiment 51. The method of any one of Embodiments 45-50, wherein the removing the solvent is at a pressure less than about 1 atm and a temperature greater than about 35°C.

[00318] Embodiment 52. The method of any one of Embodiments 45-51, wherein the method further comprises preparing a compound of structure (III):




(III)

by reacting a compound having the following structure:




or a salt thereof, with a base and a compound having the following structure:



[00319] Embodiment 53. The method of Embodiment 52, wherein the base is an amine base.

[00320] Embodiment 54. The method of Embodiment 52 or 53, wherein the base is triethylamine or diisopropylethyl amine.

[00321] Embodiment 55. The method of any one of Embodiments 52-54, wherein the method further comprises preparing the amorphous compound of structure (I):



(I)

by reacting the compound of structure (III), or a salt thereof, with a phosphoester-cleaving compound and treating with a basic solution.

[00322] Embodiment 56. The method of Embodiment 55, wherein the phosphoester-cleaving compound is trimethyl silyl bromide (TMSBr), and the basic solution comprises ammonium bicarbonate.

[00323] Embodiment 57. The method of any one of Embodiments 45-56, wherein the polymorph is a polymorph according to any one of Embodiments 1-19.

[00324] Embodiment 58. A polymorph prepared according to the method of any one of Embodiments 45-57.

[00325] Embodiment 59. A method for treating a disease associated with overexpression of a cyclin-dependent kinase (CDK) in a mammal in need thereof, the method comprising administering a therapeutically effective amount of the polymorph of any one of Embodiments 1-19 or 58, the pharmaceutical composition of any one of Embodiments 20-43, or the unit dose form of Embodiment 43 to the mammal.

[00326] Embodiment 60. The method of Embodiment 59, wherein the disease is cancer.

[00327] Embodiment 61. The method of Embodiment 60, wherein the cancer is a hematologic cancer.

[00328] Embodiment 62. The method of Embodiment 60, wherein the hematologic cancer is selected from acute myelogenous leukemia (AML), follicular lymphoma, acute lymphoblastic leukemia (ALL), chronic lymphocytic leukemia (CLL), multiple myeloma (MM), and non-Hodgkin's lymphoma.

[00329] Embodiment 63. The method of Embodiment 62, wherein the hematological cancer is acute myelogenous leukemia (AML).

[00330] Embodiment 64. The method of Embodiment 62, wherein the hematologic cancer is chronic lymphocytic leukemia (CLL).

[00331] Embodiment 65. The method of Embodiment 62, wherein the hematologic cancer is myelodysplastic syndrome (MDS).

[00332] Embodiment 66. The method of Embodiment 62, wherein the hematologic cancer is multiple myeloma (MM).

[00333] Embodiment 67. The method of Embodiment 60, wherein the cancer comprises a solid tumor.

[00334] Embodiment 68. The method of Embodiment 60 or 67, wherein the cancer is bladder cancer.

[00335] Embodiment 69. The method of Embodiment 60 or 67, wherein the cancer is lung cancer.

[00336] Embodiment 70. The method of Embodiment 60 or 67, wherein the cancer is liver cancer.

[00337] Embodiment 71. The method of Embodiment 60 or 67, wherein the cancer is prostate cancer.

[00338] Embodiment 72. The method of any one of Embodiments 60-71, wherein the polymorph, the pharmaceutical composition, or the unit dose form is administered orally.

[00339] The examples and preparations provided below further illustrate and exemplify polymorph of a compound of structure (I), and methods of preparing the same. It is to be understood that the scope of the present invention is not limited in any way by the scope of the following examples and preparations. In the following examples, and throughout the specification and claims, molecules with a single stereocenter, unless otherwise noted, exist as a racemic mixture. Those molecules with two or more stereocenters, unless otherwise noted, exist as a racemic mixture of diastereomers. Single enantiomers/diastereomers may be obtained by methods known to those skilled in the art.

EXAMPLES

[00340] Data was acquired according to the parameters listed below:

X-ray powder diffraction (XRPD):

[00341] Stoe Stadi P. Copper K α I radiation, 40kV / 40mA; Mythen 1K detector transmission mode, curved monochromator, 0.02°2 θ step size, 12 s step time, 1.5-50.5°2 θ scanning range with 1°2 θ detector step in step-scan mode. Each sample (25-40 mg of powder) was placed between two cellulose acetate foils spaced with a metal washer (0.4 mm thick, 12-mm inner diameter; "sandwich element"). The sandwich element was transferred to a sample holder (SCell) that was sealed with acetate foils. Samples were acquired in ambient air atmosphere and rotated during measurements.

TG-FTIR:

[00342] Netzsch Thermo-Microbalance TG 209 with Bruker FT-IR Spectrometer IFS28 or Vector 22; Al crucible with microhole, N₂ atmosphere, 10 K/min heating rate, 25°C to 300°C (or 350°C range).

HPLC:

[00343] The method used to detect and determine purity of compound of structure (I) and related substances (such as alvocidib) was a reverse-phase HPLC method with a gradient program and DAD detection technique. Reverse phase C₁₈ Waters X-bridge 150 mm × 4.6 mm, 3.5- μ m particle column; flow rate = 1.0 mL/min; detection wavelength = 265 nm; run time: 35.0 minutes; sample diluted in methanol; mobile phase A was 80:20 (v/v) pH 6.5 buffered aqueous acetonitrile; mobile phase B was 35:65 (v/v) pH 6.5 buffered aqueous acetonitrile; 1.0 mL/min; column temperature = 35°C. The gradient program is depicted in the following table:

Run Time (minutes)	Mobile Phase A (%)	Mobile Phase B (%)
0	100	0
8	100	0
16	66	34
24	0	100
26	0	100
26.5	100	0
35	100	0

[00344] Total Impurities (%) was calculated by summing the percentages of each individual impurity, including alvocidib. Other Impurities (%) was calculated by summing the percentages of each individual impurity, excluding alvocidib. Purity of compound of structure (I) (%) was calculated by taking the difference between 100%

and the Total Impurities (%). All individual impurities at and above 0.05% were taken for the calculation of total impurities.

^1H NMR:


[00345] Bruker DPX 300 using a frequency of 300.13 MHz, a 30° excitation pulse and a recycle delay of 1 s. 16-1024 scans were accumulated per spectrum; deuterated DMSO or D_2O was used as a solvent. Two-dimensional COSY spectra were acquired with 512 data points in the indirect dimension, an indirect time increment of 441.60 μs , 16 scans per slice and a recycle delay of 0.36 s.

Differential Scanning Calorimetry (DSC):

[00346] DSC was performed using a TA Q200/Q2000DSC from TA Instruments using a ramp method and a crimped, aluminum sample pan at 25°C . The heating rate was $10^\circ\text{C}/\text{minute}$, and the purge gas was nitrogen.

EXAMPLE 1

POLYMORPH SYNTHESIS



Scheme 1.

[00347] **Step 1:** A-1 was treated with boron tribromide in chlorobenzene. Removal of byproducts by distillation and crystallization from chlorobenzene-methanol-water resulted in A-2 as free base.

[00348] **Step 2:** A-2 (in free base form) was treated with diethyl chlorophosphate and diisopropylethylamine in *N*-methylpyrrolidone. Water was added to stop the reaction and precipitate the product. The resulting slurry was filtered, washed with water, and dried under vacuum to produce the compound A-3.

[00349] **Step 3:** A-3 was treated with trimethylsilyl bromide to deprotect A-3 and afford A-4 as a hydrobromide, which was treated with ammonium bicarbonate solution. The precipitated A-4 (*i.e.*, compound of structure (I) having an amorphous crystal structure (*i.e.*, "Form A") was filtered and dried.

[00350] **Step 4:** A-4 was suspended in a mixture of THF and water (19:1), and maleic acid was added. After stirring at room temperature, the solid was filtered and dried in a filter dryer to afford A-5. The resultant filtered solid compound A-5 was suspended in ethanol and re-slurried before an additional filtration. Filtered product (*i.e.*, A-5 as polymorph form B) was washed with ethanol and dried to afford the desired product.

[00351] Synthesis of polymorph Form B according to Scheme 1 has been conducted using 5.72 kg A-1, 4.17 kg A-2 and 2.70 kg A-3. At this scale, the yield of Step 1 was 79.9%. After recrystallization, the yield of A-3 from Step 2 was 48%, and the purity of compound A-3 thus obtained was 86% by HPLC. The combined yield of Steps 3 and 4 was 37.5%. The overall yield of the process of Scheme 1 was 15%, and the process yielded 0.90 kg of A-5.

EXAMPLE 2

STABILITY STUDY

[00352] Initial attempts to isolate the compound of structure (I) produced an amorphous form, designated "Form A." Form A showed poor stability, decomposing to compound A-2. The results of a DVS (dynamic vapor sorption) test of Form A showed that the compound was hygroscopic, taking in 12.5% of its weight of water at 80% relative humidity. After 10 days at room temperature, Form A showed 6.85% degradation to compound A-2.

[00353] The decomposition exhibited a strong temperature dependence. Samples stored at -20 °C showed no degradant trends, but a stability test of Form A at 40 °C and 75% relative humidity for one week showed accelerated decomposition. The DSC and TGA plots of Form A exhibited the characteristics predicted by hydrolytic decomposition, as shown in FIG. 2.

EXAMPLE 3

SALT SCREENING STUDY

[00354] A total of eight lattice/salt forming reagents (5 acids and 3 bases) and four solvent systems were selected for salt screening. Screening conditions are outline below in Table 1:

Table 1: Salt screening conditions

Solvent Lattice forming reagents*	A	B	C	D
	IPA	Acetone	ACN	THF/H ₂ O (19:1)
Blank	Form A	Form A + Decomposition	Form A	Form A + B
H ₂ SO ₄	Amorphous	Amorphous	Sulfate Type A	Amorphous
H ₃ PO ₄	Phosphate Type A	Phosphate Type A	Phosphate Type A + Form A	Phosphate Type A
Maleic Acid	Form B	Form B	Form B	Form B
Fumaric Acid	Form B	Form B	Form B	Form A + B
L-tartaric acid	Form B	Form B	Form B	Form B
NaOH	Decomposition	Decomposition	Decomposition	Sodium Salt A**
K ₂ CO ₃	Decomposition	Decomposition	Decomposition	Decomposition
Ca(OH) ₂	Decomposition	Decomposition	Decomposition	Calcium Salt Type A**

* the molar ratio of freebase and lattice forming reagent is 1:1.

** purity decrease (5~14 area%) was observed from HPLC results.

[00355] When the compound of structure (I) was crystallized together with maleic acid or L-tartaric acid in aqueous THF, the resulting crystal showed identical polymorph to Form B, indicating structure (I) does not form a salt with those acids. Although some amount of acid will remain in the isolated crystal with above recrystallization process for obtaining Form B, it was discovered that residual maleic acid could be removed via a re-slurry in ethanol. In contrast, the tartaric acid did not readily wash from the compound via the ethanol slurry.

EXAMPLE 4
SALT SCREENING STUDY

[00356] Two larger scale pilot batches were synthesized according to the process described above in Example 1 to confirm and optimize the process. The results of these batches are shown in Tables 2, 3, and 4, below.

Table 2: Pilot Batches of Step 1

Batch No.	Compound A-1	Compound A-2 Yield (%)	HPLC purity
1a	165.0 g	114 g (80.1%)	97.61%
2a	94.0 g	70 g (86.43%)	98.86%

Table 3: Pilot Batches of Step 2

Batch No.	Compound A-2	Compound A-3 Yield (%)	HPLC purity
2a	100 g	82.4 g (67.1%)	98.67%
2a	350 g	320 g (62.8%)	97.97%

Table 4: Pilot batches of Step 3

Batch No.	Compound A-3	Compound A-4 Yield (%)	HPLC purity	Water Content (%)
3b	220 g	54.5 g (26.9%)	99.90%	0.48%
3b	110 g	24.8 g (25.3%)	99.75%	0.38%

[00357] After removing the maleic acid from the polymorph of Form B (A-5) via a re-slurry in ethanol, the resultant polymorph was dried by a flow of nitrogen on a filter to reduce water to below 0.5%. Re-slurrying in ethanol does not increase the concentration of impurity (A-2) in the slurry. Without wishing to be bound by theory, it is thought that the formation of compound A-2 is sensitive to water included in the crystal/polymorph form after formation. That is, when water content kept low enough

following polymorph formation, hydrolysis of the phosphate moiety is prevented but the lattice forming reagent (*e.g.*, maleic acid) can be removed.

EXAMPLE 5

STABILITY STUDY

[00358] A stability study of the polymorph of the compound of structure (I) (compound A-5) with low water content sample (0.38%) showed a pronounced improvement in the compound stability (Table 5), indicating improved stability of the polymorph having the lower water content.

Table 5: Room temperature stability Form B

Storage conditions		Ambient	-20 °C
HPLC area % Form B / Compound A-2	initial	99.75% / 0.13%	
	7 days	99.89% / 0.03%	-
	30 days	99.92% / 0.03%	99.93 / 0.02%

[00359] Based on this information, it can be concluded that the polymorph of the compound of structure (I) ("Form B") formed via a THF/water polymorph transformation with maleic acid, followed by ethanol re-slurry to remove the residual maleic acid and water from the polymorph provides an unexpectedly stable compound.

EXAMPLE 5

ADDITIONAL POLYMORPH FORMATION STUDIES – POWDER X-RAY DIFFRACTION

[00360] During preliminary salt co-crystal screening (Example 3, above), it became apparent that nearly identical polymorphs were being formed despite the use of different acids or co-crystal screening agents. This is illustrated in FIG. 3, which shows maleic and nicotinic acids along with D-xylose screening studies. This experiment provided evidence that these lattice forming reagents were not forming salts or co-crystals with the compound of structure (I), but were actually forming a polymorph (*i.e.*, Form B) having the XRPD diffractogram shown in FIG. 1.

EXAMPLE 6

SINGLE CRYSTAL X-RAY DIFFRACTION

[00361] The absolute stereochemistry, the position of the phosphoric acid moiety, as well as the zwitterionic nature of the polymorph of the compound of structure (I) (Form B) were determined by a single crystal X-ray diffraction. Single crystal X-ray diffraction of Form B is shown in FIG. 4. Form B crystallizes as an anhydrous molecule without solvent inclusion. Bond distances and angles are all within the expected values.

EXAMPLE 7

FORMATION OF POLYMORPH FORM C

[00362] Two experimental conditions were explored for forming a new solid form of a compound of structure (I). A solvent mixture of acetone/water and ethanol/water were added to Form A. The experimental condition including acetone/water as the solvent was successfully reproduced on a larger scale. The diffractograms of the resultant solid form ("Form C") are shown in FIG. 5. Although the diffractograms indicate the resultant product is highly crystalline, a TG-FTIR thermogram (FIG. 6A) indicated Form C was associated with a significant amount of water. As evidenced by the thermogram, some solvent is released below the boiling point of water, but a large percentage appears to be tightly bound, which indicates that the sample is highly hydrated.

[00363] In order to determine whether Form C was stable with respect to dehydration, a sample of Form C was dried under vacuum at 40 °C for about 24 hours. After drying, the sample contained about 5.4 weight percent water, which was close to the 5.3 weight percent theoretically expected for a sesquihydrate (*see* FIG. 6B). However, this sample lost significant crystallinity upon drying, so it is not necessarily a stoichiometric hydrate. Form C was found not to be stable to drying conditions with the resultant product being poorly crystalline.

[00364] The ¹H NMR spectra of the three different forms (A, B and C) are distinguishable. All samples were recorded in DMSO-*d*₆. The chemical shifts of the amorphous Form A and the Form C sample are substantially identical to each other but differ from those of Form B (FIG. 7). The position of the proton near 7 ppm is

markedly different for the spectrum of Form B, as is the presence of the peak near 12.8 ppm for Form B (*i.e.*, which is absent in Forms A and C).

EXAMPLE 8

POWDER X-RAY DIFFRACTION SPECTRUM

[00365] Polymorph Form B was analyzed using XRPD using the parameters described above. The resultant diffractogram is shown in FIG. 1. Tabulated data generated for Form B is shown in Table 6, below:

Table 6: Tabulated data from XRPD diffractogram of Form B

D value (Å)	2θ	Intensity (relative)	Intensity (absolute)	FWHM
18.382645	4.8032	55.17	3833	0.08
9.190754	9.6155	10.48	728	0.08
8.157735	10.8365	93.39	6489	0.06
6.471747	13.6717	49.51	3440	0.08
5.956605	14.8604	73.17	5084	0.04
5.524677	16.0296	11.07	769	0.06
5.281129	16.774	30.9	2147	0.06
5.172326	17.1295	9.19	639	0.06
5.080202	17.4425	17.69	1229	0.06
4.984581	17.7798	58	4030	0.04
4.722631	18.7747	18	1250	0.06
4.654062	19.0539	25.93	1802	0.04
4.552517	19.483	32.41	2252	0.06
4.445879	19.955	100	6948	0.06
4.144506	21.4226	19.68	1367	0.04
4.079254	21.7694	22.77	1582	0.06
3.90758	22.7383	9.77	679	0.04
3.867711	22.9759	28.61	1988	0.06
3.72577	23.8638	22.65	1573	0.06

3.6868	24.1198	25.68	1785	0.04
3.616616	24.5951	48.94	3401	0.06
3.512733	25.3344	7.74	538	0.06
3.394705	26.2307	28.77	1999	0.06
3.350985	26.5791	11.43	794	0.04
3.319773	26.8337	18.57	1290	0.06
3.241092	27.4977	23.54	1635	0.08
3.142734	28.376	15.33	1065	0.06
3.055085	29.208	12.7	882	0.08
2.973274	30.0303	10.91	758	0.14
2.920812	30.5827	6.32	439	0.06
2.830324	31.5856	9.24	642	0.08
2.748258	32.5546	6.82	474	0.04
2.731945	32.7544	11.76	817	0.06
2.703955	33.1032	6.09	423	0.06
2.64826	33.8201	10.18	707	0.04
2.640271	33.9255	14.93	1037	0.06
2.60744	34.3659	16.09	1118	0.06
2.587021	34.6457	11.27	783	0.06
2.540227	35.3046	5.6	389	0.06
2.424055	37.0566	5.24	364	0.06
2.216436	40.6738	7.03	488	0.06
2.125602	42.4942	10.93	760	0.06
2.078432	43.5072	4.98	346	0.14
2.042561	44.3113	12.61	876	0.1
2.008764	45.0975	5.76	400	0.08
1.950189	46.5303	5.84	406	0.14

EXAMPLE 9
FORMULATION STABILITY STUDY

[00366] Polymorph Form B (API) was formulated into two blends for four-week stability studies. Four excipients were tested in the following mixtures:

[00367] Binary mixtures (16 mg)

1. 11.43% API + 87.57% DCP anhydrous + 1% Magnesium stearate
2. 11.03% API + 87.97% Lactose anhydrous + 1% Magnesium stearate
3. 22.86% API + 76.14% Avicel PH200 LM + 1% Magnesium stearate
4. 17.78% API + 80.22% Mannitol 200 SD + 1% Magnesium stearate
+ 1% Aerosil 200

[00368] Binary mixtures (1 mg)

1. 0.50% API + 97.50% DCP anhydrous + 1% Magnesium stearate +
1% Aerosil 200
2. 0.53% API + 97.47% Lactose anhydrous + 1% Magnesium stearate
+ 1% Aerosil 200
3. 1.18% API + 97.82% Avicel PH200 LM + 1% Magnesium stearate
4. 0.91% API + 97.09% Mannitol 200 SD + 1% Magnesium stearate +
1% Aerosil 200

[00369] The appearance results for the study performed at 20 °C and 60% relative humidity (RH) and the study performed at 40 °C and 75% RH are shown in Table 7. The initial related substance results are shown in Table 8. The related substance results after two weeks at 25 °C and 60% RH are shown in Table 9, and after four weeks at 25 °C and 60% RH are shown in Table 10. The related substance results after two weeks at 40 °C and 75% RH are shown in Table 11, and after four weeks at 40 °C and 75% RH are shown in Table 12. The content uniformity for each batch of 16 mg capsules is shown in Table 13. The content uniformity for each batch of 1 mg capsules is shown in Table 14.

Table 7.

Specification Batch Details	Appearance				
	Initial	2 Weeks 25°C/60% RH	4 Weeks 25°C/60% RH	2 Weeks 40°C/75% RH	4 Weeks 40°C/75% RH
Control	Pale yellow powder	Pale yellow powder	Pale yellow powder	Pale yellow powder with small agglomeration observed	Pale yellow powder with small agglomeration observed
Binary mixtures (16 mg)					
1	Pale yellow powder	Pale yellow powder	Pale yellow powder	Pale yellow powder	Pale yellow powder
2	Pale yellow powder	Pale yellow powder	Pale yellow powder	Pale yellow powder with small agglomeration observed	Pale yellow powder with small agglomeration observed
3	Pale yellow powder	Pale yellow powder	Pale yellow powder	Pale yellow powder with small agglomeration observed	Pale yellow powder with small agglomeration observed
4	Pale yellow powder	Pale yellow powder	Pale yellow powder	Pale yellow powder with small agglomeration observed	Pale yellow powder with small agglomeration observed
Binary mixtures (1 mg)					
1	Off-white powder	Off-white powder	Off-white powder	Off-white powder	Off-white powder
2	Off-white powder	Off-white powder	Off-white powder	Off-white powder	Off-white powder
3	Off-white powder	Off-white powder	Off-white powder	Off-white powder	Off-white powder
4	Off-white powder	Off-white powder	Off-white powder	Off-white powder	Off-white powder

Table 8.

Batch Number	Related Substance (%Area) (Reporting threshold = peaks ≥ 0.05%) Approximate Relative Retention Time (RRT)				Total Related substance (% Area)
	RRT~0.59	RRT~0.63	RRT~0.90	Alvocidib RRT~4.0	
Control	<0.05	<0.05	<0.05	<0.05	<0.05
Binary mixtures (16 mg)					
1	<0.05	<0.05	<0.05	0.10	0.10
2	<0.05	<0.05	<0.05	0.12	0.12
3	<0.05	<0.05	<0.05	0.13	0.13
4	<0.05	<0.05	<0.05	0.13	0.13
Binary mixtures (1 mg)					
1	<0.05	<0.05	<0.05	0.18	0.18
2	<0.05	0.07	<0.05	0.13	0.20
3	<0.05	<0.05	0.06	0.19	0.25
4	0.05	<0.05	0.05	0.18	0.27

Table 9.

Batch Number	Related Substance (%Area) (Reporting threshold = peaks ≥ 0.05%) Approximate Relative Retention Time (RRT)						Total Related substance (%Area)
	RRT~ 0.43	RRT~ 0.47	RRT~ 0.59	RRT~ 0.63	RRT~ 0.90	Alvocidib RRT~4.0	
API							
Control	<0.05	<0.05	<0.05	<0.05	<0.05	0.05	0.05
Binary mixtures (16 mg)							
1	<0.05	<0.05	<0.05	<0.05	<0.05	0.14	0.14
2	<0.05	<0.05	<0.05	<0.05	<0.05	0.14	0.14
3	<0.05	<0.05	<0.05	<0.05	<0.05	0.12	0.12
4	<0.05	<0.05	<0.05	<0.05	<0.05	0.12	0.12
Binary mixtures (1 mg)							
1	<0.05	<0.05	0.07	<0.05	0.10	2.33	2.50
2	0.15	0.32	0.08	0.06	0.08	1.23	1.92
3	<0.05	<0.05	0.11	<0.05	0.13	0.78	1.02
4	<0.05	0.07	0.08	<0.05	0.08	1.35	1.58

Table 10.

Batch Number	Related Substance (%Area) (Reporting threshold = peaks ≥ 0.05%) Approximate Relative Retention Time (RRT)						Total Related substance (%Area)
	RRT~ 0.43	RRT~ 0.47	RRT~ 0.59	RRT~ 0.63	RRT~ 0.90	Alvocidib RRT~4.0	
API							
Control	<0.05	<0.05	<0.05	<0.05	<0.05	0.07	0.07
Binary mixtures (16 mg)							
1	<0.05	<0.05	<0.05	<0.05	<0.05	0.24	0.24
2	<0.05	0.07	<0.05	<0.05	<0.05	0.22	0.29
3	<0.05	<0.05	<0.05	<0.05	<0.05	0.19	0.19
4	<0.05	<0.05	<0.05	<0.05	<0.05	0.21	0.21
Binary mixtures (1 mg)							
1	<0.05	<0.05	0.08	<0.05	0.12	4.18	4.38
2	0.14	0.37	0.11	0.07	0.10	2.12	2.90
3	<0.05	0.07	0.12	<0.05	0.16	1.20	1.55
4	<0.05	<0.05	0.09	<0.05	0.10	2.32	2.51

Table 11.

Batch Number	Related Substance (%Area) (Reporting threshold = peaks ≥ 0.05%) Approximate Relative Retention Time (RRT)						Total Related substance (%Area)
	RRT~ 0.43	RRT~ 0.47	RRT~ 0.59	RRT~ 0.90	RRT~ 2.06	Alvocidib RRT~4.0	
API							
Control	<0.05	<0.05	<0.05	<0.05	<0.05	0.12	0.12
Binary mixtures (16 mg)							
1	<0.05	<0.05	<0.05	<0.05	<0.05	2.57	2.57
2	<0.05	<0.05	<0.05	<0.05	<0.05	0.72	0.72
3	<0.05	<0.05	<0.05	<0.05	<0.05	0.31	0.31
4	<0.05	<0.05	<0.05	<0.05	<0.05	1.03	1.03
Binary mixtures (1 mg)							
1	<0.05	<0.05	0.15	0.06	0.12	20.42	20.75
2	0.18	0.26	0.24	0.50	0.13	14.04	15.35
3	<0.05	<0.05	0.17	0.21	0.06	4.96	5.40
4	<0.05	0.06	0.31	0.50	0.30	24.72	25.88

Table 12.

Batch Number	Related Substance (%Area) (Reporting threshold = peaks ≥ 0.05%) Approximate Relative Retention Time (RRT)						Total Related substance (%Area)
	RRT~ 0.43	RRT~ 0.47	RRT~ 0.59	RRT~ 0.90	RRT~ 2.06	Alvocidib RRT~ 4.0	
API							
Control	<0.05	<0.05	<0.05	<0.05	<0.05	0.21	0.21
Binary mixtures (16 mg)							
1	<0.05	<0.05	<0.05	<0.05	<0.05	3.73	3.73
2	<0.05	<0.05	<0.05	<0.05	<0.05	1.15	1.15
3	<0.05	<0.05	<0.05	<0.05	<0.05	0.50	0.50
4	<0.05	<0.05	<0.05	<0.05	<0.05	1.85	1.85
Binary mixtures (1 mg)							
1	<0.05	<0.05	0.12	<0.05	0.22	24.43	24.77
2	0.19	0.25	0.27	0.33	0.27	18.84	20.15
3	<0.05	<0.05	0.20	0.19	0.13	7.36	7.87
4	<0.05	<0.05	0.33	0.34	0.68	35.51	36.85

Table 13.


Binary mixture 1 (16 mg)			Binary mixture 2 (16 mg)			Binary mixture 3 (16 mg)			Binary mixture 4 (16 mg)		
Replicate	% Label Claim		Replicate	% Label Claim		Replicate	% Label Claim		Replicate	% Label Claim	
R1	97.3		R1	98.4		R1	87.4		R1	91.4	
R2	91.8		R2	99.3		R2	87.1		R2	89.2	
R3	94.9		R3	99.8		R3	88.5		R3	92.0	
R4	96.3		R4	99.1		R4	84.8		R4	94.5	
R5	94.3		R5	94.5		R5	85.9		R5	93.4	
R6	93.1		R6	95.3		R6	94.2		R6	89.7	
R7	90.6		R7	93.9		R7	87.5		R7	87.0	
R8	94.2		R8	98.3		R8	88.5		R8	89.2	
R9	95.1		R9	92.1		R9	87.0		R9	83.2	
R10	92.1		R10	89.1		R10	83.9		R10	83.5	
Min	90.6		Min	89.1		Min	83.9		Min	83.2	
Max	97.3		Max	99.8		Max	94.2		Max	94.5	
Mean	94.0		Mean	96.0		Mean	87.5		Mean	89.3	
Std. Dev.	2.1		Std. Dev.	3.6		Std. Dev.	2.8		Std. Dev.	3.8	
% RSD	2.2		% RSD	3.7		% RSD	3.2		% RSD	4.3	
AV ≤ 15.0	9.5	Pass	AV ≤ 15.0	11.1	Pass	AV ≤ 15.0	17.7	Fail	AV ≤ 15.0	18.3	Fail

Table 14.

Binary mixture 1 (1 mg)			Binary mixture 2 (1 mg)			Binary mixture 3 (1 mg)			Binary mixture 4 (1 mg)		
Replicate	% Label Claim		Replicate	% Label Claim		Replicate	% Label Claim		Replicate	% Label Claim	
R1	94.3		R1	98.0		R1	97.1		R1	97.1	
R2	91.4		R2	97.7		R2	98.4		R2	98.4	
R3	100.0		R3	98.9		R3	100.7		R3	100.7	
R4	98.7		R4	97.6		R4	96.6		R4	96.6	
R5	90.3		R5	98.3		R5	98.4		R5	98.4	
R6	96.8		R6	98.0		R6	96.9		R6	96.9	
R7	99.7		R7	95.9		R7	96.9		R7	96.9	
R8	99.4		R8	99.9		R8	96.4		R8	96.4	
R9	97.7		R9	99.0		R9	97.5		R9	97.5	
R10	95.7		R10	97.9		R10	93.9		R10	93.9	
Min	90.3		Min	95.9		Min	93.9		Min	93.9	
Max	100.0		Max	99.9		Max	100.7		Max	100.7	
Mean	96.4		Mean	98.1		Mean	97.3		Mean	97.3	
Std. Dev.	3.4		Std. Dev.	1.1		Std. Dev.	1.7		Std. Dev.	1.7	
% RSD	3.6		% RSD	1.1		% RSD	1.8		% RSD	1.8	
AV ≤ 15.0	10.3	Pass	AV ≤ 15.0	2.9	Pass	AV ≤ 15.0	5.4	Pass	AV ≤ 15.0	5.4	Pass

EXAMPLE 10

SYNTHESIS OF STRUCTURE (I)



Scheme 2.

[00370] Step 1.1: To a clean and dry, three-necked, round-bottomed flask (RBF) (3 L) was added A-1 (90 g, 0.192 mol) and chlorobenzene (774 ml) at room temperature. To the reaction flask was slowly added BBr_3 (391.5 g) at room temperature. After completion of BBr_3 addition, the temperature of the reaction mixture was slowly raised to 80-83 °C, and the reaction mixture was stirred at the same temperature for 10 hours. The reaction mixture temperature was further raised to 100-103 °C, and the reaction mixture was maintained at 100-103 °C for 5 hours. The reaction progress was monitored by TLC and HPLC. After completion of the reaction, HBr and methyl bromide was removed at room temperature by nitrogen bubbling into the reaction mixture, while maintaining the vigorous stirring. The reaction mixture was slowly quenched with a mixture of methanol (180ml)/water (90ml) (270 ml), followed by methanol (180 ml). The solvent was removed under atmospheric distillation at 25-50 °C to reach the target reaction mass volume of 12 volumes (vol). Then, the reaction mixture pH was adjusted to 3.0 ± 1 using sodium hydroxide solution (48.8 g dissolved in 135 ml of DM water) at 50-55 °C. Again, the solvent was removed under atmospheric distillation at 50-100 °C to reach the target reaction volume of 12 vol. Then, the pH of the reaction mixture was adjusted to $\text{pH } 8.1 \pm 0.2$ using sodium hydroxide solution (8.5 g dissolved in 87 ml of DM water) at 50 °C followed by slow addition of water with constant stirring at 50 °C for 1 hour. The reaction mixture was slowly allowed to come to room temperature and maintained at room temperature for 3 hours. The resulting solid was filtered and washed with a mixture of methanol (315 ml)/water (135 ml) (3 x 450 ml) followed by water (5 x 450 ml). The solid was dried in a vacuum oven at 50-55 °C for 48 hours to obtain A-2 as a yellow solid (70 g, 90%).
HPLC Purity: 99.72%.

[00371] Step 2.1A: To a clean and dry, three-necked RBF (3 L) was added A-2 (35.0 g, 0.087 mol) and DMF (245 ml), at room temperature, under nitrogen atmosphere. Then, DMAP (1.06 g, 0.0086 mol) followed by CCl_4 (66.5 g 0.434mol) were added to the reaction mixture at room temperature. To the reaction mixture di-tertiary butyl phosphite (25.5g, 0.131 mol) was added at room temperature. The reaction mixture was stirred at room temperature under nitrogen atmosphere for 24 hours. The reaction progress was monitored by HPLC. The reaction mixture was cooled to 0-5 °C, and was quenched with slow addition of DM water (1950 ml) for 30 minutes at 0-5 °C. Then,

chloroform (1627.5 ml) was added to the reaction mixture, and the reaction mixture was stirred at 0-5 °C for 10 minutes. The organic layer was separated and dried over sodium sulfate. The solvent was removed under reduced pressure, while maintaining the bath temperature below 45 °C. The resulting residue was co-distilled with toluene (4 × 175 ml). The residue was kept under high vacuum for 45 minutes to obtain A-10 as a pale yellow residue. (51.0 g, 98.5%). HPLC Purity: 91.48%.

[00372] Step 2.1B: To a clean and dry RBF (1 L) was added A-10 (51.0 g, 0.0858 mol) and acetic acid (102 ml) at room temperature. Then, 4N HCl solution in 1,4-dioxane (102 ml) was added dropwise at 25-30 °C. The reaction mixture was stirred at 25-30 °C for 40 minutes. The reaction progress was monitored by TLC. After completion of the reaction, toluene (2 × 510 ml) was added to the reaction mixture under stirring, and the reaction mixture was maintained for 5 minutes. The stirring was stopped, and the solids in the reaction mixture were allowed to settle at 25-30 °C for 5 minutes. The solvent was decanted to obtain the semi-solid. The semi-solid was co-distilled with toluene (3 × 123 ml) to obtain pale yellow solid. The resulting pale yellow solid was taken into a clean RBF, and methanol was added (123 ml) followed by dropwise addition of water (41 ml) at 25-30 °C. The reaction mixture was stirred at 25-30 °C for 2 hours to obtain pale yellow solid. The resulting solid was filtered and vacuum dried for 10 minutes to obtain A-11 as a pale yellow solid (36.5 g, 82%). HPLC Purity: 97.03%. This material was directly taken into Step 3.1 without further drying.

[00373] Step 3.1: To a clean and dry, three-necked, 500 ml RBF was added A-11 (34.0 g, 0.066 mol) and ACN (51 ml). To the reaction mixture was dropwise added ammonium bicarbonate solution (16.2 g dissolved in 170 ml of DM water) under stirring at 25-30 °C for 30 minutes. Again, ACN (51 ml) was slowly added at 25-30 °C for 30 minutes. The reaction mixture was cooled to 10-15 °C and stirred at 10-15 °C for 60 minutes. The resulting solid was filtered and washed with ACN (102 ml). The solid was dried in a vacuum oven at 25-30 °C for 16 hours to obtain A-4 as a pale yellow solid (28.5 g, 90.10%). HPLC Purity: 99.68%.

[00374] Step 4.1: To a clean and dry, 500-ml, three-necked RBF was added A-4 (7.5 g, 0.015 mol) and methanol (187.5 ml) at room temperature. To the reaction mixture was slowly added acetic acid (7.5 ml, 1.0 vol) at 50 °C, under nitrogen atmosphere. The reaction mixture was stirred at 50 °C for 1 h, under nitrogen atmosphere. The reaction

mixture was cooled to room temperature and stirred for 2 h. The solid was filtered and dried under vacuum to obtain 5.0 g A-5 (66.5%) as a pale yellow solid. HPLC Purity: 99.77%.

[00375] Several screens were conducted to identify the Step 4.1 conditions described above. In a first screen, A-4 was treated with acid (1.0 volume) and methanol (25 volumes) according to the conditions listed in Table 15 to obtain A-5 as a pale yellow solid.

Table 15.

Acid	Brief Procedure	A-4 (g) (HPLC Purity (%); A-2 (%))	A- 5 (g)	Yield (%)	HPLC Purity (%)	A-2 (%)
Acetic acid	<i>See</i> description of Step 4.1 above.	7.5 (99.68; 0.08)	5.0	66.5	99.77	0.12
Citric acid	To a clean and dry, 100-ml, three-necked RBF was added A-4 (1 g, 0.002 mol) and methanol (25 ml) at room temperature. To the reaction mixture was added citric acid (1.0 g, 2.5 eq) at 50 °C, under nitrogen atmosphere. The reaction mixture was stirred at 50 °C for 1 h, under nitrogen atmosphere. The reaction mixture was cooled to room temperature and stirred for 2 h. The solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.	1.0 (99.72; 0.09)	0.6	66.5	99.59	0.28
Propionic acid	To a clean and dry, 100-ml, three-necked RBF was added A-4 (0.8 g, 0.001 mol) and methanol (20 ml) at room temperature. To the reaction mixture was dropwise added propionic acid (0.8 ml) at 50 °C, under nitrogen atmosphere. The reaction mixture was stirred at 50 °C for 1 h, under nitrogen atmosphere. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-4 as a	0.8 (99.72; 0.09)	0.5	62.5	99.41	0.12

	pale yellow solid.					
--	--------------------	--	--	--	--	--

[00376] In another screen, A-4 was treated with acetic acid (1 volume) and a solvent (25 volumes) according to the conditions listed in Table 16 to obtain A-5 as a pale yellow solid.

Table 16.

Solvent	Brief Procedure	A-4 (g)	A-5 (g)	Yield (%)	HPLC Purity (%)	A-2 (%)
n-butanol	To a clean and dry, 100-ml, three-necked RBF was added A-4 (2 g, 0.004 mol) and n-butanol (50 ml) at room temperature. Then, acetic acid (2.0 ml) was slowly added to the reaction mixture at 50 °C, under nitrogen atmosphere. The resulting reaction mixture was stirred at 50 °C for 1 h, under nitrogen atmosphere. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.	2.0	1.0	50	99.85	0.04
Methyl ethyl ketone		2.0	1.2	60	99.48	0.25
Ethanol	To a clean and dry, 100-ml, three-necked RBF was added A-4 (2 g, 0.004 mol) and ethanol (50 ml) at room temperature. Then, acetic acid (2.0 ml) was slowly added to the reaction mixture at 50 °C, under nitrogen atmosphere. The resulting reaction mixture was stirred at 50 °C under nitrogen atmosphere for 1 h. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.	2.0	1.4	70	99.88	0.03
Ethanol (10 volumes) Acetone	To a clean and dry, 100-ml, three-necked RBF was added A-4 (1 g, 0.002 mol), ethanol (10 ml) and acetone (10 ml) at room temperature.	1.0	0.5	50	99.83	0.06

(10 volumes)	Then, acetic acid (1.0 ml) was slowly added to the reaction mixture at 50 °C, under nitrogen atmosphere. The resulting reaction mixture was stirred at 50 °C under nitrogen atmosphere for 1 h. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.					
--------------	--	--	--	--	--	--

[00377] In another screen, A-4 was treated with acetic acid (1 volume) and a solvent according to the conditions listed in Table 17 to obtain A-5 as a pale yellow solid.

Table 17.

Solvent	Brief Procedure	A-4 (g)	A-5 (g)	Yield (%)	HPLC Purity (%)	A-2 (%)
Ethanol (15 volumes) Ethyl acetate (10 volumes)	To a clean and dry, 100-ml, three-necked RBF was added A-4 (1.5 g, 0.003 mol), ethanol (22.5 ml) and ethyl acetate (15 ml) at room temperature, under nitrogen atmosphere. Then, acetic acid (1.5 ml) was slowly added at 50 °C. The resulting reaction mixture was stirred at 50 °C under nitrogen atmosphere for 1 h. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.	1.5	1.08	72	99.60	0.09
Ethanol (15 volumes) <i>i</i> -Pr ₂ O (10 volumes)		1.5	1.25	83	99.68	0.12
Ethanol (15 volumes) Water (1 volume)	To a clean and dry, 100-ml, three-necked RBF was added A-4 (1.5 g, 0.003 mol), ethanol (22.5 ml) and DM water (1.5 ml) at room temperature. Then, acetic acid (1.5 ml) was slowly added to the reaction mixture at 50 °C. The resulting reaction mixture was stirred at 50 °C	1.5	1.1	73	99.60	0.09

	for 1 h. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.					
--	---	--	--	--	--	--

[00378] In another screen, A-4 was treated with acetic acid and ethanol (25 volumes) according to the conditions listed in Table 18 to obtain A-5 as a pale yellow solid.

Table 18.

Acetic Acid	Brief Procedure	A-4 (g)	A-5 (g)	Yield (%)	HPLC Purity (%)	A-2 (%)
0.8 volumes	To a clean and dry, 100-ml, three-necked RBF was added A-4 (1.5 g, 0.003 mol) and ethanol (37.5 ml) at room temperature, under nitrogen atmosphere. Then, acetic acid (1.2 ml) was slowly added at 50 °C. The resulting reaction mixture was stirred under nitrogen atmosphere at 50 °C for 1 h. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.	1.5	1.15	76	99.70	0.12
0.6 volumes	To a clean and dry, 100-ml, three-necked RBF was added A-4 (1.5 g, 0.003 mol) and ethanol (37.5 ml) at room temperature, under nitrogen atmosphere. Then, acetic acid (0.9 ml) was slowly added at 50 °C. The resulting reaction mixture was stirred at 50 °C, under nitrogen atmosphere for 1 h. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.	1.5	1.17	78	99.73	0.09
0.4 volumes	To a clean and dry, 100-ml, three-necked RBF was added A-4 (1.5 g, 0.003 mol) and ethanol (37.5 ml) at room temperature, under nitrogen atmosphere. Then, acetic acid (0.6 ml) was slowly added at 50 °C. The	1.5	1.0	66	99.71	0.10

	resulting reaction mixture was stirred at 50 °C, under nitrogen atmosphere for 1 h. The reaction mixture was cooled to room temperature and stirred for 2 h. The resulting solid was filtered and dried under vacuum to obtain A-5 as a pale yellow solid.					
--	--	--	--	--	--	--

[00379] As an alternative to conducting Step 4.1 using the conditions cited above in Example 10, a procedure similar to that described in Step 4 of Example 1 can be used to effect polymorph conversion. Thus, in another experiment, to a clean and dry, 100-ml, three-necked RBF was added A-4 (2.0 g, 0.004 mol), THF (29 ml) and DM water (1.7 ml) at room temperature. Then, maleic acid (0.44 g) was added to the reaction mixture at room temperature. The reaction mixture was stirred at room temperature for 12 h. The resulting solid was filtered and vacuum dried. The wet solid was dissolved in ethanol (12 ml) at room temperature and was stirred for 24 h. The resulting solid was filtered, washed with ethanol (2.5 ml), vacuum dried to obtain A-5 (1.5g, 60%) as a pale yellow solid. HPLC Purity: 99.91%.

EXAMPLE 11

FORMULATION STABILITY STUDY

[00380] Polymorph Form B obtained using various Step 2.1B reaction conditions was submitted to a four-week stability study. The results of the stability study appear in Table 19. The Reference Standard is a sample obtained using the synthetic procedure described in Example 1.

Table 19.

Study No.	Step 2.1B Reaction Conditions	Stability at 40 °C ± 2 °C, 75 ± 5% RH, sealed vial, HPLC Purity of A-5 (%) / A-2 (%)				
		Initial	3 days ± 1 day	7 days ± 1 day	14 days ± 1 day	28 days ± 1 day
1	4N HCl in 1,4-dioxane	99.89/0.03	98.83/0.94	98.37/1.37	97.58/2.14	96.73/2.95
2	10% HCl in EtOAc	99.69/0.16	99.52/0.32	99.31/0.51	98.96/0.81	98.40/1.34
3	Conc. HCl in IPA	99.65/0.19	99.09/0.72	98.52/1.24	97.76/1.96	96.67/2.95

4	Acetic acid in methanol	99.77/0.09	99.64/0.23	99.42/0.40	99.24/0.58	
Reference Standard	N/A	99.90/0.03	99.91/0.06	99.81/0.13	99.70/0.24	99.41/0.53

[00381] Polymorph Form B obtained using the Step 2.1B reaction conditions described in Scheme 2 was submitted to a four-week stability study to test the effects of different packing conditions on the stability of A-5. Under Packing Condition 1, polymorph Form B was packed into an amber-colored bottle, which was closed with a rubber stopper and sealed with a flip-off aluminum cap. Under Packing Condition 2, polymorph Form B was packed in a low-density polyethylene bag (LDPE) twisted and tied. The LDPE bag was then inserted into a black, LDPE bag along with one silica gel bag, and the black, LDPE bag was heat sealed under nitrogen atmosphere. The black, LDPE bag was then inserted into a triple-laminated aluminum bag along with one silica gel bag, and the aluminum bag was heat sealed under nitrogen atmosphere. The triple-laminated aluminum bag containing polymorph Form B was kept in a high-density polyethylene container. The results of the stability study appear in Table 20.

Table 20.

Packing Condition	Stability at 40 °C ± 2 °C, 75 ± 5% RH, sealed vial, HPLC Purity of A-5 (%) / A-2 (%)				
	Initial	3 days ± 1 day	7 days ± 1 day	14 days ± 1 day	28 days ± 1 day
1	99.77/0.09	99.64/0.23	99.42/0.40	99.24/0.58	98.59/1.23
2	99.77/0.09	99.68/0.18	99.50/0.30	99.23/0.59	99.10/0.74

EXAMPLE 12

FORMULATION STABILITY STUDY

[00382] Polymorph Form B obtained using various Step 4.1 reaction conditions described in Example 10 was submitted to a four-week stability study. The results of the stability study appear in Table 21.

Table 21.

Conditions	Stability at 40 °C ± 2 °C, 75 ± 5% RH, sealed vial, HPLC Purity of A-5 (%) / A-2 (%)				
	Initial	7 days ± 1 day	14 days ± 1 day	21 days ± 1 day	28 days ± 1 day
Citric acid	99.2/0.6	96.6/3.06	93.12/6.08	91.3/8.0	89.7/9.6
Propionic acid	99.6/0.21	99.0/0.87	98.4/1.38	97.8/1.94	97.5/2.2
n-butanol	99.8/0.1	99.1/0.47	98.9/0.82	98.6/1.2	98.1/1.6
Ethanol	99.7/0.14	99.1/0.81	98.3/1.3	98.1/1.7	97.4/2.2
Ethanol + acetone	99.7/0.14	98.6/1.2	97.9/1.8	97.3/2.3	96.9/2.8
A-11 (not submitted to Step 3.1 or 4.1)	97.0/1.02	91.3/6.6	84.12/13.7	78.5/19.0	75.0/22.2
Ethanol (15 volumes) Ethyl acetate (10 volumes)	99.5/0.17	98.7/0.92	97.9/1.6	97.6/2.0	96.9/2.6
Ethanol (15 volumes) Water (1 volume)	99.8/0.04	99.6/0.25	99.3/0.41	99.2/0.58	99.1/0.73
Ethanol: AcOH (0.8 volumes)	99.7/0.14	99.3/0.46	99.0/0.68	98.6/0.98	98.6/1.1
Ethanol: AcOH (0.6 volumes)	99.7/0.11	99.1/0.6	98.5/0.99	98.3/1.3	98.1/1.7
Ethanol: AcOH (0.4 volumes)	99.7/0.14	99.3/0.51	98.5/1.1	98.3/1.4	98.0/1.7
Maleic acid	99.9/0.04	99.8/0.07	99.6/0.11	99.7/0.17	99.7/0.18
Ethanol (15 volumes) Diisopropyl ether (10 volumes)	98.9/0.71	97.6/1.86	98.3/1.5		
A-4 (not submitted to Step 4.1)	99.2/0.44	98.6/0.88	97.5/2.2		

EXAMPLE 13

EXCIPIENT COMPATIBILITY STUDY

[00383] The compatibility of binary mixtures of 95% excipient and 5% compound of structure (I) (equivalent to a 20:1 excipient:compound of structure (I) ratio) was studied. The following excipients formed part of the excipient compatibility study:

Category		Product name
Filler	D-Mannitol	PEARITOL 50C
	Anhydrous lactose	SuperTab22AN

	Microcrystalline cellulose	AvicelPH112
	Cornstarch	Cornstarch (JP)
	Anhydrous dibasic calcium phosphate	Anhydrous dibasic calcium phosphate
Disintegrant	Croscarmellose sodium	Ac-Di-Sol SD-711
	Partly pregelatinized starch	Starch1500G
	Crospovidone	KollidonCL
	Low substituted hydroxyl-propylcellulose	L-HPC(LH-21)
	Carmellose calcium	ECG505
Lubricant	Magnesium stearate	LIGAMED MF-2-V
	Sodium stearyl fumarate	PRUV
	Talc	Talc MS-P
Fluidizer	Silicon dioxide	Aerosil 200
Others	Fumaric acid	Fumaric acid NF
	Succinic acid	Succinic acid cryst EMPROVE ESSENTIAL NF JPE ACS

[00384] Samples were tested for related substances (Total Impurities, by HPLC) at 50 °C and 85% RH, and at 60 °C at two- and four-week time points. Anhydrous lactose, cornstarch, partly pregelatinized starch, low-substituted hydroxyl propyl cellulose, carmellose calcium and magnesium stearate demonstrated compatibility with the compound of structure (I). The results of the excipient compatibility study are described in Tables 22A-22P.

Table 22A.

Filler	PEARITOL 50C						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	0.05	0.05	0.07	0.15	<0.05	<0.05
0.87-0.89	N.D.	N.D.	N.D.	<0.05	0.07	<0.05	<0.05
1.00	99.64	94.79	93.91	87.82	66.13	99.41	99.02
1.35-1.37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.13	0.18	0.14	0.64	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.36	5.03	5.85	11.94	33.01	0.55	0.92
Total impurity	0.36	5.21	6.08	12.15	33.87	0.55	0.92
Δimpurity		4.85	5.72	11.79	33.51	0.19	0.56

Table 22B.

Filler	SUpertab22AN						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	0.09	N.D.	<0.05	N.D.	N.D.	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	0.09	0.09	0.10	0.15	<0.05	<0.05
0.87-0.89	N.D.	0.07	0.05	0.05	0.05	<0.05	<0.05
1.00	99.64	96.90	96.85	89.04	80.08	99.70	99.49
1.35-1.37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.09	0.14	0.23	0.65	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.36	2.76	2.88	10.55	19.04	0.25	0.45
Total impurity	0.36	3.10	3.16	10.93	19.89	0.25	0.45
Δimpurity		2.74	2.80	10.57	19.53	-0.11	0.09

Table 22C.

Filler	AvicelPH112						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	0.06	N.D.	N.D.	<0.05	N.D.	N.D.
0.48-0.51	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	0.17	0.16	0.13	0.16	<0.05	<0.05
0.87-0.89	N.D.	0.13	0.09	0.05	<0.05	0.05	0.08

1.00	99.74	96.17	96.17	84.69	77.55	99.58	99.34
1.35-1.37	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.23	0.29	0.29	0.49	N.D.	N.D.
3.36-3.45	N.D.	N.D.	<0.05	<0.05	0.05	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.26	3.21	3.22	14.82	21.69	0.33	0.53
Total impurity	0.26	3.80	3.76	15.29	22.39	0.38	0.61
Δimpurity		3.54	3.50	15.03	22.13	0.12	0.35

Table 22D.

Filler	Cornstarch(JP)						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	<0.05	N.D.	N.D.	N.D.	<0.05	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	0.12	0.12	0.07	0.09	<0.05	<0.05
0.87-0.89	N.D.	0.09	0.07	<0.05	<0.05	<0.05	0.05
1.00	99.64	96.60	96.43	93.74	90.03	99.62	99.46
1.35-1.37	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.21	0.28	0.16	0.32	N.D.	N.D.
3.36-3.45	N.D.	<0.05	<0.05	N.D.	0.06	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.36	2.93	3.05	5.99	9.47	0.30	0.46
Total impurity	0.36	3.35	3.52	6.22	9.94	0.30	0.51
Δimpurity		2.99	3.16	5.86	9.58	-0.06	0.15

Table 22E.

Filler	Anhydrous dibasic calcium phosphate						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	<0.05	<0.05	<0.05	0.06	N.D.	N.D.
0.87-0.89	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.
1.00	99.85	97.45	97.32	83.48	79.63	99.54	99.25
1.35-1.37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.08	0.12	0.18	0.35	N.D.	N.D.

3.36-3.45	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.15	2.43	2.52	16.27	19.93	0.46	0.75
Total impurity	0.15	2.51	2.64	16.45	20.34	0.46	0.75
Δimpurity		2.36	2.49	16.30	20.19	0.31	0.60

Table 22F.

Disintegrant	Ac-Di-Sol						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60Lux	120Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.26-0.27	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.38-0.41	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.4-0.45	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.
0.48-0.51	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	0.10	0.09	0.06	0.08	<0.05	<0.05
0.59	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.72	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	0.07	0.05	N.D.	N.D.	0.05	0.07
1.00	99.90	98.29	98.16	87.67	78.62	99.79	99.70
1.35-1.37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.12	0.17	0.11	0.28	N.D.	N.D.
2.37-2.39	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.10	1.39	1.51	12.17	20.98	0.13	0.20
4.75-4.91	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Total impurity	0.10	1.68	1.82	12.34	21.34	0.18	0.27
Δimpurity		1.58	1.72	12.24	21.24	0.08	0.17

Table 22G.

Disintegrant	Starch1500G						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.26-0.27	N.D.	0.09	0.08	0.09	0.09	0.07	0.06
0.38-0.41	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.

0.4-0.45	N.D.	<0.05	N.D.	<0.05	N.D.	N.D.	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	<0.05	0.05	0.05	0.09	N.D.	N.D.
0.59	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.72	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	<0.05	<0.05	<0.05	<0.05	N.D.	<0.05
1.00	99.64	98.59	98.54	94.86	90.67	99.73	99.63
1.35-1.37	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.
2.19-2.25	N.D.	0.06	0.11	0.19	0.47	N.D.	N.D.
2.37-2.39	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	<0.05	0.11	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.36	1.15	1.21	4.71	8.50	0.20	0.29
4.75-4.91	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Total impurity	0.36	1.30	1.45	5.04	9.26	0.27	0.35
Δimpurity		0.94	1.09	4.68	8.90	-0.09	-0.01

Table 22H.

Disintegrant	KollidonCL						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.26-0.27	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.38-0.41	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.4-0.45	N.D.	<0.05	N.D.	N.D.	<0.05	N.D.	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	0.19	0.33	0.10	0.21	N.D.	<0.05
0.59	N.D.	N.D.	N.D.	0.28	0.14	N.D.	N.D.
0.72	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.
0.87-0.89	N.D.	0.07	0.18	0.07	0.08	<0.05	0.07
1.00	99.62	96.41	98.26	78.75	59.59	99.66	99.48
1.35-1.37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.20	0.09	0.27	0.70	N.D.	N.D.
2.37-2.39	N.D.	N.D.	N.D.	0.11	0.23	N.D.	N.D.
3.36-3.45	N.D.	<0.05	N.D.	N.D.	0.06	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.38	2.71	1.14	18.88	38.97	0.30	0.41
4.75-4.91	N.D.	0.38	N.D.	1.50	N.D.	N.D.	N.D.
Total impurity	0.38	3.55	1.74	21.21	40.39	0.30	0.48
Δimpurity		3.17	1.36	20.83	40.01	-0.08	0.10

Table 22I.

Disintegrant	L-HPC(LH-21)						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.26-0.27	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.38-0.41	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.	N.D.
0.4-0.45	N.D.	<0.05	N.D.	N.D.	<0.05	N.D.	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	0.09	0.09	0.06	0.08	N.D.	<0.05
0.59	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.72	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	0.06	0.05	<0.05	<0.05	0.05	0.07
1.00	99.63	97.29	97.20	94.21	91.11	99.67	99.47
1.35-1.37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.19	0.24	0.17	0.31	N.D.	N.D.
2.37-2.39	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.36-3.45	N.D.	N.D.	<0.05	<0.05	0.07	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.37	2.31	2.40	5.50	8.37	0.28	0.43
4.75-4.91	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Total impurity	0.37	2.65	2.78	5.73	8.83	0.33	0.50
Δimpurity		2.28	2.41	5.36	8.46	-0.04	0.13

Table 22J.

Disintegrant	ECG505						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.26-0.27	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.38-0.41	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.4-0.45	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	0.07	0.07	0.05	0.07	N.D.	<0.05
0.59	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.72	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	0.07	0.05	<0.05	<0.05	<0.05	<0.05
1.00	99.72	98.81	98.75	96.51	94.81	99.81	99.72
1.35-1.37	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.

2.19-2.25	N.D.	0.05	0.09	0.08	0.18	N.D.	N.D.
2.37-2.39	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.28	1.00	1.04	3.33	4.89	0.17	0.22
4.75-4.91	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Total impurity	0.28	1.19	1.25	3.46	5.14	0.17	0.22
Δimpurity		0.91	0.97	3.18	4.86	-0.11	-0.06

Table 22K.

Lubricant	LIGAMED MF-2-V						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	<0.05	N.D.	N.D.	<0.05	<0.05	N.D.
0.48-0.51	N.D.	N.D.	<0.05	<0.05	0.05	<0.05	N.D.
0.57-0.59	N.D.	0.08	0.13	0.10	0.25	0.08	<0.05
0.59	N.D.	N.D.	N.D.	0.09	N.D.	N.D.	N.D.
0.68	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.76-0.77	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	0.17	0.10	0.17	0.16	N.D.	<0.05
1.00	99.88	97.61	97.55	88.71	68.97	99.74	99.77
1.17-1.18	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
1.35-1.37	N.D.	N.D.	<0.05	0.17	0.36	<0.05	N.D.
2.19-2.25	N.D.	0.09	0.13	1.00	3.24	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	0.14	0.66	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.12	2.03	2.03	9.59	26.29	0.10	0.17
4.75-4.91	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.
Total impurity	0.12	2.37	2.39	11.26	31.01	0.18	0.17
Δimpurity		2.25	2.27	11.14	30.89	0.06	0.05

Table 22L.

Lubricant	PRUV						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.

0.57-0.59	N.D.	<0.05	0.10	<0.05	0.70	0.17	<0.05
0.59	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.68	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.76-0.77	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	N.D.	N.D.	<0.05	0.20	<0.05	N.D.
1.00	99.89	99.54	97.78	98.00	7.62	99.63	99.81
1.17-1.18	N.D.	N.D.	<0.05	N.D.	N.D.	<0.05	N.D.
1.35-1.37	N.D.	N.D.	<0.05	N.D.	0.10	<0.05	N.D.
2.19-2.25	N.D.	N.D.	0.11	0.10	5.01	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	<0.05	0.57	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.11	0.43	1.97	1.80	85.74	0.11	0.16
4.75-4.91	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Total impurity	0.11	0.43	2.18	1.90	92.32	0.28	0.16
Δimpurity		0.32	2.07	1.79	92.21	0.17	0.05

Table 22M.

Lubricant	Talc MS-P						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.	N.D.
0.48-0.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	N.D.	0.05	0.19	0.05	N.D.	N.D.
0.59	N.D.	N.D.	N.D.	0.08	N.D.	N.D.	N.D.
0.68	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.
0.76-0.77	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	<0.05	<0.05	0.24	<0.05	N.D.	N.D.
1.00	99.74	99.71	99.04	79.22	96.62	99.90	99.84
1.17-1.18	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
1.35-1.37	N.D.	N.D.	N.D.	0.06	<0.05	N.D.	N.D.
2.19-2.25	N.D.	N.D.	0.10	0.70	0.21	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	0.05	0.05	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.26	0.24	0.77	19.44	3.02	0.10	0.26
4.75-4.91	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Total impurity	0.26	0.24	0.92	20.76	3.33	0.10	0.26
Δimpurity		-0.02	0.66	20.50	3.07	-0.16	0.00

Table 22N.

Fluidizer	Aerosil 200						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.4-0.45	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.	N.D.
0.57-0.59	N.D.	N.D.	N.D.	0.12	0.14	N.D.	N.D.
0.87-0.89	N.D.	N.D.	0.05	<0.05	0.05	<0.05	N.D.
1.00	99.69	99.46	99.45	84.21	77.22	99.80	99.77
2.19-2.25	N.D.	N.D.	<0.05	0.17	0.32	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.31	0.52	0.47	15.46	22.27	0.17	0.23
Total impurity	0.31	0.52	0.52	15.75	22.78	0.17	0.23
Δimpurity		0.21	0.21	15.44	22.47	-0.14	-0.08

Table 22O.

Others	Fumaric acid						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.38-0.41	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.4-0.45	N.D.	N.D.	N.D.	N.D.	61.79	N.D.	N.D.
0.48-0.51	N.D.	0.06	0.07	<0.05	<0.05	N.D.	<0.05
0.57-0.59	N.D.	0.07	0.08	0.05	0.07	N.D.	<0.05
0.72	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.76-0.77	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	0.05	0.05	<0.05	<0.05	N.D.	<0.05
1.00	99.50	98.57	98.61	96.65	34.52	99.71	99.35
1.28	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
1.35-1.37	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.
2.08	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	<0.05	0.06	0.11	0.20	N.D.	N.D.
2.37-2.39	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.49	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.16	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.36-3.45	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.
3.59	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.62-3.71	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.67-3.72(3.77-	0.50	1.22	1.14	3.13	3.31	0.29	0.57

3.79)Alvocidib							
3.89	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.27	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.61	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.70	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.75-4.91	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
5.03	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
5.14	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
5.32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
5.58	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.05	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.13	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.46	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.54	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
7.05	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
7.09	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Total impurity	0.50	1.40	1.40	3.29	65.37	0.29	0.57
Δimpurity		0.90	0.90	2.79	64.87	-0.21	0.07

Table 22P.

Others	Succinic acid						
	Initial	60°C		50°C85%		Photostability	
		2W	4W	2W	4W	60 Lux	120 Lux
RRT	%Area	%Area	%Area	%Area	%Area	%Area	%Area
0.38-0.41	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.4-0.45	N.D.	0.22	N.D.	<0.05	N.D.	N.D.	N.D.
0.48-0.51	N.D.	<0.05	N.D.	0.15	0.09	N.D.	N.D.
0.57-0.59	N.D.	0.28	0.21	2.31	2.34	<0.05	<0.05
0.72	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.76-0.77	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.87-0.89	N.D.	0.09	N.D.	0.78	0.26	<0.05	<0.05
1.00	99.50	93.38	93.23	26.27	0.41	99.53	99.16
1.28	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
1.35-1.37	N.D.	N.D.	N.D.	0.06	N.D.	N.D.	N.D.
2.08	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.19-2.25	N.D.	0.24	0.36	2.87	4.49	N.D.	N.D.
2.37-2.39	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
2.49	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.16	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
3.36-3.45	N.D.	N.D.	<0.05	0.08	0.20	N.D.	N.D.
3.59	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.

3.62-3.71	N.D.	0.19	0.19	1.05	0.36	N.D.	N.D.
3.67-3.72(3.77-3.79)Alvocidib	0.50	5.55	5.96	66.40	91.85	0.42	0.75
3.89	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.27	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.51	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.61	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.70	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
4.75-4.91	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
5.03	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
5.14	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
5.32	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
5.58	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.05	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.13	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.34	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.46	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
6.54	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
7.05	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
7.09	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
Total impurity	0.50	6.57	6.72	73.70	99.59	0.42	0.75
Δimpurity		6.07	6.22	73.20	99.09	-0.08	0.25

[00385] Samples were also tested for dissolution. Six capsules were tested for dissolution according to the parameters outlined in Table 23. Each capsule was weighed prior to dissolution testing. The stated amount of dissolution medium was transferred into each vessel in accordance with the capsule strength under analysis. The dissolution medium was allowed to equilibrate in the dissolution bath to 37 °C ± 0.5 °C. The temperature of the dissolution medium checked at the start and end of the dissolution test for each vessel. At each sample point, the solution was transferred directly into an HPLC vial.

Table 23. Dissolution Conditions

Apparatus	USP Apparatus II (Paddles)
Paddle Speed	50 ± 2 rpm
Temperature	37°C ± 0.5°C
Filters	45 µm Polyethylene (Distek) inline filters or GF-D filters
Dissolution Medium	USP pH 6.8 Phosphate buffer
Vessel Volume	900 ml
Sample Times	5, 10, 15, 30, 45 and 60 minutes
Infinity Time	15 minutes (75 minutes sampling point)
Sampling Method - 1	Automated
Infinity spin speed	Automatic (200 rpm)
Sample Volume	1.5 ml (Autosampler)
Pre-sample Rinse	10.0 ml
Recirculate Media	No
Sampling Method - 2	Manual
Infinity spin speed	200 rpm
Sampling Volume	10 ml
Transfer Volume	1.5 ml into HPLC vial for injection

[00386] The following calculations were used to calculate the sample concentration, amount released and percent released, respectively. P811 corresponds to the compound of structure (I).

$$k = \frac{A_{\text{sam}} \times W_{\text{std}} \times P}{A_{\text{std}} \times D_{\text{std}}}$$

Where:

- k = P811 concentration (mg/ml)
Astd = mean area of the P811 peak in the standard chromatograms
Wstd = weight of the reference standard (mg)
P = purity of the reference standard (expressed as a decimal)
Asam = area of the P811 peak in the sample chromatogram
Dstd = dilution of standard (ml)

$$A_n = k[M - (n - 1)(T)] + \sum_{i=1}^{n-1} C_i T$$

Where:

- A_n = cumulative amount of dissolved P811 from the first to the n^{th} transfer (mg)
- k = P811 concentration measured for the current transfer (mg/ml)
- M = initial medium volume (ml) i.e. 900 ml
- n = an index of dissolution transfers where $n=1$ for the 1st transfer (5 minutes), $n=2$ for the 2nd transfer (10 minutes), $n=3$ for the 3rd transfer (15 minutes), $n=4$ for the 4th transfer (30 minutes), $n=5$ for the 5th transfer (45 minutes), $n=6$ for the 6th transfer (60 minutes), $n=7$ for the 7th transfer (75 minutes).
- T = volume of sample removed (ml) i.e. 10.0 ml
- C_i = concentration for the i^{th} dissolution transfer (mg/ml)

For example:

At sampling point 1: $A_1 = kM$

At sampling point 2: $A_2 = k[M - T] + C_1 T$

At sampling point 3: $A_3 = k[M - 2(T)] + C_1 T + C_2 T$

$$\% \text{ Released} = \frac{A_n}{LC} \times 100$$

Where:

- A_n = cumulative amount of dissolved P811 from the first to the n^{th} transfer (mg)
- LC = label claim (16 mg, 4 mg and 1 mg)

[00387] The results of the dissolution testing are shown in FIGs. 9A-9D.

EXAMPLE 14

FORMULATIONS OF COMPOUND OF STRUCTURE (I) AND STABILITIES THEREOF

[00388] The following mixtures of the compound of structure (I) were formulated into 1-mg strength capsules, wherein the percentages are calculated on a weight/weight basis:

Formulation No.	401-01	401-02	401-03	401-04	401-05
Compound of Structure (I)	0.60%	0.60%	0.60%	0.60%	0.60%
Lactose anhydrous	97.40%	98.40%	99.40%		

Cornstarch				98.40%	99.40%
Light anhydrous silicic acid	1%				
Magnesium stearate	1%	1%		1%	

[00389] The following mixture of the compound of structure (I) was formulated into a 4-mg strength capsule, wherein the percentages are calculated on a weight/weight basis:

Formulation No.	402
Compound of Structure (I)	2.47%
Lactose anhydrous	95.53%
Light anhydrous silicic acid	1%
Magnesium stearate	1%

[00390] The following mixtures of the compound of structure (I) were formulated into 16-mg strength capsules, wherein the percentages are calculated on a weight/weight basis:

Formulation No.	401-06	401-08	401-09
Compound of Structure (I)	11.03%	11.03%	11.03%
Lactose anhydrous	86.97%	88.97%	
Cornstarch			87.97%
Light anhydrous silicic acid	1%		
Magnesium stearate	1%		1%

[00391] For manufacturing, a powder blend of compound of structure (I) and the indicated excipients were encapsulated into #4 hydroxypropylmethylcellulose (HPMC) capsules. The resulting capsules were immediate-release capsules.

[00392] Prior to encapsulation into the capsules, the drug product was made by direct blending via triturating the compound of structure (I) into the indicated excipients, followed by filling the capsules on a manual capsule filling machine in 100-capsule plates.

[00393] The capsules were packaged in aluminum blister packaging, with one capsule per blister and seven capsules per blister sheet. Three blisters on each sheet were left empty.

[00394] One-month stability testing of the compositions of the compound of structure (I) was performed at 40 °C and 75% RH, 25 °C and 60% RH or 5 °C. The compositions were tested for HPLC purity, impurities and dissolution. Tables 24-1 to 24-8 report the results of the impurity testing, and show that cornstarch-based

formulations improved capsule stability compared to Formulation Nos. 401-01 and 401-06. Tables 25-1 to 25-8 report the results of the HPLC purity assay. FIGs. 10A to 10X show the results of the dissolution testing conducted according to the dissolution procedure described in Example 13. All of the formulations submitted to dissolution testing met the criteria of Q (compound of structure (I)) = 75% within 45 minutes.

Table 24-1.

RRT	401-01							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
0.51-0.52	N.D.	0.09	0.07	N.D.	0.07	N.D.	0.05	0.05
0.63-0.64	N.D.	<0.05	<0.05	N.D.	<0.05	<0.05	N.D.	N.D.
0.83	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.88	N.D.	N.D.	0.11	0.07	N.D.	N.D.	N.D.	N.D.
1.00 (cmpd of structure (I))	100	99.91	97.13	97.38	99.73	99.61	99.88	99.87
2.2	N.D.	N.D.	0.05	<0.05	N.D.	N.D.	N.D.	N.D.
3.71-3.72 (Alvocidib)	N.D.	<0.05	2.64	2.55	0.2	0.32	0.07	0.07
Imp(%)	0	0.09	2.87	2.62	0.27	0.32	0.12	0.12

Table 24-2.

RRT	401-02							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
0.51-0.52	N.D.	0.1	0.07	N.D.	0.07	0.09	0.08	0.07
0.63	N.D.	<0.05	<0.05	<0.05	<0.05	0.05	0.05	N.D.
0.88	N.D.	N.D.	0.06	0.11	N.D.	N.D.	N.D.	N.D.
1.00 (cmpd of structure (I))	100	99.85	98.63	95	99.77	99.57	99.8	99.86
2.19	N.D.	N.D.	N.D.	0.13	N.D.	N.D.	N.D.	N.D.
3.71 (Alvocidib)	N.D.	<0.05	1.24	4.76	0.16	0.29	0.07	0.06
Imp(%)	0	0.1	1.37	5	0.23	0.43	0.2	0.13

Table 24-3.

RRT	401-03							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
0.51-0.52	N.D.	0.07	0.06	0.07	0.07	0.08	0.07	0.06
0.63-0.64	N.D.	<0.05	N.D.	<0.05	<0.05	<0.05	<0.05	<0.05
0.88	N.D.	N.D.	0.09	0.07	N.D.	N.D.	N.D.	N.D.
1.00 (cmpd of structure (I))	100	99.93	92.33	92.03	99.66	99.45	99.8	99.83
1.11	N.D.	N.D.	N.D.	0.19	N.D.	N.D.	N.D.	N.D.
2.2	N.D.	N.D.	N.D.	0.22	N.D.	N.D.	N.D.	N.D.
3.71 (Alvocidib)	N.D.	<0.05	0.15	7.42	0.27	0.47	0.13	0.11
Imp(%)	0	0.07	0.3	7.97	0.34	0.55	0.2	0.17

Table 24-4.

RRT	401-04							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
0.66	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.72	N.D.	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
1.00 (cmpd of structure (I))	100	100	99.54	99.34	99.88	99.86	99.93	99.92
3.70 (Alvocidib)	N.D.	<0.05	0.46	0.66	0.12	0.14	0.07	0.08
Imp(%)	0	0	0.46	0.66	0.12	0.14	0.07	0.08

Table 24-5.

RRT	401-05							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
0.66	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.
0.72-0.73	N.D.	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05	<0.05
1.00 (cmpd of structure (I))	100	100	99.66	99.41	99.83	99.8	99.87	99.89
3.47-3.48	N.D.	<0.05	N.D.	N.D.	<0.05	<0.05	N.D.	N.D.
3.59	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	<0.05	N.D.
3.69-3.73 (Alvocidib)	N.D.	<0.05	0.34	0.59	0.17	0.2	N.D.	0.11

3.81	N.D.	N.D.	N.D.	N.D.	N.D.	N.D.	0.13	N.D.
Imp(%)	0	0	0.34	0.59	0.17	0.2	0	0.11

Table 24-6.

RRT	401-06							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
0.89	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.
1.00 (cmpd of structure (I))	100	99.93	99.2	98.01	99.81	99.77	99.87	99.88
2.2	N.D.	N.D.	N.D.	0.05	N.D.	N.D.	N.D.	N.D.
3.71 (Alvocidib)	N.D.	0.07	0.8	1.94	0.19	0.23	0.13	0.12
Imp(%)	0	0.07	0.8	1.99	0.19	0.23	0.13	0.12

Table 24-7.

RRT	401-08							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
1.00 (cmpd of structure (I))	100	99.94	99.51	99.53	99.82	99.79	99.87	99.88
3.71 (Alvocidib)	N.D.	0.06	0.49	0.47	0.18	0.21	0.13	0.12
Imp(%)	0	0.06	0.49	0.47	0.18	0.21	0.13	0.12

Table 24-8.

RRT	401-09							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
1.00 (cmpd of structure (I))	100	99.92	99.61	98.98	99.83	99.82	99.87	99.88
2.2	N.D.	N.D.	N.D.	<0.05	N.D.	N.D.	N.D.	N.D.
3.71 (Alvocidib)	N.D.	0.08	0.39	1.02	0.17	0.18	0.13	0.12
Imp(%)	0	0.08	0.39	1.02	0.17	0.18	0.13	0.12

Table 25-1.

	401-01							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
Assay (%)	87.19	80.59	74.38	71.04	77.47	82.29	86.63	82.94

Table 25-2.

	401-02							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
Assay (%)	60.43	74.35	71.53	56.95	71.39	69.59	67.14	63.79

Table 25-3.

	401-03							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
Assay (%)	78.96	85.92	72.91	76.47	80.38	79.82	84.56	84.13

Table 25-4.

	401-04							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
Assay (%)	111.03	97.37	102.4	101.93	105.47	97.26	99.83	97.39

Table 25-5.

	401-05							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
Assay (%)	110.38	116.65	113.04	104.43	108.77	109.76	102.53	98.67

Table 25-6.

	401-06							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
Assay (%)	87.71	85.53	80.88	81.06	85.17	84.29	85.27	82.61

Table 25-7.

	401-08							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
Assay (%)	81.89	85.63	82.14	88.59	80.26	81.44	83.55	80.64

Table 25-8.

	401-09							
	Blend before encapsulation	Initial	40°C		25°C		5°C	
			2W	1M	2W	1M	2W	1M
Assay (%)	105.3	105.59	102.08	88.85	102.9	102.38	102.59	102.89

EXAMPLE 15

PHASE I, PHARMACOKINETIC (PK) AND PHARMACODYNAMIC (PD), DOSE-ESCALATION STUDY OF ORAL COMPOUND OF STRUCTURE (I) ADMINISTERED TO PATIENTS WITH ADVANCED SOLID TUMORS

[00395] Patients with advanced metastatic or progressive solid tumors who were refractory to, or intolerant of, established therapy known to provide clinical benefit for their condition were enrolled. Cohorts of 3-6 patients each received escalating doses of compound of structure (I) using a modified Fibonacci dose escalation approach. Once the optimal dose has been established, additional patients may be enrolled to confirm safety and to explore efficacy.

[00396] This is an ongoing Phase 1, open-label, dose-escalation, safety, PK and PD study. The proposed starting dose and schedule for oral compound of structure (I) was a 1-mg flat dose once daily (QD) for 14 days followed by a 7-day drug-free recovery period (each cycle = 21 days). In the absence of dose-limiting toxicities (DLTs) in the first cohort of at least 3 patients, the dose was increased using a modified Fibonacci dose escalation scheme, and BID dosing commenced according to the dose escalation schedule described in Table 28. The first patient in cohort 6 has been enrolled at 8 mg compound of structure (I) BID. The baseline demographics of the first 14 patients enrolled in the study are described in Table 26.

Table 26. Baseline Demographics (N = 14 ITT)

Median Age	65 (45-78)
Median Number of Prior Therapies	4 (1-10)
ECOG Score	ECOG 0: 1 (7%) ECOG 1: 13 (93%)
Primary Tumor Type	Bladder: 2 Pancreas: 2 Colorectal: 2 Kidney: 1 NSCLC: 1 Prostate: 1 Sarcoma: 1 Skin: 1 Thyroid: 1 Testicular: 1 Vulvar: 1
Gender	Male: 8 (57.1%) Female: 6 (42.9%)

[00397] FIG. 12 is a graph depicting completed cycles on the study through Cohort 5.

[00398] To date, there is no unexplained toxicity, and no evidence of dose-limiting diarrhea or neutropenia. The treatment-emergent adverse events of grade ≥ 3 observed thus far are reported in Table 27.

Table 27. Treatment-emergent adverse events grade ≥ 3.

MedDRA Preferred Term	Grade 3	Grade 4	Grade 5
Anemia	1 (7%)	-	-
Chest Pain	1 (7%)	-	-
Pain	1 (7%)	-	-
Swelling	1 (7%)	-	-
Malignant Pleural Effusion	1 (7%)	-	-
Haematuria	1 (7%)	-	-
Hypoxia	1 (7%)	-	-
Hypotension	1 (7%)	-	-

[00399] Sequential cohorts of 3 patients will continue to be treated with escalating doses according to Table 28 until the MTD is established.

Table 28.

Dose Level	Proposed Daily Dose	Total Daily Dose	Increment from Previous Dose ^a	No. of Patients Per Cohort
-1 ^b	1 mg QOD	N/A	-50%	3-6
1 ^c	1 mg QD	1 mg	Starting Dose	3-6
2	1 mg BID	2 mg	100%	3-6
3	2 mg BID	4 mg	100%	3-6
4	4 mg BID	8 mg	100%	3-6
5	6 mg BID	12 mg	50%	3-6
6	8 mg BID	16 mg	33%	3-6
7 ^d	11 mg BID	22 mg	33%	3-6

- a It is possible for additional and/or intermediate dose levels to be added during the course of the study.
- b Dose level -1 represents a treatment dose for patients requiring a dose reduction from the starting dose level. It will also serve as a lower dose level if the Starting Dose level is initially associated with unexpected or unacceptable toxicity. Please note that the dosing in this instance is a single morning dose every other day (QOD) (no evening doses required).
- c Please note that the dosing in Cohort 1 is a single daily (QD) morning dose (no evening dose required).
- d If clinically indicated, dose levels higher than 11 mg BID may be investigated.

[00400] If a DLT is observed in 1 of 3 patients at a given dose level, up to 3 additional patients will be enrolled and treated at that dose level. When up to 3 additional patients are added to a given dose level, if only 1 out of those 6 patients experiences a DLT, the dose will be increased to the next dose level. If ≥ 2 out of 3-6 patients at a dose level experience DLTs, the dose will be decreased to the previous (lower) dose level and 3 additional patients will be enrolled at that dose level.

[00401] If 0 or 1 patient in any of the 6 patients experience a DLT, but the next higher dose level has already been studied, then the current dose will be declared the MTD and the study will advance to the expansion cohort.

[00402] The MTD is defined as the dose at which ≤ 1 of 6 patients experience a DLT during Cycle 1 with the next higher dose having at least 2 of 3 to 6 patients experiencing a DLT during Cycle 1.

[00403] Once the MTD has been established, 20 additional patients will be enrolled at the MTD. Data collected from patients enrolled at the MTD will be used to confirm safety, explore potential biomarkers, and evaluate potential signals of compound of structure (I) activity.

[00404] All patients may continue to receive compound of structure (I) in 21-day cycles (14 days of active treatment) at the same dose given during Cycle 1 until they

experience unacceptable toxicity or unequivocal disease progression. No intra-patient escalation of the compound of structure (I) dose is permitted during the escalation phase until MTD is established.

[00405] Patients met all of the following inclusion criteria:

1. Have a histologically confirmed diagnosis of advanced metastatic or progressive solid tumor excluding tumor types with rapid cell turnover, *i.e.*, small cell cancer (lung and extra pulmonary), inflammatory breast cancer (IBC), medulloblastoma, neuroblastoma and melanoma with extensive liver metastasis ($\geq 50\%$ of the liver involved; patients with melanoma and metastasis to $< 50\%$ of the liver were eligible)
2. Be refractory to, or intolerant of, established therapy known to provide clinical benefit for their condition
3. Have one or more tumors measurable or evaluable as outlined by the Response Evaluation Criteria in Solid Tumors (RECIST) v1.1
4. Have an Eastern Cooperative Oncology Group (ECOG) performance status of ≤ 1
5. Have a life expectancy ≥ 3 months
6. Be ≥ 18 years of age
7. Have a negative pregnancy test (if female of childbearing potential)
8. Have acceptable liver function:
 - a) Bilirubin ≤ 1.5 x upper limit of normal (ULN) (unless associated with Gilbert syndrome)
 - b) Aspartate aminotransferase (AST/SGOT), alanine aminotransferase (ALT/SGPT) and alkaline phosphatase ≤ 2.5 x ULN*
* if liver metastases were present, then 3x ULN was allowed
9. Have acceptable renal function: calculated creatinine clearance ≥ 30 mL/min
10. Have acceptable hematologic status:
 - c) Granulocyte ≥ 1500 cells/mm³
 - d) Platelet count $\geq 100,000$ (plt/mm³)
 - e) Hemoglobin ≥ 8 g/dL
11. Have acceptable coagulation status:
 - f) Prothrombin time (PT) within 1.5x normal limits

- g) Activated partial thromboplastin time (aPTT) within 1.5x normal limits
12. Be nonfertile or agree to use an adequate method of contraception. Sexually active patients and their partners used an effective method of contraception (hormonal or barrier method of birth control; or abstinence) prior to study entry and for the duration of study participation and for at least 3 months (males) and 6 months (females) after the last study drug dose.
 13. Have read and signed the Institutional Review Board (IRB)-approved informed consent form (ICF) prior to any study-related procedure.

[00406] Patients meeting any one of the following exclusions criteria were prohibited from participating in the study:

1. History of congestive heart failure (CHF); cardiac disease, myocardial infarction within the past 6 months prior to Cycle 1/Day 1; left ventricular ejection fraction (LVEF) <45% by echocardiogram (ECHO), unstable arrhythmia, or evidence of ischemia on electrocardiogram (ECG) within 14 days prior to Cycle 1/Day 1
2. Have a corrected QT interval (using Fridericia's correction formula) (QTcF) of >450 msec in men and >470 msec in women
3. Have a seizure disorder requiring anticonvulsant therapy
4. Presence of symptomatic central nervous system metastatic disease or disease that requires local therapy such as radiotherapy, surgery, or increasing dose of steroids within the prior 2 weeks
5. Have severe chronic obstructive pulmonary disease with hypoxemia (defined as resting O₂ saturation of ≤90% breathing room air)
6. Have undergone major surgery within 2 weeks prior to Cycle 1/Day 1
7. Have active, uncontrolled bacterial, viral, or fungal infections, requiring systemic therapy
8. Are pregnant or nursing
9. Received treatment with radiation therapy, surgery, chemotherapy, or investigational therapy within 28 days or 5 half-lives, whichever occurs first, prior to study entry (6 weeks for nitrosoureas or mitomycin C)
10. Are unwilling or unable to comply with procedures required in this protocol

11. Have known infection with human immunodeficiency virus, hepatitis B, or hepatitis C. Patients with history of chronic hepatitis that is currently not active are eligible
12. Have a serious nonmalignant disease (eg, hydronephrosis, liver failure, or other conditions) that could compromise protocol objectives in the opinion of the Investigator and/or the Sponsor
13. Are currently receiving any other investigational agent
14. Have exhibited allergic reactions to a similar structural compound, biological agent, or formulation
15. Have malabsorption conditions (e.g., Crohn's disease) or have undergone significant surgery to the gastrointestinal tract that could impair absorption or that could result in short bowel syndrome with diarrhea due to malabsorption.

DLT was defined as any one of the following events observed in cycle 1, regardless of investigator attribution, unless there was a clear alternative explanation:

1. Grade 3 or greater febrile neutropenia
2. Grade 4 neutropenia for ≥ 7 consecutive days
3. Grade 4 thrombocytopenia or Grade 3 thrombocytopenia with clinically significant bleeding or that requires a platelet transfusion
4. Grade 3 or 4 nonhematologic AEs will be considered dose limiting, regardless of duration aside from the specific parameters described herein
5. Grade 4 nausea, vomiting, or diarrhea, regardless of duration
6. Dosing delays >1 week due to treatment-emergent adverse events (TEAEs) or related severe laboratory test values
7. Any AST and ALT elevation $>3x$ ULN (if baseline value was normal) or $\geq 3x$ the baseline value (if baseline value was abnormal) accompanied by serum bilirubin levels $>2x$ ULN
8. Any Grade ≥ 3 electrolyte disturbances (eg, hyperkalemia, hypophosphatemia, hyperuricemia) that do not resolve within <72 hours
9. Any Grade ≥ 3 elevations in creatinine
10. Any Grade 5 toxicity

[00407] Plasma PK parameters of compound of structure (I) and alvocidib were evaluated in Cohorts 1-5 at specific timepoints during the study. Blood was collected

from patients in Cohorts 1-5 according to the pharmacokinetic sampling schedule described in Table 29.

Table 29. Pharmacokinetic Sampling Schedule

Cycle No.	CYCLE 1						CYCLE 2	End of Study
	AM Day 1	PM ^a Day 1	AM Day 2	AM Day 14	PM ^a Day 14	AM Day 15	AM Day 1	
0 (pre-dose)	X		X ^b	X ^b		X ^b	X	X
0.5	X	X		X	X			
1	X	X		X	X			
2	X	X		X	X			
4	X			X				
8 ^c	X			X				

- a No evening (PM) samples were collected from patients enrolled in the first dose cohort receiving compound structure (I) as a once daily (QD) morning dose.
- b Approximately 24 hours after taking the previous days' morning dose and prior to taking current days' dose (*i.e.*, sampling on Cycle 1/Day 2 would be performed 24 hrs after taking the morning dose on Day 1 and before taking the morning dose on Day 2)
- c The 8-hour samples were collected just prior to taking that day's evening dose (for patients receiving compound of structure (I) BID), or 8 hours after taking that day's dose (for patients receiving compound of structure (I) QD).

[00408] PK parameters were estimated using standard noncompartmental methods.

Actual sample collection times were used rather than scheduled collection times.

Plasma concentrations below the limit of quantification were treated as 0. Imbedded missing plasma concentrations (*e.g.*, missing values between two observed values) were estimated using linear extrapolation. This is consistent with using the trapezoidal rule to calculate AUC. Other missing plasma concentrations were excluded from calculations to estimate PK parameters.

[00409] FIGs. 11A and 11B are graphs of plasma alvocidib concentration (ng/mL) versus time, and show the concentration of alvocidib in the plasma of patients in Cohort 1 on days 1 and 14, respectively, following daily oral QD dosing with a 1-mg strength capsule containing Formulation No. 401-01. Subject 104 showed some accumulation of alvocidib after 24 hours on day 14. Subject 102 was discontinued prior to day 14 dosing.

[00410] FIGs. 11C and 11D are graphs of plasma alvocidib concentration (ng/mL) versus time, and show the concentration of alvocidib in the plasma of patients in cohort

2 on days 1 and 14, respectively, following daily oral BID dosing with a 1-mg strength capsule containing Formulation No. 401-01. Only alvocidib was detectable at 1 mg BID, and no compound of structure (I) was detected at any timepoint for any sample. No drug was detectable (less than 1.0 ng/mL of alvocidib) by 8 hours and again at 24 hours in any subject on day 1. However, there was detectable accumulation of alvocidib on day 14 for subjects 201 and 202 (average = 2.39 ng/mL), suggesting that BID dosing helped to maintain drug levels by day 14.

[00411] FIGs. 11E and 11F are graphs of plasma alvocidib concentration (ng/mL) versus time, and show the concentration of alvocidib in the plasma of patients in Cohort 5 on days 1 and 14, respectively, following daily oral BID dosing with 6 mg of Formulation No. 401-01. Table 30 reports T_{max} , C_{max} and $AUC_{(0-24)}$ of alvocidib for patients in Cohort 5 on days 1 and 14 of cycle 1.

Table 30.


Cohort 5 6mg BID	Cycle 1					
	Day 1			Day 14		
Subject (#)	T_{max} (Hours)	C_{max} (ng/mL)	$AUC_{(0-24)}$ (ng*h/mL)	T_{max} (Hours)	C_{max} (ng/mL)	$AUC_{(0-24)}$ (ng*h/mL)
501	0.5	28	230	0.5	40.2	192
502	2	19	208	2	30.4	516
503	0.5	29.3	376	0.5	51.7	407
Mean	1.0	25.4	271.3	1.0	40.8	371.7
SD	0.9	5.6	91.3	0.9	10.7	164.9

[00412] FIG. 11G is a graph of alvocidib (ng/mL) versus cohort, and shows the mean C_{max} of alvocidib on day 1 and day 14 following daily oral QD dosing with a 1-mg strength capsule containing Formulation No. 401-01. FIG. 11H is a graph of alvocidib (ng*hr/mL) versus cohort, and shows the area under the curve (AUC) of alvocidib on day 1 (AUC_{0-8}) and day 14 (AUC_{0-8} and AUC_{0-24}) following daily oral BID dosing with a 1-mg strength capsule containing Formulation No. 401-01. There was no detectable compound of structure (I) at any timepoint. Cohort 2 showed marked increase in average C_{max} and AUC from day 1 to day 14, illustrating the impact of BID versus QD dosing. The C_{max} for Cohort 5 increased by 46% compared to Cohort 4 on day 1, and by 69% for day 14. The corresponding increase in AUC was 52% on day 1 and 30% on day 14.

[00413] FIG. 11I is a graph of mean concentration of alvocidib (nM) versus time, and shows the mean concentration of alvocidib in plasma of Cohort 5 patients over a 24-hour period. By administering alvocidib as compound of structure (I), alvocidib can be given at a lower dose over a longer time, with less toxicity and similar exposure.

EXAMPLE 16

ALTERNATIVE SYNTHESIS OF STRUCTURE (I)



Scheme 3.

[00414] A-10 was obtained according to Steps 1.1 and 2.1A, depicted in Scheme 2 and described in Example 10. A-10 (100 mg) was substantially dissolved in methanol (1 ml) at 50 °C. A solution of maleic acid (12.2 mg, 0.5 equiv) in methanol (1.5 ml) was added dropwise to the mixture of A-10 in methanol, followed by acetone (2.5 ml). The resulting reaction mixture was stirred for one hour at room temperature, and then filtered to obtain A-5 (77.4 mg) having a crystallinity of 94%. Residual maleic acid was confirmed by $^1\text{H NMR}$ (0.01 proton).

EXAMPLE 17

A PHASE II STUDY OF ORAL COMPOUND OF STRUCTURE (I) ADMINISTERED ONCE DAILY
FOR 21 DAYS TO PATIENTS WITH METASTATIC CASTRATE-RESISTANT PROSTATE
CANCER

[00415] This is a Phase 2, open-label, non-randomized, Simon 2-stage design study to establish the efficacy and safety of compound of structure (I) (*e.g.*, Form B of

compound of structure (I) taken once daily for 21 days of a 28-day cycle in patients with metastatic castration-resistant prostate cancer who have progressed on frontline treatment with androgen signaling inhibitors. A biopsy sub-study in 20 patients will enable the evaluation of tissue biomarkers in a subset of patients.

[00416] Sixty (60) patients will be enrolled. Data will be used to assess efficacy, confirm safety, and explore correlative potential biomarkers.

[00417] All patients may continue to receive compound of structure (I) in 28-day cycles (21 days of active treatment) at the same dose given during Cycle 1 until they experience unacceptable toxicity or unequivocal disease progression.

[00418] Patients must meet all of the following inclusion criteria to be eligible:

1. Male patients who also have histologically or cytologically confirmed adenocarcinoma of the prostate; AND:
 - a) Be castrate-resistant on treatment with androgen deprivation therapy (ADT) (or status post bilateral orchiectomy) and with testosterone levels of less than (<) 50 nanogram per deciliter (50ng/dL, equivalent to 1.7 nmol/L); AND:
 - b) Have radiographic progression according to Prostate Cancer Clinical Trials Working Group 3 (PCWG3) criteria, while being treated with abiraterone acetate or enzalutamide in combination with ADT
2. Be refractory to, or intolerant of, established therapy known to provide clinical benefit for their condition
3. Have one or more tumors measurable as outlined by the Response Evaluation Criteria in Solid Tumors (RECIST) v1.1
4. Willingness to undergo two (2) on-study biopsies (biopsy sub-study cohort only)
5. Have an Eastern Cooperative Oncology Group (ECOG) performance status of ≤ 1
6. Have a life expectancy ≥ 3 months
7. Be ≥ 18 years of age
8. Have acceptable liver function:
 - a) Bilirubin ≤ 1.5 x upper limit of normal (ULN) (unless associated with Gilbert syndrome)

- b) Aspartate aminotransferase (AST/SGOT), alanine aminotransferase (ALT/SGPT) and alkaline phosphatase $\leq 2.5 \times \text{ULN}^*$
*If liver metastases are present, then $3 \times \text{ULN}$ is allowed.
- 9. Have acceptable renal function: calculated creatinine clearance $\geq 30 \text{ mL/min}$
- 10. Have acceptable hematologic status:
 - a) Granulocyte $\geq 1500 \text{ cells/mm}^3$
 - b) Platelet count $\geq 100,000 \text{ (plt/mm}^3\text{)}$
 - c) Hemoglobin $\geq 8 \text{ g/dL}$
- 11. Have acceptable coagulation status:
 - a) Prothrombin time (PT) within $1.5 \times$ normal limits
 - b) Activated partial thromboplastin time (aPTT) within $1.5 \times$ normal limits
- 12. Be nonfertile or agree to use an adequate method of contraception. Sexually active patients and their partners must use an effective method of contraception (hormonal or barrier method of birth control; or abstinence) prior to study entry and for the duration of study participation and for at least 3 months (males) and 6 months (females) after the last study drug dose. Should a woman become pregnant or suspect she is pregnant while her partner is participating in this study, she should inform her treating physician immediately.
- 13. Have read and signed the Institutional Review Board (IRB)-approved informed consent form (ICF) prior to any study-related procedure. (In the event that the patient is rescreened for study participation or a protocol amendment alters the care of an ongoing patient, a new ICF must be signed.)

[00419] Patients meeting any one of these exclusion criteria will be prohibited from participating in this study:

- 1. History of congestive heart failure (CHF); cardiac disease, myocardial infarction within the past 6 months prior to Cycle 1/Day 1
- 2. Have a corrected QT interval (using Fridericia's correction formula) (QTcF) of $>450 \text{ msec}$ in men and $>470 \text{ msec}$ in women
- 3. Have a seizure disorder requiring anticonvulsant therapy
- 4. Presence of symptomatic central nervous system metastatic disease or disease that requires local therapy such as radiotherapy, surgery, or increasing dose of steroids within the prior 2 weeks

5. Have severe chronic obstructive pulmonary disease with hypoxemia (defined as resting O₂ saturation of $\leq 90\%$ breathing room air)
6. Have undergone major surgery within 2 weeks prior to Cycle 1/Day 1
7. Have active, uncontrolled bacterial, viral, or fungal infections, requiring systemic therapy
8. Are pregnant or nursing
9. Received treatment with radiation therapy, surgery, chemotherapy, or investigational therapy within 28 days or 5 half-lives, whichever occurs first, prior to study entry (6 weeks for nitrosoureas or mitomycin C)
10. Are unwilling or unable to comply with procedures required in this protocol
11. Have known infection with human immunodeficiency virus, hepatitis B, or hepatitis C. Patients with history of chronic hepatitis that is currently not active are eligible
12. Have a serious nonmalignant disease (*e.g.*, hydronephrosis, liver failure, or other conditions) that could compromise protocol objectives in the opinion of the Investigator and/or the Sponsor
13. Are currently receiving any other investigational agent
14. Have exhibited allergic reactions to a similar structural compound, biological agent, or formulation
15. Have malabsorption conditions (*e.g.*, Crohn's disease, *etc.*) or have undergone significant surgery to the gastrointestinal tract that could impair absorption or that could result in short bowel syndrome with diarrhea due to malabsorption.

[00420] Enrolled patients will receive compound of structure (I) (*e.g.*, given as a 1-mg capsule containing Formulation No. 401-01, wherein the compound of structure (I) is Form B of the compound of structure (I)), administered once daily (QD) for the first 21 days of a 28-day cycle. Patients who successfully complete a 4-week treatment cycle without evidence of significant treatment-related toxicity or progressive disease will continue to receive treatment with the same dose and dosing schedule.

[00421] Efficacy assessments will be performed based on PCWG3-modified RECIST v1.1 guidelines, to include the assessment of objective response rate (ORR), DoR, type of response (*e.g.*, complete remission, partial remission, stable disease), and time to progression. The ORR is defined as the percent of patients with CR or PR

according to PCWG3-modified RECIST v1.1 criteria, relative to the Response Evaluable population. ORR will be summarized by number and percentage of patients meeting the definition of ORR along with the corresponding exact 95% confidence intervals.

[00422] Tolerance and toxicity of oral compound of structure (I) will be assessed through evaluation of physical examinations, vital signs, laboratory parameters, AEs including DLTs, and all causes of mortality.

[00423] Incidence rates of treatment-emergent adverse events (TEAEs) will be summarized within each dose level at the Medical Dictionary for Regulatory Activities (MedDRA) preferred term and primary system organ class levels. Similar summaries will be made for subsets of AEs such as (1) those judged by the Investigator to be related to study treatment, and (2) serious adverse events (SAEs).

[00424] Other routine safety assessments (*e.g.*, clinical laboratory parameters and vital signs) will be summarized by compound of structure (I) dose level using mean, standard deviation, median, minimum, and maximum changes from baseline values.

[00425] PD parameters and assessment of potential tumor and peripheral blood biomarkers including, but not limited to, CDK9-related genes (including c-Myc) in biopsy and CTC samples; Phospho-AR; PhosphoRNAPol2 on biopsy and PBMC samples; serum PSA.

[00426] Blood will be collected from all patients for evaluation of compound of structure (I) pharmacodynamics and potential biomarkers. Biopsy samples will be taken at baseline (prior to dosing on Cycle 1/Day 1) and at the end of cycle two (2) in a subset of patients participating in the biopsy sub-study.

[00427] The most recent archived tumor tissue (primary and metastatic site(s), if available) will be requested from all patients to assess potential biomarkers.

[00428] All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification or the attached Application Data Sheet are incorporated herein by reference, in their entirety to the extent not inconsistent with the present description.


[00429] From the foregoing it will be appreciated that, although specific embodiments of the disclosure have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the disclosure.

[00430] Reference to any prior art in the specification is not an acknowledgement or suggestion that this prior art forms part of the common general knowledge in any jurisdiction or that this prior art could reasonably be expected to be combined with any other piece of prior art by a skilled person in the art.

[00431] By way of clarification and for avoidance of doubt, as used herein and except where the context requires otherwise, the term "comprise" and variations of the term, such as "comprising", "comprises" and "comprised", are not intended to exclude further additions, components, integers or steps.

CLAIMS


1. A crystalline form of a compound having structure (II):



- the crystalline form being Form B and characterized by an x-ray powder diffraction pattern comprising at least three peaks at 2-theta angles selected from the group consisting of $4.8 \pm 0.2^\circ$, $10.8 \pm 0.2^\circ$, $13.7 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$, $20.0 \pm 0.2^\circ$ and $24.6 \pm 0.2^\circ$.
2. The crystalline form of claim 1, wherein the crystalline form is substantially pure.
3. The crystalline form of claim 1 or 2, characterized by an x-ray powder diffraction pattern comprising at least four peaks at 2-theta angles selected from the group consisting of $4.8 \pm 0.2^\circ$, $10.8 \pm 0.2^\circ$, $13.7 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$, $20.0 \pm 0.2^\circ$ and $24.6 \pm 0.2^\circ$.
4. The crystalline form of any one of claims 1-3, characterized by an x-ray powder diffraction pattern comprising at least five peaks at 2-theta angles selected from the group consisting of $4.8 \pm 0.2^\circ$, $10.8 \pm 0.2^\circ$, $13.7 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$, $20.0 \pm 0.2^\circ$ and $24.6 \pm 0.2^\circ$.
5. The crystalline form of any one of claims 1-4, characterized by an x-ray powder diffraction pattern comprising peaks at the following 2-theta angles: $10.8 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$ and $20.0 \pm 0.2^\circ$.
6. The crystalline form of any one of claims 1-5, characterized by an x-ray powder diffraction pattern comprising peaks at the following 2-theta angles: $4.8 \pm 0.2^\circ$, $10.8 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$ and $20.0 \pm 0.2^\circ$.
7. The crystalline form of any one of claims 1-6, characterized by an x-ray powder diffraction pattern comprising peaks at the following 2-theta angles: $4.8 \pm 0.2^\circ$, $10.8 \pm 0.2^\circ$, $13.7 \pm 0.2^\circ$, $14.9 \pm 0.2^\circ$ and $20.0 \pm 0.2^\circ$.

8. The crystalline form of any one of claims 1-7, characterized by a differential scanning calorimetry thermogram comprising an endothermic peak at about 264 °C.
9. A pharmaceutical composition comprising a crystalline form of any one of claims 1-8 and a pharmaceutically acceptable carrier or excipient.
10. The pharmaceutical composition of claim 9, formulated for oral administration.
11. The pharmaceutical composition of claim 9 or 10, further comprising a glidant and a lubricant.
12. A pharmaceutical composition comprising:
 from about 0.5 weight percent to about 11.25 weight percent of the crystalline form of any one of claims 1-7, and
 from about 85 weight percent to about 99.5 weight percent of a pharmaceutically acceptable carrier or excipient.
13. A method for preparing the crystalline form of any one of claims 1-8, the method comprising:

contacting amorphous compound having the following structure (I):




(I)

or a tautomer or zwitterionic form thereof, with an acid in a solvent, thereby preparing the crystalline form of any one of claims 1-8.

14. A method for preparing the crystalline form of any one of claims 1-8, the method comprising:


contacting a compound having the following structure (V):




or a tautomer, salt or zwitterionic form thereof, with an acid in a solvent, thereby preparing the crystalline form of any one of claims 1-8.

15. A method for preparing the crystalline form of any one of claims 1-8, the method comprising:

(a) contacting a compound having the following structure (IV):



or a tautomer or salt thereof, with di-*tert*-butylhalophosphonate in the presence of an amine base, thereby forming a compound having the following structure (V):




or a tautomer or salt thereof; and

(b) contacting the compound having structure (V), or a tautomer or salt thereof, in an organic solvent, with an acid having at least one pK_a value that is less than about 5 and a pK_a value of greater than about 1 in an organic solvent, thereby preparing the crystalline form of any one of claims 1-8.


16. A method for preparing the crystalline form of any one of claims 1-8, the method comprising:

(a) contacting a compound having the following structure (IV):



(IV)


or a tautomer or salt thereof, with di-*tert*-butylhalophosphonate in the presence of an amine base, thereby forming a compound having the following structure (V):



(V)

or a tautomer or salt thereof;

(b) contacting the compound having structure (V), or a tautomer or salt thereof, with an acid having a pK_a value of less than about 1, thereby forming a salt of a compound having the following structure (I):



(I)

(c) contacting the salt of a compound having structure (I) with a base, thereby forming amorphous compound having structure (I); and

- (d) contacting the amorphous compound having structure (I), in an organic solvent, with an acid having at least one pK_a value that is less than about 5 and a pK_a value of greater than about 1, thereby preparing the crystalline form of any one of claims 1-8.
17. A combination comprising a crystalline form of any one of claims 1-8, and an additional therapeutic agent.
18. The combination of claim 17, further comprising a pharmaceutically acceptable carrier or excipient.




FIG. 1




FIG. 2





FIG. 3

4/46

**FIG. 4**

5/46




FIG. 5

6/46




FIG. 6A




FIG. 6B




FIG. 7




FIG. 8




FIG. 9A

10/46




FIG. 9B




FIG. 9C

12/46




FIG. 9D




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 10D




FIG. 10E




FIG. 10F




FIG. 10G




FIG. 10H




FIG. 10I




FIG. 10J




FIG. 10K

24/46




FIG. 10L

25/46




FIG. 10M

26/46




FIG. 10N

27/46




FIG. 100




FIG. 10P

29/46




FIG. 10Q

30/46




FIG. 10R




FIG. 10S




FIG. 10T




FIG. 10U

34/46




FIG. 10V




FIG. 10W




FIG. 10X

37/46




FIG. 11A




FIG. 11B

39/46

Cohort 2 (1 mg BID) C1D1 (Log scale)




FIG. 11C

40/46

Cohort 2 (1 mg BID) C1D14 (Log scale)




FIG. 11D

41/46




FIG. 11E




FIG. 11F

43/46




FIG. 11G

44/46




FIG. 11H

45/46

Compound of Structure (I) 6mg BID




FIG. 111

46/46




FIG. 12