



HU000029987T2

(19) **HU**(11) Lajstromszám: **E 029 987**(13) **T2****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 11 165011**(51) Int. Cl.: **A61K 9/00**

(2006.01)

(22) A bejelentés napja: **2007. 07. 27.**

(96) Az európai bejelentés bejelentési száma:

EP 20070165011

(97) Az európai bejelentés közzétételi adatai:

EP 2404593 A1 **2012. 01. 11.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2404593 B1 **2016. 05. 11.**

(30) Elsőbbségi adatok:

06291236	2006. 07. 28.	EP
494493	2006. 07. 28.	US
07112097	2007. 07. 09.	EP
PCT/IB2007/053441	2007. 07. 10.	WO

(73) Jogosult(ak):

Santen SAS, 91000 Evry (FR)

(74) Képviselő:

**ARINOVA Szabadalmi és Védjegy Iroda,
Budapest**

(72) Feltaláló(k):

Rabinovich-Guilatt, Laura, 60920, Kadima (IL)
Lambert, Grégory, 92290, Châtenay-Malabry (FR)
Lallemand, Frédéric, 94260, FRESNES (FR)
Philips, Betty, 92160, Antony (FR)

(54)

Kvaterner ammónium vegyületeket tartalmazó készítmény

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 2 404 593 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

11.05.2016 Bulletin 2016/19

(51) Int Cl.:

A61K 9/00 (2006.01)

(21) Application number: **11165011.5**

(22) Date of filing: **27.07.2007**

(54) **Composition containing quaternary ammonium compounds**

Zusammensetzungen mit quaternären Ammoniumverbindungen

Composition contenant des composés d'ammonium quaternaires

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE
SI SK TR**

(30) Priority: **28.07.2006 EP 06291236**

28.07.2006 US 494493

09.07.2007 EP 07112097

10.07.2007 PCT/IB2007/053441

(43) Date of publication of application:

11.01.2012 Bulletin 2012/02

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:

07787998.9 / 2 049 079

(73) Proprietor: **Santen SAS**

91000 Evry (FR)

(72) Inventors:

- **Rabinovich-Guilatt, Laura**
60920, Kadima (IL)
- **Lambert, Grégory**
92290, Châtenay-Malabry (FR)
- **Lallemand, Frédéric**
94260, FRESNES (FR)
- **Philips, Betty**
92160, Antony (FR)

(74) Representative: **Icosa**

83 avenue Denfert-Rochereau
75014 Paris (FR)

(56) References cited:

EP-A2- 0 642 782 WO-A-95/31958
WO-A-97/25975 GB-A- 633 175
US-A- 3 954 644 US-A- 5 951 988

EP 2 404 593 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] This invention relates to pharmaceutical, ophthalmic or cosmetic compositions containing quaternary ammonium compounds, more preferably to ophthalmic emulsions being useful for eye care or for the treatment of eye conditions. This invention also relates to compositions including at least one quaternary ammonium compound as cationic agent.

[0002] Quaternary ammonium compounds are organic compounds usually used as an antiseptic or antimicrobial agent. For example, benzalkonium chloride is a nitrogenous cationic surface-acting agent belonging to the quaternary ammonium group. Benzalkonium chloride is generally defined as a mixture of compounds of general formula $C_6H_5CH_2N(CH_3)_2RCl$, wherein R is a C12-C24 alkyl group.

[0003] Benzalkonium chloride, as usually provided by the manufacturers wanting to comply with the European and/or American Pharmacopeia, is a mixture of n-alkyl dimethyl benzyl ammonium chlorides of various alkyl chain lengths. For example, FeF Chemicals A/S (Denmark) supplies, under reference 8100301U (BAK USP/NF), a mixture of three alkyl dimethyl benzyl ammonium chlorides including : (1) 60-70% of C₁₂-alkyl dimethyl benzyl ammonium chloride (2) 30-40% of C₁₄-alkyl dimethyl benzyl ammonium chloride, and less than 5% of C₁₆-alkyl dimethyl benzyl ammonium chloride

[0004] Benzalkonium chloride, as a mixture of alkyl dimethyl benzyl ammonium having various alkyl chain lengths is used as preservative agent in topical ophthalmic products. Benzalkonium chloride also has cationic agent properties, and was used as cationic agents for emulsions, especially ophthalmic emulsions.

[0005] When mixtures of benzalkonium chlorides having various alkyl chain lengths are used in emulsions, they may act both as preservative agents and cationic agents.

[0006] The Applicant worked on long chain quaternary ammonium compounds, and noticed that the length of the alkyl chain was important with regards to the function performed by the quaternary ammonium compounds: acting on the length of the alkyl chain resulted in enhancing or reducing the cationic power of the quaternary ammonium compounds. Without wanting to be linked by any theory, the Applicant observed on working on oil-in-water emulsions, that long chain quaternary ammonium compounds are preferentially localized at the oil/water interface of the emulsions, resulting in (1) emulsions with higher zeta potential and (2) more stable emulsions. As quaternary ammonium may be considered as undesirable or toxic, it is thus a goal of this invention to provide cationic composition having a reduced content of quaternary ammonium compound.

[0007] The Applicant also observed that, in emulsions, quaternary ammonium compounds having long alkyl chains, for example quaternary ammonium compounds having C14-C18 alkyl chains, when compared to C12-alkyl chains, did not have a good bactericidal activity, whereas they conferred a greatest cationic power.

[0008] Moreover, the Applicant observed that long chain quaternary ammonium compounds were present preferentially at the oil/water interface of the emulsion droplets, and less in the aqueous phase. The fact that quaternary ammonium compounds may be present in the aqueous phase in a very small amount only, or not present, leads to a loss of preservative effect or poor preservative effect, as well as to less toxic emulsions.

[0009] Thus, one of the goals of this invention is to provide stable cationic emulsions comprising a reduced amount of benzalkonium chlorides, and still using said benzalkonium chlorides as a source, or the only source, of cationic agents, said emulsions being preserved or not.

Another goal of the present invention is to provide emulsions that are not toxic, even if they comprise quaternary ammonium compounds.

Preferably, the emulsions of the invention are useful for ophthalmic purposes.

[0010] In the meaning of this invention,

"Cationic emulsions" are emulsions having a positive zeta potential, preferably a zeta potential higher to 10 mV; "alkyl" means a saturated or unsaturated hydrocarbon chain; "long alkyl chain" are alkyl moieties having at least 12 carbon atoms; "quaternary ammonium compounds" refer to ammonium halides in which the nitrogen atom is substituted by only one or at least one alkyl group having at least 12 carbon atoms ; quaternary ammonium compounds also, but not exclusively, include n-alkyl dimethyl benzyl ammonium chloride also called benzalkonium chloride (hereinafter also referred to as BAK or ADBAC); n-alkyl dimethyl benzyl ammonium bromide; n-alkyl trimethyl ammonium bromide (also referred to as ATAB), n-alkyl meaning an alkyl group of at least 12 carbon atoms; "C14-alkyl ammonium halides" means ammonium halides in which the nitrogen atom of the ammonium group is substituted by at least one alkyl group having at least 14 carbon atoms.

"BAK C12" refers to benzododecinium chloride (CAS 139-07-1); "BAK C14" refers to myristalkonium chloride (CAS 139-08-2); "BAK C16" refers to cetalkonium chloride or CKC (CAS 122-18-9) ;

"ATAB C12" refers to lauryl trimethyl ammonium bromide (CAS 1119-94-4); "ATAB C14" refers to Myristil trimethyl ammonium bromide (CAS 1119-97-7); "ATAB C16" or "CTAB" refers to Cetyl trimethyl ammonium bromide (CAS 57-09-0),

"MCT" means Medium chain triglycerides; for the experimentation, Mygliol 812 (Sasol, Germany) was the MCT used; "ND" means "not determined".

[0011] The invention relates to a cationic oil-in-water emulsion comprising an ammonium halide composition as cationic

agent. The ammonium halide composition includes in a preferred embodiment only one ammonium halide, which is C16-alkyl quaternary ammonium halide.

The oil-in-water emulsion of the invention presents the advantages to be very stable and non toxic, compared to emulsions comprising benzalkonium chlorides.

By cationic oil-in water emulsion is understood an oil-in-water emulsion having a positive zeta potential. The emulsion of the invention has a positive zeta potential and is stable, which means that it keeps a positive zeta potential overtime. In a preferred embodiment, the oil-in-water emulsion according to the invention includes droplets of size 100 to 500 nm, preferably 200 to 300 nm.

[0012] In a preferred embodiment, the oil-in-water emulsion of the invention is useful for eye care or for the treatment of eye diseases or eye conditions.

In a preferred embodiment of the present invention, eye diseases or eye conditions means a dry eye condition.

A diminution of the quantity of tears produced and distributed through the lachrymal ducts or a decrease in the stability of the tear film produced, results in a condition of the eye referred to as dry eye. Dry eye conditions act to decrease visual acuity, produce discomfort, ranging from mild to intense and eventually, if allowed to remain untreated and uncorrected, result in permanent damage with degradation of the exposed ocular tissues, with a complete breakdown of corneal tissue necessitating, in the extreme, corneal transplants.

The symptoms associated with dry eye are often exacerbated with subjects using contact lens.

A dry eye condition in this context therefore refers to dry eye accompanying lacrimal fluid reduction, tear deficiency, xerosis of the eye, Sjogren's syndrome, keratoconjunctivitis sicca (KCS), atopic keratoconjunctivitis sicca (AKC), vernal keratoconjunctivitis (VKC), Stevens-Johnson syndrome, pemphigoid of the eye, marginal blepharitis, failure in eyelids closure, or sensory nerve numbness, dry eye accompanying allergic conjunctivitis, dry eye after viral conjunctivitis, dry eye after cornea surgery including laser in situ keratomileusis (LASIK), dry eye after cataract surgery, dry eye associated with contact lens wearing, or dry eye associated with VDT tasks

[0013] More preferably, the oil-in-water emulsion according to the invention comprises:

- a) an oil phase,
- b) 0.0005% to 0.1 %w/w preferably 0.001 to 0.02 % w/w of the weight of the emulsion of a composition of ammonium halides according to the invention, as described hereabove, to the weight of the emulsion
- c) surfactants,
- d) optionally antioxidants, tonicity, viscosifying, pH adjusting, buffering, preservative, solubilizers, chelating or thickener agents,
- e) water.

[0014] In an embodiment of the present invention, said composition of ammonium halides comprises at least one ammonium quaternary ammonium halide, in which the nitrogen atom of the ammonium group is substituted by only one or at least one alkyl group having at least 12 carbon atoms, said composition including:

- at least 20% in weight by weight of the total composition of ammonium halides in which the nitrogen atom is substituted by only one or at least one alkyl group having at least 14 carbon atoms, preferably 14 or 16 carbon atoms and
- more than 5%, preferably more than 7% in weight by weight of the total composition, of ammonium halides in which the nitrogen atom is substituted by only one or at least one alkyl group having at least 16 carbon atoms.

[0015] According to a preferred embodiment of this invention, the composition of ammonium halides comprises only C16-alkyl quaternary ammonium halide, more preferably ammonium chloride or bromide, in which the nitrogen atom of the ammonium group is substituted by only one or at least one alkyl group having 16 carbon atoms.

Preferably, the composition of ammonium halides comprises C16-alkylbenzyl ammonium halide, preferably C16-alkylbenzyl dimethyl ammonium halide, preferably BAK C16. According to an embodiment of the present invention, the composition of ammonium halides comprises a C16-alkyl quaternary ammonium halide, in which the nitrogen atom of the ammonium group is substituted by two or three lower alkyl groups, preferably by two or three methyl groups.

[0016] Thus, in a preferred embodiment of the present invention, said oil-in-water emulsion comprises 0.0005 to 0.1 % of C16-alkyl quaternary ammonium halide, in weight by weight of the emulsion.

[0017] According to a preferred embodiment, the emulsion of the invention comprises an oil phase, comprising mineral oil, castor oil, MCT, corn oil, olive oil, soybean oil or any suitable vegetable oil, preferably said oil phase comprises mineral oil, castor oil or MCT.

According to a preferred embodiment, the emulsion of the invention further comprises tyloxapol.

According to a preferred embodiment, the emulsion of the invention further comprises a tonicity agent such as for example glycerol, mannitol, sorbitol, sodium chloride; others surfactants such as poloxamer; and optionally at least one buffering agent such as for example citrate, phosphate, tris, borate, acetate, carbonate, borate-polyol complexes, histidine, glu-

conate and lactate.

In one preferred embodiment of the present invention, said emulsion comprises mannitol as tonicity agent.

In another preferred embodiment of the present invention, said emulsion comprises glycerol as tonicity agent. Preferably, the emulsion includes 1 to 5 % of oil phase, preferably of MCT castor oil or mineral oil, in weight by weight of the emulsion.

Preferably, the mineral oil phase is a mixture of heavy and light mineral oil.

The main droplet core is composed by 50% light mineral oil and 50% heavy mineral oil.

Mineral oil is a mixture of refined liquid saturated aliphatic (C14-C18) and cyclic hydrocarbons obtained from petroleum.

Light mineral oil is less viscous and has a lower specific gravity than heavy mineral oil. Heavy and light mineral oil are well known excipients, used in a variety of pharmaceutical formulations including oral, and topical (up to 95%) preparations. In ophthalmic ointments, mineral oil can be found as an excipient at concentrations of up to 60%

A combination of light and heavy mineral oil in ophthalmology has been recognized by the US authorities as bearing emollient properties particularly adapted to dry eye treatment (21 CFR 349).

[0018] Preferably, the emulsion includes 0.1 to 1 % of surfactants, such as tyloxapol and optionally poloxamer 188 and/or polysorbate 80 and/or tocopherol polyethylene glycol succinate and/or sorbitan monolaurate, in weight by weight of the emulsion.

[0019] Preferably, the emulsion comprises 0.3% of tyloxapol and optionally 0.1% of poloxamer 188, in weight by weight of the emulsion.

Preferably, the emulsion comprises 0.1% to 5% of tonicity agent(s), more preferably 0.5% to 4% and even more preferably 0.9% to 3.3%, in weight by weight of the emulsion.

In one embodiment, the emulsion comprises 0.1% to 5% of mannitol, more preferably 0.5% to 4% and even more preferably 0.9% to 3.3%, in weight by weight of the emulsion.

In another embodiment, the emulsion comprises 0.1 % to 2.5% of glycerol, more preferably 0.19% to 1.6%, in weight by weight of the emulsion.

[0020] In one preferred embodiment of the present invention, the emulsion comprises light and heavy mineral oil, tyloxapol, poloxamer 188, mannitol and cetalkonium chloride. Preferably, said emulsion comprises 0.5% of light mineral oil, 0.5% of heavy mineral oil, 0.3% of tyloxapol, 0.1% of poloxamer 188, 3.3% of mannitol, and 0.002% of cetalkonium chloride, in weight by weight of the emulsion.

[0021] In another preferred embodiment of the present invention, the emulsion comprises light and heavy mineral oil, tyloxapol, poloxamer 188, glycerol and cetalkonium chloride. Preferably, said emulsion comprises 0.5% of light mineral oil, 0.5% of heavy mineral oil, 0.3% of tyloxapol, 0.1% of poloxamer 188, 1.6% of glycerol, and 0.002% of cetalkonium chloride, in weight by weight of the emulsion.

[0022] According to a preferred embodiment of the invention, the emulsion is hypotonic with regards to the normal tears tonicity.

[0023] According to a preferred embodiment, the oil-in-water emulsion of the invention is less toxic than a solution comprising the same amount of the same C16-alkyl quaternary ammonium halide, in weight by weight of the solution. For example, the toxicity can be evaluated by a redness test or a Draize test as shown in the examples.

[0024] According to a preferred embodiment, the oil-in-water emulsion of the invention is less toxic than an emulsion comprising 0.01 to 0.1% of BAK in weight by weight of the emulsion, said BAK being a mixture of C12, C14 and C16-alkyl quaternary ammonium halide complying to US or European Pharmacopeia specifications.

[0025] According to a preferred embodiment, the oil-in-water emulsion of the invention does not induce redness in albino rabbit conjunctiva before administration of 9 drops of 50 μ l, said drops being administrated each 5 minutes, preferably before administration of 11 drops, and more preferably before administration of 13 drops.

[0026] According to a first embodiment, the emulsion does not contain any active principle. In this embodiment, the emulsion is particularly useful as artificial tears, or for the treatment of dry eye condition such as for example Dry Eye Syndrome or Chronic Dry Eye Disease (CDED), both clinically known as keratoconjunctivitis sicca.

According to a second embodiment, the composition of the invention contains an active principle.

In one embodiment, said active principle is selected from secretagogues such as pilocarpine or celameline, immunosuppressive agents such as natural or synthetic cyclosporins, tacrolimus or sirolimus, mucin secretagogues such as 15(S)-HETE, ecabet or diquafosol, androgen mimetics, flaxseed oil supplements, steroids, agonists of adenosine A3 receptor, squalene, vitamin A; said active principle being preferably cyclosporine.

In another embodiment, said active principle is chosen among the active substances capable of having additional or synergistic therapeutic effects for treating KCS. Preferably, said active principle can be selected in the group comprising astringents such as zinc sulfate, demulcents including cellulose derivatives, carboxymethylcellulose sodium, hydroxyethyl cellulose; hypromellose, methylcellulose, dextran 70, gelatin, polyethylene glycol 300, polyethylene glycol 400, polysorbate 80, propylene glycol, polyvinyl alcohol and povidone such as polyethylene glycol 6000, emollients such as lanolin preparations or oleaginous ingredients, vasoconstrictors such as naphazoline, ephedrine, tetrahydrozoline and phenylephrine salts.

[0027] In an embodiment of the invention, the oil-in-water emulsion is preserved.

In another embodiment of the invention, the oil-in-water emulsion is unpreserved; in an embodiment, the emulsion is packaged in unitary doses; in another embodiment, the emulsion is packaged in suitable multidose containers.

[0028] The present invention also relates to the oil-in-water emulsion as described here above, having a high viscosity, and being dispensed to the patient in the form of a gel suitable for ophthalmic use.

[0029] Another object of the present invention is also a medicament comprising the oil-in-water emulsion as described here above.

[0030] The present invention also relates to a kit for dry eye treatment comprising a first emulsion as described here above, which does not comprise an active principle, and a second emulsion which contains an active principle, preferably cyclosporine.

According to an embodiment said second emulsion comprising an active principle is an emulsion according to the present invention.

According to another embodiment, said second emulsion comprising an active principle is any emulsion suitable for ophthalmic use.

[0031] Another object of this invention is a pre-concentrate of the therapeutic oil-in-water emulsion of the invention and a process for manufacturing said pre-concentrate.

[0032] According to this invention, a pre-concentrate is defined as an emulsion having an amount of oil higher than the amount of oil of the therapeutic emulsion administered to a patient. In a first embodiment, the amount of oil in the pre-concentrate is of at least 4% v/v. In a second embodiment, the amount of oil in the pre-concentrate is of at least 8% v/v. In a third embodiment, the amount of oil in the pre-concentrate is of at least 10% v/v., preferably of at least 20 % v/v, more preferably of at least 30% v/v.

[0033] The pre-concentrate may be in a liquid form or in a gel form, or in any form suitable in view of its further dilution with water.

[0034] According to an embodiment, the pre-concentrate of ophthalmic oil-in-water emulsion according to the present invention may be sterilized, for example, by heat, such as by autoclaving, or by filtering or filtration, or by irradiation, or by gas sterilization. In another embodiment, the concentrate of the ophthalmic emulsion is prepared in an aseptic manner.

[0035] This invention also relates to a process for manufacturing a pre-concentrate of a therapeutic oil-in-water emulsion comprising the steps of emulsifying/mixing the oil phase with an aqueous phase and with surface-active component(s), wherein the optionally active principle is dissolved in the oil phase. The process for manufacturing said pre-concentrate comprises emulsifying an amount of oil with an aqueous phase and with suitable surfactants, in order to obtain an emulsion having an amount in oil higher than the amount in oil of the corresponding emulsion to be administered for therapeutic purposes.

[0036] Before beginning the manufacturing process, the therapeutic oil-in-water emulsion is designed, with a wished concentration of oil, the type of oil (suitable for ophthalmic use, such as for example mineral oil, castor oil, or MCT) the type of elements needed for emulsification such as surfactants for example, and optionally an active principle. The concentration of the concentrate is then decided, depending on the industrial volumes needed.

[0037] This invention also relates to a process for manufacturing a therapeutic oil-in-water emulsion comprising (1) manufacturing a pre-concentrate of an ophthalmic oil in water emulsion, said pre-concentrate having a content in oil of at least 4 % v/v, preferably of 10% v/v or more, more preferably of 20% v/v or more, even more preferably of 30% v/v or more by emulsifying/mixing an oil suitable for ophthalmic use selected in the group comprising mineral oil, castor oil and MCT, said oil phase containing optionally one or more active principle and a C16-alkyl quaternary ammonium halide and with surface-active component(s), with an aqueous phase and then(2) diluting one volume of the resulting pre-concentrate with 2 to 50 volumes of water.

[0038] According to an embodiment, the emulsification is such that the droplet size or the distribution of the droplet size in the pre-concentrate is about the same as the droplet size or the distribution of the droplet size of the therapeutic oil-in-water emulsion.

According to an embodiment, the diluting water may comprise additives selected from the group comprising tonicity agents, such as for example NaCl, glycerol or mannitol, viscosifying agents, buffering agents, preservatives, antioxidants or colorants.

[0039] According to an embodiment, the diluting water may also comprise a C16-alkyl quaternary ammonium halide.

[0040] Then, according to the invention, a pre-concentrate of this desired emulsion is produced by mixing the oil suitable for ophthalmic use, with an aqueous phase and with surface-active component(s); the average hydrophilic-lipophilic balance (HLB) of the surface-active component(s) may advantageously be about equal to the HLB or average HLB emulsion requirement of the oil or oils used in the present compositions.

[0041] An advantage of this invention is to produce large volumes of emulsions without having to scale-up the emulsifying process.

[0042] This invention relates to a process for manufacturing a therapeutic oil-in-water emulsion according to the invention, comprising manufacturing a concentrate according to the above-mentioned process and then diluting said concentrate, by mixing 1 volume of concentrate with 2 to 50 volumes of water, to obtain a final therapeutic emulsion

having an oil content of 5% v/v or less, preferably of 3% v/v or less, more preferably of 2% v/v or less, even more preferably of 1% v/v or less.

[0043] This invention also relates to a method for the treatment of ocular diseases or conditions consisting in the administration to a patient of an ophthalmic emulsion prepared from a pre-concentrate, according to the above described process.

[0044] The invention also relates to oil-in-water emulsions obtainable by the process of the invention, i.e. by manufacturing a concentrate including optionally an active principle, and then diluting said concentrate with 2 to 50 volumes of water, said water optionally comprising additives, such as for example tonicity agents, viscosifying agents, buffering agents, preservatives, antioxidants or colorants.

[0045] One advantage of the invention is that the oil-in-water emulsions obtained by dilution of the concentrates are formed with reduced energy input.

[0046] The following examples and figures illustrate the invention and should not be interpreted in any way as reducing the scope of this invention.

Figure 1 refers to the timing of appearance of redness in ocular surface.

Figure 2 refers to Draize Test evaluation.

Examples:

[0047] All concentrations in the emulsion formulae are expressed in weight/weight of the entire formulation, unless stated differently.

Example 1: Reduced toxicity of quaternary amines when incorporated into emulsions

Materials:

[0048]

1. Solution at 0.02% BAK (BAK Sol)

Excipients	Z01SOL472
BAK US	0.02
NaCl	0.612
Tris Buffer 5mM pH 7.1	0.069% Tris HCl 0.006% Tris Base
Water	Ad 100

2. Emulsion at 0.02%BAK (BAK Em)

Excipients	Z01EM471
Mineral oil heavy	0.500
Mineral oil light	0.500
Tyloxapol	0.300
BAK US	0.02
Tris Buffer 5mM pH 7.1	0.069% Tris HCl 0.006% Tris Base
Poloxamer 188	0.100
Glycerol	1.6
Water (up to 100)	Ad 100

EP 2 404 593 B1

3. Solution at 0.002% CKC (CKC Sol)

	Z01SOL473
CKC	0.002
NaCl	0.626
Tris Buffer 5mM pH 7.1	0.069% Tris HCl 0.006% Tris base
Water	Ad 100

4. Emulsion at 0.002%CKC (CKC Em)

Excipients	Z01EM264
Mineral oil heavy	0.500
Mineral oil light	0.500
Tyloxapol	0.300
CKC	0.002
Tris Buffer 5mM pH 7.1	0.069% Tris HCl 0.006% Tris Base
Poloxamer 188	0.100
Glycerol	1.6
Water (up to 100)	Ad 100

5. PBS

Methods:

[0049] Albino rabbits were administrated with 1 drop (50 μ l) each 5 minutes, for 15 times.

1/ Evaluation of toxicity by time of redness and DRAIZE test items analyzed at H4 and D1.

[0050] The time of the beginning of the redness in conjunctiva following the 15 times of instillations was evaluated (Figure 1). PBS did not induce any redness during all instillation period (data not shown). BAK Sol induced conjunctival redness very fast, about 10 to 15 minutes after the first instillation (after 2-3 drops). BAK Em, CKC Sol-instilled groups showed redness at about 25-35 minutes after the first instillation of eye drops (after 5-7 drops). CKC Em presented a visible redness at almost the end of the experimentation: 60 to 65 minutes after the first instillation (after 12-13 drops).

[0051] Draize Test clearly showed that at four hours (H4) after the last instillation the ocular irritation was the most important in BAK Sol-instilled group, which was higher than BAK Em and CKC Sol groups (with no difference between these two groups). BAK Sol, BAK Em, CKC Sol all showed higher ocular irritation than CKC Em, which presented no difference with PBS-instilled group (Figure 2).

[0052] One day after the administrations (D1), PBS, BAK Em, CKC Sol and CKC Em all returned to normal aspect without difference among them. But BAK Sol still induced more important ocular irritation than all other groups ($P < 0.0001$).

Example 2: Stability of the emulsions of the invention

1. Emulsions composition

[0053] Some emulsions are described below:

	Z01EM206	Z01EM209
MCT	2 %	2 %

(continued)

	Z01EM206	Z01EM209
Tyloxapol	0.3 %	0.3 %
BAK C16 (CKC)	0.02 %	0.025mM
Poloxamer 188	0.1 %	0.1 %
glycerol	2.25 %	2.25 %
water	qsp100	qsp100

Composition	Z01EM419	Z01EM264	Z01EM387	Z01EM418	Z01EM418
Light mineral oil	0.5%	0.5%	0.5%	0.5%	0.5%
Heavy mineral oil	0.5%	0.5%	0.5%	0.5%	0.5%
Tyloxapol	0.3%	0.3%	0.3%	0.3%	0.3%
Poloxamer 188	0.1%	0.1% %	-	-	-
PG	-	-	0.19%	-	-
PEG 300	-	-	0.19%	-	-
PEG 400	-	-	0.19%	-	-
Mannitol	3.3%	-	2%	2.5%	2.9%
Glycerol	-	1.6%	0.19%	0.19%	-
Cetalkonium chloride (CKC)	0.002%	0.002%	0.002%	0.002%	0.002%
Tromethamine	0.006%	0.006%	-	-	-
Tris HCl	0.071%	0.071%	-	-	-
Water	Up to 100%	Up to 100%	Up to 100%	Up to 100%	Up to 100%

Composition	Z01EM393-4	Z01EM395-6
MCT	1-2%	1-2%
Tyloxapol	0.3%	0.3%
PG	0.4%	-
Mannitol	-	0.9%
Glycerol	1%	1%
Cetalkonium chloride (CKC)	0.01%	0.01%
Water	Up to 100%	Up to 100%

2. Emulsions preparation

[0054] The oily and the water phases of the emulsion, which might contain or not an active principle, may be separately heated to an appropriate temperature. This temperature may be the same in both cases. Surfactants might be dissolved in the oil, water phase or in both. A first coarse emulsion is generated by magnetic stirring, and the droplet size is reduced by high shear mixing, high pressure homogenization, or both.

The oil-in-water emulsions of the present invention can be sterilized after preparation using heat, for example, autoclave steam sterilization.

3. Impact of chain length on emulsions characteristics

a) Emulsion droplet size

[0055] The mean diameter of the oil droplets is determined by dynamic light scattering using a High Performance Particle Sizer type HPPS 5001 (Malvern Instruments, Worcestershire, UK). Measurements are performed at 25°C following dilution of the emulsion in double distilled water.

Table 1: Emulsions droplet size values (nm)

[0056] Emulsions of Table 1 and Table 2 contain 2% MCT, 0.3% Tyloxapol and 0.1% Poloxamer 188 and 2.25% glycerol and compositions of BAK; Concentrations of BAK range from 0.001 to 0.1% in weight to the weight of the emulsion.

	0.001%	0.0025%	0.005%	0.01%	0.02%	0.04%	0.1%
BAK C12	-	-	198	263	230	225	180
BAK C14	-	204	190	190	155	238	185
BAK C16	220	210	148	180	155	188	183

b) Emulsion zeta potential

[0057] Zeta potential can be measured by a zetameter such as Zetasizer 2000, Malvern Instruments Ltd, UK. The zeta potential of the emulsion droplet surface is determined by electrophoretic mobility. Measurements are performed at 25°C following dilution at 1:250 of the emulsion in double distilled water. The electrophoretic mobility is converted into zeta potential values through the Smoluchowsky equation. The following table and graph show the evolution of the zeta potential (indicative of the surface charge) at increasing concentrations of QA. It can be observed that for more lipophilic (longer) chain lengths, positive charges are attained more rapidly and at lower concentrations, suggesting a preferential partition within the oil droplet surface.

Table 2: Emulsions zeta potential values (mV)

	0.001%	0.0025%	0.005%	0.01%	0.02%	0.04%	0.1%
BAK C12	-	-	-6.9	+4.2	+7.9	+16.8	+23.8
BAK C14	-	+11.4	+19.6	+22.9	+28.4	+39.3	+44.5
BAK C16	+16.2	+24.4	+31.4	+36.7	+44.1	+47.2	+48.9

c) Emulsion stability over time

[0058] The stability of the emulsions can be evaluated by the evolution of their aspect, with a visual score with a visual score going from 13 - best aspect to 1- total phase separation.

[0059] It can be observed from the following table that, at equimolar concentration, longer (more lipophilic) chain length QA results in more stable emulsion.

Type and conc. of QA	After preparation (TO)	Following 3 months at 40°C
0.25 mM BAK C12	12	2
0.25 mM BAK C14	13	7
0.25 mM BAK C16	13	9
0.5 mM BAK C12	10	2
0.5 mM BAK C14	13	7
0.5 mM BAK C16	11	9

Claims

1. An oil-in-water emulsion comprising:

- 1% to 5% w/w of an oil phase selected from the group consisting of medium chain triglycerides and mineral oils; and
- 0.0005% to 0.1% w/w of cetalkonium chloride, in weight to the weight of the emulsion.

2. The oil-in-water emulsion according to claim 1, further comprising poloxamer, preferably in an amount of 0.1% w/w, in weight to the weight of the emulsion.

3. The oil-in-water emulsion according to claim 1 or claim 2, further comprising tyloxapol, preferably in an amount of 0.3% w/w, in weight to the weight of the emulsion.

4. The oil-in-water emulsion according to claim 1 to 3, wherein the oil phase is medium chain triglycerides, preferably in an amount of 2% w/w, in weight to the weight of the emulsion.

5. The oil-in-water emulsion according to claim 1 to 4, wherein the amount of mineral oil is 1% w/w.

6. The oil-in-water emulsion according to claim 1 to 5, wherein the oil phase is mineral oils, said mineral oils being a mixture of heavy mineral oil and light mineral oil wherein the amount of light mineral oil is 0.5% w/w and the amount of heavy mineral oil is 0.5% w/w.

7. The oil-in-water emulsion according to claim 1 to 6, wherein the amount of cetalkonium chloride is 0.001% to 0.02% w/w, preferably 0.001%, 0.0025%, 0.005%, 0.01%, 0.02%, 0.04% or 0.1% w/w.

8. The oil-in-water emulsion according to claim 1 to 7, further comprising buffering agents and/or at least one tonicity agent, preferably mannitol or glycerol.

9. The oil-in-water emulsion according to claim 1 to 8, said emulsion having a positive zeta potential.

10. The oil-in-water emulsion according to claim 1 to 9, said emulsion having a droplet size of 100 to 500 nm.

11. The oil-in-water emulsion according to claim 1 to 10, said emulsion being hypotonic.

12. The oil-in-water emulsion according to claim 1 to 11, wherein said emulsion is in the form of a gel suitable for ophthalmic use.

13. The oil-in-water emulsion according to claim 1 to 12, wherein said emulsion further comprises an active principle, preferably cyclosporin.

14. An oil-in-water emulsion according to claim 1 to 13 comprising:

- 0.5% w/w of light mineral oil;
- 0.5% w/w of heavy mineral oil;
- 0.002% w/w of cetalkonium chloride;
- 0.3% w/w of tyloxapol; and
- 0.1% w/w of poloxamer 188.

15. An oil-in-water emulsion according to claim 1 to 13 comprising:

- 1% to 2% w/w of medium chain triglycerides;
- 0.001% to 0.02% w/w cetalkonium chloride;
- 0.3% w/w of tyloxapol; and
- 0.1% w/w of poloxamer 188.

Patentansprüche

1. Öl-in-Wasser-Emulsion, umfassend:

- 5 - 1 % bis 5 % w/w einer Ölphase, ausgewählt aus der Gruppe, bestehend aus mittelkettigen Triglyceriden und Mineralölen; und
 - 0,0005 % bis 0,1 % w/w Cetalkoniumchlorid in Gewicht auf das Gewicht der Emulsion.
- 10 **2. Öl-in-Wasser-Emulsion nach Anspruch 1, weiter umfassend Poloxamer, vorzugsweise in einer Menge von 0,1 % w/w, in Gewicht auf das Gewicht der Emulsion.**
- 3. Öl-in-Wasser-Emulsion nach Anspruch 1 oder Anspruch 2, weiter umfassend Tyloxapol, vorzugsweise in einer Menge von 0,3 % w/w, in Gewicht auf das Gewicht der Emulsion.**
- 15 **4. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 3, wobei die Ölphase mittelkettige Triglyceride ist, vorzugsweise in einer Menge von 2 % w/w, in Gewicht auf das Gewicht der Emulsion.**
- 5. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 4, wobei die Menge von Mineralöl 1 % w/w ist.**
- 20 **6. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 5, wobei die Ölphase Mineralöle ist, wobei die Mineralöle eine Mischung aus schwerem Mineralöl und leichtem Mineralöl sind, wobei die Menge von leichtem Mineralöl 0,5 % w/w und die Menge von schweren Mineralöl 0,5 % w/w ist.**
- 25 **7. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 6, wobei die Menge von Cetalkoniumchlorid 0,001 % bis 0,02 % w/w, vorzugsweise 0,001 %, 0,0025 %, 0,005 %, 0,01 %, 0,02 %, 0,04 % oder 0,1 % w/w ist.**
- 8. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 7, weiter umfassend Puffermittel und/oder mindestens ein Tonizitätsmittel, vorzugsweise Mannitol oder Glycerol.**
- 30 **9. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 8, wobei die Emulsion ein positives Zeta-Potential aufweist.**
- 10. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 9, wobei die Emulsion eine Tröpfchengröße von 100 bis 500 nm aufweist.**
- 35 **11. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 10, wobei die Emulsion hypotonsisch ist.**
- 12. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 11, wobei die Emulsion die Form eines Gels aufweist, das zur ophthalmischen Verwendung geeignet ist.**
- 40 **13. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 12, wobei die Emulsion weiter einen Wirkstoff umfasst, vorzugsweise Cyclosporin.**
- 14. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 13, umfassend**
- 45 - 0,5 % w/w leichtes Mineralöl;
 - 0,5 % w/w schweres Mineralöl;
 - 0,002 % w/w Cetalkoniumchlorid;
 - 0,3 % w/w Tyloxapol; und
 - 0,1 % w/w Poloxamer 188.
- 50 **15. Öl-in-Wasser-Emulsion nach Anspruch 1 bis 13, umfassend**
- 55 - 1 % bis 2 % w/w mittelkettige Triglyceride;
 - 0,001 % bis 0,02 % w/w Cetalkoniumchlorid;
 - 0,3 % w/w Tyloxapol; und
 - 0,1 % w/w Poloxamer 188.

Revendications

1. Une émulsion huile-dans-eau comprenant :

- 1 % à 5% en poids d'une phase huile choisie dans le groupe consistant en : triglycérides à chaîne moyenne et huiles minérales ; et
- 0.0005% à 0.1% en poids de chlorure de cétalkonium, en poids par rapport au poids de l'émulsion.

2. L'émulsion huile-dans-eau selon la revendication 1, comprenant en outre un poloxamère, de préférence dans une quantité de 0.1 % en poids par rapport au poids de l'émulsion.

3. L'émulsion huile-dans-eau selon la revendication 1 ou la revendication 2, comprenant en outre du tyloxapol, de préférence dans une quantité de 0.3% en poids par rapport au poids de l'émulsion.

4. L'émulsion huile-dans-eau selon la revendication 1 à 3, dans laquelle la phase huile est constituée de triglycérides à chaîne moyenne, de préférence dans une quantité de 2% en poids par rapport au poids de l'émulsion.

5. L'émulsion huile-dans-eau selon la revendication 1 à 4, dans laquelle la quantité d'huile minérale est de 1 % en poids.

6. L'émulsion huile-dans-eau selon la revendication 1 à 5, dans laquelle la phase huile est constituée d'huiles minérales, lesdites huiles minérales étant un mélange d'huile minérale lourde et d'huile minérale légère, dans laquelle la quantité d'huile minérale légère est de 0.5% en poids et la quantité d'huile minérale lourde est de 0.5% en poids.

7. L'émulsion huile-dans-eau selon la revendication 1 à 6, dans laquelle la quantité de chlorure de cétalkonium est de 0.001 % à 0.02% en poids, de préférence 0.001 %, 0.0025%, 0.005%, 0.01%, 0.02%, 0.04% ou 0.1% en poids.

8. L'émulsion huile-dans-eau selon la revendication 1 à 7, comprenant en outre des agents tampons et/ou au moins un agent de tonicité, de préférence mannitol ou glycérol.

9. L'émulsion huile-dans-eau selon la revendication 1 à 8, ladite émulsion ayant un potentiel zêta positif.

10. L'émulsion huile-dans-eau selon la revendication 1 à 9, ladite émulsion ayant une taille de gouttelette de 100 à 500 nm.

11. L'émulsion huile-dans-eau selon la revendication 1 à 10, ladite émulsion étant hypotonique.

12. L'émulsion huile-dans-eau selon la revendication 1 à 11, dans laquelle ladite émulsion est sous la forme d'un gel adapté pour un usage ophtalmique.


13. L'émulsion huile-dans-eau selon la revendication 1 à 12, dans laquelle ladite émulsion comprend en outre un principe actif, de préférence une cyclosporine.

14. L'émulsion huile-dans-eau selon la revendication 1 à 13 comprenant :

- 0.5% en poids d'huile minérale légère ;
- 0.5% en poids d'huile minérale lourde ;
- 0.002% en poids de chlorure de cétalkonium ;
- 0.3% en poids de tyloxapol ; et
- 0.1% en poids de poloxamère 188.


15. L'émulsion huile-dans-eau selon la revendication 1 à 13 comprenant :

- 1% to 2% en poids de triglycérides à chaîne moyenne ;
- 0.001 % à 0.02% w/w en poids de chlorure de cétalkonium ;
- 0.3% en poids de tyloxapol ; et
- 0.1% en poids de poloxamère 188.



* $P < 0.0001$ compared to BAK Sol # $P < 0.0001$ compared to CKC Em

FIG. 1



* $P < 0.0001$ compared to BAK Sol # $P < 0.0001$ compared to PBS and CKC Em

FIG. 2

KVATERNER AMMÓNIUM VEGYÜLETEKET TARTALMAZÓ KÉSZÍTMÉNY

Szabadalmi igénypontok

1. Olaj a vízben emulzió, amely a következőket tartalmazza:
 - 1 tömeg%-tól (w/w) 5 tömeg%-ig (w/w) egy olajfázist, amely a következőkből álló csoportból van választva: közepes lánc hosszúságú trigliceridek és ásványolajok; és
 - 0,0005 tömeg%-tól (w/w) 0,1 tömeg%-ig (w/w) cetalkónium-kloridot, az emulzióknak a tömegére vonatkoztatva.
2. Olaj a vízben emulzió az 1. igénypont szerint, amely továbbá tartalmaz poloxamert, előnyösen egy 0,1 tömeg% (w/w) mennyiségben, az emulzióknak a tömegére vonatkoztatva.
3. Olaj a vízben emulzió az 1. igénypont vagy 2. igénypont szerint, amely továbbá tartalmaz tiloxapolt, előnyösen egy 0,3 tömeg% (w/w) mennyiségben, az emulzióknak a tömegére vonatkoztatva.
4. Olaj a vízben emulzió az 1.-től 3.-ig igénypontok szerint, ahol az olajfázist közepes lánc hosszúságú trigliceridek képezik, előnyösen egy 2 tömeg% (w/w) mennyiségben, az emulzióknak a tömegére vonatkoztatva.
5. Olaj a vízben emulzió az 1.-től 4.-ig igénypontok szerint, ahol az ásványolaj mennyisége 1 tömeg% (w/w).
6. Olaj a vízben emulzió az 1.-től 5.-ig igénypontok szerint, ahol az olajfázist ásványolajok képezik, ahol a nevezett ásványolajokat egy keverék képezi, amely nehéz ásványolajból és könnyű ásványolajból áll, ahol a könnyű ásványolaj mennyisége 0,5 tömeg% (w/w) és a nehéz ásványolaj mennyisége 0,5 tömeg% (w/w).
7. Olaj a vízben emulzió az 1.-től 6.-ig igénypontok szerint, ahol a cetalkónium-klorid mennyisége 0,001 tömeg%-tól (w/w) 0,02 tömeg%-ig (w/w) terjed, előnyösen az 0,001 tömeg% (w/w), 0,0025 tömeg% (w/w), 0,005% tömeg% (w/w), 0,01 tömeg% (w/w), 0,02 tömeg% (w/w), 0,04 tömeg% (w/w) vagy 0,1 tömeg% (w/w).
8. Olaj a vízben emulzió az 1.-től 7.-ig igénypontok szerint, amely továbbá tartalmaz puffrelő szereket (angolul: „buffering agents”) és/vagy legalább egy tonicitási szert (angolul: „tonicity agent”), előnyösen mannitot vagy glicerint.
9. Olaj a vízben emulzió az 1.-től 8.-ig igénypontok szerint, ahol a nevezett emulzió egy pozitív zéta-potenciállal rendelkezik.
10. Olaj a vízben emulzió az 1.-től 9.-ig igénypontok szerint, ahol a nevezett emulzió egy 100 nm-től 500 nm-ig terjedő cseppmérettel rendelkezik.
11. Olaj a vízben emulzió az 1.-től 10.-ig igénypontok szerint, ahol a nevezett emulzió hipotóniás (angolul: „hypotonic”).
12. Olaj a vízben emulzió az 1.-től 11.-ig igénypontok szerint, ahol a nevezett emulzió egy gél formájában áll rendelkezésre, amely alkalmas szemészeti alkalmazásra.
13. Olaj a vízben emulzió az 1.-től 12.-ig igénypontok szerint, ahol a nevezett emulzió továbbá tartalmaz egy hatóanyagot, előnyösen ciklosporint.
14. Olaj a vízben emulzió az 1.-től 13.-ig igénypontok szerint, amely a következőket tartalmazza:

- 0,5 tömeg% (w/w) könnyű ásványolajat;
- 0,5% tömeg% (w/w) nehéz ásványolajat;
- 0,002 tömeg% (w/w) cetalkónium-kloridot;
- 0,3 tömeg% (w/w) tiloxapolt; és
- 0,1 tömeg% (w/w) poloxamer 188-t.

15. Olaj a vízben emulzió az 1.-től 13.-ig igénypontok szerint, amely a következőket tartalmazza:

- 1 tömeg%-tól (w/w) 2 tömeg%-ig (w/w) közepes lánc hosszúságú triglicerideket;
- 0,001 tömeg%-tól (w/w) 0,02 tömeg%-ig (w/w) cetalkónium-kloridot;
- 0,3 tömeg% (w/w) tiloxapolt; és
- 0,1 tömeg% (w/w) poloxamer 188-t.