

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
24 August 2006 (24.08.2006)

PCT

(10) International Publication Number
WO 2006/087223 A1(51) International Patent Classification:
C07C 209/36 (2006.01)(21) International Application Number:
PCT/EP2006/001508(22) International Filing Date:
20 February 2006 (20.02.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0302/05 21 February 2005 (21.02.2005) CH

(71) Applicant (for all designated States except US): SYNGENTA PARTICIPATIONS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WALTER, Harald** [DE/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **CORSI, Camilla** [IT/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **EHRENFREUND, Josef** [AT/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **LAMBERTH, Clemens** [DE/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **SCHNEIDER, Hermann** [DE/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH). **TOBLER, Hans** [CH/CH]; Syngenta Crop Protection AG, Schwarzwaldallee 215, CH-4058 Basel (CH).

(74) Common Representative: SYNGENTA PARTICIPATIONS AG; Intellectual Property, P.O. Box, CH-4002 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

[Continued on next page]


(54) Title: PROCESS FOR THE PRODUCTION OF 2- (2-AMINOPHENYL)-BICYCLOPROPANE DERIVATIVES




(I)



(II)



(III)



(IV)

(57) Abstract: The present invention relates to a process for the preparation of compounds of formula (I) wherein the substituents are as defined in claim 1, by a) reaction of a compound of formula (II) to form a compound of formula (III) b) reaction of that compound in the presence of a base to form a compound of formula (IV) c) conversion of that compound in the presence of a reducing agent into a compound of formula (I).

WO 2006/087223 A1



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


Process for the production of anilines

The present invention relates to a process for the preparation of 2-(2-aminophenyl)-bicyclopropanes, and to novel nitrobenzene intermediates for use in that process.

2-(2-Aminophenyl)-bicyclopropanes, such as, for example, unsubstituted 2-(2-aminophenyl)-bicyclopropane, are valuable intermediates for the preparation of ortho-bicyclopropyl-carboxanilide fungicides, such as are described, for example, in WO 03/074491.

In WO 03/074491, a process for the preparation of 2-(2-aminophenyl)-bicyclopropanes is described (see Scheme 1):

Scheme 1:



According to WO 03/074491, ketones of formula (A), wherein R³ may be, *inter alia*, unsubstituted or substituted cyclopropyl, are reacted, for example, first with bromine and methanol and then with triphenylphosphine. The compounds of formula (B) obtained are converted in a two-step reaction into compounds of formula (C) wherein Hal is bromine or iodine (first, reaction with sodium hydride, then reaction with 2-bromobenzaldehyde or

2-iodobenzaldehyde, respectively). Compounds of formula (C) can be converted into the corresponding 2-(2-halophenyl)-bicyclopropanes (D) by Kishner cyclisation, which proceeds by way of a Δ^2 -pyrazoline. For that purpose, compounds of formula (C) are reacted, with heating, with hydrazine, as a result of which the corresponding Δ^2 -pyrazolines are formed. Subsequently, potassium hydroxide is added for isomerisation, and renewed heating is carried out to remove N₂. 2-(2-Halophenyl)-bicyclopropanes (D) can be aminated in a two-step reaction to form the corresponding 2-(2-aminophenyl)-bicyclopropanes (F). For that purpose, first of all benzophenone imine, sodium tert-butanolate, tris(dibenzylideneacetone)-dipalladium (Pd₂dba₃) and racemic 2,2'-bis(diphenylphosphine)-1,1'-binaphthyl (BINAP) are added. The resulting imines (E) are reacted in the second reaction step, for example with hydroxylamine and sodium acetate, to form the corresponding 2-(2-aminophenyl)-bicyclopropanes (F).


Such a reaction procedure is not suitable, however, for the preparation of 2-(2-aminophenyl)-bicyclopropanes, especially for large-scale preparation processes, because of the costly palladium-containing catalysts and ligands, such as, for example, BINAP.

In WO 03/074491, two further routes for the preparation of 2-(2-aminophenyl)-bicyclopropanes are described. A first route is by way of nitration of bicyclopropyl-benzenes. It has been found, however, that the reaction is not workable in view of the fact that the cyclopropyl ring linked directly to the benzene ring has increased reactivity in bicyclopropyl-benzenes with respect to electrophiles. A second route is by way of application of the Simmons-Smith reaction (Zn/Cu, CH₂I₂ with ether as solvent) to 1-((E/Z)-2-cyclopropylvinyl)-2-nitrobenzenes. In that case, too, the reaction has been found to be unsuitable for the preparation of 2-(2-nitrophenyl)-bicyclopropanes, since the reactivity of the double bond is too low.

The aim of the present invention is therefore to provide a process for the preparation of 2-(2-aminophenyl)-bicyclopropanes that allows such compounds to be prepared in an economically advantageous manner in high yields and in good quality.


The present invention accordingly relates to a process for the preparation of compounds of formula I

- 3 -



wherein R_1 , R_2 and R_3 are each independently of the others hydrogen or methyl, which comprises

a) reaction of a compound of formula II




wherein R_1 , R_2 and R_3 are as defined for formula I, either

a1) with triphenylphosphine dibromide or triphenylphosphine dichloride or


a2) with RSO_2Cl , wherein R is C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, benzyl, phenyl, nitrophenyl, halophenyl or C_1 - C_6 alkylphenyl, in the presence of a base,

to form a compound of formula III



wherein X is bromine, chlorine or OSO_2R , wherein R is C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, benzyl, phenyl, nitrophenyl, halophenyl or C_1 - C_6 alkylphenyl, and R_1 , R_2 and R_3 are as defined for formula I; and

b) reaction of that compound in the presence of a base to form a compound of formula IV



wherein R_1 , R_2 and R_3 are as defined for formula 1; and


c) conversion of that compound in the presence of a reducing agent into a compound of formula I.

Ortho-bicyclopropylcarboxanilide fungicides are generally chiral molecules that occur in isomeric forms. Accordingly they exist as trans/cis isomers based on the substitution pattern of the cyclopropyl ring linked directly to the benzene ring. It is known that the fungicidal activity of compounds such as are described, for example, in WO 03/074491, can be influenced by the stereochemistry. It has been found in the case of the ortho-bicyclopropylcarboxanilide fungicides described therein that the trans isomers generally have higher fungicidal activity. The development of a process that enables the production of a marked excess of trans ortho-bicyclopropylcarboxanilide fungicides is therefore extremely desirable.


The reaction sequence described in WO 03/074491 (Scheme 1) yields a trans:cis ratio of the 2-(2-aminophenyl)-bicyclopropane isomers of about 2:1.

A further aim of the present invention is accordingly to provide a process for the preparation of 2-(2-aminophenyl)-bicyclopropanes having a significantly higher proportion of trans isomers.

The process according to the invention allows compounds of formula I




to be produced wherein R_1 , R_2 and R_3 are each independently of the others hydrogen or methyl and wherein the ratio of compounds of formula Ia (trans)



to compounds of formula Ib (cis)

- 5 -




is more than 2:1.

The alkyl groups in the definitions of the substituents may be straight-chain or branched and are, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, pentyl and hexyl and branched isomers thereof.

Halogen in the context of halophenyl is generally fluorine, chlorine, bromine or iodine.


Fluoroalkyl groups having a chain length of from 1 to 4 carbon atoms are, for example, fluoromethyl, difluoromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, 1-fluoroethyl, 2-fluoroethyl, 2-fluoroprop-2-yl, pentafluoroethyl, 2,2,3,3-tetrafluoroethyl, pentafluoroethyl or heptafluoro-n-propyl; fluoroalkyl groups are preferably trichloromethyl, fluoromethyl, dichlorofluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl or heptafluoro-n-propyl.

Compounds of formula I occur in various stereoisomeric forms, which are represented by formulae I_I, I_{II}, I_{III} and I_{IV}:




The process according to the invention includes the preparation of those stereoisomeric forms of formulae I_I , I_{II} , I_{III} and I_{IV} , wherein R_1 , R_2 and R_3 are as defined for formula I, and the preparation of mixtures of those stereoisomeric forms in any ratio.

In the context of the present invention, compounds of formula Ia (trans)




wherein R_1 , R_2 and R_3 are as defined for formula I, are understood to be compounds of formula I_I , wherein R_1 , R_2 and R_3 are as defined for formula I; compounds of formula I_{II} , wherein R_1 , R_2 and R_3 are as defined for formula I; or a mixture, in any ratio, of compounds of formula I_I , wherein R_1 , R_2 and R_3 are as defined for formula I, and compounds of formula I_{III} , wherein R_1 , R_2 and R_3 are as defined for formula I.

In the context of the present invention, compounds of formula Ia (trans)




wherein R_1 , R_2 and R_3 are as defined for formula I, are understood to be, preferably, a racemic mixture of compounds of formula I, wherein R_1 , R_2 and R_3 are as defined for formula I, and compounds of formula I_{III}, wherein R_1 , R_2 and R_3 are as defined for formula I.

In the context of the present invention, compounds of formula Ib (cis)







wherein R_1 , R_2 and R_3 are as defined for formula I, are understood to be compounds of formula I_{III}, wherein R_1 , R_2 and R_3 are as defined for formula I; compounds of formula I_{IV}, wherein R_1 , R_2 and R_3 are as defined for formula I; or a mixture, in any ratio, of compounds of formula I_{III}, wherein R_1 , R_2 and R_3 are as defined for formula I, and compounds of formula I_{IV}, wherein R_1 , R_2 and R_3 are as defined for formula I.

In the context of the present invention, compounds of formula Ib (cis)




wherein R_1 , R_2 and R_3 are as defined for formula I, are understood to be, preferably, a racemic mixture of compounds of formula I_{III}, wherein R_1 , R_2 and R_3 are as defined for formula I, and compounds of formula I_{IV}, wherein R_1 , R_2 and R_3 are as defined for formula I.

Compounds of formula IV occur in various stereoisomeric forms, which are represented by formulae IV_I, IV_{II}, IV_{III} and IV_{IV}:




The process according to the invention includes the preparation of those stereoisomeric forms of formulae IV_I, IV_{II}, IV_{III} and IV_{IV}, wherein R₁, R₂ and R₃ are as defined for formula I, and the preparation of mixtures of those stereoisomeric forms in any ratio.

In the context of the present invention, compounds of formula IVa (trans)




wherein R₁, R₂ and R₃ are as defined for formula I, are understood to be compounds of formula IV_I, wherein R₁, R₂ and R₃ are as defined for formula I; compounds of formula IV_{II}, wherein R₁, R₂ and R₃ are as defined for formula I; or a mixture, in any ratio, of compounds of formula IV_I, wherein R₁, R₂ and R₃ are as defined for formula I, and compounds of formula IV_{II}, wherein R₁, R₂ and R₃ are as defined for formula I.

In the context of the present invention, compounds of formula IVa (trans)




wherein R_1 , R_2 and R_3 are as defined for formula I, are understood to be, preferably, a racemic mixture of compounds of formula IV_i, wherein R_1 , R_2 and R_3 are as defined for formula I, and compounds of formula IV_{ii}, wherein R_1 , R_2 and R_3 are as defined for formula I.

In the context of the present invention, compounds of formula IVb (cis)



wherein R_1 , R_2 and R_3 are as defined for formula I, are understood to be compounds of formula IV_{iii}, wherein R_1 , R_2 and R_3 are as defined for formula I; compounds of formula IV_{iv}, wherein R_1 , R_2 and R_3 are as defined for formula I; or a mixture, in any ratio, of compounds of formula IV_{iii}, wherein R_1 , R_2 and R_3 are as defined for formula I, and compounds of formula IV_{iv}, wherein R_1 , R_2 and R_3 are as defined for formula I.

In the context of the present invention, compounds of formula IVb (cis)



wherein R_1 , R_2 and R_3 are as defined for formula I, are understood to be, preferably, a racemic mixture of compounds of formula IV_{iii}, wherein R_1 , R_2 and R_3 are as defined for formula I, and compounds of formula IV_{iv}, wherein R_1 , R_2 and R_3 are as defined for formula I.

In the context of the present invention, a "racemic mixture" of two enantiomers is understood to be a mixture of the two enantiomers in a ratio substantially equal to 1 :1.

The process according to the invention is suitable especially for the preparation of compounds of formula I wherein R_2 and R_3 are hydrogen.

The process according to the invention is suitable more especially for the preparation of compounds of formula I wherein R_1 , R_2 and R_3 are hydrogen.

The process according to the invention is suitable more especially for the preparation of compounds of formula I wherein R_1 is methyl and R_2 and R_3 are hydrogen.

Process Step a):

In an embodiment (a1) of the process according to the invention, in Process Step a), a compound of formula II is reacted with triphenylphosphine dibromide or triphenylphosphine dichloride.

In that embodiment, either triphenylphosphine dibromide or triphenylphosphine dichloride is added directly to the compounds of formula II, or triphenylphosphine dibromide or triphenylphosphine dichloride is generated *in situ* in the reaction mixture by the addition of bromine or chlorine in the presence of triphenylphosphane.

Suitable amounts of triphenylphosphine dibromide or triphenylphosphine dichloride for that reaction are, for example, from 1 to 3 equivalents, especially from 1 to 1.5 equivalents.

When triphenylphosphine dibromide or triphenylphosphine dichloride is generated *in situ*, an amount, for example, of from 1 to 3 equivalents, especially from 1 to 1.5 equivalents, of bromine or chlorine is suitable. Suitable amounts of triphenylphosphane for that variant of the reaction are, for example, from 1 to 3 equivalents, especially from 1 to 1.5 equivalents.

In that embodiment, the reaction can be carried out in the presence of an inert solvent. Suitable solvents are, for example, ethers, for example tetrahydrofuran or dioxane, or CH_3CN , and mixtures thereof; CH_3CN is preferred.

Temperatures are generally from -20°C to 80°C , with a range from -20°C to 25°C being preferred; special preference is given to carrying out the reaction at ambient temperature.

The reaction time for that reaction is generally from 1 to 48 hours, preferably from 1 to 18 hours.

In a further embodiment (a2) of the process according to the invention, in Process Step a), a compound of formula II is reacted in the presence of a base with RSO_2Cl , wherein R is $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ fluoroalkyl, benzyl, phenyl, nitrophenyl, halophenyl or $\text{C}_1\text{-C}_6$ alkylphenyl, especially $\text{C}_1\text{-C}_4$ alkyl, more especially methyl.

For that reaction, suitable amounts of RSO_2Cl , wherein R is $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ fluoroalkyl, benzyl, phenyl, nitrophenyl, halophenyl or $\text{C}_1\text{-C}_6$ alkylphenyl, are, for example, from 1 to 3 equivalents, especially from 1 to 1.2 equivalents.

Suitable bases are, for example, tertiary amines, such as trialkylamines, e.g. trimethylamine, triethylamine, diisopropylethylamine (Hünig's base), tri-n-butylamine, N,N-dimethylaniline or N-methylmorpholine, or inorganic bases, such as carbonates, e.g. K_2CO_3 or Na_2CO_3 , or hydroxides, e.g. NaOH or KOH, with preference being given to trialkylamines and special preference being given to triethylamine.

Suitable amounts of base for that reaction are, for example, from 1 to 3 equivalents, especially from 1 to 1.3 equivalents.


The reaction is preferably carried out in the presence of an inert solvent. Suitable solvents are, for example, dichloromethane, pyridine or ethers, for example tetrahydrofuran, and mixtures thereof, with preference being given to dichloromethane or pyridine, and special preference being given to dichloromethane.

Temperatures are generally from -20°C to 80°C , with a range from -20°C to 25°C being preferred; special preference is given to carrying out the reaction at ambient temperature.

The reaction time for that reaction is generally from 1 to 48 hours, preferably from 1 to 18 hours.

The starting compounds of formula II, wherein R_1 , R_2 and R_3 are as defined for formula I, can be prepared, for example, in accordance with the following reaction sequence (see Scheme 2):

Scheme 2:



Compounds of formula V, wherein X_1 is chlorine or bromine, are reacted with compounds of formula VI, wherein R_1 , R_2 and R_3 are as defined for formula I and R_a is C_1 - C_6 alkyl, in a two-step reaction sequence to form compounds of formula VII, wherein R_1 , R_2 and R_3 are as defined for formula I. In the first reaction step, compounds of formula V are reacted with compounds of formula VI under basic conditions, for example obtained by addition of NaH, NaOH or K_2CO_3 . After isolation of the crude product, heating in dimethyl sulfoxide (DMSO) in the presence of LiCl is carried out in the second reaction step. The resulting compounds of formula VII can be reacted to form compounds of formula II by addition of sodium borohydride in a protic solvent R_bOH , wherein R_b is C_1 - C_6 alkyl, such as, for example, isopropanol.

Compounds of formula V, wherein X_1 is chlorine or bromine, are known and are obtainable commercially.

Some of the compounds of formula VI, wherein R_1 , R_2 and R_3 are as defined for formula I and R_a is C_1 - C_6 alkyl, are known and are obtainable commercially. The remaining compounds of formula VI, wherein R_1 , R_2 and R_3 are as defined for formula I and R_a is C_1 - C_6 alkyl, can be prepared in an analogous manner to preparation processes such as are described, for

example, in Journal of Organic Chemistry 68(1), 27-34 (2003) and in Organic Preparations and Procedures International 10(5), 221-224 (1978).

Process Step b):

Suitable bases for Process Step b) are, for example, nitrogen-containing organic bases, such as, for example, tertiary amines, such as trialkylamines, e.g. trimethylamine, triethylamine, diisopropylethylamine (Hünig's Base), or tri-n-butylamine, N,N-dimethylaniline or N-methylmorpholine, piperidine, pyrrolidine, alkali metal or alkaline earth metal alcoholates, such as, for example, lithium, sodium or potassium alcoholates, especially methanolates, ethanolates or butanolates, or inorganic bases, such as hydroxides, e.g. NaOH or KOH, or hydrides, such as, for example, NaH.

Bases to which preference is given are hydroxides, especially KOH, hydrides, especially NaH, or alkali metal alcoholates, especially potassium tert-butanolate.

Suitable amounts of base for that reaction are, for example, from 1 to 3 equivalents, especially from 1.1 to 1.8 equivalents.

The reaction is preferably carried out in the presence of an inert solvent. Suitable solvents are, for example, alcohols, such as methanol, ethanol, propanol or isopropanol, or aprotic solvents, such as tetrahydrofuran, dimethylformamide, dimethylacetamide, N-methylpyrrolidone or dimethyl sulfoxide, and also mixtures thereof; dimethyl sulfoxide or dimethylformamide is especially preferred.

Temperatures are generally from 0°C to 80°C, with a range from 0°C to 25°C being preferred; special preference is given to carrying out the reaction at ambient temperature.

The reaction time for that reaction is generally from 1 to 48 hours, preferably from 1 to 18 hours.

Process Step c):

A suitable reducing agent for Process Step c) is, for example, hydrogen in the presence of a metal catalyst.

Suitable amounts of reducing agent for that reaction are, for example, from 1 to 5 equivalents, especially from 1 to 1.3 equivalents.

Suitable metal catalysts are, for example, platinum catalysts, such as, for example, platinum-carbon catalysts; palladium catalysts or rhodium catalysts, with special preference being given to platinum catalysts.

Suitable amounts of metal catalyst for that reaction are, for example, from 0.001 to 0.5 equivalent, especially from 0.01 to 0.1 equivalent.

The reaction is preferably carried out in the presence of an inert solvent. Suitable solvents are, for example, alcohols, such as methanol, ethanol, propanol or isopropanol, or aprotic solvents, such as tetrahydrofuran, tert-butyl methyl ether, dioxane or toluene, and mixtures thereof. Special preference is given to ethanol or methanol.

Temperatures are generally from 0°C to 80°C, with a range from 0°C to 25°C being preferred; special preference is given to carrying out the reaction at ambient temperature.

The reaction time for that reaction is generally from 1 to 48 hours, preferably from 1 to 6 hours.

By selecting suitable reaction conditions, the compound of formula III obtained in Reaction Step a) can be reacted to form a compound of formula IV directly, without isolation of intermediates. That reaction procedure is a particular advantage of the process according to the invention.

The process according to the invention is suitable for the preparation of compounds of formula I, wherein R_1 , R_2 and R_3 are each independently of the others hydrogen or methyl, very especially by

a) reaction of a compound of formula II, wherein R_1 , R_2 and R_3 are each independently of the others hydrogen or methyl, with RSO_2Cl , wherein R is C_1 - C_4 alkyl, in the presence of triethylamine in a temperature range of from -20°C to 25°C, using dichloromethane as solvent, to form a compound of formula III, wherein X is OSO_2 - C_1 - C_4 alkyl and R_1 , R_2 and R_3 are as defined for formula I; and

- b) reaction of that compound in the presence of a base selected from KOH, NaH and potassium tert-butanolate, in a temperature range of from -20°C to 25°C, using a solvent selected from dimethyl sulfoxide and dimethylformamide, to form a compound of formula IV, wherein R₁, R₂ and R₃ are as defined for formula I; and
- c) conversion of that compound into a compound of formula I in the presence of hydrogen and a platinum catalyst, in a temperature range from 0°C to 25°C, using ethanol as solvent.

For that preferred embodiment there are especially suitable compounds of formula I wherein R₂ and R₃ are hydrogen.

For that preferred embodiment there are very especially suitable compounds of formula I wherein R₁, R₂ and R₃ are hydrogen.

The present invention is explained in greater detail by way of the following Examples:

Example P1: Preparation of 2-(2-nitrophenyl)-bicyclopropane:

A mixture of 0.5 g of 1-cyclopropyl-3-(2-nitrophenyl)-propan-1-ol (2.26 mmol), 0.26 g of triethylamine (2.6 mmol) and 12 ml of dichloromethane is cooled to a temperature of 5°C and 0.28 g of methanesulfonic acid chloride, dissolved in 3 ml of dichloromethane, is added dropwise. The resulting mixture is stirred for 16 hours at ambient temperature. The organic phase is washed with ice-water and dried over sodium sulfate and concentrated by evaporation. 1-Cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate is obtained in the form of a crude product, which is used directly in the cyclisation.

The 1-cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate is dissolved in 15 ml of dimethyl sulfoxide and 0.17 g of potassium hydroxide (2.48 mmol) is added, and stirring is carried out for 5 hours at ambient temperature. The reaction mixture is added to ice-water. Extraction is carried out with ethyl acetate and the organic phase is dried over sodium sulfate and concentrated by evaporation. Chromatography on silica gel is carried out in order to remove by-products (eluant: ethyl acetate / hexane 1:15). After removal of the eluant, 0.28 g of 2-(2-nitrophenyl)-bicyclopropane (61 % of theory) is obtained in the form of a brownish liquid (trans:cis ratio: 4.5:1). ¹H-NMR of trans-2-(2-nitrophenyl)-bicyclopropane (CDCl₃ - ppm): 0.17/m/1H, 0.19/m/1H, 0.42/m/1H, 0.48/m/1H, 0.83/m/1H, 0.84/m/1H, 0.99/m/1H, 1.13/m/1H, 2.17/m/1H, 7.10/dd/1H, 7.25/m/1H, 7.45/m/1H, 7.78/dd/1H; ¹H-NMR of cis-2-(2-nitrophenyl)-bicyclopropane (CDCl₃ - ppm): -0.09/m/1H, 0.02/m/1H, 0.06/m/1H, 0.27/m/1H, 0.71/m/1H,

0.85/m/1H, 0.98/m/1H, 1.10/m/1H, 2.53/m/1H, 7.35/m/1H, 7.45/m/1H, 7.52/m/1H, 7.92/dd/1H.

Example P2: Preparation of 2-(2-nitrophenyl)-bicyclopropane:

A mixture of 0.5 g of 1-cyclopropyl-3-(2-nitrophenyl)-propan-1-ol (2.26 mmol), 0.26 g of triethylamine (2.6 mmol) and 12 ml of dichloromethane is cooled to a temperature of 5°C and 0.28 g of methanesulfonic acid chloride, dissolved in 3 ml of dichloromethane, is added dropwise. The resulting mixture is stirred for 16 hours at ambient temperature. The organic phase is washed with ice-water and dried over sodium sulfate and concentrated by evaporation. 1-Cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate is obtained in the form of a crude product, which is used directly in the cyclisation.

The 1-cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate is dissolved in 15 ml of dimethylformamide and 0.21 g of potassium hydroxide (3.2 mmol) is added, and stirring is carried out for 6 hours at ambient temperature. The reaction mixture is added to ice-water. Extraction is carried out with ethyl acetate, and the organic phase is dried over sodium sulfate and concentrated by evaporation. Chromatography on silica gel is carried out in order to remove by-products (eluant: ethyl acetate / hexane 1:15). After removal of the eluant, 0.28 g of 2-(2-nitrophenyl)-bicyclopropane (61 % of theory) is obtained in the form of a brownish liquid (trans:cis ratio: 4.4:1).

Example P3: Preparation of 2-(2-nitrophenyl)-bicyclopropane:

A mixture of 2.21 g of 1-cyclopropyl-3-(2-nitrophenyl)-propan-1-ol (10 mmol), 1.21 g of triethylamine (12 mmol) and 20 ml of dichloromethane is cooled to a temperature of 5°C and 1.26 g of methanesulfonic acid chloride (11 mmol), dissolved in 5 ml of dichloromethane, are added dropwise. The resulting mixture is stirred for 16 hours at ambient temperature. The organic phase is washed with ice-water and dried over sodium sulfate and concentrated by evaporation. 1-Cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate is obtained in the form of a crude product, which is used directly in the cyclisation.

0.48 g of sodium hydride (12 mmol) is introduced into 10 ml of dimethyl sulfoxide and a solution consisting of the 1-cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate and 15 ml of DMSO is added. Stirring is then carried out for 5 hours at ambient temperature. The reaction mixture is added to ice-water. Extraction is carried out with ethyl acetate, and the organic phase is dried over sodium sulfate and concentrated by evaporation.

Chromatography on silica gel is carried out in order to remove by-products (eluant: ethyl

acetate / hexane 1:15). After removal of the eluant, 0.28 g of 2-(2-nitrophenyl)-bicyclopropane (64 % of theory) is obtained in the form of a brownish liquid (trans:cis ratio: 4.1:1).

Example P4: Preparation of 2-(2-nitrophenyl)-bicyclopropane:

A mixture of 0.5 g of 1-cyclopropyl-3-(2-nitrophenyl)-propan-1-ol (2.26 mmol), 0.26 g of triethylamine (2.6 mmol) and 12 ml of dichloromethane is cooled to a temperature of 5°C and 0.28 g of methanesulfonic acid chloride, dissolved in 3 ml of dichloromethane, is added dropwise. The resulting mixture is stirred for 16 hours at ambient temperature. The organic phase is washed with ice-water and dried over sodium sulfate and concentrated by evaporation. 1-Cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate is obtained in the form of a crude product, which is used directly in the cyclisation.


The 1-cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate is dissolved in 15 ml of dimethyl sulfoxide and 0.28 g of potassium tert-butanolate (2.48 mmol) is added, and stirring is carried out for 3 hours at ambient temperature. The reaction mixture is added to ice-water.

Extraction is carried out with ethyl acetate, and the organic phase is dried over sodium sulfate and concentrated by evaporation. Chromatography on silica gel is carried out in order to remove by-products (eluant: ethyl acetate / hexane 1:15). After removal of the eluant, 0.3 g of 2-(2-nitrophenyl)-bicyclopropane (65 % of theory) is obtained in the form of a brownish liquid (trans:cis ratio: 4.7:1).

Example P5: Preparation of 2-(2-aminophenyl)-bicyclopropane:

In a hydrogenation reactor, 1 g of 2-(2-nitrophenyl)-bicyclopropane (4.9 mmol, trans:cis ratio: 4.1:1), dissolved in 20 ml of ethanol, is hydrogenated at ambient temperature using 0.1 g of 5 % platinum-carbon catalyst. After 2.5 hours and after 101 % of the amount of hydrogen theoretically required for the reduction has been taken up, the reaction is stopped. Following filtration of the reaction mixture, the solvent is removed by concentration by evaporation. 0.87 g of 2-(2-aminophenyl)-bicyclopropane (100 % of theory) is obtained in the form of a brownish liquid (trans:cis ratio: 4.4:1).

The following compounds of formula I can be prepared according to the above Examples:

Table 1: Compounds of formula I

Comp. No.	R ₁	R ₂	R ₃
A1	H	H	H
A2	CH ₃	H	H
A3	H	CH ₃	H
A4	H	H	CH ₃
A5	CH ₃	CH ₃	H
A6	CH ₃	H	CH ₃
A7	H	CH ₃	CH ₃
A8	CH ₃	CH ₃	CH ₃

The starting materials for the process of the present invention are distinguished by ease of availability and good handling properties and are moreover reasonably priced.


A further advantage of the process is that the ratio of trans isomers of formula Ia to cis isomers of formula Ib is significantly higher than described in the prior art; generally, trans:cis ratios of the prepared 2-(2-aminophenyl)-bicyclopropanes of more than 3:1 are achieved.

In accordance with the present process, compounds of formula I can be prepared in simple manner wherein the ratio of compounds of formula Ia (trans) to compounds of formula Ib (cis) is from 3:1 to 5:1.

In the process according to the invention, the trans/cis proportion of the end products of the process, the 2-(2-aminophenyl)-bicyclopropanes of formula I, is determined substantially by the trans/cis proportion of the 2-(2-nitrophenyl)-bicyclopropanes of formula IV formed when Process Step (b) is carried out. An increased proportion of trans remains substantially unchanged after Process Step (c), the reduction of the 2-(2-nitrophenyl)-bicyclopropanes to form the end products of the process, has been carried out.

The process according to the invention allows the trans proportion of compounds of formula I to be substantially increased by a further reaction step which is simple to execute.

In that especially preferred embodiment (bb) of the process according to the invention, the compounds of formula IV obtained according to Process Step (b)



wherein R_1 , R_2 and R_3 are as defined for formula 1,

bb) are isomerised in the presence of a base to form compounds of formula IV wherein the ratio of compounds of formula IVa (trans) to compounds of formula IVb (cis) is more than 6:1. Those compounds are then used in Process Step c).

The increased proportion of trans remains substantially unchanged after Process Step (c) has been carried out. That especially preferred process variant therefore yields compounds of formula I wherein the ratio of compounds of formula Ia (trans) to compounds of formula Ib (cis) is more than 6:1.

Process Step bb):

A suitable base for Process Step bb) is, for example, KOH or an alkali metal or alkaline earth metal alcoholate, such as, for example, a lithium, sodium or potassium alcoholate, especially a methanolate, ethanolate or butanolate. Special preference is given to KOH or potassium tert-butanolate, and very special preference is given to potassium tert-butanolate.

Suitable amounts of base for that reaction are, for example, from 0.3 to 3 equivalents, especially from 0.5 to 1.2 equivalents.

The reaction is preferably carried out in the presence of an inert solvent. Suitable solvents are, for example, aprotic solvents, such as tetrahydrofuran, dimethyl sulfoxide, dimethylacetamide; dimethoxyethane; dioxane or dimethylformamide, and also mixtures thereof; tetrahydrofuran is especially preferred.

In an embodiment to which very special preference is given, that reaction is carried out using potassium tert-butanolate as base and using tetrahydrofuran as solvent.

In another embodiment to which very special preference is given, that reaction is carried out using KOH as base and using dimethyl sulfoxide as solvent.

Temperatures are generally from -20°C to 80°C, with a range from -20°C to 25°C being preferred; special preference is given to carrying out the reaction at ambient temperature.

The reaction time for that reaction is generally from 0.5 to 12 hours, preferably from 1 to 3 hours.

Special preference is given to carrying out that reaction under a nitrogen atmosphere.

By selecting suitable reaction conditions, the compound of formula IV obtained in Reaction Step b) can be isomerised directly, without isolation of intermediates, to form a compound of formula IV wherein the ratio of compounds of formula IVa (trans) to compounds of formula IVb (cis) is more than 6:1 ("one-pot" method). That reaction procedure is a particular advantage of the especially preferred embodiment (bb) of the process according to the invention.

When that especially preferred embodiment (bb) of the process according to the invention is carried out as a "one-pot" method, the solvent used is more especially dimethyl sulfoxide or dimethylformamide.

The above-described especially preferred embodiment (bb) of the process according to the invention is explained in greater detail by way of the following Examples:

Example P6: Isomerisation of 2-(2-nitrophenyl)-bicyclopropane:

Under a nitrogen atmosphere, 0.5 g of potassium tert-butanolate (4.4 mmol) is added to a solution of 3 g of 2-(2-nitrophenyl)-bicyclopropane (14.7 mmol, trans:cis ratio: 3.7:1) in 100 ml of tetrahydrofuran. The resulting mixture is stirred for 1.5 hours at ambient temperature. Water is added and the reaction mixture is extracted with ethyl acetate. The organic phase is washed with a saturated sodium chloride solution and dried over sodium

sulfate and concentrated by evaporation. Chromatography on silica gel is carried out in order to remove by-products (eluant: ethyl acetate / hexane 1:10). After removal of the eluant, 2.35 g of 2-(2-nitrophenyl)-bicyclopropane (91 % of theory) are obtained in the form of a yellowish liquid (trans:cis ratio: 6.4:1).

Example P7: Isomerisation of 2-(2-nitrophenyl)-bicyclopropane:

Under a nitrogen atmosphere, 1.82 g of potassium tert-butanolate (16.1 mmol) are added to a solution of 3 g of 2-(2-nitrophenyl)-bicyclopropane (14.7 mmol, trans:cis ratio: 3.7:1) in 100 ml of tetrahydrofuran. The resulting mixture is stirred for 0.5 hours at -20°C. Water is added and the reaction mixture is extracted with ethyl acetate. The organic phase is washed with a saturated sodium chloride solution and dried over sodium sulfate and concentrated by evaporation. Chromatography on silica gel is carried out in order to remove by-products (eluant: ethyl acetate / hexane 1:10). After removal of the eluant, 2.19 g of 2-(2-nitrophenyl)-bicyclopropane (73 % of theory) are obtained in the form of a yellowish liquid (trans:cis ratio: 14.3:1).

Example P8: Preparation and isomerisation of 2-(2-nitrophenyl)-bicyclopropane ("one-pot" method):


1.18 g of 1-cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate (3.94 mmol) is dissolved in 40 ml of dimethylformamide, 0.91 g of potassium tert-butanolate (97 %, 7.88 mmol) are added and stirring is carried out for 1 hour at ambient temperature under a nitrogen atmosphere. Water is added to the reaction mixture. Extraction is carried out with ethyl acetate, and the organic phase is dried over sodium sulfate and concentrated by evaporation. Chromatography on silica gel is carried out in order to remove by-products (eluant: ethyl acetate / hexane 1:2). After removal of the eluant, 0.68 g of 2-(2-nitrophenyl)-bicyclopropane (85 % of theory) is obtained in the form of an orange-coloured liquid (trans:cis ratio: 6.5:1).

Example P9: Preparation and isomerisation of 2-(2-nitrophenyl)-bicyclopropane ("one-pot" method):

1.06 g of 1-cyclopropyl-3-(2-nitrophenyl)-propyl methanesulfonate (3.54 mmol) is dissolved in 20 ml of dimethyl sulfoxide, 1.05 g of potassium tert-butanolate (97 %, 10.62 mmol) are added, and stirring is carried out for 1.5 hours at ambient temperature under a nitrogen atmosphere. Water is added to the reaction mixture. Extraction is carried out with ethyl

acetate, and the organic phase is dried over sodium sulfate and concentrated by evaporation. Chromatography on silica gel is carried out in order to remove by-products (eluant: ethyl acetate / hexane 1:10). After removal of the eluant, 0.68 g of 2-(2-nitrophenyl)-bicyclopropane (67 % of theory) are obtained in the form of an orange-coloured liquid (trans:cis ratio: 7.7:1).

The compounds of formula III




wherein X is bromine, chlorine or OSO_2R , wherein R is $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ fluoroalkyl, benzyl, phenyl, nitrophenyl, halophenyl or $\text{C}_1\text{-C}_6$ alkylphenyl and R_1 , R_2 and R_3 are as defined for formula I, are valuable intermediates in the preparation of compounds of formula I and were developed specifically for the present process according to the invention. The present invention accordingly relates also to those compounds.

Especially valuable for the preparation of compounds of formula I are those compounds of formula III wherein X is OSO_2CH_3 .

An intermediate especially suitable for the preparation of compounds of formula I is the compound of formula III wherein X is OSO_2CH_3 and R_1 , R_2 and R_3 are hydrogen.


Preferred compounds of formula III are listed in the following Table. In the following Table, "Ph" denotes phenyl.

Table 2: Compounds of formula III




Comp. No.	R ₁	R ₂	R ₃	X
Z1.01	H	H	H	OSO ₂ CH ₃
Z1.02	CH ₃	H	H	OSO ₂ CH ₃
Z1.03	H	CH ₃	H	OSO ₂ CH ₃
Z1.04	H	H	CH ₃	OSO ₂ CH ₃
Z1.05	CH ₃	CH ₃	H	OSO ₂ CH ₃
Z1.06	CH ₃	H	CH ₃	OSO ₂ CH ₃
Z1.07	H	CH ₃	CH ₃	OSO ₂ CH ₃
Z1.08	CH ₃	CH ₃	CH ₃	OSO ₂ CH ₃
Z1.09	H	H	H	OSO ₂ CH ₂ Ph
Z1.10	CH ₃	H	H	OSO ₂ CH ₂ Ph
Z1.11	H	CH ₃	H	OSO ₂ CH ₂ Ph
Z1.12	H	H	CH ₃	OSO ₂ CH ₂ Ph
Z1.13	CH ₃	CH ₃	H	OSO ₂ CH ₂ Ph
Z1.14	CH ₃	H	CH ₃	OSO ₂ CH ₂ Ph
Z1.15	H	CH ₃	CH ₃	OSO ₂ CH ₂ Ph
Z1.16	CH ₃	CH ₃	CH ₃	OSO ₂ CH ₂ Ph
Z1.17	H	H	H	Br
Z1.18	CH ₃	H	H	Br
Z1.19	H	CH ₃	H	Br
Z1.20	H	H	CH ₃	Br
Z1.21	CH ₃	CH ₃	H	Br
Z1.22	CH ₃	H	CH ₃	Br
Z1.23	H	CH ₃	CH ₃	Br
Z1.24	CH ₃	CH ₃	CH ₃	Br
Z1.25	H	H	H	Cl
Z1.26	CH ₃	H	H	Cl
Z1.27	H	CH ₃	H	Cl
Z1.28	H	H	CH ₃	Cl
Z1.29	CH ₃	CH ₃	H	Cl
Z1.30	CH ₃	H	CH ₃	Cl
Z1.31	H	CH ₃	CH ₃	Cl
Z1.32	CH ₃	CH ₃	CH ₃	Cl


The compounds of formula IV



wherein R_1 , R_2 and R_3 are as defined for formula I and wherein the ratio of compounds of formula IVa (trans)



wherein R_1 , R_2 and R_3 are as defined for formula I, to compounds of formula IVb (cis)




wherein R_1 , R_2 and R_3 are as defined for formula I, is from 2:1 to 20:1, are valuable intermediates in the preparation of compounds of formula I and were developed specifically for the present process according to the invention. The present invention accordingly relates also to those compounds.

Especially valuable for the preparation of compounds of formula I are those compounds of formula IV wherein the ratio of compounds of formula IVa (trans) to compounds of formula IVb (cis) is from 6:1 to 20:1, especially from 6:1 to 15:1.

As intermediates for the preparation of compounds of formula I there are especially suitable compounds of formula IV wherein R_1 , R_2 and R_3 are hydrogen.


Preferred compounds of formula IV are listed in the following Table:

Table 3: Compounds of formula IV

Comp. No.	R ₁	R ₂	R ₃
Z1.1	H	H	H
Z1.2	CH ₃	H	H
Z1.3	H	CH ₃	H
Z1.4	H	H	CH ₃
Z1.5	CH ₃	CH ₃	H
Z1.6	CH ₃	H	CH ₃
Z1.7	H	CH ₃	CH ₃
Z1.8	CH ₃	CH ₃	CH ₃


What is claimed is:

1. A process for the preparation of a compound of formula I



wherein R_1 , R_2 and R_3 are each independently of the others hydrogen or methyl, which comprises

a) reaction of a compound of formula II




wherein R_1 , R_2 and R_3 are as defined for formula I, either

a1) with triphenylphosphine dibromide or triphenylphosphine dichloride or


a2) with RSO_2Cl , wherein R is C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, benzyl, phenyl, nitrophenyl, halophenyl or C_1 - C_6 alkylphenyl, in the presence of a base,

to form a compound of formula III



wherein X is OSO_2R , wherein R is C_1 - C_4 alkyl, C_1 - C_4 fluoroalkyl, benzyl, phenyl, nitrophenyl, halophenyl or C_1 - C_6 alkylphenyl, or is bromine or chlorine and R_1 , R_2 and R_3 are as defined for formula I; and

b) reaction of that compound in the presence of a base to form a compound of formula IV




- 27 -

wherein R_1 , R_2 and R_3 are as defined for formula 1; and


c) conversion of that compound in the presence of a reducing agent into a compound of formula I.

2. A process according to claim 1, which comprises, after carrying out Process Step b) and before carrying out Process Step c), isomerisation of the compound of formula IV




wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1,

bb) in the presence of a base to form a compound of formula IV wherein the ratio of the compound of formula IVa (trans)




wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, to the compound of formula IVb (cis)




wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, is more than 6:1.

3. A compound of formula III




wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1 and X is as defined for formula III in claim 1.

4. Use of a compound of formula III




wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1 and X is as defined for formula III in claim 1, in the preparation of a compound of formula I according to claim 1.


5. A compound of formula IV



wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, and wherein the ratio of the compound of formula IVa (trans)




wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, to the compound of formula IVb (cis)




wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, is from 2:1 to 20:1.


6. Use of a compound of formula IV



wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, and wherein the ratio of the compound of formula IVa (trans)




wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, to the compound of formula IVb (cis)



wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, is from 2:1 to 20:1, in the preparation of a compound of formula I according to claim 1.

7. A compound of formula IV



wherein R_1 , R_2 and R_3 are as defined for formula I in claim 1, prepared in accordance with a process according to claim 1.

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2006/001508

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07C209/36

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 03/074491 A (SYNGENTA PARTICIPATIONS AG; EHRENFREUND, JOSEF; TOBLER, HANS; WALTER,) 12 September 2003 (2003-09-12) cited in the application examples	1-7
P,A	WO 2006/015865 A (SYNGENTA PARTICIPATIONS AG; WALTER, HARALD; NEUENSCHWANDER, URS; ZEUN,) 16 February 2006 (2006-02-16) page 27 - page 29; examples	1-7

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

28 June 2006

Date of mailing of the international search report

07/07/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Menegaki, F

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2006/001508

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 03074491	A	12-09-2003	AU 2003208490 A1	16-09-2003
			BR 0308230 A	28-12-2004
			CA 2477931 A1	12-09-2003
			CN 1639128 A	13-07-2005
			EG 23424 A	12-07-2005
			EP 1480955 A1	01-12-2004
			JP 2005532271 T	27-10-2005
			MX PA04008314 A	26-11-2004
			US 2005221989 A1	06-10-2005
<hr/>				
WO 2006015865	A	16-02-2006	NONE	
<hr/>				