

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
25 August 2005 (25.08.2005)

PCT

(10) International Publication Number
WO 2005/077901 A1

(51) International Patent Classification⁷: C07D 213/40,
213/61, A01N 43/40

European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(21) International Application Number:
PCT/EP2005/002563

(22) International Filing Date: 10 February 2005 (10.02.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
04356019.2 12 February 2004 (12.02.2004) EP
04356096.0 11 June 2004 (11.06.2004) EP
60/637,120 17 December 2004 (17.12.2004) US

(71) Applicant (for all designated States except US): BAYER CROPSCIENCE SA [FR/FR]; 16 rue Jean-Marie Leclair, F-69009 LYON (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GOUOT, Jean-Marie [FR/FR]; 3 Allée des Erables, F-69450 ST CYR AU MONT D'OR (FR). GROSJEAN-COURNOYER, Marie-Claire [FR/FR]; Le Trève Collonge, Route d'Albigny, F-69250 CURIS AU MONT D'OR (FR).

(74) Agent: AGASSE, Stéphane; BAYER CROPSCIENCE SA, Patents & Licensing Department, 14/20 rue Pierre Baizet, F-69009 LYON (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FUNGICIDAL COMPOSITION COMPRISING A PYRIDYLETHYLBENZAMIDE DERIVATIVE AND A COMPOUND CAPABLE OF INHIBITING THE TRANSPORT OF ELECTRONS OF THE RESPIRATORY CHAIN IN PHYTOPATHOGENIC FUNGAL ORGANISMS

(57) Abstract: A composition comprising at least a pyridylethylbenzamide derivative of general formula (I) (a) and a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms (b) in a (a) / (b) weight ratio of from 0.01 to 20. A composition further comprising an additional fungicidal compound. A method for preventively or curatively combating the phytopathogenic fungi of crops by using this composition.

WO 2005/077901 A1

Fungicidal composition comprising a pyridylethylbenzamide derivative and a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms


5 The present invention relates to novel fungicide compositions comprising a pyridylethylbenzamide derivative and a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms. The present invention also relates to a method of combating or controlling phytopathogenic fungi by applying at a locus infested or liable to be infested such a
10 composition.

International patent application WO 01/11965 generically discloses numerous pyridylethylbenzamide derivatives. The possibility of combining one or more of these numerous pyridylethylbenzamide derivatives with known fungicidal products
15 to develop a fungicidal activity is disclosed in general terms, without any specific example or biological data.

It is always of high-interest in agriculture to use novel pesticidal mixtures showing a synergistic effect in order notably to avoid or to control the development of resistant strains to the active ingredients or to the mixtures of known active
20 ingredients used by the farmer while minimising the doses of chemical products spread in the environment and reducing the cost of the treatment.

We have now found some novel fungicidal compositions which possess the above mentioned characteristics.

25 Accordingly, the present invention relates to a composition comprising :
a) a pyridylethylbenzamide derivative of general formula (I)



in which :

- p is an integer equal to 1, 2, 3 or 4;
- q is an integer equal to 1, 2, 3, 4 or 5;
- 30 - each substituent X is chosen, independently of the others, as being halogen, alkyl or haloalkyl;

- each substituent Y is chosen, independently of the others, as being halogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl, halothioalkyl, alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or
5 benzylsulphonyl;
as to the N-oxides of the compounds thereof;
and
b) a compound capable of inhibiting the transport of electrons of the respiratory
chain in phytopathogenic fungal organisms;
10 in a (a) / (b) weight ratio of from 0.01 to 20.

In the context of the present invention, a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is chosen from the group comprising a compound capable of inhibiting reduced
15 nicotinamide-adenine dinucleotide (NADH) deshydrogenase in phytopathogenic fungal organisms, a compound capable of inhibiting succinate dehydrogenase in phytopathogenic fungal organisms and a compound capable of inhibiting mitochondrial ubiquinol:ferricytochrome-c oxidoreductase in phytopathogenic fungal organisms.

20 A compound capable of inhibiting NADH deshydrogenase in phytopathogenic fungal organisms is also known as complex I inhibitor.

A compound capable of succinate dehydrogenase in phytopathogenic fungal organisms is also known as complex II inhibitor.

25 A compound capable of inhibiting mitochondrial ubiquinol:ferricytochrome-c oxidoreductase in phytopathogenic fungal organisms is also known as complex III inhibitor.

In the context of the present invention :

- halogen means chlorine, bromine, iodine or fluorine;
30 - each of the alkyl or acyl radicals present in the molecule contains from 1 to 10 carbon atoms, preferably from 1 to 7 carbon atoms, more preferably from 1 to 5 carbon atoms, and may be linear or branched;
- each of the alkenyl or alkynyl radicals present in the molecule contains from 2 to 10
carbon atoms, preferably from 2 to 7 carbon atoms, more preferably from 2 to 5
35 carbon atoms, and may be linear or branched.

The composition according to the present invention provides a synergistic effect. This synergistic effect allows a reduction of the chemical substances spread into the environment and a reduction of the cost of the fungal treatment.

In the context of the present invention, the term "synergistic effect" is defined by Colby according to the article entitled "Calculation of the synergistic and antagonistic responses of herbicide combinations" Weeds, (1967), 15, pages 20-22.

The latter article mentions the formula:

$$E = x + y - \frac{x * y}{100}$$

in which E represents the expected percentage of inhibition of the disease for the combination of the two fungicides at defined doses (for example equal to x and y respectively), x is the percentage of inhibition observed for the disease by the compound (I) at a defined dose (equal to x), y is the percentage of inhibition observed for the disease by the compound (II) at a defined dose (equal to y). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

The composition according to the present invention comprises a pyridylethylbenzamide derivative of general formula (I). Preferably, the present invention relates to a composition comprising a pyridylethylbenzamide derivative of general formula (I) in which the different characteristics may be chosen alone or in combination as being :

- as regards p, p is 2;
- as regards q, q is 1 or 2. More preferably, q is 2;
- as regards X, X is chosen, independently of the others, as being halogen or haloalkyl. More preferably, X is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group;
- as regards Y, Y is chosen, independently of the others, as being halogen or haloalkyl. More preferably, Y is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group;

More preferably, the pyridylethylbenzamide derivative of general formula (I) present in the composition of the present invention is :

- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 1);
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-iodobenzamide (compound 2); or

- N-{2-[3,5-dichloro-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 3).

Even more preferably, the pyridylethylbenzamide derivative of general formula (I) present in the composition of the present invention is N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 1).

5

The composition according to the present invention comprises a compound capable of inhibiting the transport of electrons of the respiratory chain.

Preferably, the present invention also relates to a composition comprising a complex I inhibitor. More preferably, the present invention relates to a composition comprising a complex I inhibitor which is diflumetorin.

10

Preferably, the present invention also relates to a composition comprising a complex II inhibitor. More preferably, the present invention relates to a composition comprising a complex II inhibitor selected from the carboxamide derivatives which may for example be *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1*H*-pyrazole-4-carboxamide, *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, benodanil, carboxin, fenfuram, flutolanil, furametypr, mepronil, boscalid, oxycarboxin, thifluzamide. *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, boscalid and oxycarboxin are preferred.

15

20

Preferably, the present invention also relates to a composition comprising a complex III inhibitor. More preferably, the present invention relates to a composition comprising a complex III inhibitor selected from strobilurin derivatives, cyazofamid, fenamidone or famoxadone. Fenamidone is preferred.

25

Strobilurin derivatives are also preferred. According to the present invention, strobilurin derivatives may for example be azoxystrobin, dimoxystrobin, fluoxastrobin, kresoxim-methyl, metominostrobin, trifloxystrobin, pyraclostrobin, picoxystrobin or 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-*N*-methylacetamide. 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-*N*-methylacetamide, azoxystrobin, trifloxystrobin and fluoxastrobin are preferred.

30

35

The composition according to the present invention comprises (a) at least a pyridylethylbenzamide derivative of general formula (I) and (b) a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms in an (a) / (b) weight ratio of from 0.01 to 20; preferably of from 0.05 to 10; even more preferably, of from 0.1 to 5.

The composition of the present invention may further comprise at least one other different fungicide active ingredient (c).

The fungicidal active ingredient (c) may be selected from azaconazole, azoxystrobin, (Z)-N-[α -(cyclopropylmethoxyimino)-2,3-difluoro-6-(trifluoromethyl)benzyl]-2-phenylacetamide, 6-iodo-2-propoxy-3-propylquinazolin-4(3H)-one, benalaxyl, benomyl, benthiavalicarb, biphenyl, bitertanol, blasticidin-S, boscalid, borax, bromuconazole, bupirimate, sec-butylamine, calcium polysulfide, captafol, captan, carbendazim, carboxin, carpropamid, chinomethionat, chlorothalonil, chlozolate, copper hydroxide, copper octanoate, copper oxychloride, copper sulfate, cuprous oxide, cyazofamid, cymoxanil, cyproconazole, cyprodinil, dazomet, debacarb, dichlofluanid, dichlorophen, diclobutrazole, diclocymet, diclomezine, dicloran, diethofencarb, difenoconazole, difenzoquat metilsulfate, difenzoquat, diflumetorim, dimethirimol, dimethomorph, diniconazole, dinobuton, dinocap, diphenylamine, dithianon, dodemorph, dodemorph acetate, dodine, edifenphos, epoxiconazole, ethaboxam, etaconazole, ethirimol, ethoxyquin, etridiazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fenfuram, fenhexamid, fenciclonil, fenoxanil, fenpropidin, fenpropimorph, fentin, fentin hydroxide, fentin acetate, ferbam, ferimzone, fluazinam, fludioxonil, fluoroimide, fluoxastrobin, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpet, formaldehyde, fosetyl, fosetyl-aluminium, fuberidazole, furalaxyl, furametpyr, guazatine, guazatine acetates, hexachlorobenzene, hexaconazole, 8-hydroxyquinoline sulfate, potassium hydroxyquinoline sulfate, hymexazol, imazalil sulfate, imazalil, imibenconazole, iminoctadine, iminoctadine triacetate, ipconazole, iprobenfos, iprodione, iprovalicarb, isoprothiolane, kasugamycin, kasugamycin hydrochloride hydrate, kresoxim-methyl, mancozeb, maneb, mepanipyrim, mepronil, mercuric chloride, mercuric oxide, mercurous chloride, metalaxyl, metalaxyl-M, metam-sodium, metam, metconazole, methasulfocarb, methyl isothiocyanate, metiram, metominostrobin, mildiomyacin, myclobutanil, nabam, nickel bis(dimethyldithiocarbamate), nitrothal-isopropyl, nuarimol, octhilinone, ofurace, oleic acid, oxadixyl, oxine-copper, oxpoconazole fumarate, oxycarboxin,

pefurazoate, penconazole, pencycuron, pentachlorophenol, sodium pentachlorophenoxide, pentachlorophenyl laurate, phenylmercury acetate, sodium 2-phenylphenoxide, 2-phenylphenol, phosphorous acid, phthalide, picoxystrobin, piperalin, polyoxinspolyoxin B, polyoxin, polyoxorim, probenazole, prochloraz, 5 procymidone, propamocarb hydrochloride, propamocarb, propiconazole, propineb, prothioconazole, pyraclostrobin, pyrazophos, pyributicarb, pyrifenox, pyrimethanil, pyroquilon, quinoxyfen, quintozone, silthiofam, simeconazole, spiroxamine, sulfur, tar oils, tebuconazole, tecnazene, tetraconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, 10 triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, validamycin, vinclozolin, zineb, ziram and zoxamide.

Preferably, fungicidal active ingredient (c) is selected from captane, folpet, dodine, propineb, mancozeb, thiram, tolylfluanid, iminoctadine, dithianon, copper hydroxide, copper octanoate, copper oxychloride, copper sulfate, fosetyl-Al, 15 phosphorous acid, cymoxanil, iprovalicarb, bentiavalicarb, chlorotalonil, propamocarb, prothioconazole, tebuconazole and spiroxamine.

Where the third active ingredient (c) as defined above is present in the composition, this compound may be present in an amount of (a) : (b) : (c) weight 20 ratio of from 1 : 0.01 : 0.01 to 1 : 20 : 20; the ratios of compound (a) and compound (c) varying independently from each other. Preferably, the (a) : (b) : (c) weight ratio may be of from 1 : 0.05 : 0.05 to 1 : 10 : 10.

Following compositions may be cited to illustrate in a non-limited manner the 25 present invention : compound 1 with diflumetorin, compound 1 with *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, compound 1 with *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, compound 1 with *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, compound 30 1 with benodanil, compound 1 with carboxin, compound 1 with fenfuram, compound 1 with flutolanil, compound 1 with furametpyr, compound 1 with mepronil, compound 1 with boscalid, compound 1 with oxycarboxin, compound 1 with thifluzamide, compound 1 with cyazofamid, compound 1 with fenamidone, compound 1 with famoxadone. azoxystrobin, compound 1 with dimoxystrobin, 35 compound 1 with fluoxastrobin, compound 1 with kresoxim-methyl, compound 1 with metominostrobin, compound 1 with trifloxystrobin, compound 1 with

pyraclostrobin, compound 1 with picoxystrobin, compound 1 with 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-N-methylacetamide, compound 2 with diflumetorin, compound 2 with *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, compound 2 with *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, compound 2 with *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, compound 2 with benodanil, compound 2 with carboxin, compound 2 with fenfuram, compound 2 with flutolanil, compound 2 with furametpyr, compound 2 with mepronil, compound 2 with boscalid, compound 2 with oxycarboxin, compound 2 with thifluzamide, compound 2 with cyazofamid, compound 2 with fenamidone, compound 2 with famoxadone. azoxystrobin, compound 2 with dimoxystrobin, compound 2 with fluoxastrobin, compound 2 with kresoxim-methyl, compound 2 with metominostrobin, compound 2 with trifloxystrobin, compound 2 with pyraclostrobin, compound 2 with picoxystrobin, compound 2 with 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-N-methylacetamide, compound 3 with diflumetorin, compound 3 with *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, compound 3 with *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, compound 3 with *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, compound 3 with benodanil, compound 3 with carboxin, compound 3 with fenfuram, compound 3 with flutolanil, compound 3 with furametpyr, compound 3 with mepronil, compound 3 with boscalid, compound 3 with oxycarboxin, compound 3 with thifluzamide, compound 3 with cyazofamid, compound 3 with fenamidone, compound 3 with famoxadone. azoxystrobin, compound 3 with dimoxystrobin, compound 3 with fluoxastrobin, compound 3 with kresoxim-methyl, compound 3 with metominostrobin, compound 3 with trifloxystrobin, compound 3 with pyraclostrobin, compound 3 with picoxystrobin, compound 3 with 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-N-methylacetamide.

The composition according to the present invention may further comprise an other additional component such as an agriculturally acceptable support, carrier or filler.

In the present specification, the term "support" denotes a natural or synthetic, organic or inorganic material with which the active material is combined to make it easier to apply, notably to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support may be a solid or a liquid.
5 Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports may also be used.

The composition may also comprise other additional components. In particular, the composition may further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention may be made, for example, of polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the above compounds containing sulphate, sulphonate and phosphate functions. The presence of at least one surfactant is generally essential
10 when the active material and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content may be comprised between 5% and 40% by weight of the composition.

Additional components may also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active materials can be combined with any solid or liquid additive, which complies with the usual formulation techniques.
25

In general, the composition according to the invention may contain from 0.05 to 99% (by weight) of active material, preferably 10 to 70% by weight.

Compositions according to the present invention can be used in various forms such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable concentrate, oil miscible liquid, paste, plant rodlet, powder for dry seed treatment,
35 seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed

treatment, suspension concentrate (flowable concentrate), ultra low volume (ulv) liquid, ultra low volume (ulv) suspension, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder.

5 These compositions include not only compositions which are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions which must be diluted before they are applied to the crop.

10 The fungicidal compositions of the present invention can be used to curatively or preventively control phytopathogenic fungi of crops. Thus, according to a further aspect of the present invention, there is provided a method for preventively or curatively controlling phytopathogenic fungi of crops characterised in that a fungicidal composition as hereinbefore defined is applied to the seed, the plant and/or to the fruit
15 of the plant or to the soil in which the plant is growing or in which it is desired to grow.

The composition as used against phytopathogenic fungi of crops comprises an effective and non-phytotoxic amount of an active material of general formula (I).

The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention which is sufficient to control or destroy the
20 fungi present or liable to appear on the crops, and which does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be combated or controlled, the type of crop, the climatic conditions and the compounds included in the fungicidal composition according to the invention.

25 This amount can be determined by systematic field trials, which are within the capabilities of a person skilled in the art.

The method of treatment according to the present invention is useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or seedlings pricking out and plants or plants pricking out. This method of treatment
30 can also be useful to treat roots. The method of treatment according to the present invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruits of the concerned plant.

Among the plants that can be protected by the method according to the invention, mention may be made of cotton; flax; vine; fruit crops such as *Rosaceae*
35 *sp.* (for instance pip fruits such as apples and pears, but also stone fruits such as apricots, almonds and peaches), *Ribesioideae sp.*, *Juglandaceae sp.*, *Betulaceae sp.*,

Anacardiaceae sp., *Fagaceae sp.*, *Moraceae sp.*, *Oleaceae sp.*, *Actinidaceae sp.*,
Lauraceae sp., *Musaceae sp.* (for instance banana trees and plantains), *Rubiaceae sp.*,
Theaceae sp., *Sterculiaceae sp.*, *Rutaceae sp.* (for instance lemons, oranges and
grapefruits); leguminous crops such as *Solanaceae sp.* (for instance tomatoes),
5 *Liliaceae sp.*, *Asteraceae sp.* (for instance lettuces), *Umbelliferae sp.*, *Cruciferae sp.*,
Chenopodiaceae sp., *Cucurbitaceae sp.*, *Papilionaceae sp.* (for instance peas),
Rosaceae sp. (for instance strawberries); big crops such as *Graminae sp.* (for
instance maize, cereals such as wheat, rice, barley and triticale), *Asteraceae sp.* (for
instance sunflower), *Cruciferae sp.* (for instance colza), *Papilionaceae sp.* (for
10 instance soja), *Solanaceae sp.* (for instance potatoes), *Chenopodiaceae sp.* (for
instance beetroots); horticultural and forest crops; as well as genetically modified
homologues of these crops.

Among the plants and the possible diseases of these plants protected by the
method according to the present invention, mention may be made of :

- 15 - wheat, as regards controlling the following seed diseases: fusaria
(*Microdochium nivale* and *Fusarium roseum*), stinking smut (*Tilletia caries*, *Tilletia
controversa* or *Tilletia indica*), septoria disease (*Septoria nodorum*) and loose smut;
- wheat, as regards controlling the following diseases of the aerial parts of the
plant: cereal eyespot (*Tapesia yallundae*, *Tapesia acuiformis*), take-all
20 (*Gaeumannomyces graminis*), foot blight (*F. culmorum*, *F. graminearum*), black speck
(*Rhizoctonia cerealis*), powdery mildew (*Erysiphe graminis forma specie tritici*), rusts
(*Puccinia striiformis* and *Puccinia recondita*) and septoria diseases (*Septoria tritici* and
Septoria nodorum);
- wheat and barley, as regards controlling bacterial and viral diseases, for
25 example barley yellow mosaic;
- barley, as regards controlling the following seed diseases: net blotch
(*Pyrenophora graminea*, *Pyrenophora teres* and *Cochliobolus sativus*), loose smut
(*Ustilago nuda*) and fusaria (*Microdochium nivale* and *Fusarium roseum*);
- barley, as regards controlling the following diseases of the aerial parts of the
30 plant: cereal eyespot (*Tapesia yallundae*), net blotch (*Pyrenophora teres* and
Cochliobolus sativus), powdery mildew (*Erysiphe graminis forma specie hordei*),
dwarf leaf rust (*Puccinia hordei*) and leaf blotch (*Rhynchosporium secalis*);
- potato, as regards controlling tuber diseases (in particular *Helminthosporium
solani*, *Phoma tuberosa*, *Rhizoctonia solani*, *Fusarium solani*), mildew (*Phytophthora
35 infestans*) and certain viruses (virus Y);

- potato, as regards controlling the following foliage diseases: early blight (*Alternaria solani*), mildew (*Phytophthora infestans*);
- cotton, as regards controlling the following diseases of young plants grown from seeds: damping-off and collar rot (*Rhizoctonia solani*, *Fusarium oxysporum*) and
5 black root rot (*Thielaviopsis basicola*);
- protein yielding crops, for example peas, as regards controlling the following seed diseases: anthracnose (*Ascochyta pisi*, *Mycosphaerella pinodes*), fusaria (*Fusarium oxysporum*), grey mould (*Botrytis cinerea*) and mildew (*Peronospora pisi*);
- 10 - oil-bearing crops, for example rape, as regards controlling the following seed diseases: *Phoma lingam*, *Alternaria brassicae* and *Sclerotinia sclerotiorum*;
- corn, as regards controlling seed diseases: (*Rhizopus* sp., *Penicillium* sp., *Trichoderma* sp., *Aspergillus* sp., and *Gibberella fujikuroi*);
- flax, as regards controlling the seed disease: *Alternaria linicola*;
- 15 - forest trees, as regards controlling damping-off (*Fusarium oxysporum*, *Rhizoctonia solani*);
- rice, as regards controlling the following diseases of the aerial parts: blast disease (*Magnaporthe grisea*), bordered sheath spot (*Rhizoctonia solani*);
- leguminous crops, as regards controlling the following diseases of seeds or
20 of young plants grown from seeds: damping-off and collar rot (*Fusarium oxysporum*, *Fusarium roseum*, *Rhizoctonia solani*, *Pythium* sp.);
- leguminous crops, as regards controlling the following diseases of the aerial parts: grey mould (*Botrytis* sp.), powdery mildews (in particular *Erysiphe cichoracearum*, *Sphaerotheca fuliginea* and *Leveillula taurica*), fusaria (*Fusarium oxysporum*, *Fusarium roseum*), leaf spot (*Cladosporium* sp.), alternaria leaf spot
25 (*Alternaria* sp.), anthracnose (*Colletotrichum* sp.), septoria leaf spot (*Septoria* sp.), black speck (*Rhizoctonia solani*), mildews (for example *Bremia lactucae*, *Peronospora* sp., *Pseudoperonospora* sp., *Phytophthora* sp.);
- fruit trees, as regards diseases of the aerial parts: monilia disease (*Monilia fructigenae*, *M. laxa*), scab (*Venturia inaequalis*), powdery mildew (*Podosphaera leucotricha*);
30
- vine, as regards diseases of the foliage: in particular grey mould (*Botrytis cinerea*), powdery mildew (*Uncinula necator*), black rot (*Guignardia biwelli*) and mildew (*Plasmopara viticola*);

- beetroot, as regards the following diseases of the aerial parts: cercospora blight (*Cercospora beticola*), powdery mildew (*Erysiphe beticola*), leaf spot (*Ramularia beticola*).

5 The fungicidal composition according to the present invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists
10 in contacting one or more compounds of the present invention, or a composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

 The fungicidal composition according to the present invention may also be
15 used in the treatment of genetically modified organisms with the compounds according to the invention or the agrochemical compositions according to the invention. Genetically modified plants are plants into in which genome a heterologous gene encoding a protein of interest has been stably integrated. The expression "heterologous gene encoding a protein of interest" essentially means
20 genes which give the transformed plant new agronomic properties, or genes for improving the agronomic quality of the transformed plant.

 The dose of active material usually applied in the treatment according to the present invention is generally and advantageously between 10 and 2000 g/ha,
25 preferably between 20 and 1500 g/ha for applications in foliar treatment. The dose of active substance applied is generally and advantageously between 1 and 200 g per 100 kg of seed, preferably between 2 and 150 g per 100 kg of seed in the case of seed treatment. It is clearly understood that the doses indicated above are given as illustrative examples of the invention. A person skilled in the art will know how to
30 adapt the application doses according to the nature of the crop to be treated.

 The compositions according to the present invention may also be used fore the preparation of composition useful to curatively or preventively treat human and animal fungal diseases such as, for example, mycoses, dermatoses, trichophyton
35 diseases and candidiases or diseases caused by *Aspergillus spp.* or *Candida spp.*, for example *Aspergillus fumigatus* or *Candida albicans* respectively.

The present invention will now be illustrated with the following examples :

**Example 1 : Efficacy against *Pyrenophora teres* of a composition containing N-
5 {2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide
(Compound 1) and trifloxystrobin**

The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain
10 the desired active material concentration.

Barley plants (Express variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm tall) by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

15 After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Pyrenophora teres* spores (12 000 spores per ml). The spores are collected from a 12-day-old culture. The contaminated barley plants are incubated for 24 hours at about 20°C and at 100% relative humidity, and then for 12 days at 80% relative humidity.

20 Grading (% of efficacy) is carried out 12 days after the contamination, in comparison with the control plants.

The following table summarises the results obtained when tested compound 1 and trifloxystrobin alone and in a 1/1 weight ratio mixture.

25

	Dose (g/ha)	% Efficacy	Synergism (Colby)
Compound 1	4	20	-
	8	50	-
Trifloxystrobin	4	50	-
	8	70	-
Compound 1 + trifloxystrobin (Ratio 1/1)	4+4	85	+25
	8+8	90	+5

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

Example 2 : Efficacy against *Alternaria brassicae* (in vitro) of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and trifloxystrobin

5

The growth of *Alternaria brassicae* was followed on PDA medium containing the compounds alone or in mixtures.

The PDA medium is prepared by mixing 39 grams of PDA (Merck) in 1 liter of demineralised water. The medium is sterilized by autoclave 15 minutes at 121°C. The compounds are dissolved in acetone and added to the medium before pouring it in Petri dishes. After medium drying, the fungus is inoculated on the medium and allowed to growth at 19°C.

The percentage of efficacy is noted after 14 days of growth (measurement of the radial growth) of the fungus in comparison with a control.

15

The following table summarises the results obtained when tested compound 1 and trifloxystrobin alone and in different weight ratio mixtures.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	0.025	34	-
	0.05	45	-
	0.1	53	-
	0.25	76	-
	0.5	90	-
Trifloxystrobin	0.025	45	-
	0.05	44	-
	0.1	45	-
	0.25	53	-
	0.5	60	-
	1	65	-
	2.5	71	-
Compound 1 + trifloxystrobin (Ratio 1/1)	0.05+0.05	79	+10
	0.1+0.1	93	+19
	0.5+0.5	100	+4

Compound 1 + trifloxystrobin (Ratio 1/2)	0.05+0.1	85	+15
	0.25+0.5	100	+10
Compound 1 + trifloxystrobin (Ratio 2/1)	0.1+0.05	97	+23
	0.5+0.25	100	+5
Compound 1 + trifloxystrobin (Ratio 1/4)	0.025+0.1	75	+11
Compound 1 + trifloxystrobin (Ratio 4/1)	0.1+0.025	100	+26
Compound 1 + trifloxystrobin (Ratio 1/10)	0.05+0.5	96	+18
	0.25+2.5	100	+7

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

5 **Example 3 : Efficacy against *Alternaria brassicae* (in vitro) of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and azoxystrobin**

10 The growth of *Alternaria brassicae* was followed on PDA medium containing the compounds alone or in mixtures.

The PDA medium is prepared by mixing 39 grams of PDA (Merck) in 1 liter of demineralised water. The medium is sterilized by autoclave 15 minutes at 121°C. The compounds are dissolved in acetone and added to the medium before pouring it in Petri dishes. After medium drying, the fungus is inoculated on the medium and
15 allowed to growth at 19°C.

The percentage of efficacy is noted after 14 days of growth (measurement of the radial growth) of the fungus in comparison with a control.

20 The following table summarises the results obtained when tested compound 1 and azoxystrobin alone and in different weight ratio mixtures.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	0.025	34	-
	0.05	45	-
	0.1	53	-
	0.25	76	-
	0.5	90	-
Azoxystrobin	0.025	38	-
	0.05	41	-
	0.1	47	-
	0.25	56	-
	0.5	47	-
	1	48	-
	2.5	49	-
Compound 1 + azoxystrobin (Ratio 1/1)	0.05+0.05	70	+2
	0.1+0.1	88	+13
	0.5+0.5	100	+5
Compound 1 + azoxystrobin (Ratio 1/2)	0.05+0.1	76	+5
	0.25+0.5	95	+8
Compound 1 + azoxystrobin (Ratio 2/1)	0.1+0.05	81	+9
	0.5+0.25	98	+2
Compound 1 + azoxystrobin (Ratio 4/1)	0.1+0.025	100	+29
Compound 1 + azoxystrobin (Ratio 1/10)	0.05+0.5	86	+15
	0.25+2.5	100	+12

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

Example 4 : Efficacy against *Erysiphe graminis f. sp. tritici* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and fluoxastrobin

5 The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

 Wheat plants (Audace variety) in starter cups, sown on 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm
10 tall) by spraying with the aqueous suspension described above.

 Plants, used as controls, are treated with an aqueous solution not containing the active material.

 After 24 hours, the plants are contaminated by dusting them with *Erysiphe graminis f. sp. tritici* spores, the dusting being carried out using diseased plants.

15 Grading is carried out 7 to 14 days after the contamination, in comparison with the control plants.

 The following table summarises the results obtained when tested compound 1 and fluoxastrobin alone and in a 8:1 weight ratio mixture.

20

	Dose (g/ha)	% Efficacy	Synergism (Colby)
Compound 1	125	20	-
Fluoxastrobin	15	20	-
Compound 1 + fluoxastrobin (Ratio 8/1)	125 + 15	70	+ 34

 According to the Colby method, a synergistic effect of the mixture tested has been observed.

Example 5 : Efficacy against *Botrytis cinerea* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and kresoxim-methyl

5 The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

 Gherkin plants (Petit vert de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon Z11
10 stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

 After 24 hours, the plants are contaminated by depositing drops of an aqueous suspension of *Botrytis cinerea* spores (150,000 spores per ml) on upper surface of the leaves. The spores are collected from a 15-day-old culture and are suspended in a
15 nutrient solution composed of :

- 20 g/L of gelatine
- 50 g/L of cane sugar
- 2 g/L of NH₄NO₃
- 1 g/L of KH₂PO₄

20 The contaminated gherkin plants are settled for 5/7 days in a climatic room at 15-11°C (day/night) and at 80% relative humidity. Grading is carried out 5 to 7 days after contamination, in comparison with the control plants.

 The following table summarises the results obtained when tested compound 1
25 and kresoxim-methyl alone and in different weight ratio mixtures.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	37	0	-
	111	60	-
	333	90	-
Kresoxim-methyl	37	0	-
	111	0	-
	333	0	-
Compound 1 + kresoxim-methyl (Ratio 1/1)	111 + 111	100	40

Compound 1 + kresoxim-methyl (Ratio 1/3)	37 + 111	95	95
	111 + 333	100	40
Compound 1 + kresoxim-methyl (Ratio 1/9)	37 + 333	100	100

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

5 **Example 6 : Efficacy against *Botrytis cinerea* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and picoxystrobin**

10 The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

15 Gherkin plants (Petit vert de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon Z11 stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by depositing drops of an aqueous suspension of *Botrytis cinerea* spores (150,000 spores per ml) on upper surface of the leaves. The spores are collected from a 15-day-old culture and are suspended in a nutrient solution composed of :

- 20
- 20 g/L of gelatine
 - 50 g/L of cane sugar
 - 2 g/L of NH₄NO₃
 - 1 g/L of KH₂PO₄

25 The contaminated gherkin plants are settled for 5/7 days in a climatic room at 15-11°C (day/night) and at 80% relative humidity. Grading is carried out 5 to 7 days after contamination, in comparison with the control plants.

The following table summarises the results obtained when tested compound 1 and picoxystrobin alone and in different weight ratio mixtures.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	37	0	-
	111	60	-
	333	100	-
Picoxystrobin	37	0	-
	111	35	-
	333	40	-
Compound 1 + picoxystrobin (Ratio 1/1)	111 + 111	95	+21
Compound 1 + picoxystrobin (Ratio 1/3)	37 + 111	90	+ 55
Compound 1 + picoxystrobin (Ratio 3/1)	111 + 37	98	+ 38
Compound 1 + picoxystrobin (Ratio 1/9)	37 + 333	95	+ 55

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

5

Example 7 : Efficacy against *Botrytis cinerea* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and pyraclostrobin

10 The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

15 Gherkin plants (Petit vert de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon Z11 stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by depositing drops of an aqueous suspension of *Botrytis cinerea* spores (150,000 spores per ml) on upper surface of the

leaves. The spores are collected from a 15-day-old culture and are suspended in a nutrient solution composed of :

- 20 g/L of gelatine
- 50 g/L of cane sugar
- 5 - 2 g/L of NH_4NO_3
- 1 g/L of KH_2PO_4

The contaminated gherkin plants are settled for 5/7 days in a climatic room at 15-11°C (day/night) and at 80% relative humidity. Grading is carried out 5 to 7 days after contamination, in comparison with the control plants.

10

The following table summarises the results obtained when tested compound 1 and pyraclostrobin alone and in different weight ratio mixtures.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	12	0	-
	37	40	-
	111	92	-
Pyraclostrobin	37	0	-
	111	0	-
	333	0	-
Compound 1 + pyraclostrobin (Ratio 1/27)	12.3 + 333	100	+ 100
Compound 1 + pyraclostrobin (Ratio 1/9)	37 + 333	100	+ 60
	12.3 + 111	85	+ 85
Compound 1 + pyraclostrobin (Ratio 3/1)	37 + 12.3	55	+ 15

15

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

Example 8: Efficacy against *Botrytis cinerea* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and 2-{2-[6-(3-Chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-N-methylacetamide (Compound A)

5

The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

Gherkin plants (Petit vert de Paris variety) in starter cups, sown on a 50/50
10 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon Z11 stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by depositing drops of an aqueous
15 suspension of *Botrytis cinerea* spores (150,000 spores per ml) on upper surface of the leaves. The spores are collected from a 15-day-old culture and are suspended in a nutrient solution composed of :

- 20 g/L of gelatine
- 50 g/L of cane sugar
- 2 g/L of NH₄NO₃
- 20 - 1 g/L of KH₂PO₄

The contaminated gherkin plants are settled for 5/7 days in a climatic room at 15-11°C (day/night) and at 80% relative humidity. Grading is carried out 5 to 7 days after contamination, in comparison with the control plants.

25 The following table summarises the results obtained when tested compound 1 and compound A alone and in different weight ratio mixtures.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	37	3	-
	111	58	-
	333	75	-
Compound A	37	0	-
	111	0	-
	333	0	-

Compound 1 + compound A (Ratio 1/1)	333 + 333	100	+ 25
	111 + 111	100	+ 42
Compound 1 + compound A (Ratio 3/1)	333 + 111	100	+ 25
Compound 1 + compound A (Ratio 1/3)	111 + 333	100	+ 42
Compound 1 + compound A (Ratio 1/9)	37 + 333	75	+ 72

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

5

Example 9: Efficacy against *Erysiphe graminis* f. sp. *tritici* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and dimoxystrobin

10 The formulated compounds are diluted with water to obtain the desired active material concentration.

Wheat plants (Audace variety) in starter cups, sown on 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm tall) by spraying with the aqueous suspension described above.

15 Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by dusting them with *Erysiphe graminis* f. sp. *tritici* spores, the dusting being carried out using diseased plants.

20 Grading is carried out 7 to 14 days after the contamination, in comparison with the control plants.

The following table summarises the results obtained when tested compound 1 and dimoxystrobin alone and in a 1/8 weight ratio mixture.

	Dose (g/ha)	% Efficacy	Synergism (Colby)
Compound 1	250	50	-
Dimoxystrobin	62,5	0	-

Compound 1 + dimoxystrobin (Ratio 1/8)	250 + 62,5	95	+ 45
---	------------	----	------

According to the Colby method, a synergistic effect of the mixture tested has been observed.

5 **Example 10: Efficacy against *Botrytis cinerea* of a composition containing N-[2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2-trifluoromethylbenzamide (Compound 1) and N-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide (Compound B)**

10 The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

15 Gherkin plants (Petit vert de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon Z11 stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

20 After 24 hours, the plants are contaminated by depositing drops of an aqueous suspension of *Botrytis cinerea* spores (150,000 spores per ml) on upper surface of the leaves. The spores are collected from a 15-day-old culture and are suspended in a nutrient solution composed of :

- 20 g/L of gelatine
- 50 g/L of cane sugar
- 2 g/L of NH₄NO₃
- 1 g/L of KH₂PO₄

25 The contaminated gherkin plants are settled for 5/7 days in a climatic room at 15-11°C (day/night) and at 80% relative humidity. Grading is carried out 5 to 7 days after contamination, in comparison with the control plants.

30 The following table summarises the results obtained when tested compound 1 and compound B alone and in a 9/1 weight ratio mixture.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	37	72	-
Compound B	4	32	-
Compound 1 + compound B (Ratio 9/1)	37 + 4	100	+ 19

According to the Colby method, a synergistic effect of the mixture tested has
5 been observed.

Example 11 : Efficacy against *Sphaerotheca fuliginea* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and boscalid

10

The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

15 Gherkin plants (Vert petit de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 20°C/23°C, are treated at the 2-leaves stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

20 After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Sphaerotheca fuliginea* spores (100 000 spores per ml). The spores are collected from a contaminated plants .The contaminated gherkin plants are incubated at about 20°C/25°C and at 60/70% relative humidity.

Grading (% of efficacy) is carried out 21 days after the contamination, in comparison with the control plants.

25 The following table summarises the results obtained when tested compound 1 and boscalid alone and in a 1/1 weight ratio mixture.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	4	10	-
	8	15	-
	15	25	-
Boscalid	4	0	-
	8	10	-
	15	45	-
Compound 1 + boscalid (Ratio 1/1)	4+4	40	+30
	8+8	70	+47
	15+15	95	+36

According to the Colby method, a synergistic effect of the mixtures tested has
5 been observed.

**Example 12 : Efficacy against *Sphaerotheca fuliginea* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and N-(3',4'-dichloro-5-
10 fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1H-pyrazole-4-carboxamide (Compound C)**

The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain
15 the desired active material concentration.

Gherkin plants (Vert petit de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 20°C/23°C, are treated at the 2-leaves stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

20 After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Sphaerotheca fuliginea* spores (100 000 spores per ml). The spores are collected from a contaminated plants .The contaminated gherkin plants are incubated at about 20°C/25°C and at 60/70% relative humidity.

Grading (% of efficacy) is carried out 21 days after the contamination, in comparison with the control plants.

The following table summarises the results obtained when tested compound 1 and compound C alone and in different weight ratio mixtures.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	31.2	86	-
	15.6	18	-
	7.8	10	-
Compound C	62.5	0	-
	31.2	0	-
Compound 1 + compound C (Ratio 1/4)	15.6 + 62.5	87	40
	7.8 + 31	73	25
Compound 1 + compound C (Ratio 1/8)	7.8 + 62.5	100	59

According to the Colby method, a synergistic effect of the mixture tested has been observed.

10

Example 13 : Efficacy against *Alternaria brassicae* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethyl-benzamide (Compound 1) and fenamidone

15

The formulated compounds are diluted with water to obtain the desired active material concentration.

Radish plants (Pernot variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon stage by spraying with the aqueous suspension described above.

20

Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Alternaria brassicae* spores (40,000 spores per cm³). The spores are collected from a 12-13-day-old culture.

The contaminated radish plants are incubated for 6-7 days at about 18°C, under a humid atmosphere.

Grading is carried out 6 to 7 days after the contamination, in comparison with the control plants.

5

The following table summarises the results obtained when tested compound 1 and fenamidone alone and in a 1/27 weight ratio mixture.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	4	40	-
Fenamidone	111	55	-
Compound 1 + fenamidone (Ratio 1/27)	4 + 111	73	+ 17

10 According to the Colby method, a synergistic effect of the mixture tested has been observed.

Example 14 : Efficacy against *Botrytis cinerea* of a composition containing N-[2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl]-2-trifluoromethylbenzamide

15 **(Compound 1) N-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide (Compound D)**

The formulated compounds are diluted with water to obtain the desired active material concentration.

20 Gherkin plants (Petit vert de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 18-20°C, are treated at the cotyledon Z11 stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

25 After 24 hours, the plants are contaminated by depositing drops of an aqueous suspension of *Botrytis cinerea* spores (150,000 spores per ml) on upper surface of the leaves. The spores are collected from a 15-day-old culture and are suspended in a nutrient solution composed of :

- 20 g/L of gelatine
- 50 g/L of cane sugar

- 2 g/L of NH₄NO₃

- 1 g/L of KH₂PO₄

The contaminated gherkin plants are settled for 5/7 days in a climatic room at 15-11°C (day/night) and at 80% relative humidity. Grading is carried out 5 to 7 days after contamination, in comparison with the control plants.

The following table summarises the results obtained when tested compound 1 and compound D alone and in different weight ratio mixtures.


	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	31.2	20	-
	62.5	40	-
Compound D	250	60	-
Compound 1 + compound D (Ratio 8:1)	31,2 + 250	85	+ 17
Compound 1 + compound D (Ratio 4:1)	62.5 + 250	95	+ 19

10

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

CLAIMS

1. A composition comprising :
5 a) a pyridylethylbenzamide derivative of general formula (I)



in which :

- p is an integer equal to 1, 2, 3 or 4;
 - q is an integer equal to 1, 2, 3, 4 or 5;
 - each substituent X is chosen, independently of the others, as being halogen, alkyl or haloalkyl;
 - 10 - each substituent Y is chosen, independently of the others, as being halogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl, halothioalkyl, alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or
 - 15 benzylsulphonyl;
- as to the N-oxides of 2-pyridine thereof;
and
- b) a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms;
- 20 in a (a) / (b) weight ratio of from 0.01 to 20.

2. A composition according to claim 1, characterised in that p is 2.
3. A composition according to claim 1 or 2, characterised in that q is or 2.
- 25
4. A composition according to any of the claims 1 to 3, characterised in that X is chosen, independently of the others, as being halogen or haloalkyl.
5. A composition according to any of the claims 1 to 4, characterised in that X is
- 30 chosen independently of the others, as being a chlorine atom or a trifluoromethyl group.

6. A composition according to any of the claims 1 to 5, characterised in that Y is chosen, independently of the others, as being halogen or haloalkyl.
- 5 7. A composition according to any of the claims 1 to 6, characterised in that Y is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group.
8. A composition according to any of the claims 1 to 7, characterised in that the
10 compound of general formula (I) is :
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide;
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-iodobenzamide; or
- N-{2-[3,5-dichloro-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide .
- 15 9. A composition according to claim 8, characterised in that the compound of general formula (I) is N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide.
- 10 10. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is a compound capable of inhibiting reduced nicotinamide-adenine dinucleotide dehydrogenase in phytopathogenic fungal organisms.
- 25 11. A composition according to claim 10, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is diflumetorin.
- 30 12. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is a compound capable of inhibiting succinate dehydrogenase in phytopathogenic fungal organisms.
- 35 13. A composition according to claim 12 characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain of succinate dehydrogenase in phytopathogenic fungal organisms is *N*-[2-(1,3-dimethyl-butyl)-

phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, benodanil, carboxin, fenfuram, flutolanil, furametpyr, mepronil,
5 boscalid, oxycarboxin or thifluzamide.

14. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is a compound capable of inhibiting
10 mitochondrial ubiquinol:ferricytochrome-c oxidoreductase in phytopathogenic fungal organisms.

15. A composition according to claim 14, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain of
15 mitochondrial ubiquinol:ferricytochrome-c oxidoreductase in phytopathogenic fungal organisms is a strobilurin derivative, cyazofamid, fenamidone or famoxadone.

16. A composition according to claim 15, characterised in that the strobilurin derivative is azoxystrobin, dimoxystrobin, fluoxastrobin, kresoxim-methyl,
20 metominostrobin, trifloxystrobin, pyraclostrobin, picoxystrobin or 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-*N*-methylacetamide.

17. A composition according to any one of the claims 1 to 16 further comprising
25 a fungicidal compound (c).

18. A composition according to claim 17, characterised in that the fungicidal compound (c) is selected from captane, folpet, dodine, propineb, mancozeb, thiram, tolylfluanid, iminoctadine, dithianon, copper hydroxide, copper octanoate, copper
30 oxychloride, copper sulfate, fosetyl-Al, phosphorous acid, cymoxanil, iprovalicarb, benthiavalicarb, chlorotalonil, propamocarb, prothioconazole, tebuconazole and spiroxamine.

19. A composition according to any one of the claims 1 to 18, characterised in
35 that it further comprises an agriculturally acceptable support, carrier, filler and/or surfactant.

20. A method for preventively or curatively controlling phytopathogenic fungi of crops, characterised in that an effective and non-phytotoxic amount of a composition according to any one of the claims 1 to 19 is applied to the seed, the plant and/or to the fruit of the plant or to the soil in which the plant is growing or in which it is desired to grow.
- 5

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2005/002563

A. CLASSIFICATION OF SUBJECT MATTER		
IPC 7 C07D213/40 C07D213/61 A01N43/40		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2 899 437 A (U.S.VITAMIN & PHARMACEUTICAL CORPORATION) 11 August 1959 (1959-08-11) column 1, line 32, "fungicides"; column 2, Table I, "Py = Y"; -----	1-20
Y	WO 01/11965 A (AVENTIS CROPSCIENCE GMBH ;BRIGGS GEOFFREY GOWER (FR); STEELE CHRIS) 22 February 2001 (2001-02-22) cited in the application page 35, compound 401; claims; -----	1-20
Y	FR 2 821 718 A (AVENTIS CROPSCIENCE SA) 13 September 2002 (2002-09-13) page 5, lines 11-22; page 6, lines 4-8; page 22, lines 5-19; claims -----	1-20
	-/--	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
° Special categories of cited documents:		
A document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed		*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family
Date of the actual completion of the international search 29 June 2005		Date of mailing of the international search report 11/07/2005
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016		Authorized officer Sen, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2005/002563

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 03/041501 A (BAYER CROPSCIENCE S A ;MERCER RICHARD (FR); WEGMANN THOMAS (FR)) 22 May 2003 (2003-05-22) page 8, lines 12-33; page 12, lines 11-16 and 21-22; page 19, line 22 - page 21, line 13; claims -----	1-20

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP2005/002563

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2899437	A	11-08-1959	NONE

WO 0111965	A	22-02-2001	AT 270817 T 15-07-2004
			AU 7278500 A 13-03-2001
			BR 0013371 A 07-05-2002
			CN 1370047 A 18-09-2002
			DE 60012188 D1 19-08-2004
			WO 0111965 A1 22-02-2001
			EP 1204323 A1 15-05-2002
			ES 2220533 T3 16-12-2004
			JP 2003506465 T 18-02-2003
			MX PA02001453 A 28-01-2003
			US 6821992 B1 23-11-2004

FR 2821718	A	13-09-2002	FR 2821718 A1 13-09-2002
			AT 283630 T 15-12-2004
			DE 60202132 D1 05-01-2005
			EG 23127 A 28-04-2004
			WO 02069712 A1 12-09-2002
			EP 1365653 A1 03-12-2003
			ES 2231700 T3 16-05-2005
			JP 2004523564 T 05-08-2004
			US 2004106578 A1 03-06-2004

WO 03041501	A	22-05-2003	FR 2832031 A1 16-05-2003
			BR 0214180 A 31-08-2004
			CA 2463718 A1 22-05-2003
			CN 1585603 A 23-02-2005
			WO 03041501 A1 22-05-2003
			EP 1443821 A1 11-08-2004
			JP 2005508991 T 07-04-2005
			US 2004266729 A1 30-12-2004
