

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 February 2006 (23.02.2006)

PCT

(10) International Publication Number
WO 2006/018609 A2

(51) International Patent Classification:
A61K 31/439 (2006.01) *C07D 453/06* (2006.01)
A61K 31/55 (2006.01) *C07D 487/08* (2006.01)

(74) Agent: DAVIES, Jonathan, Mark; Reddie & Grose, 16
Theobalds Road, London WC1X 8PL (GB).

(21) International Application Number:
PCT/GB2005/003139

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ,
OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL,
SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,
VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 10 August 2005 (10.08.2005)

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language: English

Published:

— without international search report and to be republished
upon receipt of that report

(26) Publication Language: English

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


(30) Priority Data:
0418375.2 18 August 2004 (18.08.2004) GB

(71) Applicant (for all designated States except US): CAM-
BRIDGE UNIVERSITY TECHNICAL SERVICES
LIMITED [GB/GB]; The Old Schools, Trinity Lane,
Cambridge CB2 1TS (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): GRAINGER,
David, John [GB/GB]; Department of Medicine, Cam-
bridge University, Box 157, Addenbrookes Hospital,
Hill Road, Cambridge CB2 2QQ (GB). FOX, David,
John [GB/GB]; Department of Chemistry, Cambridge
University, Lensfield Road, Cambridge CB1 1EW (GB).

(54) Title: ANTI-INFLAMMATORY AGENTS



WO 2006/018609 A2

(57) Abstract: The invention provides compounds, compositions, and uses of compounds of general formula (I) or pharmaceutically acceptable salts thereof, for the preparation of a medicament intended to treat an inflammatory disorder wherein y is any integer from 1 to 8; z is any integer from 1 to 8; with the proviso that y and z cannot both equal 1; X is $-C-(Y)_k-(R')_n$ or $SO_2-(Y)_k-(R')_n$; k is 0 or 1; Y is a cycloalkyl or polycyloalkyl group (such as an adamantyl, adamantanemethyl, bicyclooctyl, cyclohexyl, cyclopropyl group); or Y is a cycloalkenyl or polycycloalkenyl group.

Anti-Inflammatory Agents

The invention relates to the use of α -aminobicyclolactams for preparing a medicament intended to prevent or treat inflammatory disorders.

Inflammation is an important component of physiological host defence. Increasingly, however, it is clear that temporally or spatially inappropriate inflammatory responses play a part in a wide range of diseases, including those with an obvious leukocyte component (such as autoimmune diseases, asthma or atherosclerosis) but also in diseases that have not traditionally been considered to involve leukocytes (such as osteoporosis or Alzheimer's disease).

The chemokines are a large family of signalling molecules with homology to interleukin-8 which have been implicated in regulating leukocyte trafficking both in physiological and pathological conditions. With more than fifty ligands and twenty receptors involved in chemokine signalling, the system has the requisite information density to address leukocytes through the complex immune regulatory processes from the bone marrow, to the periphery, then back through secondary lymphoid organs. However, this complexity of the chemokine system has at first hindered pharmacological approaches to modulating inflammatory responses through chemokine receptor blockade. It has proved difficult to determine which chemokine receptor(s) should be inhibited to produce therapeutic benefit in a given inflammatory disease.

More recently, a family of agents which block signalling by a wide range of chemokines simultaneously has been described: Reckless et al., *Biochem J.* (1999) 340:803-811. The first such agent, a peptide termed "Peptide 3", was found to inhibit leukocyte migration induced by 5 different chemokines, while leaving migration in response to other chemoattractants (such as fMLP or TGF-beta) unaltered. This peptide, and its analogs such as NR58-3.14.3 (i.e. Sequence ID No.1 c(DCys-DGln-DIle-DTrp-DLys-DGln-DLys-DPro-DAsp-DLeu-DCys)-NH₂), are collectively termed "Broad Spectrum Chemokine Inhibitors" (BSCIs). Grainger et al., *Biochem. Pharm.* 65 (2003) 1027-1034 have subsequently shown BSCIs to have potentially useful anti-inflammatory activity in a range of animal models of diseases. Interestingly, simultaneous blockade of multiple chemokines is not apparently associated with acute or chronic toxicity, suggesting this

approach may be a useful strategy for developing new anti-inflammatory medications with similar benefits to steroids but with reduced side-effects.

However, peptides and peptoid derivatives such as NR58-3.14.3, may not be optimal for use in vivo. They are quite expensive to synthesise and have relatively unfavourable pharmacokinetic and pharmacodynamic properties. For example, NR58-3.14.3 is not orally bioavailable and is cleared from blood plasma with a half-life period of less than 30 minutes after intravenous injection.

Two parallel strategies have been adopted to identify novel preparations which retain the anti-inflammatory properties of peptide 3 and NR58-3.14.3, but have improved characteristics for use as pharmaceuticals. Firstly, a series of peptide analogs have been developed, some of which have longer plasma half-lives than NR58-3.14.3 and which are considerably cheaper to synthesise. Secondly, a detailed structure: activity analysis of the peptides has been carried out to identify the key pharmacophores and design small non-peptidic structures which retain the beneficial properties of the original peptide.

This second approach yielded several structurally distinct series of compounds which retained the anti-inflammatory properties of the peptides, including 16-amino and 16-aminoalkyl derivatives of the alkaloid yohimbine, as well as a range of N-substituted 3-aminoglutarimides. (Reference: Fox et al., J Med Chem 45(2002) 360-370: WO 99/12968 and WO 00/42071.) All of these compounds are broad-spectrum chemokine inhibitors which retain selectivity over non-chemokine chemoattractants, and a number of them have been shown to block acute inflammation in vivo.

The most potent and selective of these compounds was (S)-3-(undec-10-enoyl)-aminoglutarimide (NR58,4), which inhibited chemokine-induced migration in vitro with an ED₅₀ of 5nM. However, further studies revealed that the aminoglutarimide ring was susceptible to enzymatic ring opening in serum. Consequently, for some applications (for example, where the inflammation under treatment is chronic, such as in autoimmune diseases) these compounds may not have optimal properties, and a more stable compound with similar anti-inflammatory properties may be superior.


As an approach to identifying such stable analogs, various derivatives of (S)- 3-(undec-10-enoyl)-aminoglutarimide have been tested for their stability in serum. One such derivative, the 6-deoxo analog (S)-3-(undec-10-enoyl)-tetrahydropyridin-2-one, is completely stable in human serum for at least 7 days at 37°C, but has considerably reduced potency compared with the parental molecule.

Amide derivatives of 3-aminocaprolactam have already been disclosed in the art. For example:

- Japanese patent application No. 09087331 describes 3-aminocaprolactam amide derivatives wherein the amide alkyl side chain may contain from 2 to 30 carbon atoms. These compounds have been presented as oil-gelating agents.
- US patent No. 6,395,282 describes immunogenic conjugates comprising a carrier molecule coupled to an autoinducer of a Gram negative bacteria, wherein said autoinducer can be a 3-aminocaprolactam amide derivative wherein the amide alkyl side chain may contain up to 34 carbon atoms. However, a therapeutic use is disclosed only for the conjugates and not for the isolated amide derivative.
- An article by Weiss et al. (*Research Communications in Psychology, Psychiatry and Behavior* (1992), 17(3-4), 153-159) discloses a series of 3-aminocaprolactam amide derivatives, and among others 3-hexanamido-DL- ϵ -caprolactam and 3-dodecanamido-DL- ϵ -caprolactam. These compounds are presented as having only an *in vitro* activity but no significant *in vivo* effect.

In other words, though some alkyl amide derivatives of 3-aminocaprolactam have certainly been known in the art, no actual pharmaceutical use has been described for 3-aminocaprolactam amide derivatives.

The invention provides the use of a compound of general formula (I), or a pharmaceutically acceptable salt thereof, for the preparation of a medicament intended to treat inflammatory disorder:



(I)

wherein

y is any integer from 1 to 8;

z is any integer from 1 to 8; with the proviso that y and z cannot both equal 1;

X is $-\text{CO}-(\text{Y})_k-(\text{R}^1)_n$ or $\text{SO}_2-(\text{Y})_k-(\text{R}^1)_n$;

k is 0 or 1

Y is a cycloalkyl or polycyloalkyl group (such as an adamantyl, adamantanemethyl, bicyclooctyl, cyclohexyl, cyclopropyl group);

or Y is a cycloalkenyl or polycycloalkenyl group;

each R^1 is independently selected from hydrogen or an alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl or alkylamino radical of 1 to 20 carbon atoms (for example of 5 to 20 carbon atoms, of 8 to 20 carbon atoms, of 9 to 20 carbon atoms, of 10 to 18 carbon atoms, of 12 to 18 carbon atoms, of 13 to 18 carbon atoms, of 14 to 18 carbon atoms, of 13 to 17 carbon atoms);

or each R^1 is independently selected from fluoro, chloro, bromo, iodo, hydroxy, oxyalkyl, amino, aminoalkyl or aminodialkyl radical; and

n is any integer from 1 to m, where m is the maximum number of substitutions permissible on the cyclo-group Y.


Alternatively R^1 may be selected from a peptido radical, for example having from 1 to 4 peptidic moieties linked together by peptide bonds (for example a peptido radical of 1 to 4 amino acid residues).

This class of compounds are described as α -aminobicyclolactams since the key structural features of the molecule are the lactam amide in a bicycloalkyl ring system, with an amino group attached to the carbon atom next to the lactam carbonyl group (termed the α -carbon).

The α -carbon of α -aminobicyclolactams may be asymmetric (where y and z are not equal; ie. $y > z$ in the general formula (I)) and consequently, some of the compounds according to the present invention have two possible enantiomeric forms, that is, the "R" and "S" configurations. The present invention encompasses the two enantiomeric forms and all combinations of these forms, including the racemic "RS" mixtures. With a view to simplicity, when no specific configuration is shown in the structural formulae, it should be understood that the two enantiomeric forms and their mixtures are represented.

The compounds of general formula (I) are N-substituted α -aminobicyclolactams, or their pharmaceutically acceptable salts. The N-substituent is either a carbon amide or a sulfonamide. The geometry of the carbon atom next to the carbonyl of the carbon amide or the sulfonyl group of the sulfonamide (the "key" carbon) may be important for the bioactivity of the molecule. The nature of the N-substituent may be such that the ring or rings of Y constrain the bond angles at the "key"-carbon to be essentially tetrahedral (i.e. sp³ hybrid bonds). Any substituent R¹ may be a substituent at any permissible position on the ring or rings of the cyclo-group Y. In particular it is to be noted that the invention includes compounds in which the "key carbon" is both part of the cyclo group and is itself substituted. The definition of (R¹)_n encompasses compounds of the invention with no substitution (i.e. R¹ = hydrogen), compounds of the invention with mono substitution (i.e. R¹ is not hydrogen and n = 1), and also multiple substitution (i.e. at least two R¹ groups are not hydrogen and n = 2 or more).

The invention also provides pharmaceutical compositions comprising, as active ingredient, a compound of general formula (I), or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient and/or carrier:



(I)

wherein

y is any integer from 1 to 8;

z is any integer from 1 to 8; with the proviso that y and z cannot both equal 1;

X is $-\text{CO}-(\text{Y})_k-(\text{R}^1)_n$ or $\text{SO}_2-(\text{Y})_k-(\text{R}^1)_n$;

k is 0 or 1

Y is a cycloalkyl or polycyloalkyl group (such as an adamantlyl, adamantanemethyl, bicyclooctyl, cyclohexyl, cyclopropyl group);

or Y is a cycloalkenyl or polycycloalkenyl group;

each R¹ is independently selected from hydrogen or an alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl or alkylamino radical of 1 to 20 carbon atoms (for example of 5 to 20 carbon atoms, of 8 to 20 carbon atoms, of 9 to 20 carbon atoms, of 10 to 18 carbon atoms, of 12 to 18 carbon atoms, of 13 to 18 carbon atoms, of 14 to 18 carbon atoms, of 13 to 17 carbon atoms);

or each R¹ is independently selected from fluoro, chloro, bromo, iodo, hydroxy, oxyalkyl, amino, aminoalkyl or aminodialkyl radical; and

n is any integer from 1 to m, where m is the maximum number of substitutions permissible on the cyclo-group Y.


Alternatively R¹ may be selected from a peptido radical, for example having from 1 to 4 peptidic moieties linked together by peptide bonds (for example a peptido radical of 1 to 4 amino acid residues).

By pharmaceutically acceptable salt is meant in particular the addition salts of inorganic acids such as hydrochloride, hydrobromide, hydroiodide, sulphate, phosphate, diphosphate and nitrate or of organic acids such as acetate, maleate, fumarate, tartrate, succinate, citrate, lactate, methanesulphonate, p-toluenesulphonate, palmoate and stearate. Also within the scope of the present invention, when they can be used, are the salts formed from bases such as sodium or potassium hydroxide. For other examples of pharmaceutically acceptable salts, reference can be made to "Salt selection for basic drugs", *Int. J. Pharm.* (1986), 33, 201-217.

The pharmaceutical composition can be in the form of a solid, for example powders, granules, tablets, gelatin capsules, liposomes or suppositories. Appropriate solid supports can be, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidine and wax. Other appropriate pharmaceutically acceptable excipients and/or carriers will be known to those skilled in the art.

The pharmaceutical compositions according to the invention can also be presented in liquid form, for example, solutions, emulsions, suspensions or syrups. Appropriate liquid supports can be, for example, water, organic solvents such as glycerol or glycols, as well as their mixtures, in varying proportions, in water.

The invention also provides compounds and salts thereof of general formula (I)



(I)

wherein

y is any integer from 1 to 8;

z is any integer from 1 to 8; with the proviso that y and z cannot both equal 1;

X is $-\text{CO}-(\text{Y})_k-(\text{R}^1)_n$ or $\text{SO}_2-(\text{Y})_k-(\text{R}^1)_n$;

k is 0 or 1

Y is a cycloalkyl or polycyloalkyl group (such as an adamantyl, adamantanemethyl, bicyclooctyl, cyclohexyl, cyclopropyl group);

or Y is a cycloalkenyl or polycycloalkenyl group;

each R^1 is independently selected from hydrogen or an alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl or alkylamino radical of 1 to 20 carbon atoms (for example of 5 to 20 carbon atoms, of 8 to 20 carbon atoms, of 9 to 20 carbon atoms, of 10 to 18 carbon atoms, of 12 to 18 carbon atoms, of 13 to 18 carbon atoms, of 14 to 18 carbon atoms, of 13 to 17 carbon atoms);

or each R^1 is independently selected from fluoro, chloro, bromo, iodo, hydroxy, oxyalkyl, amino, aminoalkyl or aminodialkyl radical; and

n is any integer from 1 to m, where m is the maximum number of substitutions permissible on the cyclo-group Y.

Alternatively R^1 may be selected from a peptido radical, for example having from 1 to 4 peptidic moieties linked together by peptide bonds (for example a peptido radical of 1 to 4 amino acid residues).

The invention provides compounds, compositions and uses of the compounds of general formula (I) or their pharmaceutically acceptable salts, wherein the alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl or alkylamino part of the R¹ radical is either linear or is branched but contains a linear chain of at least 8 or at least 10 carbon atoms.

The invention provides compounds, compositions and uses wherein the R¹ radical has a "key" carbon which is di-substituted with the same or different groups selected from: alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl and alkylamino radicals.

The invention provides compounds, compositions and uses wherein the "key"-carbon is chiral.

The invention provides compounds, compositions and uses wherein the "key"-carbon has sp³ hybridised bonds.

The invention provides compounds, compositions and uses wherein the "key"-carbon has essentially tetrahedral bond angles.

The compounds of general formula (I) when used in the invention, or their salts, may be such that the ring or rings of Y constrain the bond angles at the "key"-carbon to be essentially tetrahedral (i.e. sp³ hybrid bonds).

In an alternative embodiment of the invention, general formula (I) is modified such that the C3-C7 alkyl bridge -(CH₂)_y- is replaced by a bridging group independently selectable from the group consisting of alkenyl, haloalkyl, alkylamino and alkylhydroxy moieties having a carbon chain length of from 1 to 8.

The invention provides a use, composition or compound wherein y and z are the same integer, whereby the α -aminobicyclolactam is non-chiral.

The invention provides a use, composition or compound wherein y and z are not the same integer, whereby the α -aminobicyclolactam ring is chiral.

The invention provides a use, composition or compound wherein z is 3 and y is 1 or 2 or 4-8, whereby the compound contains a lactam ring which is seven membered.

The invention provides a use, composition or compound wherein z is 2 and y is 1 or 3-8, whereby the compound contains a lactam ring which is 6 membered.

In particular, preferred compounds of general formula (I) and their salts according to the present invention are selected from the group consisting of:

4-(Adamantane-1-carbonylamino)-3-oxo-2-aza-bicyclo[2.2.2]octane

5-(Adamantane-1-carbonylamino)-10-oxo-9-aza-bicyclo[3.3.2]decane

4-(2',2'-dimethyldodecanoylamino)-3-oxo-2-aza-bicyclo[2.2.2]octane

5-(2',2'-dimethyldodecanoylamino)-10-oxo-9-aza-bicyclo[3.3.2]decane

and the salts thereof.

The invention also provides the sulfonamide analogues of the exemplified compounds: i.e. the sulfonyl- α -aminobicyclolactam equivalents of the said compounds.

The invention includes compounds, compositions and uses thereof as defined, wherein the compound is in hydrated or solvated form.

The amide and sulfonamide derivatives of α -aminobicyclolactams described here are functional BSCIs. They are stable in human serum and consequently have excellent pharmacokinetic properties; they are orally bioavailable; they are highly potent broad-spectrum chemokine inhibitors *in vitro* with excellent selectivity over non-chemokine chemoattractants; they are highly potent and effective anti-inflammatory agents *in vivo* in rodent models of inflammation; their administration is not associated with any significant acute toxicity at the doses necessary to achieve a maximal therapeutic effect. Taken together, these properties suggest that amide and sulfonamide derivatives of α -aminobicyclolactams represent anti-inflammatory medications with advantages over previously described compounds.

The invention is based on a crystal structure of the head group in a highly bioactive molecule: namely a BSCI with sub-nanomolar potency *in vitro*; (S)-3-(adamantane-1-carbonylamino)azepin-2-one.

Surprisingly, this crystal structure shows that the azepan-2-one ring adopts a particular conformation likely associated with high biological activity, and demonstrates that the

torsional angle of the $-N-C-C-N-$ group of atoms (that is the bonds running from the sidechain amide through the the α -carbon to the ring lactam nitrogen) is an important determinant of bioactivity. This torsional angle can be controlled by bridging C3 and C7, since the axial hydrogens at these positions are pointing towards each other. As a result, the invention provides compounds of general formula (I), in which a bicyclic head group is generated. The number of carbon atoms in the bridge will determine the angles of the axial substituents at C3 and C7.

In comparison to the prior art the improvement of the present invention lies in the provision of a bridged aminolactam moiety allowing the $-N-C-C-N-$ torsional angles to be precisely controlled, so that a compound from the series with any particularly advantageous set of properties determined by the conformation of the aminolactam ring can be selected.

Prior art peptides (such as NR58-3.14.3) have the disadvantages that: (a) they are expensive and require solid phase synthesis (at least for the longer ones) and (b) they clear very quickly via the kidneys and (c) they are generally less potent.

The prior art aminoglutaramides are cheap, not cleared quickly via the kidneys and more potent BUT they do not show metabolic stability.

The improvement described here is a class of compounds, the N-substituted α -aminobicyclic lactams, which are even more potent and metabolically stable

According to this invention, inflammatory disorders intended to be prevented or treated by the compounds of general formula (I) or the pharmaceutically acceptable salts thereof or pharmaceutical compositions or medicaments containing them as active ingredients include notably:

- autoimmune diseases, for example such as multiple sclerosis;
- vascular disorders including stroke, coronary artery diseases, myocardial infarction, unstable angina pectoris, atherosclerosis or vasculitis, e. g., Behçet's syndrome, giant cell arteritis, polymyalgia rheumatica, Wegener's granulomatosis, Churg-Strauss syndrome vasculitis, Henoch-Schönlein purpura and Kawasaki disease;
- viral infection or replication, e.g. infections due to or replication of viruses including pox virus, herpes virus (e. g., Herpesvirus simini), cytomegalovirus (CMV) or lentivirus;

- asthma;
- osteoporosis; (low bone mineral density);
- tumor growth;
- rheumatoid arthritis;
- organ transplant rejection and/or delayed graft or organ function, e.g. in renal transplant patients;
- a disorder characterised by an elevated TNF- α level;
- psoriasis;
- skin wounds;
- disorders caused by intracellular parasites such as malaria or tuberculosis;
- allergies; or
- Alzheimer's disease.

According to this invention, further inflammatory disorders include:

- ALS;
- fibrosis (particularly pulmonary fibrosis, but not limited to fibrosis in the lung);
- the formation of adhesions (particularly in the peritoneum and pelvic region).
- antigen induced recall response
- immune response suppression

These clinical indications fall under the general definition of inflammatory disorders or disorders characterized by elevated TNF α levels.

Where legally permissible, the invention also provides a method of treatment, amelioration or prophylaxis of the symptoms of an inflammatory disease (including an adverse inflammatory reaction to any agent) by the administration to a patient of an anti-inflammatory amount of a compound, composition or medicament as claimed herein.

Administration of a medicament according to the invention can be carried out by topical, oral, parenteral route, by intramuscular injection, etc.

The administration dose envisaged for a medicament according to the invention is comprised between 0.1 mg and 10 g depending on the type of active compound used.

DEFINITIONS

The term "about" refers to an interval around the considered value. As used in this patent application, "about X" means an interval from X minus 10% of X to X plus 10% of X, and preferably an interval from X minus 5% of X to X plus 5% of X.

The use of a numerical range in this description is intended unambiguously to include within the scope of the invention all individual integers within the range and all the combinations of upper and lower limit numbers within the broadest scope of the given range. Hence, for example, the range of 1 to 20 carbon atoms specified in respect of (*inter alia*) formula I is intended to include all integers between 1 and 20 and all sub-ranges of each combination of upper and lower numbers, whether exemplified explicitly or not.

As used herein, the term "comprising" is to be read as meaning both comprising and consisting of. Consequently, where the invention relates to a "pharmaceutical composition comprising as active ingredient" a compound, this terminology is intended to cover both compositions in which other active ingredients may be present and also compositions which consist only of one active ingredient as defined.

The term "peptidic moieties" used herein is intended to include the following 20 naturally-occurring proteogenic amino acid residues:

SYMBOL:	MEANING
Ala	Alanine
Cys	Cysteine
Asp	Aspartic Acid
Glu	Glutamic Acid
Phe	Phenylalanine

Gly	Glycine
His	Histidine
Ile	Isoleucine
Lys	Lysine
Leu	Leucine
Met	Methionine
Asn	Asparagine
Pro	Proline
Gln	Glutamine
Arg	Arginine
Ser	Serine
Thr	Threonine
Val	Valine
Trp	Tryptophan
Tyr	Tyrosine

Modified and unusual amino acid residues, as well as peptido-mimetics, are also intended to be encompassed within the definition of “peptidic moieties”.

Unless otherwise defined, all the technical and scientific terms used here have the same meaning as that usually understood by an ordinary specialist in the field to which this invention belongs. Similarly, all the publications, patent applications, all the patents and all other references mentioned here are incorporated by way of reference (where legally permissible).


The following examples are presented in order to illustrate the above procedures and should in no way be considered to limit the scope of the invention.

FIGURES


Figure 1 shows the projected crystal structure of (S)-3-(adamantane-1-carbonylamino)azepin-2-one. In the Figure dark grey = C; light grey = H; white = O; and black = N. The C3 and C7 positions of the lactam ring are marked. It can be seen that the –N-C-C-N- torsional angles in the lactam ring are at or near zero.

EXAMPLES*General procedure for the synthesis of the starting compounds*


Fetzion et al. *Bull. Soc. Chim. Fr.* (1969) 194 describe the synthesis of the α -carboxylbicyclolactam (II). This compound can be converted using the sequence of reactions known in the art as a Curtius Rearrangement to generate the analogous α -aminobicyclolactam, which can subsequently be N-substituted with a range of suitable carboxylic acids, including adamantane-1-carboxylic acid or 2',2'-dimethyldodecan-1-oic acid, using a wide range of known amide coupling methodologies (such as DCC coupling), as shown below;




where X has the same meaning as in the definition of general formula (I). The method of Fetzion can be readily adapted to synthesise the α -carboxylate analogs of other members of the α -aminobicyclolactam class (such as the 9-aza-10-oxo-bicyclo[3,3,2]decane in examples 2 and 4 below). These α -carboxylates can be converted into examples of compounds under the invention using the same sequence of reactions outlined above for compound (II).

EXAMPLE 1


4-(Adamantane-1-carbonylamino)-3-oxo-2-aza-bicyclo[2.2.2]octane

EXAMPLE 2

5-(Adamantane-1-carbonylamino)-10-oxo-9-aza-bicyclo[3.3.2]decane

EXAMPLE 3

4-(2',2'-dimethyldodecanoylamino)-3-oxo-2-aza-bicyclo[2.2.2]octane

EXAMPLE 4

5-(2',2'-dimethyldodecanoylamino)-10-oxo-9-aza-bicyclo[3.3.2]decane

Pharmacological study of the products of the invention**Inhibition of MCP-1 induced leukocyte migration*****Assay principle***

The biological activity of the compounds of the current invention may be demonstrated using any of a broad range of functional assays of leukocyte migration in vitro, including

but not limited to Boyden chamber and related transwell migration assays, under-agarose migration assays and direct visualisation chambers such as the Dunn Chamber.

For example, to demonstrate the inhibition of leukocyte migration in response to chemokines (but not other chemoattractants) the 96-well format micro transwell assay system from Neuroprobe (Gaithersburg, MD, USA) has been used. In principle, this assay consists of two chambers separated by a porous membrane. The chemoattractant is placed in the lower compartment and the cells are placed in the upper compartment. After incubation for a period at 37°C the cells move towards the chemoattractant, and the number of cells in the lower compartment is proportional to the chemoattractant activity (relative to a series of controls).

This assay can be used with a range of different leukocyte populations. For example, freshly prepared human peripheral blood leukocytes may be used. Alternatively, leukocyte subsets may be prepared, including polymorphonuclear cells or lymphocytes or monocytes using methods well known to those skilled in the art such as density gradient centrifugation or magnetic bead separations. Alternatively, immortal cell lines which have been extensively validated as models of human peripheral blood leukocytes may be used, including, but not limited to THP-1 cells as a model of monocytes or Jurkat cells as model of naïve T cells.

Although a range of conditions for the assay are acceptable to demonstrate the inhibition of chemokine-induced leukocyte migration, a specific example is hereby provided.

Materials

The transwell migration systems are manufactured by Neuroprobe, Gaithersburg, MD, USA.

The plates used are ChemoTx plates (Neuroprobe 101-8) and 30 µl clear plates (Neuroprobe MP30).

Geys' Balanced Salt Solution is purchased from Sigma (Sigma G-9779).

Fatty acid-free BSA is purchased from Sigma (Sigma A-8806).

MTT, i.e. 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, is purchased from Sigma (Sigma M-5655).

RPMI-1640 without phenol red is purchased from Sigma (Sigma R-8755).

The THP-1 cell line (European Cell culture Collection) were used as the leukocyte cell population.

Test protocol

The following procedure is used for testing the invention compounds for MCP-1 induced leukocyte migration:

First, the cell suspension to be placed in the upper compartment is prepared. The THP-1 cells are pelleted by centrifugation (770 x g; 4 mins) and washed with Geys Balanced Salt Solution with 1mg/ml BSA (GBSS + BSA). This wash is then repeated, and the cells repelleted before being resuspended in a small volume of GBSS + BSA for counting, for example using a standard haemocytometer.

The volume of GBSS + BSA is then adjusted depending on the number of cells present so that the cells are at final density of 4.45×10^6 cells per ml of GBSS + BSA. This ensures that there are 100,000 THP-1 cells in each 25 μ l of the solution that will be placed in the upper chamber of the plate.

To test a single compound for its ability to inhibit MCP-1 induced migration, it is necessary to prepare two lots of cells. The suspension of THP-1 cells at 4.45×10^6 cells/ml is divided into two pots. To one pot the inhibitor under test is added at an appropriate final concentration, in an appropriate vehicle (for example at 1 μ M in not more than 1% DMSO). To the second pot an equal volume of GBSS + BSA plus vehicle as appropriate (e.g. not more than 1% DMSO) is added to act as a control.

Next, the chemoattractant solution to be placed in the lower compartment is prepared. MCP-1 is diluted in GBSS + BSA to give a final concentration of 25 ng/ml. This is divided into two pots, as for the cell suspension. To one pot, the test compound is added to the same final concentration as was added to the cell suspension, while to the other pot an equal volume of GBSS + BSA plus vehicle as appropriate (e.g. not more than 1% DMSO) is added.

Note that the volume of liquid that needs to be added to make the addition of the test compound needs to be taken into account, when establishing the final concentration of MCP-1 in the solution for the lower compartment and the final concentration of cells in the upper compartment.

Once the chemoattractant solutions for the lower wells and cell solutions for the upper chambers have been prepared, the migration chamber should be assembled. Place 29 μ l of the appropriate chemoattractant solution into the lower well of the chamber. Assays should be performed with at least triplicate determinations of each condition. Once all the lower chambers have been filled, apply the porous membrane to the chamber in accordance with the manufacturer's instructions. Finally, apply 25 μ l of the appropriate cell solution to each upper chamber. A plastic lid is placed over the entire apparatus to prevent evaporation.

The assembled chamber is incubated at 37 °C, 5% CO₂, for 2 hours. A suspension of cells in GBSS + BSA is also incubated under identical conditions in a tube: these cells will be used to construct a standard curve for determining the number of cells that have migrated to the lower chamber under each condition.

At the end of the incubation, the liquid cell suspension is gently removed from the upper chamber, and 20 μ l of ice-cold 20mM EDTA in PBS is added to the upper chamber, and the apparatus is incubated at 4°C for 15 mins. This procedure causes any cells adhering to the underside of the membrane to fall into the lower chamber.

After this incubation the filter is carefully flushed with GBSS + BSA to wash off the EDTA, and then the filter is removed.

The number of cells migrated into the lower chamber under each condition can then be determined by a number of methods, including direct counting, labelling with fluorescent or radioactive markers or through the use of a vital dye. Typically, we utilise the vital dye MTT. 3 μ l of stock MTT solution are added to each well, and then the plate is incubated at 37 °C for 1-2 hours during which time dehydrogenase enzymes within the cells convert the soluble MTT to an insoluble blue formazan product that can be quantified spectrophotometrically.

In parallel, an 8-point standard curve is set up. Starting with the number of cells added to each upper chamber (100,000) and going down in 2-fold serial dilutions in GBSS + BSA, the cells are added to a plate in 25 μ l, with 3 μ l of MTT stock solution added. The standard curve plate is incubated along side the migration plate.

At the end of this incubation, the liquid is carefully removed from the lower chambers, taking care not to disturb the precipitated formazan product. After allowing to air dry briefly, 20 μ l of DMSO is added to each lower chamber to solubilise the blue dye, and

absorbance at 595nm is determined using a 96-well plate reader. The absorbance of each well is then interpolated to the standard curve to estimate the number of cells in each lower chamber.

The MCP-1 stimulated migration is determined by subtracting the average number of cells that reached the lower compartment in wells where no MCP-1 was added from the average number of cells that reached the lower compartment where MCP-1 was present at 25ng/ml.

The impact of the test substance is calculated by comparing the MCP-1-induced migration which occurred in the presence or absence of various concentrations of the test substance. Typically, the inhibition of migration is expressed as a percentage of the total MCP-1 induced migration which was blocked by the presence of the compound. For most compounds, a dose-response graph is constructed by determining the inhibition of MCP-1 induced migration which occurs at a range of different compound concentrations (typically ranging from 1nM to 1 μ M or higher in the case of poorly active compounds). The inhibitory activity of each compound is then expressed as the concentration of compound required to reduce the MCP-1-induced migration by 50% (the ED₅₀ concentration).

Enantioselectivity


The (S)- and (R)- enantiomers of those members of the α -aminobicyclolactam series which are chiral can be synthesised to determine whether the biological activity showed enantioselectivity.

The dose-response curves for each of the compounds as inhibitors of MCP-1 induced THP-1 cell migration can be determined using the transwell migration assay.

For the application of the compounds of the present invention as anti-inflammatory agents in vivo it is preferable to use the pure enantiomer of those compounds which are chiral that showed the greater activity in the in vitro bioassay, rather than the racemic mixture of the two enantiomers or the pure enantiomer that was less active in the in vitro bioassay.

Claims

1. Use of a compound of general formula (I) or a pharmaceutically acceptable salt thereof, for the preparation of a medicament intended to treat an inflammatory disorder:



(I)

wherein

y is any integer from 1 to 8;

z is any integer from 1 to 8; with the proviso that y and z cannot both equal 1;

X is $-\text{CO}-(\text{Y})_k-(\text{R}^1)_n$ or $\text{SO}_2-(\text{Y})_k-(\text{R}^1)_n$;

k is 0 or 1

Y is a cycloalkyl or polycyloalkyl group (such as an adamantyl, adamantanemethyl, bicyclooctyl, cyclohexyl, cyclopropyl group);

or Y is a cycloalkenyl or polycycloalkenyl group;


each R^1 is independently selected from hydrogen or an alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl or alkylamino radical of 1 to 20 carbon atoms (for example of 5 to 20 carbon atoms, of 8 to 20 carbon atoms, of 9 to 20 carbon atoms, of 10 to 18 carbon atoms, of 12 to 18 carbon atoms, of 13 to 18 carbon atoms, of 14 to 18 carbon atoms, of 13 to 17 carbon atoms);

or each R^1 is independently selected from fluoro, chloro, bromo, iodo, hydroxy, oxyalkyl, amino, aminoalkyl or aminodialkyl radical; and

n is any integer from 1 to m, where m is the maximum number of substitutions permissible on the cyclo-group Y; or

alternatively R¹ is selected from a peptido radical having from 1 to 4 peptidic moieties linked together by peptide bonds.

2. A pharmaceutical composition comprising, as active ingredient, a compound of formula (I) or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient and/or carrier:



(I)

wherein

y is any integer from 1 to 8;

z is any integer from 1 to 8; with the proviso that y and z cannot both equal 1;

X is $-\text{CO}-(\text{Y})_k-(\text{R}^1)_n$ or $\text{SO}_2-(\text{Y})_k-(\text{R}^1)_n$;

k is 0 or 1

Y is a cycloalkyl or polycyloalkyl group (such as an adamantlyl, adamantanemethyl, bicyclooctyl, cyclohexyl, cyclopropyl group);

or Y is a cycloalkenyl or polycycloalkenyl group;


each R¹ is independently selected from hydrogen or an alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl or alkylamino radical of 1 to 20 carbon atoms (for example of 5 to 20 carbon atoms, of 8 to 20 carbon atoms, of 9 to 20 carbon atoms, of 10 to 18 carbon atoms, of 12 to 18 carbon atoms, of 13 to 18 carbon atoms, of 14 to 18 carbon atoms, of 13 to 17 carbon atoms);

or each R¹ is independently selected from fluoro, chloro, bromo, iodo, hydroxy, oxyalkyl, amino, aminoalkyl or aminodialkyl radical; and

n is any integer from 1 to m, where m is the maximum number of substitutions permissible on the cyclo-group Y; or

alternatively R¹ is selected from a peptido radical having from 1 to 4 peptidic moieties linked together by peptide bonds (for example a peptido radical of 1 to 4 amino acid residues).

3. A compound of general formula (I):



(I)

wherein

y is any integer from 1 to 8;

z is any integer from 1 to 8; with the proviso that y and z cannot both equal 1;

X is -CO-(Y)_k-(R¹)_n or SO₂-(Y)_k-(R¹)_n;

k is 0 or 1

Y is a cycloalkyl or polycyloalkyl group (such as an adamantlyl, adamantanemethyl, bicyclooctyl, cyclohexyl, cyclopropyl group);

or Y is a cycloalkenyl or polycycloalkenyl group;

each R¹ is independently selected from hydrogen or an alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl or alkylamino radical of 1 to 20 carbon atoms (for example of 5 to 20 carbon atoms, of 8 to 20 carbon atoms, of 9 to 20 carbon atoms, of 10 to 18 carbon atoms, of 12 to 18 carbon atoms, of 13 to 18 carbon atoms, of 14 to 18 carbon atoms, of 13 to 17 carbon atoms);

or each R¹ is independently selected from fluoro, chloro, bromo, iodo, hydroxy, oxyalkyl, amino, aminoalkyl or aminodialkyl radical; and

n is any integer from 1 to m, where m is the maximum number of substitutions permissible on the cyclo-group Y; or

alternatively R¹ is selected from a peptido radical having from 1 to 4 peptidic moieties linked together by peptide bonds.

4. Compounds, compositions and uses of the compounds of general formula (I) or their pharmaceutically acceptable salts, according to any preceding claim, wherein the alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl or alkylamino part of the R¹ radical is either linear or is branched but contains a linear chain of at least 8 or at least 10 carbon atoms.
5. Compounds, compositions and uses according to any preceding claim wherein the R¹ radical has a "key"-carbon (defined as the 2-position of a carbonyl-containing radical or the 1-position of a sulfonyl containing radical) which is di-substituted with the same or different groups selected from: alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, alkynyl and alkylamino radicals.
6. Compounds, compositions and uses according to claim 5 wherein the "key"-carbon is chiral.
7. Compounds, compositions and uses according to claim 6 wherein the "key"-carbon has sp³ hybridised bonds.
8. Compounds, compositions and uses according to claim 6 wherein the "key"-carbon has essentially tetrahedral bond angles.
9. A use according to claim 1, or a pharmaceutical composition according to claim 2, or a compound according to claim 3, wherein general formula (I) is modified such that the C3-C7 alkyl bridge -(CH₂)_y- is replaced by a bridging group independently selectable from the group consisting of alkenyl, haloalkyl, alkylamino and alkylhydroxy moieties having a carbon chain length of from 1 to 8.

10. Compounds, compositions and uses of the compounds of general formula (I), or their pharmaceutically acceptable salts, according to any preceding claim, wherein the ring or rings of Y constrain the bond angles at the "key"-carbon to be essentially tetrahedral (i.e. sp³ hybrid bonds).

11. A use according to claim 1 or a pharmaceutical composition according to claim 2, or a compound according to claim 5, wherein the compound is selected from the group consisting of:

4-(Adamantane-1-carbonylamino)-3-oxo-2-aza-bicyclo[2.2.2]octane

5-(Adamantane-1-carbonylamino)-10-oxo-9-aza-bicyclo[3.3.2]decane

4-(2',2'-dimethyldodecanoylamino)-3-oxo-2-aza-bicyclo[2.2.2]octane

5-(2',2'-dimethyldodecanoylamino)-10-oxo-9-aza-bicyclo[3.3.2]decane

and sulfonyl analogues thereof;

and pharmaceutically acceptable salts thereof.

12. A use, composition or compound according to any preceding claim wherein y and z are the same integer, whereby the α -aminobicyclolactam is non-chiral.

13. A use, composition or compound according to any of claims 1 to 11 wherein y and z are not the same integer, whereby the α -aminobicyclolactam ring is chiral.

14. A use, composition or compound according to claim 13 wherein z is 3 and y is 1 or 2 or 4-8, whereby the compound contains a lactam ring which is seven membered.

15. A use, composition or compound according to claim 13 wherein z is 2 and y is 1 or 3-8, whereby the compound contains lactam ring which is 6 membered.

16. Use of a compound of formula (I) according to claim 1 or 9 wherein the inflammatory disorder is selected from the group consisting of autoimmune diseases, vascular disorders, viral infection or replication, asthma, osteoporosis (low bone mineral density), tumor growth, rheumatoid arthritis, organ transplant rejection and/or delayed graft or organ function, a disorder characterised by an elevated TNF- α level, psoriasis, skin wounds, disorders caused by intracellular parasites, allergies, Alzheimer's disease, antigen induced

25

recall response, immune response suppression, multiple sclerosis, ALS, fibrosis, and formation of adhesions.

17. A method of treatment, amelioration or prophylaxis of the symptoms of an inflammatory disease (including an adverse inflammatory reaction to any agent) by the administration to a patient of an anti-inflammatory amount of a compound, composition or medicament as claimed in any of claims 1 to 15.




Figure 1