




(86) **Date de dépôt PCT/PCT Filing Date:** 2010/04/01
(87) **Date publication PCT/PCT Publication Date:** 2010/10/14
(45) **Date de délivrance/Issue Date:** 2017/02/28
(85) **Entrée phase nationale/National Entry:** 2011/08/19
(86) **N° demande PCT/PCT Application No.:** EP 2010/054396
(87) **N° publication PCT/PCT Publication No.:** 2010/115836
(30) **Priorités/Priorities:** 2009/04/08 (EP09157653.8);
2009/11/03 (EP09174917.6)

(51) **Cl.Int./Int.Cl.** **C07D 401/04** (2006.01),
A61K 31/4523 (2006.01), **A61K 31/454** (2006.01),
C07D 401/14 (2006.01), **C07D 413/04** (2006.01),
C07D 413/14 (2006.01), **C07D 417/14** (2006.01),
A61P 31/00 (2006.01), **A61P 35/00** (2006.01),
A61P 37/00 (2006.01)

(72) **Inventeurs/Inventors:**
GRUNDL, MARC, DE;
DOLLINGER, HORST, DE;
GIOVANNINI, RICCARDO, DE;
HOENKE, CHRISTOPH, DE;
HOFFMANN, MATTHIAS, DE;
...

(73) **Propriétaire/Owner:**

(54) **Titre : PIPERIDINES SUBSTITUEES COMME ANTAGONISTES DE CCR3**
(54) **Title: SUBSTITUTED PIPERIDINES AS CCR3 ANTAGONISTS**



(1)

(57) **Abrégé/Abstract:**

Object of the present invention are novel substituted compounds of the formula 1, (see formula I) wherein A, R¹, R², R³ and R⁴ are defined as in the description. Another object of the present invention is to provide antagonists of CCR3, more particularly to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt thereof.

(72) **Inventeurs(suite)/Inventors(continued):** KRIEGL, JAN, DE; MARTYRES, DOMNIC, DE; RAST, GEORG, DE; SEITHER, PETER, DE


(73) **Propriétaires(suite)/Owners(continued):**BOEHRINGER INGELHEIM INTERNATIONAL GMBH, DE

(74) **Agent:** FETHERSTONHAUGH & CO.

25771-1956

ABSTRACT

Object of the present invention are novel substituted compounds of the formula 1,




wherein A, R¹, R², R³ and R⁴ are defined as in the description. Another object of the present
5 invention is to provide antagonists of CCR3, more particularly to provide pharmaceutical
compositions comprising a pharmaceutically acceptable carrier and at least one of the
compounds of the present invention or a pharmaceutically acceptable salt thereof.

25771-1956

SUBSTITUTED PIPERIDINES AS CCR3 ANTAGONISTS

Object of the present invention are novel substituted compounds of the formula 1,



wherein R^1 , R^2 , R^3 and R^4 are defined as in the description. Another object of the present invention is to provide antagonists of CCR3, more particularly to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and
 10 at least one of the compounds of the present invention or a pharmaceutically acceptable salt thereof.

BACKGROUND INFORMATION

15 Chemokines are chemotactic cytokines, of molecular weight 6-15 kDa, that are released by a wide variety of cells to attract and activate, among other cell types, macrophages, T and B lymphocytes, eosinophils, basophils and neutrophils (reviewed in Luster, New Eng. J Med., 338, 436-445 (1998); Rollins, Blood, 90, 909-928 (1997); Lloyd, Curr Opin Pharmacol., 3, 443-448 (2003); Murray, Current Drug Targets., 7, 579-588 (2006); Smit, Eur J Pharmacol., 533,277-88 (2006)

There are two major classes of chemokines, CXC and CC, depending on whether the first two cysteines in the amino acid sequence are separated by a single amino acid (CXC) or are adjacent (CC). The CXC chemokines, such as interleukin-8 (IL-8), neutrophil-activating protein-2 (NAP2) and melanoma growth stimulatory activity protein (MGSA) are chemotactic primarily for neutrophils and T lymphocytes, whereas the CC chemokines, such as RANTES, MIP-1a, MIP-1, the monocyte chemotactic proteins (MCP-1, MCP-2, MCP-3, MCP-4, and MCP-5) and the eotaxins (-1,-2, and-3) are chemotactic for, among other cell types, macrophages, T lymphocytes, eosinophils, mast cells, dendritic cells, and basophils. Also in existence are the chemokines lymphotactin-1, lymphotactin-2 (both C chemokines), and fractalkine (a CXXXC chemokine) that do not fall into either of the major chemokine subfamilies.

35 The chemokines bind to specific cell-surface receptors belonging to the family of G-protein-coupled seven transmembrane-domain proteins (reviewed in Horuk, Trends Pharm. Sci., 15, 159-165 (1994); Murphy, Pharmacol Rev., 54 (2):227-229 (2002); Allen, Annu. Rev. Immunol., 25, 787-820 (2007)) which are termed "chemokine receptors." On binding their cognate ligands, chemokine receptors transduce an intracellular signal through the associated trimeric G proteins, resulting in, among other responses, a rapid increase in intracellular calcium concentration, activation of G-proteins, changes in cell shape, increased expression of cellular adhesion molecules, degranulation, promotion of cell migration, survival and proliferation. There are at least eleven human chemokine receptors that bind or respond to CC chemokines with the following characteristic patterns: CCR-1 (or "CCR-1" or "CC-CCR-1") [MIP-1a, MCP-3, MCP-4, RANTES] (Ben-Baruch, et al., Cell, 72, 415-425 (1993), Luster, New Eng. J. Med., 338, 436-445 (1998)); CCR-2A and CCR-2B (or "CCR-2A" or "CCR-2B" or "CC-CCR-2A" or "CC-CCR-2B") [MCP-1, MCP-2, MCP-3, MCP-4, MCP-5, RANTES, MIP-1a, MIP-1b, MIP-1c, MIP-1d, MIP-1e, MIP-1f, MIP-1g, MIP-1h, MIP-1i, MIP-1j, MIP-1k, MIP-1l, MIP-1m, MIP-1n, MIP-1o, MIP-1p, MIP-1q, MIP-1r, MIP-1s, MIP-1t, MIP-1u, MIP-1v, MIP-1w, MIP-1x, MIP-1y, MIP-1z, MIP-1aa, MIP-1ab, MIP-1ac, MIP-1ad, MIP-1ae, MIP-1af, MIP-1ag, MIP-1ah, MIP-1ai, MIP-1aj, MIP-1ak, MIP-1al, MIP-1am, MIP-1an, MIP-1ao, MIP-1ap, MIP-1aq, MIP-1ar, MIP-1as, MIP-1at, MIP-1au, MIP-1av, MIP-1aw, MIP-1ax, MIP-1ay, MIP-1az, MIP-1ba, MIP-1bb, MIP-1bc, MIP-1bd, MIP-1be, MIP-1bf, MIP-1bg, MIP-1bh, MIP-1bi, MIP-1bj, MIP-1bk, MIP-1bl, MIP-1bm, MIP-1bn, MIP-1bo, MIP-1bp, MIP-1bq, MIP-1br, MIP-1bs, MIP-1bt, MIP-1bu, MIP-1bv, MIP-1bw, MIP-1bx, MIP-1by, MIP-1bz, MIP-1ca, MIP-1cb, MIP-1cc, MIP-1cd, MIP-1ce, MIP-1cf, MIP-1cg, MIP-1ch, MIP-1ci, MIP-1cj, MIP-1ck, MIP-1cl, MIP-1cm, MIP-1cn, MIP-1co, MIP-1cp, MIP-1cq, MIP-1cr, MIP-1cs, MIP-1ct, MIP-1cu, MIP-1cv, MIP-1cw, MIP-1cx, MIP-1cy, MIP-1cz, MIP-1da, MIP-1db, MIP-1dc, MIP-1dd, MIP-1de, MIP-1df, MIP-1dg, MIP-1dh, MIP-1di, MIP-1dj, MIP-1dk, MIP-1dl, MIP-1dm, MIP-1dn, MIP-1do, MIP-1dp, MIP-1dq, MIP-1dr, MIP-1ds, MIP-1dt, MIP-1du, MIP-1dv, MIP-1dw, MIP-1dx, MIP-1dy, MIP-1dz, MIP-1ea, MIP-1eb, MIP-1ec, MIP-1ed, MIP-1ee, MIP-1ef, MIP-1eg, MIP-1eh, MIP-1ei, MIP-1ej, MIP-1ek, MIP-1el, MIP-1em, MIP-1en, MIP-1eo, MIP-1ep, MIP-1eq, MIP-1er, MIP-1es, MIP-1et, MIP-1eu, MIP-1ev, MIP-1ew, MIP-1ex, MIP-1ey, MIP-1ez, MIP-1fa, MIP-1fb, MIP-1fc, MIP-1fd, MIP-1fe, MIP-1ff, MIP-1fg, MIP-1fh, MIP-1fi, MIP-1fj, MIP-1fk, MIP-1fl, MIP-1fm, MIP-1fn, MIP-1fo, MIP-1fp, MIP-1fq, MIP-1fr, MIP-1fs, MIP-1ft, MIP-1fu, MIP-1fv, MIP-1fw, MIP-1fx, MIP-1fy, MIP-1fz, MIP-1ga, MIP-1gb, MIP-1gc, MIP-1gd, MIP-1ge, MIP-1gf, MIP-1gg, MIP-1gh, MIP-1gi, MIP-1gj, MIP-1gk, MIP-1gl, MIP-1gm, MIP-1gn, MIP-1go, MIP-1gp, MIP-1gq, MIP-1gr, MIP-1gs, MIP-1gt, MIP-1gu, MIP-1gv, MIP-1gw, MIP-1gx, MIP-1gy, MIP-1gz, MIP-1ha, MIP-1hb, MIP-1hc, MIP-1hd, MIP-1he, MIP-1hf, MIP-1hg, MIP-1hh, MIP-1hi, MIP-1hj, MIP-1hk, MIP-1hl, MIP-1hm, MIP-1hn, MIP-1ho, MIP-1hp, MIP-1hq, MIP-1hr, MIP-1hs, MIP-1ht, MIP-1hu, MIP-1hv, MIP-1hw, MIP-1hx, MIP-1hy, MIP-1hz, MIP-1ia, MIP-1ib, MIP-1ic, MIP-1id, MIP-1ie, MIP-1if, MIP-1ig, MIP-1ih, MIP-1ii, MIP-1ij, MIP-1ik, MIP-1il, MIP-1im, MIP-1in, MIP-1io, MIP-1ip, MIP-1iq, MIP-1ir, MIP-1is, MIP-1it, MIP-1iu, MIP-1iv, MIP-1iw, MIP-1ix, MIP-1iy, MIP-1iz, MIP-1ja, MIP-1jb, MIP-1jc, MIP-1jd, MIP-1je, MIP-1jf, MIP-1jg, MIP-1jh, MIP-1ji, MIP-1jj, MIP-1jk, MIP-1jl, MIP-1jm, MIP-1jn, MIP-1jo, MIP-1jp, MIP-1jq, MIP-1jr, MIP-1js, MIP-1jt, MIP-1ju, MIP-1jv, MIP-1jw, MIP-1jx, MIP-1jy, MIP-1jz, MIP-1ka, MIP-1kb, MIP-1kc, MIP-1kd, MIP-1ke, MIP-1kf, MIP-1kg, MIP-1kh, MIP-1ki, MIP-1kj, MIP-1kk, MIP-1kl, MIP-1km, MIP-1kn, MIP-1ko, MIP-1kp, MIP-1kq, MIP-1kr, MIP-1ks, MIP-1kt, MIP-1ku, MIP-1kv, MIP-1kw, MIP-1kx, MIP-1ky, MIP-1kz, MIP-1la, MIP-1lb, MIP-1lc, MIP-1ld, MIP-1le, MIP-1lf, MIP-1lg, MIP-1lh, MIP-1li, MIP-1lj, MIP-1lk, MIP-1ll, MIP-1lm, MIP-1ln, MIP-1lo, MIP-1lp, MIP-1lq, MIP-1lr, MIP-1ls, MIP-1lt, MIP-1lu, MIP-1lv, MIP-1lw, MIP-1lx, MIP-1ly, MIP-1lz, MIP-1ma, MIP-1mb, MIP-1mc, MIP-1md, MIP-1me, MIP-1mf, MIP-1mg, MIP-1mh, MIP-1mi, MIP-1mj, MIP-1mk, MIP-1ml, MIP-1mm, MIP-1mn, MIP-1mo, MIP-1mp, MIP-1mq, MIP-1mr, MIP-1ms, MIP-1mt, MIP-1mu, MIP-1mv, MIP-1mw, MIP-1mx, MIP-1my, MIP-1mz, MIP-1na, MIP-1nb, MIP-1nc, MIP-1nd, MIP-1ne, MIP-1nf, MIP-1ng, MIP-1nh, MIP-1ni, MIP-1nj, MIP-1nk, MIP-1nl, MIP-1nm, MIP-1nn, MIP-1no, MIP-1np, MIP-1nq, MIP-1nr, MIP-1ns, MIP-1nt, MIP-1nu, MIP-1nv, MIP-1nw, MIP-1nx, MIP-1ny, MIP-1nz, MIP-1oa, MIP-1ob, MIP-1oc, MIP-1od, MIP-1oe, MIP-1of, MIP-1og, MIP-1oh, MIP-1oi, MIP-1oj, MIP-1ok, MIP-1ol, MIP-1om, MIP-1on, MIP-1oo, MIP-1op, MIP-1oq, MIP-1or, MIP-1os, MIP-1ot, MIP-1ou, MIP-1ov, MIP-1ow, MIP-1ox, MIP-1oy, MIP-1oz, MIP-1pa, MIP-1pb, MIP-1pc, MIP-1pd, MIP-1pe, MIP-1pf, MIP-1pg, MIP-1ph, MIP-1pi, MIP-1pj, MIP-1pk, MIP-1pl, MIP-1pm, MIP-1pn, MIP-1po, MIP-1pp, MIP-1pq, MIP-1pr, MIP-1ps, MIP-1pt, MIP-1pu, MIP-1pv, MIP-1pw, MIP-1px, MIP-1py, MIP-1pz, MIP-1qa, MIP-1qb, MIP-1qc, MIP-1qd, MIP-1qe, MIP-1qf, MIP-1qg, MIP-1qh, MIP-1qi, MIP-1qj, MIP-1qk, MIP-1ql, MIP-1qm, MIP-1qn, MIP-1qo, MIP-1qp, MIP-1qq, MIP-1qr, MIP-1qs, MIP-1qt, MIP-1qu, MIP-1qv, MIP-1qw, MIP-1qx, MIP-1qy, MIP-1qz, MIP-1ra, MIP-1rb, MIP-1rc, MIP-1rd, MIP-1re, MIP-1rf, MIP-1rg, MIP-1rh, MIP-1ri, MIP-1rj, MIP-1rk, MIP-1rl, MIP-1rm, MIP-1rn, MIP-1ro, MIP-1rp, MIP-1rq, MIP-1rr, MIP-1rs, MIP-1rt, MIP-1ru, MIP-1rv, MIP-1rw, MIP-1rx, MIP-1ry, MIP-1rz, MIP-1sa, MIP-1sb, MIP-1sc, MIP-1sd, MIP-1se, MIP-1sf, MIP-1sg, MIP-1sh, MIP-1si, MIP-1sj, MIP-1sk, MIP-1sl, MIP-1sm, MIP-1sn, MIP-1so, MIP-1sp, MIP-1sq, MIP-1sr, MIP-1ss, MIP-1st, MIP-1su, MIP-1sv, MIP-1sw, MIP-1sx, MIP-1sy, MIP-1sz, MIP-1ta, MIP-1tb, MIP-1tc, MIP-1td, MIP-1te, MIP-1tf, MIP-1tg, MIP-1th, MIP-1ti, MIP-1tj, MIP-1tk, MIP-1tl, MIP-1tm, MIP-1tn, MIP-1to, MIP-1tp, MIP-1tq, MIP-1tr, MIP-1ts, MIP-1tt, MIP-1tu, MIP-1tv, MIP-1tw, MIP-1tx, MIP-1ty, MIP-1tz, MIP-1ua, MIP-1ub, MIP-1uc, MIP-1ud, MIP-1ue, MIP-1uf, MIP-1ug, MIP-1uh, MIP-1ui, MIP-1uj, MIP-1uk, MIP-1ul, MIP-1um, MIP-1un, MIP-1uo, MIP-1up, MIP-1uq, MIP-1ur, MIP-1us, MIP-1ut, MIP-1uu, MIP-1uv, MIP-1uw, MIP-1ux, MIP-1uy, MIP-1uz, MIP-1va, MIP-1vb, MIP-1vc, MIP-1vd, MIP-1ve, MIP-1vf, MIP-1vg, MIP-1vh, MIP-1vi, MIP-1vj, MIP-1vk, MIP-1vl, MIP-1vm, MIP-1vn, MIP-1vo, MIP-1vp, MIP-1vq, MIP-1vr, MIP-1vs, MIP-1vt, MIP-1vu, MIP-1vv, MIP-1vw, MIP-1vx, MIP-1vy, MIP-1vz, MIP-1wa, MIP-1wb, MIP-1wc, MIP-1wd, MIP-1we, MIP-1wf, MIP-1wg, MIP-1wh, MIP-1wi, MIP-1wj, MIP-1wk, MIP-1wl, MIP-1wm, MIP-1wn, MIP-1wo, MIP-1wp, MIP-1wq, MIP-1wr, MIP-1ws, MIP-1wt, MIP-1wu, MIP-1wv, MIP-1ww, MIP-1wx, MIP-1wy, MIP-1wz, MIP-1xa, MIP-1xb, MIP-1xc, MIP-1xd, MIP-1xe, MIP-1xf, MIP-1xg, MIP-1xh, MIP-1xi, MIP-1xj, MIP-1xk, MIP-1xl, MIP-1xm, MIP-1xn, MIP-1xo, MIP-1xp, MIP-1xq, MIP-1xr, MIP-1xs, MIP-1xt, MIP-1xu, MIP-1xv, MIP-1xw, MIP-1xx, MIP-1xy, MIP-1xz, MIP-1ya, MIP-1yb, MIP-1yc, MIP-1yd, MIP-1ye, MIP-1yf, MIP-1yg, MIP-1yh, MIP-1yi, MIP-1yj, MIP-1yk, MIP-1yl, MIP-1ym, MIP-1yn, MIP-1yo, MIP-1yp, MIP-1yq, MIP-1yr, MIP-1ys, MIP-1yt, MIP-1yu, MIP-1yv, MIP-1yw, MIP-1yx, MIP-1yy, MIP-1yz, MIP-1za, MIP-1zb, MIP-1zc, MIP-1zd, MIP-1ze, MIP-1zf, MIP-1zg, MIP-1zh, MIP-1zi, MIP-1zj, MIP-1zk, MIP-1zl, MIP-1zm, MIP-1zn, MIP-1zo, MIP-1zp, MIP-1zq, MIP-1zr, MIP-1zs, MIP-1zt, MIP-1zu, MIP-1zv, MIP-1zw, MIP-1zx, MIP-1zy, MIP-1zz, MIP-1aa, MIP-1ab, MIP-1ac, MIP-1ad, MIP-1ae, MIP-1af, MIP-1ag, MIP-1ah, MIP-1ai, MIP-1aj, MIP-1ak, MIP-1al, MIP-1am, MIP-1an, MIP-1ao, MIP-1ap, MIP-1aq, MIP-1ar, MIP-1as, MIP-1at, MIP-1au, MIP-1av, MIP-1aw, MIP-1ax, MIP-1ay, MIP-1az, MIP-1ba, MIP-1bb, MIP-1bc, MIP-1bd, MIP-1be, MIP-1bf, MIP-1bg, MIP-1bh, MIP-1bi, MIP-1bj, MIP-1bk, MIP-1bl, MIP-1bm, MIP-1bn, MIP-1bo, MIP-1bp, MIP-1bq, MIP-1br, MIP-1bs, MIP-1bt, MIP-1bu, MIP-1bv, MIP-1bw, MIP-1bx, MIP-1by, MIP-1bz, MIP-1ca, MIP-1cb, MIP-1cc, MIP-1cd, MIP-1ce, MIP-1cf, MIP-1cg, MIP-1ch, MIP-1ci, MIP-1cj, MIP-1ck, MIP-1cl, MIP-1cm, MIP-1cn, MIP-1co, MIP-1cp, MIP-1cq, MIP-1cr, MIP-1cs, MIP-1ct, MIP-1cu, MIP-1cv, MIP-1cw, MIP-1cx, MIP-1cy, MIP-1cz, MIP-1da, MIP-1db, MIP-1dc, MIP-1dd, MIP-1de, MIP-1df, MIP-1dg, MIP-1dh, MIP-1di, MIP-1dj, MIP-1dk, MIP-1dl, MIP-1dm, MIP-1dn, MIP-1do, MIP-1dp, MIP-1dq, MIP-1dr, MIP-1ds, MIP-1dt, MIP-1du, MIP-1dv, MIP-1dw, MIP-1dx, MIP-1dy, MIP-1dz, MIP-1ea, MIP-1eb, MIP-1ec, MIP-1ed, MIP-1ee, MIP-1ef, MIP-1eg, MIP-1eh, MIP-1ei, MIP-1ej, MIP-1ek, MIP-1el, MIP-1em, MIP-1en, MIP-1eo, MIP-1ep, MIP-1eq, MIP-1er, MIP-1es, MIP-1et, MIP-1eu, MIP-1ev, MIP-1ew, MIP-1ex, MIP-1ey, MIP-1ez, MIP-1fa, MIP-1fb, MIP-1fc, MIP-1fd, MIP-1fe, MIP-1ff, MIP-1fg, MIP-1fh, MIP-1fi, MIP-1fj, MIP-1fk, MIP-1fl, MIP-1fm, MIP-1fn, MIP-1fo, MIP-1fp, MIP-1fq, MIP-1fr, MIP-1fs, MIP-1ft, MIP-1fu, MIP-1fv, MIP-1fw, MIP-1fx, MIP-1fy, MIP-1fz, MIP-1ga, MIP-1gb, MIP-1gc, MIP-1gd, MIP-1ge, MIP-1gf, MIP-1gg, MIP-1gh, MIP-1gi, MIP-1gj, MIP-1gk, MIP-1gl, MIP-1gm, MIP-1gn, MIP-1go, MIP-1gp, MIP-1gq, MIP-1gr, MIP-1gs, MIP-1gt, MIP-1gu, MIP-1gv, MIP-1gw, MIP-1gx, MIP-1gy, MIP-1gz, MIP-1ha, MIP-1hb, MIP-1hc, MIP-1hd, MIP-1he, MIP-1hf, MIP-1hg, MIP-1hh, MIP-1hi, MIP-1hj, MIP-1hk, MIP-1hl, MIP-1hm, MIP-1hn, MIP-1ho, MIP-1hp, MIP-1hq, MIP-1hr, MIP-1hs, MIP-1ht, MIP-1hu, MIP-1hv, MIP-1hw, MIP-1hx, MIP-1hy, MIP-1hz, MIP-1ia, MIP-1ib, MIP-1ic, MIP-1id, MIP-1ie, MIP-1if, MIP-1ig, MIP-1ih, MIP-1ii, MIP-1ij, MIP-1ik, MIP-1il, MIP-1im, MIP-1in, MIP-1io, MIP-1ip, MIP-1iq, MIP-1ir, MIP-1is, MIP-1it, MIP-1iu, MIP-1iv, MIP-1iw, MIP-1ix, MIP-1iy, MIP-1iz, MIP-1ja, MIP-1jb, MIP-1jc, MIP-1jd, MIP-1je, MIP-1jf, MIP-1jg, MIP-1jh, MIP-1ji, MIP-1jj, MIP-1jk, MIP-1jl, MIP-1jm, MIP-1jn, MIP-1jo, MIP-1jp, MIP-1jq, MIP-1jr, MIP-1js, MIP-1jt, MIP-1ju, MIP-1jv, MIP-1jw, MIP-1jx, MIP-1jy, MIP-1jz, MIP-1ka, MIP-1kb, MIP-1kc, MIP-1kd, MIP-1ke, MIP-1kf, MIP-1kg, MIP-1kh, MIP-1ki, MIP-1kj, MIP-1kk, MIP-1kl, MIP-1km, MIP-1kn, MIP-1ko, MIP-1kp, MIP-1kq, MIP-1kr, MIP-1ks, MIP-1kt, MIP-1ku, MIP-1kv, MIP-1kw, MIP-1kx, MIP-1ky, MIP-1kz, MIP-1la, MIP-1lb, MIP-1lc, MIP-1ld, MIP-1le, MIP-1lf, MIP-1lg, MIP-1lh, MIP-1li, MIP-1lj, MIP-1lk, MIP-1ll, MIP-1lm, MIP-1ln, MIP-1lo, MIP-1lp, MIP-1lq, MIP-1lr, MIP-1ls, MIP-1lt, MIP-1lu, MIP-1lv, MIP-1lw, MIP-1lx, MIP-1ly, MIP-1lz, MIP-1ma, MIP-1mb, MIP-1mc, MIP-1md, MIP-1me, MIP-1mf, MIP-1mg, MIP-1mh, MIP-1mi, MIP-1mj, MIP-1mk, MIP-1ml, MIP-1mm, MIP-1mn, MIP-1mo, MIP-1mp, MIP-1mq, MIP-1mr, MIP-1ms, MIP-1mt, MIP-1mu, MIP-1mv, MIP-1mw, MIP-1mx, MIP-1my, MIP-1mz, MIP-1na, MIP-1nb, MIP-1nc, MIP-1nd, MIP-1ne, MIP-1nf, MIP-1ng, MIP-1nh, MIP-1ni, MIP-1nj, MIP-1nk, MIP-1nl, MIP-1nm, MIP-1nn, MIP-1no, MIP-1np, MIP-1nq, MIP-1nr, MIP-1ns, MIP-1nt, MIP-1nu, MIP-1nv, MIP-1nw, MIP-1nx, MIP-1ny, MIP-1nz, MIP-1oa, MIP-1ob, MIP-1oc, MIP-1od, MIP-1oe, MIP-1of, MIP-1og, MIP-1oh, MIP-1oi, MIP-1oj, MIP-1ok, MIP-1ol, MIP-1om, MIP-1on, MIP-1oo, MIP-1op, MIP-1oq, MIP-1or, MIP-1os, MIP-1ot, MIP-1ou, MIP-1ov, MIP-1ow, MIP-1ox, MIP-1oy, MIP-1oz, MIP-1pa, MIP-1pb, MIP-1pc, MIP-1pd, MIP-1pe, MIP-1pf, MIP-1pg, MIP-1ph, MIP-1pi, MIP-1pj, MIP-1pk, MIP-1pl, MIP-1pm, MIP-1pn, MIP-1po, MIP-1pp, MIP-1pq, MIP-1pr, MIP-1ps, MIP-1pt, MIP-1pu, MIP-1pv, MIP-1pw, MIP-1px, MIP-1py, MIP-1pz, MIP-1qa, MIP-1qb, MIP-1qc, MIP-1qd, MIP-1qe, MIP-1qf, MIP-1qg, MIP-1qh, MIP-1qi, MIP-1qj, MIP-1qk, MIP-1ql, MIP-1qm, MIP-1qn, MIP-1qo, MIP-1qp, MIP-1qq, MIP-1qr, MIP-1qs, MIP-1qt, MIP-1qu, MIP-1qv, MIP-1qw, MIP-1qx, MIP-1qy, MIP-1qz, MIP-1ra, MIP-1rb, MIP-1rc, MIP-1rd, MIP-1re, MIP-1rf, MIP-1rg, MIP-1rh, MIP-1ri, MIP-1rj, MIP-1rk, MIP-1rl, MIP-1rm, MIP-1rn, MIP-1ro, MIP-1rp, MIP-1rq, MIP-1rr, MIP-1rs, MIP-1rt, MIP-1ru, MIP-1rv, MIP-1rw, MIP-1rx, MIP-1ry, MIP-1rz, MIP-1sa, MIP-1sb, MIP-1sc, MIP-1sd, MIP-1se, MIP-1sf, MIP-1sg, MIP-1sh, MIP-1si, MIP-1sj, MIP-1sk, MIP-1sl, MIP-1sm, MIP-1sn, MIP-1so, MIP-1sp, MIP-1sq, MIP-1sr, MIP-1ss, MIP-1st, MIP-1su, MIP-1sv, MIP-1sw, MIP-1sx, MIP-1sy, MIP-1sz, MIP-1ta, MIP-1tb, MIP-1tc, MIP-1td, MIP-1te, MIP-1tf, MIP-1tg, MIP-1th, MIP-1ti, MIP-1tj, MIP-1tk, MIP-1tl, MIP-1tm, MIP-1tn, MIP-1to, MIP-1tp, MIP-1tq, MIP-1tr, MIP-1ts, MIP-1tt, MIP-1tu, MIP-1tv, MIP-1tw, MIP-1tx, MIP-1ty, MIP-1tz, MIP-1ua, MIP-1ub, MIP-1uc, MIP-1ud, MIP-1ue, MIP-1uf, MIP-1ug, MIP-1uh, MIP-1ui, MIP-1uj, MIP-1uk, MIP-1ul, MIP-1um, MIP-1un, MIP-1uo, MIP-1up, MIP-1uq, MIP-1ur, MIP-1us, MIP-1ut, MIP-1uu, MIP-1uv, MIP-1uw, MIP-1ux, MIP-1uy, MIP-1uz, MIP-1va, MIP-1vb, MIP-1vc, MIP-1vd, MIP-1ve, MIP-1vf, MIP-1vg, MIP-1vh, MIP-1vi, MIP-1vj, MIP-1vk, MIP-1vl, MIP-1vm, MIP-1vn, MIP-1vo, MIP-1vp, MIP-1vq, MIP-1vr, MIP-1vs, MIP-1vt, MIP-1vu, MIP-1vv, MIP-1vw, MIP-1vx, MIP-1vy, MIP-1vz, MIP-1wa, MIP-1wb, MIP-1wc, MIP-1wd, MIP-1we, MIP-1wf, MIP-1wg, MIP-1wh, MIP-1wi, MIP-1wj, MIP-1wk, MIP-1wl, MIP-1wm, MIP-1wn, MIP-1wo, MIP-1wp, MIP-1wq, MIP-1wr, MIP-1ws, MIP-1wt, MIP-1wu, MIP-1wv, MIP-1ww, MIP-1wx, MIP-1wy, MIP-1wz, MIP-1xa, MIP-1xb, MIP-1xc, MIP-1xd, MIP-1xe, MIP-1xf, MIP-1xg, MIP-1xh, MIP-1xi, MIP-1xj, MIP-1xk, MIP-1xl, MIP-1xm, MIP-1xn, MIP-1xo, MIP-1xp, MIP-1xq, MIP-1xr, MIP-1xs, MIP-1xt, MIP-1xu, MIP-1xv, MIP-1xw, MIP-1xx, MIP-1xy, MIP-1xz, MIP-1ya, MIP-1yb, MIP-1yc, MIP-1yd, MIP-1ye, MIP-1yf, MIP-1yg, MIP-1yh, MIP-1yi, MIP-1yj, MIP-1yk, MIP-1yl, MIP-1ym, MIP-1yn, MIP-1yo, MIP-1yp, MIP-1yq, MIP-1yr, MIP-1ys, MIP-1yt, MIP-1yu, MIP-1yv, MIP-1yw, MIP-1yx, MIP-1yy, MIP-1yz, MIP-1za, MIP-1zb, MIP-1zc, MIP-1zd, MIP-1ze, MIP-1zf, MIP-1zg, MIP-1zh, MIP-1zi, MIP-1zj, MIP-1zk, MIP-1zl, MIP-1zm, MIP-1zn, MIP-1zo, MIP-1zp, MIP-1zq, MIP-1zr, MIP-1zs, MIP-1zt, MIP-1zu, MIP-1zv, MIP-1zw, MIP-1zx, MIP-1zy, MIP-1zz, MIP-1aa, MIP-1ab, MIP-1ac, MIP-1ad, MIP-1ae, MIP-1af, MIP-1ag, MIP-1ah, MIP-1ai, MIP-1aj, MIP-1ak, MIP-1al, MIP-1am, MIP-1an, MIP-1ao, MIP-1ap, MIP-1aq, MIP-1ar, MIP-1as, MIP-1at, MIP-1au, MIP-1av, MIP-1aw, MIP-1ax, MIP-1ay, MIP-1az, MIP-1ba, MIP-1bb, MIP-1bc, MIP-1bd, MIP-1be, MIP-1bf, MIP-1bg, MIP-1bh, MIP-1bi, MIP-1bj, MIP-1bk, MIP-1bl, MIP-1bm, MIP-1bn, MIP-1bo, MIP-1bp, MIP-1bq, MIP-1br, MIP-1bs, MIP-1bt, MIP-1bu, MIP-1bv, MIP-1bw, MIP-1bx, MIP-1by, MIP-1bz, MIP-1ca, MIP-1cb, MIP-1cc, MIP-1cd, MIP-1ce, MIP-1cf, MIP-1cg, MIP-1ch, MIP-1ci, MIP-1cj, MIP-1ck, MIP-1cl, MIP-1cm, MIP-1cn, MIP-1co, MIP-1cp, MIP-1cq, MIP-1cr, MIP-1cs, MIP-1ct, MIP-1cu, MIP-1cv, MIP-1cw, MIP-1cx, MIP-1cy, MIP-1cz, MIP-1da, MIP-1db, MIP-1dc, MIP-1dd, MIP-1de, MIP-1df, MIP-1dg, MIP-1dh, MIP-1di, MIP-1dj, MIP-1dk, MIP-1dl, MIP-1dm, MIP-1dn, MIP-1do, MIP-1dp, MIP-1dq, MIP-1dr, MIP-1ds, MIP-1dt, MIP-1du, MIP-1dv, MIP-1dw, MIP-1dx, MIP-1dy, MIP-1dz, MIP-1ea, MIP-1eb, MIP-1ec, MIP-1ed, MIP-1ee, MIP-1ef, MIP-1eg, MIP-1eh, MIP-1ei, MIP-1ej, MIP-1ek, MIP-1el, MIP-1em, MIP-1en, MIP-1eo, MIP-1ep, MIP-1eq, MIP-1er, MIP-1es, MIP-1et, MIP-1eu, MIP-1ev, MIP-1ew, MIP-1ex, MIP-1ey, MIP-1ez, MIP-1fa, MIP-1fb, MIP-1fc, MIP-1fd, MIP-1fe, MIP-1ff, MIP-1fg, MIP-1fh, MIP-1fi, MIP-1fj, MIP-1fk, MIP-1fl, MIP-1fm, MIP-1fn, MIP-1fo, MIP-1fp, MIP-1fq, MIP-1fr, MIP-1fs, MIP-1ft, MIP-1fu, MIP-1fv, MIP-1fw, MIP-1fx, MIP-1fy, MIP-1fz, MIP-1ga, MIP-1gb, MIP-1gc, MIP-1gd, MIP-1ge, MIP-1gf, MIP-1gg, MIP-1gh, MIP-1gi, MIP-1gj, MIP-1gk, MIP-1gl, MIP-1gm, MIP-1gn, MIP-1go, MIP-1gp, MIP-1gq, MIP-1gr, MIP-1gs, MIP-1gt, MIP-1gu, MIP-1gv, MIP-1gw, MIP-1gx, MIP-1gy, MIP-1gz, MIP-1ha, MIP-1hb, MIP-1hc, MIP-1hd, MIP-1he, MIP-1hf, MIP-1hg, MIP-1hh, MIP-1hi, MIP-1hj, MIP-1hk, MIP-1hl, MIP-1hm, MIP-1hn, MIP-1ho, MIP-1hp, MIP-1hq, MIP-1hr, MIP-1hs, MIP-1ht, MIP-1hu, MIP-1hv, MIP-1hw, MIP-1hx, MIP-1hy, MIP-1hz, MIP-1ia, MIP-1ib, MIP-1ic, MIP-1id, MIP-1ie, MIP-1if, MIP-1ig, MIP-1ih, MIP-1ii, MIP-1ij, MIP-1ik, MIP-1il, MIP-1im, MIP-1in, MIP-1io, MIP-1ip, MIP-1iq, MIP-1ir, MIP-1is, MIP-1it, MIP-1iu, MIP-1iv, MIP-1iw, MIP-1ix, MIP-1iy, MIP-1iz, MIP-1ja, MIP-1jb, MIP-1jc, MIP-1jd, MIP-1je, MIP-1jf, MIP-1jg, MIP-1jh, MIP-1ji, MIP-1jj, MIP-1jk, MIP-1jl, MIP-1jm, MIP-1jn, MIP-1jo, MIP-1jp, MIP-1jq, MIP-1jr, MIP-1js, MIP-1jt, MIP-1ju, MIP-1jv, MIP-1jw, MIP-1jx, MIP-1jy, MIP-1jz, MIP-1ka, MIP-1kb, MIP-1kc, MIP-1kd, MIP-1ke, MIP-1kf, MIP-1kg, MIP-1kh, MIP-1ki, MIP-1kj, MIP-1kk, MIP-1kl, MIP-1km, MIP-1kn, MIP-1ko, MIP-1kp, MIP-1kq, MIP-1kr, MIP-1ks, MIP-1kt, MIP-1ku, MIP-1kv, MIP-1kw, MIP-1kx, MIP-1ky, MIP-1kz, MIP-1la, MIP-1lb, MIP-1lc, MIP-1ld, MIP-1le, MIP-1lf, MIP-1lg, MIP-1lh, MIP-1li, MIP-1lj, MIP-1lk, MIP-1ll, MIP-1lm, MIP-1ln, MIP-1lo, MIP-1lp, MIP-1lq, MIP-1lr, MIP-1ls, MIP-1lt, MIP-1lu, MIP-1lv, MIP-1lw, MIP-1lx, MIP-1ly, MIP-1lz, MIP-1ma, MIP-1mb, MIP-1mc, MIP-1md, MIP-1me, MIP-1mf, MIP-1mg, MIP-1mh, MIP-1mi, MIP-1mj, MIP-1mk, MIP-1ml, MIP-1mm, MIP-1mn, MIP-1mo, MIP-1mp, MIP-1mq, MIP-1mr, MIP-1ms, MIP-1mt, MIP-1mu, MIP-1mv, MIP-1mw, MIP-1mx, MIP-1my, MIP-1mz, MIP-1na, MIP-1nb, MIP-1nc, MIP-1nd, MIP-1ne, MIP-1nf, MIP-1ng, MIP-1nh, MIP-1ni, MIP-1nj, MIP-1nk, MIP-1nl, MIP-1nm, MIP-1nn, MIP-1no, MIP-1np, MIP-1nq, MIP-1nr, MIP-1ns, MIP-1nt, MIP-1nu, MIP-1nv, MIP-1nw, MIP-1nx, MIP-1ny, MIP-1nz, MIP-1oa, MIP-1ob, MIP-1oc, MIP-1od, MIP-1oe, MIP-1of, MIP-1og, MIP-1oh, MIP-1oi, MIP-1oj, MIP-1ok, MIP-1ol, MIP-1om, MIP-1on, MIP-1oo, MIP-1op, MIP-1oq, MIP-1or, MIP-1os, MIP-1ot, MIP-1ou, MIP-1ov, MIP-1ow, MIP-1ox, MIP-1oy, MIP-1oz, MIP-1pa, MIP-1pb, MIP-1pc, MIP-1pd, MIP-1pe, MIP-1pf, MIP-1pg, MIP-1ph, MIP-1pi, MIP-1pj, MIP-1pk, MIP-1pl, MIP-1pm, MIP-1pn, MIP-1po, MIP-1pp, MIP-1pq, MIP-1pr, MIP-1ps, MIP-1pt, MIP-1pu, MIP-1pv, MIP-1pw, MIP-1px, MIP-1py, MIP-1pz, MIP-1qa, MIP-1qb, MIP-1qc, MIP-1qd, MIP-1qe, MIP-1qf, MIP-1qg, MIP-1qh

25771-1956

2A"/"CKR-2B"or"CC-CKR-2A"/"CC-CKR-2B") [MCP-1, MCP2, MCP-3, MCP-4, MCP-5] (Charo et al., Proc. Natl. Acad. Sci. USA, 91, 2752-2756 (1994), Luster, New Eng. J. Med., 338, 436-445 (1998)); CCR3 (or"CKR-3"or"CC-CKR-3") [eotaxin-1, eotaxin-2, RANTES, MCP-3, MCP-4] (Combadiere, et al., J. Biol. Chem., 270, 16491-16494 (1995), Luster, New Eng. J. Med., 338, 436-445 (1998)); CCR-4 (or"CKR-4" or"CC-CKR-4") [TARC, MIP-1a, RANTES, MCP-1] (Power et al., J. Biol. Chem., 270, 19495-19500 (1995), Luster, New Eng. J. Med., 338, 436-445 (1998)); CCR-5 (or"CKR-5"OR"CCCKR-5") [MIP-1a, RANTES, MIP-1p] (Sanson, et al., Biochemistry, 35, 3362-3367 (1996)); CCR-6 (or"CKR-6"or "CC-CKR-6") [LARC] (Baba et al., J. Biol. Chem., 272, 14893-14898 (1997)); CCR-7 (or"CKR-7"or"CC-CKR-7") [ELC] (Yoshie et al., J. Leukoc. Biol. 62, 634-644 (1997)); CCR-8 (or"CKR-8"or"CC-CKR-8") [1-309, TARC, MIP-1p] (Napolitano et al., J. Immunol., 157, 2759-2763 (1996), Bernardini et al., Eur. J. Immunol., 28, 582-588 (1998)); CCR-10 (or"CKR-10"or"CC-CKR-10") [MCP-1, MCP-3] (Bonini et al, DNA and Cell Biol., 16, 1249-1256 (1997)) ; and CCR31 (or "CKR-11" or "CC-CKR-11") [MCP-1, MCP-2, MCP-4](Schweickart et al., J Biol Chem, 275 9550-9556 (2000)).

In addition to the mammalian chemokine receptors, the Decoy receptors CCX-CKR, D6 and DARC/Duffy as well proteins expressed by mammalian cytomegaloviruses, herpes viruses and poxviruses, exhibit binding properties of chemokine receptors (reviewed by Wells and Schwartz, Curr. Opin. Biotech., 8, 741-748 (1997); Comerford, Bioessays., 29(3):237-47 (2007)). Human CC chemokines, such as RANTES and MCP-3, can cause rapid mobilization of calcium via these virally encoded receptors. Receptor expression may be permissive for infection by allowing for the subversion of normal immune system surveillance and response to infection. Additionally, human chemokine receptors, such as CXCR-4, CCR2, CCR3, CCR5 and CCR8, can act as co receptors for the infection of mammalian cells by microbes as with, for example, the human immunodeficiency viruses (HIV).


Chemokine receptors have been implicated as being important mediators of inflammatory, infectious, and immunoregulatory disorders and diseases, including asthma and allergic diseases, as well as autoimmune pathologies such as rheumatoid arthritis, Grave's disease, chronic obstructive pulmonary disease, and atherosclerosis. For example, the chemokine receptor CCR3 is expressed among others on eosinophils, basophils, TH2 cells, alveolar macrophages, mast cells, epithelial cells, microglia cells, astrocytes and fibroblasts. CCR3 plays a pivotal role in attracting eosinophils to sites of allergic inflammation and in subsequently activating these cells. The chemokine ligands for CCR3 induce a rapid increase in intracellular calcium concentration, increased GTP exchange of G-proteins, increased ERK phosphorylation, enhanced receptor internalization, eosinophil shape change, increased expression of cellular adhesion molecules, cellular degranulation, and the promotion of migration. Accordingly, agents that inhibit chemokine receptors would be useful in such disorders and diseases. In addition, agents that inhibit chemokine receptors would also be useful in infectious diseases such as by blocking infection of CCR3 expressing cells by HIV or in preventing the manipulation of immune cellular responses by viruses such as cytomegaloviruses.

Therefore, CCR3 is an important target and antagonism of CCR3 is likely to be effective in the treatment of inflammatory, eosinophilic, immunoregulatory and infectious disorders and diseases (Wegmann, Am J Respir Cell Mol Biol., 36(1):61-67 (2007); Fryer J Clin Invest., 116(1):228-236 (2006); De Lucca, Curr Opin Drug Discov Devel., 9(4):516-524 (2006)

25771-1956

DETAILED DESCRIPTION OF THE INVENTION

Embodiment of the present invention are compounds of formula 1, wherein



A is CH₂, O or N-C₁₋₆-alkyl;

R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};
- NHR^{1.2}, NMeR^{1.2};
- NHCH₂-R^{1.3};
- NH-C₃₋₆-cycloalkyl, whereas optionally one carbon atom is replaced by a nitrogen atom, whereas the ring is optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, O-C₁₋₆-alkyl, NHSO₂-phenyl, NHCONH-phenyl, halogen, CN, SO₂-C₁₋₆-alkyl, COO-C₁₋₆-alkyl;
- a C_{9 or 10}-bicyclic-ring, whereas one or two carbon atoms are replaced by nitrogen atoms and the ring system is bound via a nitrogen atom to the basic structure of formula 1 and whereas the ring system is optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, COO-C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, NO₂, halogen, CN, NHSO₂-C₁₋₆-alkyl, methoxy-phenyl;
- a group selected from NHCH(pyridinyl)CH₂COO-C₁₋₆-alkyl, NHCH(CH₂O-C₁₋₆-alkyl)-benzoimidazolyl, optionally substituted with halogen or CN;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazole

- 5 $R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{1-6} -haloalkyl, C_{1-6} -alkylene-OH, C_{2-6} -alkenylene-OH, C_{2-6} -alkynylene-OH, $CH_2CON(C_{1-6}\text{-alkyl})_2$, $CH_2NHCONH-C_{3-6}$ -cycloalkyl, CN, CO-pyridinyl, $CONR^{1.1.1}R^{1.1.2}$, $COO-C_{1-6}$ -alkyl, $N(SO_2-C_{1-6}\text{-alkyl})(CH_2CON(C_{1-4}\text{-alkyl})_2)$ O- C_{1-6} -alkyl, O-pyridinyl, SO_2-C_{1-6} -alkyl, SO_2-C_{1-6} -alkylene-OH, SO_2-C_{3-6} -cycloalkyl, SO_2 -piperidinyl, SO_2NH-C_{1-6} -alkyl, $SO_2N(C_{1-6}\text{-alkyl})_2$, halogen, CN, CO-morpholinyl, CH_2 -pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, NHC_{1-6} -alkyl and =O;
- 10 $R^{1.1.1}$ H, C_{1-6} -alkyl, C_{3-6} -cycloalkyl, C_{1-6} -haloalkyl, $CH_2CON(C_{1-6}\text{-alkyl})_2$, CH_2CO -azetindinyl, C_{1-6} -alkylene- C_{3-6} -cycloalkyl, CH_2 -pyranyl, CH_2 -tetrahydrofuranyl, CH_2 -furanyl, C_{1-6} -alkylene-OH or thiadiazolyl, optionally substituted with C_{1-6} -alkyl;
- 15 $R^{1.1.2}$ H, C_{1-6} -alkyl, SO_2C_{1-6} -alkyl;
- 20 or $R^{1.1.1}$ and $R^{1.1.2}$ together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one N or O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, C_{1-4} -alkylene-OH, OH, =O;
- 25 or
- 30 $R^{1.1}$ is phenyl, wherein two adjacent residues are together forming a five- or six-membered carbocyclic aromatic or non-aromatic ring, optionally containing independently from each other one or two N, S, or SO_2 , replacing a carbon atom of the ring, wherein the ring is optionally substituted with C_{1-4} -alkyl or =O;
- 35 $R^{1.2}$ is selected from
- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-6} -cycloalkyl, CH_2COO-C_{1-6} -alkyl, $CONR^{1.2.1}R^{1.2.2}$, $COR^{1.2.3}$, $COO-C_{1-6}$ -alkyl, $CONH_2$, O- C_{1-6} -alkyl, halogen, CN, $SO_2N(C_{1-6}\text{-alkyl})_2$ or heteroaryl optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl;
 - heteroaryl, optionally substituted with a five- or six-membered carbocyclic non-aromatic ring containing independently from each other two N, O, S, or SO_2 , replacing a carbon atom of the ring;
 - a aromatic or non-aromatic C_9 or 10 -bicyclic-ring, whereas one or two carbon atoms are replaced by N, O or S each optionally substituted with one or two residues selected from the group consisting of $N(C_{1-6}\text{-alkyl})_2$, $CONH-C_{1-6}$ -alkyl, =O;
 - a heterocyclic non-aromatic ring, optionally substituted with pyridinyl;
 - 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with $NHCO-C_{1-6}$ -alkyl,
- 45 $R^{1.2.1}$ H, C_{1-6} -alkyl, C_{1-6} -alkylene- C_{3-6} -cycloalkyl, C_{1-4} -alkylene-phenyl, C_{1-4} -alkylene-furanyl, C_{3-6} -cycloalkyl, C_{1-4} -alkylene-O- C_{1-4} -alkyl, C_{1-6} -haloalkyl or a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O,
- 50

- S, or SO₂, replacing a carbon atom of the ring, optionally substituted with 4-cyclopropylmethyl-piperazinyl
- R^{1.2.2}** H, C₁₋₆-alkyl;
- R^{1.2.3}** a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO₂, replacing a carbon atom of the ring;
- R^{1.3}** is selected from phenyl, heteroaryl or indolyl, each optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₃₋₆-cycloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, phenyl, heteroaryl;
- R²** is selected from the group consisting of C₁₋₆-alkylene-phenyl, C₁₋₆-alkylene-naphthyl, and C₁₋₆-alkylene-heteroaryl; each optionally substituted with one, two or three residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen;
- R³** is H, C₁₋₆-alkyl;
- R⁴** is H, C₁₋₆-alkyl;
- or **R³** and **R⁴** together are forming a CH₂-CH₂ group.

Another embodiment of the present invention are compounds of formula **1**, wherein

- A** is CH₂, O or N-C₁₋₄-alkyl;
- R¹** is selected from
- NHR^{1.1}, NMeR^{1.1};
 - NHR^{1.2}, NMeR^{1.2};
 - NHCH₂-R^{1.3};
- R^{1.1}** is phenyl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-haloalkyl, C₁₋₆-alkylene-OH, C₂₋₆-alkenylene-OH, C₂₋₆-alkynylene-OH, CH₂CON(C₁₋₆-alkyl)₂, CH₂NHCONH-C₃₋₆-cycloalkyl, CN, CO-pyridinyl, CONR^{1.1.1}R^{1.1.2}, COO-C₁₋₆-alkyl, N(SO₂-C₁₋₆-alkyl)(CH₂CON(C₁₋₄-alkyl)₂) O-C₁₋₆-alkyl, O-pyridinyl, SO₂-C₁₋₆-alkyl, SO₂-C₁₋₆-alkylen-OH, SO₂-C₃₋₆-cycloalkyl, SO₂-piperidinyl, SO₂NH-C₁₋₆-alkyl, SO₂N(C₁₋₆-alkyl)₂, halogen, CN, CO-morpholinyl, CH₂-pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, NHC₁₋₆-alkyl, =O;
- R^{1.1.1}** H, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, C₁₋₆-haloalkyl, CH₂CON(C₁₋₆-alkyl)₂, CH₂CO-azetindinyl, C₁₋₆-alkylen-C₃₋₆-cycloalkyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, C₁₋₆-alkylen-OH or thiadiazolyl, optionally substituted with C₁₋₆-alkyl;
- R^{1.1.2}** H, C₁₋₆-alkyl, SO₂C₁₋₆-alkyl;
- or **R^{1.1.1}** and **R^{1.1.2}** together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one N or O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₄-alkylene-OH, OH, =O;

or

$R^{1.1}$ is phenyl, wherein two adjacent residues are together forming a five- or six-membered carbocyclic aromatic or non-aromatic ring, optionally containing independently from each other one or two N, S, or SO_2 , replacing a carbon atom of the ring, wherein the ring is optionally substituted with C_{1-4} -alkyl or $=O$;

$R^{1.2}$ is selected from

- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-6} -cycloalkyl, CH_2COO-C_{1-6} -alkyl, $CONR^{1.2.1}R^{1.2.2}$, $COR^{1.2.3}$, $COO-C_{1-6}$ -alkyl, $CONH_2$, $O-C_{1-6}$ -alkyl, halogen, CN, $SO_2N(C_{1-4}$ -alkyl) $_2$ or heteroaryl optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl;
- heteroaryl, optionally substituted with a five- or six-membered carbocyclic non-aromatic ring containing independently from each other two N, O, S, or SO_2 , replacing a carbon atom of the ring;

$R^{1.2.1}$ H, C_{1-6} -alkyl, C_{1-6} -alkylene- C_{3-6} -cycloalkyl, C_{1-4} -alkylene-phenyl, C_{1-4} -alkylene-furanyl, C_{3-6} -cycloalkyl, C_{1-4} -alkylene- $O-C_{1-4}$ -alkyl, C_{1-6} -haloalkyl or a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO_2 , replacing a carbon atom of the ring, optionally substituted with 4-cyclopropylmethyl-piperazinyl

$R^{1.2.2}$ H, C_{1-6} -alkyl;

$R^{1.2.3}$ a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO_2 , replacing a carbon atom of the ring;

$R^{1.3}$ is selected from phenyl, heteroaryl or indolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, C_{3-6} -cycloalkyl, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, phenyl, heteroaryl; preferably $R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyridinyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, C_{3-6} -cycloalkyl, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, phenyl, pyrrolidinyl;

R^2 is selected from the group consisting of C_{1-6} -alkylene-phenyl, C_{1-6} -alkylene-naphthyl, and C_{1-6} -alkylene-thiophenyl; each optionally substituted with one, two or three residues selected from the group consisting of C_{1-6} -alkyl, C_{1-6} -haloalkyl, $O-C_{1-6}$ -alkyl, $O-C_{1-6}$ -haloalkyl, halogen;

R^3 is H, C_{1-4} -alkyl;

R^4 is H, C_{1-4} -alkyl;

or R^3 and R^4 together are forming a CH_2-CH_2 group.

Another embodiment of the present invention are compounds of formula 1, wherein

A is CH_2 , O or $N-C_{1-4}$ -alkyl;

R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};

R^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-haloalkyl, C₁₋₆-alkylene-OH, C₂₋₆-alkenylene-OH, C₂₋₆-alkynylene-OH, CH₂CON(C₁₋₆-alkyl)₂, CH₂NHCONH-C₃₋₆-cycloalkyl, CN, CO-pyridinyl, CONR^{1.1.1}R^{1.1.2}, COO-C₁₋₆-alkyl, N(SO₂-C₁₋₆-alkyl)(CH₂CON(C₁₋₄-alkyl)₂) O-C₁₋₆-alkyl, O-pyridinyl, SO₂-C₁₋₆-alkyl, SO₂-C₁₋₆-alkylene-OH, SO₂-C₃₋₆-cycloalkyl, SO₂-piperidinyl, SO₂NH-C₁₋₆-alkyl, SO₂N(C₁₋₆-alkyl)₂, halogen, CN, CO-morpholinyl, CH₂-pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, NHC₁₋₆-alkyl,, =O;

R^{1.1.1} H, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, C₁₋₆-haloalkyl, CH₂CON(C₁₋₆-alkyl)₂, CH₂CO-azetindinyl, C₁₋₆-alkylene-C₃₋₆-cycloalkyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, C₁₋₆-alkylene-OH or thiadiazolyl, optionally substituted with C₁₋₆-alkyl;

R^{1.1.2} H, C₁₋₆-alkyl, SO₂C₁₋₆-alkyl;

or **R^{1.1.1}** and **R^{1.1.2}** together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one N or O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₄-alkylene-OH, OH, =O;

or

R^{1.1} is phenyl, wherein two adjacent residues are together forming a five- or six-membered carbocyclic aromatic or non-aromatic ring, optionally containing independently from each other one or two N, S, or SO₂, replacing a carbon atom of the ring, wherein the ring is optionally substituted with C₁₋₄-alkyl or =O;

R² is selected from the group consisting of C₁₋₆-alkylene-phenyl, C₁₋₆-alkylene-naphthyl, and C₁₋₆-alkylene-thiophenyl; each optionally substituted with one, two or three residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen;

R³ is H, C₁₋₄-alkyl;

R⁴ is H, C₁₋₄-alkyl;

or **R³** and **R⁴** together are forming a CH₂-CH₂ group.

Another embodiment of the present invention are compounds of formula 1, wherein

A is CH₂, O or N-C₁₋₄-alkyl;

R¹ is selected from

- NHR^{1.2}, NMeR^{1.2};

R^{1.2} is selected from

- 5
- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₃₋₆-cycloalkyl, CH₂COO-C₁₋₆-alkyl, CONR^{1.2.1}R^{1.2.2}, COR^{1.2.3}, COO-C₁₋₆-alkyl, CONH₂, O-C₁₋₆-alkyl, halogen, CN, SO₂N(C₁₋₄-alkyl)₂ or heteroaryl optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl;
 - heteroaryl, optionally substituted with a five- or six-membered carbocyclic non-aromatic ring containing independently from each other two N, O, S, or SO₂, replacing a carbon atom of the ring;
 - 10 benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolyl, each optionally substituted with one or two residues selected from the group consisting of N(C₁₋₆-alkyl)₂, CONH-C₁₋₆-alkyl, =O;
 - piperidiny, optionally substituted with pyridinyl;
 - 15 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCO-C₁₋₆-alkyl,
- 20
- R^{1.2.1} H, C₁₋₆-alkyl, C₁₋₆-alkylene-C₃₋₆-cycloalkyl, C₁₋₄-alkylene-phenyl, C₁₋₄-alkylene-furanyl, C₃₋₆-cycloalkyl, C₁₋₄-alkylene-O-C₁₋₄-alkyl, C₁₋₆-haloalkyl or a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO₂, replacing a carbon atom of the ring, optionally substituted with 4-cyclopropylmethyl-piperazinyl
- 25
- R^{1.2.2} H, C₁₋₆-alkyl;
- R^{1.2.3} a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO₂, replacing a carbon atom of the ring;
- 30
- R² is selected from the group consisting of C₁₋₆-alkylene-phenyl, C₁₋₆-alkylene-naphthyl, and C₁₋₆-alkylene-thiophenyl; each optionally substituted with one, two or three residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen;
- 35
- R³ is H, C₁₋₄-alkyl;
- R⁴ is H, C₁₋₄-alkyl;
- or R³ and R⁴ together are forming a CH₂-CH₂ group.

Another embodiment of the present invention are compounds of formula 1, wherein

- 40
- A is CH₂, O or N-C₁₋₄-alkyl;
- R¹ is selected from
- NHR^{1.2}, NMeR^{1.2};
- 45
- R^{1.2} is selected from
- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₃₋₆-cycloalkyl, CH₂COO-C₁₋₆-alkyl, CONR^{1.2.1}R^{1.2.2}, COR^{1.2.3}, COO-C₁₋₆-alkyl, CONH₂, O-C₁₋₆-alkyl, halogen, CN, SO₂N(C₁₋₄-alkyl)₂ or heteroaryl optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl;
- 50

- heteroaryl, optionally substituted with a five- or six-membered carbocyclic non-aromatic ring containing independently from each other two N, O, S, or SO₂, replacing a carbon atom of the ring;

- 5 **R^{1.2.1}** H, C₁₋₆-alkyl, C₁₋₆-alkylene-C₃₋₆-cycloalkyl, C₁₋₄-alkylene-phenyl, C₁₋₄-alkylene-furanyl, C₃₋₆-cycloalkyl, C₁₋₄-alkylene-O-C₁₋₄-alkyl, C₁₋₆-haloalkyl or a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO₂, replacing a carbon atom of the ring, optionally substituted with 4-cyclopropylmethyl-piperazinyl
- 10 **R^{1.2.2}** H, C₁₋₆-alkyl;
R^{1.2.3} a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO₂, replacing a carbon atom of the ring;
- 15 **R²** is selected from the group consisting of C₁₋₆-alkylene-phenyl, C₁₋₆-alkylene-naphthyl, and C₁₋₆-alkylene-thiophenyl; each optionally substituted with one, two or three residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen;
- 20 **R³** is H, C₁₋₄-alkyl;
R⁴ is H, C₁₋₄-alkyl;
- or **R³** and **R⁴** together are forming a CH₂-CH₂ group.
- 25

Another embodiment of the present invention are compounds of formula **1**, wherein

- 30 **A** is CH₂, O or N-C₁₋₄-alkyl;
- R¹** is selected from
- NHCH₂-R^{1.3};
- 35 **R^{1.3}** is selected from phenyl, pyrazolyl, isoxazolyl, pyridinyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₃₋₆-cycloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, phenyl, pyrrolidinyl;
- 40 **R²** is selected from the group consisting of C₁₋₆-alkylene-phenyl, C₁₋₆-alkylene-naphthyl, and C₁₋₆-alkylene-thiophenyl; each optionally substituted with one, two or three residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen;
- 45 **R³** is H, C₁₋₄-alkyl;
R⁴ is H, C₁₋₄-alkyl;
- or **R³** and **R⁴** together are forming a CH₂-CH₂ group.

50 Another embodiment of the present invention are compounds of formula **1**, wherein

- A** is CH₂, O or N-C₁₋₄-alkyl;

R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};
- NHR^{1.2}, NMeR^{1.2};
- NHCH₂-R^{1.3};
- NH-C₃₋₆-cycloalkyl, whereas optionally one carbon atom is replaced by a nitrogen atom, whereas the ring is optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, O-C₁₋₆-alkyl, NHSO₂-phenyl, NHCONH-phenyl, halogen, CN, SO₂-C₁₋₆-alkyl, COO-C₁₋₆-alkyl;
- a C_{9 or 10}-bicyclic-ring, whereas one or two carbon atoms are replaced by nitrogen atoms and the ring system is bound via a nitrogen atom to the basic structure of formula **1** and whereas the ring system is optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, COO-C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, NO₂, halogen, CN, NHSO₂-C₁₋₆-alkyl, m-methoxyphenyl;
- a group selected from NHCH(pyridinyl)CH₂COO-C₁₋₆-alkyl, NHCH(CH₂O-C₁₋₆-alkyl)-benzoimidazolyl, optionally substituted with Cl;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;

R^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₆-haloalkyl, CH₂CON(C₁₋₆-alkyl)₂, CH₂NHCONH-C₃₋₆-cycloalkyl, CN, CONR^{1.1.1}R^{1.1.2}, COO-C₁₋₆-alkyl, O-C₁₋₆-alkyl, SO₂-C₁₋₆-alkyl, SO₂-C₁₋₆-alkylen-OH, SO₂-C₃₋₆-cycloalkyl, SO₂-piperidinyl, SO₂NH-C₁₋₆-alkyl, SO₂N(C₁₋₆-alkyl)₂, halogen, CN, CO-morpholinyl, CH₂-pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, NHC₁₋₆-alkyl,, =O;

R^{1.1.1} H, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, C₁₋₆-haloalkyl, CH₂CON(C₁₋₆-alkyl)₂, CH₂CO-azetindinyl, C₁₋₆-alkylen-C₃₋₆-cycloalkyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, C₁₋₆-alkylen-OH or thiadiazolyl, optionally substituted with C₁₋₆-alkyl;

R^{1.1.2} H, C₁₋₆-alkyl, SO₂C₁₋₆-alkyl;

or **R^{1.1.1}** and **R^{1.1.2}** together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH

R^{1.2} is selected from

- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₃₋₆-cycloalkyl, CH₂COO-C₁₋₆-alkyl, CONR^{1.2.1}R^{1.2.2}, COO-C₁₋₆-alkyl, CONH₂, O-C₁₋₆-alkyl, halogen, CN, CO-pyrrolidinyl, CO-morpholinyl or heteroaryl optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolyl, each optionally substituted with one or two residues selected from the group consisting of N(C₁₋₆-alkyl)₂, CONH-C₁₋₆-alkyl, =O;
- piperidinyl, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCO-C₁₋₆-alkyl,

R^{1.2.1} H, C₁₋₆-alkyl;

$R^{1.2.2}$ H, C_{1-6} -alkyl;

$R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, C_{3-6} -cycloalkyl, O- C_{1-6} -alkyl, O- C_{1-6} -haloalkyl;

R^2 is selected from C_{1-6} -alkylene-phenyl or C_{1-6} -alkylene-naphthyl, both optionally substituted with one or two residues selected from the group consisting of C_{1-6} -alkyl, C_{1-6} -haloalkyl, O- C_{1-6} -alkyl, O- C_{1-6} -haloalkyl, halogen; or CH_2 -thiophenyl, optionally substituted with one or two residues selected from the group consisting of halogen;

R^3 is H, C_{1-4} -alkyl;

R^4 is H, C_{1-4} -alkyl;

or R^3 and R^4 together are forming a CH_2-CH_2 group.

Another embodiment of the present invention are compounds of formula 1, wherein

A is CH_2 , O or NMe;

R^1 is selected from

- $NHR^{1.1}$, $NMeR^{1.1}$;
- $NHR^{1.2}$, $NMeR^{1.2}$;
- $NHCH_2-R^{1.3}$;
- NH-cyclohexyl, optionally substituted with one or two residues selected from the group consisting of C_{1-4} -alkyl, $NHSO_2$ -phenyl, $NHCONH$ -phenyl, halogen;
- NH-pyrrolidinyl, optionally substituted with one or two residues selected from the group consisting of SO_2-C_{1-4} -alkyl, $COO-C_{1-4}$ -alkyl;
- piperidinyl, optionally substituted with one or two residues selected from the group consisting of $NHSO_2-C_{1-4}$ -alkyl, m-methoxyphenyl;
- dihydro-indolyl, dihydro-isoindolyl, tetrahydro-quinolinyl or tetrahydro-isoquinolinyl, optionally substituted with one or two residues selected from the group consisting of C_{1-4} -alkyl, $COO-C_{1-4}$ -alkyl, C_{1-4} -haloalkyl, O- C_{1-4} -alkyl, NO_2 , halogen;
- a group selected from $NHCH$ (pyridinyl) CH_2COO-C_{1-4} -alkyl, $NHCH$ (CH_2O-C_{1-4} -alkyl)-benzoimidazolyl, optionally substituted with Cl;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;

$R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of C_{1-4} -alkyl, C_{1-4} -haloalkyl, $CH_2CON(C_{1-4}-alkyl)_2$, $CH_2NHCONH-C_{3-6}$ -cycloalkyl, CN, $CONR^{1.1.1}R^{1.1.2}$, $COO-C_{1-4}$ -alkyl, O- C_{1-4} -alkyl, SO_2-C_{1-4} -alkyl, SO_2-C_{1-4} -alkylen-OH, SO_2-C_{3-6} -cycloalkyl, SO_2 -piperidinyl, SO_2NH-C_{1-4} -alkyl, $SO_2N(C_{1-4}-alkyl)_2$, halogen, CO-morpholinyl, CH_2 -pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-4} -alkyl, NHC_{1-4} -alkyl,, =O;

$R^{1.1.1}$ H, C_{1-6} -alkyl, C_{3-6} -cycloalkyl, C_{1-4} -haloalkyl, $CH_2CON(C_{1-4}-alkyl)_2$, CH_2CO -azetindinyl, C_{1-4} -alkylen- C_{3-6} -cycloalkyl, CH_2 -pyranyl, CH_2 -tetrahydrofuranyl, CH_2 -furanyl, C_{1-4} -alkylen-OH or thiadiazolyl, optionally substituted with C_{1-4} -alkyl;

$R^{1.1.2}$ H, C_{1-4} -alkyl, SO_2C_{1-4} -alkyl;

or $R^{1.1.1}$ and $R^{1.1.2}$ together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH_2OH

$R^{1.2}$ is selected from

- pyridinyl, pyridazinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of C_{1-4} -alkyl, C_{3-6} -cycloalkyl, CH_2COO-C_{1-4} -alkyl, $CONR^{1.2.1}R^{1.2.2}$, $COO-C_{1-4}$ -alkyl, $CONH_2$, $O-C_{1-4}$ -alkyl, halogen, CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-4} -alkyl;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolyl, each optionally substituted with one or two residues selected from the group consisting of $N(C_{1-4}\text{-alkyl})_2$, $CONH-C_{1-4}$ -alkyl, $=O$;
- piperidinyl, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with $NHCO-C_{1-4}$ -alkyl,

$R^{1.2.1}$ H, C_{1-4} -alkyl;

$R^{1.2.2}$ H, C_{1-4} -alkyl;

$R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C_{1-4} -alkyl, C_{3-6} -cycloalkyl, $O-C_{1-4}$ -alkyl, $O-C_{1-4}$ -haloalkyl;

R^2 is selected from C_{1-6} -alkylene-phenyl or C_{1-6} -alkylene-naphthyl, both optionally substituted with one or two residues selected from the group consisting of C_{1-4} -alkyl, C_{1-4} -haloalkyl, $O-C_{1-4}$ -haloalkyl, halogen; or CH_2 -thiophenyl, optionally substituted with one or two residues selected from the group consisting of halogen;

R^3 is H;
 R^4 is H;

or R^3 and R^4 together are forming a CH_2-CH_2 group.

Another embodiment of the present invention are compounds of formula 1, wherein

A is CH_2 , O or NMe;

R^1 is selected from

- $NHR^{1.1}$, $NMeR^{1.1}$;
- $NHR^{1.2}$, $NMeR^{1.2}$;
- $NHCH_2-R^{1.3}$;
- NH-piperidinyl, optionally substituted with pyridinyl;
- NH-cyclohexyl, optionally substituted with one or two residues selected from the group consisting of t-Bu, $NHSO_2$ -phenyl, $NHCONH$ -phenyl, F;

- NH-pyrrolidinyl, optionally substituted with one or two residues selected from the group consisting of SO₂Me, COO-t-Bu;
- piperidinyl, optionally substituted with one or two residues selected from the group consisting of NHSO₂-n-Bu, m-methoxyphenyl;
- 5 • dihydro-indolyl, dihydro-isoindolyl, tetrahydro-quinolyl or tetrahydro-isoquinolyl, optionally substituted with one or two residues selected from the group consisting of Me, COOMe, CF₃, OMe, NO₂, F, Br;
- a group selected from NHCH(pyridinyl)CH₂COOMe, NHCH(CH₂OMe)-benzoimidazolyl, optionally substituted with Cl;
- 10 • or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;

R^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, t-Bu, CF₃, CH₂CONMe₂, CH₂NHCONH-cyclohexyl, CN, CONR^{1.1.1}R^{1.1.2}, COOMe, COOEt, OMe, SO₂Me, SO₂CH₂CH₂OH, SO₂Et, SO₂-cyclopropyl, SO₂-piperidinyl, SO₂NHEt, SO₂NMeEt, F, Cl, CO-morpholinyl, CH₂-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

R^{1.1.1} H, Me, Et, t-Bu, i-Pr, cyclopropyl, CH₂-i-Pr, CH₂-t-Bu, CH(CH₃)CH₂CH₃, CH₂CHF₂, CH₂CONMe₂, CH₂CO-azetidinyl, CH₂-cyclopropyl, CH₂-cyclobutyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, CH₂CH₂OH or thiadiazolyl, optionally substituted with Me;

R^{1.1.2} H, Me, Et, SO₂Me, SO₂Et

or R^{1.1.1} and R^{1.1.2} together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH

R^{1.2} is selected from

- pyridinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, cyclopropyl, CH₂COOEt, CONR^{1.2.1}R^{1.2.2}, COOMe, COOEt, CONH₂, OMe, Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolyl, each optionally substituted with one or two residues selected from the group consisting of NMe₂, CONHMe, =O;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCOMe,

R^{1.2.1} H, Me;

R^{1.2.2} H, Me;

R^{1.3} is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, cyclopentyl, OMe, OCHF₂;

50 **R²** is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH₃, CF₃, OCF₃, F, Cl, Br, Et; or

CH₂-thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br;

5 **R**³ is H;
R⁴ is H;

or **R**³ and **R**⁴ together are forming a CH₂-CH₂ group.

10 Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

15 **R**¹ is selected from
 • NHR^{1.1}
 • NHR^{1.2},

20 **R**^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, CF₃, CH₂CONMe₂, CH₂NHCONH-cyclohexyl, CN, CONR^{1.1.1}R^{1.1.2}, COOMe, COOEt, OMe, SO₂Me, SO₂CH₂CH₂OH, SO₂Et, SO₂-cyclopropyl, SO₂-piperidinyl, SO₂NHEt, SO₂NMeEt, F, Cl, CO-morpholinyl, CH₂-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues
 25 selected from the group consisting of Me, NHMe, =O;

30 **R**^{1.1.1} H, Me, Et, t-Bu, i-Pr, cyclopropyl, CH₂-i-Pr, CH₂-t-Bu, CH(CH₃)CH₂CH₃, CH₂CHF₂, CH₂CONMe₂, CH₂CO-azetindinyl, CH₂-cyclopropyl, CH₂-cyclobutyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, CH₂CH₂OH or thiadiazolyl, optionally substituted with Me;
R^{1.1.2} H, Me, Et, SO₂Me, SO₂Et

35 or **R**^{1.1.1} and **R**^{1.1.2} together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH

40 **R**^{1.2} is selected from
 • pyridinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, cyclopropyl, CH₂COOEt, CONR^{1.2.1}R^{1.2.2}, COOMe, COOEt, CONH₂, OMe, Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;
 45 • benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolinyl, each optionally substituted with one or two residues selected from the group consisting of NMe₂, CONHMe, =O;
 • 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCOMe,

50 **R**^{1.2.1} H, Me;
R^{1.2.2} H, Me;

R² is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH₃, CF₃, OCF₃, F, Cl, Br, Et

R³ is H;

R⁴ is H.

Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};
- NHR^{1.2}, NMeR^{1.2};
- NHCH₂-R^{1.3};

R^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, CF₃, CH₂CONMe₂, CH₂NHCONH-cyclohexyl, CN, CONR^{1.1.1}R^{1.1.2}, COOMe, COOEt, OMe, SO₂Me, SO₂CH₂CH₂OH, SO₂Et, SO₂-cyclopropyl, SO₂-piperidinyl, SO₂NHEt, SO₂NMeEt, F, Cl, CO-morpholinyl, CH₂-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

R^{1.1.1} H, Me, Et, Pr, Bu, cyclopropyl, CH₂-Pr, CH₂-Bu, CH(CH₃)CH₂CH₃, CH₂CHF₂, CH₂CONMe₂, CH₂CO-azetindinyl, CH₂-cyclopropyl, CH₂-cyclobutyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, CH₂CH₂OH or thiadiazolyl, optionally substituted with Me;

R^{1.1.2} H, Me, Et, SO₂Me, SO₂Et

or **R^{1.1.1}** and **R^{1.1.2}** together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH

R^{1.2} is selected from

- pyridinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, cyclopropyl, CH₂COOEt, CONR^{1.2.1}R^{1.2.2}, COOMe, COOEt, CONH₂, OMe, Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolinyl, each optionally substituted with one or two residues selected from the group consisting of NMe₂, CONHMe, =O;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCOMe,

R^{1.2.1} H, Me;

R^{1.2.2} H, Me;

$R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, cyclopentyl, OMe, OCHF₂;

5 R^2 is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH₃, CF₃, OCF₃, F, Cl, Br, Et; or CH₂-thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br;

10 R^3 is H;
 R^4 is H;

or R^3 and R^4 together are forming a CH₂-CH₂ group.

15

Another embodiment of the present invention are compounds of formula 1, wherein

A is CH₂, O or NMe;

20 R^1 is selected from
 • NHR^{1.1}, NMeR^{1.1};

25 $R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, t-Bu, CF₃, CH₂CONMe₂, CH₂NHCONH-cyclohexyl, CN, CONR^{1.1.1}R^{1.1.2}, COOMe, COOEt, OMe, SO₂Me, SO₂CH₂CH₂OH, SO₂Et, SO₂-cyclopropyl, SO₂-piperidinyl, SO₂NHEt, SO₂NMeEt, F, Cl, CO-morpholinyl, CH₂-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

35 $R^{1.1.1}$ H, Me, Et, Bu, Pr, cyclopropyl, CH₂-Pr, CH₂-Bu, CH(CH₃)CH₂CH₃, CH₂CHF₂, CH₂CONMe₂, CH₂CO-azetindinyl, CH₂-cyclopropyl, CH₂-cyclobutyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, CH₂CH₂OH or thiadiazolyl, optionally substituted with Me;

$R^{1.1.2}$ H, Me, Et, SO₂Me, SO₂Et

40 or $R^{1.1.1}$ and $R^{1.1.2}$ together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH;

45 R^2 is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH₃, CF₃, OCF₃, F, Cl, Br, Et; or CH₂-thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br;

50 R^3 is H;
 R^4 is H;

or R^3 and R^4 together are forming a CH₂-CH₂ group.

Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};

R^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, t-Bu, CF₃, CH₂CONMe₂, CH₂NHCONH-cyclohexyl, CN, CONR^{1.1.1}R^{1.1.2}, COOMe, COOEt, OMe, SO₂Me, SO₂CH₂CH₂OH, SO₂Et, SO₂-cyclopropyl, SO₂-piperidinyl, SO₂NHEt, SO₂NMeEt, F, Cl, CO-morpholinyl, CH₂-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

R^{1.1.1} H, Me, Et, Bu, Pr, cyclopropyl, CH₂-Pr, CH₂-Bu, CH(CH₃)CH₂CH₃, CH₂CHF₂, CH₂CONMe₂, CH₂CO-azetindinyl, CH₂-cyclopropyl, CH₂-cyclobutyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, CH₂CH₂OH or thiadiazolyl, optionally substituted with Me;

R^{1.1.2} H, Me, Et, SO₂Me, SO₂Et

or **R**^{1.1.1} and **R**^{1.1.2} together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH;

R² is defined as in table 1 shown below;

R³ is H;

R⁴ is H;

Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};

R^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, t-Bu, CF₃, CH₂CONMe₂, CH₂NHCONH-cyclohexyl, CN, CONR^{1.1.1}R^{1.1.2}, COOMe, COOEt, OMe, SO₂Me, SO₂CH₂CH₂OH, SO₂Et, SO₂-cyclopropyl, SO₂-piperidinyl, SO₂NHEt, SO₂NMeEt, F, Cl, CO-morpholinyl, CH₂-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

and **R**^{1.1.1} and **R**^{1.1.2} together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH;

R^2 is defined as in table 1 shown below;;

5 R^3 is H;
 R^4 is H;

Another embodiment of the present invention are compounds of formula 1, wherein

10 **A** is CH₂, O or NMe;

R¹ is selected from
 • NHR^{1.1}, NMeR^{1.1};

15 $R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, t-Bu, CF₃, CH₂CONMe₂, CH₂NHCONH-cyclohexyl, CN, CONR^{1.1.1}R^{1.1.2}, COOMe, COOEt, OMe, F, Cl;

20 $R^{1.1.1}$ H, Me, Et, Bu, Pr, cyclopropyl, CH₂-Pr, CH₂-Bu, CH(CH₃)CH₂CH₃, CH₂CHF₂, CH₂CONMe₂, CH₂CO-azetindinyl, CH₂-cyclopropyl, CH₂-cyclobutyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, CH₂CH₂OH or thiadiazolyl, optionally substituted with Me;
 $R^{1.1.2}$ H, Me, Et, SO₂Me, SO₂Et

25 R^2 is defined as in table 1 shown below;;

R^3 is H;
 R^4 is H;

30 Another embodiment of the present invention are compounds of formula 1, wherein

A is CH₂, O or NMe;

35 **R¹** is selected from
 • NHR^{1.1}, NMeR^{1.1};

40 $R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of SO₂Me, SO₂CH₂CH₂OH, SO₂Et, SO₂-cyclopropyl, SO₂-piperidinyl, SO₂NHEt, SO₂NMeEt;

45 $R^{1.1.1}$ H, Me, Et, Bu, Pr, cyclopropyl, CH₂-Pr, CH₂-Bu, CH(CH₃)CH₂CH₃, CH₂CHF₂, CH₂CONMe₂, CH₂CO-azetindinyl, CH₂-cyclopropyl, CH₂-cyclobutyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, CH₂CH₂OH or thiadiazolyl, optionally substituted with Me;
 $R^{1.1.2}$ H, Me, Et, SO₂Me, SO₂Et

R^2 is defined as in table 1 shown below;;

50 R^3 is H;
 R^4 is H;

Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};

R^{1.1} is phenyl, optionally substituted with one residue selected from the group consisting of Me, Et, t-Bu, CF₃, CH₂CONMe₂, CH₂NHCONH-cyclohexyl, CN, CONR^{1.1.1}R^{1.1.2}, COOMe, COOEt, OMe, SO₂Me, SO₂CH₂CH₂OH, SO₂Et, SO₂-cyclopropyl, SO₂-piperidinyl, SO₂NHEt, SO₂NMeEt, F, Cl, and additionally with one residue selected from the group consisting of CO-morpholinyl, CH₂-pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

R^{1.1.1} H, Me, Et, Bu, Pr, cyclopropyl, CH₂-Pr, CH₂-Bu, CH(CH₃)CH₂CH₃, CH₂CHF₂, CH₂CONMe₂, CH₂CO-azetindinyl, CH₂-cyclopropyl, CH₂-cyclobutyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, CH₂CH₂OH or thiadiazolyl, optionally substituted with Me;

R^{1.1.2} H, Me, Et, SO₂Me, SO₂Et

R² is defined as in table 1 shown below;

R³ is H;
R⁴ is H;

Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

R¹ is selected from

- NHR^{1.2}, NMeR^{1.2};

R^{1.2} is selected from

- pyridinyl, pyridazinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, cyclopropyl, CH₂COOEt, CONR^{1.2.1}R^{1.2.2}, COOMe, COOEt, CONH₂, OMe, Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolynyl, each optionally substituted with one or two residues selected from the group consisting of NMe₂, CONHMe, =O;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCOMe,

R^{1.2.1} H, Me;
R^{1.2.2} H, Me;

R² is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH₃, CF₃, OCF₃, F, Cl, Br, Et; or CH₂-thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br;

R³ is H;

R⁴ is H;

or **R³** and **R⁴** together are forming a CH₂-CH₂ group.

Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

R¹ is selected from

- NHR^{1.2}, NMeR^{1.2};

R^{1.2} is selected from pyridinyl, pyridazinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, n-Pr, i-Pr, Bu, cyclopropyl, CH₂COOEt, CONR^{1.2.1}R^{1.2.2}, COOMe, COOEt, CONH₂, OMe, Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;

R^{1.2.1} H, Me;

R^{1.2.2} H, Me;

R² is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH₃, CF₃, OCF₃, F, Cl, Br, Et; or CH₂-thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br;

R³ is H;

R⁴ is H;

or **R³** and **R⁴** together are forming a CH₂-CH₂ group.

Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

R¹ is selected from

- NHCH₂-R^{1.3};

R^{1.3} is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, cyclopentyl, OMe, OCHF₂;

R² is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH₃, CF₃, OCF₃, F, Cl, Br, Et; or

CH₂-thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br;

5 **R**³ is H;
R⁴ is H;

or **R**³ and **R**⁴ together are forming a CH₂-CH₂ group.

10 Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;

15 **R**¹ is selected from

- NH-piperidiny, optionally substituted with pyridiny;
- NH-cyclohexyl, optionally substituted with one or two residues selected from the group consisting of t-Bu, NHSO₂-phenyl, NHCONH-phenyl, F;
- NH-pyrrolidiny, optionally substituted with one or two residues selected from the group consisting of SO₂Me, COO-t-Bu;
- 20 • piperidiny, optionally substituted with one or two residues selected from the group consisting of NHSO₂-n-Bu, m-methoxyphenyl;
- dihydro-indolyl, dihydro-isoindolyl, tetrahydro-quinoliny or tetrahydro-isoquinoliny, optionally substituted with one or two residues selected from the group consisting of Me, COOMe, CF₃, OMe, NO₂, F, Br;
- 25 • a group selected from NHCH(pyridiny)CH₂COOMe, NHCH(CH₂OMe)-benzoimidazolyl, optionally substituted with Cl;
- or 1-aminocyclopentyl, optionally substituted with Methyl-Oxadiazolyl;

30 **R**² is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH₃, CF₃, OCF₃, F, Cl, Br, Et; or CH₂-thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br;

35 **R**³ is H;
R⁴ is H;

or **R**³ and **R**⁴ together are forming a CH₂-CH₂ group.

40 Another embodiment of the present invention are compounds of formula **1**, wherein **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is and R^{1.2} is selected from

45 • pyridiny, optionally substituted with one or two residues selected from the group consisting of Me, Et, i-Pr, n-Bu, cyclopropyl, CONR^{1.2.1}R^{1.2.2}, COOMe, COOEt, CONH₂, OMe, Cl, Br CO-pyrrolidiny, CO-morpholiny or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;

• pyrrolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, COOMe, COOEt;

50 • pyrazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, cyclopropyl, COOEt, CO-pyrrolidiny;

- isoxazolyl, optionally substituted with one or two residues selected from the group consisting of t-Bu, COOEt;
- thiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, n-Pr, i-Pr, Bu, COOMe, COOEt, CH₂COOEt, CONR^{1.2.1}R^{1.2.2};
- 5 • thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of COOEt;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolyl, each optionally substituted with one or two residues selected from the group consisting of NMe₂, CONHMe, =O;
- 10 • 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCOMe,

and

- 15 R^{1.2.1} is H or Me;
R^{1.2.2} is H or Me.

Another embodiment of the present invention are compounds of formula 1, wherein **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is and R^{1.2} is selected from

- pyridinyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, i-Pr, n-Bu, CONR^{1.2.1}R^{1.2.2}, COOMe, COOEt, CONH₂, OMe, Cl, Br;
- 25 • pyrrolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, COOMe, COOEt;
- pyrazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, cyclopropyl, COOEt, CO-pyrrolidinyl;
- isoxazolyl, optionally substituted with one or two residues selected from the group consisting of t-Bu, COOEt;
- 30 • thiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, n-Pr, i-Pr, Bu, COOMe, COOEt, CONR^{1.2.1}R^{1.2.2};
- thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of COOEt;
- 35 • benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolyl, each optionally substituted with one or two residues selected from the group consisting of NMe₂, CONHMe, =O;

and

- 40 R^{1.2.1} is H or Me;
R^{1.2.2} is H or Me.

Another embodiment of the present invention are compounds of formula 1, wherein

- 45 • **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is H; R^{1.2} is pyridinyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, i-Pr, n-Bu, CONR^{1.2.1}R^{1.2.2}, COOMe, COOEt, CONH₂, OMe, Cl, Br; R^{1.2.1} is H or Me and R^{1.2.2} is H or Me.
- 50 • **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is H; R^{1.2} is pyrrolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, COOMe, COOEt; R^{1.2.1} is H or Me and R^{1.2.2} is H or Me.

- **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is H; **R**^{1.2} is pyrazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, cyclopropyl, COOEt, CO-pyrrolidinyl; **R**^{1.2.1} is H or Me and **R**^{1.2.2} is H or Me.
- 5 • **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is H; **R**^{1.2} is isoxazolyl, optionally substituted with one or two residues selected from the group consisting of t-Bu, COOE; **R**^{1.2.1} is H or Me and **R**^{1.2.2} is H or Me.
- 10 • **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is H; **R**^{1.2} is thiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, n-Pr, i-Pr, Bu, COOMe, COOEt, CONR^{1.2.1}R^{1.2.2}; **R**^{1.2.1} is H or Me and **R**^{1.2.2} is H or Me.
- 15 • **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is H; **R**^{1.2} is thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of COOEt; **R**^{1.2.1} is H or Me and **R**^{1.2.2} is H or Me.
- 20 • **A** is CH₂, O or NMe, **R**¹ is selected from NHR^{1.2}, NMeR^{1.2}; **R**² is defined as in table 1 shown below; **R**³ is H; **R**⁴ is H; **R**^{1.2} is benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinoliny, each optionally substituted with one or two residues selected from the group consisting of NMe₂, CONHMe, =O; **R**^{1.2.1} is H or Me and **R**^{1.2.2} is H or Me.

Another embodiment of the present invention are compounds of formula **1**, wherein all groups are defined as above except **R**^{1.3} is selected from

- 25 • phenyl, optionally substituted with OCHF₂;
- pyrazolyl, optionally substituted with Me or Et;
- isoxazolyl, optionally substituted with Pr;
- pyrimidinyl, optionally substituted with two OMe;
- 30 • indolyl;
- oxadiazolyl, optionally substituted with cyclopentyl.


35 Another embodiment of the present invention are compounds of formula **1**, wherein all groups are defined as above except **A** is CH₂.


Another embodiment of the present invention are compounds of formula **1**, wherein all groups are defined as above except **A** is O.


40 Another embodiment of the present invention are compounds of formula **1**, wherein all groups are defined as above except **A** is NMe.


45 Another embodiment of the present invention are compounds of formula **1**, wherein

A is CH₂, O or NMe;
R¹ is selected from











R² is selected from





R³ is H;
R⁴ is H;


or **R³** and **R⁴** together are forming a CH₂-CH₂ group.


Another embodiment of the present invention are compounds of formula **1**, wherein **A** is defined as above; **R³** is H; **R⁴** is H; and **R²** is defined as in table 1 shown below; and **R¹** is selected from














Another embodiment of the present invention are compounds of formula 1, wherein **A** is defined as above; **R**³ is H; **R**⁴ is H; and **R**² is defined as in table 1 shown below; and **R**¹ is selected from








Another embodiment of the present invention are compounds of formula 1, wherein **A** is defined as above; **R**³ is H; **R**⁴ is H; and **R**² is defined as in table 1 shown below; **R**¹ is selected from





Another embodiment of the present invention are compounds of formula 1, wherein **A** is defined as above; **R**³ is H; **R**⁴ is H; and **R**² is defined as in table 1 shown below; **R**¹ is selected from
















Table 1: $\mathbf{R}^2$ is defined as one of the groups shown below in the definitions 1 to 4:

-43-

	
Definition 3	
Definition 4	
Definition 5	
Definition 6	
Definition 7	
Definition 8	
Definition 9	
Definition 10	

25771-1956

A further embodiment of the invention are compounds of formula 1, wherein the compounds of formula 1 are present in the form of the individual optical isomers, mixtures of the individual enantiomers or racemates, preferably in the form of the enantiomerically pure compounds. Especially preferred is the R-enantiomer of the compounds of formula 1 which is the following formula **R-1**



A further embodiment of the invention are compounds of formula 1, wherein the compounds of formula 1 are present in the form of the acid addition salts thereof with pharmacologically acceptable acids as well as optionally in the form of the solvates and/or hydrates.

USED TERMS AND DEFINITIONS

Terms not specifically defined herein should be given the meanings that would be given to them by one of skill in the art in light of the disclosure and the context. As used in the specification, however, unless specified to the contrary, the following terms have the meaning indicated and the following conventions are adhered to.

In the groups, radicals, or moieties defined below, the number of carbon atoms is often specified preceding the group, for example, -C1-6 alkyl means an alkyl group or radical having 1 to 6 carbon atoms. In general, for groups comprising two or more subgroups, the last named group is the radical attachment point, for example, "thioalkyl" means a monovalent radical of the formula HS-Alk-. Unless otherwise specified below, conventional definitions of terms control and conventional stable atom valences are presumed and achieved in all formulas and groups.

In general, all tautomeric forms and isomeric forms and mixtures, whether individual geometric isomers or optical isomers or racemic or non-racemic mixtures of isomers, of a chemical structure or compound is intended, unless the specific stereochemistry or isomeric form is specifically indicated in the compound name or structure.

The term "substituted" as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valence is not exceeded, and that the substitution results in a stable compound.

By the term "optionally substituted" is meant within the scope of the invention the above-mentioned group, optionally substituted by a lower-molecular group. Examples of lower-molecular groups regarded as chemically meaningful are groups consisting of 1-200 atoms.

25771-1956

Preferably such groups have no negative effect on the pharmacological efficacy of the compounds. For example the groups may comprise:

- Straight-chain or branched carbon chains, optionally interrupted by heteroatoms, optionally substituted by rings, heteroatoms or other common functional groups.
- 5 • Aromatic or non-aromatic ring systems consisting of carbon atoms and optionally heteroatoms, which may in turn be substituted by functional groups.
- A number of aromatic or non-aromatic ring systems consisting of carbon atoms and optionally heteroatoms which may be linked by one or more carbon chains, optionally interrupted by heteroatoms, optionally substituted by heteroatoms or other common
- 10 functional groups.

The compounds disclosed herein can exist as pharmaceutically acceptable salts. The present invention includes compounds listed above in the form of salts, including acid addition salts.

15 Suitable salts include those formed with both organic and inorganic acids. Such acid addition salts will normally be pharmaceutically acceptable. However, salts of non-pharmaceutically acceptable salts may be of utility in the preparation and purification of the compound in question. Basic addition salts may also be formed and be pharmaceutically acceptable. For a more complete discussion of the preparation and selection of salts, refer to Pharmaceutical

20 Salts: Properties, Selection, and Use (Stahl, P. Heinrich. Wiley- VCHA, Zurich, Switzerland, 2002).

The term "pharmaceutically acceptable salt," as used herein, represents salts or zwitterionic forms of the compounds disclosed herein which are water or oil-soluble or dispersible and therapeutically acceptable as defined herein. The salts can be prepared during the final

25 isolation and purification of the compounds or separately by reacting the appropriate compound in the form of the free base with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, L- ascorbate, aspartate, benzoate, benzenesulfonate (besylate), bisulfate, butyrate, camphorate, camphorsulfonate, citrate, digluconate, formate,

30 fumarate, gentisate, glutarate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate (isethionate), lactate, maleate, malonate, DL-mandelate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphonate, picrate, pivalate,

35 propionate, pyroglutamate, Succinate, Sulfonate, tartrate, L-tartrate, trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, para- toluenesulfonate (p-tosylate), and undecanoate. Also, basic groups in the compounds disclosed herein can be quaternized with methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dimethyl, diethyl, dibutyl, and diamyl Sulfates; decyl, lauryl, myristyl, and steryl chlorides, bromides, and iodides; and

40 benzyl and phenethyl bromides. Examples of acids which can be employed to form pharmaceutically acceptable addition salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, and citric. Salts can also be formed by coordination of the compounds with an alkali metal or alkaline earth ion. Hence, the present invention contemplates sodium, potassium,

45 magnesium, and calcium salts of the compounds disclosed herein, and the like.

Basic addition salts can be prepared during the final isolation and purification of the compounds by reacting a carboxy group with a suitable base such as the hydroxide, carbonate, or bicarbonate of a metal cation or with ammonia or an organic primary,


50 secondary, or tertiary amine. The cations of therapeutically acceptable salts include lithium, sodium, potassium, calcium, magnesium, and aluminum, as well as nontoxic quaternary amine cations such as ammonium, tetramethylammonium, tetraethylammonium,

25771-1956


methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, dicyclohexylamine, procaine, dibenzylamine, N, N-dibenzylphenethylamine, 1-ephedrine, and N,P-dibenzylethylenediamine. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, and piperazine.

While it may be possible for the compounds of the subject invention to be administered as the raw chemical, it is also possible to present them as a pharmaceutical formulation. Accordingly, provided herein are pharmaceutical formulations which comprise one or more of certain compounds disclosed herein, or one or more pharmaceutically acceptable salts, esters, prodrugs, amides, or solvates thereof, together with one or more pharmaceutically acceptable carriers thereof and optionally one or more other ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Proper formulation is dependent upon the route of administration chosen. Any of the well-known techniques, carriers, and excipients may be used as suitable and as understood in the art; e.g., in Remington's Pharmaceutical Sciences. The pharmaceutical compositions disclosed herein may be manufactured in any manner known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.


By the term heterocyclic rings ("het") are meant five-, six- or seven-membered, saturated or unsaturated heterocyclic rings or 5-10 membered, bicyclic hetero rings which may contain one, two or three heteroatoms, selected from among oxygen, sulphur and nitrogen; the ring may be linked to the molecule by a carbon atom or, if present, by a nitrogen atom. The following are examples of five-, six- or seven-membered, saturated or unsaturated heterocyclic rings:




Unless stated otherwise, a heterocyclic ring may be provided with a keto group. Examples include:



Examples of 5-10-membered bicyclic hetero rings are pyrrolizine, indole, indolizine, isoindole, indazole, purine, quinoline, isoquinoline, benzimidazole, benzofurane, benzopyrane, benzothiazole, benzoisothiazole, pyridopyrimidine, pteridine, pyrimidopyrimidine,



Although the term heterocyclic rings includes heterocyclic aromatic groups, the term heterocyclic aromatic groups ("hetaryl") denotes five- or six-membered heterocyclic aromatic groups or 5-10 membered, bicyclic hetaryl rings which may contain one, two or three heteroatoms, selected from among oxygen, sulphur and nitrogen, which contain sufficient conjugated double bonds that an aromatic system is formed. The ring may be linked to the molecule through a carbon atom or if present through a nitrogen atom. The following are examples of five- or six-membered heterocyclic aromatic groups:



Examples of 5-10-membered bicyclic hetaryl rings include pyrrolizine, indole, indolizine, isoindole, indazole, purine, quinoline, isoquinoline, benzimidazole, benzofuran, benzopyrane, benzothiazole, benzoisothiazole, pyridopyrimidine, pteridine, pyrimidopyrimidine.

The term "halogen" as used herein means a halogen substituent selected from fluoro, chloro, bromo or iodo.

By the term "C₁₋₆-alkyl" (including those which are part of other groups) are meant branched and unbranched alkyl groups with 1 to 6 carbon atoms, and by the term "C₁₋₄-alkyl" are meant branched and unbranched alkyl groups with 1 to 4 carbon atoms. Alkyl groups with 1 to 4 carbon atoms are preferred. Examples of these include: methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *sec*-butyl, *tert*-butyl, *n*-pentyl, *iso*-pentyl, *neo*-pentyl or hexyl. The abbreviations Me, Et, *n*-Pr, *i*-Pr, *n*-Bu, *i*-Bu, *t*-Bu, etc. may optionally also be used for the above-mentioned groups. Unless stated otherwise, the definitions propyl, butyl, pentyl and hexyl include all the possible isomeric forms of the groups in question. Thus, for example, propyl includes *n*-propyl and *iso*-propyl, butyl includes *iso*-butyl, *sec*-butyl and *tert*-butyl etc.

By the term "C₁₋₆-alkylene" (including those which are part of other groups) are meant branched and unbranched alkylene groups with 1 to 6 carbon atoms and by the term "C₁₋₄-alkylene" are meant branched and unbranched alkylene groups with 1 to 4 carbon atoms. Alkylene groups with 1 to 4 carbon atoms are preferred. Examples include: methylene, ethylene, propylene, 1-methylethylene, butylene, 1-methylpropylene, 1,1-dimethylethylene, 1,2-dimethylethylene, pentylene, 1,1-dimethylpropylene, 2,2-dimethylpropylene, 1,2-dimethylpropylene, 1,3-dimethylpropylene or hexylene. Unless stated otherwise, the definitions propylene, butylene, pentylene and hexylene also include all the possible isomeric forms of the relevant groups with the same number of carbons. Thus for example propyl also includes 1-methylethylene and butylene includes 1-methylpropylene, 1,1-dimethylethylene, 1,2-dimethylethylene.

The term "C₂₋₆-alkenyl" (including those which are part of other groups) denotes branched and unbranched alkenyl groups with 2 to 6 carbon atoms and the term "C₂₋₄-alkenyl" denotes branched and unbranched alkenyl groups with 2 to 4 carbon atoms, provided that they have

at least one double bond. Preferred are alkenyl groups with 2 to 4 carbon atoms. Examples include: ethenyl or vinyl, propenyl, butenyl, pentenyl, or hexenyl. Unless otherwise stated, the definitions propenyl, butenyl, pentenyl and hexenyl include all possible isomeric forms of the groups in question. Thus, for example, propenyl includes 1-propenyl and 2-propenyl, butenyl includes 1-, 2- and 3-butenyl, 1-methyl-1-propenyl, 1-methyl-2-propenyl etc.

By the term "C₂₋₆-alkenylene" (including those which are part of other groups) are meant branched and unbranched alkenylene groups with 2 to 6 carbon atoms and by the term "C₂₋₄-alkenylene" are meant branched and unbranched alkylene groups with 2 to 4 carbon atoms. Alkenylene groups with 2 to 4 carbon atoms are preferred. Examples include: ethenylene, propenylene, 1-methylethenylene, butenylene, 1-methylpropenylene, 1,1-dimethylethenylene, 1,2-dimethylethenylene, pentenylene, 1,1-dimethylpropenylene, 2,2-dimethylpropenylene, 1,2-dimethylpropenylene, 1,3-dimethylpropenylene or hexenylene. Unless stated otherwise, the definitions propenylene, butenylene, pentenylene and hexenylene include all the possible isomeric forms of the respective groups with the same number of carbons. Thus, for example, propenyl also includes 1-methylethenylene and butenylene includes 1-methylpropenylene, 1,1-dimethylethenylene, 1,2-dimethylethenylene.

By the term "C₂₋₆-alkynyl" (including those which are part of other groups) are meant branched and unbranched alkynyl groups with 2 to 6 carbon atoms and by the term "C₂₋₄-alkynyl" are meant branched and unbranched alkynyl groups with 2 to 4 carbon atoms, provided that they have at least one triple bond. Alkynyl groups with 2 to 4 carbon atoms are preferred. Examples include: ethynyl, propynyl, butynyl, pentynyl, or hexynyl. Unless stated otherwise, the definitions propynyl, butynyl, pentynyl and hexynyl include all the possible isomeric forms of the respective groups. Thus, for example, propynyl includes 1-propynyl and 2-propynyl, butynyl includes 1-, 2- and 3-butynyl, 1-methyl-1-propynyl, 1-methyl-2-propynyl etc.

By the term "C₂₋₆-alkynylene" (including those which are part of other groups) are meant branched and unbranched alkynylene groups with 2 to 6 carbon atoms and by the term "C₂₋₄-alkynylene" are meant branched and unbranched alkylene groups with 2 to 4 carbon atoms. Alkynylene groups with 2 to 4 carbon atoms are preferred. Examples include: ethynylene, propynylene, 1-methylethynylene, butynylene, 1-methylpropynylene, 1,1-dimethylethynylene, 1,2-dimethylethynylene, pentynylene, 1,1-dimethylpropynylene, 2,2-dimethylpropynylene, 1,2-dimethylpropynylene, 1,3-dimethylpropynylene or hexynylene. Unless stated otherwise, the definitions propynylene, butynylene, pentynylene and hexynylene include all the possible isomeric forms of the respective groups with the same number of carbons. Thus for example propynyl also includes 1-methylethynylene and butynylene includes 1-methylpropynylene, 1,1-dimethylethynylene, 1,2-dimethylethynylene.

The term "C₃₋₆-cycloalkyl" (including those which are part of other groups) as used herein means cyclic alkyl groups with 3 to 8 carbon atoms, preferred are cyclic alkyl groups with 5 to 6 carbon atoms. Examples include: cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

By the term "C₁₋₆-haloalkyl" (including those which are part of other groups) are meant branched and unbranched alkyl groups with 1 to 6 carbon atoms wherein one or more hydrogen atoms are replaced by a halogen atom selected from among fluorine, chlorine or bromine, preferably fluorine and chlorine, particularly preferably fluorine. By the term

"C₁₋₄-haloalkyl" are meant correspondingly branched and unbranched alkyl groups with 1 to 4 carbon atoms, wherein one or more hydrogen atoms are replaced analogously to what was stated above. C₁₋₄-haloalkyl is preferred. Examples include: CH₂F, CHF₂, CF₃,


5

EXAMPLES

S. GENERAL SYNTHESIS PROCEDURES FOR EXAMPLES

10 The examples of the present invention are synthesized according to the following general scheme:

General synthesis scheme:







15


where "building block" and "reaction partner" are defined for the procedures A-G below in which A, R¹, R², R³ and R⁴ are defined as above. In case a building block or reaction partner is not commercially available, its synthesis is described below, commencing from a

20 commercially available starting material, and where necessary, involving intermediate(s), "I".


The term "linker" is used to mean a chemical moiety whose definition is restricted to that of R¹ and is able to bear the specific functional group as stated in order to undergo the reaction of the procedure. In procedure E, the synthesis starts from intermediate I12.2 which defines the "building block". In procedure G, the synthesis starts from example 181 or 194 which

25 define the "building block"


S.1. Procedures A – G:**Procedure A:**5 building block A1-A8**Procedure B:**10 building block B1 - B20**Procedure C:**building block C1**Procedure D:**15 building block D1 -D2

Procedure E:

112.2 (see synthesis of building block A7)

Procedure F:

5 building block F1 - F3

Procedure G:

example 181 or 194

10


S.2. Building block syntheses**S.2.1 Synthesis of building blocks of formula A**

15

Synthesis of building blocks A1 – A4

Synthesis of building block A1

Step 1:




5

4-Chloro-3-methyl benzylbromide (20 g) [synthesized according to literature: J.L. Kelley, J.A. Linn, J.W.T. Selway, *J. Med. Chem.* 1989, 32(8), 1757-1763], 4-Piperidone (22 g), K_2CO_3 (26 g) in acetonitrile (300 ml) was heated at 50°C for 14h. The suspension was filtered and the filtrate concentrated in vacuum. The residue was purified by flash chromatography (cyclohexane/ EtOAc 1:1) to yield 17.4 g of intermediate **I1**.

10

Step 2:




15

Intermediate **I1** (10 g) and D-H-Glu(O^tBu)-OMe (10.8 g) was dissolved in DMF (200 ml) and HOAc (5 ml). Then molecular sieve (1.0 g, 4Å, powder) was added and the suspension was stirred overnight. Sodium triacetoxyborohydride (37.5 g) was added and the suspension stirred until complete conversion of the intermediate formed imine was observed. A basic pH was achieved by slow addition (foam formation!) of aqueous $NaHCO_3$ solution before additional water and DCM (500 mL) was added. The organic phase was separated and the water phase extracted with DCM (500 ml). The organic phase was washed with brine, dried and concentrated in vacuum to yield 19.5 g (74% purity) of intermediate **I2**.

20

Step 3:




25

Intermediate **I2** (19.5 g, 74% purity) was dissolved in DCM (40 ml) and trifluoroacetic acid (20 ml, TFA). The solution was stirred at 25°C for 14 h before additional 40 ml TFA was added

30


and the solution continued stirring for further 7 h. The reaction mixture was then concentrated in vacuum, dissolved in toluene and concentrated again to provide 29.5 g (purity 55%) of intermediate **I3**.

5 **Step 4:**






10 Intermediate **I3** (29 g, purity 55%) was dissolved in DCM (100 ml) and DIPEA (22 ml). TBTU (15 g) was added and the solution was stirred for 30 min. Then DCM (150 ml), water (100 ml) and saturated NaHCO₃ solution (100 ml) was added, the organic phase was separated and the water phase extracted once with DCM (100 ml). The organic phase was dried and concentrated to provide an oil which was then fractionated via reversed phase HPLC. The fractions containing the building block D1 were concentrated in vacuum, then a basic pH was
15 adjusted with addition of NaHCO₃ solution and the product was extracted with DCM to provide 8.1 g of intermediate **I4**.

Step 5:




20 Intermediate **I4** (7g) was dissolved in dioxane (50 ml). 2.5 M LiOH solution (23 ml) and water (20 ml) was added and stirred at 25°C overnight. The solution was acidified with aqueous 4N HCl and then concentrated in vacuum. The residue was dissolved in water, acetonitrile and a small amount of dioxane and lyophilised to provide 10.4 g solid (71% purity) of building block
25 **A1**.

The **building blocks** were synthesized in analogy to acid **A1**:



5 Synthesis of building block A5


Step 1:



10 Acetic acid (5.7 ml,) was added to a stirred suspension of (R)-2-amino-3-hydroxy-propionic acid methyl ester hydrochloride (5.98 g) and 4-oxo-piperidine-1-carboxylic acid benzyl ester (9.86 g) in DCM (382 ml), and the conditions maintained for 70 min. To the resulting solution was then added sodium triacetoxyborohydride (28.51 g) in one portion, and the resulting suspension stirred overnight. The reaction mixture was then neutralized with a saturated aqueous solution of NaHCO₃, the organic phase concentrated under reduced pressure to 60 ml, basified (pH 8.5) with a saturated aqueous solution of NaHCO₃ and solid Na₂CO₃, MeOH (20 ml) was added and the phases separated, the aqueous one extracted twice with a DCM: MeOH 3:1 mixture, the combined organic layers dried over Na₂SO₄ and evaporated to dryness to afford the product. The crude **15** (17.58 g) was used in the next step without any further purification.

20 HPLC (R_t) = 2.48 min (method L)


Step 2:



Intermediate **I5** (8 g, 35% content) was dissolved in THF (250 ml), then triethylamine (3.48 ml), 4-dimethylamino pyridine (100 mg) and 1,1' carbonyldiimidazole (5.40 g) were added in sequence and the whole mixture stirred and refluxed under a nitrogen atmosphere for 48 h.


The solvent and the volatiles were removed under reduced pressure and the resulting residue was taken up in EtOAc (150 ml), washed with water (3 x 100 ml), 10% citric acid (100 ml) and brine (100 ml). The organic phase was dried over Na₂SO₄, filtered and evaporated to dryness to afford product (6.6 g). This was purified twice (50 g Isolute[®] silica gel cartridge, eluting in the first purification with EtOAc/n-hexane 80/20 and with 100% EtOAc in the second purification); the title compound (1.27 g).
HPLC (R_t) = 3.16 min (method G)

Step 3:



A suspension of 10% Pd/C (380 mg) in 50% aq. MeOH (4 ml) was added to a stirred solution of intermediate **I6** (3.79 g) in MeOH (28 ml). The mixture was then stirred under a positive pressure of hydrogen 14 h. The reaction mixture was then filtered over Celite[®], the cake washed with MeOH (4 x 10 ml) and the filtrate evaporated to dryness under reduced pressure to afford the desired free amine (2.2 g).
HPLC (R_t) = 0.51 min (method L)

Steps 4 + 5:



Intermediate **I7** (1.01 g) was suspended in dry acetonitrile (17.8 ml) in a nitrogen atmosphere; 4-bromomethyl-1-chloro-2-methyl-benzene (1.46 g) and K₂CO₃ (1.22 g) were added in sequence and the whole stirred overnight. The solvent was removed under reduced pressure, the residue was taken up in EtOAc (50 ml), washed with water (1 x 30 ml), the layers separated, the aqueous extracted with EtOAc (1x 50 mL) and a 3:1 DCM:MeOH mixture (50 ml). The combined organic layers were dried over Na₂SO₄, filtered and evaporated to dryness. The residue was taken up in DCM and loaded on a 10g-SCX cartridge eluting with DCM (50 ml), MeOH (50 ml), then ammonia (50 ml of a 0.5 M MeOH


solution). Evaporation under reduced pressure of the MeOH and methanolic ammonia fractions afforded the product (895 mg) under vacuum, as a mixture of the methyl and benzyl esters (4.5:1 by LC-MS).

The mixture of esters (895 mg) was suspended in THF (10 ml) and MeOH (2 ml) was then added; the solution was ice chilled, LiOH·H₂O (155 mg) and water (10 ml) were added sequentially. The reaction was allowed to warm to room temperature and the resulting solution stirred for 70 min. The reaction mixture was diluted with diethyl ether (10 ml), the phases separated; the aqueous was then acidified to pH 4 by dropwise addition of 6N HCl and freeze-dried overnight to afford the carboxylic acid-hydrochloride salt. The residue was

suspended in a 4:1 DCM : iPrOH mixture, stirred at room temperature for 10 min, then filtered on cotton, the solvent evaporated to dryness under reduced pressure to afford the product (844 mg).


HPLC (R_t) = 2.12 min (method H)

The following **building blocks** were synthesized in analogy to acid **A5**:



Synthesis of building block A7


Step 1:




To a stirred suspension of (R)-2-amino-3-tert-butoxycarbonylamino-propionic acid methyl ester hydrochloride (5.0 g) and 1-(4-chloro-3-methyl-benzyl)-piperidin-4-one (5.13 g) in DCM (200 ml), was added acetic acid (2.81 ml). After 70 min sodium triacetoxymethylborohydride (14.56 g) was added in one portion, and the resulting suspension stirred for 48 h. The reaction mixture was then diluted with a saturated aqueous solution of NaHCO₃ (1 x 150 ml), the two layers separated, the aqueous extracted with DCM (1 x 200 ml), the combined organic layers

dried over Na₂SO₄ and evaporated to dryness to afford the product. The crude was purified by flash chromatography (Biotage® SP1; 65M silica gel cartridge; gradient elution: DCM/MeOH/NH₃ from 98/2/0.2 to 80/20/2 in 12 column volumes) affording **I9** (7.0 g). HPLC (R_t) = 10.03 min (method O)

5


Step 2:

To a stirred solution of Intermediate **I9** (7.00 g in dioxane (50 ml), at room temperature, was added HCl (39.78 ml of a 4N solution in 1,4-dioxane) dropwise. MeOH (20 ml) was then added, and the reaction mixture stirred overnight. The solvents and the volatiles were removed under reduced pressure, and the residue was triturated with diethyl ether affording product **I10** (6.1 g). HPLC (R_t) = 1.89 min (method H)


15 **Step 3:**

Intermediate **I10** (6.1 g) was suspended in THF (300 ml), then triethylamine (12.41 ml), and 1,1' carbonyldiimidazole (5.99 g) were added sequentially and the reaction refluxed under a nitrogen atmosphere for 100 min. The reaction mixture was allowed to cool down, filtered and the filtrate evaporated under reduced pressure. and the residue was purified by flash column chromatography (Biotage® SP1; 65M silica gel cartridge; gradient elution: DCM/MeOH/NH₃ from 98/2/0.2 to 80/20/2 in 12 column volumes), affording a solid which was triturated with diethyl ether, to give **I11** (1.85 g)

HPLC (R_t) = 8.37 min (method O)


Step 4:

5 A stirred solution of potassium *t*-butoxide in THF (7.38 ml, 1M) at -50 °C under a nitrogen atmosphere was added dropwise to a solution of intermediate **I11** (1.80 g) in dry THF (45 ml). After 1h, a solution of iodomethane (0.46 ml) in dry THF (5ml) was added dropwise to the resulting suspension, and the temperature was allowed to warm to 0 °C over 1 h. After this time, excess iodomethane was removed under reduced pressure at room temperature, the reaction was quenched with saturated NH₄Cl(aq) (10 ml), the salts were filtered off and
 10 the filtrate diluted with EtOAc (30 mL) and brine (15 ml), the layers separated, the organic washed once more with brine (15 ml), dried over Na₂SO₄ and evaporated to dryness to afford a colourless oil, that was purified by flash column chromatography (Biotage® SP1; 65M silica gel cartridge; gradient elution: DCM/MeOH from 98/2 to 80/20 in 15 column volumes), affording product **I12** (600 mg).
 15 HPLC (R_t) = 8.92 min (method O)

Step 5:

20 Intermediate **I12** (0.9 g) was dissolved in 4N HCl (30 ml) and refluxed for 1 h. The solvent was evaporated to afford an oily residue that was taken up in acetone and evaporated to dryness to afford a white solid, which was triturated with diisopropyl ether to afford **building block A7** (0.79 g).
 25 HPLC (R_t) = 5.48 min (method O)

Building block **A8** was synthesized in analogy to **A7**:




A8

S.2.2. Synthesis of building blocks with formula B

5

Synthesis of building block B1



example 87


B1

10 To a stirred solution of example **87** (350 mg) in dioxane (10 ml) at 25°C was added LiOH(aq) (0.7 ml, 2.5 M) and water (2 ml). After 3 h, 4N HCl(aq) solution was added to adjust the pH to 1-2. The solution was concentrated in vacuum, and water and acetonitrile added and the mixture lyophilised giving **building block B1**. HPLC: 1.25 min (method D). This was used without further purification in the next step.


15

Synthesis of building blocks B2 - B15


20 The **building blocks B2-B15** were synthesized in analogy to **building block B1** from the examples 88 - 92 (**B2-B6**), 110 (**B7**), 113 (**B8**), 111 (**B9**), 112 (**B10**), 109 (**B11**), 216 (**B12**), 215 (**B13**), 195 (**B14**), 194 (**B15**)



B2














B3



B4


25

**B5****B6****B7****B8****B9****B10****B11****B12****B13****B14****B15**

5

10

15

Synthesis of building block B16**Step 1:****3-Bromo-N-methyl-5-nitro-benzamide**

To a solution of 3-bromo-5-nitro-benzoic acid (2.1 g) in DCM (50 ml) was added oxalylchloride (1.5 ml). 1 drop of DMF was added and the reaction stirred at room temperature for 2 h. The reaction was concentrated in vacuum. This was dissolved in THF and added dropwise to a pre-cooled 2M solution of methylamine in THF (22 ml) at 0°C. This was allowed to warm to room temperature and further stirred overnight. The reaction mixture was filtered and the filtrate concentrated in vacuum. The residue was dissolved in EtOAc and washed with water. The organic layer was separated, dried and concentrated in vacuum to leave the product (2.1 g). Rt 1.37min (method D).

Step 2:**5-(3-Bromo-5-nitro-phenyl)-1-methyl-1H-tetrazole**

To a solution of 3-bromo-N-methyl-5-nitro-benzamide (0.6 g) in DCM (15 ml) at -15°C was added triflic anhydride (0.6 ml) and the reaction stirred for 30 min. Sodium azide (226 mg) was added and the reaction allowed to warm to room temperature and stirred further overnight. NaHCO₃(aq) was added to neutralise the reaction and the mixture extracted into EtOAc. The organic layer was separated, dried and concentrated in vacuum and the residue purified by flash chromatography (9:1 to 2:8 cyclohexane:EtOAc) to give the title compound (290 mg). R_f = 0.51 (3:2 cyclohexane:EtOAc).

Step 3:**3-Bromo-5-(1-methyl-1H-tetrazol-5-yl)-phenylamine**

A solution of 5-(3-Bromo-5-nitro-phenyl)-1-methyl-1H-tetrazole (2.1 g) in EtOAc (100 ml) was hydrogenated on a H-cube® apparatus at 20°C, room pressure with a flow rate of 1 ml/min using a Raney-Ni cartridge. After reaction completion, the mixture was concentrated in vacuum and triturated with MeOH. The solid was filtered and further triturated and air-dried to give the title compound (290 mg). Rt 1.25 min (method D).

Step 4:**1-[1-(4-Chloro-3-methyl-benzyl)-piperidin-4-yl]-5-oxo-pyrrolidine-2-carboxylic acid [3-bromo-5-(1-methyl-1H-tetrazol-5-yl)-phenyl]-amide**

To a solution of 1-[1-(4-Chloro-3-methyl-benzyl)-piperidin-4-yl]-5-oxo-pyrrolidine-2-carboxylic acid as DIPEA salt (2.6 g) in DMF (100 ml) was added DIPEA (2.1 g), and HATU (4 g). 3-Bromo-5-(1-methyl-1H-tetrazol-5-yl)-phenylamine (1.4 g) was added and the reaction stirred overnight at rt. The reaction mixture was filtered and the filtrate partitioned between water and EtOAc. The organic layer was separated, dried and concentrated in vacuum and the residue purified by flash chromatography (DCM:MeOH 100:0 to 95:5 gradient) to give the title compound (400 mg). R_f = 0.19 (95:5 DCM:MeOH).

Step 5:**3-({1-[1-(4-Chloro-3-methyl-benzyl)-piperidin-4-yl]-5-oxo-pyrrolidine-2-carbonyl}-amino)-5-(1-methyl-1H-tetrazol-5-yl)-benzoic acid methyl ester**


5 To a solution of 1-[1-(4-Chloro-3-methyl-benzyl)-piperidin-4-yl]-5-oxo-pyrrolidine-2-carboxylic acid [3-bromo-5-(1-methyl-1H-tetrazol-5-yl)-phenyl]-amide (400 mg) in MeOH (20 ml) was added triethylamine (140 μ l), DMF (5 ml), and [1,1'-bis(diphenylphosphino)-ferrocen]palladium(II) chloride (50 mg) and the reaction stirred for 2 d at 80°C under a 3bar CO atmosphere. The reaction mixture was filtered and the filtrate concentrated in vacuum.

10 The residue was partitioned between water and EtOAc and the organic layer was then separated, dried and concentrated in vacuum. The residue was purified by flash chromatography (DCM:MeOH 100:0 to 95:5 gradient) to give the title compound (400 mg). R_f = 0.37 (95:5 DCM:MeOH).

Step 6:**3-({1-[1-(4-Chloro-3-methyl-benzyl)-piperidin-4-yl]-5-oxo-pyrrolidine-2-carbonyl}-amino)-5-(1-methyl-1H-tetrazol-5-yl)-benzoic acid**

15 To a solution of 3-({1-[1-(4-Chloro-3-methyl-benzyl)-piperidin-4-yl]-5-oxo-pyrrolidine-2-carbonyl}-amino)-5-(1-methyl-1H-tetrazol-5-yl)-benzoic acid methyl ester (270 mg) in MeOH (10 ml) at room temperature was added 4M NaOH(aq) (10 ml) and the reaction stirred overnight. The organic and aqueous layers were separated and the organic layer concentrated in vacuum. The residue was triturated in toluene and diethyl ether successively leaving the title compound (220 mg). R_t = 0.96min (method E).

20

Synthesis of building block B17**Step 1:**

30 To a stirred solution of 1-[1-(4-Chloro-3-methyl-benzyl)-piperidin-4-yl]-5-oxo-pyrrolidine-2-carboxylic acid (500 mg) in DMF (5 ml) at room temperature was added DIPEA (1.1 ml) followed by HATU (1.1 g). After 45min, intermediate I13 (0.34 g) was added and the reaction


stirred overnight at rt. Saturated brine solution was added followed by EtOAc. The organic layer was separated, dried and concentrated in vacuum and the residue purified by HPLC to give the product (557 mg). (R_t 1,50 min, method D)

5 **Step 2:**

To a stirred solution of the product of step 1 (557 mg) in dioxane (5 ml) was added lithium hydroxide (195 mg) dissolved in a sufficient amount of water. After overnight stirring, the reaction was acidified with HCl(aq) (4M), then concentrated in vacuum. The residue was purified by HPLC to afford building block B17 (155mg).

10 (R_t 1.34min, method D).

Synthesis of I13 for building block B17:



Step 1:

Ethyl 2-bromo-4-methyl-3-oxopentanoate

20

To a suspension of sodium acetate (1.9 g) in glacial acetic acid (15 ml) was added 4-methyl-3-oxo-pentanoic acid ethyl ester (2 g) and the reaction mixture cooled to 10°C. Bromine (0.7 ml) was then added dropwise. The resulting solution was allowed to warm to room temperature and stirred for 1 h. The reaction was complete (by TLC) and was concentrated in vacuum affording 3.8 g crude ethyl product which was used directly in the next synthetic step.

25

Step 2:


Ethyl 2-amino-4-isopropylthiazole-5-carboxylate (**I13**)

30

A solution of thiourea (1.1 g) in ethanol (10 mL) was brought to reflux. Ethyl 2-bromo-4-methyl-3-oxopentanoate (3.0 g) was then added and the reaction mixture refluxed further for 1 h. The reaction was complete (by TLC) and was concentrated in vacuum. Flash chromatography (100% DCM → 90:10 DCM:MeOH) afforded the product (1.1 g).

35


Synthesis of building block B18



Intermediate **I14** was synthesized according to procedure D with subsequent saponification.

Synthesis of intermediate **I16**:

5



Step 1:

10

2-chloro-6-methylnicotinic acid (9 g), of aq ammonia (44 ml), of Cu(II)SO₄ (0.9 g) and of sodium sulphide (0.32 g) were added to an autoclave and heated to 155°C overnight. The crude product was suspended in water to yield product **I15** (3.6 g). The filtrate was concentrated and again suspended in water to yield a further 1.9 g of product **I15**. HPLC: R_t = 0.37 min (method D)

15


Step 2:

20

To 50 ml of MeOH was added dropwise acetylchloride (3 ml) at rt. After 15 min, intermediate **I15** (2.3 g) was added and the mixture was stirred overnight at 50°C. After concentrating the solution, the resulting residue was suspended in acetone and then filtered and dried at 50°C in vacuum, to yield product **I16** (4.1g). HPLC: R_t = 0.91 min (method D)


Synthesis of building block B19

25




B19 was synthesized in analogy to **B17**. Synthesis of **I13a** for building block **B19** was analogous to that for **I13**.

30




Synthesis of building block B20



B20

- 5 **B20** was synthesized in analogy to **B17**. Synthesis of **I13b** for building block **B20** was analogous to that for **I13**.




I13b

10


S.2.3. Synthesis of building blocks with formula C

Synthesis of building block C1




15


- Building block A1**, as its N,N-diisopropylethylamine salt (500 mg), was suspended in dry DMF (7 ml) under inert atmosphere and TBTU (836 mg) was added, followed by N,N-diisopropylethylamine (0.53 ml). After stirring for 1 h at room temperature, hexamethyldisilazane (0.44 ml) was added and the mixture was stirred for 6 h. Further portions of TBTU (334 mg) and hexamethyldisilazane (0.22 ml) were added and the reaction was stirred for further 18 h. The solvent was evaporated under reduced pressure and residue partitioned between saturated aqueous solution of NaHCO₃ and EtOAc. The layers were separated and the aqueous phase extracted with EtOAc. The combined organic extracts were washed with brine, dried under Na₂SO₄, filtered and the solvent was evaporated under reduced pressure. The crude was purified by flash chromatography (20 g Isolute[®] silica gel cartridge; eluent: DCM/MeOH/NH₄OH 95/5/0.5) affording the title compound. (295 mg). UPLC (R_t) = 1.24 min (method M)

S.2.4. Synthesis of building blocks with formula D**5 Synthesis of building block D1****Step 1:**


10 Intermediate **I17** was prepared starting from (R)-2-amino-3-tert-butoxycarbonylamino-propionic acid methyl ester hydrochloride (10.00 g) and 4-oxo-piperidine-1-carboxylic acid phenyl ester (12.07 g), following the procedures described for the synthesis of **I9**. Product obtained 19.50 g (content 70%). HPLC (R_t) = 3.20 min (method Q)

15 Step 2:

20 Intermediate **I18** was prepared starting from **I17** (19.5 g, content 70%) and following the procedures described for the synthesis of **I10**. Product obtained 12.50 g (content 75%). HPLC (R_t) = 1.96 min (method H)


Step 3:

5 Intermediate **I19** was prepared starting from **I18** 12.50 g (content 75%) and following the procedures described for the synthesis of **I11**. Product obtained = 2.90 g. HPLC (R_t) = 2.79 min (method H)


Step 4:

10

Intermediate **I20** was prepared starting from **I19** (2.70 g) and following the procedures described for the synthesis of **I12**. Product obtained = 2.45 g (content 85%) used in the next step without any further purification. HPLC (R_t) = 3.00 min (method H)


Step 5:

To an ice-chilled solution of intermediate **120** (2.40 g, content 85%) in THF (60 ml) and water (10 ml) was added lithium hydroxide monohydrate (0.68 g, 16.30 mmol). The temperature was allowed to warm to 25°C and the resulting solution stirred until complete disappearance of the starting material (3 h). The organic solvent was evaporated under reduced pressure, the aqueous residue diluted with water 10 ml), acidified to pH 2.5 with 4N HCl. EtOAc (50 ml) was added, the phases separated, the organic layer washed with brine (20 ml), dried over Na₂SO₄ and evaporated to dryness. The residue was triturated with a 1/1 *n*-hexane/Et₂O mixture, the solid dried at 40 °C under vacuum to afford **121** (1.90 g). HPLC (*R*_t) = 2.65 min (method H)

Step 6:


Intermediate **121** (1.80 g) was dissolved in dry DMF (15 ml), and the resulting solution stirred in a nitrogen atmosphere. HATU (2.46 g) and DIPEA (3.22 ml) were added sequentially and the whole stirred at room temperature for 30 min. 3-cyclopropyl-1-methyl-1H-pyrazole-5-amine (1.02 g) was then added in one portion and the reaction mixture heated at 40 °C overnight. Extra HATU (2.27 g) and 3-cyclopropyl-1-methyl-1H-pyrazole-5-amine (0.68 g) were then added, and the heating maintained for 8h. The reaction mixture was concentrated under reduced pressure, the resulting residue diluted with water (40 ml) and EtOAc (50 ml). The layers were separated and the organic phase washed with 1M HCl (3 x 20 ml), 10% aq. Na₂CO₃ (2 x 20 ml), water (1 x 20 ml) and brine (1 x 25 ml), dried over Na₂SO₄ and evaporated to dryness, affording a solid. The residue was triturated with diisopropyl ether, affording product **122** (1.70 g). HPLC (*R*_t) = 2.85 min (method H)

Step 7:




A suspension of 10% Pd/C (211 mg) in 80% aq. ethanol (5 ml) was added to a stirred solution of intermediate **I22** (1.7 g, content 90%) in EtOH (30 ml). The whole was then stirred under a positive pressure of hydrogen. After 4h extra catalyst (100 mg) was added and the conditions maintained overnight. The reaction mixture was then filtered over Celite®, the cake carefully washed with EtOH (3 x 10 ml) and the filtrate evaporated to dryness under reduced pressure. The residue was dissolved in an 8/2 EtOAc/MeOH mixture (30 ml) and treated with HCl (6 ml of a 4N solution in 1,4 dioxane) to form the corresponding hydrochloride salt. The solvent was evaporated to dryness and the residue triturated with MTBE, the solid filtrated and dried at 40 °C under vacuum to afford the **building block D1** (1.20 g). HPLC (R_t) = 1.60 min (method H)

Synthesis of building block D2



Example **35** (1.0 g) was dissolved in MeOH (20 ml), and Pd/C was added and the mixture was hydrogenated at 50°C and 50 psi for 12 h. After this period, further Pd/C was added and the mixture was again hydrogenated at 50°C and 50 psi for 7 h. The solution was then filtered and concentrated in vacuum to give 0.62 g of **building block D2** which was used without further purification. HPLC: 0.98 min (method D)

Synthesis of building block D3


5 Building block D3 was prepared in analogy to building block D2 starting from example 135.

S.2.5 Synthesis of intermediate I12.2 (Building block for procedure E)

10 see synthesis of **building block A6**, step 4.

S.2.6. Synthesis of building blocks with formula F

15 **Synthesis of building block F1**



20 **Step1:**
2-Chloro-6-methyl-isonicotinamide (1.23 g) and N,N-dimethylformamide dimethyl acetal (1.05 ml) were dissolved in dry DMF (4 ml) under inert atmosphere and the resulting solution was heated to 90°C and stirred for 1 h. The reaction mixture was cooled to room temperature and a solution of hydrazine acetate (3.32 g) in acetic acid (8 ml) was dropwise added. The

mixture was heated to 90°C and stirred for 1 h. The solvent was removed under reduced pressure and the residue partitioned between saturated aqueous solution of NaHCO₃ and EtOAc. The organic phase was separated, washed with brine, dried under Na₂SO₄, filtered and the solvent was evaporated under reduced pressure. The crude was purified by flash column chromatography (10 g Isolute[®] silica gel cartridge; gradient elution: hexane/EtOAc from 1/1 to 3/7) affording 2-Chloro-6-methyl-4-(1H-[1,2,4]triazol-3-yl)-pyridine (0.92 g). GC (R_t) = 13.21 min (method = 3A)

Step 2:


2-Chloro-6-methyl-4-(1H-[1,2,4]triazol-3-yl)-pyridine (120 mg) and 3,4-dihydro-2H-pyran (84 µL) were suspended in dry DCM (3.5 mL) under inert atmosphere and pyridinium p-toluenesulfonate (15.5 mg) was added. The reaction mixture was stirred at room temperature for 18 h. The solution was diluted with DCM, washed with water and brine, dried under Na₂SO₄, filtered and the solvent was evaporated under reduced pressure. The crude was purified by flash column chromatography (10 g Isolute[®] silica gel cartridge; eluent: hexane/EtOAc 3/7) affording product (155 mg). UPLC (R_t) = 1.51 min (method M)

Step 3:

2-Chloro-6-methyl-4-[1-(tetrahydro-pyran-2-yl)-1H-[1,2,4]triazol-3-yl]-pyridine (33.0 mg, 0.12 mmol) and building block **C1** were dissolved in dry dioxane (1.2 mL) under inert atmosphere and palladium(II) acetate (2.12 mg), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (10.9 mg) and caesium carbonate (81.0 mg) were added subsequently. The mixture was heated to 100°C and stirred for 18 h. After cooling to room temperature the reaction mixture was diluted with DCM, washed with water and brine, dried under Na₂SO₄, filtered and the solvent was evaporated under reduced pressure. The crude was purified by flash column chromatography (10 g Isolute[®] silica gel cartridge; gradient elution: DCM/MeOH from 97/3 to 95/5) affording **building block F1** (70 mg). UPLC (R_t) = 1.75 min (method M)


Syntheses of building blocks F2 and F3

Building blocks F2 and F3 were prepared in analogy to **building block F1**



F2

UPLC (R_t) = 1.69 min
method M




F3

UPLC (R_t) = 3.10 min
method M

The chloropyridines were prepared as follows:

C1CCN(C1)c2nc3cc(Cl)nc3n2

2-Chloro-4-[1-(tetrahydro-pyran-2-yl)-1H-pyrazol-4-yl]-pyridine



Chemical reaction scheme showing the synthesis of compound **I25** from **I23** and **I24**:

I23 (3,5-dinitrobenzoic acid) reacts with **I24** (N-methyl-5-nitrobenzamide) to form **I25** (N-methyl-3-amino-5-nitrobenzamide).

The reaction scheme illustrates the synthesis of compound **I25** from **I23** and **I24**. **I23** (3,5-dinitrobenzoic acid) reacts with **I24** (N-methyl-5-nitrobenzamide) to form **I25** (N-methyl-3-amino-5-nitrobenzamide).

Step 1:


To a stirred mixture of 5-nitroisophthalic acid (20.0 g) in DCM was added oxalyl chloride (36 ml, 2M in DCM), followed by DMF (0.5 ml). The mixture was stirred for 2 h at room temperature, before volatiles were removed in vacuum. The residue was suspended in dry toluene and concentrated again to yield crude **I23** (10 g), which was used without further purification in the next step.

Step 2:


I23 (10 g) was dissolved in THF (20 ml) and added dropwise to a solution of methylamine (40 ml, 2M in THF) in THF (80 ml) under stirring at -20°C. After 20 min, the mixture was filtered, and the filtrate concentrated in vacuum. The residue was triturated with diethylether to form a solid that was filtered off and dried to yield intermediate **I24** (3.53 g)

Step 3:


A solution of intermediate **I24** (0.52 g) in MeOH (30 ml) was treated with hydrogen (50 psi) in the presence of Raney-Nickel (90 mg) at room temperature for 14 h. The mixture was then filtered and concentrated in vacuum to afford **I25**. The compound was used without further purification in the next step.
HPLC (R_t) = 0.33 min (method D)

Synthesis of N-(8-amino-4,5-dihydronaphtho[2,1-d]thiazol-2-yl)acetamide for example 10

This intermediate was synthesized according to WO2006040279.


Synthesis of N-(4-Amino-cyclohexyl)-benzenesulfonamide for example 29

This intermediate was synthesized analog to WO2001096295.

Synthesis of 3-(4-Amino-2-methyl-phenyl)-[1,3]oxazinan-2-one for example 30


This intermediate was synthesized analog to WO2005111029.

5 **Synthesis of 5-Amino-2,3-dihydro-indole-1-carboxylic acid methylamide for example 37**



This intermediate was synthesized analog to WO2008113760.

10 **Synthesis of 1-(3-Amino-benzyl)-3-cyclohexyl-urea for example 39**


**Step 1: 1-Cyclohexyl-3-(3-nitro-benzyl)-urea**

15 3-Nitro-benzyl-amine (5 g) was suspended in THF (130 ml) and cyclohexyl isocyanate (7 ml) was added dropwise at 25°C. After stirring for 2h the resulting precipitate was filtered off, washed with diethyl ether and treated with 5% aqueous NH₄OH. After filtering and drying the title compound was isolated (4.5 g, FP 168-169°C)

20 **Step 2: 1-(3-Amino-benzyl)-3-cyclohexyl-urea**


1-Cyclohexyl-3-(3-nitro-benzyl)-urea (4 g) was suspended in ethanol (70 ml) and Raney nickel (2 g) was added. Under stirring aqueous hydrazine solution (2.5 ml, 80%) was added. DMF was added (50ml) and the reaction mixture was stirred for additional 2 h and filtered.
 25 The solution was evaporated i.v. affording the title compound (3 g, R_f = 0.43 DCM/ethanol 19:1).

30 **Synthesis of trans-1-(4-Amino-cyclohexyl)-3-phenyl-urea for example 44**




This intermediate was synthesized analog to WO2005095339.

Synthesis of 1-Methanesulfonyl-pyrrolidin-3-ylamine for example 53




This intermediate was synthesized analog to WO2006118256.

Synthesis of 3-Pyridin-4-ylmethyl-phenylamine for example 66




This intermediate was synthesized analog to WO2004009546.

Synthesis of 1-(5-Chloro-1*H*-benzoimidazol-2-yl)-2-methoxy-ethylamine for example 67



This intermediate was synthesized analog to WO2004056784.

Synthesis of C-(6-Chloro-2-methyl-pyridin-3-yl)-methylamine for example 70



Step 1: 2-Methyl-6-oxo-1,6-dihydro-pyridine-3-carbonitrile

A solution of (E)-3-amino-but-2-enenitrile (100 g) and propynoic acid methyl ester in DMF (200 ml) was stirred at 25°C for 2h and refluxed for additional 1 h. Dowtherm (100 ml) was added and the temperature was raised to 230°C. DMF and MeOH were distilled off. The reaction mixture was cooled down, EtOAc (200 ml) was added and the resulting crystals were filtered off and washed with EtOAc and ether affording the title compound (44.8 g, FP>250°C).


Step 2:**6-Chloro-2-methyl-nicotinonitrile**

A mixture of 2-Methyl-6-oxo-1,6-dihydro-pyridine-3-carbonitrile (44 g) and POCl₃ (250 ml) was refluxed for 3 h. The reaction mixture was dried down in vacuum and ice followed by conc. aqueous ammonia (300 ml) was added. The aqueous layer was extracted with EtOAc. The organic layer was separated, dried and concentrated in vacuum affording the title compound (23.5 g, BP(18mbar) 119-120°C).

Step 3:**C-(6-Chloro-2-methyl-pyridin-3-yl)-methylamine**


To a solution of 6-Chloro-2-methyl-nicotinonitrile (6 g) in THF (120 ml) at 25°C was added NaBH₄ (3 g) followed by the dropwise addition of BF₃ etherate (6 ml). The reaction mixture was stirred for 3h. MeOH (100 ml) and 2N HCl (100 ml) was added and refluxed for 1h. The organic solvent was removed in vacuum and conc. NaOH (40 ml) was added followed by extraction with diethyl ether. The organic layer was separated, dried and concentrated in vacuum followed by filtration over silica gel (EtOAc / MeOH 80:20). The title compound was crystallized from MeOH/HCl (5.5 g)

Synthesis of N-(3-aminophenyl)-N-(2-(azetidin-1-yl)-2-oxoethyl)methanesulfonamide for example 75




This intermediate was synthesized analog according to WO2006114371.

Synthesis of N-(3-aminophenyl)-methylsulfonamido)-N,N-dimethylacetamide for example 76




This intermediate was synthesized analog to WO2006114371.

Synthesis of N-(3-aminophenyl)-cyclopropanesulfonamido)-N,N-dimethylacetamide for example 78




This intermediate was synthesized analog according to WO2006114371.

Synthesis of 2-[(3-Amino-phenyl)-ethanesulfonyl-amino]-*N,N*-dimethyl-acetamide for example 79




This intermediate was synthesized analog to WO2006114371.

Synthesis of ethyl 5-amino-3-methylbenzoate for example 88




2 g acid was dissolved in 22 mL ethanolic HCl and stirred at 60°C for 2 days. The mixture was then concentrated and used without further purification.

Synthesis of ethyl 3-methylaminobenzoate for example 92



2 g acid was dissolved in 22 mL ethanolic HCl and stirred at 60°C for 2 days. The mixture was then concentrated and used without further purification.

Synthesis of 4-(3,5-Dimethyl-pyrazol-1-yl)-pyridin-2-ylamine for example 102




A 10 ml microwave reaction vessel was charged with 4-Chloro-pyridinamine (0.5 g) and 3,5-dimethylpyrazole (1.12 g); hydrochloric acid (37%, 0.5 ml) and dioxane were then added. The reaction mixture was irradiated in a Microwave oven with the following parameters: potency: 150W; temperature: 120°C; time: 180 min.

The reaction mixture was cooled at room temperature and treated with diethyl ether (2 ml) and ethanol (1 ml) and sonicated during 30 min. At end of treatment the mixture was cooled

during 14 h at 4°C. A solid was filtered off and washed with diethyl ether and n-hexane to furnish the product as the hydrochloride salt (450 mg).

5 Synthesis of 5-amino-tetramethylisophthalamide (**I29**) for example 105



Step 1:

To a stirred solution of mono methyl 5-nitroisophthalate (1.0 g,) and DIPEA (1.43 ml) in DCM (30 ml) and DMF (3 ml) was added HATU (1.69 g) at rt. After 10 min, dimethylamine (4.4 ml, 2M solution in THF) was added and stirring was continued for 12 h. Water and sodium bicarbonate were added to the reaction mixture and extracted with DCM. The organic layer was separated and concentrated in vacuum. The residue was further purified by normal phase MPLC (cyclohexane: EtOAc = 80:20 to 30:70) to afford intermediate **I26** (0.91 g) HPLC (R_t) = 1.29 min (method D)

Step 2:

A mixture of **I26** (2.5 g), MeOH (20 ml) and LiOH(aq) (8 wt.%, 5.9 ml) was stirred at 50°C for 1 h. Volatiles were then removed in vacuum and the residue was acidified by treatment with 2 M HCl (aq). The mixture was lyophilized, the remaining solid suspended in water and warmed to 50°C. The resulting suspension was filtered, the solid was washed with water and petrolether and dried in vacuum at 50°C to yield **I27** (1.56 g) that was used without further purification. HPLC (R_t) = 0.27 min (method E)

Step 3:

HATU (707 mg) was added in two portions to a mixture of **I27** (400 mg), DCM (5 ml), DMF (2 ml), diisopropylamine (0.68 ml), and HOBt (227 mg) under stirring at rt. Dimethylamine (2.1 ml, 2M in THF) was added and stirring continued for 48 h. Water and sodium bicarbonate were then added and the resulting mixture was extracted with DCM. The organic layer was separated, dried (Na_2SO_4) and concentrated in vacuum. The residue was further purified by reversed phase HPLC the product containing fractions were combined volatiles removed in



vacuum and the remaining aqueous mixture extracted with DCM. The organic layer was concentrated to afford **I28** (71 mg).

Step 4:

- 5 A solution of intermediate **I28** (71 mg) in MeOH (10 ml) was treated with hydrogen in the presence of Raney-Nickel (10 mg) at rt. The mixture was then filtered and concentrated in vacuum to afford the title compound **I29** (40 mg). The compound was used without further purification in the next step.
HPLC (R_t) = 0.70 min (method D)

10

The reaction partners for example **107** and **106** were synthesized analogously as for example **105**.


Intermediate for the synthesis of example	Structure	HPLC method	R_t [min]
107		D	1.26
106		D	1.26

15


Synthesis of methyl 3-amino-5-(dimethylcarbamoyl)benzoate for example 108

- 20 A solution of intermediate **I26** (0.91 g) in MeOH (40 ml) was treated with hydrogen in the presence of Raney-Nickel (90 mg) at rt. The mixture was then filtered and concentrated in vacuum to afford the title compound (810 mg). The compound was used without further purification in the next step.
HPLC (R_t) = 0.98 min (method D)

- 25 **Synthesis of methyl 3-amino-5-methoxy benzoate for the synthesis of example 111**



- 30 To a stirred suspension of 3-amino-5-methoxy benzoic acid (500 mg) in a mixture of DCM (4.0 ml) and MeOH (2.0 ml) at 0°C was added a solution of trimethylsilyldiazomethan in hexane (3.39 ml, 2M) dropwise. After 15 min, the reaction was concentrated in vacuum. The remaining crude product was purified by reversed phase HPLC to yield the trifluoroacetate salt of title compound (280 mg).
HPLC (R_t) = 1.02 min (method D)

Synthesis of ethyl 3-amino-5-chloro benzoate for the synthesis of example 112**Step 1:**


5-tert-butylisophthalic acid (1.3 g) was dissolved in MeOH (20 ml). The resulting solution was cooled to 0°C and thionylchloride (0.45 ml) was added dropwise under stirring. After complete addition, the mixture was allowed to warm to room temperature and was stirred for additional 7 h. The mixture was then poured on an ice cold sodium bicarbonate solution and extracted repeatedly with EtOAc. The aqueous phase was acidified with HCl and again extracted with EtOAc. The combined organic layers were washed with brine and concentrated in vacuum. The residue was worked up by flash column chromatography on silica gel (DCM:MeOH = 20:1) to yield methyl 5-tert-butylisophthalate (1.3 g).

Step 2:

Under an atmosphere of argon, 5-tert-butylisophthalate (0.5 g) was dissolved in tert.-butanol (20 ml). To the resulting solution, triethylamine (0.35 ml) and diphenyl phosphoreazidate (0.56 ml) were added subsequently. The mixture was then heated to reflux for 12 h, cooled to room temperature and concentrated in vacuum. The residue was taken up in EtOAc, and subsequently washed with dilute aqueous citric acid, saturated sodium bicarbonate solution and brine. The organic layer was concentrated in vacuum and the residue was worked up by flash chromatography (petrol ether:EtOAc = 3:1) to yield methyl 3-(tert.-butoxycarbonylamino)-5-tert-butylbenzoate (0.61 g).

Step 3:

Methyl 3-(tert.-butoxycarbonylamino)-5-tert-butylbenzoate (0.42 g) was dissolved in DCM (5 ml). To the resulting solution, TFA (1 ml) was added, the mixture was stirred for 30 min at room temperature and then concentrated in vacuum. The residue was dissolved in DCM and extracted with aqueous sodium bicarbonate solution. The organic layer was separated, dried over sodium sulphate and concentrated in vacuum to yield ethyl 3-amino-5-tert-butylbenzoate (0.28 g), which was used for the synthesis of example 112 without further purification.


Synthesis of methyl 3-amino-5-chloro benzoate for the synthesis of example 113

A solution of methyl 3-chloro-5-nitrobenzoate (300 mg) in THF (50 ml) was treated with hydrogen (3 bar) in the presence of Raney-Nickel (10 mg) at rt. The mixture was then filtered

and concentrated in vacuum to afford the title compound (204 mg). The compound was used without further purification in the next step.
HPLC (R_t) = 1.36 min (method D)

5

Synthesis of 3-Methyl-5-(1-methyl-1H-tetrazol-5-yl)-phenylamine for example 114



10

Step 1:

3-Chlorocarbonyl-5-nitro-benzoic acid ethyl ester

To a solution of 5-nitroisophthalic acid monoethyl ester (80 g) in DCM (700 ml) at 25°C was added 2M oxalyl chloride solution in DCM (63 ml), and DMF (5 ml). The reaction was stirred for 2 h and then concentrated in vacuum. The residue was triturated with toluene and again concentrated in vacuum leaving the title compound (60 g). R_t = 1.40 min for derivatised methylester (method D).

15

Step 2:

N-Methyl-5-nitro-isophthalamic acid ethyl ester

To a 2M solution of methylamine in THF (256 ml) at -20°C was added a pre-cooled solution of 3-chlorocarbonyl-5-nitro-benzoic acid ethyl ester (60 g) in THF (500 ml) dropwise. This was stirred for 20 min and then filtered. The filtrate was concentrated in vacuum and the solid recrystallised from diethyl ether to give the title compound (22.5 g) as a solid. An oxalyl chloride solution in DCM (63 ml, 2M), was added with DMF (5 ml). The reaction was stirred for 2 h and then concentrated in vacuum. The residue was triturated with toluene and again concentrated in vacuum leaving the title compound (60g). R_t = 1.53min (method D).

25

Step 3:

3-(1-Methyl-1H-tetrazol-5-yl)-5-nitro-benzoic acid ethyl ester

To a 2M solution of N-methyl-5-nitro-isophthalamic acid ethyl ester (10 g) in dichloroethane (200 ml) at -20°C was added pre-cooled triflic anhydride (10 ml) dropwise. This was stirred for 30 min and then sodium azide (4.6 g) was added. This was allowed to warm to room temperature and stirred overnight. The reaction was neutralised with 5% NaHCO₃(aq) solution and the organic and aqueous layers were separated and the organic layer dried and concentrated in vacuum. Flash chromatography (100:0 to 50:50 gradient cyclohexane:EtOAc) afforded the title compound (6.5 g). R_t = 1.52 min (method D).

35

Step 4:

3-Amino-5-(1-methyl-1H-tetrazol-5-yl)-benzoic acid ethyl ester

A solution of 3-(1-Methyl-1H-tetrazol-5-yl)-5-nitro-benzoic acid ethyl ester (6.5 g) in EtOAc:ethanol (1:1, 320 ml) was hydrogenated at 20°C and 50 psi with 10% Pd/C (750 mg). After reaction completion, the mixture was filtered and the filtrate concentrated in vacuum to give the title compound (4.2 g). R_t = 1.21 min (method D).

40

45

Step 5:

[3-Amino-5-(1-methyl-1H-tetrazol-5-yl)-phenyl]-methanol

To a solution of 3-Amino-5-(1-methyl-1H-tetrazol-5-yl)-benzoic acid ethyl ester (250 mg) in THF (5 ml) at 0°C was added pre-cooled 1M LiAlH₄ solution in THF(2 ml) dropwise. This was stirred for 3 h. The reaction mixture was poured onto ice-water and extracted into EtOAc.

The organic layer was separated, dried and concentrated in vacuum. Flash chromatography (100:0 to 95:5 gradient DCM:MeOH) afforded the title compound (70 mg).
 R_f = 0.19 (DCM:MeOH 95:5).

5 **Step 6:**


3-Methyl-5-(1-methyl-1H-tetrazol-5-yl)-phenylamine

A solution of [3-Amino-5-(1-methyl-1H-tetrazol-5-yl)-phenyl]-methanol (450 mg) in MeOH (20 ml) with a few drops of conc. HCl (aq) was hydrogenated on a H-cube apparatus at 30°C, and 10bar with a flow rate of 1mL/min using a Pd/C cartridge. After reaction completion, the mixture was concentrated in vacuum and the residue basified with a dilute K_2CO_3 (aq). This was extracted with EtOAc and the organic layer was separated, dried and concentrated in vacuum. The residue was purified by flash chromatography (98:2 DCM:MeOH) to give the title compound (320 mg).

R_f = 0.46 (95:5 DCM:MeOH)


S3.2 Synthesis of aromatic-chlorides or -bromides for procedure C

Synthesis of 2-Chloro-4-(3,5-dimethyl-isoxazol-4-yl)-pyridine for example 187



The title compound was prepared from 4-bromo-3,5-dimethylisoxazole generally according to the synthetic procedure described for 2-Chloro-4-[1-(tetrahydro-pyran-2-yl)-1H-pyrazol-4-yl]-pyridine. UPLC (R_t) = 0.81 min (method N)

Synthesis of 2-Chloro-6-methyl-4-(5-methyl-[1,2,4]oxadiazol-3-yl)-pyridine for example 188



To a suspension of 2-chloro-6-methyl-isonicotinamide (0.50 g, 2.93 mmol) in dry DCM (10 ml), dry pyridine (0.26 ml) was added and the mixture was cooled to 0°C under inert atmosphere. Trifluoroacetic anhydride (0.45 ml) was added dropwise under stirring, the cooling bath was removed and the reaction mixture was stirred for 4 h. After this period other portions of pyridine (0.074 ml) and trifluoroacetic anhydride (0.12 ml) were added and the mixture stirred at room temperature for further 3 h.


The reaction was quenched with aqueous solution of $NaHCO_3$, the phases were separated and the aqueous layer extracted with DCM. The combined organic extracts were washed with brine, dried under Na_2SO_4 , filtered and the solvent was evaporated under reduced

pressure. The crude product was obtained (0.378 g) and used in next step without further purification. UPLC (R_t) = 1.65 min (method M)

2-Chloro-6-methyl-isonicotinonitrile (378 mg) was dissolved in ethanol (4 ml) and hydroxylamine hydrochloride (379 mg) was added followed by a solution of sodium carbonate (577 mg) in water (8 ml). The reaction mixture is heated to 75°C and stirred for 18 h. Ethanol was evaporated under reduced pressure and the aqueous residue extracted with EtOAc. The combined extracts were washed with water and brine, dried under Na_2SO_4 , filtered and the solvent was evaporated under reduced pressure. The crude product was obtained (401 mg) and used in next step without further purification. UPLC (R_t) = 0.67 min (method M)


2-Chloro-N-hydroxy-6-methyl-isonicotinamidine (120 mg) was suspended in trimethylorthoacetate (3 ml) under inert atmosphere and acetic acid (0.3 ml) was added. The mixture was heated to 100°C and stirred for 2 h. After cooling to room temperature the solvent was evaporated under reduced pressure and the residue partitioned between EtOAc and aqueous solution of NaHCO_3 . The organic layer was separated, washed with water and brine, dried under Na_2SO_4 , filtered and the solvent was evaporated under reduced pressure. The crude was purified by flash column chromatography (10 g Isolute[®] silica gel cartridge; gradient elution: hexane/EtOAc from 10/0 to 9/1) affording the title compound (107 mg). HPLC (R_t) = 3.17 min (method L).

Synthesis of methyl 2,6 dibromo-isonicotinate for the synthesis of example 194



In a three necked round bottom flask equipped with a mechanic stirrer and gas outlet solid citrazinic acid (15 g) and phosphorous oxybromide (45 g) were thoroughly mixed and heated to 140°C for 12 to 14 h. The resulting mixture was cooled to 0°C before MeOH (100 ml) was added carefully under vigorous stirring. The resulting mixture was then poured into a cooled (0°C) aqueous sodium carbonate solution (1M, 500ml), before Chloroform (500 ml) was added. The biphasic mixture was filtered through a paper filter, before the organic layer was separated. After filtering through charcoal, the solution was concentrated in vacuum. The residue was purified by MPLC (DCM:MeOH = 100:3 to 100:6) to yield methyl 2,6 dibromo-isonicotinate (13.7 g). HPLC (R_t) = 1.62 (method D)


Synthesis of methyl 2-bromo-6-methoxyisonicotinate for example 195



A solution of sodium methoxide (375 mg) and methyl 2,6 dibromo-isonicotinate (1.0 g) in MeOH (20 ml) was heated in a microwave oven at 130°C for 30 min. Then additional sodium methoxide (281mg) was added and heating continued for additional 15 min at 130°C. Concentrated sulfuric acid (1.86 ml) was then added to the reaction mixture and the resulting suspension was heated for 4 h at 80-85°C.

After cooling to room temperature, the mixture was poured in an ice cold aqueous sodium carbonate solution (100 mL) and extracted with DCM (100 ml). The organic layer was separated, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by MPLC (DCM:MeOH = 100:3 to 100:5) to yield 710 mg of a 70:30 mixture of desired product (497 mg) and the corresponding trimethyl cirtrazinic acid (213 mg). The mixture was used for subsequent transformations without further purification.
HPLC (R_t) = 1.66 min (method D)

Synthesis of 2-Chloro-6-methyl-4-(1-methyl-1H-tetrazol-5-yl)-pyridine for example 191



Step 1:

2-Chloro-6,N,N-trimethyl-isonicotinamide

To a solution of 2-Chloro-6-methyl-isonicotinic acid (1 g) in DMF (200 ml) at 25°C was added DIPEA (2.9 mL) and TBTU (1.8 ml). After 5 min, dimethylamine (0.75 g) was added dropwise. This was stirred overnight. The reaction was added to water and extracted with EtOAc. The organic layer was separated, dried and concentrated in vacuum. Flash chromatography (9:1 to 2:8 gradient cyclohexane:EtOAc) afforded the title compound (320mg). R_f = 0.56 (1:1 cyclohexane:EtOAc)


Step 2:

2-Chloro-6-methyl-4-(1-methyl-1H-tetrazol-5-yl)-pyridine


To a solution of 2-Chloro-6,N,N-trimethyl-isonicotinamide (2.3 g) in acetonitrile (30 ml) at -15°C was added DCM (20 ml) pre-cooled triflic anhydride (3 mL) dropwise. This was stirred for 10 min at -10°C and then sodium azide (4.6 g) was added. This was allowed to warm to room temperature and after 2h, the reaction was neutralised with a cold 5% NaHCO₃(aq) solution and extracted with EtOAc. The organic layer was separated, dried and concentrated in vacuum. Flash chromatography (100:0 to 40:60 gradient cyclohexane:EtOAc) afforded the title compound (260 mg). R_t = 1.21 min (method D).

S.4. Synthesis of examples according to procedure A to G

S.4.1. Synthesis of examples according to procedure A (examples 1 – 131)

Amide formation method A:

- 5 To a stirred solution of building block **A** (1 eq) in DMF, NMP, DCM or chloroform at room temperature is added DIPEA or TEA (2-8 eq), HATU or TBTU (1-4 eq) and HOBt (0-2 eq). After 20-45 min, amine reaction partner (1.2-2 eq) is added and the mixture stirred for an additional 40-120 min. Water and TFA is then added and the mixture is separated via reversed phase HPLC to provide the desired amides. The reaction mixture can be filtered
 10 over basic ALOX or extracted with DCM after addition of aqueous NaHCO₃ solution followed by concentrating the organic fractions in vacuum before the HPLC purification.


Synthesis of example 1:

- 15 Building block **A1** (60 mg) was dissolved in NMP (0.5 ml) and DIPEA (0.15 ml). HATU (120 mg) was added and the solution was stirred for 45 min at rt. 3-(1-methyl-1H-tetrazol-5-yl)aniline (34 mg) was added and the mixture was stirred for a further 1 h. A few drops of water and TFA were added and the mixture was separated via reversed phase HPLC. The
 20 desired fractions were collected, combined and lyophilised to yield example **1** (18 mg) as the TFA salt. HPLC: 1.3 min (method D)

Amide formation method B:


- 25 To a stirred solution of building block **A** (1 eq) in DCM at room temperature is added oxalylchloride (2-5 eq, neat or 2M solution in DCM) and the mixture stirred for 0.5-2 h and then concentrated in vacuum. Pyridine, DMAP (0-0.1 eq) and amine reaction partner (2-3 eq) is added and the mixture stirred for 1-2 h, then concentrated, dissolved in ACN, water and TFA and separated via reversed phase HPLC to give the desired amides.
- 30

Synthesis of example 2:




Building block **A1** (80 mg) was dissolved in DCM (2 ml), and a solution of oxalylchloride in DCM (0.2 ml, 2M) was added and the mixture was stirred at 25°C for 2 h and then concentrated in vacuum. Pyridine (1 ml) and 3-amino-N-methyl-5-(trifluoromethyl)benzamide hydrochloride (88 mg) is added and the mixture was further stirred for 2 h, then concentrated, dissolved in acetonitrile, water and TFA and separated via reversed phase HPLC to give of example **2** (45 mg) as the TFA salt. HPLC: 1.37 min (method D)

















Synthesis of example 3:







































Building block **A1** (80 mg) was dissolved in DCM (2 ml), oxalylchloride (0.07 ml) was added and the mixture was stirred at room temperature for 0.75 h and then concentrated in vacuum. Pyridine (1 ml), 4-dimethylaminopyridine (2 mg) and 5-Amino-1,3,4-thiadiazole-2-carboxylic acid ethyl ester (60 mg) were added and the mixture was further stirred for 1 h, then concentrated, dissolved in acetonitrile and separated via reversed phase HPLC to give example **3** (17 mg). HPLC: 1.34 min (method D)

















The following examples **4 -115** were synthesized in analogy to example **1-3** from building blocks **A1** to **A4**.







































#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
4			H, H	<i>R</i>	1.26 (D)	A	A1
5			H, H	<i>R</i>	1.36 (D)	A	A1
6			H, H	<i>R</i>	1.35 (D)	A	A3
7			H, H	<i>R</i>	1.21 (D)	A	A1
8			H, H	<i>R</i>	1.29 (D)	A	A1
9			H, H	<i>R</i>	1.35 (D)	A	A1
10			H, H	<i>R</i>	1.34 (D)	A	A1
11			H, H	<i>R</i>	1.30 (D)	A	A1



















#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
12			H, H	S	1.20 (C)	A	A1
13			H, H	R	1.39 (D)	A	A1
14			H, H	R	1.36 (D)	A	A1
15			H, H	R	1.39 (D)	A	A1
16			H, H	R	1.26 (D)	B	A1
17			H, H	R	1.55 (B)	A	A1
18			H, H	R	1.61 (B)	A	A1
19			H, H	R	1.46 (B)	A	A1
20			H, H	R	1.74 (B)	A	A1



















#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
21			H, H	R	1.61 (B)	A	A1
22			H, H	R	1.80 (B)	A	A1
23			H, H	S	1.51 (B)	A	A1
24			H, H	R	1.53 (B)	A	A1
25			H, H	R	1.25	A	A3
26			H, H	R	1.59 (B)	A	A1
27			H, H	R	1.42 (B)	A	A1
28			H, H	R	1.62 (B)	A	A1
29			H, H	R	1.58 (B)	A	A1

















#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
30			H, H	<i>R</i>	1.36 (B)	A	A1
31			H, H	<i>R</i>	1.50 (B)	A	A1
32			H, H	<i>R</i>	1.43 (B)	A	A1
33			H, H	<i>R</i>	1.54 (B)	A	A1
34			H, H	<i>R</i>	1.58 (B)	A	A1
35			H, H	<i>S</i>	1.90 (B)	A	A2
36			H, H	<i>R</i>	1.30 (B)	A	A1
37			H, H	<i>R</i>	1.52 (B)	A	A1

#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
38			H, H	S	1.71 (B)	A	A1
39			H, H	R	1.72 (B)	A	A1
40			H, H	S	1.49 (B)	A	A2
41			H, H	R	1.43 (B)	A	A1
42			H, H	S	1.64 (B)	A	A1
43			H, H	R	1.72 (B)	A	A1
44			H, H	R	1.59 (B)	A	A1
45			H, H	S	1.50 (B)	A	A2
46			H, H	S	1.63 (B)	A	A2
47			H, H	S	1.68 (B)	A	A2

















#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
48			H, H	S	1.66 (B)	A	A2
49			H, H	S	1.68 (B)	A	A2
50			H, H	S	1.59 (B)	A	A2
51			H, H	S	1.63 (B)	A	A2
52			H, H	S	1.65 (B)	A	A2
53			H, H	S	1.51 (B)	A	A2
54			H, H	R	1.75 (B)	A	A2
55			H, H	S	1.74 (B)	A	A2



















#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
56			H, H	S	1.78 (B)	A	A2
57			H, H	S	1.74 (B)	A	A2
58			H, H	S	1.72 (B)	A	A2
59			H, H	R	1.72 (B)	A	A2
60			H, H	S	1.39 (B)	A	A2
61			H, H	S	1.78 (B)	A	A2
62			H, H	S	1.72 (B)	A	A2
63			H, H	R	1.64 (B)	A	A2
64			H, H	R	1.40 (B)	A	A2













#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
65			H, H	R	1.40 (B)	A	A2
66			H, H	S	1.42 (B)	A	A2
67			H, H	S	1.56;1.64 (B)	A	A2
68			H, H	R	1.61 (B)	A	A2
69			H, H	S	1.44 (B)	A	A2
70			H, H	S	1.62 (B)	A	A2
71			H, H	S	1.72 (B)	A	A2
72			H, H	S	1.68 (B)	A	A2
73			H, H	S	1.70 (B)	A	A2











#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
74			H, H	R	1.68 (B)	A	A2
75				R	1.31 (E)	A	A4
76				R	1.31 (E)	A	A4
77				R	1.31 (E)	A	A4
78				R	1.33 (E)	A	A4
79				R	1.34 (E)	A	A4

#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
80				R	1.38 (E)	A	A4
81				R	1.22 (E)	A	A4
82				R	1.24; 1.29 (E)	A	A4
83				R	1.45 (E)	A	A4
84				R	1.23 (E)	A	A4
85				R	1.33 (E)	A	A4
86				R	1.37 (E)	A	A4
87			H, H	R	1.35 (D)	B	A1

#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
88			H, H	<i>R</i>	1.48 (D)	A	A1
89			H, H	<i>R</i>	1.37 (D)	B	A1
90			H, H	<i>R</i>	1.40 (D)	A	A1
91			H, H	<i>R</i>	1.67 (B)	A	A1
92			H, H	<i>R</i>	1.39 (D)	A	A1
93			H, H	<i>R</i>	1.26 (D)	B	A1
94			H, H	<i>R</i>	1.87 (A)	A	A1
95			H, H	<i>R</i>	1.91 (A)	A	A1

#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
96			H, H	<i>R</i>	1.66 (B)	A	A2
97			H, H	<i>S</i>	1.37 (B)	A	A2
98			H, H	<i>R</i>	1.36 (D)	B	A2
99			H, H	<i>R</i>	1.31 (D)	B	A1
100			H, H	<i>R</i>	2.07 (A)	A	A1
101			H, H	<i>R</i>	9.22 (O)	A	A1
102			H, H	<i>R</i>	9.75 (O)	A	A1
103			H, H	<i>R</i>	1.26 (D)	B	A1
104			H, H	<i>R</i>	1.35 (D)	A	A1


#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
105			H, H	<i>R</i>	1.23 (D)	B	A1
106			H, H	<i>R</i>	1.23 (D)	A	A1
107			H, H	<i>R</i>	1.20 (D)	A	A1
108			H, H	<i>R</i>	1.21 (D)	B	A1
109			H, H	<i>R</i>	1.33 (D)	A	A1
110			H, H	<i>R</i>	1.47 (D)	B	A1

#	R ²	R ¹	R ³ , R ⁴	*	R _t [min] (method)	Amide formation Method	Building block
111			H, H	<i>R</i>	1.40 (D)	A	A1
112			H, H	<i>R</i>	1.53 (D)	A	A1
113			H, H	<i>R</i>	1.48 (D)	A	A1
114			H, H	<i>R</i>	1.26 (E)	A	A1
115			H, H	<i>R</i>	1.26 (D)	A	A3

Synthesis of examples from building blocks A5 – A8

5


Example 116



To an ice-chilled stirred solution of building block **A5** (0.11 g), 3-amino-benzonitrile (0.08 g), DIPEA (0.26 ml) and DMAP (0.01 g) in dry DMF (1 ml), was added tetramethyl
 5 fluoroformamidinium hexafluorophosphate (0.1 mg). After 5 min the cooling bath was removed and the reaction mixture stirred at room temperature until complete disappearance of the starting acid.


The solvent and the volatiles were removed under reduced pressure, the thick oily residue was purified by preparative LC-MS, affording example 116 (as TFA salt) (0.02 mg). HPLC
 10 (R_t) = 8.38 min (method O)

The following examples are synthesized in analogy to the preparation of example 116:




Example	R ¹	R _t [min] (method)	Building block
117		8.27 (P)	A5
118		6.90 (P)	A5
119		8.38 (O)	A5

Example 120




Building block **A7** (0.07 g) was dissolved in dry DMF (2 ml); HATU (0.09 g) and DIPEA (0.15 mL) were added in sequence and the whole stirred at room temperature for 15 min. Then 3-pyridin-4-yl-phenylamine was added (0.04 g) and the resulting yellow solution stirred at 40 °C until complete disappearance of the starting acid. The solvent and the volatiles were removed under reduced pressure; the oily residue was purified by preparative LC-MS,
 20 affording the title compound (as TFA salt) (62 mg). HPLC (R_t) = 10.02 min, Method = F

The following examples are synthesized in analogy to the preparation of example **120**




#	R ¹	*	R _t [min] (method)	Building block
121		<i>R</i>	7.33 (P)	A7
122		<i>R</i>	6.90 (P)	A7
123		<i>R</i>	7.08 (P)	A7
124		<i>S</i>	3.05 (G)	A8
125		<i>S</i>	2.88 (G)	A8
126		<i>S</i>	2.43 (G)	A8
127		<i>S</i>	8.27 (F)	A8
128		<i>S</i>	2.72 (G)	A8
129		<i>S</i>	2.49 (G)	A8
130		<i>R</i>	2.67 (G)	A6
131		<i>R</i>	2.49 (G)	A6

S.4.2. Synthesis of examples according to procedure B (examples 132 – 191)5 **Synthesis of example 132**












10 To a stirred solution of Building block **B1** (120 mg) in DMF (2 ml) at room temperature, was added DIPEA (0.2 ml), HATU (175 mg). After 20 min, pyrrolidine (49 mg) was added and the mixture stirred for a further 1.5 h. A few drops of water and TFA were added and the mixture was separated via reversed phase HPLC. The desired fractions were collected, combined and lyophilised to yield example **132** (30 mg) as the TFA salt . HPLC: $R_t = 1.29$ min (method D)

15 The following examples have been synthesized from building block **B1** to **B18** in analogy to example 132.




















#	R^1	X	*	R_t [min] (method)	Building block
133		Cl	R	1.34 (D)	B4
134		Cl	R	1.34 (D)	B4
135		Cl	R	1.26 (D)	B1
136		Cl	R	1.35 (D)	B4







#	R ¹	X	*	R _t [min] (method)	Building block
137		Cl	<i>R</i>	1.34 (D)	B4
138		Cl	<i>R</i>	1.37 (D)	B4
139		Cl	<i>R</i>	1.32 (D)	B3
140		Cl	<i>R</i>	1.26 (D)	B1
141		Cl	<i>R</i>	1.28 (D)	B4
142		Cl	<i>R</i>	1.29 (D)	B2
143		Cl	<i>R</i>	1.36 (D)	B4
144		Cl	<i>R</i>	1.32 (D)	B4
145		Cl	<i>R</i>	1.32 (D)	B4
146		Cl	<i>R</i>	1.31 (D)	B4
147		Cl	<i>R</i>	1.24 (D)	B3


#	R ¹	X	*	R _t [min] (method)	Building block
148		Cl	<i>R</i>	1.29 (D)	B4
149		Cl	<i>R</i>	1.38 (D)	B4
150		Cl	<i>R</i>	1.26 (D)	B1
151		Cl	<i>R</i>	1.25 (D)	B4
152		Cl	<i>R</i>	1.30 (D)	B4
153		Cl	<i>R</i>	1.29 (D)	B4
154		Cl	<i>R</i>	1.34 (D)	B4
155		Cl	<i>R</i>	1.53 (B)	B5
156		Cl	<i>R</i>	1.25 (D)	B4
157		Cl	<i>R</i>	1.35 (D)	B4
158		Cl	<i>R</i>	1.39 (D)	B4

#	R ¹	X	*	R _t [min] (method)	Building block
159		Cl	<i>R</i>	1.22 (D)	B4
160		Cl	<i>R</i>	1.24 (D)	B3
161		Cl	<i>R</i>	1.29 (D)	B4
162		Cl	<i>R</i>	1.28 (D)	B4
163		Cl	<i>R</i>	1.41 (D)	B4
164		Cl	<i>R</i>	1.53 (B)	B5
165		Cl	<i>R</i>	1.59 (B)	B1
166		Cl	<i>R</i>	1.26 (D)	B6
167		Cl	<i>R</i>	1.22 (E)	B2
168		Cl	<i>R</i>	1.26 (E)	B8

#	R ¹	X	*	R _t [min] (method)	Building block
169		Cl	<i>R</i>	1.24 (E)	B8
170		Cl	<i>R</i>	1.25 (E)	B10
171		Cl	<i>R</i>	1.35 (E)	B10
172		Cl	<i>R</i>	1.30 (E)	B10
173		Cl	<i>R</i>	1.22 (E)	B9
174		Cl	<i>R</i>	1.19 (E)	B9
175		Cl	<i>R</i>	1.14 (E)	B2
176		Cl	<i>R</i>	1.24 (E)	B7
177		Cl	<i>R</i>	1.20 (E)	B7


#	R ¹	X	*	R _t [min] (method)	Building block
178		Cl	<i>R</i>	1.12 (E)	B11
179		Cl	<i>R</i>	1.18 (E)	B11
180		Cl	<i>R</i>	1.18 (E)	B11
181		Cl	<i>R</i>	1.34 (D)	B15
182		Cl	<i>R</i>	1.36 (D)	B13
183		Cl	<i>R</i>	1.29 (D)	B14
184		Cl	<i>R</i>	1.31 (D)	B12
185		Cl	<i>R</i>	1.17 (E)	B16

#	R ¹	X	*	R _t [min] (method)	Building block
186		Cl	<i>R</i>	1.24 (D)	B16
187		Cl	<i>R</i>	1.26 (D)	B16
188		Br	<i>R</i>	1.28 (D)	B18
189		Cl	<i>R</i>	1.34 (D)	B17
190		Cl	<i>R</i>	1.50 (B)	B19
191		Cl	<i>R</i>	1.34 (D)	B20


S.4.3. Synthesis of examples according to procedure C (examples 192 – 196)**Example 192**

2-Chloro-4-(3,5-dimethyl-isoxazol-4-yl)-pyridine (25 mg) and building block **C1** (25mg) were dissolved in dry dioxane (1.2 ml) under inert atmosphere and palladium(II) acetate (2 mg), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (11 mg) and caesium carbonate (81mg) were added subsequently. The mixture was heated to 100°C and stirred for 18 h. After cooling to room temperature the reaction mixture was diluted with DCM, washed with water and brine, dried under Na₂SO₄, filtered and the solvent was evaporated under reduced pressure. The crude was purified by flash column chromatography (10 g Isolute[®] silica gel cartridge; gradient elution: DCM/MeOH from 97/3 to 95/5) affording the title compound (49mg). HPLC (R_t) = 9.47 min (method O)

The following examples are synthesized in analogy to the preparation of example **192**




#	R ¹	*	R _t [min] (method)	Building block
193		R	3.37 (L)	C1
194		R	1.45 (D)	C1
195		R	1.46 (D)	C1

#	R ¹	*	R _t [min] (method)	Building block
196		<i>R</i>	1.23 (E)	C1


S.4.4. Synthesis of examples according to procedure D (examples 197 – 208)





5 Example 197




Building block **D1** (0.08 g) and 3-fluoro-4-trifluoromethoxy benzaldehyde (0.05 g) were dissolved in 25% MeOH in DCM (4 ml) and the resulting solution stirred at room temperature for 30 min. Sodium cyanoborohydride (0.02 g) was then added in one portion and the whole allowed to react overnight. The solvent was evaporated under reduced pressure, 10% aq. K₂CO₃ (3 ml) and DCM (5 ml) were added, the layers separated, the organic phase dried over Na₂SO₄ and evaporated to dryness. The resulting crude was then purified by preparative LC-MS to afford the target compound as a TFA salt (0.03 g). HPLC (R_t) = 7.01 min (method P)

The following examples are synthesized in analogy to the preparation of example **197**




#	R ¹	R ²	*	R _t [min] (method)	Building block
198			<i>R</i>	2.60 (G)	D1
199			<i>R</i>	9.30 (O)	D1

Synthesis of example 200






D2 (0.05 g) was dissolved in EtOH (1 ml) and of 4-bromobenzaldehyde (0.081 g), HOAc (0.026 g) and a small amount of molecular sieve 4 Å was added and the suspension stirred at 25°C overnight. Sodium triacetoxyborohydride (0.154 g) was added and again the mixture stirred for 1 h, concentrated and suspended in MeOH, water and TFA. After filtration the mixture was separated via reversed phase HPLC and fractions containing the product collected and lyophilised to provide example **200** (0.049 g). HPLC: 1.21 min (method D)

The following examples **201-208** were synthesized in analogy to example **200** from building block **D2** or **D3**.




#	R ²	*	R _t [min] (method)	Building block
201		<i>R</i>	1.20 (D)	D2
202		<i>R</i>	1.14 (D)	D2
203		<i>R</i>	1.21 (D)	D2
204		<i>R</i>	1.23 (D)	D2
205		<i>R</i>	1.23 (D)	D2

206		<i>R</i>	1.18 (D)	D2
207		<i>R</i>	1.18 (D)	D2
208		<i>R</i>	1.29 (D)	D3

S.4.5. Synthesis of examples according to procedure E

5

Example 209




I12.2

10 Intermediate **I12.2**, (0.20 g), 3-cyclopropyl-1-methyl-1H-pyrazole-5-amine (0.14 g), potassium *t*-butoxide (0.10 ml of a 1M THF solution) and dioxane (2 ml) were transferred to a microwave vial and heated under microwave irradiation under the following conditions: max power 150 W, T 150° C, 15 min; after the first cycle 0.05 mL of base were added and two more cycles of irradiation performed under the same conditions. The reaction mixture was
 15 evaporated to dryness, and the oily residue was purified by preparative LC-MS affording example 208 (as TFA salt) (0.09 g). HPLC (R_t) = 7.12 min (method P)

S.4.6. Synthesis of examples according to procedure F (examples 210-212)


20

Example 210



Building block **F1** (70 mg) was dissolved in ethanol (5 ml) and conc. hydrochloric acid (1.5 ml) was added. The solution was stirred at room temperature for 4 h. Ethanol was removed under reduced pressure and the residue partitioned between saturated aqueous solution of NaHCO₃ and EtOAc. The layers were separated and the aqueous phase extracted with EtOAc. The combined organic extracts were washed with brine, dried under Na₂SO₄, filtered and the solvent was evaporated under reduced pressure. The crude was purified by flash column chromatography (5 g Isolute[®] silica gel cartridge; gradient elution: DCM/MeOH from 97/3 to 95/5) affording the title compound (55 mg). HPLC (R_t) = 8.08 min (method O)


Example **211-212** was synthesized in analogy to example **210**.



#	R ¹	*	R _t [min] (method)	Building block
210		<i>R</i>	8.50(O)	F2
211		<i>R</i>	2.37(M)	F3


S.4.7. Synthesis of examples according to procedure G (examples 213-216)

Example 213



To a solution of example **181** (36 mg) in dioxane (3 ml) was added dppfPdCl₂ (2.3 mg) and butylzinc bromide (0.137 ml of a 0.5 M solution in THF) under an atmosphere of argon. The solution was stirred for 45 min at room temperature before additional dppfPdCl₂ (2 mg) and butylzinc bromide (0.07 ml of a 0.5 M solution in THF) was added and the solution was stirred for additional 20 min at rt. The reaction was quenched with a few drops of water, the resulting mixture was diluted with dioxane and filtered. The filtrate was concentrated and the crude product purified by reversed phase HPLC affording the title compound (19 mg). HPLC: 1.41 min (method D).

The following examples were synthesized in analogy to example 213



#	R ¹	*	Rt [min] (method)	Building block
214		R	1.34 (D)	example 181
215		R	1.48 (D)	example 194
216		R	1.42 (D)	example 194

5

S.5.1 Chromatographic methods (HPLC and UPLC methods)

10 Method A

Waters ZQ2000 MS, Alliance 2790 HPLC, Waters 2996 DAD (210-500 nm), Waters 2700 Autosampler

Mobile Phases:

A: Water with 0.10% TFA

15 B: Acetonitrile with 0.10% TFA

	Time in min	%A	%B	Flowrate in ml/min
	0.00	95	5	1.00
	0.10	95	5	1.00
20	3.60	2	98	1.00
	6.50	2	98	1.00
	7.00	95	5	1.00

Stationary phase: Waters, Sunfire C18, 3,5 µm, 4,6 X 50 mm.

25 Column temp: constant at 40°C.

Method B

Waters ZQ2000 MS, HP1100 HPLC + DAD (210-500nm), Gilson 215 Autosampler.

Mobile Phases:

A: Water with 0.10% TFA

B: Acetonitrile with 0.10% TFA

5	Time in min	%A	%B	Flowrate in ml/min
	0.00	95	5	1.50
	2.00	0	100	1.50
	2.50	0	100	1.5
	2.60	95	5	1.50

10

Stationary phase: Waters, Sunfire C18, 3,5 µm, 4,6 X 50 mm.

Column temp: constant at 40°C.

Method C

15 Alliance 2690/2695 HPLC, waters 996/2996 DAD (210-400 nm)

mobile phases:

A: water with 0.10% TFA

B: acetonitrile with 0.10% TFA

20	time in min	%A	%B	flow rate in ml/min
	0.00	95	5	2.50
	0.20	95	5	2.50
	1.50	2	98	2.50
	1.70	2	98	2.50
25	1.90	95	5	2.50
	2.20	95	5	2.50

Stationary phase Merck Chromolith™ Flash RP-18e, 4.6 mm x 25 mm (at 25°C).

30 **Method D**

Alliance 2690/2695 HPLC, waters 996/2996 DAD (210-400 nm)

mobile phases:

A: water with 0.10% TFA

B: acetonitrile with 0.10% TFA

35

	time in min	%A	%B	flow rate in ml/min
	0.00	95	5	2.80
	0.30	95	5	2.80
	1.60	2	98	2.80
40	1.90	2	98	2.80
	2.00	95	5	2.50

Stationary phase Merck Chromolith™ Flash RP-18e, 3 mm x 100 mm (at 25°C).

45 **Method E**

Waters ZQ MS, Alliance 2690/2695 HPLC, Waters 996/2996 Diode array detector

Mobile Phases:

A: Water with 0.10% NH3

B: Acetonitrile with 0.10% NH3

50

	Time in min	%A	%B	Flowrate in ml/min
	0.00	95	5	3.00
	0.20	95	5	3.00

25771-1956

1.50	2	98	3.00
1.90	2	98	3.00
2.00	2	98	3.00

- 5 Stationary phase: Waters, X-Bridge, C18, 3,5 µm, 4,6 X 20 mm.
Column temp: constant at 25°C.
Die Diode array detection is at the wavelength range of 210-500 nm.

Method F

- 10 Instrument: LC/MS ThermoFinnigan. HPLC Surveyor DAD, MSQ Quadrupole; column: Symmetry C8, 5 µm, 3 x 150 mm; eluent A: 90% water +10% acetonitrile+ammonium formate 10mM; eluent B= ACN 90%+10% H₂O+NH₄COOH 10mM; gradient: A(100) for 1.5 min, then to B (100) in 10 min for 1.5 min; flow rate: 1.2 mL/min; UV Detection: 254nm; Ion source: APCI

15

Method G

- Instrument: LC/MS Waters. HPLC Alliance 2695 DAD, ZQ Quadrupole; column: Sunfire MS-C18, 3.5µm, 4.6x50 mm; eluent A: water +0.1% TFA + 10% acetonitrile; eluent B: acetonitrile; gradient: A/B(80:20) , then to A/B (10:90) in 3.25 min for 0.75 min; flow rate:1.3 ml/min;; UV Detection: 254nm; Ion source: ESI.

20

Method H

- Instrument: LC/MS Waters. HPLC Alliance 2695 DAD, ZQ Quadrupole; column: Xterra™ MS-C18, 3.5µm, 4.6x50 mm; eluent A: water +0.1% TFA + 10% acetonitrile; eluent B: acetonitrile; gradient: A (100) , then to A/B (10:90) in 3.25 min for 0.75 min; flow rate:1.3 ml/min;; UV Detection: 254nm; Ion source: ESI.

25

Method L

- Instrument: LC/MS Waters. HPLC Alliance 2695 DAD, ZQ Quadrupole. Column: Xterra MS-C8, 3.5 µm, 4.6x50 mm; eluent A : water + ammonium formate 5mM + 10% acetonitrile; eluent B : acetonitrile; gradient: A 100, then to A/B (10:90) in 3.25 min for 0.75 min; flow rate: 1.3 ml/min ; UV Detection: 254nm; Ion source: ESI.

30

Method M

- Instrument: UPLC/MS Waters. Acquity UPLC® system, ZQ Quadrupole. Column: HSS C18, 1.8 µm, 2.1x50 mm; eluent A: water + ammonium formate 5mM + 10% acetonitrile; eluent B : acetonitrile; gradient: A/B (90:10), then to A/B (10:90) in 1.70 min for 0.64 min; flow rate: 0.56 ml/min ; UV Detection: 254nm; Ion source: ESI.

35

Method N

- Instrument: UPLC/MS Waters. Acquity UPLC® system, ZQ Quadrupole. Column: BEH C18, 1.7µm, 2.1x50 mm; eluent A: water +0.1% TFA + 10% acetonitrile; eluent B: acetonitrile; gradient: A/B(50:50) , then to A/B (10:90) in 1.99 min for 0.77 min; flow rate: 0.48 ml/min; UV Detection: 254nm; Ion source: ESI.

40

45

Method O

- Instrument: LC/MS ThermoFinnigan. HPLC Surveyor DAD, MSQ Quadrupole; column:SynergyHydro-RP80A, 4 µm, 4.60 x 100 mm; eluent A: 90% water +10% acetonitrile+ammonium formate 10mM; eluent B= ACN 90%+10% H₂O+NH₄COOH 10mM; gradient: A(100) for 1.5 min, then to B (100) in 10 min for 1.5 min; flow rate: 1.2 mL/min; UV Detection: 254nm; Ion source: APCI

50

Method P

Instrument: LC/MS ThermoFinnigan. HPLC Surveyor DAD, Finnigan LCQduo Ion trap;
column: Symmetry –C18, 5 μ m, 3x150 mm; eluent A: 95% water +5% acetonitrile+formic
acid 0.1%; eluent B= Acetonitrile 95%+5% water+formic acid 0.1%; gradient: A/B(95/5) for
1.5 min, then to A/B (5/95) in 10 min for 1.5 min; flow rate: 1 mL/min; UV Detection: 254nm;
Ion source: ESI

Method Q

Instrument: LC/MS Waters. HPLC Alliance 2695 DAD, ZQ Quadrupole. Column: Zorbax
Eclipse Plus C18, 3.5 μ m, 4.6x50 mm; eluent A : water + ammonium formate 5mM + 10%
acetonitrile; eluent B : acetonitrile; gradient: A 100, then to A/B (10:90) in 3.5 min for 1 min;
flow rate: 1.3 ml/min ; UV Detection: 254nm; Ion source: ESI.

S.5.2 Chromatographic methods: GC methods**Method 3A**

Instrument: GC/MS Finnigan TRACE GC, TRACE MS Quadrupole; column: Agilent DB-5MS,
25m x 0.25mm x 0.25 μ m; carriage gas: Helium; flow rate: 1 ml/min constant flow; oven
program: 50°C (hold 1 min.), to 100°C in 10°C / min., to 200°C in 20°C / min., to 300°C in
30°C / min.; Detection: TRACE MS quadrupole; Ion source: EI.

Microwave heating

Microwave apparatus type: Discover® CEM instruments, equipped with 10 and 35 mL
vessels;

PHARMACOLOGICAL PART

In another aspect, the instant invention may be used to evaluate the putative specific
agonists or antagonists of a G protein coupled receptor. The present invention is directed to
the use of these compounds in the preparation and execution of screening assays for
compounds that modulate the activity of chemokine receptors. Furthermore, the compounds
of this invention are useful in establishing or determining the binding site of other compounds
to chemokine receptors, e.g., by competitive inhibition or as a reference in an assay to
compare its known activity to a compound with an unknown activity. When developing new
assays or protocols, compounds according to the present invention could be used to test
their effectiveness.

Specifically, such compounds may be provided in a commercial kit, for example, for use in
pharmaceutical research involving the aforementioned diseases. The compounds of the
instant invention are also useful for the evaluation of putative specific modulators of the
chemokine receptors. In addition, one could utilize compounds of this invention to examine
the specificity of G protein coupled receptors that are not thought to be chemokine receptors,
either by serving as examples of compounds which do not bind or as structural variants of
compounds active on these receptors which may help define specific sites of interaction.

The CCR3 receptor binding test is based on a K562 cell line (leukemia myelogenic blast
cells) transfected with the human chemokine receptor CCR3 (hCCR3-C1 cells). The cell
membranes were prepared by disrupting the hCCR3-C1 cells by nitrogen decomposition.
The preparation was centrifuged at 400 g 4°C for 30 min. The supernatant was transferred
into fresh tubes followed by a second centrifugation at 48000 g, 4°C for 1h. The membranes
were re-suspended in the SPA incubation buffer (25mM HEPES, 25mM MgCl₂ 6xH₂O, 1mM

CaCl₂ 2xH₂O) without bovine serum albumin and homogenized by passing through a single use needle (Terumo, 23Gx1"). The membranes were stored in aliquots at -80°C.

5 The CCR3 receptor binding assay was performed in a Scintillation Proximity Assay (SPA) design with the radioligand recombinant human ¹²⁵Iodine-eotaxin-1. Cell membranes of hCCR3 C1 cells were again homogenized by passing through a single use needle (Terumo, 23Gx1") and diluted in SPA incubation buffer in suitable concentrations (0.5 – 10 µg protein/well) in 96 well microtiter plates (1450-514, Perkin Elmer). The SPA assay was set up
10 in the SPA incubation buffer with a final volume of 200µl and final concentration of 25mM HEPES, 25mM MgCl₂ 6xH₂O, 1mM CaCl₂ 2xH₂O and 0,1% bovine serum albumin . The SPA assay mixture contained 60 µl of the membrane suspension, 80 µl of Wheat Germ Agglutinin coated PVT beads (organic scintillator, GE Healthcare, RPNQ-0001) 0,2 mg/well , 40 µl of recombinant human ¹²⁵Jodine-eotaxin-1 (Biotrend), diluted in SPA buffer to a final
15 concentration of 30.000 dpm per well, and 20 µl of the test compound (dissolved in DMSO dilutions). The SPA assay mixture was incubated for 2 h at room temperature. Bound radioactivity was determined with a scintillation counter (Micro Beta "Trilux", Wallac). Included were controls for total binding (no displacer added, Bo) and non-specific binding (NSB) by adding unlabelled recombinant human Eotaxin-1 (Biotrend, Cat #300-21) or a
20 reference compound.

Determination of the affinity of a test compound was calculated by subtraction of the non-specific binding (NSB) from the total binding (Bo) or the binding in the presence of the test compound (B) at a given compound concentration. The NSB value was set to 100%
25 inhibition. The Bo-NSB value was set to 0% inhibition.

% inhibition values were obtained at a defined compound concentration, e.g. at 1µM, % inhibition of the test compound was calculated by the formula $100 - ((B - NSB) * 100 / (Bo - NSB))$. % inhibition values above 100% are founded by assay variance.
30

The dissociation constant K_i was calculated by iterative fitting of experimental data obtained at several compound concentrations over a dose range from 0.1 to 10000nM using the law of mass action based program "easy sys" (Schittkowski, Num Math 68, 129-142 (1994)).

35 The utility of the compounds in accordance with the present invention as inhibitors of chemokine receptor activity may be demonstrated by methodology known in the art, such as the assays for CCR3 ligand binding, as disclosed by Ponath et al., J. Exp. Med., 183,2437-2448 (1996) and Uguccioni et al., J. Clin. Invest., 100,11371143 (1997). Cell lines for expressing the receptor of interest include those naturally expressing the chemokine
40 receptor, such as EOL-3 or THP-1, those induced to express the chemokine receptor by the addition of chemical or protein agents, such as HL-60 or AML14.3D10 cells treated with, for example, butyric acid with interleukin-5 present, or a cell engineered to express a recombinant chemokine receptor, such as L1.2, K562, CHO or HEK-293 cells. Finally, blood or tissue cells, for example human peripheral blood eosinophils, isolated using methods as
45 described by Hansel et al., J. Immunol. Methods, 145,105-110 (1991), can be utilized in such assays. In particular, the compounds of the present invention have activity in binding to the CCR3 receptor in the aforementioned assays and inhibit the activation of CCR3 by CCR3 ligands, including eotaxin-1, eotaxin-2, eotaxin-3, MCP-2, MCP-3, MCP-4 or RANTES.

50 As used herein, "activity" is intended to mean a compound demonstrating an inhibition of 50% at 1µM or higher in inhibition when measured in the aforementioned assays. Such a result is indicative of the intrinsic activity of the compounds as inhibitor of CCR3 receptor activity.

%inhibition and Ki values are (human Eotaxin-1 at human CCR3-Rezeptor):

#	CCR3 %inh @ 1µM	CCR3 Ki (nM)
1	101	4
2	98	4
3	75	240
4	99	5
5	98	6
6	100	7
7	97	7
8	99	9
9	95	12
10	98	16
11	92	27
12	84	88
13	71	373
14	60	540
15	38	1111
16	98	20
17	100	5
18	96	7
19	98	7
20	97	9
21	96	11
22	84	13
23	95	18
24	99	22
25	91	21
26	97	27
27	96	28
28	97	29
29	94	31
30	95	47
31	93	50

#	CCR3 %inh @ 1µM	CCR3 Ki (nM)
32	82	145
33	83	150
34	80	152
35	83	152
36	80	159
37	80	175
38	78	211
39	80	227
40	75	229
41	66	395
42	63	482
43	59	523
44	63	552
45	50	713
46	44	873
47	31	1188
48	40	965
49	41	1184
50	41	1287
51	30	1615
52	28	1752
53	33	2010
54	90	89
55	78	235
56	76	254
57	74	302
58	76	221
59	66	402
60	65	430
61	57	492
62	63	507

#	CCR3 %inh @ 1µM	CCR3 Ki (nM)
63	60	518
64	60	545
65	57	602
66	81	658
67	52	700
68	48	715
69	55	749
70	46	791
71	49	810
72	48	875
73	48	922
74	46	927
75	97	30
76	97	37
77	96	38
78	94	50
79	91	52
80	95	55
81	89	111
82	87	153
83	81	223
84	77	232
85	68	350
86	65	419
91	98	18
93	95	29
94	94	52
95	94	40
96	51	670
97	50	669
98	75	202
99	91	67
100	100	7

#	CCR3 %inh @ 1µM	CCR3 Ki (nM)
101	95	45
102	100	19
103	54	1402
104	24	2399
105	99	5
106	100	3
107	99	13
108	98	4
114	99	4
115	80	18
116	93	47
117	98	17
118	94	40
119	87	81
120	95	35
121	98	21
122	95	30
123	95	34
124	93	41
125	97	58
126	89	71
127	90	81
128	87	96
129	70	334
130	91	77
131	89	85
132	102	1
133	100	3
134	101	3
135	101	3
136	100	4
137	101	4
138	101	4

#	CCR3 %inh @ 1µM	CCR3 Ki (nM)
139	101	5
140	98	6
141	94	6
142	100	17
143	99	7
144	100	7
145	99	8
146	100	8
147	99	8
148	99	10
149	99	10
150	97	10
151	95	18
152	100	12
153	99	12
154	98	14
155	98	15
156	98	15
157	99	15
158	100	16
159	96	19
160	97	19
161	97	20
162	97	20
163	99	23
164	96	24
165	90	62
166	60	905
167	98	6
168	99	4
169	99	19
170	100	6
171	99	5

#	CCR3 %inh @ 1µM	CCR3 Ki (nM)
172	95	7
173	99	13
174	99	13
175	99	6
176	100	11
177	97	24
178	86	108
179	95	34
180	97	16
182	99	4
183	75	232
184	100	8
185	98	1
186	99	1
187	99	4
188	99	6
189	100	4
190	100	6
191	100	5
192	96	6
193	99	19
196	89	95
197	44	1251
198	91	67
199	66	453
200	86	114
201	86	119
202	69	390
203	75	237
204	80	299
205	76	247
206	77	357
207	75	442

25771-1956

#	CCR3 %inh @ 1μM	CCR3 Ki (nM)
208	94	43
209	95	33
210	96	13
211	93	48
212	97	26

#	CCR3 %inh @ 1μM	CCR3 Ki (nM)
213	66	414
214	101	2

25771-1956

PHARMACEUTICAL FORMS

While it may be possible for the compounds of the subject invention to be administered as the raw chemical, it is also possible to present them as a pharmaceutical formulation.

Accordingly, provided herein are pharmaceutical formulations which comprise one or more of

- 5 certain compounds disclosed herein, or one or more pharmaceutically acceptable salts, esters, prodrugs, amides, or solvates thereof, together with one or more pharmaceutically acceptable carriers thereof and optionally one or more other ingredients. The carrier(s) must be “acceptable” in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. Proper formulation is dependent upon the route
- 10 of administration chosen. Any of the well-known techniques, carriers, and excipients may be used as suitable and as understood in the art; e.g., in Remington’s Pharmaceutical Sciences. The pharmaceutical compositions disclosed herein may be manufactured in any manner known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.
- 15 The following examples of formulations illustrate the present invention without restricting its scope:

25771-1956

EXAMPLES OF PHARMACEUTICAL FORMULATIONS

5	A)	<u>Tablets</u>	<u>per tablet</u>
		active substance 1	100 mg
		lactose	140 mg
		maize starch	240 mg
		polyvinylpyrrolidone	15 mg
		magnesium stearate	5 mg
10			500 mg

15 The finely ground active substance, lactose and some of the maize starch are mixed together. The mixture is screened, then moistened with a solution of polyvinylpyrrolidone in water, kneaded, wet granulated and dried. The granules, the remaining maize starch and the magnesium stearate are screened and mixed together. The mixture is pressed into tablets of suitable shape and size.

20	B)	<u>Tablets</u>	<u>per tablet</u>
		active substance 1	80 mg
		lactose	55 mg
		maize starch	190 mg
		microcrystalline cellulose	35 mg
		polyvinylpyrrolidone	15 mg
		sodium carboxymethyl starch	23 mg
		magnesium stearate	2 mg
25			400 mg

30 The finely ground active substance, some of the corn starch, lactose, microcrystalline cellulose and polyvinylpyrrolidone are mixed together, the mixture is screened and worked with the remaining corn starch and water to form a granulate which is dried and screened. The sodium carboxymethyl starch and the magnesium stearate are added and mixed in and the mixture is compressed to form tablets of a suitable size.

35	C)	<u>Ampoule solution</u>	
		active substance 1	50 mg
		sodium chloride	50 mg
		water for inj.	5 ml

40 The active substance is dissolved in water at its own pH or optionally at pH 5.5 to 6.5 and sodium chloride is added to make the solution isotonic. The resulting solution is filtered to remove pyrogens and the filtrate is transferred under aseptic conditions into ampoules which are then sterilised and heat-sealed. The ampoules contain 5 mg, 25 mg and 50 mg of active substance.

45	D)	<u>Metering aerosol</u>	
		active substance 1	0.005
		sorbitan trioleate	0.1
		monofluorotrichloromethane and	
		TG134a : TG227 2:1	ad 100
50			

25771-1956

The suspension is transferred into a conventional aerosol container with metering valve. Preferably 50 µl suspension are released on each actuation. The active substance may also be released in higher doses if desired (e.g. 0.02 wt.-%).

5	E)	<u>Solutions (in mg/100ml)</u>	
		active substance 1	333.3 mg
		benzalkonium chloride	10.0 mg
		EDTA	50.0 mg
		HCl (1N)	ad pH 2.4

10

This solution can be prepared in the usual way.


15	F)	<u>Inhalable powder</u>	
		active substance 1	12 µg
		lactose monohydrate	ad 25 mg

The inhalable powder is prepared in the usual way by mixing the individual ingredients.

25771-1956

CLAIMS:

1. A compound of formula 1,



5 wherein

A is CH₂, O or N-C₁₋₆-alkyl;R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};
- NHR^{1.2}, NMeR^{1.2};
- NHCH₂-R^{1.3};
- NH-C₃₋₆-cycloalkyl, whereas optionally one carbon atom is replaced by a nitrogen atom, whereas the ring is optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, O-C₁₋₆-alkyl, NHSO₂-phenyl, NHCONH-phenyl, halogen, CN, SO₂-C₁₋₆-alkyl, COO-C₁₋₆-alkyl;
- a C₉ or C₁₀-bicyclic-ring, whereas one or two carbon atoms are replaced by nitrogen atoms and the ring system is bound via a nitrogen atom to the basic structure of formula 1 and whereas the ring system is optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, COO-C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, NO₂, halogen, CN, NHSO₂-C₁₋₆-alkyl, methoxy-phenyl;
- a group selected from NHCH(pyridinyl)CH₂COO-C₁₋₆-alkyl, NHCH(CH₂O-C₁₋₆-alkyl)-benzimidazolyl, optionally substituted with halogen or CN;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazole;

25

R^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₁₋₆-haloalkyl, C₁₋₆-alkylene-OH, C₂₋₆-alkenylene-OH, C₂₋₆-alkynylene-OH, CH₂CON(C₁₋₆-alkyl)₂, CH₂NHCONH-C₃₋₆-cycloalkyl, CN, CO-pyridinyl, CONR^{1.1.1}R^{1.1.2}, COO-C₁₋₆-alkyl, N(SO₂-C₁₋₆-alkyl)(CH₂CON(C₁₋₄-alkyl)₂) O-C₁₋₆-alkyl, O-pyridinyl, SO₂-C₁₋₆-alkyl, SO₂-C₁₋₆-alkylene-OH, SO₂-C₃₋₆-cycloalkyl, SO₂-piperidinyl, SO₂NH-C₁₋₆-alkyl, SO₂N(C₁₋₆-alkyl)₂, halogen, CN, CO-morpholinyl, CH₂-pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, NHC₁₋₆-alkyl and =O;

35

R^{1.1.1} H, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, C₁₋₆-haloalkyl, CH₂CON(C₁₋₆-alkyl)₂, CH₂CO-azetindinyl, C₁₋₆-alkylene-C₃₋₆-cycloalkyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, C₁₋₆-alkylene-OH or thiadiazolyl, optionally substituted with C₁₋₆-alkyl;

40

R^{1.1.2} H, C₁₋₆-alkyl, SO₂C₁₋₆-alkyl;

or R^{1.1.1} and R^{1.1.2} together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one N or O, replacing a carbon atom of

25771-1956

the ring, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₄-alkylene-OH, OH, =O;

or

5

R^{1.1} is phenyl, wherein two adjacent residues are together forming a five- or six-membered carbocyclic aromatic or non-aromatic ring, optionally containing independently from each other one or two N, S, or SO₂, replacing a carbon atom of the ring, wherein the ring is optionally substituted with C₁₋₄-alkyl or =O;

10

R^{1.2} is selected from

15

- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₃₋₆-cycloalkyl, CH₂COO-C₁₋₆-alkyl, CONR^{1.2.1}R^{1.2.2}, COR^{1.2.3}, COO-C₁₋₆-alkyl, CONH₂, O-C₁₋₆-alkyl, halogen, CN, SO₂N(C₁₋₆-alkyl)₂ or heteroaryl optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl;

20

- heteroaryl, optionally substituted with a five- or six-membered carbocyclic non-aromatic ring containing independently from each other two N, O, S, or SO₂, replacing a carbon atom of the ring;

25

- a aromatic or non-aromatic C₉ or 10-bicyclic-ring, whereas one or two carbon atoms are replaced by N, O or S each optionally substituted with one or two residues selected from the group consisting of N(C₁₋₆-alkyl)₂, CONH-C₁₋₆-alkyl, =O;
- a heterocyclic non-aromatic ring, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCO-C₁₋₆-alkyl,

30

R^{1.2.1} H, C₁₋₆-alkyl, C₁₋₆-alkylene-C₃₋₆-cycloalkyl, C₁₋₄-alkylene-phenyl, C₁₋₄-alkylene-furanyl, C₃₋₆-cycloalkyl, C₁₋₄-alkylene-O-C₁₋₄-alkyl, C₁₋₆-haloalkyl or a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO₂, replacing a carbon atom of the ring, optionally substituted with 4-cyclopropylmethyl-piperazinyl

35

R^{1.2.2} H, C₁₋₆-alkyl;

R^{1.2.3} a five- or six-membered carbocyclic non-aromatic ring, optionally containing independently from each other one or two N, O, S, or SO₂, replacing a carbon atom of the ring;

40

R^{1.3} is selected from phenyl, heteroaryl or indolyl, each optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₃₋₆-cycloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, phenyl, heteroaryl;

45

R² is selected from the group consisting of C₁₋₆-alkylene-phenyl, C₁₋₆-alkylene-naphthyl, and C₁₋₆-alkylene-heteroaryl; each optionally substituted with one, two or three residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, O-C₁₋₆-haloalkyl, halogen;

50

R³ is H, C₁₋₆-alkyl;

R⁴ is H, C₁₋₆-alkyl;

or R³ and R⁴ together are forming a CH₂-CH₂ group.

25771-1956

2. A compound of formula 1 according to claim 1, wherein

5 A is CH₂, O or N-C₁₋₄-alkyl;

R¹ is selected from

- NHR^{1.1}, NMeR^{1.1};
- NHR^{1.2}, NMeR^{1.2};
- 10 • NHCH₂-R^{1.3};
- NH-C₃₋₆-cycloalkyl, whereas optionally one carbon atom is replaced by a nitrogen atom, whereas the ring is optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, O-C₁₋₆-alkyl, NHSO₂-phenyl, NHCONH-phenyl, halogen, CN, SO₂-C₁₋₆-alkyl, COO-C₁₋₆-alkyl;
- 15 • a C₉ or C₁₀-bicyclic-ring, whereas one or two carbon atoms are replaced by nitrogen atoms and the ring system is bound via a nitrogen atom to the basic structure of formula 1 and whereas the ring system is optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, COO-C₁₋₆-alkyl, C₁₋₆-haloalkyl, O-C₁₋₆-alkyl, NO₂, halogen, CN, NHSO₂-C₁₋₆-alkyl, m-methoxyphenyl;
- 20 • a group selected from NHCH(pyridinyl)CH₂COO-C₁₋₆-alkyl, NHCH(CH₂O-C₁₋₆-alkyl)-benzoimidazolyl, optionally substituted with Cl; or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;

25 R^{1.1} is phenyl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₁₋₆-haloalkyl, CH₂CON(C₁₋₆-alkyl)₂, CH₂NHCONH-C₃₋₆-cycloalkyl, CN, CONR^{1.1.1}R^{1.1.2}, COO-C₁₋₆-alkyl, O-C₁₋₆-alkyl, SO₂-C₁₋₆-alkyl, SO₂-C₁₋₆-alkylen-OH, SO₂-C₃₋₆-cycloalkyl, SO₂-piperidinyl, SO₂NH-C₁₋₆-alkyl, SO₂N(C₁₋₆-alkyl)₂, halogen, CN, CO-morpholinyl, CH₂-pyridinyl or a heterocyclic ring optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, NHC₁₋₆-alkyl, =O;

35 R^{1.1.1} H, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, C₁₋₆-haloalkyl, CH₂CON(C₁₋₆-alkyl)₂, CH₂CO-azetindinyl, C₁₋₆-alkylen-C₃₋₆-cycloalkyl, CH₂-pyranyl, CH₂-tetrahydrofuranyl, CH₂-furanyl, C₁₋₆-alkylen-OH or thiadiazolyl, optionally substituted with C₁₋₆-alkyl;

R^{1.1.2} H, C₁₋₆-alkyl, SO₂C₁₋₆-alkyl;

40 or R^{1.1.1} and R^{1.1.2} together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH

45 R^{1.2} is selected from

- heteroaryl, optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl, C₃₋₆-cycloalkyl, CH₂COO-C₁₋₆-alkyl, CONR^{1.2.1}R^{1.2.2}, COO-C₁₋₆-alkyl, CONH₂, O-C₁₋₆-alkyl, halogen, CN, CO-pyrrolidinyl, CO-morpholinyl or heteroaryl optionally substituted with one or two residues selected from the group consisting of C₁₋₆-alkyl;
- 50

25771-1956

- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolynyl, each optionally substituted with one or two residues selected from the group consisting of $N(C_{1-6}\text{-alkyl})_2$, $CONH-C_{1-6}\text{-alkyl}$, $=O$;
- piperidinyl, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with $NHCO-C_{1-6}\text{-alkyl}$,

$R^{1.2.1}$ H, $C_{1-6}\text{-alkyl}$;

$R^{1.2.2}$ H, $C_{1-6}\text{-alkyl}$;

$R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of $C_{1-6}\text{-alkyl}$, $C_{3-6}\text{-cycloalkyl}$, $O-C_{1-6}\text{-alkyl}$, $O-C_{1-6}\text{-haloalkyl}$;

R^2 is selected from $CH_2\text{-phenyl}$ or $CH_2\text{-naphthyl}$, both optionally substituted with one or two residues selected from the group consisting of $C_{1-6}\text{-alkyl}$, $C_{1-6}\text{-haloalkyl}$, $O-C_{1-6}\text{-alkyl}$, $O-C_{1-6}\text{-haloalkyl}$, halogen; or $CH_2\text{-thiophenyl}$, optionally substituted with one or two residues selected from the group consisting of halogen;

R^3 is H, $C_{1-4}\text{-alkyl}$;

R^4 is H, $C_{1-4}\text{-alkyl}$;

or R^3 and R^4 together are forming a $CH_2\text{-CH}_2$ group.

3. A compound of formula 1 according to claim 1 or 2, wherein

A is CH_2 , O or NMe;

R^1 is selected from

- $NHR^{1.1}$, $NMeR^{1.1}$;
- $NHR^{1.2}$, $NMeR^{1.2}$;
- $NHCH_2-R^{1.3}$;
- NH-cyclohexyl, optionally substituted with one or two residues selected from the group consisting of $C_{1-4}\text{-alkyl}$, $NHSO_2\text{-phenyl}$, $NHCONH\text{-phenyl}$, halogen;
- NH-pyrrolidinyl, optionally substituted with one or two residues selected from the group consisting of $SO_2-C_{1-4}\text{-alkyl}$, $COO-C_{1-4}\text{-alkyl}$;
- piperidinyl, optionally substituted with one or two residues selected from the group consisting of $NHSO_2-C_{1-4}\text{-alkyl}$, m-methoxyphenyl;
- dihydro-indolyl, dihydro-isindolyl, tetrahydro-quinolynyl or tetrahydro-isoquinolynyl, optionally substituted with one or two residues selected from the group consisting of $C_{1-4}\text{-alkyl}$, $COO-C_{1-4}\text{-alkyl}$, $C_{1-4}\text{-haloalkyl}$, $O-C_{1-4}\text{-alkyl}$, NO_2 , halogen;
- a group selected from $NHCH(\text{pyridinyl})CH_2COO-C_{1-4}\text{-alkyl}$, $NHCH(CH_2O-C_{1-4}\text{-alkyl})\text{-benzoimidazolyl}$, optionally substituted with Cl;
- or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;

$R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of $C_{1-4}\text{-alkyl}$, $C_{1-4}\text{-haloalkyl}$, $CH_2CON(C_{1-4}\text{-alkyl})_2$, $CH_2NHCONH-C_{3-6}\text{-cycloalkyl}$, CN, $CONR^{1.1.1}R^{1.1.2}$, $COO-C_{1-4}\text{-alkyl}$, $O-C_{1-4}\text{-alkyl}$, $SO_2-C_{1-4}\text{-alkyl}$, $SO_2-C_{1-4}\text{-alkylen-OH}$, $SO_2-C_{3-6}\text{-cycloalkyl}$, $SO_2\text{-piperidinyl}$, $SO_2NH-C_{1-4}\text{-alkyl}$, $SO_2N(C_{1-4}\text{-alkyl})_2$, halogen, CO-morpholynyl, $CH_2\text{-pyridinyl}$, or imidazolidinyl, piperidinyl, oxazinanyl,

25771-1956

pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of C₁₋₄-alkyl, NHC₁₋₄-alkyl, =O;

5 R^{1.1.1} H, C₁₋₆-alkyl, C₃₋₆-cycloalkyl, C₁₋₄-haloalkyl, CH₂CON(C₁₋₄-alkyl)₂,
CH₂CO-azetindinyl, C₁₋₄-alkylen-C₃₋₆-cycloalkyl, CH₂-pyranyl,
CH₂-tetrahydrofuranyl, CH₂-furanyl, C₁₋₄-alkylen-OH or thiadiazolyl,
optionally substituted with C₁₋₄-alkyl;
10 R^{1.1.2} H, C₁₋₄-alkyl, SO₂C₁₋₄-alkyl;

or R^{1.1.1} and R^{1.1.2} together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH₂OH

15 R^{1.2} is selected from

- pyridinyl, pyridazinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of C₁₋₄-alkyl, C₃₋₆-cycloalkyl, CH₂COO-C₁₋₄-alkyl, CONR^{1.2.1}R^{1.2.2}, COO-C₁₋₄-alkyl, CONH₂, O-C₁₋₄-alkyl, halogen, CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C₁₋₄-alkyl;
- benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolinyl, each optionally substituted with one or two residues selected from the group consisting of N(C₁₋₄-alkyl)₂, CONH-C₁₋₄-alkyl, =O;
- piperidinyl, optionally substituted with pyridinyl;
- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCO-C₁₋₄-alkyl,

30 R^{1.2.1} H, C₁₋₄-alkyl;
R^{1.2.2} H, C₁₋₄-alkyl;

35 R^{1.3} is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of C₁₋₄-alkyl, C₃₋₆-cycloalkyl, O-C₁₋₄-alkyl, O-C₁₋₄-haloalkyl;

40 R² is selected from CH₂-phenyl or CH₂-naphthyl, both optionally substituted with one or two residues selected from the group consisting of C₁₋₄-alkyl, C₁₋₄-haloalkyl, O-C₁₋₄-haloalkyl, halogen; or CH₂-thiophenyl, optionally substituted with one or two residues selected from the group consisting of halogen;

45 R³ is H;
R⁴ is H;

or R³ and R⁴ together are forming a CH₂-CH₂ group.

50 4. A compound of formula 1 according to any one of claims 1 to 3, wherein

A is CH₂, O or NMe;

25771-1956

- R^1 is selected from
- $NHR^{1.1}$, $NMeR^{1.1}$;
 - $NHR^{1.2}$, $NMeR^{1.2}$;
 - $NHCH_2R^{1.3}$;
 - 5 • NH-piperidinyl, optionally substituted with pyridinyl;
 - NH-cyclohexyl, optionally substituted with one or two residues selected from the group consisting of t-Bu, $NHSO_2$ -phenyl, $NHCONH$ -phenyl, F;
 - NH-pyrrolidinyl, optionally substituted with one or two residues selected from the group consisting of SO_2Me , COO -t-Bu;
 - 10 • piperidinyl, optionally substituted with one or two residues selected from the group consisting of $NHSO_2$ -n-Bu, m-methoxyphenyl;
 - dihydro-indolyl, dihydro-isoindolyl, tetrahydro-quinoliny or tetrahydro-isoquinoliny, optionally substituted with one or two residues selected from the group consisting of Me, $COOMe$, CF_3 , OMe , NO_2 , F, Br;
 - 15 • a group selected from $NHCH$ (pyridinyl) CH_2COOMe , $NHCH$ (CH_2OMe)-benzoimidazolyl, optionally substituted with Cl;
 - or 1-aminocyclopentyl, optionally substituted with methyl-oxadiazolyl;
- $R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, t-Bu, CF_3 , CH_2CONMe_2 , $CH_2NHCONH$ -cyclohexyl, CN, $CONR^{1.1.1}R^{1.1.2}$, $COOMe$, $COOEt$, OMe , SO_2Me , $SO_2CH_2CH_2OH$, SO_2Et , SO_2 -cyclopropyl, SO_2 -piperidinyl, SO_2NHEt , SO_2NMeEt , F, Cl, CO-morpholinyl, CH_2 -pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl, pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, $NHMe$, =O;
- 20
- $R^{1.1.1}$ H, Me, Et, t-Bu, i-Pr, cyclopropyl, CH_2 -i-Pr, CH_2 -t-Bu, $CH(CH_3)CH_2CH_3$, CH_2CHF_2 , CH_2CONMe_2 , CH_2CO -azetindinyl, CH_2 -cyclopropyl, CH_2 -cyclobutyl, CH_2 -pyranyl, CH_2 -tetrahydrofuranyl, CH_2 -furanyl, CH_2CH_2OH or thiadiazolyl, optionally substituted with Me;
- 30
- $R^{1.1.2}$ H, Me, Et, SO_2Me , SO_2Et
- or $R^{1.1.1}$ and $R^{1.1.2}$ together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH_2OH
- 35
- $R^{1.2}$ is selected from
- 40 • pyridinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, cyclopropyl, CH_2COOEt , $CONR^{1.2.1}R^{1.2.2}$, $COOMe$, $COOEt$, $CONH_2$, OMe , Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;
 - 45 • benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinoliny, each optionally substituted with one or two residues selected from the group consisting of NMe_2 , $CONHMe$, =O;
 - 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with $NHCOMe$,
- 50
- $R^{1.2.1}$ H, Me;
- $R^{1.2.2}$ H, Me;

25771-1956

$R^{1.3}$ is selected from phenyl, pyrazolyl, isoxazolyl, pyrimidinyl, indolyl or oxadiazolyl, each optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, cyclopentyl, OMe, $OCHF_2$;

5 R^2 is selected from CH_2 -phenyl or CH_2 -naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH_3 , CF_3 , OCF_3 , F, Cl, Br, Et; or CH_2 -thiophenyl, optionally substituted with one or two residues selected from the group consisting of Cl, Br,

10 R^3 is H;
 R^4 is H;

or R^3 and R^4 together are forming a CH_2-CH_2 group.

15

5. A compound of formula 1 according to any one of claims 1 to 4, wherein

A is CH_2 , O or NMe;

20 R^1 is selected from
 • $NHR^{1.1}$
 • $NHR^{1.2}$,

25 $R^{1.1}$ is phenyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Bu, CF_3 , CH_2CONMe_2 , $CH_2NHCONH$ -cyclohexyl, CN, $CONR^{1.1.1}R^{1.1.2}$, COOMe, COOEt, OMe, SO_2Me , $SO_2CH_2CH_2OH$, SO_2Et , SO_2 -cyclopropyl, SO_2 -piperidinyl, SO_2NHEt , SO_2NMeEt , F, Cl, CO-morpholinyl, CH_2 -pyridinyl, or imidazolidinyl, piperidinyl, oxazinanyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, oxadiazolyl, thiazolyl, pyridinyl,
 30 pyrimidinyl, each optionally substituted with one or two residues selected from the group consisting of Me, NHMe, =O;

35 $R^{1.1.1}$ H, Me, Et, t-Bu, i-Pr, cyclopropyl, CH_2 -i-Pr, CH_2 -t-Bu, $CH(CH_3)CH_2CH_3$, CH_2CHF_2 , CH_2CONMe_2 , CH_2CO -azetindinyl, CH_2 -cyclopropyl, CH_2 -cyclobutyl, CH_2 -pyranlyl, CH_2 -tetrahydrofuranyl, CH_2 -furanyl, CH_2CH_2OH or thiadiazolyl, optionally substituted with Me;

$R^{1.1.2}$ H, Me, Et, SO_2Me , SO_2Et

40 or $R^{1.1.1}$ and $R^{1.1.2}$ together are forming a four-, five- or six-membered carbocyclic ring, optionally containing one O, replacing a carbon atom of the ring, optionally substituted with one or two residues selected from the group consisting of CH_2OH

45 $R^{1.2}$ is selected from
 • pyridinyl, pyrrolyl, pyrazolyl, isoxazolyl, thiazolyl, thiadiazolyl, optionally substituted with one or two residues selected from the group consisting of Me, Et, Pr, Bu, cyclopropyl, CH_2COOEt , $CONR^{1.2.1}R^{1.2.2}$, COOMe, COOEt, $CONH_2$, OMe, Cl, Br CO-pyrrolidinyl, CO-morpholinyl or pyrazolyl, triazolyl, tetrazolyl, isoxazolyl, oxadiazolyl, each optionally substituted Me;
 50 • benzothiazolyl, indazolyl, dihydro-indolyl, indanyl, tetrahydro-quinolinyl, each optionally substituted with one or two residues selected from the group consisting of NMe_2 , $CONHMe$, =O;

25771-1956

- 4,5-dihydro-naphtho[2,1-d]thiazole, optionally substituted with NHCOMe,

 $R^{1.2.1}$ H, Me;

 $R^{1.2.2}$ H, Me;

5

R^2 is selected from CH_2 -phenyl or CH_2 -naphthyl, both optionally substituted with one or two residues selected from the group consisting of CH_3 , CF_3 , OCF_3 , F, Cl, Br, Et

 R^3 is H;


 R^4 is H.

6. A compound of formula 1 according to any one of claims 1 to 5, wherein


A is CH_2 , O or NMe;

R^1 is selected from


15



25771-1956




25771-1956




5

25771-1956




5

25771-1956




5

25771-1956




5

25771-1956




5

25771-1956




25771-1956



R² is selected from

5



10


R³ is H;
R⁴ is H;

or **R³** and **R⁴** together are forming a CH₂-CH₂ group.

25771-1956

7. Compounds of general formula **1** according to any one of claims 1 to 6, wherein the compounds of formula **1** are present in the form of the individual optical isomers, in the form of mixtures of the individual enantiomers or racemates, or in the form of the enantiomerically pure compounds.


8. Compounds of general formula **1** according to any one of claims 1 to 6, wherein the compounds of formula **1** are present in the form of the R-enantiomer **R-1**

**R-1.**

9. Pharmaceutical composition comprising one or more compounds of formula **1** as defined in any one of claims 1-8, and at least one pharmaceutically acceptable carrier.


10. Use of a compound of formula **1** as defined in any one of claims 1-8 as a CCR3 activity inhibitor.

11. The compound:

**1.a.**


25771-1956

12. The compound:



1.b.


13. The compound:



1.c.


5

14. The compound:



1.d.


10 15. The compound:




1.e.

25771-1956

16. The compound:




17. The compound:





5

18. The compound:



10 19. The compound:





(1)