

COMMONWEALTH of AUSTRALIA

PATENTS ACT 1952

APPLICATION FOR A STANDARD PATENT

X

We SANDOZ LTD
of 35 Lichtstrasse,
CH-4002 Basle,
Switzerland.

623078

••••

•••• hereby apply for the grant of a Standard Patent for an invention entitled:

•••• "IMPROVEMENTS IN OR RELATING TO ORGANIC COMPOUNDS"

••••

•••• which is described in the accompanying ~~provisional~~ complete specification.

Details of basic application(s):—

••••

Number

Convention Country

Date

02317/87-5

SWITZERLAND

19th June 1987

02517/87-2

SWITZERLAND


2nd July 1987

••••

••••

••••

1000420 17/06/88



255.00

The address for service is care of DAVIES & COLLISON, Patent Attorneys, of 1 Little Collins Street, Melbourne, in the State of Victoria, Commonwealth of Australia.

Dated this 17th day of June 19 88

Keith Collison

To: THE COMMISSIONER OF PATENTS

(a member of the firm of DAVIES & COLLISON for and on behalf of the Applicant).

Davies & Collison, Melbourne and Canberra.

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

DECLARATION IN SUPPORT OF CONVENTION OR
NON-CONVENTION APPLICATION FOR A PATENT

Insert title of invention

In support of the Application made for a patent for an invention
entitled: "CYCLIC PEPTOLIDES"

Insert full name(s) and address(es)
of declarant(s) being the appli-
cant(s) or person(s) authorized to
sign on behalf of an applicant
company.

+
WeJEAN KRAMER and HANS RUDOLF HAUS, BOTH OF
SANDOZ LTD., 35 Lichtstrasse, CH-4002 Basle,
Switzerland,

Cross out whichever of paragraphs
1(a) or 1(b) does not apply

do solemnly and sincerely declare as follows :-

1. (a) ~~I am~~ the applicant for the patent
We are
or (b) ~~I am~~ authorized by SANDOZ LTD.

1(a) relates to application made
by individual(s)
1(b) relates to application made
by company; insert name of
applicant company.

the applicant..... for the patent to make this declaration on ~~their~~ its behalf.

~~2. (a) I am~~ the actual inventor..... of the invention
~~We are~~

or (b) Michael Morris Dreyfuss, Paradieshofstrasse 82,
CH-4054 Basle, Switzerland; Max H. Schreier,
Oberwilerstrasse 50, CH-4054 Basle, Switzerland;
Hans Tscherter, Baselmattweg 191/13, CH-4123
Allschwil, Switzerland and Roland Wenger, Grenz-
acherweg 45, CH-4125 Riehen, Switzerland, Dr.
Alexander Haslberger, Prohausergasse 41, A-1130
Wien, Austria

~~is~~ the actual inventor..... of the invention and the facts upon which the applicant.....
~~are~~ is
~~are~~ entitled to make the application are as follows :-

the inventors have assigned the invention to the applicant

State manner in which applicant(s)
derive title from inventor(s)

Cross out paragraphs 3 and 4
for non-convention applications.
For convention applications,
insert basic country(s) followed
by date(s) and basic applicant(s).

3. The basic application..... as defined by Section 141 of the Act ~~was~~ made
in Switzerland..... on the 19th June 1987
by SANDOZ LTD.....
in Switzerland..... on the 2nd July 1987
by SANDOZ LTD.....
in on the
by

4 The basic application..... referred to in paragraph 3 of this Declaration ~~was~~
the first application..... made in a Convention country in respect of the invention the subject
of the application.

Insert place and date of signature.

Declared at Basle this 23rd day of October 1990

Signature of declarant(s) (no
attestation required)

Note Initial all alterations.

SANDOZ Ltd.

Handwritten signatures of Jean Kramer and Hans Rudolf Haus

duly authorised officers


DAVIES & COLLISON, MELBOURNE and CANBERRA.

(12) PATENT ABRIDGMENT (11) Document No. AU-B-18124/88
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 623078

- (54) Title
IMPROVEMENTS IN OR RELATING TO ORGANIC COMPOUNDS
- International Patent Classification(s)
(51)⁴ C07K 011/00 A61K 037/02 C07G 011/00 C12N 001/14
C12P 021/02
(51)⁵ C07K 011/02
- (21) Application No. : 18124/88 (22) Application Date : 17.06.88
- (30) Priority Data
- (31) Number (32) Date (33) Country
02317/87 19.06.87 CH SWITZERLAND
02517/87 02.07.87 CH SWITZERLAND
- (43) Publication Date : 22.12.88
- (44) Publication Date of Accepted Application : 07.05.92
- (71) Applicant(s)
SANDOZ LTD.
- (72) Inventor(s)
MICHAEL MORRIS DREYFUSS; MAX H. SCHREIER; HANS TSCHERTER; ROLAND WENGER; DR.
ALEXANDER HASLBERGER
- (74) Attorney or Agent
DAVIES COLLISON CAVE , 1 Little Collins Street, MELBOURNE VIC 3000
- (56) Prior Art Documents
EP 274873
EP 318680
- (57) Claim

1. A cyclic peptolide which has the structure of a cyclosporin in which one amide linkage is replaced by an ester linkage.

4. A cyclic peptolide according to Claim 3 of formula II



in which W is MeBmt, 8'-OHMeBmt or MeBmtH₂,
X is Sar or Gly,
Y is MeLeu or Leu,
Z is Leu, Ile or Val,
and A is the residue of an α -hydroxycarboxylic acid.

15. A fermentation broth obtained from the cultivation of a cyclic peptolide-producing fungal strain of the genus Cylindrotrichum Bonorden, wherein the cyclic peptolide is one of the compounds according to any one of the claims 1 to 12.

COMMONWEALTH OF AUSTRALIA

PATENT ACT 1952

COMPLETE SPECIFICATION

(ORIGINAL)

FOR OFFICE USE

623078

CLASS

INT. CLASS

Application Number:
Lodged:

Complete Specification Lodged:
Accepted:
Published:

Priority:

Related Art:

NAME OF APPLICANT: SANDOZ LTD

ADDRESS OF APPLICANT: 35 Lichtstrasse,
CH-4002 Basle,
Switzerland.

NAME(S) OF INVENTOR(S) Michael Morris DREYFUSS Hans TSCHERTER
Max H. SCHREIER Roland WENGER

ADDRESS FOR SERVICE: DAVIES & COLLISON, Patent Attorneys
1 Little Collins Street, Melbourne, 3000.

COMPLETE SPECIFICATION FOR THE INVENTION ENTITLED:


"IMPROVEMENTS IN OR RELATING TO ORGANIC COMPOUNDS"

The following statement is a full description of this invention,
including the best method of performing it known to us :-

This invention relates to novel cyclic peptolides useful as pharmaceuticals.

The term peptolide is used herein to mean a natural or synthetic compound consisting of α -hydroxy and α -amino acids joined together by both amide and ester linkages. Thus the structure obtained by replacing an amide linkage by an ester linkage in a peptide is a peptolide.

An important class of peptides is the cyclosporins, which are characterised by a cyclic structure, normally comprising 11 amino acid residues, one of which is the N-Methyl-(4R)-4-but-2E-en-1-yl-4-methyl-(L)- threonyl residue, designated MeBmt, or a derivative thereof. Many cyclosporins have pharmacological properties, particularly immunosuppressive and antiinflammatory properties. The first cyclosporin to be isolated was the naturally occurring fungal metabolite cyclosporin A, (Ciclosporin) sold commercially under the registered Trade Mark Sandimmune[®]. This compound has the structure indicated in formula I



(For a complete list of abbreviations used herein, see Table II)


By convention, the amino-acid residues of cyclosporins are

have the (L) configuration unless otherwise shown; thus in formula I the alanine at position 8 has the (D) configuration. The symbol Me before the abbreviation for an amino acid signifies that the amino acid residue is N-methylated on the nitrogen in the amide linkage.


The present invention provides a cyclic peptolide which has the structure of a cyclosporin in which one amide linkage is replaced by an ester linkage.

Preferably the cyclic peptolide is composed of one MeBmt residue or a derivative thereof, 9 other α -amino acid residues and one α -hydroxyacid residue, which is preferably located at position 8.

Preferred derivatives of MeBmt are the 8'-hydroxy derivative (8'-OHMeBmt) and the saturated dihydro derivative MeBmtH₂, having the structures shown below:




The preferred cyclic peptolides according to the invention have the structure shown in formula II




in which V is MeBmt, 8'-OHMeBmt or MeBmtH₂,
 X is Sar or Gly,
 Y is MeLeu or Leu,
 Z is Leu, Ile or Val,
 and A is the residue of an α-hydroxycarboxylic acid,

preferably of formula III




in which R is C₁₋₄ alkyl.

More preferably in formula III R is isopropyl, so that A represents



the residue of α-hydroxy isovaleric acid, abbreviated Hiv. The most preferred compound according to the invention is that in which, in formula II, W is MeBmt, X is Sar, Y is MeLeu, Z is Leu, and A is (D)Hiv. This may be represented by the full structural formula shown in formula IV



or by using the now conventional nomenclature for cyclosporins, based upon the structure of Ciclosporin (cyclosporin A) shown in formula I. This is done by listing in order each residue in the molecule which differs from that found in Ciclosporin, and adding the term "Ciclosporin". Thus the compound of formula IV may be represented as

(Thr)² (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰ - Ciclosporin,

that is, Ciclosporin in which Thr replaces Abu in position 2, Leu replaces Val in position 5, (D)Hiv replaces (D)Ala in position 8 and Leu replaces MeLeu in position 10, the other residues being identical with those in Ciclosporin.

The cyclic peptides according to the invention may be produced by cultivating a producing microorganism strain in a nutrient medium. Preferred microorganisms are fungal strains of the genus Cylindrotrichum Bonorden, in particular the strain NRRL 18230, which produces cyclic peptolides of formula II.

The strain has been isolated from a leaf sample from Waldenburg in the Swiss Jura, and a viable culture of the strain was deposited on 17 June, 1987 at the US Department of Agriculture (North Central Region, Northern Regional Research Centre), Peoria, Ill. and was given the reference number NRRL 18230. The culture may also be obtained from Sandoz Ltd., Basle, Switzerland.

The fungal strain NRRRL 18230 is a hyphomycete and when incubated at 21-24°C on 2 % malt extract/agar (=MA; 2 % malt extract, 0.4 % yeast extract, 2 % agar in demineralized water) produces aseptate or frequently 1-septate bacilliform hyaline conidia, 6-15 μ x 1.5-2.7 μ (mostly 9.5-13.5 μ) large.

The conidiogenic cells are generally cylindrical and have a conspicuous colarette; some cells appear sympodial-polyphialidic. According to the identification key of M.B. Elles (Dematiaceous Hyphomycetes; Commonwealth Mycological Institute, Kew, Surrey,

England, 1971), the strain may best be classified in the genus Cylindrotrichum Bonorden.


The fungal strain NRRL 18230 grows relatively slowly and after 10 days incubation at a temperature of 21°C forms colonies of 4-7 mm diameter with a velvety grey aerial mycelium. The optimum growth temperature is between 18°C and 27°C, and above 33°C no growth occurs. The branched and septate aerial mycelium of colonies cultivated on MA at 21°C is generally 1.5-3.5 μ (usually 2-3 μ) wide; in the substrate mycelium hyphae of up to 5.5 μ width can be observed.

The invention also provides fermentation broths obtained by cultivation of a strain of the fungal genus Cylindrotrichum Bonorden. The novel strain NRRL 18230 may be cultivated by an aerobic surface or immersion process at suitable temperatures in a variety of nutrient media containing the nutrients and minerals in usable form.

Thus, the medium should contain an assimilatable source of carbon and optionally mineral salts and growth factors. All of these constituents may be added in the form of well-defined simple compounds or in the form of complex mixtures obtained from biological sources. Cultivation is carried out according to conventional methods, and the cyclic peptolides formed during the fermentation may finally be isolated from the culture medium by the use of known chromatographic methods. The cyclic peptolides of the invention may also be obtained by the cultivation of variant or mutant strains obtained by selection or by the effect of mutation-inducing agents e.g. U.V. light, X-rays or chemical mutagens on NRRL 18230 or other strains of Cylindrotrichum Bonorden.


The cyclic peptolides of the present invention may also be prepared by synthetic or semi-synthetic methods, for example by the cyclisation of a linear peptolide or a linear peptide having an -OH terminal group in place of an -NH₂ terminal; or by the replacement of an amide linkage in a natural, synthetic or semi-synthetic cyclosporin with an ester linkage.

The total synthesis of the preferred compounds of formula II may be carried out in a manner analogous to the total synthesis of cyclosporin A and analogues as described for example in European Patent 34 567 or by R. Wenger in Transplantation Proceedings, vol. XV pp. 2230-2241 (1983). According to this method the C-protected heptapeptide having the formula V




in which Bz is the benzyl group and W, X, Y and Z are as defined above is first prepared, and this is then reacted with a tetrapeptide corresponding to the sequence 8 through 11.


This tetrapeptide, of formula VI




contains three normal peptide bonds, but has an O-terminal in place of an N-terminal since the residue at position 8 is derived from an α -hydroxy acid rather than from an α -aminoacid.

The tetrapeptide may be prepared according to the scheme shown in the following flow sheet:






in which BOC is the N-protecting group t-butyloxycarbonyl and R' is a suitable O-protecting group. Thus the reagent represented above as R' - A - OH is an OH-protected α -hydroxycarboxylic acid, which when A is of formula III, has the formula VII



where R is as defined above.

Preferably the group R' is selected from the groups



tBuSi(CH₃)₂- . The preferred compounds of formula VII may be obtained by reacting the α -hydroxy acid, in carbonyl-protected form, e.g. as the benzyl ester, with dihydrofuran, ethoxyethylene, t-butyldimethylchlorosilane or chlorodimethyl ether respectively.

R'-A-MeLeu-Leu-MeVal-W-Thr-X-Y-Z-MeLeu-Ala-OBz VIII

8 9 10 11 1 2 3 4 5 6 7

The heptapeptide of formula V and the tetrapeptide of formula VI may also be obtained by controlled hydrolysis of cyclic peptolides of formula II obtained from fermentation. This treatment with trifluoroacetic acid at low temperature cleaves the bond between residues 11 and 1 to give a linear undecapeptolide containing residues 1 (N-terminal) through 11 (C-terminal), with an ester linkage at 7-8. Alkaline hydrolysis gives the 1-7 heptapeptide and the 8-11 hydroxytetrapeptide. Semisynthetic cyclic peptolides may then be produced for example by reacting the hydroxytetrapeptide produced in this way with a synthetic heptapeptide, or vice versa, and cyclising the linear product.

For the purposes of the cyclisation reaction the peptide may if desired be in O- protected form, that is, the hydroxy groups in the 1-MeBmt or derivative thereof, and/or in the 2-threonine residue may bear protecting groups, as described in European Patent 34 567. Such O-protecting groups are then removed subsequent to ring closure by standard methods. Removal of for example a benzyl group by hydrogenation will also lead to the hydrogenation of MeBmt to MeBmtH₂, and in any case initially-produced cyclic peptolides containing a MeBmt residue at position 1 may be converted to the corresponding MeBmtH₂ compound by hydrogenation.

Accordingly the invention provides a process for the production of a cyclic peptolide of formula II, stated above, which process comprises

- a) removing the O-protecting groups from a cyclic peptolide of formula II in O-protected form;
- b) cyclising a straight chain hydroxy-endcapeptide of formula VIII, in unprotected form or O-protected on one or both of residues 1 and 2, and, when required, carrying out process step (a); and, when desired
- c) hydrogenating a cyclic peptolide of formula II thus obtained wherein W is MeBmt to obtain the corresponding cyclic peptolide wherein W is MeBmtH₂.

The cyclic peptolides of the invention exhibit pharmacological activity and may, therefore, be useful as pharmaceuticals. In particular, the compounds show immunosuppressant, anti-inflammatory and anti-parasitic activity in the following tests:

IN VITRO MODELS (1-3)

1. Interleukin 2 (IL-2) inhibition

Interleukin 2 is induced by mitogen stimulation of mouse spleen cells. Forty eight hour supernatants are collected and assayed for their content of IL-2 by use of a IL-2-dependent cell line (CTLL). The growth of these cells is assayed after 48 hours by use of an enzymatic assay which measures mitochondrial activity.

[T. Mosmann J. Immunol. Methods 65:55-63 (1983)]

The compounds of the invention have an inhibitory effect at concentrations from IC₅₀ 0.01 to approx. 0.1 ug/ml.

2. Proliferative Response of Lymphocytes to Allogeneic Stimulation

Murine Mixed Lymphocyte Reaction (MLR)

Spleen cells (0.5×10^6) from Balb/c mice (female, 8-10 weeks) are co-incubated for 5 days with 0.5×10^6 irradiated (2000 rads) or mitomycin C treated spleen cells from CBA mice (female, 8-10 weeks). The irradiated allogeneic cells induce a proliferative response in the Balb c spleen cells which can be measured by labeled precursor incorporation into the DNA. Since the stimulator cells are irradiated (or mitomycin C treated) they do not respond to the Balb/c cells with proliferation but do retain their antigenicity.

[T. Meo "Immunological Methods", L. Lefkovits and B. Pernis, Eds., Academic Press, N.Y. pp. 227-239 (1979)]

The compounds of the invention have an inhibitory effect at concentrations of from $IC_{50} = 0.0001$ to approx. 0.001 ug/ml .

3. Primary Humoral Immune Response to Sheep Red Blood Cells *in vitro*
(Mishell-Dutton Assay)

Mouse spleen cells (OF1, female, 8-10 weeks, 1×10^7) are co-cultured with sheep erythrocytes (SRBC, 3×10^7) for 3 days in 1 ml final volume in 24 well plates. Lymphocytes are harvested, washed and plated at a density of 1×10^6 cells onto soft agar with fresh antigen (SRBC). Complement (guinea pig serum) is added after a 60-90 minute incubation period and incubation is continued for another 60 minutes after which the test is evaluated by counting the plaques under the microscope. During the 3 day incubation, the lymphocytes are sensitized to the antigen (SRBC). When incubated with antigen again, B-lymphocytes secrete specific antibody which binds to the antigen in the vicinity of the secretory lymphocyte. Addition of complement causes lysis of the antibody-coated erythrocytes yielding a

plaque. Each plaque represents a single antibody-producing cell.

[R.I. Mishell & R.W. Dutton J.Exp. Med. 126:423-442 (1967)]

The suppression of plaque-forming cells is observed at concentrations of compound according to the invention of from IC_{50} 0.01 to approx. 0.1 ug/ml.

IN VIVO MODELS (4-9)

4. Formation of plaque forming cells (humoral immune response)

Female rats (OFA) are immunised by the i.v. injection of (1×10^8) sheep erythrocytes (SRBC) and treated on three consecutive days with the drugs under investigation. Spleen cell suspensions are prepared 6 days after immunisation and 1×10^6 lymphocytes are plated onto soft agar in the presence of indicator cells (SRBC) and complement. Lysis of the indicator cells due to secretion of specific antibody and presence of complement yields plaques. The number of plaques per plate are counted and corrected for the number of plaques per spleen.

[N.K. Jerne & A.A. Nordin Science 140:405 (1969); N.K. Jerne, A.A. Nordin & C. Henry (1963) In: "Cell Bound Antibodies", B. Amos & H. Koprowski, Eds., Wistar Inst. Press, Philadelphia pp. 109-125 (1963)].

The compounds according to the invention produce this effect in the rat when given orally at an ED_{50} of approx. 6.0-8.0 mg/kg.

5. Localised graft-versus-host reaction

Spleen cells (1×10^7) from 6 week old female Wistar/Furth (WF) rats are injected subcutaneously on day 0 into the left hind-paw of female (F344 x WF)FI rats weighing about 100 g. Animals are treated for 4 consecutive days and the popliteal lymph nodes are

removed and weighed on day 7. The difference in weight between the two lymph nodes is taken as the parameter for evaluating the reaction.

[W. L. Ford, W. Burr & M. Simonsen Transplantation 10:258-266 (1970)]

The compounds of the invention have an oral ED₅₀ in this test of approx. 20-30 mg/kg.

6. Freund's Adjuvant Arthritis

OFA and Wistar rats (male or female, 150 g body weight) are injected i.c. at the base of the tail or in the hind paw with 0.1 ml of mineral oil containing 0.6 mg of lyophilized heat-killed Mycobacterium smegmatis. Treatment is started on day 14, when the secondary inflammation is well developed (days 14 - 20). At the end of the experiment, the swelling of the joints is measured by means of a micro-caliper. ED₅₀ is the oral dose in mg/kg orally which reduces the swelling (primary or secondary) to half of that of the controls. For the compounds of the invention the oral ED₅₀ is up to 30 mg/kg.

[C.A. Winter & G.W. Nuss Arthritis and Rheumatism 9:394-404 (1966)]

7. Kidney allograft reaction in the rat

One kidney from a female Fisher 344 rat is transplanted onto the renal vessel of a unilaterally (left side) nephrectomized Wistar/Furth recipient rat using an end-to-end anastomosis. Ureteric anastomosis is also end-to-end. Treatment commences on the day of transplantation and is continued for 14 days. A contralateral nephrectomy is done seven days after transplantation, leaving the recipient relying on the performance of the donor kidney. Survival of the recipient is taken as the

parameter for a functional graft.

[P.C. Hiestand, et al Immunology 55 249-255 (1985)]

The compounds of the invention are effective in the rat at an oral ED₅₀ of from 6 to approx. 9 mg/kg.

8. UV Erythema test

The test is carried out on female albino guinea pigs, approx. 150 g wt. The animals are shaved on both flanks using a Braun micro razor, without causing skin irritation. For each test substance five animals are subjected to a defined intensity of UV radiation for 10 seconds on each of four areas of skin (10 mm diameter). Immediately afterwards 50 microliters of a solution of the test substance in ethanol/propylene glycol/dimethylacetamide (19:19:2 v/v) is applied to two of the irradiated areas on each animal and 50 microliters of the solvent mixture to the other two as controls. Four hours after application, the degree of erythema is estimated visually.

[Raake, W. *Arzneim.-Forsch.* 34(I) No. 4 (1984)]

The compounds of the invention are effective at a concentration of from 1 to 10%.

9. Anti-malaria test

Mice (OF1, male) are infected i.p. on day 0 with 0.2 ml of a suspension of erythrocytes containing 10⁷ cells parasitized by *Plasmodium berghei* (Strain NK 65). The test substance is administered s.c. at varying dosages using 5 to 10 mice/dose. The survival time is recorded, and the minimum effective dose (MED) calculated by comparison of survival time with that for untreated controls. For controls, survival time is approx. 7 days. The MED is the dosage at which survival time is doubled.

[L. Rane in "Chemotherapy and Drug Resistance in Malaria, ed. W. Peters, Academic Press, New York, (1979)]

In view of their immunosuppressive activity, the compounds are indicated for use in the prophylaxis and treatment of conditions and diseases requiring a suppression of the immune response, for example in the treatment of autoimmune diseases, the prevention of tissue rejection in transplantation, e.g. bone marrow and kidney transplantation. Specific autoimmune diseases, for which the compounds of the invention may be used include aplastic anaemia, pure red cell anaemia, idiopathic thrombocytopenia, systemic lupus erythematosus, polychondritis, scleroderma, Wegner's granulomatosis, dermatomyositis, chronic active hepatitis, myasthenia gravis, Steven-Johnson syndrome, psoriasis, idiopathic sprue, Crohn's disease, Graves' ophthalmopathy, sarcoidosis, multiple sclerosis, primary biliary cirrhosis, primary juvenile diabetes, posterior uveitis, interstitial pulmonary fibrosis and psoriatic arthritis.

Because of their antiinflammatory properties the compounds of the invention are indicated for use in the treatment of inflammatory conditions, particularly inflammatory states with an aetiology which includes an autoimmune component, for example the treatment of arthritis and rheumatic diseases such as chronic progressive arthritis.

In view of their anti-parasitic activity, the compounds are also indicated for use in the treatment of parasitic disease, for example schistosomiasis, filariasis, leishmaniasis, coccidioidomycosis and, in particular, malaria.

The compounds of the invention are also indicated for use in the treatment of certain skin diseases and conditions, which include, in addition to those already mentioned above, alopecia areata, urticaria, vasculitis, erythema, atopic dermatitis, eczema, cutaneous eosinophilia and angioderma.

For these indications, an indicated daily dosage is in the range from about 70 to about 3000 mg of a compound according to the invention conveniently administered in divided doses up to four times a day.

The compounds of the invention may be administered by any conventional route, in particularly orally, for example in the form of tablets or capsules, parenterally in the form of injectable solutions or suspensions or topically in the form of a lotion, cream or gel.

The present invention also provides pharmaceutical compositions containing a compound according to the invention in association with at least one pharmaceutical carrier or diluent. Such composition may be manufactured in conventional manner. Unit dosage forms contain, for example, from about 20 mg to about 1500 mg of a compound according to the invention.

The following Examples, in which all temperatures are in degrees centigrade, illustrate the invention:

Example 1

Culture of strain NRRL 18230 in Erlenmeyer flask

Starting cultures of the strain NRRL 18230 are incubated at 21° for 14 days on MA in slants. A preculture is then produced by homogenising the complete contents of one starting culture under sterile conditions, transferring to a 500 ml Erlenmeyer flask containing 200 ml nutrient solution M (2% malt extract, 0.4% yeast extract in demineralized water), and incubating on a rotating shaker at 200 rpm for 10 days at 21°.

Intermediate cultures are then obtained by transferring 20 ml of the preculture to a 500 ml Erlenmeyer flask containing 200 ml of nutrient solution M and incubating on a rotary shaker at 200 rpm for 7 days at 21°.

Finally for a production culture, 100 Erlenmeyer flasks (500 ml) each containing 200 ml of nutrient solution M are each inoculated with 20 ml of the intermediate cultures. The flasks are incubated at 21° in a rotary shaker at 200 rpm. After 10 days the 20 litres of fermentation broth are combined for the extraction of the product.

Example 2

Isolation of (Thr)² (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin

The 20 litres of culture medium obtained from Example 1 are subjected to high-shear mixing in a rod mixer (Lutz, Wertheim, Germany) to break up the cells, then extracted three times with 20 litres of ethyl acetate. The 60 litres of organic phase are combined, dried over anhydrous sodium sulphate and evaporated to dryness under vacuum, giving a residue of 17.4 g.

The residue is dissolved in 80 ml methanol and chromatographed on a 90 mm diameter column containing 1300 g Sephadex LH-20 (Pharmacia

Fine Chemicals AB, Uppsala, Sweden). Fractions of 100 ml are taken, whereby (Thr)² (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin appears in fractions 22-30. These are combined and evaporated under vacuum to give 8400 mg of a light coloured solid foam. This residue is dissolved in wet ethyl acetate and chromatographed on a 55 mm diameter column containing 500 g Kieselgel (Merck) of particle size 0.04 - 0.063 mm. The desired product is detected in fractions 9-14 (100 ml fraction size). Evaporation gives a pale yellow residue (4200 mg). Treatment with decolorizing charcoal (Merck) in diethyl ether and filtration through a thin layer of talc gives (Thr)² (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin as a white powder, recrystallized from ether m.p. 163-164° (decomp.); $[\alpha]^{20} = -186^\circ$, (c = 1 in MeOH); $[\alpha]^{20} = -223^\circ$ (c = 1 in CHCl₃).

Examples 3-7

By modification of the above chromatographic techniques, the following compounds may be isolated in minor amounts from the production fermentation broth:

Ex. No.	Compound
3	(Thr) ² (D-Hiv) ⁸ (Leu) ¹⁰ -Ciclosporin
4	(Thr) ² (Ile) ⁵ (D-Hiv) ⁸ (Leu) ¹⁰ -Ciclosporin
5	(Thr) ² (Leu) ⁴ (Leu) ⁵ (D-Hiv) ⁸ (Leu) ¹⁰ -Ciclosporin
6	(Thr) ² (Gly) ³ (Leu) ⁵ (D-Hiv) ⁸ (Leu) ¹⁰ -Cyclosporin
7	(8'-OHMeBmt) ¹ (Thr) ² (Leu) ⁵ (D-Hiv) ⁸ (Leu) ¹⁰ -Ciclosporin

The compounds have the properties shown in Table I:

Table I

Compound of Ex. No.	formula	MW calc	MW found*	mp °C	$[\alpha]^{20}$ (c=1 in MeOH)
3	$C_{63}H_{112}N_{10}O_{14}$	1233.6	1233.9	164 (dec)	-184°
4	$C_{64}H_{114}N_{10}O_{14}$	1247.7	1247.6	151.3 (dec)	-180°
5	$C_{63}H_{112}N_{10}O_{14}$	1233.6	1233.6	166.8 (dec)	-174°
6	$C_{63}H_{112}N_{10}O_{14}$	1233.6	1233.6	140.43 (dec)	-154°
7	$C_{64}H_{114}N_{14}O_{15}$	1263.7	1263	158 (sinters) 195 (dec)	-174°

* by FAB (Fast Atomic Bombardment) mass spectrometry

Example 8

Synthesis of (Thr)² (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin by cyclisation

At room temperature, 2.4 g (2.29 mmol) of the unprotected hydroxy-undecapeptide

HO-(D)Hiv-MeLeu-Leu-MeVal-MeBmt-Thr-Sar-MeLeu-Leu-MeLeu-Ala-OH,

0.84 g (6.88 mmol) of 4-dimethylaminopyridine and 0.66 g (1.5 equiv, 3.44 mmol) of N-ethyl-N-(3-dimethylamino)propyl carbodiimide are dissolved in 145 ml methylene chloride and reacted for three days with stirring and exclusion of moisture. The resulting solution is diluted with 300 ml of methylene chloride, shaken with 100 ml of water acidified to pH 2 with 1N HCl, filtered through talc and evaporated. The residue is chromatographed on 100 g of silica gel using 10% MeOH/CH₂Cl₂, to give the title compound (2.55 g, 89%). Identity with the compound of Example 2 is established by NMR spectroscopy and $[\alpha]^{20}$ (-220°, c=1 in CHCl₃).


Table II

Abbreviations

Abu	α -aminobutyric acid
Ala	alanine
BOC	t-butyloxycarbonyl
Bz	benzyl
Gly	glycine
Hiv	α -hydroxyisovaleric acid
Ile	isoleucine
Leu	leucine
MeBmt	N-methyl-(4R)-4-but-2E-en-1-yl-4-methyl-(L)-threonine
MeBmtH ₂	N-methyl-(4R)-4-but-1-yl-4-methyl-(L)-threonine
8'OHMeBmt	N-methyl-(4R)-4-(4'-hydroxybut-2E-en-1-yl)-4-methyl-(L)-threonine
Sar	sarcosine
Thr	threonine
Val	valine


THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS

1. A cyclic peptolide which has the structure of a cyclosporin in which one amide linkage is replaced by an ester linkage.
2. A cyclic peptolide according to Claim 1 which is composed of one MeBmt residue, or a derivative thereof, 9 other α -amino acid residues and one α -hydroxyacid residue.
3. A cyclic peptolide according to Claim 2 which has an α -hydroxycarboxylic acid at position 8.
4. A cyclic peptolide according to Claim 3 of formula II



in which W is MeBmt, 8'-OHMeBmt or MeBmtH₂,
X is Sar or Gly,
Y is MeLeu or Leu,
Z is Leu, Ile or Val,
and A is the residue of an α -hydroxycarboxylic acid.

5. A cyclic peptolide according to Claim 4 in which A is a group of formula III



in which R is C₁₋₄ alkyl.



6. A cyclic peptolide according to Claim 5 in which R is isopropyl.

7. (Thr)² (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin.

8. (Thr)² (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin.

9. (Thr)² (Ile)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin.

10. (Thr)² (Leu)⁴ (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin.

11. (Thr)² (Gly)³ (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin.

12. (8'-OH MeBmt)¹ (Thr)² (Leu)⁵ (D-Hiv)⁸ (Leu)¹⁰-Ciclosporin.

13. A process for the preparation of a cyclic peptolide according to any one of Claims 4-12 characterised by the step of cultivating a fungal strain of the genus Cylindrotrichum Bonorden which produces said cyclic peptolide in the presence of a nutrient medium.

14. A process for the preparation of a cyclic peptolide according to any one of Claims 7-12 characterized by the step of cultivating the strain NRRL 18230, or a mutant or derivative thereof, in the presence of a nutrient medium.

15. A fermentation broth obtained from the cultivation of a cyclic peptolide-producing fungal strain of the genus Cylindrotrichum Bonorden, wherein the cyclic peptolide is one of the compounds according to any one of the claims 1 to 12.

16. A fermentation broth obtained from the cultivation of the strain NRRL 18230, or a mutant or derivative thereof.



17. A pharmaceutical composition containing a cyclic peptolide according to any one of Claims 1-12 accompanied by a pharmaceutically acceptable diluent or carrier.

18. A biologically pure culture of the fungal strain NRRL 18230.

19. A process for the preparation of a cyclic peptolide according to any one of Claims 1-12 by the cyclization of a linear peptolide or of a peptide having a hydroxy terminal in place of an amino terminal group.

20. A process for the preparation of a cyclic peptolide of formula II, stated in Claim 4, which process comprises

- a) removing the O-protecting groups from a cyclic peptolide of formula II in O-protected form;
- b) cyclising a straight chain hydroxy-undecapeptide of formula VIII, in unprotected form or O-protected on one or both of residues 1 and 2, and, when required, carrying out process step (a); and, when desired
- c) hydrogenating a cyclic peptolide of formula II thus obtained wherein W is MeBmt to obtain the corresponding cyclic peptolide wherein W is MeBmtH₂.

DATED THIS 24th day of October, 1991

SANDOZ LTD
By Its Patent Attorneys
DAVIES & COLLISON

