



HU000034876T2

(19) **HU**(11) Lajstromszám: **E 034 876**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala**EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA

- (21) Magyar ügyszám: **E 13 703254**
- (22) A bejelentés napja: **2013. 01. 25.**
- (96) Az európai bejelentés bejelentési száma:
EP 20130703254
- (97) Az európai bejelentés közzétételi adatai:
EP 2806863 A2 **2013. 08. 01.**
- (97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2806863 B1 **2017. 08. 09.**
- (51) Int. Cl.: **A61K 31/00** (2006.01)
A61P 5/00 (2006.01)
A61P 43/00 (2006.01)
A61K 31/343 (2006.01)
- (86) A nemzetközi (PCT) bejelentési szám:
PCT/US 13/023312
- (87) A nemzetközi közzétételi szám:
WO 13112949

<p>(30) Elsőbbségi adatok:</p> <table><tr><td>201261590974 P</td><td>2012. 01. 26.</td><td>US</td></tr><tr><td>201261640067 P</td><td>2012. 04. 30.</td><td>US</td></tr><tr><td>201261650458 P</td><td>2012. 05. 22.</td><td>US</td></tr><tr><td>201261650455 P</td><td>2012. 05. 22.</td><td>US</td></tr><tr><td>201261714149 P</td><td>2012. 10. 15.</td><td>US</td></tr><tr><td>201261738985 P</td><td>2012. 12. 18.</td><td>US</td></tr><tr><td>201261738987 P</td><td>2012. 12. 18.</td><td>US</td></tr><tr><td>201361755896 P</td><td>2013. 01. 23.</td><td>US</td></tr></table>	201261590974 P	2012. 01. 26.	US	201261640067 P	2012. 04. 30.	US	201261650458 P	2012. 05. 22.	US	201261650455 P	2012. 05. 22.	US	201261714149 P	2012. 10. 15.	US	201261738985 P	2012. 12. 18.	US	201261738987 P	2012. 12. 18.	US	201361755896 P	2013. 01. 23.	US	<p>(73) Jogosult(ak): Vanda Pharmaceuticals Inc., Washington, DC 20037 (US)</p> <p>(74) Képviseelő: Danubia Szabadalmi és Jogi Iroda Kft., Budapest</p>
201261590974 P	2012. 01. 26.	US																							
201261640067 P	2012. 04. 30.	US																							
201261650458 P	2012. 05. 22.	US																							
201261650455 P	2012. 05. 22.	US																							
201261714149 P	2012. 10. 15.	US																							
201261738985 P	2012. 12. 18.	US																							
201261738987 P	2012. 12. 18.	US																							
201361755896 P	2013. 01. 23.	US																							
<p>(72) Feltaláló(k): DRESSMAN, Marlene, Michelle, Germantown, MD 20874 (US) FEENEY, John, Joseph, Olney, MD 20832 (US) LICAMELE, Louis, William, Potomac, MD 20854 (US) POLYMEROPOULOS, Mihael, H., Potomac, MD 20854 (US)</p>																									

(54) **Cirkadián ritmussal kapcsolatos rendellenességek kezelése**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmat az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 2 806 863 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
09.08.2017 Bulletin 2017/32

(51) Int Cl.:
A61K 31/00 ^(2006.01) **A61K 31/343** ^(2006.01)
A61P 5/00 ^(2006.01) **A61P 43/00** ^(2006.01)

(21) Application number: **13703254.6**

(86) International application number:
PCT/US2013/023312

(22) Date of filing: **25.01.2013**

(87) International publication number:
WO 2013/112949 (01.08.2013 Gazette 2013/31)

(54) TREATMENT OF CIRCADIAN RHYTHM DISORDERS

BEHANDLUNG VON STÖRUNGEN DER CIRCADIENEN RHYTHMIK

TRAITEMENT DE TROUBLES DU RYTHME CIRCADIEN

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(30) Priority: **26.01.2012 US 201261590974 P**
30.04.2012 US 201261640067 P
22.05.2012 US 201261650458 P
22.05.2012 US 201261650455 P
15.10.2012 US 201261714149 P
18.12.2012 US 201261738985 P
18.12.2012 US 201261738987 P
23.01.2013 US 201361755896 P

(43) Date of publication of application:
03.12.2014 Bulletin 2014/49

(73) Proprietor: **Vanda Pharmaceuticals Inc.**
Washington, DC 20037 (US)

(72) Inventors:

- **DRESSMAN, Marlene, Michelle**
Germantown, MD 20874 (US)
- **FEENEY, John, Joseph**
Olney, MD 20832 (US)
- **LICAMELE, Louis, William**
Potomac, MD 20854 (US)
- **POLYMERPOULOS, Mihael, H.**
Potomac, MD 20854 (US)

(74) Representative: **Carpmaels & Ransford LLP**
One Southampton Row
London WC1B 5HA (GB)

(56) References cited:
WO-A1-2007/137244

- **Robert L Sack ET AL: "ENTRAINMENT OF FREE-RUNNING CIRCADIAN RHYTHMS BY MELATONIN IN BLIND PEOPLE", N Engl J Med 2000;343, 10 December 2000 (2000-12-10), pages 1070-1077, XP055057908, Retrieved from the Internet:
URL: <http://www.nejm.org/doi/pdf/10.1056/NEJM200010123431503> [retrieved on 2013-03-26]**
- **OKAWA ET AL: "Circadian rhythm sleep disorders: Characteristics and entrainment pathology in delayed sleep phase and non-24 sleep-wake syndrome", SLEEP MEDICINE REVIEWS, W.B. SAUNDERS, AMSTERDAM, NL, vol. 11, no. 6, 25 October 2007 (2007-10-25), pages 485-496, XP022341295, ISSN: 1087-0792, DOI: 10.1016/J.SMRV.2007.08.001**
- **UCHIYAMA M ET AL: "Non-24-Hour Sleep-Wake Syndrome in Sighted and Blind Patients", SLEEP MEDICINE CLINICS, W.B. SAUNDERS CO, US, vol. 4, no. 2, 1 June 2009 (2009-06-01), pages 195-211, XP008161124, ISSN: 1556-407X, DOI: 10.1016/J.JSMC.2009.02.002**
- **RAJARATNAM S M ET AL: "Melatonin agonist tasimelteon (VEC-162) for transient insomnia after sleep-time shift: two randomised controlled multicentre trials", THE LANCET, LANCET LIMITED. LONDON, GB, vol. 373, no. 9662, 7 February 2009 (2009-02-07), pages 482-491, XP025913508, ISSN: 0140-6736, DOI: 10.1016/S0140-6736(08)61812-7 [retrieved on 2009-02-05]**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 806 863 B1

- HARDELAND RUEDIGER: "Tasimelteon, a melatonin agonist for the treatment of insomnia and circadian rhythm sleep disorders", CURRENT OPINION IN INVESTIGATIONAL DRUGS, THOMSON REUTERS (SCIENTIFIC) LTD, LONDON, UK, vol. 10, no. 7, 1 July 2009 (2009-07-01), pages 691-701, XP008161049, ISSN: 2040-3429
- HARDELAND R: "New approaches in the management of insomnia: Weighing the advantages of prolonged-release melatonin and synthetic melatonergic agonists", NEUROPSYCHIATRIC DISEASE AND TREATMENT, DOVE MEDICAL PRESS (NZ) LTD, NZ, vol. 5, no. 1, 1 January 2009 (2009-01-01) , pages 341-354, XP009137641, ISSN: 1176-6328 [retrieved on 2009-06-10]
- WALDEMAR OCKERT: "A new dawn in the sleep disorders pipeline?", NATURE REVIEWS DRUG DISCOVERY, vol. 11, no. 8, 20 July 2012 (2012-07-20), pages 595-596, XP055057972, ISSN: 1474-1776, DOI: 10.1038/nrd3789

Description

Field of the Invention

5 **[0001]** Embodiments of the invention relate generally to the field of circadian rhythm disorders (CRDs) and, more particularly, to the entrainment of circadian rhythms in persons afflicted with Non-24 Hour Disorder (Non-24).

Background of the Invention

10 **[0002]** The master body clock controls the timing of many aspects of physiology, behavior and metabolism that show daily rhythms, including the sleep-wake cycles, body temperature, alertness and performance, metabolic rhythms and certain hormones which exhibit circadian variation. Outputs from the suprachiasmatic nucleus (SCN) control many endocrine rhythms including those of melatonin secretion by the pineal gland as well as the control of cortisol secretion via effects on the hypothalamus, the pituitary and the adrenal glands. This master body clock, located in the SCN, spontaneously generates rhythms of approximately 24.5 hours. These non-24-hour rhythms are synchronized each day to the 24-hour day-night cycle by light, the primary environmental time cue which is detected by specialized cells in the retina and transmitted to the SCN via the retino-hypothalamic tract. Inability to detect this light signal, as occurs in most totally blind individuals, leads to the inability of the master body clock to be reset daily and maintain entrainment to a 24-hour day.

20

Non-24-Hour Disorder

25 **[0003]** Non-24, also referred to as Non-24-Hour Sleep-Wake Disorder (N24HSWD) or Non-24-Hour Disorder, is an orphan indication affecting approximately 65,000 to 95,000 people in the U.S. and 140,000 in Europe. Non-24 occurs when individuals, primarily blind with no light perception, are unable to synchronize their endogenous circadian pacemaker to the 24-hour light/dark cycle. Without light as a synchronizer, and because the period of the internal clock is typically a little longer than 24 hours, individuals with Non-24 experience their circadian drive to initiate sleep drifting later and later each day. Individuals with Non-24 have abnormal night sleep patterns, accompanied by difficulty staying awake during the day. Non-24 leads to significant impairment, with chronic effects impacting the social and occupational functioning of these individuals.

30

[0004] In addition to problems sleeping at the desired time, individuals with Non-24 experience excessive daytime sleepiness that often results in daytime napping.

35

[0005] The severity of nighttime sleep complaints and/or daytime sleepiness complaints varies depending on where in the cycle the individual's body clock is with respect to their social, work, or sleep schedule. The "free running" of the clock results in approximately a 1-4 month repeating cycle, the circadian cycle, where the circadian drive to initiate sleep continually shifts a little each day (about 15 minutes on average) until the cycle repeats itself. Initially, when the circadian cycle becomes desynchronous with the 24h day-night cycle, individuals with Non-24 have difficulty initiating sleep. As time progresses, the internal circadian rhythms of these individuals becomes 180 degrees out of synchrony with the 24h day-night cycle, which gradually makes sleeping at night virtually impossible, and leads to extreme sleepiness during daytime hours.

40

[0006] Eventually, the individual's sleep-wake cycle becomes aligned with the night, and "free-running" individuals are able to sleep well during a conventional or socially acceptable time. However, the alignment between the internal circadian rhythm and the 24-hour day-night cycle is only temporary. In addition to cyclical nighttime sleep and daytime sleepiness problems, this condition can cause deleterious daily shifts in body temperature and hormone secretion, may cause metabolic disruption and is sometimes associated with depressive symptoms and mood disorders.

45

[0007] It is estimated that 50-75% of totally blind people in the United States (approximately 65,000 to 95,000) have Non-24. This condition can also affect sighted people. However, cases are rarely reported in this population, and the true rate of Non-24 in the general population is not known.

50

[0008] The ultimate treatment goal for individuals with Non-24 is to entrain or synchronize their circadian rhythms into an appropriate phase relationship with the 24-hour day so that they will have increased sleepiness during the night and increased wakefulness during the daytime.

55

[0009] Uchiyama, M., et. al., (2009) Sleep Medicines Clinics, W.B Saunders Co, US, vol4, no. 2, 195-211, discloses the clinical characteristics of patients who have non-24-hour sleep-wake syndrome and discusses the biologic mechanisms that may underlie its development, and describes potential treatment strategies.

55

Tasimelteon

[0010] Tasimelteon is a circadian regulator which binds specifically to two high affinity melatonin receptors, Mella

(MT1R) and Mel1b (MT2R). These receptors are found in high density in the suprachiasmatic nucleus of the brain (SCN), which is responsible for synchronizing our sleep/wake cycle. Tasimelteon has been shown to improve sleep parameters in prior clinical studies, which simulated a desynchronization of the circadian clock. Tasimelteon has so far been studied in hundreds of individuals and has shown a good tolerability profile.

5

Summary of the Invention

[0011] Embodiments of the invention relate to the discovery that a composition comprising tasimelteon can be for use in the treatment of a free running circadian rhythm, in patients, including light perception impaired patients, e.g., blind patients, in whom such free running circadian rhythm manifests itself as Non-24.

10

[0012] The present invention is set out in the appended claims herein below. Disclosed is a method for determining a person's circadian rhythm (*tau*) and to the application of such methodology to the treatment of a free running circadian rhythm.

15

[0013] Further disclosed is the treatment of subjects who present with symptoms of Non-24, specifically, e.g., sleep drifting later each day, abnormal night sleep patterns, and/or difficulty staying awake during the day, leading in many cases to significant impairment, with chronic effects impacting the social and occupational functioning of these individuals, as well as possible negative health effects of chronic misalignment.

20

[0014] Also disclosed is a method of treating Non-24 in a patient suffering therefrom, said method comprising internally administering to the patient an effective amount of a melatonin agonist, e.g., tasimelteon, including, such method wherein the patient is light perception impaired (LPI) including, again without limitation, patients who have zero light perception, i.e., patients who are totally blind.

25

[0015] Further disclosed is a method of entraining a patient suffering from Non-24 to a 24 hour sleep-wake cycle in which the patient awakens at or near a target wake time following a daily sleep period, e.g. approximately 7 to 9 hours (understanding, of course, that the patient may not actually sleep during the entire sleep period), said method comprising: treating the patient by internally administering to the patient an effective amount of a melatonin agonist.

30

[0016] The melatonin agonist may be tasimelteon or an active metabolite thereof or a pharmaceutically acceptable salt of such metabolite. The melatonin agonist may have a pharmacological profile similar to tasimelteon, e.g., affinity for both MT1R and MT2R with greater affinity, e.g., 2-fold to 4-fold, greater affinity for MT2R and a t_{max} of less than 2 hours, e.g., one hour or less.

35

[0017] Also disclosed is a method for the chronic treatment of Non-24 in a person who is totally blind comprising orally administering to the person tasimelteon in an amount of 20 to 50 mg once daily about 1/2 hour to about 1-1/2 hours before a target bedtime. Patients who were previously treated with a melatonin agonist and entrained to a 24 hour circadian rhythm are maintained by ongoing daily internally administering to the patients an effective amount of a melatonin agonist, e.g., a melatonin agonist having a t_{max} of less than about 2 hours such as tasimelteon, e.g., at between about 0.5 and 1.5 hours prior to a daily sleep period of between about 7 hours and about 9 hours.

Brief Description of the Figures

40

[0018]

FIG. 1 is an example of a patient report for a patient determined not to have a free-running circadian rhythm based on aMT6s analyses.

FIG. 2 is an example of a patient report for a patient determined to have a free-running circadian rhythm based on aMT6s analyses.

45

FIG. 3 is an example of a patient report for a patient determined not to have a free-running circadian rhythm based on cortisol analyses.

FIG. 4 is an example of a patient report for a patient determined to have a free-running circadian rhythm based on cortisol analyses.

50

FIG. 5 shows a metabolic pathway of tasimelteon and several of its metabolites.

FIGS. 6-11 show plots of the effect of co-administration of tasimelteon and fluvoxamine on the concentration of, respectively, tasimelteon, the M9 metabolite, the M11 metabolite, the M12 metabolite, the M13 metabolite, and the M14 metabolite.


FIGS. 12-17 show plots of the effect of smoking on the concentration of, respectively, tasimelteon, the M9 metabolite, the M11 metabolite, the M12 metabolite, the M13 metabolite, and the M14 metabolite.

55

Detailed Description of the Invention

[0019] Tasimelteon has the chemical name: *trans*-N-[[2-(2,3-dihydrobenzofuran-4-yl)cycloprop-yl]methyl]propana-

mide, has the structure of Formula I:




Formula I

15 and is disclosed in US 5856529 and in US 20090105333.


20 **[0020]** Tasimelteon is a white to off-white powder with a melting point of about 78°C (DSC) and is very soluble or freely soluble in 95% ethanol, methanol, acetonitrile, ethyl acetate, isopropanol, polyethylene glycols (PEG-300 and PEG-400), and only slightly soluble in water. The native pH of a saturated solution of tasimelteon in water is 8.5 and its aqueous solubility is practically unaffected by pH. Tasimelteon has 2-4 times greater affinity for MT2R relative to MT1R. Its affinity (K_i) for MT1R is 0.3 to 0.4 and for MT2R, 0.1 to 0.2. Tasimelteon is useful in the practice of this invention because it is a melatonin agonist that has been demonstrated, among other activities, to entrain patients suffering from Non-24.

25 **[0021]** In related aspects, this invention relates to the use of a tasimelteon metabolite as the melatonin agonist. Tasimelteon metabolites include, for example, a phenol-carboxylic acid analog (M9) and a hydroxypropyl-phenol analog (M11). Each is formed in humans following oral administration of tasimelteon.

30 **[0022]** Specifically, aspects of the invention encompass use of tasimelteon or of compounds of Formulas II or III, including salts, solvates, and hydrates of tasimelteon or of compounds of Formula II or Formula III, in amorphous or crystalline form.



Formula II (M11)




Formula III (M9)


50 **[0023]** While depicted herein in the R-trans configuration, the invention nevertheless comprises use of stereoisomers thereof, i.e., R-cis, S-trans, and S-cis. In addition, the invention comprises use of prodrugs of tasimelteon or of compounds of Formula II or of Formula III, including, for example, esters of such compounds. The discussion that follows will refer to tasimelteon but it is to be understood that the compounds of Formula II and III are also useful in the practice of aspects of the invention.

55 **[0024]** Metabolites of tasimelteon include, for example, those described in "Preclinical Pharmacokinetics and Metabolism of BMS-214778, a Novel Melatonin Receptor Agonist" by Vachharajani et al., J. Pharmaceutical Sci., 92(4):760-772. The active metabolites of tasimelteon can also be used in the method of this invention, as can pharmaceutically acceptable salts of tasimelteon or of its active metabolites. For example, in addition to metabolites of Formula II and III, above,


metabolites of tasimelteon also include the monohydroxylated analogs M13 of Formula IV, M12 of Formula V, and M14 of Formula VI.



Formula IV



Formula V



Formula VI

40 **[0025]** Thus, it is apparent that this invention contemplates entrainment of patients suffering free running circadian rhythm to a 24 hour circadian rhythm by administration of a circadian rhythm regulator (i.e., circadian rhythm modifier) capable of phase advancing and/or entraining circadian rhythms, such as a melatonin agonist like tasimelteon or an active metabolite of tasimelteon or a pharmaceutically acceptable salt thereof. Other MT1R and MT2R agonists, i.e., melatonin agonists, can have similar effects on the master body clock. So, for example, this invention further contemplates

45 the use of melatonin agonists such as melatonin, N-[1-(2,3-dihydrobenzofuran-4-yl)pyrrolidin-3-yl]-N-ethylurea and structurally related compounds as disclosed in US 6,211,225, LY-156735 ((R)-N-(2-(6-chloro-5-methoxy-1H-indol-3-yl)propyl)acetamide) (disclosed in U.S. Patent No. 4,997,845), agomelatine (N-[2-(7-methoxy-1-naphthyl)ethyl]acetamide) (disclosed in U.S. Patent No. 5,225,442), ramelteon ((S)-N-[2-(1,6,7,8-tetrahydro-2H-indeno- [5,4-b] furan-8-yl)ethyl]propionamide), 2-phenylmelatonin, 8-M-PDOT, 2-iodomelatonin, and 6-chloromelatonin.

50 **[0026]** Additional melatonin agonists include, without limitation, those listed in U.S. Patent Application Publication No. 20050164987, specifically: TAK-375 (see Kato, K. et al. Int. J. Neuropsychopharmacol. 2000, 3 (Suppl. 1): Abst P.03.130; see also abstracts P.03.125 and P.03.127), CGP 52608 (1-(3-allyl-4-oxothiazolidine-2-ylidene)-4-methylthiosemicarbazone) (See Missbach et al., J. Biol. Chem. 1996, 271, 13515-22), GR196429 (N-[2-[2,3,7,8-tetrahydro-1H-furo(2,3-g)indol-1-yl]ethyl]acetamide) (see Beresford et al., J. Pharmacol. Exp. Ther. 1998, 285, 1239-1245), S20242 (N-[2-(7-methoxy naph-1-yl)ethyl]propionamide) (see Depres-Brummer et al., Eur. J. Pharmacol. 1998, 347, 57-66), S-23478 (see Neuropharmacology July 2000), S24268 (see Naunyn Schmiedebergs Arch. June 2003), S25150 (see Naunyn Schmiedebergs Arch. June 2003), GW-290569, luzindole (2-benzyl-N-acetyltryptamine) (see U.S. Patent No. 5,093,352), GR135531 (5-methoxycarbonylamino-N-acetyltrypt-amine) (see U.S. Patent Application Publication No. 20010047016),

Melatonin Research Compound A, Melatonin Agonist A (see IMSWorld R&D Focus August 2002), Melatonin Analogue B (see Pharmaprojects August 1998), Melatonin Agonist C (see Chem. Pharm. Bull. (Tokyo) January 2002), Melatonin Agonist D (see J. Pineal Research November 2000), Melatonin Agonist E (see Chem. Pharm. Bull. (Tokyo) February 2002), Melatonin Agonist F (see Reprod. Nutr. Dev. May 1999), Melatonin Agonist G (see J. Med. Chem. October 1993), Melatonin Agonist H (see Famaco March 2000), Melatonin Agonist I (see J. Med. Chem. March 2000), Melatonin Analog J (see Bioorg. Med. Chem. Lett. March 2003), Melatonin Analog K (see MedAd News September 2001), Melatonin Analog L, AH-001 (2-acetamido-8-methoxytetralin) (see U.S. Patent No. 5,151,446), GG-012 (4-methoxy-2-(methylene propylamide)indan) (see Drijfhout et al., Eur. J. Pharmacol. 1999, 382, 157-66), Enol-3-IPA, ML-23 (N-2,4-dinitrophenyl-5-methoxy-tryptamine) (see U.S. Patent No. 4,880,826), SL-18.1616, IP-100-9 (US 5580878), Sleep Inducing Peptide A, AH-017 (see U.S. Patent No. 5,151,446), AH-002 (8-methoxy-2-propionamido-tetralin) (see U.S. Patent No. 5,151,446), and IP-101. Metabolites, prodrugs, stereoisomers, polymorphs, hydrates, solvates, and salts of the above compounds that are directly or indirectly active can, of course, also be used in the practice of this invention.

[0027] Melatonin agonists with a MT1R and MT2R binding profile similar to that of tasimelteon, which has 2 to 4 time greater specificity for MT2R, are preferred.

[0028] Tasimelteon can be synthesized by procedures known in the art. The preparation of a 4-vinyl-2,3-dihydrobenzofuran cyclopropyl intermediate can be carried out as described in US7754902.

[0029] Pro-drugs, e.g., esters, and pharmaceutically acceptable salts can be prepared by exercise of routine skill in the art.

[0030] In patients suffering a Non-24, the melatonin and cortisol circadian rhythms and the natural day/night cycle become desynchronized. For example, in patients suffering from a free-running circadian rhythm, melatonin and cortisol acrophases occur more than 24 hours, e.g., >24.1 hours, prior to each previous day's melatonin and cortisol acrophase, respectively, resulting in desynchronization for days, weeks, or even months, depending upon the length of a patient's circadian rhythm, before the melatonin, cortisol, and day/night cycles are again temporarily synchronized.

[0031] Chronic misalignment of cortisol has been associated with metabolic, cardiac, cognitive, neurologic, neoplastic, and hormonal disorders. Such disorders include, e.g., obesity, depression, neurological impairments.

[0032] This invention shows that entrainment of the melatonin circadian rhythm is linked to entrainment of the cortisol circadian rhythm.

[0033] Thus, in one aspect, an illustrative embodiment of the invention provides a method of entraining a patient suffering from an abnormal melatonin circadian rhythm, or an abnormal cortisol circadian rhythm, to a 24 hour circadian rhythm by internally administering to the patient an effective amount of a melatonin agonist, in particular, tasimelteon or an active metabolite thereof.

[0034] In related aspects, this invention provides a method of preventing or treating disorders associated with a desynchronous melatonin or cortisol circadian rhythm, i.e., a circadian rhythm that is not synchronized with the natural day/night cycle. Such method comprises internally administering to a patient having a desynchronous melatonin or cortisol circadian rhythm an effective amount of a melatonin agonist, in particular, tasimelteon or an active metabolite thereof, as described in this specification.

[0035] The method of treating Non-24 (which includes phase advancing and/or entraining melatonin and/or cortisol circadian rhythm) in a patient suffering therefrom by internally administering an effective amount of tasimelteon as described in this specification tends to be effective more often in patients having higher amounts of endogenous melatonin. In other words, the likelihood of efficacy of treatment is related to the amount of melatonin naturally present in the patient's body.

[0036] The method of treating Non-24 (which includes phase advancing melatonin and/or cortisol circadian rhythm) in a patient suffering therefrom by internally administering an effective amount of tasimelteon as described in this specification tends to be effective more often in patients whose pre-treatment circadian rhythm (i.e., *tau*) is below a certain threshold. Such threshold can be, e.g., 25.0 hours, 24.9 hours, 24.8 hours, 24.7 hours, 24.65 hours, or 24.6 hours, such that the likelihood of efficacy of treatment is greater in the case of patients whose *tau* is below the threshold.

[0037] In accordance with this invention, a regulatory agency, a patient, a healthcare provider, or an insurance provider, or any one or more of such entities or persons, can choose a likelihood of efficacy that is sufficient to support initiation of treatment with a melatonin agonist, in particular, tasimelteon. For example, it may be decided that if the likelihood of efficacy is less than a selected threshold probability, then the patient should not be treated with the melatonin agonist.

[0038] Alternatively, such threshold probability can be used as a factor in determining whether or not to apply a heightened standard of monitoring for efficacy and/or adverse events. For example, it may be decided that if the likelihood of efficacy is less than a selected threshold probability, then the patient will be examined for signs of efficacy and/or adverse events within about 6 to 9 weeks following initiation of treatment. Such heightened monitoring can also comprise more frequent monitoring and/or decreased tolerance for lack of apparent efficacy or for occurrence of side effects. For example, if there is no or scant evidence of efficacy or if there are signs of adverse events, perhaps even minor or early signs, then the melatonin agonist treatment may be discontinued or modified. Heightened monitoring may include requiring a patient to maintain a sleep diary which would include, e.g., the patient's recordation of sleep and wake

times, frequency and duration of naps, sleep latency, duration of nighttime sleep, such recordation being, e.g., in writing, digitally, or telephonically.

[0039] Efficacy for these purposes can be determined in a number of ways, including, e.g., by determining a patient's *tau* after initiation of therapy and following at least one complete circadian cycle during which the patient has been treated, e.g., about 6 to about 9 weeks after initiation of therapy, or by examining the patient's physical or emotional health such as by subjecting the patient to a physical examination or to questioning about sleep patterns, side effects, daytime napping, general well-being.

[0040] Short of terminating treatment, it may be decided, e.g., that the patient should receive a different dose of the melatonin agonist or a different melatonin agonist, e.g., a different melatonin agonist having the pharmacological activity, i.e., MT1R and MT2R binding and relative binding affinities, and $t_{1/2}$ of tasimelteon.

[0041] The threshold probability discussed above can be correlated to a threshold concentration of melatonin in a biological sample taken from a patient. For example, melatonin levels can be directly measured in samples of blood, plasma, urine, saliva, and the melatonin concentration that corresponds to a selected threshold probability can be ascertained. The concentration of melatonin that corresponds to the selected threshold probability can be referred to as the Threshold Concentration.

[0042] Melatonin levels are generally determined (1) by measuring the amount of the primary urinary metabolite of melatonin, 6-sulphatoxymelatonin (aMT6s) collected every 2 to 8 hours over a 24 to 48 hour period, (2) by measuring melatonin levels in samples of saliva taken every 30 to 60 minutes under dim light, or (3) by measuring melatonin levels in samples of blood taken frequently, e.g., every 20 to 30 minutes. Such methods are summarized, e.g., by Benloucif et al., J Clin Sleep Med, 4(1): 66-69 (2008).

[0043] It is within the skill of the art, and therefore encompassed by this invention, to use any surrogate for melatonin concentrations or rates of production for determining the length of the melatonin rhythm, i.e., *tau*. For example, as specifically described herein, one may use amounts of aMT6s as a surrogate for amounts of melatonin and one may use the cortisol circadian rhythm or the aMT6s circadian rhythm as a melatonin circadian rhythm surrogate, i.e., the length of the circadian rhythm of cortisol can be a surrogate for the length of the circadian rhythm of aMT6s which can be a surrogate for the length of the melatonin circadian rhythm (i.e. *tau*). Alternatively or additionally, one may use cortisol as such melatonin surrogate.

[0044] In an illustrative embodiment, the amount of melatonin is indirectly measured such as by measuring the amounts of a melatonin surrogate, specifically, aMT6s in urine samples, and using such amounts to estimate acrophase and average and peak endogenous aMT6s amounts or concentrations in blood.

[0045] In an illustrative embodiment, the melatonin surrogate is the rate of aMT6s production as ascertained by measuring aMT6s in urine samples. In such case, the Threshold Concentration would actually be a rate of excretion expressed, e.g., in units of ng/hr. Such rate can be determined by measuring the concentration of aMT6s in an aliquot of urine (ng/ml) and multiplying it by volume/time (ml/hr) of the total urinary void from which the aliquot was derived, as more fully explained below. This surrogate measure is used in this illustrative embodiment for convenience only and it can readily be re-calculated as the concentration of aMT6s in urine and expressed, e.g., in ng/ml units or as the absolute amount of aMT6s in urine and expressed, e.g., in ng or mg units. Such amounts, whether expressed as excretion rates, concentrations, or weights, can also be converted into similarly expressed amounts of melatonin.

[0046] For example, a patient having a peak aMT6s production rate, i.e., excretion rate, of 1500 ng/hr in urine is a likely responder to tasimelteon. Therefore, the Threshold Concentration can be set at 1500 ng/hr aMT6s. Alternatively, the Threshold Concentration can also be set at 2000 ng/hr of urinary aMT6s (e.g., urine samples collected in 4 hour intervals and during a nighttime sleep period) or any convenient number therebetween, e.g., 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900, or 1950 ng/hr. Alternatively, the Threshold Concentration can also be set at greater than 2000 ng/hr of urinary aMT6s, e.g., 2100, 2200, 2300, 2400 or 2500 ng/hr.

[0047] A Threshold Concentration of 1500 ng/hr aMT6s is indicative of a greater than 50% probability that a given patient will respond to treatment, i.e., greater than 50% of a population of patients having a peak aMT6s concentration in urine (or the melatonin concentration that is equivalent thereto in another biological sample) are expected to respond to treatment. Based on the study results reported above, it is expected that more than about 75%, or even more than about 80% or 90% of patients will respond if they have peak aMT6s production rates in urine (or corresponding melatonin concentrations in a biological sample) of 1500 ng/hr or 2000 ng/hr.

[0048] If endogenous melatonin levels are used to predict likelihood of patient response and not for *tau* determination, then it is not necessary to determine the rate of aMT6s excretion at time points, or spans of timepoints, throughout a full day. Instead, e.g., the amount of melatonin, as inferred from aMT6s in urine, can be measured in urine collected and pooled in a single batch over a 24 hour period or even during a shorter period. Indeed, in illustrative embodiments, melatonin levels as indicated by aMT6s in urine or directly as melatonin in, e.g., blood or saliva, can be measured at given time points once or multiple times per day.

[0049] The ability to predict likelihood of response to drug is very important to healthcare providers, e.g., physicians and patients, as well as to healthcare reimbursement providers, e.g., providers of prescription drug insurance. Thus, in

one embodiment, prior to initiation of treatment of Non-24 with a melatonin agonist, e.g., tasimelteon, the patient is tested to determine his or her endogenous melatonin levels, in particular, his or her peak melatonin concentration. Such testing can be carried out using a biological sample, e.g., urine, blood, plasma, or saliva using the methodologies described above or any other methodology. Because the method of this invention provides a probability of response, the method of determining peak melatonin concentration does not require precision. It is enough that it provide an estimate within, e.g., 20%, in which case, if the Threshold Concentration is set at 2000 ng/hr urinary aMT6s, a patient would be regarded as a likely responder if the patient's peak aMT6s excretion in urine is determined to be 1600 ng/hr or higher. Even less precision, e.g., within 25% or 30%, may be acceptable. As in the case of determining *tau*, other surrogates for endogenous melatonin levels can also be used.

[0050] A further aspect of this invention arises from the fact that certain therapeutic agents are known to reduce endogenous levels of melatonin. Prominent among such agents are beta-adrenergic receptor antagonists, commonly referred to as "beta blockers", which are commonly prescribed for treatment of cardiac arrhythmias, myocardial infarction, congestive heart failure, and hypertension. Beta blockers include, e.g., alprenolol, atenolol, carvedilol, metoprolol, and propranolol, to name a few.

[0051] Thus, in one aspect, this invention comprises classifying Non-24 patients who are receiving beta blocker therapy as poor responders to melatonin agonist therapy. In this illustrative embodiment, such patients may not be subjected to a determination of peak melatonin concentration but, instead, may be treated as if their melatonin concentrations are below a Threshold Concentration. Other factors that may have an adverse effect on efficacy are NSAIDs and light.

[0052] In a related illustrative embodiment, a Non-24 patient may be directed to submit to a determination of melatonin concentration because he or she is being treated with beta blocker therapy to ascertain whether or not the beta blocker therapy is in fact causing the patient's peak melatonin level to drop below a Threshold Concentration.

[0053] In related aspects of this invention, plasma melatonin levels or beta blocker therapy, or both, are used as efficacy predictors in combination with other markers of efficacy or adverse events. So, for example, an illustrative embodiment of this invention comprises treating a patient suffering from Non-24 with tasimelteon if the patient has peak melatonin levels corresponding to 1500 ng/hr (or 2000 ng/hr) of aMT6s in urine collected during 4 hour periods or a nighttime sleep period and if the patient is positive for one or more additional efficacy markers. Incorporation of such additional efficacy marker or markers can enhance the ability of a healthcare provider to assess the likelihood that a patient suffering a non-24 hour circadian rhythm will benefit from treatment with a melatonin agonist such as tasimelteon.

[0054] In related embodiments, a computer-based system receives information about a prescription for tasimelteon and operates to associate that information with information about the patient's endogenous melatonin levels to output a report indicating a probability of efficacy or to output a report stating that a higher or lower dose of tasimelteon, e.g., < 20 mg/d or >20 mg/d, is indicated.

[0055] Patients can be diagnosed as suffering from Non-24 by estimating each patient's circadian period (*tau*). Patients whose *tau* exceeds 24 hours are diagnosed as having Non-24. Thus, in general, Non-24 patients who can benefit from treatment with tasimelteon have a *tau*, such as may be determined by analyzing the aMT6s or cortisol circadian rhythm, that is longer than 24 hours, e.g., greater than about 0.1 hours longer than 24 hours and in some cases, at least about 0.2, 0.3, 0.4 and as large as about 1.4 hours longer than 24 hours. As discussed herein, the cortisol circadian rhythm can be used in place of or in addition to the aMT6s rhythm, although cortisol circadian rhythm calculations may be slightly less precise in the sense that such data compiled from analyses of a population of patients may exhibit a larger standard deviation.

[0056] To monitor circulating melatonin cycles in a subject, it is convenient to assay for levels of the major metabolite of melatonin, which is 6-sulfatoxymelatonin (aMT6s) in urine, as its pattern of production correlates closely with circulating melatonin levels. However, this invention contemplates measurement of aMT6s levels in other bodily samples such as blood, e.g., plasma, or saliva and it also contemplates direct measurement of melatonin or of other surrogates for melatonin levels. It is within the skill of the art to correlate levels of tasimelteon or tasimelteon metabolites in other bodily samples (i.e., other than aMT6s in urine) with circulating melatonin levels. For example, the amounts of cortisol in blood or urine can be used in a manner similar to the use of aMT6s to determine *tau*.

[0057] A useful protocol for estimating *tau* in candidates for clinical testing for treatment of Non-24, which method can be applied to diagnosis of Non-24 in a given patient, is as follows:

Each subject will undergo four 48-hour urine collection sessions at nominal days 7, 14, 21, and 28. During each session, the start of the session and the time of each void will be recorded. Urine collected over periods of 4 hours (with the first 4 hour collection period of the day beginning at scheduled wake time), or about 8 hours during sleep, will be pooled (the "collection interval"); thus, subjects will have a total of 10 urine collection intervals during each 48-hour period. A study nurse will determine the volume of urine collected during each interval (urine will be transferred to a graduated cylinder) and an aliquot will be assayed for aMT6s.

For each collection interval, the start and end time of the interval will be used to determine the midpoint and duration of the interval. The start time of a given interval is defined as the last void time from the prior 4 hour (or 8 hour)

EP 2 806 863 B1

collection interval; the end time of a given interval is defined as the last void time within the collection interval.

The mass of the primary melatonin metabolite (aMT6s) excreted during the interval will be determined as the product of aMT6s concentration and volume of urine. Rate of aMT6s excretion will be determined as the mass of aMT6s excreted divided by the duration of the interval. This rate will be associated with the midpoint of the interval, referenced to the midnight preceding the start of the first interval in that session.

For example, if a collection interval on Day 27 runs from 9AM to 1PM (and the patient had a void at exactly 9AM and a void at exactly 1PM), midpoint of that interval would be assigned the value 11.0. A comparable interval on the next day of that session would be assigned a value 35.0.

To accommodate changes in the clock time due to Daylight Savings Time changes, no urine collections will occur on a day that the clock changes. For screening there will be occasions when the 4 different weeks that urine collections are conducted will span a change in the clock time. Therefore, all urine collection times will be automatically translated into local standard time for calculations and then translated back to DST for reporting purposes, if appropriate.

In certain situations, urine collections or their recording will be incomplete. The following procedures will be invoked to address this:

1. If a subject fails to timestamp a void, no action will be taken if there are multiple voids with timestamps within one interval.
2. If there is only one void in a collection interval, and the patient cannot recall the time of the void then the entire 48 hour collection period will be excluded from the analysis and the subject will be requested to collect an additional 48 hours of urine after Day 28. It would not be possible to accurately determine to which collection interval the unmarked urine belongs. Consequently, the appropriate assignment of start and stop times to all of the collection intervals would be questionable.
3. If a void is discarded by the patient but the time of the void is known, duration associated with that void (time of the void minus the time of the previous void) will be subtracted from the total duration associated with that interval. This modified duration will be used to calculate rate of aMT6s excretion. If a discarded sample is either the first or last of the samples in an interval, the midpoint of that interval will be calculated without considering that sample.
4. If fewer than 4 samples are available for one 48-hour collection session, fitting of the cosine will be compromised (inadequate degrees of freedom). Consequently, acrophase will not be determined if fewer than four samples are available.

For each session, acrophase will be determined by fitting a cosine to the data from that session using unweighted non-linear regression. Fitting will be performed using a non-linear least squares fitting algorithm. The fitting process will estimate phase shift, mesor, and amplitude and their respective standard errors; period of the cosine will be fixed to 24 hours.*

Acrophase will be determined as the phase shift modulus 24 hours.

If acrophase values are available for three or more sessions, τ will be calculated using the following procedure:

1. Acrophase will be recalculated relative to day 0 ($24 \cdot \text{start day for each session} + \text{acrophase}$).
2. These values will be regressed against start day for each session using weighted linear regression. Weighting will be by the inverse square of the standard error associated with the estimate for acrophase for each session.

* Although these subjects are presumed to have a $\tau > 24$ hours, attempts to estimate τ led to consistently poor results with multiple test datasets. Steven Lockley, Ph.D., an expert in the field uses this approach.

[0058] Thus, related to this invention is a method for determining a patient's circadian rhythm (τ) and for treating a patient with a melatonin agonist, in particular, tasimelteon, based on that patient's τ . In illustrative embodiments, the method of determining τ and treating a patient based on the patient's τ , in particular, based upon time of aMT6s acrophase, comprises steps (a) through (f), as follows:

- a) collecting at least one biological sample from the patient during each of a plurality of regular collection intervals (CIs) during at least two Collection Sessions, each Collection Session being at least 48 hours in duration;
- b) if multiple biological samples (i.e., samples of the same type) are collected during each CI, then optionally physically pooling all samples collected within a given CI and, in such case, assigning a Collection Time Point for each CI;
- c) measuring the amount (absolute or concentration) of melatonin or of a melatonin surrogate in each of the samples or pooled samples;
- d) optionally converting the amount of melatonin or melatonin surrogate at each Collection Time Point to a rate of production;

EP 2 806 863 B1

e) subjecting the amount of melatonin or melatonin surrogate or the rate of melatonin or melatonin surrogate production at each Collection Time Point to cosinor analysis to model the patient's cycle, including the acrophase, of melatonin or melatonin surrogate amount or production on each day;

f) fitting serial acrophase determinations to a weighted linear regression model in order to determine τ (τ), wherein $\tau = 24 + \text{slope}$.

[0059] While cosinor analysis is mentioned above, it will be appreciated that other methods can be used, e.g., a 2-harmonic fit analysis, in particular, for cortisol rhythm analysis.

[0060] Following such determination of τ , a patient can be treated with a melatonin agonist, e.g., tasimelteon, such as described in step (g), as follows: g) if the patient's τ is longer than 24 hours, then:

(i) projecting the patient's acrophase for each of at least 30 days following Day 2 of the final Collection Session by adding τ to the acrophase of said final Day 2 and to each day thereafter and

(ii) treating the patient by daily internally administering to the patient an effective amount of the melatonin agonist prior to sleep time, beginning on the night of the Optimal Treatment Initiation Day, or on a night within the Optimal Treatment Initiation Window, during a succeeding circadian cycle.

[0061] The Optimal Treatment Initiation Day is the day on which the patient's sleep time is expected to be closest to what it would be if the patient had a normal, i.e., 24 hour, i.e., < 24.1 hr, τ . Such day is generally the day of the night on which the patient's melatonin (or melatonin surrogate) acrophase is projected to be the optimal acrophase, i.e., the time at which acrophase would occur if the patient had a normal circadian rhythm. It is not necessary to initiate treatment precisely on the Optimal Treatment Initiation Day but it is recommended that treatment be initiated on such day or within a range of days on either side of such day, said range being referred to herein as the Optimal Treatment Initiation Window. Said window generally comprises the Optimal Treatment Initiation Day and (a) the immediately following days on which the melatonin (or surrogate) acrophase is projected to occur no later than about 3.5 hours (e.g., 3 hours, 3.5 hours or 4 hours) later than the optimal melatonin (or surrogate) acrophase and (b) the immediately preceding days on which melatonin (or surrogate) acrophase is projected to occur no earlier than 5 hours earlier than the optimal melatonin (or surrogate) acrophase.

[0062] For the sake of convenience, the Optimal Treatment Initiation Window can be conveniently defined as a set number of days before and after the projected Optimal Treatment Initiation Day, e.g., 2 days before and 2 days after, for a defined Optimal Treatment Initiation Window comprising a total of 5 days. Such window is illustrated in Figure 2 wherein the first Optimal Treatment Initiation Day is December 4, 2010 and the Optimal Treatment Initiation Window is defined for convenience as December 2, 2010 to December 6, 2010.

[0063] It will be appreciated, however, that the window can be customized as summarized above based on a given patient's τ , i.e., depending upon how fast a patient's circadian rhythm is running, such that a patient with a relatively fast-moving circadian rhythm will have a narrower optimal window than a patient with a relatively slow-moving circadian rhythm.

[0064] Normal monitoring can comprise step (h), as follows:

h) following a treatment period of at least one complete circadian cycle (based on the patient's pre-treatment τ) assessing entrainment as follows:

(i) If τ is < 24.1 hours with a 95% Confidence Interval that crosses 24.0 hours, then the patient is considered to be entrained to a 24 hour day;

(ii) If the last two acrophase estimates are within the target range, i.e., -2 to +6 hours from optimal acrophase, and the Standard Deviations of these two acrophases overlap, then, taking an additional biological sample collection and re-calculating τ based on the last three acrophase estimates (the original two + the additional) and if τ is < 24.1 hours with a 95% Confidence Interval that crosses 24.0 hours, the patient is considered to be entrained to a 24 hour day;

(iii) If $\tau \geq 24.1$ hours or the 95% Confidence Interval τ does not cross 24.0 hours, then the patient is retested.

[0065] The duration of a complete circadian cycle will vary depending upon the rate at which a given patient is free running. For example, with reference to Figure 2, a patient having a τ of 24.6 hours will complete a circadian cycle in approximately 39 days (e.g., December 4, 2010 to January 13, 2011). A patient with a slower rhythm, e.g., $\tau = 24.5$, will have a longer cycle and, conversely, a patient with a faster rhythm, e.g., $\tau = 24.7$, will have a shorter cycle.

[0066] The τ determination and treatment method generally described above can comprise any one or any combination of any two or more of the following limitations:

EP 2 806 863 B1

1. melatonin amounts are indirectly measured by measuring the amounts of a melatonin surrogate, said surrogate being aMT6s.
2. the biological sample is urine, all urine collected during a given CI is physically pooled, and the mid-point of the CI is assigned as the Collection Time Point for that CI.
- 5 3. each CI during wake time is 4 hours and sleep time is a single CI, provided that samples are not collected during the first four hour period of each Collection Session or, if collected, are not used in the determination of tau.
4. the Collection Time Point for each CI is defined as the mid-point between the time of the last urine void in the CI immediately preceding a given CI and the last urine void in the given CI.
- 10 5. there are 4 Collection Sessions.
6. there are 48 hours in each Collection Session.
7. Collection Sessions are conducted once per week.
8. the Optimal Treatment Initiation Day is the day of the night on which the melatonin or melatonin surrogate acrophase is projected to be the optimal acrophase.
- 15 9. the optimal acrophase is the time at which aMT6s acrophase is projected to be closest to and no later than about 3.5 hours prior to the patient's target wake time.
10. the Optimal Treatment Initiation Window comprises the Optimal Treatment Initiation Day and (a) the immediately following days on which the melatonin acrophase is projected to occur no later than 3 hours later than the optimal acrophase and (b) the immediately preceding days on which melatonin acrophase is projected to occur no earlier than 5 hours earlier than the optimal acrophase. In such embodiments, cortisol can be used in place of aMT6s with adjustment to account for the difference between the cortisol circadian rhythm and the aMT6s circadian rhythm.
- 20 11. treatment comprises internal administration of an effective amount of tasimelteon once per day, the time of administration being about 5 hours prior to the time of the optimal aMT6s acrophase, and wherein treatment is continued daily for at least one complete circadian cycle. In such embodiments, cortisol can be used in place of aMT6s with adjustment to account for the difference between the cortisol circadian rhythm and the aMT6s circadian rhythm.
- 25 12. the amounts of melatonin or melatonin surrogate are measured in absolute units or in concentration units.
13. the amount of melatonin or melatonin surrogate in the biological sample is determined as the product of the aMT6s concentration (mass/volume) and the volume of the biological sample.
14. the rate of melatonin or melatonin surrogate production is determined as the mass of melatonin or melatonin surrogate produced and collected during each CI divided by the duration of the CI.
- 30 15. the rate of production is expressed as g/hr.
16. no samples are collected on a day that the clock changes to or from Daylight Savings Time (DST) and, if the Collection Sessions span a change in the clock time, all Collection Time Points are translated into local standard time for calculations and then translated back to DST or standard time, as appropriate, for reporting purposes.
- 35 17. samples are collected in a sample collection container by the patient and provided to a laboratory for analysis, e.g., a diagnostic laboratory.
18. the patient records the date and time of each sample collection on a label that has been previously fixed to the collection container or that is applied to the collection container by the patient.
19. the date and time of each collection are printed onto the label by timestamp clock.
- 40 20. the biological sample is urine and melatonin amounts are indirectly measured by measuring the amounts of aMT6s and wherein if urine collections or their recordings are incomplete, then:
 - (i) if a patient fails to timestamp a void, no action is taken if there are multiple voids with timestamps within one CI;
 - 45 (ii) if there is only one void in a CI and the patient cannot recall the time of the void, then the entire 48 hour Collection Session is excluded from the analysis and an additional Collection Session is conducted;
 - (iii) if a void is discarded by the patient but the time of the void is known, the duration associated with that void (time of the void minus the time of the previous void) is subtracted from the total duration associated with that CI and the modified duration is used to calculate the rate of aMT6s production but if a discarded sample is either the first or last of the samples in a given CI, then the midpoint of that CI will be calculated without considering that sample; provided that, if fewer than 4 samples are available for any one Collection Session, acrophase will not be determined for that Collection Session.
- 50 21. in step (h), if $\tau \geq 24.1$ hours or the 95% Confidence Interval does not cross 24.0 hours, then treatment is continued and the patient is retested after a second complete circadian cycle.
- 55 22. in step (g), if the patient's τ is longer than 24 hours, e.g., $\tau \geq 24.1$ hours, the patient's acrophase is projected for each of the 90 days following Day 2 of the final Collection Session.
23. aMT6s or cortisol is extracted from pooled urine samples by solid phase extraction, the extracts are evaporated to dryness, the residue is then reconstituted with solvent, and the solution is analyzed by HPLC-MS, an antibody

binding assay, or other analytical technique.

[0067] Thus, a particular illustrative embodiment of a method of determining *tau* and thereafter treating a patient thereby determined to have a free-running circadian rhythm is as follows:

5 a) collecting and, if more than one, physically pooling urine samples from the patient during each of 9 Collection Intervals (CIs) during four weekly 48 hour collection sessions, said 9 CIs being CI2, CI3, CI4, CI5, CI6, CI7, CI8, CI9, and CI10, as follows:

10 CI1: 4 hour period beginning approximately on initiation of wake time of Day 1 of the first Collection Session;
 CI2: 4 hour period beginning at the end of CI1;
 CI3: 4 hour period beginning at the end of CI2;
 CI4: 4 hour period beginning at the end of CI3;
 CI5: Overnight, i.e., sleep time (approx 8 hours),
 15 CI6: 4 hour period beginning approximately on initiation of wake time of Day 2 of the collection session;
 CI7: 4 hour period beginning at the end of CI6;
 CI8: 4 hour period beginning at the end of CI7;
 CI9: 4 hour period beginning at the end of CI8;
 CI10: Overnight, i.e., sleep time (approx 8 hours),

20 b) (i) optionally collecting and discarding samples during CI1 and (ii) assigning the mid-point between the last void of each CI immediately preceding a given subsequent CI and the last void of the given subsequent CI as the Collection Time Point for each of CI2, CI3, CI4, CI5, CI6, CI7, CI8, CI9, and CI10;

c) measuring the amount of aMT6s or cortisol in each of the ten samples;

25 d) converting the measured amount of aMT6s or cortisol at each Collection Time Point to a rate of production;

e) subjecting the rate of aMT6s or cortisol production rate at each Collection Time Point to cosinor analysis to model the cycles, including the acrophase, of aMT6s or cortisol production on each day;

f) fitting serial acrophase determinations to a weighted linear regression model in order to determine circadian period (τ), wherein $\tau = 24 + \text{slope}$ ($p \leq 0.05$);

30 g) if the patient's τ is longer than 24 hours, then:

(i) projecting the patient's acrophase for each of the 90 days following Day 2 of the final Collection Session by adding τ to the acrophase of said final Day 2 and to each day thereafter and

35 (ii) treating the patient by daily internally administering to the patient an effective amount of tasimelteon prior to sleep time, beginning on the night of the Optimal Treatment Initiation Day, or on a different night within the Optimal Treatment Initiation Window, during the next succeeding circadian cycle

h) following a treatment period of one complete circadian cycle, assessing entrainment as follows:

40 (i) if τ is < 24.1 hours with a 95% Confidence Interval that crosses 24.0 hours, then the patient is considered to be entrained to a 24 hour day;

(ii) if the last two acrophase estimates are within the target range, i.e., -2 to +6 hours from optimal acrophase, and the Standard Deviations of these two acrophases overlap, then, taking an additional 48-hour urine collection and recalculating τ based on the last three acrophase estimates (the original two + the additional) and if *tau* is < 24.1 hours with a 95% Confidence Interval that crosses 24.0 hours, the patient is considered to be entrained to a 24 hour day;

45 (iii) if $\tau \geq 24.1$ hours or the 95% Confidence Interval does not cross 24.0 hours, then the patient is retested with an additional four 48-hour urine collection scheduled beginning 1 circadian cycle from the first collection.

50 **[0068]** It will be apparent that in the urine collection and analysis methods that may be used in the practice of aspects of this invention, it is not essential to use the entire volume of urine collected during each Collection Interval.

[0069] The method of treatment of Non-24 by internally administering an effective amount of a melatonin agonist, in particular, tasimelteon, is not dependent upon the method for diagnosing or monitoring patients. Instead, said method of treatment is useful in treating Non-24 patients regardless of how diagnosed. Similarly, other markers may be used to predict urinary aMT6s or cortisol acrophase.

55 **[0070]** Non-entrained persons, i.e., persons with a non-24 hour circadian rhythm, may exhibit symptoms of Non-24 with a clearly non-24 hour sleep period such that initiation of sleep and waking times, unless artificially interrupted, begin later each succeeding day. Other patients may exhibit less severe shifts in sleep period and a significant number may

exhibit no shift in sleep period. Such patients, particularly those who do not exhibit shift in sleep period, can be misdiagnosed as having a normal *tau* if the diagnosis is based solely on sleep and wake times. Some patients that exhibit mild or no shift in sleep period may have cyclic patterns of one or more of sleep latency, nighttime sleep duration and daytime naps. Regardless of the sleep problem, patients with non-24 hour circadian rhythms may be at risk for other circadian-related disorders, for example, metabolic disorders.

[0071] Entrainment of patients diagnosed as suffering from a non-24 hour circadian rhythm, including Non-24, can be effected by initiating internal administration of a melatonin agonist like tasimelteon or an active metabolite of tasimelteon or a pharmaceutically acceptable salt thereof, at any time or treatment can be initiated on or about a day on which the patient's melatonin acrophase (based, e.g., on urinary aMT6s acrophase) is predicted to occur about 3 to 4 hours, or about 3.5 hours, e.g., 3.25 hrs to 3.75 hrs, prior to a target wake time selected for or by a given patient. The "ideal" day for initiation of treatment can be more explicitly defined as the day when the subject's predicted acrophase is both 1) closest to 3.5 hours prior to target wake time and 2) earlier than that time. The latter qualifier makes it more likely than not that treatment initiation will occur in a phase-advance part of the phase response curve.

[0072] For example, treatment of a patient who has a target bedtime of 10:00 p.m. and a target wake time of 7:00 a.m., treatment initiation can be on a day when urinary aMT6s acrophase is predicted to occur at 3:30 am. However, treatment with tasimelteon can conveniently be initiated on a day on which melatonin acrophase, e.g., using calculated urinary aMT6s acrophase, is predicted to be between about 5.5 hours before target wake time and 2.5 hours after target wake time. Without intending to be bound to a particular theory, this flexibility is apparently owing to the unusually marked effects of such active ingredient on circadian rhythm upon initiation of treatment (e.g., phase advance by as much as about 5 hours on initial treatment).

[0073] If a marker for circulating melatonin levels other than urinary aMT6s is employed, e.g., aMT6s in plasma, then the above times would be adjusted accordingly but would nevertheless be indirectly indicative of urinary aMT6s levels.

[0074] In patients suffering Non-24, a calendar day may not be associated with an acrophase. For example, if a subject's *tau* is 24.5 hours and acrophase occurs at 23:45 (11:45 pm) on 28 August, the next acrophase is predicted to occur at 00:15 (12:15 am) on 30 August.

[0075] In addition to entraining a Non-24 patient's *tau* to 24 hours, e.g., <24.1 hours, a melatonin agonist, in particular, tasimelteon, can also increase total sleep time per day and reduce total nap time per day.

[0076] Entrainment of a patient can be determined by various methods, including by determining the patient's *tau* by the above-described or different methodologies. In addition, or alternatively, a patient's or a healthcare worker's perception of improvement can be assessed such as by use of a questionnaire. Such perception could utilize, e.g., the Clinical Global Impression of Change (CGI-C),

[0077] The CGI-C is a healthcare worker-rated assessment of change in global clinical status, defined as a sense of well-being and ability to function in daily activities. See, e.g., Lehmann E., *Pharmacopsychiatry* 1984,17:71-75. It is a 7 point rating scale whereby clinicians, physicians, or other healthcare workers rate a patient's improvement in symptoms relative to the start of the study. It is rated as: 1, very much improved; 2, much improved; 3, minimally improved; 4, no change; 5, minimally worse; 6, much worse; or 7, very much worse.

[0078] The questionnaire can be administered prior to or early following initiation of treatment, e.g., prior to Day 1 or, e.g., on Day 56 (counted from first day of treatment) and it can be re-administered later following initiation of treatment, e.g., Day 112 and/or Day 183.

[0079] Due to the cyclicity of Non-24, a patient's overall improvement should not be assessed at one time-point/visit. Consequently, the average score of CGI-C in the last two scheduled assessments (e.g., Day 112 and Day 183) can be used to evaluate the patient's overall improvement.

[0080] In addition to or as an alternative to measuring a patient's *tau* following a period of treatment and/or utilizing patient or healthcare worker assessment such as by use of the CGI-C, various sleep parameters can also be used to assess efficacy of treatment, i.e., entrainment.

[0081] For example, sleep parameters that can be assessed include one or more of Lower Quartile of Nights of nTST (LQ-nTST), Upper Quartile of Days of dTSD (UQ-dTSD), and Midpoint of Sleep Timing (MoST).

Lower Quartile of Nights of nTST (LQ-nTST)

[0082] Patients suffering from Non-24 may have trouble sleeping as a result of their sleep cycle being out of synchrony with the 24 hour clock. This leads to intervals of poor sleep followed by intervals of good sleep. Therefore, the severity of symptoms associated with Non-24 is best illustrated when isolating the worst nights of sleep and the days with the most naps. Evaluating the 25% worst nights of sleep of an individual serves as a good measure of how an individual is suffering from this circadian disease in relationship to nighttime total sleep time (nTST).

[0083] The method for calculating the LQ-nTST is described as follows. For a given individual, all non-missing values (must include > 70% of one circadian cycle for both baseline and randomized data) of nighttime total sleep time are ordered from smallest to largest. The first 25% ($\text{ceiling}(\text{number of non-missing records}) / 4$) of the records are flagged

EP 2 806 863 B1

as belonging to the lower quartile of nighttime total sleep time. The average of these values is calculated and this result is denoted LQ-nTST.

[0084] For example, assume that a subject has 21 nTST baseline records: 6.75, 6.75, 1, 1, 6.75, 1.083, 7.167, 0.833, 7.083, 7.983, 7, 7, 7.833, 7, 7.667, 7.183, 7, 7.067, 7, 7.183, and 7.

[0085] These are rank ordered and the first 25% of records are selected $[(21/4) = 6]$: 0.833, 1, 1, 1.083, 6.75, and 6.75.

[0086] Those values are averaged to obtain the subject's LQ-nTST: $(0.833 + 1 + 1 + 1.083 + 6.75 + 6.75) / 6 = 2.91$.

Upper Quartile of Days of dTSD (UQ-dTSD)

[0087] Patients suffering from Non-24 have a propensity to sleep during the day as a result of their sleep cycle being out of synchrony with a 24 hour clock including daytime napping. In contrast, they may have very little or no napping when their circadian rhythms are aligned with the 24-hour day. In order to measure the effect of this dynamic circadian disorder on daytime napping a robust assessment for measuring the worst of the daytime napping, the 25% worst days will be used for this calculation in a similar fashion as for LQ-nTST.

[0088] The method for calculating the UQ-dTSD is described as follows. For a given individual, all non-missing values of daytime total nap durations are summed for a given day and then these daily summations are rank ordered from largest to smallest (Note: days for which an individual reported no nap are recorded as zero). The first 25% (ceiling(number of non-missing records)/4) of the records are flagged as belonging to the upper quartile of daytime total sleep duration (dTSD). The average of these values is calculated and this result is denoted UQ-dTSD.

[0089] For example, assume that a subject has 26 dTSD baseline records: 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 1.083, 0, 1.083, 1.667, 1.083, 1.083, 1.083, 1.083, 1.083, and 1.083.

[0090] These are rank ordered (largest to smallest) and the first 25% of records, i.e., ceiling(26/4) = 7 records identified: 1.667, 1.083, 1.083, 1.083, 1.083, 1.083, and 1.083.

[0091] These values are averaged to obtain the subject's UQ-dTSD: $(1.667 + 1.083 + 1.083 + 1.083 + 1.083 + 1.083 + 1.083) / 7 = 1.17$.

Midpoint of Sleep Timing (MoST)

[0092] Circadian rhythm disorders, including Non-24, are characterized by a timing misalignment of the circadian rhythms to the 24-hour light-dark cycle and hence the activities that an individual is performing (e.g., attempting to sleep at night when the circadian rhythms are signaling the brain to be awake). Midpoint of sleep timing is derived from a combination of the sleep reported in both the pre- and post-sleep questionnaires. The midpoint of sleep timing over a 24 hour period (adjusted to be relative from -12 hours before bedtime until +12 hours after bedtime) can be calculated for each day. The first step in calculating the midpoint is to calculate the midpoint and weight, e.g., duration, for each sleep episode. The total 24-hour sleep time is the summation of all sleep episodes in this 24 hour period. Each of the individual sleep episodes is then assigned a weight relative to the fraction of 24 hour sleep that it contains.

[0093] A useful MoST algorithm can be summarized as follows:

1. calculate the midpoint and weight, i.e., duration, for each sleep episode in a given 24 hour period;
2. assign a weight to each sleep episode;
3. determine the average of the weighted sleep episodes; and
4. correct the average of the weighted sleep episodes for target bedtime.

[0094] More specifically, such useful algorithm may be further defined as follows:

the midpoint for each sleep episode in a 24 hour period is calculated as follows:

$$\text{Sleep Start Time} + [(\text{Sleep End Time} - \text{Sleep Start Time})/2] - 24;$$

the weight of each sleep episode is equal to the duration of sleep (as perceived or objectively measured);
the weighted value of each sleep episode is calculated as follows:

$$\text{midpoint} * (\text{weight} / \text{TST})$$

EP 2 806 863 B1

where TST is the sum of all sleep durations in the 24 hour period;
the average of the weighted sleep episodes is the sum of the weighted values of all sleep episodes divided by the
number of sleep episodes; and
the correction for target bedtime is calculated as follows:

5

24 - target bedtime + average of weighted sleep episodes.

[0095] For example, assuming an individual with a target bedtime of 10:30PM went to sleep at 10:30PM and woke up
at 6:30AM (with a self-reported total sleep time of 5 hours). Assuming, also, that he/she took a nap at 8:05PM that lasted
2 hours and 5 minutes. The mid-point of sleep timing (MoST) for that day would be 1.959559 (relative to the target
bedtime), calculated as follows.

10

Nighttime Sleep Midpoint:

15

[0096]

Sleep Start Time = Target Bedtime = targetBT = 10:30PM = 22.5

Sleep End Time = Wake Time = 6:30AM = 6.5

20

Sleep End Time (adjusted for 24 hour periodicity) = 24 + 6.5 = 30.5

Nighttime Sleep Midpoint = $[(30.5 - 22.5) / 2]$ modulus 24 = 2.5 (relative

25

to the midnight)

weight = nTST = 5 hours = 5.0

Nap Midpoint:

30

[0097]

Sleep Start Time = NapStart = 08:05PM = 20.08333

NapDuration = 02h05m = 2.083333

35

Sleep End Time = NapEnd = NapStart + NapDuration = 20.08333 +

2.083333 = 22.16667 (10:10PM)

40

Nap Midpoint = NapStart + (NapEnd - NapStart)/2 = 20.08333 +

$[(22.16667 - 20.08333) / 2] - 24 = -2.875$ (relative to the midnight)

45

weight = NapDuration = 2.083333

Weighting of Sleep Episodes

50

[0098]

TST = sum(all sleep episodes) = sum(5.0, 2.083333) = 7.083333

Weighted Nighttime Sleep = mid*(weight/TST) = 2.5 *(5/7.083333) =

55

1.7647059

$$\text{Weighted Nap Sleep} = \text{mid} * (\text{weight} / \text{TST}) = -2.875 * (2.083333 / 7.083333)$$

$$= -0.8455882$$

Average of Weighted Sleep Episodes

[0099]

$$\text{Mean of } (1.7647059, -0.8455882) = 0.4595588$$

Correction for Target Bedtime

[0100]

$$\text{Correction Amount} = 24 - \text{targetBT} = 24 - 22.5 = 1.5$$

$$\text{MoST} = 0.4595588 + 1.5 = 1.959559 \text{ (relative to the target bedtime).}$$

[0101] Under ideal circumstances in which an individual sleeps at their desired time for 7-8 hours and does not have any daytime naps the MoST will be around 3.5-4.0. In the above hypothetical example, this individual had a late afternoon or night nap which pulls the midpoint below this desired range to 1.96. Alternatively, if a patient has more morning naps then this would potentially lead to a bigger number. If the illustration were changed such that the hypothetical patient slept from 10:30 pm to 6:30 am with no naps, then the patient's MoST would be 4.0. This algorithm dynamically takes into account the information from both the nighttime sleep as well as the daytime napping. Additionally, because the weighted sleep episodes are divided by the total number of sleep episodes within a 24 hour period the derived midpoint of sleep timing will be pushed to 0 (and away from the optimal value of 3.5-4.0) as an individual's sleep becomes more fragmented. An improvement in MoST is defined as an increase in the MoST scale.

[0102] A useful clinical response scale (CRS or N24CRS) can be formed by combining the results of all of LQ-nTST, UQ-dTSD, MoST and CGI-C. In an illustrative embodiment, each assessment on the scale is scored as a 1 or 0 depending on whether the pre-specified threshold is achieved or not, as defined in the table that follows. The score for each assessment is summed with a range of 0-4. Individuals with a N24CRS score of > 3 are classified as having responded to treatment.

Non-24 Scale of Clinical Response

[0103]

Assessment	Threshold of response
LQ-nTST	>30, >40 or >45 minutes increase in average nighttime sleep duration
UQ-dTSD	>30, >40, or >45 minutes decrease in average daytime sleep duration
MoST	>20, >25 or >30 minutes increase
CGI-C	<1 or <2 from baseline

or any combination or permutation thereof. Increases and decreases in duration, and other scores in the N24CRS, may be determined by comparing baseline, which may be an average of two or more assessments, to post-treatment, which may be an average of two or more post-treatment assessments. For example, the CGI-C scoring of <=1 (or <=2) can be a comparison of baseline score, which may be a single data point or an average of two (or more) scores from assessments taken prior to or shortly after initiation of treatment, to single data point or to an average of two (or more) scores from post-treatment assessments.

[0104] In an illustrative embodiment, improvement, i.e., response to treatment, is defined as the coincident demon-

stration of:

1. shift of τ towards 24 hours and
2. a score of ≥ 3 on the above-described N24CRS.

[0105] In such embodiment, τ can be measured using any methodology including aMT6 in urine, cortisol, melatonin in blood or saliva, substantially as described above.

[0106] A score of ≥ 2 can also indicate improvement, i.e., patient response to treatment.

[0107] The data required to calculate parameters such as LQ-nTST, UQ-dTSD, and MoST, can be objectively quantified in sleep studies or, more practically, it can be collected by way of patient questionnaires that ask patients to self-assess, e.g., did the patient sleep, what time did he or she go to bed, how long did it take to fall asleep. In certain clinical studies, subjects will be required to call an Interactive Voice Response System (IVRS) twice a day starting the day after all screening assessments are completed and continue through the randomization phase for 2.5 circadian cycles or 6 months whichever is less. Subjects will call the IVRS twice, once in the morning no later than 1 hour after scheduled awakening to report nighttime sleep parameters (PSQ) and again in the evening no later than 15 minutes after the subjects daily dosing time to report the length and duration of any daytime sleep episode(s) (PreSQ). The IVRS will automatically call back any subject that fails to perform the required calls within the allocated timeframe. One of skill in the art can readily transfer this or similar methodologies to the treatment setting.

[0108] It will be appreciated, of course, that other methodologies may be used to ascertain improvement following initiation of treatment or that variations in the above-described methodologies can be employed, e.g., by utilizing other tau determination methods and/or by measuring different or additional sleep parameters.

[0109] Illustrative efficacy indicators based on the above include, e.g.:

1. Combined sleep/wake response (≥ 90 minute increase in LQ-nTST plus a 90 minute decrease in UQ-dTSD);
2. Entrainment of cortisol secretion;
3. Entrainment + 45 minute increase in LQ-nTST;
4. Entrainment + 45 minute decrease in UQ-dTSD;
5. Entrainment + ≥ 30 minutes increase in MoST;
6. Entrainment + a score of much improved or better on the CGI-C scale;
7. Increase in LQ-nTST;
8. Decrease in UQ-dTSD;
9. Improvement in MoST;
10. Improvement in CGI-C;
11. N24CRS = 4;
12. Combined sleep/wake response (≥ 45 minute increase in LQ-nTST plus a 45 minute decrease in UQ-dTSD).

[0110] In carrying out these methods of the invention, the average of multiple pre-treatment and post-treatment assessments can be used to smooth out test to test and/or day to day variability. For example, a baseline MoST can be compared to the average of two post-treatment initiation MoSTs; in this case, preferably, the difference between the two post-treatment MoSTs is less than 2 hours. If the difference is greater than about 2 hours, one or more further MoST assessments can be carried out.

[0111] If efficacy is shown, i.e., if a patient is determined to have achieved or to be moving in the direction of a normal circadian rhythm (i.e., 24 hours or up to 24.1 hours), then treatment can be continued. If efficacy is not shown, then a physician or other healthcare worker may wish to discontinue treatment or change the dose of the melatonin agonist, or otherwise alter the treatment method.

[0112] The above-described response assessment methodologies can also be utilized for diagnostic purposes. So, for example, a MoST of less than about 3.5, or less than about 3.0, or less than about 2.5 can be an indication that the patient is suffering from a free running circadian rhythm. Such diagnostic can employ one or more of the above-described parameters optionally with other diagnostic markers also being assessed. For example, the patient's MoST score in combination with a tau determination could also be or be part of a useful diagnostic for free running circadian rhythm.

[0113] Thus, in one method of treatment that comprises an aspect of this invention, a patient who presents himself or herself to a physician or other healthcare professional with symptoms of a sleep disorder, e.g., difficulty sleeping at night, frequent daytime naps, is first diagnosed by assessment of the patient's MoST, with or without other diagnostic assessments. Such patient who has a low, e.g., less than 3.5 MoST is then treated with a melatonin agonist, e.g., tasimelteon.

[0114] In Phase III clinical trials, i.e., safety and efficacy studies in humans, (SET Study), tasimelteon was demonstrated to be useful in entraining Non-24 patients to a 24 hour circadian rhythm. Specifically, patients were orally administered 20 mg tasimelteon per day for at least 12 weeks prior to re-estimating τ . Patients were selected for randomization or

EP 2 806 863 B1

open label based on baseline tau estimates. Drug was administered at about 1 hour prior to target sleep time, as determined by patients based on a 9 hour nighttime sleep period.

[0115] The SET study was an 84 patient randomized, double-masked, placebo-controlled study in patients with Non-24. The primary endpoints for this study were Entrainment of the melatonin (aMT6s) rhythm to the 24-hour clock and Clinical Response as measured by Entrainment plus a score of greater than or equal to 3 on the following N24CRS:

Non-24 Scale of Clinical Response:

[0116]

Assessment	Threshold of response
LQ-nTST	>=45 minutes increase in average nighttime sleep duration
UQ-dTSD	>=45 minutes decrease in average daytime sleep duration
MoST	>20, >25 or >30 minutes increase and a standard deviation <=2 hours during double-masked phase
CGI-C	<=2.0 from the average of Day 112 and Day 183 compared to baseline

[0117] A second study (RESET Study) was a 20 patient randomized withdrawal study designed to demonstrate the maintenance effect of 20 mg/day tasimelteon in the treatment of blind individuals with Non-24. Patients were treated with tasimelteon for at least twelve weeks during an open-label run-in phase during the SET Study. Patients who responded to tasimelteon treatment during the run-in phase were then randomized to receive either placebo or tasimelteon (20mg/day) for 2 months.

[0118] Results relating to the primary endpoint of the SET Study are summarized in Table 1A.

Table 1A. SET Study - Primary Endpoints Results:

	Tasimelteon (%)	Placebo (%)	p-value
Entrainment (aMT6s)	20.0	2.6	0.0171
Clinical Response (Entrainment ¹ + N24CRS >=3)	23.7	0.0	0.0028
Clinical Response ² (Entrainment ¹ + N24CRS >=2)	28.9	0.0	0.0006
N24CRS >=3 ²	28.9	2.9	0.0031
N24CRS >=2 ²	57.9	20.6	0.0014
NOTES: 1) Entrainment status from the randomized portion of the SET study and/or the screening portion of the RESET study 2) Sensitivity Analysis			

[0119] The SET study also assessed a number of secondary endpoints including Entrainment of cortisol rhythm and a broad range of clinical sleep and wake parameters. These parameters included improvement in the total nighttime sleep in the worst 25% of nights (LQ-nTST), decrease in the total daytime sleep duration in the worst 25% of days (UQ-dTSD) and midpoint of sleep timing (MoST) which is derived from a combination of the sleep reported for both nighttime and daytime. CGI-C is a seven-point rating scale of global functioning with lower scores indicating larger improvements.

Table 1B. SET Study - Secondary Endpoints Results

	Tasimelteon	Placebo	p-value
Entrainment (cortisol) (%)	17.5	2.6	0.0313
N24CRS (LS mean minutes)	1.77	0.67	0.0004
CGI-C ¹ (LS mean minutes)	2.6	3.4	0.0093
LQ-nTST and UQ-dTSD >=90 min ² (%)	23.8	4.5	0.0767
LQ-nTST and UQ-dTSD >= 45 min ³ (%)	31.6	8.8	0.0177
LQ-nTST (LS mean minutes)	57.0	16.8	0.0055

(continued)

	Tasimelteon	Placebo	p-value	
5	UQ-dTSD ¹ (LS mean minutes)	-46.2	-18.0	0.0050
	MoST (LS mean minutes)	34.8	14.4	0.0123
NOTES: 1) For CGI-C and UQ-dTSD smaller numbers indicate improvement. 2) For this endpoint, only subjects with significant sleep and nap problems at baseline were included. 3) Sensitivity Analysis				

[0120] The percentage of patients entrained was higher among patients on drug for two complete circadian cycles. It was also higher among patients not taking a beta blocker and lower among patients with very long tau, e.g., $\tau \geq 24.7$. Among patients on drug for at least two circadian cycles, not on beta blockers, and $\tau < 24.7$ hours, the percentage of entrained patients was approximately 85%.

[0121] The results of the SET study represent the initial data from the tasimelteon Non-24 Phase III development program and demonstrate the multiple benefits of this novel therapy in treating patients suffering from this rare circadian rhythm disorder. In the SET study, tasimelteon was demonstrated to be safe and well tolerated.

[0122] The primary endpoint of the RESET Study was the maintenance of effect as measured by entrainment of the melatonin (aMT6s) rhythm. Results relating to the primary endpoint of the RESET Study are summarized in Table 2A.

Table 2A. RESET Study - Primary Endpoint Results:

	Tasimelteon	Placebo	p-value	
25	Maintenance of entrainment (aMT6s) (%)	90.0	20.0	0.0026

[0123] The RESET study also assessed a number of secondary endpoints including maintenance of entrainment of the cortisol rhythm and a range of sleep and wake parameters including LQ-nTST (total nighttime sleep in the worst 25% of nights), UQ-dTSD (total daytime sleep duration in the worst 25% of days) and MoST (midpoint of sleep timing from both nighttime and daytime sleep). Results relating to the secondary endpoints of the RESET Study are summarized in Table 2B.

Table 2B. RESET Study - Secondary Endpoints Results:

	Tasimelteon	Placebo	Difference	p-value	
35	maintenance of entrainment (cortisol) (%)	80.0	20.0	60.0	0.0118
	LQ-nTST (LS mean minutes) ¹	-6.6	-73.8	67.2	0.0233
40	UQ-dTSD (LS mean minutes) ²	-9.6	49.8	-59.4	0.0266
	MoST (LS mean minutes) ¹	19.8	-16.2	36.0	0.0108
NOTES: 1) Higher number indicates improvement 2) Lower number indicates improvement					

[0124] From the run-in phase of the study, the rate of entrainment among tasimelteon treated patients ranged from 50% to 85% based on individual patient characteristics. In a time to relapse analysis (45 min decrement of weekly average nighttime sleep), placebo treated patients relapsed in higher numbers and at an earlier time than tasimelteon treated patients (P = 0.0907).

[0125] The RESET study demonstrates the efficacy of chronic treatment with tasimelteon in Non-24 and further supports the results of the SET study, which established the ability of tasimelteon to entrain the master body clock and significantly improve the clinical symptoms of Non-24.

[0126] For maintenance of an entrained circadian rhythm, i.e., chronic treatment, the treatment regimens described herein can be continued daily indefinitely. So, for example, tasimelteon can be administered orally, e.g., at a dose of 20 mg/day, e.g., at about 1/2 to about 1 hour prior to bedtime.

[0127] Results of clinical study also show a strong correlation between endogenous melatonin and efficacy of tasimelteon in entraining patients to a 24 hour circadian rhythm. The following table (Table 3A) compares the peak aMT6s levels

EP 2 806 863 B1

in the 24 entrained and 23 non-entrained patients.

TABLE 3A

Peak aMT6s (ng/hr) Entrained Patients	Peak aMT6s (ng/hr) Non-entrained Patients
291.05	261.68
302.40	334.34
350.92	409.12
362.07	472.99
510.60	514.14
786.85	552.77
811.80	552.90
958.89	581.95
1102.76	810.43
1205.45	846.55
1329.08	862.91
1442.48	1155.66
1502.80	1284.35
2106.44	1295.37
2211.81	1397.71
2226.06	1444.94
2287.07	1451.43
2566.27	1622.23
2706.67	1637.45
2801.31	1719.94
2891.17	1749.32
3391.00	2329.65
3867.45	2671.17
5547.22	

[0128] The average baseline aMT6s excretion rate in urine, as determined using the methodology described above, was 1814.98 ng/hr in subjects who became entrained in response to tasimelteon therapy and 1128.65 ng/hr in subjects who did not become entrained in response to tasimelteon therapy. Eleven of thirteen patients with a baseline aMT6s excretion rate > 2000 ng/hr responded to therapy. See, Table 3B.

Table 3B

Peak aMT6s (ng/hr)	All	< 1500	≥ 1500	< 2000	≥ 2000
Total	47	29	18	34	13
Entrained	24(51%)	12 (41%)	12 (67%)	13 (38%)	11 (85%)
Non-entrained	23 (49%)	17 (59%)	6 (33%)	21 (62%)	2 (15%)

[0129] Data from these studies currently available also indicate that beta blocker therapy is indirectly related to efficacy of tasimelteon, i.e., patients receiving beta blocker therapy were less likely to become entrained than patients who were not.

EP 2 806 863 B1

TABLE 4

Taking Beta Blocker	Status	
	Entrained	Non-entrained
No	24	19
Yes	0	4

5

10 **[0130]** In addition, currently available data indicate a correlation between tau as determined by assaying for aMT6s levels in urine substantially as described above and assaying for cortisol in urine substantially as described above, as shown in Table 5.

TABLE 5

Site #	Subject #	Tau (aMT6s)	CI Low	CI High	Cycle Length (Days)	Tau (Cortisol)	CI Low	CI High	Cycle Length (Days)	P Value
405	3001	23.92	23.71	24.13	N/A	23.88	23.49	24.27	n/a	0.32
410	3002	24.02	23.86	24.19	N/A	23.92	23.64	24.21	N/A	0.37
409	3003	23.97	23.77	24.17	N/A	23.94	23.75	24.12	N/A	0.37
405	3002	23.98	23.86	24.1	N/A	23.96	23.8	24.13	n/a	0.46
405	3003	23.95	23.87	24.04	N/A	23.97	23.78	24.15	n/a	0.51
424	3003	23.96	23.8	24.12	N/A	23.99	23.92	24.05	N/A	0.46
411	3001	24.02	23.77	24.26	1482	24.01	23.48	24.54	2728	0.95
426	3002	24.01	23.87	24.15	3959	24.01	23.54	24.48	3111	0.95
410	3001	24.02	23.99	24.05	N/A	24.02	23.89	24.15	1176	0.57
412	3002	23.99	23.88	24.09	N/A	24.05	23.09	25.02	468	0.84
412	3003	23.98	23.88	24.08	N/A	24.05	23.84	24.26	460	0.4
409	3002	24.08	23.99	24.17	290	24.08	23.95	24.21	287	0.11
424	3001	23.97	23.68	24.26	N/A	24.17	24.02	24.32	140	0.04
407	3003	24.33	24.21	24.44	74	24.11	23.97	24.24	225	0.08
410	3006	24.29	23.57	25.02	83	24.12	23.65	24.58	205	0.39
407	3001	24.56	24.37	24.75	43	24.13	22.89	25.37	179	0.69
401	3002	24.31	24.22	24.4	77	24.15	24.08	24.23	158	0.01
406	3002	24.41	22.66	26.16	59	24.3	24	24.6	81	0.05
421	3001	24.86	22.57	27.14	29	24.37	21.83	26.92	65	0.31
406	3003	24.48	24.07	24.9	50	24.42	24.25	24.59	58	0.01
410	3004	24.39	24.27	24.51	62	24.43	24.4	24.47	56	0.01
403	3001	24.76	23.42	26.1	32	24.44	24.06	24.82	55	0.04
419	3001	25.28	25.04	25.51	19	24.54	24.07	25.02	45	0.04
409	3001	24.52	24.41	24.63	47	24.58	24.47	24.68	42	0.01
411	3003	24.5	24.13	24.87	49	24.61	24.28	24.94	40	0.02
411	3004	24.92	24.46	25.38	27	24.74	24.15	25.34	33	0.03
403	3002	24.8	24.59	25.01	31	24.77	23.94	25.6	32	0.06

15

20

25

30

35

40

45

50

55

EP 2 806 863 B1

(continued)

Site #	Subject #	Tau (aMT6s)	CI Low	CI High	Cycle Length (Days)	Tau (Cortisol)	CI Low	CI High	Cycle Length (Days)	P Value
425	3003	24.77	23.67	25.88	32	24.86	23.91	25.81	29	0.06
425	3002	25.01	24.63	25.4	24	25.1	24.65	25.55	22	0.01

[0131] Data from clinical studies also show that CYP1A2 inhibitors and smoking both affect patient exposure to drug.

[0132] Fluvoxamine is a strong CYP1A2 inhibitor. AUC_{0-inf} for tasimelteon increased approximately 7-fold, and the C_{max} increased approximately 2-fold upon co-administration of fluvoxamine and tasimelteon, compared to tasimelteon administered alone.

[0133] Table 6 below shows the effect of co-administration of tasimelteon and fluvoxamine on tasimelteon's pharmacokinetics. Twenty-four healthy male or female subjects between the ages of 18 and 55 years of age (inclusive) who were non-smokers with a body mass index (BMI) of ≥18 and ≤35 kg/m² participated in this open-label, single-sequence study conducted at one site. On day 1, subjects were administered 5.667 mg of tasimelteon. On days 2-7, subjects were administered 50 mg of fluvoxamine. On day 8, subjects were co-administered 5.667 mg of tasimelteon and 50 mg of fluvoxamine.

TABLE 6

Analyte	Day	Cmax (ng/ml)	Tmax (h)	AUC (inf) (h×ng/mL)	t1/2(h)	CL/F (mL/min)
Tasimelteon	1	68.0 ± 28.9	0.50	102 ± 61.5	1.20 ± 0.22	107 ± 555
Tasimelteon	8	155 ± 51.1	0.50	701 ± 402	2.59 ± 0.71	189 ± 155
Geometric Mean Ratio* (%)		232.74	N/A	653.36	211.82	15.31
M12	1	31.0 ± 7.23	0.88	189 ± 90.8	3.03 ± 1.02	N/A
M12	8	30.8 ± 17.6	3.00	435 ± 109.3	7.03 ± 3.27	N/A
Geometric Mean Ratio (%)		92.74	N/A	274.81	241.02	N/A
M13	1	87.5 ± 24.4	0.50	106 ± 32.6	1.00 ± 0.30	N/A
M13	8	63.6 ± 24.6	0.50	133 ± 32.9	3.51 ± 1.18	N/A
Geometric Mean Ratio (%)*		69.31	N/A	125.05	349.81	N/A
M9	1	67.6 ± 19.1	0.50	104 ± 30.0	1.14 ± 0.29	N/A
M9	8	47.4 ± 24.2	0.75	126 ± 29.6	3.83 ± 1.34	N/A
Geometric Mean Ratio (%)*		64.94	N/A	122.56	328.02	N/A
M11	1	15.8 ± 5.40	1.00	44.5 ± 17.2	1.61 ± 0.55	N/A
M11	8	11.0 ± 3.94	1.00	55.8 ± 18.3	4.14 ± 1.44	N/A
Geometric Mean Ratio (%)*		68.71	N/A	126.03	248.35	N/A
M14	1	1.20 ± 0.40	0.75	4.54 ± 2.39	2.18 ± 0.97	N/A

EP 2 806 863 B1

(continued)

Analyte	Day	Cmax (ng/ml)	Tmax (h)	AUC (inf) (h×ng/mL)	t1/2(h)	CL/F (mL/min)
M14	8	3.20 ± 1.49	4.00	42.6 ± 27.3	4.98 ± 1.89	N/A
Geometric Mean Ratio (%)*		264.58	N/A	944.73	243.34	N/A

[0134] FIG. 5 shows a diagram of a metabolic pathway of tasimelteon. FIGS. 6-11 show plots of the effect of co-administration of tasimelteon and fluvoxamine on the concentration of, respectively, tasimelteon, the M9 metabolite, the M11 metabolite, the M12 metabolite, the M13 metabolite, and the M14 metabolite. As can be seen from FIGS. 6-11, the increase in concentration attributable to fluvoxamine co-administration was more pronounced with respect to tasimelteon and its primary metabolites (M12, M13, M14) than its secondary metabolites (M9, M11).

[0135] Table 7 below shows the effect of smoking on the concentration of tasimelteon and several of its metabolites. Smokers were defined as those smoking 10 or more cigarettes per day. Non-smokers were defined as those smoking no cigarettes per day.

TABLE 7

Analyte	Group	Cmax (ng/ml)	Tmax (h)	AUC (inf) (h×ng/mL)	t1/2(h)	CL/F (mL/min)	Vz/F (L)
Tasimelteon	Smokers	136 ± 59.5	0.75	205 ± 152	0.99 ± 0.18	2,290 ± 1,232	189 ± 94.2
Tasimelteon	Non-Smokers	239 ± 177	0.50	389 ± 429	1.18 ± 0.46	1,482 ± 1,008	133 ± 83.0
Geometric Mean Ratio* (%)		63.98	N/A	60.14	86.84	166.27	144.39
M12	Smokers	123 ± 28	1.00	526 ± 193	2.11 ± 0.67	N/A	N/A
M12	Non-Smokers	108 ± 29	1.00	679 ± 433	3.05 ± 1.73	N/A	N/A
Geometric Mean Ratio (%)		115.53	N/A	84.87	73.31	N/A	N/A
M13	Smokers	272 ± 86	0.75	329 ± 99	0.89 ± 0.26	N/A	N/A
M13	Non-Smokers	270 ± 71	0.50	337 ± 94	1.18 ± 0.50	N/A	N/A
Geometric Mean Ratio (%)*		99.49	N/A	97.31	77.51	N/A	N/A
M9	Smokers	230 ± 118	0.75	315 ± 112	1.15 ± 0.17	N/A	N/A
M9	Non-Smokers	279 ± 82.8	0.75	406 ± 75	1.38 ± 0.45	N/A	N/A
Geometric Mean Ratio (%)*		77.18	N/A	74.36	85.40	N/A	N/A
M11	Smokers	46.17 ± 11.9	1.00	124 ± 42	1.99 ± 0.85	N/A	N/A
M11	Non-Smokers	54.9 ± 15.1	1.00	154 ± 58	2.14 ± 0.94	N/A	N/A
Geometric Mean Ratio (%)*		84.50	N/A	81.84	94.13	N/A	N/A
M14	Smokers	3.72 ± 1.86	0.75	9.45 ± 11.88	1.13 ± 0.54	N/A	N/A

(continued)

Analyte	Group	C _{max} (ng/ml)	T _{max} (h)	AUC (inf) (h×ng/mL)	t _{1/2} (h)	CL/F (mL/min)	V _z /F (L)
M14	Non-Smokers	6.18 ± 3.15	0.75	22.0 ± 24.2	1.84 ± 1.22	N/A	N/A
Geometric Mean Ratio (%)*		60.17	N/A	42.98	65.09	N/A	N/A
M3	Smokers	177 ± 71.6	0.50	239 ± 44.4	3.48 ± 2.53	N/A	N/A
M3	Non-Smokers	135 ± 49.5	0.63	194 ± 64.6	4.00 ± 2.48	N/A	N/A
Geometric Mean Ratio (%)*		131.27	N/A	129.43	89.16	N/A	N/A

[0136] FIGS. 12-17 show plots of the effect of smoking on the concentration of, respectively, tasimelteon, the M9 metabolite, the M11 metabolite, the M12 metabolite, the M13 metabolite, and the M14 metabolite.

[0137] Related aspects of this invention include computer-based systems comprising means for receiving data concerning treatment-related health information, optionally transiently or indefinitely storing such information, and directly or indirectly transmitting such information to such healthcare professional or patient. Such health information can include whether or not a patient is receiving, i.e., being treated with, a CYP1A2 inhibitor, information relating to a patient's endogenous melatonin levels, information relating to a patient's endogenous cortisol levels, information relating to a patient's tau, information relating to whether or not a patient is receiving, i.e., being treated with, a beta blocker, information relating to whether or not the patient is a smoker.

[0138] Accordingly, computer implemented systems and methods using the methods described herein are provided.

[0139] For example, related to this invention is a method comprising screening patient test samples to determine melatonin levels, collecting the data, and providing the data to a patient, a health care provider or a health care manager for making a conclusion based on review or analysis of the data. In one embodiment the conclusion is provided to a patient, a health care provider or a health care manager includes transmission of the data over a network.

[0140] Melatonin level and circadian rhythm information or other patient specific information such as recited above and as described herein, may be stored in a computer readable form. Such information can also include, e.g., one or more of whether or not a patient is being treated with a CYP1A2 inhibitor, information relating to a patient's endogenous melatonin levels, information relating to a patient's endogenous cortisol levels, information relating to a patient's tau, information relating to whether or not a patient is receiving, i.e., being treated with, a beta blocker, information relating to whether or not the patient is a smoker. Such a computer system typically comprises major subsystems such as a central processor, a system memory (typically RAM), an input/output (I/O) controller, an external device such as a display screen via a display adapter, serial ports, a keyboard, a fixed disk drive via a storage interface and optionally, a disk drive operative to receive a floppy disc, a CD or DVD, or any other data storage medium. Many other devices can be connected, such as a closed or open network interface.

[0141] The computer system may be linked to a network, comprising a plurality of computing devices linked via a data link, such as a cable, telephone line, ISDN line, wireless network, optical fiber, or other suitable signal transmission medium, whereby at least one network device (e.g., computer or disk array) comprises a pattern of magnetic domains (e.g., magnetic disk) and/or charge domains (e.g., an array of DRAM cells) composing a bit pattern encoding data acquired from an assay of the invention.

[0142] The computer system can comprise code for interpreting the results of tau analyses as described herein. Thus in an exemplary embodiment, the determination of peak melatonin levels (or surrogate) and of tau results are provided to a computer where a central processor executes a computer program for determining, e.g., optimal initiation of treatment times, the likelihood of response to treatment.

[0143] Also related to this invention is use of a computer system, such as that described above, which comprises: (1) a computer including a computer processor; (2) a stored bit pattern encoding the results obtained by the melatonin analyses of the invention, which may be stored in the computer; (3) and, optionally, (4) a program for determining the likelihood of a therapeutic response.

[0144] A computer-based system for use in the methods described herein generally includes at least one computer processor (e.g., where the method is carried out in its entirety at a single site) or at least two networked computer processors (e.g., where data is to be input by a user (also referred to herein as a "client") and transmitted to a remote site to a second computer processor for analysis, where the first and second computer processors are connected by a

network, e.g., via an intranet or internet). The system can also include a user component(s) for input; and a reviewer component(s) for review of data, generated reports, and manual intervention. Additional components of the system can include a server component(s); and a database(s) for storing data (e.g., as in a database of report elements, e.g., interpretive report elements, or a relational database (RDB) which can include data input by the user and data output. The computer processors can be processors that are typically found in personal desktop computers (e.g., IBM, Dell, Macintosh), portable computers, mainframes, minicomputers, or other computing devices.

[0145] Illustrative reports which can be displayed or projected, or printed, are provided in Figures 1, 2, 3, and 4.

[0146] A networked client/server architecture can be selected as desired, and can be, for example, a classic two or three tier client server model. A relational database management system (RDMS), either as part of an application server component or as a separate component (RDB machine) provides the interface to the database.

[0147] In one example, the architecture is provided as a database-centric client/server architecture, in which the client application generally requests services from the application server which makes requests to the database (or the database server) to populate the report with the various report elements as required, particularly the interpretive report elements, especially the interpretation text and alerts. The server(s) (e.g., either as part of the application server machine or a separate RDB/relational database machine) responds to the client's requests.

[0148] The input client components can be complete, stand-alone personal computers offering a full range of power and features to run applications. The client component usually operates under any desired operating system and includes a communication element (e.g., a modem or other hardware for connecting to a network), one or more input devices (e.g., a keyboard, mouse, keypad, or other device used to transfer information or commands), a storage element (e.g., a hard drive or other computer-readable, computer-writable storage medium), and a display element (e.g., a monitor, television, LCD, LED, or other display device that conveys information to the user). The user enters input commands into the computer processor through an input device. Generally, the user interface is a graphical user interface (GUI) written for web browser applications.

[0149] The server component(s) can be a personal computer, a minicomputer, or a mainframe and offers data management, information sharing between clients, network administration and security. The application and any databases used can be on the same or different servers.

[0150] Other computing arrangements for the client and server(s), including processing on a single machine such as a mainframe, a collection of machines, or other suitable configuration are contemplated. In general, the client and server machines work together to accomplish the processing of the present invention.

[0151] Where used, the database(s) is usually connected to the database server component and can be any device which will hold data. For example, the database can be any magnetic or optical storing device for a computer (e.g., CDROM, internal hard drive, tape drive). The database can be located remote to the server component (with access via a network or modem) or locally to the server component.

[0152] Where used in the system and methods, the database can be a relational database that is organized and accessed according to relationships between data items. The relational database is generally composed of a plurality of tables (entities). The rows of a table represent records (collections of information about separate items) and the columns represent fields (particular attributes of a record). In its simplest conception, the relational database is a collection of data entries that "relate" to each other through at least one common field.

[0153] Additional workstations equipped with computers and printers may be used at point of service to enter data and, in some embodiments, generate appropriate reports, if desired. The computer(s) can have a shortcut (e.g., on the desktop) to launch the application to facilitate initiation of data entry, transmission, analysis, report receipt, as desired.

[0154] The present invention also contemplates a computer-readable storage medium (e.g. CD-ROM, memory key, flash memory card or diskette) having stored thereon a program which, when executed in a computing environment, provides for implementation of algorithms to carry out all or a portion of the results of a response likelihood assessment as described herein. Where the computer-readable medium contains a complete program for carrying out the methods described herein, the program includes program instructions for collecting, analyzing and generating output, and generally includes computer readable code devices for interacting with a user as described herein, processing that data in conjunction with analytical information, and generating unique printed or electronic media for that user.

[0155] Where the storage medium provides a program that provides for implementation of a portion of the methods described herein (e.g., the user-side aspect of the methods (e.g., data input or report receipt capabilities)), the program provides for transmission of data input by the user (e.g., via the internet or via an intranet) to a computing environment at a remote site. Processing or completion of processing of the data is carried out at the remote site to generate a report. After review of the report, and completion of any needed manual intervention, to provide a complete report, the complete report is then transmitted back to the user as an electronic document or printed document (e.g., fax or mailed paper report). The storage medium containing a program according to the invention can be packaged with instructions (e.g., for program installation or use) recorded on a suitable substrate or a web address where such instructions may be obtained. The computer-readable storage medium can also be provided in combination with one or more reagents for carrying out response likelihood assessment.

[0156] Also related to this invention are methods of generating a report based on the analyses of melatonin levels in a patient suffering from Non-24. In general, such method can comprise the steps of determining information indicative of the levels of endogenous melatonin, in a biological sample; and creating a report summarizing said information, such as by reporting whether or not a patient is being treated with a CYP1A2 inhibitor, with or without additional information.

In one illustrative embodiment of the method, said report includes one or more of an indication of whether or not a patient's melatonin levels achieve a Threshold Concentration, an indication of the patient's cortisol levels, an indication of the patient's tau, an indication of whether or not the patient is being treated with a CYP1A2 inhibitor, information relating to whether or not the patient is a smoker, and an indication of whether or not the patient is being treated with an agent that reduces endogenous melatonin such as a beta blocker.

[0157] In some embodiments, the report includes a Threshold Concentration and, optionally, the peak melatonin concentration in the patient's biological sample. In some embodiments, the report includes information relating to the co-administration of tasimelteon and a CYP1A2 inhibitor, such as information relating to increased exposure to tasimelteon that may ensue, information related reducing the dose of tasimelteon or of the CYP1A2 inhibitor or information relating to heightened monitoring. In some embodiments, the report includes information relating to the administration of tasimelteon and smoking, such as information related to decreased exposure to tasimelteon that may ensue, information relating to increasing the dose of tasimelteon or information related to monitoring for levels of tasimelteon in the blood.

[0158] Such report can further include one or more of: 1) information regarding the testing facility; 2) service provider information; 3) patient data; 4) sample data; 5) an interpretive report, which can include various information including: a) indication; b) test data, and 6) other features.

[0159] In some embodiments, the report further includes a recommendation for a treatment modality for said patient. In such aspect, the report may include information to support a treatment recommendation for said patient, e.g., a recommendation for non-treatment with a melatonin agonist or for heightened monitoring. In all aspects, the report may include a classification of a subject into a group, e.g., likely non-responders or likely responders.

[0160] In some embodiments, the report is in electronic form e.g., presented on an electronic display (e.g., computer monitor).

[0161] In some embodiments, the report is a visual report comprising:

- 1) a descriptive title
- 2) a patient identifier
- 3) the patient's target initiation of sleep time and one or more of:

(i) a graph of rate of production of melatonin or melatonin surrogate versus time for each Collection Session, the graph showing data points and the calculated circadian cycle including acrophase, each graph being annotated with the projected acrophase and Standard Error,

(ii) a graph of acrophase (time of day) vs. Day showing the projected acrophase determined for each Collection Session and the slope determined by linear regression analysis of the projected acrophase times, said graph being annotated with the length of the patient's tau, the Standard Error and the Confidence Interval expressed both as a p value and as a range of hours, and

(iii) an acrophase table showing the projected time of acrophase for 90 days following the end of the last Collection Session, said table differentially highlighting the date and time of the projected acrophase closest to the target acrophase, the optimal day for initiation of treatment and an estimated window for initiation of treatment.

[0162] Such illustrative report is provided in Fig. 1 for a subject that is not suffering Non-24 and in Fig. 2 for a patient that is suffering from N24SWD.

[0163] A person or entity who prepares a report ("report generator") may also perform the likelihood assessment. The report generator may also perform one or more of sample gathering, sample processing, and data generation, e.g., the report generator may also perform one or more of: a) sample gathering; b) sample processing; c) measuring melatonin or melatonin surrogate levels. Alternatively, an entity other than the report generator can perform one or more sample gathering, sample processing, and data generation.

[0164] For clarity, it should be noted that the term "user," which is used interchangeably with "client," is meant to refer to a person or entity to whom a report is transmitted, and may be the same person or entity who does one or more of the following: a) collects a sample; b) processes a sample; c) provides a sample or a processed sample; and d) generates data for use in the likelihood assessment. In some cases, the person(s) or entity(ies) who provides sample collection and/or sample processing and/or data generation, and the person who receives the results and/or report may be different persons, but are both referred to as "users" or "clients" herein to avoid confusion. In certain embodiments, e.g., where the methods are completely executed on a single computer, the user or client provides for data input and review of data output. A "user" can be a health professional (e.g., a clinician, a laboratory technician, or a physician).

[0165] In embodiments where the user only executes a portion of the method, the individual who, after computerized data processing according to the methods of the invention, reviews data output (e.g., results prior to release to provide a complete report, a complete, or reviews an "incomplete" report and provides for manual intervention and completion of an interpretive report) is referred to herein as a "reviewer." The reviewer may be located at a location remote to the user (e.g., at a service provided separate from a healthcare facility where a user may be located).

[0166] Where government regulations or other restrictions apply (e.g., requirements by health, malpractice or liability insurance, or policy), results, whether generated wholly or partially electronically, are subjected to a quality control routine prior to release to the user.

[0167] In another aspect, the present disclosure concerns methods of preparing a personalized pharmacologic profile for a patient by a) determining the patient's levels of endogenous melatonin or melatonin surrogate; and (b) creating a report summarizing the data and/or compiling such data with other data relevant to understanding the patient's specific pharmacologic characteristics and condition.

[0168] In accordance with the method of this invention, the dosage of tasimelteon to be administered will depend on various factors such as the characteristics of the subject being treated, e.g., the severity of disorder, responsiveness to melatonin agonists, age, weight, health, types of concurrent treatment, if any.

[0169] The above described computer-implemented methods, systems or reports, can also be applied to determination of efficacy of treatment, such as the efficacy determination methodologies described above. For example, computer-based systems can be used to record and report information relating to one or more of MoST, LQ-nTST, UQ-dTSD and CGI-C and/or to tau determinations made prior to or shortly after initiation of therapy as well as subsequent tau determinations.

[0170] By way of further illustration, related aspects of this invention include computer-based systems comprising means for receiving data concerning one or more of MoST, LQ-nTST, UQ-dTSD and CGI-C and/or to tau determinations made prior to or shortly after initiation of therapy as well as subsequent tau determinations;

a method comprising collecting data relating to one or more of MoST, LQ-nTST, UQ-dTSD and CGI-C and/or to tau determinations made prior to or shortly after initiation of therapy as well as subsequent tau determinations and providing the data to a patient, a health care provider or a health care manager for making a conclusion based on review or analysis of the data. In one embodiment the conclusion is provided to a patient, a health care provider or a health care manager includes transmission of the data over a network;

information relating to one or more of MoST, LQ-nTST, UQ-dTSD and CGI-C and/or to tau determinations made prior to or shortly after initiation of therapy as well as subsequent tau determinations stored in a computer readable form;

a computer system as described above for receiving, storing and outputting such information, optionally linked to a network and optionally comprising code for interpreting the results of efficacy assessment(s) as described herein;

a computer-readable storage medium (e.g., CD-ROM, memory key, flash memory card or diskette) having stored thereon a program which, when executed in a computing environment, provides for implementation of algorithms to carry out all or a portion of the analysis of efficacy assessments as described herein;

methods of generating a report based on the efficacy assessments as described herein, e.g., a report that includes one or more of an indication of whether or not a patient is responding to therapy.

[0171] Such information, databases, systems, methods, analyses, reports, profiles, outputs or recommendations, can be incorporated into storage media, computer systems, and networks, such as are described hereinabove with respect to other parameters, e.g., melatonin levels, circadian rhythms, cortisol levels, tau, co-treatment with CYP1A2 inhibitors, co-treatment with a beta blocker, and smoking, with or without information relating to some or all of such other parameters.

[0172] An effective dose is one that over a period of time of treatment, which may be, e.g., 1 day or multiple weeks, results in entrainment of the patient to a 24 hour circadian rhythm. Patients whose tau is reduced to 24 hours, e.g., <24.1 hrs, with a 95% confidence interval that includes 24.0 can be considered to have been entrained, although other values can also be used to define successful entrainment.

[0173] The daily dose of tasimelteon useful in entraining patients with Non-24 to a 24 hour circadian rhythm will, in general, be in the range of about 1 to about 100 mg, e.g., about 10 to about 100 or about 20 to about 50. A dose of 20 mg is typically sufficient, in particular, for individuals who are not also being administered a CYP1A2 inhibitor or a beta blocker or who are not smokers.

[0174] Similar doses may be employed when entraining a patient's cortisol circadian rhythm.

[0175] As discussed above, it has been found that co-administration of tasimelteon with CYP1A2 inhibitors unexpectedly increases the concentration of tasimelteon. This is likely a consequence of inhibition of CYP1A2-mediated conversion of tasimelteon to a metabolite.

[0176] CYP1A2 inhibitors include, for example, fluoroquinolone antibiotics, such as ciprofloxacin, SSRIs such as fluvoxamine, and calcium channel blockers such as verapamil. Accordingly, in the case that a patient is to be administered a dose of tasimelteon as part of an attempt to entrain the patient to a 24-hour circadian rhythm and that patient is also being treated with a CYP1A2 inhibitor, it may be necessary or desirable to reduce the dose of tasimelteon, the dose of the CYP1A2 inhibitor, or both. Alternatively, or in addition, it may be necessary or desirable to monitor the patient's

plasma concentration of tasimelteon or monitor the patient for an adverse reaction associated with tasimelteon.

[0177] For example, the dose of tasimelteon administered to a patient also being treated with a CYP1A2 inhibitor may be reduced to less than 20 mg per day, e.g., about 15 to about 19 mg per day, about 10 to about mg per day, or about 5 to about 10 mg per day, e.g., 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or 19 mg/day. In some cases, the dose of tasimelteon or the dose of the CYP1A2 inhibitor may be reduced to zero. In an embodiment of the invention, tasimelteon is not be used in combination with fluvoxamine. Other less strong CYP1A2 inhibitors have not been adequately studied. Tasimelteon should be administered with caution to patients taking less strong CYP1A2 inhibitors.

[0178] Aspects of the invention, as they relate to the effects of a CYP1A2 inhibitor on tasimelteon exposure, include, without limitation, the following:

treating a patient with tasimelteon wherein the patient is also being treated with a CYP1A2 inhibitor, said method comprising one or more of the following: reducing the dose of tasimelteon, reducing the dose of the CYP1A2 inhibitor, monitoring the patient's plasma concentration of tasimelteon, or monitoring the patient for an adverse reaction associated with tasimelteon;

treating a patient with tasimelteon wherein the patient is also being treated with a substance that is a known inhibitor of CYP1A2, said method comprising monitoring the patient for a potential or actual adverse event associated with increased plasma concentration of tasimelteon while the patient is being coadministered tasimelteon and the CYP1A2 inhibitor;

treating a patient suffering from a sleep disorder wherein such patient is being treated with a CYP1A2 inhibitor, the method comprising: internally administering tasimelteon to the patient in a reduced amount relative to an amount that would be administered to a patient suffering from a sleep disorder but not being treated with a CYP1A2 inhibitor; a computing device having a processor; a storage device containing information that the patient is being treated with a CYP1A2 inhibitor; an input device for inputting to either or both of the computing device or the storage device information that the patient will be prescribed a dose of tasimelteon; a computer program operable retrieve from the storage device the information that the patient is being treated with a CYP1A2 inhibitor upon inputting the information that the patient will be prescribed the dose of tasimelteon; and an output device for outputting to a user the information that the patient is being treated with a CYP1A2 inhibitor;

a computer-implemented method of treating a patient suffering from a sleep disorder, the method comprising: entering into an electronic database information related to the treatment of a patient with tasimelteon; searching, using a computing device, a medical record of the patient for information related to the current treatment of the patient with an agent other than tasimelteon; and determining, using the computing device, whether the agent other than tasimelteon is a CYP1A2 inhibitor;

a pharmaceutical composition for the treatment of a sleep disorder in an individual being treated with a CYP1A2 inhibitor, the composition comprising: a pharmaceutically-acceptable carrier; and a quantity of tasimelteon corresponding to a daily dosage of less than 20 mg.

[0179] In another embodiment, patients who are receiving a CYP1A2 inhibitor, e.g., fluvoxamine, are not treated with tasimelteon. In a related embodiment, patients are instructed not to receive, and healthcare providers are instructed not to prescribe, tasimelteon if the patient is already receiving a CYP1A2 inhibitor, e.g., fluvoxamine.

[0180] Smoking, on the other hand, has been found to increase the clearance of tasimelteon, thereby reducing patient exposure. Accordingly, administration of tasimelteon or a tasimelteon metabolite to an individual who smokes may, in some cases, require increasing the dose of tasimelteon or tasimelteon metabolite and/or reducing or eliminating the individual's smoking.

[0181] Accordingly, in the case that a patient is to be administered a dose of tasimelteon as part of an attempt to entrain the patient to a 24-hour circadian rhythm and that patient is also a smoker, it may be necessary or desirable to increase the dose of tasimelteon. Alternatively, or in addition, it may be necessary or desirable to monitor the patient's plasma concentration of tasimelteon.

[0182] For example, the dose of tasimelteon administered to a patient who also smokes may be increased to greater than 20 mg per day, e.g., 25 mg per day, 30 mg per day, 40 mg per day, 50 mg per day or even 100 mg per day.

[0183] Aspects of the invention, as they relate to the effects of smoking on tasimelteon exposure, include, without limitation, the following:

treating a patient with tasimelteon wherein the patient is a smoker, said method comprising one or more of the following: increasing a dose of tasimelteon, monitoring the patient's blood levels of tasimelteon, and instructing the patient to reduce or eliminate smoking;

treating a patient suffering from a sleep disorder wherein such patient is a smoker, the method comprising: internally administering tasimelteon to the patient in an increased amount relative to an amount that would be administered to a patient suffering from a sleep disorder who is not a smoker;

a system comprising: at least one computing device having a processor; a storage device containing information that the patient is a smoker; an input device for inputting to either or both of the computing device or the storage device information that the patient will be prescribed a dose of tasimelteon; a computer program operable retrieve from the storage device the information that the patient is a smoker upon inputting the information that the patient will be prescribed the dose of tasimelteon; and an output device for outputting to a user the information that the patient is a smoker;

a computer-implemented method of treating a patient suffering from a sleep disorder, the method comprising: entering into an electronic database information related to the treatment of a patient with tasimelteon; searching, using a computing device, a medical record of the patient for information related to whether the patient is a smoker; and determining, using the computing device, whether the patient is a smoker;

a pharmaceutical composition for the treatment of a sleep disorder in an individual who smokes, the composition comprising: a pharmaceutically-acceptable carrier; and a quantity of tasimelteon corresponding to a daily dosage of greater than 20 mg.

[0184] In general, the melatonin (MT1 and MT2 receptors) agonist, e.g., tasimelteon, is administered in a pharmaceutical formulation q.d. prior to the start of the target sleep time. It has been found that in treating Non-24, it is not necessary to administer the drug more than about 1 hour prior to the start of the target sleep time such that the drug can be administered, e.g., at about 0.5 to about 1.5 hours prior to sleep time. Administration about 1 hour prior to sleep time is convenient and useful. However, this invention also contemplates administration at earlier times in the day, e.g., about 2 hours, or about 3 hours or even about 4 hours prior to target sleep time.

[0185] The ability to administer tasimelteon as little as about one hour prior to sleep time is advantageous because it allows for avoidance of pre-sleep time soporific effects, because it allows for administration of higher doses that might have greater soporific effects, and because it allows for pharmacologic intervention at a different phase of the sleep cycle than if it were administered earlier. Without wishing to be bound to any particular theory, it appears that the ability to administer tasimelteon so close to sleep time is a function of its t_{max} , which is approximately one-half hour. Melatonin, on the other hand, which has a t_{max} of approximately 2 hours or more, is administered several hours before sleep time, which can cause premature sleepiness; to avoid this soporific effect, melatonin is sometimes administered at sub-optimal doses.

[0186] Thus, in a related aspect, this invention comprises a method of treating Non-24 patients, i.e., entraining such patients to a 24 hour circadian rhythm by internally administering an effective amount of a tasimelteon or another melatonin agonist that has a t_{max} of less than about 2 hours, e.g., less than about 1.5 hours, or even less than about 1 hour such as about one-half hour like tasimelteon. Pharmaceutical compositions can be formulated so as to alter t_{max} . Thus, e.g., use of an active pharmaceutical ingredient such as melatonin that is formulated such that its t_{max} is less than about two hours, e.g., less than about 1.5 hours, or even less than about 1 hour, to treat Non-24 is an aspect of this invention.

[0187] Pharmaceutical compositions to be used comprise a therapeutically effective amount of tasimelteon or an active metabolite of tasimelteon, or a pharmaceutically acceptable salt or other form (e.g., a solvate) thereof, together with one or more pharmaceutically acceptable excipients. The phrase "pharmaceutical composition" refers to a composition suitable for administration in medical use. It should be appreciated that the determinations of proper dosage forms, dosage amounts, and routes of administration for a particular patient are within the level of ordinary skill in the pharmaceutical and medical arts.

[0188] Administration is typically oral but other routes of administration are useful, e.g., parenteral, nasal, buccal, transdermal, sublingual, intramuscular, intravenous, rectal or vaginal. Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the compound is admixed with at least one inert pharmaceutically acceptable excipient such as (a) fillers or extenders, as for example, starches, lactose, sucrose, glucose, mannitol, and silicic acid, (b) binders, as for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, (c) humectants, as for example, glycerol, (d) disintegrating agents, as for example, agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain complex silicates, and sodium carbonate, (e) solution retarders, as for example paraffin, (f) absorption accelerators, as for example, quaternary ammonium compounds, (g) wetting agents, as for example, cetyl alcohol, and glycerol monostearate, (h) adsorbents, as for example, kaolin and bentonite, and (i) lubricants, as for example, talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, or mixtures thereof. In the case of capsules, tablets, and pills, the dosage forms may also comprise buffering agents. Solid dosage forms such as tablets, dragees, capsules, pills, and granules also can be prepared with coatings and shells, such as enteric coatings and others well known in the art. The solid dosage form also may contain opacifying agents, and can also be of such composition that they release the active compound or compounds in a certain part of the intestinal tract in a delayed manner. Examples of embedding compositions which can be used are polymeric substances and waxes. The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned excipients. Such solid dosage forms may generally contain from 1% to 95%

(w/w) of the active compound. In certain embodiments, the active compound ranges from 5% to 70% (w/w).

[0189] Solid compositions for oral administration can be formulated in a unit dosage form, each dosage containing from about 1 to about 100 mg of active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired prophylactic or therapeutic effect over the course of a treatment period, in association with the required pharmaceutical carrier. Tasimelteon can be formulated, e.g., in a unit dosage form that is a capsule having 20 mg of active in addition to excipients.

[0190] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs. In addition to the compound or composition, the liquid dosage forms may contain inert diluents commonly used in the art, such as water or other solvents, solubilizing agents and emulsifiers, as for example, ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propyleneglycol, 1,3-butyleneglycol, dimethylformamide, oils, in particular, cottonseed oil, groundnut oil, corn germ oil, olive oil, castor oil and sesame oil, glycerol, tetrahydrofurfuryl alcohol, polyethyleneglycols and fatty acid esters of sorbitan or mixtures of these substances. Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0191] The present invention can be carried out in conjunction with other treatment approaches, e.g., in combination with a second or multiple other active pharmaceutical agents, including other agents that affect insomnia, sleep-wake patterns, vigilance, depression, or psychotic episodes.

[0192] While this invention has been described in conjunction with the specific embodiments outlined above, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art or are otherwise intended to be embraced. Accordingly, the embodiments of the invention as set forth above are intended to be illustrative, not limiting.

Claims

1. Tasimelteon for use in a method of treating a patient suffering from Non 24 by entraining the patient to a 24 hour sleep-wake cycle in which the patient awakens at or near a target wake time following a daily sleep period of approximately 7 to 9 hours, wherein said method comprises internally administering to the patient 20 to 50 mg of tasimelteon, the tasimelteon is administered orally once daily 0.5 to 1.5 hours before a target sleep time and the patient is light perception impaired.
2. Tasimelteon for use according to claim 1, wherein the dose of tasimelteon is 20 mg.
3. Tasimelteon for use according to any one of the preceding claims, wherein in said method the patient is also being treated with a CYP1A2 inhibitor and wherein the method further comprises one or both of:
 - monitoring the patient's plasma concentration of tasimelteon;
 - monitoring the patient for an adverse reaction associated with tasimelteon.
4. Tasimelteon for use according to any one of the preceding claims, wherein the method is for the chronic treatment of Non-24 in a patient who is totally blind.
5. Tasimelteon for use according to any one of the preceding claims, wherein the tasimelteon is administered one hour before the target sleep time.
6. Tasimelteon for use according to any one of the preceding claims, wherein entraining the patient to a 24 hour sleep-wake cycle comprises entraining the patient's abnormal melatonin circadian rhythm and/or abnormal cortisol circadian rhythm to a 24 hour circadian rhythm.

Patentansprüche

1. Tasimelteon für die Verwendung bei einem Verfahren zum Behandeln eines Patienten, der an Non-24 leidet, durch Einkoppeln des Patienten in einen 24-stündigen Schlaf-Wach-Zyklus, bei dem der Patient bei oder nahe einer Ziel-Aufwachzeit erwacht, die einer täglichen Schlafperiode von etwa 7 bis 9 Stunden folgt, wobei das Verfahren inneres Verabreichen von 20 bis 50 mg Tasimelteon an den Patienten umfasst,

EP 2 806 863 B1

wobei das Tasimelteon einmal täglich 0,5 bis 1,5 Stunden vor einer Ziel-Einschlafzeit oral verabreicht wird und der Patient Lichtwahrnehmungs-beeinträchtigt ist.

- 5
2. Tasimelteon für die Verwendung gemäß Anspruch 1, wobei die Dosis von Tasimelteon 20 mg beträgt.
 3. Tasimelteon für die Verwendung gemäß einem der vorstehenden Ansprüche, wobei bei dem Verfahren der Patient auch mit einem CYP1A2-Hemmer behandelt wird und wobei das Verfahren ferner eines oder beides umfasst von:

10 Überwachen der Plasmakonzentration von Tasimelteon des Patienten;
Überwachen des Patienten auf eine Nebenwirkung in Verbindung mit Tasimelteon.

4. Tasimelteon für die Verwendung gemäß einem der vorstehenden Ansprüche, wobei das Verfahren für die chronische Behandlung von Non-24 bei einem Patienten, der vollständig blind ist, ist.

- 15
5. Tasimelteon für die Verwendung gemäß einem der vorstehenden Ansprüche, wobei das Tasimelteon eine Stunde vor der Ziel-Einschlafzeit verabreicht wird.

- 20
6. Tasimelteon für die Verwendung gemäß einem der vorstehenden Ansprüche, wobei das Einkoppeln des Patienten in einen 24-stündigen Schlaf-Wach-Zyklus Einkoppeln des anomalen Melatonin-Tagesrhythmus und/oder anomalen Cortisol-Tagesrhythmus des Patienten in einen 24-stündigen Tagesrhythmus umfasst.

Revendications

- 25
1. Tasimeltéon pour utilisation dans un procédé de traitement d'un patient souffrant de Non-24 par entraînement du patient dans un cycle de sommeil-veille de 24 heures dans lequel le patient se réveille à ou à proximité d'une heure de réveil cible après une période de sommeil journalière d'approximativement 7 à 9 heures, où ledit procédé comprend l'administration interne au patient de 20 à 50 mg de tasimeltéon, le tasimeltéon est administré par voie orale une fois par jour 0,5 à 1,5 heure avant une heure de sommeil cible et le patient présente une perception à la lumière altérée.
- 30

2. Tasimeltéon pour utilisation selon la revendication 1, la dose de tasimeltéon étant de 20 mg.

- 35
3. Tasimeltéon pour utilisation selon l'une quelconque des revendications précédentes, où, dans ledit procédé, le patient est en outre traité avec un inhibiteur de CYP1A2 et le procédé comprenant en outre l'un ou les deux de :

la surveillance de la concentration plasmatique de tasimeltéon du patient ;
la surveillance du patient pour une réaction indésirable associée au tasimeltéon.

- 40
4. Tasimeltéon pour utilisation selon l'une quelconque des revendications précédentes, le procédé étant pour le traitement chronique de Non-24 in chez un patient qui est totalement aveugle.



- 45
5. Tasimeltéon pour utilisation selon l'une quelconque des revendications précédentes, le tasimeltéon étant administré une heure avant l'heure de sommeil cible.

- 50
6. Tasimeltéon pour utilisation selon l'une quelconque des revendications précédentes, l'entraînement du patient dans un cycle de sommeil-veille de 24 heures comprenant l'entraînement du rythme circadien de mélatonine anormal et/ou du rythme circadien de cortisol anormal du patient vers un rythme circadien de 24 heures.
- 55

VANDA Pharmaceuticals Trial VP-VEC-162-3201
Acrophase and Tau Calculations: Screening Phase

Subject: 403-1011


Target Bedtime: 22:00



ACROPHASE			
Target Acrophase: 03:00			
November 27, 2010	03:37	December 27, 2010	03:28
November 28, 2010	03:37	December 28, 2010	03:27
November 29, 2010	03:36	December 29, 2010	03:27
November 30, 2010	03:36	December 30, 2010	03:27
December 01, 2010	03:36	December 31, 2010	03:26
December 02, 2010	03:35	January 01, 2011	03:26
December 03, 2010	03:35	January 02, 2011	03:26
December 04, 2010	03:35	January 03, 2011	03:25
December 05, 2010	03:34	January 04, 2011	03:25
December 06, 2010	03:34	January 05, 2011	03:25
December 07, 2010	03:34	January 06, 2011	03:25
December 08, 2010	03:33	January 07, 2011	03:24
December 09, 2010	03:33	January 08, 2011	03:24
December 10, 2010	03:33	January 09, 2011	03:24
December 11, 2010	03:32	January 10, 2011	03:23
December 12, 2010	03:32	January 11, 2011	03:23
December 13, 2010	03:32	January 12, 2011	03:23
December 14, 2010	03:32	January 13, 2011	03:22
December 15, 2010	03:31	January 14, 2011	03:22
December 16, 2010	03:31	January 15, 2011	03:22
December 17, 2010	03:31	January 16, 2011	03:22
December 18, 2010	03:30	January 17, 2011	03:21
December 19, 2010	03:30	January 18, 2011	03:21
December 20, 2010	03:30	January 19, 2011	03:21
December 21, 2010	03:29	January 20, 2011	03:20
December 22, 2010	03:29	January 21, 2011	03:20
December 23, 2010	03:29	January 22, 2011	03:20
December 24, 2010	03:29	January 23, 2011	03:19
December 25, 2010	03:28	January 24, 2011	03:19
December 26, 2010	03:28	January 25, 2011	03:19
January 26, 2011	03:18	February 01, 2011	03:17
January 27, 2011	03:18	February 02, 2011	03:16
January 28, 2011	03:18	February 03, 2011	03:16
January 29, 2011	03:18	February 04, 2011	03:16
January 30, 2011	03:17	February 05, 2011	03:15
January 31, 2011	03:17	February 06, 2011	03:15
February 01, 2011	03:17	February 07, 2011	03:15
February 02, 2011	03:16	February 08, 2011	03:14
February 03, 2011	03:16	February 09, 2011	03:14
February 04, 2011	03:16	February 10, 2011	03:14
February 05, 2011	03:15	February 11, 2011	03:13
February 06, 2011	03:15	February 12, 2011	03:13
February 07, 2011	03:15	February 13, 2011	03:13
February 08, 2011	03:14	February 14, 2011	03:13
February 09, 2011	03:14	February 15, 2011	03:12
February 10, 2011	03:14	February 16, 2011	03:12
February 11, 2011	03:13	February 17, 2011	03:12
February 12, 2011	03:13	February 18, 2011	03:11
February 13, 2011	03:13	February 19, 2011	03:11
February 14, 2011	03:13	February 20, 2011	03:11
February 15, 2011	03:12	February 21, 2011	03:10
February 16, 2011	03:12	February 22, 2011	03:10
February 17, 2011	03:12	February 23, 2011	03:10
February 18, 2011	03:11	February 24, 2011	03:10
February 19, 2011	03:11		
February 20, 2011	03:11		
February 21, 2011	03:10		
February 22, 2011	03:10		
February 23, 2011	03:10		
February 24, 2011	03:10		

Optimal Acrophase: blue
Visit 1 Target Days: Red: optimal (1 day before optimal acrophase); Green: possible

VANDA Pharmaceuticals Trial VP-VEC-162-3201
 Acrophase and Tau Calculations: Screening Phase
 Subject: 411-1001
 Target Bedtime: 22:00



ACROPHASE					
Target Acrophase: 03:00					
November 24, 2010	20:52	December 25, 2010	15:21	January 25, 2011	09:50
November 25, 2010	21:29	December 26, 2010	15:58	January 26, 2011	10:27
November 26, 2010	22:06	December 27, 2010	16:35	January 27, 2011	11:04
November 27, 2010	22:43	December 28, 2010	17:12	January 28, 2011	11:41
November 28, 2010	23:20	December 29, 2010	17:49	January 29, 2011	12:18
November 29, 2010	23:57	December 30, 2010	18:26	January 30, 2011	12:55
December 01, 2010	00:34	December 31, 2010	19:03	January 31, 2011	13:32
December 02, 2010	01:11	January 01, 2011	19:40	February 01, 2011	14:09
December 03, 2010	01:48	January 02, 2011	20:17	February 02, 2011	14:46
December 04, 2010	02:25	January 03, 2011	20:54	February 03, 2011	15:23
December 05, 2010	03:02	January 04, 2011	21:31	February 04, 2011	16:00
December 06, 2010	03:39	January 05, 2011	22:08	February 05, 2011	16:37
December 07, 2010	04:16	January 06, 2011	22:45	February 06, 2011	17:14
December 08, 2010	04:53	January 07, 2011	23:22	February 07, 2011	17:51
December 09, 2010	05:30	January 08, 2011	23:59	February 08, 2011	18:28
December 10, 2010	06:07	January 09, 2011	00:36	February 09, 2011	19:05
December 11, 2010	06:44	January 10, 2011	01:13	February 10, 2011	19:42
December 12, 2010	07:21	January 11, 2011	01:50	February 11, 2011	20:19
December 13, 2010	07:58	January 12, 2011	02:27	February 12, 2011	20:56
December 14, 2010	08:35	January 13, 2011	03:03	February 13, 2011	21:32
December 15, 2010	09:11	January 14, 2011	03:40	February 14, 2011	22:09
December 16, 2010	09:48	January 15, 2011	04:17	February 15, 2011	22:46
December 17, 2010	10:25	January 16, 2011	04:54	February 16, 2011	23:23
December 18, 2010	11:02	January 17, 2011	05:31	February 17, 2011	00:00
December 19, 2010	11:39	January 18, 2011	06:08	February 18, 2011	00:37
December 20, 2010	12:16	January 19, 2011	06:45	February 19, 2011	01:14
December 21, 2010	12:53	January 20, 2011	07:22	February 20, 2011	01:51
December 22, 2010	13:30	January 21, 2011	07:59	February 21, 2011	02:28
December 23, 2010	14:07	January 22, 2011	08:36	February 22, 2011	03:05
December 24, 2010	14:44	January 23, 2011	09:13	February 23, 2011	03:42

Optimal Acrophase: blue
 Visit 1 Target Days: Red: optimal (1 day before optimal acrophase); Green: possible

FIG. 3

VANDA Pharmaceuticals Trial VP-VEC-162-3203
 Acrophase Calculations
 Subject: 409-3003
 Target Bedtime: 21:00




FIG. 4
VANDA Pharmaceuticals Trial VP-VEC-162-3203
Acrophase Calculations
Subject: 410-3004
Target Bedtime: 23:00




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15







FIG. 17



REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5856529 A [0019]
- US 20090105333 A [0019]
- US 6211225 B [0025]
- US 4997845 A [0025]
- US 5225442 A [0025]
- US 20050164987 A [0026]
- US 5093352 A [0026]
- US 20010047016 A [0026]
- US 5151446 A [0026]
- US 4880826 A [0026]
- US 5580878 A [0026]
- US 7754902 B [0028]

Non-patent literature cited in the description

- **UCHIYAMA, M.** Sleep Medicines Clinics. W.B Saunders Co, 2009, vol. 4, 195-211 [0009]
- **VACHHARAJANI et al.** Preclinical Pharmacokinetics and Metabolism of BMS-214778, a Novel Melatonin Receptor Agonist. *J. Pharmaceutical Sci.*, vol. 92 (4), 760-772 [0024]
- **KATO, K. et al.** *Int. J. Neuropsychopharmacol.*, 2000, vol. 3 (1) [0026]
- **MISSBACH et al.** *J. Biol. Chem.*, 1996, vol. 271, 13515-22 [0026]
- **BERESFORD et al.** *J. Pharmacol. Exp. Ther.*, 1998, vol. 285, 1239-1245 [0026]
- **DEPRES-BRUMMER et al.** *Eur. J. Pharmacol.*, 1998, vol. 347, 57-66 [0026]
- *Neuropharmacology*, July 2000 [0026]
- *Naunyn Schmiedebergs Arch.*, June 2003 [0026]
- *IMSWorld R&D Focus*, August 2002 [0026]
- *Pharmaprojects*, August 1998 [0026]
- *Chem. Pharm. Bull.*, January 2002 [0026]
- *J. Pineal Research*, November 2000 [0026]
- *Chem. Pharm. Bull.*, February 2002 [0026]
- *Reprod. Nutr. Dev.*, May 1999 [0026]
- *J. Med. Chem.*, October 1993 [0026]
- *Famaco*, March 2000 [0026]
- *J. Med. Chem.*, March 2000 [0026]
- *Bioorg. Med. Chem. Lett.*, March 2003 [0026]
- *MedAd News*, September 2001 [0026]
- **DRIJFHOUT et al.** *Eur. J. Pharmacol.*, 1999, vol. 382, 157-66 [0026]
- **BENLOUCIF et al.** *J Clin Sleep Med*, 2008, vol. 4 (1), 66-69 [0042]
- **LEHMANN E.** *Pharmacopsychiatry*, 1984, vol. 17, 71-75 [0077]

Szabadalmi igénypontok

1. Tazimelteon nem 24 órás életritmustól (non 24) szenvedő beteg kezelésére szolgáló eljárásban történő alkalmazásra, a beteg 24 órás alvás-ébrenlét ciklusra való visszatéréssel, ahol a beteg pontosan vagy megközelítőleg egy megcélzott ébredési időben ébred fel egy kb. 7-9 órás napi alvási periódust követően, ahol az említett eljárás tartalmazza 20-50 mg tazimelteon belsőleg történő adagolását, a tazimelteon orális úton napi egyszeri adagolását 0,5 – 1,5 órával a megcélzott alvási idő előtt, és ahol a beteg fényérzékelése sérült.
2. Tazimelteon alkalmazásra az 1. igénypont szerinti, ahol a tazimelteon dózisa 20 mg.
3. Tazimelteon alkalmazásra az előző igénypontok bármelyike szerinti, ahol az eljárásban a beteg CYP1A2 gátlóval is kezeli, és ahol az eljárás tartalmazza továbbá a következő lépések egyikét vagy mindkettőt:
a beteg tazimelteon plazmakoncentrációjának a vizsgálata;
a beteg vizsgálata tazimelteonhoz kapcsolódó mellékhatásra.
4. Tazimelteon alkalmazásra az előző igénypontok bármelyike szerinti, ahol az eljárás teljesen vak beteg nem 24 órás életritmusának krónikus kezelésére szolgál.
5. Tazimelteon alkalmazásra az előző igénypontok bármelyike szerinti, ahol a tazimelteon egy órával a megcélzott alvási idő előtt adagolt.
6. Tazimelteon alkalmazásra az előző igénypontok bármelyike szerinti, ahol a betegnek 24 órás alvás-ébrenlét ciklusra való visszatérése tartalmazza a beteg abnormális melatonin cirkadián ritmusának és/vagy abnormális kortizol cirkadián ritmusának 24 órás ritmusra való átállítását.



SZTNH-100078227