



HU000026537T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 026 537**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 07 723360**(22) A bejelentés napja: **2007. 03. 19.**(51) Int. Cl.: **C07D 405/12** (2006.01)**A01N 43/40** (2006.01)**A01P 7/00** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20070723360

(97) Az európai bejelentés közzétételi adatai:

EP 2004636 A1 **2007. 10. 18.**

(86) A nemzetközi (PCT) bejelentési szám:

PCT/EP 07/002392

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2004636 B1 **2015. 04. 22.**

(87) A nemzetközi közzétételi szám:

WO 07115646

(30) Elsőbbségi adatok:

102006015468**2006. 03. 31.****DE**

(73) Jogosult(ak):

Bayer Intellectual Property GmbH, 40789
Monheim am Rhein (DE)

(72) Feltaláló(k):

JESCHKE, Peter, 51467 Bergisch Gladbach (DE)**VELTEN, Robert, 40764 Langenfeld (DE)****SCHENKE, Thomas, 51469 Bergisch Gladbach (DE)****SCHALLNER, Otto, 40789 Monheim (DE)****BECK, Michael, Edmund, 40789 Monheim (DE)****MALSAM, Olga, 51503 Rösrath (DE)****NAUEN, Ralf, 40764 Langenfeld (DE)****MÜLLER, Thomas, 63073 Offenbach am Main (DE)****ARNOLD, Christian, 40764 Langenfeld (DE)****SANWALD, Erich, 24159 Kiel (DE)****GÖRGENS, Ulrich, 40882 Ratingen (DE)**

(74) Képviselő:

dr. Fehérvári Flóra, DANUBIA Szabadalmi és
Jogi Iroda Kft., Budapest

(54)

Rovarirtóként használt szubsztituált enaminokarbonil vegyületek

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))


A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

SUBSTITUTED ENAMINOCARBONYL COMPOUNDS USED AS INSECTICIDES

The present application relates to novel substituted enaminocarbonyl compounds, to processes for their preparation and to their use for controlling animal pests, especially arthropods, in particular insects.

Substituted enaminocarbonyl compounds are already known as insecticidally active compounds (cf. EP 0 539 588 A1, DE 102004047922 A1).

This invention now provides novel compounds of the formula (I)




in which

A represents one of the radicals pyrimidinyl, pyrazolyl or 1,2,4-triazolyl which are substituted by fluorine, chlorine, bromine, cyano, nitro, C₁-C₄-alkyl (which is optionally substituted by fluorine and/or chlorine), C₁-C₃-alkylthio (which is optionally substituted by fluorine and/or chlorine), or C₁-C₃-alkylsulphonyl (which is optionally substituted by fluorine and/or chlorine),

or

A represents a radical



in which

X represents halogen, alkyl or haloalkyl,

Y represents halogen, alkyl, haloalkyl, haloalkoxy, azido or cyano,

B represents oxygen, sulphur or methylene,


R^1 represents haloalkyl, haloalkenyl, halocycloalkyl or halocycloalkylalkyl,

R^2 represents hydrogen or halogen and

R^3 represents hydrogen or alkyl.

5 Furthermore, it has been found that the novel compounds of the formula (I) are obtained when

a) compounds of the formula (II)



in which

10 B, R^2 and R^3 are as defined above

are reacted with compounds of the formula (III)




in which

A and R^1 are as defined above,

15 if appropriate in the presence of a suitable diluent and if appropriate in the presence of an acidic auxiliary (process 1), or when


b) compounds of the formula (I)



in which

A, B, R^2 and R^3 are as defined above

are reacted with compounds of the formula (IV)




5 in which

E represents a suitable leaving group such as, for example, halogen (in particular bromine, chlorine, iodine) or O-sulphonylalkyl and O-sulphonylaryl (in particular O-mesyl, O-tosyl),

10 if appropriate in the presence of a suitable diluent and if appropriate in the presence of an acid acceptor (process 2), or when


c) compounds of the formula (II)



in which

B, R^2 and R^3 are as defined above,

15 are, in a first reaction step, reacted with compounds of the formula (V)



in which

R^1 is as defined above,

20 if appropriate in the presence of a suitable diluent and if appropriate in the presence of an acidic auxiliary, and

the resulting compounds of the formula (VI)



B, R^1, R^2 and R^3 are as defined above

are then, in a second reaction step, reacted with compounds of the formula (VII)

B-CH₃-A (VII)

E and A are as defined above.

if appropriate in the presence of a suitable diluent and if appropriate in the presence of an acid acceptor (process 3).

10 Finally, it has been found that the novel compounds of the formula (I) have strongly pronounced biological properties and are suitable especially for controlling animal pests, in particular insects, arachnids and nematodes encountered in agriculture, in forests, in the protection of stored products and in the protection of materials, and also in the hygiene sector.

15 Depending, where appropriate, on the nature of the substituents, the compounds of the formula (I) may be present as geometrical and/or as optically active isomers or corresponding isomer mixtures of varying composition. The invention relates both to the pure isomers and the isomer mixtures.

The formula (I) provides a general definition of the compounds according to the
20 invention.

Preferred substituents or ranges of the radicals given in the formulae mentioned above and below are illustrated below.

A preferably represents pyrimidin-5-yl which is optionally substituted in the 2-position by halogen or C₁-C₄-alkyl, represents 1*H*-pyrazol-4-yl which is

optionally substituted in the 1-position by C₁-C₄-alkyl and in the 3-position by halogen, represents 1*H*-pyrazol-5-yl which is optionally substituted in the 2-position by halogen or C₁-C₄-alkyl, or represents 1-methyl-1,2,4-triazol-3-yl,

furthermore

- 5 A preferably represents one of the radicals 5,6-difluoropyrid-3-yl, 5-chloro-6-fluoropyrid-3-yl, 5-bromo-6-fluoropyrid-3-yl, 5-iodo-6-fluoropyrid-3-yl, 5-fluoro-6-chloropyrid-3-yl, 5,6-dichloropyrid-3-yl, 5-bromo-6-chloropyrid-3-yl, 5-iodo-6-chloropyrid-3-yl, 5-fluoro-6-bromopyrid-3-yl, 5-chloro-6-bromopyrid-3-yl, 5,6-dibromopyrid-3-yl, 5-iodo-6-bromopyrid-3-yl, 5-fluoro-6-iodopyrid-3-yl, 5-chloro-6-iodopyrid-3-yl, 5-bromo-6-iodopyrid-3-yl, 5,6-diiodopyrid-3-yl, 10 5-methyl-6-fluoropyrid-3-yl, 5-methyl-6-chloropyrid-3-yl, 5-methyl-6-bromopyrid-3-yl, 5-methyl-6-iodopyrid-3-yl, 5-difluoromethyl-6-fluoropyrid-3-yl, 5-difluoromethyl-6-chloropyrid-3-yl, 5-difluoromethyl-6-bromopyrid-3-yl or 5-difluoromethyl-6-iodopyrid-3-yl.

- 15 B preferably represents oxygen or methylene.

R¹ preferably represents in each case fluorine-substituted C₁-C₅-alkyl, C₂-C₅-alkenyl, C₃-C₅-cycloalkyl or C₃-C₅-cycloalkylalkyl.

R² preferably represents hydrogen or halogen.

R³ preferably represents hydrogen or C₁-C₃-alkyl.

- 20 A particularly preferably represents one of the radicals 2-methylpyrimidin-5-yl, 2-chloropyrimidin-5-yl, 1*H*-pyrazol-4-yl which is optionally substituted in the 1-position by methyl, ethyl, and in the 3-position by chlorine, 1*H*-pyrazol-5-yl, 2-methylpyrazol-5-yl, 1-methyl-1,2,4-triazol-3-yl,

furthermore

- 25 A particularly preferably represents one of the radicals 5-fluoro-6-chloropyrid-3-yl, 5,6-dichloropyrid-3-yl, 5-bromo-6-chloropyrid-3-yl, 5-fluoro-6-bromopyrid-

3-yl, 5-chloro-6-bromopyrid-3-yl, 5,6-dibromopyrid-3-yl, 5-methyl-6-chloropyrid-3-yl or 5-methyl-6-bromopyrid-3-yl.

B particularly preferably represents oxygen or methylene.

5 R¹ particularly preferably represents 2-fluoroethyl, 2,2-difluoroethyl or 2-fluorocyclopropyl.

R² particularly preferably represents hydrogen, fluorine, chlorine or bromine.

R³ particularly preferably represents hydrogen or methyl.

10 A very particularly preferably represents one of the radicals 2-methylpyrimidin-5-yl, 2-chloropyrimidin-5-yl, 5-fluoro-6-chloropyrid-3-yl, 5,6-dichloropyrid-3-yl or 5-fluoro-6-bromopyrid-3-yl.

B very particularly preferably represents oxygen.

R¹ very particularly preferably represents 2,2-difluoroethyl.


R² particularly preferably represents hydrogen.

R³ particularly preferably represents hydrogen.

15 In a special group of compounds of the formula (I), A represents 2-methylpyrimidin-5-yl.




In a further special group of compounds of the formula (I), A represents 2-chloropyrimidin-5-yl.




20


In a further special group of compounds of the formula (I), A represents 5-fluoro-6-chloropyrid-3-yl.




In a further special group of compounds of the formula (I), A represents 5,6-dichloropyrid-3-yl



- 5 In a further special group of compounds of the formula (I), A represents 5-fluoro-6-bromopyrid-3-yl




In a further special group of compounds of the formula (I), A represents 5-fluoro-6-iodopyrid-3-yl



10

In a further special group of compounds of the formula (I), A represents 5-chloro-6-iodopyrid-3-yl




15

Hereinbelow, a further group of preferred compounds of the formula (I) is defined in which

A represents a pyrimidin-5-yl radical which is substituted in the 2-position by halogen or halo-C₁-C₄-alkyl,

or

A represents a radical



in which

5 X represents halogen, C₁-C₄-alkyl, halo-C₁-C₄-alkyl,

Y represents halogen, halo-C₁-C₄-alkyl, halo-C₁-C₄-alkoxy, azido or cyano,

B represents oxygen, sulphur or methylene,

R¹ represents halo-C₁-C₃-alkyl, halo-C₂-C₃-alkenyl, halocyclopropyl (where halogen represents in particular fluorine or chlorine),

10 R² represents hydrogen or halogen and

R³ represents hydrogen or methyl.

A preferably represents 2-chloropyrimidin-5-yl or 2-trifluoromethylpyrimidin-5-yl,


furthermore

15 A preferably represents one of the radicals 5,6-difluoropyrid-3-yl, 5-chloro-6-fluoropyrid-3-yl, 5-bromo-6-fluoropyrid-3-yl, 5-iodo-6-fluoropyrid-3-yl, 5-fluoro-6-chloropyrid-3-yl, 5,6-dichloropyrid-3-yl, 5-bromo-6-chloropyrid-3-yl, 5-iodo-6-chloropyrid-3-yl, 5-fluoro-6-bromopyrid-3-yl, 5-chloro-6-bromopyrid-3-yl, 5,6-dibromopyrid-3-yl, 5-fluoro-6-iodopyrid-3-yl, 5-chloro-6-iodopyrid-3-yl, 5-bromo-6-iodopyrid-3-yl, 5-methyl-6-fluoropyrid-3-yl, 5-methyl-6-chloropyrid-3-yl, 5-methyl-6-bromopyrid-3-yl, 5-methyl-6-iodopyrid-3-yl, 5-difluoromethyl-6-fluoropyrid-3-yl, 5-difluoromethyl-6-chloropyrid-3-yl, 5-difluoromethyl-6-bromopyrid-3-yl, 5-difluoromethyl-6-iodopyrid-3-yl.

20

- B preferably represents oxygen or methylene.
- R¹ preferably represents in each case fluorine-substituted C₁-C₃-alkyl, C₂-C₃-alkenyl or cyclopropyl.
- R² preferably represents hydrogen or halogen (where halogen represents in particular fluorine or chlorine).
- R³ preferably represents hydrogen.
- A particularly preferably represents 2-chloropyrimidin-5-yl, 5-fluoro-6-chloropyrid-3-yl, 5,6-dichloropyrid-3-yl, 5-bromo-6-chloropyrid-3-yl, 5-fluoro-6-bromopyrid-3-yl, 5-chloro-6-bromopyrid-3-yl, 5,6-dibromopyrid-3-yl, 5-methyl-6-chloropyrid-3-yl, 5-chloro-6-iodopyrid-3-yl or 5-difluoromethyl-6-chloropyrid-3-yl.
- B particularly preferably represents oxygen.
- R¹ particularly preferably represents 2-fluoroethyl, 2,2-difluoroethyl or 2-fluorocyclopropyl.
- R² particularly preferably represents hydrogen.
- R³ particularly preferably represents hydrogen.
- A very particularly preferably represents 5-fluoro-6-chloropyrid-3-yl or 5-fluoro-6-bromopyrid-3-yl.
- B very particularly preferably represents oxygen.
- R¹ very particularly preferably represents 2,2-difluoroethyl.
- R² particularly preferably represents hydrogen.
- R³ particularly preferably represents hydrogen.


In a special group of compounds of the formula (I), R³ represents hydrogen, B represents oxygen and A represents 2-chloropyrimidin-5-yl.




In a further special group of compounds of the formula (I), R^3 represents hydrogen, B represents oxygen and A represents 5-fluoro-6-chloropyrid-3-yl,



5 In a further special group of compounds of the formula (I), R^3 represents hydrogen, B represents oxygen and A represents 5,6-dichloropyrid-3-yl




In a further special group of compounds of the formula (I), R^3 represents hydrogen, B represents oxygen and A represents 5-bromo-6-chloropyrid-3-yl




10


In a further special group of compounds of the formula (I), R^3 represents hydrogen, B represents oxygen and A represents 5-methyl-6-chloropyrid-3-yl




15 In a further special group of compounds of the formula (I), R^3 represents hydrogen, B represents oxygen and A represents 5-fluoro-6-bromopyrid-3-yl




In a further special group of compounds of the formula (I), R^3 represents hydrogen, B represents oxygen and A represents 5-chloro-6-bromopyrid-3-yl




5 In a further special group of compounds of the formula (I), R^3 represents hydrogen, B represents oxygen and A represents 5-chloro-6-iodopyrid-3-yl




In a special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents oxygen and A represents 2-chloropyrimidin-5-yl,



10 In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents oxygen and A represents 5-fluoro-6-chloropyrid-3-yl,




In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents oxygen and A represents 5,6-dichloropyrid-3-yl




15


In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents oxygen and A represents 5-bromo-6-chloropyrid-3-yl




In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents oxygen and A represents 5-methyl-6-chloropyridin-3-yl



5 In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents oxygen and A represents 5-fluoro-6-bromopyridin-3-yl




In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents oxygen and A represents 5-chloro-6-bromopyridin-3-yl




10


In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents oxygen and A represents 5-chloro-6-iodopyridin-3-yl




15 In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents methylene and A represents 2-chloropyrimidin-5-yl.




In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents methylene and A represents 5-fluoro-6-chloropyrid-3-yl,




In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents methylene and A represents 5,6-dichloropyrid-3-yl




In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents methylene and A represents 5-bromo-6-chloropyrid-3-yl



10 In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents methylene and A represents 5-methyl-6-chloropyrid-3-yl




In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents methylene and A represents 5-fluoro-6-bromopyrid-3-yl




15

In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents methylene and A represents 5-chloro-6-bromopyrid-3-yl



In a further special group of compounds of the formula (I), R^2 and R^3 represent hydrogen, B represents methylene and A represents 5-chloro-6-iodo-pyrid-3-yl



5 In a further special group of compounds of the formula (I), R^1 represents difluoromethyl, R^2 and R^3 represent hydrogen and B represents oxygen.

In a further special group of compounds of the formula (I), R^1 represents 2-fluoroethyl, R^2 and R^3 represent hydrogen and B represents oxygen.

10 In a further special group of compounds of the formula (I), R^1 represents 2,2-difluoroethyl, R^2 and R^3 represent hydrogen and B represents oxygen

In a further special group of compounds of the formula (I), R^1 represents difluoromethyl, R^2 and R^3 represent hydrogen and B represents methylene.

In a further special group of compounds of the formula (I), R^1 represents 2-fluoroethyl, R^2 and R^3 represent hydrogen and B represents methylene.

15 In a further special group of compounds of the formula (I), R^1 represents 2,2-difluoroethyl, R^2 and R^3 represent hydrogen and B represents methylene.

The general or preferred radical definitions or illustrations given above apply both to the end products and, correspondingly, to precursors and intermediates. These radical definitions can be combined with one another as desired, i.e. including combinations
20 between the respective preferred ranges.


Preference according to the invention is given to compounds of the formula (I) which contain a combination of the meanings listed above as being preferred (preferable).

Particular preference according to the invention is given to compounds of the formula (I) which contain a combination of the meanings listed above as being particularly preferred.

Very particular preference according to the invention is given to compounds of the formula (I) which contain a combination of the meanings listed above as being very particularly preferred.

If, in the process 1 according to the invention for preparing the novel compounds of the formula (I), the compound of the formula (II) is, for example, tetronic acid and the compound of the formula (III) is, for example, N-[(6-chloro-5-fluoropyridin-3-yl)methyl]-2,2-difluoroethane-1-amine, preparation process 1 can be represented by the reaction scheme I below:

Scheme I



The formula (II) provides a general definition of the compounds required as starting materials for carrying out the process 1 according to the invention.

In this formula (II), B, R² and R³ preferably represent those radicals which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as preferred substituents.

Some of the compounds of the formula (II) can be obtained commercially or by methods known from the literature (cf., for example, compounds of the general formula (II) in which B represents oxygen: tetronic acids (Said, A. Speciality Chemicals Magazine (1984), 4(4), 7-8; Rao, Y. S. Chem. Rev. (1976), 76, 625-694; Tejedor, D.; Garcia-Tellado, F. Org. Preparations and Procedures International (2004), 36, 35-59; Reviews); compounds of the general formula (II) in which B represents

sulphur: thiotetronic acids (Thomas, E. J. Special Publication - Royal Society of Chemistry (1988), 65 (Top. Med. Chem.), 284-307, Review), compounds of the general formula (II) in which B represents methylene: cyclopentane-1,3-dione (Schick, Hans; Eichhorn, Inge. Synthesis (1989), (7), 477-492, Review).

- 5 The formula (III) provides a general definition of the compounds furthermore to be used as starting materials for carrying out the process 1 according to the invention.


In formula (III), A and R¹ have the meanings already mentioned in connection with the description of the compounds of the formula (I) according to the invention.

- Some of the compounds of the formula (III) can be obtained commercially or by methods known from the literature (cf., for example, S. Patai „The Chemistry of Amino Group“, Interscience Publishers, New York, 1968; compounds of the general formula (III) in which R¹ represents hydrogen: primary amines, compounds of the general formula (III) in which R¹ represents haloalkyl, haloalkenyl or halocycloalkyl: secondary amines).

- 15 Some of the compounds of the formula (VII) are commercially available, some are known, or they can be obtained by known methods.

Routes for preparing compounds of the formula (VII) are illustrated in reaction scheme III.

Scheme III



E = Hal, for example chlorine, bromine, iodine; O-tosyl, O-mesyl.

A = as defined above, for example




Methyl-substituted compounds ($A-CH_3$), for example, can be converted by oxidation into corresponding carboxylic acids ($A-COOH$, for example 5-fluoro-6-bromonicotinic acid: F. L. Setliff, G. O. Rankin, J. Chem. Eng. Data (1972), 17, 515-516; 5-chloro-6-bromonicotinic acid and 5,6-dibromonicotinic acid: F. L. Setliff et al., J. Chem. Eng. Data (1981), 26, 332-333; 5-iodo-6-bromonicotinic acid: F. L. Setliff et al., J. Chem. Eng. Data (1978), 23, 96-97, 5-fluoro-6-iodonicotinic acid and 5-bromo-6-iodonicotinic acid: F. L. Setliff et al., J. Chem. Eng. Data (1973), 18, 449-450, 5-chloro-6-iodonicotinic acid: F. L. Setliff, J. E. Lane J. Chem. Eng. Data (1976), 21, 246-247) or carboxylic esters (for example methyl 5-methyl-6-fluoronicotinate: WO 9833772 A1, 1998; methyl 5-methyl-6-bromonicotinate: WO 9730032 A1, 1997).

The carboxylic acids ($A-COOH$) can then be converted by methods known from the literature into the corresponding hydroxymethyl compounds ($A-CH_2-OH$), which are then reacted by methods known from the literature to give activated hydroxymethyl compounds ($A-CH_2-E$, $E = O$ -tosyl, O -mesyl) or halomethyl compounds ($A-CH_2-E$, $E = Hal$). The latter can also be obtained from corresponding compounds which contain a methyl group ($A-CH_3$) using suitable halogenating agents known from the literature. The syntheses of the following compounds: 5-chloromethyl-2-methylpyrimidine (U. Eiermann et al., Chem. Ber. (1990), 123, 1885-9); 3-chloromethyl-5-bromo-6-chloropyridine, 3-bromo-5-iodo-6-chloropyridine (S. Kagabu et al., J. Pestic. Sci. (2005), 30, 409-413) may be mentioned as examples for this procedure.

Compounds of the formula (VII) in which A represents a 5,6-disubstituted pyrid-3-yl radical can also be obtained by methods known from the literature. Suitable starting materials known from the literature are, for example, the 6-halo-substituted 5-nitro- β -picolines (A-1), which can be modified by known literature procedures, as shown in reaction scheme IV.

Scheme IV



X, Y = halogen, for example fluorine, chlorine, bromine, iodine
 E = halogen, O-mesyl, O-tosyl

The reduction of the nitro group in 6-halo-substituted 5-nitro- β -picolines (A-1), for example, gives 6-halo-substituted 5-amino- β -picolines (A-2, for example 5-amino-6-chloro- β -picoline and 5-amino-6-bromo- β -picoline: Setliff, F. L. *Org. Preparations and Preparations Int.* (1971), 3, 217-222; Kagabu, S. et al. *J. Pestic. Sci.* (2005), 30, 409-413). Subsequent diazotisation and Sandmeyer reaction (C. F. H. Allen, J. R. Thirtle, *Org. Synth., Coll. Vol. III*, 1955, p. 136) allows the introduction of halogen substituents in the 5-position (A-3, for example 5-fluoro-6-chloro- β -picoline and 5-fluoro-6-bromo- β -picoline: Setliff, F. L. *Org. Preparations and Preparations Int.* (1971), 3, 217-222; 5-iodo-6-chloro- β -picoline: Kagabu, S. et al. *J. Pestic. Sci.* (2005), 30, 409-413; 5,6-dichloropicoline: Setliff, F. L.; Lane, J. E. *J. Chem. Engineering Data* (1976), 21, 246-247). The oxidation of the methyl group in the 5,6-disubstituted β -picolines (A-3) leads to the corresponding 5,6-disubstituted nicotinic acids (A-4, for example 5-fluoro-6-chloronicotinic acid and 5-fluoro-6-bromonicotinic acid: Setliff F. L., Rankin G. O. *J. Chem. Engineering Data* (1972), 17, 515-516; 5-bromo-6-fluoronicotinic acid, 5-bromo-6-chloronicotinic acid and 5-bromo-6-bromonicotinic acid: F. L. Setliff *J. Chem. Engineering Data* (1970), 15, 590-591; 5-chloro-6-bromonicotinic acid and 5-iodo-6-bromonicotinic acid: Setliff, F. L., Greene, J. S. *J. Chem. Engineering Data* (1978), 23, 96-97; also known is 5-chloro-6-trifluoromethylnicotinic acid: F. Cottet et al., *Synthesis* (2004), 10, 1619-1624), which can be converted in the presence of reducing agents into the corresponding hydroxymethylated pyridines (A-5) (for example 5-bromo-6-chloro-3-hydroxymethylpyridine: Kagabu, S. et al., *J. Pestic. Sci.* (2005), 30, 409-413).

By reduction of 6-chloro-5-nitronicotinic acid (A-4, X=Cl, Y=NO₂; Boyer, J. H.; Schoen, W., J. Am. Chem. Soc. (1956), 78, 423-425), it is possible to form 6-chloro-3-hydroxymethyl-5-nitro-pyridine (A-5, X=Cl, Y=NO₂; Kagabu, S. et al., J. Med. Chem. (2000), 43, 5003-5009), which is then reduced to 6-chloro-3-hydroxymethyl-5-aminopyridine (A-5, X=Cl, Y=NH₂; Kagabu, S. et al., J. Med. Chem. (2000), 43, 5003-5009) and, via diazotisation and reaction with hydroxylamine, converted into 6-chloro-3-hydroxymethyl-5-azidopyridine (A-5, X=Cl, Y=N₃; Kagabu, S. et al., J. Med. Chem. (2000), 43, 5003-5009). Subsequent halogenation with thionyl chloride gives 6-chloro-3-chloromethyl-5-azidopyridine (VII, X=Cl, Y=N₃, E=Cl; Kagabu, S. et al., J. Med. Chem. (2000), 43, 5003-5009).

Alternatively, halogenation of the methyl group in the 3-position of (A-3) gives compounds of the formula (VII) in which E represents halogen (for example: 3-bromomethyl-6-chloro-5-fluoro-pyridine, 3-bromomethyl-6-chloro-5-iodopyridine; Kagabu, S. et al. J. Pestic. Sci. (2005), 30, 409-413). When 6-halo-substituted 5-nitro-β-picolines (A-3; Y=NO₂) are used, there may be initial halogenation of the methyl group in the 3-position (for example 3-bromomethyl-6-chloro-5-nitro-pyridine; Kagabu, S. et al., J. Pestic. Sci. (2005), 30, 409-413). The nitro group may also optionally be reduced at a later stage in the reaction sequence.


Also known from the literature is the introduction of substituents in the 5-position (for example Y=N₃) of compounds of the formula (VII) in which E represents *N*-morpholino. This radical can subsequently very easily be replaced by halogen (E=Hal) (cf. S. Kagabu et al., J. Med. Chem. 2000, 43, 5003-5009; reaction conditions: ethyl chloroformate, tetrahydrofuran, 60°C).

In general, it is possible to replace halogen atoms in the vicinity of the pyridine nitrogen by other halogen atoms or halogenated groups such as, for example, trifluoromethyl (transhalogenation, for example: chlorine for bromine or iodine; bromine for iodine or fluorine; iodine for fluorine or trifluoromethyl). Thus, a further alternative synthesis route entails exchange of the halogen atom (for example X=Cl) in the 6-position of the pyrid-5-yl radical (for example in A-4 where X, Y=Cl; 5,6-dichloronicotinic acid: Setliff, F. L.; Lane, J. E. J. Chem. Engineering Data (1976), 21, 246-247) for another halogen atom, for example iodine or fluorine (for example: A-4

where $X=I$; 5-bromo-6-iodonicotinic acid and A-4 where $X=F$; 5-bromo-6-fluoronicotinic acid: Setliff, F. L.; Price, D. W. *J. Chem. Engineering Data* (1973), 18, 449-450). However, this transhalogenation may also optionally be carried out later in suitable compounds of the formula (I), as illustrated in reaction scheme X and by the working examples further below.

For preparing compounds of the formula (III), for example, compounds of the formula (VII) in which A and E have the meanings mentioned further above are reacted with compounds of the formula (V) in which R^1 has the meanings mentioned further above, optionally in the presence of diluents and optionally in the presence of the basic reaction auxiliaries mentioned in the description of preparation process 2 (cf. N-alkylation, Scheme V).

Scheme V



In certain cases, it is alternatively also possible to prepare compounds of the formula (III) from the corresponding aldehydes (A-CHO) and compounds of the formula (V) by reductive amination (cf. Houben-Weyl, *Methoden der Organischen Chemie* [Methods of Organic Chemistry], vol. XI/1, Georg Thieme Verlag Stuttgart, p. 602).

In general, it is advantageous to carry out the preparation process 1 according to the invention in the presence of diluents. Diluents are advantageously employed in such an amount that the reaction mixture remains readily stirrable during the entire process. Suitable diluents for carrying out the process 1 according to the invention are all inert organic solvents.

Examples which may be mentioned are: halogenated hydrocarbons, in particular chlorinated hydrocarbons, such as tetrachloroethylene, tetrachloroethane, dichloropropane, methylene chloride, dichlorobutane, chloroform, carbon tetrachloride, trichloroethane, trichloroethylene, pentachloroethane, difluorobenzene, 1,2-dichloroethane, chlorobenzene, bromobenzene, dichlorobenzene, chlorotoluene, trichlorobenzene; alcohols, such as methanol, ethanol, isopropanol, butanol; ethers, such as ethyl propyl ether, methyl tert-butyl ether, n-butyl ether, anisole, phenetole, cyclohexyl methyl ether, dimethyl ether, diethyl ether, dipropylether, diisopropyl ether, di-n-butyl ether, diisobutyl ether, diisoamyl ether, ethylene glycol dimethyl ether, tetrahydrofuran, dioxane, dichlorodiethyl ether and polyethers of ethylene oxide and/or propylene oxide; amines, such as trimethylamine, triethylamine, tripropylamine, tributylamine, N-methylmorpholine, pyridine and tetramethylenediamine; nitrated hydrocarbons, such as nitromethane, nitroethane, nitropropane, nitrobenzene, chloronitrobenzene, o-nitrotoluene; nitriles, such as acetonitrile, propionitrile, butyronitrile, isobutyronitrile, benzonitrile, m-chlorobenzonitrile and also compounds, such as tetrahydrothiophene dioxide and dimethyl sulphoxide, tetramethylene sulphoxide, dipropyl sulphoxide, benzylmethyl sulphoxide, diisobutyl sulphoxide, dibutyl sulphoxide, diisoamyl sulphoxide; sulphones, such as dimethyl sulphone, diethyl sulphone, dipropyl sulphone, dibutyl sulphone, diphenyl sulphone, dihexyl sulphone, methyl ethyl sulphone, ethyl propyl sulphone, ethyl isobutyl sulphone and pentamethylene sulphone; aliphatic, cycloaliphatic or aromatic hydrocarbons, such as pentane, hexane, heptane, octane, nonane and industrial hydrocarbons, for example white spirits with components having boiling points in the range of, for example, from 40°C to 250°C, cymene, petroleum fractions having a boiling point interval of from 70°C to 190°C, cyclohexane, methylcyclohexane, petroleum ether, ligroin, octane, benzene, toluene, chlorobenzene, bromobenzene, nitrobenzene, xylene; esters, such as methyl acetate, ethyl acetate, butyl acetate, isobutyl acetate, and also dimethyl carbonate, dibutyl carbonate, ethylene carbonate; amides, such as hexamethylenephosphoric triamide, formamide, N-methylformamide, N,N-dimethylformamide, N,N-dipropylformamide, N,N-dibutylformamide, N-methylpyrrolidine, N-methylcaprolactam, 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidine, octylpyrrolidone, octylcaprolactam, 1,3-dimethyl-2-imidazolidinedione, N-

formylpiperidine, *N,N'*-1,4-diformylpiperazine; ketones, such as acetone, acetophenone, methyl ethyl ketone, methyl butyl ketone.

It is also possible to use mixtures of the solvents and diluents mentioned for the process according to the invention.

- 5 However, preferred diluents for carrying out the process according to the invention are aromatic hydrocarbons, such as benzene, toluene, chlorobenzene, bromobenzene, nitrobenzene or xylene, in particular benzene and toluene.

The preparation of compounds of the formula (I) according to preparation process 1 is generally carried out by reacting compounds of the formula (II) with compounds of the
10 formula (III) in the presence of an acidic auxiliary in a diluent.

The reaction time is generally from 10 minutes to 48 hours.

The reaction is carried out at temperatures between -10°C and +200°C, preferably between +10°C and 180°C, particularly preferably between 20°C and 140°C. The reaction is preferably carried out under reaction conditions which allow water to be
15 separated off or to be removed, for example with the aid of a water separator or by adding suitable molecular sieves, which also allow water to be removed.

In principle, the reaction can be carried out under atmospheric pressure. The reaction is preferably carried out under atmospheric pressure or under pressures of up to 15 bar and optionally under an atmosphere of protective gas (nitrogen, helium or argon).

- 20 For carrying out the process 1 according to the invention, in general from 0.5 to 4.0 mol, preferably from 0.7 to 3.0 mol, particularly preferably from 1.0 to 2.0 mol of amino compound of the formula (III) are employed per mole of the compound of the formula (II).


Furthermore, for carrying out the process 1 according to the invention, in general
25 catalytic amounts of an acidic auxiliary are added.

Suitable acidic auxiliaries are, for example, p-toluenesulphonic acid or acetic acid.

After the reaction has gone to completion, the entire reaction mixture is concentrated. The products obtained after work-up can be purified in a customary manner by recrystallisation, distillation under reduced pressure or column chromatography (cf. also the Preparation Examples).

- 5 If, in the process 2 according to the invention for preparing the novel compounds of the formula (I), the compound of the formula (Ia) is, for example, 4-[[[6-bromo-5-chloropyridin-3-yl)methyl]amino]furan-2(5H)-one and the compound of the formula (IV) is 1-bromo-1,1-dichloroprop-1-ene, preparation process 2 can be represented by reaction scheme VI below:

10 Scheme VI



The formula (Ia) provides a general definition of the compounds required as starting materials for carrying out the process 2 according to the invention.

- 15 In this formula (Ia), A, B, R² and R³ preferably represent those radicals which have already been mentioned in connection with the description of the compounds of the general formula (I) according to the invention as preferred substituents.

The compounds of the formula (Ia) can be obtained by preparation process 1, described further above, for example by reacting compounds of the formula (II) with compounds of the formula (III) in which R¹ represents hydrogen.

- 20 The formula (IV) provides a general definition of the compounds further to be used as starting materials for carrying out the process 2 according to the invention.

In formula (IV), E and R¹ have the meanings already mentioned in connection with the description of the compounds of the formula (I) according to the invention.

Some of the compounds of the formula (IV) are commercially available, or they can be obtained by methods known from the literature (cf., for example, 1-bromo-1,1-dichloropropene: WO 8800183 A1 (1988); compounds of the formula (IV) in which E represents halogen, such as chlorine, bromine and iodine: Houben-Weyl, Methoden der Organischen Chemie, vol. V/3, Georg Thieme Verlag Stuttgart, p. 503 and vol. V/4 p. 13, 517; compounds of the formula (IV) in which E represents mesylate: Crossland, R. K., Servis, K. L. J. Org. Chem. (1970), 35, 3195; compounds of the formula (IV) in which E represents tosylate: Roos, A. T. et al., Org. Synth., Coll. Vol. I, (1941), 145; Marvel, C. S., Sekera, V. C. Org. Synth., Coll. Vol. III, (1955), 366.

10 In general, it is advantageous to carry out the preparation process 2 according to the invention in the presence of a diluent and optionally in the presence of a basic reaction auxiliary.

Diluents are advantageously employed in such an amount that the reaction mixture remains readily stirrable during the entire process. Suitable diluents for carrying out the process 2 according to the invention are all inert organic solvents.

Preferred diluents for carrying out the process 2 according to the invention are ethers, such as methyl *tert*-butyl ether, *n*-butyl ether, anisole, phenetole, cyclohexyl methyl ether, diisopropyl ether, diisobutyl ether, diisoamyl ether, ethylene glycol dimethyl ether, tetrahydrofuran, dioxane, dichlorodiethyl ether and polyethers of ethylene oxide and/or propylene oxid; amides, such as hexamethylenephosphoric triamide, formamide, *N*-methylformamide, *N,N*-dimethylformamide, *N,N*-dipropylformamide, *N,N*-dibutylformamide; aromatic hydrocarbons, such as *N*-methylbenzene, toluene, chlorobenzene, bromobenzene, nitrobenzene, xylene; ketones, such as acetone, acetophenone, methyl ethyl ketone or methyl butyl ketone.

25 It is also possible to use mixtures of the solvents and diluents mentioned for the process 2 according to the invention.

However, preferred diluents for carrying out the process 2 according to the invention are ethers, such as methyl *tert*-butyl ether or cyclic ethers, such as tetrahydrofuran and dioxane, amides, such as *N,N*-dimethylformamide, aromatic hydrocarbons, such as

benzene or toluene; ketones, such as acetone, methyl ethyl ketone or methyl butyl ketone.

Suitable for use as basic reaction auxiliaries for carrying out the process 2 according to the invention are all suitable acid binders, such as amines, in particular tertiary amines,
5 and also alkali metal and alkaline earth metal compounds.

Examples which may be mentioned are the hydroxides, hydrides, oxides and carbonates of lithium, sodium, potassium, magnesium, calcium and barium, furthermore other basic compounds, such as amidine bases or guanidine bases, such as 7-methyl-1,5,7-triazabicyclo[4.4.0]dec-5-ene (MTBD); diazabicyclo[4.3.0]nonene
10 (DBN), diazabicyclo[2.2.2]octane (DABCO), 1,8-diazabicyclo[5.4.0]undecene (DBU), cyclohexyltetraethylguanidine (CyTEG), cyclohexyltetramethylguanidine (CyTMG), *N,N,N,N*-tetramethyl-1,8-naphthalenediamine, pentamethylpiperidine, tertiary amines, such as triethylamine, trimethylamine, tribenzylamine, triisopropylamine, tributylamine, tricyclohexylamine, triamylamine, trihexylamine, *N,N*-dimethylaniline,
15 *N,N*-dimethyltoluidine, *N,N*-dimethyl-*p*-aminopyridine, *N*-methylpyrrolidine, *N*-methylpiperidine, *N*-methylimidazole, *N*-methylpyrazole, *N*-methylmorpholine, *N*-methylhexamethylenediamine, pyridine, 4-pyrrolidinopyridine, 4-dimethylaminopyridine, chinoline, α -picoline, β -picoline, isochinoline, pyrimidine, acridine, *N,N,N',N'*-tetramethylenediamine, *N,N,N',N'*-tetraethylenediamine,
20 quinoxaline, *N*-propyldiisopropylamine, *N*-ethyldiisopropylamine, *N,N'*-dimethylcyclohexylamine, 2,6-lutidine, 2,4-lutidine or triethyldiamine.

Preference is given to using hydrides of lithium or sodium.

The reaction time is generally from 10 minutes to 48 hours.


The reaction is carried out at temperatures between -10°C and +200°C, preferably
25 between +10°C and 180°C, particularly preferably between 60°C and 140°C. In principle, the reaction can be carried out under atmospheric pressure. The reaction is preferably carried out under atmospheric pressure or under pressures of up to 15 bar and optionally under an atmosphere of protective gas (nitrogen, helium or argon).

For carrying out the process 2 according to the invention, in general from 0.5 to 4.0 mol, preferably from 0.7 to 3.0 mol, particularly preferably from 1.0 to 2.0 mol of alkylating agent of the general formula (IV) are employed per mole of the compound of the formula (II).

- 5 After the reaction has gone to completion, the entire reaction mixture is concentrated. The products obtained after work-up can be purified in a customary manner by recrystallisation, distillation under reduced pressure or column chromatography (cf. also the Preparation Examples).

- 10 If, in the process 3 according to the invention for preparing the novel compounds of the formula (I), in a first reaction step, the compound of the formula (II) used is, for example, tetronic acid and the compound of the formula (V) is 2-fluoroethylamine hydrochloride, and, in a second reaction step, the resulting compound of the general formula (VI) is 4-[(2-fluoroethyl)amino]furan-2(5H)-one, which is N-alkylated with compounds of the formula (VII), for example 3-bromomethyl-5,6-dichloropyridine, the preparation process 3 can be represented by reaction scheme VII below:

Scheme VII



- 20 The formula (II) provides a general definition of the compounds required as starting materials for carrying out the process 3 according to the invention and have already been described in more detail in connection with process 1, mentioned further above.

The formula (V) provides a general definition of the compounds further to be used as starting materials for carrying out the process 3 according to the invention.

In formula (V), R^1 has the meaning already mentioned in connection with the description of the compounds of the general formula (I) according to the invention.

The amino compounds of the formula (V) are defined in a general manner, and in many cases some of them are commercially available, or they can be obtained in a manner known per se by the Leuckart-Wallach reaction (for example 2-fluoroethylamine: US-Pat. 4030994 (1977); compounds of the general formula V in which R¹ represents alkyl, primary amines: cf., for example, Houben-Weyl, Methoden der Organischen Chemie, Vol. XI/1, 4. Ed. 1957, G. Thieme Verlag, Stuttgart, p. 648; M. L. Moore in „The Leuckart Reaction“ in: Organic Reactions, vol. 5, 2. Ed. 1952, New York, John Wiley & Sons, Inc. London).

In general, it is advantageous to carry out the preparation process 3 according to the invention in the presence of diluents. Diluents are advantageously employed in such an amount that the reaction mixture remains readily stirrable during the entire process. Suitable diluents for carrying out the process 3 according to the invention are all inert organic solvents.

Preferred diluents for carrying out the process 3 according to the invention are aromatic hydrocarbons, such as benzene, toluene, chlorobenzene, bromobenzene, nitrobenzene or xylene, in particular benzene and toluene.

In the second reaction step, the compounds of the formula (VI) are N-alkylated with compounds of the formula (VII).

In general, it is advantageous to carry out the second reaction step of the preparation process 3 according to the invention in the presence of diluents and in the presence of basic reaction auxiliaries.

Diluents are advantageously employed in such an amount that the reaction mixture remains readily stirrable during the entire process. Suitable diluents for carrying out the process 3 according to the invention are all inert organic solvents.

Suitable diluents for the second reaction step are, for example, ethers, such as tetrahydrofuran or dioxane.

Suitable basic reaction auxiliaries for the second reaction step are strong bases, such as, for example, sodium hydride.

The reaction time is generally from 10 minutes to 48 hours.

The reaction is carried out at temperatures between -10°C and +200°C, preferably between +10°C and 180°C, particularly preferably between 60°C and 140°C. The reaction is preferably carried out under reaction conditions which allow water to be separated off or to be removed, for example with the aid of a water separator.

After the reaction has gone to completion, the entire reaction mixture is concentrated. The products obtained after work-up can be purified in a customary manner by recrystallisation, distillation under reduced pressure or column chromatography (cf. also the Preparation Examples).

- 10 The compounds of the formula (I) can optionally be present in different polymorphic forms or as a mixture of different polymorphic forms. Both the pure polymorphs and the polymorph mixtures are provided by the invention and can be used according to the invention.

- 15 The active compounds according to the invention, in combination with good plant tolerance and favourable toxicity to warm-blooded animals and being tolerated well by the environment, are suitable for protecting plants and plant organs, for increasing the harvest yields, for improving the quality of the harvested material and for controlling animal pests, in particular insects, arachnids, helminths, nematodes and molluscs, which are encountered in agriculture, in horticulture, in animal husbandry, in forests, in gardens and leisure facilities, in the protection of stored products and of materials, and in the hygiene sector. They may be preferably employed as plant protection agents. They are active against normally sensitive and resistant species and against all or some stages of development. The abovementioned pests include:

- 25 From the order of the Anoplura (Phthiraptera), for example, *Damalinia* spp., *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Trichodectes* spp.

From the class of the Arachnida, for example, *Acarus siro*, *Aceria sheldoni*, *Aculops* spp., *Aculus* spp., *Amblyomma* spp., *Argas* spp., *Boophilus* spp., *Brevipalpus* spp., *Bryobia praetiosa*, *Choriopetes* spp., *Dermanyssus gallinae*, *Eotetranychus* spp., *Epitrimerus pyri*, *Eutetranychus* spp., *Eriophyes* spp., *Hemitarsonemus* spp.,

Hyalomma spp., Ixodes spp., Latrodectus mactans, Metatetranychus spp., Oligonychus spp., Ornithodoros spp., Panonychus spp., Phyllocoptruta oleivora, Polyphagotarsonemus latus, Psoroptes spp., Rhipicephalus spp., Rhizoglyphus spp., Sarcoptes spp., Scorpio maurus, Stenotarsonemus spp., Tarsonemus spp., Tetranychus
5 spp., Vasates lycopersici.

From the class of the Bivalva, for example, Dreissena spp.

From the order of the Chilopoda, for example, Geophilus spp., Scutigera spp.

From the order of the Coleoptera, for example, Acanthoscelides obtectus, Adoretus spp., Agelastica alni, Agriotes spp., Amphimallon solstitialis, Anobium punctatum,
10 Anoplophora spp., Anthonomus spp., Anthrenus spp., Apogonia spp., Atomaria spp., Attagenus spp., Bruchidius obtectus, Bruchus spp., Ceuthorrhynchus spp., Cleonus mendicus, Conoderus spp., Cosmopolites spp., Costelytra zealandica, Curculio spp., Cryptorhynchus lapathi, Dermestes spp., Diabrotica spp., Epilachna spp., Faustinus cubae, Gibbium psyllodes, Heteronychus arator, Hylamorpha elegans, Hylotrupes
15 hajulus, Hypera postica, Hypothenemus spp., Lachnosterna consanguinea, Leptinotarsa decemlineata, Lissorhoptrus oryzophilus, Lixus spp., Lyctus spp., Meligethes aeneus, Melolontha melolontha, Migdolus spp., Monochamus spp., Naupactus xanthographus, Niptus hololeucus, Oryctes rhinoceros, Oryzaephilus surinamensis, Otiorrhynchus sulcatus, Oxycetonia jucunda, Phaedon cochleariae, Phyllophaga spp., Popillia
20 japonica, Premnotrypes spp., Psylliodes chrysocephala, Ptinus spp., Rhizobius ventralis, Rhizopertha dominica, Sitophilus spp., Sphenophorus spp., Stereochus spp., Symphyletes spp., Tenebrio molitor, Tribolium spp., Trogoderma spp., Tychius spp., Xylotrechus spp., Zabrus spp.

From the order of the Collembola, for example, Onychiurus armatus.

25 From the order of the Dermaptera, for example, Forficula auricularia.

From the order of the Diplopoda, for example, Blaniulus guttulatus.

From the order of the Diptera, for example, Aedes spp., Anopheles spp., Bibio hortulanus, Calliphora erythrocephala, Ceratitis capitata, Chrysomya spp.,

Cochliomyia spp., Cordylobia anthropophaga, Culex spp., Cuterebra spp., Dacus oleae, Dermatobia hominis, Drosophila spp., Fannia spp., Gastrophilus spp., Hylemyia spp., Hyppobosca spp., Hypoderma spp., Liriomyza spp., Lucilia spp., Musca spp., Nezara spp., Oestrus spp., Oscinella frit, Pegomyia hyoscyami, Phorbia spp., Stomoxys spp.,

5 Tabanus spp., Tannia spp., Tipula paludosa, Wohlfahrtia spp.

From the class of the Gastropoda, for example, Arion spp., Biomphalaria spp., Bulinus spp., Deroceras spp., Galba spp., Lymnaea spp., Oncomelania spp., Succinea spp.

From the class of the helminths, for example, Ancylostoma duodenale, Ancylostoma ceylanicum, Ancylostoma braziliensis, Ancylostoma spp., Ascaris lumbricoides, Ascaris spp., Brugia malayi, Brugia timori, Bunostomum spp., Chabertia spp., Clonorchis spp., Cooperia spp., Dicrocoelium spp., Dictyocaulus filaria, Diphylobothrium latum, Dracunculus medinensis, Echinococcus granulosus, Echinococcus multilocularis, Enterobius vermicularis, Fasciola spp., Haemonchus spp., Heterakis spp., Hymenolepis nana, Hyostrongylus spp., Loa Loa, Nematodirus spp., Oesophagostomum spp.,
10 Opisthorchis spp., Onchocerca volvulus, Ostertagia spp., Paragonimus spp., Schistosoma spp., Strongyloides fuelleborni, Strongyloides stercoralis, Strongyloides spp., Taenia saginata, Taenia solium, Trichinella spiralis, Trichinella nativa, Trichinella britovi, Trichinella nelsoni, Trichinella pseudospiralis, Trichostrongylus spp., Trichuris trichuria, Wuchereria bancrofti.

20 It is furthermore possible to control protozoa, such as Eimeria.

From the order of the Heteroptera, for example, Anasa tristis, Antestiopsis spp., Blissus spp., Calocoris spp., Campylomma livida, Cavalerius spp., Cimex spp., Creontiades dilutus, Dasynus piperis, Dichelops furcatus, Diconocoris hewetti, Dysdercus spp., Euschistus spp., Eurygaster spp., Heliopeltis spp., Horcias nobilellus,
25 Leptocoris spp., Leptoglossus phyllopus, Lygus spp., Macrops excavatus, Miridae, Nezara spp., Oebalus spp., Pentomidae, Piesma quadrata, Piezodorus spp., Psallus seriatus, Pseudacysta persea, Rhodnius spp., Sahlbergella singularis, Scolinophora spp., Stephanitis nashi, Tibraca spp., Triatoma spp.

From the order of the Homoptera, for example, Acyrthosipon spp., Aeneolamia spp.,
30 Agonosceana spp., Aleurodes spp., Aleurolobus barodensis, Aleurothrixus spp.,

- Amrasca spp., Anuraphis cardui, Aonidiella spp., Aphanostigma piri, Aphis spp., Arboridia apicalis, Aspidiella spp., Aspidiotus spp., Atanus spp., Aulacorthum solani, Bemisia spp., Brachycaudus helichrysi, Brachycolus spp., Brevicoryne brassicae, Calligypona marginata, Carneiocephala fulgida, Ceratovacuna lanigera, Cercopidae, 5 Ceroplastes spp., Chaetosiphon fragaefolii, Chionaspis tegalensis, Chlorita onukii, Chromaphis juglandicola, Chrysomphalus ficus, Cicadulina mbila, Cocomytilus halli, Coccus spp., Cryptomyzus ribis, Dalbulus spp., Dialeurodes spp., Diaphorina spp., Diaspis spp., Doralis spp., Drosicha spp., Dysaphis spp., Dysmicoccus spp., Empoasca spp., Eriosoma spp., Erythroneura spp., Euscelis bilobatus, Geococcus coffeae, 10 Homalodisca coagulata, Hyalopterus arundinis, Icerya spp., Idiocerus spp., Idioscopus spp., Laodelphax striatellus, Lecanium spp., Lepidosaphes spp., Lipaphis erysimi, Macrosiphum spp., Mahanarva fimbriolata, Melanaphis sacchari, Metcalfiella spp., Metopolophium dirhodum, Monellia costalis, Monelliopsis pecanis, Myzus spp., Nasonovia ribisnigri, Nephrotettix spp., Nilaparvata lugens, Oncometopia spp., 15 Orthezia praelonga, Parabemisia myricae, Paratrioza spp., Parlatoria spp., Pemphigus spp., Peregrinus maidis, Phenacoccus spp., Phloeomyzus passerinii, Phorodon humuli, Phylloxera spp., Pinnaspis aspidistrae, Planococcus spp., Protopulvinaria pyriformis, Pseudaulacaspis pentagona, Pseudococcus spp., Psylla spp., Pteromalus spp., Pyrrilla spp., Quadraspidiotus spp., Quesada gigas, Rastrococcus spp., Rhopalosiphum spp., 20 Saissetia spp., Scaphoides titanus, Schizaphis graminum, Selenaspis articulatus, Sogata spp., Sogatella furcifera, Sogatodes spp., Stictiocephala festina, Tenalaphara malayensis, Tinocallis caryaefoliae, Tomaspis spp., Toxoptera spp., Trialeurodes vaporariorum, Trioza spp., Typhlocyba spp., Unaspis spp., Viteus vitifolii.

- From the order of the Hymenoptera, for example, Diprion spp., Hoplocampa spp., 25 Lasius spp., Monomorium pharaonis, Vespa spp.

From the order of the Isopoda, for example, Armadillidium vulgare, Oniscus asellus, Porcellio scaber.

From the order of the Isoptera, for example, Reticulitermes spp., Odontotermes spp.

- From the order of the Lepidoptera, for example, Acronicta major, Aedia leucomelas, 30 Agrotis spp., Alabama argillacea, Anticarsia spp., Barathra brassicae, Bucculatrix

- thurberiella, Bupalus piniarius, Cacoecia podana, Capua reticulana, Carpocapsa pomonella, Cheimatomia brumata, Chilo spp., Choristoneura fumiferana, Clysia ambiguella, Cnaphalocerus spp., Earias insulana, Ephestia kuehniella, Euproctis chrysorrhoea, Euxoa spp., Feltia spp., Galleria mellonella, Helicoverpa spp., Heliothis
- 5 spp., Hofmannophila pseudospretella, Homona magnanima, Hyponomeuta padella, Laphygma spp., Lithocolletis blancardella, Lithophane antennata, Loxagrotis albicosta, Lymantria spp., Malacosoma neustria, Mamestra brassicae, Mocis repanda, Mythimna separata, Oria spp., Oulema oryzae, Panolis flammea, Pectinophora gossypiella, Phyllocnistis citrella, Pieris spp., Plutella xylostella, Prodenia spp., Pseudaletia spp.,
- 10 Pseudoplusia includens, Pyrausta nubilalis, Spodoptera spp., Thermesia gemmatilis, Tinea pellionella, Tineola bisselliella, Tortrix viridana, Trichoplusia spp.

From the order of the Orthoptera, for example, Acheta domesticus, Blatta orientalis, Blattella germanica, Gryllotalpa spp., Leucophaea maderae, Locusta spp., Melanoplus spp., Periplaneta americana, Schistocerca gregaria.

- 15 From the order of the Siphonaptera, for example, Ceratophyllus spp., Xenopsylla cheopis.

From the order of the Symphyla, for example, Scutigera immaculata.

- From the order of the Thysanoptera, for example, Baliothrips biformis, Enneothrips flavens, Frankliniella spp., Heliothrips spp., Hercinothrips femoralis, Kakothrips spp.,
- 20 Rhipiphorothrips cruentatus, Scirtothrips spp., Taeniothrips cardamoni, Thrips spp.

From the order of the Thysanura, for example, Lepisma saccharina.

- The phytoparasitic nematodes include, for example, Anguina spp., Aphelenchoides spp., Belonoaimus spp., Bursaphelenchus spp., Ditylenchus dipsaci, Globodera spp., Helicocotylenchus spp., Heterodera spp., Longidorus spp., Meloidogyne spp.,
- 25 Pratylenchus spp., Radopholus similis, Rotylenchus spp., Trichodorus spp., Tylenchorhynchus spp., Tylenchulus spp., Tylenchulus semipenetrans, Xiphinema spp.

The compounds according to the invention can optionally, at certain concentrations or application rates, also be used as herbicides, safeners, growth regulators or agents to

improve plant properties, or as microbicides, for example as fungicides, antimycotics, bactericides, viricides (including agents against viroids) or as agents against MLO (mycoplasma-like organisms) and RLO (rickettsia-like organisms). They can optionally also be employed as intermediates or precursors for the synthesis of other active compounds.

The active compounds can be converted to the customary formulations, such as solutions, emulsions, wettable powders, water- and oil-based suspensions, powders, dusts, pastes, soluble powders, soluble granules, granules for broadcasting, suspension-emulsion concentrates, natural materials impregnated with active compound, synthetic materials impregnated with active compound, fertilizers and microencapsulations in polymeric substances.

These formulations are produced in a known manner, for example by mixing the active compounds with extenders, that is liquid solvents and/or solid carriers, optionally with the use of surfactants, that is emulsifiers and/or dispersants and/or foam-formers. The formulations are prepared either in suitable plants or else before or during the application.

Suitable for use as auxiliaries are substances which are suitable for imparting to the composition itself and/or to preparations derived therefrom (for example spray liquors, seed dressings) particular properties such as certain technical properties and/or also particular biological properties. Typical suitable auxiliaries are: extenders, solvents and carriers.

Suitable extenders are, for example, water, polar and nonpolar organic chemical liquids, for example from the classes of the aromatic and non-aromatic hydrocarbons (such as paraffins, alkylbenzenes, alkylnaphthalenes, chlorobenzenes), the alcohols and polyols (which, if appropriate, may also be substituted, etherified and/or esterified), the ketones (such as acetone, cyclohexanone), esters (including fats and oils) and (poly)ethers, the unsubstituted and substituted amines, amides, lactams (such as N-alkylpyrrolidones) and lactones, the sulphones and sulfoxides (such as dimethyl sulphoxide).

If the extender used is water, it is also possible to employ, for example, organic solvents

as auxiliary solvents. Essentially, suitable liquid solvents are: aromatics such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics and chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example petroleum fractions, mineral and vegetable oils, alcohols such as butanol or glycol and also their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethyl sulphoxide, and also water.

Suitable solid carriers are:

for example, ammonium salts and ground natural minerals such as kaolins, clays, talc, chalk, quartz, attapulgit, montmorillonite or diatomaceous earth, and ground synthetic minerals, such as finely divided silica, alumina and silicates; suitable solid carriers for granules are: for example, crushed and fractionated natural rocks such as calcite, marble, pumice, sepiolite and dolomite, and also synthetic granules of inorganic and organic meals, and granules of organic material such as paper, sawdust, coconut shells, maize cobs and tobacco stalks; suitable emulsifiers and/or foam-formers are: for example, nonionic and anionic emulsifiers, such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates and also protein hydrolysates; suitable dispersants are nonionic and/or ionic substances, for example from the classes of the alcohol-POE- and/or -POP-ethers, acid and/or POP-POE esters, alkyl aryl and/or POP-POE ethers, fat- and/or POP-POE adducts, POE- and/or POP-polyol derivatives, POE- and/or POP-sorbitan- or -sugar adducts, alkyl or aryl sulphates, alkyl- or arylsulphonates and alkyl or aryl phosphates or the corresponding PO-ether adducts. Furthermore, suitable oligo- or polymers, for example those derived from vinyllic monomers, from acrylic acid, from EO and/or PO alone or in combination with, for example, (poly)alcohols or (poly)amines. It is also possible to employ lignin and its sulphonic acid derivatives, unmodified and modified celluloses, aromatic and/or aliphatic sulphonic acids and their adducts with formaldehyde.

Tackifiers such as carboxymethylcellulose natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, as well as natural phospholipids such as cephalins and lecithins, and synthetic

phospholipids, can be used in the formulations.

It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyestuffs, such as alizarin dyestuffs, azo dyestuffs and metal phthalocyanine dyestuffs, and trace nutrients such as salts of iron,
5 manganese, boron, copper, cobalt, molybdenum and zinc.

Other possible additives are perfumes, mineral or vegetable, optionally modified oils, waxes and nutrients (including trace nutrients), such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

Stabilizers, such as low-temperature stabilizers, preservatives, antioxidants, light
10 stabilizers or other agents which improve chemical and/or physical stability may also be present.

The formulations generally comprise between 0.01 and 98% by weight of active compound, preferably between 0.5 and 90%.

The active compound according to the invention can be used in its commercially
15 available formulations and in the use forms, prepared from these formulations, as a mixture with other active compounds, such as insecticides, attractants, sterilizing agents, bactericides, acaricides, nematocides, fungicides, growth-regulating substances, herbicides, safeners, fertilizers or semiochemicals.

Particularly favourable mixing components are, for example, the following
20 compounds:

Fungicides:

Inhibitors of nucleic acid synthesis

benalaxyl, benalaxyl-M, bupirimate, chiralaxyl, clozylacon, dimethirimol,
ethirimol, furalaxyl, hymexazol, metalaxyl, metalaxyl-M, ofurace, oxadixyl,
25 oxolinic acid

Inhibitors of mitosis and cell division

benomyl, carbendazim, diethofencarb, fuberidazole, penicuron, thiabendazole, thiophanat-methyl, zoxamide

Inhibitors of respiratory chain complex I

diflumetorin

5 Inhibitors of respiratory chain complex II

boscalid, carboxin, fenfuram, flutolanil, furametpyr, mepronil, oxycarboxin, penthiopyrad, thifluzamide

Inhibitors of respiratory chain complex III

10 azoxystrobin, cyazofamid, dimoxystrobin, enestrobin, famoxadone, fenamidone, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, pyraclostrobin, picoxystrobin, trifloxystrobin

Decouplers

dinocap, fluazinam

Inhibitors of ATP production

15 fentin acetate, fentin chloride, fentin hydroxide, silthiofam

Inhibitors of amino acid biosynthesis and protein biosynthesis

andoprim, blasticidin-S, cyprodinil, kasugamycin, kasugamycin hydrochloride hydrate, mepanipyrim, pyrimethanil

Inhibitors of signal transduction

20 fenpiclonil, fludioxonil, quinoxifen

Inhibitors of lipid and membrane synthesis

chlozolate, iprodione, procymidone, vinclozolin

ampropylfos, potassium-ampropylfos, edifenphos, iprobenfos (IBP), isoprothiolane, pyrazophos

tolclofos-methyl, biphenyl

iodocarb, propamocarb, propamocarb hydrochloride

Inhibitors of ergosterol biosynthesis

fenhexamid,

- 5 azaconazole, bitertanol, bromuconazole, cyproconazole, diclobutrazole, difenoconazole, diniconazole, diniconazole-M, epoxiconazole, etaconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, furconazole, furconazole-cis, hexaconazole, imibenconazole, ipconazole, metconazole, myclobutanil, paclobutrazole, penconazole, propiconazole, prothioconazole, simeconazole, 10 tebuconazole, tetraconazole, triadimefon, triadimenol, triticonazole, uniconazole, voriconazole, imazalil, imazalil sulphate, oxpoconazole, fenarimol, flurprimidole, nuarimol, pyrifenoxy, triforine, pefurazoate, prochloraz, triflumizole, viniconazole, aldimorph, dodemorph, dodemorph acetate, fenpropimorph, tridemorph, fenpropidin, spiroxamine,

- 15 naftifine, pyributicarb, terbinafine

Inhibitors of cell wall synthesis

benthiavalicarb, bialaphos, dimethomorph, flumorph, iprovalicarb, polyoxins, polyoxorim, validamycin A

Inhibitors of melanin biosynthesis

- 20 capromamid, diclofymet, fenoxanil, phthalid, pyroquilon, tricyclazole

Resistance inducers

acibenzolar-S-methyl, probenazole, tiadinil

Multisite

- 25 captan, captan, chlorothalonil, copper salts such as: copper hydroxide, copper naphthenate, copper oxychloride, copper sulphate, copper oxide, oxine-copper and Bordeaux mixture, dichlofluanid, dithianon, dodine, dodine free base,

ferbam, folpet, fluorofolpet, guazatine, guazatine acetate, iminoctadine, iminoctadine albesilate, iminoctadine triacetate, mancooper, mancozeb, maneb, metiram, metiram zinc, propineb, sulphur and sulphur preparations containing calcium polysulphide, thiram, tolylfluanid, zineb, ziram

5 Unknown mechanism

- amibromdol, benthiazol, bethoxazin, capsimycin, carvone, chinomethionat, chloropicrin, cufraneb, cyflufenamid, cymoxanil, dazomet, debacarb, diclomezine, dichlorophen, dicloran, difenzoquat, difenzoquat methyl sulphate, diphenylamine, ethaboxam, ferimzone, flumetover, flusulphamide, fluopicolide, fluoroimide, hexachlorobenzene, 8-hydroxyquinoline sulphate, irumamycin, methasulphocarb, metrafenone, methyl isothiocyanate, mildiomycin, natamycin, nickel dimethyl dithiocarbamate, nitrothal-isopropyl, octhilinone, oxamocarb, oxyfenthiin, pentachlorophenol and salts, 2-phenylphenol and salts, piperalin, propanosine-sodium, proquinazid, pyrrol nitrin, quintozone, tecloftalam, tecnazene, triazoxide, trichlamide, zarilamid and 2,3,5,6-tetrachloro-4-(methylsulphonyl)pyridine, N-(4-chloro-2-nitrophenyl)-N-ethyl-4-methylbenzenesulphonamide, 2-amino-4-methyl-N-phenyl-5-thiazolecarboxamide, 2-chloro-N-(2,3-dihydro-1,1,3-trimethyl-1H-inden-4-yl)-3-pyridinecarboxamide, 3-[5-(4-chlorophenyl)-2,3-dimethylisoxazolidin-3-yl]pyridine, cis-1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol, 2,4-dihydro-5-methoxy-2-methyl-4-[[[1-[3-(trifluoromethyl)phenyl]ethylidene]amino]oxy]methyl]phenyl]-3H-1,2,3-triazol-3-one (185336-79-2), methyl 1-(2,3-dihydro-2,2-dimethyl-1H-inden-1-yl)-1H-imidazole-5-carboxylate, 3,4,5-trichloro-2,6-pyridinedicarbonitrile, methyl 2-[[[cyclopropyl[(4-methoxyphenyl)imino]methyl]thio]methyl]-.alpha.-(methoxymethylene)benzacetate, 4-chloro-alpha-propynyloxy-N-[2-[3-methoxy-4-(2-propynyloxy)phenyl]ethyl]benzacetamide, (2S)-N-[2-[4-[[3-(4-chlorophenyl)-2-propynyl]oxy]-3-methoxyphenyl]ethyl]-3-methyl-2-[[[methylsulphonyl]amino]butanamide, 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine, 5-chloro-6-(2,4,6-trifluorophenyl)-N-[(1R)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidin-7-amine, 5-chloro-N-[(1R)-1,2-dimethylpropyl]-6-(2,4,6-

trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidin-7-amine, N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2,4-dichloronicotinamide, N-(5-bromo-3-chloropyridin-2-yl)methyl-2,4-dichloronicotinamide, 2-butoxy-6-iodo-3-propylbenzopyranon-4-one, N-((Z)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl)-2-benzacetamide, N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-formylamino-2-hydroxybenzamide, 2-[[[1-[3(1-fluoro-2-phenylethyl)oxy]phenyl]ethylidene]amino]oxy]methyl]-alpha-(methoxyimino)-N-methyl-alphaE-benzacetamide, N-{2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethyl}-2-(trifluoromethyl)benzamide, N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, N-(6-methoxy-3-pyridinyl)cyclopropanecarboxamide, 1-[(4-methoxyphenoxy)methyl]-2,2-dimethylpropyl-1H-imidazole-1-carboxylic acid, O-[1-(4-methoxyphenoxy)methyl]-2,2-dimethylpropyl-1H-imidazole-1-carbothioic acid, 2-(2-[[6-(3-chloro-2-methylphenoxy)-5-fluoropyrimidin-4-yl]oxy]phenyl)-2-(methoxyimino)-N-methylacetamide

Bactericides:

bronopol, dichlorophen, nitrapyrin, nickel dimethyldithiocarbamate, kasugamycin, oethilinone, furancarboxylic acid, oxytetracycline, probenazole, streptomycin, tecloftalam, copper sulphate and other copper preparations.

20 Insecticides/acaricides/nematicides:

Acetylcholine esterase (AChE) inhibitors

carbamates,
for example alanycarb, aldicarb, aldoxycarb, allyxycarb, aminocarb, bendiocarb, benfurcarb, bufencarb, butacarb, butocarboxim, butoxycarboxim,
25 carbaryl, carbofuran, carbosulphan, cloethocarb, dimetilan, ethiofencarb, fenobucarb, fenothiocarb, formetanate, furathiocarb, isoprocarb, metam-sodium, methiocarb, methomyl, metolcarb, oxamyl, pirimicarb, promecarb, propoxur, thiodicarb, thiofanox, trimethacarb, XMC, xylylcab, triazamate
organophosphates,

for example acephate, azamethiphos, azinphos (-methyl, -ethyl), bromophos-ethyl, bromfeninfos (-methyl), butathiofos, cadusafos, carbophenothion, chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos (-methyl/-ethyl), coumaphos, cyanofenphos, cyanophos, chlorfenvinphos, demeton-S-methyl, demeton-S-methylsulphone, dialifos, diazinon, dichlofenthion, dichlorvos/DDVP, dicrotophos, dimethoate, dimethylvinphos, dioxabenzofos, disulphoton, EPN, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitrothion, fensulphothion, fenthion, flupyrazofos, fonofos, formothion, fosmethilan, fosphiazate, heptenophos, iodofenphos, iprobenfos, isazofos, isofenphos, isopropyl O-salicylate, isoxathion, malathion, mecarbam, methacrifos, methamidophos, methidathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, parathion (-methyl/-ethyl), phenthoate, phorate, phosalone, phosmet, phosphamidon, phosphocarb, phoxim, pirimiphos (-methyl/-ethyl), profenofos, propaphos, propetamphos, prothiofos, prothoate, pyraclofos, pyridaphenthion, pyridathion, quinalphos, sebufos, sulphotep, sulprofos, tebupirimfos, temephos, terbufos, tetrachlorvinphos, thiometon, triazophos, trichlorfon, vamidothion

Sodium channel modulators / voltage-dependent sodium channel blockers

pyrethroids,
for example acrinathrin, allethrin (d-cis-trans, d-trans), beta-cyfluthrin, bifenthrin, bioallethrin, bioallethrin-S-cyclopentyl isomer, bioethanomethrin, biopermethrin, bioresmethrin, chlovaporthrin, cis-cypermethrin, cis-resmethrin, cis-permethrin, cloocythrin, cycloprothrin, cyfluthrin, cyhalothrin, cypermethrin (alpha-, beta-, theta-, zeta-), cyphenothrin, deltamethrin, empenethrin (1R isomer), esfenvalerate, etofenprox, fenfluthrin, fenpropathrin, fenpyrithrin, fenvalerate, flubrocycythrinate, flucythrinate, flufenprox, flumethrin, fluvalinate, fubfenprox, gamma-cyhalothrin, imiprothrin, kadethrin, lambda-cyhalothrin, metofluthrin, permethrin (cis-, trans-), phenothrin (1R-trans-isomer), prallethrin, profluthrin, protrifenbutate, pyresmethrin, resmethrin, RU 15525, silafluofen, tau-fluvalinate, tefluthrin, terallethrin, tetramethrin (1R isomer), tralomethrin, transfluthrin, ZXI 8901, pyrethrins (pyrethrum)

DDT

oxadiazines,

for example indoxacarb

semicarbazones,

5 for example metaflumizone (BAS3201)

Acetylcholine receptor agonists/antagonists

chloronicotinyls,

for example acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram,
nithiazine, thiacloprid, imidaclothiz, AKD-1022, thiamethoxam

10 nicotine, bensultap, cartap

Acetylcholine receptor modulators

spinosyns,

for example spinosad, spinetoram (XDE-175)

GABA-controlled chloride channel antagonists

15 organochlorines,

for example camphechlor, chlordane, endosulphan, gamma-HCH, HCH,
heptachlor, lindane, methoxychlor

fiprols,

for example acetoprole, ethiprole, fipronil, pyrafluprole, pyriprole, vaniliprole

20 Chloride channel activators

mectins,

for example abamectin, emamectin, emamectin-benzoate, ivermectin,
lepimectin, milbemycin

Juvenile hormone mimetics,

25 for example diofenolan, epofenonane, fenoxycarb, hydroprene, kinoprene,

methoprene, pyriproxifen, triprene

Ecdysone agonists/disruptors

diacylhydrazines,

for example chromafenozide, halofenozide, methoxyfenozide, tebufenozide

5 Chitin biosynthesis inhibitors

benzoylureas,

for example bistrifluron, chlofluaazuron, disflubenzuron, fluazuron,
flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron,
noviflumuron, penfluron, teflubenzuron, triflumuron

10 buprofezin

cyromazine

Oxidative phosphorylation inhibitors, ATP disruptors

diafenthiuron

organotin compounds,

15 for example azocyclotin, cyhexatin, fenbutatin-oxide

Oxidative phosphorylation decouplers acting by interrupting the H⁺-proton gradient

pyrroles,

for example chlorfenapyr

dinitrophenols,

20 for example binapacyrl, dinobuton, dinocap, DNOC

Site-I electron transport inhibitors

METIs,

for example fenazaquin, fenpyroximate, pyrimidifen, pyridaben, tebufenpyrad,
tolfenpyrad

hydramethylnon

dicofol

Site-II electron transport inhibitors

rotenone

5 Site-III electron transport inhibitors

acequinocyl, fluacrypyrim

Microbial disruptors of the insect gut membrane

Bacillus thuringiensis strains

Lipid synthesis inhibitors

10 tetronic acids,

for example spiroadiclofen, spiromesifen

tetramic acids,

for example spirotetramat

carboxamides,

15 for example flonicamid

octopaminergic agonists,

for example amitraz

Inhibitors of magnesium-stimulated ATPase,

propargite

20 nereistoxin analogues,

for example thiocyclam hydrogen oxalate, thiosultap-sodium

Ryanodin receptor agonists

benzoic acid dicarboxamides,

for example flubendiamid

anthranilamides,

5 for example rynaxypyr (3-bromo-N-{4-chloro-2-methyl-6-[(methylamino)-carbonyl]phenyl}-1-(3-chloropyridin-2-yl)-1H-pyrazole-5-carboxamide)

Biologicals, hormones or pheromones

azadirachtin, *Bacillus* spec., *Beauveria* spec., codlemone, *Metarrhizium* spec., *Paeecilomyces* spec., thuringiensin, *Verticillium* spec.

10 Active compounds with unknown or unspecific mechanisms of action

fumigants,

for example aluminium phosphide, methyl bromide, sulphuryl fluoride

antifeedants,

for example cryolite, flonicamid, pymetrozine

15 mite growth inhibitors,

for example clofentezine, etoxazole, hexythiazox

amidoflumet, benclonthiaz, benzoximate, bifenazate, bromopropylate, buprofezin, chinomethionat, chlordimeform, chlorobenzilate, chloropierin, clothiazoben, cycloprene, cyflumetofen, dicyclanil, fenoxacrim, fentrifanil, 20 flubenzimine, flufenerim, flutenzin, gossypure, hydramethylnone, japonilure, metoxadiazone, petroleum, piperonyl butoxide, potassium oleate, pyridalyl, sulphuramid, tetradifon, tetrasul, triarathene, verbutin

A mixture with other known active compounds, such as herbicides, fertilizers, growth regulators, safeners, semiochemicals, or else with agents for improving the plant 25 properties, is also possible.

When used as insecticides, the active compounds according to the invention can furthermore be present in their commercially available formulations and in the use forms, prepared from these formulations, as a mixture with synergists. Synergists are compounds which increase the action of the active compounds, without it being
5 necessary for the synergistic agent added to be active itself.

When used as insecticides, the active compounds according to the invention can furthermore be present in their commercially available formulations and in the use forms, prepared from these formulations, as mixtures with inhibitors which reduce degradation of the active compound after use in the environment of the plant, on the
10 surface of parts of plants or in plant tissues.

The active compound content of the use forms prepared from the commercially available formulations can vary within wide limits. The active compound concentration of the use forms can be from 0.00000001 to 95% by weight of active compound, preferably between 0.00001 and 1% by weight.

15 The compounds are employed in a customary manner appropriate for the use forms.

All plants and plant parts can be treated in accordance with the invention. Plants are to be understood as meaning in the present context all plants and plant populations such as desired and undesired wild plants or crop plants (including naturally occurring crop plants). Crop plants can be plants which can be obtained by conventional plant
20 breeding and optimization methods or by biotechnological and genetic engineering methods or by combinations of these methods, including the transgenic plants and including the plant cultivars protectable or not protectable by plant breeders' rights. Plant parts are to be understood as meaning all parts and organs of plants above and below the ground, such as shoot, leaf, flower and root, examples which may be
25 mentioned being leaves, needles, stalks, stems, flowers, fruit bodies, fruits, seeds, roots, tubers and rhizomes. The plant parts also include harvested material, and vegetative and generative propagation material, for example cuttings, tubers, rhizomes, offshoots and seeds.

Treatment according to the invention of the plants and plant parts with the active
30 compound combinations is carried out directly or by allowing the compounds to act on

the surroundings, habitat or storage space by the customary treatment methods, for example by immersion, spraying, evaporation, fogging, scattering, painting on and, in the case of propagation material, in particular in the case of seeds, also by applying one or more coats.

5 The mixtures according to the invention are particularly suitable for treating seed. Here, the combinations according to the invention mentioned above as preferred or particularly preferred may be mentioned as being preferred. Thus, a large part of the damage to crop plants which is caused by pests occurs as early as when the seed is attacked during storage and after the seed is introduced into the soil, during and
10 immediately after germination of the plants. This phase is particularly critical since the roots and shoots of the growing plant are particularly sensitive and even minor damage can lead to the death of the whole plant. Protecting the seed and the germinating plant by the use of suitable compositions is therefore of particularly great interest.

The control of pests by treating the seeds of plants has been known for a long time and
15 is subject-matter of continuous improvements. However, the treatment of seed entails a series of problems which cannot always be solved in a satisfactory manner. Thus, it is desirable to develop methods for protecting the seed and the germinating plant which dispense with the additional application of crop protection agents after sowing or after the emergence of the plants. It is furthermore desirable to optimize the amount of
20 active compound employed in such a way as to provide maximum protection for the seed and the germinating plant from attack by pests, but without damaging the plant itself by the active compound employed. In particular, methods for the treatment of seed should also take into consideration the intrinsic insecticidal properties of transgenic plants in order to achieve optimum protection of the seed and the
25 germinating plant with a minimum of crop protection agents being employed.

The present invention therefore in particular also relates to a method for the protection of seed and germinating plants from attack by pests, by treating the seed with a composition according to the invention. The invention likewise relates to the use of the compositions according to the invention for the treatment of seed for protecting the
30 seed and the resulting plant from pests. Furthermore, the invention relates to seed

which has been treated with a composition according to the invention so as to afford protection from pests.

One of the advantages of the present invention is that the particular systemic properties of the compositions according to the invention mean that treatment of the seed with these compositions not only protects the seed itself, but also the resulting plants after emergence, from pests. In this manner, the immediate treatment of the crop at the time of sowing or shortly thereafter can be dispensed with.

A further advantage is the synergistically increased insecticidal activity of the compositions according to the invention in comparison with the insecticidal individual active compound, which exceeds the activity to be expected of the two active compounds when applied individually. Also advantageous is the synergistically increased fungicidal activity of the compositions according to the invention in comparison with the fungicidal individual active compound, which exceeds the activity to be expected of the active compound when applied individually. This makes possible an optimization of the amount of active compound employed.

Furthermore, it must be considered as advantageous that the mixtures according to the invention can also be employed in particular in transgenic seed, the plants arising from this seed being capable of expressing a protein directed against pests. By treating such seed with the compositions according to the invention, certain pests can be controlled merely by the expression of the, for example, insecticidal protein, and additionally be protected by the compositions according to the invention against damage.

The compositions according to the invention are suitable for protecting seed of any plant variety as already mentioned above which is employed in agriculture, in the greenhouse, in forests or in horticulture. In particular, this takes the form of seed of maize, peanut, canola, oilseed rape, poppy, soya beans, cotton, beet (for example sugar beet and fodder beet), rice, sorghum and millet, wheat, barley, oats, rye, sunflower, tobacco, potatoes or vegetables (for example tomatoes, cabbage plants). The compositions according to the invention are likewise suitable for treating the seed of fruit plants and vegetables as already mentioned above. The treatment of the seed of maize, soya beans, cotton, wheat and canola or oilseed rape is of particular importance.

As already mentioned above, the treatment of transgenic seed with a composition according to the invention is also of particular importance. This takes the form of seed of plants which, as a rule, comprise at least one heterologous gene which governs the expression of a polypeptide with in particular insecticidal properties. In this context, the heterologous genes in transgenic seed may be derived from microorganisms such as *Bacillus*, *Rhizobium*, *Pseudomonas*, *Serratia*, *Trichoderma*, *Clavibacter*, *Glomus* or *Gliocladium*. The present invention is particularly suitable for the treatment of transgenic seed which comprises at least one heterologous gene originating from *Bacillus* sp. and whose gene product shows activity against the European corn borer and/or the corn root worm. It is particularly preferably a heterologous gene derived from *Bacillus thuringiensis*.

In the context of the present invention, the composition according to the invention is applied to the seed either alone or in a suitable formulation. Preferably, the seed is treated in a state which is stable enough to avoid damage during treatment. In general, the seed may be treated at any point in time between harvest and sowing. The seed usually used has been separated from the plant and freed from cobs, shells, stalks, coats, hairs or the flesh of the fruits.

When treating the seed, care must generally be taken that the amount of the composition according to the invention applied to the seed and/or the amount of further additives is chosen in such a way that the germination of the seed is not adversely affected, or that the resulting plant is not damaged. This must be borne in mind in particular in the case of active compounds which may have phytotoxic effects at certain application rates.

As already mentioned above, it is possible to treat all plants and their parts according to the invention. In a preferred embodiment, wild plant species and plant cultivars, or those obtained by conventional biological breeding methods, such as crossing or protoplast fusion, and parts thereof, are treated. In a further preferred embodiment, transgenic plants and plant cultivars obtained by genetic engineering methods, optionally in combination with conventional methods (Genetically Modified Organisms), and parts thereof are treated. The terms "parts", "parts of plants" and "plant parts" have been explained above.

Particularly preferably, plants of the plant cultivars which are in each case commercially available or in use are treated according to the invention. Plant cultivars are to be understood as meaning plants having novel properties ("traits") which have been obtained by conventional breeding, by mutagenesis or by recombinant DNA techniques. These can be cultivars, bio- or genotypes.

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the substances and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, higher quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products are possible, which exceed the effects which were actually to be expected.

The transgenic plants or plant cultivars (obtained by genetic engineering) which are preferably to be treated according to the invention include all plants which, by virtue of the genetic modification, received genetic material which imparts particularly advantageous, useful traits to these plants. Examples of such traits are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, higher quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products. Further and particularly emphasized examples of such traits are a better defence of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or viruses, and also increased tolerance of the plants to certain herbicidally active compounds. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), maize, soya beans, potatoes, sugar beet, tomatoes, peas and other vegetable varieties, cotton, tobacco, oilseed rape and also fruit plants (with the fruits apples, pears, citrus

fruits and grapes), and particular emphasis is given to maize, soya beans, potatoes, cotton, tobacco and oilseed rape. Traits that are emphasized in particular are increased defence of the plants against insects, arachnids, nematodes and slugs and snails by virtue of toxins formed in the plants, in particular those formed in the plants by the genetic material from *Bacillus thuringiensis* (for example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF and also combinations thereof) (referred to hereinbelow as "Bt plants"). Traits that are also particularly emphasized are the increased defence of the plants against fungi, bacteria and viruses by systemic acquired resistance (SAR), systemin, phytoalexins, elicitors and resistance genes and correspondingly expressed proteins and toxins. Traits that are furthermore particularly emphasized are the increased tolerance of the plants to certain herbicidally active compounds, for example imidazolinones, sulphonylureas, glyphosate or phosphinotricin (for example the "PAT" gene). The genes which impart the desired traits in question can also be present in combination with one another in the transgenic plants. Examples of "Bt plants" which may be mentioned are maize varieties, cotton varieties, soya bean varieties and potato varieties which are sold under the trade names YIELD GARD® (for example maize, cotton, soya beans), KnockOut® (for example maize), StarLink® (for example maize), Bollgard® (cotton), Nucotr® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are maize varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosate, for example maize, cotton, soya beans), Liberty Link® (tolerance to phosphinotricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulphonylureas, for example maize). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned include the varieties sold under the name Clearfield® (for example maize). Of course, these statements also apply to plant cultivars having these genetic traits or genetic traits still to be developed, which plant cultivars will be developed and/or marketed in the future.

The plants listed can be treated according to the invention in a particularly advantageous manner with the compounds of the general formula I and/or the active compound mixtures according to the invention. The preferred ranges stated above for the active compounds or mixtures also apply to the treatment of these plants. Particular

emphasis is given to the treatment of plants with the compounds or mixtures specifically mentioned in the present text.

The active compounds according to the invention act not only against plant, hygiene and stored product pests, but also in the veterinary medicine sector against animal
5 parasites (ecto- and endoparasites), such as hard ticks, soft ticks, mange mites, leaf mites, flies (biting and licking), parasitic fly larvae, lice, hair lice, feather lice and fleas. These parasites include:

From the order of the Anoplurida, for example, *Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Phthirus* spp., *Solenopotes* spp.

- 10 From the order of the Mallophagida and the suborders Amblycerina and Ischnocerina, for example, *Trimenopon* spp., *Menopon* spp., *Trinoton* spp., *Bovicola* spp., *Werneckiella* spp., *Lepikentron* spp., *Damalina* spp., *Trichodectes* spp., *Felicola* spp.

- From the order of the Diptera and the suborders Nematocerina and Brachyocerina, for example, *Aedes* spp., *Anopheles* spp., *Culex* spp., *Simulium* spp., *Eusimulium* spp.,
15 *Phlebotomus* spp., *Lutzomyia* spp., *Culicoides* spp., *Chrysops* spp., *Hybomitra* spp., *Atylotus* spp., *Tabanus* spp., *Haematopota* spp., *Philipomyia* spp., *Braula* spp., *Musca* spp., *Hydrotaea* spp., *Stomoxys* spp., *Haematobia* spp., *Morellia* spp., *Fannia* spp., *Glossina* spp., *Calliphora* spp., *Lucilia* spp., *Chrysomyia* spp., *Wohlfahrtia* spp., *Sarcophaga* spp., *Oestrus* spp., *Hypoderma* spp., *Gasterophilus* spp., *Hippobosca* spp.,
20 *Lipoptena* spp., *Melophagus* spp.

From the order of the Siphonaptera, for example, *Pulex* spp., *Ctenocephalides* spp., *Xenopsylla* spp., *Ceratophyllus* spp.

From the order of the Heteroptera, for example, *Cimex* spp., *Triatoma* spp., *Rhodnius* spp., *Panstrongylus* spp.

- 25 From the order of the Blattellidae, for example, *Blattella germanica*, *Periplaneta americana*, *Blattella germanica*, *Supella* spp.

From the subclass of the Acari (Acarina) and the orders of the Meta- and Mesostigmata, for example, *Argas* spp., *Ornithodoros* spp., *Otobius* spp., *Ixodes* spp.,

Amblyomma spp., Boophilus spp., Dermacentor spp., Haemophysalis spp., Hyalomma spp., Rhipicephalus spp., Dermanyssus spp., Raillietia spp., Pneumonyssus spp., Sternostoma spp., Varroa spp.

From the order of the Actinedida (Prostigmata) and Acaridida (Astigmata), for example, Acarapis spp., Cheyletiella spp., Ornithocheyletia spp., Myobia spp., Psorergates spp., Demodex spp., Trombicula spp., Listrophorus spp., Acarus spp., Tyrophagus spp., Caloglyphus spp., Hypodectes spp., Pterolichus spp., Psoroptes spp., Chorioptes spp., Otodectes spp., Sarcoptes spp., Notoedres spp., Knemidocoptes spp., Cytodites spp., Laminosioptes spp.

10 The active compounds of the formula (I) according to the invention are also suitable for controlling arthropods which infest agricultural productive livestock, such as, for example, cattle, sheep, goats, horses, pigs, donkeys, camels, buffalo, rabbits, chickens, turkeys, ducks, geese and bees, other pets, such as, for example, dogs, cats, caged birds and aquarium fish, and also so-called test animals, such as, for example, hamsters,
15 guinea pigs, rats and mice. By controlling these arthropods, cases of death and reduction in productivity (for meat, milk, wool, hides, eggs, honey etc.) should be diminished, so that more economic and easier animal husbandry is possible by use of the active compounds according to the invention.

The active compounds according to the invention are used in the veterinary sector and
20 in animal husbandry in a known manner by enteral administration in the form of, for example, tablets, capsules, potions, drenches, granules, pastes, boluses, the feed-through process and suppositories, by parenteral administration, such as, for example, by injections (intramuscular, subcutaneous, intravenous, intraperitoneal and the like), implants, by nasal application, by dermal use in the form, for example, of dipping or
25 bathing, spraying, pouring on and spotting on, washing and powdering, and also with the aid of moulded articles containing the active compound, such as collars, ear marks, tail marks, limb bands, halters, marking devices and the like.

When used for cattle, poultry, pets and the like, the active compounds of the formula (I) can be used as formulations (for example powders, emulsions, free-flowing
30 compositions), which comprise the active compounds in an amount of from 1 to 80%

by weight, directly or after 100 to 10 000-fold dilution, or they can be used as a chemical bath.

It has furthermore been found that the compounds according to the invention also have a strong insecticidal action against insects which destroy industrial materials.

- 5 The following insects may be mentioned as examples and as preferred - but without any limitation:

Beetles, such as *Hylotrupes bajulus*, *Chlorophorus pilosis*, *Anobium punctatum*, *Xestobium rufovillosum*, *Ptilinus pecticornis*, *Dendrobium pertinex*, *Emobius mollis*, *Priobium carpini*, *Lyctus brunneus*, *Lyctus africanus*, *Lyctus planicollis*, *Lyctus*
10 *linearis*, *Lyctus pubescens*, *Trogoxylon aequale*, *Minthes rugicollis*, *Xyleborus spec.* *Tryptodendron spec.* *Apate monachus*, *Bostrychus capucins*, *Heterobostrychus brunneus*, *Sinoxylon spec.* *Dinoderus minutus*;

Hymenopterons, such as *Sirex juveneus*, *Urocerus gigas*, *Urocerus gigas taignus*, *Urocerus augur*;

- 15 Termites, such as *Kaloterms flavicollis*, *Cryptoterms brevis*, *Heteroterms indicola*, *Reticuliterms flavipes*, *Reticuliterms santonensis*, *Reticuliterms lucifugus*, *Mastoterms darwiniensis*, *Zootermopsis nevadensis*, *Coptoterms formosanus*;

Bristletails, such as *Lepisma saccharina*.

- 20 Industrial materials in the present connection are to be understood as meaning non-living materials, such as, preferably, plastics, adhesives, sizes, papers and cardboards, leather, wood and processed wood products and coating compositions.

The ready-to-use compositions may optionally comprise further insecticides and optionally one or more fungicides.

- 25 With respect to possible additional additives, reference may be made to the insecticides and fungicides mentioned above.

The compounds according to the invention can likewise be employed for protecting objects which come into contact with saltwater or brackish water, in particular hulls,

screens, nets, buildings, moorings and signalling systems, against fouling.

Furthermore, the compounds according to the invention, alone or in combinations with other active compounds, may be employed as antifouling agents.

In domestic, hygiene and stored-product protection, the active compounds are also
5 suitable for controlling animal pests, in particular insects, arachnids and mites, which
are found in enclosed spaces such as, for example, dwellings, factory halls, offices,
vehicle cabins and the like. They can be employed alone or in combination with other
active compounds and auxiliaries in domestic insecticide products for controlling these
pests. They are active against sensitive and resistant species and against all
10 developmental stages. These pests include:

From the order of the Scorpionidea, for example, *Buthus occitanus*.

From the order of the Acarina, for example, *Argas persicus*, *Argas reflexus*, *Bryobia*
ssp., *Dermanyssus gallinae*, *Glyciphagus domesticus*, *Ornithodoros moubat*,
Rhipicephalus sanguineus, *Trombicula alfreddugesi*, *Neutrombicula autumnalis*,
15 *Dermatophagoides pteronissimus*, *Dermatophagoides forinae*.

From the order of the Araneae, for example, *Aviculariidae*, *Araneidae*.

From the order of the Opiliones, for example, *Pseudoscorpiones chelifer*,
Pseudoscorpiones cheiridium, *Opiliones phalangium*.

From the order of the Isopoda, for example, *Oniscus asellus*, *Porcellio scaber*.

20 From the order of the Diplopoda, for example, *Blaniulus guttulatus*, *Polydesmus spp.*

From the order of the Chilopoda, for example, *Geophilus spp.*

From the order of the Zygentoma, for example, *Ctenolepisma spp.*, *Lepisma*
saccharina, *Lepismodes inquilinus*.

From the order of the Blattaria, for example, *Blatta orientalis*, *Blattella germanica*,
25 *Blattella asahinai*, *Leucophaea maderae*, *Panchlora spp.*, *Parcoblatta spp.*, *Periplaneta*
australasiae, *Periplaneta americana*, *Periplaneta brunnea*, *Periplaneta fuliginosa*,

Supella longipalpa.

From the order of the Saltatoria, for example, *Acheta domesticus*.

From the order of the Dermaptera, for example, *Forficula auricularia*.

From the order of the Isoptera, for example, *Kaloterms* spp., *Reticulitermes* spp.

- 5 From the order of the Psocoptera, for example, *Lepinatus* spp., *Liposcelis* spp.

From the order of the Coleoptera, for example, *Anthrenus* spp., *Attagenus* spp., *Dermestes* spp., *Latheticus oryzae*, *Necrobia* spp., *Ptinus* spp., *Rhizopertha dominica*, *Sitophilus granarius*, *Sitophilus oryzae*, *Sitophilus zeamais*, *Stegobium paniceum*.

- 10 From the order of the Diptera, for example, *Aedes aegypti*, *Aedes albopictus*, *Aedes taeniorhynchus*, *Anopheles* spp., *Calliphora erythrocephala*, *Chrysosona pluvialis*, *Culex quinquefasciatus*, *Culex pipiens*, *Culex tarsalis*, *Drosophila* spp., *Fannia canicularis*, *Musca domestica*, *Phlebotomus* spp., *Sarcophaga carnaria*, *Simulium* spp., *Stomoxys calcitrans*, *Tipula paludosa*.

- 15 From the order of the Lepidoptera, for example, *Achroia grisella*, *Galleria mellonella*, *Plodia interpunctella*, *Tinea cloacella*, *Tinea pellionella*, *Tineola bisselliella*.

From the order of the Siphonaptera, for example, *Ctenocephalides canis*, *Ctenocephalides felis*, *Pulex irritans*, *Tunga penetrans*, *Xenopsylla cheopis*.

- 20 From the order of the Hymenoptera, for example, *Camponotus herculeanus*, *Lasius fuliginosus*, *Lasius niger*, *Lasius umbratus*, *Monomorium pharaonis*, *Paravespula* spp., *Tetramorium caespitum*.

From the order of the Anoplura, for example, *Pediculus humanus capitis*, *Pediculus humanus corporis*, *Pemphigus* spp., *Phylloera vastatrix*, *Phthirus pubis*.

From the order of the Heteroptera, for example, *Cimex hemipterus*, *Cimex lectularius*, *Rhodinus prolixus*, *Triatoma infestans*.

- 25 In the field of domestic insecticides, they are used alone or in combination with other suitable active compounds, such as phosphoric esters, carbamates, pyrethroids,


neonicotinoids, growth regulators or active compounds from other known classes of insecticides.

They are used in aerosols, pressure-free spray products, for example pump and atomizer sprays, automatic fogging systems, foggers, foams, gels, evaporator products with evaporator tablets made of cellulose or polymer, liquid evaporators, gel and membrane evaporators, propeller-driven evaporators, energy-free, or passive, evaporation systems, moth papers, moth bags and moth gels, as granules or dusts, in baits for scattering or in bait stations.

Preparation examples:

10 Process 1

4-[[[6-Chloro-5-fluoropyridin-3-yl)methyl](2,2-difluoroethyl)amino]furan-2(5H)-one




Example (1)

317 mg (1.41 mmol) of N-[(6-chloro-5-fluoropyridin-3-yl)methyl]-2,2-difluoroethylamine (III-1) are added to 173 mg (1.69 mmol) of tetronic acid in 400 μ l of acetic acid, and the mixture is stirred at room temperature for 6 hours. The reaction mixture is concentrated under reduced pressure, and the residue is then partitioned between dichloromethane and water. The aqueous phase is extracted two more times with dichloromethane. The combined organic phases are washed with 1 N aqueous sodium hydroxide solution and concentrated under reduced pressure. Purification of the residue by column chromatography on silica gel (silica gel 60 - Merck, particle size: 0.04 to 0.063 mm) using the mobile phase mixture ethyl acetate : cyclohexane (3:1) gives 316 mg (64 % of theory) of 4-[[[6-chloro-5-fluoropyridin-3-yl)methyl](2,2-difluoroethyl)amino]furan-2(5H)-one.

¹H-NMR (CD₃CN): δ [ppm] = 3.60 (td, 2 H), 4.54 (s, 2 H), 4.75 (s, 1 H), 4.80 (s, 2 H), 6.04 (tt, 1 H), 7.54 (d, 1 H), 8.15 (s, 1 H).

Process 3

4-[[[(5,6-Dichloropyridin-3-yl)methyl]](2-fluoroethyl)amino]furan-2(5H)-one



Example (2)


217 mg (1.49 mmol) of 4-[(2-fluoroethyl)amino]furan-2(5H)-one (VI-1) and 90 mg (2.24 mmol) of a 60% strength dispersion of sodium hydride in mineral oil in 100 ml of tetrahydrofuran are heated under reflux for 2 hours. After cooling to room temperature, 360 mg (1.49 mmol) of 3-(bromomethyl)-5,6-dichloropyridine are added, and the mixture is heated under reflux for a further 3 hours. After cooling to room temperature and addition of methanol, the reaction mixture is concentrated under reduced pressure. The residue is taken up in ethyl acetate, washed successively twice with 1 N aqueous hydrochloric acid, twice with 1 N aqueous sodium hydroxide solution and saturated sodium chloride solution and dried over sodium sulphate. After concentration of the organic phase under reduced pressure and purification of the residue by column chromatography on silica gel (silica gel 60 - Merck, particle size: 0.04 to 0.063 mm) using the mobile phase mixture ethyl acetate : cyclohexane (5:1), 260 mg (56 % of theory) of 4-[[[(5,6-dichloropyridin-3-yl)methyl]](2-fluoroethyl)amino]furan-2(5H)-one are obtained.

¹H-NMR (CD₃CN): δ [ppm] = 3.50 (dt, 2 H), 4.49 (s, 2 H), 4.55 (dt, 2 H), 4.64 (s, 1 H), 4.80 (s, 2 H), 7.80 (s, 1 H), 8.22 (s, 1 H).


The compounds (3) to (7) of the formula (I) listed in Table 1 below were also prepared analogous to this procedure.

Table 1

Compounds of the formula (I)



in which A =



Ex. No.	B	R ¹	R ²	R ³	X	Y	Physical data: ¹ H-NMR, δ [ppm]
3	O		H	H	Cl	Cl	CD ₃ CN, δ = 3.59 (td, 2H), 4.52 (s, 2H), 4.75 (s, 1H), 4.80 (s, 2H), 6.04 (tt, 1H), 7.80 (s, 1H), 8.22 (s, 1H)
4	O		H	H	Cl	CH ₃	CD ₃ CN, δ = 2.35 (s, 3H), 3.57 (td, 2H), 4.48 (s, 2H), 4.75 (s, 1H), 4.80 (s, 2H), 6.02 (tt, 1H), 7.54 (s, 1H), 8.11 (s, 1H)
5	O		H	H	Cl	Br	CD ₃ CN, δ = 3.60 (td, 2H), 4.50 (s, 2H), 4.75 (s, 1H), 4.80 (s, 2H), 6.04 (tt, 1H), 7.95 (s, 1H), 8.25 (s, 1H)
6	O		H	H	Br	Cl	CD ₃ CN, δ = 3.60 (td, 2H), 4.50 (s, 2H), 4.75 (s, 1H), 4.80 (s, 2H), 6.04 (tt, 1H), 7.75 (s, 1H), 8.20 (s, 1H)
7	O		H	H	Br	F	CD ₃ CN, δ = 3.61 (td, 2H), 4.53 (s, 2H), 4.75 (s, 1H), 4.80 (s, 2H), 6.04 (tt, 1H), 7.47 (d, 1H), 8.14 (s, 1H)

5 Preparation of the starting materialsCompounds of the formula (HN(R¹)-CH₂-A) (III)III-1

N-[(6-Chloro-5-fluoropyridin-3-yl)methyl]-2,2-difluoroethylamine

At 45°C, 520 mg (2.89 mmol) of 6-chloro-3-chloromethyl-5-fluoropyridine (VII-3), 1.05 ml (14.44 mmol) of 2,2-difluoroethylamine and 400 μ L (2.89 mmol) of triethylamine are stirred in 50 ml of acetonitrile for about 48 hours. After about 16 and 32 hours, in each case another 0.42 ml (5.78 mmol) of 2,2-difluoroethylamine are added. After concentration of the reaction mixture under reduced pressure, the residue is taken up in 1 N aqueous hydrochloric acid and the mixture is washed with ethyl acetate. The aqueous phase is made alkaline using 2.5 N aqueous sodium hydroxide solution and extracted repeatedly with ethyl acetate. The combined organic phase is dried over sodium sulphate and concentrated under reduced pressure, giving 370 mg (57 % of theory) of N-[(6-chloro-5-fluoropyridin-3-yl)methyl]-2,2-difluoroethylamine.

$^1\text{H-NMR}$ (CD_3CN): δ [ppm] = 2.95 (td, 2 H), 3.87 (s, 2 H), 5.87 (tt, 1 H), 7.62 (d, 1 H), 8.17 (s, 1 H).

Compounds of the formula (VI)

VI-1

4-[(2-Fluoroethyl)amino]furan-2(5H)-one

On a water separator, 550 mg (4.97 mmol) of 2-fluoroethylamine hydrochloride, 547 mg (5.47 mmol) of tetronic acid, 408 mg (4.97 mmol) of sodium acetate and 9 mg (0.05 mmol) of 4-toluenesulphonic acid in 50 ml of toluene are heated for 5 hours under reflux. The reaction mixture is concentrated under reduced pressure, the residue is taken up in ethyl acetate and the mixture is washed with 1 N aqueous sodium hydroxide solution. The aqueous phase is extracted repeatedly with ethyl acetate and the combined organic phase is dried over sodium sulphate. The organic phase is concentrated under reduced pressure and the residue is purified by recrystallisation from ethyl acetate / cyclohexane, giving 300 mg (42 % of theory) of 4-[(2-fluoroethyl)amino]furan-2(5H)-one.

$^1\text{H-NMR}$ (CD_3CN , δ , ppm) = 3.40 (dq, 2 H), 4.52 (dt, 2H), 4.61 (s, 1 H), 4.62 (s, 2 H), 5.80 (br. s, 1 H).

Compound (VI-2) was also prepared analogously to this procedure:

VI-2**4-[(2,2-Difluoroethyl)amino]furan-2(5H)-one**

¹H-NMR (CD₃CN, δ, ppm) = 3.50 (tm, 2 H), 4.65 (s, 2 H), 4.71 (s, 1 H), 5.78 (br. s, 1 H), 5.98 (tt, 1 H).

5 **Compounds of the formula (E-CH₂-A) (VII)**VII-1

(5,6-Dichloropyridin-3-yl)methanol (E = OH, A = 5,6-dichloropyrid-3-yl) (cf. R. Graf et al. J. Prakt. Chem. 1932, 134 177-87)

At 0°C, 859 ml (859 mmol) of a 1 M solution of borane-tetrahydrofuran complex in
 10 tetrahydrofuran are added dropwise to 110 g (573 mmol) of 5,6-dichloronicotinic acid
 in 250 ml of tetrahydrofuran. The mixture is warmed to room temperature and stirred
 at this temperature for 3 hours. After cooling to 0°C, the reaction mixture is made
 alkaline with saturated aqueous potassium carbonate solution, most of the
 tetrahydrofuran is removed using a rotary evaporator and the residue is extracted
 15 repeatedly with ethyl acetate. The combined organic phases are washed with water and
 saturated aqueous sodium chloride solution and dried over sodium sulphate.
 Concentration of the organic phase under reduced pressure and purification of the
 residue by column chromatography on silica gel (silica gel 60 - Merck, particle size:
 0.04 to 0.063 mm) using the mobile phase mixture ethyl acetate : cyclohexane (1:2)
 20 gives 62 g (61 % of theory) of (5,6-dichloropyridin-3-yl)methanol.

¹H-NMR (CD₃CN): δ [ppm] = 3.31 (t, 1 H), 4.60 (d, 2 H), 7.85 (s, 1 H), 8.26 (s, 1 H)

Compound (VII-5) from Table 3 was also prepared analogously to the procedure for
 the compound (VII-1).

VII-2

25 **3-Bromomethyl-5,6-dichloropyridine** (E = Br, A = 5,6-dichloropyrid-3-yl) (cf. WO
 2000046196 A1)

At 0°C, 16.40 g (62.52 mmol) of triphenylphosphine and 11.66 g (65.50 mmol) of *N*-bromosuccinimide are added to a solution of 10.60 g (59.55 mmol) of (5,6-dichloropyridin-3-yl)methanol (VII-1) in 100 ml of dichloromethane. After 2 hours, the reaction mixture is substantially concentrated and the residue is purified by column chromatography on silica gel (silica gel 60 - Merck, particle size: 0.04 to 0.063 mm) using the mobile phase mixture ethyl acetate : cyclohexane (1:5). This gives 12.4 g (86 % of theory) of 3-bromomethyl-5,6-dichloropyridine.

¹H-NMR (CD₃CN): δ [ppm] = 4.53 (s, 2 H), 7.97 (s, 1 H), 8.35 (s, 1 H)

The compounds (VII-6) to (VII-8) from Table 3 were also prepared analogously to the procedure for the compound (VII-2).

VII-3

6-Chloro-3-chloromethyl-5-fluoropyridine (E = Cl, A = 6-chloro-5-fluoropyrid-3-yl)

1.00 g (6.87 mmol) of 6-chloro-5-fluoro-3-methylpyridine (cf. F. L. Setliff, Organic Preparations and Procedures International 1971, 3, 217-222), 1.01 g (7.56 mmol) of *N*-chlorosuccinimide and 0.11 g (0.69 mmol) of 2,2'-azobis(2-methylpropanenitrile) in 100 ml of chlorobenzene are boiled under reflux for 2 days. After about 16 and 32 hours, in each case a further 1.01 g (7.56 mmol) of *N*-chlorosuccinimide and 0.11 g (0.69 mmol) of 2,2'-azobis(2-methylpropanenitrile) are added. The reaction mixture is washed with saturated aqueous sodium sulphite solution and sodium bicarbonate solution and then dried over sodium sulphate and concentrated under reduced pressure. Column chromatography of the residue on silica gel (silica gel 60 - Merck, particle size: 0.04 to 0.063 mm) using the mobile phase mixture ethyl acetate : cyclohexane (1:20) gives 0.65 g (53 % of theory) 6-chloro-3-chloromethyl-5-fluoropyridine.

¹H-NMR (CD₃CN): δ [ppm] = 4.68 (s, 2 H), 7.69 (d, 1 H), 8.27 (s, 1 H)

VII-4

6-Bromo-3-bromomethyl-5-fluoropyridine (E = Br, A = 6-bromo-5-fluoropyrid-3-yl)





10.00 g (52.63 mmol) of 6-bromo-5-fluoro-3-methylpyridin (cf. F. L. Setliff, Organic Preparations and Procedures International 1971, 3, 217-222), 12.18 g (68.42 mmol) of *N*-bromosuccinimide and 0.86 g (5.26 mmol) of 2,2'-azobis(2-methylpropanenitrile) in 1 l of chlorobenzene are boiled under reflux for 6 hours. The reaction mixture is washed with saturated aqueous sodium sulphite solution and sodium bicarbonate solution and then dried over sodium sulphate and concentrated under reduced pressure. Column chromatography of the residue on silica gel (silica gel 60 - Merck, particle size: 0.04 to 0.063 mm) using the mobile phase mixture ethyl acetate : cyclohexane (1:20) gives 6.0 g (42 % of theory) of 6-bromo-3-chloromethyl-5-fluoropyridine.

¹H-NMR (CD₃CN): δ [ppm] = 4.55 (s, 2 H), 7.65 (d, 1 H), 8.27 (s, 1 H)

Further compounds (VII-5) to (VII-8) of the formula (VII) are listed in Table 3 below.

Table 3

E-CH₂-A (VII)

Ex. No.	E	A	Physical Data ^{a)}
VII-5	OH		3.30 (t, 1H), 4.59 (d, 2H), 7.83 (s, 1H), 8.26 (s, 1H)
VII-6	Br		2.37 (s, 3H), 4.52 (s, 2H), 7.70 (s, 1H), 8.24 (s, 1H)
VII-7	Br		4.52 (s, 2H), 8.10 (s, 1H), 8.38 (s, 1H)
VII-8	Br		4.52 (d, 2H), 7.92 (s, 1H), 8.35 (s, 1H)

^{a)} ¹H-NMR (CD₃CN), δ [ppm]

Biological examplesExample No. 1**Myzas test (MYZUPE spray treatment)**

Solvent: 78 parts by weight of acetone
 5 1.5 parts by weight of dimethylformamide

Emulsifier: 0.5 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

- 10 Discs of Chinese cabbage leaves (*Brassica pekinensis*) which are infested by all stages of the green peach aphid (*Myzus persicae*) are sprayed with a preparation of active compound of the desired concentration.

After the desired period of time, the effect in % is determined. 100% means that all aphids have been killed; 0% means that none of the aphids have been killed.

- 15 In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in g/ha	Kill rate in % after 5 days
Example 1	500	100
Example 2	500	100
Example 3	500	100
Example 4	500	100
Example 5	500	100
Example 6	500	100
Example 7	500	100

Example No.2

Nilaparvata lugens test (NILALU hydroponic treatment)

Solvent: 78 parts by weight of acetone
1.5 parts by weight of dimethylformamide

Emulsifier: 0.5 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

The preparation of active compound is pipetted into water. The stated concentration refers to the amount of active compound per volume unit of water (mg/l = ppm). The
10 water is then infected with the brown planthopper (*Nilaparvata lugens*).

After the desired period of time, the effect in % is determined. 100% means that all planthoppers have been killed; 0% means that none of the planthoppers have been killed.

In this test, for example, the following compounds of the Preparation Examples show
15 good activity: see table

Example	Active compound concentration in g/ha	Kill rate in % after 7 days
Example 2	100	80
Example 3	100	100

Example No. 3**Bemisia tabaci (BEMITA spray treatment)**

Solvent: 78 parts by weight of acetone
1.5 parts by weight of dimethylformamide

20 Emulsifier: 0.5 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

5 Discs of cotton leaves (*Gossypium hirsutum*) which are infested by larvae of the whitefly (*Bemisia tabaci*) are sprayed with a preparation of active compound of the desired concentration.

After the desired period of time, the effect in % is determined. 100% means that all whiteflies have been killed; 0% means that none of the whiteflies have been killed.

10 In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in g/ha	Kill rate in % after 7 days
Example 1	500	100

Example No. 4

Meloidogyne test (MELGIN spray treatment)

Solvent: 80 parts by weight of acetone

15 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

Containers are filled with sand, active compound solution, *Meloidogyne incognita* egg/larvae suspension and lettuce seeds. The lettuce seeds germinate and the plants develop. On the roots, galls are formed.

20 After the desired period of time, the nematocidal activity is determined by the formation of galls in %. 100% means that no galls were found; 0% means that the number of galls on the treated plants corresponds to that of the untreated controls.

In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in ppm	Kill rate in % after 14 days
Example 2	20	100

Example No. 5

Myzus persicae test, hydroponic treatment (MYZUPE sys.)

- 5 Solvent: 7 parts by weight of dimethylformamide
- Emulsifier: 2 parts by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

- 10 The preparation of active compound is mixed with water. The stated concentration refers to the amount of active compound per volume unit of water (mg/l = ppm). The treated water is filled into containers housing a pea plant (*Pisum sativum*) which is then infected with the green peach aphid (*Myzus persicae*).

- After the desired period of time, the kill in % is determined. 100% means that all
- 15 aphids have been killed; 0% means that none of the aphids have been killed.

In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in ppm	Kill rate in % after 6 days
Example 2	20	100

Example No. 6

Aphis gossypii test (APHIGO)

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with emulsifier-containing water to the desired concentration.

Cotton leaves (*Gossypium hirsutum*) which are heavily infested by the cotton aphid (*Aphis gossypii*) are treated by being dipped into the preparation of active compound of the desired concentration.

10 After the desired period of time, the kill in % is determined. 100% means that all aphids have been killed; 0% means that none of the aphids have been killed.

In this test, for example, the following compounds of the Preparation Examples show good activity; see table

Example	Active compound concentration in ppm	Kill rate in % after 6 days
Example 2	100	95

Example No. 7

15 **Myzus persicae test; (MYZUPE G)**

Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

Cabbage plants (*Brassica oleracea*) which are heavily infested by the green peach aphid (*Myzus persicae*) are watered with a preparation of active compound of the desired concentration.

After the desired period of time, the kill in % is determined. 100% means that all aphids have been killed; 0% means that none of the aphids have been killed.

In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in ppm	Kill rate in % after 10 days
Example 2	4	98

Example No. 8

Aphis gossypii test; (APHIGO G)

10 Solvent: 7 parts by weight of dimethylformamide

Emulsifier: 2 parts by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

15 Cotton plants (*Gossypium hirsutum*) which are heavily infested by the cotton aphid (*Aphis gossypii*) are watered with a preparation of active compound of the desired concentration.

After the desired period of time, the kill in % is determined. 100% means that all aphids have been killed; 0% means that none of the aphids have been killed.

20 In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in ppm	Kill rate in % after 10 days
Example 2	4	100

Example No. 9**Myzus persicae test (soil application)**

Solvent: 4 parts by weight of acetone

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

The preparation of active compound is mixed with soil. The stated concentration refers to the amount of active compound per volume unit of soil (mg/l = ppm). The treated
10 soil is filled into pots and planted with a bell pepper plant (*Capsicum annuum*). After one week, the plant is infected with the green peach aphid (*Myzus persicae*).

After the desired period of time, the kill in % is determined. 100% means that all aphids have been killed; 0% means that none of the aphids have been killed.

- In this test, for example, the following compounds of the Preparation Examples show
15 good activity; see table

Example	Active compound concentration in ppm	Kill rate in % after 35 days
Example 4	0.25	90

Example No. 10**Ctenocephalides felis; oral (CTECFE)**

Solvent: dimethyl sulphoxide

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of water. Part of the concentrate is diluted with citrated cattle blood, and the desired concentration is prepared.

- 20 unfed adult fleas (*Ctenocephalides felis*) are placed into a chamber whose top and bottom ends are closed with gauze. A metal cylinder whose bottom end is closed with parafilm is placed onto the chamber. The cylinder contains the blood/active compound preparation, which can be taken up by the fleas through the parafilm membrane. The blood is warmed to 37°C, but the flea chamber is at room temperature.

- After the desired period of time, the kill in % is determined. 100% means that all fleas have been killed; 0% means that none of the fleas have been killed.

In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in ppm	Kill rate in % after 2 days
Example 1	100	80
Example 3	100	80
Example 4	100	80
Example 6	100	80

Example No. 11

Lucilia cuprina test (LUCICU)

- 15 Solvent: dimethyl sulphoxide

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of water, and the concentrate is diluted with water to the desired concentration.

- Containers containing horse meat treated with the preparation of active compound of the desired concentration are populated with *Lucilia cuprina* larvae.

After the desired period of time, the kill in % is determined. 100% means that all larvae have been killed; 0% means that none of the larvae have been killed.

In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in ppm	Kill rate in % after 2 days
Example 1	100	100
Example 7	100	100

5 Example No. 12

Boophilus microplus test (BOOPMI injection)

Solvent: dimethyl sulphoxide

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amount of solvent, and the concentrate is diluted
10 with solvent to the desired concentration.


The solution of active compound is injected into the abdomen (*Boophilus microplus*), and the animals are transferred into dishes and kept in a temperature-controlled room.

After the desired period of time, the effect in % is determined. 100% means that no tick has laid any fertile eggs.

15 In this test, for example, the following compounds of the Preparation Examples show good activity: see table

Example	Active compound concentration in µg/animal	Kill rate in % after 7 days
Example 2	20	100

1. (I) képletű vegyületek



(I)

ahol

A jelentése pirimidinil-, pirazolil- vagy 1,2,4-triazolilcsoport egyike, amely szubsztituálva van fluor-, klór-, brómatommal, ciano-, nitro-, C_1 - C_4 -alkilcsoporttal (amely adott esetben fluor- és/vagy klóratommal szubsztituált) C_1 - C_3 -alkiltiocsoporttal (amely adott esetben fluor- és/vagy klóratommal szubsztituált) vagy C_1 - C_3 -alkilszulfonil-csoporttal (amely adott esetben fluor- és/vagy klóratommal szubsztituált), vagy

A jelentése



csoport, ahol

X jelentése halogénatom, alkil- vagy halogénalkil-csoport,

Y jelentése halogénatom, alkil-, halogénalkil-, halogénalkoxi-, azido- vagy cianocsoport,

B jelentése oxigénatom, kénatom vagy metilénecsoport,

R^1 jelentése halogénalkil-, halogénalkenil-, halogéncikloalkil- vagy halogéncikloalkilalkil-csoport,


R^2 jelentése hidrogén- vagy halogénatom, és

R^3 jelentése hidrogénatom vagy alkilcsoport.

2. Az 1. igénypont szerinti (I) képletű vegyületek, ahol

A jelentése pirimidin-5-il-csoport, amely 2-helyzetben halogénatommal vagy halogén-(C_1 - C_4 alkil)-csoporttal van szubsztituálva, vagy

A jelentése



csoport, ahol

X jelentése halogénatom, C_1 - C_4 -alkil- vagy halogén-(C_1 - C_4 -alkil)-csoport,

Y jelentése halogénatom, C_1 - C_4 -alkil-, halogén-(C_1 - C_4 -alkil)-, halogén-(C_1 - C_4 -alkoxi)-, azido- vagy cianocsoport,

- B jelentése oxigén-, kénatom vagy metilénecsoport,
 R^1 jelentése halogén-(C_1 - C_3 -alkil)-, halogén-(C_2 - C_3 -alkenil)-, halogénciklopropil-csoport,
 R^2 jelentése hidrogén- vagy halogénatom, és
 R^3 jelentése hidrogénatom vagy alkilcsoport.


3. A 2. igénypontban meghatározott (I) képletű vegyületek, ahol R^1 meghatározásában a halogénatom kifejezés fluor- vagy klóratomot jelent.

4. Készítmény, azzal jellemezve, hogy az legalább egy, az 1-3. igénypontok bármelyike szerinti (I) képletű vegyületet és szokásos töltőanyagokat és/vagy felületaktív anyagokat tartalmaz.

5. Eljárás kártevők irtására, azzal jellemezve, hogy egy, az 1-3. igénypontok bármelyike szerinti (I) képletű vegyületet vagy egy, a 4. igénypont szerinti készítményt hagyunk hatni a kártevőkön és/vagy ezek tartózkodási helyén.


6. Az 1-3. igénypontok bármelyike szerinti (I) képletű vegyületek vagy a 4. igénypont szerinti készítmények alkalmazása kártevők irtására.

7. Eljárás az 1. igénypontban meghatározott (I) képletű vegyületek előállítására, amelynek során (II) képletű vegyületeket



ahol

B, R^2 és R^3 jelentése az 1. igénypontban megadott,
 (III) képletű vegyületekkel



ahol

A és R^1 jelentése az 1. igénypontban megadott,
 reagáltatunk adott esetben megfelelő hígítószer jelenlétében és adott esetben egy savas segédanyag jelenlétében.

8. A 7. igénypont szerinti eljárás, ahol a hígítószer benzol, toluol, klórbenzol, brómbenzol, nitrobenzol és xilol közül van kiválasztva.

9. A 7. vagy 8. igénypont szerinti eljárás, ahol a savas segédanyag p-toluolszulfonsav vagy ecetsav.

10. A 7. igénypont szerinti eljárás, ahol az alkalmazott (II) képletű vegyület tetronsav és az alkalmazott (III) képletű vegyület N-[(6-klór-5-fluorpiridin-3-il)metil]-2,2-difluor-étán-1-amin.

11. (III) képletű vegyület



ahol

A jelentése 5-fluor-6-klórpírid-3-il-, 5,6-diklórpírid-3-il-, 5-bróm-6-klórpírid-3-il-, 5-fluor-6-brómpírid-3-il-, 5-klór-6-brómpírid-3-il-, 5,6-dibrómpírid-3-il-, 5-metil-6-klórpírid-3-il- vagy 5-metil-6-brómpírid-3-il-csoport egyike és

R¹ jelentése minden esetben fluoratommal szubsztituált C₁-C₃-alkil-, C₂-C₃-alkenil-, C₂-C₃-cikloalkil- vagy C₃-C₃-cikloalkilalkil-csoport.

12. A 11. igénypont szerinti vegyület, azaz N-[(6-klór-5-fluorpiridin-3-il)metil]-2,2-difluoretán-1-amin.