



(51) International Patent Classification:

C07D 413/14 (2006.01) C07D 273/00 (2006.01)

C07D 413/10 (2006.01) A61P 33/00 (2006.01)

C07D 417/14 (2006.01) A61K 38/15 (2006.01)

(21) International Application Number:

PCT/US2019/031158

(22) International Filing Date:

07 May 2019 (07.05.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/669,623 10 May 2018 (10.05.2018) US

(71) Applicant: ZOETIS SERVICES LLC [US/US]; 10 Sylvan Way, Parsippany, NJ 07054 (US).


(72) Inventors: MADDUX, Todd; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US). BEDORE, Matthew, W.; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US). JOHNSON, Paul, D.; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US).

49007 (US). **RESPONDEK, Tomasz**; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US). **SHEEHAN, Susan, M. K.**; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US). **KYNE, Graham, M.**; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US). **EWIN, Richard, Andrew**; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US). **VAIRAGOUNDAR, Rajendran**; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US). **CURTIS, Michael, P.**; c/o Zoetis Services LLC, 333 Portage Street, Kalamazoo, MI 49007 (US).

(74) Agent: MISIAK, Paul, M. et al.; c/o Zoetis Services LLC, 10 Sylvan Way, Parsippany, NJ 07054 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

(54) Title: ENDOPARASITIC DEPSIPEPTIDES



(57) **Abstract:** The present invention provides cyclic depsipeptides of Formula (1), stereoisomers thereof, and veterinary acceptable salts thereof wherein each of R^1 , R^2 , R^3 , R^4 , L_1 , and L_2 are as defined herein. The present invention also contemplates compositions, methods of treatment, and uses as a medicament to treat an animal for an endoparasitic infection with a Formula (1) compound.



SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to the identity of the inventor (Rule 4.17(i))*
- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*

ENDOPARASITIC DEPSIPEPTIDES

Field of the Invention

The present invention is directed to new endoparasitic depsipeptide compounds with improved activity against endoparasites. The invention is also directed to compositions comprising the compounds, methods and uses of the compounds for eradicating, controlling, treating and preventing a parasite infestation and/or infection in animals. The compounds of the invention may be administered to animals, particularly non-human animals, to prevent or treat parasitic infections.

Background

Animals, such as non-human mammals, for example, companion animals and livestock, are often susceptible to parasite infestations. These parasites may be endoparasites including for example, helminthiasis which is most frequently caused by a group of parasitic worms described as nematodes or roundworms. These parasites cause severe economic losses in pigs, sheep, horses, and cattle as well as affecting companion animals (e.g. cats and dogs). Other parasites include those which occur in the gastrointestinal tract of animals and humans include *Ancylostoma*, *Necator*, *Ascaris*, *Strongyloides*, *Trichinella*, *Capillaria*, *Toxocara*, *Toxascaris*, *Trichuris*, *Enterobius* and parasites which are found in the blood or other tissues and organs such as filarial worms and the extra intestinal stages of *Strongyloides*, *Toxocara* and *Trichinella*.

One type of endoparasite which seriously harms animals is *Dirofilaria immitis*, also known as heartworm. Other filarial endoparasites include *Dirofilaria repens* and *Dirofilaria honkongensis*, which can also infect humans. The most common hosts are dogs and cats but other mammals such as ferrets and raccoons may also be infected. Heartworms go through several life stages before they become adults infecting the pulmonary artery of the host mammal, particularly canine. The worms require the mosquito as an intermediate host to complete their life cycle. The period between the initial infection when the dog is bitten by a mosquito and the maturation of the worms into adults living in the heart and pulmonary arteries is six to seven months in dogs and is known as the

"prepatent period". L3 larvae migrate during blood feeding of the mosquito to the tip of the mosquito's mouth parts (labium), leave the mosquito and are deposited on the skin of the dog where they then migrate through the bite wound into the host. Most L3 larvae molt to fourth-stage larvae (L4's) in canine subcutaneous tissues within 1-3 days after infection. Then, they migrate to the muscles of the chest and abdomen, and 45 to 60 days after infection, molt to the fifth stage (L5, immature adult). Between 75 and 120 days after infection, these immature heartworms then enter the bloodstream and are carried through the heart to reside in the pulmonary artery. Around seven months after infection, *Dirofilaria immitis* adults reach maturity and sexually reproduce in the pulmonary arteries and right ventricle. Adult males are around 15cm in length, and females are around 25cm in length and their normal life span as adults is calculated to be about 5 years.

Heartworm infection is a severe and life-threatening disease. Canine heartworm infection is preventable and prophylaxis treatment is a priority in heartworm endemic areas. Treatment of mature heartworm infection with an adulticide (e.g. melarsomine dihydrochloride) is costly and can cause serious adverse side effects, thus prevention by monthly administration of drugs that interrupt larvae development is widely used. The goal of marketed heartworm preventive therapies in dogs is to prevent the development of the parasite to adult heartworms by interrupting the *Dirofilaria immitis* life cycle post-infection. The macrocyclic lactones (MLs, e.g. ivermectin, eprinomectin, milbemycin oxime, moxidectin, and selamectin) are the most commonly used chemoprophylaxis agents and are administered at monthly or six-month intervals. These drugs have been effective against *Dirofilaria immitis* infective third-stage larvae (L3) deposited by the mosquito as well as maturing fourth-stage larvae (L4). When administered monthly, MLs kill L3 and L4 larvae acquired within the previous 30 days, and thus prevent disease caused by adult worms. MLs can also be used monthly in infected dogs to suppress reproduction in adult worms and remove microfilariae, thereby reducing transmission and gradually causing the attrition of adult worms (*Vet. Parasitol.* 2005 Oct 24 133(2-3) 197-206).

In recent years, an increased number of lack of efficacy cases have been reported, in which dogs develop mature heartworm infections despite receiving monthly prophylactic doses of macrocyclic lactone drugs. For example,

Atkins *et al.*, (*Vet. Parasitol.* 206 (2014) 106-113) recently reported that an increasing number of cases of dogs that tested heartworm antigen positive while receiving heartworm preventive medication which suggests that some populations of *Dirofilaria immitis* have developed selectional resistance to heartworm preventives (American Heartworm Society, 2010. Heartworm Preventive Resistance. Is it Possible, vol. 37. Bulletin of the American Heartworm Society, pp. 5.). Thus, there is an ongoing need to develop new anthelmintic agents with improved activity against *Dirofilaria immitis* and other endoparasites.


Various parasiticides exist in the art for treating endoparasites infections in animals. In addition to the macrocyclic lactones, cyclic depsipeptides with antiparasitic activity are known. PF1022a, a 24-membered cyclooctadepsipeptide isolated from the fungus *Afvelia sterilia* by Sasaki *et al* (*J Antibiotics* 45: 692-697 (1992)), has been found to exhibit broad anthelmintic activity against a variety of endoparasites *in vivo* with low toxicity. These compounds are described, for example, in U.S. Patent Nos. 5,514,773; 5,747,448; 5,646,244; 5,874,530; among others, which are incorporated herein by reference. Emodepside is a semi-synthetic analog of PF1022a containing a morpholine group at the *para* position of the aryl ring in the phenyl lactate groups. Emodepside is a potent anthelmintic used in combination with praziquantel in the product Profender® for the treatment of parasitic worms in cats and dogs. However, the antiparasitic activity of PF 1022a and emodepside is not satisfactory for the treatment of certain parasites, especially for the control of *Dirofilaria immitis* in mammals to prevent the establishment of heartworm disease. Thus, there is a need in the art for more effective antiparasitic agents for treatment and protection of animals against internal parasites including nematodes and filarial worms such as heartworm.

Summary of the Invention

The invention provides novel and inventive cyclic depsipeptide compounds with selective anthelmintic activity against L3 and L4 staged endoparasites, particularly, *Dirofilaria immitis* L4 (DiL4). In addition, the invention provides compositions comprising the depsipeptide compounds and methods and uses for

the treatment and prevention of parasitic infection and possibly infestation of animals using the compounds.

In one aspect, the present invention provides symmetrical (*i.e.*, L₁ and L₂ are the same; bis) and asymmetric (*i.e.*, L₁ is absent) cyclic depsipeptides of Formula (1)



(1)

wherein

L₁ and L₂ are the same and are C₁-C₄alkyl, C₃-C₆cycloalkyl, C₁-C₄alkylaryl, C₁-C₄alkylheteroaryl or C₁-C₄alkylheterocycle; or L₁ is absent; and wherein the cycloalkyl is a 3- to 6-membered carbocyclic ring; the aryl is a 6-membered monocyclic aromatic ring or a 9- to 10-membered fused aromatic ring; the heteroaryl is a 5- or 6-member monocyclic aromatic ring or a 9- to 10-member fused aromatic ring wherein each mono- or fused-heteroaryl ring contains at least one heteroatom selected from N, O, and S; and the heterocycle is a 4- to 6-member monocyclic saturated or partially saturated ring or a 9- to 10-member fused saturated or partially saturated ring, each mono- or fused-heterocyclic ring contains at least one heteroatom selected from N, O and S; and wherein the heteroaryl and heterocycle rings are linked to the L₁ and L₂ C₁-C₄alkyl or L₂ C₁-C₄alkyl by a ring carbon atom;

and wherein the L₁ and L₂ alkyl group of the alkylcycloalkyl, alkylaryl, alkylheteroaryl and alkylheterocycle can be substituted with an R^c and R^d substituent each individually and separately selected from the group consisting of hydroxyl, halo, C₁-C₆alkyl, C₃-C₆cycloalkyl, C₁-C₆alkoxy, C₁-C₄haloalkyl, -NR^aR^b, pyrazolyl, imidazolyl, pyrrolidinyl, and morpholinyl; and when L₁ is not absent, then each R^c is the same and each R^d is the same;

and wherein the L₁ and L₂ cycloalkyl, aryl, heteroaryl, and heterocycle group can be substituted with (R)_n; and when L₁ is not absent, then each (R)_n is the same; and each R substituent is independently selected from the group consisting of C₁-C₆alkyl optionally substituted with at least one substituent selected from hydroxyl, C₃-C₆cycloalkyl and -OR^a; halo, oxo, cyano, hydroxyl, -OR⁵, -NR^aR^b, C₁-C₄haloalkyl, C₁-C₄haloalkoxy, -S(O)_pR^a, C₁-C₆alkoxy, C₃-C₆cycloalkyl, -C(O)NR^aR^b, -C(O)R^a, -C(O)OR^a, -NR^aC(O)R^b, pyrazolyl, imidazolyl, pyridinyl, pyrimidinyl, piperidinyl, morpholinyl, pyrrolidinyl, dihydropyrimidinyl, and phenyl; and wherein the phenyl is optionally substituted with at least one substituent selected from C₁-C₃alkyl, -CF₃, halo, and hydroxyl; and wherein n is the integer 0, 1, 2, or 3;

R¹, R², R³, and R⁴ are each separately and independently H and C₁-C₆alkyl;

R^a and R^b are each separately H and C₁-C₆alkyl; and

R⁵ is C₃-C₆cycloalkyl and C₁-C₃alkylC₃-C₆cycloalkyl;


p is the integer 0, 1, or 2;

stereoisomers thereof, and veterinary acceptable salts thereof.

In another aspect of the invention, is a composition comprising a Formula (1) compound. In yet another aspect, the Formula (1) composition further comprises a veterinary acceptable excipient. In another aspect of the invention, is a method of treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (1) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (1) compound for treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (1) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (1) compound for preparing a medicament for the use in treating a parasitic infection in an animal.

In one aspect of the invention, R¹, R², R³, and R⁴ are the same and are each C₁-C₆alkyl. In another aspect, R¹, R², R³, and R⁴ are the same and are each methyl, ethyl, propyl, isopropyl, isobutyl, and n-butyl. Preferably, each of R¹, R², R³, and R⁴ are the same and are each ethyl, isopropyl or isobutyl. More preferred, R¹, R², R³, and R⁴ are the same and are each isopropyl or isobutyl. Most preferred, R¹, R², R³, and R⁴ are the same and are each isobutyl.


In one aspect of the invention, is a Formula (1) compound that is a Formula (1A) compound



stereoisomers thereof, and veterinary acceptable salts thereof.

In another aspect of the invention, is a composition comprising a Formula (1A) compound. In yet another aspect, the Formula (1A) composition further comprises a veterinary acceptable excipient. In another aspect of the invention, is a method of treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (1A) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (1A) compound for treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (1A) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (1A) compound for preparing a medicament for the use in treating a parasitic infection in an animal.

In another aspect of the invention, is a Formula (1) compound that is a Formula (2A) compound,



stereoisomers thereof, and veterinary acceptable salts thereof.

In another aspect of the invention, is a composition comprising a Formula (2A) compound. In yet another aspect, the Formula (2A) composition further comprises a veterinary acceptable excipient. In another aspect of the invention, is a method of treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (2A) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (2A) compound for treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (2A) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (2A) compound for preparing a medicament for the use in treating a parasitic infection in an animal.

In one aspect of the invention, when L_1 and L_2 are present, they are the same, i.e., a symmetrical bis depsipeptide compound of Formula (1). In another aspect, L_1 is absent, i.e., a non-symmetrical depsipeptide compound of Formula (1).

In one aspect of the invention, L_1 and L_2 are the same and are each a C_1 alkyl/ C_3 - C_6 cycloalkyl (-C- C_3 - C_6 cycloalkyl); a C_1 alkylaryl (-C-aryl); a - C_1 alkylheterocycle (-C-heterocycle); or a - C_1 alkylheteroaryl (-C-heteroaryl); or L_1 is absent; and wherein each cycloalkyl, aryl, heterocycle, or heteroaryl group can be substituted with $(R)_n$ as described herein; and each -C- of the alkyl group can be substituted with R^c and R^d , as described herein.

In one aspect of the invention, L_1 and L_2 are the same and are each a C_1 alkyl/ C_3 - C_6 cycloalkyl (-C- C_3 - C_6 cycloalkyl); or L_1 is absent. In another aspect, the L_1 and L_2 cycloalkyls are the same and are -C-cyclopropyl, -C-cyclobutyl, -C-cyclopentyl, or -C-cyclohexyl; or L_1 is absent. The cycloalkyl groups of L_1 and L_2 or L_2 can be substituted with $(R)_n$ as described herein; and each alkyl group of L_1 and L_2 or L_2 can be substituted with R^c and R^d , as described herein. In Formula (3A), Ring A and Ring B is the L_1 and L_2 cycloalkyl group, respectively. In Formula (4A), Ring B is the L_2 cycloalkyl group.


In one aspect of the invention, L_1 and L_2 are the same and are each a C_1 alkylaryl (-C-aryl); or L_1 is absent. In another aspect, the L_1 and L_2 aryls are the same and are each -C-phenyl or -C-naphthyl; or L_1 is absent. In yet another aspect, the L_1 and L_2 aryls are each -C-phenyl; or L_1 is absent. The aryl groups of L_1 and L_2 or L_2 can be substituted with $(R)_n$, as described herein; and each alkyl group of L_1 and L_2 or L_2 can be substituted with R^c and R^d , as described herein.

In one aspect of the invention, L₁ and L₂ are the same and are each a C₁alkylheterocycle (-C-heterocycle); or L₁ is absent. In another aspect, the L₁ and L₂ heterocycles are the same and are each oxetanyl, thietanyl, azetidiny, tetrahydrofuranyl, tetrahydrothiophenyl, pyrrolidinyl, tetrahydropyranyl, tetrahydropyridinyl, tetrahydrothiopyranyl, piperidinyl, 1,4-dioxanyl, morpholinyl, piperazinyl, azathianyl, or 3,4-dihydro-2H-pyranyl; or L₁ is absent. In yet another aspect, the L₁ and L₂ heterocycles are the same and are each oxetanyl, cyclohexyl, azetidiny, pyrrolidinyl, tetrahydropyranyl, piperidinyl, or morpholinyl; or L₁ is absent. In yet another aspect, the L₁ and L₂ heterocycles are the same and are each pyrrolidinyl, tetrahydropyranyl, piperidinyl, or morpholinyl; or L₁ is absent. In yet another aspect, the L₁ and L₂ heterocycles are the same and are each morpholinyl; or L₁ is absent. The heterocycle groups of L₁ and L₂ or L₂ can be substituted with (R)_n as described herein; and each alkyl group of L₁ and L₂ or L₂ can be substituted with R^c and R^d, as described herein. In Formula (3A), Ring A and Ring B is the L₁ and L₂ heterocycle group, respectively. In Formula (4A), Ring B is the L₂ heterocycle group.

In one aspect of the invention, L₁ and L₂ are the same and are each a C₁alkylheteroaryl (-C-heteroaryl); or L₁ is absent. In another aspect, the L₁ and L₂ heteroaryls are the same and are each pyrrolyl, furanyl, thiophenyl, pyrazolyl, imidazolyl, isoxazolyl, oxazolyl, isothiazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, dihydropyridinyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, quinolinyl, isoquinolinyl, benzofuranyl, indazolyl, benzothiophenyl, indolyl, benzimidazolyl, pyrazolo[1,5-a]pyridinyl, pyrazolo[1,5-a]pyrimidinyl, 1H-pyrrolo[3,2-b]pyridinyl, indazolebenzo[d][1,3]diazolyl, benzo[d]thiazolyl, furo[2,3-b]pyridinyl, 2,3-dihydrothieno[3,4-b][1,4]dioxinyl, 3,4-dihydro-2H-pyrano[2,3-b]pyridinyl, dihydro-1H-pyrrolo[3,2-b]pyridine, dihydro-1H-pyrrolo[3,2-b]pyridinyl, oxazolo[5,4-b]pyridinyl, or benzo[d]1,3 diaxolyl; or L₁ is absent. In yet another aspect, the L₁ and L₂ heteroaryls are the same and are each thiophenyl, thiazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, oxazolyl, isoxazolyl, isothiazolyl, or furanyl; or L₁ is absent. In yet another aspect, the L₁ and L₂ heteroaryls are the same and are each thiophenyl, thiazolyl, pyrazolyl, pyridinyl, pyridazinyl, pyrazinyl, or pyrimidinyl; or L₁ is absent. In yet another aspect, the L₁ and L₂ heteroaryls are the same and are each pyrazolyl, pyridinyl, pyridazinyl, pyrazinyl, and pyrimidyl; or L₁ is absent. In yet another aspect, the L₁ and L₂ heteroaryls are the same and are each pyrazolyl, pyridinyl, or


pyrimidinyl; or L_1 is absent. In yet another aspect, the L_1 and L_2 heteroaryls are the same and are each pyridazinyl; or L_1 is absent. In yet another aspect, the L_1 and L_2 heteroaryls are the same and are each thiazole; or L_1 is absent. In yet another aspect of the invention, the L_1 and L_2 heteroaryls are the same and are each pyrazole, or L_1 is absent. In yet another aspect, the L_1 and L_2 heteroaryls are the same and are each pyridine; or L_1 is absent. In yet another aspect, the L_1 and L_2 heteroaryls are the same and are each pyrimidine; or L_1 is absent. In yet another aspect, the L_1 and L_2 heteroaryls are the same and are each pyrazinyl; or L_1 is absent. The heteroaryl groups of L_1 and L_2 or L_2 can be substituted with $(R)_n$ as described herein; and each alkyl group of L_1 and L_2 or L_2 can be substituted with R^c and R^d , as described herein. In Formula (3A), Ring A and Ring B is the L_1 and L_2 heteroaryl group, respectively. In Formula (4A), Ring B is the L_2 heteroaryl group.

In one aspect of the invention, is a Formula (1A) compound that is a Formula (1A1) compound



stereoisomers thereof, and veterinary acceptable salts thereof. Each of $(R)_n$, R^c , and R^d are as described herein; and each $(R)_n$ is the same, each R^c is the same, and each R^d is the same; stereoisomers thereof, and veterinary acceptable salts thereof.


In one aspect of the invention, is a Formula (2A) compound that is a Formula (2A1) compound,



Each of $(R)_n$, R^c , and R^d are as described herein; stereoisomers thereof, and veterinary acceptable salts thereof.


In another aspect of the invention, is a composition comprising a Formula (1A1) or (2A1) compound. In yet another aspect, the Formula (1A1) or (2A1) composition further comprises a veterinary acceptable excipient. In another aspect of the invention, is a method of treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (1A1) or (2A1) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (1A1) or (2A1) compound for treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (1A1) or (2A1) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (1A1) or (2A1) compound for preparing a medicament for the use in treating a parasitic infection in an animal.

In another aspect of the invention are Formula (1A) compounds that are Formula (3A) compounds,



Each of Ring A, Ring B, $(R)_n$, R^c , and R^d are as described herein. Ring A and Ring B are the same, and each $(R)_n$ is the same, each R^c is the same, and each R^d is the same; stereoisomers thereof, and veterinary acceptable salts thereof.

In yet another aspect of the invention, is a Formula (2A) compound that is a Formula (4A) compound



Each of Ring B, $(R)_n$, R^c , and R^d are as described herein; stereoisomers thereof, and veterinary acceptable salts thereof.

In another aspect of the invention, is a composition comprising a Formula (3A) or (4A) compound. In yet another aspect, the Formula (3A) or (4A) composition further comprises a veterinary acceptable excipient. In another aspect of the invention, is a method of treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (3A) or (4A) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (3A) or (4A) compound for treating a parasitic infection in an animal by administering a therapeutic amount of a Formula (3A) or (4A) compound to the animal in need thereof. In yet another aspect, is the use of a Formula (3A) or (4A) compound for preparing a medicament for the use in treating a parasitic infection in an animal.

As described herein, Ring A and Ring B are the L_1 and L_2 cycloalkyl, heterocycle, or heteroaryl groups of Formula (1), respectively; or L_1 is absent. Ring A and Ring B of Formula (3A) are the same; and Ring B of Formula (4A); are each cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, oxetanyl, thietanyl, azetidiny, tetrahydrofuranyl, tetrahydrothiophenyl, pyrrolidinyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidinyl, dioxanyl, morpholinyl, piperazinyl, azathianyl, 3,4-dihydro-2H-pyranyl, 4,5,6,7-tetrahydropyrazolo[1,5-a]pyridinyl, 4,5-

tetrahydropyrazolo[1,5-a]pyrimidine, pyrrolyl, furanyl, thiophenyl, pyrazolyl, imidazolyl, isoxazolyl, oxazolyl, isothiazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, dihydropyridinyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, quinolonyl, isoquinolonyl, benzofuranyl, indazolyl, benzothiophenyl, indolyl, triazinyl, benzimidazolyl, tetrahydropyranyl, pyrazolo[1,5-a]pyridinyl, pyrazolo[1,5-a]pyrimidinyl, 1H-pyrrolo[3,2-b]pyridinyl, indazolebenzo[d][1,3]diazolyl, benzo[d]thiazolyl, furo[2,3-b]pyridinyl, 2,3-dihydrothieno[3,4-b][1,4]dioxinyl, 3,4-dihydro-2H-pyrano[2,3-b]pyridinyl, oxazolo[5,4-b]pyridinyl, or benzo[d]1,3 diaxolyl. In another aspect, Ring A and Ring B of Formula (3A) are the same; and Ring B of Formula (4A); are each pyrrolidinyl, tetrahydropyranyl, piperidinyl, morpholinyl, thiadiazolyl, thiophenyl, thiazolyl, pyrazolyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, oxazolyl, isoxazolyl, isothiazolyl, or furanyl. In yet another aspect, Ring A and B are the same and are each morpholinyl, thiophenyl, thiazolyl, pyrazolyl, pyridinyl, pyridazinyl, pyrazinyl, or pyrimidinyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are each pyrazolyl, thiazolyl, thiadiazolyl, isothiazolyl, pyridinyl, pyridazinyl, pyrazinyl, and pyrimidyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are each pyrazolyl, pyrazinyl, isothiazolyl, thiazolyl, pyridinyl, pyridazinyl, and pyrimidinyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are each pyrazolyl, isothiazolyl, thiazolyl, pyridinyl, pyrazinyl, or pyrimidinyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are each isothiazolyl, thiazolyl, pyridinyl, pyrazinyl, or pyrimidinyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are pyrazolyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are isothiazolyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are thiazolyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are pyridinyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are pyrazinyl. In yet another aspect, Ring A and B of Formula (3A) are the same; and Ring B of Formula (4A); are pyrimidinyl.

In one aspect of the invention, R of (R)_n is independently selected from the group consisting of C₁-C₆alkyl optionally substituted with at least one substituent selected from hydroxyl, C₃-C₆cycloalkyl, and -OR^a; halo, oxo, cyano, hydroxyl,

-OR⁵, -NR^aR^b, C₁-C₄haloalkyl, C₁-C₄haloalkoxy, -S(O)_pR^a, C₁-C₆alkoxy, C₃-C₆cycloalkyl, -C(O)NR^aR^b, -C(O)R^a, -C(O)OR^a, -NR^aC(O)R^b, pyrazolyl, imidazolyl, pyridinyl, pyrimidinyl, piperidinyl, morpholinyl, pyrrolidinyl, dihydropyrimidinyl, and phenyl; and wherein the phenyl is optionally substituted with at least one substituent selected from C₁-C₃alkyl, -CF₃, halo, and hydroxyl; and wherein n of (R)_n is the integer 0, 1, 2, or 3; and wherein each (R)_n of Formula (1A1) and Formula (3A) are the same. In another aspect, R of (R)_n is independently selected from the group consisting of C₁-C₆alkyl optionally substituted with at least one substituent selected from hydroxyl, OR^a, and C₃-C₆cycloalkyl; halo, cyano, hydroxyl, -NR^aR^b, C₁-C₄haloalkyl, C₁-C₄haloalkoxy, -S(O)_pR^a, C₁-C₆alkoxy, -OR⁵, C₃-C₆cycloalkyl, morpholinyl, and phenyl optionally substituted with methyl, -CF₃, halo, and hydroxyl; and wherein n is the integer 0, 1, 2, or 3; and wherein each (R)_n of Formula (1A1) and Formula (3A) are the same. In yet another aspect, R of (R)_n is independently selected from the group consisting of C₁-C₆alkyl optionally substituted with at least one substituent selected from hydroxyl and C₃-C₆cycloalkyl; halo, cyano, hydroxyl, -NR^aR^b, C₁-C₄haloalkyl, C₁-C₄haloalkoxy, -S(O)_pR^a, C₁-C₆alkoxy, C₃-C₆cycloalkyl, morpholinyl, and phenyl optionally substituted with methyl, -CF₃, halo, and hydroxyl; and wherein n is the integer 0, 1, 2, or 3; and wherein each (R)_n of Formula (1A1) and Formula (3A) are the same. In yet another aspect, R of (R)_n is independently selected from the group consisting of methyl, ethyl, propyl, isopropyl, isobutyl, cyclopropyl, cyclobutyl, -C-cyclopropyl, halo, -SCH₃, -S(O)₂CH₃, cyano, hydroxyl, -C-CF₃, -CF₃, -C-CF₂, -NR^aR^b, methoxy, ethoxy, isopropoxy, -O-C-CF₃, -O-CF₂, -COH, morpholinyl, and phenyl; and wherein the phenyl can be further substituted with methyl; and wherein n of (R)_n is the integer 0, 1, 2, or 3; and wherein each (R)_n of Formula (1A1) and Formula (3A) are the same. In yet another aspect, R of (R)_n is independently selected from the group consisting of methyl, ethyl, isopropyl, cyclopropyl, -C-cyclopropyl, halo, -SCH₃, cyano, hydroxyl, -C-CF₃, -CF₂, -CF₃, -N(CH₃)₂, methoxy, ethoxy, -OCF₃, -OCF₂, and morpholinyl; and wherein n of (R)_n is the integer 0, 1, 2, or 3; and wherein each (R)_n of Formula (1A1) and Formula (3A) are the same.

In one aspect of the invention, n of (R)_n of Formula (1A1) and (3A) is the same integer; and n of (R)_n of Formula (2A) and (4A); is the integer 0, 1, 2, or 3. In another aspect, n is the integer 0. In yet another aspect, n is the integer 1. In yet another aspect, n is the integer 2. In yet another aspect, n is the integer 3.

In one aspect of the invention, p is the integer 0, 1, or 2. In another aspect, p is the integer 0. In yet another aspect, p is the integer 1. In yet another aspect, p is the integer 2.

In one aspect of the invention, R^a and R^b are each separately selected from H, methyl, ethyl, propyl, isopropyl, and iso-butyl. In another aspect of the invention, R^a and R^b are each separately H, methyl, ethyl, and isopropyl. In another aspect, R^a and R^b are each separately H, methyl, and ethyl.

In one aspect of the invention, R^c and R^d are each independently and separately selected from the group consisting of H, hydroxyl, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-CF_3$, cyclopropyl, $-NR^aR^b$, pyrazolyl, pyrrolidinyl, and morpholinyl; and wherein each R^c of Formula (1A1) and Formula (3A) are the same and each R^d of Formula (1A1) and Formula (3A) are the same. In another aspect of the invention, R^c and R^d are each independently and separately selected from the group consisting of H, hydroxyl, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NR^aR^b$, $-CF_3$, cyclopropyl, pyrazolyl, pyrrolidinyl, and morpholinyl; and wherein each R^c of Formula (1A1) and Formula (3A) are the same and each R^d of Formula (1A1) and Formula (3A) are the same.

In yet another aspect of the invention, R^c and R^d are each independently and separately selected from H, F, Cl, C_1 - C_6 alkyl, methoxy, ethoxy, isopropoxy, $-CF_3$, hydroxyl, $-N(CH_3)_2$, $-N(CH_2CH_3)_2$, cyclopropyl, pyrazolyl, and morpholine; and wherein each R^c of Formula (1A1) and Formula (3A) are the same and each R^d of Formula (1A1) and Formula (3A) are the same. In another aspect, R^c and R^d are each independently and separately selected from H, F, Cl, methyl, ethyl, isopropyl, hydroxyl, $-CF_3$, methoxy, ethoxy, isopropoxy, cyclopropyl, and $-N(CH_2CH_3)_2$; and wherein each R^c of Formula (1A1) and Formula (3A) are the same and each R^d of Formula (1A1) and Formula (3A) are the same. In yet another aspect, R^c and R^d are each independently and separately selected from H, F, methyl, ethyl, hydroxyl, $-CF_3$, methoxy, ethoxy, and $-N(CH_2CH_3)_2$; and wherein each R^c of Formula (1A1) and Formula (3A) are the same and each R^d of Formula (1A1) and Formula (3A) are the same. In yet another aspect, R^c and R^d are each independently and separately selected from H, F, methyl, ethyl, hydroxyl, methoxy, and ethoxy; and wherein each R^c of Formula (1A1) and Formula (3A) are the same and each R^d of Formula (1A1) and Formula (3A) are the same. In yet another aspect, R^c and R^d are each independently and separately selected from H, F, methyl, hydroxyl, and methoxy; and wherein each R^c of Formula (1A1) and Formula (3A) are the same and each R^d

of Formula (1A1) and Formula (3A) are the same. In yet another aspect, R^c and R^d are each independently and separately selected from H, hydroxyl, and methoxy; and wherein each R^c of Formula (1A1) and Formula (3A) are the same and each R^d of Formula (1A1) and Formula (3A) are the same.

In one aspect of the invention, R^5 is cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, C₁-cyclopropyl, C₂-cyclopropyl, C₁-cyclobutyl, C₂-cyclobutyl, C₂-cyclopentyl, C₂-cyclohexyl, C₃-cyclopropyl, or C₃-cyclobutyl. In another aspect, R^5 is cyclopropyl, cyclobutyl, C₁-cyclopropyl, C₂-cyclopropyl, C₁-cyclobutyl, or C₂-cyclobutyl. In yet another aspect, R^5 is cyclopropyl, cyclobutyl, C₁-cyclopropyl, or C₁-cyclobutyl. In yet another aspect, R^5 is cyclopropyl, cyclobutyl, or C₁-cyclopropyl. In yet another aspect, R^5 is cyclopropyl. In yet another aspect, R^5 is cyclobutyl. In yet another aspect, R^5 is C₁-cyclopropyl.

In yet another aspect of the invention are Formula (1A1) compounds selected from the group consisting of those compounds described in Table 1, stereoisomers thereof, and veterinary acceptable salts thereof. In another aspect, is a composition comprising a compound described in Table 1. In another aspect, is a method of treating a parasitic infection in an animal in need thereof by administering a therapeutically effective amount of a Table 1 compound to the animal. In another aspect, is the use of a compound described in Table 1 to prepare a medicament for the treatment of a parasitic infection in an animal.

In another aspect of the invention are Formula (1A1) compounds selected from the group consisting of Examples: (1-5), (1-11), and (1-18), stereoisomers thereof, and veterinary acceptable salts. Preferred (1A1) compounds are Examples (1-5) and (1-11), stereoisomers thereof, and veterinary acceptable salts thereof. In another aspect, is a composition comprising one of these Formula (1A1) compounds, stereoisomers thereof, and veterinary acceptable salts thereof. In another aspect, the composition further comprises a veterinary acceptable excipient. In another aspect, is a method of treating a parasitic infection in an animal in need thereof by administering a therapeutically effective amount of one of these Formula (1A1) compounds to the animal. In another aspect, is the use of one of these Formula (1A1) compounds to prepare a medicament for the treatment of a parasitic infection in an animal.

In yet another aspect of the invention are Formula (2A1) compounds selected from the group consisting of those compounds described in Table 2, stereoisomers

thereof, and veterinary acceptable salts thereof. In another aspect, is a composition comprising a compound described in Table 2. In another aspect, is a method of treating a parasitic infection in an animal in need thereof by administering a therapeutically effective amount of a Table 2 compound to the animal. In another aspect, is the use of a compound described in Table 2 to prepare a medicament for the treatment of a parasitic infection in an animal.

In another aspect of the invention are Formula (2A1) compounds selected from the group consisting of Examples: (2-1), (2-4), (2-36), (2-37), (2-38), and (2-42), stereoisomers thereof, and veterinary acceptable salts thereof; and veterinary acceptable salts thereof. Preferred (2A1) compounds are Examples (2-4), (2-36), and (2-42), stereoisomers thereof, and veterinary acceptable salts thereof. In another aspect, is a composition comprising one of these Formula (2A1) compounds, stereoisomers thereof, and veterinary acceptable salts thereof. In another aspect, the composition further comprises a veterinary acceptable excipient. In another aspect, is a method of treating a parasitic infection in an animal in need thereof by administering a therapeutically effective amount of one of these Formula (2A1) compounds to the animal. In another aspect, is the use of one of these Formula (2A1) compounds to prepare a medicament for the treatment of a parasitic infection in an animal.

In yet another aspect of the invention are Formula (3A) compounds selected from the group consisting of those compounds described in Table 3, stereoisomers thereof, and veterinary acceptable salts thereof. In another aspect, is a composition comprising a compound described in Table 3. In another aspect, is a method of treating a parasitic infection in an animal in need thereof by administering a therapeutically effective amount of a Table 3 compound to the animal. In another aspect, is the use of a compound described in Table 3 to prepare a medicament for the treatment of a parasitic infection in an animal.

In yet another aspect of the invention is a Formula (3A) compound selected from the group consisting of Examples: (3-4), (3-8), (3-12), (3-17), (3-18), (3-21), (3-24), (3-31), (3-44), (3-49), (3-50), (3-56), (3-57), (3-60), (3-69), (3-87), (3-110), (3-112), (3-118), (3-120), (3-121), (3-124), (3-125), (3-132), (3-133), (3-148), (3-155), (3-156), (3-157), (3-165), (3-173), (3-176), (3-178), (3-193), (3-194), (3-207), (3-209), (3-215), (3-217), (3-218), (3-220), (3-223), (3-225), (3-228), (3-238), (3-240), (3-243), (3-252), (3-253), (3-254), (3-259), (3-261), (3-268), (3-273), (3-274), (3-275), (3-282),

(3-283), (3-284), (3-285), (3-286), (3-287), (3-288), (3-289), (3-290), (3-291), (3-292), (3-293), (3-294), (3-295), (3-296), (3-297), (3-307), (3-308), (3-309), (3-310), (3-313), (3-318), (3-319), (3-321), (3-322), (3-323), (3-324), (3-326), (3-329), (3-332), (3-333), and (3-334), stereoisomers thereof, and veterinary acceptable salts thereof.

Preferred (3A) compounds are Examples (3-4), (3-44), (3-56), (3-69), (3-87), (3-110), (3-118), (3-121), (3-124), (3-125), (3-133), (3-157), (3-165), (3-173), (3-207), (3-225), (3-261), (3-284), (3-286), (3-287), (3-293), (3-296), (3-324), (3-326), and (3-332), stereoisomers thereof, and veterinary acceptable salts thereof. In yet another aspect of the invention is a composition comprising one of these Formula (3A) compounds. In yet another aspect, the composition further comprises a veterinary acceptable excipient. In another aspect, is a method of treating a parasitic infection in an animal in need thereof by administering a therapeutically effective amount of one of these Formula (3A) compounds to the animal. In another aspect, is the use of one of these Formula (3A) compounds to prepare a medicament for the treatment of a parasitic infection in an animal.

In yet another aspect of the invention are Formula (4A) compounds selected from the group consisting of those compounds described in Table 4, stereoisomers thereof, and veterinary acceptable salts thereof. In another aspect, is a composition comprising a compound described in Table 4. In another aspect, is a method of treating a parasitic infection in an animal in need thereof by administering a therapeutically effective amount of a Table 4 compound to the animal. In another aspect, is the use of a compound described in Table 4 to prepare a medicament for the treatment of a parasitic infection in an animal.

In yet another aspect of the invention, is a Formula (4A) compound selected from the group consisting of Examples: (4-18), (4-19), (4-21), (4-30), (4-32), (4-34), (4-40), (4-50), (4-63), (4-75), (4-105), (4-107), (4-109), (4-118), (4-126), (4-139), and (4-141), stereoisomers thereof, and veterinary acceptable salts thereof. Preferred Formula (4A) compounds are selected from the group of Examples: (4-18), (4-19), and (4-21), stereoisomers thereof, and veterinary acceptable salts thereof. In yet another aspect of the invention is a composition comprising one of the Formula (4A) compounds. In yet another aspect, the composition further comprises a veterinary acceptable excipient. In another aspect, is a method of treating a parasitic infection in an animal in need thereof by administering a therapeutically effective amount of one of these Formula (4A) compounds to the animal. In another aspect, is the use

of one of these Formula (4A) compounds to prepare a medicament for the treatment of a parasitic infection in an animal.

In yet another aspect, is a Formula (1) compound that is at least 10x selective against *D.immitis* L4 (i.e., DiMf/DiL4). In yet another aspect of the invention is a composition comprising a Formula (1) compound that is at least 10x selective against *D.immitis* L4 and a veterinary acceptable excipient. In yet another aspect of the invention is a composition comprising a therapeutically effective amount of a Formula (1) compound that is at least 10x selective over *D.immitis* L4. In yet another aspect of the invention is a composition comprising a therapeutically effective amount of a Formula (1) compound that is at least 10x selective over *D.immitis* L4 and a veterinary acceptable excipient. In another aspect, the Formula (1) compounds that are at least 10x selective include Example #'s: (1-18), (2-1), (2-4), (2-37), (2-38), (3-8), (3-31), (3-50), (3-58), (3-60), (3-69), (3-112), (3-121), (3-132), (3-133), (3-173), (3-176), (3-207), (3-209), (3-218), (3-228), (3-238), (3-240), (3-252), (3-259), (3-274), (3-275), (3-286), (3-288), (3-295), (3-296), (3-297), (3-308), (3-318), (3-319), (3-322), (3-323), (3-329), (3-333), (3-334), (4-18), (4-21), (4-34), (4-50), (4-75), (4-105), (4-107), (4-118), and (4-126).

In yet another aspect, is a Formula (1) compound that is at least 100x selective against *D.immitis* L4 (i.e., DiMf/DiL4). In yet another aspect of the invention is a composition comprising a Formula (1) compound that is at least 100x selective against *D.immitis* L4 and a veterinary acceptable excipient. In yet another aspect of the invention is a composition comprising a therapeutically effective amount of a Formula (1) compound that is at least 100x selective over *D.immitis* L4. In yet another aspect of the invention is a composition comprising a therapeutically effective amount of a Formula (1) compound that is at least 100x selective over *D.immitis* L4 and a veterinary acceptable excipient. In another aspect, the Formula (1) compounds that are at least 100x selective include Formula (1A1) compounds: (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(3-methoxybenzyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (1-5) and (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(2-methoxybenzyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (1-11); and Formula (2A1) compounds: (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((diethylamino)(3,4-

difluorophenyl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (2-36) and (3S,6R,9S,12R,15S,18R,21S,24S)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-[[4-[(S)-morpholino-[4-(trifluoromethoxy)phenyl]-methyl]phenyl]methyl]-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,5,8,11,14,17,20,23-octaone (2-42); and Formula (3A) compounds:

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiophen-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-4),

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((3-methylthiophen-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-44),

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiophen-3-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-56),

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-chlorothiophen-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-87),

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((3-methoxythiophen-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-110),

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-fluoro-6-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-118),

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylthiophen-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-124),

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-bromothiophen-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-125),

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((5-methylfuran-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-157),

(3-165). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-ethoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-165), (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(1-fluoro-1-(6-methoxypyridin-2-yl)ethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-225), (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(1-(2-methoxypyridin-3-yl)ethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-261), (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-cyclopropylisoxazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-284), (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-(difluoromethyl)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-287), (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-fluoro-2-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-293), (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,5-difluoropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-324), and (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,3-difluoropyridin-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-326); (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-chloro-2-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-332); and a Formula (4A) compound: (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((2,4-dimethyl-1H-pyrrol-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclo-tetracosan-2,5,8,11,14,17,20,23-octaone; stereoisomers thereof, and veterinary acceptable salts thereof.

In yet another aspect of the invention, are Formula (2A1) compounds (2-1), (2-9), and (2-38); or Formula (3A) compounds (3-4), (3-8), (3-14), (3-17), (3-18), (3-21), (3-24), (3-36), (3-45), (3-50), (3-52), (3-56), (3-57), (3-58), (3-69), (3-73), (3-87), (3-112), (3-119), (3-120), (3-124), (3-126), (3-155), (3-193), (3-215), (3-253), (3-268), (3-273), (3-274), (3-283), (3-288), (3-290), (3-291), (3-294), (3-299), (3-303), (3-305), (3-306), (3-307), (3-310), (3-319), (3-321), (3-323), (3-324), (3-329), (3-333), and (3-334); or Formula (4A) compounds (4-30), (4-31), (4-32), (4-34), (4-40), (4-52), (4-63), (4-75), (4-105), (4-107), (4-109), (4-118), and (4-139); stereoisomers thereof, and veterinary acceptable salts thereof; all of which have HcL3 MED values $\leq 1\mu\text{M}$.

The compounds of the invention are intended to encompass racemic mixtures, specific stereoisomers, regio-isomers, and tautomeric forms of the Formula (1) compounds.

Detailed Description

Definitions

For purposes of the present invention, as described and claimed herein, the following terms and phrases are defined as follows:

“Additional veterinary agent(s)” as used herein, refers to other veterinary or veterinary compounds or products that provide a therapeutically effective amount of said agents that are useful for the treatment of a parasitic infection in an animal, as described herein.

“Alkoxy”, as used herein, refers to -O-alkyl, wherein the term alkyl is defined below. Non-limiting alkoxy examples include: methoxy, ethoxy, propoxy, isopropoxy, n-butoxy, n-pentoxo, 1-methylbutoxy, 1-ethylpropoxy, and the like.

“Alkyl”, as used herein, unless otherwise indicated, refers to saturated monovalent hydrocarbon alkane radicals of the general formula $\text{C}_n\text{H}_{2n+1}$. The alkane radical may be straight or branched and may be unsubstituted or substituted. For example, the term “C₁-C₆alkyl” refers to a monovalent, straight or branched aliphatic group containing 1 to 6 carbon atoms. In addition, the term “C₁-C₄alkyl” refers to a monovalent, straight or branched aliphatic group containing 1 to 4 carbon atoms, and is also described as C₁ (-C-), C₂ (-C-C-), C₃ (-C-C-C-), and C₄ (-C-C-C-C-) for the straight aliphatic group. Terms like -C-C-, -C-C-C- and the like, are synonymous with -CH₂CH₂-, -CH₂CH₂CH₂-, and the like, respectively. Non-exclusive examples of C₁-C₆ alkyl groups include, but are not limited to methyl, ethyl, propyl, isopropyl, sec-

butyl, t-butyl, n-propyl, n-butyl, i-butyl, s-butyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, neopentyl, 3,3-dimethylpropyl, 2-methylpentyl, hexyl, and the like. The alkyl group may be attached to the chemical moiety by any one of the carbon atoms of the aliphatic chain. Alkyl groups can be substituted as described herein. Further when used in compound words such as alkylphenyl, said alkyl group has the same meaning as herein defined and may be attached to the chemical moiety by any one of the carbon atoms of the aliphatic chain. Non-limiting examples include:

C₁alkylaryl which is -CH₂aryl (e.g., -CH₂phenyl); C₁alkylC₃cycloalkyl which is -CH₂cyclopropyl; C₁alkylheteroaryl which can be -CH₂pyrazole, -CH₂pyridine, -CH₂pyrimidine, and the like.

"Animal(s)", as used herein, unless otherwise indicated, refers to an individual animal that is a mammal or bird. Specifically, mammal refers to a vertebrate animal that is human and non-human, which are members of the taxonomic class Mammalia. Non-exclusive examples of non-human mammals include companion animals and livestock. Non-exclusive examples of a companion animal include: dog, cat, llama, and horse. Preferred companion animals are dog, cat, and horse. More preferred is dog. Non-exclusive examples of livestock include: swine, camel, rabbits, goat, sheep, deer, elk, bovine (cattle), and bison. Preferred livestock is cattle and swine. Specifically, bird refers to a vertebrate animal of the taxonomic class Aves. Birds are feathered, winged, bipedal, endothermic, and egg-laying. Non-exclusive examples of bird include, poultry (e.g., chicken, turkey, duck, and geese), all of which are also referred to herein as fowl.

"Aryl", as described herein, refers to a monovalent aromatic carbocyclic group of from 6 to 10 carbon atoms having a single ring or multiple fused rings. Aryl groups include, but are not limited to, phenyl, biphenyl, and naphthyl. Further when used in compound words such as alkylaryl, said alkyl and aryl groups have the same meaning as herein defined and may be attached to the chemical moiety by any one of the carbon atoms of the aliphatic chain. For example, C₀alkylaryl is aryl, (e.g., C₀alkylphenyl is phenyl), C₁alkylaryl is -CH₂aryl (e.g., -CH₂phenyl), C₂alkylaryl is -CH₂CH₂aryl (e.g., -CH₂CH₂phenyl), and the like. Aryls can be substituted as described herein.

"Chiral", as used herein, unless otherwise indicated, refers to the structural characteristic of a molecule that makes it impossible to superimpose it on its mirror image, (e.g., "R" and "S" enantiomers).

"Comprise(s)", as used herein, refers to an inclusive meaning, i.e., that it will be taken to mean an inclusion of not only the listed components it directly references, but also other non-specified components or elements. This rationale will also be used when the term "comprised" or "comprising" is used in relation to one or more Steps in a method or process. Contains is herein construed as being synonymous to comprise. The term "consisting of", and/or "consisting essentially of" has a non-inclusive meaning.

"Compounds of the present invention", as used herein, unless otherwise indicated, refers to compounds of Formula (1), stereoisomers thereof, and veterinary acceptable salts thereof. The phrase also refers to the sub-genus formulas of Formula (1) including: Formula (1A), (1A1), (2A), (2A1), (3A), and (4A), stereoisomers thereof, and veterinary acceptable salts thereof. Preferred compounds are Formula (3A) compounds.

"Cycloalkyl", as used herein, unless otherwise indicated, includes fully saturated or partially saturated carbocyclic alkyl moieties. Non-limiting examples of partially saturated cycloalkyls include: cyclopropene, cyclobutene, cycloheptene, cyclooctene, cyclohepta-1,3-diene, and the like. Preferred cycloalkyls are 3- to 6-membered saturated monocyclic rings including cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. The cycloalkyl group may be attached to the chemical moiety by any one of the carbon atoms within the carbocyclic ring. Cycloalkyl groups can be substituted with at least one substituent, as described herein. Further when used in compound words such as alkylcycloalkyl, said alkyl and cycloalkyl groups have the same meaning as herein defined and may be attached to the chemical moiety by any one of the carbon atoms of the aliphatic chain. Examples of C₁-C₆alkylC₃-C₆cycloalkyl include, methylcyclopropane (C₁alkylC₃cycloalkyl or -CH₂cyclopropane), ethylcyclopropane (C₂alkylC₃cycloalkyl or -CH₂CH₂cyclopropane), methylcyclobutane (C₁alkylC₄cycloalkyl or -CH₂cyclobutane), ethylcyclobutane (C₂alkylC₄cycloalkyl or -CH₂CH₂cyclobutane), methylcyclohexane (C₁alkylC₆cycloalkyl or -CH₂cyclohexane), and the like. Cycloalkyl moieties can be substituted as described herein.

"Halogen" or "halo", as used herein, unless otherwise indicated, refers to fluorine (F), chlorine (Cl), bromine (Br) and iodine (I). Preferred halo are fluorine, chlorine, and bromine. Further, when used in compound words such as "haloalkyl" or "haloalkoxy", said alkyl and alkoxy may be partially or fully substituted with

halogen atoms which may be the same or different and said alkyl and alkoxy groups have the same meaning as above and may be attached to the chemical moiety by any one of the carbon atoms of the aliphatic chain. Examples of haloalkyl include F_3C- , F_2CH- , $ClCH_2-$, CF_3CH_2- and CF_2CCl_2- , and the like. The term "haloalkoxy" is defined analogously to the term "haloalkyl". Examples of haloalkoxy include CF_3O- , Cl_3CH_2O- , $HCF_2CH_2CH_2O-$ and CF_3CH_2O- , and the like.

"Heteroaryl", as used herein, unless otherwise indicated, refers to a 5- to 6-membered aromatic monocyclic ring or an 9- to 10-membered fused aromatic ring where said monocyclic- and fused-ring contains one or more heteroatoms each independently selected from N, O, or S, preferably from one to four heteroatoms. Non-exclusive examples of monocyclic heteroaryls include pyrrolyl, furanyl, thiophenyl, pyrazolyl, imidazolyl, triazolyl, triazinyl, tetrazolyl, thiazolyl, isoxazolyl, isothiazolyl, oxazolyl, oxadiazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, dihydropyridinyl, and the like. Non-exclusive examples of fused heteroaryls include: benzofuranyl, oxazolo[5,4-b]pyridine, indolyl, 3,4-dihydro-2H-pyrrano[2,3-b]pyridine, 2,3-dihydrothieno[3,4-b]dioxine, benzo[d][1,3]dioxole, furo[2,3-b]pyridine, benzothiophenyl, benzimidazolyl, indazolyl, quinolinyl, isoquinolinyl, cinnolinyl, quinazolinyl, benzotriazolyl, quinoxaline, benzo[d]thiazolyl, thieno[2,3-c]pyridine, thieno[3,2-b]pyridine, benzo[1,2,5]thiadiazole, pyrazolo[1,5-a]pyridine, pyrazolo[1,5-a]pyrimidine, dihydro-1H-pyrrolo[3,2-b]pyridine, oxazolo[5,5-b]pyridine, and the like. The heteroaryl ring is attached to the chemical moiety by any one of the carbon atoms within the monocyclic or fused ring. Further when used in compound words such as alkylheteroaryl, said alkyl and heteroaryl groups have the same meaning as herein defined and may be attached to the chemical moiety by any one of the carbon atoms of the aliphatic chain. For example, C_1 alkylheteroaryl is $-CH_2$ heteroaryl, C_2 alkylheteroaryl is $-CH_2CH_2$ heteroaryl, and the like. Heteroaryls can be substituted as described herein.

"Heterocycle", as used herein, unless otherwise indicated, refers to a partially saturated or saturated 4- to 6-membered monocyclic ring or a 9- to 10-membered fused ring, each containing one or more heteroatoms independently selected from N, O, or S, preferably from one to four heteroatoms. Non-exclusive examples of heterocycle include oxetane, azetidine, thiatane, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, morpholine, thiomorpholine, tetrahydropyran, piperidine, piperazine, tetrahydropyridine, 2H-azirine, 2,3-dihydro-azete,

tetrahydrocyclopentapyrazole, 3,4-dihydro-2H-pyrrole, imidazolidine, oxazolidine, isoxazolidine, tetrahydropyrimidinyl, tetrahydropyridinyl, and the like. The heterocyclic ring is attached to the chemical moiety by any one of the carbon atoms within the ring. Further when used in compound words such as alkylheterocycle, said alkyl and heterocycle groups have the same meaning as herein defined and may be attached to the chemical moiety by any one of the carbon atoms of the aliphatic chain. For example, C₁alkylheterocycle is -CH₂heterocycle, C₂alkylheterocycle is -CH₂CH₂heterocycle, and the like. Heterocycles can be substituted as described herein.

“Parasite(s)”, as used herein, unless otherwise indicated, refers to endoparasites and ectoparasites. Endoparasites are parasites that live within the body of its host and include helminths (e.g., trematodes, cestodes, and nematodes) and protozoa. Ectoparasites are organisms of the Arthropoda phylum (e.g., arachnids, insects, and crustaceans (e.g., copepods-sea lice) which feed through or upon the skin of its host. Preferred arachnids are of the order Acarina, e.g., ticks and mites. Preferred insects are midges, fleas, mosquitos, biting flies (stable fly, horn fly, blow fly, horse fly, and the like), bed bugs, and lice. Preferred compounds of the present invention can be used for the treatment of parasites, i.e., treatment of a parasitic infection or infestation. A preferred parasite is an endoparasite. A preferred endoparasite is heart worm (*D.immitis*).

“Percent” (%), as used herein, refers to individual percent values. When referring to % in liquids (volume/volume% or v/v%) like an aqueous organic solvent, the % is the volume % of the solvent in the total volume of the solution. When referring to % for solids in liquids (weight/volume% or w/v%), the % value is construed to be the weight of the solid in the total volume of the solution and refers to the number of grams of solute in 100mL of solution. When referring to solids (weight% or w/w%) refers to the weight (mass) of one component relative to the total weight (mass) of the solid composition.

“Protecting group” or “Pg”, as used herein, unless otherwise indicated, refers to a substituent that is commonly employed to block or protect an amine on the compound thereby protecting its functionality while allowing for the reaction of other functional groups on the compound. Non-exclusive examples of an amine-protecting group include: acyl groups (e.g., formyl, acetyl, chloroacetyl, trichloro-acetyl, o-nitrophenylacetyl, o-nitrophenoxyacetyl, trifluoroacetyl, acetoacetyl, 4-chlorobutyryl,

isobutyryl, *o*-nitrocinnamoyl, picolinoyl, acylisothiocyanate, aminocaproyl, benzoyl, and the like), acyloxy groups (e.g., 1-*tert*-butyloxycarbonyl (Boc), methoxycarbonyl, 9-fluorenyl-methoxycarbonyl, 2,2,2-trifluoroethoxycarbonyl, 2-trimethylsilylethoxycarbonyl, vinyloxycarbonyl, allyloxycarbonyl, 1,1 –dimethyl-propynyloxycarbonyl, benzyloxy-carbonyl, *p*-nitrobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, and the like), diphenylmethane, and benzylcarbamates.

“Substituted”, as used herein, refers to a substituent that is bonded to the chemical moiety in place of a hydrogen atom. Common substituents of the invention for alkyl, cycloalkyl, aryl, heterocycle, heteroaryl moieties, and the like, are as described herein, and include, for example, -NO₂, -CN, alkyl, halo, alkoxy, haloalkyl, haloalkoxy, cycloalkyl, hydroxy, amino, alkylamino, dialkylamino, and the like. Substituents can be bonded to any carbon in the aliphatic chain or carbocyclic, aryl, heterocyclic, or heteroaryl ring system. The substituent can also be bonded to any accepting nitrogen and/or sulfur atom.

"Therapeutically effective amount", as used herein, refers to an amount of the active agent (i.e., Formula (1) compound) that (i) treat the particular parasitic infection or infestation, (ii) attenuates, ameliorates, or eliminates one or more symptoms of the particular parasitic infection or infestation, or (iii) prevents or delays the onset of one or more symptoms of the particular parasitic infection or infestation described herein. The active agent may be in a composition sufficient to elicit the desired biological response to the target parasite(s) after administration of the composition to the animal, as measured by methods known in the art and/or described in the examples herein. In some instances, an "effective amount" of the active agent in the composition will provide an efficacy of at least 70% against the target parasite compared to an untreated control. In other instances, "an effective amount" of the active agent will provide an efficacy of at least 80%, or at least 85% compared to untreated controls. More typically, "an effective amount" of the active agent will provide an efficacy of at least 90%, at least 93%, at least 95% or at least 97% against the target parasite. In certain instances, including the prevention of *Dirofilaria immitis*, the term "effective amount" may provide efficacy as high as 100%. As is understood in the art, a therapeutically effective amount may be in one or more doses, i.e., a single dose or multiple doses may be required to achieve the desired treatment endpoint, for example, (i) treat the particular parasitic infection or infestation, (ii) attenuates, ameliorates, or eliminates one or more symptoms of the

particular parasitic infection or infestation, or (iii) prevents or delays the onset of one or more symptoms of the particular parasitic infection or infestation described herein. The therapeutically effective amount will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

"Treatment", "treating", and the like, as used herein, unless otherwise indicated, refers to reversing, alleviating, or inhibiting the parasitic infection, infestation, or condition. As used herein, these terms also encompass, depending on the condition of the animal, preventing the onset of a disorder or condition, or of symptoms associated with a disorder or condition, including reducing the severity of a disorder or condition or symptoms associated therewith prior to affliction with said infection or infestation or after said infection or infestation. As such, compounds of the invention prevent heartworm in an animal by killing the L3 and L4 *D. immitis* before it can morph into an adult worm. Thus, treatment can refer to administration of the compounds of the present invention to an animal that is not at the time of administration afflicted with the infection or infestation. Treating also encompasses preventing the recurrence of an infection or infestation or of symptoms associated therewith as well as references to "control" (e.g., kill, repel, expel, incapacitate, deter, eliminate, alleviate, minimize, and eradicate).

"Veterinary acceptable" as used herein, unless otherwise indicated, indicates that the substance or composition must be compatible chemically and/or toxicologically, with the other ingredients comprising a formulation, composition, and/or the animal being treated therewith. The term "veterinary" acceptable has the same meaning as that recited for "pharmaceutical" acceptable.

The Formula (1) compounds of the invention are 24-membered cyclic depsipeptide compounds which have potent activity against endoparasites such as nematodes and filarial worms (microfilarial and larval stages) and also in some cases against ectoparasites such as fleas and ticks. In one aspect of the invention is a cyclic depsipeptide of Formula (1), or a veterinaryly acceptable salt thereof. Surprisingly, it has been found that addition of a methylene linker between the aryl ring in one or both of the phenyl lactate groups in the molecule versus the parent cyclic depsipeptide PF1022 and emodepside, improves the selectivity and activity of the compounds against parasites, particularly, endoparasites. This improvement provides selectivity against the L3 and L4 larvae

and the actual microfilaria by killing the L3 and L4 larvae before metamorphosis into the next lifecycle change, the animal is healthier and does not need to succumb to the killing of microfilaria or adult worms which can cause emboli and ultimately death to the host animal. Furthermore, it has been surprisingly found that substitution of the compounds of Formula (1) with certain L₁ and L₂ groups also significantly improves the in vitro metabolic stability of the compounds of the invention compared with PF1022 and emodepside. Thus, the compounds of the invention have been found to have significantly improved metabolic stability and equal or significantly improved efficacy against endoparasites including *Dirofilaria immitis* microfilaria and/or L3 and L4 larvae and/or *Haemonchus contortus* larvae. In some aspects, the compounds of Formula (1) with certain substituents will also exhibit improved activity against ectoparasites.

Compounds of the present invention may be synthesized by synthetic routes that include processes analogous to those well known in the chemical arts, particularly in light of the description contained herein. The starting materials are generally available from commercial sources such as Aldrich Chemicals (Milwaukee, Wis.) or are readily prepared using methods well known to those skilled in the art (e.g., prepared by methods generally described in Louis F. Fieser and Mary Fieser, "Reagents for Organic Synthesis", 1; 19, Wiley, New York (1967, 1999 ed.), or Beilsteins Handbuch der organischen Chemie, 4, Aufl. ed. Springer-Verlag, Berlin, including supplements (also available via the Beilstein online database)). For illustrative purposes, the reaction schemes depicted below demonstrate potential routes for synthesizing compounds of the present invention, and key intermediates. For a more detailed description of the individual reaction Steps, see the Examples section below. A skilled artisan will appreciate that other suitable starting materials, reagents, and synthetic routes may be used to synthesize the compounds of the present invention and a variety of derivatives thereof. Further, many of the compounds prepared by the methods described below can be further modified in light of this disclosure using conventional chemistry well known to the skilled artisan.

The depsipeptides of the present invention described herein, include one or more chiral centers which results in a theoretical number of optically active isomers. Where compounds of the invention include *n* chiral centers, the compounds may comprise up to 2^{*n*} optical isomers. The present invention encompasses the specific

enantiomers or diastereomers, and mixtures thereof, of each compound. The optically active forms can be prepared by, for example, resolution of the racemic forms by selective crystallization techniques, by synthesis from optically active precursors, by chiral synthesis, by chromatographic separation using a chiral stationary phase or by enzymatic resolution.

For illustrative purposes, the reaction schemes depicted below demonstrate potential routes for synthesizing key intermediates and compounds of the present invention. For a more detailed description of the individual reaction steps, see the Examples section below. Those skilled in the art will appreciate that other suitable starting materials, reagents, and synthetic routes may be used to synthesize the intermediates and compounds of the present invention and a variety of derivatives thereof. Further, many of the compounds prepared by the methods described below can be further modified in light of this disclosure using conventional chemistry. Schemes 1-9 outline the general procedures useful for the preparation and isolation of compounds of the present invention. It is to be understood, however, that the invention, as fully described herein and as recited in the claims, is not intended to be limited by the details of the following schemes or modes of preparation. In the preparation of compounds of the present invention, protection of remote functionality of intermediates from undesired reactions can be accomplished with a protecting group.

In the preparation of compounds of the present invention, protection of remote functionality of intermediates from undesired reactions can be accomplished with a protecting group. The term "protecting group" or "Pg" refers to a substituent that is commonly employed to block or protect a particular functionality while reacting other functional groups on the compound. For example, an amine-protecting group is a substituent attached to an amine that blocks or protects the amine-functionality of the compound or intermediate. Suitable amine protecting groups include: 1-*tert*-butyloxycarbonyl (Boc), acyl groups including: formyl, acetyl, chloroacetyl, trichloroacetyl, *o*-nitrophenylacetyl, *o*-nitrophenoxyacetyl, trifluoroacetyl, acetoacetyl, 4-chlorobutyryl, isobutyryl, *o*-nitrocinnamoyl, picolinoyl, acylisothiocyanate, aminocaproyl, benzoyl, and the like; and acyloxy groups including: methoxycarbonyl, 9-fluorenyl-methoxycarbonyl, 2,2,2-trifluoroethoxycarbonyl, 2-trimethylsilylethoxycarbonyl, vinyloxycarbonyl, allyloxycarbonyl, 1,1 –dimethyl-propynyloxycarbonyl, benzyloxy-carbonyl, *p*-nitrobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, and

the like. Similarly, diphenylmethane and benzylcarbamates can be used as amine protecting groups. Suitable protecting groups and their respective uses are readily determined by one skilled in the art. For a general description of protecting groups and their use, see T. W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, 1991.

The compounds of the invention are intended to encompass racemic mixtures, specific stereoisomers and tautomeric forms of the compound. Another aspect of the invention is a salt form of the compound of the invention.

The compounds of present invention may also be present in different solid forms such as different crystalline forms or in the form of an amorphous solid. The present invention encompasses different crystalline forms as well as amorphous forms of the inventive compounds.

In addition, the compounds of the invention may exist as hydrates or solvates, in which a certain stoichiometric amount of water or a solvent is associated with the molecule in the crystalline form. The hydrates and solvates of the compounds of formula (I) are also the subject of the invention.

In addition to the neutral compounds of Formula (1), salt forms of the compounds are also active against endoparasites. The term "veterinary acceptable salt" is used throughout the specification to describe any salts of the compounds that are acceptable for administration for veterinary applications, and which provides the active compound upon administration.

In cases where compounds are sufficiently basic or acidic to form stable non-toxic acid or base salts, the compounds may be in the form of a veterinary or agriculturally acceptable salt. Veterinary acceptable salts include those derived from veterinary or agriculturally acceptable inorganic or organic bases and acids. Suitable salts include those comprising alkali metals such as lithium, sodium or potassium, alkaline earth metals such as calcium, magnesium and barium. Salts comprising transition metals including, but not limited to, manganese, copper, zinc and iron are also suitable. In addition, salts comprising ammonium cations (NH_4^+) as well as substituted ammonium cations, in which one or more of the hydrogen atoms are replaced by alkyl or aryl groups are encompassed by the invention.

Salts derived from inorganic acids including, but not limited to, hydrohalide acids (HCl, HBr, HF, HI), sulfuric acid, nitric acid, phosphoric acid, and the like are particularly suitable. Suitable inorganic salts also include, but not limited to, bicarbonate, and carbonate salts. In some embodiments, examples of veterinary and agriculturally acceptable salts are organic acid addition salts formed with organic acids including, but not limited to, maleate, dimaleate, fumarate, tosylate, methanesulfonate, acetate, citrate, malonate, tartarate, succinate, benzoate, ascorbate, α -ketoglutarate, and α -glycerophosphate. Of course, other acceptable organic acids may be used.

Alkali metal (for example, sodium, potassium or lithium) or alkaline earth metal (for example calcium) salts of the compounds can also be made by reacting a sufficiently acidic residue on the compounds with a hydroxide of the alkali metal or alkaline earth metal.

Veterinarily acceptable salts may be obtained using standard procedures well known in the art, for example by reacting a sufficiently basic compound such as an amine with a suitably acid functional group present in the compound, or by reacting a suitable acid with a suitably basic functional group on the compound of the invention.

The compounds of Formula (1) may be prepared by processes adapted from those described in US Patent Nos. 5,514,773; 5,747,448; 5,874,530; 5,856,436; 6,033,879; 5,763,221; 6,329,338, 5,116,815; 6,468,966; 6,369,028; 5,777,075; and 5,646,244. In addition, various synthetic methods for cyclic depsipeptides have been reported in the chemical literature (see Luttenberg *et al.*, *Tetrahedron* 68 (2012), 2068-2073; Byung H. Lee, *Tetrahedron Letters*, 1997, 38 (5), 757-760; Scherkenbeck *et al.*, *Eur. J. Org. Chem.*, 2012, 1546-1553; *Biosci. Biotech. Biochem.*, 1994, 58(6), 1193-1194; and Scherkenbeck *et al.*, *Tetrahedron*, 1995, 51(31), 8459-8470) It will be understood by those skilled in the art that certain functional groups in the compounds and intermediates may be unprotected or protected by suitable protecting groups, as taught by Greene and Wuts, *Protective Groups in Organic Synthesis*, John Wiley and Sons, Inc., 4th edition 2006. Further, it will be apparent to those skilled in the art that the compounds and intermediates may be isolated by standard aqueous work-up conditions and optionally purified. For example, the compounds or intermediates

may be purified by chromatographic methods or crystallized to yield the desired product in suitable purity.

One skilled in the art will recognize that, in some cases, after the introduction of a given reagent as it is depicted in the schemes, it may be necessary to perform additional routine synthetic Steps not described in detail to complete the synthesis of Formula (1) compounds.

The present invention includes all veterinary acceptable isotopically-labelled Formula (1) compounds wherein one or more atoms are replaced by atoms having the same atomic number, but an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes suitable for inclusion in the compounds of the present invention include isotopes of hydrogen, such as ^2H and ^3H , carbon, such as ^{11}C , ^{13}C and ^{14}C , chlorine, such as ^{36}Cl , fluorine, such as ^{18}F , iodine, such as ^{123}I and ^{125}I , nitrogen, such as ^{13}N and ^{15}N , oxygen, such as ^{15}O , ^{17}O and ^{18}O , and sulphur, such as ^{35}S .

The skilled person will appreciate that the compounds of the present invention could be made by methods other than those herein described as incorporated herein by reference, by adaptation of the methods herein described and/or adaptation of methods known in the art, for example the art described herein, or using standard textbooks such as "Comprehensive Organic Transformations - A Guide to Functional Group Transformations", RC Larock, Wiley-VCH (1999 or later editions).

The Formula (1) compounds are useful as antiparasitic agents, therefore, another aspect of the present invention is a veterinary composition comprising a therapeutically effective amount of a Formula (1) compound, stereoisomers thereof, and at least one veterinary acceptable excipient. The compounds of the present invention (including the compositions and processes used therein) may also be used in the manufacture of a medicament for the therapeutic applications described herein.

The compound of the present invention can be administered alone or in a formulation appropriate to the specific use envisaged, the particular species of host animal being treated and the parasite involved. Generally, it will be administered as a formulation in association with at least one veterinary acceptable excipient. The term "excipient" is used herein to describe any ingredient (e.g., carrier, diluents, and the like) other than the compound of the present invention or any additional veterinary (e.g., antiparasitic) agent. The choice of excipient will to a large extent

depend on factors such as the particular mode of administration, the effect of the excipient on solubility and stability, and the nature of the dosage form. In addition to the excipient(s), the amount of the compound of the present invention that is administered and the dosage regimen for treating a condition or disorder with the compound depends on a variety of factors, including the age, weight, sex and medical condition of the animal, the severity of the disease, the route and frequency of administration, and thus may vary widely.

In another aspect, the veterinary composition comprises a Formula (1) compound with at least one veterinary acceptable excipient. The concentration range will vary depending on the composition (e.g., oral, topical, or injectable). For an oral dose, the range of active (i.e., compound of the present invention) is about 0.1 to 50 mg/kg, preferably from about 0.2 to 25 mg/kg, and even more preferably from about 0.25 to 10mg/kg, and most preferably from about 0.5 to 7 mg/kg or 1-5 mg/kg. For a topical solution, the range of active is about 0.1 to 1000 mg/mL, and preferably from about 0.5 to 500 mg/mL, and more preferably from about 1 to 250 mg/mL, and even more preferably from about 2 to 200 mg/mL. Depending upon the final volumes of the topical solution(s), the concentration of the active can change from that described above. Generally, injectable doses tend to be, but not always, lower in concentration.

The formulations can be prepared using conventional dissolution and mixing procedures. Such compositions and methods for their preparation may be found, for example, in 'Remington's Veterinary Sciences', 19th Edition (Mack Publishing Company, 1995; and "Veterinary Dosage Forms: Tablets, Vol. 1", by H. Lieberman and L. Lachman, Marcel Dekker, N.Y., 1980 (ISBN 0-8247-6918-X).

A typical formulation is prepared by mixing a Formula (1) compound with at least one veterinary acceptable excipient. Suitable excipients are well known to those skilled in the art and include materials such as carbohydrates, waxes, water soluble and/or swellable polymers, starches, hydrophilic or hydrophobic materials, gelatin, oils, solvents, water, and the like. The particular excipient(s) will depend upon the means and purpose for which the compound of the present invention is being applied. Solvents are generally selected based on solvents recognized by persons skilled in the art as safe to be administered to an animal. The formulations may also include one or more buffers, stabilizing agents, surfactants, wetting agents, lubricating agents, emulsifiers, suspending agents, preservatives, antioxidants,

opaquing agents, glidants, processing aids, colorants, sweeteners, flavoring agents and other known additives to provide an elegant presentation of the drug (i.e., a compound of the present invention or veterinary composition thereof) or aid in the manufacturing of the veterinary product (i.e., medicament). The compound of the present invention will typically be formulated into veterinary dosage forms to provide an easily controllable dosage form for administration.

The methods by which the compound of the present invention may be administered include oral, topical, and injectable (e.g., parenteral and subcutaneous) administration. The particular route selected by the practitioner depends upon factors such as the physicochemical properties of the therapeutic agent, the condition of the host and economics. In certain cases, it is convenient and efficient to administer veterinary medicines orally by placing the therapeutic agent in a solid or liquid matrix that is suitable for oral delivery. These methods include chewable drug-delivery formulations. The problem associated with administering oral formulations to animals is that the therapeutic agent often provides an unpleasant taste, aroma, or texture, which causes the animals to reject the composition. This is further exacerbated by compositions that are hard and difficult to swallow.

The compound of the present invention can be administered orally by capsule, bolus, tablet, powders, lozenges, chews, multi and nanoparticulates, gels, solid solution, films, sprays, or liquid form. This is a preferred method of administration and as such it is desirable to develop the compound for oral administration. Such formulations may be employed as fillers in soft or hard capsules, soft or hard palatable chews, which typically comprise at least one veterinary acceptable excipient, for example, water, ethanol, polyethylene glycol, N-methylpyrrolidone, propylene glycol, methylcellulose, or a suitable oil, and one or more emulsifying agents, flavorants, and/or suspending agents. Liquid forms include suspensions, solutions, syrups, drenches and elixirs. Liquid formulations may also be prepared by the reconstitution of a solid, for example, from a sachet. Oral drenches are commonly prepared by dissolving or suspending the compound of the present invention in a suitable medium (e.g. triethylene glycol, benzyl alcohol, and the like). The compound of the present invention can also be formulated with a food substance, e.g., a dietary admixture (food pellets or powder for birds).

The compound of the present invention can be administered topically to the skin or mucosa, that is dermally or transdermally. This is another preferred method

of administration and as such it is desirable to develop the compound of the present invention to be suited to such formulations, for example liquid forms. Typical formulations for this purpose include pour-on, spot-on, multi-spot-on, stripe-on, comb-on, roll-on, dip, spray, mousse, shampoo, powder formulation, gels, hydrogels, lotions, solutions, creams, ointments, dusting powders, dressings, foams, films, skin patches, wafers, implants, sponges, fibers, bandages and micro emulsions. Liposomes may also be used. Typical excipients include alcohol, water, mineral oil, liquid petrolatum, white petrolatum, glycerin, N-methyl formamide, glycol monomethyl ethers, polyethylene glycol, propylene glycol, and the like. Penetration enhancers may be incorporated - see, for example, J Pharm Sci, 88 (10), 955-958 by Finnin and Morgan (October 1999). Pour-on or spot-on formulations may be prepared by dissolving the active ingredients in an acceptable liquid excipient such as butyl digol, liquid paraffin or a non-volatile ester, optionally with the addition of a volatile component such as propan-2-ol or a glycol ether. Alternatively, pour-on, spot-on or spray formulations can be prepared by encapsulation, to leave a residue of active agent on the surface of the animal, this effect may ensure that the compound of the present invention has increased persistence of action and is more durable, for example it may be more water-fast. Topical formulations contemplated herein can comprise from about 0.1 mg/kg to 50 mg/kg of a compound of the present invention, and more preferably from about 1 mg/kg to 10 mg/kg of a compound of the present invention, and even more preferably, from 1mg/kg to 5 mg/kg.

The compounds of the present invention can also be administered topically via a support matrix for example, a synthetic or natural resin, plastic, cloth, leather, or other such polymeric system in the shape of a collar or ear tag. Said collar or ear tag may be coated, impregnated, layered, by any means so as to provide a veterinary acceptable amount of a compound of the present invention alone, or with at least one veterinary acceptable excipient, and optionally an additional antiparasitic agent, or veterinary acceptable salts thereof. Such formulations are prepared in a conventional manner in accordance with standard medicinal or veterinary practice. Further, these formulations will vary with regard to the weight of active compound contained therein, depending on the species of host animal to be treated, the severity and type of infection or infestation, and the body weight of the animal. The volume of the applied composition can be from about 0.2 mL/kg to 5 mL/kg and preferably from about 1 mL/kg to 3mL/kg.

Agents may be added to the formulations of the present invention to improve the persistence of such formulations on the surface of the animal to which they are applied, for example to improve their persistence on the coat of the animal. It is particularly preferred to include such agents in a formulation which is to be applied as a pour-on or spot-on formulation. Examples of such agents include acrylic copolymers and in particular fluorinated acrylic copolymers. A particular suitable reagent is the trademark reagent "Foraperle" (Redline Products Inc, Texas, USA). Certain topical formulations may include unpalatable additives to minimize oral exposure.

The compounds of the present invention can also be administered by injection. Injectable (e.g., subcutaneous and parenteral) formulations may be prepared in the form of a sterile solution, which may contain other substances, for example enough salts or glucose to make the solution isotonic with blood. Acceptable liquid excipients include vegetable oils such as sesame oil, glycerides such as triacetin, esters such as benzyl benzoate, isopropyl myristate and fatty acid derivatives of propylene glycol, as well as organic solvents such as pyrrolidin-2-one and glycerol formal. The formulations are prepared by dissolving or suspending compounds of the present invention alone or with at least one additional antiparasitic agent in the liquid excipient such that the final formulation contains from about 0.01 to 30% by weight of the active ingredients.

Suitable devices for injectable administration include needle (including micro needle) injectors, needle-free injectors and infusion techniques. Injectable formulations are typically aqueous solutions which may contain excipients such as salts, carbohydrates and buffering agents (preferably to a pH of from 3 to 9), but, for some applications, they may be more suitably formulated as a sterile non-aqueous solution or as a dry powder form to be used in conjunction with a suitable vehicle such as sterile, pyrogen-free water. The preparation of injectable formulations under sterile conditions, for example, by lyophilization, may readily be accomplished using standard veterinary techniques well known to those skilled in the art. The solubility of a compound of the present invention used in the preparation of an injectable solution may be increased by the use of appropriate formulation techniques, such as the incorporation of solubility-enhancing agents.

Administration of the compound of the instant invention is contemplated to be once a month. However, an extended duration formulation may allow for dosing once every 2, 3, 4, 5, or 6 months.

Such formulations are prepared in a conventional manner in accordance with standard medicinal or veterinary practice. Further, these formulations will vary with regard to the weight of active compound contained therein, depending on the species of host animal to be treated, the severity and type of infection or infestation, and the body weight of the animal.

The composition of the present invention may be administered alone, as described above, or in combination with at least one other additional antiparasitic agent to form a multi-component parasiticide giving an even broader spectrum of veterinary utility. Thus, the present invention also envisions a combination veterinary composition comprising an effective amount of the compound of the present invention in combination with at least one other additional antiparasitic agent and can further comprise at least one veterinary acceptable excipient.

The following list of additional antiparasitic agents together with which the compound of the present invention can be used is intended to illustrate the possible combinations, but not to impose any limitation. Non-limiting examples of additional antiparasitic agents include: amitraz, aminoacetonitriles, albendazole, cambendazole, fenbendazole, flubendazole, thiabendazole, mebendazole, cyclic octadepsipeptides, oxfendazole, oxibendazole, paraherquamide, parbendazole, piperazines, praziquantel, thiabendazole, tetramisole, triclabendazole, levamisole, pyrantel (including the salt forms - pamoate, citrate, and tartrate), oxantel, morantel, abamectin, doramectin, emamectin, eprinomectin, ivermectin, moxidectin, selamectin, dimadectin, latidectin, lepimectin, milbemycin, milbemycin oxime, demiditraz, emodepside, fipronil, methoprene, diethylcarbamazine, hydroprene, kinoprene, lufenuron, metaflumizone, niclosamide, permethrin, pyrethrins, pyriproxyfen, closantel, clorsulon, novaluron, fluazuron, spinosad, sarolaner ((S)-1-(5'-(5-(3,5-dichloro-4-fluorophenyl)-5-(trifluoromethyl)-4,5-dihydroisoxazol-3-yl)-3'H-spiro[azetidine-3,1'-isobenzofuran]-1-yl)-2-(methylsulfonyl)-ethan-1-one), fluralaner (4-(5-(3,5-dichlorophenyl)-5-(trifluoromethyl)-4,5-dihydro-isoxazol-3-yl)-2-methyl-N-(2-oxo-2-((2,2,2-trifluoroethyl)amino)-ethyl)benzamide), afoxolaner (4-(5-(3-chloro-5-(trifluoromethyl)phenyl)-5-(trifluoromethyl)-4,5-dihydro-isoxazol-3-yl)-N-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-1-naphthamide), lotilaner (3-methyl-N-{2-oxo-2-

[(2,2,2-trifluoroethyl)amino]ethyl}-5-[(5S)-5-(3,4,5-trichlorophenyl)-5-(trifluoromethyl)-4,5-dihydro-1,2-oxazol-3-yl]thiophene-2-carboxamide); and mixtures thereof.

Preferred additional antiparasitic agents include moxidectin, doramectin, selamectin, abamectin, milbemycin, milbemycin oxime, pyrantel, praziquatel, sarolaner, afoxolaner, lotilaner, fluralaner, and levamisole.

The veterinary composition for application to an animal may be packaged in a variety of ways depending upon the method used for administering the compound of the present invention or combination, thereof. Generally, an article for distribution includes a container having deposited therein the veterinary composition in an appropriate form. Suitable containers are well-known to those skilled in the art and include materials such as bottles (plastic and glass), sachets, ampoules, plastic bags, metal cylinders, and the like. The container may also include a tamper-proof assemblage to prevent indiscreet access to the contents of the package. In addition, the container has deposited thereon a label that describes the contents of the container. The label may also include appropriate warnings.

The compounds of the present invention (including the compositions and processes used therein) may also be used in the manufacture of a medicament for the therapeutic applications described herein.

The compounds of the present invention, stereoisomers thereof, and compositions comprising a therapeutically effective amount of a Formula (1) compound and at least one veterinary acceptable excipient are useful as ectoparasitocides for the control and treatment of infections or infestations manifested by said ectoparasite in an animal. The compounds of the present invention have utility as an ectoparasiticide, in particular, as an acaricide and insecticide. They may, in particular, be used in the fields of veterinary medicine, livestock husbandry and the maintenance of public health: against acarids, insects, and copepods which are parasitic upon vertebrates, particularly warm-blooded vertebrates, including companion animals, livestock, and fowl and cold-blooded vertebrates like fish.

The compounds of the present invention are useful for the treatment of parasitic worms categorized as cestodes (tapeworms), nematodes (roundworms) and trematodes (flatworms or flukes). The gastrointestinal roundworms include, for example, *Ostertagia ostertagi* (including inhibited larvae), *O. lyrata*, *Haemonchus placei*, *H. similis*, *H. contortus*, *Toxocara canis*, *T. leonina*, *T. cati*, *Trichostrongylus*

axei, *T. colubriformis*, *T. longispicularis*, *Cooperia oncophora*, *C. pectinata*, *C. punctata*, *C. surnabada* (syn. *mcmasteri*), *C. spatula*, *Ascaris suum*, *Hyostrongylus rubidus*, *Bunostomum phlebotomum*, *Capillaria bovis*, *B. trigonocephalum*, *Strongyloides papillosus*, *S. ransomi*, *Oesophagostomum radiatum*, *O. dentatum*, *O. columbianum*, *O. quadrispinulatum*, *Trichuris* spp., and the like. Other parasites include: hookworms (e.g., *Ancylostoma caninum*, *A. tubaeforme*, *A. braziliense*, *Uncinaria stenocephala*); lungworms (e.g., *Dictyocaulus viviparus* and *Metastrongylus* spp.); eyeworms (e.g., *Thelazia* spp.); parasitic stage grubs (e.g., *Hypoderma bovis*, *H. lineatum*, *Dermatobia hominis*); kidneyworms (e.g., *Stephanurus dentatus*); screw worm (e.g., *Cochliomyia hominivorax* (larvae)); filarial nematodes of the super-family *Filarioidea* and the *Onchocercidae* Family. Non-limiting examples of filarial nematodes within the *Onchocercidae* Family include the genus *Brugia* spp. (i.e., *B. malayi*, *B. pahangi*, *B. timori*, and the like), *Wuchereria* spp. (i.e., *W. bancrofti*, and the like), *Dirofilaria* spp. (*D. immitis*, *D. repens*, *D. ursi*, *D. tenuis*, *D. spectans*, *D. lutrae*, and the like), *Dipetalonema* spp. (i.e., *D. reconditum*, *D. repens*, and the like), *Onchocerca* spp. (i.e., *O. gibsoni*, *O. gutturosa*, *O. volvulus*, and the like), *Elaeophora* spp. (*E. bohmi*, *E. elaphi*, *E. poeli*, *E. sagitta*, *E. schneideri*, and the like), *Mansonella* spp. (i.e., *M. ozzardi*, *M. perstans*, and the like), and *Loa* spp. (i.e., *L. loa*). In another aspect of the invention, the compound of the present invention is useful for treating endoparasiticide infection from filarial nematodes within the genus *Dirofilaria* (i.e., *D. immitis*, *D. repens*, *D. ursi*, *D. tenuis*, and the like). The Formula (1) compounds are preferably endoparasitics.

The compounds of the present invention can also be used against ectoparasites, alone or in combination with at least one additional antiparasitic agent. Some non-limiting examples of ectoparasites include: ticks (e.g., *Ixodes* spp., (e.g., *I. ricinus*, *I. hexagonus*), *Rhipicephalus* spp., (e.g., *R. sanguineus*), *Boophilus* spp., *Amblyomma* spp. (e.g., *A. maculatum*, *A. triste*, *A. parvum*, *A. cajennense*, *A. ovale*, *A. oblongoguttatum*, *A. aureolatum*, *A. cajennense*), *Hyalomma* spp., *Haemaphysalis* spp., *Dermacentor* spp. (e.g., *D. variabilis*, *D. andersoni*, *D. marginatus*), *Ornithodoros* spp., and the like); mites (e.g., *Dermanyssus* spp., *Sarcoptes* spp., (e.g., *S. scabiei*), *Psoroptes* spp., (e.g., *P. bovis*), *Otodectes* spp., *Chorioptes* spp., *Demodex* spp., (e.g., *D. folliculorum*, *D. canis*, and *D. brevis*) and the like); chewing and sucking lice (e.g., *Damalinia* spp., *Linognathus* spp., *Cheyletiella* spp., *Haematopinus* spp., *Solenoptes* spp., *Trichodectes* spp., *Felicola*

spp., and the like); fleas (*e.g.*, *Siphonaptera spp.*, *Ctenocephalides spp.*, and the like); biting flies, midges, and mosquitos (*e.g.*, *Tabanidae spp.*, *Haematobia spp.*, *Musca spp.*, *Stomoxys spp.*, *Dematobia spp.*, *Cochliomyia spp.*, *Simuliidae spp.*, *Ceratopogonidae spp.*, *Psychodidae spp.*, *Aedes spp.*, *Culex spp.*, *Anopheles spp.*, and the like); bed bugs (*e.g.*, insects within the genus *Cimex* and family *Cimicidae*); and grubs (*e.g.*, *Hypoderma bovis*, *H. lineatum*); and copepods (*e.g.*, sea lice within the Order *Siphonostomatoida*, including genera *Lepeophtheirus* and *Caligus*).

The compounds of the present invention and compositions comprising compounds of the present invention in conjunction with at least one other antiparasitic agent are of particular value in the control of ectoparasites and endoparasites which are injurious to, or spread or act as vectors of diseases in companion animals, livestock, birds, and fish. The ectoparasites and endoparasites which can be treated with a combination of a Formula (1) compound and an additional antiparasitic agent include those as herein before described.

Any of the compounds of the present invention, or a suitable combination of a compound of the present invention and optionally, with at least one additional antiparasitic agent may be administered directly to the animal and/or indirectly by applying it to the local environment in which the animal dwells (such as bedding, enclosures, and the like). Direct administration includes contacting the skin, fur, or feathers of a subject animal with the compound(s), or by feeding or injecting the compounds into the animal.

The Formula (1) compound, stereoisomers thereof, veterinary acceptable salts thereof, and combinations with at least one additional antiparasitic agent, as described herein, are of value for the treatment and control of the various lifecycle stages of parasites including egg, nymph, larvae, juvenile and adult stages.

The present invention also relates to a method of administering a compound of the present invention alone or in combination with at least one additional antiparasitic agent to animals in good health comprising the application to said animal to reduce or eliminate the potential for human parasitic infection or infestation from parasites carried by the animal and to improve the environment in which the animals inhabit.

The present invention also relates to a method of administering a compound of the present invention alone or in combination with at least one additional antiparasitic agent, and optionally at least one veterinary acceptable excipient, to a

human in good or poor health comprising the application to said human to reduce or eliminate the potential for human parasitic infection or infestation from parasites carried by the human and to improve the environment in which the human inhabits.

The present invention also relates to a method of administering a compound of the present invention alone or in combination with at least one additional antiparasitic agent, and optionally at least one veterinary acceptable excipient, to a plant or soil to prevent parasitic infection to the plant.

The reactions set forth below were done generally under a positive pressure of argon or nitrogen or with a drying tube, at ambient temperature (unless otherwise stated), in anhydrous solvents, and the reaction flasks were fitted with rubber septa for the introduction of substrates and reagents via syringe. Glassware was oven dried and/or heat dried. Analytical thin layer chromatography (TLC) was performed using glass-backed silica gel 60 F 254 precoated plates and eluted with appropriate solvent ratios (v/v). Reactions were assayed by TLC or LCMS and terminated as judged by the consumption of starting material. Visualization of the TLC plates was done with UV light (254 nm wavelength) or with an appropriate TLC visualizing solvent and activated with heat. Flash column chromatography (Still et al., J. Org. Chem. 43, 2923, (1978) was performed using silica gel (RediSep Rf) or various MPLC systems, such as Biotage or ISCO purification system.

Conventional methods and/or techniques of separation and purification known to one of ordinary skill in the art can be used to isolate the compounds of the present invention, as well as the various intermediates related thereto. Such techniques will be well-known to one of ordinary skill in the art and may include, for example, all types of chromatography (e.g., high pressure liquid chromatography (HPLC), column chromatography using common adsorbents such as silica gel, and thin-layer chromatography (TLC), recrystallization, and differential (*i.e.*, liquid-liquid) extraction techniques.

The compound structures in the examples below were confirmed by one or more of the following methods: proton magnetic resonance spectroscopy, and mass spectroscopy. Proton magnetic resonance (¹H NMR) spectra were determined using a Bruker spectrometer operating at a field strength of 400 megahertz (MHz). Chemical shifts are reported in parts per million (ppm) downfield from an internal tetramethylsilane standard. Mass spectra (MS) data were obtained using Agilent mass spectrometer with atmospheric pressure chemical ionization. Method: Acquity

UPLC with chromatography performed on a Waters BEH C18 column (2.1 x 50 mm, 1.7 μ m) at 50°C. The mobile phase was a binary gradient of acetonitrile (containing 0.1% trifluoroacetic acid) and water (5-100%).


Certain aspects of the present invention are illustrated by the following Examples. It is to be understood, however, that the aspects of the invention are not limited to the specific details of these Examples, as other variations thereof will be known, or apparent in light of the instant disclosure, to one of ordinary skill in the art.

Compounds of this invention can exist as one or more stereoisomers. One skilled in the art will appreciate that one stereoisomer may be more active and/or may exhibit beneficial effects when enriched relative to the other stereoisomer(s) or when separated from the other stereoisomer(s). Additionally, the skilled artisan knows how to separate, enrich, and/or to selectively prepare said stereoisomers. The compounds of the invention may be present as a mixture of stereoisomers.


EXAMPLES

The following examples were prepared according to the Schemes and Preparations as presented herein.

PF1022a: (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-dibenzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone




Emodepside: (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-morpholinobenzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone



In the Schemes and Examples described below, the following catalysts/reactants and miscellaneous abbreviations include: room temperature (RT); dichloromethane (DCM); methanol (MeOH); dimethylformamide (DMF); ethyl acetate (EtOAc); propylphosphonic anhydride (T₃P); acetonitrile (MeCN or ACN); ethanol (EtOH); tributyltin hydride (HSnBu₃); tributyltin chloride (Bu₃SnCl); sodium borohydride (NaBH₄); sodium nitrite (NaNO₂); sulfuric acid (H₂SO₄); zinc chloride (ZnCl₂); tetrahydrofuran (THF); cesium carbonate (CsCO₃); denatured alcohol or industrial methylated spirits (IMS); trifluoromethyltrimethylsilane (CF₃TMS); 4-dimethylaminopyridine (DMAP); t-butyloxycarbonyl (BOC, boc); palladium (Pd); N,N-diisopropylethylamine (DIPEA); 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDCI); methoxymethyl (MOM); sodium azide (NaN₃); tetra-n-butylammonium fluoride (TBAF); benzyl bromide (BnBr); tetrakis(triphenylphosphine)palladium (0) (Pd(PPh₃)₄); FurCat (succinimide-containing Pd catalyst: Pd(N-succ)Br(P(2-Fu)₃)₂); isopropylmagnesium chloride lithium chloride (iPrMgCl-LiCl); hour (h, hr); minute(s) (min); bis(triphenylphosphine)palladium(II) dichloride (PdCl₂(PPh₃)₂), Ghosez Reagent (1-chloro-N,N,2-trimethyl-1-propenylamine; Dess-Martin (periodinane); iodomethane (MeI); saturated (sat); aqueous (aq); equivalents (eq or equiv); 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC); hydrochloric acid (HCl); dichloromethane (DCM); dichloroethane (DCE); N-methylmorpholine (NMM); palladium on carbon (Pd-C or Pd/C); triphenylphosphine (PPh₃); di-tert-butyl azodicarboxylate; trifluoroacetic acid (TFA); 1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate (HATU); hydrogen bromide (HBr); acetic acid (AcOH); p-toluenesulfonic acid (PTSA); triethylsilane (Et₃SiH); copper iodide (CuI); potassium iodide (KI); methylene chloride (CH₂Cl₂); triethanolamine (TEA); methyl tert-butyl ether (MTBE); tetramethylsilane (TMS); trimethylsilyl cyanide (TMSCN); diethylaminosulfur trifluoride (DAST); t-butyl nitrite

(TBN or tBuONO); sodium hydride (NaH); tetrakis(triphenylphosphine)palladium (0) ($\text{Pd}(\text{TPP})_4$); tetra-n-butylammonium fluoride (nBu_4F); di-tert-butyl azodicarboxylate (DBAD); and phosphorus tribromide (PBr_3).

Scheme 1: Synthetic Scheme for Preparation of Monomers M1, M2, M3 and M4.



In Scheme 1, synthesis of monomers **M1**, **M2**, **M3** and **M4** was carried out by procedures well known in the literature. (Journal of Organic Chemistry, 79(17), 8491-8497; 2014; Organic Letters, 15(24), 6132-6135; 2013; ChemBioChem, 9(8), 1235-1242; 2008) These monomers were used in a stepwise synthesis of the PF1022a core as shown in subsequent schemes. In addition, the allyl protected monomers, allyl (S)-2-hydroxypropanoate and allyl (R)-2-hydroxy-3-(4-nitrophenyl)propanoate and allyl (R)-2-hydroxy-3-phenylpropanoate were also prepared according to the following literature procedures: Faming Zhuanli Shenqing, 101962323, 02Feb2011 and Journal of Organic Chemistry, 67(4), 1061-1070; 2002). These allyl protected monomers were used in an alternative route which also provided Bis-Iodo PF1022a using the same coupling conditions described below and the requisite chemistry to remove the allyl protecting group as described in T. W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, 1991.

Synthesis of Boc-Methyl-L-Leucine (**M1**):

Sodium hydride (65.16g, 1.63mol) was added portion-wise over 1.25 hours to a mixture of Boc-L-leucine(125g, 0.50mol) and iodomethane (160mL, 2.50mol) in THF (2L) cooled to 0°C. The temperature was maintained below 5°C during the addition and then allowed to warm up to room temperature and stirred for 2.5 days. The reaction mixture was cooled to 0°C and quenched with water (2L); the temperature was maintained below 5°C during the addition and then allowed to warm up to room temperature. The aqueous layer was extracted with EtOAc (2 × 750mL), then the aqueous was acidified to pH 5 with 10% aqueous citric acid solution, extracted with EtOAc (3×1L), dried over MgSO₄, filtered and the solvent removed *in vacuo* (50°C) and azeotroped with DCM. The two batches were combined to yield 236.09 g, **M1**, 94% yield. ¹H NMR (CDCl₃, 300 MHz): δ 4.85 (t, 0.5H), 4.61 (dd, 0.5H), 2.81 (s, 1.5H), 2.78 (s, 1.5H), 1.77-1.65 (m, 2H), 1.57-1.51 (m, 1H), 1.45 (s, 9H), 0.95-0.92 (m, 6H). Chiral analysis: 99.5% e.e. by GC.

Synthesis of Benzyl-L-Lactate (**M2**):

L-Lactic acid (467g, 5.2mol, anhydrous) was dissolved in DMF (1L). Cesium carbonate (847g, 2.6mol, 0.5eq) was added with stirring, followed by benzyl bromide (886g, 5.18mol, 0.99eq) over about 45 minutes. The mixture was stirred at room

temperature for 4 days, after which time analysis showed the reaction to be complete. Heterogeneous mixture was filtered through Celite®; the filter cake was washed with ethyl acetate (2 x 500mL). The filtrate was extracted with water (2 x 500mL), saturated aqueous Na₂CO₃ (4 x 500mL) and brine (2 x 500mL), then concentrated *in vacuo* to yield a thick orange oil, **M2**, 775g, 80%. This was purified by dry flash chromatography using EtOAc/heptane (0% to 10%) as eluent. Appropriate fractions were combined and concentrated *in vacuo* to yield benzyl-L-lactate, 385g, 41%. ¹H NMR (CDCl₃, 300 MHz): δ 7.40-7.35 (m, 5H), 5.21 (s, 2 H), 4.37-4.27 (m, 1H), 2.79 (d, 1H), 1.43 (d, 3H).

Synthesis of benzyl (R)-2-hydroxy-3-(4-((methoxycarbonyl)amino)phenyl)propanoate (**M3**):

Step 1:

NaNO₂ (147.6g, 2.13mol) in water (600mL) was added drop-wise to a mixture of 4-nitro-D-phenylalanine, (150g, 0.71mol) in 1M H₂SO₄ (aqueous, 900 mL), water (750mL) and acetone (2.25L) maintaining a temperature between -5°C and -3°C. The mixture was stirred at -5°C for 1.5 hours then allowed to warm to room temperature and stirred for 3 days. The mixture was concentrated *in vacuo* to remove the acetone and the mixture was extracted with EtOAc (5x~375mL), the combined organics were dried over MgSO₄ filtered and the solvent removed *in vacuo* to yield a yellow solid that was recrystallised from isopropyl acetate (~2.5 volumes), to yield p-nitrophenyl-D-lactic acid (68.38g, 45%) as a yellow solid; a second crop 6.25g (combined yield 50%) was obtained from the residue by slurrying in DCM. ¹H NMR (CDCl₃, 300 MHz): δ 8.18 (d, 2H), 7.45 (d, 2H), 4.59-4.54 (m, 1H), 3.31 (dd, 1H), 3.10 (dd, 1H). No chiral analysis was performed.

Step 2:

Cs₂CO₃ (123.3g, 378mmol) was added to a mixture of p-nitrophenyl-d-lactic acid (156.75g, 742mmol) in DMF (700mL) and stirred for 15 minutes, gas was evolved and the heterogeneous solution cleared yielding a homogeneous brown solution (a mild exotherm was observed). Benzyl bromide (124.39g, 727mmol) in DMF (140mL) was added at room temperature over about 5 minutes; a precipitate formed and the mixture was stirred for about 20 hours at room temperature. The mixture was filtered through Celite® and the filter cake was washed with EtOAc

(2×500mL), the filtrate was washed with water (2×500mL), saturated aqueous NaHCO₃ (3×500mL), and then brine (500mL). Heptane (about 1L) was added to the organic layer and the resultant off white precipitate was filtered off, further crops were obtained from the mother liquors after concentration and trituration with heptanes. The batches were dried at 40°C for about 18 hours and combined to yield an off white solid (197.6, 88% yield). ¹H NMR (CDCl₃, 300 MHz): δ 8.04 (m, 2H), 7.45-7.25 (m, 7H), 5.20 (dd, 2H), 4.55-4.48 (m, 1H), 3.25-3.00 (dd, 2H), 2.86 (d, 1H). UPLC (CSH_C18, Short acid 2-95%): 0.73 min. No mass ion observed.

Step 3:

Iron powder (256g, 4.59mol) was added to a stirred mixture of (R)-benzyl 2-hydroxy-3-(4-nitrophenyl)propanoate (197.5g, 0.66mol), ammonium chloride (455g, 8.52mol), EtOAc (1L), water (900mL), IMS (100mL) heated at 60°C. The reaction became darker in colour and was stirred for 18 hours at 50°C. The mixture was cooled, filtered through Celite® and the filter cake was washed with EtOAc (about 600mL). The layers were separated, and the organic layer was washed with water (3×500mL). The organic layer was extracted with aqueous 1 M HCl (4×250mL); then the acidic extracts were basified with saturated aqueous NaHCO₃ until about pH 8-9. This was extracted into EtOAc (1L, then 2×500mL); the combined organic layers were concentrated *in vacuo* to yield a solid which was azeotroped with toluene, EtOAc, then DCM. The residue was dried at 50°C to yield an off white solid (146.5g, 82%). ¹H NMR (CDCl₃, 300 MHz): δ 7.42-7.30 (m, 5H), 6.91 (d, 2H), 6.56 (d, 2H), 5.13 (s, 2H), 4.42 (t, 1H), 3.54 (s, br 1H), 3.08-2.80 (m, 2H). UPLC (CSH_C18, Short acid 2-95%): 0.32 min, 271.43 Da [M+H]⁺.

Step 4:


Methyl chloroformate (24.5mL, 316mmol, 1.5eq.), was added drop-wise to a stirred mixture of (R)-benzyl 3-(4-aminophenyl)-2-hydroxypropanoate (57.30g, 211mmol), EtOAc (375mL) and saturated aqueous NaHCO₃ (300mL, 5.2vol) at room temperature over about 20 minutes (a mild exotherm to about 25°C was observed along with gas evolution). The mixture was stirred for a further 10 minutes, the organic layer was removed and washed with brine (200mL), dried over Na₂SO₄ filtered and concentrated *in vacuo* to yield **M3** as a white solid (73.48g, 99% yield) ¹H NMR (CDCl₃, 300 MHz): δ 7.42-7.31 (m, 5H), 7.26-7.13 (m, 2H), 7.12-7.01 (m, 2H),


6.70 (s, 1H), 5.13-5.22 (m, 2H), 4.49-4.43 (m, 1H), 3.74 (s, 3H), 3.06 (dd, 1H), 2.93 (dd, 1H), 2.82 (d, 1H). UPLC (CSH_C18, Short acid 2-95%): 0.64 min, 330.4 Da [M+H]⁺.


Synthesis of benzyl (R)-2-hydroxy-3-phenylpropanoate (**M4**)

Cs₂CO₃ (97.5, 300 mmol) was added to a mixture of (R)-2-hydroxy-3-phenylpropanoic acid (100g, 602 mmol) in DMF (700mL) and stirred for 15 minutes, gas was evolved and the heterogenous solution cleared yielding a homogenous brown solution (a mild exotherm was observed). Benzyl bromide (102.0 g, 602mmol) in DMF (100mL) was added at room temperature over about 5 minutes; a precipitate formed and the mixture was stirred for about 20 hours at room temperature. The mixture was filtered through Celite® and the filter cake was washed with EtOAc (2×400mL), the filtrate was washed with water (2×400mL), saturated aqueous NaHCO₃ (3×400mL), and then brine (300mL). Heptane (about 1L) was added to the organic layer and the resultant off white precipitate was filtered off, further crops were obtained from the mother liquors after concentration and trituration with heptanes. The batches were dried at 40°C for about 18 hours and combined to yield an off white solid (**M4**, 151 g, 89% yield). ¹H NMR (CDCl₃, 300 MHz): δ 8.04 (m, 2H), 7.45-7.25 (m, 7H), 5.20 (dd, 2H), 4.55-4.48 (m, 1H), 3.25-3.00 (dd, 2H), 2.86 (d, 1H).

Scheme 2: Route for Stepwise Synthesis of Bis-Iodo PF1022a







As shown in Scheme 2, dimers **D1** and **D2** are prepared using standard amide bond forming methods and ester bond forming methods followed by standard protecting group removal steps. Tetramer **T1**, is formed from the reaction of **D1** with **D2** using routine amide bond forming methods. **T1** is then selectively deprotected to provide either of two tetramers, **T1-1** and **T1-2**, which when coupled together provide the corresponding linear octadepsipeptide, **O1**. Sequential deprotection and cyclization yielded the appropriately aryl functionalized cyclic octadepsipeptide, **C1**. Conversion to the advanced intermediate Bis-Iodo PF1022a was accomplished by performing a two-step sequence that included protecting group removal followed by routine sequence to convert the amino group to an iodo group. It will be appreciated by ones skilled in the art that mono-Iodo PF1022a may be readily obtained from the route shown in Scheme 2 through the appropriate use of **M3** and **M4**.

Synthesis of (R)-1-(benzyloxy)-3-(4-((methoxycarbonyl)amino)phenyl)-1-oxopropan-2-yl N-(tert-butoxycarbonyl)-N-methyl-L-leucinate (**D1**):

Step 1:

EDC.HCl (134.8g, 703mmol), was added in one portion to a mixture of (R)-benzyl 2-hydroxy-3-(4-((methoxycarbonyl)amino)phenyl)propanoate (178.2g, 541mmol), Boc-Methyl-L-Leucine (146.0g, 595mmol), DMAP (6.60g, 54mmol), 4-methylmorpholine (154.6mL, 406mmol) and DCM (3.4L). EDC.HCl slowly dissolves forming an orange solution; a mild exotherm was controlled by cooling after about 15 minutes. After 5 hours the reaction was washed with water (2×2 L), 10% aqueous citric acid solution (2×1L), 5% aqueous citric acid solution (1×1L), saturated aqueous NaHCO₃, water (1L), dried over MgSO₄, filtered and the solvent removed *in vacuo* to yield an oil 296.1g, 98%. ¹H NMR (CDCl₃, 400 MHz): δ 7.44-6.95 (m, 9H), 6.67-6.45 (m, 1H), 5.36-4.64 (m, 4H), 4.14-3.63 (m, 3H), 3.22-2.92 (m, 2H), 2.74-2.50 (m, 3H), 1.69-1.32 (m, 12H), 1.01-0.79 (m, 6H). UPLC (CSH_C18, Short acid, 2-95%): 1.06 min, 457.6 Da, [M-Boc+H]⁺.

Step 2:

Pd-C 10% w/w (20.7g, 19mmol) was washed into a solution of (R)-1-(benzyloxy)-3-(4-((methoxycarbonyl)amino)phenyl)-1-oxopropan-2-yl N-(tert-butoxycarbonyl)-N-methyl-L-leucinate (296g, 532mmol) in industrial methylated spirits (IMS, 3L) with toluene (about 80mL). The mixture was left to stir for about 18 hours under hydrogen (1 atm) for 2 days. The mixture filtered through Celite®, the filter cake was washed with IMS (about 500mL) and the mixture concentrated *in vacuo*. The residue was purified by filtering through a plug of silica eluting with 40% EtOAc in heptanes and concentrated *in vacuo* to yield **D1**. (232.72g, 91%). ¹H NMR (CDCl₃, 300 MHz): δ 7.35-7.25 (m, 2H), 7.17-7.11 (m, 2H), 6.94 (s, br 1H), 5.27-5.18 (m, 1H), 4.73-4.67 (m, 1H), 3.76 (s, 3H), 3.20-3.05 (m, 2H), 2.80-2.70 (m, 3H), 1.76-1.40 (m, 11H), 0.95-0.85 (M, 6H), 96.72% pure w/w by HNMR with 0.34% DCM and 1.60% EtOAc, equivalent to 225.08g, 90.6%. UPLC (CSH_C18, Short acid 2-95%): 0.86 min, 367.6 Da [M-Boc+H]⁺.

Synthesis of Synthesis of (R)-1-(benzyloxy)-1-oxopropan-2-yl N-(tert-butoxycarbonyl)-N-methyl-L-leucinate (**D2**):

Step 1:

Triphenylphosphine (192g, 732mmol), Boc-Methyl-L-Leucine (165g, 672.6mmol), and benzyl-L-Lactate (120g, 666mmol) were dissolved in tetrahydrofuran (700mL). The resulting solution was cooled to 0°C, then di-tertbutylazodicarboxylate (192g, 834mmol, 1.25eq.) was added portionwise, maintaining internal temperature < 5°C. Once addition complete, mixture was allowed to stir and warm to room temperature overnight, during which time an off-white precipitate had formed. Heptane (500mL) was added, and the resulting mixture filtered through Celite®. The cake was washed with heptane (2x100mL). The filtrate was concentrated *in vacuo* to yield a thick orange oil (602g). Heptane (500mL) was added, and the mixture was stirred vigorously for about 1 hour, resulting in further white precipitate forming. This was removed by filtration; the cake was washed with heptane (2x300mL). The resulting filtrate was loaded directly onto silica (2kg), and eluted with EtOAc/heptane (1% to 10%). First fraction (72.6g) still contained triphenylphosphine oxide; this was recolumned (about 700g silica, about 10% loading), using EtOAc/heptane (0% to 6%), to yield 68.2g (25.1%) of Boc-MeLeu-DLac-OBn. The second fraction from the initial column was concentrated *in vacuo* to yield 100.1g (combined = 168.3g, 62%) of (R)-1-(benzyloxy)-1-oxopropan-2-yl N-(tert-butoxycarbonyl)-N-methyl-L-leucinate as a straw coloured oil. ¹H NMR (CDCl₃): δ 7.46-7.30 (m, 5H), 5.21-5.08 (m, 3H), 4.99-4.70 (ddd, 1H), 2.73 (d, 3H), 1.73-1.40 (m, 18H), 0.92 (t, 6H). UPLC (CSH_C18, Short acid 2-95%): 1.08 min, 308.5 Da [M-Boc+H]⁺.

Step 2:

(R)-1-(benzyloxy)-1-oxopropan-2-yl N-(tert-butoxycarbonyl)-N-methyl-L-leucinate (100.1g, 243mmol) was dissolved in dichloromethane (575mL, 5.75vol), and subsequently cooled to 0°C. Trifluoroacetic acid (150mL, 1.96mol, 8eq., 1.5vol) was then introduced. UPLC data after 4 hours indicated that reaction was about 50% complete. Further portion of trifluoroacetic acid (75mL, 980mmol, 4eq., 0.75vol) was added. UPLC indicated not complete after 7 hours, so mixture was stirred overnight for about 16 hours. Toluene (300mL) was added, and the mixture concentrated *in vacuo* to yield a thick orange oil, **D2**, ¹H NMR shows residual toluene and trifluoroacetic acid. ¹H NMR (CDCl₃, 300 Mhz): 9.67-9.15 (br s, 1H), 8.65-8.10 (br s, 1H), 7.47-7.26 (m, 5H), 5.29-5.09 (m, 3H), 3.97-3.85 (m, 1H), 1.92-1.78 (m, 1H), 1.77-1.66 (m, 2H), 1.59-1.54 (s, 3H), 0.98-0.90 (m, 6H).

UPLC (CSH_C18, Short acid 2-95%): 0.43 min, 308.5 Da [M+H]⁺.

Synthesis of (R)-1-(benzyloxy)-1-oxopropan-2-yl N-((R)-2-((N-(tert-butoxycarbonyl)-N-methyl-L-leucyl)oxy)-3-(4-((methoxycarbonyl)amino)phenyl)propanoyl)-N-methyl-L-leucinate (**T1**):

D1 (193g, 458mmol) and **D2** (225g, 482mmol) were stirred in dichloromethane (2L). HATU (250g, 658mmol, 1.36eq.) was added, followed by diisopropylethylamine (375mL, 2150mmol) through a dropping funnel. Temperature of reaction was maintained with the use of an external ice-bath. The mixture was allowed to stir overnight at room temperature. The reaction mixture was washed with water (3x2L), aqueous citric acid (10%, 2x1L), saturated aqueous NaHCO₃ (2x1L), and water (1L). The organic phase was dried over MgSO₄, filtered and concentrated *in vacuo* to yield an orange oil. This was purified by dry flash chromatography using EtOAc/heptane as eluent (0% to 30%), to yield **T1** (313 g, 83 %).

¹H NMR (CDCl₃, 300 MHz): δ 7.41-7.26 (m, 7H), 7.20-7.10 (m, 2H), 6.59 (s, 1H), 5.42-4.60 (m, 6H), 3.82-3.71 (m, 3H), 3.10-2.96 (m, 2H), 2.93-2.71 (m, 6H), 1.77-1.36 (m, 18H), 1.01-0.83 (m, 12H). UPLC (CSH_C18, Short acid, 2-95%): 1.14 min, 656.9 Da [M-Boc+H]⁺.

Synthesis of (R)-1-(benzyloxy)-1-oxopropan-2-yl N-((R)-3-(4-((methoxycarbonyl)amino)phenyl)-2-((methyl-L-leucyl)oxy)propanoyl)-N-methyl-L-leucinate (**T1-1**):

T1 (48.7g, 64.4mmol) was dissolved in dichloromethane (300mL, 6vol). The resulting solution was cooled to 0°C with an external ice-bath. TFA (75mL, 980mmol, 1.5vol) was added to the reaction mixture and the mixture stirred until complete. Toluene (300mL) was added, and the mixture concentrated *in vacuo* yield a thick orange oil, 65.3g, about 125% (containing ~16g of TFA). Used without further purification. ¹H NMR (CDCl₃, 300 MHz): δ 7.40-7.23 (m, 9H), 6.73 (s, 1H), 5.50-5.42 (m, 1H), 5.29-5.08 (m, 4H), 3.87-3.83 (m, 1H), 3.79-3.77 (m, 3H), 3.06 (d, 2H), 3.01-2.90 (m, 3H), 2.70-2.64 (m, 3H), 2.36 (m, 3H), 1.76-1.42 (m, 6H), 1.39-1.25 (m, 2H), 1.04-0.96 (m, 1H), 0.94-0.87 (m, 6H), 0.79-0.76 (m, 6H). UPLC (CSH_C18, Short acid 2-95%): 0.65 min, 656.91 Da [M+H]⁺.

Synthesis of (6S,9R,12S,15R)-6,12-diisobutyl-9-(4-((methoxycarbonyl)amino)-benzyl)-2,2,5,11,15-pentamethyl-4,7,10,13-tetraoxo-3,8,14-trioxa-5,11-diazaheptadecan-16-oic acid (**T1-2**):

Pd-C (10% w/w, 2.5g, 2.3mmol) was washed into a solution of **T1** (48.3g, 63.9mmol) in IMS (500 mL) with toluene (about 20mL) and stirred under hydrogen (1atm) for 4 hours, filtered and the solvent removed *in vacuo* and then azeotroped with DCM to yield Boc-MeLeu-DCbmPheLac-MeLeu-DLac-OH as an off-white foam 48.87g, **T1-2**, 115% yield (used crude in subsequent reaction). ¹H NMR (CDCl₃, 400 MHz): δ 7.05-7.29 (m, 4H), 5.70 (t, 1H), 4.65-5.45 (m, 5H), 3.75 (d, 3H), 2.68-3.19 (m, 8H), 2.30-2.34 (m, 1H), 1.38-1.74 (m, 17H), 1.10-1.29 (m, 1H), 0.83-0.99 (m, 13H). UPLC (CSH_C18, Short acid 2-95%): 0.97 min, 610.8 Da [M-^tBu+H]⁺, 664.8 Da [M-H]⁻.

Synthesis of dimethyl (((((2S,5R,8S,11R,14S,17R,20S,23R)-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octaoxo-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,14-diyl)bis(methylene))bis(4,1-phenylene))dicarbamate (**C1**):

Step 1:

T1-1 (49.3g, actual mass 65.3g, contains ~ 16 g trifluoroacetic acid, 64.4mmol) and **T1-2** (42.5g, actual mass 48.5g, contains residual solvent from hydrogenation) were dissolved in dichloromethane (500mL). The mixture was cooled with an external ice-bath. HATU (36.4g, 95.7mmol, 1.5eq.) was added, followed by diisopropylethylamine (70mL, 400mmol, 6.3eq.). The resulting mixture was stirred for 20 hours. Water (1L) was added to the reaction mixture, and vigorously stirred for 0.83 hours. The layers were separated, and the organic phase washed with aq. citric acid (10%, 3x700mL), saturated aqueous NaHCO₃ (1x700 mL), and brine (1x500mL), then dried over MgSO₄, filtered and concentrated *in vacuo* directly onto Celite® (150g) and silica gel (20g). The material was dry loaded onto a column of silica (1kg) and purified by dry flash chromatography using EtOAc/heptane (0-50%) as eluent. This afforded, after concentration *in vacuo*, **O1**, 85 g, 101 % (some trace heptane observable in ¹H NMR spectrum). ¹H NMR (CDCl₃, 300 MHz): δ 7.06-7.34 (m, 13H), 6.61 (s, 1H), 4.58-5.46 (m, 10H), 3.73-3.76 (m, 6H), 2.66-3.16 (m, 15H),

1.16-1.69 (m, 28H), 0.83-0.98 (m, 24H). UPLC (CSH_C18, Long acid, 2-95%): retentiontime of 3.53 minutes, 1204.5 Da, [M-Boc+H]⁺.

Step 2:

O1 (199.5 g, 153 mmol) was dissolved in IMS (400 mL, 2 vol) and charged to the 2 L autoclave. 5% Pd/C (2 g) was added as a paste in toluene. The autoclave was sealed and charged to 150 psi hydrogen pressure. After stirring for 5 hours at room temperature, 16% starting material remained so the autoclave was recharged to 150 psi and allowed to stir at room temperature overnight. The reaction was complete by UPLC, so the autoclave was vented and the catalyst removed by filtration through Whatman GF/A media, washing with IMS (50 mL) and ethyl acetate (200 mL). The solvent was removed on the rotary to leave a white foam, **O1-1** (175 g, 94% yield). ¹H NMR (CDCl₃, 300 MHz): δ 7.06-7.34 (m, 8H), 4.58-5.46 (m, 10H), 3.73-3.76 (m, 6H), 2.66-3.16 (m, 15H), 1.16-1.69 (m, 28H), 0.83-0.98 (m, 24H). UPLC (Long acid, 2-95%): RT 3.36 min, 1113.5 Da, [M-Boc+H]⁺.

Step 3:

O1-1 (245 g, 202 mmol) was dissolved in DCM (1.5 L, 6 vol) and cooled to <10°C (ice/water bath). Trifluoroacetic acid (370 mL, 551 g, 1.5 vol) was added and the solution allowed to warm to room temperature and stirred overnight. The solvent was removed on the rotary and azeotroped with toluene (2 x 500 mL). The product was obtained as a thick orange oil **O1-2** (359.5 g, overweight – estimated purity from NMR is 65%, giving calculated mass of 234 g, 94% yield). ¹H NMR (CDCl₃, 300 MHz): δ 11.91 (s, TFA), 7.1-7.35 (m, 8H), 4.6-5.65 (m, 8H), 3.79 (s, 6H), 2.6-3.2 (m, 18H), 1.1-1.9 (m, 16H), 0.65-1.07 (m, 28H).

Step 4:

A solution of **O1-2** (359.5g, 65% purity, 0.190mol) in DCM (2.5L) was added slowly over 1.5 hours to a solution of propylphosphonic anhydride (50% solution in EtOAc, 670mL, 1.126mol) and diisopropylethylamine (392mL, 2.25mol) in DCM (2.5L), maintaining the temperature between 20 and 22°C. When the addition was complete, the solution was allowed to stir for a further 20 minutes and the reaction was checked for completion by UPLC. The solvent was removed on the rotary and replaced with EtOAc (5L). The solution was washed with potassium hydrogen sulphate solution (1M, 2x1L), 3% sodium carbonate solution (2x1.5L) and brine (1L).

The organic layer was dried over MgSO_4 and filtered through Fisherbrand QL100 paper twice. The solvent was removed to leave a yellow foam (233g). This was recrystallised from ethanol (660mL) and water (130mL) and the product was washed with ethanol/water (600mL, 3:1). The product was dried on the filter bed and further dried on the rotary at 60°C. The product, **C1**, was obtained as a white solid (159.2g, 77% yield). ^1H NMR (CDCl_3 , 300 MHz): δ 7.3 (m, 4H), 7.15 (m, 4H), 4.4-5.67 (m, 8H), 3.76 (s, 6H), 2.7-3.1 (m, 16H), 1.16-1.8 (m, 16H), 0.73-1.1 (m, 28H). UPLC (Short acid 2-95%): 1.08 min, 1095.5 Da $[\text{M}+\text{H}]^+$.

Synthesis of Bis-Iodo PF1022a:

Step 1:


C1 (151.5g, 0.138mol) was added in portions to a 33% solution of HBr in acetic acid (750mL) and stirred at 35°C for 6.5 hours. The reaction mixture was cooled to 15°C (ice/water bath) and ice/water (1.75L) was added. A solution of 10% sodium carbonate was added carefully (about 2.5L) to basify the mixture to pH 4-5. The mixture was extracted with ethyl acetate (2L). The organic layer was separated and washed with saturated aqueous NaHCO_3 (2x1.5L), and brine (1L). The organic layer was dried over Na_2SO_4 and evaporated to a crisp foam (143g). The foam was recrystallised from 2-propanol (600mL) and dried to leave the product, **C1-1**, as a free flowing white powder (117.0g, 87% yield). ^1H NMR (CDCl_3 , 300 MHz): δ 6.99 (m, 4H), 6.58 (m, 4H), 4.4-5.6 (m, 8H), 3.62 (br s, 4H), 2.7-3.1 (m, 16H), 1.16-1.9 (m, 14H), 0.73-1.1 (m, 28H). UPLC (Long acid 2-95%): 2.79 min, 980.3 Da $[\text{M}+\text{H}]^+$.

Step 2:

To a solution of **C1-1** (10g, 10.2mmol) in CH_3CN (125mL) and added pTSA (7.1g, 40.8mmol, 4.0eq) at room temperature and then cooled to -2°C using ice-salt bath. A 1.5M solution of NaNO_2 (1.70g, 24.5mmol, 2.4eq) in water (15mL) was slowly added over a period of 7 minutes while maintaining temperature around -1°C. After 15 minutes, CuI (8.7g, 46.0mmol, 4.5eq) was added over 5 minutes and rinsed with water (20ml). Resulting brown mixture was stirred at about -1°C for 30 minutes, and then stirred at about -2 to 5°C for 30 minutes without cooling. Reaction was cooled to 2°C by adding ice into reaction and quenched with 25% sodium thiosulphate solution (100mL). Organic layer was separated and aqueous layer

extracted with ethyl acetate (2x75ml). Combined organic solution was washed with water (100mL) and saturated NaHCO_3 (75mL), brine (100mL), dried over Mg_2SO_4 and concentrated under vacuum to get solid 16g. Obtained solid dissolved in DCM (30mL), adsorbed on silica gel (60g) and filter through silica gel bed (100g) using 50% EtOAc/hexane. Organic solution was concentrated under vacuum to a solid and dried under vacuum at 40°C for overnight to get Bis-Iodo PF1022a, (7.5 g, 62 %). ^1H NMR (600 MHz, CDCl_3) δ : 0.68-0.98 (m, 29H), 1.24-1.7 (m, 13 H), 2.67-3.07 (m, 16 H), 4.36-4.43 (m, 1H), 4.96-5.63 (m, 7 H), 6.98-7.05 (m, 4H), 7.60-7.69 (m, 4H). LC-MS (m/z): $[\text{M}^+]$ = 1200

Scheme 3: Route for Preparation of Bis- CH_2Cl PF1022a:



As shown in Scheme 3, conversion of Bis-Iodo PF1022a to Bis-Aldehyde PF1022a is accomplished by a three step sequence. Palladium catalyzed carbonylation in an atmosphere of carbon monoxide to install the aldehyde that is then reduced with a hydride reagent to form Bis- CH_2OH PF1022a that is readily

converted using standard functional group to the Bis-CH₂Cl PF1022a intermediate. In the preparation of Bis-Aldehyde PF1022a a small but appreciable amount (2%-20%) of 4-(((2R,5S,8R,11S,14R,17S,20R,23S)-14-benzyl-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octaoxo-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2-yl)methyl)benzaldehyde (Mono-Aldehyde PF1022a) is also formed. Mono-Aldehyde PF1022a may be separated from Bis-Aldehyde PF1022a using standard reverse phase HPLC methods. The mono-aldehyde PF1022a was also used as a starting material to prepare some of the compounds presented in Tables 1-4, according to Scheme 8.

Synthetic Procedures for the Preparation of Bis-CH₂Cl PF1022a (Scheme 3):

Step 1:

A 2L pressure reactor was charged with Bis-Iodo PF1022a (130.0g, 101.7mmol), HSi(octyl)₃ (225g, 610.1mmol), DMF (700mL), TEA (131mL, 938.8mmol) and Pd(TPP)₄ (5.6g, 4.8mmol) and then purged with nitrogen and carbon monoxide. Reaction was heated to 65°C under carbon monoxide (55 psi) and stirred for 2 hours. Reaction was cooled to 22°C, slowly diluted with ice-water (2.5 L). Solid was collected, washed with water (2x250 mL) hexane (2x200 mL) and dried under vacuum at 50°C for overnight to get crude Bis-Aldehyde PF1022a as light brown solid (130g).

Step 2:

A 3L round bottom flask charged with crude Bis-Aldehyde PF1022a (130 g) and methanol (1100 mL) and then cooled to -35°C using acetonitrile/dry-ice bath. To the mixture was slowly added NaBH₄ (5.0g, 130mmol) while maintaining temperature below -30°C and stirred between -35 and -37°C for 1 hour. Reaction was slowly quenched with saturated NH₄Cl (100mL), brought to 10°C and concentrated to approximately 3/4 volume under vacuum at 40°C. Mixture was diluted with water (150 mL), extracted with ethyl acetate (3x 150 mL). Organic solution was dried over MgSO₄, filtered and concentrated under vacuum at 45°C to a dark brown solid. Solid was heated in 1:2 mixture of EtOAc/Hexane (200 ml) at 60°C and stirred at room temperature for 4 hours. Solid was collected, washed with 25% ethyl acetate in hexane (2x 50 mL), hexane (50 mL) and dried under vacuum to get


Bis-CH₂OH PF1022a (103.5 g, 91% yield for 2 steps). ¹H NMR (600 MHz, CDCl₃) δ: 0.75-1.17 (m, 26H), 1.23-1.96 (m, 16H), 2.70-2.92 (m, 9H), 3.01-3.23 (m, 7H), 4.45-4.53 (m, 1H), 4.65-4.74 (m, 4H), 5.03-5.14 (m, 1H), 5.19-5.72 (m, 6H) 7.20-7.36 (m, 8H). LC-MS (*m/z*): [M+H] = 1009 and [(M+23)] = 1031

Step 3:


A 1L 3 neck round bottom flask was charged with Bis-CH₂OH PF1022a (98g, 90.30 mmol) and DCM (300mL) and then cooled to 10°C with external ice-water bath. Thionyl chloride (25 ml, 343.2 mmol) was added over 7 minutes, cooling bath removed and mixture stirred at room temperature. After 1 hour, reaction was concentrated under vacuum at 35°C to a syrup, diluted with heptane (2x 75 mL) and concentrated to a solid. Brown solid was dissolved in ethyl acetate (250 ml) washed with aqueous Na₂CO₃ (150 mL), 50 mL saturated Na₃CO₃ solution diluted with water (100 mL) to pH 7.8. Organic layer was separated and aqueous layer extracted with ethyl acetate (2x100 mL). Combined organic solution was dried over MgSO₄, and concentrated under vacuum to a brown solid. Crude product in ethyl acetate (300 ml) was stirred with activated carbon (12g) for 4 hours at room temperature, filtered through celite bed and concentrated to solid. Obtained crude material in MTBE (110mL) was heated to reflux, diluted with hexane (50ml) and stirred at room temperature for overnight. Solid was collected, washed with 20% MTBE/hexane (50mL), hexane (50mL) and dried under vacuum at 50°C to get BisCH₂Cl PF1022a as off-white solid (98 g, 98% yield). ¹H NMR (600 MHz, CDCl₃) δ: 0.75-1.17 (m, 26H), 1.25-1.95 (m, 16H), 2.70-2.90 (m, 9H), 3.04-3.26 (m, 7H), 4.44-4.57 (m, 1H), 4.54-4.62 (m, 4H), 5.05-5.13 (m, 1H), 5.24-5.74 (m, 6H) 7.20-7.39 (m, 8H). LC-MS (*m/z*): [M+H] = 1045, [M+2] = 1046, [M+3] = 1047 and [M+4] = 1047.

Scheme 4: Grignard Process [4.1 - 4.4]


[4.1]




and [4.2]



and [4.3]



and [4.4]



In Scheme 4, R* represents the L₁ and L₂ aryl, cycloalkyl, heteroaryl, and heterocycle moieties, each of which can be substituted as described herein; and R' can represent H or an alkyl group (e.g., methyl).

As shown in Scheme 4, Bis-Iodo PF1022a will undergo metal halogen exchange when treated with iPrMgCl-LiCl to give a depsipeptide Grignard reagent that can be reacted with a variety of electrophiles, such as aldehydes or ketones, to provide compounds **1a** and **1b**. The reaction yields the mono-substituted product, **1b**, in minor amounts for most such reactions. Compounds **1a** and **1b** can be deoxygenated by treatment with triethylsilane in TFA or triethylsilane with PdCl₂ to give compounds of the type **2a** and **2b**. Compounds **1a** and **1b** can be separated by preparatory HPLC so as to prepare compounds **2a**, **2d**, and **2e** from **1a**; and **2b**, **2c**, and **2f** from **1b**; as shown in Scheme 4. Alternatively, a mixture of **1a** and **1b** can be reacted to provide mixtures of **2a** and **2b**, **2d** and **2c**, and **2e** and **2f**, respectively, which can then be separated by preparatory HPLC.


Preparation of Examples 3-7, 3-9, 4-6, and 4-8 (Scheme 4).

(3-7). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(hydroxy(6-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone.

(3-9). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-(hydroxy(6-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone.

(4-6). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-(hydroxy(6-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone.

(4-8). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-18-(4-((6-methoxypyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone




Step 1:

Bis-Iodo PF1022a (200 mg, 0.16 mmol) was dissolved in THF (6 ml) in an atmosphere of nitrogen. The solution was cooled to -78°C using a dry ice/ $i\text{PrOH}$ bath. An excess of Turbo Grignard ($i\text{PrMgCl}:\text{LiCl}$, 1.3M in THF, 0.90mL, 1.17mmol,) was added and the mixture was stirred at -78°C for 10 minutes. The dry ice bath was then replaced with an ice water bath (0°C). The solution was allowed to stir at 0°C for 20 minutes. A THF solution (2 ml) of 6-methoxynicotinaldehyde (137 mg, 1.0 mmol) that had been cooled to 0°C was then added to the Grignard mixture. The solution was stirred for an additional 30 minutes at 0°C before the reaction was quenched with AcOH (0.150 ml). The reaction mixture was added to EtOAc (25 ml) and the organic phase was washed with water (20 ml), dried over sodium sulfate and concentrated to give viscous oil which was dissolved in acetonitrile (3 ml) and purified by reverse phase HPLC to give the products: Example (3-7), 115mg; and Example (4-6), 39mg.

Step 2:

Example (3-7) (100 mg, 0.07 mmol) was dissolved in 1,2-DCE (4 ml). TFA (0.5 ml) and triethylsilane (0.5 ml, 4.3 mmol) were then added and the solution was heated at 90°C in an atmosphere of nitrogen for 6 hours. The solution was concentrated and purified using reverse phase HPLC to give Example (3-9) (51 mg) product. From steps 1 and 2, a minor amount of Example (4-8); 13mg is also obtained.

When compounds such as **1a** and **1b** contain an R' substituent, such as a methyl (Scheme 4.4), treatment with triethylsilane and TFA provide an intermediate alkene from alcohol dehydration. The crude intermediate can then be reduced to the alkanes **2e** and **2f** by further addition of Pd/C.



Preparation of Example (3-207)


To a solution of Bis-Iodo PF1022a (441.9 mg, 0.3679 mmol) in THF (26.5 mL) at -78°C was added $i\text{PrMgCl} \cdot \text{LiCl}$ complex in THF (1.3 M, 2.8 mL) dropwise over 2 minutes and the mixture was allowed to stir for 5 minutes at this temperature. The solution was warmed at 0°C in an ice bath and after 30 minutes a solution of 1-(6-methoxy-3-pyridyl)ethanone (1.0 g, 6.62 mmol) in THF (2.95 mL) was added all at once down the sides of the flask before the mixture was quickly warmed to rt. The reaction mixture was stirred at this temperature for 1 hour before being quenched with saturated aqueous NH_4Cl and extracted with EtOAc. The combined organics were washed with brine, dried, and concentrated. The residue was purified via reverse phase HPLC to provide 83 mg (18%) of the bis-substituted product and 92

mg (22%) of mono-substituted product.

To a solution of (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis[[4-[1-hydroxy-1-(6-methoxy-3-pyridyl)ethyl]phenyl]methyl]-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetrazacyclotetracosane-2,5,8,11,14,17,20,23-octaone (35.0 mg, 0.0280 mmol) in 1,2-dichloroethane (1.0 mL) was added TFA (0.3 mL, 4 mmol) followed by triethylsilane (0.3 mL, 2 mmol) in a MW vial. The reaction mixture was heated at 90°C for 1 hour and then cooled to room temperature. To the cooled reaction mixture was added Pd/C (30 mg) and gas evolution proceeded. After 30 minutes the crude mixture was filtered through Celite and the cake washed with DCM. The combined filtrates were concentrated and by purified reverse phase HPLC to afford 18 mg (52%) of Example (3-207) as a white powder.

Compounds **1a** and **1b** can also undergo deoxyfluorination with reagents such as DAST to provide compounds such as **2c** and **2d** (Scheme 4.3).


Preparation of Example (3-252)



A solution of (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis({4-[hydroxy(5-methoxypyrazin-2-yl)methyl]phenyl}methyl)-4,10,12,16,22,24-hexamethyl-3,9,15,21-tetrakis(2-methylpropyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,5,8,11,14,17,20,23-octaone **2** (200 mg, 0.16 mmol) in DCM (10 mL) was cooled to -78°C and a solution of DAST (0.32 mL, 2.39 mmol) in DCM (2 mL) was added slowly. The resulting mixture was warmed to room temperature and stirred for 16 hours. The reaction mixture was quenched with ice-water and extracted with 10% methanol in dichloromethane. The combined organic layer was washed with brine,

dried over sodium sulphate, filtered and concentrated under reduced pressure. Crude material was purified by prep-HPLC to afford Example (3-252) (5 mg, 3%) as an off white solid.

Scheme 5: Reverse Grignard or Halogen-Metal Exchange Process




In Scheme 5, R^* represents the L_1 and L_2 aryl, cycloalkyl, heteroaryl, and heterocycle moieties, each of which can be substituted as described herein; and R' is H.

As shown in Scheme 5, an aryl, cycloalkyl, heterocycle, or heteroaryl halide (such as iodide or bromide) will undergo metal halogen exchange when treated with $i\text{PrMgCl-LiCl}$ to give a Grignard reagent that can be reacted with Bis-Aldehyde PF1022a to provide compounds **1a** and **1b** (Scheme 4). The reaction yields the mono-substituted product, **1b**, in minor amounts for most such reactions. Compounds **1a** and **1b** can be deoxygenated by treatment with triethylsilane in TFA or triethylsilane with PdCl_2 to give compounds of the type **2a** and **2b** (Scheme 4).

Preparation of (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-Bis-[4-(5-cyclopropyl-[1,3,4]oxadiazol-2-ylmethyl)-benzyl]-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-

hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraaza-cyclotetracosan-2,5,8,11,14,17,20,23-octaone (3-330) in view of Scheme 5.




A stirred solution of 2-bromo-5-cyclopropyl-[1,3,4]oxadiazole (941 mg, 4.98 mmol) in dry THF (5 ml) was cooled to -78°C and $i\text{PrMgCl} \cdot \text{LiCl}$ (1.3M, 3 ml, 3.984 mmol) was added drop-wise. The reaction mixture was allowed to stir at -10°C for 40 minutes. A solution of Bis-Aldehyde PF1022a (500 mg, 0.498 mmol) in dry THF (1 ml) was added to the mixture in one portion and allowed to stir at -10°C for 1.5 hours followed by 2 hours at room temperature. The reaction mixture was quenched with saturated NH_4Cl solution and extracted with ethyl acetate (2 x 150 ml). The combined organic layer was washed with water, brine, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford crude (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-Bis-{4-[(5-cyclopropyl-[1,3,4]oxadiazol-2-yl)-hydroxy-methyl]-benzyl}-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraaza-cyclotetracosan-2,5,8,11,14,17,20,23-octaone as a gum.

To a solution of the above intermediate (crude, 750 mg, 3.065mmol) in DCE (5 ml) were added triethyl silane (5 ml) and TFA (5 ml) at 23°C . The resulting mixture was heated at 100°C for 1 hour under microwave irradiation. The reaction mass was

poured into saturated NaHCO₃ solution and extracted with DCM (2 x 100ml). The combined organic layers were washed with water, brine, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford crude compound (3-330) (650mg) as a gum. The crude material was purified by preparatory HPLC to afford the desired Example (3-330) (35 mg) as an off white solid.

Scheme 6. Stille and Suzuki Coupling Process



In Scheme 6, R* represents the L₁ and L₂ aryl, cycloalkyl, heterocycle, and heteroaryl, moieties, each of which can be substituted as described herein. "Pd" represents a palladium catalyst.


As shown in Scheme 6, compounds like **2a** can be accessed directly from Bis-CH₂Cl PF1022a via palladium catalyzed cross coupling reactions such as in the Stille or Suzuki reaction. In this case, Bis-CH₂Cl PF1022a is treated with an organoboronic acid, organoboronate ester, or organostannane in the presence of a palladium catalyst with or without base to provide the desired products after heating. The organostannane reagents and organoboronic acids/esters are either commercially available or prepared by standard methods found in the literature (European Journal of Organic Chemistry (2014), 24, 5153-5157; Journal of the American Chemical Society (2002), 12, 8001); Organic Letters (2012), 14(2), 502-505; Journal of Medicinal Chemistry (2005), 48(22), 7080-7083; and PCT Application publications WO2006/108591, WO2003/087102, and WO2012/177714; and from commercially available starting materials.

For Examples 3-50 and 4-40, to a stirred solution of TMP (137 mg, 0.967 mmol) in THF (1 ml), 2.3M n-BuLi in THF (0.37 mL, 0.86 mmol) was added at 0°C, then stirred for 30 minutes. Furan-3-carbonitrile (100 mg, 1.074 mmol) in THF (1 ml) was added at -78°C and stirred at same temperature for 2 hours. Then Bu₃SnCl (0.321 ml, 1.182 mmol) in THF (1 mL) was added at -78°C and the reaction mixture was allowed to stir at room temperature for 15 hours. The reaction was quenched with saturated NH₄Cl solution and extracted with DCM (2 X 15 ml), combined organic layer was washed with brine solution (5 ml), dried over Na₂SO₄ filtered and concentrated. Crude was purified by column chromatography used on silica gel eluting with 20 DCM-Hexane to afford crude stannane monomer, 2-(tributylstannyl)furan-3-carbonitrile, as a light yellow liquid.

For Example 3-209, to a stirred solution of 2-methoxy-4-methylthiazole (75 mg, 0.581 mmol) in THF (5 mL) at -78°C, n-BuLi (0.30 mL) was added and the reaction mixture was stirred for 1 hour at that temperature. Then tributyltin chloride (0.18 ml, 0.64 mmol) was added at -78°C and the reaction mixture was stirred for 2 hours at room temperature. Water (10 mL) was added to the reaction mixture and extracted with ethyl acetate (2 x 20 mL). Total organic part was washed with brine solution, dried (Na₂SO₄) and evaporated at 25°C to afford the stannane monomer, 2-methoxy-4-methyl-5-(tributylstannyl)thiazole, as a dark brown liquid.


For Example 3-238, to a stirred solution of 3-bromo-2-(trifluoromethyl)pyridine (400 mg, 1.77 mmol) in THF (15 mL) at -78°C, n-BuLi (1.8 mL, 1.1 equiv) was added and the reaction mixture was stirred for 1 hour at that temperature. Then tributyltin chloride (0.52 ml, 1.947 mmol, 1.1 equiv) was added at -78°C and the reaction mixture was allowed to stir for 1.5 hours at room temperature. Water (10 mL) was added to the reaction mixture and extracted with ethyl acetate (2 x 20 mL). Total organic part was washed with brine solution, dried (Na₂SO₄) and evaporated to afford the crude stannane monomer, 3-(tributylstannyl)-2-(trifluoromethyl)pyridine, as a dark liquid.

Preparation of (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((5-methoxypyrazin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone, Example (3-33) using the Stille process shown below. Similar compounds, as described in Tables 1-4, were also readily prepared using the Stille process.



To a stirred solution of Bis-CH₂Cl PF1022a (150 mg, 0.14 mmol) and 2-methoxy-5-(tributylstannyl)pyrazine (0.40 g, 1.0 mmol) in p-dioxane (5 ml) was added Pd(PPh₃)₄ (75 mg, 0.065 mmol). The solution was heated at 100°C in an atmosphere of N₂ for 5 hours. The reaction mixture was cooled to room temperature and filtered through Celite. The filtrate was concentrated to a viscous black oil that was dissolved in a small volume of DMSO and purified by reverse phase HPLC to give Example (3-33).


Preparation of (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis[[4-[[2-(difluoromethyl)-3-pyridyl]methyl]phenyl]methyl]-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetrazacyclotetracosane-2,5,8,11,14,17,20,23-octaone, Example (3-290) using the Suzuki process shown below. Similar compounds, as described in Tables 1-4, were also readily prepared using the Suzuki process.



To a solution of Bis-CH₂Cl PF1022a (120 mg, 0.11 mmol) in EtOH:Toluene (4 mL, 1:1) was added 2-(difluoromethyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (120 mg, 0.46 mmol), NaHCO₃ (1 mL of 2M sol.) and Pd(dppf)₂Cl₂ (15 mg, 0.014mmol). The reaction mixture was purged with N₂ and heated at 100°C for 15 minutes in a microwave. After cooling at room temperature the reaction mixture

was diluted with EtOAc/water and the organic phase was separated and concentrated under reduced pressure. The residue was purified by reverse phase chromatography to provide 88 mg (62%) of Example (3-290) as a white solid.


Scheme 7. Process for Preparing Compounds **2g** and **2h**



In Scheme 7, R^* represents the L_1 and L_2 aryl, cycloalkyl, heteroaryl, and heterocycle moieties, each of which can be substituted as described herein; and R' is H. "Nuc" represents a nucleophilic functional group, e.g., alcohols, acyclic and cyclic amines, heterocycles (e.g., morpholine), heteroaryls (e.g., pyrazole), and any nucleophilic organometallic cycloalkyl, heterocycle, aryl, or heteroaryl group, such as organolithiums or Grignard reagents.

In Scheme 7, compounds **1a** and **1b** (Scheme 4) are treated individually with phosphorous tribromide to convert the hydroxyl group to a bromo group using conditions readily found in the literature. The bromo group is readily displaced by nucleophiles (Nuc) to form compounds of type **2g** and **2h** that are described in Tables 1-4.

Preparation of Compound (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3,4-difluorophenyl)(methoxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone (2-37)




Step 1:

Compound **1** (300 mg, 0.275 mmol) was dissolved in DCM (5 ml) and the solution was cooled to 0°C with an ice bath. PBr_3 (0.27 g, 1.0 mmol) was then added and the solution was allowed to stir 1 hour at 0°C and for 2 hours at room temperature. The mixture was diluted with methylene chloride (25 ml) and ice water was added (20 ml). The layers were mixed then allowed to separate. The organic phase was collected, dried over sodium sulfate, then concentrated to give the crude product as an amber amorphous glassy solid. The solid was used as is in Step 2.

Step 2:

To the crude bromide, **2**, (150 mg, 0.13 mmol) was added MeOH (2 ml) and the solution was stirred at 45°C for 3 hours. The mixture was directly purified by reverse phase HPLC to give Example (2-37) (42 mg, 0.038 mmol) as a white solid after lyophilization.


Scheme 8: Organometallic Process for Preparing Compounds from Bis-Iodo PF1022a



In Scheme 8, R^* represents the L_1 and L_2 aryl, cycloalkyl, heteroaryl, and heterocycle moieties, each of which can be substituted as described herein. "Nuc" represents a nucleophilic functional group as defined above.

As shown in Scheme 8 compounds having the general structures **4a** and **4b** are prepared from **1a** and **1b** (when $R' = H$) via a two-step process. In the first step the alcohol **1a** or **1b** is treated with an oxidizing reagent, such as Dess-Martin periodinane, to yield a ketone (**2i** and **2j**) that will react with a Grignard reagent to

give compounds of the type **4a** and **4b** in the second step; e.g., Example (4-26) can be prepared.




Step 1:

Compound **4-6** (50 mg, 0.046 mmol) was dissolved in DCM (3 ml) and Dess-Martin periodane (42 mg, 1 mmol) was added to the stirring solution. The solution was stirred at room temperature for 3 hours. The reaction mixture was diluted with DCM (10 ml) and cooled to 0°C. Then a 1M aqueous solution of sodium bisulfite (2 ml) was added and the mixture was stirred at 0°C for 0.5 hours. The organic phase was collected and washed with water (2 x 3 ml). The organic phase was dried with sodium sulfate then concentrated to give crude compound 5 that was used without further purification.


Step 2:

Compound **5** (39 mg, 0.035 mmol) was dissolved in anhydrous THF (5 ml) and the solution was cooled to 0°C before MeMgBr (3 M in THF, 0.5 ml, 1.5 mmol) was added. The solution was then warmed to room temperature and stirred for 30 minutes. The solution was cooled again to 0°C and quenched by the addition of AcOH (0.2 ml). The solution was then diluted with EtOAc (15 ml), washed with water (1 x 5 ml) and concentrated to provide crude Example (4-26). Purification was done using reverse phase HPLC to give Example (4-26) as a white solid after lyophilization.

Scheme 9: Process for Preparing Pyrrole Containing Compounds




As shown in Scheme 9, pyrroles were reacted with BisCH₂Cl PF1022a under basic conditions to give bis-C-linked pyrroles as products. (R)_n is as defined herein. It will be understood that monoCH₂Cl PF1022a may be reacted similarly with pyrroles to give the mono-C-linked pyrroles as products; e.g., Example (3-29) can be prepared using this scheme, as shown below.



To (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis[[4-(chloromethyl)phenyl]-methyl]-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetrazacyclotetracosane-2,5,8,11,14,17,20,23-octaone (50.0 mg, 0.048 mmol) was added 2,4-dimethyl-1H-pyrrole (5.0 equiv., 0.24 mmol), potassium bicarbonate (24 mg), and potassium iodide (4.0 equiv, 0.191 mmol) followed by acetonitrile (1.00 mL). The reaction mixture was then heated at 50°C for 48 hours. The reaction was cooled and filtered through celite and washed with DCM. The crude material was absorbed onto celite and purified via reverse phase

chromatography with a gradient of 60-95% acetonitrile in H₂O. Like fractions were combined and lyophilized overnight to provide the product, Example (3-29), (20 mg, 32%), a purplish solid.

In addition, other C-linked heterocyclic compounds, for Example (3-5) and (3-14), can be prepared using the following procedural Steps.



Step 1:

Pre-mixed tetrabutylammonium fluoride (4 eq) and TMSCN (4 eq) in acetonitrile were heated at 40°C for 1 hour, after that addition of depsibenzylchloride (500 mg) was added at room temperature and stirred for 16 hours. After aqueous workup, the residue was purified by preparative HPLC to give the bis benzyl nitrile.

Step 2:

A mixture of bis benzyl nitrile (40 mg), thiosemicarbazide (5 eq) and TFA was stirred at 90°C for 16 hours. After cooling to room temperature, the mixture was given an aqueous work up and purified by preparatory HPLC to give the bis aminothiadiazole Example (3-5).

Step 3:

Aminothiadiazaole (3-5; 23.5 mg), *t*-BuONO (10 eq), THF, 0°C to room temperature, stirred for 16 hours. The solvent was evaporated and the residue purified by preparatory HPLC to give the bis thiadiazaole Example (3-14).

Similarly, the tetrazole Example (3-6) can be prepared.

Mass data ((ESI-MS m/z $[M+H]^+$), unless defined otherwise, is presented in each of Tables 1-4. The compounds described in the upper section of each of Tables 1-4 are compounds wherein the DiL4 data was ≤ 1 nM, the compounds in the lower section of each of Tables 1-4 had DiL4 data > 1 nM. The upper and lower sections of Tables 1-4 are separated by the shaded row. Examples with an asterisk (*) have HcL3 MED values ≤ 1 μ M. Examples with a carrot (^) have HcL3 MED values > 1 and ≤ 3.3 μ M. Compounds were named using Perkin Elmer ChemBioDraw Ultra 2014, Version 14.0.0.117; CambridgeSoft Corporation.

The following Formula (1A1) compounds described in Table 1 were prepared in accordance with the schemes and examples described herein. Each tabulated Formula (1A1) compound is a bis-substitution; *i.e.*, each of $(R)_n$ are the same; each of R^c are the same, and each of R^d are the same, and are as described in the Table.




Table 1. Formula (1A1) Compounds

Ex#	R^c	R^d	n	R	Mass
1-5	H	H	1	3-methoxy	1190
1-11	H	H	1	2-methoxy	1190
1-18	H	1H-pyrazol-1-yl	2	3,4-difluoro	1334
1-1	H	H	1	4-methoxy	1190

1-2	H	H	1	4-F	1165
1-3	H	H	1	4-Cl	1198
1-4	H	H	1	4-CN	1180
1-6	H	H	1	3-F	1165
1-7	H	H	0	--	1129
1-8	H	H	2	2-methoxy; 4-F	1198
1-9	H	-OH	1	4-methoxy	1222
1-10	H	-OH	1	2-methoxy	1222
1-12	H	-OH	2	3-F; 4-F	1233
1-13	H	H	2	3-F; 4-F	1201
1-14	H	-OH	1	4-F	1197
1-15	H	-OH	1	4-(1H-pyrazol-1-yl)	1294
1-16	H	-OH	1	4-(1H-imidazol-1-yl)	1294
1-17	H	H	1	2-methylsulfonyl	1286
1-19	H	H	1	4-(1H-pyrazol-1-yl)	1262
1-20	H	H	1	4-(1H-imidazol-1-yl)	1262
1-21	H	H	1	4-(1,2-dihydropyrimidin-5-yl)	1290
1-22	H	H	1	4-(1,6-dihydropyrimidin-5-yl)	1290
1-23	-OH	-CF ₃	1	3-isopropyl	1382
1-24	H	H	1	4-(pyridin-4-yl)	1284
1-25	-OH	-CF ₃	0	--	1297
1-26	H	H	2	3-fluoro; 4-methoxy	1225
1-27	H	H	1	4-methoxy	1175
1-28	H	H	1	4-methylsulfonyl	1286
1-29	H	-OH	1	4-(pyrimidin-5-yl)	1318
1-30	H	-OH	1	4-(pyrimidin-2-yl)	1318

The following compound names and example #'s refer to the upper section compounds of Table 1.

(1-5). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(3-methoxybenzyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(1-11). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(2-methoxybenzyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and

(1-18). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3,4-difluorophenyl)(1H-pyrazol-1-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone, stereoisomers thereof, and veterinary acceptable salts thereof.

NMR data (^1H NMR: 400 MHz, DMSO- d_6 , δ ppm) for the named Table 1 compounds are shown below:

(1-5). δ : 0.64-1.04 (m, 26 H), 1.14-1.79 (m, 16 H), 2.64-3.12 (m, 16 H), 3.70 (s, 6 H), 3.87 (s, 4 H), 4.38-5.72 (m, 8 H), 6.72-6.77 (m, 6 H), 7.12-7.25 (m, 10 H);

(1-11). δ : 0.63-0.98 (m, 26 H) 1.09-1.78 (m, 16 H) 2.66-3.07 (m, 20 H) 3.75-3.77 (m, 3 H) 3.85 (s, 3 H) 4.37 - 4.45 (m, 1 H) 4.99 - 5.72 (m, 7 H) 6.80 - 7.23 (m, 16 H); and

(1-18). δ : 0.63 - 1.10 (m, 26 H) 1.14 - 1.80 (m, 16 H) 2.63 - 3.13 (m, 16 H) 4.34 - 4.47 (m, 1 H) 4.97 - 5.76 (m, 7 H) 6.31 (s, 2 H) 6.91 - 6.97 (m, 2 H) 6.97 - 7.04 (m, 2 H) 7.07 - 7.22 (m, 6 H) 7.32 (br d, $J=7.82$ Hz, 4 H) 7.37 - 7.49 (m, 2 H) 7.55 (s, 2 H) 7.67 - 7.79 (m, 2 H).

The following Formula (2A1) compounds described in Table 2 were prepared in accordance with the schemes and examples described herein. Each of $(R)_n$, R^c , and R^d are as described in the Table.




Table 2. Formula (2A1) Compounds

Ex#	R^c	R^d	n	R	Mass
2-1*	H	H	0	--	1039
2-4^	H	H	1	4-F	1057

2-36	H	-N(CH ₂ CH ₃) ₂	2	3-F; 4-F	1146
2-37 [^]	H	-OCH ₃	2	3-F; 4-F	1105
2-38 [*]	H	-OCH ₂ CH ₃	2	3-F; 4-F	1119
2-42	H	morpholine	1	4-OCF ₃	1208
2-2	H	H	1	4-methoxy	1069
2-3	H	-OH	1	4-OCF ₃	1139
2-5	H	H	1	4-Cl	1074
2-6	H	H	1	4-CN	1064
2-7	H	H	1	3-F	1057
2-8	H	H	1	4-OCF ₃	1123
2-9 [*]	H	H	2	2-methoxy; 4-F	1087
2-10	H	-OH	1	4-methoxy	1085
2-11	H	-OH	1	2-methoxy	1085
2-12 [^]	H	H	1	2-methoxy	1069
2-13	H	-OH	1	4-OH	1071
2-14	H	H	2	3-F; 4-F	1075
2-15	H	-OH	1	3-OCF ₃	1139
2-16	H	-OH	1	4-Cl	1090
2-17	H	-OH	1	4-(1H-imidazol-1-yl)	1121
2-18	H	-OH	1	4-F	1073
2-19	H	-OH	1	4-(1H-pyrazol-1-yl)	1121
2-20	-OH	-CF ₃	1	4-methoxy	1153
2-21	H	-OH	1	3-(pyrimidin-5-yl)	1133
2-22	H	-OH	1	3-(pyrimidin-2-yl)	1133
2-23	H	H	1	4-(1H-pyrazol-1-yl)	1105
2-24	H	F	1	4-Cl	1092
2-25	H	H	1	4-(1H-imidazol-1-yl)	1105
2-26	-CH ₃	-OH	1	4-OCF ₃	1153
2-27	H	H	1	3-(1,6-dihydropyrimidin-2-yl)	1119
2-28	H	H	1	3-(1,2-dihydropyrimidin-5-yl)	1119
2-29	-OH	-CF ₃	1	3-isopropyl	1165
2-30	H	F	1	4-F	1075

2-31	H	-OH	1	3-(pyridin-4-yl)	1132
2-32	H	H	1	3-(pyridin-4-yl)	1116
2-33	H	-OH	1	3-OCF ₃	1139
2-34	H	-OH	1	2-OCF ₃	1139
2-35	H	-OH	1	4-OCF ₃	1139
2-39	-OH	-CF ₃	0	--	1123
2-40	H	H	2	3-F; 4-methoxy	1087
2-41	H	imidazolyl	1	4-OCF ₃	1189
2-43	H	pyrrolidin-1-yl	2	3-F; 4-F	1144

The following compound names and example #'s refer to the upper section compounds of Table 2.

(2-1). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-benzylbenzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(2-4). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-(4-fluorobenzyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(2-36). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((diethylamino)(3,4-difluorophenyl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(2-37). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3,4-difluorophenyl)(methoxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(2-38). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3,4-difluorophenyl)(ethoxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and

(2-42). (3S,6R,9S,12R,15S,18R,21S,24S)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-[[4-[(S)-morpholino-[4-(trifluoromethoxy)phenyl]methyl]phenyl]methyl]-1,7,13,19-tetraoxa-4,10,16,22-tetrazacyclotetracosane-2,5,8,11,14,17,20,23-octaone; stereoisomers thereof, and veterinary acceptable salts thereof.

NMR data (1H NMR: 400 MHz, DMSO-d₆, δ ppm) for the named Table 2 compounds are shown below:

(2-1). Not available

(2-4). Not available

(2-36). δ: 0.64 - 1.79 (m, 48 H) 2.58 - 3.22 (m, 21 H) 4.32 - 4.51 (m, 1 H) 4.96 - 5.82 (m, 7 H) 7.18 - 7.90 (m, 12 H);

(2-37). δ: 0.53 - 1.76 (m, 42 H) 2.58 - 3.18 (m, 16 H) 3.25 (s, 3 H) 4.29 - 4.59 (m, 1 H) 4.92 - 5.89 (m, 8 H) 7.03 - 7.51 (m, 12 H);

(2-38). δ: 0.65 - 1.82 (m, 45 H) 2.63 - 3.14 (m, 16 H) 3.35 - 3.47 (q, 3 H) 4.35 - 4.47 (m, 1 H) 4.98 - 5.79 (m, 8 H) 7.14 - 7.42 (m, 12 H); and

(2-42). δ: 0.64 - 1.79 (m, 42 H) 2.60 - 3.12 (m, 21 H) 3.58 - 4.57 (br, 5 H) 4.96 - 5.78 (m, 7 H) 7.20 - 7.67 (m, 13 H).

The following Formula (3A) compounds described in Table 3 were prepared in accordance with the schemes and examples described herein. Each tabulated Formula (3A) compound is a bis-substitution; *i.e.*, Ring A is the same as Ring B, each of (R)_n are the same; each of R^c are the same, and each of R^d are the same, and are as described in the Table.
















































Table 3. Formula (3A) Compounds












Ex#	Rings A and B with (R) _n	R ^c	R ^d	Mass
3-4*		H	H	1142
3-8*		H	H	1109
3-12 [^]		H	H	1144
3-17*		H	H	1144
3-18*		H	H	1111
3-21*		H	H	1168
3-24*		H	H	1137
3-31 [^]		H	H	1191











3-44 [^]		H	H	1170
3-49 [^]		H	H	1166
3-50 [*]		H	H	1159
3-56 [*]		H	H	1142
3-57 [*]		H	H	1111
3-58 [*]		H	H	1172
3-60		H	H	1159
3-69 [*]		H	H	1191
3-87 [*]		H	H	1210
3-110		H	H	1202
3-112 [*]		H	H	1212












3-118		H	H	1227
3-120*		H	H	1173
3-121^		H	H	1220
3-124*		H	H	1170
3-125		H	H	1299
3-132		H	-OH	1331
3-133		H	H	1220
3-148^		H	H	1111
3-155*		H	H	1172
3-156		H	H	1143
3-157		H	H	1137












3-165		H	H	1220
3-173 [^]		H	H	1109
3-176 [^]		H	H	1218
3-178		H	H	1304
3-193 [*]		H	H	1193
3-194		H	H	1167
3-207		H	-CH ₃	1220
3-209		H	H	1191
3-215 [*]		H	H	1192
3-217 [^]		H	H	1193
3-218		H	H	1193














3-220		F	-CH ₃	1255
3-223		H	H	1159
3-225		-CH ₃	F	1255
3-228		H	H	1230
3-238		H	H	1267
3-240		H	H	1188
3-243		H	H	1137
3-252		F	-CH ₃	1257
3-253*		H	H	1165
3-254		H	H	1143
3-259^A		H	-OCH ₃	1223













3-261		H	-CH ₃	1220
3-268*		H	H	1189
3-273*		H	H	1159
3-274*		H	-CH ₃	1201
3-275^		H	H	
3-282		H	H	1191
3-283*		H	H	1218
3-284		H	H	1191
3-285		H	F	1179
3-286		H	H	1201
3-287		H	H	1231













3-288*		H	H	1172
3-289^		H	H	1214
3-290*		H	H	1231
3-291*		H	-CH ₃	1172
3-292^		H	H	1302
3-293		H	H	1227
3-294*		H	H	1161
3-295^		H	H	1231
3-296^		H	H	1212
3-297		H	H	1263













3-307*		H	H	1181
3-308^		H	H	1223
3-309^		H	H	1167
3-310*		H	H	1189
3-313		H	H	1167
3-318		H	H	1269
3-319*		H	H	1199
3-321*		H	H	1233
3-322		H	H	1195
3-323*		H	H	1167
3-324*		H	H	1195













3-326		H	H	1203
3-329*		H	H	1286
3-332		H	H	1260
3-333*		H	H	1267
3-334*		H	H	1252
3-1		H	H	1166
3-2		H	H	1137
3-3^		H	H	1131
3-5		H	H	1176
3-6		H	H	1113
3-7		H	-OH	1224
3-9		H	H	1192












3-10		H	-OH	1178
3-11		H	-OH	1210
3-13		H	H	1139
3-14*		H	H	1146
3-15		H	-OH	1232
3-16		H	-OH	1358
3-19		H	H	1273
3-20		H	-OH	1392
3-22		H	H	1290
3-23		H	H	1146
3-25		H	H	1204
3-26		H	H	1139
3-27^A		H	H	1131














3-28		H	-OH	1264
3-29		H	H	1164
3-30		H	-OH	1242
3-32		H	H	1141
3-33		H	H	1194
3-34		-OH	-CF ₃	1310
3-35		H	H	1181
3-36*		H	H	1296
3-37		H	H	1200
3-38 [^]		H	H	1172
3-39		H	H	1172
3-40		H	H	1296













3-41		H	H	1228
3-42		H	H	1244
3-43		H	H	1210
3-45*		H	H	1144
3-46		H	H	1192
3-47		H	H	1204
3-48		H	-OH	1174
3-51		H	H	1137
3-52*		H	H	1200
3-53		H	-OH	1143
3-54		H	-OH	1204
3-55		H	-OH	1232













3-59		H	H	1139
3-61		H	H	1324
3-62		H	-OH	1278
3-63		H	-OH	1141
3-64		H	-OH	1171
3-65		H	-OH	1305
3-66		H	-OH	1204
3-67		H	H	1113
3-68		H	H	1252
3-70		H	H	1172
3-71		H	H	1198
3-72		H	-OH	1191











3-73*		H	H	1133
3-74		H	H	1248
3-75		H	-OH	1299
3-76^		H	H	1172
3-77		H	H	1220
3-78		H	-OH	1232
3-79		H	-OH	1284
3-80		H	-OH	1244
3-81^		H	H	1167
3-82		H	H	1279
3-83		H	-OH	1290
3-84		H	-OH	1244












3-85		H	H	1294
3-86		H	-OH	1328
3-88		H	-OH	1328
3-89		H	-OH	1242
3-90		H	H	1198
3-91		H	H	1310
3-92		H	-OH	1232
3-93		H	-OH	1234
3-94		H	-OH	1223
3-95		H	H	1267
3-96		H	-OH	1230













3-97		H	-OH	1204
3-98		H	H	1356
3-99		H	-OH	1167
3-100		H	-OH	1342
3-101		H	-OH	1199
3-102		H	H	1210
3-103		H	-OH	1204
3-104		H	-OH	1177
3-105		H	-OH	1242
3-106		H	-OH	1169
3-107		H	H	1218
3-108		H	H	1200
3-109		H	H	1302













3-111		H	H	1274
3-113		H	-OH	1230
3-114		H	H	1212
3-115^		H	H	1252
3-116		H	H	1262
3-117		H	H	1296
3-119*		H	H	1209
3-122		H	H	1286
3-123		H	H	1228
3-126*		H	-OH	1205
3-127		H	-OH	1311
3-128		H	-OH	1229











3-129		H	-OH	1260
3-130		H	-OH	1296
3-131		H	-OH	1294
3-134		H	H	1327
3-135		H	H	1327
3-136		H	H	1224
3-137		H	H	1314
3-138		H	H	1270
3-139		H	-OH	1328
3-140		H	H	1279
3-141		H	H	1189
3-142		H	H	1202












3-143		H	H	1220
3-144		H	H	1191
3-145		H	-OH	1223
3-146		H	H	1139
3-147		H	H	1290
3-149		H	-OH	1175
3-150		H	H	1244
3-151		H	-OH	1252
3-152		H	-OH	1234
3-153		H	H	1258
3-154		H	H	1265













3-158		H	H	1202
3-159^		H	H	1172
3-160		H	H	1135
3-161		H	H	1224
3-162		H	H	1213
3-163		H	H	1260
3-164		H	H	1263
3-166		H	H	1244
3-167		H	H	1265
3-168		H	H	1194
3-169		H	H	1264













3-170		H	H	1273
3-171		H	H	1191
3-172		H	H	1131
3-174		H	H	1255
3-175		H	H	1159
3-177		H	H	1133
3-179		H	H	1299
3-180		H	H	1189
3-181		H	H	1226
3-182		H	H	1232
3-183		H	H	1221
3-184		H	H	1159











3-185		H	H	1272
3-186		H	H	1191
3-187		H	H	1133
3-188		H	H	1191
3-189		H	H	1197
3-190		H	H	1211
3-191		H	H	1220
3-192^A		H	H	1191
3-195		H	H	1290
3-196		H	H	1250
3-197		H	H	1193
3-198		H	H	1202












3-199		H	H	1272
3-200		H	H	1244
3-201		H	H	1221
3-202		H	H	1159
3-203		H	H	1193
3-204		H	H	1267
3-205		H	H	1232
3-206		H	H	1246
3-208		H	H	1248
3-210		H	H	1177
3-211 [^]		H	H	1221
3-212		H	H	1179













3-213		H	H	1189
3-214		H	H	1205
3-216		H	H	1230
3-219		H	H	1253
3-221		H	H	1274
3-222		H	H	1133
3-224		H	H	1253
3-226		H	H	1173
3-227		-CH ₃	F	1181
3-229		H	H	1250
3-230		H	H	1209
3-231		H	H	1139


3-232		H	H	1237
3-233		H	H	1137
3-234		H	H	1161
3-235		H	H	1211
3-236		H	H	1172
3-237		H	H	1218
3-239		H	H	1144
3-241		H	H	1145
3-242		H	H	1316
3-244		H	H	1189
3-245		H	H	1194
3-246		H	H	1137

3-247		H	H	1344
3-248		H	H	1200
3-249		H	H	1137
3-250		H	H	1195
3-251		H	H	1218
3-255		H	H	1214
3-256		H	H	1288
3-257		H	H	1115
3-258		H	H	1228
3-260		H	H	1221
3-262		H	H	1171
3-263		H	H	1228

3-264		H	H	1115
3-265		H	H	1248
3-266		H	H	1248
3-267		H	H	1163
3-269		H	H	1274
3-270		H	H	1191
3-271		H	H	1189
3-272		H	H	1191
3-276		H	H	1238
3-277		H	H	1220
3-278		H	H	1165

3-279		H	H	1193
3-280		H	H	1137
3-281		H	H	1209
3-298^		H	H	1139
3-299*		H	H	1229
3-300^		H	H	1217
3-301^		H	H	1109
3-302		H	F	1227
3-303*		H	H	1193
3-304		H	H	1189
3-305*		H	H	1193

3-306*		H	H	1211
3-311		H	H	1243
3-312		H	H	1181
3-314		H	H	1269
3-315		H	H	1301
3-316		H	H	1211
3-317		H	H	1211
3-320		H	H	1231
3-325		H	H	1195
3-327		H	H	1159
3-328		H	H	1213
3-330		H	-OH	1204

3-331		H	H	1200
-------	---	---	---	------

The following compound names and example #'s refer to the upper section compounds of Table 3.

- (3-4). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiophen-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-8). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(furan-2-ylmethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-12). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiazol-5-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-17). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiazol-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-18). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(oxazol-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-21). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3,5-dimethylisoxazol-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-24). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((1-methyl-1H-pyrazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-31). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((2-methoxypyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

- (3-44). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((3-methylthiophen-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-49). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1,3-dimethyl-1H-pyrazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-50). 2,2'-((((2R,5S,8R,11S,14R,17S,20R,23S)-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octaoxo-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,14-diyl)bis(methylene))bis(4,1-phenylene))bis(methylene))bis(furan-3-carbonitrile);
- (3-56). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiophen-3-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-57). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(oxazol-4-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-60). 4,4'-((((2R,5S,8R,11S,14R,17S,20R,23S)-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octaoxo-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,14-diyl)bis(methylene))bis(4,1-phenylene))bis(methylene))bis(furan-2-carbonitrile);
- (3-69). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((6-methoxypyridin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-87). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-chlorothiophen-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-110). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((3-methoxythiophen-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-112). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-chlorothiazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

- (3-118). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-fluoro-6-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-120). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methyl-1,2,3-thiadiazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-121). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((6-methoxy-2-methylpyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-124). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylthiophen-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-125). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-bromothiophen-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-132). (3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-((R)-(3-bromothiophen-2-yl)(hydroxy)methyl)benzyl)-18-(4-((S)-(3-bromothiophen-2-yl)(hydroxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-133). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((6-methoxy-4-methylpyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-148). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(oxazol-5-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-155). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylthiazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-156). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(isothiazol-5-ylmethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

- (3-157). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((5-methylfuran-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-165). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-ethoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-173). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(furan-3-ylmethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-176). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-(dimethylamino)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-178). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-morpholinopyrimidin-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-193). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((3-methoxypyrazin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-194). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-fluoropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-207). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(1-(6-methoxypyridin-3-yl)ethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-209). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((2-methoxy-4-methylthiazol-5-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-215). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((3-methoxypyridin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-217). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((4-methoxypyrimidin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

- (3-218). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((6-methoxypyrazin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-220). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(1-fluoro-1-(2-methoxypyridin-3-yl)ethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-223). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylpyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-225). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(1-fluoro-1-(6-methoxypyridin-2-yl)ethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-228). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((4-isopropyl-1,2,3-thiadiazol-5-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-238). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-(trifluoromethyl)pyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-240). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,6-dimethylpyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-243). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((5-methyl-1H-pyrazol-4-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-252). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(1-fluoro-1-(5-methoxypyrazin-2-yl)ethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-253). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1-ethyl-1H-pyrazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

- (3-254). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(isothiazol-4-ylmethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-259). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-hydroxypyridin-3-yl)(methoxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-261). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(1-(2-methoxypyridin-3-yl)ethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-268). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1-cyclopropyl-1H-pyrazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-273). (3S,6R,9R,12R,15R,18R,21S,24S)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis[[4-[(2-methyl-3-pyridyl)methyl]phenyl]methyl]-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,5,8,11,14,17,20,23-octaone;
- (3-274). (3R,6R,9R,12R,15S,18S,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((5-methoxypyrazin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-275). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1-(cyclopropylmethyl)-1H-pyrazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-282). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((4-methoxypyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-283). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((4-isopropylpyrimidin-5-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-284). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-cyclopropylisoxazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (3-285). (3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-((R)-fluoro(thiazol-2-yl)methyl)benzyl)-18-(4-((S)-fluoro(thiazol-2-yl)methyl)benzyl)-3,9,15,21-

tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-286). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1-(2-fluoroethyl)-1H-pyrazol-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-287). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-(difluoromethyl)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-288). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylisothiazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-289). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-cyclopropylpyrimidin-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-290). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-(difluoromethyl)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-291). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((R)-1-(thiazol-2-yl)ethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-292). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-morpholinopyridin-4-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-293). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-fluoro-2-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-294). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylpyrimidin-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-295). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-(difluoromethyl)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-296). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-cyclopropylpyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and

(3-297). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-(difluoromethoxy)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone,

(3-307). 3,3'-((((2R,5S,8R,11S,14R,17S,20R,23S)-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octaoxo-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,14-diyl)bis(methylene))bis(4,1-phenylene))bis(methylene))diisonicotinonitrile;

(3-308). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-(methylthio)pyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-309). (3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-((3-fluoropyridin-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-((3-methylpyridin-4-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-310). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-ethylpyrimidin-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-313). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-fluoropyridin-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-318). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-(trifluoromethyl)pyrimidin-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-319). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-chloropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-321). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-(difluoromethyl)pyrimidin-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-322). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-fluoro-4-methylpyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-323). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-fluoropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-324). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,5-difluoropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-326). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,3-difluoropyridin-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-329). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-bromopyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-332). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-chloro-2-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-333). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-(trifluoromethyl)pyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3-334). (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,3-dimethoxypyridin-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone, stereoisomers thereof, and veterinary acceptable salts thereof.

NMR data (¹H NMR: 400 MHz, DMSO-d₆, δ ppm) for the named Table 3 compounds are shown below:

(3-4). δ: 6.8-7.4 (m, 14H), 5.0-5.8 (m, 8H), (4.4 (m, 1H)), 4.1 (s, 4H), 2.6-3.2 (m, 16H), 1.1-1.8 (m, 13H), 0.6-1.0 (m, 30H);

(3-8). δ: 7.5 (s, 2H), 7.1-7.4 (m, 8H), 6.35 (s, 2H), 6.1 (s, 2H), 5.0-5.8 (m, 8H), 4.4 (m, 0.68H), 3.9 (s, 4H), 2.7-3.1 (m, 16H), 1.1-1.8 (m, 18H), 0.6-1.0 (m, 28H);

- (3-12). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.66 - 3.06 (m, 16H), 4.17 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 7.18 - 7.26 (m, 8H), 7.70 (brs, 2H), 8.92 (s, 2H);
- (3-17). δ : 0.70 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.67 - 3.04 (m, 16H), 4.30 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 7.26 - 7.28 (m, 8H), 7.57(d, J= 2.8 Hz, 2H), 7.70 (d, J= 2.8 Hz, 2H);
- (3-18). δ : 0.70 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.67 - 3.04 (m, 16H), 4.09 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 7.12 (brs, 2H), 7.18 - 7.28 (m, 8H), 7.98 (brs, 2H);
- (3-21). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 1.99 (s, 6H), 2.31 (s, 6H), 2.66 - 3.02 (m, 16H), 3.65 (s, 4H), 4.40-4.42 (m, 1H), 5.02-5.69 (m, 7H), 7.06-7.23 (m, 8H);
- (3-24). δ : 7.2-7.35 (m, 6H), 7.1-7.2 (m, 4H), 5.9-5.95 (m, 2H), 5.0-5.8 (m, 8H), 4.4 (m, 0.8H), 4.0 (s, 4H), 3.65 (s, 6H), 2.7-3.1 (m, 15H), 1.1-1.8 (m, 16H), 0.6-1.0 (m, 29H);
- (3-31). δ : 0.62 - 1.79 (m, 42 H) 2.65 - 3.10 (m, 16 H) 3.80 - 3.90 (m, 10 H) 4.36 - 4.46 (m, 1 H) 4.99 - 5.74 (m, 7 H) 6.84 - 6.96 (m, 2 H) 7.09 - 7.17 (m, 4 H) 7.18 - 7.27 (m, 4 H) 7.37 - 7.47 (m, 2 H) 7.96 - 8.06 (m, 2 H);
- (3-44). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.12 (s, 6H), 2.66 - 3.06 (m, 16H), 4.01 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 6.82-6.83 (m, 2H), 7.11 - 7.22 (m, 10H);
- (3-49). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.04 (s, 6H), 2.66 - 3.06 (m, 16H), 3.56 (s, 6H), 3.90 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 9H), 7.11 - 7.25 (m, 8H);
- (3-50). δ : 0.67 - 0.94 (m, 26H), 1.23 - 1.71 (m, 16H), 2.67 - 3.05 (m, 16H), 4.15 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 6.89 (brs, 2H), 7.15-7.18 (m, 4H), 7.27-7.30 (m, 4H), 7.78 (brs, 2H);
- (3-56). δ : 0.68 - 0.95 (m, 26H), 1.23 - 1.69 (m, 16H), 2.67 - 3.05 (m, 16H), 3.90 (s, 4H), 4.40 - 4.42 (m, 1H), 5.03 - 5.70 (m, 7H), 6.91-7.43 (m, 14H);
- (3-57). δ : 0.70 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.67 - 3.04 (m, 16H), 3.79 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.73 (m, 7H), 7.17 - 7.21 (m, 8H), 7.79-7.81 (m, 2H), 8.26 (s, 2H);
- (3-58). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.39 (s, 6H), 2.66 - 3.06 (m, 16H), 3.99 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.69 (m, 7H), 7.11 - 7.21 (m, 8H), 8.78 (s, 2H);

- (3-60). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.66 - 3.02 (m, 16H), 4.05 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 6.45 (brs, 2H), 7.17 - 7.36 (m, 8H), 7.52 (brs, 2H);
- (3-69). δ : 0.61 - 0.99 (m, 27 H), 1.15 - 1.76 (m, 15 H), 2.64 - 2.94 (m, 12 H), 2.96 - 3.11 (m, 3 H), 3.82 (s, 6 H), 3.95 (s, 4 H), 4.36 - 4.48 (m, 1 H), 4.99 - 5.54 (m, 6 H), 5.63 - 5.75 (m, 2 H), 6.61 (d, J = 8.2 Hz, 2 H) 6.73 - 6.82 (m, 2 H), 7.13 - 7.30 (m, 1 H), 7.23 (s, 7 H), 7.54 - 7.64 (m, 2 H);
- (3-87). δ : 0.68 - 0.95 (m, 26H), 1.23 - 1.70 (m, 16H), 2.67 - 3.05 (m, 16H), 4.07 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.73 (m, 7H), 7.01(d, J= 5.2Hz, 2H), 7.15-7.17(m, 4H), 7.24-7.27 (m, 4H), 7.48-7.50 (m, 2H);
- (3-110). δ : 0.67-0.94 (m, 27H), 1.23-1.68 (m, 15H), 2.67-3.10 (m, 16H), 3.77 (s, 6H), 3.91 (s, 4H), 4.39-4.44 (m, 1H), 5.01-5.70 (m, 7H), 6.96-6.99 (m, 2H), 7.10-7.25 (m, 10H);
- (3-112). δ : 0.68 - 0.95 (m, 26H), 1.24 - 1.70 (m, 16H), 2.67 - 3.05 (m, 16H), 4.12 (s, 4H), 4.40 - 4.42 (m, 1H), 5.04 - 5.70 (m, 7H), 7.17-7.27 (m, 8H), 8.98 (s, 2H);
- (3-118). δ : 0.68 - 0.95 (m, 26H), 1.24 - 1.69 (m, 16H), 2.67 - 3.05 (m, 16H), 3.87 (s, 4H), 3.89 (s, 6H), 4.40 - 4.42 (m, 1H), 5.04 - 5.70 (m, 7H), 7.17-7.24 (m, 8H), 7.46-7.50 (m, 2H), 7.88 (s, 2H);
- (3-120). δ : 0.70 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.60 (s, 6H), 2.67 - 3.04 (m, 16H), 4.29 (s, 4H), 4.40 - 4.42 (m, 1H), 5.03 - 5.71 (m, 7H), 7.20-7.28 (m, 8H);
- (3-121). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.26 (s, 6H), 2.66 - 3.02 (m, 16H), 3.76 (s, 6H), 3.84 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 6.54-6.56 (m, 2H), 7.02-7.20 (m, 8H), 7.38-7.40 (m, 2H);
- (3-124). δ 0.65-0.94 (m, 27H), 1.02-1.80 (m, 15H), 2.12 (s, 6H), 2.67-3.10 (m, 16H), 4.03 (bs, 4H), 4.42-4.46 (m, 1H), 5.02-5.69 (m, 7H), 6.65 (s, 2H), 6.83-6.87 (m, 2H), 7.11-7.23 (m, 8H);
- (3-125). δ 0.68-0.94 (m, 27H), 1.23-1.75 (m, 15H), 2.68-3.08 (m, 16H), 4.07 (s, 4H), 4.40-4.45 (m, 1H), 5.02-5.73 (m, 7H), 7.04-7.27 (m, 10H), 7.50-7.51 (m, 2H);
- (3-132). δ 0.68-0.94 (m, 26H), 1.20-1.78 (m, 16H), 2.66-3.10 (m, 16H), 4.36-4.42 (m, 1H), 5.02-5.72 (m, 7H), 5.88-5.91(m, 2H), 6.40-6.50 (m, 2H), 6.99 (d, J = 5 Hz, 2H), 7.25-7.38 (m, 8H), 7.54-7.60 (m, 2H);
- (3-133). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.08 (s, 6H), 2.66 - 3.02 (m, 16H), 3.79 (s, 6H), 3.86 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 6.61 (s, 2H), 7.04-7.22 (m, 8H), 7.89 (s, 2H);

- (3-148). δ : 0.70 – 0.94 (m, 26H), 1.23 - 1.68 (m, 16H), 2.67 - 3.04 (m, 16H), 4.00 (s, 4H), 4.40 - 4.42 (m, 1H), 5.03 - 5.71 (m, 7H), 6.88 (s, 2H), 7.16-7.25 (m, 8H), 8.21 (s, 2H);
- (3-155). δ : 8.76-8.95 (m, 2H), 7.02-7.35 (m, 8H), 4.95-5.81 (m, 10H), 4.36-4.45 (m, 1H), 4.02-4.12 (m, 4H), 2.59-3.19 (m, 16H), 2.26-2.42 (m, 6H), 1.10-1.80 (m, 16H), 0.62-1.01 (m, 26H);
- (3-156). δ : 0.68 - 0.95 (m, 26H), 1.21 - 1.70 (m, 16H), 2.68 - 3.05 (m, 16H), 4.28 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 7.18 - 7.30 (m, 10H), 8.40-8.42 (m, 2H);
- (3-157). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.17 (s, 6H), 2.66 - 3.06 (m, 16H), 3.85 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.69 (m, 7H), 5.91-5.93 (m, 4H), 7.12 - 7.23 (m, 8H);
- (3-165). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 22H), 2.66 - 3.06 (m, 16H), 3.82 (s, 4H), 4.29 (q, J = 7 Hz, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.69 (m, 7H), 6.85-6.89 (m, 2H), 7.15 - 7.23 (m, 8H), 7.43-7.46 (m, 2H), 7.97-7.99 (m, 2H);
- (3-173). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.66 - 3.06 (m, 16H), 3.69 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 6.28-6.29 (m, 2H), 7.11 - 7.24 (m, 8H), 7.43-7.44 (m, 2H), 7.55 (brs, 2H);
- (3-176). δ : 0.68 – 0.95 (m, 26H), 1.24 – 1.69 (m, 16H), 2.67 - 3.05 (m, 28H), 3.95 (s, 4H), 4.40 - 4.42 (m, 1H), 5.04 - 5.72 (m, 7H), 6.83-6.87 (m, 2H), 7.08-7.10 (m, 4H), 7.19-7.31 (m, 6H), 8.08-8.09 (m, 2H);
- (3-178). δ : 8.16-8.40 (m, 4H), 7.03-7.48 (m, 8H), 4.99-5.82 (m, 8H), 4.33-4.44 (m, 1H), 3.70-3.86 (m, 4H), 3.3-3.5 (m, 4H), 2.62-3.14 (m, 16H), 1.02-1.88 (m, 16H), 0.61-0.97 (m, 27H);
- (3-193). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.66 - 3.06 (m, 16H), 3.89 (s, 6H), 4.04 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 7.13 - 7.21 (m, 8H), 8.07-8.09 (m, 4H);
- (3-194). δ : 8.21-8.68 (m, 4H), 7.48-7.81 (m, 2H), 7.12-7.42 (m, 8H), 4.96-5.80 (m, 8H), 4.32-4.55 (m, 1H), 3.76-4.08 (m, 4H), 2.63-3.17 (m, 16H), 1.07-1.79 (m, 16H), 0.62-1.01 (m, 26H);
- (3-207). δ : 0.61 - 1.01 (m, 26 H), 1.13 - 1.77 (m, 16 H), 2.64 - 3.10 (m, 16 H), 3.75 (s, 6 H), 4.12 (m, 2 H), 4.37 - 4.46 (m, 1 H), 4.98 - 5.74 (m, 8 H), 6.72 (d, 2 H), 7.15 - 7.28 (m, 8 H), 7.54 (d, 2 H), 8.02 - 8.11 (m, 2 H);

- (3-209). δ : 0.68 – 0.95 (m, 26H), 1.24 – 1.69 (m, 16H), 2.17 (s, 6H), 2.67 - 3.05 (m, 16H), 3.90 (s, 10H), 4.40 - 4.42 (m, 1H), 5.04 - 5.69 (m, 7H), 7.11-7.13 (m, 4H), 7.22-7.25 (m, 4H);
- (3-215). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 16H), 2.67 - 3.05 (m, 16H), 3.76 (s, 6H), 4.00 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.68 (m, 7H), 7.08-7.21 (m, 10H), 7.32-7.35 (m, 2H), 8.00-8.02 (m, 2H);
- (3-217). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 16H), 2.67 - 3.05 (m, 16H), 3.88 (s, 6H), 4.05 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.68 (m, 7H), 6.74-6.76 (m, 2H), 7.23-7.25 (m, 8H), 8.39-8.40 (m, 2H);
- (3-218). δ : 0.69 -0.94 (m, 26H), 1.23 -1.70 (m, 16H), 2.67 - 3.05 (m, 16H), 3.87 (s, 6H), 4.00 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.23-7.25 (m, 8H), 8.09-8.12 (m, 4H);
- (3-220). δ : 0.65 -0.94 (m, 26H), 1.23 -1.68 (16H), 2.01 (s, 3H), 2.07 (3H), 2.67 -3.04 (m, 16H), 3.67 (s, 6H), 4.41-4.43 (m, 1H), 5.03 -5.70 (m, 7 H), 7.05 -7.08 (m, 2H), 7.25 -7.29 (m, 8H), 7.85 -7.87 (m, 2 H), 8.10 -8.15 (m, 2H);
- (3-223). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 16H), 2.16 (s, 6H), 2.67 - 3.05 (m, 16H), 3.96 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.68 (m, 7H), 7.06-7.24 (m, 10H), 8.28-8.30 (m, 4H);
- (3-225). δ : 0.63 - 0.92 (m, 26 H), 1.23 - 1.69 (m, 16 H), 1.98 (s, 3 H), 2.04 (s, 3 H), 2.67 - 3.04 (m, 16 H), 3.83 (s, 6 H), 4.38 - 4.43 (m, 1 H), 5.01 - 5.73 (m, 7 H), 6.72 (d, $J = 8.2$ Hz, 2 H), 7.09 (d, $J = 7.0$ Hz, 2 H), 7.30 - 7.31 (m, 4 H), 7.40 - 7.42 (m, 4 H), 7.71 (t, $J = 7.8$ Hz, 2 H);
- (3-228). δ : 7.15-7.45 (m, 8H), 4.94-5.77 (m, 8H), 4.37-4.53 (m, 1H), 4.24-4.35 (m, 4H), 3.24-3.49 (m, 2H), 2.60-3.12 (m, 15H), 1.14-1.84 (m, 28H), 0.61-1.07 (m, 26H);
- (3-238). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 16H), 2.67 - 3.05 (m, 16H), 4.15 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.07-7.26 (m, 8H), 7.65-7.67(m, 2H), 7.80-7.82 (m, 2H), 8.60-8.61 (m, 2H);
- (3-240). δ : 0.69-0.95 (m, 26H), 1.20-1.74 (m, 16H), 2.60-3.07 (m, 28H), 4.10 (s, 4H), 4.39-5.72 (m, 8H), 7.14 (d, 4H), 7.28 (d, 4H), 7.69 (d, 2H), 8.17 (d, 2H);
- (3-243). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 16H), 2.05 (s, 6H), 2.67 - 3.05 (m, 16H), 3.66 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.07-7.20 (m, 10H), 12.25 (brs, 2H);

(3-252). δ : 0.64 - 0.92 (m, 26 H), 1.19 - 1.57 (m, 16 H), 2.00 (s, 3 H), 2.06 (s, 3 H), 2.67 - 3.05 (m, 16 H), 3.90 (s, 6 H), 4.42 - 4.48 (m, 1 H), 5.02 - 5.72 (m, 7 H), 7.31 - 7.33 (m, 8 H), 8.28 - 7.32 (m, 4 H);

(3-253). δ : 0.64 - 0.99 (m, 27 H), 1.09 - 1.77 (m, 21 H), 2.64 - 2.94 (m, 13 H), 3.03 (br d, 3 H), 3.93 - 4.05 (m, 8 H), 4.99 - 5.77 (m, 7 H), 5.60 - 5.76 (m, 1 H), 5.90 - 5.98 (m, 2 H), 7.06 - 7.19 (m, 4 H), 7.24 - 7.36 (m, 6 H);

(3-254). δ : 0.69 - 0.94 (m, 26H), 1.23 - 1.68 (m, 16H), 2.67 - 3.05 (m, 16H), 4.01 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.18-7.25 (m, 8H), 8.42 (s, 2H), 8.67 (s, 2H);

(3-259). Not available

(3-261). δ : 0.67 - 0.93 (m, 26H), 1.23 - 1.68 (22 H), 2.67 - 2.88 (s, 12H), 2.99 - 3.02 (m, 4H), 3.81 (s, 6H), 4.30 - 4.42 (m, 3H), 5.03 - 5.67 (m, 7H), 6.91 - 6.94 (m, 2H), 7.12 - 7.21 (m, 8H), 7.51 - 7.54 (m, 2H), 7.99 - 8.01 (m, 2H);

(3-268). δ : 0.67-1.01 (m, 34H), 1.20-1.75 (m, 16H), 2.68-3.08 (m, 16H), 3.35-3.40 (m, 2H), 4.07 (s, 4H), 4.39-5.71 (m, 8H), 5.91 (s, 2H), 7.16 (d, 4H), 7.23-7.28 (m, 6H);

(3-273). δ : 0.69 - 0.94 (m, 26H), 1.23 - 1.68 (m, 16H), 2.36 (s, 6H), 2.67 - 3.05 (m, 16H), 3.94 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.06-7.25 (m, 10H), 7.43-7.45 (m, 2H), 8.28-8.29 (m, 2H);

(3-274). δ : 0.67 - 0.92 (m, 26 H), 1.23 - 1.66 (m, 22 H), 2.67 - 3.00 (m, 16 H), 3.85 (s, 6 H), 4.26 - 4.28 (m, 2 H), 4.37 - 4.43 (m, 1 H), 5.02 - 5.69 (m, 7 H), 7.22 (brs, 8 H), 8.11 (brs, 2 H), 8.23 (s, 2 H);

(3-275). δ : 0.22-0.28 (m, 4H), 0.38-0.44 (m, 4H), 0.69-1.01 (m, 26H), 1.06-1.15 (m, 2H), 1.21-1.76 (m, 16H), 2.68-3.07 (m, 16H), 3.86 (d, 4H), 4.00 (s, 4H), 4.40-5.70 (m, 8H), 5.88-5.91 (m, 2H), 7.14 (d, 4H), 7.24-7.28 (d, 4H), 7.30-7.33 (m, 2H);

(3-282). Not available

(3-283). δ : 0.69-1.08 (m, 38H), 1.18-1.74 (m, 16H), 2.66-3.07 (m, 16H), 3.19-3.27 (m, 2H), 4.02 (s, 4H), 4.38-5.69 (m, 8H), 7.11 (d, 4H), 7.23-7.28 (m, 4H), 8.56 (s, 2H), 9.00 (s, 2H);

(3-284). δ : 0.61 - 0.97 (m, 34H), 1.23 - 1.93 (m, 18H), 2.67 - 3.05 (m, 16H), 4.01 (s, 4H), 4.40 - 4.42 (m, 1H), 5.08 - 5.70 (m, 7H), 5.91-5.93 (m, 2H), 7.18-7.27 (m, 8H);

(3-285). δ : 0.69 - 0.94 (m, 26H), 1.23 - 1.68 (m, 16H), 2.67 - 3.05 (m, 16H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 6.95 (d, J = 44 Hz, 2H), 7.41 (brs, 8H), 7.86 (brs, 4H);

(3-286). δ : 0.67-0.96 (m, 26H), 1.18-1.76 (m, 16H), 2.68-3.07 (m, 16H), 3.72 (s, 4H), 4.28-5.72 (m, 16H), 7.14 (d, 4H), 7.19-7.27 (m, 6H), 7.50 (s, 2H);

(3-287). δ : 0.66-0.94 (m, 26H), 1.15-1.76 (m, 16H), 2.67-3.07 (m, 16H), 4.02 (s, 4H), 4.39-5.73 (m, 8H), 6.90 (t, 2H), 7.19-7.28 (m, 8H), 7.61 (d, 2H), 7.80 (d, 2H), 8.58 (s, 2H);

(3-288). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 16H), 2.17 (s, 6H), 2.67 - 3.05 (m, 16H), 4.15 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.18-7.29 (m, 8H), 8.30 (s, 2H);

(3-289). δ : 0.66-1.05 (m, 34H), 1.18-1.73 (m, 16H), 2.19-2.26 (m, 2H), 2.68-3.06 (m, 16H), 4.11 (s, 4H), 4.39-5.70 (m, 8H), 7.16 (d, 4H), 7.23-7.28 (m, 4H), 8.45-8.48 (m, 2H), 8.84 (s, 2H);

(3-290). δ : 0.67-0.97 (m, 26H), 1.17-1.76 (m, 16H), 2.68-3.07 (m, 16H), 4.16 (s, 4H), 4.39-5.71 (m, 8H), 6.96-7.28 (m, 10H), 7.49-7.53 (m, 2H), 7.69 (d, 2H), 8.52-8.54 (m, 2H);

(3-291). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 22H), 2.67 - 3.05 (m, 16H), 4.40 - 4.42 (m, 1H), 4.51-4.53 (m, 2H), 5.02 - 5.70 (m, 7H), 7.27 (brs, 8H), 7.54-7.56 (m, 2H), 7.69-7.71 (m, 2H);

(3-292). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 16H), 2.67 - 3.05 (m, 16H), 3.37-3.39 (m, 8H), 3.65-3.68 (m, 8H), 3.81 (s, 4H), 4.40 - 4.42 (m, 1H), 5.07 - 5.70 (m, 7H), 6.47-6.48 (m, 2H), 6.73 (s, 2H), 7.16-7.23 (m, 8H), 7.97-7.98 (m, 2H);

(3-293). δ : 0.67-0.95 (m, 26H), 1.19-1.76 (m, 16H), 2.68-3.06 (m, 16H), 3.83-3.86 (m, 10H), 4.39-5.72 (m, 8H), 7.16-7.26 (m, 8H), 7.38-7.43 (m, 2H), 7.98-8.00 (m, 2H);

(3-294). δ : 0.68-1.01 (m, 26H), 1.18-1.74 (m, 16H), 2.36 (s, 6H), 2.68-3.07 (m, 16H), 3.98 (s, 4H), 4.39-5.71 (m, 8H), 7.12 (d, 4H), 7.24-7.28 (m, 4H), 8.49-8.52 (m, 2H), 8.90 (s, 2H);

(3-295). δ : 0.68-1.00 (m, 26H), 1.18-1.73 (m, 16H), 2.67-3.07 (m, 16H), 4.14 (s, 4H), 4.38-5.71 (m, 8H), 7.13-7.40 (m, 10H), 7.56 (d, 2H), 8.53 (s, 2H), 8.62 (d, 2H);

(3-296). δ : 0.68-1.06 (m, 34H), 1.18-1.74 (m, 16H), 2.21-2.30 (m, 2H), 2.68-3.07 (m, 16H), 4.18 (s, 4H), 4.39-5.71 (m, 8H), 7.15 (d, 4H), 7.24-7.39 (m, 6H), 7.73-7.80 (m, 2H), 8.37 (d, 2H);

(3-297). δ : 0.67-0.96 (m, 26H), 1.18-1.76 (m, 16H), 2.68-3.07 (m, 16H), 3.90 (s, 4H), 4.39-5.71 (m, 8H), 7.14-7.27 (m, 10H), 7.52-7.89 (m, 4H), 8.09-8.13 (m, 2H);

- (3-307). δ : 0.64 - 1.01 (m, 27 H), 1.15 - 1.74 (m, 15 H), 2.60 - 2.97 (m, 13 H), 2.97 - 3.10 (m, 3 H), 3.86 - 4.24 (m, 4 H), 4.31 - 4.49 (m, 1 H), 4.95 - 5.80 (m, 9 H), 7.13 - 7.36 (m, 8 H), 7.76 - 7.96 (m, 2 H), 8.48 - 9.01 (m, 2 H), 8.56 - 8.86 (m, 2 H);
- (3-308). δ : 0.64 - 1.03 (m, 27 H), 1.17 - 1.74 (m, 15 H), 2.51 - 2.56 (m, 3 H), 2.60 - 2.96 (m, 13 H), 2.96 - 3.12 (m, 3 H), 3.90 (s, 4 H), 4.41 (br dd, 1 H), 4.96 - 5.77 (m, 8 H), 7.01 - 7.16 (m, 6 H), 7.16 - 7.28 (m, 4 H), 7.35-7.45 (m, 2 H), 8.30-8.40 (m, 2 H);
- (3-309). δ : 0.65 - 1.01 (m, 27 H), 1.12 - 1.74 (m, 15 H), 2.63 - 2.98 (m, 13 H), 2.98 - 3.09 (m, 3 H), 3.92 - 4.10 (m, 4 H), 4.34 - 4.49 (m, 1 H), 4.95 - 5.76 (m, 7 H), 7.18 (br d, 4 H), 7.26 (br d, 4 H), 7.34 (t, 2 H), 8.28 - 8.41 (m, 2 H), 8.44 - 8.57 (m, 2 H);
- (3-310). δ : 0.62 - 1.03 (m, 28 H), 1.04 - 1.14 (m, 6 H), 1.17 - 1.80 (m, 16 H), 2.59 - 2.96 (m, 17 H), 2.96 - 3.11 (m, 3 H), 3.89 - 4.06 (m, 4 H), 4.33 - 4.48 (m, 1 H), 4.98 - 5.75 (m, 7 H), 7.03 - 7.19 (m, 4 H), 7.19 - 7.33 (m, 4 H), 8.48 - 8.57 (m, 2 H), 8.91 - 9.01 (m, 2 H);
- (3-313). δ : 0.41 - 1.00 (m, 27 H), 1.06 - 1.80 (m, 15 H), 2.52 - 3.17 (m, 1 H), 2.58 - 2.96 (m, 12 H), 2.96 - 3.14 (m, 3 H), 4.00 (s, 4 H), 4.36 - 4.48 (m, 1 H), 4.98 - 5.77 (m, 7 H), 6.87 - 7.04 (m, 2 H), 7.11 - 7.50 (m, 10 H), 7.82 - 7.96 (m, 2 H);
- (3-318). δ : 0.51 - 1.02 (m, 27 H), 1.09 - 1.77 (m, 15 H), 2.59 - 3.14 (m, 15 H), 4.07 - 4.24 (m, 4 H), 4.34 - 4.48 (m, 1 H), 4.99 - 5.76 (m, 8 H), 7.12 (br d, 4 H), 7.20 - 7.36 (m, 4 H), 8.96 - 9.08 (m, 2 H), 9.25 - 9.37 (m, 2 H);
- (3-319). δ : 0.61 - 1.02 (m, 27 H), 1.15 - 1.79 (m, 15 H), 2.61 - 3.12 (m, 16 H), 4.32 - 4.52 (m, 1 H), 4.96 - 5.77 (m, 7 H), 7.07 - 7.30 (m, 8 H), 7.31 - 7.45 (m, 2 H), 7.66 - 7.82 (m, 2 H), 8.21 - 8.34 (m, 2 H);
- (3-321). δ : 0.67 - 1.01 (m, 27 H), 1.16 - 1.74 (m, 15 H), 2.62 - 3.09 (m, 16 H), 4.34 - 4.48 (m, 1 H), 4.98 - 5.75 (m, 8 H), 7.00 - 7.36 (m, 10 H), 8.80 - 8.92 (m, 2 H), 9.12 - 9.26 (m, 2 H);
- (3-322). δ : 0.64 - 0.97 (m, 27 H), 1.16 - 1.75 (m, 15 H), 2.20 (s, 6 H), 2.63 - 3.09 (m, 16 H), 3.95 (s, 4 H), 4.36 - 5.71 (m, 8 H), 6.99 (s, 2 H), 7.05 - 7.11 (m, 4 H), 7.18 - 7.30 (m, 4 H), 7.94 - 7.80 (m, 2 H);
- (3-323). δ : 0.63 - 0.97 (m, 27 H), 1.14 - 1.75 (m, 15 H), 2.65 - 3.10 (m, 16 H), 3.94 (s, 4 H), 4.37 - 5.74 (m, 8 H), 7.12 - 7.18 (m, 4 H), 7.21 - 7.27 (m, 4 H), 7.27 - 7.33 (m, 2 H), 7.77 - 7.86 (m, 2 H), 8.05 - 8.11 (m, 2 H);
- (3-324). δ : 0.62 - 0.98 (m, 27 H), 1.13 - 1.76 (m, 15 H), 2.24 (s, 6 H), 2.64 - 3.10 (m, 16 H), 3.89 (s, 4 H), 4.37 - 5.74 (m, 8 H), 7.10 - 7.18 (m, 4 H), 7.21 - 7.28 (m, 4 H), 7.61 - 7.68 (m, 2 H), 7.88 (s, 2 H);

(3-326). δ : 0.62 - 0.97 (m, 27 H), 1.13 - 1.76 (m, 15 H), 2.64 - 3.13 (m, 16 H), 4.07 (s, 4 H), 4.37 - 5.75 (m, 8 H), 7.17 - 7.24 (m, 4 H), 7.24 - 7.32 (m, 6 H), 7.92 - 7.98 (m, 2 H);

(3-329). δ : 0.69 - 0.94 (m, 26H), 1.23 - 1.68 (m, 16H), 2.67 - 3.04 (m, 16H), 4.03 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.13-7.15 (m, 4H), 7.24-7.27 (m, 4H), 7.39-7.42 (m, 2H), 7.67-7.69 (m, 2H), 8.24-8.26 (m, 2H);

(3-332). δ : 0.64 - 0.99 (m, 27 H) 1.16 - 1.77 (m, 15 H) 2.66 - 2.95 (m, 13 H) 2.98 - 3.11 (m, 3 H) 3.79 - 3.91 (m, 10 H) 4.34 - 4.47 (m, 1 H) 5.00 - 5.75 (m, 7 H) 7.13 - 7.30 (m, 8 H) 7.49 (br s, 2 H) 8.02 - 8.12 (m, 2 H)

(3-333). δ : 0.60 - 1.05 (m, 27 H) 1.16 - 1.74 (m, 15 H) 2.64 - 2.96 (m, 13 H) 2.97 - 3.15 (m, 3 H) 4.11 - 4.22 (m, 4 H) 4.48 (s, 1 H) 4.95 - 5.76 (m, 7 H) 7.09 (m, 4 H) 7.20 - 7.34 (m, 4 H) 7.72 (m, 2 H) 8.62 - 8.79 (m, 4 H); and

(3-334). δ : 0.67 - 0.97 (m, 27 H), 1.13 - 1.76 (m, 15 H), 2.64 - 3.09 (m, 16 H), 3.61 - 3.71 (m, 6 H), 3.80 - 3.95 (m, 10 H), 4.34 - 4.49 (m, 1 H), 4.98 - 5.74 (m, 7 H), 6.69 - 6.80 (m, 2 H), 7.08 - 7.28 (m, 8 H), 7.71 - 7.81 (m, 2 H).

The following Formula (4A) compounds described in Table 4 were prepared in accordance with the schemes and examples described herein. Each of (R)_n, R^c and R^d are as described in the Table.





















































Table 4. Formula (4A) Compounds














Ex#	Ring B with (R) _n	R ^c	R ^d	Mass
-----	------------------------------	----------------	----------------	------













4-18		H	H	1079
4-19		H	H	1056
4-21		H	-OH	1095
4-30*		H	H	1030
4-32*		H	H	1029
4-34*		H	H	1059
4-40*		H	H	1054
4-50^		H	H	1045
4-63*		H	H	1043
4-75*		H	H	1080
4-105*		H	H	1075













4-107*		H	H	1081
4-109*		H	H	1058
4-118*		H	H	1084
4-126^		H	H	1084
4-139*		H	H	1054
4-141^		H	H	
4-1		H	-OH	1059
4-2		H	-OH	1056
4-3^		H	H	1057
4-4		H	H	1043
4-5^		H	H	1045
4-6		H	-OH	1086













4-7		H	H	1031
4-8		H	H	1070
4-9		H	-OH	1063
4-10		H	-OH	1061
4-11		H	Cl	1091
4-12		H	-OH	1135
4-13		H	-OH	1170
4-14		H	-OH	1095
4-15		H	-OH	1106
4-16		H	H	1090
4-17		H	H	1083
4-20		H	-OH	1106














4-22 [^]		H	H	1070
4-23		H	H	1071
4-24		-OH	-CF ₃	1129
4-25		H	H	1090
4-26		-OH	-CH ₃	1100
4-27		-OH	-CH ₂ CH ₃	1114
4-28		-CH(CH ₃) ₂	-OH	1128
4-29			-OH	1126
4-31 [*]		H	H	1046
4-33		H	H	1080
4-35 [^]		H	H	1046
4-36 [^]		H	H	1046
4-37		H	H	1070
4-38		H	H	1076












4-39		H	H	1076
4-41		H	-OH	1061
4-42 [^]		H	H	1057
4-43		H	H	1054
4-44		H	H	1074
4-45		H	-OH	1137
4-46		H	-OH	1075
4-47		H	-OH	1046
4-48		H	-OH	1045
4-49		H	-OH	1090
4-51		H	H	1059
4-52 [*]		H	H	1030
4-53 [^]		H	-OH	1060













4-54		H	H	1044
4-55		H	H	1113
4-56^A		H	-OH	1076
4-57		H	H	1060
4-58		H	-OH	1127
4-59		H	-OH	1076
4-60		H	-OH	1076
4-61		H	H	1100
4-62		H	H	1108
4-64		H	H	1100
4-65		H	-OH	1070
4-66^A		H	H	1060

4-67		H	-OH	1091
4-68		H	-OH	1124
4-69		H	-OH	1145
4-70		H	-OH	1097
4-71		H	-OH	1089
4-72		H	H	1121
4-73		H	H	1111
4-74		H	H	1152
4-76		H	-OH	1138
4-77^A		H	-OH	1096
4-78		H	H	1122
4-79		H	H	1073

4-80		H	-OH	1090
4-81		H	H	1131
4-82		H	-OH	1091
4-83^A		H	H	1044
4-84		H	-OH	1147
4-85		H	-OH	1097
4-86		H	H	1122
4-87		H	-OH	1119
4-88		H	-OH	1058
4-89		H	H	1106
4-90^A		H	H	1060
4-91^A		H	H	1058

4-92		H	H	1130
4-93		H	H	1114
4-94^		H	H	1060
4-95		H	-OH	1116
4-96		H	-OH	1074
4-97		H	-OH	1063
4-98		H	H	1080
4-99		H	-OH	1096
4-100		H	-OH	1080
4-101^		H	H	1054
4-102		H	H	1083
4-103		H	H	1129
4-104		H	-OH	1060

4-106		H	-OH	1122
4-108		H	-OH	1138
4-110		H	-OH	1089
4-111^		H	H	1081
4-112		H	-OH	1089
4-113		H	H	1122
4-114		H	H	1088
4-115^		H	H	1074
4-116^		H	H	1061
4-117		H	H	1105
4-119		H	H	1117

4-120		H	H	1088
4-121		H	H	1105
4-122		H	-OH	1077
4-123		H	-OH	1130
4-124		H	-OH	1121
4-125		H	H	1060
4-127		H	H	1138
4-128		H	H	1138
4-129		H	H	1086
4-130		H	H	1109
4-131		H	H	1084
4-132		H	H	1075

4-133		H	H	1044
4-134		H	H	1119
4-135^		H	H	1030
4-136		H	H	1125
4-137		H	-OH	1062
4-138		H	H	1103
4-140		H	H	1106
4-142		H	H	1069
4-143		H	H	1067

The following compound names and example #'s refer to the upper section compounds of Table 4.

(4-18). (3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-((1H-indazol-6-yl)methyl)benzyl)-18-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

- (4-19). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((2,4-dimethyl-1H-pyrrol-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (4-21). (3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-(benzofuran-3-yl(hydroxy)methyl)benzyl)-18-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (4-30). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-(oxazol-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (4-32). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-(furan-2-ylmethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (4-34). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-((3-methylthiophen-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (4-40). 2-(4-(((2R,5S,8R,11S,14R,17S,20R,23S)-14-benzyl-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octa-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2-yl)methyl)benzyl)furan-3-carbonitrile;
- (4-50). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-(thiophen-3-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (4-63). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-((1-methyl-1H-pyrazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (4-75). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3-chlorothiophen-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
- (4-105). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-18-(4-((3-methoxythiophen-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(4-107). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((4-chlorothiazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(4-109). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3,5-dimethylisoxazol-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(4-118). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-18-(4-((6-methoxy-2-methylpyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(4-126). (3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-18-(4-((6-methoxy-4-methylpyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(4-139). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-methylpyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and

(4-141). (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylisothiazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; stereoisomers thereof, and veterinary acceptable salts thereof.

NMR data (¹H NMR: 400 MHz, DMSO-d₆, δ ppm) for the named Table 4 compounds are shown below:

(4-18). δ: 0.56 - 1.85 (m, 42 H) 2.62 - 3.19 (m, 16 H) 4.05 (s, 2 H) 3.94-4.44 (m, 1 H) 4.97 - 5.83 (m, 9 H) 6.97 (m, 1 H) 7.15 - 7.39 (m, 10 H) 7.65 (m, 1 H) 7.99 (s, 1 H);

(4-19). δ: 0.62 - 1.02 (m, 26 H), 1.13 - 1.78 (m, 16 H), 1.81 - 1.91 (m, 3 H), 2.00 - 2.13 (m, 3 H), 2.63 - 3.18 (m, 16 H), 3.70 (s, 2 H), 4.42 (m, 1 H), 4.95 - 5.82 (m, 7 H), 7.02 - 7.41 (m, 10 H), 9.93 - 10.06 (m, 1 H);

(4-21). δ: 0.65 - 0.99 (m, 27 H), 1.14 - 1.80 (m, 15 H), 2.47 - 2.62 - 2.99 (m, 13 H), 2.99 - 3.15 (m, 3 H), 4.38 - 4.46 (m, 2 H), 4.98 - 5.77 (m, 8 H), 5.88 - 5.93 (m, 1 H), 7.07 - 7.17 (m, 1 H), 7.17 - 7.35 (m, 9 H), 7.37 - 7.44 (m, 2 H), 7.51 (d, J = 8.31 Hz, 1 H), 7.78 - 7.83 (m, 1 H);

- (4-30). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.66 - 3.02 (m, 16H), 4.09 (s, 2H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.12-7.31 (m, 10H), 7.98-7.99 (m, 1H);
- (4-32). δ : 0.70 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.67 - 3.04 (m, 16H), 3.93 (s, 2H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 6.07 (brs, 1H), 6.34-6.35 (m, 1H), 7.13 - 7.16 (m, 2H), 7.22-7.31 (m, 7H), 7.50 (brs, 1H);
- (4-34). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.12 (s, 3H), 2.66 - 3.06 (m, 16H), 4.01 (s, 2H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 6.82-6.83 (m, 1H), 7.11 - 7.31 (m, 10H);
- (4-40). δ : 0.70 - 0.94 (m, 26H), 1.23 - 1.71 (m, 16H), 2.67 - 3.05 (m, 16H), 4.15 (s, 2H), 4.40 - 4.42 (m, 1H), 5.04 - 5.70 (m, 7H), 6.87 (s, 1H), 7.16-7.30 (m, 9H), 7.79 (s, 1H);
- (4-50). δ : 0.68 - 0.95 (m, 26H), 1.23 - 1.69 (m, 16H), 2.67 - 3.05 (m, 16H), 3.90 (s, 2H), 4.40 - 4.42 (m, 1H), 5.03 - 5.70 (m, 7H), 6.91-7.43 (m, 12H);
- (4-63). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.66 - 3.06 (m, 16H), 3.65 (s, 3H), 3.98 (s, 2H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 5.92 (brs, 1H), 7.11 - 7.31 (m, 10H);
- (4-75). δ : 0.68 - 0.95 (m, 26H), 1.23 - 1.70 (m, 16H), 2.67 - 3.05 (m, 16H), 4.07 (s, 2H), 4.40 - 4.42 (m, 1H), 5.02 - 5.73 (m, 7H), 7.01-7.50 (m, 11H);
- (4-105). δ : 0.68-0.94 (m, 27H), 1.20-1.80 (m, 15H), 2.66-3.10 (m, 16H), 3.77 (s, 3H), 3.91 (s, 2H), 4.38-4.44 (m, 1H), 5.02-5.75 (m, 7H), 6.97-7.31 (m, 11H);
- (4-107). δ : 0.68 - 0.95 (m, 26H), 1.24 - 1.70 (m, 16H), 2.67 - 3.05 (m, 16H), 4.12 (s, 2H), 4.40 - 4.42 (m, 1H), 5.04 - 5.70 (m, 7H), 7.17-7.30 (m, 9H), 8.98 (s, 1H);
- (4-109). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 1.99 (s, 3H), 2.30 (s, 3H), 2.67 - 3.05 (m, 16H), 3.65 (s, 2H), 4.40 - 4.42 (m, 1H), 5.09 - 5.67 (m, 7H), 7.06-7.08 (m, 2H), 7.22-7.31 (m, 7H);
- (4-118). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.26 (s, 3H), 2.66 - 3.02 (m, 16H), 3.76 (s, 3H), 3.84 (s, 2H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 6.54-6.56 (m, 1H), 7.02-7.40 (m, 10H);
- (4-126). δ : 0.68 - 0.94 (m, 26H), 1.23 - 1.70 (m, 16H), 2.08 (s, 3H), 2.66 - 3.02 (m, 16H), 3.79 (s, 3H), 3.86 (s, 2H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 6.61 (s, 1H), 7.04-7.31 (m, 9H), 7.89 (s, 1H);
- (4-139). δ : 0.68 - 0.95 (m, 26H), 1.23 - 1.72 (m, 16H), 2.37 (s, 3H), 2.66 - 3.06 (m, 18H), 3.94 (s, 4H), 4.40 - 4.42 (m, 1H), 5.02 - 5.71 (m, 7H), 7.06 - 7.31 (m, 10H), 7.43-7.46 (m, 1H), 8.28-8.29 (m, 1H); and

(4-141). δ : 0.69 -0.94 (m, 26H), 1.23 -1.68 (m, 16H), 2.17 (s, 3H), 2.67 - 3.05 (m, 16H), 4.15 (s, 2H), 4.40 - 4.42 (m, 1H), 5.02 - 5.70 (m, 7H), 7.18-7.31 (m, 9H), 8.30 (s, 1H).

Biological

Heartworm infection, caused by the endoparasite *Dirofilaria immitis* (*D. immitis*), can be a severe and life-threatening disease in animals such as dogs and cats. Heartworm has a complicated life cycle involving several life stages before they mature into adults that will eventually infect the pulmonary artery of the host animal. Heartworm transmission also requires the mosquito to act as an intermediate host to complete this life cycle. For example, the beginning of the heartworm life cycle and transmission process involves a mosquito biting a previously infected dog and ingesting blood containing heartworm microfilariae (larva stage 1). Within the mosquito, the microfilariae will molt into infective larva stage 3 (L3) worms over a two week period. Once the mosquito bites another dog, infective L3 worms will move through the bite wound to enter the host and migrate into the tissues where they will begin molting into larva stage 4 (L4) worms, usually within 1 to 3 days post infection. Subsequently, L4 worms will continue their migration through tissues and molt into sexually immature or “adolescent” adults (larva stage 5, immature adult), approximately 50-70 days post infection. Sexually mature worms will eventually migrate to the heart and lungs of the dog, as early as 70 days post infection. Approximately 6-7 months post infection *D. immitis* adults reach maturity and sexually reproduce in the pulmonary artery leading to microfilaria (MF) production and circulation in the blood of the dog, thus completing the heartworm life cycle.

The most commonly used heartworm preventatives are the macrocyclic lactones (MLs) such as ivermectin, moxidectin and selamectin. These agents are administered on a monthly basis whereby they kill *D. immitis* L3 and L4 worms acquired by the host within the previous 30 days. Their primary action is to disrupt the heartworm life cycle by killing L3 and L4 worms thus preventing adult formation and subsequent disease. While very effective at preventing heartworm disease, owners are advised to test dogs for existing heartworm infections (i.e. heartworm positive dogs) prior to starting treatment with MLs due to their potential to kill circulating microfilariae. A rapid decrease in the numbers of microfilariae in the blood can lead to hypersensitivity-type reactions and circulatory shock (e.g.

anaphylaxis), presumably due to dead or dying microfilariae. These potential adverse effects can be life-threatening to the dog and as such are presented as caution statements on many ML product labels. Therefore, the discovery of a novel heartworm preventative that would selectively target L3 and L4 stage worms versus microfilariae would offer a potential safety advantage. By not killing circulating microfilariae in heartworm positive dogs, a targeted treatment would prevent the adverse effects known to occur with other heartworm preventatives that lack *D. immitis* stage selectivity.

To identify novel heartworm preventatives, compounds were screened for nematocidal activity using *in vitro* motility assays. The compounds described herein have demonstrated nematocidal activity against either *Dirofilaria immitis* (Larva stage 4 (*DiL4*)) and/or *Dirofilaria immitis* microfilaria (*DiMF*)) as determined by reductions in nematode motility either by paralysis or death. Active and selective (*DiL4* vs. *DiMF* potency) example compounds were subsequently evaluated in heartworm positive dog studies to correlate the *in vitro* selectivity profile with *in vivo* effects on circulating microfilariae.

The *in vitro* (*DiL4* and *DiMF*) and *in vivo* (heartworm positive dog studies) biological activity against *Dirofilaria immitis* of the compounds of the invention can be measured using the test methods described below.

Dirofilaria immitis, Microfilaria (*DiMf*) *In vitro* Assay

Compounds were dissolved and serially diluted in DMSO. Aliquots were spotted to the empty wells of assay plates. Media and microfilariae of *Dirofilaria immitis* were added to each well to dilute the test compounds to the desired concentrations. Assay plates were incubated for approximately 72 hours, and the larvae in each well were observed microscopically for drug effect. Microfilariae in each well were assessed subjectively for survival or paralysis, and results were reported as Minimum Effective Dose (MED).

Dirofilaria immitis, L4 stage (*DiL4*) *In vitro* Assay

Compounds were initially dissolved in DMSO. The stock concentration was subsequently diluted in basal media and serially diluted to give a concentration response curve starting at 100 μM (≥ 11 total concentrations). Following the serial dilution, compound solution was transferred to an assay plate (384-well) where

D.immitis L4 larvae (10/well) that have been molted from L3 larvae *in vitro* were subsequently added. Assay plates were observed at 72 hours for drug effect. Each compound was evaluated for decrease in L4 motility by subjective visual assessment and endpoint data were recorded as minimally effective concentration (MEC) in μM following the incubation period. Examples with DiL4 MED data $\leq 1\text{nM}$ are presented in Tables 1-4 (*i.e.*, upper sections). Examples with data $>1\text{nM}$ are also presented in Tables 1-4 (*i.e.*, lower sections).

Heartworm Positive Dog Studies

Dogs with pre-existing heartworm infections, via surgical transplantation were used for these studies. To confirm that the dogs had circulating microfilariae, blood samples were taken from each dog and examined for microfilariae by using the modified Knott's method. All dog cohorts included in the studies exhibited average microfilariae counts of at least 15,000 MF/mL of the blood (pre-dose). On approximately Day -7, dogs were randomly allocated to treatments (based on Day -7 MF counts) and pens according to a randomized complete block design with one-way structure. Dogs were fasted overnight prior to dosing and fed immediately following dosing of the test articles. Compounds were administered by point dosing in oral liquid-filled capsules on Day 0 (3 dogs/treatment group). Blood samples were collected to measure MF counts on Days 0 (pre-dose and 2 hours post-dose), 1, 7, 21 and 28. Clinical observations were conducted by a suitably experienced Veterinarian on days -7, 0 (immediately prior to treatment, 1-2 hours post-treatment), 1 and 2 whereby any abnormal clinical signs were documented using standard veterinary medical terminology. Additionally, general health observations were conducted throughout the study including (but not limited to) general physical appearance and behavior, abnormalities of food and water consumption, vomiting/regurgitation, appearance of urine and feces and any sign of MF anaphylaxis.

The reference depsipeptide, emodepside, has equivalent *in vitro* potency against DiL4 (30 nM) and DiMF (30 nM) when assessed in the nematode motility assays and therefore represents a non-selective compound. As such, emodepside was evaluated for effects against circulating microfilariae in a heartworm positive dog study. At a dose of 1 mg/kg (po), emodepside demonstrated a rapid killing of circulating microfilariae, reducing the average MF counts from 61,000 MF/mL of

blood pre-dose to 8,300 MF/mL of blood at 2 hours post-dose (87% decrease; Table 5). Moreover, all treated dogs exhibited adverse effects consistent with dead or dying circulating microfilariae (e.g. lethargy, tremors, ataxia, hyper-salivation) within 2 hours of emodepside administration. All dogs were treated with dexamethasone, epinephrine and valium such that they were able to recover from their symptoms.

In contrast to emodepside, the compounds described herein demonstrate larval stage selectivity (i.e. *DiL4* vs. *DiMF* potency) as shown in Table 5. Emodepside (1mg/kg) and Example (3-9; 3mg/kg) were progressed into a heartworm positive dog study to correlate the in vitro selectivity profile with in vivo effects on circulating microfilariae. Both compounds were administered orally. Selectivity is shown as a ratio of *Dimf*/*DiL4*. The *Dimf* and *DiL4* MED for emodepside 10nM. There was no selectivity (1x) between these larval stages with emodepside. In contrast, For Example (3-9), the *Dimf* and *DiL4* MED was 60nM and 3nM, respectively; thereby providing a 20x larval selectivity ratio. Additionally, there were no adverse effects observed following the administration of Example (3-9), consistent with a lack of microfilariae kill in the blood. Average mean MF counts were 18,000 MF/mL of blood pre-dose and remained elevated throughout the study where it averaged 24,000 MF/mL of blood on day 21 of the dog study. Moreover, there were no adverse clinical observations noted throughout the study, consistent with a lack of potency against circulating MF in the dogs. These data highlight the importance of generating larval stage selective compounds as novel heartworm preventatives. Specifically targeting the L4 stage versus the MF stage *D. immitis* worms provides an improved safety profile when administering preventatives to dogs with an already active adult heartworm infection. Indeed, the compounds described herein exhibit improved selectivity ratios whereby they also offer improved safety profiles over older preventatives and reference depsipeptides such as emodepside.

Table 5. Heartworm positive dog studies evaluating effects on circulating microfilariae (MF) following compound administration.

Compound		Day 0 Pre- dose	Day 0, Post- dose, 2 hours	Day 1	Day 7	Day 14	Day 21
Emodepside	Mean MF/mL	61,000	8,300	4,500	8,800	N.D.	N.D.
	% Reduction		87%	93%	86%	N.D.	N.D.
Example 3-9	Mean MF/mL	18,000	20,000	23,000	25,000	22,000	24,000

	% Reduction		-11%	-28%	-50%	-22%	-33%
--	-------------	--	------	------	------	------	------

N.D. = Not Determined; Microfilariae raw counts are shown as mean MF/mL (3 dogs/treatment) of blood. Percent reductions were calculated from counts on day of assessment compared back with pre-dose (day 0) levels. Negative percent reductions are a reflection of natural variability across time due to active adult heartworm infection and continued microfilariae production.

Haemonchus contortus, Larvae Stage 3 (HcL3) *In vitro* Assay

The *Haemonchus contortus* L3 strain was obtained from the University of Georgia and is a relatively recent multiple-resistant field isolate (International Journal for Parasitology 37 (2007) 795-804). Compounds were dissolved and serially diluted in DMSO. Aliquots were spotted to the empty wells of assay plates. Media and third stage larvae of *Haemonchus contortus* were added to each well to dilute the test compounds to the desired concentrations. Assay plates were incubated for approximately 96 hours, and the larvae in each well were observed microscopically for drug effect. Larvae in each well were assessed subjectively for survival or paralysis, and results were reported as Minimum Effective Dose (MED). Examples with HcL3 MED data $\leq 1 \mu\text{M}$ (Ex#*) and data $1 > x \leq 3.3 \mu\text{M}$ (Ex#^) are presented in Tables 1-4.

Summary worm data is presented in Table 6 for those compounds with a DiL4 MED (nM) value of $<1\text{nM}$. In addition, DiMf MED (nM), HcL3 MED (μM) data, and DiMf//DiL4 specificity is also shown.

Table 5. DiL4, Dimf, Dimf/DiL4, and HcL3 data

Ex.#	DiL4 (nM)	DiMf (nM)	DiMf/DiL4	HcL3 (μM)	Ex.#	DiL4 (nM)	DiMf (nM)	DiMf/DiL4	HcL3 (μM)
(1-5)	1	500	500	>100	(3-274)	0.3	10	33.3	1
(1-11)	1	300	300	>10	(3-275)	1	10	10	1.8
(1-18)	<0.1	3	>30	>100	(3-282)	1	5	5	30
(2-1)	1	20	20	0.65	(3-283)	0.5	3	6	1
(2-4)	0.7	50	71.4	3.3	(3-284)	0.5	100	200	>10
(2-36)	0.2	30	150	1	(3-285)	0.5	3	6	5.7
(2-37)	1	20	20	3.3	(3-286)	0.5	30	60	>10
(2-38)	0.6	30	50	0.5	(3-287)	0.5	50	100	>10
(2-42)	0.3	300	1000	>10	(3-288)	0.1	3	30	0.2
(3-4)	0.1	20	200	0.5	(3-289)	0.5	3	6	3.3
(3-8)	0.3	10	33.3	1	(3-290)	0.1	0.3	3	0.2
(3-12)	1	0.7	0.7	3.3	(3-291)	0.1	0.3	3	0.3
(3-17)	0.3	0.3	1	0.5	(3-268)	0.5	2	4	1
(3-18)	0.1	0.2	2	1	(3-273)	0.3	2	6.7	1


(3-21)	<0.03	0.1	>3.3	0.3	(3-274)	0.3	10	33.3	1
(3-24)	0.03	0.1	3.3	1	(3-275)	1	10	10	1.8
(3-31)	0.2	5	25	2.4	(3-282)	1	5	5	30
(3-44)	0.1	100	1000	1.8	(3-283)	0.5	3	6	1
(3-49)	1	5	5	3.3	(3-284)	0.5	100	200	>10
(3-50)	0.1	1	10	0.3	(3-285)	0.5	3	6	5.7
(3-56)	0.5	100	200	0.5	(3-286)	0.5	30	60	>10
(3-57)	0.3	0.3	1	1	(3-287)	0.5	50	100	>10
(3-58)	0.2	5	25	1	(3-288)	0.1	3	30	0.2
(3-60)	1	50	50	>10	(3-289)	0.5	3	6	3.3
(3-69)	0.7	30	42.9	0.5	(3-290)	0.1	0.3	3	0.2
(3-87)	0.3	100	333	0.3	(3-291)	0.1	0.3	3	0.3
(3-110)	0.1	30	300	>100	(3-292)	0.3	1	3.3	1.8
(3-112)	0.03	1	33.3	1	(3-293)	0.3	50	167	5.7
(3-118)	1	100	100	>100	(3-294)	0.1	0.1	1	1
(3-120)	0.1	0.4	4	1	(3-295)	0.3	5	16.7	1.8
(3-121)	0.05	3	60	1.8	(3-296)	0.5	30	60	1.8
(3-124)	1	100	100	0.3	(3-297)	1	30	30	>10
(3-125)	<0.2	30	>150	>100	(3-307)	0.1	0.3	3	1
(3-132)	0.4	10	25	>10	(3-308)	1	10	10	3.3
(3-133)	0.3	20	66.7	57	(3-309)	0.55	3	5.5	3.3
(3-148)	0.3	1	3.3	3	(3-310)	1	0.3	0.3	1
(3-155)	0.02	0.1	5	0.5	(3-313)	0.55	3	5.5	0.55
(3-156)	0.1	0.5	5	5.6	(3-318)	0.03	0.3	10	0.55
(3-157)	1	100	100	>100	(3-319)	0.055	1.73	31.5	0.55
(3-165)	1	200	200	>100	(3-321)	0.02	0.05	2.5	0.3
(3-173)	<0.1	3	>30	3.3	(3-322)	0.3	10	33.3	10
(3-176)	0.3	4	13.3	1.2	(3-323)	0.1	1	10	0.3
(3-178)	1	1	1	>100	(3-324)	0.1	10	100	1
(3-193)	0.4	2	5	1	(3-326)	0.3	30	100	10
(3-194)	1	1	1	>100	(3-329)	0.1	1	10	--
(3-207)	1	50	50	>100	(3-332)	0.5	300	548	--
(3-209)	0.5	10	20	>100	(3-333)	0.2	3	17	--
(3-215)	<0.2	0.7	>3.5	1	(3-334)	0.3	3	10	0.1
(3-217)	<0.7	3	>4.3	3.3	(4-18)	1	50	50	>10
(3-218)	0.8	10	12.5	>23	(4-19)	1	100	100	>10
(3-220)	0.7	5	7.1	>100	(4-21)	1	50	50	>10
(3-223)	0.7	5	7.1	5.7	(4-30)	1	1	1	1
(3-225)	0.7	300	429	>100	(4-32)	1	3	3	1
(3-228)	0.5	10	20	>100	(4-34)	0.3	10	33.3	0.5
(3-238)	0.1	2	20	>5	(4-40)	0.2	0.5	2.5	0.5
(3-240)	1	20	20	5.7	(4-50)	1	10	10	1.8
(3-243)	1	1	1	7.5	(4-63)	1	1	1	1
(3-252)	0.7	20	28.6	>42	(4-75)	0.5	10	20	1
(3-253)	1	0.3	0.3	1	(4-105)	0.2	5	25	0.3
(3-254)	1	3	3	10	(4-107)	0.2	3	15	0.3
(3-259)	0.5	10	20	3.3	(4-109)	0.4	1	2.5	1
(3-261)	<0.1	30	>300	>100	(4-118)	0.1	1	10	0.3
(3-268)	0.5	2	4	1	(4-126)	0.2	3	15	1.8
(3-273)	0.3	2	6.7	1	(4-139)	0.4	1	2.5	1

					(4-141)	1	3	3	1.8
--	--	--	--	--	---------	---	---	---	-----

Claims

We Claim:

1. A compound of Formula (1) compound



wherein

L_1 and L_2 are the same and are C_1 - C_4 alkyl, C_3 - C_6 cycloalkyl, C_1 - C_4 alkylaryl, C_1 - C_4 alkylheteroaryl or C_1 - C_4 alkylheterocycle; or L_1 is absent; and wherein the cycloalkyl is a 3- to 6-membered carbocyclic ring; the aryl is a 6-membered monocyclic aromatic ring or a 9- to 10-membered fused aromatic ring; the heteroaryl is a 5- or 6-member monocyclic aromatic ring or a 9- to 10-member fused aromatic ring wherein each mono- or fused-heteroaryl ring contains at least one heteroatom selected from N, O, and S; and the heterocycle is a 4- to 6-member monocyclic saturated or partially saturated ring or a 9- to 10-member fused saturated or partially saturated ring, each mono- or fused-heterocyclic ring contains at least one heteroatom selected from N, O and S; and wherein the heteroaryl and heterocycle rings are linked to the L_1 and L_2 C_1 - C_4 alkyl or L_2 C_1 - C_4 alkyl by a ring carbon atom;

and wherein the L_1 and L_2 alkyl group of the alkylcycloalkyl, alkylaryl, alkylheteroaryl and alkylheterocycle can be substituted with an R^c and R^d substituent each individually and separately selected from the group consisting of hydroxyl, halo, C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_1 - C_6 alkoxy, C_1 - C_4 haloalkyl, $-NR^aR^b$, pyrazolyl, imidazolyl, pyrrolidinyl, and morpholinyl; and when L_1 is not absent, then each R^c is the same and each R^d is the same;

and wherein the L_1 and L_2 cycloalkyl, aryl, heteroaryl, and heterocycle groups can be substituted with $(R)_n$; and when L_1 is not absent, then each $(R)_n$ is the same; and each R substituent is independently selected from the group consisting of C_1 - C_6 alkyl optionally substituted with at least one substituent selected from hydroxyl, C_3 - C_6 cycloalkyl and $-OR^a$; halo, oxo, cyano, hydroxyl, $-OR^5$, $-NR^aR^b$, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, $-S(O)_pR^a$, C_1 - C_6 alkoxy, C_3 - C_6 cycloalkyl, $-C(O)NR^aR^b$, $-C(O)R^a$, $-C(O)OR^a$, $-NR^aC(O)R^b$, pyrazolyl, imidazolyl, pyridinyl, pyrimidinyl, piperidinyl, morpholinyl, pyrrolidinyl, dihydropyrimidinyl, and phenyl; and wherein the phenyl is optionally substituted with at least one substituent selected from C_1 - C_3 alkyl, $-CF_3$, halo, and hydroxyl; and wherein n is the integer 0, 1, 2, or 3;

R^1 , R^2 , R^3 , and R^4 are each separately and independently H and C_1 - C_6 alkyl;

R^a and R^b are each separately H and C_1 - C_6 alkyl; and


R^5 is C_3 - C_6 cycloalkyl and C_1 - C_3 alkyl- C_3 - C_6 cycloalkyl;

p is the integer 0, 1, or 2;

stereoisomers thereof, and veterinary acceptable salts thereof.

2. A composition comprising a compound of Formula 1 of Claim 1 and wherein the composition further comprises a veterinary acceptable excipient.

3. A compound of Formula (1) of Claim 1 that is a compound of Formula (1A1)




wherein each $(R)_n$ is the same and is independently selected from the group consisting of C_1 - C_6 alkyl optionally substituted with at least one substituent selected from hydroxyl and C_3 - C_6 cycloalkyl; halo, cyano, hydroxy, $-NR^aR^b$, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, $-S(O)_pR^a$, C_1 - C_6 alkoxy, C_3 - C_6 cycloalkyl, morpholinyl, and phenyl

optionally substituted with methyl, $-\text{CF}_3$, halo, and hydroxyl; and wherein n is the integer 0, 1, 2, or 3;

R^c and R^d are each independently selected from the group consisting of H, hydroxyl, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-\text{NR}^a\text{R}^b$, $-\text{CF}_3$, cyclopropyl, pyrazolyl, pyrrolidinyl, and morpholinyl; and wherein each R^c is the same and each R^d is the same; stereoisomers thereof, and veterinary acceptable salts thereof.

4. A compound of Formula (1) of Claim 1 that is a compound of Formula (3A)



wherein Ring A and Ring B are the same and are selected from the group consisting of a 3- to 6-membered cyloalkyl ring selected from cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; or a 4- to 6-membered heterocycle ring selected from oxetanyl, azetidiny, piperidiny, pyrrolidinyl, morpholinyl, tetrahydropyranyl, tetrahydropyrimidinyl, tetrahydropyridinyl, and tetrahydropyranyl; or a 5- or 6-membered heteroaryl ring selected from thiophenyl, furanyl, thiazolyl, oxazolyl, pyrrolyl, pyrazolyl, imidazolyl, pyridinyl, thiadiazolyl, isothiazolyl, triazinyl, pyrimidinyl, pyrazinyl, pyridazinyl, isoxazolyl, tetrazolyl, oxadiazolyl, 1,2-dihydropyridinyl, or triazolyl; or a 9- to 10-membered fused heteroaryl ring selected from the group consisting of quinolinyl, isoquinolinyl, benzofuranyl, 2,3-dihydrothieno[3,4-b][1,4]dioxine, 3,4-dihydro-2H-pyrano[2,3-b]pyridinyl, oxazolo[5,4-b]pyridinyl, pyrazolo[1,5-a]pyridinyl, pyrazolo[1,5-a]pyrimidine, 4,5,6,7-tetrahydropyrazolo[1,5-a]pyridinyl, 4,5-tetrahydropyrazolo[1,5-a]pyrimidine, 1H-pyrrolo[3,2-b]pyridinyl, indazolyl, furo[2,3-b]pyridinyl, and benzo[d][1,3]dioxole; and wherein each $(\text{R})_n$ is the same and is independently selected from the group consisting of C_1 - C_6 alkyl optionally substituted with at least one substituent selected from hydroxyl and C_3 - C_6 cycloalkyl; halo, cyano, hydroxy, $-\text{NR}^a\text{R}^b$, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, $-\text{S}(\text{O})_p\text{R}^a$, C_1 - C_6 alkoxy,

C₃-C₆cycloalkyl, morpholinyl, and phenyl optionally substituted with methyl, -CF₃, halo, and hydroxyl; and wherein n is the integer 0, 1, 2, or 3;

R^c and R^d are each independently selected from the group consisting of H, hydroxyl, halo, C₁-C₆alkyl, C₁-C₆alkoxy, -NR^aR^b, -CF₃, cyclopropyl, pyrazolyl, pyrrolidinyl, and morpholinyl; and wherein each R^c is the same and each R^d is the same; stereoisomers thereof, and veterinary acceptable salts thereof.

5. The compound of Claim 4 wherein Ring A and Ring B are the same and are selected from the group consisting of thiophenyl, furanyl, thiazolyl, oxazolyl, pyrazolyl, pyridinyl, thiadiazolyl, isothiazolyl, triazinyl, pyrimidinyl, pyrazinyl, pyridazinyl, isoxazolyl, benzofuranyl, indazolyl, and tetrahydropyranyl; stereoisomers thereof, and veterinary acceptable salts thereof.

6. The compound of Claim 5 wherein Ring A and Ring B are the same and are selected from the group consisting of pyrazolyl, isothiazolyl, thiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, and pyridazinyl; and wherein each (R)_n is the same and is independently selected from the group consisting of methyl, ethyl, isopropyl, propyl, isobutyl, fluoro, chloro, bromo, cyano, hydroxyl, -NH₂, -NHCH₃, -N(CH₃)₂, -N(CH₂CH₃)₂, -NHC(O)CH₃, methoxy, ethoxy, isopropoxy, oxo, -CF₃, -CHF₂, -CH₂CH₂F, -CH₂CF₃, cyclopropyl, cyclobutyl, -CH₂cyclopropyl, -OCHF₂, -OCF₃, -OCH₂CF₃, -O-cycloalkyl, -O-cyclobutyl, -O-CH₂-cyclopropyl, -C-O-CH₃, -C-C-O-CH₃, -COH, -C(CH₃)₂OH, -C(O)CH₃, -SCH₃, -S(O)₂CH₃, -C(O)N(CH₃)₂, -C(O)OC(CH₃)₃, -C(O)C(CH₃)₂, pyrazolyl, imidazolyl, pyrimidinyl, pyridinyl, 1,2-dihydropyrimidinyl, 1,6-dihydropyrimidinyl, piperidinyl, pyrrolidinyl, morpholinyl, and phenyl and wherein the phenyl is optionally substituted with methyl; stereoisomers thereof, and veterinary acceptable salts thereof.

7. The compound of Claim 6 wherein Ring A and B are the same and are selected from the group consisting of pyrazolyl, pyrazinyl, isothiazolyl, thiazolyl, pyridinyl, pyridazinyl, and pyrimidinyl.

8. The compound of Claim 6 or 7 wherein each (R)_n is the same and is independently selected from the group consisting of methyl, ethyl, isopropyl, fluoro, chloro, bromo, cyano, hydroxyl, -N(CH₃)₂, methoxy, ethoxy, isopropoxy, -CHF₂,

-CH₂CH₂F, -CF₃, cyclopropyl, -CH₂cyclopropyl, -OCHF₂, OCF₃, -C(O)CH₃, -SCH₃, and morpholinyl; and wherein R^c and R^d are each independently selected from the group consisting of H, hydroxyl, methyl, methoxy, ethoxy, -N(CH₂CH₃)₂, pyrazolyl, and morpholinyl; and wherein each R^c is the same and each R^d is the same; stereoisomers thereof, and veterinary acceptable salts thereof.

9. A compound of Claim 4 selected from the group consisting of:

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiophen-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(furan-2-ylmethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiazol-5-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiazol-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(oxazol-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3,5-dimethylisoxazol-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((1-methyl-1H-pyrazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((2-methoxypyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((3-methylthiophen-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1,3-dimethyl-1H-pyrazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
2,2'-((((2R,5S,8R,11S,14R,17S,20R,23S)-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octa-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,14-diyl)bis(methylene))bis(4,1-phenylene))bis(methylene))bis(furan-3-carbonitrile);
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(thiophen-3-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(oxazol-4-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
4,4'-((((2R,5S,8R,11S,14R,17S,20R,23S)-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octa-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,14-diyl)bis(methylene))bis(4,1-phenylene))bis(methylene))bis(furan-2-carbonitrile);
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((6-methoxypyridin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-chlorothiophen-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((3-methoxythiophen-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-chlorothiazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-fluoro-6-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methyl-1,2,3-thiadiazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((6-methoxy-2-methylpyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylthiophen-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-bromothiophen-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-((R)-(3-bromothiophen-2-yl)(hydroxy)methyl)benzyl)-18-(4-((S)-(3-bromothiophen-2-yl)(hydroxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((6-methoxy-4-methylpyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-(oxazol-5-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylthiazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(isothiazol-5-ylmethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((5-methylfuran-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-ethoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(furan-3-ylmethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-(dimethylamino)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-morpholinopyrimidin-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((3-methoxypyrazin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-fluoropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(1-(6-methoxypyridin-3-yl)ethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((2-methoxy-4-methylthiazol-5-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((3-methoxypyridin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((4-methoxypyrimidin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((6-methoxypyrazin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(1-fluoro-1-(2-methoxypyridin-3-yl)ethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylpyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(1-fluoro-1-(6-methoxypyridin-2-yl)ethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((4-isopropyl-1,2,3-thiadiazol-5-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-(trifluoromethyl)pyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,6-dimethylpyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((5-methyl-1H-pyrazol-4-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-(1-fluoro-1-(5-methoxypyrazin-2-yl)ethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1-ethyl-1H-pyrazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(isothiazol-4-ylmethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-hydroxypyridin-3-yl)(methoxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(1-(2-methoxypyridin-3-yl)ethyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1-cyclopropyl-1H-pyrazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9R,12R,15R,18R,21S,24S)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis[[4-[(2-methyl-3-pyridyl)methyl]phenyl]methyl]-1,7,13,19-tetraoxa-4,10,16,22-tetrazacyclotetracosane-2,5,8,11,14,17,20,23-octaone;

(3R,6R,9R,12R,15S,18S,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((5-methoxypyrazin-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1-(cyclopropylmethyl)-1H-pyrazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((4-methoxypyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-((4-isopropylpyrimidin-5-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3-cyclopropylisoxazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-((R)-fluoro(thiazol-2-yl)methyl)benzyl)-18-(4-((S)-fluoro(thiazol-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((1-(2-fluoroethyl)-1H-pyrazol-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-(difluoromethyl)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylisothiazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-cyclopropylpyrimidin-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-(difluoromethyl)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((R)-1-(thiazol-2-yl)ethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-morpholinopyridin-4-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-fluoro-2-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylpyrimidin-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-(difluoromethyl)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-cyclopropylpyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and
(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-(difluoromethoxy)pyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone,
3,3'-((((2R,5S,8R,11S,14R,17S,20R,23S)-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octa-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosane-2,14-diyl)bis(methylene))bis(4,1-phenylene))bis(methylene))diisonicotinonitrile;
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-(methylthio)pyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-((3-fluoropyridin-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-((3-methylpyridin-4-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-ethylpyrimidin-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-fluoropyridin-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-(trifluoromethyl)pyrimidin-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-chloropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((4-(difluoromethyl)pyrimidin-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((6-fluoro-4-methylpyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-fluoropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;


(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,5-difluoropyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,3-difluoropyridin-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2-bromopyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((5-chloro-2-methoxypyridin-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
 (3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-(trifluoromethyl)pyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and
 (3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((2,3-dimethoxypyridin-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone,
 stereoisomers thereof, and veterinary acceptable salts thereof.

10. A compound of Formula (1) of Claim 1 that is a compound of Formula (2A1) or Formula (4A)



wherein Ring B is selected from the group consisting of pyrrolyl, thiophenyl, furanyl, thiazolyl, oxazolyl, pyrazolyl, pyridinyl, isothiazolyl, isoxazolyl, benzofuranyl and indazolyl; and

wherein each (R)_n is the same and is independently selected from the group consisting of methyl, ethyl, isopropyl, propyl, isobutyl, fluoro, chloro, bromo, cyano, hydroxyl, -NH₂, -NHCH₃, -N(CH₃)₂, -N(CH₂CH₃)₂, -NHC(O)CH₃, methoxy, ethoxy, isopropoxy, oxo, -CF₃, -CHF₂, -CH₂CH₂F, -CH₂CF₃, cyclopropyl, cyclobutyl, -CH₂cyclopropyl, -OCHF₂, -OCF₃, -OCH₂CF₃, -O-cycloalkyl, -O-cyclobutyl, -O-C-cyclopropyl, -C-O-CH₃; -C-C-O-CH₃, -COH, -C(CH₃)₂OH, -C(O)CH₃, -SCH₃, -S(O)₂CH₃, -C(O)N(CH₃)₂, -C(O)OC(CH₃)₃, -C(O)C(CH₃)₂, pyrazolyl, imidazolyl, pyrimidinyl, pyridinyl, 1,2-dihydropyrimidinyl, 1,6-dihydropyrimidinyl, piperidinyl, pyrrolidinyl, morpholinyl, and phenyl and wherein the phenyl is optionally substituted with methyl; and wherein R^c and R^d are each independently H, methyl, methoxy, and hydroxyl; stereoisomers thereof, and veterinary acceptable salts thereof.

11. A compound of Claim 10 that is a Formula (2A1) compound selected from the group consisting of:

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-benzylbenzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-(4-fluorobenzyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((diethylamino)(3,4-difluorophenyl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3,4-difluorophenyl)(methoxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3,4-difluorophenyl)(ethoxy)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and

(3S,6R,9S,12R,15S,18R,21S,24S)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-[[4-[(S)-morpholino]-[4-

(trifluoromethoxy)phenyl]methyl]phenyl]methyl]-1,7,13,19-tetraoxa-4,10,16,22-tetrazacyclotetracosane-2,5,8,11,14,17,20,23-octaone; stereoisomers thereof, and veterinary acceptable salts thereof;

or a Formula (4A) compound selected from the group consisting of:

(3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-((1H-indazol-6-yl)methyl)benzyl)-18-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((2,4-dimethyl-1H-pyrrol-3-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-(4-(benzofuran-3-yl(hydroxy)methyl)benzyl)-18-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-(oxazol-2-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-(furan-2-ylmethyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-((3-methylthiophen-2-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

2-(4-(((2R,5S,8R,11S,14R,17S,20R,23S)-14-benzyl-5,11,17,23-tetraisobutyl-4,8,10,16,20,22-hexamethyl-3,6,9,12,15,18,21,24-octaoxo-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2-yl)methyl)benzyl)furan-3-carbonitrile;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-(thiophen-3-ylmethyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-18-(4-((1-methyl-1H-pyrazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;

(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3-chlorothiophen-2-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-18-(4-((3-methoxythiophen-2-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((4-chlorothiazol-5-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-18-(4-((3,5-dimethylisoxazol-4-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-18-(4-((6-methoxy-2-methylpyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-6-benzyl-3,9,15,21-tetraisobutyl-18-(4-((6-methoxy-4-methylpyridin-3-yl)methyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((2-methylpyridin-3-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-4,6,10,16,18,22-hexamethyl-12,24-bis(4-((4-methylisothiazol-5-yl)methyl)benzyl)-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; stereoisomers thereof, and veterinary acceptable salts thereof;
or a Formula (1A1) compound of Claim 3 selected from the group consisting of
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(3-methoxybenzyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone;
(3S,6R,9S,12R,15S,18R,21S,24R)-3,9,15,21-tetraisobutyl-6,18-bis(4-(2-methoxybenzyl)benzyl)-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone; and

(3S,6R,9S,12R,15S,18R,21S,24R)-6,18-bis(4-((3,4-difluorophenyl)(1H-pyrazol-1-yl)methyl)benzyl)-3,9,15,21-tetraisobutyl-4,10,12,16,22,24-hexamethyl-1,7,13,19-tetraoxa-4,10,16,22-tetraazacyclotetracosan-2,5,8,11,14,17,20,23-octaone, stereoisomers thereof, and veterinary acceptable salts thereof.

12. A composition comprising a compound of any one of Claims 3 to 11, stereoisomers thereof, and veterinary acceptable salts thereof, and wherein the composition further comprises a veterinary acceptable excipient.

13. A method of treating or preventing a parasitic infection in an animal comprising administering a therapeutically effective amount of a compound of any one of Claims 1 to 11, stereoisomers thereof, and veterinary acceptable salts thereof, to an animal in need thereof.

14. The method of Claim 13 wherein the parasitic infection is an endoparasitic infection and the animal is a companion animal; and the compound is administered orally, topically, or parenterally.

15. Use of a compound of any one of Claims 1 to 11, stereoisomer thereof, and veterinary acceptable salt thereof, to prepare a medicament for treating an endoparasitic infection in a companion animal.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2019/031158

A. CLASSIFICATION OF SUBJECT MATTER INV. C07D413/14 C07D413/10 C07D417/14 C07D273/00 A61P33/00 A61K38/15 ADD. According to International Patent Classification (IPC) or to both national classification and IPC														
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) C07D A61P A61K Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, WPI Data														
C. DOCUMENTS CONSIDERED TO BE RELEVANT <table border="1"> <thead> <tr> <th>Category*</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>X</td> <td>WO 2016/187534 A1 (MERIAL INC [US]) 24 November 2016 (2016-11-24) claims 1-33; compounds 19-6 -----</td> <td>1-15</td> </tr> <tr> <td>A</td> <td>WO 99/66794 A1 (BAYER AG [DE]; JESCHKE PETER [DE] ET AL.) 29 December 1999 (1999-12-29) claims 1-2; compounds 15-18 -----</td> <td>1-15</td> </tr> <tr> <td>X</td> <td>WO 98/55469 A1 (BAYER AG [DE]; DYKER HUBERT [DE] ET AL.) 10 December 1998 (1998-12-10) claims 1-10; compound 37 ----- -/-</td> <td>1-15</td> </tr> </tbody> </table>			Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	X	WO 2016/187534 A1 (MERIAL INC [US]) 24 November 2016 (2016-11-24) claims 1-33; compounds 19-6 -----	1-15	A	WO 99/66794 A1 (BAYER AG [DE]; JESCHKE PETER [DE] ET AL.) 29 December 1999 (1999-12-29) claims 1-2; compounds 15-18 -----	1-15	X	WO 98/55469 A1 (BAYER AG [DE]; DYKER HUBERT [DE] ET AL.) 10 December 1998 (1998-12-10) claims 1-10; compound 37 ----- -/-	1-15
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
X	WO 2016/187534 A1 (MERIAL INC [US]) 24 November 2016 (2016-11-24) claims 1-33; compounds 19-6 -----	1-15												
A	WO 99/66794 A1 (BAYER AG [DE]; JESCHKE PETER [DE] ET AL.) 29 December 1999 (1999-12-29) claims 1-2; compounds 15-18 -----	1-15												
X	WO 98/55469 A1 (BAYER AG [DE]; DYKER HUBERT [DE] ET AL.) 10 December 1998 (1998-12-10) claims 1-10; compound 37 ----- -/-	1-15												
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.														
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family														
Date of the actual completion of the international search 16 July 2019		Date of mailing of the international search report 06/08/2019												
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Sáez Díaz, R												

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2019/031158

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MAKOTO OHYAMA ET AL: "Structure-Activity Relationship of Anthelmintic Cyclooctadepsipeptides", BIOSCI. BIOTECH. BIOCHEM., vol. 75, no. 7, 23 July 2011 (2011-07-23), pages 1354-1363, XP055350719, ISSN: 0916-8451, DOI: 10.1271/bbb.110129 table 1; compounds 27-31, 37-43, 55-59 -----	1-15
E	WO 2019/108591 A1 (ZOETIS SERVICES LLC [US]) 6 June 2019 (2019-06-06) claims 1-15; tables 1a, 1b, 1c, 2-12 -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2019/031158

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2016187534	A1	24-11-2016	AR 104714 A1 09-08-2017
		AU 2016264635 A1 14-12-2017	
		BR 112017024773 A2 06-11-2018	
		CA 2986478 A1 24-11-2016	
		CL 2017002936 A1 16-03-2018	
		CN 107835818 A 23-03-2018	
		CO 2017011738 A2 09-02-2018	
		EP 3298027 A1 28-03-2018	
		JP 2018516904 A 28-06-2018	
		KR 20180002886 A 08-01-2018	
		PH 12017502106 A1 28-05-2018	
		RU 2017144495 A 20-06-2019	
		TW 201706287 A 16-02-2017	
		US 2017022253 A1 26-01-2017	
		US 2018362589 A1 20-12-2018	
		UY 36683 A 30-12-2016	
		WO 2016187534 A1 24-11-2016	
WO 9966794	A1	29-12-1999	AT 229269 T 15-12-2002
		AU 751711 B2 22-08-2002	
		BR 9912199 A 10-04-2001	
		CA 2336129 A1 29-12-1999	
		CN 1306394 A 01-08-2001	
		DE 19828043 A1 30-12-1999	
		DK 1089625 T3 24-03-2003	
		EP 1089625 A1 11-04-2001	
		ES 2186371 T3 01-05-2003	
		HU 0102808 A2 28-11-2001	
		JP 4404484 B2 27-01-2010	
		JP 2002518415 A 25-06-2002	
		NZ 509105 A 20-12-2002	
		PT 1089625 E 30-04-2003	
		US 6630569 B1 07-10-2003	
		WO 9966794 A1 29-12-1999	
WO 9855469	A1	10-12-1998	AU 735424 B2 05-07-2001
		BR 9810412 A 22-08-2000	
		CA 2292698 A1 10-12-1998	
		CN 1259129 A 05-07-2000	
		EP 0986549 A1 22-03-2000	
		HK 1028605 A1 02-08-2002	
		JP 2002502398 A 22-01-2002	
		NZ 501478 A 25-05-2001	
		US 6355615 B1 12-03-2002	
		WO 9855469 A1 10-12-1998	
WO 2019108591	A1	06-06-2019	NONE