



(22) Date de dépôt/Filing Date: 1999/11/23

(41) Mise à la disp. pub./Open to Public Insp.: 2000/06/22

(45) Date de délivrance/Issue Date: 2013/11/19

(30) Priorité/Priority: 1998/12/22 (US60/113,345)

(51) Cl.Int./Int.Cl. *C12N 15/40* (2006.01),
A61K 31/70 (2006.01), *A61K 39/12* (2006.01),
A61K 48/00 (2006.01), *C07K 14/08* (2006.01),
C12N 15/63 (2006.01), *C12N 15/86* (2006.01),
C12N 5/10 (2006.01), *C12N 7/04* (2006.01)


(72) Inventeurs/Inventors:
CALVERT, JAY GREGORY, US;
SHEPPARD, MICHAEL GEORGE, US;
WELCH, SIAO-KUN WAN, US

(73) Propriétaire/Owner:
ZOETIS P LLC, US

(74) Agent: TORYS LLP

(54) Titre : CLONE D'ADNC INFECTIEUX DU VIRUS NORD-AMERICAIN DU SYNDROME DYSGENESIQUE
RESPIRATOIRE PORCIN (SDRP) ET SES UTILISATIONS

(54) Title: AN INFECTIOUS CDNA CLONE OF NORTH AMERICAN PORCINE REPRODUCTIVE AND RESPIRATORY
SYNDROME (PRRS) VIRUS AND USES THEREOF



(57) Abrégé/Abstract:

The invention provides isolated polynucleotide molecules, including plasmids; viral vectors; and transfected host cells that comprise a DNA sequence encoding an infectious RNA sequence encoding a North American PRRS virus; and also North American PRRS

(57) Abrégé(suite)/Abstract(continued):

viruses encoded thereby. The invention further provides isolated infectious RNA molecules encoding a North American PRRS virus. The invention also provides isolated polynucleotide molecules, infectious RNA molecules, viral vectors, and transfected host cells encoding genetically-modified North American PRRS viruses; and genetically-modified North American PRRS viruses encoded thereby. The invention also provides vaccines comprising such plasmids, RNA molecules, viral vectors, and North American PRRS viruses, and methods of using these vaccines in swine and in other animals. Also provided are isolated polynucleotide molecules, viral vectors, and transfected host cells that comprise a nucleotide sequence encoding a peptide of a North American PRRS virus. These viral vectors and transfected host cell lines are useful in providing peptides to compensate for mutated peptide coding sequences of DNA sequences encoding genetically-modified North American PRRS viruses so that functional virions can be generated.

**AN INFECTIOUS cDNA CLONE OF NORTH AMERICAN PORCINE REPRODUCTIVE AND
RESPIRATORY SYNDROME (PRRS) VIRUS AND USES THEREOF**

Abstract

The invention provides isolated polynucleotide molecules, including plasmids; viral vectors; and transfected host cells that comprise a DNA sequence encoding an infectious RNA sequence encoding a North American PRRS virus; and also North American PRRS viruses encoded thereby. The invention further provides isolated infectious RNA molecules encoding a North American PRRS virus. The invention also provides isolated polynucleotide molecules, infectious RNA molecules, viral vectors, and transfected host cells encoding genetically-modified North American PRRS viruses; and genetically-modified North American PRRS viruses encoded thereby. The invention also provides vaccines comprising such plasmids, RNA molecules, viral vectors, and North American PRRS viruses, and methods of using these vaccines in swine and in other animals. Also provided are isolated polynucleotide molecules, viral vectors, and transfected host cells that comprise a nucleotide sequence encoding a peptide of a North American PRRS virus. These viral vectors and transfected host cell lines are useful in providing peptides to compensate for mutated peptide coding sequences of DNA sequences encoding genetically-modified North American PRRS viruses so that functional virions can be generated.

PC10278A

-1-

**AN INFECTIOUS cDNA CLONE OF NORTH AMERICAN PORCINE REPRODUCTIVE AND
RESPIRATORY SYNDROME (PRRS) VIRUS AND USES THEREOF**

Field of the Invention

5 The present invention is in the field of animal health and is directed to infectious cDNA clones of positive polarity RNA viruses and the construction of vaccines, in particular, swine vaccines, using such cDNA clones.

Background of the Invention

10 Porcine reproductive and respiratory syndrome (PRRS) is a new disease of swine, first described in 1987 in North America and in 1990 in Europe. The disease has since spread to Asia and affects most of the major swine producing countries of the world. Primary symptoms are reproductive problems in sows and gilts, including late term abortions, stillbirths and mummies, and litters of small weak pigs which are born viremic and often fail to survive.
15 In addition, the syndrome manifests itself as a respiratory disease in young pigs which spreads horizontally and causes fever, lethargy, labored breathing, loss of appetite, slow growth, and occasionally death, often in association with other respiratory pathogens. The disease furthermore can be transmitted to sows and gilts via the semen of infected boars, either naturally or by artificial insemination. For these, and other reasons, PRRS has proven
20 to be a difficult disease to control and therefore one of the most economically damaging diseases to the swine industry.

 The causative agent of PRRS is the PRRS virus, which exists as two genetically and serologically distinct types (Murtaugh, M.P. *et al.*, 1995, Arch-Virol. 140, 1451-1460; Suarez, P. *et al.*, 1996, Virus Research 42:159-165). The two types are believed to have first entered
25 swine populations independently, one in North America and the other in Europe, in the 1980's, from unknown biological reservoirs, possibly of rodent or avian origin. The European type, represented by the prototype "Lelystad Virus", was isolated and sequenced in the Netherlands in 1991 (Terpstra, C. *et al.*, 1991, Vet. Quart. 13:131-136; Wensvoort, G. *et al.*, 1991, Vet. Quart. 13:121-130; Wensvoort, G. *et al.*, WO 92/213751992 (PCT/NL92/00096), 1992;
30 Meulenbergh, J.J.M. *et al.*, 1993, Virol. 192:62-72).

 Both the North American PRRS virus and the European PRRS virus are classified within the family *Arteriviridae*, which also includes equine arteritis virus, lactate dehydrogenase-elevating virus, and simian haemorrhagic fever virus. The arteriviruses are in turn placed within the order *Nidovirales*, which also includes the coronaviruses and
35 toroviruses. The nidoviruses are enveloped viruses having genomes consisting of a single strand of positive polarity RNA. The genomic RNA of a positive-stranded RNA virus fulfills the

50204-11

-2-

dual role in both storage and expression of genetic information. No DNA is involved in replication or transcription in nidoviruses. The reproduction of nidoviral genomic RNA is thus a combined process of genome replication and mRNA transcription. Moreover, some proteins are translated directly from the genomic RNA of nidoviruses. The molecular biology of the family *Arteriviridae* has recently been reviewed by Snijder and Meulenberg (Snijder, E.J. and Meulenberg, J.J.M., 1998, Journal of General Virology 79:961-979).

Currently available commercial vaccines against PRRS are either conventional modified live virus (cell culture, attenuated) or conventional killed (inactivated cell culture preparations of virulent virus). Several of these vaccines have been criticized based on safety and/or efficacy concerns. The development of a second generation of PRRS vaccines, based upon specific additions, deletions, and other modifications to the PRRS genome, is therefore highly desirable. However, since the PRRS viruses do not include any DNA intermediates during their replication, such vaccines have thus far awaited the construction of full-length cDNA clones of PRRS viruses for manipulation by molecular biology techniques at the DNA level. Very recently, a full-length infectious cDNA clone of the European PRRS virus has been reported (Meulenberg, J.J.M. *et al.*, 1998, *supra*; Meulenberg, J.J.M. *et al.*, 1988, J. Virol. 72, 380-387.).

50204-11

2a

Summary of the Invention

The subject invention provides an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is SEQ ID NO:1 or a sequence having at least about 93% nucleotide sequence identity thereto.

The subject invention further provides an isolated infectious RNA molecule encoded by the isolated polynucleotide molecule recited above, and isolated infectious RNA molecules having at least about 93% nucleotide sequence identity thereto, which isolated infectious RNA molecules each encode a North American PRRS virus.

The subject invention further provides the above-recited isolated polynucleotide molecule encoding the infectious RNA molecule in the form of a vector such as a plasmid.

The subject invention further provides a viral vector comprising a DNA encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is SEQ ID NO:1 or is a DNA sequence having at least about 93% nucleotide sequence identity thereto.

The subject invention further provides a transfected host cell comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is SEQ ID NO:1 or is a DNA sequence having at least about 93% nucleotide sequence identity thereto, which transfected host cell is capable of expressing the encoded North American PRRS virus.

64680-1720

2b

The subject invention further relates to a method for making a genetically modified North American PRRS virus, which method comprises mutating a DNA sequence of the present invention encoding an infectious RNA molecule
5 encoding the North American PRRS virus by any one or a combination of replacing, deleting, and adding one or more nucleotides, and expressing the genetically modified North American PRRS virus therefrom subsequent to said mutation.

The subject invention further relates to an
10 isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a genetically modified North American PRRS virus. In a preferred embodiment, the PRRS virus is genetically modified by any one or a combination of replacing, deleting, and adding one
15 or more nucleotides such that when it infects a porcine animal it is: a) unable to produce PRRS in the animal, and b) able to elicit an effective immunoprotective response against infection by a PRRS virus in the animal. In a particular embodiment, the DNA sequence is SEQ ID NO:1 or a
20 sequence having at least about 93% nucleotide sequence identity thereto, except for that it contains one or more mutations that genetically disable the encoded PRRS virus in its ability to produce PRRS.

The subject invention further relates to an
25 isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of
30 SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559

64680-1720

2c

of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1.

The subject invention further relates to a transfected host cell comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, which transfected host cell is capable of expressing the encoded North American PRRS virus.

The subject invention further relates to a plasmid capable of directly transfecting a suitable host cell and completely cellularly expressing a *Nidovirales* virus from the suitable host cell so transfected, the plasmid comprising a) a DNA sequence corresponding to (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, wherein said DNA sequence encodes an infectious RNA molecule encoding the *Nidovirales* virus; and b) a promoter capable of transcribing said infectious RNA molecule in said suitable host cell.

64680-1720

2d

The subject invention further relates to an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is derived by genetic modification by replacing, deleting, and/or adding one or more nucleotides to (i) SEQ ID NO:1; or (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine, wherein the DNA sequence encoding the infectious RNA molecule has at least 85% identity to SEQ ID NO:1.

The subject invention further relates to a transfected host cell comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is derived by genetic modification by replacing, deleting, and/or adding one or more nucleotides to (i) SEQ ID NO:1; or (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; wherein the DNA sequence encoding the infectious RNA molecule has at least 85% identity to SEQ ID NO:1, and which transfected host cell is capable of expressing the encoded North American PRRS virus.

The subject invention further relates to an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence comprises (i)

64680-1720

2e

SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, and wherein said DNA sequence further comprises one or more further nucleotide sequences that each encode a heterologous antigenic epitope, and wherein each heterologous antigenic epitope is capable of inducing an effective immunoprotective response against a particular pathogen corresponding to said heterologous antigenic epitope in a mammal or a bird.

The subject invention further relates to a transfected host cell comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence comprises (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, and wherein said DNA sequence further comprises one or more further nucleotide sequences that each encode a heterologous antigenic epitope, and wherein each heterologous antigenic epitope is capable of inducing an effective immunoprotective response against a particular pathogen corresponding to said heterologous antigenic epitope in a mammal or a bird.

The subject invention further relates to an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a North

64680-1720

2f

American PRRS virus, wherein said DNA sequence comprises (i) the sequence as set forth in SEQ ID NO:1; or (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the
5 nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1 and wherein said DNA
10 sequence further comprises one or more further nucleotide sequences that each encode detectable heterologous antigenic epitope.

The subject invention further relates to an isolated polynucleotide molecule comprising a DNA sequence
15 encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to
20 nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; and wherein said DNA sequence lacks one or more sequences in (i) or (ii) that encode a detectable antigenic epitope, and
25 wherein said DNA sequence has at least 85% identity to SEQ ID NO:1.

The subject invention further relates to an isolated polynucleotide molecule comprising one or more nucleotide sequences that each encode a peptide encoded by a
30 North American PRRS virus, wherein the genome of said North American PRRS virus is the same as an RNA molecule corresponding to (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and

64680-1720

2g

including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a
5 thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1.

The subject invention further relates to a transfected host cell comprising one or more nucleotide sequences that each encode a peptide encoded by a North
10 American PRRS virus, wherein each said nucleotide sequence is the same as an RNA molecule corresponding to (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to
15 nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine, or (iii) a sequence having at least 85% identity to SEQ ID NO:1, which transformed cell is capable of expressing said peptide
20 or peptides.

The subject invention further relates to a method for generating a functional virion of a North American PRRS virus encoded by an RNA sequence corresponding to (i) SEQ ID NO:1; (ii) the sequence beginning with and including
25 nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii)
30 a sequence having at least 85% identity to SEQ ID NO:1, and wherein said method comprises further transfecting the transfected host cell according to claim 15 with an

64680-1720

2h

infectious RNA molecule, plasmid or viral vector encoding the North American PRRS virus.

The subject invention further relates to a method for generating a functional virion of a North American PRRS virus encoded by an RNA sequence corresponding to (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, and wherein said method comprises transfecting a suitable cell with an infectious RNA molecule, plasmid or viral vector encoding the North American PRRS virus and with a helper virus that expresses the North American PRRS virus peptide or sequences of said North American PRRS virus, and separating North American PRRS virions expressed by the cell from said helper virus.

The subject invention further relates to a completely cellular method for preparing a genetically modified *Nidovirales* virus that is capable of eliciting an immunoprotective response in a mammal or a bird vaccinated therewith, wherein said method comprises (a) obtaining an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule which encodes a wild-type *Nidovirales* virus, and wherein said DNA sequence is (i) SEQ ID NO:1; or (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; (b)

64680-1720

2i

(b) genetically mutating the DNA sequence encoding the infectious RNA molecule which encodes the wild-type *Nidovirales* virus so as to obtain an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule which encodes a genetically modified *Nidovirales* virus having at least 85% identity to SEQ ID NO:1, and wherein said genetically modified *Nidovirales* virus remains able to elicit an effective immunoprotective response against infection by the wild-type *Nidovirales* virus in a mammal or a bird; and (c) completely cellularly expressing the genetically modified *Nidovirales* virus from the isolated polynucleotide molecule so obtained.

The subject invention further relates to a vaccine for protecting a porcine animal from infection by a PRRS virus, which vaccine comprises a genetically modified North American PRRS virus encoded by an infectious RNA molecule encoded by the isolated polynucleotide molecule defined herein, or said infectious RNA molecule, or said polynucleotide molecule in the form of a plasmid, or a viral vector comprising the polynucleotide molecule defined herein that is able to elicit an effective immunoprotective response against infection by a PRRS virus, in an amount effective to produce immunoprotection against infection by a PRRS virus; and a carrier acceptable for veterinary use.

The subject invention further relates to a vaccine for protecting a mammal or a bird from infection by a pathogen, which vaccine comprises a genetically modified North American PRRS virus encoded by an infectious RNA molecule encoded by the isolated polynucleotide molecule defined herein, or said infectious RNA molecule, or a plasmid comprising said polynucleotide molecule, in an amount effective to produce immunoprotection against infection by a pathogen from which the heterologous

64680-1720

2j

antigenic epitope or epitopes thereof or encoded thereby are derived; and a carrier acceptable for pharmaceutical or veterinary use.

The subject invention further relates to a vaccine
5 for protecting a porcine animal from infection by a PRRS virus and from infection by a swine pathogen other than a North American PRRS virus, which vaccine comprises a genetically modified North American PRRS virus encoded by an infectious RNA molecule encoded by the isolated
10 polynucleotide molecule defined herein, or said infectious RNA molecule, in an amount effective to produce immunoprotection against infection by the PRRS virus and against infection by the swine pathogen from which the heterologous antigenic epitope or epitopes thereof or
15 encoded thereby are derived; and a carrier acceptable for veterinary use.

The subject invention further relates to a vaccine for protecting a porcine animal from infection by a PRRS virus, which vaccine comprises a genetically modified North
20 American PRRS virus encoded by an infectious RNA molecule encoded by the isolated polynucleotide molecule defined herein, or said infectious RNA molecule or said isolated polynucleotide in the form of a plasmid; or a viral vector comprising said polynucleotide molecule in an amount
25 effective to produce immunoprotection against infection by the PRRS virus; and a carrier acceptable for veterinary use.

The subject invention further relates to a vaccine for protecting a porcine animal from infection by a PRRS virus, which vaccine comprises a genetically modified North
30 American PRRS virus encoded by an infectious RNA molecule encoded by the isolated polynucleotide molecule defined herein, or said infectious RNA molecule, or said isolated

64680-1720

2k

polynucleotide in the form of a plasmid; or a viral vector comprising said polynucleotide molecule in an amount effective to produce immunoprotection against infection by the PRRS virus; and a carrier acceptable for veterinary use.

5 The subject invention further relates to a genetically modified *Nidovirales* virus that is capable of eliciting an immunoprotective response in a mammal or a bird vaccinated therewith, which genetically modified *Nidovirales* virus is prepared by obtaining an isolated polynucleotide
10 molecule comprising a DNA sequence corresponding to (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of
15 an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, and expressing the genetically modified *Nidovirales* virus from the isolated polynucleotide molecule so obtained.

20 The subject invention further relates to a vaccine for protecting a mammal or a bird from infection by a *Nidovirales* virus, which vaccine comprises the genetically modified *Nidovirales* virus defined in claim 25 in an amount effective to elicit an effective immunoprotective response
25 against the wild-type *Nidovirales* virus in a mammal or a bird vaccinated therewith, and a carrier acceptable for pharmaceutical or veterinary use.

50204-11

-3-

The subject invention further provides an isolated infectious RNA molecule encoded by the isolated polynucleotide molecule recited above, and isolated infectious RNA molecules homologous thereto, which isolated infectious RNA molecules each encode a genetically modified North American PRRS virus, disabled in its ability to produce PRRS.

5 The subject invention further provides a genetically modified North American PRRS virus encoded by an infectious RNA molecule as recited above, which genetically modified North American PRRS virus is disabled such that when it infects a porcine animal it is unable to produce PRRS in the animal, yet is able to elicit an effective immunoprotective response against infection by a PRRS virus in the animal.

10 The subject invention further provides a viral vector comprising a DNA sequence encoding an infectious RNA molecule encoding a genetically modified North American PRRS virus as recited above.

15 The subject invention further provides a transfected host cell comprising a DNA sequence encoding an infectious RNA molecule encoding a genetically modified North American PRRS virus as recited above.

20 The subject invention further provides a vaccine for protecting a porcine animal from infection by a PRRS virus, which vaccine comprises a genetically modified North American PRRS virus as recited above; an infectious RNA molecule as recited above encoding the genetically modified North American PRRS virus; an isolated polynucleotide molecule recited above, in the form of a plasmid, encoding the genetically modified North American PRRS virus; or the above-recited viral vector encoding the genetically modified North American

50204-11

-4-

PRRS virus; in an amount effective to produce immunoprotection against infection by a PRRS virus; and a carrier acceptable for veterinary use.

The subject invention further provides a method for protecting a porcine animal from infection by a PRRS virus, which comprises vaccinating the animal with an amount of the
5 above-recited vaccine that is effective to produce immunoprotection against infection by a PRRS virus.

The present invention further provides any of the aforementioned polynucleotide molecules further comprising a nucleotide sequence encoding a heterologous antigenic epitope, as well as corresponding infectious RNA molecules, vectors, transfected host cells,
10 genetically modified North American PRRS viruses, vaccines, and methods of administering such vaccines to mammals and/or birds. Such heterologous antigenic epitopes can be from any antigenic epitope, presently known in the art, or to be determined in the future. In a non-limiting embodiment, such antigenic epitope is from a pathogen capable of pathogenically infecting a bird or mammal other than porcine animal, for example a human, and is capable of
15 inducing an effective immunoprotective response against said pathogen. In another non-limiting embodiment, such antigenic epitope is from a swine pathogen other than a North American PRRS virus and is capable of inducing an effective immunoprotective response against said swine pathogen. In another non-limiting embodiment, such antigenic epitope is a detectable antigenic epitope.

20 The present invention further provides any of the aforementioned polynucleotide molecules, but which lack one or more detectable antigenic epitopes, as well as corresponding infectious RNA molecules, vectors, transfected host cells, genetically modified North American PRRS viruses, vaccines, and methods of administering such vaccines to mammals and/or birds.

25 The subject invention further provides an isolated polynucleotide molecule comprising one or more nucleotide sequences that encode a peptide encoded by a North American PRRS virus, wherein the genome sequence of said North American PRRS virus is SEQ ID NO:1 or a sequence having at least about 93% nucleotide sequence identity thereto.

30 The subject invention further provides a transfected host cell comprising one or more nucleotide sequences that encode a peptide encoded by a North American PRRS virus, wherein the genome sequence of said North American PRRS virus is SEQ ID NO:1 or a sequence having at least about 93% nucleotide sequence identity thereto.

35 The subject invention further provides a genetically modified *Nidovirales* virus that is able to elicit an effective immunoprotective response in a mammal or a bird vaccinated therewith, which genetically modified *Nidovirales* virus is prepared by obtaining an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule

50204-11

-5-

encoding a wild-type *Nidovirales* virus, genetically mutating the DNA sequence so as to obtain an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a genetically modified *Nidovirales* virus which virus is unable to produce a pathogenic infection yet is able to elicit an effective immunoprotective response against infection by the wild-type *Nidovirales* virus in a mammal or a bird, and expressing the genetically modified *Nidovirales* virus from the isolated polynucleotide molecule so obtained.

The subject invention further provides a completely cellular method for preparing a genetically modified *Nidovirales* virus that is capable of eliciting an immunoprotective response in a mammal or a bird vaccinated therewith, which method comprises obtaining an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a wild-type *Nidovirales* virus, genetically mutating the DNA so as to obtain an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a genetically modified *Nidovirales* virus which virus is unable to produce a pathogenic infection yet able to elicit an effective immunoprotective response against infection by the wild-type *Nidovirales* virus in a mammal or a bird, and completely cellularly expressing the genetically modified *Nidovirales* virus from the isolated polynucleotide molecule so obtained.

The subject invention further provides a vaccine for protecting a mammal or a bird from infection by a *Nidovirales* virus, which vaccine comprises a genetically modified *Nidovirales* virus as described above in an amount effective to elicit an effective immunoprotective response against the wild-type *Nidovirales* virus in a mammal or a bird vaccinated therewith, and a carrier acceptable for pharmaceutical or veterinary use.

Brief Description of the Figures

FIGURE 1: Cloning strategy for construction of full-length infectious cDNA clone of North American PRRS virus, pT7P129A. Arrowheads represent T7 promoter sequences.

FIGURE 2: Serum viremia following infection with P129A or recombinant PRRS virus rP129A-1. Determined by plaque assay on MARC-145 cells. The lower limit of detection is 5 pfu/ml (or 0.7 on the log scale).

FIGURE 3: Anti-PRRS virus serum antibody following infection with P129A or recombinant PRRS virus rP129A-1. Determined by HerdChek PRRS ELISA assay (IDEXX (Westbrook, Maine, USA)).

Detailed Description of the Invention

Production and manipulation of the isolated polynucleotide molecules described herein are within the skill in the art and can be carried out according to recombinant techniques described, among other places, in Maniatis, *et al.*, 1989, Molecular Cloning, A

50204-11

-6-

Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Ausubel, et al., 1989, Current Protocols In Molecular Biology, Greene Publishing Associates & Wiley Interscience, NY; Sambrook, et al., 1989, Molecular Cloning: A Laboratory Manual, 2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Innis et al. (eds), 1995, PCR Strategies, Academic Press, Inc., San Diego; and Erlich (ed), 1992, PCR Technology, Oxford University Press, New York.

A. *Isolated Polynucleotide Molecules and RNA Molecules Encoding a North American PRRS Virus, and Isolated Polynucleotide Molecules and RNA Molecules Encoding Genetically Modified North American PRRS Viruses:*

The subject invention provides isolated polynucleotide molecules comprising DNA sequences that encode infectious RNA molecules that encode a North American PRRS virus. The present invention further provides isolated polynucleotide molecules comprising DNA sequences that encode infectious RNA molecules that encode genetically modified North American PRRS viruses.

In particular, the subject invention provides an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule that encodes a North American PRRS virus, wherein said DNA sequence is SEQ ID NO:1 or a sequence homologous thereto. In a preferred embodiment, the present invention provides an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule that encodes a North American PRRS virus, wherein said DNA sequence is the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, except that the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine. Said DNA sequence encodes an infectious RNA molecule that is the RNA genome of the North American PRRS isolate P129.

It is understood that terms herein referring to nucleic acid molecules such as "isolated polynucleotide molecule", "nucleotide sequence", "open reading frame (ORF)", and the like, unless otherwise specified, include both DNA and RNA molecules and include both single-stranded and double-stranded molecules. Also, when reference to a particular sequence from the "Sequence Listing" section of the subject application is made, it is intended, unless otherwise specified, to refer to both the DNA of the "Sequence Listing", as well as RNA corresponding to the DNA sequence, and includes sequences complementary to the DNA and RNA sequences. In such contexts in this application, "corresponding to" refers to sequences of DNA and RNA that are identical to one another but for the fact that the RNA

sequence contains uracil in place of thymine and the backbone of the RNA molecule contains ribose instead of deoxyribose.

For example, SEQ ID NO:1 is a DNA sequence corresponding to the RNA genome of a North American PRRS virus. Thus, a DNA sequence complementary to the DNA sequence
5 set forth in SEQ ID NO:1 is a template for, *i.e.* is complementary to or "encodes", the RNA genome of the North American PRRS virus (*i.e.*, RNA that encodes the North American PRRS virus). Nonetheless, a reference herein to SEQ ID NO:1 includes both the RNA sequence corresponding to SEQ ID NO:1 and a DNA sequence complementary to SEQ ID NO:1.

Furthermore, when reference is made herein to sequences homologous to a
10 sequence in the Sequence Listing, it is to be understood that sequences homologous to a sequence corresponding to the sequence in the Sequence Listing and sequences homologous to a sequence complementary to the sequence in the Sequence Listing are also included.

An "infectious RNA molecule", for purposes of the present invention, is an RNA
15 molecule that encodes the necessary elements for viral replication, transcription, and translation into a functional virion in a suitable host cell, provided, if necessary, with a peptide or peptides that compensate for any genetic modifications, *e.g.* sequence deletions, in the RNA molecule.

An "isolated infectious RNA molecule" refers to a composition of matter comprising
20 the aforementioned infectious RNA molecule purified to any detectable degree from its naturally occurring state, if such RNA molecule does indeed occur in nature. Likewise, an "isolated polynucleotide molecule" refers to a composition of matter comprising a polynucleotide molecule of the present invention purified to any detectable degree from its naturally occurring state, if any.

For purposes of the present invention, the nucleotide sequence of a second
25 polynucleotide molecule (either RNA or DNA) is "homologous" to the nucleotide sequence of a first polynucleotide molecule where the nucleotide sequence of the second polynucleotide molecule encodes the same polyaminoacid as the nucleotide sequence of the first polynucleotide molecule as based on the degeneracy of the genetic code, or when it encodes
30 a polyaminoacid that is sufficiently similar to the polyaminoacid encoded by the nucleotide sequence of the first polynucleotide molecule so as to be useful in practicing the present invention. For purposes of the present invention, a polynucleotide molecule is useful in practicing the present invention where it can be used as a diagnostic probe to detect the presence of the North American PRRS virus in a fluid or tissue sample of an infected pig, *e.g.*
35 by standard hybridization or amplification techniques. It is to be understood that the polyaminoacid encoded by the nucleotide sequence of the polynucleotide molecule can

comprise a group of two or more polyaminoacids. Generally, the nucleotide sequence of a second polynucleotide molecule is homologous to the nucleotide sequence of a first polynucleotide molecule if it has at least about 70% nucleotide sequence identity to the nucleotide sequence of the first polynucleotide molecule as based on the BLASTN algorithm
5 (National Center for Biotechnology Information, otherwise known as NCBI, (Bethesda, Maryland, USA) of the United States National Institute of Health). Preferably, a homologous nucleotide sequence has at least about 75% nucleotide sequence identity, even more preferably at least about 85% nucleotide sequence identity. Since the genetic code is degenerate, a homologous nucleotide sequence can include any number of "silent" base
10 changes, *i.e.* nucleotide substitutions that nonetheless encode the same amino acid. A homologous nucleotide sequence can further contain non-silent mutations, *i.e.* base substitutions, deletions, or additions resulting in amino acid differences in the encoded polyaminoacid, so long as the sequence remains at least about 70% identical to the polyaminoacid encoded by the first nucleotide sequence or otherwise is useful for practicing
15 the present invention. Homologous nucleotide sequences can be determined by comparison of nucleotide sequences, for example by using BLASTN, above. Alternatively, homologous nucleotide sequences can be determined by hybridization under selected conditions. For example, the nucleotide sequence of a second polynucleotide molecule is homologous to SEQ ID NO:1 if it hybridizes to the complement of SEQ ID NO:1 under moderately stringent
20 conditions, *e.g.*, hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.2xSSC/0.1% SDS at 42°C (see Ausubel *et al.* above), or conditions which will otherwise result in hybridization of sequences that encode a North American PRRS virus as defined below. In another embodiment, a second nucleotide sequence is homologous to SEQ ID NO:1 if it hybridizes to the complement of SEQ
25 ID NO:1 under highly stringent conditions, *e.g.* hybridization to filter-bound DNA in 0.5 M NaHPO₄, 7% SDS, 1 mM EDTA at 65°C, and washing in 0.1xSSC/0.1% SDS at 68°C (Ausubel *et al.*, above).

It is furthermore to be understood that the isolated polynucleotide molecules and the isolated RNA molecules of the present invention include both synthetic molecules and
30 molecules obtained through recombinant techniques, such as by *in vitro* cloning and transcription.

As used herein, the term "PRRS" encompasses disease symptoms in swine caused by a PRRS virus infection. Examples of such symptoms include, but are not limited to, abortion in pregnant females, and slow growth, respiratory difficulties, loss of appetite, and
35 mortality in young pigs. As used herein, a PRRS virus that is "unable to produce PRRS" refers to a virus that can infect a pig, but which does not produce any disease symptoms

normally associated with a PRRS infection in the pig, or produces such symptoms, but to a lesser degree, or produces a fewer number of such symptoms, or both.

The terms "porcine" and "swine" are used interchangeably herein and refer to any animal that is a member of the family Suidae such as, for example, a pig. "Mammals" include
5 any warm-blooded vertebrates of the Mammalia class, including humans.

The term "PRRS virus", as used herein, unless otherwise indicated, means any strain of either the North American or European PRRS viruses.

The term "North American PRRS virus" means any PRRS virus having genetic characteristics associated with a North American PRRS virus isolate, such as, but not limited
10 to the PRRS virus that was first isolated in the United States around the early 1990's (see, e.g., Collins, J.E., *et al.*, 1992, *J. Vet. Diagn. Invest.* 4:117-126); North American PRRS virus isolate MN-1b (Kwang, J. *et al.*, 1994, *J. Vet. Diagn. Invest.* 6:293-296); the Quebec IAF-exp91 strain of PRRS (Mardassi, H. *et al.*, 1995, *Arch. Virol.* 140:1405-1418); and North American PRRS virus isolate VR 2385 (Meng, X.-J. *et al.*, 1994, *J. Gen. Virol.* 75:1795-1801). Genetic
15 characteristics refers to genomic nucleotide sequence similarity and aminoacid sequence similarity shared by North American PRRS virus strains. For purposes of the present invention, a North American PRRS virus is a virus that is encoded by an RNA sequence the same as or homologous to SEQ ID NO:1, wherein the term "homologous" is as defined previously. Thus, strains of North American PRRS viruses have, preferably, at least about
20 70% genomic nucleotide sequence identity with SEQ ID NO:1, and more preferably at least about 75% genomic nucleotide sequence identity with SEQ ID NO:1, at least about 85% genomic nucleotide sequence identity with SEQ ID NO:1 being even more preferred.

The term "European PRRS virus" refers to any strain of PRRS virus having the genetic characteristics associated with the PRRS virus that was first isolated in Europe
25 around 1991 (see, e.g., Wensvoort, G., *et al.*, 1991, *Vet. Q.* 13:121-130). "European PRRS virus" is also sometimes referred to in the art as "Lelystad virus".

Unless otherwise indicated, a North American PRRS virus is "useful in practicing the present invention" if its characteristics are within the definition of a North American PRRS virus set forth herein. For example, a virus encoded by one of the isolated polynucleotide
30 molecules of the present invention is a "North American PRRS virus useful in practicing the present invention" if it, e.g., has genetic characteristics associated with a North American PRRS virus.

Other polyaminoacids are "useful in practicing the present invention", e.g., peptides encoded by polynucleotide sequences homologous to North American PRRS virus ORFs, if
35 they can compensate for an RNA molecule encoding a genetically modified PRRS virus,

deficient in a gene essential for expressing functional PRRS virions, in a transfected host cell so that functional PRRS virions can be generated by the cell.

The term "open reading frame", or "ORF", as used herein, means the minimal nucleotide sequence required to encode a particular PRRS virus protein without an
5 intervening stop codon.

Terms such as "suitable host cell" and "appropriate host cell", unless otherwise indicated, refer to cells into which RNA molecules (or isolated polynucleotide molecules or viral vectors comprising DNA sequences encoding such RNA molecules) of the present invention can be transformed or transfected. "Suitable host cells" for transfection with such
10 RNA molecules, isolated polynucleotide molecules, or viral vectors, include mammalian, particularly porcine, and avian cells, and are described in further detail below.

A "functional virion" is a virus particle that is able to enter a cell capable of hosting a PRRS virus, and express genes of its particular RNA genome (either an unmodified genome or a genetically modified genome as described herein) within the cell. Cells capable of
15 hosting a PRRS virus include porcine alveolar macrophage cells and MARC 145 monkey kidney cells. Other mammalian or avian cells, especially other porcine cells, may also serve as suitable host cells for PRRS virions.

The isolated polynucleotide molecules of the present invention encode North American PRRS viruses that can be used to prepare live, killed, or attenuated vaccines using
20 art-recognized methods for protecting swine from infection by a PRRS virus, as described in further detail below. These isolated polynucleotide molecules are also useful as vectors for delivering heterologous genes into mammals, including swine, or birds, as is also described in detail below. Furthermore, these isolated polynucleotide molecules are useful because they can be mutated using molecular biology techniques to encode genetically-modified North
25 American PRRS viruses useful, *inter alia*, as vaccines for protecting swine from PRRS infection. Such genetically-modified North American PRRS viruses, as well as vaccines comprising them, are also described in further detail below.

Accordingly, the subject invention further provides a method for making a genetically modified North American PRRS virus, which method comprises mutating the DNA sequence
30 encoding an infectious RNA molecule which encodes the North American PRRS virus as described above, and expressing the genetically modified North American PRRS virus using a suitable expression system. A North American PRRS virus, either wild-type or genetically modified, can be expressed from an isolated polynucleotide molecule using suitable expression systems generally known in the art, examples of which are described in this
35 application. For example, the isolated polynucleotide molecule can be in the form of a

-11-

plasmid capable of expressing the encoded virus in a suitable host cell *in vitro*, as is described in further detail below.

The term "genetically modified", as used herein and unless otherwise indicated, means genetically mutated, *i.e.* having one or more nucleotides replaced, deleted and/or added. Polynucleotide molecules can be genetically mutated using recombinant techniques known to those of ordinary skill in the art, including by site-directed mutagenesis, or by random mutagenesis such as by exposure to chemical mutagens or to radiation, as known in the art. In one embodiment, genetic modification of the North American PRRS virus of the present invention renders the virus unable to replicate effectively, or reduces its ability to replicate effectively, in a bird or mammal in which the wild-type virus otherwise can effectively replicate. In another embodiment, the genetically modified North American PRRS virus of the present invention remains able to replicate effectively in birds or mammals infected therewith. "Effective replication" means the ability to multiply and produce progeny viruses (virions) in an infected animal, *i.e.* the ability to "productively infect" an animal.

The subject invention further provides an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule which encodes a genetically modified North American PRRS virus that is unable to produce PRRS in a porcine animal, wherein the DNA sequence encoding the infectious RNA molecule encoding said North American PRRS virus is SEQ ID NO:1 or a sequence homologous thereto, except that it contains one or more mutations that genetically disable the encoded PRRS virus in its ability to produce PRRS. "Genetically disabled" means that the PRRS virus is unable to produce PRRS in a swine animal infected therewith.

In one embodiment, the genetically modified North American PRRS virus disabled in its ability to cause PRRS is able to elicit an effective immunoprotective response against infection by a PRRS virus in a swine animal. Accordingly, the subject invention also provides an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule which encodes a North American PRRS virus that is genetically modified such that when it infects a porcine animal it: a) is unable to produce PRRS in the animal, and b) is able to elicit an effective immunoprotective response against infection by a PRRS virus in the animal, wherein the DNA sequence encoding said North American PRRS virus is SEQ ID NO:1 or a sequence homologous thereto, except that it contains one or more mutations that genetically disable the encoded PRRS virus in its ability to produce PRRS.

The term "immune response" for purposes of this invention means the production of antibodies and/or cells (such as T lymphocytes) that are directed against, or assist in the decomposition or inhibition of, a particular antigenic epitope or particular antigenic epitopes. The phrases "an effective immunoprotective response", "immunoprotection", and like terms,

-12-

for purposes of the present invention, mean an immune response that is directed against one or more antigenic epitopes of a pathogen so as to protect against infection by the pathogen in a vaccinated animal. For purposes of the present invention, protection against infection by a pathogen includes not only the absolute prevention of infection, but also any detectable
5 reduction in the degree or rate of infection by a pathogen, or any detectable reduction in the severity of the disease or any symptom or condition resulting from infection by the pathogen in the vaccinated animal as compared to an unvaccinated infected animal. An effective immunoprotective response can be induced in animals that have not previously been infected with the pathogen and/or are not infected with the pathogen at the time of vaccination. An
10 effective immunoprotective response can also be induced in an animal already infected with the pathogen at the time of vaccination.

An "antigenic epitope" is, unless otherwise indicated, a molecule that is able to elicit an immune response in a particular animal or species. Antigenic epitopes are proteinaceous molecules, i.e. polyaminoacid sequences, optionally comprising non-protein groups such as
15 carbohydrate moieties and/or lipid moieties.

The term "pathogenically infecting" used herein refers to the ability of a pathogen to infect an animal and cause a disease in the animal. As an example, a PRRS virus is capable of pathogenically infecting a porcine animal since it can cause PRRS in swine. However, although a PRRS virus may be able to infect, either productively or non-productively, a bird or
20 another mammal, such as a human, it does not pathogenically infect any animal other than a porcine animal since it does not cause any disease in animals other than porcine animals.

The genetically modified North American PRRS viruses encoded by the above-described isolated polynucleotide molecules are, in one embodiment, able to elicit an effective immunoprotective response against infection by a PRRS virus. Such genetically modified
25 North American PRRS viruses are preferably able to elicit an effective immunoprotective response against any strain of PRRS viruses, including both European and North American strains.

In one embodiment, the mutation or mutations in the isolated polynucleotide molecule encoding the genetically disabled North American PRRS virus are non-silent and occur in one
30 or more open reading frames of the nucleotide sequence encoding the North American PRRS virus; i.e., the mutation or mutations occur in one or more of the sequences within the nucleotide sequence encoding the North American PRRS virus that are the same as or homologous to ORFs 1a, 1b, 2, 3, 4, 5, 6, or 7 of SEQ ID NO:1. In another embodiment, the mutation or mutations occur in one or more noncoding regions of the North American PRRS
35 virus genome, such as, for example, in the leader sequence of the North American PRRS virus genome; i.e., the mutation or mutations occur within the sequence that is the same as or

homologous to the sequence of nucleotides 1-191 of SEQ ID NO:1. In the same isolated polynucleotide molecule, mutations can occur in both coding and noncoding regions.

As used herein, unless otherwise indicated, "noncoding regions" of the nucleotide sequence encoding the North American PRRS virus refer to those sequences of RNA that are not translated into a protein and those sequences of cDNA that encode such RNA sequences. Coding regions refer to those sequences of RNA from which North American PRRS virus proteins are expressed, and also refer to cDNA that encodes such RNA sequences. Likewise, "ORFs" refer both to RNA sequences that encode North American PRRS virus proteins and to cDNA sequence encoding such RNA sequences.

Determining suitable locations for a mutation or mutations that will encode a North American PRRS virus that is genetically disabled so that it is unable to produce PRRS yet remains able to elicit an effective immunoprotective response against infection by a PRRS virus can be made based on the SEQ ID NO:1 provided herein. One of ordinary skill can refer to the sequence of the infectious cDNA clone of North American PRRS virus provided by this invention, make sequence changes which will result in a mutation, and test the viruses encoded thereby both for their ability to produce PRRS in swine, and to elicit an effective immunoprotective response against infection by a PRRS virus. In so doing, one of ordinary skill can refer to techniques known in the art and also those described and/or exemplified herein.

For example, an ORF of the sequence encoding the infectious RNA molecule encoding the North American PRRS virus can be mutated and the resulting genetically modified North American PRRS virus tested for its ability to cause PRRS. The ORF of a North American PRRS virus encodes proteins as follows: ORF 1a encodes a polyprotein comprising protease function; ORF 1b encodes a polyprotein comprising replicase (RNA polymerase) and helicase functions; ORFs 2, 3, and 4 encode small membrane glycoproteins; ORF 5 encodes a major envelope glycoprotein; ORF 6 encodes a nonglycosylated integral membrane protein; and ORF 7 encodes a nucleocapsid protein. Genetic mutations of one or more of these ORFs can be used in preparing the genetically modified North American PRRS viruses described *infra*.

The subject invention also provides an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus that is genetically modified such that it comprises one or more heterologous antigenic epitopes, wherein the DNA sequence encoding the RNA molecule encoding the North American PRRS virus is SEQ ID NO:1 or a sequence homologous thereto, and further comprising one or more additional nucleotide sequences that each encode a heterologous

antigenic epitope, and wherein each heterologous antigenic epitope is capable of inducing an effective immunoprotective response against a particular pathogen in a mammal or a bird.

A pathogen against which an effective immunoprotective response can be induced by means of the above recited aspect of the present invention is any pathogen, such as a virus,
 5 bacteria, fungus, or protozoan, capable of causing a disease in a mammal or bird, which pathogen comprises or has associated therewith one or more antigenic epitopes which can be used to induce an effective immunoprotective response against the pathogen in the mammal or bird.

The term "heterologous antigenic epitope" for purposes of the present invention
 10 means an antigenic epitope, as defined above, not normally found in a wild-type North American PRRS virus. A nucleotide sequence encoding a heterologous antigenic epitope can be inserted into a North American PRRS viral genome using known recombinant techniques. Antigenic epitopes useful as heterologous antigenic epitopes for the present invention include additional North American PRRS virus antigenic epitopes, antigenic epitopes from European
 15 PRRS viruses, antigenic epitopes from swine pathogens other than PRRS viruses, or antigenic epitopes from pathogens that pathogenically infect birds or mammals other than swine, including humans. Sequences encoding such antigenic epitopes are known in the art or are provided herein. For example, a second North American PRRS virus envelope protein, encoded by North American PRRS ORF 5 described herein, can be inserted into a DNA
 20 sequence encoding an RNA molecule encoding a North American PRRS virus of the present invention to generate a genetically modified North American PRRS virus comprising an additional envelope protein as a heterologous antigenic epitope. Such a genetically modified North American PRRS virus can be used to induce a more effective immunoprotective response against PRRS viruses in a porcine animal vaccinated therewith.

25 Examples of an antigenic epitope from a swine pathogen other than a North American PRRS virus include, but are not limited to, an antigenic epitope from a swine pathogen selected from the group consisting of European PRRS, porcine parvovirus, porcine circovirus, a porcine rotavirus, swine influenza, pseudorabies virus, transmissible gastroenteritis virus, porcine respiratory coronavirus, classical swine fever virus, African swine fever virus,
 30 encephalomyocarditis virus, porcine paramyxovirus, *Actinobacillus pleuropneumoni*, *Bacillus anthraci*, *Bordetella bronchiseptica*, *Clostridium haemolyticum*, *Clostridium perfringens*, *Clostridium tetani*, *Escherichia coli*, *Erysipelothrix rhusiopathiae*, *Haemophilus parasuis*, *Leptospira spp.*, *Mycoplasma hyopneumoniae*, *Mycoplasma hyorhinis*, *Pasteurella haemolytica*, *Pasteurella multocida*, *Salmonella choleraesuis*, *Salmonella typhimurium*,
 35 *Streptococcus equismilis*, and *Streptococcus suis*. Nucleotide sequences encoding antigenic epitopes from the aforementioned swine pathogens are known in the art and can be obtained

64680-1480

15

from public gene databases.

If the heterologous antigenic epitopes are antigenic epitopes from one or more other swine pathogens, then the isolated polynucleotide molecule can further contain one or more mutations that genetically disable the encoded PRRS virus in its ability to produce PRRS. Such isolated polynucleotide molecules and the viruses they encode are useful for preparing vaccines for protecting swine against the swine pathogen or pathogens from which the heterologous antigenic epitopes are derived.

In a preferred embodiment, the genetically modified North American PRRS is able to elicit an effective immunoprotective response against infection by a PRRS virus in a porcine animal. Such isolated polynucleotide molecules and the viruses they encode are useful for preparing dual-function vaccines for protecting swine against infection by both a North American PRRS virus and the swine pathogen or pathogens from which the heterologous antigenic epitopes are derived. In another preferred embodiment, the genetically modified North American PRRS virus useful in a dual-function vaccine is genetically disabled.

The isolated polynucleotide molecules of the present invention comprising nucleotide sequences encoding heterologous antigenic epitopes can be prepared as described above based on the sequence encoding a North American PRRS virus described herein using known techniques in molecular biology.

In a further preferred embodiment, a heterologous antigenic epitope of the genetically modified North American PRRS virus of the present invention is a detectable antigenic epitope. Such isolated polynucleotide molecules and the North American PRRS viruses they encode are useful, *inter alia*, for studying PRRS infections in swine, determining successfully vaccinated swine, and/or for distinguishing vaccinated swine from swine infected by a wild-type PRRS virus. Preferably, such isolated polynucleotide molecules further contain one or more mutations that genetically disable the encoded PRRS virus in its ability to produce PRRS, and more preferably are able to elicit an effective immunoprotective response in a porcine animal against infection by a PRRS virus.

Heterologous antigenic epitopes that are detectable, and the sequences that encode them, are known in the art. Techniques for detecting such antigenic epitopes are also known in the art and include serological detection of antibody specific to the heterologous antigenic epitope by means of, for example, Western blot, ELISA, or fluorescently labeled antibodies capable of binding to the antibodies specific to the heterologous antigenic epitope. Techniques for serological detection useful in practicing the present invention can be found in texts recognized in the art, such as Coligan, J.E., *et al.* (eds), 1998, Current Protocols in Immunology, John Willey & Sons, Inc.

50204-11

-16-

Alternatively, the heterologous antigenic epitope itself can be detected by, for example, contacting samples that potentially comprise the antigenic epitope with fluorescently-labeled antibodies or radioactively-labeled antibodies that specifically bind to the antigenic epitopes.

5 The present invention further provides an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule which encodes a genetically modified North American PRRS virus that detectably lacks a North American PRRS virus antigenic epitope, wherein the DNA sequence encoding the RNA molecule encoding the North American PRRS virus is SEQ ID NO:1 or a sequence homologous thereto, except that it lacks
10 one or more nucleotide sequences encoding a detectable North American PRRS virus antigenic epitope. Such isolated polynucleotide molecules are useful for distinguishing between swine infected with a recombinant North American PRRS virus of the present invention and swine infected with a wild-type PRRS virus. For example, animals vaccinated with killed, live or attenuated North American PRRS virus encoded by such an isolated
15 polynucleotide molecule can be distinguished from animals infected with wild-type PRRS based on the absence of antibodies specific to the missing antigenic epitope, or based on the absence of the antigenic epitope itself: If antibodies specific to the missing antigenic epitope, or if the antigenic epitope itself, are detected in the animal, then the animal was exposed to and infected by a wild-type PRRS virus. Means for detecting antigenic epitopes and
20 antibodies specific thereto are known in the art, as discussed above. Preferably, such an isolated polynucleotide molecule further contains one or more mutations that genetically disable the encoded PRRS virus in its ability to produce PRRS. More preferably, the encoded virus remains able to elicit an effective immunoprotective response against infection by a PRRS virus.

25

B. Plasmids Encoding a North American PRRS Virus or a Genetically Modified North American PRRS Virus:

The present invention also provides any of the above-described isolated polynucleotide molecules in the form of a plasmid capable of expressing the North American
30 PRRS virus encoded thereby.

Plasmids of the present invention can express the encoded North American PRRS virus outside of a living organism, to produce North American PRRS viruses of the invention useful, *inter alia*, for preparing vaccines. In one embodiment, a plasmid of the present invention capable of expressing a North American PRRS virus outside of a living organism is
35 a plasmid wherein transcription of viral RNA therefrom occurs *in vitro* (*i.e.* extracellularly); the resulting viral RNA molecule is transfected into a suitable host cell using known mechanisms

-17-

of transfection, such as electroporation, lipofection (in some cases using a commercially available reagent, such as Lipofectin™ (Life Technologies Inc., Rockville, Maryland, USA)), or DEAE dextran mediated transfection. Other methods of transfection are known in the art and can be employed in the present invention. An example of such a plasmid for *in vitro* transcription of North American PRRS viral RNA is the plasmid pT7P129A (ATCC Accession No. 203488). Any promoter useful for *in vitro* transcription can be used in such plasmids of this invention. T7 is one such promoter, but other promoters can be used, such as an SP6 promoter or a T3 promoter. The sequences of such promoters can be artificially synthesized or cloned from commercially available plasmids. Suitable plasmids for preparing such plasmids capable of expressing North American PRRS virus include, but are not limited to, general purpose cloning vector plasmids such as pCR2.1 (Invitrogen, Carlsbad, California, USA), pBR322, and pUC18. A nucleotide sequence of the present invention encoding the North American PRRS virus can be inserted into any of these plasmids using known recombinant techniques. Other plasmids into which the polynucleotide molecules of the present invention can be inserted will be recognized by those of ordinary skill in the art.

Suitable conditions for *in vitro* transcription of viral RNA from any of the above-described recombinant plasmids comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus depends on the type of plasmid, for example, its particular promoter, and can be ascertained by one of ordinary skill in the art. For example, if a plasmid of the present invention is based on a pCR2.1 plasmid comprising a T7 promoter, then an example of suitable conditions for *in vitro* transcription includes reacting the plasmid with T7 RNA polymerase and ribonucleotides in a standard buffer and incubating the reaction at 37°C for about 30 minutes. In some cases, commercial kits are available for transcribing RNA from a particular plasmid, and such kits can be used in the present invention. The reaction mixture following transcription can be directly transfected into a suitable host cell without purification, or the transcribed North American PRRS virus RNA can be purified by known RNA purification techniques, for example by organic (e.g. phenol) extraction and alcohol (e.g. ethanol or isopropanol) precipitation, prior to transfection.

Practically any mammalian or avian cell culture can be transfected with the North American PRRS virus RNA obtained as described above in order to generate a first round of North American PRRS virions. An example of cells which one might find particularly useful because of their ready availability and ease of use are BHK (baby hamster kidney) cells. However, if one wishes to generate a cell culture capable of sustained production of North American PRRS virions, then porcine alveolar macrophage cells or MARC-145 cells (Kim, H.S., *et al.*, *supra*) are preferred since these cells excrete high levels of new generation PRRS virions subsequent to PRRS virus infection. Other cell lines derived from the MA-104

cell line may also be used for sustained generation of North American PRRS virions of the present invention. Primary porcine alveolar macrophage cells can be obtained by lung lavages from pigs, and the MARC-145 monkey kidney cell line can be obtained from the National Veterinary Services Laboratories otherwise known as NVSL (Ames, Iowa, USA).

5 In another embodiment, a plasmid capable of expressing a North American PRRS virus of the present invention outside of a living organism is a plasmid which is transfected into a suitable host cell, for example by electroporation or lipofection, transcription of the infectious RNA molecule and expression of the North American PRRS virus therefrom occurring within the transfected host cell. The transfected host cell therefore generates North
10 American PRRS virions. Such a completely cellular method has heretofore never been disclosed or suggested for any virus within the order of *Nidovirales*. Because of possible cryptic splicing and termination sequences present in the RNA genome of viruses of the *Nidovirales* order, a completely cellular method of expressing a *Nidovirales* virus was believed unlikely. Cryptic sequences include RNA splice donor and splice acceptor sequences, which
15 could cause inappropriate splicing of the RNA transcript, as well as polyadenylation sequences, which could cause premature termination by the cellular RNA polymerase II. The present invention demonstrates, however, that the presence of such sequences in a plasmid comprising a cDNA clone of a Nidovirus does not prevent the plasmid's ability to express the Nidovirus when the plasmid is directly transfected into a suitable host cell.

20 Accordingly, the subject invention also provides plasmids and a completely cellular method for expressing a *Nidovirales* virus, wherein the plasmid comprises: a) a DNA sequence encoding an infectious RNA molecule encoding the *Nidovirales* virus; and b) a promoter capable of transcribing said encoding sequence in a cell, wherein said promoter is in operative association with the DNA sequence encoding the infectious RNA molecule. The
25 method comprises transfecting a suitable host cell with such a plasmid, subjecting the transfected host cell to conditions suitable for expression of gene sequences transfected therein, and collecting the expressed *Nidovirales* virus therefrom. An example of a plasmid suitable for completely cellular expression of North American PRRS virus outside of a living organism is the plasmid pCMV-S-P129 (ATCC Accession No. 203489). In a preferred
30 embodiment, the promoter of such a plasmid is a CMV promoter. In a preferred embodiment, a plasmid of the invention suitable for a completely cellular method of expressing a *Nidovirales* virus comprises a eukaryotic promoter, such as a CMV promoter, immediately upstream and adjacent to the nucleotide sequence encoding the *Nidovirales* virus. In a preferred embodiment, the nucleotide sequence encoding the *Nidovirales* virus encodes a
35 PRRS virus, either European or North American. Other examples of *Nidovirales* viruses that can be expressed by means of the above-described completely cellular method include other

Arteriviruses such as, equine arteritis virus, lactate dehydrogenase-elevating virus, and simian haemorrhagic fever virus; viruses that are members of the genus *Coronaviridae*, such as, but not limited to, feline infectious peritonitis virus, feline enteric coronavirus, canine coronavirus, bovine coronavirus, porcine respiratory coronavirus, turkey coronavirus, porcine transmissible gastroenteritis virus, human coronavirus, murine hepatitis virus, and avian infectious bronchitis virus; and members of the genus *Toroviridae*, such as, but not limited to, Berne virus, Breda virus, and human torovirus. Thus, plasmids suitable for completely cellular expression comprising a nucleotide sequence encoding one of these viruses are also encompassed by the present invention.

10 Suitable plasmids that can be used to prepare recombinant plasmids of the present invention for completely cellular expression outside of a living organism of a *Nidovirales* virus, such as a PRRS virus, include virtually any plasmid useful for transfection and expression in eukaryotic cells. An examples of a plasmid suitable for preparing recombinant plasmids of the present invention for completely cellular expression of a *Nidovirales* virus is the plasmid
15 pCMVbeta (Clontech, Palo Alto, California, USA). Other plasmids which are able to transfect and express genes in eukaryotic cells which can be used to prepare plasmids of the present invention include, but are not limited to, pcDNA3.1, pRc/RSV, and pZeoSV2 (all from Invitrogen); and pCMV-Sport3 and pSV-Sport1 (both from Life Technologies Inc.). However, almost any eukaryotic expression vector will work for the present invention. Constructs based
20 on cosmids can also be used for completely cellular *ex vivo* expression of a *Nidovirales* virus.

 Suitable host cells for the completely cellular method of the present invention for expressing PRRS virus include porcine alveolar macrophage cells and the MARC-145 cells, described above. Methods of transfecting these cells with a plasmid are basically the same as those methods for transfecting cells with viral RNA described above. Such methods
25 include, but are not limited to, electroporation, lipofection, DEAE dextran mediated transfection, and calcium phosphate coprecipitation.

 Once host cells, such as porcine alveolar macrophage cells or a MARC-145 cells, have been transfected according to the subject invention, either with viral RNA or with a plasmid comprising a nucleotide sequence encoding a virus, then the cells can be frozen at
30 about -80°C or below for storage for up to several years. For longer periods of time, *i.e.* decades, storage in liquid nitrogen is preferred. If relatively frequent use of the encoded virus is envisioned, then cells hosting the virus can also be maintained (unfrozen) in culture using known techniques, for shorter periods of time. Moreover, viral particles excreted by such cells can be stored frozen at about -80°C or below as a source of virus. Transfection of such cell
35 lines with the polynucleotide molecule encoding the virus can be confirmed if desired, for example, by testing exhausted medium excreted by the cell line for a PRRS virus antigen

using an immunofluorescent antibody test. Antibodies which are specific for PRRS virus antigens are known in the art (see, e.g., Collins, E.J., *et al.*, WO 93/03760 March 4, 1993).

In another embodiment, a plasmid of the present invention comprising a nucleotide sequence encoding a North American PRRS virus is suitable for *in vivo* expression of the
5 North American PRRS virus, *i.e.* expression in a living organism. Plasmids which can be used for preparing recombinant plasmids for *in vivo* expression of a North American PRRS virus include, but are not limited to the plasmids capable of transfecting eukaryotic cells described above, such as pCMVbeta.

Animals that can be transfected with plasmids of the present invention include
10 mammals and birds. If the animal is other than a porcine animal, for example, a mallard duck, then the plasmid can comprise a nucleotide sequence encoding a North American PRRS virus comprising further antigenic epitopes from pathogens which are capable of pathogenically infecting the animal; in such a case, the plasmid will encode a North American PRRS virus serving as a vector for transporting epitopes into the animal. If the animal is a
15 porcine animal, then the plasmid can usefully encode any of the North American PRRS viruses described herein, including the genetically-modified North American PRRS viruses described herein.

*C. Viral Vectors Encoding a North American PRRS Virus, Including Viral Vectors
20 Encoding Genetically Modified North American PRRS Viruses:*

The present invention also provides viral vectors comprising a DNA sequence encoding an infectious RNA molecule encoding any of the North American PRRS viruses described herein, including the genetically-modified North American PRRS viruses described herein. Such viral vectors are useful for transfecting eukaryotic cells for production of PRRS
25 viruses of the present invention outside of a living organism, or for transfecting swine, or other mammals, or avians, with the sequence encoding the North American PRRS virus, for *in vivo* expression of the North American PRRS virus therein.

Some examples of viruses that can be used as vectors for preparing the viral vectors of the present invention include, but are not limited to, swine viruses such as, but not limited
30 to, swine pox virus, pseudorabies virus, or African swine fever virus. Such swine viruses can be obtained from The National Veterinary Services Laboratories (Ames, Iowa, USA) of the United States Department of Agriculture; the American Type Culture Collection, otherwise known as the ATCC (Manassas, Virginia, USA); and other known sources. Recombinant viral vectors based on suitable swine viruses such as the aforementioned swine viruses are useful
35 for transfecting swine animals with a nucleotide sequence encoding a North American PRRS virus of the present invention.

Viral vectors comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus of the present invention based on these and other viruses can be prepared using known recombinant techniques described in texts such as those cited previously in this application.

5

D. Transfected Host cells Encoding or a Genetically Modified North American PRRS Viruses:

The present invention also provides transfected host cells that comprise a DNA sequence encoding an infectious RNA molecule encoding any of the North American PRRS viruses described herein, including the genetically-modified North American PRRS viruses described herein, which transfected host cells are capable of expressing the North American PRRS virus. Such transfected host cells are useful for producing North American PRRS viruses of the present invention. Examples of transfected host cells of the present invention include the transfected porcine alveolar macrophage cells and the transfected MARC-145 cells described above.

Other transfected host cells of the invention include, but are not limited to, transfected MA-104 cells and other derivatives of MA-104 cells that are transfected; transfected Baby Hamster Kidney (BHK) cells; transfected Chinese Hamster Ovary (CHO) cells; and African Green Monkey kidney cells other than MA-104 cells or MARC-145 cells, such as VERO cells; that are transfected.

E. North American PRRS Viruses, Including Genetically Modified North American PRRS Viruses:

The present invention also provides North American PRRS viruses as described herein, including genetically-modified North American PRRS viruses as described herein, expressed and/or encoded by any of the above-described isolated polynucleotide molecules, RNA molecules, plasmids, viral vectors, or transfected host cells.

In certain situations, for example where the North American PRRS virus is to be used in a vaccine for swine and the North American PRRS virus has not been genetically modified as described above so as to be unable to cause PRRS, it is desirable to treat the North American PRRS virus, for example by inactivating or attenuating it, so that it is unable to cause PRRS in swine to which it is administered. Known methods can be used to inactivate a North American PRRS virus of the present invention so that it is unable to cause PRRS in an animal. Examples of such methods include, but are not limited to, treatment with formaldehyde, BEI (binary ethyleneimine), or BPL (beta-propiolactone). Methods of attenuation are also known in the art, and such methods can be used to attenuate a North

American PRRS virus of the present invention. A North American PRRS virus of the present invention can, for example, be attenuated by serial passage in cell culture.

If a North American PRRS virus of the present invention is for use in an animal other than a porcine animal, or if it has been genetically modified as described herein so that it is
5 unable to produce PRRS in a porcine animal, then it is not necessary to treat the virus as described in the preceding paragraph prior to using it in a vaccine.

F. Vaccines and Uses Thereof:

The present invention also provides vaccines comprising North American PRRS
10 viruses, including genetically modified North American PRRS viruses disabled in their ability to produce PRRS in a swine animal as described herein; infectious RNA molecules and plasmids encoding such North American PRRS viruses as described herein; and viral vectors encoding such North American PRRS viruses and isolated RNA molecules as described herein. The invention also provides methods for protecting animals from infection comprising
15 vaccination with such vaccines.

In a preferred embodiment, the subject invention provides a vaccine comprising a genetically modified North American PRRS virus comprising one or more heterologous antigenic epitopes as described herein, an infectious RNA molecule encoding such a genetically modified North American PRRS virus, or a plasmid as described herein encoding
20 such a genetically modified North American PRRS virus, in an amount effective to elicit an immunoprotective response against infection by the pathogen or pathogens from which the heterologous antigenic epitope(s) are derived, and a carrier acceptable for pharmaceutical or veterinary use.

Such vaccines can be used to protect from infection a mammal or a bird capable of
25 being pathogenically infected by the pathogen or pathogens from which the heterologous antigenic epitope(s) are derived. Accordingly, the subject invention also provides a method for protecting a mammal or a bird from infection by a pathogen, which comprises vaccinating the mammal or bird with an amount of the vaccine described in the preceding paragraph effective to elicit an immunoprotective response in the mammal or bird from infection by the
30 pathogen.

In a further preferred embodiment, the vaccine comprises a genetically modified North American PRRS virus, or an infectious RNA molecule or plasmid encoding such a genetically modified North American PRRS virus, comprising or encoding one or more heterologous antigenic epitopes from a swine pathogen other than a North American PRRS virus. These
35 vaccines are useful for protecting swine from infection by the swine pathogen or pathogens from which the heterologous antigenic epitope(s) are derived. If such a vaccine comprises the

50204-11

-23-

genetically modified North American PRRS virus, the genetic modification of the North American PRRS virus preferably renders the virus unable to cause PRRS in swine. In another preferred embodiment, the genetically modified North American PRRS virus in the vaccine is able to elicit an immunoprotective response against infection by a PRRS virus, thus providing a dual-vaccine for swine, protecting swine from infection by the swine pathogen or pathogens from which the heterologous antigenic epitope(s) are derived as well as from infection by a PRRS virus. If the vaccine comprises an infectious RNA molecule or a plasmid encoding a genetically-modified North American PRRS virus comprising one or more heterologous antigenic epitopes from another swine pathogen, then the sequence encoding the infectious RNA molecule encoding the genetically modified PRRS virus preferably comprises one or more further mutations that genetically disable the encoded North American PRRS virus so that it is unable to cause PRRS. In another preferred embodiment, the encoded genetically modified, disabled North American PRRS virus is able to elicit an immunoprotective response against a PRRS infection in a swine animal, thus providing a dual-vaccine for swine, able to protect swine from infection by the swine pathogen or pathogens from which the heterologous antigenic epitope(s) are derived as well as from infection by a PRRS virus. All of these vaccines also further comprise a carrier acceptable for veterinary use.

The present invention further provides a method for protecting a porcine animal from infection by a swine pathogen or pathogens other than a North American PRRS virus and, optionally, for simultaneously protecting the swine animal from infection by a PRRS virus, which comprises vaccinating the animal with an amount of a vaccine described in the preceding paragraph effective to elicit an immunoprotective response in the swine animal from an infection by the swine pathogen or pathogens and optionally, by a PRRS virus.

Vaccines of the present invention can be formulated following accepted convention to include acceptable carriers for animals, including humans (if applicable), such as standard buffers, stabilizers, diluents, preservatives, and/or solubilizers, and can also be formulated to facilitate sustained release. Diluents include water, saline, dextrose, ethanol, glycerol, and the like. Additives for isotonicity include sodium chloride, dextrose, mannitol, sorbitol, and lactose, among others. Stabilizers include albumin, among others. Other suitable vaccine vehicles and additives, including those that are particularly useful in formulating modified live vaccines, are known or will be apparent to those skilled in the art. See, e.g., Remington's Pharmaceutical Science, 18th ed., 1990, Mack Publishing.

Vaccines of the present invention can further comprise one or more additional immunomodulatory components such as, e.g., an adjuvant or cytokine, among others. Non-

50204-11

-24-

limiting examples of adjuvants that can be used in the vaccine of the present invention include the RIBI^{*} adjuvant system (Ribi Inc., Hamilton, MT), alum, mineral gels such as aluminum hydroxide gel, oil-in-water emulsions, water-in-oil emulsions such as, e.g., Freund's complete and incomplete adjuvants, Block copolymer (CytRx, Atlanta GA), QS-21^{*} (Cambridge Biotech Inc., Cambridge MA), SAF-M^{*} (Chiron, Emeryville CA), AMPHIGEN[®] adjuvant, saponin, Quil A^{*} or other saponin fraction, monophosphoryl lipid A, and Avridine lipid-amine adjuvant. Non-limiting examples of oil-in-water emulsions useful in the vaccine of the invention include modified SEAM62^{*} and SEAM 1/2^{*} formulations. Modified SEAM62 is an oil-in-water emulsion containing 5% (v/v) squalene (Sigma), 1% (v/v) SPAN[®] 85 detergent (ICI Surfactants), 0.7% (v/v) TWEEN[®] 80 detergent (ICI Surfactants), 2.5% (v/v) ethanol, 200 µg/ml Quil A, 100 µg/ml cholesterol, and 0.5% (v/v) lecithin. Modified SEAM 1/2 is an oil-in-water emulsion comprising 5% (v/v) squalene, 1% (v/v) SPAN[®] 85 detergent, 0.7% (v/v) Tween 80 detergent, 2.5% (v/v) ethanol, 100 µg/ml Quil A, and 50 µg/ml cholesterol. Other immunomodulatory agents that can be included in the vaccine include, e.g., one or more interleukins, interferons, or other known cytokines.

Vaccines of the present invention can optionally be formulated for sustained release of the virus, infectious RNA molecule, plasmid, or viral vector of the present invention. Examples of such sustained release formulations include virus, infectious RNA molecule, plasmid, or viral vector in combination with composites of biocompatible polymers, such as, e.g., poly(lactic acid), poly(lactic-co-glycolic acid), methylcellulose, hyaluronic acid, collagen and the like. The structure, selection and use of degradable polymers in drug delivery vehicles have been reviewed in several publications, including A. Domb *et al.*, 1992, *Polymers for Advanced Technologies* 3: 279-292.

Additional guidance in selecting and using polymers in pharmaceutical formulations can be found in texts known in the art, for example M. Chasin and R. Langer (eds), 1990, "Biodegradable Polymers as Drug Delivery Systems" in: Drugs and the Pharmaceutical Sciences, Vol. 45, M. Dekker, NY.

Alternatively, or additionally, the virus, plasmid, or viral vector can be microencapsulated to improve administration and efficacy. Methods for microencapsulating antigens are well-known in the art, and include techniques described, e.g., in U.S. Patent 3,137,631; U.S. Patent 3,959,457; U.S. Patent 4,205,060; U.S. Patent 4,606,940; U.S. Patent 4,744,933; U.S. Patent 5,132,117; and International Patent Publication WO 95/28227.

Liposomes can also be used to provide for the sustained release of virus, plasmid, or viral vector. Details concerning how to make and use liposomal formulations can be found in, among other places, U.S. Patent 4,016,100; U.S. Patent 4,452,747; U.S. Patent 4,921,706;

*Trade-mark

50204-11

-25-

U.S. Patent 4,927,637; U.S. Patent 4,944,948; U.S. Patent 5,008,050; and U.S. Patent 5,009,956.

An effective amount of any of the above-described vaccines can be determined by conventional means, starting with a low dose of virus, plasmid or viral vector, and then
5 increasing the dosage while monitoring the effects. An effective amount may be obtained after a single administration of a vaccine or after multiple administrations of a vaccine. Known factors can be taken into consideration when determining an optimal dose per animal. These include the species, size, age and general condition of the animal, the presence of other drugs in the animal, and the like. The actual dosage is preferably chosen after consideration of the results
10 from other animal studies.

One method of detecting whether an adequate immune response has been achieved is to determine seroconversion and antibody titer in the animal after vaccination. The timing of vaccination and the number of boosters, if any, will preferably be determined by a doctor or veterinarian based on analysis of all relevant factors, some of which are described above.

15 The effective dose amount of virus, infectious RNA molecule, plasmid, or viral vector, of the present invention can be determined using known techniques, taking into account factors that can be determined by one of ordinary skill in the art such as the weight of the animal to be vaccinated. The dose amount of virus of the present invention in a vaccine of the present invention preferably ranges from about 10^1 to about 10^9 pfu (plaque forming units), more
20 preferably from about 10^2 to about 10^8 pfu, and most preferably from about 10^3 to about 10^7 pfu. The dose amount of a plasmid of the present invention in a vaccine of the present invention preferably ranges from about 0.1 μ g to about 100mg, more preferably from about 1 μ g to about 10mg, even more preferably from about 10 μ g to about 1mg. The dose amount of an infectious RNA molecule of the present invention in a vaccine of the present invention preferably ranges
25 from about 0.1 μ g to about 100mg, more preferably from about 1 μ g to about 10mg, even more preferably from about 10 μ g to about 1mg. The dose amount of a viral vector of the present invention in a vaccine of the present invention preferably ranges from about 10^1 pfu to about 10^9 pfu, more preferably from about 10^2 pfu to about 10^8 pfu, and even more preferably from about 10^3 to about 10^7 pfu. A suitable dosage size ranges from about 0.5 ml to about 10 ml, and more
30 preferably from about 1 ml to about 5 ml.

The present invention further provides a method of preparing a vaccine comprising a North American PRRS virus, infectious RNA molecule, plasmid, or viral vector described herein, which method comprises combining an effective amount of one of the North American PRRS virus, infectious RNA molecule, plasmid, or viral vector of the present invention, with a carrier
35 acceptable for pharmaceutical or veterinary use.

It is to be understood that the term "North American PRRS viruses of the present invention" and like terms, unless otherwise indicated, include any of the genetically modified North American PRRS viruses described herein as well as the unmodified North American PRRS virus described herein encoded by SEQ ID NO:1 or a sequence homologous thereto.

5

G. Isolated Polynucleotide Molecules and Transfected Host cells Encoding North American PRRS Virus Peptides, and Methods for Making Functional North American PRRS Virions:

The present invention also provides an isolated polynucleotide molecule comprising
10 one or more nucleotide sequences that encode a peptide encoded by a North American PRRS virus, wherein the genome sequence of said North American PRRS virus is the same as or homologous to an RNA molecule corresponding to SEQ ID NO:1. As used herein, terms such as "North American PRRS virus peptide" mean a peptide that is expressed by a North American PRRS virus. Such a peptide can be, but is not necessarily, specific to North
15 American PRRS viruses.

In a preferred embodiment, an isolated polynucleotide molecule of the present invention encoding a North American PRRS virus peptide comprises a sequence or sequences independently selected from SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, and a sequence
20 homologous to any of said sequences. Such isolated polynucleotide molecules are useful in plasmids or for preparing viral vectors for transfecting a suitable host cell to create a "helper" cell comprising one or more nucleotide sequences that encode a peptide encoded by a North American PRRS virus, wherein the genome sequence of said North American PRRS virus is the same as or homologous to an RNA sequence corresponding to SEQ ID NO:1, which
25 helper cell is useful in the preparation of functional virions of genetically modified North American PRRS viruses of the present invention, which viruses have been genetically modified as described above so that they are missing from their RNA genome the sequence(s) encoding the peptide or peptides encoded by the helper cell.

Accordingly, the subject invention also includes plasmids and viral vectors comprising
30 one or more nucleotide sequences encoding a peptide encoded by a North American PRRS virus, wherein the genome sequence of said North American PRRS virus is the same as or homologous to an RNA sequence corresponding to SEQ ID NO:1. Such plasmids of the invention can be based on those plasmids described above useful for preparing plasmids comprising a nucleotide sequence encoding a North American PRRS virus. Such viral
35 vectors of the invention can be based on, for example, retrovirus vectors or adeno-associated

viral vectors. These plasmids and viral vectors are useful for preparing the helper cells described in the preceding paragraph.

The present invention also thus provides a helper cell, *i.e.*, a transfected host cell comprising one or more nucleotide sequences that encode a peptide encoded by a North American PRRS virus, wherein the genome sequence of said North American PRRS virus is the same as or homologous to an RNA sequence corresponding to SEQ ID NO:1. In preferred embodiments, the transfected host cell comprises a nucleotide sequence or sequences independently selected from SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9 and a sequence homologous to any of said sequences. These helper cells are useful as described above for providing peptides for completing infectious RNA molecules deficient in sequences encoding the peptide(s) encoded by the helper cell so that functional virions of a genetically modified North American PRRS virus can be generated by the cell.

Suitable cells for this aspect of the invention include the cells described above which are suitable for expressing North American PRRS viruses, such as the porcine alveolar macrophage cells and the MARC-145 cells. However, practically any mammalian or avian cell can be used. As discussed above, if one wishes to obtain a transfected host cell capable of reinfection by PRRS virions, and thus able to generate multiple generations of functional genetically modified North American PRRS virions, porcine alveolar macrophage cells and MARC-145 cells are preferred.

The subject invention thus further provides a method for generating a functional virion of a genetically modified North American PRRS virus encoded by an RNA sequence corresponding to SEQ ID NO:1 or a sequence homologous thereto comprising one or more mutations that disable one or more peptide coding sequences, which method comprises transfecting a helper cell as described in the preceding paragraph with an infectious RNA molecule, plasmid, or viral vector encoding the genetically modified North American PRRS virus, which helper cell comprises a nucleotide sequence or sequences encoding the North American PRRS virus peptide or peptides of the disabled peptide coding sequence(s) of the genetically modified North American PRRS virus.

The subject invention further provides a method for generating a functional virion of a genetically modified North American PRRS virus encoded by an RNA sequence corresponding to SEQ ID NO:1 or a sequence homologous thereto comprising one or more mutations in one or more peptide coding sequences, which method comprises transfecting a suitable cell with both an infectious RNA molecule, plasmid, or viral vector encoding the genetically modified North American PRRS virus and a helper virus that expresses in the cell the North American PRRS virus peptide or peptides of the mutated peptide coding sequence

or sequences of the modified North American PRRS virus, and separating genetically modified North American PRRS virions from the helper virus. Methods of separation are known in the art and include use of conditional lethal helper viruses that distinguish (kill) the helper virus under certain conditions wherein the expressed North American PRRS virus is not killed. Other known methods of separation include physical methods of separation subsequent to expression of both helper virions and North American PRRS virions, for example by density gradient centrifugation. Suitable cells for this method of the invention include the cells capable of infection by a PRRS virus, such as porcine alveolar macrophage cells and MARC-145 cells. Other cells which are able to be infected by a PRRS virus are described previously in this application, for example the MA-104 cell line or other cell lines derived from the MA-104 cell line. An example of a helper virus is a PRRS virus, preferably a North American PRRS virus isolate, such as P129. Any virus, either wild-type, isolated, or recombinant, that expresses PRRS peptides can be used as a helper virus.

Infectious RNA molecules, plasmids, and viral vectors that encode a genetically modified North American PRRS virus useful in either of the above methods for generating a functional virion of a genetically modified North American PRRS virus are those infectious RNA molecules, plasmids, and viral vectors described in this application.

H. *Immunoprotective Genetically modified Nidovirales Viruses and Vaccines Comprising Them:*

The subject invention further provides a genetically modified *Nidovirales* virus that is capable of eliciting an immunoprotective response in a mammal or a bird vaccinated therewith, which genetically modified *Nidovirales* virus is prepared by obtaining an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a wild-type *Nidovirales* virus, genetically mutating the DNA sequence encoding the infectious RNA molecule encoding the wild-type *Nidovirales* virus so as to obtain an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a genetically modified *Nidovirales* virus which virus is able to elicit an effective immunoprotective response against infection by the wild-type *Nidovirales* virus in a mammal or a bird, and expressing the genetically modified *Nidovirales* virus from the isolated polynucleotide molecule so obtained.

The subject invention further provides a method for preparing a genetically modified *Nidovirales* virus that is capable of eliciting an immunoprotective response in a mammal or a bird vaccinated therewith, which method comprises obtaining an isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a wild-type *Nidovirales* virus, genetically mutating the DNA sequence encoding the infectious RNA molecule encoding the wild-type *Nidovirales* virus so as to obtain an isolated polynucleotide

50204-11

-29-

molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a genetically modified *Nidovirales* virus which virus is able to elicit an effective immunoprotective response against infection by the wild-type *Nidovirales* virus in a mammal or a bird, and expressing the genetically modified *Nidovirales* virus from the isolated polynucleotide molecule so obtained.

DNA sequences encoding infectious RNA molecule encoding a *Nidovirales* virus include the DNA sequences which are the same as or homologous to SEQ ID NO:1 described herein, which encode a North American PRRS virus. DNA sequences encoding infectious RNA molecules encoding *Nidovirales* viruses other than a North American PRRS virus are known in the art and can be obtained from known sources, for example the genetic databases cited above. Other examples of some DNA sequences encoding infectious RNA molecules encoding *Nidovirales* viruses which can be used in the subject invention include the DNA sequence encoding European PRRS virus described in Meulenbergh, J.J.M., *et al.*, 1993, *supra*; the DNA sequence encoding equine arteritis virus described in van Dinten, L.D., *et al.*, 1997, *Proc. Natl. Acad. Sci. USA*, 94(3):991-6; and the DNA sequence encoding equine arteritis virus described in den Boon, J.A., *et al.*, 1991, *J. Virol.* 65(6):2910-20.

Nidovirales

viruses which can be genetically modified by means of the present invention can be any *Nidovirales* virus for which a coding DNA sequence is obtained. Some examples of *Nidovirales* viruses are described in previous sections of the present application.

Once a DNA sequence encoding an infectious RNA molecule encoding a *Nidovirales* virus is obtained, isolated polynucleotide molecules, for example plasmids, comprising the DNA sequence encoding an infectious RNA molecule encoding a *Nidovirales* virus can be synthesized using recombinant techniques described above.

The sequences encoding the infectious RNA molecule encoding a *Nidovirales* virus can be genetically mutated as described above for the North American PRRS virus so as to obtain *Nidovirales* viruses comprising genetic modifications such as an inability to cause disease in an animal infected therewith, inclusion of heterologous antigenic epitopes that are able to elicit an immunoprotective response in an animal, inclusion of heterologous antigenic epitopes that are detectable, and deletion of detectable antigenic epitopes. Such genetic modifications are described above. Genetic mutations so as to encode genetically modified *Nidovirales* viruses can occur in coding and/or noncoding regions of the DNA sequence encoding the infectious RNA molecule encoding the *Nidovirales* virus. In one embodiment, one or more genetic mutations occur in an ORF encoding a viral peptide of the *Nidovirales* virus.

50204-11

-30-

"Wild-type *Nidovirales* virus", as used herein, means a *Nidovirales* virus whose genome has not intentionally been genetically mutated, such as a *Nidovirales* virus occurring in nature and isolates thereof.

The subject invention further provides a vaccine for protecting a mammal or a bird from infection by a *Nidovirales* virus, which comprises a genetically modified *Nidovirales* virus as described above in an amount effective to elicit an effective immunoprotective response against the wild-type *Nidovirales* virus in a mammal or a bird vaccinated therewith, and a carrier acceptable for pharmaceutical or veterinary use. These vaccines of the present invention can be formulated using the techniques for formulating vaccines described above.

The following examples are provided to merely illustrate aspects of the subject invention. They are not intended, and should not be construed, to limit the invention set forth in the claims and more fully described herein.

Examples

Example I. Preparation of an Infectious cDNA Clone of a North American PRRS Virus Isolate.

Source of PRRS virus and MARC-145 cells: A North American PRRS virus isolate designated P129 was obtained from Drs. Gregory W. Stevenson, William G. Van Alstine, and Charles L. Kanitz of Purdue University's Animal Disease Diagnostic Laboratory in West Lafayette, Indiana. The P129 virus was originally isolated in the autumn of 1995 from a swine herd in southern Indiana experiencing a severe PRRS outbreak. This farm had no previous history of PRRS problems or PRRS vaccination. The P129 isolate was more virulent than several other field isolates from the same time period and geographic area, in that it produced more severe and more consistent respiratory disease in young pigs. The virus was initially isolated on primary porcine alveolar macrophage (the natural host cell), and subsequently passaged on MARC-145 cells (Kim, H.S. *et al.*, 1993, Arch. Virol. 133:477-483). Genes encoding structural proteins of P129 were found to be homologous to corresponding known North American PRRS gene sequences.

The MARC-145 cell line that was used to propagate PRRS viruses is a clone of the MA-104 Rhesus Macaque Monkey Kidney cell line. The MARC-145 cells were obtained from the National Veterinary Services Laboratories (NVSL, Ames, Iowa) of the USDA. These cells have been tested and found negative for mycoplasmas and for common porcine extraneous agents. MARC-145 cells are routinely grown at 37C in OptiMEM* (Life Technologies Inc.) with 2% fetal bovine serum and antibiotics.

Five biological clones were plaque purified from the P129 virus stock, and these were designated P129A through P129E. Plaque purification was carried out by infecting

*Trade-mark

50204-11

-31-

monolayers of MARC-145 cells with P129 virus, adding an overlay of OptiMEM containing 1.25% SeaPlaque* agarose (FMC BioProducts), 2% fetal bovine serum, and antibiotics. Plaques were clearly visible following incubation for 7 days, when 5 well-isolated plaques were picked and passaged onto fresh MARC-145 monolayers. When cytopathic effect (virus induced cell death) became apparent the progeny virus from each of these cultures was subjected to another round of plaque purification. One well-isolated plaque from each of the five clones was picked and expanded to produce large stocks. The 5 clones were tested for virulence in young pigs, either individually (clones A and E) or in combination (clones B-D, or clones A-E). In all cases, the plaque purified virus replicated well in pigs and caused clinical disease. The severity of clinical symptoms was less than that caused by the uncloned P129 virus, even when all five clones were used together. P129A was chosen for sequencing, and was used in subsequent molecular manipulations.

Determination of the genome sequence of P129A: Plaque purified virus P129A was used for sequence determination after 10 serial passages from the pig (including two plaque purifications and one subsequent passage). SEQ ID NO:1 shows the cDNA sequence corresponding to the P129A RNA genome. The genome is 15,395 nucleotides in length (excluding the polyadenosine tail), begins with ATGACGTA, and ends with CCGCAATT. A typical polyadenosine tail of 55 residues is also provided in SEQ ID NO:1.

For the structural genes of P129A (ORFs 2 through 7), which comprise the 3' 20% of the genome, various PCR primers were chosen based on several partial cDNA sequences of other North American PRRS virus isolates available in the public DNA sequence database GenBank (for example PRU00153). Purified viral RNA was reverse transcribed into cDNA using reverse transcriptase and random hexamer primers. This cDNA was then used in PCR with gene-specific primers. PCR products were excised from gels and T/A cloned into plasmid pCR2.1 (Invitrogen). For each primer pair, multiple plasmids (from independent PCR reactions) were DNA sequenced. Sequences were assembled using the Seqman program from the Lasergene package (DNASTAR, Inc). This permitted completing the sequence of positions 11,992 through 15,347 of the P129A genome.

Also in the GenBank database are a series of short sequences (approximately 218 nucleotides total) which comprise a portion of the ORF 1b gene of several isolates of PRRS virus. One of these (PPSSEQB) was used to design PCR primers (forward 5'-ACAGTTTGGTGATCTATG-3' (SEQ ID NO:10), corresponding to positions 9063-9080; reverse 5'-CAGATTCAGATGTTCAA-3' (SEQ ID NO:11), corresponding to positions 9252-9268). These amplified a 206 nucleotide fragments, which includes 171 nucleotides of new sequence from the P129A ORF1b gene, corresponds to positions 9081 to 9251. A new forward primer was designed within this region (5'-ACCTCGTGCTGTATGCCGAATCTC-3' *Trade-mark

(SEQ ID NO:12), positions 9201-9224), and a matching primer was designed within ORF1b immediately upstream of ORF2 (5'-TCAGGCCTAAAGTTGGTTCAATGA-3' (SEQ ID NO:13), positions 12,027-12,050). These primers were used in RT-PCR to amplify a 2850 nucleotide fragment of ORF1b, corresponding to positions 9201-12,050 of the P129A genome.

5 During RT-PCR amplification of ORF5 of another North American field isolate of PRRS virus, a minor band was seen which was smaller than the expected size. This was sequenced and found to have limited homology with ORF1a of Lelystad virus (resulting from false priming). New primers within this region were chosen to amplify P129A (forward 5'-GATGACTGGGCTACTGACGAGGAT-3' (SEQ ID NO:14), corresponding to positions 1587-10 1610; reverse 5'-AGAGCGGCTGGGATGACACTG-3' (SEQ ID NO:15), corresponding to positions 1877-1897). In addition to the product of 311 nucleotides (266 nucleotides of new P129A sequence between the primers corresponding to positions 1611-1876), a larger minor PCR product of 701 nucleotides was cloned and sequenced (656 nucleotides of new P129A sequence between the primers corresponding to positions 1611-2264). The larger band 15 results from false priming of the reverse primer at positions 2265-2269.

The extreme 5' end of the genome of P129A was determined by 5' RACE (rapid amplification of cDNA ends) using a commercially available kit (Life Technologies Inc). Two nested reverse primers were chosen from within the known ORF1a sequence ("RACE2" 5'-CCGGGGAAGCCAGACGATTGAA-3' (SEQ ID NO:16), positions 1917-1938; and "RACE3" 20 5'-AGGGGGAGCAAAGAAGGGGTCATC-3' (SEQ ID NO:17), positions 1733-1756). RACE2 was used to prime cDNA synthesis, while RACE3 was used in PCR. The resulting PCR products were cloned and sequenced. The two longest products ended at precisely the same base (position 1 in SEQ ID NO:1).

The large gap between known sequence in ORF1a and ORF1b was bridged using 25 long RT-PCR. Two new primers were used (forward 5'-AGCACGCTCTGGTGCAACTG-3' (SEQ ID NO:18), positions 1361-1380; reverse 5'-GCCGCGGCGTAGTATTTCAG-3' (SEQ ID NO:19), positions 9420-9438). The resulting 8078 nucleotide RT-PCR product was cloned and sequenced.

The extreme 3' end of the genome of P129A was determined by ligating the 3' and 5' 30 ends of the viral RNA together and using RT-PCR to amplify the junction fragment. The resulting junction fragments were cloned and sequenced. Briefly, RNA extracted from pelleted virions was treated with tobacco acid pyrophosphatase (to remove 5' cap structures), then self-ligated with T4 RNA ligase (both from Epicentre Technologies). The primers used were 5'-CGCGTCACAGCATCACCTCAG-3' (SEQ ID NO:20) (forward, positions 15,218-35 15,239) and either 5'-CGGTAGGTTGGTTAACACATGAGTT-3' (SEQ ID NO:21) (reverse, positions 656-680) or 5'-TGGCTCTTCGGGCCTATAAAATA-3' (SEQ ID NO:22) (reverse,

Creation of an infectious full-length cDNA clone of P129A: A full-length infectious cDNA clone of P129A, designated pT7P129A, was assembled from four overlapping cloned RT-PCR products. The four RT-PCR products were first T/A cloned into plasmid pCR2.1 (Invitrogen) and transfected into *Escherichia coli* strain DH5-alpha. Bacterial colonies were screened, and those which contained inserts of the expected sizes in the "T7 to M13" orientation were chosen for sequencing and further manipulation. All four cloned RT-PCR products contained one or more non-silent mutations (deviations from the consensus nucleotide sequence for P129A of SEQ ID NO:1 which would result in a change in amino acid sequence of the encoded ORFs). These non-silent mutations (save one at position 12,622 in ORF 2) were repaired by subcloning fragments from other cloned RT-PCR products. The four repaired subgenomic clones were assembled into a full-length clone in a stepwise manner, using available restriction sites (see Figure 1). The 5' and 3' ends of the cDNA corresponding to the P129A genome in pT7P129A were modified by the addition of a T7 promoter and appropriate restriction endonuclease sites. The construction of pT7P129A is described in further detail in the following paragraphs:

The 5' end of the genome (positions 1-1756), generated by 5'-RACE and cloned into pCR2.1 as described above, was modified to include a T7 promoter immediately upstream of the cDNA corresponding to the P129A genome and a *PacI* site for future cloning. A 3-way ligation was performed using the 1216 bp *DsaI* - *BseRI* fragment of this plasmid (containing bases 27-1242 of P129A), the 4407 bp *BseRI* - *XbaI* fragment of the same plasmid (containing bases 1243-1756 of P129A and the entire plasmid vector up to the *XbaI* site), and the following synthetic double-stranded adapter (SEQ ID NOS:23 and 24):

35 The predicted transcript from the T7 promoter includes a single “G” residue from the promoter immediately upstream of the first “A” of the viral genome. A non-silent mutation at position 1230 (A to G) was repaired by replacing the 906 bp *AatII* - *SacII* fragment (bases

740-1645) with the same fragment from another clone. This plasmid was designated "pT7R3A-2".

The 8078 nucleotide PCR product described above was used to cover bases 1361-9438 of the P129A genome. A 58 bp deletion (positions 2535-2592) and 7 non-silent point mutations were corrected by subcloning fragments from other cloned RT-PCR products, yielding plasmid "pGAP10B-4". The 7837 bp *Bsu36I* - *SpeI* fragment from this plasmid was ligated to the 5482 bp *Bsu36I* - *SpeI* fragment from pT7R3A-2. The resulting plasmid "pT7RG" contains the first 9438 bases of the P129A genome behind the T7 promoter.

The 2850 nucleotide fragment of ORF1b described above (genome positions 9201-12,050) was corrected to repair non-silent mutations and designated "p1B3A-2". The 2682 bp *NdeI* - *SpeI* fragment of this plasmid was ligated to the 13,249 bp *NdeI* - *SpeI* fragment of pT7RG to yield plasmid "pT71A1B", which contains the first 12,050 bases of the P129A genome.

The fourth and final fragment of the P129A genome was derived by RT-PCR of ORFs 2 through 7, including the 3' non-translated region and a portion of the polyA tail. The forward primer was 5'-ACTCAGTCTAAGTGCTGGAAAGTTATG-3' (SEQ ID NO:25) (positions 11,504-11,530) and the reverse primer was 5'-GGGATTTAAATATGCATTTTTTTTTTTTTTTTTTTTAATTGCGGCCGCATGGTTCTCG-3' (SEQ ID NO:26). The reverse primer contains the last 22 bases of the P129A genome (positions 15,374-15,395), a polyA tail of 21 bases, an *NsiI* site (ATGCAT) and a *SwaI* site (ATTTAAAT). Non-silent point mutations and a single base deletion were repaired by subcloning fragments from other clones. An additional non-silent point mutation at position 12,622 (A to G) was inadvertently introduced at this stage. This results in a change from glutamine to arginine near the C-terminus of the ORF2 protein (amino acid residue 189 of the 256 amino acids in ORF2, which does not affect the overlapping ORF 3). This mutation had no apparent influence on viral growth, in cell culture or in pigs, and was not repaired. This mutation served as a genetic marker to distinguish virus derived from the cDNA clone from possible contamination with parental P129A or other PRRS viruses. The plasmid was designated "p2_7D-4". The structural genes of P129A were added to the rest of the genome by ligating the 3678 bp *Eco47III* - *SpeI* fragment of p2_7D-4 to the 15,635 bp *Eco47III* - *SpeI* fragment of pT71A1B.

This yields the final construct "pT7P129A", which comprises cDNA corresponding almost identically to the entire genome of P129A (however, with only a 21 base polyA tail, as opposed to 55 base polyA tail) behind a T7 promoter, cloned into the pCR2.1 vector between unique restriction enzyme sites (*PacI* and *SwaI*). The total length of pT7P129A is 19,313 bp, and it is stable in *E. coli* strain DH5-alpha. pT7P129A contains an A to G non-silent point

50204-11

-35-

mutation at position 12,622 that results in an arginine at position 189 of ORF2 rather than a glutamine (as is encoded by SEQ ID NO:1) and a silent C to T mutation at position 1559. Neither of these mutations affected viral growth under the conditions examined, both in cell culture and in pigs. For example, pT7P129A was used for *in vitro* transcription and the resulting RNA transcripts produced live North American PRRS virus when transfected into MARC-145 cells, thus demonstrating that this full-length clone is infectious.

In vitro transcription and transfection of RNA transcripts: In plasmid pT7P129A there are two T7 promoters in tandem upstream of the viral genome. One of these is positioned immediately upstream of the viral genome and was built into the PCR primer as described above. The other is present in the pCR2.1 cloning vector and is located outside of the multiple cloning site (initiating transcription 44 bases upstream of the viral genome). *PacI* was used to cut between these T7 promoters prior to *in vitro* transcription to generate a transcript that is closer to authentic viral RNA (a single extra G immediately upstream the viral genome, as opposed to 44 extra bases from the distal T7 promoter). In addition, pT7P129A was cut with *SwaI* prior to *in vitro* transcription. The resulting run-off transcripts include a 21 base long polyA tail and nine non-PRRS nucleotides, including an *NsiI* site (which was not used to linearize the plasmid, since the site also occurs once in the viral genome). The digested plasmid was purified by phenol extraction and ethanol precipitation prior to use.

A commercial kit (T7 Cap-Scribe^{*}, Boehringer Mannheim) was used for *in vitro* transcription. The DNA pellet from above, containing about 0.6 µg of *PacI*/*SwaI* digested pT7P129A, was resuspended in 20 µl of T7 Cap-Scribe buffer/T7 polymerase and incubated at 37°C for 30 minutes. A portion of the reaction was analyzed by agarose gel electrophoresis and shown to contain full-length RNA transcripts in addition to the expected DNA bands of 15,445 bp and 3868 bp. The *in vitro* transcription reaction was used fresh, immediately following incubation, without purification. Freshly confluent monolayers of MARC-145 cells were washed once in OptiMEM (without serum), and covered with 1 ml per 35mm well of OptiMEM (without serum) containing 500 µg/ml DEAE dextran (molecular weight approx. 500,000, Pharmacia Biotech). *In vitro* transcription reaction (15 µl) was added immediately. After 1 hour at 37°C, the transfection mixture was removed, monolayers were washed once with PBS and overlaid with 1.25% SeaPlaque agarose (FMC corporation) in OptiMEM with 2% fetal bovine serum and antibiotics. After 5 days at 37°C, a single plaque was visible. This virus was designated "rP129A-1" and was expanded on MARC-145 cells and characterized in cell culture and in pigs. Subsequent transfections of *in vitro* transcribed RNA from pT7P129A, using both DEAE dextran and electroporation, have yielded many additional plaques.

Characterization of recombinant virus rP129A-1: There are no apparent differences in growth kinetics, yield, or plaque morphology between cDNA-derived recombinant virus

*Trade-mark

rP129A-1 and its non-recombinant parent P129A. As discussed above, there are two differences in nucleotide sequence between the coding sequence of pT7P129A and the consensus sequence of P129A (shown in SEQ ID NO:1). Firstly, at position 1559 pT7P129A contains a T, whereas P129A contains a C (this is a silent mutation). Secondly, at position 12,622 pT7P129A contains a G, whereas P129A contains an A (this is the glutamine to arginine change in ORF2 described above). In order to rule out the possibility that rP129A-1 is actually a non-recombinant PRRS virus contaminant, RT-PCR and sequencing were performed on the regions surrounding these two differences. In the case of both genetic markers, rP129A-1 was identical to plasmid pT7P129A and different from parental virus P129A, thus confirming that rP129A-1 is derived from the infectious cDNA clone.

Characterization of recombinant virus rP129A-1 in pigs: The cDNA-derived virus rP129A-1 was compared to its non-recombinant parent P129A for its ability to infect and cause clinical disease in young pigs. Three groups of 10 pigs each from a PRRS-negative herd were infected at 4 weeks of age with either P129A, rP129A-1, or mock-infected with cell culture medium. Clinical signs, rectal temperatures, and body weights were monitored. Blood was collected on days 0, 2, 6, 10, and 13 post-infection for determination of serum viremia (by plaque assay on MARC-145 cells, Figure 2) and serum antibody (by ELISA using HerdChek PRRS from IDEXX, Figure 3). Gross and microscopic lesions of the lung were observed upon necropsy. There were no significant differences between the two virus-infected groups, indicating that rP129A-1 replicates in pigs and causes clinical disease which is quantitatively and qualitatively similar to its non-recombinant parent virus.

Example II. Deletion of ORF7 (Nucleocapsid Gene) from the North American PRRS Virus; Preparation of a Negatively-Marked, Replication-Defective Vaccine Thereby.

The viral nucleocapsid gene (ORF7) was partially deleted from an infectious cDNA clone of the PRRS virus of the present invention. The resulting recombinant modified PRRS virus would be expected to be replication-defective in pigs. This recombinant modified PRRS virus could be used as a vaccine to induce an immune response to the other PRRS virus proteins without the risks of clinical disease, spread to non-vaccinated animals, or reversion to virulence associated with attenuated live vaccines. In addition to being very safe, such a vaccine virus would also be "negatively marked", in the sense that it would allow exposure to field isolates of PRRS virus to be determined serologically, even in the presence of antibody to the vaccine virus. Antibodies to the ORF7 protein are commonly found in the sera of PRRS virus-infected pigs, whereas pigs vaccinated with an ORF7-deleted PRRSV would lack antibodies to the ORF7 protein.

Deletion of ORF7 from an infectious clone was accomplished as follows: Plasmid p2_7D-4 (see Figure 1) was used as template in PCR to amplify the 5' and 3' flanking regions upstream and downstream of ORF7. The upstream flank forward primer 5'-ATTAGATCTTGCCACCATGGTGGGGAAATGCTTGAC-3' (SEQ ID NO:27) (which binds to
 5 genome positions 13,776-13,791 near the beginning of ORF5 and contains additional restriction sites which are irrelevant to the current cloning) and the upstream flank reverse primer 5'-CTTTACGCGTTTGCTTAAGTTATTTGGCGTATTTGACAAGGTTTAC-3' (SEQ ID NO:28) (which binds to genome positions 14,857-14,902 at the junction of ORFs 6 and 7) amplified a fragment of 1147 bp. The reverse primer introduced *MluI* and *Afl* sites and a
 10 single base change at position 14,874, destroying the ATG start codon for ORF7 without altering the tyrosine encoded in the overlapping ORF6. For the downstream flank, the forward primer 5'-CAACACGCGTCAGCAAAAGAAAAAGAAGGGG-3' (SEQ ID NO:29) (positions 14,884-14,914 near the 5' end of ORF7, introduced an *MluI* site) and reverse primer 5'-GCGCGTTGGCCGATTCATTA-3' (SEQ ID NO:30) (downstream of the viral genome in the
 15 pCR2.1 plasmid) amplified a 462 bp fragment. A 3-way ligation was performed, using the 611 bp *Bst*EII-*MluI* fragment of the upstream flank PCR product, the 575 bp *MluI*-*SpeI* fragment of the downstream flank PCR product, and the 6653 bp *Bst*EII-*SpeI* fragment from plasmid p2_7D-4 (all fragments were gel purified following digestion). The resulting plasmid p2_7Ddelta7+7023 was deleted in the first seven amino acids of ORF7, and lacks a functional
 20 ATG start codon. Two new restriction sites which are absent in both the viral genome and the plasmid backbone, *Afl* and *MluI*, have been inserted to facilitate directional cloning of foreign genes into the space previously occupied by the 5' end of ORF7.

The changes made in p2_7Ddelta7+7023 were incorporated into a full-length genomic clone by ligating the 3683 bp *Eco*47III-*SpeI* fragment of p2_7Ddelta7+7023 with the 15,214 bp
 25 *Eco*47III-*SpeI* fragment of pCMV-S-p129. The resulting plasmid pCMV-S-p129delta7+7023 was used to transfect cells.

Since nucleocapsid is essential for viral growth, it is necessary to provide this protein in order to allow generation and replication of an ORF7-deficient PRRS virus. This can be accomplished using a helper virus or a complementing cell line, for example. ORF7-
 30 expressing MARC-145 cell lines could be created by stably transfecting cells with a plasmid containing both the ORF7 gene from P129A and the neomycin resistance gene. After selecting for neomycin resistance using the antibiotic G418, single-cell colonies could then be expanded and characterized. Clonal MARC-145-derived cell lines that are positive for ORF7 expression by both immunofluorescence and RT-PCR could be transfected with RNA from
 35 pT7P129delta7 in order to generate ORF7-deficient P129 virus.

Similar strategies can be used to generate PRRS viruses deficient in other structural genes (ORFs 2, 3, 4, 5, or 6), or deficient in all or portions of non-structural genes 1a and 1b. In addition, multiple deletions can be engineered into a single PRRS virus, and these can be grown in complementing cells which provide all necessary functions. Such gene-deficient PRRS viruses are likely to be either partially or completely attenuated in pigs, making them useful as vaccines against PRRS. They can also be used to distinguish vaccinated animals from animals infected with a wild-type PRRS virus as discussed above and/or as vectors for vaccinating animals with epitopes of other porcine pathogens (see Example III, below).

Example III. Insertion of Heterologous Genes into the North American PRRS Virus Genome; use of PRRS Virus as a Vector, and a Positively-Marked North American PRRS Virus.

In Example II, above, *Afl*III and *Mlu*I restriction enzyme sites were inserted into the region formerly occupied by the 5' end of ORF7. These sites are absent in the P129A genome and in the pCR2.1 and pCMV plasmids, and can be used in the directional cloning of foreign (heterologous) genes into the viral genome for expression. Potential leader-junction sites for transcription of the ORF7 subgenomic RNA at positions 14,744-14,749 (ATAACC) and 14,858-14,863 (TAAACC) are not affected by deletion of the ORF7 coding sequence, and can function in transcription of a foreign gene. Foreign (heterologous) genes can include genes from other PRRS virus isolates or genotypes, and/or genes from other non-PRRS pathogens, either pathogens that infect swine or pathogens that infect mammals other than swine, or avians.


In addition, these foreign genes (or portions thereof) can provide antigenic epitopes which are not normally found in swine. Such epitopes can be used to "positively mark" a vaccine, so that successful vaccination can be monitored serologically, even in the presence of antibody to field or conventional vaccine strains of PRRS virus. A positive marker needs not be a separate expression cassette. An antigenic epitope can be fused to a structural gene of the PRRS virus. For example, the upstream flank reverse primer described in Example I, above, can be extended in such a way as to add a carboxyl-terminal fusion of a non-PRRS virus antigenic epitope to the ORF6 membrane protein. The presence of antibody to this epitope in swine indicates successful vaccination.

Example IV. Cellular Expression of a PRRS Virus by Direct Transfection of cDNA into Cells.

The eukaryotic expression vector pCMV-MC1 (SEQ ID NO:31) was derived from the commercially available plasmid pCMVbeta (Clontech) by replacing the LacZ coding sequence

between two *Not* I sites with a linker containing *Not* I, *EcoR* V, *Avr* II, *Bgl* II, *Cla* I, *Kpn* I, *Pac* I, *Nhe* I, *Swa* I, *Sma* I, *Spe* I and *Not* I sites. Modification of the human CMV immediate early promoter was accomplished by substituting the sequence between the *Sac* I and the second *Not* I sites of pCMV-MC1 with a synthetic linker (shown below). The linker contains a half site for *Sac* I following by *Pac* I, *Spe* I and a half site for *Not* I. After annealing the two single stranded oligonucleotides, the linker was cloned into pCMV-MC1 between the *Sac* I and *Not* I sites, and a selected clone was designated pCMV-S1. The *Spe* I site of pCMV-S1 could not be cut, possibly due to a mistake in the oligo sequence. Therefore, the fragment between *Pac* I and *Hind* III in pCMV-S1 was replaced with *Pac* I (at position 877) - *Hind* III (at position 1162) fragment from pCMV-MC1. Thus, a *Spe* I site was regained. This final construct (pCMV-S) was used to clone the full length P129 genome.


Linker sequence (SEQ ID NOS:32 and 33):



20 The sequence immediately upstream of the 5' end of the P129 genome was modified to contain proper spacing and a convenient restriction enzyme site (*Pac* I). This was done by designing appropriate PCR primers (SEQ ID NOS:34 and 35) for amplification from pT7P129. After digestion with *Pac* I and *Aat* II, this PCR fragment was subcloned into the *Pac* I and *Aat* II sites of pT7RG (Fig. 1). The resulting plasmid was designated pT7RG-deltaT7.

25 The final construction was completed by subcloning the viral sequences from pT7RG-deltaT7 at the *Pac* I and *Nde* I sites into pT7P129, creating pT7P129-deltaT7. The full length P129 genome was digested from pT7P129-deltaT7 at *Pac* I and *Spe* I and transferred into pCMV-S at the *Pac* I and *Spe* I sites. This constructed was named pCMV-S-P129.

30 The sequence of the region of modification between the CMV promoter TATA box and the 5' end of the P129 sequence in pCMV-S-P129 is shown in SEQ ID NO:36 and schematically presented below:



To test the use of the CMV promoter to initiate PRRS virus infection in cells, pCMV-S-P129 plasmid DNA (0.5 or 1.0 µg) was transfected into MARC-145 cells by lipofection using Lipofectamine™ (Life Technologies Inc.). PRRS virus specific cytopathic effect was observed after transfection and the presence of PRRS virus antigen was determined by the immunofluorescent antibody test.

PRRS virus was generated efficiently from pCMV-S-P129, and the progeny virus could be passaged on MARC-145 cells. This demonstrates that a PRRS virus infection can be initiated directly from a plasmid cDNA encoding a PRRS virus, without an *in vitro* transcription step. Furthermore, pCMV-S-P129 generated a greater amount of progeny virus compared to plasmids wherein the 3' end of the pCMV promoter was not immediately in front of the start of the sequence encoding the North American PRRS virus.

Example V. Deletion of ORF4 from the North American PRRS Virus; Preparation of a Replication-Defective Vaccine Thereby.

A portion of the gene for ORF4, which encodes a membrane glycoprotein, was deleted from an infectious cDNA clone of the PRRS virus of the present invention. The resulting recombinant modified PRRS virus is expected to be replication-defective in pigs and to induce an immune response to the other PRRS virus proteins without the risks of clinical disease, spread to non-vaccinated animals, or reversion to virulence associated with attenuated live vaccines.

Deletion of ORF4 from an infectious clone was accomplished as follows. Plasmid p2_7D-4 (see Figure 1) was used as template in PCR to amplify the 5' and 3' flanking regions upstream and downstream of ORF4. The upstream flank forward primer was 5'-AGGTCGACGGCGGCAATTGGTTTCACCTAGAGTGGCTGCGTCCCTTCT-3' (SEQ ID NO:37). This primer binds to genome positions 13194-13241, near the beginning of ORF4, and introduces a mutation at position 13225 which destroys the ATG start codon of ORF4 without altering the overlapping amino acid sequence of ORF3. The upstream flank reverse primer was 5'-TCTTAAGCATTGGCTGTGATGGTGATATAC-3' (SEQ ID NO:38). This primer binds to genome positions 13455-13477 within the ORF4 coding region, downstream of ORF3, and introduces an *Afl*I site. For the downstream flanking region, the forward primer was 5'-CTTCTTAAGTCCACGCGTTTTCTTCTTGCCTTTTCTATGCTTCT-3' (SEQ ID NO:39). This primer binds to genome positions 13520-13545 in the middle of ORF4, and introduces *Afl*I and *Mlu*I sites for directional cloning of foreign genes. The reverse primer was 5'-TGCCCGGTCCCTTGCCTCT3' (SEQ ID NO:40). This primer binds to genome positions 14981-14999 in the ORF7 coding sequence. A three-way ligation was performed using the *Sal*I-*Afl*I fragment of the upstream flank PCR product, the *Afl*I-*Bst*EII fragment of the

downstream flank PCR product, and the *Sa*II-*Bst*EI fragment from plasmid p2_7D-4. All fragments were gel-purified following digestion. The resulting plasmid p2_7D-4delta4N has 42 bases of the central portion of ORF4 deleted and replaced with a 15 base artificial cloning site. The cloning site contains two restriction sites (*A*flI and *M*luI) that are absent from both
5 the viral genome and the plasmid backbone. These can be used to facilitate directional cloning of foreign genes into the space previously occupied by ORF4. The cloning site also contains a stop codon (TAA) that is in frame with ORF4 and further assures that functional ORF4 protein is not produced.

It was found that a more extensive deletion of the ORF4 coding sequence could be
10 made without interfering with expression of the downstream ORF5 envelope gene. In this case a shorter downstream flanking region was amplified by PCR using the same template and reverse primer, and using forward primer 5'-GTTTACGCGTCGCTCCTTGGTGGTTCG-3' (SEQ ID NO:41). This primer binds to genome positions 13654-13669 near the 3' end of ORF4, and contains an *A*flI site. Two-way ligation between the *A*flI-*Bst*EI fragment of the
15 downstream flank PCR product and the *A*flI-*Bst*EI fragment from plasmid p2_7D-4delta4N yielded the new plasmid p2_7D-4delta4NS. This plasmid has 176 bases of the ORF4 coding sequence deleted and replaced with the 15 base cloning site.

The changes made in p2_7D-4delta4N and p2_7Ddelta4NS were incorporated into the full-length genomic clone by replacing the *Bsr*GI-*Spe*I fragment from pCMV-S-P129 with
20 the modified *Bsr*GI-*Spe*I fragments from p2_7D-4delta4N and p2_7D-4delta4NS. The resulting plasmids pCMV-S-P129delta4N and pCMV-S-P129delta4NS were used to transfect cells.

In contrast to pCMV-S-P129, transfection of MARC-145 cells with plasmids pCMV-S-P129delta4N or pCMV-S-P129delta4NS did not result in viral plaques or fluorescent foci.
25 Individual transfected cells could be seen to be producing the ORF7 nucleocapsid protein, suggesting that the ORF4 gene product is not required for RNA replication or expression of viral genes, but is essential for release of infectious progeny virus. Since deletion of ORF4 is lethal to virus replication, it is necessary to provide this protein. This can be accomplished by using a complementing cell line. We created ORF4-expressing MARC-145 cell lines by stable
30 transfecting cells with a plasmid containing both ORF4 and the neomycin resistance gene. After selection for neomycin resistance using the antibiotic G418, single-cell colonies were expanded and characterized. After transfection with pCMV-S-P129delta4NS, three ORF4-expressing cell clones yielded live virus that could be propagated in these cells but not in MARC-145 cells. One of these, MARC400E9, was further characterized. Immunofluorescent
35 staining for viral nucleocapsid in MARC400E9 cells transfected with plasmid pCMV-S-P129delta4NS was positive.

Example VI. Use of Gene-Deleted PRRS Virus as a Vector for Expression of Foreign Genes.

In order to determine whether heterologous genes can be expressed from a gene-deleted PRRS virus, we inserted a copy of green fluorescent protein (GFP) into the site of ORF4 deletion in plasmid pCMV-S-P129delta4N. The GFP gene was amplified from a commercially available plasmid vector using PCR primers that introduced an *Afl*I site at the 5' end of the gene and an *Mlu*I site at the 3' end of the gene. The resulting plasmid pCMV-S-P129delta4N-GFP was used to transfect MARC-145 and MARC400E9 cells. As anticipated, MARC-145 cells did not support replication of the ORF4-deleted virus. Single green cells resulting from primary transfection were seen under the UV scope, indication that GFP was being expressed, but the virus did not spread to neighboring cells and no CPE was observed. In contrast, large green foci were observed in transfected MARC400E9 cells. Visible plaques formed and merged, destroying the monolayer. These results indicate that foreign genes can be expressed from ORF4-deleted PRRS virus, and that increasing the size of the viral genome by 692 bases (4.5%) does not interfere with packaging of the viral RNA into infectious particles.

Example VII. Use of Replication-Competent PRRS Virus as a Vector for Expression of Foreign Genes.

In order to determine whether heterologous genes can be expressed from a replication-competent PRRS virus, we inserted a copy of GFP into the region between ORF1b and ORF2. Since the leader/junction (L/J) sequence for ORF2 lies within ORF1b, this L/J sequence was used to drive expression of the GFP gene and a copy of the ORF6 L/J sequence was inserted downstream from GFP to drive expression of ORF2.

Plasmid p2_7D-4 (see Figure 1) was used as template in PCR to amplify the 5' and 3' flanking regions upstream and downstream of the insertion site. The upstream flank forward primer was 5'-AACAGAAGAGTTGTCTGGGTCCAC-3' (SEQ ID NO:42). This primer binds to genome positions 11699-11721 in ORF1b. The upstream flank reverse primer was 5'-GCTTTGACGCGTCCCCACTTAAGTTCAATTCAGGCCTAAAGTTGGTTCA-3' (SEQ ID NO:43). This primer binds to genome positions 12031-12055 in ORF1b and adds *Afl*II and *Mlu*I sites for directional cloning of foreign genes between ORF1b and ORF2. The downstream flank forward primer was 5'-GCGACGCGTGTTCCGTGGCAACCCCTTTAACCAGAGTTTCAGCGGAACAATGAAATGGG GTCTATACAAAGCCTCTTCGACA-3' (SEQ ID NO:44). This primer binds to genome positions 12056-12089 in ORF2 and contains an *Mlu*I site followed by the 40 bases that

64680-1480

43

precede the start of ORF6 (containing the ORF6 L/J sequence). The downstream flank reverse primer was 5'-AACAGAACGGCACGATACACCACAAA-3' (SEQ ID NO:45). This primer binds to genome positions 13819-13844 in ORF5. A three-way ligation was performed using the *Eco47III-MluI* fragment of the upstream flank PCR product, the *MluI-BsrGI* fragment from the downstream flank PCR product, and the *Eco47III-BsrGI* fragment from pCMV-S-P129. The resulting plasmid pCMV-S-P129-1bMCS2 contains the entire P129 genome with a cloning site and an additional L/J site between ORF1b and ORF2. The plasmid produces functionally normal virus when transfected into MARC-145 cells.

The GFP gene from a commercially available plasmid was PCR amplified using primers that add an *Afl* site to the 5' end and an *MluI* site to the 3' end of the gene. After digestion of the PCR fragment and pCMV-S-P129-1bMCS2 with *Afl* and *MluI*, the insert was ligated into the vector to yield plasmid pCMV-S-P129-1bGFP2. This plasmid produced green plaques when transfected into MARC-145 cells. The resulting virus could be passaged onto MARC-145 cells and continued to produce green plaques when observed under the UV scope. Thus, foreign genes may be expressed from replication-competent PRRS virus vectors. The P129-1bGFP2 virus contains a genome which is 774 bases (5%) longer than that of its P129 parent, yet it is packaged normally.

DEPOSIT OF BIOLOGICAL MATERIALS

The following biological material was deposited with the American Type Culture Collection (ATCC) at 10801 University Blvd., Manassas, Virginia, 20110-2209, USA, on November 19, 1998 and were assigned the following accession numbers:

<u>Plasmid</u>	<u>Accession No.</u>
plasmid pT7P129A	203488
plasmid pCMV-S-P129	203489

25

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

30

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: PFIZER PRODUCTS INC.
- (ii) TITLE OF INVENTION: INFECTIOUS CDNA CLONE OF NORTH AMERICAN
PORCINE REPRODUCTIVE AND RESPIRATORY
SYNDROME (PRRS) VIRUS AND USES THEREOF
- (iii) NUMBER OF SEQUENCES: 45
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: SMART & BIGGAR
 - (B) STREET: P.O. BOX 2999, STATION D
 - (C) CITY: OTTAWA
 - (D) STATE: ONT
 - (E) COUNTRY: CANADA
 - (F) ZIP: K1P 5Y6
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: ASCII (text)
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: CA 2,290,220
 - (B) FILING DATE: 23-NOV-1999
 - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 60/113,345
 - (B) FILING DATE: 22-DEC-1998
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: SMART & BIGGAR
 - (B) REGISTRATION NUMBER:

(C) REFERENCE/DOCKET NUMBER: 64680-1184

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (613)-232-2486

(B) TELEFAX: (613)-232-8440

(2) INFORMATION FOR SEQ ID NO.: 1:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 15450

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA

corresponding to North American Porcine

Reproductive And Respiratory Syndrome (PRRS) Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 1:

ATGACGTATA GGTGTTGGCT CTATGCCACG GCATTTGTAT TGTCAGGAGC TGTGACCATT 60
GGCACAGCCC AAAACTTGCT GCACGGAAAA CGCCCTTCTG TGACAGCCTT CTCAGGGGA 120
GCTTAGGGGT CTGTCCCTAG CACCTTGCTT CTGGAGTTGC ACTGCTTTAC GGTCTCTCCA 180
CCCCTTTAAC CATGTCTGGG ATACTTGATC GGTGCACGTG CACCCCAAT GCCAGGGTGT 240
TTATGGCGGA GGGCCAAGTC TACTGCACAC GATGTCTCAG TGCACGGTCT CTCCTTCCTC 300
TGAATCTCCA AGTTCCTGAG CTTGGGGTGC TGGGCCTATT TTATAGGCCC GAAGAGCCAC 360
TCCGGTGGAC GTTGCCACGT GCATTCCCCA CTGTCGAGTG CTCCCCCGCC GGGGCCTGCT 420
GGCTTTCTGC GATCTTTCCA ATTGCACGAA TGACCAGTGG AACCTGAAC TTTCAACAAA 480
GAATGGTGCG GGTGTCAGCT GAGATTTACA GAGCCGGCCA ACTCACCCCT GCAGTTTTGA 540
AGGCTCTACA AGTTTATGAA CGGGGTTGTC GCTGGTACCC CATTGTCGGA CCTGTCCCTG 600
GAGTGGCCGT TCACGCCAAC TCCCTACATG TGAGTGACAA ACCTTTCCCG GGAGCAACTC 660
ATGTGTTAAC CAACCTACCG CTCCCGCAGA GGCCAAGCC TGAAGACTTT TGCCCTTTTG 720

AGTGTGCTAT GGCTGACGTC TATGACATTA GCCATGACGC CGTCATGTAT GTGGCCAGAG 780
 GGAAAGTCTC CTGGGCCCCT CGTGGCGGGG ATGAAGTGAA ATTTGAAACC GTCCCCGAAG 840
 AGTTGAAGTT GATTGCGAAC CGACTCCACA TCTCCTTCCC GCCCCACCAC GCAGTGGACA 900
 TGTCTGAGTT TGCCTTCATA GCCCCTGGGA GTGGTGTCTC CTTGCGGGTC GAGCACCAAC 960
 ACGGTTGCCT TCCCGCTGAT ACTGTCCCTG AAGGGAACTG CTGGTGGTGC TTGTTTGA CT 1020
 TGCTCCCACC GGAAGTTCAG AATAAAGAAA TTCGCCGTGC TAACCAATTT GGCTATCAAA 1080
 CCAAGCATGG TGTCCCTGGC AAGTACCTAC AGCGGAGGCT GCAAGTTAAT GGTCTCCGAG 1140
 CAGTGACTGA TACAGATGGA CCTATTGTCTG TACAGTACTT CTCTGTTAGG GAGAGTTGGA 1200
 TCCGCCACTT CAGACTGGCG GAAGAACCTA GCCTCCCTGG GTTTGAAGAC CTCCTCAGAA 1260
 TAAGGGTAGA GCCTAATACG TCGCCATTGG GTGGCAAGGG TGAAAAAATC TTCCGGTTTG 1320
 GCAGTCACAA GTGGTACGGT GCTGGAAAGA GAGCAAGGAG AGCACGCTCT GGTGCAACTG 1380
 CCACGGTCGC TCACTGCGCT TTGCCCCTC GCGAAGCCCA GCAGGCCAAG AAGCTCGAGG 1440
 TTGCCAGCGC CAACAGGGCT GAGCATCTCA AGTACTATTC CCCGCCTGCC GACGGGAACT 1500
 GTGGTTGGCA CTGCATTTCC GCCATTACCA ACCGGATGGT GAATTCCAAA TTTGAAACCA 1560
 CTCTTCCCGA GAGAGTGAGA CCTTCAGATG ACTGGGCTAC TGACGAGGAT CTTGTGAATA 1620
 CCATCCAAAT CCTCAGGCTC CCCGCGGCCT TGGACAGGAA CGGTGCTTGT GCTGGCGCCA 1680
 AGTACGTGCT CAAGCTGGAA GGTGAGCACT GGACCGTCTC TGTGACCCCT GGGATGACCC 1740
 CTTCTTTGCT CCCCCTTGAA TGTGTTTCAGG GTTGTTGTGA GCATAAGAGC GGTCTTGGTT 1800
 TCCCAGACGT GGTCGAAGTT TCCGGATTTG ACCCTGCCTG TCTTGACCGA CTTGCTGAGA 1860
 TAATGCACTT ACCTAGCAGT GTCATCCCAG CTGCTCTGGC CGAGATGTCC GACGACTTCA 1920
 ATCGTCTGGC TTCCCCGGCC GCCACTGTGT GGACTGTTTC GCAATTCTTT GCCCGCCACA 1980
 GAGGAGGAGA GCATCCTGAC CAGGTGTGCT TAGGGAAAAT TATCAACCTT TGTCAGGTGA 2040
 TTGAGGAATG CTGCTGTTCC CGGAACAAAG CCAACCGGGC TACCCCGGAA GAGGTTGCGG 2100
 CAAAAGTTGA CCAGTACCTC CGTGGTGCAG CAAGCCTTGG AGAATGCTTG GCCAAGCTTG 2160
 AGAGGGCTCG CCCGCCGAGC GCGATGGACA CCTCCTTTGA TTGGAATGTT GTGCTTCCTG 2220
 GGGTTGAGAC GGCGGATCAG ACAACCAAAC AGCTCCATGT CAACCAGTGC CGCGCTCTGG 2280
 TTCCTGTCGT GACTCAAGAG CCTTTGGACA GAGACTCGGT CCCTCTGACC GCCTTCTCGC 2340
 TGTCCAATTG CTACTACCCT GCACAAGGTG ACGAGGTCCG TCACCGTGAG AGGCTAAACT 2400
 CCGTGCTCTC TAAGTTGGAG GGGGTTGTTC GTGAGGAATA TGGGCTCACG CCAACTGGAC 2460

CTGGCCCGCG ACCCGCACTG CCGAACGGGC TCGACGAGCT TAAAGACCAG ATGGAGGAGG 2520
ATCTGCTGAA ATTAGTCAAC GCCCAGGCAA CTTCAGAAAT GATGGCCTGG GCAGCCGAGC 2580
AGGTTGATCT AAAAGCTTGG GTCAAAAATT ACCCACGGTG GACACCGCCA CCCCCTCCAC 2640
CAAGAGTTCA GCCTCGAAAA ACGAAGTCTG TCAAGAGCTT GCTAGAGAAC AAGCCTGTCC 2700
CTGCTCCGCG CAGGAAGGTC AGATCTGATT ATGGCAGCCC GATTTTGATG GGCGACAATG 2760
TTCCTAACGG TTGGGAAGAT TCGACTGTTG GTGGTCCCCT TGACCTTTCG GCACCATCCG 2820
AGCCGATGAC ACCTCTGAGT GAGCCTGTAC TTATTTCCAG GCCAGTGACA TCTTTGAGTG 2880
TGCCGGCCCC AGTTCCTGCA CCGCGTAGAG CTGTGTCTCG ACCGATGACG CCCTCGAGTG 2940
AGCCAATTTT TGTGTCTGCA CTGCGACACA AATTTGAGCA GGTGGAAAAA GCAAATCTGG 3000
CGGCAGCAGC GCCGATGTAC CAGGACGAAC CCTTAGATTT GTCTGCATCC TCACAGACTG 3060
AATATGGGGC TTCTCCCCTA ACACCACCGC AGAACGTGGG CATTCTGGAG GTAAGGGGGC 3120
AAGAAGCTGA GGAAGTTCTG AGTGAAATCT CGGATATTCT GAATGATACC AACCTGCAC 3180
CTGTGTCATC AAGCAGCTCC CTGTCAAGTG TTAGGATCAC ACGCCCAAAA TACTCAGCTC 3240
AAGCCATTAT CGACTTGGGC GGGCCCTGCA GTGGGCACCT CCAAAGGGAA AAAGAAGCAT 3300
GCCTCCGCAT CATGCGTGAG GCTTGTGATG CGGCCAAGCT TAGTGACCCT GCCACGCAGG 3360
AATGGCTTTC TCGCATGTGG GATAGGGTGG ACATGCTGAC TTGGCGCAAC ACGTCTGCTT 3420
ACCAGGCGTT TCGCACCTTA GATGGCAGGT TTGGGTTTCT CCCAAAGATG ATACTCGAGA 3480
CGCCGCCGCC CTACCCGTGT GGGTTTGTGA TGTTGCCTCA CACCCCTGCA CCTTCCGTGA 3540
GTGCAGAGAG CGACCTTACC ATCGGTTTCTG TCGCCACTGA AGATATTCCA CGCATCCTCG 3600
GGAAAATAGA AAATACCGGT GAGATGATCA ACCAGGGACC CTTGGCATCC TCTGAGGAAG 3660
AACCGGTATA CAACCAACCT GCCAAAGACT CCCGGATATC GTCGCGGGGG TCTGACGAGA 3720
GCACAGCAGC TCCGTCCGCA GGTACAGGTG GCGCCGGCTT ATTTACTGAT TTGCCACCTT 3780
CAGACGGCGT AGATGCGGAC GGTGGGGGGC CGTTGCAGAC GGTAAGAAAG AAAGCTGAAA 3840
GGCTCTTCGA CCAATTGAGC CGTCAGGTTT TTAACCTCGT CTCCCATCTC CCTGTTTTCT 3900
TCTCACACCT CTTCAAATCT GACAGTGGTT ATTCTCCGGG TGATTGGGGT TTTGCAGCTT 3960
TACTCTATT TTGCCTCTTT TTGTGTTACA GCTACCCATT CTTGGGTTTC GTTCCCCTCT 4020
TGGGTGTATT TTCTGGGTCT TCTCGGCGTG TGCGCATGGG GGTTTTTGGC TGCTGGCTGG 4080
CTTTTGCTGT TGGCCTGTTC AAGCCTGTGT CCGACCCAGT CGGCACTGCT TGTGAGTTTG 4140
ACTCGCCAGA GTGTAGGAAC GTCCTTCATT CTTTGTAGCT TCTCAAACCT TGGGACCCTG 4200

TTCGCAGCCT TGTGTGGGC CCCGTCGGTC TCGGTCTTGC CATTCTTGGC AGGTTACTGG 4260
 GCGGGGCACG CTACATCTGG CATTTTTTGC TTAGGCTTGG CATTGTTGCA GATTGTATCT 4320
 TGGCTGGAGC TTATGTGCTT TCTCAAGGTA GGTGTAAAAA GTGCTGGGGA TCTTGTATAA 4380
 GAACTGCTCC TAATGAAATC GCCTTCAACG TGTTCCTTTT TACACGTGCG ACCAGGTCGT 4440
 CACTCATCGA CCTGTGCGAT CGGTTTTGTG CGCCAAAAGG CATGGACCCC ATTTTCCTCG 4500
 CCACTGGGTG GCGTGGGTGC TGGACCGGCC GAAGTCCCAT TGAGCAACCC TCTGAAAAAC 4560
 CCATCGCGTT CGCCCAGTTG GATGAAAAGA GGATTACGGC TAGAACTGTG GTCGCTCAGC 4620
 CTTATGATCC TAATCAAGCC GTAAAGTGCT TGCGGGTGTT ACAGGCGGGT GGGGCGATGG 4680
 TGGCCGAGGC AGTCCCAAAA GTGGTCAAAG TTTCTGCTAT TCCATTCCGA GCCCCCTTTT 4740
 TTCCCACCGG AGTGAAAGTT GATCCCGAGT GCAGGATCGT GGTCGACCCC GATACTTTTA 4800
 CTACAGCCCT CCGGTCTGGT TACTCTACCA CAAACCTCGT CCTTGGTGTG GGGGACTTTG 4860
 CCCAGCTGAA TGGACTAAAG ATCAGGCAAA TTTCCAAGCC TTCGGGAGGA GGCCACACC 4920
 TCATTGCTGC CCTGCATGTT GCCTGCTCGA TGGCGTTGCA CATGCTTGCT GGGGTTTATG 4980
 TAACTTCAGT GGGGTCTTGC GGTGCCGGCA CCAACGATCC ATGGTGCACT AATCCGTTTG 5040
 CCGTTCCTGG CTACGGACCA GGCTCTCTCT GCACGTCCAG ATTGTGCATC TCCCAACATG 5100
 GCCTTACCCT GCCCTTGACA GCACTTGTGG CGGGATTCGG TCTTCAGGAA ATCGCCTTGG 5160
 TCGTTTTGAT TTTCGTTTCC ATCGGAGGCA TGGCTCATAG GTTGAGTTGT AAGGCTGATA 5220
 TGCTGTGCAT CTTACTTGCA ATCGCCAGCT ATGTTTGGGT ACCCCTTACC TGGTTGCTTT 5280
 GTGTGTTTCC TTGTTGGTTG CGCTGGTTCT CTTTGCACCC CCTCACCATC CTATGGTTGG 5340
 TGTTTTTCTT GATTTCTGTA AATATGCCTT CGGGAATCTT GGCCGTGGTG TTATTGGTTT 5400
 CTCTTTGGCT TTTGGGACGT TATACTAACA TTGCTGGTCT TGTCACCCCC TATGATATTC 5460
 ATCATTACAC CAGTGGCCCC CGCGGTGTTG CCGCCTTAGC TACCGCACCA GATGGAACCT 5520
 ACTTGGCTGC CGTCCGCCGC GCTGCGTTGA CTGGTCGCAC CATGCTGTTC ACCCCGTCTC 5580
 AGCTTGGGTC CCTTCTTGAG GGCGCTTTCA GAACTCGAAA GCCCTCACTG AACACCGTCA 5640
 ATGTGGTTGG GTCCTCCATG GGCTCTGGTG GAGTGTTTAC CATCGACGGG AAAATTAGGT 5700
 GCGTGA CTGC CGCACATGTC CTTACGGGTA ATTCGGCTAG GGTTTCCGGA GTCGGCTTCA 5760
 ATCAAATGCT TGACTTTGAT GTGAAAGGGG ACTTCGCCAT AGCTGATTGC CCGAATTGGC 5820
 AAGGAGCTGC TCCCAAGACC CAATTCTGCG AGGATGGATG GGCTGGCCGT GCCTATTGGC 5880
 TGACATCCTC TGGCGTCGAA CCCGGTGTTA TTGGGAATGG ATTCGCCTTC TGCTTCACCG 5940

CGTGCGGCGA TTCCGGGTCC CCAGTGATCA CCGAAGCTGG TGAGCTTGTC GGCGTTCACA 6000
 CAGGATCAAA TAAACAAGGA GGTGGCATCG TCACGCGCCC TTCAGGCCAG TTTTGTAACG 6060
 TGGCACCCAT CAAGCTGAGC GAATTAAGTG AATTCTTTGC TGGACCCAAG GTCCCGCTCG 6120
 GTGATGTGAA GGTGGCAGC CACATAATTA AAGATACGTG CGAAGTACCT TCAGATCTTT 6180
 GCGCCTTGCT TGCTGCCAAA CCTGAACTGG AGGGAGGCCT CTCCACCGTC CAACTTCTGT 6240
 GTGTGTTTTT CCTACTGTGG AGAATGATGG GACATGCCTG GACGCCCTTG GTTGCTGTGG 6300
 GGTTTTTCAT TCTGAATGAG GTTCTCCCAG CTGTCCTGGT TCGGAGTGTT TTCTCCTTTG 6360
 GGATGTTTGT GCTATCTTGG CTCACACCAT GGTCTGCGCA AGTTCTGATG ATCAGGCTTC 6420
 TAACAGCAGC TCTTAACAGG AACAGATGGT CACTTGCCTT TTACAGCCTT GGTGCGGTGA 6480
 CCGGTTTTGT CGCAGATCTT GCGGCAACTC AAGGGCACCC GTTGCAGGCA GTAATGAATT 6540
 TGAGCACCTA TGCCTTCCTG CCTCGGATGA TGGTTGTGAC CTCACCAGTC CCAGTGATTG 6600
 CGTGTGGTGT TGTGCACCTA CTTGCCATCA TTTTGTA CTT GTTCAAGTAC CGCGGCCTGC 6660
 ACAATGTTCT TGTGTTGAT GGAGCGTTTT CTGCAGCTTT CTTCTTGCGA TACTTTGCCG 6720
 AGGGAAAGTT GAGGGAAGGG GTGTCGCAAT CCTGCGGAAT GAATCATGAG TCATTAAGT 6780
 GTGCCCTCGC TATGAGACTC AATGACGAGG ACTTGGA CTT CTTTACGAAA TGGACTGATT 6840
 TTAAGTGCTT TGTTTCTGCG TCCAACATGA GGAATGCAGC AGGCCAATTC ATCGAGGCTG 6900
 CCTATGCAAA AGCACTTAGA ATTGAACTTG CCCAGTTGGT GCAGGTTGAT AAGGTTGAG 6960
 GTACTTTGGC CAAGCTTGAG GCTTTTGCTG ATACCGTGGC ACCCCAACCTC TCGCCCGGTG 7020
 ACATTGTTGT TGCTCTTGGC CACACGCTG TTGGCAGCAT CTTGACCTA AAGGTTGGTG 7080
 GTACCAAGCA TACTCTCAA GTCATTGAGA CCAGAGTCCT TGCCGGGTCC AAAATGACCG 7140
 TGGCGCGCGT CGTTGACCCA ACCCCCACGC CCCCACCGC ACCCGTGCCC ATCCCCCTCC 7200
 CACCGAAAGT TCTAGAGAAT GGTCCCAACG CCTGGGGGGA TGGGGACCGT TTGAATAAGA 7260
 AGAAGAGGCG TAGGATGGAA ACCGTCGGCA TCTTTGTCAT GGGTGGGAAG AAGTACCAGA 7320
 AATTTTGGGA CAAGAATTCC GGTGATGTGT TTTACGAGGA GGTCCATGAC AACACAGATG 7380
 CGTGGGAGTG CCTCAGAGTT GGTGACCCTG CCGACTTTGA CCCTGAGAAG GGAAGTCTGT 7440
 GTGGGCATAC TACTATTGAA GATAAGGATT ACAAAGTCTA CGCCTCCCCA TCTGGCAAGA 7500
 AGTTCCTGGT CCCCCTCAAC TCAGAGAGCG GAAGAGCCCA ATGGGAAGCT GCAAAGCTTT 7560
 CCGTGGAGCA GGCCCTTGGC ATGATGAATG TCGACGGTGA ACTGACGGCC AAAGAAGTGG 7620
 AGAAACTGAA AAGAATAATT GACAACTTC AGGGCCTGAC TAAGGAGCAG TGTTTAAACT 7680

GCTAGCCGCC AGCGGCTTGA CCCGCTGTGG TCGCGGCGGC TTGGTTGTTA CTGAGACAGC 7740
GGTAAAAATA GTCAAATTTT ACAACCGGAC TTTCACCCTA GGGCCTGTGA ATTTAAAAGT 7800
GGCCAGTGAG GTTGAGCTGA AAGACGCGGT CGAGCACAAC CAACACCCGG TTGCAAGACC 7860
GGTTGACGGT GGTGTTGTGC TCCTGCGTTC CGCAGTTCCT TCGCTTATAG ATGTCCTGAT 7920
CTCCGGTGCT GACGCATCTC CTAAGTTACT CGCTCGTCAC GGGCCGGGGA AACTGGGAT 7980
CGATGGCACG CTTTGGGACT TTGAGGCCGA GGCCACCAAA GAGGAAATTG CACTCAGTGC 8040
GCAAATAATA CAGGCTTGTG ACATTAGGCG CGGCGACGCA CCTGAAATTG GTCTCCCTTA 8100
CAAGCTGTAC CCTGTTAGGG GCAACCCTGA GCGGGTAAAA GGAGTTTTAC AGAATAACAAG 8160
GTTTGGAGAC ATACCTTACA AAACCCCCAG TGACACTGGA AGCCCAGTGC ACGCGGCTGC 8220
CTGCCTCACG CCCAATGCCA CTCCGGTGAC TGATGGGCGC TCTGTCTTGG CTACTACCAT 8280
GCCCTCCGGT TTTGAATTGT ATGTACCGAC CATTCCAGCG TCTGTCCTTG ATTATCTTGA 8340
CTCTAGGCCT GACTGCCCCA AACAGTTGAC AGAGCACGGC TGTGAGGATG CCGCATTGAG 8400
AGACCTCTCC AAGTATGACT TGTCCACCCA AGGCTTTGTT TTACCTGGGG TTCTTCGCCT 8460
TGTGCGTAAG TACCTGTTTG CCCATGTGGG TAAGTGCCCG CCCGTTTCATC GGCCTTCCAC 8520
TTACCCTGCC AAGAATTCTA TGGCTGGAAT AAATGGGAAC AGGTTTCCAA CCAAGGACAT 8580
TCAGAGCGTC CCTGAAATCG ACGTTCTGTG CGCACAGGCC GTGCGAGAAA ACTGGCAAAC 8640
TGTTACCCCT TGTACCCTCA AGAAACAGTA TTGTGGGAAG AAGAAGACTA GGACAATACT 8700
CGGCACCAAT AATTTTCATTG CGTTGGCCCA CCGGGCAGCG TTGAGTGGTG TCACCCAGGG 8760
CTTCATGAAA AAGGCGTTTA ACTCGCCCAT CGCCCTCGGG AAAAACAAAT TTAAGGAGCT 8820
ACAGACTCCG GTCTTAGGCA GGTGCCTTGA AGCTGATCTT GCATCCTGTG ATCGATCCAC 8880
ACCTGCAATT GTCCGCTGGT TTGCCGCCAA TCTTCTTTAT GAACTTGCCT GTGCTGAAGA 8940
GCACCTACCG TCGTACGTGC TGAAGTGTG CCATGACCTA TTGGTCACGC AGTCCGGCGC 9000
AGTGACTAAG AGGGGTGGCC TATCGTCTGG CGACCCGATC ACTTCTGTGT CTAACACCAT 9060
TTACAGCTTG GTGATATATG CACAGCACAT GGTGCTTAGT TACTTTAAAA GTGGTCACCC 9120
TCATGGCCTT CTGTTCTTAC AAGACCAGCT GAAGTTCGAG GACATGCTCA AAGTCCAACC 9180
CCTGATCGTC TATTCGGACG ACCTCGTGCT GTATGCCGAA TCTCCCACCA TGCCGAACCTA 9240
CCACTGGTGG GTCGAACATC TGAATTTGAT GCTGGGTTTT CAGACGGACC CAAAGAAGAC 9300
AGCCATAACG GACTCGCCAT CATTTCTAGG CTGTAGGATA ATAAATGGAC GCCAGCTAGT 9360
CCCCAACCGT GACAGGATCC TCGCGGCCCT CGCTTACCAT ATGAAGGCAA GCAATGTTTC 9420

TGAATACTAC GCCGCGGCGG CTGCAATACT CATGGACAGC TGTGCTTGTT TAGAGTATGA 9480
TCCTGAATGG TTTGAAGAGC TTGTGGTTGG GATAGCGCAG TGCGCCCGCA AGGACGGCTA 9540
CAGCTTTCCC GGCCCGCCGT TCTTCTTGTC CATGTGGGAA AACTCAGAT CCAATCATGA 9600
GGGAAGAAG TCCAGAATGT GCGGGTATTG CGGGGCCCCG GCTCCGTACG CCACTGCCTG 9660
TGGCCTCGAC GTCTGTATTT ACCACACCCA CTTCCACCAG CATTGTCCAG TCATAATCTG 9720
GTGTGGCCAC CCGGCTGGTT CTGGTTCTTG TAGTGAGTGC AAACCCCCC TAGGGAAAGG 9780
CACAAGCCCT CTAGATGAGG TGTTAGAACA AGTCCCGTAT AAGCCTCCAC GGACTGTAAT 9840
CATGCATGTG GAGCAGGGTC TCACCCCTCT TGACCCAGGC AGATACCAGA CTCGCCGCGG 9900
ATTAGTCTCC GTTAGGCGTG GCATCAGAGG AAACGAAGTT GACCTACCAG ACGGTGATTA 9960
TGCTAGCACC GCCCTACTCC CCACTTGTAAG AGAGATCAAC ATGGTCGCTG TCGCCTCTAA 10020
TGTGTTGCGC AGCAGGTTCA TCATCGGTCC GCCCGGTGCT GGGAAAACAT ACTGGCTCCT 10080
TCAGCAGGTC CAGGATGGTG ATGTCATTTA CACACCGACT CACCAGACCA TGCTCGACAT 10140
GATTAGGGCT TTGGGGACGT GCCGGTTCAA CGTCCCAGCA GGTACAACGC TGCAATTCCC 10200
TGCCCCCTCC CGTACCGGCC CGTGGGTTCG CATCCTGGCC GCGGGTTGGT GTCCTGGTAA 10260
GAATTCCTTC CTGGATGAAG CAGCGTATTG TAATCACCTT GATGTCTTGA GGCTCCTTAG 10320
CAAAACCACC CTTACCTGTC TGGGAGACTT CAAACAACCTC CACCCAGTGG GTTTTGATTC 10380
TCATTGCTAT GTTTTGTACA TCATGCCTCA GACCCAGTTG AAGACCATCT GGAGATTCGG 10440
ACAGAACATC TGTGATGCCA TCCAACCAGA TTACAGGGAC AAACCTGTGT CCATGGTCAA 10500
CACAACCCGT GTAACCTACA TGGAAAAACC TGTCAAGTAT GGGCAAGTCC TCACCCCTTA 10560
CCACAGGGAC CGAGAGGACG GCGCCATCAC AATTGACTCC AGTCAAGGCG CCACATTTGA 10620
TGTGGTTACA CTGCATTTGC CCACTAAAGA TTCACTCAAC AGGCAAAGAG CCCTTGTTGC 10680
TATCACCAGG GCAAGACATG CTATCTTTGT GTATGACCCA CACAGGCAAT TGCAGAGCAT 10740
GTTTGATCTT CCTGCGAAGG GCACACCCGT CAACCTCGCA GTGCACCGTG ATGAGCAGCT 10800
GATCGTACTG GATAGAAATA ATAAAGAATG CACAGTTGCT CAGGCTATAG GCAACGGAGA 10860
TAAATTCAGG GCCACCGACA AGCGCGTTGT AGATTCTCTC CGCGCCATTT GTGCTGATCT 10920
GGAAGGGTCG AGCTCCCCGC TCCCAAGGT CGCACACAAC TTGGGATTTT ATTTCTCACC 10980
TGATTTGACA CAGTTTGCTA AACTCCCGGT AGACCTTGCA CCCCCTGGC CCGTGGTGAC 11040
AACCAGAAC AATGAAAAGT GGCCGGATCG GCTGGTTGCC AGCCTTCGCC CTGTCCATAA 11100
GTATAGCCGT GCGTGCATTG GTGCCGGCTA TATGGTGGGC CCCTCGGTGT TTCTAGGCAC 11160

CCCTGGGGTC GTGTCATACT ACCTCACAAA ATTTGTCAAG GGCGAGGCTC AAGTGCTTCC 11220
GGAGACAGTC TTCAGCACCG GCCGAATTGA GGTGGATTGC CGGGAGTATC TTGATGACAG 11280
GGAGCGAGAA GTTGCTGAGT CCCTCCCACA TGCCTTCATT GGCGACGTCA AAGGCACCAC 11340
CGTTGGGGGA TGTCATCATG TCACCTCCAA ATACCTTCCG CGCTTCCTTC CCAAGGAATC 11400
AGTCGCGGTA GTCGGGGTTT CGAGCCCCGG GAAAGCCGCA AAAGCAGTGT GCACATTGAC 11460
GGATGTGTAC CTCCCAGACC TTGAGGCCTA CCTCCACCCA GAGACTCAGT CTAAGTGCTG 11520
GAAAGTTATG TTGGACTTCA AGGAAGTTCG ACTGATGGTC TGGAAAGACA AGACGGCCTA 11580
TTTCCAACCTT GAAGGCCGCT ATTTACCTG GTATCAGCTT GCAAGCTACG CCTCGTACAT 11640
CCGTGTTCTT GTCAACTCCA CGGTGTATCT GGACCCCTGC ATGGGCCCTG CCCTTTGCAA 11700
CAGAAGAGTT GTCGGGTCCA CCCATTGGGG AGCTGACCTC GCAGTCACCC CTTATGATTA 11760
CGGTGCTAAA ATCATCTTGT CTAGCGCTTA CCATGGTGAA ATGCCTCCTG GATACAAGAT 11820
TCTGGCGTGC GCGGAGTTCT CGCTCGACGA CCCAGTCAAG TACAAACACA CCTGGGGTTT 11880
TGAATCGGAT ACAGCGTATC TGTATGAGTT CACCGGAAAC GGTGAGGACT GGGAGGATTA 11940
CAATGATGCG TTTCGTGCGC GCCAGAAAGG GAAAATTTAT AAGGCCACTG CTACCAGCAT 12000
GAAGTTTTAT TTTCCCCCGG GCCCCGTCAT TGAACCAACT TTAGGCCTGA ATTGAAATGA 12060
AATGGGGTCT ATACAAAGCC TCTTCGACAA AATTGGCCAG CTTTTTGTGG ATGCTTTCAC 12120
GGAATTTTTG GTGTCCATTG TTGATATCAT CATATTTTTG GCCATTTTGT TTGGCTTCAC 12180
CATCGCCGGT TGGCTGGTGG TCTTTTGCAT CAGATTGGTT TGCTCCGCGG TATTCCGTGC 12240
GCGCCCTGCC ATTCACCCTG AGCAATTACA GAAGATCCTA TGAGGCCTTT CTTTCTCAGT 12300
GCCGGGTGGA CATTCCCACC TGGGGGGTAA AACACCCTTT GGGGATGTTT TGGCACCATA 12360
AGGTGTCAAC CCTGATTGAT GAAATGGTGT CGCGTCGAAT GTACCGCATC ATGGAAAAAG 12420
CAGGGCAAGC TGCCTGGAAA CAGGTGGTGA GCGAGGCTAC GCTGTCTCGC ATTAGTAGTT 12480
TGGATGTGGT GGCTCATTTT CAACATCTTG CCGCCATTGA AGCCGAGACC TGTAATATT 12540
TGGCTTCTCG ACTGCCCATG CTACACAACC TGCGCATGAC AGGGTCAAAT GTAACCATAG 12600
TGTATAATAG CACTTTAAAT CAGGTGTTTG CTATTTTTCC AACCCTGGT TCCCGGCCAA 12660
AGCTTCATGA TTTTCAGCAA TGGCTAATAG CTGTACATTC CTCCATATTT TCCTCTGTTG 12720
CAGCTTCTTG TACTCTTTTT GTTGTGCTGT GGTGCGGGT TCCAATGCTA CGTACTGTTT 12780
TTGGTTTCCG CTGGTTAGGG GCAATTTTTC TTTCGAACTC ATGGTGAATT ACACGGTGTG 12840
TCCACCTTGC CTCACCCGAC AAGCAGCCGC TGAGGTCCTT GAACCCGGTA GGTCTCTTTG 12900

GTGCAGGATA GGGCATGACC GATGTGGGGA GGACGATCAC GACGAACTAG GGTTTCATGGT 12960
TCCGCCTGGC CTCTCCAGCG AAAGCCACTT GACCAGTGTT TACGCCTGGT TGGCGTTCCT 13020
GTCCTTCAGC TACACGGCCC AGTTCCATCC CGAGATATTT GGGATAGGGA ACGTGAGTGA 13080
AGTTTATGTT GACATCAAGC ACCAATTCAT CTGCGCCGTT CATGACGGGC AGAACACCAC 13140
CTTGCCTCGC CATGACAATA TTTCAGCCGT ATTTAGACC TACTATCAAC ATCAGGTCGA 13200
CGGCGGCAAT TGGTTTCACC TAGAATGGCT GCGTCCCTTC TTTTCCTCTT GGTGTTTTT 13260
AAATGTTTCG TGGTTTCTCA GGCGTTCGCC TGCAAGCCAT GTTTCAGTTC GAGTCTTTCA 13320
GACATCAAAA CCAACACTAC CGCAGCATCA GGCTTTGTTG TCCTCCAGGA CATCAGCTGC 13380
CTTAGGCATG GCGACTCGTC CTTTCCGACG ATTCGCAAAA GCTCTCAATG CCGCACGGCG 13440
ATAGGGACAC CCGTGTATAT CACCATCACA GCCAATGTGA CAGATGAGAA TTAATTACAT 13500
TCTTCTGATC TCCTCATGCT TTCTTCTTGC CTTTTCTATG CTTCTGAGAT GAGTGAAAAG 13560
GGATTCAAGG TGGTGTTTGG CAATGTGTCA GGCATCGTGG CTGTGTGTGT CAACTTTACC 13620
AGCTACGTCC AACATGTCAA AGAGTTTACC CAACGCTCCT TGGTGGTCGA TCATGTGCGG 13680
CTGCTTCATT TCATGACACC TGAGACCATG AGGTGGGCAA CCGTTTTAGC CTGTCTTTTT 13740
GCCATCCTAC TGGCAATTTG AATGTTCAAG TATGTTGGGG AAATGCTTGA CCGCGGGCTG 13800
TTGCTCGCGA TTGCTTTCTT TGTGGTGTAT CGTGCCGTTT TGTTTTGCTG TGCTCGGCAG 13860
CGCCAACAGC AGCAGCAGCT CTCATTTTCA GTTGATTTAT AACTTGACGC TATGTGAGCT 13920
GAATGGCACA GATTGGCTGG CAGAAAAATT TGATTGGGCA GTGGAGACTT TTGTCATCTT 13980
TCCCGTGTTG ACTCACATTG TTTCCTATGG TGCACTCACC ACCAGCCATT TCCTTGACAC 14040
AGTTGGTCTG GTTACTGTGT CCACCGCCGG GTTTTATCAC GGGCGGTATG TCTTGAGTAG 14100
CATCTACGCG GTCTGTGCTC TGGCTGCGTT GATTTGCTTC GTTATTAGGC TTGCGAAGAA 14160
CTGCATGTCC TGGCGCTACT CTTGTACCAG ATATACCAAC TTCCTTCTGG AACTAAGGG 14220
CAGACTCTAT CGTTGGCGGT CGCCCGTTAT CATAGAAAAA GGGGGTAAGG TTGAGGTCGA 14280
AGGTCACCTG ATCGACCTCA AAAGAGTTGT GCTTGATGGT TCCGTGGCAA CCCCTTTAAC 14340
CAGAGTTTCA GCGGAACAAT GGGGTCGTCT CTAGACGACT TTTGCCATGA TAGCACGGCT 14400
CCACAAAAGG TGCTTTTGGC GTTTTCCATT ACCTACACGC CAGTAATGAT ATATGCTCTA 14460
AAGGTAAGTC GCGGCCGACT ACTAGGGCTT CTGCACCTTT TGATCTTTCT GAATTGTGCT 14520
TTTACCTTCG GGTACATGAC ATTCGAGCAC TTTCAGAGCA CAAATAGGGT CGCGCTCACT 14580
ATGGGAGCAG TAGTTGCACT TCTTTGGGGG GTGTACTCAG CCATAGAAAC CTGGAAATTC 14640

```

ATCACCTCCA GATGCCGTTT GTGCTTGCTA GGCCGCAAGT ACATTCTGGC CCCTGCCCAC 14700
CACGTCGAAA GTGCCGCGGG CTTTCATCCG ATTGCGGCAA ATGATAACCA CGCATTTGTC 14760
GTCCGGCGTC CCGGCTCCAC TACGGTTAAC GGCACATTGG TGCCCGGGTT GAAAAGCCTC 14820
GTGTTGGGTG GCAGAAAAGC TGTAAACAG GGAGTGGTAA ACCTTGTC AAATGCCAAA 14880
TAACAACGGC AAGCAGCAAA AGAAAAAGAA GGGGAATGGC CAGCCAGTCA ATCAGCTGTG 14940
CCAGATGCTG GGTAATAATCA TCGCCCAGCA AAACCAGTCC AGAGGCAAGG GACCGGGCAA 15000
GAAAAGTAAG AAGAAAAACC CGGAGAAGCC CCATTTTCCT CTAGCGACCG AAGATGACGT 15060
CAGGCATCAC TTCACCCCTG GTGAGCGGCA ATTGTGTCTG TCGTCGATCC AGACTGCCTT 15120
TAACCAGGGC GCTGGAACCT GTACCCTGTC AGATTCAGGG AGGATAAGTT AACTGTGGA 15180
GTTTAGTTTG CCGACGCATC AACTGTGCG CCTGATCCGC GTCACAGCAT CACCCTCAGC 15240
ATGATGGGCT GGCATTCTTT AGGCACCTCA GTGTCAGAAT TGGAAGAATG TGTGGTGGAT 15300
GGCACTGATT GACATTGTGC CTCTAAGTCA CCTATTCAAT TAGGGCGACC GTGTGGGGGT 15360
AAAATTTAAT TGGCGAGAAC CATGCGGCCG CAATTAAAAA AAAAAAAAAA AAAAAAAAAA 15420
AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA                                     15450

```

(2) INFORMATION FOR SEQ ID NO.: 2:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 7494

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA of Open

Reading Frame 1a of North American PRRS Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 2:

```

ATGTCTGGGA TACTTGATCG GTGCACGTGC ACCCCAATG CCAGGGTGTT TATGGCGGAG 60
GGCCAAGTCT ACTGCACACG ATGTCTCAGT GCACGGTCTC TCCTTCCTCT GAATCTCCAA 120
GTTCTGAGC TTGGGGTGCT GGGCCTATTT TATAGGCCCG AAGAGCCACT CCGGTGGACG 180

```

TTGCCACGTG CATTCCCCAC TGTCGAGTGC TCCCCGCGG GGGCCTGCTG GCTTTCTGCG 240
ATCTTTCCAA TTGCACGAAT GACCAGTGGA AACCTGAACT TTCAACAAAG AATGGTGCGG 300
GTTGCAGCTG AGATTTACAG AGCCGGCCAA CTCACCCCTG CAGTTTTGAA GGCTCTACAA 360
GTTTATGAAC GGGGTTGTCTG CTGGTACCCC ATTGTCGGAC CTGTCCCTGG AGTGGCCGTT 420
CACGCCAACT CCCTACATGT GAGTGACAAA CCTTTCCCGG GAGCAACTCA TGTGTTAACC 480
AACCTACCGC TCCCGCAGAG GCCCAAGCCT GAAGACTTTT GCCCTTTTGA GTGTGCTATG 540
GCTGACGTCT ATGACATTAG CCATGACGCC GTCATGTATG TGGCCAGAGG GAAAGTCTCC 600
TGGGCCCCCTC GTGGCGGGGA TGAAGTGAAA TTTGAAACCG TCCCCGAAGA GTTGAAGTTG 660
ATTGCGAACC GACTCCACAT CTCCTTCCCG CCCACCACG CAGTGGACAT GTCTGAGTTT 720
GCCTTCATAG CCCCTGGGAG TGGTGTCTCC TTGCGGGTCG AGCACCAACA CGGTTGCCTT 780
CCCGCTGATA CTGTCCCTGA AGGGAAGTGC TGGTGGTGCT TGTTTGACTT GCTCCCACCG 840
GAAGTTCAGA ATAAAGAAAT TCGCCGTGCT AACCAATTTG GCTATCAAAC CAAGCATGGT 900
GTCCCTGGCA AGTACCTACA GCGGAGGCTG CAAGTTAATG GTCTCCGAGC AGTGAAGTAT 960
ACAGATGGAC CTATTGTCGT ACAGTACTTC TCTGTTAGGG AGAGTTGGAT CCGCCACTTC 1020
AGACTGGCGG AAGAACCTAG CCTCCCTGGG TTTGAAGACC TCCTCAGAAT AAGGGTAGAG 1080
CCTAATACGT CGCCATTGGG TGGCAAGGGT GAAAAAATCT TCCGGTTTGG CAGTCACAAG 1140
TGGTACGGTG CTGGAAAGAG AGCAAGGAGA GCACGCTCTG GTGCAACTGC CACGGTCGCT 1200
CACTGCGCTT TGCCCGCTCG CGAAGCCCAG CAGGCCAAGA AGCTCGAGGT TGCCAGCGCC 1260
AACAGGGCTG AGCATCTCAA GTACTATTCC CCGCCTGCCG ACGGGAAGTGTG TGGTTGGCAC 1320
TGCATTTCCG CCATTACCAA CCGGATGGTG AATTCCAAAT TTGAAACCAC TCTTCCCGAG 1380
AGAGTGAGAC CTTGAGATGA CTGGGCTACT GACGAGGATC TTGTGAATAC CATCCAAATC 1440
CTCAGGCTCC CCGCGGCCTT GGACAGGAAC GGTGCTTGTTG CTGGCGCCAA GTACGTGCTC 1500
AAGCTGGAAG GTGAGCACTG GACCGTCTCT GTGACCCCTG GGATGACCCC TTCTTTGCTC 1560
CCCCTTGAAT GTGTTGAGG TTGTTGTGAG CATAAGAGCG GTCTTGTTTT CCCAGACGTG 1620
GTCGAAGTTT CCGGATTTGA CCCTGCCTGT CTTGACCGAC TTGCTGAGAT AATGCACTTA 1680
CCTAGCAGTG TCATCCCAGC TGCTCTGGCC GAGATGTCCG ACGACTTCAA TCGTCTGGCT 1740
TCCCCGGCCG CCACTGTGTG GACTGTTTCG CAATTCTTTG CCCGCCACAG AGGAGGAGAG 1800
CATCCTGACC AGGTGTGCTT AGGGAAAATT ATCAACCTTT GTCAGGTGAT TGAGGAATGC 1860
TGCTGTTCCC GGAACAAAGC CAACCGGGCT ACCCCGGAAG AGGTTGCGGC AAAAGTTGAC 1920

CAGTACCTCC GTGGTGCAGC AAGCCTTGGA GAATGCTTGG CCAAGCTTGA GAGGGCTCGC 1980
 CCGCCGAGCG CGATGGACAC CTCCTTTGAT TGGAAATGTTG TGCTTCCTGG GGTTGAGACG 2040
 GCGGATCAGA CAACCAAACA GCTCCATGTC AACCAGTGCC GCGCTCTGGT TCCTGTCGTG 2100
 ACTCAAGAGC CTTTGGACAG AGACTCGGTC CCTCTGACCG CCTTCTCGCT GTCCAATTGC 2160
 TACTACCCTG CACAAGGTGA CGAGGTCCGT CACCGTGAGA GGCTAAACTC CGTGCTCTCT 2220
 AAGTTGGAGG GGGTTGTTCG TGAGGAATAT GGGCTCACGC CAACTGGACC TGGCCCGCGA 2280
 CCCGCACTGC CGAACGGGCT CGACGAGCTT AAAGACCAGA TGGAGGAGGA TCTGCTGAAA 2340
 TTAGTCAACG CCCAGGCAAC TTCAGAAATG ATGGCCTGGG CAGCCGAGCA GGTTGATCTA 2400
 AAAGCTTGGG TCAAAAATTA CCCACGGTGG ACACCGCCAC CCCCTCCACC AAGAGTTCAG 2460
 CCTCGAAAAA CGAAGTCTGT CAAGAGCTTG CTAGAGAACA AGCCTGTCCC TGCTCCGCGC 2520
 AGGAAGGTCA GATCTGATTA TGGCAGCCCG ATTTTGATGG GCGACAATGT TCCTAACGGT 2580
 TGGGAAGATT CGACTGTTGG TGGTCCCCTT GACCTTTCGG CACCATCCGA GCCGATGACA 2640
 CCTCTGAGTG AGCCTGTACT TATTTCCAGG CCAGTGACAT CTTTGAGTGT GCCGGCCCCA 2700
 GTTCCTGCAC CGCGTAGAGC TGTGTCTCGA CCGATGACGC CCTCGAGTGA GCCAATTTTT 2760
 GTGTCTGCAC TGCACACAA ATTTGAGCAG GTGGAAAAAG CAAATCTGGC GGCAGCAGCG 2820
 CCGATGTACC AGGACGAACC CTTAGATTG TCTGCATCCT CACAGACTGA ATATGGGGCT 2880
 TCTCCCCTAA CACCACCGCA GAACGTGGGC ATTCTGGAGG TAAGGGGGCA AGAAGCTGAG 2940
 GAAGTTCTGA GTGAAATCTC GGATATTCTG AATGATACCA ACCCTGCACC TGTGTCATCA 3000
 AGCAGCTCCC TGTCAAGTGT TAGGATCACA CGCCCAAAT ACTCAGCTCA AGCCATTATC 3060
 GACTTGGGCG GGCCCTGCAG TGGGCACCTC CAAAGGGAAA AAGAAGCATG CCTCCGCATC 3120
 ATGCGTGAGG CTTGTGATGC GGCCAAGCTT AGTGACCCTG CCACGCAGGA ATGGCTTTCT 3180
 CGCATGTGGG ATAGGGTGGA CATGCTGACT TGGCGCAACA CGTCTGCTTA CCAGGCGTTT 3240
 CGCACCTTAG ATGGCAGGTT TGGGTTTCTC CCAAAGATGA TACTCGAGAC GCCGCCGCC 3300
 TACCCGTGTG GGTTTGTGAT GTTGCCTCAC ACCCCTGCAC CTTCCGTGAG TGCAGAGAGC 3360
 GACCTTACCA TCGGTTTCAGT CGCCACTGAA GATATTCCAC GCATCCTCGG GAAAATAGAA 3420
 AATACCGGTG AGATGATCAA CCAGGGACCC TTGGCATCCT CTGAGGAAGA ACCGGTATAC 3480
 AACCAACCTG CCAAAGACTC CCGGATATCG TCGCGGGGGT CTGACGAGAG CACAGCAGCT 3540
 CCGTCCGCAG GTACAGGTGG CGCCGGCTTA TTTACTGATT TGCCACCTTC AGACGGCGTA 3600
 GATGCGGACG GTGGGGGGCC GTTGCAGACG GTAAGAAAGA AAGCTGAAAG GCTCTTCGAC 3660

CAATTGAGCC GTCAGGTTTT TAACCTCGTC TCCCATCTCC CTGTTTTCTT CTCACACCTC 3720
TTCAAATCTG ACAGTGGTTA TTCTCCGGGT GATTGGGGT TTGCAGCTTT TACTCTATTT 3780
TGCTCTTTT TGTGTTACAG CTACCCATTC TTCGGTTTCG TTCCCCTCTT GGGTGTATTT 3840
TCTGGGTCTT CTCGGCGTGT GCGCATGGGG GTTTTTGGCT GCTGGCTGGC TTTTGCTGTT 3900
GGCCTGTTCA AGCCTGTGTC CGACCCAGTC GGCCTGCTT GTGAGTTTGA CTCGCCAGAG 3960
TGTAGGAACG TCCTTCATTC TTTTGAGCTT CTCAAACCTT GGGACCCTGT TCGCAGCCTT 4020
GTTGTGGGCC CCGTCGGTCT CGGTCTTGCC ATTCTTGGCA GGTTACTGGG CGGGGCACGC 4080
TACATCTGGC ATTTTTTGCT TAGGCTTGGC ATTGTTGCAG ATTGTATCTT GGCTGGAGCT 4140
TATGTGCTTT CTCAAGGTAG GTGTAAAAAG TGCTGGGGAT CTTGTATAAG AACTGCTCCT 4200
AATGAAATCG CCTTCAACGT GTTCCCTTTT ACACGTGCGA CCAGGTCGTC ACTCATCGAC 4260
CTGTGCGATC GGTTTTGTGC GCCAAAAGGC ATGGACCCCA TTTTCCTCGC CACTGGGTGG 4320
CGTGGGTGCT GGACCGGCCG AAGTCCCATT GAGCAACCCT CTGAAAAACC CATCGCGTTC 4380
GCCCAGTTGG ATGAAAAGAG GATTACGGCT AGAACTGTGG TCGCTCAGCC TTATGATCCT 4440
AATCAAGCCG TAAAGTGCTT GCGGGTGTTA CAGGCGGGTG GGGCGATGGT GGCCGAGGCA 4500
GTCCCAAAG TGGTCAAAGT TTCTGCTATT CCATTCCGAG CCCCTTTTTT TCCCACCGGA 4560
GTGAAAGTTG ATCCCGAGTG CAGGATCGTG GTCGACCCCG ATACTTTTAC TACAGCCCTC 4620
CGGTCTGGTT ACTCTACCAC AAACCTCGTC CTTGGTGTGG GGGACTTTGC CCAGCTGAAT 4680
GGACTAAAGA TCAGGCAAAT TTCCAAGCCT TCGGGAGGAG GCCCACACCT CATTGCTGCC 4740
CTGCATGTTG CCTGCTCGAT GCGGTTGCAC ATGCTTGCTG GGGTTTATGT AACTTCAGTG 4800
GGGTCTTGCG GTGCCGGCAC CAACGATCCA TGGTGCACTA ATCCGTTTGC CGTTCCTGGC 4860
TACGGACCAG GCTCTCTCTG CACGTCCAGA TTGTGCATCT CCCAACATGG CCTTACCCTG 4920
CCCTTGACAG CACTTGTGGC GGGATTCGGT CTTCAGGAAA TCGCCTTGGT CGTTTTGATT 4980
TTCGTTTCCA TCGGAGGCAT GGCTCATAGG TTGAGTTGTA AGGCTGATAT GCTGTGCATC 5040
TACTTGCAA TCGCCAGCTA TGTTTGGGTA CCCCTTACCT GGTTGCTTTG TGTGTTTCCT 5100
TGTTGGTTGC GCTGGTTCTC TTTGCACCCC CTCACCATCC TATGGTTGGT GTTTTTCTTG 5160
ATTTCTGTAA ATATGCCTTC GGGAATCTTG GCCGTGGTGT TATTGGTTTC TCTTTGGCTT 5220
TTGGGACGTT ATACTAACAT TGCTGGTCTT GTCACCCCT ATGATATTCA TCATTACACC 5280
AGTGGCCCCC GCGGTGTTGC CGCCTTAGCT ACCGCACCAG ATGGAACCTA CTTGGCTGCC 5340
GTCCGCCGCG CTGCGTTGAC TGGTCGCACC ATGCTGTTCA CCCCGTCTCA GCTTGGGTCC 5400

CTTCTTGAGG GCGCTTTCAG AACTCGAAAG CCCTCACTGA ACACCGTCAA TGTGGTTGGG 5460
 TCCTCCATGG GCTCTGGTGG AGTGTTCCACC ATCGACGGGA AAATTAGGTG CGTGACTGCC 5520
 GCACATGTCC TTACGGGTAA TTCGGCTAGG GTTTCGGAG TCGGCTTCAA TCAAATGCTT 5580
 GACTTTGATG TGAAAGGGGA CTTCGCCATA GCTGATTGCC CGAATTGGCA AGGAGCTGCT 5640
 CCCAAGACCC AATTCTGCGA GGATGGATGG GCTGGCCGTG CCTATTGGCT GACATCCTCT 5700
 GCGTCGAAC CCGGTGTTAT TGGGAATGGA TTCGCCTTCT GCTTCACCGC GTGCGGCGAT 5760
 TCCGGGTCCC CAGTGATCAC CGAAGCTGGT GAGCTTGTCG GCGTTCACAC AGGATCAAAT 5820
 AAACAAGGAG GTGGCATCGT CACGCGCCCT TCAGGCCAGT TTTGTAACGT GGCACCCATC 5880
 AAGCTGAGCG AATTAAGTGA ATTCTTTGCT GGACCCAAGG TCCCGCTCGG TGATGTGAAG 5940
 GTTGGCAGCC ACATAATTAA AGATACGTGC GAAGTACCTT CAGATCTTTG CGCCTTGCTT 6000
 GCTGCCAAAC CTGAACTGGA GGGAGGCCTC TCCACCGTCC AACTTCTGTG TGTGTTTTTC 6060
 CTACTGTGGA GAATGATGGG ACATGCCTGG ACGCCCTTGG TTGCTGTGGG GTTTTTTCATT 6120
 CTGAATGAGG TTCTCCCAGC TGTCCTGGTT CGGAGTGTTT TCTCCTTTGG GATGTTTGTG 6180
 CTATCTTGGC TCACACCATG GTCTGCGCAA GTTCTGATGA TCAGGCTTCT AACAGCAGCT 6240
 CTTAACAGGA ACAGATGGTC ACTTGCCTTT TACAGCCTTG GTGCGGTGAC CGGTTTTGTC 6300
 GCAGATCTTG CGGCAACTCA AGGGCACCCG TTGCAGGCAG TAATGAATTT GAGCACCTAT 6360
 GCCTTCCTGC CTCGGATGAT GGTTGTGACC TCACCAGTCC CAGTGATTGC GTGTGGTGTT 6420
 GTGCACCTAC TTGCCATCAT TTTGTACTTG TTCAAGTACC GCGGCCTGCA CAATGTTCTT 6480
 GTTGGTGATG GAGCGTTTTT TGCAGCTTTC TTCTTGCGAT ACTTTGCCGA GGGAAAGTTG 6540
 AGGGAAGGGG TGTCGCAATC CTGCGGAATG AATCATGAGT CATTAACTGG TGCCCTCGCT 6600
 ATGAGACTCA ATGACGAGGA CTTGGACTTC CTTACGAAAT GGACTGATTT TAAGTGCTTT 6660
 GTTTCTGCGT CCAACATGAG GAATGCAGCA GGCCAATTCA TCGAGGCTGC CTATGCAAAA 6720
 GCACTTAGAA TTGAACTTGC CCAGTTGGTG CAGGTTGATA AGGTTGAGG TACTTTGGCC 6780
 AAGCTTGAGG CTTTTGCTGA TACCGTGGCA CCCCAACTCT CGCCCGGTGA CATTGTTGTT 6840
 GCTCTTGGCC ATACGCCTGT TGGCAGCATC TTCGACCTAA AGGTTGGTGG TACCAAGCAT 6900
 ACTCTCCAAG TCATTGAGAC CAGAGTCCTT GCCGGGTCCA AAATGACCGT GGCGCGCGTC 6960
 GTTGACCCAA CCCCCACGCC CCCACCCGCA CCCGTGCCCA TCCCCCTCCC ACCGAAAGTT 7020
 CTAGAGAATG GTCCCAACGC CTGGGGGGAT GGGGACCGTT TGAATAAGAA GAAGAGGCGT 7080
 AGGATGGAAA CCGTCGGCAT CTTTGTCATG GGTGGGAAGA AGTACCAGAA ATTTTGGGAC 7140

AAGAATTCCG GTGATGTGTT TTACGAGGAG GTCCATGACA ACACAGATGC GTGGGAGTGC 7200
CTCAGAGTTG GTGACCCTGC CGACTTTGAC CCTGAGAAGG GAACTCTGTG TGGGCATACT 7260
ACTATTGAAG ATAAGGATTA CAAAGTCTAC GCCTCCCCAT CTGGCAAGAA GTTCCTGGTC 7320
CCCGTCAACT CAGAGAGCGG AAGAGCCCAA TGGGAAGCTG CAAAGCTTTC CGTGGAGCAG 7380
GCCCTTGGCA TGATGAATGT CGACGGTGAA CTGACGGCCA AAGAAGTGGA GAAACTGAAA 7440
AGAATAATTG ACAAACCTCA GGCCTGACT AAGGAGCAGT GTTTAAACTG CTAG 7494

(2) INFORMATION FOR SEQ ID NO.: 3:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 4392

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA of

Open Reading Frame 1b of North American PRRS Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 3:

GGAGCAGTGT TTAAACTGCT AGCCGCCAGC GGCTTGACCC GCTGTGGTCG CGGCGGCTTG 60
GTTGTTACTG AGACAGCGGT AAAAATAGTC AAATTTACACA ACCGGACTTT CACCCTAGGG 120
CCTGTGAATT TAAAAGTGGC CAGTGAGGTT GAGCTGAAAG ACGCGGTCGA GCACAACCAA 180
CACCCGGTTG CAAGACCGGT TGACGGTGGT GTTGTGCTCC TCGGTTCCGC AGTTCCTTCG 240
CTTATAGATG TCCTGATCTC CGGTGCTGAC GCATCTCCTA AGTTACTCGC TCGTCACGGG 300
CCGGGGAACA CTGGGATCGA TGGCACGCTT TGGGACTTTG AGGCCGAGGC CACCAAAGAG 360
GAAATTGCAC TCAGTGCGCA AATAATACAG GCTTGTGACA TTAGGCGCGG CGACGCACCT 420
GAAATTGGTC TCCCTTACAA GCTGTACCCT GTTAGGGGCA ACCCTGAGCG GTTAAAAGGA 480
GTTTTACAGA ATACAAGGTT TGGAGACATA CCTTACAAA CCCCCAGTGA CACTGGAAGC 540
CCAGTGCACG CGGCTGCCTG CCTCACGCCC AATGCCACTC CGGTGACTGA TGGGCGCTCT 600
GTCTTGGCTA CTACCATGCC CTCCGGTTTT GAATTGTATG TACCGACCAT TCCAGCGTCT 660

GTCCTTGATT ATCTTGACTC TAGGCCTGAC TGCCCCAAAC AGTTGACAGA GCACGGCTGT 720
GAGGATGCCG CATTGAGAGA CCTCTCCAAG TATGACTTGT CCACCCAAGG CTTTGTTTTTA 780
CCTGGGGTTC TTCGCCTTGT GCGTAAGTAC CTGTTTGCCC ATGTGGGTAA GTGCCCCGCC 840
GTTTCATCGGC CTTCCACTTA CCCTGCCAAG AATTCTATGG CTGGAATAAA TGGGAACAGG 900
TTTCCAACCA AGGACATTCA GAGCGTCCCT GAAATCGACG TTCTGTGCGC ACAGGCCGTG 960
CGAGAAAAC TGGCAAAC TGT TACCCCTTGT ACCCTCAAGA AACAGTATTG TGGGAAGAAG 1020
AAGACTAGGA CAATACTCGG CACCAATAAT TTCATTGCGT TGGCCCACCG GGCAGCGTTG 1080
AGTGGTGTCA CCCAGGGCTT CATGAAAAAG GCGTTTAACT CGCCCATCGC CCTCGGGAAA 1140
AACAAATTTA AGGAGCTACA GACTCCGGTC TTAGGCAGGT GCCTTGAAGC TGATCTTGCA 1200
TCCTGTGATC GATCCACACC TGCAATTGTC CGCTGGTTTG CCGCCAATCT TCTTTATGAA 1260
CTTGCCTGTG CTGAAGAGCA CCTACCGTCG TACGTGCTGA ACTGCTGCCA TGACCTATTG 1320
GTCACGCAGT CCGGCGCAGT GACTAAGAGG GGTGGCCTAT CGTCTGGCGA CCCGATCACT 1380
TCTGTGTCTA ACACCATTTA CAGCTTGGTG ATATATGCAC AGCACATGGT GCTTAGTTAC 1440
TTTAAAAGTG GTCACCCTCA TGGCCTTCTG TTCCTACAAG ACCAGCTGAA GTTCGAGGAC 1500
ATGCTCAAAG TCCAACCCCT GATCGTCTAT TCGGACGACC TCGTGCTGTA TGCCGAATCT 1560
CCCACCATGC CGAACTACCA CTGGTGGGTC GAACATCTGA ATTTGATGCT GGGTTTTTCAG 1620
ACGGACCCAA AGAAGACAGC CATAACGGAC TCGCCATCAT TTCTAGGCTG TAGGATAATA 1680
AATGGACGCC AGCTAGTCCC CAACCGTGAC AGGATCCTCG CGGCCCTCGC TTACCATATG 1740
AAGGCAAGCA ATGTTTCTGA ATACTACGCC GCGGCGGCTG CAATACTCAT GGACAGCTGT 1800
GCTTGTTTAG AGTATGATCC TGAATGGTTT GAAGAGCTTG TGGTTGGGAT AGCGCAGTGC 1860
GCCCCAAGG ACGGCTACAG CTTTCCCGGC CCGCCGTTCT TCTTGTCAT GTGGGAAAAA 1920
CTCAGATCCA ATCATGAGGG GAAGAAGTCC AGAATGTGCG GGTATTGCGG GGCCCCGGCT 1980
CCGTACGCCA CTGCCTGTGG CCTCGACGTC TGTATTTACC ACACCCACTT CCACCAGCAT 2040
TGTCCAGTCA TAATCTGGTG TGGCCACCCG GCTGGTTCTG GTTCTTGTAG TGAGTGCAAA 2100
CCCCCCTAG GGAAAGGCAC AAGCCCTCTA GATGAGGTGT TAGAACAAGT CCCGTATAAG 2160
CCTCCACGGA CTGTAATCAT GCATGTGGAG CAGGGTCTCA CCCCTCTTGA CCCAGGCAGA 2220
TACCAGACTC GCCGCGGATT AGTCTCCGTT AGGCGTGGCA TCAGAGGAAA CGAAGTTGAC 2280
CTACCAGACG GTGATTATGC TAGCACC GCCC CTACTCCCCA CTTGTAAAGA GATCAACATG 2340
GTCGCTGTCG CCTCTAATGT GTTGCGCAGC AGGTTTCATCA TCGGTCCGCC CGGTGCTGGG 2400

AAAACATACT GGCTCCTTCA GCAGGTCCAG GATGGTGATG TCATTTACAC ACCGACTCAC 2460
 CAGACCATGC TCGACATGAT TAGGGCTTTG GGGACGTGCC GGTTC AACGT CCCAGCAGGT 2520
 ACAACGCTGC AATTCCTGC CCCCTCCCGT ACCGGCCCGT GGGTTCGCAT CCTGGCCGGC 2580
 GGTGGTGTC CTGGTAAGAA TTCCTTCCTG GATGAAGCAG CGTATTGTAA TCACCTTGAT 2640
 GTCTTGAGGC TCCTTAGCAA AACCACCCTT ACCTGTCTGG GAGACTTCAA ACAACTCCAC 2700
 CCAGTGGGTT TTGATTCTCA TTGCTATGTT TTTGACATCA TGCCTCAGAC CCAGTTGAAG 2760
 ACCATCTGGA GATTCGGACA GAACATCTGT GATGCCATCC AACCAGATTA CAGGGACAAA 2820
 CTTGTGTCCA TGGTCAACAC AACCCGTGTA ACCTACATGG AAAAACCTGT CAAGTATGGG 2880
 CAAGTCCTCA CCCCTTACCA CAGGGACCGA GAGGACGGCG CCATCACAAT TGA CTCCAGT 2940
 CAAGGCGCCA CATTTGATGT GGT TACACTG CATTTGCCCA CTAAAGATTC ACTCAACAGG 3000
 CAAAGAGCCC TTGTTGCTAT CACCAGGGCA AGACATGCTA TCTTTGTGTA TGACCCACAC 3060
 AGGCAATTGC AGAGCATGTT TGATCTTCCT GCGAAGGGCA CACCCGTCAA CCTCGCAGTG 3120
 CACCGTGATG AGCAGCTGAT CGTACTGGAT AGAAATAATA AAGAATGCAC AGTTGCTCAG 3180
 GCTATAGGCA ACGGAGATAA ATTCAGGGCC ACCGACAAGC GCGTTGTAGA TTCTCTCCGC 3240
 GCCATTTGTG CTGATCTGGA AGGGTCGAGC TCCCCGCTCC CCAAGGTCGC ACACAACTTG 3300
 GGATTTTATT TCTCACCTGA TTTGACACAG TTTGCTAAAC TCCCGGTAGA CCTTGCACCC 3360
 CACTGGCCCG TGGTGACAAC CCAGAACAAT GAAAAGTGGC CGGATCGGCT GGTGCCAGC 3420
 CTTGCCCCTG TCCATAAGTA TAGCCGTGCG TGCATTGGTG CCGGCTATAT GGTGGGCCCC 3480
 TCGGTGTTTC TAGGCACCCC TGGGGTCGTG TCATACTACC TCACAAAATT TGTCAAGGGC 3540
 GAGGCTCAAG TGCTTCCGGA GACAGTCTTC AGCACCGGCC GAATTGAGGT GGATTGCCGG 3600
 GAGTATCTTG ATGACAGGGA GCGAGAAGTT GCTGAGTCCC TCCCACATGC CTTCA TTGGC 3660
 GACGTCAAAG GCACCACCGT TGGGGGATGT CATCATGTCA CCTCCAAATA CCTTCCGCGC 3720
 TTCCTTCCCA AGGAATCAGT CGCGGTAGTC GGGGTTTCGA GCCCCGGGAA AGCCGCAAAA 3780
 GCAGTGTGCA CATTGACGGA TGTGTACCTC CCAGACCTTG AGGCCTACCT CCACCCAGAG 3840
 ACTCAGTCTA AGTGCTGGAA AGTTATGTTG GACTTCAAGG AAGTTCGACT GATGGTCTGG 3900
 AAAGACAAGA CGGCCTATTT CCAACTTGAA GGCCGCTATT TCACCTGGTA TCAGCTTGCA 3960
 AGCTACGCCT CGTACATCCG TGTTCCTGTC AACTCCACGG TGTATCTGGA CCCCTGCATG 4020
 GGCCCTGCCC TTTGCAACAG AAGAGTTGTC GGGTCCACCC ATTGGGGAGC TGACCTCGCA 4080
 GTCACCCCTT ATGATTACGG TGCTAAAATC ATCTTGCTA GCGCTTACCA TGGTGAAATG 4140

CCTCCTGGAT ACAAGATTCT GGCGTGCGCG GAGTTCTCGC TCGACGACCC AGTCAAGTAC 4200
 AAACACACCT GGGGTTTTGA ATCGGATACA GCGTATCTGT ATGAGTTCAC CGGAAACGGT 4260
 GAGGACTGGG AGGATTACAA TGATGCGTTT CGTGCGCGCC AGAAAGGGAA AATTTATAAG 4320
 GCCACTGCTA CCAGCATGAA GTTTTATTTT CCCCCGGGCC CCGTCATTGA ACCAACTTTA 4380
 GGCCTGAATT GA 4392

(2) INFORMATION FOR SEQ ID NO.: 4:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 771

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA of

Open Reading Frame 2 of North American PRRS Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 4:

ATGAAATGGG GTCTATACAA AGCCTCTTCG ACAAATTTGG CCAGCTTTTT GTGGATGCTT 60
 TCACGGAATT TTTGGTGTCC ATTGTTGATA TCATCATATT TTTGGCCATT TTGTTTGGCT 120
 TCACCATCGC CGGTTGGCTG GTGGTCTTTT GCATCAGATT GGTTTGCTCC GCGGTATTCC 180
 GTGCGCGCCC TGCCATTACAC CCTGAGCAAT TACAGAAGAT CCTATGAGGC CTTTCTTTCT 240
 CAGTGCCGGG TGGACATTCC CACCTGGGGG GTAAAACACC CTTTGGGGAT GTTTTGGCAC 300
 CATAAGGTGT CAACCCTGAT TGATGAAATG GTGTCGCGTC GAATGTACCG CATCATGGAA 360
 AAAGCAGGGC AAGCTGCCTG GAAACAGGTG GTGAGCGAGG CTACGCTGTC TCGCATTAGT 420
 AGTTTGGATG TGGTGGCTCA TTTTCAACAT CTTGCCGCCA TTGAAGCCGA GACCTGTAAA 480
 TATTTGGCTT CTCGACTGCC CATGCTACAC AACCTGCGCA TGACAGGGTC AAATGTAACC 540
 ATAGTGTATA ATAGCACTTT AAATCAGGTG TTTGCTATTT TTCCAACCCC TGGTTCCCGG 600
 CCAAAGCTTC ATGATTTTCA GCAATGGCTA ATAGCTGTAC ATTCCTCCAT ATTTTCCTCT 660
 GTTGCAGCTT CTTGTACTCT TTTTGTGTG CTGTGGTTGC GGGTTCCAAT GCTACGTACT 720

GTTTTTGGTT TCCGCTGGTT AGGGGCAATT TTTCTTTCGA ACTCATGGTG A 771

(2) INFORMATION FOR SEQ ID NO.: 5:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 765

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA of Open

Reading Frame 3 of North American PRRS Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 5:

ATGGCTAATA GCTGTACATT CCTCCATATT TTCCTCTGTT GCAGCTTCTT GTACTCTTTT 60
TGTTGTGCTG TGGTTGCGGG TTCCAATGCT ACGTACTGTT TTTGGTTTCC GCTGGTTAGG 120
GGCAATTTTT CTTTCGAACT CATGGTGAAT TACACGGTGT GTCCACCTTG CCTCACCCGA 180
CAAGCAGCCG CTGAGGTCCT TGAACCCGGT AGGTCTCTTT GGTGCAGGAT AGGGCATGAC 240
CGATGTGGGG AGGACGATCA CGACGAACTA GGGTTCATGG TTCCGCCTGG CCTCTCCAGC 300
GAAAGCCACT TGACCAGTGT TTACGCCTGG TTGGCGTTCC TGTCCTTCAG CTACACGGCC 360
CAGTTCCATC CCGAGATATT TGGGATAGGG AACGTGAGTG AAGTTTATGT TGACATCAAG 420
CACCAATTCA TCTGCGCCGT TCATGACGGG CAGAACACCA CCTTGCCTCG CCATGACAAT 480
ATTTAGCCG TATTTAGAC CTACTATCAA CATCAGGTCG ACGGCGGCAA TTGGTTTCAC 540
CTAGAATGGC TCGTCCCTT CTTTTCCTCT TGGTTGGTTT TAAATGTTTC GTGGTTTCTC 600
AGGCGTTCGC CTGCAAGCCA TGTTTCAGTT CGAGTCTTTC AGACATCAAA ACCAACAATA 660
CCGCAGCATC AGGCTTTGTT GTCCTCCAGG ACATCAGCTG CCTTAGGCAT GGCGACTCGT 720
CCTTTCCGAC GATTCGCAAA AGCTCTCAAT GCCGCACGGC GATAG 765

(2) INFORMATION FOR SEQ ID NO.: 6:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 537

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA of Open

Reading Frame 4 of North American PRRS Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 6:

ATGGCTGCGT CCCTTCTTTT CCTCTTGGTT GGTTTTAAAT GTTTCGTGGT TTCTCAGGCG 60
TTCGCCTGCA AGCCATGTTT CAGTTCGAGT CTTTCAGACA TCAAAACCAA CACTACCGCA 120
GCATCAGGCT TTGTTGTCCT CCAGGACATC AGCTGCCTTA GGCATGGCGA CTCGTCCTTT 180
CCGACGATTC GCAAAGCTC TCAATGCCGC ACGGCGATAG GGACACCCGT GTATATCACC 240
ATCACAGCCA ATGTGACAGA TGAGAATTAC TTACATTCTT CTGATCTCCT CATGCTTTCT 300
TCTTGCCTTT TCTATGCTTC TGAGATGAGT GAAAAGGGAT TCAAGGTGGT GTTTGGCAAT 360
GTGTCAGGCA TCGTGGCTGT GTGTGTCAAC TTTACCAGCT ACGTCCAACA TGTCAAAGAG 420
TTTACCCAAC GCTCCTTGGT GGTCGATCAT GTGCGGCTGC TTCATTTTCAT GACACCTGAG 480
ACCATGAGGT GGGCAACCGT TTTAGCCTGT CTTTTTGCCA TCCTACTGGC AATTTGA 537

(2) INFORMATION FOR SEQ ID NO.: 7:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 603

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA of Open

Reading Frame 5 of North American PRRS Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 7:

```

ATGTTGGGGA AATGCTTGAC CGCGGGCTGT TGCTCGCGAT TGCTTTCTTT GTGGTGTATC 60
GTGCCGTTCT GTTTTGCTGT GCTCGGCAGC GCCAACAGCA GCAGCAGCTC TCATTTTCAG 120
TTGATTTATA ACTTGACGCT ATGTGAGCTG AATGGCACAG ATTGGCTGGC AGAAAAATTT 180
GATTGGGCAG TGGAGACTTT TGTCATCTTT CCCGTGTTGA CTCACATTGT TTCCTATGGT 240
GCACTCACCA CCAGCCATTT CCTTGACACA GTTGGTCTGG TTAGTGTC CACCGCCGGG 300
TTTTATCACG GGCGGTATGT CTTGAGTAGC ATCTACGCGG TCTGTGCTCT GGCTGCGTTG 360
ATTTGCTTCG TTATTAGGCT TGCGAAGAAC TGCATGTCCT GGCGCTACTC TTGTACCAGA 420
TATACCAACT TCCTTCTGGA CACTAAGGGC AGACTCTATC GTTGGCGGTC GCCCGTTATC 480
ATAGAAAAAG GGGGTAAGGT TGAGGTCGAA GGTACCTGA TCGACCTCAA AAGAGTTGTG 540
CTTGATGGTT CCGTGGCAAC CCCTTTAACC AGAGTTTCAG CGGAACAATG GGGTCGTCTC 600
TAG 603

```

(2) INFORMATION FOR SEQ ID NO.: 8:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 525

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA of Open

Reading Frame 6 of North American PRRS Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 8:

```

ATGGGGTCGT CTCTAGACGA CTTTGGCCAT GATAGCACGG CTCCACAAAA GGTGCTTTTG 60
GCGTTTTCCA TTACCTACAC GCCAGTAATG ATATATGCTC TAAAGGTAAG TCGCGGCCGA 120
CTACTAGGGC TTCTGCACCT TTTGATCTTT CTGAATTGTG CTTTACCTT CGGGTACATG 180
ACATTCGAGC ACTTTCAGAG CACAAATAGG GTCGCGCTCA CTATGGGAGC AGTAGTTGCA 240
CTTCTTTGGG GGGTGTACTC AGCCATAGAA ACCTGGAAAT TCATCACCTC CAGATGCCGT 300

```

TTGTGCTTGC TAGGCCGCAA GTACATTCTG GCCCCTGCCC ACCACGTCGA AAGTGCCGCG 360
GGCTTTCATC CGATTGCGGC AAATGATAAC CACGCATTTG TCGTCCGGCG TCCCGGCTCC 420
ACTACGGTTA ACGGCACATT GGTGCCCCGGG TTGAAAAGCC TCGTGTGGG TGGCAGAAAA 480
GCTGTTAAAC AGGGAGTGGT AAACCTTGTC AAATATGCCA AATAA 525

(2) INFORMATION FOR SEQ ID NO.: 9:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 372

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: cDNA of Open

Reading Frame 7 of North American PRRS Virus

Genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 9:

ATGCCAAATA ACAACGGCAA GCAGCAAAAG AAAAAGAAGG GGAATGGCCA GCCAGTCAAT 60
CAGCTGTGCC AGATGCTGGG TAAATCATC GCCCAGCAAA ACCAGTCCAG AGGCAAGGGA 120
CCGGGCAAGA AAAGTAAGAA GAAAACCCG GAGAAGCCCC ATTTTCCTCT AGCGACCGAA 180
GATGACGTCA GGCATCACTT CACCCCTGGT GAGCGGCAAT TGTGTCTGTC GTCGATCCAG 240
ACTGCCTTTA ACCAGGGCGC TGGAACCTGT ACCCTGTCAG ATTCAGGGAG GATAAGTTAC 300
ACTGTGGAGT TTAGTTTGCC GACGCATCAT ACTGTGCGCC TGATCCGCGT CACAGCATCA 360
CCCTCAGCAT GA 372

(2) INFORMATION FOR SEQ ID NO.: 10:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 18

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

forward strand, used for determining cDNA

corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 10:

ACAGTTTGGT GATCTATG

18

(2) INFORMATION FOR SEQ ID NO.: 11:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 17

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

reverse strand, used for determining cDNA

corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 11:

CAGATTCAGA TGTTCAA

17

(2) INFORMATION FOR SEQ ID NO.: 12:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 24

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer used

for determining cDNA corresponding to North
American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 12:

ACCTCGTGCT GTATGCCGAA TCTC

24

(2) INFORMATION FOR SEQ ID NO.: 13:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 24

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer, used

for determining cDNA corresponding to North
American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 13:

TCAGGCCTAA AGTTGGTTCA ATGA

24

(2) INFORMATION FOR SEQ ID NO.: 14:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 24

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

forward strand, used for determining cDNA
corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 14:

GATGACTGGG CTACTGACGA GGAT

24

(2) INFORMATION FOR SEQ ID NO.: 15:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 21

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

reverse strand, used for determining cDNA

corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 15:

AGAGCGGCTG GGATGACACT G

21

(2) INFORMATION FOR SEQ ID NO.: 16:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 22

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer, used

for determining cDNA corresponding to North

American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 16:

CCGGGGAAGC CAGACGATTG AA

22

(2) INFORMATION FOR SEQ ID NO.: 17:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 24

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer, used

for determining cDNA corresponding to North

American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 17:

AGGGGGAGCA AAGAAGGGGT CATC

24

(2) INFORMATION FOR SEQ ID NO.: 18:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 20

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer

forward strand, used for determining cDNA

corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 18:

AGCACGCTCT GGTGCAACTG

20

(2) INFORMATION FOR SEQ ID NO.: 19:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 19

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

reverse strand, used for determining cDNA

corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 19:

GCCGCGGCGT AGTATTCAG

19

(2) INFORMATION FOR SEQ ID NO.: 20:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 22

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

forward strand, used for determining cDNA

corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 20:

CGCGTCACAG CATCACCTC AG

22

(2) INFORMATION FOR SEQ ID NO.: 21:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 25

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Primer,
reverse strand, used for determining cDNA
corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 21:
CGGTAGGTTG GTTAACACAT GAGTT 25

(2) INFORMATION FOR SEQ ID NO.: 22:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 23

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Primer,
reverse strand, used for determining cDNA
corresponding to North American PRRS virus genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 22:
TGGCTCTTCG GGCCTATAAA ATA 23

(2) INFORMATION FOR SEQ ID NO.: 23:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 56

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Strand of
synthetic doublestranded adapter used in pT7P129A.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 23:

CTAGATTAAT TAATACGACT CACTATAGGG ATGACGTATA GGTGTTGGCT CTATGC 56

(2) INFORMATION FOR SEQ ID NO.: 24:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 56

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Strand of

synthetic double-stranded adapter used in

pT7P129A.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 24:

TAATTAATTA TGCTGAGTGA TATCCCTACT GCATATCCAC AACCGAGATA CGGTGC 56

(2) INFORMATION FOR SEQ ID NO.: 25:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 27

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

forward strand, used in preparing pT7P129A.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 25:

ACTCAGTCTA AGTGCTGGAA AGTTATG 27

(2) INFORMATION FOR SEQ ID NO.: 26:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 59

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

reverse strand, used in preparing pT7P129A.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 26:

GGGATTAAA TATGCATTTT TTTTTTTTTT TTTTTTTAAT TCGGGCCGCA TGGTTCTCG 59

(2) INFORMATION FOR SEQ ID NO.: 27:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 36

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Primer,

forward strand, used for synthesizing region

flanking North American PRRS virus ORF7 upstream.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 27:

ATTAGATCTT GCCACCATGG TGGGGAAATG CTTGAC

36

(2) INFORMATION FOR SEQ ID NO.: 28:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 46

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

64680-1480

75

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Primer,
reverse strand, used for synthesizing region
flanking North American PRRS virus ORF7 upstream.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 28:

CTTTACGCGT TTGCTTAAGT TATTTGGCGT ATTTGACAAG GTTTAC

46

(2) INFORMATION FOR SEQ ID NO.: 29:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 31

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Primer,
forward strand, used for synthesizing region
FLANKING NORTH AMERICAN PRRS VIRUS ORF7
downstream.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 29:

CAACACGCGT CAGCAAAAGA AAAAGAAGGG G

31

(2) INFORMATION FOR SEQ ID NO.: 30:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 20

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Primer,
reverse strand, used for synthesizing region
FLANKING NORTH AMERICAN PRRS VIRUS ORF7
downstream.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 30:
GCGCGTTGGC CGATTCATTA 20

(2) INFORMATION FOR SEQ ID NO.: 31:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 37

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Upper primer
used in preparing pCMV-S-P129.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 31:
CTCGTTAATT AAACCGTCAT GACGTATAGG TGTTGGC 37

(2) INFORMATION FOR SEQ ID NO.: 32:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 3796

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Plasmid

(ix) FEATURE:

(D) OTHER INFORMATION: DESCRIPTION OF PLASMID: PCMV-MC1

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 32:
GAATTCGAGC TTGCATGCCT GCAGGTCGTT ACATAACTTA CGGTAAATGG CCCGCCTGGC 60

TTCGGTGTAG GTCGTTCGCT CCAAGCTGGG CTGTGTGCAC GAACCCCCCG TTCAGCCCGA 1860
 CCGCTGCGCC TTATCCGGTA ACTATCGTCT TGAGTCCAAC CCGGTAAGAC ACGACTTATC 1920
 GCCACTGGCA GCAGCCACTG GTAACAGGAT TAGCAGAGCG AGGTATGTAG GCGGTGCTAC 1980
 AGAGTTCTTG AAGTGGTGGC CTAACACGG CTACACTAGA AGGACAGTAT TTGGTATCTG 2040
 CGCTCTGCTG AAGCCAGTTA CCTTCGGAAA AAGAGTTGGT AGCTCTTGAT CCGGCAAACA 2100
 AACCACCGCT GGTAGCGGTG GTTTTTTTGT TTGCAAGCAG CAGATTACGC GCAGAAAAAA 2160
 AGGATCTCAA GAAGATCCTT TGATCTTTTC TACGGGGTCT GACGCTCAGT GGAACGAAAA 2220
 CTCACGTAA GGGATTTTGG TCATGAGATT ATCAAAAAGG ATCTTCACCT AGATCCTTTT 2280
 AAATTAAAAA TGAAGTTTAA AATCAATCTA AAGTATATAT GAGTAAACTT GGTCTGACAG 2340
 TTACCAATGC TTAATCAGTG AGGCACCTAT CTCAGCGATC TGTCTATTTT GTTCATCCAT 2400
 AGTTGCCTGA CTCCCCGTCG TGTAAGATAAC TACGATACGG GAGGGCTTAC CATCTGGCCC 2460
 CAGTGCTGCA ATGATACCGC GAGACCCACG CTCACCGGCT CCAGATTTAT CAGCAATAAA 2520
 CCAGCCAGCC GGAAGGGCCG AGCGCAGAAG TGGTCCTGCA ACTTTATCCG CCTCCATCCA 2580
 GTCTATTAAT TGTTGCCGGG AAGCTAGAGT AAGTAGTTCG CCAGTTAATA GTTTGCGCAA 2640
 CGTTGTTGCC ATTGCTACAG GCATCGTGGT GTCACGCTCG TCGTTTGGTA TGGCTTCATT 2700
 CAGCTCCGGT TCCCAACGAT CAAGGCGAGT TACATGATCC CCCATGTTGT GCAAAAAAGC 2760
 GGTTAGCTCC TTCGGTCCTC CGATCGTTGT CAGAAGTAAG TTGGCCGCAG TGTTATCACT 2820
 CATGGTTATG GCAGCACTGC ATAATTCTCT TACTGTCATG CCATCCGTAA GATGCTTTTC 2880
 TGTGACTGGT GAGTACTCAA CCAAGTCATT CTGAGAATAG TGTATGCGGC GACCGAGTTG 2940
 CTCTTGCCCG GCGTCAATAC GGGATAATAC CGCGCCACAT AGCAGAACTT TAAAAGTGCT 3000
 CATCATTGGA AAACGTTCTT CGGGGCGAAA ACTCTCAAGG ATCTTACCGC TGTTGAGATC 3060
 CAGTTCGATG TAACCCACTC GTGCACCCAA CTGATCTTCA GCATCTTTTA CTTTCACCAG 3120
 CGTTTCTGGG TGAGCAAAAA CAGGAAGGCA AAATGCCGCA AAAAAGGGAA TAAGGGCGAC 3180
 ACGGAAATGT TGAATACTCA TACTCTTCCT TTTTCAATAT TATTGAAGCA TTTATCAGGG 3240
 TTATTGTCTC ATGAGCGGAT ACATATTTGA ATGTATTTAG AAAAATAAAC AAATAGGGGT 3300
 TCCGCGCACA TTTCCCCGAA AAGTGCCACC TGACGTCTAA GAAACCATTA TTATCATGAC 3360
 ATTAACCTAT AAAAATAGGC GTATCACGAG GCCCTTTCGT CTCGCGCGTT TCGGTGATGA 3420
 CGGTGAAAAC CTCTGACACA TGCAGCTCCC GGAGACGGTC ACAGCTTGTC TGTAAGCGGA 3480
 TGCCGGGAGC AGACAAGCCC GTCAGGGCGC GTCAGCGGGT GTTGGCGGGT GTCGGGGCTG 3540

GCTTAACTAT GCGGCATCAG AGCAGATTGT ACTGAGAGTG CACCATATGC GGTGTGAAAT 3600
ACCGCACAGA TGCGTAAGGA GAAAATACCG CATCAGGCGC CATTCGCCAT TCAGGCTGCG 3660
CAACTGTTGG GAAGGGCGAT CGGTGCGGGC CTCTTCGCTA TTACGCCAGC TGGCGAAAGG 3720
GGGATGTGCT GCAAGGCGAT TAAGTTGGGT AACGCCAGGG TTTTCCCAGT CACGACGTTG 3780
TAAAACGACG GCCAGT 3796

(2) INFORMATION FOR SEQ ID NO.: 33:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 22

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Strand from

synthetic linker used in the preparation of

pCMV-S-P129.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 33:

CGTTAATTAA ACCGACTAGT GC

22

(2) INFORMATION FOR SEQ ID NO.: 34:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 30

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence: Strand from

synthetic linker used in the preparation of

pCMV-S-P129.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 34:

TCGAGCAATT AATTGGCTG ATCACGCCGG

30

(2) INFORMATION FOR SEQ ID NO.: 35:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 24

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Lower primer

used in preparing pCMV-S-P129.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 35:

CGGGGACGGT TTCAAATTC ACTT

24

(2) INFORMATION FOR SEQ ID NO.: 36:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 50

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence: Portion of

plasmid pCMV-S-P129, in 5'to 3' direction,

immediately prior to P129 genome.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 36:

TATATAAGCA GAGCTCGTTA ATTAAACCGT CATGACGTAT AGGTGTTGGC

50

(2) INFORMATION FOR SEQ ID NO.: 37:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 48

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence:Primer,

forward, used for synthesizing upstream flanking

REGION TO ORF4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 37:

AGGTCGACGG CGGCAATTGG TTTCACCTAG AGTGGCTGCG TCCCTTCT

48

(2) INFORMATION FOR SEQ ID NO.: 38:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 30

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence:Primer,

reverse, used for synthesizing upstream flanking

REGION TO ORF4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 38:

TCTTAAGCAT TGGCTGTGAT GGTGATATAC

30

(2) INFORMATION FOR SEQ ID NO.: 39:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 44

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence:Primer,

froward, used for synthesizing downstream flanking

REGION TO ORF4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 39:

CTTCTTAAGT CCACGCGTTT TCTTCTTGCC TTTTCTATGC TTCT

44

(2) INFORMATION FOR SEQ ID NO.: 40:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 19

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence:Primer,

reverse, used for synthesizing downstream flanking

REGION TO ORF4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 40:

TGCCCCGGTCC CTTGCCTCT

19

(2) INFORMATION FOR SEQ ID NO.: 41:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 26

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence:Primer,

forward, used for synthesizing downstream flanking

REGION TO ORF4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 41:

GTTTACGCGT CGCTCCTTGG TGGTCG

26

(2) INFORMATION FOR SEQ ID NO.: 42:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 23

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence:Primer,

forward, used for synthesizing upstream flanking

REGION TO INSERTION SITE BETWEEN ORF1B AND ORF2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 42:

AACAGAAGAG TTGTCGGGTC CAC

23

(2) INFORMATION FOR SEQ ID NO.: 43:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 49

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial
Sequence:Primer,

reverse, used for synthesizing upstream flanking

REGION TO INSERTION SITE BETWEEN ORF1B AND ORF2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 43:

GCTTTGACGC GTCCCCACTT AAGTTCAATT CAGGCCTAAA GTTGGTTCA 49

(2) INFORMATION FOR SEQ ID NO.: 44:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 82

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence:Primer,

forward, used for synthesizing downstream flanking

REGION TO INSERTION SITE BETWEEN ORF1B AND ORF2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 44:

GCGACGCGTG TTCCGTGGCA ACCCCTTTAA CCAGAGTTTC AGCGGAACAA TGAAATGGGG 60
TCTATACAAA GCCTCTTCGA CA 82

(2) INFORMATION FOR SEQ ID NO.: 45:

(i) SEQUENCE CHARACTERISTICS

(A) LENGTH: 26

(ii) MOLECULAR TYPE: DNA

(A) ORGANISM: Artificial Sequence

(ix) FEATURE:

(D) OTHER INFORMATION: Description of Artificial

Sequence:Primer,

reverse, used for synthesizing downstream flanking

REGION TO INSERTION SITE BETWEEN ORF1B AND ORF2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.: 45:

AACAGAACGG CACGATACAC CACAAA

CLAIMS:

1. An isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a PRRS virus, wherein said DNA sequence is (i) SEQ ID NO:1;
5 (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1
10 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1.
2. An isolated infectious RNA molecule encoded by the isolated polynucleotide molecule according to claim 1, which infectious RNA molecule encodes a PRRS virus.
- 15 3. An isolated polynucleotide molecule according to claim 1 in the form of a plasmid.
4. A transfected host cell comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence is (i) SEQ ID
20 NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID
25 NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, which transfected host cell is capable of expressing the encoded North American PRRS virus.
5. A plasmid capable of directly transfecting a
30 suitable host cell and completely cellularly expressing a

North American PRRS virus from the suitable host cell so transfected, the plasmid comprising

(a) a DNA sequence that is (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1
5 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence
10 having at least 85% identity to SEQ ID NO:1, wherein said DNA sequence encodes an infectious RNA molecule encoding the North American PRRS virus; and

(b) a promoter capable of transcribing said infectious RNA molecule in said suitable host cell.

15 6. A completely cellular method for generating a North American PRRS virus, which method comprises transfecting a suitable host cell with the plasmid according to claim 5 encoding the North American PRRS virus and obtaining North American PRRS virus generated by the transfected host cell.

20 7. An isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus that is genetically modified such that it comprises one or more heterologous antigenic epitopes, wherein said DNA sequence comprises (i) SEQ ID
25 NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID
30 NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, and wherein said

DNA sequence further comprises one or more further nucleotide sequences that each encode a heterologous antigenic epitope, and wherein each heterologous antigenic epitope is capable of inducing an effective immunoprotective response against a particular pathogen corresponding to said heterologous antigenic epitope in a mammal or a bird.

8. An isolated polynucleotide molecule according to claim 7 in the form of a plasmid.

9. The isolated polynucleotide molecule according to claim 7 or claim 8 wherein the particular pathogen is a swine pathogen.

10. A transfected host cell comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence comprises (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1, and wherein said DNA sequence further comprises one or more further nucleotide sequences that each encode a heterologous antigenic epitope, and wherein each heterologous antigenic epitope is capable of inducing an effective immunoprotective response against a particular pathogen corresponding to said heterologous antigenic epitope in a mammal or a bird.

11. An isolated polynucleotide molecule comprising a DNA sequence encoding an infectious RNA molecule encoding a North American PRRS virus, wherein said DNA sequence comprises (i) the sequence as set forth in SEQ ID NO:1; or

(ii) the sequence beginning with and including nucleotide 1 through and including nucleotide 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the
5 nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1 and wherein said DNA sequence further comprises one or more further nucleotide sequences that each encode a detectable
10 heterologous antigenic epitope.

12. A method for generating an infectious a North American PRRS virus encoded by an infectious RNA molecule that is (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide
15 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or (iii) a sequence having at least 85% identity to SEQ ID NO:1,
20 and wherein said method comprises further transfecting the transfected host cell according to claim 10 with the infectious RNA molecule according to claim 2, or the plasmid according to claim 5, or a viral vector comprising the polynucleotide molecule according to claim 1 or claim 11
25 encoding the North American PRRS virus.

13. A method for generating an infectious North American PRRS virus encoded by an infectious RNA sequence that is (i) SEQ ID NO:1; (ii) the sequence beginning with and including nucleotide 1 through and including nucleotide
30 15,416 of SEQ ID NO:1, wherein the nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1 is a guanine instead of an adenine and the nucleotide corresponding to nucleotide

1,559 of SEQ ID NO:1 is a thymine instead of a cytosine; or
(iii) a sequence having at least 85% identity to SEQ ID NO:1,
and wherein said method comprises transfecting a suitable
cell with the infectious RNA molecule according to claim 2,
5 or the plasmid according to claim 5, or a viral vector
comprising the polynucleotide molecule according to claim 1 or
claim 11, encoding the North American PRRS virus and with a
helper virus that expresses the North American PRRS virus
peptide or sequences of said North American PRRS virus, and
10 separating North American PRRS virions expressed by the cell
from said helper virus.

14. A completely cellular method for preparing a
genetically modified North American PRRS virus that is
capable of eliciting an immunoprotective response in a mammal
15 or a bird vaccinated therewith, wherein said method comprises

(a) obtaining an isolated polynucleotide molecule
comprising a DNA sequence encoding an infectious RNA
molecule which encodes a wild-type North American PRRS, and
wherein said DNA sequence is (i) SEQ ID NO:1; or (ii) the
20 sequence beginning with and including nucleotide 1 through
and including nucleotide 15,416 of SEQ ID NO:1, wherein the
nucleotide corresponding to nucleotide 12,622 of SEQ ID NO:1
is a guanine instead of an adenine and the nucleotide
corresponding to nucleotide 1,559 of SEQ ID NO:1 is a thymine
25 instead of a cytosine; or (iii) a sequence having at least
85% identity to SEQ ID NO:1;

(b) genetically mutating the DNA sequence
encoding the infectious RNA molecule which encodes the wild-
type North American PRRS virus so as to obtain an isolated
30 polynucleotide molecule comprising a DNA sequence encoding an
infectious RNA molecule which encodes a genetically modified
North American PRRS virus having at least 85% identity to SEQ

ID NO:1, and wherein said genetically modified North American PRRS virus remains able to elicit an effective immunoprotective response against infection by the wild-type North American PRRS virus in a mammal or a bird; and

5 (c) completely cellularly expressing the genetically modified North American PRRS virus from the isolated polynucleotide molecule so obtained.

15. A vaccine for protecting a mammal or a bird from infection by a pathogen, which vaccine comprises a
10 genetically modified North American PRRS virus encoded by an infectious RNA molecule encoded by the isolated polynucleotide molecule defined in claim 7, or said infectious RNA molecule, or a plasmid comprising said polynucleotide molecule, in an amount effective to produce
15 immunoprotection against infection by a pathogen from which the heterologous antigenic epitope or epitopes thereof or encoded thereby are derived; and a carrier acceptable for pharmaceutical or veterinary use.

16. An isolated polynucleotide molecule comprising a
20 DNA sequence encoding an infectious RNA molecule wherein said DNA sequence is at least 85% identical to the sequence of SEQ ID NO:1.





FIG. 1

FIG. 2**FIG. 3**