

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2014/093518 A1

(43) International Publication Date

19 June 2014 (19.06.2014)

(51) International Patent Classification:

A01N 25/12 (2006.01) *A01N 47/02* (2006.01)
A01N 53/00 (2006.01) *A01P 7/00* (2006.01)
A01N 43/36 (2006.01)

(21) International Application Number:

PCT/US2013/074447

(22) International Filing Date:

11 December 2013 (11.12.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/736,151	12 December 2012 (12.12.2012)	US
12197917.3	18 December 2012 (18.12.2012)	EP
61/897,018	29 October 2013 (29.10.2013)	US

(71) Applicant: **BASF CORPORATION** [US/US]; 100 Park Avenue, Florham Park, New Jersey 07932 (US).

(72) Inventor: **RICHARDSON, Ronald O.**; 3568 Tree Court Industrial Blvd., St. Louis, Missouri 63122 (US).

(74) Agents: **GOFF, Christopher M.** et al.; Armstrong Teasdale LLP, 7700 Forsyth Blvd., Suite 1800, St. Louis, Missouri 63105-1847 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2014/093518 A1

(54) Title: SOLID AGROFORMULATIONS FOR PREPARING NEAR MICRO-EMULSION AQUEOUS PESTICIDES

(57) Abstract: Disclosed are extruded pesticide granules that when mixed with water form a near micro-emulsion that is highly stable. The extruded pesticide granules include at least one pesticide active ingredient, a non-ionic surfactant, and a carrier, which is desirably urea powder or urea pearl. Processes for making the extruded pesticide granules are also disclosed.

SOLID AGROFORMULATIONS FOR PREPARING NEAR MICRO-EMULSION AQUEOUS PESTICIDES

FIELD OF THE DISCLOSURE

[0001] The present disclosure relates to solid agroformulations and methods of preparing the same. More particularly, the present disclosure relates to extruded pesticide granules including a urea carrier and at least one pesticide active agent that when mixed with water form long term stable near micro-emulsions suitable for pesticide application. The present disclosure also relates to methods of preparing and using the solid agroformulations.

BACKGROUND OF THE DISCLOSURE

[0002] Crop and pest protection agents have conventionally been formulated in solid or liquid compositions, usually in the form of a concentrate for ease of handling and transportation. The concentrate is generally diluted with water by the user before application. Many liquid formulations in the form of emulsifiable or near-emulsifiable concentrates may contain a very high proportion of organic solvents (often up to 80 percent), which are increasingly coming under scrutiny for their potential negative effect on the environment.

[0003] Water-based suspension concentrates, which are another conventionally available form, are often viscous giving rise to handling problems and loss of active ingredient through retention in the packaging. Solid formulations, which may also be commercially available, can also have disadvantages; the more common granules and powders in particular can be difficult to measure but more importantly can be dusty and pose inhalation hazards for the formulator and the user. Tablets have not been used extensively because they are often slow to dissolve. In addition, solid formulations have been found generally to possess a lower biological activity than liquid formulations. Also, with unsophisticated mixing techniques at the site of use, the tendency of solid forms not to disperse immediately can cause not only clogging of spray equipment with undispersed

formulation, but also an inadequate application of active ingredient to the crop or area to be treated.

[0004] Based on the foregoing, there is a need for fast-dispersing, solid pesticide formulations which have improved handling characteristics and enhanced biological activity over conventional forms, to satisfy both environmental concerns and provide an effective pesticidal product with commercially acceptable levels of stability for long term use.

SUMMARY OF THE DISCLOSURE

[0005] The present disclosure is directed to solid agroformulations, and in particular extruded pesticide granules, and liquid aqueous pesticide formulations including these granules. The extruded pesticide granules include a urea carrier in combination with a non-ionic surfactant, at least one pesticide active ingredient, and a small amount of residual water. This novel combination of components, when extruded as described herein, rapidly dissolves in room temperature water with gentle agitation to form a near micro-emulsion that has only a very slight haze that is substantially resistant to settling out of materials. In many embodiments, the extruded pesticide granules as described herein will dissolve in two minutes or less in room temperature water with gentle agitation. Because of these significant dissolution properties and the formation of a near micro-emulsion and retardation of settling for many days, the extruded pesticide granules described herein may be easily mixed and used “on site” thus improving the usefulness of the agroformulation.

[0006] The present disclosure is further directed to an extruded pesticide granular composition comprising urea, a non-ionic surfactant, a pesticide active ingredient, and water.

[0007] The present disclosure is further directed to an extruded pesticide granule comprising from about 70wt.% to about 80wt.% urea pearls, from about 5wt.% to about 10wt.% of a non-ionic surfactant having a molecular weight of from about 14,000 to about 15,000 Daltons, from about 5wt.% to about 10wt.% pesticide active ingredient, and less than 3% water.

[0008] The present disclosure is further directed to a process for preparing an extruded pesticide granule. The process comprises melting a non-ionic surfactant, dissolving a pesticide active into the molten non-ionic surfactant, introducing powdered urea, adding water, and mixing to form a damp powder, extruding the damp powder to form an extrudate, and drying the extrudate to form the extruded pesticide granule.

[0009] It has been unexpectedly found that extruded pesticide granules that form highly desirable and stable near micro-emulsions in water can be suitably produced with one or more pesticide active agents in combination with a urea-based carrier material and a non-ionic surfactant. Surprisingly, by incorporating urea as the carrier material, the extruded pesticide granule has excellent dissolution/dispersion characteristics in water and forms a near micro-emulsion with only a very slight haze that is stable for many days upon mixing. Additionally, a suitable amount of the extruded pesticide granule can be incorporated into water in two minutes or less with only gentle agitation. Importantly, either water-soluble or water-insoluble, or a combination of both, pesticide active ingredients can be used in the disclosed extruded pesticide granules.

[0010] Additionally, the extruded pesticide granules and methods of the present disclosure provide numerous other advantages, including improved biological efficacy and lower dose rates for controlling pests; ability to be easily prepared to be organic compound-free; and ability to be easily transported to a work site and mixed with water onsite for easy and convenient application. Also, through the use of the specific components described herein for preparing the extruded pesticide granules, an approximate 66% reduction in the amount of water used during extrusion can be obtained without any negative impact on the resulting product. This has a positive environmental impact.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0011] The extruded pesticide granules of the present disclosure provide for an easily mixable pesticide granule that dissolves/suspends in water at suitable pesticide concentrations to form a near micro-emulsion that is stable for many days or even a week. The extruded pesticide granules include one or more pesticide active ingredients in

combination with a non-ionic surfactant and urea, optionally in the form or urea pearls. This unique combination of components is easily extruded to form the extruded pesticide granules that have numerous desirable characteristics and uses. In many embodiments of the present disclosure, the extruded pesticide granules are organic solvent-free extruded pesticide granules.

[0012] These and other optional elements or limitations of the extruded pesticide granules and methods of the present disclosure are described in detail hereinafter.

[0013] The term “pesticide” as used herein refers to at least one active substance selected from the group of the fungicides, insecticides, nematicides, herbicides, safeners, growth regulators and combinations thereof. Preferred pesticides for use within the scope of the present disclosure include fungicides, insecticides, and herbicides.

[0014] Numerical ranges as used herein are intended to include every number and subset of numbers within that range, whether specifically disclosed or not. Further, these numerical ranges should be construed as providing support for a claim directed to any number or subset of numbers in that range. For example, a disclosure of from 1 to 10 should be construed as supporting a range of from 2 to 8, from 3 to 7, from 5 to 6, from 1 to 9, from 3.6 to 4.6, from 3.5 to 9.9, and so forth.

[0015] All combinations of method or process steps as used herein can be performed in any order, unless otherwise specified or clearly implied to the contrary by the context in which the referenced combination is made.

[0016] The extruded pesticide granules and corresponding manufacturing methods and uses of the present disclosure can comprise, consist of, or consist essentially of the essential elements and limitations of the disclosure as described herein, as well as any additional or optional ingredients, components, or limitations described herein or otherwise useful in pesticide applications.

Pesticide Active Ingredient

[0017] The extruded pesticide granules described herein include at least one pesticide active ingredient. In some embodiments, the extruded pesticide granules will

include two, three or more pesticide active ingredients. Suitable pesticide active ingredients include both water-soluble (pesticide has a solubility in water of at least 10 g/L, preferably at least 25 g/L, and in particular at least 35 g/L) and water-insoluble (pesticide has a solubility in water of up to 10 g/L, including up to 2 g/L, and in particular up to 0.5 g/L at 20C) pesticide active ingredients, although the pesticide active ingredient should be substantially or completely soluble in the non-ionic surfactant described herein such that there is no milling required of the pesticide active ingredient. Insecticide active ingredients are particularly preferred within the scope of the present disclosure.

[0018] Suitable pesticide active ingredients can be found, for example, in the Pesticide Manual, 15th Ed. (2009), The British Crop Protection Council, London. Examples of pesticides may be selected from the following list (A to L are fungicides):

A) Respiration inhibitors

Inhibitors of complex III at Qo site (e.g. strobilurins): azoxystrobin, coumethoxystrobin, cou moxystrobin, dimoxystrobin, enestroburin, fenaminstrobin, fenoxy strobin/flufenoxystrobin, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, picoxy strobin, pyraclostrobin, pyrametostrobin, pyraoxystrobin, trifloxystrobin, 2-[2-(2,5-dimethyl-phenoxy methyl)-phenyl]-3-methoxy-acrylic acid methyl ester and 2-(2-(3-(2,6-dichlorophenyl)-1-methyl-allylideneamino oxymethyl)-phenyl)-2-methoxyimino-N-methyl-acetamide, pyribencarb, triclopyri carb/chlorodincarb, famoxadone, fenamidone;

Inhibitors of complex III at Qi site: cyazofamid, amisulbrom, [(3S,6S,7R,8R)-8-benzyl-3-[(3-acetoxy-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl]2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[(3-(acetoxymethoxy)-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[(3-isobutoxycarbonyloxy-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate, [(3S,6S,7R,8R)-8-benzyl-3-[(3-(1,3-benzodioxol-5-ylmethoxy)-4-methoxy-pyridine-2-carbonyl)amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl] 2-methylpropanoate;

Inhibitors of complex II (e. g. carboxamides): benodanil, bixafen, boscalid, carboxin, fenfuram, fluopyram, flutolanil, fluxapyroxad, furametpyr, isopyrazam, mepronil, oxycarboxin, penflufen, penthiopyrad, sedaxane, tecloftalam, thifluzamide, N-(4'-trifluoromethylthiobiphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, N-(2-(1,3,3-trimethyl-butyl)-phenyl)-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide, N-[9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide;

Other respiration inhibitors (e.g. complex I, uncouplers): diflumetorim, (5,8-difluoroquinazolin-4-yl)-{2-[2-fluoro-4-(4-trifluoromethylpyridin-2-yloxy)-phenyl]-ethyl}-amine; nitrophenyl derivates: binapacryl, dinobuton, dinocap, fluazinam; ferimzone; organometal compounds: fentinsalts, such as fentin-acetate, fentin chloride or fentin hydroxide; ametoctradin; and silthiofam;

B) Sterol biosynthesis inhibitors (SBI fungicides)

C14 demethylase inhibitors (DMI fungicides): triazoles: azaconazole, bitertanol, bromuconazole, cyeconazole, difenoconazole, diniconazole, diniconazole-M, epiconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imibenconazole, ipconazole, metconazole, myclobutanil, oxoconazole, paclobutrazole, penconazole, propiconazole, prothioconazole, simeconazole, tebuconazole, tetraconazole, triadimefon, triadimenol, triticonazole, uniconazole; imidazoles: imazalil, pefurazoate, prochloraz, triflumizol; pyrimidines, pyridines and piperazines: fenarimol, nuarimol, pyrifenoxy, triforine;

Delta14-reductase inhibitors: aldimorph, dodemorph, dodemorph-acetate, fenpropimorph, tridemorph, fenpropidin, piperalin, spiroxamine;

Inhibitors of 3-keto reductase: fenhexamid;

C) Nucleic acid synthesis inhibitors

Phenylamides or acyl amino acid fungicides: benalaxyl, benalaxyl-M, kiralaxyl, metalaxyl, metalaxyl-M (mefenoxam), ofurace, oxadixyl;

Others: hymexazole, octrilinone, oxolinic acid, bupirimate, 5-fluorocytosine, 5-fluoro-2-(ptolylmethoxy) pyrimidin-4-amine, 5-fluoro-2-(4-fluorophenylmethoxy)pyrimidin-4-amine;

D) Inhibitors of cell division and cytoskeleton

Tubulin inhibitors, such as benzimidazoles, thiophanates: benomyl, carbendazim, fuberidazole, thiabendazole, thiophanate-methyl; triazolopyrimidines: 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo[1,5-a]pyrimidine

Other cell division inhibitors: diethofencarb, ethaboxam, pencycuron, fluopicolide, zoxamide, metrafenone, pyriofenone;

E) Inhibitors of amino acid and protein synthesis

Methionine synthesis inhibitors (anilino-pyrimidines): cyprodinil, mepanipyrim, pyrimethanil;

Protein synthesis inhibitors: blasticidin-S, kasugamycin, kasugamycin hydrochloride-hydrate, mildiomycin, streptomycin, oxytetracyclin, polyoxine, validamycin A;

F) Signal transduction inhibitors

MAP / histidine kinase inhibitors: fluoroimid, iprodione, procymidone, vinclozolin, fenpiclonil, fludioxonil;

G protein inhibitors: quinoxyfen;

G) Lipid and membrane synthesis inhibitors

Phospholipid biosynthesis inhibitors: edifenphos, iprobenfos, pyrazophos, isoprothiolane;

Lipid peroxidation: dicloran, quintozene, tecnazene, tolclofos-methyl, biphenyl, chloroneb, etridiazole;

Phospholipid biosynthesis and cell wall deposition: dimethomorph, flumorph, mandipropamid, pyrimorph, benthiavalicarb, iprovalicarb, valifenalate and N-(1-(1-(4-cyano-phenyl)- ethanesulfonyl)-but-2-yl) carbamic acid-(4-fluorophenyl) ester;

Compounds affecting cell membrane permeability and fatty acids: propamocarb, propamocarb-hydrochlorid

H) Inhibitors with Multi Site Action

Inorganic active substances: Bordeaux mixture, copper acetate, copper hydroxide, copperoxychloride, basic copper sulfate, sulfur;

Thio- and dithiocarbamates: ferbam, mancozeb, maneb, metam, metiram, propineb, thiram, zineb, ziram;

Organochlorine compounds (e.g. phthalimides, sulfamides, chloronitriles): anilazine, chlorothalonil, captafol, captan, folpet, dichlofluanid, dichlorophen, flusulfamide, hexachlorobenzene, pentachlorphenole and its salts, phthalide, tolylfluanid, N-(4-chloro-2-nitro-phenyl)-Nethyl-4-methyl-benzenesulfonamide;

Guanidines and others: guanidine, dodine, dodine free base, guazatine, guazatine-acetate, iminoctadine, iminoctadine-triacetate, iminoctadine-tris(albesilate), dithianon;

I) Cell wall synthesis inhibitors

inhibitors of glucan synthesis: validamycin, polyoxin B; melanin synthesis inhibitors: pyroquilon, tricyclazole, carpropamid, dicyclomet, fenoxanil;

J) Plant defense inducers

Acibenzolar-S-methyl, probenazole, isotianil, tiadinil, prohexadione-calcium; phosphonates: fosetyl, fosetyl-aluminum, phosphorous acid and its salts;

K) Unknown mode of action

Bronopol, chinomethionat, cyflufenamid, cymoxanil, dazomet, debacarb, diclomezine, difenoquat, difenoquat-methylsulfate, diphenylamin, fenpyrazamine, flumetover,

flusulfamide, flutianil, methasulfocarb, nitrappyrin, nitrothal-isopropyl, oxin-copper, proquinazid, tebufloquin, tecloftalam, triazoxide, 2-butoxy-6-iodo-3-propylchromen-4-one, N-(cyclopropylmethoxyimino-(6-difluoro-methoxy-2,3-difluoro-phenyl)-methyl)-2-phenyl acetamide, N'-(4-(4-chloro-3-trifluoromethyl-phenoxy)-2,5-dimethyl-phenyl)-N-ethyl-N-methyl formamidine, N'-(4-(4-fluoro-3-trifluoromethyl-phenoxy)-2,5-dimethyl-phenyl)-N-ethyl-N-methyl formamidine, N'-(2-methyl-5-trifluoromethyl-4-(3-trimethylsilanyl-propoxy)-phenyl)-N-ethyl formamidine, N'-(5-difluoromethyl-2-methyl-4-(3-trimethylsilanyl-propoxy)-phenyl)-Nethyl formamidine, 2-{1-[2-(5-methyl-3-trifluoromethyl-pyrazole-1-yl)-acetyl]-piperidin-4-yl}-thiazole-4-carboxylic acid methyl-(1,2,3,4-tetrahydro-naphthalen-1-yl)-amide, 2-{1-[2-(5-methyl-3-trifluoromethyl-pyrazole-1-yl)-acetyl]-piperidin-4-yl}-thiazole-4-carboxylic acid methyl-(R)-1,2,3,4-tetrahydro-naphthalen-1-yl-amide, 1-[4-[4-[5-(2,6-difluorophenyl)-4,5-dihydro-3-isoxazolyl]-2-thiazolyl]-1-piperidinyl]-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, methoxy-acetic acid 6-tert-butyl-8-fluoro-2,3-dimethyl-quinolin-4-yl ester, N-Methyl-2-{1-[(5-methyl-3-trifluoro methyl-1H-pyrazol-1-yl)-acetyl]-piperidin-4-yl}-N-[(1R)-1,2,3,4-tetrahydronaphthalen-1-yl]-4-thiazolecarboxamide, 3-[5-(4-methylphenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine, 3-[5-(4-chloro-phenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine (pyrisoxazole), N-(6-methoxy-pyridin-3-yl) cyclopropanecarboxylic acid amide, 5-chloro-1-(4,6-dimethoxy-pyrimidin-2-yl)-2-methyl-1H-benzoimidazole, 2-(4-chlorophenyl)-N-[4-(3,4-dimethoxy-phenyl)-isoxazol-5-yl]-2-prop-2-ynyl-oxo-acetamide;

L) Antifungal biocontrol agents, plant bioactivators:

Ampelomyces quisqualis (e.g. AQ 10® from Intrachem Bio GmbH & Co. KG, Germany), Aspergillus flavus (e.g. AFLAGUARD® from Syngenta, CH), Aureobasidium pullulans (e.g. BOTECTOR® from bio-ferm GmbH, Germany), Bacillus pumilus (e.g. NRRL Accession No. B-30087 in SONATA® and BALLAD®

Plus from AgraQuest Inc., USA), Bacillus subtilis (e.g. isolate NRRL-Nr. B-21661 in RHAPSODY®, SERENADE® MAX and SERENADE® ASO from AgraQuest Inc., USA), Bacillus subtilis var. amyloliquefaciens FZB24 (e.g. TAEGRO® from Novozyme Biologicals, Inc., USA), Candida oleophila I-82 (e.g. ASPIRE® from Ecogen Inc., USA), Candida saitoana (e.g. BIOCURE® (in mixture with lysozyme) and BIOCOAT® from

Micro Flo Company, USA (BASF SE) and Arysta), Chitosan (e.g. ARMOUR-ZEN from BotriZen Ltd., NZ), Clonostachys rosea f. catenulata, also named Gliocladium catenulatum (e.g. isolate J1446:PRESTOP® from Verdera, Finland), Coniothyrium minitans (e.g. CONTANS® from Prophyta, Germany), Cryphonectria parasitica (e.g. Endothia parasitica from CNICM, France), Cryptococcus albidus (e.g. YIELD PLUS® from Anchor Bio-Technologies, South Africa), Fusarium oxysporum (e.g. BIOFOX® from S.I.A.P.A., Italy, FUSACLEAN® from Natural Plant Protection, France), Metschnikowia fructicola (e.g. SHEMER® from Agrogreen, Israel), Microdochium dimerum (e.g. ANTIBOT® from Agrauxine, France), Phlebiopsis gigantea (e.g. ROTSOP® from Verdera, Finland), Pseudozyma flocculosa (e.g. SPORODEX® from Plant Products Co. Ltd., Canada), Pythium oligandrum DV74 (e.g. POLYVERSUM® from Remeslo SSRO, Biopreparaty, Czech Rep.), Reynoutria sachalinensis (e.g. REGALIA® from Marrone BioInnovations, USA), Talaromyces flavus V117b (e.g. PROTUS® from Prophyta, Germany), Trichoderma asperellum SKT-1 (e.g. ECO-HOPE® from Kumiai Chemical Industry Co., Ltd., Japan), T. atroviride LC52 (e.g. SENTINEL® from Agrimm Technologies Ltd, NZ), T. harzianum T-22 (e.g. PLANTSHIELD® der Firma BioWorks Inc., USA), T. harzianum TH 35 (e.g. ROOT PRO® from Mycontrol Ltd., Israel), T. harzianum T-39 (e.g. TRICHODEX® and TRICHODERMA 2000® from Mycontrol Ltd., Israel and Makhteshim Ltd., Israel), T. harzianum and T. viride (e.g. TRICHOPEL from Agrimm Technologies Ltd, NZ), T. harzianum ICC012 and T. viride ICC080 (e.g. REMEDIER® WP from Isagro Ricerca, Italy), T. polysporum and T. harzianum (e.g. BINAB® from BINAB Bio-Innovation AB, Sweden), T. stromaticum (e.g. TRICOVAB® from C.E.P.L.A.C., Brazil), T. virens GL-21 (e.g. SOILGARD® from Certis LLC, USA), T. viride (e.g. TRIECO® from Ecosense Labs. (India) Pvt. Ltd., In dien, BIO-CURE® F from T. Stanes & Co. Ltd., Indien), T. viride TV1 (e.g. T. viride TV1 from Agribiotec srl, Italy), Ulocladium oudemansii HRU3 (e.g. BOTRY-ZEN® from Botry-Zen Ltd, NZ);

M) Growth regulators

Abscisic acid, amidochlor, ancyimidol, 6-benzylaminopurine, brassinolide, butralin, chlormequat (chlormequat chloride), choline chloride, cyclanilide, daminozide, dikegulac, dimethipin, 2,6-dimethylpuridine, ethephon, flumetralin, flurprimidol, fluthiacet, forchlorfenuron, gibberellic acid, inabenfide, indole-3-acetic acid, maleic hydrazide,

mefluidide, mepiquat (mepiquat chloride), naphthaleneacetic acid, N-6-benzyladenine, paclobutrazol, prohexadione (prohexadionecalcium), prohydrojasmon, thidiazuron, triapenthenol, tributyl phosphorotriethioate, 2,3,5-tri-iodobenzoic acid, trinexapac-ethyl and uniconazole;

N) Herbicides

Acetamides: acetochlor, alachlor, butachlor, dimethachlor, dimethenamid, flufenacet, mefenacet, metolachlor, metazachlor, napropamide, naproanilide, pethoxamid, pretilachlor, propachlor, thenylchlor;

Amino acid derivatives: bilanafos, glyphosate, glufosinate, sulfosate;

Aryloxyphenoxypropionates: clodinafop, cyhalofop-butyl, fenoxaprop, fluazifop, haloxyfop, metamifop, propaquizafop, quizalofop, quizalofop- P-tefuryl;

Bipyridyls: diquat, paraquat;

(thio)carbamates: asulam, butylate, carbetamide, desmedipham, dimepiperate, eptam (EPTC), esprocarb, molinate, orbencarb, phenmedipham, prosulfocarb, pyributicarb, thiobencarb, triallate;

Cyclohexanediones: butoxydim, clethodim, cycloxydim, profoxydim, sethoxydim, tepraloxymid, tralkoxydim;

Dinitroanilines: benfluralin, ethalfluralin, oryzalin, pendimethalin, prodiamine, trifluralin;

Diphenyl ethers: acifluorfen, aclonifen, bifenoxy, diclofop, ethoxyfen, fomesafen, lactofen, oxyfluorfen;

Hydroxybenzonitriles: bomoxynil, dichlobenil, ioxynil;

Imidazolinones: imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr;

Phenoxy acetic acids: clomeprop, 2,4-dichlorophenoxyacetic acid (2,4-D), 2,4-DB, dichlorprop, MCPA, MCPA-thioethyl, MCPB, Mecoprop;

Pyrazines: chloridazon, flufenpyr-ethyl, fluthiacet, norflurazon, pyridate;

Pyridines: aminopyralid, clopyralid, diflufenican, dithiopyr, fluridone, fluroxypyr, picloram, picolinafen, thiazopyr;

Sulfonyl ureas: amidosulfuron, azimsulfuron, bensulfuron, chlorimuron-ethyl, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethoxysulfuron, flazasulfuron, flucetosulfuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metazosulfuron, metsulfuron-methyl, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxsulfuron, triflusulfuron, tritosulfuron, 1-((2-chloro-6-propyl-imidazo[1,2-b]pyridazin-3-yl)sulfonyl)-3-(4,6-dimethoxy-pyrimidin-2-yl)urea;

Triazines: ametryn, atrazine, cyanazine, dimethametryn, ethiozin, hexazinone, metamitron, metribuzin, prometryn, simazine, terbutylazine, terbutryn, triaziflam;

Ureas: chlorotoluron, daimuron, diuron, fluometuron, isoproturon, linuron, methabenzthiazuron, tebuthiuron;

Other acetolactate synthase inhibitors: bispyribac-sodium, cloransulam-methyl, diclosulam, florasulam, flucarbazone, flumetsulam, metosulam, ortho-sulfamuron, penoxsulam, propoxy carbazole, pyribambenz-propyl, pyribenzoxim, pyriftalid, pyriminobac-methyl, pyrimisulfan, pyrithiobac, pyroxasulfone, pyroxulam;

Others: amicarbazone, aminotriazole, anilofos, beflubutamid, benazolin, bencarbazone, benfluresate, benzofenap, bentazone, benzobicyclon, bicyclopyrone, bromacil, bromobutide, butafenacil, butamifos, cafenstrole, carfentrazone, cinidon-ethyl, chlorthal, cinmethylin, clomazone, cumyluron, cyprosulfamide, dicamba, difenzoquat, diflufenopyr, Drechslera monoceras, endothal, ethofumesate, etobenzanid, fenoxasulfone, fentrazamide, flumiclorac-pentyl, flumioxazin, flupoxam, flurochloridone, flurtamone, indanofan, isoxaben, isoxaflutole, lenacil, propanil, propyzamide, quinclorac, quinmerac, mesotrione, methyl arsenic acid, naptalam, oxadiargyl, oxadiaxon, oxaziclofene, pentoxazone, pinoxaden, pyraclonil, pyraflufen-ethyl, pyrasulfotole, pyrazoxyfen, pyrazolynate,

quinoclamine, saflufenacil, sulcotrione, sulfentrazone, terbacil, tefuryltrione, tembotrione, thiencarbazone, topramezone, (3-[2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-5 3,6-dihydro-2Hpyrimidin-1-yl)-phenoxy]-pyridin-2-yloxy)-acetic acid ethyl ester, 6-amino-5-chloro-2-cyclopropyl-pyrimidine-4-carboxylic acid methyl ester, 6-chloro-3-(2-cyclopropyl-6-methylphenoxy)-pyridazin-4-ol, 4-amino-3-chloro-6-(4-chlorophenyl)-5-fluoro-pyridine-2-carboxylic acid, 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)-pyridine-2-carboxylic acid methyl ester, and 4-amino-3-chloro-6-(4-chloro-3-dimethylamino-2-fluoro-phenyl)-pyridine-2- carboxylic acid methyl ester.

O) Insecticides

Organo(thio)phosphates: acephate, azamethiphos, azinphos-methyl, chlorpyrifos, chlorpyrifos-methyl, chlorgenvinphos, diazinon, dichlorvos, dicrotophos, dimethoate, disulfoton, ethion, fenitrothion, fenthion, isoxathion, malathion, methamidophos, methidathion, methylparathion, mevinphos, monocrotophos, oxydemeton-methyl, paraoxon, parathion, phentoate, phosalone, phosmet, phosphamidon, phorate, phoxim, pirimiphos-methyl, profenofos, prothiofos, sulprophos, tetrachlorvinphos, terbufos, triazophos, trichlorfon;

Carbamates: alanycarb, aldicarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, fenoxy carb, furathiocarb, methiocarb, methomyl, oxamyl, pirimicarb, propoxur, thiodicarb, triazamate;

Pyrethroids: allethrin, bifenthrin, cyfluthrin, cyhalothrin, cyphenothrin, cypermethrin, alphacypermethrin, beta-cypermethrin, zeta-cypermethrin, deltamethrin, esfenvalerate, etofenprox, fenpropathrin, fenvalerate, imiprothrin, lambda-cyhalothrin, permethrin, prallethrin, pyrethrin I and II, resmethrin, silafluofen, tau-fluvalinate, tefluthrin, tetramethrin, tralomethrin, transfluthrin, profluthrin, dimefluthrin;

Insect growth regulators: a) chitin synthesis inhibitors: benzoylureas: chlorfluazuron, cyramazin, diflubenzuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, teflubenzuron, triflumuron; buprofezin, diofenolan, hexythiazox, etoxazole, clofentazine; b) ecdysone antagonists: halofenozide, methoxyfenozide, tebufenozide,

azadirachtin; c) juvenoids: pyriproxyfen, methoprene, fenoxy carb; d) lipid biosynthesis inhibitors: spirodiclofen, spiromesifen, spirotetramat;

Nicotinic receptor agonists/antagonists compounds: clothianidin, dinotefuran, flupyradifurone, imidacloprid, thiamethoxam, nitenpyram, acetamiprid, thiacloprid, 1-(2-chloro-thiazol-5-ylmethyl)-2-nitrimino-3,5-dimethyl-[1,3,5]triazinane;- GABA antagonist compounds: endosulfan, ethiprole, fipronil, vaniliprole, pyrafluprole, pyriproxyfen, 5-amino-1-(2,6-dichloro-4-methyl-phenyl)-4-sulfinamoyl-1H-pyrazole-3-carbothioic acid amide;

Macrocyclic lactone insecticides: abamectin, emamectin, milbemectin, lepimectin, spinosad, spinetoram;

Mitochondrial electron transport inhibitor (METI) I acaricides: fenazaquin, pyridaben, tebufenpyrad, tolfenpyrad, flufenim;

METI II and III compounds: acequinocyl, fluacyprin, hydramethylnon;

Uncouplers: chlorfenapyr;

Oxidative phosphorylation inhibitors: cyhexatin, diafenthiuron, fenbutatin oxide, propargite;

Moult disruptor compounds: cryomazine;

Mixed function oxidase inhibitors: piperonyl butoxide;

Sodium channel blockers: indoxacarb, metaflumizone;

Others: benclothiaz, bifenazate, cartap, flonicamid, pyridalyl, pymetrozine, sulfur, thiocyclam, flubendiamide, chlorantraniliprole, cyazypyr (HG-W86), cyenopyrafen, flupyrazofos, cyflumetofen, amidoflumet, imicyafos, bistrifluron, dinotefuran, and pyrifluquinazon.

[0019] Preferred pesticide active ingredients are pyraclostrobin, metconazole, alpha-cypermethrin, boscalid, dinotefuran, Chlorfenapyr, epoxiconazol and combinations thereof. In some embodiments, a preferred pesticide active ingredient will be a pesticide

active ingredient that is substantially, or wholly, soluble in the molten non-ionic surfactant described below.

[0020] The pesticide active ingredient may be present in the extruded pesticide granules in an amount of from about 1wt.% to about 25wt.%, including from about 5wt.% to about 25wt.%, including from about 10wt.% to about 25wt.%, including from about 10wt.% to about 20wt.%, including from about 15wt.% to about 20wt.%. In one specific embodiment, the pesticide active ingredient may be present in the extruded pesticide granules in an amount of about 20wt.%. In other specific embodiments, the pesticide active ingredient may be present in an amount of from about 1wt.% to about 50wt.%, including from about 10wt.% to about 50wt.%.

Non-Ionic Surfactant

[0021] The extruded pesticide granules of the present disclose additionally include at least one non-ionic surfactant, which acts as a lubricant during the extrusion process described herein and swells when it becomes wet. The non-ionic surfactant is desirably a non-ionic amphiphilic polyalkoxylate surfactant free of ionic groups that has a melting point of 55°C or more. The polyalkoxylate is amphiphilic, which usually means that it has surfactant properties and lowers the surface tension of water. Usually, the polyalkoxylate is obtainable by alkoxylation using alkyleneoxides, such as C2-C6-alkylene oxide, preferably ethylene oxide, propylene oxide, or butylene oxide. Examples of suitable polyalkoxylates are block polymers or compounds such as alcohols, alkylphenols, amines, amides, arylphenols, fatty acids or fatty acid esters which have been alkoxylated with 1 to 50 equivalents.

[0022] The non-ionic surfactant may have a melting point of at least 35°C, preferably at least 43°C, more preferably at least 48°C and in particular at least 50°C and in particular at least 55°C.

[0023] The non-ionic surfactant is usually substantially soluble in water at 20°C. Desirably, the solubility in water of the non-ionic surfactant is at least 3 wt%, more preferably at least 7 wt%, and in particular at least 10 wt%.

[0024] The molecular weight of the non-ionic surfactant is typically in the range of from about 5,000 to about 50,000 Daltons, desirably from about 2,000 to about 35,000 Daltons, and desirably from about 5,000 to about 20,000 Daltons. In one particular embodiment, the molecular weight of the non-ionic surfactant is from about 14,000 Daltons to about 15,000 Daltons.

[0025] The non-ionic surfactant is desirably a block polymer, which may contain a hydrophilic block and a hydrophobic block. Suitable block polymers are block polymers of the A-B or A-B-A type comprising blocks of polyethylene oxide and polypropylene oxide, or of the A-B-C type comprising alkanol, polyethylene oxide and polypropylene oxide. Desirably, the non-ionic surfactant is a block polymer comprising at least one polyethoxylate block and at least one poly-C3-C5-alkoxylate block (e.g. polypropoxylate or polybutoxylate). In particular, the non-ionic surfactant may be a triblock polymer of A-B-A type comprising a polyethoxylate type A block and a poly-C3-C5-alkoxylate block (preferably polypropoxylate) type B block.

[0026] One particularly preferred non-ionic surfactant for use in the extruded pesticide granules of the present disclosure is Pluronic F127 (BASF Germany).

[0027] The non-ionic surfactant may be present in the extruded pesticide granules in an amount of from about 5wt.% to about 20wt.%, including from about 5wt.% to about 10wt.%, including from about 7wt.% to about 10wt.%.

[0028] In some desirable embodiments, the weight ratio of non-ionic surfactant to pesticide active ingredient is about 1:1.

Solid Carrier

[0029] The extruded pesticide granules of the present disclosure include one or more solid carriers. Some suitable solid carriers or fillers are mineral earths, e.g. silicates, silica gels, talc, kaolins, limestone, lime, chalk, clays, dolomite, diatomaceous earth, bentonite, calcium sulfate, magnesium sulfate, magnesium oxide; polysaccharides, e.g. cellulose, methylcellulose, starch; fertilizers, e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate; products of vegetable origin, e.g. cereal meal, tree bark

meal, wood meal, nutshell meal; sugars, e.g. mono- or di saccharides, and mixtures thereof. Some useful solid carriers are mono- or di-saccharides, polysaccharides, and mixtures thereof.

[0030] A particularly suitable and desirable solid carrier is urea, which may optionally be in the form of urea powder or urea pearls (commercially available from Cole-Parmer or Acros Organics). Urea is highly soluble in water, and cools down upon dissolution in water to improve the overall solubility of components through its interaction with the non-ionic surfactant described above.

[0031] The carrier material, and in many desirable embodiments the urea carrier material, is generally present in the extruded pesticide granules in an amount of at least 50wt.%, including at least 60wt.%, including at least 70wt.%, including at least 80wt.%. In some embodiments, the urea carrier material may be present in the extruded pesticide granules in an amount of from about 50wt.% to about 95wt.%, including from about 70wt.% to about 90wt.%, including from about 70wt.% to about 80wt.%, and including about 80wt.%. In some other embodiments, the urea carrier material may be present in the extruded pesticide granules in an amount of from about 40wt.% to about 80wt.%.

Solvent

[0032] The extruded pesticide granules of the present disclosure generally include a small amount solvent that is utilized during the extrusion process. Generally, the vast majority of the solvent utilized in the extrusion process is removed in the drying of the granules step; however, it is generally not possible or always desirable to completely remove all of the solvent and, as such, the dried granules will contain a small trace of solvent in many embodiments.

[0033] Suitable solvents and liquid carriers for use in the extrusion processes described herein are water and organic solvents, such as mineral oil fractions of medium to high boiling point, e.g. kerosene, diesel oil; oils of vegetable or animal origin; aliphatic, cyclic and aromatic hydrocarbons, e. g. toluene, paraffin, tetrahydronaphthalene, alkylated naphthalenes; alcohols, e.g. ethanol, propanol, butanol, benzylalcohol, cyclohexanol; glycols; DMSO; ketones, e.g. cyclohexanone; esters, e.g. lactates, carbonates, fatty acid

esters, 15 gamma-butyrolactone; fatty acids; phosphonates; amines; amides, e.g. N-methylpyrrolidone, fatty acid dimethylamides; and mixtures thereof. A particularly suitable and desirable solvent is water. When water is used as the solvent, the resulting extruded pesticide granule is substantially or completely organic compound-free.

[0034] Typically, the extruded pesticide granules will include up to 5 wt% solvents and liquid carriers (e.g. water), desirably up to 3 wt%, and in particular up to 1wt%. In some embodiments, the extruded pesticide granules will include less than 0.8wt.%, including less than 0.5wt.%, including less than 0.2wt.% solvent (water).

Optional Components/Auxiliaries

[0035] The extruded pesticide granules may optionally comprise other components or auxiliaries to facilitate the manufacturing or processing characteristics of the granules, or otherwise improve one or more properties thereof. Some suitable auxiliaries are additional surfactants, dispersants, emulsifiers, wetters, adjuvants, solubilizers, penetration enhancers, protective colloids, adhesion agents, thickeners, humectants, repellents, attractants, feeding stimulants, compatibilizers, bactericides, anti-freezing agents, anti-foaming agents, colorants, tackifiers, effervescents, and binders.

[0036] Suitable effervescent is a combination of a hydrogen carbonate and an organic acid, such as a combination of citric acid and potassium hydrogencarbonate. Examples of the hydrogen carbonate include sodium hydrogen carbonate, potassium hydrogen carbonate and ammonium hydrogen carbonate. Examples of the organic acid include citric acid, succinic acid, malic acid, lactic acid, tartaric acid, fumaric acid and maleic acid. The organic acid is preferably used in an amount of 0.5 percent by weight to 20 percent by weight and, particularly, 1 percent by weight to 10 percent by weight based on the whole weight. The organic acid may be used alone or as a mixture of two or more of them. The hydrogencarbonate can be preferably used in an amount of 0.25 times to 2 times by molar ratio of the amount of the organic acid.

[0037] Suitable surfactants are surface-active compounds, such as anionic, cationic, nonionic and amphoteric surfactants, polyelectrolytes, and mixtures thereof. Preferred surfactants are anionic surfactants. Such surfactants can be used as emulsifier,

dispersant, solubilizer, wetter, penetration enhancer, protective colloid, or adjuvant. Examples of surfactants are listed in McCutcheon's, Vol.1: Emulsifiers & Detergents, McCutcheon's Directories, Glen Rock, USA, 2008 (International Ed. or North American Ed.).

[0038] Suitable anionic surfactants are alkali, alkaline earth or ammonium salts of sulfonates, sulfates, phosphates, carboxylates, and mixtures thereof. Examples of sulfonates are alkylarylsulfonates, diphenylsulfonates, alpha-olefin sulfonates, lignine sulfonates, sulfonates of fatty acids and oils, sulfonates of ethoxylated alkylphenols, sulfonates of alkoxylation arylphenols, sulfonates of condensed naphthalenes, sulfonates of dodecyl- and tridecylbenzenes, sulfonates of naphthalenes and alkyl naphthalenes, sulfosuccinates or sulfosuccinamates. Examples of sulfates are sulfates of fatty acids and oils, of ethoxylated alkylphenols, of alcohols, of ethoxylated alcohols, or of fatty acid esters. Examples of phosphates are phosphate esters. Examples of carboxylates are alkyl carboxylates, and carboxylated alcohol or alkylphenol ethoxylates.

[0039] Suitable nonionic surfactants are alkoxylates, N-substituted fatty acid amides, amine oxides, esters, sugar-based surfactants, polymeric surfactants, and mixtures thereof. Examples of alkoxylates are compounds such as alcohols, alkylphenols, amines, amides, arylphenols, fatty acids or fatty acid esters which have been alkoxylated with 1 to 50 equivalents. Ethylene oxide and/or propylene oxide may be employed for the alkoxylation, preferably ethylene oxide. Examples of N-substituted fatty acid amides are fatty acid glucamides or fatty acid alkanolamides. Examples of esters are fatty acid esters, glycerol esters or monoglycerides. Examples of sugar-based surfactants are sorbitans, ethoxylated sorbitans, sucrose and glucose esters or alkylpolyglucosides. Examples of polymeric surfactants are homo- or copolymers of vinylpyrrolidone, vinylalcohols, or vinylacetate.

[0040] Suitable cationic surfactants are quaternary surfactants, for example quaternary ammonium compounds with one or two hydrophobic groups, or salts of long-chain primary amines. Suitable amphoteric surfactants are alkylbetrains and imidazolines. Suitable polyelectrolytes are polyacids or polybases. Examples of polyacids are alkali salts

of polyacrylic acid or polyacid comb polymers. Examples of polybases are polyvinylamines or polyethyleneamines.

[0041] Suitable adjuvants are compounds, which have a negligible or even no pesticidal activity themselves, and which improve the biological performance of the compound I on the target. Examples are surfactants, mineral or vegetable oils, and other auxiliaries. Further examples are listed by Knowles, Adjuvants and additives, Agrow Reports DS256, T&F Informa UK, 2006, chapter 5.

[0042] Preferred adjuvants are non-ionic surfactants selected from alkoxylates. Examples of alkoxylates are compounds such as alcohols, alkylphenols, amines, amides, arylphenols, fatty acids or fatty acid esters which have been alkoxylated with 1 to 50 equivalents. Ethylene oxide and/or propylene oxide may be employed for the alkoxylation, preferably ethylene oxide. More preferred adjuvants are linear or branched, aliphatic C6-C20-alkanols, which have been alkoxylated with ethylene oxide and optionally with propylene oxide.

[0043] Suitable thickeners are polysaccharides (e.g. xanthan gum, carboxymethylcellulose), anorganic clays (organically modified or unmodified), polycarboxylate thickeners, and silicates.

[0044] Suitable bactericides are bronopol and isothiazolinone derivatives such as alkylisothiazolinones and benzisothiazolinones.

[0045] Suitable anti-freezing agents are ethylene glycol, propylene glycol, urea and glycerin.

[0046] Suitable anti-foaming agents are silicones, long chain alcohols, and salts of fatty acids.

Manufacture of Extruded Pesticide Granules

[0047] The extruded pesticide granules described herein and capable forming a near micro-emulsion upon introduction into water and gentle agitation, can be manufactured using a conventional low pressure extrusion apparatus, including a low

pressure dome extruder. In some embodiments, high pressure extruders may also be utilized within the scope of the present disclosure to produce suitable extruded pesticide granules. To prepare an extrudable mix for extrusion and subsequent drying, the surfactant may be first melted at elevated temperature to provide a molten surfactant composition. A suitable temperature may be, for example, about 90C. Into the molten surfactant is added the pesticide active ingredient, or pesticide active ingredient. The pesticide active ingredient (or active ingredients, if there is more than one) is mixed into the molten surfactant until the pesticide active ingredient is dissolved into the surfactant and a substantially homogeneous mixture is obtained.

[0048] Once a substantially homogeneous mixture of the molten surfactant and pesticide active ingredient is obtained, the carrier material (desirably urea in the form of powdered urea or urea pearls in many embodiments) is mixed in to prepare a substantially homogeneous mixture. Once this is completed, the desired amount of water is added and the resulting mixture thoroughly mixed and kneaded. Generally, from about 4 grams to about 10 grams, including from about 4 grams to about 8 grams, and desirably about 6 grams of water is added for each 100 grams of non-ionic surfactant, pesticide active ingredient, and carrier material (urea), to achieve the desired moisture content for extrusion of the mixture. This amount of water produces a damp powder that can then be suitably extruded through a low pressure dome extruder using a 1mm dome screen, or similar dome screen size depending upon the desired granule size. The resulting granules may then be dried using conventional means to produce the extruded pesticide granules that have a high solubility in water due to the formation of a near micro-emulsion.

[0049] These dried extruded pesticide granules can be used to prepare an aqueous tank mix, in which a pesticide in near micro-emulsion form wherein the particle size is generally below 1.0 micrometer, by introducing the extruded granules into water and applying gentle agitation. In many embodiments, the particle size will be less than 0.8 micrometers, or even less than 0.7 micrometers, or even less than 0.5 micrometers.

[0050] The mixing of the extruded pesticide granules and water may be done directly in a spraying apparatus containing a tank, or multiple tanks. The mixing may be done at a temperature of from about 5C to about 50C, desirably from about 10C to about

30C. The mixing may be done by adding the solid composition to the water already present in a tank and agitating.

[0051] The methods of the present disclosure further include methods of controlling phytopathogenic fungi and/or undesired plant growth and/or undesired attack by insects or mites and/or for regulating the growth of plants, where the extruded pesticide granules obtained by the methods as described herein are allowed to act upon the particular pests, their habitat or the plants to be protected from the particular pest, the soil and/or on undesired plants and/or the useful plants and/or their habitat.

[0052] A user may apply the extruded pesticide granules after preparing an aqueous tank mix usually from a pre-dosage device, a knapsack sprayer, a spray tank, a spray plane, or an irrigation system. In some embodiments, the agrochemical composition is made up with water, buffer, and/or further auxiliaries to the desired application concentration and the ready-to-use spray composition is obtained.

[0053] The disclosure furthermore relates to a method for controlling phytopathogenic fungi and/or undesired vegetation and/or undesired attack by insects or mites and/or for regulating the growth of plants, where the concentrate according to the invention or the emulsion according to the invention is allowed to act on the respective pests, their environment or on the crop plants to be protected from the respective pests, on the soil and/or on undesired plants and/or on the crop plants and/or their environment. In general, the therapeutic treatment of humans and animals is excluded from the method for controlling phytopathogenic fungi and/or undesired vegetation and/or undesired attack by insects or mites and/or for regulating the growth of plants.

[0054] When employed in crop protection, the application rates of the pesticides amount is from 0.001 to 2 kg per ha, preferably from 0.005 to 2 kg per ha, especially preferably from 0.05 to 0.9 kg per ha and in particular from 0.1 to 0.75 kg per ha, depending on the nature of the desired effect. In treatment of plant propagation materials such as seeds, e. g. by dusting, coating or drenching seed, amounts of active substance of from 0.1 to 1000 g, preferably from 1 to 1000 g, more preferably from 1 to 100 g and most preferably from 5 to 100 g, per 100 kg of plant propagation material (preferably seed) are

generally required. When used in the protection of materials or stored products, the amount of active substance applied depends on the kind of application area and on the desired effect. Amounts customarily applied in the protection of materials are 0.001 g to 2 kg, preferably 0.005 g to 1 kg, of active substance per cubic meter of treated material.

[0055] Various types of oils, wetters, adjuvants, fertilizers or micronutrients and further pesticides (for example herbicides, insecticides, fungicides, growth regulators, safeners) may be added to the emulsion in the form of a premix or optionally only shortly before use (tank mix). These agents can be admixed to the compositions according to the disclosure at a weight ratio of from 1:100 to 100:1, preferably from 1:10 to 10:1.

[0056] The user applies the composition according to the disclosure usually from a predosage device, a knapsack sprayer, a spray tank, a spray plane, or an irrigation system. Usually, the agrochemical composition is made up with water, buffer, and/or further auxiliaries to the desired application concentration and the ready-to-use spray liquor or the agrochemical composition according to the disclosure is thus obtained. Usually, 20 to 2000 liters, preferably 50 to 400 liters, of the ready-to-use spray liquor are applied per hectare of agricultural useful area.

[0057] The present embodiments are to be considered in all respects as illustrative and not restrictive and that all changes and equivalents also come within the description of the present disclosure. The following non-limiting examples will further illustrate the extruded pesticide compositions and methods of the present disclosure.

EXAMPLES

[0058] The following examples illustrate specific embodiments and/or features of the extruded pesticide granules of the present disclosure. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present disclosure, as many variations thereof are possible without departing from the spirit and scope of the disclosure. In addition to the specific components set forth in the Examples, each dried extruded pesticide granule described included a small amount of residual water, typically less than 2wt.%, and more typically less than 1wt.%.

Example 1

[0059] In this Example, extruded pesticide granules including one pesticide active agent suitable for preparing a near micro-emulsion in water were prepared.

[0060] 20.00 grams of Pluronic F-127 non-ionic surfactant was melted at 90°C to prepare a molten surfactant. To this molten surfactant was added 20.06 grams of alpha-cypermethrin and the resulting mixture stirred to provide a substantially homogeneous mixture of pesticide active and surfactant. To this substantially homogenous mixture was added 159.94 grams of urea (98% pearls) and the resulting blend kneaded well to produce a substantially homogenous mixture of pesticide active, surfactant, and urea. To this homogeneous mixture was added 12.0 grams of water and the resulting blend was mixed well. This produced a damp powder extrudable composition.

[0061] The damp powder was then extruded using a low pressure dome extruder having a 1mm dome screen at a temperature of about 81°F. The resulting granules were dried at room temperature on a glass plate. The resulting dried granules had a specific gravity of about 0.5, and a cylindrical shape (about 1.2mm x about 3mm). The granules were about 10.03wt.% pesticide active, about 10.0wt.% non-ionic surfactant, and about 79.97wt.% urea. The extruded granules can be added directly to water with gentle agitation to form a near micro-emulsion.

Example 2

[0062] In this Example, extruded pesticide granules including two pesticide active agents suitable for preparing a near micro-emulsion in water were prepared.

[0063] 20.00 grams of Pluronic F-127 non-ionic surfactant was melted at 80°C to prepare a molten surfactant. To this molten surfactant was added 20.24 grams of alpha-cypermethrin and the resulting mixture stirred to provide a substantially homogeneous mixture of pesticide active and surfactant. To this substantially homogenous mixture was added 119.31 grams of urea (98% pearls) and the resulting blend kneaded well until cool to produce a substantially homogeneous mixture of pesticide active, surfactant, and urea. To this homogeneous mixture was uniformly blended in 40.44 grams of dinotefuran. To this

uniformly blended mixture was added 6.0 grams of water and the resulting blend was mixed well. This produced a damp powder extrudable composition.

[0064] The damp powder was then extruded using a low pressure dome extruder having a 0.8 mm dome screen at a temperature of about 81°F. The resulting granules were dried at room temperature on a glass plate. The resulting dried granules had a specific gravity of about 0.5, and a cylindrical shape (about 1.2mm x about 3mm). The granules were about 10.12wt.% alpha cypermethrin, about 10.0wt.% non-ionic surfactant, about 20.22wt.% dinotefuran, and about 59.65wt.% urea. The extruded granules can be added directly to water with gentle agitation to form a near micro-emulsion.

Example 3

[0065] In this Example, extruded pesticide granules including one pesticide active agent suitable for preparing a near micro-emulsion in water were prepared.

[0066] 20.00 grams of Pluronic F-127 non-ionic surfactant was melted at 80°C to prepare a molten surfactant. To this molten surfactant was added 20.53 grams of chlорfenapyr and the resulting mixture stirred to provide a substantially homogeneous mixture of pesticide active and surfactant. To this substantially homogenous mixture was added 159.47 grams of urea (98% pearls) and the resulting blend kneaded well to produce a substantially homogeneous mixture of pesticide active, surfactant, and urea. To this homogeneous mixture was added 6.0 grams of water and the resulting blend was mixed well. This produced a damp powder extrudable composition.

[0067] The damp powder was then extruded using a low pressure dome extruder having a 0.8 mm dome screen at a temperature of about 81°F. The resulting granules were dried at room temperature on a glass plate. The resulting dried granules had a specific gravity of about 0.5, and a cylindrical shape (about 1.2mm x about 3mm). The granules were about 10.27wt.% pesticide active, about 10.0wt.% non-ionic surfactant, and about 79.73wt.% urea. The extruded granules can be added directly to water with gentle agitation to form a near micro-emulsion.

Example 4

[0068] In this Example, extruded pesticide granules including one pesticide active agent suitable for preparing a near micro-emulsion in water were prepared.

[0069] 40.00 grams of Pluronic F-127 non-ionic surfactant was melted at 80°C to prepare a molten surfactant. To this molten surfactant was added 41.67 grams of permethrin and the resulting mixture stirred to provide a substantially homogeneous mixture of pesticide active and surfactant. To this substantially homogenous mixture was added 118.33 grams of urea (98% pearls) and the resulting blend kneaded well to produce a substantially homogenous mixture of pesticide active, surfactant, and urea. To this homogeneous mixture was added 12.0 grams of water and the resulting blend was mixed well. This produced a damp powder extrudable composition.

[0070] The damp powder was then extruded using a low pressure dome extruder having a 1mm dome screen at a temperature of about 81°F. The resulting granules were dried at room temperature on a glass plate. The resulting dried granules had a specific gravity of about 0.5, and a cylindrical shape (about 1.2mm x about 3mm). The granules were about 20.83wt.% pesticide active, about 20.0wt.% non-ionic surfactant, and about 59.17wt.% urea. The extruded granules can be added directly to water with gentle agitation to form a near micro-emulsion.

Example 5

[0071] In this Example, extruded pesticide granules including one pesticide active agent suitable for preparing a near micro-emulsion in water were prepared.

[0072] 20.00 grams of Pluronic F-127 non-ionic surfactant was melted at 80°C to prepare a molten surfactant. To this molten surfactant was added 22.54 grams of fipronil and the resulting mixture stirred to provide a substantially homogeneous mixture of pesticide active and surfactant. To this substantially homogenous mixture was added 157.46 grams of urea (98% pearls) and the resulting blend kneaded well to produce a substantially homogenous mixture of pesticide active, surfactant, and urea. To this

homogeneous mixture was added 12.0 grams of water and the resulting blend was mixed well. This produced a damp powder extrudable composition.

[0073] The damp powder was then extruded using a low pressure dome extruder having a 0.8 mm dome screen at a temperature of about 81°F. The resulting granules were dried at room temperature on a glass plate. The resulting dried granules had a specific gravity of about 0.5, and a cylindrical shape (about 1.2mm x about 3mm). The granules were about 11.27wt.% pesticide active, about 10.0wt.% non-ionic surfactant, and about 78.73wt.% urea. The extruded granules can be added directly to water with gentle agitation to form a near micro-emulsion.

Example 6

[0074] In this Example, the biological residual efficacy of extruded pesticide granules of the present disclosure was tested and compared to the efficacy of Phantom Termiticide-Insecticide and a control. Specifically, extruded pesticide granules as prepared in Example 3 and including about 10wt.% chlорfenапyr and extruded pesticide granules as prepared in Example 3 and including about 20wt.% chlорfenапyr we evaluated and compared to the Phantom and control (control was no chemical treatment).

[0075] The pesticides were tested against German cockroaches (*Blattella germanica*), bed bugs (*Cimex lectularius*), confused flour beetles (*Tribolium castaneum*), and house crickets (*Acheta domesticus*). This testing was a residual test on unpainted plywood and painted plywood (Behr white semi-gloss exterior paint). The insects were confined on the treated surfaces with various cups depending on the insect. Four replications were used with ten insects for each replication. The knockdown and mortality were recorded over time, as shown in the following Tables 1-4. Each test formulation was diluted with water to a 0.5wt.% active level and applied with a DeVibiss atomizer set at 10 psi from 12 inches at the rate of 1 gallon of product per 1000 square feet. Both the extruded pesticide granules of the present disclosure tested form a near micro-emulsion in water. The tests were discontinued when the control mortality was above 20%.

Table 1
Percent Mortality of German Cockroaches at Time in Days

Product	Surface	Aged Day 1				Aged Days 28			
		1	2	3	5	1	2	3	5
Phantom SC	Unpainted	3.3	76.7	100.0	--	66.7	100.0	--	--
Chlorfenapyr 10% WMG		16.7	96.7	100.0	--	50.0	96.7	100.0	--
Chlorfenapyr 20% WMG		50.0	100.0	100.0	--	83.3	100.0	--	--
Control		3.3	6.7	10.0	--	0.0	0.0	3.3	--
Phantom SC	Painted	3.3	66.7	100.0	100.0	6.7	63.3	93.3	--
Chlorfenapyr 10% WMG		8.3	50.0	90.0	100.0	3.3	50.0	93.3	--
Chlorfenapyr 20% WMG		6.7	63.3	96.7	100.0	6.7	53.3	93.3	--
Control		0.0	3.3	3.3	10.0	0.0	6.7	10.0	66.7

Table 2
Percent Mortality of Bed Bugs at Time in Days

Product	Surface	Aged Day 1				Aged Days 28			
		1	2	3	5	7	1	2	3
Phantom SC	Unpainted	0.0	10.0	10.0	30.7	41.1	3.3	10.0	26.7
Chlorfenapyr 10% WMG		16.1	55.2	64.5	71.5	87.3	23.3	33.3	43.3
Chlorfenapyr 20% WMG		3.3	33.3	43.3	76.7	83.3	16.7	36.7	50.0
Control		0.0	0.0	3.3	3.3	6.7	0.0	3.3	16.7
Phantom SC	Painted	0.0	6.7	20.0	40.0	53.3	12.9	--	--
Chlorfenapyr 10% WMG		30.0	46.7	50.0	83.3	96.7	42.2	--	--
Chlorfenapyr 20% WMG		0.0	23.3	33.3	86.7	93.3	24.6	--	--
Control		0.0	0.0	3.3	6.7	16.7	23.3	26.7	--

Table 3
Percent Mortality of Confused Flour Beetles at Time in Days

Product	Surface	Aged Day 1				Aged Days 28			
		1	2	3	5	1	2	3	5
Phantom SC	Unpainted	33.3	90.0	--	--	83.3	100.0	--	--
Chlorfenapyr 10% WMG		58.7	95.8	--	--	96.7	100.0	--	--
Chlorfenapyr 20% WMG		83.6	100.0	--	--	93.3	100.0	--	--
Control		0.0	0.0	--	--	0.0	0.0	0.0	--
Phantom SC	Painted	26.7	90.0	--	--	20.0	93.3	93.3	100.0
Chlorfenapyr 10% WMG		12.4	70.9	--	--	50.0	86.7	90.0	100.0
Chlorfenapyr 20% WMG		37.0	96.7	--	--	46.7	96.7	96.7	100.0
Control		0.0	0.0	--	--	0.0	0.0	6.7	13.3

Table 4
Percent Mortality of House Crickets at Time in Days

Product	Surface	Aged Day 1				Aged Days 28			
		1	2	3	5	1	2	3	5
Phantom SC	Unpainted	70.0	90.0	--	--	70.8	95.0	--	--
Chlorfenapyr 10% WMG		90.0	100.0	--	--	51.7	90.0	--	--
Chlorfenapyr 20% WMG		100.0	100.0	--	--	95.0	100.0	--	--
Control		5.0	20.0	--	--	0.0	10.0	20.0	--
Phantom SC	Painted	90.0	100.0	--	--	20.0	85.0	85.0	--
Chlorfenapyr 10% WMG		91.7	100.0	--	--	90.8	100.0	--	-
Chlorfenapyr 20% WMG		90.0	100.0	--	--	39.2	100.0	--	--
Control		0.0	0.0	--	--	9.2	14.2	14.2	19.2

[0076] As shown in the Tables 1-4 above, for cockroaches all of the treatments were effective (>90%) of both surfaces at 1 and 28 days after treatment. For bed bugs, the bed bugs were not controlled (not >90%) by Phantom SC. Both of the extruded pesticide granule formulation of the present disclosure performed well at 1 day after treatment. For the confused flour beetles, all of the treatments worked on both surfaces at day 1, with the exception of the 10% extruded pesticide granule on painted wood; however at 28 days after treatment, all of the products worked well on both surfaces. For the house crickets, all of the treatments worked on both surfaces at day 1. At day 30, the extruded pesticide granule formulations worked well on both surfaces while the Phantom SC worked well on the unpainted wood and did not work well on the painted wood.

Example 7

[0077] In this Example, a contact test for biological efficacy of extruded pesticide granules of the present disclosure was tested and compared to the efficacy of Phantom Termiticide-Insecticide and a control. Specifically, extruded pesticide granules as prepared in Example 3 and including about 10wt.% chlorfenapyr and extruded pesticide granules as prepared in Example 3 and including about 20wt.% chlorfenapyr we evaluated and compared to the Phantom and control (control was no chemical treatment).

[0078] The pesticides were tested against German cockroaches (*Blattella germanica*) and bed bugs (*Cimex lectularius*). This testing was a contact test. The insects were confined in cups, treated and moved to clean cups. Four replications were used with ten insects for each replication. The knockdown and mortality were recorded over time as

shown in Tables 1-2. Each test formulation was diluted with water to a 0.25% or a 0.5wt.% active level and applied with a DeVibiss atomizer set at 10 psi from 12 inches at the rate of 1 gallon of product per 1000 square feet. Both the extruded pesticide granules of the present disclosure tested form a near micro-emulsion in water.

Table 1 Percent Mortality of German Cockroaches at Time				
Product	Rate	Hours		
		4	24	
Phantom SC	0.25 %	2.5	100.0	
Chlorfenapyr 10% WMG	0.25 %	0.0	100.0	
Chlorfenapyr 20% WMG	0.25 %	0.0	100.0	
Phantom SC	0.5%	7.5	97.5	
Chlorfenapyr 10% WMG	0.5%	22.5	100.0	
Chlorfenapyr 20% WMG	0.5%	12.5	100.0	
Control	--	0.0	0.0	

Table 2 Percent Mortality of Bed Bugs at Time							
Product	Rate	Reps	Hours		Days		
			4	24	2	3	5
Phantom SC	0.25 %	3	7.2	17.4	26.8	44.3	68.1
Chlorfenapyr 10% WMG	0.25 %	4	0.0	14.1	33.2	54.8	76.4
Chlorfenapyr 20% WMG	0.25 %	4	0.0	15.0	27.5	45.0	50.0
Control	--	1	0.0	0.0	0.0	0.0	0.0

[0079] As shown in Tables 1-2, for cockroaches all three of the product were effective (>90%) at both dilution rates at 24 hours after treatment. For bed bugs, the bed bugs were not controlled (not >90%) by the Phantom or the extruded pesticide granules. At 5 days after treatment the Phantom and the 10% extruded pesticide granule had the best control.

WHAT IS CLAIMED IS:

1. An extruded pesticide granular composition comprising urea, a non-ionic surfactant, a pesticide active ingredient, and water.
2. The extruded composition of claim 1 wherein the urea is in the form of urea pearls.
3. The extruded composition of any of claims 1-2 wherein the non-ionic surfactant is a block copolymer having a molecular weight of from about 10,000 to about 15,000 Daltons.
4. The extruded composition of any of claims 1-3 wherein the pesticide active ingredient is alpha cypermethrin.
5. The extruded composition of any of claims 1-4 wherein the pesticide active ingredient is dinotefuran.
6. The extruded composition of any of claims 1-5 wherein the pesticide active ingredient is water-soluble.
7. The extruded composition of any of claims 1-6 wherein the urea is present in the composition in an amount of from about 50wt.% to about 95wt.%.
8. The extruded composition of any of claims 1-7 wherein the urea is present in the composition in an amount of from about 70wt.% to about 80wt.%.
9. The extruded composition of any of claims 1-8 wherein the non-ionic surfactant is present in an amount of from about 5wt.% to about 20wt.%.
10. The extruded composition of any of claims 1-9 wherein the pesticide active ingredient is present in an amount of from about 5wt.% to about 20wt.%.
11. The extruded composition of any of claims 1-10 wherein the composition comprises less than 1% water.

12. The extruded composition of any of claims 1-11 wherein the weight ratio of non-ionic surfactant to pesticide active ingredient is from about 5:1 to about 1:1.25.

13. The extruded composition of any of claims 1-12 wherein the non-ionic surfactant is a block polymer comprising at least one polyethoxylate block and at least one poly-C3-C5-alkoxylate block.

14. The extruded composition of any of claims 1-13 wherein the non-ionic surfactant is a block polymer including a hydrophilic block and a hydrophobic block.

15. A method for controlling phytopathogenic fungi and/or undesired vegetation and/or undesired attack by insects or mites and/or for regulating the growth of plants, where the extruded pesticide granular composition of any of claims 1-13 is allowed to act on the respective pests, their environment or on the crop plants to be protected from the

5 respective pests, on the soil and/or on undesired plants and/or on the crop plants and/or their environment.

16. A process for preparing an extruded pesticide granule comprising melting a non-ionic surfactant, dissolving a pesticide active into the molten non-ionic surfactant, introducing urea, adding water, and mixing to form a damp powder, extruding the damp powder to form an extrudate, and drying the extrudate to form the extruded pesticide

5 granule.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/074447

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A01N25/12 A01N53/00 A01N43/36 A01N47/02 A01P7/00
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>DATABASE CA [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; 12 May 1984 (1984-05-12), KOMAKI, TOMOO ET AL: "Urea fertilizer containing agrochemicals", XP002696654, retrieved from STN Database accession no. 1978:1627 abstract & JP 52 030577 B (SUMITOMO CHEMICAL CO., LTD., JAPAN) 9 August 1977 (1977-08-09) -/-</p>	1-16

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
14 February 2014	24/02/2014
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Habermann, Jörg

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/074447

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	<p>-& "Urea formulations - contg polyoxyethylene surfactant for agrichemical use", WPI WORLD PATENT INFORMATION DERWENT, vol. 37, no. 72, 31 December 1977 (1977-12-31), XP002085209,</p> <p>-----</p>	
X	EP 0 501 798 A1 (DU PONT [US]) 2 September 1992 (1992-09-02)	1-16
Y	examples 1-4 page 19, line 23 - line 24 claims 1-4 page 18, line 30 - line 37	1-16
Y	----- US 4 289 525 A (PASARELA NUNZIO R ET AL) 15 September 1981 (1981-09-15) example 5	1-16
Y	----- CA 1 209 363 A1 (BASF WYANDOTTE CORP) 12 August 1986 (1986-08-12) examples 1-3 claim 9	1-16
Y	----- US 4 776 881 A (MOSTAFA MOHAMED A [US]) 11 October 1988 (1988-10-11) examples 1, 2	1-16
Y	----- EP 0 638 235 A1 (SHELL INT RESEARCH [NL]) 15 February 1995 (1995-02-15) examples 3-6 tables 2, 3	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2013/074447

Patent document cited in search report	Publication date	Patent family member(s)		Publication date	
EP 0501798	A1 02-09-1992	AT 154868 T AU 661686 B2 BR 9205703 A CA 2105399 A1 CN 1065181 A DE 69220662 D1 DE 69220662 T2 DK 0575489 T3 EP 0501798 A1 EP 0575489 A1 ES 2104907 T3 GR 3024900 T3 IE 920640 A1 IL 101098 A JP H06505493 A PL 170484 B1 RU 2096955 C1 TR 28893 A US 5474971 A WO 9215197 A1 ZA 9201526 A			15-07-1997 03-08-1995 17-05-1994 02-09-1992 14-10-1992 07-08-1997 02-01-1998 16-02-1998 02-09-1992 29-12-1993 16-10-1997 30-01-1998 09-09-1992 05-12-1996 23-06-1994 31-12-1996 27-11-1997 17-07-1997 12-12-1995 17-09-1992 30-08-1993
US 4289525	A 15-09-1981	AU 532784 B2 AU 5928480 A CA 1126047 A1 US 4289525 A		13-10-1983 22-01-1981 22-06-1982 15-09-1981	
CA 1209363	A1 12-08-1986	NONE			
US 4776881	A 11-10-1988	AU 599991 B2 AU 7473987 A BR 8703332 A CA 1303375 C NZ 220891 A US 4776881 A		02-08-1990 07-01-1988 15-03-1988 16-06-1992 26-02-1990 11-10-1988	
EP 0638235	A1 15-02-1995	AT 178187 T BR 9403172 A DE 69417490 D1 DE 69417490 T2 DK 0638235 T3 EP 0638235 A1 EP 0893058 A1 ES 2132285 T3 GR 3029839 T3 IL 110536 A JP 3556280 B2 JP H07145002 A KR 100343332 B1 SG 43112 A1 TW 453853 B US 5709871 A US 5837651 A		15-04-1999 11-04-1995 06-05-1999 23-09-1999 11-10-1999 15-02-1995 27-01-1999 16-08-1999 30-07-1999 27-12-1998 18-08-2004 06-06-1995 10-01-2005 17-10-1997 11-09-2001 20-01-1998 17-11-1998	