

(54) Title
Dual variable domain immunoglobulins and uses thereof

(51) International Patent Classification(s)
C07K 16/00 (2006.01) **A61P 37/04** (2006.01)
A61K 39/00 (2006.01) **A61P 37/06** (2006.01)
A61K 39/395 (2006.01) **C07K 16/46** (2006.01)
A61K 49/00 (2006.01) **C12N 1/00** (2006.01)
A61K 51/00 (2006.01) **C12N 1/15** (2006.01)
A61P 11/06 (2006.01) **C12N 1/19** (2006.01)
A61P 17/06 (2006.01) **C12N 1/21** (2006.01)
A61P 21/02 (2006.01) **C12N 5/10** (2006.01)
A61P 23/00 (2006.01) **C12N 9/96** (2006.01)
A61P 23/02 (2006.01) **C12N 15/62** (2006.01)
A61P 25/18 (2006.01) **C12N 15/63** (2006.01)
A61P 25/20 (2006.01) **C12P 21/02** (2006.01)
A61P 25/24 (2006.01) **C40B 30/04** (2006.01)
A61P 29/00 (2006.01) **G01N 33/53** (2006.01)
A61P 31/00 (2006.01)

(21) Application No: **2011293253** (22) Date of Filing: **2011.08.25**

(87) WIPO No: **WO12/027570**

(30) Priority Data

(31) Number (32) Date (33) Country
61/377,117 **2010.08.26** **US**

(43) Publication Date: **2012.03.01**

(44) Accepted Journal Date: **2014.12.11**

(71) Applicant(s)
AbbVie Inc.

(72) Inventor(s)
Ghayur, Tariq; Salfeld, Jochen G.; McPherson, Michael J.

(74) Agent / Attorney
Spruson & Ferguson, L 35 St Martins Tower 31 Market St, Sydney, NSW, 2000

(56) Related Art
WO 2010/006060 A2 (ABBOTT LABORATORIES) 14 January 2010
US 2009/0311253 A1 (GHAYUR, et al.) 17 December 2009
WO 2011/059755 A2 (ABBOTT LABORATORIES) 19 May 2011



(51) International Patent Classification:
A61K 39/00 (2006.01) *C07K 16/00* (2006.01)

(71) Applicant (for all designated States except US): **ABBOTT LABORATORIES** [US/US]; 100 Abbott Park Road, Abbott Park, Illinois 60064 (US).

(21) International Application Number:
 PCT/US2011/049147

(72) Inventors; and
 (75) Inventors/Applicants (for US only): **GHAYUR, Tariq** [US/US]; 1014 Washington Street, Holliston, Massachusetts 01746 (US). **SALFELD, Jochen, G.** [US/US]; 177 Old Westboro Road, North Grafton, Massachusetts 01536 (US). **MCPHERSON, Michael, J.** [US/US]; 287 Pillsbury Road, Ashby, Massachusetts 01431 (US).

(22) International Filing Date:
 25 August 2011 (25.08.2011)

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:
 61/377,117 26 August 2010 (26.08.2010) US

(74) Agent: **MCNEILL, Rebecca, M.**; Finnegan, Henderson, Farabow, Garrett & Dummer, L.L.P., 901 New York Avenue, NW, Washington, D.C. 20001-4413 (US).

[Continued on next page]

(54) Title: DUAL VARIABLE DOMAIN IMMUNOGLOBULINS AND USES THEREOF

Figure 1



(57) Abstract: Engineered multivalent and multispecific binding proteins, methods of making, and their uses in the prevention, diagnosis, and/or treatment of disease are provided.



(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,

ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

— with amended claims (Art. 19(1))

(88) Date of publication of the international search report:

10 May 2012

Date of publication of the amended claims:

19 July 2012

DUAL VARIABLE DOMAIN IMMUNOGLOBULINS AND USES THEREOF**Cross Reference to Related Applications**

This application is a non-provisional application claiming priority to U.S. Provisional Application Serial No. 61/377,117, filed August 26, 2010, the entire content of which is hereby
5 incorporated by reference.

Field

Multivalent and multispecific binding proteins, methods of making, and specifically to their uses in the, diagnosis, prevention and/or treatment of acute and chronic inflammatory diseases, cancer, and other diseases are provided.

10

Background

Engineered proteins, such as multispecific antibodies capable of binding two or more antigens are known in the art. Such multispecific binding proteins can be generated using cell fusion, chemical conjugation, or recombinant DNA techniques.

Bispecific antibodies have been produced using quadroma technology (see Milstein, C. and A.C. Cuello (1983) Nature 305(5934):537-40) based on the somatic fusion of two different
15 hybridoma cell lines expressing murine monoclonal antibodies (mAbs) with the desired specificities of the bispecific antibody. Because of the random pairing of two different immunoglobulin (Ig) heavy and light chains within the resulting hybrid-hybridoma (or quadroma) cell line, up to ten different Ig species are generated, of which only one is the
20 functional bispecific antibody. The presence of mis-paired by-products, and significantly reduced production yields, means sophisticated purification procedures are required.

Bispecific antibodies can also be produced by chemical conjugation of two different mAbs (see Staerz, U.D., et al. (1985) Nature 314(6012): 628-31). This approach does not yield
25 homogeneous preparation. Other approaches have used chemical conjugation of two different mAbs or smaller antibody fragments (see Brennan, M., et al. (1985) Science 229(4708): 81-3).

Another method used to produce bispecific antibodies is the coupling of two parental antibodies with a hetero-bifunctional crosslinker, but the resulting bispecific antibodies suffer from significant molecular heterogeneity because reaction of the crosslinker with the parental
30 antibodies is not site-directed. To obtain more homogeneous preparations of bispecific antibodies two different Fab fragments have been chemically crosslinked at their hinge cysteine residues in

a site-directed manner (see Glennie, M.J., et al. (1987) *J. Immunol.* 139(7): 2367-75). But this method results in Fab'2 fragments, not full IgG molecule.

A wide variety of other recombinant bispecific antibody formats have been developed (see Kriangkum, J., et al. (2001) *Biomol. Eng.* 18(2): 31-40). Amongst them tandem single-chain Fv molecules and diabodies, and various derivatives thereof, are the most widely used. Routinely, construction of these molecules starts from two single-chain Fv (scFv) fragments that recognize different antigens (see Economides, A.N., et al. (2003) *Nat. Med.* 9(1): 47-52). Tandem scFv molecules (taFv) represent a straightforward format simply connecting the two scFv molecules with an additional peptide linker. The two scFv fragments present in these tandem scFv molecules form separate folding entities. Various linkers can be used to connect the two scFv fragments and linkers with a length of up to 63 residues (see Nakanishi, K., et al. (2001) *Ann. Rev. Immunol.* 19: 423-74). Although the parental scFv fragments can normally be expressed in soluble form in bacteria, it is, however, often observed that tandem scFv molecules form insoluble aggregates in bacteria. Hence, refolding protocols or the use of mammalian expression systems are routinely applied to produce soluble tandem scFv molecules. In a recent study, *in vivo* expression by transgenic rabbits and cattle of a tandem scFv directed against CD28 and a melanoma-associated proteoglycan was reported (see Gracie, J.A., et al. (1999) *J. Clin. Invest.* 104(10): 1393-401). In this construct, the two scFv molecules were connected by a CH1 linker and serum concentrations of up to 100 mg/L of the bispecific antibody were found. Various strategies including variations of the domain order or using middle linkers with varying length or flexibility were employed to allow soluble expression in bacteria. A few studies have now reported expression of soluble tandem scFv molecules in bacteria (see Leung, B.P., et al. (2000) *J. Immunol.* 164(12): 6495-502; Ito, A., et al. (2003) *J. Immunol.* 170(9): 4802-9; Karni, A., et al. (2002) *J. Neuroimmunol.* 125(1-2): 134-40) using either a very short Ala3 linker or long glycine/serine-rich linkers. In another recent study, phage display of a tandem scFv repertoire containing randomized middle linkers with a length of 3 or 6 residues was employed to enrich for those molecules that are produced in soluble and active form in bacteria. This approach resulted in the isolation of a tandem scFv molecule with a 6 amino acid residue linker (see Arndt, M. and J. Krauss (2003) *Methods Mol. Biol.* 207: 305-21). It is unclear whether this linker sequence represents a general solution to the soluble expression of tandem scFv molecules. Nevertheless, this study demonstrated that phage display of tandem scFv molecules in combination with directed mutagenesis is a powerful tool to enrich for these molecules, which can be expressed in bacteria in an active form.

Bispecific diabodies (Db) utilize the diabody format for expression. Diabodies are produced from scFv fragments by reducing the length of the linker connecting the VH and VL domain to approximately 5 residues (see Peipp, M. and T. Valerius (2002) Biochem. Soc. Trans. 30(4): 507-11). This reduction of linker size facilitates dimerization of two polypeptide chains by crossover pairing of the VH and VL domains. Bispecific diabodies are produced by expressing, 5 two polypeptide chains with, either the structure VHA-VLB and VHB-VLA (VH-VL configuration), or VLA-VHB and VLB-VHA (VL-VH configuration) within the same cell. A large variety of different bispecific diabodies have been produced in the past and most of them are expressed in soluble form in bacteria. However, a recent comparative study demonstrates that 10 the orientation of the variable domains can influence expression and formation of active binding sites (see Mack, M. et al.(1995) Proc. Natl. Acad. Sci. U S A 92(15): 7021-5). Nevertheless, soluble expression in bacteria represents an important advantage over tandem scFv molecules. However, since two different polypeptide chains are expressed within a single cell inactive homodimers can be produced together with active heterodimers. This necessitates the 15 implementation of additional purification steps in order to obtain homogenous preparations of bispecific diabodies. One approach to force the generation of bispecific diabodies is the production of knob-into-hole diabodies (see Holliger, P., T. Prospero, and G. Winter (1993) Proc. Natl. Acad. Sci. U S A 90(14): 6444-8.18). This approach was demonstrated for a bispecific diabody directed against HER2 and CD3. A large knob was introduced in the VH domain by 20 exchanging Val37 with Phe and Leu45 with Trp and a complementary hole was produced in the VL domain by mutating Phe98 to Met and Tyr87 to Ala, either in the anti-HER2 or the anti-CD3 variable domains. By using this approach the production of bispecific diabodies could be increased from 72% by the parental diabody to over 90% by the knob-into-hole diabody. Importantly, production yields only slightly decrease as a result of these mutations. However, a 25 reduction in antigen-binding activity was observed for several constructs. Thus, this rather elaborate approach requires the analysis of various constructs in order to identify those mutations that produce heterodimeric molecule with unaltered binding activity. In addition, such approach requires mutational modification of the immunoglobulin sequence at the constant region, thus creating non-native and non-natural form of the antibody sequence, which may result in increased 30 immunogenicity, poor *in vivo* stability, as well as undesirable pharmacokinetics.

Single-chain diabodies (scDb) represent an alternative strategy for improving the formation of bispecific diabody-like molecules (see Holliger, P. and G. Winter (1997) Cancer Immunol. Immunother. 45(3-4): 128-30; Wu, A.M., et al. (1996) Immunotechnology 2(1): p. 21-36). Bispecific single-chain diabodies are produced by connecting the two diabody-forming 35 polypeptide chains with an additional middle linker with a length of approximately 15 amino acid

residues. Consequently, all molecules with a molecular weight corresponding to monomeric single-chain diabodies (50-60 kDa) are bispecific. Several studies have demonstrated that bispecific single chain diabodies are expressed in bacteria in soluble and active form with the majority of purified molecules present as monomers (see Holliger, P. and G. Winter (1997) *Cancer Immunol. Immunother.* 45(3-4): 128-30; Wu, A.M., et al. (1996) *Immunotechnol.* 2(1): 21-36; Pluckthun, A. and P. Pack (1997) *Immunotechnol.* 3(2): 83-105; Ridgway, J.B., et al. (1996) *Protein Engin.* 9(7): 617-21). Thus, single-chain diabodies combine the advantages of tandem scFvs (all monomers are bispecific) and diabodies (soluble expression in bacteria).

More recently diabodies have been fused to Fc to generate more Ig-like molecules, named di-diabodies (see Lu, D., et al. (2004) *J. Biol. Chem.* 279(4): 2856-65). In addition, multivalent antibody construct comprising two Fab repeats in the heavy chain of an IgG and capable of binding four antigen molecules has been described (see WO 0177342A1, and Miller, K., et al. (2003) *J. Immunol.* 170(9): 4854-61).

There is a need in the art for improved multivalent binding proteins capable of binding two or more antigens. U.S. Patent Application Serial No. 1 1/507,050 provides a novel family of binding proteins capable of binding two or more antigens with high affinity, which are called dual variable domain immunoglobulins (DVD-Ig™). Novel binding proteins capable of binding two or more antigens are also provided.

Summary

Multivalent binding proteins capable of binding two or more antigens are provided. A novel family of binding proteins capable of binding two or more antigens with high affinity are also provided.

In an embodiment the invention provides a binding protein comprising first and second polypeptide chains, each independently comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein

- VD1 is a first variable domain;
- VD2 is a second variable domain;
- C is a constant domain;
- X1 is a linker;
- X2 is an Fc region;
- n is 0 or 1;

wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second

polypeptide chains form a second functional target binding site, and wherein the binding protein is capable of binding Tumor Necrosis Factor (TNF) and Nerve Growth Factor (NGF), wherein the binding protein comprises any one of:

DVD1459 (comprising SEQ ID NOs: 52 and 53); DVD1460 (comprising SEQ ID NOs: 54 and 55); DVD1461 (comprising SEQ ID NOs: 56 and 57); DVD1462 (comprising SEQ ID NOs: 58 and 59); DVD1463 (comprising SEQ ID NOs: 60 and 61); DVD1464 (comprising SEQ ID NOs: 62 and 63); DVD1477 (comprising SEQ ID NOs: 64 and 65); DVD1478 (comprising SEQ ID NOs: 66 and 67); DVD1485 (comprising SEQ ID NOs: 68 and 69); DVD1486 (comprising SEQ ID NOs: 70 and 71); DVD1501 (comprising SEQ ID NOs: 136 and 137); DVD1502 (comprising SEQ ID NOs: 138 and 139); DVD1503 (comprising SEQ ID NOs: 140 and 141); DVD1504 (comprising SEQ ID NOs: 142 and 143); DVD1505 (comprising SEQ ID NOs: 144 and 145); DVD1506 (comprising SEQ ID NOs: 146 and 147); DVD1519 (comprising SEQ ID NOs: 148 and 149); DVD1520 (comprising SEQ ID NOs: 150 and 151); DVD1527 (comprising SEQ ID NOs: 152 and 153); DVD1528 (comprising SEQ ID NOs: 154 and 155); DVD1543 (comprising SEQ ID NOs: 220 and 221); DVD1544 (comprising SEQ ID NOs: 222 and 223); DVD1545 (comprising SEQ ID NOs: 224 and 225); DVD1546 (comprising SEQ ID NOs: 226 and 227); DVD1547 (comprising SEQ ID NOs: 228 and 229); DVD1548 (comprising SEQ ID NOs: 230 and 231); DVD1561 (comprising SEQ ID NOs: 232 and 233); DVD1562 (comprising SEQ ID NOs: 234 and 235); DVD1569 (comprising SEQ ID NOs: 236 and 237); and DVD1570 (comprising SEQ ID NOs: 238 and 239).

In one embodiment a binding protein comprising a polypeptide chain, wherein the polypeptide chain comprises VD1 -(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first variable domain, VD2 is a second variable domain, C is a constant domain, XI represents an amino acid or polypeptide, X2 represents an Fc region and n is 0 or 1 is provided. In an embodiment the VD1 and VD2 in the binding protein are heavy chain variable domains. In another embodiment, the heavy chain variable domain is a murine heavy chain variable domain, a human heavy chain variable domain, a CDR grafted heavy chain variable domain, or a humanized heavy chain variable domain. In yet another, embodiment VD1 and VD2 are capable of binding the same antigen. In another embodiment VD1 and VD2 are capable of binding different antigens. In still another embodiment, C is a heavy chain constant domain. For example, XI is a linker with the proviso that XI is not CHI. For example, XI is AKTTPKLEEGEFSEAR (SEQ ID NO: 1);

AKTTPKLEEGEFSEARV (SEQ ID NO: 2); AKTTPKLG (SEQ ID NO: 3); SAKTTPKLG (SEQ ID NO: 4); SAKTTP (SEQ ID NO: 5); RADAAP (SEQ ID NO: 6); RADAAPT (SEQ ID NO: 7); RADAAGGPGS (SEQ ID NO: 8); RADAAGGPGS₄ (SEQ ID NO: 9); SAKTTPKLEEGEFSEARV (SEQ ID NO: 10); ADAAP (SEQ ID NO: 11); ADAAPT (SEQ ID NO: 12); TVAAP (SEQ ID NO: 13); TVAAPSVFIFPP (SEQ ID NO: 14); QPKAAP (SEQ ID NO: 15); QPKAAPSVTLFPP (SEQ ID NO: 16); AKTTP (SEQ ID NO: 17); AKTTPSVTLFPP (SEQ ID NO: 18); AKTTAP (SEQ ID NO: 19); AKTTAPSVYPLAP (SEQ ID NO: 20); ASTKGP (SEQ ID NO: 21); ASTKGPSVFPLAP (SEQ ID NO: 22); GGGGSGGGGSGGGGS (SEQ ID NO: 23); GENKVEYAPALMALS (SEQ ID NO: 24); GPAKELTPLKEAKVS (SEQ ID NO: 25); or GHEAAAVMQVQYPAS (SEQ ID NO: 26) In an embodiment, X2 is an Fc region. In another embodiment, X2 is a variant Fc region.

In an embodiment the binding protein disclosed herein comprises a polypeptide chain, wherein the polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region.

In an embodiment, VD1 and VD2 in the binding protein are light chain variable domains. In an embodiment, the light chain variable domain is a murine light chain variable domain, a human light chain variable domain, a CDR grafted light chain variable domain, or a humanized light chain variable domain. In one embodiment VD1 and VD2 are capable of binding the same antigen. In another embodiment VD1 and VD2 are capable of binding different antigens. In an embodiment, C is a light chain constant domain. In another embodiment, X1 is a linker with the proviso that X1 is not CL. In an embodiment, X1 is AKTTPKLEEGEFSEAR (SEQ ID NO: 1); AKTTPKLEEGEFSEARV (SEQ ID NO: 2); AKTTPKLG (SEQ ID NO: 3); SAKTTPKLG (SEQ ID NO: 4); SAKTTP (SEQ ID NO: 5); RADAAP (SEQ ID NO: 6); RADAAPT (SEQ ID NO: 7); RADAAGGPGS (SEQ ID NO: 8); RADAAGGPGS₄ (SEQ ID NO: 9); SAKTTPKLEEGEFSEARV (SEQ ID NO: 10); ADAAP (SEQ ID NO: 11); ADAAPT (SEQ ID NO: 12); TVAAP (SEQ ID NO: 13); TVAAPSVFIFPP (SEQ ID NO: 14); QPKAAP (SEQ ID NO: 15); QPKAAPSVTLFPP (SEQ ID NO: 16); AKTTP (SEQ ID NO: 17); AKTTPSVTLFPP (SEQ ID NO: 18); AKTTAP (SEQ ID NO: 19); AKTTAPSVYPLAP (SEQ ID NO: 20); ASTKGP (SEQ ID NO: 21); ASTKGPSVFPLAP (SEQ ID NO: 22); GGGGSGGGGSGGGGS (SEQ ID NO: 23); GENKVEYAPALMALS (SEQ ID NO: 24); GPAKELTPLKEAKVS (SEQ ID NO: 25); or GHEAAAVMQVQYPAS (SEQ ID NO: 26). In an embodiment, the binding protein does not comprise X2.

In an embodiment, both the variable heavy and variable light chain comprise the same linker. In another embodiment, the variable heavy and variable light chain comprise different linkers. In another embodiment, both the variable heavy and variable light chain comprise a short (about 6 amino acids) linker. In another embodiment, both the variable heavy and variable light chain comprise a long (greater than 6 amino acids) linker. In another embodiment, the variable heavy chain comprises a short linker and the variable light chain comprises a long linker. In another embodiment, the variable heavy chain comprises a long linker and the variable light chain comprises a short linker.

In an embodiment the binding protein disclosed herein comprises a polypeptide chain, wherein said polypeptide chain comprises $VD1-(X1)_n-VD2-C-(X2)_n$, wherein VD1 is a first light chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant domain, X1 is a linker with the proviso that it is not CL, and X2 does not comprise an Fc region.

In another embodiment a binding protein comprising two polypeptide chains, wherein said first polypeptide chain comprises $VD1-(X1)_n-VD2-C-(X2)_n$, wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region; and said second polypeptide chain comprises $VD1-(X1)_n-VD2-C-(X2)_n$, wherein VD1 is a first light chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant domain, X1 is a linker (optionally which is not CL), and X2 does not comprise an Fc region is provided. In a particular embodiment, the binding protein (i.e., Dual Variable Domain immunoglobulin (DVD-Ig)) comprises four polypeptide chains wherein the first two polypeptide chains comprises $VD1-(X1)_n-VD2-C-(X2)_n$, respectively wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region; and the second two polypeptide chain comprises $VD1-(X1)_n-VD2-C-(X2)_n$ respectively, wherein VD1 is a first light chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant domain, X1 is a linker (optionally which is not CL), and X2 does not comprise an Fc region. Such a binding protein has four antigen binding sites.

In another embodiment the binding proteins disclosed herein are capable of binding one or more targets. Accordingly, in some embodiments, the binding proteins comprise at least two variable domain sequences (e.g., VD1 and VD2) capable of binding at least two different targets. In some embodiments, VD1 and VD2 are independently chosen. Therefore, in some embodiments, VD1 and VD2 comprise the same SEQ ID NO and, in other embodiments, VD1 and VD2 comprise different SEQ ID NOS.

In another embodiment the binding proteins disclosed herein bind one or more targets. In an embodiment, the target is a cytokine, a cell surface protein, an enzymes, or a receptor. In another embodiment, the binding protein is capable of modulating a biological function of one or more targets. In another embodiment, the binding protein is capable of neutralizing one or more targets. In one embodiment, the binding protein is capable of binding cytokines. In some 5 embodiments, the cytokines are lymphokines, monokines, polypeptide hormones, receptors, or tumor markers. In some embodiments, the DVD-Ig is capable of binding two or more of the following: Tumor Necrosis Factor (TNF), Nerve Growth Factor (NGF), Sclerostin (SOST), Prostaglandin E2 (PGE2), Lysophosphatidic Acid (LPA) (see also Table 2). In a specific 10 embodiment the binding protein is capable of binding TNF (seq. 2) and NGF; TNF (seq. 3) and NGF; TNF (seq. 4) and NGF; TNF (seq. 5) and NGF; TNF (seq. 6) and NGF; TNF (seq. 2) and SOST; TNF (seq. 3) and SOST; TNF (seq. 4) and SOST; TNF (seq. 5) and SOST; TNF (seq. 6) and SOST; TNF (seq. 2) and PGE2; TNF (seq. 3) and PGE2; TNF (seq. 4) and PGE2; TNF (seq. 5) and PGE2; TNF (seq. 6) and PGE2; TNF (seq. 1) and LPA; TNF (seq. 2) and LPA; TNF (seq. 3) and LPA; TNF (seq. 4) and LPA; TNF (seq. 5) and LPA; or TNF (seq. 6) and LPA. In an 15 embodiment, the DVD-Ig contains one or more VH chains that comprise a short linker and one or more VL chains that contain a long linker. In an embodiment, the DVD-Ig contains one or more VH chains that comprise a long linker and one or more VL chains that contain a short linker. In an embodiment, the -Ig contains one or more VH chains that comprise a long linker and one or 20 more VL chains that contain a long linker.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 52 or SEQ ID NO. 54; and a light chain amino acid sequence of SEQ ID NO. 53 or SEQ ID NO. 55. In an embodiment, the binding protein capable of binding TNF (seq. 2) and NGF comprises a heavy chain amino acid sequence 25 of SEQ ID NO. 52 and a light chain amino acid sequence of SEQ ID NO. 53. In another embodiment, the binding protein capable of binding TNF (seq. 2) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 54 and a light chain amino acid sequence of SEQ ID NO. 55.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and NGF 30 comprises a heavy chain amino acid sequence of SEQ ID NO. 56 or SEQ ID NO. 58; and a light chain amino acid sequence of SEQ ID NO. 57 or SEQ ID NO. 59. In an embodiment, the binding protein capable of binding TNF (seq. 3) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 56 and a light chain amino acid sequence of SEQ ID NO. 57. In another embodiment, the binding protein capable of binding TNF (seq. 3) and NGF has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 58 and a light chain amino acid sequence of SEQ ID NO: 59.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and NGF comprises a heavy chain amino acid of SEQ ID NO. 60 or SEQ ID NO.62; and a light chain amino acid sequence of SEQ ID NO. 61 or SEQ ID NO.63. In an embodiment, the binding protein capable of binding TNF (seq. 4) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 60 and a light chain amino acid sequence of SEQ ID NO:61. In another embodiment, the binding protein capable of binding TNF (seq. 4) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 62 and a light chain amino acid sequence of SEQ ID NO: 63.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 64 or SEQ ID NO. 66; and a light chain amino acid sequence of SEQ ID NO. 65 or SEQ ID NO. 67. In an embodiment, the binding protein capable of binding TNF (seq. 5) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 64 and a light chain amino acid sequence of SEQ ID NO: 65. In another embodiment, the binding protein capable of binding TNF (seq. 5) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 66 and a light chain amino acid sequence of SEQ ID NO: 67.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 68 or SEQ ID NO. 70; and a light chain amino acid sequence of SEQ ID NO. 69 or SEQ ID NO. 71. In an embodiment, the binding protein capable of binding TNF (seq. 6) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 68 and a light chain amino acid sequence of SEQ ID NO: 69. In another embodiment, the binding protein capable of binding TNF (seq. 6) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 70 and a light chain amino acid sequence of SEQ ID NO: 71.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 72 or SEQ ID NO. 74; and a light chain amino acid sequence of SEQ ID NO. 73 or SEQ ID NO. 75. In an embodiment, the binding protein capable of binding TNF (seq. 2) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 72 and a light chain amino acid sequence of SEQ ID NO: 73. In another embodiment, the binding protein capable of binding TNF (seq. 2) and SOST has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 74 and a light chain amino acid sequence of SEQ ID NO: 75.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 76 or SEQ ID NO. 78; and a light chain amino acid sequence of SEQ ID NO. 77 or SEQ ID NO. 79. In an embodiment, the binding protein capable of binding TNF (seq. 3) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 76 and a light chain amino acid sequence of SEQ ID NO: 77. In another embodiment, the binding protein capable of binding TNF (seq. 3) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 78 and a light chain amino acid sequence of SEQ ID NO: 79.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 80 or SEQ ID NO. 82; and a light chain amino acid sequence of SEQ ID NO. 81 or SEQ ID NO. 83. In an embodiment, the binding protein capable of binding TNF (seq. 4) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 80 and a light chain amino acid sequence of SEQ ID NO: 81. In another embodiment, the binding protein capable of binding TNF (seq. 4) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 82 and a light chain amino acid sequence of SEQ ID NO: 83.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 84 or SEQ ID NO. 86; and a light chain amino acid sequence of SEQ ID NO. 85 or SEQ ID NO. 87. In an embodiment, the binding protein capable of binding TNF (seq. 5) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 84 and a light chain amino acid sequence of SEQ ID NO: 85. In another embodiment, the binding protein capable of binding TNF (seq. 5) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 86 and a light chain amino acid sequence of SEQ ID NO: 87.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 88 or SEQ ID NO. 90; and a light chain amino acid sequence of SEQ ID NO. 89 or SEQ ID NO. 91. In an embodiment, the binding protein capable of binding TNF (seq. 6) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 88 and a light chain amino acid sequence of SEQ ID NO: 89. In another embodiment, the binding protein capable of binding TNF (seq. 6) and SOST has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 90 and a light chain amino acid sequence of SEQ ID NO: 91.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 92 or SEQ ID NO. 94; and a light chain amino acid sequence of SEQ ID NO. 93 or SEQ ID NO. 95. In an embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 92 and a light chain amino acid sequence of SEQ ID NO: 93. In another embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 94 and a light chain amino acid sequence of SEQ ID NO: 95.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 96 or SEQ ID NO. 98; and a light chain amino acid sequence of SEQ ID NO. 97 or SEQ ID NO. 99. In an embodiment, the binding protein capable of binding TNF (seq. 3) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 96 and a light chain amino acid sequence of SEQ ID NO: 97. In another embodiment, the binding protein capable of binding TNF (seq. 3) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 98 and a light chain amino acid sequence of SEQ ID NO: 99.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 100 or SEQ ID NO. 102; and a light chain amino acid sequence of SEQ ID NO. 101 or SEQ ID NO. 103. In an embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 100 and a light chain amino acid sequence of SEQ ID NO: 101. In another embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 102 and a light chain amino acid sequence of SEQ ID NO: 103.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 104 or SEQ ID NO. 106; and a light chain amino acid sequence of SEQ ID NO. 105 or SEQ ID NO. 107. In an embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 104 and a light chain amino acid sequence of SEQ ID NO: 105. In another embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 106 and a light chain amino acid sequence of SEQ ID NO: 107.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 108 or SEQ ID NO. 110; and a
5 light chain amino acid sequence of SEQ ID NO. 109 or SEQ ID NO. 111. In an embodiment, the binding protein capable of binding TNF (seq. 6) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 108 and a light chain amino acid sequence of SEQ ID NO: 109. In another embodiment, the binding protein capable of binding TNF (seq. 6) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 110 and a light
10 chain amino acid sequence of SEQ ID NO: 111.

In an embodiment, the binding protein capable of binding TNF (seq. 1) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 112 or SEQ ID NO. 114; and a
light chain amino acid sequence of SEQ ID NO. 113 or SEQ ID NO. 115. In an embodiment, the binding protein capable of binding TNF (seq. 1) and LPA comprises a heavy chain amino acid
15 sequence of SEQ ID NO. 112 and a light chain amino acid sequence of SEQ ID NO: 113. In another embodiment, the binding protein capable of binding TNF (seq. 1) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 114 and a light chain amino acid sequence of SEQ ID NO: 115.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and LPA
20 comprises a heavy chain amino acid sequence of SEQ ID NO. 116 or SEQ ID NO. 118; and a light chain amino acid sequence of SEQ ID NO. 117 or SEQ ID NO. 119. In an embodiment, the binding protein capable of binding TNF (seq. 2) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 116 and a light chain amino acid sequence of SEQ ID NO: 117. In another embodiment, the binding protein capable of binding TNF (seq. 2) and LPA has a reverse
25 orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 118 and a light chain amino acid sequence of SEQ ID NO: 119.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and LPA
comprises a heavy chain amino acid sequence of SEQ ID NO. 120 or SEQ ID NO. 122; and a
light chain amino acid sequence of SEQ ID NO. 121 or SEQ ID NO. 123. In an embodiment, the
30 binding protein capable of binding TNF (seq. 3) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 120 and a light chain amino acid sequence of SEQ ID NO: 121. In another embodiment, the binding protein capable of binding TNF (seq. 3) and LPA has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 122 and a light chain amino acid sequence of SEQ ID NO: 123.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 124 or SEQ ID NO. 126; and a light chain amino acid sequence of SEQ ID NO. 125 or SEQ ID NO. 127. In an embodiment, the binding protein capable of binding TNF (seq. 4) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 124 and a light chain amino acid sequence of SEQ ID NO: 125. In another embodiment, the binding protein capable of binding TNF (seq. 4) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 126 and a light chain amino acid sequence of SEQ ID NO: 127.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 128 or SEQ ID NO. 130; and a light chain amino acid sequence of SEQ ID NO. 129 or SEQ ID NO. 131. In an embodiment, the binding protein capable of binding TNF (seq. 5) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 128 and a light chain amino acid sequence of SEQ ID NO: 129. In another embodiment, the binding protein capable of binding TNF (seq. 5) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 130 and a light chain amino acid sequence of SEQ ID NO: 131.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 132 or SEQ ID NO. 134; and a light chain amino acid sequence of SEQ ID NO. 133 or SEQ ID NO. 135. In an embodiment, the binding protein capable of binding TNF (seq. 6) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 132 and a light chain amino acid sequence of SEQ ID NO: 133. In another embodiment, the binding protein capable of binding TNF (seq. 6) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 134 and a light chain amino acid sequence of SEQ ID NO: 135.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 136 or SEQ ID NO. 138; and a light chain amino acid sequence of SEQ ID NO. 137 or SEQ ID NO. 139. In an embodiment, the binding protein capable of binding TNF (seq. 2) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 136 and a light chain amino acid sequence of SEQ ID NO: 137. In another embodiment, the binding protein capable of binding TNF (seq. 2) and NGF has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 138 and a light chain amino acid sequence of SEQ ID NO: 139.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 140 or SEQ ID NO. 142; and a
5 light chain amino acid sequence of SEQ ID NO. 141 or SEQ ID NO. 143. In an embodiment, the binding protein capable of binding TNF (seq. 3) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 140 and a light chain amino acid sequence of SEQ ID NO: 141. In another embodiment, the binding protein capable of binding TNF (seq. 3) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 142 and a light
10 chain amino acid sequence of SEQ ID NO: 143.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 144 or SEQ ID NO. 146; and a
light chain amino acid sequence of SEQ ID NO. 145 or SEQ ID NO. 147. In an embodiment, the binding protein capable of binding TNF (seq. 4) and NGF comprises a heavy chain amino acid
15 sequence of SEQ ID NO. 144 and a light chain amino acid sequence of SEQ ID NO: 145. In another embodiment, the binding protein capable of binding TNF (seq. 4) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 146 and a light chain amino acid sequence of SEQ ID NO: 147.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and NGF
20 comprises a heavy chain amino acid sequence of SEQ ID NO. 148 or SEQ ID NO. 150; and a light chain amino acid sequence of SEQ ID NO. 149 or SEQ ID NO. 151. In an embodiment, the binding protein capable of binding TNF (seq. 5) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 148 and a light chain amino acid sequence of SEQ ID NO: 149. In another embodiment, the binding protein capable of binding TNF (seq. 5) and NGF has a reverse
25 orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 150 and a light chain amino acid sequence of SEQ ID NO: 151.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and NGF
comprises a heavy chain amino acid sequence of SEQ ID NO. 152 or SEQ ID NO. 154; and a
light chain amino acid sequence of SEQ ID NO. 153 or SEQ ID NO. 155. In an embodiment, the
30 binding protein capable of binding TNF (seq. 6) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 152 and a light chain amino acid sequence of SEQ ID NO: 153. In another embodiment, the binding protein capable of binding TNF (seq. 6) and NGF has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 154 and a light chain amino acid sequence of SEQ ID NO: 155.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 156 or SEQ ID NO. 158; and a
5 light chain amino acid sequence of SEQ ID NO. 157 or SEQ ID NO. 159. In an embodiment, the binding protein capable of binding TNF (seq. 2) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 156 and a light chain amino acid sequence of SEQ ID NO: 157. In another embodiment, the binding protein capable of binding TNF (seq. 2) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 158 and a
10 light chain amino acid sequence of SEQ ID NO: 159.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 160 or SEQ ID NO. 162; and a light chain amino acid sequence of SEQ ID NO. 161 or SEQ ID NO. 163. In an embodiment, the binding protein capable of binding TNF (seq. 3) and SOST comprises a heavy chain amino acid
15 sequence of SEQ ID NO. 160 and a light chain amino acid sequence of SEQ ID NO: 161. In another embodiment, the binding protein capable of binding TNF (seq. 3) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 162 and a light chain amino acid sequence of SEQ ID NO: 163.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and SOST
20 comprises a heavy chain amino acid sequence of SEQ ID NO. 164 or SEQ ID NO. 166; and a light chain amino acid sequence of SEQ ID NO. 165 or SEQ ID NO. 167. In an embodiment, the binding protein capable of binding TNF (seq. 4) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 164 and a light chain amino acid sequence of SEQ ID NO: 165. In another embodiment, the binding protein capable of binding TNF (seq. 4) and SOST has a
25 reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 166 and a light chain amino acid sequence of SEQ ID NO: 167.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and SOST
30 comprises a heavy chain amino acid sequence of SEQ ID NO. 168 or SEQ ID NO. 170; and a light chain amino acid sequence of SEQ ID NO. 169 or SEQ ID NO. 171. In an embodiment, the binding protein capable of binding TNF (seq. 5) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 168 and a light chain amino acid sequence of SEQ ID NO: 169. In another embodiment, the binding protein capable of binding TNF (seq. 5) and SOST has a

reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 170 and a light chain amino acid sequence of SEQ ID NO: 171.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 172 or SEQ ID NO. 174; and a light chain amino acid sequence of SEQ ID NO. 173 or SEQ ID NO. 175. In an embodiment, the binding protein capable of binding TNF (seq. 6) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 172 and a light chain amino acid sequence of SEQ ID NO: 173. In another embodiment, the binding protein capable of binding TNF (seq. 6) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 174 and a light chain amino acid sequence of SEQ ID NO: 175.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 176 or SEQ ID NO. 178; and a light chain amino acid sequence of SEQ ID NO. 177 or SEQ ID NO. 179. In an embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 176 and a light chain amino acid sequence of SEQ ID NO: 177. In another embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 178 and a light chain amino acid sequence of SEQ ID NO: 179.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 180 or SEQ ID NO. 182; and a light chain amino acid sequence of SEQ ID NO. 181 or SEQ ID NO. 183. In an embodiment, the binding protein capable of binding TNF (seq. 3) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 180 and a light chain amino acid sequence of SEQ ID NO: 181. In another embodiment, the binding protein capable of binding TNF (seq. 3) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 182 and a light chain amino acid sequence of SEQ ID NO: 183.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 184 or SEQ ID NO. 186; and a light chain amino acid sequence of SEQ ID NO. 185 or SEQ ID NO. 187. In an embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 184 and a light chain amino acid sequence of SEQ ID NO: 185. In another embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 186 and a light chain amino acid sequence of SEQ ID NO: 187.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 188 or SEQ ID NO. 190; and a
5 light chain amino acid sequence of SEQ ID NO. 189 or SEQ ID NO. 191. In an embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 188 and a light chain amino acid sequence of SEQ ID NO: 189. In another embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 190 and a light
10 chain amino acid sequence of SEQ ID NO: 191.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 192 or SEQ ID NO. 194; and a light chain amino acid sequence of SEQ ID NO. 193 or SEQ ID NO. 195. In an embodiment, the
15 binding protein capable of binding TNF (seq. 6) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 192 and a light chain amino acid sequence of SEQ ID NO: 193. In another embodiment, the binding protein capable of binding TNF (seq. 6) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 194 and a light chain amino acid sequence of SEQ ID NO: 195.

In an embodiment, the binding protein capable of binding TNF (seq. 1) and LPA
20 comprises a heavy chain amino acid sequence of SEQ ID NO. 196 or SEQ ID NO. 198; and a light chain amino acid sequence of SEQ ID NO. 197 or SEQ ID NO. 199. In an embodiment, the binding protein capable of binding TNF (seq. 1) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 196 and a light chain amino acid sequence of SEQ ID NO: 197. In another embodiment, the binding protein capable of binding TNF (seq. 1) and LPA has a reverse
25 orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 198 and a light chain amino acid sequence of SEQ ID NO: 199.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and LPA
30 comprises a heavy chain amino acid sequence of SEQ ID NO. 200 or SEQ ID NO. 202; and a light chain amino acid sequence of SEQ ID NO. 201 or SEQ ID NO. 203. In an embodiment, the binding protein capable of binding TNF (seq. 2) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 200 and a light chain amino acid sequence of SEQ ID NO: 201. In another embodiment, the binding protein capable of binding TNF (seq. 2) and LPA has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 202 and a light chain amino acid sequence of SEQ ID NO: 203.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 204 or SEQ ID NO. 206; and a light chain amino acid sequence of SEQ ID NO. 205 or SEQ ID NO. 207. In an embodiment, the binding protein capable of binding TNF (seq. 3) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 204 and a light chain amino acid sequence of SEQ ID NO: 205. In another embodiment, the binding protein capable of binding TNF (seq. 3) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 206 and a light chain amino acid sequence of SEQ ID NO: 207.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 208 or SEQ ID NO. 210; and a light chain amino acid sequence of SEQ ID NO. 209 or SEQ ID NO. 211. In an embodiment, the binding protein capable of binding TNF (seq. 4) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 208 and a light chain amino acid sequence of SEQ ID NO: 209. In another embodiment, the binding protein capable of binding TNF (seq. 4) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 210 and a light chain amino acid sequence of SEQ ID NO: 211.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 212 or SEQ ID NO. 214; and a light chain amino acid sequence of SEQ ID NO. 213 or SEQ ID NO. 215. In an embodiment, the binding protein capable of binding TNF (seq. 5) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 212 and a light chain amino acid sequence of SEQ ID NO: 213. In another embodiment, the binding protein capable of binding TNF (seq. 5) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 214 and a light chain amino acid sequence of SEQ ID NO: 215.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 216 or SEQ ID NO. 218; and a light chain amino acid sequence of SEQ ID NO. 217 or SEQ ID NO. 219. In an embodiment, the binding protein capable of binding TNF (seq. 6) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 216 and a light chain amino acid sequence of SEQ ID NO: 217. In another embodiment, the binding protein capable of binding TNF (seq. 6) and LPA has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 218 and a light chain amino acid sequence of SEQ ID NO: 219.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 220 or SEQ ID NO. 222; and a
5 light chain amino acid sequence of SEQ ID NO. 221 or SEQ ID NO. 223. In an embodiment, the binding protein capable of binding TNF (seq. 2) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 220 and a light chain amino acid sequence of SEQ ID NO: 221. In another embodiment, the binding protein capable of binding TNF (seq. 2) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 222 and a light
10 chain amino acid sequence of SEQ ID NO: 223.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 224 or SEQ ID NO. 226; and a
light chain amino acid sequence of SEQ ID NO. 225 or SEQ ID NO. 227. In an embodiment, the
15 binding protein capable of binding TNF (seq. 3) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 224 and a light chain amino acid sequence of SEQ ID NO: 225. In another embodiment, the binding protein capable of binding TNF (seq. 3) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 226 and a light
chain amino acid sequence of SEQ ID NO: 227.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and NGF
20 comprises a heavy chain amino acid sequence of SEQ ID NO. 228 or SEQ ID NO. 230; and a light chain amino acid sequence of SEQ ID NO. 229 or SEQ ID NO. 231. In an embodiment, the binding protein capable of binding TNF (seq. 4) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 228 and a light chain amino acid sequence of SEQ ID NO: 229. In another embodiment, the binding protein capable of binding TNF (seq. 4) and NGF has a reverse
25 orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 230 and a light chain amino acid sequence of SEQ ID NO: 231.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and NGF
comprises a heavy chain amino acid sequence of SEQ ID NO. 232 or SEQ ID NO. 234; and a
light chain amino acid sequence of SEQ ID NO. 233 or SEQ ID NO. 235. In an embodiment, the
30 binding protein capable of binding TNF (seq. 5) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 232 and a light chain amino acid sequence of SEQ ID NO: 233. In another embodiment, the binding protein capable of binding TNF (seq. 5) and NGF has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 234 and a light chain amino acid sequence of SEQ ID NO: 235.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 236 or SEQ ID NO. 238; and a light chain amino acid sequence of SEQ ID NO. 237 or SEQ ID NO. 239. In an embodiment, the binding protein capable of binding TNF (seq. 6) and NGF comprises a heavy chain amino acid sequence of SEQ ID NO. 236 and a light chain amino acid sequence of SEQ ID NO: 237. In another embodiment, the binding protein capable of binding TNF (seq. 6) and NGF has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 238 and a light chain amino acid sequence of SEQ ID NO: 239.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 240 or SEQ ID NO. 242; and a light chain amino acid sequence of SEQ ID NO. 241 or SEQ ID NO. 243. In an embodiment, the binding protein capable of binding TNF (seq. 2) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 240 and a light chain amino acid sequence of SEQ ID NO: 241. In another embodiment, the binding protein capable of binding TNF (seq. 2) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 242 and a light chain amino acid sequence of SEQ ID NO: 243.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 244 or SEQ ID NO. 246; and a light chain amino acid sequence of SEQ ID NO. 245 or SEQ ID NO. 247. In an embodiment, the binding protein capable of binding TNF (seq. 3) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 244 and a light chain amino acid sequence of SEQ ID NO: 245. In another embodiment, the binding protein capable of binding TNF (seq. 3) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 246 and a light chain amino acid sequence of SEQ ID NO: 247.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 248 or SEQ ID NO. 250; and a light chain amino acid sequence of SEQ ID NO. 249 or SEQ ID NO. 251. In an embodiment, the binding protein capable of binding TNF (seq. 4) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 248 and a light chain amino acid sequence of SEQ ID NO: 249. In another embodiment, the binding protein capable of binding TNF (seq. 4) and SOST has a

reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 250 and a light chain amino acid sequence of SEQ ID NO: 251.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 252 or SEQ ID NO. 254; and a
5 light chain amino acid sequence of SEQ ID NO. 253 or SEQ ID NO. 255. In an embodiment, the binding protein capable of binding TNF (seq. 5) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 252 and a light chain amino acid sequence of SEQ ID NO: 253. In another embodiment, the binding protein capable of binding TNF (seq. 5) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 254 and a
10 light chain amino acid sequence of SEQ ID NO: 255.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and SOST comprises a heavy chain amino acid sequence of SEQ ID NO. 256 or SEQ ID NO. 258; and a
light chain amino acid sequence of SEQ ID NO. 257 or SEQ ID NO. 259. In an embodiment, the binding protein capable of binding TNF (seq. 6) and SOST comprises a heavy chain amino acid
15 sequence of SEQ ID NO. 256 and a light chain amino acid sequence of SEQ ID NO: 257. In another embodiment, the binding protein capable of binding TNF (seq. 6) and SOST has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 258 and a light chain amino acid sequence of SEQ ID NO: 259.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2
20 comprises a heavy chain amino acid sequence of SEQ ID NO. 260 or SEQ ID NO. 262; and a light chain amino acid sequence of SEQ ID NO. 261 or SEQ ID NO. 263. In an embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 260 and a light chain amino acid sequence of SEQ ID NO: 261. In another embodiment, the binding protein capable of binding TNF (seq. 2) and PGE2 has a reverse
25 orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 262 and a light chain amino acid sequence of SEQ ID NO: 263.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and PGE2
comprises a heavy chain amino acid sequence of SEQ ID NO. 264 or SEQ ID NO. 266; and a
light chain amino acid sequence of SEQ ID NO. 265 or SEQ ID NO. 267. In an embodiment, the
30 binding protein capable of binding TNF (seq. 3) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 264 and a light chain amino acid sequence of SEQ ID NO: 265. In another embodiment, the binding protein capable of binding TNF (seq. 3) and PGE2 has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 266 and a light chain amino acid sequence of SEQ ID NO: 267.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 268 or SEQ ID NO. 270; and a light chain amino acid sequence of SEQ ID NO. 269 or SEQ ID NO. 271. In an embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 268 and a light chain amino acid sequence of SEQ ID NO: 269. In another embodiment, the binding protein capable of binding TNF (seq. 4) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 270 and a light chain amino acid sequence of SEQ ID NO: 271.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 272 or SEQ ID NO. 274; and a light chain amino acid sequence of SEQ ID NO. 273 or SEQ ID NO. 275. In an embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 272 and a light chain amino acid sequence of SEQ ID NO: 273. In another embodiment, the binding protein capable of binding TNF (seq. 5) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 274 and a light chain amino acid sequence of SEQ ID NO: 275.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 276 or SEQ ID NO. 278; and a light chain amino acid sequence of SEQ ID NO. 277 or SEQ ID NO. 279. In an embodiment, the binding protein capable of binding TNF (seq. 6) and PGE2 comprises a heavy chain amino acid sequence of SEQ ID NO. 276 and a light chain amino acid sequence of SEQ ID NO: 277. In another embodiment, the binding protein capable of binding TNF (seq. 6) and PGE2 has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 278 and a light chain amino acid sequence of SEQ ID NO: 279.

In an embodiment, the binding protein capable of binding TNF (seq. 1) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 280 or SEQ ID NO. 282; and a light chain amino acid sequence of SEQ ID NO. 281 or SEQ ID NO. 283. In an embodiment, the binding protein capable of binding TNF (seq. 1) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 280 and a light chain amino acid sequence of SEQ ID NO: 281. In another embodiment, the binding protein capable of binding TNF (seq. 1) and LPA has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 282 and a light chain amino acid sequence of SEQ ID NO: 283.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 284 or SEQ ID NO. 286; and a
5 light chain amino acid sequence of SEQ ID NO. 285 or SEQ ID NO. 287. In an embodiment, the binding protein capable of binding TNF (seq. 2) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 284 and a light chain amino acid sequence of SEQ ID NO: 285. In another embodiment, the binding protein capable of binding TNF (seq. 2) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 286 and a light
10 chain amino acid sequence of SEQ ID NO: 287.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 288 or SEQ ID NO. 290; and a
light chain amino acid sequence of SEQ ID NO. 289 or SEQ ID NO. 291. In an embodiment, the
15 binding protein capable of binding TNF (seq. 3) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 288 and a light chain amino acid sequence of SEQ ID NO: 289. In another embodiment, the binding protein capable of binding TNF (seq. 3) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 290 and a light chain amino acid sequence of SEQ ID NO: 291.

In an embodiment, the binding protein capable of binding TNF (seq. 4) and LPA
20 comprises a heavy chain amino acid sequence of SEQ ID NO. 292 or SEQ ID NO. 294; and a light chain amino acid sequence of SEQ ID NO. 293 or SEQ ID NO. 295. In an embodiment, the binding protein capable of binding TNF (seq. 4) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 292 and a light chain amino acid sequence of SEQ ID NO: 293. In another embodiment, the binding protein capable of binding TNF (seq. 4) and LPA has a reverse
25 orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 294 and a light chain amino acid sequence of SEQ ID NO: 295.

In an embodiment, the binding protein capable of binding TNF (seq. 5) and LPA
comprises a heavy chain amino acid sequence of SEQ ID NO. 296 or SEQ ID NO. 298; and a
light chain amino acid sequence of SEQ ID NO. 297 or SEQ ID NO. 299. In an embodiment, the
30 binding protein capable of binding TNF (seq. 5) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 296 and a light chain amino acid sequence of SEQ ID NO: 297. In another embodiment, the binding protein capable of binding TNF (seq. 5) and LPA has a reverse

orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 298 and a light chain amino acid sequence of SEQ ID NO: 299.

In an embodiment, the binding protein capable of binding TNF (seq. 6) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 300 or SEQ ID NO. 302; and a light chain amino acid sequence of SEQ ID NO. 301 or SEQ ID NO. 303. In an embodiment, the binding protein capable of binding TNF (seq. 6) and LPA comprises a heavy chain amino acid sequence of SEQ ID NO. 300 and a light chain amino acid sequence of SEQ ID NO: 301. In another embodiment, the binding protein capable of binding TNF (seq. 6) and LPA has a reverse orientation and comprises a heavy chain amino acid sequence of SEQ ID NO. 302 and a light chain amino acid sequence of SEQ ID NO: 303.

In another embodiment, the binding protein comprises a polypeptide chain, wherein said polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein; VD1 is a first heavy chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second heavy chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a heavy chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n is an Fc region, wherein said (X2)_n is either present or absent. In an embodiment, the Fc region is absent from the binding protein.

In another embodiment, the binding protein comprises a polypeptide chain, wherein said polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein, VD1 is a first light chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second light chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a light chain constant domain; (X1)_n is a linker with the proviso that it is not CL, wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region, wherein said (X2)_n is either present or absent. In an embodiment, (X2)_n is absent from the binding protein.

In another embodiment the binding protein comprises first and second polypeptide chains, wherein said first polypeptide chain comprises a first VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first heavy chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second heavy chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a heavy chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n is an Fc region, wherein said (X2)_n is either present or absent; and wherein said second

polypeptide chain comprises a second VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first light chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second light chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a light chain constant domain; (X1)_n is a linker (optionally which is not CL), wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region, wherein said (X2)_n is either present or absent. In another embodiment, the binding protein comprises two first polypeptide chains and two second polypeptide chains. In yet another embodiment, (X2)_n is absent from the second polypeptide. In still another embodiment, the Fc region, if present in the first polypeptide is a native sequence Fc region or a variant sequence Fc region. In still another embodiment, the Fc region is from an IgG1, IgG2, IgG3, IgG4, IgA, IgM, IgE, or IgD.

In another embodiment, the binding protein is a DVD-Ig capable of binding two antigens comprising four polypeptide chains, wherein, first and third polypeptide chains comprise VD1-(X1)_n-VD2-C-(X2)_n, wherein, VD1 is a first heavy chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second heavy chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a heavy chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n is an Fc region, wherein said (X2)_n is either present or absent; and wherein second and fourth polypeptide chains comprise VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first light chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second light chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a light chain constant domain; (X1)_n is a linker (optionally which is not CL), wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region, wherein said (X2)_n is either present or absent.

A method of making a DVD-Ig binding protein by preselecting the parent antibodies is also provided. In an embodiment, the method of making a Dual Variable Domain Immunoglobulin capable of binding two antigens comprising the steps of a) obtaining a first parent antibody or antigen binding portion thereof, capable of binding a first antigen; b) obtaining a second parent antibody or antigen binding portion thereof, capable of binding a second antigen; c) constructing first and third polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein, VD1 is a first heavy chain variable domain obtained from said first parent antibody or antigen binding portion thereof; VD2 is a second heavy chain variable domain obtained from said second parent antibody or antigen binding portion thereof; C is a heavy chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and

(X2)_n is an Fc region, wherein said (X2)_n is either present or absent; d) constructing second and fourth polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein, VD1 is a first light chain variable domain obtained from said first parent antibody or antigen binding portion thereof; VD2 is a second light chain variable domain obtained from said second parent antibody or antigen binding thereof; C is a light chain constant domain; (X1)_n is a linker (optionally which is not CL), wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region, wherein said (X2)_n is either present or absent; e) expressing said first, second, third and fourth polypeptide chains; such that a Dual Variable Domain Immunoglobulin capable of binding said first and said second antigen is generated.

10 In still another embodiment, a method of generating a Dual Variable Domain Immunoglobulin capable of binding two antigens with desired properties comprising the steps of a) obtaining a first parent antibody or antigen binding portion thereof, capable of binding a first antigen and possessing at least one desired property exhibited by the Dual Variable Domain Immunoglobulin; b) obtaining a second parent antibody or antigen binding portion thereof,
15 capable of binding a second antigen and possessing at least one desired property exhibited by the Dual Variable Domain Immunoglobulin; c) constructing first and third polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein; VD1 is a first heavy chain variable domain obtained from said first parent antibody or antigen binding portion thereof; VD2 is a second heavy chain variable domain obtained from said second parent antibody or antigen binding
20 portion thereof; C is a heavy chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n is an Fc region, wherein said (X2)_n is either present or absent; d) constructing second and fourth polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein; VD1 is a first light chain variable domain obtained from said first parent antibody or antigen binding portion thereof; VD2 is a second light
25 chain variable domain obtained from said second parent antibody or antigen binding portion thereof; C is a light chain constant domain; (X1)_n is a linker (optionally which is not CL), wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region, wherein said (X2)_n is either present or absent; e) expressing said first, second, third and fourth polypeptide chains; such that a Dual Variable Domain Immunoglobulin capable of binding said
30 first and said second antigen with desired properties is generated is provided.

In one embodiment, the VDI of the first and second polypeptide chains disclosed herein are obtained from the same parent antibody or antigen binding portion thereof. In another embodiment, the VDI of the first and second polypeptide chains disclosed herein are obtained from different parent antibodies or antigen binding portions thereof. In another embodiment, the

VD2 of the first and second polypeptide chains disclosed herein are obtained from the same parent antibody or antigen binding portion thereof. In another embodiment, the VD2 of the first and second polypeptide chains disclosed herein are obtained from different parent antibodies or antigen binding portions thereof.

5 In one embodiment the first parent antibody or antigen binding portion thereof, and the second parent antibody or antigen binding portion thereof, are the same antibody. In another embodiment the first parent antibody or antigen binding portion thereof, and the second parent antibody or antigen binding portion thereof, are different antibodies.

10 In one embodiment the first parent antibody or antigen binding portion thereof, binds a first antigen and the second parent antibody or antigen binding portion thereof, binds a second antigen. In a particular embodiment, the first and second antigens are the same antigen. In another embodiment, the parent antibodies bind different epitopes on the same antigen. In another embodiment the first and second antigens are different antigens. In another embodiment, the first parent antibody or antigen binding portion thereof, binds the first antigen with a potency different
15 from the potency with which the second parent antibody or antigen binding portion thereof, binds the second antigen. In yet another embodiment, the first parent antibody or antigen binding portion thereof, binds the first antigen with an affinity different from the affinity with which the second parent antibody or antigen binding portion thereof, binds the second antigen.

20 In another embodiment the first parent antibody or antigen binding portion thereof, and the second parent antibody or antigen binding portion thereof, are a human antibody, a CDR grafted antibody, or a humanized antibody. In an embodiment, the antigen binding portions are Fab fragments, F(ab')₂ fragments, bivalent fragments comprising two Fab fragments linked by a disulfide bridge at the hinge region, Fd fragments comprising VH and CH1 domains, Fv fragments comprising the VL and VH domains of a single arm of an antibody, dAb fragments,
25 isolated complementarity determining regions (CDR), single chain antibodies, or diabodies.

30 In another embodiment the binding protein possesses at least one desired property exhibited by the first parent antibody or antigen binding portion thereof, or the second parent antibody or antigen binding portion thereof. Alternatively, the first parent antibody or antigen binding portion thereof and the second parent antibody or antigen binding portion thereof possess at least one desired property exhibited by the Dual Variable Domain Immunoglobulin. In an embodiment, the desired property is selected from one or more antibody parameters. In another embodiment, the antibody parameters are antigen specificity, affinity to antigen, potency, biological function, epitope recognition, stability, solubility, production efficiency,

immunogenicity, pharmacokinetics, bioavailability, tissue cross reactivity, or orthologous antigen binding. In an embodiment the binding protein is multivalent. In another embodiment, the binding protein is multispecific. The multivalent and or multispecific binding proteins described herein have desirable properties particularly from a therapeutic standpoint. For instance, the multivalent and or multispecific binding protein may (1) be internalized (and/or catabolized) faster than a bivalent antibody by a cell expressing an antigen to which the antibodies bind; (2) be an agonist antibody; and/or (3) induce cell death and/or apoptosis of a cell expressing an antigen which the multivalent antibody is capable of binding to. The "parent antibody" which provides at least one antigen binding specificity of the multivalent and or multispecific binding proteins may be one which is internalized (and/or catabolized) by a cell expressing an antigen to which the antibody binds; and/or may be an agonist, cell death-inducing, and/or apoptosis-inducing antibody, and the multivalent and or multispecific binding protein as described herein may display improvement(s) in one or more of these properties. Moreover, the parent antibody may lack any one or more of these properties, but may be endowed with them when constructed as a multivalent binding protein as described herein.

In another embodiment the binding protein has an on rate constant (K_{on}) to one or more targets of: at least about $10^2 M^{-1} s^{-1}$; at least about $10^3 M^{-1} s^{-1}$; at least about $10^4 M^{-1} s^{-1}$; at least about $10^5 M^{-1} s^{-1}$; and at least about $10^6 M^{-1} s^{-1}$, as measured by surface plasmon resonance. In an embodiment, the binding protein has an on rate constant (K_{on}) to one or more targets between $10^2 M^{-1} s^{-1}$ and $10^3 M^{-1} s^{-1}$; between $10^3 M^{-1} s^{-1}$ and $10^4 M^{-1} s^{-1}$; between $10^4 M^{-1} s^{-1}$ and $10^5 M^{-1} s^{-1}$; or between $10^5 M^{-1} s^{-1}$ and $10^6 M^{-1} s^{-1}$, as measured by surface plasmon resonance.

In another embodiment the binding protein has an off rate constant (K_{off}) for one or more targets of: at most about $10^{-3} s^{-1}$; at most about $10^{-4} s^{-1}$; at most about $10^{-5} s^{-1}$; and at most about $10^{-6} s^{-1}$, as measured by surface plasmon resonance. In an embodiment, the binding protein has an off rate constant (K_{off}) to one or more targets of $10^{-3} s^{-1}$ to $10^{-4} s^{-1}$; of $10^{-4} s^{-1}$ to $10^{-5} s^{-1}$; or of $10^{-5} s^{-1}$ to $10^{-6} s^{-1}$, as measured by surface plasmon resonance.

In another embodiment the binding protein has a dissociation constant (K_D) to one or more targets of: at most about $10^{-7} M$; at most about $10^{-8} M$; at most about $10^{-9} M$; at most about $10^{-10} M$; at most about $10^{-11} M$; at most about $10^{-12} M$; and at most $10^{-13} M$. In an embodiment, the binding protein has a dissociation constant (K_D) to its targets of $10^{-7} M$ to $10^{-8} M$; of $10^{-8} M$ to $10^{-9} M$; of $10^{-9} M$ to $10^{-10} M$; of $10^{-10} M$ to $10^{-11} M$; of $10^{-11} M$ to $10^{-12} M$; or of $10^{-12} M$ to $10^{-13} M$.

In another embodiment, the binding protein described herein is a conjugate further comprising an agent, wherein said agent is an immunoadhesion molecule, an imaging agent, a

therapeutic agent, or a cytotoxic agent. In an embodiment, the imaging agent is a radiolabel, an enzyme, a fluorescent label, a luminescent label, a bioluminescent label, a magnetic label, or biotin. In another embodiment, the radiolabel is: ^3H , ^{14}C , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I , ^{177}Lu , ^{166}Ho , or ^{153}Sm . In yet another embodiment, the therapeutic or cytotoxic agent is an anti-metabolite, an alkylating agent, an antibiotic, a growth factor, a cytokine, an anti-angiogenic agent, an anti-mitotic agent, an anthracycline, toxin, or an apoptotic agent.

In another embodiment, the binding protein described herein is a crystallized binding protein and exists as a crystal. In an embodiment, the crystal is a carrier-free pharmaceutical controlled release crystal. In yet another embodiment, the crystallized binding protein has a greater half life in vivo than the soluble counterpart of said binding protein. In still another embodiment, the crystallized binding protein retains biological activity.

In another embodiment, the binding protein described herein is glycosylated. For example, the glycosylation is a human glycosylation pattern.

An isolated nucleic acid encoding any one of the binding proteins disclosed herein is also provided. A further embodiment provides a vector comprising the isolated nucleic acid disclosed herein wherein said vector is pcDNA; pTT (Durocher et al., *Nucleic Acids Research* 2002, Vol 30, No.2); pTT3 (pTT with additional multiple cloning site; pEFBOS (Mizushima, S. and Nagata, S., (1990) *Nucleic acids Research* Vol 18, No. 17); pBV; pJV; pcDNA3.1 TOPO; pEF6 TOPO; or pBJ. In an embodiment, the vector is a vector disclosed in US Patent Application Serial No. 61/021,282.

In another aspect a host cell is transformed with the vector disclosed herein. In an embodiment, the host cell is a prokaryotic cell. In another embodiment, the host cell is E.Coli. In a related embodiment the host cell is a eukaryotic cell. In another embodiment, the eukaryotic cell is a protist cell, an animal cell, a plant cell, or a fungal cell. In yet another embodiment, the host cell is a mammalian cell including, but not limited to, CHO, COS; NS0, SP2, PER.C6 or a fungal cell such as *Saccharomyces cerevisiae*; or an insect cell such as Sf9.

In an embodiment, two or more DVD-Igs, e.g., with different specificities, are produced in a single recombinant host cell. For example, the expression of a mixture of antibodies has been called Oligoelomics™ Merus B.V., The Netherlands); U.S. Patent Nos. 7,262,028; 7,429,486.

A method of producing a binding protein disclosed herein comprising culturing any one of the host cells also disclosed herein in a culture medium under conditions sufficient to produce the binding protein is also provided. In an embodiment, 50%-75% of the binding protein

produced by this method is a dual specific tetravalent binding protein. In a particular embodiment, 75%-90% of the binding protein produced by this method is a dual specific tetravalent binding protein. In a particular embodiment, 90%-95% of the binding protein produced is a dual specific tetravalent binding protein.

5 One embodiment provides a composition for the release of a binding protein wherein the composition comprises a formulation that in turn comprises a crystallized binding protein, as disclosed herein, and an ingredient, and at least one polymeric carrier. For example, the polymeric carrier is: poly (acrylic acid), poly (cyanoacrylates), poly (amino acids), poly (anhydrides), poly (depsipeptide), poly (esters), poly (lactic acid), poly (lactic-co-glycolic acid)
10 or PLGA, poly (b-hydroxybutyrate), poly (caprolactone), poly (dioxanone); poly (ethylene glycol), poly ((hydroxypropyl) methacrylamide, poly [(organo)phosphazene], poly (ortho esters), poly (vinyl alcohol), poly (vinylpyrrolidone), maleic anhydride- alkyl vinyl ether copolymers, pluronic polyols, albumin, alginate, cellulose and cellulose derivatives, collagen, fibrin, gelatin, hyaluronic acid, oligosaccharides, glycaminoglycans, sulfated polysaccharides, blends and
15 copolymers thereof. For example, in some embodiments the ingredient is albumin, sucrose, trehalose, lactitol, gelatin, hydroxypropyl- β - cyclodextrin, methoxypolyethylene glycol, or polyethylene glycol. Another embodiment provides a method for treating a mammal comprising the step of administering to the mammal an effective amount of the composition disclosed herein.

A pharmaceutical composition comprising a binding protein, as disclosed herein and a
20 pharmaceutically acceptable carrier is also provided. In a further embodiment the pharmaceutical composition comprises at least one additional therapeutic agent for treating a disorder. For example, the additional agent is: a therapeutic agent, an imaging agent, a cytotoxic agent, an angiogenesis inhibitor (including but not limited to an anti-VEGF antibody or a VEGF-trap), a kinase inhibitor (including but not limited to a KDR and a TIE-2 inhibitor), a co-stimulation
25 molecule blocker (including but not limited to anti-B7.1, anti-B7.2, CTLA4-Ig, anti-CD20), an adhesion molecule blocker (including but not limited to an anti-LFA-1 antibody, an anti-E/L selectin antibody, a small molecule inhibitor), an anti-cytokine antibody or functional fragment thereof (including but not limited to an anti-IL-18, an anti-TNF, and an anti-IL-6/cytokine receptor antibody), methotrexate, cyclosporin, rapamycin, FK506, a detectable label or reporter, a
30 TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, an erythropoietin, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, a radiopharmaceutical, an antidepressant, an antipsychotic, a stimulant, an

asthma medication, a beta agonist, an inhaled steroid, an epinephrine or analog, a cytokine, or a cytokine antagonist.

A method for treating a human subject suffering from a disorder in which the target, or targets, capable of being bound by the binding protein disclosed herein is detrimental, comprising administering to the human subject a binding protein disclosed herein such that the activity of the target, or targets in the human subject is inhibited and one of more symptoms is alleviated or treatment is achieved is also provided. For example, the disorder is selected from the group comprising arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's granulomatosis, Henoch-Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison's disease, sporadic polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella associated arthropathy, spondyloarthropathy, atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous candidiasis, giant cell arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, Acquired Immunodeficiency Disease Syndrome, Acquired Immunodeficiency Related Diseases, Hepatitis B, Hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-inflammatory interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease,

systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated lung disease, Sjögren's disease associated lung disease, ankylosing spondylitis associated lung disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced interstitial lung disease, fibrosis, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated hypoglycaemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute immune disease associated with organ transplantation, chronic immune disease associated with organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS, glomerulonephritides, microscopic vasculitis of the kidneys, lyme disease, discoid lupus erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all subtypes), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic sclerosis, Sjögren's syndrome, Takayasu's disease/arteritis, autoimmune thrombocytopaenia, idiopathic thrombocytopaenia, autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary myxoedema, phacogenic uveitis, primary vasculitis, vitiligo acute liver disease, chronic liver diseases, alcoholic cirrhosis, alcohol-induced liver injury, choleosatis, idiosyncratic liver disease, Drug-Induced hepatitis, Non-alcoholic Steatohepatitis, allergy and asthma, group B streptococci (GBS) infection, mental disorders (e.g., depression and schizophrenia), Th2 Type and Th1 Type mediated diseases, acute and chronic pain (different forms of pain), and cancers such as lung, breast, stomach, bladder, colon, pancreas, ovarian, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma), Abetalipoproteinemia, Acrocyanosis, acute and chronic parasitic or infectious processes, acute leukemia, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinomas, aerial ectopic beats, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allograft rejection, alpha-1- antitrypsin deficiency, amyotrophic lateral sclerosis, anemia, angina pectoris, anterior horn cell degeneration, anti cd3 therapy, antiphospholipid syndrome, anti-receptor hypersensitivity reactions, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, arteriosclerosis, arteriovenous fistula, ataxia, atrial fibrillation (sustained or paroxysmal), atrial flutter, atrioventricular block, B cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt's lymphoma, Burns, cardiac arrhythmias, cardiac stun syndrome, cardiac tumors,

cardiomyopathy, cardiopulmonary bypass inflammation response, cartilage transplant rejection, cerebellar cortical degenerations, cerebellar disorders, chaotic or multifocal atrial tachycardia, chemotherapy associated disorders, chronic myelocytic leukemia (CML), chronic alcoholism, chronic inflammatory pathologies, chronic lymphocytic leukemia (CLL), chronic obstructive pulmonary disease (COPD), chronic salicylate intoxication, colorectal carcinoma, congestive heart failure, conjunctivitis, contact dermatitis, cor pulmonale, coronary artery disease, Creutzfeldt-Jakob disease, culture negative sepsis, cystic fibrosis, cytokine therapy associated disorders, Dementia pugilistica, demyelinating diseases, dengue hemorrhagic fever, dermatitis, dermatologic conditions, diabetes, diabetes mellitus, diabetic atherosclerotic disease, Diffuse Lewy body disease, dilated congestive cardiomyopathy, disorders of the basal ganglia, Down's Syndrome in middle age, drug- induced movement disorders induced by drugs which block CNS dopamine receptors, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottitis, epstein-barr virus infection, erythromelalgia, extrapyramidal and cerebellar disorders, familial hemophagocytic lymphohistiocytosis, fetal thymus implant rejection, Friedreich's ataxia, functional peripheral arterial disorders, fungal sepsis, gas gangrene, gastric ulcer, glomerular nephritis, graft rejection of any organ or tissue, gram negative sepsis, gram positive sepsis, granulomas due to intracellular organisms, hairy cell leukemia, Hallerorden-Spatz disease, hashimoto's thyroiditis, hay fever, heart transplant rejection, hemachromatosis, hemodialysis, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, hemorrhage, hepatitis (A), His bundle arrhythmias, HIV infection/HIV neuropathy, Hodgkin's disease, hyperkinetic movement disorders, hypersensitivity reactions, hypersensitivity pneumonitis, hypertension, hypokinetic movement disorders, hypothalamic-pituitary-adrenal axis evaluation, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody mediated cytotoxicity, Asthenia, infantile spinal muscular atrophy, inflammation of the aorta, influenza a, ionizing radiation exposure, iridocyclitis/uveitis/optic neuritis, ischemia- reperfusion injury, ischemic stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, kidney transplant rejection, legionella, leishmaniasis, leprosy, lesions of the corticospinal system, lipedema, liver transplant rejection, lymphedema, malaria, malignant Lymphoma, malignant histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic diseases, migraine headache, mitochondrial multi.system disorder, mixed connective tissue disease, monoclonal gammopathy, multiple myeloma, multiple systems degenerations (Mencel Dejerine-Thomas Shi-Drager and Machado-Joseph), myasthenia gravis, mycobacterium avium intracellulare, mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disorders, nasopharyngeal carcinoma, neonatal chronic lung disease, nephritis, nephrosis, neurodegenerative diseases, neurogenic I muscular atrophies, neutropenic fever, non- hodgkins lymphoma, occlusion of the abdominal aorta and its branches, occlusive

arterial disorders, okt3 therapy, orchitis/epididymitis, orchitis/vasectomy reversal procedures, organomegaly, osteoporosis, pancreas transplant rejection, pancreatic carcinoma, paraneoplastic syndrome/hypercalcemia of malignancy, parathyroid transplant rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral atherosclerotic disease, peripheral
5 vascular disorders, peritonitis, pernicious anemia, pneumocystis carinii pneumonia, pneumonia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy, and skin changes syndrome), post perfusion syndrome, post pump syndrome, post-MI cardiomyopathy syndrome, preeclampsia, Progressive supranucleo Palsy, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon and disease, Raynaud's disease, Refsum's disease,
10 regular narrow QRS tachycardia, renovascular hypertension, reperfusion injury, restrictive cardiomyopathy, sarcomas, scleroderma, senile chorea, Senile Dementia of Lewy body type, seronegative arthropathies, shock, sickle cell anemia, skin allograft rejection, skin changes syndrome, small bowel transplant rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degenerations, streptococcal myositis, structural lesions of the cerebellum,
15 Subacute sclerosing panencephalitis, Syncope, syphilis of the cardiovascular system, systemic anaphalaxis, systemic inflammatory response syndrome, systemic onset juvenile rheumatoid arthritis, T-cell or FAB ALL, Telangiectasia, thromboangitis obliterans, thrombocytopenia, toxicity, transplants, trauma/hemorrhage, type III hypersensitivity reactions, type IV
hypersensitivity, unstable angina, uremia, urosepsis, urticaria, valvular heart diseases, varicose
20 veins, vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal infections, viral encephalitis/aseptic meningitis, vital-associated hemaphagocytic syndrome, Wernicke- Korsakoff syndrome, Wilson's disease, xenograft rejection of any organ or tissue, acute coronary syndromes, acute idiopathic polyneuritis, acute inflammatory demyelinating
polyradiculoneuropathy, acute ischemia, adult Still's disease, alopecia areata, anaphylaxis, anti-
25 phospholipid antibody syndrome, aplastic anemia, arteriosclerosis, atopic eczema, atopic dermatitis, autoimmune dermatitis, autoimmune disorder associated with streptococcus infection, autoimmune enteropathy, autoimmune hearing loss, autoimmune lymphoproliferative syndrome (ALPS), autoimmune myocarditis, autoimmune premature ovarian failure, blepharitis, bronchiectasis, bullous pemphigoid, cardiovascular disease, catastrophic antiphospholipid
30 syndrome, celiac disease, cervical spondylosis, chronic ischemia, cicatricial pemphigoid, clinically isolated syndrome (cis) with risk for multiple sclerosis, conjunctivitis, childhood onset psychiatric disorder, chronic obstructive pulmonary disease (COPD), dacryocystitis, dermatomyositis, diabetic retinopathy, diabetes mellitus, disk herniation, disk prolaps, drug
induced immune hemolytic anemia, endocarditis, endometriosis, endophthalmitis, episcleritis,
35 erythema multiforme, erythema multiforme major, gestational pemphigoid, Guillain-Barré syndrome (GBS), hay fever, Hughes syndrome, idiopathic Parkinson's disease, idiopathic

interstitial pneumonia, IgE-mediated allergy, immune hemolytic anemia, inclusion body myositis, infectious ocular inflammatory disease, inflammatory demyelinating disease, inflammatory heart disease, inflammatory kidney disease, IPF/UIP, iritis, keratitis, keratoconjunctivitis sicca, Kussmaul disease or Kussmaul-Meier disease, Landry's paralysis, Langerhan's cell histiocytosis, livedo
5 reticularis, macular degeneration, microscopic polyangiitis, morbus bechterev, motor neuron disorders, mucous membrane pemphigoid, multiple organ failure, myasthenia gravis, myelodysplastic syndrome, myocarditis, nerve root disorders, neuropathy, non-A non-B hepatitis, optic neuritis, osteolysis, ovarian cancer, pauciarticular JRA, peripheral artery occlusive disease (PAOD), peripheral vascular disease (PVD), peripheral artery disease (PAD), phlebitis,
10 polyarteritis nodosa (or periarteritis nodosa), polychondritis, polymyalgia rheumatica, poliosis, polyarticular JRA, polyendocrine deficiency syndrome, polymyositis, polymyalgia rheumatica (PMR), post-pump syndrome, primary Parkinsonism, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma), prostatitis, pure red cell aplasia, primary adrenal insufficiency, recurrent neuromyelitis optica, restenosis, rheumatic heart disease, sapho
15 (synovitis, acne, pustulosis, hyperostosis, and osteitis), scleroderma, secondary amyloidosis, shock lung, scleritis, sciatica, secondary adrenal insufficiency, silicone associated connective tissue disease, sneddon-wilkinson dermatosis, spondylitis ankylosans, Stevens-Johnson syndrome (SJS), systemic inflammatory response syndrome, temporal arteritis, toxoplasmic retinitis, toxic epidermal necrolysis, transverse myelitis, TRAPS (tumor necrosis factor receptor, type 1 allergic
20 reaction, type II diabetes, urticaria, usual interstitial pneumonia (UIP), vasculitis, vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), wet macular degeneration, wound healing, yersinia and salmonella associated arthropathy.

In an embodiment, diseases that can be treated or diagnosed with the compositions and methods disclosed herein include, but are not limited to, primary and metastatic cancers,
25 including carcinomas of breast, colon, rectum, lung, oropharynx, hypopharynx, esophagus, stomach, pancreas, liver, gallbladder and bile ducts, small intestine, urinary tract (including kidney, bladder and urothelium), female genital tract (including cervix, uterus, and ovaries as well as choriocarcinoma and gestational trophoblastic disease), male genital tract (including prostate, seminal vesicles, testes and germ cell tumors), endocrine glands (including the thyroid, adrenal, and pituitary glands), and skin, as well as hemangiomas, melanomas, sarcomas
30 (including those arising from bone and soft tissues as well as Kaposi's sarcoma), tumors of the brain, nerves, eyes, and meninges (including astrocytomas, gliomas, glioblastomas, retinoblastomas, neuromas, neuroblastomas, Schwannomas, and meningiomas), solid tumors arising from hematopoietic malignancies such as leukemias, and lymphomas (both Hodgkin's and
35 non-Hodgkin's lymphomas).

The DVD-IgS disclosed herein may also treat one or more of the following diseases:

Acute coronary syndromes, Acute Idiopathic Polyneuritis, Acute Inflammatory Demyelinating
 Polyradiculoneuropathy, Acute ischemia, Adult Still's Disease, Alopecia areata, Anaphylaxis,
 Anti-Phospholipid Antibody Syndrome, Aplastic anemia, Arteriosclerosis, Atopic eczema,
 5 Atopic dermatitis, Autoimmune dermatitis, Autoimmune disorder associated with Streptococcus
 infection, Autoimmune hearingloss, Autoimmune Lymphoproliferative Syndrome (ALPS),
 Autoimmune myocarditis, autoimmune thrombocytopenia (AITP), Blepharitis, Bronchiectasis,
 Bullous pemphigoid, Cardiovascular Disease, Catastrophic Antiphospholipid Syndrome, Celiac
 Disease, Cervical Spondylosis, Chronic ischemia, Cicatricial pemphigoid, Clinically isolated
 10 Syndrome (CIS) with Risk for Multiple Sclerosis, Conjunctivitis, Childhood Onset Psychiatric
 Disorder, Chronic obstructive pulmonary disease (COPD), Dacryocystitis, dermatomyositis,
 Diabetic retinopathy, Diabetes mellitus, Disk herniation, Disk prolaps, Drug induced immune
 hemolytic anemia, Endocarditis, Endometriosis, endophthalmitis, , Episcleritis, Erythema
 multiforme, erythema multiforme major, Gestational pemphigoid, Guillain-Barré Syndrome
 15 (GBS), Hay Fever, Hughes Syndrome , Idiopathic Parkinson's Disease, idiopathic interstitial
 pneumonia, IgE-mediated Allergy, Immune hemolytic anemia, Inclusion Body Myositis,
 Infectious ocular inflammatory disease, Inflammatory demyelinating disease, Inflammatory heart
 disease, Inflammatory kidney disease, IPF/UIP, Iritis, Keratitis, Keratoconjunctivitis sicca, Kussmaul
 disease or Kussmaul-Meier Disease, Landry's Paralysis, Langerhan's Cell Histiocytosis, Livedo
 20 reticularis, Macular Degeneration, malignancies, Microscopic Polyangiitis, Morbus Bechterev,
 Motor Neuron Disorders, Mucous membrane pemphigoid, Multiple Organ failure, Myasthenia
 Gravis, Myelodysplastic Syndrome, Myocarditis, Nerve Root Disorders, Neuropathy, Non-A
 Non-B Hepatitis, Optic Neuritis, Osteolysis, Ovarian cancer, Pauciarticular JRA, peripheral
 artery occlusive disease (PAOD), peripheral vascular disease (PVD), peripheral artery disease
 25 (PAD), Phlebitis, Polyarteritis nodosa (or periarteritis nodosa), Polychondritis, Polymyalgia
 Rheumatica, Poliosis, Polyarticular JRA, Polyendocrine Deficiency Syndrome, Polymyositis,
 polymyalgia rheumatica (PMR), Post-Pump Syndrome, primary parkinsonism, prostate and rectal
 cancer and hematopoietic malignancies (leukemia and lymphoma), Prostatitis, Pure red cell
 aplasia, Primary Adrenal Insufficiency, Recurrent Neuromyelitis Optica, Restenosis, Rheumatic
 30 heart disease, SAPHO (synovitis, acne, pustulosis, hyperostosis, and osteitis), Scleroderma,
 Secondary Amyloidosis, Shock lung, Scleritis, Sciatica, Secondary Adrenal Insufficiency,
 Silicone associated connective tissue disease, Sneddon-Wilkinson Dermatositis, spondylitis
 ankylosans, Stevens-Johnson Syndrome (SJS), Systemic inflammatory response syndrome,
 Temporal arteritis, toxoplasmic retinitis, toxic epidermal necrolysis, Transverse myelitis, TRAPS
 35 (Tumor Necrosis Factor Receptor, Type I allergic reaction, Type II Diabetes, Urticaria, Usual

interstitial pneumonia (UIP), Vasculitis, Vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), Wet macular degeneration, and Wound healing.

In an embodiment, the binding proteins disclosed herein or antigen-binding portions thereof, are used to treat cancer or in the prevention of metastases from the tumors described
 5 herein either when used alone or in combination with radiotherapy and/or other chemotherapeutic agents.

In another aspect, a method of treating a patient suffering from a disorder comprising the step of administering any one of the binding proteins disclosed herein before, concurrent, or after the administration of a second agent, as discussed herein is provided. In a particular embodiment
 10 the second agent is budenoside, epidermal growth factor, corticosteroids, cyclosporin, sulfasalazine, aminosalicylates, 6-mercaptopurine, azathioprine, metronidazole, lipoxygenase inhibitors, mesalamine, olsalazine, balsalazide, antioxidants, thromboxane inhibitors, IL-1 receptor antagonists, anti-IL-1 β mAbs, anti-IL-6 or IL-6 receptor mAbs, growth factors, elastase inhibitors, pyridinyl-imidazole compounds, antibodies or agonists of TNF, LT, IL-1, IL-2, IL-6,
 15 IL-7, IL-8, IL-12, IL-13, IL-15, IL-16, IL-18, IL-23, EMAP-II, GM-CSF, FGF, and PDGF, antibodies of CD2, CD3, CD4, CD8, CD-19, CD25, CD28, CD30, CD40, CD45, CD69, CD90 or their ligands, methotrexate, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, ibuprofen, corticosteroids, prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, IRAK, NIK, IKK, p38,
 20 MAP kinase inhibitors, IL-1 β converting enzyme inhibitors, TNF α converting enzyme inhibitors, T-cell signalling inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors, soluble p55 TNF receptor, soluble p75 TNF receptor, sIL-1RI, sIL-1RII, sIL-6R, antiinflammatory cytokines, IL-4, IL-10, IL-11, IL-13, or TGF β .

In a particular embodiment the pharmaceutical compositions disclosed herein are administered to the patient by at least one mode selected from parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular,
 intracartilaginous, intracavitary, intracelical, intracerebellar, intracerebroventricular, intracolonic,
 intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac,
 30 intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal, buccal, sublingual, intranasal, and transdermal.

In one embodiment, at least one anti-idiotypic antibody to at least one binding protein disclosed herein is provided. The anti-idiotypic antibody includes any protein or peptide containing molecule that comprises at least a portion of an immunoglobulin molecule such as, but not limited to, at least one complementarily determining region (CDR) of a heavy or light chain or a ligand binding portion thereof, a heavy chain or light chain variable region, a heavy chain or light chain constant region, a framework region, or any portion thereof, that can be incorporated into a binding protein disclosed herein.

Brief Description of the Drawings

Figure 1A is a schematic representation of Dual Variable Domain immunoglobulin (DVD-Ig) constructs and shows the strategy for generation of a DVD-Ig from two parent antibodies;

Figure 1B, is a schematic representation of constructs DVD1-Ig, DVD2-Ig, and two chimeric mono-specific antibodies from hybridoma clones 2D13.E3 (anti-IL-1 α) and 13F5.G5 (anti-IL-1 β).

Detailed Description

Multivalent and/or multispecific binding proteins capable of binding two or more antigens are provided. Specifically, dual variable domain immunoglobulins (DVD-Ig), and pharmaceutical compositions thereof, as well as nucleic acids, recombinant expression vectors and host cells for making such DVD-Igs are provided. Methods of using the DVD-Igs provided herein to detect specific antigens, either in vitro or in vivo are also provided.

Unless otherwise defined herein, scientific and technical terms used herein shall have the meanings that are commonly understood by those of ordinary skill in the art. The meaning and scope of the terms should be clear, however, in the event of any latent ambiguity, definitions provided herein take precedent over any dictionary or extrinsic definition. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. In this application, the use of "or" means "and/or" unless stated otherwise. Furthermore, the use of the term "including", as well as other forms, such as "includes" and "included", is not limiting. Also, terms such as "element" or "component" encompass both elements and components comprising one unit and elements and components that comprise more than one subunit unless specifically stated otherwise.

Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid

chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques provided herein are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless
5 otherwise indicated. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclatures used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques are
10 used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

Select terms are defined below:

The term "polypeptide" as used herein, refers to any polymeric chain of amino acids. The terms "peptide" and "protein" are used interchangeably with the term polypeptide and also refer
15 to a polymeric chain of amino acids. The term "polypeptide" encompasses native or artificial proteins, protein fragments and polypeptide analogs of a protein sequence. A polypeptide may be monomeric or polymeric. Use of "polypeptide" herein is intended to encompass polypeptide and fragments and variants (including fragments of variants) thereof, unless otherwise contradicted by context. For an antigenic polypeptide, a fragment of polypeptide optionally contains at least
20 one contiguous or nonlinear epitope of polypeptide. The precise boundaries of the at least one epitope fragment can be confirmed using ordinary skill in the art. The fragment comprises at least about 5 contiguous amino acids, such as at least about 10 contiguous amino acids, at least about 15 contiguous amino acids, or at least about 20 contiguous amino acids. A variant of polypeptide is as described herein.

The term "isolated protein" or "isolated polypeptide" is a protein or polypeptide that by virtue of its origin or source of derivation is not associated with naturally associated components that accompany it in its native state; is substantially free of other proteins from the same species;
25 is expressed by a cell from a different species; or does not occur in nature. Thus, a polypeptide that is chemically synthesized or synthesized in a cellular system different from the cell from which it naturally originates will be "isolated" from its naturally associated components. A
30 protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well known in the art.

The term "recovering" as used herein, refers to the process of rendering a chemical species such as a polypeptide substantially free of naturally associated components by isolation, e.g., using protein purification techniques well known in the art.

5 "Biological activity" as used herein, refers to any one or more inherent biological properties of a molecule (whether present naturally as found in vivo, or provided or enabled by recombinant means). Biological properties include but are not limited to binding receptor; induction of cell proliferation, inhibiting cell growth, inductions of other cytokines, induction of apoptosis, and enzymatic activity. Biological activity also includes activity of an Ig molecule.

10 The terms "specific binding" or "specifically binding", as used herein, in reference to the interaction of an antibody, a protein, or a peptide with a second chemical species, mean that the interaction is dependent upon the presence of a particular structure (e.g., an antigenic determinant or epitope) on the chemical species; for example, an antibody recognizes and binds to a specific protein structure rather than to proteins generally. If an antibody is specific for epitope "A", the presence of a molecule containing epitope A (or free, unlabeled A), in a reaction containing
15 labeled "A" and the antibody, will reduce the amount of labeled A bound to the antibody.

The term "antibody", as used herein, broadly refers to any immunoglobulin (Ig) molecule comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains, or any functional fragment, mutant, variant, or derivation thereof, which retains the essential epitope binding features of an Ig molecule. Such mutant, variant, or derivative antibody formats are
20 known in the art. Nonlimiting embodiments of which are discussed below.

In a full-length antibody, each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain
25 constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3,
30 CDR3, FR4. Immunoglobulin molecules can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG 1, IgG2, IgG 3, IgG4, IgA 1 and IgA2) or subclass.

The term "Fc region" is used to define the C-terminal region of an immunoglobulin heavy chain, which may be generated by papain digestion of an intact antibody. The Fc region

may be a native sequence Fc region or a variant Fc region. The Fc region of an immunoglobulin generally comprises two constant domains, a CH2 domain and a CH3 domain, and optionally comprises a CH4 domain. Replacements of amino acid residues in the Fc portion to alter antibody effector function are known in the art (Winter, *et al.* US Patent Nos 5,648,260 and 5,624,821).

5 The Fc portion of an antibody mediates several important effector functions e.g., cytokine induction, ADCC, phagocytosis, complement dependent cytotoxicity (CDC) and half-life/clearance rate of antibody and antigen-antibody complexes. In some cases these effector functions are desirable for therapeutic antibody but in other cases might be unnecessary or even deleterious, depending on the therapeutic objectives. Certain human IgG isotypes, particularly

10 IgG1 and IgG3, mediate ADCC and CDC via binding to FcγRs and complement C1q, respectively. Neonatal Fc receptors (FcRn) are the critical components determining the circulating half-life of antibodies. In still another embodiment at least one amino acid residue is replaced in the constant region of the antibody, for example the Fc region of the antibody, such that effector functions of the antibody are altered. The dimerization of two identical heavy chains

15 of an immunoglobulin is mediated by the dimerization of CH3 domains and is stabilized by the disulfide bonds within the hinge region (Huber *et al.* *Nature*; 264: 415-20; Thies *et al.* 1999 *J Mol Biol*; 293: 67-79.). Mutation of cysteine residues within the hinge regions to prevent heavy chain-heavy chain disulfide bonds will destabilize dimerization of CH3 domains. Residues responsible for CH3 dimerization have been identified (Dall'Acqua 1998 *Biochemistry* 37: 9266-73.).

20 Therefore, it is possible to generate a monovalent half-Ig. Interestingly, these monovalent half Ig molecules have been found in nature for both IgG and IgA subclasses (Seligman 1978 *Ann Immunol* 129: 855-70; Biewenga *et al.* 1983 *Clin Exp Immunol* 51: 395-400). The stoichiometry of FcRn: Ig Fc region has been determined to be 2:1 (West *et al.* 2000 *Biochemistry* 39: 9698-708), and half Fc is sufficient for mediating FcRn binding (Kim *et al.* 1994 *Eur J Immunol*; 24: 542-548.). Mutations to disrupt the dimerization of CH3 domain may not have greater adverse effect on its FcRn binding as the residues important for CH3 dimerization are located on the inner

25 interface of CH3 β sheet structure, whereas the region responsible for FcRn binding is located on the outside interface of CH2-CH3 domains. However the half Ig molecule may have certain advantage in tissue penetration due to its smaller size than that of a regular antibody. In one

30 embodiment at least one amino acid residue is replaced in the constant region of the binding protein, for example the Fc region, such that the dimerization of the heavy chains is disrupted, resulting in half DVD Ig molecules. The anti-inflammatory activity of IgG is completely dependent on sialylation of the N-linked glycan of the IgG Fc fragment. The precise glycan requirements for anti-inflammatory activity has been determined, such that an appropriate IgG1

35 Fc fragment can be created, thereby generating a fully recombinant, sialylated IgG1 Fc with greatly enhanced potency (Anthony, R.M., *et al.* (2008) *Science* 320:373-376).

The term "antigen-binding portion" of an antibody (or simply "antibody portion"), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen. It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Such antibody embodiments may also be

5 bispecific, dual specific, or multi-specific formats; specifically binding to two or more different antigens. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment comprising the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments

10 linked by a disulfide bridge at the hinge region; (iii) a Fd fragment comprising the VH and CH1 domains; (iv) a Fv fragment comprising the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.*, (1989) *Nature* 341:544-546, Winter *et al.*, PCT publication WO 90/05144 A1), which comprises a single variable domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a

15 synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. Other forms of single chain antibodies, such as diabodies are

20 also encompassed. Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see *e.g.*, Holliger, P., *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R.J., *et al.* (1994)

25 *Structure* 2:1121-1123). Such antibody binding portions are known in the art (Kontermann and Dubel eds., *Antibody Engineering* (2001) Springer-Verlag, New York, 790 pp. (ISBN 3-540-41354-5). In addition single chain antibodies also include "linear antibodies" comprising a pair of tandem Fv segments (VH-CH1-VH-CH1) which, together with complementary light chain polypeptides, form a pair of antigen binding regions (Zapata *et al.* *Protein Eng.* 8(10):1057-1062

30 (1995); and US Patent No. 5,641,870).

The term "multivalent binding protein" is used throughout this specification to denote a binding protein comprising two or more antigen binding sites. In an embodiment, the multivalent binding protein is engineered to have the three or more antigen binding sites, and is generally not a naturally occurring antibody. The term "multispecific binding protein" refers to a binding

35 protein capable of binding two or more related or unrelated targets. Dual variable domain (DVD)

binding proteins provided herein comprise two or more antigen binding sites and are tetravalent or multivalent binding proteins. In certain embodiments, the binding proteins are monospecific, i.e., capable of binding one antigen or multispecific, i.e. capable of binding two or more antigens. DVD binding proteins comprising two heavy chain polypeptides and two light chain polypeptides are referred to as DVD-Igs. Each half of a DVD-Ig comprises a heavy chain polypeptide, and a light chain polypeptide, and two antigen binding sites. Each binding site comprises a heavy chain variable domain and a light chain variable domain with a total of 6 CDRs involved in antigen binding per antigen binding site.

The term "bispecific antibody", as used herein, refers to full-length antibodies that are generated by quadroma technology (see Milstein, C. and A.C. Cuello, *Nature*, 1983. 305(5934): p. 537-40), by chemical conjugation of two different monoclonal antibodies (see Staerz, U.D., et al., *Nature*, 1985. 314(6012): p. 628-31), or by knob-into-hole or similar approaches which introduces mutations in the Fc region (see Holliger, P., T. Prospero, and G. Winter, *Proc Natl Acad Sci U S A*, 1993. 90(14): p. 6444-8.18), resulting in multiple different immunoglobulin species of which only one is the functional bispecific antibody. By molecular function, a bispecific antibody binds one antigen (or epitope) on one of its two binding arms (one pair of HC/LC), and binds a different antigen (or epitope) on its second arm (a different pair of HC/LC). By this definition, a bispecific antibody has two distinct antigen binding arms (in both specificity and CDR sequences), and is monovalent for each antigen it binds to.

The term "dual-specific antibody", as used herein, refers to full-length antibodies that can bind two different antigens (or epitopes) in each of its two binding arms (a pair of HC/LC) (see PCT publication WO 02/02773). Accordingly a dual-specific binding protein has two identical antigen binding arms, with identical specificity and identical CDR sequences, and is bivalent for each antigen it binds to.

A "functional antigen binding site" of a binding protein is one that is capable of binding a target antigen. The antigen binding affinity of the antigen binding site is not necessarily as strong as the parent antibody from which the antigen binding site is derived, but the ability to bind antigen must be measurable using any one of a variety of methods known for evaluating antibody binding to an antigen. Moreover, the antigen binding affinity of each of the antigen binding sites of a multivalent antibody herein need not be quantitatively the same.

The term "cytokine" is a generic term for proteins released by one cell population, which act on another cell population as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines

are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor-alpha and -beta; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF-alpha; platelet-growth factor; placental growth factor, transforming growth factors (TGFs) such as TGF-alpha and TGF-beta; insulin-like growth factor-1 and -11; erythropoietin (EPO); osteoinductive factors; interferons such as interferon-alpha, -beta and -gamma colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-15, IL-18, IL-21, IL-22, IL-23, IL-33; a tumor necrosis factor such as TNF-alpha or TNF-beta; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

The term "linker" is used to denote polypeptides comprising two or more amino acid residues joined by peptide bonds and are used to link one or more antigen binding portions. Such linker polypeptides are well known in the art (see *e.g.*, Holliger, P., *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R.J., *et al.* (1994) *Structure* 2:1121-1123). Exemplary linkers include, but are not limited to, AKTTPKLEEGEFSEAR (SEQ ID NO: 1); AKTTPKLEEGEFSEARV (SEQ ID NO: 2); AKTTPKLGG (SEQ ID NO: 3); SAKTTPKLGG (SEQ ID NO: 4); SAKTTP (SEQ ID NO: 5); RADAAP (SEQ ID NO: 6); RADAAPTVS (SEQ ID NO: 7); RADAAAAGGPGS (SEQ ID NO: 8); RADAAAA(G₄S)₄ (SEQ ID NO: 9); SAKTTPKLEEGEFSEARV (SEQ ID NO: 10); ADAAP (SEQ ID NO: 11); ADAAPTVSIFPP (SEQ ID NO: 12); TVAAP (SEQ ID NO: 13); TVAAPSVFIFPP (SEQ ID NO: 14); QPKAAP (SEQ ID NO: 15); QPKAAPSVTLFPP (SEQ ID NO: 16); AKTTPP (SEQ ID NO: 17); AKTTPPSVTPLAP (SEQ ID NO: 18); AKTTAP (SEQ ID NO: 19); AKTTAPSVYPLAP (SEQ ID NO: 20); ASTKGP (SEQ ID NO: 21); ASTKGPSVFPLAP (SEQ ID NO: 22); GGGGSGGGGSGGGGS (SEQ ID NO: 23); GENKVEYAPALMALS (SEQ ID NO: 24); GPAKELTPLKEAKVS (SEQ ID NO: 25); and GHEAAAVMQVQYPAS (SEQ ID NO: 26).

An immunoglobulin constant domain refers to a heavy or light chain constant domain. Human IgG heavy chain and light chain constant domain amino acid sequences are known in the art.

The term "monoclonal antibody" or "mAb" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigen. Furthermore, in contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each mAb is directed against a single determinant on the antigen. The modifier "monoclonal" is not to be construed as requiring production of the antibody by any particular method.

The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies provided herein may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo), for example in the CDRs and in particular CDR3. However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described further in Section II C, below), antibodies isolated from a recombinant, combinatorial human antibody library (Hoogenboom H.R. (1997) TIB Tech. 15:62-70; Azzazy H., and Highsmith W.E. (2002) Clin. Biochem. 35:425-445; Gavilondo J.V., and Larrick J.W. (2002) BioTechniques 29:128-145; Hoogenboom H., and Chames P. (2000) Immunology Today 21:371-378), antibodies isolated from an animal (e.g., a mouse) that is transgenic for human immunoglobulin genes (see, Taylor, L. D., et al. (1992) Nucl. Acids Res. 20:6287-6295; Kellermann S-A. and Green L.L. (2002) Current Opinion in Biotechnology 13:593-597; Little M. et al. (2000) Immunology Today 21:364-370) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies are subjected to in vitro mutagenesis (or, when an animal transgenic for human Ig sequences is used, in vivo somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related

to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire in vivo.

An "affinity matured" antibody is an antibody with one or more alterations in one or more CDRs thereof which result an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). Exemplary affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by procedures known in the art. Marks et al. *BidlTechnology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas et al. *Proc Nat. Acad. Sci, USA* 91:3809-3813 (1994); Schier et al. *Gene* 169:147- 155 (1995); Yelton et al. *J. Immunol.* 155:1994-2004 (1995); Jackson et al., *J. Immunol.* 154(7):3310-9 (1995); Hawkins et al, *J. Mol. BioL* 226:889-896 (1992) and selective mutation at selective mutagenesis positions, contact or hypermutation positions with an activity enhancing amino acid residue as described in US patent US 6914128B1.

The term "chimeric antibody" refers to antibodies which comprise heavy and light chain variable region sequences from one species and constant region sequences from another species, such as antibodies having murine heavy and light chain variable regions linked to human constant regions.

The term "CDR-grafted antibody" refers to antibodies which comprise heavy and light chain variable region sequences from one species but in which the sequences of one or more of the CDR regions of VH and/or VL are replaced with CDR sequences of another species, such as antibodies having murine heavy and light chain variable regions in which one or more of the murine CDRs (e.g., CDR3) has been replaced with human CDR sequences.

The term "humanized antibody" refers to antibodies which comprise heavy and light chain variable region sequences from a non-human species (e.g., a mouse) but in which at least a portion of the VH and/or VL sequence has been altered to be more "human-like", i.e., more similar to human germline variable sequences. One type of humanized antibody is a CDR-grafted antibody, in which human CDR sequences are introduced into non-human VH and VL sequences to replace the corresponding nonhuman CDR sequences. Also "humanized antibody" is an antibody or a variant, derivative, analog or fragment thereof which immunospecifically binds to an antigen of interest and which comprises a framework (FR) region having substantially the amino acid sequence of a human antibody and a complementary determining region (CDR) having substantially the amino acid sequence of a non-human antibody. As used herein, the term

"substantially" in the context of a CDR refers to a CDR having an amino acid sequence at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the amino acid sequence of a non-human antibody CDR. A humanized antibody comprises substantially all of at least one, and typically two, variable domains (Fab, Fab', F(ab')₂, FabC, Fv) in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin (i.e., donor antibody) and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. In an embodiment, a humanized antibody also comprises at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. In some embodiments, a humanized antibody contains both the light chain as well as at least the variable domain of a heavy chain. The antibody also may include the CH1, hinge, CH2, CH3, and CH4 regions of the heavy chain. In some embodiments, a humanized antibody only contains a humanized light chain. In some embodiments, a humanized antibody only contains a humanized heavy chain. In specific embodiments, a humanized antibody only contains a humanized variable domain of a light chain and/or humanized heavy chain.

The terms "Kabat numbering", "Kabat definitions" and "Kabat labeling" are used interchangeably herein. These terms, which are recognized in the art, refer to a system of numbering amino acid residues which are more variable (*i.e.* hypervariable) than other amino acid residues in the heavy and light chain variable regions of an antibody, or an antigen binding portion thereof (Kabat *et al.* (1971) *Ann. NY Acad. Sci.* 190:382-391 and, Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). For the heavy chain variable region, the hypervariable region ranges from amino acid positions 31 to 35 for CDR1, amino acid positions 50 to 65 for CDR2, and amino acid positions 95 to 102 for CDR3. For the light chain variable region, the hypervariable region ranges from amino acid positions 24 to 34 for CDR1, amino acid positions 50 to 56 for CDR2, and amino acid positions 89 to 97 for CDR3.

As used herein, the term "CDR" refers to the complementarity determining region within antibody variable sequences. There are three CDRs in each of the variable regions of the heavy chain and the light chain, which are designated CDR1, CDR2 and CDR3, for each of the variable regions. The term "CDR set" as used herein refers to a group of three CDRs that occur in a single variable region capable of binding the antigen. The exact boundaries of these CDRs have been defined differently according to different systems. The system described by Kabat (Kabat *et al.*, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, Md. (1987) and (1991)) not only provides an unambiguous residue numbering system applicable to any variable region of an antibody, but also provides precise residue boundaries defining the

three CDRs. These CDRs may be referred to as Kabat CDRs. Chothia and coworkers (Chothia & Lesk, *J. Mol. Biol.* 196:901-917 (1987) and Chothia et al., *Nature* 342:877-883 (1989)) found that certain sub-portions within Kabat CDRs adopt nearly identical peptide backbone conformations, despite having great diversity at the level of amino acid sequence. These sub-portions were designated as L1, L2 and L3 or H1, H2 and H3 where the "L" and the "H" designates the light chain and the heavy chains regions, respectively. These regions may be referred to as Chothia CDRs, which have boundaries that overlap with Kabat CDRs. Other boundaries defining CDRs overlapping with the Kabat CDRs have been described by Padlan (*FASEB J.* 9:133-139 (1995)) and MacCallum (*J Mol Biol* 262(5):732-45 (1996)). Still other CDR boundary definitions may not strictly follow one of the herein systems, but will nonetheless overlap with the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. The methods used herein may utilize CDRs defined according to any of these systems, although certain embodiments use Kabat or Chothia defined CDRs.

As used herein, the term "framework" or "framework sequence" refers to the remaining sequences of a variable region minus the CDRs. Because the exact definition of a CDR sequence can be determined by different systems, the meaning of a framework sequence is subject to correspondingly different interpretations. The six CDRs (CDR-L1, -L2, and -L3 of light chain and CDR-H1, -H2, and -H3 of heavy chain) also divide the framework regions on the light chain and the heavy chain into four sub-regions (FR1, FR2, FR3 and FR4) on each chain, in which CDR1 is positioned between FR1 and FR2, CDR2 between FR2 and FR3, and CDR3 between FR3 and FR4. Without specifying the particular sub-regions as FR1, FR2, FR3 or FR4, a framework region, as referred by others, represents the combined FR's within the variable region of a single, naturally occurring immunoglobulin chain. As used herein, a FR represents one of the four sub-regions, and FRs represents two or more of the four sub-regions constituting a framework region.

As used herein, the term "germline antibody gene" or "gene fragment" refers to an immunoglobulin sequence encoded by non-lymphoid cells that have not undergone the maturation process that leads to genetic rearrangement and mutation for expression of a particular immunoglobulin. (See, e.g., Shapiro et al., *Crit. Rev. Immunol.* 22(3): 183-200 (2002); Marchalonis et al., *Adv Exp Med Biol.* 484:13-30 (2001)). One of the advantages provided by various embodiments provided herein stems from the recognition that germline antibody genes are more likely than mature antibody genes to conserve essential amino acid sequence structures

characteristic of individuals in the species, hence less likely to be recognized as from a foreign source when used therapeutically in that species.

As used herein, the term "neutralizing" refers to counteracting the biological activity of an antigen when a binding protein specifically binds the antigen. In an embodiment, the neutralizing binding protein binds the cytokine and reduces its biological activity by at least about 20%, 40%, 60%, 80%, 85% or more.


The term "activity" includes activities such as the binding specificity and affinity of a binding protein for two or more antigens.

The term "epitope" includes any polypeptide determinant capable of specific binding to an immunoglobulin or T-cell receptor. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three dimensional structural characteristics, and/or specific charge characteristics. An epitope is a region of an antigen that is bound by an antibody. In certain embodiments, an antibody is said to specifically bind an antigen when it recognizes its target antigen in a complex mixture of proteins and/or macromolecules. Antibodies are said to "bind to the same epitope" if the antibodies cross-compete (one prevents the binding or modulating effect of the other). In addition structural definitions of epitopes (overlapping, similar, identical) are informative, but functional definitions are often more relevant as they encompass structural (binding) and functional (modulation, competition) parameters.


The term "surface plasmon resonance", as used herein, refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore® system (BIAcore International AB, a GE Healthcare company, Uppsala, Sweden and Piscataway, NJ). For further descriptions, see Jönsson, U., *et al.* (1993) *Ann. Biol. Clin.* 51:19-26; Jönsson, U., *et al.* (1991) *Biotechniques* 11:620-627; Johnsson, B., *et al.* (1995) *J. Mol. Recognit.* 8:125-131; and Johnsson, B., *et al.* (1991) *Anal. Biochem.* 198:268-277.

The term " K_{on} ", as used herein, is intended to refer to the on rate constant for association of a binding protein (e.g., an antibody) to the antigen to form the, e.g., antibody/antigen complex as is known in the art. The " K_{on} " also is known by the terms "association rate constant", or "ka", as used interchangeably herein. This value indicating the binding rate of an antibody to its target

antigen or the rate of complex formation between an antibody and antigen also is shown by the equation below:



The term " K_{off} ", as used herein, is intended to refer to the off rate constant for dissociation, or "dissociation rate constant", of a binding protein (e.g., an antibody) from the, e.g., antibody/antigen complex as is known in the art. This value indicates the dissociation rate of an antibody from its target antigen or separation of Ab-Ag complex over time into free antibody and antigen as shown by the equation below:



The term " K_D ", as used herein, is intended to refer to the "equilibrium dissociation constant", and refers to the value obtained in a titration measurement at equilibrium, or by dividing the dissociation rate constant (k_{off}) by the association rate constant (k_{on}). The association rate constant, the dissociation rate constant and the equilibrium dissociation constant are used to represent the binding affinity of an antibody to an antigen. Methods for determining association and dissociation rate constants are well known in the art. Using fluorescence-based techniques offers high sensitivity and the ability to examine samples in physiological buffers at equilibrium. Other experimental approaches and instruments such as a BIAcore® (biomolecular interaction analysis) assay can be used (e.g., instrument available from BIAcore International AB, a GE Healthcare company, Uppsala, Sweden). Additionally, a KinExA® (Kinetic Exclusion Assay) assay, available from Sapidyne Instruments (Boise, Idaho) can also be used.

"Label" and "detectable label" mean a moiety attached to a specific binding partner, such as an antibody or an analyte, e.g., to render the reaction between members of a specific binding pair, such as an antibody and an analyte, detectable, and the specific binding partner, e.g., antibody or analyte, so labeled is referred to as "detectably labeled." Thus, the term "labeled binding protein" as used herein, refers to a protein with a label incorporated that provides for the identification of the binding protein. In an embodiment, the label is a detectable marker that can produce a signal that is detectable by visual or instrumental means, e.g., incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I , ^{177}Lu , ^{166}Ho , or ^{153}Sm); chromogens, fluorescent labels

(e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase, luciferase, alkaline phosphatase); chemiluminescent markers; biotinyl groups; predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags); and magnetic agents, such as gadolinium chelates. Representative examples of labels commonly employed for immunassays include moieties that produce light, e.g., acridinium compounds, and moieties that produce fluorescence, e.g., fluorescein. Other labels are described herein. In this regard, the moiety itself may not be detectably labeled but may become detectable upon reaction with yet another moiety. Use of "detectably labeled" is intended to encompass the latter type of detectable labeling.

The term "conjugate" refers to a binding protein, such as an antibody, chemically linked to a second chemical moiety, such as a therapeutic or cytotoxic agent. The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials. In an embodiment, the therapeutic or cytotoxic agents include, but are not limited to, pertussis toxin, taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. When employed in the context of an immunoassay, the conjugate antibody may be a detectably labeled antibody used as the detection antibody.

The terms "crystal" and "crystallized" as used herein, refer to a binding protein (e.g., an antibody), or antigen binding portion thereof, that exists in the form of a crystal. Crystals are one form of the solid state of matter, which is distinct from other forms such as the amorphous solid state or the liquid crystalline state. Crystals are composed of regular, repeating, three-dimensional arrays of atoms, ions, molecules (e.g., proteins such as antibodies), or molecular assemblies (e.g., antigen/antibody complexes). These three-dimensional arrays are arranged according to specific mathematical relationships that are well-understood in the field. The fundamental unit, or building block, that is repeated in a crystal is called the asymmetric unit. Repetition of the asymmetric unit in an arrangement that conforms to a given, well-defined crystallographic symmetry provides the "unit cell" of the crystal. Repetition of the unit cell by regular translations in all three dimensions provides the crystal. See Giege, R. and Ducruix, A. Barrett, *Crystallization of Nucleic Acids and Proteins, a Practical Approach*, 2nd ea., pp. 20 1-16, Oxford University Press, New York, New York, (1999)."

The term "polynucleotide" means a polymeric form of two or more nucleotides, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

5 The term "isolated polynucleotide" shall mean a polynucleotide (e.g., of genomic, cDNA, or synthetic origin, or some combination thereof) that, by virtue of its origin, the "isolated polynucleotide" is not associated with all or a portion of a polynucleotide with which the "isolated polynucleotide" is found in nature; is operably linked to a polynucleotide that it is not linked to in nature; or does not occur in nature as part of a larger sequence.

10 The term "vector", is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated
15 into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification,
20 "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, embodiments include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

25 The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences. "Operably linked" sequences include both expression control sequences that are contiguous with the gene of interest
30 and expression control sequences that act in *trans* or at a distance to control the gene of interest. The term "expression control sequence" as used herein refers to polynucleotide sequences which are necessary to effect the expression and processing of coding sequences to which they are ligated. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and

polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term "control sequences" is intended to include components whose presence is essential for expression and processing, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

10 "Transformation", refers to any process by which exogenous DNA enters a host cell. Transformation may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the host cell being transformed and may include, but is not limited to, viral infection, 15 electroporation, lipofection, and particle bombardment. Such "transformed" cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome. They also include cells which transiently express the inserted DNA or RNA for limited periods of time.

The term "recombinant host cell" (or simply "host cell"), is intended to refer to a cell into 20 which exogenous DNA has been introduced. In an embodiment, the host cell comprises two or more (e.g., multiple) nucleic acids encoding antibodies, such as the host cells described in US Patent No. 7,262,028, for example. Such terms are intended to refer not only to the particular subject cell, but also to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may 25 not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein. In an embodiment, host cells include prokaryotic and eukaryotic cells selected from any of the Kingdoms of life. In another embodiment, eukaryotic cells include protist, fungal, plant and animal cells. In another embodiment, host cells include but are not limited to the prokaryotic cell line E.Coli; mammalian cell lines CHO, HEK 293, COS, NS0, SP2 30 and PER.C6; the insect cell line Sf9; and the fungal cell *Saccharomyces cerevisiae*.

Standard techniques may be used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection). Enzymatic reactions and purification techniques may be performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and

procedures may be generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. See e.g., Sambrook et al. *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)).

5 “Transgenic organism”, as known in the art, refers to an organism having cells that contain a transgene, wherein the transgene introduced into the organism (or an ancestor of the organism) expresses a polypeptide not naturally expressed in the organism. A “transgene” is a DNA construct, which is stably and operably integrated into the genome of a cell from which a transgenic organism develops, directing the expression of an encoded gene product in one or
10 more cell types or tissues of the transgenic organism.

The term “regulate” and “modulate” are used interchangeably, and, as used herein, refers to a change or an alteration in the activity of a molecule of interest (e.g., the biological activity of a cytokine). Modulation may be an increase or a decrease in the magnitude of a certain activity or function of the molecule of interest. Exemplary activities and functions of a molecule include, but
15 are not limited to, binding characteristics, enzymatic activity, cell receptor activation, and signal transduction.

Correspondingly, the term “modulator” is a compound capable of changing or altering an activity or function of a molecule of interest (e.g., the biological activity of a cytokine). For example, a modulator may cause an increase or decrease in the magnitude of a certain activity or
20 function of a molecule compared to the magnitude of the activity or function observed in the absence of the modulator. In certain embodiments, a modulator is an inhibitor, which decreases the magnitude of at least one activity or function of a molecule. Exemplary inhibitors include, but are not limited to, proteins, peptides, antibodies, peptibodies, carbohydrates or small organic molecules. Peptibodies are described, e.g., in WO01/83525.

25 The term “agonist”, refers to a modulator that, when contacted with a molecule of interest, causes an increase in the magnitude of a certain activity or function of the molecule compared to the magnitude of the activity or function observed in the absence of the agonist. Particular agonists of interest may include, but are not limited to, polypeptides, nucleic acids, carbohydrates, or any other molecules that bind to the antigen.

30 The term “antagonist” or “inhibitor”, refer to a modulator that, when contacted with a molecule of interest causes a decrease in the magnitude of a certain activity or function of the molecule compared to the magnitude of the activity or function observed in the absence of the antagonist. Particular antagonists of interest include those that block or modulate the biological

or immunological activity of of the antigen. Antagonists and inhibitors of antigens may include, but are not limited to, proteins, nucleic acids, carbohydrates, or any other molecules, which bind to the antigen.

As used herein, the term "effective amount" refers to the amount of a therapy which is sufficient to reduce or ameliorate the severity and/or duration of a disorder or one or more symptoms thereof, prevent the advancement of a disorder, cause regression of a disorder, prevent the recurrence, development, onset or progression of one or more symptoms associated with a disorder, detect a disorder, or enhance or improve the prophylactic or therapeutic effect(s) of another therapy (e.g., prophylactic or therapeutic agent).

"Patient" and "subject" may be used interchangeably herein to refer to an animal, such as a mammal, including a primate (for example, a human, a monkey, and a chimpanzee), a non-primate (for example, a cow, a pig, a camel, a llama, a horse, a goat, a rabbit, a sheep, a hamster, a guinea pig, a cat, a dog, a rat, a mouse, a whale), a bird (e.g., a duck or a goose), and a shark. Preferably, the patient or subject is a human, such as a human being treated or assessed for a disease, disorder or condition, a human at risk for a disease, disorder or condition, a human having a disease, disorder or condition, and/or human being treated for a disease, disorder or condition.

The term "sample", as used herein, is used in its broadest sense. A "biological sample", as used herein, includes, but is not limited to, any quantity of a substance from a living thing or formerly living thing. Such living things include, but are not limited to, humans, mice, rats, monkeys, dogs, rabbits and other animals. Such substances include, but are not limited to, blood (e.g., whole blood), plasma, serum, urine, amniotic fluid, synovial fluid, endothelial cells, leukocytes, monocytes, other cells, organs, tissues, bone marrow, lymph nodes and spleen.

"Component," "components," and "at least one component," refer generally to a capture antibody, a detection or conjugate antibody, a control, a calibrator, a series of calibrators, a sensitivity panel, a container, a buffer, a diluent, a salt, an enzyme, a co-factor for an enzyme, a detection reagent, a pretreatment reagent/solution, a substrate (e.g., as a solution), a stop solution, and the like that can be included in a kit for assay of a test sample, such as a patient urine, serum or plasma sample, in accordance with the methods described herein and other methods known in the art. Thus, in the context of the present disclosure, "at least one component," "component," and "components" can include a polypeptide or other analyte as above, such as a composition comprising an analyte such as polypeptide, which is optionally immobilized on a solid support,

such as by binding to an anti-analyte (e.g., anti-polypeptide) antibody. Some components can be in solution or lyophilized for reconstitution for use in an assay.

“Control” refers to a composition known to not analyte (“negative control”) or to contain analyte (“positive control”). A positive control can comprise a known concentration of analyte. “Control,” “positive control,” and “calibrator” may be used interchangeably herein to refer to a composition comprising a known concentration of analyte. A “positive control” can be used to establish assay performance characteristics and is a useful indicator of the integrity of reagents (e.g., analytes).

“Predetermined cutoff” and “predetermined level” refer generally to an assay cutoff value that is used to assess diagnostic/prognostic/therapeutic efficacy results by comparing the assay results against the predetermined cutoff/level, where the predetermined cutoff/level already has been linked or associated with various clinical parameters (e.g., severity of disease, progression/nonprogression/improvement, etc.). While the present disclosure may provide exemplary predetermined levels, it is well-known that cutoff values may vary depending on the nature of the immunoassay (e.g., antibodies employed, etc.). It further is well within the ordinary skill of one in the art to adapt the disclosure herein for other immunoassays to obtain immunoassay-specific cutoff values for those other immunoassays based on this disclosure. Whereas the precise value of the predetermined cutoff/level may vary between assays, correlations as described herein (if any) should be generally applicable.

“Pretreatment reagent,” e.g., lysis, precipitation and/or solubilization reagent, as used in a diagnostic assay as described herein is one that lyses any cells and/or solubilizes any analyte that is/are present in a test sample. Pretreatment is not necessary for all samples, as described further herein. Among other things, solubilizing the analyte (e.g., polypeptide of interest) may entail release of the analyte from any endogenous binding proteins present in the sample. A pretreatment reagent may be homogeneous (not requiring a separation step) or heterogeneous (requiring a separation step). With use of a heterogeneous pretreatment reagent there is removal of any precipitated analyte binding proteins from the test sample prior to proceeding to the next step of the assay.

“Quality control reagents” in the context of immunoassays and kits described herein, include, but are not limited to, calibrators, controls, and sensitivity panels. A “calibrator” or “standard” typically is used (e.g., one or more, such as a plurality) in order to establish calibration (standard) curves for interpolation of the concentration of an analyte, such as an antibody or an analyte. Alternatively, a single calibrator, which is near a predetermined

positive/negative cutoff, can be used. Multiple calibrators (i.e., more than one calibrator or a varying amount of calibrator(s)) can be used in conjunction so as to comprise a “sensitivity panel.”

“Risk” refers to the possibility or probability of a particular event occurring either presently or at some point in the future. “Risk stratification” refers to an array of known clinical risk factors that allows physicians to classify patients into a low, moderate, high or highest risk of developing a particular disease, disorder or condition.

“Specific” and “specificity” in the context of an interaction between members of a specific binding pair (e.g., an antigen (or fragment thereof) and an antibody (or antigenically reactive fragment thereof)) refer to the selective reactivity of the interaction. The phrase “specifically binds to” and analogous phrases refer to the ability of antibodies (or antigenically reactive fragments thereof) to bind specifically to analyte (or a fragment thereof) and not bind specifically to other entities.

“Specific binding partner” is a member of a specific binding pair. A specific binding pair comprises two different molecules, which specifically bind to each other through chemical or physical means. Therefore, in addition to antigen and antibody specific binding pairs of common immunoassays, other specific binding pairs can include biotin and avidin (or streptavidin), carbohydrates and lectins, complementary nucleotide sequences, effector and receptor molecules, cofactors and enzymes, enzyme inhibitors and enzymes, and the like. Furthermore, specific binding pairs can include members that are analogs of the original specific binding members, for example, an analyte-analog. Immunoreactive specific binding members include antigens, antigen fragments, and antibodies, including monoclonal and polyclonal antibodies as well as complexes, fragments, and variants (including fragments of variants) thereof, whether isolated or recombinantly produced.

“Variant” as used herein means a polypeptide that differs from a given polypeptide (e.g., IL-18, BNP, NGAL or HIV polypeptide or anti-polypeptide antibody) in amino acid sequence by the addition (e.g., insertion), deletion, or conservative substitution of amino acids, but that retains the biological activity of the given polypeptide (e.g., a variant IL-18 can compete with anti-IL-18 antibody for binding to IL-18). A conservative substitution of an amino acid, i.e., replacing an amino acid with a different amino acid of similar properties (e.g., hydrophilicity and degree and distribution of charged regions) is recognized in the art as typically involving a minor change. These minor changes can be identified, in part, by considering the hydrophobic index of amino acids, as understood in the art (see, e.g., Kyte et al., *J. Mol. Biol.* 157: 105-132 (1982)). The

hydropathic index of an amino acid is based on a consideration of its hydrophobicity and charge. It is known in the art that amino acids of similar hydropathic indexes can be substituted and still retain protein function. In one aspect, amino acids having hydropathic indexes of ± 2 are substituted. The hydrophilicity of amino acids also can be used to reveal substitutions that would result in proteins retaining biological function. A consideration of the hydrophilicity of amino acids in the context of a peptide permits calculation of the greatest local average hydrophilicity of that peptide, a useful measure that has been reported to correlate well with antigenicity and immunogenicity (see, e.g., U.S. Pat. No. 4,554,101). Substitution of amino acids having similar hydrophilicity values can result in peptides retaining biological activity, for example immunogenicity, as is understood in the art. In one aspect, substitutions are performed with amino acids having hydrophilicity values within ± 2 of each other. Both the hydrophobicity index and the hydrophilicity value of amino acids are influenced by the particular side chain of that amino acid. Consistent with that observation, amino acid substitutions that are compatible with biological function are understood to depend on the relative similarity of the amino acids, and particularly the side chains of those amino acids, as revealed by the hydrophobicity, hydrophilicity, charge, size, and other properties. "Variant" also can be used to describe a polypeptide or fragment thereof that has been differentially processed, such as by proteolysis, phosphorylation, or other post-translational modification, yet retains its biological activity or antigen reactivity, e.g., the ability to bind to IL-18. Use of "variant" herein is intended to encompass fragments of a variant unless otherwise contradicted by context.

I. Generation of binding proteins

Dual Variable Domain binding proteins capable of binding one or more targets and methods of making the same are provided. In an embodiment, the binding protein comprises a polypeptide chain, wherein said polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first variable domain, VD2 is a second variable domain, C is a constant domain, X1 represents an amino acid or polypeptide, X2 represents an Fc region and n is 0 or 1. The binding proteins can be generated using various techniques. Expression vectors, host cell and methods of generating the binding protein are provided.

A. Generation of parent monoclonal antibodies

The variable domains of the binding proteins provided herein can be obtained from parent antibodies, including polyclonal and mAbs capable of binding antigens of interest. These antibodies may be naturally occurring or may be generated by recombinant technology.

MABs can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, mAbs can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al. , *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced. Hybridomas are selected, cloned and further screened for desirable characteristics, including robust hybridoma growth, high antibody production and desirable antibody characteristics, as discussed in Example 1 below. Hybridomas may be cultured and expanded *in vivo* in syngeneic animals, in animals that lack an immune system, e.g., nude mice, or in cell culture *in vitro*. Methods of selecting, cloning and expanding hybridomas are well known to those of ordinary skill in the art. In a particular embodiment, the hybridomas are mouse hybridomas. In another embodiment, the hybridomas are produced in a non-human, non-mouse species such as rats, sheep, pigs, goats, cattle or horses. In another embodiment, the hybridomas are human hybridomas, in which a human non-secretory myeloma is fused with a human cell expressing an antibody capable of binding a specific antigen.

Recombinant mAbs are also generated from single, isolated lymphocytes using a procedure referred to in the art as the selected lymphocyte antibody method (SLAM), as described in U.S. Patent No. 5,627,052, PCT Publication WO 92/02551 and Babcock, J.S. *et al.* (1996) *Proc. Natl. Acad. Sci. USA* 93:7843-7848. In this method, single cells secreting antibodies of interest, e.g., lymphocytes derived from an immunized animal, are identified, and, heavy- and light-chain variable region cDNAs are rescued from the cells by reverse transcriptase-PCR and these variable regions can then be expressed, in the context of appropriate immunoglobulin constant regions (e.g., human constant regions), in mammalian host cells, such as COS or CHO cells. The host cells transfected with the amplified immunoglobulin sequences, derived from *in vivo* selected lymphocytes, can then undergo further analysis and selection *in vitro*, for example by panning the transfected cells to isolate cells expressing antibodies to the antigen of interest. The amplified immunoglobulin sequences further can be manipulated *in vitro*, such as by *in vitro* affinity maturation methods such as those described in PCT Publication WO 97/29131 and PCT Publication WO 00/56772.

Monoclonal antibodies are also produced by immunizing a non-human animal comprising some, or all, of the human immunoglobulin locus with an antigen of interest. In an

embodiment, the non-human animal is a XENOMOUSE transgenic mouse, an engineered mouse strain that comprises large fragments of the human immunoglobulin loci and is deficient in mouse antibody production. *See, e.g.*, Green et al. *Nature Genetics* 7:13-21 (1994) and United States Patents Nos. 5,916,771, 5,939,598, 5,985,615, 5,998,209, 6,075,181, 6,091,001, 5 6,114,598 and 6,130,364. See also WO 91/10741, published July 25,1991, WO 94/02602, published February 3, 1994, WO 96/34096 and WO 96/33735, both published October 31, 1996, WO 98/16654, published April 23, 1998, WO 98/24893, published June 11, 1998, WO 98/50433, published November 12, 1998, WO 99/45031, published September 10, 1999, WO 99/53049, published October 21, 1999, WO 00 09560, published February 24, 2000 and WO 10 00/037504, published June 29, 2000. The XENOMOUSE transgenic mouse produces an adult-like human repertoire of fully human antibodies, and generates antigen-specific human monoclonal antibodies. The XENOMOUSE transgenic mouse contains approximately 80% of the human antibody repertoire through introduction of megabase sized, germline configuration YAC fragments of the human heavy chain loci and x light chain loci. See Mendez et al., *Nature* 15 15:146-156 (1997), Green and Jakobovits *J. Exp. Med.* 188:483-495 (1998).

In vitro methods also can be used to make the parent antibodies, wherein an antibody library is screened to identify an antibody having the desired binding specificity. Methods for such screening of recombinant antibody libraries are well known in the art and include methods described in, for example, Ladner et al. U.S. Patent No. 5,223,409; Kang et al. PCT Publication 20 No. WO 92/18619; Dower et al. PCT Publication No. WO 91/17271; Winter et al. PCT Publication No. WO 92/20791; Markland et al. PCT Publication No. WO 92/15679; Breitling et al. PCT Publication No. WO 93/01288; McCafferty et al. PCT Publication No. WO 92/01047; Garrard et al. PCT Publication No. WO 92/09690; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum Antibod Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-25 1281; McCafferty et al., *Nature* (1990) 348:552-554; Griffiths et al. (1993) *EMBO J* 12:725-734; Hawkins et al. (1992) *J Mol Biol* 226:889-896; Clackson et al. (1991) *Nature* 352:624-628; Gram et al. (1992) *PNAS* 89:3576-3580; Garrard et al. (1991) *Bio/Technology* 9:1373-1377; Hoogenboom et al. (1991) *Nuc Acid Res* 19:4133-4137; and Barbas et al. (1991) *PNAS* 88:7978-7982, US patent application publication 20030186374, and PCT Publication No. WO 97/29131.

30 Parent antibodies can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles that carry the polynucleotide sequences encoding them. In a particular, such phage can be utilized to display antigen-binding domains expressed from a repertoire or combinatorial antibody library (e. g., human or murine). Phage expressing an antigen binding

domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods include those disclosed in Brinkman et al., *J. Immunol. Methods* 182:41-50 (1995); Ames et al., *J. Immunol. Methods* 184:177-186 (1995); Kettleborough et al., *Eur. J. Immunol.* 24:952-958 (1994); Persic et al., *Gene* 187 9-18 (1997); Burton et al., *Advances in Immunology* 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Pat. Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780, 225; 5,658,727; 5,733,743 and 5,969,108.

As described in the herein references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies including human antibodies or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., *BioTechniques* 12(6):864-869 (1992); and Sawai et al., *AJRI* 34:26-34 (1995); and Better et al., *Science* 240:1041-1043 (1988). Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Pat. 4,946,778 and 5,258, 498; Huston et al., *Methods in Enzymology* 203:46-88 (1991); Shu et al., *PNAS* 90:7995-7999 (1993); and Skerra et al., *Science* 240:1038-1040 (1988).

Alternative to screening of recombinant antibody libraries by phage display, other methodologies known in the art for screening large combinatorial libraries can be applied to the identification of parent antibodies. One type of alternative expression system is one in which the recombinant antibody library is expressed as RNA-protein fusions, as described in PCT Publication No. WO 98/31700 by Szostak and Roberts, and in Roberts, R.W. and Szostak, J.W. (1997) *Proc. Natl. Acad. Sci. USA* 94:12297-12302. In this system, a covalent fusion is created between an mRNA and the peptide or protein that it encodes by *in vitro* translation of synthetic mRNAs that carry puromycin, a peptidyl acceptor antibiotic, at their 3' end. Thus, a specific mRNA can be enriched from a complex mixture of mRNAs (e.g., a combinatorial library) based on the properties of the encoded peptide or protein, e.g., antibody, or portion thereof, such as binding of the antibody, or portion thereof, to the dual specificity antigen. Nucleic acid sequences

encoding antibodies, or portions thereof, recovered from screening of such libraries can be expressed by recombinant means as described herein (*e.g.*, in mammalian host cells) and, moreover, can be subjected to further affinity maturation by either additional rounds of screening of mRNA-peptide fusions in which mutations have been introduced into the originally selected sequence(s), or by other methods for affinity maturation *in vitro* of recombinant antibodies, as described herein.

In another approach the parent antibodies can also be generated using yeast display methods known in the art. In yeast display methods, genetic methods are used to tether antibody domains to the yeast cell wall and display them on the surface of yeast. In particular, such yeast can be utilized to display antigen-binding domains expressed from a repertoire or combinatorial antibody library (*e.g.*, human or murine). Examples of yeast display methods that can be used to make the parent antibodies include those disclosed in Wittrup, et al. U.S. Patent No. 6,699,658.

The antibodies described herein can be further modified to generate CDR grafted and humanized parent antibodies. CDR-grafted parent antibodies comprise heavy and light chain variable region sequences from a human antibody wherein one or more of the CDR regions of V_H and/or V_L are replaced with CDR sequences of murine antibodies capable of binding antigen of interest. A framework sequence from any human antibody may serve as the template for CDR grafting. However, straight chain replacement onto such a framework often leads to some loss of binding affinity to the antigen. The more homologous a human antibody is to the original murine antibody, the less likely the possibility that combining the murine CDRs with the human framework will introduce distortions in the CDRs that could reduce affinity. Therefore, in an embodiment, the human variable framework that is chosen to replace the murine variable framework apart from the CDRs have at least a 65% sequence identity with the murine antibody variable region framework. In an embodiment, the human and murine variable regions apart from the CDRs have at least 70% sequence identify. In a particular embodiment, that the human and murine variable regions apart from the CDRs have at least 75% sequence identity. In another embodiment, the human and murine variable regions apart from the CDRs have at least 80% sequence identity. Methods for producing such antibodies are known in the art (see EP 239,400; PCT publication WO 91/09967; U.S. Pat. Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, *Molecular Immunology* 28(4/5):489-498 (1991); Studnicka et al., *Protein Engineering* 7(6):805-814 (1994); Roguska et al., *PNAS* 91:969-973 (1994)), and chain shuffling (U.S. Pat. No. 5,565,352); and anti-idiotypic antibodies.

Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from

the non-human species and framework regions from a human immunoglobulin molecule. Known human Ig sequences are disclosed, e.g., [www.ncbi.nlm.nih.gov/entrez-query.fcgi](http://www.ncbi.nlm.nih.gov/entrez/query.fcgi); www.atcc.org/phage/hdb.html; www.sciquest.com/; www.abcam.com/; www.antibodyresource.com/onlinecomp.html;

5 www.public.iastate.edu/about.pedro/research_tools.html; www.mgen.uni-heidelberg.de/SD/IT/IT.html; www.whfreeman.com/immunology/CH-05/kuby05.htm; www.library.thinkquest.org/12429/Immune/Antibody.html;

www.hhmi.org/grants/lectures/1996/vlab/; www.path.cam.ac.uk/about.mrc7/m-ikeimages.html; www.antibodyresource.com/; mcb.harvard.edu/BioLinks/Immunology.html; www.immunologylink.com/; pathbox.wustl.edu/about.hcenter/index.html;

10 www.biotech.ufl.edu/about.hcl/; www.pebio.com/pa/340913/340913.html; www.nal.usda.gov/awic/pubs/antibody/; www.m.ehime-u.ac.jp/about.yasuhito/Elisa.html;

www.biodesign.com/table.asp; www.icnet.uk/axp/facs/davies/links.html; www.biotech.ufl.edu/about.fccl/protocol.html; www.isac-net.org/sites_geo.html; aximtl.imt.uni-marburg.de/about.rek/AEP-Start.html; baserv.uci.kun.nl/about.jraats/links1.html;

15 www.recab.uni-hd.de/immuno.bme.nwu.edu/; www.mrc-cpe.cam.ac.uk/imt-doc/public/INTRO.html; www.ibt.unam.mx/vir/V_mice.html; imgt.cnusc.fr:8104/;

www.biochem.ucl.ac.uk/about.martin/abs/index.html; antibody.bath.ac.uk/; abgen.cvm.tamu.edu/lab/wwwabgen.html; www.unizh.ch/about.honegger/AHOseminar/Slide01.html; www.cryst.bbk.ac.uk/about.ubcg07s/;

20 www.nimr.mrc.ac.uk/CC/caewg/caewg.htm; www.path.cam.ac.uk/about.mrc7/humanisation/TAHHP.html; www.ibt.unam.mx/vir/structure/stat_aim.html;

www.biosci.missouri.edu/smithgp/index.html; www.cryst.bioc.cam.ac.uk/about.fmolina/Webpages/Pept/spottech.html; www.jerini.de/fr/roducts.htm; www.patents.ibm.com/ibm.html. Kabat et al., Sequences of Proteins of Immunological Interest, U.S. Dept. Health (1983). Such imported sequences can be used to reduce immunogenicity or reduce, enhance or modify binding, affinity, on-rate, off-rate, avidity, specificity, half-life, or any other suitable characteristic, as known in the art.

Framework residues in the human framework regions may be substituted with the corresponding residue from the CDR donor antibody to alter, e.g., improve, antigen binding.

30 These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Pat. No. 5,585,089; Riechmann et al., Nature 332:323

35 (1988). Three-dimensional immunoglobulin models are commonly available and are familiar to

those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from the consensus and import sequences so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding. Antibodies can be humanized using a variety of techniques known in the art, such as but not limited to those described in Jones et al., *Nature* 321:522 (1986); Verhoeyen et al., *Science* 239:1534 (1988)), Sims et al., *J. Immunol.* 151: 2296 (1993); Chothia and Lesk, *J. Mol. Biol.* 196:901 (1987), Carter et al., *Proc. Natl. Acad. Sci. U.S.A.* 89:4285 (1992); Presta et al., *J. Immunol.* 151:2623 (1993), Padlan, *Molecular Immunology* 28(4/5):489-498 (1991); Studnicka et al., *Protein Engineering* 7(6):805-814 (1994); Roguska. et al. , *PNAS* 91:969-973 (1994); PCT publication WO 91/09967, PCT/: US98/16280, US96/18978, US91/09630, US91/05939, US94/01234, GB89/01334, GB91/01134, GB92/01755; WO90/14443, WO90/14424, WO90/14430, EP 229246, EP 592,106; EP 519,596, EP 239,400, U.S. Pat. Nos. 5,565,332, 5,723,323, 5,976,862, 5,824,514, 5,817,483, 5,814,476, 5,763,192, 5,723,323, 5,766,886, 5,714,352, 6,204,023, 6,180,370, 5,693,762, 5,530,101, 5,585,089, 5,225,539; 4,816,567.

20 **B. Criteria for selecting parent monoclonal antibodies**

In an embodiment, parent antibodies are selected with at least one or more properties desired in the DVD-Ig molecule. In an embodiment, the desired property is selected from one or more antibody parameters. In another embodiment, the antibody parameters are antigen specificity, affinity to antigen, potency, biological function, epitope recognition, stability, solubility, production efficiency, immunogenicity, pharmacokinetics, bioavailability, tissue cross reactivity, or orthologous antigen binding.

B1. Affinity to Antigen

The desired affinity of a therapeutic mAb may depend upon the nature of the antigen, and the desired therapeutic end-point. In an embodiment, monoclonal antibodies have higher affinities ($K_d = 0.01 - 0.50$ pM) when blocking a cytokine-cytokine receptor interaction as such interaction are usually high affinity interactions (e.g., <pM - <nM ranges). In such instances, the mAb affinity for its target should be equal to or better than the affinity of the cytokine (ligand) for its receptor. On the other hand, mAb with lesser affinity (> nM range) could be

therapeutically effective e.g., in clearing circulating potentially pathogenic proteins, e.g., monoclonal antibodies that bind to, sequester, and clear circulating species of A- β amyloid. In other instances, reducing the affinity of an existing high affinity mAb by site-directed mutagenesis or using a mAb with lower affinity for its target could be used to avoid potential side-effects e.g., a high affinity mAb may sequester/neutralize all of its intended target, thereby completely depleting/eliminating the function(s) of the targeted protein. In this scenario, a low affinity mAb may sequester/neutralize a fraction of the target that may be responsible for the disease symptoms (the pathological or over-produced levels), thus allowing a fraction of the target to continue to perform its normal physiological function(s). Therefore, it may be possible to reduce the K_d to adjust dose and/or reduce side-effects. The affinity of the parental mAb might play a role in appropriately targeting cell surface molecules to achieve desired therapeutic outcome. For example, if a target is expressed on cancer cells with high density and on normal cells with low density, a lower affinity mAb will bind a greater number of targets on tumor cells than normal cells, resulting in tumor cell elimination via ADCC or CDC, and therefore might have therapeutically desirable effects. Thus selecting a mAb with desired affinity may be relevant for both soluble and surface targets.

Signaling through a receptor upon interaction with its ligand may depend upon the affinity of the receptor-ligand interaction. Similarly, it is conceivable that the affinity of a mAb for a surface receptor could determine the nature of intracellular signaling and whether the mAb may deliver an agonist or an antagonist signal. The affinity-based nature of mAb-mediated signaling may have an impact of its side-effect profile. Therefore, the desired affinity and desired functions of therapeutic monoclonal antibodies need to be determined carefully by in vitro and in vivo experimentation.

The desired K_d of a binding protein (e.g., an antibody) may be determined experimentally depending on the desired therapeutic outcome. In an embodiment, parent antibodies with affinity (K_d) for a particular antigen equal to, or better than, the desired affinity of the DVD-Ig for the same antigen are selected. The antigen binding affinity and kinetics are assessed by Biacore or another similar technique. In one embodiment, each parent antibody has a dissociation constant (K_d) to its antigen of: at most about 10⁻⁷ M; at most about 10⁻⁸ M; at most about 10⁻⁹ M; at most about 10⁻¹⁰ M; at most about 10⁻¹¹ M; at most about 10⁻¹² M; or at most 10⁻¹³ M. First parent antibody from which VD1 is obtained and second parent antibody from which VD2 is obtained may have similar or different affinity (K_D) for the respective antigen. Each parent antibody has an on rate constant (K_{on}) to the antigen of: at least about 10²M⁻¹s⁻¹; at least about 10³M⁻¹s⁻¹; at least about 10⁴M⁻¹s⁻¹; at least about 10⁵M⁻¹s⁻¹; or at least about 10⁶M⁻¹s⁻¹, as

measured by surface plasmon resonance. The first parent antibody from which VD1 is obtained and the second parent antibody from which VD2 is obtained may have similar or different on rate constant (K_{on}) for the respective antigen. In one embodiment, each parent antibody has an off rate constant (K_{off}) to the antigen of: at most about $10^{-3}s^{-1}$; at most about $10^{-4}s^{-1}$; at most about $10^{-5}s^{-1}$; or at most about $10^{-6}s^{-1}$, as measured by surface plasmon resonance. The first parent antibody from which VD1 is obtained and the second parent antibody from which VD2 is obtained may have similar or different off rate constants (K_{off}) for the respective antigen.

B2. Potency

The desired affinity/potency of parental monoclonal antibodies will depend on the desired therapeutic outcome. For example, for receptor-ligand (R-L) interactions the affinity (k_d) is equal to or better than the R-L k_d (pM range). For simple clearance of a pathologic circulating protein, the k_d could be in low nM range e.g., clearance of various species of circulating A- β peptide. In addition, the k_d will also depend on whether the target expresses multiple copies of the same epitope e.g a mAb targeting conformational epitope in A β oligomers.

Where VDI and VD2 bind the same antigen, but distinct epitopes, the DVD-Ig will contain 4 binding sites for the same antigen, thus increasing avidity and thereby the apparent k_d of the DVD-Ig. In an embodiment, parent antibodies with equal or lower k_d than that desired in the DVD-Ig are chosen. The affinity considerations of a parental mAb may also depend upon whether the DVD-Ig contains four or more identical antigen binding sites (i.e; a DVD-Ig from a single mAb). In this case, the apparent k_d would be greater than the mAb due to avidity. Such DVD-Igs can be employed for cross-linking surface receptor, increase neutralization potency, enhance clearance of pathological proteins etc.

In an embodiment parent antibodies with neutralization potency for specific antigen equal to or better than the desired neutralization potential of the DVD-Ig for the same antigen are selected. The neutralization potency can be assessed by a target-dependent bioassay where cells of appropriate type produce a measurable signal (i.e. proliferation or cytokine production) in response to target stimulation, and target neutralization by the mAb can reduce the signal in a dose-dependent manner.

B3. Biological functions

Monoclonal antibodies can perform potentially several functions. Some of these functions are listed in Table 1. These functions can be assessed by both in vitro assays (e.g., cell-based and biochemical assays) and in vivo animal models.

Table 1: Some Potential Applications For Therapeutic Antibodies

Target (Class)	Mechanism of Action (target)
Soluble (cytokines, other)	Neutralization of activity (e.g., a cytokine) Enhance clearance (e.g., A β oligomers) Increase half-life (e.g., GLP 1)
Cell Surface (Receptors, other)	Agonist (e.g., GLP1 R; EPO R; etc.) Antagonist (e.g., integrins; etc.) Cytotoxic (CD 20; etc.)
Protein deposits	Enhance clearance/degradation (e.g., A β plaques, amyloid deposits)

MABs with distinct functions described in the examples herein in Table 1 can be selected to achieve desired therapeutic outcomes. Two or more selected parent monoclonal antibodies can then be used in DVD-Ig format to achieve two distinct functions in a single DVD-Ig molecule. For example, a DVD-Ig can be generated by selecting a parent mAb that neutralizes function of a specific cytokine, and selecting a parent mAb that enhances clearance of a pathological protein. Similarly, we can select two parent monoclonal antibodies that recognize two different cell surface receptors, one mAb with an agonist function on one receptor and the other mAb with an antagonist function on a different receptor. These two selected monoclonal antibodies each with a distinct function can be used to construct a single DVD-Ig molecule that will possess the two distinct functions (agonist and antagonist) of the selected monoclonal antibodies in a single molecule. Similarly, two antagonistic monoclonal antibodies to cell surface receptors each blocking binding of respective receptor ligands (e.g., EGF and IGF) can be used in a DVD-Ig format. Conversely, an antagonistic anti-receptor mAb (e.g., anti-EGFR) and a neutralizing anti-soluble mediator (e.g., anti-IGF1/2) mAb can be selected to make a DVD-Ig.

B4. Epitope Recognition

Different regions of proteins may perform different functions. For example specific regions of a cytokine interact with the cytokine receptor to bring about receptor activation whereas other regions of the protein may be required for stabilizing the cytokine. In this instance one may select a mAb that binds specifically to the receptor interacting region(s) on the cytokine and thereby block cytokine-receptor interaction. In some cases, for example certain chemokine receptors that bind multiple ligands, a mAb that binds to the epitope (region on chemokine receptor) that interacts with only one ligand can be selected. In other instances, monoclonal antibodies can bind to epitopes on a target that are not directly responsible for physiological functions of the protein, but binding of a mAb to these regions could either interfere with physiological functions (steric hindrance) or alter the conformation of the protein such that the

protein cannot function (mAb to receptors with multiple ligand which alter the receptor conformation such that none of the ligand can bind). Anti-cytokine monoclonal antibodies that do not block binding of the cytokine to its receptor, but block signal transduction have also been identified (e.g., 125-2H, an anti-IL-18 mAb).

5 Examples of epitopes and mAb functions include, but are not limited to, blocking Receptor-Ligand (R-L) interaction (neutralizing mAb that binds R-interacting site); steric hindrance resulting in diminished or no R-binding. An Ab can bind the target at a site other than a receptor binding site, but still interferes with receptor binding and functions of the target by inducing conformational change and eliminate function (e.g., Xolair), binding to R but block
10 signaling (125-2H).

 In an embodiment, the parental mAb needs to target the appropriate epitope for maximum efficacy. Such epitope should be conserved in the DVD-Ig. The binding epitope of a mAb can be determined by several approaches, including co-crystallography, limited proteolysis of mAb-antigen complex plus mass spectrometric peptide mapping (Legros V. et al 2000 Protein Sci.
15 9:1002-10), phage displayed peptide libraries (O'Connor KH et al 2005 J Immunol Methods. 299:21-35), as well as mutagenesis (Wu C. et al . 2003 J Immunol 170:5571-7).

B5. Physicochemical and pharmaceutical properties

 Therapeutic treatment with antibodies often requires administration of high doses, often several mg/kg (due to a low potency on a mass basis as a consequence of a typically large
20 molecular weight). In order to accommodate patient compliance and to adequately address chronic disease therapies and outpatient treatment, subcutaneous (s.c.) or intramuscular (i.m.) administration of therapeutic mAbs is desirable. For example, the maximum desirable volume for s.c. administration is ~1.0 mL, and therefore, concentrations of >100 mg/mL are desirable to limit the number of injections per dose. In an embodiment, the therapeutic antibody is administered in
25 one dose. The development of such formulations is constrained, however, by protein-protein interactions (e.g., aggregation, which potentially increases immunogenicity risks) and by limitations during processing and delivery (e.g., viscosity). Consequently, the large quantities required for clinical efficacy and the associated development constraints limit full exploitation of the potential of antibody formulation and s.c. administration in high-dose regimens. It is apparent
30 that the physicochemical and pharmaceutical properties of a protein molecule and the protein solution are of utmost importance, e.g., stability, solubility and viscosity features.

B5.1. Stability

A "stable" antibody formulation is one in which the antibody therein essentially retains its physical stability and/or chemical stability and/or biological activity upon storage. Stability can be measured at a selected temperature for a selected time period. In an embodiment, the antibody in the formulation is stable at room temperature (about 30°C) or at 40°C for at least 1 month and/or stable at about 2-8°C. for at least 1 year for at least 2 years. Furthermore, in an embodiment, the formulation is stable following freezing (to, e.g., -70°C) and thawing of the formulation, hereinafter referred to as a "freeze/thaw cycle." In another example, a "stable" formulation may be one wherein less than about 10% and less than about 5% of the protein is present as an aggregate in the formulation.

A DVD-Ig stable in vitro at various temperatures for an extended time period is desirable. One can achieve this by rapid screening of parental mAbs stable in vitro at elevated temperature, e.g., at 40°C for 2-4 weeks, and then assess stability. During storage at 2-8°C, the protein reveals stability for at least 12 months, e.g., at least 24 months. Stability (% of monomeric, intact molecule) can be assessed using various techniques such as cation exchange chromatography, size exclusion chromatography, SDS-PAGE, as well as bioactivity testing. For a more comprehensive list of analytical techniques that may be employed to analyze covalent and conformational modifications see Jones, A. J. S. (1993) Analytical methods for the assessment of protein formulations and delivery systems. In: Cleland, J. L.; Langer, R., editors. Formulation and delivery of peptides and proteins, 1st edition, Washington, ACS, pg. 22-45; and Pearlman, R.; Nguyen, T. H. (1990) Analysis of protein drugs. In: Lee, V. H., editor. Peptide and protein drug delivery, 1st edition, New York, Marcel Dekker, Inc., pg. 247-301.

Heterogeneity and aggregate formation: stability of the antibody may be such that the formulation may reveal less than about 10%, and, in an embodiment, less than about 5%, in another embodiment, less than about 2%, or, in an embodiment, within the range of 0.5% to 1.5% or less in the GMP antibody material that is present as aggregate. Size exclusion chromatography is a method that is sensitive, reproducible, and very robust in the detection of protein aggregates.

In addition to low aggregate levels, the antibody must, in an embodiment, be chemically stable. Chemical stability may be determined by ion exchange chromatography (e.g., cation or anion exchange chromatography), hydrophobic interaction chromatography, or other methods such as isoelectric focusing or capillary electrophoresis. For instance, chemical stability of the antibody may be such that after storage of at least 12 months at 2-8°C the peak representing unmodified antibody in a cation exchange chromatography may increase not more than 20%, in

an embodiment, not more than 10%, or, in another embodiment, not more than 5% as compared to the antibody solution prior to storage testing.

In an embodiment, the parent antibodies display structural integrity; correct disulfide bond formation, and correct folding: Chemical instability due to changes in secondary or tertiary structure of an antibody may impact antibody activity. For instance, stability as indicated by activity of the antibody may be such that after storage of at least 12 months at 2-8°C the activity of the antibody may decrease not more than 50%, in an embodiment not more than 30%, or even not more than 10%, or in an embodiment not more than 5% or 1% as compared to the antibody solution prior to storage testing. Suitable antigen-binding assays can be employed to determine antibody activity.

B5.2. Solubility

The "solubility" of a mAb correlates with the production of correctly folded, monomeric IgG. The solubility of the IgG may therefore be assessed by HPLC. For example, soluble (monomeric) IgG will give rise to a single peak on the HPLC chromatograph, whereas insoluble (e.g., multimeric and aggregated) will give rise to a plurality of peaks. A person skilled in the art will therefore be able to detect an increase or decrease in solubility of an IgG using routine HPLC techniques. For a more comprehensive list of analytical techniques that may be employed to analyze solubility (see Jones, A. G. *Dep. Chem. Biochem. Eng., Univ. Coll. London, London, UK*. Editor(s): Shamlou, P. Ayazi. *Process. Solid-Liq. Suspensions* (1993), 93-117. Publisher: Butterworth-Heinemann, Oxford, UK and Pearlman, Rodney; Nguyen, Tue H, *Advances in Parenteral Sciences* (1990), 4 (*Pept. Protein Drug Delivery*), 247-301). Solubility of a therapeutic mAb is critical for formulating to high concentration often required for adequate dosing. As outlined herein, solubilities of >100 mg/mL may be required to accommodate efficient antibody dosing. For instance, antibody solubility may be not less than about 5 mg/mL in early research phase, in an embodiment not less than about 25 mg/mL in advanced process science stages, or in an embodiment not less than about 100 mg/mL, or in an embodiment not less than about 150 mg/mL. It is obvious to a person skilled in the art that the intrinsic properties of a protein molecule are important the physico-chemical properties of the protein solution, e.g., stability, solubility, viscosity. However, a person skilled in the art will appreciate that a broad variety of excipients exist that may be used as additives to beneficially impact the characteristics of the final protein formulation. These excipients may include: (i) liquid solvents, cosolvents (e.g., alcohols such as ethanol); (ii) buffering agents (e.g., phosphate, acetate, citrate, amino acid buffers); (iii) sugars or sugar alcohols (e.g., sucrose, trehalose, fructose, raffinose, mannitol, sorbitol, dextran); (iv) surfactants (e.g., polysorbate 20, 40, 60, 80, poloxamers); (v) isotonicity

modifiers (e.g., salts such as NaCl, sugars, sugar alcohols); and (vi) others (e.g., preservatives, chelating agents, antioxidants, chelating substances (e.g., EDTA), biodegradable polymers, carrier molecules (e.g., HSA, PEGs)

5 Viscosity is a parameter of high importance with regard to antibody manufacture and antibody processing (e.g., diafiltration/ultrafiltration), fill-finish processes (pumping aspects, filtration aspects) and delivery aspects (syringeability, sophisticated device delivery). Low viscosities enable the liquid solution of the antibody having a higher concentration. This enables the same dose may be administered in smaller volumes. Small injection volumes inhere the advantage of lower pain on injection sensations, and the solutions not necessarily have to be
10 isotonic to reduce pain on injection in the patient. The viscosity of the antibody solution may be such that at shear rates of 100 (1/s) antibody solution viscosity is below 200 mPa s, in an embodiment below 125 mPa s, in another embodiment below 70 mPa s, and in yet another embodiment below 25 mPa s or even below 10 mPa s.

B 5.3. Production efficiency

15 The generation of a DVD-Ig that is efficiently expressed in mammalian cells, such as Chinese hamster ovary cells (CHO), will in an embodiment require two parental monoclonal antibodies which are themselves expressed efficiently in mammalian cells. The production yield from a stable mammalian line (i.e., CHO) should be above about 0.5g/L, in an embodiment above about 1g/L, and in another embodiment in the range of about 2 to about 5 g/L or more
20 (Kipriyanov SM, Little M. 1999 Mol Biotechnol. 12:173-201; Carroll S, Al-Rubeai M. 2004 Expert Opin Biol Ther. 4:1821-9).

Production of antibodies and Ig fusion proteins in mammalian cells is influenced by several factors. Engineering of the expression vector via incorporation of strong promoters, enhancers and selection markers can maximize transcription of the gene of interest from an
25 integrated vector copy. The identification of vector integration sites that are permissive for high levels of gene transcription can augment protein expression from a vector (Wurm et al, 2004, Nature Biotechnology, 2004, Vol/Iss/Pg. 22/11 (1393-1398)). Furthermore, levels of production are affected by the ratio of antibody heavy and light chains and various steps in the process of protein assembly and secretion (Jiang et al. 2006, Biotechnology Progress, Jan-Feb 2006, vol. 22,
30 no. 1, p. 313-8).

B 6. Immunogenicity

Administration of a therapeutic mAb may result in certain incidence of an immune response (i.e., the formation of endogenous antibodies directed against the therapeutic mAb). Potential elements that might induce immunogenicity should be analyzed during selection of the parental monoclonal antibodies, and steps to reduce such risk can be taken to optimize the parental monoclonal antibodies prior to DVD-Ig construction. Mouse-derived antibodies have been found to be highly immunogenic in patients. The generation of chimeric antibodies comprised of mouse variable and human constant regions presents a logical next step to reduce the immunogenicity of therapeutic antibodies (Morrison and Schlom, 1990). Alternatively, immunogenicity can be reduced by transferring murine CDR sequences into a human antibody framework (reshaping/CDR grafting/humanization), as described for a therapeutic antibody by Riechmann et al., 1988. Another method is referred to as "resurfacing" or "vencering", starting with the rodent variable light and heavy domains, only surface-accessible framework amino acids are altered to human ones, while the CDR and buried amino acids remain from the parental rodent antibody (Roguska et al., 1996). In another type of humanization, instead of grafting the entire CDRs, one technique grafts only the "specificity-determining regions" (SDRs), defined as the subset of CDR residues that are involved in binding of the antibody to its target (Kashmiri et al., 2005). This necessitates identification of the SDRs either through analysis of available three-dimensional structures of antibody-target complexes or mutational analysis of the antibody CDR residues to determine which interact with the target. Alternatively, fully human antibodies may have reduced immunogenicity compared to murine, chimeric or humanized antibodies.

Another approach to reduce the immunogenicity of therapeutic antibodies is the elimination of certain specific sequences that are predicted to be immunogenic. In one approach, after a first generation biologic has been tested in humans and found to be unacceptably immunogenic, the B-cell epitopes can be mapped and then altered to avoid immune detection. Another approach uses methods to predict and remove potential T-cell epitopes. Computational methods have been developed to scan and to identify the peptide sequences of biologic therapeutics with the potential to bind to MHC proteins (Desmet et al., 2005). Alternatively a human dendritic cell-based method can be used to identify CD4⁺ T-cell epitopes in potential protein allergens (Stickler et al., 2005; S.L. Morrison and J. Schlom, *Important Adv. Oncol.* (1990), pp. 3-18; Riechmann, L., Clark, M., Waldmann, H. and Winter, G. "Reshaping human antibodies for therapy." *Nature* (1988) 332: 323-327; Roguska-M-A, Pedersen-J-T, Henry-A-H, Searle-S-M, Roja-C-M, Avery-B, Hoffee-M, Cook-S, Lambert-J-M, Blättler-W-A, Rees-A-R, Guild-B-C. A comparison of two murine mAbs humanized by CDR-grafting and variable domain

resurfacing. Protein engineering, {Protein-Eng}, 1996, vol. 9, p. 895-904; Kashmiri-Syed-V-S, De-Pascalis-Roberto, Gonzales-Noreen-R, Schlom-Jeffrey. SDR grafting--a new approach to antibody humanization. Methods (San Diego Calif.), {Methods}, May 2005, vol. 36, no. 1, p. 25-34; Desmet-Johan, Meersseman-Geert, Boutonnet-Nathalie, Pletinckx-Jurgen, De-Clercq-
5 Krista, Debulpap-Maja, Bracckman-Tessa, Lasters-Ignace. Anchor profiles of HLA-specific peptides: analysis by a novel affinity scoring method and experimental validation. Proteins, 2005, vol. 58, p. 53-69; Stickler-M-M, Estell-D-A, Harding-F-A. CD4+ T-cell epitope determination using unexposed human donor peripheral blood mononuclear cells. Journal of immunotherapy 2000, vol. 23, p. 654-60.)

10 **B 7. In vivo efficacy**

To generate a DVD-Ig molecule with desired in vivo efficacy, it is important to generate and select mAbs with similarly desired in vivo efficacy when given in combination. However, in some instances the DVD-Ig may exhibit in vivo efficacy that cannot be achieved with the combination of two separate mAbs. For instance, a DVD-Ig may bring two targets in close
15 proximity leading to an activity that cannot be achieved with the combination of two separate mAbs. Additional desirable biological functions are described herein in section B 3. Parent antibodies with characteristics desirable in the DVD-Ig molecule may be selected based on factors such as pharmacokinetic $t_{1/2}$; tissue distribution; soluble versus cell surface targets; and target concentration- soluble/density –surface.

20 **B 8. In vivo tissue distribution**

To generate a DVD-Ig molecule with desired in vivo tissue distribution, in an embodiment parent mAbs with similar desired in vivo tissue distribution profile must be selected. Alternatively, based on the mechanism of the dual-specific targeting strategy, it may at other times not be required to select parent mAbs with the similarly desired in vivo tissue distribution
25 when given in combination. For instance, in the case of a DVD-Ig in which one binding component targets the DVD-Ig to a specific site thereby bringing the second binding component to the same target site. For example, one binding specificity of a DVD-Ig could target pancreas (islet cells) and the other specificity could bring GLP1 to the pancreas to induce insulin.

B 9. Isotype

30 To generate a DVD-Ig molecule with desired properties including, but not limited to, Isotype, Effector functions and the circulating half-life, in an embodiment parent mAbs with appropriate Fc-effector functions depending on the therapeutic utility and the desired therapeutic

end-point are selected. There are five main heavy-chain classes or isotypes some of which have several sub-types and these determine the effector functions of an antibody molecule. These effector functions reside in the hinge region, CH2 and CH3 domains of the antibody molecule. However, residues in other parts of an antibody molecule may have effects on effector functions as well. The hinge region Fc-effector functions include: (i) antibody-dependent cellular cytotoxicity, (ii) complement (C1q) binding, activation and complement-dependent cytotoxicity (CDC), (iii) phagocytosis/clearance of antigen-antibody complexes, and (iv) cytokine release in some instances. These Fc-effector functions of an antibody molecule are mediated through the interaction of the Fc-region with a set of class-specific cell surface receptors. Antibodies of the IgG1 isotype are most active while IgG2 and IgG4 having minimal or no effector functions. The effector functions of the IgG antibodies are mediated through interactions with three structurally homologous cellular Fc receptor types (and sub-types) (FcγR1, FcγR2 and FcγR3). These effector functions of an IgG1 can be eliminated by mutating specific amino acid residues in the lower hinge region (e.g., L234A, L235A) that are required for FcγR and C1q binding. Amino acid residues in the Fc region, in particular the CH2-CH3 domains, also determine the circulating half-life of the antibody molecule. This Fc function is mediated through the binding of the Fc-region to the neonatal Fc receptor (FcRn) which is responsible for recycling of antibody molecules from the acidic lysosomes back to the general circulation.

Whether a mAb should have an active or an inactive isotype will depend on the desired therapeutic end-point for an antibody. Some examples of usage of isotypes and desired therapeutic outcome are listed below:

- a) If the desired end-point is functional neutralization of a soluble cytokine then an inactive isotype may be used;
- b) If the desired out-come is clearance of a pathological protein an active isotype may be used;
- c) If the desired out-come is clearance of protein aggregates an active isotype may be used;
- d) If the desired outcome is to antagonize a surface receptor an inactive isotype is used (Tysabri, IgG4; OKT3, mutated IgG1);
- e) If the desired outcome is to eliminate target cells an active isotype is used (Herceptin, IgG1 (and with enhanced effector functions); and

- f) If the desired outcome is to clear proteins from circulation without entering the CNS an IgM isotype may be used (e.g., clearing circulating Ab peptide species).

The Fc effector functions of a parental mAb can be determined by various in vitro methods well known in the art.

- 5 As discussed, the selection of isotype, and thereby the effector functions will depend upon the desired therapeutic end-point. In cases where simple neutralization of a circulating target is desired, for example blocking receptor-ligand interactions, the effector functions may not be required. In such instances isotypes or mutations in the Fc-region of an antibody that eliminate effector functions are desirable. In other instances where elimination of target cells is
10 the therapeutic end-point, for example elimination of tumor cells, isotypes or mutations or defucosylation in the Fc-region that enhance effector functions are desirable (Presta GL, *Adv. Drug Delivery Rev.* 58:640-656, 2006; Satoh M., Iida S., Shitara K. *Expert Opinion Biol. Ther.* 6:1161-1173, 2006). Similarly, depending up on the therapeutic utility, the circulating half-life of an antibody molecule can be reduced/prolonged by modulating antibody-FcRn interactions by
15 introducing specific mutations in the Fc region (Dall'Acqua WF, Kiener PA, Wu H. *J. Biol. Chem.* 281:23514-23524, 2006; Petkova SB., Akilesh S., Sproule TJ. et al. *Internat. Immunol.* 18:1759-1769, 2006; Vaccaro C., Bawdon R., Wanjie S et al. *PNAS* 103:18709-18714, 2007).

- The published information on the various residues that influence the different effector functions of a normal therapeutic mAb may need to be confirmed for DVD-Ig. It may be possible
20 that in a DVD-Ig format additional (different) Fc-region residues, other than those identified for the modulation of monoclonal antibody effector functions, may be important.

- Overall, the decision as to which Fc-effector functions (isotype) will be critical in the final DVD-Ig format will depend up on the disease indication, therapeutic target, desired therapeutic end-point and safety considerations. Listed below are exemplary appropriate heavy
25 chain and light chain constant regions including, but not limited to:

- IgG1 – allotype: G1mz
- IgG1 mutant – A234, A235
- IgG2 – allotype: G2m(n-)
- Kappa – Km3
- 30 ○ Lambda

Fc Receptor and C1q Studies: The possibility of unwanted antibody-dependent cell-mediated cytotoxicity (ADCC) and complement-dependent cytotoxicity (CDC) by antibody complexing to any overexpressed target on cell membranes can be abrogated by the (for example, L234A, L235A) hinge-region mutations. These substituted amino acids, present in the IgG1 hinge region of mAb, are expected to result in diminished binding of mAb to human Fc receptors (but not FcRn), as FcγR binding is thought to occur within overlapping sites on the IgG1 hinge region. This feature of mAb may lead to an improved safety profile over antibodies containing a wild-type IgG. Binding of mAb to human Fc receptors can be determined by flow cytometry experiments using cell lines (e.g., THP-1, K562) and an engineered CHO cell line that expresses FcγRIIb (or other FcγRs). Compared to IgG1 control monoclonal antibodies, mAb show reduced binding to FcγRI and FcγRIIa whereas binding to FcγRIIb is unaffected. The binding and activation of C1q by antigen/IgG immune complexes triggers the classical complement cascade with consequent inflammatory and/or immunoregulatory responses. The C1q binding site on IgGs has been localized to residues within the IgG hinge region. C1q binding to increasing concentrations of mAb was assessed by C1q ELISA. The results demonstrate that mAb is unable to bind to C1q, as expected when compared to the binding of a wildtype control IgG1. Overall, the L234A, L235A hinge region mutation abolishes binding of mAb to FcγRI, FcγRIIa and C1q but does not impact the interaction of mAb with FcγRIIb. This data suggests that in vivo, mAb with mutant Fc will interact normally with the inhibitory FcγRIIb but will likely fail to interact with the activating FcγRI and FcγRIIa receptors or C1q.

Human FcRn binding: The neonatal receptor (FcRn) is responsible for transport of IgG across the placenta and to control the catabolic half-life of the IgG molecules. It might be desirable to increase the terminal half-life of an antibody to improve efficacy, to reduce the dose or frequency of administration, or to improve localization to the target. Alternatively, it might be advantageous to do the converse that is, to decrease the terminal half-life of an antibody to reduce whole body exposure or to improve the target-to-non-target binding ratios. Tailoring the interaction between IgG and its salvage receptor, FcRn, offers a way to increase or decrease the terminal half-life of IgG. Proteins in the circulation, including IgG, are taken up in the fluid phase through micropinocytosis by certain cells, such as those of the vascular endothelia. IgG can bind FcRn in endosomes under slightly acidic conditions (pH 6.0–6.5) and can recycle to the cell surface, where it is released under almost neutral conditions (pH 7.0–7.4). Mapping of the Fc-region-binding site on FcRn80, 16, 17 showed that two histidine residues that are conserved across species, His310 and His435, are responsible for the pH dependence of this interaction. Using phage-display technology, a mouse Fc-region mutation that increases binding to FcRn and extends the half-life of mouse IgG was identified (see Victor, G. et al.; Nature Biotechnology

(1997), 15(7), 637-640). Fc-region mutations that increase the binding affinity of human IgG for FcRn at pH 6.0, but not at pH 7.4, have also been identified (see Dall'Acqua William F, et al., Journal of Immunology (2002), 169(9), 5171-80). Moreover, in one case, a similar pH-dependent increase in binding (up to 27-fold) was also observed for rhesus FcRn, and this resulted in a
5 twofold increase in serum half-life in rhesus monkeys compared with the parent IgG (see Hinton, Paul R. et al., Journal of Biological Chemistry (2004), 279(8), 6213-6216). These findings indicate that it is feasible to extend the plasma half-life of antibody therapeutics by tailoring the interaction of the Fc region with FcRn. Conversely, Fc-region mutations that attenuate interaction with FcRn can reduce antibody half-life.

10 **B.10 Pharmacokinetics (PK)**

To generate a DVD-Ig molecule with desired pharmacokinetic profile, in an embodiment parent mAbs with the similarly desired pharmacokinetic profile are selected. One consideration is that immunogenic response to monoclonal antibodies (i.e., HAHA, human anti-human antibody response; HACA, human anti-chimeric antibody response) further complicates the
15 pharmacokinetics of these therapeutic agents. In an embodiment, monoclonal antibodies with minimal or no immunogenicity are used for constructing DVD-Ig molecules such that the resulting DVD-Igs will also have minimal or no immunogenicity. Some of the factors that determine the PK of a mAb include, but are not limited to, Intrinsic properties of the mAb (VH amino acid sequence); immunogenicity; FcRn binding and Fc functions.

20 The PK profile of selected parental monoclonal antibodies can be easily determined in rodents as the PK profile in rodents correlates well with (or closely predicts) the PK profile of monoclonal antibodies in cynomolgus monkey and humans. The PK profile is determined as described in Example section 1.2.2.3.A.

25 After the parental monoclonal antibodies with desired PK characteristics (and other desired functional properties as discussed herein) are selected, the DVD-Ig is constructed. As the DVD-Ig molecules contain two antigen-binding domains from two parental monoclonal antibodies, the PK properties of the DVD-Ig are assessed as well. Therefore, while determining the PK properties of the DVD-Ig, PK assays may be employed that determine the PK profile based on functionality of both antigen-binding domains derived from the 2 parent monoclonal
30 antibodies. The PK profile of a DVD-Ig can be determined as described in Example 1.2.2.3.A. Additional factors that may impact the PK profile of DVD-Ig include the antigen-binding domain (CDR) orientation; Linker size; and Fc / FcRn interactions. PK characteristics of parent

antibodies can be evaluated by assessing the following parameters: absorption, distribution, metabolism and excretion.

Absorption: To date, administration of therapeutic monoclonal antibodies is via parenteral routes (e.g., intravenous [IV], subcutaneous [SC], or intramuscular [IM]). Absorption of a mAb into the systemic circulation following either SC or IM administration from the interstitial space is primarily through the lymphatic pathway. Saturable, presystemic, proteolytic degradation may result in variable absolute bioavailability following extravascular administration. Usually, increases in absolute bioavailability with increasing doses of monoclonal antibodies may be observed due to saturated proteolytic capacity at higher doses. The absorption process for a mAb is usually quite slow as the lymph fluid drains slowly into the vascular system, and the duration of absorption may occur over hours to several days. The absolute bioavailability of monoclonal antibodies following SC administration generally ranges from 50% to 100%. In the case of a transport-mediating structure at the blood-brain barrier targeted by the DVD-Ig construct, circulation times in plasma may be reduced due to enhanced trans-cellular transport at the blood brain barrier (BBB) into the CNS compartment, where the DVD-Ig is liberated to enable interaction via its second antigen recognition site.

Distribution: Following IV administration, monoclonal antibodies usually follow a biphasic serum (or plasma) concentration-time profile, beginning with a rapid distribution phase, followed by a slow elimination phase. In general, a biexponential pharmacokinetic model best describes this kind of pharmacokinetic profile. The volume of distribution in the central compartment (V_c) for a mAb is usually equal to or slightly larger than the plasma volume (2-3 liters). A distinct biphasic pattern in serum (plasma) concentration versus time profile may not be apparent with other parenteral routes of administration, such as IM or SC, because the distribution phase of the serum (plasma) concentration-time curve is masked by the long absorption portion. Many factors, including physicochemical properties, site-specific and target-oriented receptor mediated uptake, binding capacity of tissue, and mAb dose can influence biodistribution of a mAb. Some of these factors can contribute to nonlinearity in biodistribution for a mAb.

Metabolism and Excretion: Due to the molecular size, intact monoclonal antibodies are not excreted into the urine via kidney. They are primarily inactivated by metabolism (e.g., catabolism). For IgG-based therapeutic monoclonal antibodies, half-lives typically ranges from hours or 1-2 days to over 20 days. The elimination of a mAb can be affected by many factors, including, but not limited to, affinity for the FcRn receptor, immunogenicity of the mAb, the

degree of glycosylation of the mAb, the susceptibility for the mAb to proteolysis, and receptor-mediated elimination.

B.11 Tissue cross-reactivity pattern on human and tox species

Identical staining pattern suggests that potential human toxicity can be evaluated in tox species. Tox species are those animal in which unrelated toxicity is studied.

The individual antibodies are selected to meet two criteria. (1) Tissue staining appropriate for the known expression of the antibody target. (2) Similar staining pattern between human and tox species tissues from the same organ.

Criterion 1: Immunizations and/or antibody selections typically employ recombinant or synthesized antigens (proteins, carbohydrates or other molecules). Binding to the natural counterpart and counterscreen against unrelated antigens are often part of the screening funnel for therapeutic antibodies. However, screening against a multitude of antigens is often unpractical. Therefore tissue cross-reactivity studies with human tissues from all major organs serve to rule out unwanted binding of the antibody to any unrelated antigens.

Criterion 2: Comparative tissue cross reactivity studies with human and tox species tissues (cynomolgus monkey, dog, possibly rodents and others, the same 36 or 37 tissues are being tested as in the human study) help to validate the selection of a tox species. In the typical tissue cross-reactivity studies on frozen tissues sections therapeutic antibodies may demonstrate the expected binding to the known antigen and/or to a lesser degree binding to tissues based either on low level interactions (unspecific binding, low level binding to similar antigens, low level charge based interactions, etc.). In any case the most relevant toxicology animal species is the one with the highest degree of coincidence of binding to human and animal tissue.

Tissue cross reactivity studies follow the appropriate regulatory guidelines including EC CPMP Guideline III/5271/94 "Production and quality control of mAbs" and the 1997 US FDA/CBER "Points to Consider in the Manufacture and Testing of Monoclonal Antibody Products for Human Use". Cryosections (5 μ m) of human tissues obtained at autopsy or biopsy were fixed and dried on object glass. The peroxidase staining of tissue sections was performed, using the avidin-biotin system. FDA's Guidance "*Points to Consider in the Manufacture and Testing of Monoclonal Antibody Products for Human Use*". Relevant references include Clarke J 2004, Boon L. 2002a, Boon L. 2002b, Ryan A 1999.

Tissue cross reactivity studies are often done in two stages, with the first stage including cryosections of 32 tissues (typically: Adrenal Gland, Gastrointestinal Tract, Prostate, Bladder, Heart, Skeletal Muscle, Blood Cells, Kidney, Skin, Bone Marrow, Liver, Spinal Cord, Breast, Lung, Spleen, Cerebellum, Lymph Node, Testes, Cerebral Cortex, Ovary, Thymus, Colon, Pancreas, Thyroid, Endothelium, Parathyroid, Ureter, Eye, Pituitary, Uterus, Fallopian Tube and Placenta) from one human donor. In the second phase a full cross reactivity study is performed with up to 38 tissues (including adrenal, blood, blood vessel, bone marrow, cerebellum, cerebrum, cervix, esophagus, eye, heart, kidney, large intestine, liver, lung, lymph node, breast mammary gland, ovary, oviduct, pancreas, parathyroid, peripheral nerve, pituitary, placenta, prostate, salivary gland, skin, small intestine, spinal cord, spleen, stomach, striated muscle, testis, thymus, thyroid, tonsil, ureter, urinary bladder, and uterus) from 3 unrelated adults. Studies are done typically at minimally two dose levels.

The therapeutic antibody (i.e., test article) and isotype matched control antibody may be biotinylated for avidin-biotin complex (ABC) detection; other detection methods may include tertiary antibody detection for a FITC (or otherwise) labeled test article, or precomplexing with a labeled anti-human IgG for an unlabeled test article.

Briefly, cryosections (about 5 μm) of human tissues obtained at autopsy or biopsy are fixed and dried on object glass. The peroxidase staining of tissue sections is performed, using the avidin-biotin system. First (in case of a precomplexing detection system), the test article is incubated with the secondary biotinylated anti-human IgG and developed into immune complex. The immune complex at the final concentrations of 2 and 10 $\mu\text{g}/\text{mL}$ of test article is added onto tissue sections on object glass and then the tissue sections were reacted for 30 minutes with a avidin-biotin-peroxidase kit. Subsequently, DAB (3,3'-diaminobenzidine), a substrate for the peroxidase reaction, was applied for 4 minutes for tissue staining. Antigen-Sepharose beads are used as positive control tissue sections.

Any specific staining is judged to be either an expected (e.g., consistent with antigen expression) or unexpected reactivity based upon known expression of the target antigen in question. Any staining judged specific is scored for intensity and frequency. Antigen or serum competition or blocking studies can assist further in determining whether observed staining is specific or nonspecific.

If two selected antibodies are found to meet the selection criteria – appropriate tissue staining, matching staining between human and toxicology animal specific tissue – they can be selected for DVD-Ig generation.

The tissue cross reactivity study has to be repeated with the final DVD-Ig construct, but while these studies follow the same protocol as outline herein, they are more complex to evaluate because any binding can come from any of the two parent antibodies, and any unexplained binding needs to be confirmed with complex antigen competition studies.

5 It is readily apparent that the complex undertaking of tissue crossreactivity studies with a multispecific molecule like a DVD-Ig is greatly simplified if the two parental antibodies are selected for (1) lack of unexpected tissue cross reactivity findings and (2) for appropriate similarity of tissue cross reactivity findings between the corresponding human and toxicology animal species tissues.

10 B.12 Specificity and selectivity

To generate a DVD-Ig molecule with desired specificity and selectivity, one needs to generate and select parent mAbs with the similarly desired specificity and selectivity profile.

Binding studies for specificity and selectivity with a DVD-Ig can be complex due to the four or more binding sites, two each for each antigen. Briefly, binding studies using ELISA,
15 BIAcore. KinExA or other interaction studies with a DVD-Ig need to monitor the binding of one, two or more antigens to the DVD-Ig molecule. While BIAcore technology can resolve the sequential, independent binding of multiple antigens, more traditional methods including ELISA or more modern techniques like KinExA cannot. Therefore careful characterization of each parent antibody is critical. After each individual antibody has been characterized for specificity,
20 confirmation of specificity retention of the individual binding sites in the DVD-Ig molecule is greatly simplified.

It is readily apparent that the complex undertaking of determining the specificity of a DVD-Ig is greatly simplified if the two parental antibodies are selected for specificity prior to being combined into a DVD-Ig.

25 Antigen-antibody interaction studies can take many forms, including many classical protein protein interaction studies, including ELISA (Enzyme linked immunosorbent assay), Mass spectrometry, chemical cross linking, SEC with light scattering, equilibrium dialysis, gel permeation, ultrafiltration, gel chromatography, large-zone analytical SEC, micropreparative ultracentrifugation (sedimentation equilibrium), spectroscopic methods, titration
30 microcalorimetry, sedimentation equilibrium (in analytical ultracentrifuge), sedimentation velocity (in analytical centrifuge), surface plasmon resonance (including BIAcore). Relevant references include "Current Protocols in Protein Science", John E. Coligan, Ben M. Dunn, David

W. Speicher, Paul T. Wingfield (eds.) Volume 3, chapters 19 and 20, published by John Wiley & Sons Inc., and references included therein and "Current Protocols in Immunology", John E. Coligan, Barbara E. Bierer, David H. Margulies, Ethan M. Shevach, Warren Strober (eds.) published by John Wiley & Sons Inc and relevant references included therein.

5 Cytokine Release in Whole Blood: The interaction of mAb with human blood cells can be investigated by a cytokine release assay (Wing, M. G. *Therapeutic Immunology* (1995), 2(4), 183-190; "Current Protocols in Pharmacology", S.J. Enna, Michael Williams, John W. Ferkany, Terry Kenakin, Paul Moser, (eds.) published by John Wiley & Sons Inc; Madhusudan, S. *Clinical Cancer Research* (2004), 10(19), 6528-6534; Cox, J. *Methods* (2006), 38(4), 274-282; Choi, I. European Journal of Immunology (2001), 31(1), 94-106). Briefly, various concentrations of mAb are incubated with human whole blood for 24 hours. The concentration tested should cover a wide range including final concentrations mimicking typical blood levels in patients (including but not limited to 100 ng/ml – 100µg/ml). Following the incubation, supernatants and cell lysates were analyzed for the presence of IL-1R α , TNF- α , IL-1 β , IL-6 and IL-8. Cytokine concentration profiles generated for mAb were compared to profiles produced by a negative human IgG control and a positive LPS or PHA control. The cytokine profile displayed by mAb from both cell
10 supernatants and cell lysates was comparable to control human IgG. In an embodiment, the monoclonal antibody does not interact with human blood cells to spontaneously release inflammatory cytokines.

20 Cytokine release studies for a DVD-Ig are complex due to the four or more binding sites, two each for each antigen. Briefly, cytokine release studies as described herein measure the effect of the whole DVD-Ig molecule on whole blood or other cell systems, but can resolve which portion of the molecule causes cytokine release. Once cytokine release has been detected, the purity of the DVD-Ig preparation has to be ascertained, because some co-purifying cellular
25 components can cause cytokine release on their own. If purity is not the issue, fragmentation of DVD-Ig (including but not limited to removal of Fc portion, separation of binding sites etc.), binding site mutagenesis or other methods may need to be employed to deconvolute any observations. It is readily apparent that this complex undertaking is greatly simplified if the two parental antibodies are selected for lack of cytokine release prior to being combined into a DVD-
30 Ig.

B.13 Cross reactivity to other species for toxicological studies

In an embodiment, the individual antibodies selected with sufficient cross-reactivity to appropriate tox species, for example, cynomolgus monkey. Parental antibodies need to bind to

orthologous species target (i.e. cynomolgus monkey) and elicit appropriate response (modulation, neutralization, activation). In an embodiment, the cross-reactivity (affinity/potency) to orthologous species target should be within 10-fold of the human target. In practice, the parental antibodies are evaluated for multiple species, including mouse, rat, dog, monkey (and other non-human primates), as well as disease model species (i.e. sheep for asthma model). The acceptable cross-reactivity to tox species from the parental monoclonal antibodies allows future toxicology studies of DVD-Ig-Ig in the same species. For that reason, the two parental monoclonal antibodies should have acceptable cross-reactivity for a common tox species therefore allowing toxicology studies of DVD-Ig in the same species.

10 Parent mAbs may be selected from various mAbs capable of binding specific targets and well known in the art. These include, but are not limited to anti-TNF antibody (US Patent No. 6,258,562), anti-IL-12 and/or anti-IL-12p40 antibody (US Patent No. 6,914,128); anti-IL-18 antibody (US 2005/0147610 A1), anti-C5, anti-CBL, anti-CD147, anti-gp120, anti-VLA-4, anti-CD11a, anti-CD18, anti-VEGF, anti-CD40L, anti CD-40 (e.g., see WO2007124299) anti-Id, anti-15 ICAM-1, anti-CXCL13, anti-CD2, anti-EGFR, anti-TGF-beta 2, anti-HGF, anti-cMet, anti DLL-4, anti-NPR1, anti-PLGF, anti-ErbB3, anti-E-selectin, anti-Fact VII, anti-Her2/neu, anti-F gp, anti-CD11/18, anti-CD14, anti-ICAM-3, anti-RON, anti CD-19, anti-CD80 (e.g., see WO2003039486, anti-CD4, anti-CD3, anti-CD23, anti-beta2-integrin, anti-alpha4beta7, anti-CD52, anti-HLA DR, anti-CD22 (e.g., see US Patent NO: 5,789,554), anti-CD20, anti-MIF, anti-20 CD64 (FcR), anti-TCR alpha beta, anti-CD2, anti-Hep B, anti-CA 125, anti-EpCAM, anti-gp120, anti-CMV, anti-gpIIbIIIa, anti-IgF, anti-CD25, anti-CD33, anti-HLA, anti-IGF1,2, anti IGFR, anti-VNRIIntegrin, anti-IL-1alpha, anti-IL-1beta, anti-IL-1 receptor, anti-IL-2 receptor, anti-IL-4, anti-IL-4 receptor, anti-IL5, anti-IL-5 receptor, anti-IL-6, anti- IL-6R, RANKL, NGF, DKK, alphaVbeta3, IL-17A, anti-IL-8, anti-IL-9, anti-IL-13, anti-IL-13 receptor, anti-IL-17, and anti-IL-25 23; IL-23p19; (see Presta LG. 2005 Selection, design, and engineering of therapeutic antibodies J Allergy Clin Immunol. 116:731-6 and <http://www.path.cam.ac.uk/~mrc7/humanisation/antibodies.html>).

Parent mAbs may also be selected from various therapeutic antibodies approved for use, in clinical trials, or in development for clinical use. Such therapeutic antibodies include, but are not limited to, rituximab (Rituxan®, IDEC/Genentech/Roche) (see for example U. S. Pat. No. 5,736,137), a chimeric anti-CD20 antibody approved to treat Non-Hodgkin's lymphoma; HuMax-CD20, an anti-CD20 currently being developed by Genmab, an anti-CD20 antibody described in U.S. Pat. No. 5,500,362, AME-133 (Applied Molecular Evolution), hA20 (Immunomedics, Inc.), HumaLYM (Intracel), and PRO70769 (PCT/US2003/040426, entitled

"Immunoglobulin Variants and Uses Thereof"), trastuzumab (Herceptin®, Genentech) (see for example U.S. Pat. No. 5,677,171), a humanized anti-Her2/neu antibody approved to treat breast cancer; pertuzumab (rhuMab-2C4, Omnitarg®), currently being developed by Genentech; an anti-Her2 antibody described in U.S. Pat. No. 4,753,894; cetuximab (Erbix®), Imclone (U.S. Pat. No. 4,943,533; PCT WO 96/40210), a chimeric anti-EGFR antibody in clinical trials for a variety of cancers; ABX-EGF (U.S. Pat. No. 6,235,883), currently being developed by Abgenix-Immunex-Amgen; HuMax-EGFr (U.S. Ser. No. 10/172,317), currently being developed by Genmab; 425, EMD55900, EMD62000, and EMD72000 (Merck KGaA) (U.S. Pat. No. 5,558,864; Murthy et al. 1987, Arch Biochem Biophys. 252(2):549-60; Rodeck et al., 1987, J Cell Biochem. 35(4):315-20; Kettleborough et al., 1991, Protein Eng. 4(7):773-83); ICR62 (Institute of Cancer Research) (PCT WO 95/20045; Modjtahedi et al., 1993, J. Cell Biophys. 1993, 22(1-3):129-46; Modjtahedi et al., 1993, Br J Cancer. 1993, 67(2):247-53; Modjtahedi et al., 1996, Br J Cancer, 73(2):228-35; Modjtahedi et al., 2003, Int J Cancer, 105(2):273-80); TheraCIM hR3 (YM Biosciences, Canada and Centro de Immunologia Molecular, Cuba (U.S. Pat. No. 5,891,996; U.S. Pat. No. 6,506, 883; Mateo et al, 1997, Immunotechnology, 3(1):71-81); mAb-806 (Ludwig Institute for Cancer Research, Memorial Sloan-Kettering) (Jungbluth et al. 2003, Proc Natl Acad Sci USA. 100(2):639-44); KSB-102 (KS Biomedix); MR1-1 (IVAX, National Cancer Institute) (PCT WO 0162931A2); and SC100 (Scancell) (PCT WO 01/88138); alemtuzumab (Campath®, Millenium), a humanized mAb currently approved for treatment of B-cell chronic lymphocytic leukemia; muromonab-CD3 (Orthoclone OKT3®), an anti-CD3 antibody developed by Ortho Biotech/Johnson & Johnson, ibritumomab tiuxetan (Zevalin®), an anti-CD20 antibody developed by IDEC/Schering AG, gemtuzumab ozogamicin (Mylotarg®), an anti-CD33 (p67 protein) antibody developed by Celltech/Wyeth, alefacept (Amevive®), an anti-LFA-3 Fc fusion developed by Biogen), abciximab (ReoPro®), developed by Centocor/Lilly, basiliximab (Simulect®), developed by Novartis, palivizumab (Synagis®), developed by Medimmune, infliximab (Remicade®), an anti-TNFalpha antibody developed by Centocor, adalimumab (Humira®), an anti-TNFalpha antibody developed by Abbott, Humicade®, an anti-TNFalpha antibody developed by Celltech, golimumab (CNTO-148), a fully human TNF antibody developed by Centocor, etanercept (Enbrel®), an p75 TNF receptor Fc fusion developed by Immunex/Amgen, lenercept, an p55TNF receptor Fc fusion previously developed by Roche, ABX-CBL, an anti-CD147 antibody being developed by Abgenix, ABX-IL8, an anti-IL8 antibody being developed by Abgenix, ABX-MA1, an anti-MUC18 antibody being developed by Abgenix, Pentumomab (R1549, 90Y-muHMFG1), an anti-MUC1 in development by Antisoma, Therex (R1550), an anti-MUC1 antibody being developed by Antisoma, AngioMab (AS1405), being developed by Antisoma, HuBC-1, being developed by Antisoma, Thioplatin (AS1407) being developed by Antisoma, Antegren® (natalizumab), an

anti-alpha-4-beta-1 (VLA-4) and alpha-4-beta-7 antibody being developed by Biogen, VLA-1 mAb, an anti-VLA-1 integrin antibody being developed by Biogen, LTBR mAb, an anti-lymphotoxin beta receptor (LTBR) antibody being developed by Biogen, CAT-152, an anti-TGF- β 2 antibody being developed by Cambridge Antibody Technology, ABT 874 (J695), an
5 anti-IL-12 p40 antibody being developed by Abbott, CAT-192, an anti-TGF β 1 antibody being developed by Cambridge Antibody Technology and Genzyme, CAT-213, an anti-Eotaxin1 antibody being developed by Cambridge Antibody Technology, LymphoStat-B[®] an anti-Blys antibody being developed by Cambridge Antibody Technology and Human Genome Sciences Inc., TRAIL-R1mAb, an anti-TRAIL-R1 antibody being developed by Cambridge Antibody
10 Technology and Human Genome Sciences, Inc., Avastin[®] bevacizumab, rhuMAB-VEGF, an anti-VEGF antibody being developed by Genentech, an anti-HER receptor family antibody being developed by Genentech, Anti-Tissue Factor (ATF), an anti-Tissue Factor antibody being developed by Genentech, Xolair[®] (Omalizumab), an anti-IgE antibody being developed by Genentech, Raptiva[®] (Efalizumab), an anti-CD11a antibody being developed by Genentech and
15 Xoma, MLN-02 Antibody (formerly LDP-02), being developed by Genentech and Millenium Pharmaceuticals, HuMax CD4, an anti-CD4 antibody being developed by Genmab, HuMax-IL15, an anti-IL15 antibody being developed by Genmab and Amgen, HuMax-Inflam, being developed by Genmab and Medarex, HuMax-Cancer, an anti-Heparanase I antibody being developed by Genmab and Medarex and Oxford GcoSciences, HuMax-Lymphoma, being
20 developed by Genmab and Amgen, HuMax-TAC, being developed by Genmab, IDEC-131, and anti-CD40L antibody being developed by IDEC Pharmaceuticals, IDEC-151 (Clenoliximab), an anti-CD4 antibody being developed by IDEC Pharmaceuticals, IDEC-114, an anti-CD80 antibody being developed by IDEC Pharmaceuticals, IDEC-152, an anti-CD23 being developed by IDEC Pharmaceuticals, anti-macrophage migration factor (MIF) antibodies being developed
25 by IDEC Pharmaceuticals, BEC2, an anti-idiotypic antibody being developed by Imclone, IMC-1C11, an anti-KDR antibody being developed by Imclone, DC101, an anti-flk-1 antibody being developed by Imclone, anti-VE cadherin antibodies being developed by Imclone, CEA-Cide[®] (labetuzumab), an anti-carcinoembryonic antigen (CEA) antibody being developed by Immunomedics, LymphoCide[®] (Epratuzumab), an anti-CD22 antibody being developed by
30 Immunomedics, AFP-Cide, being developed by Immunomedics, MyelomaCide, being developed by Immunomedics, LkoCide, being developed by Immunomedics, ProstaCide, being developed by Immunomedics, MDX-010, an anti-CTLA4 antibody being developed by Medarex, MDX-060, an anti-CD30 antibody being developed by Medarex, MDX-070 being developed by Medarex, MDX-018 being developed by Medarex, Osidem[®] (IDM-1), and anti-Her2 antibody
35 being developed by Medarex and Immuno-Designed Molecules, HuMax[®]-CD4, an anti-CD4 antibody being developed by Medarex and Genmab, HuMax-IL15, an anti-IL15 antibody being

developed by Medarex and Genmab, CNTO 148, an anti-TNF α antibody being developed by
 Medarex and Centocor/J&J, CNTO 1275, an anti-cytokine antibody being developed by
 Centocor/J&J, MOR101 and MOR102, anti-intercellular adhesion molecule-1 (ICAM-1)
 (CD54) antibodies being developed by MorphoSys, MOR201, an anti-fibroblast growth factor
 5 receptor 3 (FGFR-3) antibody being developed by MorphoSys, Nuvion® (visilizumab), an anti-
 CD3 antibody being developed by Protein Design Labs, HuZAF®, an anti-gamma interferon
 antibody being developed by Protein Design Labs, Anti- α 5 β 1 Integrin, being developed by
 Protein Design Labs, anti-IL-12, being developed by Protein Design Labs, ING-1, an anti-Ep-
 CAM antibody being developed by Xoma, Xolair® (Omalizumab) a humanized anti-IgE
 10 antibody developed by Genentech and Novartis, and MLN01, an anti-Beta2 integrin antibody
 being developed by Xoma. In another embodiment, the therapeutics include KRN330 (Kirin);
 huA33 antibody (A33, Ludwig Institute for Cancer Research); CNTO 95 (alpha V integrins,
 Centocor); MEDI-522 (alpha V β 3 integrin, Medimmune); volociximab (alpha V β 1 integrin,
 Biogen/PDL); Human mAb 216 (B cell glycosolated epitope, NCI); BiTE MT103 (bispecific
 15 CD19 x CD3, Medimmune); 4G7xH22 (Bispecific BcellxFcgammaR1, Medarex/Merck KGa);
 rM28 (Bispecific CD28 x MAPG, US Patent No. EP1444268); MDX447 (EMD 82633)
 (Bispecific CD64 x EGFR, Medarex); Catumaxomab (removab) (Bispecific EpCAM x anti-
 CD3, Trion/Fres); Ertumaxomab (bispecific HER2/CD3, Fresenius Biotech); oregovomab
 (OvaRex) (CA-125, ViRexx); Rencarex® (WX G250) (carbonic anhydrase IX, Wilex); CNTO
 20 888 (CCL2, Centocor); TRC105 (CD105 (endoglin), Tracon); BMS-663513 (CD137 agonist,
 Bristol Myers Squibb); MDX-1342 (CD19, Medarex); Siplizumab (MEDI-507) (CD2,
 Medimmune); Ofatumumab (Humax-CD20) (CD20, Genmab); Rituximab (Rituxan) (CD20,
 Genentech); velvuzumab (hA20) (CD20, Immunomedics); Epratuzumab (CD22, Amgen);
 lumiliximab (IDEC 152) (CD23, Biogen); muromonab-CD3 (CD3, Ortho); HuM291 (CD3 fc
 25 receptor, PDL Biopharma); HeFi-1, CD30, NCI); MDX-060 (CD30, Medarex); MDX-1401
 (CD30, Medarex); SGN-30 (CD30, Seattle Genentics); SGN-33 (Lintuzumab) (CD33, Seattle
 Genentics); Zanolimumab (HuMax-CD4) (CD4, Genmab); HCD122 (CD40, Novartis); SGN-40
 (CD40, Seattle Genentics); Campath1h (Alemtuzumab) (CD52, Genzyme); MDX-1411 (CD70,
 Medarex); hLL1 (EPB-1) (CD74.38, Immunomedics); Galiximab (IDEC-144) (CD80, Biogen);
 30 MT293 (TRC093/D93) (cleaved collagen, Tracon); HuLuc63 (CS1, PDL Pharma); ipilimumab
 (MDX-010) (CTLA4, Bristol Myers Squibb); Tremelimumab (Ticilimumab, CP-675,2)
 (CTLA4, Pfizer); HGS-ETR1 (Mapatumumab) (DR4 TRAIL-R1 agonist, Human Genome
 Science /Glaxo Smith Kline); AMG-655 (DR5, Amgen); Apomab (DR5, Genentech); CS-1008
 (DR5, Daiichi Sankyo); HGS-ETR2 (Iexatumumab) (DR5 TRAIL-R2 agonist, HGS); Cetuximab
 35 (Erbix) (EGFR, Imclone); IMC-11F8, (EGFR, Imclone); Nimotuzumab (EGFR, YM Bio);
 Panitumumab (Vectabix) (EGFR, Amgen); Zalutumumab (HuMaxEGFr) (EGFR, Genmab);

CDX-110 (EGFRvIII, AVANT Immunotherapeutics); adecatumumab (MT201) (Epcam , Merck); edrecolomab (Panorex, 17-1A) (Epcam , Glaxo/Centocor); MORAb-003 (folate receptor a, Morphotech); KW-2871 (ganglioside GD3, Kyowa); MORAb-009 (GP-9, Morphotech); CDX-1307 (MDX-1307) (hCGb, Celldex); Trastuzumab (Herceptin) (HER2, Celldex); Pertuzumab (rhuMAb 2C4) (HER2 (DI), Genentech); apolizumab (HLA-DR beta chain, PDL Pharma); AMG-479 (IGF-1R, Amgen); anti-IGF-1R R1507 (IGF1-R, Roche); CP 751871 (IGF1-R, Pfizer); IMC-A12 (IGF1-R, Imclone); BIIB022 (IGF-1R , Biogen); Mik-beta-1 (IL-2Rb (CD122), Hoffman LaRoche); CNTO 328 (IL6, Centocor); Anti-KIR (1-7F9) (Killer cell Ig-like Receptor (KIR), Novo); Hu3S193 (Lewis (y), Wyeth, Ludwig Institute of Cancer Research); hCBE-11 (LTBR, Biogen); HuHMF1 (MUC1, Antisoma/NCI); RAV12 (N-linked carbohydrate epitope, Raven); CAL (parathyroid hormone-related protein (PTH-rP), University of California); CT-011 (PD1, CureTech); MDX-1106 (ono-4538) (PD1, Medarex/Ono); MAb CT-011 (PD1, Curetech); IMC-3G3 (PDGFRa, Imclone); bavituximab (phosphatidylserine, Peregrine); huJ591 (PSMA, Cornell Research Foundation); muJ591 (PSMA, Cornell Research Foundation); GC1008 (TGFb (pan) inhibitor (IgG4), Genzyme); Infliximab (Remicade) (TNFa, Centocor); A27.15 (transferrin receptor, Salk Institute, INSERN WO 2005/111082); E2.3 (transferrin receptor, Salk Institute); Bevacizumab (Avastin) (VEGF, Genentech); HuMV833 (VEGF, Tsukuba Research Lab-WO/2000/034337, University of Texas); IMC-18F1 (VEGFR1, Imclone); IMC-1121 (VEGFR2, Imclone).

20 C. Construction of binding protein molecules

The dual variable domain immunoglobulin (DVD-Ig) molecule is designed such that two different light chain variable domains (VL) from the two different parent monoclonal antibodies are linked in tandem directly or via a short linker by recombinant DNA techniques, followed by the light chain constant domain. Similarly, the heavy chain comprises two different heavy chain variable domains (VH) linked in tandem, followed by the constant domain CH1 and Fc region (Figure 1A).

The variable domains can be obtained using recombinant DNA techniques from a parent antibody generated by any one of the methods described herein. In an embodiment, the variable domain is a murine heavy or light chain variable domain. In another embodiment, the variable domain is a CDR grafted or a humanized variable heavy or light chain domain. In an embodiment, the variable domain is a human heavy or light chain variable domain.

In one embodiment the first and second variable domains are linked directly to each other using recombinant DNA techniques. In another embodiment the variable domains are linked via a

Linker sequences may also be derived from other proteins such as Ig-like proteins, (e.g. TCR, FcR, KIR); G/S based sequences (e.g., G4S (SEQ ID NO: 27) or G4S repeats); hinge region-derived sequences; and other natural sequences from other proteins.

In an embodiment a constant domain is linked to the two linked variable domains using recombinant DNA techniques. In an embodiment, sequence comprising linked heavy chain variable domains is linked to a heavy chain constant domain and sequence comprising linked light chain variable domains is linked to a light chain constant domain. In an embodiment, the constant domains are human heavy chain constant domain and human light chain constant domain respectively. In an embodiment, the heavy chain is further linked to an Fc region. The Fc region may be a native sequence Fc region, or a variant Fc region. In another embodiment, the Fc region is a human Fc region. In another embodiment the Fc region includes Fc region from IgG1, IgG2, IgG3, IgG4, IgA, IgM, IgE, or IgD.

In another embodiment two heavy chain polypeptides and two light chain polypeptides are combined to form a DVD-Ig molecule. Table 2 lists amino acid sequences of VH and VL regions of exemplary antibodies for targets useful for treating disease, e.g., for treating cancer. In an embodiment, the DVD Ig comprises at least two of the VH and/or VL regions listed in Table 2, in any orientation. In some embodiments, VD1 and VD2 are independently chosen. Therefore, in some embodiments, VD1 and VD2 comprise the same SEQ ID NO and, in other embodiments, VD1 and VD2 comprise different SEQ ID NOS.

The VH and VL domain sequences provided below comprise complementary determining region (CDR) and framework sequences that are either known in the art or readily discernable using methods known in the art. In some embodiments, one or more of these CDR and/or framework sequences are replaced, without loss of function, by other CDR and/or framework sequences from binding proteins that are known in the art to bind to the same antigen.

Table 2: List of Amino Acid Sequences of VH and VL regions of Antibodies for Generating DVD-Igs

SEQ ID No.	ABT Unique ID	Protein Region	Sequence
			123456789012345678901234567890123456
28	AB017VH	VH TNF (seq. 1)	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPFKGLEWVSAITWNSGHIDYADSVGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGLVTVSS
29	AB017VL	VL TNF (seq. 1)	DIQMTQSPFSSLSASVGDRTTTCRASQGIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFSGSGSGTDFTLTISLQPEDVATYYCQRYNRAPYTFGQGTKVEIKR

SEQ ID No.	ABT Unique ID	Protein Region	Sequence
			123456789012345678901234567890123456
30	AB020VH	VH NGF	QVQLQESGGGLVVKPSETLSLTCTVSGFSLIGYDLNWRQP PGKGLEWIGIHWGDCGTTDYN SAVKSRVTISKDTSKNQFSL KLSSVTAADTAVYYCARGGYWYATSYFYFDYWGQGLVTVS S
31	AB020VL	VL NGF	DIQMTQSPSSLSASVGDRTVITCRASQSIENLNWYQCKP GKAPKLLIYYTSRFSHSGVPSRFSGSGSGTDFTTISSLOP EDIATYYCQOEHTLPYTFGGGTKLEIKR
32	AB022VH	VH SOST	EVQLQQSGPELVTPGASVKISCKASGYTFTDHYMSWVKQS HGKSLIEWIGDINPYSGETTYNQKFKGATLTVDKSSSIAY MEIRGLTSEDSAVYYCARDYDASPFAYWGQGLVTVSA
33	AB022VL	VL SOST	DVQMIQSPSSLSASLGDIVTMTCPQASQGTINLNWFQCKP GKAPKLLIYGSNLEDGVPSPRFSGSRVGTDFTLTISSELD EDLATYFCLQHSYLPYTFGGGTKLEIKR
34	AB048VH	VH PGE2	EVQLVQSGAEVKKPGASVKVCKASGYTFTKYWLGWVRQA PGQGLEWMDIYPGYDYTHYNEKFKDRVTLTDTSTSTAY MELRSLRSDDTAVYYCARSDGSSTYWGQGLVTVSS
35	AB048VL	VL PGE2	DVLTQTPLSLPVTPEPASI SCTSSQNI VHSNGNTYLEW YLQKPGQSPQLLIYKVSNRFRSGVDPDRFSGSGSGTDFTLKI SRVEAEDLVGVYCFQVSHVYPYTFGGGTKVEIKR
36	AB213VH	VH TNF (seq. 2)	QVQLKESGGGLVAPSQSLTCTVSGFSLTDYGVNWRQP PGKGLEWLGMIWDCGSTDYDSTLKSRLSISKDNSKSKQIFL KMNSLQTDATARYYCAREWHHGEVAYWGQGLVTVSA
37	AB213VL	VL TNF (seq. 2)	DIVMTQSHKFMSTTVGDRVSTTCASQAVSSAVAWYQCKP GQSPKLLIYWASTRHTGVPDRFTGSGSVTDFTLTHNIQA EDLALYYCQHYSTPFTFGSGTKLEIKR
38	AB214VH	VH TNF (seq. 3)	EVKLEESGGGLVQPGGSMKLSVAVSGFI FSNHMMNWRQS PEKGLEWVAEIRSKSINSATHYAESVKGRFTISRDDSKSA VYLQMI DLRTEDTGVYYCSRNYGSTYDYWGQGLTTLVSS
39	AB214VL	VL TNF (seq. 3)	DILLTQSPAILSVSPGERVSEFCRASQFVGGSSIHVYQCRT NGSPRLLIKYASESMSGIPSRFSGSGSGTDFTLTINTVES EDIADYYCQESHVWPFPTFGSGTNLEVKR
40	AB215VH	VH TNF (seq. 4)	QVQLVESGGGVVQPGSRSLRSCAASGFTFSSYAMHWVRQA PGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLY LQMNSLRAEDTAVYYCARDRGLAAGNYYYYGMDVWGQGT TVTVSS
41	AB215VL	VL TNF (seq. 4)	EIVLTQSPATLSLSPGERATLSCRASQSVSYLAWYQCKP GQAPRLLIYDASNRAATGIPARFSGSGSGTDFTLTISSELP EDFAVYYCQQRSNWPPFTFGGPKVDIKR
42	AB216VH	VH LPA	QVQLQSGAEIVRPGTSVKVCKASGYGFINYLIEWIKQR PGQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAY MHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGLTTLV SS
43	AB216VL	VL LPA	DVVMTQTPSLPVS LGDQASISCTSGQSLVHNGNTYLHW YLQKPGQSPKLLIYKVSNFLSGVDPDRFSGSGSGTDFTLKI SRVEAEDLVGVYFCSQSTHFPFTFGGTKLEIKR
44	AB217VH	VH TNF (seq. 5)	QVQLVESGGGVVQPGSRSLRSCAASGFTFSSYDMHWVRQA PGKGLEWVAVIWSGSIKYYADSVKGRFTISRDNKNTLY LQMNSLRAEDTAVYYCAREVESAMGGFYNGMDVWGQGT TVTVSS
45	AB217VL	VL TNF (seq. 5)	DIQMTQSPSSLSASVGDRTVITCRASQGI RIDLGWYQCKP GKAPKRLIYAATLQSGVPSRFSGSGSGTEFTTISLQ EDFAVYYCQKHSYPLTFGGGTKVEIKR
46	AB218VH	VH TNF (seq. 6)	EVQLVESGGGLIQPGGSLRSCAASGFTVSRNYMSWVRQA PGKGLEWVSVIYSGDRYYADSVKGRFTISRDNKNTLYL QMNSLRAEDTAVYYCARGEGGFYWGQGLVTVSS
47	AB218VL	VL TNF (seq. 6)	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQCKP GQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISSLQ EDFAVYYCQYNYWWTFGGPKVEIKR

Detailed description of specific DVD-Ig molecules capable of binding specific targets, and methods of making the same, is provided in the Examples section below.

D. Production of binding proteins

Binding proteins provided herein may be produced by any of a number of techniques
5 known in the art. For example, expression from host cells, wherein expression vector(s) encoding the heavy and light chains is (are) transfected into a host cell by standard techniques. The various forms of the term "transfection" are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, *e.g.*, electroporation, calcium-phosphate precipitation, DEAE-dextran transfection and the like.
10 Although it is possible to express the binding proteins provided herein in either prokaryotic or eukaryotic host cells, binding proteins are expressed in eukaryotic cells, for example, mammalian host cells, because such eukaryotic cells (and in particular mammalian cells) are more likely than prokaryotic cells to assemble and secrete a properly folded and immunologically active binding protein.

15 Exemplary mammalian host cells for expressing the recombinant binding proteins include Chinese Hamster Ovary (CHO cells) (including dhfr- CHO cells, described in Urlaub and Chasin, (1980) *Proc. Natl. Acad. Sci. USA* 77:4216-4220, used with a DHFR selectable marker, *e.g.*, as described in R.J. Kaufman and P.A. Sharp (1982) *Mol. Biol.* 159:601-621), NS0 myeloma cells, COS cells, SP2 and PER.C6 cells. When recombinant expression vectors encoding binding
20 proteins are introduced into mammalian host cells, the binding proteins are produced by culturing the host cells for a period of time sufficient to allow for expression of the binding proteins in the host cells or secretion of the binding proteins into the culture medium in which the host cells are grown. Binding proteins can be recovered from the culture medium using standard protein purification methods.

25 In an exemplary system for recombinant expression, a recombinant expression vector encoding both the heavy chain and the light chain is introduced into dhfr- CHO cells by calcium phosphate-mediated transfection. Within the recombinant expression vector, the heavy and light chain genes are each operatively linked to CMV enhancer/AdMLP promoter regulatory elements to drive high levels of transcription of the genes. The recombinant expression vector also carries
30 a DHFR gene, which allows for selection of CHO cells that have been transfected with the vector using methotrexate selection/amplification. The selected transformant host cells are cultured to allow for expression of the heavy and light chains and intact binding protein is recovered from the culture medium. Standard molecular biology techniques are used to prepare the recombinant

expression vector, transfect the host cells, select for transformants, culture the host cells and recover the binding protein from the culture medium. Still further a method of synthesizing a binding protein by culturing a host cell in a suitable culture medium until a binding protein is synthesized is provided. The method can further comprise isolating the binding protein from the
5 culture medium.

An important feature of DVD-Ig is that it can be produced and purified in a similar way as a conventional antibody. The production of DVD-Ig results in a homogeneous, single major product with desired dual-specific activity, without any sequence modification of the constant region or chemical modifications of any kind. Other previously described methods to generate
10 "bi-specific", "multi-specific", and "multi-specific multivalent" full length binding proteins do not lead to a single primary product but instead lead to the intracellular or secreted production of a mixture of assembled inactive, mono-specific, multi-specific, multivalent, full length binding proteins, and multivalent full length binding proteins with combination of different binding sites. As an example, based on the design described by Miller and Presta (PCT publication
15 WO2001/077342(A1), there are 16 possible combinations of heavy and light chains. Consequently only 6.25% of protein is likely to be in the desired active form, and not as a single major product or single primary product compared to the other 15 possible combinations. Separation of the desired, fully active forms of the protein from inactive and partially active forms of the protein using standard chromatography techniques, typically used in large scale
20 manufacturing, is yet to be demonstrated.

Surprisingly the design of the "dual-specific multivalent full length binding proteins" leads to a dual variable domain light chain and a dual variable domain heavy chain which assemble primarily to the desired "dual-specific multivalent full length binding proteins".

At least 50%, at least 75% and at least 90% of the assembled, and expressed dual
25 variable domain immunoglobulin molecules are the desired dual-specific tetravalent protein. This embodiment particularly enhances commercial utility. Therefore, a method to express a dual variable domain light chain and a dual variable domain heavy chain in a single cell leading to a single primary product of a "dual-specific tetravalent full length binding protein" is provided.

Methods of expressing a dual variable domain light chain and a dual variable domain
30 heavy chain in a single cell leading to a "primary product" of a "dual-specific tetravalent full length binding protein", where the "primary product" is more than 50% of all assembled protein, comprising a dual variable domain light chain and a dual variable domain heavy chain are provided.

Methods of expressing a dual variable domain light chain and a dual variable domain heavy chain in a single cell leading to a single "primary product" of a "dual-specific tetravalent full length binding protein", where the "primary product" is more than 75% of all assembled protein, comprising a dual variable domain light chain and a dual variable domain heavy chain are provided.

Methods of expressing a dual variable domain light chain and a dual variable domain heavy chain in a single cell leading to a single "primary product" of a "dual-specific tetravalent full length binding protein", where the "primary product" is more than 90% of all assembled protein, comprising a dual variable domain light chain and a dual variable domain heavy chain are provided.

II. Derivatized binding proteins

One embodiment provides a labeled binding protein wherein the binding protein is derivatized or linked to another functional molecule (e.g., another peptide or protein). For example, a labeled binding protein can be derived by functionally linking a binding protein provided herein (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody (e.g., a bispecific antibody or a diabody), a detectable agent, a cytotoxic agent, a pharmaceutical agent, and/or a protein or peptide that can mediate association of the binding protein with another molecule (such as a streptavidin core region or a polyhistidine tag).

Useful detectable agents with which a binding protein provided herein may be derivatized include fluorescent compounds. Exemplary fluorescent detectable agents include fluorescein, fluorescein isothiocyanate, rhodamine, 5-dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin and the like. A binding protein may also be derivatized with detectable enzymes, such as alkaline phosphatase, horseradish peroxidase, glucose oxidase and the like. When a binding protein is derivatized with a detectable enzyme, it is detected by adding additional reagents that the enzyme uses to produce a detectable reaction product. For example, when the detectable agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. a binding protein may also be derivatized with biotin, and detected through indirect measurement of avidin or streptavidin binding.

Another embodiment provides a crystallized binding protein and formulations and compositions comprising such crystals. In one embodiment the crystallized binding protein has a

greater half-life *in vivo* than the soluble counterpart of the binding protein. In another embodiment the binding protein retains biological activity after crystallization.

Crystallized binding proteins may be produced according to methods known in the art and as disclosed in WO 02072636.

5 Another embodiment provides a glycosylated binding protein wherein the binding protein or antigen-binding portion thereof comprises one or more carbohydrate residues. Nascent *in vivo* protein production may undergo further processing, known as post-translational modification. In particular, sugar (glycosyl) residues may be added enzymatically, a process known as glycosylation. The resulting proteins bearing covalently linked oligosaccharide side chains are
10 known as glycosylated proteins or glycoproteins. Antibodies are glycoproteins with one or more carbohydrate residues in the Fc domain, as well as the variable domain. Carbohydrate residues in the Fc domain have important effect on the effector function of the Fc domain, with minimal effect on antigen binding or half-life of the antibody (R. Jefferis, *Biotechnol. Prog.* **21** (2005), pp. 11–16). In contrast, glycosylation of the variable domain may have an effect on the antigen
15 binding activity of the antibody. Glycosylation in the variable domain may have a negative effect on antibody binding affinity, likely due to steric hindrance (Co, M.S., et al., *Mol. Immunol.* (1993) 30:1361- 1367), or result in increased affinity for the antigen (Wallick, S.C., et al., *Exp. Med.* (1988) 168:1099-1109; Wright, A., et al., *EMBO J.* (1991) 10:2717 2723).

One aspect is directed to generating glycosylation site mutants in which the O- or N-
20 linked glycosylation site of the binding protein has been mutated. One skilled in the art can generate such mutants using standard well-known technologies. Glycosylation site mutants that retain the biological activity but have increased or decreased binding activity are other embodiments.

In still another embodiment, the glycosylation of the binding protein or antigen-binding
25 portion thereof is modified. For example, an aglycosylated binding protein can be made (i.e., the binding protein lacks glycosylation). Glycosylation can be altered to, for example, increase the affinity of the binding protein for antigen. Such carbohydrate modifications can be accomplished by, for example, altering one or more sites of glycosylation within the binding protein sequence. For example, one or more amino acid substitutions can be made that result in elimination of one
30 or more variable region glycosylation sites to thereby eliminate glycosylation at that site. Such aglycosylation may increase the affinity of the binding protein for antigen. Such an approach is described in further detail in PCT Publication WO2003016466A2, and U.S. Pat. Nos. 5,714,350 and 6,350,861.

Additionally or alternatively, a modified binding protein can be made that has an altered type of glycosylation, such as a hypofucosylated binding protein having reduced amounts of fucosyl residues (see Kanda, Yutaka et al., *Journal of Biotechnology* (2007), 130(3), 300-310) or binding protein having increased bisecting GlcNAc structures. Such altered glycosylation patterns have been demonstrated to increase the ADCC ability of antibodies. Such carbohydrate modifications can be accomplished by, for example, expressing the binding proteins in a host cell with altered glycosylation machinery. Cells with altered glycosylation machinery have been described in the art and can be used as host cells in which to express the recombinant binding proteins to thereby produce binding proteins with altered glycosylation. See, for example, Shields, R. L. et al. (2002) *J. Biol. Chem.* 277:26733-26740; Umana et al. (1999) *Nat. Biotech.* 17:176-1, as well as, European Patent No: EP 1,176,195; PCT Publications WO 03/035835; WO 99/54342 80.

Protein glycosylation depends on the amino acid sequence of the protein of interest, as well as the host cell in which the protein is expressed. Different organisms may produce different glycosylation enzymes (eg., glycosyltransferases and glycosidases), and have different substrates (nucleotide sugars) available. Due to such factors, protein glycosylation pattern, and composition of glycosyl residues, may differ depending on the host system in which the particular protein is expressed. Glycosyl residues may include, but are not limited to, glucose, galactose, mannose, fucose, n-acetylglucosamine and sialic acid. In an embodiment, the glycosylated binding protein comprises glycosyl residues such that the glycosylation pattern is human.

It is known to those skilled in the art that differing protein glycosylation may result in differing protein characteristics. For instance, the efficacy of a therapeutic protein produced in a microorganism host, such as yeast, and glycosylated utilizing the yeast endogenous pathway may be reduced compared to that of the same protein expressed in a mammalian cell, such as a CHO cell line. Such glycoproteins may also be immunogenic in humans and show reduced half-life *in vivo* after administration. Specific receptors in humans and other animals may recognize specific glycosyl residues and promote the rapid clearance of the protein from the bloodstream. Other adverse effects may include changes in protein folding, solubility, susceptibility to proteases, trafficking, transport, compartmentalization, secretion, recognition by other proteins or factors, antigenicity, or allergenicity. Accordingly, a practitioner may choose a therapeutic protein with a specific composition and pattern of glycosylation, for example glycosylation composition and pattern identical, or at least similar, to that produced in human cells or in the species-specific cells of the intended subject animal.

Expressing glycosylated proteins different from that of a host cell may be achieved by genetically modifying the host cell to express heterologous glycosylation enzymes. Using techniques known in the art a practitioner may generate binding proteins or antigen-binding portions thereof exhibiting human protein glycosylation. For example, yeast strains have been genetically modified to express non-naturally occurring glycosylation enzymes such that glycosylated proteins (glycoproteins) produced in these yeast strains exhibit protein glycosylation identical to that of animal cells, especially human cells (U.S patent applications 20040018590 and 20020137134 and PCT publication WO2005100584 A2).

In addition to the binding proteins, certain embodiments are also directed to anti-idiotypic (anti-Id) antibodies specific for such binding proteins. An anti-Id antibody is an antibody, which recognizes unique determinants generally associated with the antigen-binding region of another antibody. The anti-Id can be prepared by immunizing an animal with the binding protein or a CDR containing region thereof. The immunized animal will recognize, and respond to the idiotype determinants of the immunizing binding protein and produce an anti-Id antibody. It is readily apparent that it may be easier to generate anti-idiotypic antibodies to the two or more parent antibodies incorporated into a DVD-Ig molecule; and confirm binding studies by methods well recognized in the art (e.g., BIAcore, ELISA) to verify that anti-idiotypic antibodies specific for the idiotype of each parent antibody also recognize the idiotype (e.g., antigen binding site) in the context of the DVD-Ig. The anti-idiotypic antibodies specific for each of the two or more antigen binding sites of a DVD-Ig provide ideal reagents to measure DVD-Ig concentrations of a human DVD-Ig in patient serum; DVD-Ig concentration assays can be established using a "sandwich assay ELISA format" with an antibody to a first antigen binding regions coated on the solid phase (e.g., BIAcore chip, ELISA plate etc.), rinsed with rinsing buffer, incubation with the serum sample, another rinsing step and ultimately incubation with another anti-idiotypic antibody to the another antigen binding site, itself labeled with an enzyme for quantitation of the binding reaction. In an embodiment, for a DVD-Ig with more than two different binding sites, anti-idiotypic antibodies to the two outermost binding sites (most distal and proximal from the constant region) will not only help in determining the DVD-Ig concentration in human serum but also document the integrity of the molecule *in vivo*. Each anti-Id antibody may also be used as an "immunogen" to induce an immune response in yet another animal, producing a so-called anti-anti-Id antibody.

Further, it will be appreciated by one skilled in the art that a protein of interest may be expressed using a library of host cells genetically engineered to express various glycosylation enzymes, such that member host cells of the library produce the protein of interest with variant

glycosylation patterns. A practitioner may then select and isolate the protein of interest with particular novel glycosylation patterns. In an embodiment, the protein having a particularly selected novel glycosylation pattern exhibits improved or altered biological properties.

III. Uses of DVD-Ig

- 5 Given their ability to bind to two or more antigens the binding proteins provided herein can be used to detect the antigens (*e.g.*, in a biological sample, such as serum or plasma), using a conventional immunoassay, such as an enzyme linked immunosorbent assays (ELISA), an radioimmunoassay (RIA) or tissue immunohistochemistry. The DVD-Ig is directly or indirectly labeled with a detectable substance to facilitate detection of the bound or unbound DVD-Ig.
- 10 Suitable detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate,
- 15 rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; and examples of suitable radioactive material include ^3H , ^{14}C , ^{35}S , ^{90}Y , $^{99\text{m}}\text{Tc}$, ^{111}In , ^{125}I , ^{131}I , ^{177}Lu , ^{166}Ho , or ^{153}Sm .

- In an embodiment, the binding proteins are capable of neutralizing the activity of the antigens both *in vitro* and *in vivo*. Accordingly, such DVD-Igs can be used to inhibit antigen
- 20 activity, *e.g.*, in a cell culture containing the antigens, in human subjects or in other mammalian subjects having the antigens with which a binding protein cross-reacts. In another embodiment, a method for reducing antigen activity in a subject suffering from a disease or disorder in which the antigen activity is detrimental is provided. In some embodiments, the binding proteins can be administered to a human subject for therapeutic purposes.

- 25 As used herein, the term "a disorder in which antigen activity is detrimental" is intended to include diseases and other disorders in which the presence of the antigen in a subject suffering from the disorder has been shown to be or is suspected of being either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Accordingly, a disorder in which antigen activity is detrimental is a disorder in which reduction
- 30 of antigen activity is expected to alleviate the symptoms and/or progression of the disorder. Such disorders may be evidenced, for example, by an increase in the concentration of the antigen in a biological fluid of a subject suffering from the disorder (*e.g.*, an increase in the concentration of antigen in serum, plasma, synovial fluid, *etc.* of the subject). Non-limiting examples of disorders

that can be treated with the binding proteins provided herein include those disorders discussed below and in the section pertaining to pharmaceutical compositions.

The DVD-Igs provided herein may bind one antigen or multiple antigens. Such antigens include, but are not limited to, the targets listed in the following databases, which databases are incorporated herein by reference. These target databases include those listings:

Therapeutic targets (<http://xin.cz3.nus.edu.sg/group/cjttd/ttd.asp>);

Cytokines and cytokine receptors (<http://www.cytokinewebfacts.com/>,
<http://www.copewithcytokines.de/cope.cgi>, and

[http://cmbi.bjmu.edu.cn/cmbidata/cgi/CGF_Database/cytokine.medic.kumamoto-](http://cmbi.bjmu.edu.cn/cmbidata/cgi/CGF_Database/cytokine.medic.kumamoto-u.ac.jp/CFC/indexR.html)
10 [u.ac.jp/CFC/indexR.html](http://cmbi.bjmu.edu.cn/cmbidata/cgi/CGF_Database/cytokine.medic.kumamoto-u.ac.jp/CFC/indexR.html));

Chemokines (<http://cytokine.medic.kumamoto-u.ac.jp/CFC/CK/Chemokine.html>);

Chemokine receptors and GPCRs (<http://csp.medic.kumamoto-u.ac.jp/CSP/Receptor.html>,
<http://www.gpcr.org/7tm/>);

Olfactory Receptors (<http://senselab.med.yale.edu/senselab/ORDB/default.asp>);

15 Receptors (<http://www.iuphar-db.org/iuphar-rd/list/index.htm>);

Cancer targets (<http://cged.hgc.jp/cgi-bin/input.cgi>);

Secreted proteins as potential antibody targets (<http://spd.cbi.pku.edu.cn/>);

Protein kinases (<http://spd.cbi.pku.edu.cn/>), and

Human CD markers (http://content.labvelocity.com/tools/6/1226/CD_table_final_locked.pdf) and
20 (Zola H, 2005 CD molecules 2005: human cell differentiation molecules Blood, 106:3123-6).

DVD-Igs are useful as therapeutic agents to simultaneously block two different targets to enhance efficacy/safety and/or increase patient coverage. Such targets may include soluble targets (TNF) and cell surface receptor targets (VEGFR and EGFR). It can also be used to induce redirected cytotoxicity between tumor cells and T cells (Her2 and CD3) for cancer therapy, or
25 between autoreactive cell and effector cells for autoimmune disease or transplantation, or between any target cell and effector cell to eliminate disease-causing cells in any given disease.

In addition, DVD-Ig can be used to trigger receptor clustering and activation when it is designed to target two different epitopes on the same receptor. This may have benefit in making agonistic and antagonistic anti-GPCR therapeutics. In this case, DVD-Ig can be used to target two different epitopes (including epitopes on both the loop regions and the extracellular domain) on one cell for clustering/signaling (two cell surface molecules) or signaling (on one molecule). Similarly, a DVD-Ig molecule can be designed to trigger CTLA-4 ligation, and a negative signal by targeting two different epitopes (or 2 copies of the same epitope) of CTLA-4 extracellular domain, leading to down regulation of the immune response. CTLA-4 is a clinically validated target for therapeutic treatment of a number of immunological disorders. CTLA-4/B7 interactions negatively regulate T cell activation by attenuating cell cycle progression, IL-2 production, and proliferation of T cells following activation, and CTLA-4 (CD152) engagement can down-regulate T cell activation and promote the induction of immune tolerance. However, the strategy of attenuating T cell activation by agonistic antibody engagement of CTLA-4 has been unsuccessful since CTLA-4 activation requires ligation. The molecular interaction of CTLA-4/B7 is in "skewed zipper" arrays, as demonstrated by crystal structural analysis (Stamper 2001 Nature 410:608). However none of the currently available CTLA-4 binding reagents have ligation properties, including anti-CTLA-4 mAbs. There have been several attempts to address this issue. In one case, a cell member-bound single chain antibody was generated, and significantly inhibited allogeneic rejection in mice (Hwang 2002 JI 169:633). In a separate case, artificial APC surface-linked single-chain antibody to CTLA-4 was generated and demonstrated to attenuate T cell responses (Griffin 2000 JI 164:4433). In both cases, CTLA-4 ligation was achieved by closely localized member-bound antibodies in artificial systems. While these experiments provide proof-of-concept for immune down-regulation by triggering CTLA-4 negative signaling, the reagents used in these reports are not suitable for therapeutic use. To this end, CTLA-4 ligation may be achieved by using a DVD-Ig molecule, which target two different epitopes (or 2 copies of the same epitope) of CTLA-4 extracellular domain. The rationale is that the distance spanning two binding sites of an IgG, approximately 150-170Å, is too large for active ligation of CTLA-4 (30-50 Å between 2 CTLA-4 homodimer). However the distance between the two binding sites on DVD-Ig (one arm) is much shorter, also in the range of 30-50 Å, allowing proper ligation of CTLA-4.

Similarly, DVD-Ig can target two different members of a cell surface receptor complex (e.g., IL-12R alpha and beta). Furthermore, DVD-Ig can target CR1 and a soluble protein/pathogen to drive rapid clearance of the target soluble protein/pathogen.

Additionally, DVD-Igs provided herein can be employed for tissue-specific delivery (target a tissue marker and a disease mediator for enhanced local PK thus higher efficacy and/or lower toxicity), including intracellular delivery (targeting an internalizing receptor and a intracellular molecule), delivering to inside brain (targeting transferrin receptor and a disease mediator for crossing the blood-brain barrier). DVD-Ig can also serve as a carrier protein to deliver an antigen to a specific location via binding to a non-neutralizing epitope of that antigen and also to increase the half-life of the antigen. Furthermore, DVD-Ig can be designed to either be physically linked to medical devices implanted into patients or target these medical devices (see Burke, Sandra E.; Kuntz, Richard E.; Schwartz, Lewis B., Zotarolimus eluting stents. *Advanced Drug Delivery Reviews* (2006), 58(3), 437-446; Surface coatings for biological activation and functionalization of medical devices, Hildebrand, H. F.; Blanchemain, N.; Mayer, G.; Chai, F.; Lefebvre, M.; Boschini, F., *Surface and Coatings Technology* (2006), 200(22-23), 6318-6324; Drug/ device combinations for local drug therapies and infection prophylaxis, Wu, Peng; Grainger, David W., *Biomaterials* (2006), 27(11), 2450-2467; Mediation of the cytokine network in the implantation of orthopedic devices., Marques, A. P.; Hunt, J. A.; Reis, Rui L., *Biodegradable Systems in Tissue Engineering and Regenerative Medicine* (2005), 377-397). Briefly, directing appropriate types of cell to the site of medical implant may promote healing and restoring normal tissue function. Alternatively, inhibition of mediators (including but not limited to cytokines), released upon device implantation by a DVD-Ig coupled to or target to a device is also provided. For example, Stents have been used for years in interventional cardiology to clear blocked arteries and to improve the flow of blood to the heart muscle. However, traditional bare metal stents have been known to cause restenosis (re-narrowing of the artery in a treated area) in some patients and can lead to blood clots. Recently, an anti-CD34 antibody coated stent has been described which reduced restenosis and prevents blood clots from occurring by capturing endothelial progenitor cells (EPC) circulating throughout the blood. Endothelial cells are cells that line blood vessels, allowing blood to flow smoothly. The EPCs adhere to the hard surface of the stent forming a smooth layer that not only promotes healing but prevents restenosis and blood clots, complications previously associated with the use of stents (Aoji et al. 2005 *J Am Coll Cardiol.* 45(10):1574-9). In addition to improving outcomes for patients requiring stents, there are also implications for patients requiring cardiovascular bypass surgery. For example, a prosthetic vascular conduit (artificial artery) coated with anti-EPC antibodies would eliminate the need to use arteries from patients legs or arms for bypass surgery grafts. This would reduce surgery and anesthesia times, which in turn will reduce coronary surgery deaths. DVD-Ig are designed in such a way that it binds to a cell surface marker (such as CD34) as well as a protein (or an epitope of any kind, including but not limited to proteins, lipids and polysaccharides) that has been coated on the implanted device to facilitate the cell recruitment. Such approaches can also be applied to

other medical implants in general. Alternatively, DVD-IgS can be coated on medical devices and upon implantation and releasing all DVD-IgS from the device (or any other need which may require additional fresh DVD-Ig, including aging and denaturation of the already loaded DVD-Ig) the device could be reloaded by systemic administration of fresh DVD-Ig to the patient, where
 5 the DVD-Ig is designed to binds to a target of interest (a cytokine, a cell surface marker (such as CD34) etc.) with one set of binding sites and to a target coated on the device (including a protein, an epitope of any kind, including but not limited to lipids, polysaccharides and polymers) with the other. This technology has the advantage of extending the usefulness of coated implants.

A. Use of DVD-IgS in various diseases

10 DVD-Ig molecules provided herein are also useful as therapeutic molecules to treat various diseases. Such molecules may bind one or more targets involved in a specific disease. Examples of such targets in various diseases are described below.

A1. Human Autoimmune and Inflammatory Response

Many proteins have been implicated in general autoimmune and inflammatory responses,
 15 including C5, CCL1 (I-309), CCL11 (eotaxin), CCL13 (mcp-4), CCL15 (MIP-1d), CCL16 (HCC-4), CCL17 (TARC), CCL18 (PARC), CCL19, CCL2 (mcp-1), CCL20 (MIP-3a), CCL21 (MIP-2), CCL23 (MPIF-1), CCL24 (MPIF-2 / eotaxin-2), CCL25 (TECK), CCL26, CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (mcp-3), CCL8 (mcp-2), CXCL1, CXCL10 (IP-10), CXCL11 (I-TAC / IP-9), CXCL12 (SDF1), CXCL13, CXCL14, CXCL2, CXCL3, CXCL5
 20 (ENA-78 / LIX), CXCL6 (GCP-2), CXCL9, IL13, IL8, CCL13 (mcp-4), CCR1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CX3CR1, IL8RA, XCR1 (CCXCR1), IFNA2, IL10, IL13, IL17C, IL1A, IL1B, IL1F10, IL1F5, IL1F6, IL1F7, IL1F8, IL1F9, IL22, IL5, IL8, IL9, LTA, LTB, MIF, SCYE1 (endothelial Monocyte-activating cytokine), SPP1, TNF, TNFSF5, IFNA2, IL10RA, IL10RB, IL13, IL13RA1, IL5RA, IL9, IL9R, ABCF1, BCL6, C3, C4A,
 25 CEBPB, CRP, ICEBERG, IL1R1, IL1RN, IL8RB, LTB4R, TOLLIP, FADD, IRAK1, IRAK2, MYD88, NCK2, TNFAIP3, TRADD, TRAF1, TRAF2, TRAF3, TRAF4, TRAF5, TRAF6, ACVR1, ACVR1B, ACVR2, ACVR2B, ACVRL1, CD28, CD3E, CD3G, CD3Z, CD69, CD80, CD86, CNR1, CTLA4, CYSLTR1, FCER1A, FCER2, FCGR3A, GPR44, HAVCR2, OPRD1, P2RX7, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10, BLR1, CCL1, CCL2,
 30 CCL3, CCL4, CCL5, CCL7, CCL8, CCL11, CCL13, CCL15, CCL16, CCL17, CCL18, CCL19, CCL20, CCL21, CCL22, CCL23, CCL24, CCL25, CCR1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CX3CL1, CX3CR1, CXCL1, CXCL2, CXCL3, CXCL5, CXCL6, CXCL10, CXCL11, CXCL12, CXCL13, CXCR4, GPR2, SCYE1, SDF2, XCL1, XCL2, XCR1,

AMH, AMHR2, BMPR1A, BMPR1B, BMPR2, C19orf10 (IL27w), CER1, CSF1, CSF2, CSF3, DKFZp451J0118, FGF2, GFII, IFNA1, IFNB1, IFNG, IGF1, IL1A, IL1B, IL1R1, IL1R2, IL2, IL2RA, IL2RB, IL2RG, IL3, IL4, IL4R, IL5, IL5RA, IL6, IL6R, IL6ST, IL7, IL8, IL8RA, IL8RB, IL9, IL9R, IL10, IL10RA, IL10RB, IL11, IL11RA, IL12A, IL12B, IL12RB1, IL12RB2, 5 IL13, IL13RA1, IL13RA2, IL15, IL15RA, IL16, IL17, IL17R, IL18, IL18R1, IL19, IL20, KITLG, LEP, LTA, LTB, LTB4R, LTB4R2, LTBR, MIF, NPPB, PDGFB, TBX21, TDGF1, TGFA, TGFB1, TGFB1I1, TGFB2, TGFB3, TGFB1, TGFB1R1, TGFB1R2, TGFB1R3, TII1L, TNF, TNFRSF1A, TNFRSF1B, TNFRSF7, TNFRSF8, TNFRSF9, TNFRSF11A, TNFRSF21, TNFSF4, TNFSF5, TNFSF6, TNFSF11, VEGF, ZFPM2, and RNF110 (ZNF144). In one aspect, 10 DVD-Igs capable of binding one or more of the targets listed herein are provided.

DVD Igs capable of binding the following pairs of targets to treat inflammatory disease are contemplated: TNF and IL-17A; TNF and RANKL; TNF and VEGF; TNF and SOST (seq. 1); TNF and DKK; TNF and alphaVbeta3; TNF and NGF; TNF and IL-23p19; TNF and IL-6; TNF and SOST (seq. 2); TNF and IL-6R; TNF and CD-20; TNF and LPA; TNF and PGE2; IgE 15 and IL-13 (seq. 1); IL-13 (seq. 1) and IL23p19; IgE and IL-4; IgE and IL-9 (seq. 1); IgE and IL-9 (seq. 2); IgE and IL-13 (seq. 2); IL-13 (seq. 1) and IL-9 (seq. 1); IL-13 (seq. 1) and IL-4; IL-13 (seq. 1) and IL-9 (seq. 2); IL-13 (seq. 2) and IL-9 (seq. 1); IL-13 (seq. 2) and IL-4; IL-13 (seq. 2) and IL-23p19; IL-13 (seq. 2) and IL-9 (seq. 2); IL-6R and VEGF; IL-6R and IL-17A; IL-6R and RANKL; IL-17A and IL-1beta (seq. 1); IL-1beta (seq. 1) and RANKL; IL-1beta (seq. 1) and 20 VEGF; RANKL and CD-20; IL-1alpha and IL-1beta (seq. 1); IL-1alpha and IL-1beta (seq. 2) (see Examples 2.1 to 2.40).

A2. Asthma

Allergic asthma is characterized by the presence of eosinophilia, goblet cell metaplasia, epithelial cell alterations, airway hyperreactivity (AHR), and Th2 and Th1 cytokine expression, 25 as well as elevated serum IgE levels. It is now widely accepted that airway inflammation is the key factor underlying the pathogenesis of asthma, involving a complex interplay of inflammatory cells such as T cells, B cells, eosinophils, mast cells and macrophages, and of their secreted mediators including cytokines and chemokines. Corticosteroids are the most important anti-inflammatory treatment for asthma today, however their mechanism of action is non-specific and 30 safety concerns exist, especially in the juvenile patient population. The development of more specific and targeted therapies is therefore warranted. There is increasing evidence that IL-13 in mice mimics many of the features of asthma, including AHR, mucus hypersecretion and airway fibrosis, independently of eosinophilic inflammation (Finotto et al., *International Immunology* (2005), 17(8), 993-1007; Padilla et al., *Journal of Immunology* (2005), 174(12), 8097-8105).

IL-13 has been implicated as having a pivotal role in causing pathological responses associated with asthma. The development of anti-IL-13 mAb therapy to reduce the effects of IL-13 in the lung is an exciting new approach that offers considerable promise as a novel treatment for asthma. However other mediators of differential immunological pathways are also involved in asthma pathogenesis, and blocking these mediators, in addition to IL-13, may offer additional therapeutic benefit. Such target pairs include, but are not limited to, IL-13 and a pro-inflammatory cytokine, such as tumor necrosis factor- α (TNF- α). TNF- α may amplify the inflammatory response in asthma and may be linked to disease severity (McDonnell, et al., *Progress in Respiratory Research* (2001), 31(New Drugs for Asthma, Allergy and COPD), 247-250.). This suggests that blocking both IL-13 and TNF- α may have beneficial effects, particularly in severe airway disease. In another embodiment the DVD-Ig binds the targets IL-13 and TNF α and is used for treating asthma.

Animal models such as OVA-induced asthma mouse model, where both inflammation and AHR can be assessed, are known in the art and may be used to determine the ability of various DVD-Ig molecules to treat asthma. Animal models for studying asthma are disclosed in Coffman, et al., *Journal of Experimental Medicine* (2005), 201(12), 1875-1879; Lloyd, et al., *Advances in Immunology* (2001), 77, 263-295; Boyce et al., *Journal of Experimental Medicine* (2005), 201(12), 1869-1873; and Snibson, et al., *Journal of the British Society for Allergy and Clinical Immunology* (2005), 35(2), 146-52. In addition to routine safety assessments of these target pairs specific tests for the degree of immunosuppression may be warranted and helpful in selecting the best target pairs (see Luster et al., *Toxicology* (1994), 92(1-3), 229-43; Descotes, et al., *Developments in biological standardization* (1992), 77 99-102; Hart et al., *Journal of Allergy and Clinical Immunology* (2001), 108(2), 250-257).

Based on the rationale disclosed herein and using the same evaluation model for efficacy and safety other pairs of targets that DVD-Ig molecules can bind and be useful to treat asthma may be determined. In an embodiment, such targets include, but are not limited to, IL-13 and IL-1beta, since IL-1beta is also implicated in inflammatory response in asthma; IL-13 and cytokines and chemokines that are involved in inflammation, such as IL-13 and IL-9; IL-13 and IL-4; IL-13 and IL-5; IL-13 and IL-25; IL-13 and TARC; IL-13 and MDC; IL-13 and MIF; IL-13 and TGF- β ; IL-13 and LHR agonist; IL-13 and CL25; IL-13 and SPRR2a; IL-13 and SPRR2b; and IL-13 and ADAM8. DVD-Igs capable of binding one or more targets involved in asthma are also provided. In some embodiments the targets are CSF1 (MCSF), CSF2 (GM-CSF), CSF3 (GCSF), FGF2, IFNA1, IFNB1, IFNG, histamine and histamine receptors, IL1A, IL1B, IL2, IL3, IL4, IL5, IL6, IL7, IL8, IL9, IL10, IL11, IL12A, IL12B, IL13, IL14, IL15, IL16, IL17, IL18, IL19, KITLG,

PDGFB, IL2RA, IL4R, IL5RA, IL8RA, IL8RB, IL12RB1, IL12RB2, IL13RA1, IL13RA2, IL18R1, TSLP, CCL1, CCL2, CCL3, CCL4, CCL5, CCL7, CCL8, CCL13, CCL17, CCL18, CCL19, CCL20, CCL22, CCL24, CX3CL1, CXCL1, CXCL2, CXCL3, XCL1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CX3CR1, GPR2, XCR1, FOS, GATA3, JAK1, JAK3, 5 STAT6, TBX21, TGFB1, TNF, TNFSF6, YY1, CYSLTR1, FCER1A, FCER2, LTB4R, TB4R2, LTBR, or Chitinase.

A3. Rheumatoid arthritis

Rheumatoid arthritis (RA), a systemic disease, is characterized by a chronic inflammatory reaction in the synovium of joints and is associated with degeneration of cartilage and erosion of juxta-articular bone. Many pro-inflammatory cytokines including TNF, 10 chemokines, and growth factors are expressed in diseased joints. Systemic administration of anti-TNF antibody or sTNFR fusion protein to mouse models of RA was shown to be anti-inflammatory and joint protective. Clinical investigations in which the activity of TNF in RA patients was blocked with intravenously administered infliximab (Harriman G, Harper LK, 15 Schaible TF. 1999 Summary of clinical trials in rheumatoid arthritis using infliximab, an anti-TNF α treatment. *Ann Rheum Dis* 58 Suppl 1:161-4), a chimeric anti-TNF mAb, has provided evidence that TNF regulates IL-6, IL-8, MCP-1, and VEGF production, recruitment of immune and inflammatory cells into joints, angiogenesis, and reduction of blood levels of matrix metalloproteinases-1 and -3. A better understanding of the inflammatory pathway in rheumatoid 20 arthritis has led to identification of other therapeutic targets involved in rheumatoid arthritis. Promising treatments such as interleukin-6 antagonists (IL-6 receptor antibody MRA, developed by Chugai, Roche (see Nishimoto, Norihiro et al., *Arthritis & Rheumatism* (2004), 50(6), 1761-1769), CTLA4Ig (abatacept, Genovese Mc et al 2005 Abatacept for rheumatoid arthritis refractory to tumor necrosis factor alpha inhibition. *N Engl J Med.* 353:1114-23.), and anti-B cell 25 therapy (rituximab, Okamoto H, Kamatani N. 2004 Rituximab for rheumatoid arthritis. *N Engl J Med.* 351:1909) have already been tested in randomized controlled trials over the past year. Other cytokines have been identified and have been shown to be of benefit in animal models, including interleukin-15 (therapeutic antibody HuMax-IL₁₅, AMG 714 see Baslund, Bo et al., *Arthritis & Rheumatism* (2005), 52(9), 2686-2692), interleukin-17, and interleukin-18, and clinical trials of 30 these agents are currently under way. Dual-specific antibody therapy, combining anti-TNF and another mediator, has great potential in enhancing clinical efficacy and/or patient coverage. For example, blocking both TNF and VEGF can potentially eradicate inflammation and angiogenesis, both of which are involved in pathophysiology of RA. Blocking other pairs of targets involved in RA including, but not limited to, TNF and IL-18; TNF and IL-12; TNF and IL-23; TNF and IL-

Ibeta; TNF and MIF; TNF and IL-17; and TNF and IL-15 with specific DVD Igs is also contemplated. In addition to routine safety assessments of these target pairs, specific tests for the degree of immunosuppression may be warranted and helpful in selecting the best target pairs (see Luster et al., *Toxicology* (1994), 92(1-3), 229-43; Descotes, et al., *Developments in biological standardization* (1992), 77 99-102; Hart et al., *Journal of Allergy and Clinical Immunology* (2001), 108(2), 250-257). Whether a DVD Ig molecule will be useful for the treatment of rheumatoid arthritis can be assessed using pre-clinical animal RA models such as the collagen-induced arthritis mouse model. Other useful models are also well known in the art (see Brand DD., *Comp Med.* (2005) 55(2):114-22). Based on the cross-reactivity of the parental antibodies for human and mouse orthologues (e.g., reactivity for human and mouse TNF, human and mouse IL-15 etc.) validation studies in the mouse CIA model may be conducted with "matched surrogate antibody" derived DVD-Ig molecules; briefly, a DVD-Ig based on two (or more) mouse target specific antibodies may be matched to the extent possible to the characteristics of the parental human or humanized antibodies used for human DVD-Ig construction (similar affinity, similar neutralization potency, similar half-life etc.).

A4. SLE

The immunopathogenic hallmark of SLE is the polyclonal B cell activation, which leads to hyperglobulinemia, autoantibody production and immune complex formation. The fundamental abnormality appears to be the failure of T cells to suppress the forbidden B cell clones due to generalized T cell dysregulation. In addition, B and T-cell interaction is facilitated by several cytokines such as IL-10 as well as co-stimulatory molecules such as CD40 and CD40L, B7 and CD28 and CTLA-4, which initiate the second signal. These interactions together with impaired phagocytic clearance of immune complexes and apoptotic material, perpetuate the immune response with resultant tissue injury. The following targets may be involved in SLE and can potentially be used for DVD-Ig approach for therapeutic intervention: B cell targeted therapies: CD-20, CD-22, CD-19, CD28, CD4, CD80, HLA-DRA, IL10, IL2, IL4, TNFRSF5, TNFRSF6, TNFSF5, TNFSF6, BLR1, HDAC4, HDAC5, HDAC7A, HDAC9, ICOSL, IGBP1, MS4A1, RGS1, SLA2, CD81, IFNB1, IL10, TNFRSF5, TNFRSF7, TNFSF5, AICDA, BLNK, GALNAC4S-6ST, HDAC4, HDAC5, HDAC7A, HDAC9, IL10, IL11, IL4, INHA, INHBA, KLF6, TNFRSF7, CD28, CD38, CD69, CD80, CD83, CD86, DPP4, FCER2, IL2RA, TNFRSF8, TNFSF7, CD24, CD37, CD40, CD72, CD74, CD79A, CD79B, CR2, IL1R2, ITGA2, ITGA3, MS4A1, ST6GAL1, CD1C, CHST10, HLA-A, HLA-DRA, and NT5E.; co-stimulatory signals: CTLA4 or B7.1/B7.2; inhibition of B cell survival: BlyS, BAFF; Complement inactivation: C5; Cytokine modulation: the key principle is that the net biologic response in any tissue is the result

of a balance between local levels of proinflammatory or anti-inflammatory cytokines (see Sfikakis PP et al 2005 *Curr Opin Rheumatol* 17:550-7). SLE is considered to be a Th-2 driven disease with documented elevations in serum IL-4, IL-6, IL-10. In some embodiments the DVD Igs are capable of binding IL-4, IL-6, IL-10, IFN- α , or TNF- α . Combination of targets discussed herein will enhance therapeutic efficacy for SLE which can be tested in a number of lupus preclinical models (see Peng SL (2004) *Methods Mol Med.*;102:227-72). Based on the cross-reactivity of the parental antibodies for human and mouse orthologues (e.g., reactivity for human and mouse CD20, human and mouse Interferon alpha etc.) validation studies in a mouse lupus model may be conducted with "matched surrogate antibody" derived DVD-Ig molecules; briefly, a DVD-Ig based two (or more) mouse target specific antibodies may be matched to the extent possible to the characteristics of the parental human or humanized antibodies used for human DVD-Ig construction (similar affinity, similar neutralization potency, similar half-life etc.).

A5. Multiple sclerosis

Multiple sclerosis (MS) is a complex human autoimmune-type disease with a predominantly unknown etiology. Immunologic destruction of myelin basic protein (MBP) throughout the nervous system is the major pathology of multiple sclerosis. MS is a disease of complex pathologies, which involves infiltration by CD4+ and CD8+ T cells and of response within the central nervous system. Expression in the CNS of cytokines, reactive nitrogen species and costimulator molecules have all been described in MS. Of major consideration are immunological mechanisms that contribute to the development of autoimmunity. In particular, antigen expression, cytokine and leukocyte interactions, and regulatory T-cells, which help balance/modulate other T-cells such as Th1 and Th2 cells, are important areas for therapeutic target identification.

IL-12 is a proinflammatory cytokine that is produced by APC and promotes differentiation of Th1 effector cells. IL-12 is produced in the developing lesions of patients with MS as well as in EAE-affected animals. Previously it was shown that interference in IL-12 pathways effectively prevents EAE in rodents, and that *in vivo* neutralization of IL-12p40 using an anti-IL-12 mAb has beneficial effects in the myelin-induced EAE model in common marmosets.

TWEAK is a member of the TNF family, constitutively expressed in the central nervous system (CNS), with pro-inflammatory, proliferative or apoptotic effects depending upon cell types. Its receptor, Fn14, is expressed in CNS by endothelial cells, reactive astrocytes and neurons. TWEAK and Fn14 mRNA expression increased in spinal cord during experimental autoimmune encephalomyelitis (EAE). Anti-TWEAK antibody treatment in myelin

oligodendrocyte glycoprotein (MOG) induced EAE in C57BL/6 mice resulted in a reduction of disease severity and leukocyte infiltration when mice were treated after the priming phase.

One embodiment pertains to DVD Ig molecules capable of binding one or more, for example two, targets. In some embodiments, the targets are IL-12, TWEAK, IL-23, CXCL13, CD40, CD40L, IL-18, VEGF, VLA-4, TNF, CD45RB, CD200, IFN γ , GM-CSF, FGF, C5, CD52, or CCR2. An embodiment includes a dual-specific anti-IL-12/TWEAK DVD Ig as a therapeutic agent beneficial for the treatment of MS.

Several animal models for assessing the usefulness of the DVD-Ig molecules to treat MS are known in the art (see Steinman L, et al., (2005) Trends Immunol. 26(11):565-71; Lublin FD, et al., (1985) Springer Semin Immunopathol.8(3):197-208; Genain CP, et al., (1997) J Mol Med. 75(3):187-97; Tuohy VK, et al., (1999) J Exp Med. 189(7):1033-42; Owens T, et al., (1995) Neurol Clin.13(1):51-73; and Hart BA, et al., (2005) J Immunol 175(7):4761-8. Based on the cross-reactivity of the parental antibodies for human and animal species orthologues (e.g., reactivity for human and mouse IL-12, human and mouse TWEAK etc.) validation studies in the mouse EAE model may be conducted with "matched surrogate antibody" derived DVD-Ig molecules; briefly, a DVD-Ig based on to (or more) mouse target specific antibodies may be matched to the extent possible to the characteristics of the parental human or humanized antibodies used for human DVD-Ig construction (similar affinity, similar neutralization potency, similar half-life etc.). The same concept applies to animal models in other non-rodent species, where a "matched surrogate antibody" derived DVD-Ig would be selected for the anticipated pharmacology and possibly safety studies. In addition to routine safety assessments of these target pairs specific tests for the degree of immunosuppression may be warranted and helpful in selecting the best target pairs (see Luster et al., Toxicology (1994), 92(1-3), 229-43; Descotes, et al., Developments in biological standardization (1992), 77 99-102; Jones R. 2000 Rovelizumab (ICOS Corp). IDrugs.3(4):442-6).

A6. Sepsis

The pathophysiology of sepsis is initiated by the outer membrane components of both gram-negative organisms (lipopolysaccharide [LPS], lipid A, endotoxin) and gram-positive organisms (lipoteichoic acid, peptidoglycan). These outer membrane components are able to bind to the CD14 receptor on the surface of monocytes. By virtue of the recently described toll-like receptors, a signal is then transmitted to the cell, leading to the eventual production of the proinflammatory cytokines tumor necrosis factor-alpha (TNF-alpha) and interleukin-1 (IL-1). Overwhelming inflammatory and immune responses are essential features of septic shock and

play a central part in the pathogenesis of tissue damage, multiple organ failure, and death induced by sepsis. Cytokines, especially tumor necrosis factor (TNF) and interleukin (IL-1), have been shown to be critical mediators of septic shock. These cytokines have a direct toxic effect on tissues; they also activate phospholipase A2. These and other effects lead to increased concentrations of platelet-activating factor, promotion of nitric oxide synthase activity, promotion of tissue infiltration by neutrophils, and promotion of neutrophil activity.

The treatment of sepsis and septic shock remains a clinical conundrum, and recent prospective trials with biological response modifiers (i.e. anti-TNF, anti-MIF) aimed at the inflammatory response have shown only modest clinical benefit. Recently, interest has shifted toward therapies aimed at reversing the accompanying periods of immune suppression. Studies in experimental animals and critically ill patients have demonstrated that increased apoptosis of lymphoid organs and some parenchymal tissues contribute to this immune suppression, anergy, and organ system dysfunction. During sepsis syndromes, lymphocyte apoptosis can be triggered by the absence of IL-2 or by the release of glucocorticoids, granzymes, or the so-called 'death' cytokines: tumor necrosis factor alpha or Fas ligand. Apoptosis proceeds via auto-activation of cytosolic and/or mitochondrial caspases, which can be influenced by the pro- and anti-apoptotic members of the Bcl-2 family. In experimental animals, not only can treatment with inhibitors of apoptosis prevent lymphoid cell apoptosis; it may also improve outcome. Although clinical trials with anti-apoptotic agents remain distant due in large part to technical difficulties associated with their administration and tissue targeting, inhibition of lymphocyte apoptosis represents an attractive therapeutic target for the septic patient. Likewise, a dual-specific agent targeting both inflammatory mediator and a apoptotic mediator, may have added benefit. One embodiment pertains to DVD Igs capable of binding one or more targets involved in sepsis, in an embodiment two targets. In some embodiments, the targets are TNF, IL-1, MIF, IL-6, IL-8, IL-18, IL-12, IL-23, FasL, LPS, Toll-like receptors, TLR-4, tissue factor, MIP-2, ADORA2A, CASP1, CASP4, IL-10, IL-1B, NFKB1, PROC, TNFRSF1A, CSF3, CCR3, IL1RN, MIF, NFKB1, PTAFR, TLR2, TLR4, GPR44, HMOX1, midkine, IRAK1, NFKB2, SERPINA1, SERPINE1, or TREM1. The efficacy of such DVD Igs for sepsis can be assessed in preclinical animal models known in the art (see Buras JA, et al.,(2005) Nat Rev Drug Discov. 4(10):854-65 and Calandra T, et al., (2000) Nat Med. 6(2):164-70).

A7. Neurological disorders

A7.1. Neurodegenerative Diseases

Neurodegenerative diseases are either chronic in which case they are usually age-dependent or acute (e.g., stroke, traumatic brain injury, spinal cord injury, etc.). They are characterized by progressive loss of neuronal functions (neuronal cell death, demyelination), loss of mobility and loss of memory. Emerging knowledge of the mechanisms underlying chronic neurodegenerative diseases (e.g., Alzheimer's disease) show a complex etiology and a variety of factors have been recognized to contribute to their development and progression e.g., age, glycemic status, amyloid production and multimerization, accumulation of advanced glycation-end products (AGE) which bind to their receptor RAGE (receptor for AGE), increased brain oxidative stress, decreased cerebral blood flow, neuroinflammation including release of inflammatory cytokines and chemokines, neuronal dysfunction and microglial activation. Thus these chronic neurodegenerative diseases represent a complex interaction between multiple cell types and mediators. Treatment strategies for such diseases are limited and mostly constitute either blocking inflammatory processes with non-specific anti-inflammatory agents (e.g., corticosteroids, COX inhibitors) or agents to prevent neuron loss and/or synaptic functions. These treatments fail to stop disease progression. Recent studies suggest that more targeted therapies such as antibodies to soluble A- β peptide (including the A- β oligomeric forms) can not only help stop disease progression but may help maintain memory as well. These preliminary observations suggest that specific therapies targeting more than one disease mediator (e.g., A- β and a pro-inflammatory cytokine such as TNF) may provide even better therapeutic efficacy for chronic neurodegenerative diseases than observed with targeting a single disease mechanism (e.g., soluble A- β alone) (see C.E. Shepherd, et al, *Neurobiol Aging*. 2005 Oct 24; Nelson RB., *Curr Pharm Des*. 2005;11:3335; William L. Klein.; *Neurochem Int*. 2002 ;41:345; Michelle C Janelisins, et al., *J Neuroinflammation*. 2005 ;2:23; Soloman B., *Curr Alzheimer Res*. 2004;1:149; Igor Klyubin, et al., *Nat Med*. 2005;11:556-61; Arancio O, et al., *EMBO Journal* (2004) 1-10; Bornemann KD, et al., *Am J Pathol*. 2001;158:63; Deane R, et al., *Nat Med*. 2003;9:907-13; and Eliezer Masliah, et al., *Neuron*. 2005;46:857).

The DVD-Ig molecules provided herein can bind one or more targets involved in Chronic neurodegenerative diseases such as Alzheimers. Such targets include, but are not limited to, any mediator, soluble or cell surface, implicated in AD pathogenesis, e.g., AGE (S100 A, amphoterin), pro-inflammatory cytokines (e.g., IL-1), chemokines (e.g., MCP 1), molecules that inhibit nerve regeneration (e.g., Nogo, RGM A), molecules that enhance neurite growth (neurotrophins) and molecules that can mediate transport at the blood brain barrier (e.g.,

transferrin receptor, insulin receptor or RAGE). The efficacy of DVD-Ig molecules can be validated in pre-clinical animal models such as the transgenic mice that over-express amyloid precursor protein or RAGE and develop Alzheimer's disease-like symptoms. In addition, DVD-Ig molecules can be constructed and tested for efficacy in the animal models and the best therapeutic DVD-Ig can be selected for testing in human patients. DVD-Ig molecules can also be employed for treatment of other neurodegenerative diseases such as Parkinson's disease. Alpha-Synuclein is involved in Parkinson's pathology. A DVD-Ig capable of targeting alpha-synuclein and inflammatory mediators such as TNF, IL-1, MCP-1 can prove effective therapy for Parkinson's disease and are also contemplated.

10 A7.2 Neuronal Regeneration and Spinal Cord Injury

Despite an increase in knowledge of the pathologic mechanisms, spinal cord injury (SCI) is still a devastating condition and represents a medical indication characterized by a high medical need. Most spinal cord injuries are contusion or compression injuries and the primary injury is usually followed by secondary injury mechanisms (inflammatory mediators e.g., cytokines and chemokines) that worsen the initial injury and result in significant enlargement of the lesion area, sometimes more than 10-fold. These primary and secondary mechanisms in SCI are very similar to those in brain injury caused by other means e.g., stroke. No satisfying treatment exists and high dose bolus injection of methylprednisolone (MP) is the only used therapy within a narrow time window of 8 h post injury. This treatment, however, is only intended to prevent secondary injury without causing any significant functional recovery. It is heavily criticized for the lack of unequivocal efficacy and severe adverse effects, like immunosuppression with subsequent infections and severe histopathological muscle alterations. No other drugs, biologics or small molecules, stimulating the endogenous regenerative potential are approved, but promising treatment principles and drug candidates have shown efficacy in animal models of SCI in recent years. To a large extent the lack of functional recovery in human SCI is caused by factors inhibiting neurite growth, at lesion sites, in scar tissue, in myelin as well as on injury-associated cells. Such factors are the myelin-associated proteins NogoA, OMgp and MAG, RGM A, the scar-associated CSPG (Chondroitin Sulfate Proteoglycans) and inhibitory factors on reactive astrocytes (some semaphorins and ephrins). However, at the lesion site not only growth inhibitory molecules are found but also neurite growth stimulating factors like neurotrophins, laminin, L1 and others. This ensemble of neurite growth inhibitory and growth promoting molecules may explain that blocking single factors, like NogoA or RGM A, resulted in significant functional recovery in rodent SCI models, because a reduction of the inhibitory influences could shift the balance from growth inhibition to growth promotion. However,

recoveries observed with blocking a single neurite outgrowth inhibitory molecule were not complete. To achieve faster and more pronounced recoveries either blocking two neurite outgrowth inhibitory molecules, e.g., Nogo and RGM A, or blocking an neurite outgrowth inhibitory molecule and enhancing functions of a neurite outgrowth enhancing molecule, e.g.,

5 Nogo and neurotrophins, or blocking a neurite outgrowth inhibitory molecule, e.g., Nogo and a pro-inflammatory molecule e.g., TNF, may be desirable (see McGee AW, et al. (2003) Trends Neurosci. 26:193; Marco Domeniconi, et al. (2005) J. Neurol. Sci. 233:43; Milan Makwanal, et al. (2005) FEBS J. 272:2628; Barry J. Dickson (2002) Science 298:1959; Felicia Yu Hsuan Teng, et al. (2005) J. Neurosci. Res. 79:273; Tara Karnezis, et al. (2004) Nature Neuroscience 7:736;

10 Gang Xu, et al. (2004) J. Neurochem. 91:1018).

In one aspect, DVD-Igs capable of binding target pairs such as NgR and RGM A; NogoA and RGM A; MAG and RGM A; OMGp and RGM A; RGM A and RGM B; CSPGs and RGM A; aggrecan, midkine, neurocan, versican, phosphacan, Te38 and TNF- α ; A β globulomer-specific antibodies combined with antibodies promoting dendrite & axon sprouting are provided. Dendrite

15 pathology is a very early sign of AD and it is known that NOGO A restricts dendrite growth. One can combine such type of ab with any of the SCI-candidate (myelin-proteins) Ab. Other DVD-Ig targets may include any combination of NgR-p75, NgR-Troy, NgR-Nogo66 (Nogo), NgR-Lingo, Lingo-Troy, Lingo-p75, MAG or Ompg. Additionally, targets may also include any mediator, soluble or cell surface, implicated in inhibition of neurite e.g Nogo, Ompg, MAG, RGM A,

20 semaphorins, ephrins, soluble A-b, pro-inflammatory cytokines (e.g., IL-1), chemokines (e.g., MIP 1a), molecules that inhibit nerve regeneration. The efficacy of anti-nogo / anti-RGM A or similar DVD-Ig molecules can be validated in pre-clinical animal models of spinal cord injury. In addition, these DVD-Ig molecules can be constructed and tested for efficacy in the animal models and the best therapeutic DVD-Ig can be selected for testing in human patients. In addition, DVD-

25 Ig molecules can be constructed that target two distinct ligand binding sites on a single receptor e.g., Nogo receptor which binds three ligand Nogo, Ompg, and MAG and RAGE that binds A-b and S100 A. Furthermore, neurite outgrowth inhibitors e.g., nogo and nogo receptor, also play a role in preventing nerve regeneration in immunological diseases like multiple sclerosis. Inhibition of nogo-nogo receptor interaction has been shown to enhance recovery in animal models of

30 multiple sclerosis. Therefore, DVD-Ig molecules that can block the function of one immune mediator eg a cytokine like IL-12 and a neurite outgrowth inhibitor molecule eg nogo or RGM may offer faster and greater efficacy than blocking either an immune or an neurite outgrowth inhibitor molecule alone.

In general, antibodies do not cross the blood brain barrier (BBB) in an efficient and relevant manner. However, in certain neurologic diseases, e.g., stroke, traumatic brain injury, multiple sclerosis, etc., the BBB may be compromised and allows for increased penetration of DVD-Igs and antibodies into the brain. In other neurological conditions, where BBB leakage is not occurring, one may employ the targeting of endogenous transport systems, including carrier-mediated transporters such as glucose and amino acid carriers and receptor-mediated transcytosis-mediating cell structures/receptors at the vascular endothelium of the BBB, thus enabling trans-BBB transport of the DVD-Ig. Structures at the BBB enabling such transport include but are not limited to the insulin receptor, transferrin receptor, LRP and RAGE. In addition, strategies enable the use of DVD-Igs also as shuttles to transport potential drugs into the CNS including low molecular weight drugs, nanoparticles and nucleic acids (Coloma MJ, et al. (2000) *Pharm Res.* 17(3):266-74; Boado RJ, et al. (2007) *Bioconjug. Chem.* 18(2):447-55).

A8. Oncological disorders

Monoclonal antibody therapy has emerged as an important therapeutic modality for cancer (von Mehren, M., et al. (2003) *Annu. Rev. Med.* 54:343-69). Antibodies may exert antitumor effects by inducing apoptosis, redirected cytotoxicity, interfering with ligand-receptor interactions, or preventing the expression of proteins that are critical to the neoplastic phenotype. In addition, antibodies can target components of the tumor microenvironment, perturbing vital structures such as the formation of tumor-associated vasculature. Antibodies can also target receptors whose ligands are growth factors, such as the epidermal growth factor receptor. The antibody thus inhibits natural ligands that stimulate cell growth from binding to targeted tumor cells. Alternatively, antibodies may induce an anti-idiotypic network, complement-mediated cytotoxicity, or antibody-dependent cellular cytotoxicity (ADCC). The use of dual-specific binding proteins that target two separate tumor mediators will likely give additional benefit compared to a mono-specific therapy.

In another embodiment, the DVD Ig is capable of binding VEGF and phosphatidylserine; VEGF and ErbB3; VEGF and PLGF; VEGF and ROBO4; VEGF and BSG2; VEGF and CDCP1; VEGF and ANPEP; VEGF and c-MET; HER-2 and ERB3; HER-2 and BSG2; HER-2 and CDCP1; HER-2 and ANPEP; EGFR and CD64; EGFR and BSG2; EGFR and CDCP1; EGFR and ANPEP; IGF1R and PDGFR; IGF1R and VEGF; IGF1R and CD20; CD20 and CD74; CD20 and CD30; CD20 and DR4; CD20 and VEGFR2; CD20 and CD52; CD20 and CD4; HGF and c-MET; HGF and NRP1; HGF and phosphatidylserine; ErbB3 and IGF1R; ErbB3 and IGF1,2; c-Met and Her-2; c-Met and NRP1; c-Met and IGF1R; IGF1,2 and PDGFR; IGF1,2 and CD20; IGF1,2 and IGF1R; IGF2 and EGFR; IGF2 and HER2; IGF2 and CD20; IGF2 and VEGF; IGF2

and IGF1R; IGF1 and IGF2; PDGFRa and VEGFR2; PDGFRa and PLGF; PDGFRa and VEGF; PDGFRa and c-Met; PDGFRa and EGFR; PDGFRb and VEGFR2; PDGFRb and c-Met; PDGFRb and EGFR; RON and c-Met; RON and MTSP1; RON and MSP; RON and CDCP1; VGFR1 and PLGF; VGFR1 and RON; VGFR1 and EGFR; VEGFR2 and PLGF; VEGFR2 and NRP1; VEGFR2 and RON; VEGFR2 and DLL4; VEGFR2 and EGFR; VEGFR2 and ROBO4; VEGFR2 and CD55; LPA and S1P; EPHB2 and RON; CTLA4 and VEGF; CD3 and EPCAM; CD40 and IL6; CD40 and IGF; CD40 and CD56; CD40 and CD70; CD40 and VEGFR1; CD40 and DR5; CD40 and DR4; CD40 and APRIL; CD40 and BCMA; CD40 and RANKL; CD28 and MAPG; CD80 and CD40; CD80 and CD30; CD80 and CD33; CD80 and CD74; CD80 and CD2; CD80 and CD3; CD80 and CD19; CD80 and CD4; CD80 and CD52; CD80 and VEGF; CD80 and DR5; CD80 and VEGFR2; CD22 and CD20; CD22 and CD80; CD22 and CD40; CD22 and CD23; CD22 and CD33; CD22 and CD74; CD22 and CD19; CD22 and DR5; CD22 and DR4; CD22 and VEGF; CD22 and CD52; CD30 and CD20; CD30 and CD22; CD30 and CD23; CD30 and CD40; CD30 and VEGF; CD30 and CD74; CD30 and CD19; CD30 and DR5; CD30 and DR4; CD30 and VEGFR2; CD30 and CD52; CD30 and CD4; CD138 and RANKL; CD33 and FTL3; CD33 and VEGF; CD33 and VEGFR2; CD33 and CD44; CD33 and DR4; CD33 and DR5; DR4 and CD137; DR4 and IGF1,2; DR4 and IGF1R; DR4 and DR5; DR5 and CD40; DR5 and CD137; DR5 and CD20; DR5 and EGFR; DR5 and IGF1,2; DR5 and IGFR, DR5 and HER-2, and EGFR and DLL4. Other target combinations include one or more members of the EGF/erb-2/erb-3 family. Other targets (one or more) involved in oncological diseases that DVD Igs may bind include, but are not limited to: CD52, CD20, CD19, CD3, CD4, CD8, BMP6, IL12A, IL1A, IL1B, IL2, IL24, INHA, TNF, TNFSF10, BMP6, EGF, FGF1, FGF10, FGF11, FGF12, FGF13, FGF14, FGF16, FGF17, FGF18, FGF19, FGF2, FGF20, FGF21, FGF22, FGF23, FGF3, FGF4, FGF5, FGF6, FGF7, FGF8, FGF9, GRP, IGF1, IGF2, IL12A, IL1A, IL1B, IL2, INHA, TGFA, TGFB1, TGFB2, TGFB3, VEGF, CDK2, FGF10, FGF18, FGF2, FGF4, FGF7, IGF1R, IL2, BCL2, CD164, CDKN1A, CDKN1B, CDKN1C, CDKN2A, CDKN2B, CDKN2C, CDKN3, GNRH1, IGFBP6, IL1A, IL1B, ODZ1, PAWR, PLG, TGFB11, AR, BRCA1, CDK3, CDK4, CDK5, CDK6, CDK7, CDK9, E2F1, EGFR, ENO1, ERBB2, ESR1, ESR2, IGFBP3, IGFBP6, IL2, INSL4, MYC, NOX5, NR6A1, PAP, PCNA, PRKCQ, PRKD1, PRL, TP53, FGF22, FGF23, FGF9, IGFBP3, IL2, INHA, KLK6, TP53, CHGB, GNRH1, IGF1, IGF2, INHA, INSL3, INSL4, PRL, KLK6, SHBG, NR1D1, NR1H3, NR1I3, NR2F6, NR4A3, ESR1, ESR2, NR0B1, NR0B2, NR1D2, NR1H2, NR1H4, NR1I2, NR2C1, NR2C2, NR2E1, NR2E3, NR2F1, NR2F2, NR3C1, NR3C2, NR4A1, NR4A2, NR5A1, NR5A2, NR6A1, PGR, RARB, FGF1, FGF2, FGF6, KLK3, KRT1, APOC1, BRCA1, CHGA, CHGB, CLU, COL1A1, COL6A1, EGF, ERBB2, ERK8, FGF1, FGF10, FGF11, FGF13, FGF14, FGF16, FGF17, FGF18, FGF2, FGF20, FGF21, FGF22, FGF23, FGF3, FGF4, FGF5, FGF6, FGF7, FGF8, FGF9, GNRH1, IGF1, IGF2,

IGFBP3, IGFBP6, IL12A, IL1A, IL1B, IL2, IL24, INHA, INSL3, INSL4, KLK10, KLK12,
 KLK13, KLK14, KLK15, KLK3, KLK4, KLK5, KLK6, KLK9, MMP2, MMP9, MSMB, NTN4,
 ODZ1, PAP, PLAU, PRL, PSAP, SERPINA3, SHBG, TGFA, TIMP3, CD44, CDH1, CDH10,
 CDH19, CDH20, CDH7, CDH9, CDH1, CDH10, CDH13, CDH18, CDH19, CDH20, CDH7,
 5 CDH8, CDH9, ROBO2, CD44, ILK, ITGA1, APC, CD164, COL6A1, MTSS1, PAP, TGFB111,
 AGR2, AIG1, AKAP1, AKAP2, CANT1, CAV1, CDH12, CLDN3, CLN3, CYB5, CYC1,
 DAB2IP, DES, DNCL1, ELAC2, ENO2, ENO3, FASN, FLJ12584, FLJ25530, GAGEB1,
 GAGEC1, GGT1, GSTP1, HIP1, HUMCYT2A, IL29, K6HF, KAI1, KRT2A, MIB1, PART1,
 PATE, PCA3, PIAS2, PIK3CG, PPID, PR1, PSCA, SLC2A2, SLC33A1, SLC43A1, STEAP,
 10 STEAP2, TPM1, TPM2, TRPC6, ANGPT1, ANGPT2, ANPEP, ECGF1, EREG, FGF1, FGF2,
 FIGF, FLT1, JAG1, KDR, LAMA5, NRP1, NRP2, PGF, PLXDC1, STAB1, VEGF, VEGFC,
 ANGPTL3, BAI1, COL4A3, IL8, LAMA5, NRP1, NRP2, STAB1, ANGPTL4, PECAM1, PF4,
 PROK2, SERPINF1, TNFAIP2, CCL11, CCL2, CXCL1, CXCL10, CXCL3, CXCL5, CXCL6,
 CXCL9, IFNA1, IFNB1, IFNG, IL1B, IL6, MDK, EDG1, EFNA1, EFNA3, EFNB2, EGF,
 15 EPHB4, FGFR3, HGF, IGF1, ITGB3, PDGFA, TEK, TGFA, TGFB1, TGFB2, TGFB1, CCL2,
 CDH5, COL18A1, EDG1, ENG, ITGAV, ITGB3, THBS1, THBS2, BAD, BAG1, BCL2,
 CCNA1, CCNA2, CCND1, CCNE1, CCNE2, CDH1 (E-cadherin), CDKN1B (p27Kip1),
 CDKN2A (p16INK4a), COL6A1, CTNNB1 (b-catenin), CTSB (cathepsin B), ERBB2 (Her-2),
 ESR1, ESR2, F3 (TF), FOSL1 (FRA-1), GATA3, GSN (Gelsolin), IGFBP2, IL2RA, IL6, IL6R,
 20 IL6ST (glycoprotein 130), ITGA6 (a6 integrin), JUN, KLF5, KRT19, MAP2K7 (c-Jun), MKI67
 (Ki-67), NGFB (NGF), NGFR, NME1 (NM23A), PGR, PLAU (uPA), PTEN, SERPINB5
 (maspin), SERPINE1 (PAI-1), TGFA, THBS1 (thrombospondin-1), TIE (Tie-1), TNFRSF6 (Fas),
 TNFSF6 (FasL), TOP2A (topoisomerase Iia), TP53, AZGP1 (zinc-a-glycoprotein), BPAG1
 (plectin), CDKN1A (p21Wap1/Cip1), CLDN7 (claudin-7), CLU (clusterin), ERBB2 (Her-2),
 25 FGF1, FLRT1 (fibronectin), GABRP (GABAa), GNAS1, ID2, ITGA6 (a6 integrin), ITGB4 (b 4
 integrin), KLF5 (GC Box BP), KRT19 (Keratin 19), KRTHB6 (hair-specific type II keratin),
 MACMARCKS, MT3 (metallothionectin-III), MUC1 (mucin), PTGS2 (COX-2), RAC2
 (p21Rac2), S100A2, SCGB1D2 (lipophilin B), SCGB2A1 (mammaglobin 2), SCGB2A2
 (mammaglobin 1), SPRR1B (Spr1), THBS1, THBS2, THBS4, and TNFAIP2 (B94), RON, c-Met,
 30 CD64, DLL4, PLGF, CTLA4, phosphatidylserine, ROBO4, CD80, CD22, CD40, CD23, CD28,
 CD80, CD55, CD38, CD70, CD74, CD30, CD138, CD56, CD33, CD2, CD137, DR4, DR5,
 RANKL, VEGFR2, PDGFR, VEGFR1, MTSP1, MSP, EPHB2, EPHA1, EPHA2, EpCAM,
 PGE2, NKG2D, LPA, SIP, APRIL, BCMA, MAPG, FLT3, PDGFR alpha, PDGFR beta, ROR1,
 PSMA, PSCA, SCD1, or CD59.

IV. Pharmaceutical Compositions

Pharmaceutical compositions comprising a binding protein disclosed herein and a pharmaceutically acceptable carrier are also provided. The pharmaceutical compositions comprising binding proteins disclosed herein are for use in, but not limited to, diagnosing, 5 detecting, or monitoring a disorder, in preventing, treating, managing, or ameliorating of a disorder or one or more symptoms thereof, and/or in research. In a specific embodiment, a composition comprises one or more binding proteins disclosed herein. In another embodiment, the pharmaceutical composition comprises one or more binding proteins and one or more prophylactic or therapeutic agents other than binding proteins provided herein for treating a 10 disorder. In an embodiment, the prophylactic or therapeutic agents are known to be useful for or having been or currently being used in the prevention, treatment, management, or amelioration of a disorder or one or more symptoms thereof. In accordance with these embodiments, the composition may further comprise of a carrier, diluent or excipient.

The binding proteins provided herein can be incorporated into pharmaceutical 15 compositions suitable for administration to a subject. Typically, the pharmaceutical composition comprises a binding protein and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Examples of pharmaceutically acceptable carriers include one or 20 more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In some embodiments, isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride, are included in the composition. Pharmaceutically acceptable carriers may further comprise minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or 25 effectiveness of the binding protein or antigen-binding portion.

Various delivery systems are known and can be used to administer one or more binding proteins provided herein or the combination of one or more binding proteins and a prophylactic agent or therapeutic agent useful for preventing, managing, treating, or ameliorating a disorder or one or more symptoms thereof, e.g., encapsulation in liposomes, microparticles, microcapsules, 30 recombinant cells capable of expressing the binding protein or antigen-binding fragment, receptor-mediated endocytosis (see, e. g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of administering a prophylactic or therapeutic agents include, but are not limited to, parenteral administration (e.g., intradermal, intramuscular, intraperitoneal, intravenous and subcutaneous), epidurala

administration, intratumoral administration, and mucosal administration (e.g., intranasal and oral routes). In addition, pulmonary administration can be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent. See, e.g., U.S. Pat. Nos. 6,019,968; 5,985,320; 5,985,309; 5,934,272; 5,874,064; 5,855,913; 5,290,540; and 4,880,078; and PCT Publication Nos. WO 92/19244; WO 97/32572; WO 97/44013; WO 98/31346; and WO 99/66903. In one embodiment, a binding protein, combination therapy, or a composition disclosed herein is administered using Alkermes AIR® pulmonary drug delivery technology (Alkermes, Inc., Cambridge, Mass.). In a specific embodiment, prophylactic or therapeutic agents disclosed herein are administered intramuscularly, intravenously, intratumorally, orally, 5 intranasally, pulmonary, or subcutaneously. The prophylactic or therapeutic agents may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local.

15 In an embodiment, specific binding of antibody-coupled carbon nanotubes (CNTs) to tumor cells in vitro, followed by their highly specific ablation with near-infrared (NIR) light can be used to target tumor cells. For example, biotinylated polar lipids can be used to prepare stable, biocompatible, noncytotoxic CNT dispersions that are then attached to one or two different neutralite avidin-derivatized DVD-Igs directed against one or more tumor antigens (e.g., CD22) 20 (Chakravarty, P. et al. (2008) Proc. Natl. Acad. Sci. USA 105:8697-8702.

In a specific embodiment, it may be desirable to administer the prophylactic or therapeutic agents provided herein locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion, by injection, or by means of an implant, said implant being of a porous or non-porous material, including membranes and 25 matrices, such as sialastic membranes, polymers, fibrous matrices (e.g., Tissuel®), or collagen matrices. In one embodiment, an effective amount of one or more binding proteins provided herein is administered locally to the affected area to a subject to prevent, treat, manage, and/or ameliorate a disorder or a symptom thereof. In another embodiment, an effective amount of one or more binding proteins provided herein is administered locally to the affected area in 30 combination with an effective amount of one or more therapies (e.g., one or more prophylactic or therapeutic agents) other than a binding protein provided herein to prevent, treat, manage, and/or ameliorate a disorder or one or more symptoms thereof.

In another embodiment, the prophylactic or therapeutic agent can be delivered in a controlled release or sustained release system. In one embodiment, a pump may be used to

achieve controlled or sustained release (see Langer, *supra*; Sefton, 1987, *CRC Crit. Ref. Biomed. Eng.* 14:20; Buchwald et al., 1980, *Surgery* 88:507; Saudek et al., 1989, *N. Engl. J. Med.* 321:574). In another embodiment, polymeric materials can be used to achieve controlled or sustained release of the therapies provided herein (see e.g., *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Fla. (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, *J., Macromol. Sci. Rev. Macromol. Chem.* 23:61; see also Levy et al., 1985, *Science* 228:190; Doring et al., 1989, *Ann. Neurol.* 25:351; Howard et al., 1989, *J. Neurosurg.* 71:105); U.S. Pat. No. 5,679,377; U.S. Pat. No. 5,916,597; U.S. Pat. No. 5,912,015; U.S. Pat. No. 5,989,463; U.S. Pat. No. 5,128,326; PCT Publication No. WO 99/15154; and PCT Publication No. WO 99/20253. Examples of polymers used in sustained release formulations include, but are not limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl alcohol), polyacrylamide, poly(ethylene glycol), polylactides (PLA), poly(lactide-co-glycolides) (PLGA), and polyorthoesters. In an embodiment, the polymer used in a sustained release formulation is inert, free of leachable impurities, stable on storage, sterile, and biodegradable. In yet another embodiment, a controlled or sustained release system can be placed in proximity of the prophylactic or therapeutic target, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138 (1984)).

Controlled release systems are discussed in the review by Langer (1990, *Science* 249:1527-1533). Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more therapeutic agents provided herein. See, e.g., U.S. Pat. No. 4,526,938, PCT publication WO 91/05548, PCT publication WO 96/20698, Ning et al., 1996, "Intratumoral Radioimmunotherapy of a Human Colon Cancer Xenograft Using a Sustained-Release Gel," *Radiotherapy & Oncology* 39:179-189, Song et al., 1995, "Antibody Mediated Lung Targeting of Long-Circulating Emulsions," *PDA Journal of Pharmaceutical Science & Technology* 50:372-397, Cleek et al., 1997, "Biodegradable Polymeric Carriers for a bFGF Antibody for Cardiovascular Application," *Pro. Int'l. Symp. Control. Rel. Bioact. Mater.* 24:853-854, and Lam et al., 1997, "Microencapsulation of Recombinant Humanized Monoclonal Antibody for Local Delivery," *Proc. Int'l. Symp. Control Rel. Bioact. Mater.* 24:759-760.

In a specific embodiment, where the composition is a nucleic acid encoding a prophylactic or therapeutic agent, the nucleic acid can be administered *in vivo* to promote expression of its encoded prophylactic or therapeutic agent, by constructing it as part of an

appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see U. S. Pat. No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see, e.g., Joliot et al., 1991, Proc. Natl. Acad. Sci. USA 88:1864-1868). Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression by homologous recombination.

A pharmaceutical composition provided herein is formulated to be compatible with its intended route of administration. Examples of routes of administration include, but are not limited to, parenteral, e.g., intravenous, intradermal, subcutaneous, oral, intranasal (e.g., inhalation), transdermal (e.g., topical), transmucosal, and rectal administration. In a specific embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous, subcutaneous, intramuscular, oral, intranasal, or topical administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection.

If the compositions provided herein are to be administered topically, the compositions can be formulated in the form of an ointment, cream, transdermal patch, lotion, gel, shampoo, spray, aerosol, solution, emulsion, or other form well-known to one of skill in the art. See, e.g., Remington's Pharmaceutical Sciences and Introduction to Pharmaceutical Dosage Forms, 19th ed., Mack Pub. Co., Easton, Pa. (1995). In an embodiment, for non-sprayable topical dosage forms, viscous to semi-solid or solid forms comprising a carrier or one or more excipients compatible with topical application and having a dynamic viscosity greater than water are employed. Suitable formulations include, without limitation, solutions, suspensions, emulsions, creams, ointments, powders, liniments, salves, and the like, which are, if desired, sterilized or mixed with auxiliary agents (e.g., preservatives, stabilizers, wetting agents, buffers, or salts) for influencing various properties, such as, for example, osmotic pressure. Other suitable topical dosage forms include sprayable aerosol preparations wherein the active ingredient, in an embodiment, in combination with a solid or liquid inert carrier, is packaged in a mixture with a pressurized volatile (e.g., a gaseous propellant, such as freon) or in a squeeze bottle. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well-known in the art.

If the method provided herein comprises intranasal administration of a composition, the composition can be formulated in an aerosol form, spray, mist or in the form of drops. In particular, prophylactic or therapeutic agents can be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant (e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas). In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges (composed of, e.g., gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

If the method comprises oral administration, compositions can be formulated orally in the form of tablets, capsules, cachets, gels, solutions, suspensions, and the like. Tablets or capsules can be prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone, or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose, or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc, or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well-known in the art. Liquid preparations for oral administration may take the form of, but not limited to, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives, or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol, or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring, and sweetening agents as appropriate. Preparations for oral administration may be suitably formulated for slow release, controlled release, or sustained release of a prophylactic or therapeutic agent(s).

The method provided herein may comprise pulmonary administration, e.g., by use of an inhaler or nebulizer, of a composition formulated with an aerosolizing agent. See, e.g., U.S. Pat. Nos. 6,019,968; 5,985,320; 5,985,309; 5,934,272; 5,874,064; 5,855,913; 5,290,540; and 4,880,078; and PCT Publication Nos. WO 92/19244; WO 97/32572; WO 97/44013; WO 98/31346; and WO 99/66903. In a specific embodiment, a binding protein, combination therapy, and/or composition provided herein is administered using Alkermes AIR® pulmonary drug delivery technology (Alkermes, Inc., Cambridge, Mass.).

The method provided herein may comprise administration of a composition formulated for parenteral administration by injection (e. g., by bolus injection or continuous infusion). Formulations for injection may be presented in unit dosage form (e.g., in ampoules or in multi-dose containers) with an added preservative. The compositions may take such forms as
5 suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle (e.g., sterile pyrogen-free water) before use.

The methods may additionally comprise of administration of compositions formulated as
10 depot preparations. Such long acting formulations may be administered by implantation (e.g., subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compositions may be formulated with suitable polymeric or hydrophobic materials (e.g., as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives (e.g., as a sparingly soluble salt).

In some embodiments, the methods encompass administration of compositions
15 formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine,
20 procaine, etc.

Generally, the ingredients of compositions are supplied either separately or mixed
together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the mode of administration is infusion, composition can be dispensed with an
25 infusion bottle containing sterile pharmaceutical grade water or saline. Where the mode of administration is by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

In some embodiments, one or more of the prophylactic or therapeutic agents, or
pharmaceutical compositions provided herein is packaged in a hermetically sealed container such
30 as an ampoule or sachette indicating the quantity of the agent. In one embodiment, one or more of the prophylactic or therapeutic agents, or pharmaceutical compositions provided herein is supplied as a dry sterilized lyophilized powder or water free concentrate in a hermetically sealed container and can be reconstituted (e.g., with water or saline) to the appropriate concentration for

administration to a subject. In an embodiment, one or more of the prophylactic or therapeutic agents or pharmaceutical compositions provided herein is supplied as a dry sterile lyophilized powder in a hermetically sealed container at a unit dosage of at least 5 mg, at least 10 mg, at least 15 mg, at least 25 mg, at least 35 mg, at least 45 mg, at least 50 mg, at least 75 mg, or at least 100 mg. The lyophilized prophylactic or therapeutic agents or pharmaceutical compositions provided herein should be stored at between 2° C. and 8° C. in its original container and the prophylactic or therapeutic agents, or pharmaceutical compositions provided herein should be administered within 1 week, e.g., within 5 days, within 72 hours, within 48 hours, within 24 hours, within 12 hours, within 6 hours, within 5 hours, within 3 hours, or within 1 hour after being reconstituted.

In an alternative embodiment, one or more of the prophylactic or therapeutic agents or pharmaceutical compositions provided herein are supplied in liquid form in a hermetically sealed container indicating the quantity and concentration of the agent. In an embodiment, the liquid form of the administered composition is supplied in a hermetically sealed container at least 0.25 mg/ml, at least 0.5 mg/ml, at least 1 mg/ml, at least 2.5 mg/ml, at least 5 mg/ml, at least 8 mg/ml, at least 10 mg/ml, at least 15 mg/kg, at least 25 mg/ml, at least 50 mg/ml, at least 75 mg/ml or at least 100 mg/ml. The liquid form should be stored at between 2° C. and 8° C. in its original container.

The binding proteins provided herein can be incorporated into a pharmaceutical composition suitable for parenteral administration. In an embodiment, the binding protein or antigen-binding portions will be prepared as an injectable solution containing 0.1-250 mg/ml binding protein. The injectable solution can be composed of either a liquid or lyophilized dosage form in a flint or amber vial, ampule or pre-filled syringe. The buffer can be L-histidine (1-50 mM), optimally 5-10mM, at pH 5.0 to 7.0 (optimally pH 6.0). Other suitable buffers include but are not limited to, sodium succinate, sodium citrate, sodium phosphate or potassium phosphate.

Sodium chloride can be used to modify the toxicity of the solution at a concentration of 0-300 mM (optimally 150 mM for a liquid dosage form). Cryoprotectants can be included for a lyophilized dosage form, principally 0-10% sucrose (optimally 0.5-1.0%). Other suitable cryoprotectants include trehalose and lactose. Bulking agents can be included for a lyophilized dosage form, principally 1-10% mannitol (optimally 2-4%). Stabilizers can be used in both liquid and lyophilized dosage forms, principally 1-50 mM L-Methionine (optimally 5-10 mM). Other suitable bulking agents include glycine, arginine, can be included as 0-0.05% polysorbate-80 (optimally 0.005-0.01%). Additional surfactants include but are not limited to polysorbate 20 and BRIJ surfactants. In some embodiments, the pharmaceutical composition comprising the binding proteins provided herein is prepared as an injectable solution for parenteral administration, can further comprise an agent useful as an adjuvant, such as those used to increase the absorption, or

dispersion of a therapeutic protein (e.g., antibody). A particularly useful adjuvant is hyaluronidase, such as Hylenex® (recombinant human hyaluronidase). Addition of hyaluronidase in the injectable solution improves human bioavailability following parenteral administration, particularly subcutaneous administration. It also allows for greater injection site volumes (i.e. greater than 1 ml) with less pain and discomfort, and minimum incidence of injection site reactions. (see WO2004078140, and US2006104968).

The compositions provided herein may be in a variety of forms. These include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (e.g., injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders, liposomes and suppositories. The form chosen depends on the intended mode of administration and therapeutic application. Typical compositions are in the form of injectable or infusible solutions, such as compositions similar to those used for passive immunization of humans with other antibodies. The chosen mode of administration is parenteral (e.g., intravenous, subcutaneous, intraperitoneal, intramuscular). In an embodiment, the binding protein is administered by intravenous infusion or injection. In another embodiment, the binding protein is administered by intramuscular or subcutaneous injection.

Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating the active compound (i.e., antibody or antibody portion) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated herein, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated herein. In the case of sterile, lyophilized powders for the preparation of sterile injectable solutions, the methods of preparation are vacuum drying and spray-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including, in the composition, an agent that delays absorption, for example, monostearate salts and gelatin.

The binding proteins provided herein can be administered by a variety of methods known in the art, although for many therapeutic applications, in an embodiment, the route/mode of administration is subcutaneous injection, intravenous injection or infusion. As will be appreciated

by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. In certain embodiments, the active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. See, e.g., *Sustained and Controlled Release Drug Delivery Systems*, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

10 In certain embodiments, a binding protein provided herein may be orally administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer 15 a compound provided herein by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

Supplementary active compounds can also be incorporated into the compositions. In certain embodiments, a binding protein provided herein is coformulated with and/or 20 coadministered with one or more additional therapeutic agents that are useful for treating disorders with binding protein provided herein. For example, a binding protein provided herein may be coformulated and/or coadministered with one or more additional antibodies that bind other targets (e.g., antibodies that bind other cytokines or that bind cell surface molecules). Furthermore, one or more binding proteins provided herein may be used in combination with two 25 or more of the foregoing therapeutic agents. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

In certain embodiments, a binding protein is linked to a half-life extending vehicle known in the art. Such vehicles include, but are not limited to, the Fc domain, polyethylene glycol, and dextran. Such vehicles are described, e.g., in U.S. Application Serial No. 09/428,082 30 and published PCT Application No. WO 99/25044.

In a specific embodiment, nucleic acid sequences encoding a binding protein provided herein or another prophylactic or therapeutic agent are administered to treat, prevent, manage, or

ameliorate a disorder or one or more symptoms thereof by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment, the nucleic acids produce their encoded binding protein or prophylactic or therapeutic agent that mediates a prophylactic or therapeutic effect.

5 Any of the methods for gene therapy available in the art can be used. For general reviews of the methods of gene therapy, see Goldspiel et al., 1993, *Clinical Pharmacy* 12:488-505; Wu and Wu, 1991, *Biotherapy* 3:87-95; Tolstoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596; Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, 1993, *Ann. Rev. Biochem.* 62:191-217; May, 1993, *TIBTECH* 11(5):155-215. Methods commonly known in the
10 art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); and Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990). Detailed description of various methods of gene therapy are disclosed in US20050042664 A1.

The binding proteins provided herein are useful in treating various diseases wherein the
15 targets that are recognized by the binding proteins are detrimental. Such diseases include, but are not limited to, rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host
20 disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's granulomatosis, Henoch-Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia,
25 infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison's disease, sporadic, polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, seronegative
30 arthropathy, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella associated arthropathy, spondyloarthropathy, atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile

pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous
 candidiasis, giant cell arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis,
 Acquired Immunodeficiency Disease Syndrome, Acquired Immunodeficiency Related Diseases,
 Hepatitis B, Hepatitis C, common varied immunodeficiency (common variable
 5 hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature
 ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-inflammatory
 interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial
 lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis
 associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease,
 10 systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated
 lung disease, Sjögren's disease associated lung disease, ankylosing spondylitis associated lung
 disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced
 interstitial lung disease, fibrosis, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic
 pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty
 15 arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid
 hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated
 hypoglycaemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute
 immune disease associated with organ transplantation, chronic immune disease associated with
 organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis
 20 type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS,
 glomerulonephritides, microscopic vasculitis of the kidneys, lyme disease, discoid lupus
 erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all
 subtypes), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue
 disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute
 25 rheumatic fever, rheumatoid spondylitis, Still's disease, systemic sclerosis, Sjögren's syndrome,
 Takayasu's disease/arteritis, autoimmune thrombocytopaenia, idiopathic thrombocytopaenia,
 autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism
 (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary myxoedema, phacogenic
 uveitis, primary vasculitis, vitiligo acute liver disease, chronic liver diseases, alcoholic cirrhosis,
 30 alcohol-induced liver injury, choleosatis, idiosyncratic liver disease, Drug-Induced hepatitis,
 Non-alcoholic Steatohepatitis, allergy and asthma, group B streptococci (GBS) infection, mental
 disorders (*e.g.*, depression and schizophrenia), Th2 Type and Th1 Type mediated diseases, acute
 and chronic pain (different forms of pain), and cancers such as lung, breast, stomach, bladder,
 colon, pancreas, ovarian, prostate and rectal cancer and hematopoietic malignancies (leukemia
 35 and lymphoma), Abetalipoproteinemia, Acrocyanosis, acute and chronic parasitic or infectious
 processes, acute leukemia, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML),

acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinomas, aerial ectopic beats, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allograft rejection, alpha-1- antitrypsin deficiency, amyotrophic lateral sclerosis, anemia, angina pectoris, anterior horn cell degeneration, anti cd3
5 therapy, antiphospholipid syndrome, anti-receptor hypersensitivity reactions, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, arteriosclerosis, arteriovenous fistula, ataxia, atrial fibrillation (sustained or paroxysmal), atrial flutter, atrioventricular block, B cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt's lymphoma, Burns, cardiac arrhythmias, cardiac stun syndrome, cardiac tumors,
10 cardiomyopathy, cardiopulmonary bypass inflammation response, cartilage transplant rejection, cerebellar cortical degenerations, cerebellar disorders, chaotic or multifocal atrial tachycardia, chemotherapy associated disorders, chronic myelocytic leukemia (CML), chronic alcoholism, chronic inflammatory pathologies, chronic lymphocytic leukemia (CLL), chronic obstructive pulmonary disease (COPD), chronic salicylate intoxication, colorectal carcinoma, congestive
15 heart failure, conjunctivitis, contact dermatitis, cor pulmonale, coronary artery disease, Creutzfeldt-Jakob disease, culture negative sepsis, cystic fibrosis, cytokine therapy associated disorders, Dementia pugilistica, demyelinating diseases, dengue hemorrhagic fever, dermatitis, dermatologic conditions, diabetes, diabetes mellitus, diabetic atherosclerotic disease, Diffuse Lewy body disease, dilated congestive cardiomyopathy, disorders of the basal ganglia, Down's
20 Syndrome in middle age, drug- induced movement disorders induced by drugs which block CNS dopamine receptors, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottitis, epstein-barr virus infection, erythromelalgia, extrapyramidal and cerebellar disorders, familial hematophagocytic lymphohistiocytosis, fetal thymus implant rejection, Friedreich's ataxia, functional peripheral arterial disorders, fungal sepsis, gas gangrene, gastric ulcer,
25 glomerular nephritis, graft rejection of any organ or tissue, gram negative sepsis, gram positive sepsis, granulomas due to intracellular organisms, hairy cell leukemia, Hallerorden-Spatz disease, hashimoto's thyroiditis, hay fever, heart transplant rejection, hemachromatosis, hemodialysis, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, hemorrhage, hepatitis (A), His bundle arrhythmias, HIV infection/HIV neuropathy, Hodgkin's disease,
30 hyperkinetic movement disorders, hypersensitivity reactions, hypersensitivity pneumonitis, hypertension, hypokinetic movement disorders, hypothalamic-pituitary-adrenal axis evaluation, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody mediated cytotoxicity, Asthenia, infantile spinal muscular atrophy, inflammation of the aorta, influenza a, ionizing radiation exposure, iridocyclitis/uveitis/optic neuritis, ischemia- reperfusion injury, ischemic
35 stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, kidney transplant rejection, legionella, leishmaniasis, leprosy, lesions of the corticospinal system,

lipedema, liver transplant rejection, lymphedema, malaria, malignant Lymphoma, malignant histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic, migraine headache, mitochondrial multi.system disorder, mixed connective tissue disease, monoclonal gammopathy, multiple myeloma, multiple systems degenerations (Mencel Dejerine- Thomas Shi-
 5 Drager and Machado-Joseph), myasthenia gravis, mycobacterium avium intracellulare, mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disorders, nasopharyngeal carcinoma, neonatal chronic lung disease, nephritis, nephrosis, neurodegenerative diseases, neurogenic I muscular atrophies , neutropenic fever, non-hodgkins lymphoma, occlusion of the abdominal aorta and its branches, occlusive arterial
 10 disorders, okt3 therapy, orchitis/epididymitis, orchitis/vasectomy reversal procedures, organomegaly, osteoporosis, pancreas transplant rejection, pancreatic carcinoma, paraneoplastic syndrome/hypercalcemia of malignancy, parathyroid transplant rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral atherosclerotic disease, peripheral vascular disorders, peritonitis, pernicious anemia, pneumocystis carinii pneumonia, pneumonia,
 15 POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy, and skin changes syndrome), post perfusion syndrome, post pump syndrome, post-MI cardiomyopathy syndrome, preeclampsia, Progressive supranucleo Palsy, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon and disease, Raynaud's disease, Refsum's disease, regular narrow QRS tachycardia, renovascular hypertension, reperfusion injury, restrictive
 20 cardiomyopathy, sarcomas, scleroderma, senile chorea, Senile Dementia of Lewy body type, seronegative arthropathies, shock, sickle cell anemia, skin allograft rejection, skin changes syndrome, small bowel transplant rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degenerations, streptococcal myositis, structural lesions of the cerebellum, Subacute sclerosing panencephalitis, Syncope, syphilis of the cardiovascular system, systemic
 25 anaphalaxis, systemic inflammatory response syndrome, systemic onset juvenile rheumatoid arthritis, T-cell or FAB ALL, Telangiectasia, thromboangitis obliterans, thrombocytopenia, toxicity, transplants, trauma/hemorrhage, type III hypersensitivity reactions, type IV hypersensitivity, unstable angina, uremia, urosepsis, urticaria, valvular heart diseases, varicose veins, ,vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal
 30 infections, vital encephalitis/aseptic meningitis, vital-associated hemaphagocytic syndrome, Wernicke- Korsakoff syndrome, Wilson's disease, xenograft rejection of any organ or tissue. (see Peritt et al. PCT publication No. WO2002097048A2, Leonard et al., PCT publication No. WO9524918 A1, and Salfeld et al., PCT publication No. WO00/56772A1).

The DVD-Igs provided herein may also treat one or more of the following diseases:

35 Acute coronary syndromes, Acute Idiopathic Polyneuritis, Acute Inflammatory Demyelinating

Polyradiculoneuropathy, Acute ischemia, Adult Still's Disease, Alopecia areata, Anaphylaxis,
 Anti-Phospholipid Antibody Syndrome, Aplastic anemia, Arteriosclerosis, Atopic eczema,
 Atopic dermatitis, Autoimmune dermatitis, Autoimmune disorder associated with Streptococcus
 infection, Autoimmune hearingloss, Autoimmune Lymphoproliferative Syndrome (ALPS),
 5 Autoimmune myocarditis, autoimmune thrombocytopenia (AITP), Blepharitis, Bronchiectasis,
 Bullous pemphigoid, Cardiovascular Disease, Catastrophic Antiphospholipid Syndrome, Celiac
 Disease, Cervical Spondylosis, Chronic ischemia, Cicatricial pemphigoid, Clinically isolated
 Syndrome (CIS) with Risk for Multiple Sclerosis, Conjunctivitis, Childhood Onset Psychiatric
 Disorder, Chronic obstructive pulmonary disease (COPD), Dacryocystitis, dermatomyositis,
 10 Diabetic retinopathy, Diabetes mellitus, Disk herniation, Disk prolaps, Drug induced immune
 hemolytic anemia, Endocarditis, Endometriosis, endophthalmitis, , Episcleritis, Erythema
 multiforme, erythema multiforme major, Gestational pemphigoid, Guillain-Barré Syndrome
 (GBS), Hay Fever, Hughes Syndrome , Idiopathic Parkinson's Disease, idiopathic interstitial
 pneumonia, IgE-mediated Allergy, Immune hemolytic anemia, Inclusion Body Myositis,
 15 Infectious ocular inflammatory disease, Inflammatory demyelinating disease, Inflammatory heart
 disease, Inflammatory kidney disease, IPF/UIP, Iritis, Keratitis, Keratoconjunctivitis sicca, Kussmaul
 disease or Kussmaul-Meier Disease, Landry's Paralysis, Langerhan's Cell Histiocytosis, Livedo
 reticularis, Macular Degeneration, malignancies, Microscopic Polyangiitis, Morbus Bechterev,
 Motor Neuron Disorders, Mucous membrane pemphigoid, Multiple Organ failure, Myasthenia
 20 Gravis, Myelodysplastic Syndrome, Myocarditis, Nerve Root Disorders, Neuropathy, Non-A
 Non-B Hepatitis, Optic Neuritis, Osteolysis, Ovarian cancer, Pauciarticular JRA, peripheral
 artery occlusive disease (PAOD), peripheral vascular disease (PVD), peripheral artery disease
 (PAD), Phlebitis, Polyarteritis nodosa (or periarteritis nodosa), Polychondritis, Polymyalgia
 Rheumatica, Poliosis, Polyarticular JRA, Polyendocrine Deficiency Syndrome, Polymyositis,
 25 polymyalgia rheumatica (PMR), Post-Pump Syndrome, primary parkinsonism, prostate and rectal
 cancer and hematopoietic malignancies (leukemia and lymphoma), Prostatitis, Pure red cell
 aplasia, Primary Adrenal Insufficiency, Recurrent Neuromyelitis Optica, Restenosis, Rheumatic
 heart disease, SAPHO (synovitis, acne, pustulosis, hyperostosis, and osteitis), Scleroderma,
 Secondary Amyloidosis, Shock lung, Scleritis, Sciatica, Secondary Adrenal Insufficiency,
 30 Silicone associated connective tissue disease, Sneddon-Wilkinson Dermatitis, spondylitis
 ankylosans, Stevens-Johnson Syndrome (SJS), Systemic inflammatory response syndrome,
 Temporal arteritis, toxoplasmic retinitis, toxic epidermal necrolysis, Transverse myelitis, TRAPS
 (Tumor Necrosis Factor Receptor, Type 1 allergic reaction, Type II Diabetes, Urticaria, Usual
 interstitial pneumonia (UIP), Vasculitis, Vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-
 35 Harada syndrome (VKH syndrome), Wet macular degeneration, and Wound healing.

The binding proteins provided herein can be used to treat humans suffering from autoimmune diseases, in particular those associated with inflammation, including, rheumatoid arthritis, spondylitis, allergy, autoimmune diabetes, autoimmune uveitis. In an embodiment, the binding proteins or antigen-binding portions thereof, are used to treat rheumatoid arthritis, Crohn's disease, multiple sclerosis, insulin dependent diabetes mellitus and psoriasis.

In an embodiment, diseases that can be treated or diagnosed with the compositions and methods provided herein include, but are not limited to, primary and metastatic cancers, including carcinomas of breast, colon, rectum, lung, oropharynx, hypopharynx, esophagus, stomach, pancreas, liver, gallbladder and bile ducts, small intestine, urinary tract (including kidney, bladder and urothelium), female genital tract (including cervix, uterus, and ovaries as well as choriocarcinoma and gestational trophoblastic disease), male genital tract (including prostate, seminal vesicles, testes and germ cell tumors), endocrine glands (including the thyroid, adrenal, and pituitary glands), and skin, as well as hemangiomas, melanomas, sarcomas (including those arising from bone and soft tissues as well as Kaposi's sarcoma), tumors of the brain, nerves, eyes, and meninges (including astrocytomas, gliomas, glioblastomas, retinoblastomas, neuromas, neuroblastomas, Schwannomas, and meningiomas), solid tumors arising from hematopoietic malignancies such as leukemias, and lymphomas (both Hodgkin's and non-Hodgkin's lymphomas).

In an embodiment, the binding proteins provided herein or antigen-binding portions thereof, are used to treat cancer or in the prevention of metastases from the tumors described herein either when used alone or in combination with radiotherapy and/or other chemotherapeutic agents.

The binding proteins provided herein, or antigen binding portions thereof, may be combined with agents that include but are not limited to, antineoplastic agents, radiotherapy, chemotherapy such as DNA alkylating agents, cisplatin, carboplatin, anti-tubulin agents, paclitaxel, docetaxel, taxol, doxorubicin, gemcitabine, gemzar, anthracyclines, adriamycin, topoisomerase I inhibitors, topoisomerase II inhibitors, 5-fluorouracil (5-FU), leucovorin, irinotecan, receptor tyrosine kinase inhibitors (e.g., erlotinib, gefitinib), COX-2 inhibitors (e.g., celecoxib), kinase inhibitors, and siRNAs.

A binding proteins provided herein also can be administered with one or more additional therapeutic agents useful in the treatment of various diseases.

A binding protein provided herein can be used alone or in combination to treat such diseases. It should be understood that the binding proteins can be used alone or in combination

with an additional agent, e.g., a therapeutic agent, said additional agent being selected by the skilled artisan for its intended purpose. For example, the additional agent can be a therapeutic agent art-recognized as being useful to treat the disease or condition being treated by the binding proteins disclosed herein. The additional agent also can be an agent that imparts a beneficial
5 attribute to the therapeutic composition e.g., an agent which effects the viscosity of the composition.

It should further be understood that the combinations provided herein are those combinations useful for their intended purpose. The agents set forth below are illustrative for purposes and not intended to be limited. The combinations provided herein can be the binding
10 proteins and at least one additional agent selected from the lists below. The combinations can also include more than one additional agent, e.g., two or three additional agents if the combination is such that the formed composition can perform its intended function.

Combinations to treat autoimmune and inflammatory diseases are non-steroidal anti-inflammatory drug(s) also referred to as NSAIDS which include drugs like ibuprofen. Other
15 combinations are corticosteroids including prednisolone; the well known side-effects of steroid use can be reduced or even eliminated by tapering the steroid dose required when treating patients in combination with the DVD Igs provided herein. Non-limiting examples of therapeutic agents for rheumatoid arthritis with which a binding protein provided herein can be combined include the following: cytokine suppressive anti-inflammatory drug(s) (CSAIDs); antibodies to or
20 antagonists of other human cytokines or growth factors, for example, TNF, LT, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-15, IL-16, IL-18, IL-21, IL-23, interferons, EMAP-II, GM-CSF, FGF, and PDGF. Binding proteins provided herein, or antigen binding portions thereof, can be combined with antibodies to cell surface molecules such as CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD80 (B7.1), CD86 (B7.2), CD90, CTLA or their ligands including
25 CD154 (gp39 or CD40L).

Combinations of therapeutic agents may interfere at different points in the autoimmune and subsequent inflammatory cascade; examples include TNF antagonists like chimeric, humanized or human TNF antibodies, Adalimumab, (PCT Publication No. WO 97/29131), CA2 (RemicadeTM), CDP 571, and soluble p55 or p75 TNF receptors, derivatives, thereof,
30 (p75TNFR1gG (EnbrelTM) or p55TNFR1gG (Lenercept), and also TNF α converting enzyme (TACE) inhibitors; similarly IL-1 inhibitors (Interleukin-1-converting enzyme inhibitors, IL-1RA etc.) may be effective for the same reason. Other combinations include Interleukin 11. Yet another combination include key players of the autoimmune response which may act parallel to, dependent on or in concert with IL-12 function; especially are IL-18 antagonists including IL-18

antibodies or soluble IL-18 receptors, or IL-18 binding proteins. It has been shown that IL-12 and IL-18 have overlapping but distinct functions and a combination of antagonists to both may be most effective. Yet another combination are non-depleting anti-CD4 inhibitors. Yet other combinations include antagonists of the co-stimulatory pathway CD80 (B7.1) or CD86 (B7.2) including antibodies, soluble receptors or antagonistic ligands.

The binding proteins provided herein may also be combined with agents, such as methotrexate, 6-MP, azathioprine sulphasalazine, mesalazine, olsalazine chloroquine/hydroxychloroquine, pencillamine, aurothiomalate (intramuscular and oral), azathioprine, cochlincine, corticosteroids (oral, inhaled and local injection), beta-2 adrenoreceptor agonists (salbutamol, terbutaline, salmeteral), xanthines (theophylline, aminophylline), cromoglycate, nedocromil, ketotifen, ipratropium and oxitropium, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNF- α or IL-1 (e.g., IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TNF α converting enzyme (TACE) inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (e.g., soluble p55 or p75 TNF receptors and the derivatives p75TNFRlgG (EnbrelTM and p55TNFRlgG (Lenercept)), sIL-1RI, sIL-1RII, sIL-6R), antiinflammatory cytokines (e.g., IL-4, IL-10, IL-11, IL-13 and TGF β), celecoxib, folic acid, hydroxychloroquine sulfate, rofecoxib, etanercept, infliximab, naproxen, valdecoxib, sulfasalazine, methylprednisolone, meloxicam, methylprednisolone acetate, gold sodium thiomalate, aspirin, triamcinolone acetonide, propoxyphene napsylate/apap, folate, nabumetone, diclofenac, piroxicam, etodolac, diclofenac sodium, oxaprozin, oxycodone hcl, hydrocodone bitartrate/apap, diclofenac sodium/misoprostol, fentanyl, anakinra, human recombinant, tramadol hcl, salsalate, sulindac, cyanocobalamin/fa/pyridoxine, acetaminophen, alendronate sodium, prednisolone, morphine sulfate, lidocaine hydrochloride, indomethacin, glucosamine sulf/chondroitin, amitriptyline hcl, sulfadiazine, oxycodone hcl/acetaminophen, olopatadine hcl, misoprostol, naproxen sodium, omeprazole, cyclophosphamide, rituximab, IL-1 TRAP, MRA, CTLA4-IG, IL-18 BP, anti-IL-18, Anti-IL15, BIRB-796, SCIO-469, VX-702, AMG-548, VX-740, Roflumilast, IC-485, CDC-801, and Mesopram. Combinations include methotrexate or leflunomide and in moderate or severe rheumatoid arthritis cases, cyclosporine.

Nonlimiting additional agents which can also be used in combination with a binding protein to treat rheumatoid arthritis include, but are not limited to, the following: non-steroidal anti-inflammatory drug(s) (NSAIDs); cytokine suppressive anti-inflammatory drug(s) (CSAIDs); CDP-571/BAY-10-3356 (humanized anti-TNF α antibody; Celltech/Bayer); cA2/infliximab

5 (chimeric anti-TNF α antibody; Centocor); 75 kdTNFR-IgG/etanercept (75 kD TNF receptor-IgG fusion protein; Immunex; see *e.g.*, *Arthritis & Rheumatism* (1994) Vol. 37, S295; *J. Invest. Med.* (1996) Vol. 44, 235A); 55 kdTNF-IgG (55 kD TNF receptor-IgG fusion protein; Hoffmann-LaRoche); IDEC-CE9.1/SB 210396 (non-depleting primatized anti-CD4 antibody; IDEC/SmithKline; see *e.g.*, *Arthritis & Rheumatism* (1995) Vol. 38, S185); DAB 486-IL-2 and/or

10 DAB 389-IL-2 (IL-2 fusion proteins; Seragen; see *e.g.*, *Arthritis & Rheumatism* (1993) Vol. 36, 1223); Anti-Tac (humanized anti-IL-2R α ; Protein Design Labs/Roche); IL-4 (anti-inflammatory cytokine; DNAX/Schering); IL-10 (SCH 52000; recombinant IL-10, anti-inflammatory cytokine; DNAX/Schering); IL-4; IL-10 and/or IL-4 agonists (*e.g.*, agonist antibodies); IL-1RA (IL-1 receptor antagonist; Synergen/Amgen); anakinra (Kineret[®]/Amgen); TNF-bp/s-TNF (soluble

15 TNF binding protein; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S284; *Amer. J. Physiol. - Heart and Circulatory Physiology* (1995) Vol. 268, pp. 37-42); R973401 (phosphodiesterase Type IV inhibitor; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S282); MK-966 (COX-2 Inhibitor; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S81); Iloprost (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9

20 (supplement), S82); methotrexate; thalidomide (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S282) and thalidomide-related drugs (*e.g.*, Celgen); leflunomide (anti-inflammatory and cytokine inhibitor; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S131; *Inflammation Research* (1996) Vol. 45, pp. 103-107); tranexamic acid (inhibitor of plasminogen activation; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9

25 (supplement), S284); T-614 (cytokine inhibitor; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S282); prostaglandin E1 (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S282); Tenidap (non-steroidal anti-inflammatory drug; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S280); Naproxen (non-steroidal anti-inflammatory drug; see *e.g.*, *Neuro Report* (1996) Vol. 7, pp. 1209-1213); Meloxicam (non-

30 steroidal anti-inflammatory drug); Ibuprofen (non-steroidal anti-inflammatory drug); Piroxicam (non-steroidal anti-inflammatory drug); Diclofenac (non-steroidal anti-inflammatory drug); Indomethacin (non-steroidal anti-inflammatory drug); Sulfasalazine (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S281); Azathioprine (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S281); ICE inhibitor (inhibitor of the enzyme

35 interleukin-1 β converting enzyme); zap-70 and/or lck inhibitor (inhibitor of the tyrosine kinase zap-70 or lck); VEGF inhibitor and/or VEGF-R inhibitor (inhibitors of vascular endothelial cell

growth factor or vascular endothelial cell growth factor receptor; inhibitors of angiogenesis);
 corticosteroid anti-inflammatory drugs (e.g., SB203580); TNF-convertase inhibitors; anti-IL-12
 antibodies; anti-IL-18 antibodies; interleukin-11 (see e.g., *Arthritis & Rheumatism* (1996) Vol.
 39, No. 9 (supplement), S296); interleukin-13 (see e.g., *Arthritis & Rheumatism* (1996) Vol. 39,
 5 No. 9 (supplement), S308); interleukin -17 inhibitors (see e.g., *Arthritis & Rheumatism* (1996)
 Vol. 39, No. 9 (supplement), S120); gold; penicillamine; chloroquine; chlorambucil;
 hydroxychloroquine; cyclosporine; cyclophosphamide; total lymphoid irradiation; anti-thymocyte
 globulin; anti-CD4 antibodies; CD5-toxins; orally-administered peptides and collagen; lobenzarit
 disodium; Cytokine Regulating Agents (CRAs) HP228 and HP466 (Houghten Pharmaceuticals,
 10 Inc.); ICAM-1 antisense phosphorothioate oligo-deoxynucleotides (ISIS 2302; Isis
 Pharmaceuticals, Inc.); soluble complement receptor 1 (TP10; T Cell Sciences, Inc.); prednisone;
 orgotein; glycosaminoglycan polysulphate; minocycline; anti-IL2R antibodies; marine and
 botanical lipids (fish and plant seed fatty acids; see e.g., DeLuca *et al.* (1995) *Rheum. Dis. Clin.
 North Am.* 21:759-777); auranofin; phenylbutazone; meclofenamic acid; flufenamic acid;
 15 intravenous immune globulin; zileuton; azaribine; mycophenolic acid (RS-61443); tacrolimus
 (FK-506); sirolimus (rapamycin); amiprilose (therafectin); cladribine (2-chlorodeoxyadenosine);
 methotrexate; bcl-2 inhibitors (see Bruncko, Milan *et al.*, *Journal of Medicinal Chemistry* (2007),
 50(4), 641-662); antivirals and immune modulating agents.

In one embodiment, the binding protein or antigen-binding portion thereof, is
 20 administered in combination with one of the following agents for the treatment of rheumatoid
 arthritis: small molecule inhibitor of KDR, small molecule inhibitor of Tie-2; methotrexate;
 prednisone; celecoxib; folic acid; hydroxychloroquine sulfate; rofecoxib; etanercept; infliximab;
 leflunomide; naproxen; valdecoxib; sulfasalazine; methylprednisolone; ibuprofen; meloxicam;
 methylprednisolone acetate; gold sodium thiomalate; aspirin; azathioprine; triamcinolone
 25 acetate; propoxyphene napsylate/apap; folate; nabumetone; diclofenac; piroxicam; etodolac;
 diclofenac sodium; oxaprozin; oxycodone hcl; hydrocodone bitartrate/apap; diclofenac
 sodium/misoprostol; fentanyl; anakinra, human recombinant; tramadol hcl; salsalate; sulindac;
 cyanocobalamin/fa/pyridoxine; acetaminophen; alendronate sodium; prednisolone; morphine
 sulfate; lidocaine hydrochloride; indomethacin; glucosamine sulfate/chondroitin; cyclosporine;
 30 amitriptyline hcl; sulfadiazine; oxycodone hcl/acetaminophen; olopatadine hcl; misoprostol;
 naproxen sodium; omeprazole; mycophenolate mofetil; cyclophosphamide; rituximab; IL-1
 TRAP; MRA; CTLA4-IG; IL-18 BP; IL-12/23; anti-IL 18; anti-IL 15; BIRB-796; SCIO-469; VX-
 702; AMG-548; VX-740; Roflumilast; IC-485; CDC-801; and mesopram.

Non-limiting examples of therapeutic agents for inflammatory bowel disease with which a binding protein provided herein can be combined include the following: budenoside; epidermal growth factor; corticosteroids; cyclosporin, sulfasalazine; aminosalicylates; 6-mercaptopurine; azathioprine; metronidazole; lipoxigenase inhibitors; mesalamine; olsalazine; balsalazide; 5 antioxidants; thromboxane inhibitors; IL-1 receptor antagonists; anti-IL-1 β mAbs; anti-IL-6 mAbs; growth factors; elastase inhibitors; pyridinyl-imidazole compounds; antibodies to or antagonists of other human cytokines or growth factors, for example, TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-15, IL-16, IL-17, IL-18, EMAP-II, GM-CSF, FGF, and PDGF. Binding proteins provided herein, or antigen binding portions thereof, can be combined with antibodies to cell 10 surface molecules such as CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD90 or their ligands. The binding proteins provided herein, or antigen binding portions thereof, may also be combined with agents, such as methotrexate, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, 15 complement inhibitors, adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNF α or IL-1 (e.g., IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TNF α converting enzyme inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine 20 receptors and derivatives thereof (e.g., soluble p55 or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-6R) and antiinflammatory cytokines (e.g., IL-4, IL-10, IL-11, IL-13 and TGF β) and bcl-2 inhibitors.

Examples of therapeutic agents for Crohn's disease in which a binding protein can be combined include the following: TNF antagonists, for example, anti-TNF antibodies, 25 Adalimumab (PCT Publication No. WO 97/29131; HUMIRA), CA2 (REMICADE), CDP 571, TNFR-Ig constructs, (p75TNFRIgG (ENBREL) and p55TNFRIgG (LENERCEPT)) inhibitors and PDE4 inhibitors. Binding proteins provided herein, or antigen binding portions thereof, can be combined with corticosteroids, for example, budenoside and dexamethasone. Binding proteins provided herein or antigen binding portions thereof, may also be combined with agents such as 30 sulfasalazine, 5-aminosalicylic acid and olsalazine, and agents which interfere with synthesis or action of proinflammatory cytokines such as IL-1, for example, IL-1 β converting enzyme inhibitors and IL-1ra. Binding proteins provided herein or antigen binding portion thereof may also be used with T cell signaling inhibitors, for example, tyrosine kinase inhibitors 6-mercaptopurines. Binding proteins provided herein, or antigen binding portions thereof, can be 35 combined with IL-11. Binding proteins provided herein, or antigen binding portions thereof, can

be combined with mesalamine, prednisone, azathioprine, mercaptopurine, infliximab, methylprednisolone sodium succinate, diphenoxylate/atrop sulfate, loperamide hydrochloride, methotrexate, omeprazole, folate, ciprofloxacin/dextrose-water, hydrocodone bitartrate/apap, tetracycline hydrochloride, fluocinonide, metronidazole, thimerosal/boric acid,
 5 cholestyramine/sucrose, ciprofloxacin hydrochloride, hyoscyamine sulfate, meperidine hydrochloride, midazolam hydrochloride, oxycodone hcl/acetaminophen, promethazine hydrochloride, sodium phosphate, sulfamethoxazole/trimethoprim, celecoxib, polycarbophil, propoxyphene napsylate, hydrocortisone, multivitamins, balsalazide disodium, codeine phosphat/apap, colesevelam hcl, cyanocobalamin, folic acid, levofloxacin, methylprednisolone,
 10 natalizumab and interferon-gamma

Non-limiting examples of therapeutic agents for multiple sclerosis with which binding proteins provided herein can be combined include the following: corticosteroids; prednisolone; methylprednisolone; azathioprine; cyclophosphamide; cyclosporine; methotrexate; 4-aminopyridine; tizanidine; interferon- β 1a (AVONEX; Biogen); interferon- β 1b (BETASERON; Chiron/Berlex); interferon α -n3) (Interferon Sciences/Fujimoto), interferon- α (Alfa Wassermann/J&J), interferon β 1A-IF (Serono/Inhale Therapeutics), Peginterferon α 2b (Enzon/Schering-Plough), Copolymer 1 (Cop-1; COPAXONE; Teva Pharmaceutical Industries, Inc.); hyperbaric oxygen; intravenous immunoglobulin; clabribine; antibodies to or antagonists of other human cytokines or growth factors and their receptors, for example, TNF, LT, IL-1, IL-2,
 15 IL-6, IL-7, IL-8, IL-23, IL-15, IL-16, IL-18, EMAP-II, GM-CSF, FGF, and PDGF. Binding proteins provided herein can be combined with antibodies to cell surface molecules such as CD2, CD3, CD4, CD8, CD19, CD20, CD25, CD28, CD30, CD40, CD45, CD69, CD80, CD86, CD90 or their ligands. Binding proteins provided herein, may also be combined with agents, such as methotrexate, cyclosporine, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs,
 20 for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNF α or IL-1 (e.g., IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TACE inhibitors, T-cell signaling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine,
 25 azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (e.g., soluble p55 or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-6R), antiinflammatory cytokines (e.g., IL-4, IL-10, IL-13 and TGF β) and bcl-2 inhibitors.

Examples of therapeutic agents for multiple sclerosis in which binding proteins provided herein can be combined include interferon- β , for example, IFN β 1a and IFN β 1b; copaxone,

corticosteroids, caspase inhibitors, for example inhibitors of caspase-1, IL-1 inhibitors, TNF inhibitors, and antibodies to CD40 ligand and CD80.

The binding proteins provided herein, may also be combined with agents, such as alemtuzumab, dronabinol, Unimed, daclizumab, mitoxantrone, xaliproden hydrochloride, fampridine, glatiramer acetate, natalizumab, sinnabidol, a-immunokine NNSO3, ABR-215062, AnergiX.MS, chemokine receptor antagonists, BBR-2778, calagualine, CPI-1189, LEM (liposome encapsulated mitoxantrone), THC.CBD (cannabinoid agonist) MBP-8298, mesopram (PDE4 inhibitor), MNA-715, anti-IL-6 receptor antibody, neurovax, pirfenidone allotrap 1258 (RDP-1258), sTNF-R1, talampanel, teriflunomide, TGF-beta2, tiplimotide, VLA-4 antagonists (for example, TR-14035, VLA4 Ultrahaler, Antegran-ELAN/Biogen), interferon gamma antagonists, IL-4 agonists.

Non-limiting examples of therapeutic agents for Angina with which binding proteins provided herein can be combined include the following: aspirin, nitroglycerin, isosorbide mononitrate, metoprolol succinate, atenolol, metoprolol tartrate, amlodipine besylate, diltiazem hydrochloride, isosorbide dinitrate, clopidogrel bisulfate, nifedipine, atorvastatin calcium, potassium chloride, furosemide, simvastatin, verapamil hcl, digoxin, propranolol hydrochloride, carvedilol, lisinopril, spironolactone, hydrochlorothiazide, enalapril maleate, nadolol, ramipril, enoxaparin sodium, heparin sodium, valsartan, sotalol hydrochloride, fenofibrate, ezetimibe, bumetanide, losartan potassium, lisinopril/hydrochlorothiazide, felodipine, captopril, bisoprolol fumarate.

Non-limiting examples of therapeutic agents for Ankylosing Spondylitis with which binding proteins provided herein can be combined include the following: ibuprofen, diclofenac and misoprostol, naproxen, meloxicam, indomethacin, diclofenac, celecoxib, rofecoxib, Sulfasalazine, Methotrexate, azathioprine, minocyclin, prednisone, etanercept, infliximab.

Non-limiting examples of therapeutic agents for Asthma with which binding proteins provided herein can be combined include the following: albuterol, salmeterol/fluticasone, montelukast sodium, fluticasone propionate, budesonide, prednisone, salmeterol xinafoate, levalbuterol hcl, albuterol sulfate/ipratropium, prednisolone sodium phosphate, triamcinolone acetonide, beclomethasone dipropionate, ipratropium bromide, azithromycin, pirbuterol acetate, prednisolone, theophylline anhydrous, methylprednisolone sodium succinate, clarithromycin, zafirlukast, formoterol fumarate, influenza virus vaccine, methylprednisolone, amoxicillin trihydrate, flunisolide, allergy injection, cromolyn sodium, fexofenadine hydrochloride, flunisolide/menthol, amoxicillin/clavulanate, levofloxacin, inhaler assist device, guaifenesin,

dexamethasone sodium phosphate, moxifloxacin hcl, doxycycline hyclate, guaifenesin/d-methorphan, p-ephedrine/cod/chlorphenir, gatifloxacin, cetirizine hydrochloride, mometasone furoate, salmeterol xinafoate, benzonatate, cephalixin, pe/hydrocodone/chlorphenir, cetirizine hcl/pseudoephed, phenylephrine/cod/promethazine, codeine/promethazine, cefprozil,
 5 dexamethasone, guaifenesin/pseudoephedrine, chlorpheniramine/hydrocodone, nedocromil sodium, terbutaline sulfate, epinephrine, methylprednisolone, metaproterenol sulfate.

Non-limiting examples of therapeutic agents for COPD with which binding proteins provided herein can be combined include the following: albuterol sulfate/ipratropium, ipratropium bromide, salmeterol/fluticasone, albuterol, salmeterol xinafoate, fluticasone
 10 propionate, prednisone, theophylline anhydrous, methylprednisolone sodium succinate, montelukast sodium, budesonide, formoterol fumarate, triamcinolone acetonide, levofloxacin, guaifenesin, azithromycin, beclomethasone dipropionate, levalbuterol hcl, flunisolide, ceftriaxone sodium, amoxicillin trihydrate, gatifloxacin, zafirlukast, amoxicillin/clavulanate, flunisolide/menthol, chlorpheniramine/hydrocodone, metaproterenol sulfate, methylprednisolone,
 15 mometasone furoate, p-ephedrine/cod/chlorphenir, pirbuterol acetate, p-ephedrine/loratadine, terbutaline sulfate, tiotropium bromide, (R,R)-formoterol, TgAAT, Cilomilast, Roflumilast.

Non-limiting examples of therapeutic agents for HCV with which binding proteins provided herein can be combined include the following: Interferon-alpha-2a, Interferon-alpha-2b, Interferon-alpha con1, Interferon-alpha-n1, Pegylated interferon-alpha-2a, Pegylated interferon-alpha-2b,
 20 alpha-2b, ribavirin, Peginterferon alfa-2b + ribavirin, Ursodeoxycholic Acid, Glycyrrhizic Acid, Thymalfasin, Maxamine, VX-497 and any compounds that are used to treat HCV through intervention with the following targets: HCV polymerase, HCV protease, HCV helicase, HCV IRES (internal ribosome entry site).

Non-limiting examples of therapeutic agents for Idiopathic Pulmonary Fibrosis with
 25 which binding proteins provided herein can be combined include the following: prednisone, azathioprine, albuterol, colchicine, albuterol sulfate, digoxin, gamma interferon, methylprednisolone sod succ, lorazepam, furosemide, lisinopril, nitroglycerin, spironolactone, cyclophosphamide, ipratropium bromide, actinomycin d, alteplase, fluticasone propionate, levofloxacin, metaproterenol sulfate, morphine sulfate, oxycodone hcl, potassium chloride,
 30 triamcinolone acetonide, tacrolimus anhydrous, calcium, interferon-alpha, methotrexate, mycophenolate mofetil, Interferon-gamma-1 β .

Non-limiting examples of therapeutic agents for Myocardial Infarction with which binding proteins provided herein can be combined include the following: aspirin, nitroglycerin,

metoprolol tartrate, enoxaparin sodium, heparin sodium, clopidogrel bisulfate, carvedilol, atenolol, morphine sulfate, metoprolol succinate, warfarin sodium, lisinopril, isosorbide mononitrate, digoxin, furosemide, simvastatin, ramipril, tenecteplase, enalapril maleate, torsemide, retavase, losartan potassium, quinapril hcl/mag carb, bumetanide, alteplase, 5 enalaprilat, amiodarone hydrochloride, tirofiban hcl m-hydrate, diltiazem hydrochloride, captopril, irbesartan, valsartan, propranolol hydrochloride, fosinopril sodium, lidocaine hydrochloride, eptifibatide, cefazolin sodium, atropine sulfate, aminocaproic acid, spironolactone, interferon, sotalol hydrochloride, potassium chloride, docusate sodium, dobutamine hcl, alprazolam, pravastatin sodium, atorvastatin calcium, midazolam hydrochloride, 10 meperidine hydrochloride, isosorbide dinitrate, epinephrine, dopamine hydrochloride, bivalirudin, rosuvastatin, ezetimibe/simvastatin, avasimibe, cariporide.

Non-limiting examples of therapeutic agents for Psoriasis with which binding proteins provided herein can be combined include the following: small molecule inhibitor of KDR, small molecule inhibitor of Tie-2, calcipotriene, clobetasol propionate, triamcinolone acetonide, 15 halobetasol propionate, tazarotene, methotrexate, fluocinonide, betamethasone diprop augmented, fluocinolone acetonide, acitretin, tar shampoo, betamethasone valerate, mometasone furoate, ketoconazole, pramoxine/fluocinolone, hydrocortisone valerate, flurandrenolide, urea, betamethasone, clobetasol propionate/emoll, fluticasone propionate, azithromycin, hydrocortisone, moisturizing formula, folic acid, desonide, pimecrolimus, coal tar, diflorasone 20 diacetate, etanercept folate, lactic acid, methoxsalen, hc/bismuth subgal/znox/resor, methylprednisolone acetate, prednisone, sunscreen, halcinonide, salicylic acid, anthralin, clo cortolone pivalate, coal extract, coal tar/salicylic acid, coal tar/salicylic acid/sulfur, desoximetasone, diazepam, emollient, fluocinonide/emollient, mineral oil/castor oil/na lact, mineral oil/peanut oil, petroleum/isopropyl myristate, psoralen, salicylic acid, soap/tribromsalan, 25 thimerosal/boric acid, celecoxib, infliximab, cyclosporine, alefacept, efalizumab, tacrolimus, pimecrolimus, PUVA, UVB, sulfasalazine.

Non-limiting examples of therapeutic agents for Psoriatic Arthritis with which binding proteins provided herein can be combined include the following: methotrexate, etanercept, rofecoxib, celecoxib, folic acid, sulfasalazine, naproxen, leflunomide, methylprednisolone 30 acetate, indomethacin, hydroxychloroquine sulfate, prednisone, sulindac, betamethasone diprop augmented, infliximab, methotrexate, folate, triamcinolone acetonide, diclofenac, dimethylsulfoxide, piroxicam, diclofenac sodium, ketoprofen, meloxicam, methylprednisolone, nabumetone, tolmetin sodium, calcipotriene, cyclosporine, diclofenac sodium/misoprostol, fluocinonide, glucosamine sulfate, gold sodium thiomalate, hydrocodone bitartrate/apap,

ibuprofen, risedronate sodium, sulfadiazine, thioguanine, valdecoxib, alefacept, efalizumab and bcl-2 inhibitors.

Non-limiting examples of therapeutic agents for Restenosis with which binding proteins provided herein can be combined include the following: sirolimus, paclitaxel, everolimus,
5 tacrolimus, Zotarolimus, acetaminophen.

Non-limiting examples of therapeutic agents for Sciatica with which binding proteins provided herein can be combined include the following: hydrocodone bitartrate/apap, rofecoxib, cyclobenzaprine hcl, methylprednisolone, naproxen, ibuprofen, oxycodone hcl/acetaminophen, celecoxib, valdecoxib, methylprednisolone acetate, prednisone, codeine phosphate/apap,
10 tramadol hcl/acetaminophen, metaxalone, meloxicam, methocarbamol, lidocaine hydrochloride, diclofenac sodium, gabapentin, dexamethasone, carisoprodol, ketorolac tromethamine, indomethacin, acetaminophen, diazepam, nabumetone, oxycodone hcl, tizanidine hcl, diclofenac sodium/misoprostol, propoxyphene napsylate/apap, asa/oxycod/oxycodone ter,
15 ibuprofen/hydrocodone bit, tramadol hcl, etodolac, propoxyphene hcl, amitriptyline hcl, carisoprodol/codeine phos/asa, morphine sulfate, multivitamins, naproxen sodium, orphenadrine citrate, temazepam.

Examples of therapeutic agents for SLE (Lupus) in which binding proteins provided herein can be combined include the following: NSAIDS, for example, diclofenac, naproxen, ibuprofen, piroxicam, indomethacin; COX2 inhibitors, for example, Celecoxib, rofecoxib,
20 valdecoxib; anti-malarials, for example, hydroxychloroquine; Steroids, for example, prednisone, prednisolone, budenoside, dexamethasone; Cytotoxics, for example, azathioprine, cyclophosphamide, mycophenolate mofetil, methotrexate; inhibitors of PDE4 or purine synthesis inhibitor, for example Cellcept. Binding proteins provided herein, may also be combined with agents such as sulfasalazine, 5-aminosalicylic acid, olsalazine, Imuran and agents which interfere
25 with synthesis, production or action of proinflammatory cytokines such as IL-1, for example, caspase inhibitors like IL-1 β converting enzyme inhibitors and IL-1ra. Binding proteins provided herein may also be used with T cell signaling inhibitors, for example, tyrosine kinase inhibitors; or molecules that target T cell activation molecules, for example, CTLA-4-IgG or anti-B7 family antibodies, anti-PD-1 family antibodies. Binding proteins provided herein, can be combined with
30 IL-11 or anti-cytokine antibodies, for example, fonotolizumab (anti-IFN γ antibody), or anti-receptor receptor antibodies, for example, anti-IL-6 receptor antibody and antibodies to B-cell surface molecules. Binding proteins provided herein or antigen binding portion thereof may also be used with LJP 394 (abetimus), agents that deplete or inactivate B-cells, for example, Rituximab (anti-CD20 antibody), lymphotoxin-B (anti-BlyS antibody), TNF antagonists, for

example, anti-TNF antibodies, Adalimumab (PCT Publication No. WO 97/29131; HUMIRA), CA2 (REMICADE), CDP 571, TNFR-Ig constructs, (p75TNFRIG (ENBREL) and p55TNFRIG (LENERCEPT)) and bcl-2 inhibitors, because bcl-2 overexpression in transgenic mice has been demonstrated to cause a lupus like phenotype (see Marquina, Regina et al., Journal of Immunology (2004), 172(11), 7177-7185), therefore inhibition is expected to have therapeutic effects.

The pharmaceutical compositions provided herein may include a "therapeutically effective amount" or a "prophylactically effective amount" of a binding protein provided herein. A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the binding protein may be determined by a person skilled in the art and may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the binding protein to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the binding protein or antigen-binding portion, are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

Dosage regimens may be adjusted to provide the optimum desired response (*e.g.*, a therapeutic or prophylactic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms provided herein are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

An exemplary, non-limiting range for a therapeutically or prophylactically effective amount of a binding protein provided herein is 0.1-20 mg/kg, for example, 1-10 mg/kg. It is to be

noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

V. Diagnostics

The disclosure herein also provides diagnostic applications. This is further elucidated below.

10 A. Method of Assay

The present disclosure also provides a method for determining the presence, amount or concentration of an analyte (or a fragment thereof) in a test sample using at least one DVD-Ig as described herein. Any suitable assay as is known in the art can be used in the method. Examples include, but are not limited to, immunoassay, such as sandwich immunoassay (e.g., monoclonal, polyclonal and/or DVD-Ig sandwich immunoassays or any variation thereof (e.g., monoclonal/DVD-Ig, DVD-Ig/polyclonal, etc.), including radioisotope detection (radioimmunoassay (RIA)) and enzyme detection (enzyme immunoassay (EIA) or enzyme-linked immunosorbent assay (ELISA) (e.g., Quantikine ELISA assays, R&D Systems, Minneapolis, MN)), competitive inhibition immunoassay (e.g., forward and reverse), fluorescence polarization immunoassay (FPIA), enzyme multiplied immunoassay technique (EMIT), bioluminescence resonance energy transfer (BRET), and homogeneous chemiluminescent assay, etc. In a SELDI-based immunoassay, a capture reagent that specifically binds an analyte (or a fragment thereof) of interest is attached to the surface of a mass spectrometry probe, such as a pre-activated protein chip array. The analyte (or a fragment thereof) is then specifically captured on the biochip, and the captured analyte (or a fragment thereof) is detected by mass spectrometry. Alternatively, the analyte (or a fragment thereof) can be eluted from the capture reagent and detected by traditional MALDI (matrix-assisted laser desorption/ionization) or by SELDI. A chemiluminescent microparticle immunoassay, in particular one employing the ARCHITECT® automated analyzer (Abbott Laboratories, Abbott Park, IL), is an example of a preferred immunoassay.

30 Methods well-known in the art for collecting, handling and processing urine, blood, serum and plasma, and other body fluids, are used in the practice of the present disclosure, for instance, when a DVD-Ig as described herein is employed as an immunodiagnostic reagent and/or in an analyte immunoassay kit. The test sample can comprise further moieties in addition to the

analyte of interest, such as antibodies, antigens, haptens, hormones, drugs, enzymes, receptors, proteins, peptides, polypeptides, oligonucleotides and/or polynucleotides. For example, the sample can be a whole blood sample obtained from a subject. It can be necessary or desired that a test sample, particularly whole blood, be treated prior to immunoassay as described herein, e.g., with a pretreatment reagent. Even in cases where pretreatment is not necessary (e.g., most urine samples), pretreatment optionally can be done (e.g., as part of a regimen on a commercial platform).

The pretreatment reagent can be any reagent appropriate for use with the immunoassay and kits provided herein. The pretreatment optionally comprises: (a) one or more solvents (e.g., methanol and ethylene glycol) and optionally, salt, (b) one or more solvents and salt, and optionally, detergent, (c) detergent, or (d) detergent and salt. Pretreatment reagents are known in the art, and such pretreatment can be employed, e.g., as used for assays on Abbott TDx, AxSYM®, and ARCHITECT® analyzers (Abbott Laboratories, Abbott Park, IL), as described in the literature (see, e.g., Yatscoff et al., *Abbott TDx Monoclonal Antibody Assay Evaluated for Measuring Cyclosporine in Whole Blood*, *Clin. Chem.* 36: 1969-1973 (1990), and Wallemacq et al., *Evaluation of the New AxSYM Cyclosporine Assay: Comparison with TDx Monoclonal Whole Blood and EMIT Cyclosporine Assays*, *Clin. Chem.* 45: 432-435 (1999)), and/or as commercially available. Additionally, pretreatment can be done as described in Abbott's U.S. Pat. No. 5,135,875, European Pat. Pub. No. 0 471 293, U.S. Provisional Pat. App. 60/878,017, filed December 29, 2006, and U.S. Pat. App. Pub. No. 2008/0020401 (incorporated by reference in its entirety for its teachings regarding pretreatment). The pretreatment reagent can be a heterogeneous agent or a homogeneous agent.

With use of a heterogeneous pretreatment reagent, the pretreatment reagent precipitates analyte binding protein (e.g., protein that can bind to an analyte or a fragment thereof) present in the sample. Such a pretreatment step comprises removing any analyte binding protein by separating from the precipitated analyte binding protein the supernatant of the mixture formed by addition of the pretreatment agent to sample. In such an assay, the supernatant of the mixture absent any binding protein is used in the assay, proceeding directly to the capture step.

With use of a homogeneous pretreatment reagent there is no such separation step. The entire mixture of test sample and pretreatment reagent are contacted with a labeled specific binding partner for analyte (or a fragment thereof), such as a labeled anti-analyte antibody (or an antigenically reactive fragment thereof). The pretreatment reagent employed for such an assay typically is diluted in the pretreated test sample mixture, either before or during capture by the first specific binding partner. Despite such dilution, a certain amount of the pretreatment reagent

is still present (or remains) in the test sample mixture during capture. According to certain embodiments, the labeled specific binding partner can be a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof).

In a heterogeneous format, after the test sample is obtained from a subject, a first mixture
5 is prepared. The mixture contains the test sample being assessed for an analyte (or a fragment thereof) and a first specific binding partner, wherein the first specific binding partner and any analyte contained in the test sample form a first specific binding partner-analyte complex. Preferably, the first specific binding partner is an anti-analyte antibody or a fragment thereof. The first specific binding partner can be a DVD-Ig (or a fragment, a variant, or a fragment of a variant
10 thereof) as described herein. The order in which the test sample and the first specific binding partner are added to form the mixture is not critical. Preferably, the first specific binding partner is immobilized on a solid phase. The solid phase used in the immunoassay (for the first specific binding partner and, optionally, the second specific binding partner) can be any solid phase known in the art, such as, but not limited to, a magnetic particle, a bead, a test tube, a microtiter
15 plate, a cuvette, a membrane, a scaffolding molecule, a film, a filter paper, a disc and a chip.

After the mixture containing the first specific binding partner-analyte complex is formed, any unbound analyte is removed from the complex using any technique known in the art. For example, the unbound analyte can be removed by washing. Desirably, however, the first specific binding partner is present in excess of any analyte present in the test sample, such that all analyte
20 that is present in the test sample is bound by the first specific binding partner.

After any unbound analyte is removed, a second specific binding partner is added to the mixture to form a first specific binding partner-analyte-second specific binding partner complex. The second specific binding partner is preferably an anti-analyte antibody that binds to an epitope on analyte that differs from the epitope on analyte bound by the first specific binding partner.
25 Moreover, also preferably, the second specific binding partner is labeled with or contains a detectable label as described above. The second specific binding partner can be a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) as described herein.

Any suitable detectable label as is known in the art can be used. For example, the detectable label can be a radioactive label (such as ^3H , ^{125}I , ^{35}S , ^{14}C , ^{32}P , and ^{33}P), an
30 enzymatic label (such as horseradish peroxidase, alkaline peroxidase, glucose 6-phosphate dehydrogenase, and the like), a chemiluminescent label (such as acridinium esters, thioesters, or sulfonamides; luminol, isoluminol, phenanthridinium esters, and the like), a fluorescent label (such as fluorescein (e.g., 5-fluorescein, 6-carboxyfluorescein, 3'-6-carboxyfluorescein, 5(6)-

carboxyfluorescein, 6-hexachloro-fluorescein, 6-tetrachlorofluorescein, fluorescein isothiocyanate, and the like)), rhodamine, phycobiliproteins, R-phycoerythrin, quantum dots (e.g., zinc sulfide-capped cadmium selenide), a thermometric label, or an immuno-polymerase chain reaction label. An introduction to labels, labeling procedures and detection of labels is found in Polak and Van Noorden, *Introduction to Immunocytochemistry*, 2nd ed., Springer Verlag, N.Y. (1997), and in Haugland, *Handbook of Fluorescent Probes and Research Chemicals* (1996), which is a combined handbook and catalogue published by Molecular Probes, Inc., Eugene, Oregon. A fluorescent label can be used in FPIA (see, e.g., U.S. Patent Nos. 5,593,896, 5,573,904, 5,496,925, 5,359,093, and 5,352,803). An acridinium compound can be used as a detectable label in a homogeneous or heterogeneous chemiluminescent assay (see, e.g., Adamczyk et al., *Bioorg. Med. Chem. Lett.* 16: 1324-1328 (2006); Adamczyk et al., *Bioorg. Med. Chem. Lett.* 4: 2313-2317 (2004); Adamczyk et al., *Bioorg. Med. Chem. Lett.* 14: 3917-3921 (2004); and Adamczyk et al., *Org. Lett.* 5: 3779-3782 (2003)).

A preferred acridinium compound is an acridinium-9-carboxamide. Methods for preparing acridinium 9-carboxamides are described in Mattingly, *J. Biolumin. Chemilumin.* 6: 107-114 (1991); Adamczyk et al., *J. Org. Chem.* 63: 5636-5639 (1998); Adamczyk et al., *Tetrahedron* 55: 10899-10914 (1999); Adamczyk et al., *Org. Lett.* 1: 779-781 (1999); Adamczyk et al., *Bioconjugate Chem.* 11: 714-724 (2000); Mattingly et al., *In Luminescence Biotechnology: Instruments and Applications*; Dyke, K. V. Ed.; CRC Press: Boca Raton, pp. 77-105 (2002); Adamczyk et al., *Org. Lett.* 5: 3779-3782 (2003); and U.S. Pat. Nos. 5,468,646, 5,543,524 and 5,783,699. Another preferred acridinium compound is an acridinium-9-carboxylate aryl ester. An example of an acridinium-9-carboxylate aryl ester is 10-methyl-9-(phenoxycarbonyl)acridinium fluorosulfonate (available from Cayman Chemical, Ann Arbor, MI). Methods for preparing acridinium 9-carboxylate aryl esters are described in McCapra et al., *Photochem. Photobiol.* 4: 1111-21 (1965); Razavi et al., *Luminescence* 15: 245-249 (2000); Razavi et al., *Luminescence* 15: 239-244 (2000); and U.S. Patent No. 5,241,070. Further details regarding acridinium-9-carboxylate aryl ester and its use are set forth in US 2008-0248493.

Chemiluminescent assays (e.g., using acridinium as described above or other chemiluminescent agents) can be performed in accordance with the methods described in Adamczyk et al., *Anal. Chim. Acta* 579(1): 61-67 (2006). While any suitable assay format can be used, a microplate chemiluminometer (Mithras LB-940, Berthold Technologies U.S.A., LLC, Oak Ridge, TN) enables the assay of multiple samples of small volumes rapidly.

The order in which the test sample and the specific binding partner(s) are added to form the mixture for chemiluminescent assay is not critical. If the first specific binding partner is

detectably labeled with a chemiluminescent agent such as an acridinium compound, detectably
labeled first specific binding partner-analyte complexes form. Alternatively, if a second specific
binding partner is used and the second specific binding partner is detectably labeled with a
chemiluminescent agent such as an acridinium compound, detectably labeled first specific
5 binding partner-analyte-second specific binding partner complexes form. Any unbound specific
binding partner, whether labeled or unlabeled, can be removed from the mixture using any
technique known in the art, such as washing.

Hydrogen peroxide can be generated in situ in the mixture or provided or supplied to the
mixture (e.g., the source of the hydrogen peroxide being one or more buffers or other solutions
10 that are known to contain hydrogen peroxide) before, simultaneously with, or after the addition of
an above-described acridinium compound. Hydrogen peroxide can be generated in situ in a
number of ways such as would be apparent to one skilled in the art.

Upon the simultaneous or subsequent addition of at least one basic solution to the
sample, a detectable signal, namely, a chemiluminescent signal, indicative of the presence of
15 analyte is generated. The basic solution contains at least one base and has a pH greater than or
equal to 10, preferably, greater than or equal to 12. Examples of basic solutions include, but are
not limited to, sodium hydroxide, potassium hydroxide, calcium hydroxide, ammonium
hydroxide, magnesium hydroxide, sodium carbonate, sodium bicarbonate, calcium hydroxide,
calcium carbonate, and calcium bicarbonate. The amount of basic solution added to the sample
20 depends on the concentration of the basic solution. Based on the concentration of the basic
solution used, one skilled in the art can easily determine the amount of basic solution to add to
the sample.

The chemiluminescent signal that is generated can be detected using routine techniques
known to those skilled in the art. Based on the intensity of the signal generated, the amount of
25 analyte in the sample can be quantified. Specifically, the amount of analyte in the sample is
proportional to the intensity of the signal generated. The amount of analyte present can be
quantified by comparing the amount of light generated to a standard curve for analyte or by
comparison to a reference standard. The standard curve can be generated using serial dilutions or
solutions of known concentrations of analyte by mass spectroscopy, gravimetric methods, and
30 other techniques known in the art. While the above is described with emphasis on use of an
acridinium compound as the chemiluminescent agent, one of ordinary skill in the art can readily
adapt this description for use of other chemiluminescent agents.

Analyte immunoassays generally can be conducted using any format known in the art, such as, but not limited to, a sandwich format. Specifically, in one immunoassay format, at least two binding proteins are employed to separate and quantify analyte, such as human analyte, or a fragment thereof in a sample. More specifically, the at least two binding proteins bind to different epitopes on an analyte (or a fragment thereof) forming an immune complex, which is referred to as a "sandwich." Generally, in the immunoassays one or more binding proteins can be used to capture the analyte (or a fragment thereof) in the test sample (these binding proteins are frequently referred to as a "capture" antibody or "capture" antibodies) and one or more binding proteins can be used to bind a detectable (namely, quantifiable) label to the sandwich (these binding proteins are frequently referred to as the "detection antibody," the "detection antibodies," the "conjugate," or the "conjugates"). Thus, in the context of a sandwich immunoassay format, a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) as described herein can be used as a capture antibody, a detection antibody, or both. For example, one DVD-Ig having a domain that can bind a first epitope on an analyte (or a fragment thereof) can be used as a capture antibody and/or another DVD-Ig having a domain that can bind a second epitope on an analyte (or a fragment thereof) can be used as a detection antibody. In this regard, a DVD-Ig having a first domain that can bind a first epitope on an analyte (or a fragment thereof) and a second domain that can bind a second epitope on an analyte (or a fragment thereof) can be used as a capture antibody and/or a detection antibody. Alternatively, one DVD-Ig having a first domain that can bind an epitope on a first analyte (or a fragment thereof) and a second domain that can bind an epitope on a second analyte (or a fragment thereof) can be used as a capture antibody and/or a detection antibody to detect, and optionally quantify, two or more analytes. In the event that an analyte can be present in a sample in more than one form, such as a monomeric form and a dimeric/multimeric form, which can be homomeric or heteromeric, one DVD-Ig having a domain that can bind an epitope that is only exposed on the monomeric form and another DVD-Ig having a domain that can bind an epitope on a different part of a dimeric/multimeric form can be used as capture antibodies and/or detection antibodies, thereby enabling the detection, and optional quantification, of different forms of a given analyte. Furthermore, employing DVD-Igs with differential affinities within a single DVD-Ig and/or between DVD-Igs can provide an avidity advantage. In the context of immunoassays as described herein, it generally may be helpful or desired to incorporate one or more linkers within the structure of a DVD-Ig. When present, optimally the linker should be of sufficient length and structural flexibility to enable binding of an epitope by the inner domains as well as binding of another epitope by the outer domains. In this regard, if a DVD-Ig can bind two different analytes and one analyte is larger than the other, desirably the larger analyte is bound by the outer domains.

Generally speaking, a sample being tested for (for example, suspected of containing) analyte (or a fragment thereof) can be contacted with at least one capture antibody (or antibodies) and at least one detection antibody (which can be a second detection antibody or a third detection antibody or even a successively numbered antibody, e.g., as where the capture and/or detection antibody comprise multiple antibodies) either simultaneously or sequentially and in any order. For example, the test sample can be first contacted with at least one capture antibody and then (sequentially) with at least one detection antibody. Alternatively, the test sample can be first contacted with at least one detection antibody and then (sequentially) with at least one capture antibody. In yet another alternative, the test sample can be contacted simultaneously with a capture antibody and a detection antibody.

In the sandwich assay format, a sample suspected of containing analyte (or a fragment thereof) is first brought into contact with at least one first capture antibody under conditions that allow the formation of a first antibody/analyte complex. If more than one capture antibody is used, a first capture antibody/analyte complex comprising two or more capture antibodies is formed. In a sandwich assay, the antibodies, i.e., preferably, the at least one capture antibody, are used in molar excess amounts of the maximum amount of analyte (or a fragment thereof) expected in the test sample. For example, from about 5 μ g to about 1 mg of antibody per mL of buffer (e.g., microparticle coating buffer) can be used.

Competitive inhibition immunoassays, which are often used to measure small analytes because binding by only one binding protein is required, comprise sequential and classic formats. In a sequential competitive inhibition immunoassay a capture antibody to an analyte of interest is coated onto a well of a microtiter plate or other solid support. When the sample containing the analyte of interest is added to the well, the analyte of interest binds to the capture antibody. After washing, a known amount of labeled (e.g., biotin or horseradish peroxidase (HRP)) analyte is added to the well. A substrate for an enzymatic label is necessary to generate a signal. An example of a suitable substrate for HRP is 3,3',5,5'-tetramethylbenzidine (TMB). After washing, the signal generated by the labeled analyte is measured and is inversely proportional to the amount of analyte in the sample. In a classic competitive inhibition immunoassay an antibody to an analyte of interest is coated onto a solid support (e.g., a well of a microtiter plate). However, unlike the sequential competitive inhibition immunoassay, the sample and the labeled analyte are added to the well at the same time. Any analyte in the sample competes with labeled analyte for binding to the capture antibody. After washing, the signal generated by the labeled analyte is measured and is inversely proportional to the amount of analyte in the sample.

Optionally, prior to contacting the test sample with the at least one capture antibody (for example, the first capture antibody), the at least one capture antibody can be bound to a solid support, which facilitates the separation of the first antibody/analyte (or a fragment thereof) complex from the test sample. The substrate to which the capture antibody is bound can be any suitable solid support or solid phase that facilitates separation of the capture antibody-analyte complex from the sample.

Examples include a well of a plate, such as a microtiter plate, a test tube, a porous gel (e.g., silica gel, agarose, dextran, or gelatin), a polymeric film (e.g., polyacrylamide), beads (e.g., polystyrene beads or magnetic beads), a strip of a filter/membrane (e.g., nitrocellulose or nylon), microparticles (e.g., latex particles, magnetizable microparticles (e.g., microparticles having ferric oxide or chromium oxide cores and homo- or hetero-polymeric coats and radii of about 1-10 microns). The substrate can comprise a suitable porous material with a suitable surface affinity to bind antigens and sufficient porosity to allow access by detection antibodies. A microporous material is generally preferred, although a gelatinous material in a hydrated state can be used. Such porous substrates are preferably in the form of sheets having a thickness of about 0.01 to about 0.5 mm, preferably about 0.1 mm. While the pore size may vary quite a bit, preferably the pore size is from about 0.025 to about 15 microns, more preferably from about 0.15 to about 15 microns. The surface of such substrates can be activated by chemical processes that cause covalent linkage of an antibody to the substrate. Irreversible binding, generally by adsorption through hydrophobic forces, of the antigen or the antibody to the substrate results; alternatively, a chemical coupling agent or other means can be used to bind covalently the antibody to the substrate, provided that such binding does not interfere with the ability of the antibody to bind to analyte. Alternatively, the antibody can be bound with microparticles, which have been previously coated with streptavidin (e.g., DYNAL® Magnetic Beads, Invitrogen, Carlsbad, CA) or biotin (e.g., using Power-Bind™-SA-MP streptavidin-coated microparticles (Seradyn, Indianapolis, IN)) or anti-species-specific monoclonal antibodies. If necessary, the substrate can be derivatized to allow reactivity with various functional groups on the antibody. Such derivatization requires the use of certain coupling agents, examples of which include, but are not limited to, maleic anhydride, N-hydroxysuccinimide, and 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide. If desired, one or more capture reagents, such as antibodies (or fragments thereof), each of which is specific for analyte(s) can be attached to solid phases in different physical or addressable locations (e.g., such as in a biochip configuration (see, e.g., U.S. Pat. No. 6,225,047; Int'l Pat. App. Pub. No. WO 99/51773; U.S. Pat. No. 6,329,209; Int'l Pat. App. Pub. No. WO 00/56934, and U.S. Pat. No. 5,242,828). If the capture reagent is attached to a mass spectrometry probe as the solid support, the amount of analyte bound to the probe can be

detected by laser desorption ionization mass spectrometry. Alternatively, a single column can be packed with different beads, which are derivatized with the one or more capture reagents, thereby capturing the analyte in a single place (see, antibody-derivatized, bead-based technologies, e.g., the xMAP technology of Luminex (Austin, TX)).

5 After the test sample being assayed for analyte (or a fragment thereof) is brought into contact with the at least one capture antibody (for example, the first capture antibody), the mixture is incubated in order to allow for the formation of a first antibody (or multiple antibody)-analyte (or a fragment thereof) complex. The incubation can be carried out at a pH of from about 4.5 to about 10.0, at a temperature of from about 2°C to about 45°C, and for a period from at least
10 about one (1) minute to about eighteen (18) hours, preferably from about 1 to about 24 minutes, most preferably for about 4 to about 18 minutes. The immunoassay described herein can be conducted in one step (meaning the test sample, at least one capture antibody and at least one detection antibody are all added sequentially or simultaneously to a reaction vessel) or in more than one step, such as two steps, three steps, etc.

15 After formation of the (first or multiple) capture antibody/analyte (or a fragment thereof) complex, the complex is then contacted with at least one detection antibody under conditions which allow for the formation of a (first or multiple) capture antibody/analyte (or a fragment thereof)/second detection antibody complex. While captioned for clarity as the "second" antibody (e.g., second detection antibody), in fact, where multiple antibodies are used for capture
20 and/or detection, the at least one detection antibody can be the second, third, fourth, etc. antibodies used in the immunoassay. If the capture antibody/analyte (or a fragment thereof) complex is contacted with more than one detection antibody, then a (first or multiple) capture antibody/analyte (or a fragment thereof)/(multiple) detection antibody complex is formed. As
25 with the capture antibody (e.g., the first capture antibody), when the at least one (e.g., second and any subsequent) detection antibody is brought into contact with the capture antibody/analyte (or a fragment thereof) complex, a period of incubation under conditions similar to those described above is required for the formation of the (first or multiple) capture antibody/analyte (or a fragment thereof)/(second or multiple) detection antibody complex. Preferably, at least one
30 detection antibody contains a detectable label. The detectable label can be bound to the at least one detection antibody (e.g., the second detection antibody) prior to, simultaneously with, or after the formation of the (first or multiple) capture antibody/analyte (or a fragment thereof)/(second or multiple) detection antibody complex. Any detectable label known in the art can be used (see discussion above, including of the Polak and Van Noorden (1997) and Haugland (1996) references).

The detectable label can be bound to the antibodies either directly or through a coupling agent. An example of a coupling agent that can be used is EDAC (1-ethyl-3-(3-dimethylaminopropyl) carbodiimide, hydrochloride), which is commercially available from Sigma-Aldrich, St. Louis, MO. Other coupling agents that can be used are known in the art.

5 Methods for binding a detectable label to an antibody are known in the art. Additionally, many detectable labels can be purchased or synthesized that already contain end groups that facilitate the coupling of the detectable label to the antibody, such as CPSP-Acrininium Ester (i.e., 9-[N-tosyl-N-(3-carboxypropyl)]-10-(3-sulfopropyl)acridinium carboxamide) or SPSP-Acrininium Ester (i.e., N10-(3-sulfopropyl)-N-(3-sulfopropyl)-acridinium-9-carboxamide).

10 The (first or multiple) capture antibody/analyte/(second or multiple) detection antibody complex can be, but does not have to be, separated from the remainder of the test sample prior to quantification of the label. For example, if the at least one capture antibody (e.g., the first capture antibody) is bound to a solid support, such as a well or a bead, separation can be accomplished by removing the fluid (of the test sample) from contact with the solid support. Alternatively, if the at
15 least first capture antibody is bound to a solid support, it can be simultaneously contacted with the analyte-containing sample and the at least one second detection antibody to form a first (multiple) antibody/analyte/second (multiple) antibody complex, followed by removal of the fluid (test sample) from contact with the solid support. If the at least one first capture antibody is not
20 bound to a solid support, then the (first or multiple) capture antibody/analyte/(second or multiple) detection antibody complex does not have to be removed from the test sample for quantification of the amount of the label.

After formation of the labeled capture antibody/analyte/detection antibody complex (e.g., the first capture antibody/analyte/second detection antibody complex), the amount of label in the complex is quantified using techniques known in the art. For example, if an enzymatic label is
25 used, the labeled complex is reacted with a substrate for the label that gives a quantifiable reaction such as the development of color. If the label is a radioactive label, the label is quantified using appropriate means, such as a scintillation counter. If the label is a fluorescent label, the label is quantified by stimulating the label with a light of one color (which is known as the “excitation wavelength”) and detecting another color (which is known as the “emission
30 wavelength”) that is emitted by the label in response to the stimulation. If the label is a chemiluminescent label, the label is quantified by detecting the light emitted either visually or by using luminometers, x-ray film, high speed photographic film, a CCD camera, etc. Once the amount of the label in the complex has been quantified, the concentration of analyte or a fragment thereof in the test sample is determined by appropriate means, such as by use of a

standard curve that has been generated using serial dilutions of analyte or a fragment thereof of known concentration. Other than using serial dilutions of analyte or a fragment thereof, the standard curve can be generated gravimetrically, by mass spectroscopy and by other techniques known in the art.

5 In a chemiluminescent microparticle assay employing the ARCHITECT® analyzer, the conjugate diluent pH should be about 6.0 +/- 0.2, the microparticle coating buffer should be maintained at about room temperature (i.e., at from about 17 to about 27 °C), the microparticle coating buffer pH should be about 6.5 +/- 0.2, and the microparticle diluent pH should be about 7.8 +/- 0.2. Solids preferably are less than about 0.2%, such as less than about 0.15%, less than
10 about 0.14%, less than about 0.13%, less than about 0.12%, or less than about 0.11%, such as about 0.10%.

FPIAs are based on competitive binding immunoassay principles. A fluorescently labeled compound, when excited by a linearly polarized light, will emit fluorescence having a degree of polarization inversely proportional to its rate of rotation. When a fluorescently labeled tracer-
15 antibody complex is excited by a linearly polarized light, the emitted light remains highly polarized because the fluorophore is constrained from rotating between the time light is absorbed and the time light is emitted. When a "free" tracer compound (i.e., a compound that is not bound to an antibody) is excited by linearly polarized light, its rotation is much faster than the corresponding tracer-antibody conjugate produced in a competitive binding immunoassay. FPIAs
20 are advantageous over RIAs inasmuch as there are no radioactive substances requiring special handling and disposal. In addition, FPIAs are homogeneous assays that can be easily and rapidly performed.

In view of the above, a method of determining the presence, amount, or concentration of analyte (or a fragment thereof) in a test sample is provided. The method comprises assaying the
25 test sample for an analyte (or a fragment thereof) by an assay (i) employing (i') at least one of an antibody, a fragment of an antibody that can bind to an analyte, a variant of an antibody that can bind to an analyte, a fragment of a variant of an antibody that can bind to an analyte, and a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) that can bind to an analyte, and (ii') at least one detectable label and (ii) comprising comparing a signal generated by the
30 detectable label as a direct or indirect indication of the presence, amount or concentration of analyte (or a fragment thereof) in the test sample to a signal generated as a direct or indirect indication of the presence, amount or concentration of analyte (or a fragment thereof) in a control or calibrator. The calibrator is optionally part of a series of calibrators, in which each of the calibrators differs from the other calibrators by the concentration of analyte.

The method can comprise (i) contacting the test sample with at least one first specific binding partner for analyte (or a fragment thereof) comprising an antibody, a fragment of an antibody that can bind to an analyte, a variant of an antibody that can bind to an analyte, a fragment of a variant of an antibody that can bind to an analyte, or a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) that can bind to an analyte so as to form a first specific binding partner/analyte (or fragment thereof) complex, (ii) contacting the first specific binding partner/analyte (or fragment thereof) complex with at least one second specific binding partner for analyte (or fragment thereof) comprising a detectably labeled anti-analyte antibody, a detectably labeled fragment of an anti-analyte antibody that can bind to analyte, a detectably labeled variant of an anti-analyte antibody that can bind to analyte, a detectably labeled fragment of a variant of an anti-analyte antibody that can bind to analyte, or a detectably labeled DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) so as to form a first specific binding partner/analyte (or fragment thereof)/second specific binding partner complex, and (iii) determining the presence, amount or concentration of analyte in the test sample by detecting or measuring the signal generated by the detectable label in the first specific binding partner/analyte (or fragment thereof)/second specific binding partner complex formed in (ii). A method in which at least one first specific binding partner for analyte (or a fragment thereof) and/or at least one second specific binding partner for analyte (or a fragment thereof) is a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) as described herein can be preferred.

Alternatively, the method can comprise contacting the test sample with at least one first specific binding partner for analyte (or a fragment thereof) comprising an antibody, a fragment of an antibody that can bind to an analyte, a variant of an antibody that can bind to an analyte, a fragment of a variant of an antibody that can bind to an analyte, or a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) and simultaneously or sequentially, in either order, contacting the test sample with at least one second specific binding partner, which can compete with analyte (or a fragment thereof) for binding to the at least one first specific binding partner and which is a detectably labeled analyte, a detectably labeled fragment of analyte that can bind to the first specific binding partner, a detectably labeled variant of analyte that can bind to the first specific binding partner, or a detectably labeled fragment of a variant of analyte that can bind to the first specific binding partner. Any analyte (or a fragment thereof) present in the test sample and the at least one second specific binding partner compete with each other to form a first specific binding partner/analyte (or fragment thereof) complex and a first specific binding partner/second specific binding partner complex, respectively. The method further comprises determining the presence, amount or concentration of analyte in the test sample by detecting or measuring the signal generated by the detectable label in the first specific binding partner/second

specific binding partner complex formed in (ii), wherein the signal generated by the detectable label in the first specific binding partner/second specific binding partner complex is inversely proportional to the amount or concentration of analyte in the test sample.

5 The above methods can further comprise diagnosing, prognosticating, or assessing the efficacy of a therapeutic/prophylactic treatment of a patient from whom the test sample was obtained. If the method further comprises assessing the efficacy of a therapeutic/prophylactic treatment of the patient from whom the test sample was obtained, the method optionally further comprises modifying the therapeutic/prophylactic treatment of the patient as needed to improve efficacy. The method can be adapted for use in an automated system or a semi-automated system.

10 With regard to the methods of assay (and kit therefor), it may be possible to employ commercially available anti-analyte antibodies or methods for production of anti-analyte as described in the literature. Commercial supplies of various antibodies include, but are not limited to, Santa Cruz Biotechnology Inc. (Santa Cruz, CA), GenWay Biotech, Inc. (San Diego, CA), and R&D Systems (RDS; Minneapolis, MN).

15 Generally, a predetermined level can be employed as a benchmark against which to assess results obtained upon assaying a test sample for analyte or a fragment thereof, e.g., for detecting disease or risk of disease. Generally, in making such a comparison, the predetermined level is obtained by running a particular assay a sufficient number of times and under appropriate conditions such that a linkage or association of analyte presence, amount or concentration with a particular stage or endpoint of a disease, disorder or condition or with particular clinical indicia
20 can be made. Typically, the predetermined level is obtained with assays of reference subjects (or populations of subjects). The analyte measured can include fragments thereof, degradation products thereof, and/or enzymatic cleavage products thereof.

In particular, with respect to a predetermined level as employed for monitoring disease
25 progression and/or treatment, the amount or concentration of analyte or a fragment thereof may be “unchanged,” “favorable” (or “favorably altered”), or “unfavorable” (or “unfavorably altered”). “Elevated” or “increased” refers to an amount or a concentration in a test sample that is higher than a typical or normal level or range (e.g., predetermined level), or is higher than another reference level or range (e.g., earlier or baseline sample). The term “lowered” or “reduced” refers
30 to an amount or a concentration in a test sample that is lower than a typical or normal level or range (e.g., predetermined level), or is lower than another reference level or range (e.g., earlier or baseline sample). The term “altered” refers to an amount or a concentration in a sample that is

altered (increased or decreased) over a typical or normal level or range (e.g., predetermined level), or over another reference level or range (e.g., earlier or baseline sample).

The typical or normal level or range for analyte is defined in accordance with standard practice. Because the levels of analyte in some instances will be very low, a so-called altered level or alteration can be considered to have occurred when there is any net change as compared to the typical or normal level or range, or reference level or range, that cannot be explained by experimental error or sample variation. Thus, the level measured in a particular sample will be compared with the level or range of levels determined in similar samples from a so-called normal subject. In this context, a "normal subject" is an individual with no detectable disease, for example, and a "normal" (sometimes termed "control") patient or population is/are one(s) that exhibit(s) no detectable disease, respectively, for example. Furthermore, given that analyte is not routinely found at a high level in the majority of the human population, a "normal subject" can be considered an individual with no substantial detectable increased or elevated amount or concentration of analyte, and a "normal" (sometimes termed "control") patient or population is/are one(s) that exhibit(s) no substantial detectable increased or elevated amount or concentration of analyte. An "apparently normal subject" is one in which analyte has not yet been or currently is being assessed. The level of an analyte is said to be "elevated" when the analyte is normally undetectable (e.g., the normal level is zero, or within a range of from about 25 to about 75 percentiles of normal populations), but is detected in a test sample, as well as when the analyte is present in the test sample at a higher than normal level. Thus, inter alia, the disclosure provides a method of screening for a subject having, or at risk of having, a particular disease, disorder, or condition. The method of assay can also involve the assay of other markers and the like.

Accordingly, the methods described herein also can be used to determine whether or not a subject has or is at risk of developing a given disease, disorder or condition. Specifically, such a method can comprise the steps of:

(a) determining the concentration or amount in a test sample from a subject of analyte (or a fragment thereof) (e.g., using the methods described herein, or methods known in the art); and

(b) comparing the concentration or amount of analyte (or a fragment thereof) determined in step (a) with a predetermined level, wherein, if the concentration or amount of analyte determined in step (a) is favorable with respect to a predetermined level, then the subject is determined not to have or be at risk for a given disease, disorder or condition. However, if the concentration or amount of analyte determined in step (a) is unfavorable with respect to the

predetermined level, then the subject is determined to have or be at risk for a given disease, disorder or condition.

Additionally, provided herein is method of monitoring the progression of disease in a subject. Optimally the method comprising the steps of:

5 (a) determining the concentration or amount in a test sample from a subject of analyte;

(b) determining the concentration or amount in a later test sample from the subject of analyte; and

(c) comparing the concentration or amount of analyte as determined in step (b) with the concentration or amount of analyte determined in step (a), wherein if the concentration or amount
10 determined in step (b) is unchanged or is unfavorable when compared to the concentration or amount of analyte determined in step (a), then the disease in the subject is determined to have continued, progressed or worsened. By comparison, if the concentration or amount of analyte as determined in step (b) is favorable when compared to the concentration or amount of analyte as determined in step (a), then the disease in the subject is determined to have discontinued,
15 regressed or improved.

Optionally, the method further comprises comparing the concentration or amount of analyte as determined in step (b), for example, with a predetermined level. Further, optionally the method comprises treating the subject with one or more pharmaceutical compositions for a period of time if the comparison shows that the concentration or amount of analyte as determined in step
20 (b), for example, is unfavorably altered with respect to the predetermined level.

Still further, the methods can be used to monitor treatment in a subject receiving treatment with one or more pharmaceutical compositions. Specifically, such methods involve providing a first test sample from a subject before the subject has been administered one or more pharmaceutical compositions. Next, the concentration or amount in a first test sample from a
25 subject of analyte is determined (e.g., using the methods described herein or as known in the art). After the concentration or amount of analyte is determined, optionally the concentration or amount of analyte is then compared with a predetermined level. If the concentration or amount of analyte as determined in the first test sample is lower than the predetermined level, then the subject is not treated with one or more pharmaceutical compositions. However, if the
30 concentration or amount of analyte as determined in the first test sample is higher than the predetermined level, then the subject is treated with one or more pharmaceutical compositions for a period of time. The period of time that the subject is treated with the one or more

pharmaceutical compositions can be determined by one skilled in the art (for example, the period of time can be from about seven (7) days to about two years, preferably from about fourteen (14) days to about one (1) year).

During the course of treatment with the one or more pharmaceutical compositions,
5 second and subsequent test samples are then obtained from the subject. The number of test samples and the time in which said test samples are obtained from the subject are not critical. For example, a second test sample could be obtained seven (7) days after the subject is first administered the one or more pharmaceutical compositions, a third test sample could be obtained two (2) weeks after the subject is first administered the one or more pharmaceutical
10 compositions, a fourth test sample could be obtained three (3) weeks after the subject is first administered the one or more pharmaceutical compositions, a fifth test sample could be obtained four (4) weeks after the subject is first administered the one or more pharmaceutical compositions, etc.

After each second or subsequent test sample is obtained from the subject, the
15 concentration or amount of analyte is determined in the second or subsequent test sample is determined (e.g., using the methods described herein or as known in the art). The concentration or amount of analyte as determined in each of the second and subsequent test samples is then compared with the concentration or amount of analyte as determined in the first test sample (e.g., the test sample that was originally optionally compared to the predetermined level). If the
20 concentration or amount of analyte as determined in step (c) is favorable when compared to the concentration or amount of analyte as determined in step (a), then the disease in the subject is determined to have discontinued, regressed or improved, and the subject should continue to be administered the one or pharmaceutical compositions of step (b). However, if the concentration or amount determined in step (c) is unchanged or is unfavorable when compared to the
25 concentration or amount of analyte as determined in step (a), then the disease in the subject is determined to have continued, progressed or worsened, and the subject should be treated with a higher concentration of the one or more pharmaceutical compositions administered to the subject in step (b) or the subject should be treated with one or more pharmaceutical compositions that are different from the one or more pharmaceutical compositions administered to the subject in step
30 (b). Specifically, the subject can be treated with one or more pharmaceutical compositions that are different from the one or more pharmaceutical compositions that the subject had previously received to decrease or lower said subject's analyte level.

Generally, for assays in which repeat testing may be done (e.g., monitoring disease progression and/or response to treatment), a second or subsequent test sample is obtained at a

period in time after the first test sample has been obtained from the subject. Specifically, a second test sample from the subject can be obtained minutes, hours, days, weeks or years after the first test sample has been obtained from the subject. For example, the second test sample can be obtained from the subject at a time period of about 1 minute, about 5 minutes, about 10
5 minutes, about 15 minutes, about 30 minutes, about 45 minutes, about 60 minutes, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, about 10 hours, about 11 hours, about 12 hours, about 13 hours, about 14 hours, about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours, about 21 hours, about 22 hours, about 23 hours, about 24 hours, about 2 days, about 3 days, about 4 days,
10 about 5 days, about 6 days, about 7 days, about 2 weeks, about 3 weeks, about 4 weeks, about 5 weeks, about 6 weeks, about 7 weeks, about 8 weeks, about 9 weeks, about 10 weeks, about 11 weeks, about 12 weeks, about 13 weeks, about 14 weeks, about 15 weeks, about 16 weeks, about 17 weeks, about 18 weeks, about 19 weeks, about 20 weeks, about 21 weeks, about 22 weeks, about 23 weeks, about 24 weeks, about 25 weeks, about 26 weeks, about 27 weeks, about 28
15 weeks, about 29 weeks, about 30 weeks, about 31 weeks, about 32 weeks, about 33 weeks, about 34 weeks, about 35 weeks, about 36 weeks, about 37 weeks, about 38 weeks, about 39 weeks, about 40 weeks, about 41 weeks, about 42 weeks, about 43 weeks, about 44 weeks, about 45 weeks, about 46 weeks, about 47 weeks, about 48 weeks, about 49 weeks, about 50 weeks, about 51 weeks, about 52 weeks, about 1.5 years, about 2 years, about 2.5 years, about 3.0 years, about
20 3.5 years, about 4.0 years, about 4.5 years, about 5.0 years, about 5.5 years, about 6.0 years, about 6.5 years, about 7.0 years, about 7.5 years, about 8.0 years, about 8.5 years, about 9.0 years, about 9.5 years or about 10.0 years after the first test sample from the subject is obtained.

When used to monitor disease progression, the above assay can be used to monitor the progression of disease in subjects suffering from acute conditions. Acute conditions, also known
25 as critical care conditions, refer to acute, life-threatening diseases or other critical medical conditions involving, for example, the cardiovascular system or excretory system. Typically, critical care conditions refer to those conditions requiring acute medical intervention in a hospital-based setting (including, but not limited to, the emergency room, intensive care unit, trauma center, or other emergent care setting) or administration by a paramedic or other field-
30 based medical personnel. For critical care conditions, repeat monitoring is generally done within a shorter time frame, namely, minutes, hours or days (e.g., about 1 minute, about 5 minutes, about 10 minutes, about 15 minutes, about 30 minutes, about 45 minutes, about 60 minutes, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, about 10 hours, about 11 hours, about 12 hours, about 13 hours, about 14 hours,
35 about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours,

about 21 hours, about 22 hours, about 23 hours, about 24 hours, about 2 days, about 3 days, about 4 days, about 5 days, about 6 days or about 7 days), and the initial assay likewise is generally done within a shorter timeframe, e.g., about minutes, hours or days of the onset of the disease or condition.

5 The assays also can be used to monitor the progression of disease in subjects suffering from chronic or non-acute conditions. Non-critical care or, non-acute conditions, refers to conditions other than acute, life-threatening disease or other critical medical conditions involving, for example, the cardiovascular system and/or excretory system. Typically, non-acute conditions include those of longer-term or chronic duration. For non-acute conditions, repeat
10 monitoring generally is done with a longer timeframe, e.g., hours, days, weeks, months or years (e.g., about 1 hour, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, about 10 hours, about 11 hours, about 12 hours, about 13 hours, about 14 hours, about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours, about 21 hours, about 22 hours, about 23 hours, about 24 hours,
15 about 2 days, about 3 days, about 4 days, about 5 days, about 6 days, about 7 days, about 2 weeks, about 3 weeks, about 4 weeks, about 5 weeks, about 6 weeks, about 7 weeks, about 8 weeks, about 9 weeks, about 10 weeks, about 11 weeks, about 12 weeks, about 13 weeks, about 14 weeks, about 15 weeks, about 16 weeks, about 17 weeks, about 18 weeks, about 19 weeks, about 20 weeks, about 21 weeks, about 22 weeks, about 23 weeks, about 24 weeks, about 25
20 weeks, about 26 weeks, about 27 weeks, about 28 weeks, about 29 weeks, about 30 weeks, about 31 weeks, about 32 weeks, about 33 weeks, about 34 weeks, about 35 weeks, about 36 weeks, about 37 weeks, about 38 weeks, about 39 weeks, about 40 weeks, about 41 weeks, about 42 weeks, about 43 weeks, about 44 weeks, about 45 weeks, about 46 weeks, about 47 weeks, about 48 weeks, about 49 weeks, about 50 weeks, about 51 weeks, about 52 weeks, about 1.5 years, about 2 years, about 2.5 years, about 3.0 years, about 3.5 years, about 4.0 years, about 4.5 years, about 5.0 years, about 5.5 years, about 6.0 years, about 6.5 years, about 7.0 years, about 7.5
25 years, about 8.0 years, about 8.5 years, about 9.0 years, about 9.5 years or about 10.0 years), and the initial assay likewise generally is done within a longer time frame, e.g., about hours, days, months or years of the onset of the disease or condition.

30 Furthermore, the above assays can be performed using a first test sample obtained from a subject where the first test sample is obtained from one source, such as urine, serum or plasma. Optionally, the above assays can then be repeated using a second test sample obtained from the subject where the second test sample is obtained from another source. For example, if the first test sample was obtained from urine, the second test sample can be obtained from serum or

plasma. The results obtained from the assays using the first test sample and the second test sample can be compared. The comparison can be used to assess the status of a disease or condition in the subject.

Moreover, the present disclosure also relates to methods of determining whether a
5 subject predisposed to or suffering from a given disease, disorder or condition will benefit from treatment. In particular, the disclosure relates to analyte companion diagnostic methods and products. Thus, the method of "monitoring the treatment of disease in a subject" as described herein further optimally also can encompass selecting or identifying candidates for therapy.

Thus, in particular embodiments, the disclosure also provides a method of determining
10 whether a subject having, or at risk for, a given disease, disorder or condition is a candidate for therapy. Generally, the subject is one who has experienced some symptom of a given disease, disorder or condition or who has actually been diagnosed as having, or being at risk for, a given disease, disorder or condition, and/or who demonstrates an unfavorable concentration or amount of analyte or a fragment thereof, as described herein.

The method optionally comprises an assay as described herein, where analyte is assessed
15 before and following treatment of a subject with one or more pharmaceutical compositions (e.g., particularly with a pharmaceutical related to a mechanism of action involving analyte), with immunosuppressive therapy, or by immunoabsorption therapy, or where analyte is assessed following such treatment and the concentration or the amount of analyte is compared against a
20 predetermined level. An unfavorable concentration or amount of analyte observed following treatment confirms that the subject will not benefit from receiving further or continued treatment, whereas a favorable concentration or amount of analyte observed following treatment confirms that the subject will benefit from receiving further or continued treatment. This confirmation assists with management of clinical studies, and provision of improved patient care.

It goes without saying that, while certain embodiments herein are advantageous when
25 employed to assess a given disease, disorder or condition as discussed herein, the assays and kits can be employed to assess analyte in other diseases, disorders and conditions. The method of assay can also involve the assay of other markers and the like.

The method of assay also can be used to identify a compound that ameliorates a given
30 disease, disorder or condition. For example, a cell that expresses analyte can be contacted with a candidate compound. The level of expression of analyte in the cell contacted with the compound can be compared to that in a control cell using the method of assay described herein.

B. Kit

A kit for assaying a test sample for the presence, amount or concentration of an analyte (or a fragment thereof) in a test sample is also provided. The kit comprises at least one component for assaying the test sample for the analyte (or a fragment thereof) and instructions for
5 assaying the test sample for the analyte (or a fragment thereof). The at least one component for assaying the test sample for the analyte (or a fragment thereof) can include a composition comprising an anti-analyte DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof), which is optionally immobilized on a solid phase.

The kit can comprise at least one component for assaying the test sample for an analyte
10 by immunoassay, e.g., chemiluminescent microparticle immunoassay, and instructions for assaying the test sample for an analyte by immunoassay, e.g., chemiluminescent microparticle immunoassay. For example, the kit can comprise at least one specific binding partner for an analyte, such as an anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof that can bind to the analyte, a variant thereof that can bind to the analyte, or a fragment of a variant that
15 can bind to the analyte) or an anti-analyte DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof), either of which can be detectably labeled. Alternatively or additionally, the kit can comprise detectably labeled analyte (or a fragment thereof that can bind to an anti-analyte, monoclonal/polyclonal antibody or an anti-analyte DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof)), which can compete with any analyte in a test sample for binding
20 to an anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof that can bind to the analyte, a variant thereof that can bind to the analyte, or a fragment of a variant that can bind to the analyte) or an anti-analyte DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof), either of which can be immobilized on a solid support. The kit can comprise a calibrator or control, e.g., isolated or purified analyte. The kit can comprise at least one container (e.g.,
25 tube, microtiter plates or strips, which can be already coated with a first specific binding partner, for example) for conducting the assay, and/or a buffer, such as an assay buffer or a wash buffer, either one of which can be provided as a concentrated solution, a substrate solution for the detectable label (e.g., an enzymatic label), or a stop solution. Preferably, the kit comprises all components, i.e., reagents, standards, buffers, diluents, etc., which are necessary to perform the
30 assay. The instructions can be in paper form or computer-readable form, such as a disk, CD, DVD, or the like.

Any antibodies, such as an anti-analyte antibody or an anti-analyte DVD-Ig, or tracer can incorporate a detectable label as described herein, such as a fluorophore, a radioactive moiety, an enzyme, a biotin/avidin label, a chromophore, a chemiluminescent label, or the like, or the kit can

include reagents for carrying out detectable labeling. The antibodies, calibrators and/or controls can be provided in separate containers or pre-dispensed into an appropriate assay format, for example, into microtiter plates.

Optionally, the kit includes quality control components (for example, sensitivity panels, calibrators, and positive controls). Preparation of quality control reagents is well-known in the art and is described on insert sheets for a variety of immunodiagnostic products. Sensitivity panel members optionally are used to establish assay performance characteristics, and further optionally are useful indicators of the integrity of the immunoassay kit reagents, and the standardization of assays.

The kit can also optionally include other reagents required to conduct a diagnostic assay or facilitate quality control evaluations, such as buffers, salts, enzymes, enzyme co-factors, enzyme substrates, detection reagents, and the like. Other components, such as buffers and solutions for the isolation and/or treatment of a test sample (e.g., pretreatment reagents), also can be included in the kit. The kit can additionally include one or more other controls. One or more of the components of the kit can be lyophilized, in which case the kit can further comprise reagents suitable for the reconstitution of the lyophilized components.

The various components of the kit optionally are provided in suitable containers as necessary, e.g., a microtiter plate. The kit can further include containers for holding or storing a sample (e.g., a container or cartridge for a urine sample). Where appropriate, the kit optionally also can contain reaction vessels, mixing vessels, and other components that facilitate the preparation of reagents or the test sample. The kit can also include one or more instruments for assisting with obtaining a test sample, such as a syringe, pipette, forceps, measured spoon, or the like.

If the detectable label is at least one acridinium compound, the kit can comprise at least one acridinium-9-carboxamide, at least one acridinium-9-carboxylate aryl ester, or any combination thereof. If the detectable label is at least one acridinium compound, the kit also can comprise a source of hydrogen peroxide, such as a buffer, a solution, and/or at least one basic solution. If desired, the kit can contain a solid phase, such as a magnetic particle, bead, test tube, microtiter plate, cuvette, membrane, scaffolding molecule, film, filter paper, disc or chip.

C. Adaptation of Kit and Method

The kit (or components thereof), as well as the method of determining the presence, amount or concentration of an analyte in a test sample by an assay, such as an immunoassay as

described herein, can be adapted for use in a variety of automated and semi-automated systems (including those wherein the solid phase comprises a microparticle), as described, e.g., in U.S. Patent Nos. 5,089,424 and 5,006,309, and as commercially marketed, e.g., by Abbott Laboratories (Abbott Park, IL) as ARCHITECT®.

5 Some of the differences between an automated or semi-automated system as compared to a non-automated system (e.g., ELISA) include the substrate to which the first specific binding partner (e.g., an anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof, a variant thereof, or a fragment of a variant thereof) or an anti-analyte DVD-Ig (or a fragment thereof, a variant thereof, or a fragment of a variant thereof) is attached; either way, sandwich formation
10 and analyte reactivity can be impacted), and the length and timing of the capture, detection and/or any optional wash steps. Whereas a non-automated format, such as an ELISA, may require a relatively longer incubation time with sample and capture reagent (e.g., about 2 hours), an automated or semi-automated format (e.g., ARCHITECT®, Abbott Laboratories) may have a relatively shorter incubation time (e.g., approximately 18 minutes for ARCHITECT®). Similarly,
15 whereas a non-automated format, such as an ELISA, may incubate a detection antibody, such as the conjugate reagent, for a relatively longer incubation time (e.g., about 2 hours), an automated or semi-automated format (e.g., ARCHITECT®) may have a relatively shorter incubation time (e.g., approximately 4 minutes for the ARCHITECT®).

Other platforms available from Abbott Laboratories include, but are not limited to,
20 AxSYM®, IMx® (see, e.g., U.S. Pat. No. 5,294,404), PRISM®, EIA (bead), and Quantum™ II, as well as other platforms. Additionally, the assays, kits and kit components can be employed in other formats, for example, on electrochemical or other hand-held or point-of-care assay systems. The present disclosure is, for example, applicable to the commercial Abbott Point of Care (i-STAT®, Abbott Laboratories) electrochemical immunoassay system that performs sandwich
25 immunoassays. Immunosensors and their methods of manufacture and operation in single-use test devices are described, for example in, U.S. Patent No. 5,063,081, U.S. Pat. App. Pub. No. 2003/0170881, U.S. Pat. App. Pub. No. 2004/0018577, U.S. Pat. App. Pub. No. 2005/0054078, and U.S. Pat. App. Pub. No. 2006/0160164.

In particular, with regard to the adaptation of an analyte assay to the I-STAT® system,
30 the following configuration is preferred. A microfabricated silicon chip is manufactured with a pair of gold amperometric working electrodes and a silver-silver chloride reference electrode. On one of the working electrodes, polystyrene beads (0.2 mm diameter) with immobilized anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof, a variant thereof, or a fragment of a variant thereof) or anti-analyte DVD-Ig (or a fragment thereof, a variant thereof, or a

fragment of a variant thereof), are adhered to a polymer coating of patterned polyvinyl alcohol over the electrode. This chip is assembled into an I-STAT® cartridge with a fluidics format suitable for immunoassay. On a portion of the wall of the sample-holding chamber of the cartridge there is a layer comprising a specific binding partner for an analyte, such as an anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof, a variant thereof, or a fragment of a variant thereof that can bind the analyte) or an anti-analyte DVD-Ig (or a fragment thereof, a variant thereof, or a fragment of a variant thereof that can bind the analyte), either of which can be detectably labeled. Within the fluid pouch of the cartridge is an aqueous reagent that includes p-aminophenol phosphate.

10 In operation, a sample suspected of containing an analyte is added to the holding chamber of the test cartridge, and the cartridge is inserted into the I-STAT® reader. After the specific binding partner for an analyte has dissolved into the sample, a pump element within the cartridge forces the sample into a conduit containing the chip. Here it is oscillated to promote formation of the sandwich. In the penultimate step of the assay, fluid is forced out of the pouch and into the
15 conduit to wash the sample off the chip and into a waste chamber. In the final step of the assay, the alkaline phosphatase label reacts with p-aminophenol phosphate to cleave the phosphate group and permit the liberated p-aminophenol to be electrochemically oxidized at the working electrode. Based on the measured current, the reader is able to calculate the amount of analyte in the sample by means of an embedded algorithm and factory-determined calibration curve.

20 It further goes without saying that the methods and kits as described herein necessarily encompass other reagents and methods for carrying out the immunoassay. For instance, encompassed are various buffers such as are known in the art and/or which can be readily prepared or optimized to be employed, e.g., for washing, as a conjugate diluent, microparticle diluent, and/or as a calibrator diluent. An exemplary conjugate diluent is ARCHITECT®
25 conjugate diluent employed in certain kits (Abbott Laboratories, Abbott Park, IL) and containing 2-(N-morpholino)ethanesulfonic acid (MES), a salt, a protein blocker, an antimicrobial agent, and a detergent. An exemplary calibrator diluent is ARCHITECT® human calibrator diluent employed in certain kits (Abbott Laboratories, Abbott Park, IL), which comprises a buffer containing MES, other salt, a protein blocker, and an antimicrobial agent. Additionally, as
30 described in U.S. Patent Application No. 61/142,048 filed December 31, 2008, improved signal generation may be obtained, e.g., in an I-Stat cartridge format, using a nucleic acid sequence linked to the signal antibody as a signal amplifier.

EXEMPLIFICATION

Example 1: Design, Construction, and Analysis of a DVD-Ig

Example 1.1: Assays Used to Identify and Characterize Parent Antibodies and DVD-Ig

The following assays were used throughout the Examples to identify and characterize
5 parent antibodies and DVD-Ig, unless otherwise stated.

Example 1.1.1: Assays Used To Determine Binding and Affinity of Parent Antibodies and DVD-Ig for Their Target Antigen(s)

Example 1.1.1A: Direct Bind ELISA

Enzyme Linked Immunosorbent Assays to screen for antibodies that bind a desired target
10 antigen were performed as follows. High bind ELISA plates (Corning Costar # 3369, Acton, MA)
were coated with 100 μ L/well of 10 μ g/ml of desired target antigen (R&D Systems, Minneapolis,
MN) or desired target antigen extra-cellular domain / FC fusion protein (R&D Systems,
Minneapolis, MN) or monoclonal mouse anti-polyHistidine antibody (R&D Systems # MAB050,
Minneapolis, MN) in phosphate buffered saline (10X PBS, Abbott Bioresearch Center, Media
15 Prep# MPS-073, Worcester, MA) overnight at 4°C. Plates were washed four times with PBS
containing 0.02% Tween 20. Plates were blocked by the addition of 300 μ L/well blocking
solution (non-fat dry milk powder, various retail suppliers, diluted to 2% in PBS) for 1/2 hour at
room temperature. Plates were washed four times after blocking with PBS containing 0.02%
Tween 20.

20 Alternatively, one hundred microliters per well of 10 μ g/ml of Histidine (His) tagged
desired target antigen (R&D Systems, Minneapolis, MN) was added to ELISA plates coated with
monoclonal mouse anti-polyHistidine antibody as described above and incubated for 1 hour at
room temperature. Wells were washed four times with PBS containing 0.02% Tween 20.

25 One hundred microliters of antibody or DVD-Ig preparations diluted in blocking solution
as described above was added to the desired target antigen plate or desired target antigen / FC
fusion plate or the anti-polyHistidine antibody / His tagged desired target antigen plate prepared
as described above and incubated for 1 hour at room temperature. Wells were washed four times
with PBS containing 0.02% Tween 20.

30 One hundred microliters of 10ng/mL goat anti-human IgG –FC specific HRP conjugated
antibody (Southern Biotech # 2040-05, Birmingham, AL) was added to each well of the desired
target antigen plate or anti-polyHistidine antibody / Histidine tagged desired target antigen plate.

Alternatively, one hundred microliters of 10 ng/mL goat anti-human IgG –kappa light chain specific HRP conjugated antibody (Southern Biotech # 2060-05 Birmingham, AL) was added to each well of the desired target antigen / FC fusion plate and incubated for 1 hour at room temperature. Plates were washed 4 times with PBS containing 0.02% Tween 20.

- 5 One hundred microliters of enhanced TMB solution (Neogen Corp. #308177, K Blue, Lexington, KY) was added to each well and incubated for 10 minutes at room temperature. The reaction was stopped by the addition of 50 μ L 1N sulphuric acid. Plates were read spectrophotometrically at a wavelength of 450 nm.

Table 3 contains a list of the antigens used in the Direct Bind Assay.

- 10 Table 4 contains the binding data expressed as EC50 in nM for those antibodies and DVD-Ig constructs tested in the Direct Bind ELISA assay.

In the Direct Bind ELISA, binding was sometimes not observed, probably because the antibody binding site on the target antigen was either “masked” or the antigen is “distorted” when coated to the plastic surface. The inability of a DVD-Ig to bind its target may also be due to steric limitation imposed on DVD-Ig by the Direct Bind ELISA format. The parent antibodies and DVD-Igs that did not bind in the Direct Bind ELISA format bound to target antigen in other ELISA formats, such as FACS, Biacore or bioassay. Non-binding of a DVD-Ig was also restored by adjusting the linker length between the two variable domains of the DVD-Ig, as shown previously.

20

Example 1.1.1.B: Capture ELISA

ELISA plates (Nunc, MaxiSorp, Rochester, NY) are incubated overnight at 4°C with anti-human Fc antibody (5 μ g/ml in PBS, Jackson ImmunoResearch, West Grove, PA). Plates are washed three times in washing buffer (PBS containing 0.05% Tween 20), and blocked for 1 hour at 25°C in blocking buffer (PBS containing 1% BSA). Wells are washed three times, and serial dilutions of each antibody or DVD-Ig in PBS containing 0.1% BSA are added to the wells and incubated at 25°C for 1 hour. The wells are washed three times, and biotinylated antigen (2nM) is added to the plates and incubated for 1 hour at 25°C. The wells are washed three times and incubated for 1 hour at 25°C with streptavidin-HRP (KPL #474-3000, Gaithersburg, MD). The wells are washed three times, and 100 μ l of ULTRA-TMB ELISA (Pierce, Rockford, IL) is added per well. Following color development the reaction is stopped with 1N HCL and absorbance at 450nm is measured.

30

Example 1.1.1.C: Affinity Determination Using BIACORE Technology**Table 3: Reagent Used in Biacore Analyses**

Assay	Antigen	Vendor Designation	Vendor	Catalog #
	NGF	Recombinant Human β -NGF	R&D systems	256-GF
	TNF α	Recombinant Human TNF- α /TNFSF1A	R&D systems	210-TA
	SOST	Recombinant Human SOST	R&D systems	1406-ST

ECD = Extracellular Domain

5 /FC = antigen/IgG FC domain fusion protein

BIACORE Methods:

The BIACORE assay (Biacore, Inc, Piscataway, NJ) determines the affinity of antibodies or DVD-Ig with kinetic measurements of on-rate and off-rate constants. Binding of antibodies or DVD-Ig to a target antigen (for example, a purified recombinant target antigen) is determined by surface plasmon resonance-based measurements with a Biacore® 1000 or 3000 instrument (Biacore® AB, Uppsala, Sweden) using running HBS-EP (10 mM HEPES [pH 7.4], 150 mM NaCl, 3 mM EDTA, and 0.005% surfactant P20) at 25° C. All chemicals are obtained from Biacore® AB (Uppsala, Sweden) or otherwise from a different source as described in the text.

15 For example, approximately 5000 RU of goat anti-mouse IgG, (Fc γ), fragment specific polyclonal antibody (Pierce Biotechnology Inc, Rockford, IL) diluted in 10 mM sodium acetate (pH 4.5) is directly immobilized across a CM5 research grade biosensor chip using a standard amine coupling kit according to manufacturer's instructions and procedures at 25 μ g/ml. Unreacted moieties on the biosensor surface are blocked with ethanolamine. Modified carboxymethyl dextran surface in flowcell 2 and 4 is used as a reaction surface. Unmodified carboxymethyl dextran without goat anti-mouse IgG in flow cell 1 and 3 is used as the reference surface. For kinetic analysis, rate equations derived from the 1:1 Langmuir binding model are fitted simultaneously to association and dissociation phases of all eight injections (using global fit analysis) with the use of Biaevaluation 4.0.1 software. Purified antibodies or DVD-Ig are diluted in HEPES-buffered saline for capture across goat anti-mouse IgG specific reaction surfaces. Antibodies or DVD-Ig to be captured as a ligand (25 μ g/ml) are injected over reaction matrices at a flow rate of 5 μ l/min. The association and dissociation rate constants, k_{on} ($M^{-1}s^{-1}$) and k_{off} (s^{-1}) are determined under a continuous flow rate of 25 μ l/min. Rate constants are derived by making kinetic binding measurements at different antigen concentrations ranging from 10 – 200 nM. The equilibrium dissociation constant (M) of the reaction between antibodies or DVD-Igs and the target antigen is then calculated from the kinetic rate constants by the following formula: $K_D =$

20
25
30

k_{off}/k_{on} . Binding is recorded as a function of time and kinetic rate constants are calculated. In this assay, on-rates as fast as $10^6 M^{-1}s^{-1}$ and off-rates as slow as $10^{-6} s^{-1}$ can be measured.

Table 4: BIACORE Analysis of Parental Antibodies and DVD Constructs

Parent Antibody or DVD-Ig ID	N-Terminal Variable Domain (VD)	C-Terminal Variable Domain (VD)	k_{on}	k_{off}	K_D
			($M^{-1}s^{-1}$)	(s^{-1})	(M)
AB213	TNF (seq 2)		3.23E+06	1.08E-04	3.35E-11
AB022	SOST		9.68E+06	3.96E-04	4.08E-11
DVD1454	SOST		1.00E+07	8.60E-04	8.40E-11
DVD1454		TNF (seq 2)	2.30E+05	4.90E-05	2.10E-10
AB213	TNF (seq 2)		3.23E+06	1.08E-04	3.35E-11
AB020	NGF		6.06E+05	4.08E-06	6.73E-12
DVD1460	NGF		2.30E+06	2.00E-06	8.50E-13
DVD1460		TNF (seq 2)	1.40E+05	3.60E-05	2.60E-10
AB213	TNF (seq 2)		3.23E+06	1.08E-04	3.35E-11
DVD1466	LPA	TNF (seq 2)	1.60E+05	6.30E-05	3.90E-10
AB017	TNF (seq 1)		1.61E+06	1.29E-04	7.99E-11
DVD1471	TNF (seq 1)	LPA	2.20E+06	3.80E-05	1.80E-11
DVD1472	LPA	TNF (seq 1)	2.90E+05	1.60E-04	5.50E-10
AB217	TNF (seq 5)		1.85E+06	5.04E-05	2.72E-11
DVD1473	TNF (seq 5)	PGE2	2.60E+06	4.40E-05	1.70E-11
AB217	TNF (seq 5)		1.85E+06	5.04E-05	2.72E-11
AB022	SOST		9.68E+06	3.96E-04	4.08E-11
DVD1475	TNF (seq 5)		2.70E+06	3.60E-05	1.30E-11
DVD1475		SOST	8.10E+05	7.30E-04	9.00E-10
AB217	TNF (seq 5)		1.85E+06	5.04E-05	2.72E-11
AB020	NGF		6.06E+05	4.08E-06	6.73E-12
DVD1477	TNF (seq 5)		2.50E+06	3.80E-05	1.50E-11
DVD1477		NGF	1.60E+05	<1.0E-06	<6.3E-12
DVD1478	NGF		2.40E+06	6.00E-06	2.50E-12
DVD1478		TNF (seq 5)	1.50E+05	1.50E-05	1.00E-10
AB217	TNF (seq 5)		1.85E+06	5.04E-05	2.72E-11
DVD1479	TNF (seq 5)	LPA	2.40E+06	4.10E-05	1.70E-11
DVD1480	LPA	TNF (seq 5)	1.30E+05	3.40E-05	2.60E-10
AB218	TNF (seq 6)		1.79E+06	5.15E-05	2.88E-11
DVD1481	TNF (seq 6)	PGE2	2.60E+06	2.40E-05	8.90E-12
DVD1482	PGE2	TNF (seq 6)	1.60E+05	1.50E-05	9.20E-11
AB218	TNF (seq 6)		1.79E+06	5.15E-05	2.88E-11
AB022	SOST		9.68E+06	3.96E-04	4.08E-11
DVD1483	TNF (seq 6)		2.80E+06	3.30E-05	1.20E-11
DVD1483		SOST	1.10E+06	8.30E-04	7.80E-10
AB218	TNF (seq 6)		1.79E+06	5.15E-05	2.88E-11
AB020	NGF		6.06E+05	4.08E-06	6.73E-12
DVD1485	TNF (seq 6)		2.10E+06	3.60E-05	1.70E-11
DVD1485		NGF	1.00E+05	<1.0E-06	<1.0E-11
AB218	TNF (seq 6)		1.79E+06	5.15E-05	2.88E-11
DVD1487	TNF (seq 6)	LPA	1.80E+06	2.60E-05	1.40E-11

Parent Antibody or DVD-Ig ID	N-Terminal Variable Domain (VD)	C-Terminal Variable Domain (VD)	k_{on}	k_{off}	K_D
			(M-1s-1)	(s-1)	(M)
DVD1488	LPA	TNF (seq 6)	2.30E+05	1.90E-05	8.40E-11
AB213		TNF (seq 2)	3.23E+06	1.08E-04	3.35E-11
DVD1490	PGE2	TNF (seq 2)	5.00E+05	6.10E-05	1.2E-10
AB213		TNF (seq 2)	3.23E+06	1.08E-04	3.35E-11
AB022		SOST	9.68E+06	3.96E-04	4.08E-11
DVD1495	TNF (seq 2)		2.50E+06	5.40E-05	2.10E-11
DVD1495		SOST	4.40E+04	9.50E-04	2.20E-08
AB213		TNF (seq 2)	3.23E+06	1.08E-04	3.35E-11
AB020		NGF	6.06E+05	4.08E-06	6.73E-12
DVD1502	NGF		2.50E+06	<1.0E-06	<4.0E-13
DVD1502		TNF (seq 2)	1.80E+05	5.50E-05	3.00E-10
AB213		TNF (seq 2)	3.23E+06	1.08E-04	3.35E-11
DVD1508	LPA	TNF (seq 2)	2.50E+05	5.00E-05	2.00E-10
AB017		TNF (seq 1)	1.61E+06	1.29E-04	7.99E-11
DVD1514	LPA	TNF (seq 1)	3.30E+05	6.70E-05	2.00E-10
AB217		TNF (seq 5)	1.85E+06	5.04E-05	2.72E-11
DVD1515	TNF (seq 5)	PGE2	3.10E+06	3.70E-05	1.20E-11
AB217		TNF (seq 5)	1.85E+06	5.04E-05	2.72E-11
AB022		SOST	9.68E+06	3.96E-04	4.08E-11
DVD1517	TNF (seq 5)		2.60E+06	4.70E-05	1.80E-11
DVD1517		SOST	8.90E+04	5.00E-04	5.50E-09
AB217		TNF (seq 5)	1.85E+06	5.04E-05	2.72E-11
AB020		NGF	6.06E+05	4.08E-06	6.73E-12
DVD1519	TNF (seq 5)		2.10E+06	4.20E-05	2.00E-11
DVD1519		NGF	1.10E+05	<1.0E-06	<1.0E-11
DVD1520	NGF		2.70E+06	9.30E-06	3.40E-12
DVD1520		TNF (seq 5)	1.10E+06	4.40E-06	3.80E-12
AB218		TNF (seq 6)	1.79E+06	5.15E-05	2.88E-11
DVD1523	TNF (seq 6)	PGE2	2.20E+06	2.00E-05	9.10E-12
DVD1524	PGE2	TNF (seq 6)	1.00E+05	1.70E-06	1.70E-11
AB218		TNF (seq 6)	1.79E+06	5.15E-05	2.88E-11
AB022		SOST	9.68E+06	3.96E-04	4.08E-11
DVD1525	TNF (seq 6)		2.00E+06	3.00E-05	1.50E-11
DVD1525		SOST	9.10E+04	8.90E-04	9.80E-09
AB218		TNF (seq 6)	1.79E+06	5.15E-05	2.88E-11
AB020		NGF	6.06E+05	4.08E-06	6.73E-12
DVD1527	TNF (seq 6)		2.30E+06	4.00E-05	1.70E-11
DVD1527		NGF	1.00E+05	<1.0E-06	<1.0E-11
DVD1528	NGF		2.40E+06	7.30E-06	3.00E-12
DVD1528		TNF (seq 6)	1.70E+05	9.40E-06	5.60E-11
AB218		TNF (seq 6)	1.79E+06	5.15E-05	2.88E-11
DVD1529	TNF (seq 6)	LPA	2.50E+06	2.60E-05	1.00E-11
DVD1530	LPA	TNF (seq 6)	8.70E+04	1.60E-06	1.90E-11

Example 1.1.2: Assays Used To Determine the Functional Activity Of Parent Antibodies And DVD-Ig

Example 1.1.2.A: Cytokine Bioassay

The ability of an anti-cytokine or an anti-growth factor parent antibody or DVD-Ig
5 containing anti-cytokine or anti-growth factor sequences to inhibit or neutralize a target cytokine
or growth factor bioactivity is analyzed by determining the inhibitory potential of the antibody or
DVD-Ig. For example, the ability of an anti-IL-4 antibody to inhibit IL-4 mediated IgE
production may be used. For example, human naive B cells are isolated from peripheral blood,
respectively, buffy coats by Ficoll-paque density centrifugation, followed by magnetic separation
10 with MACS beads (Miltenyi Biotec, Bergisch Gladbach, Germany) specific for human sIgD
FITC labeled goat F(ab)₂ antibodies followed by anti-FITC MACS beads. Magnetically sorted
naive B cells are adjusted to 3×10^5 cells per ml in XV15 and plated out in 100 μ l per well of 96-
well plates in a 6 x 6 array in the center of the plate, surrounded by PBS filled wells during the 10
days of culture at 37° C in the presence of 5% CO₂. One plate each is prepared per antibody to be
15 tested, comprising 3 wells each of un-induced and induced controls and quintuplicate repeats of
antibody titrations starting at 7 μ g/ml and running in 3-fold dilution down to 29 ng/ml final
concentrations added in 50 μ l four times concentrated pre-dilution. To induce IgE production,
rhIL-4 at 20 ng/ml plus anti-CD40 monoclonal antibody (Novartis, Basel, Switzerland) at 0.5
 μ g/ml final concentrations in 50 μ l each are added to each well, and IgE concentrations are
20 determined at the end of the culture period by a standard sandwich ELISA method.

Example 1.1.2.B: Cytokine Release Assay

The ability of a parent antibody or DVD-Ig to cause cytokine release is analyzed.
Peripheral blood is withdrawn from three healthy donors by venipuncture into heparized
vacutainer tubes. Whole blood is diluted 1:5 with RPMI-1640 medium and placed in 24-well
25 tissue culture plates at 0.5 mL per well. The anti-cytokine antibodies (e.g., anti-IL-4) are diluted
into RPMI-1640 and placed in the plates at 0.5 mL/well to give final concentrations of 200, 100,
50, 10, and 1 μ g/mL. The final dilution of whole blood in the culture plates is 1:10. LPS and PHA
are added to separate wells at 2 μ g/mL and 5 μ g/mL final concentration as a positive control for
cytokine release. Polyclonal human IgG is used as negative control antibody. The experiment is
30 performed in duplicate. Plates are incubated at 37°C at 5% CO₂. Twenty-four hours later the
contents of the wells are transferred into test tubes and spun for 5 minutes at 1200 rpm. Cell-free
supernatants are collected and frozen for cytokine assays. Cells left over on the plates and in the
tubes are lysed with 0.5 mL of lysis solution, and placed at -20°C and thawed. 0.5 mL of medium
is added (to bring the volume to the same level as the cell-free supernatant samples) and the cell
35 preparations are collected and frozen for cytokine assays. Cell-free supernatants and cell lysates

are assayed for cytokine levels by ELISA, for example, for levels of IL-8, IL-6, IL-1 β , IL-1RA, or TNF- α .

Example 1.1.2.C: Cytokine Cross-Reactivity Study

The ability of an anti-cytokine parent antibody or DVD-Ig directed to a cytokine(s) of
5 interest to cross react with other cytokines is analyzed. Parent antibodies or DVD-Ig are
immobilized on a Biacore biosensor matrix. An anti-human Fc mAb is covalently linked via free
amine groups to the dextran matrix by first activating carboxyl groups on the matrix with 100mM
N-hydroxysuccinimide (NHS) and 400mM N-Ethyl-N'-(3-dimethylaminopropyl)-carbodiimide
hydrochloride (EDC). Approximately 50 μ L of each antibody or DVD-Ig preparation at a
10 concentration of 25 μ g/mL, diluted in sodium acetate, pH 4.5, is injected across the activated
biosensor and free amines on the protein are bound directly to the activated carboxyl groups.
Typically, 5000 Resonance Units (RU's) are immobilized. Unreacted matrix EDC-esters are
deactivated by an injection of 1 M ethanolamine. A second flow cell is prepared as a reference
standard by immobilizing human IgG1/K using the standard amine coupling kit. SPR
15 measurements are performed using the CM biosensor chip. All antigens to be analyzed on the
biosensor surface are diluted in HBS-EP running buffer containing 0.01% P20.

To examine the cytokine binding specificity, excess cytokine of interest (100nM, e.g.,
soluble recombinant human) is injected across the anti-cytokine parent antibody or DVD-Ig
immobilized biosensor surface (5 minute contact time). Before injection of the cytokine of
20 interest and immediately afterward, HBS-EP buffer alone flows through each flow cell. The net
difference in the signals between the baseline and the point corresponding to approximately 30
seconds after completion of cytokine injection are taken to represent the final binding value.
Again, the response is measured in Resonance Units. Biosensor matrices are regenerated using
10mM HCl before injection of the next sample where a binding event is observed, otherwise
25 running buffer was injected over the matrices. Human cytokines (e.g., IL-1 α , IL-1 β , IL-2, IL-3,
IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-15, IL-16, IL-17, IL-18, IL-19,
IL-20, IL-22, IL-23, IL-27, TNF- α , TNF- β , and IFN- γ , for example) are also simultaneously
injected over the immobilized mouse IgG1/K reference surface to record any nonspecific binding
background. By preparing a reference and reaction surface, Biacore can automatically subtract
30 the reference surface data from the reaction surface data in order to eliminate the majority of the
refractive index change and injection noise. Thus, it is possible to ascertain the true binding
response attributed to an anti-cytokine antibody or DVD-Ig binding reaction.

When a cytokine of interest is injected across immobilized anti-cytokine antibody,
significant binding is observed. 10mM HCl regeneration completely removes all non-covalently

associated proteins. Examination of the sensorgram shows that immobilized anti-cytokine antibody or DVD-Ig binding to soluble cytokine is strong and robust. After confirming the expected result with the cytokine of interest, the panel of remaining recombinant human cytokines is tested, for each antibody or DVD-Ig separately. The amount of anti-cytokine antibody or DVD-Ig bound or unbound cytokine for each injection cycle is recorded. The results from three independent experiments are used to determine the specificity profile of each antibody or DVD-Ig. Antibodies or DVD-Ig with the expected binding to the cytokine of interest and no binding to any other cytokine are selected.

Example 1.1.2.D: Tissue Cross Reactivity

10 Tissue cross reactivity studies are done in three stages, with the first stage including cryosections of 32 tissues, second stage including up to 38 tissues, and the 3rd stage including additional tissues from 3 unrelated adults as described below. Studies are done typically at two dose levels.

Stage 1: Cryosections (about 5 μ m) of human tissues (32 tissues (typically: Adrenal Gland, Gastrointestinal Tract, Prostate, Bladder, Heart, Skeletal Muscle, Blood Cells, Kidney, Skin, Bone Marrow, Liver, Spinal Cord, Breast, Lung, Spleen, Cerebellum, Lymph Node, Testes, Cerebral Cortex, Ovary, Thymus, Colon, Pancreas, Thyroid, Endothelium, Parathyroid, Ureter, Eye, Pituitary, Uterus, Fallopian Tube and Placenta) from one human donor obtained at autopsy or biopsy) are fixed and dried on object glass. The peroxidase staining of tissue sections is performed, using the avidin-biotin system.

Stage 2: Cryosections (about 5 μ m) of human tissues 38 tissues (including adrenal, blood, blood vessel, bone marrow, cerebellum, cerebrum, cervix, esophagus, eye, heart, kidney, large intestine, liver, lung, lymph node, breast mammary gland, ovary, oviduct, pancreas, parathyroid, peripheral nerve, pituitary, placenta, prostate, salivary gland, skin, small intestine, spinal cord, spleen, stomach, striated muscle, testis, thymus, thyroid, tonsil, ureter, urinary bladder, and uterus) from 3 unrelated adults obtained at autopsy or biopsy) are fixed and dried on object glass. The peroxidase staining of tissue sections is performed, using the avidin-biotin system.

Stage 3: Cryosections (about 5 μ m) of cynomolgus monkey tissues (38 tissues (including adrenal, blood, blood vessel, bone marrow, cerebellum, cerebrum, cervix, esophagus, eye, heart, kidney, large intestine, liver, lung, lymph node, breast mammary gland, ovary, oviduct, pancreas, parathyroid, peripheral nerve, pituitary, placenta, prostate, salivary gland, skin, small intestine, spinal cord, spleen, stomach, striated muscle, testis, thymus, thyroid, tonsil, ureter, urinary

bladder, and uterus) from 3 unrelated adult monkeys obtained at autopsy or biopsy) are fixed and dried on object glass. The peroxidase staining of tissue sections is performed, using the avidin-biotin system.

The antibody or DVD-Ig is incubated with the secondary biotinylated anti-human IgG and developed into immune complex. The immune complex at the final concentrations of 2 and 10 $\mu\text{g}/\text{mL}$ of antibody or DVD-Ig is added onto tissue sections on object glass and then the tissue sections are reacted for 30 minutes with a avidin-biotin-peroxidase kit. Subsequently, DAB (3,3'-diaminobenzidine), a substrate for the peroxidase reaction, is applied for 4 minutes for tissue staining. Antigen-Sepharose beads are used as positive control tissue sections. Target antigen and human serum blocking studies serve as additional controls. The immune complex at the final concentrations of 2 and 10 $\mu\text{g}/\text{mL}$ of antibody or DVD-Ig is pre-incubated with target antigen (final concentration of 100 $\mu\text{g}/\text{ml}$) or human serum (final concentration 10%) for 30 minutes, and then added onto the tissue sections on object glass and then the tissue sections are reacted for 30 minutes with a avidin-biotin-peroxidase kit. Subsequently, DAB (3,3'-diaminobenzidine), a substrate for the peroxidase reaction, is applied for 4 minutes for tissue staining.

Any specific staining is judged to be either an expected (e.g., consistent with antigen expression) or unexpected reactivity based upon known expression of the target antigen in question. Any staining judged specific is scored for intensity and frequency. The tissue staining between stage 2 (human tissue) and stage 3 (cynomolgus monkey tissue) is either judged to be similar or different.

Example 1.1.2.D: Neutralization of huTNF α Neutralization of huTNF α

L929 cells were grown to a semi-confluent density and harvested using 0.05% trypsin (Gibco#25300). The cells were washed with PBS, counted and resuspended at $1\text{E}6$ cells/mL in assay media containing 4 $\mu\text{g}/\text{mL}$ actinomycin D. The cells were seeded in a 96-well plate (Costar#3599) at a volume of 50 μL and $5\text{E}4$ cells/well. The DVD-IgTM and control IgG were diluted to a 4x concentration in assay media and serial 1:3 dilutions were prepared. The huTNF α was diluted to 400 pg/mL in assay media. An antibody sample (200 μL) was added to the huTNF α (200 μL) in a 1:2 dilution scheme and allowed to incubate for 0.5 hour at room temperature.

The DVD-IgTM / huTNF α solution was added to the plated cells at 100 μL for a final concentration of 100 pg/mL huTNF α and 25 nM – 0.00014 nM DVD-IgTM. The plates were incubated for 20 hours at 37°C, 5 % CO₂. To quantitate viability, 100 μL was removed from the wells and 10 μL of WST-1 reagent (Roche cat# 11644807001) was added. Plates were incubated under assay conditions for 3.5 hours, centrifuged at 500 xg and 75 μL supernatant transferred to

an ELISA plate (Costar cat#3369). The plates were read at OD 420-600 nm on a Spectromax 190 ELISA plate reader. An average EC50 from several assays is included in Table 5.

Table 5: HuTNF α Neutralization Assay With huTNF α Parent Antibody and DVD-Ig Constructs

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	N-terminal VD TNF α Neutralization Assay EC50 nM	C-terminal VD TNF α Neutralization Assay EC50 nM
AB017	TNF (seq 1)			0.0461
AB213	TNF (seq 2)			0.124
AB217	TNF (seq 5)			0.0762
AB218	TNF (seq 6)			0.045
DVD1448	PGE2	TNF (seq 2)	-	1.388
DVD1454	SOST	TNF (seq 2)	-	0.951
DVD1466	LPA	TNF (seq 2)	-	29.92
DVD1471	TNF (seq 1)	LPA	0.0153	-
DVD1472	LPA	TNF (seq 1)	-	0.732
DVD1473	TNF (seq 5)	PGE2	0.011	-
DVD1474	PGE2	TNF (seq 5)	-	0.273
DVD1475	TNF (seq 5)	SOST	0.002	-
DVD1476	SOST	TNF (seq 5)	-	0.139
DVD1477	TNF (seq 5)	NGF	0.009	-
DVD1478	NGF	TNF (seq 5)	-	0.329
DVD1479	TNF (seq 5)	LPA	0.012	-
DVD1480	LPA	TNF (seq 5)	-	0.183
DVD1481	TNF (seq 6)	PGE2	0.011	-
DVD1482	PGE2	TNF (seq 6)	-	1.453
DVD1483	TNF (seq 6)	SOST	0.012	-
DVD1484	SOST	TNF (seq 6)	-	0.219
DVD1485	TNF (seq 6)	NGF	0.011	-
DVD1486	NGF	TNF (seq 6)	-	0.363
DVD1487	TNF (seq 6)	LPA	0.007	-
DVD1488	LPA	TNF (seq 6)	-	0.894
DVD1490	PGE2	TNF (seq 2)	-	0.283
DVD1495	TNF (seq 2)	SOST	0.0038	-
DVD1496	SOST	TNF (seq 2)	-	0.085
DVD1501	TNF (seq 2)	NGF	0.1952	-
DVD1502	NGF	TNF (seq 2)	-	0.4596
DVD1507	TNF (seq 2)	LPA	0.1327	-
DVD1508	LPA	TNF (seq 2)	-	0.234
DVD1514	LPA	TNF (seq 1)	-	0.407
DVD1516	PGE2	TNF (seq 5)	-	8.709
DVD1517	TNF (seq 5)	SOST	0.0012	-
DVD1518	SOST	TNF (seq 5)	-	0.89
DVD1519	TNF (seq 5)	NGF	0.0079	-
DVD1520	NGF	TNF (seq 5)	-	7.21

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	N-terminal VD TNF α Neutralization Assay EC50 nM	C-terminal VD TNF α Neutralization Assay EC50 nM
DVD1522	LPA	TNF (seq 5)	-	4.179
DVD1524	PGE2	TNF (seq 6)	-	0.963
DVD1525	TNF (seq 6)	SOST	0.0029	-
DVD1526	SOST	TNF (seq 6)	-	0.059
DVD1527	TNF (seq 6)	NGF	0.0029	-
DVD1528	NGF	TNF (seq 6)	-	0.197
DVD1530	LPA	TNF (seq 6)	-	0.571

All DVD-Ig proteins containing VDs from AB017, AB213, AB217, and AB218 in either the N-terminal or C-terminal position showed neutralization in the TNF α inhibition assay.

Example 1.1.2.E: Inhibition of NGF in TF-1 Cell Proliferation bioassay

TF-1 cells were cultured in RPMI 1640 (Invitrogen) +10% Fetal Bovine Serum (Hyclone) +L-glutamine (Invitrogen) + rhu GM-CSF (R&D Systems). TF-1 cells were serum starved for 24 hours in RPMI 1640 + L-glutamine at 1×10^5 cells per mL and incubated overnight at 37°C, 5% CO₂. The day of the experiment, TF-1 cells were plated in opaque walled 96-well plates at 2.5×10^4 cells per well in a 100 μ L volume + assay media (RPMI-1640 +L-glutamine + 4% FBS) The cells were stimulated by adding NGF/DVD-Ig or antibody thereto. The DVD-IgTM and control IgG were diluted to a 4x concentration in assay media and serial 1:5 dilutions were performed. The huNGF was diluted to 8 ng/mL in assay media. The DVD-IgTM (50 μ l) and huNGF (50 μ L) solutions were added to the plate for a final concentration of 2 ng/mL huNGF and 25 nM – 0.000003 nM DVD-IgTM. The plates were incubated for 72 hours at 37°C, 5 % CO₂. To quantitate viability, a Cell Titer Glo kit (Promega cat# TB288) was used (100 μ l of solution added to each well following manufacturer's instructions). The plates were read by measuring luminescence on a Spectromax 190 ELISA plate reader. An average EC50 from several assays is included in Table 6.

Table 6: NGF Inhibition Assay With NGF Parent Antibodies and DVD-Ig Constructs

Parent Antibody or DVD-Ig ID	N-Terminal Variable Domain (VD)	C-Terminal Variable Domain (VD)	N-Terminal VD NGF Inhibition Assay EC50 nM	C-Terminal VD NGF Inhibition Assay EC50 nM
AB020	NGF		0.003	
DVD1460	NGF	TNF (seq 2)	1.695	-
DVD1477	TNF (seq 5)	NGF	-	0.038
DVD1478	NGF	TNF (seq 5)	0.019	-

Parent Antibody or DVD-Ig ID	N-Terminal Variable Domain (VD)	C-Terminal Variable Domain (VD)	N-Terminal VD NGF Inhibition Assay EC50 nM	C-Terminal VD NGF Inhibition Assay EC50 nM
DVD1485	TNF (seq 6)	NGF	-	0.053
DVD1486	NGF	TNF (seq 6)	0.076	-
DVD1501	TNF (seq 2)	NGF	-	ND
DVD1502	NGF	TNF (seq 2)	0.028	-
DVD1519	TNF (seq 5)	NGF	-	1.946
DVD1520	NGF	TNF (seq 5)	0.025	-
DVD1527	TNF (seq 6)	NGF	-	0.350
DVD1528	NGF	TNF (seq 6)	0.018	-

All DVD-Ig proteins containing VDs from AB020 in either the N-terminal or C-terminal position showed neutralization in the NGF inhibition assay.

Example 1.2.2.F: Inhibition of Sclerostin Activity in the Wnt-1/Luciferase Double Stable HEK Clone #14

5 HEK 293A cells were stably transfected with TopFlash plasmid (TCF reporter plasmid, Cell Signaling catalog #21-170, lot# 26217) and infected with Wnt-1 lentivirus (Origene Cat# SC303644), resulting in clones that co-express luciferase and Wnt-1. One double stable clone (#14) has been maintained in culture medium: DMEM (Invitrogen Cat#11965-092) with 10% Qualified FBS (Invitrogen Cat#26140-079), Pen-Strep (Invitrogen Cat#15140-122), L-glutamine 10 (Invitrogen Cat#25030-081 2mM final), Sodium Pyruvate (Invitrogen Cat#11360-070 final 1mM) and 5µg/ml Puromycin (Invivogen Cat#ant-pr-1) in T75 flasks until 80-90% confluent on day of assay. Assay is performed in assay medium: culture medium without puromycin. Human Sclerostin (Abbott PR-1261069 Lot#1769536 1.06mg/mL) was aliquoted into 10µl and stored frozen at -80°C. On day 1, clone #14 cells were plated at 10,000 cells per well in 50ul assay 15 medium in black-sided, clear bottomed tissue culture treated 96 well plates (Costar #3603) and incubated at 37°C overnight (20-24 hours). On day 2, the Sclerostin stock is diluted to 200nM (4X) in the assay medium. Anti-Sclerostin antibodies were diluted to 4X (typically 600nM) in assay medium. Media was removed and replaced with 50µl/well of fresh assay medium. Cells were next incubated with 25µl of Sclerostin at 200nM (4X) for 1 hour. Anti-Sclerostin antibodies 20 (4X conc, 25µl) were then added to cells and plates were incubated overnight at 37°C (20-24 hours). The final volume was 100µl. On day 3, cells were washed once with 200µl of PBS (RT). A Promega Luciferase Kit #E1501 was used for cell lysis and Luciferase read out. Briefly, 5X cell lysis reagent (Promega, cat #E153A) was diluted with milliQ water to 1X and 20µl was added to each well. To ensure a complete lysis, plate was rotated 500rpm for 20 minutes. 100µl 25 of Luciferase assay reagent (1vial cat #E151A substrate + 10ml cat #E152A assay buffer) was

added to each well. The plate were read on a TopCount machine (Program: Luciferase 96, Assay 17: 1 sec/well read). Table 7 represents those antibodies or DVD-Igs that were able to inhibit SOST at 150 nM.

5 **Table 7: SOST Inhibition Assay With SOST Parent Antibodies and DVD-Ig Constructs**

Parent Antibody or DVD-Ig ID	N-Terminal Variable Domain (VD)	C-Terminal Variable Domain (VD)	N-Terminal VD SOST Inhibition Assay	C-Terminal VD SOST Inhibition Assay
AB022	SOST		+	
DVD145 4	TNF (seq 4)	SOST		-
DVD147 5	TNF (seq 5)	SOST		+
DVD147 6	SOST	TNF (seq 5)	+	
DVD148 3	TNF (seq 6)	SOST		+/-
DVD148 4	SOST	TNF (seq 6)	+	
DVD149 5	TNF (seq 2)	SOST		-
DVD149 6	SOST	TNF (seq 2)	+	
DVD151 7	TNF (seq 5)	SOST		-
DVD151 8	SOST	TNF (seq 5)	+	
DVD152 6	SOST	TNF (seq 6)	+	

All DVD-Igs containing VDs from AB022 in either the N-terminal or C-terminal position showed neutralization in the SOST inhibition assay.

Example 1.1.2.G: Growth Inhibitory Effect of a Tumor Receptor Monoclonal Antibody or DVD-Igs In Vitro

10 Tumor receptor monoclonal antibodies or DVD-Igs diluted in D-PBS-BSA (Dulbecco's phosphate buffered saline with 0.1%BSA) 20µL are added to human tumor cells at final concentrations of 0.01 µg/mL-100 µg/mL in 180uL. The plates are incubated at 37 °C in a humidified, 5% CO₂ atmosphere for 3 days. The number of live cells in each well is quantified using MTS reagents according to the manufacturer's instructions (Promega, Madison, WI) to
 15 determine the percent of tumor growth inhibition. Wells without antibody treatment are used as controls of 0% inhibition whereas wells without cells are considered to show 100% inhibition.

Example 1.1.2.H: Tumoricidal Effect of A Parent or DVD-Ig Antibody In Vitro

Parent antibodies or DVD-Ig that bind to target antigens on tumor cells may be analyzed for tumoricidal activity. Briefly, parent antibodies or DVD-Ig are diluted in D-PBS-BSA (Dulbecco's phosphate buffered saline with 0.1%BSA) and added to human tumor cells at final concentrations of 0.01 µg/mL to 100 µg/mL 200µL. The plates are incubated at 37 °C in a humidified, 5% CO₂ atmosphere for 3 days. The number of live cells in each well is quantified using MTS reagents according to the manufacturer's instructions (Promega, Madison, WI) to determine the percent of tumor growth inhibition. Wells without antibody treatment are used as controls of 0% inhibition whereas wells without cells were considered to show 100% inhibition.

For assessment of apoptosis, caspase-3 activation is determined by the following protocol: antibody-treated cells in 96 well plates are lysed in 120 µl of 1x lysis buffer (1.67mM Hepes, pH 7.4, 7mM KCl, 0.83mM MgCl₂, 0.11mM EDTA, 0.11mM EGTA, 0.57% CHAPS, 1mM DTT, 1x protease inhibitor cocktail tablet; EDTA-free; Roche Pharmaceuticals, Nutley, NJ) at room temperature with shaking for 20 minutes. After cell lysis, 80 µl of a caspase-3 reaction buffer (48mM Hepes, pH 7.5, 252mM sucrose, 0.1% CHAPS, 4mM DTT, and 20 µM Ac-DEVD-AMC substrate; Biomol Research Labs, Inc., Plymouth Meeting, PA) is added and the plates are incubated for 2 hours at 37°C. The plates are read on a 1420 VICTOR Multilabel Counter (Perkin Elmer Life Sciences, Downers Grove, IL) using the following settings: excitation= 360/40, emission= 460/40. An increase of fluorescence units from antibody-treated cells relative to the isotype antibody control-treated cells is indicative of apoptosis.

Example 1.1.2.I: Inhibition of Cell Proliferation by Parent Antibody and DVD-Ig Constructs

U87-MG human glioma tumor cells are plated at 2,000 cells/well in 100 µl in 96-well dishes in RPMI medium supplemented with 5% fetal bovine serum, and incubated at 37°C, 5% CO₂ overnight. The following day the cells are treated with serial dilutions of antibody or DVD-Igs (0.013 nM to 133 nM dose range), and incubated at 37 °C in a humidified, 5% CO₂ atmosphere for 5 days. Cell survival/proliferation is measured indirectly by assessing ATP levels using an ATPlite kit (Perkin Elmer, Waltham, MA) according to the manufacturer's instructions.

Example 1.1.2.J: VEGF Parent Antibody and DVD-Ig Constructs Prevent VEGF₁₆₅**30 Interaction with VEGFR1**

ELISA plates (Nunc, MaxiSorp, Rochester, NY) are incubated overnight at 4°C with 100 µl PBS containing recombinant VEGFR1 extra-cellular domain-Fc fusion protein (5µg/ml, R&D

systems, Minneapolis, MN). Plates are washed three times in washing buffer (PBS containing 0.05% Tween 20), and blocked for 1 hour at 25°C in blocking buffer (PBS containing 1% BSA). Serial dilutions of each antibody/DVD-Ig in PBS containing 0.1% BSA are incubated with 50µl of 2nM biotinylated VEGF for 1 hour at 25°C. The antibody/DVD-Ig-biotinylated VEGF mixtures (100µl) are then added to the VEGFR1-Fc coated wells and incubated at 25°C for 10 minutes. The wells are washed three times, and then incubated for 1 hour at 25°C with 100µl of streptavidin-HRP (KPL #474-3000, Gaithersburg, MD). The wells are washed three times, and 100µl of ULTRA-TMB ELISA (Pierce, Rockford, IL) are added per well. Following color development the reaction is stopped with 1N HCL and absorbance at 450nm is measured.

10 **Example 1.1.2.K: Inhibition of Receptor Phosphorylation by Parent Antibodies or DVD-Ig Constructs *In Vitro***

Human carcinoma cells are plated in 96-well plates at 40,000 cells/well in 180µl serum-free medium (DMEM+ 0.1% BSA), and incubated overnight at 37°C, 5% CO₂. Costar EIA plates (Lowell, MA) are coated with 100 µl/well of receptor capture Ab (4µg/ml final concentration), and incubated overnight at room temperature while shaking. The following day, receptor antibody-coated ELISA plates are washed (three times with PBST = 0.05% Tween 20 in PBS, pH 7.2 - 7.4), and 200µl blocking solution is added (1% BSA, 0.05% NaN₃ in PBS, pH 7.2 - 7.4.) to block for 2 hours at room temperature on a rocker. Human tumor cells are co-incubated with antibodies or DVD-Igs and ligand. Monoclonal antibodies or DVD-Igs diluted in D-PBS-BSA (Dulbecco's phosphate buffered saline with 0.1%BSA) are added to human carcinoma cells at final concentrations of 0.01 µg/mL-100 µg/mL. Growth factors are simultaneously added to the cells at concentrations of 1-100ng/mL (200µL), and cells are incubated at 37°C in a humidified, 5% CO₂ atmosphere for 1 hour. Cells are lysed in 120µl/well of cold cell extraction buffer (10 mM Tris, pH 7.4, 100 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM NaF, 1 mM sodium orthovanadate, 1% Triton X-100, 10% Glycerol, 0.1% SDS, and protease inhibitor cocktail), and incubated at 4°C for 20 minutes with shaking. Cell lysates (100µl) are added to the ELISA plate, and incubated overnight at 4°C with gentle shaking. The following day, ELISA plates are washed, and 100 µl/well of pTyr-HRP detection Ab is added (p-IGF1R ELISA kit, R&D System # DYC1770, Minneapolis, MN), and plates are incubated for 2 hours at 25°C in the dark. Plates are developed to determine phosphorylation per the manufacturer's instructions.

Example 1.1.2.L: Inhibition Of VEGFR2 (KDR) Phosphorylation By VEGF Parent Antibody And DVD-Ig Constructs

NIH3T3 cells expressing human VEGFR2 (KDR) are plated at 20,000 cells/well (100 μ l) in 96-well plates in DMEM supplemented with 10% FBS. The following day, the cells are washed twice with DMEM and serum-starved for three hours in DMEM without FBS. Anti-VEGF parent antibody or DVD-Igs (at final concentrations of 67 nM, 6.7 nM and 0.67 nM) diluted in DMEM with 0.1%BSA are pre-incubated with recombinant human VEGF₁₆₅ (50ng/ml) for 1 hour at 25°C. These antibody/DVD-Ig and VEGF mixtures are then added to the cells, and the plates are incubated at 37°C in a humidified, 5% CO₂ atmosphere for 10 minutes. Cells are washed twice with ice cold PBS and lysed by addition of 100 μ l/ well of Cell Lysis Buffer (Cell Signaling, Boston, MA) supplemented with 0.1% NP40. Duplicate samples are pooled and 170 μ l is added to wells of ELISA plates previously coated with anti-VEGFR2 antibody (R&D systems, AF357, Minneapolis, MN) and incubated at 25°C with gentle shaking for two hours. The wells are washed five times with washing buffer (PBS containing 0.05% Tween 20), and incubated with 50 μ l of of 1:2000 dilution of biotinylated anti-phosphotyrosine antibody (4G10; Millipore, Billerica, MA) for 1 hour at 25°C. The wells are washed five times with PBS containing 0.05% Tween 20, and then incubated for 1 hour at 25°C with streptavidin-HRP (KPL #474-3000, Gaithersburg, MD). The wells are washed three times with streptavidin-HRP (KPL #474-3000, Gaithersburg, MD)). The wells are washed three times with PBS containing 0.05% Tween 20, and 100 μ l of ULTRA-TMB ELISA (Pierce, Rockford, IL) are added per well. Following color development the reaction is stopped with 1N HCL and absorbance at 450nm was measured.

Example 1.1.2.M: Efficacy Of A DVD-Ig On The Growth Of Human Carcinoma Subcutaneous Flank Xenografts

A-431 human epidermoid carcinoma cells are grown *in vitro* to 99% viability, 85% confluence in tissue culture flasks. SCID female mice (Charles Rivers Labs, Wilmington, MA) at 19-25 grams are injected subcutaneously into the right flank with 1 x 10⁶ human tumor cells (1:1 matrigel) on study day 0. Administration (IP, QD, 3x/ week) of human IgG control or DVD-Ig was-initiated after mice are size matched into groups of mice with mean tumor volumes of approximately 200 to 320 mm³. The tumors are measured twice a week starting on approximately day 10 post tumor cell injection.

Example 1.1.2.N: Binding of Monoclonal Antibodies to the Surface of Human Tumor Cell Lines as Assessed by Flow Cytometry

Stable cell lines overexpressing a cell-surface antigen of interest or human tumor cell lines were harvested from tissue culture flasks and resuspended in phosphate buffered saline (PBS) containing 5% fetal bovine serum (PBS/FBS). Prior to staining, human tumor cells were incubated on ice with (100 μ l) human IgG at 5 μ g/ml in PBS/FCS. 1-5 x10⁵ cells were incubated with antibody or DVD-Ig (2 μ g/mL) in PBS/FBS for 30-60 minutes on ice. Cells were washed twice and 100 μ l of F(ab')₂ goat anti human IgG, Fcy- phycoerythrin (1:200 dilution in PBS) (Jackson ImmunoResearch, West Grove, PA, Cat.#109-116-170) was added. After 30 minutes incubation on ice, cells were washed twice and resuspended in PBS/FBS. Fluorescence was measured using a Becton Dickinson FACSCalibur (Becton Dickinson, San Jose, CA).

Example 1.1.2.O: Binding of Parent Receptor Antibody and DVD-Ig Constructs to the Surface of Human Tumor Cell Lines as Assessed by Flow Cytometry

Stable cell lines overexpressing cell-surface receptors or human tumor cell lines are harvested from tissue culture flasks and resuspended in Dulbecco's phosphate buffered saline (DPBS) containing 1% fetal calf serum (DPBS/FCS). 1-5 x10⁵ cells are incubated with 100 μ L antibodies or DVD-Igs (10 μ g/mL) in DPBS/FCS for 30-60 minutes on ice. Cells are washed twice and 50 μ l of goat anti-human IgG-phycoerythrin (1:50 dilution in DPBS/BSA) (Southern Biotech Associates, Birmingham, AL cat#2040-09) is added. After 30-45 minutes incubation on ice, cells are washed twice and resuspended in 125 μ L/well 1% formaldehyde in DPBS/FCS. Fluorescence was measured using a Becton Dickinson LSRII (Becton Dickinson, San Jose, CA).

Example 1.2: Generation Of Parent Monoclonal Antibodies to a Human Antigen of Interest

Parent mouse mAbs able to bind to and neutralize a human antigen of interest and a variant thereof are obtained as follows:

Example 1.2.A: Immunization Of Mice With a Human Antigen of Interest

Twenty micrograms of recombinant purified human antigen (e.g., IGF1,2) mixed with complete Freund's adjuvant or Immuneeasy adjuvant (Qiagen, Valencia, CA) is injected subcutaneously into five 6-8 week-old Balb/C, five C57B/6 mice, and five AJ mice on Day 1. On days 24, 38, and 49, twenty micrograms of recombinant purified human antigen variant mixed with incomplete Freund's adjuvant or Immuneeasy adjuvant is injected subcutaneously into the

same mice. On day 84 or day 112 or day 144, mice are injected intravenously with 1 µg recombinant purified human antigen of interest.

Example 1.2.B: Generation of a Hybridoma

Splenocytes obtained from the immunized mice described in Example 1.2.A are fused
5 with SP2/O-Ag-14 cells at a ratio of 5:1 according to the established method described in Kohler,
G. and Milstein (1975) Nature, 256:495 to generate hybridomas. Fusion products are plated in
selection media containing azaserine and hypoxanthine in 96-well plates at a density of 2.5×10^6
spleen cells per well. Seven to ten days post fusion, macroscopic hybridoma colonies are
observed. Supernatant from each well containing hybridoma colonies is tested by ELISA for the
10 presence of antibody to the antigen of interest (as described in Example 1.1.1.A). Supernatants
displaying antigen-specific activity are then tested for activity (as described in the assays of
Example 1.1.2), for example, the ability to neutralize the antigen of interest in a bioassay such as
that described in Example 1.1.2.I).

Example 1.2.C: Identification And Characterization Of Parent Monoclonal Antibodies to a 15 Human Target Antigen of Interest

Example 1.2.C.1: Analyzing Parent Monoclonal Antibody Neutralizing Activity

Hybridoma supernatants are assayed for the presence of parent antibodies that bind an
antigen of interest, generated according to Examples 1.2.A and 1.2.B, and are also capable of
binding a variant of the antigen of interest ("antigen variant"). Supernatants with antibodies
20 positive in both assays are then tested for their antigen neutralization potency, for example, in the
cytokine bioassay of Example 1.1.2.I. The hybridomas producing antibodies with IC_{50} values in
the bioassay less than 1000pM, in an embodiment, less than 100pM are scaled up and cloned by
limiting dilution. Hybridoma cells are expanded into media containing 10% low IgG fetal bovine
serum (Hyclone #SH30151, Logan, UT). On average, 250 mL of each hybridoma supernatant
25 (derived from a clonal population) is harvested, concentrated and purified by protein A affinity
chromatography, as described in Harlow, E. and Lane, D. 1988 "Antibodies: A Laboratory
Manual". The ability of purified mAbs to inhibit the activity of its target antigen is determined,
for example, using the cytokine bioassay as described in Example 1.1.2.I.

Example 1.2.C.2: Analyzing Parent Monoclonal Antibody Cross-Reactivity To Cynomolgus Target Antigen Of Interest

To determine whether the selected mAbs described herein recognize cynomolgus antigen of interest, BIACORE analysis is conducted as described herein (Example 1.1.1.G) using
5 recombinant cynomolgus target antigen. In addition, neutralization potencies of mAbs against recombinant cynomolgus antigen of interest may also be measured in the cytokine bioassay (Example 1.1.2.I). MAbs with good cyno cross-reactivity (in an embodiment, within 5-fold of reactivity for human antigen) are selected for future characterization.

**Example 1.2.D: Determination Of The Amino Acid Sequence Of The Variable Region For
10 Each Murine Anti-Human Monoclonal Antibody**

Isolation of the cDNAs, expression and characterization of the recombinant anti-human mouse mAbs is conducted as follows. For each amino acid sequence determination, approximately 1×10^6 hybridoma cells are isolated by centrifugation and processed to isolate total RNA with Trizol (Gibco BRL/Invitrogen, Carlsbad, CA.) following manufacturer's instructions.
15 Total RNA is subjected to first strand DNA synthesis using the SuperScript First-Strand Synthesis System (Invitrogen, Carlsbad, CA) per the manufacturer's instructions. Oligo(dT) is used to prime first-strand synthesis to select for poly(A)⁺ RNA. The first-strand cDNA product is then amplified by PCR with primers designed for amplification of murine immunoglobulin variable regions (Ig-Primer Sets, Novagen, Madison, WI). PCR products are resolved on an
20 agarose gel, excised, purified, and then subcloned with the TOPO Cloning kit into pCR2.1-TOPO vector (Invitrogen, Carlsbad, CA) and transformed into TOP10 chemically competent E. coli (Invitrogen, Carlsbad, CA). Colony PCR is performed on the transformants to identify clones containing insert. Plasmid DNA is isolated from clones containing insert using a QIAprep Miniprep kit (Qiagen, Valencia, CA). Inserts in the plasmids are sequenced on both strands to
25 determine the variable heavy or variable light chain DNA sequences using M13 forward and M13 reverse primers (Fermentas Life Sciences, Hanover MD). Variable heavy and variable light chain sequences of the mAbs are identified. In an embodiment, the selection criteria for a panel of lead mAbs for next step development (humanization) includes the following:

- The antibody does not contain any N-linked glycosylation sites (NXS), except from the
30 standard one in CH2
- The antibody does not contain any extra cysteines in addition to the normal cysteines in every antibody

- The antibody sequence is aligned with the closest human germline sequences for VH and VL and any unusual amino acids should be checked for occurrence in other natural human antibodies
- N-terminal Glutamine (Q) is changed to Glutamic acid (E) if it does not affect the activity of the antibody. This will reduce heterogeneity due to cyclization of Q
- Efficient signal sequence cleavage is confirmed by Mass Spectrophotometry. This can be done with COS cell or 293 cell material
- The protein sequence is checked for the risk of deamidation of Asn that could result in loss of activity
- The antibody has a low level of aggregation
- The antibody has solubility >5-10 mg/ml (in research phase); >25 mg/ml
- The antibody has a normal size (5-6 nm) by Dynamic Light Scattering (DLS)
- The antibody has a low charge heterogeneity
- The antibody lacks cytokine release (see Example 1.1.2.B)
- The antibody has specificity for the intended cytokine (see Example 1.1.2.C)
- The antibody lacks unexpected tissue cross reactivity (see Example 1.1.2.D)
- The antibody has similarity between human and cynomolgus tissue cross reactivity (see Example 1.1.2.D)

Example 1.2.2: Recombinant Humanized Parent Antibodies

20 Example 1.2.2.1: Construction And Expression Of Recombinant Chimeric Anti Human Parent Antibodies

The DNA encoding the heavy chain constant region of murine anti-human parent mAbs is replaced by a cDNA fragment encoding the human IgG1 constant region containing 2 hinge-region amino acid mutations by homologous recombination in bacteria. These mutations are a leucine to alanine change at position 234 (EU numbering) and a leucine to alanine change at position 235 (Lund et al., 1991, J. Immunol., 147:2657). The light chain constant region of each of these antibodies is replaced by a human kappa constant region. Full-length chimeric antibodies are transiently expressed in COS cells by co-transfection of chimeric heavy and light chain cDNAs ligated into the pBOS expression plasmid (Mizushima and Nagata, Nucleic Acids Research 1990, Vol 18, pg 5322). Cell supernatants containing recombinant chimeric antibody are purified by Protein A Sepharose chromatography and bound antibody is eluted by addition of acid buffer. Antibodies are neutralized and dialyzed into PBS.

The heavy chain cDNA encoding a chimeric mAb is co-transfected with its chimeric light chain cDNA (both ligated in the pBOS vector) into COS cells. Cell supernatant containing

recombinant chimeric antibody is purified by Protein A Sepharose chromatography and bound antibody is eluted by addition of acid buffer. Antibodies are neutralized and dialyzed into PBS.

The purified chimeric anti-human parent mAbs are then tested for their ability to bind (by Biacore) and for functional activity, e.g., to inhibit the cytokine induced production of IgE as described in Examples 1.1.1.G and 1.1.2.B. Chimeric mAbs that maintain the activity of the parent hybridoma mAbs are selected for future development.

Example 1.2.2.2: Construction And Expression Of Humanized Anti Human Parent Antibodies

Example 1.2.2.2.A: Selection Of Human Antibody Frameworks

Each murine variable heavy and variable light chain gene sequence is separately aligned against 44 human immunoglobulin germline variable heavy chain or 46 germline variable light chain sequences (derived from NCBI Ig Blast website at <http://www.ncbi.nlm.nih.gov/igblast/retrieveig.html>.) using Vector NTI software.

Humanization is based on amino acid sequence homology, CDR cluster analysis, frequency of use among expressed human antibodies, and available information on the crystal structures of human antibodies. Taking into account possible effects on antibody binding, VH-VL pairing, and other factors, murine residues are mutated to human residues where murine and human framework residues are different, with a few exceptions. Additional humanization strategies are designed based on an analysis of human germline antibody sequences, or a subgroup thereof, that possessed a high degree of homology, i.e., sequence similarity, to the actual amino acid sequence of the murine antibody variable regions.

Homology modeling is used to identify residues unique to the murine antibody sequences that are predicted to be critical to the structure of the antibody combining site, the CDRs. Homology modeling is a computational method whereby approximate three dimensional coordinates are generated for a protein. The source of initial coordinates and guidance for their further refinement is a second protein, the reference protein, for which the three dimensional coordinates are known and the sequence of which is related to the sequence of the first protein. The relationship among the sequences of the two proteins is used to generate a correspondence between the reference protein and the protein for which coordinates are desired, the target protein. The primary sequences of the reference and target proteins are aligned with coordinates of identical portions of the two proteins transferred directly from the reference protein to the target protein. Coordinates for mismatched portions of the two proteins, e.g., from residue

mutations, insertions, or deletions, are constructed from generic structural templates and energy refined to insure consistency with the already transferred model coordinates. This computational protein structure may be further refined or employed directly in modeling studies. The quality of the model structure is determined by the accuracy of the contention that the reference and target proteins are related and the precision with which the sequence alignment is constructed.

For the murine mAbs, a combination of BLAST searching and visual inspection is used to identify suitable reference structures. Sequence identity of 25% between the reference and target amino acid sequences is considered the minimum necessary to attempt a homology modeling exercise. Sequence alignments are constructed manually and model coordinates are generated with the program Jackal (see Petrey, D. et al. (2003) Proteins 53 (Suppl. 6): 430–435).

The primary sequences of the murine and human framework regions of the selected antibodies share significant identity. Residue positions that differ are candidates for inclusion of the murine residue in the humanized sequence in order to retain the observed binding potency of the murine antibody. A list of framework residues that differ between the human and murine sequences is constructed manually. Table 8 shows the framework sequences chosen for this study.

Table 8: Sequence Of Human IgG Heavy Chain Constant Domain And Light Chain Constant Domain

Protein	SEQ ID NO	Sequence
		123456789012345678901234567890123456789012345678901
Wild type hlgG1 constant region	48	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSW NSCALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYI CNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPELGGPSV FLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTIKAKGQPREPQVYITLPPSREEMTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSF FLYSKLTVDKSRWQQGNVVFSCVMHEALHNHYTQKSLSLSP GK
Mutant hlgG1 constant region	49	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYI CNVNHKPSNTKVDKKVEPKSCDKHTHTCPPCPAPEAAGGPSV FLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTIKAKGQPREPQVYITLPPSREEMTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSF FLYSKLTVDKSRWQQGNVVFSCVMHEALHNHYTQKSLST.SP GK
Ig kappa constant region	50	TVAAPSVFIFPPSDEQLKSGTASVVCLLNFFYPREAKVQWK VDNALQSGNSQESVTEQDSKSTYLSSTLTLSKADYEKHK VYACEVTHQGLSSPVTKSFNRGEC
Ig Lambda constant region	51	QPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAW KADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSRR SYSCQVTHEGSTVEKTVAPTECS

20

The likelihood that a given framework residue would impact the binding properties of the antibody depends on its proximity to the CDR residues. Therefore, using the model structures, the residues that differ between the murine and human sequences are ranked according to their distance from any atom in the CDRs. Those residues that fell within 4.5 Å of any CDR atom are identified as most important and are recommended to be candidates for retention of the murine residue in the humanized antibody (i.e., back mutation).

In silico constructed humanized antibodies are constructed using oligonucleotides. For each variable region cDNA, 6 oligonucleotides of 60-80 nucleotides each are designed to overlap each other by 20 nucleotides at the 5' and/or 3' end of each oligonucleotide. In an annealing reaction, all 6 oligonucleotides are combined, boiled, and annealed in the presence of dNTPs. DNA polymerase I, Large (Klenow) fragment (New England Biolabs #M0210, Beverly, MA.) is added to fill-in the approximately 40bp gaps between the overlapping oligonucleotides. PCR is performed to amplify the entire variable region gene using two outermost primers containing overhanging sequences complementary to the multiple cloning site in a modified pBOS vector (Mizushima, S. and Nagata, S. (1990) *Nucleic Acids Res.* 18: 17). The PCR products derived from each cDNA assembly are separated on an agarose gel and the band corresponding to the predicted variable region cDNA size is excised and purified. The variable heavy region is inserted in-frame onto a cDNA fragment encoding the human IgG1 constant region containing 2 hinge-region amino acid mutations by homologous recombination in bacteria. These mutations are a leucine to alanine change at position 234 (EU numbering) and a leucine to alanine change at position 235 (Lund et al. (1991) *J. Immunol.* 147:2657). The variable light chain region is inserted in-frame with the human kappa constant region by homologous recombination. Bacterial colonies are isolated and plasmid DNA extracted. cDNA inserts are sequenced in their entirety. Correct humanized heavy and light chains corresponding to each antibody are co-transfected into COS cells to transiently produce full-length humanized anti-human antibodies. Cell supernatants containing recombinant chimeric antibody are purified by Protein A Sepharose chromatography and bound antibody is eluted by addition of acid buffer. Antibodies are neutralized and dialyzed into PBS.

Example 1.2.2.3: Characterization Of Humanized Antibodies

The ability of purified humanized antibodies to inhibit a functional activity is determined, e.g., using the cytokine bioassay as described in Examples 1.1.2.A. The binding affinities of the humanized antibodies to recombinant human antigen are determined using surface plasmon resonance (Biacore®) measurement as described in Example 1.1.1.B. The IC₅₀ values from the bioassays and the affinity of the humanized antibodies are ranked. The

humanized mAbs that fully maintain the activity of the parent hybridoma mAbs are selected as candidates for future development. The top 2-3 most favorable humanized mAbs are further characterized.

Example 1.2.2.3.A: Pharmacokinetic Analysis Of Humanized Antibodies

5 Pharmacokinetic studies are carried out in Sprague-Dawley rats and cynomolgus monkeys. Male and female rats and cynomolgus monkeys are dosed intravenously or subcutaneously with a single dose of 4mg/kg mAb and samples are analyzed using antigen capture ELISA, and pharmacokinetic parameters are determined by noncompartmental analysis. Briefly, ELISA plates are coated with goat anti-biotin antibody (5 mg/ml, 4°C, overnight),
10 blocked with Superblock (Pierce), and incubated with biotinylated human antigen at 50 ng/ml in 10% Superblock TTBS at room temperature for 2 hours. Serum samples are serially diluted (0.5% serum, 10% Superblock in TTBS) and incubated on the plate for 30 minutes at room temperature. Detection is carried out with HRP-labeled goat anti human antibody and concentrations are determined with the help of standard curves using the four parameter logistic
15 fit. Values for the pharmacokinetic parameters are determined by non-compartmental model using WinNonlin software (Pharsight Corporation, Mountain View, CA). Humanized mAbs with good pharmacokinetics profile (T1/2 is 8-13 days or better, with low clearance and excellent bioavailability 50-100%) are selected.

Example 1.2.2.3.B: Physicochemical And In Vitro Stability Analysis Of Humanized

20 **Monoclonal Antibodies**

Size exclusion chromatography

Antibodies are diluted to 2.5 mg/mL with water and 20 mL is analyzed on a Shimadzu HPLC system using a TSK gel G3000 SWXL column (Tosoh Bioscience, cat# k5539-05k). Samples are eluted from the column with 211 mM sodium sulfate, 92 mM sodium phosphate, pH
25 7.0, at a flow rate of 0.3 mL/minutes. The HPLC system operating conditions are the following:

Mobile phase: 211 mM Na₂SO₄, 92 mM Na₂HPO₄*7H₂O, pH 7.0

Gradient: Isocratic

Flow rate: 0.3 mL/minute

Detector wavelength: 280 nm

30 Autosampler cooler temp: 4°C

Column oven temperature: Ambient

Run time: 50 minutes

Table 9 contains purity data of parent antibodies and DVD-Ig constructs expressed as percent monomer (unaggregated protein of the expected molecular weight) as determined by the above protocol.

5 **Table 9: Purity of Parent Antibodies and DVD-Ig Constructs as Determined by Size Exclusion Chromatography**

Parent Antibody or DVD-Ig ID	N-Terminal Variable Domain (VD)	C-Terminal Variable Domain (VD)	% Monomer (purity)
AB017	TNF (seq 1)		97.5
AB020	NGF		88.2
AB022	SOST		93.2
AB048	PGE2		100
AB213	TNF (seq 2)		100
AB214	TNF (seq 3)		100
AB215	TNF (seq 4)		99.9
AB216	LPA		100
AB217	TNF (seq 5)		100
AB218	TNF (seq 6)		100
DVD1448	PGE2	TNF (seq 2)	76.4
DVD1454	SOST	TNF (seq 2)	88.8
DVD1460	NGF	TNF (seq 2)	87.6
DVD1466	LPA	TNF (seq 2)	100
DVD1471	TNF (seq 1)	LPA	90.5
DVD1472	LPA	TNF (seq 1)	96.8
DVD1473	TNF (seq 5)	PGE2	94.4
DVD1474	PGE2	TNF (seq 5)	67.9
DVD1475	TNF (seq 5)	SOST	91.6
DVD1476	SOST	TNF (seq 5)	76.7
DVD1477	TNF (seq 5)	NGF	96.7
DVD1478	NGF	TNF (seq 5)	89.2
DVD1479	TNF (seq 5)	LPA	95.3
DVD1480	LPA	TNF (seq 5)	100
DVD1481	TNF (seq 6)	PGE2	97.5
DVD1482	PGE2	TNF (seq 6)	80.7
DVD1483	TNF (seq 6)	SOST	90.8
DVD1484	SOST	TNF (seq 6)	76
DVD1485	TNF (seq 6)	NGF	89.9
DVD1486	NGF	TNF (seq 6)	92.2
DVD1487	TNF (seq 6)	LPA	90.9
DVD1488	LPA	TNF (seq 6)	100
DVD1490	PGE2	TNF (seq 2)	97.2
DVD1495	TNF (seq 2)	SOST	98.8
DVD1496	SOST	TNF (seq 2)	72.8
DVD1502	NGF	TNF (seq 2)	96.2
DVD1508	LPA	TNF (seq 2)	100
DVD1513	TNF (seq 1)	LPA	73.9

Parent Antibody or DVD-Ig ID	N-Terminal Variable Domain (VD)	C-Terminal Variable Domain (VD)	% Monomer (purity)
DVD1514	LPA	TNF (seq 1)	96.3
DVD1515	TNF (seq 5)	PGE2	97.2
DVD1517	TNF (seq 5)	SOST	95.4
DVD1518	SOST	TNF (seq 5)	78.4
DVD1519	TNF (seq 5)	NGF	100
DVD1520	NGF	TNF (seq 5)	95.2
DVD1523	TNF (seq 6)	PGE2	97.5
DVD1524	PGE2	TNF (seq 6)	98.1
DVD1525	TNF (seq 6)	SOST	96.5
DVD1526	SOST	TNF (seq 6)	70.5
DVD1527	TNF (seq 6)	NGF	98.2
DVD1528	NGF	TNF (seq 6)	93.3
DVD1529	TNF (seq 6)	LPA	81.3
DVD1530	LPA	TNF (seq 6)	100

SDS-PAGE

Antibodies are analyzed by sodium dodecyl sulfate - polyacrylamide gel electrophoresis (SDS-PAGE) under both reducing and non-reducing conditions. Adalimumab lot AFP04C is used as a control. For reducing conditions, the samples are mixed 1:1 with 2X tris glycine SDS-PAGE sample buffer (Invitrogen, cat# LC2676, lot# 1323208) with 100 mM DTT, and heated at 60°C for 30 minutes. For non-reducing conditions, the samples are mixed 1:1 with sample buffer and heated at 100°C for 5 minutes. The reduced samples (10 mg per lane) are loaded on a 12% pre-cast tris-glycine gel (Invitrogen, cat# EC6005box, lot# 6111021), and the non-reduced samples (10 mg per lane) are loaded on an 8%-16% pre-cast tris-glycine gel (Invitrogen, cat# EC6045box, lot# 6111021). SeeBlue Plus 2 (Invitrogen, cat# LC5925, lot# 1351542) is used as a molecular weight marker. The gels are run in a XCell SureLock mini cell gel box (Invitrogen, cat# EI0001) and the proteins are separated by first applying a voltage of 75 to stack the samples in the gel, followed by a constant voltage of 125 until the dye front reached the bottom of the gel. The running buffer used is 1X tris glycine SDS buffer, prepared from a 10X tris glycine SDS buffer (ABC, MPS-79-080106). The gels are stained overnight with colloidal blue stain (Invitrogen cat# 46-7015, 46-7016) and destained with Milli-Q water until the background is clear. The stained gels are then scanned using an Epson Expression scanner (model 1680, S/N DASX003641).

Sedimentation Velocity Analysis

Antibodies are loaded into the sample chamber of each of three standard two-sector carbon epon centerpieces. These centerpieces have a 1.2 cm optical path length and are built with

sapphire windows. PBS is used for a reference buffer and each chamber contained 140 μ L. All samples are examined simultaneously using a 4-hole (AN-60Ti) rotor in a Beckman ProteomeLab XL-I analytical ultracentrifuge (serial # PL106C01).

Run conditions are programmed and centrifuge control is performed using ProteomeLab (v5.6). The samples and rotor are allowed to thermally equilibrate for one hour prior to analysis (20.0 \pm 0.1 $^{\circ}$ C). Confirmation of proper cell loading is performed at 3000 rpm and a single scan is recorded for each cell. The sedimentation velocity conditions are the following:

Sample Cell Volume: 420 mL
Reference Cell Volume: 420 mL
10 Temperature: 20 $^{\circ}$ C
Rotor Speed: 35,000 rpm
Time: 8:00 hours
UV Wavelength: 280 nm
Radial Step Size: 0.003 cm
15 Data Collection: One data point per step without signal averaging.
Total Number of Scans: 100

LC-MS molecular weight measurement of intact antibodies

Molecular weight of intact antibodies are analyzed by LC-MS. Each antibody is diluted to approximately 1 mg/mL with water. An 1100 HPLC (Agilent) system with a protein microtrap (20 (Michrom Bioresources, Inc, cat# 004/25109/03) is used to desalt and introduce 5 mg of the sample into an API Qstar pulsar i mass spectrometer (Applied Biosystems). A short gradient is used to elute the samples. The gradient is run with mobile phase A (0.08% FA, 0.02% TFA in HPLC water) and mobile phase B (0.08% FA and 0.02% TFA in acetonitrile) at a flow rate of 50 mL/minute. The mass spectrometer is operated at 4.5 kvolts spray voltage with a scan range from 25 2000 to 3500 mass to charge ratio.

LC-MS molecular weight measurement of antibody light and heavy chains

Molecular weight measurement of antibody light chain (LC), heavy chain (HC) and deglycosylated HC are analyzed by LC-MS. Antibody is diluted to 1 mg/mL with water and the sample is reduced to LC and HC with a final concentration of 10 mM DTT for 30 minutes at 30 37 $^{\circ}$ C. To deglycosylate the antibody, 100 mg of the antibody is incubated with 2 mL of PNGase F, 5 mL of 10% N-octylglucoside in a total volume of 100 mL overnight at 37 $^{\circ}$ C. After deglycosylation the sample is reduced with a final concentration of 10 mM DTT for 30 minutes at 37 $^{\circ}$ C. An Agilent 1100 HPLC system with a C4 column (Vydac, cat# 214TP5115, S/N

060206537204069) is used to desalt and introduce the sample (5 mg) into an API Qstar pulsar i mass spectrometer (Applied Biosystems). A short gradient is used to elute the sample. The gradient is run with mobile phase A (0.08% FA, 0.02% TFA in HPLC water) and mobile phase B (0.08% FA and 0.02% TFA in acetonitrile) at a flow rate of 50 mL/minute. The mass spectrometer is operated at 4.5 kvolts spray voltage with a scan range from 800 to 3500 mass to charge ratio.

Peptide mapping

Antibody is denatured for 15 minutes at room temperature with a final concentration of 6 M guanidine hydrochloride in 75 mM ammonium bicarbonate. The denatured samples are reduced with a final concentration of 10 mM DTT at 37°C for 60 minutes, followed by alkylation with 50 mM iodoacetic acid (IAA) in the dark at 37°C for 30 minutes. Following alkylation, the sample is dialyzed overnight against four liters of 10 mM ammonium bicarbonate at 4°C. The dialyzed sample is diluted to 1 mg/mL with 10 mM ammonium bicarbonate, pH 7.8 and 100 mg of antibody is either digested with trypsin (Promega, cat# V5111) or Lys-C (Roche, cat# 11 047 825 001) at a 1:20 (w/w) trypsin/Lys-C:antibody ratio at 37°C for 4 hrs. Digests are quenched with 1 mL of 1 N HCl. For peptide mapping with mass spectrometer detection, 40 mL of the digests are separated by reverse phase high performance liquid chromatography (RPHPLC) on a C18 column (Vydac, cat# 218TP51, S/N NE9606 10.3.5) with an Agilent 1100 HPLC system. The peptide separation is run with a gradient using mobile phase A (0.02% TFA and 0.08% FA in HPLC grade water) and mobile phase B (0.02% TFA and 0.08% FA in acetonitrile) at a flow rate of 50 mL/minutes. The API QSTAR Pulsar i mass spectrometer is operated in positive mode at 4.5 kvolts spray voltage and a scan range from 800 to 2500 mass to charge ratio.

Disulfide Bond Mapping

To denature the antibody, 100 mL of the antibody is mixed with 300 mL of 8 M guanidine HCl in 100 mM ammonium bicarbonate. The pH is checked to ensure that it is between 7 and 8 and the samples are denatured for 15 minutes at room temperature in a final concentration of 6 M guanidine HCl. A portion of the denatured sample (100 mL) is diluted to 600 mL with Milli-Q water to give a final guanidine-HCl concentration of 1 M. The sample (220 mg) is digested with either trypsin (Promega, cat # V5111, lot# 22265901) or Lys-C (Roche, cat# 11047825001, lot# 12808000) at a 1:50 trypsin or 1:50 Lys-C: antibody (w/w) ratios (4.4 mg enzyme: 220 mg sample) at 37°C for approximately 16 hours. An additional 5 mg of trypsin or Lys-C is added to the samples and digestion is allowed to proceed for an additional 2 hours at 37°C. Digestions are stopped by adding 1 mL of TFA to each sample. Digested samples are

separated by RPHPLC using a C18 column (Vydac, cat# 218TP51 S/N NE020630-4-1A) on an Agilent HPLC system. The separation is run with the same gradient used for peptide mapping using mobile phase A (0.02% TFA and 0.08% FA in HPLC grade water) and mobile phase B (0.02% TFA and 0.08% FA in acetonitrile) at a flow rate of 50 mL/minute. The HPLC operating conditions are the same as those used for peptide mapping. The API QSTAR Pulsar i mass spectrometer is operated in positive mode at 4.5 kvolts spray voltage and a scan range from 800 to 2500 mass-to-charge ratio. Disulfide bonds are assigned by matching the observed MWs of peptides with the predicted MWs of tryptic or Lys-C peptides linked by disulfide bonds.

Free sulfhydryl determination

The method used to quantify free cysteines in an antibody is based on the reaction of Ellman's reagent, 5,5'-dithio-bis(2-nitrobenzoic acid) (DTNB), with sulfhydryl groups (SH) which gives rise to a characteristic chromophoric product, 5-thio-(2-nitrobenzoic acid) (TNB). The reaction is illustrated in the formula:



The absorbance of the TNB⁻ is measured at 412 nm using a Cary 50 spectrophotometer. An absorbance curve is plotted using dilutions of 2-mercaptoethanol (b-ME) as the free SH standard and the concentrations of the free sulfhydryl groups in the protein are determined from absorbance at 412 nm of the sample.

The b-ME standard stock is prepared by a serial dilution of 14.2 M b-ME with HPLC grade water to a final concentration of 0.142 mM. Then standards in triplicate for each concentration are prepared. Antibody is concentrated to 10 mg/mL using an amicon ultra 10,000 MWCO centrifugal filter (Millipore, cat# UFC801096, lot# L3KN5251) and the buffer is changed to the formulation buffer used for adalimumab (5.57 mM sodium phosphate monobasic, 8.69 mM sodium phosphate dibasic, 106.69 mM NaCl, 1.07 mM sodium citrate, 6.45 mM citric acid, 66.68 mM mannitol, pH 5.2, 0.1% (w/v) Tween). The samples are mixed on a shaker at room temperature for 20 minutes. Then 180 mL of 100 mM Tris buffer, pH 8.1 is added to each sample and standard followed by the addition of 300 mL of 2 mM DTNB in 10 mM phosphate buffer, pH 8.1. After thorough mixing, the samples and standards are measured for absorption at 412 nm on a Cary 50 spectrophotometer. The standard curve is obtained by plotting the amount of free SH and OD₄₁₂ nm of the b-ME standards. Free SH content of samples are calculated based on this curve after subtraction of the blank.

Weak Cation Exchange Chromatography

Antibody is diluted to 1 mg/mL with 10 mM sodium phosphate, pH 6.0. Charge heterogeneity is analyzed using a Shimadzu HPLC system with a WCX-10 ProPac analytical column (Dionex, cat# 054993, S/N 02722). The samples are loaded on the column in 80% mobile phase A (10 mM sodium phosphate, pH 6.0) and 20% mobile phase B (10 mM sodium phosphate, 500 mM NaCl, pH 6.0) and eluted at a flow rate of 1.0 mL/minute.

Oligosaccharide Profiling

Oligosaccharides released after PNGase F treatment of antibody are derivatized with 2-aminobenzamide (2-AB) labeling reagent. The fluorescent-labeled oligosaccharides are separated by normal phase high performance liquid chromatography (NPHPLC) and the different forms of oligosaccharides are characterized based on retention time comparison with known standards.

The antibody is first digested with PNGaseF to cleave N-linked oligosaccharides from the Fc portion of the heavy chain. The antibody (200 mg) is placed in a 500 mL Eppendorf tube along with 2 mL PNGase F and 3 mL of 10% N-octylglucoside. Phosphate buffered saline is added to bring the final volume to 60 mL. The sample is incubated overnight at 37°C in an Eppendorf thermomixer set at 700 RPM. Adalimumab lot AFP04C is also digested with PNGase F as a control.

After PNGase F treatment, the samples are incubated at 95°C for 5 minutes in an Eppendorf thermomixer set at 750 RPM to precipitate out the proteins, then the samples are placed in an Eppendorf centrifuge for 2 minutes at 10,000 RPM to spin down the precipitated proteins. The supernatant containing the oligosaccharides are transferred to a 500 mL Eppendorf tube and dried in a speed-vac at 65°C.

The oligosaccharides are labeled with 2AB using a 2AB labeling kit purchased from Prozyme (cat# GKK-404, lot# 132026). The labeling reagent is prepared according to the manufacturer's instructions. Acetic acid (150 mL, provided in kit) is added to the DMSO vial (provided in kit) and mixed by pipeting the solution up and down several times. The acetic acid/DMSO mixture (100 mL) is transferred to a vial of 2-AB dye (just prior to use) and mixed until the dye is fully dissolved. The dye solution is then added to a vial of reductant (provided in kit) and mixed well (labeling reagent). The labeling reagent (5 mL) is added to each dried oligosaccharide sample vial, and mixed thoroughly. The reaction vials are placed in an Eppendorf thermomixer set at 65°C and 700-800 RPM for 2 hours of reaction.

After the labeling reaction, the excess fluorescent dye is removed using GlycoClean S Cartridges from Prozyme (cat# GKI-4726). Prior to adding the samples, the cartridges are washed with 1 mL of milli-Q water followed with 5 washes of 1 mL 30% acetic acid solution. Just prior to adding the samples, 1 mL of acetonitrile (Burdick and Jackson, cat# AH015-4) is added to the cartridges.

After all of the acetonitrile passed through the cartridge, the sample is spotted onto the center of the freshly washed disc and allowed to adsorb onto the disc for 10 minutes. The disc is washed with 1 mL of acetonitrile followed by five washes of 1 mL of 96% acetonitrile. The cartridges are placed over a 1.5 mL Eppendorf tube and the 2-AB labeled oligosaccharides are eluted with 3 washes (400 μ L each wash) of milli Q water.

The oligosaccharides are separated using a Glycosep N HPLC (cat# GKI-4728) column connected to a Shimadzu HPLC system. The Shimadzu HPLC system consisted of a system controller, degasser, binary pumps, autosampler with a sample cooler, and a fluorescent detector.

Stability at Elevated Temperatures

The buffer of antibody is either 5.57 mM sodium phosphate monobasic, 8.69 mM sodium phosphate dibasic, 106.69 mM NaCl, 1.07 mM sodium citrate, 6.45 mM citric acid, 66.68 mM mannitol, 0.1% (w/v) Tween, pH 5.2; or 10 mM histidine, 10 mM methionine, 4% mannitol, pH 5.9 using Amicon ultra centrifugal filters. The final concentration of the antibodies is adjusted to 2 mg/mL with the appropriate buffers. The antibody solutions are then filter sterilized and 0.25 mL aliquots are prepared under sterile conditions. The aliquots are left at either -80°C, 5°C, 25°C, or 40°C for 1, 2 or 3 weeks. At the end of the incubation period, the samples are analyzed by size exclusion chromatography and SDS-PAGE.

The stability samples are analyzed by SDS-PAGE under both reducing and non-reducing conditions. The procedure used is the same as described herein. The gels are stained overnight with colloidal blue stain (Invitrogen cat# 46-7015, 46-7016) and destained with Milli-Q water until the background is clear. The stained gels are then scanned using an Epson Expression scanner (model 1680, S/N DASX003641). To obtain more sensitivity, the same gels are silver stained using silver staining kit (Owl Scientific) and the recommended procedures given by the manufacturer is used.

Dynamic Scanning Fluorimetry

The DVD-Igs were dialysed in 10mM citrate 10mM phosphate buffer, pH 6.0 to get a final concentration of 1 mg/ml. Triplicates were run for each DVD-Ig. For each sample, 27 μ l of

the DVD was added in a well of a 96 well plate and mixed with 3 μ l of 4X diluted SYPRO Orange dye (Invitrogen). The dye was supplied in DMSO at a concentration of 5000X and was diluted to the working concentration of 4X in water. The plate was centrifuged for 30 seconds to ensure that both the dye and the protein settled to the bottom of the wells and complete mixing was ensured by gentle aspiration by a pipette tip. The plate was then sealed with an adhesive film.

A real time PCR (Applied Biosciences, 7500 Series) was used to measure the change in fluorescence intensities with temperature. The plate was heated from 25° C to 95° C at a temperature ramp rate of approximately 0.5°C/minute and emission fluorescence was collected using TAMRA filter. The data was exported to Microsoft Excel and plotted as temperature vs fluorescence for each DVD-Ig. Onset of melting was noted as the temperature where the thermogram rises above the baseline fluorescence. SYPRO Orange is a hydrophobic dye and preferentially binds to the exposed hydrophobic residues in an unfolded protein molecule. Hence the onset of unfolding temperature, as measured by an increase in fluorescence is an indication of the thermal stability of the DVD-Ig. The unfolding temperature for the DVD-Igs can be found in Table 10.

Table 10: Thermal Stability of Parent Antibodies and CDR-grafted DVD-Ig Constructs as Determined by Dynamic Scanning Fluorimetry

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	Onset of melting (rank) (deg C)
DVD1244	TNF (seq 1)	LPA	51.3
DVD1248	TNF (seq 5)	SOST	54.0
DVD1249	SOST	TNF (seq 5)	54.3
DVD1256	TNF (seq 6)	SOST	56
DVD1257	SOST	TNF (seq 6)	57
DVD1258	TNF (seq 6)	NGF	59.5
DVD1259	NGF	TNF (seq 6)	51.7
DVD1261	LPA	TNF (seq 6)	54.7
DVD1475	TNF (seq 5)	SOST	49.0
DVD1476	SOST	TNF (seq 5)	53.0
DVD1483	TNF (seq 6)	SOST	52.7
DVD1484	SOST	TNF (seq 6)	49.3
DVD1485	TNF (seq 6)	NGF	54.7
DVD1488	LPA	TNF (seq 6)	52.3
DVD1513	TNF (seq 1)	LPA	50.7
DVD1518	SOST	TNF (seq 5)	52.7
DVD1525	TNF (seq 6)	SOST	55.3
DVD1526	SOST	TNF (seq 6)	52.7
DVD1527	TNF (seq 6)	NGF	55.7

DVD1528	NGF	TNF (seq 6)	47.3
---------	-----	-------------	------

Most DVD-Igs showed an unfolding temperature >50. This DVD-Ig profile is similar to that observed for parent antibodies.

Solubility determination

- 5 DVD-Ig candidates were dialyzed in 15mM His, pH 6.0 followed by concentrating them up to 50 µl in centricons with a 30K cutoff. Solubility was visually confirmed by absence of precipitation after storage at 4°C and quantitatively determined by UV absorbance measurement at 280nm.

Table 11: Solubility of DVD-Ig Constructs

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	Solubility(rank) 15mM His, pH 6	
			Appearance	mg/ml
DVD1244	TNF (seq 1)	LPA	clear	>62
DVD1248	TNF (seq 5)	SOST	clear	>103
DVD1249	SOST	TNF (seq 5)	clear	>115
DVD1256	TNF (seq 6)	SOST	clear	>95
DVD1257	SOST	TNF (seq 6)	clear	>80
DVD1258	TNF (seq 6)	NGF	clear	>125
DVD1259	NGF	TNF (seq 6)	clear	>92
DVD1261	LPA	TNF (seq 6)	clear	>52
DVD1475	TNF (seq 5)	SOST	ppt	84
DVD1476	SOST	TNF (seq 5)	clear	>78
DVD1483	TNF (seq 6)	SOST	clear	>97
DVD1484	SOST	TNF (seq 6)	clear	>78
DVD1485	TNF (seq 6)	NGF	ppt	26
DVD1488	LPA	TNF (seq 6)	clear	>27
DVD1513	TNF (seq 1)	LPA	clear	>81
DVD1518	SOST	TNF (seq 5)	clear	>21
DVD1525	TNF (seq 6)	SOST	clear	>107
DVD1526	SOST	TNF (seq 6)	ppt	32
DVD1527	TNF (seq 6)	NGF	ppt	51
DVD1528	NGF	TNF (seq 6)	ppt	59

- 10 Most DVD-Igs showed clear appearance and could be concentrated to greater than 25 mg/ml. This DVD-Ig profile is similar to that observed for parent antibodies.

Example 1.2.2.3.C: Efficacy Of A Humanized Monoclonal Antibody By Itself Or In Combination With Chemotherapy On The Growth Of Human Carcinoma Xenografts

- 15 Human cancer cells are grown *in vitro* to 99% viability, 85% confluence in tissue culture flasks. SCID female or male mice (Charles Rivers Labs) at 19-25 grams, are ear tagged and

shaved. Mice are then inoculated subcutaneously into the right flank with 0.2 ml of 2×10^6 human tumor cells (1:1 matrigel) on study day 0. Administration (IP, Q3D/ week) of vehicle (PBS), humanized antibody, and/or chemotherapy is initiated after mice are size matched into separate cages of mice with mean tumor volumes of approximately 150 to 200 mm³. The tumors are measured by a pair of calipers twice a week starting on approximately day 10 post inoculation and the tumor volumes calculated according to the formula $V = L \times W^2/2$ (V: volume, mm³; L: length, mm; W: width, mm). Reduction in tumor volume is seen in animals treated with mAb alone or in combination with chemotherapy relative to tumors in animals that received only vehicle or an isotype control mAb.

10 **Example 1.2.2.3.D: FACS Based Redirected Cytotoxicity (rCTL) Assay**

Human CD3+ T cells were isolated from previously frozen isolated peripheral blood mononuclear cells (PBMC) by a negative selection enrichment column (R&D Systems, Minneapolis, MN; Cat.#HTCC-525). T cells were stimulated for 4 days in flasks (vent cap, Corning, Acton, MA) coated with 10µg/mL anti-CD3 (OKT-3, eBioscience, Inc., San Diego, CA) and 2µg/mL anti-CD28 (CD28.2, eBioscience, Inc., San Diego, CA) in D-PBS (Invitrogen, Carlsbad, CA) and cultured in 30U/mL IL-2 (Roche) in complete RPMI 1640 media (Invitrogen, Carlsbad, CA) with L-glutamine, 55mM β-ME, Pen/Strep, 10% FBS). T cells were then rested overnight in 30U/mL IL-2 before using in assay. DoHH2 or Raji target cells were labeled with PKH26 (Sigma-Aldrich, St. Louis, MO) according to manufacturer's instructions. RPMI 1640 media (no phenol, Invitrogen, Carlsbad, CA) containing L-glutamine and 10% FBS (Hyclone, Logan, UT) was used throughout the rCTL assay. (See Dreier *et al.* (2002) Int J Cancer 100:690).

Effector T cells (E) and targets (T) were plated at a final cell concentration of 10^5 and 10^4 cells/well in 96-well plates (Costar #3799, Acton, MA), respectively to give an E:T ratio of 10:1. DVD-Ig molecules were diluted to obtain concentration-dependent titration curves. After an overnight incubation cells are pelleted and washed with D-PBS once before resuspending in FACS buffer containing 0.1% BSA (Invitrogen, Carlsbad, CA), 0.1% sodium azide and 0.5µg/mL propidium iodide (BD) in D-PBS. FACS data was collected on a FACS Canto II machine (Becton Dickinson, San Jose, CA) and analyzed in Flowjo (Treestar). The percent live targets in the DVD-Ig treated samples divided by the percent total targets (control, no treatment) was calculated to determine percent specific lysis. IC50s were calculated in Prism (Graphpad).

A CD3 / CD20 DVD-Ig was tested for redirected toxicity and showed in vitro tumor killing with an IC50 = 325pM. The sequence of this CD3 / CD20 DVD-Ig was disclosed in US Patent Application Serial No. 20070071675.

Example 1.4: Generation of a DVD-Ig

DVD-Ig molecules capable of binding two antigens are constructed using two parent monoclonal antibodies, one against human antigen A, and the other against human antigen B, selected as described herein.

5 Example 1.4.1: Generation Of A DVD-Ig Having Two Linker Lengths

A constant region containing $\mu 1$ Fc with mutations at 234, and 235 to eliminate ADCC/CDC effector functions is used. Four different anti-A/B DVD-Ig constructs are generated: 2 with short linker and 2 with long linker, each in two different domain orientations: V_A-V_B-C and V_B-V_A-C (see Table 11). The linker sequences, derived from the N-terminal sequence of
 10 human C1/Ck or CH1 domain, are as follows:

For DVDAB constructs:

light chain (if anti-A has λ): Short linker: QPKAAP (SEQ ID NO: 15); Long linker:
 QPKAAPSVTLFPP (SEQ ID NO: 16)

light chain (if anti-A has κ): Short linker: TVAAP (SEQ ID NO: 13); Long linker:
 15 TVAAPSVFIFPP (SEQ ID NO: 14)

heavy chain ($\gamma 1$): Short linker: ASTKGP (SEQ ID NO: 21); Long linker:
 ASTKGPSVFPLAP (SEQ ID NO: 22)

For DVDBA constructs:

light chain (if anti-B has λ): Short linker: QPKAAP (SEQ ID NO: 15); Long linker:
 20 QPKAAPSVTLFPP (SEQ ID NO: 16)

light chain (if anti-B has κ): Short linker: TVAAP (SEQ ID NO: 13); Long linker:
 TVAAPSVFIFPP (SEQ ID NO: 14)

heavy chain ($\gamma 1$): Short linker: ASTKGP (SEQ ID NO: 21); Long linker:
 ASTKGPSVFPLAP (SEQ ID NO: 22)

25 Heavy and light chain constructs are subcloned into the pBOS expression vector, and expressed in COS cells, followed by purification by Protein A chromatography. The purified materials are subjected to SDS-PAGE and SEC analysis.

Table 12 describes the heavy chain and light chain constructs used to express each anti-A/B DVD-Ig protein.

Table 12: Anti-A/B DVD-Ig Constructs

DVD-Ig protein	Heavy chain construct	Light chain construct
DVDABSL	DVDABHC-SL	DVDABLC-SL
DVDABLL	DVDABHC-LL	DVDABLC-LL
DVDBASL	DVDBAHC-SL	DVDBALC-SL
DVDBALL	DVDBAHC-LL	DVDBALC-LL

Example 1.4.2: Molecular cloning of DNA constructs for DVDABSL and DVDABLL

To generate heavy chain constructs DVDABHC-LL and DVDABHC-SL, VH domain of A antibody is PCR amplified using specific primers (3' primers contain short/long liner sequence for SL/LL constructs, respectively); meanwhile VH domain of B antibody is amplified using specific primers (5' primers contains short/long liner sequence for SL/LL constructs, respectively). Both PCR reactions are performed according to standard PCR techniques and procedures. The two PCR products are gel-purified, and used together as overlapping template for the subsequent overlapping PCR reaction. The overlapping PCR products are subcloned into Srf I and Sal I double digested pBOS-hC γ 1,z non-a mammalian expression vector (Abbott) by using standard homologous recombination approach.

To generate light chain constructs DVDABLC-LL and DVDABLC-SL, VL domain of A antibody is PCR amplified using specific primers (3' primers contain short/long liner sequence for SL/LL constructs, respectively); meanwhile VL domain of B antibody is amplified using specific primers (5' primers contains short/long liner sequence for SL/LL constructs, respectively). Both PCR reactions are performed according to standard PCR techniques and procedures. The two PCR products are gel-purified, and used together as overlapping template for the subsequent overlapping PCR reaction using standard PCR conditions. The overlapping PCR products are subcloned into Srf I and Not I double digested pBOS-hC κ mammalian expression vector (Abbott) by using standard homologous recombination approach. Similar approach has been used to generate DVDBASL and DVDBALL as described below:

Example 1.4.3: Molecular cloning of DNA constructs for DVDBASL and DVDBALL

To generate heavy chain constructs DVDBAHC-LL and DVDBAHC-SL, VH domain of antibody B is PCR amplified using specific primers (3' primers contain short/long liner sequence

for SL/LL constructs, respectively); meanwhile VH domain of antibody A is amplified using specific primers (5' primers contains short/long liner sequence for SL/LL constructs, respectively). Both PCR reactions are performed according to standard PCR techniques and procedures. The two PCR products are gel-purified, and used together as overlapping template
5 for the subsequent overlapping PCR reaction using standard PCR conditions. The overlapping PCR products are subcloned into Srf I and Sal I double digested pBOS-hC γ 1,z non-a mammalian expression vector (Abbott) by using standard homologous recombination approach.

To generate light chain constructs DVDBALC-LL and DVDBALC-SL, VL domain of antibody B is PCR amplified using specific primers (3' primers contain short/long liner sequence
10 for SL/LL constructs, respectively); meanwhile VL domain of antibody A is amplified using specific primers (5' primers contains short/long liner sequence for SL/LL constructs, respectively). Both PCR reactions are performed according to standard PCR techniques and procedures. The two PCR products are gel-purified, and used together as overlapping template for the subsequent overlapping PCR reaction using standard PCR conditions. The overlapping
15 PCR products are subcloned into Srf I and Not I double digested pBOS-hCk mammalian expression vector (Abbott) by using standard homologous recombination approach.

Example 1.4.4: Construction and Expression of Additional DVD-Ig

Example 1.4.4.1: Preparation of DVD-Ig vector constructs

Parent antibody amino acid sequences for specific antibodies, which recognize specific
20 antigens or epitopes thereof, for incorporation into a DVD-Ig can be obtained by preparation of hybridomas as described above or can be obtained by sequencing known antibody proteins or nucleic acids. In addition, known sequences can be obtained from the literature. The sequences can be used to synthesize nucleic acids using standard DNA synthesis or amplification technologies and assembling the desired antibody fragments into expression vectors, using
25 standard recombinant DNA technology, for expression in cells.

For example, nucleic acid codons were determined from amino acids sequences and oligonucleotide DNA was synthesized by Blue Heron Biotechnology, Inc. (www.blueheronbio.com) Bothell, WA USA. The oligonucleotides were assembled into 300-2,000 base pair double-stranded DNA fragments, cloned into a plasmid vector and sequence-
30 verified. Cloned fragments were assembled using an enzymatic process to yield the complete gene and subcloned into an expression vector. (See 7,306,914; 7,297,541; 7,279,159; 7,150,969; 20080115243; 20080102475; 20080081379; 20080075690; 20080063780; 20080050506; 20080038777; 20080022422; 20070289033; 20070287170; 20070254338; 20070243194;

20070225227; 20070207171; 20070150976; 20070135620; 20070128190; 20070104722;
20070092484; 20070037196; 20070028321; 20060172404; 20060162026; 20060153791;
20030215458; 20030157643).

A group of pHybE vectors (US Patent Application Serial No. 61/021,282) were used for
parental antibody and DVD-Ig cloning. V1, derived from pJP183; pHybE-hCg1,z,non-a V2, was
used for cloning of antibody and DVD heavy chains with a wildtype constant region. V2, derived
from pJP191; pHybE-hCk V2, was used for cloning of antibody and DVD light chains with a
kappa constant region. V3, derived from pJP192; pHybE-hCl V2, was used for cloning of
antibody and DVDs light chains with a lambda constant region. V4, built with a lambda signal
peptide and a kappa constant region, was used for cloning of DVD light chains with a lambda-
kappa hybrid V domain. V5, built with a kappa signal peptide and a lambda constant region, was
used for cloning of DVD light chains with a kappa-lambda hybrid V domain. V7, derived from
pJP183; pHybE-hCg1,z,non-a V2, was used for cloning of antibody and DVD heavy chains with
a (234,235 AA) mutant constant region.

Referring to Table 13, a number of vectors were used in the cloning of the parent
antibodies and DVD-Ig VH and VL chains.

Table 13: Vectors Used to Clone Parent Antibodies and DVD-Igs

ID	Heavy chain vector	Light chain vector
DVD1447	V1	V2
DVD1448	V1	V2
DVD1449	V1	V2
DVD1450	V1	V2
DVD1451	V1	V2
DVD1452	V1	V2
DVD1453	V1	V2
DVD1454	V1	V2
DVD1455	V1	V2
DVD1456	V1	V2
DVD1457	V1	V2
DVD1468	V1	V2
DVD1459	V1	V2
DVD1460	V1	V2
DVD1461	V1	V2
DVD1462	V1	V2
DVD1463	V1	V2
DVD1464	V1	V2
DVD1465	V1	V2
DVD1466	V1	V2
DVD1467	V1	V2

ID	Heavy chain vector	Light chain vector
DVD1468	V1	V2
DVD1469	V1	V2
DVD1470	V1	V2
DVD1471	V1	V2
DVD1472	V1	V2
DVD1473	V1	V2
DVD1474	V1	V2
DVD1475	V1	V2
DVD1476	V1	V2
DVD1477	V1	V2
DVD1478	V1	V2
DVD1479	V1	V2
DVD1480	V1	V2
DVD1481	V1	V2
DVD1482	V1	V2
DVD1483	V1	V2
DVD1484	V1	V2
DVD1485	V1	V2
DVD1486	V1	V2
DVD1487	V1	V2
DVD1488	V1	V2
DVD1489	V1	V2
DVD1490	V1	V2
DVD1491	V1	V2
DVD1492	V1	V2
DVD1493	V1	V2
DVD1494	V1	V2
DVD1495	V1	V2
DVD1496	V1	V2
DVD1497	V1	V2
DVD1498	V1	V2
DVD1499	V1	V2
DVD1500	V1	V2
DVD1501	V1	V2
DVD1502	V1	V2
DVD1503	V1	V2
DVD1504	V1	V2
DVD1505	V1	V2
DVD1506	V1	V2
DVD1507	V1	V2
DVD1508	V1	V2
DVD1509	V1	V2
DVD1510	V1	V2
DVD1511	V1	V2
DVD1512	V1	V2
DVD1513	V1	V2

ID	Heavy chain vector	Light chain vector
DVD1514	V1	V2
DVD1515	V1	V2
DVD1516	V1	V2
DVD1517	V1	V2
DVD1518	V1	V2
DVD1519	V1	V2
DVD1520	V1	V2
DVD1521	V1	V2
DVD1522	V1	V2
DVD1523	V1	V2
DVD1524	V1	V2
DVD1525	V1	V2
DVD1526	V1	V2
DVD1527	V1	V2
DVD1528	V1	V2
DVD1529	V1	V2
DVD1530	V1	V2
DVD1531	V1	V2
DVD1532	V1	V2
DVD1533	V1	V2
DVD1534	V1	V2
DVD1535	V1	V2
DVD1536	V1	V2
DVD1537	V1	V2
DVD1538	V1	V2
DVD1539	V1	V2
DVD1540	V1	V2
DVD1541	V1	V2
DVD1542	V1	V2
DVD1543	V1	V2
DVD1544	V1	V2
DVD1545	V1	V2
DVD1546	V1	V2
DVD1547	V1	V2
DVD1548	V1	V2
DVD1549	V1	V2
DVD1550	V1	V2
DVD1551	V1	V2
DVD1552	V1	V2
DVD1553	V1	V2
DVD1554	V1	V2
DVD1555	V1	V2
DVD1556	V1	V2
DVD1557	V1	V2
DVD1558	V1	V2
DVD1559	V1	V2

ID	Heavy chain vector	Light chain vector
DVD1560	V1	V2
DVD1561	V1	V2
DVD1562	V1	V2
DVD1563	V1	V2
DVD1564	V1	V2
DVD1565	V1	V2
DVD1566	V1	V2
DVD1567	V1	V2
DVD1568	V1	V2
DVD1569	V1	V2
DVD1570	V1	V2
DVD1571	V1	V2
DVD1572	V1	V2

Example 1.4.4.2: Transfection And Expression In 293 Cells

Expression of the reference antibodies and DVD-Igs was accomplished by transiently cotransfecting HEK293 (EBNA) cells with plasmids containing the corresponding light-chain (LC) and heavy-chain (HC) nucleic acids. HEK293 (EBNA) cells were propagated in Freestyle 293 media (Invitrogen, Carlsbad CA) at a 0.5L-scale in flasks (2L Corning Cat# 431198) shaking in a CO₂ incubator (8% CO₂, 125 RPM, 37°C). When the cultures reached a density of 1x10⁶ cells/ml, cells were transfected with transfection complex. Transfection complex was prepared by first mixing 150µg LC-plasmid and 100µg HC-plasmid together in 25ml of Freestyle media, followed by the addition of 500ul PEI stock solution [stock solution: 1 mg/ml (pH 7.0) Linear 25kDa PEI, Polysciences Cat# 23966]. The transfection complex was mixed by inversion and allowed to incubate at room temperature for 10 minutes prior to being added to the cell culture. Following transfection, cultures continued to be grown in the CO₂ incubator (8% CO₂, 125 RPM, 37°C). Twenty-four hours after transfection, the culture was supplemented with 25ml of a 10% Tryptone N1 solution (Organo Technie, La Courneuve France Cat# 19553). Nine days after transfection, cells were removed from the cultures by centrifugation (16,000 g, 10 minutes), and the retained supernatant was sterile filtered (Millipore HV Durapore Stericup, 0.45µm) and placed at 4°C until initiation of the purification step.

Each antibody or DVD-Ig was individually purified using a disposable 1ml packed column (packed by Orochem Technologies) containing MabSelect SuRe resin (GE Healthcare). Columns were pre-equilibrated in PBS and then loaded with the harvested 0.55L samples overnight (15 hours) at 1 ml/minute with the flow-through being recirculated back into the feed container. Following the loading step, columns were washed with 20ml PBS and protein was eluted by feeding elution buffer [50mM Citric acid pH 3.5] at 4 ml/min and collecting fractions (1 ml) in tubes already containing 0.2ml of 1.5M Tris pH 8.2 (bringing the final pH to

approximately 6.0). Fractions containing antibody were pooled based on the chromatograms and dialyzed into the final storage buffer [10mM citric acid, 10mM Na₂HPO₄, pH 6.0]. Following dialysis, samples were filtered through a 0.22um Steriflip (Millipore) and the protein concentration was determined by absorbance [Hewlett Packard 8453 diode array spectrophotometer]. SDS-PAGE analysis was performed on analytical samples (both reduced and non-reduced) to assess final purity, verify the presence of appropriately sized heavy- and light-chain bands, and confirm the absence of significant amounts of free (e.g., uncomplexed) light chain (in the non-reduced samples).

Table 14 contains the yield data for parent antibodies or DVD-Ig constructs expressed as milligrams per liter in 293 cells.

Table 14: Transient Expression in Yields of Parent Antibodies and DVD-Ig Constructs in 293 Cells

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	Expression Yield (mg/L)
DVD1447	TNF (seq 2)	PGE2	0.014
DVD1448	PGE2	TNF (seq 2)	1.06
DVD1453	TNF (seq 2)	SOST	0.05
DVD1454	SOST	TNF (seq 2)	0.184
DVD1459	TNF (seq 2)	NGF	0
DVD1460	NGF	TNF (seq 2)	0.24
DVD1465	TNF (seq 2)	LPA	0
DVD1466	LPA	TNF (seq 2)	0
DVD1471	TNF (seq 1)	LPA	0.34
DVD1472	LPA	TNF (seq 1)	0.66
DVD1473	TNF (seq 5)	PGE2	0.14
DVD1474	PGE2	TNF (seq 5)	2
DVD1475	TNF (seq 5)	SOST	21.4
DVD1476	SOST	TNF (seq 5)	18.2
DVD1477	TNF (seq 5)	NGF	3.6
DVD1478	NGF	TNF (seq 5)	1.32
DVD1479	TNF (seq 5)	LPA	1.6
DVD1480	LPA	TNF (seq 5)	6
DVD1481	TNF (seq 6)	PGE2	3
DVD1482	PGE2	TNF (seq 6)	5.6
DVD1483	TNF (seq 6)	SOST	21.8
DVD1484	SOST	TNF (seq 6)	19.8
DVD1485	TNF (seq 6)	NGF	8.4
DVD1486	NGF	TNF (seq 6)	2.1
DVD1487	TNF (seq 6)	LPA	2.6

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	Expression Yield (mg/L)
DVD1488	LPA	TNF (seq 6)	8.6
DVD1489	TNF (seq 2)	PGE2	0
DVD1490	PGE2	TNF (seq 2)	0.6
DVD1495	TNF (seq 2)	SOST	2.2
DVD1496	SOST	TNF (seq 2)	5
DVD1501	TNF (seq 2)	NGF	0.056
DVD1502	NGF	TNF (seq 2)	2.4
DVD1507	TNF (seq 2)	LPA	0.05
DVD1508	LPA	TNF (seq 2)	3.28
DVD1513	TNF (seq 1)	LPA	17.72
DVD1514	LPA	TNF (seq 1)	0.8
DVD1515	TNF (seq 5)	PGE2	1
DVD1516	PGE2	TNF (seq 5)	4.98
DVD1517	TNF (seq 5)	SOST	3.54
DVD1518	SOST	TNF (seq 5)	8.78
DVD1519	TNF (seq 5)	NGF	0.4
DVD1520	NGF	TNF (seq 5)	0.86
DVD1521	TNF (seq 5)	LPA	0
DVD1522	LPA	TNF (seq 5)	1.34
DVD1523	TNF (seq 6)	PGE2	2
DVD1524	PGE2	TNF (seq 6)	2.56
DVD1525	TNF (seq 6)	SOST	23.8
DVD1526	SOST	TNF (seq 6)	14.6
DVD1527	TNF (seq 6)	NGF	11
DVD1528	NGF	TNF (seq 6)	11.6
DVD1529	TNF (seq 6)	LPA	5.2
DVD1530	LPA	TNF (seq 6)	5.2

Example 1.4.5: Characterization and Lead Selection of A/B DVD-Igs

The binding affinities of anti-A/B DVD-Igs are analyzed on Biacore against both protein A and protein B. The tetravalent property of the DVD-Ig is examined by multiple binding studies on Biacore. Meanwhile, the neutralization potency of the DVD-Igs for protein A and protein B are assessed by bioassays, respectively, as described herein. The DVD-Ig molecules that best retain the affinity and potency of the original parent mAbs are selected for in-depth physicochemical and bio-analytical (rat PK) characterizations as described herein for each mAb. Based on the collection of analyses, the final lead DVD-Ig is advanced into CHO stable cell line development, and the CHO-derived material is employed in stability, pharmacokinetic and efficacy studies in cynomolgus monkey, and preformulation activities.

Example 2: Generation and Characterization of Dual Variable Domain Immunoglobulins (DVD-Ig)

Dual variable domain immunoglobulins (DVD-Ig) using parent antibodies with known amino acid sequences were generated by synthesizing polynucleotide fragments encoding DVD-Ig variable heavy and DVD-Ig variable light chain sequences and cloning the fragments into a pHybC-D2 vector according to Example 1.4.4.1. The DVD-Ig constructs were cloned into and expressed in 293 cells as described in Example 1,4.4.2. The DVD-Ig protein was purified according to standard methods. Functional characteristics were determined according to the methods described in Example 1.1.1 and 1.1.2 as indicated. DVD-Ig VH and VL chains for the DVD-Igs disclosed herein are provided below.

Example 2.1: Generation of TNF (seq. 2) and NGF DVD-Igs with Linker Set 2

Table 15

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
52	DVD1459H	AB213VH	AB020VH	QVQLKESGPGGLVAPSQSLSTICTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSDYDSTLKSRLSIS KDNSKSQLFLKMNLSLQDDTARYYCAREWHHGPVA YWGQGLVTVSA A STKGPQVQLQESGPGLVKPSSET LSLCTVSGFSLIGYDLNWIROPFGKLEWIGI IW GDGTTDYN SAVKSRVTISKDTSKNQFSLKLSSVTA ADTAVYYCARGGYWYATSYFDYWGQGLVTVSS
53	DVD1459L	AB213VL	AB020VL	DIVMTQSHKFMSTTVGDRVSTICKASQAVSSAVAW YQKPGQSPKLLIYWASTRHTGVDRFTGSGSVTD FTLTIHNLQAEDLALYYCQQHYSTPFTFGSGTKLE IKRT V A A PS V F I FPPIQMTQSPSSLSASVGDVRT ITCRASQSI SNLNWYQKPGKAPKLLIYYTSRFH SGVPSRFSGSGSDTFTFTISSLPEDIATYYCQQ EHTLPYTFGQGTKLEIKR
54	DVD1460H	AB020VH	AB213VH	QVQLQESGPGLVKPSSETLSLCTVSGFSLIGYDLN WIRQPPGKLEWIGI IWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSSVTAADTAVYYCARGGYWYATS YYFDYWGQGLVTVSS A STKGPQVQLKESGPGLVA PSQSLSTICTVSGFSLTDYGVN WVRQPPGKLEWL GMIWGDGSDYDSTLKSRLSISKDNSKSQLFLKMN LSLQDDTARYYCAREWHHGPVAYWGQGLVTVSA
55	DVD1460L	AB020VL	AB213VL	DIQMTQSPSSLSASVGDVRTITCRASQSI SNLNW YQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSDT FTFTISSLPEDIATYYCQQEHTLPYTFGQGTKLE IKRT V A A PS V F I FPPIVMTQSHKFMSTTVGDRVS ITCKASQAVSSAVAWYQKPGQSPKLLIYWASTRH TGVPDRFTGSGSVTFTLTIHNLQAEDLALYYCQQ HYSTPFTFGSGTKLEIKR

Example 2.2: Generation of TNF (seq. 3) and NGF DVD-Igs with Linker Set 2

Table 16

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
56	DVD1461H	AB214VH	AB020VH	EVKLEESGGGLVQPGGSMKLSVCVASGFIFSNHWMN WVRQSPKGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSksAVYLQMTDLRTEDTGVYYCSRNYGS TYDYWGQGTTLTVSS ASTKGP QVQLQESGPGLVKPK SETLSLTCTVSGFSLIGYDLNWIQPPGKLEWIG IIWGDGTTDYNsAVKSRVTISKDTskNQFSLKLS VTAADTAVYYCARGGYWYATSYFDYWGQGTTLTV SS
57	DVD1461L	AB214VL	AB020VL	DILLTQSPAILSvSPGERVsfSCRASQFVGSSIHw YQRTNGSPRLLIKYASESMGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHsWPFTFGSGTNLE VKR VAAPSVFI FPDIQMTQSPSSLSASVgDRVT ITCRASQSI SNLNWYQqKPGKAPKLLIYYTSRFH SGVPSRFSGSGSGTDFTFTISSLQPEDiatYYCQ EHTLPYTFGQGTKLEIKR
58	DVD1462H	AB020VH	AB214VH	QVQLQESGPGLVKPKSETLSLTCTVSGFSLIGYDLN WIQPPGKLEWIGIIWGDGTTDYNsAVKSRVTIS KDTskNQFSLKLSVTAADTAVYYCARGGYWYATS YYFDYWGQGTTLTVSS ASTKGP EVKLEESGGGLVQ PGGSMKLSVCVASGFIFSNHWMNWVRQSPKGLEWV AEIRSKSINSATHYAESVKGRFTISRDDSksAVYL QMTDLRTEDTGVYYCSRNYGStYDYWGQGTTLTV SS
59	DVD1462L	AB020VL	AB214VL	DIQMTQSPSSLSASVgDRVTITCRASQSI SNLNW YQqKPGKAPKLLIYYTSRFHsGVPSRFSGSGSGTD FTFTISSLQPEDiatYYCQEHtLPYTFGQGTKLE IKR VAAPSVFI FPDILLTQSPAILSvSPGERVsf SCRASQFVGSSIHwYQRTNGSPRLLIKYASESM SGIPSRFSGSGSGTDFTLSINTVESEDIADYYCQ ESHsWPFTFGSGTNLEVKR

Example 2.23: Generation of TNF (seq. 4) and NGF DVD-Igs with Linker Set 2

Table 17

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
60	DVD1463H	AB215VH	AB020VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMH WVRQAPGNGLEWVAFMSYDGSNRYAKDSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDRGIAAG GNYYYYGMDVWGQGTFTTVSS ASTKGP QVQLQESG PGLVKPSETLSLTCTVSGFSLIGYDLNWRQPPGK GLEWIGLIWGDGTTDYN SAVKSRVTISKDTSKNQF SLKLSVTAADTAVYYCARGGYWYATSYFDYWGQ GTLTVSS
61	DVD1463L	AB215VL	AB020VL	EIVLTQSPATLSLSPGERATLSCRASQSVSYLAW YQQKPGQAPRLLIYDASNRAITGIPARFSGSGSGTD FTLTISSLEPEDFAVYYCQQRSNWPPFTFGPGTKV DIKRT VAA PSVFI FP DIQMTIQSPSSLSASVGD RVTTTCRASQSI SNNLNWYQQKPKAPKLLIYTSRF HSGVPSRFSGSGSGTDFTFTISSLOPEDIAVYYCQ QEHTLPYTFGQGTKLEIKR
62	DVD1464H	AB020VH	AB215VH	QVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLN WIRQPPGKGLEWIGLIWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSVTAADTAVYYCARGGYWYATS YFDYWGQGTTLTVSS ASTKGP QVQLVESGGGVVQ PGRSLRLSCAASGFTFSSYAMHWVRQAPGNGLEWV AFMSYDGSNRYAKDSVKGRFTISRDN SKNTLYLQM NSLRAEDTAVYYCARDRGIAAGGNYYYYGMDVWGQ GTTTVSS
63	DVD1464L	AB020VL	AB215VL	DIQMTQSPSSLSASVGDRTTTCRASQSI SNNLNW YQQKPGKAPKLLIYTSRFHSGVPSRFSGSGSGTD FTFTISSLPEDIAVYYCQEHTLPYTFGQGTKLE IKRT VAA PSVFI FP EIVLTQSPATLSLSPGERAT LSCRASQSVSYLAWYQQKPCQAPRLLIYDASNRA ITGIPARFSGSGSGTDFTLTISSLEPEDFAVYYCQ RSNWPFTFGPGTKVDIKR

Example 2.4: Generation of TNF (seq. 5) and NGF DVD-Igs with Linker Set 2

Table 18

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
64	DVD1477H	AB217VH	AB020VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYDMH WVRQAPGKGLEWVAVIWSGSIKYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAREVESAMG GFYNGMDVWGQGT T V T V S S A S T K G P Q V Q L Q E S G P GLVKPSETLSLTCTVSGFSLIGYDLNWI R Q P P G K G LEWIGI IWGDGTTDYN SAVKSRVTISKDTSKNQFS LKLSSVTAADTAVYYCARGGYWYATSY Y F D Y W G Q G TLVTVSS
65	DVD1477L	AB217VL	AB020VL	DIQMTQSPSSLSASVGDRVTITCRASQGIRIDLGW YQQKPGKAPKRLIYAAS TLQSGVPSRFSGSGSGT F E I F T I S S L Q P E D F A S Y Y C L Q H K S Y P L T F G G G T K V E I K R T V A A P S V F I F P P D I Q M T Q S P S S L S A S V G D R V T ITCRASQSI S N N L N W Y Q Q K P G K A P K L L I Y Y T S R F H SGVPSRFSGSGSGTDF T F T I S S L Q P E D I A T Y Y C Q Q E H T L P Y T F G Q G T K L E I K R
66	DVD1478H	AB020VH	AB217VH	QVQLQESGPGLVK P S E T L S L T C T V S G F S L I G Y D L N WIRQPPGKGLEWIGI IWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSSVTAADTAVYYCARGGYWYATS YYFDYWGQGT L V T V S S A S T K G P Q V Q L V E S G G G V V Q PGRSLRLSCAASGFTFSSYDMHWVRQAPGKGLEWV AVIWSGSIKYYADSVKGRFTISRDN SKNTLYLQ M NSLRAEDTAVYYCAREVESAMGGFYNGMDVWGQ G TTVTVSS
67	DVD1478L	AB020VL	AB217VL	DIQMTQSPSSLSASVGDRVTITCRASQSI S N N L N W YQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGT D F T F T I S S L Q P E D I A T Y Y C Q Q E H T L P Y T F G Q G T K L E I K R T V A A P S V F I F P P D I Q M T Q S P S S L S A S V G D R V T ITCRASQGI R I D L G W Y Q Q K P G K A P K R L I Y A A S T L Q SGVPSRFSGSGSGT F E I F T I S S L Q P E D F A S Y Y C L Q H K S Y P L T F G G G T K V E I K R

Example 2.5: Generation of TNF (seq. 6) and NGF-Igs with Linker Set 2

Table 19

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
68	DVD1485H	AB218VH	AB020VH	EVQLVESGGGLIQPGGSLRRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARGEGGFYWGQGLVTVSSASTKGPQVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNWIRQPPGKGLEWIGI I WGDGTTDYN SAVKSRVTISKDT SKNQFSLKLSSVTAADTAVYYCARGGYWYATSYFDYWGQGLVTVSS
69	DVD1485L	AB218VL	AB020VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLQSEDFAVYYCQQYNYWWTFGQGTKVEIKRTVAAPSVFIFPPDIQMTQSPSSLSASVGDRTITCRASQSI SNNLNWYQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTDFTFTISLQPED IATYYCQQEHTL P YTFGQGTKLEIKR
70	DVD1486H	AB020VH	AB218VH	QVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNWIRQPPGKGLEWIGI I WGDGTTDYN SAVKSRVTISKDT SKNQFSLKLSSVTAADTAVYYCARGGYWYATSYFDYWGQGLVTVSSASTKGPVQLVESGGGLIQPGGSLRRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARGEGGFYWGQGLVTVSS
71	DVD1486L	AB020VL	AB218VL	DIQMTQSPSSLSASVGDRTITCRASQSI SNNLNWYQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTDFTFTISLQPED IATYYCQQEHTL P YTFGQGTKLEIKRTVAAPSVFIFPPEIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLQSEDFAVYYCQQYNYWWTFGQGTKVEIKR

Example 2.6: Generation of TNF (seq. 2) and SOST DVD-Igs with Linker Set 2

Table 20

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
72	DVD1453H	AB213VH	AB022VH	QVQLKESGPGLVAPSQSLSTICTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSTDYDSTLKSRLSIS KDNSKSIIFLKMNSLQDDTARYYCAREWHHGPVA YWGQGLVTVSA AASTKGPEVQLQOSGPELVT PGAS VKISCKASGYTFTDHYMSWVKOSHGKSLEWIGDIN PYSGETTYNQKFKGTATLTVDKSSSIAYMEIRGLT SEDSAVYYCARDYDASPFAYWGQGLVTVSA
73	DVD1453L	AB213VL	AB022VL	DIVMTQSHKFMSTTVGDRVSI TC KASQAVSSAVAW YQKPGQSPKLLIYWASTRHTGVPDRFTGSGSVTD FTLTIHNLQAEDLALYYCQQHYSTPFTFGSGTKLE IKR TVAAPSVFI FPDPVQMIQSPSSLSASLGDIVT MTCQASQGTSINLNWFQKPKAPKLLIYGSSNLE DGVPSRFSGSRYGTDFTLTISSELEDELATYFCLQ HSYLPYTFGGGKLEIKR
74	DVD1454H	AB022VH	AB213VH	EVQLQOSGPELVT PGAS VKISCKASGYTFTDHYMS WVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGLVTVSA AASTKGPQVQLKESGPGLVAPS QSLSTICTVSGFSLTDYGVN WVRQPPGKGLEWLGMI WGDGSTDYDSTLKSRLSISKDNSKSIIFLKMNSL QDDTARYYCAREWHHGPVAYWGQGLVTVSA
75	DVD1454L	AB022VL	AB213VL	DVQMIQSPSSLSASLGDIVT MTCQASQGTSINLNW FQKPKAPKLLIYGSSNLE DGVPSRFSGSRYGTD FTLTISSELEDELATYFCLQHSYLPYTFGGGKLE IKR TVAAPSVFI FPDIVMTQSHKFMSTTVGDRVS ITCKASQAVSSAVAWYQKPGQSPKLLIYWASTRH TGVPDRFTGSGSVTDFTLTIHNLQAEDLALYYCQQ HYSTPFTFGSGTKLEIKR

Example 2.7: Generation of TNF (seq. 3) and SOST DVD-Igs with Linker Set 2

Table 21

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
76	DVD1455H	AB214VH	AB022VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGLWVAEIRSKSINSATHYAESVKGRF TISRDDSksAVYLQMTDLRTEdTGvYYCSrNYGS TYDYWGQGTTLTVSS ASTKGPEVQLQQSGPELVTP GASVKLSCKASGYTFDHYMSWVKOSHGKSLEWIG DINPYSGETTYNQKFKGTATLTVDKSSSIAYMEIR GLTSEDSAVYYCARDYDASPFAYWGQGTTLTVSA
77	DVD1455L	AB214VL	AB022VL	DILLTQSPAILSVSPGERVSVFSCRASQFVGSSIH YQORTNGSPRLLIKYASESMGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKRT VAAPSVFI FPPDVQMIQSPSSLSASLGDIVT MTCQASQGTSINLNWFQQKPGKAPKLLIYGSSNLE DGVPSRFSGSRYGTDFTLTISSELEDEDLATYFCLQ HSYLPYTFGGGTKLEIKR
78	DVD1456H	AB022VH	AB214VH	EVQLQQSGPELVTPGASVKISCKASGYTFDHYMS WVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGTTLTVSA ASTKGPEV KLEESGGGLVQPG GSMKLSCVASGFIFSNHWMNWVRQSPKGLWVAE IRSKSINSATHYAESVKGRFTISRDDSksAVYLQ MtdLRTEdTGvYYCSrNYGStYDYWGQGTTLTVSS
79	DVD1456L	AB022VL	AB214VL	DVQMIQSPSSIASLGDIVTMTTCQASQGTSINLNW FQQKPGKAPKLLIYGSSNLEDGVPSRFSGSRYGTD FTLTISSELEDEDLATYFCLQHSYLPYTFGGGTKLE IKRT VAAPSVFI FPPDILLTQSPAILSVSPGERVS FSCRASQFVGSSIHVYQORTNGSPRLLIKYASESM SGIPSRFSGSGSGTDFTLSINTVESEDIADYYCQ ESHWPFTFGSGTNLEVKR

Example 2.8: Generation of TNF(seq. 4) and SOST DVD-Igs with Linker Set 2

Table 22

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
80	DVD1457H	AB215VH	AB022VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFFSSYAMHWVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDRGIAAGGNYYYGMDVWGQGTITVTVSS ASTKGP EVQLQDSGPELVTPGASVKISCKASGYTFTDHYMSWVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLTVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASPFAYWGQGTITVTVSA
81	DVD1457L	AB215VL	AB022VL	EIVLTQSPATLSLSPGERATLSCRASQSVSYLAWYQQKPGQAPRLLIYDASNRTGIPARFSGSGSGTDFTLTISLSLEPEDEFAVYYCQQRSNWPPFTFGPGTKVDIKR TVAAPSVFI FPDPVQMISPSLSASLGDIVTMTQASQGTSLNINWFQKPKAPKLLIYGSSNLEDGVPSRFSGSRYGTDFTLTISLSLEDEDLATYFCLQHSYLPYTFGGGKLEIKR
82	DVD1458H	AB022VH	AB215VH	EVQLQDSGPELVTPGASVKISCKASGYTFTDHYMSWVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLTVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASPFAYWGQGTITVTVSA ASTKGP QVQLVESGGGVVQPGRSLRLSCAASGFTFFSSYAMHWVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDRGIAAGGNYYYGMDVWGQGTITVTVSS
83	DVD1458L	AB022VL	AB215VL	DVQMISPSLSASLGDIVTMTQASQGTSLNINWFQKPKAPKLLIYGSSNLEDGVPSRFSGSRYGTDFTLTISLSLEDEDLATYFCLQHSYLPYTFGGGKLEIKR TVAAPSVFI FPPEIVLTQSPATLSLSPGERATLSCRASQSVSYLAWYQQKPGQAPRLLIYDASNRTGIPARFSGSGSGTDFTLTISLSLEPEDEFAVYYCQQRSNWPPFTFGPGTKVDIKR

Example 2.9: Generation of TNF (seq. 5) and SOST DVD-Igs with Linker Set 2

Table 23

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
84	DVD1475H	AB217VH	AB022VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSYDMH WVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAREVESAMG GFYYNGMDVWGQGTITVTVSS ASTKGP EVQLQSSGP ELVTPGASVKISCKASGYTFTHYMSWVKOSHGKS LEWIGDINPYSGETTYNQKFKGTATLTVDKSSSIA YMEIRGLTSEDSAVYYCARDYDASPFAYWGQGT ITVSA
85	DVD1475L	AB217VL	AB022VL	DIQMTQSPSSLSASVGDRVTITCRASQGRIDLGW YQQKPGKAPKRLIYAASLTQSGVPSRFSGSGSGTE FIFTISSLQPEDFASYCLQHKSYPLTFGGGTKVE IKRT VAAPSVFI FPDPVQMISPSLSASLGDIVT MTCQASQGTSINLNWFQQKPGKAPKLLIYSSNLE DGVPSRFSGSRYGTDFTLTISLLEDEDLATYFCLQ HSYLPYTFGGGTKLEIKR
86	DVD1476H	AB022VH	AB217VH	EVQLQSSGPELVTPGASVKISCKASGYTFTHYMS WVKQSHCKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGTITVTVS AASTKGP QVQLVESGGGVVQPG RSLRLSCAASGFTFSYDMHWVRQAPGKGLEWVAV IWSDGSIKYYADSVKGRFTISRDNKNTLYLQMN SLRAEDTAVYYCAREVESAMGGFYNGMDVWGQGT ITVSS
87	DVD1476L	AB022VL	AB217VL	DVQMIQSPSSLSASLGDIVTMTTCQASQGTSINLNW FQQKPGKAPKLLIYSSNLEDGVPSRFSGSRYGTD FTLTISLLEDEDLATYFCLQHSYLPYTFGGGTKLE IKRT VAAPSVFI FPDIQMTQSPSSLSASVGDRVT ITCRASQGRIDLGWYQQKPGKAPKRLIYAASLTQ SGVPSRFSGSGSGTEFIFTISSLQPEDFASYCLQ HKSYPYTFGGGTKVEIKR

Example 2.10: Generation of TNF (seq. 6) and SOST DVD-Igs with Linker Set 2

Table 24

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
88	DVD1483H	AB218VH	AB022VH	EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFQDYWGQGLVTVSS ASTKGPEVQLQQSGPELVTPGASVKISCKASGYTF TDHYMSWVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLTVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASPFAYWGQGLVTVSA
89	DVD1483L	AB218VL	AB022VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQKQPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISISLQSEDFAVYYCQQYNYWWTFGQGTKVEIKR TVAAPSVFI FPDPVQMIQSPSSLSASLGDIVTMTCCASQGTINLNWFQQKPGKAPKLLIYGSSNLEDGVP SRFRSGSRYGTDFTLTISLSEDEDLATYFCLQHSYLPYTFGGGTKLEIKR
90	DVD1484H	AB022VH	AB218VH	EVQLQQSGPELVTPGASVKISCKASGYTFTDHYMSWVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLTVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASPFAYWGQGLVTVSA ASTKGPEVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFQDYWGQGLVTVSS
91	DVD1484L	AB022VL	AB218VL	DVQMIQSPSSLSASLGDIVTMTCCASQGTINLNWFQQKPGKAPKLLIYGSSNLEDGVP SRFRSGSRYGTDFTLTISLSEDEDLATYFCLQHSYLPYTFGGGTKLEIKR TVAAPSVFI FPPEIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQKQPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLQSEDFAVYYCQQYNYWWTFGQGTKVEIKR

Example 2.11: Generation of TNF (seq. 2) and PGE2 with Linker Sct 2

Table 25

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				1234567890123456789012345678901234
92	DVD1447H	AB213VH	AB048VH	QVQLKESGPGGLVAPSQSL SITCTVSGFSLTDYGV NWVRQPPGKGLEWLGMIWGDGSTDYDSTLKSRLS ISKDNSKSLFLKMNSLQTDDTARYYCAREWHHG PVAYWQGGLVTVSA ASTKGP EVQLVQSGAEVKK PGASVKVSCASGYTFTKYWLGWVROAPGOGLEW MGDIYPGYDYTHYNEKFKDRVTLTDTSTSTAYM ELRSLRSDDTAVYYCARSDGSSTYWGQGLVTVS S
93	DVD1447L	AB213VL	AB048VL	DIVMTQSHKFMSTTVGDRV SITCKASQAVSSAVA WYQQKPGQSPKLLIYWASTRHTGVPDRFTGSGSV TDFTLTIHNLQAEDLALYYCQQHYSTPFTFGSGT KLEIKR TVAAPSVFI FPDVLMTQTPLSLPVT EPASISCTSSQNIVHSNGNTYLEWYLQKPGQSPQ LLIYKVSNRFSGVPDRFSGSGSDTDFTLKISRVE AEDVGVYYCFQVSHVPTFGGCTKVEIKR
94	DVD1448H	AB048VH	AB213VH	EVQLVQSGAEVKKPGASVKVSCASGYTFTKYWL GWVROAPGQGLEWMGDIYPGYDYTHYNEKFKDRV TLTDTSTSTAYMELRSLRSDDTAVYYCARSDGS STYWGQGLVTVSS ASTKGP QVQLKESGPGGLVAP SQSL SITCTVSGFSLTDYGVNWVRQPPGKGLEWL GMIWGDGSTDYDSTLKSRLSISKDNSKSLFLKM NSLQTDDTARYYCAREWHHG PVAYWQGGLVTVS A
95	DVD1448L	AB048VL	AB213VL	DVLMTQTPLSLPVTGPGEPAISICTSSQNIVHSNG NTYLEWYLQKPGQSPQLLIYKVSNRFSGVPDRFS GSGSDTDFTLKISRVEAEDVGVYYCFQVSHVPT FGGCTKVEIKR TVAAPSVFI FPDVLMTQSHKFM STTVGDRV SITCKASQAVSSAVAWYQQKPGQSPK LLIYWASTRHTGVPDRFTGSGSVTDFTLTIHNLQ AEDLALYYCQQHYSTPFTFGSGTKLEIKR

Example 2.12: Generation of TNF (seq. 3) and PGE2 DVD-Igs with Linker Set 2

Table 26

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
96	DVD1449H	AB214VH	AB048VH	EVKLEESGGGLVQPGGSMKLSCVASGFI FSNHWMN WVRQSPKGLWVAEIRSKSINSATHYAESVKGKF TISRDDSKSAVYLQMTDLRTEDTGVIYCSRNYGGS TYDYWGQGTTLTVSS ASTKGPEVQLVQSGAEVKKP GASVKVSKASGYTFTKYWLGWVRQAPGQGLEWVG DIYPGYDYTHYNEKFKDRVLTLDTSTSTAYMELR SLRSDDTAVYYCARSDGSSTYWGQGTTLTVSS
97	DVD1449L	AB214VL	AB048VL	DILLTQSPAILLSVSPGERVVSFSCRASQFVGSSTHW YQORTNGSPRLLIKYASEMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHVSWPFTFGSGTNLE VKRT VAAPSVFI FPDVLMTQTPLSLPVTPGEPAS ISCTSSQNI VHSNGNTYLEWYLQKPGQSPQLLIYK VSNRFSGVDPDRFSGSGSGTDFTLTKISRVEAEDVGV YYCFQVSHVPTYTFGGGKVEIKR
98	DVD1450H	AB048VH	AB214VH	EVQLVQSGAEVKKPGASVKVSKASGYTFTKYWLG WVRQAPGQGLEWVGDIYPGYDYTHYNEKFKDRVTL TDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTY WGQGTTLTVSS ASTKGPEV KLEESGGGLVQPGGSM KLSVCVASGFI FSNHWMNWVRQSPKGLWVAEIRS KSINSATHYAESVKGFTISRDDSKSAVYLQMTDL RTEDTGVIYCSRNYGSTYDYWGQGTTLTVSS
99	DVD1450L	AB048VL	AB214VL	DVLMTQTPLSLPVTPGEPASISCTSSQNI VHSNGN TYLEWYLQKPGQSPQLLIYKSNRFSGVDPDRFSGS SGTDFTLTKISRVEAEDVGVYYCFQVSHVPTYTFGG GKVEIKRT VAAPSVFI FPDILLTQSPAILLSVSP GERVVSFSCRASQFVGSSTHWYQORTNGSPRLLIKY ASEMSGIPSRFSGSGSGTDFTLSINTVESEDIAD YYCQESHVSWPFTFGSGTNLEVKR

Example 2.13: Generation of TNF (seq. 4) and PGE2 DVD-Igs with Linker Set 2

Table 27

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
100	DVD1451H	AB215VH	AB048VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMH WVRQAPGNGLLEWVAFMSYDGSNKYAKDSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDRGIAAG GNYYYYGMDVWGQGT TTVT VSS ASTKGP EVQLVQSG AEVKKPGASVKVSKASGYTFKYWLGWVRQAPGQ GLEWMGDIYPGYDYTHYNEKFKDRVTLTDTSTST AYMELRSLRSDDTAVYYCARSDGSSTYWGQGT LV VSS
101	DVD1451L	AB215VL	AB048VL	EIVLTQSPATLSLSPGERATLSCRASQSVYSYLAW YQQKPGQAPRLLIYDASNRAIGIPARFSGSGGTD FTLTISSLEPEDFAVYYCQQRSNWPPFTFGPGTKV DIKRT VAAPSVFI FPDVLMTQTPLSLPVT P GEPA SISCTSSQNIVHSNGNTYLEWYLQKPGQSPQLLIY KVSNRFSGV PDR FSGSGSGTDFTLKISRVEAEDVG VYYCFQVSHVPYTFGGGTKVEIKR
102	DVD1452H	AB048VH	AB215VH	EVQLVQSGAEVKKPGASVKVSKASGYTFKYWLG WVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTL TDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTY WGQGT LV VSS ASTKGP QVQLVESGGGVVQPGRSL RLSCLASGFTFSSYAMHWVRQAPGNGLLEWVAFMSY DGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRA EDTAVYYCARDRGIAAGGNYYYYGMDVWGQGT TVT VSS
103	DVD1452L	AB048VL	AB215VL	DVLMTQTPLSLPVT P GEPA S ISCTSSQNIVHSNGN TYLEWYLQKPGQSPQLLIYKVSNRFSGV PDR FSGS GSGTDFTLKISRVEAEDVG V YYCFQVSHVPYTFGG GTKVEIKR VAAPSVFI FPPEI V LVTQSPATLSLSP GERATLSCRASQSVYSYLAWYQQKPGQAPRLLIYD ASNRAIGIPARFSGSGGTDFTLTISSLEPEDFAV YYCQQRSNWPPFTFGPGTKVDIKR

Example 2.14: Generation of TNF (seq. 5) and PGE2 DVD-Igs with Linker Set 2

Table 28

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
104	DVD1473H	AB217VH	AB048VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSYDMH WVRQAPGKGLEWVAVIWSGSIKYYADSVKGRFTI SRDNSKNTLYLQMNLSRAEDTAVYYCAREVESAMG GFYNGMDVWGQGTITVTVSS ASTKGP EVQLVQSGA EVKKGASVKVSKASGYTFTKYWLGWVROAPGQG LEWMGDIYPGYDYTHYNEKFKORVTLTDTSTSTA YMLRSLRSDDTAVYYCARSDGSSTYWGQGTITVTV SS
105	DVD1473L	AB217VL	AB048VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGW YQQKPGKAPKRLIYAASLTQSGVPSRFSGSGSGTE FIFTISSLQPEDFASYCYLQHKSYPLTFGGGTKVE IKR TVAAPSVFIFPP DVLMTQTPLSLPVTPGEPAS ISCTSSQNIIVHSNGNTYLEWYLQKPGQSPQLLIYK VSNRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGV YYCFQVSHVPTYFGGGTKVEIKR
106	DVD1474H	AB048VH	AB217VH	EVQLVQSGAEVKKPGASVKVSKASGYTFTKYWLW WVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTL TDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTY WGQGTITVTVSS ASTKGP QVQLVESGGGVVQPGRSL RLSCAASGFTFSYDMHWVROAPGKGLEWVAVIWS DGSIKYYADSVKGRFTISRDNKNTLYLQMNLSRA EDTAVYYCAREVESAMGGFYNGMDVWGQGTITVTV SS
107	DVD1474L	AB048VL	AB217VL	DVLMTQTPLSLPVTPGEPASISCTSSQNIIVHSNGN TYLEWYLQKPGQSPQLLIYKVSNRFSGVPDRFSGS GSGTDFTLKISRVEAEDVGVYYCFQVSHVPTYFGG GTKVEIKR TVAAPSVFIFPP DIQMTQSPSSLSASV GDRVTITCRASQGIRIDLGWYQQKPGKAPKRLIYA ASTLQSGVPSRFSGSGSGTEFIFTISSLQPEDFAS YYCQLQHKSYPLTFGGGTKVEIKR

Example 2.15: Generation of TNF (seq. 6) and PGE2 DVD-Igs with Linker Set 2

Table 29

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
108	DVD1481H	AB218VH	AB048VH	EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRTYYADSVKGRFTISRDNKNTLYLQMNLSLRAEDTAVYYCARGEGGFQDYWGQGLVTVSS ASTKGP EVQLVQSGAEVKKPGASVKVSCKASGYTFYKYWLGWVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTDTSTSTAYMELRSLRSDTAVYYCARSDGSSTYWGQGLVTVSS
109	DVD1481L	AB218VL	AB048VL	EIVMTQSPATLSVSPGERATLSCRASQSVSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISSLQSEDFAVYYCQQYNYWWTFGGQTKVEIKR TVAAPSVFI FPDVLMTQTPLSLPVTPGEPASISCTSSQNIIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFSGVDPDRFSGSGSGTEFTLKISRVEAEDVGVVYCFQVSHVPPYTFGGQTKVEIKR
110	DVD1482H	AB048VH	AB218VH	EVQLVQSGAEVKKPGASVKVSCKASGYTFYKYWLGWVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTDTSTSTAYMELRSLRSDTAVYYCARSDGSSTYWGQGLVTVSS ASTKGP EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRTYYADSVKGRFTISRDNKNTLYLQMNLSLRAEDTAVYYCARGEGGFQDYWGQGLVTVSS
111	DVD1482L	AB048VL	AB218VL	DVLMTQTPLSLPVTPGEPASISCTSSQNIIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFSGVDPDRFSGSGSGTEFTLKISRVEAEDVGVVYCFQVSHVPPYTFGGQTKVEIKR TVAAPSVFI FPPEIVMTQSPATLSVSPGERATLSCRASQSVSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISSLQSEDFAVYYCQQYNYWWTFGGQTKVEIKR

Example 2.16: Generation of TNF (seq. 2) and LPA DVD-Igs with Linker Set 2

Table 30

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
112	DVD1465H	AB213VH	AB216VH	QVQLKESGPGLVAPSQSL SITCTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSTDYDSTLKSRLSIS KDNSKSQLFLKMNSLQTD TARYYCAREWHHGPVA YWGQGLVTVSAASTKGPQVQLQQSGAELVRPGTS VKVSKASCYGFINYLIEWIKQRPQGLEWIGLIN PGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLT SEDSAVYFCARRFGYYGSGNYFDYWQGTTLTVSS
113	DVD1465L	AB213VL	AB216VL	DIVMTQSHKFMSTTVGDRV SITCKASQAVSSAVAW YQKPGQSPKLLIYWASTRHTGVPDRFTGSGSVTD FTLTIHNLQAEDLALYYCQHYSTPFTFGSGTKLE IKRTVAAPSVEIFPPDVMTQTPLSLPVSLGDQAS ISCTSGQSLVHINGNTYLHWYLQKPGQSPKLLIYK VSNLFSGVDPDRFSGSGSTDFTLKISRVEAEDLGV YFCSQSTHFPFTFGTGKLEIKR
114	DVD1466H	AB216VH	AB213VH	QVQLQQSGAELVRPGTSVKVSKASCYGFINYLIE WIKQRPQGLEWIGLINPGSDYTNYNENFKGKATL TADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGS GNYFDYWQGTTLTVSSASTKGPQVQLKESGPGLV APSQSL SITCTVSGFSLTDYGVN WVRQPPGKGLEW LGMIWGDGSTDYDSTLKSRLSISKDNSKSQLFLKM NSLQTD TARYYCAREWHHGPVAYWGQGLVTVSA
115	DVD1466L	AB216VL	AB213VL	DVVM TQTPLSLPVSLGDQASISCTSGQSLVHINGN TYLHWYLQKPGQSPKLLIYK VSNLFSGVDPDRFSGS GSGTDFTLKISRVEAEDLGVYFCSQSTHFPFTFGT GTKLEIKRTVAAPSVEIFPPDIVMTQSHKFMSTTV GDRV SITCKASQAVSSAVAWYQKPGQSPKLLIYW ASTRHTGVPDRFTGSGSVTDFTLTIHNLQAEDLAL YYCQHYSTPFTFGSGTKLEIKR

Example 2.17: Generation of TNF (seq. 3) and LPA DVD-Igs with Linker Set 2

Table 31

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
116	DVD1467H	AB214VH	AB216VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSKSAVYLQMTDLRTEDTGVYYCSRNYGGS TYDYWGQGTTLTVSS ASTKGP QVQLQQSGAELVRF GTSVKVSCKASGYGFINYLIEWIKORPGGLEWIG LINPGSDYTNYNENFKGKATLTADKSSSTAYMHL SLTSEDSAVYFCARREGYYGSNYFDYWGQGTTLT VSS
117	DVD1467L	AB214VL	AB216VL	DILLTQSPAILSVSPGERVSFSCRASQFVGS SIHWYQRTNGSPRLLIKYASEMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKR TVAAPSVFI FPPDVVMTQTPFLSLPVS LGDQASISCTSGQSLVHINGNTYLHWYLQKPGQSPKLLIYK VSNLFSGVPDRFSGSGSGTDFTLKISRVEAEDLGV YFCSQSTHFPFTFGTGKLEIKR
118	DVD1468H	AB216VH	AB214VH	QVQLQQSGAELVRFPTSVKVSCKASGYGFINYLIE WIKORPGGLEWIGLINPGSDYTNYNENFKGKATL TADKSSSTAYMHLSSLTSEDSAVYFCARREGYYGS GNYFDYWGQGTTLTVSS ASTKGP EVKLEESGGGLV QPGGSMKLSCVASGFIFSNHWMNWVRQSPKGGLEW VAEIRSKSINSATHYAESVKGRFTISRDDSKSAVY LQMTDLRTEDTGVYYCSRNYGGSYDYWGQGTTLT VSS
119	DVD1468L	AB216VL	AB214VL	DVVMTQTPFLSLPVS LGDQASISCTSGQSLVHINGN TYLHWYLQKPGQSPKLLIYK VSNLFSGVPDRFSGSGSGTD FTLKISRVEAEDLGVYFCSQST HFPFTFGTGKLEIKR TVAAPSVFI FPPDILLTQSPAILSVSP GERVSFSCRASQFVGS SIHWYQRTNGSPRLLIKY ASEMSGIPSRFSGSGSGTD FTLSINTVESEDIAD YYCQESHWPFTFGSGTNLEVKR

Example 2.18: Generation of TNF (seq. 4) and LPA DVD-Igs with Linker Set 2

Table 32

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
120	DVD1469H	AB215VH	AB216VH	QVQLVESGGGVVQPGRSLRSLSCAASGFTFSYAMHWVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDRGIAAGGNYYYGMDVWGQGT TTVTVSS ASTKGP QVQLQQSGAELVRPQTSVKVSKASGYGFINYLIEWIKORPGQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGGTTLTVSS
121	DVD1469L	AB215VL	AB216VL	EIVLTQSPATLSLSPGERATLSCRASQSVSYLAWYQQKFGQAPRLLIYDASNRAATGIPARFSGSGSGTDFTLTISLSLEPEDEFAVYYCQQRSNWPPFTFGPGTKVDIKR TVAAPS VFI FP PDVMTQTPLSLPVSLGDAQSISCTSGQSLVHNGNTYLHWYLQKPGQSPKLLIYKVSINLFSGVDPDRFSGSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPTFGTGTKLEIKR
122	DVD1470H	AB216VH	AB215VH	QVQLQQSGAELVRPQTSVKVSKASGYGFINYLIEWIKORPGQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGGTTLTVSS ASTKGP QVQLVESGGGVVQPGRSLRSLSCAASGFTFSYAMHWVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDRGIAAGGNYYYGMDVWGQGT TTVTVSS
123	DVD1470L	AB216VL	AB215VL	DVVTQTPLSLPVSLGDAQSISCTSGQSLVHNGNTYLHWYLQKPGQSPKLLIYKVSINLFSGVDPDRFSGSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPTFGTGTKLEIKR TVAAPS VFI FP PEIVLTQSPATLSLSPGERATLSCRASQSVSYLAWYQQKFGQAPRLLIYDASNRAATGIPARFSGSGSGTDFTLTISLSLEPEDEFAVYYCQQRSNWPPFTFGPGTKVDIKR

Example 2.19: Generation of TNF (seq. 1) and LPA DVD-Igs with Linker Set 2

Table 33

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
124	DVD1471H	AB017VH	AB216VH	EVQLVESGGGLVQPGKSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVGRFTISRDNAKNSLYLQMNLSRAEDTAVYYCAKVSYLSTASSLDYWGQGTLLVTVSS ASTKG QVQLQSGAELVPGTSVKVSCKASGYGFINYLIEWIKORPGGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGTLLTVSS
125	DVD1471L	AB017VL	AB216VL	DIQMTQSPSSLSASVGRVTITCRASQGIIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNRAPYTFGQGTKVEIKR TVAAPS VFI FP PDVVMQTPLSLPVSLGDAQASISCTSGQSLVHINGNTYLNHWYLNKPKGQSPKLLIYKVSINLFSGVPPDRFSGSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR
126	DVD1472H	AB216VH	AB017VH	QVQLQSGAELVPGTSVKVSCKASGYGFINYLIEWIKORPGGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGTLLTVSS ASTKG PEVQLVESGGGLVQPGKSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVGRFTISRDNAKNSLYLQMNLSRAEDTAVYYCAKVSYLSTASSLDYWGQGTLLTVSS
127	DVD1472L	AB216VL	AB017VL	DVVMQTPLSLPVSLGDAQASISCTSGQSLVHINGNTYLNHWYLNKPKGQSPKLLIYKVSINLFSGVPPDRFSGSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR TVAAPS VFI FP PDVVMQTQSPSSLSASVGRVTITCRASQGIIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNRAPYTFGQGTKVEIKR

Example 2.20: Generation of TNF (seq. 5) and LPA DVD-Igs with Linker Set 2

Table 34

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
128	DVD1479H	AB217VH	AB216VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSYDMHWVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGGFYNGMDVWGQGTITVTVSS ASTKGP QVQLQQSGAELVLRPGTSVKVSKASGYGFINYLIWIKQRPQGQLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARREFGYGSGNYFDYWGGTTLTVSS
129	DVD1479L	AB217VL	AB216VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGWYQQKPGKAPKRLIYAASLTQSGVPSRFSGSGSGTEFIFTISSLQPEDFASYYCLQHKSYPLTFGGGKVEIKR TVAAPSVFIFPP DVVMITQPLSLPVSLGDQASISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVSINLFSGVPDFRFSGSGSGTEFTLKISRVEAEDLGVYFCSQSTHEFFFTFGTGTKLEIKR
130	DVD1480H	AB216VH	AB217VH	QVQLQQSGAELVLRPGTSVKVSKASGYGFINYLIWIKQRPQGQLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARREFGYGSGNYFDYWGGTTLTVSS ASTKGP QVQLVESGGGVVQPGRSLRLSCAASGFTFSYDMHWVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGGFYNGMDVWGQGTITVTVSS
131	DVD1480L	AB216VL	AB217VL	DVVMITQPLSLPVSLGDQASISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVSINLFSGVPDFRFSGSGSGTEFTLKISRVEAEDLGVYFCSQSTHEFFFTFGTGTKLEIKR TVAAPSVFIFPP DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGWYQQKPGKAPKRLIYAASLTQSGVPSRFSGSGSGTEFIFTISSLQPEDFASYYCLQHKSYPLTFGGGKVEIKR

Example 2.21: Generation of TNF (seq. 6) and LPA DVD-Igs with Linker Set 2

Table 35

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
132	DVD1487H	AB218VH	AB216VH	EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYADSVKGRFTISRDNKNTLYLQMNLSLRAEDTAVYYCARGEGGFQYWGQGTLLVTVSS ASTKGP QVQLQQSGAELVLRPGTSVKVSKASGYGFINYLIEWIKORPGQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGGTLLTVSS
133	DVD1487L	AB218VL	AB216VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLQSEDFAVYYCQQYNYWWTFGQGTKVEIKR TVAA PSVFI FP PDVMTQTPLSLPVSLGDAQSISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVSNFLFSGVPDRFSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR
134	DVD1488H	AB216VH	AB218VH	QVQLQQSGAELVLRPGTSVKVSKASGYGFINYLIEWIKORPGQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGGTLLTVSS ASTKGP EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYADSVKGRFTISRDNKNTLYLQMNLSLRAEDTAVYYCARGEGGFQYWGQGTLLVTVSS
135	DVD1488L	AB216VL	AB218VL	DVVTQTPLSLPVSLGDAQSISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVSNFLFSGVPDRFSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR TVAA PSVFI FP PEIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLQSEDFAVYYCQQYNYWWTFGQGTKVEIKR

Example 2.22: Generation of TNF (seq. 2) and NGF DVD-Igs with Linker Set 3

Table 36

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
136	DVD1501H	AB213VH	AB020VH	QVQLKESGPGGLVAPSQSLSTICTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSTDYDSTLKSRLSIS KDNSKSQLFLKMNSLQDDTARYYCAREWHHGPVA YWGQGLVTVSA ASTKGPSVFPLAPQVQLQESGPG LVKPSETLSLTCTVSGFSLIGYDLNWIRQPPGKGL EWIGI IWGDGTTDYN SAVKSRVTISKDTSKNQFSL KLSSVTAADTAVYYCARGGYWYATSYFFDYWGQGT LTVSS
137	DVD1501L	AB213VL	AB020VL	DIVMTQSHKFMSTTVGDRVSI TCKASQAVSSAVAW YQQKPGQSPKLLIYWASTRHTGVPDRFTGSGSVTD FTLTIHNLQAEDLALYYCQQHYSTPFTFGSGTKLE IKR TVAAP DIQMTQSPSSLSASVGDRTITCRASQ SISNNLNWYQQKPGKAPKLLIYYTSRFHSGVPSRF SGSGSGTDFTFTISSLPEDIATYYCQEHLPYTFGQGT KLEIKR
138	DVD1502H	AB020VH	AB213VH	QVQLQESGPGGLVVKPSETLSLTCTVSGFSLIGYDLN WIRQPPGKGLEWIGI IWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSSVTAADTAVYYCARGGYWYATS YFFDYWGQGLVTVSS ASTKGPSVFPLAPQVQLKE SGPGLVAPSQSLSTICTVSGFSLTDYGVNWVRQPP GKGLEWLGMIWGDGSTDYDSTLKSRLSISKDNSKS QIFLKMNSLQDDTARYYCAREWHHGPVAYWGQGT LTVSA
139	DVD1502L	AB020VL	AB213VL	DIQMTQSPSSLSASVGDRTITCRASQSISNNLNW YQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTD FTFTISSLPEDIATYYCQEHLPYTFGQGT IKR TVAAP DIQMTQSHKFMSTTVGDRVSI TCKASQ AVSSAVAWYQQKPGQSPKLLIYWASTRHTGVPDRF TGSGSVTDFTLTIHNLQAEDLALYYCQQHYSTPFT FGSGTKLEIKR

Example 2.23: Generation of TNF (seq. 3) and NGF DVD-Igs with Linker Set 3

Table 37

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
140	DVD1503H	AB214VH	AB020VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSKSAVYLQMTDLRTEDTGVIYCSRNYGS TYDYWGQGTTLTVSS ASTKGPSVFPLAPQVQLQES GPGLVKPSETLSLTCTVSGFSLIGYDLNWIROP KGLEWIGLIWGDGTTDYN SAVKSRVTISKDTSKNQ FSLKLSVTAADTAVYYCARGGYWYATSYFDYWG QGTTLTVSS
141	DVD1503L	AB214VL	AB020VL	DILLTQSFALLSVSPGERVSFSCRASQFVGS SIHWYQRTNGSPRLLIKYASEMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKRT VAAP DIQMTQSPSSLSASVGDRTITCRASQ SISNNLNWYQKPGKAPKLLIYYTSRFHSGVPSRF SGSGSGTDFTFTISSLPEDIATYYCQEHLPYTFGQGT KLEIKR
142	DVD1504H	AB020VH	AB214VH	QVQLQESGPGLVKPSETLSLTCTVSGFSLIGYDLN WIRQPPKGLEWIGLIWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSVTAADTAVYYCARGGYWYATS YFDYWGQGTTLTVSS ASTKGPSVFPLAPEVKLEE SGGGLVQPGGSMKLSCVASGFIFSNHWMNWVRQSP EKGLEWVAEIRSKSINSATHYAESVKGRFTISRDD SKSAVYLQMTDLRTEDTGVIYCSRNYGSTYDYWG QGTTLTVSS
143	DVD1504L	AB020VL	AB214VL	DIQMTQSPSSLSASVGDRTITCRASQSISNNLNW YQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTD FTFTISSLPEDIATYYCQEHLPYTFGQGT KLEIKR VAAP DILLTQSFALLSVSPGERVSFSCRASQ FVGS SIHWYQRTNGSPRLLIKYASEMSGIPSRF SGSGSGTDFTLSINTVESEDIADYYCQESHWPFT FGSGTNLEVKR

Example 2.24: Generation of TNF (seq. 4) and NGF DVD-Igs with Linker Set 3

Table 38

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
144	DVD1505H	AB215VH	AB020VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMH WVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDRGIAAC GNYYYYGMDVWGQGT'TVTVSS ASTKGPSVFPLAPQ VQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNW IRQPPGKLEWIGI IWGDGTTDYN SAVKSRVTISK DTSKNQFSKLSSVTAADTAVYYCARGGYWYATSY YFDYWQGT'LVTVSS
145	DVD1505L	AB215VL	AB020VL	EIVLTQSPFATLSLSPGERATLSCRASQSVYSYLAW YQQKPGQAPRLLIYDASN RATGIPARFSGSGSGTD FTLTISSLEPEDEFAVYYCQQRSNWPPFTFGPGTKV DIKR TVAAP DIQMTQSPSSLSASVGDRTITCRAS QSISNNLNWYQQKPGKAPKLLIYYTSRFHSGVPSR FSGSGSGTDFTFTISSLQPEDATYYCQEH TLPY TFGQGTKLEIKR
146	DVD1506H	AB020VH	AB215VH	QVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLN WIRQPPGKLEWIGI IWGDGTTDYN SAVKSRVTIS KDTSKNQFSKLSSVTAADTAVYYCARGGYWYATS YFDYWCQGT'LVTVSS ASTKGPSVFPLAPQ VQLVE SGGGVVQPGRSLRLSCAASGFTFSSYAMH WVRQAP GNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDN SK NTLYLQMNSLRAEDTAVYYCARDRGIAAGNYYYY GMDVWGQGT'TVTVSS
147	DVD1506L	AB020VL	AB215VL	DIQMTQSPSSLSASVGDRTITCRASQSISNNLNW YQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTD FTFTISSLQPEDATYYCQEH TLPYTFGQGTKLE IKR TVAAP EIVLTQSPFATLSLSPGERATLSCRASQ SVYSYLAWYQQKPGQAPRLLIYDASN RATGIPARF SGSGSGTDFTLTISSLEPEDEFAVYYCQQRSNWPPF TFGPGTKVDIKR

Example 2.25: Generation of TNF (seq. 5) and NGF DVD-Igs with Linker Set 3

Table 39

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
148	DVD1519H	AB217VH	AB020VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYDMH WVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAREVESAMG GFYYNGMDVWGQGTITVTVSS ASTKGPSVFPLAPQV QLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNWI RQPPGKLEWIGI IWGDGTTDYN SAVKSRVTISKD TSKNQFSLKLSVTAADTAVYYCARGGYWYATSY FDYWGQGLTVTVSS
149	DVD1519L	AB217VL	AB020VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGW YQQKPGKAPKRLIYAASLTQSGVPSRFSGSGSGTE FIFTISSLQPEDFASYCYLQHKSYPLTFGGGTKVE IKRT VAAP DIQMTQSPSSLSASVGDRTITCRASQ SISNNLNWYQQKPGKAPKLLIYYTSRFHSGVPSRF SGSGSGTDFITFTISSLQPEDIATYYCQEQHTLPYT FGQGTKLEIKR
150	DVD1520H	AB020VH	AB217VH	QVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLN WIRQPPGKLEWIGI IWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSVTAADTAVYYCARGGYWYATS YYFDYWGQGLTVTVSS ASTKGPSVFPLAPQVQLVE SGGGVVQPGRSLRLSCAASGFTFSSYDMHWVRQAP GKGLEWVAVIWSDGSIKYYADSVKGRFTISRDNK NTLYLQMNSLRAEDTAVYYCAREVESAMGGFYNG MDVWGQGTITVTVSS
151	DVD1520L	AB020VL	AB217VL	DIQMTQSPSSLSASVGDRTITCRASQISNNLNW YQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTD FTFTISSLQPEDIATYYCQEQHTLPYTFGQGTKLE IKRT VAAP DIQMTQSPSSLSASVGDRTITCRASQ GIRIDLGWYQQKPGKAPKRLIYAASLTQSGVPSRF SGSGSGTEFI FTISSLQPEDFASYCYLQHKSYPLT FGGGTKVEIKR

Example 2.26: Generation of TNF (seq. 6) and NGF-Igs with Linker Set 3

Table 40

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
152	DVD1527H	AB218VH	AB020VH	EVQLVESGGGLIQPGGSLRRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDNSKNTLYIQMNSLRAEDTAVYYCARGEGGFYWGQGLTIVTVSS ASTKGPSVFFPLAP QVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNWIRQPPGKLEWIGIITWGDGTTDYN SAVKSRVTISKDTSKNQFSLKLSVTAADTAVYYCARGGYWYATSYFFDYWGQGLTVTVSS
153	DVD1527L	AB218VL	AB020VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRATGLPARFSCGSGSGTEFTLTISLQSEDFAVYYCQQYNYWWTFCGQTKVEIKR TVAAP DIQMTQSPSSLSASVGDRTITCRASQISNNLNWYQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTDFFTTISLQPEDIATYYCQQEHTLPYTFGQGTKLEIKR
154	DVD1528H	AB020VH	AB218VH	QVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNWIRQPPGKLEWIGIITWGDGTTDYN SAVKSRVTISKDTSKNQFSLKLSVTAADTAVYYCARGGYWYATSYFFDYWGQGLTIVTVSS ASTKGPSVFFPLAPE VQLVESGGGLIQPGGSLRRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDNSKNTLYIQMNSLRAEDTAVYYCARGEGGFYWGQGLTVTVSS
155	DVD1528L	AB020VL	AB218VL	DIQMTQSPSSLSASVGDRTITCRASQISNNLNWYQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTDFFTTISLQPEDIATYYCQQEHTLPYTFGQGTKLEIKR TVAAPE IIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRATGLPARFSCGSGSGTEFTLTISLQSEDFAVYYCQQYNYWWTFCGQTKVEIKR

Example 2.27: Generation of TNF (seq. 2) and SOST DVD-Igs with Linker Set 3

Table 41

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
156	DVD1495H	AB213VH	AB022VH	QVQLKESGPGLVAPSQSL SITCTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSDYDSTLKSRLSIS KDNSKSI FLKMNSLQDDTARYYCAREWHHG PVA YWGQGLVTVVSAASTKGPSVFPPLAPEVQLQQSGPE LVT PGASVKISCKASGYTFTDHYMSWVKOSHGKSL EWIGDINPYSGETTYNQKFKGTATLTVDKSSSIAY MEIRGLTSEDSAVYYCARDYDASPFAYWGQGLV TVSA
157	DVD1495L	AB213VL	AB022VL	DIVMTQSHKFMSTTVGDRVSI TCKASQAVSSAVAW YQQKPGQSPKLLIYWASTRHTGVPDRFTGSGSVTD FTLTIHNLQAEDLALYYCQQHYSTPFTFGSGTKLE IKRTVAAPDVQMIQSPSSLSASLGDIVTMTQASQ GTSINLWVWFQQKPGKAPKLLIYGSSNLEDGVPSPRF SGSRYGTDFTLTISSLEDEDLATYFCLQHSYLPYT FGGGTKLEIKR
158	DVD1496H	AB022VH	AB213VH	EVQLQQSGPELVTPGASVKISCKASGYTFTDHYMS WVKQSHGKSL EWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASE FAYWGQGLVTVVSAASTKGPSVFPPLAPVQLKESG PGLVAPSQSL SITCTVSGFSLTDYGVN WVRQPPGK GLEWLGMIWGDGSDYDSTLKSRLSISKDNSKSI FLKMNSLQDDTARYYCAREWHHG PVA YWGQGLV TVSA
159	DVD1496L	AB022VL	AB213VL	DVQMIQSPSSLSASLGDIVTMTQASQGTSINLWV FQQKPGKAPKLLIYGSSNLEDGVPSPRFSGSRYGTD FTLTISSLEDEDLATYFCLQHSYLPYTFGGGTKLE IKRTVAAPDIVMTQSHKFMSTTVGDRVSI TCKASQ AVSSAVAWYQQKPGQSPKLLIYWASTRHTGVPDRF TGSGSVTDFTLTIHNLQAEDLALYYCQQHYSTPFT FGSGTKLEIKR

Example 2.28: Generation of TNF (seq. 3) and SOST DVD-Igs with Linker Set 3

Table 42

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
160	DVD1497H	AB214VII	AB022VH	EVKLEESGGGLVQPGGSMKLSVCVASGFIFSNHWMN WVRQSPKGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSKSAVYLMQMDLRTEDTGVYYCSRNYGGS TYDYWGQGTTLTVSS ASTKGPSVFPLAPEVQLQQS GPELVTPGASVKISCKASGYFTFDHYMSWVKQSHG KSLEWIGDINPYSGETTYNQKFKGTATLTVDKSSS IAYMEIRGLTSEDSAVYYCARDYDASPFAYWGQG TLVTVSA
161	DVD1497L	AB214VL	AB022VL	DILLTQSPAILSVSFGERVSVFSCRASQFVGSIIHW YQORTNGSPRLLIKYASESMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKR TVAAP DVQMIQSPSSLSASLGDIVTMTQASQ GTSINLNWFQKPKAPKLLIYGSSNLEDGVPSRF SGSRYGTDFTLTISSELEDEDLATYFCLQHSYLPYT FGGGTKLEIKR
162	DVD1498H	AB022VH	AB214VH	EVQLQQSGPELVTPGASVKISCKASGYFTFDHYMS WVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGTTLVTVS ASTKGPSVFPLAPEV KLEESG GGLVQPGGSMKLSVCVASGFIFSNHWMNWVRQSPK GLEWVAEIRSKSINSATHYAESVKGRFTISRDDSK SAVYLMQMDLRTEDTGVYYCSRNYGGSYDYWGQG TTLTVSS
163	DVD1498L	AB022VL	AB214VL	DVQMIQSPSSLSASLGDIVTMTQASQGTINLNW FQKPKAPKLLIYGSSNLEDGVPSRFSGSRYGTD FTLTISSELEDEDLATYFCLQHSYLPYTFGGGTKLE IKR TVAAP DILLTQSPAILSVSFGERVSVFSCRASQ FVGSIIHWYQORTNGSPRLLIKYASESMSGIPSRF SGSGSGTDFTLSINTVESEDIADYYCQESHWPFT FGSGTNLEVKR

Example 2.29: Generation of TNF(seq. 4) and SOST DVD-Igs with Linker Set 3

Table 43

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
164	DVD1499H	AB215VH	AB022VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMH WVRQAPGNGLLEWVAFMSYDGSNKYAKDSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDRGIAAG GNYYYYGMDVWGQGTITVTVSS ASTKGPSVFPLAPE VQLQQSGPELVTPGASVKISCKASGYTFTDHYMSW VKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLT VDKSSSIAYMEIRGLTSEDSAVYYCARDYDASPF AYWGQGTITVTVSA
165	DVD1499L	AB215VL	AB022VL	EIVLTQSPATLSLSPGERATLSCRASQSVSYLAW YQQKPGQAPRLLIYDASNRTGIPARFSGSGSGTD FTLTISSLEPEDFAVYYCQQRSNWPPFTEGPGTKV DIKRT VAA PDVQMIQSPSSLSASLGDIVTMTQAS QGTSINLNWFQQKPGKAPKLLIYGSSNLEDGVPSR FSGSRYGTDFTLTISSLEDEDLATYFCLQHSYLPY TFGGGTKLEIKR
166	DVD1500H	AB022VH	AB215VH	EVQLQQSGPELVTPGASVKISCKASGYTFTDHYMS WVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGTITVTVS AASTKGPSVFPLAP QVQLVESG GGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAPGN GLEWVAFMSYDGSNKYAKDSVKGRFTI SRDNSKNT LYLQMNSLRAEDTAVYYCARDRGIAAGGNYYYYGM DVWGQGTITVTVSS
167	DVD1500L	AB022VL	AB215VL	DVQMIQSPSSLSASLGDIVTMTQASQGTSINLNW FQQKPGKAPKLLIYGSSNLEDGVPSRFSGSRYGTD FTLTISSLEDEDLATYFCLQHSYLPYTFGGGTKLE IKRT VAA PEIVLTQSPATLSLSPGERATLSCRASQ SVSYLAWYQQKPGQAPRLLIYDASNRTGIPARF SGSGSGTDFTLTISSLEPEDFAVYYCQQRSNWPPF TFGPGTKVDIKR

Example 2.30: Generation of TNF (seq. 5) and SOST DVD-Igs with Linker Set 3

Table 44

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
168	DVD1517H	AB217VH	AB022VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFESSYDMHWVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGGFYNGMDVWGQGTITVTVSS ASTKGPSVFPLAPEVQLQQSGPELVTPGASVKISCKASGYTFTDHYMSWKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASPFAVWGQTLVTVSA
169	DVD1517L	AB217VL	AB022VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGWYQQKPGKAPKRLIYAASLTQSGVPSRFSGSGSGTEFIFTISLQPEDFASYICLQHKSYPLTFGGGTKVEIKR TVAAPDVQMIQSPSSLSASLGDIVTMTQASQGTSLNLNWFQQKPGKAPKLLIYGSSNLEDGVPSRFGSRYGTDFTLTISLLEDEDLATYFCLQHSYLPYTFGGGTKLEIKR
170	DVD1518H	AB022VH	AB217VH	EVQLQQSGPELVTPGASVKISCKASGYTFTDHYMSWVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLVTKSSSIAYMEIRGLTSEDSAVYYCARDYDASPFAVWGQTLVTVSA ASTKGPSVFPLAPQVQLVESGGVVQPGRSLRLSCAASGFTFESSYDMHWVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGGFYNGMDVWGQGTITVTVSS
171	DVD1518L	AB022VL	AB217VL	DVQMIQSPSSLSASLGDIVTMTQASQGTSLNLNWFQQKPGKAPKLLIYGSSNLEDGVPSRFGSRYGTDFTLTISLLEDEDLATYFCLQHSYLPYTFGGGTKLEIKR TVAAPDIQMTQSPSSLSASVGDRTITCRASQGIRIDLGWYQQKPGKAPKRLIYAASLTQSGVPSRFGSGSGGTEFIFTISLQPEDFASYICLQHKSYPLTFGGGTKVEIKR

Example 2.31: Generation of TNF (seq. 6) and SOST DVD-Igs with Linker Set 3

Table 45

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
172	DVD1525H	AB218VH	AB022VH	EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARGEGGF DYWGQGLVTVSSASTKGPSVFPLAPEVQLQSGPELVTPGASVKISCKASGYTFTDHYMSWVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLTVDKSSSIAYMEIRGLTSEDSAVYYCARD DYDASPFA YWGQGLVTVSS
173	DVD1525L	AB218VL	AB022VL	EIVMTQSPATLSVSPGERATLSCRASQSVSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLSQSEDFAVYYCQQYNYWWTFGGQTKVEIKRTVAAPDVQMIQSPSSLSASLGDIVTMTCAASQTSINLNWFQKPGKAPKLLIYGSSNLEDGVPSRFSGSRYGTDFTLTISLSEDEDLATYFCLQHSYLPYTFGGTKLEIKR
174	DVD1526H	AB022VH	AB218VH	EVQLQSGPELVTPGASVKISCKASGYTFTDHYMSWVKQSHGKSI EWIGDINPYSGETTYNQKFKGTATLTVDKSSSIAYMEIRGLTSEDSAVYYCARD DYDASPFA YWGQGLVTVSSASTKGPSVFPLAPEVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARGEGGF DYWGQGLVTVSS
175	DVD1526L	AB022VL	AB218VL	DVQMIQSPSSLSASLGDIVTMTCAASQTSINLNWFQKPGKAPKLLIYGSSNLEDGVPSRFSGSRYGTDFTLTISLSEDEDLATYFCLQHSYLPYTFGGQTKLEIKRTVAAP EIVMTQSPATLSVSPGERATLSCRASQSVSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLSQSEDFAVYYCQQYNYWWTFGGQTKVEIKR

Example 2.32: Generation of TNF (seq. 2) and PGE2 with Linker Set 3

Table 46

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				1234567890123456789012345678901234
176	DVD1489H	AB213VH	AB048VH	QVQLKESGPGLVAPSQSL SITCTVSGFSLTDYGV NWVRQPPGKGLEWLGMIWGDGSTDYDSTLKSRLS ISKDNSKSIIFLKMNSLQTD TARYCAREWHHG PVAYWGQGLVTVSA ASTKGPSVFPLA PEVQLVQ SGAEVKKPGASVKVSKASGYTFTKYWLGWVRQA PGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTDT STSTAYMELRSLRSDDTAVYYCARSDGSSTYWGQ GTLVTVSS
177	DVD1489L	AB213VL	AB048VL	DI VMTQSHKFMSTTVGDRVSI TCKASQAVSSAVA WYQQKPGQSPKLLIYWAS TRHTGVPDRFTGSGSV TDFTLTIHNLQAEDLALY YCQQHYSTPFTFGSGT KLEIKRT VAAP DVLMTQTPLSLPVT PGEPA SISC TSSQNIVHSNGNTYLEWYLQKPGQSPQLLIYKVS NRFSGVPRDFTGSGSGTDFTL KISRVEAEDVGVY YCFQVSHVPYTFGGGKVEIKR
178	DVD1490H	AB048VH	AB213VH	EVQLVQSGAEVKKPGASVKVSKASGYTFTKYWL GWVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRV TLTDTSTSTAYMELRSLRSDDTAVYYCARSDGS STYWGQGLVTVSS ASTKGPSVFPLA PQVQLKES GPGLVAPSQSL SITCTVSGFSLTDYGVNWVRQPP GKGLEWLGMIWGDGSTDYDSTLKSRLSISKDNSK SQIFLKMNSLQTD TARYCAREWHHG PVAYWGQ GTLVTVSA
179	DVD1490L	AB048VL	AB213VL	DVLMTQTPLSLPVT PGEPA SICTSSQNIVHSNG NTYLEWYLQKPGQSPQLLIYKVS NRFSGVPRDFS GSGSGTDFTL KISRVEAEDVGVYCFQVSHVPYT FGGGKVEIKRT VAAP DI VMTQSHKFMSTTVGDR VSI TCKASQAVSSAVAWYQQKPGQSPKLLIYWAS TRHTGVPDRFTGSGSVTDFTLTIHNLQAEDLALY YCQQHYSTPFTFGSGTKLEIKR

Example 2.33: Generation of TNF (seq. 3) and PGE2 DVD-Igs with Linker Set 3

Table 47

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
180	DVD1491H	AB214VH	AB048VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSKSAVYLQMTDLRTEDTGVYYCSRNYYS TYDYWGQGTTLTVSS ASTKGPSVFPLAPE VLVQS GAEVKKPGASVKVSCKASGYTFKYWLGWVRQAPG QGLEWMGDIYPGYDYTHYNEKFKDRVTLTDTSTS TAYMELRSLRSDDTAVYYCARSDGSSTYWGQGTTLV TVSS
181	DVD1491L	AB214VL	AB048VL	DILLTQSPAILSVSPGERVSFSCRASQFVGSIIHW YQORTNGSPRLLIKYASESMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHSWPFTFGSGTNLE VKR TVAAP DVLMTQTPLSLPVTPEGPASISCTSSQ NIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFSG VPDRFSGSGSGTDFTLKISRVEAEDVGVYYCFQVS HVPYTFGGGTKVEIKR
182	DVD1492H	AB048VH	AB214VH	EVQLVQSGAEVKKPGASVKVSCKASGYTFKYWLG WVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTL TDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTY WGQGTTLVTVSS ASTKGPSVFPLAPE VKLEESGGGL VQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGLE WVAEIRSKSINSATHYAESVKGRFTISRDDSKSAV YLQMTDLRTEDTGVYYCSRNYYS TYDYWGQGTTL TVSS
183	DVD1492L	AB048VL	AB214VL	DVLMTQTPLSLPVTPEGPASISCTSSQNIVHSNGN TYLEWYLQKPGQSPQLLIYKVSNRFSGVPDRFSGS GSGTDFTLKISRVEAEDVGVYYCFQVSHVPYTFGG GTKVEIKR TVAAP DILLTQSPAILSVSPGERVSFS CRASQFVGSIIHWYQORTNGSPRLLIKYASEMSG IPSRFSGSGSGTDFTLSINTVESEDIADYYCQESH SWPFTFGSGTNLEVKR

Example 2.34: Generation of TNF (seq. 4) and PGE2 DVD-Igs with Linker Set 3

Table 48

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
184	DVD1493H	AB215VH	AB048VH	QVQLVESGGGVVQPGRSLRLSCLASGFTFSSYAMHWVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDRLAAGGNYYYGMDVWGQGTITVTVSSASTKGPSVFPLAPEVQLVQSGAEVKKPGASVKVSCKASGYTFITKYWLGWVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTYWGQGTITVTVSS
185	DVD1493L	AB215VL	AB048VL	EIVLTQSPATLSLSPGERATLSCRASQSVYSYLAWYQQKPGQAPRLLIYDASNRTGIPARFSGSGSGTDFTLTITSSLEPEDFAVYYCQQRSNWPPFTFGPGTKVDIKRTVAAPDVLMTQITPLSLPVTPEPASISCTSSQNIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFSGVPDRFSGSGSGTDFTLTKISRVEAEDVGVYYCFQVSHVPYTFGGGTKVEIKR
186	DVD1494H	AB048VH	AB215VH	EVQLVQSGAEVKKPGASVKVSCKASGYTFITKYWLGWVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTYWGQGTITVTVSSASTKGPSVFPLAPQVQLVESGGGVVQPGRSLRLSCLASGFTFSSYAMHWVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDRLAAGGNYYYGMDVWGQGTITVTVSS
187	DVD1494L	AB048VL	AB215VL	DVLMTQITPLSLPVTPEPASISCTSSQNIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFSGVPDRFSGSGSGTDFTLTKISRVEAEDVGVYYCFQVSHVPYTFGGGTKVEIKRTVAAPDVLMTQITPLSLPVTPEPASISCTSSQNIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFSGVPDRFSGSGSGTDFTLTKISRVEAEDVGVYYCFQVSHVPYTFGGGTKVEIKR

Example 2.35: Generation of TNF (seq. 5) and PGE2 DVD-Igs with Linker Set 3

Table 49

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
188	DVD1515H	AB217VH	AB048VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYDMH WVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAREVESAMG GFYYNGMDVWGQGTTVTVSS ASTKGPSVFPLA PEV QLVQSGAEVKKPGASVKVSCKASGYTF TKY WLGWV RQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTT TDTSTAYMELRSLRSDDTAVYYCARSDGSSTYWG QGT LV TVSS
189	DVD1515L	AB217VL	AB048VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGW YQQKPGKAPKRLIYAAS T LQSGVPSRFSGSGSGTE FI FT ISS LQ PEDFASYYCLQHKSYPLTFGGG TK VE IKR TVAAP DVLM TQ TPLSLPVT P GE P ASISCTSSQ NIVHSNGNTYLEWYLQKPGQSPQLLIYKVS NR FSG VPDRFSGSGSGTDFTLKI S RVEAEDVGVYYCFQV S HVPYTFGGG TK VEIKR
190	DVD1516H	AB048VH	AB217VH	EVQLVQSGAEVKKPGASVKVSCKASGYTF TKY WLG WVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTL TDTSTAYMELRSLRSDDTAVYYCARSDGSSTY WGQGT LV TVSS ASTKGPSVFPLA PQVQLVESGGGV VQPGRSLRLSCAASGFTFSSYDMHWVRQAPGKGLE WVAVIWSDGSIKYYADSVKGRFTISRDN SK NTLYL QMN SL RAEDTAVYYCAREVESAMGGFYNGMDVWG QGT TV TVSS
191	DVD1516L	AB048VL	AB217VL	DVLM TQ TPLSLPVT P GE P ASISCTSSQNI V HSNGN TYLEWYLQKPGQSPQLLIYKVS NR FSGVPDRFSGS GSGTDFTLKI S RVEAEDVGVYYCFQVSHVPYTFGG G TK VEIKR TVAAP DIQMTQSPSSLSASVGDRTIT CRASQGIRIDLGWYQQKPGKAPKRLIYAAS T LQSG VPSRFSGSGSGTEFI FT ISS LQ PEDFASYYCLQHK SYPLTFGGG TK VEIKR

Example 2.36: Generation of TNF (seq. 6) and PGE2 DVD-Igs with Linker Set 3

Table 50

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
192	DVD1523H	AB218VH	AB048VH	EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFQDYWGQGLVTVVSSASTKGPSVFFLAPEVQLVQSGAEVKKPGASVKVSKASGYTFTKYWLGWVRQAPGQGLEWMDIYPGYDYTHYNEKFKDRVLTDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTYWGQGLVTVVSS
193	DVD1523L	AB218VL	AB048VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISISIQSEDFAVYYCQQYNYWWTFGQGTKVEIKRRTVAAPDVLMTQTPLSLPVTPEPASI SCTSSQNI VHSNGNTYLEWYLQKPGQSPQLLIYKVS NRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCFQVSHVPYTFGGGTKVEIKR
194	DVD1524H	AB048VH	AB218VH	EVQLVQSGAEVKKPGASVKVSKASGYTFTKYWLGWVRQAPGQGLEWMDIYPGYDYTHYNEKFKDRVLTDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTYWGQGLVTVVSSASTKGPSVFFLAPEVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFQDYWGQGLVTVVSS
195	DVD1524L	AB048VL	AB218VL	DVLMTQTPLSLPVTPEPASI SCTSSQNI VHSNGNTYLEWYLQKPGQSPQLLIYKVS NRFSGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCFQVSHVPYTFGGGTKVEIKRRTVAAPEIIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISISIQSEDFAVYYCQQYNYWWTFGQGTKVEIKR

Example 2.37: Generation of TNF (seq. 2) and LPA DVD-Igs with Linker Sct 3

Table 51

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
196	DVD1507H	AB213VH	AB216VH	QVQLKESGPGLVAPSSQLSITCTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSTDYDSTLKSRLSIS KDNSKSOIFLKMNSLQDDTARYYCAREWHHGPVA YWGQGLVTVSA AASKGFSVFP LAPQVQLQSGAE LVRPGTSVKVSCASGYGFINYLIEWIKORPGOGL EWIGLINPGSDYTNYNENFKGKATLTADKSSSTAY MHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQG TTLTVSS
197	DVD1507L	AB213VL	AB216VL	DIVMTQSHKFMSTTVGDRVSITCKASQAVSSAVAW YQKPKGQSPKLLIYWASTRHTGVPDRFTGSGSVTD FTLTIHNLQAEDLALYCCQHYSTPFTFGSGTKLE IKRT VAAP DVVMTQTPPLSLFVSLGDQASISCTSGQ SLVHINGNTYLHWYLQKPGQSPKLLIYKVSNFLFSG VPDRFSGSGSGTDFTLKISRVEAEDLGVFCSQST HFPFTFGTKLEIKR
198	DVD1508H	AB216VH	AB213VH	QVQLQSGAELVLRPGTSVKVSCASGYGFINYLIE WIKORPGQGLEWIGLINPGSDYTNYNENFKGKATL TADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYS GNYFDYWGQGTTLTVSS AASKGFSVFP LAPQVQLK ESGPGLVAPSSQLSITCTVSGFSLTDYGVNWRQP PGKLEWLGMIWGDGSTDYDSTLKSRLSISKDNSK SOIFLKMNSLQDDTARYYCAREWHHGPVAYWGQ TLVTVSA
199	DVD1508L	AB216VL	AB213VL	DVVMTQTPPLSLPVSIGDQASISCTSGQSLVHINGN TYLHWYLQKPGQSPKLLIYKVSNFLFSGVPDRFSGS SGTDFTLKISRVEAEDLGVFCSQSTHFPFTFGT GKLEIKRT VAAP DLVMTQSHKFMSTTVGDRVSIT CKASQAVSSAVAWYQKPGQSPKLLIYWASTRHTG VPDRFTGSGSVTDFTLTIHNLQAEDLALYCCQHY STPFTFGSGTKLEIKR

Example 2.38: Generation of TNF (seq. 3) and LPA DVD-Igs with Linker Set 3

Table 52

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
200	DVD1509H	AB214VH	AB216VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGLEWVAEIRSKSINSATHYAESVKGRF TISRDDS KSAVYLQMTDLRTEDTG VYYCSRNYGS TYDYWGQGTTLTVSSASTKGPSVFPLAPQVQLQQS GAELVREPGTSVKVSCASGYGFINYLIEWIKORPG QGLEWIGLINPGSDYTNYNENFKGKATLTADKSSS TAYMHLSSLTSEDSAVYFCARREFGYGSGNYFDYW GQGTTLTVSS
201	DVD1509L	AB214VL	AB216VL	DILLTQSPALLSVSPGERVSVFSCRASQFVGSIIHW YQRTNGSPRLLIKYASESMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKRTVAAPDVVMTQTPLSLPVLGDAQASISCTSGQ SLVHINGNTYLHWYLQKPGQSPKLLIYKVSNFLFSG VPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQST HFPFTFGTGTKLEIKR
202	DVD1510H	AB216VH	AB214VH	QVQLQSGAELVREPGTSVKVSCASGYGFINYLIE WIKORPGQGLEWIGLINPGSDYTNYNENFKGKATL TADKSSSTAYMHLSSLTSEDSAVYFCARREFGYGS GNYFDYWGQGTTLTVSSASTKGPSVFPLAPEVKLE ESGGGLVQPGGSMKLSCVASGFIFSNHWMNWVRQS PEKGLEWVAEIRSKSINSATHYAESVKGRFTISR DSKSAVYLQMTDLRTEDTG VYYCSRNYGTYDYW GQGTTLTVSS
203	DVD1510L	AB216VL	AB214VL	DVVMTQTPLSLPVLGDAQASISCTSGQSLVHINGN TYLHWYLQKPGQSPKLLIYKVSNFLFSGVPDRFSGS GSGTDFTLKISRVEAEDLGVYFCSQSTHFPFTFGT GTKLEIKRTVAAPDILLTQSPALLSVSPGERVSVF SCRASQFVGSIIHWYQRTNGSPRLLIKYASEMSG IPSRFSGSGSGTDFTLSINTVESEDIADYYCQESH SWPFTFGSGTNLEVKR

Example 2.39: Generation of TNF (seq. 4) and LPA DVD-Igs with Linker Set 3

Table 53

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
204	DVD1511H	AB215VH	AB216VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSYAMH WVRQAPGNGLWVAFMSYDGSNKYAKDSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDRIAAG GNYYYGMDVWGQGTITVTVSS ASTKGPSVFP LAPQ VQLQQSGAELVRPGTSVKVSKASGYGFINYLIEW IKORPGQGLEWIGLINPGSDYTNYNENFKGKATLT ADKSSSTAYMHLSSLTSEDSAVYFCARFQYGGSG NYFDYWGGQGTTLTVSS
205	DVD1511L	AB215VL	AB216VL	EIVLTQSPATLSLSPGERATLSCRASQSVSYLAW YQQKPGQAPRLLIYDASNRATGIPAREFSGSGSGTD FTLTISISLEPEDEFAVYYCQQRSNWPPFTFGPGTKV DIKRT VAA PDVMTQTPLSLPVSLGDAQISCTSG QSLVHINGNTYLHWYLQKPGQSPKLLIYKVSNFLFS GVPDRFSGSGSGTDFTLTKISRVEAEDLGVFCSQS THFPFTFGTGTKLEIKR
206	DVD1512H	AB216VH	AB215VH	QVQLQQSGAELVRPGTSVKVSKASGYGFINYLIE WIKORPGQGLEWIGLINPGSDYTNYNENFKGKATL TADKSSSTAYMHLSSLTSEDSAVYFCARFQYGGSG GNYFDYWGGQGTTLTVSS ASTKGPSVFP LAPQVQLV ESGGGVVQPGRSLRLSCAASGFTFSYAMHWVRQA PGNGLWVAFMSYDGSNKYAKDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARDRIAAGGNY YGMVWGQGTITVTVSS
207	DVD1512L	AB216VL	AB215VL	DVMTQTPLSLPVSLGDAQISCTSGQSLVHIN NTYLHWYLQKPGQSPKLLIYKVSNFLFSGVPDRFSGS SGTDFTLTKISRVEAEDLGVFCSQSTHFPFTFGT GTKLEIKRT VAA PEIVLTQSPATLSLSPGERATLS CRASQSVSYLAWYQQKPGQAPRLLIYDASNRATG IPAREFSGSGSGTDFTLTISISLEPEDEFAVYYCQQR SNWPPFTFGPGTKVDIKR

Example 2.40: Generation of TNF (seq. 1) and LPA DVD-Igs with Linker Set 3

Table 54

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
208	DVD1513H	AB017VH	AB216VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSV EGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGTTLVTVSS ASTKGPSVFP LAPQVQLQSGAELVRPGETSVKVSCKASGYGFINYLIEWIKORPGGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYGS GNYFDYWGQGTTLVTVSS
209	DVD1513L	AB017VL	AB216VL	DIQMTQSPSSLSASVGDRTITCRASQGIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFRSGSGSGTDFLTISSLQPEDVATYYCQRYNRAPYTFGQGTKVEIKR TVAAP DVVMTQTPLSLPVSLGDOASISCTSGQSLVHINGNTYLVHWYLPKPGQSPKLLIYKVSNFLFSGVPDFRFGSGSGTDFTLTKISRVEAEDLG VYFCSQSTHFPFTFGTGTKLEIKR
210	DVD1514H	AB216VH	AB017VH	QVQLQQSGAELVRPGETSVKVSCKASGYGFINYLIEWIKORPGGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYGS GNYFDYWGQGTTLVTVSS ASTKGPSVFP LAPQVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSV EGRFTISRDNKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGTTLVTVSS
211	DVD1514L	AB216VL	AB017VL	DVVMTQTPLSLPVSLGDOASISCTSGQSLVHINGNTYLVHWYLPKPGQSPKLLIYKVSNFLFSGVPDFRFGSGSGTDFTLTKISRVEAEDLG VYFCSQSTHFPFTFGTGTKLEIKR TVAAP DIQMTQSPSSLSASVGDRTITCRASQGIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFRFGSGSGTDFLTISSLQPEDVATYYCQRYNRAPYTFGQGTKVEIKR

Example 2.41: Generation of TNF (seq. 5) and LPA DVD-Igs with Linker Set 3

Table 55

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
212	DVD1521H	AB217VH	AB216VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYDMH WVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAREVESAMG GFYNGMDVWGQGT TT TVSS ASTKGPSVFPLAPQV QLQQSGAELVLRPGTSVKVSCASGYGFINYLIEWI KQRPQGLEWIGLINPGSDYTNYNENFKGKATLTA DKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGN YFDYWQGT TT TVSS
213	DVD1521L	AB217VL	AB216VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGW YQQKPGKAPKRLIYAAS T LQSGVPSRFSGSGSGTE FIFTISSLQPEDFASYCLQHKSYPLTFGGGTKVE IKRT VAAP DVVM T QTPLSLPVSLGDAQASISCTSGQ SLVHINGNTYLHWYLQKPGQSPKLLIYKVSNFLFSG VPDRFSGSGSGTDF TL KISRVEADLG V YFCSQST HFPFTFGTGT K LEIKR
214	DVD1522H	AB216VH	AB217VH	QVQLQQSGAELVLRPGTSVKVSCASGYGFINYLIE WIKQRPQGLEWIGLINPGSDYTNYNENFKGKATL TADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGS GNYFDYWQGT TT TVSS ASTKGPSVFPLAPQVQLV ESGGGVVQPGRSLRLSCAASGFTFSSYDMHWVRQA PGKGLEWVAVIWSDGSIKYYADSVKGRFTI SRDNS KNTLYLQMNSLRAEDTAVYYCAREVESAMGGFYIN GMDVWGQGT TT TVSS
215	DVD1522L	AB216VL	AB217VL	DVVM T QTPLSLPVSLGDAQASISCTSGQSLVHINGN TYLHWYLQKPGQSPKLLIYKVSNFLFSGVPDRFSGS GSGTDF TL KISRVEADLG V YFCSQSTHFPFTFGT GT K LEIKRT VAAP DIQMTQSPSSLSASVGDRTIT CRASQGIRIDLGWYQQKPGKAPKRLIYAAS T LQSG VPSRFSGSGSGTEFIFTISSLQPEDFASYCLQHK SYPLTFGGGTKVEIKR

Example 2.42: Generation of TNF (seq. 6) and LPA DVD-Igs with Linker Set 3

Table 56

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
216	DVD1529H	AB218VH	AB216VH	EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFQYWGQGT LTVSSASTKGPSVFFPLAPQVQLQQSGAELV RPTSVKVSCKASGYGFINYLIWIKQRPGQGLEW IGLINPGSDYTNYNENFKGKATLTADKSSSTAYMH LSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGT TTVSS
217	DVD1529L	AB218VL	AB216VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLSQSEDFAVYYCQQYNYWWTFGQGTKVEIKR TVAAPDVVMTQTPLSLPVSLGDAQSISCTSGQS LVHNGNTYLVHWYLPKPGQSPKLLIYKVSNFLSGV PDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTH FPFTEGTGKLEIKR
218	DVD1530H	AB216VH	AB218VH	QVQLQQSGAELVRPPTSVKVSCKASGYGFINYLIWIKQRPGQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMH LSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGT TTVSSASTKGPSVFFPLAPEVQLV ESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQA PGKGLEWVSVIYSGDRYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFQYWGQGT LTVSS
219	DVD1530L	AB216VL	AB218VL	DVVMTQTPLSLPVSLGDAQSISCTSGQSLVHNGNTYLVHWYLPKPGQSPKLLIYKVSNFLSGV PDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTH FPFTEGTGKLEIKR TVAAPEIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISLSQSEDFAVYYCQQYNYWWTFGQGTKVEIKR

Example 2.43: Generation of TNF (seq. 2) and NGF DVD-Igs with Linker Set 4

Table 57

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
220	DVD1543H	AB213VH	AB020VH	QVQLKESGPGPLVAPSQSLSTICTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSTDYDSTLKSRLSIS KDNSKSQLFLKMNSLQTDARTARYYCAREWHHGPVA YWGQGLVTVSAASTKGPSVFPLAPQVQLQESGPG LVKPSSETLSLTCTVSGFSLIGYDLNWIROPKGL EWIGI IWGDGTTDYN SAVKSRVTISKDTSKNQFSL KLSSVTAADTAVYYCARGGYWYATSYFDYWGQGT LTVSS
221	DVD1543L	AB213VL	AB020VL	DIVMTQSHKFMSTTVGDRVSICTKASQAVSSAVAW YQKPGQSPKLLIYWASTRHTGVPDRFTGSGSVTD FTLTIHNLQAEDLALYYCQQHYSYTFPTFGSGTKLE IKRTVAAPSVFI FPPDIQMTQSPSSLSASVDRVT ITCRASQSI SNNLNWYQKPGKAPKLLIYYTSRFRH SGVPSRFSGSGSDTFTFTISSLPEDIATYYCQ EHTLPYTFGQGTKLEIKR
222	DVD1544H	AB020VH	AB213VH	QVQLQESGPGPLVLPKPSSETLSLTCTVSGFSLIGYDLN WIRQPPGKGLEWIGI IWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSSVTAADTAVYYCARGGYWYATS YFDYWGQGLVTVSSASTKGPSVFPLAPQVQLKE SGPGLVAPSQSLSTICTVSGFSLTDYGVN WVRQPP GKGLEWLGMIWGDGSTDYDSTLKSRLSISKDNSKS QIFLKMNSLQTDARTARYYCAREWHHGPVAYWGQGT LTVSA
223	DVD1544L	AB020VL	AB213VL	DIQMTQSPSSLSASVDRVTITCRASQSI SNNLNW YQKPGKAPKLLIYYTSRFRHSGVPSRFSGSGSDT FTFTISSLPEDIATYYCQHEHTLPYTFGQGTKLE IKRTVAAPSVFI FPPDIVMTQSHKFMSTTVGDRVS ITCKASQAVSSAVAWYQKPGQSPKLLIYWASTRH TGVPDRFTGSGSVTDFTLTIHNLQAEDLALYYCQ HYSTPFTFGSGTKLEIKR

Example 2.44: Generation of TNF (seq. 3) and NGF DVD-Igs with Linker Set 4

Table 58

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
224	DVD1545H	AB214VH	AB020VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSKSAVYLQMTDLRTEDTGVYYCSRNYGS TYDYWGQGTTLTVSS ASTKGPSVFPLAP QVQLQES GPGLVKPSETLSLTCTVSGFSLIGYDLNWIROPPG KGLEWIGIIWGDGTTDYN SAVKSRVTISKDTSKNQ FSLKLSVTAADTAVYYCARGGYWYATSYFDYWG QGTTLTVSS
225	DVD1545L	AB214VL	AB020VL	DILLTQSPAILSVPGERVSVFSCRASQFVGSIIHW YQRTNGSPRLLIKYASEMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKRT VAAPSVFI FPPDIQMTQSPSSLSASVGDRTV ITCRASQSI SNNLNWYQQKPKAPKLLIYYTSRFH SGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQ EHTLPYTFGQGTKLEIKR
226	DVD1546H	AB020VH	AB214VH	QVQLQESGPGPKVLPSETLSLTCTVSGFSLIGYDLN WIRQPPKGLEWIGIIWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSVTAADTAVYYCARGGYWYATS YYFDYWGQGTTLTVSS ASTKGPSVFPLAPE VKLEE SGGGLVQPGGSMKLSCVASGFIFSNHWMNWVRQSP EKGLEWVAEIRSKSINSATHYAESVKGRFTISRDD SKSAVYLQMTDLRTEDTGVYYCSRNYGSTYDYWG QGTTLTVSS
227	DVD1546L	AB020VL	AB214VL	DIQMTQSPSSLSASVGDRTVITCRASQSI SNNLNW YQQKPKAPKLLIYYTSRFHSGVPSRFSGSGSGTD FTFTISSLPEDIATYYCQEHHTLPYTFGQGTKLE IKRT VAAPSVFI FPPDILLTQSPAILSVPGERVS FSCRASQFVGSIIHWYQRTNGSPRLLIKYASESM SGIPSRFSGSGSGTDFTLSINTVESEDIADYYCQE SHSWPFTFGSGTNLEVKR

Example 2.45: Generation of TNF (seq. 4) and NGF DVD-Igs with Linker Set 4

Table 59

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
228	DVD1547H	AB215VH	AB020VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMH WVRQAPGNGLWVAFMSYDGSNKYAKDSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDRGIAAG GNYYYYGMDVWGQGTITVTVSS ASTKGPSVFPLAPQ VQLQESGPGLVKPSETLSLTCTVSGFSLIGYDLNW IRQPPGKLEWIGI IWGDGTTDYN SAVKSRVTISK DTSKNQFSLKLSVTAADTAVYYCARGGYWYATSY YFDYWGQGT LVTVSS
229	DVD1547L	AB215VL	AB020VL	EIVLTQSPATLSLSPGERATLSCRASQSVYSYLAW YQQKPGQAPRLLIYDASN RATGIPAREFSGSGSGTD FTLTISSLEPEDFAVYYCQQRSNWPPFTFGPGTKV DIKRT VAAAPSVFI FPDIQMTQSPSSLSASVGD RVTITCRASQSI SNLNWYQQKPGKAPKLLIYYTSRF HSGVPSRFSGSGSGTDFTFTISSLQPEDIA TYYCQ QEHTLPYTFGQGTKLEIKR
230	DVD1548H	AB020VH	AB215VH	QVQLQESGPGLVKPSETLSLTCTVSGFSLIGYDLN WIRQPPGKLEWIGI IWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSVTAADTAVYYCARGGYWYATS YFDYWGQGT LVTVSS ASTKGPSVFPLAPQ VQLVE SGGGVVQPGRSLRLSCAASGFTFSSYAMHWVRQAP GNGLWVAFMSYDGSNKYAKDSVKGRFTISRDN SK NTLYLQMNSLRAEDTAVYYCARDRGIAAGGNYYYY GMDVWGQGTITVTVSS
231	DVD1548L	AB020VL	AB215VL	DIQMTQSPSSLSASVGD RVTITCRASQSI SNLNW YQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTD FTFTISSLQPEDIA TYYCQEHTLPYTFGQGTKLE IKRT VAAAPSVFI FPPEI VLTQSPATLSLSPGERAT LSCRASQSVYSYLAWYQQKPGQAPRLLIYDASNRA TGIPAREFSGSGSGTDFTLTISSLEPEDFAVYYCQ RSNWP PFTFGPGTKVDIKR

Example 2.46: Generation of TNF (seq. 5) and NGF DVD-Igs with Linker Set 4

Table 60

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
232	DVD1561H	AB217VH	AB020VH	QVQLVESGGGVVQGRSLRLSCAASGFTFSYDMH WVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAREVESAMG GFYNGMDVWGQGTITVTVSS ASTKGPSVFPLAPQV QLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNWI RQPPGKLEWIGLIWGDGTTDYN SAVKSRVTISKD TSKNQFSLKLSVTAADTAVYYCARGGYWATSYY FDYWGQGLTVTVSS
233	DVD1561L	AB217VL	AB020VL	DIQMTQSPSSLSASVGDRTITCRASQGI RIDLGW YQQKPGKAPKRLIYAAS TLQSGVPSRFSGSGSGTE FIFTISLQPEDFASY YCLOHKS YPLTFGGG TKVE IKR TVAAPS VFI FPP DIQMTQSPSSLSASVGDRTI ITCRASQSI SNNLNWYQQKPGKAPKLLIYYTSRFH SGVPSRFSGSGSGTDFFTISSLOPEDIATYYCQQ EHTLPYTFGQGTKLEIKR
234	DVD1562H	AB020VH	AB217VH	QVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLN WIRQPPGKLEWIGLIWGDGTTDYN SAVKSRVTIS KDTSKNQFSLKLSVTAADTAVYYCARGGYWATS YYFDYWGQGLTVTVSS ASTKGPSVFPLAPQV QLVE SGGGVVQGRSLRLSCAASGFTFSYDMHWVRQAP GKLEWVAVIWSDGSIKYYADSVKGRFTISRDN SK NTLYLQMNSLRAEDTAVYYCAREVESAMGGFYNG MDVWGQGTITVTVSS
235	DVD1562L	AB020VL	AB217VL	DIQMTQSPSSLSASVGDRTITCRASQSI SNNLNW YQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTD FTFTISSLOPEDIATYYCQEHTLPYTFGQGTKLE IKR TVAAPS VFI FPP DIQMTQSPSSLSASVGDRTI ITCRASQGI RIDLGWYQQKPGKAPKRLIYAAS TLQ SGVPSRFSGSGSGTEFIFTISSLOPEDFASY YCLO HKS YPLTFGGG TKVEIKR

Example 2.47: Generation of TNF (seq. 6) and NGF-Igs with Linker Set 4

Table 61

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
236	DVD1569H	AB218VH	AB020VH	EVQLVESGGGLIQPGGSLRRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFYWGQGLVTVSSASTKGPSVFPLAPQVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNWIRQPPGKGLEWIGLIWGDGTTDYN SAVKSRVTISKDTSKNQFSLKLSVTAADTAVYYCARGGYWYATSYFFDYWGQGLVTVSS
237	DVD1569L	AB218VL	AB020VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISSSLQSEDFAVYYCQYNYWWTFGQGTKVEIKRTVAAPSVEIFPPDIQMTQSPSSLSASVGDRTVITCRASQSVSSNLN WYQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQEHTLPYTFGQGTKLEIKR
238	DVD1570H	AB020VH	AB218VH	QVQLQESGPGLVKPSSETLSLTCTVSGFSLIGYDLNWIRQPPGKGLEWIGLIWGDGTTDYN SAVKSRVTISKDTSKNQFSLKLSVTAADTAVYYCARGGYWYATSYFFDYWGQGLVTVSSASTKGPSVFPLAPEVQLVESGGGLIQPGGSLRRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFYWGQGLVTVSS
239	DVD1570L	AB020VL	AB218VL	DIQMTQSPSSLSASVGDRTVITCRASQSVSSNLN WYQQKPGKAPKLLIYYTSRFHSGVPSRFSGSGSGTDFTFTISSLPEDIATYYCQQEHTLPYTFGQGTKLEIKRTVAAPSVEIFPPEIVMTQSPATLSVSPGERATLSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISSSLQSEDFAVYYCQYNYWWTFGQGTKVEIKR

Example 2.48: Generation of TNF (seq. 2) and SOST DVD-Igs with Linker Set 4

Table 62

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
240	DVD1537H	AB213VH	AB022VH	QVQLKESGPGLVAPSQSLSICTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSTDYDSTLKSRLSIS KDNSKSIIFLKMNSLQTDDTARYYCAREWHHGPPVA YWGQGLVTVSA AASTKGPSVFFPLAPE VQLQQSGPE LVTPGASVKISCKASGYFTFDHYMSWVKQSHGKSL EWIGDINPYSGETTYNQKFKGTATLTVDKSSSIAY MEIRGLTSEDSAVYYCARDYDASPFAYWGQGLV TVSA
241	DVD1537L	AB213VL	AB022VL	DIVMTQSHKFMSTTVGDRVSICTCKASQAVSSAVAW YQKPGQSPKLLIYWASTRHTGVPDRFTGSGSVTD FTLTIHNLQAEDLALYYCQGHYSTPFTFGSGTKLE IKR TVAAPSVFIFPPD VQMIQSPSSLSASLGDIVT MTCQASQGT SINLNWFQKPGKAPKLLIYSSNLE DGVPSRFSGSRYGTDFTLTISSLEDEDLATYFCLQ HSYLPYTFGGGTKLEIKR
242	DVD1538H	AB022VH	AB213VH	EVQLQQSGPELVTPGASVKISCKASGYFTFDHYMS WVKQSHGKSLWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGLVTVSA AASTKGPSVFFPLAP QVQLKESG PGLVAPSQSLSICTVSGFSLTDYGVNWRQPPGK GLEWLGMIWGDGSTDYDSTLKSRLSISKDNSKSI FLKMNSLQTDDTARYYCAREWHHGPPVAYWGQGLV TVSA
243	DVD1538L	AB022VL	AB213VL	DVQMIQSPSSLSASLGDIVTMTQASQGT SINLNW FQKPGKAPKLLIYSSNLEDGVPSRFSGSRYGTD FTLTISSLEDEDLATYFCLQHSYLPYTFGGGTKLE IKR TVAAPSVFIFPPD IVMTQSHKFMSTTVGDRVS ICTCKASQAVSSAVAWYQKPGQSPKLLIYWASTRH TGVPDRFTGSGSVTDFTLTIHNLQAEDLALYYCQ GHYSTPFTFGSGTKLEIKR

Example 2.49: Generation of TNF (seq. 3) and SOST DVD-Igs with Linker Set 4

Table 63

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
244	DVD1539H	AB214VH	AB022VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSKSAVYLQMTDLRTEDTGVYYCSRNYGGS TYDYWGQGTTLTVSS ASTKGPSVFPLAPEVQLQGS GPELVTPGASVKISCKASGYFTFDHYMSWVKQSHG KSLEWIGDINPYSGETTYNQKFKGTATLTVDKSSS IAYMEIRGLTSEDSAVYYCARDYDASPFAYWGQG TLTVSA
245	DVD1539L	AB214VL	AB022VL	DILLTQSPAILSVSFGERVVSFCRASQFVGSIIHW YQRTNGSPRLLIKYASEMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKR VAAPSVFIFPPDVQMIQSESSLSASLGDIVT MTCQASQGTSINLNWFQQKPGKAPKLLIYGSSNLE DGVPSRFSGSRYGTDFTLTISSELEDEDLATYFCLQ HSYLPYTFGGGTKLEIKR
246	DVD1540H	AB022VH	AB214VH	EVQLQQSGPELVTPGASVKISCKASGYFTFDHYMS WVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGTTLTVSA ASTKGPSVFPLAPEV KLEESG GGLVQPGGSMKLSCVASGFIFSNHWMNWVRQSPK GLEWVAEIRSKSINSATHYAESVKGRFTISRDDSK SAVYLQMTDLRTEDTGVYYCSRNYGGSYDYWGQG TTLTVSS
247	DVD1540L	AB022VL	AB214VL	DVQMIQSPSSLSASLGDIVTMTTCQASQGTSINLNW FQQKPGKAPKLLIYGSSNLEEDGVPSRFSGSRYGTD FTLTISSELEDEDLATYFCLQHSYLPYTFGGGTKLE IKR VAAPSVFIFPPDILLTQSPAILSVSFGERVS FSCRASQFVGSIIHWYQRTNGSPRLLIKYASESM SGIPSRFSGSGSGTDFTLSINTVESEDIADYYCQE SHSWPFTFGSGTNLEVKR

Example 2.50: Generation of TNF(seq. 4) and SOST DVD-Igs with Linker Set 4

Table 64

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
240	DVD1541H	AB215VH	AB022VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFS SYAMH WVRQAPGNGLLEWVAFMSYDGSNKYAKDSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDRIAAG GNYYYYGMDVWGQGT TTVTVSS ASTKGPSVFPLAPE VQLQQSGPELVT PGASVKISCKASGYTFTDHYMSW VKQSHGKSLEWIGDINPYSGETTYNQKFKGTATLT VDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP AYWGQGT LVTVSA
249	DVD1541L	AB215VL	AB022VL	EIVLTQSPATLSLSPGERATLSCRASQSVSYLAW YQQKPGQAPRLLIYDASNRA TGI PARFSGSGSGTD FTLTISSLEPEDEFAVYYCQQRSNWPPFTFGPGTKV DIKRT VAAAPSVFIFPP DVQMIQSPSSLSASLGDIV TMTQASQGTSINLNWFQKPGKAPKLLIYGSSNL EDGVPSRFSGSRYGTDFTLTISSLEDEDLATYFCL QHSYLPYTFGGGTKLEIKR
250	DVD1542H	AB022VH	AB215VH	EVQLQQSGPELVT PGASVKISCKASGYTFTDHYMS WVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGT LVTVSA ASTKGPSVFPLAP QVQLVESG GGVVQPGRSLRLSCAASGFTFS SYAMHWVRQAPGN GLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNT LYLQMNSLRAEDTAVYYCARDRIAAGGNYYYYGM DVWGQGT TTVTVSS
251	DVD1542L	ABC22VL	AB215VL	DVQMIQSPSSLSASLGDIVTMTQASQGTSINLNW FQKPGKAPKLLIYGSSNLEDGVPSRFSGSRYGTD FTLTISSLEDEDLATYFCLQHSYLPYTFGGGTKLE IKRT VAAAPSVFIFPPE EIVLTQSPATLSLSPGERAT LSCRASQSVSYLAWYQQKPGQAPRLLIYDASNRA TGI PARFSGSGSGTDFTLTISSLEPEDEFAVYYCQ RSNWPPFTFGPGTKVDIKR

Example 2.51: Generation of TNF (seq. 5) and SOST DVD-Igs with Linker Set 4

Table 65

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
252	DVD1559H	AB217VH	AB022VH	QVQLVESGGGVVQPRSLRLSCAASGFTFSYDMH WVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCAREVESAMG GFYNGMDVWGQGTITVTVSS ASTKGPSVFPLA PEV QLQQSGPELVTPGASVKISCKASGYFTFDHYMSWV KQSHGKSLEWIGDINPYSGETTYNQKFKGTATLTV DKSSSIAYMEIRGLTSEDSAVYYCARDYDASPFA YWGQGLVTVSA
253	DVD1559L	AB217VL	AB022VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGW YQQKPGKAPKRLIYAASLTQSGVPSRFSGSGSGTE FIFTISSLQPEDFASYCYLQHKSYPLTFGGGTKVE IKR TVAAPSVFIFPP DVQMIQSPSSLSASLGDIVT MTCQASQGTSLNINWFQQKPGKAPKLLIYGSSNLE DGVPSRFSGSRYGTDFTLTISSLEDEDLATYFCLQ HSYLPYTFGGGTKLEIKR
254	DVD1560H	AB022VH	AB217VH	EVQLQQSGPELVTPGASVKISCKASGYFTFDHYMS WVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAYWGQGLVTVSA ASTKGPSVFPLAP QVQLVESG GGVVQPRSLRLSCAASGFTFSYDMHWVRQAPGK GLEWVAVIWSDGSIKYYADSVKGRFTISRDNKNT LYLQMNSLRAEDTAVYYCAREVESAMGGFYNGMD VWGQGTITVTVSS
255	DVD1560L	AB022VL	AB217VL	DVQMIQSPSSLSASLGDIVTMTTCQASQGTSLNINW FQQKPGKAPKLLIYGSSNLEDGVPSRFSGSRYGTD FTLTISSLEDEDLATYFCLQHSYLPYTFGGGTKLE IKR TVAAPSVFIFPP DIQMTQSPSSLSASVGDRT ITCRASQGIRIDLGWYQQKPGKAPKRLIYAASLTQ SGVPSRFSGSGSGTEFIFTISSLQPEDFASYCYLQ HKSYPYTFGGGTKVEIKR

Example 2.52: Generation of TNF (seq. 6) and SOST DVD-Igs with Linker Set 4

Table 66

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
256	DVD1567H	AB218VH	AB022VH	EVQLVESGGGLIQPGGSLRRLSCAASGFTVSRNYMS WVRQAPGKGLEWVSVIYSGDRITYYADSVKGRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCARGEGGFYD WGQGLTLVTVSSASTKGPSVFPLAPEVQLQSGPELV TPGASVKISCKASGYTFTDHYMSWVKQSHGKSLEW IGDINPYSGETTYNQKFKGTATLTVDKSSSIAYME IRGLTSEDSAVYYCARDYDASPFAWGGQGLVTV SA
257	DVD1567L	AB218VL	AB022VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAW YQQKPGQAPRLLIHGASIRATGLPARFSGSGSGTE FTLTISLQSEDFAVYYCQYNYWTFGQGTKVEI KRTVAAPSVEIFPPDVQMIQSPSSLSASLGDIVTM TCQASQGTSLNINWFQKPKGKAPKLLIYGSSNLED GVPSRFSGSRYGTDFTLTISLSEDEDLATYFCLQH SYLPYTFGGGTKLEIKR
258	DVD1568H	AB022VH	AB218VH	EVQLQDSGPELVTPGASVKISCKASGYTFTDHYMS WVKQSHGKSLEWIGDINPYSGETTYNQKFKGTATL TVDKSSSIAYMEIRGLTSEDSAVYYCARDYDASP FAWGGQGLVTVSSASTKGPSVFPLAPEVQLVESG GGLIQPGGSLRRLSCAASGFTVSRNYMSWVRQAPGK GLEWVSVIYSGDRITYYADSVKGRFTISRDNKNTL YLQMNSLRAEDTAVYYCARGEGGFYDYGQGLVTV SS
259	DVD1568L	ABC22VL	AB218VL	DVQMIQSPSSLSASLGDIVTMTQASQGTSLNINW FQKPKGKAPKLLIYGSSNLEDGVPSRFSGSRYGTD FTLTISLSEDEDLATYFCLQHSYLPYTFGGGTKLE IKRTVAAPSVEIFPPEIIVMTQSPATLSVSPGERAT LSCRASQSVSSNLAWYQQKPGQAPRLLIHGASIRA TGLPARFSGSGSGTEFTLTISLQSEDFAVYYCQY NYWTFGQGTKVEIKR

Example 2.53: Generation of TNF (seq. 2) and PGE2 with Linker Set 4

Table 67

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				1234567890123456789012345678901234
260	DVD1531H	AB213VH	AB048VH	QVQLKESGPGLVAPSQSLSI TCTVSGFSLTDYGV NWVRQPPGKLEWLGMIWGDGSTDYDSTLKSRLS ISKDNSKSI FLKMNSLQTD TARYYCAREWHHG PVAYWGQGLVTVSA ASTKGPSVFPLAPEVQLVQ SGAEVKKPGASVKVCKASGYTFTKYWLGWVRQA PGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTDT STSTAYMELRSLRSDDTAVYYCARSDGSSTYWGQ GTLVTVSS
261	DVD1531L	AB213VL	AB048VL	DIVMTQSHKFMSTTVGDRVSI TCKASQAVSSAVA WYQQKPGQSPKLLIYWASTRHTGVPDRFTGSGSV TDFTLTIHNLQAEDLALYCCQHYSTPFTFGSGT KLEIKR TVAAPSVFIFPPD VLMTQTPLSLPVT EPASISCTSSQNI VHSNGNTYLEWYLQKPGQSPQ LLIYKVSNRFSGVPDRFSGSGSDFTLTKISRVE AEDVGVYYCFQVSHVPYTFGGGKVEIKR
262	DVD1532H	AB048VH	AB213VH	EVQLVQSGAEVKKPGASVKVCKASGYTFTKYWL GWVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRV TLTDTSTSTAYMELRSLRSDDTAVYYCARSDGS STYWGQGLVTVSS ASTKGPSVFPLAPQVQLKES GGLVAPSQSLSI TCTVSGFSLTDYGVNWVRQPP GKLEWLGMIWGDGSTDYDSTLKSRLSISKDNSK SQIFLKMNSLQTD TARYYCAREWHHG PVAYWGQ GTLVTVSA
263	DVD1532L	AB048VL	AB213VL	DVLMTQTFPLSLPVTTPGEPASISCTSSQNI VHSNG NTYLEWYLQKPGQSPQLLIYKVSNRFSGVPDRFS GSGSDFTLTKISRVEAEDVGVYYCFQVSHVPYT FGGKVEIKR TVAAPSVFIFPPD IVMTQSHKFM STTVGDRVSI TCKASQAVSSAVAWYQQKPGQSPK LLIYWASTRHTGVPDRFTGSGSVTDFTLTIHNLQ AEDLALYCCQHYSTPFTFGSGTKLEIKR

Example 2.54: Generation of TNF (seq. 3) and PGE2 DVD-Igs with Linker Set 4

Table 68

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
264	DVD1533H	AB214VH	AB048VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSKSAVYLQMTDLRTEDTGVIYCSRNYYS TYDYWGQGTTLTVSSASTKGPSVFPLAPEVQLVQS GAEVKKPGASVKVSKASGYFTTKYWLGWVRQAPG QGLEWMGDIYPGYDYTHYNEKFKDRVTLTIDTSTS TAYMELRSLRSDDTAVYYCARSDGSSTYWGQGTLV TVSS
265	DVD1533L	AB214VL	AB048VL	DILLTQSPAILSVSPGERVSFSCRASQFVGSIIHW YQORTNGSPRLLIKYASEMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKR TVAAPSVEIFPPD VLMTQTPLSLPVTPEGPAS ISCTSSQNIVHNSNGNTYLEWYLQKPGQSPQLLIYK VSNRFSGVDPDRFSGSGSGTDFTLTKISRVEAEDVGV YYCFQVSHVPYTFGGTKVEIKR
266	DVD1534H	AB048VH	AB214VH	EVQLVQSGAEVKKPGASVKVSKASGYFTTKYWL WVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTL TTDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTY WGQGTLVTVSSASTKGPSVFPLAPEVKLEESGGGL VQPGGSMKLSCVASGFIFSNHWMNWRQSPKGLE WVAEIRSKSINSATHYAESVKGRFTISRDDSKSAV YLQMTDLRTEDTGVIYCSRNYGTYDYWGQGTTL TVSS
267	DVD1534L	AB048VL	AB214VL	DVLMTQTPLSLPVTPEGPASISCTSSQNIVHNSNGN TYLEWYLQKPGQSPQLLIYKVSNRFSGVDPDRFSGS GSGTDFTLTKISRVEAEDVGVYYCFQVSHVPYTFGG GTKVEIKR TVAAPSVEIFPPD DILLTQSPAILSVSP GERVSFSCRASQFVGSIIHWYQORTNGSPRLLIKY ASEMSGIPSRFSGSGSGTDFTLSINTVESEDIAD YYCQESHWPFTFGSGTNLEVKR

Example 2.55: Generation of TNF (seq. 4) and PGE2 DVD-Igs with Linker Set 4

Table 69

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
268	DVD1535H	AB215VH	AB048VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYAMH WVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDRGIAAG GNYYYYGMDVWGQGT TV TVSS ASTKGPSVFPLAPE VQLVQSGAEVKKPGASVKVCKASGYTFTKYWLGW VRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTLT TDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTYW GQGT LV TVSS
269	DVD1535L	AB215VL	AB048VL	EIVLTQSPATLSLSPGERATLSCRASQSVYSYLAW YQQKPGQAPRLLIYDASNRTGIPARFSGSGSGTD FTLT ISSLEPE DFAVYYCQQRSNWPPFTFGPGTKV DIKRT VVAAPSVFIFPP DVLMTQTPLSLPVTPGEP ASISCTSSQNIVHNSNGNTYLEWYLQKPGQSPQLLIY KVSNRFSGV PDR FSGSGSGTDFTLTKISRVEAEDVG VYYCFQVSHVPYTFGGTKVEIKR
270	DVD1536H	AB048VH	AB215VH	EVQLVQSGAEVKKPGASVKVCKASGYTFTKYWLG WVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTL TDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTY WGQGT LV TVSS ASTKGPSVFPLAP QVQLVESGGGV VQPGRSLRLSCAASGFTFSSYAMHWVRQAPGNGLE WVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYL QMNSLRAEDTAVYYCARDRGIAAGGNYYYYGMDVW GQGT TV TVSS
271	DVD1536L	AB048VL	AB215VL	DVLMTQTPLSLPVTPGEPASISCTSSQNIVHNSNG NTYLEWYLQKPGQSPQLLIYKVSNRFSGV PDR FSGS GSGTDFTLTKISRVEAEDVGYYCFQVSHVPYTFGG GKVEIKR VVAAPSVFIFPPE IIVLTQSPATLSLSP GERATLSCRASQSVYSYLAWYQQKPGQAPRLLIYD ASNRTGIPARFSGSGSGTDFTLT ISSLEPE DFAV YYCQQRSNWPPFTFGPGTKVDIKR

Example 2.56: Generation of TNF (seq. 5) and PGE2 DVD-Igs with Linker Set 4

Table 70

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
272	DVD1557H	AB217VH	AB048VH	QVQLVESGGGVVQPCRSLRLSCLASGFTFSYDMHWVRQAPGKGLWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGGFYNGMDVWGQGTITVTVSS ASTKGPSVFPLA PEVQLVQSGAEVKKPGASVKVSCKASGYTFTKYWLGWVRRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTYWGQGTITVTVSS
273	DVD1557L	AB217VL	AB048VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGWYQKPKGKAPKRLIYAASLTQSGVPSRFSGSGSGTEFIFTISLQPEDFASYCLQHKSYPLTFGGGKVEIKR TVAAPSVFI FPDVLMTQTPLSLPVTPGEPASISCTSSQNIIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCFQVSHVPYTFGGGKVEIKR
274	DVD1558H	AB048VH	AB217VH	EVQLVQSGAEVKKPGASVKVSCKASGYTFTKYWLGWVRQAPGQGLEWMGDIYPGYDYTHYNEKFKDRVTLTDTSTSTAYMELRSLRSDDTAVYYCARSDGSSTYWGQGTITVTVSS ASTKGPSVFPLA PQVQLVESGGGVVQPCRSLRLSCLASGFTFSYDMHWVRQAPGKGLWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGGFYNGMDVWGQGTITVTVSS
275	DVD1558L	AB048VL	AB217VL	DVIMTQTPLSLPVTPGEPASISCTSSQNIIVHSNGNTYLEWYLQKPGQSPQLLIYKVSNRFGVPPDRFSGSGSGTDFTLKISRVEAEDVGVYYCFQVSHVPYTFGGGKVEIKR TVAAPSVFI FPDQMTQSPSSLSASVGDRTITCRASQGIRIDLGWYQKPKGKAPKRLIYAASLTQSGVPSRFSGSGSGTEFIFTISLQPEDFASYCLQHKSYPLTFGGGKVEIKR

Example 2.57: Generation of TNF (seq. 6) and PGE2 DVD-Igs with Linker Set 4

Table 71

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
276	DVD1565H	AB218VH	AB048VH	EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMS WVRQAPGKGLEWVSVIYSGDRTYYADSVKGRFTIS RDNSKNTLYLQMNLSLRAEDTAVYYCARGEGGFQYD WGQGLVTVSSASTKGPSVFFLAPEVQLVQSGAEVK KPGASVKVSKASGYFTFTKYWLGWVRQAPGQGLEW MGDIYPGYDYTHYNEKFKDRVTLTDTSTSTAYME LRSLSRSDDTAVYYCARSDGSSTYWGQGLVTVSS
277	DVD1565L	AB218VL	AB048VL	EIVMTQSPATLSVSPGERATLSCRASQSVSSNLAW YQKPGQAPRLLIHGASIRATGLPARFSGSGSGTE FTLTISSLOSEDFAVYYCQYNYWWTFGQGTKVEI KR TVAAPSVFIFPP DVLMTQTPLSLPVTPGEPASI SCTSSQNIIVHSNGNTYLEWYLQKPGQSPQLLIYKV SNRFGVDPDRFSGSGSGTDFTLKISRVEAEDVGVY YCFQVSHVPTYFGGGTKVEIKR
278	DVD1566H	AB048VH	AB218VH	EVQLVQSGAEVKKPGASVKVSKASGYFTFTKYWLG WVRQAPGQGLEWMDIYPGYDYTHYNEKFKDRVTL TDTSTSTAYMELRSLSRSDDTAVYYCARSDGSSTY WGQGLVTVSSASTKGPSVFFLAPEVQLVESGGGL IQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLE WVSVIYSGDRYYADSVKGRFTISRDNKNTLYLQ MNSLRAEDTAVYYCARGEGGFQYWGQGLVTVSS
279	DVD1566L	AB048VL	AB218VL	DVLMTQTPLSLPVTPGEPASISCTSSQNIIVHSNGN TYLEWYLQKPGQSPQLLIYKVSNRFGVDPDRFSGS GSGTDFTLKISRVEAEDVGVYCFQVSHVPTYFGG GTKVEIKR TVAAPSVFIFPP EIVMTQSPATLSVSP GERATLSCRASQSVSSNLAWYQKPGQAPRLLIHG ASIRATGLPARFSGSGSGTEFTLTISSLQSEDFAV YYCQYNYWWTFGQGTKVEIKR

Example 2.58: Generation of TNF (seq. 2) and LPA DVD-Igs with Linker Set 4

Table 72

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
280	DVD1549H	AB213VH	AB216VH	QVQLKESGPGFLVAPSQSL SITCTVSGFSLTDYGVN WVRQPPGKGLEWLGMIWGDGSDYDSTLKSRLSIS KDNSKSIQIFLKMNSLQDDTARYYCAREWHHGPVA YWGQGLVTVSA ASTKGPSVFPLAPQ VQLQSGAE LVRPGTSVKVSKASGYGFINYLIEWIKORPGOGL EWIGLINPGSDYTNYNENFKGKATLTADKSSSTAY MHLSSLTSEDSAVYFCARFRGYGSGNYFDYWGQG TTLTVSS
281	DVD1549L	AB213VL	AB216VL	DIVMTQSHKFMSTTVGDRVSITCKASQAVSSAVAW YQQKPGQSPKLLIYWASTRHTGVPDRFTGSGSVTD FTLTIHNLQAEDLALYYCQHYSTPFTFGSGTKLE IKRT VAAPSVEIFPPD VVMTQTPFLSLPVS LGDQAS ISCTSGQSLVHINGNTYLHWYLQKPGQSPKLLIYK VSNLFSGVDRFSGSGSGTDFTLTKISRVEADLGV YFCSQSTHFPFTFGTKLEIKR
282	DVD1550H	AB216VH	AB213VH	QVQLQSGAELVLRPGTSVKVSKASGYGFINYLIE WIKORPGGLEWIGLINPGSDYTNYNENFKGKATL TADKSSSTAYMHLSSLTSEDSAVYFCARFRGYGSG GNYFDYWGQTTTLTVSS ASTKGPSVFPLAPQ VQLK ESGPGFLVAPSQSL SITCTVSGFSLTDYGVN WVRQP PGKGLEWLGMIWGDGSDYDSTLKSRLSISKDNSK SIQIFLKMNSLQDDTARYYCAREWHHGPVAYWGQG TLVTVSA
283	DVD1550L	AB216VL	AB213VL	DVVMTQTPFLSLPVS LGDQASISCTSGQSLVHINGN TYLHWYLQKPGQSPKLLIYK VSNLFSGVDRFSGS GSGTDFTLTKISRVEADLGVYFCSQSTHFPFTFGT GTKLEIKRT VAAPSVEIFPPD IVMTQSHKFMSTTV GDRVSITCKASQAVSSAVAWYQQKPGQSPKLLIYW ASTRHTGVPDRFTGSGSVTDFTLTIHNLQAEDLAL YYCQHYSTPFTFGSGTKLEIKR

Example 2.59: Generation of TNF (seq. 3) and LPA DVD-Igs with Linker Set 4

Table 73

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
284	DVD1551H	AB214VH	AB216VH	EVKLEESGGGLVQPGGSMKLSCVASGFIFSNHWMN WVRQSPKGLEWVAEIRSKSINSATHYAESVKGRF TISRDDSKSAVYLQMTDLRTEDTGVYYCSRNYGS TYDYWGQGTTLTVSS ASTKGPSVFPLA PQVQLQQS GAELVLRPGTSVKVSKASGYGFINYLIEWIKRPG QGLEWIGLINPGSDYTNYNENFKGKATLTADKSSS TAYMHLSSLTSEDSAVYFCARREGYYGSGNYFDYW GQGTTLTVSS
285	DVD1551L	AB214VL	AB216VL	DILLTQSPAILSVSPGERVSVFSCRASQFVGSIIHW YQRTNGSPRLLIKYASEMSGIPSRFSGSGSGTD FTLSINTVESEDIADYYCQESHWPFTFGSGTNLE VKR TVAAPSVEIFPP DVVMVTQTPLSLPVSLGDQAS ISCTSGQSLVHINGNTYLHWYLQKPGQSPKLLIYK VSNLFSGVPDFRFGSGSGTDFTLKI SRVEAEDLGV YFCSQSTHFPTFGTGTGLEIKR
286	DVD1552H	AB216VH	AB214VH	QVQLQQSGAELVLRPGTSVKVSKASGYGFINYLIE WIKRPGQGLEWIGLINPGSDYTNYNENFKGKATL TADKSSSTAYMHLSSLTSEDSAVYFCARREGYYGS GNYFDYWQGTTLTVSS ASTKGPSVFPLA PEVKLE ESGGGLVQPGGSMKLSCVASGFIFSNHWMNWVRQS PEKGLEWVAEIRSKSINSATHYAESVKGRFTISR DSKSAVYLQMTDLRTEDTGVYYCSRNYGSTYDYW GQGTTLTVSS
287	DVD1552L	AB216VL	AB214VL	DVVMVTQTPLSLPVSLGDQASISCTSGQSLVHINGN TYLHWYLQKPGQSPKLLIYKVSINLFSGVPDFRFGS GSGTDFTLKI SRVEAEDLGVYFCSQSTHFPTFGT GTGLEIKR TVAAPSVEIFPP DILLTQSPAILSVSP GERVSVFSCRASQFVGSIIHWYQRTNGSPRLLIKY ASEMSGIPSRFSGSGSGTDFTLSINTVESEDIAD YYCQESHWPFTFGSGTNLEVKR

Example 2.60: Generation of TNF (seq. 4) and LPA DVD-Igs with Linker Set 4

Table 74

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
288	DVD1553H	AB215VH	AB216VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSYAMHWVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDRIAAGGNYYYGMDVWGQGTITVTVSS ASTKGPSVFPLAPQVQLQSGAELVLRPGTSVKVCSKASGYGFINYLIEWIKQRPQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARF'GYGSGNYFDYWGQGTITLVSS
289	DVD1553L	AB215VL	AB216VL	EIVLTQSPATLSLSPGERATLSCRASQSVYSYLAWYQQKPGQAPRLLIYDASNRTGIPARFSGSGSGTDFTLTISISLEPEDEFAVYYCQQRSNWPPFTFGEGTKVDIKR TVAAPSVFIFPPD VVMTQTPLSLPVSLGDOASISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVSINLFSGVPDFRFSGSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR
290	DVD1554H	AB216VH	AB215VH	QVQLQSGAELVLRPGTSVKVCSKASGYGFINYLIEWIKQRPQGLEWIGLINPGSDYTNYNENFKGKATLTADRSSSTAYMHLSSLTSEDSAVYFCARF'GYGSGNYFDYWGQGTITLVSS ASTKGPSVFPLAPQVQLVESGGGVVQPGRSLRLSCAASGFTFSYAMHWVRQAPGNGLEWVAFMSYDGSNKYAKDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARDRIAAGGNYYYGMDVWGQGTITVTVSS
291	DVD1554L	AB216VL	AB215VL	DVVMTQTPLSLPVSLGDOASISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVSINLFSGVPDFRFSGSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR TVAAPSVFIFPPEI VLTQSPATLSLSPGERATLSCRASQSVYSYLAWYQQKPGQAPRLLIYDASNRTGIPARFSGSGSGTDFTLTISISLEPEDEFAVYYCQQRSNWPPFTFGPGTKVDIKR

Example 2.61: Generation of TNF (seq. 1) and LPA DVD-Igs with Linker Set 4

Table 75

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence 12345678901234567890123456789012345
292	DVD1555H	AB017VH	AB216VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGTLLVTVSSASTKGPSVFPLAPQVQLQSGAELVRPGETSVKVSCKASGYGFINYLIEWIKQRPQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGTLLTVSS
293	DVD1555L	AB017VL	AB216VL	DIQMTQSPSSLSASVGRVTITCRASQGIIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNRAPYTFGQGTKVEIKR TVAAPSVFIFPPD VVMTQTPPLSLPVSLGDAQASISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVSNFLFSGVPDFRFGSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR
294	DVD1556H	AB216VH	AB017VH	QVQLQQSGAELVRPGETSVKVSCKASGYGFINYLIEWIKQRPQGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGTLLTVSSASTKGPSVFPLAPEVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGTLLTVSS
295	DVD1556L	AB216VL	AB017VL	DVVMTQTPPLSLPVSLGDAQASISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVSNFLFSGVPDFRFGSGSGTDFTLTKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR TVAAPSVFIFPPDI QMTQSPSSLSASVGRVTITCRASQGIIRNYLAWYQQKPGKAPKLLIYAASLTQSGVPSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNRAPYTFGQGTKVEIKR

Example 2.62: Generation of TNF (seq. 5) and LPA DVD-Igs with Linker Set 4

Table 76

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
296	DVD1563H	AB217VH	AB216VH	QVQLVESGGGVVQPGRSLRLSCAASGFTFSYDMHWVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGFFYYNGMDVWGQGTITVTVSS ASTKGPSVFPLAPQV QLQQSGAELVLRPGTSVKVSCKASGYGFINYLIEWIKORPGGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGTITLVSS ASTKGPSVFPLAPQVQLV ESGGGVVQPGRSLRLSCAASGFTFSYDMHWVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGFFYYNGMDVWGQGTITVTVSS
297	DVD1563L	AB217VL	AB216VL	DIQMTQSPSSLSASVGDRTITCRASQGIRIDLGWYQQKPKGKAPKRLIYAASLTQSGVPSRFSGSGSGTEFIFTISSLPEDFASYYCLQHKSYPLTFGGGTKVEIKR TVAAPSVFI FPDVMVTQTPLSLPVSLGDAQSISCTSGQSLVHNGNTYLHWYLQKPGQSPKLLIYKVSNFLFSGVPSRFSGSGSGTEFTLKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR
298	DVD1564H	AB216VH	AB217VH	QVQLQSGAELVLRPGTSVKVSCKASGYGFINYLIEWIKORPGGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYYGSGNYFDYWGQGTITLVSS ASTKGPSVFPLAPQVQLV ESGGGVVQPGRSLRLSCAASGFTFSYDMHWVRQAPGKGLEWVAVIWSDGSIKYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAREVESAMGFFYYNGMDVWGQGTITVTVSS
299	DVD1564L	AB216VL	AB217VL	DVVMVTQTPLSLPVSLGDAQSISCTSGQSLVHNGNTYLHWYLQKPGQSPKLLIYKVSNFLFSGVPSRFSGSGSGTEFTLKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR TVAAPSVFI FPDVMVTQTPLSLPVSLGDAQSISCTSGQSLVHNGNTYLHWYLQKPGQSPKLLIYKVSNFLFSGVPSRFSGSGSGTEFTLKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR TVAAPSVFI FPDVMVTQTPLSLPVSLGDAQSISCTSGQSLVHNGNTYLHWYLQKPGQSPKLLIYKVSNFLFSGVPSRFSGSGSGTEFTLKISRVEAEDLGVYFCSQSTHFPFTFGTGTKLEIKR

Example 2.63: Generation of TNF (seq. 6) and LPA DVD-Igs with Linker Set 4

Table 77

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
300	DVD1571H	AB218VH	AB216VH	EVQLVESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFQDWGQGT LVTVSS ASTKGPSVFPLAPQVQLQSGAELV RPTSVKVSCKASGYGFINYLEWIKQRPGGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYGGNYFDYWGQGT
301	DVD1571L	AB218VL	AB216VL	EIVMTQSPATLSVSPGERATLSCRASQSVSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISSLQSEDFAVYYCQYNYWWTFGQGTKVEIKR VAAPSVFIFPPD VVMTQTPLSLPVSLGDQASISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVS NLFSGVDPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTHFPFTFGTKLEIKR
302	DVD1572H	AB216VH	AB218VH	QVQLQSGAELV RPTSVKVSCKASGYGFINYLEWIKQRPGGLEWIGLINPGSDYTNYNENFKGKATLTADKSSSTAYMHLSSLTSEDSAVYFCARRFGYGGNYFDYWGQGTTLTVSS ASTKGPSVFPLAPEVQLV ESGGGLIQPGGSLRLSCAASGFTVSRNYMSWVRQAPGKGLEWVSVIYSGDRYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARGEGGFQDWGQGT LVTVSS
303	DVD1572L	AB216VL	AB218VL	DVVMTQTPLSLPVSLGDQASISCTSGQSLVHINGNTYLNHWYLQKPGQSPKLLIYKVS NLFSGVDPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTHFPFTFGTKLEIKR VAAPSVFIFPPE IIVMTQSPATLSVSPGERATLSCRASQSVSNLAWYQKPGQAPRLLIHGASIRATGLPARFSGSGSGTEFTLTISSLQSEDFAVYYCQYNYWWTFGQGTKVEIKR

Example 2.64: Cloning Vector Sequences Used to Clone Parent Antibody and DVD-Ig Sequences

Table 78

5

SEQ ID NO	Vector name	Nucleotide sequences
304	V1	123456789012345678901234567890123456789012345678901 GCCTCGACCAAGGCCCATCGGTCTTCCCCCTGGCACCCCTCTCCAAGAGC ACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCC GAACCGGTGACCGTGTCTGTGGAACCTCAGGCCCTGACCAGCGCGTGCAC ACCTTCCCGGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGT GTGACCGTGCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTG AATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCT TGTGACAAAACCTACACATGCCACCGTGGCCAGCACCTGAACCTCTGGGG GGACCGTCAGTCTTCTTCCCCCAAACCAAGGACACCCCTCATGATC TCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGAC CCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCC AAGACAAAGCCCGGGGAGGAGTACAACAGCACGTACCGTGTGGTCAAGT GTCCCTCACCGTCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC AAGGTCTCCAACAAGCCCTCCAGCCCCATCGAGAAACCATCTCCGAAA GCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCTGCCCCATCCCGC GAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGCTTC TATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGGAAAC AACTACAAGACCAGCCTCCCGTGGTGGACTCCGACGGCTCTTCTTCTC TACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCACAGGGAACGTC TCATGCTCCGTGATGCATGAGGCTCTGCACAACCCTACAGCAGAAGAGC CTCTCCCTGTCTCCGGGTAATGAGCGGCCGCTCGAGGCCGGCAAGGCCGG ATCCCCGACCTCGACCTCTGGCTAATAAAGGAAATTTATTTTCATTGCAA TAGTGTGTGGAAATTTTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGG CAAATCATTTGGTCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCC CCGCCCCGGACGAACTAAACCTGACTACGACATCTCTGCCCTTCTTCGCG GGGCAGTGCATGTAATCCCTTCAGTTGGTGGTACAACCTGCCAACTGGGC CCTGTTCCACATGTGACACGGGGGGGACCAACACAAAGGGGTTCTTGA CTGTAGTTGACATCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTG GCTTTTCACTGGAGCAGACTTTCAGTCTGTGGACTGCAACACAACATTTG CCTTTATGTGTAACCTTGGCTGAAGCTCTTACACCAATGCTGGGGGACAT GTACCTCCAGGGGCCAGGAAGACTACGGGAGGCTACACCAACATCAATC AGAGGGGCTGTCTAGCTACCCATAAGCCGACCCCTCAAGAGGGCATTAGCA ATAGTGTTTATAAGCCCCCTTGTAAACCTAAACGGGTAGCATATGCTTC CCGGGTAGTAGTATATACTATCCAGACTAACCCCTAATCAATAGCATATGT TACCCAACGGGAAGCATATGCTATCGAATTAGGGTTAGTAAAAGGCTCTA AGGAACAGCGATATCTCCACCCCATGAGCTGTCACGGTTTTTATTTACATG GGGTCAGGATTCACGAGGGTAGTGAACCATTTTAGTCAAAAGGGCAGTGG CTGAAGATCAAGGAGCGGGCAGTGAACCTCTCTGAATCTTCGCCTGCTTCT TCATTCCTTTCGTTAGCTAATAAGAATAACTGCTGAGTTGTGAACAGTAA GGTGTATGTGAGGTGCTCGAAAACAAGGTTTCAGGTGACGCCCCAGAATA AAATTTGGACGGGGGGTTCAGTGGTGGCATGTGCTATGACACCAATATAA CCCTCACAACCCCTTGGGCAATAAATACTAGTGTAGGAATGAAACATTTCT GAATATCTTAAACAATAGAAATCCATGGGGTGGGACAAAGCCGTAAGACT GGATGTCCATCTCACACGAATTTATGGCTATGGGCAACACATAATCCTAGT GCAATATGATACTGGGTTATTAAGATGTGTCAGGCAAGGACCAAGACA GGTGAACCATGTTGTACACTTATTTGTAACAAGGGGAAAGAGAGTGGAC GCCGACAGCAGCGACTCCACTGGTTGTCTTAACACCCCGAAAATTTAA CGGGGCTCCACGCAATGGGGCCATAAACAAGACAAAGTGGCCACTCTTT TTTTGAATTTGGAGTGGGGGCACGCGTCAAGCCCCACAGCCGCCCTG CGTTTTGGACTGTAAAATAAGGGTGTAA'AACT'GGCTGATTGTAACCCC GCTAACCACTGGGTCAAACCACTTGGCCAAAACCACTAATGGCACCCC GGGGAATACCTGCATAAGTAGTGGGGCGGCAAGATAGGCGGATGCTG TGCGATCTGGAGGACAAATTACACACACTTGGCCTGAGCGCCAAGCACAG GGTGTGTGGTCCCTCATATTCAGAGGTGCGTGAAGAGCAGGTGGGCTAATG TTGCCATGGGTAGCATATACTACCAATAATCTGGATAGCATATGCTATCC TAATCTAATCTGGGTAGCATAGGCTATCCTAATCTAATCTATCTGGGTAGC ATGCTATCCTAATCTAATCTGGGTAGTATATGCTATCCTAATTTATATCT CGGTAGCATAGGCTATCCTAATCTAATCTGGGTAGCATATGCTATCCTAA

SEQ ID NO	Vector name	Nucleotide sequences 123456789012345678901234567890123456789012345678901
		TCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATG CTATCCTAATAGAGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGG TAGCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATAT CTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCT AATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATA TGCTATCCTAATTTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTG GGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAAT CTGTATCCGGGTAGCATATGCTATCCTCATATAGCTGCAAACATGACA ATTTTCTTGAAGACGAAAGGGCCCTCGTGATACGCCTATTTTATAGGTTAA TGTCATGATAATAATGGTTTCTTAGACGTCAGGTGGCCTTTTCGGGGAAA TGTGCGCGGAACCCCTATTTGTTATTTTTCTAAATACATTCAAATATGTA TCCGCTCATGAGACAATAACCCGTGATAAATGCTTCAATAATATTGAAAAG GAAGAGTATGAGTATCAACATTTCCGTGTCGCCCTTATCCCTTTTTTGC GGCATTTTGCCTTCTGTTTTTGTCTCACCCAGAAACGCTGGTGAAGTAAA AGATGCTGAAGATCAGTTGGGTGCACGAGTGGTTACATCGAATGGATCT CAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAGAACGTTTTTCCAAT GATGAGCACTTTTTAAAGTTCTGCTATGTGGCGCGGTATTTATCCCGTGTGA CGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCAGAATGACTT GGTGAGTACTCACCACTCACAGAAAAGCATCTTACGGTATGGCATGACAGT AAGAGAATTATGCAGTGTGCCATAACCATGAGTGATAACACTGCGGCCAA CTTACTTCTGACACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCA CAACATGGGGGATCATGTAACCTGCCTTGATCGTTGGGAACCGGAGCTGAA TGAAGCCATACCAACGACGAGCGTGACACCAGATGCTGTCAGCAATGGC AACACGTTGCGCAAACTATTAACCTGGCGAACTACTTACTCTAGCTTCCCG GCAACAATTAATAGACTGGATGGAGGCGGATAAAGTTGACAGGACCCTTCT GCGCTCGGCCCTTCCGGCTGGCTGGTTTATGCTGATAAATCTGGAGCCGG TGAGCGTGGGTCTCGCGGTATCATGACGACTGGGGCAGATGCTGACAAAGC CTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGA ACGAAATAGACAGATCGCTGAGATAGGTGCCCTCACTGATTAAGCATTGGTA ACTGTGACAGCAAGTTTACTCATATATACTTTAGATTGATTTAAAACCTTCA TTTTAAATTTAAAAGGATCTAGGTGAAGATCCTTTTTTGATCAATCTATGAC CAAAATCCCTTAACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCTAGA AAAGATCAAAGGATCTTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTG CTTGCAAAACAAAAAACACCCGTAACAGCCGTGTTTTGTTGCGGGATCA AGAGCTACCAACTCTTTTTCCGAAGGTAACCTGGCTTCAGCAGACCGCAGAT ACCAAATACTGTTCTTCTAGTGTAGCCGTAGTTAGGCCACTTCAAGAA CTCTGTAGCACCGCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGC TGCTGCCAGTGGCGATAAGTCTGTCTTACCGGGTTGGACTCAAGACGATA GTTACCAGGATAAGGCGCAGCGGTCGGGCTGAACGGGGGGTTCTGTCACACA GCCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACTTACCGCTGA GCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGCGGACAGGTATCC GGTAAAGCGCAGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGG AAACGCTGATCTTTATAGTCTGTGCGGTTTTCGCCACTCTGACTTGA GCGTCGATTTTTTGATGCTCGTCAGGGGGCGGAGCCTATGGAATAACGC CAGCAACGCGGCCCTTTTTACGGTTCTGGCCTTTTGTGGCCTTTTGTCTCA CATGTTCTTTCCCTGCGTTATCCCTGATTCTGTGGATAACCGTATTACCGC CTTTGTAGTGAGCTGATACCGCTCGCCGACCGCAACCGAGCGCAGCGA GTCAGTGAGCGAGGAAGCGGAAGAGCGCCCAATACGCAACCCCTTCCC CGCGCGTTGGCCGATTCAATTAATGCAGCTGGCAGACAGGTTTCCCGACTG GAAAGCGGGCAGTGAAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTA GGCACCCAGGCTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAAT TGTGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATTACGC CAAGCTCTAGCTAGAGGTCGAGTCCCTCCCCAGCAGGCAGAAGTATGCAAAA GCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCCA TCCCGCCCTAACTCCGCCAGTTCCGCCCATTTCTCCGCCCTATGGCTGAC TAATTTTTTTTATTTATGTCAGAGCCGAGCCGCTCGGCTCTGAGCTAT TCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCTAGGCTTTTGCAAAAAGC TTTGCAAGATGGATAAAGTTTTAAACAGAGAGGAATCTTTGCAGCTAATG GACCTTCTAGGCTTGAAGGAGTGGGAATTTGGCTCCGGTGCCTGAGT GGCAGAGCGCACATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGGTCGG CAATTGAACCGGTCCTAGAGAGGTTGGCGGGGGTAACTGGGAAAGTGA TGCTGTACTGGCTCCGCTTTTTTCCCGAGGTTGGGGGAGAACCCTATAT AAGTGCAGTAGTCGCCGTGAACGTTCTTTTTCCGAACGGGTTTCCGCCAG

SEQ ID NO	Vector name	Nucleotide sequences
		<p>123456789012345678901234567890123456789012345678901</p> <p>AACACAGGTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGG TTATGGCCCTTGCGTGCCTTGAATTACTTCCACCTGGCTGCAGTACGTGAT TCTTGATCCCGAGCTTCGGGTGGAAGTGGGTGGGAGAGTTCAGGCCCTTG CGCTTAAGGAGCCCTTCGCCTCGTGTGTTGAGTTGAGGCCCTGGCCTGGGCG CTGGGGCCGCGCGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGTGC TTTCGATAAGTCTCTAGCCATTTAAAATTTTGTATGACCTGCTGCGACGCT TTTTTTCTGGCAAGATAGTCTTTGTAATGCGGGCAAGATCTCACACTGG TATTTCCGTTTTTTGGGGCCGGGCGGGGACGGGGCCGTGCGTCCCAGCG CACATGTTTCGGCGAGGCGGGGCTGCCAGCGCGGCCACCGAGAATCGGACG GGGTAGTCTCAAGCTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCG GTGTATCGCCCGCCCTGGCGGCAAGGCTGGCCCGGCTGGCCCGTTCG GTGAGCGAAAGATGGCCGCTTCGCGCCCTGCTGCAGGGAGCTCAAAATG GAGGACCGGGCGCTCGGGAGAGCGGGCGGGTGTGAGTACCACACAAAGGAA AAGGCCCTTCCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCG GCGCCCTCCAGGACCTCGATTAGTTCTCGAGCTTTTGGAGTACCGTCTG TTTAGGTTGGGGGAGGGTTTTATGCGATGGAGTTTCCCACACTGAGTG GGTGGAGACTGAAGTAGCCAGCTTGGCACTTGATGTAATTCTCCTTGGGA ATTTGCCCTTTTGTAGTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGT GGTTCAAAGTTTTTTCTCCATTTCCAGGTGTGAGGAAATCTCTAGAG ATCCCTCGACCTCGAGATCCATTGTGCCCGGGCGCCACCATGGAGTTTGGG CTGAGCTGGCTTTTTCTTGTGCGGATTTTAAAGGTGTCCAGTGC</p>
305	V2	<p>ACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTG AAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGA GAGGCCAAAAGTACAGTGGAAAGGTGGATAACCCCTCCAAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGC AGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCC TCGAAGTCAACCAACAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAAC AAGGGAGAGTGTGAGCGGCCGCTCGAGGCCGGCAAGGCCGATCCCCCGA CCTCGACCTCTGGCTAATAAAGGAAATTTATTTTCATTGCAATAGTGTGT GGAATTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAAAATCATT TGGTCCGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCCCGCCCGG ACCAACTAAACCTGACTACGACATCTCTGCCCTTCTTCGCGGGCAGTGC ATGTAATCCCTTCACTTGGTGGTACAACCTGCCAATGGGCCCTGTTCCA CATGTGACACGGGGGGGACCAACACAAAGGGGTTCTCTGACTGTAGTTG ACATCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTGGCTTTTCATC CTGGAGCAGACTTTCAGTCTGTGGACTGCAACACAACATTTGCCTTTATGT GTAACCTTGGCTGAAGCTCTTACACCAATGCTGGGGGACATGTACCTCCC AGGGGCCAGGAAGACTACGGGAGGCTACACCAACGTCATCAGAGGGGCC TGTGTAGCTACCGATAAGCGGACCCTCAAGAGGGCATTAGCAATAGTGT ATAAGGCCCTTTGTTAACCTTAAACGGGTAGCATATGCTTCCCGGTAGT AGTATAFACTATCCAGACTAACCTAATTCAATAGCATATGTTACCCAACG GGAAGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCTAAGGAACAGC GATATCTCCACCCATGAGCTGTCACGGTTTTATTTACATGGGGTACAGGA TTCCACGAGGGTAGTGAACCATTTTAGTACAAGGGCAGTGGCTGAAGATC AAGGAGCGGGCAGTGAACCTCTCTGAATCTTCGCCTGCTTCTTCAATCTCC TTCGTTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAAGGTGTATGT GAGGTGCTCGAAAACAAGGTTTCAGGTGACGCCCCAGAAATAAAATTTGGA CGGGGGTTCAGTGGTGGCATTGTGCTATGACACCAATATAACCCTCACAA ACCCTTGGGGCAATAAATACTAGTGTAGGAATGAAACATTTCTGAATATCTT TAACAAATAGAAATCCATGGGGTGGGGACAAGCCGTAAGACTGGATGTCCA TCTCACAGAAATTTATGGCTATGGGCAACACATAATCCTAGTGCATATGA TACTGGGGTTATTAAGATGTGTCCAGGCAGGACCAAGACAGGTGAACCA TGTGTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGACCGGACAGC AGCGGACTCCACTGGTTGTCTTAACACCCCGAAAATTAACGGGGCTCC ACGCCAATGGGGCCATAAACAAGACAAGTGGCCACTCTTTTTTTTGGAA TTGTGGAGTGGGGGCACGCGTCAGCCCCACAGCCCGCCTGCGGTTTTTGG ACTGTAATAAAGGTTGTAATAACTTGGCTGATTGTAACCCCGGTAACAC TGCGGTCAAACCACTTGGCCACAAAACCACTAATGGCACCCCGGGGAATAC CTGCATAAGTAGGTGGCGGGCCAAGATAGGGCCGCGATTGCTGCGATCTG GAGGACAAATACACACACTTGGCCTGAGCGCAAGCACAGGGTTGTGG TCCTCATATTCAGGTCGCTGAGAGCACGGTGGGCTAATGTGCCATGG GTACCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATA TCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCC</p>

SEQ ID NO	Vector name	Nucleotide sequences
		1234567890123456789012345678901234567890123456789012345678901 TAATCTAATCTGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCAT AGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCT GGGTAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATGCTATCCTAA TAGAGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATATA CTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCTGGGTAGC ATATGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATAT CTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCT AATTTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATA TGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCG GGTAGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGAAATTTTCCTTG AAGACGAAAGGGCCTCGTGATACGCCCTATTTTATAGGTGAATGACATGAT AATAATGGTTTCTTAGACGTCAGGTGGCACTTTTTCGGGGAAATGTGCGCGG AACCCCTATTTGTTATTTTTCTAAATACATTCAAATATGTATCCGCTCAT GAGACAATAACCCGTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTAT GAGTATTCACACATTTCCGTGTCCGCCCTTATTCCTTTTTTTCGGCATTTTG CCTTCCGTTTTTGCTCACCCAGAAACCGCTCGTGAAAGTAAAAGATGCTGA AGATCAGTTGGGTGCACGAGTGGGTACATCGAAGTGGATCTCAACAGCGG TAAGATCCTTGAGAGTTTTCGCCCCGAAGAAGCTTTTCCAATGATGAGCAC TTTTAAAGTTCCTGCTATGTGGGCGGTATTTACCGGTGTAAGCAGCGGCA AGAGCAACTCGGTCCGCGCATACACTATTTTCAGAATGACTTGGTTGAGTA CTCACCAGTACAGAAAAGCATTTACGGATGGCATGACAGTAAGAGAATT ATGACAGTGTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCT GACAACGATCGGAGACCSAAGGAGCTAACCGTTTTTGCACAACATGGG GGATCATGTAACCTCGCCTTGATCGTTGGGAACCGGAGCTGAAGCCAT ACCAAACGACGAGCGTGACACCAGGATGCCGTCAGCAATGGCAACAACGTT GCGCAAACTATTAAGTGGCGAAGTACTTACTCTAGCTTCCCGCAACAAAT AATAGACTGGATGGAGCGGATAAAGTTGCAGGACCCTTCTGCGCTCGGC CTTTCCGGCTGGCTGGTTTTATTGCTGATAAATCTEGAGCCGGTGAAGCTGG GTCTCGCGGTATCATTTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTAT CGTAGTTATCTACAGACGGGGAGTCAGGCAACTATGGATGAACGAAATAG ACAGATCGCTGAGATAGGTGCCCTCACTGATTAAGCATGGTAAGTGTGAGA CCAAGTTTACTATATATACTTTAGATTGATTTAAAACCTCAATTTTTAAT TAAAAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCC TTAACGTGAGTTTTCGTTCACCTGAGCGTCAAGACCCCGTAGAAAAGATCAA AGGATCTTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTGCTTGCAAAAC AAAAAAACCCGCTACACAGCGGTGGTTTTGTTGCCGGATCAAGGATACC AACTCTTTTTCCGAAGGTAAGTGGCTTCCAGCAGAGCGCAGATACCAAAATAC TGTTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAAGTCTGTAGC ACCGCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAG TGCCGATAAGTCCGTCTTACCAGGTTGGACTCAAGACGATACCGGGA TAAGCGCAGCGGTCCGGCTGAACGGGGGGTTGCTGCACACAGCCCAGCTT GGAGCGAACGACCTACACCGAAGTGAATACCTACAGCGTGAAGTATGAGA AAGCGCCACGCTTCCCGAAGGGAGAAAGGGCGGACAGGTATCCGGAAGCGG CAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGAAACCGCTG GTATCTTTATAGTCTGTGCGGTTTTCGCCACCTCTGACTTGAGCGTCCGATT TTTGTGATGCTCGTCAGGGGGCGGAGCCTATGGAAAACGCCAGCAACGC GGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCTTTTTGCTCACATGTTCTT TCCGTGCTTATCCCTGATCTGTGGATAACCGTATTACCGCCTTTGAGTG AGCTGATACCGCTCGCCGACGCGAAGCAGCCGAGCGCAGGTCAGTGAG CGAGGAAGCGGAAGAGCGCCCAATACGCAAAACCGCTTCCCCGCGGTTG GCCGATTCATTAATGCAGCTGGCAGCAGAGTTTCCCGACTGAAAGCGGG CAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCCA GGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGGAAATGTGAGCGG ATAACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTCTA GCTAGAGGTGAGTCCCTCCCCAGCAGGCAGAAGTATGCAAGCATGCATC TCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCCATCCCCGCC TAACCTCGCCCGTTCGGCCCATTTCTCCGCCCATGGCTACTAATTTTTTT TTATTTATGCAGAGGCGAGGCGCCTCGGCTCTGAGCTATTCCAGAAGT AGTGAGGAGGCTTTTTGGAGGCTAGGCTTTTGCAAAAAGCTTTGCAAAAG ATGGATAAAGTTTTAAACAGAGAGGAATCTTTGCAGCTAATGGACCTTCTA GGCTTTGAAAGGAGTGGGAATTGGCTCCCGTGCCCGTCACTGGCAGCGG CACATCCCCACAGTCCCGGAGAAGTTGGGGGGAGGGTCCGGCAATTGAAC CGGTGCTAGAGAAGGTGGCGGGGGTAACTGGGAAAGTATGCTGCTGTA

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901 CTGGCTCCGCCTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGT AGTCGCGTGAACGTTCCTTTTTTCGCAACGGTTCGCCGCCAGAACACAGGT AAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGGTATGGCCC TTGCGTGCCCTGAATTACTTCCACCTGGCTGCAGTACGTGATTCCTGATCC CGAGCTTCGGGTGGAAGTGGTGGGAGAGTTCGAGGCTTGCCTTAAGG AGCCCTTCGCTCGTGTGAGTTGAGGCTGGCCTGGGCGCTGGGGCCG CCGCTGCGAATCTGGTGGCACCTTCGCGCTGTCTCGCTCTTCGATAA GTCTCTAGCCATTTAAAATTTTGATGACCTGCTGCGACGCTTTTTTCTG GCAAGATAGTCTTGTAAATGCGGGCCAAGATCTGCACACTGGTATTCGGT TTTTGGGGCCGCGGGCGGCGACGGGGCCGTCGCTCCAGCGCACATGTTTC GCGGAGCGGGGCTGCGAGCGCGGCCACCAGAGAATCGACGAAAGGGCTC TCAAGCTGGCCGCTGCTCTGGTGCCTGGCCTCGCGCCGCGTGTATCGC CCCGCCCTGGGCGCAAGGCTGGCCCGGTGCGCACAGTTGCGTGAGCGGA AAGATGGCCGCTTCCCGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCG GCGCTCGGGAGAGCGGGCGGGTGAATCACCCACACAAAGCAAGGGCCTT TCCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCGGGCGCCGTC CAGGCACCTCGATTAGTTCGAGCTTTTGGAGTACGTGCTTTTAGGTTG GGGGAGGGGTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGAC TGAAGTTAGGCAGCTTGGCACTTGATGTAATTCCTTGAATTTGCCCT TTTTGTGTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGTGGTCAAAG TTTTTTCTTCCATTTGAGGTGTCGTGAGGAATTCCTAGAGATCCCTCGA CCTCGAGATCCATTGTGCCGGGCGCACCATGGACATGCGCGTCCCGCCC AGCTGCTGGCCCTGCTGCTGCTGTTTCCCGGCTCGCGATG
306	V3	CAACCAAGGCTGCCCTCGGTCACTCTGTTCCCGCCCTCCTCTGAGGAG CTTCAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGAATTCACCCG GGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCGTCAAGGCGGGA GTGGAGACCACCACCCTCCAAACAAGCAACAAGTACGCGGCCAGC AGCTACTGAGCTGACGCCCTGAGCAGTGAAGTCCCACAGAAGTACAGC TGCCAGGTACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCTACA GAATGTTTCATGAGCGGCCGCTCGAGGCCGCAAGGCCGATCCCCGACCT CGACCTCTGGCTAATAAAGGAAATTTATTTTCATTGCAATAGTGTGTTGGA ATTTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGCAATCATTTGG TCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCCCCGCCCGGACG AACTAAACCTGACTACGACATCTCTGCCCTTCTTCGCGGGCAGTGCATG TAATCCCTTCAGTTGGTGGTACAACCTGCCAAGTGGGCCCTGTTCCACAT CTGACACGGGGGGGACCAAAACAAGGGTCTCTGACTGAGTTGACA TCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTGGCTTTTATCCTG GAGCAGACTTTGCACTGTGTGACTGCAACACAACATTCCTTTATGTGTA ACTCTTGGCTGAAGCTCTTACACCAATGCTGGGGGACATGTACCTCCAGG GGCCAGGAAGACTACGGGAGGCTACACCACGTCATTAAGGACCGCTGT GTAGCTACCGATAAGCGGACCCTCAAGAGGGCATAGCAATAGTGTTTATA AGGCCCTTGTAAACCTAAACGGGTAGCATATGCTTCCCGGGTAGTAGT ATATACTATCCAGACTAACCCTAATTCATAGCATATGTTACCCAACGGGA AGCATATGCTATCGAATTAGGTTAGTAAAAGGGTCCTAAGGAACAGCGAT ATCTCCACCCCATGAGCTGTACGGTTTTATTTACATGGGGTCAGGATTC CACGAGGCTAGTGAACATTTTAGTCACAAGGGCAGTGGCTGAAGATCAAG GAGCGGGCAGTGAACCTCTCTGAATCTTCGCTGCTTCTTCACTCTCTTC GTTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAAGGTGTATGTGAG GTGCTCGAAAACAAGGTTTCAGGTGACGCCCCCAAGATAAAATTTGGACGG GGGGTTCACTGGTGGCATTGTGCTATGACACCAATATAACCTCACAAACC CCTTGGGCAATAAATACTAGTGTAGGAATGAAACATTTGAATATCTTTAA CAATAGAAATCCAATGGGGTGGGGACAAGCCGTAAGACTGGATGTCATCT CACACGAATTTATGCTATGGGCAACACATAATCCTAGTGCAATATGATAC TGGGGTTATTAAGATGTGTCCAGGCAGGGACCAAGACAGGTGAACCATGT TGTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGACGCCGACAGCAGC GGACTCCACTGGTGTCTCTAACACCCCGAAAATTAACCGGGGCTCCACG CCAATGGGGCCCAAAACAAGACAAGTGGCCACTCTTTTTTGAATTTG TGGAGTGGGGGCACGCGTCAAGCCCCACACGCCGCTTGGGTTTTGGACT GTAAAATAAGGGTGTAAATACTTGGCTGATGTAAACCCGCTAACCACTGC GGTCAAACCACTTGCCCAAAAACCACTAATGGCACCCCGGGGAATACCTG CATAAGTAGTGGCGGGCCAAGATAGGGGGCGGATTCGCTGCGTAGGAG GACAAATTACACACACTTGGCCTGAGCGCCAAGCACAGGGTGTGGTCC TCATATTCACGAGTGCCTGAGAGCACGGTGGGCTAATGTTGCCATGGGTA

SEQ ID NO	Vector name	Nucleotide sequences 123456789012345678901234567890123456789012345678901
		GCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCT GGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAA TCTATATCTGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATAGG CTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCTGGG TAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATGCTATCCTAATAG AGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATATACTA CCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCTGGGTAGCATA TGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTG GGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAAT TTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGC TATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCGGGT AGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGAATTTTCTTGAAG ACGAAAGGGCCTCGTGATACGCCATTTTTTATAGGTTAATGTCATGATAAT AATGGTTTCTTAGACGTCAGGTGGCACTTTCCGGGAAATGTCCGGGAAC CCTATTTGTTTTATTTTTCTAAATACATTCAAATATGATCCCGCTCATGAG ACAATAACCCTGATAAATGCTTCAATAATATTGAAAAGGAAGAGTATGAG TATTCACATTTCCGTGTCCGCCCTTATCCCTTTTTTCCGGCATTTTCCT TCCTGTTTTGCTCACCAGAAACGCTGGTGAAGTAAAGATGCTGAAGA TCAGTTGGGTGCACGACTGGGTACATCCGACTGCATCTCAACAGCGGTAA GATCCTTGAGAGTTTTCCGCCGGAAGAACGTTTTTCCAATGATGAGCACTTT TAAAGTTCTGCTATGTGGCGCGTATTATCCCGTGTGACGCCGGGCAAGA GCAACTCGGTCCGCCATACACTATTCTCAGAATGACTTGGTTGAGTACTC ACCAGTACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTAAG CAGTGTCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGAC AACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGA TCATGTAACCTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACC AAACGACGAGCGTACACCACGATGCCTGCAGCAATGGCAAGAGATTTGGG CAAACTATTAACGGGCAACTACTTACTCTAGCTTCCCGGCAACAATTAAT AGACTGGATGGAGCGGATAAAGTTGCAGGACCCTTCTGCGCTCGGCCCT TCCGGCTGGCTGGTTTTATTGCTGATAAATCTGGAGCCCGTACCCGTGGGTC TCCCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCTCCCGTATCGT AGTTATCIACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACA GATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAGACCA AGTTTACTCATATATACTTTAGATTGATTTAAAACCTTCATTTTTAATTTAA AAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTA ACGTGAGTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGG ATCTTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTGCTTGCAAAACAAA AAAACCACCGCTACCAGCGGTGGTTTTGTTTTGCCGGATCAAGAGCTACCAAC TCTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGATACCAAACTACTGT TCTTCTAGTTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCC GCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGG CGATAAGTCTGTCTTACCAGGTTGGACTCAAGACGATAGTTACCAGGATAA GGCGCAGCGGTCCGGCTGAACGGGGGGTTCGTGCACACAGCCAGCTTGGGA GCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAAGTATGATAAAG CGCCACGCTTCCGAAGGGGAGAAAGGGGACAGGTATCCGGTAAGCGGCAG GGTCCGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGAAACGCCITGGTA TCTTTATAGTCTGTGGGTTTTCCGCCACCTCTGACTTGAAGCGTCAATTTTT GTGATGCTCGTCAGGGGGGGGAGCCTATGGAAAACGCCAGCAACCGCGGC CTTTTTACGGTTCTCGCCCTTTTCTGCGCTTTTCTGCTCACATGTTCTTTCC TGCGTTATCCCCTGATCTGTGGATAACCGTATTACCAGCTTTGAGTGAGC TGATACCGCTCGCCGACCGCAACGACCGAGCGCAGCGAGTCAAGTGAAGCA GGAAGCGGAAGAGCGCCCAATACGCAACCGCCTCTCCCGCGGTTGGCC GATTCATTAATGCAGCTGGCAGCACAGGTTTTCCCGACTGGAAAAGCGGGCAG TGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCAGGC TTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTTGTGAGCGGATA ACAATTTACACAGGAACAGCTATGACCATGATTACGCCAAGCTCTAGCT AGAGGTGAGTCCCTCCCCAGCAGGCAGAGTATGCAAGCATGCAATCTCA ATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCATCCCGCCCTAA CTCGCCCAAGTTCCGCCATTTCTCCGCCCATGGCTGACTAATTTTTTTTA ITTATGACAGGGCCGAGGCCCTCGGCCCTCTGAGCTATTCCAGAAGTAGT CAGGAGGCTTTTTTGGAGGCTTAGGCTTTTGCAAAAAGCTTTGCAAAAGATG GATAAAGTTTTAAACAGAGAGGAATCTTTGACGCTAATGGACCTTCTAGGT CTTGAAAAGGAGTGGGAATGGCTCCGGTGCCTGCTCAGTGGCAGAGCCGAC

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901 ATCGCCACAGTCCCCGAGAAGTTGGGGGAGGGGTCCGCAATTGAACCGG TGCCTAGAGAAGTGGCGGGGTAAACTGGGAAAGTATGTCGTGFACTG GCTCCGCCTTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGT CGCGTGAACGTTCTTTTTCGCAACGGGTTTGCCGCCAGAACACAGGTAAG TGCCGTGTGTGGTTCGCCGGGCTGGCCTCTTTACGGGTATGGCCCTTG CGTGCCTTGAATTACTCCACCTGGCTGCAGTACGTGATCTTGATCCCGA GCTTCGGGTGGAACTGGGTGGGACAGTTCGAGGCCCTTGGCTTAAGGAGC CCCTTCGCCTCGTGTGAGTTGAGGCTGGCCTGGGCGCTGGGGCCGCCG CGTGGGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGCTTCGATAAGTC TCTAGCCATTTAAAATTTTGTATGACCTGCTGCGACGCTTTTTTTTCGGCA AGAAGTCTTTGTAATGCGGGCCAAGATCTGCACACTGGTATTTCCGTTTTT TGGGGCCGGGGCCGACGGGGCCCGTGGCTCCAGCGCACATGTTCCGGC GAGGCGGGGCTCGAGCGCGCCACCGAGAATCGGAGGGGGTAGTCTCA AGCTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCCGGTGTATCGCCCC GCCCTGGGCGGCAAGGCTGGCCCGGTGGCACCAAGTTGGCTGAGCGGAAAG ATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGAGGACGGGGC CTCGGGAGAGCGGGCGGGTCACTCACCCACACAAAGGAAAAGGGCCTTCC GTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCGGGCCCGCTCCAG GCACCTCGAATAGTCTCGAGCTTTTGGAGTACGTGCTTTAGTTGGGG GGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACTGA AGTTAGGCCAGCTTGGCAGTTGATGTAATTCCTTGAATTTGCCCTTTT TGAGTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGTGGTTCAAAGTTT TTTTCTCCATTCAGGTGTCTGAGGAATTCCTTAGAGATCCCTCGACCT CGAGATCCATTGTGCCGGGGCCACCATGACTTGGACCCCACTCCTCTTC CTCACCTCCTCTCCACTGCACAGGAAGCTTATCG
307	V4	ACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTG AAATCTGSAACTGCCTCTGTGTGTGCCTGCTGAATAACTTCTATCCAGA GAGGCCAAAGTACAGTGGAAAGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGC AGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCC TGCGAAGTCACCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAAC AGGGGAGAGTGTGAGCGGGCGCTCGAGGCGGCAAGGCCGATCCCGGA CCTCGACCTCTGGCTAATAAAGGAAATTTATTTTCATTGCAATAGTGTGTT GGAATTTTTTGTCTCTCTCACFCGGAAGGACATATGGGAGGGCAAATCATT TGGTCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCCCCGCCCGG ACCAACTAAACCTGACTACGACATCTCTGCCCTTCTTCGCGGGGAGTGC ATGTAATCCCTCAGTTGGTGGTACAACCTTGCCAACCTGGGCCCTGTTCCA CATGTGACACGGGGGGGACCAAACACAAGGGGTTCTCTGACTGTAGTTG ACATCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTGGCTTTCATC CTGGAGCAGACTTTCAGTCTGTGGACTGCAACACAACACTGCTTATGT GTAACCTTTGGCTGAAGCTCTTACACCAATGCTGGGGGACATGTACCTCCC AGGGGCCAGGAAGACTACGGGAGGCTACACCAACGTCATCAGAGGGGCC TGTGTAGCTACCSATAAGCGGACCCTCAAGAGGGCATFAGCAATAGTGT ATAAGGCCCCCTTGTAAACCTAAACGGGTAGCATATGCTTCCCGGGTAGT AGTATATACTATCCAGACTAACCCCTAATCAATAGCATATGTTACCCAACG GGAAGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCCCTAAGGAACAGC GATATCTCCACCCCATGAGCTGTACGGTTTTATTTACATGGGGTCAGGA TTCCAGAGGGTAGTGAACCATTTTAGTCAAGGGCAGTGGCTGAAGATC AAGGAGCGGGCAGTGAACCTCCTGAATCTTCGCCCTGCTTCTTCAATCTCC TTCGTTAGCTAAATAGAATAACTGCTGAGTGTGTAACAGTAACCTTATGT GAGGTGCTCGAAACAAGGTTTCAGGTGACGCCCCCAAGATAAAATTTGGA CGGGGGTTTCAGTGGTGGCATGTGCTATGACACCAATATAACCCCTCAAA ACCCCTTGGCAATAAATACTAGTGTAGGAATGAAACATCTGAAATATCTT TAACAATAGAAATCCATGGGGTGGGGACAAGCCGTAAAGACTGGATGTCCA TCTCACACGAATTTATGGCTATGGGCAACACATAATCCTAGTGCATATGA TACTGGGGTTATTAAGATGTGTCCCAGGCAGGGACCAAGACGGTGAACCA TGTGTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGACCGCCGACAGC AGCGGACTCCACTGGTTGTCTTAACACCCCGAAAAATTAACGGGGCTCC ACCCAATGGGGCCATAAACAAGACAAGTGGCCACTCTTTTTTTTGGAAA TTGTGGAGTGGGGGCAGCGCTCAGCCCCACAGCCGCCCTGCGGTTTTGG ACTGTAAAATAAGGGTGAATAAAGTGGCTGATTTGTAACCCCGTAACTAC TGCGCTCAAACCACTTGGCCACAAAACCACTAATGGCACCCCGGGGAATAC CTGCATAAGTAGTGGGGGGCAAGATAGGGGCGGATGCTGCGATCTG

SEQ ID NO	Vector name	Nucleotide sequences
		<p>123456789012345678901234567890123456789012345678901</p> <p>GAGGACAAATTACACACACTTGGCCTGAGCGCCAAGCACAGGGTTGTTGG TCCTCATATTCACGAGGTGCGCTGAGAGCACGGTGGGCTAATGTTGCCATGG GTAGCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATA TCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCC TAATCTATATCTGGGTAGTATATGCTATCCIAATTTATATCTGGGTAGCAT AGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCT GGGTAGTATATGCTATCCTAATCTGATCCGGGTAGCATATGCTATCCTAA TAGAGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATATA CTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCTGGGTAGC ATATGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATAT CTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCT AATTTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATA TGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATCTGATCCG GGTAGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGAATTTTCTTG AAGACGAAAGGGCTCGTGATACGCTATTTTTATAGGTTAATGTCATGAT AATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGG AACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGATCCGCTCAT GAGACAATAACCCGTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTAT GAGTATTCACCATTTCCGTGTCGCCCTTATTCCTTTTTGCGCGCTTTG CCTTCTGTTTTTGGCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGA AGATCAGTTGGGTGCACGAGTGGGTACATCGAAGTGGATCTCAACAGCGG TAAGATCCTTGAGAGTTTTCGCCCCGAAGAAGCTTTTCCAATGATGAGCAC TTTTAAAGTTCTGCTATGTTGGCGCGGTATTTATCCCGTATGACGCCGGCA AGAGCAACTCGGTGCGCGCATACACTATTCTCAGAATGACTTGGTTGAGTA CTCACAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATT ATGCAAGTGTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCT GACAACGATCGGAGGACCGAAGGAGCTAACCGTTTTTTGACACAACATGGG GGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCAT ACCAAACGACGAGCGTGACACCAGGATGCCGTCAGCAATGGCAACAAGTT GCGCAAACTATTAAGTGGCGAAGTACTTACTCTAGCTTCCCGGCAACAATT AATACACTCGATGAGGCGGATAAAGTTGCAAGGACCATTTCGCTGCGGC CCTTCCGGCTGGCTGGTTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGG GTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTAT CGTAGTTATCTACAGACGGGGAGTCAGGCAACTATGGATGAACGAAATAG ACAGATCGCTGAGATAGGTGCTCACTGATTAAGCATGGTAAGTCTCAGA CCAAGTTTACTCATATATACTTTAGATTGATTTAAAACCTCAATTTTTAATT TAAAAGGATCTAGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCC TTAACGTGAGTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAA AGGATCTTCTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCTTGCAAAC AAAAAAACCACCGCTACCAGCGGTGTTTTGTTTGGCGGATCAAGACTACC AACTCTTTTTCCGAAGGTAACGGCTTCCAGCAGAGCGCAGATACCAAAATAC TGTTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGC ACCGCTACATACTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAG TGGCGATAAGTCTGTCTTACCAGGTTGGACTCAAGACGATAGTTACCAGGA TAAGGCGCAGCGGTGCGGCTGAACGGGGGGTTGTTGCACACAGCCAGCTT GGAGCGAACGACCTACACCGAAGTACGATACCTACAGCGTGAGCTATGAGA AAGCGCCACGCTTCCCAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGG CAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGAAAACGCTG GTATCTTTATAGTCTGTGCGGTTTTCGCCACCTCTGACTTGAGCGTGCATT TTTGTGATGCTCGTCAGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGC GGCCTTTTTACGGTTCCCTGGCCTTTTTGCTGGCCTTTTTGCTCACATGTTCTT TCCTGCGTTATCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTG AGCTGATACCGCTCGCCGAGCCGAACGACCGAGCGCAGCGAGTCACTGAG CGAGGAAAGCGAAGAGCGCCCAATACGCAACCGCCTTCCCGCGCGTTG GCCGATTCATTAATGCAGCTGGCAGCAGAGTTTCCCGACTGGAAAGCGGG CAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCCA GGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGGAATTTGTGAGCGG ATAACAATTTACACAGGAAACAGCTATCACCATGATACGCCAAGCTCTA GCTAGAGGTGAGTCCCTCCCGAGCAGGCGAAGTATGCAAAGCATGCATC TCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCCATCCCGCCCT TAACCTCGCCAGTTCCGCCCCATTCTCCGCCCATGGCTGACTAATTTTTT TTATTTATGCAGAGGCGGAGGCGCCTCGGCTCTGAGCTATTCAGAAGT AGTGAGGAGGCTTTTTTGGAGCCTAGGCTTTTGAAAAAGCTTTGCAAG</p>

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901 ATGGATAAAAGTTTTAAACAGAGAGGAATCTTTGCAGCTAATGGACCTTCTA GGTCTTGAAAGCAGTGGGAATTGGCTCCGGTGCCCGTCAGTGGGCAGAGCG CACATCGCCACAGTCCCGGAGAAGTTGGGGGAGGGGTCGGCAATGAAC CGGTGCTAGACAAGGTGGCGCGGGTAAACTGGGAAAGTGATGTCGTGTA CTGGCTCCGCCCTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGT AGTCGCGGTGAACGTTCTTTTTTCGCAACGGGTTTGGCCGACAGAACAGGT AAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGGTATGGCCC TTGGCTGCCTTGAATTACTTCCACTGCGTCCAGTACGTGATTCTTGATCC CGAGCTTCGGGTGGAAAGTGGGTGGGAGAGTTCGAGGCCCTTGCCTTAAGG AGCCCTTCGCTCGTGTGAGTTGAGGCTGGCCTGGGCGCTGGGGCCG CCGCGTGCGAATCTGGTGGCACCTTCGCGCTGTCTCGCTGCTTTTCGATAA GTCTC'AGCCATTTAAAATTTTTGATGACCTGCTGCCAGCCTTTTTTTCTG GCAAGATAGTCTTTGAAAAGCGGGCCAAGATCTGCACACTGGTATTTCCGT TTTTGGGGCCGGGGCGGACGGGGCCCGTGGTCCAGCGCACATGTTTC GCGGAGCGGGGCTTGGGAGCGCGGCCACCGAGAATCGGACGGGGGTAGTC TCAAGCTGGCCGGCCTGCTCTGGTGGCTGGCTTCGGCCCGCGTGTATCGC CCCGCCCTGGGCGGAAGGCTGGCCCGGTCCGCACCAAGTTCGCTGAGCGGA AAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGG GCGCTCCGGAGAGCGGGCGGGTGGTGCAGTCAACCACACAAAGGAAAGGGCTT TCCGTCCTCAGCCGTGCTTTCATGTGACTCCACGGAGTACCGGGCGCCGTC CAGGCACCTCGATTAGTTC'CGAGCTTTTGAGTACGTGCTTTTAGGTTG GGGGGAGGGTTTTATGCGATGGAGTTTCCCACACTGAGTGGGTGGAGAC TGAAGT'FAGGCCAGTTGGCAC'TGATGTAAT'CTCCT'CGAGAAAGGGCTT TTTTGGATTTGGATCTTGETTCA'TTCTCAAGCCTCAGACAGTGGTCAAAG TTTTTTCTTCCATTT'GAGGTGTCGTGAGGAAT'CTC'TAGAGATCCCTCGA CCTCGAGATCCATTTGTGCCGGGGCGCACCATGACTTGGACCCCACTCCTCT TCCTCACCTCCTCCTCCACTGCACAGGAAGCTTATCG
308	V5	CAACCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCCTCCTCTGAGGAG CTTCAAGCCAACAAGGCCACACTGGTGTGTC'ATAAGT'GACTTCTACCCG GGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCGTCAAGGCGGGA GTGGAGACCACCACCCTCCAAACAAGCAACAACAGTACCGGCCAGC AGCTACCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCCAGGAAAGTACAGC TGCCAGTCA'CGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCTACA GAATGTT'CATGAGCGCCGCTCGAGGCCGGCAAGGCCGATCCCCGACCT CGACCTCTGGCTAATAAAGGAAATTTATTTT'CAT'GCAATAGTGTGTTGGA ATTTTTGTGTC'CTCACTCGGAAGGACATATGGGAGGCAAAATCATTTGG TCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCCCCGCCCCGACG AACTAAACCTGACTACGACATCTCTGCCCTTCTTCGCGGGGACAGTGCATG TAATCCCT'CACTGGTGGTACAACTTGCCAACTGGGCCCTGT'TCCACAT GTGACACGGGGGGGACCAAACAAGGGGTCTCTGACTGTAGTGCAGACA TCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTGGCTTTCATCCTG GAGCAGACTTTGCAGTCTGTGGACTGCAACACAACATTCGCTTTATGTGTA ACTCT'FGGTGAAGCTTACACCAATGCTGGGGACATGTACCTCCAGG GGCCAGGAAGACTACGGGAGGCTACACCAACGTCAATCAGAGGGGCTGT GTAGCTACCGATAAGCGGACCCCAAGAGGGCATTAGCAATAGTGTTTATA AGGCCCCCTTGT'AAACCCTAAACGGGTAGCATATGCTTCCCGGGTAGTAGT ATATACTATCCAGACTAACCC'AAATCAATAGCATATGTTACCCAACGGGA AGCATATGCTATCGAATTAGGGT'AGTAAAAGGGTCC'AAAGGAACAGCGAT ATCTCCACCCCATGAGCTGT'CACGGTTTTATTTACATGGGGTCAGGATTC CACGAGGGTAGTGAAACATTTTAGT'CAACAAGGGCAGTGGCTGAAGATCAAG GAGCGGGCAGTGA'ACTCTCCTGAATCTTCGCTGCTTCT'CATTTCTCCTTC GTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAAGGTGTATGTGAG GTGCTCGAAAACAAGGTTTCAGGTGACGCCCCCAAGAAATAAATTTGGACGG GGGGTT'CACTGGTGGCATTGTGCTATGACACCAATATAACCCCTCACAAACC CCTTGGGCAATAAATACTAGTGTAGGAATGAAACATTT'GAATATCTTTAA CAATAGAAATCCATGGGGTGGGGACAAGCCGTAAGACTGGATGTCATCT CACACGAATTTATGGCTATGGGCAACACATAATCCTAGT'GCAATATGATAC TGGGGTTATTAAGATGTGTC'CCAGGCAGGGACCAAGACAGGTGAACCATGT TGTTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGACGCCGACAGCAGC GGACTCCACTGGTTGTCTTAACACCCCGAAAATTAACCGGGGCTCCACG CCAATGGGGCCCA'AAAACAAGACAAGTGGCCACTCTTTTTTTGAAATTTG TGGAGTGGGGCCACCGCTCAGCCCCACACGCCCGCTTGGGTTTTTCCACT GTAAAATAAGGGT'GTAATAACTTGGCTGATGTAAACCCGCTAACCACTGC

SEQ ID NO	Vector name	Nucleotide sequences
		<p>1234567890123456789012345678901234567890123456789012345678901</p> <p>GGTCAAACCACTTGCCACAAAACCACTAATGGCACCCCGGGGAATACCTG CATAAGTAGGTGGGGGGCCAAAGATAGGGGGCGGATTGCTGCGATCTGGAG GACAAATTACACACACTTGCGCCTGAGCGCCAAGCACAGGGTTGTTGGTCC TCATATTCACGAGGTGCGTGTAGAGCACGGTGGGCTAATGTTGCCATGGGTA GCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCT GGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAA TCTATATCTGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATAGG CTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCTGGG TAGTATATGCTATCCTAATCTGATCCGGGTAGCATATGCTATCCTAATAG AGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATATACTA CCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCTGGGTAGCATA TGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTG GGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAAT TTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCT TATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATCTGATCCGGGT AGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGAATTTCTTGAAG ACGAAAGGGCCTCCTGATACGCTATTTTTATAGGTTAATGTCATGATAAT AATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGAAATGTGCGCGGAAC CCCTATTTGTTATTTTTCTAAATACATCAAAATATGATCCGCTCATGAG ACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGATGAG TATTCACATTTCCGTGTGCGCCTTATTCCTTTTTTGGGCGATTTTTGCT TCCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGA TCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGGTAA GATCCTTGAGAGTTTTGCGCCCGAAGAAGTTTTTCCAATGATGAGCACTTT TAAAGTCTGCTATGTGGCGGTATTATCCCGTGTGACGCCGGCAAGA GCAACTCGGTGCGCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTC ACCAGTCACAGAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTTATG CAGTGTGCCATAACCATGAGTGATAAACACTGCGGCCAACTTACTTCTGAC AACGATCGGAGGACCGAAGGAGCTAACCGTTTTTTGCACAACATGGGGGA TCATGTAACCTCGCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACC AAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCAACAACGTTGCG CAAACATAAATGCGCAACTACTTACTCTAGCTTCCCGCAACAATTAAT AGACTGGATGGAGCGGATAAAGTTGCAGGACCCTTCTGCGCTCGGCCCT TCCGGCTGGCTGGTTTATGTGATAAATCTGGAGCCGTGACCGTGGGTC TCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCCTATCGT AGTTATCTACACGCGGGGAGTCAGGCAACTATGGATGAAAGATCAAGACA GATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGCTAATCTCAGACCA AGTTTACTCATATACTTTAGATTGATTTAAAACCTCAATTTTTAATTTAA AAGGATCTAGGTGAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTA ACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAGG ATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCTTGCAAAACAAA AAAACCAACCGCTACCGAGCGGTGGTTTGTGTTGCCGGATCAAGAGCTACCAAC TCTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGATACCAAACTACTG TCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCC GCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGG CGATAAGTCTGCTTACCGGTTGGACTCAAGACGATAGTTACCGGATAA GGCGCAGCGGTGCGGCTGAACGGGGGGTTCGTGCACACAGCCAGCTTGA CGAAGACCTACACCGAAGTACGATACCTACAGCGTACGATGAGAAAG CGCCACGCTTCCGGAAGGAGAAAGCGGACAGGTATCCGGTAAAGCGCAG GGTCGGAACAGGAGAGCGCACGAGGAGCTTCCAGGGGAAACGCTGGTA TCTTTATAGTCTGCTGGGTTTCGCCACCTCTGACTTGAGCGTTCGATTTTT GTGATGCTCGTCAGGGGGCGGAGCCTATGAAAAACGCCAGCAACCGCGC CTTTTTACGGTCTGGCCTTTTGTGGCCTTTTGTCTACATGTTCTTTCC TGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGC TGATAACGCTCGCCGACCGCAACGACCGAGCGCAGCGAGT/CAGT/GAGCGA GGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCCCGCGCTTGGCC GATTCATTAATGACGCTGGCAGCAGGTTTCCCGAGTATGCAAGCATGCTCA TGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCAGGC TTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTTGTGAGCGGATA ACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTCTAGCT AGAGGTCGAGTCCCTCCCAGCAGGCAGAAAGTATGCAAGCATGCTCA ATTAGTCAGCAACCATAGTCCCGCCCTAATCCGCCATCCCCCCTAA CTCCGCCAGTTCCGCCATTCGCCCCATGGCTGACTAATTTTTTTTA</p>

SEQ ID NO	Vector name	Nucleotide sequences 123456789012345678901234567890123456789012345678901
		TTTATGCAGAGGCCGAGGCCGCTCGGCCCTCTGAGCTATTCAGAAGTAGT GAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTTTGCAAAAGATG GATAAAGTTTTAAACAGAGAGGAATCTTTGCAGCTAATCCACCTTCTAGGT CTTGAAAGGAGTGGGAATTGGCTCCCGGTGCCCGTCAGTGGGCAGAGCGCAC ATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGGTCCGCAATTGAACCGG TGCCTAGAGAAGTGGCGCGGGGTAACCTGGGAAAGTGATGTCGTGTACTG GCTCCGCCTTTTTCCCAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGT CGCCGTGAACGTTCTTTTTCCGAACGGCTTTGCCGCCAGAACACAGGTAAG TGCCGTGTGTGGTTCCCGCGGGCCTGGCCCTCTTACGGGTATGGCCCTTG CGTGCCTTGAATTAATCCACCTGGCTGCAGTACGTGATCTTGATCCCGA GCTTCGGGTTGGAGTGGGTGGGAGAGTTTCGAGGCCTTTCGAGGAGC CCGTTTCGCTCGTGTGAGTTGAGGCTGGCCCTGGCGCTGGGGCCCGG CGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGCTTTCGATAAGTC TCTAGCCATTTAAAATTTTTGATGACCTGCTGCGACGCTTTTTTCTGGCA AGATAGTCTTTGTAATGCGGGCCAAGATCTGCACACTGGTATTCGTTTTT TGGGGCCGCGGGCGGGCACGGGGCCCGTGGCTCCAGCGCACATGTTCCGGC GAGGCGGGCCCTCGAGCGCGGCCACCGAGAATCGGACGGGGGTAGTCTCA AGCTGGCCGGCCTGCTCTGGTGCCTGGCCCTCGCGCCCGCTGTATCGCCCC GTCCTGGGCGGCAAGGCTGGCCCGGTTCGGCACCAAGTTGGTGGAGGAAAG ATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAAAGGAGGACGGGGC CTCGGGAGAGCGGGCGGGTGAATCACCCACACAAAGGAAAGGGCCCTTCC GTCCTCAGCCGTCGCTTCAATGACTCCACGGAGTACCGGGCCCGTCCAG GCACCTCSATTAGTTCTCGAGCTTTTGGAGTACGTGCTGTTAGGTGGGG GGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACTGA AGTTAGGCCAGCTTGGCACTTGTGTAATCTCCTTGGAAATTTGCCCTTTT TGAGTTGGATCTTGGTTCATCTCAAGCCTCAGACAGTGGTTCAAAGTTT TTTTCTCCATTCAGGTGTGCTGAGGAATTTCTCTAGACTCCCTCGACCT CGAGATCCATTGTGCCGGGGCCACCATGGACATGCGCGTGGCCGGCCAG CTGCTGGGCCTGCTGCTGCTGTTTCCCCGGCTCGCGATGC
309	V7	GCCTCGACCAAGGGCCATCGGTCTTCCCCCTGGCACCTCCTCCAAGAGC ACCTCTGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCC GAACCGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCGCGGGTGCAC ACCTTCCCGGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTG GTGACCGTGCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTG AATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCCAATCT TGTGACAAAACCTCACACATGCCACCGTCCCGAGCACCTGAGCGGGGG GGACCGTCAGTCTTCTCTTCCCCCAAACCAAGCACACCTCATGATC TCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAGAC CCTGAGGTCAAGTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCC AAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTGCTGGTGCAGC GTCCTCACCGTCTTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC AAGTCTCCAACAAGCCCTCCAGCCCCATCGAGAAAACCATCTCCAAA GCCAAAGGGCAGCCCGAGAACCACAGGTGTACACCTGCCCCATCCCGC GAGGAGATGACCAAGAACCAGTTCAGCCTGACCTGCCGTGTCAGGCTTTC TATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC AACTACAAGACCACGCCTCCCGTGTGACTCCGACGGCTCCTTCTCCTC TACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGCTTC TCATGCTCCGTGATGCATGAGGCTCTGCACAACCCTAGACGCAGAGAGC CTCTCCCTGTCTCCGGTAAATGAGCGGCCGCTCGAGGCCGGCAAGGCCG ATCCCCGACCTCGACCTCTGGCTAATAAAGGAAATTTATTTTCATGCAA TAGTGTGTTGGAATTTTTGTGTCTCTCACTCGGAAGGCATATGGGAGG CAAATCATTGGTCGAGATCCCTCGGAGATCTTAGCTAGAGGATCGATCC CCGCCCCGACGAACTAAACCTGACTACGACATCTCTGCCCTTCTTCGCG GGGCAGTGCATGTAATCCCTTCAGTTGGTGGTACAACCTGCCAATGGGC CCTGTTCACATGTGACACGGGGGGGACCAACACAAGGGGTTCTCTGA CTGTAGTTGACATCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTG GCTTTTCATCCCTGGAGCAGACTTTGCACTCTGTGGACTGCAACACAACATG CCTTTATGTGTAACCTTTGGCTGAAGCTCTTACACCAATGCTGGGGGACAT GTACCTCCCAGGGGGCCAGGAAGACTACGGGAGGCTACACCAACGTCATC AGAGGGCCCTGTGTAGCTACCGATAAGCGGACCTCAAGAGGCGATTAGCA ATAGTGTTTATAAGGCCCCCTTGTAAACCTAAACGGGTAGCATCTCTTC CCGGTAGTAGTATATACTATCCAGACTAACCTAATTCMAATAGCATATCT TACCAACGGGAAGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCCTA

SEQ ID NO	Vector name	Nucleotide sequences 123456789012345678901234567890123456789012345678901
		AGGAACAGCGATATCTCCCACCCCATGAGCTGTCACGGTTTTATTTACATG GGGTCAGGATTCCACGAGGGTAGTGAACCATTTTAGTCACAAGGGCAGTGG CTGAAGATCAAGGAGCGGGCAGTGAACCTCCTGAATCTTCGCCTGCTTCT ICATTCCTCCTCGTTAGCTAATAGAACTGCTGAGTTGTGAACAGTAA GGTGTATGTGAGGTGCTCGAAAAACAAGGTTTCAGGTGACGCCCCAGAATA AAATTTGGACGGGGGTTTCAGTGGTGGCATGTGTCTATGACACCAATATA CCCTCACAACCCCTTGGGCAATAAATACTAGTGTAGGAATGAAACATTTCT CAATATCTTTAACAATAGAAATCCATGGGGTGGGACAAAGCCGTAAAGACT GGATGTCATCTCACACGAATTTATGGCTATGGGCAACACATAATCCTAGT GCAATATGATACTGGGTTATTAAGATGTGTCCAGGCAGGGACCAAGACA GGTGAACCATGTTGTTACACTCTATTTGTAACAAGGGGAAAGAGTGGAC GCCGACAGCAGCGGACTCCACTGGTTGCTCTAACACCCCCGAAAAATAAA CGGGGCTCCACGCCAATGGGGCCATAAACAAAGACAAGTGGCCACTCTTT TTTTGAATTTGTGGAGTGGGGGCACGGCTCAGCCCCACACGCGCCCTG CGGTTTTGGACTGTAAAAATAAGGGTGTAACTTGGCTGATTTGTAACCCC GCTAACCACTGCGGTCAAACCCTTGCCCCAAAAACCCTAATGGCACCCC GGGGAATACCTGCATAAGTAGGTGGGCGGGCCAAGATAGGGGCGCGATTCG TCGGATCTGGAGGACAAATTACACACACTTGGCCTGAGCGCCAAGCACAG GGTGTGGTCTCATATTCACGAGGTGCTGAGAGCACGGTGGGCTAATG TTGCCATGGGTAGCATATACTACCCAAATATCTGGATAGCATATGCTATCC TAATCTAIATCTGGGTAGCATAGGCTATCCATCTATATCTGGGTAGCAT ATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATTTATATCT GGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTAATA TCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATG CTATCCTAATAGAGATAGGGTAGTATATGCTATCCTAATTTATATCTGGG TAGCATAIACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATAT CTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGCATATCCTAATCT AATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATA TGCTATCCTAATTTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTG GGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAAT CTGTATCCGGGTAGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGA ATTTTCTTGAAGACGAAAGGGCTCGTGATACGCCTATTTTATAGCTTAA TGTCATGATAAATAAGTTTCTTAGACGTCAGGTGGCCTTTTCGGGGAAA TGTGCGCGGAACCCCTATTTGTTATTTTTCTAAATACATTCAAATATGTA ICCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATGAAAAAG GAAGAGTATGAGTATCAACATTTCCGTGTCGCCCTTATCCTTTTTTGC GGCATTTGCTTCTGTTTTGCTCACCAGAAACCGTGGTGAAGTAAA AGATCCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAAGCTGATCT CAACAGCGGTAGATCCTTGAGAGTTTTCGCCCCGAAGACGTTTTTCCAAT GATGAGCACTTTAAAGTTCTGCTATGTGGCGCGGTATATCCCGTGTGA CGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTTCTCAGAATGACTT GGTGAGTACTCACCAGTACAGAAAAGCATCTTACGGATGGCATGACAGT AAGAGAATTATGCACTGCTGCCATAACCATGAGTGATAACACTGCGGCCAA CTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCCCTTTTTTGCA CAACATGGGGGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAA TGAAGCCATACCAAACGACGAGCGTGACACCACGATGCTGACGCAATGGC AACACGTTGCGCAAACTATTAACCTGGCGAACTACTTACTCTAGCTTCCCG GCAACAATTAATAGACTGGATGGAGGCGGATAAAGTTGACAGGACCACTTCT CGGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGCTGATAAATCTGGACCGG TGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAAGCC CTCCTGATCGTGTATCTACACGACGGGGAGTCAGGCAACTATGGATGA ACGAAATAGACAGATCGCTGAGATAGGCTGCTCACTGATTAAGCATGGTA ACTGTCAGACCAAGTTTACTCATATATACTTTAGATTGATTTAAAAGTTCA TTTTTAATTTAAAAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGAC CAAAATCCCTTAACGTGAGTTTTCGTTCCACTGAGCGTCAGACCCCGTAGA AAAGATCAAAGSATCTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTG CTTGCAAAACAAAAACCACCGCTACCAGCGGTGGTTTTGTTCCCGGATCA AGAGCTACCAACTCTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGAT ACCAAATACTGTTCTTAGTGTAGCCGTAGTGGCCACCACTTCAAGAA CTCTGTAGCACCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGC TGCTGCCAGTGGCGATAAGTCTGTCTTACCAGGTTGAGTCAAGACGATA GTTACCAGATAAGGCGCAGCGCTCGGCTGAACGGGGGTTCGTGCACACA GCCCAGCTTGGAGCGAACGACCTACACCGAAGTGGATACCTACAGCGTGA

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901 GCTATGAGAAAGCGCCACGCTTCCCAGAGGGAGAAAGGCCGACAGGTATCC GGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGG AAACGCCTGGTATCTTTATAGTCTGTCCGGTTTCGCCACCTCTGACTTGA GCGTCGATTTTTGTGATGCTCGTCAGGGGGCGGAGCCTATGGAAAAACGC CAGCAACCGCGGCTTTTTACGGTTCCTGGCCTTTTGCTGCCTTTTGTCTCA CATGTTCTTCCCTGCGTTATCCCTGATTCTGTGGATAACCGTATTACCGC CTTTGAGTGAGCTGATAACCGCTCGCCGACCCGAACCGACCCGACCGGA GTCAGTGAGCGAGGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCC CGCGCGTTGGCCGATTCAATTAATGCAGCTGGCAGACAGGTTTCCCAGCTG GAAAGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAA GGCACCCCAAGGCTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAAAT TGTGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATTACGC CAAGCTCTAGCTAGAGGTCGAGTCCCTCCCCAGCAGGCGAAGTATGCAAA GCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCTAACTCCGCCCA TCCCGCCCTAACTCCGCCAGTTCCGCCCATTTCTCCGCCCTGTGTGTGAC TAAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCTCCGCCCTCTGAGCTAT TCCAGAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGC TTTGCAAAGATGCATAAAGTTTTAAACAGAGAGGAATCTTTGCAGCTAATG GACCTTCTAGGCTTTGAAAGGAGTGGGAATFGGCTCCGGTGCCTCGTAGTG GGCAGAGCGCACATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGTTCGG CAATGAACCGGTGCTTAGAGAGGTTGGCGCGGGGTAAGTGGGAAAGTGA TGTCTGTACTGGCTCCGCCTTTTTCCCGAGGGTGGGGGAGAACCCTATAT AAGTGCAGTAGTCGCCGTGAACGTTCTTTTTGCAACGGTTTGCGCCAG AACACAGGTAAGTGCCGTGTGTGGTTCCCGCGGCCTGGCCTCTTTACGGG TTATGGCCCTTGCGTGCCTTGAATTAATTAATTAATTAATTAATTAATTAAT TCTTGATCCCGAGCTTCCGGTTGGAAGTGGTGGGAGAGTTCCGAGGCCTTG CGCTTAAGGAGCCCTTCGCCTCGTGTGTTGAGTTGAGGCTGGCCTGGCCGG CTGGGGCCCGCGGTCCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGC TTTGATAAGTCTTAGCCATTTAAAATTTTTGATGACCTGCTGCGACGCT TTTTTTCTGGCAAGATAGTCTTGTAATGCGGGCCAAGTCTGCACACTGG TATTTCTGTTTTTGGGCCCGGGCGGGCGGACGGGGCCCGTGCCTCCAGCG CACATGTTCCGGCAGGCGGGGCTGCGAGCGCGGCCACCGAGAATCCGACG GGGTAGTCTCAAGCTGGCCGCCCTGCTCTGGTGCCTGGCCTCGCGCCGCC GTGTATCGCCCCCCTGGGGCGCAAGGCTGGCCCGTGGCACCAGTTGC CTGAGCGAAAGATGGCCGCTTCCGGCCCTGCTGCAGGGAGCTCAAAATG GAGGACCGGGCGCTCGGGAGAGCGGGCGGGTGAGTACCACACAAAAGGAA AAGGGCCTTTCCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGACTACCG GCGCGCTCCAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTCGTC TTTAGGTTGGGGGAGGGGTTTTATGCGATGGAGTTTCCCACACTGAGTG GGTGGAGACTGAAGTTAGGCCAGCTTGGCACTTGATGTAATTTCTCCTTGA ATTTGCCCTTTTTGAGTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGT CGTTCAAAGTTTTTTCTCCATTTAGGTTGTCGTGAGGAATTTCTTAGAG ATCCCTCGACCTCGAGATCCATTTGTGCCGGGGCGCCACCATGGAGTTTGGG CTGAGCTGGCTTTTCTTGTCCGATTTTAAAAGGTGTCAGTGC

The present disclosure incorporates by reference in their entirety techniques well known in the field of molecular biology and drug delivery. These techniques include, but are not limited to, techniques described in the following publications:

- Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993);
- 5 Ausubel, F.M. et al. eds., *Short Protocols In Molecular Biology* (4th Ed. 1999) John Wiley & Sons, NY. (ISBN 0-471-32938-X).
- Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984);
- Giege, R. and Ducruix, A. Barrett, *Crystallization of Nucleic Acids and Proteins, a Practical*
- 10 *Approach*, 2nd ed., pp. 20 1-16, Oxford University Press, New York, New York, (1999);
- Goodson, in *Medical Applications of Controlled Release*, vol. 2, pp. 115-138 (1984);
- Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981);
- Harlow et al. , *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed.
- 15 1988);
- Kabat et al., *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, Md. (1987) and (1991);
- Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242;
- 20 Kontermann and Dubel eds., *Antibody Engineering* (2001) Springer-Verlag. New York. 790 pp. (ISBN 3-540-41354-5).
- Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990);
- Lu and Weiner eds., *Cloning and Expression Vectors for Gene Function Analysis* (2001) BioTechniques Press. Westborough, MA. 298 pp. (ISBN 1-881299-21-X).
- 25 *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Fla. (1974);
- Old, R. W. & S.B. Primrose, *Principles of Gene Manipulation: An Introduction To Genetic Engineering* (3d Ed. 1985) Blackwell Scientific Publications, Boston. *Studies in Microbiology*; V.2:409 pp. (ISBN 0-632-01318-4).
- 30 Sambrook, J. et al. eds., *Molecular Cloning: A Laboratory Manual* (2d Ed. 1989) Cold Spring Harbor Laboratory Press, NY. Vols. 1-3. (ISBN 0-87969-309-6).
- Sustained and Controlled Release Drug Delivery Systems*, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978
- Winnacker, E.L. *From Genes To Clones: Introduction To Gene Technology* (1987) VCH
- 35 Publishers, NY (translated by Horst Ibelgauf). 634 pp. (ISBN 0-89573-614-4).

Incorporation by Reference

The contents of all cited references (including literature references, patents, patent applications, and websites) that maybe cited throughout this application are hereby expressly incorporated by reference in their entirety for any purpose, as are the references cited therein.

- 5 The practice of the present disclosure will employ, unless otherwise indicated, conventional techniques of immunology, molecular biology and cell biology, which are well known in the art.

Equivalents

- 10 The disclosure may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting. Scope of the disclosure is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are therefore intended to be embraced herein.

15

CLAIMS:

1. A binding protein comprising first and second polypeptide chains, each independently comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first variable domain;

VD2 is a second variable domain;

C is a constant domain;

X1 is a linker;

X2 is an Fc region;

n is 0 or 1;

wherein the VD1 domains on the first and second polypeptide chains form a first functional target binding site and the VD2 domains on the first and second polypeptide chains form a second functional target binding site, and wherein the binding protein is capable of binding Tumor Necrosis Factor (TNF) and Nerve Growth Factor (NGF), wherein the binding protein comprises any one of:

DVD1459 (comprising SEQ ID NOs: 52 and 53); DVD1460 (comprising SEQ ID NOs: 54 and 55); DVD1461 (comprising SEQ ID NOs: 56 and 57); DVD1462 (comprising SEQ ID NOs: 58 and 59); DVD1463 (comprising SEQ ID NOs: 60 and 61); DVD1464 (comprising SEQ ID NOs: 62 and 63); DVD1477 (comprising SEQ ID NOs: 64 and 65); DVD1478 (comprising SEQ ID NOs: 66 and 67); DVD1485 (comprising SEQ ID NOs: 68 and 69); DVD1486 (comprising SEQ ID NOs: 70 and 71); DVD1501 (comprising SEQ ID NOs: 136 and 137); DVD1502 (comprising SEQ ID NOs: 138 and 139); DVD1503 (comprising SEQ ID NOs: 140 and 141); DVD1504 (comprising SEQ ID NOs: 142 and 143); DVD1505 (comprising SEQ ID NOs: 144 and 145); DVD1506 (comprising SEQ ID NOs: 146 and 147); DVD1519 (comprising SEQ ID NOs: 148 and 149); DVD1520 (comprising SEQ ID NOs: 150 and 151); DVD1527 (comprising SEQ ID NOs: 152 and 153); DVD1528 (comprising SEQ ID NOs: 154 and 155); DVD1543 (comprising SEQ ID NOs: 220 and 221); DVD1544 (comprising SEQ ID NOs: 222 and 223); DVD1545 (comprising SEQ ID NOs: 224 and 225); DVD1546 (comprising SEQ ID NOs: 226 and 227); DVD1547 (comprising SEQ ID NOs: 228 and 229); DVD1548 (comprising SEQ ID NOs: 230 and 231); DVD1561 (comprising SEQ ID NOs: 232 and 233); DVD1562 (comprising SEQ ID NOs: 234 and 235); DVD1569 (comprising SEQ ID NOs: 236 and 237); and DVD1570 (comprising SEQ ID NOs: 238 and 239).

2. The binding protein according to claim 1, comprising two first and two second polypeptide chains, and four functional target binding sites.

3. The binding protein according to any one of claims 1-2, wherein the binding protein is capable of binding TNF and NGF, and is capable of binding to TNF with a dissociation constant (K_D) of at most about 3.00×10^{-10} M as measured by surface plasmon resonance, and/or is capable of binding to NGF with a dissociation constant (K_D) of at most about 3.40×10^{-12} M as measured by surface plasmon resonance and/or capable of inhibiting NGF with an EC_{50} of at most about 1.946 nM as measured with an NGF inhibition assay.

4. The binding protein according to any one of claims 1-3, wherein

- (a) the Fc region is a variant sequence Fc region,
- (b) the Fc region is an Fc region from an IgG1, IgG2, IgG3, IgG4, IgA, IgM, IgE, or IgD, and/or
- (c) the binding protein is a crystallized binding protein.

5. A binding protein conjugate comprising the binding protein according to any one of claims 1-4, the binding protein conjugate further comprising an immunoadhesion molecule, an imaging agent, a therapeutic agent, or a cytotoxic agent,

wherein the imaging agent is optionally selected from the group consisting of a radiolabel, an enzyme, a fluorescent label, a luminescent label, a bioluminescent label, a magnetic label, and biotin, wherein the radiolabel is optionally selected from the group consisting of: ^3H , ^{14}C , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I , ^{177}Lu , ^{166}Ho , and ^{153}Sm , or

wherein the therapeutic or cytotoxic agent is optionally selected from the group consisting of an anti-metabolite, an alkylating agent, an antibiotic, a growth factor, a cytokine, an anti-angiogenic agent, an anti-mitotic agent, an anthracycline, toxin, and an apoptotic agent.

6. An isolated nucleic acid encoding the binding protein amino acid sequence according to any one of claims 1-5.

7. A vector comprising the isolated nucleic acid according to claim 6, wherein the vector is optionally selected from the group consisting of pcDNA, pTT, pTT3, pEFBOS, pBV, pJV, pcDNA3.1 TOPO, pEF6, pHybE, TOPO, and pBJ.

8. An isolated host cell comprising the vector according to claim 7, wherein the host cell is optionally selected from the group consisting of a prokaryotic cell, *Escherichia coli*, a eukaryotic cell, a protist cell, an animal cell, a plant cell, a fungal cell, a yeast cell, an Sf9 cell, a mammalian cell, an avian cell, an insect cell, a CHO cell, and a COS cell.
9. A method of producing a binding protein, comprising culturing the host cell of claim 8 in culture medium under conditions sufficient to produce the binding protein.
10. A pharmaceutical composition comprising the binding protein of any one of claims 1-5, and a pharmaceutically acceptable carrier.
11. The pharmaceutical composition of claim 10, further comprising at least one additional therapeutic agent, wherein the additional therapeutic agent is optionally selected from the group consisting of an imaging agent, a cytotoxic agent, an angiogenesis inhibitor, a kinase inhibitor, a co-stimulation molecule blocker, an adhesion molecule blocker, an anti-cytokine antibody or functional fragment thereof, methotrexate, cyclosporin, rapamycin, FK506, a detectable label or reporter, a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, an erythropoietin, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, a radiopharmaceutical, an antidepressant, an antipsychotic, a stimulant, an asthma medication, a beta agonist, an inhaled steroid, an epinephrine or analog, a cytokine, and a cytokine antagonist.
12. An in vitro method of determining the presence, amount, or concentration of at least one target or fragment thereof in a test sample by an immunoassay,
 - wherein the immunoassay comprises contacting the test sample with at least one binding protein and at least one detectable label,
 - wherein the at least one binding protein comprises the binding protein of any one of claims 1-5.
13. The method of claim 12, further comprising:
 - (a) contacting the test sample with the at least one binding protein, wherein the binding protein binds to an epitope on the target or fragment thereof so as to form a first complex;

(b) contacting the complex with the at least one detectable label, wherein the detectable label binds to the binding protein or an epitope on the target or fragment thereof that is not bound by the binding protein to form a second complex; and

(c) detecting the presence, amount, or concentration of the target or fragment thereof in the test sample based on the signal generated by the detectable label in the second complex, wherein the presence, amount, or concentration of the target or fragment thereof is directly correlated with the signal generated by the detectable label.

14. The method of claim 12, further comprising:

(a) contacting the test sample with the at least one binding protein, wherein the binding protein binds to an epitope on the target or fragment thereof so as to form a first complex;

(b) contacting the complex with the at least one detectable label, wherein the detectable label competes with the target or fragment thereof for binding to the binding protein so as to form a second complex; and

(c) detecting the presence, amount, or concentration of the target or fragment thereof in the test sample based on the signal generated by the detectable label in the second complex, wherein the presence, amount, or concentration of the target or fragment thereof is indirectly correlated with the signal generated by the detectable label.

15. A kit for assaying an in vitro test sample for the presence, amount, or concentration of a target or fragment thereof, the kit comprising (a) instructions for assaying the test sample for the target or fragment thereof and (b) at least one binding protein comprising the binding protein of any one of claims 1-5.

16. Use of the binding protein according to any one of claims 1-5 in the manufacture of a medicament for treating a subject for a disease or a disorder.

17. The use according to claim 16, wherein the disorder is arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's granulomatosis, Henoch-

Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, Addison's disease, sporadic, polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella associated arthropathy, atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous candidiasis, giant cell arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, acquired immunodeficiency related diseases, hepatitis B, hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-inflammatory interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease, systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated lung disease, Sjögren's disease associated lung disease, ankylosing spondylitis associated lung disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced interstitial lung disease, fibrosis, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated hypoglycaemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute immune disease associated with organ transplantation, chronic immune disease associated with organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS, glomerulonephritides, microscopic vasculitis of the kidneys, lyme disease, discoid lupus erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all subtypes), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic sclerosis, Sjögren's syndrome, Takayasu's disease/arteritis, autoimmune

thrombocytopaenia, idiopathic thrombocytopaenia, autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary myxoedema, phacogenic uveitis, primary vasculitis, vitiligo acute liver disease, chronic liver diseases, alcoholic cirrhosis, alcohol-induced liver injury, choleosatatis, idiosyncratic liver disease, drug-induced hepatitis, non-alcoholic steatohepatitis, allergy and asthma, group B streptococci (GBS) infection, mental disorders, depression, schizophrenia, Th2 Type and Th1 Type mediated diseases, acute and chronic pain, pain, cancers, lung cancer, breast cancer, stomach cancer, bladder cancer, colon cancer, pancreatic cancer, ovarian cancer, prostate cancer, rectal cancer, hematopoietic malignancies, leukemia, lymphoma, Abetalipoproteinemia, acrocyanosis, acute and chronic parasitic or infectious processes, acute leukemia, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinomas, aerial ectopic beats, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allograft rejection, alpha-1-antitrypsin deficiency, amyotrophic lateral sclerosis, anemia, angina pectoris, anterior horn cell degeneration, anti cd3 therapy, antiphospholipid syndrome, anti-receptor hypersensitivity reactions, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, arteriosclerosis, arteriovenous fistula, ataxia, atrial fibrillation (sustained or paroxysmal), atrial flutter, atrioventricular block, B cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt's lymphoma, burns, cardiac arrhythmias, cardiac stun syndrome, cardiac tumors, cardiomyopathy, cardiopulmonary bypass inflammation response, cartilage transplant rejection, cerebellar cortical degenerations, cerebellar disorders, chaotic or multifocal atrial tachycardia, chemotherapy associated disorders, chronic myelocytic leukemia (CML), chronic alcoholism, chronic inflammatory pathologies, chronic lymphocytic leukemia (CLL), chronic obstructive pulmonary disease (COPD), chronic salicylate intoxication, colorectal carcinoma, congestive heart failure, conjunctivitis, contact dermatitis, cor pulmonale, coronary artery disease, Creutzfeldt-Jakob disease, culture negative sepsis, cystic fibrosis, cytokine therapy associated disorders, dementia pugilistica, demyelinating diseases, dengue hemorrhagic fever, dermatitis, dermatologic conditions, diabetes, diabetes mellitus, diabetic atherosclerotic disease, diffuse Lewy body disease, dilated congestive cardiomyopathy, disorders of the basal ganglia, Down's syndrome in middle age, drug-induced movement disorders induced by drugs which block CNS dopamine receptors, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottitis, Epstein-Barr virus infection, erythromelalgia, extrapyramidal and cerebellar disorders, familial hematosphagocytic lymphohistiocytosis, fetal thymus implant rejection, Friedreich's ataxia, functional peripheral arterial disorders, fungal sepsis, gas gangrene, gastric ulcer, glomerular nephritis, graft rejection of any organ or tissue, gram negative sepsis, gram positive sepsis, granulomas

due to intracellular organisms, hairy cell leukemia, Hallervorden-Spatz disease, Hashimoto's thyroiditis, hay fever, heart transplant rejection, hemachromatosis, hemodialysis, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, hemorrhage, hepatitis A, His bundle arrhythmias, HIV infection/HIV neuropathy, Hodgkin's disease, hyperkinetic movement disorders, hypersensitivity reactions, hypersensitivity pneumonitis, hypertension, hypokinetic movement disorders, hypothalamic-pituitary-adrenal axis evaluation, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody mediated cytotoxicity, Asthenia, infantile spinal muscular atrophy, inflammation of the aorta, influenza a, ionizing radiation exposure, iridocyclitis, uveitis, optic neuritis, ischemia- reperfusion injury, ischemic stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, kidney transplant rejection, legionella, leishmaniasis, leprosy, lesions of the corticospinal system, lipedema, liver transplant rejection, lymphedema, malaria, malignant lymphoma, malignant histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic, migraine headache, mitochondrial multi-system disorder, mixed connective tissue disease, monoclonal gammopathy, multiple myeloma, multiple systems degenerations, mycobacterium avium intracellulare, mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disorders, nasopharyngeal carcinoma, neonatal chronic lung disease, nephritis, nephrosis, neurodegenerative diseases, neurogenic muscular atrophies, neutropenic fever, non-hodgkins lymphoma, occlusion of the abdominal aorta and its branches, occlusive arterial disorders, okt3 therapy, orchitis/epididymitis, orchitis/vasectomy reversal procedures, organomegaly, osteoporosis, pancreas transplant rejection, pancreatic carcinoma, paraneoplastic syndrome/hypercalcemia of malignancy, parathyroid transplant rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral atherosclerotic disease, peripheral vascular disorders, peritonitis, pernicious anemia, pneumocystis carinii pneumonia, pneumonia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy, and skin changes syndrome), post perfusion syndrome, post pump syndrome, post-MI cardiomyopathy syndrome, preeclampsia, progressive supranucleo palsy, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon and disease, Raynaud's disease, Refsum's disease, regular narrow QRS tachycardia, renovascular hypertension, reperfusion injury, restrictive cardiomyopathy, sarcomas, scleroderma, senile chorea, senile dementia of Lewy body type, seronegative arthropathies, shock, sickle cell anemia, skin allograft rejection, skin changes syndrome, small bowel transplant rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degenerations, streptococcal myositis, structural lesions of the cerebellum, subacute sclerosing panencephalitis, syncope, syphilis of the cardiovascular system, systemic anaphalaxis, systemic inflammatory response syndrome, systemic onset juvenile rheumatoid arthritis, T-cell or FAB ALL, telangiectasia, thromboangitis obliterans, thrombocytopenia, toxicity, transplants,

trauma/hemorrhage, type III hypersensitivity reactions, type IV hypersensitivity, unstable angina, uremia, urosepsis, valvular heart diseases, varicose veins, vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal infections, viral encephalitis/aseptic meningitis, viral-associated hemaphagocytic syndrome, Wernicke-Korsakoff syndrome, Wilson's disease, xenograft rejection of any organ or tissue, acute coronary syndromes, acute idiopathic polyneuritis, acute inflammatory demyelinating polyradiculoneuropathy, acute ischemia, adult Still's disease, anaphylaxis, anti-phospholipid antibody syndrome, aplastic anemia, atopic eczema, atopic dermatitis, autoimmune dermatitis, autoimmune disorder associated with streptococcus infection, autoimmune enteropathy, autoimmune hearing loss, autoimmune lymphoproliferative syndrome (ALPS), autoimmune myocarditis, autoimmune premature ovarian failure, blepharitis, bronchiectasis, bullous pemphigoid, cardiovascular disease, catastrophic antiphospholipid syndrome, celiac disease, cervical spondylosis, chronic ischemia, cicatricial pemphigoid, clinically isolated syndrome (cis) with risk for multiple sclerosis, childhood onset psychiatric disorder, dacryocystitis, dermatomyositis, diabetic retinopathy, disk herniation, disk prolapse, drug induced immune hemolytic anemia, endometriosis, endophthalmitis, episcleritis, erythema multiforme, erythema multiforme major, gestational pemphigoid, Guillain-Barré syndrome (GBS), Hughes syndrome, idiopathic Parkinson's disease, idiopathic interstitial pneumonia, IgE-mediated allergy, immune hemolytic anemia, inclusion body myositis, infectious ocular inflammatory disease, inflammatory demyelinating disease, inflammatory heart disease, inflammatory kidney disease, IPF/UIP, iritis, keratitis, keratoconjunctivitis sicca, Kussmaul disease or Kussmaul-Meier disease, Landry's paralysis, Langerhan's cell histiocytosis, livedo reticularis, macular degeneration, microscopic polyangiitis, morbus bechtereiv, motor neuron disorders, mucous membrane pemphigoid, multiple organ failure, myasthenia gravis, myelodysplastic syndrome, myocarditis, nerve root disorders, neuropathy, non-A non-B hepatitis, optic neuritis, osteolysis, pauciarticular JRA, peripheral artery occlusive disease (PAOD), peripheral vascular disease (PVD), peripheral artery disease (PAD), phlebitis, polyarteritis nodosa, periarteritis nodosa, polychondritis, poliosis, polyarticular JRA, polyendocrine deficiency syndrome, polymyositis, polymyalgia rheumatica (PMR), primary Parkinsonism, prostatitis, pure red cell aplasia, primary adrenal insufficiency, recurrent neuromyelitis optica, restenosis, rheumatic heart disease, sapho (synovitis, acne, pustulosis, hyperostosis, and osteitis), secondary amyloidosis, shock lung, scleritis, sciatica, secondary adrenal insufficiency, silicone associated connective tissue disease, sneddon-wilkinson dermatosis, spondilitis ankylosans, Stevens-Johnson syndrome (SJS), temporal arteritis, toxoplasmic retinitis, toxic epidermal necrolysis, transverse myelitis, TRAPS (tumor necrosis factor receptor, type 1 allergic reaction, type II diabetes, urticaria, usual interstitial pneumonia (UIP), vasculitis,

vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), wet macular degeneration, or wound healing.

18. The use according to claim 16 or 17, wherein the medicament is formulated for subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracerebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal, buccal, sublingual, intranasal, or transdermal administration.

19. A method of treating a subject for a disease or disorder, the method comprising administering to said subject a therapeutically effective amount of a binding protein according to any one of claims 1 to 5.

20. The method according to claim 19 wherein wherein the disorder is arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's granulomatosis, Henoch-Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, Addison's disease, sporadic, polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella associated arthropathy, atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous candidiasis, giant cell arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, acquired

immunodeficiency related diseases, hepatitis B, hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-inflammatory interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease, systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated lung disease, Sjögren's disease associated lung disease, ankylosing spondylitis associated lung disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced interstitial lung disease, fibrosis, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated hypoglycaemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute immune disease associated with organ transplantation, chronic immune disease associated with organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS, glomerulonephritides, microscopic vasculitis of the kidneys, lyme disease, discoid lupus erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all subtypes), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic sclerosis, Sjörgren's syndrome, Takayasu's disease/arteritis, autoimmune thrombocytopaenia, idiopathic thrombocytopaenia, autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary myxoedema, phacogenic uveitis, primary vasculitis, vitiligo acute liver disease, chronic liver diseases, alcoholic cirrhosis, alcohol-induced liver injury, choleosatatis, idiosyncratic liver disease, drug-induced hepatitis, non-alcoholic steatohepatitis, allergy and asthma, group B streptococci (GBS) infection, mental disorders, depression, schizophrenia, Th2 Type and Th1 Type mediated diseases, acute and chronic pain, pain, cancers, lung cancer, breast cancer, stomach cancer, bladder cancer, colon cancer, pancreatic cancer, ovarian cancer, prostate cancer, rectal cancer, hematopoietic malignancies, leukemia, lymphoma, Abetalipoproteinemia, acrocyanosis, acute and chronic parasitic or infectious processes, acute leukemia, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinomas, aerial ectopic beats, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic

rhinitis, allograft rejection, alpha-1-antitrypsin deficiency, amyotrophic lateral sclerosis, anemia, angina pectoris, anterior horn cell degeneration, anti cd3 therapy, antiphospholipid syndrome, anti-receptor hypersensitivity reactions, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, arteriosclerosis, arteriovenous fistula, ataxia, atrial fibrillation (sustained or paroxysmal), atrial flutter, atrioventricular block, B cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt's lymphoma, burns, cardiac arrhythmias, cardiac stun syndrome, cardiac tumors, cardiomyopathy, cardiopulmonary bypass inflammation response, cartilage transplant rejection, cerebellar cortical degenerations, cerebellar disorders, chaotic or multifocal atrial tachycardia, chemotherapy associated disorders, chronic myelocytic leukemia (CML), chronic alcoholism, chronic inflammatory pathologies, chronic lymphocytic leukemia (CLL), chronic obstructive pulmonary disease (COPD), chronic salicylate intoxication, colorectal carcinoma, congestive heart failure, conjunctivitis, contact dermatitis, cor pulmonale, coronary artery disease, Creutzfeldt-Jakob disease, culture negative sepsis, cystic fibrosis, cytokine therapy associated disorders, dementia pugilistica, demyelinating diseases, dengue hemorrhagic fever, dermatitis, dermatologic conditions, diabetes, diabetes mellitus, diabetic atherosclerotic disease, diffuse Lewy body disease, dilated congestive cardiomyopathy, disorders of the basal ganglia, Down's syndrome in middle age, drug-induced movement disorders induced by drugs which block CNS dopamine receptors, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottitis, Epstein-Barr virus infection, erythromelalgia, extrapyramidal and cerebellar disorders, familial hemophagocytic lymphohistiocytosis, fetal thymus implant rejection, Friedreich's ataxia, functional peripheral arterial disorders, fungal sepsis, gas gangrene, gastric ulcer, glomerular nephritis, graft rejection of any organ or tissue, gram negative sepsis, gram positive sepsis, granulomas due to intracellular organisms, hairy cell leukemia, Hallervorden-Spatz disease, Hashimoto's thyroiditis, hay fever, heart transplant rejection, hemochromatosis, hemodialysis, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, hemorrhage, hepatitis A, His bundle arrhythmias, HIV infection/HIV neuropathy, Hodgkin's disease, hyperkinetic movement disorders, hypersensitivity reactions, hypersensitivity pneumonitis, hypertension, hypokinetic movement disorders, hypothalamic-pituitary-adrenal axis evaluation, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody mediated cytotoxicity, Asthenia, infantile spinal muscular atrophy, inflammation of the aorta, influenza a, ionizing radiation exposure, iridocyclitis, uveitis, optic neuritis, ischemia- reperfusion injury, ischemic stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, kidney transplant rejection, legionella, leishmaniasis, leprosy, lesions of the corticospinal system, lipedema, liver transplant rejection, lymphoderma, malaria, malignant lymphoma, malignant histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic, migraine headache, mitochondrial multi-

system disorder, mixed connective tissue disease, monoclonal gammopathy, multiple myeloma, multiple systems degenerations, mycobacterium avium intracellulare, mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disorders, nasopharyngeal carcinoma, neonatal chronic lung disease, nephritis, nephrosis, neurodegenerative diseases, neurogenic muscular atrophies, neutropenic fever, non-hodgkins lymphoma, occlusion of the abdominal aorta and its branches, occlusive arterial disorders, okt3 therapy, orchitis/epididymitis, orchitis/vasectomy reversal procedures, organomegaly, osteoporosis, pancreas transplant rejection, pancreatic carcinoma, paraneoplastic syndrome/hypercalcemia of malignancy, parathyroid transplant rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral atherosclerotic disease, peripheral vascular disorders, peritonitis, pernicious anemia, pneumocystis carinii pneumonia, pneumonia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy, and skin changes syndrome), post perfusion syndrome, post pump syndrome, post-MI cardiomyopathy syndrome, preeclampsia, progressive supranuclear palsy, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon and disease, Raynaud's disease, Refsum's disease, regular narrow QRS tachycardia, renovascular hypertension, reperfusion injury, restrictive cardiomyopathy, sarcomas, scleroderma, senile chorea, senile dementia of Lewy body type, seronegative arthropathies, shock, sickle cell anemia, skin allograft rejection, skin changes syndrome, small bowel transplant rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degenerations, streptococcal myositis, structural lesions of the cerebellum, subacute sclerosing panencephalitis, syncope, syphilis of the cardiovascular system, systemic anaphylaxis, systemic inflammatory response syndrome, systemic onset juvenile rheumatoid arthritis, T-cell or FAB ALL, telangiectasia, thromboangitis obliterans, thrombocytopenia, toxicity, transplants, trauma/hemorrhage, type III hypersensitivity reactions, type IV hypersensitivity, unstable angina, uremia, urosepsis, valvular heart diseases, varicose veins, vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal infections, viral encephalitis/aseptic meningitis, viral-associated hemophagocytic syndrome, Wernicke-Korsakoff syndrome, Wilson's disease, xenograft rejection of any organ or tissue, acute coronary syndromes, acute idiopathic polyneuritis, acute inflammatory demyelinating polyradiculoneuropathy, acute ischemia, adult Still's disease, anaphylaxis, anti-phospholipid antibody syndrome, aplastic anemia, atopic eczema, atopic dermatitis, autoimmune dermatitis, autoimmune disorder associated with streptococcus infection, autoimmune enteropathy, autoimmune hearing loss, autoimmune lymphoproliferative syndrome (ALPS), autoimmune myocarditis, autoimmune premature ovarian failure, blepharitis, bronchiectasis, bullous pemphigoid, cardiovascular disease, catastrophic antiphospholipid syndrome, celiac disease, cervical spondylosis, chronic ischemia, cicatricial pemphigoid, clinically isolated syndrome (cis) with risk for multiple sclerosis, childhood onset

psychiatric disorder, dacryocystitis, dermatomyositis, diabetic retinopathy, disk herniation, disk prolapse, drug induced immune hemolytic anemia, endometriosis, endophthalmitis, episcleritis, erythema multiforme, erythema multiforme major, gestational pemphigoid, Guillain-Barré syndrome (GBS), Hughes syndrome, idiopathic Parkinson`s disease, idiopathic interstitial pneumonia, IgE-mediated allergy, immune hemolytic anemia, inclusion body myositis, infectious ocular inflammatory disease, inflammatory demyelinating disease, inflammatory heart disease, inflammatory kidney disease, IPF/UIP, iritis, keratitis, keratoconjunctivitis sicca, Kussmaul disease or Kussmaul-Meier disease, Landry's paralysis, Langerhan's cell histiocytosis, livedo reticularis, macular degeneration, microscopic polyangiitis, morbus bechtereve, motor neuron disorders, mucous membrane pemphigoid, multiple organ failure, myasthenia gravis, myelodysplastic syndrome, myocarditis, nerve root disorders, neuropathy, non-A non-B hepatitis, optic neuritis, osteolysis, pauciarticular JRA, peripheral artery occlusive disease (PAOD), peripheral vascular disease (PVD), peripheral artery disease (PAD), phlebitis, polyarteritis nodosa, periarteritis nodosa, polychondritis, poliosis, polyarticular JRA, polycendocrine deficiency syndrome, polymyositis, polymyalgia rheumatica (PMR), primary Parkinsonism, prostatitis, pure red cell aplasia, primary adrenal insufficiency, recurrent neuromyelitis optica, restenosis, rheumatic heart disease, sapho (synovitis, acne, pustulosis, hyperostosis, and osteitis), secondary amyloidosis, shock lung, scleritis, sciatica, secondary adrenal insufficiency, silicone associated connective tissue disease, sneddon-wilkinson dermatosis, spondylitis ankylosans, Stevens-Johnson syndrome (SJS), temporal arteritis, toxoplasmic retinitis, toxic epidermal necrolysis, transverse myelitis, TRAPS (tumor necrosis factor receptor, type 1 allergic reaction, type II diabetes, urticaria, usual interstitial pneumonia (UIP), vasculitis, vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), wet macular degeneration, or wound healing.


AbbVie Inc.

Patent Attorneys for the Applicant/Nominated Person

SPRUSON & FERGUSON

Figure 1

A



B

