



HU000031957T2



(19) **HU**

(11) Lajstromszám: **E 031 957**

(13) **T2**

MAGYARORSZÁG
Szellemi Tulajdon Nemzeti Hivatala

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 10 817520**

(51) Int. Cl.: **A61K 39/106**

(2006.01)

(22) A bejelentés napja: **2010. 09. 16.**

A61K 39/108

(2006.01)

A61P 31/04

(2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20100817520

(86) A nemzetközi (PCT) bejelentési szám:

PCT/SE 10/050996

(97) Az európai bejelentés közzétételi adatai:

EP 2477649 A1 2011. 03. 24.

(87) A nemzetközi közzétételi szám:

WO 11034495

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2477649 B1 2016. 11. 23.

(30) Elsőbbségi adatok:

272351 P 2009. 09. 16. US

(73) Jogosult(ak):

**MSD Wellcome Trust Hilleman Laboratories
Pvt Ltd., New Delhi 110062 (IN)**

(72) Feltaláló(k):

HOLMGREN, Jan, S-426 74 Västra Frölunda (SE)

(74) Képviselő:

LEBENS, Michael, S-523 99 Hökerum (SE)

Olle Bäcklund, Spanga

(54) **Kolera és enterotoxigenikus E. coli diarrhoea elleni vakcina**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

EP 2 477 649 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
23.11.2016 Bulletin 2016/47

(51) Int Cl.:
A61K 39/106 (2006.01) **A61K 39/108** (2006.01)
A61P 31/04 (2006.01)

(21) Application number: 10817520.9

(86) International application number:
PCT/SE2010/050996

(22) Date of filing: 16.09.2010

(87) International publication number:
WO 2011/034495 (24.03.2011 Gazette 2011/12)

(54) **VACCINE AGAINST CHOLERA AND ENTEROTOXIGENIC E. COLI (ETEC) DIARRHEA**
IMPFSTOFF GEGEN CHOLERA UND ENTEROTOXIGENE E. COLI (ETEC)-DIARRHÖE
VACCIN CONTRE LA DIARRHÉE DUE AU CHOLÉRA ET A E.COLI ENTÉROTOXIGÈNE (ETEC)

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO SE SI SK SM TR**

(30) Priority: 16.09.2009 US 272351 P

(43) Date of publication of application:
25.07.2012 Bulletin 2012/30

(73) Proprietor: **MSD Wellcome Trust Hilleman
Laboratories Pvt Ltd.**
New Delhi 110062 (IN)

(72) Inventors:
• **HOLMGREN, Jan**
S-426 74 Västra Frölunda (SE)
• **LEBENS, Michael**
S-523 99 Hökerum (SE)

(74) Representative: **Novitas Patent AB**
P.O. Box 55557
102 04 Stockholm (SE)

(56) References cited:
WO-A1-00/67784 WO-A1-94/01533
WO-A2-2009/049013 US-A- 5 330 753
US-A- 5 882 653 US-A- 5 942 242
US-A1- 2006 099 229 US-B1- 6 203 799
US-B2- 7 270 961

- **SU L CHIANG AND JOHN J MEKALANOS:**
"Construction of a *Vibrio cholerae* vaccine candidate using transposon delivery and FLP recombinase-mediated excision", *INFECTION AND IMMUNITY, AMERICAN SOCIETY FOR MACROBIOLOGY, USA*, vol. 68, no. 11, 1 January 2000 (2000-01-01), pages 6391-6397, XP008153886, ISSN: 0019-9567, DOI: 10.1128/IAI.68.11
- **S. G RIJPKEMA ET AL:** "Assessing clonality of *Vibrio cholerae* Inaba isolates by characterization of nonsense mutations in *wbeT*", *JOURNAL OF MEDICAL MICROBIOLOGY*, vol. 53, no. 11, 1 November 2004 (2004-11-01), pages 1105-1107, XP055048436, ISSN: 0022-2615, DOI: 10.1099/jmm.0.45744-0
- **KANUNGO S ET AL:** "Immune responses following one and two doses of the reformulated, bivalent, killed, whole-cell, oral cholera vaccine among adults and children in Kolkata, India: A randomized, placebo-controlled trial", *VACCINE*, ELSEVIER LTD, GB, vol. 27, no. 49, November 2009 (2009-11), pages 6887-6893, XP026722082, ISSN: 0264-410X, DOI: 10.1016/J.VACCINE.2009.09.008 [retrieved on 2009-09-15]
- **SANCHEZ J ET AL:** "Virulence factors, pathogenesis and vaccine protection in cholera and ETEC diarrhea", *CURRENT OPINION IN IMMUNOLOGY*, ELSEVIER, OXFORD, GB, vol. 17, no. 4, 1 August 2005 (2005-08-01), pages 388-398, XP027787152, ISSN: 0952-7915 [retrieved on 2005-08-01]

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 477 649 B1

- HOLMGREN J ET AL: "An oral B subunit: whole cell vaccine against cholera", VACCINE, ELSEVIER LTD, GB, vol. 10, no. 13, 1 January 1992 (1992-01-01), pages 911-914, XP023710802, ISSN: 0264-410X, DOI: 10.1016/0264-410X(92)90324-D [retrieved on 1992-01-01]
- SHAMSUZZAMAN S ET AL: "Robust gut associated vaccine-specific antibody-secreting cell responses are detected at the mucosal surface of Bangladeshi subjects after immunization with an oral killed bivalent *V. cholerae* O1/O139 whole cell cholera vaccine: Comparison with other mucosal and systemic responses", VACCINE, ELSEVIER LTD, GB, vol. 27, no. 9, 25 February 2009 (2009-02-25), pages 1386-1392, XP026194432, ISSN: 0264-410X, DOI: 10.1016/J.VACCINE.2008.12.041 [retrieved on 2009-01-13]
- ANH ET AL: "Safety and immunogenicity of a reformulated Vietnamese bivalent killed, whole-cell, oral cholera vaccine in adults", VACCINE, ELSEVIER LTD, GB, vol. 25, no. 6, 20 December 2006 (2006-12-20), pages 1149-1155, XP005809564, ISSN: 0264-410X, DOI: 10.1016/J.VACCINE.2006.09.049
- HOLMGREN J ET AL: "New cholera vaccines", VACCINE, ELSEVIER LTD, GB, vol. 7, no. 2, 1 April 1989 (1989-04-01), pages 94-96, XP023798738, ISSN: 0264-410X, DOI: 10.1016/0264-410X(89)90042-X [retrieved on 1989-04-01]
- CLEMENS J D ET AL: "Field trial of oral cholera vaccines in Bangladesh: evaluation of anti-bacterial and anti-toxic breast-milk immunity in response to ingestion of the vaccines", VACCINE, ELSEVIER LTD, GB, vol. 8, no. 5, 1 October 1990 (1990-10-01), pages 469-472, XP023712189, ISSN: 0264-410X, DOI: 10.1016/0264-410X(90)90248-K [retrieved on 1990-10-01]
- JERTBORN M ET AL: "Safety and immunogenicity of an oral recombinant cholera B subunit-whole cell vaccine in Swedish volunteers", VACCINE, ELSEVIER LTD, GB, vol. 10, no. 2, 1 January 1992 (1992-01-01), pages 130-132, XP023710419, ISSN: 0264-410X, DOI: 10.1016/0264-410X(92)90030-N [retrieved on 1992-01-01]
- LYCKE N ET AL: "Strong biotype and serotype cross-protective antibacterial and antitoxic immunity in rabbits after cholera infection", MICROBIAL PATHOGENESIS, ACADEMIC PRESS LIMITED, NEW YORK, NY, US, vol. 1, no. 4, 1 August 1986 (1986-08-01), pages 361-371, XP023313147, ISSN: 0882-4010, DOI: 10.1016/0882-4010(86)90068-9 [retrieved on 1986-08-01]
- TAYLOR ET AL: "Evaluation of a bivalent (CVD 103-HgR/CVD 111) live oral cholera vaccine in adult volunteers from the United States and Peru.", INFECTION AND IMMUNITY, vol. 65, no. 9, 1 September 1997 (1997-09-01), pages 3852-3856, XP055048355, ISSN: 0019-9567
- TAYLOR ET AL: "Expanded safety and immunogenicity of a bivalent, oral, attenuated cholera vaccine, CVD 103-HgR plus CVD 111, in United States military personnel stationed in Panama.", INFECTION AND IMMUNITY, vol. 67, no. 4, 1 April 1999 (1999-04-01), pages 2030-2034, XP055048356, ISSN: 0019-9567
- MANNING ET AL: "Molecular cloning and expression in *Escherichia coli* K-12 of the O antigens of the Inaba and Ogawa serotypes of the *Vibrio cholerae* O1 lipopolysaccharides and their potential for vaccine development.", INFECTION AND IMMUNITY, vol. 53, no. 2, 1 August 1986 (1986-08-01), pages 272-277, XP055048384, ISSN: 0019-9567
- MICHAEL LEBENS ET AL: "Construction of novel vaccine strains co-expressing the Inaba and Ogawa serotype antigens", VACCINE, vol. 29, no. 43, 30 July 2011 (2011-07-30), pages 7505-7513, XP028306514, ISSN: 0264-410X, DOI: 10.1016/J.VACCINE.2011.06.121 [retrieved on 2011-07-20]
- CHIANG ET AL: 'Construction of a *Vibrio cholerae* vaccine candidate using transposon delivery and FLP recombinase-mediated excision' INFECTION AND IMMUNITY vol. 68, no. 11, 2000, pages 6391 - 6397, XP008153886
- 'PRODUCT MONOGRAPH, DUKORAL(R), ATCC Code: J07AE01', [Online] 2007, XP008159292 Retrieved from the Internet: <URL:https://www.vaccineshoppecanada.com/se_cure/pdfs/ca/dukoral_e.pdf>
- DUTTA ET AL: 'Spread of cholera with newer clones of *Vibrio cholerae* O1 El Tor, serotype inaba, in India' JOURNAL OF CLINICAL MICROBIOLOGY vol. 44, no. 9, 2006, pages 3391 - 3393, XP008153884 & DATABASE UNIPROT [Online] 04 April 2006 EBIDatabase accession no. Q27HR8
- TOBIAS ET AL: 'Construction of non-toxic *Escherichia coli* and *Vibrio cholerae* strains expressing high and immunogenic levels of enterotoxigenic *E. coli* colonization factor I fimbriae' VACCINE vol. 26, no. 6, 2008, pages 743 - 752, XP022425413
- STROEHER ET AL: 'Serotype conversion in *Vibrio cholerae* O1' PNAS vol. 89, no. 7, 1992, pages 2566 - 2570, XP008153868

Description**Technical field the invention**

5 [0001] The present invention relates to the field of vaccines, in particular vaccines against cholera and enterotoxigenic *E. coli* (ETEC) diarrhea.

Background to the invention

10 [0002] Cholera remains a major health problem in large parts of the world. This is also true for ETEC, which is the main cause of diarrheal disease in developing countries as well as in travelers to these countries. In many developing countries effective water and sanitary measures for control of cholera and other enteric infections are currently impossible, and in this context, vaccines have an important role to play. In order to do so however, they need to be effective, readily accessible and above all cheap. There is also a medical need and a very substantial commercial market for use of cholera and especially ETEC vaccines in travelers.

15 [0003] One approach has been the development of oral killed whole cell vaccines. Dukoral™ is an oral cell vaccine (OCV) with demonstrated up to 90% efficacy against cholera and also a significant efficacy against Enterotoxigenic *Escherichia coli* (ETEC)-induced diarrhea. It comprises 3 different *V. cholerae* strains in four different formulations (two heat-killed and two formalin-killed) and in addition recombinantly produced cholera toxin B subunit (rCTB). The rCTB component contributes significantly to the efficacy against cholera and is solely responsible for the observed protection against ETEC diarrhea due to its ability to induce cross-neutralizing antibodies against the cholera toxin (CT)-like *E. coli* heat-labile toxin (LT). However, rCTB is acid-labile and thus the vaccine (which needs to be given in two doses) must be administered with a bicarbonate buffer.

20 [0004] Despite Dukoral™ being the only internationally licensed OCV, copies of this vaccine with or without the CTB component are currently being marketed in developing countries - Vietnam, India and China. The OCV made in Vietnam and India (which lacks the CTB component) contains the same 4 bacterial components as in Dukoral plus a fifth formalin-killed *V. cholerae* strain of serogroup 0139.

25 [0005] Protective immunity against cholera elicited by OCVs is mainly if not exclusively based on mucosal production of antibodies against cell wall lipopolysaccharide O1 (O1 LPS) and for the CTB-containing Dukoral vaccine also antitoxin antibodies in the intestine.

30 [0006] From the above it is evident that the present state of the art for production of cholera/ETEC vaccine is far from simple, and although already effective, a real contribution to making a cholera vaccine more accessible would be to rationalize the composition of the formulation at several levels.

35 [0007] The necessity to include several different *Vibrio cholerae* strains in killed whole cell vaccines such as Dukoral™ arises from the need to represent several different antigenic variants of *Vibrio cholerae* in the vaccine. All protective strains in the currently used vaccines are of the O1 serogroup which until 1993 was the only one of more than 200 identified serogroups known to cause epidemic cholera and is still the dominant serogroup. However, the O1 serogroup has two variants called the Ogawa and Inaba serotypes that differ in the methylation of the terminal sugar of the O-antigen of the surface lipopolysaccharide (LPS). Serotype switching is known to occur in which the Ogawa serotype organism can give rise to Inaba organisms. The reverse switch however is rare.

40 [0008] Although immunization with especially Inaba but also Ogawa serotype can give rise to antibodies cross-reacting with the other serotypes it also gives rise to serotype specific antibodies that contribute significantly to protection. Thus, an effective vaccine should induce not only cross reactive but also serotype-specific antibodies against both Inaba and Ogawa serotype variants.

45 [0009] The serotype switch is known to be related to a mutation in a single gene (*wbeT*). Any mutation that inactivates this gene results in a switch from the Ogawa to the Inaba serotype. Mutations that can reverse such an event are predictably much more uncommon although a switch from the Inaba to the Ogawa serotype can easily be achieved by provision of the relevant gene *in trans*. The gene involved (*wbeT*, also denoted *rfbT*) encodes a methyl transferase that methylates the terminal perosamine residue in the O-antigen polysaccharide repeating unit. Mutations in this gene that lead to the Inaba serotype are almost invariably insertions, deletions or base changes that introduce a nonsense codon.

50 [0010] A third O1 variant known as Hikojima has also been documented to occur in the wild. Hikojima is characterized by that it expresses both the Ogawa and Inaba determinants on its surface and agglutinates with antisera specific for both types. The Hikojima phenotype is extremely rare and is considered in the literature to be an unstable transitional form.

55 [0011] Chiang and Mekalonos (Infection and Immunity, vol. 68, no. 11, 1 January 2000 (2000-01-01), pages 6391-6397, XP008153886, ISSN: 0019-9567, DOI: 10.1128/IAI.68.11) report construction of a *Vibrio cholerae* vaccine candidate using transposon delivery and FLP recombinase mediate excision. A motile Inaba+ vaccine candidate, Peru-2, was converted into nonmotile Ogawa+ by the manipulation.

[0012] Rijkema et al. (Journal of Medical Microbiology, vol. 53, no. 11, 1 November 2004 (2004-11-01), pages

1105-1107, XP055048436, ISSN: 0022-2615, DOI: 10.1099/jmm.0.45744-0) report analysis of *WbeT* mutations in clinical *Vibrio cholerae* isolates.

[0013] Kanungo et al. (Vaccine, vol. 27, no. 49, 16 November 2009 (2009-11-16), pages 6887-6893, XP026722082, ISSN: 0264-410X, DOI: 10.1016/J.Vaccine.2009.09.008) report a randomized placebo-controlled trial of a reformulated, bivalent, killed, whole-cell, oral cholera vaccine conducted among adults and children in Kolkata, India.

[0014] Sanchez and Holmgren (Current Opinion in Immunology, vol. 17, no. 4, 1 August 2005 (2005-08-01), pages 388-398, XP027787152, ISSN: 0952-7915) review virulence factors, pathogenesis and vaccine protection in cholera and ETEC diarrhea.

[0015] With this in mind the inventors have set out to engineer a single vaccine strain of *V. cholerae* that would effectively replace the three currently used strains.

[0016] Thus, it is an object of the invention to provide an efficient vaccine against cholera and/or ETEC diarrhea, with simplified formulation and with lower production costs and that also ideally produces protective immunity after single administration.

15 **Summary of the invention**

[0017] The present invention describes the construction, method of manufacture, formulation and medical-preventive use of a novel cholera and/or ETEC vaccine. Not all aspects disclosed herein are part of the presently claimed invention. The claimed subject matter is limited to that specified in the appended claims.

20 [0018] Throughout this text, in line with established scientific practice, the designation "*wbeT*" (in italics) denotes the gene, whereas the designation "*WbeT*" (in italics) denotes a protein coded for by a *wbeT* gene.

[0019] In a **first aspect**, a vaccine comprising a *Vibrio cholerae* O1 cell, characterized in that said cell comprises O1 antigens of both Ogawa and Inaba serotypes is provided, wherein the vaccine comprises multiple *Vibrio cholerae* O1 cells that comprise O1 antigens of both Ogawa and Inaba serotypes, and wherein on average, 10-90 % of the O1 antigens of said cells are of the Ogawa-serotype.

[0020] More preferably, 10-70 % of the O1 antigen expressed by the cells is of the Ogawa-serotype. Yet more preferably, 10-50 % of the O1 antigen expressed by the cells is of the Ogawa-serotype. Still more preferably, 10-40 % of the O1 antigen expressed by the cells is of the Ogawa-serotype. Most preferably, 10-30 % of the O1 antigen expressed by the cells is of the Ogawa-serotype.

30 [0021] The cell of the vaccine may further comprise one or more ETEC colonization factor (CF) protein(s), such as CFA/I, CS2 or CS5, wherein said CF protein(s) is/are expressed either as single, double or hybrid fimbriae.

[0022] Preferably, said vaccine does not contain any further immunologically active whole cells in addition to *Vibrio cholerae* O1 cells that comprise O1 antigens of both Ogawa and Inaba serotypes.

[0023] Preferably, the vaccine is for oral administration. Preferably, the cell in the vaccine is formalin-inactivated.

35 [0024] Preferably, the cell is a genetically modified cell, preferably a genetically modified cell according the seventh or eighth aspects of the invention (see below).

[0025] In a **second aspect** a vaccine according to the first aspect is provided, for use in preventive immunization, preferably for use in preventive immunization against cholera and/or enterotoxigenic *Escherichia coli*-infection (ETEC).

40 [0026] In a **third aspect** (not part of the claimed invention), a method for inducing preventive immunity is described, comprising administering a vaccine according to the first aspect to a subject to be immunized. Preferably, the preventive immunity is against cholera and/or enterotoxigenic *Escherichia coli*-infection (ETEC). Also preferably, the administration is done orally.

45 [0027] In a **fourth aspect** (not part of the claimed invention as such), a DNA-construct is disclosed, comprising DNA coding for a *WbeT*-protein having at least 70 % sequence identity to SEQ ID NO: 6 (more preferably at least 80 % identity, even more preferably at least 90 % identity, yet more preferably at least 95 % identity and most preferably at least 99 % identity) operatively coupled to a promoter suitable for inducing protein expression in a *Vibrio cholerae* O1 host cell, characterized in that the coded *WbeT*-protein comprises sequence modifications in relation to SEQ ID NO: 6 that reduce the enzymatic activity of the coded protein relative to the enzymatic activity of a protein with a sequence identical to SEQ ID NO: 6.

50 [0028] Preferably, the sequence modifications comprise a substitution of the serine residue in position 158 of SEQ ID NO: 6, more preferably a substitution of the serine residue in position 158 of SEQ ID NO: 6 to glycine, proline, threonine, phenylalanine or tryptophan.

55 [0029] In a **fifth aspect** (not part of the claimed invention as such), a DNA-construct is disclosed, comprising DNA coding for a *WbeT*-protein having at least 70 % sequence identity to SEQ ID NO: 6 (more preferably at least 80 % identity, even more preferably at least 90 % identity, yet more preferably at least 95 % identity and most preferably at least 99 % identity) operatively coupled to a promoter suitable for inducing protein expression in a *Vibrio cholerae* O1 host cell, characterized in that the promoter is suitable for inducing the expression of the coded *WbeT*-protein in a *Vibrio cholerae* O1 host cell initially of Inaba phenotype (i.e. host cell which is Inaba prior to transformation by the DNA construct)

to such level of transgenic *WbeT* protein expression as to allow simultaneous expression of both Inaba and Ogawa antigens by the host cell.

[0030] Preferably, the promoter of the above aspects is an inducible promoter, such as a *tac* or *lac* promoter.

[0031] Preferably, the DNA construct of the above aspects is a plasmid vector capable of replication in a host cell or a vector capable of chromosomal integration in a host cell.

[0032] Preferably, the DNA-construct according to the above aspects further comprises a selectable marker, more preferably a positive selectable marker such as antibiotic resistance gene or a metabolic selectable marker.

[0033] In a **sixth aspect** (not part of the claimed invention as such), a DNA construct for homologous recombination in a *Vibrio Cholerae* O1 host is described, characterized in that the construct is adapted to modifying the endogenous *wbeT* gene of the host by means of homologous recombination. Preferably, the DNA-construct according to the sixth aspect further comprises a selectable marker, more preferably a positive selectable marker such as antibiotic resistance gene or a metabolic selectable marker.

[0034] In a **seventh aspect** (not part of the claimed invention as such), a *Vibrio cholerae* O1 cell simultaneously expressing both Inaba and Ogawa antigens is disclosed, characterized in that

a. the endogenous *wbeT*-gene of the host cell or the protein coded thereof is inactive;

b. the cell comprises a recombinant DNA-construct inducing expression of *WbeT* enzyme activity; and wherein

c. the level of transgenic *WbeT* enzyme activity is such that the cell simultaneously expresses Inaba and Ogawa antigens.

[0035] Preferably, the recombinant DNA-construct of the above aspects is a DNA construct according to the fourth to fifth aspects.

[0036] In an **eighth aspect**, a *Vibrio cholerae* O1 cell simultaneously expressing both Inaba and Ogawa antigens is provided, characterized in that

a. the cell comprises an endogenous *wbeT*-gene; and

b. the cell comprises a recombinant DNA-construct capable of modulating the expression level of endogenous *wbeT* gene or the enzymatic activity of the product thereof; and wherein

c. the modulated level of *WbeT* enzyme activity is such that the cell simultaneously expresses Inaba and Ogawa antigens, wherein 10-90 % of the O1 antigen expressed by the cell is of the Ogawa-serotype.

[0037] Preferably, the recombinant DNA-construct of the above aspect is a DNA construct according to the sixth aspect.

[0038] Preferably, 10-90 % of the O1 antigen expressed by the cell of the above aspects is of the Ogawa-serotype. More preferably, 10-70 % of the O1 antigen expressed by the cell of the above aspects is of the Ogawa-serotype. Yet more preferably, 10-50 % of the O1 antigen expressed by the cell of the above aspects is of the Ogawa-serotype. Still more preferably, 10-40 % of the O1 antigen expressed by the cell of the above aspects is of the Ogawa-serotype. Most preferably, 10-30 % of the O1 antigen expressed by the cell of the above aspects is of the Ogawa-serotype.

[0039] Preferably, the cell of the above aspects further expresses one or more ETEC colonization factor (CF) protein(s), such as CFA/I, CS2 or CS5, wherein said CF protein(s) is/are expressed either as single, double or hybrid fimbriae.

[0040] In a **ninth aspect**, a method for manufacturing a vaccine is provided, comprising the steps of:

providing a *Vibrio cholerae* O1 cell comprising O1 antigens of both Ogawa and Inaba serotypes; and

killing said cell.

[0041] Preferably, the killing is performed by formalin-treatment or by heat-treatment.

[0042] Preferably, the cell is a cell according to the seventh or the eighth aspects.

[0043] In a **tenth aspect** (not part of the claimed invention), the present disclosure also describes a kit for use in vaccination formulated as a one-unit composition, whereby the composition is presented in one part of the kit and instructions for use in another part.

Detailed description of the invention**Vaccine comprising cells with O1 antigens of both Ogawa and Inaba serotypes**

5 [0044] Although *V. cholerae* of serogroup 0139 can also cause cholera, >98% of all cholera worldwide is caused by *V. cholerae* O1. The O1 serogroup has two subtypes/serotypes - Ogawa and Inaba. Serotype switching from the Ogawa to the Inaba subtype occurs at a relatively high frequency whereas the reciprocal conversion is rare. The basis of the serotype switch is a mutation in the LPS synthetic pathway that leads to change in the structure of the O1 antigen. An effective vaccine need to include both Ogawa and Inaba strains in its composition since their LPS are serologically distinct with both shared and distinct epitopes contributing to protection.

10 [0045] A vaccine comprising *Vibrio cholerae* O1 cells expressing O1 antigens of both Ogawa and Inaba serotypes is disclosed, and has the advantage of simplifying production because it obviates the need to use distinct cells for each of the phenotypes in the manufacture of the vaccine. By enabling the use of single type of cell, the manufacture of the vaccine is also simplified, since only one type of inactivation treatment is needed.

15 [0046] Immunization with vaccine of the invention based on single *V. cholerae* strain that express different amounts of both Ogawa and Inaba serotypes gives rise to cross-reactive as well as type-specific antibodies to both Ogawa and Inaba antigens (see example 1).

20 [0047] On average, 10-90 % of the O1 antigens of the cells are of the Ogawa-serotype. The Inaba-antigen is preferably present in a higher amount (meaning more than 50 %) than the Ogawa-antigen, as the Inaba-antigen can elicit a certain level of cross-serotype protection against Ogawa, whereas Ogawa-antigen can only elicit protection against itself.

25 [0048] The efficacy of the vaccine against ETEC may further be improved by incorporation of the feature that the cells further express one or more ETEC colonization factor (CF) protein(s), such as CFA/I, CS2 or CS5, wherein said CF protein(s) is/are expressed either as single, double or hybrid fimbriae (see below for details).

30 [0049] It is preferable, that the above vaccine does not contain any further immunologically active whole cells in addition to *Vibrio cholerae* O1 cells that comprise O1 antigens of both Ogawa and Inaba serotypes. However, the above vaccine may further comprise recombinant CTB in a manner similar to Dukoral™.

35 [0050] The vaccine is preferably for oral administration, but may also be administered by injection. Preferably, the vaccine comprises *Vibrio cholerae* O1 cells that comprise O1 antigens of both Ogawa and Inaba serotypes that are formalin-inactivated.

40 [0051] Preferably, the vaccine comprises genetically modified *Vibrio cholerae* O1 cells, preferably such genetically modified *Vibrio cholerae* O1 cells as described below.

45 [0052] The vaccine of the invention may be a vaccine composition comprising one or more pharmaceutically acceptable, excipients, carriers, diluents and adjuvants.

50 [0053] The formulation of vaccine compositions according to the invention is well known to persons skilled in the art. Suitable pharmaceutically acceptable carriers and/or diluents include any and all conventional solvents, dispersion media, fillers, solid carriers, aqueous solutions, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art, and is described, by way of example, in Remington's Pharmaceutical Sciences, 18th Edition, Mack Publishing Company, Pennsylvania, USA. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the pharmaceutical compositions of the present invention is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

55 [0054] The vaccine composition for oral use may preferably comprise 10^8 - 10^{14} cells/ml, more preferably 10^{10} - 10^{12} cell per ml, and most preferably about 10^{11} cells/ml.

60 [0055] The vaccine composition for oral use may be formulated as a foodstuff, beverage or a feed supplement (when for use in immunizing animals).

65 [0056] The vaccine composition may comprise an adjuvant known in the art, or may lack any adjuvants.

Use of the vaccine

50 [0057] The present invention discloses use of the above vaccine in preventive immunization, preferably against cholera and/or enterotoxic *Escherichia coli*-infection (ETEC).

55 [0058] Preferably, the vaccine is administered orally or sublingually.

[0059] The administration may also be done by injection.

[0060] The vaccine is preferably used to immunize humans and other mammals, such as pets (cats, dogs and the like) or farm animals (such as cows, horses, sheep, goats, pigs and the like).

[0061] Preferably, the vaccine is administered orally at 10^8 - 10^{14} cells per dose, more preferably 10^{10} - 10^{12} cells per dose, and most preferably about 10^{11} cells per dose.

[0062] The immunization protocol may consist of a single administration or may comprise two or more administrations.

In a preferred embodiment, the initial immunization protocol to induce protective immunity comprises a first administration and a second administration, separated in time by at least 7 days but by no more than about 2 months. After the initial immunization protocol, protective immunity may be maintained as long as desired by booster administrations occurring with less than 3 year intervals, preferably less than 2 year intervals. It is may be preferable that a booster administration does not take place before at least 1 year has elapsed from the first administration.

Method of manufacture for a vaccine

[0063] A method for manufacturing a vaccine is disclosed, comprising the steps of:

- 10 a. providing a *Vibrio cholerae* O1 cell comprising O1 antigens of both Ogawa and Inaba serotypes; and
- b. killing said cell, such as by formalin treatment or by heat treatment.

[0064] Preferably, the killing is performed by formalin-treatment. Preferably the cell is a genetically modified cell, preferably such as described below.

[0065] Besides having the advantage enabling the use of a single inactivation (killing) method, the vaccine may be manufactured using standard protocols known e.g. from the manufacture of Dukoral™.

Genetically modified cells useful for vaccine manufacture and DNA constructs for obtaining such cells

[0066] The *Vibrio cholerae* cells comprising O1 antigens of both Ogawa and Inaba serotypes comprised in the vaccine could in principle be obtained from a naturally occurring strain having a Hikojima phenotype. However, to the best knowledge of the inventors such strains are very rare and no such strains are presently available to the public. In the literature, such natural strains have also been described as unstable, which renders them less promising for industrial production of vaccines.

[0067] Thus, the inventors have derived *V. cholerae* cells that express O1 antigens of both Ogawa and Inaba serotypes by way of genetic engineering and have obtained novel strains with stable Hikojima phenotype. Cells derived in this manner also have the advantage that any desired strain (such as a known and well-characterized vaccine strain) can be used as a starting point, substantially simplifying the production and streamlining the experiments needed for GMP production and regulatory approval.

[0068] The inventors demonstrate herein that the key parameter for obtaining the desired Hikojima phenotype is to obtain a suitable level of *WbeT* enzyme activity. By suitable in this context is meant that the *WbeT* enzyme activity level of the cell is not so low that the cells have an essentially pure Inaba phenotype and not so high that the cells have an essentially pure Ogawa phenotype.

[0069] In the context of the present invention, 10-90% of the O1 antigens on the cells are of the Ogawa type (with remainder consequently of the Inaba type). More preferably, 10-80% of the O1 antigens on the cells are of the Ogawa type, yet more preferably 10-50%, still more preferably 10-40% and most preferably 20-30%.

[0070] As shown in the Examples below, a suitable Hikojima phenotype as outlined above is obtainable by several distinct strategies utilizing recombinant DNA technology:

- 45 a) A mutant *WbeT* protein having low enzymatic activity may be expressed at high levels in a host having Inaba phenotype;
- b) A *WbeT* protein having high enzymatic activity may be expressed at low levels in an Inaba host; or
- c) The endogenous *WbeT* gene of an Ogawa host may be mutated e.g. by means of homologous recombination to render the resulting protein to have suitably reduced activity.
- 50 d) The endogenous *WbeT* gene of an Inaba host may be replaced or modified e.g. by means of homologous recombination to render the protein produced by the gene to have suitably increased activity.

[0071] The present disclosure discloses cells obtained by each of the above strategies, as well as DNA-constructs suitable for obtaining cells by each of the above strategies. However, as specified in the appended claims, only certain embodiments of cells are part of the presently claimed invention as such.

[0072] From the teachings herein it is apparent to the skilled person that achieving the desired level of *WbeT* expression by the strategies outlined above can be realized in many different ways. For instance, levels of expression of a *WbeT* transgene may be modulated by using an inducible promoter (such as *cat*, *lac* or *tac*) whereby the level of expression

may be modulated during culture of the host cells by adjusting the level of the inducer that the host cells are exposed to.

[0073] Alternatively, several weak and strong constitutional promoters are also known and may be used in conjunction with a suitably modified *WbeT* protein. A weak promoter can be used to constitutionally induce a very low level of expression of a highly active (such as wild-type; SEQ ID NO: 6) *WbeT* protein. Conversely, a strong constitutional promoter can be used to induce a high level of expression of a *WbeT* protein having low activity (such as a mutated *WbeT* protein, preferably such as described below).

[0074] Both plasmids and chromosomally integrated *wbeT* transgenes may be used in the cells to achieve the desired phenotype.

[0075] Many different mutations of the *WbeT*-protein can potentially result in a suitably active protein, and such mutated variants can readily be obtained by the skilled person using methods well known in the arts by mere routine experimentation, on the basis of the teachings herein. Whether a cell of the desired phenotype is obtained by expressing the mutated *WbeT* protein or not can readily be analyzed by the skilled person e.g. using the methods disclosed in Example 5. The inventors have identified serine 158 in the *WbeT* protein (SEQ ID NO: 6) as a suitable residue to modulate activity. Thus, the mutations preferably comprise a substitution on serine 158, more preferably substitution of serine 158 to glycine, proline, valine, leucine, alanine, threonine, methionine, tryptophan, arginine or phenylalanine. Most preferably, the serine 158 is substituted by glycine, proline, threonine, phenylalanine or tryptophan.

ETEC colonization factor (CF) protein(s)

[0076] As is evident from the above, the vaccine of the invention (or rather the cells on which the vaccine is based upon) may also comprise other enhanced features besides the combined expression of O1 antigens of both Inaba and Ogawa serotypes. In particular, the cells may express one or more ETEC colonization factor (CF) proteins, such as CFA/I, CS2 or CS5, wherein said CF protein(s) is/are expressed either as single, double, or hybrid fimbriae, as demonstrated in Example 7. Inclusion of such CF proteins in the cells of the vaccine is especially useful for inducing protective immunity against ETEC.

[0077] The expression "comprising" as used herein should be understood to include, but not be limited to, the stated items.

[0078] The invention is further illustrated by the following examples, which are to be seen as non-limiting.

Example 1. Preparation and testing of a vaccine comprising *Vibrio cholerae* cells comprising O1 antigens of both Ogawa and Inaba serotypes

[0079] It was tested whether a vaccine comprising *V. cholerae* bacteria of strain JS1569 (Inaba) that had been genetically modified to express O1 antigens of both Ogawa and Inaba serotypes would give rise to an antibody response with a different proportion of antibodies reacting with Ogawa and Inaba LPS in ELISA as compared to the antibody response after immunization with the parent JS1569 Inaba strain.

[0080] The bacteria from example 2 (see below) were formalin-killed and used for immunization. Formalin-killing of bacteria and oral immunizations and assay methods were performed as previously described (Nygren E, Li BL, Holmgren J, Attridge SR. Infect Immun. 2009 Aug;77(8):3475-84). Briefly, Balb/c mice were immunized in 3 rounds at 2-week intervals with two daily doses of 3×10^8 formalin-killed cells (together with an adjuvant for the *WbeT* strain), and one week after the last immunization the mice were sacrificed and serum collected and tested for combined IgG/IgM antibody titers in ELISA plates coated with either Inaba or Ogawa LPS.

[0081] The results are presented in the table below and show that in contrast to the parent JS1569 Inaba strain which gave rise to an antibody response with a slightly higher anti-Inaba than anti-Ogawa titer, the JS1569/wild type *WbeS158S* vaccine gave rise to an antibody response with a much higher anti-Ogawa than anti-Inaba titer, although it still gave rise also to modest formation of specific anti-Inaba antibodies:

| Immune serum | Inaba/Ogawa titers (Ratio) |
|-------------------------------------|----------------------------|
| To JS1569 | 10290/5060 (2:1) |
| Same absorbed with Ogawa | 2940/160 (18:1) |
| To JS1569/wild type <i>WbeS158S</i> | 36000/365000 (1:10) |
| Same absorbed with Inaba | 1600/49000 (1:30) |
| Same absorbed with Ogawa | 1700/2500 (1:1.5) |

[0082] These findings were confirmed when immunizations were given subcutaneously without adjuvant. In marked

difference to immunization with the parent JS1569 Inaba strain and more similar to immunization with the Ogawa A457 reference strain, immunization with JS1569 WbeT gave rise to immune serum with a strong proportion of Ogawa specific antibodies, as shown in the table below.

| 5 | Immune serum | Inaba titer | Ogawa titer | Ratio Inaba/Ogawa |
|----|----------------------------|-------------|-------------|-------------------|
| 10 | JS1569 Inaba | 480 | 260 | 1.8:1 |
| | JS1569 abs with Ogawa A457 | 180 | 60 | 3:1 |
| | JS1569 abs with 1569 WbeT | 240 | 60 | 4:1 |
| | A457 Ogawa | 660 | 1940 | 1:1.9 |
| | A457 abs with Inaba 1569 | 240 | 1020 | 1:4.3 |
| | A457 abs with JS1569 WbeT | 180 | 40 | 1.8:1 |
| | JS1569 WbeT | 420 | 1580 | 1:3.8 |
| | WbeT abs with Ogawa A457 | 150 | 60 | 2.5:1 |
| | WbeT abs with Inaba JS1569 | 180 | 920 | 1:5.1 |
| | WbeT abs with JS1569 WbeT | 120 | 100 | 1.2:1 |

Example 2: Genetically modified *Vibrio cholerae* cells expressing O1 antigens of both Ogawa and Inaba serotypes by expression of mutated *WbeT*-protein: Plasmid-based expression of mutated *WbeT*-protein

25 [0083] In a single entry in GenBank of the *wbeT* gene from a Hikojima strain there is a mutation converting a serine to proline at position 158 of the protein (S158P) although the same mutation has been described in a strain identified as being of the Inaba serotype (GenBank accession numbers FJ619106 and DQ401028 respectively). Having amplified the wild-type *wbeT* gene from the O1 El Tor Ogawa strain VX44945 with primers *wbeT1* EcoRI (SEQ ID NO: 1 5'-CCCGGTCTCGAATTCTGCATCTGCAAGTTGATTCTGTATG-3') and *wbeT2* HindIII (SEQ ID NO: 2 5'-CCCGGTCTCAAGCTTATAGTGAACCTCTTCGGAAATGTCTG-3'), it was digested with Eco31I and cloned into an expression vector derived from pAF1 () in which the cloned gene was placed under the control of the powerful synthetic tac promoter that had been digested with EcoRI and HindIII. The sequence of the gene was confirmed by DNA sequencing of the plasmid with primers *wbe1* (SEQ ID NO: 3 5'-CTGCATCTGCAAGTTGATTCTGTATG-3') and *wbe2* (SEQ ID NO: 4 5'-ATAGTGAACCTCTTCGGAAATGTCTG-3').

30 [0084] The DNA sequence of the wild-type *wbeT* gene is shown in SEQ ID NO: 5 whereas the wild-type protein is shown in SEQ ID NO: 6.

35 [0085] The full sequence of the plasmid (pML-*wbeT* tac) expressing the wild-type *wbeT* is shown in SEQ ID NO: 7.

40 [0086] In order to construct the mutant library of *wbeT* carrying mutations at amino acid position 158 of the gene product, oligonucleotides *wbeT m3* (SEQ ID NO: 8 5'-GCGCGCCAGAACCTGGCTATTTTAACC-3') and *wbeT m1* (SEQ ID NO: 9 5'-GGGGGTTCGAAGTTATGAGTTGATAATAGGGTGNNBTCATTATTTCAAAAAAATACA GACATAGCAGATAAGGTTAAAAATAGCCAAGTTCTGGCGCGC-3') were synthesized. The two oligonucleotides were mixed in equimolar quantities and allowed to anneal at room temperature overnight. Full length double-stranded DNA was made by extension of the short *wbeT m3* primer using T4 DNA polymerase in the presence of excess deoxyribonucleotide triphosphates. The resulting fragment was digested with Bsp119I and Van91I and ligated into pML-*wbeT* tac (SEQ ID NO: 7) digested with the same enzymes. The ligated DNA was used to transform commercially obtained electro-competent *E. coli* strain DH12S (Invitrogen). After incubation without antibiotic selection a small aliquot of the cells were spread onto a selective LB agar plate supplemented with ampicillin (100 µg/ml). The rest of the cells were diluted to 25 ml with fresh LB broth. Ampicillin was added to a final concentration of 100 µg/ml and the culture was grown overnight at 37° C in order to obtain a clone library.

45 [0087] Aliquots of the resulting culture were supplemented with glycerol to a final concentration of 17% and stored at -70° C. Other aliquots were used to prepare plasmid DNA.

50 [0088] The colonies obtained on the LB agar plate were picked onto a fresh plate and colonies were cultured to prepare plasmid DNA. The plasmids were sequenced in order to determine whether the *wbeT* genes carried mutations. Mutants of *wbeT* obtained from the library are the following: S158G, S158P, S158V, S158I, S158L, S158A, S158T, S158M, S158W, S158R, S158C and S158F. Additionally the wild-type gene and a gene with the stop signal TGA at position 158 were isolated.

55 [0089] The different plasmids were isolated and used to transform the O1 classical Inaba strain JS1569. This strain

has a mutant *wbeT* gene with the glycine (GGA) at position 219 of the protein being changed to a stop codon (TGA) resulting in a truncated and inactive product (SEQ ID NO: 10 and SEQ ID NO: 11).

[0090] There are other polymorphisms that do not appear to have any significance.

[0091] The different strains generated by the introduction of the different recombinant plasmids expressed different levels of the Ogawa antigen when grown under inducing conditions (in the presence of 1mM IPTG). The phenotype was assessed on the basis of agglutination assays and in some cases using inhibition ELISA (see Example 5 for description of materials and methods). The wild type gene gave rise to almost total serotype switching whereas others (such as S158P and S158G) gave slight but detectable agglutination with Ogawa-specific antiserum as well as agglutination with an Inaba-specific antiserum (and therefore conferred a Hikojima serotype). Some mutants had no detectable activity with Ogawa-specific antiserum (S158I and S158C) and yet others gave intermediate agglutination (S158T, S158F and S158W).

[0092] The results demonstrate unambiguously that mutations and specifically mutations at position 158 of the *wbeT* gene product result in proteins with altered enzyme activity. At present there is no reliable assay to directly quantitatively determine the levels of enzyme activity of these mutants compared to the wild-type, but the relevant end result can readily be evaluated as in Example 5. In summary, all except S158C and S158I were able to complement the Inaba phenotype of the host strain to some extent.

Example 3: Genetically modified *Vibrio cholerae* cells expressing O1 antigens of both Ogawa and Inaba serotypes by expression of mutated *WbeT*-protein: Chromosomal insertion of mutant *wbeT*

[0093] The truncated chromosomal *wbeT* gene in the strain JS1569 was substituted with the mutant genes generated in example 2. The relevant mutated genes were amplified with primers *wbeT1* BlgII (SEQ ID NO: 1) and *wbeT2* BgIII (SEQ ID NO: 2). Amplified fragments were digested with BgIII and ligated into the suicide vector pMT-SUICIDE (SEQ ID NO: 12) which had been digested with BamHI. This is a small R6K-based suicide vector constructed in this laboratory by M. Lebens which carries the chloramphenicol resistance gene and the origin of transfer (oriT) from the broad host-range plasmid RP4 that allows the plasmid to be conjugally transferred to *V. cholerae* strain with the aid of a helper plasmid (pNJ5000; Grinter NJ, Gene. 1983 JanFeb;21(1-2):133-43).

[0094] In the clones that were generated, the *wbeT* genes (S158G and the wild-type) were both inserted with the cloned genes in the opposite orientation to the *cat* gene. The sequence of such vector is exemplified by SEQ ID NO: 13 (carrying wild-type *wbeT* gene; the construct for S158G is identical save for the nucleotides coding for *WbeT* residue 158).

[0095] The resulting plasmids were mated into strain JS1569 and selected on the basis of chloramphenicol and rifampicin resistance. Since the plasmid has no counter-selection for loss of the plasmid, its insertion into the chromosome by homologous recombination results in tandem copies of the *wbeT* gene separated by the plasmid. Depending upon where the recombination occurred, the clones had different phenotypes (see Example 5).

[0096] The strain that had received the wild-type gene (1342) had a clear Hikojima phenotype. Inhibition ELISA showed that it expressed only 15% of the Ogawa LPS present on the surface of the strain that received the S158G mutant. The latter strain (1356) was in effect an Ogawa strain that agglutinated strongly with Ogawa-specific antiserum, but not at all with the Inaba-specific antiserum.

[0097] The strains were however very stable; they retained their LPS serotypes and remained chloramphenicol resistant even in the absence of selection, indicating that the plasmid was not readily lost.

[0098] PCR and sequencing using the *wbe1* and 2 primers (SEQ ID NO: 3 and 4, respectively) showed that there were two genes in the strains that varied at sites of variation between the gene present in the host and that which was introduced. Amplification and sequencing with primers *wbeTfor* 87> (SEQ ID NO 14: 5'-CGGTGCAAACGTTGGAACCTT-TCTG-3') and *wbeT rev* 51< (SEQ ID NO 15: 5'-GGAAAACAATGCCATCCAAATTGCG-3') that only allow amplification if there are tandem copies of the *wbeT* gene successfully amplified the 3' end of the proximal gene (from amino acid 87) and the 5' end of the distal gene up to amino acid 51 and the plasmid in between. Sequencing using the *wbeTfor* 87> primer showed that in the strain 1342 the *wbeT* gene adjacent to the native promoter was the truncated host gene. The distal gene had the wild-type sequence but no promoter. This arrangement led to the Hikojima phenotype since the wild-type gene was being expressed at extremely low levels from a cryptic promoter.

[0099] In the Ogawa strain 1356 the arrangement was different. The recombination had resulted in the native *wbeT* gene being expressed from the native promoter and the mutant S158G gene being placed distally and therefore having no promoter recognizable at all. Both copies of the gene appear to have lost the stop codon at position 219, but this mutation had no apparent influence on the phenotype.

Example 4: Genetically modified *Vibrio cholerae* cells expressing O1 antigens of both Ogawa and Inaba serotypes by expressing low levels of native *WbeT*-protein

[0100] In conjunction with the experiments on mutant *wbeT* described in Example 2 it was noted that a control plasmid carrying the wild-type *wbeT* was able to partially complement the mutant gene in strain JS1569 even when it was not induced. This resulted in a Hikojima serotype even in the presence of the wild type gene demonstrating that the phenotype can also be achieved by limiting the levels of expression; in this case by keeping the *tac* promoter repressed and allowing only breakthrough expression that occurs in the absence of inducer.

[0101] In Example 3, the clone 1342 had a chromosomally integrated wild-type gene expressed from a cryptic promoter, which resulted in a Hikojima serotype. These results confirm that the expression of the wild-type gene at very low levels can result in the Hikojima serotype.

Example 5: Characterization of the phenotype of genetically modified *Vibrio cholerae* cells

[0102] *V. cholerae* bacteria of strain JS1569 (Inaba) that had been modified to contain plasmids encoding for either the wild-type *WbeT* methylase protein (strain JS1569/S158S) or a mutated *wbeT* gene encoding for *WbeT* protein with a mutation in position 158 from S to G (JS1569/S158G) or from S to A (JS1569/S158A) were grown on LB agar plates, and single colonies were tested for agglutination by antibodies specific for Inaba and Ogawa O antigens, respectively.

[0103] These antibodies were obtained after first immunizing rabbits with purified Ogawa and Inaba LPS, respectively and then extensively absorbing the sera with formalin killed bacteria of the heterologous serotype in order to remove cross-reactive antibodies. After absorption, Ogawa-specific antiserum gave strong agglutination of reference *V. cholerae* cells of the Ogawa serotype but could not agglutinate Inaba cells and vice versa for the Inaba specific serum.

[0104] Agglutination tests were performed by standard method. Briefly, a single colony from a fresh plate of the tested strain was suspended in 50-100 μ l physiological saline buffer, and 10 μ l of the suspension was placed on a microscope slide. Then 10 μ l of appropriately diluted specific antiserum was added and mixed with the cells by tilting the slide backwards and forwards for up to 5 minutes until agglutination was clearly visible. Each assay was compared with negative and positive controls consisting of cells from reference Inaba and Ogawa strains.

[0105] Additionally, a control for spontaneous agglutination in which serum was replaced by buffer was performed for each strain tested. Results are shown in the table below and show that the JS1569/S158S strain containing plasmid encoding for wild-type *WbeT* protein had completely switched in serotype from Inaba to Ogawa, the JS1569/S158G strain with a *WbeT* 158S to G mutation expressed strong Ogawa but also detectable Inaba reactivity, and JS1569/S158A with a *WbeT* 158S to A mutation had only marginal Ogawa reactivity.

| Strain | <i>WbeT</i> encoding Plasmid | Agglutination with anti-Ogawa antibody | Agglutination with anti-Inaba antibody |
|----------------|------------------------------|--|--|
| JS1569 Inaba | None | - | +++ |
| Cairo 50 Ogawa | None | +++ | - |
| A457 Ogawa | None | +++ | - |
| JS1569/S158S | <i>WbeT</i> S158S wild-type | +++ | (+) |
| JS1569/S158G | <i>WbeT</i> S158G | ++ | ++ |
| JS1569/S158A | <i>WbeT</i> S158A | (+) | +++ |

[0106] These results were confirmed when formalin-killed bacteria of the same strains were tested for agglutination with the same sera. They were also confirmed and extended when the formalin-killed bacteria were tested for their quantitative expression of Inaba and Ogawa antigens on the bacterial surface using an ELISA-inhibition method. The method was performed as follows: High binding Greiner Bio-one plates were coated with Ogawa LPS by incubating overnight with 100 μ l per well of a solution of 5 μ g/ml purified Ogawa LPS in PBS. Starting with 200 microliter of OD₆₀₀ 1.00 formalin killed bacteria; seven serial five-fold dilutions were made (down to 1:15625) in PBS with an eighth blank tube containing no cells. 150 microliter of each dilution was mixed together with an equal amount of appropriately diluted anti-Ogawa serum in PBS, 0.2% BSA. The samples were incubated at room temperature for 1 hour without shaking. The coated plates were washed twice with PBS and blocked with 200 μ l/well of PBS, 0.1% BSA for 30 minutes at 37° C. The cells were removed from the suspensions by centrifugation for 5 minutes at 20,000 x g and 100 microliter of the supernatants were added to the blocked the plate(s). A blank containing PBS, 0.1% BSA with neither cells nor anti-

Ogawa serum was included in all plates and all samples were run in duplicate. Plates were incubated for 1 hour at room temperature and then washed three times with PBS, 0.05% Tween 20. 100 microliter of appropriately diluted goat anti-rabbit IgG in PBS, 0.1% BSA, 0.05% Tween 20 was added to each well and incubated for 1 hour at room temperature. The plate was washed three times with PBS, 0.05% Tween 20 before adding the substrate solution of 0.1% orthophe-nylenediamine (OPD) and 0.012% H₂O₂ in 0.1M citrate pH4.5. The absorbance was read at 490nm after 10min.

[0107] The bacterial dilution resulting in 50% inhibition was extrapolated and also the percentage inhibition of the absorbance at bacterial dilution of 1:25 with the results shown in the table below. The results show that the JS1569/S158S strain containing a *wbeT* gene encoding for wild-type *WbeT* could efficiently inhibit the specific anti-Ogawa serum, whereas the strains containing *wbeT* genes with single mutations could also inhibit the anti-Ogawa serum with intermediate activity (JS1569/S158G) or just detectably (JS1569/S158A):

| Strain | Dilution for 50% inhibition | % inhibition at dilution 1:25 |
|--------------|-----------------------------|-------------------------------|
| JS1569 Inaba | << 1:1 | 0 |
| A457 Ogawa | 1:60 | 65 |
| JS1569/S158G | 1:70 | 70 |
| JS1569/S158S | 1:1 | 20 |

[0108] Based on the plasmids encoding for *WbeT* and S158G, respectively, two recombinant derivative strains of JS1569, 1356 (wild-type *WbeT*) and 1342 (S158G *WbeT*), were obtained where the mutated *wbeT* gene had been stably inserted into the chromosome, giving unexpected but stable phenotypes. For an explanation and description of these strains see examples above. These strains and appropriate reference strains were subjected to colony blots for assaying Ogawa expression. Strains containing mutated *wbeT* genes were patched out on an LB agar plate together with Inaba and Ogawa control strains and grown overnight at 37° C. A nitrocellulose membrane moistened with PBS was applied on the plate with the grown out colonies and was left for 15 minutes at room temperature. The membrane was left to dry on a piece of paper for 5 minutes before it was blocked twice for 20 minutes in 10 ml of 1% Bovine serum albumin (BSA) in PBS at room temperature. The blocking liquid was discarded and replaced with an appropriate dilution of the anti-Ogawa serum in 10 ml in PBS containing 0.1% BSA and 0.05% Tween 20. The membrane was incubated at room temperature on a rocking table for 2 hours. The membrane was then washed three times with PBS containing 0.05% Tween20 before adding goat anti-rabbit IgG conjugated to horseradish peroxidase (Jackson ImmunoResearch Laboratories Inc.) in 0.1% BSA in PBS with 0.05%Tween20 and incubating for 2 hours at room temperature with gentle agitation.

[0109] Following a further three washes in PSB, 0.05%Tween 20 and a single wash with PBS alone the membrane was developed for 15 minutes with 0.05% 4-chloro-1-naphthol and 0.015% H₂O₂ in 20 mM Tris -HCl pH 7.5 containing 500 mM NaCl and 16.7% methanol. It was then washed thoroughly with tap water and was left to dry on a piece of paper. A digital photo was taken of the developed membrane and the staining density measured with a computer system.

[0110] Results in the table show that strain 1356 expressed almost as much Ogawa antigen as the Ogawa reference strain whereas strain 1342 expressed substantially lesser amounts of Ogawa antigen. These findings were further confirmed when formalin-killed preparations of the strains were tested for quantitative expression of Ogawa antigen by inhibition-ELISA done as described above as also shown in the table:

| Strain | Dot blot density units/mm ² | Dilution for 50% inhibition |
|--------------|--|-----------------------------|
| A 457 Ogawa | 12700 | 1:60 |
| JS1569 Inaba | 0 | <<1:1 |
| 1356 | 10000 | 1:80 |
| 1342 | 4500 | 1:7 |

Example 6: Genetic modification of the endogenous *wbeT* gene to obtain Hikojima serotype

[0111] Although the strains presented in the above examples are stable and clearly have the desired phenotypes, an alternative manner of obtaining a Hikojima phenotype is to perform true gene substitutions in the endogenous gene. Suitable mutations (as disclosed above) at position 158 of the endogenous, active *wbeT* gene product also result in diminished activity and therefore a Hikojima serotype. This will result in a range of mutants with differing levels of Ogawa expression that can be tested for Hikojima expression as in Example 5 and in immunization experiments for the optimal

immunogenic properties.

[0112] As mentioned, the pMT-SUICIDE vector lacks a suitable counter-selection gene and it has proved difficult isolate derivatives that have lost the plasmid and retain the desired gene and phenotype. For this reason, a new suicide vector was constructed carrying the *sacB* gene from *Bacillus subtilis*, allowing strains that have lost the plasmid to be isolated by selection on plates containing sucrose since expression of the *sacB* gene product in gram-negative bacteria is lethal and only colonies derived from cells that have lost the plasmid due to homologous recombination between the two copies of the *wbeT* gene will survive. The plasmid, pMT Suicide/sacB (SEQ ID NO: 16) is much smaller than comparable plasmids with the same functions and is much easier to use.

[0113] The wild-type and a number of mutant *wbeT* genes have been cloned into the new vector and will be mated into the strain JS1569 as before, as well as to other strains such as a suitable Ogawa strain.

[0114] In order to optimize the chances of obtaining a correct clone the gene arrangement of the partial diploids that resulting from the transconjugation experiments will be analysed before a candidate strain is chosen for taking further for selection of strains that have recombined out the plasmid.

15 Example 7: Expression of hybrid CFA/I+CS2

[0115] It has recently been shown that *E. coli* TOP10 and *V. cholerae* JS1569 strains can express one of the major colonization factor of ETEC, i.e. CFA/I fimbriae, on their surface (Tobias J, Lebens M, Bölin I, Wiklund G, Svennerholm AM. Vaccine. 2008 Feb 6;26(6):743-52). Following electroporation of a CFA/I-containing expression vector into the above strains, the surface expression was detected by dot blot assay.

[0116] It has also recently been shown that *E. coli* TOP10 is capable of expressing a hybrid fimbriae containing the major subunits of both CFA/I and CS2- i.e. an additional major colonization factor of ETEC (Tobias J, Svennerholm AM, Holmgren J, Lebens M. Appl Microbiol Biotechnol. 2010 Jul;87(4))

[0117] Therefore, we examined the feasibility of expression of the same hybrid fimbriae on *V. cholerae* JS1569. The strain was electroporated with the expression vector pJT-CFA/I-CotA as described (Tobias et al. 2008, *supra*; Tobias J, Holmgren J, Hellman M, Nygren E, Lebens M, Svennerholm AM. Vaccine. 2010 Aug 20. [Epub ahead of print]) Fifty colonies (clones) were then streaked on two LB plates supplemented with chloramphenicol (12.5 µg/ml) and IPTG (1 mM), followed by over-night incubation at 37°C. Specific monoclonal antibodies (MAb) 1:6 against CFA/I and MAb 10:3 against CS2 were then applied in colony blot assay (Tobias et al, 2010, *supra*) to examine the surface expression of the hybrid CFA/I-CotA fimbrie on *V. cholerae* JS1569. The blots were then developed and showed positive signal of surface expression of the hybrid CFA/I-CS2 fimbriae on all tested fifty clones of *V. cholerae* JS1569.

[0118] Thus it is possible to combine the expression of antigenic hybrid fimbriae in the same cells that have been engineered to express O1 antigens of both Ogawa and Inaba serotypes.

35 SEQUENCE LISTING

[0119]

<110> Gotovax AB Holmgren, Jan Lebens, Michael

40 <120> Vaccine against cholera and enterotoxigenic *E. coli* (ETEC) diarrhea

<130> P40903745PCT00

45 <150> US 61/272351
<151> 2009-09-16

<160> 16

50 <170> PatentIn version 3.5

<210> 1

<211> 41

<212> DNA

55 <213> *Vibrio cholerae*

<400> 1

cccggtctcg aattcctgca tctgcaagtt gattctgtat g

<210> 2
<211> 40
<212> DNA
<213> Vibrio cholerae

5
<400> 2
cccggtctca agcttatagt gaactctcg gaaatgtctg 40

10
<210> 3
<211> 26
<212> DNA
<213> Vibrio cholerae

15
<400> 3
ctgcatctgc aagttgattc tgtatg 26

20
<210> 4
<211> 25
<212> DNA
<213> Vibrio cholerae

<400> 4
atagtgaact cttcgaaat gtctg 25

25
<210> 5
<211> 953
<212> DNA
<213> Vibrio cholerae

30
<220>
<221> CDS
<222> (67)..(927)

35
<400> 5

40

45

50

55

EP 2 477 649 B1

cctgcattctg caagttgatt ctgtatgtta tttttacgc taatattttt taaaatttag 60
 gtagta atg aaa cat cta ata aaa aac tat gta caa aaa tta att aaa 108
 Met Lys His Leu Ile Lys Asn Tyr Val Gln Lys Leu Ile Lys
 1 5 10
 aca gag ctt gat gct att cag tca aag tct gtt cat gat aat cga aac 156
 Thr Glu Leu Asp Ala Ile Gln Ser Lys Ser Val His Asp Asn Arg Asn
 15 20 25 30
 ttc att tac aat gga gag ttt tta att ctt gaa agc gaa ttt gga tgg 204
 Phe Ile Tyr Asn Gly Glu Phe Leu Ile Leu Glu Ser Glu Phe Gly Trp
 35 40 45
 cat tgt ttt ccc aga gtg cag ttg aac cat gct tta agc tac aaa aac 252
 His Cys Phe Pro Arg Val Gln Leu Asn His Ala Leu Ser Tyr Lys Asn
 15 50 55 60
 cca aac ttt gat tta ggt atg cgt cac tgg att gtt aat cat tgt aag 300
 Pro Asn Phe Asp Leu Gly Met Arg His Trp Ile Val Asn His Cys Lys
 65 70 75
 cat gac acc act tat att gat atc ggt gca aac gtt gga act ttc tgt 348
 His Asp Thr Thr Tyr Ile Asp Ile Gly Ala Asn Val Gly Thr Phe Cys
 80 85 90
 gga atc gct gct cgt cat att aca caa gga aaa att ata gcg ata gaa 396
 Gly Ile Ala Ala Arg His Ile Thr Gln Gly Lys Ile Ile Ala Ile Glu
 25 95 100 105 110
 cca ctc aca gaa atg gaa aat agt att agg atg aat gtt caa tta aat 444
 Pro Leu Thr Glu Met Glu Asn Ser Ile Arg Met Asn Val Gln Leu Asn
 115 120 125
 aat cca ctg gtt gag ttt cat cat ttt ggc tgt gca ata ggt gag aat 492
 Asn Pro Leu Val Glu Phe His His Phe Gly Cys Ala Ile Gly Glu Asn
 130 135 140
 gaa ggg gaa aat att ttc gaa gtt tat gag ttt gat aat agg gtg tca 540
 Glu Gly Glu Asn Ile Phe Glu Val Tyr Glu Phe Asp Asn Arg Val Ser
 35 145 150 155
 tca tta tat ttt caa aaa aat aca gac ata gca gat aag gtt aaa aat 588
 Ser Leu Tyr Phe Gln Lys Asn Thr Asp Ile Ala Asp Lys Val Lys Asn
 160 165 170
 agc caa gtt ctg gtt aga aag tta agt agt tta gat ata tcg cct act 636
 Ser Gln Val Leu Val Arg Lys Leu Ser Ser Leu Asp Ile Ser Pro Thr
 175 180 185 190
 aac tct gta gtt ata aaa att gat gct gaa ggc gca gaa ata gag ata 684
 Asn Ser Val Val Ile Lys Ile Asp Ala Glu Gly Ala Glu Ile Glu Ile
 45 195 200 205
 tta aac cag att tac gaa ttc aca gaa aag cat aat gga att gaa tat 732
 Leu Asn Gln Ile Tyr Glu Phe Thr Glu Lys His Asn Gly Ile Glu Tyr
 210 215 220
 tat att tgc ttt gaa ttt gca atg ggt cat ata cag agg tct aat aga 780
 Tyr Ile Cys Phe Glu Phe Ala Met Gly His Ile Gln Arg Ser Asn Arg
 225 230 235
 act ttt gat gag att ttt aac ata ata aac tca aaa ttc gga agt aag 828
 Thr Phe Asp Glu Ile Phe Asn Ile Ile Asn Ser Lys Phe Gly Ser Lys
 55 240 245 250

EP 2 477 649 B1

gca tat ttt att cat cca tta tca tcc gct gaa cat cct gag ttt aat 876
Ala Tyr Phe Ile His Pro Leu Ser Ser Ala Glu His Pro Glu Phe Asn
255 260 265 270

5 aaa gca acg cag gat att aat ggg aat atc tgt ttt aaa tat gta tca 924
Lys Ala Thr Gln Asp Ile Asn Gly Asn Ile Cys Phe Lys Tyr Val Ser
275 280 285

taa aataatttaa tatattccgt atgtca 953

10 <210> 6
<211> 286
<212> PRT
<213> Vibrio cholerae

15 <400> 6

20

25

30

35

40

45

50

55

Met Lys His Leu Ile Lys Asn Tyr Val Gln Lys Leu Ile Lys Thr Glu
 1 5 10 15

5 Leu Asp Ala Ile Gln Ser Lys Ser Val His Asp Asn Arg Asn Phe Ile
 20 25 30

10 Tyr Asn Gly Glu Phe Leu Ile Leu Glu Ser Glu Phe Gly Trp His Cys
 35 40 45

15 Phe Pro Arg Val Gln Leu Asn His Ala Leu Ser Tyr Lys Asn Pro Asn
 50 55 60

Phe Asp Leu Gly Met Arg His Trp Ile Val Asn His Cys Lys His Asp
 65 70 75 80

20 Thr Thr Tyr Ile Asp Ile Gly Ala Asn Val Gly Thr Phe Cys Gly Ile
 85 90 95

25 Ala Ala Arg His Ile Thr Gln Gly Lys Ile Ile Ala Ile Glu Pro Leu
 100 105 110

30 Thr Glu Met Glu Asn Ser Ile Arg Met Asn Val Gln Leu Asn Asn Pro
 115 120 125

Leu Val Glu Phe His His Phe Gly Cys Ala Ile Gly Glu Asn Glu Gly
 130 135 140

35 Glu Asn Ile Phe Glu Val Tyr Glu Phe Asp Asn Arg Val Ser Ser Leu
 145 150 155 160

40 Tyr Phe Gln Lys Asn Thr Asp Ile Ala Asp Lys Val Lys Asn Ser Gln
 165 170 175

45 Val Leu Val Arg Lys Leu Ser Ser Leu Asp Ile Ser Pro Thr Asn Ser
 180 185 190

50

55

Val Val Ile Lys Ile Asp Ala Glu Gly Ala Glu Ile Glu Ile Leu Asn
195 200 205

5 Gln Ile Tyr Glu Phe Thr Glu Lys His Asn Gly Ile Glu Tyr Tyr Ile
210 215 220

10 Cys Phe Glu Phe Ala Met Gly His Ile Gln Arg Ser Asn Arg Thr Phe
225 230 235 240

15 Asp Glu Ile Phe Asn Ile Ile Asn Ser Lys Phe Gly Ser Lys Ala Tyr
245 250 255

Phe Ile His Pro Leu Ser Ser Ala Glu His Pro Glu Phe Asn Lys Ala
260 265 270

20 Thr Gln Asp Ile Asn Gly Asn Ile Cys Phe Lys Tyr Val Ser
275 280 285

<210> 7

<211> 4858

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic plasmid

30 <400> 7

35

40

45

50

55

| | |
|---|-----|
| aggtggcact tttccccggaa atgtgcgcgg aacccttatt tgtttatttt tctaaataca | 60 |
| ttcaaatatg tatccgctca tgagacaata accctgataa atgcttcaat aatccggatt | 120 |
| 5 gaaaaaggaa gagtatgagt attcaacatt tccgtgtcgc ccttattccc tttttgcgg | 180 |
| cattttgcct tcctgtttt gctcacccag aaacgctggt gaaagtaaaa gatgctgaag | 240 |
| 10 atcagttggg tgcacgagtg ggttacatcg aactggatct caacagcggt aagatcctg | 300 |
| agagtttcg cccccaaagaa cgttttccaa tgatgagcac ttttaaagtt ctgctatgtg | 360 |
| 15 gcgccgtatt atcccgttt gacgccggc aagagcaact cggtcgccgc atacactatt | 420 |
| ctcagaatga cttgggttag tactcaccag tctcagaaaa gcatcttacg gatggcatga | 480 |
| cagtaagaga attatgcagt gctgccataa ccatgagtga taacactgcg gccaaacttac | 540 |
| 20 ttctgacaac gatcggagga ccgaaggagc taaccgctt tttgcacaac atgggggatc | 600 |
| atgtaactcg ccttgatcgt tggaaaccgg agctgaatga agccataccca aacgacgagc | 660 |
| gtgacaccac gatgcctgta gcaatggcaa caacgttgcg caaactatta actggcgaac | 720 |
| 25 tacttactct agcttcccg caacaattaa tagactggat ggaggcggat aaagttgcag | 780 |
| gaccacttct gcgctcgccc cttccggctg gctggttat tgctgataaa tctggagccg | 840 |
| gtgagcgtgg gtctcgccgt atcattgcag cactggggcc agatggtaag ccctcccgta | 900 |

30

35

40

45

50

55

| | | |
|----|--|------|
| | tcgttagttat ctacacgacg gggagtctagg caactatgga tgaacgaaat agacagatcg | 960 |
| | ctgagatagg tgcctcaactg attaagcatt ggttaactgtc agaccaagtt tactcatata | 1020 |
| 5 | tacttttagat tgatTTaaaa ctTCatTTT aattttaaaag gatcttaggtg aagatcctt | 1080 |
| | ttgataatct catgaccaaa atcccttaac gtgagTTTc gttccactca gcgtcagacc | 1140 |
| | ccgttagaaaa gatcaaagga tcttctttag atcCTTTTt tctgcgcgt aatctgtgt | 1200 |
| 10 | tgcaaacaaa aaaaccaccg ctaccagcgg tggTTTgttT gcccgtacaa gagctaccaa | 1260 |
| | ctCTTTTcc gaaggtaact ggcttcagca gagcgcagat accaaatact gtccttctag | 1320 |
| | tgttagccgt aTTaggccac cacttcaaga actctgttagc accgcctaca tacctcgctc | 1380 |
| 15 | tgctaattcct gttaccagtg gctqctgcca gtggcgataa gtcgtgtctt accggggttgg | 1440 |
| | actcaagacg atagttaccg gataaggcgc agcggtcggg ctgaacgggg gttcgtgca | 1500 |
| | cacagccccag cttggagcga acgacctaca ccgaactgag atacctacag cgtgagcatt | 1560 |
| 20 | gagaaaagcgc cacgcttccc gaagggagaa aggcggacag gtatccgt aacggcaggg | 1620 |
| | tcggaacagg acagcgcacg agggagctc cagggggaaa cgcctggat otTTatagtc | 1680 |
| | ctgtcgggtt tcgcccaccc tgacttgagc gtcgattttt gtgtatgtcg tcaggggggc | 1740 |
| 25 | ggagcctatg gaaaatctt cctgcgttat cccctgattc tgtggataac cgtattaccc | 1800 |
| | cctttgagtg agctgacgcc agcaacgcgg ccttttacg gttctggcc ttttgctggc | 1860 |
| | cTTTgctca catgttctt cctgcgttat cccctgattc tgtggataac cgtattaccc | 1920 |
| 30 | cctttgagtg agctgatacc gtcgcccga gccgaacgac cgagcgcagc gagtcagtga | 1980 |
| | gcgaggaagc ggaagagcgc ctgatgcggt atTTTctct tacgcattctg tgcggatTT | 2040 |
| | cacaccgcatt atggtgcaact ctcagttacaa tctgctctga tgccgcatac ttaagccagt | 2100 |
| 35 | atacactccg cttatcgctac gtgactgggt catggctgcg cccgcacacc cgccaaacacc | 2160 |
| | cgctgacgcgc ccctgacggg cttgtctgtc cccggcatcc gtttacagac aagctgtgac | 2220 |
| | cgtctccggg agctgcattgt gtcagaggTT ttcaccgtca tcaccgaaac gcgcgaggca | 2280 |
| 40 | ggatcccggaa cgccagcaag acgtagccca gcgctgcggc cagcttgcac ttgcgcgtaa | 2340 |
| | ctcacattaa ttgcgttgcg ctcactgccc gctttccagt cgggaaacct gtcgtgccag | 2400 |
| | ctgcattaaat gaatcgccca acgcgcgggg agaggcgggt tgcttattgg gcccagggt | 2460 |
| 45 | ggTTTTCTT ttcaccaggT agacgggcaaa cagctgatttgc cccttcaccg cctggccctg | 2520 |
| | agagagttgc agcaagcggcgt ccacgctggT ttgcggcagc aggcaaaaat cctgtttgt | 2580 |
| | ggTggTTaac ggcgggatAT aacatgagct gtcttcggta tcgtctatc ccactaccga | 2640 |
| 50 | gatatccgcac ccaacgcgcac gcccggactc ggtaatggcg cgcattgcgc ccagcgcacat | 2700 |
| | ctgatcgTTg gcaaccagca tcgcagtggg aacgatgccc tcattcagca tttgcattgt | 2760 |
| | ttgttggaaaa ccggacatgg cactccagtc gccttcccgT tccgctatcg gtcgaatttg | 2820 |
| 55 | attgcgagtg agatattttat gccagccagc cagacgcaga cgccggaga cagaacttaa | 2880 |

EP 2 477 649 B1

agagttcaact ataagcttag cccgcctaata gggcgggctt tttttctcg aggacgtc 4858

5 <210> 8
 <211> 28
 <212> DNA
 <213> Vibrio cholerae

10 <400> 8
 ggcgcgcaga acttggctat ttttaacc 28

15 <210> 9
 <211> 104
 <212> DNA
 <213> Vibrio cholerae

20 <220>
 <221> misc_feature
 <222> (36)..(37)
 <223> n denotes a, t, c or g

25 <220>
 <221> misc_feature
 <222> (38)..(38)
 <223> b denotes c or g or t

<400> 9

30 ggggggttcga agtttatgag tttgataata gggtgnnbtc attatatttt caaaaaaata 60
 cagacatagc agataagggtt aaaaatagcc aagttctggc gcgc 104

35 <210> 10
 <211> 934
 <212> DNA
 <213> Vibrio cholerae

40 <220>
 <221> CDS
 <222> (22)..(678)

45 <400> 10

50

55

EP 2 477 649 B1

| | |
|---|-----|
| attatttaaa ttgaggtagt a atg aaa cat cta ata aaa aac tat gta caa Met Lys His Leu Ile Lys Asn Tyr Val Gln 1 5 10 | 51 |
| 5 aaa tta att aaa aca gag ctt gat gct att cag tca aag tct gtt cat Lys Leu Ile Lys Thr Glu Leu Asp Ala Ile Gln Ser Lys Ser Val His 15 20 25 | 99 |
| 10 gat aat cga aac ttc att tac aat gga gag ttt tta att ctt gaa agc Asp Asn Arg Asn Phe Ile Tyr Asn Gly Glu Phe Leu Ile Leu Glu Ser 30 35 40 | 147 |
| 15 gaa ttt gga ttg cat tgt ttt ccc aga gtg cag ttg aac cat gct tta Glu Phe Gly Leu His Cys Phe Pro Arg Val Gln Leu Asn His Ala Leu 45 50 55 | 195 |
| 20 agc tac aaa aac cca aac ttt gat tta ggt atg cgt cac tgg att gtt Ser Tyr Lys Asn Pro Asn Phe Asp Leu Gly Met Arg His Trp Ile Val 60 65 70 | 243 |
| 25 | |
| 30 | |
| 35 | |
| 40 | |
| 45 | |
| 50 | |
| 55 | |

| | |
|--|-----|
| aat cat tgt aag cat gac acc act tat att gat atc ggt gca aac gtt | 291 |
| Asn His Cys Lys His Asp Thr Thr Tyr Ile Asp Ile Gly Ala Asn Val | |
| 75 80 85 90 | |
| 5 gga act ttc tgt gga atc gct gct cgt cat ata cac caa gga aaa att | 339 |
| Gly Thr Phe Cys Gly Ile Ala Ala Arg His Ile His Gln Gly Lys Ile | |
| 95 100 105 | |
| 10 ata gcg ata gaa cca ctc aca gaa atg gaa aat agt att agg atg aat | 387 |
| Ile Ala Ile Glu Pro Leu Thr Glu Met Glu Asn Ser Ile Arg Met Asn | |
| 110 115 120 | |
| 15 gtt caa tta aat aat cca cta gtt gag ttt cat cat ttt ggc tgt gca | 435 |
| Val Gln Leu Asn Asn Pro Leu Val Glu Phe His His Phe Gly Cys Ala | |
| 125 130 135 | |
| ata ggt gag aat gaa ggg gaa aat att ttc gaa gtt tat gag ttt gat | 483 |
| Ile Gly Glu Asn Glu Gly Glu Asn Ile Phe Glu Val Tyr Glu Phe Asp | |
| 140 145 150 | |
| 20 aat agg gtg tca tca tta tat ttt aaa aaa aat aca gac ata gca gat | 531 |
| Asn Arg Val Ser Ser Leu Tyr Phe Lys Lys Asn Thr Asp Ile Ala Asp | |
| 155 160 165 170 | |
| 25 aag gtt aaa aat agc caa gtt ctg gtt aga aag tta agt agt tta gat | 579 |
| Lys Val Lys Asn Ser Gln Val Leu Val Arg Lys Leu Ser Ser Leu Asp | |
| 175 180 185 | |
| 30 ata tcg cct act aac tct gta gtt ata aaa att gat gct gaa ggc gca | 627 |
| Ile Ser Pro Thr Asn Ser Val Val Ile Lys Ile Asp Ala Glu Gly Ala | |
| 190 195 200 | |
| gaa ata gag ata tta aac cag att tac gaa ttc aca gaa aag cat aat | 675 |
| Glu Ile Glu Ile Leu Asn Gln Ile Tyr Glu Phe Thr Glu Lys His Asn | |
| 205 210 215 | |
| 35 tga attgaatatt atatttgctt tgaatttgca atgggtcata tacagaggc | 728 |
| taatagaact tttgatgaga ttttaacat aataaactca aaattcggaa gtaaggcata | 788 |
| 40 ttttattcat ccattatcat ccgctgaaca tcctgagtt aataaagcaa cgcaaggat | 848 |
| taatggaaat atctgtttt aatatgtatc ataaaataat ttaatatatt ccgtatgtca | 908 |
| ttgcaagttc aacagacatt tcgaga | 934 |
| 45 <210> 11 | |
| <211> 218 | |
| <212> PRT | |
| <213> Vibrio cholerae | |
| 50 <400> 11 | |

Met Lys His Leu Ile Lys Asn Tyr Val Gln Lys Leu Ile Lys Thr Glu
 1 5 10 15

5 Leu Asp Ala Ile Gln Ser Lys Ser Val His Asp Asn Arg Asn Phe Ile
 20 25 30

10 Tyr Asn Gly Glu Phe Leu Ile Leu Glu Ser Glu Phe Gly Leu His Cys
 35 40 45

15 Phe Pro Arg Val Gln Leu Asn His Ala Leu Ser Tyr Lys Asn Pro Asn
 50 55 60

20 Phe Asp Leu Gly Met Arg His Trp Ile Val Asn His Cys Lys His Asp
 65 70 75 80

25 Thr Thr Tyr Ile Asp Ile Gly Ala Asn Val Gly Thr Phe Cys Gly Ile
 85 90 95

30 Ala Ala Arg His Ile His Gln Gly Lys Ile Ile Ala Ile Glu Pro Leu
 100 105 110

35 Thr Glu Met Glu Asn Ser Ile Arg Met Asn Val Gln Leu Asn Asn Pro
 115 120 125

40 Leu Val Glu Phe His His Phe Gly Cys Ala Ile Gly Glu Asn Glu Gly
 130 135 140

45 Glu Asn Ile Phe Glu Val Tyr Glu Phe Asp Asn Arg Val Ser Ser Leu
 145 150 155 160

50 Tyr Phe Lys Lys Asn Thr Asp Ile Ala Asp Lys Val Lys Asn Ser Gln
 165 170 175

55 Val Leu Val Arg Lys Leu Ser Ser Leu Asp Ile Ser Pro Thr Asn Ser
 180 185 190

Val Val Ile Lys Ile Asp Ala Glu Gly Ala Glu Ile Glu Ile Leu Asn
 195 200 205

60 Gln Ile Tyr Glu Phe Thr Glu Lys His Asn
 210 215

65 <210> 12
 <211> 1954
 <212> DNA
 <213> Artificial

70 <220>

<223> Synthetic plasmid

<400> 12

5 agtaatacga ctcactagtg ggcagatctt cgaatgcac ggcgcacccg tacgtctcga 60
 ggaattcctg caggatatct ggatccacga agcttcccat ggtgacgtca ccggttctag 120
 ataccttaggt gagctctggt accctctagt caaggcctgt cagccgttaa gtgttcctgt 180
 gtcactgaaa attgctttga gaggctctaa gggcttctca gtgcgttaca tccctggctt 240
 gttgtccaca accgttaaac cttaaaagct ttaaaagcct tatatattct tttttttctt 300
 ataaaaactta aaaccttaga ggctatttaa gttgctgatt tatattaatt ttattgttca 360

20

25

30

35

40

45

50

55

| | | | | | | | |
|-------------|-------------|-------------|------------|-------------|-------------|------------|------|
| aacatgagag | cttagtacgt | gaaacatgag | agcttagtac | gttagccatg | agagcttagt | 420 | |
| acgttagcca | tgagggttta | gttcgttaaa | catgagagct | tagtacgtta | aacatgagag | 480 | |
| 5 | cttagtacgt | gaaacatgag | agcttagtac | gtactatcaa | cagggtgaac | tgctgatctt | 540 |
| cagatctccg | cttgccctca | tctgttacgc | cggcggtagc | cggccagcct | cgcagagcag | 600 | |
| 10 | gattcccggtt | gagcaccgccc | aggtgcgaat | aaggagacgt | gaagaaggaa | caccgcctcg | 660 |
| cgggtgggccc | tacttcacct | atcctgccc | gctgacgccc | ttggatacac | caaggaaagt | 720 | |
| 15 | ctacacgaac | cctttggcaa | aatcctgtat | atcgtgcgaa | aaaggatgga | tataccgaaa | 780 |
| aaatcgctat | aatgaccccg | aagcagggtt | atgcagcgga | aaagcgctgc | ttccctgctg | 840 | |
| 20 | ttttgtggaa | tatctaccga | ctggaaacag | gcaaattgcag | gaaattactg | aactgagggg | 900 |
| acaggcgaga | gatctggcct | aggccgaccg | aataaatacc | tgtgacggaa | gatcacttcg | 960 | |
| 25 | cagaataaaat | aaatcctggt | gtccctgtt | ataccggaa | gccctgggccc | aactttggc | 1020 |
| gaaaatgaga | cgttgatcg | cacgtaagag | gttccaactt | tcaccataat | gaaataagat | 1080 | |
| 30 | cactaccggg | cgtatTTTT | gagttgtcga | gatTTTcagg | agctaaggaa | gctaaaatgg | 1140 |
| agaaaaaaat | cactggatat | accaccgtt | atatatccc | atggcatcgt | aaagaacatt | 1200 | |
| ttgaggcatt | tcagtcagtt | gctcaatgta | cctataacca | gaccgttcag | ctggatatta | 1260 | |
| 35 | cggccttttt | aaagaccgta | aagaaaaata | agcacaagtt | ttatccggcc | tttattcaca | 1320 |
| ttcttgcccg | cctgatgaat | gctcatccgg | aattacgtat | ggcaatgaaa | gacggtgagc | 1380 | |
| tggtgatatg | ggatagtgtt | cacccttgtt | acaccgtttt | ccatgagcaa | actgaaacgt | 1440 | |
| 40 | tttcatcgct | ctggagtgaa | taccacgacg | atttccggca | gtttctacac | atatattcgc | 1500 |
| aagatgtggc | gtgttacggt | gaaaacctgg | cctatttccc | taaagggttt | attgagaata | 1560 | |
| tgttttcgt | ctcagccaat | ccctgggtga | gtttcaccag | ttttgattt | aacgtggcca | 1620 | |
| 45 | atatggacaa | cttcttcgccc | cccgTTTca | ccatgggcaa | atattatacg | caaggcgaca | 1680 |
| aggtgctgat | gccgctggcg | attcaggttc | atcatgccgt | ttgtgatggc | ttccatgtcg | 1740 | |
| gcagaatgct | taatgaatta | caacagtact | gcgatgagtg | gcagggcggg | gcgtaatttt | 1800 | |
| 50 | tttaaggcag | ttattggtgc | ccataaacgc | ctggttgcta | cgcctgaata | agtgataata | 1860 |
| agcgatgaa | tggcagaaat | tcgaaagcaa | attcgacccg | gtcgtcgggt | cagggcaggg | 1920 | |
| tcgttaaata | gccgcttatg | tctattgctg | gttt | | | 1954 | |

<210> 13

<211> 2948

<212> DNA

<213> Artificial

<220>

<223> Synthetic vector construct with wild-type wbeT insert

<220>
<221> misc_feature
<222> (151)..(1011)
<223> wbeT wild-type CDS insert in reverse orientation

5

<400> 13

10

15

20

25

30

35

40

45

50

55

| | | |
|----|---|------|
| | agtaatacga ctcactatgt ggcagatctt cgaatgcac | 60 |
| 5 | gca gcgccaccc tacgtctcgaaattcctg caggatatct ggatctata | 120 |
| | gata tggatcttggaaatgtct gttgaacttgcaatgacata | 180 |
| | cgagaatata ttAAatttatt ttatgataca tattttAAac agatattccc | |
| 10 | attaatatcc tgcgttgctt tattttAAactc aggatgttca gcggtgata atggatgaat | 240 |
| | aaaatatgcc ttacttccga attttgagtt tattatgttaaaatctcat caaaagttct | 300 |
| | attagacctc tgtatatgac ccattgcaaa ttcaaaagcaa atataatatt caattccatt | 360 |
| 15 | atgctttct gtgaattcgt aaatctggtt taatatctct atttctgcgc cttcagcatc | 420 |
| | aattttata actacagagt tagtaggcga tataatctaaa ctacttaact ttctaaaccag | 480 |
| | aacttggcta tttttAACCT tatctgctat gtctgttattt ttttggaaat ataatgatga | 540 |
| 20 | caccctatta tcaaactcat aaacttcgaa aatattttcc ctttcattct cacctattgc | 600 |
| | acagccaaaa tggatggaaact caactatgtgg attatTTAAT tgaacattca tcctaaatact | 660 |
| | atTTTCCATT tctgtgagtg gttctatcgc tataattttt ctttggtaa tatgacgagc | 720 |
| 25 | agcgattcca cagaaagttc caacgtttgc accgatatac atataatgtgg tgcgtatgctt | 780 |
| | acaatgatta acaatccagt gacgcatacc taaatcaaag tttgggtttt tggatgttAA | 840 |
| | agcatggttc aactgcactc tgggaaaaca atgcccattcca aattcgcttt caagaattaa | 900 |
| 30 | aaactctcca ttgtAAatga agtttcgatt atcatgaaca gactttgact gaatagcatc | 960 |
| | aagctctgtt ttaattaatt tttgtacata gtttttattt agatgtttca tactacctca | 1020 |
| | atTTTAAATA atattagcgt aaaaaataac atacagaatc aacttgcaga tgcagagatc | 1080 |
| 35 | cacgaagctt cccatggta cgtcaccggc tctagatacc taggtgagct ctggtaaccct | 1140 |
| | ctagtcaagg cctgtcagcc gttaaatgtt cctgtgtcac tgaaaattgc tttgagaggc | 1200 |
| | tctaaaggct tctcagtgcg ttacatccct ggcttgggtt ccacaaccgt taaacctaa | 1260 |
| 40 | aagctttaaa agccttatattt attttttttt ttcttataaa actttAAacc tttagaggcta | 1320 |
| | tttaagttgc tgatttatattt gttcaaacat gagagcttag tacgtgaaac | 1380 |
| | atgagagctt agtacgttag ccatgagagc tttagtacgtt agccatgagg gtttagttcg | 1440 |
| 45 | ttaaacatga gagcttagta cgttAAacat gagagcttag tacgtgaaac atgagagctt | 1500 |
| | agtacgtact atcaacaggt tgaactgctg atcttcagat ctccgcttgc cctcatctgt | 1560 |
| 50 | tacgcccggcg gtagccggcc agcctcgcaag agcaggattc ccgttggca ccggccagggt | 1620 |
| | cgaataaggg acagtgaaga aggaacaccc gctcgccgggt gggctactt cacctatcct | 1680 |
| | gcccggctga cgccgttggta tacaccaagg aaagtctaca cgaaccctt ggcaaaatcc | 1740 |
| 55 | tgtatatcgt gcgaaaaagg atggatatac cgaaaaatc gctataatga ccccgaagca | 1800 |

| | | |
|----|---|--|
| 5 | gggttatgca gcggaaaagc gctgcttccc tgctgttttgc tggaaatatct accgactgga aacaggcaaa tgcagggaaat tactgaactg aggggacagg cgagagatct ggcctaggcc gaccgaataa atacctgtga cggaagatca cttcgccagaa taaataaaatc ctggtgtccc tggtgatacc gggaaagccct gggccaaactt ttggcgaaaa tgagacgttg atcggcacgt aagagggttcc aactttcacc ataatgaaat aagatcacta ccgggcgtat ttttgagtt 10 gtcgagattt tcaggagcta aggaagctaa aatggagaaa aaaatcactg gatataccac cggtgatata tcccaatggc atcgtaaaga acattttag gcatttcagt cagttgctca atgtacctat aaccagaccg ttcaagctgga tattacggcc tttttaaaga ccgtaaagaa aaataaagcac aagtttatac cggcccttat tcacattctt gcccgcctga tgaatgctca 20 tccggaatta cgtatggcaa tgaaagacgg tgagctggtg atatggata gtgttcaccc ttgttacacc gttttccatg agcaaactga aacgtttca tcgctctgga gtgaataccca cgacgatttc cggcagtttc tacacatata ttgcgaagat gtggcgtgtt acggtgaaaa cctggcctat ttccctaaag gttttattga gaatatgttt ttgcgtctcag ccaatccctg 25 ggtgagtttc accagtttg atttaaacgt ggccaatatg gacaacttct tcgccccgtt ttcaccatgg gcaaataattta tacgcaaggc gacaagggtgc tgatgcccgt ggcgattcag gttcatcatg ccgtttgtga tggcttccat gtcggcagaa tgcttaatga attacaacag 30 tactgcgatg agtggcaggg cggggcgtaa ttttttaag gcagttattg gtgcccataa acgcctgggtt gctacgcctg aataagtgtat aataagcggg tgaatggcag aaattcgaaa 35 gcaaattcga cccggtcgtc gtttcagggc agggtcgtta aatagccgt tatgtctatt gtgggtt 2948 | 1860 1920 1980 2040 2100 2160 2220 2280 2340 2400 2460 2520 2580 2640 2700 2760 2820 2880 2940 2948 |
| 40 | <210> 14 <211> 24 <212> DNA <213> Vibrio cholerae | |
| 45 | <400> 14 cggtgcaaac gtttcaactt tctg 24 | |
| 50 | <210> 15 <211> 25 <212> DNA <213> Vibrio cholerae | |
| 55 | <400> 15 ggaaaaacaat gccatccaaa ttgc 25 | |

<220>
<223> Synthetic construct

<400> 16

5

10

15

20

25

30

35

40

45

50

55

| | | |
|----|--|------|
| | ttcgatattt ttagttctt taggcccgt a | 60 |
| 5 | aacaaaagag gaaaatagac cagttgcaat ccaaacgaga gtctaata | 120 |
| | aaagtaaatac gcgccgggtt gttactgata aagcaggcaaa gacctaata | 180 |
| | caaagtgtat actttggcgt cacccttac atatttttagg tctttttta ttgtgcgtaa | 240 |
| 10 | ctaaacttgcc atcttcaaac aggagggctg gaagaagcag accgctaaca cagtacataa | 300 |
| | aaaaggagac atgaacgatg aacatcaaaa agtttgcaaa acaagcaaca gtattaacct | 360 |
| | ttactaccgc actgctggca ggaggcgaa ctcaagcggt tgcaagaa acgaacccaa | 420 |
| 15 | agccatataa ggaaacatac ggcatttccc atattacacg ccatgatatg ctgcaaatcc | 480 |
| | ctgaacagca aaaaaatgaa aaatatcaag ttccctgaatt cgattcgtcc acaattaaaa | 540 |
| | atatctttc tgcaaaaaggc ctggacgtt gggacagctg gccattacaa aacgctgacg | 600 |
| 20 | gcactgtcgc aaactatcac ggctaccaca tcgtctttgc attagccgga gatcctaaaa | 660 |
| | atgcggatga cacatcgatt tacatgttct atcaaaaagt cggcgaaact tctattgaca | 720 |
| | gctggaaaaa cgctggccgc gtctttaag acagcgacaa attcgatgca aatgattcta | 780 |
| 25 | tcctaaaaga ccaaacacaa gaatggtcag gttcagccac atttacatct gacggaaaaa | 840 |
| | tccgttatt ctacactgat ttctccgtt aacattacgg caaacaaca ctgacaactg | 900 |
| | cacaagttaa cgtatcagca tcagacagct ctttgaacat caacgggttta gaggattata | 960 |
| 30 | aatcaatctt tgacgggttac ggaaaaacgt atcaaaaatgt acagcagttc atcgatgaag | 1020 |
| | gcaactacag ctcaggcgac aaccatacgc tgagagatcc tcactacgtt gaagataaaag | 1080 |
| | gccacaaaata ctttgttattt gaagcaacaa ctggaaactga agatggctac caaggcgaag | 1140 |
| 35 | aatctttattt taacaaagca tactatggca aaagcacatc attcttcgtt caagaaagtc | 1200 |
| | aaaaacttctt gcaaagcgat aaaaaacgca cggctgagtt agcaaacggc gctctcggtt | 1260 |
| 40 | tgattgagct aaacgatgat tacacactga aaaaagtgtt gaaaccgctt attgcattctt | 1320 |
| | acacagtaaac agatgaaattt gaacgcgcga acgtctttaa aatgaacggc aaatggtacc | 1380 |
| | tgttcaactga ctcccgccgtt tcaaaaatgtt cgttgcgtt cattacgtt aacgatattt | 1440 |
| 45 | acatgcttgg ttatgtttctt aattctttaa ctggccata caagccgtt aacaaaactgt | 1500 |
| | gccttgggtt aaaaatggat cttgatccta acgttgcgtt ctttacttac tcacacttcg | 1560 |
| | ctgtacctca agcgaaagga aacaatgtcg tgattacaag ctatatgaca aacagaggat | 1620 |
| 50 | tctacgcaga caaacaatca acgtttgcgc caagcttcctt gctgaacatc aaaggcaaga | 1680 |
| | aaacatctgt tgcataaagac agcatccttgc aacaaggaca attaacagtt aacaaataaa | 1740 |
| | aacgcaaaag aaaaatgccgtt ttgaggccgtt ttgtgcgtt ctttccgtt ggaggttataa | 1800 |
| 55 | attgacgata tgatcgttac tacgactcac tagtggccgtt atcttcgtt gcatcgccgc | 1860 |

| | | |
|----|--|------|
| | caccgtacgt ctcgaggaat tcctgcagga tatctggatc cacgaagctt cccatggtga | 1920 |
| 5 | cgtcacccgt tctagatacc taggtgagct ctggtaccct ctagtcaagg cctgtcagcc | 1980 |
| | gttaagtgtt cctgtgtcac tgaaaattgc tttgagaggc tctaagggct tctcagtgcg | 2040 |
| | ttacatccct ggcttgttgt ccacaaccgt taaaccttaa aagctttaaa agccttatat | 2100 |
| 10 | attcttttt ttcttataaa actaaaaacc ttagaggcta tttaagttgc tgatttatat | 2160 |
| | taattttatt gttcaaaacat gagagcttag tacgtgaaac atgagagctt agtacgttag | 2220 |
| | ccatgagagc ttagtacgtt agccatgagg gtttagttcg ttaaacatga gagcttagta | 2280 |
| 15 | cgttaaacat gagagcttag tacgtgaaac atgagagctt agtacgtact atcaacaggt | 2340 |
| | tgaactgctg atcttcagat ctccgcttgc cctcatctgt tacgcccccg gtagccggcc | 2400 |
| | agcctcgcag agcaggattc ccgtttagca ccgccaggtg cgaataaggg acagtgaaga | 2460 |
| 20 | aggaacaccc gctcgccgg gggctactt cacctatcct gcccggctga cgccgttgg | 2520 |
| | tacaccaagg aaagtctaca cgaacccttt ggcaaaatcc tgtatatcgt gcgaaaaagg | 2580 |
| | atggatatac cgaaaaaaatc gctataatga ccccgaagca gggttatgca gcggaaaaagc | 2640 |
| 25 | gctgcttccc tgctgtttt tggaatatct accgactgga aacaggcaaa tgcagggaaat | 2700 |
| | tactgaactg aggggacagg cgagagatct ggcctaggcc gaccgaataa atacctgtga | 2760 |
| | cggaagatca cttcgcagaa taaataaatac ctgggtgtccc tggtgataacc ggaaagccct | 2820 |
| 30 | gggccaactt tgccggaaaa tgagacgttg atcggcacgt aagaggttcc aactttcacc | 2880 |
| | ataatgaaat aagatcacta ccggcgtat ttttgagtt gtcgagattt tcaggagcta | 2940 |
| | aggaagctaa aatggagaaa aaaatcactg gatataccac cgttgatata tcccaatggc | 3000 |
| 35 | atcgtaaaga acatttttag gcatttca cagttgctca atgtacctat aaccagaccg | 3060 |
| | ttcagctgga tattacggcc tttttaaaga ccgtaaagaa aaataagcac aagttttatc | 3120 |
| 40 | cggcctttat tcacattctt gcccgcctga tgaatgctca tccggatta cgtatggcaa | 3180 |
| | tgaaagacgg tgagctggtg atatggata gtgttcaccc ttgttacacc gttttccatg | 3240 |
| | agcaaactga aacgaaaaatca tcgctctgga gtgaatacca cgacgatttc cggcagttc | 3300 |
| 45 | tacacatata ttgcgaagat gtggcgtgtt acggtaaaaa cctggcctat ttccctaaag | 3360 |
| | ggtttattga gaatatgttt ttcgtctca ccaatccctg ggtgagttc accagtttg | 3420 |
| | atttaaacgt gccaatatg gacaacttct tcggccccgt tttcaccatg ggcaaatatt | 3480 |
| 50 | atacgcaagg cgacaagggtg ctgatgccgc tggcgattca gttcatcat gccgtttgt | 3540 |
| | atggcttcca tgctggcaga atgcttaatg aattacaaca gtactgcgt gagtggcagg | 3600 |
| | gcggggcgtta attttttaa ggcagttatt ggtgcccata aacgcctggc tgctacgcct | 3660 |
| 55 | gaataagtga taataagcgg atgaatggca gaaa | 3694 |

Claims

1. A vaccine comprising a *Vibrio cholerae* O1 cell, **characterized in that** said cell comprises O1 antigens of both Ogawa and Inaba serotypes, wherein the vaccine comprises multiple *Vibrio cholerae* O1 cells that comprise O1 antigens of both Ogawa and Inaba serotypes, and wherein on average, 10-90 % of the O1 antigens of the cells are of the Ogawa-serotype.

5 2. A vaccine according to any of the preceding claims, with the proviso that said vaccine does not contain any further immunologically active whole cells in addition to *Vibrio cholerae* O1 cells that comprise O1 antigens of both Ogawa and Inaba serotypes.

10 3. A vaccine according to any of the preceding claims, wherein the vaccine is for oral administration.

15 4. A vaccine according to any of the preceding claims, wherein the cell is formalin-inactivated.

5 5. A vaccine according to any of the preceding claims, wherein said cell is as defined in any of claims 10-11.

10 6. A vaccine according to any of the preceding claims, for use in preventive immunization.

20 7. A vaccine according to any of the preceding claims, for use in preventive immunization against cholera.

25 8. A vaccine according to any of the preceding claims, wherein the cell further comprises one or more ETEC colonization factor (CF) protein(s), such as CFA/I, CS2 or CS5, wherein said CF protein(s) is/are expressed either as single, double or hybrid fimbriae.

30 9. A vaccine according to claim 8, for use in preventive immunization against enterotoxigenic *Escherichia coli*-infection (ETEC).

35 10. A *Vibrio cholerae* O1 cell simultaneously expressing both Inaba and Ogawa antigens **characterized in that**

30 a. the cell comprises an endogenous *wbeT*-gene; and

35 b. the cell comprises a recombinant DNA-construct capable of modulating the expression level of endogenous *wbeT* gene or the enzymatic activity of the product thereof;

40 wherein said construct is adapted to modifying the endogenous *wbeT* gene of the host by means of homologous recombination and wherein

45 c. the modulated level of *WbeT* enzyme activity is such that the cell simultaneously expresses Inaba and Ogawa antigens, wherein 10-90 % of the O1 antigen expressed by the cell is of the Ogawa-serotype.

11. A *Vibrio cholerae* O1 cell according to claim 10, wherein the cell further expresses one or more ETEC colonization factor (CF) protein(s), such as CFA/I, CS2 or CS5, wherein said CF protein(s) is/are expressed either as single, double or hybrid fimbriae.

12. A method for manufacturing a vaccine, comprising the steps of:

45 a. providing a *Vibrio cholerae* O1 cell comprising O1 antigens of both Ogawa and Inaba serotypes; and
b. killing said cell.

13. A method according to claim 12, wherein the killing is performed by formalin-treatment.

50 14. A method according to any of claims 12-13 wherein the cell is a cell according to any of claims 10-11.

Patentansprüche

55 1. Impfstoff, umfassend eine *Vibrio cholerae* 01-Zelle, **dadurch gekennzeichnet, dass** die Zelle O1 Antigene der Serotypen Ogawa und Inaba umfasst, wobei der Impfstoff mehrere *Vibrio cholerae* 01-Zellen umfasst, die Antigene der Serotypen Ogawa und Inaba umfassen, und wobei durchschnittlich 10 bis 90 % der O1-Antigene der Zellen vom Serotyp Ogawa sind.

2. Impfstoff nach einem der vorhergehenden Ansprüche, mit der Vorgabe, dass der Impfstoff zusätzlich zu *Vibrio cholerae* O1 Zellen, die Antigene der Serotypen Ogawa und Inaba umfassen, keine weiteren immunologisch aktiven ganzen Zellen enthält.

5 3. Impfstoff nach einem der vorhergehenden Ansprüche, wobei der Impfstoff für eine orale Verabreichung bestimmt ist.

4. Impfstoff nach einem der vorhergehenden Ansprüche, wobei die Zelle mit Formalin inaktiviert ist.

10 5. Impfstoff nach einem der vorhergehenden Ansprüche, wobei die Zelle der Definition nach einem der Ansprüche 10 bis 11 entspricht.

6. Impfstoff nach einem der vorhergehenden Ansprüche zur Verwendung bei einer Schutzimpfung.

15 7. Impfstoff nach einem der vorhergehenden Ansprüche zur Verwendung bei einer Schutzimpfung gegen Cholera.

8. Impfstoff nach einem der vorhergehenden Ansprüche, wobei die Zelle ferner mindestens ein ETEC-Besiedelungsfaktor(CF)-Protein umfasst, wie beispielsweise CFA/I, CS2 oder CS5, wobei das CF-Protein bzw. die CF-Proteine als Einzel-, Doppel- oder Hybridfimbrien exprimiert ist bzw. sind.

20 9. Impfstoff nach Anspruch 8 zur Verwendung bei einer Schutzimpfung gegen eine Infektion mit enterotoxischen *Escherichia coli* (ETEC).

10. *Vibrio cholerae* 01-Zelle, die simultan sowohl Inaba- als auch Ogawa-Antigene exprimiert, **dadurch gekennzeichnet, dass**

25 a. die Zelle ein endogenes wbeT-Gen aufweist; und

b. die Zelle ein rekombinantes DNA-Konstrukt aufweist, das zum Modulieren des Expressionsgrades eines endogenen wbeT-Gens oder der enzymatischen Aktivität von dessen Produkt in der Lage ist; wobei das Konstrukt angepasst ist, um das endogene wbeT-Gen des Wirts durch homologe Rekombination zu modifizieren;

30 und wobei

c. das modulierte Niveau der Aktivität des WbeT-Enzyms derart ist, dass die Zelle simultan Inaba- und Ogawa-Antigene exprimiert, wobei 10 bis 90 % des von der Zelle exprimierten 01-Antigens vom Serotyp Ogawa sind.

35 11. *Vibrio cholerae* 01-Zelle nach Anspruch 10, wobei die Zelle ferner mindestens ein ETEC-Besiedelungsfaktor(CF)-Protein exprimiert, wie beispielsweise CFA/I, CS2 oder CS5, wobei das CF-Protein bzw. die CF-Proteine als Einzel-, Doppel- oder Hybridfimbrien exprimiert ist bzw. sind.

12. Verfahren zum Herstellen eines Impfstoffs, umfassend die Schritte:

40 a. Bereitstellen einer *Vibrio cholerae* 01-Zelle, umfassend 01-Antigene der Serotypen Ogawa und Inaba; und

b. Töten der Zelle.

13. Verfahren nach Anspruch 12, wobei das Töten durch eine Behandlung mit Formalin durchgeführt wird.

45 14. Verfahren nach einem der Ansprüche 12 bis 13, wobei es sich bei der Zelle um eine Zelle nach einem der Ansprüche 10 bis 11 handelt.

50 **Revendications**

1. Vaccin comprenant une cellule de *Vibrio cholerae* O1, **caractérisé en ce que** ladite cellule comprend des antigènes O1 des sérotypes Ogawa et Inaba, le vaccin comprenant de multiples cellules de *Vibrio cholerae* O1 qui comprennent des antigènes O1 des sérotypes Ogawa et Inaba, et dans lequel, en moyenne, 10 à 90 % des antigènes O1 des cellules sont du sérotype Ogawa.

55 2. Vaccin selon l'une quelconque des revendications précédentes, avec pour restriction que ledit vaccin ne contient pas d'autres cellules immunologiquement actives que les cellules de *Vibrio cholerae* O1 comprenant des antigènes

O1 des sérotypes Ogawa et Inaba.

3. Vaccin selon l'une quelconque des revendications précédentes, dans lequel le vaccin est destiné à une administration orale.

5

4. Vaccin selon l'une quelconque des revendications précédentes, dans lequel la cellule est inactivée par la formaline.

5. Vaccin selon l'une quelconque des revendications précédentes, dans lequel la cellule est une cellule selon l'une quelconque des revendications 10 et 11.

10

6. Vaccin selon l'une quelconque des revendications précédentes, destiné à une immunisation préventive.

7. Vaccin selon l'une quelconque des revendications précédentes, destiné à une immunisation préventive contre le choléra.

15

8. Vaccin selon l'une quelconque des revendications précédentes, dans lequel la cellule comprend en outre une ou plusieurs protéines facteurs de colonisation (CF) d'ETEC, telles que CFA/I, CS2 ou CS5, lesdites protéines CF étant exprimées sous la forme de fimbriae simples, doubles ou bien hybrides.

20

9. Vaccin selon la revendication 8, destiné à une immunisation préventive contre une infection par *Escherichia coli* entérotoxinogène (ECET).

10. Cellule de *Vibrio cholerae* O1 exprimant simultanément les antigènes Inaba et Ogawa, **caractérisée en ce que**

25

a. la cellule comprend un gène *wbeT* endogène ; et

b. la cellule comprend une construction d'ADN recombinant apte à moduler le niveau d'expression du gène *wbeT* endogène ou l'activité enzymatique de son produit ;

ladite construction étant conçue pour modifier le gène *wbeT* endogène de l'hôte au moyen d'une recombinaison homologue et dans laquelle

30

c. le niveau modulé de l'activité enzymatique de *WbeT* est tel que la cellule exprime simultanément les antigènes Inaba et Ogawa, 10 à 90 % de l'antigène O1 exprimé par la cellule étant du sérotype Ogawa.

35

11. Cellule de *Vibrio cholerae* O1 selon la revendication 10, dans lequel la cellule exprime en outre une ou plusieurs protéines facteurs de colonisation (CF) d'ETEC, telles que CFA/I, CS2 ou CS5, lesdites protéines CF étant exprimées sous la forme de fimbriae simples, doubles ou bien hybrides.

40

12. Procédé de production d'un vaccin, comprenant les étapes consistant à :

a. disposer d'une cellule de *Vibrio cholerae* O1 comprenant des antigènes O1 des sérotypes Ogawa et Inaba ; et

b. inactiver ladite cellule.

13. Procédé selon la revendication 12, dans lequel l'inactivation s'effectue par traitement à la formaline.

45

14. Procédé selon l'une quelconque des revendications 12 et 13, dans lequel la cellule est une cellule selon l'une quelconque des revendications 10 et 11.

50

55

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61272351 B [0119]

Non-patent literature cited in the description

- CHIANG ; MEKALONOS. *Infection and Immunity*, 01 January 2000, vol. 68, ISSN 0019-9567, 6391-6397 [0011]
- RIJPKEMA et al. *Journal of Medical Microbiology*, 01 November 2004, vol. 53 (11), ISSN 0022-2615, 1105-1107 [0012]
- KANUNGO et al. *Vaccine*, 16 November 2009, vol. 27 (49), ISSN 0264-410X, 6887-6893 [0013]
- SANCHEZ ; HOLMGREN. *Current Opinion in Immunology*, 01 August 2005, vol. 17 (4), ISSN 0952-7915, 388-398 [0014]
- Remington's Pharmaceutical Sciences. Mack Publishing Company [0053]
- NYGREN E ; LI BL ; HOLMGREN J ; ATTRIDGE SR. *Infect Immun.*, August 2009, vol. 77 (8), 3475-84 [0080]
- TOBIAS J ; LEBENS M ; BÖLIN I ; WIKLUND G ; SVENNERHOLM AM. *Vaccine*, 06 February 2008, vol. 26 (6), 743-52 [0115]
- TOBIAS J ; SVENNERHOLM AM ; HOLMGREN J ; LEBENS M. *Appl Microbiol Biotechnol.*, July 2010, vol. 87 (4) [0116]
- TOBIAS J ; HOLMGREN J ; HELLMAN M ; NYGREN E ; LEBENS M ; SVENNERHOLM AM. *Vaccine*, 20 August 2010 [0117]

SZABADALMI IGÉNYPONTOK

1. Egy *Vibrio cholerae* O1 sejtet tartalmazó vakcina, azzal jellemzve, hogy a sejt az Ogawa és Inaba szerotípusok O1 antigénjeit egyaránt tartalmazza, ahol a vakcina több, az Ogawa és Inaba szerotípusok O1 antigénjeit egyaránt tartalmazó *Vibrio cholerae* O1 sejteket tartalmaz, és ahol átlagban, a sejtek O1 antigénjainak 10-90 %-a Ogawa-szerotípus.
2. Az előző igénypontok bármelyike szerinti vakcina, azzal a feltételel, hogy a vakcina nem tartalmaz semmilyen további immunológiaiag aktív teljes sejtet az Ogawa és Inaba szerotípusok O1 antigénjeit egyaránt tartalmazó *Vibrio cholerae* O1 sejtekben kívül.
3. Az előző igénypontok bármelyike szerinti vakcina, ahol a vakcina orális adagolásra szolgál.
4. Az előző igénypontok bármelyike szerinti vakcina, ahol a sejt formalinnal inaktivált.
5. Az előző igénypontok bármelyike szerinti vakcina, ahol a sejt a 10-11. igénypontok bármelyikében megadott.
6. Az előző igénypontok bármelyike szerinti vakcina, megelőző immunizációra történő alkalmazásra.
7. Az előző igénypontok bármelyike szerinti vakcina, kolera elleni megelőző immunizációra történő alkalmazásra.
8. Az előző igénypontok bármelyike szerinti vakcina, ahol a sejt tartalmaz továbbá egy vagy több ETEC kolonizációs faktor (CF) protein(ek)t, például CFA/I-t, CS2-t vagy CS5-t, ahol a CF protein(ek) egyszeres, dupla vagy hibrid fimbriaként fejeződnek ki.



SZTNH-100004278

9. A 8. igénypont szerinti vakcina, enterotoxigenikus *Escherichia coli*-fentőzés (ETEC) elleni megelőző immunizációra történő alkalmazásra.

10. Inaba és Ogawa antigéneket egyidejűleg kifejező *Vibrio cholerae* O1 sejt azzal jellemzve, hogy

- a. a sejt egy endogén *wbeT*-gént tartalmaz; és
- b. a sejt egy rekombináns DNS-konstruktot tartalmaz, mely az endogén *wbeT* gén expressziós szintjét vagy a termékének enzimaktivitását modulálni képes; ahol a konstrukt a gazda endogén *wbeT* génjének homológ rekombinációval történő módosítására kialakított, és ahol
- c. a *WbeT* enzimaktivitás modulációs szintje olyan, hogy a sejt egyidejűleg fejez ki Inaba és Ogawa antigéneket, ahol a sejt által kifejezett O1 antigén 10-90 %-a Ogawa-szerotípusú.

11. A 10. igénypont szerinti *Vibrio cholerae* O1 sejt, ahol a sejt kifejez továbbá egy vagy több ETEC kolonizációs faktor (CF) proteinet, például CFA/I-t, CS2-t vagy CS5-t, ahol a CF protein(ek) egyszeres, dupla vagy hibrid fimbriáként fejeződnek ki.

12. Eljárás egy vakcina előállítására, melynek során:

- a. az Ogawa és Inaba szerotípusok O1 antigénjeit egyaránt tartalmazó *Vibrio cholerae* O1 sejyet hozunk létre; és
- b. a sejyet előírjük.

13. A 12. igénypont szerinti eljárás, ahol az előlést formalinos kezeléssel végezzük.

14. A 12-13. igénypontok bármelyike szerinti eljárás, ahol a sejt a 10-11. igénypontok bármelyike szánni sejt.
