

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3494962 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **A 61 K 9/12 (2006.01)** **A 61 K 31/495 (2006.01)** **A 61 K 31/519 (2006.01)**
A 61 P 9/10 (2006.01) **A 61 P 11/02 (2006.01)** **A 61 P 11/06 (2006.01)**
A 61 P 17/06 (2006.01) **A 61 P 27/02 (2006.01)** **A 61 P 37/02 (2006.01)**
A 61 P 37/06 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2021-06-21**
- (80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2021-05-26**
- (86) Europæisk ansøgning nr.: **19152629.2**
- (86) Europæisk indleveringsdag: **2015-09-15**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-06-12**
- (30) Prioritet: **2014-09-15 GB 201416274** **2015-03-19 GB 201504662**
- (62) Stamansøgningsnr: **17207623.4**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV
MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Verona Pharma PLC, One Central Square, Cardiff, CF10 1FS, Storbritannien**
- (72) Opfinder: **SPARGO, Peter Lionel, Nash House, Beech Drive, Harbledown Park, Harbledown, Canterbury, Kent
CT2 8NR, Storbritannien**
FRENCH, Edward James, 28 Norman Road, Canterbury, Kent CT1 3LX, Storbritannien
HAYWOOD, Phillip A, Goldings, Silver Street, Anstey, Buntingford, Hertfordshire SG9 0DH, Storbritannien
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **Flydende inhalationsformulering omfattende RPL554**
- (56) Fremdragne publikationer:
WO-A1-2014/037727
**L. CALZETTA ET AL: "Effect of the Mixed Phosphodiesterase 3/4 Inhibitor RPL554 on Human Isolated
Bronchial Smooth Muscle Tone", JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS, vol.
346, no. 3, 13 June 2013 (2013-06-13) , pages 414-423, XP055202034, DOI: 10.1124/jpet.113.204644**
**C. COMPTON ET AL: "The Novartis view on emerging drugs and novel targets for the treatment of chronic
obstructive pulmonary disease", PULMONARY PHARMACOLOGY & THERAPEUTICS, vol. 26, no. 5, 1 October
2013 (2013-10-01), pages 562-573, XP055121337, ISSN: 1094-5539, DOI: 10.1016/j.pupt.2013.05.009**
**FRANCIOSI LUI G ET AL: "Efficacy and safety of RPL554, a dual PDE3 and PDE4 inhibitor, in healthy
volunteers and in patients with asthma or chronic obstructive pulmonary disease: findings from four clinical
trials.", THE LANCET. RESPIRATORY MEDICINE NOV 2013, vol. 1, no. 9, November 2013 (2013-11), pages 714-
727, XP009177640, ISSN: 2213-2600**

DESCRIPTION


FIELD OF THE INVENTION

[0001] The present invention relates to a liquid pharmaceutical composition comprising a respiratory drug.

BACKGROUND OF THE INVENTION

[0002] The toxicokinetic properties of liquid pharmaceutical compositions are unpredictable. It is important that an active pharmaceutical ingredient (API) is correctly formulated so that a safe, effective and controlled dosage is provided when the composition is delivered to a patient. This is particularly the case for inhaled compositions. Furthermore, liquid pharmaceutical compositions must have reliable long term stability to ensure that the dosage profile of the composition will be maintained after storage. This avoids the administration of incorrect dosages. Pharmaceutical compositions must also be formulated in such a way that administration is not unpleasant for a patient, for instance as regards taste and acidity.

[0003] RPL554 (9,10-Dimethoxy-2-(2,4,6-trimethylphenylimino)-3-(N-carbamoyl-2-aminoethyl)-3,4,6,7-tetrahydro-2H-pyrimido[6,1-a]isoquinolin-4-one) is a dual PDE3/PDE4 inhibitor and is described in WO 00/58308 (and is also sometimes referred to as RPL554). As a combined PDE3/PDE4 inhibitor, RPL554 has both anti-inflammatory and bronchodilatory activity and is useful in the treatment of respiratory disorders such as asthma and chronic obstructive pulmonary disease (COPD). The structure of RPL554 is shown below.



[0004] It is often preferable to administer RPL554 by inhalation because of its efficacy in the treatment of respiratory disorders. An effective method of administration is nebulisation. Franciosi *et al.* disclose a solution of RPL554 in a citrate-phosphate buffer (Efficacy and safety of RPL554, a dual PDE3 and PDE4 inhibitor, in healthy volunteers and in patients with asthma or chronic obstructive pulmonary disease: findings from four clinical trials, *The Lancet: Respiratory Medicine* 11/2013; 1(9):714-27. DOI: 10.1016/S2213-2600(13)70187-5).

[0005] Calzetta *et al.*, *J Pharmacol Exp Ther* 346:414-423, 2013 describes the effect of the mixed PDE 3/4 inhibitor RPL554 on human isolated bronchial smooth muscle tone. Compton *et al.*,

Pulmonary Pharmacology & Therapeutics 26 (2013) 562-573 describes treatments for COPD.

SUMMARY OF THE INVENTION

[0006] It is a surprising finding of the present invention that the optimal composition for inhalation comprising RPL554 is a liquid pharmaceutical composition comprising a suspension of particles of RPL554. Suspensions of RPL554 have been found to have highly desirable properties for a clinical setting. A number of benefits have been found to be associated with a liquid pharmaceutical composition which comprises a diluent and a suspension of particles of RPL554 or a pharmaceutically acceptable salt thereof.

[0007] The liquid pharmaceutical composition of the invention has greatly improved properties over previous formulations, for instance when compared with a solution formulation of RPL554. Regarding toxicokinetic properties, the suspension formulation has been found to have delayed release characteristics compared to those of a solution. This can reduce the number of treatments that must be administered to a patient. The suspension has also been found to allow a much greater dosage of RPL554 to be administered, as can be seen from plasma $C_{\max}$ and AUC values. This can also reduce the frequency of administrations required.

[0008] The liquid pharmaceutical composition of the invention also shows excellent stability, with compositions showing no degradation after 12 months at ambient conditions (25°C), nor after 6 months at accelerated conditions (40°C). Degradation is observed in a comparable solution composition under the same conditions. The compositions may show no degradation after 12 months at ambient conditions (25°C/60%RH), nor after 6 months at accelerated conditions (40°C/75%RH).

[0009] Accordingly, the present invention provides a liquid pharmaceutical composition comprising a diluent and a suspension of particles of 9,10-dimethoxy-2-(2,4,6-trimethylphenylimino)-3-(N-carbamoyl-2-aminoethyl)-3,4,6,7-tetrahydro-2H-pyrimido[6,1-a]isoquinolin-4-one (RPL554) or a pharmaceutically acceptable salt thereof, wherein: the particles of RPL554 have a particle size distribution with a Dv_{50} (median particle size by volume) value of from 0.2 μm to 5 μm ; and the composition comprises a phosphate buffer at a concentration of from 0 mg/mL to 5 mg/mL.

[0010] The invention also provides a liquid pharmaceutical composition according to the invention for use in the treatment of the human or animal body. A liquid pharmaceutical composition according to the invention may be used in the treatment or prevention of a disease or condition selected from asthma, allergic asthma, hay fever, allergic rhinitis, bronchitis, emphysema, bronchiectasis, chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), steroid resistant asthma, severe asthma, paediatric asthma, cystic fibrosis, lung fibrosis, pulmonary fibrosis, interstitial lung disease, skin disorders, atopic dermatitis, psoriasis, ocular inflammation, cerebral ischaemia, inflammatory diseases and auto-immune diseases.

[0011] Typically, said disease or condition is asthma or COPD, more typically COPD.

[0012] Related to the invention is a method of treating or preventing a disease or condition as defined herein in a subject, which method comprises administering to said subject an effective amount of a liquid pharmaceutical composition according according to the invention.

BRIEF DESCRIPTION OF THE FIGURES

[0013]

Figure 1 - The provocation concentration (PC) of acetylcholine which cumulatively produced a 50% increase in total airway resistance (Acetylcholine RL PC50) in animals treated with saline or RPL554 solution formulation: 2 and 5 h time point: Acetylcholine RL PC50 was log transformed for the purposes of statistical analysis. Post hoc t-test of the untransformed data showed differences between groups vs saline. *Non parametric analysis did show a significant difference between 2 h (1 mg/mL RPL554 solution, $P = 0.0190$ unadjusted) and 5 h (2.5 mg/mL solution, $P = 0.0095$ unadjusted) vs saline group. N values = 6, 4, 4; 4, 4 left to right. (Note 5 h control not plotted).

Figure 2 - Acetylcholine RL PC50 in animals treated with vehicle or the RPL554 suspension formulation: 2 and 5 h time point: Acetylcholine RL PC50 was log transformed for the purposes of statistical analysis (upper panel). Non parametric analysis did show a significant difference between 2 h (10 mg/mL RPL554 suspension, $P = 0.0268$ unadjusted) vs vehicle control group. Re-analysis of the data by combining the vehicle data (2 + 5 h), revealed an overall significant bronchoprotection ($P=0.0016$) which was reflected by a significant degree of bronchoprotection with 2.5 mg/mL RPL554 suspension (2.52 fold (1-5.72), $P < 0.05$ and 10 mg/mL RPL554 suspension (4.67 fold (2-11), $*P < 0.05$) vs vehicle. At the 5 h time point, RPL554 10 mg/mL suspension caused a significant degree of bronchoprotection (2.77 fold (1.3-6.0), $*P < 0.05$). (N=8,4,4,3, 6, left to right) (lower panel).

Figure 3 shows mean concentrations of RPL554 in plasma from healthy human volunteers following administration of either (i) the suspension formulation or (ii) the solution formulation by nebuliser at a nominal dose-level of about 1.5 mg.

DETAILED DESCRIPTION OF THE INVENTION

[0014] Typically, the particles of RPL554 are micronized particles.

[0015] The particles have a particle size distribution with a Dv_{50} value of from 0.2 μm to 5 μm . Typically, the particles of RPL554 have a particle size distribution with a Dv_{50} value of from about 0.7 μm to about 3.0 μm . For instance, the particles of RPL554 may have a particle size distribution with a Dv_{50} value of from 0.9 μm to 1.7 μm , or alternatively from 1.7 μm to 2.7 μm . Often, the particles of RPL554 have a particle size distribution with a Dv_{50} value of from about 1.1 μm to about 2.6 μm .

[0016] The Dv50 value is the median particle size for a volume distribution. Thus, half the volume of the particles is comprised in particles having diameters of less than the Dv50 value and half the volume of the particles is comprised in particles having diameters of greater than the Dv50 value. This is a well known manner in which to describe particle size distributions.

[0017] The particles typically have a particle size distribution with a Dv10 value of from about 0.4 μm to about 1.0 μm . The particles typically have a particle size distribution with a Dv90 value of from about 2.0 μm to about 4.0 μm . The Dv10 value reflects the particle diameter where 10% of the volume of the sample is in particles having a particle diameter less than the Dv10 value. The Dv90 value reflects the particle diameter where 90% of the volume of the sample is in particles having a particle diameter less than the Dv90 value.

[0018] The technique used to measure the Dv50 value is typically laser diffraction. For instance, the RPL554 particles typically have a particle size distribution with a Dv50 value of from 0.2 μm to 5 μm as measured by laser diffraction. The particle size distribution analysis can be performed by laser diffraction using the Malvern Spraytec in conjunction with a wet dispersion cell. Typically, the instrument parameters for the Malvern Spraytec are as follows:

- particle - standard opaque particle;
- refractive index Particle - 1.50;
- refractive index (imaginary) - 0.50;
- density of particle - 1.00;
- refractive index of dispersant - 1.33;
- controller unit - 1000RPM;
- measurement type - timed;
- initial sampling time - 30s;
- obscuration - 20% - 30%;
- dispersant - 1% Polysorbate 20 in deionised water.

[0019] The particles of RPL554 may be produced by any pharmaceutically acceptable size reduction process or particle size controlled production process. For instance, the particles may be produced by spray-drying a solution of RPL554, or by controlled crystallisation, or by size reduction of a solid form of RPL554, for example by air jet milling, mechanical micronisation or media milling.

[0020] The concentration of particles of RPL554 in the liquid pharmaceutical composition is typically from about 0.01 mg/mL to about 40 mg/mL. More typically, the concentration of particles of RPL554 in the liquid pharmaceutical composition is from about 0.1 mg/mL to about 20 mg/mL. For instance, the concentration of particles of RPL554 in the liquid pharmaceutical composition may be from about 0.01 mg/mL to about 5 mg/mL, or from 0.1 mg/mL to 5 mg/mL. The concentration of particles of RPL554 in the liquid pharmaceutical composition may for instance be from 0.1 mg/mL to 6 mg/mL.

[0021] The pH of the suspension may be from about 6 to about 8 or from about 6.5 to about 7.

[0022] The liquid pharmaceutical composition may further comprise one or more surfactants. The surfactants are pharmaceutically acceptable surfactants. The surfactants may be non-ionic surfactants, anionic surfactants, cationic surfactants or zwitterionic surfactants. Preferably, the one or more surfactants are selected from one or more non-ionic surfactants.

[0023] The one or more surfactants are typically selected from polyoxyethylene glycol alkyl ethers, polyoxypropylene glycol alkyl ethers, glucoside alkyl ethers, polyoxyethylene glycol octylphenol ethers, polyoxyethylene glycol alkylphenol ethers, glycerol alkyl esters, polyoxyethylene glycol sorbitan alkyl esters (polysorbates), sorbitan alkyl esters, cocamide MEA, cocamide DEA, dodecyldimethylamine oxide, block copolymers of polyethylene glycol and polypropylene glycol (poloxamers) and polyethoxylated tallow amine (POEA).

[0024] Preferably, the one or more surfactants are selected from polyoxyethylene glycol sorbitan alkyl esters, for instance polysorbate 20 (polyoxyethylene (20) sorbitan monolaurate), polysorbate 40 (polyoxyethylene (20) sorbitan monopalmitate), polysorbate 60 (polyoxyethylene (20) sorbitan monostearate) and polysorbate 80 (polyoxyethylene (20) sorbitan monooleate), and sorbitan alkyl esters, for instance sorbitan monolaurate, sorbitan monopalmitate, sorbitan monostearate, sorbitan tristearate and sorbitan monooleate.

[0025] The total concentration of the one or more surfactants is typically from about 0.01 mg/mL to about 2 mg/mL. The composition often comprises two or more surfactants, for instance a polysorbate and a Span surfactant.

[0026] The liquid pharmaceutical composition may further comprise one or more buffers. The buffers are pharmaceutically acceptable buffers. The buffers may be any buffers suitable for use in a liquid pharmaceutical composition suitable for inhalation. The one or more buffers are typically selected from citrate or phosphate buffers. Citrate buffers include citric acid, sodium citrate and mixtures thereof. Phosphate buffers include phosphoric acid, monosodium phosphate, dibasic sodium phosphate and mixtures thereof.

[0027] The composition comprises a phosphate buffer at a concentration of from 0 mg/mL to 5 mg/mL. Typically, the composition does not comprise a phosphate buffer. For instance, the composition does not comprise a buffer.

[0028] The liquid pharmaceutical composition typically further comprises a tonicity adjuster. The tonicity adjuster may be any pharmaceutically acceptable tonicity adjuster. Examples of tonicity adjusters include simple non toxic salts such as alkali metal halides, for instance sodium chloride and potassium iodide. Typically, the tonicity adjuster is sodium chloride.

[0029] The concentration of the tonicity adjuster will depend on the amount required to reach the desired tonicity, for instance isotonicity with the body or lungs. The concentration of the tonicity adjuster is typically from about 2 mg/mL to about 8 mg/mL, and more typically from about 3.5 mg/mL to 6 mg/mL.

[0030] The liquid pharmaceutical composition may comprise other components. Alternatively,

other components may be excluded. In some cases, the liquid pharmaceutical composition does not comprise a β_2 -adrenergic receptor agonist or a muscarinic receptor antagonist (or does not comprise greater than 0.1 wt% total of a β_2 -adrenergic receptor agonist and a muscarinic receptor antagonist).

[0031] The diluent may be any pharmaceutically acceptable diluent. The diluent is suitable for administration by inhalation. Examples of suitable diluents include water, ethanol and glycerol. The diluent is preferably water. The diluent is preferably sterile.

[0032] Typically, the liquid pharmaceutical composition is suitable for administration by nebulizer.

[0033] The invention also provides a nebulizer comprising a liquid pharmaceutical composition according to the invention. The nebulizer is typically loaded with the liquid pharmaceutical composition. The nebulizer typically comprises from about 1 mL to about 200 mL, more typically from 1 mL to 20 mL of the liquid pharmaceutical composition.

[0034] Nebulizers use compressed air to aerosolise a liquid pharmaceutical composition into an aerosol that is inhaled into a subject's respiratory tract. Examples of nebulizers include a soft mist nebulizer, a vibrating mesh nebulizer, a jet nebulizer and an ultrasonic wave nebulizer. Suitable nebulizer devices include the Philips I-neb™ (Philips), the Pari LC Sprint (Pari GmbH), the AERx^{RTM} Pulmonary Delivery System (Aradigm Corp.) and the Pari LC Plus Reusable Nebulizer (Pari GmbH).

[0035] The invention also provides a liquid pharmaceutical composition for use in the treatment of the human or animal body. The composition is as defined herein. The liquid pharmaceutical composition is typically for use in the treatment of the human or animal body, wherein treatment comprises administration by inhalation.

[0036] The compositions of the invention allow for delayed release of RPL554 into the bloodstream. The plasma concentration of RPL554 after a certain time is therefore increased relative to the use of known compositions of RPL554. Typically, the blood plasma concentration of RPL554 is greater than or equal to 1 ng/mL at a time greater than or equal to four hours after a nebulised dose of the composition is administered. The nebulised dose is typically from 0.02 mg/kg to 0.6 mg/kg. For example, the nebulised dose may be from 0.2 mg/kg to 0.6 mg/kg. For instance, the invention provides a liquid pharmaceutical composition as defined herein for use in the treatment of the human or animal body, wherein the blood plasma concentration of RPL554 is greater than or equal to 1 ng/mL at a time greater than or equal to four hours after a nebulised dose of from 0.2 mg/kg to 0.6 mg/kg is administered via inhalation of the nebulised composition of the invention. The blood plasma concentration of RPL554 is typically greater than or equal to 1 ng/mL at a time greater than or equal to three, four or five hours after a nebulised dose of the composition is administered.

[0037] Related to the invention is a method of treating or preventing a disease or condition as defined herein in a subject, which method comprises administering to said subject an effective amount of a liquid pharmaceutical composition as defined herein. Typically, the method is for

treating a disease or condition. The disease or condition is typically selected from asthma, allergic asthma, hay fever, allergic rhinitis, bronchitis, emphysema, bronchiectasis, chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), steroid resistant asthma, severe asthma, paediatric asthma, cystic fibrosis, lung fibrosis, pulmonary fibrosis, interstitial lung disease, skin disorders, atopic dermatitis, psoriasis, ocular inflammation, cerebral ischaemia and auto-immune diseases. Preferably the method comprises treating asthma or chronic obstructive pulmonary disease (COPD), more preferably COPD.

[0038] An effective amount of RPL554 is typically from about 0.01 mg/kg to 50 mg/kg for a single dose. An effective amount of RPL554 is often from about 0.01 mg/kg to 1 mg/kg for a single dose. For instance, an effective amount may be a dose of from about 0.1 mg to about 500 mg, or from about 0.1 mg to 100 mg, preferably from about 0.1 mg to about 6 mg. An effective amount may be from 0.1 mg to 12 mg. A single dose of RPL554 may be from 0.5 mg to 3 mg, for instance about 1.5 mg. Doses may be administered daily. For instance, the dose of RPL554 may be from 0.01 mg/kg/day to 50 mg/kg/day, typically from 0.01 mg/kg/day to 10 mg/kg/day or from 0.01 mg/kg/day to 1 mg/kg/day. These doses are typically the nominal dose emitted from the inhaler. The liquid pharmaceutical composition may be administered once, twice or three times a day, or may be administered twice, three times, four times or five times a week. The composition may be administered as often as required by the patient.

[0039] An effective amount may be a dose of from about 0.1 mg to about 500 mg, or from about 0.1 mg to 100 mg, preferably from about 0.5 mg to about 6 mg. An effective amount may be from 0.1 mg to 12 mg. A single dose of RPL554 may be from 0.5 mg to 24 mg, which may be administered by nebuliser. A single dose of RPL554 may be from 0.5 mg to 6 mg, for instance about 1.5 mg. Doses may be administered once, twice or three times daily. For instance, the dose of RPL554 may be from 0.1 mg/day to 50 mg/day, typically from 1.5 mg/day to 18 mg/day. These doses are typically the nominal dose emitted from the inhaler. The liquid pharmaceutical composition may be administered once, twice or three times a day, or may be administered twice, three times, four times or five times a week. The composition may be administered as often as required by the patient.

[0040] It is a surprising finding of the present invention that a suspension formulation of RPL554 allows for high doses of RPL554 to be administered to a patient without a concomitant increase in cardiovascular liability. The liquid pharmaceutical composition is therefore suitable for use in the treatment of patients suffering from cardiovascular disorders or at risk from suffering cardiovascular disorders.

[0041] For example, the composition may be used to treat a disease or condition as defined above in a patient suffering from cardiovascular vulnerability. Patients suffering from cardiovascular vulnerability are typically those at increased risk of adverse events arising from tachycardia or hypertension. More typically, patients suffering from cardiovascular vulnerability are those suffering from a pre-existing heart condition, or a condition which would be aggravated by tachycardia or hypertension. Such patients include those suffering from coronary artery disease, cardiomyopathy, hypertension, hypertensive heart disease, heart failure, arrhythmia, cardiac dysrhythmias, endocarditis, myocarditis, valvular heart disease and stroke. Also included are patients who have suffered a heart attack.

[0042] The invention is described in more detail by the following Examples.

EXAMPLES

Reference Example 1 - formulation and stability of RPL554 suspension formulation

[0043] Liquid pharmaceutical compositions were produced with the formulations set out Tables 1 and 2 below.

Table 1 - formulation of variant 1a (0.4 mg/mL)

Constituent	Function	Concentration (mg/mL)
RPL554 (micronized)	Active compound	0.4
Polysorbate 20 (Tween 20)	Surfactant	0.50
Sorbitan Monolaurate (Span 20)	Surfactant	0.05
Monosodium Phosphate Monohydrate	Buffer	6.58
Dibasic Sodium Phosphate Anhydrous	Buffer	6.80
Sodium Chloride	Tonicity Adjuster	4.80
Water	Diluent	Q.S.

Table 2 - formulation of variant 1b (20.0 mg/mL)

Constituent	Function	Concentration (mg/mL)
RPL554 (micronized)	Active compound	20.0
Polysorbate 20 (Tween 20)	Surfactant	0.50
Sorbitan Monolaurate (Span 20)	Surfactant	0.05
Monosodium Phosphate Monohydrate	Buffer	6.58
Dibasic Sodium Phosphate Anhydrous	Buffer	6.80
Sodium Chloride	Tonicity Adjuster	4.80
Water	Diluent	Q.S.

[0044] Assay testing was performed in duplicate. The key method parameters are as follows:

- mobile phase: acetonitrile:water:TFA (45:55:0.1);
- column: Waters X-Bridge phenyl, 3.5 µm, 150 x 4.6 mm;
- flow rate: 1.5 ml/min;
- injection volume: 10 µL;
- detection: UV @ 254 nm;
- runtime: 6 minutes; and
- sample and standard concentration: 0.1 mg/mL.

[0045] Variant 1a was stored under conditions of either (i) a temperature of 25°C and a relative humidity (RH) of 60% or (ii) a temperature of 40°C and a RH of 75%. The appearance, pH, assay, impurities and particle size distribution (PSD) were measured initially, at 1 month, at 2 months and at 6 months (6 months only for the 25°C and RH of 60% conditions). The amount of impurities were measured by HPLC with the relative retention time (RRT) of the impurity indicated.

[0046] The results are shown in Tables 3 and 4 below.

Table 3 - Stability of variant 1a under conditions: 25°C/60% RH

Test	Initial	1 month	2 months	6 months
Appearance	Light yellow suspension free from agglomerates	Light yellow suspension free from agglomerates	Light yellow suspension free from agglomerates	Light yellow suspension free from agglomerates
pH	6.55	6.59	6.68	6.51
Assay (mg/g)	0.372	0.374	0.373	0.372
Impurities (% area)				
Total	1.8	1.8	1.6	1.9
Greatest	0.490 (RRT 0.28)	0.533 (RRT 0.26)	0.429 (RRT 0.26)	0.463 (RRT 0.26)
Second greatest	0.401 (RRT 1.13)	0.408 (RRT 1.12)	0.389 (RRT 1.12)	0.432 (RRT 1.12)
PSD (Dv50, µm)	2.282	2.069	2.089	2.133

Table 4 - Stability of variant 1a under conditions: 40°C/75% RH

Test	Initial	1 month	2 months
Appearance	Light yellow suspension free from agglomerates	Light yellow suspension free from agglomerates	Light yellow suspension free from agglomerates
pH	6.55	6.91	6.67
Assay (mg/g)	0.372	0.374	0.377
Impurities (% area)			
Total	1.8	1.7	1.6
Greatest	0.490 (RRT 0.28)	0.445 (RRT 0.26)	0.428 (RRT 0.26)
Second greatest	0.401 (RRT 1.13)	0.409 (RRT 1.12)	0.380 (RRT 1.12)
PSD (Dv50, µm)	2.28	2.10	2.14

[0047] Variant 1b was stored under conditions of either (i) a temperature of 25°C and a relative humidity (RH) of 60% or (ii) a temperature of 40°C and a RH of 75%. The appearance, pH, impurities and particle size distribution (PSD) were measured initially, at 1 month, and at 2 months. The amount of impurities were measured by HPLC with the relative retention time (RRT) of the impurity indicated.

[0048] The results are shown in Tables 5 and 6 below.

Table 5 - Stability of variant 1b under conditions: 25°C/60% RH

Test	Initial	1 month	2 months
Appearance	Yellow suspension free from agglomerates	Yellow suspension free from agglomerates	Yellow suspension free from agglomerates
pH	6.55	6.67	6.68
Assay (mg/g)	18.72	18.86	18.54
Impurities (% area)			
Total	1.2	1.2	1.2
Greatest	0.399 (RRT 1.13)	0.401 (RRT 1.12)	0.401 (RRT 1.12)
Second greatest	0.306 (RRT 1.11)	0.328 (RRT 1.11)	0.329 (RRT 1.11)
PSD (Dv50, µm)	2.11	2.37	2.05

Table 6 - Stability of variant 1b under conditions: 40°C/75% RH

Test	Initial	1 month	2 months
Appearance	Yellow suspension free from agglomerates	Yellow suspension free from agglomerates	Yellow suspension free from agglomerates
pH	6.55	6.76	6.3
Assay (mg/g)	18.72	18.99	19.01
Impurities (% area)			
Total	1.2	1.2	1.2
Greatest	0.399 (RRT 1.13)	0.394 (RRT 1.12)	0.388 (RRT 1.13)
Second greatest	0.306 (RRT 1.11)	0.308 (RRT 1.11)	0.322 (RRT 1.11)
PSD (Dv50, µm)	2.11	2.09	2.13

[0049] As can be seen from Tables 3 to 6, the liquid pharmaceutical suspension composition shows excellent long term stability with no significant variation in pH or the amount of impurities

present. This stability is even observed after 2 months at 40°C.

Comparative Example 1 - formulation and stability of RPL554 solution formulation

[0050] A similar stability test was performed for the solution formulation of RPL554. A 1.0 mg/ml solution of RPL554 in an aqueous citrate/phosphate buffer solution at a pH of approximately 3.2 was held under conditions of a temperature of 25°C and a relative humidity (RH) of 60% for six months. The appearance, pH, assay and impurities present were measured initially and at 6 months. The results are shown in Table 7 below.

Table 7 - Stability of RPL554 solution under conditions: 25°C/60% RH

Test	Initial	6 months
Appearance	A clear, slightly coloured solution. Free from visible contamination	A clear, slightly coloured solution. Free from visible contamination.
pH	3.04	3.00
Assay	0.98 mg/mL	0.98 mg/mL
Impurities by HPLC		
RRT 0.85	0.19%	0.19%
RRT 0.87	0.15%	0.42%
RRT 1.03	< 0.10%	< 0.10%
RRT 1.09	0.20%	0.16%
RRT 1.12	0.54%	0.57%
RRT 1.14	0.14%	< 0.10%
RRT 1.15	0.11%	< 0.10%

[0051] While there is no change in the appearance of the solution, there is a significant increase in the amount of the impurity with the RRT value of 0.87. This impurity has been identified as a hydrolysis product of RPL554. The solution of RPL554 is less stable than the suspension formulation.

Reference Example 2 - particle size distribution

[0052] The particle size distribution of a sample of micronized RPL554 suitable for use in nebulized formulations was evaluated using laser diffraction. The particle size distribution analysis was performed by laser diffraction using the Malvern Spraytec in conjunction with a wet dispersion cell.

[0053] The results are as follows: Dv10 = 0.69 µm; Dv50 = 1.35 µm; Dv90 = 2.5 µm.

Reference Example 3 - formulations

[0054] RPL554 suspension and solution formulations were prepared as follows. These formulations were used in Examples 4 to 6.

Formulation constituents (Suspension)	20 mg/ml RPL554 Suspension		
	Constituent	Amount	Concentration (mg/mL)
	RPL554 - Micronised	2.12g ^a	20 mg/mL
	Wetting Solution	10.0 mL	N/A
	Buffer Solution	To 100mL	N/A
Vehicle = as above, omitting RPL554			
	RPL554 Buffer Solution Constituent	Amount (g)	Concentration (mg/mL)
	Monosodium phosphate monohydrate	32.9	6.58
	Dibasic sodium phosphate anhydrous	34.0	6.80
	Sodium chloride	24.0	4.80
	Water for Injection	To 5000ml	N/A
	RPL554 Wetting Solution Constituent	Amount (g)	Concentration (mg/mL)
	Polysorbate 20 (Tween 20)	10.0	5.00
	Sorbitan monolaurate (Span 20)	1.0	0.50
	RPL554 Buffer Solution	To 2000 mL	N/A
Method of preparation (Suspension)	RPL554 Buffer Solution: The required amount of vehicle excipients were weighed out into a suitable container and made up to the required volume with water for injection. The excipients were magnetically stirred for 10 minutes until the buffer salts were fully dissolved. The pH was measured, recorded and adjusted if necessary to pH 7.0 ± 0.3 using HCl/NaOH and filtered through a 0.22 µm filter.		
	RPL554 Wetting Solution: The required amount of polysorbate 20 was weighed into a suitable container and the required amount of sorbitan monolaurate was added. The wetting solution was magnetically stirred for 10 minutes, then transferred to a larger container with buffer solution and made up to the final volume with the buffer solution. The solution was magnetically stirred for 10 minutes and filtered through a 0.22 µm filter.		
	Vehicle: The required volume of RPL554 Wetting Solution was measured into a suitable container and made to the final volume with		

	RPL554 Wetting Solution Constituent	Amount (g)	Concentration (mg/mL)
	RPL554 Buffer Solution, mixed with a magnetic stirrer for 10 minutes. The pH was measured, recorded and adjusted if necessary to pH 7.0 ± 0.3 using HCl/NaOH		
	20 mg/mL RPL554 Suspension: The required amount of RPL554 was weighed into a small beaker and the required amount of Wetting Solution was added. The suspension was mixed using a Silverson mixer fitted with a 5/8" high sheer screen for approximately 1 minute at 8000 rpm. The mixture was transferred to a new clean, dry beaker rinsing with Buffer Solution. The required weight was made up with Buffer Solution and mixed using a Silverson mixer fitted with a 1" high sheer screen for approximately 1 minute at 8000 rpm. The pH was measured, recorded and adjusted if necessary to pH 7.0 ± 0.3 using HCl/NaOH. The solution was transferred into the final container. For the lower concentrations of RPL554 suspensions, the required amount of 20 mg/mL formulation was weighed out and diluted to the required volume with vehicle using a syringe fitted with a 0.22 µm filter. The formulation was gently mixed on a magnetic stirrer for 5 minutes.		
Formulation constituents (Solution)	1 mg/mL (pH 3.5) RPL554 Solution		
	Constituent	Amount	
	RPL554	100 mg	
	0.1M Citric acid solution	22.5 mL	
	0.2M Dibasic sodium phosphate	9.7 mL	
	dodecahydrate solution		
	0.9% (w/w) Saline solution	67.8 mL	
	1.5 mg/mL (pH 2.5) RPL554 Solution		
	Constituent	Amount	
	RPL554	150 mg	
	0.1M Citric acid solution	45 mL	
	0.2M Dibasic sodium phosphate	5 mL	
	dodecahydrate solution		
	0.9% (w/w) Saline solution	50 mL	
	Vehicle: As above, omitting RPL554		
Method of preparation	RPL554 0.1M Citric Acid Solution: The required amount of citric acid was weighed into a suitable container, made up to the required volume		

	1.5 mg/mL (pH 2.5) RPL554 Solution		
	Constituent	Amount	
(Solution)	with 0.9% saline and magnetically stirred until fully dissolved.		
	RPL554 0.2M Dibasic Sodium Phosphate Solution: The required amount of dibasic sodium phosphate anhydrous was weighed into a suitable container, made up to the required volume with 0.9% saline and magnetically stirred until fully dissolved.		
	Vehicle: The required volume of Citric Acid Solution, Dibasic Sodium Phosphate Solution and 0.9% saline were measured into a suitable container and mixed thoroughly. The pH was measured and adjusted if necessary (pH 3.5 (±0.3) or pH 2.5 (±0.2)) by drop wise addition of further citric acid or sodium phosphate solution as required.		
	RPL554 Solution: The required amount of test material was weighed into a suitable container, made up to the final volume with Vehicle and magnetically stirred until fully dissolved. The pH was measured and adjusted if necessary.		
aIncludes a 6% overage for expected manufacturing losses			

Reference Example 4 - toxicokinetic profile in rats

Summary

[0055] Han Wistar rats (5/sex/group) were dosed by nose-only inhalation once daily at target doses of 2.4, 8.4 or 21.6 mg/kg/day of an RPL554 suspension according to the invention for 7 consecutive days (groups 2, 3 and 4). An additional 5 rats/sex/group received the vehicle suspension and acted as controls (group 1). The duration of dose administration was 240 minutes on each day. At the end of the treatment period, all surviving animals were euthanized and necropsied. Satellite animals (3/sex/group) were similarly dosed and bled on Day 7 for toxicokinetic analysis. An additional group of satellite animals (3/sex) were dosed by nose-only inhalation once daily at a target dose of 2.4 mg/kg/day RPL554 solution for comparison with the RPL554 suspension and bled on Day 7 for toxicokinetic analysis (group 5). Mean aerosol concentrations and achieved doses are presented in Table 8 below.

Table 8 - Mean Aerosol Concentrations and Achieved Doses

Group	Type of Formulation	Target exposure level (µg/L)	Achieved exposure level (µg/L)	Target dose (mg/kg/day)	Achieved dose (mg/kg/day)	Formulation concentration (mg/mL)
1	Vehicle Suspension	0	0.00	0	0	0
2	Suspension	13	18.10	2.4	3.27	1.5
3	Suspension	47	55.18	8.4	9.96	4.5

Group	Type of Formulation	Target exposure level (µg/L)	Achieved exposure level (µg/L)	Target dose (mg/kg/day)	Achieved dose (mg/kg/day)	Formulation concentration (mg/mL)
4	Suspension	121	141.51	21.6	25.49	12
5	Solution	13	14.47	2.4	2.62	1.6

Toxicokinetic results

[0056] The toxicokinetic profile of the RPL554 suspension was measured at different doses (groups 2, 3 and 4). This was compared with the toxicokinetic profile of the known RPL554 solution (group 5). The dosages administered to group 2 and group 5 are similar. Group 1 is the control group.

[0057] The toxicokinetic parameters of RPL554 in suspension (invention) or solution (comparative) form were evaluated on day 7 of 1 week of daily inhalation exposure of the RPL554 compositions to rats. The results are shown in Tables 10 and 11 below. The parameters measured are maximum concentration of RPL554 in the blood plasma of the rats (C_{max}), the time taken to reach C_{max} (T_{max}) and the area under the dose curve after 8 hours (AUC_{8h}).

Table 10 - toxicokinetic results for male rats

Group	Achieved dose (mg/kg/day)	C_{max} (ng/mL)	T_{max} (hours, median)	AUC_{8h} (ng.h/mL)
1 (vehicle)	0	0	0	0
2 (suspension)	3.27	10.5	4.25	41.9
3 (suspension)	9.96	44.1	4.25	172
4 (suspension)	25.49	94.1	4.083	372
5 (solution)	2.62	53.4	4.083	139

Table 11 - toxicokinetic results for female rats

Group	Achieved dose (mg/kg/day)	C_{max} (ng/mL)	T_{max} (hours, median)	AUC_{8h} (ng.h/mL)
1 (vehicle)	0	0	0	0
2 (suspension)	3.27	19.9	4.25	79.1
3 (suspension)	9.96	133	4.25	478
4 (suspension)	25.49	573	4.25	2030
5 (solution)	2.62	52.0	4.083	153

[0058] These results demonstrate that it is possible to reach substantially higher $C_{\max}$ values using the liquid pharmaceutical composition (e.g groups 3 and 4) compared with a solution composition (group 5). The solubility limited dose feasible from a solution of RPL554 prevents such $C_{\max}$ values being achieved with a solution composition. The suspension compositions are also shown to have a delayed release, with greater $T_{\max}$ values observed for groups 2 to 4.

Reference Example 5 - toxicokinetic profile in dogs

[0059] The toxicokinetic profile of RPL554 suspensions were evaluated for beagle dogs. The study design was as set out in Table 12 below.

Table 12 - study design, doses and nominal concentrations

Phase	Dose	Treatment	Formulation	Target dose of RPL554 (mg/kg/day)	Target aerosol concentration* $\mu\text{g/L}$	Nominal concentration (mg/mL)
1	1	RPL554	Suspension	1	19.8	1.5
1	2	RPL554	Suspension	3	59.4	4.5
1	3	RPL554	Suspension	9	178	16
2		RPL554	Suspension	9	178	16
3		RPL554	Suspension and solution	0.5	10	1.5 (suspension) 1.0 (solution)
* Aerosol concentration calculated assuming an exposure duration of 120 minutes and a body weight of 12 kg						

[0060] Animals received the test substance, RPL554, by inhalation administration for 3 days at each dose in the variable dose phase (Phase 1), with at least 2 days washout period between doses. In the constant dose phase (Phase 2), animals received the test substance for 7 consecutive days. In the crossover phase (Phase 3), animals received the test substance as a single exposure to the suspension formulation followed by 2 days off-dose followed by a single exposure to the comparative solution formulation.

Toxicokinetic results

[0061] Following 7 days of dosing at 9.82 mg/kg/day with the RPL554 suspension (group 2), the $C_{\max}$ values for males and females were 116 ng/mL and 95.9 ng/mL, respectively, and the AUC_{6h} values for the males and females were 389 ng.h/mL and 289 ng.h/mL, respectively.

[0062] After a single exposure of RPL554 suspension at 0.395 mg/kg, the $C_{\max}$ values for the male and females were 13.4 ng/mL and 12.7 ng/mL, respectively, and the AUC_{6h} values for the

male and females were 33.9 ng.h/mL and 31.5 ng.h/mL, respectively.

[0063] After a single exposure of RPL554 solution at 0.543 mg/kg, the $C_{\max}$ values for the male and females were 40.9 ng/mL and 34.2 ng/mL, respectively, and the AUC_{6h} values for the male and females were 74.6 ng.h/mL and 62.8 ng.h/mL, respectively.

[0064] Here it is observed that it is possible to achieve high $C_{\max}$ and AUC values using the RPL554 suspension (e.g. 116 ng/mL and 95.9 ng/mL).

[0065] The toxicokinetic results also confirm that the suspension formulation gives a delayed release compared to the solution formulation. This can be seen from the data from the phase 3 experiment (separate single doses of the suspension formulation). These data are presented in Table 13 below, where the solution formulation (target dose 0.5 mg/kg/day, actual 0.543 mg/kg/day) is compared with the suspension formulation (target dose 0.5 mg/kg/day, actual 0.395 mg/kg/day). Table 13 shows plasma concentrations (ng/mL) over time following treatment with a single dose of the suspension or a single dose of the solution. The two treatments were performed two days apart. Data for all three dogs are presented: one male dog (61) and two female dogs (62 and 64). BLQ refers to plasma concentrations of less than 1.00 ng/mL.

Table 13 - RPL554 plasma concentration values (ng/mL) for 3 dogs following a dose with an RPL554 suspension and an RPL554 solution

Dog	61 (M)		62 (F)		64 (F)	
Time (hours)	Suspension	Solution	Suspension	Solution	Suspension	Solution
0	BLQ	BLQ	BLQ	BLQ	BLQ	BLQ
0.017	11	39.7	9.83	35.2	13.5	33.1
0.25	13.4	40.9	11.6	26.5	13.7	26.5
0.667	10.4	13	7.26	15.7	10.8	13
1.667	4.57	4.84	2.51	3.31	4.99	3.15
3	2.88	1.23	1.87	1.99	2.42	1.21
4	1.81	BLQ	1.4	BLQ	1.74	BLQ

[0066] The data in Table 13 demonstrates that the suspension formulation of RPL554 leads to a more delayed release profile than the solution formulation of RPL554. For instance, the plasma concentration of RPL554 is below the detectable limit 4 hours after inhalation of the solution formulation for all dogs, whereas at least 1.4 ng/mL of RPL554 is present after 4 hours in the blood plasma for all dogs following inhalation of the suspension.

Reference Example 6 - lung function in guinea pigs

Introduction

[0067] RPL554 causes bronchoprotection against intravenously administered spasmogenic agents when administered as dry power or when prepared as a solution in acidified (pH 2.5) saline.

[0068] This functional antagonism (1.5 doubling dilution ie 2.8 fold) against methacholine has also been verified in human asthmatic subjects (Franciosi *et al.*, 2013). This Example evaluated the bronchoprotection afforded by RPL554 when presented as the new suspension formulation for nebulisation.

Methodology

Drug exposure

[0069] Male Dunkin Hartley guinea-pigs (300-400 g) were placed in a custom built aerosol chamber and exposed to either vehicle (0.9% saline pH 2.5), RPL554 (2.5 mg/mL solution) or RPL554 (1 mg/mL solution) for a 15 min period. Aerosols were generated using an Ultrasonic Nebuliser (Ultraneb 99) and a flow rate of 1 L/min was used to direct the nebulized solution to an aerosol chamber which contained 4 arms that allows animals to be restrained with their upper respiratory tract (ie nose) protruding into the aerosol chamber.

[0070] After this 15 min exposure period, animals were allowed to recover and then prior to the 2 h and 5 h post drug exposure, animals were anaesthetised with urethane (1.5 g/kg) and a midline incision made to expose the trachea. A tracheostomy was performed, and a tracheal cannula tied in place and attached to a pneumotachograph which in turn was connected to a Validyne pressure transducer (+ 2 cmH₂O) for the detection of flow. Breath by breath changes in airflow were measured using a Lung Function Recording system (LFR, Version 9, Mumed UK) and displayed in real time on a PC. The flow signal was integrated to give a measure of tidal volume.

[0071] A cannula was inserted into the thoracic cavity between the 3rd and 5th intercostal space and connected to the negative side of a Validyne pressure transducer (+ 20 cmH₂O). The positive side of the pressure transducer was connected to the side of the pneumotachograph proximal to the animal, in order to obtain a measure of transpulmonary pressure (TPP: difference between mouth and thoracic pressure). The lung function parameters, total airway resistance (RL; cmH₂O.s/L) and dynamic lung compliance (mL/cmH₂O) was derived from each measure of flow, tidal volume and TPP by the method of integration. The carotid artery was cannulated for the measurement of blood pressure, respectively. Cumulative concentration-effect curves were established and the provocation concentration (PC) of acetylcholine which cumulatively produced a 50% increase in RL (PC₅₀).

Acetylcholine exposure

[0072] Bronchoconstriction to acetylcholine (0.25 to 16 mg/mL, 4 sec exposure) was monitored 2 h and 5 h post administration of a solution of RPL554 (1 and 2.5 mg/mL) or vehicle (acidified saline pH 3.5 or pH 2.5). Aerosols of acetylcholine were generated with an Aeroneb® Lab Nebulizer (Aerogen Inc). The inflow was directed either via the pneumotachograph for the measurement of respiratory lung mechanics, or shunted through the nebulizer for the purposes of delivering drug directly into the lung (respiratory mechanics not measured during the 4 sec exposure period).

[0073] In a separate series of experiments, lung function was performed 2 and 5 h post inhalation of the new suspension formulation of RPL554 (2.5 mg/mL and 10 mg/mL) and the vehicle.

Results

Baseline parameters

[0074] Baseline total lung resistance (RL), dynamic lung compliance (C_{dyn}), mean arterial blood pressure (BP) and nebulizer output were measured in animals previously exposed to vehicle control (saline pH 2.5), and solution formulations of RPL554 (1 mg/mL or 2.5 mg/mL, Figure 1) and to new suspension formulations of RPL554 (2.5 mg/mL and 10 mg/mL, Figure 2). Baseline respiratory and cardiovascular parameters were made prior to acetylcholine exposure and were monitored 2 and 5 h post saline or RPL554 exposure in anaesthetised guinea-pigs.

Lung function parameters

[0075] In general RL represents changes in airway diameter in the central airways following the activation of muscarinic receptors on airway smooth muscles in this region of the lung. Acetylcholine was chosen for its short duration of action.

1. RPL554 Solution Formulation

[0076] Acetylcholine administered to anaesthetized guinea-pigs by the inhaled route caused a dose dependent bronchoconstrictor response as determined by changes in RL measured 2 h and 5 h post saline and RPL554 (1 mg/mL, 2.5 mg/mL solution) exposure.

1.1 - Analysis of Acetylcholine RL PC50

[0077] Post hoc analysis of the arithmetic mean demonstrated a significant bronchoprotection at the 2 h point for RPL554 1 mg/mL solution (mean diff = 3.5 (0.11 - 8.83) mg/mL, P = 0.0445) and 2.5 mg/mL at the 2 h (4.7 (0.4 - 9) mg/mL, P = 0.0357) and 5 h time point (3.4 (0.9 - 6) mg/mL, P

= 0.0133). The degree of bronchoprotection was less evident at 5 h for 1 mg/mL RPL554 solution (2.5 (-0.17-5.2) mg/mL, $P = 0.0628$).

[0078] A significant difference between 2 h (1 mg/mL RPL554, $P = 0.0190$ unadjusted) and 5 h (2.5 mg/mL, $P = 0.0095$ unadjusted) RPL554 solution vs saline group was observed (Figure 1).

2. New Suspension Formulation of RPL554

[0079] Acetylcholine administered to anaesthetized guinea-pigs by the inhaled route caused a dose dependent bronchoconstriction as determined by changes in RL measured 2 h and 5 h post saline and RPL554 suspension (2.5 mg/mL and 10 mg/mL) exposure. Dose dependent reduction in baseline lung compliance (Cdyn) was also simultaneously recorded following increasing concentrations of acetylcholine 2 h and 5 h post saline or RPL554 suspension (2.5 mg/mL and 10 mg/mL) exposure.

2.1 - Analysis of Acetylcholine RL PC50

[0080] Analysis of variance of the log2 transformed data for Acetylcholine RL PC50 revealed an overall drug effect at the 2 h ($P = 0.0238$). For the 2 h time group, this was reflected by a significant difference between the 10 mg/mL RPL554 suspension and vehicle group (Mean fold difference = 2.67 (1.12 - 6.36), $P < 0.05$). Despite the greater mean difference, between RPL554 (10 mg/mL suspension) versus vehicle at the 5 h time point 14 fold shift (0.5-400), the wide confidence interval is the reason why statistical difference was not achieved (Figure 2, upper panel).

[0081] The vehicle data for the 2 and 5 h time points were combined and re-analyzed (Figure 2 lower panel). There was an overall significant treatment effect ($P=0.0016$) which was reflected by a significant degree of bronchoprotection with 2.5 mg/mL RPL554 suspension (2.52 fold (1-5.72), $P < 0.05$ and 10 mg/mL RPL554 (4.67 fold (2-11), $P < 0.05$). At the 5 h time point, RPL554 10 mg/mL suspension caused a significant degree of bronchoprotection (2.77 fold (1.3-6.0) (Figure 2, lower panel).

Summary

[0082] RPL554 inhibited bronchoconstriction induced by aerosolized administered acetylcholine when measured 2 and 5 h post drug exposure. Both the RPL554 solution formulations (1 mg/mL, 2.5 mg/mL) and the new RPL554 suspension formulations (2.5 mg/mL, 10 mg/mL) produced a statistically significant bronchoprotection.

Reference Example 7 - pharmacokinetic study in humans

[0083] A phase I, randomised, double-blind, placebo-controlled, study was performed to assess the safety, tolerability and pharmacokinetics of single inhaled doses of RPL554 suspension formulation, administered by nebuliser to healthy male subjects aged 18 to 50 inclusive.

[0084] The nominal dose of suspension formulation was approximately 1.5 mg, being essentially equivalent to the 0.018 mg/kg dose previously shown to be well-tolerated in clinical studies with the solution formulation.

[0085] The suspension formulation used was a sterile suspension for nebulisation containing 1.5 mg micronised RPL554 drug substance in 5 mL phosphate buffered saline solution at pH 7, with surfactants (Tween 20 and Span 20) (as described in Example 1 above, but with a different concentration of RPL554). The dosing of the suspension was performed using a standard Jet nebuliser (PARI LC Sprint).

[0086] Blood samples (4mL at each time point) were collected at appropriate time intervals after administration. Samples were collected by venepuncture or via indwelling cannula in the forearm into lithium heparin tubes and will be immediately chilled (ice bath). The blood was centrifuged within 15 minutes of collection. The plasma was separated in a refrigerated centrifuge (about 4°C) at 1100g for 15 minutes and transferred into polypropylene tubes. The concentration of the active was then measured.

[0087] The mean results are presented in Figure 3, which shows the mean concentration curves for the subjects after administration of the solution or suspension formulations. The results for the solution formulation were derived from a previous clinical study (The Lancet Respiratory Medicine Volume 1, No.9, p.714-727, November 2013) which used a sterile solution for nebulisation containing RPL554 dissolved in citrate-phosphate buffered saline at pH 3.2. The quantity administered via nebuliser was such that the nominal RPL554 dose was 0.018 mg/kg (~1.26 mg for a 70kg person).

[0088] Mean pharmacokinetic parameters measured during the study are presented in Table 14 below.

Table 14

Formulation	C _{max} (pg/ml)	AUC _{inf} (pg.h/ml)	Half-life (h)
Solution	870	1833	3
Suspension	393	3795	10

[0089] The values in Table 14 demonstrate that the suspension formulation has a substantially improved pharmacokinetic profile compared with the comparative solution formulation. In particular, the area under the curve (AUC) and the half-life are both significantly increased. This demonstrates that the suspension formulation provides an unexpectedly improved way by which RPL554 may be administered by inhalation.

Reference Example 8 - twelve months stability study

[0090] Three batches of the suspension formulation of the invention were made with the compositions set out in Table 15 below.

Table 15

Constituent	Concentration (mg/mL)
RPL554 (heat treated API)	0.5, 2.5 and 10
Polysorbate 20 (Tween 20)	0.5
Sorbitan monolaurate (Span 20)	0.05
Monosodium phosphate monohydrate	6.58
Dibasic sodium phosphate anhydrous	6.80
Sodium chloride	4.80
Water for injection	To weight

Analytical testing

[0091] Analytical testing was performed after 1, 3, 6 and 12 months storage of the formulations at 25°C and 60% RH. The 12 months results are described below.

Appearance

[0092] Appearance testing was performed visually on a single sample. The placebo sample was found to be a clear, colourless solution. The 0.5mg/mL strength samples were a light yellow suspension free from visible agglomerates while the 2.5mg/mL and 10mg/mL strength samples appeared to be yellow suspensions free from visible agglomerates.

pH

[0093] The pH determination was performed on a single sample. The results are shown in Table 16.

Table 16

Batch Number	Strength	pH
BN011/14	0.5mg/mL	6.656
BN010/14	2.5mg/mL	6.652
BN009/14	10mg/mL	6.655
BN008/14	Placebo	6.661

[0094] All three active formulations and placebo were found to have a pH of 6.7 at the twelve month time-point. No change was observed from the initial time point.

Assay

[0095] Assay testing was performed in duplicate. The key method parameters are summarised as follows:

Mobile phase	Acetonitrile: water: TFA 45:55:0.1
Column	Waters X-Bridge phenyl, 3.5 µm, 150 × 4.6 mm, (Part Number 186003335)
Column temperature	40°C
Flow rate	1.5 ml/min
Injection volume	10 µl
Detection	UV, 254 nm
Runtime	6 minutes
Autosampler temperature	Ambient
Standard Concentration	0.1 mg/ml

[0096] The results of the assay are shown in Table 17.

Table 17				
Batch Number	Strength	Prep 1	Prep 2	Mean
BN011/14	0.5mg/mL	0.5180	0.5178	0.52
BN010/14	2.5mg/mL	2.6541	2.6407	2.65
BN009/14	10mg/mL	10.4219	10.4926	10.46

[0097] The assay results for the active suspensions of each strength at the twelve month time point did not show any change from initial time point.

Related substances

[0098] Related substances determination was performed in duplicate. The key method parameters are summarised as follows.

Mobile phase	A - Purified water/Acetonitrile/TFA (95/5/0.1).
	B - Acetonitrile/Water/TFA (95/5/0.1).

Column	X-Bridge Phenyl 4.6 x 150 mm 3.5 µm particle size (ex - Waters; #186003335)		
Column temperature	30°C		
Autosampler Temperature	Ambient		
Flow rate	1.0 ml/min		
Injection volume	10 µl		
Detection	UV, 254 nm		
Gradient	Time (min)	%A	%B
	0	100	0
	2	100	0
	15	0	100
	25	0	100
	27	100	0
	37	100	0
Diluent	Acetonitrile : Water (50 : 50)		
Sample Concentration	0.2mg/ml		
Standard Concentration	2 µg/ml		

[0099] The results from the related substances determination of the suspensions are presented in Table 18.

Table 18 - Results from the related substances analysis of RPL554 suspensions (as %LC)

Batch BN011/14 - (0.5mg/mL) - 25°C/60%RH			
RRT	%LC		Mean
	T1	T2	
0.78	0.0776	0.0764	0.08
0.86	0.1028	0.1046	0.10
0.92	0.2084	0.2101	0.21
0.93	0.1875	0.1846	0.19
1.09	0.0548	0.0562	0.06
1.11	0.4130	0.4142	0.41
Total			1.05
Batch BN010/14-(2.5mg/mL) - 25°C/60%RH			
RRT	%LC		Mean
	T1	T2	
0.77	0.0779	0.0792	0.08
0.85	0.1041	0.1051	0.10

Batch BN010/14-(2.5mg/mL) - 25°C/60%RH			
RRT	%LC		Mean
	T1	T2	
0.92	0.2184	0.2227	0.22
0.93	0.1768	0.1746	0.18
1.11	0.4226	0.4270	0.42
Total			1.00
Batch BN009/14 - (10mg/mL) - 25°C/60%RH			
RRT	%LC		Mean
	T1	T2	
0.78	0.0769	0.0789	0.08
0.86	0.1016	0.1031	0.10
0.92	0.2056	0.2194	0.21
0.93	0.1819	0.1684	0.18
1.11	0.4137	0.4250	0.42
Total			0.99

[0100] The related substances results for the active suspensions of each strength at the twelve month time-point were similar to the initial time point.

Particle size distribution (PSD)

[0101] The Particle Size Distribution method was performed using a Spraytec with the settings detailed below.

Optical Properties: Particle: RI (real): 1.50 (standard opaque particle); RI (imaginary): 0.50; Density: 1.00; Dispersant (water); Dispersant RI: 1.33

Dispersant used for sample analysis: 1% polysorbate 20 in deionised water

Controller unit settings: Stirrer speed 1000rpm

Measurement time: Sampling time: 30s; Background time: 10s

[0102] The results are shown in Table 19.

Table 19 - Results from the Particle Size Determination by Laser Diffraction of RPL554 suspensions (in µm)

Batch Number	Strength	Dv10	Dv50	Dv90
BN011/14	0.5 mg/mL	0.7946	1.577	3.168
BN010/14	2.5mg/mL	0.7872	1.589	3.119
BN009/14	10mg/mL	0.8012	1.566	2.976

[0103] The Particle Size Distribution profiles were similar for all 3 strengths at the twelve month time-point and did not show any change compared to the initial time point.

Microscopic evaluation

[0104] Microscopic evaluation was performed by visually assessing the formulations using a G3 microscope. No aggregates were observed in any of the batches tested, although some large loose agglomerates were occasionally seen for the 2.5mg/mL and 10mg/mL suspensions. The evaluation determined that there was no change compared to the initial time point.

Delivery rate and total delivered dose

[0105] Delivery rate and total delivered dose was performed on 3 vials per batch. The suspensions were dispensed and delivered using a PARI LC PLUS® Nebuliser / PARI TurboBOY S® compressor combination. The results from the delivery rate and total delivered dose are presented in Table 20.

Table 20 - Results from the delivery rate and total dose delivered determination using PARI LC PLUS nebuliser

Batch BN011/14 0.5mg/mL			
Replicate number	T1	T2	T3
Initial weight (g)	36.8175	36.7852	36.8030
Final weight (g)	32.6399	32.7369	32.6272
Delivered mass (g)	4.1776	4.0483	4.1758
Total Delivery Time (s)	900	1020	960
Delivery Rate (mg/min) (over the first min)	0.136	0.130	0.137
Mean Delivery Rate (mg/min)	0.1		
Assay* (mg)	1.179	1.144	1.198
Mean Assay* (mg)	1.2		
% Efficiency	54.50	54.56	55.39
Mean % Efficiency	54.8		

Batch BN010/14 2.5mg/mL			
Replicate number	T1	T2	T3
Initial weight (g)	37.0977	38.7543	37.2942
Final weight (g)	32.8590	34.9311	32.9187
Delivered mass (g)	4.2387	4.3632	4.3755
Total Delivery Time (s)	1020	1020	1020
Delivery Rate (mg/min) (over the first min)	0.303	0.229	0.335
Mean Delivery Rate (mg/min)	0.3		
Assay* (mg)	6.212	6.684	6.565
Mean Assay* (mg)	6.5		
% Efficiency	55.36	57.86	56.68
Mean % Efficiency	56.6		
Batch BN009/14 10mg/mL			
Replicate number	T1	T2	T3
Initial weight (g)	36.9365	38.4396	37.0922
Final weight (g)	32.6608	34.2031	32.7858
Delivered mass (g)	4.2757	4.2365	4.3064
Total Delivery Time (s)	1080	1140	1080
Delivery Rate (mg/min) (over the first min)	1.326	1.095	1.221
Mean Delivery Rate (mg/min)	1.2		
Assay* (mg)	24.279	24.168	24.705
Mean Assay* (mg)	24.4		
% Efficiency	54.30	54.55	54.86
Mean % Efficiency	54.6		
* assay is total delivered % Efficiency is total actual dose delivered/theoretical dose delivered (calculated using delivered mass and formulation strength from assay)			

[0106] The total delivered dose and delivery rate for all strengths were similar to the initial time point. The delivery efficiency of all the suspensions was consistent with previous time points.

Aerodynamic Particle Size Distribution (APSD)

[0107] Aerodynamic Particle Size Distribution (APSD) determination was performed on 3 vials per

batch per condition using the Next Generation Impactor (NGI). The suspensions were dispensed and delivered using a PARI LC PLUS® Nebuliser / PARI TurboBOY S® compressor combination. The results from the APSD determination are presented in Table 21. The results were input into the CITDAS program (Version 3.10) to calculate the fine particle dose using a cut-off value of 5µm and the results obtained from the calculations are also presented Table 21.

Table 21 - Results from the Aerodynamic Particle Size Distribution by NGI (as mg RPL554) using PARI LC PLUS nebuliser

Batch BN011/14 - (0.5mg/mL) 25°C/60%RH				
Stage	NGI 1	NGI 2	NGI 3	Mean
Throat	0.05495	0.09324	0.08583	0.08
Stage 1	0.13433	0.12798	0.12692	0.13
Stage 2	0.26114	0.18827	0.22637	0.23
Stage 3	0.40347	0.29208	0.39028	0.36
Stage 4	0.49538	0.41178	0.46929	0.46
Stage 5	0.23507	0.24668	0.29097	0.26
Stage 6	0.07792	0.06313	0.08504	0.08
Stage 7	0.01105	0.01100	0.01525	0.01
MOC	0.00034	0.00169	0.00151	0.00
Sum	1.67365	1.43585	1.69147	1.60
Delivered Mass (g)	4.2864	4.1465	4.2623	4.23
FPD ≤ 5µm*	0.735	0.667	0.785	0.7
FPD/delivered mass (mg/g)	0.171	0.161	0.184	0.2
FPF (FPD as % total dose) ≤ 5µm*	43.9	46.4	46.4	45.6
GSD*	2.0	2.2	2.0	2.1
MMAD*	5.3	5.0	5.1	5.2
Batch BN010/14 - (2.5mg/mL) 25°C/60%RH				
Stage	NGI 1	NGI 2	NGI 3	Mean
Throat	0.35731	0.43867	0.47848	0.42
Stage 1	0.71616	1.04125	0.93292	0.90
Stage 2	1.28121	1.58441	1.43900	1.43
Stage 3	1.98259	2.21051	1.95403	2.05
Stage 4	2.41709	2.37565	2.23993	2.34
Stage 5	1.69766	1.45824	1.39429	1.52
Stage 6	0.54927	0.43922	0.37542	0.45
Stage 7	0.07994	0.07293	0.06443	0.07
MOC	0.00163	0.00151	0.00540	0.00
Sum	9.08286	9.62239	8.88390	9.20

Batch BN010/14 - (2.5mg/mL) 25°C/60%RH				
Stage	NGI 1	NGI 2	NGI 3	Mean
Delivered Mass (g)	4.3881	4.5659	4.3137	4.42
FPD $\leq 5\mu\text{m}^*$	4.356	3.945	3.702	4.0
FPD/delivered mass (mg/g)	0.993	0.864	0.858	0.9
FPF (FPD as % total dose) $\leq 5\mu\text{m}^*$	48.0	41.0	41.7	43.5
GSD*	2.1	2.1	2.1	2.1
MMAD*	5.0	5.7	5.5	5.4
Batch BN009/14 - (10mg/mL) 25°C/60%RH				
Stage	NGI 1	NGI 2	NGI3	Mean
Throat	1.82141	1.46478	1.70097	1.66
Stage 1	2.99033	3.28692	3.75069	3.34
Stage 2	5.51909	6.88069	7.01532	6.47
Stage 3	8.46593	10.48679	10.09839	9.68
Stage 4	9.42726	10.16201	9.36013	9.65
Stage 5	6.30967	4.85282	4.67990	5.28
Stage 6	1.84396	0.98008	1.00599	1.28
Stage 7	0.28335	0.13666	0.12275	0.18
MOC	0.00383	0.00710	0.00108	0.00
Sum	36.66483	38.25785	37.73522	37.55
Delivered Mass (g)	4.3787	4.6309	4.4044	4.47
FPD $\leq 5\mu\text{m}^*$	16.313	14.318	13.490	14.7
FPD/delivered mass (mg/g)	3.726	3.092	3.063	3.3
FPF (FPD as % total dose) $\leq 5\mu\text{m}^*$	44.5	37.4	35.7	39.2
GSD*	2.0	1.9	1.9	2.0
MMAD*	5.3	5.9	6.1	5.8
* calculated using CITDAS on unrounded data. Abbreviations: FPD: fine particle dose; FPF: fine particle fraction; GSD: geometric size distribution; MMAD: mass median aerodynamic diameter; MOC: micro-orifice collector.				

12 months stability

[0108] Tables 22 (for 0.5 mg/mL), 23 (for 2.5 mg/mL) and 24 (for 10 mg/mL) below combine the

12 months data described above with the corresponding data for the 1, 3 and 6 months time points for the RPL554 suspension formulations of the invention.

[0109] (Note: Impurity peaks marked with * in Tables 22, 23 and 24 were observed as a doublet but had been observed as a single peak at the initial time point. This is probably due to a new column being used which gave better separation and is not believed to be a sign of degradation.)

Summary

[0110] RPL554 Suspensions for Nebulisation 0.5 mg/mL (BN011/14), 2.5 mg/mL (BN010/14) and 10 mg/mL (BN009/14) and associated placebo (BN008/14) were manufactured and placed on stability at 25°C/60%RH and were tested at the 12 month time-point.

[0111] The results showed that there was essentially no change in any of the batches tested at the 12 month time point and confirmed that the RPL554 suspensions for nebulisation were stable for 12 months at 25°C/60%RH.

Table 22 (0.5 mg/mL) Test	Initial	1 month	3 month	6 month	12 month
Storage condition	N/A	25°C/60%RH	25°C/60%RH	25°C/60%RH	25°C/60%RH
Appearance	Pale yellow suspensions free from visible agglomerates	Pale yellow suspensions free from visible agglomerates	Pale yellow suspensions free from visible agglomerates	Pale yellow suspensions free from visible agglomerates	Pale yellow suspensions free from visible agglomerates
pH	6.7	6.7	6.6	6.6	6.7
Mean Assay (n=2) (mg/mL)	0.51	0.51	0.52	0.52	0.52
Mean Impurities (n=2) (%LC)					
Total	0.98	1	0.89	0.95	1.05
Greatest	0.39 (RRT 1.11)	0.23 (RRT 0.92)	0.38 (RRT 1.10)	0.38 (RRT 1.10)	0.41 (RRT 1.11)
Second greatest	0.23 (RRT 0.92)	0.20 (RRT 1.11)* 0.18 (RRT 1.10)*	0.22 (RRT 0.92)	0.22 (RRT 0.93)	0.21 (RRT 0.92)
PSD (µm)	Dv10= 0.7646	Dv10= 0.7721	Dv10= 0.7823	Dv10= 0.8036	Dv10= 0.7946
	Dv50= 1.534	Dv50= 1.542	Dv50= 1.56	Dv50= 1.585	Dv50= 1.577

Table 22 (0.5 mg/mL) Test	Initial	1 month	3 month	6 month	12 month
Storage condition	N/A	25°C/60%RH	25°C/60%RH	25°C/60%RH	25°C/60%RH
	Dv90= 2.999	Dv90= 2.991	Dv90= 3.084	Dv90= 3.158	Dv90= 3.168
Mean FPD (mg)	0.4	0.6	0.6	0.6	0.7
Mean FPD/mass delivered (n=3) (mg/g)	0.1	0.2	0.1	0.1	0.2
Mean MMAD (n=3)	5.4	5.7	5.4	5.6	5.2
Mean GSD (n=3)	2	2	2.1	2	2.1
Mean vial fill weight (n=5) (g)	5.0 (n=10)	5.1	5.1	5.1	5.1
Mean Total delivered dose (n=3) (mg)	1.1	1.1	1.2 (n=5)	1	1.2
Mean Delivery rate (n=3) (mg/min) (over first two min)	0.1	0.1	0.1 (n=5)	0.1	0.1
Mean % Efficiency (n=3)	53.8	55.5	56.0 (n=5)	51.4	54.8
Table 23 (2.5 mg/mL) Test	Initial	1 month	3 month	6 month	12 month
Storage condition	N/A	25°C/60%RH	25°C/60%RH	25°C/60%RH	25°C/60%RH
Appearance	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates
pH	6.7	6.7	6.7	6.6	6.7

Table 23 (2.5 mg/mL) Test	Initial	1 month	3 month	6 month	12 month
Storage condition	N/A	25°C/60%RH	25°C/60%RH	25°C/60%RH	25°C/60%RH
Mean Assay (n=2) (mg/g)	2.6	2.57	2.64	2.63	2.65
Mean Impurities (n=2) (%LC)					
Total	1.02	0.92	0.89	0.98	1
Greatest	0.41 (RRT 1.10)	0.23 (RRT 0.92)	0.39 (RRT 1.10)	0.41 (RRT 1.10)	0.42 (RRT 1.11)
Second greatest	0.23 (RRT 0.92)	0.20 (RRT 1.11)* 0.18 (RRT 1.10)*	0.22 (RRT 0.93)	0.23 (RRT 0.93)	0.22 (RRT 0.92)
PSD (µm)	Dv10= 0.7964	Dv10= 0.7981	Dv10=0.7943	Dv10=0.7851	Dv10=0.7872
	Dv50= 1.586	Dv50= 1.569	Dv50= 1.561	Dv50= 1.583	Dv50= 1.589
	Dv90= 3.075	Dv90= 3	Dv90=2.98	Dv90=3.099	Dv90=3.119
Mean FPD (mg)	3	3.1	3.4	3.5	4
Mean FPD/mass delivered (n=3) (mg/g)	0.7	0.7	0.8	0.8	0.9
Mean MMAD (n=3)	6.0	6.1	5.8	6	5.4
Mean GSD (n=3)	1.9	1.9	1.9	2	2.1
Mean Vial Content Uniformity (n=10)	Pass EP 2.9.6 and 2.9.40	N/A	N/A	N/A	N/A
Mean vial fill weight (n=5) (g)	5.1 (n=10)	5.2	5.2	5.3	5.3
Mean Total delivered dose (n=3)	6	6.2	4.5	5.7	6.5

Table 23 (2.5 mg/mL) Test	Initial	1 month	3 month	6 month	12 month
Storage condition	N/A	25°C/60%RH	25°C/60%RH	25°C/60%RH	25°C/60%RH
(mg)					
Mean Delivery rate (n=3) (mg/min) (over the first min)	0.3	0.3	0.2	0.3	0.3
Mean % Efficiency (n=3)	56.3	55.5	41.2	55	56.6
Table 24 (10 mg/mL) Test	Initial	1 month	3 month	6 month	12 month
Storage condition	N/A	25°C/60%RH	25°C/60%RH	25°C/60%RH	25°C/60%RH
Appearance	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates
pH	6.7	6.7	6.7	6.6	6.7
Mean Assay (n=2) (mg/mL)	10.24	10.16	10.28	10.23	10.46
Mean Impurities (n=2) (%LC)					
Total	0.98	0.92	0.89	0.97	0.99
Greatest	0.40 (RRT 1.11)	0.23 (RRT 0.92)	0.39 (RRT 1.10)	0.41 (RRT 1.10)	0.42 (RRT 1.11)
Second greatest	0.23 (RRT 0.92)	0.20 (RRT 1.11)* 0.18 (RRT 1.10)*	0.22 (RRT 0.93)	0.22 (RRT 0.93)	0.21 (RRT 0.92)
PSD (µm)	Dv10= 0.7884	Dv10= 0.7877	Dv10= 0.7954	Dv10= 0.7756	Dv10= 0.8012
	Dv50= 1.564	Dv50= 1.55	Dv50= 1.545	Dv50= 1.558	Dv50= 1.566
	Dv90= 3.02	Dv90= 2.965	Dv90=2.909	Dv90=3.059	Dv90=2.976

Table 24 (10 mg/mL) Test	Initial	1 month	3 month	6 month	12 month
Storage condition	N/A	25°C/60%RH	25°C/60%RH	25°C/60%RH	25°C/60%RH
Mean FPD (mg)	12.2	12.7	14.2	13.3	14.7
Mean FPD/mass delivered (n=3) (mg/g)	2.9	3	3.1	2.2 (n=2)	3.3
Mean MMAD (n=3)	6.1	5.9	6	5.7 (n=2)	5.8
Mean GSD (n=3)	1.9	1.9	1.9	2.0 (n=2)	2
Mean Vial Content Uniformity (n=10)	Pass EP 2.9.6 and 2.9.40	N/A	N/A	N/A	N/A
Mean vial fill weight (n=5) (g)	5.2 (n=10)	5.1	5.1	5.1	5.1
Mean Total delivered dose (n=3) (mg)	25.5	23.4	26.4	22.7	24.4
Mean Delivery rate (n=3) (mg/min) (over the first min)	1.3	1.4	1.4	1.2	1.2
Mean % Efficiency (n=3)	57.3	52.8	56.7	53.5	54.6

Reference Example 9 - six month analysis at accelerated conditions (40°C/75% RH)

[0112] Accelerated stability studies at 40°C/75% RH were also continued to the sixth month. The results of these are shown in Table 25 (for 0.5 mg/mL), Table 26 (for 2.5 mg/mL) and Table 27 (for 10 mg/mL).

[0113] (As noted in Example 8, impurity peaks marked with * were observed as a doublet but had been observed as a single peak at the initial time point. This is probably due to a new column being used which gave better separation and is not believed to be a sign of degradation.)

[0114] The results showed that there was essentially no change in any of the batches tested at the 6 month time point and confirmed that the RPL554 suspensions for nebulisation have excellent stability, being stable for 6 months at 40°C/75%RH.

Table 25 (0.5 mg/mL) Test	Initial	1 month	3 month	6 month
Storage condition	N/A	40°C/75%RH	40°C/75%RH	40°C/75%RH
Appearance	Pale yellow suspensions free from visible agglomerates	Pale yellow suspensions free from visible agglomerates	Pale yellow suspensions free from visible agglomerates	Pale yellow suspensions free from visible agglomerates
pH	6.7	6.7	6.6	6.6
Mean Assay (n=2) (mg/mL)	0.51	0.51	0.52	0.52
Mean Impurities (n=2) (%LC)				
Total	0.98	0.9	0.9	0.98
Greatest	0.39 (RRT 1.11)	0.23 (RRT 0.92)	0.37 (RRT 1.10)	0.37 (RRT 1.10)
Second greatest	0.23 (RRT 0.92)	0.20 (RRT 1.11)* 0.18 (RRT 1.10)*	0.22 (RRT 0.92)	0.22 (RRT 0.93)
PSD (µm)	Dv10= 0.7646	Dv10= 0.7902	Dv10= 0.7749	Dv10= 0.8324
	Dv50= 1.534	Dv50= 1.56	Dv50= 1.586	Dv50= 1.573
	Dv90= 2.999	Dv90= 3.026	Dv90= 3.335	Dv90= 2.899
Mean FPD (mg)	0.4	0.6	0.7	0.6
Mean FPD/mass delivered (n=3) (mg/g)	0.1	0.2	0.2	0.1
Mean MMAD (n=3)	5.4	5.8	5.3	5.5
Mean GSD (n=3)	2	1.9	2	2.1
Mean Vial Content Uniformity (n=10)	Pass EP 2.9.6 and 2.9.40	N/A	N/A	N/A
Mean vial fill weight (n=5) (g)	5.0 (n=10)	5.1	5.1	5.1
Mean Total				

Table 25 (0.5 mg/mL) Test	Initial	1 month	3 month	6 month
Storage condition	N/A	40°C/75%RH	40°C/75%RH	40°C/75%RH
delivered dose (n=3) (mg)	1.1	1.1	1.2	1.1
Mean Delivery rate (n=3) (mg/min) (over first two min)	0.1	0.1	0.1	0.1
Mean % Efficiency (n=3)	53.8	55.1	54.3	53.4
Table 26 (2.5 mg/mL) Test	Initial	1 month	3 month	6 month
Storage condition	N/A	40°C/75%RH	40°C/75%RH	40°C/75%RH
Appearance	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates
pH	6.7	6.7	6.6	6.6
Mean Assay (n=2) (mg/mL)	2.6	2.58	2.64	2.63
Mean Impurities (n=2) (%LC)				
Total	1.02	0.92	0.9	0.99
Greatest	0.41 (RRT 1.10)	0.23 (RRT 0.92)	0.38 (RRT 1.10)	0.40 (RRT 1.10)
Second greatest	0.23 (RRT 0.92)	0.20 (RRT 1.11)* 0.18 (RRT 1.10)*	0.22 (RRT 0.93)	0.23 (RRT 0.93)
PSD (µm)	Dv10= 0.7964	Dv10= 0.7861	Dv10= 0.7994	Dv10= 0.7997
	Dv50= 1.586	Dv50= 1.563	Dv50= 1.577	Dv50= 1.614
	Dv90= 3.075	Dv90= 3.023	Dv90=3.058	Dv90=3.183
Mean FPD (mg)	3	3.2	3.6	3.6
Mean FPD/mass delivered (n=3) (mg/g)	0.7	0.8	0.8	0.8
Mean MMAD (n=3)	6	6	5.7	5.9
Mean GSD (n=3)	1.9	1.9	2	2
Mean Vial				

Table 26 (2.5 mg/mL) Test	Initial	1 month	3 month	6 month
Storage condition	N/A	40°C/75%RH	40°C/75%RH	40°C/75%RH
Content Uniformity (n=10)	Pass EP 2.9.6 and 2.9.40	N/A	N/A	N/A
Mean vial fill weight (n=5) (g)	5.1 (n=10)	5.3	5.3	5.3
Mean Total delivered dose (n=3) (mg)	6	6.1	5.6	6.4
Mean Delivery rate (n=3) (mg/min) (over the first min)	0.3	0.3	0.3	0.4
Mean % Efficiency (n=3)	56.3	56.1	50.5	56.3
Table 27 (10 mg/mL) Test	Initial	1 month	3 month	6 month
Storage condition	N/A	40°C/75%RH	40°C/75%RH	40°C/75%RH
Appearance	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates	Yellow suspension free from visible agglomerates
pH	6.7	6.7	6.6	6.6
Mean Assay (n=2) (mg/mL)	10.24	10.1	10.41	10.37
Mean Impurities (n=2) (%LC)				
Total	0.98	0.97	0.89	0.98
Greatest	0.40 (RRT 1.11)	0.23 (RRT 0.92)	0.38 (RRT 1.10)	0.40 (RRT 1.10)
Second greatest	0.23 (RRT 0.92)	0.20 (RRT 1.11)* 0.18 (RRT 1.10)*	0.23 (RRT 0.93)	0.23 (RRT 0.93)
PSD (µm)	Dv10= 0.7884	Dv10= 0.7906	Dv10= 0.7882	Dv10= 0.7757
	Dv50= 1.564	Dv50= 1.554	Dv50= 1.565	Dv50= 1.553
	Dv90= 3.02	Dv90= 2.971	Dv90= 3.024	Dv90= 3.015
Mean FPD (mg)	12.2	13.1	14.5	13.4
Mean FPD/mass delivered (n=3)	2.9	3.1	3.4	3.1

Table 27 (10 mg/mL) Test	Initial	1 month	3 month	6 month
Storage condition	N/A	40°C/75%RH	40°C/75%RH	40°C/75%RH
(mg/g)				
Mean MMAD (n=3)	6.1	5.8	5.8	5.7
Mean GSD (n=3)	1.9	1.9	2	2
Mean Vial Content Uniformity (n=10)	Pass EP 2.9.6 and 2.9.40	N/A	N/A	N/A
Mean vial fill weight (n=5) (g)	5.2 (n=10)	5.2	5.1	5.1
Mean Total delivered dose (n=3) (mg)	25.5	24.1	26.9	22.8
Mean Delivery rate (n=3) (mg/min) (over the first min)	1.3	1.3	1.2	1.3
Mean % Efficiency (n=3)	57.3	54.3	56.2	51.6

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO0058308A \[0003\]](#)

Non-patent literature cited in the description

- **LANCET**Respiratory Medicine 11, 2013, vol. 1, 9714-27 [\[0004\]](#)
- **CALZETTA et al.**J Pharmacol Exp Ther, 2013, vol. 346, 414-423 [\[0005\]](#)
- **COMPTON et al.**Pulmonary Pharmacology & Therapeutics, 2013, vol. 26, 562-573 [\[0005\]](#)
- The Lancet Respiratory Medicine, 2013, vol. 1, 9714-727 [\[0087\]](#)

PATENTKRAV

1. Flydende farmaceutisk sammensætning, der er egnet til indgivelse ved inhalation, og som omfatter en fortynder og en suspension af partikler af 9,10-dimethoxy-2-(2,4,6-trimethylphenylimino)-3-(N-carbamoyl-2-aminoethyl)-3,4,6,7-tetrahydro-2H-pyrimido[6,1-a]isoquinolin-4-on (RPL554) eller et farmaceutisk acceptabelt salt deraf, hvor:

partiklerne af RPL554 har en partikelstørrelsesfordeling med en Dv50 (medianpartikelstørrelse efter volumen) værdi på fra 0,2 µm til 5 µm; og

sammensætningen omfatter en phosphatbuffer ved en koncentration på fra 0 mg/mL til 5 mg/mL.

2. Flydende farmaceutisk sammensætning ifølge krav 1, hvor sammensætningen ikke omfatter en phosphatbuffer.

3. Flydende farmaceutisk sammensætning ifølge krav 1 eller krav 2, hvor sammensætningen ikke omfatter en buffer.

4. Flydende farmaceutisk sammensætning ifølge et hvilket som helst af de foregående krav, hvor partiklerne af RPL554 har en partikelstørrelsesfordeling med en Dv50-værdi på fra 0,7 µm til 2,5 µm.

5. Flydende farmaceutisk sammensætning ifølge et hvilket som helst af de foregående krav, hvor koncentrationen af partikler af RPL554 i den flydende farmaceutisk sammensætning er fra 0,01 mg/mL til 40 mg/mL.

6. Flydende farmaceutisk sammensætning ifølge et hvilket som helst af de foregående krav, der endvidere omfatter ét eller flere tensider.

7. Flydende farmaceutisk sammensætning ifølge krav 6, hvor det ene eller flere tensider er udvalgt fra ét eller flere ikke-ioniske tensider.

8. Flydende farmaceutisk sammensætning ifølge krav 6 eller krav 7, hvor det ene eller flere tensider er udvalgt fra polyoxyethylenglycolsorbitanalkylestere og sorbitanalkylestere.

9. Flydende farmaceutisk sammensætning ifølge et hvilket som helst af de foregående krav, hvor sammensætningen endvidere omfatter et

tonicitetsjusteringsmiddel, eventuelt hvor tonicitetsjusteringsmidlet er natriumchlorid.

10. Flydende farmaceutisk sammensætning ifølge et hvilket som helst af de foregående krav, hvor fortyndingsmidlet er vand.

5 **11.** Flydende farmaceutisk sammensætning ifølge et hvilket som helst af de foregående krav, der er egnet til indgivelse via forstøver.

12. Forstøver, der omfatter en sammensætning ifølge et hvilket som helst af de foregående krav.

10 **13.** Flydende farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 11 til anvendelse i behandling af menneske- eller dyrekroppen.

15 **14.** Flydende farmaceutisk sammensætning ifølge et hvilket som helst af kravene 1 til 11 til anvendelse i behandling eller forebyggelse af en sygdom eller tilstand udvalgt fra astma, allergisk astma, høfeber, allergisk rhinitis, bronchitis, emfysem, bronkiektasi, kronisk obstruktiv lungesygdom (COPD), respiratorisk distress-syndrom hos voksne (ARDS), steroidresistent astma, svær astma, pædiatrisk astma, cystisk fibrose, lungefibrose, pulmonal fibrose, interstitiel lungesygdom, hudlidelser, atopisk dermatitis, psoriasis, okulær inflammation, cerebral iskæmi, inflammatoriske sygdomme og autoimmune sygdomme.

20 **15.** Flydende farmaceutisk sammensætning til anvendelse ifølge krav 14, hvor sygdommen eller tilstanden er kronisk obstruktiv lungesygdom (COPD).

DRAWINGS




Figure 1




Figure 2




Figure 3