

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 2922877 T3**

-
- (51) Int.Cl.: *C 07 K 19/00 (2006.01)* *A 61 K 38/00 (2006.01)* *A 61 K 38/07 (2006.01)*
A 61 K 38/08 (2006.01) *A 61 K 38/26 (2006.01)* *A 61 K 38/29 (2006.01)*
C 07 K 1/107 (2006.01) *C 07 K 5/065 (2006.01)* *C 07 K 7/06 (2006.01)*
C 09 D 101/28 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2019-01-02**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-09-05**
- (86) Europæisk ansøgning nr.: **13857269.8**
- (86) Europæisk indleveringsdag: **2013-11-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2015-09-30**
- (86) International ansøgning nr.: **US2013071067**
- (87) Internationalt publikationsnr.: **WO2014081864**
- (30) Prioritet: **2012-11-20 US 201261728667 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Eumederis Pharmaceuticals, Inc., 7515 Guinevere Drive, Sugar Land, Texas 77479, USA**
- (72) Opfinder: **NESTOR, John, J., 725 Lynwood Drive, Encinitas, CA 92024, USA**
- (74) Fuldmægtig i Danmark: **Pure Ideas Ltd, Stratford-upon-Avon, Warwickshire, CV37 6QB, Storbritannien**
- (54) Benævnelse: **FORBEDREDE PEPTIDLÆGEMIDLER**
- (56) Fremdragne publikationer:
WO-A1-00/78302
WO-A1-02/098446
WO-A2-2008/052043
WO-A2-2012/158962
US-A1- 2003 202 981
US-B2- 7 390 788
KODA Y ET AL: "Synthesis and in vitro evaluation of a library of modified endomorphin 1 peptides", BIOORGANIC & MEDICINAL CHEMISTRY, PERGAMON, GB, vol. 16, no. 11, 1 June 2008 (2008-06-01), pages 6286-6296, XP022700358, ISSN: 0968-0896, DOI: 10.1016/J.BMC.2008.04.020 [retrieved on 2008-04-15]
SNEZANA SAVIC ET AL: "From conventional towards new - natural surfactants in drug delivery systems design: current status and perspectives", EXPERT OPINION ON DRUG DELIVERY, vol. 7, no. 3, 1 March 2010 (2010-03-01), pages 353-369, XP055267174, GB ISSN: 1742-5247, DOI: 10.1517/17425240903535833
ATTILA KERESZTES ET AL: "Recent Advances in Endomorphin Engineering", CHEMMEDCHEM, vol. 5, no. 8, 2 August 2010 (2010-08-02), pages 1176-1196, XP055161044, ISSN: 1860-7179, DOI: 10.1002/cmdc.201000077
COSTANTINO L ET AL: "Peptide-derivatized biodegradable nanoparticles able to cross the blood-brain barrier", JOURNAL OF CONTROLLED RELEASE, ELSEVIER, AMSTERDAM, NL, vol. 108, no. 1, 2 November 2005 (2005-

Fortsættes ...

11-02), pages 84-96, XP027664360, ISSN: 0168-3659 [retrieved on 2005-11-02]

EGLETON ET AL.: "Development of Neuropeptide Drugs that Cross the Blood-Brain Barrier", JOURNAL OF THE AMERICAN SOCIETY FOR EXPERIMENTALNEUROTHERAPEUTICS, XX, XX, vol. 2, no. 1, 1 January 2005 (2005-01-01) , pages 44-53, XP005871740, ISSN: 1545-5343, DOI: 10.1602/NEURORX.2.1.44

SANTONICOLA, M. GABRIELLA ET AL.: 'Binding of alkyl polyglucoside surfactants to bacteriorhodopsin and its relation to protein stability' BIOPHYSICAL JOURNAL vol. 94, no. 9, 2008, pages 3647 - 3658, XP025961760

ZHOU, BIHONG ET AL.: 'Small surfactant-like peptides can drive soluble proteins into active aggregates'

MICROBIAL CELL FACTORIES vol. 11, no. 10, 18 January 2012, pages 1 - 8, XP021118466


DESCRIPTION

FIELD OF THE INVENTION


[0001] Covalently modified peptide and protein analogs allow for improved pharmaceutical properties of peptide and/or protein-based therapeutics.

SUMMARY

[0002] The present invention provides a peptide product comprising a surfactant X covalently attached to a peptide, the peptide comprising a linker amino acid U and at least one other amino acid:



wherein X is



wherein:

1. A is a hydrophobic group which is a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group, or a steroid nucleus-containing moiety; and
2. B is a melibiose or melibiouronic acid group covalently attached to the peptide via the linker amino acid U.


[0003] Peptide and/or protein pharmaceuticals suffer from several limitations in their use in medicine (Nestor, J.J., Jr. (2007) Comprehensive Medicinal Chemistry II 2: 573-601) - short duration of action, poor bioavailability, and lack of receptor subtype selectivity. In addition, peptides and/or proteins are unstable in formulations, being subject to aggregation. In some instances, aggregation of peptides and/or proteins leads to the development of an immunological response to both native and foreign peptides or proteins. WO2012/158962 describes covalently modified peptides and/or proteins said to have improved pharmaceutical properties.

[0004] Described herein is a method and reagents for covalently modifying peptides and/or proteins to generate products with improved pharmaceutical properties. In some instances, covalently modified peptides and/or proteins allow for improved stability, bioavailability, selectivity, and duration of effect in peptide and/or protein-based therapeutics.


[0005] Described herein are certain covalently modified peptides and/or proteins with improved pharmaceutical properties. In some instances, these covalently modified peptides and/or proteins allow for improved stability, bioavailability, selectivity, and duration of effect in peptide and/or protein-based therapeutics.

[0006] In some cases, the covalently modified peptides and/or proteins described herein are attached to glycoside surfactants. In one case, the covalently modified peptides and/or proteins are attached to alkyl glycosides. In one case, the covalently modified peptides and/or proteins are attached to an alkyl glycoside surfactant wherein the peptide and/or protein is attached to the glycoside in the surfactant and the glycoside is then attached to a hydrophobic and/or alkyl group. Described herein, are reagents and intermediates for the covalent modification of peptides and/or proteins through the incorporation of surfactants such as alkyl glycosides.

[0007] Described herein are peptide products comprising a surfactant X, covalently attached to a peptide, the peptide comprising a linker amino acid U and at least one other amino acid:



wherein X is



wherein

1. A is a hydrophobic group; and
2. B is a hydrophilic group covalently attached to the peptide via a linker amino acid U.

[0008] In some embodiments, the peptide product is synthesized by reaction of a functionalized surfactant with the peptide as described herein. In some embodiments, the peptide product is synthesized by reaction of a functionalized surfactant with a reversibly-protected linker amino acid as described herein followed by reaction with one or more amino acids to form a surfactant-modified peptide product. In some embodiments, U is a terminal amino acid of the peptide. In some embodiments, U is a non-terminal amino acid of the peptide.

[0009] In some embodiments, A is a substituted or unsubstituted C_1 - C_{30} alkyl chain, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group or a steroid nucleus containing moiety. In some embodiments, A is a substituted or unsubstituted C_8 - C_{20} alkyl chain, a substituted or unsubstituted 1-alkoxyaryl group, a substituted or unsubstituted 1-aralkyl group, or a steroid nucleus containing moiety. In some embodiments, A is a substituted or unsubstituted C_{10} - C_{20} alkyl chain, a substituted or unsubstituted 1-alkoxyaryl

group, a substituted or unsubstituted 1-alkyl group, or a steroid nucleus containing moiety.

[0010] In some embodiments, B is a polyol. In some embodiments, the polyol is a saccharide. In some embodiments, the saccharide is a monosaccharide, a disaccharide, or a polysaccharide. In some embodiments, the saccharide is selected from glucose, mannose, maltose, a glucuronic acid, galacturonic acid, diglucuronic acid and maltouronic acid.

[0011] In some embodiments, the surfactant is a 1-alkyl glycoside class surfactant.

[0012] In some embodiments, the hydrophilic group in the surfactant is attached to the peptide via an amide bond.

[0013] In some embodiments of the peptide product, the surfactant is comprised of 1-eicosyl beta-D-glucuronic acid, 1-octadecyl beta-D-glucuronic acid, 1-hexadecyl beta-D-glucuronic acid, 1-tetradecyl beta-D-glucuronic acid, 1-dodecyl beta-D-glucuronic acid, 1-decyl beta-D-glucuronic acid, 1-octyl beta-D-glucuronic acid, 1-eicosyl beta-D-diglucuronic acid, 1-octadecyl beta-D-diglucuronic acid, 1-hexadecyl beta-D-diglucuronic acid, 1-tetradecyl beta-D-diglucuronic acid, 1-dodecyl beta-D-diglucuronic acid, 1-decyl beta-D-diglucuronic acid, 1-octyl beta-D-diglucuronic acid, or functionalized 1-eicosyl beta-D-glucose, 1-octadecyl beta-D-glucose, 1-hexadecyl beta-D-glucose, 1-tetradecyl beta-D-glucose, 1-dodecyl beta-D-glucose, 1-decyl beta-D-glucose, 1-octyl beta-D-glucose, 1-eicosyl beta-D-maltoside, 1-octadecyl beta-D-maltoside, 1-hexadecyl beta-D-maltoside, 1-dodecyl beta-D-maltoside, 1-decyl beta-D-maltoside, 1-octyl beta-D-maltoside, and the like, and the peptide product is prepared by formation of a linkage between the aforementioned groups and a group on the peptide (e.g., a -COOH group in the aforementioned groups and an amino group of the peptide).


[0014] In some embodiments, a combination of a hydrophilic with a hydrophobic group generates a surfactant. In some embodiments, the surfactant is an ionic surfactant. In some embodiments, the surfactant is a non-ionic surfactant. In some embodiments, the hydrophobic group is a substituted or unsubstituted C₁-C₃₀ alkyl chain or an aralkyl chain. In some embodiments the hydrophobic group is a chain having mixed hydrophobic and hydrophilic properties, for example a polyethylene glycol (PEG) group.

[0015] In some embodiments, the linker amino acid is a natural D- or L- amino acid. In some embodiments, the linker amino acid is an unnatural amino acid. In some embodiments, the linker amino acid is selected from Lys, Cys, Orn, Asp, Glu or an unnatural amino acid, comprising a functional group used for covalent attachment to the surfactant X. In some embodiments, the linker amino acid is selected from Lys, Cys, Orn, or an unnatural amino acid, comprising a functional group used for covalent attachment to the surfactant X. In some embodiments, the functional group used for covalent attachment to a surfactant is -NH₂, -SH, -OH, -N₃, haloacetyl, a -(CH₂)_m-maleimide or an acetylenic group, wherein m is 1-10.

[0016] In some embodiments, the peptide is an opioid peptide. In some embodiments, the peptide and/or protein product contains a covalently linked alkyl glycoside. In some of such

embodiments, the peptide and/or protein product contains a covalently linked alkyl glycoside that is a 1-O-alkyl glucuronic acid of alpha or beta configuration. In some of such embodiments, the peptide and/or protein product comprises a covalently linked alkyl glycoside that is a 1-O-alkyl glucuronic acid and the alkyl chain is a C₁ to C₂₀ alkyl chain.

[0017] In some embodiments, the peptide product has a structure of Formula IA:



wherein:

each of aa₁, aa₂, aa₃, aa₄, and aa₅ is independently absent, a D- or L- natural or unnatural amino acid, an N-alkylated amino acid, an N-acetylated amino acid, a CαR³ amino acid, a Ψ-amino acid, or a linker amino acid U covalently attached to the surfactant X;


Z is -OH, -NH₂ or -NHR³;

each R³ is independently substituted or unsubstituted C₁-C₁₂ branched or straight chain alkyl, a PEG chain of less than 10 Da, or substituted or unsubstituted aralkyl chain;

provided that one, or at least one of aa₁, aa₂, aa₃, aa₄, and aa₅ is the linker amino acid U covalently attached to the surfactant X;

and further provided that not all of aa₁, aa₂, aa₃, aa₄, and aa₅ are absent.

[0018] In one aspect, provided herein is a peptide product having a structure of Formula II:




wherein:

each of aa₁, aa₂, aa₃, aa₄, and aa₅ is independently absent, a D- or L- natural or unnatural amino acid, an N-alkylated amino acid, an N-acetylated amino acid, a CαR³ amino acid, a Ψ-amino acid, or a linker amino acid U covalently attached to a surfactant X;

Z is -OH, -NH₂ or -NHR³;

each R³ is independently substituted or unsubstituted C₁-C₁₂ branched or straight chain alkyl, a PEG chain of less than 10 Da, or substituted or unsubstituted aralkyl chain; and

X is



wherein

1. A is a hydrophobic group; and
2. B is a hydrophilic group covalently attached to the peptide via a linker amino acid U;

provided that one, or at least one of aa₁, aa₂, aa₃, aa₄, and aa₅ is the linker amino acid U covalently attached to the surfactant X;

and further provided that not all of aa₁, aa₂, aa₃, aa₄, and aa₅ are absent.

[0019] In one aspect, provided herein is a peptide product that has a structure of Formula III:



wherein:

aa₁ is Tyr, Dmt, N-R³-Tyr, N-R³-Dmt, N-(R³)₂-Tyr, or N-(R³)₂-Dmt;

aa₂ is Pro, D-Arg, D-Cit, D-U(X), D-Ala, Tic, or Tic(Ψ[CH₂-NH]);


aa₃ is Phe, Trp, Tmp, D-or L-Nal(1), D-or L-Nal(2), CαMePhe, or Ψ-Phe;

aa₄ is Phe, Tmp, D-or L-Nal(1), D-or L-Nal(2), U(X), D- or L-CαMeU(X);

aa₅ is absent or Pro, Aib, U(X), D- or L-CαMeU(X); and

U is a dibasic natural or unnatural amino acid, a natural or unnatural amino acid comprising a thiol, an unnatural amino acid comprising a -N₃ group, an unnatural amino acid comprising an acetylenic group, or an unnatural amino acid comprising a -NH-C(=O)-CH₂-Br or a -(CH₂)_m-maleimide, wherein m is 1-10, used for covalent attachment to the surfactant X.

[0020] In some embodiments of peptide products of Formula I, Formula II or Formula III described above and herein, X has the structure:



Formula V

wherein:

A is a substituted or unsubstituted C_1 - C_{30} alkyl chain, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group, or a steroid nucleus containing moiety;

R^{1b} , R^{1c} , and R^{1d} are each, independently at each occurrence, H, a protecting group, a saccharide, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group;


W^1 is independently, at each occurrence, $-CH_2-$, $-CH_2-O-$, $-(C=O)-$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$, or $-CH_2-S-$;

W^2 is $-O-$, or $-S-$;

R^2 is a bond, C_2 - C_4 -alkene, C_2 - C_4 -alkyne, or $-(CH_2)_m$ -maleimide; and


m is 1-10.

[0021] In some embodiments of peptide products of Formula I, Formula II or Formula III described above and herein, X has the structure:




[0022] Accordingly, in the embodiment described above, R^2 is a bond.

[0023] In some embodiments of peptide products of Formula I, Formula II or Formula III described above and herein, X has the structure:



[0024] In some embodiments of peptide products of Formula I, Formula II or Formula III described above and herein, X has the structure:



OR^{1c}

Formula V

wherein:

A is a substituted or unsubstituted C₁-C₃₀ alkyl group, or a steroid nucleus containing moiety;


R^{1b}, R^{1c}, and R^{1d} are each, independently at each occurrence, H, a protecting group, a saccharide or a substituted or unsubstituted C₁-C₃₀ alkyl group;

W¹ is -(C=O)-NH-;

W² is -O-;

R² is a bond.

[0025] In some embodiments of peptide products of Formula I, Formula II or Formula III described above and herein, X has the structure:



wherein:

A is a substituted or unsubstituted C₁-C₃₀ alkyl group;

R^{1b}, R^{1c}, and R^{1d} are H;

W¹ is -(C=O)-NH-;

W² is -O-; and


R² is a bond.

[0026] In some embodiments of peptide products of Formula I, Formula II or Formula III described above and herein, X is as described above and A is a substituted or unsubstituted C₁-C₂₀ alkyl group.

[0027] In some embodiments described above and herein, R^{1a} is a saccharide. In some embodiments, the saccharide is a galactose. In certain embodiments, the saccharide is an alpha-linked galactose. In other embodiments, the saccharide is alpha-linked galactopyranose,

beta-linked galactopyranose, alpha-linked galactofuranose, or beta-linked galactofuranose.

[0028] Also contemplated herein are alternate embodiments wherein X in Formula I has the structure:



[0029] For instance, in an exemplary embodiment of the structure of X described above, W¹ is -S-, R² is a C₁-C₃₀ alkyl group, W² is S, R^{1a} is a bond between W² and a suitable moiety of an amino acid residue U within the peptide (e.g., a thiol group in a cysteine residue of the peptide forms a thioether with X).

[0030] In another exemplary alternate embodiment of the structure of X described above, W¹ is -O-, R² is a C₁-C₃₀ alkyl group, W² is O, R^{1a} is a bond between W² and a suitable moiety of an amino acid residue U within the peptide (e.g., a hydroxyl group in a serine or threonine residue of the peptide forms an ether with X).

[0031] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 149**) wherein:

U is a dibasic natural or unnatural amino acid;

X is a surfactant of the 1-alkyl glycoside class wherein 1-alkyl is substituted or unsubstituted C₁-C₂₀ alkyl or a substituted or unsubstituted alkoxyaryl substituent;

Z is NH₂;

aa₁ is Tyr, Dmt, Nα-Me-Tyr, Na-Me-Tyr, N,Nα-diMe-Tyr, or N,Nα-diMe-Dmt;

aa₂ is Pro, D-Arg, D-Cit, D-U(X), D-Ala, Tic, or Tic(Ψ[CH₂-NH]);

aa₃ is Phe, Trp, Tmp, D-or L-Nal(1), D-or L-Nal(2), CaMePhe, or Ψ-Phe;

aa₄ is Phe, Tmp, D-or L-Nal(1), D-or L-Nal(2), U(X), D- or L-CcMeU(X);

aa₅ is absent or Pro, Aib, U(X), D- or L-CaMeU(X).

[0032] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 150**) wherein:

X is comprised of 1-alkyl glucuronic acid or 1-alkyl diglucuronic acid;

Z is NH₂;

aa₁ is Dmt;

aa₂ is Pro, D-Lys(X), Tic, or Tic(Ψ[CH₂-NH]);

aa₃ is Phe, Tmp, D-or L-Nal(1), D-or L-Nal(2), or Ψ-Phe;

aa₄ is Phe, D-or L-Nal(1), D-or L-Nal(2), or Lys(X);

aa₅ is absent or Pro, Aib, Lys(X), D- or L-CaMeLys(X).

[0033] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 151**) wherein:

aa₁ is Dmt;

aa₂ is Pro;

aa₃ is Phe, or Tmp;

aa₄ is Phe, or Lys(X);

aa₅ is absent or Aib, Lys(X), D- or L-CaMeLys(X).

[0034] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 152**) wherein:

aa₁ is Dmt;

aa₂ is Pro;

aa₃ is Phe, or Tmp;

aa₄ is Phe, or Lys(X);

aa₅ is absent or Lys(X).

[0035] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 153**) wherein:

U is a dibasic natural or unnatural amino acid;

X is a surfactant of the 1-alkyl glycoside class wherein the 1-alkyl group of the 1-alkyl glycoside is substituted or unsubstituted C₁₋₂₀ alkyl or a substituted or unsubstituted alkoxyaryl substituent;

aa₁ is Tyr, Dmt, N-R³-Tyr, N-R³-Tyr, N-(R³)₂-Tyr, or N-(R³)₂-Dmt;

aa₂ is D-Arg, D-Cit, or D-U(X);

aa₃ is Phe, Trp, D-or L-Nal(1), D-or L-Nal(2), or Tmp;

aa₄ is Phe, Tmp, D-or L-Nal(1), D-or L-Nal(2), U(X), D- or L-CaMeU(X);

aa₅ is absent, Pro, or U(X).

[0036] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 154**) wherein:

X is a surfactant of the 1-alkyl glucuronic acid or 1-alkyl diglucuronic acid class;

aa₁ is Dmt, Nα-Me-Dmt, or N,Nα-Me-Dmt;

aa₂ is D-Arg, D-Cit, D-Lys(X), or D-Orn(X);

aa₃ is Phe, or Tmp;

aa₄ is Phe, Tmp, Lys(X), or Orn(X);

aa₅ is absent or Pro, Lys(X), or Orn(X).

[0037] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 155**) wherein:

U is a dibasic natural or unnatural amino acid;

X is a surfactant of the 1-alkyl glycoside class wherein 1-alkyl is substituted or unsubstituted C_{1-C20} alkyl or a substituted or unsubstituted alkoxyaryloxy substituent;

Z is NH₂;

aa₁ is Tyr, Dmt, N-R³-Tyr, N-R³-Dmt, N-(R³)₂-Tyr, or N-(R³)₂-Dmt;

aa₂ is Tic, or Tic(Ψ [CH₂-NH]);

aa₃ is Ψ -Phe when aa₂ is Tic(Ψ [CH₂-NH]);

aa₄ is Phe, Tmp, D-or L-Nal(1), D-or L-Nal(2), or U(X);

aa₅ is absent, Pro, Aib, or U(X).

[0038] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 156**) wherein:

X is a surfactant of the 1-alkyl glucuronic acid or 1-alkyl diglucuronic acid class;

aa₁ is Tyr, Dmt, N α -Me-Tyr, N α -alkyl-Dmt, N,N α -diMe-Tyr, or N,N α -Me-Dmt;

aa₂ is Tic, or Tic(Ψ [CH₂-NH]);

aa₃ is Ψ -Phe when aa₂ is Tic(Ψ [CH₂-NH]), Phe, or TMP;

aa₄ is Phe, Tmp, D-or L-Nal(1), D-or L-Nal(2), Lys(X), or Orn(X);

aa₅ is absent or Aib, Lys(X), or Orn(X).

[0039] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 157**) wherein:

X is comprised of 1-alkyl glucuronic acid or 1-alkyl diglucuronic acid;

aa₂ is Tic, or Tic(Ψ [CH₂-NH]);

aa₃ is Phe or Ψ -Phe;

aa₄ is Lys(X);

aa₅ is absent.

[0040] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 158**) wherein:

X is comprised of 1-alkyl glucuronic acid;

aa₂ is Tic:

aa₃ is Phe;

aa₄ is Lys(X);

aa₅ is absent.

[0041] In some embodiments, a peptide product of Formula III is a product (**SEQ. ID. NO. 159**) wherein:

X is comprised of a 1-alkyl glucuronic acid selected from 1-methyl beta-D-glucuronic acid, 1-octyl beta-D-glucuronic acid, 1-dodecyl beta-D-glucuronic acid,

1-tetradecyl beta-D-glucuronic acid, 1-hexadecyl beta-D-glucuronic acid, and 1-octadecyl beta-D-glucuronic acid;

aa₂ is Tic;

aa₃ is Phe;

aa₄ is Lys(X);

aa₅ is absent.

[0042] In some embodiments, a peptide product of Formula III is H-Dmt-Tic-Phe-Lys(N-epsilon-1-methyl beta-D-glucuronyl)-NH₂. (**SEQ. ID. NO. 78**)

[0043] In some embodiments, a peptide product of Formula III is H-Dmt-Tic-Phe-Lys(N-epsilon-1-octyl beta-D-glucuronyl)-NH₂. (**SEQ. ID. NO. 80**)

[0044] In some embodiments, a peptide product of Formula III is H-Dmt-Tic-Phe-Lys(N-epsilon-1-dodecyl beta-D-glucuronyl)-NH₂. (**SEQ. ID. NO. 79**)

[0045] In some embodiments, a peptide product of Formula III is H-Dmt-Tic-Phe-Lys(N-epsilon-1-tetradecyl beta-D-glucuronyl)-NH₂. (**SEQ. ID. NO. 160**)

[0046] In some embodiments, a peptide product of Formula III is H-Dmt-Tic-Phe-Lys(N-epsilon-1-hexadecyl beta-D-glucuronyl)-NH₂. (**SEQ. ID. NO. 82**)

[0047] In some embodiments, a peptide product of Formula III is H-Dmt-Tic-Phe-Lys(N-epsilon-1-octadecyl beta-D-glucuronyl)-NH₂. (**SEQ. ID. NO. 83**)

[0048] In some embodiments, the peptide product is biologically active.

[0049] In a specific embodiment, provided herein is a compound selected from compounds of Table 1 in Figure 1.

[0050] Also provided herein is a pharmaceutical composition comprising a therapeutically effective amount of a peptide product of Formula I, II or III, or pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier or excipient.

[0051] Provided herein is a method of treating pain comprising administration of a therapeutically effective amount of a peptide product of Formula I, II or III or compounds of Table 1 in Figure 1.

[0052] A method for improving the pharmaceutical and medicinal behavior of a peptide, thereby improving its duration of action, bioavailability or its stability in formulation, comprising covalent attachment of a surfactant X to the peptide, wherein X is as described herein.

[0053] In some embodiments, a peptide product comprising a covalently linked alkyl glycoside described herein is an analog of Leu-enkephalin. In some of such embodiments, the peptide product contains a covalently linked 1-O-alkyl β -D-glucuronic acid and the peptide is an analog of Leu-enkephalin.

[0054] In some embodiments, a peptide product comprising a covalently linked alkyl glycoside described herein is an analog of opioid peptide DPDPE (Akiyama K, et al. (1985) Proc Natl Acad Sci USA 82:2543-7). In some of such embodiments, the peptide product contains a covalently linked 1-O-alkyl glucuronic acid and the peptide is an analog of opioid peptide DPDPE.

[0055] In some embodiments, a peptide product comprising a covalently linked alkyl glycoside described herein is an analog of the D-amino acid containing natural product peptides dermorphin (Melchiorri, P. and Negri, L. (1996) Gen Pharmacol 27: 1099-1107) or deltorphin (Erspamer, V., et al. (1989) Proc Natl Acad Sci USA 86:5188-92). In some of such embodiments, the peptide product contains a covalently linked 1-O-alkyl β -D-glucuronic acid and the peptide is an analog of dermorphin or deltorphin.


[0056] In some embodiments, a peptide product comprising a covalently linked alkyl glycoside is an analog of endomorphin-1 or -2. In some of such embodiments, the peptide product comprises a covalently linked 1-O-alkyl β -D-glucuronic acid and the peptide is an analog of endomorphin (Lazarus, L.H. and Okada, Y. (2012) Expert Opin Ther Patents 22: 1-14).

[0057] In some embodiments, a peptide product comprising a covalently linked alkyl glycoside described herein is an analog of opioid peptide dynorphin (James, I.F., et al. (1982) Life Sci 31:1331-4). In some of such embodiments, the peptide product contains a covalently linked 1-O-alkyl β -D-glucuronic acid and the peptide is an analog of dynorphin.

[0058] In some embodiments side chain functional groups of two different amino acid residues are linked to form a cyclic lactam. For example, in some embodiments, a Lys side chain forms a cyclic lactam with the side chain of Glu. In some embodiments such lactam structures are reversed and are formed from a Glu and a Lys. Such lactam linkages in some instances are known to stabilize alpha helical structures in peptides (Condon, S.M., et al. (2002) Bioorg Med Chem 10: 731-736).


[0059] In some embodiments side chain functional groups of two different amino acid residues with -SH containing side chains are linked to form a cyclic disulfide. For example, in some embodiments, two penicillamine or two Cys side chains may be linked to constrain the conformation of a peptide (Akiyama K, et al. (1985) Proc Natl Acad Sci USA 82:2543-7) in order to yield greater duration of action or greater receptor selectivity.

[0060] In a specific embodiment, the peptide products of Formula I, Formula II or Formula III, described above and herein have the following structure:



wherein A is a C₁-C₂₀ alkyl chain as described in Table 1 of Figure 1, R' is a peptide as described in Table 1 of Figure 1, W² of Formula V is -O-, and W¹ of Formula V is -(C=O)NH- and is part of an amide linkage to the peptide R'. In some of such embodiments, A is a C₆-C₂₀ alkyl chain. In some of such embodiments, A is a C₁-C₁₀ alkyl chain. In some of such embodiments, A is a C₁₂-C₂₀ alkyl chain. In some of such embodiments, A is a C₁₂-C₁₈ alkyl chain.


[0061] In embodiments described above, an amino moiety of an amino acid and/or a peptide R' (e.g., an amino group of an amino acid residue such as a lysine, or a lysine within the peptide R') is used to form a covalent linkage with a compound of structure:




(Formula A),


wherein A is a C₁-C₂₀ alkyl chain as described above and in Table 1 of Figure 1.

[0062] In such cases, the amino acid residue having an amino moiety (e.g., a Lysine within the peptide R') which is used to form a covalent linkage to the compound of Formula A described above, is a linker amino acid U which is attached to a surfactant X. Accordingly, as one example, Lys(C12) of Table 1 of Figure 1 has the following structure:






[0063] Also described are peptide products of Formula I derived from maltouronic acid-based surfactants prepared by covalent linkage of a peptide to either or both carboxylic acid groups. Thus, as one example, peptides in Table 1 of Figure 1 comprise a lysine linker amino acid bonded to a maltouronic acid based surfactant X and having a structure:



[0064] Also described are peptide products of Formula I-A derived from melibiouronic acid-based surfactants through binding at the carboxylic acid function. Thus, as one example, peptides in Table 1 of Figure 1 or Table 2 of Figure 2 or Table 3 of Figure 3 comprise a lysine linker amino acid bonded to a melibiouronic acid based surfactant X and having a structure:



[0065] It will be understood that in one embodiment, compounds of Formula I are prepared by attaching a lysine to a group X, followed by attachment of additional amino acid residues and/or peptides are attached to the lysine-X compound to obtain compounds of Formula I. It will be understood that other natural or non-natural amino acids described herein are also

suitable for attachment to the surfactant X and are suitable for attaching additional amino acid/peptides to obtain compounds of Formula I. It will be understood that in another embodiment, compounds of Formula I are prepared by attaching a full length or partial length peptide to a group X, followed by optional attachment of additional amino acid residues and/or peptides are attached to obtain compounds of Formula I.

[0066] Also described herein is a pharmaceutical composition comprising a therapeutically effective amount of a peptide product described above, or acceptable salt thereof, and at least one pharmaceutically acceptable carrier or excipient. In some cases, the carrier is an aqueous-based carrier. In some cases, the carrier is a nonaqueous-based carrier. In some cases, the nonaqueous-based carrier is a hydrofluoroalkane-like solvent comprising sub-micron anhydrous α -lactose, or other excipients.

[0067] Described herein is the reaction of an amino acid and/or a peptide comprising a linker amino acid U bearing a nucleophile, and a group X comprising a leaving group or a functional group that can be activated to contain a leaving group, for example a carboxylic acid, or any other reacting group, thereby allowing for covalent linkage of the amino acid and/or peptide to a surfactant X via the linker amino acid U to provide a peptide product of Formula I.


[0068] Also described herein is the reaction of an amino acid and/or a peptide comprising a linker amino acid U bearing a leaving group or a functional group that can be activated to contain a leaving group, for example a carboxylic acid, or any other reacting group, and a group X comprising a nucleophilic group, thereby allowing for covalent linkage of the amino acid and/or peptide to a surfactant X via the linker amino acid U to provide a peptide product of Formula I.

[0069] It will be understood that, in one case, Compounds of Formula I are prepared by reaction of a linker amino acid U with X, followed by addition of further residues to U to obtain the peptide product of Formula I. It will be understood that in an alternative case, Compounds of Formula I are prepared by reaction of a suitable peptide comprising a linker amino acid U with X, followed by optional addition of further residues to U, to obtain the peptide product of Formula I.

[0070] Further provided herein are certain intermediates and/or reagents that are suitable for synthesis of peptide products described herein. In certain embodiments, such intermediates and/or reagents are functionalized surfactants that allow for covalent linkage with a peptide. In certain embodiments, such intermediates are functionalized 1-alkyl glycoside surfactants that allow for covalent linkage with a peptide. In certain embodiments, such intermediates are functionalized 1-alkyl glycoside surfactants linked to reversibly-protected linker amino acids that allow for covalent linkage with other amino acids to form a peptide. It will be understood that a suitably functionalized surfactant is covalently coupled to a peptide via reaction with an appropriately matched functional group that is on a linker amino acid.

[0071] In some embodiments, intermediates suitable for synthesis of peptide products

described herein are compounds of Formula IV. Accordingly, provided herein are intermediates and/or compounds of Formula IV:



Formula IV

wherein:

R^{1a} is independently at each occurrence a bond, H, a protecting group, a natural or unnatural amino acid, a substituted or unsubstituted C_1 - C_{30} alkyl hydrophobic group, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group, or a steroid nucleus containing moiety;

R^{1b} , R^{1c} , and R^{1d} are each independently at each occurrence a bond, H, a protecting group, a natural or unnatural amino acid, a substituted or unsubstituted C_1 - C_{30} alkyl hydrophobic group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group;

W^1 is $-CH_2-$, $-CH_2-O-$, $-(C=O)-$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$, or $-CH_2-S-$;

W^2 is $-O-$, $-CH_2-$ or $-S-$;

R^2 is H, a protecting group, a natural or unnatural amino acid, a substituted or unsubstituted C_1 - C_{30} alkyl hydrophobic group, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group, $-NH_2$, $-SH$, C_2 - C_4 -alkene, C_2 - C_4 -alkyne, $-NH(C=O)-CH_2-Br$, $-(CH_2)_m$ -maleimide, or $-N_3$;

n is 1, 2 or 3; and

m is 1-10.

[0072] In some embodiments, each natural or unnatural amino acid is independently a reversibly protected or free linker amino acid. In some of such embodiments, the linker amino acid is a reversibly protected or free lysine.

[0073] In some embodiments of Formula IV,

n is 1;

W^1 is $-(C=O)-$;

R^{1a} is a substituted or unsubstituted C_1 - C_{30} alkyl hydrophobic group, a substituted or unsubstituted 1-alkoxyaryl group, or a substituted or unsubstituted 1-aralkyl group,

R^2 is a reversibly-protected lysine of D- or L-configuration.

[0074] In some embodiments of Formula IV,

n is 1;

W^1 is $-(C=O)-$;

R^{1a} is a substituted or unsubstituted C_8 - C_{30} alkyl hydrophobic group, a substituted or unsubstituted 1-alkoxyaryl group, or a substituted or unsubstituted 1-aralkyl group,

R^2 is a reversibly protected lysine of D- or L- configuration.

[0075] In some of such embodiments, R^{1a} is an octyl, decyl, dodecyl, tetradecyl, or hexadecyl group.

[0076] In some embodiments of Formula IV,

n is 1;

W^1 is $-(C=O)-NH-$ or $-(C=O)-O-$;

R^2 is a substituted or unsubstituted C_1 - C_{30} alkyl hydrophobic group, a substituted or unsubstituted 1-alkoxyaryl group, or a substituted or unsubstituted 1-aralkyl group,

R^{1a} is a reversibly protected serine or threonine of D- or L- configuration.

[0077] In some of such embodiments, R^2 is an octyl, decyl, dodecyl, tetradecyl or hexadecyl group.

[0078] In some embodiments of Formula IV,

n is 1;

m is 1-6;

W^1 is $-CH_2-$;

R^{1a} is a C_1 - C_{30} alkyl hydrophobic group, a 1-alkoxyaryl group, or a 1-aralkyl group,

R^2 is $-N_3$, NH_2 , $-C_2$ -alkyne, $-(CH_2)_m$ -maleimide, $NH-(C=O)-CH_2-Br$, or $NH-(C=O)-CH_2-I$.

[0079] In some embodiments of Formula IV,

n is 1;

W^1 is $-(C=O)-O-$;

R^2 is H,

R^{1a} is a substituted or unsubstituted C_1 - C_{30} alkyl hydrophobic group.

[0080] In some embodiments of Formula IV, R^2 is attached to the peptide or is an amino acid residue in the peptide. In some of such embodiments, R^2 is a reversibly protected or free lysine.

[0081] In some embodiments of Formula IV, n is 1. In some embodiments of Formula IV, n is 2, and a first glycoside is attached to a second glycoside via a bond between W^2 of the first glycoside and any one of OR^{1b} , OR^{1c} or OR^{1d} of the second glycoside.

[0082] In some embodiments of Formula IV, n is 3, and a first glycoside is attached to a second glycoside via a bond between W^2 of the first glycoside and any one of OR^{1b} , OR^{1c} or OR^{1d} of the second glycoside, and the second glycoside is attached to a third glycoside via a bond between W^2 of the second glycoside and any one of OR^{1b} , OR^{1c} or OR^{1d} of the third glycoside.

[0083] Provided herein is a method for synthesizing a peptide product described above, comprising sequential steps of


1. (a) coupling a compound of Formula IV to the peptide; and
2. (b) optionally deprotecting the coupled peptide of step (a).

[0084] In some embodiments, the deprotecting comprises the use of mild acid and or mild base treatments. In some embodiments of the methods, the deprotecting comprises the use of strong acids.

[0085] In some embodiments, the method further comprises the steps of chromatography, desalting of intermediates by reversed-phase, high-performance liquid chromatography or ion exchange chromatography of intermediates.

[0086] Provided herein is a method for improving the pharmaceutical and medicinal behavior of a peptide, thereby improving its duration of action, bioavailability or its stability in formulation, comprising covalent attachment of a surfactant X to the peptide, wherein:

X is



wherein

A is a hydrophobic group; and

B is a hydrophilic group covalently attached to the peptide via a linker amino acid.

[0087] In some embodiments, A is a substituted or unsubstituted C₁-C₃₀ alkyl chain, a substituted or unsubstituted alkoxyaryl group or a substituted or unsubstituted aralkyl group.

[0088] In some embodiments, A is a substituted or unsubstituted C₈-C₂₀ alkyl chain, a substituted or unsubstituted 1-alkoxyaryl group or a substituted or unsubstituted 1-aralkyl group.

[0089] In some embodiments, A is a substituted or unsubstituted C₁₀-C₂₀ alkyl chain, a substituted or unsubstituted 1-alkoxyaryl group or a substituted or unsubstituted 1-aralkyl group.

[0090] In some embodiments, B is a functionalized polyol.

[0091] In some embodiments, the polyol is a saccharide,


[0092] In some embodiments, the saccharide is a monosaccharide, a disaccharide, or a polysaccharide.

[0093] In some embodiments, the saccharide is selected from glucuronic acid, galacturonic acid, diglucuronic acid and maltouronic acid.

[0094] The methods of synthesis described above are suitable for synthesis of all compounds described herein, including compounds of Formula I, II, III, 2-1-1, 2-III, 2-V, 2-VI, 2-VII, 3-I-A, 3-III-A, 3-III-B, or 3-V, and compounds in Table 1, Table 2, Table 3 and Table 4 provided in Figure 1, Figure 2, Figure 8 and Figure 9 respectively.

[0095] Provided herein is a method for improving the pharmaceutical and medicinal behavior of a peptide, thereby improving its duration of action, bioavailability or its stability in formulation, comprising covalent attachment of a surfactant X to the peptide, wherein:

X is



wherein

A is a substituted or unsubstituted C₁-C₂₀ alkyl chain, a substituted or unsubstituted alkoxyaryl group or a substituted or unsubstituted aralkyl group; and

B is a saccharide covalently attached to the peptide via a linker amino acid.

[0096] In some embodiments, the surfactant is a 1-alkyl glycoside class surfactant.

[0097] In some embodiments, the hydrophilic group in the surfactant is attached to the peptide via an amide bond.

[0098] In some embodiments, the surfactant is composed of 1-hexadecyl-beta-D-glucuronic acid, 1-tetradecyl-beta-D-glucuronic acid, 1-dodecyl-beta-D-glucuronic acid, 1-decyl-beta-D-glucuronic acid, 1-octyl-beta-D-glucuronic acid, 1-hexadecyl-beta-D-diglucuronic acid, 1-tetradecyl-beta-D-diglucuronic acid, 1-dodecyl-beta-D-diglucuronic acid, 1-decyl-beta-D-diglucuronic acid, 1-octyl-beta-D-diglucuronic acid.

[0099] Provided herein is a method of treating pain in an individual in need thereof comprising administration of a therapeutically effective amount of a peptide product described herein, or a compound of Formula IV to an individual in need thereof.

[0100] Also provided herein is a covalently modified peptide and/or protein product comprising a hydrophilic group as described herein; and a hydrophobic group covalently attached to the hydrophilic group. In specific embodiments, the covalently modified peptide and/or protein product comprises a hydrophilic group that is a saccharide and a hydrophobic group that is a C₁-C₂₀ alkyl chain or an aralkyl chain.

BRIEF DESCRIPTION OF THE FIGURES

[0101]

Figure 1. Table 1 in Figure 1 depicts compounds that were prepared by methods described herein. The specification provides sequences for SEQ. ID. Nos. 1 and 149-169. Additionally, Table 1 of Figure 1 provides SEQ. ID Numbers for compounds EU-A101 to EU-A199 and EU-

A600 to EU-A649 having SEQ. ID. NOs. 2-148, and SEQ. ID. NO. 645 respectively, as shown in Table 1 of Figure 1. Compounds in Table 1 of Figure 1, and their respective SEQ. ID. NOs. shown in Table 1 of Figure 1 are hereby incorporated into the specification as filed.

Figure 2. Table 2 in Figure 2 depicts compounds that were prepared by methods described herein. The specification provides sequences for SEQ. ID. Nos. 170-174 and SEQ. ID. NOs. 283-302. Additionally, Table 2 of Figure 2 provides SEQ. ID. Numbers for compounds EU-201 to EU-299 and EU-900 to EU-908 having SEQ. ID. NOs. 175-282 respectively, as shown in Table 2 of Figure 2. Compounds in Table 2 of Figure 2, and their respective SEQ. ID. NOs. shown in Table 2 of Figure 2 are hereby incorporated into the specification as filed.

Figure 3. Figure 3 has two panels. The **top panel** in Figure 3 illustrates the interaction between PTH 1-34 or PTHrP 1-34 and the PTH R1 receptor based on the x-ray crystal structure (Piozak, A.A., et al. (2009) J Biol Chem 284:28382-391) of the extracellular domain of the receptor. The critical hydrophobic interactions of residues 23' (W or F), 24' (L) and 28' (L or I) are illustrated. The surfactant modification on the modified peptides described herein, in some instances, replace these critical hydrophobic interactions. The **bottom panel** in Figure 3 shows the structural comparison between the sequence of PTH 1-34 and PTHrP 1-34 and illustrates the strong regions of identity and homology in the sequences and peptide helical conformation.

Figure 4. Figure 4 shows the cAMP responses of human cells in culture (SaOS2) to stimulation by a representative peptide product described herein, EU-232. The ordinate (vertical axis) shows the response as a percentage of the maximal response shown to the internal assay standard, i.e., PTHrP. The data illustrates a super-agonistic response to EU-232.

Figure 5. Figure 5 shows the response of human cells in culture (SaOS2) to treatment with various doses of EU-285 (a coded sample of human PTHrP). The ordinate (vertical axis) shows the cAMP response as a percentage of the maximal response of the internal assay standard, i.e., PTHrP.

Figure 6. Blood phosphate levels in rat serum were tested at various time points after subcutaneous dosing rats with saline (G1), 80 micrograms per kg of PTH (G2), 80 micrograms per kg of EU-232 (G3) or 320 micrograms per kg of EU-232 (G4). This surfactant modified analog, EU-232, demonstrates prolonged duration of action, as evidenced by the maximal statistically significant effect, which is seen at the last time point in the assay (i.e., 5 hrs post dosing). See Example 6.

Figure 7. Blood calcium levels in rat serum were tested at various time points after subcutaneous dosing in rats with saline (G1), 80 micrograms per kg of PTH (G2), 80 micrograms per kg of EU-232 (G3) or 320 micrograms per kg of EU-232 (G4). No groups were statistically significantly different from control (G1). Further, the maximally effective dose and time point for EU-232 (G4; at 5hrs) shows no elevation and thus no indications of a propensity for hypercalcemia at a maximally effective dose. See Example 6.

Figure 8. Table 3 in Figure 8 depicts compounds that were prepared by methods described

herein. The specification provides sequences for SEQ. ID. Nos. 303-305 and SEQ. ID. Nos. 619 -644. Additionally, Table 3 of Figure 8 provides SEQ. ID Numbers for compounds EU-A300 to EU-A425 having SEQ. ID. NOs. 306-431 respectively, as shown in Table 3 of Figure 8. Compounds in Table 3 of Figure 8, and their respective SEQ. ID. NOs. shown in Table 3 of Figure 8 are hereby incorporated into the specification as filed.

Figure 9. Table 4 in Figure 9 depicts compounds that were prepared by methods described herein. The specification provides SEQ. ID. Nos. 303-305 and SEQ. ID. Nos. 619 - 644. Additionally, Table 4 of Figure 9 provides SEQ. ID Numbers for compounds EU-A426 to EU-A599 having SEQ. ID. NOs. 432-618 respectively, as shown in Table 4 of Figure 9. Additionally, Table 4 of Figure 9 provides SEQ. ID Numbers for compounds EU-A700 to EU-A1174 having SEQ. ID. NOs 646-1120. Compounds in Table 4 of Figure 9, and their respective SEQ. ID. NOs. shown in Table 4 of Figure 9 are hereby incorporated into the specification as filed.

Figure 10. Figure 10 illustrates the x-ray crystal structure (Runge, S., et al. (2008) J Biol Chem 283: 11340-7) of the binding site of the extracellular domain of the GLP-1 receptor and illustrates critical hydrophobic binding elements of the receptor and the ligand exendin-4 (Val^{19*}, Phe^{22*}, Trp^{25*}, Leu^{26*}) which are mimicked and replaced by the hydrophobic 1'-alkyl portion of the surfactant on the peptides of the invention.

Figure 11. Figure 11 shows cellular assay data of a representative compound, EU-A178, acting on cells containing the mu opioid receptor (MOP) and acting as an antagonist on the delta opioid receptor (DOP) in competition with 30nM DPDPE. The representative compound shows potent, full agonistic behavior in the MOP agonist assay and highly potent action as a pure antagonist on the DOP receptor.

Figure 12. Figure 12 illustrates the *in vivo* blood glucose response in db/db mice upon s.c. administration of the listed amount of test compounds of the invention (EU-A993 and EU-A1023) at times t=0, 7 hrs.

Figure 13. Figure 13 illustrates examples of the detailed structure of some compounds of the invention and their linkage through the epsilon amino function of a Lys residue, in this case at position 24, to examples of mono and disaccharide surfactants modified according to one method of the invention.

Figure 14. Figure 14 illustrates the structure of EU-A937, an example of the types of structures of the invention.

Figure 15. Figure 15 illustrates the concentrations of compounds EU-A993, EU-A1023 and the native hormone GLP-1 (7-36) with time during incubation in human plasma. This data illustrates the prolonged half life and protection from proteolysis for the compounds of the invention.

DETAILED DESCRIPTION

[0102] Provided herein are modified peptides and/or proteins that comprise a peptide and/or protein covalently attached to a hydrophilic group, a "head" (e.g., a polyol, (e.g., a saccharide)); the hydrophilic group is covalently attached to a hydrophobic group, a "tail", thereby generating a surfactant. In some embodiments, use of hydrophobic-linked glycoside surfactant (e.g., alkyl glycoside) moieties for covalent modification of the peptides or proteins, prolongs the duration of action of the peptides or proteins by multiple mechanisms, including formation of depots of the drug at the site of administration in the body and binding to hydrophobic carrier proteins. In some embodiments, incorporation of steric hindrance into peptide and/or protein structure can prevent approach of proteases to the peptide and/or protein product and thereby prevent proteolysis. In some embodiments, surfactant modification (e.g., covalent attachment of alkyl glycoside class of surfactants) of peptides and/or proteins as described herein, increases the transport across mucosal barriers. Accordingly, the modifications of the peptides and/or proteins described herein provide desirable benefits including and not limited to, protection from proteolysis, and slowed movement from the site of administration, thereby leading to prolonged pharmacokinetic behavior (e.g., prolongation of circulating $t_{1/2}$) and improved transmucosal bioavailability.

[0103] In some embodiments, interaction of the improved peptides and/or proteins with their receptors is modified in beneficial ways by the truncation of the sequence, introduction of constraint, and/or the incorporation of steric hindrance. Described herein are novel alkyl glycoside reagents that allow for incorporation of both rigidity and steric hindrance in the modified peptides and/or proteins. In some embodiments, steric hindrance confers receptor selectivity to the modified peptides and/or proteins described herein. In some embodiments, steric hindrance provides protection from proteolysis.

[0104] It has been recognized that activation of G Protein-coupled Receptors (GPCRs) cause the activation of multiple second messenger systems. Activated GPCRs also recruit arrestins, which had been thought to be responsible for stopping GPCR signaling and for internalization of the GPCR. Recently it has been recognized that arrestins themselves are a mediator of various signaling pathways, so that ligands which do not recruit arrestin or which only recruit arrestin can have important biological effects through "biased" signaling (Shenoy, S. K. and Lefkowitz, R.J.(2011) Trends Pharmacol Sci 32: 521-533). Incorporation of surfactants into the peptides of the invention, with the steric hindrance and altered presentation caused by surfactant modification, may exhibit such surprising and beneficial behavior. Evidence for this behavior is shown by peptides EU-A770 to EU-A773 in example 5.

[0105] Proteins and peptides undergo numerous physical and chemical changes that may affect potency and safety. Among these are aggregation, which includes dimerization, trimerization, and the formation of higher-order aggregates such as amyloids. Aggregation is a key issue underlying multiple potentially deleterious effects for peptide and/or protein-based therapeutics, including loss of efficacy, altered pharmacokinetics, reduced stability or product shelf life, and induction of undesirable immunogenicity. Bioavailability and pharmacokinetics of

a self-associating peptide can be influenced by aggregate size and the ease of disruption of the noncovalent intermolecular interactions at the subcutaneous site (Maji, S.K., et al. (2008) PLoS Biol 6: e17). In some instances peptides can aggregate into subcutaneous depots that disassociate with $t_{1/2}$ of 30 or more days. Such slow dissolution can lead to favorable effects such as delivery for one month from a single sc injection, causes such a low blood concentration that the peptide appears inactive in vivo. Thus hydrophobic aggregation can appear to totally preclude a peptide's bioavailability and effectiveness (Clodfelter, D.K., et al. (1998) Pharm Res 15: 254-262).

[0106] Aggregation has been associated with increased immunogenicity of the administered peptide and/or protein therapeutic. One means to avoid this problem is to work with solutions of lower concentration, however concentrated peptide and protein solutions are desirable in some instances for ease of administration. In some instances, adding polyols (e.g., mono- or oligosaccharides) or alkyl glycosides to the peptide and/or protein solutions during the course of purification and concentration, reduces or eliminates aggregation, providing greater efficiency in the manufacturing process, and providing a final product which has less immunogenic potential.

[0107] The FDA and other regulatory agencies have increased their scrutiny of aggregation, especially because of this potential linkage to undesirable immunogenicity. The immunogenicity of a self-associating peptide can be influenced by the formation of aggregates as a result of noncovalent intermolecular interactions. For example, interferon has been shown to aggregate resulting in an antibody response (Hermeling, S., et al. (2006) J Pharm Sci 95: 1084-1096). An antibody response to erythropoietin produced "pure red cell aplasia", a potentially life threatening side effect, in a number of patients receiving recombinant EPO (Casadevall, N., et al. (2002) N Engl J Med 346: 469-475) following a change in formulation that altered the serum albumin source and concentration. Insulin loses activity due to protein aggregation upon agitation at temperatures above those found in refrigerated storage (Pezron, I., et al. (2002) J Pharm Sci 91: 1135-1146, Sluzky, V., et al. (1991) Proc Natl Acad Sci U S A 88: 9377-9381). Monoclonal antibody based therapeutics are subject to inactivation as a result of protein aggregation (King, H.D., et al. (2002) J Med Chem 45: 4336-4343). Highly concentrated monoclonal antibody formulations pose stability, manufacturing, and delivery challenges related to the potential of those antibodies to aggregate. Enzymes may also lose activity as a result of aggregation. For example thermal inactivation of urokinase is reported to occur via aggregation (Porter, W.R., et al. (1993) Thromb Res 71: 265-279).

[0108] Protein stabilization during lyophilization has also posed problems. Protein therapeutics frequently lose biological activity after lyophilization and reconstitution as a result of aggregate formation and precipitation. In some instances, addition of reconstitution additives (including, for example, sulfated polysaccharides, polyphosphates, amino acids, polyethylene glycol (PEG) and various surfactants (Zhang, M.Z., et al. (1995) Pharm Res 12: 1447-1452, Vrkljan, M., et al. (1994) Pharm Res 11: 1004-1008) reduces aggregation. In some cases, a combination of alcohols, or other organic solvents, is used for solubilization. Trifluoroethanol has an effect on maintaining peptide structure and it has been used in mixtures to stabilize

various peptides (Roccatano, D., et al. (2002) *Proc Natl Acad Sci U S A* 99: 12179-12184). There is a danger that such agents may have a harsh effect on mucosal tissue, causing patient discomfort or local toxic effects. US Patent 7,390,788 and US Patent 7,425,542 describe the use of alkyl glycosides as stabilizers because of their gentle, non-ionic detergent-like effect. However covalent incorporation of alkyl glycosides into a peptide and/or protein structure itself has not been described heretofore.

[0109] Often naturally occurring oligosaccharides that are covalently attached to proteins do not have surfactant character. In some embodiments, peptide and/or protein products described herein have a covalently attached saccharide and an additional hydrophobic group that confers surfactant character to the modified peptides, thereby allowing for tunability of bioavailability, immunogenicity, and/or pharmacokinetic behavior of the surfactant-modified peptides.

[0110] Proteins and peptides modified with oligosaccharides are described in, for example, Jensen, K.J. and Brask, J. (2005) *Biopolymers* 80: 747-761, through incorporation of saccharide or oligosaccharide structures using enzymatic (Gijssen, H.J., et al. (1996) *Chem Rev* 96: 443-474; Sears, P. and Wong, C.H. (1998) *Cell Mol Life Sci* 54: 223-252; Guo, Z. and Shao, N. (2005) *Med Res Rev* 25: 655-678) or chemical approaches (Urge, L., et al. (1992) *Biochem Biophys Res Commun* 184: 1125-1132; Salvador, L.A., et al. (1995) *Tetrahedron* 51: 5643-5656; Kihlberg, J., et al. (1997) *Methods Enzymol* 289: 221-245; Gregoriadis, G., et al. (2000) *Cell Mol Life Sci* 57: 1964-1969; Chakraborty, T.K., et al. (2005) *Glycoconj J* 22: 83-93; Liu, M., et al. (2005) *Carbohydr Res* 340: 2111-2122; Payne, R.J., et al. (2007) *J Am Chem Soc* 129: 13527-13536; Pedersen, S.L., et al. (2010) *Chembiochem* 11: 366-374). Peptides as well as proteins have been modified by glycosylation (Filira, F., et al. (2003) *Org Biomol Chem* 1: 3059-3063); (Negri, L., et al. (1999) *J Med Chem* 42: 400-404); (Negri, L., et al. (1998) *Br J Pharmacol* 124: 1516-1522); Rocchi, R., et al. (1987) *Int J Pept Protein Res* 29: 250-261; Filira, F., et al. (1990) *Int J Biol Macromol* 12: 41-49; Gobbo, M., et al. (1992) *Int J Pept Protein Res* 40: 54-61; Urge, L., et al. (1992) *Biochem Biophys Res Commun* 184: 1125-1132; Djedaini-Pilard, F., et al. (1993) *Tetrahedron Lett* 34: 2457 - 2460; Drouillat, B., et al. (1997) *Bioorg Med Chem Lett* 7: 2247-2250; Lohof, E., et al. (2000) *Angew Chem Int Ed Engl* 39: 2761-2764; Gruner, S.A., et al. (2001) *Org Lett* 3: 3723-3725; Pean, C., et al. (2001) *Biochim Biophys Acta* 1541: 150-160; Filira, F., et al. (2003) *Org Biomol Chem* 1: 3059-3063; Grotenbreg, G.M., et al. (2004) *J Org Chem* 69: 7851-7859; Biondi, L., et al. (2007) *J Pept Sci* 13: 179-189; Koda, Y., et al. (2008) *Bioorg Med Chem* 16: 6286-6296; Lowery J.J., et al. (2011) *J Pharmacol Exptl Therap* 336: 767-78; Yamamoto, T., et al. (2009) *J Med Chem* 52: 5164-5175).

[0111] However, the aforementioned attempts do not describe an additional hydrophobic group attached to the peptide-linked oligosaccharide. Accordingly, provided herein are modified peptides and/or proteins that incorporate a hydrophobic group attached to a saccharide and/or oligosaccharide that is covalently attached to the peptide and/or protein and that allow for tunability of bioavailability, immunogenicity and pharmacokinetic behaviour. Accordingly, also provided herein are surfactant reagents comprising an oligosaccharide and a hydrophobic

group, that allow for modification of peptide and/or proteins.

[0112] Provided herein is the use of saccharide-based surfactants in covalent linkage to a peptide for improvement of peptide and/or protein properties. In some embodiments, surfactant modification (e.g., covalent attachment of alkyl glycoside class of surfactants) of peptides and/or proteins as described herein, increases the transport across mucosal barriers. In some embodiments, covalent attachment of a surfactant to a peptide and/or protein product prevents aggregation of the peptide and/or protein.

[0113] The surfactant-modified peptides and/or proteins described herein overcome limitations of peptide pharmaceuticals including and not limited to short duration of action, poor bioavailability, aggregation, immunogenicity and lack of receptor subtype specificity through the covalent incorporation of surfactants such as alkyl glycosides as novel peptide and protein modifiers.

[0114] In certain instances, the effects of surfactants are beneficial with respect to the physical properties or performance of pharmaceutical formulations, but are irritating to the skin and/or other tissues and in particular are irritating to mucosal membranes such as those found in the nose, mouth, eye, vagina, rectum, buccal or sublingual areas. Additionally, in some instances, surfactants denature proteins thus destroying their biological function. Since surfactants exert their effects above the critical micelle concentration (CMC), surfactants with low CMC's are desirable so that they may be utilized with effectiveness at low concentrations or in small amounts in pharmaceutical formulations. Accordingly, in some embodiments, surfactants (e.g., alkyl glycosides) suitable for peptide modifications described herein have the CMC's less than about 1 mM in pure water or in aqueous solutions. By way of example only, certain CMC values for alkyl glycosides in water are: Octyl maltoside 19.5 mM; Decyl maltoside 1.8 mM; Dodecyl- β -D-maltoside 0.17 mM; Tridecyl maltoside 0.03 mM; Tetradecyl maltoside 0.01 mM; Sucrose dodecanoate 0.3 mM. It will be appreciated that a suitable surfactant could have a higher or lower CMC depending on the peptide and /or protein that is modified. As used herein, "Critical Micelle Concentration" or "CMC" is the concentration of an amphiphilic component (alkyl glycoside) in solution at which the formation of micelles (spherical micelles, round rods, lamellar structures etc.) in the solution is initiated. In certain embodiments, the alkyl glycosides dodecyl, tridecyl and tetradecyl maltoside or glucoside as well as sucrose dodecanoate, tridecanoate, and tetradecanoate are possess lower CMC's and are suitable for peptide and/or protein modifications described herein.

Opioid peptides and analogs

[0115] In some embodiments, a peptide therapeutic class amenable to the methods of peptide modifications described herein is that of the peptide opioids. This class derives from the endogenous peptide opioids which have a very broad range of functions in the body, carried out through binding to the mu (MOR), delta (DOR), and kappa (KOR) opioid receptors (Schiller, P.W. (2005) AAPS J 7: E560-565). Of most interest is their role in modulating and, in

particular, suppressing of transmission and perception of pain signals. In the development of such agents the central side effects (respiratory suppression, place preference indicating reward from self administration) are a major concern, so peripherally acting agents would be attractive (Stein, C., et al. (2009) Brain Res Rev 60: 90-113). Studies from a number of labs have suggested that the optimal class of agents will have mu opioid receptor agonism with the possibility of delta receptor antagonism (Schiller, P.W. (2010) Life Sci 86: 598-603).

[0116] Although the endomorphins (Janecka, A., et al. (2007) Curr Med Chem 14: 3201-3208) are primarily mu receptor specific, judicious modification of the framework can result in molecules with both mu and delta selectivity (Lazarus, L.H. and Okada, Y. (2012) Expert Opin Ther Patents 22: 1-14; Keresztes, A., et al. (2010) ChemMedChem 5: 1176-96). Derived from the dermorphin family are the DALDA class of analogs (Schiller, P.W. (2010) Life Sci 86: 598-603). Also derived from the dermorphin family are the TIPP family of peptides (Schiller, P.W., et al., (1999) Biopolymers 51:411-25). Described herein are certain opioid peptides that are covalently attached to a saccharide group of an alkyl-glycoside surfactant and have improved pharmaceutical properties.

[0117] Some of the exemplary synthetic peptide analogs described herein are derived from endomorphins and some are derived from dermorphin, two classes of native peptide opioid sequences. In one aspect, the present peptide analogs of the native sequences are endorphin related sequences such as illustrated by EU-A101 to EU-A115. In another aspect, the peptide analogs are dermorphin-related sequences such as EU-A107, EU-A108 and EU-A120 to EU-A133. A related class of opioid peptide analogs is illustrated by sequences EU-A134 to EU-A142 wherein a Tic residue replaces the D-Ala residue seen in the dermorphin structure (TIPP family). An additional class of specialized linkage is shown when Tic is replaced by Tic(Ψ [CH₂-NH]) as this requires a complementary residue Ψ -Phe to complete the linkage.

[0118] In some embodiments, a surfactant-modified peptide product has amino acid sequences corresponding to the general Formula III:



wherein:

aa₁ is Tyr, Dmt, N-alkyl-Tyr, N-alkyl-Dmt, N-dialkyl-Tyr, N-dialkyl-Dmt and the like;

aa₂ is Pro, D-Arg, D-Cit, D-U(X), D-Ala, Tic, Tic(Ψ [CH₂-NH]);

aa₃ is Phe, Trp, Tmp, D- or L-Nal(1), D- or L-Nal(2), CaMePhe, Ψ -Phe;

aa₄ is Phe, Tmp, D- or L-Nal(1), D- or L-Nal(2), U(X), D- or L-CaMeU(X);

aa₅ is absent or Pro, Aib, U(X), D- or L-CaMeU(X)

alkyl or dialkyl is independently a substituted or unsubstituted C₁-C₁₀ branched or straight


chain, or substituted or unsubstituted aralkyl chain;

U is a linking amino acid;

X is a functionalized surfactant linked to the side chain of U;

Z is -OH or NH₂.

[0119] In some specific embodiments of Formula III, X has the structure:



wherein:

A is a substituted or unsubstituted C₁-C₃₀ alkyl group;

R^{1b}, R^{1c}, and R^{1d} are H;

W¹ is -(C=O)-NH-;

W² is -O-; and

R² is a bond.

[0120] In one embodiment, N-alkyl is N-methyl; U is a dibasic amino acid, such as Lys or Orn; X is a modified nonionic detergent of the 1-alkyl glycoside class wherein alkyl is C₁-C₂₀ alkyl or an alkoxyaryl substituent wherein the glycosidic linkage to the saccharide ring is through -O- or other heteroatom (e.g., S or N); Z is NH₂.

[0121] In another embodiment, the 1-alkyl group in the 1-alkyl-glycoside is substituted or unsubstituted C₁-C₁₆alkyl; U is Lys, Z is NH₂;

aa₁ is Tyr, Dmt, Nα-Me-Tyr, Nα-Me-Dmt;

aa₂ is Pro;

aa₃ is Phe, Trp, Tmp;

aa₄ is Phe, Lys(X);

aa₅ is absent or Lys(X).

[0122] In a further embodiment, 1-alkyl group in the 1-alkylglycoside is substituted or unsubstituted C₁-C₂₀alkyl; Z is NH₂ ;

aa₁ is Tyr, Dmt;

aa₂ is D-Arg, D-Lys(X), Tic, Tic(Ψ[CH₂-NH]);

aa₃ is Phe, Trp, Tmp, CaMePhe, Ψ-Phe;

aa₄ is Phe, Tmp, Lys(X);

aa₅ is absent or Pro, Aib, Lys(X), D- or L-CaMeLys(X).

[0123] The endomorphin class parent structures are :

Endomorphin 1 - Tyr-Pro-Trp-Phe-NH₂

Endomorphin 2 - Tyr-Pro-Phe-Phe-NH₂

[0124] Certain analog family substitutions are as shown below:

Position 1	Position 2	Position 3	Position 4	Position 5
Tyr	Pro	Phe	Phe	-NH ₂
Dmt		Trp	Tmp	Lys(X)-NH ₂
N-Alkyl-Tyr		Tmp	Lys(X)	Aib-NH ₂
N-Alkyl-Dmt		CaMePhe	CaMeLys(X)	Pro-NH ₂
N-dialkyl-Tyr		D- or L-Nal(1)	Pro	
N-dialkyl-Dmt		D- or L-Nal(2)		

X = 1-alkyl glucuronyl or 1-alkyl mannouronyl substituted with 1-(C₁-C₂₀ alkyl, aryloxyalkyl, and the like)

[0125] Certain sequences contain the following amino acids:

Position 1	Position 2	Position 3	Position 4	Position 5
Tyr	Pro	Phe	Phe	-NH ₂
Dmt		Trp	Tmp	Lys(X)-NH ₂
		Tmp	Lys(X)	

X = 1-alkyl glucuronyl or 1-alkyl mannouronyl substituted with 1-(C₁-C₂₀ alkyl, aryloxyalkyl, and the like)

[0126] In specific embodiments, analogs with attached surfactants include and are not limited to:

	Position 1	Position 2	Position 3	Position 4
EU-A101	Dmt	Pro	Tmp	Lys(C1-glucuronyl)-NH ₂
EU-A102	Dmt	Pro	Tmp	Lys(C8-glucuronyl)-NH ₂
EU-A103	Dmt	Pro	Tmp	Lys(C12-glucuronyl)-NH ₂
EU-A105	Dmt	Pro	Tmp	Phe-Lys(C1-glucuronyl)-NH ₂
EU-A106	Dmt	Pro	Tmp	Phe-Lys(C12-glucuronyl)-NH ₂
EU-A162	Dmt	Pro	Phe	Lys(C1-glucuronyl)-NH ₂
EU-A163	Dmt	Pro	Phe	Lys(C8-glucuronyl)-NH ₂
EU-A164	Dmt	Pro	Phe	Lys(C12-glucuronyl)-NH ₂
EU-A189	Dmt	Pro	Phe	Phe-Lys(C1-glucuronyl)-NH ₂
EU-A190	Dmt	Pro	Phe	Phe-Lys(C12-glucuronyl)-NH ₂
X = 1-alkyl glucuronyl or 1-alkyl mannouronyl substituted with 1-(C ₁ -C ₂₀ alkyl, aryloxyalkyl, and the like)				

[0127] The Dermorphin class parent structure is:

Parent dermorphin - Tyr-D-Ala-Phe-Gly-Tyr-Pro-Ser-NH₂

[0128] Certain analog family structures are as shown below:

Position 1	Position 2	Position 3	Position 4	
Tyr	D-Arg	Phe	Phe	NH ₂
Dmt	D-Lys(X)	Trp	Tmp	Lys(X)-NH ₂
N-Alkyl-Tyr	D-Ala	Tmp	Lys(X)	Pro-NH ₂
N-Alkyl-Dmt	Tic	CaMePhe	CaMeLys(X)	Aib-NH ₂
N-dialkyl-Tyr	Tic-(Ψ[CH ₂ -NH])	Ψ-Phe		
N-dialkyl-Dmt		D-or L-Nal(1)	D- or L-Nal(1)	D- or L-Nal(1)
		D-or L-Nal(2)	D- or L-Nal(2)	D- or L-Nal(2)
X = 1-alkyl glucuronyl or 1-alkyl mannouronyl substituted with 1-(C ₁ -C ₂₀ alkyl, aryloxyalkyl, and the like)				

[0129] In some embodiments, certain analogs suitable for attachment of surfactants include and are not limited to:

Position 1	Position 2	Position 3	Position 4	
Tyr	D-Arg	Phe	Phe	NH ₂
Dmt	D-Lys(X)	Trp	Tmp	Lys(X)-NH ₂

Position 1	Position 2	Position 3	Position 4	
	Tic	Tmp	Lys(X)	Pro-NH ₂
	Tic-(Ψ[CH ₂ -NH])	Ψ-Phe		Aib-NH ₂
		CaMePhe		
		D- or L-Nal(1)	D- or L-Nal(1)	
		D- or L-Nal(2)	D- or L-Nal(2)	

X = 1-alkyl glucuronyl or 1-alkyl mannouronyl substituted with 1-(C₁₋₂₀ alkyl, aryloxyalkyl, and the like)

[0130] In specific embodiments, analogs suitable for attachment of surfactants include and are not limited to:

	Position 1	Position 2	Position 3	Position 4
EU-A107	Dmt	D-Lys(C1-glucuronyl)	Tmp	Phe-NH ₂
EU-A108	Dmt	D-Lys(C12-glucuronyl)	Tmp	Phe-NH ₂
EU-A120	Dmt	D-Lys(C8-glucuronyl)	Nal(1)	Phe-NH ₂
EU-A121	Dmt	D-Lys(C12-glucuronyl)	Nal(1)	Phe-NH ₂
EU-A122	Dmt	D-Lys(C16-glucuronyl)	Nal(1)	Phe-NH ₂
EU-A123	Dmt	D-Lys(C18-glucuronyl)	Nal(1)	Phe-NH ₂
EU-A124	Dmt	D-Lys(C20-glucuronyl)	Nal(1)	Phe-NH ₂
EU-A125	Dmt	D-Lys(C22-glucuronyl)	Nal(1)	Phe-NH ₂
EU-A126	Dmt	D-Lys(C24-glucuronyl)	Nal(1)	Phe-NH ₂
EU-A178	Dmt	Tic	Phe	Lys(C1-glucuronyl)-NH ₂
EU-A179	Dmt	Tic	Phe	Lys(C12-glucuronyl)-NH ₂
EU-A180	Dmt	Tic	Phe	Lys(C8-glucuronyl)-NH ₂
EU-A181	Dmt	Tic	Phe	Lys(C10-glucuronyl)-NH ₂
EU-A182	Dmt	Tic	Phe	Lys(C16-glucuronyl)-NH ₂
EU-A183	Dmt	Tic	Phe	Lys(C18-glucuronyl)-NH ₂
EU-A184	Dmt	Tic	Phe	Lys(C20-glucuronyl)-NH ₂
EU-A189	Dmt	Tic	Phe	Phe-Lys(C1-glucuronyl)-

	Position 1	Position 2	Position 3	Position 4
				NH ₂
EU-A190	Dmt	Tic	Phe	Phe-Lys(C12-glucuronyl)-NH ₂
EU-A191	Dmt	Tic	Phe	Phe-Lys(C8-glucuronyl)-NH ₂
EU-A600	Dmt	Tic	Phe	Lys(C1-glucuronyl)-Aib-NH ₂
EU-A601	Dmt	Tic	Phe	Lys(C8-glucuronyl)-Aib-NH ₂
EU-A603	Dmt	Tic	Phe	Lys(C12-glucuronyl)-Aib-NH ₂
EU-A615	Dmt	Tic	Phe	D-Lys(C1-glucuronyl)-Aib-NH ₂
EU-A616	Dmt	Tic	Phe	D-Lys(C8-glucuronyl)-Aib-NH ₂
EU-A618	Dmt	Tic	Phe	D-Lys(C12-glucuronyl)-NH ₂
EU-A619	Dmt	Tic	Phe	D-Lys(C16-glucuronyl)-NH ₂
EU-A620	Dmt	Tic	Phe	Lys(C1-glucuronyl)-NHCH ₂ Ph
EU-A621	Dmt	Tic	Phe	Lys(C8-glucuronyl)-NHCH ₂ Ph
EU-A623	Dmt	Tic	Phe	Lys(C12-glucuronyl)-NHCH ₂ Ph
EU-A624	Dmt	Tic	Phe	Lys(C16-glucuronyl)-NHCH ₂ Ph
EU-A639	Dmt	Tic	Phe	D-Lys(C1-glucuronyl)-NHCH ₂ Ph
EU-A642	Dmt	Tic	Phe	D-Lys(C12-glucuronyl)-NHCH ₂ Ph
EU-A648	Dmt	Tic	Phe	Lys(C16-glucuronyl)-NH ₂
EU-649	Dmt	Tic	Phe	Lys(C14-glucuronyl)-NH ₂
X = 1-alkyl glucuronyl or 1-alkyl mannouronyl substituted with 1-(C ₁ -C ₂₀ alkyl, aryloxyalkyl, and the like)				

[0131] Described herein are peptide chains substituted in a suitable position by the substitution of the analogs claimed herein by acylation on a linker amino acid, at for example, the ϵ -position of Lys, with fatty acids such as octanoic, decanoic, dodecanoic, tetradecanoic, hexadecanoic, octadecanoic, 3-phenylpropanoic acids and the like, with saturated or unsaturated alkyl chains (Nestor, J.J., Jr. (2009) Current Medicinal Chemistry 16: 4399 - 4418; Zhang, L and Bulaj, G. (2012) Curr Med Chem 19: 1602-18). Non-limiting, illustrative examples of such analogs are:

H-Dmt-Tic-Phe-Lys(N-epsilon-acetyl)-NH₂, (SEQ. ID. NO. 161)

H-Dmt-Tic-Phe-Lys(N-epsilon-dodecanoyl)-NH₂, (SEQ. ID. NO. 162)

H-Dmt-Tic-Phe-Lys(N-epsilon-tetradecanoyl)-NH₂, (SEQ. ID. NO. 163)

H-Dmt-Tic-Phe-Lys(N-epsilon-(gamma-glutamyl)-N-alpha-dodecanoyl)-NH₂, (SEQ. ID. NO. 164)

H-Dmt-Tic-Phe-Lys(N-epsilon-(gamma-glutamyl)-N-alpha-tetradecanoyl)-NH₂, (SEQ. ID. NO. 165)

H-Dmt-Tic-Phe-Lys(N-epsilon-acetyl)-NH-benzyl, (SEQ. ID. NO. 166)

H-Dmt-Tic-Phe-Lys(N-epsilon-dodecanoyl)-NH-benzyl (SEQ. ID. NO. 167), and the like.

[0132] In other cases described herein the peptide chain may be substituted in a suitable position by reaction on a linker amino acid, for example the sulfhydryl of Cys, with a spacer and a hydrophobic moiety such as a steroid nucleus, for example a cholesterol moiety. In some of such embodiments, the modified peptide further comprises one or more PEG chains. Non-limiting examples of such molecules are:

H-Dmt-Tic-Phe-Cys(S-(3-(PEG4-aminoethylacetamide-cholesterol)))-NH₂, (SEQ. ID. NO. 168)

H-Dmt-Tic-Phe-Cys(S-(3-(PEG4-aminoethylacetamide-cholesterol)))-NH-benzyl (SEQ. ID. NO. 169), and the like.

[0133] The compounds of Formula I, Formula II, or Formula III are assayed for mu opioid receptor activity in a cellular assay (MOP in agonist and antagonist mode), and for delta2 opioid receptor activity in cellular assay (DOP in agonist and antagonist mode) as described in Example 12.

[0134] In certain embodiments, as shown in Example 12, a compound having a pure MOP agonist activity along with a pure DOP antagonistic activity, is a suitable profile for clinical applications. Also contemplated within the scope of the disclosure herein are compounds that have low solubility and low apparent in vitro potency but exhibit prolonged duration of action (pharmacodynamic action) in vivo.


[0135] Contemplated within the scope of embodiments presented herein are peptide products of Formula I, Formula II or Formula III, wherein the peptide product comprises one, or, more than one surfactant groups (e.g., group X having the structure of Formula I). In one embodiment, a peptide product of Formula I, Formula II or Formula III, comprises one

surfactant group. In another embodiment, a peptide product of Formula I, Formula II or Formula III, comprises two surfactant groups. In yet another embodiment, a peptide product of Formula I, Formula II or Formula III, comprises three surfactant groups.

PTH peptides and analogs


[0136] Also provided herein, in some embodiments, are reagents and intermediates for synthesis of modified peptides and/or proteins (e.g., modified PTH, PTHrP, or the like) through the incorporation of surfactants.

[0137] Provided herein, in some embodiments, are peptide products comprising a surfactant X, covalently attached to a peptide, the peptide comprising a linker amino acid U and at least one other amino acid:



Formula 2-I-A

wherein the surfactant X is a group of Formula 2-1:



Formula 2-I

wherein:

R^{1a} is independently, at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group, or a steroid nucleus containing moiety;

R^{1b} , R^{1c} , and R^{1d} are each, independently at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group;

W^1 is independently, at each occurrence, $-CH_2-$, $-CH_2-O-$, $-(C=O)-$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$, or $-CH_2-S-$;

W^2 is $-O-$, $-CH_2-$, or $-S-$;

R^2 is independently, at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl

group, $-\text{NH}_2$, $-\text{SH}$, $\text{C}_2\text{-C}_4$ -alkene, $\text{C}_2\text{-C}_4$ -alkyne, $-\text{NH}(\text{C}=\text{O})\text{-CH}_2\text{-Br}$, $-(\text{CH}_2)_m\text{-maleimide}$, or $-\text{N}_3$;

n is 1, 2 or 3; and

m is 1-10;

the peptide is selected from Formula 2-II:

$\text{aa}_1\text{-Val}_2\text{-aa}_3\text{-Glu}_4\text{-aa}_5\text{-aa}_6\text{-aa}_7\text{-aa}_8\text{-His}_9\text{-aa}_{10}\text{-aa}_{11}\text{-aa}_{12}\text{-aa}_{13}\text{-aa}_{14}\text{-aa}_{15}\text{-aa}_{16}\text{-aa}_{17}\text{-}$

$\text{aa}_{18}\text{-aa}_{19}\text{-aa}_{20}\text{-aa}_{21}\text{-aa}_{22}\text{-aa}_{23}\text{-aa}_{24}\text{-aa}_{25}\text{-aa}_{26}\text{-Z}$ Formula 2-II (SEQ. ID. NO.

170)

wherein:

Z is OH , or $-\text{NH-R}^3$,

R^3 is H , a substituted or unsubstituted $\text{C}_1\text{-C}_{12}$ alkyl, or a PEG chain of less than 10 Da;

aa_1 is Aib, Ac5c, or Deg;

aa_3 is Aib, Ac4c, or Deg;

aa_5 is His, or Ile;

aa_6 is Gln, or Cit;

aa_7 is Leu, or Phe;

aa_8 is Leu, or Nle;

aa_{10} is Asp, Asn, Gln, Glu, Cit, Ala, or Aib;

aa_{11} is Arg, or hArg;

aa_{12} is Gly, Glu, Lys, Ala, Aib, or Ac5c;

aa_{13} is Lys, or Arg;

aa_{14} is Ser, His, Trp, Phe, Leu, Arg, Lys, Glu, or Nal(2);

aa_{15} is Ile, Leu, or Aib;

aa_{16} is Gln, Asn, Glu, Lys, Ser, Cit, Aib, or U;

aa_{17} is Asp, Ser, Aib, Ac4c, Ac5c, or U;

aa_{18} is absent or Leu, Gln, Cit, Aib, Ac5c, Lys, Glu or U;

aa_{19} is absent or Arg, Glu, Aib, Ac4c, Ac5c, or U;

aa₂₀ is absent or Arg, Glu, Lys, Aib, Ac4c, Ac5c, or U;

aa₂₁ is absent or Arg, Val, Aib, Ac5C, Deg, or U;

aa₂₂ is absent or Phe, Glu, Aib, Ac5C, Lys, or U;

aa₂₃ is absent or Leu, Phe, Trp, or U;

aa₂₄ is absent or His, Arg, or U;

aa₂₅ is absent or His, Lys, or U and

aa₂₆ is absent or Aib, Ac5c, Lys;


U is a natural or unnatural amino acid comprising a functional group used for covalent attachment to the surfactant X;

wherein any two of aa₁-aa₂₆ are optionally cyclized through their side chains to form a lactam linkage; and

provided that one, or at least one of aa₁₆ - aa₂₆ is the linker amino acid U covalently attached to X.

[0138] In some embodiments, n is 1. In some embodiments, n is 2, and a first glycoside is attached to a second glycoside via a bond between W² of the first glycoside and any one of OR^{1b}, OR^{1c} or OR^{1d} of the second glycoside. In some embodiments, n is 3, and a first glycoside is attached to a second glycoside via a bond between W² of the first glycoside and any one of OR^{1b}, OR^{1c} or OR^{1d} of the second glycoside, and the second glycoside is attached to a third glycoside via a bond between W² of the second glycoside and any one of OR^{1b}, OR^{1c} or OR^{1d} of the third glycoside.

[0139] In one embodiment, compounds of Formula I-A are compounds wherein X has the structure:



Formula 2-I

wherein:

R^{1a} is H, a protecting group, a substituted or unsubstituted C₁-C₃₀ alkyl group, or a steroid nucleus containing moiety;

R^{1b} , R^{1c} , and R^{1d} are each, independently at each occurrence, H, a protecting group, or a substituted or unsubstituted C_1 - C_{30} alkyl group;


W^1 is independently, at each occurrence, $-CH_2-$, $-CH_2-O-$, $-(C=O)$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$, or $-CH_2-S-$;

W^2 is $-O-$ or $-S-$;

R^2 is a bond, C_2 - C_4 -alkene, C_2 - C_4 -alkyne, or $-(CH_2)_m$ -maleimide; and

m is 1-10.


[0140] In another embodiment, compounds of Formula I-A are compounds wherein X has the structure:



[0141] Accordingly, in the embodiment described above, R^2 is a bond.


[0142] For instance, in an exemplary embodiment of the structure of X described above, W^1 is $C(=O)NH-$, R^2 is a bond between W^1 and an amino acid residue U within the peptide (e.g., an amino group in the sidechain of a lysine residue present in the peptide).

[0143] In a further embodiment, compounds of Formula I-A are compounds wherein X has the structure:



[0144] For instance, in an exemplary embodiment of the structure of X described above, W^1 is CH_2- and R^2 is an alkyl-linked maleimide functional group on X and R^2 is attached to a suitable moiety of an amino acid residue U within the peptide (e.g., a thiol group in a cysteine residue of the peptide forms a thioether with the maleimide on X).

[0145] In yet another embodiment, compounds of Formula I-A are compounds wherein X has the structure:



Formula 2-I

wherein:

R^{1a} is H, a protecting group, a substituted or unsubstituted C₁-C₃₀ alkyl group, or a steroid nucleus containing moiety;


R^{1b}, R^{1c}, and R^{1d} are each, independently at each occurrence, H, a protecting group, or a substituted or unsubstituted C₁-C₃₀ alkyl group;

W¹ is -(C=O)-NH-;

W² is -O-;

R² is a bond.

[0146] In an additional embodiment, compounds of Formula I-A are compounds wherein X has the structure:



Formula 2-I

wherein:

R^{1a} is a substituted or unsubstituted C₁-C₃₀ alkyl group;

R^{1b}, R^{1c}, and R^{1d} are H;

W¹ is -(C=O)-NH-;


W² is -O-; and

R² is a bond.

[0147] In some embodiments described above and herein, R^{1a} is a substituted or unsubstituted C_1 - C_{30} alkyl group.

[0148] In some embodiments described above and herein, R^{1a} is a substituted or unsubstituted C_6 - C_{20} alkyl group.

[0149] Also contemplated herein are alternate embodiments wherein X in Formula 2-I-A has the structure:



[0150] For instance, in an exemplary embodiment of the structure of X described above, W^1 is -S-, R^2 is a C_1 - C_{30} alkyl group, W^2 is S, R^{1a} is a bond between W^2 and a suitable moiety of an amino acid residue U within the peptide (e.g., a thiol group in a cysteine residue of the peptide forms a thioether with X).

[0151] In another exemplary alternate embodiment of the structure of X described above, W^1 is -O-, R^2 is a C_1 - C_{30} alkyl group, W^2 is O, R^{1a} is a bond between W^2 and a suitable moiety of an amino acid residue U within the peptide (e.g., a hydroxyl group in a serine or threonine residue of the peptide forms an ether with X).

[0152] In some embodiments, U is used for covalent attachment to X and is a dibasic natural or unnatural amino acid, a natural or unnatural amino acid comprising a thiol, an unnatural amino acid comprising a $-N_3$ group, an unnatural amino acid comprising an acetylenic group, or an unnatural amino acid comprising a $-NH-C(=O)-CH_2-Br$ or a $-(CH_2)_m$ -maleimide, wherein m is 1-10.

[0153] In some embodiments of the peptide product, the surfactant is a 1-alkyl glycoside class surfactant. In some embodiments of the peptide product, the surfactant is attached to the peptide via an amide bond.

[0154] In some embodiments of the peptide product, the surfactant X is comprised of 1-eicosyl beta-D-glucuronic acid, 1-octadecyl beta-D-glucuronic acid, 1-hexadecyl beta-D-glucuronic acid, 1-tetradecyl beta-D-glucuronic acid, 1-dodecyl beta-D-glucuronic acid, 1-decyl beta-D-glucuronic acid, 1-octyl beta-D-glucuronic acid, 1-eicosyl beta-D-diglucuronic acid, 1-octadecyl beta-D-diglucuronic acid, 1-hexadecyl beta-D-diglucuronic acid, 1-tetradecyl beta-D-diglucuronic acid, 1-dodecyl beta-D-diglucuronic acid, 1-decyl beta-D-diglucuronic acid, 1-octyl beta-D-diglucuronic acid, or functionalized 1-eicosyl beta-D-glucose, 1-octadecyl beta-D-glucose, 1-hexadecyl beta-D-glucose, 1-tetradecyl beta-D-glucose, 1-dodecyl beta-D-glucose,

1-decyl beta-D-glucose, 1-octyl beta-D-glucose, 1-eicosyl beta-D-maltoside, 1-octadecyl beta-D-maltoside, 1-hexadecyl beta-D-maltoside, 1-dodecyl beta-D-maltoside, 1-decyl beta-D-maltoside, 1-octyl beta-D-maltoside, and the like, and the peptide product is prepared by formation of a linkage between the aforementioned groups and a group on the peptide (e.g., a -COOH group in the aforementioned groups and an amino group of the peptide).

[0155] In some embodiments of the peptide product, U is a terminal amino acid of the peptide. In some embodiments of the peptide product, U is a non-terminal amino acid of the peptide. In some embodiments of the peptide product, U is a natural D- or L- amino acid. In some embodiments of the peptide product, U is an unnatural amino acid. In some embodiments of the peptide product, U is selected from Lys, Cys, Orn, or an unnatural amino acid comprising a functional group used for covalent attachment to the surfactant X.

[0156] In some embodiments of the peptide product, the functional group used for covalent attachment of the peptide to the surfactant X is -NH₂, -SH, -OH, -N₃, haloacetyl, a -(CH₂)_m-maleimide (wherein m is 1-10), or an acetylenic group.

[0157] In some embodiments side chain functional groups of two different amino acid residues are linked to form a cyclic lactam. For example a Lys₁₄ side chain may form a cyclic lactam with the side chain of Glu₁₈ or a Lys₁₈ may form a lactam with the side chain of a Glu₂₂. In some embodiments such lactam structures are reversed and are formed from a Glu₁₄ and a Lys₁₈, for example. Such lactam linkages, in some instances, stabilize alpha helical structures in peptides (Condon, S.M., et al. (2002) Bioorg Med Chem 10: 731-736).

[0158] In some embodiments, the peptide product comprising a covalently linked alkyl glycoside is a covalently modified PTH or analog thereof. In some of such embodiments, the peptide product contains a covalently linked 1-O-alkyl β-D-glucuronic acid and the peptide is an analog of PTH.

[0159] In some embodiments, a peptide product comprising a covalently linked alkyl glycoside is a covalently modified PTHrP, or analog thereof. In some of such embodiments, the peptide product comprises a covalently linked 1-O-alkyl β-D-glucuronic acid and the peptide is an analog of PTHrP.

[0160] In some embodiments, the peptide product has the structure of Formula 2-III:

aa₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Lcu₇- Nlc₈- His₉- Gln₁₀- hArg₁₁- aa₁₂- Arg₁₃- aa₁₄- Ile₁₅-
aa₁₆- aa₁₇- aa₁₈- aa₁₉- aa₂₀- aa₂₁- aa₂₂- aa₂₃- aa₂₄- aa₂₅- aa₂₆-Z

Formula 2-III

(SEQ. ID. NO. 171)

wherein:

Z is OH or -NH₂;

aa₁ is Aib, or Ac5c;

aa₁₂ is Ala, Glu, Lys, Aib, or Ac5c;

aa₁₄ is Trp, Phe, Lys, Glu or Nal(2);

aa₁₆ is Gln, Asn, Glu, Lys, Cit, or U(X);

aa₁₇ is Asp, Ser, Aib, Ac4c, Ac5C or U(X);

aa₁₈ is absent or Leu, Gln, Aib, Lys, Glu or U(X);

aa₁₉ is absent or Arg, Glu, Aib, Ac4c or Ac5c;

aa₂₀ is absent or Arg, Glu, Lys, Aib, Ac4c, Ac5c;

aa₂₁ is absent or Arg, Val, Aib, Ac5C, or Deg;

aa₂₂ is absent or Phe, Glu, Lys or U(X);

aa₂₃ is absent or Leu, Phe, Trp or U(X);

aa₂₄ is absent or His, Arg, or U(X);

aa₂₅ is absent or His, Lys, or U(X); and

aa₂₆ is absent or Aib, Ac5c;

wherein any two of aa₁-aa₂₆ are optionally cyclized through their side chains to form a lactam linkage; and

provided that one, or at least one of aa₁₆, aa₁₇, aa₁₈, aa₂₂, aa₂₃, aa₂₄ or aa₂₅ is the linker amino acid U covalently attached to X.

[0161] In some embodiments of Formula 2-III, U is any linker amino acid described herein. In some embodiments, the compound of Formula 2-III is a compound wherein aa₁₂ and aa₁₆ are cyclized through their side chains to form a lactam linkage. In some embodiments, the compound of Formula 2-III is a compound wherein aa₁₆ and aa₂₀ are cyclized through their side chains to form a lactam linkage.

[0162] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅- Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-dodecyl beta-D-glucuronyl)₁₈-NH₂. (**SEQ. ID. NO. 180**)

[0163] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅- Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-alkyl beta-D-glucuronyl)₁₈-Aib₁₉-NH₂, wherein alkyl is dodecyl, tetradecyl, hexadecyl, or octadecyl. (**SEQ. ID. NO. 283**)

[0164] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-alkyl beta-D-glucuronyl)₁₈-Ac4c₁₉-NH₂, wherein alkyl is dodecyl, tetradecyl, hexadecyl, or octadecyl. (**SEQ. ID. NO. 284**)

[0165] In some embodiments, the peptide product has the structure:

Ac5c₁-Val₂-Aib₃-Glu₄-Ile₄-Gln₆-Leu₇-Nle₈-His₉-Gln₁₀-hArg₁₁-Ala₁₂-Arg₁₃-Trp₁₄-Ile₁₅-Glu*₁₆-Aib₁₇-Lys(N-epsilon-1'-alkyl beta-D-glucuronyl)₁₈-Aib₁₉-Lys*₂₀-NH₂, wherein Glu*₁₆ and Lys*₂₀ are linked through their sidechains by a lactam and alkyl is dodecyl, tetradecyl, hexadecyl, or octadecyl. (**SEQ. ID. NO. 285**)

[0166] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Glu*₁₂- Arg₁₃-Trp₁₄- Ile₁₅- Lys*₁₆- Aib₁₇-Lys(N-epsilon-1'-alkyl beta-D-glucuronyl)₁₈-Aib₁₉-NH₂, wherein Glu*₁₂ and Lys*₁₆ are linked through their sidechains by a lactam and alkyl is dodecyl, tetradecyl, hexadecyl, or octadecyl. (**SEQ. ID. NO. 286**)

[0167] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Phe₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-alkyl beta-D-glucuronyl)₁₈-Aib₁₉-NH₂, wherein alkyl is dodecyl, tetradecyl, hexadecyl, or octadecyl. (**SEQ. ID. NO. 287**)

[0168] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-alkyl beta-D-glucuronyl)₁₈-Aib₁₉-Aib₂₀-NH₂, wherein alkyl is dodecyl, tetradecyl, hexadecyl, or octadecyl. (**SEQ. ID. NO. 288**)

[0169] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Glu*₁₂- Arg₁₃-Trp₁₄- Ile₁₅- Lys*₁₆- Aib₁₇-Lys(N-epsilon-1'-alkyl beta-D-glucuronyl)₁₈-Aib₁₉-Aib₂₀-NH₂, wherein Glu*₁₂ and Lys*₁₆ are linked through their sidechains by a lactam and alkyl is dodecyl, tetradecyl, hexadecyl, or octadecyl. (**SEQ. ID. NO. 289**)

[0170] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-dodecyl beta-D-glucuronyl)₁₈-Aib₁₉-NH₂. (**SEQ. ID. NO. 207**)

[0171] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-tetradecyl beta-D-glucuronyl)₁₈-Aib₁₉-NH₂. (**SEQ. ID. NO. 260**)

[0172] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-hexadecyl beta-D-glucuronyl)₁₈-Aib₁₉-NH₂. (**SEQ. ID. NO. 261**)

[0173] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-octadecyl beta-D-glucuronyl)₁₈-Aib₁₉-NH₂. (**SEQ. ID. NO. 262**)

[0174] In some embodiments, the peptide product has the structure:


Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-dodecyl beta-D-glucuronyl)₁₈-Aib₁₉-Aib₂₀-NH₂. (**SEQ. ID. NO. 275**)

[0175] In some embodiments, the peptide product has the structure:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-1'-hexadecyl beta-D-glucuronyl)₁₈-Aib₁₉-Aib₂₀-NH₂. (**SEQ. ID. NO. 277**)


[0176] In some embodiments, the peptide product is a biologically active peptide product that binds to the PTH receptor (PTHR1).

[0177] In a specific embodiment, the peptide products of Formula 2-I-A described above and herein have the following structure:




wherein R^{1a} is a C₁-C₂₀ alkyl chain as described in Table 2 of Figure 2, R' is a peptide as described in Table 2 of Figure 2, W² of Formula I-A is -O-, and W¹ of Formula 2-I-A is (C=O)NH- and is part of an amide linkage to the peptide R'. In some of such embodiments, R^{1a} is a C₆-C₂₀ alkyl chain. In some of such embodiments, R^{1a} is a C₈-C₂₀ alkyl chain. In some of such embodiments, R^{1a} is a C₈-C₂₀ alkyl chain. In some of such embodiments, R^{1a} is a C₈-C₁₈ alkyl chain. In some of such embodiments, R^{1a} is a C₈-C₁₆ alkyl chain.

[0178] In embodiments described above, an amino moiety of an amino acid and/or a peptide R' (e.g., an amino group of an amino acid residue such as a Lysine, or a lysine within the peptide R') is used to form a covalent linkage with a compound of structure:




wherein R^{1a} is a C₁-C₂₀ alkyl chain as described in Table 2 of Figure 2.

[0179] In such cases, the amino acid having an amino moiety (e.g., a Lysine residue within the peptide R') which is used to form a covalent linkage to the compound A described above, is a linker amino acid U which is attached to a surfactant X having the structure of Formula 2-A. Accordingly, as one example, Lys(C12) of Table 2 of Figure 2 has the following structure:



[0180] Also contemplated within the scope of the embodiments presented herein are peptide products of Formula 2- I-A derived from maltouronic acid-based surfactants through binding at either or both carboxylic acid functions. Thus, as one example, peptides in Table 2 of Figure 2 comprise a lysine linker amino acid bonded to a maltouronic acid based surfactant X and having a structure:



[0181] It will be understood that in one embodiment, compounds of Formula 2-I-A are prepared by attaching a lysine to a group X, followed by attachment of additional amino acid residues and/or peptides are attached to the lysine-X compound to obtain compounds of Formula 2-I-A. It will be understood that other natural or non-natural amino acids described herein are also suitable for attachment to the surfactant X and are suitable for attaching additional amino acid/peptides to obtain compounds of Formula 2-I-A. It will be understood that in another embodiment, compounds of Formula 2-I-A are prepared by attaching a full length or partial length peptide to a group X, followed by optional attachment of additional amino acid residues and/or peptides are attached to obtain compounds of Formula 2-I-A.

[0182] In a specific embodiment, provided herein are compounds selected from compounds of Table 2 in Figure 2.

[0183] Also provided herein are pharmaceutical compositions comprising a therapeutically effective amount of a peptide product described above, or acceptable salt thereof, and at least one pharmaceutically acceptable carrier or excipient.

[0184] In some embodiments of the pharmaceutical compositions, the carrier is an aqueous-based carrier. In some embodiments of the pharmaceutical compositions, the carrier is a nonaqueous-based carrier. In some embodiments of the pharmaceutical compositions, the nonaqueous-based carrier is a hydrofluoroalkane-like solvent comprising sub-micron anhydrous α -lactose or other excipients.

[0185] Contemplated within the scope of embodiments presented herein is the reaction of an amino acid and/or a peptide comprising a linker amino acid U bearing a nucleophile, and a group X comprising a leaving group or a functional group that can be activated to contain a leaving group, for example a carboxylic acid, or any other reacting group, thereby allowing for covalent linkage of the amino acid and/or peptide to a surfactant X via the linker amino acid U to provide a peptide product of Formula 2-I-A.

[0186] Also contemplated within the scope of embodiments presented herein is the reaction of an amino acid and/or a peptide comprising a linker amino acid U bearing a leaving group or a functional group that can be activated to contain a leaving group, for example a carboxylic acid, or any other reacting group, and a group X comprising a nucleophilic group, thereby allowing for covalent linkage of the amino acid and/or peptide to a surfactant X via the linker amino acid U to provide a peptide product of Formula 2-I-A.

[0187] It will be understood that, in one embodiment, Compounds of Formula 2-I-A are prepared by reaction of a linker amino acid U with X, followed by addition of further residues to U to obtain the peptide product of Formula 2-I-A. It will be understood that in an alternative embodiment, Compounds of Formula 2-I-A are prepared by reaction of a suitable peptide comprising a linker amino acid U with X, followed by optional addition of further residues to U, to obtain the peptide product of Formula 2-I-A.

[0188] Provided herein are methods of treating hypoparathyroidism comprising administering to a subject in need thereof a therapeutically effective amount of a peptide product described above. In some embodiments, the hypoparathyroidism is associated with bone mass reduction.

[0189] Also provided herein are methods of stimulating bone repair or favoring the engraftment of a bone implant comprising administering to a subject in need thereof a therapeutically effective amount of a peptide product described above.

[0190] Also provided herein is a covalently modified PTH or PTHrP peptide or analog thereof, comprising a hydrophilic group as described herein; and a hydrophobic group covalently

attached to the hydrophilic group. In specific embodiments, the covalently modified peptide and/or protein product comprises a hydrophilic group that is a saccharide and a hydrophobic group that is a C₁-C₂₀ alkyl chain or an aralkyl chain.

[0191] In one embodiment, provided is a method for chemically modifying a molecule by covalent linkage to a surfactant to increase or sustain the biological action of the composition or molecule, for example, receptor binding or enzymatic activity. In some embodiments, the molecule is a peptide. The method additionally can include further modification comprising covalent attachment of the molecule in the composition to a polymer such as polyethylene glycol.

[0192] In another embodiment, provided is a method of reducing or eliminating immunogenicity of a peptide and/or protein drug by covalently linking the peptide chain to at least one alkyl glycoside wherein the alkyl has from 1 to 30 carbon atoms.

[0193] Also provided is a method of treating hypoparathyroidism, osteoporosis, osteopenia, post-menopausal osteoporosis, Paget's disease, glucocorticoid induced osteoporosis, old age osteoporosis, humoral hypercalcemia, or the like comprising administering a drug composition comprising a peptide covalently linked to at least one alkyl glycoside and delivered to a vertebrate, wherein the alkyl has from 1 to 30 carbon atoms, or further in the range of 6 to 16 carbon atoms, and wherein covalent linkage of the alkyl glycoside to the peptide increases the stability, bioavailability and/or duration of action of the drug.

[0194] In some embodiments, the covalently modified peptides and/or proteins are covalently modified PTH or PTHrP, or analogs thereof, which are modified to improve their pharmaceutical and medical properties by covalent modification with alkyl glycoside surfactant moieties. These surfactant-modified analogs have increased steric hindrance that hinder proteolysis, slows uptake and slows clearance from the body.

[0195] Some studies show that approximately 50 percent of patients suffering from osteoporosis discontinue oral bisphosphonate therapy within the first year. Among patients who discontinue these treatments, many do so because of side effects including intolerance. Although truncated recombinant parathyroid hormone 1-34 (rhPTH1-34) is available commercially as a bone anabolic agent (Brixen, K.T., et al. (2004) Basic Clin Pharmacol Toxicol 94: 260-270; Dobnig, H. (2004) Expert Opin Pharmacother 5: 1153-1162) as teriparatide (Lilly), poor compliance is a major problem. More recently an antibody (denosumab, Amgen) against the ligand controlling osteoclast function has been approved, but it has significant known side effects (serious skin infections, observed cases of osteonecrosis of the jaw and significant suppression of bone remodeling), some of which may have as yet unclear long-term effects.

Bone Structure

[0196] The architecture of bone in man is maintained and elaborated by the coordinated

function of osteoclasts, which cause bone resorption, and osteoblasts, which lay down new bone matrix. Bone is an important depot for storage of calcium (a critical signaling ion) in the body and a decrease in ambient extracellular Ca level causes an increase in PTH secretion in via the Ca-sensing receptors on the parathyroid cellular membrane. PTH binds to its receptor (PTHr1), present on osteoblasts cell membranes, leading to expression of the "ligand of receptor activator of nuclear factor- κ B" (RANKL). RANKL binds to its receptor, RANK, on osteoclasts precursors, stimulating their differentiation and proliferation (Boyce, B.F. and Xing, L. (2007) *Arthritis Res Ther* 9 Suppl 1: S1). This leads to bone resorption, mobilization of calcium from the bone. PTH also acts to increase renal tubular Ca reabsorption and indirectly to enhance intestinal Ca absorption via its stimulatory action on renal 1- α cholecalciferol hydroxylase (increasing circulating calcitriol). Both actions serve to provide a longer term increase in circulating calcium ion.

[0197] The native form of human parathyroid hormone (hPTH) is an 84-amino acid peptide that plays an important role in the maintenance of Calcium homeostasis in mammals (Rosen, C.J. and Bilezikian, J.P. (2001) *J Clin Endocrinol Metab* 86: 957-964). A structurally-related but independent hormone, parathyroid hormone-related protein (PTHrP), plays a paracrine role, focused on bone growth in local tissue. Both hormones bind to the same receptor on osteoblasts, PTHr1, and cause activation of multiple signaling pathways, including that regulated by increased cAMP levels.

[0198] However intermittent presence of PTH(1-34) leads only to stimulation of osteoblasts, lack of RANKL expression, and increased bone density. PTH(1-34) exhibits potent anabolic effects on the skeleton when given exogenously by intermittent injection. A small group of patients received teriparatide by daily sc injections for 6-24 months (Reeve, J., et al. (1980) *Br Med J* 280: 1340-1344) and paired bone biopsies revealed substantial increases in iliac trabecular bone volume, with evidence of new bone formation and a suggestion that there was a dissociation between bone formation and resorption rates. Numerous studies have confirmed improvements in bone tissue after daily injections of PTH analogs (Hodsman, A.B., et al. (2005) *Endocr Rev* 26: 688-703; Cheng, Z., et al. (2009) *J Bone Miner Res* 24: 209-220). A review of the literature supports the observation that architectural improvements do occur within the skeleton after daily teriparatide injections, in contrast to the skeletal architecture observed after therapy with antiresorptive agents, which act mainly by inhibition of osteoblastic activity to reduce bone turnover, thus preserving rather than building new bone.

[0199] Continuous rather than intermittent administration of exogenous PTH(1-34) results in bone absorption. Thus treatment with infusions of PTH(1-34) for less than 6 hrs result in bone density increases but infusion for over 8 hr or longer results in bone resorption (Frolik, C.A., et al. (2003) *Bone* 33: 372-379). The prolonged administration of PTH(1-34) causes the expression of RANKL, activating RANK on osteoblast precursors, thus stimulating their differentiation and proliferation (Boyce, B.F. and Xing, L. (2007) *Arthritis Res Ther* 9 Suppl 1: S1). This observation has led to the current treatment paradigm, once daily administration of PTH(1-34) by subcutaneous injection. More recently, studies with infusion of PTH(1-34) for one day and withdrawal for one week showed important gains in bone density (Etoh, M. and

Yamaguchi, A. (2010) *J Bone Miner Metab*, 28: 641-9)). Teriparatide has a Cmax of 10 minutes and a half life of 19 minutes (Frolik, C.A., et al. (2003) *Bone* 33: 372-379). A more efficacious PTH analog might be expected to give more substantive bone density increase.

[0200] During toxicology studies with PTH(1-34), and with PTH(1-84), it was observed that a substantial percentage of the rats developed osteosarcomas, beginning at around 20 months (Tashjian, A.H., Jr. and Goltzman, D. (2008) *J Bone Miner Res* 23: 803-811). No treatment-related sarcomas are reported in human trials with recombinant PTH(1-34), teriparatide (Forteo®). However, treatment with current therapy is limited to <2yrs of continuous daily subcutaneous injection.

PTH and PTHrP

[0201] In some embodiments, the methods and compositions described herein comprise the use of PTH and/or PTHrP peptides and/or proteins and/or analogs thereof. All of the biological activity of intact human PTH (hPTH1-84) resides in the N-terminal sequence; most clinical studies have used the 34-amino acid peptide hPTH(1-34), known as teriparatide. The first two amino acids are obligatory for biological activity, and it appears that the bone anabolic properties are fully maintained by the foreshortened fragment hPTH(1-31) or its cyclized lactam (Whitfield, J.F. and Morley, P. (1995) *Trends Pharmacol Sci* 16: 382-386). More recently studies of sequences as short as hPTH(1-11) have shown activity and can be further modified to decrease their EC₅₀ to the low nM range (Shimizu, M., et al. (2000) *J Biol Chem* 275: 21836-21843; Shimizu, N., et al. (2004) *J Bone Miner Res* 19: 2078-2086).

[0202] Studies of the interaction of the PTH and PTHrP with the PTH1R have indicated that the ligands each have two binding regions, one in the N-terminal 1-14 region and a second in the C-terminal 15-34 region. The 1-14 portion has a more locally ordered structure and interacts with the 7-transmembrane region of the receptor while the 15-34 region is alpha helical and interacts with the extracellular, N-terminal extension of the receptor. While the N-terminal region of these peptides appears to have the primary role of receptor activation through this juxtamembrane region interaction, the C-terminal helical region has important binding interactions (Figure 2.) that give rise to higher potency of the ligand (Gardella, T.J., et al. (1994) *Endocrinology* 135: 1186-1194; Luck, M.D., et al. (1999) *Mol Endocrinol* 13: 670-680) through interaction with the extracellular region of the receptor (Dean, T., et al. (2006) *J Biol Chem* 281: 32485-32495; Potetinova, Z., et al. (2006) *Biochemistry* 45: 11113-11121). This leads to a two domain model of binding that also has been extended to other members of the family of class B GPCRs (Holtmann, M.H., et al. (1995) *J Biol Chem* 270: 14394-14398; Bergwitz, C., et al. (1996) *J Biol Chem* 271: 26469-26472; Runge, S., et al. (2003) *J Biol Chem* 278: 28005-28010).

[0203] Accordingly, provided herein are covalently modified peptides or peptide analogs that have a sequence that allows for binding to PTH1R. Covalent modification of ligands in PTH/PTHrP class relies on the development of N-terminal ligand sequences that interact with

the juxtamembrane portion of the PTHR1 coupled to a C-terminal region that has been truncated and simplified through the use of a surfactant moiety (e.g., a 1-alkyl-glucuronic acid moiety such as the glucose-derived 1-alkylglucuronic acid). Since much of the C-terminal region interaction involves general hydrophobic interaction with the hydrophobic channel (Pioszak, A.A. and Xu, H.E. (2008) *Proc Natl Acad Sci U S A* 105: 5034-5039 ; Pioszak, A.A., et al. (2009) *J Biol Chem* 284: 28382-28391) in the extracellular region of the receptor, the covalent modifications described herein allow for increased binding interaction through low specificity interaction.

[0204] An example of the improved cellular stimulation by a surfactant modified peptide described herein is illustrated in **Figure 4**. Thus substantially greater cAMP output (125%; super-agonistic stimulation) is shown by cells stimulated with doses of EU-232 (**Figure 5**.) than by the internal standard, human PTHrP. Similarly, a coded sample of human PTHrP (**Figure 6**.; EU-285) achieves only 100% of the maximal stimulation of the internal assay standard, human PTHrP. EU-232 is modified with a 1-dodecyl β -D-glucouronic acid moiety in the C-terminal region. Importantly, shorter peptide chains or peptide chains of this size unmodified with a 1-dodecyl β -D-glucouronic acid moiety can be expected to show decreased efficacy.

[0205] Similarly, the in vivo response to this surfactant modified analog EU-232 shows high potency and prolonged duration of action (**Figure 7**.). Blood phosphate levels were tested at various time points after dosing rats with saline (G1), 80 micrograms per kg of PTH (G2), 80 micrograms per kg of EU-232 (G3) or 320 micrograms per kg of EU-232 (G4). EU-232 demonstrates prolonged duration of action in that the maximal statistically significant effect is seen at the last time point in the assay (5 hrs post dosing) and most animals show an increasing biological effect at that timepoint. In contrast PTH has a half-life of 30 minutes in the body and in a similar in vivo assay has a duration of biological effect of about 1 hr. Thus the duration of effect of the surfactant-modified compound is remarkably longer than that for the native hormone.

[0206] In some embodiments, the surfactant modified analogs described herein displays prolonged duration of action by comparison to the unmodified peptide. In some embodiments, the duration of action is longer compared to the unmodified peptide by 10 mins, 30 mins, 45 mins, 1 h, 1.5 h, 2 h, 2.5 h, 3 h, 4 h, 5 h, 5.5 h, 6 h, 6.5 h, 7 h, 7.5 h, 8 h, 8.5 h, 9 h, 9.5 h, 10 h, 11 h, 12 h, 13 h, 14 h, 15 h, 16 h, 20 h, 24 h, 36 h, or 48 h. In some embodiments, the duration of action is 1.5 times, 2 times, 2.5 times, 3 times, 3.5 times, 4 times, 4.5 times, 5 times, 6 times, 7 times, 8 times, 9 times, or 10 times longer compared to the the unmodified peptide.

[0207] As shown in **Figure 7**, Blood calcium levels were tested at various time points after dosing rats with saline (G1), 80 micrograms per kg of PTH (G2), 80 micrograms per kg of EU-232 (G3) or 320 micrograms per kg of EU-232 (G4). No groups were statistically significantly different from control (G1). Importantly, the maximally effective dose and time point for EU-232 (G4; at 5hrs) shows no elevation and thus no indications of a propensity for hypercalcemia at a maximally effective dose. Hypercalcemia is an important side effect see following

administration of PTH 1-34 and of potent analogs of PTH, so this modification has demonstrated an ability to both increase desirable potency on target as well as diminished propensity for undesired side effects.

[0208] The improvements, described above and in the figures, associated with surfactant modification to yield the peptides described herein have significant implications for their use in medicine. Such molecules are suitable for use by once daily, or less frequent, administration to give enhanced biological results compared to treatment with short-acting native hormones such as PTH ($T_{1/2}$ 30min by s.c. injection) or PTHrP. Surfactant-modified peptides such as EU-232 may be expected to show greater biological effect when administered by intranasal insufflation due to the well-known effects of surfactants on nasal bioavailability.

[0209] Accordingly, in some embodiments, surfactant-modified peptide products described herein reduce the occurrence of proteolytic degradation. In some embodiments, covalent modification of PTH and/or PTHrP and/or analogs thereof, allows for decreased cost of production of therapeutics, and provide favorable pharmaceutical properties due to the presence of the covalently attached surfactant moiety. In some embodiments, surfactant modified PTH and/or PTHrP described herein prolong the PK and duration of action (PD) behavior of the resulting ligands compared to other known peptide ligands (such as those of Dean, et al. (Dean, T., et al. (2006) J Biol Chem 281: 32485-32495)) that lack such covalent modifications. Also contemplated within the scope of embodiments presented herein is long term and safe administration of surfactant modified PTH and/or PTHrP and/or analogs thereof.

[0210] In some instances, the N-terminal binding region ends at about residue 14 and the helical region encompasses residue 16 onward. Thus a ligand optimized in the 1-14 region with a 1-alkylglucuronic acid modification in the 15 onward region will have high specific binding (N-terminus) with high potency (1-alkyl modification). Use of an α -helical stabilizing substitution (Kaul, R. and Balaram, P. (1999) Bioorg Med Chem 7: 105-117) in the C-terminal region leads to higher helical content and higher potency. Commonly used α -helical stabilizers are Ala and the class of 1,1-dialkyl amino acids such as Aib, Ac4c, Ac5c and the like (see definitions below). Minimization of the PTH structure led to shortened analogs (Shimizu, M., et al. (2000) J Biol Chem 275: 21836-21843) wherein constrained α -helical stabilizers led to important potency increases. For example, substitution of Aib into position 1 and 3 of PTH1-14 and PTH1-11 analogs led to increased potency (Shimizu, N., et al. (2001) J Biol Chem 276: 49003-49012). Incorporation of more hindered α -helical stabilizers in position 1 lead to further potency increases (Shimizu, N., et al. (2004) J Bone Miner Res 19: 2078-2086). Substitution of Aib into various positions of PTH1-34 analogs also led to improvements in potency, particularly substitutions at positions 12 and 13 (Peggion, E., et al. (2003) Biopolymers 68: 437-457). However it was shown that Aib in positions 1 and 3 of the simple PTH1-11 sequence was not acceptable (Barazza, A., et al. (2005) J Pept Res 65: 23-35). Thus, in some embodiments, the α -helical content of PTH and/or PTHrP is a determinant of peptide product stability (Marx, U.C., et al. (1995) J Biol Chem 270: 15194-15202; Schievano, E., et al. (2000) Biopolymers 54: 429-447).

[0211] In some embodiments, the use of a long side chain, such as in the Nε-(1'-dodecyl beta-D-glucuronyl)-lysine in covalent peptide modifications described herein, destabilizes an α-helix. Accordingly, also contemplated within the scope of embodiments presented herein are modifications that comprise α-helical stabilizers. Thus in some embodiments, surfactant modified peptide products described herein comprise a helix stabilizer (e.g., in the position just N-Terminal and/or just C-terminal of the surfactant substitution). In some embodiments, an α-helix stabilizer is located at position 12 in the PTH and/or PTHrP chain. By way of example only, the table below describes EU-212 to EU-282 certain surfactant modified peptide products (EU-212 to EU-282) that comprise an α-helix stabilizer.

[0212] In one aspect, the peptides that are covalently modified and are suitable for methods described herein are truncated analogs of PTHrP and/or the related hormone PTH, including and not limited to:

hPTH(1-34): Ser₁-Val₂-Ser₃-Glu₄-Ile₅-Gln₆-Leu₇-Met₈-His₉-Asn₁₀-Leu₁₁-Gly₁₂-Lys₁₃-His₁₄-Leu₁₅-Asn₁₆-Ser₁₇-Met₁₈-Glu₁₉-Arg₂₀-Val₂₁-Glu₂₂-Trp₂₃-Leu₂₄-Arg₂₅-Lys₂₆-Lys₂₇-Leu₂₈-Gln₂₉-Asp₃₀-Val₃₁-His₃₂-Asn₃₃-Phe₃₄-OH; (**SEQ. ID. NO. 290**) or

hPTHrP(1-34): Ala₁-Val₂-Ser₃-Glu₄-His₅-Gln₆-Leu₇-Leu₈-His₉-Asp₁₀-Lys₁₁-Gly₁₂-Lys₁₃-Ser₁₄-Leu₁₅-Gln₁₆-Asp₁₇-Leu₁₈-Arg₁₉-Arg₂₀-Arg₂₁-Phe₂₂-Phe₂₃-Leu₂₄-His₂₅-His₂₆-Leu₂₇-Ile₂₈-Ala₂₉-Glu₃₀-Ile₃₁-His₃₂-Thr₃₃-Ala₃₄-OH (**SEQ. ID. NO. 291**)

[0213] In some embodiments, a peptide product described herein has the structure of Formula 2-V:

aa₁-aa₂-aa₃-aa₄-aa₅-aa₆-aa₇-aa₈-aa₉-aa₁₀-aa₁₁-aa₁₂-aa₁₃-aa₁₄-aa₁₅-aa₁₆-aa₁₇-aa₁₈-
aa₁₉-aa₂₀-aa₂₁-aa₂₂-aa₂₃-aa₂₄-aa₂₅-aa₂₆-aa₂₇-aa₂₈-aa₂₉-aa₃₀-aa₃₁-aa₃₂-aa₃₃-aa₃₄-aa₃₅-
aa₃₆-Z FORMULA 2-V (**SEQ. ID. NO. 172**)

wherein:

Z is OH, or -NH-R³, wherein R³ is H or C₁-C₁₂ alkyl; or a PEG chain of less than 10Da.

aa₁ is Ala, Ser, Val, Pro, Aib, Ac5c, or Deg;

aa₂ is Val;

aa₃ is Ser, Ala, Aib, Ac4c, or Deg;

aa₄ is Glu;

aa₅ is His, or Ile;

aa₆ is Gln, or Cit;

aa₇ is Leu, or Phe;

aa₈ is Leu, Met, or Nle;

aa₉ is His;

aa₁₀ is Asp, Asn, Gln, Glu, Cit, Ala, or Aib;

aa₁₁ is Lys, Leu, Ile, Arg, or hArg;

aa₁₂ is Gly, Ala, Glu, Lys, Aib, or Ac5c;

aa₁₃ is Lys, or Arg;

aa₁₄ is Ser, His, Trp, Phe, Leu, Arg, Lys, Glu, Nal(2), or cyclized to position aa₁₈;

aa₁₅ is Ile, Leu, Aib;

aa₁₆ is Gln, Asn, Glu, Lys, Ser, Cit, Aib, Ac5c, U(X);

aa₁₇ is Asp, Ser, Aib, Ac4c, Ac5c, U(X), Z;

aa₁₈ is absent, Leu, Gln, Cit, Aib, Ac5c, Lys, Glu, or U(X), or cyclized to position aa₁₄;

aa₁₉ is absent, Arg, Glu, Aib, Ac4c, Ac5c, or U(X);

aa₂₀ is absent, Arg, Glu, Lys, Aib, Ac4c, Ac5c, or U(X);

aa₂₁ is absent, Arg, Val, Aib, Ac5c, Deg, or U(X);

aa₂₂ is absent, Phe, Glu, Aib, Ac5c, Lys, U(X), or cyclized to position aa₁₈ or aa₂₆;

aa₂₃ is absent, or Leu, Phe, Trp, or U(X);

aa₂₄ is absent, His, Arg, Leu, Aib, Ac5c or U(X);

aa₂₅ is absent, His, Lys, Arg, or U(X);

aa₂₆ is absent, His, Lys, Arg, Aib, Ac5c or cyclized to position aa₂₂;

aa₂₇ is absent, Leu, or Lys;

aa₂₈ is absent, Ile, or Leu;

aa₂₉ is absent, Ala, Gln, Cit, or Aib;

aa₃₀ is absent, Glu, Asp, or Aib;

aa₃₁ is absent, Ile, Val, Aib, Ac5C, or U(X);

aa₃₂ is absent, His, Aib, Ac5C, or U(X);

aa₃₃ is absent, Thr, Asn, Aib, Ac5C, or U(X);

aa₃₄ is absent, Ala, Phe, Aib, Ac5C, or U(X);

aa₃₅ is absent, Aib, Ac5C, or U(X);

aa₃₆ is absent, Aib, Ac5C, or U(X);

U is a linking amino acid; and

X is a surfactant-linked to the side chain of U;

wherein any two of aa₁-aa₃₆ are optionally cyclized through their side chains to form a lactam linkage; and

provided that one, or least one of aa₁ - aa₃₆ is U.

[0214] In some embodiments, a peptide product described herein comprises aa₁-aa₂₀ of Formula V as described above (**SEQ. ID. NO. 292**). In some embodiments, a peptide product described herein comprises aa₁-aa₁₉ of Formula V as described above (**SEQ. ID. NO. 293**).

[0215] In specific embodiments, the linking amino acid U, is a diamino acid like Lys or Orn, X is a modified surfactant from the 1-alkyl glycoside class linked to U, and Z is OH, or -NH-R³, wherein R³ is H or C₁-C₁₂; or a PEG chain of less than 10Da.

[0216] In some embodiments, a peptide product described herein has the structure of Formula 2-VI:

aa₁- Val₂- aa₃- Glu₄- aa₅ aa₆- aa₇- aa₈- His₉- aa₁₀- aa₁₁- aa₁₂- aa₁₃- aa₁₄- aa₁₅- aa₁₆- aa₁₇- aa₁₈-

aa₁₉- aa₂₀- aa₂₁- aa₂₂- aa₂₃- aa₂₄- aa₂₅- aa₂₆-Z Formula 2-VI (**SEQ. ID. NO. 173**)

wherein:

Z is OH, or -NH-R³, wherein R³ is H, C₁-C₁₂ alkyl or a PEG chain of less than 10 Da;

aa₁ is Aib, Ac5c, Deg;

aa₃ is Aib, Ac4c, Deg;

aa₈ is His, Ile;

aa₆ is Gln, Cit;

aa₇ is Leu, Phe;

aa₈ is Leu, Nle;

aa₁₀ is Asp, Asn, Gln, Glu, Cit, Ala, Aib;

aa₁₁ is Arg, hArg;

aa₁₂ is Gly, Ala, Glu, Lys, Aib, Ac5c;

aa₁₃ is Lys, Arg;

aa₁₄ is Ser, His, Trp, Phe, Leu, Arg, Lys, Nal(2);

aa₁₅ is Ile, Leu, Aib;

aa₁₆ is Gln, Asn, Glu, Lys, Ser, Cit, Aib, U(X);

aa₁₇ is Asp, Ser, Aib, Ac4c, Ac5c, U(X);

aa₁₈ is absent or Leu, Gln, Cit, Aib, Ac5c, Lys, Glu, U(X);

aa₁₉ is absent or Arg, Glu, Aib, Ac4c, Ac5c, U(X);

aa₂₀ is absent or Arg, Glu, Lys, Aib, Ac4c, Ac5c, U(X);

aa₂₁ is absent or Arg, Val, Aib, Ac5C, Deg U(X);

aa₂₂ is absent or Phe, Glu, Lys or U(X);

aa₂₃ is absent or Leu, Phe, Trp or U(X);

aa₂₄ is absent or Leu, His, Arg, or U(X);

aa₂₅ is absent or His, Lys, or U(X) and

aa₂₆ is absent or Aib, Ac5c;

U is a linking amino acid;

X is a modified surfactant from the 1-alkyl glycoside class linked to U, wherein the 1-alkyl group is substituted or unsubstituted C₁-C₂₀ alkyl or substituted or unsubstituted C₁-C₂₀ aralkyl;

provided that one, or least one of aa₁ - aa₂₆ is U.

[0217] In some embodiments, a peptide product described herein has the structure of Formula 2-VII

aa₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- aa₁₂- Arg₁₃- aa₁₄- Ile₁₅-

aa₁₆- aa₁₇- aa₁₈- aa₁₉- aa₂₀- aa₂₁- aa₂₂- aa₂₃- aa₂₄- aa₂₅- aa₂₆-Z Formula 2-VII

(SEQ. ID. NO. 174)

wherein:

Z is OH or -NH₂;

aa₁ is Aib, Ac5c;

aa₁₂ is Ala, Glu, Lys, Aib, Ac5c;

aa₁₄ is Trp, Phe, Nal(2);

aa₁₆ is Gln, Asn, Glu, Lys, Cit, U(X);

aa₁₇ is Asp, Ser, Aib, Ac4c, Ac5c, U(X);

aa₁₈ is absent or Leu, Gln, Aib, U(X);

aa₁₉ is absent or Arg, Glu, Aib, Ac4c, Ac5c;

aa₂₀ is absent or Arg, Glu, Lys, Aib, Ac4c, Ac5c;

aa₂₁ is absent or Arg, Val, Aib, Ac5C, Deg;

aa₂₂ is absent or Phe, Glu, Lys or U(X);

aa₂₃ is absent or Leu, Phe, Trp or U(X);

aa₂₄ is absent or His, Arg, or U;

aa₂₅ is absent or His, Lys, or U and

aa₂₆ is absent or Aib, Ac5c;

U is a linking amino acid; and

X is a modified surfactant from the 1-alkyl glycoside class linked to U, wherein the 1-alkyl group is substituted or unsubstituted C₁-C₂₀ alkyl or substituted or unsubstituted C₁-C₂₀ aralkyl;

provided that one, or least one of aa₁ - aa₂₆ is U.

[0218] In some embodiments, the peptide product has the structure of Formula 2-III:

aa₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- aa₁₂- Arg₁₃- aa₁₄- Ile₁₅-

aa₁₆- aa₁₇- aa₁₈- aa₁₉- aa₂₀- aa₂₁- aa₂₂- aa₂₃- aa₂₄- aa₂₅- aa₂₆-Z Formula 2-III

(SEQ. ID. NO. 171)

wherein:

Z is OH or -NH₂;

aa₁ is Aib, or Ac5c;

aa₁₂ is Ala, Aib, Glu, Lys, or Ac5c;

aa₁₄ is Trp, Phe, Lys, Glu or Nal(2);

aa₁₆ is Gln, Asn, Glu, Lys, Cit, or U(X);

aa₁₇ is Asp, Ser, Aib, Ac4c, Ac5c, or U(X);

aa₁₈ is absent or Leu, Gln, Aib, Lys, Glu or U(X);

aa₁₉ is absent or Arg, Glu, Aib, Ac4c, or Ac5c;

aa₂₀ is absent or Arg, Glu, Lys, Aib, Ac4c, Ac5c;

aa₂₁ is absent or Arg, Val, Aib, Ac5C, or Deg;

aa₂₂ is absent or Phe, Glu, Lys or U(X);

aa₂₃ is absent or Leu, Phe, Trp or U(X);

aa₂₄ is absent or His, Arg, or U(X);


aa₂₅ is absent or His, Lys, or U(X); and

aa₂₆ is absent or Aib, Ac5c;

wherein any two of aa₁-aa₂₆ are optionally cyclized through their side chains to form a lactam linkage; and

provided that one, or at least one of aa₁₆, aa₁₇, aa₁₈, aa₂₂, aa₂₃, aa₂₄ or aa₂₅ is the linker amino acid U covalently attached to X.

[0219] In a specific embodiment of Formula 2-III above, X has the structure:



Formula 2-I

wherein:

R^{1a} is a substituted or unsubstituted C₁-C₃₀ alkyl group;

R^{1b}, R^{1c}, and R^{1d} are H;

W¹ is -(C=O)-NH-;

W² is -O-; and

R² is a bond.

[0220] In some of the embodiments described above, R^{1a} is a C₁-C₂₀ alkyl group, a C₁-C₁₈ alkyl group, a C₁-C₁₆ alkyl group, or C₁-C₁₂ alkyl group. In some embodiments of Formula 2-III, U is any linker amino acid described herein.

[0221] Contemplated within the scope of embodiments presented herein are peptide products of Formula 2-I-A, Formula 2-III, Formula 2-V, Formula 2-VI or Formula 2-VII, wherein the peptide product comprises one, or, more than one surfactant groups (e.g., group X having the structure of Formula I). In one embodiment, a peptide product of Formula 2-I-A, Formula 2-III, Formula 2-V, Formula 2-VI or Formula 2-VII comprises one surfactant group. In another embodiment, a peptide product of Formula 2-I-A, Formula 2-III, Formula 2-V, Formula 2-VI or Formula 2-VII comprises two surfactant groups. In yet another embodiment, a peptide product of Formula 2-I-A, Formula 2-III, Formula 2-V, Formula 2-VI or Formula 2-VII comprises three surfactant groups.


[0222] Table 2 in Figure 2 illustrates certain examples of peptides that are suitable for covalent linkage with surfactants as described herein.

[0223] Recognized herein is the importance of certain portions of **SEQ. ID. NO. 170** for the treatment of conditions associated with bone loss and/or hyperparathyroidism including, and not limited to, osteoporosis, osteopenia, post-menopausal osteoporosis, Paget's disease, glucocorticoid-induced osteoporosis, inflammatory bone loss, fixation of implants, osteonecrosis of the jaw, stem cell proliferation, old age osteoporosis, humoral hypercalcemia, or the like.

[0224] Accordingly, provided herein is a method of treating conditions associated with bone loss (e.g., osteoporosis) and/or hyperparathyroidism in an individual in need thereof comprising administration of a therapeutically effective amount of a PTH analog comprising amino acid residues aa₁-aa₁₇ of **SEQ. ID. NO. 170** to the individual in need thereof.

[0225] In a further embodiment, provided herein is a method of treating conditions associated with bone loss (e.g., osteoporosis) and/or hyperparathyroidism in an individual in need thereof comprising administration of a therapeutically effective amount of a PTH analog comprising amino acid residues aa₁-aa₁₈ of **SEQ. ID. NO. 170** to the individual in need thereof.

[0226] In another embodiment, provided herein is a method of treating conditions associated with bone loss (e.g., osteoporosis) and/or hyperparathyroidism in an individual in need thereof comprising administration of a therapeutically effective amount of a PTH analog comprising amino acid residues aa₁-aa₁₉ of **SEQ. ID. NO. 170** to the individual in need thereof.



Formula 2-I

wherein:

R^{1a} is independently, at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group, or a steroid nucleus containing moiety;

R^{1b} , R^{1c} , and R^{1d} are each, independently at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group;

W^1 is independently, at each occurrence, $-CH_2-$, $-CH_2-O-$, $-(C=O)$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$, or $-CH_2-S-$;


W^2 is $-O-$, $-CH_2-$ or $-S-$;

R^2 is independently, at each occurrence, a bond to U, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group, $-NH_2$, $-SH$, C_2 - C_4 -alkene, C_2 - C_4 -alkyne, $-NH(C=O)-CH_2-Br$, $-(CH_2)_m$ -maleimide, or $-N_3$;

n is 1, 2 or 3; and

m is 1-10.

[0236] In a specific embodiment, the said PTH analog is modified with a surfactant, X having the structure:



Formula 2-I

wherein:

R^{1a} is a substituted or unsubstituted C_1 - C_{30} alkyl group;

R^{1b} , R^{1c} , and R^{1d} are H;

W^1 is $-(C=O)-NH-$;

W^2 is $-O-$; and

R^2 is a bond.

[0237] In some of the embodiments described above, R^{1a} is a C_1 - C_{20} alkyl group, a C_8 - C_{20} alkyl group, C_{12} - C_{18} alkyl group or C_{14} - C_{18} alkyl group.

[0238] Modifications at the amino or carboxyl terminus may optionally be introduced into the peptides (e.g., PTH or PTHrP) (Nestor, J.J., Jr. (2009) *Current Medicinal Chemistry* 16: 4399 - 4418). For example, the peptides can be truncated or acylated on the N-terminus to yield peptides analogs exhibiting low efficacy, partial agonist and antagonist activity, as has been seen for some peptides (Gourlet, P., et al. (1998) *Eur J Pharmacol* 354: 105-111, Gozes, I. and Furman, S. (2003) *Curr Pharm Des* 9: 483-494). For example, deletion of the first 6 residues of bPTH yields antagonistic analogs (Mahaffey, J.E., et al. (1979) *J Biol Chem* 254: 6496-6498; Goldman, M.E., et al. (1988) *Endocrinology* 123: 2597-2599) and a similar operation on peptides described herein generates potent antagonistic analogs. Other modifications to the N-terminus of peptides, such as deletions or incorporation of D-amino acids such as D-Phe also can give potent and long acting agonists or antagonists when substituted with the modifications described herein such as long chain alkyl glycosides. Such agonists and antagonists also have commercial utility and are within the scope of contemplated embodiments described herein.

[0239] Also contemplated within the scope of embodiments presented herein is N-terminal truncation of PTH (e.g. 7-34 residue analogs) or PTHrP thereby providing inverse agonists (Gardella, T.J., et al. (1996) *Endocrinology* 137: 3936-3941) or antagonists. In some embodiments, inverse agonists and/or antagonists of PTH and/or PTHrP are useful for treatment of "humoral hypercalcemia" associated with a wide range of tumors.

[0240] Also contemplated within the scope of embodiments described herein are surfactants covalently attached to peptide analogs, wherein the native peptide is modified by acetylation, acylation, PEGylation, ADP-ribosylation, amidation, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-link formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, (Nestor, J.J., Jr. (2007) *Comprehensive Medicinal Chemistry II*

2: 573-601, Nestor, J.J., Jr. (2009) *Current Medicinal Chemistry* 16: 4399 - 4418, Creighton, T.E. (1993), Wold, F. (1983) *Posttranslational Covalent Modification of Proteins* 1-12, Seifter, S. and England, S. (1990) *Methods Enzymol* 182: 626-646, Rattan, S.I., et al. (1992) *Ann N Y Acad Sci* 663: 48-62).

[0241] Also contemplated within the scope of embodiments described herein are peptides that are branched or cyclic, with or without branching. Cyclic, branched and branched circular peptides result from post-translational natural processes and are also made by suitable synthetic methods. In some embodiments, any peptide product described herein comprises a peptide analog described above that is then covalently attached to an alkyl-glycoside surfactant moiety.

[0242] Also contemplated within the scope of embodiments presented herein are peptide chains that are substituted in a suitable position by the modification of the analogs claimed herein, e.g., by acylation on a linker amino acid, at for example the ϵ -position of Lys, with fatty acids such as octanoic, decanoic, dodecanoic, tetradecanoic, hexadecanoic, octadecanoic, 3-phenylpropanoic acids and the like, or with saturated or unsaturated alkyl chains (Zhang, L. and Bulaj, G. (2012) *Curr Med Chem* 19: 1602-1618). Non-limiting, illustrative examples of such analogs are:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-dodecanoyl)₁₈-Aib₁₉-NH₂, (**SEQ. ID. NO. 294**)

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-dodecanoyl)₁₈-NH₂, (**SEQ. ID. NO. 295**)

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-palmitoyl)₁₈-Aib₁₉-NH₂, (**SEQ. ID. NO. 296**)

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Phe₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-dodecanoyl)₁₈-Aib₁₉-NH₂, (**SEQ. ID. NO. 297**)

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-tetradecanoyl)₁₈-Aib₁₉-NH₂, (**SEQ. ID. NO. 298**)

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-tetradecanoyl)₁₈-Aib₁₉-Aib₂₀-NH₂, (**SEQ. ID. NO. 299**)

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Lys(N-epsilon-(Nalpha-dodecanoyl-L-glutamyl))₁₈-Aib₁₉-NH₂, (**SEQ. ID. NO. 300**) and the like.

[0243] In other embodiments, a peptide chain is substituted in a suitable position by reaction

on a linker amino acid, for example the sulfhydryl of Cys, with a spacer and a hydrophobic moiety such as a steroid nucleus, for example a cholesterol moiety. In some of such embodiments, the modified peptide further comprises one or more PEG chains. Non-limiting examples of such molecules are:

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Cys(S-(3-(PEG4-aminoethylacetamide-Cholesterol)))₁₈-Aib₁₉-NH₂, (**SEQ. ID. NO. 301**)

Ac5c₁- Val₂- Aib₃- Glu₄- Ile₅-Gln₆- Leu₇- Nle₈- His₉- Gln₁₀- hArg₁₁- Ala₁₂- Arg₁₃- Trp₁₄-Ile₁₅- Gln₁₆- Aib₁₇-Cys(S-(3-(PEG4-aminoethylacetamide-Cholesterol)))₁₈-NH₂, (**SEQ. ID. NO. 302**) and the like.

[0244] In some embodiments, the modified peptides described herein displays improved protection from proteolysis when compared to the unmodified peptide. In some embodiments, the proteolysis of the modified peptides described herein is reduced by 10-90%, 10-20%, 10-50%, 20-80%, 20-40%, 20-30%, 30-60%, 30-50%, 30-40%, 10-30%, 15-30%, 15-45%, 50-75%, 60-75%, 70-90%, or 80-100% when compared with the unmodified peptide. In some embodiments, the proteolysis of the modified peptides described herein is reduced by 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 100%, greater than 5%, greater than 10%, greater than 15%, greater than 20%, greater than 25%, greater than 30%, greater than 35%, greater than 40%, greater than 45%, greater than 50%, greater than 55%, greater than 60%, greater than 65%, greater than 70%, greater than 75%, greater than 80%, greater than 85%, greater than 90%, greater than 95%, greater than 98%, at least 2%, at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% when compared with the unmodified peptide.

[0245] In some embodiments, the modified peptides described herein display arrestin-biased agonism. In other embodiments, the modified peptides described herein display arrestin-biased antagonism.

[0246] In some embodiments, any of the modified peptides described herein displays improved receptor binding potency and/or cellular activation potency compared to the unmodified peptide. In some embodiments, the receptor binding potency and/or cellular activation potency of the modified peptides described herein is increased by 10-90%, 10-20%, 10-50%, 20-80%, 20-40%, 20-30%, 30-60%, 30-50%, 30-40%, 10-30%, 15-30%, 15-45%, 50-75%, 60-75%, 70-90%, or 80-100% when compared with the unmodified peptide. In some embodiments, the receptor binding potency and/or cellular activation potency of the modified peptides described herein is increased by 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 100%, 200%, 300%, 400%,


500%, 600%, 700%, 800\$, 900%, 1000%, greater than 5%, greater than 10%, greater than 15%, greater than 20%, greater than 25%, greater than 30%, greater than 35%, greater than 40%, greater than 45%, greater than 50%, greater than 55%, greater than 60%, greater than 65%, greater than 70%, greater than 75%, greater than 80%, greater than 85%, greater than 90%, greater than 95%, greater than 98%, at least 2%, at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% when compared with the unmodified peptide.

[0247] In some embodiments, any of the modified peptides described herein displays improved cellular activation potency compared to the unmodified peptide. In some embodiments, the cellular activation potency of the modified peptides described herein is increased by 10-90%, 10-20%, 10-50%, 20-80%, 20-40%, 20-30%, 30-60%, 30-50%, 30-40%, 10-30%, 15-30%, 15-45%, 50-75%, 60-75%, 70-90%, or 80-100% when compared with the unmodified peptide. In some embodiments, the cellular activation potency of the modified peptides described herein is increased by 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 100%, 200%, 300%, 400%, 500%, 600%, 700%, 800\$, 900%, 1000%, greater than 5%, greater than 10%, greater than 15%, greater than 20%, greater than 25%, greater than 30%, greater than 35%, greater than 40%, greater than 45%, greater than 50%, greater than 55%, greater than 60%, greater than 65%, greater than 70%, greater than 75%, greater than 80%, greater than 85%, greater than 90%, greater than 95%, greater than 98%, at least 2%, at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% when compared with the unmodified peptide.

GLP peptides and analogs


[0248] Also provided herein, in some embodiments, are reagents and intermediates for synthesis of modified peptides and/or proteins (e.g., modified GLP-1, glucagon, analogs of glucagon or GLP-1, or the like) through the incorporation of surfactants.


[0249] Provided herein, in some embodiments, are peptide products comprising a surfactant X, covalently attached to a peptide, the peptide comprising a linker amino acid U and at least one other amino acid:



Formula 3-I-A

wherein the surfactant X is a group of Formula I:





Formula 3-I

wherein:

R^{1a} is independently, at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group;

R^{1b} , R^{1c} , and R^{1d} are each, independently at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group;

W^1 is independently, at each occurrence, $-CH_2-$, $-CH_2-O-$, $-(C=O)$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$, or $-CH_2-S-$;

W^2 is $-O-$, $-CH_2-$ or $-S-$;

R^2 is independently, at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group, $-NH_2$, $-SH$, C_2 - C_4 -alkene, C_2 - C_4 -alkyne, $-NH(C=O)-CH_2-Br$, $-(CH_2)_m$ -maleimide, or $-N_3$;

n is 1, 2 or 3; and

m is 1-10;

the peptide is selected from Formula 3-II:

aa₁-aa₂-aa₃-aa₄-aa₅-aa₆-aa₇-aa₈-aa₉-aa₁₀- aa₁₁-aa₁₂-aa₁₃-aa₁₄-aa₁₅-aa₁₆-aa₁₇-aa₁₈-

aa₁₉-aa₂₀- aa₂₁-aa₂₂-aa₂₃-aa₂₄-aa₂₅-aa₂₆-aa₂₇-aa₂₈-aa₂₉-aa₃₀-aa₃₁-aa₃₂-aa₃₃-aa₃₄-aa₃₅-

aa₃₆-aa₃₇-Z

Formula 3-II (SEQ. ID. NO. 303)

wherein:

Z is OH, or $-NH-R^3$, wherein R^3 is H or C_1 - C_{12} substituted or unsubstituted alkyl, or a PEG chain of less than 10 Da;

aa₁ is His, N-Ac-His, pGlu-His, or $N-R^3$ -His;

aa₂ is Ser, Ala, Gly, Aib, Ac4c or Ac5c;

aa₃ is Gln, or Cit;

aa₄ is Gly, or D-Ala;

aa₈ is Thr, or Ser;

aa₆ is Phe, Trp, F2Phe, Me2Phe, or Nal2;

aa₇ is Thr, or Ser;

aa₈ is Ser, or Asp;

aa₉ is Asp, or Glu;

aa₁₀ is Tyr, Leu, Met, Nal2, Bip, or Bip2EtMeO;

aa₁₁ is Ser, Asn, or U;

aa₁₂ is Lys, Glu, Ser, Arg, or U;

aa₁₃ is absent or Tyr, Gln, Cit, or U;

aa₁₄ is absent or Leu, Met, Nle, or U;

aa₁₅ is absent or Asp, Glu, or U;

aa₁₆ is absent or Ser, Gly, Glu, Aib, Ac5c, Lys, Arg, or U;

aa₁₇ is absent or Arg, hArg, Gln, Glu, Cit, Aib, Ac4c, Ac5c, or U;

aa₁₈ is absent or Arg, hArg, Ala, Aib, Ac4c, Ac5c, or U;

aa₁₉ is absent or Ala, Val, Aib, Ac4c, Ac5c, or U;

aa₂₀ is absent or Gln, Lys, Arg, Cit, Glu, Aib, Ac4c, Ac5c, or U;

aa₂₁ is absent or Asp, Glu, Leu, Aib, Ac4c, Ac5c, or U;

aa₂₂ is absent or Phe, Trp, Nal2, Aib, Ac4c, Ac5c, or U

aa₂₃ is absent or Val, Ile, Aib, Ac4c, Ac5c, or U;

aa₂₄ is absent or Gln, Ala, Glu, Cit, or U;

aa₂₅ is absent or Trp, Nal2, or U;

aa₂₆ is absent or Leu, or U;

aa₂₇ is absent or Met, Val, Nle, Lys, or U;

aa₂₈ is absent or Asn, Lys, or U;

aa₂₉ is absent or Thr, Gly, Aib, Ac4c, Ac5c, or U;

aa₃₀ is absent or Lys, Aib, Ac4c, Ac5c, or U;

aa₃₁ is absent or Arg, Aib, Ac4c, Ac5c, or U;

aa₃₂ is absent or Asn, Aib, Ac4c, Ac5c, or U;

aa₃₃ is absent or Arg, Aib, Ac4c, Ac5c, or U;

aa₃₄ is absent or Asn, Aib, Ac4c, Ac5c, or U;

aa₃₅ is absent or Asn, Aib, Ac4c, Ac5c, or U;

aa₃₆ is absent or Ile, Aib, Ac4c, Ac5C, or U;

aa₃₆ is absent or Ala, Aib, Ac4c, Ac5C, or U;

aa₃₇ absent or U;


U is a natural or unnatural amino acid comprising a functional group used for covalent attachment to the surfactant X;

wherein any two of aa₁-aa₃₇ are optionally cyclized through their side chains to form a lactam linkage; and

provided that one, or at least one of aa₁₁ - aa₃₇ is the linker amino acid U covalently attached to X.

[0250] In some embodiments, n is 1. In some embodiments, n is 2, and a first glycoside is attached to a second glycoside via bond between W² of the first glycoside and any one of OR^{1b}, OR^{1c} or OR^{1d} of the second glycoside. In some embodiments, n is 3, and a first glycoside is attached to a second glycoside via bond between W² of the first glycoside and any one of OR^{1b}, OR^{1c} or OR^{1d} of the second glycoside, and the second glycoside is attached to a third glycoside via bond between W² of the second glycoside and any one of OR^{1b}, OR^{1c} or OR^{1d} of the third glycoside.

[0251] In one embodiment, compounds of Formula I-A are compounds wherein X has the structure:



Formula 3-I

wherein:

R^{1a} is H, a protecting group, a substituted or unsubstituted C_1 - C_{30} alkyl group, or a steroid nucleus containing moiety;

R^{1b} , R^{1c} , and R^{1d} are each, independently at each occurrence, H, a protecting group, or a substituted or unsubstituted C_1 - C_{30} alkyl group;


W^1 is independently, at each occurrence, $-CH_2-$, $-CH_2-O-$, $-(C=O)$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$, or $-CH_2-S-$;

W^2 is $-O-$, $-S-$;

R^2 is a bond, C_2 - C_4 -alkene, C_2 - C_4 -alkyne, or $-(CH_2)_m$ -maleimide; and

m is 1-10.


[0252] In another embodiment, compounds of Formula I-A are compounds wherein X has the structure:



[0253] Accordingly, in the embodiment described above, R^2 is a bond.


[0254] For instance, in an exemplary embodiment of the structure of X described above, W^1 is $-C(=O)NH-$, R^2 is a bond between W^1 and an amino acid residue U within the peptide (e.g., an amino group in the sidechain of a lysine residue present in the peptide).

[0255] In a further embodiment, compounds of Formula I-A are compounds wherein X has the structure:



[0256] For instance, in an exemplary embodiment of the structure of X described above, W^1 is CH_2- and R^2 is an alkyl-linked maleimide functional group on X and R^2 is attached to a suitable moiety of an amino acid residue U within the peptide (e.g., a thiol group in a cysteine residue of the peptide forms a thioether with the maleimide on X).

[0257] In yet another embodiment, compounds of Formula I-A are compounds wherein X has the structure:



Formula 3-I

wherein:

R^{1a} is H, a protecting group, a substituted or unsubstituted C₁-C₃₀ alkyl group, or a steroid nucleus containing moiety;


R^{1b}, R^{1c}, and R^{1d} are each, independently at each occurrence, H, a protecting group, or a substituted or unsubstituted C₁-C₃₀ alkyl group;

W¹ is -(C=O)-NH-;

W² is -O-;

R² is a bond.

[0258] In an additional embodiment, compounds of Formula I-A are compounds wherein X has the structure:



Formula 3-I

wherein:

R^{1a} is a substituted or unsubstituted C₁-C₃₀ alkyl group;

R^{1b}, R^{1c}, and R^{1d} are H;

W¹ is -(C=O)-NH-;


W² is -O-; and

R^2 is a bond.

[0259] In some embodiments described above and herein, R^{1a} is a substituted or unsubstituted C_1 - C_{30} alkyl group.

[0260] In some embodiments described above and herein, R^{1a} is a substituted or unsubstituted C_6 - C_{20} alkyl group.

[0261] Also contemplated herein are alternate embodiments wherein X in Formula 3-I-A has the structure:



[0262] For instance, in an exemplary embodiment of the structure of X described above, W^1 is -S-, R^2 is a C_1 - C_{30} alkyl group, W^2 is S, R^{1a} is a bond between W^2 and a suitable moiety of an amino acid residue U within the peptide (e.g., a thiol group in a cysteine residue of the peptide forms a thioether with X).

[0263] In another exemplary embodiment of the structure of X described above, W^1 is -O-, R^2 is a C_1 - C_{30} alkyl group, W^2 is O, R^{1a} is a bond between W^2 and a suitable moiety of an amino acid residue U within the peptide (e.g., a hydroxyl group in a serine or threonine residue of the peptide forms an ether with X).

[0264] In some embodiments, U is used for covalent attachment to X and is a dibasic natural or unnatural amino acid, a natural or unnatural amino acid comprising a thiol, an unnatural amino acid comprising a $-N_3$ group, an unnatural amino acid comprising an acetylenic group, or an unnatural amino acid comprising a $-NH-C(=O)-CH_2-Br$ or a $-(CH_2)_m$ -maleimide, wherein m is 1-10.

[0265] In some embodiments of the peptide product, the surfactant is a 1-alkyl glycoside class surfactant. In some embodiments of the peptide product, the surfactant is attached to the peptide via an amide bond.

[0266] In some embodiments of the peptide product, the surfactant X is comprised of 1-eicosyl beta-D-glucuronic acid, 1-octadecyl beta-D-glucuronic acid, 1-hexadecyl beta-D-glucuronic acid, 1-tetradecyl beta D-glucuronic acid, 1-dodecyl beta D-glucuronic acid, 1-decyl beta-D-glucuronic acid, 1-octyl beta-D-glucuronic acid, 1-eicosyl beta-D-diglucuronic acid, 1-octadecyl

beta-D-diglucuronic acid, 1-hexadecyl beta-D-diglucuronic acid, 1-tetradecyl beta-D-diglucuronic acid, 1-dodecyl beta-D-diglucuronic acid, 1-decyl beta-D-diglucuronic acid, 1-octyl beta-D-diglucuronic acid, or functionalized 1-ecosyl beta-D-glucose, 1-octadecyl beta-D-glucose, 1-hexadecyl beta-D-glucose, 1-tetradecyl beta-D-glucose, 1-dodecyl beta-D-glucose, 1-decyl beta-D-glucose, 1-octyl beta-D-glucose, 1-eicosyl beta-D-maltoside, 1-octadecyl beta-D-maltoside, 1-hexadecyl beta-D-maltoside, 1-dodecyl beta-D-maltoside, 1-decyl beta-D-maltoside, 1-octyl beta-D-maltoside, and the like, and the peptide product is prepared by formation of a linkage between the aforementioned groups and a group on the peptide (e.g., a -COOH group in the aforementioned groups and an amino group of the peptide).

[0267] In some embodiments of the peptide product, U is a terminal amino acid of the peptide. In some embodiments of the peptide product, U is a non-terminal amino acid of the peptide. In some embodiments of the peptide product, U is a natural D- or L- amino acid. In some embodiments of the peptide product, U is an unnatural amino acid. In some embodiments of the peptide product, U is selected from Lys, Cys, Orn, or an unnatural amino acid comprising a functional group used for covalent attachment to the surfactant X.

[0268] In some embodiments of the peptide product, the functional group used for covalent attachment of the peptide to the surfactant X is -NH₂, -SH, -OH, -N₃, haloacetyl, a -(CH₂)_m-maleimide (wherein m is 1-10), or an acetylenic group.

[0269] In some embodiments side chain functional groups of two different amino acid residues are linked to form a cyclic lactam. For example, in some embodiments, a Lys side chain forms a cyclic lactam with the side chain of Glu. In some embodiments such lactam structures are reversed and are formed from a Glu and a Lys. Such lactam linkages in some instances are known to stabilize alpha helical structures in peptides (Condon, S.M., et al. (2002) *Bioorg Med Chem* 10: 731-736; Murage, E.N., et al (2008) *Bioorg Med Chem* 16: 10106-12; Murage, E.N., et al. (2010) *J Med Chem* 53: 6412-20). In some embodiments cysteine residues may be linked through disulfide formation in order to accomplish a similar form of conformational restriction and assist in the formation of helical structures (Li, Y., et al. (2011) *Peptides* 32: 1400-1407. In some embodiments side chain functional groups of two different amino acid residues are linked to form a heterocycle generated through a "click reaction" between side chain azide and alkyne functional groups in order to achieve a similar form of conformational restriction and stabilized helical conformations (Le Chevalier Isaad A., et al. (2009) *J Peptide Sci* 15: 451-4).

[0270] In some embodiments, the peptide product comprising a covalently linked alkyl glycoside is a covalently modified glucagon or analog thereof. In some of such embodiments, the peptide product contains a covalently linked 1-O-alkyl β-D-glucuronic acid and the peptide is an analog of glucagon.

[0271] In some embodiments, a peptide product comprising a covalently linked alkyl glycoside is a covalently modified GLP-1, or analog thereof. In some of such embodiments, the peptide product comprises a covalently linked 1-O-alkyl β-D-glucuronic acid and the peptide is an

analog of GLP-1.

[0272] In some embodiments, the peptide product of Formula I-A has the structure of Formula III-A

aa₁-aa₂-aa₃-aa₄-aa₅-aa₆-aa₇-aa₈-aa₉-aa₁₀- aa₁₁-aa₁₂-aa₁₃-aa₁₄-aa₁₅-aa₁₆-aa₁₇-aa₁₈-aa₁₉-aa₂₀-

aa₂₁-aa₂₂-aa₂₃-aa₂₄-aa₂₅-aa₂₆-aa₂₇-aa₂₈-aa₂₉ -Z Formula 3-III-A (SEQ. ID. NO.

304)

wherein:

Z is OH, or -NH-R³, wherein R³ is H, or C₁-C₁₂ substituted or unsubstituted alkyl, or a PEG chain of less than 10 Da;

aa₁ is His, N-Ac-His, pGlu-His, or N-R³-His;

aa₂ is Ser, Ala, Gly, Aib, Ac4c, or Ac5c;

aa₃ is Gln, or Cit;

aa₄ is Gly, or D-Ala;

aa₅ is Thr, or Ser;

aa₆ is Phe, Trp, F2Phe, Me2Phe, or Nal2;

aa₇ is Thr, or Ser;

aa₈ is Ser, or Asp;

aa₉ is Asp, or Glu;

aa₁₀ is Tyr, Leu, Met, Nal2, Bip, or Bip2EtMeO;

aa₁₁ is Ser, Asn, or U;

aa₁₂ is Lys, Glu, Ser, Arg, or U(X);

aa₁₃ is absent or Tyr, Gln, Cit, or U(X);

aa₁₄ is absent or Leu, Met, Nle, or U(X);

aa₁₅ is absent or Asp, Glu, or U(X);

aa₁₆ is absent or Ser, Gly, Glu, Aib, Ac5c, Lys, Arg, or U(X);

aa₁₇ is absent or Arg, hArg, Gln, Glu, Cit, Aib, Ac4c, Ac5c, or U(X);

aa₁₈ is absent or Arg, hArg, Ala, Aib, Ac4c, Ac5c, or U(X);

aa₁₉ is absent or Ala, Val, Aib, Ac4c, Ac5c, or U(X);

aa₂₀ is absent or Gln, Lys, Arg, Cit, Glu, Aib, Ac4c, Ac5c, or U(X);

aa₂₁ is absent or Asp, Glu, Leu, Aib, Ac4c, Ac5c, or U(X);

aa₂₂ is absent or Phe, Trp, Nal2, Aib, Ac4c, Ac5c, or U(X);

aa₂₃ is absent or Val, Ile, Aib, Ac4c, Ac5c, or U(X);

aa₂₄ is absent or Gln, Ala, Glu, Cit, or U(X);

aa₂₅ is absent or Trp, Nal2, or U(X);

aa₂₆ is absent or Leu, or U(X);

aa₂₇ is absent or Met, Val, Nle, Lys, or U(X);

aa₂₈ is absent or Asn, Lys, or U(X);

aa₂₉ is absent or Thr, Gly, Aib, Ac4c, Ac5c, or U(X);

wherein any two of aa₁-aa₂₉ are optionally cyclized through their side chains to form a lactam linkage; and

provided that one, or at least one of aa₁₆, aa₁₇, aa₁₈, aa₁₉, aa₂₀, aa₂₁, aa₂₂, aa₂₃, aa₂₄, aa₂₅, aa₂₆, aa₂₇, aa₂₈ or aa₂₉ is the natural or unnatural amino acid U covalently attached to X.

[0273] In some embodiments, the peptide product of Formula I-A has the structure of Formula 3-III-B:

His₁-aa₂-aa₃-Gly₄-Thr₅-aa₆-Thr₇-Ser₈-Asp₉-aa₁₀-aa₁₁-aa₁₂-aa₁₃-aa₁₄-aa₁₅-aa₁₆-aa₁₇-aa₁₈-aa₁₉-
aa₂₀-aa₂₁-aa₂₂-aa₂₃-Z

Formula 3-III-B (SEQ. ID. NO. 305)

wherein:

Z is OH, or -NH-R³, wherein R³ is H or substituted or unsubstituted C₁-C₁₂ alkyl; or a PEG chain of less than 10Da;

aa₂ is Ser, Ala, Gly, Aib, Ac4c, or Ac5c;

aa₃ is Gln, or Cit;

aa₆ is Phe, Trp, F2Phe, Me2Phe, MePhe, or Nal2;

aa₁₀ is Tyr, Leu, Met, Nal2, Bip, or Bip2EtMeO;

aa₁₁ is Ser, Asn, or U(X);

aa₁₂ is Lys, Glu, Ser, or U(X);

aa₁₃ is absent or Tyr, Gln, Cit, or U(X);

aa₁₄ is absent or Leu, Met, Nle, or U(X);

aa₁₅ is absent or Asp, Glu, or U(X);

aa₁₆ is absent or Ser, Gly, Glu, Aib, Ac4c, Ac5c, Lys, R, or U(X);

aa₁₇ is absent or Arg, hArg, Gln, Glu, Cit, Aib, Ac4c, Ac5c, or U(X);

aa₁₈ is absent or Arg, hArg, Ala, Aib, Ac4c, Ac5c, or U(X);

aa₁₉ is absent or Ala, Val, Aib, Ac4c, Ac5c, or U(X);

aa₂₀ is absent or Gln, Lys, Arg, Cit, Glu, Aib, Ac4c, Ac5c, or U(X);

aa₂₁ is absent or Asp, Glu, Leu, Aib, Ac4c, Ac5c, or U(X);

aa₂₂ is absent or Phe, Aib, Ac4c, Ac5c, or U(X)

aa₂₃ is absent or Val, Ile, Aib, Ac4c, Ac5c, or U(X);

wherein any two of aa₁-aa₂₃ are optionally cyclized through their side chains to form a lactam linkage; and

provided that one, or at least one of aa₁₆, aa₁₇, aa₁₈, aa₁₉, aa₂₀, aa₂₁, aa₂₂, aa₂₃ or aa₂₄ is the natural or unnatural amino acid U covalently attached to X.

[0274] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, U is any linker amino acid described herein.

[0275] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, U is any linker amino acid described herein. In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, aa₁₂ is lysine. In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, aa₁₄ is leucine.

[0276] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, aa₁₂ is lysine. In some embodiments of Formula I-A, III-A, III-B or Formula V, aa₁₄ is leucine.

[0277] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, aa₁₈ is a lysine residue attached to X.

[0278] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, aa₁₇ is a homo Arginine (hArg) residue.

[0279] In some embodiments of Formula I-A, III-A, III-B or Formula V, aa₁₇ is a glycine residue.

[0280] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, aa₂ is an Aib or Ac4c residue.

[0281] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide comprises one or more Aib residues.

[0282] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, peptide comprises one or more Aib residues at the C-terminus.

[0283] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 619**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Se-Lys₁₂-Tyr₁₃- Leu₁₄-Asp₁₅-Aib₁₆-aa₁₇-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₁₈-aa₁₉-NH₂; wherein

aa₂ is Aib or Ac4c;

aa₁₇ is Arg, hArg or Gln;

aa₁₉ is Aib, Ac4c or Ac5c; and

alkyl is a C_s to C₂₀ linear alkyl chain.

[0284] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 620**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃- Leu₁₄-Asp₁₅-Aib₁₆-aa₁₇-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₁₈-aa₁₉-aa₂₀-NH₂; wherein

aa₂ is Aib or Ac4c,

aa₁₇ is Arg, hArg or Gln,

aa₁₉ and aa₂₀ are individually Aib, Ac4c or Ac5c; and

alkyl is a C₈ to C₂₀ linear alkyl chain.

[0285] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 621**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-aa₁₆-aa₁₇-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₁₈-aa₁₉-NH₂; wherein

aa₂ is Aib or Ac4c;

aa₁₆ is Aib or Ac4c;

aa₁₇ is Arg, hArg or Gln;

aa₁₉ is Aib, Ac4c or Ac5c; and

alkyl is a C₈ to C₂₀ linear alkyl chain.

[0286] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, aa₁₆ and aa₂₀ are cyclized to form a lactam linkage.

[0287] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure: (**SEQ. ID. NO. 622**)

His₁-aa₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-aa₁₆-aa₁₇-Ala₁₈-Ala₁₉-aa₂₀-Glu₂₁-Phe₂₂-Ile₂₃-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₂₄-Trp₂₅-Leu₂₆-aa₂₇-Asn₂₈-Thr₂₉-NH₂; wherein

aa₂ is Aib or Ac4c;

aa₁₆ and aa₂₀ are each individually either Lys or Glu and are cyclized through their side chains to form a lactam linkage;

aa₁₇ is Arg, hArg or Gln;

aa₂₇ is Met or Nle; and

alkyl is a C₈-C₂₀ linear alkyl chain.

[0288] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 623**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-cyclic(Glu₁₆-Gln₁₇-Ala₁₈-Ala₁₉-Lys₂₀)-Glu₂₁-Phe₂₂-Ile₂₃-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₂₄-Trp₂₅-Leu₂₆-Met₂₇-Asn₂₈-aa₂₉-NH₂; wherein aa₂ is Aib or Ac4c, aa₂₉ is Thr, Aib, Ac4c, or Ac5c, and the 1'-alkyl group is selected from dodecyl, tetradecyl, hexadecyl, or octadecyl; and the side chains on the amino acids in position 16 and 20 are cyclized to form a side chain lactam.

[0289] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, aa₁₂ and aa₁₆ are cyclized to form a lactam linkage.

[0290] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 624**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-aa₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-aa₁₆-aa₁₇-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₁₈-aa₁₉-aa₂₀-NH₂; wherein

aa₂ is Aib or Ac4c;

aa₁₂ and aa₁₆ are each individually either Lys or Glu and are cyclized through their side chains to form a lactam linkage;

aa₁₇ is Arg, or hArg;

aa₁₉ and aa₂₀ are individually either Aib, Ac4c or Ac5c; and

alkyl is a C₈-C₂₀ linear alkyl chain.

[0291] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 625**):

His₁-Ac4c₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-cyclo(Glu₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Lys₁₆)-aa₁₇-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₁₈-Aib₁₉-Aib₂₀-NH₂; wherein

aa₁₂ and aa₁₆ are cyclized through their side chains to form a lactam linkage;

aa₁₇ is Arg or hArg; and

alkyl is a C₁₂, C₁₄, C₁₆, or C₁₈ linear alkyl chain.

[0292] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 626**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-aa₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-aa₁₆-aa₁₇-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₁₈-aa₁₉-aa₂₀-NH₂;

wherein

aa₁₂ and aa₁₆ are each individually either Lys or Glu

and aa₁₂ and aa₁₆ are cyclized through their side chains to form a lactam linkage;

aa₁₇ is Arg or hArg; aa₁₉ and aa₂₀ are individually either Aib, Ac4c or Ac5c; and the 1'-alkyl group is selected from dodecyl, tetradecyl, hexadecyl, or octadecyl.

[0293] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 627**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-aa₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Ser₁₆-Aib₁₇-Lys(N-omega-1'-dodecyl beta-D-glucuronyl)₁₈-aa₁₉-NH₂; wherein aa₂ is Aib or Ac4c, aa₆ is Me2Phe, MePhe, or Phe; and aa₁₉ is Aib, Ac4c, or Ac5c.

[0294] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 628**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-aa₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Ser₁₆-aa₁₇-Lys(N-omega-1'-dodecyl beta-D-glucuronyl)₁₈-aa₁₉-aa₂₀-NH₂; wherein aa₂ is Aib or Ac4c, aa₆ is Me2Phe, MePhe, or Phe; aa₁₇ is Arg or hArg, and aa₁₉ or aa₂₀ is Aib, Ac4c, or Ac5c.

[0295] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 629**):

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-cyclo(Glu₁₆-Arg₁₇-Ala₁₈-Ala₁₉-Lys₂₀)-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₂₁-Phe₂₂-aa₂₃-NH₂; wherein aa₂₃ is Aib, Ac4c, or Ac5c and the 1'-alkyl group is selected from dodecyl, tetradecyl, hexadecyl, or octadecyl..

[0296] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide product has the structure (**SEQ. ID. NO. 630**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-aa₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-aa₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-aa₁₆-aa₁₇-aa₁₈-Ala₁₉-aa₂₀-Lys(N-omega-1'-alkyl beta-D-glucuronyl)₂₁-Phe₂₂-aa₂₃-NH₂;

wherein

aa₂ is Aib or Ac4c;

aa₆ is Me2Phe, MePhe, or Phe;

aa₁₂ and aa₁₆ are each individually either Lys or Glu;

and aa₁₆ and aa₂₀ are cyclized through their side chains to form a lactam linkage;

aa₁₇ is Arg, hArg or Gln;

aa₁₈ is Aib or Ala;

aa₂₃ is Aib, Ac4c, or Ac5c and the 1'-alkyl group is selected from dodecyl, tetradecyl, hexadecyl, or octadecyl.

[0297] In some embodiments of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, the peptide

product has the structure (**SEQ. ID. NO. 631**):

His₁-aa₂-Gln₃-Gly₄-Thr₅-aa₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-aa₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-aa₁₆-aa₁₇-
Lys(N-omega-1'-alkyl beta-D-glucuronyl)₁₈-aa₁₉-aa₂₀-NH₂;

wherein

aa₂ is Aib or Ac4c;

aa₆ is Phe;

aa₁₂ and aa₁₆ are each individually either Lys or Glu; and aa₁₂ and aa₁₆ are cyclized through their side chains to form a lactam linkage;

aa₁₇ is Arg or hArg;


aa₁₉ is Aib, Ac4c, or Ac5c;

aa₂₀ is Aib, Ac4c, or Ac5c and the and the 1'-alkyl group is selected from dodecyl, tetradecyl, hexadecyl, or octadecyl.

[0298] In some embodiments, for any compound of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, X is comprised of a dodecyl alkyl chain.


[0299] In some embodiments, the peptide product is a biologically active peptide product that binds to the GLP1R and/or to the GLCR.


[0300] In a specific embodiment, the peptide products of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V, described above and herein have the following structure:



wherein R^{1a} is a C₁-C₂₀ alkyl chain as described in Table 1 of Figure 1, R' is a peptide as described in Table 3 of Figure 8 and Table 4 of Figure 9, W² of Formula I-A is -O-, and W¹ of Formula I-A is -(C=O)NH- and is part of an amide linkage to the peptide R'. In some of such embodiments, R^{1a} is a C₆-C₂₀ alkyl chain. In some of such embodiments, R^{1a} is a C₈-C₂₀ alkyl chain. In some of such embodiments, R^{1a} is a C₁₂-C₂₀ alkyl chain. In some of such embodiments, R^{1a} is a C₁₂-C₁₆ alkyl chain.


[0301] In embodiments described above, an amino moiety of an amino acid and/or a peptide R' (e.g., an amino group of an amino acid residue such as a Lysine, or a lysine residue within the peptide R') is used to form a covalent linkage with a compound of structure:






wherein R^{1a} is a C₁-C₂₀ alkyl chain as described above and in Table 3 of Figure 8 and Table 4 of Figure 9.

[0302] In such cases, the amino acid residue having an amino moiety (e.g., a Lysine within the peptide R') which is used to form a covalent linkage to the compound A described above, is a linker amino acid U which is attached to a surfactant X having the structure of Formula A. Accordingly, as one example, Lys(C12) of Table 3 of Figure 8 and Table 4 of Figure 9 has the following structure:



[0303] Also contemplated within the scope of the embodiments presented herein are peptide products of Formula 3-I-A derived from maltouronic acid-based surfactants through binding at either or both carboxylic acid functions. Thus, as one example, peptides in Table 3 of Figure 8 and Table 4 of Figure 9 comprise a lysine linker amino acid bonded to a maltouronic acid based surfactant X and having a structure:



[0304] It will be understood that in one embodiment, compounds of Formula 3-I-A are prepared by attaching a lysine to a group X, followed by attachment of additional amino acid residues and/or peptides are attached to the lysine-X compound to obtain compounds of Formula 3-I-A. It will be understood that other natural or non-natural amino acids described herein are also suitable for attachment to the surfactant X and are suitable for attaching

additional amino acid/peptides to obtain compounds of Formula 3-I-A. It will be understood that in another embodiment, compounds of Formula 3-I-A are prepared by attaching a full length or partial length peptide to a group X, followed by optional attachment of additional amino acid residues and/or peptides are attached to obtain compounds of Formula 3-I-A.

[0305] In a specific embodiment, provided herein are compounds selected from compounds of Table 3 in Figure 8 and Table 4 in Figure 9.

[0306] Also provided herein are pharmaceutical compositions comprising a therapeutically effective amount of a peptide product described above, or acceptable salt thereof, and at least one pharmaceutically acceptable carrier or excipient.

[0307] In some embodiments of the pharmaceutical compositions, the carrier is an aqueous-based carrier. In some embodiments of the pharmaceutical compositions, the carrier is a nonaqueous-based carrier. In some embodiments of the pharmaceutical compositions, the nonaqueous-based carrier is a hydrofluoroalkane-like solvent that may comprise sub-micron anhydrous α -lactose or other excipients.

[0308] Contemplated within the scope of embodiments presented herein is the reaction of an amino acid and/or a peptide comprising a linker amino acid U bearing a nucleophile, and a group X comprising a bearing a leaving group or a functional group that can be activated to contain a leaving group, for example a carboxylic acid, or any other reacting group, thereby allowing for covalent linkage of the amino acid and/or peptide to a surfactant X via the linker amino acid U to provide a peptide product of Formula 3-I-A.

[0309] Also contemplated within the scope of embodiments presented herein is the reaction of an amino acid and/or a peptide comprising a linker amino acid U bearing a bearing a leaving group or a functional group that can be activated to contain a leaving group, for example a carboxylic acid, or any other reacting group, and a group X comprising a nucleophilic group, thereby allowing for covalent linkage of the amino acid and/or peptide to a surfactant X via the linker amino acid U to provide a peptide product of Formula I-A.

[0310] It will be understood that, in one embodiment, Compounds of Formula 3-I-A are prepared by reaction of a linker amino acid U with X, followed by addition of further residues to U to obtain the peptide product of Formula 3-I-A. It will be understood that in an alternative embodiment, Compounds of Formula 3-I-A are prepared by reaction of a suitable peptide comprising a linker amino acid U with X, followed by optional addition of further residues to U, to obtain the peptide product of Formula 3-I-A.

[0311] Provided herein are methods for treating conditions associated with insulin resistance comprising administration of a compound of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V.

[0312] Provided herein are methods for treating diabetes, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, wound healing, insulin resistance, hyperglycemia,

hyperinsulinemia, metabolic syndrome, diabetic complications, elevated blood levels of free fatty acids or glycerol, hyperlipidemia, obesity, hypertriglyceridemia, atherosclerosis, acute cardiovascular syndrome, infarction, ischemic reperfusion or hypertension, comprising administering a therapeutically effective amount of a peptide product described above and herein to an individual in need thereof.

[0313] Provided herein are methods of reducing weight gain or inducing weight loss comprising administering to a subject in need thereof a therapeutically effective amount of a peptide product described above and herein to an individual in need thereof.

[0314] Provided herein are methods for treating mammalian conditions characterized by obesity-linked insulin resistance or the metabolic syndrome comprising administering to a subject in need thereof a weight loss-inducing or insulin-sensitizing amount of a peptide product described above and herein to an individual in need thereof.

[0315] In some embodiments, the condition to be treated is the metabolic syndrome (Syndrome X). In some embodiments, the condition to be treated is diabetes. In some embodiments, the condition to be treated is hyperlipidemia. In some embodiments, the condition to be treated is hypertension. In some embodiments, the condition to be treated is vascular disease including atherosclerosis, or the systemic inflammation characterized by elevated C reactive protein.

[0316] In some embodiments of the methods, the effective amount of the peptide product for administration is from about 0.1 µg/kg/day to about 100.0 µg/kg/day, or from 0.01 µg/kg/day to about 1 mg/kg/day or from 0.1 µg/kg/day to about 50 mg/kg/day.

[0317] Provided herein are methods of treating the metabolic syndrome, or its component diseases, comprising administering to a subject in need thereof a therapeutically effective amount of a peptide product described above. In some embodiments, the metabolic syndrome condition has progressed to diabetes.

[0318] Also provided herein is a covalently modified GLCR and/or GLP1R binding peptide or analog thereof, comprising a hydrophilic group as described herein; and a hydrophobic group covalently attached to the hydrophilic group. In specific embodiments, the covalently modified peptide and/or protein product comprises a hydrophilic group that is a saccharide and a hydrophobic group that is a C₁-C₂₀ alkyl chain or an aralkyl chain.

Insulin resistance

[0319] The risks associated with prolonged hyperglycemia include an increased risk of microvascular complications, sensory neuropathy, myocardial infarction, stroke, macrovascular mortality, and all-cause mortality. Type 2 diabetes is also linked causally with obesity, an additional global epidemic. At least \$232 billion were spent on treatment and prevention of

diabetes worldwide in 2007, with three quarters of that amount spent in industrialized countries on the treatment of long-term complications and on general care, such as efforts to prevent micro and macrovascular complications. In 2007, estimated indirect costs of diabetes (disability, lost productivity, and premature death due to diabetes) to the United States economy were \$58 billion.

[0320] Obesity leads to insulin resistance, a decreased ability of the cells in the body to react to insulin stimulation through decreased numbers of insulin receptors and a decreased coupling of those receptors to critical intracellular signaling systems. The obese state further leads to the "metabolic syndrome", a constellation of diseases (insulin resistance, hypertension, atherosclerosis, et al.) with very large healthcare consequences. If insulin resistance is diagnosed early enough, overt type 2 diabetes can be prevented or delayed, with lifestyle interventions aimed at reducing calorie intake and body fat and through drug treatment to normalize glycemic control. Despite treatment guidelines recommending early, aggressive intervention, many patients fail to reach targets for glycemic control. Many factors contribute to the failure to manage type 2 diabetes successfully including psychosocial and economic influences and shortcomings in the efficacy, convenience and tolerability profiles of available antidiabetic drugs. The peptide and/or protein products described herein are designed to overcome these shortcomings.

Incretin effect

[0321] The "incretin effect" is used to describe the phenomenon whereby a glucose load delivered orally produces a much greater insulin secretion than the same glucose load administered intravenously. This effect is mediated by at least two incretin hormones secreted by intestinal L-cells. Glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide 1 (GLP-1) were identified as incretins and it is thought that healthy individuals may derive up to 70% of their prandial insulin secretory response from the incretin effect.

[0322] Normally the incretin peptides are secreted as needed, in response to ingested nutrients, and have a short plasma half-life due to degradation by dipeptidyl peptidase IV (DPP-4) enzyme. In people with type 2 diabetes, pancreatic responsiveness to GLP-1 is impaired, but the insulin-secretory response can be restored with pharmacologic doses of human GLP-1 (Kieffer, T.J., et al. (1995) *Endocrinology* 136: 3585-3596). In addition, GLP-1 promotes beta-cell neogenesis and preservation (Aaboe, K., et al. (2008) *Diabetes Obes Metab* 10: 994-1003). GLP-1 has additional beneficial effects such as on cardiac function: for example it improves left ventricular function (Sokos, G.G., et al. (2006) *J Card Fail* 12: 694-699) in human subjects. GLP-1 also slows gastric emptying in humans and reduces appetite (Toft-Nielsen, M.B., et al. (1999) *Diabetes Care* 22: 1137-1143).

[0323] Treatment of diabetes patients with metabolically stable and long-acting analogs of GLP-1 is described in, for example, Drab, S.R. (2010) *Pharmacotherapy* 30: 609-624, suffers from issues related to convenience of use and side effects such as nausea, risk of pancreatitis

and thyroid carcinoma. GLP-1 analogs provide glucose-dependent stimulation of insulin secretion and lead to a reduced risk of hypoglycemia. In addition, while a number of the current treatments for diabetes cause weight gain, as described below, GLP-1 analogs induce satiety and a mild weight loss. Accordingly, in some embodiments, provided herein are GLP-1 analogs that are long acting and are administered at low doses thereby reducing side-effects associated with current treatments.

[0324] A number of peptide gut hormones are known to modulate appetite (Sanger, G.J. and Lee, K. (2008) *Nat Rev Drug Discov* 7: 241-254). Several peptides are derived from tissue-specific, enzymatic processing (prohormone convertases; PCs) of the preproglucagon gene product: e.g. glucagon, GLP-1, glucagon-like peptide-2 (GLP-2), glicentin and oxyntomodulin (OXM) (Drucker, D.J. (2005) *Nat Clin Pract Endocrinol Metab* 1: 22-31; Sinclair, E.M. and Drucker, D.J. (2005) *Physiology (Bethesda)* 20: 357-365). GLP-1, GLP-2, glicentin and OXM are co-secreted from L-cells in the gut in response to feeding. Preproglucagon is alternatively processed (PC2) to produce glucagon in the alpha cells in the pancreatic islets. The structure of OXM is essentially glucagon with a C-terminal extension of 8 residues.

[0325] In addition to the stimulation of insulin biosynthesis and of glucose-dependent insulin secretion, GLP-1 and its stable mimetics (e.g. Byetta) also cause modest weight loss in animal models (Mack, C.M., et al. (2006) *Int J Obes (Lond)* 30: 1332-1340) and in Type 2 diabetic patients (DeFronzo, R.A., et al. (2005) *Diabetes Care* 28: 1092-1100; Buse, J.B., et al. (2010) *Diabetes Care* 33: 1255-1261). Glucagon infusion reduces food intake in man (Geary, N., et al. (1992) *Am J Physiol* 262: R975-980), while continuous glucagon treatment of adipose tissue also promotes lipolysis (Heckemeyer, C.M., et al. (1983) *Endocrinology* 113: 270-276) and weight loss (Salter, J.M., et al. (1960) *Metabolism* 9: 753-768; Chan, E.K., et al. (1984) *Exp Mol Pathol* 40: 320-327). Glucagon has wide-ranging effects on energy metabolism (Heppner, K.M., et al. (2010) *Physiol Behav*). Glucagon, or analogs, can be used in a diagnostic mode for temporary paralysis of the intestinal tract. Thus at least two of the products from PC processing of the preproglucagon protein are linked to satiety and metabolic effects.

[0326] In rodents, repeated intraperitoneal administration of OXM, a third product of preproglucagon, has been associated with reduced white adipose tissue and a reduction in weight compared with controls (Dakin, C.L., et al. (2004) *Endocrinology* 145: 2687-2695). Oxm reduced food intake by 19.3% during an intravenous infusion administration to normal-weight humans and this effect continues for more than 12 hr. after infusion (Cohen, M.A., et al. (2003) *J Clin Endocrinol Metab* 88: 4696-4701). Treatment of volunteers over a 4 week period resulted in a sustained satiety effect and weight loss reflecting a decrease in body fat (Wynne, K., et al. (2005) *Diabetes* 54: 2390-2395).

[0327] OXM is structurally homologous with GLP-1 and glucagon, and activates both the glucagon receptor (GCGR) and the GLP-1 receptor (GLP1R), but with 10 to 100 fold less potency than the eponymous ligands. In addition, study of OXM interactions with GLP1R suggest it might have different effects on beta-arrestin recruitment compared to GLP-1 (Jorgensen, R., et al. (2007) *J Pharmacol Exp Ther* 322: 148-154), thus acting as a "biased"

ligand. A unique receptor for OXM was sought for a number of years, but has not yet been elucidated and it is assumed to act through the GLP1R and GCGR pathways. Accordingly, provided herein are methods for surfactant modification of gut peptides that allow for induction of satiety, weight loss, alleviation of insulin resistance and/or delay in progression of pre-diabetes to diabetes.

GLP-1

[0328] In view of the complex and interacting behavior of the products of the preproglucagon protein on satiety and metabolism described above, workers from multiple groups have studied the structure activity relationships on GLP-1 and glucagon structure. Residues throughout the sequences were shown to accept replacement. For example, replacement by Ala is well accepted in the N-terminal region of GLP-1, especially at 2, 3, 5, 8, 11, and 12 (Adelhorst, K., et al. (1994) J Biol Chem 269: 6275-6278).

[0329] It was shown that chimeric analogs with the ability to bind to GLP1R and GLCR could be achieved by grafting C-terminal residues from GLP-1 onto the N-terminus of glucagon (Hjorth, S.A., et al. (1994) J Biol Chem 269: 30121-30124). The residue at position 3 (acidic Glu in GLP1 or neutral Gln in Glucagon or OXM) reduces the affinity of glucagon (Runge, S., et al. (2003) J Biol Chem 278: 28005-28010) or OXM (Pocai, A., et al. (2009) Diabetes 58: 2258-2266) for the GLP1R. The effect on metabolic profile of animals treated with stabilized analogs of GLP-1 or glucagon or OXM with Gln in position 3 was studied (Day, J.W., et al. (2009) Nat Chem Biol 5: 749-757; Druce, M.R., et al. (2009) Endocrinology 150: 1712-1722; Pocai, A., et al. (2009) Diabetes 58: 2258-2266). These analogs were designed to have agonistic action on both GLP1R and on GCGR (Day, J.W., et al. US 2010/0190701 A1).

[0330] Chimeric analogs should have the desirable effects of the parent hormones acting on their receptors, and therefore similar to the effects of OXM, which apparently acts on both GLP-1R and GLCR: glucose-dependent insulin secretion and satiety, coupled with lipolysis and increased burning of fat due to glucagon. The analogs were shown to cause the desired effects of decreased weight and increased burning of fat. Such a profile would be attractive in the treatment of obesity, but a major challenge in obesity treatment is compliance. Although currently known full length analogs of glucagon and OXM, respectively, with affinity for both GLP-1R and GLCR can result in weight loss, these analogs are not optimized for the high bioavailability, pharmaceutical properties, and convenient delivery to patients that are necessary for optimal drug treatment regimens. Accordingly, provided herein are analogs of gut peptides (e.g., GLP, OXM, glucagon or the like) that allow for high bioavailability and/or long lasting effects for improved therapeutic outcome in treatment of conditions such as obesity and/or diabetes and/or the metabolic syndrome.

[0331] Additional factors for optimized treatment of the metabolic syndrome and diabetes with OXM-like molecules relate to the duration of treatment and the amount of glucagon action. For example, continuous treatment with analogs that activate GLP-1 and glucagon receptors (the

OXM pharmacological profile) can result in very large and rapid loss of fat mass (Day, J.W., et al. (2009) *Nat Chem Biol* 5: 749-757), but it can also cause the loss of lean muscle mass (Kosinski, J.R., et al. (2012) *Obesity* (Silver Spring): doi: 10.1038/oby.2012.67), which is unfavorable for a pharmaceutical in this class. For example, in the research article by Kosinski, J.R., et al., the natural hormone Oxm is administered continuously for 14 days from an Alzet minipump and results in a decrease of 30% in fat mass, but also caused a 7% decrease in lean mass (muscle).

[0332] Glucagon action is known to stimulate glycogenolysis, lipolysis and the increased burning of fat, but can also have catabolic effects on muscle. A successful treatment using an agent that combines GLP-1 and glucagon action (the OXM profile) will need to optimally cause the satiety and potentiated glucose-dependent insulin secretion of a GLP-1 analog with a judicious amount of glucagon action (fat burning). In addition, intermittent use of such an agent will provide the desired clinical profile of moderate, continuous weight loss, through loss of fat mass, with minimized loss of lean mass. Provided herein are molecules with a desirable combination of GLP-1 and OXM action as well as a tunable pharmacokinetic/pharmacodynamic profile to allow optimum use in therapy (for example in the metabolic syndrome, diabetes, obesity, and the like).

[0333] In one embodiment, the compounds of Formula 3-I-A, 3-III-A, 3-III-B and 3-V are designed to provide either glucagon-like activity or GLP-1 like activity. In a further embodiment, the compounds of Formula 3-I-A, 3-III-A, 3-III-B and 3-V provide tunable activity. For example, in one instance, the peptide products described herein (e.g., compounds in Table 3 of Figure 8 and Table 4 of Figure 9) have an EC₅₀ of less than about 500 nM, preferably less than about 50 nM, more preferably less than about 20 nM at receptors for both glucagon, and GLP-1. In another instance, the peptide products described herein (e.g., compounds in Table 3 of Figure 8 and Table 4 of Figure 9) are more potent (e.g., EC₅₀ of less than 10 nM, preferably less than 5 nM, more preferably about 1 nM) for the GLP-1 receptor and less potent for the glucagon receptor (e.g., EC₅₀ of less than 50 nM, preferably less than about 20 nM, more preferably about 5 nM) for the glucagon receptor. This tunability of biological activity allows for some retention of a judicious amount of glucagon action, thereby allowing for fat burning to occur, while also retaining the beneficial effects of potentiated glucose-dependent insulin secretion. OXM is structurally homologous with GLP-1 and glucagon, and activates both the glucagon receptor (GCGR) and the GLP-1 receptor (GLP1R). Accordingly, in some embodiments, the compounds of Formula 3-I-A, 3-III-A, 3-III-B and 3-V provide a tunable OXM-like biological activity. In some specific embodiments, the peptide products described herein comprise a peptide having amino acid residues 1-17 of GLP-1 and/or analogs thereof (e.g., analogs comprising modified non-natural amino acid replacements as described herein, cyclized lactam linkages as described herein, surfactant modifications as described herein, or a combination thereof). In some other embodiments, the peptide products described herein comprise a peptide having amino acid residues 1-16 of GLP-1 and/or analogs thereof (e.g., analogs comprising modified non-natural amino acid replacements as described herein, cyclized lactam linkages as described herein, surfactant modifications as described herein, or a combination thereof). In additional embodiments, the peptide products described herein comprise a peptide

having amino acid residues 1-18 of GLP-1 and/or analogs thereof (e.g., analogs comprising modified non-natural amino acid replacements as described herein, cyclized lactam linkages as described herein, surfactant modifications as described herein, or a combination thereof). Additionally the peptide products described herein comprise one or more residues (e.g., Aib, Ac4C) which provide helix stabilization of the designed compounds of Formula 3-I-A, 3-III-A, 3-III-B and 3-V, and compounds in Table 3 of Figure 8 and Table 4 of Figure 9.

[0334] It is believed that the glucagon subfamily of ligands bind to their receptors in a two domain mode common to a number of the class B of receptors (secretin class, G Protein-coupled Receptors (GPCR)). For GLP-1 it is felt that there is a N-terminal region of from residue 1 to about residue 16 which binds to the tops of the transmembrane helices (juxtomembrane region) and a helical C-terminal region from 17 to 31 which binds to the large, extracellular, N-terminal extension (ECD) of the receptor. The binding of these ligands focuses on the fact that N-terminally truncated analogs of these peptide ligands can still retain substantial binding affinity and selectivity for just the isolated ECD region of the receptor. Therefore it has been suggested that the N-terminal region is responsible for receptor activation while the C-terminal region is responsible for binding. It recently has been shown that short, N-terminal analogs of GLP-1 can be both potent binders as well as receptor activators (Mapelli, C., et al. (2009) *J Med Chem* 52: 7788-7799; Haque, T.S., et al. (2010) *Peptides* 31: 950-955; Haque, T.S., et al. (2010) *Peptides* 31: 1353-1360).

[0335] In addition, study of an x-ray crystal structure (Runge, S., et al. (2008) *J Biol Chem* 283: 11340-7) of the N-terminal region of the GLP1R with a truncated antagonist analogs of the GLP-1 mimic, exendin-4 (Byetta), bound in this region show that a critical ligand-binding region in the ECD is of high hydrophobicity (**Figure 10**). The sequence of exendin-4 beyond Glu15 interacts as an amphiphilic helix with this very hydrophobic region (Val^{19*}, Phe^{22*}, Trp^{25*}, Leu^{26*}). In one embodiment, truncated N-terminal fragments of GLP-1 or glucagon are modified to bind to GLCR and are covalently linked to a surfactant. The hydrophobic 1'-alkyl portion of the surfactant mimics and replaces the C-terminal region of the native hormone ligand and increases the peptides potency, efficacy, and duration of action. In addition, such analogs have major advantages due to their smaller size, which reduces their complexity, synthesis costs, and susceptibility to proteolysis. In addition smaller peptides are more readily absorbed through the nasal mucosa or gut enterocyte barrier.

[0336] Hypoglycemia is a condition of low blood sugar that can be life-threatening and is increasingly seen as more aggressive treatment of elevated blood sugar by intensive insulin treatment is being used in more patients. Hypoglycemia is seen when blood glucose levels drop too low to provide enough energy to the brain and muscles for the body's activities. Glucagon can be used to treat this condition and does so by stimulating the liver to break down glycogen to generate glucose and cause the blood glucose levels to rise toward the normal value. Analogs of glucagon that retain the ability to activate the GLCR may be used to achieve this desirable effect on blood glucose levels.

[0337] Analogs of GLP-1 that activate the GLP1R stimulate the production and, in the

presence of elevated blood glucose levels, release of insulin from the pancreas. This action results in efficient control and normalization of blood glucose levels, as seen with current products such as exenatide (Byetta®). In addition, such products appear to produce a decreased appetite and slow the movement of food from the stomach. Thus they are effective in treatment of diabetes through multiple mechanisms. Analogs that combine the effects of glucagon and GLP-1 that activate both the GLCR and the GLP1R may offer a benefit in the treatment of diabetes through a concerted action to suppress appetite, release insulin in a glucose-dependent fashion, assist in the protection from hypoglycemia and accelerate the burning of fat.

[0338] Such methods for treating hyperglycemia, including diabetes, diabetes mellitus type I, diabetes mellitus type II, or gestational diabetes, either insulin-dependent or non-insulin dependent, are expected to be useful in reducing complications of diabetes including nephropathy, retinopathy and vascular disease. Applications in cardiovascular disease encompass microvascular as well as macrovascular disease (Davidson, M.H., (2011) Am J Cardiol 108[suppl]:33B-41B; Gejl, M., et al. (2012) J Clin Endocrinol Metab 97:doi:10.1210/jc.2011-3456), and include treatment for myocardial infarction. Such methods for reducing appetite or promoting loss of body weight are expected to be useful in reducing body weight, preventing weight gain, or treating obesity of various causes, including drug-induced obesity, and reducing complications associated with obesity including vascular disease (coronary artery disease, stroke, peripheral vascular disease, ischemia reperfusion, etc.), hypertension, onset of diabetes type II, hyperlipidemia and musculoskeletal diseases.

[0339] As used herein, the term glucagon or GLP-1 analogs includes all pharmaceutically acceptable salts or esters thereof.

[0340] In one aspect, the peptides that are covalently modified and are suitable for methods described herein are truncated analogs of glucagon and/or the related hormone GLP-1, including and not limited to:

Glucagon:

His₁-Ser₂- Gln₃-Gly₄-Thr₅ Phe₆- Thr₇-Ser₈- Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-
Ser₁₆-Arg₁₇-Arg₁₈-Ala₁₉-Gln₂₀-Asp₂₁-Phe₂₂-Val₂₃-Gln₂₄-Trp₂₅-Leu₂₆-Met₂₇-Asn₂₈-Thr₂₉-
(SEQ. ID. NO. 632)

Oxyntomodulin:

His₁-Ser₂- Gln₃-Gly₄-Thr₅ Phe₆- Thr₇-Ser₈- Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-
Ser₁₆-Arg₁₇-Arg₁₈-Ala₁₉-Gln₂₀-Asp₂₁-Phe₂₂-Val₂₃-Gln₂₄-Trp₂₅-Leu₂₆-Met₂₇-Asn₂₈-Thr₂₉-
Lys₃₀-Arg₃₁-Asn₃₂-Arg₃₃-Asn₃₄-Asn₃₅-Ile₃₆-Ala₃₇ (SEQ. ID. NO. 633)

GLP-1 (using glucagon numbering):

His₁-Ala₂- Glu₃-Gly₄-Thr₅ Phe₆- Thr₇-Ser₈- Asp₉-Val₁₀-Ser₁₁-Ser₁₂-Tyr₁₃-Leu₁₄-Glu₁₅-
Gly₁₆-Gln₁₇-Ala₁₈-Ala₁₉-Lys₂₀-Glu₂₁-Phe₂₂-Ile₂₃-Ala₂₄-Trp₂₅-Leu₂₆-Val₂₇-Lys₂₈-Gly₂₉-Arg₃₀-
(SEQ. ID. NO. 634)

[0341] In some embodiments, a peptide product described herein has the structure of Formula

3-V:

aa₁-aa₂-aa₃-aa₄-aa₅-aa₆-aa₇-aa₈-aa₉-aa₁₀- aa₁₁-aa₁₂-aa₁₃-aa₁₄-aa₁₅-aa₁₆-aa₁₇-aa₁₈-aa₁₉-aa₂₀- aa₂₁-

aa₂₂-aa₂₃-aa₂₄-aa₂₅-aa₂₆-aa₂₇-aa₂₈-aa₂₉-aa₃₀-aa₃₁-aa₃₂-aa₃₃-aa₃₄-aa₃₅-aa₃₆-aa₃₇-Z FORMULA

3-V (SEQ. ID. NO. 635)

wherein:

U is a linking amino acid;

X is a surfactant-linked to the side chain of U;

Z is OH, or -NH-R³, wherein R³ is H or C₁-C₁₂ substituted or unsubstituted alkyl;

aa₁ is His, N-Ac-His, pGlu-His or N-R³-His;

aa₂ is Ser, Ala, Gly, Aib, Ac4c or Ac5c;

aa₃ is Gln, or Cit;

aa₄ is Gly, or D-Ala;

aa₅ is Thr, or Ser;

aa₆ is Phe, Trp, F2Phe, Me2Phe, or Nal(2);

aa₇ is Thr, or Ser;

aa₈ is Ser, or Asp;

aa₉ is Asp, or Glu;

aa₁₀ is Tyr, Leu, Met, Nal(2), Bip, or Bip2EtMeO;

aa₁₁ is Ser, Asn, or U(X);

aa₁₂ is Lys, Glu, Ser, Arg, or U(X);

aa₁₃ is absent, Tyr, Gln, Cit, or U(X);

aa₁₄ is absent, Leu, Met, Nle, or U(X);

aa₁₅ is absent, Asp, Glu, or U(X);

aa₁₆ is absent, Ser, Gly, Glu, Aib, Ac5c, Lys, Arg, or U(X);

aa₁₇ is absent, Arg, hArg, Gln, Glu, Cit, Aib, Ac4c, Ac5c, or U(X);

aa₁₈ is absent, Arg, hArg, Ala, Aib, Ac4c, Ac5c, or U(X);

aa₁₉ is absent, Ala, Val, Aib, Ac4c, Ac5c, or U(X);

aa₂₀ is absent, Gln, Lys, Arg, Cit, Glu, Aib, Ac4c, Ac5c, or U(X);

aa₂₁ is absent, Asp, Glu, Leu, Aib, Ac4c, Ac5c, or U(X);

aa₂₂ is absent, Phe, Trp, Nal(2), Aib, Ac4c, Ac5c, or U(X);

aa₂₃ is absent, Val, Ile, Aib, Ac4c, Ac5c, or U(X);

aa₂₄ is absent, Gln, Ala, Glu, Cit, or U(X);

aa₂₅ is absent, Trp, Nal(2), or U(X);

aa₂₆ is absent, Leu, U(X);

aa₂₇ is absent, Met, Val, Nle, Lys, or U(X);

aa₂₈ is absent, Asn, Lys, or U(X);

aa₂₉ is absent, Thr, Gly, Aib, Ac4c, Ac5c, or U(X);

aa₃₀ is absent, Lys, Aib, Ac4c, Ac5c, or U(X);

aa₃₁ is absent, Arg, Aib, Ac4c, Ac5c, or U(X);

aa₃₂ is absent, Asn, Aib, Ac4c, Ac5c, or U(X);

aa₃₃ is absent, Arg, Aib, Ac5c, or U(X);

aa₃₄ is absent, Asn, Aib, Ac4c, Ac5c, or U(X);

aa₃₅ is absent, Asn, Aib, Ac4c, Ac5c, or U(X);

aa₃₆ is absent, Ile, Aib, Ac4c, Ac5C, or U(X);


aa₃₆ is absent, Ala, Aib, Ac4c, Ac5C, or U(X);

aa₃₇ absent or U(X);

provided that one, or at least one of aa₁₁ - aa₃₇ is U(X).

[0342] In specific embodiments, the linking amino acid U, is a diamino acid like Lys or Orn, X is a modified surfactant from the 1-alkyl glycoside class linked to U, and Z is OH, or -NH-R₂, wherein R³ is H or C₁-C₁₂, or a PEG chain of less than 10Da.

[0343] In some embodiments, a peptide product described herein has the structure of Formula III-B:



wherein:

Z is OH, or -NH-R^3 , wherein R^3 is H or substituted or unsubstituted $\text{C}_1\text{-C}_{12}$ alkyl; or a PEG chain of less than 10Da;

aa_2 is Ser, Ala, Gly, Aib, Ac4c, or Ac5c;

aa_3 is Gln, or Cit;

aa_6 is Phe, Trp, F2Phe, Me2Phe, MePhe, or Nal2;

aa_{10} is Tyr, Leu, Met, Nal2, Bip, or Bip2EtMeO;

aa_{11} is Ser, Asn, or U;

aa_{12} is Lys, Glu, Ser or U(X);

aa_{13} is absent or Tyr, Gln, Cit, or U(X);

aa_{14} is absent or Leu, Met, Nle, or U(X);

aa_{15} is absent or Asp, Glu, or U(X);

aa_{16} is absent or Ser, Gly, Glu, Aib, Ac4c, Ac5c, Lys, R, or U(X);

aa_{17} is absent or Arg, hArg, Gln, Glu, Cit, Aib, Ac4c, Ac5c, or U(X);

aa_{18} is absent or Arg, hArg, Ala, Aib, Ac4c, Ac5c, or U(X);

aa_{19} is absent or Ala, Val, Aib, Ac4c, Ac5c, or U(X);

aa_{20} is absent or Gln, Lys, Arg, Cit, Glu, Aib, Ac4c, Ac5c, or U(X);

aa_{21} is absent or Asp, Glu, Leu, Aib, Ac4c, Ac5c, or U(X);


aa_{22} is absent or Phe, Aib, Ac4c, Ac5c, or U(X)

aa_{23} is absent or Val, Ile, Aib, Ac4c, Ac5c, or U(X);

wherein any two of $\text{aa}_1\text{-aa}_{23}$ are optionally cyclized through their side chains to form a lactam linkage; and

provided that one, or at least one of aa_{16} , aa_{17} , aa_{18} , aa_{19} , aa_{20} , aa_{21} , aa_{22} , aa_{23} or aa_{24} is the natural or unnatural amino acid U covalently attached to X.

[0344] In some specific embodiments of Formula 3-III-A, Formula 3-III-B and Formula 3-V, X has the structure:



Formula 3-I

wherein:

R^{1a} is a substituted or unsubstituted C₁-C₃₀ alkyl group;

R^{1b}, R^{1c}, and R^{1d} are H;

W¹ is -(C=O)-NH-;

W² is -O-; and

R² is a bond.

[0345] In some of the embodiments described above, R^{1a} is a C₁-C₂₀ alkyl group, a C₈-C₂₀ alkyl group, C₁₂-18 alkyl group or C₁₄-C₁₈ alkyl group.

[0346] In some embodiments of Formula 3-III-B, U is any linker amino acid described herein. Table 3 of Figure 8 and Table 4 of Figure 9 illustrate certain examples of peptides that covalently linked with surfactants as described herein.

[0347] Contemplated within the scope of embodiments presented herein are peptide products of Formula 3-I-A, Formula 3-III-A, Formula 3-III-B or Formula 3-V, wherein the peptide product comprises one, or, more than one surfactant groups (e.g., group X having the structure of Formula 3-1). In one embodiment, a peptide product of Formula 3-I-A, Formula 3-III-A, Formula 3-III-B or Formula 3-V, comprises one surfactant group. In another embodiment, a peptide product of Formula 3-I-A, Formula 3-III-A, Formula 3-III-B or Formula 3-V, comprises two surfactant groups. In yet another embodiment, a peptide product of Formula 3-I-A, Formula 3-III-A, Formula 3-III-B or Formula 3-V, comprises three surfactant groups.

[0348] Recognized herein is the importance of certain portions of SEQ. ID. NO. 632 for the treatment of conditions associated with insulin resistance and/or cardiovascular conditions. Accordingly, provided herein is a method of treating diabetes in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₁₇ of SEQ. ID. NO. 632 to the individual in need thereof.

[0349] In a further embodiment, provided herein is a method of treating diabetes in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₁₈ of **SEQ. ID. NO. 632** to the individual in need thereof.

[0350] In another embodiment, provided herein is a method of treating diabetes in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₁₉ of **SEQ. ID. NO. 632** to the individual in need thereof.

[0351] In another embodiment, provided herein is a method of treating diabetes in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₂₀ of **SEQ. ID. NO. 632** to the individual in need thereof.

[0352] In an additional embodiment, the administration of the said glucagon analog described above causes weight loss.

[0353] Recognized herein is the importance of certain portions of **SEQ. ID. NO. 303** for the treatment of conditions associated with insulin resistance and/or cardiovascular conditions. Accordingly, provided herein is a method of treating diabetes in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₁₇ of **SEQ. ID. NO. 303** to the individual in need thereof.


[0354] In a further embodiment, provided herein is a method of treating diabetes in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₁₈ of **SEQ. ID. NO. 303** to the individual in need thereof.

[0355] In another embodiment, provided herein is a method of treating diabetes in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₁₉ of **SEQ. ID. NO. 303** to the individual in need thereof.

[0356] In another embodiment, provided herein is a method of treating diabetes in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₂₀ of **SEQ. ID. NO. 303** to the individual in need thereof.

[0357] In an additional embodiment, the administration of the said glucagon analog described above causes weight loss.

[0358] In any of the embodiments described above, the said glucagon analog is modified with a surfactant X of Formula 3-I:



wherein:

R^{1a} is independently, at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group, or a steroid nucleus containing moiety;

R^{1b} , R^{1c} , and R^{1d} are each, independently at each occurrence, a bond, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group;

W^1 is independently, at each occurrence, $-CH_2-$, $-CH_2-O-$, $-(C=O)-$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$, or $-CH_2-S-$;


W^2 is $-O-$, $-CH_2-$ or $-S-$;

R^2 is independently, at each occurrence, a bond to U, H, a substituted or unsubstituted C_1 - C_{30} alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group, $-NH_2$, $-SH$, C_2 - C_4 -alkene, C_2 - C_4 -alkyne, $-NH(C=O)-CH_2-Br$, $-(CH_2)_m$ -maleimide, or $-N_3$;

n is 1, 2 or 3; and

m is 1-10.

[0359] In a specific embodiment, the said glucagon analog is modified with a surfactant, X having the structure:



wherein:

R^{1a} is a substituted or unsubstituted C_1 - C_{30} alkyl group;

R^{1b} , R^{1c} , and R^{1d} are H;

W^1 is $-(C=O)-NH-$;

W^2 is $-O-$; and

R^2 is a bond.

[0360] In some of the embodiments described above, R^{1a} is a C_1 - C_{20} alkyl group, a C_8 - C_{20} alkyl group, C_{12} - C_{18} alkyl group or C_{14} - C_{18} alkyl group.

[0361] Also provided herein is a method of treating a cardiovascular disease in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa_1 - aa_{17} of **SEQ. ID. NO. 632** to the individual in need thereof.

[0362] Also provided herein is a method of treating a cardiovascular disease in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa_1 - aa_{18} of **SEQ. ID. NO. 632** to the individual in need thereof.

[0363] Also provided herein is a method of treating a cardiovascular disease in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa_1 - aa_{19} of **SEQ. ID. NO. 632** to the individual in need thereof.

[0364] Also provided herein is a method of treating a cardiovascular disease in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa_1 - aa_{20} of **SEQ. ID. NO. 632** to the individual in need thereof.

[0365] In some cases for the embodiments described above, the said glucagon analog is administered when the cardiovascular disease is associated with an ischemic event.

[0366] Also provided herein is a method of treating a cardiovascular disease in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa_1 - aa_{17} of **SEQ. ID. NO. 303** to the individual in need thereof.


[0367] Also provided herein is a method of treating a cardiovascular disease in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₁₈ of **SEQ. ID. NO. 303** to the individual in need thereof.

[0368] Also provided herein is a method of treating a cardiovascular disease in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₁₉ of **SEQ. ID. NO. 303** to the individual in need thereof.

[0369] Also provided herein is a method of treating a cardiovascular disease in an individual in need thereof comprising administration of a therapeutically effective amount of a glucagon analog comprising amino acid residues aa₁-aa₂₀ of **SEQ. ID. NO. 303** to the individual in need thereof.

[0370] In some cases for the embodiments described above, the said glucagon analog is administered when the cardiovascular disease is associated with an ischemic event.

[0371] In any of the embodiments described above, the said glucagon analog is modified with a surfactant X of Formula 3-I:



wherein:

R^{1a} is independently, at each occurrence, a bond, H, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted alkoxyaryl group, a substituted or unsubstituted aralkyl group, or a steroid nucleus containing moiety;

R^{1b}, R^{1c}, and R^{1d} are each, independently at each occurrence, a bond, H, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted aralkyl group;

W¹ is independently, at each occurrence, -CH₂-, -CH₂-O-, -(C=O)-, -(C=O)-O-, -(C=O)-NH-, -(C=S)-, -(C=S)-NH-, or -CH₂-S-;

W² is -O-, -CH₂- or -S-;


R² is independently, at each occurrence, a bond to U, H, a substituted or unsubstituted C₁-C₃₀ alkyl group, a substituted or unsubstituted alkoxyaryl group, or a substituted or unsubstituted

aralkyl group, $-\text{NH}_2$, $-\text{SH}$, $\text{C}_2\text{-C}_4\text{-alkene}$, $\text{C}_2\text{-C}_4\text{-alkyne}$, $-\text{NH}(\text{C}=\text{O})\text{-CH}_2\text{-Br}$, $-(\text{CH}_2)_m$ -maleimide, or $-\text{N}_3$;

n is 1, 2 or 3; and

m is 1-10.

[0372] In a specific embodiment, the said glucagon analog is modified with a surfactant, X having the structure:



Formula 3-I

wherein:

R^{1a} is a substituted or unsubstituted $\text{C}_1\text{-C}_{30}$ alkyl group;

R^{1b} , R^{1c} , and R^{1d} are H;

W^1 is $-(\text{C}=\text{O})\text{-NH-}$;

W^2 is $-\text{O-}$; and

R^2 is a bond.

[0373] In some of the embodiments described above, R^{1a} is a $\text{C}_1\text{-C}_{20}$ alkyl group, a $\text{C}_8\text{-C}_{20}$ alkyl group, $\text{C}_{12}\text{-C}_{18}$ alkyl group or $\text{C}_{14}\text{-C}_{18}$ alkyl group.

[0374] Modifications at the amino or carboxyl terminus may optionally be introduced into the peptides (e.g., glucagon or GLP-1) (Nestor, J.J., Jr. (2009) Current Medicinal Chemistry 16: 4399 - 4418). For example, the peptides can be truncated or acylated on the N-terminus to yield peptides analogs exhibiting low efficacy, partial agonist and antagonist activity, as has been seen for some peptides (Gourlet, P., et al. (1998) Eur J Pharmacol 354: 105-111, Gozes, I. and Furman, S. (2003) Curr Pharm Des 9: 483-494)). For example, deletion of the first 6 residues of bPTH yields antagonistic analogs (Mahaffey, J.E., et al. (1979) J Biol Chem 254: 6496-6498; Goldman, M.E., et al. (1988) Endocrinology 123: 2597-2599) and a similar operation on peptides described herein generates potent antagonistic analogs. Other modifications to the N-terminus of peptides, such as deletions or incorporation of D-amino

acids such as D-Phe also can give potent and long acting agonists or antagonists when substituted with the modifications described herein such as long chain alkyl glycosides. Such agonists and antagonists also have commercial utility and are within the scope of contemplated embodiments described herein.

[0375] Also contemplated within the scope of embodiments described herein are surfactants covalently attached to peptide analogs, wherein the native peptide is modified by acetylation, acylation, PEGylation, ADP-ribosylation, amidation, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-link formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, (Nestor, J.J., Jr. (2007) *Comprehensive Medicinal Chemistry II* 2: 573-601, Nestor, J.J., Jr. (2009) *Current Medicinal Chemistry* 16: 4399 - 4418, Creighton, T.E. (1993), Wold, F. (1983) *Posttranslational Covalent Modification of Proteins* 1-12, Seifter, S. and England, S. (1990) *Methods Enzymol* 182: 626-646, Rattan, S.I., et al. (1992) *Ann N Y Acad Sci* 663: 48-62). Also contemplated within the scope of embodiments described herein are peptides that are branched or cyclic, with or without branching. Cyclic, branched and branched circular peptides result from post-translational natural processes and are also made by suitable synthetic methods. In some embodiments, any peptide product described herein comprises a peptide analog described above that is then covalently attached to an alkyl-glycoside surfactant moiety.

[0376] Also contemplated within the scope of embodiments presented herein are peptide chains substituted in a suitable position by the substitution of the analogs claimed herein by acylation on a linker amino acid, at for example the ϵ -position of Lys, with fatty acids such as octanoic, decanoic, dodecanoic, tetradecanoic, hexadecanoic, octadecanoic, 3-phenylpropanoic acids and the like, with saturated or unsaturated alkyl chains (Zhang, L. and Bulaj, G. (2012) *Curr Med Chem* 19: 1602-1618). Non-limiting, illustrative examples of such analogs are:

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Ser₁₆-Arg₁₇-Lys(N-epsilon-dodecanoyl)₁₈-Aib₁₉-NH₂, (**SEQ. ID. NO. 636**)

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Ser₁₆-Arg₁₇-Lys(N-epsilon-tetradecanoyl)₁₈-Ac₄C₁₉-NH₂, (**SEQ. ID. NO. 637**)

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Ser₁₆-Arg₁₇-Lys(N-epsilon-hexadecanoyl)₁₈-Aib₁₉-NH₂, (**SEQ. ID. NO. 638**)

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Aib₁₆-

Arg₁₇-Lys(N-epsilon-dodecanoyl)₁₈-NH₂, (**SEQ. ID. NO. 639**)

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Aib₁₆-
Arg₁₇-Lys(N-epsilon-tetradecanoyl)₁₈-NH₂, (**SEQ. ID. NO. 640**)

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Aib₁₆-
Arg₁₇-Lys(N-epsilon-hexadecanoyl)₁₈-NH₂, (**SEQ. ID. NO. 641**)

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Ser₁₆-
Arg₁₇-Lys(N-epsilon-(gamma-glutamyl)-N-alpha-tetradecanoyl)₁₈-Aib₁₉-NH₂ (**SEQ. ID. NO. 642**), and the like.

[0377] In further embodiments, a peptide chain is optionally substituted in a suitable position by reaction on a linker amino acid, for example the sulfhydryl of Cys, with a spacer and a hydrophobic moiety such as a steroid nucleus, for example a cholesterol moiety. In some of such embodiments, the modified peptide further comprises one or more PEG chains. Non-limiting examples of such molecules are:

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-Lys₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-Aib₁₆-
Arg₁₇-Cys(S-(3-(PEG4-aminoethylacetamide-Cholesterol)))₁₈-Aib₁₉-NH₂ (**SEQ. ID. NO. 643**),

His₁-Aib₂-Gln₃-Gly₄-Thr₅-Phe₆-Thr₇-Ser₈-Asp₉-Tyr₁₀-Ser₁₁-cyclo(Glu₁₂-Tyr₁₃-Leu₁₄-Asp₁₅-
Lys₁₆)-Arg₁₇-Cys(S-(3-(PEG4-aminoethylacetamide-Cholesterol)))₁₈-NH₂ (**SEQ. ID. NO. 644**).

[0378] Aside from the twenty standard amino acids, there are a vast number of "nonstandard amino acids" or unnatural amino acids that are known to the art and that may be incorporated in the compounds described herein, as described above. Other nonstandard amino acids are modified with reactive side chains for conjugation (Gauthier, M.A. and Klok, H.A. (2008) Chem Commun (Camb) 2591-2611; de Graaf, A.J., et al. (2009) Bioconjug Chem 20: 1281-1295). In one approach, an evolved tRNA/ tRNA synthetase pair and is coded in the expression plasmid by the amber suppressor codon (Deiters, A, et al. (2004). Bio-org. Med. Chem. Lett. 14, 5743-5). For example, p-azidophenylalanine was incorporated into peptides and then reacted with a functionalized surfactant, or a PEG polymer having an acetylene moiety in the presence of a reducing agent and copper ions to facilitate an organic reaction known as "Huisgen [3+2] cycloaddition." A similar reaction sequence using the reagents described herein containing an acetylene modified alkyl glycoside or PEG modified glycoside will result in PEGylated or alkyl glycoside modified peptides. For peptides of less than about 50 residues, standard solid phase synthesis is used for incorporation of said reactive amino acid residues at the desired position in the chain. Such surfactant-modified peptides and/or proteins offer a different spectrum of pharmacological and medicinal properties than peptides modified by PEG incorporation alone.

[0379] The skilled artisan will appreciate that numerous permutations of the peptide analogs are possible and, provided that an amino acid sequence has an incorporated surfactant moiety, will possess the desirable attributes of surfactant modified peptide products described herein.

Certain Definitions

[0380] As used in the specification, "a" or "an" means one or more. As used in the claim(s), when used in conjunction with the word "comprising," the words "a" or "an" mean one or more. As used herein, "another" means at least a second or more.

[0381] As used herein, the one- and three-letter abbreviations for the various common amino acids are as recommended in Pure Appl. Chem. 31, 639-645 (1972) and 40, 277-290 (1974) and comply with 37 CFR § 1.822 (55 FR 18245, May 1, 1990). The abbreviations represent L-amino acids unless otherwise designated as D- or DL. Certain amino acids, both natural and non-natural, are achiral, e.g., glycine, α -amino-isobutyric acid (Aib). All peptide sequences are presented with the N-terminal amino acid on the left and the C-terminal amino acid on the right.


[0382] An "alkyl" group refers to an aliphatic hydrocarbon group. Reference to an alkyl group includes "saturated alkyl" and/or "unsaturated alkyl", i.e., an alkene or an alkyne. The alkyl group, whether saturated or unsaturated, includes branched, straight chain, or cyclic groups. An alkyl group is optionally substituted with substituents including and not limited to oxo, halogen, aryl, cycloalkyl, hydrophobic natural product such as a steroid, an aralkyl chain (including alkoxyaryl), alkyl chain containing an acyl moiety, or the like. In some embodiments, an alkyl group is linked to the N α -position of a residue (e.g., Tyr or Dmt) in a peptide. This class is referred to as N-alkyl and comprises straight or branched alkyl groups from C₁-C₁₀, or an aryl substituted alkyl group such as benzyl, phenylethyl and the like. In some embodiments, an alkyl moiety is a 1-alkyl group that is in glycosidic linkage (typically to the 1-position of, for example, glucose) to the saccharide moiety. Such a 1-alkyl group is a C₁-C₃₀ alkyl group.

[0383] An "aryl" group refers to an aromatic ring wherein each of the atoms forming the ring is a carbon atom. Aryl rings described herein include rings having five, six, seven, eight, nine, or more than nine carbon atoms. Aryl groups are optionally substituted with substituents selected from halogen, alkyl, acyl, alkoxy, alkylthio, sulfonyl, dialkyl-amino, carboxyl esters, cyano or the like. Examples of aryl groups include, but are not limited to phenyl, and naphthalenyl.

[0384] The term "acyl" refers to a C₁-C₂₀ acyl chain. This chain may comprise a linear aliphatic chain, a branched aliphatic chain, a chain containing a cyclic alkyl moiety, a hydrophobic natural product such as a steroid, an aralkyl chain, or an alkyl chain containing an acyl moiety.

[0385] The term "steroid nucleus" refers to the core of steroids comprising an arrangement of

four fused rings designated A, B, C and D as shown below:



Examples of steroid nucleus containing moieties include, and are not limited to, cholesterol and the like.

[0386] As used herein, a "therapeutic composition" can comprise an admixture with an aqueous or organic carrier or excipient, and can be compounded, for example, with the usual nontoxic, pharmaceutically acceptable carriers for tablets, pellets, capsules, lyophilizates, suppositories, solutions, emulsions, suspensions, or other form suitable for use. The carriers, in addition to those disclosed above, can include alginate, collagen, glucose, lactose, mannose, gum acacia, gelatin, mannitol, starch paste, magnesium trisilicate, talc, corn starch, keratin, colloidal silica, potato starch, urea, medium chain length triglycerides, dextrans, and other carriers suitable for use in manufacturing preparations, in solid, semisolid, or liquid form. In addition, auxiliary stabilizing, thickening or coloring agents can be used, for example a stabilizing dry agent such as triulose.

[0387] As used herein, a "pharmaceutically acceptable carrier" or "therapeutic effective carrier" is aqueous or nonaqueous (solid), for example alcoholic or oleaginous, or a mixture thereof, and can contain a surfactant, emollient, lubricant, stabilizer, dye, perfume, preservative, acid or base for adjustment of pH, a solvent, emulsifier, gelling agent, moisturizer, stabilizer, wetting agent, time release agent, humectant, or other component commonly included in a particular form of pharmaceutical composition. Pharmaceutically acceptable carriers are well known in the art and include, for example, aqueous solutions such as water or physiologically buffered saline or other solvents or vehicles such as glycols, glycerol, and oils such as olive oil or injectable organic esters. A pharmaceutically acceptable carrier can contain physiologically acceptable compounds that act, for example, to stabilize or to increase the absorption of specific inhibitor, for example, carbohydrates, such as glucose, sucrose or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins or other stabilizers or excipients.

[0388] The pharmaceutical compositions can also contain other pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such "substances" include, but are not limited to, pH adjusting and buffering agents, tonicity adjusting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, etc. Additionally, the peptide, or variant thereof, suspension may include lipid-protective agents which protect lipids against free-radical and lipid-peroxidative damages on storage. Lipophilic free-radical quenchers, such as alpha-tocopherol and water-soluble iron-specific chelators, such as ferrioxamine, are suitable.

[0389] As used herein, a "surfactant" is a surface active agent that modifies interfacial tension of water. Typically, surfactants have one lipophilic and one hydrophilic group or region in the molecule. Broadly, the group includes soaps, detergents, emulsifiers, dispersing and wetting

agents, and several groups of antiseptics. More specifically, surfactants include stearyltriethanolamine, sodium lauryl sulfate, sodium taurocholate, laurylaminopropionic acid, lecithin, benzalkonium chloride, benzethonium chloride and glycerin monostearate; and hydrophilic polymers such as polyvinyl alcohol, polyvinylpyrrolidone, polyethyleneglycol (PEG), carboxymethylcellulose sodium, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose and hydroxypropylcellulose or alkyl glycosides. In some embodiments, a surfactant is a non-ionic surfactant (e.g., an alkyl glycoside surfactant). In some embodiments, a surfactant is an ionic surfactant.

[0390] As used herein, "alkyl glycoside" refers to any sugar joined by a linkage to any hydrophobic alkyl, as is known in the art. The hydrophobic alkyl can be chosen of any desired size, depending on the hydrophobicity desired and the hydrophilicity of the saccharide moiety. In one aspect, the range of alkyl chains is from 1 to 30 carbon atoms; or from 6 to 16 carbon atoms.

[0391] As used herein, "saccharide" is inclusive of monosaccharides, oligosaccharides or polysaccharides in straight chain or ring forms. Oligosaccharides are saccharides having two or more monosaccharide residues. Some examples of the many possible saccharides suitable for use in functionalized form include glucose, galactose, maltose, melibiose, maltotriose, maltotetraose, sucrose, trehalose or the like.

[0392] As used herein, "sucrose esters" are sucrose esters of fatty acids. Sucrose esters can take many forms because of the eight hydroxyl groups in sucrose available for reaction and the many fatty acid groups, from acetate on up to larger, more bulky fats that can be reacted with sucrose. This flexibility means that many products and functionalities can be tailored, based on the fatty acid moiety used. Sucrose esters have food and non-food uses, especially as surfactants and emulsifiers, with growing applications in pharmaceuticals, cosmetics, detergents and food additives. They are biodegradable, non-toxic and mild to the skin.

[0393] As used herein, a "suitable" alkyl glycoside means one that is nontoxic and nonionic. In some instances, a suitable alkyl glycoside reduces the immunogenicity or aggregation and increases the bioavailability of a compound when it is administered with the compound via the ocular, nasal, nasolacrimal, sublingual, buccal, inhalation routes or by injection routes such as the subcutaneous, intramuscular, or intravenous routes. Suitable compounds can be determined using the methods known to the art and those set forth in the examples.

[0394] A "linker amino acid" is any natural or unnatural amino acid that comprises a reactive functional group (de Graaf, A.J., et al. (2009) *Bioconjug Chem* 20: 1281-1295) that is used for covalent linkage with the functionalized surfactant. By way of example, in some embodiments, a linker amino acid is Lys, or Orn having a reactive functional group $-NH_2$; or Cys, having a reactive functional group $-SH$; or Asp or Glu, having a reactive functional group $-C(=O)-OH$. By way of example, in some other embodiments, a linker amino acid is any amino acid having a reactive functional group such as $-OH$, $-N_3$, haloacetyl or an acetylenic group that is used for formation of a covalent linkage with a suitably functionalized surfactant.

[0395] As used herein, a "functionalized surfactant" is a surfactant comprising a reactive group suitable for covalent linkage with a linker amino acid. By way of example, in some embodiments, a functionalized surfactant comprises a carboxylic acid group (e.g., at the 6-position of a monosaccharide) as the reactive group suitable for covalent linkage with a linker amino acid. By way of example, in some embodiments, a functionalized surfactant comprises a -NH₂ group, a -N₃ group, an acetylenic group, a haloacetyl group, a -O-NH₂ group, or a -(CH₂)_m-maleimide group, e.g., at the 6-position of a monosaccharide (as shown in Scheme 6), that allows for covalent linkage with a suitable linker amino acid. In some embodiments, a functionalized surfactant is a compound of Formula IV as described herein.




[0396] As used herein, the term "peptide" is any peptide comprising two or more amino acids. The term peptide includes short peptides (e.g., peptides comprising between 2 - 14 amino acids), medium length peptides (15-50) or long chain peptides (e.g., proteins). The terms peptide, medium length peptide and protein may be used interchangeably herein. As used herein, the term "peptide" is interpreted to mean a polymer composed of amino acid residues, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof linked via peptide bonds, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof. Synthetic peptides can be synthesized, for example, using an automated peptide synthesizer.








[0397] Peptides may contain amino acids other than the 20 gene encoded amino acids. "Peptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, and are well-known to those of skill in the art. It will be appreciated that in some embodiments, the same type of modification is present in the same or varying degree at several sites in a given peptide. Also, a given peptide, in some embodiments, contains more than one type of modifications. Modifications occur anywhere in a peptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini.

[0398] Accordingly, also contemplated within the scope of embodiments described herein are surfactants covalently attached to peptides that are modified, including, for example, modification by, acetylation, acylation, PEGylation, ADP-ribosylation, amidation, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-link formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, (Nestor, J.J., Jr. (2007) Comprehensive Medicinal Chemistry II 2: 573-601, Nestor, J.J., Jr. (2009) Current Medicinal Chemistry 16: 4399 - 4418, Creighton, T.E. (1993), Wold, F. (1983) Posttranslational Covalent


Modification of Proteins 1-12, Seifter, S. and England, S. (1990) *Methods Enzymol* 182: 626-646, Rattan, S.I., et al. (1992) *Ann N Y Acad Sci* 663: 48-62). Also contemplated within the scope of embodiments described herein are peptides that are branched or cyclic, with or without branching. Cyclic, branched and branched circular peptides result from post-translational natural processes and are also made by suitable synthetic methods.

[0399] The term peptide includes peptides or proteins that comprise natural and unnatural amino acids or analogs of natural amino acids. As used herein, peptide and/or protein "analogs" comprise non-natural amino acids based on natural amino acids, such as tyrosine analogs, which includes para-substituted tyrosines, ortho-substituted tyrosines, and meta substituted tyrosines, wherein the substituent on the tyrosine comprises an acetyl group, a benzoyl group, an amino group, a hydrazine, an hydroxyamine, a thiol group, a carboxy group, a methyl group, an isopropyl group, a C₂-C₂₀ straight chain or branched hydrocarbon, a saturated or unsaturated hydrocarbon, an O-methyl group, a polyether group, a halogen, a nitro group, or the like. Examples of Tyr analogs include 2,4-dimethyl-tyrosine (Dmt), 2,4-diethyl-tyrosine, O-4-allyl-tyrosine, 4-propyl-tyrosine, C α -methyl-tyrosine and the like. Examples of lysine analogs include ornithine (Orn), homo-lysine, C α -methyl-lysine (CMeLys), and the like. Examples of phenylalanine analogs include, but are not limited to, meta-substituted phenylalanines, wherein the substituent comprises a methoxy group, a C₁-C₂₀ alkyl group, for example a methyl group, an allyl group, an acetyl group, or the like. Specific examples include, but are not limited to, 2,4,6-trimethyl-L-phenylalanine (Tmp), O-methyl-tyrosine, 3-(2-naphthyl)alanine (Nal(2)), 3-(1-naphthyl)alanine (Nal(1)), 3-methyl-phenylalanine, 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid (Tic), fluorinated phenylalanines, isopropyl-phenylalanine, p-azidophenylalanine, p-acyl-phenylalanine, p-benzoyl-phenylalanine, p-iodo-phenylalanine, p-bromophenylalanine, p-amino-phenylalanine, and isopropyl-phenylalanine, and the like. Among the vast array of nonstandard or unnatural amino acids known to the art and used in peptide analog design are C-alpha-disubstituted amino acids such as Aib, C α -diethylglycine (Deg), aminocyclopentane-1-carboxylic acid (Ac4c), aminocyclopentane-1-carboxylic acid (Ac5c), and the like. Such amino acids frequently lead to a restrained structure, often biased toward an alpha helical structure (Kaul, R. and Balaram, P. (1999) *Bioorg Med Chem* 7: 105-117). Additional examples of such unnatural amino acids useful in analog design are homo-arginine (Har), and the like. Substitution of reduced amide bonds in certain instances leads to improved protection from enzymatic destruction or alters receptor binding. By way of example, incorporation of a Tic-Phe dipeptide unit with a reduced amide bond between the residues (designated as Tic- Ψ [CH₂-NH]- Ψ -Phe) reduces enzymatic degradation. Accordingly, also contemplated within the scope of embodiments described herein are surfactants covalently attached to peptides that comprise modified amino acids and/or peptide analogs described above. Certain non-natural amino acids are shown below.



2,6-dimethyl-L-tyrosine
(Dmt)2,4,6-trimethyl-L-phenylalanine
(Tmp)2-(1-naphthyl-L-alanine
(Nal(1))2-(2-naphthyl-L-alanine
(Nal(2))1,2,3,4-tetrahydroisoquinoline-
3-carboxylic acid
(Tic)alpha-amino isobutyric acid
(Aib)2,2-diethylglycine
(Deg)2-aminocyclobutane-
1-carboxylic acid
(Ac4c)aminocyclopentane-
1-carboxylic acid
(Ac5c)

2-L-biphenyl-alanine (Bip)

2-L-(2'-ethyl,4'-methoxy)-
biphenyl-alanine
(Bip2EtMeO)

[0400] As used herein, "opioid peptides" are short sequences of amino acids that bind to opioid receptors in the body. In some embodiments, opioid peptides are endogenous peptides, such as, for example, endorphins, enkephalins, endomorphins, dermorphins or the like. In some embodiments, opioid peptides are derived from endogenous opioid peptides (e.g., pseudopeptides, constrained peptides, alpha-methyl analogs, or the like). In some embodiments, opioid peptides are exogenous and/or synthetic and comprise modified amino acids and/or unnatural amino acids that mimic the effects of opioid peptides.

[0401] As used herein, the term "variant" is interpreted to mean a peptide that differs from a reference peptide, but retains essential properties. A typical variant of a peptide differs in amino acid sequence from another, reference peptide. Generally, differences are limited so that the sequences of the reference peptide and the variant are closely similar overall and, in many regions, identical. A variant and reference peptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. Non-naturally occurring variants of peptides may be made by mutagenesis techniques, by direct synthesis, and by other suitable recombinant methods.

[0402] Reference now will be made in detail to various embodiments and particular applications of the covalently modified peptides and/or proteins described herein. While the covalently modified peptides and/or proteins will be described in conjunction with the various embodiments and applications, it will be understood that such embodiments and applications are exemplary and are not intended to limit the scope of the embodiments described herein.

Peptides

[0403] There are many important roles played by peptides in the body and some commercial opportunities have been exploited (Nestor, J.J., Jr. (2009) *Current Medicinal Chemistry* 16: 4399 - 4418; Stevenson, C.L. (2009) *Curr Pharm Biotechnol* 10: 122-137). However even these recognized targets (Tyndall, J.D., et al. (2005) *Chem Rev* 105: 793-826) and products continue to suffer from deficiencies in duration of action and bioavailability. In some embodiments, the improved peptides described herein provide longer duration of action and/or bioavailability and/or therapeutic efficacy compared to currently available commercial products. Some illustrative examples of peptides that represent attractive commercial targets for analog design (agonists and antagonists) for clinical development include, for example, members of the Class B, G Protein-Coupled Receptor (GPCR) ligands and related peptides ("Secretin family"): Secretin, Parathyroid Hormone (PTH), Parathyroid Hormone-related Protein (PTHrP), Glucagon, Glucagon Like Protein-1 and -2 (GLP-1, GLP-2), Glucose-dependent Insulinotropic Peptide (GIP), Oxyntomodulin, Pituitary Adenylate Cyclase-Activating Peptide (PACAP), Vasoactive Intestinal Peptide (VIP), Amylin (and analogs such as pramlintide, devalintide, et al.), Calcitonin (and analogs such as salmon calcitonin, elcatonin, et al.), calcitonin gene-related peptide (CGRP), Adrenomedullin, Corticotrophin-Releasing Factor family (CRF, Xerecept; Urocortin), and the like, including synthetic analogs thereof which would be improved as clinical products through further modification by the methods described herein. Also contemplated within the scope of embodiments presented herein are Opioid peptide families; such peptides would benefit from the methods of peptide modification described herein to give increased duration of action and increased specificity. By way of example analogs of the endomorphin, dynorphin, enkephalin, dermorphin, casomorphin, et al. peptide families offer attractive therapeutic approaches to the treatment of pain, addiction, et al. Additional attractive peptide targets for modification to yield peptide products described herein are the

hypothalamic hormones, for example gonadotrophin hormone-releasing hormone and its analogs (for example nafarelin, goserelin, triptorelin, leuprorelin, fertirelin, histrelin, buserelin, ganirelix, cetrorelix, degarelix, deslorelin and the like), adrenocorticotrophin, somatostatin (for example octreotide, lanreotide, valpreotide et al.), thyrotropin-releasing hormone, growth hormone-releasing hormone, and neurotensin. A further example of attractive commercial targets are the pituitary hormones and analogs such as vasopressin (desmopressin, and the like), oxytocin and analogs, thyroid hormone stimulating hormone, prolactin, growth hormone, luteinizing hormone, follicular-stimulating hormone, alpha melanocyte-stimulating hormone analogs (melanotan analogs), and the like, as well as their analogs. Growth factors are an important class of molecules that may be advantageously modified using the methods described herein to yield improved pharmaceutical candidates, for example insulin (and analogs such as Lispro, Levemir, glargin, et al.), insulin-like growth factor-I (IGF-I or Somatomedin-C), Nerve Growth Factor (NGF), Fibroblast Growth Factor (FGF; FGF-18, FGF-20, FGF-21, and the like), Keratinocyte Growth Factor (KGF) and Vascular Endothelial Growth Factor (VEGF), and the like. Especially attractive targets are those peptides which control gut function and appetite, but which have short duration of action (some of which are mentioned above), including but not limited to, ghrelin, Pancreatic Peptide, Peptide YY, Neuropeptide Y, Cholecystokinin (Sincalide, et al.), Melanocortin, and the like. Additional targets benefiting from the methods described herein are the proinflammatory adipose tissue products relating to obesity, including Leptin (and related analogs such as OB-3 peptide), adipokines, adiponectin, Chemerin, visfatin, nesfatin, resistin, tumor necrosis factor alpha, chemokines, monocyte chemotactic protein-1 (MCP-1), omentin, interleukins, and the like. Important targets that would benefit from improved pharmaceutical and pharmacodynamic behavior are proteins that control immune function, among which are given the following examples, which are not meant to be limiting, merely illustrative: members of the interferon family (interferons-alpha, -beta, -gamma, -kappa, -omega, the IL-10 cytokine family, including IL-10, IL-19, IL-20, IL-22, IL-24, IL-26, and the like), Thymopentin, Thymosin alpha1, and the like. Important peptide products which control the circulatory, or blood clotting would be improved by the methods described herein whereby increased duration of action would be achieved, for example, bivalirudin (Angiomax), eptifibatide (Integrelin), atrial natriuretic peptides (ANP, Ularitide), brain natriuretic peptide, c type natriuretic peptide, b-type natriuretic peptide (nesiritide), angiotensin, Angiostatin, Rotigaptide, thrombospondins and the like. Peptides that stimulate stem cell proliferation and differentiation are important agents that could be improved by the modifications described herein. For example, erythropoietin, hematide, thrombopoietin, macrophage colony-stimulating factor (M-CSF), leukemia inhibitory factor (LIF), interleukin-6 (IL-6), granulocyte colony-stimulating factor (G-CSF), granulocyte-macrophage-colony-stimulating factor (GM-CSF), and the like are contemplated as peptides that are suitable for covalent attachment to a surfactant (e.g., an alkyl glycoside surfactant). Neurotrophic factors are another class of small proteins that would greatly benefit from modifications described herein to increase duration of action and efficacy. For example, Glial cell-derived neurotrophic factor (GDNF) and family (neurturin, artemin, persephin), neurotrophins such as nerve growth factor (NGF), BDNF and the like. Proinflammatory and pain-causing peptides are important peptide targets that would be improved by modifications described herein. For example, inhibitors of bradykinin (or its release, e.g. ecallantide) and substance P especially antagonists,

offer important therapeutic targets (icatibant, et al.) are suitable for covalent attachment of surfactants (e.g. alkyl-glycoside surfactants). Inhibitors of viral fusion (Fuzeon), protein maturation (protease inhibitors) or integration suffer from short duration of action. Modification of such inhibitors via covalent attachment of a surfactant (e.g., an alkyl-glycoside surfactant) will allow for longer duration of action. Many proteases have been found to be involved in disease and inhibitors of their action have reached the clinic (Abbenante, G. and Fairlie, D.P. (2005) *Med Chem* 1: 71-104), but would be further improved by the modifications described herein and such targets are also contemplated within the scope of the embodiments described herein. Additional natural peptide products and analogs thereof such as conotoxin peptides for pain, antimicrobial defensins and the like also suffer from lack of bioavailability and short duration of action in physiological fluid and therefore would benefit from the peptide modifications described herein. Those skilled in the art will recognize many additional commercially important peptides that are amenable to modifications described herein and provide increased duration of action and bioavailability, and such peptides are also contemplated within the scope of the present disclosure.

[0404] Modifications at the amino or carboxyl terminus may optionally be introduced into the present peptides (Nestor, J.J., Jr. (2009) *Current Medicinal Chemistry* 16: 4399 - 4418). For example, the present peptides can be truncated or acylated on the N-terminus to yield peptides exhibiting low efficacy, partial agonist and antagonist activity, as has been seen for some peptides (Gourlet, P., et al. (1998) *Eur J Pharmacol* 354: 105-111, Gozes, I. and Furman, S. (2003) *Curr Pharm Des* 9: 483-494). Other modifications to the N-terminus of peptides, such as deletions or incorporation of D-amino acids such as D-Phe also can give potent and long acting agonists or antagonists when substituted with the modifications described herein such as long chain alkyl glycosides. Such agonists and antagonists also have commercial utility and are within the scope of contemplated embodiments described herein.

[0405] Aside from the twenty standard amino acids, there are a vast number of "nonstandard amino acids" or unnatural amino acids that are known to the art and that may be incorporated in the compounds described herein, as described above. Other nonstandard amino acids are modified with reactive side chains for conjugation (Gauthier, M.A. and Klok, H.A. (2008) *Chem Commun (Camb)* 2591-2611; de Graaf, A.J., et al. (2009) *Bioconjug Chem* 20: 1281-1295). In one approach, an evolved tRNA/ tRNA synthetase pair and is coded in the expression plasmid by the amber suppressor codon (Deiters, A, et al. (2004). *Bio-org. Med. Chem. Lett.* 14, 5743-5). For example, p-azidophenylalanine was incorporated into peptides and then reacted with a functionalized surfactant, or a PEG polymer having an acetylene moiety in the presence of a reducing agent and copper ions to facilitate an organic reaction known as "Huisgen [3+2] cycloaddition." A similar reaction sequence using the reagents described herein containing an acetylene modified alkyl glycoside or PEG modified glycoside will result in PEGylated or alkyl glycoside modified peptides. For peptides of less than about 50 residues, standard solid phase synthesis is used for incorporation of said reactive amino acid residues at the desired position in the chain. Such surfactant-modified peptides and/or proteins offer a different spectrum of pharmacological and medicinal properties than peptides modified by PEG incorporation alone.

Intermediates

[0406] In one embodiment provided herein are intermediates and/or reagents comprising a surfactant moiety and a reactive functional group capable of forming a bond with a reactive functional group on a natural or unnatural amino acid. These intermediates and/or reagents allow for improvement in the bioavailability and pharmaceutical, pharmacokinetic and/or pharmacodynamic behavior of peptides and/or proteins of use in human and animal disease. Covalent attachment of such intermediates and/or reagents via a functional group on a side chain of an amino acid, for example on an epsilon-amino function of Lys, the sulfhydryl of Cys, or at the amino or carboxy terminus of the peptide and/or protein target allows for synthesis of the peptide products described herein. In specific embodiments, non-ionic surfactant moieties are mono or disaccharides with an O-alkyl glycosidic substitution, said glycosidic linkage being of the alpha or beta configuration. In specific embodiments, O-alkyl chains are from C₁-C₂₀ or from C₆-C₁₆ alkyl chains.

[0407] In another embodiment provided herein are intermediates and/or reagents comprising a non-ionic surfactant moiety with certain alkyl glycosidic linkage that mimic O-alkyl glycosidic linkages and a reactive functional group capable of forming a bond with a reactive functional group on a natural or unnatural amino acid. Such intermediates and/or reagents contain S-linked alkyl chains or N-linked alkyl chains and have altered chemical and/or enzymatic stability compared to the O-linked alkyl glycoside-linked products.

[0408] In some embodiments, an intermediate and/or reagent provided herein is a compound wherein the hydrophilic group is a modified glucose, galactose, maltose, glucuronic acid, diglucuronic acid, maltouronic acid or the like. In some embodiments, the hydrophilic group is glucose, maltose, glucuronic acid, diglucuronic acid or maltouronic acid and the hydrophobic group is a C₁-C₂₀ alkyl chain or an aralkyl chain. In some embodiments the glycosidic linkage to the hydrophobic group is of an alpha configuration and in some the linkage is beta at the anomeric center on the saccharide.

[0409] In some embodiments, the hydrophilic group is glucose, maltose, glucuronic acid, diglucuronic acid or maltouronic acid and the hydrophobic group is a C₁-C₂₀ alkyl or aralkyl chain.

[0410] In some embodiments, an intermediate and/or reagent provided herein comprises a surfactant containing a reactive functional group that is a carboxylic acid group, an amino group, an azide, an aldehyde, a maleimide, a sulfhydryl, a hydroxylamino group, an alkyne or the like.

[0411] In some embodiments, the intermediate and/or reagent is an O-linked alkyl glycoside with one of the hydroxyl functions modified to be a carboxylic acid or amino functional group. In some embodiments, the reagent is a 1-O-alkyl glucuronic acid of alpha or beta configuration


and the alkyl chain is from C₁ to C₂₀ in length. In some of such embodiments, the alkyl group is from C₆ to C₁₈ in length. In some of such embodiments, the alkyl group is from C₆ to C₁₆ in length.

[0412] In some embodiments, the reagent comprises a 1-O-alkyl diglucuronic acid of alpha or beta configuration and the alkyl chain is from C₁ to C₂₀ in length. In some of such embodiments, the alkyl group is from C₆ to C₁₆ in length.


[0413] In some embodiments, the reagent is an S-linked alkyl glycoside of alpha or beta configuration with one of the hydroxyl functions modified to be a carboxylic acid or amino functional group.

[0414] In some embodiments, the reagent is an N-linked alkyl glycoside of alpha or beta configuration with one of the hydroxyl functions modified to be a carboxylic acid or amino functional group.

[0415] In yet another embodiment provided herein are peptide and/or protein products containing a covalently linked alkyl glycoside with properties acceptable for use in human and animal disease. Scheme 1 lists exemplary non-ionic surfactants that can be modified to yield the reagents and/or intermediates that are useful for synthesis of surfactant-modified peptide products described herein.





1-Octyl beta-D-melibioside



Scheme 1. Examples of commercially-available non-ionic surfactants of the alkyl glycoside class


[0416] In some embodiments, the covalently modified peptides and/or proteins described herein incorporate a surfactant moiety into the peptide structure. In specific embodiments, the covalently modified peptides and/or proteins described herein incorporate a non-ionic surfactant of the alkyl, alkoxyaryl, or aralkyl glycoside class. Alkyl glycosides, for example in the mono and disaccharide classes, are important commodities and are widely used in the food, service and cleaning industries. Thus their production on commercially significant scale has been the subject of extensive study. Both enzymatic and chemical processes are available for their production at very low cost (Park, D.W., et al. (2000) *Biotechnology Letters* 22: 951-956). These alkyl glycosides can be modified further to generate the intermediates for the synthesis of the covalently modified peptides and/or proteins described herein. Thus it is known that 1-dodecyl beta-D-glucoside is preferentially oxidized on the 6-position to yield the corresponding glucuronic acid analog in high yield when using the unprotected material and platinum black catalyst in the presence of oxygen (van Bekkum, H. (1990) *Carbohydrates as Organic Raw Materials* 289-310). Additional chemoselective methods for oxidation of the primary alcohol at the 6 position of alkyl glucosides are available. For example, use of catalytic amounts of 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) with stoichiometric amounts of the organic oxidant [bis(acetoxy)iodo]benzene (BAIB) (De Mico, A., et al. (1997) *J Org Chem* 1997: 6974-6977) gave outstanding yields of nucleoside-5'-carboxylic acids (Epp, J.B. and Widlanski, T.S. (1999) *J Org Chem* 64: 293-295) by oxidation of the primary hydroxyl. This oxidation is chemoselective for the primary hydroxyl even when the other, secondary hydroxyls are unprotected (Codee, J.D., et al. (2005) *J Am Chem Soc* 127: 3767-3773). In a similar manner, 1-dodecyl β -D-glucopyranoside and 1-tetradecyl β -D-glucopyranoside were oxidized to the corresponding uronic acids (1-dodecyl β -D-glucuronic acid, 1-tetradecyl β -D-glucuronic acid) by oxidation with TEMPO using KBr and sodium hypochlorite as stoichiometric oxidant (Milkereit, G., et al. (2004) *Chem Phys Lipids* 127: 47-63) in water. A mild oxidation procedure using (diacetoxyiodo)benzene (DAIB aka BAIB) is given in the Examples. Certain of these glucuronic acid intermediates are commercially available (for example octyl β -D-glucuronic acid; Carbosynth, MO 07928) and, as indicated, a broad range are subject to preparation by routine methods (Schamann, M. and Schafer, H.J. (2003) *Eur J Org Chem* 351-358; Van den Bos, L.J., et al. (2007) *Eur J Org Chem* 3963-3976) or upon request. Scheme 2 illustrates, as examples, certain functionalized surfactant intermediates comprising a -COOH group as a reactive functional group that are used to prepare the intermediates and/or reagents described herein.






Scheme 2. Examples of alkyl melibiouronic, diglucuronic and glucuronic acid class reagents.


[0417] Similarly, aralkyl glycosides (including alkoxyaryl) can form the basis for closely related nonionic surfactant reagents. For example, 4-alkoxyphenyl β -D-glucopyranosides are readily synthesized by the reaction of 4-alkoxyphenols with penta-O-acetyl β -D-glucose in the presence of boron trifluoride etherate. Subsequent deacetylation using trimethylamine in methanol/water and selective oxidation as described above and in the examples, yields the alkoxyaryl glucuronic acid reagents suitable for forming the reagents and peptides described herein ((Smits, E., et al. (1996) *J Chem Soc, Perkin Trans I* 2873-2877; Smits, E., et al. (1997) *Liquid Crystals* 23: 481-488).



Scheme 3. Illustrative members of aralkyl or alkoxyaryl surfactant moiety.

[0418] The glucuronic acid class of intermediate is readily activated by standard coupling agents for linkage to an amino acid side chain, e.g. that of Lys. Thus Fmoc-Lys-O-Tms (trimethylsilyl=TMS) can be reacted with octyl β -D-glucuronic acid in the presence of a coupling agent and the O-Tms protecting group can then be hydrolyzed on aqueous workup to yield Fmoc-Lys(1-octyl β -D-glucuronamide) as shown in Scheme 4. This reagent can be used for incorporation into the solid phase synthesis of peptides, using standard coupling protocols, when it is desired to incorporate the surfactant moiety near the N-terminal region of the molecule. The secondary hydroxyl groups can be left unprotected, due to the very much higher reactivity of the Lys amino functional group or they can be protected by peracetylation. If an acetyl protected form is used, the acetyl protecting groups can be removed in high yield by treatment with either MeOH/NaOMe or by MeOH/Et₃N. Scheme 4 illustrates preparation of the reagents described herein.






Scheme 4. Example of a preparation of a reagent

[0419] In some embodiments, reagents and/or intermediates for the preparation of the biologically active peptide products described herein comprise a family of surfactant-modified linker amino acids for incorporation into synthetic peptide products. Thus in one embodiment, peptide products described herein are synthesized in a linear fashion wherein a functionalized surfactant is attached to a reversibly-protected linker amino acid via a functional group on a side chain of a linker amino acid (e.g., an amino group of a lysine residue) to yield a proprietary reagent (as shown in Scheme 4.) which can be incorporated into the growing peptide chain and then the remaining peptide is synthesized by attachment of further amino acids to the cysteine residue. Protecting group suitable for synthesis of modified peptides and/or protein described herein are described in, for example, T. W. Green, P. G. M. Wuts, *Protective Groups in Organic Synthesis*, Wiley-Interscience, New York, 1999, 503-507, 736-739.

[0420] In another embodiment, peptide products described herein are synthesized by covalent attachment of a functionalized surfactant to a full-length peptide via a suitable functional group on a linker amino acid that is in the peptide chain.

[0421] Alternatively a functionalized surfactant can be added to a linker amino acid side chain which has been deprotected during the course of the solid phase synthesis of the peptide. As an example, an alkyl glucuronide group can be added directly to a linker amino acid side chain (e.g., a deprotected Lys side chain) during the solid phase synthesis of the peptide. For example, use of Fmoc-Lys(Alloc)-OH as a subunit provides orthogonal protection that can be removed while the peptide is still on the resin. Thus deprotection of the Lys side chain using Pd/thiobarbital, Pd/1,3-dimethyl barbituric acid (DMBA) or other alloc deprotection recipe allows exposure of the amino group for coupling with the acyl protected or unprotected 1-octyl beta-

D-glucuronic acid unit or for side chain lactam formation. Final deprotection with a low % $\text{CF}_3\text{CO}_2\text{H}$ (TFA) cleavage cocktail will then deliver the desired product. Although the glycosidic linkage is labile to strong acid, the experience here and by others is that it is relatively stable to low % TFA cleavage conditions. Alternatively, acyl protection (e.g. acetyl, Ac; benzoyl, Bz) or trialkylsilyl protection on the saccharide OH functional groups may be used to provide increased protection to the glycosidic linkage. Subsequent deprotection by base ($\text{NH}_2\text{NH}_2/\text{MeOH}$; NH_3/MeOH , NaOMe/MeOH) yields the desired deprotected product. Scheme 4 illustrates reagents described herein. Scheme 5 illustrates peptide intermediates described herein.



Scheme 5. Illustrative example of a peptide intermediate.


[0422] Additional reagents are generated by modification of the 6-position functional group to give varied means of linkage to amino acid side chain functional groups, as shown below in Scheme 6. Thus amino substitution can be used for linkage to Asp or Glu side chains. Azido or alkyne substitution can be used for linkage to unnatural amino acids containing the complementary acceptor for Huisgen 3+2 cycloaddition (Gauthier, M.A. and Klok, H.A. (2008) *Chem Commun (Camb)* 2591-2611). Aminoxy or aldehyde functional groups can be used to link to aldehyde (i.e. oxime linkage) or to amino functions (i.e. reductive alkylation), respectively. The maleimide or $-\text{NH}-(\text{C}=\text{O})-\text{CH}_2-\text{Br}$ functional group can bind chemo selectively with a Cys or other SH functional group. These types of linkage strategies are advantageous when used in conjunction with the reagents described herein. Interconversion of functional groups is widely practiced in organic synthesis and comprehensive lists of multiple routes to each of the functional group modifications listed herein are available (Larock, R.C. (1999) "Comprehensive Organic Transformations", VCH Publishers, New York).

[0423] Thus, for example, the primary hydroxyl on position 6 of octyl 1- β -D-glucoside is converted to the azide by activation and displacement with an azide anion, reactions such as reactions used in carbohydrate chemistry (e.g. by tosylation followed by NaN_3). The corresponding azide is reduced to the amino function by reduction with thiolacetic acid in pyridine (Elofsson, M., et al. (1997) *Tetrahedron* 53: 369-390) or by similar methods of amino group generation (Stangier, P., et al. (1994) *Liquid Crystals* 17: 589-595). Approaches to the

acetylene, aminoxy, and aldehyde moieties are best carried out on the triacetoxy form, available from the commercially available glucoside by treatment with AC_2O , followed by mild hydrolysis of the primary amine. This 6-hydroxy form can be selectively oxidized to the aldehyde, or activated as a tosylate or triflate and displaced by NH_2OH or by sodium acetylide. The maleimide linkage can be through a carbon linkage as shown or, preferably though an O or amide linkage, again by displacement of the activated hydroxyl or coupling of the glucuronic acid derivative to an amino linked maleimide reagent, well known in the art. Additional functional group interconversions are well known to those of average skill in the art of medicinal chemistry and are within the scope of the embodiments described herein.

[0424] Also contemplated within the scope of synthetic methods described herein are surfactants wherein the saccharide and hydrophobic chain are covalently attached via an alpha glycosidic linkage. Synthetic routes to predominantly α -linked glycosides are well known in the art and typically originate with the peracetyl sugar and use acidic catalysis (e.g. SnCl_4 , BF_3 or HCl) to effect the α -glycosylation (Cudic, M. and Burstein, G.D. (2008) *Methods Mol Biol* 494: 187-208, Vill, V., et al. (2000) *Chem Phys Lipids* 104: 75-91,). Similar synthetic routes exist for disaccharide glycosides (von Minden, H.M., et al. (2000) *Chem Phys Lipids* 106: 157-179,). Functional group interconversions then proceed as above to lead to the 6-carboxylic acid, et al. for generation of the corresponding α -linked reagents.

[0425] Scheme 6 lists certain compounds and reagents useful in the synthesis of the covalently modified peptides and/or proteins described herein. Standard nomenclature using single letter abbreviations for amino acids are used.




Scheme 6. Additional reagent examples.

[0426] Many alkyl glycosides can be synthesized by known procedures, as described, e.g., in (Rosevear, P., et al. (1980) *Biochemistry* 19: 4108-4115, Li, Y.T., et al. (1991) *J Biol Chem* 266: 10723-10726) or Koeltzow and Urfer, J. *Am. Oil Chem. Soc.*, 61:1651-1655 (1984), U.S. Pat. No. 3,219,656 and U.S. Pat. No. 3,839,318 or enzymatically, as described, e.g., in (Li, Y.T., et al. (1991) *J Biol Chem* 266: 10723-10726, Gopalan, V., et al. (1992) *J Biol Chem* 267: 9629-9638). O-alkyl linkages to natural amino acids such as Ser can be carried out on the Fmoc-Ser-OH using peracetylglucose to yield $\text{N}\alpha$ -Fmoc-4-O-(2,3,4,6-tetra-O-acetyl- β -D-

glucopyranosyl)-L-serine. This material is selectively deprotected at the primary carbon atom (position 6) and selectively oxidized using TEMPO/BAIB as described above to yield the corresponding 6-carboxyl function which may be coupled to lipophilic amines to generate a new class of nonionic surfactant and reagents (Scheme 7).

Scheme 7. Alternative example of nonionic surfactant reagent.



[0427] The linkage between the hydrophobic alkyl and the hydrophilic saccharide can include, among other possibilities, a glycosidic, thioglycosidic, amide (Carbohydrates as Organic Raw Materials, F. W. Lichtenthaler ed., VCH Publishers, New York, 1991), ureide (Austrian Pat. 386,414 (1988); Chem. Abstr. 110:137536p (1989); see Gruber, H. and Greber, G., "Reactive Sucrose Derivatives" in Carbohydrates as Organic Raw Materials, pp. 95-116) or ester linkage (Sugar Esters: Preparation and Application, J. C. Colbert ed., (Noyes Data Corp., New Jersey), (1974)).

[0428] Examples from which useful alkyl glycosides can be chosen for modification to the reagents or for the formulation of the products described herein, include: alkyl glycosides, such as octyl-, nonyl-, decyl-, undecyl-, dodecyl-, tridecyl-, tetradecyl-, pentadecyl-, hexadecyl-, heptadecyl-, and octadecyl-D-maltoside, -glucoside or -sucroside (i.e., sucrose ester) (synthesized according to Koeltzow and Urfer; Anatrache Inc., Maumee, Ohio; Calbiochem, San Diego, Calif.; Fluka Chemie, Switzerland); alkyl thiomaltosides, such as heptyl, octyl, dodecyl-, tridecyl-, and tetradecyl- β -D-thiomaltoside (synthesized according to Defaye, J. and Pederson, C., "Hydrogen Fluoride, Solvent and Reagent for Carbohydrate Conversion Technology" in Carbohydrates as Organic Raw Materials, 247-265 (F. W. Lichtenthaler, ed.) VCH Publishers, New York (1991); Ferenci, T., J. Bacteriol, 144:7-11 (1980)); alkyl thioglucosides, such as 1-dodecyl- or 1-octyl-thio α - or β -D-glucopyranoside (Anatrache, Inc., Maumee, Ohio; see Saito, S. and Tsuchiya, T. Chem. Pharm. Bull. 33:503-508 (1985)); alkyl thiosucroses (synthesized according to, for example, Binder, T. P. and Robyt, J. F., Carbohydr. Res. 140:9-20 (1985)); alkyl maltotriosides (synthesized according to Koeltzow and Urfer); long chain aliphatic carbonic acid amides of sucrose amino-alkyl ethers; (synthesized according to Austrian Patent 382,381 (1987); Chem. Abstr., 108:114719 (1988) and Gruber and Greber pp. 95-116); derivatives of palatinose and isomaltamine linked by amide linkage to an alkyl chain (synthesized according to Kunz, M., "Sucrose-based Hydrophilic Building Blocks as Intermediates for the Synthesis of Surfactants and Polymers" in Carbohydrates as Organic Raw Materials, 127-153); derivatives of isomaltamine linked by urea to an alkyl chain (synthesized according to Kunz); long chain aliphatic carbonic acid ureides of sucrose amino-

alkyl ethers (synthesized according to Gruber and Greber, pp. 95-116); and long chain aliphatic carbonic acid amides of sucrose amino-alkyl ethers (synthesized according to Austrian Patent 382,381 (1987), Chem. Abstr., 108:114719 (1988) and Gruber and Greber, pp. 95-116).

[0429] Some preferred glycosides include the saccharides maltose, sucrose, glucose and galactose linked by glycosidic or ester linkage to an alkyl chain of 6, 8, 10, 12, or 14 carbon atoms, e.g., hexyl-, octyl-, decyl-, dodecyl- and tetradecyl- maltoside, sucroside, glucoside and galactoside. In the body these glycosides are degraded to non-toxic alcohol or fatty acid and an oligosaccharide or saccharide. The above examples are illustrative of the types of alkyl glycosides to be used in the methods claimed herein, however the list is not intended to be exhaustive.

[0430] Generally, these surfactants (e.g., alkyl glycosides) are optionally designed or selected to modify the biological properties of the peptide, such as to modulate bioavailability, half-life, receptor selectivity, toxicity, biodistribution, solubility, stability, e.g. thermal, hydrolytic, oxidative, resistance to enzymatic degradation, and the like, facility for purification and processing, structural properties, spectroscopic properties, chemical and/or photochemical properties, catalytic activity, redox potential, ability to react with other molecules, e.g., covalently or noncovalently, and the like.

Surfactants

[0431] The term "surfactant" comes from shortening the phrase "surface active agent". In pharmaceutical applications, surfactants are useful in liquid pharmaceutical formulations in which they serve a number of purposes, acting as emulsifiers, solubilizers, and wetting agents. Emulsifiers stabilize the aqueous solutions of lipophilic or partially lipophilic substances. Solubilizers increase the solubility of components of pharmaceutical compositions increasing the concentration which can be achieved. A wetting agent is a chemical additive which reduces the surface tension of a fluid, inducing it to spread readily on a surface to which it is applied, thus causing even "wetting" of the surface with the fluids. Wetting agents provide a means for the liquid formulation to achieve intimate contact with the mucous membrane or other surface areas with which the pharmaceutical formulation comes in contact. Thus surfactants may be useful additives for stabilization of the formulation of the peptide products described herein as well as for the modification of the properties of the peptide itself.

[0432] In specific embodiments, alkyl glycosides which are synthetically accessible, e.g., the alkyl glycosides dodecyl, tridecyl and tetradecyl maltoside or glucoside as well as sucrose dodecanoate, tridecanoate, and tetradecanoate are suitable for covalent attachment to peptides as described herein. Similarly, the corresponding alkylthioglycosides are stable, synthetically accessible surfactants which are acceptable for formulation development.

[0433] A wide range of physical and surfactant properties can be achieved by appropriate modification of the hydrophobic or hydrophilic regions of the surfactant (e.g., the alkyl glycoside). For example, a study comparing the bilayer activity of dodecyl maltoside (DM) with that of dodecyl glucoside (DG) found that of DM to be more than three times higher than that of DG, despite having the same length of hydrophobic tail (Lopez, O., et al. (2002) *Colloid Polym Sci* 280: 352-357). In this particular instance the identity of the polar region (disaccharide vs monosaccharide) influences surfactant behavior. In the case of a surfactant linked to a peptide, e.g. the peptide products described herein, the peptide region also may contribute hydrophobic or hydrophilic character to the overall molecule. Thus tuning of the physical and surfactant properties may be used to achieve the particular physical and pharmaceutical properties suitable for the individual peptide targets.

PEG modification

[0434] In some embodiments, surfactant-modified peptide products described herein are further modified to incorporate one or more PEG moieties (Veronese, F.M. and Mero, A. (2008) *BioDrugs* 22: 315-329). In some instances, incorporation of large PEG chains prevents filtration of the peptide in the glomeruli in the kidney into the dilute urine forming there (Nestor, J.J., Jr. (2009) *Current Medicinal Chemistry* 16: 4399 - 4418, Caliceti, P. and Veronese, F.M. (2003) *Adv Drug Deliv Rev* 55: 1261-1277). In some embodiments, an optional PEG hydrophilic chain allows for balancing the solubility and physical properties of the peptides or proteins that have been rendered hydrophobic by the incorporation of the longer chain alkyl glycoside moiety.

[0435] PEGylation of a protein can have potentially negative effects as well. It is widely recognized that PEGylation and other polymeric modifications can cause a substantial loss of biological activity for many proteins and this may relate to ligands for specific classes of receptors. In such instances there may be a benefit to reversible PEGylation (Peleg-Shulman, T., et al. (2004) *J Med Chem* 47: 4897-4904, Greenwald, R.B., et al. (2003) *Adv Drug Deliv Rev* 55: 217-250, Roberts, M.J. and Harris, J.M. (1998) *J Pharm Sci* 87: 1440-1445).

[0436] In addition, the increased molecular mass may prevent penetration of physiological barriers other than the glomerular membrane barrier. For example, it has been suggested that high molecular weight forms of PEGylation may prevent penetration to some tissues and thereby reduce therapeutic efficacy. In addition, high molecular weight may prevent uptake across mucosal membrane barriers (nasal, buccal, vaginal, oral, rectal, lung delivery). However delayed uptake may be highly advantageous for administration of stable molecules to the lung, substantially prolonging the duration of action. The peptide and/or protein products described herein have increased transmucosal bioavailability and this will allow longer chain PEG modifications to be used in conjunction with the surfactant modification with the achievement of commercially significant bioavailability following intranasal or other transmucosal route.

[0437] In some embodiments, long chain PEG polymers, and short chain PEG polymers are

suitable for modification of the proteins and peptides described herein. Administration of treatments for diabetes by inhalation is a new approach for drug delivery and the lung has a highly permeable barrier (e.g. Exubera). For this application, delayed penetration of the lung barrier, preferred forms of PEGylation are in the lower molecular weight range of C₁₀ to C₄₀₀ (roughly 250 to 10,000Da). Thus while a primary route to prolongation by PEG is the achievement of an "effective molecular weight" above the glomerular filtration cut-off (greater than 68kDa), use of shorter chains may be a route for prolongation of residence in the lung for treatment of lung diseases and other respiratory conditions. Thus PEG chains of about 500 to 3000 Da are of sufficient size to slow the entry into the peripheral circulation, but insufficient to cause them to have a very prolonged circulation time. In some embodiments, PEGylation is applied to give increased local efficacy to the lung tissue with reduced potential for systemic side effects for the covalently modified peptides and/or proteins described herein. In some of such embodiments, PEG chains in the range from about 750 to about 1500 Da are referred collectively as "PEG1K."

[0438] In addition, other polymers may be used in conjunction with the compounds of described herein in order to optimize their physical properties. For example poly(2-ethyl 2-oxazoline) conjugates have variable hydrophobicity and sufficient size to enhance duration of action (Mero, A., et al. (2008) J Control Release 125: 87-95). Linkage of such a polymer to a saccharide yields a class of surfactant suitable for use in modification of peptides and/or proteins described herein.

[0439] Polyethylene glycol chains are functionalized to allow their conjugation to reactive groups on the peptide and/or protein chain. Typical functional groups allow reaction with amino, carboxyl or sulfhydryl groups on the peptide through the corresponding carboxyl, amino or maleimido groups (and the like) on the polyethylene glycol chain. In an embodiment, PEG comprises a C₁₀-C₃₀₀₀ chain. In another embodiment, PEG has a molecular weight above 40,000 Daltons. In yet another embodiment, PEG has a molecular weight below 10,000 Daltons. PEG as a protein modification is well known in the art and its use is described, for example, in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192; and 4,179,337.

[0440] A non-traditional type of PEG chain is modified to be amphiphilic in nature. That is it has both the hydrophilic PEG structure but is modified to contain hydrophobic regions such as fatty acid esters and other hydrophobic components. See for example (Miller, M.A., et al. (2006) Bioconjug Chem 17: 267-274) ; Ekwuribe, et al. US 6,309,633; Ekwuribe, et al. US 6,815,530; Ekwuribe, et al. US 6,835,802). Although these amphiphilic PEG conjugates to proteins were originally developed to increase oral bioavailability they were relatively ineffective in this role. However the use of such amphiphilic PEG conjugates with amphipathic peptides will give significantly prolonged residence in the lung to extend the useful biological activity of these pharmaceuticals. The preferred PEG chains are in the molecular weight range of 500 to 3000Da. Detailed descriptions of the methods of synthesis of these conjugates is given in the references above.

[0441] A PEG entity itself does not have a functional group to be attached to a target molecular, such as a peptide. Therefore, to create PEG attachment, a PEG entity must be functionalized first, then a functionalized attachment is used to attach the PEG entity to a target molecule, such as a peptide (Greenwald, R.B., et al. (2003) *Adv Drug Deliv Rev* 55: 217-250, Veronese, F.M. and Pasut, G. (2005) *Drug Discov Today* 10: 1451-1458, Roberts, M.J., et al. (2002) *Adv Drug Deliv Rev* 54: 459-476). In one embodiment, site-specific PEGylation can be achieved through Cys substitution on a peptide molecule. The target peptide can be synthesized by solid phase synthesis, recombinant means, or other means, as described herein.

[0442] Thus in some embodiments, a peptide product described herein comprises a Lys or other reactive residue modified with an alkyl glycoside and specific PEGylation on at least one Cys residue, a Lys residue or other reactive amino acid residue elsewhere in the molecule.

[0443] In another embodiment, a Lys or other residue with a nucleophilic side chain may be used for incorporation of the PEG residue. This may be accomplished through the use of an amide or carbamate linkage to a PEG-carboxyl or PEG-carbonate chain. See for example as described (Veronese, F.M. and Pasut, G. (2005) *Drug Discov Today* 10: 1451-1458). An alternative approach is to modify the Lys side chain amino function through attachment of an SH containing residue, such as mercaptoacetyl, mercaptopropionyl (CO-CH₂-CH₂-CH₂-SH), and the like. Alternatively the PEG chain may be incorporated at the C-Terminus as an amide during the course of the synthesis. Additional methods for attaching PEG chains utilize reaction with the side chains of His and Trp. Other similar methods of modifying the peptide chain to allow attachment of a PEG chain are known in the art (Roberts, M.J., et al. (2002) *Adv Drug Deliv Rev* 54: 459-476).

Formulations

[0444] In one embodiment, the covalently modified peptides or proteins as disclosed herein are provided in a formulation that further reduces, prevents, or lessens peptide and/or protein association or aggregation in the composition, for example, reduces peptide and/or protein self-association or self-aggregation, or reduces association or aggregation with other peptides or proteins when administered to the subject.

[0445] Self-Association at high protein concentration is problematic in therapeutic formulations. For example, self-association increases the viscosity of a concentrated monoclonal antibody in aqueous solution. Concentrated insulin preparations are inactivated by self aggregation. These self associating protein interactions, particularly at high protein concentration, reduce, modulate or obliterate biological activity of many therapeutics (Clodfelter, D.K., et al. (1998) *Pharm Res* 15: 254-262). Therapeutic proteins formulated at high concentrations for delivery by injection or other means can be physically unstable or become insoluble as a result of these protein interactions.

[0446] A significant challenge in the preparation of peptide and protein formulations is to develop manufacturable and stable dosage forms. Physical stability properties, critical for processing and handling, are often poorly characterized and difficult to predict. A variety of physical instability phenomena are encountered such as association, aggregation, crystallization and precipitation, as determined by protein interaction and solubility properties. This results in significant manufacturing, stability, analytical, and delivery challenges. Development of formulations for peptide and protein drugs requiring high dosing (on the order of mg/kg) are required in many clinical situations. For example, using the SC route, approximately <1.5 mL is the allowable administration volume. This may require >100 mg/mL protein concentrations to achieve adequate dosing. Similar considerations exist in developing a high-concentration lyophilized formulation for monoclonal antibodies. In general, higher protein concentrations permit smaller injection volume to be used which is very important for patient comfort, convenience, and compliance. The surfactant-modified compounds described herein are designed to minimize such aggregation events and may be further facilitated through the use of small amounts of surfactants as herein described.

[0447] Because injection is an uncomfortable mode of administration for many people, other means of administering peptide therapeutics have been sought. Certain peptide and protein therapeutics may be administered, for example, by intranasal, buccal, oral, vaginal, inhalation, or other transmucosal administration. Examples are nafarelin (Synarel®) and calcitonin which are administered as commercial nasal spray formulations. The covalently modified peptides and/or proteins described herein are designed to facilitate such transmucosal administration and such formulations may be further facilitated through the use of small amounts of surfactants as described herein.

[0448] Typical formulation parameters include selection of optimum solution pH, buffer, and stabilizing excipients. Additionally, lyophilized cake reconstitution is important for lyophilized or powdered formulations. A further and significant problem comprises changes in viscosity of the protein formulation upon self-association. Changes in viscosity can significantly alter delivery properties e.g., in spray (aerosol) delivery for intranasal, pulmonary, or oral cavity sprays. Furthermore, increased viscosity can make injection delivery by syringe or iv line more difficult or impossible.

[0449] Many attempts to stabilize and maintain the integrity and physiological activity of peptides have been reported. Some attempts have produced stabilization against thermal denaturation and aggregation, particularly for insulin pump systems. Polymeric surfactants are described (Thurrow, H. and Geisen, K. (1984) *Diabetologia* 27: 212-218; Chawla, A.S., et al. (1985) *Diabetes* 34: 420-424). The stabilization of insulin by these compounds was believed to be of a steric nature. Among other systems used are saccharides (Arakawa, T. and Timasheff, S.N. (1982) *Biochemistry* 21: 6536-6544), osmolytes, such as amino acids (Arakawa, T. and Timasheff, S.N. (1985) *Biophys J* 47: 411-414), and water structure breakers, such as urea (Sato, S., et al. (1983) *J Pharm Sci* 72: 228-232). These compounds exert their action by modulating the intramolecular hydrophobic interaction of the protein or peptide.

[0450] Various peptides, peptides, or proteins are described herein and may be modified with any of the covalently bound surfactant reagents described herein. Advantageously, the peptide modifications described herein comprise covalent attachment of a surfactant that comprises both hydrophilic (e.g. saccharide) and hydrophobic (e.g. alkyl chain) groups, thereby allowing for stabilization of the peptide in physiological conditions. In some embodiments, covalent linkage of a moiety comprising a hydrophilic group and hydrophobic group (e.g. a glycoside surfactant) to a peptide, and/or protein described herein eliminates the need for modifying the amino acid sequence of the peptide, and/or protein to enhance stability (e.g., reduce aggregation).

[0451] In some embodiments, the formulations comprise at least one drug comprising a peptide modified with a surfactant derived reagent described herein and in formulation additionally may be associated with a surfactant, wherein the surfactant is further comprised of, for example, a saccharide, an alkyl glycoside, or other excipient and can be administered in a format selected from the group consisting of a drop, a spray, an aerosol, a lyophilizate, a spray dried product, an injectable, and a sustained release format. The spray and the aerosol can be achieved through use of the appropriate dispenser and may be administered by intranasal, transbuccal, inhalation or other transmucosal route. The lyophilizate may contain other compounds such as mannitol, saccharides, submicron anhydrous α -lactose, gelatin, biocompatible gels or polymers. The sustained release format can be an ocular insert, erodible microparticulates, hydrolysable polymers, swelling mucoadhesive particulates, pH sensitive microparticulates, nanoparticles/latex systems, ion-exchange resins and other polymeric gels and implants (Ocusert, Alza Corp., California; Joshi, A., S. Ping and K. J. Himmelstein, Patent Application WO 91/19481). Significant oral bioavailability is also achievable.

[0452] The peptide and protein modifications described herein mitigate and, in some cases, may eliminate the need for organic solvents. Trehalose, lactose, and mannitol and other saccharides have been used to prevent aggregation. Aggregation of an anti-IgE humanized monoclonal antibody was minimized by formulation with trehalose at or above a molar ratio in the range of 300:1 to 500:1 (excipient:protein). However, the powders were excessively cohesive and unsuitable for aerosol administration or exhibited unwanted protein glycation during storage (Andya, J.D., et al. (1999) Pharm Res 16: 350-358). Each of the additives discovered have limitations as additives to therapeutics including xenobiotic metabolism, irritation or toxicity, or high cost. Contemplated for use with the covalently modified peptides and/or proteins described herein are excipients that are effective, non-irritating and non-toxic, do not require xenobiotic metabolism since they are comprised of the natural sugars, fatty acids, or long chain alcohols, and which may also be used to minimize aggregation in aqueous solutions or upon aqueous reconstitution of dried peptide and/or protein formulations in situ by physiologic aqueous reconstitution by aqueous body fluids such as plasma or saliva.

[0453] Other formulation components could include buffers and physiological salts, non-toxic protease inhibitors such as aprotinin and soybean trypsin inhibitor, alpha-1-antitrypsin, and protease-inactivating monoclonal antibodies, among others. Buffers could include organics such as acetate, citrate, gluconate, fumarate, malate, polylysine, polyglutamate, chitosan,

dextran sulfate, etc. or inorganics such as phosphate, and sulfate. Such formulations may additionally contain small concentrations of bacteriostatic agents like benzyl alcohol, and the like.

[0454] Formulations suitable for intranasal administration also comprise solutions or suspensions of the modified peptides and/or protein products described herein in an acceptable evaporating solvents such as hydrofluoroalkanes. Such formulations are suitable for administration from metered dose inhalers (MDI) and have advantages of lack of movement from site of administration, low irritation and absence of need for sterilization. Such formulations may also contain acceptable excipients or bulking agents such as submicron anhydrous α -lactose.

[0455] In yet another aspect, the covalently modified peptides and/or proteins described herein exhibit increased shelf-life. As used herein, the phrase "shelf life" is broadly described as the length of time a product may be stored without becoming unsuitable for use or consumption. The "shelf life" of the composition described herein, can also indicate the length of time that corresponds to a tolerable loss in quality of the composition. The compositional shelf life as used herein is distinguished from an expiration date; "shelf life" relates to the quality of the composition described herein, whereas "expiration date" relates more to manufacturing and testing requirements of the composition. For example, a composition that has passed its "expiration date" may still be safe and effective, but optimal quality is no longer guaranteed by the manufacturer.

[0456] In some embodiments, the modified peptides described herein have greater bioavailability compared to the unmodified peptide. In some embodiments, the bioavailability is greater for the modified peptides compared to the unmodified peptide when administered by nasal routes. In some embodiments, the bioavailability is greater for the modified peptides compared to the unmodified peptide when administered by buccal. In some embodiments, the bioavailability is greater for the modified peptides compared to the unmodified peptide when administered by oral routes. In some embodiments, the bioavailability is greater for the modified peptides compared to the unmodified peptide when administered by vaginal routes. In some embodiments, the bioavailability is greater for the modified peptides compared to the unmodified peptide when administered by intravenous routes. In some embodiments, the bioavailability is greater for the modified peptides compared to the unmodified peptide when administered by subcutaneous routes. In some embodiments, the bioavailability of the modified peptides described herein have between 10-90%, 20-80%, 30-60%, 10-30%, 15-30%, 15-45%, 50-75 %, 60-75%, 70-90%, 80-100%, 100-1000%, 100-500%, 50-200%, or 100-200% greater bioavailability when compared with the unmodified peptide. In some embodiments, the bioavailability of the modified peptides described herein have at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 100%, at least 120%, at least 140%, at least 160%, at least 180%, at least 200%, at least 220%, at least 240%, at least 260%, at least 280%, at least

300%, at least 400%, at least 500%, at least 600%, at least 700%, at least 800%, at least 900%, at least 1000%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 60%, about 70%, about 80%, about 90%, about 100%, about 120%, about 140%, about 160%, about 180%, about 200%, about 220%, about 240%, about 260%, about 280%, about 300%, about 400%, about 500%, about 600%, about 700%, about 800%, about 900% or about 1000% greater bioavailability when compared with the unmodified peptide.

Methods

[0457] In one aspect, provided herein are methods of administering to a subject in need thereof an effective amount of the therapeutic compositions described herein. As used herein, "therapeutically effective amount" is interchangeable with "effective amount" for purposes herein, and is determined by such considerations as are known in the art. The amount must be effective to achieve a desired drug-mediated effect in the treated subjects suffering from the disease thereof. A therapeutically effective amount also includes, but is not limited to, appropriate measures selected by those skilled in the art, for example, improved survival rate, more rapid recovery, or amelioration, improvement or elimination of symptoms, or other acceptable biomarkers or surrogate markers.

[0458] The compositions described herein are delivered to a vertebrate subject in need of treatment including but not limited to, for example, a human. Moreover, depending on the condition being treated, these therapeutic compositions may be formulated and administered systemically or locally. Techniques for formulation and administration may be found in the latest edition of "Remington's Pharmaceutical Sciences" (Mack Publishing Co, Easton Pa.). Suitable routes may, for example, include oral or transmucosal administration; such as intranasal; buccal; ocular, vaginal; rectal; as well as parenteral delivery, including intramuscular, subcutaneous, intravenous, or intraperitoneal administration.

[0459] In some embodiments of the methods, the effective amount of the peptide product for administration is from about 0.1 $\mu\text{g/kg/day}$ to about 100.0 $\mu\text{g/kg/day}$, or from 0.01 $\mu\text{g/kg/day}$ to about 1 mg/kg/day or from 0.1 $\mu\text{g/kg/day}$ to about 50 mg/kg/day . In some embodiments, the peptide product is administered parenterally. In some embodiments, the peptide product is administered subcutaneously. In some embodiments, the method of administration of the peptide product is nasal insufflation.

[0460] It will be understood, however, that the specific dose level and frequency of dosage for any particular subject in need of treatment may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and duration of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

[0461] Described herein is a method for chemically modifying a molecule by covalent linkage to a surfactant to increase or sustain the biological action of the composition or molecule, for example, receptor binding or enzymatic activity. In some embodiments, the molecule is a peptide. The method additionally can include further modification comprising covalent attachment of the molecule in the composition to a polymer such as polyethylene glycol.

[0462] The method(s) includes all aspects of the compositions described herein including but not limited to compositions which reduce or eliminate immunogenicity of peptide and/or protein drugs, are non-irritating, have anti-bacterial or anti-fungal activity, have increased stability or bioavailability of a drug, decrease the bioavailability variance of that drug, avoid first pass liver clearance and reduce or eliminate any adverse effects. As used herein, the term "immunogenicity" is the ability of a particular substance or composition or agent to provoke an immunological response. The immunogenicity of the covalently modified peptides and/or proteins described herein is confirmed by methods known in the art.

[0463] Also provided is a method of administering a drug composition comprising a peptide covalently linked to at least one alkyl glycoside and delivered to a vertebrate, wherein the alkyl has from 1 to 30 carbon atoms, or further in the range of 6 to 16 carbon atoms, and the alkyl glycoside increases the stability, bioavailability and/or duration of action of the drug.

[0464] In another embodiment, provided is a method of reducing or eliminating immunogenicity of a peptide and/or protein drug by covalently linking the peptide chain to at least one alkyl glycoside wherein the alkyl has from 1 to 30 carbon atoms.

Methods of treatment

[0465] Described herein, in some embodiments are methods for treatment of pain, including post-operative or chronic pain, comprising administration of a surfactant-modified peptide and/or protein product described herein (e.g., a peptide product of Formula I, II or III) to individuals in need thereof. In some of such embodiments, the opioid analogs described herein are not addictive or habit forming, and/or are administered in lower dosages compared to current medications (e.g., codeine) and are longer lasting compared to current medications.

[0466] Provided herein, in some embodiments are methods for prevention and/or treatment of conditions associated with hypoparathyroidism and/or decreases in bone mass density comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) to individuals in need thereof. In some embodiments, the conditions characterized by decreases in bone mass density include, and are not limited to, osteoporosis, osteopenia, post-menopausal osteoporosis, Paget's disease, glucocorticoid induced osteoporosis, old age osteoporosis, humoral hypercalcemia, or the like.

[0467] In some embodiments, provided herein are methods for treatment of

hypoparathyroidism comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) to individuals in need thereof. In some embodiments, the hypoparathyroidism is associated with decrease in bone mass density.

[0468] Further provided herein are methods for stimulating bone repair and/or favoring engraftment of a bone implant comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) to individuals in need thereof.

[0469] In yet further embodiments, provided herein are methods for increasing bone density and/or reducing incidence of fractures (e.g., vertebrae fractures, hip fractures, or the like) comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) to individuals in need thereof.

[0470] In some embodiments, provided herein are methods for treatment of humoral hypercalcemia comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) to individuals in need thereof. In some embodiments, humoral hypercalcemia is associated with tumors. In some of such embodiments, the peptide product (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is an inverse agonist or an antagonist of PTH or PTHrP.

[0471] In some embodiments of the methods described above, the peptide and/or protein that is covalently attached to a surfactant is PTH, PTHrP, or an analog thereof. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered prophylactically and delays occurrence of any condition associated with loss of bone density, including and not limited to osteoporosis, osteopenia, post-menopausal osteoporosis, Paget's disease, glucocorticoid induced osteoporosis, old age osteoporosis, or the like. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered therapeutically and delays progression of any condition associated with loss of bone density, including and not limited to osteoporosis, osteopenia, post-menopausal osteoporosis, Paget's disease, glucocorticoid induced osteoporosis, humoral hypercalcemia, or the like. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered prophylactically and/or therapeutically and delays progression of osteopenia to osteoporosis. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered prophylactically and/or therapeutically and reduces or halts further loss of bone density, thereby stabilizing disease.

[0472] In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered parenterally. In some

embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered subcutaneously. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered by nasal insufflation.

[0473] In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) has a longer duration of action compared to a pharmaceutical comprising currently known therapeutics (e.g., recombinant PTH, bisphosphonates, antibody Denosumab, or the like). In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered for longer period of time (e.g., > two years) compared to a pharmaceutical comprising currently known therapeutics (e.g., recombinant PTH, bisphosphonates, antibody Denosumab, or the like) while reducing or ameliorating side-effects (e.g., osteonecrosis in jaw, skin infections or the like) associated with currently known therapeutics (e.g., recombinant PTH, bisphosphonates, antibody Denosumab, or the like). In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is an agonist of PTH or PTHrP. In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is an antagonist of PTH or PTHrP. In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is an inverse agonist of PTH or PTHrP.

[0474] Provided herein, in some embodiments are methods for prevention and/or treatment of conditions associated with decreases in insulin sensitivity comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) to individuals in need thereof. In some embodiments, the conditions characterized by decreases in insulin sensitivity include, and are not limited to, the metabolic syndrome, obesity-related insulin resistance, hypertension, systemic inflammation associated with high C reactive protein, diabetes, or the like.

[0475] Also provided is a method of treating conditions associated with insulin resistance including and not limited to obesity, the metabolic syndrome, type 2 diabetes, hypertension, atherosclerosis or the like, comprising administering a drug composition comprising a peptide covalently linked to at least one alkyl glycoside and delivered to a vertebrate, wherein the alkyl has from 1 to 30 carbon atoms, or further in the range of 6 to 18 carbon atoms (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V), and wherein covalent linkage of the alkyl glycoside to the peptide increases the stability, bioavailability and/or duration of action of the drug.

[0476] Also provided herein are methods for treatment of insulin resistance comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or

Formula 3-V) to individuals in need thereof. In some embodiments, the insulin resistance is associated with the metabolic syndrome (Syndrome X) and/or diabetes.

[0477] Further provided herein are methods for stimulating resensitization of the body to insulin comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g. a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) to individuals in need thereof.

[0478] In yet further embodiments, provided herein are methods for increasing insulin sensitivity through weight loss, comprising administration of a therapeutically effective amount of a surfactant-modified peptide and/or protein product described herein (e.g. a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V and in Table 1 of Figure 1 and Table 2 of Figure 2) to individuals in need thereof.

[0479] Also provided herein are methods of treating diabetes or prediabetes comprising administering to a subject in need thereof a therapeutically effective amount of a peptide product described above and herein and in Table 1 of Figure 1 and Table 2 of Figure 2 to an individual in need thereof.

[0480] Provided herein are methods for treating or delaying the progression or onset of conditions selected from diabetes, diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, insulin resistance, hyperglycemia, hyperinsulinemia, metabolic syndrome, diabetic complications, elevated blood levels of free fatty acids or glycerol, hyperlipidemia, obesity, hypertriglyceridemia, atherosclerosis, acute cardiovascular syndrome, infarction, ischemic reperfusion a hypertension, comprising administering a therapeutically effective amount of a peptide product described herein and in Table 1 of Figure 1 and Table 2 of Figure 2 to an individual in need thereof. In an additional embodiment, provided herein are methods for treating delays in wound healing comprising administering a therapeutically effective amount of a peptide product described herein and in Table 1 of Figure 1 and Table 2 of Figure 2 to an individual in need thereof.

[0481] In one embodiment said condition to be treated is diabetes. In one embodiment said condition to be treated is insulin resistance. In one embodiment said condition to be treated is the metabolic syndrome. In one embodiment said effective amount of said peptide is from about 0.1 µg/kg/day to about 100.0 µg/kg/day.

[0482] In one embodiment the method of administration is parenteral. In one embodiment the method of administration is per oral. In one embodiment the method of administration is subcutaneous. In one embodiment the method of administration is nasal insufflation.

[0483] Further provided herein is a method of reducing weight gain or inducing weight loss comprising administering a therapeutically effective amount of a peptide product described herein and in Table 1 of Figure 1 and Table 2 of Figure 2 to an individual in need thereof. In some embodiments, the weight gain is associated with metabolic syndrome.

[0484] Provided herein is a method of treating hypoglycemia comprising administering a therapeutically effective amount of a peptide product described herein and in Table 1 of Figure 1 and Table 2 of Figure 2 to an individual in need thereof.

[0485] Also provided herein are methods for treatment of diabetes comprising administering a therapeutically effective amount of a peptide product described herein and in Table 1 of Figure 1 and Table 2 of Figure 2 to an individual in need thereof and at least one additional therapeutic agent; wherein said therapeutic agent is selected from an antidiabetic agent, an anti-obesity agent, a satiety agent, an anti-inflammatory agent, an anti-hypertensive agent, an anti-atherosclerotic agent and a lipid-lowering agent.

[0486] In some embodiments of the methods described above, the peptide and/or protein that is covalently attached to a surfactant is a glucagon or GLP-1 peptide, or an analog thereof. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered prophylactically and delays occurrence of any condition associated with insulin resistance, including and not limited to the metabolic syndrome, hypertension, diabetes, type 2 diabetes, gestational diabetes, hyperlipidemia, atherosclerosis, systemic inflammation or the like. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered therapeutically and delays progression of any condition associated with the metabolic syndrome, hypertension, diabetes, type 2 diabetes, gestational diabetes, hyperlipidemia, atherosclerosis, systemic inflammation or the like. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered prophylactically and/or therapeutically and delays progression of insulin resistance to diabetes. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered prophylactically and/or therapeutically and reduces or halts further loss of insulin resistance, thereby stabilizing disease.

[0487] In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered parenterally. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered subcutaneously. In some embodiments, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered by nasal insufflation.

[0488] In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) has a longer duration of action compared to a pharmaceutical comprising currently known therapeutics (e.g., exenatide, metformin or the like).

Combination therapy with PTH analogs

[0489] In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered in combination with a bone resorption inhibitor including, a bisphosphonate, (e.g. alendronate) or strontium salt; or a substance with estrogen-like effect, e.g. estrogen; or a selective estrogen receptor modulator, e.g. raloxifene, tamoxifene, droloxifene, toremifene, idoxifene, or levormeloxifene; or a calcitonin-like substance, e.g. calcitonin; or a vitamin D analog; or a calcium salt. The therapeutic agents are optionally administered simultaneously, or sequentially in any order. By way of example, in some embodiments of the methods described above, a first regimen of the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered to an individual in need thereof, and the regimen is followed by a second regimen of bisphosphonate therapy. By way of example, in some other embodiments, a first regimen of the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 2-I-A, 2-III, 2-V, 2-VI or 2-VII) is administered to an individual in need thereof, followed by a drug holiday, followed by a second regimen of estrogen receptor modulators.

Combination therapy with GLP analogs

[0490] In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered in combination with other methods of treatment of the metabolic syndrome selected from the group comprising an antidiabetic agent, an anti-obesity agent, an anti-hypertensive agent, an anti-atherosclerotic agent and a lipid-lowering agent. By way of example, efficacious antidiabetic agents suitable for administration in combination with a surfactant-modified peptide and/or protein product described herein include a biguanide, a sulfonylurea, a glucosidase inhibitor a PPAR γ agonist, a PPAR α/γ dual agonist, an $\alpha 2$ inhibitor, a DPP4 inhibitor, an insulin sensitizer, a GLP-1 analog, insulin and a meglitinide. Additional examples include metformin, glyburide, glimepiride, glipiride, glipizide, chlorpropamide, gliclazide, acarbose, miglitol, pioglitazone, troglitazone, rosiglitazone, muraglitazar, insulin, G1-262570, isaglitazone, JTT-501, NN-2344, L895 645, YM-440, R-119702, A19677, repaglinide, nateglinide, KAD 1129, AR-HO 39242, GW-40 I 5 44, KRP2 I 7, AC2993, LY3 I 5902, NVP-DPP-728A and saxagliptin.

[0491] In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered in combination with other methods of treatment of the metabolic syndrome selected from the group of efficacious anti-obesity agents. By way of example, efficacious anti-obesity agents suitable for administration with the peptide products described herein include beta 3 adrenergic agonist, a lipase inhibitor, a serotonin (and dopamine) reuptake inhibitor, a thyroid receptor beta compound, a CB-1 antagonist, a NPY-Y2 and a NPY-Y4 receptor agonist and an anorectic agent. Specific members of these classes comprise orlistat, AfL-962, A19671, L750355, CP331648, sibutramine, topiramate, axokine, dexamphetamine, phentermine,

phenylpropanolamine, rimonabant (SRI 417164), and mazindol.

[0492] In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered in combination with other methods of treatment of the metabolic syndrome selected from the group of efficacious lipid-lowering agents. By way of example, efficacious lipid-lowering agents suitable for administration with the peptide products described herein include agents selected from the group consisting of an MTP inhibitor, cholesterol ester transfer protein, an HMG CoA reductase inhibitor, a squalene synthetase inhibitor, a fibrin acid derivative, an upregulator of LDL receptor activity, a lipoxigenase inhibitor, and an ACAT inhibitor. Specific examples from these classes comprise pravastatin, lovastatin, simvastatin, atorvastatin, cerivastatin, fluvastatin, nisvastatin, visastatin, fenofibrate, gemfibrozil, clofibrate, avasimibe, TS-962, MD-700, CP -52941 4, and LY295 427.

[0493] In some embodiments of the methods described above, the surfactant-modified peptide and/or protein (e.g., a peptide product of Formula 3-I-A, 3-III-A, 3-III-B or Formula 3-V) is administered in combination with peptide hormones, and analogs thereof, that are known to exhibit pro-satiety effects in animal models and in man. Contemplated within the scope of embodiments presented herein is a combination of the peptide products described herein and long-acting satiety agents for treatment of obesity. Examples of such peptide satiety agents include GLP-1, pancreatic polypeptide (PP), cholecystokinin (CCK), peptide YY (PYY), amylin, calcitonin, OXM, neuropeptide Y (NPY), and analogs thereof (Bloom, S.R., et al. (2008) *Mol Interv* 8: 82-98; Field, B.C., et al. (2009) *Br J Clin Pharmacol* 68: 830-843).

[0494] Also contemplated within the scope of embodiments presented herein are methods for treatment of obesity comprising administration of peptide products described herein in combination with peptide hormones including and not limited to leptin, ghrelin and CART (cocaine- and amphetamine-regulated transcript) analogs and antagonists.

[0495] Additional peptide products in the body are associated with fat cells or the obese state (adipokines) and are known to have proinflammatory effects (Gonzalez-Periz, A. and Claria, J. (2010) *Scientific WorldJournal* 10: 832-856). Such agents will have additional favorable actions when used in combination with the peptide products described herein. Examples of agents that offer a beneficial effect when used in combination with the peptide products described herein include analogs and antagonists of adiponectin, chemerin, visfatin, nesfatin, omentin, resistin, TNFalpha, IL-6 and obestatin.

Dosing

[0496] The covalently modified peptides and/or proteins described herein may be administered in any amount to impart beneficial therapeutic effect in a number of disease states. In some embodiments, covalently modified peptides and/or proteins described herein are useful in the treatment of inflammation. In an embodiment, compounds presented herein impart beneficial

activity in the modulation of post-operative or chronic pain. In an embodiment, the present peptides are administered to a patient at concentrations higher or lower than that of other forms of treatment which modulate pain. In yet another embodiment, the present peptides are administered with other compounds to produce synergistic therapeutic effects.

[0497] Representative delivery regimens include oral, parenteral (including subcutaneous, intramuscular and intravenous injection), rectal, buccal (including sublingual), transdermal, inhalation ocular and intranasal. An attractive and widely used method for delivery of peptides entails subcutaneous injection of a controlled-release injectable formulation. In some embodiments, covalently modified peptides and/or proteins described herein are useful for subcutaneous, intranasal and inhalation administration.

[0498] The selection of the exact dose and composition and the most appropriate delivery regimen will be influenced by, inter alia, the pharmacological properties of the selected peptide, the nature and severity of the condition being treated, and the physical condition and mental acuity of the recipient. Additionally, the route of administration will result in differential amounts of absorbed material. Bioavailabilities for administration of peptides through different routes are particularly variable, with amounts from less than 1% to near 100% being seen. Typically, bioavailability from routes other than intravenous, intraperitoneal or subcutaneous injection are 50% or less.

[0499] In general, covalently modified peptides and/or proteins described herein, or salts thereof, are administered in amounts between about 0.001 and 20 mg/kg body weight per day, between about 0.01 and 10 mg/kg body weight per day, between about 0.1 and 1000 µg/kg body weight per day, or between about 0.1 to about 100 µg/kg body weight per day. Routes of administration vary. For example, covalently modified opioid peptides and/or proteins described herein, or salts thereof, are administered in amounts between about 0.1 and 1000 µg/kg body weight per day, or between about 0.1 to about 100 µg/kg body weight per day, by subcutaneous injection. By way of example, for a 50 kg human female subject, the daily dose of active ingredient is from about 5 to about 5000 µg, or from about 5 to about 5000 µg by subcutaneous injection. Different doses will be needed, depending on the route of administration, the compound potency, the pharmacokinetic profile and the applicable bioavailability observed, and the active agent and the disease being treated. In an alternate embodiment where the administration is by inhalation, the daily dose is from 1000 to about 20,000 µg, twice daily. In other mammals, such as horses, dogs, and cattle, higher doses may be required. This dosage may be delivered in a conventional pharmaceutical composition by a single administration, by multiple applications, or via controlled release, as needed to achieve the most effective results.

[0500] Pharmaceutically acceptable salts retain the desired biological activity of the parent peptide without toxic side effects. Examples of such salts are (a) acid addition salts formed with inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; and salts formed with organic acids such as, for example, acetic acid, trifluoroacetic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric

acid, malic acid, ascorbic acid, benzoic acid, tannic acid, pamoic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acids, naphthalene disulfonic acids, polygalacturonic acid and the like; (b) base addition salts or complexes formed with polyvalent metal cations such as zinc, calcium, bismuth, barium, magnesium, aluminum, copper, cobalt, nickel, cadmium, and the like; or with an organic cation formed from N,N'-dibenzylethylenediamine or ethylenediamine; or (c) combinations of (a) and (b), e.g., a zinc tannate salt and the like.

[0501] Also contemplated, in some embodiments, are pharmaceutical compositions comprising as an active ingredient covalently modified peptides and/or proteins described herein, or pharmaceutically acceptable salt thereof, in admixture with a pharmaceutically acceptable, non-toxic carrier. As mentioned above, such compositions may be prepared for parenteral (subcutaneous, intramuscular or intravenous) administration, particularly in the form of liquid solutions or suspensions; for oral or buccal administration, particularly in the form of tablets or capsules; for intranasal administration, particularly in the form of powders, nasal drops, evaporating solutions or aerosols; for inhalation, particularly in the form of liquid solutions or dry powders with excipients, defined broadly; and for rectal or transdermal administration.

[0502] The compositions may conveniently be administered in unit dosage form and may be prepared by any of the methods well-known in the pharmaceutical art, for example as described in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, Pa., (1985). Formulations for parenteral administration may contain as excipients sterile water or saline, alkylene glycols such as propylene glycol, polyalkylene glycols such as polyethylene glycol, saccharides, oils of vegetable origin, hydrogenated naphthalenes, serum albumin nanoparticles (as used in Abraxane™, American Pharmaceutical Partners, Inc. Schaumburg IL), and the like. For oral administration, the formulation can be enhanced by the addition of bile salts or acylcarnitines. Formulations for nasal administration may be solid or solutions in evaporating solvents such as hydrofluorocarbons, and may contain excipients for stabilization, for example, saccharides, surfactants, submicron anhydrous α -lactose or dextran, or may be aqueous or oily solutions for use in the form of nasal drops or metered spray. For buccal administration typical excipients include sugars, calcium stearate, magnesium stearate, pregelatinated starch, and the like.

[0503] When formulated for nasal administration, the absorption across the nasal mucous membrane may be further enhanced by surfactants, such as for example, glycocholic acid, cholic acid, taurocholic acid, ethocholic acid, deoxycholic acid, chenodeoxycholic acid, dehydrocholic acid, glycodeoxycholic acid, cyclodextrins and the like in an amount in the range between about 0.1 and 15 weight percent, between about 0.5 and 4 weight percent, or about 2 weight percent. An additional class of absorption enhancers reported to exhibit greater efficacy with decreased irritation is the class of alkyl maltosides, such as tetradecylmaltoside (Arnold, J.J., et al. (2004) J Pharm Sci 93: 2205-2213, Ahsan, F., et al. (2001) Pharm Res 18: 1742-1746) and references therein.

[0504] When formulated for delivery by inhalation, a number of formulations offer advantages. Adsorption of the active peptide to readily dispersed solids such as diketopiperazines (for

example Technosphere particles; (Pfutzner, A. and Forst, T. (2005) Expert Opin Drug Deliv 2: 1097-1106) or similar structures gives a formulation which results in a rapid initial uptake of the therapeutic agent. Lyophilized powders, especially glassy particles, containing the active peptide and an excipient are useful for delivery to the lung with good bioavailability, for example, see Exubera® (inhaled insulin by Pfizer and Aventis Pharmaceuticals Inc.). Additional systems for delivery of peptides by inhalation are described (Mandal, T.K., Am. J. Health Syst. Pharm. 62: 1359-64 (2005)).

[0505] Delivery of covalently modified peptides and/or proteins described herein to a subject over prolonged periods of time, for example, for periods of one week to one year, may be accomplished by a single administration of a controlled release system containing sufficient active ingredient for the desired release period. Various controlled release systems, such as monolithic or reservoir-type microcapsules, depot implants, polymeric hydrogels, osmotic pumps, vesicles, micelles, liposomes, transdermal patches, iontophoretic devices and alternative injectable dosage forms may be utilized for this purpose. Localization at the site to which delivery of the active ingredient is desired is an additional feature of some controlled release devices, which may prove beneficial in the treatment of certain disorders.

[0506] One form of controlled release formulation contains the peptide or its salt dispersed or encapsulated in a slowly degrading, non-toxic, non-antigenic polymer such as copoly(lactic/glycolic) acid, as described in the pioneering work of Kent, Lewis, Sanders, and Tice, U.S. Pat. No. 4,675,189. The compounds, or their salts, may also be formulated in cholesterol or other lipid matrix pellets, or silastomer matrix implants. Additional slow release, depot implant or injectable formulations will be apparent to the skilled artisan. See, for example, Sustained and Controlled Release Drug Delivery Systems, J. R. Robinson ed., Marcel Dekker, Inc., New York, 1978, and R. W. Baker, Controlled Release of Biologically Active Agents, John Wiley & Sons, New York, 1987.

[0507] An additional form of controlled-release formulation comprises a solution of a biodegradable polymer, such as copoly(lactic/glycolic acid) or block copolymers of lactic acid and PEG, in a bioacceptable solvent, which is injected subcutaneously or intramuscularly to achieve a depot formulation. Mixing of the peptides described herein with such a polymeric formulation is suitable to achieve very long duration of action formulations.

[0508] The covalently modified peptides and/or proteins described herein and the reagents for the synthesis thereof are more particularly described in the following examples which are intended as illustrative only since numerous modifications and variations therein will be apparent to those of ordinary skill in the art.

EXAMPLES

Example 1: Reagents - N- α -Fmoc, N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine

[0509] In an oven-dried 250 mL Erlenmeyer flask is placed 1-octyl β -D-glucuronic acid (Carbosynth Ltd., 3.06 g, 10 mmol), 50 mL anhydrous DMF, and anhydrous HOBt (1.62 g, 12 mmol). A chilled (4°C) solution of DCC (2.48g, 12 mmol) in 50 mL of DMF is added, with stirring, and the reaction is allowed to proceed for 5 min. The copious white precipitate of N, N'-dicyclohexylurea is filtered on a fritted glass funnel and the filtrate is added to a solution of N- α -Fmoc-L-lysine (3.68g, 10 mmol) in 25 ml anhydrous DMF. The reaction is allowed to proceed for 25 min with warming to room temp or until the ninhydrin color is very faint. The reaction mixture is filtered, stripped to dryness and crystallized from MeOH/Et₂O by dissolution in MeOH and slow dilution to the cloud point with Et₂O, followed by refrigeration. Further purification can be achieved by silica gel chromatography using a solvent gradient from EtOAc to EtOAc/EtOH/AcOH.

[0510] In a similar manner, but substituting N- α -Boc-L-lysine is obtained N- α -Boc,N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine, suitable for N-terminal incorporation and cleavage to a free N-Terminus. In a similar manner, but substituting N- α -Ac-L-lysine is obtained N- α -Ac,N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine, suitable for incorporation at the N-terminus of a peptide with a blocked N-terminus. In a similar manner, but substituting the appropriate amount of N- α -Fmoc-L-ornithine is obtained N- α -Fmoc,N- δ -(1-octyl β -D-glucuronide-6-yl)-L-ornithine. In a similar manner but substituting other N-mono-protected diamino acids one obtains the corresponding reagents. Alternatively, use of a transient Me₃Si ester protecting group during the coupling and without preactivation of the 1-octyl β -D-glucuronic acid provides a facile route to the formation of the reagents. The transient Me₃Si ester is produced by reaction of the Fmoc-Lys-OH with an equimolar amount of N,O-bis(trimethylsilyl)acetamide in dichloromethane (CH₂Cl₂). The organic layer contains the desired reagent as a solution in CH₂Cl₂ ready for coupling with the 1-alkyl glucuronide as above. The filtered reaction mixture is washed with aqueous NaHSO₄ to hydrolyze the Me₃Si ester, dried over MgSO₄ and solvent is removed.

[0511] Similarly, but using peracetyl or perbenzoyl 1-octyl β -D-glucuronic acid one obtains the Ac, or Bz protected form of the reagents (e.g. 2,3,4-trisacetyl 1-octyl β -D-glucuronic acid, and the like, formed by treatment with Ac₂O). Such reagents have increased stability during acid cleavage from the resin and are used when instability during deprotection is detected, see (Kihlberg, J., et al. (1997) *Methods Enzymol* 289: 221-245) and references therein. Final deprotection of such products is carried out by base-catalyzed transesterification after cleavage, by use of MeOH/NH₃, MeOH/NaOMe, MeOH/NH₂NH₂, as described above.

Example 2: Synthetic Peptide Analogs

[0512] In general, peptide synthesis methods involve the sequential addition of protected amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid and any reactive side chain group are protected. This protected amino acid is

then either attached to an inert solid support, or utilized in solution, and the next amino acid in the sequence, also suitably protected, is added under conditions amenable to formation of the amide linkage. After all the desired amino acids have been linked in the proper sequence, protecting groups and any solid support are removed to afford the crude peptide. The peptide is desalted and purified chromatographically.

[0513] A preferred method of preparing the analogs of the physiologically active truncated peptides, having fewer than about fifty amino acids, involves solid phase peptide synthesis. In this method the α -amino ($N\alpha$) functions and any reactive side chains are protected by acid- or base-sensitive groups. The protecting group should be stable to the conditions of peptide linkage formation, while being readily removable without affecting the extant peptide chain. Suitable α -amino protecting groups include, but are not limited to t-butoxycarbonyl (Boc), benzyloxycarbonyl (Cbz), o-chlorobenzyloxycarbonyl, biphenylisopropylloxycarbonyl, t-amyloxycarbonyl (Amoc), isobornyloxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, o-nitrophenylsulfenyl, 2-cyano-t-butoxycarbonyl, 9-fluorenyl-methoxycarbonyl (Fmoc) and the like, preferably Boc or more preferably, Fmoc. Suitable side chain protecting groups include, but are not limited to: acetyl, benzyl (Bzl), benzyloxymethyl (Bom), Boc, t-butyl, o-bromobenzyloxycarbonyl, t-butyl, t-butyldimethylsilyl, 2-chlorobenzyl (Cl-z), 2,6-dichlorobenzyl, cyclohexyl, cyclopentyl, isopropyl, pivalyl, tetrahydropyran-2-yl, tosyl (Tos), 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf), trimethylsilyl and trityl. A preferred $N\alpha$ -protecting group for synthesis of the compounds is the Fmoc group. Preferred side chain protecting groups are O-t-Butyl group for Glu, Tyr, Thr, Asp and Ser; Boc group for Lys and Trp side chains; Pbf group for Arg; Trt group for Asn, Gln, and His. For selective modification of a Lys residue, orthogonal protection with a protecting group not removed by reagents that cleave the Fmoc or t-butyl based protecting groups is preferred. Preferred examples for modification of the Lys side chain include, but are not limited to, those removed by hydrazine but not piperidine; for example 1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)-3-methylbutyl (ivDde) or 1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethyl (Dde) and allyloxycarbonyl (Alloc).

[0514] The Fmoc-Lys(ivDde) or Fmoc-Lys(Dde) protecting group scheme is preferred in cases where side chain lactam formation is desired (Houston, M.E., Jr., et al. (1995) J Pept Sci 1: 274-282; Murage, E.N., et al. (2010) J Med Chem), since in this case Fmoc-Glu(O-Allyl) and Fmoc-Lys(Alloc) can be incorporated and used to provide transient protection, then deprotected for lactam formation while the Lys(Dde) protecting group remains for later removal and reaction with the functionalized surfactant. The side chain lactam between acidic and basic residue (e.g. Glu and Lys) is carried out after removal of the allyl-based protection by activation of the carboxyl side chain function with N,N'-diisopropylcarbodiimide (DIC)/1-hydroxybenzotriazole (HOBt) or 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate (HBTU)/ N,N-di-isopropylethylamine (DIEA), using standard protocols well known to the art.

[0515] In solid phase synthesis, the C-terminal amino acid is first attached to a suitable resin support. Suitable resin supports are those materials which are inert to the reagents and reaction conditions of the stepwise condensation and deprotection reactions, as well as being

insoluble in the media used. Examples of commercially available resins include styrene/divinylbenzene resins modified with a reactive group, e.g., chloromethylated co-poly-(styrene-divinylbenzene), hydroxymethylated co-poly-(styrene-divinylbenzene), and the like. Benzylated, hydroxymethylated phenylacetamidomethyl (PAM) resin is preferred for the preparation of peptide acids. When the C-terminus of the compound is an amide, preferred resins are p-methylbenzhydrylamino-co-poly(styrene-divinyl-benzene) resin, a 2,4 dimethoxybenzhydrylamino-based resin ("Rink amide"), 4-Hydroxymethylphenoxyacetyl aminomethyl resin (HMP Am) and the like. An especially preferred support for the synthesis of larger peptides are commercially available resins containing PEG sequences grafted onto other polymeric matrices, such as the Rink Amide-PEG and PAL-PEG-PS resins (Applied Biosystems) or similar resins designed for peptide amide synthesis using the Fmoc protocol. Thus in certain cases it is desirable to have an amide linkage to a PEG chain. In those cases it is convenient to link an N-Fmoc-amino-PEG-carboxylic acid to the amide forming resin above (e.g. Rink amide resin and the like). The first amino acid of the chain can be coupled as an N-Fmoc-amino acid to the amino function of the PEG chain. Final deprotection will yield the desired Peptide-NH-PEG-CO-NH₂ product.

[0516] Attachment to the PAM resin may be accomplished by reaction of the Na⁺ protected amino acid, for example the Boc-amino acid, as its ammonium, cesium, triethylammonium, 1,5-diazabicyclo-[5.4.0]undec-5-ene, tetramethylammonium, or similar salt in ethanol, acetonitrile, N,N-dimethylformamide (DMF), and the like, preferably the cesium salt in DMF, with the resin at an elevated temperature, for example between about 40° and 60 °C, preferably about 50 °C, for from about 12 to 72 hours, preferably about 48 hours. This will eventually yield the peptide acid product following acid cleavage or an amide following aminolysis.

[0517] The Na-Boc-amino acid may be attached to the benzhydrylamine or HMP Am resin by means of, for example, a DIC/HOBt mediated coupling for from about 2 to about 24 hours, preferably about 2 hours at a temperature of between about 10° and 50 °C, preferably 25 °C in a solvent such as CH₂Cl₂ or DMF, preferably CH₂Cl₂.

[0518] For Boc-based protocols, the successive coupling of protected amino acids may be carried out by methods well known in the art, typically in an automated peptide synthesizer. Following neutralization with triethylamine, DIEA, N-methylmorpholine (NMM), collidine, or similar base, each protected amino acid is introduced in approximately about 1.5 to 2.5 fold molar excess and the coupling carried out in an inert, nonaqueous, polar solvent such as CH₂Cl₂, DMF, N-methylpyrrolidone (NMP), N,N-dimethylacetamide (DMA), or mixtures thereof, preferably in dichloromethane at ambient temperature. For Fmoc-based protocols no acid is used for deprotection but a base, preferably DIEA or NMM, is usually incorporated into the coupling mixture. Couplings are typically done in DMF, NMP, DMA or mixed solvents, preferably DMF. Representative coupling agents are N,N'-dicyclohexylcarbodiimide (DCC), DIC or other carbodiimide, either alone or in the presence of HOBt, O-acyl ureas, benzotriazol-1-yl-oxytris(pyrrolidino)phosphonium hexafluorophosphate (PyBop), N-hydroxysuccinimide, other N-hydroxyimides, or oximes. Alternatively, protected amino acid active esters (e.g. p-

nitrophenyl, pentafluorophenyl and the like) or symmetrical anhydrides may be used. Preferred coupling agents are of the aminium/uronium (alternative nomenclatures used by suppliers) class such as HBTU, O-(7-azabenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU), 2-(6-Chloro-1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate (HCTU), and the like.

[0519] A preferred method of attachment to the Fmoc-PAL-PEG-PS resin may be accomplished by deprotection of the resin linker with 20% piperidine in DMF, followed by reaction of the N- α -Fmoc protected amino acid, about a 5 fold molar excess of the N- α -Fmoc-amino acid, using HBTU: DIEA (1:2) in DMF in a microwave-assisted peptide synthesizer with a 5min, 75° max coupling cycle.

[0520] For this Fmoc-based protocol in the microwave-assisted peptide synthesizer, the N- α -Fmoc amino acid protecting groups are removed with 20% piperidine in DMF containing 0.1M 1-hydroxybenzotriazole (HOBt), in a double deprotection protocol for 30 sec and then for 3min with a temperature maximum set at 75°C. HOBt is added to the deprotection solution to reduce aspartimide formation. Coupling of the next amino acid then employs a five-fold molar excess using HBTU:DIEA (1:2) with a 5min, 75° max. double-coupling cycle.

[0521] At the end of the solid phase synthesis the fully protected peptide is removed from the resin. When the linkage to the resin support is of the benzyl ester type, cleavage may be effected by means of aminolysis with an alkylamine or fluoroalkylamine for peptides with an alkylamide C-terminus, or by ammonolysis with, for example, ammonia/methanol or ammonia/ethanol for peptides with an unsubstituted amide C-terminus, at a temperature between about -10° and 50° C, preferably about 25° C, for between about 12 and 24 hours, preferably about 18 hours. Peptides with a hydroxy C-terminus may be cleaved by HF or other strongly acidic deprotection regimen or by saponification. Alternatively, the peptide may be removed from the resin by transesterification, e.g., with methanol, followed by aminolysis or saponification. The protected peptide may be purified by silica gel or reverse-phase HPLC.

[0522] The side chain protecting groups may be removed from the peptide by treating the aminolysis product with, for example, anhydrous liquid hydrogen fluoride in the presence of anisole or other carbonium ion scavenger, treatment with hydrogen fluoride/pyridine complex, treatment with tris(trifluoroacetyl)boron and trifluoroacetic acid, by reduction with hydrogen and palladium on carbon or polyvinylpyrrolidone, or by reduction with sodium in liquid ammonia, preferably with liquid hydrogen fluoride and anisole at a temperature between about - 10° and +10° C, preferably at about 0°C, for between about 15 minutes and 2 hours, preferably about 1.5 hours.

[0523] For peptides on the benzhydrylamine type resins, the resin cleavage and deprotection steps may be combined in a single step utilizing liquid hydrogen fluoride and anisole as described above or preferably through the use of milder cleavage cocktails. For example, for the PAL-PEG-PS resin, a preferred method is through the use of a double deprotection protocol in the microwave-assisted peptide synthesizer using one of the mild cleavage cocktails

known in the art, such as TFA/water/tri-iso-propylsilane/3,6-dioxa-1,8-octanedithiol (DODT) (92.5/2.5/2.5/2.5) for 18min at 38°C each time. Cleavage of alkyl glycoside containing materials have shown survival of the alkyl glycoside linkage using protocols with TFA/water ratios in the 9/1 to 19/1 range. A typical cocktail is 94% TFA; 2% EDT; 2% H₂O; 2% TIS. Typically the fully deprotected product is precipitated and washed with cold (-70° to 4°C) diethylether, dissolved in deionized water and lyophilized.

[0524] The peptide solution may be desalted (e.g. with BioRad AG-3® anion exchange resin) and the peptide purified by a sequence of chromatographic steps employing any or all of the following types: ion exchange on a weakly basic resin in the acetate form; hydrophobic adsorption chromatography on underivatized co-poly(styrene-divinylbenzene), e.g. Amberlite® XAD; silica gel adsorption chromatography; ion exchange chromatography on carboxymethylcellulose; partition chromatography, e.g. on Sephadex® G-25; counter-current distribution; supercritical fluid chromatography; or HPLC, especially reversed-phase HPLC on octyl- or octadecylsilylsilica (ODS) bonded phase column packing.

[0525] Also provided herein are processes for preparing covalently modified peptides and/or proteins described herein and pharmaceutically acceptable salts thereof, which processes comprise sequentially condensing protected amino acids on a suitable resin support, removing the protecting groups and resin support, and purifying the product, to afford analogs of the physiologically active truncated homologs and analogs of the covalently modified peptides and/or proteins described herein. In some embodiments, covalently modified peptides and/or proteins described herein incorporate alkyl glycoside modifications as defined above.

[0526] Another aspect relates to processes for preparing covalently modified peptides and/or proteins described herein and pharmaceutically acceptable salts thereof, which processes comprise the use of microwave-assisted solid phase synthesis-based processes or standard peptide synthesis protocols to sequentially condense protected amino acids on a suitable resin support, removing the protecting groups and resin support, and purifying the product, to afford analogs of the physiologically active peptides, as defined above.

Example 3: N-Terminal endomorphin-1 analog - AcLys(1-octyl β-D-glucuronide-6-yl)endomorphin 1 (Ac-Lys(1-octyl β-glucuronide-6-yl)-Tyr-Pro-Trp-Phe-NH₂ (AcLys(1-octyl β-D-glucuronide-6-yl)endomorphin 1)

[0527] As described above, Fmoc-Tyr(*t*-Bu)-Pro-Trp(Boc)-Phe-NH-Rink amide MBHA resin is prepared as described in (Koda, Y., et al. (2008) Bioorg Med Chem 16: 6286-6296). The resin is N α -deprotected with piperidine/DMF solution, washed with solvent and coupled with 2 equivalents of N- α -Ac,N- ϵ -(1-Octyl β-D-glucuronide-6-yl)-L-lysine using a standard coupling mixture (e.g. HBTU/DIPEA). Following completion of the coupling, the peptide is deprotected and cleaved from the resin using a deprotection mixture (95% TFA/5%H₂O or DODT). The solvent is removed by a stream of nitrogen and the crude peptide is precipitated with cold

Et₂O, collected, dissolved in 20% acetonitrile and lyophilized. Purification is by reversed phase hplc in a mobile phase containing a gradient from H₂O to H₂O/acetonitrile using a 0.1% TFA or NH₄OAc buffer system. The mixture is subjected to multiple lyophilizations from H₂O.

Example 4: N-Terminal endomorphin-1 analog - AcLys(1-octyl β-D-glucuronide-6-yl)endomorphin 1 (Ac-Lys(1-octyl β-D-glucuronide-6-yl)-Tyr-Pro-Trp-Phe-NH₂ (AcLys(1-octyl β-D-glucuronide-6-yl)endomorphin 1)

[0528] As described above, Ac-Lys(Boc)-Tyr(*t*-Bu)-Pro-Trp(Boc)-Phe-NH-Rink amide MBHA resin is prepared as described in. Following completion of the synthesis, the peptide is deprotected and cleaved from the resin using a deprotection mixture (95% TFA/5% H₂O or DODT). The solvent is removed by a stream of nitrogen and the crude peptide is precipitated with cold Et₂O, collected, dissolved in 20% acetonitrile and lyophilized. The peptide, containing a deprotected Lys ε-amino function is coupled with 2 equivalents of 1-octyl β-D-glucuronic acid using a standard coupling mixture (e.g. HBTU/DIPEA) in DMF or similar anhydrous aprotic solvent. The solvent is removed in vacuo and the product is lyophilized from 20% acetonitrile or H₂O. Purification is by reversed phase hplc in a mobile phase containing a gradient from H₂O to H₂O/acetonitrile using a 0.1% TFA or NH₄OAc buffer system. The mixture is subjected to multiple lyophilizations from H₂O.

Example 5: 2',6'-dimethyl-L-tyrosyl-prolyl-2',4',6'-trimethyl-L-phenylalanyl-Nε-(1'-octyl β-D-glucuronyl-L-lysine amide (EU-A102).

[0529] A 0.3 mmol sample of Fmoc-Rink-Amide resin (0.5 mmol/g) was coupled with the following sequence of amino acids using a standard DIC/HOBt solid phase coupling protocol (3 equivalent of DIC/HOBt and amino acid): Fmoc-L-lysine(Nε-Alloc); Fmoc-2',4',6'-trimethyl-L-phenylalanine; Fmoc-L-proline; Fmoc-2',4'-dimethyl-L-tyrosine.

[0530] The sample of 2',6'-dimethyl-L-tyrosyl-prolyl-2',4',6'-trimethyl-L-phenylalanyl-Nε-(Alloc)-L-lysine amide resin was deprotected on the Lys-Nε position by incubation with Pd(PPh₃)₄ (0.5 eq) and DMBA (20 eq) in DMF/CH₂Cl₂ (1:1) overnight in the dark at room temperature. Following washing by DMF/CH₂Cl₂, the Lys side chain was acylated with 1'-octyl β-D-glucuronic acid (Carbosynth) in DMF/CH₂Cl₂ through the use of DIC/HOBt. Completion of the coupling was checked by ninhydrin and the product was washed extensively with CH₂Cl₂.

[0531] The product resin (0.77g) was submitted to final deprotection and cleavage from the resin by treatment with the cleavage cocktail (94% TFA; 2% EDT; 2% H₂O; 2% TIS) for a period of 240 min at room temperature. The mixture was treated with Et₂O, to precipitate the product and washed extensively with Et₂O to yield 290 mg of crude title peptide product after drying in

vacuo.

[0532] Purification was carried out in two batches by reversed-phase (C18) hplc. The crude peptide was loaded on a 4.1x25 cm hplc column at a flow rate of 15 mL/min (15% organic modifier; acetic acid buffer) and eluted with a gradient from 15-45% buffer B in 60min at 50°C. The product fraction was lyophilized to yield 100 mg of the title product peptide with a purity >96% by analytical hplc/mass spectrometry (M+1 peak = 911.87). The overall synthesis yield was calculated at 18%.

[0533] In a similar manner, but using the reagents 1'-methyl β -D-glucuronic acid and 1'-dodecyl β -D-glucuronic acid, were prepared the corresponding N ϵ -(1'-methyl β -D-glucuronyl)-L-lysine⁴ (EU-A101) and N ϵ -(1'-dodecyl β -D-glucuronyl)-L-lysine⁴ (EUA-103) analogs of the title compound.

[0534] Analysis was done by HPLC/mass spectrometry in positive ion mode using the eluent gradients given in the table below.

Compound Name	Position 4 N ϵ	Molecular Wt expected	Molecular Wt found	HPLC (min; elution)
EU-A101	Me	812.95	812.80	13.6 [a]
EU-A102	n-octyl	911.16	910.87	12.9 [b]
EU-A103	n-dodecyl	967.27	966.53	12.5 [c]

HPLC gradients in 0.1% TFA [a.] 20 to 50% CH₃CN over 30 min.
 [b.] 35 to 65% CH₃CN over 30 min. [c.] 40 to 75% CH₃CN over 20 min.
 HPLC on Phenomenex Luna C18 5micron 250x4.6 mm

Example 6: **2',6'-dimethyl-L-tyrosyl-prolyl-2',4',6'-trimethyl-L-phenylalanyl-L-phenylalanyl-N ϵ -(1'-dodecyl β -D-glucuronyl)-L-lysine amide (EU-A106).**

[0535] In a similar manner to that given for the solid phase synthesis and lysine side chain modification in example 5, given above, but using 1-dodecyl β -D-glucuronic acid (Milkereit, G., et al. (2004) Chem Phys Lipids 127: 47-63) for lysine N ϵ -acylation, was prepared the title peptide as a crude product. Following reversed phase hplc purification as above one obtains the title product as a white powder, of 96.1% purity by hplc/mass spectrometry (M+1 = 1114.8).

[0536] In a similar manner, but using the reagent 1'-methyl β -D-glucuronic acid, was prepared the corresponding N ϵ -(1'-methyl β -D-glucuronyl)-L-lysine⁵ (EUA-105) analog of the title compound.

[0537] Analysis was done by HPLC/mass spectrometry in positive ion mode using the

gradient eluents given in the table below.

Compound Name	Position 5 Nε	Molecular Wt expected	Molecular Wt found	HPLC (min; elution)
EU-A105	Me	960.12	959.60	8.8 [d]
EU-A106	n-dodecyl	1114.44	1113.80	11.6 [e]
HPLC gradients in 0.1% TFA [d.] 30 to 60% CH ₃ CN over 20 min. [e.] 45 to 75% CH ₃ CN over 20 min. HPLC on Phenomenex Luna C18 5micron 250x4.6 mm				

Example 7: 2',6'-dimethyl-L-tyrosyl-Nε-(1'-dodecyl β-D-glucuronyl)-D-lysyl-2',4',6'-trimethyl-L-phenylalanyl-L-phenylalanine-amide (EU-A108).

[0538] In a similar manner as that given in Example 6 was prepared the title peptide as a white powder of 95.8% purity by hplc/mass spectrometry (M= 1017.07).

[0539] In a similar manner, but using the reagent 1'-methyl β-D-glucuronic acid, was prepared the corresponding Nε-(1'-methyl β-D-glucuronyl)-L-lysine (EU-A107) analog of the title compound.

[0540] Analysis was done by HPLC/mass spectrometry in positive ion mode using the eluent gradients given in the table below.

Compound Name	Position 2 Nε	Molecular Wt expected	Molecular Wt found	HPLC (min; elution)
EU-A107	Me	862.01	862.73	11.6 [f]
EU-A108	n-dodecyl	1017.32	1017.07	11.9 [g]
HPLC gradients in 0.1% TFA [f.] 25 to 55% CH ₃ CN over 20 min. [g.] 50 to 80% CH ₃ CN over 20 min. HPLC on Phenomenex Luna C18 5micron 250x4.6 mm				

Example 8: 2',6'-dimethyl-L-tyrosyl-L-1,2,3,4-tetrahydroisoquinoline-3-carbonyl-L-phenylalanyl-Nε-(1'-methyl β-D-glucuronyl)-L-lysyl-amide (EU-A178).

[0541] In a similar manner as that given in Example 6 was prepared the title peptide as a white powder of 95.5% purity by hplc/mass spectrometry (M= 832.33).

[0542] In a similar manner, but using the reagent 1'-dodecyl β-D-glucuronic acid, was prepared the corresponding Nε-(1'-dodecyl β-D-glucuronyl)-L-lysine analog (EU-A179) of the

title compound. In a similar manner, but using the corresponding 1-alkyl glucuronic acid reagents are made the corresponding peptides of the invention, EU-A180, EU-A181, EU-A182, EU-A183, EU-A184, EU-A185, and the like.

[0543] Analysis was done by HPLC/mass spectrometry in positive ion mode using the eluent gradients given in the table below.

Compound Name	Position 4 Nε	Molecular Wt expected	Molecular Wt found	HPLC (min; elution)
EU-A178	Me	832.91	832.33	11.2 [h]
EU-A179	n-dodecyl	987.23	986.47	10.7 [f]
EU-A180	n-octyl	931.12	930.67	10.1 [i]
HPLC gradients in 0.1% TFA [f.] 25 to 55% CH ₃ CN over 20 min. [h.] 20 to 50% CH ₃ CN over 20 min. [i.] 35 to 65% CH ₃ CN over 20 min HPLC on Phenomenex Luna C18 5micron 250x4.6 mm				

Example 9: 2',6'-dimethyl-L-tyrosyl-L-1,2,3,4-tetrahydroisoquinoline-3-carbonyl-L-phenylalanyl-L-phenylalanyl-Nε-(1'-methyl β-D-glucuronyl)-L-lysyl-amide (EU-A189).

[0544] In a similar manner as that given in Example 6 was prepared the title peptide as a white powder of 98.99% purity by hplc/mass spectrometry (M= 979.53).

[0545] In a similar manner, but using the reagent 1'-dodecyl β-D-glucuronic acid, was prepared the corresponding Nε-(1'-dodecyl β-D-glucuronyl)-L-lysine (EU-A190) in position 5 analog of the title compound.

[0546] Analysis was done by HPLC/mass spectrometry in positive ion mode using the eluent gradients given in the table below.

Compound Name	Position 5 Nε	Molecular Wt expected	Molecular Wt found	HPLC (min; elution)
EU-A189	Me	980.09	979.53	10.9 [e]
EU-A190	n-dodecyl	1134.41	1134.53	12.5 [e]
HPLC gradients in 0.1% TFA [e.] 45 to 75% CH ₃ CN over 20 min. HPLC on Phenomenex Luna C18 5micron 250x4.6 mm				

Example 10: Additional analogs of the invention.

[0547] In a similar manner as that given in Example 6, but using the corresponding 1-alkyl β-D-

glucuronic acid reagent are prepared the additional peptides of the invention as white powders of greater than 95% purity by hplc/mass spectrometry.

[0548] Analysis is done by HPLC/mass spectrometry in positive ion mode using the appropriate eluent gradients such as those given in the table below.

Compound Name	Position 3 or 4 Nε	Molecular Wt expected	Molecular Wt found	HPLC (min; elution)
EU-A600	Me	918.02	918.00	10.7 [h]
EU-A601	n-octyl	1016.23	1015.87	10.1 [i]
EU-A615	Me	832.91	832.67	11.0 [h]
EU-A620	Me	923.06	922.73	9.5 [d]
EU-A639	Me	923.06	922.80	11.5 [f]
HPLC gradients in 0.1% TFA [d.] 30 to 60% CH ₃ CN over 20 min. [e.] 45 to 75% CH ₃ CN over 20 min. [h.] 20 to 50% CH ₃ CN over 20 min. [f.] 25 to 55% CH ₃ CN over 20 min. [i.] 35 to 65% CH ₃ CN over 20 min HPLC on Phenomenex Luna C18 5micron 250x4.6 mm				

Example 11. General oxidation method for uronic acids

[0549] To a solution of 1-dodecyl β-D-glucopyranoside (Carbosynth) [2.0g, 5.74 mmol] in 20 mL of acetonitrile and 20 mL of DI water was added (diacetoxyiodo)benzene (Fluka) [4.4 g, 13.7 mmol] and TEMPO (SigmaAldrich) [0.180g, 1.15 mmol]. The resulting mixture was stirred at room temperature for 20 h. The reaction was followed by mass spectrometry (for example LCQ ESI) and upon completion, the reaction mixture was diluted with water and lyophilized to dryness to give 1.52 g (crude yield 73.1%) of the crude product, 1-dodecyl β-D-glucuronic acid, as a white powder, which was used directly for the solid phase synthesis without further purification. In a like manner, but using the corresponding 1-tetradecyl, 1-hexadecyl, and 1-octadecyl β-D-glucopyranosides (purchased from Anatrace, Maumee, OH) were prepared the desired alkyl saccharide uronic acids used to make the products and reagents described herein. This product was previously prepared by an alternative process using NaOCl as oxidant and also has been used for longer alkyl groups. For longer alkyl groups 1,4-dioxane was used instead of acetonitrile and the temperature was raised as high as 30°C.

In a like manner, but using, for example, the corresponding 1-octyl, 1-decyl, 1-undecyl, 1-tetradecyl, 1-hexadecyl, and 1-octadecyl glycosides (purchased from Anatrace, Maumee, OH) were prepared the desired 1-alkyl saccharide uronic acids which were used to make the products and reagents described herein. In a like manner, but using, for example, the corresponding 1-octyl, 1-decyl, 1-undecyl, 1-tetradecyl, 1-hexadecyl, and 1-octadecyl β-D-melibiosides or β-D-maltosides (purchased from Anatrace, Maumee, OH) were prepared the desired 1-alkyl disaccharide uronic acids which were used to make the products and reagents described herein. In each case the corresponding alpha glycosides can be treated similarly.

Example 12: Cellular assay of the compounds.

[0550] Compounds were weighed precisely in an amount of approximately 2 mg and assayed in standard cellular assays by the contract research organization Cerep, Inc. (Pullman, WA) as executed at their subsidiary, Cerep SA (Le Bois l'Eveque, France). The readout is the amount of cAMP generated in the cells treated with the test compounds, in either agonist or antagonist mode. The assays used were the mu opioid receptor cellular assay (MOP in agonist and antagonist mode), the delta2 opioid receptor cellular assay (DOP in agonist and antagonist mode) and the kappa opioid receptor cellular assay (KOP in agonist and antagonist mode). The assays used are described in Wang, J.B., et al. (1994) FEBS Lett 338: 217-222, Law, P.Y. and Loh, H.H. (1993) Mol Pharmacol 43: 684-693, and Avidor-Reiss, T., et al. (1995) FEBS Lett 361: 70-74. The stimulant for the cells in the DOR antagonist assay was 3×10^{-8} M DPDPE, a well-accepted DOR literature standard.

[0551] For the series of compounds EU-A101 to EU-A103, where the hydrophobic portion of the surfactant (1-alkyl glucuronic acid) varies in length from C1 to C12, the character of the receptor selectivity and activation (see below) varies from full agonist (C1) to pure antagonist (C12). No activity was seen in cells used for DOP or KOP assays, thus showing selectivity for the mu opioid receptor as agonists. This behavior demonstrates the ability of the modifications described herein to vary the fundamental properties of the receptor interactions. Modifications elsewhere in the molecule (e.g., using amino acid analogs) are used to further modify the potency and character of the interaction of the drug candidates.

Compound Name	Structure	MOP agonist EC_{50} (nM)	MOP antagonist IC_{50} (nM)	Characterization
EU-A101	Me	13	nc	pure agonist
EU-A102	n-octyl	100	100	partial agonist
EU-A103	n-dodecyl	Nc	86	pure antagonist
EU-A107	Me	60	nc	Agonist

[0552] DOP antagonistic activity was assessed by inhibition of the stimulatory cAMP response of 3×10^{-8} M DPDPE.

Compound Name	Structure	MOP agonist EC_{50} (nM)	DOP antagonist IC_{50} (M)	Characterization
EU-A178	Me	15	$<10^{-10}$	Pure MOP Ag; pure DOP antag
EU-A179	n-dodecyl	§	NT	Pure MOP Ag; pure DOP antag
EU-A180	n-octyl	36	$<10^{-10}$	Pure MOP Ag; pure DOP antag

Compound Name	Structure	MOP agonist EC ₅₀ (nM)	DOP antagonist IC ₅₀ (M)	Characterization
EU-A189	Me	41	NT	Pure MOP Ag; pure DOP antag
EU-A190	n-dodecyl	§	NT	Pure MOP Ag; pure DOP antag
EU-A600	Me	110	<10E-10	Pure MOP Ag; pure DOP antag
EU-A601	n-octyl	§	<10E-10	Pure MOP Ag; pure DOP antag
EU-A615	Me	410	3.1E-10	Pure MOP Ag; pure DOP antag
EU-A620	Me	480	<10E-10	Pure MOP Ag; pure DOP antag
EU-A639	Me	280	<10E-10	Pure MOP Ag; pure DOP antag
§ indicates solubility issues during dissolution, NT means not tested, nc means not calculable				

Example 13: In vivo assay of the compounds

[0553] λ -Carrageenan-induced Inflammatory Pain Model: Adult male Sprague-Dawley rats were briefly anesthetized with isoflurane, and unilateral hind paw inflammatory pain was induced by injecting 3.5 g/100 μ l of λ -carrageenan. A λ -carrageenan injection into the left hind paw produces redness and edema. These alterations were observed within 6 h after induction of inflammation and peaked by 24 h. Further, the changes were confined to the carrageenan treated paw. Paired t test showed a significant increase in paw volume ($p < 0.0001$) and decrease in paw pressure threshold ($p < 0.001$) following carrageenan treatment.

[0554] Changes in Paw Pressure Thresholds (PPT), a measurement of nociception and antinociception, was evaluated using a digital Randall-Selitto method. Briefly, animals are gently restrained and incremental pressure (maximum 350 g) was applied onto the dorsal surface of the hindpaw. The pressure required to elicit paw withdrawal, PPT, was determined. Paw-thickness, the indicator of inflammation, was measured using digital calipers. 24 h after induction of inflammation, rats received either saline, positive controls (DAMGO (8-32 μ g) or fentanyl (0.4-0.8 μ g)) or test peptides in 100 μ l volume i.p.l. The antinociception was measured at different time points (5 to 120 min or 5hr to 24 hr) after drug administration. Baseline measures were taken before administering drug to confirm the development of inflammation and nociception. Eu-A178, for example, showed statistically significant increases in PPT beyond 30 min.

Example 14: Uses of the compounds

[0555] The covalently modified peptides and/or proteins described herein are useful for the prevention and treatment of a variety of diseases depending on which class is being considered. For example, covalently modified peptides and/or proteins described herein are indicated for the prophylaxis and therapeutic treatment of chronic and acute pain and other MOR- or DOR-related disease states. Applications for sunburn, pruritus, cancer, immune function, inflammation, cardiovascular disease also have been documented (Lazarus, L.H., et al. (2012) Expert Opin Ther Patents 22: 1-14).

[0556] Representative delivery regimens include oral, parenteral (including subcutaneous, intramuscular and intravenous injection), rectal, buccal (including sublingual), transdermal, inhalation ocular and intranasal. An attractive and widely used method for delivery of peptides entails subcutaneous injection of a controlled release injectable formulation. Other administration routes for the application of the covalently modified peptides and/or proteins described herein are oral, subcutaneous, intranasal and inhalation administration.

Example 15. Pharmaceutical usage for treatment of pain.

[0557] A human patient with acute or chronic pain is treated with EU-A178 by intranasal administration (200µL) from a standard atomizer used in the art of a solution of the pharmaceutical agent in physiological saline containing from 0.5 to 10 mg/mL of the pharmaceutical agent and containing standard excipients such as benzyl alcohol. The treatment is repeated as necessary for the alleviation of pain. Alternatively a solution of EU-A178, and selected excipients, in an evaporating solvent containing such as a hydrofluoroalkane is administered intranasally by MDI as needed to alleviate pain. Alternatively an aqueous solution of EU-A178, with selected excipients, is administered by subcutaneous injection as needed to alleviate pain.

[0558] The effect of treatment is determined from evaluation of patients, including quality of life questionnaires. Pain scales are based on self-report, observational (behavioral), or physiological data. Some pain scales suitable for use in clinical setting include Alder Hey Triage Pain Score, Brief Pain Inventory (BPI), Dallas Pain Questionnaire, Dolorimeter Pain Index (DPI), McGill Pain Questionnaire (MPQ), Numerical 11 point box (BS-11), Numeric Rating Scale (NRS-11), Roland-Morris Back Pain Questionnaire, Visual analog scale (VAS) or the like.

[0559] In a similar manner, administration of an adjusted amount by transbuccal, intravaginal, inhalation, subcutaneous, intravenous, intraocular, or oral routes is tested to determine relief from pain.

Example 2-1: Reagents - N- α -Fmoc, N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine

[0560] In an oven-dried 250 mL Erlenmeyer flask is placed 1-octyl β -D-glucuronic acid (Carbosynth Ltd., 3.06 g, 10 mmol), 50 mL anhydrous DMF, and anhydrous 1-hydroxybenzotriazole (1.62 g, 12 mmol). A chilled (4°C) solution of N, N'-dicyclohexylcarbodiimide (2.48g, 12 mmol) in 50 mL of DMF is added, with stirring, and the reaction is allowed to proceed for 5 min. The copious white precipitate of N, N'-dicyclohexylurea is filtered on a fritted glass funnel and the filtrate is added to a solution of N- α -Fmoc-L-lysine (3.68g, 10 mmol) in 25 ml anhydrous DMF. The reaction is allowed to proceed for 25 min with warming to room temp or until the ninhydrin color is very faint. The reaction mixture is filtered, stripped to dryness and crystallized from MeOH/Et₂O by dissolution in MeOH and slow dilution to the cloud point with Et₂O, followed by refrigeration. Further purification can be achieved by silica gel chromatography using a solvent gradient from EtOAc to EtOAc/EtOH/AcOH.

[0561] In a similar manner, but substituting N- α -Boc-L-lysine is obtained N- α -Boc,N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine, suitable for N-terminal incorporation and cleavage to a free N-Terminus. In a similar manner, but substituting N- α -Ac-L-lysine is obtained N- α -Ac, N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine, suitable for incorporation at the N-terminus of a peptide with a blocked N-terminus. In a similar manner, but substituting the appropriate amount of N- α -Fmoc-L-ornithine is obtained N- α -Fmoc,N- δ -(1-octyl β -D-glucuronide-6-yl)-L-ornithine. In a similar manner but substituting other N-mono-protected diamino acids one obtains the corresponding reagents. Alternatively, use of a transient Me₃Si ester protecting group during the coupling and without preactivation of the 1-octyl β -D-glucuronic acid provides a facile route to the formation of the reagents. The transient Me₃Si ester is produced by reaction of the Fmoc-Lys-OH with an equimolar amount of N,O-bis(trimethylsilyl)acetamide in dichloromethane (CH₂Cl₂). The organic layer contains the desired reagent as a solution in CH₂Cl₂ ready for coupling with the 1-alkyl glucuronide as above. The filtered reaction mixture is washed with aqueous NaHSO₄ to hydrolyze the Me₃Si ester, dried over MgSO₄ and solvent is removed.

[0562] Similarly, but using peracetyl or perbenzoyl 1-octyl β -D-glucuronic acid one obtains the Ac, or Bz protected form of the reagents (e.g. 2,3,4-trisacetyl 1-octyl β -D-glucuronic acid, and the like, formed by treatment with Ac₂O). Such reagents have increased stability during acid cleavage from the resin and are used when instability during deprotection is detected, see (Kihlberg, J., et al. (1997) *Methods Enzymol* 289: 221-245) and references therein. Final deprotection of such products is carried out by base-catalyzed transesterification after cleavage, by use of MeOH/NH₃, MeOH/NaOMe, MeOH/NH₂NH₂, as described above.

Example 2-2: Synthetic Peptide Analogs

[0563] In general, peptide synthesis methods involve the sequential addition of protected

amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid and any reactive side chain group are protected. This protected amino acid is then either attached to an inert solid support, or utilized in solution, and the next amino acid in the sequence, also suitably protected, is added under conditions amenable to formation of the amide linkage. After all the desired amino acids have been linked in the proper sequence, protecting groups and any solid support are removed to afford the crude peptide. The peptide is desalted and purified chromatographically.

[0564] A preferred method of preparing the analogs of the physiologically active truncated peptides, having fewer than about fifty amino acids, involves solid phase peptide synthesis. In this method the α -amino ($N\alpha$) functions and any reactive side chains are protected by acid- or base-sensitive groups. The protecting group should be stable to the conditions of peptide linkage formation, while being readily removable without affecting the extant peptide chain. Suitable α -amino protecting groups include, but are not limited to t-butyloxycarbonyl (Boc), benzyloxycarbonyl (Cbz), o-chlorobenzyloxycarbonyl, biphenylisopropylloxycarbonyl, t-amyloxycarbonyl (Amoc), isobornyloxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, o-nitrophenylsulfenyl, 2-cyano-t-butoxycarbonyl, 9-fluorenyl-methoxycarbonyl (Fmoc) and the like, preferably Boc or more preferably, Fmoc. Suitable side chain protecting groups include, but are not limited to: acetyl, benzyl (Bzl), benzyloxymethyl (Bom), Boc, t-butyl, o-bromobenzyloxycarbonyl, t-butyl, t-butylidimethylsilyl, 2-chlorobenzyl (Cl-z), 2,6-dichlorobenzyl, cyclohexyl, cyclopentyl, isopropyl, pivalyl, tetrahydropyran-2-yl, tosyl (Tos), 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf), trimethylsilyl and trityl. A preferred $N\alpha$ -protecting group for synthesis of the compounds is the Fmoc group. Preferred side chain protecting groups are O-t-Butyl group for Glu, Tyr, Thr, Asp and Ser; Boc group for Lys and Trp side chains; Pbf group for Arg; Trt group for Asn, Gln, and His. For selective modification of a Lys residue, orthogonal protection with a protecting group not removed by reagents that cleave the Fmoc or t-butyl based protecting groups is preferred. Preferred examples for modification of the Lys side chain include, but are not limited to, those removed by hydrazine but not piperidine; for example 1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)-3-methylbutyl (ivDde) or 1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethyl (Dde) and allyloxycarbonyl (Alloc).

[0565] The Fmoc-Lys(ivDde) or Fmoc-Lys(Dde) protecting group scheme is preferred in cases where side chain lactam formation is desired (Houston, M.E., Jr., et al. (1995) J Pept Sci 1: 274-282; Murage, E.N., et al. (2010) J Med Chem), since in this case Fmoc-Glu(O-Allyl) and Fmoc-Lys(Alloc) can be incorporated and used to provide transient protection, then deprotected for lactam formation while the Lys(Dde) protecting group remains for later removal and reaction with the functionalized surfactant. The side chain lactam between acidic and basic residue (e.g. Glu and Lys) is carried out after removal of the allyl-based protection by activation of the carboxyl side chain function with N,N'-diisopropylcarbodiimide (DIC)/1-hydroxybenzotriazole (HOBt) or 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate (HBTU)/ N,N-di-isopropylethylamine (DIEA), using standard protocols well known to the art.

[0566] In solid phase synthesis, the C-terminal amino acid is first attached to a suitable resin

support. Suitable resin supports are those materials which are inert to the reagents and reaction conditions of the stepwise condensation and deprotection reactions, as well as being insoluble in the media used. Examples of commercially available resins include styrene/divinylbenzene resins modified with a reactive group, e.g., chloromethylated co-poly-(styrene-divinylbenzene), hydroxymethylated co-poly-(styrene-divinylbenzene), and the like. Benzylated, hydroxymethylated phenylacetamidomethyl (PAM) resin is preferred for the preparation of peptide acids. When the C-terminus of the compound is an amide, a preferred resin is p-methylbenzhydrylamino-co-poly(styrene-divinylbenzene) resin, a 2,4-dimethoxybenzhydrylamino-based resin ("Rink amide"), 4-Hydroxymethylphenoxyacetyl aminomethyl resin (HMP Am) and the like. An especially preferred support for the synthesis of larger peptides are commercially available resins containing PEG sequences grafted onto other polymeric matrices, such as the Rink Amide-PEG and PAL-PEG-PS resins (Applied Biosystems) or similar resins designed for peptide amide synthesis using the Fmoc protocol. Thus in certain cases it is desirable to have an amide linkage to a PEG chain. In those cases it is convenient to link an N-Fmoc-amino-PEG-carboxylic acid to the amide forming resin above (e.g. Rink amide resin and the like). The first amino acid of the chain can be coupled as an N-Fmoc-amino acid to the amino function of the PEG chain. Final deprotection will yield the desired Peptide-NH-PEG-CO-NH₂ product.

[0567] Attachment to the PAM resin may be accomplished by reaction of the Na⁺ protected amino acid, for example the Boc-amino acid, as its ammonium, cesium, triethylammonium, 1,5-diazabicyclo-[5.4.0]undec-5-ene, tetramethylammonium, or similar salt in ethanol, acetonitrile, N,N-dimethylformamide (DMF), and the like, preferably the cesium salt in DMF, with the resin at an elevated temperature, for example between about 40° and 60 °C, preferably about 50 °C, for from about 12 to 72 hours, preferably about 48 hours. This will eventually yield the peptide acid product following acid cleavage or an amide following aminolysis.

[0568] The Na-Boc-amino acid may be attached to the benzhydrylamine or HPM Am resin by means of, for example, an N,N'-diisopropylcarbodiimide (DIC)/1-hydroxybenzotriazole (HOBt) mediated coupling for from about 2 to about 24 hours, preferably about 2 hours at a temperature of between about 10° and 50 °C, preferably 25 °C in a solvent such as CH₂Cl₂ or DMF, preferably CH₂Cl₂.

[0569] For Boc-based protocols, the successive coupling of protected amino acids may be carried out by methods well known in the art, typically in an automated peptide synthesizer. Following neutralization with triethylamine, N,N-diisopropylethylamine (DIEA), N-methylmorpholine (NMM), collidine, or similar base, each protected amino acid is introduced in approximately about 1.5 to 2.5 fold molar excess and the coupling carried out in an inert, nonaqueous, polar solvent such as CH₂Cl₂, DMF, N-methylpyrrolidone (NMP), N,N-dimethylacetamide (DMA), or mixtures thereof, preferably in dichloromethane at ambient temperature. For Fmoc-based protocols no acid is used for deprotection but a base, preferably DIEA or NMM, is usually incorporated into the coupling mixture. Couplings are typically done in DMF, NMP, DMA or mixed solvents, preferably DMF. Representative coupling agents are N,N'-

dicyclohexylcarbodiimide (DCC), N,N'-diisopropyl-carbodiimide (DIC) or other carbodiimide, either alone or in the presence of HOBt, O-acyl ureas, benzotriazol-1-yl-oxytris(pyrrolidino)phosphonium hexafluorophosphate (PyBop), N-hydroxysuccinimide, other N-hydroxyimides, or oximes. Alternatively, protected amino acid active esters (e.g. p-nitrophenyl, pentafluorophenyl and the like) or symmetrical anhydrides may be used. Preferred coupling agents are of the aminium/uronium (alternative nomenclatures used by suppliers) class such as 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate (HBTU), O-(7-azabenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU), 2-(6-Chloro-1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate (HCTU), and the like.

[0570] A preferred method of attachment to the Fmoc-PAL-PEG-PS resin may be accomplished by deprotection of the resin linker with 20% piperidine in DMF, followed by reaction of the N- α -Fmoc protected amino acid, about a 5 fold molar excess of the N- α -Fmoc-amino acid, using HBTU: di-isopropylethylamine (DIEA) (1:2) in DMF in a microwave-assisted peptide synthesizer with a 5min, 75° max coupling cycle.

[0571] For this Fmoc-based protocol in the microwave-assisted peptide synthesizer, the N- α -Fmoc amino acid protecting groups are removed with 20% piperidine in DMF containing 0.1M 1-hydroxybenzotriazole (HOBt), in a double deprotection protocol for 30 sec and then for 3min with a temperature maximum set at 75°C. HOBt is added to the deprotection solution to reduce aspartimide formation. Coupling of the next amino acid then employs a five-fold molar excess using HBTU:DIEA (1:2) with a 5min, 75° max. double-coupling cycle.

[0572] At the end of the solid phase synthesis the fully protected peptide is removed from the resin. When the linkage to the resin support is of the benzyl ester type, cleavage may be effected by means of aminolysis with an alkylamine or fluoroalkylamine for peptides with an alkylamide C-terminus, or by ammonolysis with, for example, ammonia/methanol or ammonia/ethanol for peptides with an unsubstituted amide C-terminus, at a temperature between about -10° and 50° C, preferably about 25° C, for between about 12 and 24 hours, preferably about 18 hours. Peptides with a hydroxy C-terminus may be cleaved by HF or other strongly acidic deprotection regimen or by saponification. Alternatively, the peptide may be removed from the resin by transesterification, e.g., with methanol, followed by aminolysis or saponification. The protected peptide may be purified by silica gel or reverse-phase HPLC.

[0573] The side chain protecting groups may be removed from the peptide by treating the aminolysis product with, for example, anhydrous liquid hydrogen fluoride in the presence of anisole or other carbonium ion scavenger, treatment with hydrogen fluoride/pyridine complex, treatment with tris(trifluoroacetyl)boron and trifluoroacetic acid, by reduction with hydrogen and palladium on carbon or polyvinylpyrrolidone, or by reduction with sodium in liquid ammonia, preferably with liquid hydrogen fluoride and anisole at a temperature between about - 10° and +10° C, preferably at about 0°C, for between about 15 minutes and 2 hours, preferably about 1.5 hours.

[0574] For peptides on the benzhydrylamine type resins, the resin cleavage and deprotection steps may be combined in a single step utilizing liquid hydrogen fluoride and anisole as described above or preferably through the use of milder cleavage cocktails. For example, for the PAL-PEG-PS resin, a preferred method is through the use of a double deprotection protocol in the microwave-assisted peptide synthesizer using one of the mild cleavage cocktails known in the art, such as TFA/water/tri-iso-propylsilane/3,6-dioxa-1,8-octanedithiol (DODT) (92.5/2.5/2.5/2.5) for 18min at 38°C each time. Cleavage of alkyl glycoside containing materials have shown survival of the alkyl glycoside linkage using protocols with TFA/water ratios in the 9/1 to 19/1 range. A typical cocktail is 94% TFA; 2% EDT; 2% H₂O; 2% TIS. Typically the fully deprotected product is precipitated and washed with cold (-70° to 4°C) Et₂O, dissolved in deionized water and lyophilized.

[0575] The peptide solution may be desalted (e.g. with BioRad AG-3® anion exchange resin) and the peptide purified by a sequence of chromatographic steps employing any or all of the following types: ion exchange on a weakly basic resin in the acetate form; hydrophobic adsorption chromatography on underivatized co-poly(styrene-divinylbenzene), e.g. Amberlite® XAD; silica gel adsorption chromatography; ion exchange chromatography on carboxymethylcellulose; partition chromatography, e.g. on Sephadex® G-25; counter-current distribution; supercritical fluid chromatography; or HPLC, especially reversed-phase HPLC on octyl- or octadecylsilylsilica (ODS) bonded phase column packing.

[0576] Also provided herein are processes for preparing covalently modified peptides and/or proteins described herein and pharmaceutically acceptable salts thereof, which processes comprise sequentially condensing protected amino acids on a suitable resin support, removing the protecting groups and resin support, and purifying the product, to afford analogs of the physiologically active truncated homologs and analogs of the covalently modified peptides and/or proteins described herein. In some embodiments, covalently modified peptides and/or proteins described herein incorporate alkyl glycoside modifications as defined above.

[0577] Another aspect relates to processes for preparing covalently modified peptides and/or proteins described herein and pharmaceutically acceptable salts thereof, which processes comprise the use of microwave-assisted solid phase synthesis-based processes or standard peptide synthesis protocols to sequentially condense protected amino acids on a suitable resin support, removing the protecting groups and resin support, and purifying the product, to afford analogs of the physiologically active peptides, as defined above.

Example 2-3. General oxidation method for uronic acids

[0578] To a solution of 1-dodecyl β-D-glucopyranoside (Carbosynth) [2.0g, 5.74 mmol] in 20 mL of acetonitrile and 20 mL of DI water was added (diacetoxyiodo)benzene (Fluka) [4.4 g, 13.7 mmol] and TEMPO (SigmaAldrich) [0.180g, 1.15 mmol]. The resulting mixture was stirred at room temperature for 20 h. The reaction was followed by mass spectrometry (for example

LCQ ESI) and upon completion, the reaction mixture was diluted with water and lyophilized to dryness to give 1.52 g (crude yield 73.1%) of the crude product, 1-dodecyl β -D-glucuronic acid, as a white powder, which was used directly for the solid phase synthesis without further purification. This product was previously prepared by an alternative process using NaOCl as oxidant, as described in the specification, and also has been used for longer alkyl groups. In a like manner, but using the corresponding 1-tetradecyl, 1-hexadecyl, and 1-octadecyl β -D-glucopyranosides (purchased from Anatrace, Maumee, OH) were prepared the desired alkyl saccharide uronic acids used to make the products and reagents described herein. For longer alkyl groups 1,4-dioxane was used instead of acetonitrile and the temperature was raised as high as 30°C.

[0579] In a like manner, but using, for example, the corresponding 1-octyl, 1-decyl, 1-undecyl, 1-tetradecyl, 1-hexadecyl, and 1-octadecyl glycosides (purchased from Anatrace, Maumee, OH) were prepared the desired 1-alkyl saccharide uronic acids which were used to make the products and reagents described herein. In a like manner, but using, for example, the corresponding 1-octyl, 1-decyl, 1-undecyl, 1-tetradecyl, 1-hexadecyl, and 1-octadecyl β -D-melibiosides or β -D-maltosides (purchased from Anatrace, Maumee, OH) were prepared the desired 1-alkyl disaccharide uronic acids which were used to make the products and reagents described herein. In each case the corresponding alpha glycosides can be treated similarly.

Example 2-4: Preparation of PTHrP analog EU-204.

[0580] A sample of Fmoc- Ac5c-Val-Aib-Glu-Ile-Gln-Leu-Nle-His-Gln-Arg-Ala-Arg-Trp-Ile-Gln-Lys(Alloc)-Rink amide resin was deprotected on the Lys-N-epsilon position by incubation with Pd(PPh₃)₄ (0.5 eq) and DMBA (20 eq) in DMF/ CH₂Cl₂ (1:1) overnight in the dark at room temperature. Following washing by DMF/ CH₂Cl₂, the Lys side chain was acylated with 1'-octyl dodecyl β -D-glucuronic acid (Carbosynth) in DMF/ CH₂Cl₂ through the use of DIC/HOBt. Completion of the coupling was checked by ninhydrin and the product was washed extensively with CH₂Cl₂.

[0581] The product resin was submitted to final deprotection and cleavage from the resin by treatment with the cleavage cocktail (94% TFA; 2% EDT; 2% H₂O; 2% TIS) for a period of 240 min at room temperature. The mixture was treated with Et₂O, to precipitate the product and washed extensively with Et₂O to yield the crude title peptide product after drying in vacuo.

[0582] Purification was carried out in two batches by reversed phase (C18) hplc. The crude peptide was loaded on a 4.1x25 cm hplc column at a flow rate of 15 mL/min (15% organic modifier; acetic acid buffer) and eluted with a gradient from 15-45% buffer B in 60min at 50°C. The product fraction was lyophilized to yield the title product peptide with a purity >97% by analytical hplc(12.0 min; 35-65% CH₃CN in 0.1% TFA)/mass spectrometry (M+1 peak = 2473.9).

[0583] The corresponding 1-methyl, 1-octyl, 1-decyl, 1-dodecyl, 1-tetradecyl, 1-hexadecyl, 1-octadecyl and 1-eicosyl analogs are prepared using the corresponding glucuronic acids, prepared as described above. Alternatively, the 1-alkyl glucuronyl, or other uronic acylated analogs, may be prepared by initial purification of the deprotected or partially deprotected peptide followed by acylation by the desired uronic acid reagent.

[0584] Analysis was done by HPLC/mass spectrometry in positive ion mode using the eluent gradients given in the table below.

Compound Name	Molecular Wt Expected	Molecular Wt Found	HPLC (min; elution)
EU-201	2278.63	2278.14	14.1 [a]
EU-202	2376.86	2376.80	11.2 [b]
EU-203	2432.97	2432.40	14.1 [c]
EU-204	2472.00	2471.86	12.0 [d]
EU-205	2343.87	2344.26	8.0 [e]
EU-207	2557.12	2557.06	13.2 [c]
EU-232	2642.23	2642.14	12.6 [c]
EU-251	2941.56	2942.26	13.1 [c]
EU-260	2967.61	2966.66	13.9 [c]
EU-283	2881.47	2882.26	11.0 [c]
EU-284	3640.39	3640.00	11.9 [c]
EU-286	2670.47	2669.86	6.3 [e]
EU-287	2698.47	2697.74	8.4 [e]
EU-288	2726.47	2726.26	10.9 [f]
HPLC gradients in 0.1% TFA [a.] 20 to 50% CH ₃ CN over 30 min. [b.] 25 to 55% CH ₃ CN over 20 min. [c.] 30 to 60% CH ₃ CN over 20 min. [d.] 35 to 65% CH ₃ CN over 20 min. [e.] 40 to 70% CH ₃ CN over 20 min. [f.] 45 to 75% CH ₃ CN over 20 min. HPLC on Phenomenex Luna C18 5micron 250x4.6 mm.			

Example 2-5: Cellular assay of the compounds.

[0585] Compounds were weighed precisely in an amount of approximately 1 mg and assayed in standard cellular assays (cerep SA). The readout is the amount of cAMP generated in the cells treated with the test compounds, in either agonist or antagonist mode. The PTH1 cellular

assay used is described in Orloff, J.J., et al. (1992) Endocrinol 131: 1603-1611.

[0586] For the series of compounds EU-201 to EU-203, where the hydrophobic portion of the surfactant (1-alkyl glucuronic acid) varies in length from C1 to C12, the cellular response increases in potency and efficacy with the increased chain length. All of the analogs were agonists. Further substitutions led to molecules with an EC₅₀ similar to PTH1-34, but with super-agonistic activity (e.g. EU-232) and such molecules have important applications in medicine. Additional analogs are designed to have very prolonged duration of action in vivo (that is EU-286, EU-287, and EU-288). In this assay, PTHrP (coded sample) had an EC₅₀ of 2.9nM and a maximal response of 100% while the internal standard, PTH had EC₅₀ of 1.4nM and maximal response of 105%. Compounds were dissolved in water and diluted in assay buffer containing 1% bovine serum albumin. The following table shows potency and efficacy of certain peptide products described herein.

Compound Name	Structure	EC50 (nM)	Maximal response (% PTHrP)	Characterization
EU-201	1-Me	40	115	agonist
EU-202	1-octyl	25	118	agonist
EU-203	1-dodecyl	25	129	agonist
EU-204	1-dodecyl	20	135	agonist
EU-205	1-dodecyl	40	115	agonist
EU-207	1-dodecyl	40	110	agonist
EU-232	1-dodecyl	2.4	145	agonist
EU-251	1-dodecyl	4.3	135	agonist
EU-260	1-dodecyl	2.2	120	agonist
EU-283	1-dodecyl	5.2	100	agonist
EU-284	1-dodecyl	62	110	agonist
EU-287	1-hexadecyl	<1	128	agonist
EU-288	1-octadecyl	<1	130	agonist

[0587] When tested in antagonist mode, with added PTHrP, the maximal effects were even greater (to 146% of PTHrP maximal for the C-12 compound, EU-203). This behavior demonstrates the ability of the modification described herein to vary the fundamental properties of the receptor interactions. Modifications elsewhere in the molecule can be used to further modify the potency and character of the interaction of the drug candidates.

Example 2-6: In vivo assay of the compounds.

[0588] Following the method of Frolik, C.A., et al. (2003) Bone 33: 372-379, 20 male rats from Sino-British SIPPR /BK Lab Animal Ltd were acclimated to standard laboratory conditions for a period of 7 days. After acclimation, the animals were sorted by age into groups of 5. Each animal in a group was treated with a single sc injection of either vehicle or test agent.

[0589] Animals in two test groups were treated with 80mcg/animal of huPTH1-34 (Bachem) or 80mcg/animal of EU-232. A fourth group was treated with EU-232 at 320mcg/animal. Blood samples were collected via retro-orbital vein at 0.5, 1, 2, 4, and 5 hrs. post-injection and blood samples were stored on ice prior to centrifugation and testing for blood PO₄ and Ca levels.

[0590] In the 80mcg groups (PTH and EU-232) there was a transient but not statistically significant decrease in blood PO₄ levels in response to PTH or EU-232 and PO₄ levels did not further diminish after 1 hour. In response to treatment with EU-232, the blood PO₄ levels decreased to statistically significantly lower levels with time and the maximal decrease (25-35% decrease from time 0 hr. level) was seen at the 5 hr. time point indicating a potent and prolonged duration of action for EU-232. No groups showed a statistically different blood Ca level compared to vehicle at any time point, thus there was no indication of a propensity for hypercalcemia following dosing.

[0591] In a similar manner, the analogs described herein (including compounds of Table 1 in Figure 1) are tested to evaluate their potency and duration of action in vivo.

[0592] The covalently modified peptides and/or proteins described herein are useful for the prevention and treatment of a variety of diseases. PTHR1 agonists are effective in the treatment of bone density diseases such as postmenopausal or senile osteoporosis, hypoparathyroidism, osteopenia, implant fixation, and certain metastatic tumors. Antagonistic analogs are suitable for treatment of hypercalcemia, especially as related to hyperparathyroidism or hypercalcemia of malignancy. PTH and PTHrP agonists can be used to mobilize proliferation of haematopoietic stem cells (HSC) in bone marrow in vivo or in vitro for use in bone marrow transplant and in disease syndromes related to low blood cell concentrations. Expansion post-transplant is an attractive application as well. Since many cells in the blood originate from HSCs, a wide range of applications is possible. Suitably labeled surfactant modified peptides can be used as diagnostic probes.

[0593] Representative delivery regimens include oral, parenteral (including subcutaneous, intramuscular and intravenous injection), rectal, buccal (including sublingual), transdermal, inhalation ocular and intranasal. An attractive and widely used method for delivery of peptides entails subcutaneous injection of a controlled release injectable formulation. Other administration routes for the application of the covalently modified peptides and/or proteins described herein are subcutaneous, intranasal and inhalation administration.

Example 2-7. Pharmaceutical usage for treatment of osteoporosis.

[0594] A human patient, with evidence of osteoporosis or osteopenia is treated with EU-204 by intranasal administration (200 μ L) from a standard atomizer used in the art of a solution of the pharmaceutical agent in physiological saline containing from 0.5 to 10 mg/mL of the pharmaceutical agent and containing standard excipients such as benzyl alcohol. The treatment is repeated as necessary for the alleviation of symptoms such as bone pain, osteopenia, low bone density, or fractures. In a similar manner, a solution of EU-204, and selected excipients, in an evaporating solvent containing such as a hydrofluoroalkane is administered intranasally by metered dose inhaler (MDI) as needed to stimulate bone accretion. The effect of treatment is measured by use of standard tests, including the Bone Mineral Density test (BMD test).

[0595] All of the compounds described in Table 1 of Figure 1 are tested using a similar protocol.

[0596] In a similar manner, administration of an adjusted amount by transbuccal, intravaginal, inhalation, subcutaneous, intravenous, intraocular, or oral routes is tested to determine the level of stimulation of PTHR1 on cells in the body, and to determine therapeutic effects.

Example 3-1: Reagents - N- α -Fmoc, N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine

[0597] In an oven-dried 250 mL Erlenmeyer flask is placed 1-octyl β -D-glucuronic acid (Carbosynth Ltd., 3.06 g, 10 mmol), 50 mL anhydrous DMF, and anhydrous 1-hydroxybenzotriazole (1.62 g, 12 mmol). A chilled (4°C) solution of N,N'-dicyclohexylcarbodiimide (2.48g, 12 mmol) in 50 mL of DMF is added, with stirring, and the reaction is allowed to proceed for 5 min. The copious white precipitate of N, N'-dicyclohexylurea is filtered on a fritted glass funnel and the filtrate is added to a solution of N- α -Fmoc-L-lysine (3.68g, 10 mmol) in 25 ml anhydrous DMF. The reaction is allowed to proceed for 25 min with warming to room temp or until the ninhydrin color is very faint. The reaction mixture is filtered, stripped to dryness and crystallized from MeOH/Et₂O by dissolution in MeOH and slow dilution to the cloud point with Et₂O, followed by refrigeration. Further purification can be achieved by silica gel chromatography using a solvent gradient from EtOAc to EtOAc/EtOH/AcOH.

[0598] In a similar manner, but substituting N- α -Boc-L-lysine is obtained N- α -Boc,N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine, suitable for N-terminal incorporation and cleavage to a free N-Terminus. In a similar manner, but substituting N- α -Ac-L-lysine is obtained N- α -Ac,N- ϵ -(1-octyl β -D-glucuronide-6-yl)-L-lysine, suitable for incorporation at the N-terminus of a peptide with a blocked N-terminus. In a similar manner, but substituting the appropriate amount of N- α -Fmoc-L-ornithine is obtained N- α -Fmoc,N- δ -(1-octyl β -D-glucuronide-6-yl)-L-ornithine. In a similar manner but substituting other N-mono-protected diamino acids one obtains the corresponding reagents. Alternatively, use of a transient Me₃Si ester protecting group during the coupling and without preactivation of the 1-octyl β -D-glucuronic acid provides a facile route to the formation

of the reagents. The transient Me₃Si ester is produced by reaction of the Fmoc-Lys-OH with an equimolar amount of N,O-bis(trimethylsilyl)acetamide in dichloromethane (CH₂Cl₂). The organic layer contains the desired reagent as a solution in CH₂Cl₂ ready for coupling with the 1-alkyl glucuronide as above. The filtered reaction mixture is washed with aqueous NaHSO₄ to hydrolyze the Me₃Si ester, dried over MgSO₄ and solvent is removed.

[0599] Similarly, but using peracetyl or perbenzoyl 1-octyl β-D-glucuronic acid one obtains the Ac, or Bz protected form of the reagents (e.g. 2,3,4-trisacetyl 1-octyl β-D-glucuronic acid, and the like, formed by treatment with Ac₂O). Such reagents have increased stability during acid cleavage from the resin and are used when instability during deprotection is detected, see (Kihlberg, J., et al. (1997) *Methods Enzymol* 289: 221-245) and references therein. Final deprotection of such products is carried out by base-catalyzed transesterification after cleavage, by use of MeOH/NH₃, MeOH/NaOMe, MeOH/NH₂NH₂, as described above.

Example 3-2: Synthetic Peptide Analogs

[0600] In general, peptide synthesis methods involve the sequential addition of protected amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid and any reactive side chain group are protected. This protected amino acid is then either attached to an inert solid support, or utilized in solution, and the next amino acid in the sequence, also suitably protected, is added under conditions amenable to formation of the amide linkage. After all the desired amino acids have been linked in the proper sequence, protecting groups and any solid support are removed to afford the crude peptide. The peptide is desalted and purified chromatographically.

[0601] A preferred method of preparing the analogs of the physiologically active truncated peptides, having fewer than about fifty amino acids, involves solid phase peptide synthesis. In this method the α-amino (Nα) functions and any reactive side chains are protected by acid- or base-sensitive groups. The protecting group should be stable to the conditions of peptide linkage formation, while being readily removable without affecting the extant peptide chain. Suitable α-amino protecting groups include, but are not limited to t-butyloxycarbonyl (Boc), benzyloxycarbonyl (Cbz), o-chlorobenzyloxycarbonyl, biphenylisopropylloxycarbonyl, t-amyloxycarbonyl (Amoc), isobornyloxycarbonyl, α,α-dimethyl-3,5-dimethoxybenzyloxycarbonyl, o-nitrophenylsulfonyl, 2-cyano-t-butyloxycarbonyl, 9-fluorenyl-methoxycarbonyl (Fmoc) and the like, preferably Boc or more preferably, Fmoc. Suitable side chain protecting groups include, but are not limited to: acetyl, benzyl (Bzl), benzyloxymethyl (Bom), Boc, t-butyl, o-bromobenzyloxycarbonyl, t-butyl, t-butyldimethylsilyl, 2-chlorobenzyl (Cl-z), 2,6-dichlorobenzyl, cyclohexyl, cyclopentyl, isopropyl, pivalyl, tetrahydropyran-2-yl, tosyl (Tos), 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf), trimethylsilyl and trityl. A preferred Nα-protecting group for synthesis of the compounds is the Fmoc group. Preferred side chain protecting groups are O-t-Butyl group for Glu, Tyr, Thr, Asp and Ser; Boc group for Lys and Trp side chains; Pbf group for Arg; Trt group for Asn, Gln, and His. For selective modification of a

Lys residue, orthogonal protection with a protecting group not removed by reagents that cleave the Fmoc or t-butyl based protecting groups is preferred. Preferred examples for modification of the Lys side chain include, but are not limited to, those removed by hydrazine but not piperidine; for example 1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)-3-methylbutyl (ivDde) or 1-(4,4-dimethyl-2,6-dioxocyclohex-1-ylidene)ethyl (Dde) and allyloxycarbonyl (Alloc). The Fmoc-Lys(ivDde) or Fmoc-Lys(Dde) protecting group scheme is preferred in cases where side chain lactam formation is desired (Houston, M.E., Jr., et al. (1995) *J Pept Sci* 1: 274-282; Murage, E.N., et al. (2010) *J Med Chem*), since in this case Fmoc-Glu(O-Allyl) and Fmoc-Lys(Alloc) can be incorporated and used to provide transient protection, then deprotected for lactam formation while the Lys(Dde) protecting group remains for later removal and reaction with the functionalized surfactant.

[0602] The Fmoc-Lys(ivDde) or Fmoc-Lys(Dde) protecting group scheme is preferred in cases where side chain lactam formation is desired (Houston, M.E., Jr., et al. (1995) *J Pept Sci* 1: 274-282; Murage, E.N., et al. (2010) *J Med Chem*), since in this case Fmoc-Glu(O-Allyl) and Fmoc-Lys(Alloc) can be incorporated and used to provide transient protection, then deprotected for lactam formation while the Lys(Dde) protecting group remains for later removal and reaction with the functionalized surfactant. The side chain lactam between acidic and basic residue (e.g. Glu and Lys) is carried out after removal of the allyl-based protection by activation of the carboxyl side chain function with N,N'-diisopropylcarbodiimide (DIC)/1-hydroxybenzotriazole (HOBt) or 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate (HBTU)/ N,N-di-isopropylethylamine (DIEA), using standard protocols well known to the art.

[0603] In solid phase synthesis, the C-terminal amino acid is first attached to a suitable resin support. Suitable resin supports are those materials which are inert to the reagents and reaction conditions of the stepwise condensation and deprotection reactions, as well as being insoluble in the media used. Examples of commercially available resins include styrene/divinylbenzene resins modified with a reactive group, e.g., chloromethylated co-poly-(styrene-divinylbenzene), hydroxymethylated co-poly-(styrene-divinylbenzene), and the like. Benzylated, hydroxymethylated phenylacetamidomethyl (PAM) resin is preferred for the preparation of peptide acids. When the C-terminus of the compound is an amide, a preferred resin is p-methylbenzhydrylamino-co-poly(styrene-divinylbenzene) resin, a 2,4 dimethoxybenzhydrylamino-based resin ("Rink amide"), and the like. An especially preferred support for the synthesis of larger peptides are commercially available resins containing PEG sequences grafted onto other polymeric matrices, such as the Rink Amide-PEG and PAL-PEG-PS resins (Applied Biosystems) or similar resins designed for peptide amide synthesis using the Fmoc protocol. Thus in certain cases it is desirable to have an amide linkage to a PEG chain. In those cases it is convenient to link an N-Fmoc-amino-PEG-carboxylic acid to the amide forming resin above (e.g. Rink amide resin and the like). The first amino acid of the chain can be coupled as an N-Fmoc-amino acid to the amino function of the PEG chain. Final deprotection will yield the desired Peptide-NH-PEG-CO-NH₂ product.

[0604] Attachment to the PAM resin may be accomplished by reaction of the Na protected

amino acid, for example the Boc-amino acid, as its ammonium, cesium, triethylammonium, 1,5-diazabicyclo-[5.4.0]undec-5-ene, tetramethylammonium, or similar salt in ethanol, acetonitrile, N,N-dimethylformamide (DMF), and the like, preferably the cesium salt in DMF, with the resin at an elevated temperature, for example between about 40° and 60 °C, preferably about 50 °C, for from about 12 to 72 hours, preferably about 48 hours. This will eventually yield the peptide acid product following acid cleavage or an amide following aminolysis.

[0605] The Na-Boc-amino acid may be attached to the benzhydrylamine or HMP Am resin by means of, for example, an N,N'-diisopropylcarbodiimide (DIC)/1-hydroxybenzotriazole (HOBt) mediated coupling for from about 2 to about 24 hours, preferably about 2 hours at a temperature of between about 10° and 50 °C, preferably 25 °C in a solvent such as CH₂Cl₂ or DMF, preferably CH₂Cl₂.

[0606] For Boc-based protocols, the successive coupling of protected amino acids may be carried out by methods well known in the art, typically in an automated peptide synthesizer. Following neutralization with triethylamine, N,N-di-isopropylethylamine (DIEA), N-methylmorpholine (NMM), collidine, or similar base, each protected amino acid is introduced in approximately about 1.5 to 2.5 fold molar excess and the coupling carried out in an inert, nonaqueous, polar solvent such as CH₂Cl₂, DMF, N-methylpyrrolidone (NMP), N,N-dimethylacetamide (DMA), or mixtures thereof, preferably in dichloromethane at ambient temperature. For Fmoc-based protocols no acid is used for deprotection but a base, preferably DIEA or NMM, is usually incorporated into the coupling mixture. Couplings are typically done in DMF, NMP, DMA or mixed solvents, preferably DMF. Representative coupling agents are N,N'-dicyclohexylcarbodiimide (DCC), N,N'-diisopropyl-carbodiimide (DIC) or other carbodiimide, either alone or in the presence of HOBt, O-acyl ureas, benzotriazol-1-yl-oxytris(pyrrolidino)phosphonium hexafluorophosphate (PyBop), N-hydroxysuccinimide, other N-hydroxyimides, or oximes. Alternatively, protected amino acid active esters (e.g. p-nitrophenyl, pentafluorophenyl and the like) or symmetrical anhydrides may be used. Preferred coupling agents are of the aminium/uronium (alternative nomenclatures used by suppliers) class such as 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate (HBTU), O-(7-azabenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU), 2-(6-Chloro-1H-benzotriazole-1-yl)-1,1,3,3-tetramethylaminium hexafluorophosphate (HCTU), and the like.

[0607] A preferred method of attachment to the Fmoc-PAL-PEG-PS resin may be accomplished by deprotection of the resin linker with 20% piperidine in DMF, followed by reaction of the N- α -Fmoc protected amino acid, about a 5 fold molar excess of the N- α -Fmoc-amino acid, using HBTU: di-isopropylethylamine (DIEA) (1:2) in DMF in a microwave-assisted peptide synthesizer with a 5min, 75° max coupling cycle.

[0608] For this Fmoc-based protocol in the microwave-assisted peptide synthesizer, the N- α -Fmoc amino acid protecting groups are removed with 20% piperidine in DMF containing 0.1M 1-hydroxybenzotriazole (HOBt), in a double deprotection protocol for 30 sec and then for 3min with a temperature maximum set at 75°C. HOBt is added to the deprotection solution to reduce

aspartimide formation. Coupling of the next amino acid then employs a five-fold molar excess using HBTU:DIEA (1:2) with a 5min, 75° max. double-coupling cycle.

[0609] At the end of the solid phase synthesis the fully protected peptide is removed from the resin. When the linkage to the resin support is of the benzyl ester type, cleavage may be effected by means of aminolysis with an alkylamine or fluoroalkylamine for peptides with an alkylamide C-terminus, or by ammonolysis with, for example, ammonia/methanol or ammonia/ethanol for peptides with an unsubstituted amide C-terminus, at a temperature between about -10° and 50° C, preferably about 25° C, for between about 12 and 24 hours, preferably about 18 hours. Peptides with a hydroxy C-terminus may be cleaved by HF or other strongly acidic deprotection regimen or by saponification. Alternatively, the peptide may be removed from the resin by transesterification, e.g., with methanol, followed by aminolysis or saponification. The protected peptide may be purified by silica gel or reverse-phase HPLC.

[0610] The side chain protecting groups may be removed from the peptide by treating the aminolysis product with, for example, anhydrous liquid hydrogen fluoride in the presence of anisole or other carbonium ion scavenger, treatment with hydrogen fluoride/pyridine complex, treatment with tris(trifluoroacetyl)boron and trifluoroacetic acid, by reduction with hydrogen and palladium on carbon or polyvinylpyrrolidone, or by reduction with sodium in liquid ammonia, preferably with liquid hydrogen fluoride and anisole at a temperature between about - 10° and +10° C, preferably at about 0°C, for between about 15 minutes and 2 hours, preferably about 1.5 hours.

[0611] For peptides on the benzhydrylamine type resins, the resin cleavage and deprotection steps may be combined in a single step utilizing liquid hydrogen fluoride and anisole as described above or preferably through the use of milder cleavage cocktails. For example, for the PAL-PEG-PS resin, a preferred method is through the use of a double deprotection protocol in the microwave-assisted peptide synthesizer using one of the mild cleavage cocktails known in the art, such as TFA/water/tri-iso-propylsilane/3,6-dioxo-1,8-octanedithiol (DODT) (92.5/2.5/2.5/2.5) for 18min at 38°C each time. Cleavage of alkyl glycoside containing materials have shown survival of the alkyl glycoside linkage using protocols with TFA/water ratios in the 9/1 to 19/1 range. A typical cocktail is 94% TFA; 2% EDT; 2% H₂O; 2% TIS. Typically the fully deprotected product is precipitated and washed with cold (-70° to 4°C) Et₂O, dissolved in deionized water and lyophilized.

[0612] The peptide solution may be desalted (e.g. with BioRad AG-3® anion exchange resin) and the peptide purified by a sequence of chromatographic steps employing any or all of the following types: ion exchange on a weakly basic resin in the acetate form; hydrophobic adsorption chromatography on underivatized co-poly(styrene-divinylbenzene), e.g. Amberlite® XAD; silica gel adsorption chromatography; ion exchange chromatography on carboxymethylcellulose; partition chromatography, e.g. on Sephadex® G-25; counter-current distribution; supercritical fluid chromatography; or HPLC, especially reversed-phase HPLC on octyl- or octadecylsilylsilica (ODS) bonded phase column packing.

[0613] Also provided herein are processes for preparing covalently modified peptides and/or proteins described herein and pharmaceutically acceptable salts thereof, which processes comprise sequentially condensing protected amino acids on a suitable resin support, removing the protecting groups and resin support, and purifying the product, to afford analogs of the physiologically active truncated homologs and analogs of the covalently modified peptides and/or proteins described herein. In some embodiments, covalently modified peptides and/or proteins described herein incorporate alkyl glycoside modifications as defined above. Another aspect relates to processes for preparing covalently modified peptides and/or proteins described herein and pharmaceutically acceptable salts thereof, which processes comprise the use of microwave-assisted solid phase synthesis-based processes or standard peptide synthesis protocols to sequentially condense protected amino acids on a suitable resin support, removing the protecting groups and resin support, and purifying the product, to afford analogs of the physiologically active peptides, as defined above.

Example 3-3. General oxidation method for uronic acids

[0614] To a solution of 1-dodecyl β -D-glucopyranoside (Carbosynth) [2.0g, 5.74 mmol] in 20 mL of acetonitrile and 20 mL of DI water was added (diacetoxyiodo)benzene (Fluka) [4.4 g, 13.7 mmol] and TEMPO (SigmaAldrich) [0.180g, 1.15 mmol]. The resulting mixture was stirred at room temperature for 20 h. The reaction was followed by mass spectrometry (for example LCQ ESI) and upon completion, the reaction mixture was diluted with water and lyophilized to dryness to give 1.52 g (crude yield 73.1%) of the crude product, 1-dodecyl β -D-glucuronic acid, as a white powder, which was used directly for the solid phase synthesis without further purification. This product was previously prepared by an alternative process using NaOCl as oxidant, as described in the specification, and also has been used for longer alkyl groups. In a similar manner are prepared the desired alkyl saccharide uronic acids used to make the products and reagents described herein.

[0615] In a like manner, but using the corresponding 1-tetradecyl, 1-hexadecyl, and 1-octadecyl β -D-glucopyranosides (purchased from Anatrace, Maumee, OH) were prepared the desired 1-alkyl saccharide uronic acids which were used to make the products and reagents described herein. For longer alkyl groups 1,4-dioxane was used instead of acetonitrile and the temperature was raised as high as 30°C.

[0616] In a like manner, but using, for example, the corresponding 1-octyl, 1-decyl, 1-undecyl, 1-tetradecyl, 1-hexadecyl, and 1-octadecyl glycosides (purchased from Anatrace, Maumee, OH) were prepared the desired 1-alkyl saccharide uronic acids which were used to make the products and reagents described herein. In a like manner, but using, for example, the corresponding 1-octyl, 1-decyl, 1-undecyl, 1-tetradecyl, 1-hexadecyl, and 1-octadecyl β -D-melibiosides or β -D-maltosides (purchased from Anatrace, Maumee, OH) were prepared the desired 1-alkyl disaccharide uronic acids which were used to make the products and reagents described herein. In each case the corresponding alpha glycosides can be treated similarly

Example 3-4: Preparation of analog EU-A387.

[0617] A sample of Fmoc-His-Aib-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Bip-Ser-Lys-Tyr-Leu-Glu-Ser-Lys(Alloc)-Rink amide resin was prepared by sequential addition of N-alpha-Fmoc protected amino acids as described in Example 1 and deprotected on the Lys-N-epsilon position by incubation with Pd(PPh₃)₄ (0.5 eq) and DMBA (20 eq) in DMF/ CH₂Cl₂ (1:1) overnight in the dark at room temperature. Following washing by DMF/ CH₂Cl₂, the Lys side chain was acylated with 1'-dodecyl β-D-glucuronic acid in DMF/ CH₂Cl₂ through the use of DIC/HOBt. Completion of the coupling was checked by ninhydrin and the product was washed extensively with CH₂Cl₂.

[0618] The product resin was submitted to final deprotection and cleavage from the resin by treatment with the cleavage cocktail (94% TFA; 2% EDT; 2% H₂O; 2% TIS) for a period of 240 min at room temperature. The mixture was treated with Et₂O, to precipitate the product and washed extensively with Et₂O to yield the crude title peptide product after drying in vacuo.

[0619] Purification was carried out in two batches by reversed phase (C18) hplc. The crude peptide was loaded on a 4.1x25 cm hplc column at a flow rate of 15 mL/min (15% organic modifier; acetic acid buffer) and eluted with a gradient from 15-45% buffer B in 60min at 50°C. The product fraction was lyophilized to yield the title product peptide with a purity 98.03% by analytical hplc (18.6 min; 30-60% CH₃CN in 0.1% TFA)/mass spectrometry (M+1 peak = 2382.14).

Example 3- 5: Preparation of analog EU-A1024.

[0620] A sample of Boc-His(Trt)-Aib-Gln(Trt)-Gly-Thr(tBu)-Phe-Thr(tBu)-Ser(tBu)-Asp(OtBu)-Tyr(tBu)-Ser(tBu)-Lys(Boc)-Tyr(tBu)-Leu-Asp(OtBu)-Glu(O-Allyl)-Gln(Trt)-Ala-Ala-Lys(Alloc)-Glu(O-tBu)-Phe-Ile-Lys(Dde)-Trp(Boc)-Leu-Leu-Gln(Trt)-Thr(tBu)-HMP amide resin (started from Fmoc-Thr(tBu)-HMP Am Resin, substitution 0.45mmol/g) was prepared by sequential addition of N-alpha-Fmoc protected amino acids as described in Example 1. The allyl based side chains on Glu and Lys were deprotected by incubation with Pd(PPh₃)₄ (0.5 eq) and DMBA (20 eq) in DMF/ CH₂Cl₂ (1:1) overnight in the dark at room temperature. The resin was washed with 0.5% DIEA in DMF (twice), 0.5% sodium diethyldithiocarbamate in DMF (twice) and DMF/ CH₂Cl₂ until a light yellow resin was obtained. The side chain lactam linkage was formed by coupling the Glu and Lys with DIC/HOBT (5 equivalents) in DMF. The reaction was checked for completeness with ninhydrin and recoupled if necessary. Following washing by DMF/CH₂Cl₂, the Lys side chain was deprotected by incubation with 5% hydrazine hydrate in DMF (10 equivalents) twice, in each case for 15 min. Following washing by DMF/ CH₂Cl₂ the side chain amino group of the deprotected Lys residue was reacted with 1'-tetradecyl β-D-

melibiouronic acid in DMF/ CH₂Cl₂ through the use of DIC/HOBt. Completion of the coupling was checked by ninhydrin and the product was washed extensively with CH₂Cl₂. Any couplings which were not complete by ninhydrin were rerun. In general, 10-12 g of peptide product resin was obtained from a 2mmole synthesis.

[0621] The product resin is submitted to final deprotection and cleavage from the resin by treatment with the cleavage cocktail (94% TFA; 2% EDT; 2% H₂O; 2% TIS) for a period of 240 min at room temperature. The mixture was treated with Et₂O, to precipitate the product and washed extensively with Et₂O to yield the crude title peptide product after drying in vacuo. In general, 5 to 8g crude product peptide was obtained.

[0622] Purification was carried out in two batches by reversed phase (C18) hplc. The crude peptide (1-1.5g) was loaded on a 4.1x25 cm hplc column at a flow rate of 15 mL/min (15% organic modifier; 0.1%TFA buffer) and eluted with a gradient from 35-55% buffer B in 70min at room temperature. Repurification of the less pure fractions was done for the fractions with a purity of >70%. The product fraction was lyophilized to yield the title product peptide with a purity 98.7% by analytical hplc (10.3 min; 45-75% CH₃CN in 0.1% TFA)/mass spectrometry (1317.67, +3 charged; 1976.13, +2 charged; molecular weight 3950.44). In a similar manner were prepared the other analogs of the invention, the characterization of which is illustrated below.

[0623] The corresponding 1-methyl and 1-octyl analogs of the title compound are prepared in a similar manner, but using the reagents 1'-methyl β -D-glucuronic acid and 1'-octyl β -D-glucuronic acid (Carbosynth). In a like manner, but using the corresponding 1-octyl, 1-decyl, 1-undecyl, 1-tetradecyl, 1-hexadecyl, and 1-octadecyl β -D-glucouronic acids (prepared as described above) were prepared the products of the invention. In a like manner, but using the corresponding 1-octyl, 1-decyl, 1-undecyl, 1-tetradecyl, 1-hexadecyl, and 1-octadecyl β -D-melibiouronic or β -D-maltouronic (prepared as described above) were prepared the products and reagents described herein. Alternatively, the 1-alkyl glucuronyl, or other uronic acylated analogs, may be prepared by initial purification of the deprotected or partially deprotected peptide followed by acylation by the desired uronic acid reagent.

[0624] Analysis and characterization was done by HPLC/mass spectrometry in positive ion mode using the eluent gradients given in the table below. Analysis and characterization was done by HPLC/mass spectrometry in positive ion mode using the eluent gradients given in the table below.

Compound Name	Molecular Wt Expected	Molecular Wt Found	HPLC (min; elution)
EU-A387	2379.66	2380.14	18.6[b]
EU-A388	2393.69	2393.74	16.0 [a]
EU-A391	2317.62	2318.26	11.2 [b]
EU-A455	2988.36	2988.00	11.5 [b]

Compound Name	Molecular Wt Expected	Molecular Wt Found	HPLC (min; elution)
EU-A474	2570.86	2570.54	11.3 [b]
EU-A478	2459.75	2459.74	11.1 [b]
EU-A484	2544.86	2545.06	9.6 [b]
EU-A501	2904.2	2903.34	7.9 [b]
EU-A502	2776.07	2776.14	8.0 [b]
EU-A503	2704.98	2704.40	8.0 [b]
EU-A504	2548.80	2548.00	9.1 [b]
EU-A505	2392.61	2392.40	10.5 [b]
EU-A506	2305.53	2305.06	10.7 [b]
EU-A507	3763.23	3762.66	9.0 [b]
EU-A521	2303.56	2303.60	8.2 [c]
EU-A522	2315.60	2315.60	14.2 [d]
EU-A523	2615.94	2616.00	8.1 [b]
EU-A524	2459.75	2459.74	12.7 [d]
EU-A525	2459.75	2459.06	6.0 [c]
EU-A526	2473.75	2473.60	12.7 [d]
EU-A527	2390.64	2390.40	14.6 [d]
EU-A529	2546.83	2546.80	9.5 [b]
EU-A531	2546.83	2546.80	9.5 [b]
EU-A532	2559.00	2558.66	9.6 [b]
EU-A533	2560.96	2560.66	9.5 [b]
EU-A534	2544.99	2544.94	9.7 [b]
EU-A535	2573.05	2574.00	12.0 [b]
EU-A536	2602.96	2603.46	14.3 [b]
EU-A538	2516.99	2516.40	10.3 [b]
EU-A539	2657.20	2656.80	10.8 [b]
EU-A540	2685.20	2684.94	9.8 [c]
EU-A541	2713.20	2712.80	13.0 [c]
EU-A544	2631.94	2632.26	10.8 [b]
EU-A546	2687.94	2688.80	9.1 [c]
EU-A549	2388.67	2388.66	6.3 [e]
EU-A551	2444.67	2445.20	11.4 [e]
EU-A552	2472.67	2473.14	10.7 [f]
EU-A554	2560.86	2560.40	10.3 [c]

Compound Name	Molecular Wt Expected	Molecular Wt Found	HPLC (min; elution)
EU-A556	2616.86	2616.40	11.7 [e]
EU-A557	2644.86	2645.74	10.4 [f]
EU-A560	2570.86	2571.06	8.3 [c]
EU-A562	2626.86	2626.66	9.9 [e]
EU-A563	2654.86	2655.06	8.7 [f]
EU-A565	2542.80	2542.54	9.5 [c]
EU-A567	2598.80	2599.06	12.0 [e]
EU-A568	2626.80	2626.54	10.1 [f]

HPLC gradients in 0.1% TFA

[a.] 35 to 65% CH₃CN over 30 min.

[b.] 30 to 60% CH₃CN over 20 min.

[c.] 35 to 65% CH₃CN over 20 min.

[d.] 25 to 55% CH₃CN over 20 min.

[e.] 40 to 70% CH₃CN over 20 min.

[f.] 45 to 75% CH₃CN over 20 min.

HPLC on Phenomenex Luna C18 5micron 250x4.6 mm.

Additional compounds synthesized and analyzed as described above are :

Compound Name	Molecular Wt Expected	Molecular Wt Found	HPLC (min; elution)
EU-A570	2656.16	2656.00	10.4 [b]
EU-A571	2684.16	2683.34	11.2 [c]
EU-A575	2670.16	2670.94	11.8 [b]
EU-A576	2698.16	2697.20	11.2 [c]
EU-A580	2668.20	2667.20	12.3 [b]
EU-A581	2696.20	2695.46	11.1 [c]
EU-A592	2724.20	2724.58	9.9 [e]
EU-A595	2682.20	2682.40	9.7 [c]
EU-A596	2710.20	2710.46	10.4 [c]
EU-A597	2738.20	2738.18	10.9 [e]
EU-A721	2461.85	2461.74	10.3 [b]
EU-A722	2475.85	2475.34	10.8 [b]
EU-A723	2459.88	2459.86	7.7 [c]
EU-A724	2473.88	2473.34	11.1 [b]
EU-A725	2471.92	2472.00	10.8 [b]
EU-A726	2557.03	2556.80	11.0 [b]
EU-A727	2485.92	2485.74	10.9 [b]

Compound Name	Molecular Wt Expected	Molecular Wt Found	HPLC (min; elution)
EU-A728	2513.92	2513.86	10.6 [c]
EU-A729	2541.92	2541.86	9.7 [e]
EU-A730	2569.92	2569.74	9.4 [f]
EU-A731	2425.88	2425.32	10.6 [d]
EU-A732	2476.95	2476.40	9.4 [c]
EU-A733	2381.83	2382.02	11.4 [b]
EU-A734	2616.09	2616.18	11.4 [b]
EU-A750	1611.89	1611.56	9.4 [c]
EU-A751	1625.89	1625.35	9.7 [c]
EU-A752	1709.93	1709.41	11.7 [g]
EU-A753	1637.84	1637.46	12.5 [b]
EU-A754	1651.84	1651.18	9.9 [c]
EU-A755	1711.93	1711.46	10.3 [h]
EU-A756	1671.98	1671.37	9.8 [e]
EU-A757	1770.02	1769.17	14.9 [g]
EU-A770	3333.61	3334.65	9.5 [b]
EU-A771	3678.25	3677.96	11.3 [c]
EU-A772	3762.25	3763.35	14.9 [e]
EU-A773	3790.31	3791.31	11.1 [f]
EU-A774	3475.74	3477.15	10.5 [d]
EU-A775	3820.38	3821.52	10.4 [c]
EU-A776	3904.38	3905.76	10.2 [f]
EU-A777	3932.44	3933.69	9.7 [f]
EU-A792	3793.43	3793.52	8.9 [f]
EU-A793	3821.43	3821.60	10.8 [f]
EU-A794	3849.43	3848.78	10.5 [g]
EU-A945	3777.18	3777.54	12.2 [e]
EU-A948	3861.18	3862.14	13.0 [f]
EU-A993	3759.22	3759.00	9.3 [f]
EU-A994	3787.22	3787.44	11.4 [f]
EU-A995	3815.22	3815.52	14.5 [g]
EU-A996	3843.22	3843.12	12.3 [g]
EU-A999	3935.35	3934.66	13.4 [e]
EU-A1011	3854.21	3854.38	14.2 [c]

Compound Name	Molecular Wt Expected	Molecular Wt Found	HPLC (min; elution)
EU-A1017	3896.26	3895.77	12.3 [c]
EU-A1023	3921.35	3921.15	10.5 [e]
EU-A1024	3950.44	3949.95	10.3 [f]
EU-A1026	4005.55	4004.64	12.8 [g]
EU-A1029	3939.31	3939.06	10.7 [e]
EU-A1032	4023.51	4025.22	13.9 [f]
EU-A1035	3840.21	3838.21	9.8 [e]
EU-A1041	3882.26	3880.50	9.6 [e]
EU-A1044	3966.46	3965.46	12.6 [f]
EU-A1167	3731.15	3731.42	8.9 [f]
EU-A1168	3745.15	3745.20	10.8 [i]
EU-A1173	3978.49	3977.00	12.9 [f]

HPLC gradients in 0.1% TFA
[a] 35 to 65% CH₃CN over 30 min.
[b] 30 to 60% CH₃CN over 20 min.
[c] 35 to 65% CH₃CN over 20 min.
[d] 25 to 55% CH₃CN over 20 min.
[e] 40 to 70% CH₃CN over 20 min.
[f] 45 to 75% CH₃CN over 20 min.
[g] 50 to 80% CH₃CN over 20 min.
[h] 10 to 40% CH₃CN over 20 min.
[i] 30 to 90% CH₃CN over 20 min.
HPLC on was carried out on a Phenomenex Luna C18 5micron 250x4.6 mm analytical column.

Example 3-6: Cellular assay of the compounds.

[0625] Compounds were weighed precisely in an amount of approximately 1 mg and assayed in standard cellular assays (Cerep SA). The readout is the amount of cAMP generated in the cells treated with the test compounds, in either agonist or antagonist mode. The assay used was the stimulation of cAMP levels in the glucagon (human, cloned into CHO cells) and GLP-1 (murine cell line) cellular assays. The assays are described in Chicchi, G.G., et al. (1997) J Biol Chem 272: 7765-7769 and Runge, S., et al. (2003) Br J Pharmacol 138: 787-794.

[0626] For compound EU-A391 the GLCR cellular response does not change and the GLP1R cellular response rises steeply with and EC50 of 420nM

Compound	Structure	EC ₅₀ GLP-1 R (nM)	EC ₅₀ murine glucagon R (nM)
EU-A391	1-dodecyl	420	n.c.
EU-A455	1-dodecyl	59	770
EU-A474	1-dodecyl	3000	n.c.
EU-A478	1-dodecyl	n.c.	n.c.
EU-A484	1-dodecyl	n.c.	n.c.
EU-A501	1-dodecyl	20000	12000
EU-A502	1-dodecyl	9400	n.c.
EU-A503	1-dodecyl	n.c.	n.c.
EU-A504	1-dodecyl	3100	1100
EU-A505	1-dodecyl	8500	6100
EU-A506	1-dodecyl	4600	1300
EU-A507	1-dodecyl	18	1
EU-A521	1-dodecyl	n.c.	n.c.
EU-A522	1-dodecyl	n.c.	9000
EU-A523	1-dodecyl	n.c.	n.c.
EU-A524	1-dodecyl	n.c.	n.c.
EU-A525	1-dodecyl	n.c.	n.c.
EU-A526	1-dodecyl	n.c.	n.c.
EU-A527	1-dodecyl	n.c.	5000
EU-A529	1-dodecyl	n.c.	7000
EU-A531	1-dodecyl	2100	1100
EU-A532	1-dodecyl	5000	2600
EU-A533	1-dodecyl	770	780
EU-A534	1-dodecyl	290	1900
EU-A535	1-tetradecyl	§4800	2100
EU-A536	1-hexadecyl	>10000	4400
EU-A538	1-dodecyl	270	n.c.
EU-A539	1-dodecyl	860	2300
EU-A540	1-tetradecyl	n.c.	8800
EU-A541	1-hexadecyl	800	5000
n.c. means EC50 not calculable § means superagonist			

A further series of cellular assays were carried out using standard cellular assays (DiscoverRx, LeadHunter assays) using readout of cAMP stimulation or arrestin activation. Compounds were

weighed precisely in an amount of approximately 1 mg and shipped to DiscoverX for dilution and assay. The assay used were for the glucagon (human, cloned into CHO cells) and GLP-1 (human, cloned into CHO cells) receptors in cellular assays.

Compound	EC ₅₀ cAMP GLP-1 R (nM)	EC ₅₀ arrestin GLP-1 R (nM)	EC ₅₀ cAMP glucagon R (nM)	EC ₅₀ arrestin glucagon R (nM)
EU-A507	0.01	9	0.02	100
EU-A534	87		1100	
EU-A538	55		3500	
EU-A750	>1000		>1000	
EU-A751	>1000		>1000	
EU-A752	146		>1000	
EU-A753	>1000		>1000	
EU-A754	360		>1000	
EU-A755	486		471	
EU-A756	611		>1000	
EU-A757	6.7		>1000	
EU-A770	0.01	2.3	0.5	>100
EU-A771	0.07	14.2	0.4	>100
EU-A772	0.07	8.4	5.4	>100
EU-A773	0.08	8.3	1.5	>100
EU-A774	0.009	6.8	0.15	22.7
EU-A775	0.16	17	0.3	33.6
EU-A776	1.2	>100	6.5	>100
EU-A777	0.1	34.5	0.6	73
EU-A792	<0.05	27.9	<0.05	
EU-A793	<0.05	23.8	<0.05	
EU-A794	0.05	59.4	0.18	
EU-A945	<0.05	9.4	0.06	
EU-A948	0.08	25.6	9.1	
EU-A992	0.009		0.019	
EU-A993	<0.05	12.3	<0.05	
EU-A994	0.05	10.2	<0.05	
EU-A995	0.035	59.5	0.15	
EU-A996	0.05	>100	0.87	
EU-A999	0.05		0.015	
EU-A1011	0.16		0.51	

Compound	EC ₅₀ cAMP GLP-1 R (nM)	EC ₅₀ arrestin GLP-1 R (nM)	EC ₅₀ cAMP glucagon R (nM)	EC ₅₀ arrestin glucagon R (nM)
EU-A1017	0.44		0.10	
EU-A1023	0.028		0.035	
EU-A1029	0.019		0.06	
EU-A1032	0.03		5.4	
EU-A1035	0.02		0.19	
EU-A1041	0.02		0.13	
EU-A1044	0.07		0.57	
EU-A1167	0.013		0.019	
EU-A1168	0.07		0.14	

Example 3-7: In vivo assay of compounds - db/db mice

[0627] The 60 female db/db B6BKS (D) Leprdb/J (strain 000697) mice for this study were approximately 8 to 9 weeks old at arrival (Jackson Laboratory, Bar Harbor, Maine). The mice were randomized by weight and two treatment groups of 8 female mice each were administered the test articles, EU-A994, EU-A995, or EU-A1026, at dose levels of 100 or 300 nmoles/kg. One group of 8 female mice served as the vehicle control and received the vehicle, 0.2% BSA in saline, pH 7.4. One additional group of 8 female mice received the positive control article, liraglutide, at a dose level of 50 nmoles/kg. The test articles, vehicle, and positive control article were administered on Day 1, at approximately 0, 7, and 24 hours during the study via subcutaneous injection at a dose volume of 6 mL/kg.

[0628] Clinical observations were conducted at receipt, prior to randomization, and daily from Days 1 to 5. Body weights were measured and recorded at receipt, prior to randomization, and daily from Days 1 to 5. Food consumption was measured and recorded daily from Days 1 to 5. Blood samples for glucose analysis were collected pretest (Day -3) and at 0, 1, 2, 4, 8, 10, 24, 48, 72, and 96 hours following the first dose on Day 1. At study termination all animals were euthanized and the carcasses were discarded without further evaluation.

[0629] Significant body weight changes were noted against vehicle for liraglutide and high dose EU-A994 and high dose EU-A1026 on Days 2 and 3 and low dose EU-A1026 on Days 3 and 4. In the food consumption analysis, liraglutide-treated animals were significantly different from vehicle on Days 1 and 2, high dose EU-A994 at Day 1, low dose EU-A995 at Day 1 and 2, high dose EU-A994 at Day 1, and low and high dose of EU-A1026 at Day 2 were significantly different from Liraglutide. Glucose levels for Liraglutide at 10 hours and high dose EU-A994 at 10 and 24 hours were significantly different from vehicle. Low dose EU-A995 and EU-A1026 at

10 hours were significantly different from liraglutide (Figure 5). In a like manner, other analogs from the series were tested for effects on blood glucose, body weight and food consumption.

Compound	Dose (nmole/kg)	db/db mice - Mean Blood Glucose (mg/dL)			
		0 hr	10hr	24hr	48hr
liraglutide	50	595	265	347	377
EU-A994	300	554	242	209	385
EU-A995	300	503	471	493	527
EU-A1026	300	531	399	438	505
liraglutide	50	528	286	404	477
EU-A993	250	503	305	324	426
EU-A995	250	539	584	548	546
EU-A1023	50	507	310	355	375
EU-A1023	250	508	264	194	375
liraglutide	50	442	174	244	439
EU-A992	250	405	104	112	130
EU-A1167	250	379	171	117	113
EU-A1168	250	380	133	169	164

Example 3-8: In vivo assay of compounds

[0630] Sixty (60) diet induced obese C57BL/6J male mice are received from JAX labs at 14 wks of age. The mice are ear notched for identification and housed in individually and positively ventilated polycarbonate cages with HEPA filtered air at density of one mouse per cage. The animal room is lighted entirely with artificial fluorescent lighting, with a controlled 12 h light/dark cycle. The normal temperature and relative humidity ranges in the animal rooms are $22 \pm 4^{\circ}\text{C}$ and $50 \pm 15\%$, respectively. Filtered tap water, acidified to a pH of 2.8 to 3.1, and high fat diet (60 kcal %) are provided ad libitum.

[0631] Following a 2 week acclimation, 40 mice are chosen based on desired body weight range and mice are randomized into groups (n=10) as below. Group 1. Vehicle treated; Group 2. Low dose test cmpd; Group 3. Mid dose test cmpd; Group 4. High dose test cmpd. Mice are dosed via SC daily for 28 days. Body weights and cage side observations are recorded daily. Food and water intake will be recorded weekly. Mice undergo NMR measurements for determining whole body fat and lean composition on days 1 (pre dose) and 26. On days 0, 14 and 27, mice are fasted overnight for an oral glucose tolerance test. Next day, the first blood sample is collected via tail nick (t=0). Mice are then administered a bolus of 1.0 g/kg glucose. Blood samples are obtained via tail nick at 5, 30, 60 and 120 min after glucose and plasma glucose will be immediately determined using a glucometer.

[0632] Sacrifice and tissue collection: Mice are sacrificed on day 29. Terminal blood is processed to serum/plasma and aliquots are sent for analysis of glucose, insulin and lipid profile. Body composition is determined by NMR. The optimal compound profile shows decreased glucose excursion in the OGTT, decreased basal insulin secretion, with potentiated glucose-dependent insulin secretion, decreased weight gain, decreased fat mass but minimal effects on lean mass.

Example 3-9: Plasma protease stability

[0633] Bioanalytical Method Development Summary I. Instruments used, API-4000 Mass Spectrometer, ESI positive, MRM Scan; Shimadzu HPLC/CTC Autosampler with ACE C8 column (2.1x50mm, 5µm), Mobile Phase A: 0.1% Formic acid, 5mM NH₄OAc in water, Mobile Phase B: 0.1% Formic acid in CH₃CN, 10µL of sample was injected, II. Standard and QC Sample Preparation: i. 1 mg/mL stock solution was prepared in DMSO/CH₃CN (1/1), ii. Make standard working solutions in 50%CH₃CN with the stock solution. The concentrations of the working solutions were 20, 50, 100, 200, 500, 1000, 2000, 5000, 10000 and 20000 ng/mL, iii. Spiked 10µL of the working solutions into 90µL blank plasma and vortex them, iv. Add 300µL of internal standard solution (Verapamil, 20ng/mL in 100%CH₃CN), vortex and centrifuge, v. Transfer the supernatant into an HPLC injection plate for loading on the HPLC column, vi. The standard samples were at 2, 5, 10, 20, 50, 100, 200, 500, 1000 and 2000 ng/mL. The QC samples were at 5 (LQC), 50 (MQC), and 500 (HQC) ng/mL. For the plasma stability study, samples of EU-A993, EU-A1023 and human GLP-1 (7-36, Bachem) were prepared in human plasma (approximately 6-20ng/mL or similar concentrations above the limit of quantitation) and sampled at time points t=0, 0.5, 1, 2, 4, 8hrs during incubation at 37degC. Samples were treated with internal standard solution (100% CH₃CN) as described above in order to precipitate the proteins and supernatants were loaded into an injection plate to load them onto the hplc column for quantitation by mass spectrometry. A plot of compound signal (amount) versus time shows rapid decrease in G1P-1 (7-36) and essentially no change over 8 hrs in the amount of intact compound in the human plasma for the compounds of the invention (see Figure 15).

Example 3-10: Uses of the compounds

[0634] The covalently modified peptides and/or proteins described herein are useful for the prevention and treatment of a variety of diseases related to obesity, the metabolic syndrome, cardiovascular disease and diabetes. Suitably labeled surfactant modified peptides can be used as diagnostic probes.

[0635] Representative delivery regimens include oral, parenteral (including subcutaneous, intramuscular and intravenous injection), rectal, buccal (including sublingual), transdermal,

inhalation ocular and intranasal. An attractive and widely used method for delivery of peptides entails subcutaneous injection of a controlled release injectable formulation. Other administration routes for the application of the covalently modified peptides and/or proteins described herein are subcutaneous, intranasal and inhalation administration.

Example 3-11. Pharmaceutical usage for treatment of insulin resistance.

[0636] A human patient, with evidence of insulin or metabolic syndrome is treated with EU-A596 by intranasal administration (200 μ L) from a standard atomizer used in the art of a solution of the pharmaceutical agent in physiological saline containing from 0.5 to 10 mg/mL of the pharmaceutical agent and containing standard excipients such as benzyl alcohol. The treatment is repeated as necessary for the alleviation of symptoms such as obesity, elevated blood glucose and the like. In a similar manner, a solution of EU-A596, and selected excipients, in an evaporating solvent containing such as a hydrofluoroalkane is administered intranasally by metered dose inhaler (MDI) as needed to reduce insulin resistance. The effect of treatment is determined using standard tests including measurement of blood glucose levels, Body Mass Index, and/or body weight and/or measurement of waist to hip ratios.

[0637] In a similar manner, administration of an adjusted amount by transbuccal, intravaginal, inhalation, subcutaneous, intravenous, intraocular, or oral routes is tested to determine level of stimulation of GLP1R and/or GLCR on cells in the body and to determine therapeutic effects.

SEQUENCES

[0638] Table 1 in Figure 1 depicts compounds that were prepared by methods described herein. The specification provides sequences for SEQ. ID. Nos. 1 and 149-169. Additionally, Table 1 of Figure 1 provides SEQ. ID Numbers for compounds EU-A101 to EU-A199 and EU-A600 to EU-A649 having SEQ. ID. NOs. 2-148, and SEQ. ID. NO. 645 respectively, as shown in Table 1 of Figure 1. Compounds in Table 1 of Figure 1, and their respective SEQ. ID. NOs. shown in Table 1 of Figure 1 are hereby incorporated into the specification as filed.

[0639] Table 2 in Figure 2 depicts compounds that were prepared by methods described herein. The specification provides sequences for SEQ. ID. Nos. 170-174 and SEQ. ID. NOs. 283-302. Additionally, Table 2 of Figure 2 provides SEQ. ID. Numbers for compounds EU-201 to EU-299 and EU-900 to EU-908 having SEQ. ID. NOs. 175-282 respectively, as shown in Table 2 of Figure 2. Compounds in Table 2 of Figure 2, and their respective SEQ. ID. NOs. shown in Table 2 of Figure 2 are hereby incorporated into the specification as filed.

[0640] Table 3 in Figure 8 depicts compounds that were prepared by methods described herein. The specification provides sequences for SEQ. ID. Nos. 303-305 and SEQ. ID. Nos. 619 -644. Additionally, Table 3 of Figure 8 provides SEQ. ID Numbers for compounds EU-A300

to EU-A425 having SEQ. ID. NOs. 306-431 respectively, as shown in Table 3 of Figure 8. Compounds in Table 3 of Figure 8, and their respective SEQ. ID. NOs. shown in Table 3 of Figure 8 are hereby incorporated into the specification as filed.

[0641] Table 4 in Figure 9 depicts compounds that were prepared by methods described herein. The specification provides SEQ. ID. Nos. 303-305 and SEQ. ID. Nos. 619 -644. Additionally, Table 4 of Figure 9 provides SEQ. ID Numbers for compounds EU-A426 to EU-A599 having SEQ. ID. NOs. 432-618 respectively, as shown in Table 4 of Figure 9. Additionally, Table 4 of Figure 9 provides SEQ. ID Numbers for compounds EU-A700 to EU-A1174 having SEQ. ID. NOs 646-1120. Compounds in Table 4 of Figure 9, and their respective SEQ. ID. NOs. shown in Table 4 of Figure 9 are hereby incorporated into the specification as filed.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2012158962A \[0003\]](#)
- [US7390788B \[0108\]](#)
- [US7425542B \[0108\]](#)
- [US20100190701A1 \[0329\]](#)
- [FR18245 \[0381\]](#)
- [US3219656A \[0426\]](#)
- [US3839318A \[0426\]](#)
- [AT386414 \[0427\]](#)
- [AT382381 \[0428\] \[0428\]](#)
- [US4640835A \[0439\]](#)
- [US4496689A \[0439\]](#)
- [US4301144A \[0439\]](#)
- [US4670417A \[0439\]](#)
- [US4791192A \[0439\]](#)
- [US4179337A \[0439\]](#)
- [US6309633B \[0440\]](#)
- [US6815530B \[0440\]](#)
- [US6835802B \[0440\]](#)
- [WO9119481A \[0451\]](#)

- [US4675189A](#) [\[0506\]](#)

Non-patent literature cited in the description

- NESTOR, J.J., JR.Comprehensive Medicinal Chemistry II, 2007, vol. 2, 573-601 [\[0003\]](#) [\[0240\]](#) [\[0375\]](#) [\[0398\]](#)
- AKIYAMA K et al.Proc Natl Acad Sci USA, 1985, vol. 82, 2543-7 [\[0054\]](#) [\[0059\]](#)
- MELCHIORRI, P.NEGRI, L.Gen Pharmacol, 1996, vol. 27, 1099-1107 [\[0055\]](#)
- ERSPAMER, V. et al.Proc Natl Acad Sci USA, 1989, vol. 86, 5188-92 [\[0055\]](#)
- LAZARUS, L.H.OKADA, Y.Expert Opin Ther Patents, 2012, vol. 22, 1-14 [\[0056\]](#) [\[0116\]](#)
- JAMES, I.F. et al.Life Sci, 1982, vol. 31, 1331-4 [\[0057\]](#)
- CONDON, S.M. et al.Bioorg Med Chem, 2002, vol. 10, 731-736 [\[0058\]](#) [\[0157\]](#) [\[0269\]](#)
- PIOZAK, A.A. et al.J Biol Chem, 2009, vol. 284, 28382-391 [\[0101\]](#)
- RUNGE, S. et al.J Biol Chem, 2008, vol. 283, 11340-7 [\[0101\]](#) [\[0335\]](#)
- SHENOY, S. K.LEFKOWITZ, R.J.Trends Pharmacolog Sci, 2011, vol. 32, 521-533 [\[0104\]](#)
- MAJI, S.K. et al.PLoS Biol, 2008, vol. 6, e17- [\[0105\]](#)
- CLODFELTER, D.K. et al.Pharm Res, 1998, vol. 15, 254-262 [\[0105\]](#) [\[0445\]](#)
- HERMELING, S. et al.J Pharm Sci, 2006, vol. 95, 1084-1096 [\[0107\]](#)
- CASADEVALL, N. et al.N Engl J Med, 2002, vol. 346, 469-475 [\[0107\]](#)
- PEZRON, I. et al.J Pharm Sci, 2002, vol. 91, 1135-1146 [\[0107\]](#)
- SLUZKY, V. et al.Proc Natl Acad Sci U S A, 1991, vol. 88, 9377-9381 [\[0107\]](#)
- KING, H.D. et al.J Med Chem, 2002, vol. 45, 4336-4343 [\[0107\]](#)
- PORTER, W.R. et al.Thromb Res, 1993, vol. 71, 265-279 [\[0107\]](#)
- ZHANG, M.Z. et al.Pharm Res, 1995, vol. 12, 1447-1452 [\[0108\]](#)
- VRKLJAN, M. et al.Pharm Res, 1994, vol. 11, 1004-1008 [\[0108\]](#)
- ROCCATANO, D. et al.Proc Natl Acad Sci U S A, 2002, vol. 99, 12179-12184 [\[0108\]](#)
- JENSEN, K.J.BRASK, J.Biopolymers, 2005, vol. 80, 747-761 [\[0110\]](#)
- GIJSEN, H.J. et al.Chem Rev, 1996, vol. 96, 443-474 [\[0110\]](#)
- SEARS, P.WONG, C.H.Cell Mol Life Sci, 1998, vol. 54, 223-252 [\[0110\]](#)
- GUO, Z.SHAO, N.Med Res Rev, 2005, vol. 25, 655-678 [\[0110\]](#)
- URGE, L. et al.Biochem Biophys Res Commun, 1992, vol. 184, 1125-1132 [\[0110\]](#) [\[0110\]](#)
- SALVADOR, L.A. et al.Tetrahedron, 1995, vol. 51, 5643-5656 [\[0110\]](#)
- KIHLEBERG, J. et al.Methods Enzymol, 1997, vol. 289, 221-245 [\[0110\]](#) [\[0511\]](#) [\[0562\]](#) [\[0599\]](#)
- GREGORIADIS, G. et al.Cell Mol Life Sci, 2000, vol. 57, 1964-1969 [\[0110\]](#)
- CHAKRABORTY, T.K. et al.Glycoconj J, 2005, vol. 22, 83-93 [\[0110\]](#)
- LIU, M. et al.Carbohydr Res, 2005, vol. 340, 2111-2122 [\[0110\]](#)
- PAYNE, R.J. et al.J Am Chem Soc, 2007, vol. 129, 13527-13536 [\[0110\]](#)

- PEDERSEN, S.L. et al.Chembiochem, 2010, vol. 11, 366-374 [0110]
- FILIRA, F. et al.Org Biomol Chem, 2003, vol. 1, 3059-3063 [0110] [0110]
- NEGRI, L. et al.J Med Chem, 1999, vol. 42, 400-404 [0110]
- NEGRI, L. et al.Br J Pharmacol, 1998, vol. 124, 1516-1522 [0110]
- ROCCHI, R. et al.Int J Pept Protein Res, 1987, vol. 29, 250-261 [0110]
- FILIRA, F. et al.Int J Biol Macromol, 1990, vol. 12, 41-49 [0110]
- GOBBO, M. et al.Int J Pept Protein Res, 1992, vol. 40, 54-61 [0110]
- DJEDAINI-PILARD, F. et al.Tetrahedron Lett, 1993, vol. 34, 2457-2460 [0110]
- DROUILLAT, B. et al.Bioorg Med Chem Lett, 1997, vol. 7, 2247-2250 [0110]
- LOHOF, E. et al.Angew Chem Int Ed Engl, 2000, vol. 39, 2761-2764 [0110]
- GRUNER, S.A. et al.Org Lett, 2001, vol. 3, 3723-3725 [0110]
- PEAN, C. et al.Biochim Biophys Acta, 2001, vol. 1541, 150-160 [0110]
- GROTENBREG, G.M. et al.J Org Chem, 2004, vol. 69, 7851-7859 [0110]
- BIONDI, L. et al.J Pept Sci, 2007, vol. 13, 179-189 [0110]
- KODA, Y. et al.Bioorg Med Chem, 2008, vol. 16, 6286-6296 [0110] [0527]
- LOWERY J.J. et al.J Pharmacol Exptl Therap, 2011, vol. 336, 767-78 [0110]
- YAMAMOTO, T. et al.J Med Chem, 2009, vol. 52, 5164-5175 [0110]
- SCHILLER, P.W.AAPS J, 2005, vol. 7, E560-565 [0115]
- STEIN, C. et al.Brain Res Rev, 2009, vol. 60, 90-113 [0115]
- SCHILLER, P.W.Life Sci, 2010, vol. 86, 598-603 [0115] [0116]
- JANECKA, A. et al.Curr Med Chem, 2007, vol. 14, 3201-3208 [0116]
- KERESZTES, A. et al.ChemMedChem, 2010, vol. 5, 1176-96 [0116]
- SCHILLER, P.W. et al.Biopolymers, 1999, vol. 51, 411-25 [0116]
- NESTOR, J.J., JR.Current Medicinal Chemistry, 2009, vol. 16, 4399-4418 [0131] [0240] [0374] [0375] [0398] [0404] [0434]
- ZHANG, LBULAJ, G.Curr Med Chem, 2012, vol. 19, 1602-18 [0131]
- BRIKEN, K.T. et al.Basic Clin Pharmacol Toxicol, 2004, vol. 94, 260-270 [0195]
- DOBNIG, H.Expert Opin Pharmacother, 2004, vol. 5, 1153-1162 [0195]
- BOYCE, B.F.XING, L.Arthritis Res Ther, 2007, vol. 9, 11- [0196] [0199]
- ROSEN, C.J.BILEZIKIAN, J.P.J Clin Endocrinol Metab, 2001, vol. 86, 957-964 [0197]
- REEVE, J. et al.Br Med J, 1980, vol. 280, 1340-1344 [0198]
- HODSMAN, A.B. et al.Endocr Rev, 2005, vol. 26, 688-703 [0198]
- CHENG, Z. et al.J Bone Miner Res, 2009, vol. 24, 209-220 [0198]
- FROLIK, C.A. et al.Bone, 2003, vol. 33, 372-379 [0199] [0199] [0588]
- ETOH, M.YAMAGUCHI, A.J Bone Miner Metab, 2010, vol. 28, 641-9 [0199]
- TASHJIAN, A.H., JR.GOLTZMAN, D.J Bone Miner Res, 2008, vol. 23, 803-811 [0200]
- WHITFIELD, J.F.MORLEY, P.Trends Pharmacol Sci, 1995, vol. 16, 382-386 [0201]
- SHIMIZU, M. et al.J Biol Chem, 2000, vol. 275, 21836-21843 [0201] [0210]
- SHIMIZU, N. et al.J Bone Miner Res, 2004, vol. 19, 2078-2086 [0201] [0210]
- GARDELLA, T.J. et al.Endocrinology, 1994, vol. 135, 1186-1194 [0202]
- LUCK, M.D. et al.Mol Endocrinol, 1999, vol. 13, 670-680 [0202]
- DEAN, T. et al.J Biol Chem, 2006, vol. 281, 32485-32495 [0202] [0209]
- POTETINOVA, Z. et al.Biochemistry, 2006, vol. 45, 11113-11121 [0202]
- HOLTMANN, M.H. et al.J Biol Chem, 1995, vol. 270, 14394-14398 [0202]

- **BERGWITZ, C. et al.**J Biol Chem, 1996, vol. 271, 26469-26472 [0202]
- **RUNGE, S. et al.**J Biol Chem, 2003, vol. 278, 28005-28010 [0202] [0329]
- **PIOSZAK, A.A.XU, H.E.**Proc Natl Acad Sci U S A, 2008, vol. 105, 5034-5039 [0203]
- **PIOSZAK, A.A. et al.**J Biol Chem, 2009, vol. 284, 28382-28391 [0203]
- **KAUL, R.BALARAM, P.**Bioorg Med Chem, 1999, vol. 7, 105-117 [0210] [0399]
- **SHIMIZU, N. et al.**J Biol Chem, 2001, vol. 276, 49003-49012 [0210]
- **PEGGION, E. et al.**Biopolymers, 2003, vol. 68, 437-457 [0210]
- **BARAZZA, A. et al.**J Pept Res, 2005, vol. 65, 23-35 [0210]
- **MARX, U.C. et al.**J Biol Chem, 1995, vol. 270, 15194-15202 [0210]
- **SCHIEVANO, E. et al.**Biopolymers, 2000, vol. 54, 429-447 [0210]
- **NESTOR, J.J., JR.**Current Medicinal Chemistry, 2009, vol. 16, 4399-4418 [0238] [0403]
- **GOURLET, P. et al.**Eur J Pharmacol, 1998, vol. 354, 105-111 [0238] [0374] [0404]
- **GOZES, I.FURMAN, S.**Curr Pharm Des, 2003, vol. 9, 483-494 [0238] [0374] [0404]
- **MAHAFFEY, J.E. et al.**J Biol Chem, 1979, vol. 254, 6496-6498 [0238] [0374]
- **GOLDMAN, M.E. et al.**Endocrinology, 1988, vol. 123, 2597-2599 [0238] [0374]
- **GARDELLA, T.J. et al.**Endocrinology, 1996, vol. 137, 3936-3941 [0239]
- **WOLD, F.**Posttranslational Covalent Modification of Proteins, 1983, 1-12 [0240] [0375] [0398]
- **SEIFTER, S.ENGLARD, S.**Methods Enzymol, 1990, vol. 182, 626-646 [0240] [0375] [0398]
- **RATTAN, S.I. et al.**Ann N Y Acad Sci, 1992, vol. 663, 48-62 [0240] [0375] [0398]
- **ZHANG, L.BULAJ, G.**Curr Med Chem, 2012, vol. 19, 1602-1618 [0242] [0376]
- **MURAGE, E.N. et al.**Bioorg Med Chem, 2008, vol. 16, 10106-12 [0269]
- **MURAGE, E.N. et al.**J Med Chem, 2010, vol. 53, 6412-20 [0269]
- **LI, Y. et al.**Peptides, 2011, vol. 32, 1400-1407 [0269]
- **LE CHEVALIER ISAAD A. et al.**J Peptide Sci, 2009, vol. 15, 451-4 [0269]
- **KIEFFER, T.J. et al.**Endocrinology, 1995, vol. 136, 3585-3596 [0322]
- **AABOE, K. et al.**Diabetes Obes Metab, 2008, vol. 10, 994-1003 [0322]
- **SOKOS, G.G. et al.**J Card Fail, 2006, vol. 12, 694-699 [0322]
- **TOFT-NIELSEN, M.B. et al.**Diabetes Care, 1999, vol. 22, 1137-1143 [0322]
- **DRAB, S.R.**Pharmacotherapy, 2010, vol. 30, 609-624 [0323]
- **SANGER, G.J.LEE, K.**Nat Rev Drug Discov, 2008, vol. 7, 241-254 [0324]
- **DRUCKER, D.J.**Nat Clin Pract Endocrinol Metab, 2005, vol. 1, 22-31 [0324]
- **SINCLAIR, E.M.DRUCKER, D.J.**Physiology (Bethesda), 2005, vol. 20, 357-365 [0324]
- **MACK, C.M. et al.**Int J Obes (Lond), 2006, vol. 30, 1332-1340 [0325]
- **DEFRONZO, R.A. et al.**Diabetes Care, 2005, vol. 28, 1092-1100 [0325]
- **BUSE, J.B. et al.**Diabetes Care, 2010, vol. 33, 1255-1261 [0325]
- **GEARY, N. et al.**Am J Physiol, 1992, vol. 262, R975-980 [0325]
- **HECKEMEYER, C.M. et al.**Endocrinology, 1983, vol. 113, 270-276 [0325]
- **SALTER, J.M. et al.**Metabolism, 1960, vol. 9, 753-768 [0325]
- **CHAN, E.K. et al.**Exp Mol Pathol, 1984, vol. 40, 320-327 [0325]
- **HEPPNER, K.M. et al.**Physiol Behav, 2010, [0325]
- **DAKIN, C.L. et al.**Endocrinology, 2004, vol. 145, 2687-2695 [0326]
- **COHEN, M.A. et al.**J Clin Endocrinol Metab, 2003, vol. 88, 4696-4701 [0326]

- WYNNE, K. et al. Diabetes, 2005, vol. 54, 2390-2395 [0326]
- JORGENSEN, R. et al. J Pharmacol Exp Ther, 2007, vol. 322, 148-154 [0327]
- ADELHORST, K. et al. J Biol Chem, 1994, vol. 269, 6275-6278 [0328]
- HJORTH, S.A. et al. J Biol Chem, 1994, vol. 269, 30121-30124 [0329]
- POCAI, A. et al. Diabetes, 2009, vol. 58, 2258-2266 [0329] [0329]
- DAY, J.W. et al. Nat Chem Biol, 2009, vol. 5, 749-757 [0329] [0331]
- DRUCE, M.R. et al. Endocrinology, 2009, vol. 150, 1712-1722 [0329]
- KOSINSKI, J.R. et al. Obesity Silver Spring 2012 0000 [0331]
- MAPELLI, C. et al. J Med Chem, 2009, vol. 52, 7788-7799 [0334]
- HAQUE, T.S. et al. Peptides, 2010, vol. 31, 950-955 [0334]
- HAQUE, T.S. et al. Peptides, 2010, vol. 31, 1353-1360 [0334]
- DAVIDSON, M.H. Am J Cardiol, 2011, vol. 108, 33B-41B [0338]
- GEJL, M. et al. J Clin Endocrinol Metab, 2012, vol. 97, [0338]
- GAUTHIER, M.A. KLOK, H.A. Chem Commun (Camb), 2008, 2591-2611 [0378] [0405] [0422]
- DE GRAAF, A.J. et al. Bioconjug Chem, 2009, vol. 20, 1281-1295 [0378] [0394] [0405]
- DEITERS, A et al. Bio-org. Med. Chem. Lett., 2004, vol. 14, 5743-5 [0378] [0405]
- Pure Appl. Chem., 1972, vol. 31, 639-645 [0381]
- PURE APPL. CHEM., 1974, vol. 40, 277-290 [0381]
- STEVENSON, C.L. Curr Pharm Biotechnol, 2009, vol. 10, 122-137 [0403]
- TYNDALL, J.D. et al. Chem Rev, 2005, vol. 105, 793-826 [0403]
- ABBENANTE, G. FAIRLIE, D.P. Med Chem, 2005, vol. 1, 71-104 [0403]
- PARK, D.W. et al. Biotechnology Letters, 2000, vol. 22, 951-956 [0416]
- VAN BEKKUM, H. Carbohydrates as Organic Raw Materials, 1990, 289-310 [0416]
- DE MICO, A. et al. J Org Chem 1997, 1997, 6974-6977 [0416]
- EPP, J.B. WIDLANSKI, T.S. J Org Chem, 1999, vol. 64, 293-295 [0416]
- CODEE, J.D. et al. J Am Chem Soc, 2005, vol. 127, 3767-3773 [0416]
- MILKEREIT, G. et al. Chem Phys Lipids, 2004, vol. 127, 47-63 [0416] [0535]
- SCHAMANN, M. SCHAFER, H.J. Eur J Org Chem, 2003, 351-358 [0416]
- VAN DEN BOS, L.J. et al. Eur J Org Chem, 2007, 3963-3976 [0416]
- SMITS, E. et al. J Chem Soc, Perkin Trans, 1996, vol. I, 2873-2877 [0417]
- SMITS, E. et al. Liquid Crystals, 1997, vol. 23, 481-488 [0417]
- T. W. GREENP. G. M. WUTS Protective Groups in Organic Synthesis Wiley-Interscience 1999 0000 vol. 503-507, 736-739 [0419]
- LAROCK, R.C. Comprehensive Organic Transformations VCH Publishers 1999 0000 [0422]
- ELOFSSON, M. et al. Tetrahedron, 1997, vol. 53, 369-390 [0423]
- STANGIER, P. et al. Liquid Crystals, 1994, vol. 17, 589-595 [0423]
- CUDIC, M. BURSTEIN, G.D. Methods Mol Biol, 2008, vol. 494, 187-208 [0424]
- VILL, V. et al. Chem Phys Lipids, 2000, vol. 104, 75-91 [0424]
- MINDEN, H.M. et al. Chem Phys Lipids, 2000, vol. 106, 157-179 [0424]
- ROSEVEAR, P. et al. Biochemistry, 1980, vol. 19, 4108-4115 [0426]
- LI, Y.T. et al. J Biol Chem, 1991, vol. 266, 10723-10726 [0426] [0426]
- KOELTZOWURFER J. Am. Oil Chem. Soc., 1984, vol. 61, 1651-1655 [0426]


- **GOPALAN, V. et al.** J Biol Chem, 1992, vol. 267, 9629-9638 [\[0426\]](#)
- Carbohydrates as Organic Raw Materials VCH Publishers 19910000 [\[0427\]](#)
- Chem. Abstr., 1989, vol. 110, 137536- [\[0427\]](#)
- **GRUBER, H. GREBER, G.** Reactive Sucrose Derivatives Carbohydrates as Organic Raw Materials, 95-116 [\[0427\]](#)
- **SUGAR ESTERS** Preparation and Application Noyes Data Corp. 19740000 [\[0427\]](#)
- Hydrogen Fluoride, Solvent and Reagent for Carbohydrate Conversion Technology **DEFAYE, J. PEDERSON, C.** Carbohydrates as Organic Raw Materials VCH Publishers 19910000247-265 [\[0428\]](#)
- **FERENCI, T.J.** Bacteriol, 1980, vol. 144, 7-11 [\[0428\]](#)
- **SAITO, S. TSUCHIYA, T.** Chem. Pharm. Bull., 1985, vol. 33, 503-508 [\[0428\]](#)
- **BINDER, T. P. ROBYT, J. F.** Carbohydr. Res., 1985, vol. 140, 9-20 [\[0428\]](#)
- Chem. Abstr., 1988, vol. 108, 114719- [\[0428\]](#) [\[0428\]](#)
- **GRUBER GREBER** CHEM. ABSTR., 95-116 [\[0428\]](#) [\[0428\]](#)
- **KUNZ, M.** Sucrose-based Hydrophilic Building Blocks as Intermediates for the Synthesis of Surfactants and Polymers Carbohydrates as Organic Raw Materials, 127-153 [\[0428\]](#)
- **LOPEZ, O. et al.** Colloid Polym Sci, 2002, vol. 280, 352-357 [\[0433\]](#)
- **VERONESE, F.M. MERO, A.** BioDrugs, 2008, vol. 22, 315-329 [\[0434\]](#)
- **CALICETI, P. VERONESE, F.M.** Adv Drug Deliv Rev, 2003, vol. 55, 1261-1277 [\[0434\]](#)
- **PELEG-SHULMAN, T. et al.** J Med Chem, 2004, vol. 47, 4897-4904 [\[0435\]](#)
- **GREENWALD, R.B. et al.** Adv Drug Deliv Rev, 2003, vol. 55, 217-250 [\[0435\]](#) [\[0441\]](#)
- **ROBERTS, M.J. HARRIS, J.M.** J Pharm Sci, 1998, vol. 87, 1440-1445 [\[0435\]](#)
- **MERO, A. et al.** J Control Release, 2008, vol. 125, 87-95 [\[0438\]](#)
- **MILLER, M.A. et al.** Bioconj Chem, 2006, vol. 17, 267-274 [\[0440\]](#)
- **VERONESE, F.M. PASUT, G.** Drug Discov Today, 2005, vol. 10, 1451-1458 [\[0441\]](#) [\[0443\]](#)
- **ROBERTS, M.J. et al.** Adv Drug Deliv Rev, 2002, vol. 54, 459-476 [\[0441\]](#) [\[0443\]](#)
- **THUROW, H. GEISEN, K.** Diabetologia, 1984, vol. 27, 212-218 [\[0449\]](#)
- **CHAWLA, A.S. et al.** Diabetes, 1985, vol. 34, 420-424 [\[0449\]](#)
- **ARAKAWA, T. TIMASHEFF, S.N.** Biochemistry, 1982, vol. 21, 6536-6544 [\[0449\]](#)
- **ARAKAWA, T. TIMASHEFF, S.N.** Biophys J, 1985, vol. 47, 411-414 [\[0449\]](#)
- **SATO, S. et al.** J Pharm Sci, 1983, vol. 72, 228-232 [\[0449\]](#)
- **ANDYA, J.D. et al.** Pharm Res, 1999, vol. 16, 350-358 [\[0452\]](#)
- Remington's Pharmaceutical Sciences Mack Publishing Co [\[0458\]](#)
- **BLOOM, S.R. et al.** Mol Interv, 2008, vol. 8, 82-98 [\[0493\]](#)
- **FIELD, B.C. et al.** Br J Clin Pharmacol, 2009, vol. 68, 830-843 [\[0493\]](#)
- **GONZALEZ-PERIZ, A. CLARIA, J.** Scientific World Journal, 2010, vol. 10, 832-856 [\[0495\]](#)
- Remington's Pharmaceutical Sciences Mack Publishing Company 19850000 [\[0502\]](#)
- **ARNOLD, J.J. et al.** J Pharm Sci, 2004, vol. 93, 2205-2213 [\[0503\]](#)
- **AHSAN, F. et al.** Pharm Res, 2001, vol. 18, 1742-1746 [\[0503\]](#)
- **PFUTZNER, A. FORST, T.** Expert Opin Drug Deliv, 2005, vol. 2, 1097-1106 [\[0504\]](#)
- **MANDAL, T.K.** Am. J. Health Syst. Pharm., 2005, vol. 62, 1359-64 [\[0504\]](#)
- Sustained and Controlled Release Drug Delivery Systems Marcel Dekker, Inc. 19780000 [\[0506\]](#)
- **R. W. BAKER** Controlled Release of Biologically Active Agents John Wiley &

Sons19870000 [0506]

- **HOUSTON, M.E., JR. et al.**J Pept Sci, 1995, vol. 1, 274-282 [0514] [0565] [0601] [0602]
- **MURAGE, E.N. et al.**J Med Chem, 2010, [0514] [0565] [0601] [0602]
- **WANG, J.B. et al.**FEBS Lett, 1994, vol. 338, 217-222 [0550]
- **LAW, P.Y.LOH, H.H.**Mol Pharmacol, 1993, vol. 43, 684-693 [0550]
- **AVIDOR-REISS, T. et al.**FEBS Lett, 1995, vol. 361, 70-74 [0550]
- **LAZARUS, L.H. et al.**Expert Opin Ther Patents, 2012, vol. 22, 1-14 [0555]
- **ORLOFF, J.J. et al.**Endocrinol, 1992, vol. 131, 1603-1611 [0585]
- **CHICCHI, G.G. et al.**J Biol Chem, 1997, vol. 272, 7765-7769 [0625]
- **RUNGE, S. et al.**Br J Pharmacol, 2003, vol. 138, 787-794 [0625]


Patentkrav

- 5 **1.** Peptidprodukt, som omfatter et tensid X, der er kovalent bundet til et peptid, idet peptidet omfatter en linker-aminosyre U og mindst en anden aminosyre:



Formel I

hvor X er



10 hvor:

A er en hydrofob gruppe, som er en substitueret eller ikke-substitueret C₁-C₃₀-alkylgruppe, en substitueret eller ikke-substitueret alkoxyarylgruppe, en substitueret eller ikke-substitueret aralkylgruppe eller en steroidkerneholdig del; og

15 B er en melibiose eller melibiouronsyregruppe, der er kovalent bundet til peptidet via linker-aminosyren U.

20 **2.** Peptidprodukt ifølge krav 1, hvor peptidet er et opioidt, PTH-, PTHrP-, glucagon- eller GLP-peptid med en naturligt eller ikke-naturligt forekommende aminosyresekvens eller -struktur.

3. Peptidprodukt ifølge krav 1, som har en struktur med formelen II:

25 $aa_1-aa_2-aa_3-aa_4-aa_5-Z$ Formel II


hvor:

30 hver af aa₁, aa₂, aa₃, aa₄ og aa₅ uafhængigt er fraværende, en naturlig eller unaturlig D- eller L-aminosyre, en N-alkyleret aminosyre, en N-acetyleret aminosyre, en CaR³-aminosyre, en Ψ-aminosyre eller en linker-aminosyre U, der er kovalent bundet til et tensid X;

Z er -OH, -NH₂ eller -NHR³;

hver R³ uafhængigt er substitueret eller ikke-substitueret, forgrenet eller
 ligekædet C₁-C₁₂-alkyl, en PEG-kæde på mindre end 10 Da eller en
 substitueret eller ikke-substitueret aralkylgruppe; og

5 X er



hvor:

A er en hydrofob gruppe, som er en substitueret eller ikke-substitueret C₁-C₃₀-
 alkylgruppe, en substitueret eller ikke-substitueret alkoxyarylgruppe, en
 10 substitueret eller ikke-substitueret aralkylgruppe eller en steroidkerneholdig
 del; og

B er en melibiose eller melibiouronsyregruppe, der er kovalent bundet til
 peptidet via en linker-aminosyre U;

idet det forudsættes, at en eller mindst en af aa₁, aa₂, aa₃, aa₄ og aa₅ er en
 15 linker-aminosyre U, der er kovalent bundet til tensidet X; og
 idet det endvidere forudsættes, at ikke alle af aa₁, aa₂, aa₃, aa₄ og aa₅ er
 fraværende.

20 **4. Peptidprodukt ifølge krav 3, hvor peptidproduktet har en struktur med formlen III:**

aa₁-aa₂-aa₃-aa₄-aa₅-Z

Formel III (SEQ. ID. NO.1)

hvor:

25 aa₁ er Tyr, Dmt, N-R³-Tyr, N-R³-Dmt, N-(R³)₂-Tyr eller N-(R³)₂-Dmt;
 aa₂ er Pro, D-Arg, D-U(X), D-Ala, D-Cit, Tic eller Tic(Ψ[CH₂-NH]);
 aa₃ er Phe, Trp, Tmp, D- eller L-Nal(1), D- eller L-Nal(2), CaMePhe eller Ψ-
 Phe;


30 aa₄ er Phe, Tmp, D- eller L-Nal(1), D- eller L-Nal(2), U(X) eller D- eller L-
 CaMeU(X);

aa₅ er fraværende eller Pro, Aib, U(X) eller D- eller L-CaMeU(X); og
 U er en linker-aminosyre, der er kovalent bundet til tensidet X, og er en
 naturlig eller unaturlig dibasisk eller disur aminosyre, en naturlig eller unaturlig

aminosyre, der omfatter en thiol, en unaturlig aminosyre, der omfatter en $-N_3$ -gruppe, en unaturlig aminosyre, der omfatter en acetylengruppe, eller en unaturlig aminosyre, der omfatter en $-NH-C(=O)-CH_2-Br-$ eller en $-(CH_2)_m$ -maleimidgruppe, hvor m er 1-10.

5

5. Peptidprodukt ifølge krav 3 eller 4, hvor tensidet X har strukturen:




hvor:

- 10 A er en substitueret eller ikke-substitueret C_1 - C_{30} -alkylgruppe, en substitueret eller ikke-substitueret alkoxyarylgruppe, en substitueret eller ikke-substitueret aralkylgruppe eller en steroidkerneholdig del;
 R^{1b} , R^{1c} og R^{1d} hver uafhængigt i hvert tilfælde er H, en beskyttelsesgruppe eller en substitueret eller ikke-substitueret C_1 - C_{30} -alkylgruppe;
- 15 W^1 uafhængigt i hvert tilfælde er $-CH_2-$, $-CH_2-O-$, $-(C=O)-$, $-(C=O)-O-$, $-(C=O)-NH-$, $-(C=S)-$, $-(C=S)-NH-$ eller $-CH_2-S-$;
 W^2 er $-O-$ eller $-S-$;
 R^2 er en binding, C_2 - C_4 -alken, C_2 - C_4 -alkyn eller $-(CH_2)_m$ -maleimid;
m er 1-10; og
- 20 n er 2.

6. Peptidprodukt ifølge krav 5, hvor tensidet X har strukturen:

(a)



25

hvor:

A er en substitueret eller ikke-substitueret C₁-C₃₀-alkylgruppe eller en steroidkerneholdig del;


R^{1b}, R^{1c} og R^{1d} hver uafhængigt i hvert tilfælde er H, en beskyttelsesgruppe eller en substitueret eller ikke-substitueret C₁-C₃₀-alkylgruppe;

5 W¹ er -(C=O)-NH-;

W² er -O-; og

R² er en binding; eller

(b)



Formel V

10 hvor:

A er en substitueret eller ikke-substitueret C₁-C₃₀- eller C₁-C₂₀-alkylgruppe;

R^{1b}, R^{1c} og R^{1d} er H;

W¹ er -(C=O)-NH-;

W² er -O-; og

15 R² er en binding.

7. Peptidprodukt ifølge et hvilket som helst af kravene 3 til 6, hvor:

(a) U er en naturlig eller unaturlig dibasisk aminosyre; (**SEQ. ID. NO. 149**)

20 X er et tensid i 1-alkylglycosid-klassen, hvor 1-alkyl er substitueret eller ikke-substitueret C₁-C₂₀-alkyl eller en substitueret eller ikke-substitueret aralkylgruppe;

Z er NH₂;

25 aa₁ er Tyr, Dmt, Nα-Me-Tyr, Nα-Me-Dmt, N,Nα-diMe-Tyr eller N,Nα-diMe-Dmt;

aa₂ er Pro, D-Arg, D-U(X), D-Ala, D-Cit, Tic eller Tic(Ψ[CH₂-NH]);

aa₃ er Phe, Trp, Tmp, D- eller L-Nal(1), D- eller L-Nal(2), CaMePhe eller Ψ-Phe;

30 aa₄ er Phe, Tmp, D- eller L-Nal(1), D- eller L-Nal(2), U(X) eller D- eller L-CaMeU(X); og

aa₅ er fraværende eller Pro, Aib, U(X) eller D- eller L-CaMeU(X); eller

(b) X omfatter 1-alkylmelibiouronsyre; (**SEQ. ID. NO. 150**)

Z er NH₂;

aa₁ er Dmt;

5 aa₂ er Pro, D-Lys(X), Tic eller Tic(Ψ[CH₂-NH]);

aa₃ er Phe, Tmp, D- eller L-Nal(1), D- eller L-Nal(2) eller Ψ-Phe;

aa₄ er Phe, D- eller L-Nal(1), D- eller L-Nal(2) eller Lys(X); og

aa₅ er fraværende eller Pro, Aib, Lys(X) eller D- eller L-CaMeLys(X); eller

(c) aa₁ er Dmt; (**SEQ. ID. NO. 151**)

10 aa₂ er Pro;

aa₃ er Phe eller Tmp;

aa₄ er Phe eller Lys(X); og

aa₅ er fraværende eller Aib, Lys(X) eller D- eller L-CaMeLys(X); eller

(d) aa₁ er Dmt; (**SEQ. ID. NO. 152**)

15 aa₂ er Pro;

aa₃ er Phe eller Tmp;

aa₄ er Phe eller Lys(X); og

aa₅ er fraværende eller Lys(X); eller

(e) U er en naturlig eller unaturlig dibasisk aminosyre; (**SEQ. ID. NO. 153**)

20 X er et tensid i 1-alkylglycosid-klassen, hvor 1-alkyl er substitueret eller ikke-substitueret C₁-C₂₀-alkyl eller en substitueret eller ikke-substitueret aralkylgruppe;

aa₁ er Tyr, Dmt, N-R³-Tyr, N-R³-Dmt, N-(R³)₂-Tyr eller N-(R³)₂-Dmt;

aa₂ er D-Arg, D-Cit eller D-U(X);

25 aa₃ er Phe, Trp, D- eller L-Nal(1), D- eller L-Nal(2) eller Tmp;

aa₄ er Phe, Tmp, D- eller L-Nal(1), D- eller L-Nal(2), U(X) eller D- eller L-CaMeU(X); og

aa₅ er fraværende eller Pro eller U(X); eller

(f) X omfatter 1-alkylmelibiouronsyre; (**SEQ. ID. NO. 154**)

30 aa₁ er Dmt, Nα-Me-Dmt eller N,Nα-diMe-Dmt;

aa₂ er D-Arg, D-Cit, D-Lys(X) eller D-Orn(X);

aa₃ er Phe eller Tmp;

aa₄ er Phe, Tmp, Lys(X) eller Orn(X); og

aa₅ er fraværende eller Pro, Lys(X) eller Orn(X); eller

(g) U er en naturlig eller unaturlig dibasisk aminosyre; (**SEQ. ID. NO. 155**)

X er et tensid i 1-alkylglycosid-klassen, hvor 1-alkyl er substitueret eller ikke-substitueret C₁-C₂₀-alkyl eller en substitueret eller ikke-substitueret aralkylgruppe;

5 Z er NH₂;

aa₁ er Tyr, Dmt, N-R³-Tyr, N-R³-Dmt, N-(R³)₂-Tyr eller N-(R³)₂-Dmt;

aa₂ er Tic eller Tic(Ψ[CH₂-NH]);

aa₃ er Phe, Tmp eller Ψ-Phe, når aa₂ er Tic(Ψ[CH₂-NH]);

aa₄ er Phe, Tmp, D- eller L-Nal(1), D- eller L-Nal(2) eller U(X); og

10 aa₅ er fraværende eller Pro, Aib eller U(X); eller

(h) X omfatter 1-alkylmelibiouronsyre; (**SEQ. ID. NO. 156**)

aa₁ er Tyr, Dmt, Nα-Me-Tyr, Nα-Me-Dmt, N,Nα-diMe-Tyr eller N,Nα-diMe-Dmt;

aa₂ er Tic eller Tic(Ψ[CH₂-NH]);

15 aa₃ er Phe, Tmp eller Ψ-Phe, når aa₂ er Tic(Ψ[CH₂-NH]);

aa₄ er Phe, Tmp, D- eller L-Nal(1), D- eller L-Nal(2), Lys(X) eller Orn(X); og

aa₅ er fraværende eller Aib, Lys(X) eller Orn(X); eller

(i) X omfatter 1-alkylmelibiouronsyre; (**SEQ. ID. NO. 157**)

aa₂ er Tic eller Tic(Ψ[CH₂-NH]);

20 aa₃ er Phe eller Ψ-Phe;

aa₄ er Lys(X); og

aa₅ er fraværende; eller

(j) X omfatter 1-alkylmelibiouronsyre, såsom 1-eicosyl-beta-D-

melibiouronsyre, 1-octadecyl-beta-D-melibiouronsyre, 1-hexadecyl-beta-D-

25 melibiouronsyre, 1-tetradecyl-beta-D-melibiouronsyre, 1-dodecyl-beta-D-

melibiouronsyrer, 1-decyl-beta-D-melibiouronsyre eller 1-octyl-beta-D-

melibiouronsyre; (**SEQ. ID. NO. 158**)

aa₂ er Tic;

aa₃ er Phe;

30 aa₄ er Lys(X); og

aa₅ er fraværende.

8. Peptidprodukt ifølge et hvilket som helst af de foregående krav, hvor melibiose- eller melibiouronsyregruppen er bundet til peptidet via en amidbinding.

9. Peptidprodukt ifølge et hvilket som helst af de foregående krav, hvor tensidet X omfatter et 1-alkylglycosid, såsom 1-eicosyl-beta-D-melibiouronsyre, 1-octadecyl-beta-D-melibiouronsyre, 1-hexadecyl-beta-D-melibiouronsyre, 1-tetradecyl-beta-D-melibiouronsyre, 1-dodecyl-beta-D-melibiouronsyre, 1-decyl-beta-D-melibiouronsyre, 1-octyl-beta-D-melibiouronsyre eller funktionaliseret 1-eicosyl-beta-D-melibiose, 1-octadecyl-beta-D-melibiose, 1-hexadecyl-beta-D-melibiose, 1-tetradecyl-beta-D-melibiose, 1-dodecyl-beta-D-melibiose, 1-decyl-beta-D-melibiose eller 1-octyl-beta-D-melibiose.

10. Farmaceutisk sammensætning, der omfatter en terapeutisk effektiv mængde af et peptidprodukt ifølge et hvilket som helst af de foregående krav eller et farmaceutisk acceptabelt salt deraf og mindst en farmaceutisk acceptabel bærer eller excipients.

11. Fremgangsmåde til forbedring af varigheden af virkningen, biotilgængeligheden eller stabiliteten af et peptid, hvilken fremgangsmåde omfatter kovalent binding af et tensid X til peptidet, hvor X er som defineret i et hvilket som helst af kravene 1 til 9.

12. Peptidprodukt ifølge et hvilket som helst af kravene 2 til 9 til anvendelse ved behandling af smerte, hvor peptidproduktets peptid er et opioidt peptid med en naturligt eller ikke-naturligt forekommende aminosyresekvens eller -struktur.

13. Peptidprodukt ifølge et hvilket som helst af kravene 1 til 9, der udviser forbedret varighed af virkningen, biotilgængelighed, stabilitet, receptorbindingskraft eller -selektivitet eller celleaktiveringskraft sammenlignet med det ikke-modificerede peptid, der ikke er kovalent bundet til tensidet X.

14. Peptidprodukt ifølge krav 2, der udviser arrestinpåvirket agonisme eller antagonisme sammenlignet med det ikke-modificerede peptid, der ikke er kovalent bundet til tensidet X, hvor peptidet er et glucagon- eller GLP-peptid med en naturligt

eller ikke-naturligt forekommende aminosyresekvens eller -struktur.

DRAWINGS

Figure 1

Table 1

Compound	SEQ. ID. NO.	1	2	3	4	
EU-A101	2	Dmt	Pro	Tmp	Lys(C1-Glucuronyl)	NH2
EU-A102	3	Dmt	Pro	Tmp	Lys(C8-Glucuronyl)	NH2
EU-A103	4	Dmt	Pro	Tmp	Lys(C12-Glucuronyl)	NH2
EU-A105	5	Dmt	Pro	Tmp	Phe	Lys(C1-Glucuronyl)#
EU-A106	6	Dmt	Pro	Tmp	Phe	Lys(C12-Glucuronyl)#
EU-A107	7	Dmt	D-Lys(C1-Glucuronyl)	TMP	Phe	NH2
EU-A108	8	Dmt	D-Lys(C12-Glucuronyl)	TMP	Phe	NH2
EU-A109	9	Dmt	D-Lys(C8-Glucuronyl)	TMP	Phe	NH2
EU-A110	10	Dmt	D-Lys(C10-Glucuronyl)	TMP	Phe	NH2
EU-A111	11	Dmt	D-Lys(C16-Glucuronyl)	TMP	Phe	NH2
EU-A112	12	Dmt	D-Lys(C18-Glucuronyl)	TMP	Phe	NH2
EU-A113	13	Dmt	D-Lys(C20-Glucuronyl)	TMP	Phe	NH2
EU-A114	14	Dmt	D-Lys(C22-Glucuronyl)	TMP	Phe	NH2
EU-A115	15	Dmt	D-Lys(C24-Glucuronyl)	TMP	Phe	NH2
EU-A116	16	Dmt	D-Lys(OPEG2k-Glucuronyl)	TMP	Phe	NH2
EU-A117	17	Dmt	D-Lys(OPEG30k-Glucuronyl)	TMP	Phe	NH2
EU-A118	18	Dmt	D-Lys(C1-Glucuronyl)	Nal(1)	Phe	NH2
EU-A119	19	Dmt	D-Lys(C12-Glucuronyl)	Nal(1)	Phe	NH2
EU-A120	20	Dmt	D-Lys(C8-Glucuronyl)	Nal(1)	Phe	NH2
EU-A121	21	Dmt	D-Lys(C10-Glucuronyl)	Nal(1)	Phe	NH2
EU-A122	22	Dmt	D-Lys(C16-Glucuronyl)	Nal(1)	Phe	NH2
EU-A123	23	Dmt	D-Lys(C18-Glucuronyl)	Nal(1)	Phe	NH2
EU-A124	24	Dmt	D-Lys(C20-Glucuronyl)	Nal(1)	Phe	NH2

Figure 1(continued)

Table 1 (continued)

Compound	SEQ. ID. NO.	1	2	3	4	
EU-A125	25	Dmt	D-Lys(C22-Glucuronyl)	Nal(1)	Phe	NH2
EU-A126	26	Dmt	D-Lys(C24-Glucuronyl)	Nal(1)	Phe	NH2
EU-A127	27	Dmt	D-Lys(OPEG2k-Glucuronyl)	Nal(1)	Phe	NH2
EU-A128	28	Dmt	D-Lys(OPEG30k-Glucuronyl)	Nal(1)	Phe	NH2
EU-A129	29	Dmt	D-Lys(C1-Glucuronyl)	TMP	Lys	NH2
EU-A130	30	Dmt	D-Lys(C12-Glucuronyl)	TMP	Lys	NH2
EU-A131	31	Dmt	D-Lys(C8-Glucuronyl)	TMP	Lys	NH2
EU-A132	32	Dmt	D-Lys(C10-Glucuronyl)	TMP	Lys	NH2
EU-A133	33	Dmt	D-Lys(C16-Glucuronyl)	TMP	Lys	NH2
EU-A134	34	Dmt	D-Lys(C18-Glucuronyl)	TMP	Lys	NH2
EU-A135	35	Dmt	D-Lys(C20-Glucuronyl)	TMP	Lys	NH2
EU-A136	36	Dmt	D-Lys(C22-Glucuronyl)	TMP	Lys	NH2
EU-A137	37	Dmt	D-Lys(C24-Glucuronyl)	TMP	Lys	NH2
EU-A138	38	Dmt	D-Lys(OPEG2k-Glucuronyl)	TMP	Lys	NH2
EU-A139	39	Dmt	D-Lys(OPEG30k-Glucuronyl)	TMP	Lys	NH2
EU-A140	40	Dmt	D-Lys(C1-Glucuronyl)	Nal(2)	Lys	NH2
EU-A141	41	Dmt	D-Lys(C12-Glucuronyl)	Nal(2)	Lys	NH2
EU-A142	42	Dmt	D-Lys(C8-Glucuronyl)	Nal(2)	Lys	NH2
EU-A143	43	Dmt	D-Lys(C10-Glucuronyl)	Nal(2)	Lys	NH2
EU-A144	44	Dmt	D-Lys(C16-Glucuronyl)	Nal(2)	Lys	NH2
EU-A145	45	Dmt	D-Lys(C18-Glucuronyl)	Nal(2)	Lys	NH2
EU-A146	46	Dmt	D-Lys(C20-Glucuronyl)	Nal(2)	Lys	NH2
EU-A147	47	Dmt	D-Lys(C22-Glucuronyl)	Nal(2)	Lys	NH2
EU-A148	48	Dmt	D-Lys(C24-Glucuronyl)	Nal(2)	Lys	NH2

Figure 1(continued)

Table 1 (continued)

Compound	SEQ. ID. NO.	1	2	3	4	
EU-A149	49	Dmt	D-Lys(OPEG2k-Glucuronyl)	Nal(2)	Lys	NH2
EU-A150	50	Dmt	D-Lys(OPEG30k-Glucuronyl)	Nal(2)	Lys	NH2
EU-A151	51	Dmt	D-Lys(C1-Glucuronyl)	Phe	Lys	NH2
EU-A152	52	Dmt	D-Lys(C12-Glucuronyl)	Phe	Lys	NH2
EU-A153	53	Dmt	D-Lys(C8-Glucuronyl)	Phe	Lys	NH2
EU-A154	54	Dmt	D-Lys(C10-Glucuronyl)	Phe	Lys	NH2
EU-A155	55	Dmt	D-Lys(C16-Glucuronyl)	Phe	Lys	NH2
EU-A156	56	Dmt	D-Lys(C18-Glucuronyl)	Phe	Lys	NH2
EU-A157	57	Dmt	D-Lys(C20-Glucuronyl)	Phe	Lys	NH2
EU-A158	58	Dmt	D-Lys(C22-Glucuronyl)	Phe	Lys	NH2
EU-A159	59	Dmt	D-Lys(C24-Glucuronyl)	Phe	Lys	NH2
EU-A160	60	Dmt	D-Lys(OPEG2k-Glucuronyl)	Phe	Lys	NH2
EU-A161	61	Dmt	D-Lys(OPEG30k-Glucuronyl)	Phe	Lys	NH2
EU-A162	62	Dmt	Pro	Phe	Lys(C1-Glucuronyl)	NH2
EU-A163	63	Dmt	Pro	Phe	Lys(C8-Glucuronyl)	NH2
EU-A164	64	Dmt	Pro	Phe	Lys(C12-Glucuronyl)	NH2
EU-A165	65	Dmt	Pro	Phe	D-Lys(C1-Glucuronyl)	NH2
EU-A166	66	Dmt	Pro	Phe	D-Lys(C8-Glucuronyl)	NH2
EU-A167	67	Dmt	Pro	Phe	D-Lys(C12-Glucuronyl)	NH2
EU-A168	68	Dmt	Pro	Phe	Lys(C1-Glucuronyl)	Pro#
EU-A169	69	Dmt	Pro	Phe	Lys(C8-Glucuronyl)	Pro#
EU-A170	70	Dmt	Pro	Phe	Lys(C12-Glucuronyl)	Pro#
EU-A171	71	Dmt	Pro	Phe	MeLys(C1-Glucuronyl)	NH2
EU-A172	72	Dmt	Pro	Phe	MeLys(C8-Glucuronyl)	NH2

Figure 1(continued)

Table 1 (continued)

Compound	SEQ. ID. NO.	1	2	3	4	
EU-A173	73	Dmt	Pro	Phe	MeLys(C12- Glucuronyl)	NH2
EU-A174	74	Dmt	Pro	Phe	Phe	Lys(C1- Glucuronyl)#
EU-A175	75	Dmt	Pro	Phe	Phe	Lys(C12- Glucuronyl)#
EU-A176	76	Dmt	Pro	Phe	Phe	MeLys(C1- Glucuronyl)#
EU-A177	77	Dmt	Pro	Phe	Phe	MeLys(C12- Glucuronyl)#
EU-A178	78	Dmt	Tic	Phe	Lys(C1- Glucuronyl)	NH2
EU-A179	79	Dmt	Tic	Phe	Lys(C12- Glucuronyl)	NH2
EU-A180	80	Dmt	Tic	Phe	Lys(C8- Glucuronyl)	NH2
EU-A181	81	Dmt	Tic	Phe	Lys(C10- Glucuronyl)	NH2
EU-A182	82	Dmt	Tic	Phe	Lys(C16- Glucuronyl)	NH2
EU-A183	83	Dmt	Tic	Phe	Lys(C18- Glucuronyl)	NH2
EU-A184	84	Dmt	Tic	Phe	Lys(C20- Glucuronyl)	NH2
EU-A185	85	Dmt	Tic	Phe	Lys(C22- Glucuronyl)	NH2
EU-A186	86	Dmt	Tic	Phe	Lys(C24- Glucuronyl)	NH2
EU-A187	87	Dmt	Tic	Phe	Lys(OPEG2k- Glucuronyl)	NH2
EU-A188	88	Dmt	Tic	Phe	Lys(OPEG30 k- Glucuronyl)	NH2
EU-A189	89	Dmt	Tic	Phe	Phe	Lys(C1- Glucuronyl)#
EU-A190	90	Dmt	Tic	Phe	Phe	Lys(C12- Glucuronyl)#
EU-A191	91	Dmt	Tic	Phe	Phe	Lys(C8- Glucuronyl)#
EU-A192	92	Dmt	Tic	Phe	Phe	Lys(C10- Glucuronyl)#
EU-A193	93	Dmt	Tic	Phe	Phe	Lys(C16- Glucuronyl)#
EU-A194	94	Dmt	Tic	Phe	Phe	Lys(C18- Glucuronyl)#
EU-A195	95	Dmt	Tic	Phe	Phe	Lys(C20- Glucuronyl)#

Figure 1(continued)

Table 1 (continued)

Compound	SEQ. ID. NO.	1	2	3	4	
EU-A196	96	Dmt	Tic	Phe	Phe	Lys(C22-Glucuronyl)#
EU-A197	97	Dmt	Tic	Phe	Phe	Lys(C24-Glucuronyl)#
EU-A198	98	Dmt	Tic	Phe	Phe	Lys(OPEG2k-Glucuronyl)#
EU-A199	99	Dmt	Tic	Phe	Phe	Lys(OPEG30k-Glucuronyl)#
EU-A600	100	Dmt	Tic	Phe	Lys(C1-Glucuronyl)	Aib#
EU-A601	101	Dmt	Tic	Phe	Lys(C8-Glucuronyl)	Aib#
EU-A602	102	Dmt	Tic	Phe	Lys(C10-Glucuronyl)	Aib#
EU-A603	103	Dmt	Tic	Phe	Lys(C12-Glucuronyl)	Aib#
EU-A604	104	Dmt	Tic	Phe	Lys(C16-Glucuronyl)	Aib#
EU-A605	105	Dmt	Tic	Phe	Lys(C1-Glucuronyl)	Ac5c#
EU-A606	106	Dmt	Tic	Phe	Lys(C8-Glucuronyl)	Ac5c#
EU-A607	107	Dmt	Tic	Phe	Lys(C10-Glucuronyl)	Ac5c#
EU-A608	108	Dmt	Tic	Phe	Lys(C12-Glucuronyl)	Ac5c#
EU-A609	109	Dmt	Tic	Phe	Lys(C16-Glucuronyl)	Ac5c#
EU-A610	110	Dmt	Tic	Phe	D-Lys(C1-Glucuronyl)	Aib#
EU-A611	111	Dmt	Tic	Phe	D-Lys(C8-Glucuronyl)	Aib#
EU-A612	112	Dmt	Tic	Phe	D-Lys(C10-Glucuronyl)	Aib#
EU-A613	113	Dmt	Tic	Phe	D-Lys(C12-Glucuronyl)	Aib#
EU-A614	114	Dmt	Tic	Phe	D-Lys(C16-Glucuronyl)	Aib#
EU-A615	115	Dmt	Tic	Phe	D-Lys(C1-Glucuronyl)	NH2
EU-A616	116	Dmt	Tic	Phe	D-Lys(C8-Glucuronyl)	NH2
EU-A617	117	Dmt	Tic	Phe	D-Lys(C10-Glucuronyl)	NH2
EU-A618	118	Dmt	Tic	Phe	D-Lys(C12-Glucuronyl)	NH2
EU-A619	119	Dmt	Tic	Phe	D-Lys(C16-Glucuronyl)	NH2

Figure 1(continued)

Table 1 (continued)

Compound	SEQ. ID. NO.	1	2	3	4	
EU-A620	120	Dmt	Tic	Phe	Lys(C1- Glucuronyl)	NH-CH2-Ph
EU-A621	121	Dmt	Tic	Phe	Lys(C8- Glucuronyl)	NH-CH2-Ph
EU-A622	122	Dmt	Tic	Phe	Lys(C10- Glucuronyl)	NH-CH2-Ph
EU-A623	123	Dmt	Tic	Phe	Lys(C12- Glucuronyl)	NH-CH2-Ph
EU-A624	124	Dmt	Tic	Phe	Lys(C16- Glucuronyl)	NH-CH2-Ph
EU-A625	125	Dmt	Tic	Phe	Lys(C1- Glucuronyl)	NH-CH2-Ph-F
EU-A626	126	Dmt	Tic	Phe	Lys(C8- Glucuronyl)	NH-CH2-Ph-F
EU-A627	127	Dmt	Tic	Phe	Lys(C10- Glucuronyl)	NH-CH2-Ph-F
EU-A628	128	Dmt	Tic	Phe	Lys(C12- Glucuronyl)	NH-CH2-Ph-F
EU-A629	129	Dmt	Tic	Phe	Lys(C16- Glucuronyl)	NH-CH2-Ph-F
EU-A630	130	Dmt	D-Lys(C1- Glucuronyl)	Phe	Aib	NH2
EU-A631	131	Dmt	D-Lys(C8- Glucuronyl)	Phe	Aib	NH2
EU-A632	132	Dmt	D-Lys(C10- Glucuronyl)	Phe	Aib	NH2
EU-A633	133	Dmt	D-Lys(C12- Glucuronyl)	Phe	Aib	NH2
EU-A634	134	Dmt	D-Lys(C16- Glucuronyl)	Phe	Aib	NH2
EU-A635	135	Dmt	Pro	Phe	Lys(C1- Glucuronyl)	Aib#
EU-A636	136	Dmt	Pro	Phe	Lys(C8- Glucuronyl)	Aib#
EU-A637	137	Dmt	Pro	Phe	Lys(C10- Glucuronyl)	Aib#
EU-A638	138	Dmt	Pro	Phe	Lys(C12- Glucuronyl)	Aib#
EU-A639	139	Dmt	Tic	Phe	D-Lys(C1- Glucuronyl)	NH-CH2-Ph
EU-A640	140	Dmt	Tic	Phe	D-Lys(C8- Glucuronyl)	NH-CH2-Ph
EU-A641	141	Dmt	Tic	Phe	D-Lys(C10- Glucuronyl)	NH-CH2-Ph
EU-A642	142	Dmt	Tic	Phe	D-Lys(C12- Glucuronyl)	NH-CH2-Ph
EU-A643	143	Dmt	Tic	Phe	D-Lys(C16- Glucuronyl)	NH-CH2-Ph

Figure 1(continued)

Table 1 (continued)

Compound	SEQ. ID. NO.	1	2	3	4	
EU-A644	144	Dmt	Tic	Phe	D-Lys(C1-Glucuronyl)	NH-CH ₂ -Ph-F
EU-A645	145	Dmt	Tic	Phe	D-Lys(C8-Glucuronyl)	NH-CH ₂ -Ph-F
EU-A646	146	Dmt	Tic	Phe	D-Lys(C10-Glucuronyl)	NH-CH ₂ -Ph-F
EU-A647	147	Dmt	Tic	Phe	D-Lys(C12-Glucuronyl)	NH-CH ₂ -Ph-F
EU-A648	148	Dmt	Tic	Phe	D-Lys(C16-Glucuronyl)	NH-CH ₂ -Ph-F
EU-A649	645	Dmt	Tic	Phe	Lys(C14-glucuronyl)-	NH ₂

Where used, # means C-terminal amide

Figure 2

Table 2

SEQ. ID. NO.	1	5	10	15																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								</
--------------	---	---	----	----	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	----

Figure 2 (Continued)

Table 2 (Continued)

	SEQ. ID. NO.	1			5				10			15					20					
EU-229	204	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	Lys(C16)	Aib-#		
EU-230	205	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	Lys(C12)	Aib-#			
EU-231	206	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Lys(C12)	Aib-#			
EU-232	207	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Aib	Lys(C12)	Aib-#		
EU-233	208	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	F	I	Q	Lys(C12)	Aib-#			
EU-234	209	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	F	I	Q	Aib	Lys(C12)	Aib-#		
EU-235	210	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	Lys(C12)	Aib-#			
EU-236	211	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	Aib	Lys(C12)	Aib-#		
EU-237	212	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	Aib	R	F	I	Q	Lys(C12)	Aib-#			
EU-238	213	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	Aib	R	F	I	Q	Aib	Lys(C12)	Aib-#		
EU-239	214	Ac5c	V	Aib	E	H	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Lys(C12)	Aib-#			
EU-240	215	Ac5c	V	Aib	E	H	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Aib	Lys(C12)	Aib-#		
EU-241	216	Ac5c	V	Aib	E	H	Q	L	Nle	H	Q	hR	A	R	F	I	Q	Lys(C12)	Aib-#			
EU-242	217	Ac5c	V	Aib	E	H	Q	L	Nle	H	Q	hR	A	R	F	I	Q	Aib	Lys(C12)	Aib-#		
EU-243	218	Ac5c	V	Aib	E	H	Q	L	L	H	Q	hR	A	R	W	I	Q	Lys(C12)	Aib-#			
EU-244	219	Ac5c	V	Aib	E	H	Q	L	L	H	Q	hR	A	R	W	I	Q	Aib	Lys(C12)	Aib-#		
EU-245	220	Ac5c	V	Aib	E	H	Q	L	L	H	Q	hR	A	R	F	I	Q	Lys(C12)	Aib-#			
EU-246	221	Ac5c	V	Aib	E	H	Q	L	L	H	Q	hR	A	R	F	I	Q	Aib	Lys(C12)	Aib-#		
EU-247	222	Ac5c	V	Aib	E	H	Q	L	L	H	Q	hR	Aib	R	W	I	Q	Lys(C12)	Aib-#			
EU-248	223	Ac5c	V	Aib	E	H	Q	L	L	H	Q	hR	Aib	R	W	I	Q	Aib	Lys(C12)	Aib-#		
EU-249	224	Ac5c	V	Aib	E	H	Q	L	L	H	Q	hR	Aib	R	F	I	Q	Lys(C12)	Aib-#			
EU-250	225	Ac5c	V	Aib	E	H	Q	L	L	H	Q	hR	Aib	R	F	I	Q	Aib	Lys(C12)	Aib-#		
EU-251	226	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Aib	Lys(C12)	Aib-#		
EU-252	227	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	L	Lys(C12)	R	Aib-#
EU-253	228	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	L	Lys(C14)	R	Aib-#
EU-254	229	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	L	Lys(C16)	R	Aib-#
EU-255	230	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	Aib	Lys(C12)	R	Aib-#
EU-256	231	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	Aib	Lys(C14)	R	Aib-#
EU-257	232	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	Aib	Lys(C16)	R	Aib-#
EU-257	232	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	Ac5c	Lys(C12)	R	Aib-#

Figure 2 (Continued)
Table 2 (Continued)

SEQ. ID. NO.	1			5			10				15				20	)						
EU-258	233	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	Ac5c	Lys(C14)	R	Aib-#
EU-259	234	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	Ac5c	Lys(C16)	R	Aib-#
EU-260	235	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	L	Lys(C12)	R	Ac5c-#
EU-261	236	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	L	Lys(C14)	R	Ac5c-#
EU-262	237	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	D	L	Lys(C16)	R	Ac5c-#
EU-263	238	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	L	Lys(C14)	R	Ac5c-#
EU-264	239	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	L	Lys(C16)	R	Ac5c-#
EU-265	240	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	L	Lys(C12)	R	Ac5c-#
EU-266	241	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	L	Lys(C14)	R	Aib-#
EU-267	242	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	L	Lys(C16)	R	Aib-#
EU-268	243	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	Aib	Lys(C12)	R	Aib-#
EU-269	244	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	Aib	Lys(C14)	R	Aib-#
EU-270	245	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	Aib	Lys(C16)	R	Aib-#
EU-271	246	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	Ac5c	Lys(C12)	R	Aib-#
EU-272	247	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	Ac5c	Lys(C14)	R	Aib-#
EU-273	248	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	Ac5c	Lys(C16)	R	Aib-#
EU-274	249	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	A	R	W	I	Q	D	L	Lys(C12)	R	Aib-#
EU-275	250	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	A	R	W	I	Q	D	L	Lys(C14)	R	Aib-#
EU-276	251	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	A	R	W	I	Q	D	L	Lys(C16)	R	Aib-#
EU-277	252	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	Aib	R	W	I	Q	D	L	Lys(C12)	R	Aib-#
EU-278	253	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	Aib	R	W	I	Q	D	L	Lys(C14)	R	Aib-#
EU-279	254	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	Aib	R	W	I	Q	D	L	Lys(C16)	R	Aib-#
EU-280	255	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	Aib	R	W	I	N	D	L	Lys(C12)	R	Aib-#
EU-281	256	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	Aib	R	W	I	N	D	L	Lys(C14)	R	Aib-#
EU-282	257	Ac5c	V	Aib	E	I	Q	L	L	H	Q	hR	Aib	R	W	I	N	D	L	Lys(C16)	R	Aib-#
EU-283	258	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	K*	I	Q	D	E*	Lys(C12)	R	Aib-#
EU-284	259	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	Aib	R	W	I	Q	D	K*	R	R	E*-F-L- Lys(C12)-Aib-#
EU-286	260	Ac5c	V	Aib	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Aib	Lys(C14)	Aib-#		

Figure 2 (Continued)
Table 2 (Continued)

SEQ. ID. NO.	1			5			10			15					20					
EU-287	261	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Alb	Lys(C16)	Alb-#
EU-288	262	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Alb	Lys(C18)	Alb-#
EU-289	263	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	F	I	Q	Alb	Lys(C12)	Alb-#
EU-290	264	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	F	I	Q	Alb	Lys(C14)	Alb-#
EU-291	265	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	F	I	Q	Alb	Lys(C16)	Alb-#
EU-292	266	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	F	I	Q	Alb	Lys(C18)	Alb-#
EU-293	267	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	Nal2	I	Q	Alb	Lys(C12)	Alb-#
EU-294	268	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	Nal2	I	Q	Alb	Lys(C14)	Alb-#
EU-295	269	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	Nal2	I	Q	Alb	Lys(C16)	Alb-#
EU-296	270	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	Nal2	I	Q	Alb	Lys(C18)	Alb-#
EU-297	271	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	E*	R	W	I	K*	Alb	Lys(C12)	Alb-#
EU-298	272	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	E*	R	W	I	K*	Alb	Lys(C14)	Alb-#
EU-299	273	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	E*	R	W	I	K*	Alb	Lys(C16)	Alb-#
EU-900	274	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	E*	R	W	I	K*	Alb	Lys(C18)	Alb-#
EU-901	275	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Alb	Lys(C12)	Alb-#
EU-902	276	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Alb	Lys(C14)	Alb-#
EU-903	277	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Alb	Lys(C16)	Alb-#
EU-904	278	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	A	R	W	I	Q	Alb	Lys(C18)	Alb-#
EU-905	279	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	E*	R	W	I	K*	Alb	Lys(C12)	Alb-#
EU-906	280	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	E*	R	W	I	K*	Alb	Lys(C14)	Alb-#
EU-907	281	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	E*	R	W	I	K*	Alb	Lys(C16)	Alb-#
EU-908	282	Ac5c	V	Alb	E	I	Q	L	Nle	H	Q	hR	E*	R	W	I	K*	Alb	Lys(C18)	Alb-#

hR refers to homo-arginine

Figure 4

AGONIST EFFECT OF COMPOUND EU-232
AT THE HUMAN PTH1 RECEPTOR

$EC_{50} = 2.4E-09$ M
 $A = 0.0$ $D = 127.3$




Figure 5

AGONIST EFFECT OF COMPOUND EU-285
AT THE HUMAN PTH1 RECEPTOR

$EC_{50} = 2.9E-09 \text{ M}$

$A = 0.0$ $D = 90.5$




Figure 6




Figure 7




Figure 8 (Continued)

Table 3 (Continued)

SEQ. ID. NO.	1	5	10	15	20
EU-A328	H Aib	Q G T F T S D	Bip	L	Lys(C8)-#
EU-A329	H Aib	Q G T F T S D	Bip	L	Lys(C12)-#
EU-A330	H Aib	Q G T F T S D	Bip	L	Lys(C16)-#
EU-A331	H Aib	Q G T F T S D	NaI(2)	L	Lys(C8)-#
EU-A332	H Aib	Q G T F T S D	NaI(2)	L	Lys(C12)-#
EU-A333	H Aib	Q G T F T S D	NaI(2)	L	Lys(C16)-#
EU-A334	H Aib	Q G T 26FF T S D	Bip2Et4MeO	L	Lys(C8)-#
EU-A335	H Aib	Q G T 26FF T S D	Bip2Et4MeO	L	Lys(C12)-#
EU-A336	H Aib	Q G T 26FF T S D	Bip2Et4MeO	L	Lys(C16)-#
EU-A337	H Aib	Q G T F T S D	L	L	Lys(C8)-#
EU-A338	H Aib	Q G T F T S D	L	L	Lys(C12)-#
EU-A339	H Aib	Q G T F T S D	L	L	Lys(C16)-#
EU-A340	H Aib	Q G T F T S D	Bip	L	Lys(C8)-#
EU-A341	H Aib	Q G T F T S D	Bip	L	Lys(C12)-#
EU-A342	H Aib	Q G T F T S D	Bip	L	Lys(C16)-#
EU-A343	H Aib	Q G T F T S D	NaI(2)	L	Lys(C8)-#
EU-A344	H Aib	Q G T F T S D	NaI(2)	L	Lys(C12)-#
EU-A345	H Aib	Q G T F T S D	NaI(2)	L	Lys(C16)-#
EU-A346	H Aib	Q G T 26FF T S D	Bip2Et4MeO	L	Lys(C8)-#
EU-A347	H Aib	Q G T 26FF T S D	Bip2Et4MeO	L	Lys(C12)-#
EU-A348	H Aib	Q G T 26FF T S D	Bip2Et4MeO	L	Lys(C16)-#
EU-A349	H Aib	Q G T 26MeF T S D	Bip	L	Lys(C8)-#
EU-A350	H Aib	Q G T 26MeF T S D	Bip	L	Lys(C12)-#
EU-A351	H Aib	Q G T 26MeF T S D	Bip	L	Lys(C16)-#
EU-A352	H Aib	Q G T MeF T S D	L	L	Lys(C8)-#
EU-A353	H Aib	Q G T MeF T S D	L	L	Lys(C12)-#
EU-A354	H Aib	Q G T MeF T S D	L	L	Lys(C16)-#
EU-A355	H Aib	Q G T MeF T S D	Bip	L	Lys(C8)-#
EU-A356	H Aib	Q G T MeF T S D	Bip	L	Lys(C12)-#

Figure 8 (Continued)

Table 3 (Continued)

SEQ. ID. NO.	1			5				10					15			20
EU-A357	363	H	Aib	Q	G	T	MeF	T	S	D	Bip	S	K	Y	L	
EU-A358	364	H	Aib	Q	G	T	MeF	T	S	D	Nal(2)	S	K	Y	L	
EU-A359	365	H	Aib	Q	G	T	MeF	T	S	D	Nal(2)	S	K	Y	L	
EU-A360	366	H	Aib	Q	G	T	MeF	T	S	D	Nal(2)	S	K	Y	L	
EU-A361	367	H	Aib	Q	G	T	MeF	T	S	D	Bip2E4MeO	S	K	Y	L	
EU-A362	368	H	Aib	Q	G	T	MeF	T	S	D	Bip2E4MeO	S	K	Y	L	
EU-A363	369	H	Aib	Q	G	T	MeF	T	S	D	Bip2E4MeO	S	K	Y	L	
EU-A364	370	H	Aib	Q	G	T	F	T	S	D	V	S	K	Y	L	
EU-A365	371	H	Aib	Q	G	T	F	T	S	D	L	S	K	Y	L	
EU-A366	372	H	Aib	Q	G	T	F	T	S	D	L	S	K	Y	L	
EU-A367	373	H	Aib	Q	G	T	F	T	S	D	L	S	K	Y	L	
EU-A368	374	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A369	375	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A370	376	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A371	377	H	Aib	Q	G	T	F	T	S	D	Nal(2)	S	K	Y	L	
EU-A372	378	H	Aib	Q	G	T	F	T	S	D	Nal(2)	S	K	Y	L	
EU-A373	379	H	Aib	Q	G	T	F	T	S	D	Nal(2)	S	K	Y	L	
EU-A374	380	H	Aib	Q	G	T	26FF	T	S	D	Bip2E4MeO	S	K	Y	L	
EU-A375	381	H	Aib	Q	G	T	26FF	T	S	D	Bip2E4MeO	S	K	Y	L	
EU-A376	382	H	Aib	Q	G	T	26FF	T	S	D	Bip2E4MeO	S	K	Y	L	
EU-A377	383	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A378	384	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A379	385	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A380	386	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A381	387	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A382	388	H	Aib	Q	G	T	F	T	S	D	Bip	S	K	Y	L	
EU-A383	389	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	
EU-A384	390	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	
EU-A385	391	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	
EU-A386	392	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	

Figure 8 (Continued)

Table 3 (Continued)

SEQ. ID. NO.	1				5					10					15			20
EU-A387	393	H	Alb	Q	G	T	F	T	S	D	Bip	S	K	Y	L	S	Lys(C12)-#	
EU-A388	394	H	Alb	Q	G	T	MeF	T	S	D	Bip	S	K	Y	L	S	Lys(C12)-#	
EU-A389	395	H	Alb	Q	G	T	MeF	T	S	D	Bip	S	Alb	Y	L	S	Lys(C12)-#	
EU-A390	396	H	Alb	Q	G	T	MeF	T	S	D	Bip	S	K	Y	L	Alb	Lys(C12)-#	
EU-A391	397	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	Alb	Lys(C12)-#	
EU-A392	398	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	L	D	Lys(C12)-#	
EU-A393	399	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	Lys(C12)-#					
EU-A394	400	H	Alb	Q	G	T	MeF	T	S	D	Nal(2)	Alb	Lys(C12)-#					
EU-A395	401	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	Lys(C12)	Alb-#
EU-A396	402	H	Alb	Q	G	T	MeF	T	S	D	Nal(2)	S	K	Y	L	D	Lys(C12)	Alb-#
EU-A397	403	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	L	D	Lys(C12)-#	
EU-A398	404	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	L	D	Lys(C12)	Alb-#
EU-A399	405	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	Lys(C12)	Alb-#
EU-A400	406	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	R	Lys(C12) Alb-#
EU-A401	407	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	L	D	R	Lys(C12) Alb-#
EU-A402	408	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	L	D	R	Lys(C12) Ac5c-#
EU-A403	409	H	Alb	Q	G	T	MeF	T	S	D	Nal(2)	Alb	K	Y	L	E	R	Lys(C12)-#
EU-A404	410	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Har	Lys(C8) Alb-#
EU-A405	411	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Har	Lys(C12) Alb-#
EU-A406	412	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Har	Lys(C16) Alb-#
EU-A407	413	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	Lys(C12)-#			
EU-A408	414	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	Lys(C12)	-NHet		
EU-A409	415	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	Lys(C12)	Alb-#		
EU-A410	416	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	Lys(C12)	Alb		
EU-A411	417	H	Alb	Q	G	T	MeF	T	S	D	Y	Alb	K	Y	Lys(C12)	Ac5c-#		
EU-A412	418	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	Alb	Lys(C8)-#
EU-A413	419	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	Alb	Lys(C16)-#
EU-A414	420	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Alb	Lys(C8)-#
EU-A415	421	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Alb	Lys(C12)-#
EU-A416	422	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Alb	Lys(C16)-#

Figure 8 (Continued)
Table 3 (Continued)

	SEQ. ID. NO.	1				5				10						15				20	
	EU-A417	423	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C8)	Alb-#
	EU-A418	424	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C12)	Alb-#
	EU-A419	425	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C16)	Alb-#
	EU-A420	426	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C8)	Ac5c-#
	EU-A421	427	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C12)	Ac5c-#
	EU-A422	428	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C16)	Ac5c-#
	EU-A423	429	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C8)	Ac5c-#
	EU-A424	430	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C12)	Ac5c-#
	EU-A425	431	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Alb	Lys(C16)	Ac5c-#

MeF means C-alphaMe-Phe; Har means homoArg; 2,6F-F means 2',6'-difluoro-Phe; Bip2E14MeO means 2'-ethyl-4'-MeO-biphenylalanine.

Lys(C12) means N-epsilon-(1'-dodecyl beta-D-glucuronyl)-L-lysine and other C numbers mean the corresponding 1'-alkyl glucoronide.

means amide C-terminus.

Figure 9

Table 4

SEQ. ID. NO.	1			5			10			15					20				25																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
EU-A426	432	H	Alb	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	S	R	Alb	Lys(C8)	Alb-#																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		</

Figure 9 (Continued)

Table 4 (Continued)

SEQ. ID. NO.	1			5		10		15			20		25											
EU-A448	463	H	Aib	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	E*	R	Aib	Ala	K*	Lys(C12)	F	Aib#
EU-A449	464	H	Aib	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	E*	R	Aib	Ala	K*	Lys(C16)	F	Aib#
EU-A450	465	H	Aib	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	E*	R	Aib	Ala	K*	Lys(C8)	F	Ac5c#
EU-A451	466	H	Aib	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	K*	R	Aib	Ala	K*	Lys(C12)	F	Ac5c#
EU-A452	467	H	Aib	Q	G	T	MeF	T	S	D	Y	S	K	Y	L	D	K*	R	Aib	Ala	K*	Lys(C16)	F	Ac5c#
EU-A447	468	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	R	Aib	Ala	K*	Lys(C8)	F	Aib#
EU-A448	469	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	R	Aib	Ala	K*	Lys(C12)	F	Aib#
EU-A449	470	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	R	Aib	Ala	K*	Lys(C16)	F	Aib#
EU-A450	471	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	R	Aib	Ala	K*	Lys(C8)	F	Ac5c#
EU-A451	472	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	K*	R	Aib	Ala	K*	Lys(C12)	F	Ac5c#
EU-A452	473	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	K*	R	Aib	Ala	K*	Lys(C16)	F	Ac5c#
EU-A453	474	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	Lys(C12)	F	Aib#
EU-A454	475	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	Lys(C16)	F	Aib#
EU-A455	476	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	R	A	A	K*	Lys(C12)	F	Aib#
EU-A456	477	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	R	A	A	K*	Lys(C16)	F	Aib#
EU-A457	478	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	R	A	A	K*	Lys(C12)	F	Ac5c#
EU-A458	479	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	R	A	A	K*	Lys(C16)	F	Ac5c#
EU-A459	480	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	Lys(C12)	F	Ac5c#
EU-A460	481	H	Aib	Q	G	T	F	T	S	D	Y	S	R	Y	L	D	E*	R	A	A	K*	Lys(C12)	F	Aib#
EU-A461	482	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Aib	Lys(C12)	Aib#				
EU-A462	483	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C12)	Aib#				
EU-A463	484	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C12)	Ac5c#				
EU-A464	485	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C8)	Aib#				
EU-A465	486	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C16)	Aib#				
EU-A466	487	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Aib	Lys(C12)	Aib#				
EU-A467	488	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Aib	Lys(C8)	Aib#				
EU-A468	489	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Aib	Lys(C1)	Aib#				
EU-A469	490	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Aib	Lys(C16)	Aib#				
EU-A470	491	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Aib	Lys(C16)	Aib#				
EU-A471	492	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Aib	Lys(C12)	Aib#				
EU-A472	493	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Aib	Lvs(C8)	Aib#				

Figure 9 (Continued)
Table 4 (Continued)

SEQ. ID. NO.	1	5	10	15	20	25
EU-A473	494	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C16) Alb#				
EU-A474	495	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) Alb#				
EU-A475	496	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) Alb#				
EU-A476	497	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) AcSc#				
EU-A477	498	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) #				
EU-A478	499	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) #				
EU-A479	500	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C16) #				
EU-A480	501	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) #				
EU-A481	502	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) #				
EU-A482	503	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C16) #				
EU-A483	504	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) #				
EU-A484	505	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) #				
EU-A485	506	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C16) #				
EU-A486	507	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) #				
EU-A487	508	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) #				
EU-A488	509	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C16) #				
EU-A489	510	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) #				
EU-A490	511	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) #				
EU-A491	512	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C16) #				
EU-A492	513	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) #				
EU-A493	514	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) #				
EU-A494	515	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C16) #				
EU-A495	516	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) #				
EU-A496	517	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) #				
EU-A497	518	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C16) #				
EU-A498	519	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C8) #				
EU-A499	520	H Alb Q G T F T S D Y S K* Y L D E* R Lys(C12) #				

Figure 9 (Continued)

Table 4 (Continued)

SEQ. ID. NO.	1	5	10	15	20	25
EU-A501	521	H S Q G T F T S D Y S K Y L D S R R A Q Lys(C12) Alb #				
EU-A502	522	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A503	523	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A504	524	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A505	525	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A506	526	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A507	527	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A508	528	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A509	529	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A510	530	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A511	531	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A512	532	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A513	533	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A514	534	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A515	535	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A516	536	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A517	537	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A518	538	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A519	539	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A520	540	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A521	541	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A522	542	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A523	543	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A524	544	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A525	545	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A526	546	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A527	547	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A528	548	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A529	549	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				
EU-A530	550	H S Q G T F T S D Y S K Y L D S R R A Lys(C12) Alb #				

Figure 9 (Continued)
Table 4 (Continued)

SEQ. ID. NO.	1			5				10				15					20				25
EU-A532	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C12)	Ac		
EU-A533	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	hAg	Lys(C12)	Ab		
EU-A534	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ab	R	Lys(C12)	Ab		
EU-A535	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C14)	Ab		
EU-A536	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C16)	Ab		
EU-A537	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C18)	Ab		
EU-A538	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ab	Q	Lys(C12)	Ab		
EU-A539	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	Lys(C12)	A	#	
EU-A540	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	Lys(C14)	A	#	
EU-A541	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	Lys(C16)	A	#	
EU-A542	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	Lys(C18)	A	#	
EU-A543	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	Lys(C20)	A	#	
EU-A544	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C12)	Ab	#	
EU-A545	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C14)	Ab	#	
EU-A546	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C16)	Ab	#	
EU-A547	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C18)	Ab	#	
EU-A548	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	R	Lys(C20)	Ab	#	
EU-A549	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ab	Lys(C12)	#			
EU-A550	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ab	Lys(C14)	#			
EU-A551	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ab	Lys(C16)	#			
EU-A552	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ab	Lys(C18)	#			
EU-A553	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ab	Lys(C20)	#			
EU-A554	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Ab	Lys(C12)	Ab	Ab	#
EU-A555	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Ab	Lys(C14)	Ab	Ab	#
EU-A556	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Ab	Lys(C16)	Ab	Ab	#
EU-A557	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Ab	Lys(C18)	Ab	Ab	#
EU-A558	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Ab	Lys(C20)	Ab	Ab	#
EU-A559	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	S	Ab	Lys(C22)	Ab	Ab	#
EU-A560	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	Lys(C12)	Ab	#	
EU-A561	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	Lys(C14)	Ab	#	
EU-A562	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	Lys(C16)	Ab	#	
EU-A563	H	Ab	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	Lys(C18)	Ab	#	

Figure 9 (Continued)

Table 4 (Continued)

SEQ. ID. NO.	1	5	10	15	20	25																	
EU-A564	583	H	Ab	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	R	Lys(C20)	Ab	#		
EU-A565	584	H	Ab	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	Q	Lys(C12)	Ab	#		
EU-A566	585	H	Ab	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	Q	Lys(C14)	Ab	#		
EU-A567	586	H	Ab	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	Q	Lys(C16)	Ab	#		
EU-A568	587	H	Ab	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	Q	Lys(C18)	Ab	#		
EU-A569	588	H	Ab	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	Q	Lys(C20)	Ab	#		
EU-A570	589	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C12)	Ab	Ab	#	
EU-A571	590	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C14)	Ab	Ab	#	
EU-A572	591	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C16)	Ab	Ab	#	
EU-A573	592	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C18)	Ab	Ab	#	
EU-A574	593	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C20)	Ab	Ab	#	
EU-A575	594	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	hAg	Lys(C12)	Ab	Ab	#	
EU-A576	595	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	hAg	Lys(C14)	Ab	Ab	#	
EU-A577	596	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	hAg	Lys(C16)	Ab	Ab	#	
EU-A578	597	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	hAg	Lys(C18)	Ab	Ab	#	
EU-A579	598	H	Ab	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	hAg	Lys(C20)	Ab	Ab	#	
EU-A580	599	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	R	Lys(C12)	Ab	Ab	#	
EU-A581	600	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	R	Lys(C14)	Ab	Ab	#	
EU-A582	601	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	R	Lys(C16)	Ab	Ab	#	
EU-A583	602	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	R	Lys(C18)	Ab	Ab	#	
EU-A584	603	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	R	Lys(C20)	Ab	Ab	#	
EU-A585	604	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	hAg	Lys(C12)	Ab	Ab	#	
EU-A586	605	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	hAg	Lys(C14)	Ab	Ab	#	
EU-A587	606	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	hAg	Lys(C16)	Ab	Ab	#	
EU-A588	607	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	hAg	Lys(C18)	Ab	Ab	#	
EU-A589	608	H	Ac4c	Q	G	T	F	T	S	D	Y	S	K*	Y	L	D	E*	hAg	Lys(C20)	Ab	Ab	#	
EU-A590	609	H	Ac4c	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C12)	Ab	Ab	#	
EU-A591	610	H	Ac4c	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C14)	Ab	Ab	#	
EU-A592	611	H	Ac4c	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C16)	Ab	Ab	#	
EU-A593	612	H	Ac4c	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C18)	Ab	Ab	#	
EU-A594	613	H	Ac4c	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	R	Lys(C20)	Ab	Ab	#	
EU-A595	614	H	Ac4c	Q	G	T	F	T	S	D	Y	S	E*	Y	L	D	K*	hAg	Lys(C12)	Ab	Ab	#	

Figure 9 (Continued)

Table 4 (Continued)

SEQ. ID. NO.	1	5	10	15	20	25
EU-A596 615	H	AcAc Q G T F T S D Y S E* Y L D K* hAg Lys(C14) Ab #				
EU-A597 616	H	AcAc Q G T F T S D Y S E* Y L D K* hAg Lys(C16) Ab #				
EU-A598 617	H	AcAc Q G T F T S D Y S E* Y L D K* hAg Lys(C18) Ab #				
EU-A599 618	H	AcAc Q G T F T S D Y S E* Y L D K* hAg Lys(C20) Ab #				

MeF means C-alphaMe-Phe.

Lys(C12) means N-epsilon-(1'-dodecyl beta-D-glucuronyl)-L-lysine and other C numbers mean the corresponding 1'-alkyl glucuronide. The pair of amino acids E*, K* or K*, E* separated by 4 residues denotes a side chain lactam linkage formed between the side chain functional groups on these amino acids.

means amide C-terminus.

SEQ ID. NO.	1	5	10	15	20	25
EU-A700 646	H	Alb Q G T F T S D Y S K Y L D Al _b Lys(C12) b Al _b #				
EU-A701 647	H	Alb Q G T F T S D Y S K Y L D Al _b Lys(C14) b Al _b #				
EU-A702 648	H	Alb Q G T F T S D Y S K Y L D Al _b Lys(C16) b Al _b #				
EU-A703 649	H	Alb Q G T F T S D Y S K Y L D Al _b Lys(C18) b Al _b #				
EU-A704 650	H	Alb Q G T F T S D Y S K Y L D Al _b Lys(C20) b Al _b #				
EU-A705 651	H	Alb Q G T F T S D Y S K Y L D Al _b hAg Lys(C12) b Al _b #				
EU-A706 652	H	Alb Q G T F T S D Y S K Y L D Al _b hAg Lys(C14) b Al _b #				

Figure 9 (Continued)

[illegible]

Figure 9 (Continued)

Table 4 (Continued)

SEQ ID. NO.	1	5	10	15	20	25
EU- A724	H	Q G T	F T S D Y	K Y L D	hArg	Lys(C12)
EU- A725	H	Q G T	F T S D Y	K Y L D	Arg	Lys(C12)
EU- A726	H	Q G T	F T S D Y	K Y L D	Arg	Lys(C12)
EU- A727	H	Q G T	F T S D Y	K Y L D	hArg	Lys(C12)
EU- A728	H	Q G T	F T S D Y	K Y L D	hArg	Lys(C14)
EU- A729	H	Q G T	F T S D Y	K Y L D	hArg	Lys(C16)
EU- A730	H	Q G T	F T S D Y	K Y L D	hArg	Lys(C18)
EU- A731	H	Q G T	F T S D E*	K Y K D S	Arg	Lys(C12)
EU- A732	H	Q G T	F T S D E*	K Y K D S	hArg	Lys(C14)
EU- A733	H	Q G T	F T S D Y	K Y L D S	K*	Lys(C12)
EU- A734	H	Q G T	F T S D Y	K Y L D S	K*	Lys(C12)
EU- A735	H	Q G T	F T S D Y	K Y L D S	E*	Lys(C12)
EU- A736	H	Q G T	F T S D Y	K Y L D S	E*	Lys(C14)
EU- A737	H	Q G T	F T S D Y	K Y L D S	E*	Lys(C16)
EU- A738	H	Q G T	F T S D Y	K Y L D S	E*	Lys(C18)
EU- A739	H	Q G T	F T S D Y	K Y L D S	E*	Lys(C20)
EU- A740	H	Q G T	F T S D Y	K Y L D S	E*	Lys(C12)

Figure 9 (Continued)

Table 4 (Continued)

[illegible]

Figure 9 (Continued)
Table 4 (Continued)

	SEQ ID. NO.	1				5					10				15					20				25					
EU-A758	H	aMePr _o	Q	G	T	aMeF	T	S	D	Y	Lys(C8)	#																	
EU-A759	H	aMePr _o	Q	G	T	aMeF	T	S	D	Y	Lys(16)	#																	
EU-A760	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C8)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A761	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C12)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A762	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C14)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A763	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C16)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A764	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C18)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A765	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C20)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A766	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C8)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A767	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C14)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A768	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C16)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A769	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C20)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A770	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	K	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A771	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C12)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A772	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C18)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A773	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C18)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A774	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	R	A	K	D	F	V	Q	W	L	L	N	T#

Figure 9 (Continued)

Table 4 (Continued)

SEQ ID. NO.	1				5					10					15				20					25				T#	
EU-A775	721	H	G	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	R	A	Lys(C12)	D	F	V	Q	W	L	L	N	T#
EU-A776	722	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C18)	D	F	V	Q	W	L	L	N	T#
EU-A777	723	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C18)	D	F	V	Q	W	L	L	N	T#
EU-A778	724	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C8)	D	F	V	Q	W	L	L	N	T#
EU-A779	725	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C12)	D	F	V	Q	W	L	L	N	T#
EU-A780	726	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C14)	D	F	V	Q	W	L	L	N	T#
EU-A781	727	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C16)	D	F	V	Q	W	L	L	N	T#
EU-A782	728	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C18)	D	F	V	Q	W	L	L	N	T#
EU-A783	729	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C20)	D	F	V	Q	W	L	L	N	T#
EU-A784	730	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C8)	D	F	V	Q	W	L	L	N	T#
EU-A785	731	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C12)	D	F	V	Q	W	L	L	N	T#
EU-A786	732	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C14)	D	F	V	Q	W	L	L	N	T#
EU-A787	733	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C16)	D	F	V	Q	W	L	L	N	T#
EU-A788	734	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C18)	D	F	V	Q	W	L	L	N	T#
EU-A789	735	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	A	Lys(C20)	D	F	V	Q	W	L	L	N	T#
EU-A790	736	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C8)	A	A	K	E	F	I	A	W	L	N	T#
EU-A791	737	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C12)	A	A	K	E	F	I	A	W	L	N	T#

Figure 9 (Continued)

Table 4 (Continued)

	SEQ ID. NO.	1				5					10					15				20			25										
EU- A792	738	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C14)	A	A	K	E	F	I	A	W	L	L	N	T#	
EU- A793	739	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C16)	A	A	K	E	F	I	A	W	L	L	N	T#	
EU- A794	740	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C18)	A	A	K	E	F	I	A	W	L	L	N	T#	
EU- A795	741	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C20)	A	A	K	E	F	I	A	W	L	L	N	T#	
EU- A796	742	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C8)	W	L	L	M	N	T#
EU- A797	743	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C12)	W	L	L	M	N	T#
EU- A798	744	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C14)	W	L	L	M	N	T#
EU- A799	745	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C16)	W	L	L	M	N	T#
EU- A800	746	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C18)	W	L	L	M	N	T#
EU- A801	747	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C20)	W	L	L	M	N	T#
EU- A802	748	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C8)	W	L	L	L	N	T#
EU- A803	749	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C12)	W	L	L	L	N	T#
EU- A804	750	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C14)	W	L	L	L	N	T#
EU- A805	751	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C16)	W	L	L	L	N	T#

Figure 9 (Continued)

Table 4 (Continued)

	SEQ ID. NO.	1								10					15						20				25					
EU- A806	752	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C18)	W	L	L	N	T#
EU- A807	753	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C20)	W	L	L	N	T#
EU- A808	754	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(C8)	W	L	L	N	T#
EU- A809	755	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(C12)	W	L	L	N	T#
EU- A810	756	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(C14)	W	L	L	N	T#
EU- A811	757	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(C15)	W	L	L	N	T#
EU- A812	758	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(C18)	W	L	L	N	T#
EU- A813	759	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(C20)	W	L	L	N	T#
EU- A814	760	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C8)	A	A	R	E	F	I	A	W	L	L	N	T#
EU- A815	761	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C12)	A	A	R	E	F	I	A	W	L	L	N	T#
EU- A816	762	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C14)	A	A	R	E	F	I	A	W	L	L	N	T#
EU- A817	763	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C16)	A	A	R	E	F	I	A	W	L	L	N	T#
EU- A818	764	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C18)	A	A	R	E	F	I	A	W	L	L	N	T#

Figure 9 (Continued)

[illegible]

Figure 9 (Continued)

[illegible]

Figure 9 (Continued)

Table 4 (Continued)

	SEQ ID, NO.	1				5				10						15					20										
EU-A851		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C14)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A852		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C16)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A853		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C18)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A854		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C20)	A	A	R	E	F	I	A	W	L	L	N	T#
EU-A855		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C8)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A855		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C12)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A857		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C14)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A858		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C16)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A859		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C18)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A860		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Lys(C20)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A861		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ai _b	Lys(C8)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A862		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ai _b	Lys(C12)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A863		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ai _b	Lys(C14)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A864		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ai _b	Lys(C16)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A865		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ai _b	Lys(C18)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A866		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ai _b	Lys(C20)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A867		H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Ai _b	Lys(C8)	A	A	R	E	F	I	A	W	L	L	N	T#

Figure 9 (Continued)

	SEQ ID. NO.	1							10					15					20				25								
EU-A885	831	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Q	A	A	R	E	F	I	Lys(C8)	W	L	L	N	T#	
EU-A886	832	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Q	A	A	R	E	F	I	Lys(C12)	W	L	L	N	T#
EU-A887	833	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Q	A	A	R	E	F	I	Lys(C14)	W	L	L	N	T#
EU-A888	834	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Q	A	A	R	E	F	I	Lys(C16)	W	L	L	N	T#
EU-A889	835	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Q	A	A	R	E	F	I	Lys(C18)	W	L	L	N	T#
EU-A890	836	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	Q	A	A	R	E	F	I	Lys(C20)	W	L	L	N	T#
EU-A891	837	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(C8)	D	F	V	Q	W	L	L	N	T#
EU-A892	838	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(12)	D	F	V	Q	W	L	L	N	T#
EU-A893	839	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(C14)	D	F	V	Q	W	L	L	N	T#
EU-A894	840	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(C16)	D	F	V	Q	W	L	L	N	T#
EU-A895	841	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(C18)	D	F	V	Q	W	L	L	N	T#
EU-A896	842	H	G	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(C20)	D	F	V	Q	W	L	L	N	T#
EU-A897	843	H	Alb	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(C12)	D	F	V	Q	W	L	L	N	T#
EU-A898	844	H	Alb	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(C14)	D	F	V	Q	W	L	L	N	T#
EU-A899	845	H	Alb	Q	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	A	R	R	A	Lys(C16)	D	F	V	Q	W	L	L	N	T#

Figure 9 (Continued)

	SEQ ID. NO.	1			5				10					15					20					25				
EU-A900	846	H	Alb	Q	G	T				S	K	Y	L	D	A	R	R	A	Lys(C15)	D	F	V	Q	W	L	L	N	T#
EU-A901	847	H	Alb	Q	G	T				S	K	Y	L	D	A	R	R	A	Lys(C20)	D	F	V	Q	W	L	L	N	T#
EU-A902	848	H	G	Q	G	T				S	K	Y	L	D	A	R	R	A	Q	D	F	V	Lys(C8)	W	L	L	N	T#
EU-A903	849	H	G	Q	G	T				S	K	Y	L	D	A	R	R	A	Q	D	F	V	Lys(C12)	W	L	L	N	T#
EU-A904	850	H	G	Q	G	T				S	K	Y	L	D	A	R	R	A	Q	D	F	V	Lys(C14)	W	L	L	N	T#
EU-A905	851	H	G	Q	G	T				S	K	Y	L	D	A	R	R	A	Q	D	F	V	Lys(C16)	W	L	L	N	T#
EU-A906	852	H	G	Q	G	T				S	K	Y	L	D	A	R	R	A	Q	D	F	V	Lys(C18)	W	L	L	N	T#
EU-A907	853	H	G	Q	G	T				S	K	Y	L	D	A	R	R	A	Q	D	F	V	Lys(C20)	W	L	L	N	T#
EU-A908	854	H	Alb	Q	G	T				S	K	Y	L	D	Alb	Lys(C18)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A909	855	H	Alb	Q	G	T				S	K	Y	L	D	Alb	Lys(C8)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A910	856	H	Alb	Q	G	T				S	K	Y	L	D	Alb	Lys(C12)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A911	857	H	Alb	Q	G	T				S	K	Y	L	D	Alb	Lys(C14)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A912	858	H	Alb	Q	G	T				S	K	Y	L	D	Alb	Lys(C16)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A913	859	H	Alb	Q	G	T				S	K	Y	L	D	Alb	Lys(C18)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A914	860	H	Alb	Q	G	T				S	K	Y	L	D	Alb	Lys(C20)	A	A	K	E	F	I	A	W	L	L	N	T#

Figure 9 (Continued)

Table 4 (Continued)

[illegible]

Figure 9 (Continued)

Table 4 (Continued)

[illegible]

Figure 9 (Continued)

SEQ ID. NO.	1				5				10			15			20			25													
EU-A959	905	H	G		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C16)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A960	906	H	G		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C18)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A961	907	H	G		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C20)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A962	908	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C8)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A963	909	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C12)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A964	910	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C14)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A965	911	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C16)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A966	912	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C18)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A967	913	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(C20)	A	A	K	E	F	I	A	W	L	L	Q	T#
EU-A968	914	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Al _b	Lys(C8)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A969	915	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Al _b	Lys(C12)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A970	916	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Al _b	Lys(C14)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A971	917	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Al _b	Lys(C16)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A972	918	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Al _b	Lys(C18)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A973	919	H	Alb		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	Al _b	Lys(C20)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A974	920	H	G		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	R	A	Lys(C8)	D	F	V	Q	W	L	L	Q	T#
EU-A975	921	H	G		Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	R	R	A	Lys(C12)	D	F	V	Q	W	L	L	Q	T#

Figure 9 (Continued)

	SEQ ID. NO.	1	G	Q	T	F					10	S	K	Y	L	D	E	R	A		20	D	F	V	Q	W	L	L	Q	#
	EU- A976	922	H	G	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C1 4)	D	F	V	Q	W	L	L	Q	T#
	EU- A977	923	H	G	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C1 6)	D	F	V	Q	W	L	L	Q	T#
	EU- A978	924	H	G	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C1 8)	D	F	V	Q	W	L	L	Q	T#
	EU- A979	925	H	G	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C2 0)	D	F	V	Q	W	L	L	Q	T#
	EU- A980	926	H	Alib	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C8)	D	F	V	Q	W	L	L	Q	T#
	EU- A981	927	H	Alib	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C1 2)	D	F	V	Q	W	L	L	Q	T#
	EU- A982	928	H	Alib	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C1 4)	D	F	V	Q	W	L	L	Q	T#
	EU- A983	929	H	Alib	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C1 6)	D	F	V	Q	W	L	L	Q	T#
	EU- A984	930	H	Alib	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C1 8)	D	F	V	Q	W	L	L	Q	T#
	EU- A985	931	H	Alib	Q	G	T					Y	S	K	Y	L	D	E	R	A	Lys(C2 0)	D	F	V	Q	W	L	L	Q	T#
	EU- A986	932	H	Alib	Q	G	T					Y	S	K	Y	L	D	E*	Q	A	K*	E	F	I	Lys(C8) ,	W	L	M	Q	T#
	EU- A987	933	H	Alib	Q	G	T					Y	S	K	Y	L	D	E*	Q	A	K*	E	F	I	Lys(C12)	W	L	M	Q	T#
	EU- A988	934	H	Alib	Q	G	T					Y	S	K	Y	L	D	E*	Q	A	K*	E	F	I	Lys(C14)	W	L	M	Q	T#
	EU- A989	935	H	Alib	Q	G	T					Y	S	K	Y	L	D	E*	Q	A	K*	E	F	I	Lys(C16)	W	L	M	Q	T#
	EU- A990	936	H	Alib	Q	G	T					Y	S	K	Y	L	D	E*	Q	A	K*	E	F	I	Lys(C18)	W	L	M	Q	T#

Figure 9 (Continued)

Table 4 (Continued)

SEQ ID. NO.	1					5					10					15				20				25				
EU-A991	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(C20)	W	L	M	Q	T#
EU-A992	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(C8)	W	L	L	Q	T#
EU-A993	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(C12)	W	L	L	Q	T#
EU-A994	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(C14)	W	L	L	Q	T#
EU-A995	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(C16)	W	L	L	Q	T#
EU-A996	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(C18)	W	L	L	Q	T#
EU-A997	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(C20)	W	L	L	Q	T#
EU-A998	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(M8)	W	L	L	Q	T#
EU-A999	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(M12)	W	L	L	Q	T#
EU-A1000	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(M14)	W	L	L	Q	T#
EU-A1001	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(M16)	W	L	L	Q	T#
EU-A1002	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F I	Lys(M18)	W	L	L	Q	T#

Figure 9 (Continued)

Table 4 (Continued)

	SEQ ID. NO.	1				5				10						15						20					25					
EU-A100 3	949	H	Alb			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(M20)	W	L	L	Q	T#
EU-A100 4	950	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(M8)	W	L	L	Q	T#
EU-A100 5	951	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(M12)	W	L	L	Q	T#
EU-A100 6	952	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(M14)	W	L	L	Q	T#
EU-A100 7	953	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(M16)	W	L	L	Q	T#
EU-A100 8	954	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(M18)	W	L	L	Q	T#
EU-A100 9	955	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(M20)	W	L	L	Q	T#
EU-A101 0	956	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M8)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A101 1	957	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M12)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A101 2	958	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M14)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A101 3	959	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M16)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A101	960	H	G			Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M18)	A	A	K	E	F	I	A	W	L	L	N	T#

Figure 9 (Continued)

Table 4 (Continued)

	SEQ ID. NO.	1				5				10					15					20				25						
4																														
EU-A101 5	961	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M20)	A	A	K	E	F	I	A	W	L	L	N	T#
EU-A101 6	962	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M8)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A101 7	963	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M12)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A101 8	964	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M14)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A101 9	965	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M16)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A102 0	966	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M18)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A102 1	967	H	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(M20)	A	A	R	E	F	I	A	W	L	L	Q	T#
EU-A102 2	968	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(M8)	W	L	L	Q	T#
EU-A102 3	969	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(Met 2)	W	L	L	Q	T#
EU-A102 4	970	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(Met 4)	W	L	L	Q	T#
EU-A102 5	971	H	Alb	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(Met 6)	W	L	L	Q	T#

Figure 9 (Continued)

Table 4 (Continued)

	SEQ ID. NO.	1								5						10					15						20			25						T#
EU- A102 6	972	H	Alb	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(Met 8)	W	L	L	Q						T#
EU- A102 7	973	H	Alb	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(Met 0)	W	L	L	Q						T#
EU- A102 8	974	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(Met)	W	L	L	Q						T#
EU- A102 9	975	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(Met 2)	W	L	L	Q						T#
EU- A103 0	976	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(Met 4)	W	L	L	Q						T#
EU- A103 1	977	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(Met 6)	W	L	L	Q						T#
EU- A103 2	978	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(Met 8)	W	L	L	Q						T#
EU- A103 3	979	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Q	A	A	R	E	F	I	Lys(Met 0)	W	L	L	Q						T#
EU- A103 4	980	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(Met8)	A	A	K	E	F	I	A	W	L	L	N						T#
EU- A103 5	981	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(Met1 2)	A	A	K	E	F	I	A	W	L	L	N						T#
EU- A103 6	982	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(Met1 4)	A	A	K	E	F	I	A	W	L	L	N						T#
EU- A103	983	H	G	G	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E	Lys(Met1 6)	A	A	K	E	F	I	A	W	L	L	N						T#

Figure 9 (Continued)

Table 4 (Continued)

SEQ ID. NO.	1	5	10	15	20	25	T#
7							
EU-A103 8	H G	Q G T F	T S D Y	K Y L D E	A A K E F I A	W L L N	T#
EU-A103 9	H G	Q G T F	T S D Y	K Y L D E	A A K E F I A	W L L N	T#
EU-A104 0	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#
EU-A104 1	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#
EU-A104 2	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#
EU-A104 3	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#
EU-A104 4	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#
EU-A104 5	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#
EU-A104 6	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#
EU-A104 7	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#
EU-A104 8	H G	Q G T F	T S D Y	K Y L D E	A A R E F I A	W L L Q	T#

Figure 9 (Continued)

Table 4 (Continued)

SEQ ID. NO.	1			5					10			15			20			25	
EU-A1049	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C15)	#							
EU-A1050	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C18)	#							
EU-A1051	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C20)	#							
EU-A1052	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C8)	#							
EU-A1053	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C12)	#							
EU-A1054	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C14)	#							
EU-A1055	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C15)	#							
EU-A1056	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C18)	#							
EU-A1057	H	cMePr ₀	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C20)	#							
EU-A1058	H	Alb	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C8)	#							
EU-A1059	H	Alb	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C12)	#							
EU-A106	H	Alb	Q	G	T	cMeF	T	S	D	EIBi _p	Lys(C14)	#							

Figure 9 (Continued)
Table 4 (Continued)

	SEQ ID. NO.	1				S					10			15		20		25	
0																			
EU- A106 1	1007	H	Alb	Q	G	T	αMeF	T	S	D	EiBi P	Lys(C16)	#						
EU- A105 2	1008	H	Alb	Q	G	T	αMeF	T	S	D	EiBi P	Lys(C18)	#						
EU- A105 3	1009	H	Alb	Q	G	T	αMeF	T	S	D	EiBi P	Lys(C20)	#						
EU- A106 4	1010	H	Alb	Q	G	T	αMeF _F	T	S	D	EiBi P	Lys(C8)	#						
EU- A106 5	1011	H	Alb	Q	G	T	αMeF _F	T	S	D	EiBi P	Lys(C12)	#						
EU- A106 6	1012	H	Alb	Q	G	T	αMeF _F	T	S	D	EiBi P	Lys(C14)	#						
EU- A105 7	1013	H	Alb	Q	G	T	αMeF _F	T	S	D	EiBi P	Lys(C16)	#						
EU- A105 8	1014	H	Alb	Q	G	T	αMeF _F	T	S	D	EiBi P	Lys(C18)	#						
EU- A105 9	1015	H	Alb	Q	G	T	αMeF _F	T	S	D	EiBi P	Lys(C20)	#						
EU- A107 0	1016	H	αMePr ₀	Q	G	T	αMeF	T	S	D	EiBi P	MeLys(C ₈)	#						
EU- A107 1	1017	H	αMePr ₀	Q	G	T	αMeF	T	S	D	EiBi P	MeLys(C ₁₂)	#						

Figure 9 (Continued)

Table 4 (Continued)

[illegible]

Figure 9 (Continued)

[illegible]

Figure 9 (Continued)

Table 4 (Continued)

[illegible]

Figure 9 (Continued)

Table 4 (Continued)

[illegible]

Figure 9 (Continued)

Table 4 (Continued)

SEQ ID. NO.	1				5					10				15			20		25
2																			
EU- A115 3	H	αMePr 0	Q	G	T	αMeF F	T	S	D	Bip	Me20	#							
EU- A115 4	H	Alb	Q	G	T	αMeF	T	S	D	Bip	Me8	#							
EU- A115 5	H	Alb	Q	G	T	αMeF	T	S	D	Bip	Me12	#							
EU- A115 6	H	Alb	Q	G	T	αMeF	T	S	D	Bip	Me14	#							
EU- A115 7	H	Alb	Q	G	T	αMeF	T	S	D	Bip	Me16	#							
EU- A115 8	H	Alb	Q	G	T	αMeF	T	S	D	Bip	Me18	#							
EU- A115 9	H	Alb	Q	G	T	αMeF	T	S	D	Bip	Me20	#							
EU- A116 0	H	Alb	Q	G	T	αMeF F	T	S	D	Bip	Me8	#							
EU- A116 1	H	Alb	Q	G	T	αMeF F	T	S	D	Bip	Me12	#							
EU- A116 2	H	Alb	Q	G	T	αMeF F	T	S	D	Bip	Me14	#							
EU- A116 3	H	Alb	Q	G	T	αMeF F	T	S	D	Bip	Me16	#							

Figure 9 (Continued)
Table 4 (Continued)

C with a numeral means 1-alkyl beta-D-glucouronyl acyl group, while the numeral denotes the chain length.

MeLys = C-alpha-methyl-lysine, EtBip = 2'-ethyl-biphenylalanine; alphaMeF = C-alpha-methyl-2-fluoro-Phe

M is maltoside and Me is melibioside-based surfactant, while the numeral is the carbon chain length of the 1-alkyl substitution. Starred means alpha linked glycoside. For example Me* means alpha-melibioside.

Lys(C12) means N-epsilon-(1'-dodecyl beta-D-glucuronyl)-L-lysine and other C numbers mean the corresponding 1'-alkyl glucuronide. The pair of amino acids E*, K* or K*, E* separated by 4 residues denotes a side chain lactam linkage formed between the side chain functional groups on these amino acids.

means amide C-terminus.

Figure 10




Figure 11




Figure 12




Figure 13





Figure 14

H-His¹-Aib-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Tyr¹⁰-Ser-Lys-Tyr-Leu-Asp-



Glu^{*}-Gln-Ala-Ala-Lys^{20*}-Glu-Phe-Ile-Lys-Trp-Leu-Leu-Gln-Thr²⁹-NH₂

EU- A992	938	H	Aib	Q	G	T	F	T	S	D	Y	S	K	Y	L	D	E*	Q	A	A	K*	E	F	I	Lys(C8)	W	L	L	Q	T#
-------------	-----	---	-----	---	---	---	---	---	---	---	---	---	---	---	---	---	----	---	---	---	----	---	---	---	-------------	---	---	---	---	----

Figure 15

