




(86) Date de dépôt PCT/PCT Filing Date: 2002/05/17
 (87) Date publication PCT/PCT Publication Date: 2002/11/21
 (45) Date de délivrance/Issue Date: 2010/10/05
 (85) Entrée phase nationale/National Entry: 2003/12/16
 (86) N° demande PCT/PCT Application No.: CA 2002/000720
 (87) N° publication PCT/PCT Publication No.: 2002/091854
 (30) Priorité/Priority: 2001/05/17 (US09/858,602)

(51) Cl.Int./Int.Cl. *A23L 1/09* (2006.01),
A23G 3/00 (2006.01), *A23G 3/34* (2006.01),
A23L 1/00 (2006.01), *A23L 1/0532* (2006.01),
A23L 1/054 (2006.01), *A23L 1/48* (2006.01),
A23P 1/16 (2006.01), *C12N 9/26* (2006.01)
 (72) Inventeurs/Inventors:
 JAMPEN, STEPHAN, CA;
 SWAIN, ROBERT, CA
 (73) Propriétaire/Owner:
 CITADELLE COOPERATIVE DE PRODUCTEURS DE
 SIROP D'ERABLE, CA
 (74) Agent: BLAKE, CASSELS & GRAYDON LLP

(54) Titre : SIROP D'ERABLE TRAITE AUX ENZYMES, STABLE A L'ENTREPOSAGE, PRODUITS A BASE D'ERABLE
 PUR ET PRODUITS MELANGES
 (54) Title: ENZYME TREATED MAPLE SYRUP, SHELF STABLE, PURE MAPLE BASED PRODUCTS, AND BLENDED
 PRODUCTS

**Viscosity of a 55:45 Enzyme Treated to Non-Enzyme Treated
 Blend of Maple Syrup at Various Concentrations at 25°C**



(57) **Abrégé/Abstract:**

A method for producing shelf stable, maple syrup products is provided, which comprises adding a sucrose-cleaving enzyme, for example invertase, to maple syrup and incubating the resulting solution. A shelf stable, thick, pourable pure maple syrup is provided

(57) **Abrégé(suite)/Abstract(continued):**

that has the viscosity of common table syrup. Also provided is a shelf stable, spreadable maple syrup product with a consistency similar to that of clear honey, which may be used as a spread, or sweetener or a topping. Further, a shelf stable, spreadable pure maple butter product is provided, with the consistency of churned honey which may be used as a spread, sweetener, topping, or snack. The enzyme treated maple syrup may also be added to other products to increase the shelf stability of the products, by preventing the visible crystallization or separation of the products.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
21 November 2002 (21.11.2002)

PCT


(10) International Publication Number
WO 02/091854 A3

- (51) International Patent Classification⁷: A23L 1/09, C12N 9/26, A23L 1/48, A23G 3/00
- (21) International Application Number: PCT/CA02/00720
- (22) International Filing Date: 17 May 2002 (17.05.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/858,602 17 May 2001 (17.05.2001) US
- (63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 09/858,602 (CIP)
Filed on 17 May 2002 (17.05.2002)
- (71) Applicant (for all designated States except US): 2002872 ONTARIO LIMITED INC. [CA/CA]; 222 Bay Street, Suite 1800, Toronto, Ontario M5K 1H6 (CA).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): SWAIN, Robert [CA/CA]; 51 Lynngrove Avenue, Toronto, Ontario M8X 1M7 (CA). JAMPEN, Stephan [CA/CA]; 2 Lockyer Road, Guelph, Ontario N1G 1K1 (CA).
- (74) Agent: BLAKE, CASSELS & GRAYDON LLP; Box 25, Commerce Court West, Toronto, Ontario M5L 1A9 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,

[Continued on next page]

(54) Title: ENZYME TREATED MAPLE SYRUP, SHELF STABLE, PURE MAPLE BASED PRODUCTS, AND BLENDED PRODUCTS

Viscosity of a 55:45 Enzyme Treated to Non-Enzyme Treated Blend of Maple Syrup at Various Concentrations



(57) Abstract: A method for producing shelf stable, maple syrup products is provided, which comprises adding a sucrose-cleaving enzyme, for example invertase, to maple syrup and incubating the resulting solution. A shelf stable, thick, pourable pure maple syrup is provided that has the viscosity of common table syrup. Also provided is a shelf stable, spreadable maple syrup product with a consistency similar to that of clear honey, which may be used as a spread, or sweetener or a topping. Further, a shelf stable, spreadable pure maple butter product is provided, with the consistency of churned honey which may be used as a spread, sweetener, topping, or snack. The enzyme treated maple syrup may also be added to other products to increase the shelf stability of the products, by preventing the visible crystallization or separation of the products.



WO 02/091854 A3

WO 02/091854 A3



GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

(88) Date of publication of the international search report:
10 April 2003

Published:

— *with international search report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**ENZYME TREATED MAPLE SYRUP,
SHELF STABLE, PURE MAPLE BASED PRODUCTS, AND BLENDED
PRODUCTS**

5

Field of the Invention

The present invention relates to enzyme treated maple syrup. The enzyme treatment increases the shelf stability of the maple syrup, and other products to which the enzyme treated maple syrup is added, by preventing the visible crystallization or separation of the product. The invention also relates to shelf stable, pure, maple based products, for example (1) shelf stable, pourable, thick maple syrup with a similar viscosity to common table syrup, (2) shelf stable, spreadable maple syrup product with the consistency of clear honey, and (3) shelf stable, spreadable maple butter with the consistency of churned honey. This invention also relates to an enzyme treated maple syrup that when added to products prevents the visible crystallization or separation of the products.

20

Background of the Invention

The sap of maple trees forms the basis of maple-based products, including maple syrup, maple sugar and maple confection products. Maple syrup is obtained by the concentration of maple sap, a low solids (low sugar) solution obtained from the maple tree, to a solids content of 66 percent (66 degrees Brix). The production of maple syrup and related products is highly regulated in Canada and the United States, such that all additives are prohibited. As a result, maple producers cannot simply use additives such as stabilizers if they wish to label their product as a "pure maple-based product" or as "pure maple syrup".

30

To produce maple syrup, the sap from maple tree is concentrated to 66 degrees Brix (at 68 degrees Fahrenheit; 20 degrees Celsius) to be considered as "pure maple syrup" by U.S. and Canadian law. About 40 liters of maple sap are needed to make 1 liter of maple syrup. During the evaporation process, the sap is heated which induces chemical changes that give maple syrup its characteristic color and flavor. These chemical

changes include non-enzymatic browning and caramelization reactions (Edson, 1910; Hayward and Pederson, 1946).

Concentration can be achieved through simple boiling in an open kettle or using more
5 advanced evaporation techniques such as vacuum pan evaporators. Various other
methods are known for concentrating maple sap. For example, U.S. patent 5,389,209
to Paquette discloses a method of boiling the sap under normal pressure, then heating
the sap to below boiling temperature and using an air circulating column to further
evaporate the water. Reverse osmosis can be used to pre-concentrate the sap to about
10 20-25 degrees Brix. An alternate concentration step, which involves the addition of
heat, must be used to complete the concentration to 66 degrees Brix such that the
characteristic maple flavor is created (North American Maple Producers Manual,
Bulletin 856, chapter 7).

15 Concentrating to a higher level (over 67.5 degrees Brix) will result in crystallization of
the sucrose, the main sugar found in maple sap, within the container while in storage.
Conversely, a maple syrup of lower Brix (under 64.5 degrees Brix) would spoil
(ferment) while in storage. Crystallization occurs because the main sugar in maple
syrup is sucrose (90-100 percent), with the rest being glucose (0 to 10 percent) (North
20 American Maple Syrup Producers Manual, Bulletin 856, Appendix 2).

It is the crystallization behavior of maple syrup at a higher concentration that allows for
the production of other maple-based products such as maple sugar. These products are
obtained by concentrating maple syrup past 66 degrees Brix. At these elevated Brix
25 levels, a supersaturated sucrose solution is made. Therefore, if this solution is cooled,
crystallization will occur. Depending on the rate of cooling and/or whether agitation is
present during the cooling process, characteristic maple-based products are obtained.
Slow cooling without agitation results in crystals that are very large, often termed "rock
candy". More rapid cooling, but again without agitation, will result in smaller crystals
30 but the product has a very gritty mouthfeel. When a highly supersaturated maple syrup
solution (85 to 90 degrees Brix) is cooled very rapidly without agitation, a non-

crystalline glass-like solid is obtained (North American Maple Syrup Producers Manual, Bulletin 856, chapter 9).

5 Conversely, if a supersaturated (84 to 85 degrees Brix; or 12 to 13 degrees Celsius above the boiling point of water) maple syrup solution is cooled rapidly with high-speed agitation, very small crystals are obtained with the resultant product being paste-like in consistency and being spreadable. This product is known as maple butter or maple cream. Currently there is no pure maple-based product with this paste-like consistency that is shelf stable.

10

Crystallization of the sugars found in maple syrup is random and occurs spontaneously at higher Brix levels. Because the crystallization behavior of concentrated maple sap is difficult to control, only a select few products have been developed. A common complaint with pure maple syrup is that it is too thin, especially when warmed. These comments are made when pure maple syrup is compared to common table syrups such as *Log Cabin*TM, *Mrs. Butterworth's*TM and *Aunt Jemima*TM. Currently, there is no natural or pure maple syrup which has the viscosity similar to that of common table syrup and that is shelf stable.

20 Additionally, at a concentration of 66 degrees Brix, maple syrup may be too runny (thin) to be used in an application such as a honey like spread. Currently, there is no natural or pure maple-based product which has the consistency and/or appearance similar to that of clear honey and that is shelf stable.

25 Clear honey is a transparent high viscosity 3.9 Pascal seconds sugar syrup with a moisture content of 17 to 19 percent (81 to 83 degrees Brix solids) (Rheological Methods in Food Process Engineering, Steffe, J.F. 1996, pp 82, 26 and 367). This material remains in a clear state, without crystallization for extended periods of time. The main sugar components in honey are the monosaccharides glucose and fructose.
30 These sugars are present in maple syrup in small amounts.

As previously described, a more viscous solution similar to that of clear honey, having a unique maple flavor, is attainable by further concentrating maple syrup to a higher Brix level (81 to 82 degrees Brix for example). However, when this supersaturated solution is cooled, crystallization occurs rapidly such that the clear viscous solution is not preserved. This occurs because the main sugar in maple syrup is the disaccharide sucrose, which crystallizes much more readily than glucose and fructose.

Pure glucose and fructose blends are available commercially and are known as invert sugars. It is known to use invert sugar (glucose/fructose) when making artificial maple-based products. It is also known that invert sugar tends to retard crystallization in maple-based products. However, simply adding invert sugar can lead to loss of natural maple flavor. Additionally, the resultant maple-based product may no longer be labeled or considered as pure under the Canadian and U.S. legal standards.

Invert sugars have also been used in making imitation maple syrup or syrup substitutes. U.S. patent 3,878,306 to Garstick discloses an imitation maple syrup made from various sugars and artificial flavorings. U.S. patent 4,938,989 to Steeves and McKelvey provides a maple syrup substitute which contains maple syrup, maple flavor, fructose and glucose and white sugar. Again, these products could not be considered pure maple-based products.

It is known in the art that sucrose can be cleaved into its constituent sugars, glucose and fructose by use of an acid such as L-tartaric acid (cream of tartar). However, the use of organic acids leads to products that have very poor flavor profiles and unacceptable appearances. A further challenge is that the acid would have to be removed, after it has cleaved the sucrose. This step would also remove important flavor components.

The difficulty in making stable high viscosity maple syrups extends to other maple-based products. For example, maple butter (also called maple cream) separates into two layers if not stored at temperatures below 0 degrees Celsius or 32 degrees Fahrenheit. A dilute syrup layer forms on top and a solid crystalline mass forms underneath. Maple butter is made by heating maple syrup to 11 to 13 degrees Celsius

above the boiling point of water (83 to 85 degrees Brix), and cooling rapidly while stirring.

Many viscosity-altering hydrocolloid substances are commonly used in food products,
5 and are known in the art as stabilizing products or 'stabilizers'. Two commonly used stabilizers are carageenan and xanthan. Stabilizers are added to prevent phase separation in liquids or solids such as ice cream and peanut butter. However, these stabilizers have not been used to date in maple-based products. Carrageenan is
10 extracted from seaweed while xanthan is derived from an industrial fermentation of a bacteria.

Accordingly, there is a need for pure maple-based products that are shelf stable, and that can extend the scope of the use of maple syrup and related products. There is a need for enzyme treated maple syrup. The enzyme treatment increases the shelf
15 stability of the maple syrup, and products to which the enzyme treated maple syrup is added. There is also a need for shelf stable, pure maple based products, for example, (1) there is a need for a shelf stable, pourable, thick maple syrup with the viscosity of common table syrup, (2) there is a need for a shelf stable, spreadable, non-crystalline maple syrup product with the consistency of clear honey and, (3) there is also a need
20 for a shelf stable, spreadable maple butter with the consistency of churned honey.

Further, there is a need for an enzyme treated maple syrup that when added to products prevents the visible crystallization or separation of the products.

25 **Summary of the Invention**

As uses herein, the term maple syrup refers to concentrated or unconcentrated sap of the botanical genus *Acer*. It is in the scope of this invention that the unconcentrated maple sap may be used directly.

30

The main aspect of the present invention is the enzyme treatment of maple syrup. The enzyme treatment cleaves the main sugar in maple syrup (sucrose) to yield glucose and

fructose. Products containing a high concentration of sucrose are not shelf stable and will crystallize, whereas products containing a similarly high concentration of the combination of glucose and fructose, and proportionally less sucrose, will not crystallize or separate as readily.

5

The present invention provides methods for the enzymatic treatment of maple syrup, and products produced thereof. For example, the invention provides: (1) an enzyme treated maple syrup product which is a pure maple syrup product, which when added to other products prevents visible crystallization and separation of the products, (2) a shelf
10 stable, thick, pourable maple syrup with the viscosity of common table syrup, (3) a shelf stable, spreadable, non-crystalline maple syrup product with a consistency similar to that of clear honey, wherein the product is preferably transparent or translucent, (4) a shelf stable maple-based product with high Brix value, for example, shelf stable maple butter with a similar consistency to churned honey.

15

The following aspect relates to the general method of enzyme treating maple syrup.

According to one object of the present invention, there is provide a method of producing an enzyme treated maple syrup, comprising: (a) adding a sucrose-cleaving
20 enzyme to a maple syrup; and (b) incubating the maple syrup to produce the enzyme treated maple syrup. Optionally, the method further comprises the step of concentrating the enzyme treated maple syrup. Optionally, the method further comprises the step of removing or inactivating the sucrose-cleaving enzyme.

Optionally, the sucrose-cleaving enzyme is invertase. Further, the maple syrup may be
25 diluted to about 55 to 66 degrees Brix prior to the step of adding a sucrose-cleaving enzyme. The invention also provides the enzyme treated maple syrup produced by this method.

The following aspects relate to the shelf stable, thick, pourable, pure maple syrup with
30 the viscosity of common table syrup.

According to another object of the present invention, there is provided a method for producing a shelf stable, thick, pourable, pure maple syrup product with a viscosity of common table syrup, comprising: a) adding a sucrose-cleaving enzyme to a maple syrup; b) incubating the maple syrup to produce an enzyme treated maple syrup; c) 5 adding an untreated maple syrup to the enzyme treated maple syrup at a predetermined ratio of enzyme treated maple syrup to untreated maple syrup, to produce a maple syrup blend of untreated maple syrup and enzyme treated maple syrup; and d) concentrating the maple syrup blend to produce the shelf stable, thick and pure maple syrup product with the viscosity of common table syrup. Optionally, the ratio of enzyme treated 10 maple syrup to untreated maple syrup is from about 30:70 to 60:40 by percentage of sugars present, and preferably about 40:60. Optionally the maple syrup product is concentrated to between about 72 and 76 degrees Brix. The invention also provides the shelf stable, thick pourable, pure maple syrup product made by this method.

15 According to another object of the present invention, there is provided a method for producing a shelf stable, thick, pourable, pure maple syrup product with a viscosity of common table syrup, comprising: a) adding a sucrose-cleaving enzyme to a maple syrup; b) incubating the maple syrup to produce an enzyme treated maple syrup; c) concentrating the enzyme treated maple syrup; and d) adding an untreated maple syrup 20 to the concentrated enzyme treated maple syrup at a predetermined ratio of enzyme treated maple syrup to untreated maple syrup, to produce a maple syrup blend of untreated maple syrup and enzyme treated maple syrup, wherein the maple syrup blend is the shelf stable, thick, pourable and pure maple syrup product with the viscosity of common table syrup. Optionally, the ratio of enzyme treated maple syrup to untreated 25 maple syrup is from about 30:70 to 60:40 by percentage of sugars present, and preferably about 40:60. Optionally, the enzyme-treated maple syrup is between about 84 and 89 degrees Brix, and the maple syrup product has a final concentration of between about 72 and 76 degrees Brix. The invention also provides the thick, shelf stable maple syrup made by this method.

30

According to another object of the present invention, there is provided a method for producing a shelf stable, thick, pourable, pure maple syrup product with a viscosity of

common table syrup, comprising: a) adding a sucrose-cleaving enzyme to a maple syrup; b) incubating the maple syrup to produce an enzyme treated maple syrup; c) inactivating or removing the sucrose-cleaving enzyme; and d) concentrating the enzyme treated maple syrup to produce the shelf stable, thick pourable and pure maple syrup product with the viscosity of common table syrup. Optionally, the maple syrup product is concentrated to between about 72 and 76 degrees Brix. The invention also provides the thick maple syrup made by this method.

Optionally, the method to produce shelf stable, thick, pourable, pure maple syrup with the viscosity of common table syrup utilizes invertase. Optionally, the method comprises the additional step of diluting the maple syrup to about 55 to 66 degrees Brix prior to the step of adding the sucrose-cleaving enzyme. Optionally, the sucrose-cleaving enzyme is removed by filtration or inactivated by heat treatment. However, due to requirements of various regulatory agencies, it may be necessary to remove the enzyme in order to meet applicable regulatory standards for pure maple syrup. Optionally, the method includes an additional step monitoring the ratio of sucrose to fructose and glucose, and removing or inactivating the sucrose-cleaving enzyme at a predetermined sucrose to fructose and glucose ratio, followed by the step of concentrating. Optionally, the viscosity of the maple syrup product is between about 1200 and 2100 centipoise at 2.5 rpm as determined by a Brockfield Viscometer at 22⁰C, and preferably between about 1300 and 1900 centipoise.

According to another aspect of the invention, there is provided a shelf stable, thick, pourable and pure maple syrup product having a Brix measurement of between about 70 and about 76 degrees and a viscosity of between about 1200 and 2100 centipoise (and preferably between 1300 and 1900) at 2.5 rpm as determined by a Brockfield Viscometer at 22⁰C.

According to another aspect of the present invention, there is provided the use of the shelf stable, thick, pourable, pure maple syrup as a topping, sweetener or ingredient in a food.

The following aspects relate to the shelf stable maple-based product with the consistency of clear honey.

According to another aspect of the present invention there is provided a method for producing a shelf stable, spreadable maple syrup product with a consistency of clear honey, comprising: a) adding a sucrose-cleaving enzyme to a maple syrup; b) incubating the maple syrup to produce an enzyme treated maple syrup; c) adding an untreated maple syrup to the enzyme treated maple syrup at a predetermined ratio of enzyme treated maple syrup to untreated maple syrup, to produce a maple syrup blend of untreated maple syrup and enzyme treated maple syrup; and d) concentrating the maple syrup blend to produce the shelf stable, spreadable maple syrup product with the consistency of clear honey. Optionally, the ratio of enzyme treated maple syrup to untreated maple syrup is from about 45:55 to 60:40 by percentage of sugars present, preferably about 55:45 by percentage of sugars present. Optionally, the maple syrup product has a final concentration of between about 79 and 83 degrees Brix. There is also provided a shelf stable, spreadable maple syrup product with the consistency of clear honey made by this method.

According to another aspect, there is provided a method for producing a shelf stable, spreadable maple syrup product with the consistency of clear honey, comprising: a) adding a sucrose-cleaving enzyme to a maple syrup; b) incubating the maple syrup to produce an enzyme treated maple syrup; c) inactivating or removing the sucrose-cleaving enzyme; and d) concentrating the enzyme treated maple syrup to produce the shelf stable, spreadable maple syrup product with the consistency of clear honey. Optionally, the maple syrup product has a final concentration of between about 79 and 83 degrees Brix. There is also provided a shelf stable, spreadable maple syrup product with the consistency of clear honey made by this method.

Optionally, the sucrose-cleaving enzyme is invertase. Optionally, the method comprises the additional step of diluting the maple syrup to about 55 to 66 degrees Brix prior to the step of adding the sucrose-cleaving enzyme. Further, the sucrose-cleaving enzyme can be removed by filtration or inactivated by heat treatment. Due to

requirements of various regulatory agencies, it may be necessary to remove the enzyme in order to meet applicable regulatory standards for pure maple-based products. In addition, the method may comprise the additional step of monitoring the ratio of sucrose to fructose and glucose, and removing or inactivating the sucrose-cleaving enzyme at a predetermined sucrose to fructose and glucose ratio, followed by the step of concentrating.

There is also provided a shelf stable, spreadable maple syrup product with the consistency of clear honey having a Brix measurement of between about 72 and 90 degrees, preferably between 79 and 83 degrees.

According to another aspect of the present invention, there is provided the use of the shelf stable, spreadable maple syrup product with the consistency of clear honey as a topping, sweetener or ingredient in a food.

The following aspects relate to shelf stable maple butter.

Another object of the invention is to provide a method of producing shelf stable maple butter which comprises, (1) combining maple syrup with an enzyme treated maple syrup, or one or more stabilizers, or a combination of both to form a maple syrup solution, and (2) concentrating the maple syrup solution. The stabilizer can be any or all of one or more viscosity altering hydrocolloids. Optionally, air or an inert gas, for example nitrogen, can be added after concentrating the solution. Optionally, the viscosity altering hydrocolloid is xanthan or carageenan or both, at a concentration of about 0.1 to 1 percent. Optionally the ratio of pure maple-based product stabilizer to maple syrup is 20:80 to 25:75 by percentage of sugars present. Optionally the resulting maple butter is concentrated to between 83 degrees Brix and 87 degrees Brix.

Another object of the present invention is to provide a method of producing a shelf stable maple butter, comprising: (a) adding a sucrose-cleaving enzyme to a maple syrup; (b) incubating the maple syrup to produce an enzyme treated maple syrup; (c) inactivating or removing the sucrose cleaving enzyme; and (d) concentrating the

enzyme treated maple syrup to produce the shelf stable maple butter. Optionally, the sucrose-cleaving enzyme is invertase. Optionally, the maple syrup is diluted to about 55 to 66 degrees Brix prior to the step of adding the sucrose cleaving enzyme. The sucrose-cleaving enzyme may be removed by filtration or inactivated by heat treatment.

5 Due to requirements of various regulatory agencies, it may be necessary to remove the enzyme in order to meet applicable regulatory standards for pure maple-based products. Further, the method may comprise the additional step of monitoring the ratio of sucrose to fructose and glucose, and removing or inactivating the sucrose cleaving enzyme at a predetermined sucrose to fructose and glucose ratio, followed by the step of
10 concentrating. Also provided is the maple butter produced by this method.

Another object of the invention is to provide a shelf stable maple butter with the consistency of churned honey. Another object is to provide a stabilized maple butter, which is shelf stable, can be a pure maple-based product.

15

The following aspects relate to improving the flavor.

Another object of the invention is to provide a method of making a maple-based product with improved flavor, for example maple butter, comprising (1) concentrating
20 maple syrup, and (2) adding air or an inert gas, for example nitrogen, after concentrating. Optionally, air can be added during the cooling step. Optionally, the air or the inert gas is added to a final concentration of 1% to 15%, most preferably to a final concentration of 12 percent.

25 Another object of the invention is to provide the improved flavor maple-based product.

The following aspects relate to enzyme treated maple syrup that when added to products, prevents the visible crystallization or separation of the product.

30 Another object of the present invention is to provide a method of producing an enzyme treated maple syrup, that when added to a maple based product functions to prevent the visible crystallization or separation of the maple based product, comprising (1) adding

a sucrose-cleaving enzyme, for example, invertase, to maple syrup and (2) incubating the maple syrup to produce the enzyme treated maple syrup. Optionally, the method could include the further steps of concentrating the enzyme treated maple syrup and removing the sucrose-cleaving enzyme post-incubation. The method could also include
5 the optional step of diluting the maple syrup to between about 55-66 degrees Brix prior to the addition of the sucrose-cleaving enzyme.

Another object of the present invention is to provide an enzyme treated maple syrup that is a pure maple-based product, and that when added to a maple based product
10 functions to prevent the visible crystallization and separation of the maple based product.

Another object of the present invention is to provide a blended product comprising an enzyme treated maple syrup and a second product. Also the invention provides an
15 enzyme treated maple syrup that when added to a maple based product prevents the crystallization and separation of the maple based product. Also provided is a blended product comprising a maple-based product and an enzyme treated maple syrup.

According to a further object of the invention there is provided a method of using a
20 stabilized maple-based product as a spread, sweetener, or a topping, or on its own as a snack food product.

Another object of the invention is to provide a method for stabilizing a maple-based product comprising, (1) combining maple syrup with an enzyme treated maple syrup, or
25 one or more stabilizers, or a combination of both to form a maple syrup solution, and (2) optionally, concentrating the maple syrup solution. The stabilizer can be any or all of one or more viscosity altering hydrocolloids. Optionally, air or an inert gas, for example nitrogen, can be added after concentrating the solution. In another object, a method is provided to stabilize a product by combining the product with an enzyme
30 treated maple syrup.

Another object of the invention is to provide a shelf stable maple-based product and a stabilized maple-based product.

According to a further object of the invention there is provided a method of using a
5 stabilized maple-based product as an ingredient in the manufacture of a pure maple-based product.

Detailed Description of the Drawings

10 Figure 1 is a graph showing the viscosity profile of a 55:45 enzyme treated to non-enzyme treated blend of maple syrup at various concentrations at 25°C.

Figure 2 is a graph showing the viscosity profile of a 55:45 enzyme treated to non-treated blend of maple syrup at various concentrations at 40°C.

15

Detailed Description of the Invention

The present invention provides an enzyme treated maple syrup that increases the shelf stability of the maple syrup. The enzyme treatment increases the shelf stability of other
20 maple-based products to which the enzyme treated maple syrup is added, by preventing the visible crystallization or separation of the product. The enzyme treatment cleaves the main sugar in maple syrup (sucrose) to yield glucose and fructose. Products containing a high concentration of sucrose are not shelf stable and will crystallize or separate, whereas products containing a similarly high concentration of the
25 combination of glucose and fructose, with proportionally lower sucrose, will not crystallize or separate as readily. Therefore, the enzyme treatment provides shelf stable, pure maple based products.

The present invention provides a shelf stable, thick, pourable, pure maple syrup which
30 has the flow properties similar to that of common table syrup. The pure maple syrup of the present invention is thicker than the pure maple syrup of the current state-of-the-art.

Therefore, the thick, pure maple syrup of the present invention is less runny, especially when warm.

5 The present invention also provides a spreadable maple syrup product which is shelf stable, having a consistency similar to that of clear honey. The maple syrup product may be used in a similar manner as honey: as a spread on bread products, topping for other food products such as ice cream or other desserts, or as a sweetener in home cooking and/or in commercial food products. It is also suitable for use in pure maple-based products.

10

The present invention also provides a shelf stable maple butter which is spreadable and has a similar consistency to churned honey. The maple butter may be used: as a spread on bread products, topping for other food products such as waffles, pancakes, ice cream or other desserts, or as a sweetener in home cooking and/or in commercial food
15 products, for example, filling for chocolates and other confections.

20

The present invention also provides an enzyme treated maple syrup that when added to a product increases the shelf stability of the product by preventing the visible crystallization or separation of the product. The product can be a maple syrup product or other product, for example, fondants, fudge and other crystalline candy. This
20 enzyme treated maple syrup has the advantage of being a pure maple-based product.

25

There are several advantages to the present invention: (1) the invention provides for maple-based products with a high Brix value that are shelf stable, (2) the products are
25 pure maple-based products, and are labeled as "pure"; (3) the products taste like maple based products, (4) the method to produce the products is easy and relatively inexpensive and (5) the enzyme treated maple syrup can be added to other products, to increase the shelf stability of other products.

30

Definitions

"About" when used herein in relation to Brix measurement means ± 2 units of measure, when used in relation to % sugar concentration, % air, % viscosity or % stabilizer
5 means $\pm 20\%$, when used in relation to % enzyme treated maple syrup, % maple syrup, % untreated maple syrup, ratio of enzyme treated maple syrup to untreated maple syrup and % stabilizer means $\pm 10\%$.

"Brix" when used herein means the refractometric sucrose value determined in
10 accordance with the 'International Scale of Refractive Indices of Sucrose Solutions' and to which the applicable corrections for temperature and other solids have been made. The Brix value would be determined in accordance with the refractometric method outlined in the 'Official Methods of Analysis of the Association of Analytical
15 Chemists'. In the case of maple syrup, the Brix value, as measured by refractometer or hydrometer, essentially equals the amount of sugar present because most (about 98 %) of the solids are sugars.

"Crystallization" when used to describe the shelf stable, spreadable maple syrup
20 product with the consistency of clear honey and the shelf stable, thick pourable maple syrup with the viscosity of common table syrup means crystals that are visible.

"Enzyme treated maple syrup" when used herein means a maple syrup that has been treated with a sucrose cleaving-enzyme.

25 "Maple-based product" when used herein means a product that contains maple syrup or maple sap.

"Maple butter" when used herein means the creamy smooth textured product obtained
30 by the rapid cooling of a supersaturated maple syrup solution under high amounts of agitation. Maple butter is also known as maple cream. It is a crystallized product wherein the crystals are not visible without the use of magnification.

"Maple syrup" when used herein means the concentrated or unconcentrated sap of trees of the botanical genus *Acer*.

5 "Pourable" when used herein means having a viscosity such that the product flows from a vessel, similar to the viscosity of common table syrup.

"Pure maple-based product" when used herein means a maple-based product that contains only maple syrup, enzyme treated maple syrup or a combination of enzyme treated maple syrup and untreated maple syrup.

10

"Separation" when used to describe the maple butter means the formation of two or more distinct layers within the product.

15 "Shelf stable" when used herein in reference to the thick, pourable maple syrup with the viscosity of common table syrup and maple syrup product with the consistency of clear honey means that the maple syrup product is substantially free from crystallization for a period of at least 3 months. "Shelf stable" when used herein in reference to a maple butter product means that the maple butter is substantially free from separation for a period of at least 1 month.

20

"Spreadable" when used herein means distributable over a surface, in a manner similar to clear honey or churned honey.

25 "Stabilizer" when used herein means a substance that when added to a food product, inhibits crystallization, or prevents phase separation, or both.

A "sucrose-cleaving enzyme" when used herein is a hydrolysis enzyme which preferentially cleaves the beta-D-fructofuranoside linkage between the glucose and fructose molecules that make up sucrose.

30

"Untreated maple syrup" when used herein is maple syrup that has not been treated with a sucrose cleaving enzyme.

"Visible" when used herein means visible to the naked eye, without the use of magnification.

5 To make the high viscosity maple syrup of the present invention, any grade of maple syrup may be used as the starting material. Preferably, the maple syrup is high grade. The maple syrup is placed in a sanitary vessel such as a sterile agitated incubation tank and may optionally be diluted in order to optimize reaction conditions. Preferably, the maple syrup is diluted using sterile deionized water to a final Brix content of between
10 about 55-66 degrees Brix. Alternatively, maple sap which has been concentrated to about a final Brix content of between about 55 and 66 degrees Brix may be used in place of maple syrup.

An enzyme which cleaves sucrose into glucose and fructose is added to the maple
15 syrup. Preferably an invertase enzyme is used. There are two main groups of invertase enzymes: 1) alpha-glucosidases which are also known as maltase, glucoinvertase, glucosidosucrase, maltase-glucoamylase, lysosomal alpha glucosidase, and acid maltase and 2) beta fructofuranosidases which are also known as invertase, saccharase and beta-fructosidase. Invertase is available commercially from Sigma
20 Chemicals (Grade V: Practical from baker's yeast) and is used in the baking industry to control the amount of surface browning in bread and cookies. Invertase is added in the amount of 0.05 % of the final weight of the diluted mixture. Invertase is sucrose specific and self-terminating in that once all of the substrate sucrose is cleaved, nothing else will be hydrolyzed.

25 The solution may be adjusted to the optimal pH for invertase (pH 4.6) through known pH adjustment methods. Preferably, to maximize the natural maple flavor, the pH is not adjusted. The reaction still proceeds at the natural pH of maple syrup (pH 6.8), but the time necessary to complete the hydrolysis of sucrose to glucose and fructose
30 increases when the pH is not adjusted. At pH 6.8, conversion takes approximately 7 days.

The incubation temperature is preferably between 15 and 35 degrees Celsius and most preferably, the incubation should take place at room temperature (20-23 degrees Celsius) under continuous gentle agitation. Because of the low Brix, sanitary practices should be used when sampling and in further processing due to the potential for
5 microbial growth.

The cleavage of sucrose to glucose and fructose can be monitored using various known methods. The preferred method is monitoring the optical rotation of the sugars. As the hydrolysis progresses, the muta-rotation decreases, and is negative when complete. A
10 second method encompasses monitoring the hydrolysis using HPLC techniques methods (AOAC Official Method 977.20, Separation of Sugars in Honey). Glucose, fructose and sucrose can be separated by a carbohydrate column (Waters Inc., carbohydrate analysis column, part no. 84038) using a mobile phase of 83:17 acetonitrile: water. With corresponding standards, the three sugars can be identified
15 and quantified. Hydrolysis is deemed complete after negligible amounts of sucrose are detectable.

Once the reaction is complete, preferably, the invertase enzyme can be removed or inactivated from the mixture. The advantages of removing the enzyme from the syrup
20 are to remove the visible haze and to remove the protein source which could initiate the Maillard browning reaction upon heating in the evaporation step. The Maillard reaction potentially creates bitter flavors, which would be objectionable in this product.

The enzyme can be removed by known means, including precipitation, hydrolysis and
25 filtration or can be inactivated by known means such as heating (> 90 degrees Celsius). Preferably, the invertase is removed by filtration through a filter of pore size of less than 1µm.

It is also within the scope of this invention that the enzyme be immobilized onto a resin
30 bead that is then placed into a reaction column. The maple syrup is flushed through the column on a continuous basis, allowing the reaction to occur as the maple syrup passes over the resin beads onto which the enzyme had been attached. In such a case, the

enzyme does not have to be removed through alternate means such as filtration as it remains bound to the resin beads in the column.

Once the enzyme is removed or inactivated, the syrup can be concentrated to any
5 concentration, depending on the viscosity or consistency that is desired in the product. It is preferably concentrated to between 72 and 90 degrees Brix, depending on the final product. For example, it is concentrated to (1) between 72 and 76 degrees Brix for the thick, pourable maple syrup product that has a viscosity of common table syrup, (2)
10 between 79 and 83 degrees Brix for the spreadable maple syrup product which has the consistency of clear honey, and (3) between about 83 and 86 degrees Brix for the spreadable maple butter product with a consistency of churned butter. It is within the scope of the invention that other products can be made by this method. The maple butter product can also be blended with untreated maple syrup for the spreadable maple butter with a consistency of churned honey. The concentration may be accomplished
15 using known methods such as heating in an open kettle and vacuum pan evaporators. Concentration of the syrup is most preferably accomplished by heating the syrup to high temperature for short periods of time and flashing off the appropriate quantities of water to reach the desired Brix level. Low temperatures for longer periods of time with vacuum can also be used. Low temperature long time concentration is best carried out
20 between 50 and 80 degrees Celsius, more preferably from 50 to 65 degrees Celsius and most preferably at 65 degrees Celsius and under vacuum (about 0.8 bar). The high viscosity maple syrup product produced in this manner may then be used by a consumer after the usual packaging steps, or it may be packaged and provided to the consumer. The maple syrup products can be used by pouring the product over foods, as
25 a sweetener or topping by adding the product to foods and as an ingredient in pure maple based products. The enzyme-treated maple syrup product can also be used to increase the shelf life of products, by preventing the visible crystallization or separation of the product when added to it.

30 To further improve the flavor profile, the dilute enzyme treated syrup is preferably blended with maple syrup of any grade prior to concentration. Optimum blends are 30:70, 40:60, 50:50, 55:45 and 60:40 (enzyme treated to non-enzyme treated syrup).

The blended syrup can then be concentrated as described above. Alternatively, the enzyme treated maple syrup can be concentrated prior to blending with untreated maple syrup. The maple flavor profile can be tailored to suit the retail climate by using different grades of maple syrup with varying flavor profiles in the incubation step and
5 the blending step. The color of the final product can be controlled in a similar manner.

To make the shelf stable maple-based product of the present invention, for example maple butter, any grade of maple syrup may be used as the starting material. Preferably, the maple syrup is high grade. The maple syrup is combined with either (1)
10 a small quantity of viscosity altering hydrocolloid stabilizer, preferably 0.1% carageenan or 0.1% xanthan, which has been first dispersed in a small quantity of water, or (2) an enzyme treated maple syrup, preferably in a blend ratio of 20 to 25% enzyme treated maple syrup to 75 to 80% untreated maple syrup, or (3) one or more viscosity altering hydrocolloid stabilizers and enzyme treated maple syrup. The
15 enzyme treated maple syrup is produced by adding enzyme to maple syrup and incubating the maple syrup.

The resultant blend is then concentrated, preferably to a concentration of between 83 and 87 degrees Brix, and most preferably to a concentration of 86 degrees Brix. This
20 concentration may be accomplished using known methods such as heating in an open kettle and vacuum pan evaporation. Concentration of the resultant blend is most preferably accomplished by heating the syrup to high temperature for short periods of time in a plate heat exchanger, followed by flashing off the appropriate amounts of water to obtain the desired Brix content. Low temperatures for longer periods of time
25 with vacuum can also be used. The hot concentrated product is then transferred to a scraped surface heat exchanger where it is cooled rapidly under high amounts of agitation. The maple butter produced in this manner may then be provided to the consumer. The maple butter flavor profile can be tailored to suit the retail climate by using different grades of maple syrup with varying flavor profiles in the blending step.
30 The color of the final product can be controlled in a similar manner.

To further improve the flavor profile of the maple-based product, for example, maple butter, air or an inert gas (for example nitrogen) can be whipped, for example, into the 84 degree Brix concentrated blend of maple syrup, enzyme treated maple syrup and hydrocolloid stabilizer, while cooling under high or rapid agitation. Preferably, a blend
5 of pure maple syrup and enzyme treated maple syrup with a sugar ratio of 75 to 25 (pure maple syrup:enzyme treated maple syrup) is concentrated to 84 degrees Brix. Optionally, 0.15 xanthan can be added to the blend prior to concentration. After concentration, and during cooling, air can be whipped into the product to a final concentration of 1% to 15%, preferably between 9% to 12% air, most preferably to a
10 final concentration of 12% air.

The present invention is described in more detail by reference to the following specific examples, which are not to be construed as limiting.

15 Example 1: Addition of Invert Sugar to Maple Syrup

The first stabilization technique evaluated was simple addition of invert sugar (glucose and fructose) to maple syrup and then concentrating to 82 degrees Brix on a hot plate. Four different levels of invert sugar were added: 1, 6, 10, and 20 percent.
20 Crystallization was evident in all samples within three weeks of storage at room temperature. Interestingly, the more invert sugar added, the longer the time before crystallization was evident. However, with increasing invert sugar content, the maple flavor decreased substantially. The type of invert sugars was also evaluated. Three different invert sugars, solid invert, medium invert and high fructose corn syrup, were
25 added at 20 percent of the total volume to 66 degrees Brix maple syrup and concentrated to 80 to 82 degrees Brix. Crystallization appeared in each of the samples within about three weeks.

Example 2: Addition of Cream of Tartar to Maple Syrup

30

Cream of tartar (L-Tartaric Acid, Sigma Chemicals, Oakville, ON, Canada.) is an accepted way of converting sucrose to glucose and fructose (North American Maple

Producers Manual s.7). Three levels of tartaric acid were evaluated: 0.02, 0.08 and 0.10 percent. Tartaric acid was added to the 66 degrees Brix maple syrup. The solution was heated to boiling, as it was thought that the heat added in concentrating the syrup to 80-82 degrees Brix would be sufficient to cleave a portion of the sucrose to
5 glucose and fructose. However, the resultant samples were very bitter in taste and had a distinct hazy appearance. All samples also crystallized after a two-day period at room temperature.

Example 3: Enzymatic Cleavage of Sucrose to Glucose and Fructose with pH

10 Adjustment

The enzyme invertase (Sigma Chemicals, Oakville ON, Canada) was used to cleave sucrose to glucose and fructose. The optimal pH for this specific invertase enzyme is 4.5. However, the natural pH of maple syrup is 6.8. In the first trial, 1 kg of maple
15 syrup was placed in a sterile container. The pH of the maple syrup was adjusted to 5.0 using tartaric acid, closer to the optimal pH of the enzyme. To this, enzyme was added (1 g enzyme per 500 g syrup). In order to assist in the dispersion of the enzyme, the maple syrup was diluted to 55 degrees Brix using sterile deionized water. The solution was divided into two portions with one portion incubated at room temperature (22 to 23
20 degrees Celsius) and the other portion incubated at 33 degrees Celsius. After a 6 day incubation period the sucrose, glucose and fructose content was determined using HPLC methods (AOAC Official Method 977.20, Separation of Sugars in Honey). At both temperatures, all of the sucrose had been converted to glucose and fructose. A filtration step was then carried out to remove the enzyme (Whatman No. 42, Slow, Fine
25 Crystalline Material; Fisher Canada, Napean, ON, Canada). The syrup was placed in a beaker atop a hot plate and heated to boiling. Concentration of this syrup to 82 Brix resulted in a product that was substantially free of crystals, and was shelf stable at room temperature for at least 2 months.

30

Example 4: Enzymatic Cleavage of Sucrose to Glucose and Fructose without pH Adjustment

5 In an effort to reduce the effect of pH adjustment on the flavor of the product, the activity of the enzyme invertase was evaluated in maple syrup at the normal pH of maple syrup (pH 6.8) as described in Example 3. The incubation was carried out at room temperature, and allowed to proceed for 11 days. HPLC analysis of the sample showed that all of the sucrose had been converted to glucose and fructose. After
10 filtration and concentrating to 82 degrees Brix, the resultant syrup had improved flavor and clarity. However, a more intense maple flavor may be desired. The flavor can be improved by using a more continuous evaporation/concentration system such as a plate heat exchanger with a flash system.

15 Example 5: Flavor Improvements by Blending

To increase the maple flavor profile of the maple syrup at higher Brix, treated syrup (as described in Example 4) was blended with untreated maple syrup. Four blend ratios have been evaluated: 45:55, 50:50, 55:45, 60:40 (ratio of enzyme treated maple syrup
20 to untreated maple syrup). The concentration to 82 degrees Brix was carried out under vacuum (0.8 bar) and at a lower temperature (60 degrees Celsius). The resultant flavor profile was much improved over product produced in examples 3 and 4. Also, a shelf-life of more than 8 months is attainable with the 55:45 enzyme treated to pure maple syrup blend. Other blend ratios may be possible, for example 20 percent, but
25 this would result in a shelf life of only about three weeks. For extended shelf life, a pure (100 percent) treated syrup could be concentrated as in Example 4, but the flavor and texture would be less desirable.

Example 6: Shelf Life of Maple-based product at 75, 80, 83 and 87 Degrees Brix

30 Using a blend of 55:45 enzyme treated to non-enzyme treated syrup, further shelf life evaluations were carried out, evaluating the influence of maple syrup concentration. A

blend of 55:45 enzyme treated to non enzyme treated syrup was concentrated using an experimental continuous evaporation system. The system was set-up to concentrate the product to 87 degrees Brix. A portion of this product was then diluted using deionized water to 75, 80 and 83 degrees Brix. Resultant samples were stored in the dark at room temperature (20-25 degrees Celsius). Shelf life studies of the four samples (75, 80, 83, 87 degrees Brix) show no visual signs of crystallization.

Example 7: Viscosity

The viscosity of the maple syrup was determined at various concentrations. A 55:45 enzyme treated to non enzyme treated syrup blend was concentrated to 87 degrees Brix using an experimental continuous evaporation system. A sample of this product was then diluted to 75, 80 and 83 degrees Brix. The viscosity of the maple syrup was measured at each of the seven concentrations (66.5, 70, 73.5, 75, 80, 83, 87 degrees Brix) on a Carri-Med CLS² 500 Rheometer (TA instruments, New Castle, DE). A shear rate sweep (0 to 643 reciprocal seconds) was performed on each of the samples using a 2 cm, 4 degree cone and plate geometry. Temperature of the sample was controlled to 25 and 40 degrees Celsius. Because of the high viscosity of the samples, all samples were pre-warmed to 80 degrees Celsius prior to testing. Samples were then placed onto the testing apparatus and allowed to cool to the test temperature. The average viscosity of three runs is presented in Table 1 and shown in Figure 1. The average viscosity at 40°C is presented in Table 2 and shown in Figure 2.

Table 1 Viscosity of Maple Syrup at Various Concentrations determined at 40 Reciprocal Seconds at 25°C.

Concentration	Viscosity, Pa.s
75	1.48
80	11.8
83	61.5
87	487

WO 02/091854

PCT/CA02/00720

Table 2 Viscosity of a 40°C Maple Syrup Blend (55:45 ETS:Pure Maple Syrup) as a Function of Concentration

Concentration	Viscosity (Pa · s)
66.5	0.05868
70	0.1145
73.5	0.262
75	0.3799
80	2.266
87	9.754

Example 8: Blend methods of manufacture of the shelf stable, thick, pourable maple syrup product with a viscosity of common table syrup.

The method is based on the creation of the proper sugar ratio within the final product.

A known amount of maple syrup at 66 degrees Brix was blended with previously concentrated enzyme treated maple syrup such that both the final concentration and the proportions of each of the sugars present was corrected to yield a stable product. To obtain a 40:60 sugar ratio, the enzyme treated maple syrup was concentrated to between about 84 to 89 degrees Brix, preferably to about 87 degrees Brix to get a final Brix of about 72 to 76 degrees Brix, preferably about 73 degrees Brix, once blended with the pure maple syrup at 66 degrees Brix.

Example 9: Partial treat then concentrate method of manufacture of the shelf stable, thick, pourable maple syrup with a viscosity of common table syrup.

The method is based on the creation of the proper sugar ratio within the final product.

Another way to obtain specific sugar ratio in the maple syrup is to control the conversion of sucrose to glucose and fructose. Sucrose is converted to glucose and fructose by the use of the enzyme invertase. As the reaction progresses, the amount of sucrose in the vessel decreases directly in proportion to the increase in the amount of glucose and fructose. Therefore, the reaction can be stopped at any desired sucrose:glucose and fructose ratio. Termination of the reaction is achieved by the

inactivation or removal of the enzyme. In this application, this is accomplished by heat treatment of the treated syrup or by filtration. Once the reaction is terminated, the syrup is concentrated by accepted methods to the desired final concentration, which corresponds to the viscosity of common table syrups.

5

Example 10. Blend maple syrup and unconcentrated ETS (enzyme treated syrup) followed by concentration method of manufacture of the shelf stable, thick, pourable maple syrup product with a viscosity of common table syrup

10 The following method of manufacture of a shelf stable, thick, pourable maple syrup has been devised. The method is based on the creation of the proper sugar ratio within the final product.

Another method of obtaining a shelf stable, thick, pourable maple syrup is to blend 15 pure maple syrup with enzyme treated maple syrup in the appropriate proportions such that the final product has the correct sugar ratio followed by concentrating the whole blend to the desired final concentration using applicable evaporation technologies. The preferred sugar ratio would be 30:70 or 40:60 ETS:untreated maple syrup.

20 Experiment 11: Viscosity of Common Table Syrups and Concentrated Maple Syrup

The viscosity of various table syrups was determined using a Brockfield Model HAT Viscometer at 22°C. The viscosity was measured at five rotational speeds (0.5, 1, 2.5, 5, 10, 20 50, and 100) with two different spindles attached (#3 and #4). The following 25 results were obtained.

Table 3: Apparent Viscosity of Common Table Syrups

Syrup	Apparent Viscosity*, Centipoise (Cp)
Aunt Jemima™	1500
Log Cabin™	1800
Mrs Butterworth's™	1900

* Apparent Viscosity at 2.5 RPM, average between the two spindles

30

Sugar solutions have a specific viscosity at a given concentration. In order to make a shelf stable, thick, pourable maple syrup that has a viscosity of common table syrups, the viscosity of various maple sugar concentrations were measured. Maple syrup at 66 degrees Brix was blended with a high viscosity maple syrup (86 degrees Brix, 55:45 ETS:Pure blend) such that the resultant Brix of the sugar solution was 72, 74, 76, 78, and 80. The viscosity was then measured using the same protocol as for the table syrups. The following results were obtained.

10

Table 4: Apparent Viscosity of Sugar Solutions at Various Concentrations

15

Sugar Solution Concentration, Degrees Brix	Apparent Viscosity*, Centipoise (Cp)
72	1000
74	1500
76	3000
78	4800
80	12600

20

By comparing the viscosity data of Table 2 and Table 3, it was determined that in order to create a maple syrup that has a similar viscosity to that of common table syrup, the maple syrup blend needed to be concentrated to about 72 to 76 degrees Brix, preferably to about 72-74 degrees Brix.

Experiment 12: Determination Sugar Blend Ratio Needed to Create a Shelf Stable, Pourable Thick Maple Syrup with a Viscosity of Common Table Syrup.

30

The stability of a thick maple syrup is dependent on the glucose, fructose and sucrose ratio. As previously described, concentrated sucrose solutions crystallizes much more readily in comparison to solutions of glucose, fructose or glucose and fructose mixtures at similar concentrations. Therefore, to obtain a product that has extended shelf-life, the proper blend of glucose, fructose and sucrose must be attained.

35

Five different blends of glucose and fructose (enzyme treated maple syrup) and sucrose (pure maple syrup) were made, corresponding to the following ratios: 10:90, 20:80, 30:70, 40:60 and 50:50 enzyme treated syrup: pure maple syrup. All samples were monitored on a regular basis, making observations on the appearance of any crystals.

5 Crystals were observed in the 10:90 and 20:80 samples after five days. At 6 months no crystals were observed in the other three samples.

Example 13: Maple Butter Containing Stabilizers

10 To test the efficacy of two stabilizers, a small-scale scraped surface heat exchanger (Sweden, Soft Serve Ice Cream Maker) was used to cool concentrated maple syrup solutions (86 degrees Brix) under high amounts of agitation. The following stabilizer levels of each stabilizer were evaluated: carageenan 0.05, 0.1 and 0.5 percent; xanthan 0.05, 0.1 and 0.33 percent. The stabilizer was first dispersed in a small quantity of

15 water which was then blended with the unconcentrated maple syrup. Concentration to 86 degrees Brix was carried out in an open vat heated by steam. The hot concentrated product was then transferred into sealed containers that were subsequently cooled quiescently to 4 degrees Celsius. The cooled product was then poured into the scraped surface heat exchanger and agitated for 6 minutes. Product was drawn off the machine

20 into 500 mL clear plastic containers. Samples from each stabilizer type and from each stabilizer concentration were monitored for stability as a function of time. All samples performed well, with limited amounts of separation being observed. The samples containing 0.1 percent carageenan did not show any signs of separation for 14 months. Similarly, the samples containing 0.1 percent xanthan also showed no signs of

25 separation at 14 months.

Example 14: Stabilization of Maple-based products Using Enzyme treated Maple Syrup

Enzyme treated maple syrup was produced by enzyme-treating maple syrup in the

30 manner described in Example 3. This enzyme treated maple syrup was added to untreated maple syrup to produce maple butter. Various blends of ratios of maple syrup with enzyme treated maple syrup were evaluated to test the stabilization effect of

enzyme treated maple syrup to maple butter. The two blend ratios used in this experiment were 20% enzyme treated maple syrup to 80% maple syrup, and 25% enzyme treated maple syrup to 75% maple syrup. The resultant blends were concentrated to 86 degrees Brix and cooled rapidly under high agitation. A very smooth creamy maple butter product was obtained in both 20 and 25 percent enzyme treated maple syrup to maple syrup blends. Both products did not shown any signs of separation for 12 months.

Example 15: Maple Butter with Air or Inert Gas and Enzyme Treated Maple Syrup

10

A sensory evaluation of maple butter made in example 9 indicated that the sugar intensity could be considered too strong for some people. Also, the product was somewhat difficult to spread. To increase product spreadability and to create a more appealing flavor profile, maple butter was made by concentrating to only 84 degrees Brix and whipping air into the product while cooling. To aid in the retention of the air and with the spreadability, 0.01 percent xanthan was also added to the blend before concentration. Five different amounts of air were incorporated into the maple butter after crystallizing (rapid cooling with agitation), corresponding to 9, 15, 26, 35 and 37 percent over-run in the sample.

20

Stability of products varied with the amount of air incorporated into the product. Air bubble destabilization (bubble growth) was visible after about 1 week of storage at room temperature for samples containing more than 15 percent air. Samples containing more than 15 percent air also exhibited phase separation after approximately 3 weeks of storage at room temperature.

25

Maple syrup product, containing 25 percent enzyme treated maple syrup, 75 percent maple syrup and 0.01 percent xanthan was concentrated to 85 degrees Brix. When 12 percent air was added in the rapid cooling and agitation process, air bubble destabilization was minimal, and phase separation was not detected, after 1 month of storage at room temperature. Optionally, inert gas, for example nitrogen, may be used.

30

While the invention has been particularly shown and described with a reference to a preferred embodiment, it will be understood by those skilled in the art that various changes in form and detail may be made therein without departing from the spirit and scope of the invention.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A method of producing an enzyme treated maple syrup, comprising:
 - (a) adding a sucrose-cleaving enzyme to a maple syrup; and
 - (b) incubating the maple syrup to produce the enzyme treated maple syrup.
2. The method of claim 1 which further comprises the step of concentrating the enzyme treated maple syrup.
3. The method of claim 1 or 2 which further comprises the step of removing or inactivating the sucrose-cleaving enzyme.
4. The method of any one of claims 1 to 3 wherein the sucrose-cleaving enzyme is invertase.
5. The method of any one of claims 1 to 4 wherein the maple syrup is diluted to 55 to 66 degrees Brix prior to the step of adding a sucrose-cleaving enzyme.
6. A method for producing a shelf stable, thick, pourable, pure maple syrup product with a viscosity of common table syrup, comprising:
 - a) adding a sucrose-cleaving enzyme to a maple syrup;
 - b) incubating the maple syrup to produce an enzyme treated maple syrup;
 - c) adding an untreated maple syrup to the enzyme treated maple syrup at a predetermined ratio of enzyme treated maple syrup to untreated maple syrup, to produce a maple syrup blend of untreated maple syrup and enzyme treated maple syrup; and
 - d) concentrating the maple syrup blend to produce the shelf stable, thick, pourable, pure maple syrup product with the viscosity of common table syrup.

7. A method for producing a shelf stable, thick, pourable, pure maple syrup product with a viscosity of common table syrup, comprising:

a) adding a sucrose-cleaving enzyme to a maple syrup;

b) incubating the maple syrup to produce an enzyme treated maple syrup;

c) concentrating the enzyme treated maple syrup; and

d) adding an untreated maple syrup to the enzyme treated maple syrup at a predetermined ratio of enzyme treated maple syrup to untreated maple syrup, to produce a maple syrup blend of untreated maple syrup and enzyme treated maple syrup, wherein the maple syrup blend is the shelf stable, thick, pourable pure maple syrup product with the viscosity of common table syrup.

8. A method for producing a shelf stable, thick, pourable, pure maple syrup product with a viscosity of common table syrup, comprising:

a) adding a sucrose-cleaving enzyme to a maple syrup;

b) incubating the maple syrup to produce an enzyme treated maple syrup;

c) inactivating or removing the sucrose-cleaving enzyme; and

d) concentrating the enzyme treated maple syrup to produce the shelf stable, thick pourable, pure maple syrup product with the viscosity of common table syrup.

9. The method of any one of claims 6 to 8 where the sucrose-cleaving enzyme is invertase.

10. The method of any one of claims 6 to 8, which comprises the additional step of diluting the maple syrup to 55 to 66 degrees Brix prior to the step of adding the sucrose-cleaving enzyme.

11. The method of claim 8, wherein the sucrose-cleaving enzyme is removed by filtration.

12. The method of claim 8, wherein the sucrose-cleaving enzyme is inactivated by heat treatment.

13. The method of claim 8 which comprises the additional step of monitoring the ratio of sucrose to fructose and glucose, and removing or inactivating the sucrose- cleaving enzyme at a predetermined sucrose to fructose and glucose ratio, followed by the step of concentrating.
14. The method of any one of claims 6 to 8 wherein the viscosity of the maple syrup product is between 1200 and 2100 centipoise at 2.5 rpm as determined by a Brockfield Viscometer at 22°C.
15. The method of claim 14 wherein the viscosity of the maple syrup is between 1300 and 1900 centipoise at 2.5 rpm as determined by a Brockfield Viscometer at 22°C.
16. The method of claim 6 or 7 wherein the ratio of enzyme treated maple syrup to untreated maple syrup is from 30: 70 to 60: 40 by percentage of sugars present.
17. The method of claim 16 where the ratio of enzyme-treated maple syrup to untreated maple syrup is 40: 60 by percentage of sugars present.
18. The method of claim 7 wherein the enzyme-treated maple syrup is between 84 and 89 degrees Brix, and the maple syrup product has a final concentration of between 72 and 76 degrees Brix.
19. The method of claim 18 wherein the ratio of enzyme treated maple syrup to untreated maple syrup is 40: 60, and the enzyme treated maple syrup is concentrated to between 84 to 89 degrees Brix, and the product has a final concentration of between 72 and 76 degrees Brix.
20. The method of any one of claims 6 to 8, wherein the maple syrup product is concentrated to between 72 and 76 degrees Brix.
21. The method of any one of claims 6 to 8, wherein the maple syrup product is concentrated to between 72 and 74 degrees Brix.
22. The method of claim 21, wherein the maple syrup product is concentrated by a method selected from the group consisting of flash evaporation and under vacuum using low heat of less than 65 degrees Celsius.

23. A shelf stable, thick, pourable, pure maple syrup product made by the method of any one of claims 1 to 5.
24. A shelf stable, thick, pourable, pure maple syrup product made by the method of any one of claims 6 to 22.
25. A shelf stable, thick, pourable, pure maple syrup product having a Brix measurement of between 70 and 76 degrees and a viscosity of between 1200 and 2100 centipoise at 2.5 rpm as determined by a Brockfield Viscometer at 22°C.
26. A shelf stable, thick, pourable, pure maple syrup product of claim 26, having a Brix measurement of between 72 and 74 degrees and a viscosity of between 1300 and 1900 centipoise at 2.5 rpm as determined by a Brockfield Viscometer at 22°C.
27. Use of the shelf stable, thick, pourable and pure maple syrup product of claim 25 as a topping, sweetener or ingredient in a food.
28. A method for producing a shelf stable, spreadable maple syrup product with a consistency of clear honey, comprising:
- a) adding a sucrose-cleaving enzyme to a maple syrup;
 - b) incubating the maple syrup to produce an enzyme treated maple syrup;
 - c) adding an untreated maple syrup to the enzyme treated maple syrup at a predetermined ratio of enzyme treated maple syrup to untreated maple syrup, to produce a maple syrup blend of untreated maple syrup and enzyme treated maple syrup; and
 - d) concentrating the maple syrup blend to produce the shelf stable, spreadable maple syrup product with the consistency of clear honey.
29. A method for producing a shelf stable, spreadable maple syrup product with the consistency of clear honey, comprising:
- a) adding a sucrose-cleaving enzyme to a maple syrup;

- b) incubating the maple syrup to produce an enzyme treated maple syrup;
 - c) inactivating or removing the sucrose-cleaving enzyme; and
 - d) concentrating the enzyme treated maple syrup to produce the shelf stable, spreadable maple syrup product with the consistency of clear honey.
30. The method of claim 28 or 29 wherein the sucrose-cleaving enzyme is invertase.
31. The method of claim 28 or 29, which comprises the additional step of diluting the maple syrup to 55 to 66 degrees Brix prior to the step of adding the sucrose-cleaving enzyme.
32. The method of claim 29, wherein the sucrose-cleaving enzyme is removed by filtration.
33. The method of claim 29, wherein the sucrose-cleaving enzyme is inactivated by heat treatment.
34. The method of claim 29 which comprises the additional step of monitoring the ratio of sucrose to fructose and glucose, and removing or inactivating the sucrose-cleaving enzyme at a predetermined sucrose to fructose and glucose ratio, followed by the step of concentrating.
35. The method of claim 28 wherein the ratio of enzyme treated maple syrup to untreated maple syrup is from 45: 55 to 60: 40 by percentage of sugars present.
36. The method of claim 35 where the ratio of enzyme-treated maple syrup to untreated maple syrup is 55: 45 by percentage of sugars present.
37. The method of claim 28 or 29 wherein the maple syrup product has a final concentration of between 72 and 90 degrees Brix.
38. The method of claim 37 wherein the maple syrup product has a final concentration of between 79 and 83 degrees Brix.

39. The method of claim 38, wherein the maple syrup product is concentrated by a method selected from the group consisting of flash evaporation or under vacuum using low heat of less than 65 degrees Celsius.
40. A shelf stable, spreadable maple syrup product with the consistency of clear honey made by the method of any one of claims 28 to 39.
41. A shelf stable, spreadable maple syrup product with the consistency of clear honey having a Brix measurement of between 72 and 90 degrees.
42. A shelf stable, spreadable maple syrup product with the consistency of clear honey having a Brix measurement of between 79 and 83 degrees.
43. A method of producing a shelf stable maple butter with the consistency of churned honey, comprising:
- a) combining a maple syrup with an enzyme treated maple syrup, one or more stabilizers, or a combination of both to form a maple syrup solution; and
 - b) concentrating the maple syrup solution to form shelf stable maple butter.
44. The method of claim 43 wherein the enzyme treated maple syrup is made according to the method of claim 1.
45. The method of claim 43 wherein the stabilizer is a viscosity altering hydrocolloid substance.
46. The method of claim 45 wherein the stabilizer is selected from the group consisting of xanthan and carageenan.
47. The method of claim 45 wherein the concentration of stabilizer is between 0.01 and 1 percent.
48. The method of claim 45 wherein the concentration of stabilizer is 0.1 percent.

49. The method of claim 44 wherein the ratio of enzyme treated maple syrup to maple syrup is from 20: 80 to 25: 75 by percentage of sugars present.
50. The method of claim 43 wherein the maple butter is concentrated to between 83 degrees Brix and 87 degrees Brix.
51. A method of producing a shelf stable maple butter, comprising:
- (a) adding a sucrose-cleaving enzyme to a maple syrup;
 - (b) incubating the maple syrup to produce an enzyme treated maple syrup;
 - (c) inactivating or removing the sucrose cleaving enzyme; and
 - (d) concentrating the enzyme treated maple syrup to produce the shelf stable maple butter.
52. The method of claim 51 wherein the sucrose-cleaving enzyme is invertase.
53. The method of claim 51 which comprises the additional step of diluting the maple syrup to 55 to 66 degrees Brix prior to the step of adding the sucrose cleaving enzyme.
54. The method of claim 51 wherein the sucrose-cleaving enzyme is removed by filtration.
55. The method of claim 51 wherein the sucrose-cleaving enzyme is inactivated by heat treatment.
56. The method of claim 51 which comprises the additional step of monitoring the ratio of sucrose to fructose and glucose, and removing or inactivating the sucrose cleaving enzyme at a predetermined sucrose to fructose and glucose ratio, followed by the step of concentrating.
57. The maple butter made by the method of claim 43 or 51.
58. A shelf stable maple butter that is a pure maple-based product.
59. A method of producing an enzyme treated maple syrup, that when added to a product functions to prevent the crystallization and separation of another product, comprising:

- a) adding a sucrose-cleaving enzyme to a maple syrup;
- b) incubating the maple syrup to produce the enzyme treated maple syrup; and
- c) adding the enzyme treated maple syrup to the other product.

60. The method of claim 59, wherein the product to which the enzyme treated maple syrup is added is a maple based product.

61. The method of claim 59 which further comprises the step of concentrating the enzyme treated maple syrup.

62. The method of claim 59 or 60 which further comprises the step of removing or inactivating the sucrose-cleaving enzyme.

63. The method of any one of claims 59 to 61 wherein the sucrose-cleaving enzyme is invertase.

64. The method of any one of claims 59 to 62 wherein the maple syrup is diluted to 55 to 66 degrees Brix prior to the step of adding a sucrose-cleaving enzyme.

65. An enzyme treated maple syrup that when added to another product prevents the crystallization and separation of the product.

66. A blended product comprising an enzyme treated maple syrup and a second product.

67. An enzyme treated maple syrup that when added to a maple based product prevents the crystallization and separation of the maple based product.

68. A blended product comprising a maple based product and an enzyme treated maple syrup.

69. A method of stabilizing a maple-based product, comprising:

- a) combining a maple syrup with an enzyme treated maple syrup, one or more stabilizers, or a combination of both to form a maple syrup solution; and

b) adding the maple syrup solution to another product.

70. The method of claim 69 which further comprises the step of concentrating the maple syrup solution.

71. The method of claim 69 wherein the enzyme treated maple syrup is made according to the method of claim 1.

72. The method of claim 69 wherein the stabilizer is a viscosity altering hydrocolloid substance.

73. The method of claim 70 further comprising the addition of air or an inert gas after concentrating the solution.

74. A stabilized maple-based product made by the method of claim 69.

75. A method of stabilizing a product, comprising combining the product with an enzyme treated maple syrup.

76. A method of making a maple-based product with an improved flavor profile comprising:

(a) adding enzyme treated maple syrup to maple syrup;

(b) concentrating the maple syrup; and

(c) adding air or an inert gas after concentrating the maple syrup.

77. The method of claim 76 wherein the air or inert gas is added after concentrating the maple syrup solution.

78. The method of claim 76 or 77 wherein air or inert gas is added to a final concentration of 1 to 15 percent air to 99 to 85 percent maple syrup.

79. The method of claim 76 or 77 wherein air or inert gas is added to a final concentration of 12 percent.

80. A maple-based product with an improved flavor profile made by the method of any one of claims 76 to 79.

81. Use of the product of any one of claims 57, 58, 65-68, 79 or 80 as an ingredient in pure maple-based products, which comprises adding the product to a pure maple-based product during manufacture of the pure maple-based product.

82. A method for producing a shelf stable, spreadable maple syrup composition which comprises:

- (a) adding a sucrose-cleaving enzyme to maple syrup;
- (b) incubating the resulting maple syrup solution; and
- (c) concentrating the resulting maple syrup solution.

83. The method of claim 82 which additionally comprises the step of removing the sucrose-cleaving enzyme.

84. The method of claim 83 where the sucrose-cleaving enzyme is removed by filtration.

85. The method of any one of claims 82 to 84, wherein the sucrose-cleaving enzyme is invertase.

86. The method of any one of claims 82 to 85, wherein the maple syrup is diluted to 55 degrees Brix prior to the step of adding a sucrose-cleaving enzyme.

87. A method for producing a shelf stable, spreadable maple syrup composition which comprises:

- (d) diluting maple syrup to 55 degrees Brix;
- (e) adding a sucrose-cleaving enzyme to the maple syrup; where the sucrose-cleaving enzyme is invertase;
- (f) incubating the resulting maple syrup solution; and
- (g) concentrating the resulting maple syrup solution.

88. The method of claim 87 which comprises the additional step of blending enzyme-treated maple syrup with untreated maple syrup prior to the step of concentrating the resulting maple syrup solution.
89. The method of claim 88 wherein the ratio of enzyme treated maple syrup to untreated maple syrup is from 45:55 to 60:40 by percentage of sugars present.
90. The method of claim 89 wherein the ratio of enzyme-treated maple syrup to untreated maple syrup is 55:45 by percentage of sugars present.
91. The method of any one of claims 87 to 90, wherein the resulting maple syrup solution is concentrated to between 75 degrees Brix and 90 degrees Brix.
92. The method of any one of claims 87 to 90, wherein the resulting maple syrup solution is concentrated to between 77 and 87 degrees Brix.
93. The method of any one of claims 87 to 90, wherein the resulting maple syrup solution is concentrated to between 82 and 85 degrees Brix.
94. The method of any one of claims 87 to 90, wherein the resulting maple syrup solution is concentrated to 82 degrees Brix.
95. The method of claim 87 wherein the resulting maple syrup solution is concentrated under vacuum using low heat of less than 65 degrees Celsius.
96. The shelf stable maple syrup composition made by the method of claim 82, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
97. The shelf stable maple syrup composition made by the method of any one of claims 87 to 90, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
98. The shelf stable maple syrup composition made by the method of claim 91, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.

99. The shelf stable maple syrup composition made by the method of claim 92, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
100. The shelf stable maple syrup composition made by the method of claim 93, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
101. The shelf stable maple syrup composition made by the method of claim 94, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
102. A shelf stable, spreadable maple syrup composition having a Brix measurement of between 75 and 90 degrees, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
103. A shelf stable, spreadable maple syrup composition having a Brix measurement of between 77 and 87 degrees, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
104. A shelf stable, spreadable maple syrup composition, having a Brix measurement of between 82 and 85 degrees, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
105. A shelf stable, spreadable maple syrup composition, having a Brix measurement of 82 degrees, wherein the maple syrup composition is free from crystallization for a period of at least 3 months.
106. The method of using the composition of claim 97 as a spread, which comprises spreading the composition on a food product.
107. The method of using the composition of claim 96 as a sweetener which comprises adding the composition to a food product.
108. The method of using the composition of claim 96 as a topping for a food product, which comprises applying the composition to a food product.

109. The method of using the composition of claim 96 as an ingredient in pure maple-based products, which comprises adding the composition to a pure maple-based product during manufacture of the pure maple-based food product.

110. A method of stabilizing a pure maple-based product, comprising combining untreated maple syrup with a sucrose-cleaving enzyme treated maple syrup, in a ratio suitable for forming a stabilized, pure maple-based product.

111. The method of claim 110 which further comprises the step of concentrating the combined untreated maple syrup and sucrose-cleaving enzyme treated maple syrup to form the stabilized, pure, maple-based product.

112. The method of claim 110 wherein the sucrose-cleaving enzyme treated maple syrup is made according to a method comprising the steps of: (a) adding a sucrose-cleaving enzyme to a maple syrup; and (b) incubating the maple syrup to produce the sucrose-cleaving enzyme treated maple syrup.

113. A stabilized, pure maple-based product made by the method of claim 110.

114. A method of producing a shelf stable pure maple butter, comprising:

- (a) combining a maple syrup with a sucrose-cleaving enzyme treated maple syrup in a ratio suitable for forming a shelf stable maple butter; and
- (b) concentrating the product of step (a) to form pure shelf table maple butter.

115. The method of claim 114 wherein the sucrose-cleaving enzyme treated maple syrup is made according to a method comprising the steps of: (a) adding a sucrose-cleaving enzyme to a maple syrup; and (b) incubating the maple syrup to produce the sucrose-cleaving enzyme treated maple syrup.

116. The method of claim 114 wherein the ratio of sucrose-cleaving enzyme treated maple syrup to maple syrup is from 20:80 to 25:75 by percentage of sugars present.

117. The method of claim 51 which comprises the additional step of monitoring the ratio of sucrose to fructose and glucose during step (b), and conducting step (c) at a predetermined sucrose to fructose and glucose ratio.

118. The method of using a sucrose-cleaving enzyme treated maple syrup as an ingredient in pure maple-based products, which comprises adding the sucrose cleaving enzyme treated maple syrup to pure maple product during manufacture of the pure maple-based product in an amount sufficient to stabilize the pure maple-based product.

119. A method of producing a shelf stable pure maple based product, the method comprising:

- (a) adding a sucrose-cleaving enzyme to a maple syrup;
- (b) incubating the maple syrup to produce the enzyme treated maple syrup;
- (c) adding the enzyme treated maple syrup to a maple base product in a ratio suitable for forming a shelf-stable pure maple-based product; and
- (d) concentrating the product of step (c) to produce the shelf stable pure maple based product.

120. The method of claim 119 which further comprise the step of removing or inactivating the sucrose-cleaving enzyme prior to the step of concentrating.

121. The method of claim 119 wherein the sucrose-cleaving enzyme is invertase.

122. The method of claim 120 wherein the sucrose-cleaving enzyme is removed by filtration.

123. The method of claim 120 wherein the sucrose-cleaving enzyme is inactivated by heat treatment.

124. The method of claim 120 which comprises the additional step of monitoring the ratio of sucrose to fructose and glucose during step (b) and removing or inactivating the sucrose-cleaving enzyme at a predetermined sucrose to fructose and glucose ratio.

125. The method of claim 119 wherein the product of step (c) is concentrated to a Brix level of between 72 degrees Brix and 90 degrees Brix.

126. The method of claim 119 wherein the maple syrup product is concentrated to between 83 and 87 degrees Brix.

127. The method of claim 119, wherein the maple syrup product is concentrated by a method selected from the group consisting of flash evaporation and under vacuum using low heat of between 50 and 65 degrees Celsius.

128. The method of claim 112 wherein the maple syrup solution is concentrated to between 83 degrees Brix and 87 degrees Brix.

129. The method of using the product of claim 113 as a spread, sweetener or topping which comprises adding the product to food.

130. The method of claim 110 wherein the ratio of sucrose-cleaving enzyme treated maple syrup to untreated maple syrup is between 20:80 and 25:75 by percentage of sugars present.

131. The method of claim 110 wherein the pure, stabilized maple-base product contains a ratio of glucose and fructose to sucrose, of between 20:80 to 25:75 by percentage of sugars present.

132. The method of claim 112 which comprises the additional step of diluting the maple syrup to 55 to 66 degrees Brix prior to step (a).

133. The method of claim 112 which further comprises the step of removing or inactivating the sucrose-cleaving enzyme prior to combining the untreated maple syrup with the sucrose-cleaving enzyme treated maple syrup.

134. The method of claim 115 which further comprises the step of removing or inactivating the sucrose-cleaving enzyme prior to combining the maple syrup with the sucrose-cleaving enzyme treated maple syrup.

135. The method of claim 134 wherein the sucrose-cleaving enzyme is removed by filtration.

136. The method of claim 134 wherein the sucrose-cleaving enzyme is inactivated by heat treatment.

137. The method of claim 115 which comprises the additional step of diluting the maple syrup to 55 to 66 degrees Brix prior to the step of adding the sucrose cleaving enzyme.

138. The method of claim 114 wherein the shelf stable pure maple butter contains a ratio of glucose and fructose to sucrose of between 20:80 to 25:75 by percentage of sugars present.
139. The method of claim 117 wherein the predetermined ratio of sucrose to fructose and glucose is between 75:25 to 80:20 by percentage of sugars present.
140. The method of claim 119 wherein the ratio of enzyme treated maple syrup to maple-based product is between 20:80 to 25:75 by percentage of sugars present.
141. The method of claim 119 wherein the shelf stable pure maple-based product contains a ratio of glucose and fructose to sucrose of between 20:80 to 25:75 by percentage of sugars present.
142. The method of claim 119 which comprises the additional step of diluting the maple syrup to 55 to 66 degrees Brix prior to the step of adding the sucrose cleaving enzyme.
143. The method of claim 132 wherein the sucrose-cleaving enzyme is removed by filtration.
144. The method of claim 132 wherein the sucrose-cleaving enzyme is inactivated by heat treatment.
145. The method of claim 133 which comprise the additional step of monitoring the ratio of sucrose to fructose and glucose during step (b) and removing or inactivating the sucrose-cleaving enzyme at a predetermined sucrose to fructose and glucose ratio.
146. The method of claim 76 wherein the enzyme treated maple syrup is made according to the method of claim 1.
147. Use of the product of any one of claims 23, 24, 57, 58, 65-68, 74 or 80 as a spread, sweetener or topping which comprises adding the product to food.
148. Use of the product of any one of claims 23, 24, 57, 58, 65-68, 74 or 80 as a snack food.
149. Use of the shelf stable, spreadable maple syrup product with the consistency of clear honey of any one of claims 40 to 42 as a topping, sweetener or ingredient in a food.

Viscosity of a 55:45 Enzyme Treated to Non-Enzyme Treated Blend of Maple Syrup at Various Concentrations at 25°C




Figure 1.

2/2




Figure 2.

Viscosity of a 55:45 Enzyme Treated to Non-Enzyme Treated Blend of Maple Syrup at Various Concentrations at 25°C

