

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 2500730 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **G 01 N 33/68 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-05-28**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-03-07**
- (86) Europæisk ansøgning nr.: **12164660.8**
- (86) Europæisk indleveringsdag: **2007-12-21**
- (87) Den europæiske ansøgnings publiceringsdag: **2012-09-19**
- (30) Prioritet: **2006-12-22 DK 200601709** **2006-12-22 US 876838 P**
2007-06-29 DK 200700956 **2007-06-29 US 947074 P**
- (62) Stamansøgningsnr: **07858108.9**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR**
- (73) Patenthaver: **Hvidovre Hospital, Kettegårds Allé 30, 2650 Hvidovre, Danmark**
- (72) Opfinder: **Eugen-Olsen, Jesper, Vedbendvej 14, 2900 Hellerup, Danmark**
Haugaard, Steen B, Langelandsvej 32, 4, 2000 Frederiksberg, Danmark
ANDERSEN, Ove, Hvilevej 13, 2900 Hellerup, Danmark
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Rued Langgaards Vej 8, 2300 København S, Danmark**
- (54) Benævnelse: **Opløselig urokinase-plasminogen-aktivatorreceptor (suPAR) som diagnostisk markør for en lavgrad-inflammation**
- (56) Fremdragne publikationer:
WO-A-03/031650
OSTROWSKI S R ET AL: "Soluble Urokinase Receptor Levels in Plasma during 5 Years of Highly Active Antiretroviral Therapy in HIV-1-Infected Patients", J ACQUIR IMMUNE DEFIC SYNDR, vol. 35, no. 4, 1 April 2004 (2004-04-01), pages 337-342, XP002446705,
OSTROWSKI S R ET AL: "Plasma levels of intact and cleaved urokinase receptor decrease in HIV-1-infected patients initiating highly active antiretroviral therapy", SCAND J IMMUNOL, vol. 63, no. 6, June 2006 (2006-06), pages 478-486, XP002446704, -& OSTROWSKI S R ET AL: "High plasma levels of intact and cleaved soluble urokinase receptor reflect immune activation and are independent predictors of mortality in HIV-1-infected patients.", J ACQUIR IMMUNE DEFIC SYNDR, vol. 39, no. 1, 1 May 2005 (2005-05-01), pages 23-31, XP002488617,
SLOAND E M ET AL: "A role for plasma urokinase plasminogen activator receptor (uPAR) in thrombosis in paroxysmal nocturnal hemoglobinuria (PNH)", BLOOD, vol. 104, no. 11, Part 1, November 2004 (2004-11), page 84A, XP008099893, & 46TH ANN MEETING AM SOC HEMATOL; SAN DIEGO, CA, USA; DEC 04 -07, 2004
ANDERSEN O ET AL: "Soluble urokinase plasminogen activator receptor is a marker of dysmetabolism in HIV-infected patients receiving highly active antiretroviral therapy.", J MED VIROL, vol. 80, no. 2, 20 December 2007 (2007-12-20), pages 209-216, XP008094314,
DANESH JOHN ET AL: "Low grade inflammation and coronary heart disease: Prospective study and updated

Fortsættes ...

meta-analyses", BRITISH MEDICAL JOURNAL, LONDON, GB, vol. 321, no. 7255, 22 July 2000 (2000-07-22), pages 199-204, XP002211089, ISSN: 0267-0623

FALOIA EMANUELA ET AL: "Inflammation as a Link between Obesity and Metabolic Syndrome", JOURNAL OF NUTRITION AND METABOLISM, vol. 100, no. 7, 1 January 2012 (2012-01-01), pages 1863-7, XP055093895, ISSN: 2090-0724, DOI: 10.1161/01.CIR.0000127953.98131.ED

J. EUGEN-OLSEN ET AL: "Circulating soluble urokinase plasminogen activator receptor predicts cancer, cardiovascular disease, diabetes and mortality in the general population", JOURNAL OF INTERNAL MEDICINE, vol. 268, no. 3, 15 September 2010 (2010-09-15), pages 296-308, XP055140843, ISSN: 0954-6820, DOI: 10.1111/j.1365-2796.2010.02252.x

TOLDI GERGELY ET AL: "Soluble urokinase plasminogen activator receptor (suPAR) in the assessment of inflammatory activity of rheumatoid arthritis patients in remission.", CLINICAL CHEMISTRY AND LABORATORY MEDICINE : CCLM / FESCC FEB 2013, vol. 51, no. 2, February 2013 (2013-02), pages 327-332, ISSN: 1434-6621

ISTVÁ• N IVANCSÁĹ ET AL: "Relationship of Circulating Soluble Urokinase Plasminogen Activator Receptor (suPAR) Levels to Disease Control in Asthma and Asthmatic Pregnancy", PLOS ONE, vol. 8, no. 4, 2 April 2013 (2013-04-02), page e60697, XP055403571, DOI: 10.1371/journal.pone.0060697

YANFENG ZHAO ET AL: "Plasma Soluble Urokinase Receptor Level Is Correlated with Podocytes Damage in Patients with IgA Nephropathy", PLOS ONE, vol. 10, no. 7, 13 July 2015 (2015-07-13), page e0132869, XP055403573, DOI: 10.1371/journal.pone.0132869

AZIZ GUMUS ET AL: "Soluble urokinase-type plasminogen activator receptor is a novel biomarker predicting acute exacerbation in COPD", INTERNATIONAL JOURNAL OF CHRONIC OBSTRUCTIVE PULMONARY DISEASE, 13 February 2015 (2015-02-13), page 357, XP055403565, DOI: 10.2147/COPD.S77654

ENOCSSON HELENA ET AL: "Soluble urokinase plasminogen activator receptor-A valuable biomarker in systemic lupus erythemato", CLINICA CHIMICA ACTA, vol. 444, February 2015 (2015-02), pages 234-241, XP029149608, ISSN: 0009-8981, DOI: 10.1016/J.CCA.2015.02.031

STIG LYNGBAEK ET AL: "CRP and suPAR are differently related to anthropometry and subclinical organ damage", INTERNATIONAL JOURNAL OF CARDIOLOGY, vol. 167, no. 3, 1 August 2013 (2013-08-01), pages 781-785, XP055093897, ISSN: 0167-5273, DOI: 10.1016/j.ijcard.2012.03.040

DESCRIPTION

FIELD OF THE INVENTION

[0001] The invention is directed to a method of diagnosing low-grade inflammation in an apparently healthy mammalian subject. The invention is further directed to a method of predicting the risk of developing a low-grade inflammation-related disease in a mammalian subject.

BACKGROUND OF THE INVENTION

[0002] Metabolic syndrome is characterized by a combination of high blood pressure, insulin resistance with increased glucose production and decreased glucose utilization, central (abdominal) obesity, dyslipidemia (namely elevated triglycerides and reduced high-density lipoprotein cholesterol), and increased IL-6 and TN- α production.

[0003] Metabolic syndrome has emerged as an important cluster of risk factors for development of atherosclerotic diseases, e.g. prevalent and incident atherothrombotic cardiovascular disease (CVD), coronary heart disease, (CHD), stroke, peripheral arterial disease, chronic obstructive lung disease, as well as type 2 diabetes. Stimuli such as over-nutrition, physical inactivity and ageing may result in cytokine hyper-secretion and eventually lead to insulin resistance and diabetes in some, but not all, human subjects. According to the clinical criteria developed by the Adult Treatment Panel III (ATP III) Guidelines, it is estimated that 1 out of 4 adults living in the United States merits the diagnosis of metabolic syndrome; however, not all obese individuals develop this syndrome. Cardiovascular disease and Type 2 diabetes are leading causes of death and illness in developed countries, and these chronic diseases are becoming a dominating health problem worldwide. Early diagnosis of metabolic syndrome and prediction of early diagnosis of future cardiovascular events and diabetes would greatly improve the chances of successful prophylactic and therapeutic treatment.

[0004] Low-grade inflammation is a triggering factor of metabolic syndrome. Currently, blood levels of pro-inflammatory cytokines, hypersensitive C-reactive protein (hs-CRP), interleukin 6 (IL-6), tumor necrosis factor alpha (TNF- α), serum amyloid A (SAA), fibrinogen and albumin; plasma viscosity; erythrocyte sedimentation rate (ESR); and leukocyte count are used to indicate the presence of low-grade inflammation and thereby predict the risk of developing metabolic syndrome, associated with vascular risk factors and with prevalent and incident atherothrombotic cardiovascular disease (CVD), coronary heart disease (CHD), stroke, peripheral arterial disease, chronic obstructive lung disease and type 2 diabetes.

[0005] Although apparently healthy individuals develop metabolic syndrome, it has, in recent years, become evident that HIV-infected individuals on treatment can also develop metabolic

syndrome. The widespread implementation of Highly Active Anti-Retroviral Therapy (HAART) in western countries, from 1996 onwards, has increased the prevalence of long-living HIV-infected patients with suppressed HIV RNA and fairly high CD4 counts. However, adverse effects of HAART, in particular the administration of protease inhibitors, lead to a form of Metabolic Syndrome, similar to that observed in non-HIV infected obese individuals, including insulin resistance, fat redistribution, dyslipidemia and other risk factors of cardiovascular disease. There is thus a need to develop early diagnostic and predictive tools to prevent an increase in the morbidity and mortality of HIV-infected patients on HAART who display this cluster of dysmetabolic phenotypes. An individual's immunological status impacts his/her metabolic status, which is reflected in the level of immunological markers, e.g. TNF- α and IL-6. Such immunological markers have been shown to correlate inversely with symptoms of metabolic syndrome in HIV-infected patients as well as in HIV-negative subjects; however, data are not consistent and more stable and stronger predictive markers are needed.

[0006] Urokinase-type Plasminogen Activator Receptor (uPAR, CD87) is the cellular receptor for urokinase (uPA), and is expressed by most leukocytes, including monocytes, macrophages, neutrophils and platelets. uPAR is an activation antigen in monocytes and T cells. uPAR may be shed from the cell surface, generating a soluble form of the receptor (suPAR) lacking the GPI-anchor. The shedding mechanism is poorly understood but may occur by cleavage of the GPI-anchor catalyzed by a GPI-specific phospholipase D. Soluble forms of uPAR (suPAR) have been identified in cell culture supernatants and in diverse biological fluids such as tumor ascites, cystic fluid, serum, plasma and urine. The cellular origin of circulating suPAR is not known. Many, if not all, cells which express uPAR also shed soluble forms of the receptor when cultured *in vitro*.

[0007] Svendsen et al (2006) Surgical Infections 7, 463-471 disclosed that elevated preoperative concentrations of suPAR were associated with the development of pneumonia following surgery for colorectal cancer.

[0008] WO 2005/116077 disclosed the use of a ligand that binds to a binary uPA-uPAR complex, but which does not inhibit uPA-uPAR binding, in the treatment of a large number of diseases, including various cancers and cardiovascular diseases (such as atherosclerosis).

[0009] Ostrowski et al (2004) J. Acquir. Immune Defic. Syndr. 35, 337-342, Ostrowski et al (2006) Scand. J. Immunol. 63, 478-486 and Ostrowski et al (2005) J. Acquir. Immune Defic. Syndr. 39, 23-31 disclose studies into the suPAR levels in HIV-infected patients, and conclude that the level correlates with inflammation in such patients. It decreases when the patients are treated with HAART.

SUMMARY OF THE INVENTION

[0010] The invention provides a method for diagnosing low-grade inflammation in accordance with Claim 1.

[0011] The invention also provides a method for predicting the risk of developing a low-grade inflammation-related disease in accordance with Claim 2.

[0012] The measurement in step (b) of either of these methods may furthermore be compared with a reference value wherein an increase in the level of said marker indicates the grade of low-grade inflammation. In all embodiments of the invention, it is preferable for the mammalian subject not to have been given a blood transfusion in the 24 hours (more preferably, 48 hours or one week) preceding the taking of the sample for the *in vitro* measurement of suPAR level.

[0013] The subject, in a specific embodiment, may be a human subject. The sample from step (a) may undergo processing before step (b) is carried out. For example, it might be frozen and thawed, diluted, concentrated, stabilised, filtered or treated with preservative.

[0014] The reference value is a standard of levels of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in a population. Alternatively, the reference value is obtained by measuring the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR from a sample from a subject with a known level of low-grade inflammation and/or metabolic syndrome. In a particular embodiment, the reference level is ≤ 3.0 ng per ml using the assays known in the art, e.g., the suPARnostic™ kit sold by Virogates A/S, Denmark. The level of said marker in a subject having low-grade inflammation and/or metabolic syndrome is increased at least about 10% as compared to said reference sample. In more specific embodiments, the level of said marker is increased at least about 25%, 50%, 75% and/or above 100%.

[0015] Also disclosed is a method for predicting the risk of developing a metabolic syndrome-related disease and/or low-grade inflammation-related disease, in particular, cancer and/or the risk of mortality within 1 year, 3 years, 5 years, 10 years, 1-10 years, 1-5 years or 5-10 years in a mammalian subject, particularly, a human subject, comprising:

1. (a) performing an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples comprising a biological fluid derived from said subject, and
2. (b) using the measurement values obtained as a factor to predict the risk of developing a metabolic syndrome-related disease and/or low-grade inflammation-related disease, the risk of developing cancer and/or the risk of mortality within e.g., 1, 3, 5, 10 or between 1-10 years, 1-15 years, 5-10 years.

[0016] By the 'risk of mortality' we mean the risk of general mortality, rather than the risk of dying from a particular disease.

[0017] As above, the reference value is a standard of levels of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in a population. Alternatively, the reference value is obtained by measuring the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR from a sample from a subject free of low-grade inflammation-related disease and/or metabolic syndrome-related disease.

[0018] The level of said marker is increased at least about 10% and/or 1 ng/ml, or between about 1-20 ng/ml in said sample as compared to said reference sample. In specific embodiments, the level of marker may be increased at least about 2 ng/ml, 3 ng/ml, 3.5 ng/ml, 5 ng/ml, 8 ng/ml, 10 ng/ml, 15, ng/ml, 20 ng/ml or in the following ranges: (1-3 ng/ml, 1-5 ng/ml, 3-5 ng/ml, 5-10 ng/ml, 10-15 ng/ml, 15-20 ng/ml). In a particular embodiment, an increased risk of developing any of the said diseases or general mortality within any of the periods defined above is diagnosed if the suPAR level is more than about 3.0 ng/ml (or 4, 5 or 6 ng/ml) in men and more than about 3.5 ng/ml (or 4.5, 5.5 or 6.5) in women.

[0019] There is also disclosed a method for monitoring the treatment of low-grade inflammation in a mammalian subject comprising:

1. (a) isolating a sample from a subject undergoing treatment for low-grade inflammation (e.g. anti-inflammatory treatment, cancer);
2. (b) measuring the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), (ii) and/or D2D3 cleavage products of suPAR,

wherein a decrease in the level of said marker used in step (b) is an indication of progression of treatment.

[0020] In a particular embodiment, the level of said marker is decreased at least about 10% and /or 1 ng/ml and preferably from about 1.3-3 ng/ml in said sample as compared to said reference sample during the course of said treatment. Alternatively, the level of said marker may be decreased at least about 25%, 50%, 75%, and/or above 100% or between at least about 10% to about 100%.

[0021] According to the above methods, said low-grade inflammation-related disease may be either cardiovascular disease, chronic obstructive lung disease or diabetes type 2, Alzheimer's disease, chronic eczema, asthma bronchiale, or autoimmune disease. In particular, an inflammation-related disease may be cancer (e.g., lung, breast, leukemia, prostate).

[0022] According to the above methods, said biological fluid is preferably one of blood, blood serum, blood plasma or urine.

[0023] According to one embodiment of the above methods, said subject is either human immunodeficiency virus-infected and receiving Highly Active Anti-Retroviral Treatment

(HAART) or is apparently healthy. If the patient is HIV-infected, then preferably, at least in the context of assessing the risk of mortality, the patient does not have AIDS. By 'healthy' we mean that the patient has not been diagnosed with a life-threatening condition (such as cancer, cardiovascular disease or chronic obstructive lung disease), an auto-immune disease or diabetes type 2.

[0024] In the various embodiments of the invention, the patient may be over 40 years of age, for example over 50 or over 60, and may be under 70 years of age, for example under 60 or under 50. The patient may be Caucasian.

[0025] According to one embodiment of the above methods, said one or more samples are derived from said subject at one or more points in time. Optionally, said one or more measurements are compared to assess the efficacy of a prophylactic course of action and/or treatment.

[0026] According to the above methods, said in vitro measurement is preferably an ELISA assay. Suitable ELISAs for suPAR levels are disclosed in Mizukami et al (1995) Blood 86, 203-211.

[0027] According to the above methods, in step b) one or more of the following additional measurements derived from the subject selected from the group consisting of: age, gender, ethnicity, smoker, non-smoker, systolic/diastolic blood pressure, cholesterol, triglycerides and waist circumference measurement, may be used to diagnose low-grade inflammation.

[0028] The method may further comprise conducting a glucose tolerance test, assaying TNF-alpha, IL-6, leukocyte count, hsCRP, fasting plasma (Blood-) glucose, blood pressure, cholesterol, C-reactive protein and/or triglyceride levels.

[0029] The disclosure is further directed to methods and compositions for treatment of metabolic syndrome and/or low-grade inflammation. Specifically, there is a method comprising administering to a subject in need thereof an amount of a substance that decreases the level of suPAR sufficiently to treat said low-grade inflammation and/or metabolic syndrome. Preferably, the plasma level is reduced to below 6 ng/ml, more preferably to below 5, 4 or 3 ng/ml. The substance that is administered may be an anti-suPAR antibody (e.g., a monoclonal or polyclonal antibody). The substance may alternatively be a suPAR analog or derivative capable of binding uPA or a suPAR cleavage product such as one or more of the D2D3 isoforms. The composition may be a pharmaceutical composition for use in treatment or prevention of low-grade inflammation and/or metabolic syndrome comprising a substance that lowers the level of suPAR in a mammalian subject.

[0030] In a related aspect, there is the use of a substance for preparing a therapeutic composition for the treatment or prevention of low-grade inflammation and/or metabolic syndrome, wherein said substance lowers the level of suPAR in a mammalian subject.

[0031] In a specific aspect, there is a computer readable medium having computer logic stored therein configured to cause, when executed on a data processing system, the data processing system to compute risk and/or change in risk of a low-grade inflammation-related disease, a metabolic syndrome-related disease, cancer and/or mortality within e.g., 1, 3, 5, 10 or between 1-10 years, 1-5 years, 5-10 years comprising:

1. (a) logic for receiving one or more measurements derived from a mammalian subject wherein at least one measurement is an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject;
2. (b) logic for analysis of one or more of said measurements of step (a); and
3. (c) logic for computing risk and/or change in risk of said low-grade inflammation-related disease and/or said metabolic syndrome-related disease (e.g., cardiovascular disease, chronic obstructive lung disease and diabetes type 2), cancer (e.g., breast cancer or lung cancer), and/or mortality in e.g., 1 year, 3 years, 5 years, 10 years, 1-10 years, 1-5 years, 5-10 years based on analysis in step (b) of said measurements.

[0032] In a particular aspect, the logic in (a) comprises a logic for receiving two or more measurement derived from a human subject wherein at least one measurement is an *in vitro* measurement of the level of marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject. If the data processing system is used to compute risk of developing diabetes, the logic may further receive measurements of blood glucose levels derived from a mammalian subject; if the data processing system is used to compute risk of developing cardiovascular disease, the logic may further receive measurements of blood pressure, HDL-cholesterol, LDL-cholesterol, total cholesterol, hsCRP, and/or triglycerides; if the data processing system is used to compute risk of mortality in a mammalian subject in e.g., 1 year, 3 years, 5 years, 10 years, 1-5 years, 1-10 years or 5-10 years the logic may further receive measurements of brain natriuretic peptide concentrations.

[0033] The data processing system in said computer readable medium may be further caused to monitor progression of treatment of low-grade inflammation-related disease and/or metabolic syndrome-related disease from said change in risk.

[0034] The invention further encompasses a computer program product, comprising the computer readable medium of the present invention. A "computer program product" is used to generally refer to media such as a hard drive and removable storage drive and storage unit. In one embodiment, the software can be stored in a computer program product and loaded into the computer using a removable storage drive, hard drive and communications interface.

[0035] The disclosure is further directed to a computer for determining the risk and/or change in risk of developing a metabolic syndrome-related disease, low-grade inflammation-related disease, cancer and/or mortality within e.g., 1 year, 3 years, 5 years, 10 years or between 1

and 10 years, 1 and 5 years, or 5 and 10 years comprising

1. (a) a computer-readable data storage medium configured to store a data storage material encoded with computer-readable data, wherein said data comprises one or more measurements derived from one of more samples from a mammalian subject, wherein at least one measurement is an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR;
2. (b) a storage medium having stored therein instructions for processing said computer-readable data;
3. (c) a central-processing unit responsive to said instructions coupled to said storage medium and to said computer-readable data storage medium for processing said computer-machine readable data into risk measurement and treatment progression values; and
4. (d) an output device coupled to said central-processing unit of (c) for outputting said risk and/or change in risk values.

[0036] In a particular aspect, the computer-readable data storage medium (a) of the computer comprises data comprising two or more measurements derived from a human subject wherein at least one measurement is an *in vitro* measurement of the level of marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject. If the computer is used to compute risk of developing diabetes, the data may further comprise measurements of blood glucose levels from a mammalian subject; if the computer is used to compute risk of developing cardiovascular disease, the data may further comprise measurements of blood pressure, HDL-cholesterol, LDL-cholesterol, total cholesterol and/or triglycerides; if the computer is used to compute risk of mortality in a mammalian subject within e.g., 1 year, 3 years, 5 years, 10 years or 1-10 years, 1-5 years, or 5-10 years, the data may further comprise measurements of brain natriuretic peptide concentrations.

[0037] In a specific aspect, there is a client computer for providing access to a risk computation system for determining the risk and/or change in risk of developing a metabolic syndrome-related disease, low-grade inflammation-related disease (e.g., cancer) and/or risk of mortality in a mammalian subject within, e.g., 1 year, 3 years, 5 years, 10 years or between 1-10 years, 1-5 years, 5-10 years, the client computer comprising:

1. (a) an input device for receiving one or more measurements derived from a mammalian subject wherein at least one measurement is an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject;
2. (b) a communication interface for communicating the received one or more measurements to a server computer configured to analyse said one or more

- measurements of step (a); and compute a risk value; and
3. (c) a communications interface for receiving, from the server computer a computed risk value.

[0038] In a particular aspect, the input device (a) of the client computer receives two or more measurements derived from a human subject wherein at least one measurement is an in vitro measurement of the level of marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject. If the client computer is used to compute risk of developing diabetes, the input device (a) may further receive measurements of blood glucose levels from a mammalian subject; if the client computer is used to compute risk of developing cardiovascular disease, the input device (a) may further receive measurements of blood pressure, HDL-cholesterol, LDL-cholesterol, total cholesterol and/or triglycerides; if the client computer is used to compute risk of developing cancer (e.g., breast cancer or lung cancer); if the client computer is used to compute risk of mortality in a mammalian subject within e.g., 1 year, 3 years, 5 years, 10 years or between 1-10 years, 1-5 years, 5-10 years, the input device (a) may further receive measurements of NT-proBNP.

[0039] In yet another aspect, there is a computer based method for determining the risk of developing a metabolic syndrome-related disease, low-grade inflammation-related disease (e.g., cancer) and/or risk of mortality in a mammalian subject, e.g., within 1 year, 3 years, 5 years, 10 years or between 1-10 years, 1-5 years or 5-10 years comprising:

1. (a) receiving under computer control in vitro measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject;
2. (b) analyzing under computer control said in vitro measurement received in (a)
3. (c) calculating under computer control a risk value for developing a metabolic syndrome-related disease, low-grade inflammation-related disease (e.g., cancer) and/or risk of mortality in a mammalian subject within 5 years, 10 years or between 5-10 years using analyzed measurements obtained in (b).

BRIEF DESCRIPTION OF THE DRAWINGS

[0040]

Figure 1 shows box-plots showing fasting serum suPAR concentration in HIV infected patients receiving HAART with lipodystrophy (LIPO) and without lipodystrophy (NONLIPO). Box and whisker plots show the 10th, 25th, 50th (median), 75th and 90th percentiles. Extreme cases of individual variables are indicated by asterisks.

Figure 2 shows correlation between suPAR and measures of glucose metabolism and insulin sensitivity. **Figure 2A** shows correlation between suPAR and insulin stimulated nonoxidative glucose metabolism (NOGM). ●, lipodystrophic HIV-infected patients; Δ, nonlipodystrophic HIV-infected patients. Linear correlation line, correlation coefficient and significance level are indicated. **Figure 2B** shows correlations between plasma suPAR level versus HOMA-IR (homeostasis model assessment insulin resistance index; Matthews et al. 1985) in HIV-infected patients on HAART (highly active antiretroviral therapy), who either displayed lipodystrophy (LIPO, n=18) [●]; or was free of lipodystrophy (NONLIPO, n=18) [Δ]. Further, a group of HIV-infected patients naïve to HAART (NAÏVE, n=7) [], was included in the analysis. Linear regression lines including all patients in analysis are presented. LIPO and NONLIPO (n=36) p-suPAR vs. HOMA-IR $r=0.58$, $p<0.001$ (age corrected $r=0.49$, $p=0.003$). All (n=43) p-suPAR vs. HOMA-IR $r=0.49$, $p<0.001$ (age corrected $r=0.45$, $p=0.003$). **Figure 2C** shows correlations between suPAR versus 2 hours plasma glucose after a standard oral glucose (75 gram) challenge in HIV-infected patients on HAART. ●, lipodystrophic HIV-infected patients; Δ, nonlipodystrophic HIV-infected patients. Linear correlation line, correlation coefficient and significance level are indicated. **Figure 2D** shows correlations between suPAR versus a composite measure of insulin sensitivity (ISI composite) obtained during an oral glucose tolerance test in HIV-infected patients on HAART. ●, lipodystrophic HIV-infected patients; Δ, nonlipodystrophic HIV-infected patients. Linear correlation line, correlation coefficient and significance level are indicated.

Figure 3 shows linear correlation between fasting plasma suPAR versus overnight fasting urine suPAR corrected for urine creatinine in a sub-sample of 24 HIV-infected patients, where both scales are log transformed. The strength of the correlation is given as R^2 .

Figure 4 shows circadian concentrations of suPAR in intervals of 20 min as percentage of the mean (100%) from n=5 HIV-infected patients receiving HAART, who had suppressed HIVRNA. Upper plot shows individual diurnal profiles. Lower plot shows mean $\pm$ SD of the same patients.

Figure 5 shows the correlation of suPAR levels with overall survival. **Figure 5A** shows correlations between suPAR, age and sex. Boxplot with lines correspond to women, and dots with men. **Figure 5B** shows survival according to suPAR quartiles. Thin line represents low suPAR (N=650, suPAR below 3,620 ng/ml), medium thin line N=651, suPAR between 3,622-4,266), medium thick line, N=651, suPAR between 4,269-4,912. Thickest line represents 650 individuals with highest suPAR (> 4.915). Difference between groups significant (Log Rank, $p<0.001$).

Figure 6 shows the correlation of suPAR levels with smoking status and exercise. **Figure 6A** shows suPAR levels among men according to smoking status. Boxes with horizontal lines represent smokers, white boxes occasional-smokers and boxes with vertical lines represent non-smokers. Mean suPAR levels were significantly higher among male smokers (4.57 (sd1.45) ng/ml, N=580) compared to occasional smokers (3,64 (sd 0.86), N=56), $P<0.001$, Mann-Whitney U-test for all age groups. The suPAR level among male non-smokers was mean 3.72 (std 1.22), N=674) which was significantly lower than smokers (Mann-Whitney $p<0.001$)

but not different compared to occasional-smokers ($p=0.8$). Black line indicates median suPAR and boxes 25% quartiles above and below median. **Figure 6B** shows suPAR levels among women according to smoking status and age. Boxes with horizontal lines represent smokers; white boxes represent occasional-smokers; and boxes with vertical lines represent non-smokers. Female smokers ($N=511$) had significantly higher suPAR levels compared to occasional smokers ($N=50$) and non-smokers ($N=731$), $p<0.001$. No difference between occasional-smokers and non-smokers. Black line indicates median suPAR and boxes 25% quartiles above- and below-median. **Figure 6C** shows the relationship between suPAR and exercise in the different age-groups. Age-group 1 consisted of individuals that did not exercise regularly ($N=539$), age-group 2 of individuals did low-to moderate exercise (taking walk etc) ($N=1454$), age-group 3 of individuals exercised on a regularly basis (e.g. jogging) ($N=546$) and age-group 4 consisted of professional sportsmen ($N=17$).

Figure 7 shows the risk of developing a CVD as a competing risk model. Each line represents a suPAR quartile, with the first quartile being the lowest suPAR.

Figure 8 shows a risk analysis of diabetes according to median suPAR at inclusion. Upper continuous line (with high risk) represents individuals with suPAR above-median, while lower line (dotted line) individuals with suPAR below-median.

Figure 9 shows the relationship of suPAR levels to a number of components of metabolic syndrome in a subject, the components being one or more of: Elevated waist circumference: Men - Equal to or greater than 40 inches (102 cm) and Women - Equal to or greater than 35 inches (88 cm); Elevated triglycerides: Equal to or greater than 150 mg/dL; Reduced HDL ("good") cholesterol: Men - Less than 40 mg/dL, Women - Less than 50 mg/dL; Elevated blood pressure: Equal to or greater than 130/85 mm Hg; Elevated fasting glucose: Equal to or greater than 100 mg/dL

Figure 10 shows the relationship of suPAR levels and number of components (as in figure 9) of metabolic syndrome of a subject according to sex.

Figure 11 shows the risk of developing an IHD during follow-up. Fat line indicates individuals with suPAR above-median (4.03 ng/ml). Thin line indicates individuals with suPAR below-median. The difference was statistically significantly ($p<0.001$, log-rank test).

Figure 12 shows ROC analysis of variables significant for development of CVD in multivariate analysis.

Figure 13 shows Kaplan Meier analysis of time to cancer diagnosis. The thin upper line represents the lowest suPAR quartile, the one below (median thin line) represents the second lowest suPAR quartile, the medium thick line represents the second highest suPAR quartile and the fattest (bottom line) represents the highest suPAR quartile. $P<0.001$, log rank test.

Figure 14 shows Kaplan Meier analysis of risk of developing cancer over time. The thin upper line represents the lowest suPAR quartile, the one below (median thin line) represents the second lowest suPAR quartile, the medium thick line represents the second highest suPAR

quartile and the fattest (bottom line) represents the highest suPAR quartile.

Figure 15 shows Kaplan Meier analysis of risk of developing breast cancer over time. The thin upper line represents the lowest 1st suPAR quartile, the one below (median thin line) represents the second lowest 2nd suPAR quartile, the medium thick line represents the second highest 3rd suPAR quartile and the fattest (bottom line) represents the highest 4th suPAR quartile.

Figure 16 shows Kaplan Meier analysis of risk of developing lung cancer over time. The thin upper line represents the 1st suPAR quartile (lowest), the one below (median thin line) represents the 2nd suPAR quartile (second lowest), the medium thick line represents the 3rd suPAR quartile (second highest) and the fattest (bottom line) represents the 4th suPAR quartile (highest).

Figure 17 shows Kaplan Meier analysis of risk of developing leukemia over time. Lower continuous line (with high risk) represents individuals with suPAR above-median, while upper line (dotted line) individuals with suPAR below-median.

Figure 18 shows Kaplan Meier analysis of risk of developing cancer, excluding breast or lung cancer, over time. Lower continuous line (with high risk) represents individuals with suPAR above-median, while upper line (dotted line) individuals with suPAR below-median.

Figure 19 (Example 13) shows that full length suPAR, D₁D₂D₃(1-277), and the chemotactic active suPAR fragment, D₂D₃(88-277), were purified by immune-affinity chromatography using antibodies that specifically recognized either D₁ (antibodies R3 and R5) or D₃ (VG-1 antibody). TAG Copenhagen synthesized the amino acid sequences SRSRY and ATVYSRSRY. A microtiter plate was coated with either purified D₁D₂D₃(1-277) (30ng/ml), purified D₂D₃(88-277) (30ng/ml), SRSRY (50ng/ml), ATVYSRSRY (50ng/ml) or a mix of D₁D₂D₃(1-277) and D₂D₃(88-277) (58ng/ml). The wells were coated in triplicates and incubated with the antibodies. The antibodies was detected with a horse-radish peroxidase-conjugated goat-anti-mouse antibody, which was developed by the addition of the 3,3',5,5'-tetramethylbenzidine (TMB) substrate and stopped by the application of H₂SO₄. An ELISA reader measured the absorbance at 450 nm.

Figure 20. Purified full length suPAR, D₁D₂D₃(1-277) and purified chymotrypsin-cleaved suPAR fragment, D₂D₃(88-277) were subjected to immuno-blotting analysis using both the antibodies of the present invention. The VG-1 antibody recognized both full length suPAR, D₁D₂D₃(1-277) and the cleaved form, D₂D₃(88-277). However, the VG-2 antibody showed not to be applicable for the western blot procedure. D₁D₂D₃(1-277) was loaded in lane 2, 4 and 6, while D₂D₃(88-277) was loaded in lane 1, 3 and 5. Lane 1 and 2 were incubated with a D1 specific antibody), lane 3 and 4 with a D3 specific antibody) and lane 5, 6 and 7 were incubated with VG-1.

Figure 21: purified suPAR (4.6 µg/ml) was digested with chymotrypsin in concentrations between 1 and 300 ng/ml for 4 or 24 hours, or analyzed directly without digestion (lane 4),

separated in size by SDS-PAGE and then blotted to a PVDF membrane by western blotting. When digesting with 40 ng/ml of chymotrypsin for 4 hours only a small portion of full length suPAR was digested (lane 1). When using 40 ng/ml for 24 hours the suPAR was only partially digested (lane 2). When using 100 ng/ml for 24 hours, completely digested full length suPAR was obtained (lane 3). MagicMark was used as a protein ladder (lane 5).

Figure 22: Silver staining of D2D3(88-277) (lane 1), D1D2D3 (lane 2), a sample of albumin (lane 3) and SeeBlue (lane 4) following affinity chromatography.

Figure 23. Graphs of the upregulated cytokines. Relative concentration (The mean of each sample was divided with the mean of the mock samples) in the samples stimulated with 100 ng/ml D1D2D3₍₁₋₂₇₇₎ or D2D3₍₈₈₋₂₇₇₎.

DETAILED DESCRIPTION OF THE INVENTION

[0041] The terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0042] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either both of those included limits are also included in the invention.

[0043] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described.

[0044] It must be noted that as used herein and in the appended claims, the singular forms "a," "and" and "the" include plural references unless the context clearly dictates otherwise.

[0045] The invention is based on the unexpected observation that soluble uPAR (suPAR) is present in elevated levels in serum, plasma and urine of individuals with low-grade inflammation and in individuals with metabolic disorders. The level of suPAR is shown to vary independently of body mass index (BMI), and thus provides a more useful marker of low-grade inflammation and metabolic syndrome than currently available markers. Furthermore suPAR

levels measured in a biological fluid derived from an individual are shown to be a strong predictive marker for risk of development of diseases that can originate from low-grade inflammation and metabolic disorders, including cardiovascular disease, chronic obstructive lung disease, diabetes type 2, and developing cancer (e.g. lung cancer, leukemia, prostate and breast cancer and/or the risk of mortality within e.g., 1 year, 3 years, 5 years, 10 years or between 1-5 years, 5 years-10 years, 1-10 years).

[0046] As defined herein, "low grade inflammation" is characterized by (1) elevated levels of one or more of a pro-inflammatory cytokine (e.g., IL-6, TNF alpha, IFN gamma), serum amyloid A (SAA), orosomucoid, fibrinogen, gamma-globulin; and/or (2) increased plasma viscosity, erythrocyte sedimentation rate (ESR) and/or leukocyte count; and/or (3) decreased levels of serum albumin as compared to a reference value (see, for example, Kolb et al., 2005, *Diabetologia* 48:1038-1050); and/or (4) increased acute phase response as measured for example by hypersensitive C-reactive protein (hs-CRP) (e.g. an hs-CRP level of 1.5-3.5 times, preferably about twice, the normal level); all in the absence of observable inflammation. In a particular embodiment, low grade inflammation is characterized by elevation of one or more of the mentioned markers, in the absence of observable inflammation. In particular, low-grade inflammation can be characterised as an hs-CRP level of higher than 1mg/l (particularly more than 3mg/l), for example as measured using the particle enhanced immunoturbidimetry assay (Roche Diagnostics) described below.

[0047] As defined herein, "metabolic syndrome" is characterized by a combination of high blood pressure, insulin resistance with increased glucose production and decreased glucose utilization, central (abdominal) obesity, dyslipidemia namely elevated triglycerides, LDL-cholesterol and total cholesterol, and reduced high-density lipoprotein cholesterol, and "low grade inflammation" as stated above. The broadly accepted criteria for diagnosis of metabolic syndrome, provided by the National Cholesterol Education Program (NCEP) Adult Treatment Panel III (ATP III) with minor modifications, are as follows:

Elevated waist circumference: Men - Equal to or greater than 40 inches (102 cm) and Women

- Equal to or greater than 35 inches (88 cm);

Elevated triglycerides: Equal to or greater than 150 mg/dL;

Reduced HDL ("good") cholesterol: Men - Less than 40 mg/dL, Women - Less than 50 mg/dL;

Elevated blood pressure: Equal to or greater than 130/85 mm Hg;

Elevated fasting glucose: Equal to or greater than 100 mg/dL. Accordingly, in a particular embodiment of the invention, a subject is diagnosed with metabolic syndrome if the subject exhibits 3 or more of the above listed 5 components.

In a more particular embodiment, metabolic syndrome is defined according to the International Diabetes Federation definition as central obesity (waist men/women > 94/80 cm) together with at least two of the following: elevated serum triglycerides >1.7 mmol/l (150 mg/dl), reduced serum high-density lipoprotein cholesterol (<1.03 mmol/l (40 mg/dl), elevated plasma glucose

(>5.6 mmol/l (100 mg/dl) or elevated BP (systolic BP >130 mm Hg or diastolic BP > 85 mm Hg).

[0048] Decreasing the suPAR level may be achieved by decreasing the rate of suPAR production or by decreasing the stability of suPAR produced, e.g. by clearing it from the body with an anti-suPAR anti-body.

1. suPAR and/or its cleavage products for use as a marker:

[0049] The protein suPAR (NCBI Accession no. AAK31795 and isoforms of the receptor, NP_002650, Q03405, NP_002650, NP_001005376) is the soluble portion of Urokinase-type Plasminogen Activator Receptor (uPAR), which is released by cleavage of the GPI anchor of membrane-bound uPAR. suPAR is a family of glycosylated proteins consisting of full length suPAR (277 amino acids (1-277)) and suPAR fragments D1 (1-83), and D2D3 (84-277) generated by urokinase cleavage or human airway trypsin-like protease, D1 (1-87) and D2D3 (88-277) generated by MMP cleavage, D1 (1-89) and D2D3 (90-277) also generated by urokinase cleavage or human airway trypsin-like protease, D1 (1-91) and D2D3 (92-277) generated by cleavage by plasmin. Continuous and discontinuous epitopes present in the protein suPAR and its cleavage products may be used to monitor their presence and abundance in a biological fluid by immunodetection with mono- or poly-clonal antibodies. Thus, antibodies directed to accessible epitopes common to suPAR and its cleavage products (e.g. D2D3) can be used to detect both suPAR and its cleavage products in a biological fluid.

[0050] The use of suPAR as a marker has the advantage that it is more stable than other markers of the immune response, e.g. TNF- α and IL-6. For example, suPAR levels in healthy individuals are known to be stable throughout the day (Sier CF, et al., 1999, Lab Invest 79:717-722). Similarly, circadian changes in plasma concentration of suPAR in HIV-infected patients on stable HAART are shown to be very limited (Example 1). Thus, suPAR measurements based on a biological fluid derived from a subject will be valid, independent of whether the subject is fasting or not, and largely independent of the time schedule for sampling. This is not the case for alternative markers such as TNF- α and IL-6, which show substantial circadian fluctuations. A flexible sampling schedule and the stability of suPAR molecules in stored samples of biological fluid is a great advantage for its use as a marker in the clinical setting.

2. Biological samples suitable for detection of suPAR or its cleavage products as a marker

[0051] suPAR, or its cleavage products (e.g., D2D3), can be used as a marker for diagnostic, prognostic and predictive purposes by measuring the level of suPAR or its cleavage products in a biological fluid derived from a mammalian subject, particularly, a human subject or patient, as illustrated in the examples herein. suPAR and its cleavage products are present in all biological fluids derived from a mammalian subject, particularly, a human subject, including cerebrospinal

fluid, plasma, serum, blood, urine, semen, saliva and sputum.

[0052] Where the biological sample is urine, the measurements may be based on the urine suPAR/creatinine value from a subject, since this value is known to be highly correlated to the concentration of suPAR in a serum (plasma) sample derived from the same subject. Similarly the urine suPAR/creatinine values of an HIV patient on HAART is highly correlated to the concentration of suPAR in serum (plasma) samples derived from the same subject (Example 1). Thus, urine samples may also be employed for the measurement of suPAR, or its cleavage products, in a subject (including an HIV patient on HAART), where the measured level in urine is normalized for protein content (e.g. using creatinine). These normalized values may be employed as a marker for the diagnostic, prognostic or predictive purposes of the present invention.

3. Detection and quantitation of suPAR and its cleavage products

[0053] Accurate methods for measuring the level of suPAR and/or its cleavage products in a biological fluid derived from a subject, include immunodetection methods e.g. Enzyme-Linked ImmunoSorbent Assay (ELISA), which are particularly suitable as such methods are relatively cheap and simple to perform in the clinical setting. ELISAs can be adapted to analyze both small and large numbers of samples, and include both an ELISA plate format with wells coated with suPAR specific antibodies, or adapted to a dipstick format incorporating all components of the ELISA assay. Additionally, suPAR levels can be measured by proteomic approaches such as western blot, Luminex, MALDI-TOF, HPLC and automated immune analyzer platforms such as Bayer Centaur, Abbott Architect, Abbott AxSym, Roche COBAS and the Axis Shield Afinion.

4. suPAR as a marker for diagnosis of low-grade inflammation and metabolic syndrome in A) HIV-infected patients on HAART treatment, and B) apparently healthy subjects

[0054] Low-grade inflammation, as defined above, is a sub-clinical state in an apparently healthy subject, who exhibits no observable inflammation, yet in whom immune activation is detectable, although at a level far below that seen in acute infections. hsCRP (C-reactive protein), measured using available sensitive tests, is currently used as a marker for low-grade inflammation in a subject.

[0055] Markers for the diagnosis of low-grade inflammation in a subject have particular importance, since this sub-clinical condition seems to be linked to subsequent development of low-grade inflammation-related diseases including type-2 diabetes and CVD. Diagnosis of low-grade inflammation in a population would identify those members whose risk of developing low-grade inflammation related-diseases should be monitored further.

1. **A)** HIV-infected patients on HAART, who display a cluster of dysmetabolic features, have elevated levels of suPAR compared to treatment-matched, non-lipodystrophic HIV-

infected patients (Example 1). Levels of suPAR in these HIV patient groups was positively correlated with leukocyte, lymphocyte, and monocyte numbers, all of which are blood cells of particular importance for the immune activation and inflammatory status of a patient. These data, and the observed strong positive-correlation between suPAR and TNF- α , identify suPAR as a novel strong diagnostic marker of low-grade inflammation in HIV-infected patients.

suPAR levels in these HIV patient groups was also strongly correlated with respectively, fat distribution, insulin sensitivity and lipidemia, identifying suPAR as a marker of metabolic syndrome in these patients. Use of SuPAR as a marker is also shown to provide a stronger prediction of metabolic syndrome than the alternative markers, TNF- α and IL-6.

2. **B)** White blood cell (WBC, i.e. leukocyte) count is a marker for immune activation and low-grade inflammation in the general population. In the general human population, comprising the Monica10 cohort, the suPAR level and the WBC count was highly correlated, and shown to be independent of age and gender (Example 4, Table 9; Example 5). These data establish suPAR as a novel marker of low-grade inflammation in the general population that includes apparently healthy individuals. Furthermore suPAR provides a valuable marker for the purpose of staging the progress of low-grade inflammation in a subject. suPAR has application as a marker to monitor the risk of onset of low-grade inflammation-related diseases, since suPAR is shown to be a marker for the diseases Type 2 Diabetes and CVD. Cancer is an additional low-grade inflammation-related disease for which suPAR is a useful marker. This is consistent with the observation that leucocytes (white blood cell count), are a well established marker of inflammation, linked to cancer development.

[0056] The metabolic syndrome is defined by the following 5 components: waist circumference, fasting glucose level, low HDL-cholesterol, high plasma triglyceride, increased diastolic and/or systolic blood pressure; according to the NCEP ATPIII 2001 definition with the 2004 glucose modification;(Grundy SM et al., 2005, Circulation 112:2735-2752).

[0057] In the general human population, comprising the Monica10 cohort, suPAR levels were positively correlated with the number of components of metabolic syndrome detected in an individual (Example 6). This was the case for both men and women in the tested population, and the association was independent of the subject's age. These data establish suPAR as a novel marker of metabolic syndrome in the general population that includes apparently healthy individuals.

4. suPAR as a marker for predicting the risk for an apparently healthy subject of developing Diabetes type 2.

[0058] suPAR is shown to be an independent predictor of the risk of developing diabetes

(Example 5). In both univariate and multivariate analysis, high suPAR, age (older), sex (being male) and having the metabolic syndrome were significantly associated with an increased risk of developing diabetes. Independent of age, sex and the metabolic syndrome, a 1 ng/mL increase in suPAR was associated with a 28% increased risk of developing diabetes. The use of suPAR as a marker for predicting the risk of developing diabetes will allow individuals at risk to improve their health chances by initiating a prophylactic course of action at a sufficiently early time point.

[0059] suPAR was also shown to correlate significantly with an increasing fasting glucose (Table 9). Example 5 demonstrates that suPAR strongly predicts type 2 diabetes. From these observations, it is deduced that suPAR most likely is a strong marker of insulin resistance, which is an underlying cause of the metabolic syndrome and type 2 diabetes mellitus (Grundy SM et al 2005, *Circulation* 112:2735-2752).

[0060] As prediabetes, in the form of e.g. impaired glucose tolerance, precedes type 2 diabetes and is more prevalent in individuals exhibiting the metabolic syndrome than individuals not having this syndrome, it is most likely that suPAR is both a strong marker and strong predictor of prediabetes and impaired glucose tolerance in the general population. In support of suPAR being a strong marker of insulin resistance is that this particular association has been demonstrated in HIV-infected individuals on a stable HAART regime using gold standard measurement of insulin sensitivity (Example 1, Figure 2A). Also a more simple measure of insulin sensitivity namely HOMA-IR was associated with suPAR (Figure 2B). Moreover, the observed correlation between plasma suPAR level and the 2-hour plasma glucose during an oral glucose tolerance test in HIV-infected individuals on a stable HAART regime (Figure 2C) confirms suPAR as a strong marker of impaired glucose tolerance.

5. suPAR as a marker for predicting the risk for an apparently healthy subjects of developing Cardiovascular Disease (CVD) and Ischemic Heart Disease (IHD).

[0061] suPAR is shown to be a strong, independent risk marker of development of a CVD event in apparently healthy individuals, that is not, or only weakly, correlated to classical predictors of CVD development (Example 4). Independent of sex, age, smoking status and metabolic syndrome, a 1 ng increase in suPAR was associated with a 15.4% increased risk of developing a CVD.

[0062] suPAR level is a significant predictor of the risk of developing an IHD in individuals who have not previously experienced an Acute Myocardial Infarct (Example 7). SuPAR predicted the risk of IHD independently of other classical markers of IHD. In an age-adjusted analysis, a 1 ng/ml increase in suPAR was associated with an increased risk of IHD of 21%.

[0063] suPAR as a marker for prediction of the risk of developing both CVC and IHD also gives an individual at risk the chance to initiate a prophylactic course of action at a time point that is sufficiently early as to lower their risk.

6. suPAR as a marker for predicting the risk for an apparently healthy subjects of developing Cancer.

[0064] suPAR is shown to be a strong, independent risk marker of development of cancer in apparently healthy individuals (Example 10). Independent of sex and age, a 1 ng increase in suPAR is associated with a hazard ratio of 1.34 increased risk of developing overall cancer: Significant associations were found within the following cancer subgroups breast cancer, leukemia, prostate cancer and lung cancer.

[0065] suPAR as a marker for prediction of the risk of developing cancer also gives an individual at risk the chance to initiate a prophylactic course of action at a time point that is sufficiently early as to lower their risk. The effect of the risk intervention can be monitored using suPAR levels. An effective intervention will lower the suPAR level.

7. suPAR as a predictor of survival and efficacy of life style intervention

[0066] suPAR levels in apparently healthy individuals are demonstrated to be strongly associated with mortality, independent of other markers (Example 3). In multivariate analysis, suPAR was the second strongest marker of mortality, after age.

[0067] The life style of an apparently healthy individual is shown to impact their suPAR levels significantly. In the case of smoking habit, smokers significantly impact their suPAR level and their survival prospects. Similarly, exercise habit has a significant impact on suPAR levels in an individual, where the level of exercise is associated with lower suPAR levels. It is thus apparent that regular suPAR measurements on a biological fluid derived from a mammalian subject, particularly, a human subject, would allow the physician to guide his subject with respect to his life expectancy based on current life style, and the efficacy of life-style changes taken in the course of prophylactic course of action, e.g. a physical exercise program may be proposed by a physician. Similarly, regular suPAR measurements would allow the physician to monitor the efficacy of medication; to monitor patient compliance in taking the prescribed medication, and to monitor the efficacy of a course of treatment to prevent or treat Metabolic Syndrome related diseases, including IHD, CVD and Type 2 Diabetes.

8. suPAR as a Marker for Monitoring Progression of Treatment of Metabolic Syndrome and/or Low-Grade Inflammation

[0068] suPAR levels may also be used to monitor treatment of metabolic syndrome and/or low-grade inflammatory disease. A suPAR measurement may be made before treatment begins. During treatment, suPAR measurements may be taken at various time points. A

decrease of at least about 10% and/or 1 ng/ml would be indicative of the effectiveness of a treatment protocol. suPAR levels may further be measured at various time points post-treatment.

9. Treatment of Metabolic Syndrome and/or Low-Grade Inflammation

[0069] Substances that decrease the level of suPAR in a mammalian subject may be used to treat metabolic syndrome and/or low-grade inflammation. In a particular aspect, the substance may be an anti suPAR monoclonal or polyclonal antibody. Such antibodies may be obtained using methods known in the art. Various hosts may be used and include but are not limited to goats, rabbits, rats, mice, humans, and others. These hosts may be immunized by injection with suPAR or peptide fragments having a minimum length of 5-10 amino acids with immunogenic properties (see e.g., WO90/12091). Various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, Pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum* are especially preferable in humans. The following steps may be used to obtain said antibody: (a) optionally conjugating uPAR or an immunogenic part thereof to a carrier protein; (b) immunizing a host animal with said polypeptide or polypeptide-carrier protein conjugate of step (a) with an adjuvant and (c) obtaining antibody from said immunized host animal. Preferably, any antibody that is used binds to full length suPAR, not just to fragments thereof, such as the D2D3 fragment.

[0070] Monoclonal antibodies to the said receptor or receptor peptides may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. See, e.g., Kohler, et al., 1975, Nature 256: 495-497; Kozbor et al., 1985, J. Immunol. Methods 81: 31-42; Cote et al., 1983, Proc. Natl. Acad. Sci. USA 80: 2026-2030; Cole et al., 1984, Mol. Cell Biol. 62: 109-120. Specifically, the method comprises the following steps: (a) immunizing an animal with an immunogenic receptor peptide; (b) isolating antibody producing cells from the animal; (c) fusing the antibody producing cells with immortalized cells in culture to form monoclonal antibody-producing hybridoma cells; (d) culturing the hybridoma cells; and (e) isolating from the culture monoclonal antibodies which bind to said polypeptide.

[0071] Antigenic specificity is conferred by variable domains and is independent of the constant domains, as is known from experiments involving the bacterial expression of antibody fragments, all containing one or more variable domains. These molecules include Fab-like molecules (Better et al (1988) Science 240, 1041); Fv molecules (Skerra et al (1988) Science 240, 1038); single-chain Fv (ScFv) molecules where the V_H and V_L partner domains are linked via a flexible oligopeptide (Bird et al (1988) Science 242, 423; Huston et al (1988) Proc. Natl. Acad. Sci. USA 85, 5879) and single domain antibodies (dAbs) comprising isolated V domains (Ward et al (1989) Nature 341, 544). A general review of the techniques involved in the

synthesis of antibody fragments which retain their specific binding sites is to be found in Winter & Milstein (1991) Nature 349, 293-299. By "ScFv molecules" we mean molecules wherein the V_H and V_L partner domains are linked via a flexible oligopeptide. These molecules may be used in the present invention.

[0072] Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between the polypeptide(s) and its specific antibody.

[0073] Other substances that may be used to obtain agents that decrease suPAR levels include but are not limited to a D2D3 cleavage product and/or another suPAR fragment. The substance may also include but is not limited to a suPAR analog or derivative capable of binding uPA. In a specific aspect, such substances include but are not limited to 1) uPAR having an amino acid sequence containing a substitution, a deletion, a truncation and/or an addition, 2) chemically modified uPAR such as different degrees of glycosylation and 3) synthetic peptides containing part of the uPAR sequence. The suPAR analogues or fragments are large enough to generate antibodies that recognize human suPAR, uPAR and/or suPAR/uPAR fragments. Such analogues will generally have at least 60%, preferably at least 70%, 80%, 90%, 95% or 99% sequence identity with the most similar region of human suPAR.

[0074] Formulations of the substances into pharmaceutical compositions is well known in the art, and is further described in Gennaro (ed.), 2000, Remington: The Science and Practice of Pharmacy, 20th ed., Lippincott, Williams & Wilkins (2000); and Ansel et al., 1999, Pharmaceutical Dosage Forms and Drug Delivery Systems, 7th ed., Lippincott Williams & Wilkins Publishers.

[0075] Such a composition typically contains from about 0.1 to 90% by weight (such as about 1 to 20% or about 1 to 10%) of the polypeptide or antibody of the invention in a pharmaceutically accepted carrier.

[0076] Various liquid and powder formulations can be prepared by conventional methods for inhalation into the lungs of the mammal to be treated.

[0077] Injectable formulations of the compositions can contain various carriers such as vegetable oils, dimethylacetamide, dimethylformamide, ethyl lactate, ethyl carbonate, isopropyl myristate, ethanol, polyols (glycerol, propylene glycol, liquid polyethylene glycol, and the like). For intravenous injections, water soluble versions of the compounds can be administered by the drip method, whereby a pharmaceutical formulation containing the antifungal agent and a physiologically acceptable excipient is infused. Physiologically acceptable excipients can include, for example, 5% dextrose, 0.9% saline, Ringer's solution or other suitable excipients. Intramuscular preparations, e.g., a sterile formulation of a suitable soluble salt form of the compounds, can be dissolved and administered in a pharmaceutical excipient such as Water-

for-Injection, 0.9% saline, or 5% glucose solution. A suitable insoluble form of the compound can be prepared and administered as a suspension in an aqueous base or a pharmaceutically acceptable oil base, such as an ester of a long chain fatty acid (e.g., ethyl oleate).

[0078] The optimal percentage of the therapeutic agent in each pharmaceutical formulation varies according to the formulation itself and the therapeutic effect desired in the specific pathologies and correlated therapeutic regimens.

[0079] Conventional methods, known to those of ordinary skill in the art of medicine, can be used to administer compositions to the patient. These include but are not limited to subcutaneous, intrapulmonary, transmucosal, intraperitoneal, intrauterine, sublingual, intrathecal, or intramuscular routes) by using standard methods. In addition, the pharmaceutical formulations can be administered to the patient via injectable depot routes of administration such as by using 1-, 3-, or 6-month depot injectable or biodegradable materials and methods.

[0080] Regardless of the route of administration, the substance of the present invention is typically administered at a daily dosage of about 0.01 mg to about 30 mg/kg of body weight of the patient (e.g., 1 mg/kg to 5 mg/kg). The pharmaceutical formulation can be administered in multiple doses per day, if desired, to achieve the total desired daily dose.

[0081] Conventional methods, known to those of ordinary skill in the art of medicine, can be used to administer the pharmaceutical formulation(s) to the patient. The pharmaceutical compositions can be administered alone, or in combination with other therapeutic agents or interventions. Specifically, the compositions may further comprise a plurality of agents.

10. suPAR based tools for predicting Metabolic Syndrome-related diseases, low-grade inflammatory-related diseases, cancer and/or mortality within 10 years.

[0082] A combination of factors, one of which being suPAR, are predictive of the risk of developing CVD (including IHD and AMI), Diabetes type 2, cancer and/or mortality within 10 years. For example, stepwise backward logistic regression identifies those factors that have the highest statistical significance in determining the outcome of a large human population, and their risk of developing any one of these diseases. The combination of prediction factors, or criteria, comprises a model that may be employed in the form of an algorithm for predicting the individual health risks of a subject, as set out in Examples 8 and 9. Other algorithms can be implemented for calculating the risk of mortality or development of the diseases. For example, in an alternative embodiment, a risk analysis algorithm can include suPAR and age, or suPAR, age and gender, or suPAR, age and smoking. Using these factors as the input values of the algorithm, the output score may then be the different risk percentages over the next 5 years, 10 years or 5 years-10 years for the different diseases. Each disease may thus obtain a specific risk percentage, e.g. such as:

Diabetes type 2	risk 22%
CVD	risk 19%
Cancer	risk 36%
Mortality	risk 33%

[0083] To further estimate the specific risk for a person, the following parameters could e.g. be measured and used as input values in a further calculation of risk:

For Diabetes type 2	fasting blood glucose
For CVD	blood pressure, HDL-cholesterol, LDL-cholesterol, total cholesterol, triglycerides
For cancer	smoking, age
For mortality	NT-proBNP concentrations

[0084] N-terminal fragment of brain-type natriuretic peptide In the context of diagnosing the risk of developing cancer, the invention may have particular utility in relation to women who belong to a family with hereditary breast cancer and/or who have been shown to have a genetic marker associated with cancer, such as the BRCA1 and BRCA2 mutations.

[0085] The described combination of prediction factors may be used in the manufacture of a computer readable medium for entering determined prediction factor measurements and means for calculating and providing the risk of developing a Metabolic Syndrome-related diseases and/or low-grade inflammatory-related diseases e.g. CVD (including IHD and AMI), Diabetes type 2, developing cancer and/or mortality within one year, three years, five years, 10 years or 1-5 years, 1-10 years, five years-ten years.

[0086] In order to reach a prediction for a given patient, the physician performs a number of measurements, including the measurement of suPAR in a sample, e.g. biological fluid, derived from the patient, and enters the determined values into an computer logic for receiving test information comprising the level or relative amount of a protein that is suPAR and/or a D2D3 cleavage products thereof in a biological fluid derived from a mammalian subject, particularly, a human subject, and one or more further prediction factor parameters obtained from said mammalian subject, particularly, a human subject, executing computer logic for analysis of the level or relative amount of said protein in said biological fluid, and said one or more parameter, wherein the computer logic analyses the entered determined values and computes the risk of developing a Metabolic Syndrome-related disease, low-grade inflammatory-related disease, developing cancer and/or mortality within one year, three years, five years, 10 years or one-five years, one-ten years, five years-ten years.

[0087] Use of a computer comprising said computer readable medium would allow the

physician to predict the risk of developing CVD (including IHD and AMI) and/or Diabetes type 2 in a mammalian subject, particularly, a human subject; as well as monitor the efficacy of medication, and patient compliance in taking the prescribed medication, in the course of treatment to prevent or treat Metabolic syndrome-related diseases, including CVD (including IHD and AMI), Diabetes type 2.

[0088] The use of a computer comprising said computer readable medium may also allow the physician to predict the risk of developing cancer and/or mortality within one year, three years, five years, 10 years or one-five years, one-ten years, five years-ten years in a mammalian, particularly, a human subject.

[0089] One such computer that may be used is a client computer e.g. in a doctor's surgery for providing access to a risk computation system for determining the risk and/or change in risk of developing a metabolic syndrome-related disease, low-grade inflammation-related disease, cancer and/or mortality within e.g., one year, five years, ten years, 1-5 years, 1-10 years or five years-ten years comprising:

1. (a) a computer-readable data storage medium configured to store a data storage material encoded with computer-readable data, wherein said data comprises measurements derived from one of more samples from a mammalian subject, wherein at least one measurement is an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR;
2. (b) a storage medium having stored therein instructions for processing said computer-readable data;
3. (c) a central-processing unit responsive to said instructions coupled to said storage medium and to said computer-readable data storage medium for processing said computer-machine readable data into risk measurement and treatment progression values; and
4. (d) an output device coupled to said central-processing unit of (c) for outputting said risk and/or change in risk values.

[0090] Alternatively, the client computer comprises: an input device for receiving one or more measurements derived from a mammalian subject wherein at least one measurement is an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject; a communication interface for communicating the received one or more measurements to a server computer configured to analyse said one or more measurements of step (a); and compute a risk value; and a communications interface for receiving, from the server computer a computed risk value.

[0091] The output device may include any device or circuitry suitable for outputting data. Examples of such output devices include devices for presenting data in a user-readable way,

such as a display, a printer, and/or the like. Alternatively or additionally, the output device may include a communications interface for communicating the data to one or more other devices, e.g., computers. For example, the communications interface may include a serial port, a USB port, a parallel port, a short-range wireless communications interface, e.g., an infrared (e.g. IrDa) port, a Bluetooth transceiver, or the like. Further examples of interface circuits include a network card, a DSL modem, a gateway computer, a wireless communications circuit for cellular telecommunications, or the like. Yet further examples of output devices include devices or circuitry for writing the data on a computer-readable medium, such as a disk drive, a memory card interface, and/or the like.

[0092] For example, the data processing system to process computer-readable data may comprise a server computer or other data processing system having access to the communications interface. In some embodiments, the functionality of the risk computation system described herein may be distributed among a plurality of computers, e.g., computers connected via a computer network, e.g. a local area network, a wide area network, an Internet, or the like.

[0093] The input device may include any device or circuitry suitable for receiving data, such as a user interface, e.g. a keyboard, a graphical user interface, a pointing device, and/or the like. Other examples of input devices include communications interfaces as described herein and devices or circuitry for reading data from a storage medium, such as a disk drive, a memory card reader, and/or the like.

EXAMPLES

Example 1: SuPAR is a marker of low-grade inflammation and metabolic syndrome (dysmetabolism) in HIV infected patients

[0094] HIV infected individuals on HAART have recently been found to develop dysmetabolism, which is a form of metabolic syndrome similar to that seen in uninfected individuals, but within a shorter timeframe. The following study on HIV patients demonstrates the utility of suPAR as a marker for metabolic syndrome, and shows this marker to have a stronger predictive power than the markers, IL-6 and TNF- α for metabolic syndrome in this patient group.

[0095] Patient groups. A cohort of 36 HIV-infected male patients on HAART had estimated overnight fasting suPAR (Andersen et al., 2003, Metabolism 52:1343-1353). This cohort comprises 18 patients with lipodystrophy (LIPO), a marker of the metabolic syndrome in HIV patients, and 18 without any manifestations of metabolic syndrome (NONLIPO) as judged by a clinical investigation and a questionnaire, both of which have been described by (Andersen et al., 2003, Metabolism 52:1343-1353). All patients had received HAART for at least 12 months

prior to examination, had a stable weight, and were free of any acute infection episode within three months of examination. All participants were Caucasians except two NONLIPO (Black Africans). Subjects gave their written informed consent and the protocol was approved by the Ethical Committee in Copenhagen, Denmark, and performed in accordance with the Helsinki Declaration II.

[0096] Protocol: Data obtained from hyperinsulinemic ($40 \text{ mU} \cdot \text{m}^{-2} \cdot \text{min}^{-1}$) euglycemic (5.0 mM) clamps including glucose tracer technique and indirect calorimetry provided estimates of insulin sensitivity (R_d) and nonoxidative glucose disposal rate (NOGM) from the 18 LIPO and the 18 NONLIPO patients, as described by (Haugaard et al. 2005, *Eur J. Endocrinol.* 152:103-112). Five HIV-infected male patients on HAART had determined circadian plasma suPAR by collecting venous blood continuously during 20 min periods (6 ml/h) for 24 hours starting at 0800 in the morning after an overnight fast (Constant blood withdrawal pump type 3003, SWEMED LAB, International AB, V. Frölunda, Sweden). The samples were drawn from a catheter (inserted in the left antecubital vein) into unpreserved test tubes, and allowed to coagulate for 1 hour at 25°C , and thereafter centrifuged at 4°C . Serum was stored at -80°C for later analysis. These 5 patients had meals served at 0900, 1230, and 1800. A snack was provided at 1500 and at 2100. They were not allowed to sleep before bedtime at 2230. They were allowed to walk around within the hospital area, however, exercise, as such, was prohibited.

[0097] To test whether plasma suPAR in HIV-infected patients on stable HAART may correlate with urine suPAR, as has been demonstrated previously in HIV negative individuals (Sier et al., 1999, *Lab Invest* 79:717-722), a sub-sample of 24 of the 36 patients had given overnight fasting urine. The effect of differences in dilution of the urine on suPAR levels was corrected with the amount of creatinine, as described previously (Sier et al., 1999, *Lab Invest* 79:717-722). Urine creatinine was measured as described (Mustjoki et al., 2000, *Cancer Res* 60:7126-7132).

[0098] suPAR measurement: suPAR levels were determined by ELISA assay as follows: Nunc maxisorp ELISA-plates (Nunc, Roskilde, Denmark) were coated overnight at 4°C with a monoclonal rat anti-suPAR antibody (VG-1, ViroGates A/S, Copenhagen, Denmark, $3 \text{ } \mu\text{g/ml}$, $100 \text{ } \mu\text{l/well}$). Plates were blocked with PBS buffer + 1% BSA and 0.1% Tween20, 1 hour at RT, and washed 3 times with PBS buffer containing 0.1% Tween20. $85 \text{ } \mu\text{l}$ dilution buffer (100 mm phosphate, 97.5 mm NaCl, 10 gL^{-1} bovine serum albumin (BSA, Fraction V, Boehringer Mannheim, Penzberg, Germany), 50 U mL^{-1} heparin sodium salt (Sigma Chemical Co., St. Louis, MO), 0.1% (v/v) Tween-20, pH 7.4) containing $1.5 \text{ } \mu\text{g/ml}$ mouse-anti suPAR-HRP labeled antibody (VG-2-HRP, ViroGates) and $15 \text{ } \mu\text{l}$ plasma (or urine) sample was added in duplicates to the ELISA plate. After 1 hour of incubation at 37°C , plates were washed 10 times with PBS buffer + 0.1% Tween20 and $100 \text{ } \mu\text{l/well}$ HRP substrate added (Substrate reagent pack, R&D Systems Minneapolis, Minnesota). The color reaction was stopped after 30 min using $50 \text{ } \mu\text{l} / \text{well}$ 1 M H_2SO_4 and measured at 450 nm . The intra-assay variation was 9.6% and the inter-assay variation 12.4%. Fasting plasma suPAR was measured in all 36 HIV-infected

patients on HAART.

[0099] Other Assays: Measurement of plasma glucose, insulin, C-peptide, free fatty acids, cholesterol, triglyceride including immunological and hematological parameters were performed as described by (Andersen et al., 2003, *Metabolism* 52:1343-1353; Haugaard et al., 2005, *Eur J Endocrinol* 152:103-112; Lihn et al., 2003, *Am J Physiol Endocrinol Metab* 285:E1072-E1080). Body composition was estimated by dual energy x-ray absorptiometry (DEXA) scanning (XR-36; Norland Medical System, Fort Atkinson, WI) and regional fat distribution was determined as previously described (Andersen et al., 2003, *Metabolism* 52:1343-1353).

[0100] Calculations: Total glucose disposal rate (Rd) was calculated using Steele's non-steady-state equations adapted for labeled glucose infusates (Hother-Nielsen et al., 1996, *Metabolism* 45:82-91). Distribution volume of glucose was taken as 200-ml/kg body wt and pool fraction as 0.65. Rate of glucose oxidation (GOX) was calculated from validated equations (Frayn, 1983, *J Appl Physiol* 55:628-634). NOGM was calculated as the difference between Rd and GOX.

[0101] Statistical analysis: Data are presented as means $\pm$ SEM, and as medians and interquartile ranges when distributions were skewed. Analysis of variance (ANOVA) was performed to compare distribution of data between LIPO and NONLIPO. As age was slightly increased in LIPO compared with NONLIPO age was adjusted for by calculation of partial correlations. If data distribution was skewed, data were log transformed before applying ANOVA. Pearson correlation coefficient (r) and Spearman correlation coefficient (p) were applied to estimate associations between variables, when appropriate. A linear stepwise multiple regression model was used to test robustness of suPAR as an explanatory variable for dysmetabolic parameters. Calculations were performed by SPSS (SPSS ver. 12.0; SPSS Inc., Chicago, Illinois, USA). Two-sided P values less than 0.05 were defined as statistically significant. A trend was noted if P was greater or equal to 0.05 but less than 0.2.

Results:

A. suPAR levels are tightly correlated with low-grade inflammation and metabolic syndrome in HIV patients:

[0102] suPAR was measurable in all 36 patients, with a median plasma suPAR level of 2.52 ng/ml (range 0.38-19.25). As suPAR showed a significant positive correlation with age ($r=0.60$, $P<0.001$), comparisons between study groups and correlation analysis were corrected for age (Table 1 and Table 2). The median plasma suPAR was increased 2-fold in LIPO compared to NONLIPO ($P<0.005$), although the difference attenuated after adjustment for age (Figure 1, Table 1).

Table 1. Anthropometry, lipidemia, insulin sensitivity, and immunology of study groups

	LIPO	NONLIPO	P
Number	18	18	
Age (yr)	50 (2)	43 (2)	<0.05
BMI (kg/m ²)	24.7 (0.6)	22.6 (0.8)	<0.05
Total fat mass (kg)	16 (1)	13 (2)	0.13
Total lean mass (kg)	61 (2)	54 (2)	<0.05 ^a
Limb fat (%)	36 (1)	46 (2)	<0.0001 ^c
Rd _{clamp} (mg min ⁻¹ kg _{FFM} ⁻¹)	5.9 (0.5)	8.9 (0.6)	<0.001 ^b
NOGM _{clamp} (mg min ⁻¹ kg _{FFM} ⁻¹)	2.5 (0.4)	4.9 (0.5)	<0.005 ^a
GOX _{clamp} (mg min ⁻¹ kg _{FFM} ⁻¹)	3.3 (0.2)	4.0 (0.3)	0.056 ^a
fP-insulin (mmol/L)	77 (11)	32 (4)	<0.001 ^b
fP-glucose (mmol/L)	4.9 (0.1)	4.8 (0.1)	ns
fP-total cholesterol (mmol/L)	6.2 (0.4)	4.8 (0.2)	<0.01 ^a
fP-nonHDL cholesterol (mmol/L)	5.2 (0.4)	3.8 (0.2)	<0.01 ^a
fP-TG (log10 mmol/L)	0.47 (0.08)	0.30 (0.06)	0.09
fP-suPAR (log10 ng/mL)	0.56 (0.07)	0.20 (0.09)	<0.005 [#]
fP-TNF-α (log10 pg/mL)	0.42 (0.06)	0.32 (0.07)	ns
fP-IL-6 (log 10 pg/mL)	0.54 (0.06)	0.46 (0.06)	ns
Hemoglobin (mmol/L)	8.8 (0.2)	8.6 (0.2)	ns
Leukocytes (10 ⁹ /L)	7.2 (6.0-9.0)	5.6 (4.6-6.5)	<0.05
Neutrophils (10 ⁹ /L)	3.4 (2.5-4.8)	3.1 (2.1-4.3)	ns
Lymphocytes (10 ⁹ /L)	2.1 (0.2)	1.6 (0.1)	<0.05 ^a
Monocytes (10 ⁹ /L)	0.57 (0.06)	0.43 (0.06)	0.14
CD4 (cells/μL)	427 (45)	352 (46)	ns
HIV-RNA (copies/mL)	<20 (<20, 70)	<20 (<20, <20)	ns
CD4 nadir (cells/μL)	130 (25)	123 (32)	ns
HIV-RNA peak (log copies/mL)	4.85 (0.33)	5.05 (0.15)	ns
Duration of HIV-infection (mo)	99 (14)	72 (11)	0.16

	LIPO	NONLIPO	P
Duration of NRTI therapy (mo)	47 (7)	42 (6)	ns
Duration of PI therapy (mo)	32 (4)	25 (4)	0.16
CDC classification: number A/B/C	6/6/6	4/5/9	ns

Mean (SEM) or median (interquartile range); ^a, P<0.05; ^b, P<0.01; ^c, P<0.001, following adjustment for age (yr); [#], P=0.06 after adjustment for age; Limb fat (%), (limb fat mass + trunk fat mass)/trunk fat mass; Rd is glucose disposal as estimated by a hyperinsulinemic euglycemic clamp using glucose tracer technique, Rd minus GOX (glucose oxidation rate) is non-oxidative glucose metabolism rate (NOGM). fP, fasting plasma; CDC classification: categorized according to the Centers for Disease Control and Prevention (CDC) system of classification clinical stages A, B, and C, where C indicates AIDS diagnosis.

[0103] LIPO patients were slightly older on average than NONLIPO and had slightly higher BMI, attributable to an increased lean body mass, whereas total fat mass did not differ significantly between study groups (Table 1). The percentage of limb fat was significantly reduced in LIPO patients compared to NONLIPO, consistent with lipodystrophy in LIPO patients. Several dysmetabolic features were apparent in LIPO patients, despite their fasting glucose being normal. LIPO patients displayed lower insulin stimulated glucose disposal both in the oxidative and non-oxidative pathway compared to NONLIPO. Fasting insulin, non-HDL-cholesterol, total leukocyte, lymphocyte and monocytes counts were increased in LIPO patients. Plasma TNF- α and plasma IL-6 were increased in LIPO patients, although the differences were not statistically significant. Duration of HIV infection, duration of HAART and CD4 cell number did not differ significantly between study groups. HIV-RNA was fully suppressed in both study groups (Table 1). Categorizing patients according to the Centers for Disease Control and Prevention (CDC) system of classification clinical stages A, B, and C, where C indicates AIDS diagnosis, did not reveal significant differences between LIPO and NONLIPO patients (Table 1).

[0104] All LIPO and NONLIPO patients were treated with nucleoside reverse transcriptase inhibitors (NRTIs) as part of HAART; which were respectively: lamivudine (83%, 83%), zidovudine (39%, 50%), stavudine (56%, 39%), and didanosine (6%, 11%). HIV-1 protease inhibitors (PIs) were used by 89% of LIPO and NONLIPO patients, that is indinavir (44%, 22%), ritonavir (22%, 39%), nelfinavir (22%, 22%), and saquinavir (17%, 11%), respectively.

[0105] Univariate correlation analyses revealed that suPAR correlated with the following parameters related to dysmetabolism:

- 1) suPAR correlated highly significantly with measures of fat distribution, insulin sensitivity and lipidemia independently of age in all patients (n=36, Figure 2 and Table 2)
- 2) suPAR correlated strongly and positively with TNF- α linking suPAR to a key mediator

of inflammation. suPAR was surprisingly found to be a stronger predictor than TNF- α of fat distribution, insulin sensitivity and total cholesterol (Table 2 and Table 3)

3. 3) suPAR correlated positively with leukocytes, lymphocytes, and monocytes establishing association to blood cells of importance for the immune activation and inflammatory status of the patient (Table 4).

Table 2. Linear correlation coefficients between suPAR and parameters of fat distribution, insulin sensitivity, lipidemia and immunology

n=36	suPAR r-value	P
Limb fat (%)	-0.48	<0.005 ^b
Body mass index	0.19	ns
Total fat mass (% of BW)	0.11	ns
Rd _{clamp}	-0.58	<0.001 ^a
NOGM _{clamp}	-0.66	<0.001 ^b
GOX _{clamp}	0.03	ns
fP-Insulin	0.58	<0.001 ^b
fP-glucose	-0.11	ns
Total cholesterol	0.54	<0.001 ^a
Non-HDL cholesterol	0.51	<0.005 ^a
Triglyceride	0.22	0.19
TNF- α	0.54	<0.001 ^b
IL-6	0.26	0.13
CD4	0.06	ns
Duration of HIV-infection	0.20	ns
Duration of NRTI therapy	0.07	ns
Duration of PI therapy	0.14	ns
Nadir CD4	-0.20	ns
HIV-RNA peak	-0.24	0.17
Abbreviations are given in Table 1; ^a , P<0.05; ^b , P<0.01 following adjustment for age (yr); suPAR, triglyceride, TNF- α , IL-6 and HIV-RNA peak were log-transformed to obtain a normal distribution of these parameters		

Table 3. Linear correlation coefficients between TNF- α and IL-6 versus parameters of fat distribution, insulin sensitivity, lipidemia and immunology

n=36	TNF- α r-value	P	IL-6	P r-value
Limb fat (%)	-0.34	<0.05	0.08	ns
Rd _{clamp}	-0.25	0.15	-0.13	ns
NOGM _{clamp}	-0.17	ns	-0.33	<0.05

n=36	TNF- α r-value	P	IL-6	P r-value
Total cholesterol	0.44	<0.01 ^a	0.14	ns
Non-HDL cholesterol	0.49	<0.005 ^a	0.07	ns
Triglyceride	0.53	<0.001 ^c	-0.03	ns
Abbreviations are given in Table 1; ^a , P<0.05; ^c , P<0.001, following adjustment for age (yr); TNF- α , IL-6, triglyceride and relative increase in CD4 were log-transformed to obtain a normal distribution of these parameters				

Table 4. Univariate correlations (Spearman ρ) between suPAR and blood cells, which are activators of immune and inflammatory response

n=36	ρ
Correlations between suPAR and:	
Leukocytes	0.48**
Neutrophils	0.18
Lymphocytes	0.51**
Monocytes	0.34*
* P<0.05; ** P<0.01	

[0106] Multiple regression analyses were undertaken to demonstrate that suPAR is a robust marker of key parameters of dysmetabolism. Non-oxidative glucose metabolism remained strongly correlated to suPAR ($R^2=0.43$, $P<0.001$) in a model where suPAR, age, TNF- α , IL-6, fasting triglyceride and non-HDL cholesterol were included. Most of the variation of non-HDL-cholesterol could be explained by fasting triglyceride and suPAR ($R^2=0.69$, $P<0.0001$) in a model where also age, TNF- α , and IL-6 were included as explanatory variables. Moreover, suPAR remained a significant marker of Rd ($R^2=0.33$, $P<0.001$) in a model, which included suPAR, age, TNF- α and IL-6.

[0107] Figure 3 shows that fasting plasma suPAR and urine suPAR are highly correlated in HIV-infected patients on stable HAART. Since urine suPAR is shown to be a robust estimate of plasma suPAR, the level of suPAR can be performed on urine as well as plasma samples from such individuals. Urine suPAR corrected for urine creatinine has previously been shown to correlate positively and strongly with plasma suPAR in healthy individuals, and that diurnal changes in urine suPAR are small (Sier et al. 1999, Lab Invest 79:717-722).

[0108] The present data indicate that the circadian changes in plasma concentration of suPAR in HIV-infected patients on stable HAART are limited (Figure 4).

[0109] Conclusion: The data shows that HIV-infected patients on HAART, who display a cluster of dysmetabolic features, are those with relatively high suPAR compared to treatment

matched non-lipodystrophic HIV-infected patients. The fact that suPAR levels are strongly correlated with, respectively, the number of white blood cells, TNF- α , fat distribution, insulin sensitivity and lipidemia, emphasizes that suPAR is a significant link between immune constitution and metabolism in these patients. The present data show that suPAR is a stronger marker of metabolic syndrome in this patient group than TNF- α and IL-6.

[0110] The strong positive correlation between suPAR and TNF- α , which was observed in the present example, show that suPAR is a novel marker of a low-grade inflammatory state in HIV-infected patients on stable HAART regime, and hence diagnostic for this condition. Moreover, it was observed in the present study that suPAR correlated positively with leukocytes, lymphocytes and monocytes, which are themselves markers and effectors of low-grade inflammation. The density of uPAR at the cell surface of these blood cells is high, and much of plasma suPAR is thought to originate from these cells, which would be consistent with the present data.

[0111] Previous epidemiology studies have shown total leukocyte count and lymphocyte count to be inversely correlated with insulin resistance, and to be associated with an increased risk of developing diabetes type 2. It is suggested that chronic activation of the immune system may play a role in the pathogenesis of type 2 diabetes. The data in the present LIPO/NONLIPO HIV study show that suPAR is a novel marker of both low-grade inflammation and metabolic syndrome, supporting the conclusion that low-grade inflammation is a driving factor in the development of metabolic syndrome and thereby of diseases associated the metabolic syndrome, including type 2 diabetes and cardiovascular diseases in HIV patients.

Example 2. suPAR carries prediction value in healthy individuals

[0112] suPAR levels are shown to be a predictor of overall life expectancy.

[0113] Population group: The MONICA 10 cohort: In 1982 to 1984, 4807 individuals aged 30, 40, 50 or 60 years of age selected randomly from the population living in the vicinity of Glostrup University Hospital, Denmark, were invited to participate in a population survey. In 1993 to 1994, 4444 were re-invited and 2656 came and participated in subsequent investigations and gave blood. 2605 plasma samples from 1993/94 were available for the study.

[0114] SuPAR measurement: SuPAR was measured using the suPARnostic™ kit (ViroGates, Copenhagen, Denmark), as described in Example 1. Briefly, the kit comprises an ELISA plate pre-coated with capture monoclonal antibody. According to the protocol, an HRP-labelled detection monoclonal antibody was added to the sample dilution buffer. Then 25 μ l of plasma sample was mixed with 225 μ l of dilution buffer and 100 μ l aliquots (in duplicate) were aliquots into the wells of the ELISA plate and incubated for one hour. The ELISA plate was then washed, and then 50 μ l substrate was added for 20 minutes and reaction stopped with 50 μ l 0.5M H₂SO₄. Plates were measured at 450 nM. The kit intra-assay variation was 2.75% and

inter-assay variation was 9.17%. 2605 samples were measured. The kit standard curve was validated to measure suPAR levels between 0.6 to 22.0 ng/ml. Three samples gave suPAR levels below (2) or above (1) the validated range and were excluded from the results.

[0115] Statistical analysis: Survival was analysed using univariate and multivariate Cox regression and Kaplan-Meier analysis. Differences between groups was analysed using the Mann-Whitney test. The statistical software SPSS version 13 was used for analysis. A p-value < 0.05 was considered significant.

Results:

[0116] suPAR levels as a predictor of overall survival: Median suPAR in the 2602 individuals was 4.03 ng/ml plasma (range 1.3 - 19.9). SuPAR levels increased with age and were higher among women (N=1292, median suPAR 4.26, range 1.9 - 19.9) compared to men (N= 1310, median suPAR 3.84, range 1.3 -17.8), $p < 0.001$ (Figure 5A).

[0117] Overall survival: during the >10 year follow-up, 391 individuals died and 13 were lost to follow-up. Using cox-regression analysis, high suPAR was significantly associated with increased risk of mortality during follow-up (N=2602, RH=1.30 per ng suPAR increase, 95%CI: 1.25 - 1.34). suPAR was associated with survival in both men (N=1310, RH=1.41, 95%CI: 1.35-1.47) and women (1292, RH=1.23, 95%CI: 1.16-1.31).

[0118] Univariate and multivariate analysis: Other markers that were associated with survival was age (RH per year older=1.10, 95%CI: 1.09-1.14), LDL cholesterol (RH=1.002 per unit increase, 95%CI: 1.001-1.002), non-smoking (RH=0.79, 95% CI: 0.72-0.88) and sex (RH=0.613 for women compared to men, 95%CI: 0.50-0.75). When all markers significant in univariate analysis were included in multivariate analysis, suPAR remained a significant and independent predictor of mortality (Table 5, RH per ng suPAR increase 1.22, 95%CI: 1.67-1.28).

Table 5: Age-adjusted multivariate analysis

	Variables in the Equation				
	Wald	Sig.	Exp(B)	95.0% CI for Exp(B)	
				Lower	Upper
SuPAR (per ng/ml increase)	73.5	.000	1.219	1.165	1.275
Sex	28.7	.000	1.780	1.441	2.197
Age (per year older)	248.1	.000	1.102	1.088	1.115
Smoking regularly	36.3	.000	1.884	1.534	2.315
LDL-cholesterol (mmol/l)	0.75	.385	.955	.861	1.060

[0119] Kaplan-Meier analysis: In order to visualize the impact of suPAR on survival, suPAR values were divided into 3 equally sized groups; a low suPAR group (N=867, suPAR range 1.34 - 3.590 ng/ml), a medium suPAR group (N=868, range 3.591 - 4.56) and a high suPAR group (N=867, range 4.57 - 19.4). Kaplan-Meier analysis showed significant differences in survival between the 3 groups, with increasing mortality with increasing suPAR levels ($p < 0.001$). Similarly, this was the case when dividing into 4 groups (Figure 5).

[0120] Conclusion: suPAR levels in apparently healthy individuals are demonstrated to be strongly associated with mortality, independent of other markers. According to the Wald value obtained in multivariate analysis (Table 5) suPAR was the second strongest marker of mortality, after age. Thus, for a given age, sex and smoking status, suPAR significantly predicts the risk of mortality.

Example 3. suPAR as a marker for the efficacy of intervention

[0121] This example demonstrates the influence of life-style intervention on suPAR levels and thereby predicted disease risk of an individual.

[0122] Population group: MONICA 10 cohort as defined above in Example 2.

[0123] Methods: SuPAR was measured as described in Example 2.

[0124] Statistics: Comparison between groups was made using the Mann-Whitney test. Survival was analyzed using Cox regression analysis.

Results:

[0125] SuPAR is significantly associated an individuals smoking habit and with survival in both smokers and non-smokers: Of the 2602 individuals, 1091 were smokers, 1405 non-smokers and 106 occasional-smokers. Smokers had significantly higher suPAR levels compared to non-smokers and occasional-smokers (both $p > 0.001$). There was no difference in suPAR levels between non-smokers and occasional-smokers ($p = 0.7$).

[0126] Smoking and suPAR levels among men: Male smokers had higher suPAR levels compared to non-smokers for all age groups ($p < 0.001$, Figure 6A). Among male smokers (N=580), 133 died during follow-up. suPAR was significantly associated with survival (Cox regression RH= 1.32, 95%CI: 1.24-1.41). Among male non-smokers men (N=674), 100 died during follow-up. SuPAR was significantly associated with survival (RH=1.73, 95%CI: 1.57-1.90). 56 men were occasional-smokers and 6 died during follow-up. RH was 2.16, 95%CI: 0.88 - 5.30).

[0127] Smoking and suPAR levels among women: Female smokers had higher suPAR levels compared to non-smokers for all age groups ($p < 0.001$, Figure 6B). Female smokers ($N=511$) had significantly higher suPAR than female occasional-smokers ($N=56$) ($p < 0.001$), Mann-Whitney. Mean suPAR among female smokers was 5.11 (std 1.38) compared to female occasional-smokers (mean 3.99 ng/ml), std 0.93).

[0128] No difference in suPAR levels was observed between occasional-smokers ($N=50$) and non-smokers ($N=731$), Mann-Whitney $p=0.79$. Female smokers had higher suPAR (mean 5.11 ng/ml, std 1.38) compared to non-smokers ($N=731$, mean 4.07 std 1.14), $p < 0.001$. Cox regression for female smokers showed that suPAR was significantly associated with survival. Among the 511 women, 72 died during follow-up. High suPAR was associated with decreased survival, RH=1.26, 95%CI 1.10 - 1.45.

[0129] Among 731 female non-smokers, 73 died during follow-up. High suPAR was associated with decreased survival, RH=1.21, 95%CI 1.11 - 1.31. 50 women were occasional-smokers and 5 died during follow-up, RH=2.91, 95%CI 1.10-7.69.

[0130] Exercise and suPAR: All individuals in the MONICA cohort were asked about their exercise habits. Individuals were grouped into four groups; Group 1 consisted of individuals reporting they did not exercise regularly ($N=539$), group 2 of individuals doing low-to moderate exercise (taking walk etc) ($N=1454$), group 3 of individuals exercising on a regularly basis (e.g. jogging) ($N=546$) and group 4 consisted of professional sportsmen ($N=17$). As exercise is age dependent, Figure 6C shows suPAR levels among the different exercise groups according to age. In all age groups, except for 71-year old individuals, significantly lower suPAR levels were observed for individuals who exercised regularly compared to individuals who did not exercise ($p < 0.004$ for all analysis). Correlation analysis adjusted for age and sex showed a significant negative correlation between exercise and suPAR ($p < 0.001$, $r = -0.12$).

[0131] Conclusion: suPAR is shown to be a marker that changes with a change of lifestyle, namely individuals that do not smoke or have stopped smoking have significantly lower suPAR levels than individuals that continue to smoke. suPAR is elevated by a number of diverse diseases, which induce immune activation and, a high suPAR level is shown here to predict increased mortality.

[0132] suPAR is also shown to be a dynamic marker of life-style changes associated with exercise and physical fitness. Individuals that have high suPAR can lower their suPAR levels by exercise, an observation that was significant for all age individuals below 71 years of age, which in turn extends their life expectancy.

Example 4: SuPAR is a risk marker for development of cardiovascular disease (CVD)

[0133] CVD is the most common cause of disease in the world. For example, in Sweden, non-communicable conditions account for 83% of all deaths; 41% of total deaths are due to

cardiovascular diseases (CVD), 26% to cancer and about 7% to external causes (intentional and unintentional injuries).

[0134] Population group: Monica10 cohort as defined in Example 2.

[0135] Methods: suPAR measurement as in Example 2.

[0136] Statistics: One minus the Kaplan Meier estimator is shown for time until death. The cumulative incidence (probability of experiencing an event) is shown for time CVD. The cumulative incidence of CVD is estimated in a competing risks model with CVD, censoring due to death and censoring due to other reasons as the three competing causes. All analyses are stratified according to the quartiles of the distribution of suPAR. The quartile cut-offs were generated in the original 2602 patients and is 3.361 ng/ml (25%), 4.029 ng/ml (50%) and 4.910 ng/ml (75%). Follow-up on CVD event was available for analysis until 31st of December 2001.

Results:

[0137] At the time of blood sampling, 208 individuals had already experienced a CVD, and were excluded from the analysis of risk of development of CVD. Those who had experienced a CVD before blood sampling had higher suPAR levels compared to those who had not experienced a CVD at time of blood sampling (Mann-Whitney, $p < 0.001$). During the follow-up, 318 developed a CVD event, of which 70 events were an acute myocardial infarct. The risk of developing a CVD according to suPAR quartiles is shown as a competing risk model in Figure 7.

[0138] suPAR is an independent predictor of risk of CVD: In multivariate analysis, data was available for analysis for a total of 2394 individuals, of whom 316 developed a CVD event. Using continuous suPAR values, suPAR remained a significant predictor of risk of CVD, when age, sex, smoking status and waist circumference are known (Table 6).

Table 6: Multivariate analysis of risk of developing CVD

	Variables in the Equation				
	Wall	Sig.	Exp(B)	95.0% CI for Exp(B)	
				Lower	Upper
SuPAR (ng/ml)	19.290	.000	1.153	1.082	1.229
Sex	6.623	.010	1.400	1.083	1.808
Age	162.778	.000	1.082	1.069	1.095
Waist (cm)	13.207	.000	1.018	1.008	1.028
Smoking regularly	17.102	.000	1.629	1.293	2.053

[0139] LDL-cholesterol was significantly associated with a risk of development of a CVD event in univariate Cox analysis, however it is not a significant predictor of CVD in multivariate analysis (Table 7).

Table 7: Multivariate analysis including LDL-cholesterol (mmol/l)

	Wald	Sig.	Exp(B)	95.0% CI for Exp(B)	
				Lower	Upper
suPAR_ng_ml	20.950	.000	1.158	1.088	1.233
Sex	6.510	.011	1.397	1.081	1.807
Age (per year older)	146.721	.000	1.079	1.066	1.093
Waist (cm)	14.740	.000	1.020	1.010	1.030
smoking	16.466	.000	1.619	1.283	2.044
LDI-cholesterol (mmol/l)	1.564	.211	1.073	.961	1.198

[0140] The metabolic syndrome was also independently associated with the development of CVD. When including the metabolic syndrome in the model, suPAR still remained a highly significant and independent predictor of other CVD events (multivariate analysis in Table 8).

Table 8. Multivariate analysis including the metabolic syndrome, defined according to the NCEP ATPIII 2001 definition with the 2004 glucose modification.

	Wald	Sig.	Exp(B)	95.0% for Exp(B)	
				Lower	Upper
suPAR_ng_ml	19.542	.000	1.154	1.083	1.229
sex	21.346	.000	1.719	1.366	2.164
Age (per year older)	169.495	.000	1.083	1.070	1.096
Smoking regularly	14.865	.000	1.574	1.250	1.981
Metabolic syndrome	7.212	.007	1.438	1.103	1.875

[0141] In a partial correlation analysis, controlling for sex and age, suPAR showed little or no correlation to other known markers of risk of CVD, including body mass index (BMI), LDL-cholesterol and triglycerides (Table 9).

Table 9. Partial correlation between suPAR and known risk markers of CVD adjusted for sex and age.

Control Variables			
SEX & AGE	SuPAR (ng/ml)	Correlation	1.000
		Significance (2-tailed)	

Control Variables			
		df	0
	Regular Smoker (yes or no)	Correlation	-.383
		Significance (2-tailed)	.000
		df	2550
	White blood cells	Correlation	.355
		Significance (2-tailed)	.000
		df	2550
	LDL-cholesterol	Correlation	-.003
		Significance (2-tailed)	.896
		df	2550
	Triglyceride	Correlation	.097
		Significance (2-tailed)	.000
		df	2550
	Waist circumference	Correlation	.027
		Significance (2-tailed)	.172
		df	2550
	Systolic blood pressure	Correlation	.042
		Significance (2-tailed)	.032
		df	2550
	Fasting glucose	Correlation	.081
		Significance (2-tailed)	.000
		df	2550

[0142] Conclusion. suPAR is shown to be a surprisingly strong and independent risk marker of development of a CVD event in apparently healthy individuals. Independent of sex, age, smoking status and the metabolic syndrome, a 1 ng/ml increase in suPAR levels was associated with a 15.4% increased risk (CI95: 8.4 to 22.8%, $P < 0.001$) of developing a CVD during the follow-up. Interestingly, suPAR is not, or only weakly, correlated to classical predictors of CVD development, strengthening the strong and independent role of suPAR as a

risk marker for CVD.

Example 5: The plasma suPAR level is diagnostic of low-grade inflammation and a independent predictor of development of Diabetes type 2

[0143] The number of individuals that develop diabetes (type 2 diabetes) has increased in the last decades. There is a higher prevalence of diabetes in overweight individuals. However, not all overweight individuals develop diabetes. Thus, there is a need for markers to predict the risk of an individual developing diabetes independently of BMI.

[0144] Population group: Monica10 cohort as defined in Example 2.

Methods: suPAR determination as in Example 2; Metabolic syndrome was defined according to the NCEP ATPIII 2001 with the glucose modification of 2004. At baseline, 425 (16.3%) had the metabolic syndrome.

[0145] Results: At the time of blood sampling of the Monica10 cohort, 34 individuals had been diagnosed with diabetes according to ICD-8 and -10 codes. These individuals had significantly higher suPAR levels compared to those who did not have diabetes at time of inclusion (Mann-Whitney, $p < 0.001$). During the follow-up period, 65 out of 2568 individuals, who had not previously been diagnosed with diabetes I or II, went on to develop diabetes.

[0146] When the original cohort (2602 individuals) was divided into 2 groups, based on median suPAR, significantly more individuals having suPAR levels above the median went on to develop diabetes (log rank test, $p < 0.001$, inverse Kaplan Meier plot, Figure 8).

[0147] Multivariate analysis of risk of diabetes: Smoking was not associated with a risk of developing diabetes. In univariate analysis, high suPAR, age (older), sex (being a man) and having a high BMI were significantly associated with an increased risk of developing diabetes. In multivariate analysis (Table 10), these latter factors all remained significantly associated with the risk of developing diabetes.

Table 10. Multivariate analysis of univariate significant predictors of development of diabetes

	Variables in the Equation				
	Wald	Sig.	Exp(B)	95.0% CI for Exp	(B)
				Lower	Upper
Systolic blood pressure	6.895	.009	1.019	1.005	1.034
SuPAR (ng/ml)	19.551	.000	1.247	1.131	1.376
sex	9.547	.002	2.326	1.362	3.974
Age (per year older)	6.510	.011	1.037	1.008	1.066

	Variables in the Equation				
	Wald	Sig.	Exp(B)	95.0% CI for Exp	(B)
				Lower	Upper
BMI	26.968	.000	1.138	1.084	1.195

[0148] Leukocyte count as predictor of development of diabetes mellitus - the strength of suPAR: Number of White blood cells (WBC), i.e. leukocytes, which is accepted as a marker for immune activation and low-grade inflammation in the general population, has been shown to significantly predict development of diabetes mellitus (Vozarova B et al., 2002, Diabetes 51:455-461). WBC count showed a strong positive correlation to suPAR level both univariate ($r=0.34$, $P<0.0001$) and after correction for gender and age (partial correlation $r=0.35$, $P<0.0001$) in the MONICA 10 cohort. Thus, suPAR may be regarded as a marker for immune activation and low-grade inflammation in the general population. Substituting suPAR by WBC count in Table 10, WBC was associated with an increased risk of developing diabetes mellitus independent of age, sex and metabolic syndrome of 13% per 10^9 WBC per Liter (95% CI: 0% to 26%, $P<0.05$). However, taking both suPAR and WBC values, WBC was not-significant ($P=0.$), whereas suPAR remained strongly associated with an increased risk of developing diabetes (27% per 1 ng/mL increase in plasma suPAR, 95% CI: 13% to 43%, $P<0.0001$). Thus, suPAR is a stronger predictor of diabetes than WBC.

[0149] Conclusion. These data show that suPAR is an independent predictor of the risk of developing diabetes. When age, sex and the metabolic syndrome is known, a 1 ng/ml increase in suPAR was associated with a 28% increased risk of developing diabetes (95%CI 15-43%). In addition, these data provide evidence that suPAR correlate strongly with WBC count independently of age and gender, thus establishing suPAR as a marker of immune activation and low-grade inflammation in the general population. Excluding individuals who had baseline blood sugar levels above 6.9 mM or used insulin did not change the significance of suPAR in relation to development of diabetes

Example 6: SuPAR is a marker of the metabolic syndrome

[0150] Population group: Monica 10 cohort as defined in Example 2.

[0151] Methods: suPAR determination as in Example 2. The metabolic syndrome was defined according to the NCEP ATPIII 2001 definition with the 2004 glucose modification (Grundy SM et al., 2005, Circulation 112: 2735 - 2752). The distribution of the 2602 individuals into 5 metabolic classes, identified as having none or 1-5 components of the metabolic syndrome is shown in Table 11. The 5 components are: Elevated waist circumference: Men - Equal to or greater than 40 inches (102 cm) and Women - Equal to or greater than 35 inches

(88 cm); Elevated triglycerides: Equal to or greater than 150 mg/dL; Reduced HDL ("good") cholesterol: Men - Less than 40 mg/dL, Women - Less than 50 mg/dL; Elevated blood pressure: Equal to or greater than 130/85 mm Hg; Elevated fasting glucose: Equal to or greater than 100 mg/dL, (based on the definition provided by (NCEP) Adult Treatment Panel III (ATP III), with minor modifications comprising).

Table 11: Distribution of metabolic syndrome and suPAR values among 2602 individuals

Number of Metabolic syndrome components	Frequency	Percent	Mean suPAR (ng/ml)
0	748	29.2	3.97
1	816	32.7	4.27
2	548	21.8	4.47
3	315	11.4	4.52
4	136	4.6	4.65
5	39	0.4	4.80
Total	2602	100.0	

[0152] Results: SuPAR levels increased as the number of components of metabolic syndrome in an individual increased. Individuals with no components of the metabolic syndrome (0) had significantly lower suPAR levels compared to individuals that had one or more components (Table 11 and Figure 9), $p < 0.001$ Mann Whitney u test. This was the case for both men and women (Figure 10). In univariate Pearson correlation suPAR (log transformed) correlated positively with an increasing number of metabolic components ($r = 0.17$, $P < 0.0001$), a correlation, which remained highly significant after correction for age and gender (partial correlation $r = 0.13$, $P < 0.0001$).

[0153] Conclusion: SuPAR is a strong marker of the metabolic syndrome independent of age and gender. suPAR was also shown to correlate significantly with an increasing fasting glucose (Table 9). In Example 5 it was demonstrated that suPAR strongly predicts diabetes type 2. From these observations it is deduced that suPAR is a strong marker of insulin resistance, which is an underlying cause of the metabolic syndrome and type 2 diabetes mellitus.

Example 7. SuPAR as a marker of Ischemic Heart Disease as a component of CVD

[0154] SuPAR is shown to be a predictive marker of the future development of Ischemic Heart Disease (IHD), classified as a component of CVD.

[0155] Population group: Monica 10 cohort as defined in Example 2

[0156] Methods: suPAR determination as in Example 2.

[0157] Results: Of the 2534 individuals without a previous IHD, 115 developed IHD during the follow-up. Multivariate cox regression analysis showed that a 1 ng/ml suPAR increase, independent of sex, age, smoking status and presence of the metabolic syndrome, was associated with a 16.7% (CI95: 4,8 to 29,9%, $P=0.005$) increased risk for developing IHD. Of the 115 individuals that developed an IHD, 9 had previously developed diabetes. When these 9 individuals were excluded from the analysis, suPAR still remained significantly and independently associated with risk of IHD.

[0158] Age-adjusted multivariate Cox regression analysis for risk of IHD: In an age adjusted Cox regression analysis, a 1 ng/ml increase in suPAR was associated with an increased risk of IHD of 21% ($RH=1.21$, 95%CI: 1.09-1.35), having the metabolic syndrome increased risk with 61% ($RH=1.61$, 95%CI 1.00-2.80), being female compared to male decreased risk with 61% ($RH=0.39$, 95%CI 0.25-0.60), being a non-smoker compared to smoker decreased risk with 33% ($RH=0.67$, 95%CI 0.54-0.84), and an increase in LDL cholesterol (per 10 mM increase) with a 3.5% increase in risk ($RH=1.035$, 95%CI: 1.02-1.05).

[0159] Kaplan Meier Analysis: In order to carry out Kaplan-Meier analysis, individuals were divided into 2 groups based on the median suPAR value of the 2602 individuals included in study (4.03 ng) (Figure 11).

[0160] Conclusion: The results indicate that the suPAR level is a significant predictor of the risk of developing an IHD, as a component of CVD, in individuals who have not previously experienced an AMI. SuPAR predicted the risk of IHD independently of other classical markers of IHD.

Example 8: Algorithm for the prediction of the risk of developing CVD or Diabetes mellitus type II

[0161] A combination of factors, one of which being suPAR, are predictive of the risk of developing CVD or Type II Diabetes. In one embodiment stepwise backward logistic regression was employed to identify those factors that have the highest statistical significance in predicting the outcome of a large number of human subjects, and the risk of developing each of these diseases. The combination of predictive factors or criteria provides a model employed in the form of an algorithm for predicting the health risks of an individual subject.

Stepwise, backwards regression analysis

[0162] The following univariate parameters were significantly associated with the future development of a CVD: suPAR (RH (RH =Hazard ratio) per ng increase 1.23 (95%CI: 1.18-1.29)), Hypertension ($RH = 2.45$ (95%CI: 1.93-3.10)), Asthma ($RH = 1.60$ (95%CI: 1.12-2.30)),

Diabetes (RH = 3.19 (95%CI: 2.01-5.08)), snaps drinking (RH per glass increase 1.03 (95%CI: 1.01-1.04)), smoking (RH = 1.19 (95%CI: 1.08-1.34)), red blood cell count (RH=1.05 (95%CI: 1.02-1.08)), Presence of the metabolic syndrome (RH =1.73 (95%CI: 1.33-2.25)), Total cholesterol (RH per mmol/l increase =1.002 (95%CI: 1.001-1.003)), LDL- cholesterol (RH per mmol/l increase 1.003 (95%CI: 1.001-1.004)), Triglycerides (RH per mmol/l increase 1.001 (95%CI: 1.000-1.002)), Diastolic blood pressure (RH per si (Si pressure unit) increase 1.04 (95%CI: 1.03-1.06)), Systolic blood pressure (RH per si increase 1.036 (95%CI: 1.030-1.041)), Age (RH per year increase 1.08 (95%CI: 1.07-1.10)), Fasting glucose level (RH = 1.02 (95%CI: 1.015-1.026)), Receiving asthma treatment (RH = 1.56 (95%CI: 1.07-2.29)), Sex (men vs women, RH = 1.62 (95%CI: 1.30-2.03)).

[0163] Multivariate analysis: All the univariate significant data were then entered into a multivariate analysis. The multivariate analysis is shown in Table 12. Those factors significant in Table 12 were then entered into a new multivariate analysis (Table 13). The different parameter's ability to predict the event of CVD within 8 years from blood sampling were analysed using receiver operating characteristic curves (ROC), as shown in Figure 12. According to the area under curve (AUC), age, followed by suPAR were the best predictors of the future development of CVD (Table 14). Combining information from all the parameters that were significant in multivariate analysis will improve the predictive accuracy for development of a future CVD. Identifying the linear combination of the parameters yielding the highest AUC can be carried out using the method by Xiong et al., 2004, Med. Decision Making; 24:659-69 based on Fisher's Z-transformation. This way, additional data can be added to the algorithm, also parameters not yet discovered or included in the current model to increase the combined AUC. Furthermore, data can also be omitted from the model if not available. For example, if fasting glucose data are not available, the model can be used with the available data. The result of the model is a risk output, which can be in % risk or as a calculated age (which can then be compared to the age of the patient). If the risk is higher than normal, the calculated age will be higher then the individual's age.

Table 12: Multivariate analysis per 1 unit change of parameters significantly associated with risk of development of CVD in univariate analysis.

	Wald	Sig.	Exp(B)	95.0% CI for Exp(B)	
				Lower	Upper
Diastolic blood pressure	7.539	.006	1.027	1.008	1.047
SuPAR (ng/ml)	11.342	.001	1.135	1.054	1.221
Sex	12.078	.001	1.629	1.238	2.146
Age	89.715	.000	1.074	1.059	1.091
Asthma	.114	.735	1.129	.559	2.278
Alcohol consumption	1.146	.284	1.014	.989	1.039
Treatment for asthma	.683	.409	1.368	.650	2.882

	Wald	Sig.	Exp(B)	95.0% CI for Exp(B)	
				Lower	Upper
Smoking	17.159	.000	1.340	1.167	1.541
LDL-cholesterol	.080	.778	1.000	.997	1.004
Triglycerides	.538	.463	1.001	.999	1.002
Total cholesterol	.067	.796	1.000	.997	1.004
Leucocytes	5.288	.021	1.008	1.001	1.015
Red blood cell count	.014	.906	1.002	.968	1.037
Diabetes	1.585	.208	1.501	.797	2.833
Hypertension	4.096	.043	1.340	1.009	1.779
Systolic blood pressure	1.255	.263	1.006	.995	1.017
Fasting glucose level	3.039	.081	1.009	.999	1.019

Table 13. Multivariate analysis of parameters significantly associated with CVD from Table 12.

	Wald	df	Sig.	Exp(B)	95.0% CI for Exp(B)	
					Lower	Upper
Diastolic blood pressure	26.545	1	.000	1.033	1.020	1.045
SuPAR (ng/ml)	15.132	1	.000	1.145	1.070	1.227
Sex	16.752	1	.000	"1.629	1.289	2.053
Age	160.881	1	.000	1.082	1.069	1.096
Smoking	17.426	1	.000	1.290	1.601	1.516
Leucocytes	4.131	1	.042	1.007	1.000	1.013
Hypertension	9.908	1	.002	1.536	1.176	2.008

Table 14. Area under curve for the data corresponding to Figure 12

	Area Under the Curve				
Test Result Variable(s)	Area	Std. Error(a)	Asymptotic Sig.(b)	Asymptotic 95% Confidence Interval	
				Lower Bound	Upper Bound
suPAR (ng/ml)	.620	.017	.000	.587	.653
Hypertension	.432	.018	.000	.397	.467
Smoking status	.451	.017	.005	.417	.485
Sex	.446	.017	.002	.413	.480
Age	.705	.015	.000	.677	.734

Test Result Variable(s)	Area Under the Curve				
	Area	Std. Error(a)	Asymptotic Sig.(b)	Asymptotic 95% Confidence Interval	
				Lower Bound	Upper Bound
Leucocytes	.546	.017	.008	.512	.580
Diastolic blood pressure	.616	.017	.000	.583	.648

Example 9: Computer implemented assessment of the risk of mortality and/or development of CVD, Diabetes type 2, and/or cancer

[0164] Development of a statistical algorithm for assessment of a patient's risk of mortality and/or developing CVD, cancer or type 2 diabetes within a specified time in which suPAR is a parameter is set out below. The calculation can be executed in a standard computer program like a Microsoft Excel spreadsheet.

[0165] In the present example, the algorithm is based on the data set of covariates determined for subjects in the MONICO10 cohort as set out in Examples 2 and 10, and it has been applied separately for the four different endpoints, time until CVD, cancer, diabetes and death. The algorithm builds on a so-called Proportional Hazards Model, which is a statistical model developed in the statistical field of Survival Analysis. The Baseline Hazard Function is chosen to be a power function, and it yields Weibull distributed event times. The underlying time scale is age, and the event times are left-truncated and right-censored.

[0166] The algorithm produces separate estimates of the risk of acquiring CVD, cancer or diabetes or of dying within a specified time, as a function of an individual's characteristics (from hereon termed "covariates"). These risk estimates are obtained by inserting estimated effects of the covariates obtained from the model in the following expression:

$$P = 1 - \exp(-(\exp(a \cdot \log(\text{age} + t)) - \exp(a \cdot \log(\text{age}))) \cdot \exp(z \cdot \text{beta} - a \cdot \log(b)))$$

[0167] where a, b and beta are estimates of the shape and scale parameters and the effect of the covariates, respectively, obtained from fitting the data to the model, e.g. by using Maximum Likelihood Estimation.

[0168] The variables "age", t and z contain the covariates from a (possibly fictitious) individual for whom the risk of acquiring CVD, cancer or diabetes or of dying can be estimated. P is the probability of the individual acquiring CVD, cancer or diabetes or of dying within t years, "age" is the current age of the individual, and z is a vector containing all other covariates included in the model to calculate the probability.

[0169] The risk estimates depend on which covariates are included in the model, on the endpoint and on the estimates of a , b and β . The core covariates for each risk assessment are as follows:

Time until CVD: Age, sex, suPAR, smoker/occasional or non-smoker, systemic/diastolic blood pressure, waist circumference, HDL-cholesterol, LDL-cholesterol;

Time until cancer: Age, sex, smoker/occasional or non-smoker, suPAR;

Time until diabetes: Age, sex, fasting blood glucose, suPAR;

Time until death: Age, sex, smoker/occasional or non-smoker, suPAR, systemic/diastolic blood pressure, waist circumference.

[0170] Furthermore, in addition to the covariates shown above, the model allows for further covariates to be added e.g. (total -cholesterol, triglyceride, hs-CRP, NTproBNP, BMI, physical activity, alcohol intake and genetic polymorphisms associated with risk of disease). Interactions between the covariates may be included in the model as well.

[0171] In the following, four examples are given, one for each endpoint. Algorithms with fewer covariates are easily included in the model by omitting z 's and β 's, and interactions are included by introducing more z 's and β 's. In all examples, the individual considered is a 58 year old man with a suPAR measurement of 3.0 ng/ml. The risk that is assessed is the risk of experiencing an event within 10 years. The working range for each parameter is given by the 95% confidence interval, which is the interval from the estimate minus twice its standard error to the estimate plus twice its standard error ($[\text{est.}-2*\text{s.e.}; \text{est.}+2*\text{s.e.}]$). In the examples, a standard error (s.e.) provided (in parenthesis) each time an estimate is shown.

[0172] Time until CVD: The estimates are as follows: $\beta_1=0.13$ (0.03) (one unit increase of suPAR concentration), $\beta_2=0.38$ (0.14) (male vs. female), $\beta_3=-0.14$ (0.33) (occasional vs. no smoking), $\beta_4=0.66$ (0.12) (regular vs. no smoking), $\beta_5=0.007$ (0.005) (one mm Hg increase in systolic blood pressure), $\beta_6=0.029$ (0.009) (one mm Hg increase in diastolic blood pressure), $\beta_7=0.009$ (0.006) (one cm increase in waist measurement), $\beta_8=0.64$ (1.58) (one unit increase of HDL-cholesterol) and $\beta_9=0.50$ (0.58) (one unit increase of LDL-cholesterol). The logarithm of the shape parameter is $\log(a)=1.75$ (0.08), and the logarithm of the scale parameter is $\log(b)=5.34$ (0.17).

[0173] According to the formula, if the man is a non-smoker with a blood pressure of 160/120, a waist circumference of 100 cm, an HDL measurement of 0.14 and an LDL measurement of 0.4, he has a risk of $P=45.3\%$ of getting CVD within 10 years. Had he been 48 years of age, his risk would have been 23.4%. His risk of getting CVD within 5 years is $P=21.8\%$. The corresponding risk of a female is 33.8% within 10 years. Had the man had a suPAR measurement of 2.0 ng/ml then he would have had a probability of 41.1% of getting CVD within

10 years, and if it was 4.0 ng/ml it would have been 49.8%. If the man would start smoking regularly, he would increase his risk of getting CVD within 10 years to 68.9%. If the man would bring his blood pressure down to 120/80, he would lower his risk of getting CVD within 10 years to 13.4%. If he could decrease his waist by 5 cm, he would lower his risk to 43.9%. If he could increase his HDL measurement to 0.15, he would obtain a risk of 45.5%, and if he could lower his LDL measurement to 0.3 he would obtain a risk of 43.7%.

[0174] Time until cancer: The estimates are as follows: $\beta_1=0.25$ (0.05) (one unit increase of suPAR concentration), $\beta_2=0.08$ (0.17) (male vs. female), $\beta_3=0.04$ (0.52) (occasional vs. no smoking) and $\beta_4=0.62$ (0.19) (regular vs. no smoking). The logarithm of the shape parameter is $\log(a)=4.83$ (0.07), and the logarithm of the scale parameter is $\log(b)=1.73$ (0.10).

[0175] According to the formula, if the man is a non-smoker, he has a risk of $P=4.2\%$ of getting cancer within 10 years. Had he been 48 years of age, his risk would have been 1.9%. His risk of getting cancer within 5 years is $P=1.7\%$. The corresponding risk of a female is 3.9% within 10 years. Had the man had a suPAR measurement of 2.0 ng/ml then he would have had a probability of 3.3% of getting cancer within 10 years, and if it was 4.0 ng/ml it would have been 5.3%. If the man would start smoking regularly, he would increase his risk of getting cancer within 10 years to 7.6%.

[0176] Time until diabetes: The estimates are as follows: $\beta_1=0.22$ (0.06) (one unit increase of suPAR concentration), $\beta_2=0.79$ (0.27) (male vs. female) and $\beta_3=0.45$ (0.03) (one unit increase of fasting glucose). The logarithm of the shape parameter is $\log(a)=5.81$ (0.32), and the logarithm of the scale parameter is $\log(b)=1.45$ (0.20).

[0177] According to the formula, if the man has a fasting glucose of 5, he has a risk of $P=2.3\%$ of getting diabetes within 10 years. Had he been 48 years of age, his risk would have been 1.3%. His risk of getting diabetes within 5 years is $P=1.0\%$. The corresponding risk of a female is 1.0% within 10 years. Had the man had a suPAR measurement of 2.0 ng/ml then he would have had a probability of 1.8% of getting diabetes within 10 years, and if it was 4.0 ng/ml it would have been 2.8%. If the man were able to lower his fasting glucose level to 4, his risk of getting diabetes would decrease to 1.5%.

[0178] Time until death. The estimates are as follows: $\beta_1=0.20$ (0.03) (one unit increase of suPAR concentration), $\beta_2=0.49$ (0.12) (male vs. female), $\beta_3=-0.12$ (0.33) (occasional vs. no smoking), $\beta_4=0.72$ (0.11) (regular vs. no smoking), $\beta_5=-0.001$ (0.004) (one mmHg increase in systolic blood pressure), $\beta_6=0.021$ (0.007) (one mm Hg increase in diastolic blood pressure) and $\beta_7=0.002$ (0.005) (one cm increase in waist measurement). The logarithm of the shape parameter is $\log(a)=2.01$ (0.06), and the logarithm of the scale parameter is $\log(b)=4.89$ (0.08).

[0179] According to the formula, if the man is a non-smoker with a blood pressure of 160/120 and a waist circumference of 100 cm, he has a risk of $P=16.9\%$ of dying within 10 years. Had

he been 48 years of age, his risk would have been 5.9%. His risk of dying within 5 years is $P=6.7\%$. The corresponding risk of a female is 10.7% of dying within 10 years. Had the man had a suPAR measurement of 2.0 ng/ml then he would have had a probability of 14.1% of dying within 10 years, and if it was 4.0 ng/ml it would have been 20.1%. If the man would start smoking regularly, he would increase his risk of dying within 10 years to 31.5%. If the man would bring his blood pressure down to 120/80, he would lower his risk of dying within 10 years to 7.7%. If he could decrease his waist by 5 cm, he would lower his risk to 16.7%.

Example 10 SuPAR levels is a risk marker of future cancer development

[0180] Aim of study. Previous studies have shown that suPAR levels are predictive of prognosis in certain types of cancer patients. However, it has never been investigated if suPAR levels are predictive of future development of cancer in apparently healthy individuals without cancer. Thus, we have investigated whether suPAR predicts the development of cancer.

Subjects and methods:

[0181] The MONICA 10 cohort: In 1982 to 1984, 4807 individuals aged 30, 40, 50 or 60 years of age selected randomly from the population living in the vicinity of Glostrup University Hospital, Denmark, were invited to participate in a population survey. In 1993 to 1994, 3785 were re-invited and 2656 came and participated in subsequent investigations and gave blood. 2605 plasma samples obtained from June 1993 to December 94 were available for the study. The 2605 participants completed a self-administrative questionnaire. Based on self-reported physical activity, participants were divided into 3 groups: subjects without physical activity (20%), subjects with light physical activity (daily walking, bicycling or less than half an hour sport activity each day (55%), and subjects exercising more than half an hour every day (21%). 4% did not fill out questionnaire concerning exercise. 1091 participants were daily smokers (42%), 106 (4%) reported occasional smoking and 1405 (54%) were non-smokers. Alcohol intake was grouped into two groups based on whether the participants drank more or less than the Danish Health Ministries recommendation (21 alcohol units (1 unit =12 g alcohol) per week for men and 14 for women). Information of cancer was obtained through the Danish National Health Register. BMI was calculated as the weight divided by the squared height (kg/m^2). The metabolic syndrome was defined according to the International Diabetes Federation definition as central obesity (waist men/women > 94/80 cm) together with at least two of following: elevated serum triglycerides >1.7 mmol/l (150 mg/dl), reduced serum high-density lipoprotein cholesterol (<1.03 mmol/l (40 mg/dl), elevated plasma glucose (>5.6 mmol/l (100 mg/dl) or elevated BP (systolic BP >130 mmHg or diastolic BP > 85 mmHg).

[0182] Exercise: Based on self-reported physical activity, the subjects were divided into four groups: 1), subjects without physical activity, 2), subjects with light physical activity (daily walking, frequently bicycling or less than half an hour of sport activity each day), 3), subjects

participating in sport activities for more than half an hour each day and 4), professional athletes.

[0183] High sensitive CRP determinations (hsCRP): Plasma concentrations was determined using the particle enhanced immunoturbidimetry assay (Roche/Hitachi; Roche Diagnostics, Basel, Switzerland) with a range of 0.1-20 mg/l and lowest detection limit 0.03 mg/l. hsCRP was measured in 2510 samples and the median hsCRP was 1.81 (range 0.12 - 98.45 mg/ml).

[0184] SuPAR measurements: SuPAR was measured using the suPARnostic™ kit (ViroGates, Copenhagen, Denmark). The kit was developed to measure disease progression in HIV-1 infected individuals. Briefly, the kit comes with catching monoclonal antibody pre-coated plates and a HRP-labelled detection monoclonal antibody which is added to the sample dilution buffer. Then 25 µl of plasma sample is mixed with 225 µl of dilution buffer and 100 µl in duplicates are added to the pre-coated plate and incubated for one hour. Following wash, 50 µl substrate was added for 20 minutes and the reaction stopped with 50 µl 0.5 M H₂SO₄. Plates were measured at 450 nM with reference 630 nM. 8 plates per working day (312 samples) were measured. The samples were labelled with a number and the technician was unaware of identity of patient sample. The intra-assay variation was 2.75% and inter-assay variation was 9.17%. 2605 samples were measured. The kit standard curve is validated to measure suPAR levels between 0.6 to 22.0 ng/ml. Three samples gave suPAR levels below (2) or above (1) the validated range of the assay and were excluded from the analysis. Cancer data were obtained from the National Health register by a statistician with no knowledge of the individuals suPAR levels. Cancer was diagnosed using the International Classification of Diseases (ICD)-8 and 10 codes. If ICD code was C500:C509, the diagnosis was set as breast cancer. C340:C349 was set to lung cancer. During the follow-up, 28 developed colorectal, 5 developed neoplasms, 37 developed lung cancer, 4 melanoma, 23 breast cancer, one cervix, 3 endometrium, six ovary, 12 prostate, 9 bladder, 9 leukemia. The MONICA study was approved by the local ethical committee.

[0185] Statistics: Kaplan Meier survival analysis was carried out on suPAR quartiles and tested for difference between quartiles using the log rank test. Cox regression analysis was used to determine hazard ratios and independence of suPAR. A $p < 0.05$ was considered significant.

[0186] Results: 66 individuals had a cancer diagnosis before blood sampling and were excluded from analysis. During the follow-up, 136 developed cancer.

[0187] Kaplan Meier analysis on time to cancer diagnosis. The cohort was divided into 4 equally sized groups based on suPAR levels; a low suPAR group (N=650, suPAR range 1.34 - 3.4 ng/ml), a medium low suPAR group (N=651, range 3.4 - 4.0), a medium high suPAR group (N=651, range 4.0 - 4.9) and a high suPAR group (N=650, range 4.9 - 19.9). Individuals that had a cancer diagnosis were censored from the analysis and a Kaplan Meier analysis on time to cancer diagnosis is shown in Figure 14 (log rank $p < 0.001$).

[0188] Age adjusted Cox regression analysis on time to cancer diagnosis: In order to address whether suPAR was an independent predictor of cancer, multivariate analysis was carried out. In age- and sex adjusted Cox regression analysis, a 1 ng suPAR increase (suPAR ranged between 1.3 and 19.9 ng/ml) was associated with an increased Hazard ratio (HR) of developing cancer of 1.34 (95%CI: 1.32 - 1.44%). Univariate (Age adjusted) Cox analysis was carried out for a number of possible predictors of risk of cancer, including sex (P=0.8), diastolic and systolic blood pressure (p=0.5 and 0.3, respectively), metabolic syndrome (p=0.3), diabetes (p=0.9), exercise (p=0.02), smoking (p<0.001), haemoglobin (p=0.9), LDL cholesterol (p=0.14), HDL cholesterol (P=0.4), total cholesterol (P=0.06), triglycerides (P=0.5), fasting glucose level (P=0.5), hsCRP (P=0.4) and BMI (P=0.04). In age-adjusted cox regression multivariate analysis, predictors that were significant or carried a trend in age-adjusted univariate analysis were included. Including suPAR, BMI, Exercise, total cholesterol and smoking, only suPAR and smoking predicted the development of Cancer (Table 15). **Breast cancer (subgroup analysis).** During the follow-up, 22 women developed breast cancer. In univariate analysis, a 1 ng/ml suPAR increase was associated with a RH of 1.37 (95%CI: 1.15 - 1.62), p<0.001). Adjusting for age did not alter this increased risk (age adjusted RH =1.35, 95%CI: 1.13 - 1.61) for developing breast cancer. Smoking, exercise or lipids were not significantly associated with the future risk of developing breast cancer. Only women developed breast cancer. Kaplan Meier analysis on suPAR quartiles (based on the 2602 individuals and excluding individuals with any form of cancer at baseline) showed that individuals with lowest suPAR (thin line Figure 15) had significantly less risk of developing breast cancer compared to individuals in the highest suPAR quartile (fattest line Figure 15).

TABLE 15

suPAR0123	Total N	N of Events	Censored	
			N	Percent
0.00	642	2	640	99.7%
1.00	638	6	632	99.1%
2.00	631	4	627	99.4%
3.00	625	10	615	98.4%
Overall	2536	22	2514	99.1%

Overall Comparisons-TABLE 16

	Chi-Square	df	Sig.
Log Rank (Mantel-Cox)	5.389	1	0.020

[0189] Lung cancer: During the follow-up, 36 individuals developed lung cancer. In age-adjusted multivariate cox regression analysis, suPAR (RH=1.49, 95%CI: 1.32-1.69 per ng/ml increase), sex (men had an increased risk of 3.1 (95%CI: 1.45-6.56) and smoking (daily smokers had an increased risk of 5.16 (95%CI 2.21-12.04) were significantly associated with the risk of developing lung cancer. When looking at the wald value (Table 17), suPAR was the best variable to explain future development of lung cancer.

[0190] Kaplan Meier analysis on suPAR quartiles (based on the 2602 individuals and excluding individuals with any form of cancer at baseline) showed that individuals with lowest suPAR (thin line Figure 16 below), median low suPAR (second-thinnest line), median high suPAR (second fattest line) and highest suPAR (fattest line) had significantly different risk of developing lung cancer ($p < 0.001$, log rank test) compared to individuals in the highest suPAR quartile.

TABLE 17-Case Processing Summary

suPAR0123	Total N	N of Events	Censored	
			N	Percent
,00	642	1	641	99.8%
1,00	638	5	633	99.2%
2,00	631	9	622	98.6%
3,00	625	21	604	96.6%
Overall	2536	36	2500	98.6%

TABLE 18-Overall Comparisons

	Chi-Square	df	Sig.
Log Rank (Mantel-Cox)	25.167	1	.000

[0191] Leukemia: During follow-up, 9 individuals developed leukemia. In univariate analysis, suPAR (RH=1.45, 95%CI: 1.15 - 1.82 per ng increase) was significantly associated with risk of developing leukaemia. Neither age (RH=1.06, 95%CI: 0.99 - 1.14), sex (RH=0.28 for being a man, 95%CI 0.6-1.2) or smoking (RH=2.85, 95%CI:0.7-11.4) was significantly associated with risk of developing leukemia. Due to the small number of events (9 developed leukaemia), Kaplan Meier plot was carried out using median suPAR on the 2602 individuals excluding those who had cancer at baseline (See Figure 17).

Table 19 - Leukemia

suPAR0123	Total N	N of Events	Censored	
			N	Percent
Low suPAR	642	0	642	100.0%
Median low suPAR	638	1	637	99.8%
Median high suPAR	631	2	629	99.7%
High suPAR	625	6	619	99.0%
Overall	2536	9	2527	99.6%

Table 20 - Overall Comparisons

	Chi-Square	df	Sig.
Log Rank (Mantel-Cox)	8.882	1	.003

[0192] Prostate cancer: During follow-up, 10 individuals developed prostate cancer. SuPAR tended to predict development with a RH of 1.31 per ng suPAR/ml increase (95%CI: 0.98 - 1.75). Number of events in each suPAR quartile shown below (p=0.12)

Table 21-Prostate Cancer

suPAR012 3	Total N	N of Events	Censored	
			N	Percent
,00	642	1	641	99,8%
1,00	638	2	636	99,7%
2,00	631	3	628	99,5%
3,00	625	4	621	99,4%
Overall	2536	10	2526	99,6%

[0193] Discussion: The results indicate that suPAR is a highly significant and independent predictor of future risk of developing cancer in individuals who do not have cancer at time of blood sampling. According to the wald-value, suPAR was the strongest predictor of development of overall cancer (see Table 22). Subgroup analysis showed, independent of other predictors, including age, sex and smoking, that increased suPAR levels at baseline increases significantly the risk of developing lung, breast and leukaemia and a trend was observed with regard to prostate-cancer. With regard to minor cancer forms (excluding breast and lung cancer) suPAR was significantly associated with the development of non-lung/non-breast cancer. The minor cancer forms included colorectal, neoplasms, melanoma, cervix, endometrium, ovary, prostate and bladder cancers.

Table 22-All Cancers

	Wald	Sig.	Exp(B)	95,0% CI for Exp(B)	
				Lower	Upper
SuPAR (ng/ml)	20.1	0.000	1.27	1.15	1.41
sex	0.1	0.825	1.04	0.73	1.48
smoking	7.7	0.006	1.70	1.17	2.48
Total cholesterol	3.5	0.060	1.00	0.99	1.00
Exercise	3.4	0.065	0.77	0.587	1.02
BMI	2.1	0.145	0.97	0.928	1.01

Table 23. Age-adjusted cox regression multivariate analysis on development of over-all cancer (N=136)

	Wald	Sig.	Exp(B)	95.0% CI for	
				Lower	Upper
SuPAR (per ng/ml increase)	39.535	0.000	1.492	1.317	1.690

	Wald	Sig.	Exp(B)	95.0% CI for	Exp(B)
				Lower	Upper
AGE (per year older)	24.377	0.000	1.098	1.058	1.140
Sex (men vs. women)	8.540	0.003	3.083	1.449	6.561
Smoking (daily vs non-daily smokers)	14.385	0.000	5.157	2.209	12.038

[0194] As noted above, Kaplan Meier analysis of time to overall-cancer diagnosis is shown in Figure 14. The thin upper line represents the lowest suPAR quartile, the one below (median thin line) represents the second lowest suPAR quartile, the medium thick line represents the second highest suPAR quartile and the fattest (bottom line) represents the highest suPAR quartile. $P < 0.001$, log rank test. SuPAR and risk of developing non-lung/non-breast cancer. When excluding lung and breast cancer, 66 individuals developed other forms of cancer. According to median suPAR levels, those with suPAR above median had significantly higher chance of developing non-lung/non-breast cancer during the follow-up (Figure 18).

Example 11. Further studies on the Use of SuPAR, as a Predictor of cardiovascular disease and mortality in apparently healthy individuals

Materials and methods

[0195] The MONICA 10 cohort: In 1982 to 1984, 4807 individuals aged 30, 40, 50 or 60 years of age selected randomly from the population living in the vicinity of Glostrup University Hospital, Denmark, were invited to participate in a population survey. In 1993 to 1994, 3785 were re-invited and 2656 came and participated in subsequent investigations and gave blood. 2605 plasma samples obtained from June 1993 to December 94 were available for the study. The 2605 participants completed a self-administrative questionnaire.

[0196] Based on self-reported physical activity, participants were divided into 4 groups: subjects without physical activity (20%), subjects with light physical activity (daily walking, bicycling or less than half an hour sport activity each day (55%), subjects exercising more than half an hour every day (21%) and professional athletes ($N=17$). 4% did not fill out questionnaire concerning exercise. 1091 participants were daily smokers (42%), 106 (4%) reported occasional smoking and 1405 (54%) were non-smokers. When entered into the statistical models, occasional smokers were grouped with non-smokers. Alcohol intake was grouped into two groups based on whether the participants drank more or less than the Danish Health Ministries recommendation (21 alcohol units (1 unit =12 g alcohol) per week for men and 14 for women). Information of CVD was obtained in December 2001 through the Danish National

Health Register and the data has previously been validated. A composite CVD endpoint of CV death, non-fatal myocardial infarction and non-fatal stroke were used for statistical analysis because prior occurrences of these endpoints were recorded at baseline. Median CVD follow-up time was 7.6 years. During the follow-up, 316 developed CVD of which 291 were non-fatal. Of the 316, 118 developed an IHD of whom 70 were diagnosed with AMI. After a median of 11.0 years of follow-up (December 2005), information on mortality was obtained through the Civil Registration System. During the follow-up, 388 died, and 13 were lost to follow-up. BMI was calculated as the weight divided by the squared height (kg/m^2).

[0197] Blood pressure was measured sitting after 5 minutes of rest using a mercury sphygmomanometer and the mean of two measurements was reported.

[0198] Exercise: Based on self-reported physical activity, the subjects were divided into four groups: 1), subjects without physical activity, 2), subjects with light physical activity (daily walking, frequently bicycling or less than half an hour of sport activity each day), 3), subjects participating in sport activities for more than half an hour each day and 4), professional athletes. The study was approved by the local ethical committee and the subjects gave informed consent.

[0199] SuPAR measurements: SuPAR was measured using the suPARnostic™ kit (ViroGates, Copenhagen, Denmark). The kit was developed to measure disease progression in HIV-1 infected individuals. Briefly, the kit comes with catching monoclonal antibody pre-coated plates and a HRP-labelled detection monoclonal antibody which is added to the sample dilution buffer. Then 25 μl of plasma sample is mixed with 225 μl of dilution buffer and 100 μl in duplicates are added to the pre-coated plate and incubated for one hour. Following wash, 50 μl substrate was added for 20 minutes and the reaction stopped with 50 μl 0.5M H_2SO_4 . Plates were measured at 450 nM with reference 630 nM. The samples were labelled with a number and the technician was unaware of identity of patient sample. The intra-assay variation was 2.75% and inter-assay variation was 9.17%. 2605 samples were measured. The kit standard curve is validated to measure suPAR levels between 0.6 to 22.0 ng/ml. Three samples gave suPAR levels below (2) or above (1) the validated range of the assay and were excluded from the analysis. There was no correlation between suPAR levels and the date of plasma sampling (samples were obtained between June 1993 and December 1994) indicating the time of sample freezing did not influence on the plasma level of suPAR ($\rho = 0.001$, $p = 0.96$).

[0200] Statistics: The prognostic significance of suPAR was investigated for two different outcomes. These were the time that passed from the blood sample in 1993-94 until 1) diagnosis of CVD and 2) death. After stratifying suPAR in four groups according to its quartiles, a Kaplan-Meier plot was drawn for the time until death. For the time until CVD, a competing risks analysis was carried out with CVD and death as the competing risks, and the estimate of the time until CVD was drawn. Cox's proportional hazards model was used for the analysis yielding hazard ratios for quantification of effect size. First, the effect of suPAR was estimated in different models, all including age and gender. To investigate whether the effects of suPAR on the outcomes were confounded, three groups of variables, termed Lifestyle, Lipids, and

Cardiovascular, were included separately and jointly, and the effect of suPAR was estimated in these models. In the analysis, the following variables were entered as continuous: age, suPAR, total cholesterol, HDL cholesterol, triglycerides, systolic blood pressure, diastolic blood pressure, BMI and waist circumference. The following variables were entered as categorical: gender, smoking (def), exercise (def) and alcohol (def). Second, adjusted for age and gender, hazard ratios are calculated for all other variables.

[0201] Using multiple regression analyses, relative hazards were adjusted for age and gender, calculating the standardized regression quotient for lifestyle measures (daily exercise, alcohol consumption and smoking status), lipids (total cholesterol, HDL cholesterol and triglycerides) and metabolic risk factors (Diastolic and systolic blood pressure, BMI, waist).

[0202] Results: SuPAR was quantified in 2602 individuals. Median suPAR was 4.03 ng/ml plasma (range 1.3 - 19.9). SuPAR levels increased with age and were higher among women (N=1292, median suPAR 4.26, range 1.9 - 19.9) compared to men (N= 1310, median suPAR 3.84, range 1.3 -17.8), $p<0.001$.

[0203] SuPAR predicts CVD independent of classical cardiovascular markers: At the time of blood sampling, 209 individuals had suffered a cardiovascular event and were excluded from the CVD analysis. During the follow-up, 316 developed a cardiovascular event. In age and sex-adjusted Cox regression analysis, a 1 ng/ml suPAR increase was associated with a 19 % increased risk (95%CI: 12 - 25%) of developing a cardiovascular event. To determine whether suPAR was an independent marker of CVD, the influence of classical cardiovascular risk markers on the hazard ratio (HR) of suPAR was determined (see Table 24).

Table 24. The relative hazard per ng suPAR increase adjusted for other risk factors

	CVD	Mortality
	HR (95%CI)	HR (95%CI)
SuPAR	N=2379	N=2602
	1.19 (1.12 - 1.25)	1.24 (1.19 - 1.29)
Lifestyle (Smoking, exercise, alcohol)	N=2324	N=2537
	1.15 (1.08 - 1.22)	1.21 (1.16 - 1.27)
Lipids (Total cholesterol, HDL cholesterol, triglycerides)	N= 2378	N=2601
	1.19 (1.12 -1.25)	1.24 (1.19 - 1.29)
CVD risk factors (Diastolic and systolic blood pressure, BMI, waist)	N=2379	N=2602
	1.18 (1.12 - 1.25)	1.25 (1.19 - 1.30)
All	N=2323	N=2536
	1.13 (1.06 - 1.22)	1.22 (1.16 - 1.28)

[0204] As shown in Table 24, the effect on suPAR remained largely unaltered when including the cardiovascular risk markers and it may be concluded that suPAR is an independent

predictor of CVD.

[0205] In order to directly compare the predictive values of suPAR and classical risk markers on the future risk of developing CVD, the change of one standard deviation (SD) were used. As shown in Table 25, suPAR was the second best predictor (after diastolic blood pressure) for the assessment of a future CVD event.

Table 25. Hazard ratios of a 1 SD change for risk of developing CVD and mortality.

	CVD HR (95%CI)	Total mortality
suPAR	1.26 (1.17 - 1.36)	1.34 (1.27 - 1.41)
HDL	0.92 (0.81 - 1.04)	0.96 (0.86 - 1.07)
Total cholesterol	1.07 (0.96 - 1.20)	0.95 (0.85 - 1.05)
Triglycerides	1.10 (1.00 - 1.21)	1.03 (0.93 - 1.15)
Systemic blood pressure	1.07 (0.96 - 1.20)	1.15 (1.04 - 1.28)
Diastolic blood pressure	1.41 (1.26 - 1.58)	1.14 (1.03 - 1.26)
Body Mass Index	1.05 (0.93 - 1.18)	0.93 (0.84 - 1.04)
Waist	1.22 (1.08 - 1.38)	1.07 (0.95 - 1.20)

SuPAR levels are predictive of longevity

[0206] During the 11-year follow-up, 388 died (150 women and 238 men) and 13 were lost to follow-up. When dividing the cohort into quartiles based on suPAR values, a significant difference in survival was observed between all quartiles.

[0207] In Cox regression analysis, a 1 ng/ml higher suPAR was associated with a 24% increased risk of mortality during follow-up (95%CI: 19 - 29%). In multivariate analysis (Table 24), the risk was largely unaltered by the addition of other mortality risk factors. In the direct comparison with other measured and registered variables, a 1 SD change in suPAR was the strongest marker of mortality. To determine whether the effect of suPAR was most pronounced in the early or late stage of life, a subgroup analysis for the four age-groups was carried out. The RH of mortality among 41 year olds was 1.47 (95%CI: 1.30 - 1.67 per 1 ng/ml suPAR increase), for 51 year olds 1.53 (95%CI: 1.31 - 1.79), for 61 year olds 1.36 (95%CI: 1.23 - 1.49), and for 71 year olds 1.12 (95%CI: 1.05-1.20).

SuPAR levels are correlated to lifestyle

[0208] Of the 2602 individuals, 1091 were daily smokers, 1511 were non-smokers or non-regular smokers (106 of the 1511). In both men and women, daily smoking was associated with significantly elevated suPAR levels. Non-smoking men (N=730) had a mean suPAR level of

3.72 ng/ml (SD 1.20) which was significantly lower than smoking men (N= 580, mean suPAR = 4.57, SD 1.45 ng/ml), $p<0.001$. Similarly, Non-smoking women (N=781) had a mean suPAR of 4.07 (SD 1.13) and smoking women (N=511) of 5.11 (SD 1.38), $p<0.001$. Among the 1511 non-smokers, 614 reported being previous daily smokers. Previous smokers had significantly lower suPAR levels compared to individuals that continued smoking (4.0 vs 4.8 ng/ml, $p<0.001$). The risk of CVD was significant among both non-smokers (age and sex-adjusted, RH=1.11 per ng/ml increase (95%CI: 1.01 - 1.23) and smokers (RH=1.21, 95%CI: 1.10-1.31). With regard to total mortality, the HR among non-smokers was 1.17 (95%CI: 1.10-1.25) and among daily smokers 1.31 (95%CI: 1.21-1.42) per ng suPAR increase.

[0209] It was also determined whether uPAR levels were influenced by exercise. Using the Mann Whitney test, it was found that individuals that exercised more than 30 minutes per day had significantly lower suPAR levels than individuals reporting not to exercise, an observation that was significant among the 41 ($p<0.001$), 51 ($p<0.001$), and 61 ($p=0.01$) year olds.

[0210] The results indicate that the plasma level of suPAR predicts future cardiovascular disease. The risk increase per ng/ml increase in suPAR was not decreased by adding classical markers of CVD, indicating that suPAR is a novel and independent marker of CVD.

Example 12: From healthy to disease, a composite analysis

Materials and methods

[0211] The MONICA 10 cohort as described above.

[0212] To determine the ability of suPAR to predict the risk of developing a disease (CVD, T2DM, cancer or dying) in healthy individuals, we excluded the 220 individuals who had a previous diagnosed event of CVD, T2DM, or cancer at baseline. In the 2382 remaining individuals, age and sex-adjusted Cox regression analysis, a 1 quartile increase in suPAR was associated with a 59% increased risk of developing the composite endpoint consisting of the diseases CVD, T2D, cancer or death (95%CI: 47-72%, Table 26). After adjustment for other markers, suPAR remained a significant predictor of disease with a HR of 41% (95%CI: 29-57 %)(Table 26). Individuals within the highest suPAR quartile were at highest risk of developing disease ($P<0.001$).

[0213] Table 26, below, shows the hazard ratio for inter-quartile range of each variable (except smoking, exercise and alcohol which were entered as grouped variables). The inter-quartile ranges are shown in Table 1. Individuals with history of the disease investigated were excluded from the analysis. All hazard ratios were adjusted for gender and age.

Table 26

Variable	Groups/unit	CVD	Diabetes	Cancer	Mortality	Composite
suPAR	per 1.55 ng/ml	1.35 (1.23-1.49)	1.46 (1.20-1.76)	1.59 (1.39-1.81)	1.39 (1.31-1.48)	1.59 (1.46-1.73)
Smoking	No	1	1	1	1	1
	Occasional	0.89 (0.47-1.69)	0.58 (0.08-4.34)	1.99 (0.95-4.16)	0.86 (0.45-1.62)	1.02 (0.59-1.76)
	Regular	1.78 (1.42-2.24)	0.86 (0.43-1.71)	2.58 (1.85-3.60)	2.11 (1.72-2.59)	2.15 (1.76-2.63)
Exercise	Inactive	1	Not analyzed*	1	1	1
	Light active	0.65 (0.49-0.86)	Not analyzed*	0.72 (0.49-1.05)	0.56 (0.44-0.71)	0.36 (0.11-1.16)
	Highly active	0.51 (0.36-0.73)	Not analyzed*	0.42 (0.25-0.73)	0.39(0.28-0.52)	0.52 (0.17-1.64)
	Professionals	0.41 (0.06-2.20)	Not analyzed*	1.25 (0.91-1.71)	1.6 (0.5-5.11)	0.83 (0.26-2.62)
Alcohol	Below recommended	1	1	1	1	1
Intake	Above recommended	1.48 (1.12-1.95)	2.25 (1.09-4.65)	1.22 (0.82-1.83)	1.63 (1.28-2.08)	1.52 (1.19-1.93)
HDL	Per 0.53 mmol/l	0.90 (0.77-1.05)	0.92 (0.18-4.70)	0.88 (0.71-1.09)	0.95 (0.83-1.10)	0.74 (0.46-1.10)
Triglycerides Per 0.83 mmol/l		1.08 (1.00-1.17)	1.28 (1.16-1.40)	1.29 (1.03-1.64)	1.03 (0.94-1.12)	1.07 (0.99-1.15)
Total cholesterol per 1.4 mmol/l		1.10 (0.95-1.26)	0.89 (0.64-1.24)	0.77 (0.50-1.20)	0.94 (0.82-1.07)	1.03 (0.91-1.16)
Systolic blood pressure		1.60 (1.39-1.85)	1.74 (1.28-2.36)	1.76(1.19-2.61)	1.19 (1.04-1.37)	1.34 (1.18-1.52)
Diastolic blood pressure		1.32 (1.21-1.44)	1.85 (1.34-2.56)	1.96 (1.31-2.93)	1.12 (1.03-1.21)	1.26 (1.10-1.44)

Variable	Groups/unit	CVD	Diabetes	Cancer	Mortality	Composite
BMI	Per 5.27 kg/m ²	1.07 (0.93-1.23)	2.07 (1.48-2.87)	0.78 (0.63-0.97)	0.92 (0.80-1.05)	0.98 (0.88-1.11)
Waist	Per 18 cm	1.33 (1.11-1.59)	2.55 (1.61-4.04)	0.89 (0.68-1.16)	1.10(0.93-1.30)	1.16 (0.99-1.37)
Blood glucose	Per 0.6 mmol/l	1.11 (1.05-1.16)	8,43 (3.72-19.1)	1.01 (0.92-1.10)	1.09 (1.05-1.13)	1.23 (1.10-1.38)
hsCRP	Per 3.17 mg/ml	1.59 (1.31-1.94)	1.12 (1.01-1.24)	1.23 (0.96-1.57)	1.58 (1.35-1.84)	1.11 (1.07-1.14)
Leucocytes	Per 2.2x10 ⁹ cells/l	1.35 (1.17-1.56)	1.37 (0.98-1.93)	1.22 (1.03-1.46)	1.37 (1.23-1.53)	1.37 (1.24-1.52)
*Not analysed due to too few endpoints						

Example 13 Antibodies for the quantification of suPAR in a sample from an individual.

[0214] A method for measuring suPAR and suPAR fragments in a biological sample from an individual employs the antibodies (VG-1 and VG-2) having the following characteristics:

- are able to recognize and bind full length suPAR
- are able to recognize and bind the D₂D₃(88-277) suPAR fragment obtained by cleavage with chymotrypsin
- do not recognize or bind the chemotactic amino acid sequence SRSRY, which is displayed through cleavage of suPAR with chymotrypsin
- do not recognize or bind the amino acid sequence ATVYSRSRY, which is located from amino acids 84 to 92 in the linker region between D₁ and D₂D₃

Description of the VG-1 and VG-2 antibody specificity

[0215] The specificity of antibodies used in the ELISA, suPARnostic® (commercially available from Virogates A/S, Denmark) namely VG-1 (coating antibody) and VG-2 (detection antibody), are shown in figure 19 and 20. These monoclonal antibodies, raised against full length human

suPAR in rat (VG-1) and mouse (VG-2), specifically recognize full length suPAR, $D_1D_2D_3(1-277)$. The antibodies also specifically recognize the suPAR fragment, $D_2D_3(88-277)$, which is obtained by cleavage with chymotrypsin. The antibodies do not recognize the synthesized amino acid sequence SRSRY, which corresponds to the chemotactic active residues 88 to 92 of $D_2D_3(88-277)$ that is displayed at the amino-terminal end following cleavage with chymotrypsin (figure 19). Neither do the antibodies recognize the synthesized amino acid sequence ATVYSRSRY, which corresponds to the amino-terminal residues 84 to 92 that can be obtained by cleavage with uPA. However, a polyclonal antibody mixture obtained by immunization of rabbits with full length suPAR was able to recognize - though yielding low intensities - the peptide sequence ATVYSRSRY. Furthermore, a polyclonal antibody obtained from rabbits immunized with the SRSRY peptide sequence recognized both the SRSRY and the ATVYSRSRY sequences.

[0216] In conclusion, it has been shown that the antibodies (VG-1 and VG-2) both recognize and bind full length suPAR and the suPAR fragment $D_2D_3(88-277)$ obtained by cleavage with chymotrypsin. However, the antibodies do not recognize or bind to the chemotactic active sequence SRSRY or the amino-terminal end AVTYRSRY, which can be obtained by cleavage with uPA.

[0217] Antibodies of the VG-1 and VG-2 type are suitable for use in any of these methods.


Example 14: suPAR is an inducer of inflammation


[0218] Elevated suPAR levels are shown to be diagnostic of a pro-inflammatory condition and to predict the risk of disease. If suPAR induces the pro-inflammatory condition, removal of suPAR e.g. by administering anti-suPAR antibodies, would lower the inflammation of the individual.

[0219] The purpose of the present example was to investigate whether suPAR induces the sub-clinical pro-inflammatory condition or whether it reflects a pro-inflammatory condition. In the present example, purified full-length suPAR ($D_1D_2D_3(1-277)$) as well as the D_2D_3 fragment of suPAR ($D_2D_3(88-277)$), was added to freshly drawn human blood and 10 relevant cytokines were measured at different time points following stimulation.

Materials and methods

[0220] An overview of the experimental flow is shown below





CHO-cell line production of suPAR D1D2D3₍₁₋₂₇₇₎

[0221] The CHO cell line was stably transfected with recombinant human full length uPAR gene.

[0222] The CHO was grown in 90% Dulbecco's Modified Eagle's Medium with GlutaMAX™ I, Sodium Pyruvate (D-MEM, Invitrogen Cat. No. 31966-021) and 10% FBS at 37°C in a humidified 5% CO₂ incubator supplemented with 50 Units/ml penicillin and 50µg/ml streptomycin (Sigma, Cat No. P4333) and 100 nM methotrexate (MTX, Sigma-Aldrich Cat. No. A6770) as a selective agent ensuring that only stable transfected cells would dominate the population.

D1D2D3₍₁₋₂₇₇₎ purification from cell culture medium by affinity chromatography

[0223] A VG-1 anti-suPAR column (ViroGates, Denmark) was equilibrated with 20 ml PBS before the CHO-cell-line media containing recombinant human suPAR was applied at a volumetric pump rate of 1.5 ml/min. After sample application, the column was washed with 20 ml PBS pH 7.4 to remove unbound proteins and medium components, especially albumin, the most abundant protein in the bovine serum growth supplement. Finally, the suPAR bound by immunoaffinity to VG-1 was eluted with 0.1 M citrate buffer pH 3.0 (Sigma-Aldrich Cat. No. C1909) to break the non-covalent bonds and elute suPAR fragments only. The eluates were collected in 2.5 ml volumes and buffer exchanged into 0.1 M PBS pH 7.4 using the PD-10 columns as previously described and kept either at -20°C until further analysis.

Chymotrypsin digestion of D1D2D3₍₁₋₂₇₇₎ to D2D3₍₈₈₋₂₇₇₎ and D1₍₁₋₈₇₎

[0224] To generate D2D3₍₈₈₋₂₇₇₎ from full length suPAR, the eluates were treated with chymotrypsin (Chymotrypsin sequencing grade Roche Applied Biosciences (Cat. No. 11 418 467 001) which cleaves between the amino acids 87 and 88). 3.5 mL suPAR eluate at a concentration of 9.2 µg/mL suPAR was mixed with 3.5 mL digestion buffer consisting of 0.1% CHAPS and 0.1 M Tris.HCl pH 8.1 giving a final buffer concentration of 0.05% CHAPS and 0.05% Tris-HCl and a final suPAR concentration of 4.6 µg/ml. The solution was aliquoted in eppendorf tubes in volumes of 250 µl and chymotrypsin added to yield the desired final concentration ranging from 1 to 300 ng/mL chymotrypsin. PBS was added to give a final volume of 500 µl. After 4 or 24 hours of incubation at 37°C the enzymatic reaction was terminated by freezing the relevant sample at -20°C. The resulting products were size separated by SDS-PAGE and then transferred to PVDF membrane by western blotting (figure 21).

Additional affinity chromatography using D3 and D1 binding antibodies, respectively Generation of pure D1D2D3

[0225] After the primary affinity purification by the VG-1 column, the eluate was buffer exchanged into a PBS buffer pH 7.4 and applied to a D1 antibody column to remove any D₂D₃, giving a pure D1D2D3 eluate. This was buffer exchanged into PBS pH 7.4 and the concentration was determined using the suPARnostic ELISA kit.

Generation of pure D2D3 suPAR fragment

[0226] When preparing an eluate consisting of D2D3, the procedure required additional steps. Preceding the primary purification of suPAR, and a buffer exchange into PBS, an enzymatic digestion with chymotrypsin was employed as described above yielding the fragments D₁, D2D3 and possibly still some D1D2D3. In order to purify D2D3 from the remaining suPAR fragments, the anti-D1 antibody column was used for negative selection, whereby the flowthrough containing D2D3 and chymotrypsin was collected. This was run through the VG-1 column one more time. Finally, the eluate was buffer exchanged into PBS pH 7.4 and the concentration determined. Purity of the obtained D1D2D3 and D2D3 fragments were visualized using silver staining (figure 22) and validated as suPAR molecules using western blotting (figure 20) and showed that the correct, pure, compounds had been prepared.

[0227] In order to compare and normalize the effects of adding suPAR to blood, the mock sample (negative control) was prepared by adding citrate elution buffer to the column without prior loading of suPAR containing sample and then buffer exchanging the eluate (with no

suPAR) into PBS pH 7.4, to mimic the buffer of the suPAR samples. If the elution would cause any bonds to break, either internally in the antibody or between the antibody and the sepharose beads, and contaminate the suPAR eluates, the mock samples would contain the same background as the suPAR eluates

Exposure of D1D2D3 and D2D3₍₈₈₋₂₇₇₎ to whole blood

[0228] A 27-year-old healthy female was used as blood donor and no food was ingested 2 hours prior blood sampling in either of the assays. Blood was collected by venipuncture into seven sterile 10 ml lithium-heparin coated tubes (Venoject Li-Heparin tubes, Terumo Cat. No. VT-100SHL), which were to be used for the experiment, and two 4 ml EDTA tubes that were immediately sent for blood data analysis at the Clinical Biochemical Department, Unit 339, Hvidovre Hospital. The concentration of the respective blood cells was determined by enzyme based optical cytometry (employing light scatter) using a Bayer Advia® 120 Hematology Analyzer and the distinction between CD4 and CD8 T-cells by flow cytometry with fluorescence labelled antibodies (Center for Disease Control recommended 4 colour method with gating on CD45⁺-lymphocytes. Neutrophil concentration was in the 1st, 2nd and 3rd experiment 4.0, 4.3 and 3.1 * 10⁹ cells/L respectively, eosinophils 0.02, 0.05 and 0.07 * 10⁹ cells/L, monocytes 0.3, 0.5 and 0.4 * 10⁹ cells/L and lymphocytes (unspecified) 1.5, 1.7 and 1.9 * 10⁹ cells/L. Basophil concentration was 0.01 * 10⁹ cells/L in all the experiments. The overall concentration of leukocytes varied from 5.5, 6.6 to 5.8 * 10⁹ cells/L.

Analysis of cytokine profile by Luminex multiplex

[0229] Plasma was thawed, kept on ice and analyzed following the manufacturer's instructions for the Cytokine 10-plex AB Bead Kit, hu (BioSource™, Invitrogen, Cat. No. LHC0001) or 10-plex Cytokine Ultrasensitive™ AB Bead Kit, hu (BioSource™, Invitrogen, Cat. No. LHC6004) on the Luminex®xMAP system (Luminex Corporation) using the software Starstation 2.0 (Applied Cytometry Systems). If intra-assay variation exceeded 20%, the sample was re-measured. Based on the intra-assay variation of the assay, an increase of a markers was considered biologically significant if the analyse in question exceeded the control (mock) by more than 50%.

Results

[0230] Three independent experiments were carried out where the whole blood samples were stimulated 5 min, 3 h, 6 h or 24 h with 100 ng/ml D1D2D3₍₁₋₂₇₇₎, D2D3₍₈₈₋₂₇₇₎ or with the mock eluate. Samples from each of the experiments were measured in duplicates.

[0231] GM-CSF, IL-2, IL-4 and IL-5 were not significantly induced by D1D2D3 or by D2D3. With regard to GM-CSF and IL-5, the obtained results were below the lower limit of quantification. Table 27 show representative results from three independent experiments.

Table 27: Cytokine concentrations (pg/ml) of whole blood samples stimulated 5 min, 3 h, 6 h or 24 h with either 100 ng/ml D1D2D3₍₁₋₂₇₇₎, D2D3₍₈₈₋₂₇₇₎ or with the mock eluate.

GM-CSF	Assay results			
	5 min	3h	6h	24h
Mock (1.)	0,59	0,59	0,59	0,59
100 ng/ml D1D2D3	0,59	0,59	0,59	0,59
100 nh/ml D2D3 ₍₈₈₋₂₇₇₎	0,59	0,59	0,59	0,59
IL-2	Assay results			
	5 min	3h	6h	24h
Monk(1.)	0,59	0,59	0,92	3,27
100 ng/ml D1D2D3	0,59	0,59	0,92	3,89
100 ng/ml D2D3 ₍₈₈₋₂₇₇₎	0,59	0,59	1,61	3,13
IL-4	Assay results			
	5 min	3h	6h	24h
Mock(1.)	1,23	1,87	4,26	8,08
100 ng/ml D1D2D3	1,07	2,62	4,79	12,95
100 ng/ml D2D3 ₍₈₈₋₂₇₇₎	1,26	2,40	3,09	7,35
IL-5	Assay results			
	5 min	3h	6h	24h
Mock (1.)	0,60	0,60	0,60	0,60
100 ng/ml D1D2D3	0,60	0,60	0,60	0,60
100 ng/ml D2D3 ₍₈₈₋₂₇₇₎	0,60	0,60	0,60	0,60

[0232] In contrast, the pro-inflammatory cytokines, INF-gamma, IL-1 beta, IL-6, IL-8 and TNF-alfa were up-regulated by D1D2D3₍₁₋₂₇₇₎ while D2D3 had less effect although TNF-alfa upregulation was observed (figure 23)

Conclusions

[0233] This example showed that full length suPAR is an inducer of certain cytokines as well as the chemokine IL-8. As can be seen in table 28 below, the induction in the early phase (up to 6 hours after stimulation) was observed with regard to IL-1 beta, IL-6, IL-8 and TNF-alfa while no significant upregulation was observed of the other 6 cytokines investigated. The early upregulated cytokines are all pro-inflammatory markers. As often seen when adding a pro-inflammatory marker, the induced effect is counteracted by the late induction (after 24 hours) of IL-10, which was also the case with regard to 24-hour stimulation with D1D2D3.

[0234] Based on these results, we conclude that full length suPAR is an active inducer of the inflammatory pathways.

Table 28: Schematic overview of the measured cytokines and chemokine (IL-8), source- and target cells and function of cytokines on target cells.

	Function	Pro- or anti inflammatory	D2D3 early	D2D3 late	D1D2D3 early	D1D2D3 late
GM-CSF	Inhibits apoptosis of target cells, proliferation and differentiation into granulocytic and macrophage lineages, functional activation of a wide range of cells	Neither	No effect	No effect	No effect	No effect
IFN-γ	Up-regulates class I and II MHC expression on macrophages. Activates macrophages, NK and T cells. Ig class switch to IgG2a of activated B cells. Inhibit viral-, bacterial and parasitical replications.		No effect	No effect	No effect	Up-regulated
IL-1β	Induction of other cytokines. Costimulates activation of T cells with IL-2. Activates NK cells. B cell maturation and proliferation. Fibroblast proliferation. Increase expression of adhesion molecules on endothelial cells. Mediator of acute-phase response. Pyrogenic. Induce inflammation. Stimulate hepatocytes to produce acute-phase proteins.	Pro-: participate in the acute-phase response and synergize to mediate inflammation, fever, shock, and death.	No effect	No effect	Up-regulated	Up-regulated
IL-2	Activation, maturation and proliferation of T, B, NK cells, monocytes and macrophages. Stimulates cytolytic activity of activated T cells, large granular lymphocytes and monocytes. Stimulates T cell motility.		No effect	No effect	No effect	No effect
IL-4	Proliferation and differentiation of B cells. Activation to IgG1 and IgE synthesis. Activation of monocytes. Macrophage up-regulation of MHC Class II. Expansion of Th ₂ subset. Mast cell development and proliferation.	Anti-: reduces endotoxin-induced TNF and IL-1 production	No effect	No effect	No effect	No effect
IL-5	Eosinophil activation, differentiation, proliferation and		No effect	No effect	No effect	No effect

	chemotaxis. Basophil activation. Proliferation and differentiation of B cells. IgA synthesis of B cells.					
--	--	--	--	--	--	--

IL-6	Proliferation and differentiation of B cells into plasma cells. Antibody production of plasma cells. Differentiation of T cells into cytotoxic T cells. Stimulates production of acute phase response proteins from the liver. Mediates and regulates inflammatory responses.	Pro-: participate in the acute-phase response and synergize to mediate inflammation, shock, and death. Anti-: inhibits TNF production	No effect	No effect	Up-regulated	Up-regulated
IL-8	Chemotactic for neutrophils and macrophages	Pro-: is one of the first cytokines to be released in inflammation	No effect	No effect	Up regulated	Up-regulated
IL-10	Inhibits production of pro-inflammatory cytokines by monocytes, macrophages and mast cells. Inhibits IL-2 production of T cells. Inhibits antigen-specific activation of T cells and production of cytokines. Growth and differentiation factor for B cells. Inhibits monocyte and macrophage MCH class II and costimulatory molecule expression, and NO production.	Anti-: suppresses lymphocyte functions and regulates production of pro-inflammatory cytokines	No effect	Up-regulated	No effect	Up-regulated
TNF-α	Mediates host response to Gram-negative bacteria and other infectious agents. Death of tumor cells. CAM and cytokine expression of macrophages. Increase expression of adhesion molecules on endothelial cells. Induce inflammation.	Pro-: participate in the acute-phase response and synergize to mediate inflammation, shock, and death.	No effect	Border-line Up-regulated	Up-regulated	Up-regulated

Abbreviations: DC: dendritic cells; NK: natural killer

[0235] Summary of aspects of the disclosure

1. A method for diagnosing low-grade inflammation and/or metabolic syndrome in a mammalian subject, in particular, a human subject, comprising:

1. (a) performing an *in vitro* measurement of the level of a marker in the form of soluble urokinase plasminogen activator receptor (suPAR), in one or more samples comprising a biological fluid derived from said subject, and
2. (b) using the one or more measurement value obtained as a factor in the diagnosis of low-grade inflammation related disease and/or metabolic syndrome.

2. A method for predicting the risk of developing a metabolic syndrome-related disease and/or low-grade inflammation-related disease and/or cancer and/or the risk of mortality within ten

years in a mammalian subject, in particular, a human subject, comprising:

1. (a) performing an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), (ii) and/or D2D3 cleavage products of suPAR in one or more samples comprising a biological fluid derived from said subject, and
 2. (b) using the measurement values obtained as a factor to predict the risk of developing a metabolic syndrome-related disease and/or low-grade inflammation-related disease and/or cancer and/or the risk of mortality within ten years.
3. A method of predicting the risk of developing a metabolic syndrome-related disease and/or low-grade inflammation-related disease according to aspect 2, wherein said disease is selected from the group consisting of cardiovascular disease, chronic obstructive lung disease, eczema, obesity, autoimmune disease, cancer and diabetes type 2.
 4. A method according to aspect 1 or 2, wherein said biological fluid is blood, blood serum, blood plasma or urine.
 5. A method according to aspect 1, wherein said subject is Human immunodeficiency virus-infected and receiving Highly Active Anti-Retroviral Treatment.
 6. A method according to aspect 1 or 2, wherein said subject is apparently healthy.
 7. A method according to aspect 1 or 2, wherein more than one said sample comprising a biological fluid is derived from said subject at more than one points in time.
 8. A method according to aspect 7, further comprising comparing the predicted risk at one or more time points for assessing the efficacy of a prophylactic course of action and/or treatment.
 9. A method according to aspect 7, wherein said prophylactic course of action comprises physical exercise.
 10. A method according to aspect 1 or 2, wherein said *in vitro* measurement is an ELISA assay.
 11. A method according to aspect 1 or 2, further comprising in step b) said one or more measurement value is compared with a reference value for the prediction of the clinical outcome.
 12. A method according to aspect 1, where in step b) one or more of age, gender, ethnicity, smoker/non-smoker status, and waist circumference measurement are additionally used to diagnose low-grade inflammation.
 13. A method for detecting the level of low-grade inflammation and/or metabolic syndrome in a mammalian subject comprising:
 1. (a) isolating a sample from a subject;
 2. (b) measuring the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in said sample

of (a);

3. (c) comparing the measurement obtained in step (b) with a reference value

wherein an increase in the level of said marker in said sample indicates increased low-grade inflammation and/or the presence of metabolic syndrome.

14. The method according to aspect 13, wherein said reference value is a standard of levels of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in a population.

15. The method according to aspect 13, wherein said reference value is obtained by measuring the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) a D2D3 cleavage products of suPAR from a sample from a subject free of low-grade inflammation and/or metabolic syndrome.

16. The method according to aspect 13, wherein the level of said marker is increased at least about 10% as compared to said reference sample.

17. The method according to aspect 13, wherein said method further comprises conducting a glucose tolerance test, assaying TNF-alpha, IL-6, leukocyte count, fasting plasma-glucose or fasting blood-glucose, blood pressure, cholesterol, C-reactive protein and/or triglyceride levels.

18. A method for predicting the risk of developing a metabolic syndrome-related disease, low-grade inflammation-related disease and/or cancer and/or the risk of mortality within ten years in a subject, comprising

1. (a) isolating a sample from a subject;
2. (b) measuring the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), (ii) and/or a D2D3 cleavage products of suPAR;
3. (c) comparing the measurement obtained in step (b) with a reference value, wherein an increase in the levels of said marker indicates an increased risk of low-grade inflammation, metabolic syndrome, cancer and/or the risk of mortality within ten years.

19. The method according to aspect 18, wherein the level of said marker is increased at least about 10% and /or 1 ng/ml in said sample as compared to said reference sample.

20. The method according to aspect 18, wherein the disease is diabetes.

21. The method according aspect 20, wherein step (b) further comprises measuring glucose levels.

22. The method according to aspect 18, wherein the disease is cardiovascular disease.

23. The method according aspect 22, wherein step (b) further comprises measuring blood pressure, HDL-cholesterol, LDL-cholesterol, total cholesterol and/or triglycerides.

24. The method according to aspect 18, wherein the disease is cancer.

25. The method according to aspect 24, wherein the cancer is breast cancer, leukemia,

prostate cancer or lung cancer.

26. The method according aspect 25, wherein step (b) further comprises determining age and smoking.

27. The method according to aspect 18, wherein increased risk of mortality within ten years is measured.

28. The method according to aspect 27, wherein step (b) further comprises measuring brain natriuretic peptide concentrations.

29. A method for monitoring the treatment of low-grade inflammation and/or metabolic syndrome in a mammalian subject comprising:

1. (a) isolating a sample from a subject undergoing treatment for low-grade inflammation and/or metabolic syndrome;
2. (b) measuring the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), (ii) and/or D2D3 cleavage products of suPAR,

wherein a decrease in the level of said marker used in step (b) is an indication of progression of treatment.

30. The method according to aspect 29, wherein the level of said marker is decreased at least about 10% and/or 1 ng/ml.

31. A method for treating low-grade inflammation and/or metabolic syndrome in a mammalian subject comprising administering to a subject in need thereof an amount of a substance that decreases the level of suPAR effective to treat said low-grade inflammation and/or metabolic syndrome.

32. The method according to aspect 31, wherein said substance is an anti-suPAR antibody.

33. The method according to aspect 32, wherein said anti-suPAR antibody is a monoclonal or polyclonal antibody.

34. The method according to aspect 31, wherein said substance is a suPAR analog or derivative capable of binding uPA.

35. The method according to aspect 31, wherein said substance is a D2D3 cleavage product.

36. A pharmaceutical composition for use in treatment or prevention of low-grade inflammation and/or metabolic syndrome comprising a substance that decreases the level of suPAR in a mammalian subject.

37. Use of a substance for preparing a therapeutic composition for the treatment or prevention of low-grade inflammation and/or metabolic syndrome, wherein said substance decreases the level of suPAR in a mammalian subject.

38. A computer readable medium having computer logic stored therein configured to cause,

when executed on a data processing system, the data processing system to compute risk and/or change in risk of a low-grade inflammation-related disease, a metabolic syndrome-related disease, cancer and/or mortality within ten years comprising:

1. (a) logic for receiving one or more measurements derived from a mammalian subject wherein at least one measurement is an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject;
2. (b) logic for analysis of one or more of said measurements of step (a); and
3. (c) logic for computing risk and/or change in risk of said low-grade inflammation-related disease, a said metabolic syndrome-related disease, said cancer and/or said mortality within ten years based on analysis in step (b) of said measurements.

39. The computer readable medium of aspect 38, wherein the logic in (a) receives two or more measurement derived from a human subject wherein at least one measurement is an *in vitro* measurement of the level of marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject.

40. The computer readable medium of aspect 38, wherein the data processing system is used to compute risk of developing diabetes.

41. The computer readable medium of aspect 40, wherein the logic further receives measurements of blood glucose levels from said mammalian subject.

42. The computer readable medium of aspect 38, wherein the data processing system is used to compute risk of developing cardiovascular disease.

43. The computer readable medium of aspect 42, wherein the logic further receives measurements of blood pressure, HDL-cholesterol, LDL-cholesterol, total cholesterol and/or triglycerides.

44. The computer readable medium of aspect 38, wherein the data processing system is used to compute risk of developing cancer.

45. The computer readable medium of aspect 44, wherein the cancer is breast cancer, leukemia, prostate cancer or lung cancer.

46. The computer readable medium of aspect 38, wherein the data processing system is used to compute risk of mortality in a mammalian subject within 1 year, 3 years, 5 years, 10 years, 1 year-10 years, 1-5 years, 5-10 years.

47. The computer readable medium of aspect 46, wherein the logic further receives measurements of brain natriuretic peptide concentrations.

48. A computer program product, comprising the computer readable medium of aspect 38.

49. A computer for determining the risk and/or change in risk of developing a metabolic

syndrome-related disease and/or low-grade inflammation-related disease comprising

1. (a) a computer-readable data storage medium configured to store a data storage material encoded with computer-readable data, wherein said data comprises measurements derived from one of more samples from a mammalian subject, wherein at least one measurement is an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR;
2. (b) a storage medium having stored therein instructions for processing said computer-readable data;
3. (c) a central-processing unit responsive to said instructions coupled to said storage medium and to said computer-readable data storage medium for processing said computer-machine readable data into risk measurement and treatment progression values; and
4. (d) an output device coupled to said central-processing unit of (c) for outputting said risk and/or change in risk values.

50. The computer of aspect 49, wherein the computer-readable data storage medium (a) comprises data comprising two or more measurements derived from a human subject wherein at least one measurement is an *in vitro* measurement of the level of marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject.

51. The computer of aspect 49, wherein the computer is used to compute risk of developing diabetes.

52. The computer of aspect 49, wherein the computer is used to compute risk of developing diabetes and the data in the computer storage medium further comprises measurements of blood glucose levels from said mammalian subject.

53. The computer of aspect 49, wherein the computer is used to compute risk of developing cardiovascular disease.

54. The computer of aspect 49, wherein the computer is used to compute risk of developing cardiovascular disease and the data in the computer storage medium further comprises measurements of blood pressure, HDL-cholesterol, LDL-cholesterol, total cholesterol, C-reactive protein and/or triglycerides.

55. The computer of aspect 49, wherein the computer is used to compute risk of developing cancer (e.g., breast cancer, leukemia, prostate cancer or lung cancer).

56. The computer of aspect 49, wherein the computer is used to compute risk of mortality in a mammalian subject within 1 year, 3 years, 5 years, 10 years, 1 year-10 years, 1 year-5 years, 5 years-10 years.

57. The computer of aspect 49, wherein the computer is used to compute risk of mortality in a mammalian subject within 10 years and the data in the computer storage medium further

comprise measurements of brain natriuretic peptide concentrations.

58. A client computer for providing access to a risk computation system for determining the risk and/or change in risk of developing a metabolic syndrome-related disease and/or low-grade inflammation-related disease, the client computer comprising:

1. (a) an input device for receiving one or more measurements derived from a mammalian subject wherein at least one measurement is an *in vitro* measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject;
2. (b) a communication interface for communicating the received one or more measurements to a server computer configured to analyse said one or more measurements of step (a); and compute a risk value; and
3. (c) a communications interface for receiving, from the server computer a computed risk value.

59. The computer of aspect 58, wherein the input device (a) receives two or more measurements derived from a human subject wherein at least one measurement is an *in vitro* measurement of the level of marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject.

60. The computer of aspect 60, wherein said computer is used to compute risk of developing diabetes.

61. The computer of aspect 59, wherein said computer is used to compute risk of developing diabetes and wherein the input device (a) further receives measurements of blood glucose levels from a mammalian subject.

62. The computer of aspect 59, wherein said computer is used to compute risk of developing cardiovascular disease.

63. The computer of aspect 59, wherein said computer is used to compute risk of developing cardiovascular disease and wherein the input device (a) further receives measurements of blood pressure, HDL-cholesterol, LDL-cholesterol, total cholesterol and/or triglycerides.

64. The computer of aspect 59, wherein said computer is used to compute risk of developing cancer (e.g., breast cancer, prostate cancer, leukemia or lung cancer).

65. The computer of aspect 59, wherein said computer is used to compute risk of developing cancer (e.g., breast cancer, prostate cancer, leukemia or lung cancer) and wherein said the input device (a) further receives data relating to smoking and age.

66. The computer of aspect 59, wherein said computer is used to compute risk of mortality in a mammalian subject within 1-10 years.

67. The computer of aspect 60, wherein said computer is used to compute risk of mortality in a

mammalian subject within 1-10 years and the input device (a) may further receive measurements of brain natriuretic peptide concentrations.

68. A computer based method for determining the risk of developing a metabolic syndrome-related disease, low-grade inflammation-related disease (e.g., cancer) and/or risk of mortality in a mammalian subject comprising

1. (a) receiving under computer control in vitro measurement of the level of a marker in the form of (i) soluble urokinase plasminogen activator receptor (suPAR), and/or (ii) D2D3 cleavage products of suPAR in one or more samples derived from said subject;
2. (b) analyzing under computer control said in vitro measurement received in (a)
3. (c) calculating under computer control a risk value for developing a metabolic syndrome-related disease, low-grade inflammation-related disease (e.g., cancer) and/or risk of mortality in a mammalian subject within 1 year, 3 years, 5 years, 10 years or between 1-5 years, 1-10 years, or 5-10 years using analyzed measurements obtained in (b).

68. A method of reducing the risk of a healthy individual developing cardiovascular disease, chronic obstructive lung disease, eczema, obesity, autoimmune disease, cancer, metabolic syndrome or diabetes type 2 comprising the steps of:

1. (a) determining the level of suPAR in the individual; and
2. (b) deciding on the appropriate treatment for the patient at least partly in the light of the said level of suPAR.

69. A method of reducing the risk of a healthy individual developing cardiovascular disease, chronic obstructive lung disease, eczema, obesity, autoimmune disease, cancer, metabolic syndrome or type 2 diabetes comprising the step of reducing the plasma level of the individual's suPAR.

70. A method according to Aspect 70 wherein the plasma level is reduced to below 6 ng/ml.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2005116077A \[0008\]](#)

- [WO9012091A](#) [0069]

Non-patent literature cited in the description

- **SVENDSEN et al.** Surgical Infections, 2006, vol. 7, 463-471 [0007]
- **OSTROWSKI et al.** J. Acquir. Immune Defic. Syndr., 2004, vol. 35, 337-342 [0009]
- **OSTROWSKI et al.** Scand. J. Immunol., 2006, vol. 63, 478-486 [0009]
- **OSTROWSKI et al.** J. Acquir. Immune Defic. Syndr., 2005, vol. 39, 23-31 [0009]
- **MIZUKAMI et al.** Blood, 1995, vol. 86, 203-211 [0026]
- **KOLB et al.** Diabetologia, 2005, vol. 48, 1038-1050 [0046]
- **SIER CF et al.** Lab Invest, 1999, vol. 79, 717-722 [0050]
- **GRUNDY SM et al.** Circulation, 2005, vol. 112, 2735-2752 [0056] [0059]
- **KOHLER et al.** Nature, 1975, vol. 256, 495-497 [0070]
- **KOZBOR et al.** J. Immunol. Methods, 1985, vol. 81, 31-42 [0070]
- **COTE et al.** Proc. Natl. Acad. Sci. USA, 1983, vol. 80, 2026-2030 [0070]
- **COLE et al.** Mol. Cell Biol., 1984, vol. 62, 109-120 [0070]
- **BETTER et al.** Science, 1988, vol. 240, 1041- [0071]
- **SKERRA et al.** Science, 1988, vol. 240, 1038- [0071]
- **BIRD et al.** Science, 1988, vol. 242, 423- [0071]
- **HUSTON et al.** Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 5879- [0071]
- **WARD et al.** Nature, 1989, vol. 341, 544- [0071]
- **WINTERMILSTEIN** Nature, 1991, vol. 349, 293-299 [0071]
- Remington: The Science and Practice of Pharmacy Lippincott, Williams & Wilkins 2000 0000 [0074]
- **ANSEL et al.** Pharmaceutical Dosage Forms and Drug Delivery Systems Lippincott Williams & Wilkins Publishers 1999 0000 [0074]
- **ANDERSEN et al.** Metabolism, 2003, vol. 52, 1343-1353 [0095] [0095] [0099] [0099]
- **HAUGAARD et al.** Eur J. Endocrinol., 2005, vol. 152, 103-112 [0096]
- **SIER et al.** Lab Invest, 1999, vol. 79, 717-722 [0097] [0097] [0107]
- **MUSTJOKI et al.** Cancer Res, 2000, vol. 60, 7126-7132 [0097]
- **HAUGAARD et al.** Eur J Endocrinol, 2005, vol. 152, 103-112 [0099]
- **LIHN et al.** Am J Physiol Endocrinol Metab, 2003, vol. 285, E1072-E1080 [0099]
- **HOTHER-NIELSEN et al.** Metabolism, 1996, vol. 45, 82-91 [0100]
- **FRAYNJ** Appl Physiol, 1983, vol. 55, 628-634 [0100]
- **VOZAROVA B et al.** Diabetes, 2002, vol. 51, 455-461 [0148]
- **GRUNDY SM et al.** Circulation, 2005, vol. 112, 2735-2752 [0151]
- **XIONG et al.** Med. Decision Making, 2004, vol. 24, 659-69 [0163]

Patentkrav

1. Fremgangsmåde til diagnosticering af lavgrad inflammation i et tilsyneladende raskt pattedyrindivid, såsom et menneske-individ, omfattende:

5 (a) udføre en *in vitro*-måling af niveauet af en markør i form af opløselig urokinase-plasminogen-aktivatorreceptor (suPAR) og/eller D2D3-kløvningsprodukter af suPAR, i én eller flere prøver indeholdende en biologisk væske opnået fra individet, og

(b) anvende den ene eller flere målte værdier opnået som en faktor i diagnosen af lavgrad inflammation.

10

2. Fremgangsmåde til forudsigelse af risikoen for at udvikle en lavgrad-inflammation-relateret sygdom, i et pattedyr-individ, såsom et menneske-individ, omfattende at:

15 (a) udføre en *in vitro*-måling af niveauet af en markør i form af (i) opløselig urokinase-plasminogen-aktivatorreceptor (suPAR), og/eller (ii) D2D3 kløvningsprodukter af suPAR i én eller flere prøver indeholdende en biologisk væske opnået fra individet, og

(b) anvende den målte værdi opnået som en faktor til at forudse risikoen for at udvikle en lavgrad-inflammations-relateret sygdom.

20

3. Fremgangsmåde ifølge krav 1 eller 2, hvor den biologiske væske er blodserum, blodplasma eller urin.

25 **4.** Fremgangsmåde ifølge krav 1 eller 2, hvor mere end én af prøverne omfatter et biologisk væske der er blevet afledt fra individet ved mere end et tidspunkt.

5. Fremgangsmåde ifølge krav 4, yderligere sammenligne niveauet af markøren ved ét eller flere tidspunkter for at bedømme virkningen af et profylaktisk forløb og/eller behandling.

30

6. Fremgangsmåde ifølge krav 5, hvor det profylaktiske forløb omfatter fysisk motion.

7. Fremgangsmåde ifølge krav 5 eller 6, hvor et fald i niveauet af markøren på 5 mindst 10% og/eller 1 ng/ml angiver virkning.

8. Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor *in vitro*-målingen er et ELISA-assay.

10 **9.** Fremgangsmåde ifølge et hvilket som helst af de foregående krav, hvor fremgangsmåden yderligere omfatter at udføre en glucosetolerencetest, eller teste TNF-alfa, IL-6, leukocyttælling, fastende plasma-glucose eller fastende blod-glucose, blodtryk, kolesterol, C-reaktiv protein- og/eller triglyceridniveauer.

DRAWINGS




Figure 2A.




Figure 2B




Figure 2 C.




Figure 2 D




Figure 3




Figure 4. [upper plot and lower plot]




Figure 5A




Figure 5B




Figure 6A




Figure 6B




Figure 6C




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19





Figure 20



5

Figure 21




Figure 22




Figure 23
A




Figure 23
B




Figure 23
C




Figure 23
D




Figure 23
E




Figure 23
F

