

(19)



(11)

EP 3 094 654 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
11.03.2020 Bulletin 2020/11

(51) Int Cl.:
C07K 16/30 (2006.01) **A61P 35/00** (2006.01)
C07K 16/18 (2006.01) **A61K 39/00** (2006.01)

(21) Application number: **15701267.5**

(86) International application number:
PCT/US2015/010219

(22) Date of filing: **06.01.2015**

(87) International publication number:
WO 2015/108719 (23.07.2015 Gazette 2015/29)

(54) TARGETING CLPTM1L FOR TREATMENT AND PREVENTION OF CANCER

ABZIELUNG AUF CLPTM1L ZUR BEHANDLUNG UND PRÄVENTION VON KREBS

CIBLAGE DE CLPTM1L POUR LE TRAITEMENT ET LA PRÉVENTION DU CANCER

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(74) Representative: **J A Kemp LLP**
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(30) Priority: **14.01.2014 US 201461927330 P**

(43) Date of publication of application:
23.11.2016 Bulletin 2016/47

(73) Proprietor: **The Medical College of Wisconsin, Inc.**
Milwaukee, WI 53226 (US)

(72) Inventor: **JAMES, Michael, A.**
Big Bend, WI 53103 (US)

(56) References cited:

- **MICHAEL A. JAMES ET AL: "Functional Characterization of CLPTM1L as a Lung Cancer Risk Candidate Gene in the 5p15.33 Locus", PLOS ONE, vol. 7, no. 6, 4 June 2012 (2012-06-04), page e36116, XP055175210, DOI: 10.1371/journal.pone.0036116**
- **NIZHENHUA ET AL: "CLPTM1L Is Overexpressed in Lung Cancer and Associated with Apoptosis", PLOS ONE, vol. 7, no. 12, December 2012 (2012-12), XP055175473, cited in the application**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 3 094 654 B1

Description

[0001] This invention was made with government support under Grant No. U19 CA128147 awarded by the National Institutes of Health. The government has certain rights in the invention.

FIELD OF THE INVENTION

[0002] The present disclosure generally relates to compounds and compositions for targeting *CLPTM1L* polypeptides and methods for using such compounds and compositions to treat or prevent a cancer or other condition associated with *CLPTM1L* protein dysfunction (e.g., pathogenic production, modification, or function). In particular, the present disclosure relates to therapeutic agents comprising *CLPTM1L* targeting moieties and to methods of administering such therapeutic agents to a subject to treat or prevent a disease or condition associated with *CLPTM1L* protein dysfunction such as non-small cell lung cancer. Therapeutic agents, pharmaceutical compositions, and methods are described herein.

BACKGROUND

[0003] Cancer is a disease that begins with mutation of oncogenes and tumor suppressor genes. Mutation of these critical genes allows for a cancer cell to evolve and ultimately results in pathogenic replication (a loss of normal regulatory control leading to excessive cell proliferation) of various given types of cells found in the human body. Tumor formation, tumor survival, and cancer metastasis require anchorage-independent growth and protection from genotoxin-induced apoptosis and anoikis, a programmed cell death mechanism associated with detachment of tumor cells from an extracellular substrate. Tumor cells require protection from apoptosis and anoikis to invade surrounding tissue and to undergo metastasis.

[0004] James et al (2012) PLoS ONE 7(6):e36116 refers to functional characterization of *CLPTM1L* as a lung cancer risk candidate gene in the 5p15.33 locus. Zhenhua et al (2012) PLoS ONE 7(12):e52598 refers to *CLPTM1L* being overexpressed in lung cancer and associated with apoptosis.

[0005] There remains a need in the art for methods for treating or preventing cancer and, in particular, for methods which slow or curb tumor growth and prevent metastasis.

BRIEF SUMMARY OF THE INVENTION

[0006] The present invention provides a *CLPTM1L*-targeting agent, wherein the *CLPTM1L*-targeting agent is a monoclonal antibody that binds specifically to a peptide having an amino acid sequence of SEQ ID NO:6 or SEQ ID NO:10. In a first aspect, described herein is a compound comprising a *CLPTM1L*-targeting agent selected from the group consisting of an antibody, a peptide, an aptamer, or a fragment thereof. The *CLPTM1L*-targeting agent is a monoclonal antibody specific for at least a portion of a *CLPTM1L* polypeptide. The *CLPTM1L* polypeptide can have the amino acid sequence set forth in SEQ ID NO:2 or a portion thereof. The monoclonal antibody can bind specifically to a peptide having an amino acid sequence selected from the group consisting of SEQ ID NO:4-25. The monoclonal antibody can be a monoclonal anti-*CLPTM1L* clone selected from the group consisting of 6-1, 6-2, 10-1, 10-2, and 10-3. The monoclonal antibody can be specific to a *CLPTM1L* polypeptide having one or more of the mutations set forth in Table 2. In some cases, the *CLPTM1L*-targeting agent is a peptide having an amino acid sequence selected from the group consisting of SEQ ID NO:4-25.

[0007] In another aspect, described herein is a pharmaceutical composition comprising a compound provided herein and a pharmaceutically acceptable carrier.

[0008] In yet another aspect, described herein is an agent according to a compound or composition provided herein for use as a medicament. Further described is use of such an agent in the manufacture of a medicament.

[0009] In a further aspect, described herein is an article of manufacture comprising a pharmaceutical composition of the invention and instructions for administration to a human subject.

[0010] In another aspect, described herein is a method for treating or preventing a tumor in a subject. The method typically comprises administering a therapeutically effective amount of a compound comprising a *CLPTM1L*-targeting agent to a subject in need thereof, whereby the tumor is treated or prevented in the subject. In some cases, the *CLPTM1L*-targeting agent is selected from the group consisting of an antibody, a peptide, and aptamer, or a fragment thereof. The *CLPTM1L*-targeting agent can be a monoclonal antibody specific for at least a portion of a *CLPTM1L* polypeptide. The *CLPTM1L* polypeptide can have the amino acid sequence set forth in SEQ ID NO:2 or a portion thereof. The monoclonal antibody can be specific to a peptide having an amino acid sequence selected from the group consisting of SEQ ID NO:4-25. The monoclonal antibody can be specific to a *CLPTM1L* polypeptide having one or more of the mutations set forth in Table 2. The *CLPTM1L*-targeting agent can be a peptide having an amino acid sequence selected from the group consisting of SEQ ID NO:4-25.

[0011] In some cases, the tumor is a solid tumor selected from the group consisting of glioblastoma, sarcoma, carcinoma, and lymphoma. In some cases, the tumor is associated with a cancer or pre-neoplastic lesion selected from the group consisting of lung cancer, pancreatic cancer, prostate cancer, skin cancer, bladder cancer, kidney cancer, ovarian cancer, colon cancer, colorectal cancer, breast cancer, cervical cancer, brain cancer, esophageal cancer, and stomach cancer, or a pre-neoplastic lesion thereof. The tumor can be associated with lung cancer or a pre-neoplastic lesion thereof. The tumor can exhibit resistance to a chemotherapeutic agent. The chemotherapeutic agent can be selected from the group consisting of cisplatin and gemcitabine. In some cases, the compound is administered with a pharmaceutically acceptable carrier.

[0012] In yet another aspect, described herein is a method of treating or preventing a disease or condition associated with over-expression or inappropriate expression of a nucleic acid sequence encoding CLPTM1L in a subject. The method typically comprises administering a compound comprising a CLPTM1L-targeting agent to a cell, tissue, or organ of a subject at risk of, diagnosed as having, or exhibiting a symptom of the disease or condition. The disease or condition can be a cancer, tumor, (e.g., solid tumor) or a pre-neoplastic lesion. The tumor, cancer, or pre-neoplastic lesion can be selected from the group consisting of lung cancer, pancreatic cancer, glioblastoma, prostate cancer, skin cancer, bladder cancer, kidney cancer, ovarian cancer, colon cancer, colorectal cancer, breast cancer, cervical cancer, brain cancer, esophageal cancer, stomach cancer, lymphoma, chronic leukemia, and acute leukemia. The tumor can exhibit resistance to a chemotherapeutic agent. The chemotherapeutic agent can be selected from the group consisting of cisplatin and gemcitabine. The CLPTM1L-targeting agent can be selected from the group consisting of an antibody, a peptide, an aptamer, or a fragment thereof. The CLPTM1L-targeting agent can be a monoclonal antibody specific for at least a portion of a CLPTM1L polypeptide. The CLPTM1L polypeptide can have the amino acid sequence set forth in SEQ ID NO:2 or a portion thereof. In some cases, the monoclonal antibody is specific to a peptide having an amino acid sequence selected from the group consisting of SEQ ID NO:4-25. The monoclonal antibody can be specific to a CLPTM1L polypeptide having one or more of the mutations set forth in Table 2. The CLPTM1L-targeting agent can be a peptide having an amino acid sequence selected from the group consisting of SEQ ID NO:4-25. The compound can be administered with a pharmaceutically acceptable carrier.

[0013] These and other features, aspects, and advantages of the present invention will become better understood from the description that follows. In the description, reference is made to the accompanying drawings, which form a part hereof and in which there is shown by way of illustration, not limitation, embodiments of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0014] The present invention will be better understood and features, aspects and advantages other than those set forth above will become apparent when consideration is given to the following detailed description thereof. The detailed description makes reference to the following drawings, wherein:

FIG. 1 presents structural data demonstrating the presence of two separate globular domains separated by a small disordered region (amino acid residues 142-162) within the larger extracellular region.

FIG. 2 presents (A) a summary of CLPTM1L predicted structure including orientation of the human sequence of CLPTM1L within the plasma membrane, two extracellular globular domains (ECR1 and ECR2), the interstitial disordered region, transmembrane regions, and identified glycosylated cell surface peptides (in blue) and predicted sites N-glycosylation sites (green). Monoclonal antibodies generated against CLPTM1L peptides targeting each ECR are indicated in parentheses. Image was generated using the Protter application (Omasits et al., Bioinformatics 30(6):884-6 (2014)). (B) A splicing variant of CLPTM1L, lacking extracellular region 1 and some transmembrane domains changing the predicted structure. The expression pattern of this isoform is not known.

FIG. 3 depicts selection of epitopes to raise monoclonal antibodies specific for human CLPTM1L protein. The focus region for monoclonal antibody design included the ordered regions of the predicted extracellular domain. Glycosylation sites were excluded.

FIG. 4 demonstrates CLPTM1L localization. (A) Western blotting of cell surface and whole cell protein lysates for CLPTM1L, transmembrane receptor EGFR and nuclear protein Cyclin D1. Monoclonal antibodies 6-1, 10-2, and 10-3 were used for western blotting. (B) Immunofluorescent staining of non-permeabilized A549 lung tumor cells using DAPI dye to stain cell nuclei and polyclonal antibody to the predicted extracellular N-terminus of human CLPTM1L followed by a secondary FITC conjugated antibody to detect CLPTM1L. A non-specific antibody was used as a control. Staining indicates punctuate plasma membrane localization of the protein. (C) Western blotting for CLPTM1L in H838 cells with monoclonal antibodies. (D) Immunofluorescent staining of non-permeabilized A549 cells with monoclonal CLPTM1L antibodies 6-1 (1000X) and 10-2 (400X). Monoclonal antibodies 6-1 and 10-2 were

used for immunofluorescence.

FIG. 5 presents graphs and images demonstrating that CLPTM1L expression is increased in pancreatic tumor tissue over normal pancreatic ductal epithelia and that higher expression is correlated with poor survival. (A) Immunohistochemical scoring of staining with antibody specific to CLPTM1L in pancreatic tumor vs. normal ductal epithelial tissues in 31 patients. (B) Kaplan-Meier survival curves stratified into tertiles of CLPTM1L expression. (C) Representative staining of tumor.

FIG. 6 demonstrates that targeting CLPTM1L in tumor cells modulates Bcl-xL and Akt survival signaling and sensitizes cells to killing with cisplatin. (A) Treating A549 cells with polyclonal anti-CLP antibody (1:1000 = 1 μ g/mL) reduced accumulation of anti-apoptotic molecules Bcl-xL and total Akt. (B) Anti-CLPTM1L significantly increased sensitivity to killing upon cisplatin treatment for 24 hours ($p < 0.05$). (C) Treatment with monoclonal anti-CLP (10-2) increased sensitivity of pancreatic adenocarcinoma cells to killing by gemcitabine (500 μ M for 72 hours). (D) Treatment of pancreatic adenocarcinoma cells with monoclonal anti-CLP (10-2) reduced accumulation of phosphorylated Akt.

FIG. 7 presents images and graphs demonstrating that treatment with anti-CLP antibody (Ab) inhibits anchorage-independent growth of human lung tumor and bronchial epithelial cells *in vitro*. Anchorage-independent growth of human lung tumor cells, as measured by colony formation in 0.4% top agar over a 0.8% bottom agar layer, was inhibited by 55%-65% by anti-CLPTM1L treatment compared to solvent control treated cells ($p < 0.05$). Treatments were administered one day after plating and again one week later. EGF treatment induced anchorage independent growth in Beas2B human bronchial epithelial cells, which was inhibited by 76% upon treatment with anti-CLP Ab.

FIG. 8 demonstrates colony growth inhibition of pancreatic tumor cells by anti-CLPTM1L antibodies. (A) Panc-1 pancreatic tumor cells were plated on 6-well plates in triplicate at 1000 cells per well in DMEM/F12 w/ 10% FBS. After plating, the indicated antibodies were added to the culture media at 5 μ g/mL. (B) Colonies per well were counted after 5 days in culture following fixation in methanol and staining with 0.5% crystal violet. ** $p = 0.005$.

FIG. 9 presents xenograft models for inhibition of the growth of established human lung tumors. (A) Mean tumor volume in athymic nude mice injected subcutaneously with A549 tumor cells and treated with control ascites, polyclonal anti-CLPTM1L, or monoclonal anti-CLPTM1L clones after tumor establishment on days 0, 5, and 12. Error bars represent standard error of the mean. (B) Tumor volumes of individual tumors at day 17. Error bars represent standard deviations. * $p < 0.05$, ** $p < 0.005$.

FIG. 10 summarizes characterization of *in vitro* anti-tumor effects of monoclonal anti-CLPTM1L antibodies.

DETAILED DESCRIPTION OF THE INVENTION

[0015] The present invention is based at least in part on the Inventors' discovery that CLPTM1L is commonly over-expressed in non-small cell lung cancer (NSCLC) and that it protects lung tumor cells from genotoxic apoptosis. In particular, depletion of CLPTM1L robustly and significantly inhibited tumorigenesis *in vivo* in multiple models. Cisplatin Resistance Related Protein-9 (CRR9), otherwise known as Cleft-Lip and Palate Transmembrane Protein-Like Protein 1 (CLPTM1L), is located at chromosome 5p15.33 as defined by multiple Genome Wide Association (GWA) studies. The CLPTM1L gene lies within a locus on chromosome 5 that is frequently gained in copy number early in lung cancer and that is associated by genotype with lung cancer susceptibility. For example, genetic variants near and within the CLPTM1L gene are associated with lung cancer, cervical cancer, ovarian cancer, pancreatic cancer, bladder cancer, glioma, prostate cancer, basal cell carcinoma, and melanoma. Although the mechanism of action remains to be fully elucidated, it is believed that CLPTM1L is involved in Bcl-xL survival protein accumulation. Indeed, the Inventors previously demonstrated that CLPTM1L protects chemotherapeutically treated tumor cells from genotoxin-induced apoptosis and that CLPTM1L is required for anchorage independent growth and for Ras-driven lung tumorigenesis.

[0016] The Inventors further discovered that CLPTM1L localizes to the plasma membrane, and that polyclonal antibodies targeting the N-terminal region of the CLPTM1L protein, which is predicted to be extracellular, elicit comparable phenotypes to those elicited by RNA interference-mediated depletion of CLPTM1L; specifically, Bcl-xL and Akt inhibition, chemosensitization, and anchorage dependence. Accordingly, the present invention relates to immunoglobulins, compositions, and methods of using such compositions for targeting human CLPTM1L for the prevention or treatment of solid tumors and cancers and for chemosensitization of tumor cells.

[0017] As used herein, the term "solid tumor" refers to an abnormal mass of tissue that usually does not contain cysts or liquid areas. Non-cancerous tumors are described as "benign," while cancerous tumors are described as "malignant." Different types of solid tumors are named for the particular cells that form them, for example, sarcomas formed from

connective tissue cells (such as bone cartilage, fat), carcinomas formed from epithelial tissue cells (such as breast, colon, pancreas), and lymphomas formed from lymphatic tissue cells (such as lymph nodes, spleen, thymus). Treatment of all types of solid tumors is within the scope of this invention.

[0018] The human CLPTM1L gene (SEQ ID NO:1; Genbank ID AAH25305.1) encodes a 538 amino acid polypeptide (SEQ ID NO:2; UniProt ID: Q96KA5.1). Gene products of human *CLPTM1L* include a primary mRNA (Genbank ID BC016399.1; SEQ ID NO:3) and two additional predicted transcript splice variants (Ensembl transcript IDs: ENST00000320927 and ENST00000507807). These predicted splice variants encode proteins of 502 and 369 amino acids, respectively. Splice variant prediction methods are available at useast.ensembl.org/info/docs/genbuild/genome_annotation.html on the World Wide Web.

Compositions of the Invention

[0019] In one aspect, described herein are compositions comprising CLPTM1L-targeting agent having specificity for at least a portion of a CLPTM1L polypeptide. As described herein, compositions of the present invention are useful as therapeutic agents and pharmaceutical compositions for a variety of clinical applications. In particular, described herein are therapeutic compositions comprising one or more CLPTM1L-targeting agent including antibodies, antibody fragments, immunoconjugates, bispecific antibodies, trispecific antibodies, and chimeric antigen receptors having specificity for (*i.e.*, targeted to) one or more CLPTM1L epitopes. Also described are compositions comprising one or more CLPTM1L-targeting peptides. As used herein, a CLPTM1L-targeting agent is a molecule or complex that specifically binds to a CLPTM1L polypeptide or to one or more epitopes of a CLPTM1L polypeptide or fragment thereof and/or competes for binding at one or more epitopes of a CLPTM1L polypeptide or fragment thereof.

[0020] As used herein, the term "polypeptide" is intended to encompass a singular "polypeptide" as well as plural "polypeptides," and refers to a molecule composed of monomers (amino acids) linearly linked by amide bonds (also known as peptide bonds). The term "polypeptide" refers to any chain or chains of two or more amino acids, and does not refer to a specific length of the product. Thus, peptides, dipeptides, tripeptides, oligopeptides, "protein," "amino acid chain," or any other term used to refer to a chain or chains of two or more amino acids, are included within the definition of "polypeptide," and the term "polypeptide" may be used instead of, or interchangeably with, any of these terms. The term "polypeptide" is also intended to refer to the products of post-expression modifications of the polypeptide, including without limitation glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, or modification by non-naturally occurring amino acids. A polypeptide may be derived from a natural biological source or produced by recombinant technology, but is not necessarily translated from a designated nucleic acid sequence. It may be generated in any manner, including by chemical synthesis.

[0021] As used herein, the terms "antibody" and "antibodies" are synonymous with "immunoglobulin" and "immunoglobulins," and the terms are used interchangeably herein. The terms "antibody" and "antibodies" include whole immunoglobulins including, without limitation, polyclonal antibodies or monoclonal antibodies (mAbs). The term antibody encompasses various broad classes of polypeptides that can be distinguished biochemically. Those skilled in the art will appreciate that heavy chains are classified as gamma, mu, alpha, delta, or epsilon (γ , μ , α , δ , ϵ) with some subclasses among them (*e.g.*, $\gamma 1$ - $\gamma 4$). It is the nature of this chain that determines the "class" of the antibody as IgG, IgM, IgA, IgG, or IgE, respectively. The immunoglobulin subclasses (isotypes) *e.g.*, IgG₁, IgG₂, IgG₃, IgG₄, IgG₅, etc. are well characterized and are known to confer functional specialization.

[0022] Antibodies appropriate for the present invention also include antibody fragments or modified products thereof, provided that they can be suitably used in the present invention. Appropriate antibody fragments comprise at least one variable domain of an immunoglobulin, such as single variable domains Fv (Skerra & Pluckthun, Science 240:1038-41 (1988)), scFv (Bird et al., Science 242:423-26 (1988); Huston et al., Proc. Natl. Acad. Sci. USA 85:5879-83 (1988)), Fab, (Fab')₂ or other proteolytic fragments. The terms "antibody" and "antibodies" further include chimeric antibodies; human and humanized antibodies; recombinant and engineered antibodies, conjugated antibodies, and fragments thereof. Humanized antibodies are antibodies wherein the complementarity determining regions (CDRs) of an antibody from a mammal other than human (*e.g.*, a mouse antibody) are transferred into the CDRs of human antibodies. Chimeric and humanized antibodies can be made according to standard protocols such as those disclosed in U.S. Pat. No. 5,565,332. Other antibody formats are described in, for example, "Antibody Engineering," McCafferty et al. (Eds.) (IRL Press 1996). Also encompassed in the invention are CLPTM1L-targeting immunoglobulins that have been conjugated or bound in some manner to various molecules including, without limitation, polyethylene glycol (PEG), radioactive substances, and drugs. Such conjugated antibodies can be obtained by chemically modifying a CLPTM1L-targeting immunoglobulin. Methods for obtaining conjugated antibodies are known and available in the art.

[0023] The antibodies described herein may be polyclonal or monoclonal antibodies. Preferably, the CLPTM1L-targeting immunoglobulins are monoclonal. Methods of producing polyclonal and monoclonal antibodies are known in the art and described generally, *e.g.*, in U.S. Pat. No. 6,824,780. Typically, the antibodies of the present invention are produced recombinantly, using vectors and methods available in the art, as described further below. Human antibodies

may also be generated by *in vitro* activated B cells (see U.S. Pat. Nos. 5,567,610 and 5,229,275). Human antibodies may also be produced in transgenic animals (e.g., mice) that are capable of producing a full repertoire of human antibodies in the absence of endogenous immunoglobulin production. For example, it has been described that the homozygous deletion of the antibody heavy-chain joining region (J_H) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array into such germ-line mutant mice results in the production of human antibodies upon antigen challenge. See, e.g., Jakobovits et al., Proc. Natl. Acad. Sci. USA, 90:2551 (1993); Jakobovits et al., Nature, 362:255-258 (1993); Bruggemann et al., Year in Immuno., 7:33 (1993); U.S. Pat. Nos. 5,545,806, 5,569,825, 5,591,669; U.S. Pat. No. 5,545,807; and WO 97/17852. Such animals may be genetically engineered to produce human antibodies comprising a polypeptide described herein.

[0024] The source of the antibodies described herein is not particularly restricted in the present invention; however, the antibodies are preferably derived from mammals, and more preferably derived from humans. Monoclonal antibodies appropriate for the present invention can be prepared by standard hybridoma methods. For example, standard hybridoma methods employ differential binding assays to ensure that the resulting monoclonal antibodies are specific for a CLPTM1L polypeptide and do not show cross-reactivity between related proteins. Alternatively, monoclonal antibodies appropriate for the present invention can be prepared using antibody engineering methods such as phage display. Methods for obtaining highly specific antibodies from antibody phage display libraries are known in the art, and several phage antibody libraries are commercially available from, for example, MorphoSys (Martinsried, Germany), Cambridge Antibody Technology (Cambridge UK) and Dyax (Cambridge Mass.). Suitable phage display methods are described, for example, in U.S. Pat. Nos. 6,300,064 and 5,969,108, and in "Antibody Engineering," McCafferty et al. (Eds.) (IRL Press 1996)). Once the antibody heavy and light chain genes are recovered from the phage antibodies, antibodies in any suitable format may be prepared for use according to the present invention, e.g., whole antibodies, Fab fragments, scFv, etc.

[0025] Antibodies disclosed herein may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine, donkey, rabbit, goat, guinea pig, camel, llama, horse, or chicken antibodies.

[0026] Polyclonal antibodies can be prepared by may also be prepared using traditional animal-based methods. For example, an appropriate animal can be immunized using a polypeptide immunogen (e.g., peptide of CLPTM1L). Polypeptide antibody titers in the immunized animal can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized polypeptide. Antibodies specific to the antigen can be isolated from the mammal (e.g., from the blood) and further purified by techniques known to those practicing in the art including, for example, protein A chromatography to obtain the IgG fraction. In some cases, at an appropriate time after immunization (e.g., when the antibody titers are highest) antibody-producing cells can be obtained from the animal and used to prepare monoclonal antibodies.

[0027] As used herein, the terms "epitope" or "antigenic determinant" refer to a site on an antigen (e.g., on CLPTM1L) to which an immunoglobulin or antibody specifically binds. Generally, an epitope includes at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 consecutive or non-consecutive amino acids in a unique spatial conformation. See, e.g., Epitope Mapping Protocols in Methods in Molecular Biology, Vol. 66, G. E. Morris, Ed. (1996). As used herein, the terms "specific binding," "selective binding," "selectively binds," and "specifically binds," refer to an antibody binding to an epitope via its antigen-binding domain, and that the binding entails some complementarity between the antigen-binding domain and the epitope. Generally, an antibody specifically or selectively binds with an affinity (generally represented by the dissociation constant K_D) of approximately less than 10^{-7} M, such as approximately less than 10^{-8} M, 10^{-9} M, or 10^{-10} M, or lower. As used herein, the term "affinity" denotes the strength of the sum total of non-covalent interactions between a single binding site of a molecule (e.g., a peptide, polypeptide, or antibody) and its binding partner (e.g., a target or an antigen). Unless indicated otherwise, as used herein, "binding affinity" refers to intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (e.g., between a peptide and its target, or between an antibody and its antigen). The terms " K_D " and " K_d " are synonymous and refer to the dissociation equilibrium constant of a particular molecule X-binding partner Y interaction. Affinities of antibodies can be readily determined using methods known in the art such as surface plasmon resonance. Other conventional techniques for determining antibody affinities are known in the art, such as those described by Scatchard et al. (Ann. N.Y. Acad. Sci. USA 51:660 (1949)). Binding properties of an antibody to antigens, cells or tissues thereof may generally be determined and assessed using immunodetection methods including, for example, immunofluorescence-based assays, such as immuno-histochemistry (IHC), and/or fluorescence-activated cell sorting (FACS).

[0028] In exemplary embodiments, antibodies of the present invention bind to CLPTM1L with a dissociation equilibrium constant (K_D) of less than approximately 10^{-7} M, such as approximately less than 10^{-8} M, 10^{-9} M, or 10^{-10} M, or lower.

[0029] Table 1 presents amino acid sequences of epitopes useful for producing CLPTM1L-targeted monoclonal antibodies. The "start" and "end" positions are numbered relative to numbered relative to the amino acid sequence of human CLPTM1L set forth as SEQ ID NO:2 (UniProt ID: Q96AK5-1).

Table 1. Peptides and Epitopes For Monoclonal Antibody Production

Start position	End position	Epitope sequences	SEQ ID NO:
49	60	RRPKLQLSVYTT	4
68	79	ENNIDLVLNVED	5
78	89	EDFDVESKFERT	6
112	123	HAGVLPWHDGKQ	7
186	197	DGSSLPADVHRY	8
195	206	HRYMKMIQLGKT	9
233	244	TELPLTVSYDKV	10
263	274	QQFGFSEKDADE	11
33	42	TRPCSGDANC	12
49	60	RRPKLQLSVYTT	13
68	78	ENNIDLVLNVE	14
80	89	FDVESKFERT	15
112	123	HAGVLPWHDGKQ	16
131	140	TTYMVPKPEE	17
141	150	INLLTGESDT	18
147	156	ESDTQQIEAE	19
157	167	KKPTSALDEPV	20
186	195	DGSSLPADVH	21
195	206	HRYMKMIQLGKT	22
235	244	LPLTVSYDKV	23
255	265	MQDAVYSLQQF	24
266	274	GFSEKDADE	25

[0030] As described in greater detail in the Examples section, epitopes presented in the shaded boxes of Table 1 were initially selected for monoclonal antibody production using the following criteria: (a) epitopes that reside within the predicted surface-exposed globular domains of CLPTM1L (e.g., residues at positions 32-284); (b) epitopes that avoid predicted glycosylation sites (e.g., residues at positions 91, 101, and 229); (c) epitopes that avoid the predicted disordered region (e.g., positions 141-162); and (d) epitopes having amino acid sequences that score highly on the basis of hydrophilicity, folding potential, and antigen presentation.

[0031] Monoclonal antibodies can be obtained by hybridoma technology, which is the process of producing hybrid cell lines by fusing an antibody-producing B cell with a myeloma cell that can grow in tissue culture. The resulting hybridoma line produces a monoclonal antibody of a single specificity.

[0032] In exemplary aspects, a composition described herein comprises a CLPTM1L-specific immunoglobulin (e.g., polyclonal or monoclonal antibody) that is chemically or structurally modified, conjugated, chimerized, humanized, or otherwise engineered antibody. For example, compositions of the invention include a CLPTM1L-specific antibody modified as an immunoconjugate. As used herein, the term "immunoconjugate" refers to a therapeutic agent that comprises (1) an antibody that binds to an antigen (e.g., a cancer cell antigen) with high specificity, (2) an effector molecule, and (3) a linker. The effector molecule can be selected from the group consisting of therapeutic agents, prodrugs, peptides, proteins, enzymes, viruses, lipids, biological response modifiers, pharmaceutical agents, and polyethylene glycol (PEG). Preferably, the effector molecule has anti-cancer cell activity. See, e.g., Smaglo et al., Nat. Rev. Clin. Oncol. 11(11):637-48 (2014).

[0033] Appropriate linkers include, without limitation, can be a thioester bond, a disulfide bond, a hydrazone bond, or a peptide. Preferably, the linker operates to ensure that the effector does not separate from the antibody during transit and will reliably release the effector to a targeted cancer cell or tumor stroma. See, for review, Smaglo et al., Nature Reviews Clin. Oncol. 11:637-48 (2014). Without being bound by any particular mechanism or mode of action, it is believed that immunoconjugates are particularly effective anticancer agents because they possess the potent anticancer effects of a therapeutic agent (e.g., chemotherapeutic) as well as the highly specific cancer targeting properties of a cancer cell antigen-specific monoclonal antibody. In exemplary aspects, immunoconjugates of the present invention comprise a

therapeutic agent selected from a pharmacologic agent, radioisotope, and toxin; a CLPTM1L-specific monoclonal or polyclonal antibody moiety; and a linker.

[0034] In some cases, a CLPTM1L-specific immunoglobulin of the present invention is a bispecific monoclonal antibody (BsMAb, BsAb). As used herein, the terms "bispecific monoclonal antibody," "BsMAb," and "BsAb" are used interchangeably and refer to an engineered immunoglobulin comprising two different monoclonal antibodies or fragments thereof. Accordingly, bispecific antibodies are antibodies that have binding specificities for at least two different epitopes. Exemplary bispecific antibodies may bind to two different epitopes of a single antigen. Other such antibodies may combine a first antigen binding site with a binding site for a second antigen. For example, a BsMAb can be engineered to simultaneously bind an immune cell (e.g., a T cell, a B cell, a monocyte, a macrophage, a neutrophil, a dendritic cell, a phagocyte, a natural killer cell, an eosinophil, a basophil, a mast cell) and a cancer cell antigen (e.g., CLPTM1L) to target and kill the cancer cell.

[0035] A bispecific antibody of the invention can further comprise a human or a humanized Fc fragment such as a human IgG Fc fragment. CLPTM1L-specific immunoglobulins can be humanized or subjected to Fc modification according to any appropriate methodology. As used herein, the term "humanized antibody" generally refers to a human antibody that has one or more amino acid residues introduced into it from a source that is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization is traditionally performed following the method of Winter and co-workers (Jones et al., *Nature*, 321:522-525 (1986); Reichmann et al., *Nature*, 332:323-327 (1988); Verhoeven et al., *Science*, 239:1534-1536 (1988)), by substituting import hypervariable region sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567) wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g., F(ab')₂ bispecific antibodies). WO 96/16673 describes a bispecific anti-ErbB2/anti-FcγRIII antibody and U.S. Pat. No. 5,837,234 discloses a bispecific anti-ErbB2/anti-FcγRI antibody. A bispecific anti-ErbB2/Fcα antibody is shown in WO98/02463. U.S. Pat. No. 5,821,337 teaches a bispecific anti-ErbB2/anti-CD3 antibody. Traditional production of full length bispecific antibodies is based on the co-expression of two immunoglobulin heavy chain-light chain pairs, where the two chains have different specificities (Millstein et al., *Nature*, 305:537-539 (1983)).

[0036] As used herein, the term "Fc" fragment refers to the carboxy-terminal portions of two heavy (H) chains held together by disulfide bonds. The effector functions of antibodies are determined by sequences in the Fc region, which region is also the part recognized by Fc receptors (FcR) found on certain types of cells. Fc regions can be modified to improve or reduce binding to effector regions. See, e.g., Vincent and Zurini, *Biotechnol. J.* 7(12):1444-50 (2012). Other examples of Fc engineering include modifying the half-life of immunoglobulin G (IgG). For example, antibody fragments having an increased *in vivo* half-life comprising a salvage receptor binding epitope residues are described in U.S. Pat. No. 5,869,046. Furthermore, engineered pH-dependent antigen binding can be applied to enhance the recycling of IgG via FcRn, enabling binding to additional target molecules. For review, see Kaneko and Niwa, *Biodrugs* 25(1):1-11 (2011). Other techniques for the production of antibody fragments will be apparent to the skilled practitioner.

[0037] Without being bound by any particular mechanism or mode of action, it is believed that bispecific monoclonal antibodies exhibit increased antibody-dependent cell-mediated cytotoxicity (Chames et al., *mAbs* 1:6, 539-547 (2009)). Accordingly, an effective dose of a CLPTM1L-targeted bispecific monoclonal antibody of the present invention can be lower than the effective dose of a conventional monoclonal CLPTM1L-specific antibody. In some cases, an effective dose of such a bispecific antibody is one or more orders of magnitude lower than that of a conventional monoclonal antibody. For example, an effective dose of a CLPTM1L-targeted bispecific monoclonal antibody can be around 0.01 mg·m⁻²·d⁻¹ (milligrams per square meter body surface area per day).

[0038] Bispecific antibodies may also be used to localize cytotoxic agents to cancer cells. These antibodies possess a cancer/tumor-binding arm and an arm that binds a cytotoxic agent (e.g., saporin, anti-interferon-α, vinca alkaloid, ricin A chain, methotrexate, or radioactive isotope hapten).

[0039] Bispecific antibodies also include antibodies having more than two valencies. For example, bispecific antibodies include trifunctional hybrid antibodies which, as used herein, are antibodies capable of inducing antibody-mediated cytotoxicity (via interactions with tumor-associated antigens on target cancer cells) and T-cell mediated cytotoxicity (via recognition of cytotoxic T lymphocytes) as well as anti-tumor immunologic memory (via interactions of a Fc region with markers on accessory cells such as macrophages). See Tutt et al., *J. Immunol.* 147: 60 (1991). For example, bi- and tri-specific antibodies against CD16, CD19, and CD22 have been shown to activate natural killer cell activation against B-cell leukemia (Gleason et al., *Mol. Cancer Ther.* 11(12):2674-84 (2012)).

[0040] Tri-specific antibodies (also known as trifunctional hybrid antibodies) comprise light and heavy chains originating from parental mouse IgG2a and rat IgG2b monoclonal antibodies, each having different antigen binding properties that provide additive tumor killing capabilities through the efficient recruitment of macrophages and NK cells and efficient co-stimulation of T cells through direct contact with accessory cells such as macrophages. By forming a tri-cell complex comprising a tumor cell, T cell, and accessory cell, trifunctional hybrid antibodies induce coordinated signaling events

required for efficient tumor cell destruction. Among those coordinated signaling events are immune effector mechanisms regulating destructive processes such as phagocytosis and perforin-mediated necrosis. For example, tri-specific antibodies targeting chorionic embryonic antigen (CEA) and the T-cell co-receptors CD3 and CD28 have been shown to recruit T lymphocytes to kill CEA positive tumors (Wang et al., J Biochem. 135(4):555-65 (2004)).

[0041] In some embodiments, antibodies of the invention are modified with respect to effector function, e.g., so as to enhance antigen-dependent cell-mediated cytotoxicity (ADCC) and/or complement dependent cytotoxicity (CDC) of the antibody. This may be achieved by introducing one or more amino acid substitutions in an Fc region of the antibody. Alternatively or additionally, cysteine residue(s) may be introduced in the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and ADCC. See Caron et al., J. Exp Med. 176:1191-1195 (1992). Homodimeric antibodies with enhanced anti-infection activity may also be prepared using heterobifunctional cross-linkers as described in Wolff et al., Cancer Research 53:2560-2565 (1993). Alternatively, an antibody can be engineered which has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., Anti-Cancer Drug Design 3:219-230 (1989).

[0042] In other cases, a CLPTM1L-specific antibody of the present invention is embodied in a chimeric antigen receptor T cell. As used herein, the term "chimeric antigen receptor T cell" refers to a genetically engineered antibody-T cell chimera that comprises a chimeric antigen receptor (CAR). As described herein, chimeric antigen receptor T cells of the present invention are therapeutic agents having the antigen specificity of a CLPTM1L-specific antibody and the poly-functionality and potency of cellular immunity. Techniques for chimeric antigen receptor T cell therapies are known and available in the art. See, e.g., Kenderian et al., Cancer Res. 74(22):6383-9 (2014).

[0043] In exemplary embodiments, CLPTM1L-specific immunoglobulins provided herein are modified to possess altered binding properties or increased affinity to the target antigen by, for example, affinity improvement or affinity maturation. As used herein, the term "affinity-matured antibody" refers to an antibody or a fragment thereof with one or more amino acid substitutions in a variable region, which results in improved affinity of the antibody for an antigen, as compared to a parent (starting) antibody which does not possess those substitutions. Antibody affinity enhancement techniques are known and available in the art. For example, random mutagenesis (Groves et al., J. Immunol. Methods, 313:129-39, 2006) and site-directed mutagenesis methods (Barbas et al., Proc. Natl. Acad. Sci. USA, 91:3809-13, 1994) are known. In some cases, these mutagenesis methods are combined with, for example, *in vitro* display-based technologies such as phage or ribosome display to generate libraries of variants for subsequent screens (Almagro and Strohl, Antibody Engineering: Humanization, Affinity Maturation and Selection Methods. 307-327. In: Therapeutic Monoclonal Antibodies: From Bench to Clinic. Ed. Zhiqiang An. John Wiley & Sons, Inc. 2009). Preferably, a modified antibody obtained by affinity improvement or affinity maturation as described herein will exhibit 2-fold, 5-fold, 10-fold, 100-fold, 200-fold, 300-fold, 400-fold, 500-fold, 600-fold, 700-fold, 800-fold, 900-fold, 1000-fold, 2000-fold, 3000-fold, 4000-fold, 5000-fold, 10000-fold or more improved activity for the target antigen compared to a starting antibody. For example, Kaneko and Zurini review optimizing therapeutic antibodies.

[0044] In some cases, an immunoglobulin of the invention specifically targets a CLPTM1L polypeptide having one or more mutations. Frameshift and missense mutations of CLPTM1L polypeptides that may be targeted may include, but are not limited to, those set forth in Table 2. The amino acid residue positions are numbered relative to the amino acid sequence of human CLPTM1L set forth as SEQ ID NO:2. Nucleotide positions for coding sequence mutations are numbered relative to the nucleotide sequence of human CLPTM1L set forth as SEQ ID NO:1. Somatic CLPTM1L mutations have been found in tumors of the autonomic ganglia, breast, central nervous system, endometrium, kidney, large intestine, liver, lung, ovary, prostate, skin, stomach, and upper respiratory tract. Each of the known CLPTM1L mutations is rare (<3% of tumors tested), and the functional effects of these mutations are unknown.

Table 2. Known Frameshift and Missense Mutations in Human Tumors.

Position	CDS Mutation	AA Mutation	Type
206	c.616delA	p.T206fs*17	Deletion - Frameshift
61	c.181A>G	p.T61A	Substitution - Missense
72	c.214G>A	p.D72N	Substitution - Missense
74	c.221T>C	p.V74A	Substitution - Missense
105	c.313T>C	p.Y105H	Substitution - Missense
108	c.323T>A	p.I108N	Substitution - Missense
110	c.329T>C	p.L110P	Substitution - Missense

EP 3 094 654 B1

(continued)

Position	CDS Mutation	AA Mutation	Type
113	c.338C>A	p.A113D	Substitution - Missense
132	c.394A>G	p.T132A	Substitution - Missense
151	c.452A>C	p.Q151 P	Substitution - Missense
159	c.476C>T	p.P159L	Substitution - Missense
175	c.524C>T	p.A175V	Substitution - Missense
196	c.586C>T	p.R196W	Substitution - Missense
196	c.587G>A	p.R196Q	Substitution - Missense
216	c.646G>A	p.D216N	Substitution - Missense
221	c.661C>T	p.R221C	Substitution - Missense
250	c.749G>A	p.R250H	Substitution - Missense
266	c.796G>A	p.G266R	Substitution - Missense
313	c.939G>T	p.K313N	Substitution - Missense
319	c.957C>G	p.I319M	Substitution - Missense
320	c.959G>A	p.G320D	Substitution - Missense
322	c.965C>T	p.S322F	Substitution - Missense
353	c.1058C>T	p.A353V	Substitution - Missense
377	c.1128_1129CC>AA	p.L377M	Substitution - Missense
390	c.1168G>A	p.E390K	Substitution - Missense
396	c.1187A>G	p.Y396C	Substitution - Missense
400	c.1198G>T	p.A400S	Substitution - Missense
405	c.1214C>T	p.S405L	Substitution - Missense
430	c.1289C>A	p.S430Y	Substitution - Missense
431	c.1291T>C	p.W431R	Substitution - Missense
465	c.1394C>T	p.P465L	Substitution - Missense
490	c.1469C>T	p.T490M	Substitution - Missense
493	c.1477C>T	p.R493W	Substitution - Missense
498	c.1493G>A	p.R498Q	Substitution - Missense
500	c.1498G>A	p.D500N	Substitution - Missense
517	c.1550A>G	p.K517R	Substitution - Missense
529	c.1587G>C	p.E529D	Substitution - Missense
532	c.1595C>T	p.T532M	Substitution - Missense
533	c.1597C>T	p.R533W	Substitution - Missense
538	c.1612G>A	p.D538N	Substitution - Missense
* CDS=coding sequence			

[0045] In another aspect, described herein is a nanoparticle conjugated to one or more CLPTM1L-specific immunoglobulins. Generally, antibody-conjugated nanoparticles are able to bind to targets with high affinity and cross biological barriers (e.g., blood-brain barrier (BBB)) and, therefore, are advantageous for molecular and cellular targeting. Accordingly, an antibody-conjugated nanoparticle described herein is useful for a variety of biomedical applications such as

in vivo diagnosis, clinical therapies, targeted drug delivery, gene therapy, cell labeling/tracking, and molecular imaging. In some cases, nanoparticles are lipid-based vehicles (e.g., liposomes, solid lipid nanoparticles, micelles); polymer carriers, such as hydrogels, polymersomes, dendrimers, and nanofibers; metallic nanoparticles (e.g., gold, silver, titanium); carbon structures (e.g., nanotubes, nanohorns, nanodiamonds (NDs), grapheme); or inorganic particles (e.g., silica). See, e.g., Chow and Ho, *Sci Transl Med* 5:216rv4 (2013); Montenegro et al., *Adv. Drug Delivery Rev.* 65:677-688 (2013).

[0046] In a further aspect, the present invention provides compositions comprising an immunoglobulin that targets a CLPTM1L polypeptide and at least one pharmaceutically acceptable carrier, diluent, or excipient. As used herein, the phrase "pharmaceutically acceptable" refers to those agents, compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and animals without excessive toxicity, irritation, allergic response, or other problems or complications, commensurate with a reasonable benefit/risk ratio.

[0047] In another aspect, compositions described herein include peptides comprised of part of the CLPTM1L amino acid sequence. For example, described herein is a peptide having the amino acid sequence of SEQ ID NO:4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 (see Table 1). Additional CLPTM1L peptides include peptides selected according to the above-defined epitope properties. Peptides may represent 8-22 amino acid length portions of SEQ ID NO. 2. In some cases, a CLPTM1L peptide may be embodied as a therapeutic agent. For example, a CLPTM1L peptide can be embodied in a peptide vaccine, a ligand blocking peptide, or a competitive inhibitory peptide. Such agents are expected to be useful therapeutic agents for treating or preventing lung cancer. Peptide vaccines are particularly amenable to prophylactic approaches. It is further expected that a CLPTM1L epitope, a CLPTM1L immunoglobulin, or any other CLPTM1L-targeting therapeutic agent will be useful in the treatment or prevention of many cancer types including, without limitation, lung, cervical, myeloblastic leukemia, pancreatic, bladder, glioma, prostate, basal cell carcinoma, and melanoma.

[0048] In some cases, a composition described herein is an aptamer having specificity for a CLPTM1L target molecule. As used herein, the term "aptamer" refers to a structured oligonucleotide-based or peptide-based molecule having high affinity and specificity for a specific target molecule (e.g., proteins, phospholipids, iron channels, nucleic acids, whole cells) due to unique structural features that restrict the aptamer to particular conformations. Oligonucleotide aptamers such as ribonucleic acid (RNA) and single-strand deoxyribonucleic acid (ssDNA) aptamers. Peptide aptamers are combinatorial protein molecules that comprise a short peptide region attached at both ends to a protein scaffold. Methods for designing and identifying therapeutic aptamers having high affinity and specificity for a CLPTM1L target molecule are known in the art. See, e.g., Rohloff et al., *Molecular Therapy Nucleic Acids* 3:e201 (2014); Zhu et al., *Theranostics* 4(9):931-944 (2014); Ellington and Szostak, *Nature* 346:818-822 (1990).

[0049] Compositions provided herein can comprise at least one pharmaceutically acceptable diluent, excipient, or carrier. As used herein, "pharmaceutically acceptable carrier" refers to any and all solvents, dispersion media, coatings, surfactants, antioxidants, preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, absorption delaying agents, salts, preservatives, drugs, drug stabilizers, gels, binders, excipients, disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, such like materials and combinations thereof, as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, pp. 1289-1329). Except insofar as any conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated.

[0050] Regardless of the route of administration selected, therapeutic agents of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, can be formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

[0051] In certain embodiments, a pharmaceutical composition provided herein comprises one or more CLPTM1L-targeting agents, a chemotherapeutic agent, and a pharmaceutically acceptable carrier. Chemotherapeutic agents include, without limitation, platinum-based agents, such as cisplatin, gemcitabine, and carboplatin; nitrogen mustard alkylating agents; nitrosourea alkylating agents, such as carmustine (BCNU) and other alkylating agents; antimetabolites, such as methotrexate; purine analog antimetabolites; pyrimidine analog antimetabolites, such as fluorouracil (5-FU) and gemcitabine; hormonal antineoplastics, such as goserelin, leuprolide, and tamoxifen; natural antineoplastics, such as taxanes (e.g., docetaxel and paclitaxel), aldesleukin, interleukin-2, etoposide (VP-16), interferon .alpha., and tretinoin (ATRA); antibiotic natural antineoplastics, such as bleomycin, dactinomycin, daunorubicin, doxorubicin, and mitomycin; and vinca alkaloid natural antineoplastics, such as vinblastine and vincristine.

[0052] In other embodiments, compositions of the invention provided herein can additionally comprise one or more other biologically active substances including, without limitation, therapeutic drugs or pro-drugs such as chemotherapeutic agents other than those identified above, scavenger compounds, antibiotics, antiviral agents, antifungal agents, anti-inflammatory agents, vasoconstrictors and anticoagulants, antigens useful for cancer vaccine applications or corresponding pro-drugs.

Methods of Using Compositions of the Invention

[0053] In one aspect, described herein are methods of treating or preventing a disease or condition in a subject by inhibiting CLPTM1L. For example, described herein are methods comprising administering to a subject in need thereof an immunoglobulin or a composition comprising an immunoglobulin that targets and specifically binds to CLPTM1L. As used herein, the term "subject" refers to an individual having, suspected of having, or susceptible to having a disease or condition associated with CLPTM1L protein dysfunction (e.g., pathogenic production, modification, or function) or for which there is a genetic association with *CLPTM1L* (e.g., a disease or condition associated with a gain in *CLPTM1L* locus copy number or a genotype for susceptibility to the disease or disorder). By "subject" or "individual" or "animal" or "patient" or "mammal," is meant any subject, particularly a mammalian subject, for whom diagnosis, prognosis, or therapy is desired. Mammalian subjects include humans, domestic animals, farm animals, and zoo, sport, or pet animals such as dogs, cats, guinea pigs, rabbits, rats, mice, horses, cattle, cows, and so on.

[0054] As used herein, the terms "treating," "treat," and "treatment" refer to the management and care of a patient for the purpose of combating the disease, condition, or disorder. The terms embrace both preventative, i.e., prophylactic, and palliative treatments. In some cases, the term "treated" refers to any beneficial effect on progression of a disease or condition. Beneficial effects can include reversing, alleviating, inhibiting the progress of, preventing, or reducing the likelihood of the disease or condition to which the term applies or one or more symptoms or manifestations of such a disease or condition. Where the disease or condition is a tumor, cancer, or cancer-associated condition, treating can refer to the management and care of a patient for the purpose of combating cancer, and can include reversing, alleviating, inhibiting the progress of, preventing, reducing the size of, or reducing the likelihood of, or lessening the severity of any aspect of the cancer or cancer-associated condition (e.g., metastasis, tumor growth). A therapeutic beneficial effect on the health and well-being of a patient includes, but is not limited to: (1) curing the cancer; (2) slowing the progress of the cancer; (3) causing the tumor to regress; or (4) alleviating one or more symptoms of the cancer. As used herein, the terms "preventing" and "prevent" refer not only to a complete prevention of a certain disease or condition, but also to partially or substantially attenuating, reducing the risk of, or delaying the development or recurrence of the disease or condition to which the term applies.

[0055] In some cases the methods described herein are directed to treating or preventing a tumor cancer in a subject by administering a composition provided herein. In an exemplary aspect, the method treats or prevent a solid tumor in the subject receiving administration of a composition comprising a CLPTM1L-targeting agent to a subject in need thereof. As used herein, the term "cancer" includes, without limitation, solid tumors and blood-borne tumors (e.g., leukemias). The term cancer includes, without limitation, diseases of the skin, tissues, organs, bone, cartilage, blood, and vessels. The term further encompasses both primary and metastatic cancers. In some cases, therefore, methods of treating or preventing cancer as described herein include methods of inhibiting, retarding, or preventing growth of a tumor or tumor cells. A subject in need thereof may include, for example, a subject who has been diagnosed with a tumor, including a pre-cancerous tumor, a cancer (e.g., a non-small cell lung cancer), or a subject who has been treated, including subjects that have been refractory to previous treatment.

[0056] In some cases, solid tumors are considered to be a "refractory" or "resistant" solid tumor, meaning that the solid tumor does not respond to treatment. The tumor may be resistant at the outset of treatment or it may develop resistance during treatment. While any and all tumors that are susceptible to treatment and/or prophylactic administration of a CLPTM1L-targeting agent described herein are of course within the scope of this disclosure, it is anticipated that the compositions and methods provided herein will be particularly useful in the treatment of refractory tumors.

[0057] The methods described herein are appropriate for treating or preventing any type of disease or condition associated with CLPTM1L protein dysfunction (e.g., pathogenic production, modification, or function) or any disease or condition for which there is a genetic association with CLPTM1L (e.g., a disease or condition associated with a gain in *CLPTM1L* locus copy number or a genotype for susceptibility to the disease or disorder). In an exemplary aspect, a method described herein is for the treatment or prevention of a cancer, a tumor (e.g., a solid tumor), or a pre-neoplastic lesion (e.g., pre-cancerous lesion). Examples of cancers appropriate for methods of treating or preventing as described herein include, without limitation, lung cancer, pancreatic cancer, prostate cancer, skin cancer, bladder cancer, kidney cancer, ovarian cancer, colon cancer, colorectal cancer, breast cancer, cervical cancer, brain cancer, esophageal cancer, and stomach cancer. Solid tumors can be associated with cancers and pre-neoplastic lesions of the lung, pancreas, prostate, breast, kidney, and other sarcomas, carcinomas, and glioblastomas. Other diseases or conditions appropriate for methods of treating or preventing as described herein include, without limitation, lymphomas, chronic leukemia, and acute leukemia.

[0058] In some cases, a method described herein can be practiced to treat or prevent a disease or condition in a subject, where the disease or condition exhibits chemotherapeutic drug resistance. For example, a subject can be diagnosed or identified as having a disease or condition such as cancer that exhibits resistance to a chemotherapeutic agent such as cisplatin or gemcitabine. CLPTM1L has been found to be highly expressed in cisplatin resistant ovarian tumor cell lines. Moreover, CLPTM1L appears to be anti-apoptotic under genotoxic conditions. Examples of cancers for

which treatment involves administering cisplatin include lung cancer, colorectal cancer, non-small cell lung cancer (NSCLC), bronchioloalveolar cell lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous melanoma, intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, anal region cancer, stomach cancer, gastric cancer, colon cancer, breast cancer, uterine cancer, fallopian tube carcinoma, endometrial carcinoma, cervical carcinoma, Hodgkin's Disease, esophagus cancer, small intestine cancer, endocrine system cancer, thyroid gland cancer, parathyroid gland cancer, adrenal gland cancer, soft tissue sarcoma, prostate cancer, bladder cancer, kidney cancer, renal cell carcinoma, renal pelvis carcinoma, mesothelioma, hepatocellular cancer, biliary cancer, chronic leukemia, acute leukemia, lymphocytic lymphoma, CNS neoplasm, spinal axis cancer, brain stem glioma, glioblastoma multiforme, astrocytoma, schwannoma, ependymoma, medulloblastoma, meningioma, squamous cell carcinoma and pituitary adenoma tumors or tumor metastases.

[0059] In some cases, a method described herein can be practiced to treat or prevent a cancer associated with a mutation, translocation, amplification, or deletion of at least a portion of at least one of the following: K-Ras, H-Ras, N-Ras, RASSF1, PI3KCA, PTEN, EGFR, FGFR1, PDGFRA, BRAF, AKL, ROS1, BCL-x, BIM, BAD, BAX, AKT, and mTOR. In an exemplary embodiment, a method described herein can be practiced to treat or prevent a cancer for which at least one of these genes suspected of being or demonstrated to be regulated by CLPTM1L.

[0060] In some cases, a method described herein can be practiced to treat or prevent a cancer previously or currently being subjected to radiation therapy. As used herein, the term "radiation therapy" refers to any manner of treatment of solid tumors and cancers with ionizing radiation and includes, without limitation, external beam radiotherapy, stereotatic radiotherapy, virtual simulation, 3-dimensional conformal radiotherapy, intensity-modulated radiotherapy, ionizing particle therapy, and radioisotope therapy.

[0061] In other cases, a method described herein can be practiced to treat or prevent a cancer previously or currently being subjected to chemotherapy using a chemotherapeutic agent such as, for example, an alkylating agent, a cross-linking agent, an anti-metabolite, an antibiotic, a topoisomerase inhibitor, or a mitotic inhibitor. As used herein, the term "chemotherapeutic agent" refers to any substance that, when administered in a therapeutically effective amount to a patient suffering from a tumor or cancer, has a therapeutic beneficial effect on the health and well-being of the patient. A therapeutic beneficial effect on the health and well-being of a patient includes, but is not limited to: (1) curing the cancer; (2) slowing the progress of the cancer; (3) causing the tumor to regress; or (4) alleviating one or more symptoms of the cancer. As used herein, a chemotherapeutic agent also includes any substance that, when administered in a prophylactic amount to a patient afflicted with a solid tumor or who has been rendered substantially free of cancer as the result of one or more therapeutic treatment regimes, has a beneficial effect on the health and well-being of the patient.

[0062] The method also can be practiced to treat or prevent a cancer if use of such a therapeutic agent is anticipated. For example, methods described herein are useful to treat or prevent a cancer for which targeting CLPTM1L sensitizes the cancer cells to DNA damage-induced apoptosis. As used herein, the terms "sensitize" and "sensitized" refers to cells made more responsive to an agent, a treatment, an environment, a stimulus, or a condition. For example, a "sensitized" cell can exhibit an increase in the magnitude of said response or an increase in any type of response. In an exemplary aspect, cancer cells exhibit increased apoptosis when sensitized to DNA damage by targeting CLPTM1L according to a method described herein.

[0063] In some cases, a method described herein can be practiced to treat or prevent a cancer suspected of being or demonstrated to be refractory to one or more chemotherapeutics such as, for example, an alkylating agent, a cross-linking agent, an anti-metabolite, an antibiotic, a topoisomerase inhibitor, or a mitotic inhibitor. For example, a method described herein can be appropriate to treat or prevent a cancer for which targeting CLPTM1L using an immunoglobulin sensitizes the cancer cells to cytotoxic treatment with a cross-linking agent and a topoisomerase inhibitor.

[0064] In exemplary aspects, a method of treating or preventing cancer as described herein comprises administering to a subject in need thereof two or more CLPTM1L-specific immunoglobulins. The two or more CLPTM1L-specific immunoglobulins can be administered simultaneously or sequentially. For example, the method can comprise administering two or more CLPTM1L-specific immunoglobulins, where the immunoglobulins are selected for capacity to increase sensitivity of CLPTM1L-expressing tumor cells to a chemotherapeutic agent (e.g., cisplatin, gemcitabine) and/or for an enhanced capacity for cytotoxicity toward CLPTM1L-expressing tumor cells. The combination can comprise antibodies having specificity to different polypeptides or to different epitopes on the same polypeptide. Such combination therapies have been shown to improve tumor responses and reverse resistance of tumor cells to a single agent. See, e.g., Weiner et al., *Nature Rev. Immunol.* 10:317-327 (2010).

[0065] In another aspect, described herein is a method of treating or preventing cancer in a subject, where the method comprises administering an immunoconjugate. As described herein, immunoconjugates are effective cytotoxic and anti-cancer therapeutic agents.

[0066] In a further aspect, described herein is a method of treating or preventing cancer in a subject, where the method comprises administering activated T cells to a subject in need thereof. Such activated T cell therapy methods generally comprise *ex-vivo* activation and expansion of T cells. See, for review, Slaney et al., *Cancer Res.* 74:7168-7174 (2014). In some cases, T cells are obtained from a subject, then purified, expanded, and activated *ex vivo*. Activation comprises

culturing purified and expanded T cells in the presence of a cancer cell-specific or tumor specific antigen (e.g., a monoclonal antibody provided herein). The resulting activated T cells are provided back to the subject. T cells activated in this manner have enhanced immunostimulatory capabilities. In exemplary embodiments, T cells obtained from a subject in need of treatment according to a method provided herein are cultured in the presence of one or more CLPTM1L-specific monoclonal antibodies (e.g., mAbs 6-1, 10-2, and 10-3).

[0067] Methods of treating or preventing cancer as described herein can be practiced at any appropriate time. In some cases, a composition comprising a CLPTM1L-targeting agent as provided herein is administered to a subject following diagnosis of a cancer or a pre-neoplastic lesion (e.g., pre-cancerous lesion) in, for example, a biological sample from the subject. In other cases, a composition comprising a CLPTM1L-targeting agent as provided herein is administered to a subject following identification of copy number gain or expression of *CLPTM1L* in a biological sample from the subject, where the level of expression is above a threshold that is empirically determined to constitute cancer risk.

[0068] Determination of such a threshold includes analysis of current and future data correlating *CLPTM1L* expression in normal tissue with incidence of a cancer. With respect to lung tissue, a threshold that is empirically determined to constitute cancer risk can be at least 2-fold greater average expression for one or more test lung tissue samples relative to the average expression in a healthy sample taken from surrounding tissue. In some cases, treatment may be indicated if it is determined that expression of *CLPTM1L* is higher in a test sample (e.g., tissue suspected of comprising cancer cells) compared to tissue surrounding the sampled tissue. In other cases, treatment may be indicated if it is determined that an increased amount of CLPTM1L is present on a tumor cell surface. Such an increased level of CLPTM1L can be detected by, for example, flow cytometry in circulating tumor cells or in solid tumors. A composition comprising a CLPTM1L-targeting agent as provided herein can be administered to a subject upon identification of one or more indicators or risk factors for the development of cancer or following identification of a genotype associated with a cancer in a biological sample from the subject. To prevent or slow tumor formation, a composition comprising a CLPTM1L-targeting agent as provided herein is administered to a subject prior to or in the absence of a cancer or pre-neoplastic lesion. In such cases, the composition is administered as a preventative agent.

[0069] Treatment or prevention according to a method described herein can occur before, during, or after the subject is treated by surgery, radiation, and/or chemotherapy. In some cases, treatment according to a method provided herein prior to chemo- or radiotherapy may improve the outcome of the conventional therapy. In an exemplary aspect, a composition comprising a CLPTM1L-targeting agent as provided herein is administered to a subject concurrently with one or more other treatments or preventative measures such as radiotherapy, chemotherapy, or surgery.

[0070] A composition comprising a CLPTM1L-targeting agent as provided herein is administered to a subject by any method that achieves the intended purpose or is deemed appropriate by those of skill in the art. For example, a composition of the present invention can be administered as a pharmaceutical, and may be administered systemically or locally via oral or parenteral administration. As used herein, the term "administration" includes oral and parenteral administration. Oral administration includes, for example, administration of oral agents. Such oral agents include, for example, granules, powders, tablets, capsules, solutions, emulsions, and suspensions. Parenteral administration includes, for example, administration of injections. Such injections include, for example, subcutaneous injections, intramuscular injections, and intraperitoneal injection. In some cases, intravenous injections such as drip infusions, intramuscular injections, intraperitoneal injections, subcutaneous injections, suppositories, enemas, oral enteric tablets, or the like can be selected.

[0071] Appropriate modes of administration can be determined based on the physical location of a tumor or tumors in the subject's body. In exemplary aspects, a composition comprising a CLPTM1L-targeting agent as provided herein is administered to a subject having a diagnosis of lung cancer or a pre-cancerous lesion, where the composition is administered orally or intravenously. Alternatively, a composition comprising a CLPTM1L-targeting agent can be administered locally to an intended area of treatment. For example, a composition comprising a CLPTM1L-targeting agent can be administered by local injection during surgery.

[0072] Compositions can be administered to a subject in need thereof in dosage unit form where each discrete dosage unit contains a predetermined quantity of an active ingredient or compound that was calculated to elicit a desirable therapeutic effect when administered with, in some cases, a pharmaceutically acceptable carrier.

[0073] A therapeutically effective dose relates to the amount of a compound which is sufficient to improve the symptoms, for example a treatment, healing, prevention or improvement of such conditions. In exemplary aspects, a therapeutically effective amount or dose is an amount such that free antibody is present in the blood. For dosage determinations, it can be advantageous to assess toxicity and therapeutic efficacy of a compound in cell cultures or in experimental animals. For example, the LD₅₀ (i.e., the dose lethal to 50% of the population) and ED₅₀ (i.e., the dose therapeutically effective in 50% of the population) can be determined. From these calculations, dosage ranges for use in humans can be formulated. Dosage ranges can vary depending on factors such as mode of administration. A therapeutically effective amount of a pharmaceutical composition provided herein can range from about 0.001 to 100 mg of antibody per kg body weight of the subject (e.g., about 0.01 to 100 mg/kg body weight; about 0.1 to 40 mg/kg body weight; about 1 to 20 mg/kg body weight).

[0074] In some cases, an appropriate dose for a monoclonal antibody can be from 0.005 mg/kg up to a maximum

tolerated dose. In some cases, an appropriate dose of a pharmaceutical composition as provided herein can be determined according to body surface area of a subject, calculated using the subject's height and weight, to whom the composition will be administered. In such cases, a dose can be provided as a particular amount of the composition per m² (e.g., mg/m²). In some cases, an appropriate dose can be between approximately 10 mg/m² and approximately 40 mg/m² of a monoclonal antibody. When converted to milligrams (mg) per kilogram (kg) of a subject's body weight, a dose of 15 mg/m² is the same as about 0.4 mg/kg. See Freireich et al., *Cancer Chemotherapy Rep.* 50(4):219-244 (1966). Additional information about dosage calculation can be found in Center for Drug Evaluation and Research, Center for Biologics Evaluation and Research (2002), *Estimating the safe starting dose in clinical trials for therapeutics in adult healthy volunteers*, U.S. Food and Drug Administration, Rockville, Maryland, USA.

[0075] It will be understood that mass of pharmaceutical composition comprising a monoclonal antibody as provided herein can refer to mass of the antibody plus a delivery agent or pharmaceutically acceptable carrier, if applicable. In some cases, dosages and dosage ranges appropriate for a composition provided herein can be determined using pharmacokinetic data (i.e., drug metabolism and clearance). As used herein, "pharmacokinetics" refers to the process by which a drug or pharmaceutical composition is absorbed, distributed, metabolized, and excreted from the body.

[0076] Clinicians, physicians, and other health care professionals can administer a composition to a subject in need thereof according to a method described herein by a physician or other health professional. In some cases, a single administration of the composition may be sufficient. In other cases, more than one administration of the composition is performed at various intervals (e.g., once per week, twice per week, daily, monthly) or according to any other appropriate treatment regimen. The duration of treatment can be a single dose or periodic multiple doses for as long as administration of a composition provided herein is tolerated by the subject.

[0077] Any appropriate method can be practiced to determine, detect, or monitor a subject's response to treatment according to a method described herein. As used herein, "determining a subject's response to treatment" refers to the assessment of the results of a therapy in a subject in response to administration of a composition provided herein or to treatment according to a method described herein. For example, a subject's condition can be monitored continuously or evaluated at appropriate time intervals (e.g., at regular or irregular time points) to detect and/or monitor any changes in disease progression (e.g., change in tumor size) as an indicator of the subject's response to a composition comprising an RNAi-inducing construct targeted to *CLPTM1L*. In some cases, tumors can be measured to detect or monitor any change in, for example, tumor size or tumor growth rate (e.g., tumor expansion or shrinkage, inhibited or accelerated tumor growth rate). For example, detection methods such as computed tomography (CT), magnetic resonance imaging (MRI) scanning, and x-ray (e.g., chest x-ray) can be used. In some cases, ultrasound examinations can be used to detect and measure tumor regression or to detect progression of lesions. In other cases, evaluation of a tumor or pre-neoplastic lesion can involve cytology or histology of, for example, biopsy samples. For solid tumors, evaluation of a subject's response to treatment as described herein can include assessing RECIST ("Response Evaluation Criteria in Solid Tumors"). RECIST criteria can be used to evaluate a subject's response to the therapy used to treat their disease or condition. See, for review, Therasse et al., *J. Natl. Cancer Inst.* 92:205-16, 2000.

[0078] In some cases, biomarkers (e.g., mRNA, protein) can be used to detect or monitor the efficacy of a treatment or prevention method described herein. In an exemplary aspect, use of a biomarker can comprises a) administering a composition provided herein; b) determining the levels of a biomarker according to the present invention in one or more biological samples taken from the subject at different time points (before, during and/or after administration); and c) comparing the determinations made for the biological samples obtained during a particular phase of treatment and comparing them to controls or to levels determined for the subject's samples obtained at different phases of treatment. For bone and blood tumors, evaluation using biomarkers can include detecting or monitoring expression levels for one or more tumor markers and assessing hematologic indicators including, for example, mean platelet volume, platelet counts, leukocyte counts, and hemoglobin level. Other indicators or "efficacy markers" of a positive outcome following administration of a composition comprising a *CLPTM1L*-specific agent according to a method described herein can include (1) reduced *CLPTM1L* transcript and/or protein levels in tumors or target tissues and (2) reduced phosphorylated Akt or Bcl-xL protein levels in tumors or target tissues. These efficacy markers can be determined by biopsy, aspirate, or lavage followed by, in an exemplary aspect, an appropriate diagnostic test such as PCR, Western blotting, immunohistochemistry using specific antibodies. A positive result for any of the outcome criteria or evaluation methods described herein is indicative of the method's efficacy for treating or preventing the subject's disease or condition.

[0079] Indicators of a positive response to administration of a composition comprising a *CLPTM1L*-targeting agent according to a method described herein can include, for example, a significant decrease in *CLPTM1L* transcript and/or *CLPTM1L* protein levels in tumors or target tissues relative to pre-treatment levels or to untreated samples (e.g., expression reduced to approximately 60%-80% of expression in a control sample). In some cases, an indicator of a positive response to administration of a composition comprising a *CLPTM1L*-targeting agent can be significantly reduced levels of phosphorylated Akt and/or Bcl-xL proteins in tumors or target tissues relative to pre-treatment levels or to untreated samples or a significant increase in apoptosis in tumors or target tissues relative to pre-treatment levels or to untreated samples. According to RECIST criteria, a partial response to treatment can be indicated by at least a 30% decrease in

the sum of the longest diameter of a target lesions, taking as reference the baseline sum longest diameter, and a complete or substantially complete response to treatment can be indicated by the complete or nearly complete disappearance of all target lesions relative to measurements obtained for the subject prior to treatment (*i.e.*, baseline measurement) or relative to a control or a comparative decrease in disease progression. In some cases, response to treatment is evaluated relative to one or more subjects who were not administered a composition described herein. Other parameters for evaluating a subject's response to treatment according to a method described herein include detecting a comparative decrease in metastatic growth; detecting any improvement in RECIST criteria for solid tumors; documenting short-term or long-term survival; documenting disease-free survival; detecting increased or decreased expression of tumor markers; detecting hematologic changes for blood and bone cancers; detecting or monitoring positive or negative responses to radiotherapy and/or chemotherapy; and detecting an increase or decrease in recurrence of the treated disease or condition. In some cases, a subject treated according to a method described herein may exhibit signs of stable disease, where there is neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease, taking as reference the smallest sum longest diameter since the treatment started (Therasse *et al.*, *supra*).

[0080] The determination of the response of a subject to a specific therapy can be determined using any assessment criterion used in oncology and known by persons skilled in the art. Assessment parameters useful for describing progression of a disease include: disease-free progression which, as used herein, describes the ratio of subjects in complete remission who have not had disease relapse during the time period under study; objective response, which, describes the ratio of subjects treated in whom a complete or partial response is observed; tumor control, which, relates to the ratio of people treated in whom a complete response, partial response, minor response or stable disease ≥ 6 months is observed; progression-free survival which, as used herein, is defined as the time from the beginning of the treatment until the first measurement of cancer growth. In a preferred aspect, the response of a subject is determined by means of a parameter selected from time to progression and survival. In an exemplary aspect, a subject's response to a treatment or preventative method described herein should be statistically significant. The determination of whether a response is statistically significant can be carried out using statistical evaluation tools such as confidence intervals, determination of the p value, Student's t-test, Mann-Whitney test, etc. Preferred confidence intervals are at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%. Preferably, p values are 0.2, 0.1, or 0.05.

[0081] While the subject is being treated, the health of the subject may be monitored by measuring one or more of the relevant indices at predetermined times during a 24-hour period. All aspects of the treatment, including supplements, amounts, times of administration and formulation, can be optimized based on the results of such monitoring. A patient can be periodically reevaluated to determine the extent of improvement by measuring the same parameters. For example, a first such reevaluation can occur at the end of an appropriate length of time (*e.g.*, about 2 weeks, about 4 weeks, about 8 weeks, or more) following the onset of therapy, and subsequent reevaluations can occur at appropriate intervals (*e.g.*, every four to eight weeks during therapy and then every 3, 6, 9, or more months) thereafter.

[0082] The articles "a" and "an" are used herein to refer to one or to more than one (*i.e.*, to at least one) of the grammatical object of the article. By way of example, "an antibody" means one antibody or more than one antibody. As such, the terms "a" (or "an"), "one or more," and "at least one" are used interchangeably herein.

[0083] While the present invention is susceptible to various modifications and alternative forms, specific embodiments thereof have been shown by way of example in the drawings and are herein described in detail.

[0084] The invention will be more fully understood upon consideration of the following nonlimiting Examples.

EXAMPLES

Example 1 - Structural Analysis of CLPTM1L

[0085] We generated a basic structural model of CLPTM1L that includes six transmembrane domains, extracellular globular domains (ECR1 and ECR2), the disordered region, and N-glycosylated cell-surface peptides identified by mass spectrometry (FIG. 2A). There is a putative alternatively spliced isoform of CLPTM1L lacking ECR1 and exons 1, 2, and 9 (FIG. 2B). Glycosylated (N-linked) amino acid residues are predicted at positions 91, 101, and 229 of the human CLPTM1L protein sequence (SEQ ID NO:2). Mass spectrometry evidence of glycosylation has been reported for amino acid residue 91 of human CLPTM1L (data not shown) and for residue 229 (Chen *et al.*, J Proteome Res. 2009;8:651-61). For a plasma membrane protein, this is highly indicative of extracellular orientation of the glycosylated region. It is predicted that this extracellular region comprises two separate globular domains separated by a small disordered region (amino acid residues 142-162) within the larger extracellular region (FIG. 1).

[0086] Although our analysis of CLPTM1L suggested the possibility of plasma membrane localization and cell surface exposure, subcellular localization of CLPTM1L had not been well studied and plasma membrane localization was not previously reported. One study suggested that CLPTM1L may be localized to mitochondria. See Ni *et al.*, PLoS One 7:e52598, 2012. The potential for CLPTM1L exposure at the surface of tumor cells has significant relevance to the ability to target this protein with exogenous agents.

[0087] Since monoclonal antibodies are definable, consistent, reproducible, and capable of being manipulated, we contracted the production of a panel of monoclonal antibodies targeting the human CLPTM1L protein. Design of peptide epitopes against which to raise monoclonal antibodies took several factors into consideration. We designed the epitopes to reside within the cell surface exposed globular domains of CLPTM1L (residues 32-284, avoiding glycosylation sites (91,101, 229), avoiding the disordered region (141-162), and prioritizing amino acid residues scoring highly on the basis of hydrophilicity, folding potential, and antigen presentation. Eight epitopes, each 11 amino acids in length, were selected for monoclonal antibody production (FIG. 3; see also shaded boxes of Table 1). While polyclonal antibodies may be generated against peptides of 8-22 amino acids, epitopes of 9-11 amino acids are typically chosen for specific monoclonal antibody production. Other potential epitopes selected bases on similar parameters are presented in the unshaded boxes of Table 1.

[0088] To confirm the localization of CLPTM1L at the surface of tumor cells, we isolated cell-surface proteins from A549 human lung tumor cells by biotinylation and extraction with a streptavidin resin. Western analysis of whole cell and cell-surface lysates demonstrated a prevalence of CLPTM1L at the cell-surface (FIG. 4A). EGFR, used as a cell-surface localized control, was detected at the cell-surface at a lesser intensity than CLPTM1L relative to whole cell lysate signal. Cyclin D1, used as a nuclear protein control, was not detected at the cell surface. Immunofluorescence staining of non-permeabilized A549 cells for CLPTM1L revealed a punctate cell-surface staining pattern (FIG. 4B).

[0089] Our panel of monoclonal antibodies targeting the extracellular globular domains of CLPTM1L was tested for detection of CLPTM1L and immunofluorescent labeling of surface CLPTM1L. All antibodies strongly detected the 62-kilodalton CLPTM1L protein by western blot (FIG. 4C; some data not shown). All antibodies resulted in a cell-surface staining pattern when used for immunofluorescence on non-permeabilized A549 cells, with antibodies targeting ECR2 exhibiting a more distinct punctate staining (FIG. 4D; some data not shown).

Example 2 - *In Vitro* Analysis of CLPTM1L Monoclonal Antibodies

[0090] To determine if CLPTM1L expression was associated with K-Ras driven pancreatic cancer treated with adjuvant therapy, we utilized available pancreatic tissue microarrays on 31 patients using matched survival data. Staining intensity of tumors that were subjected CLPTM1L immunohistochemistry (IHC) were scored by three pathologists and the scores averaged. Staining was uniformly negative in normal ductal epithelia, while positive in 94% (29/31) of tumor cells ($r=0.78$, $p<5\times 10^{-10}$) (FIG. 5A). Tumors were scored as negative (0), weak (+1), moderate (+2) or strong (+3). Independent scores were averaged. Tumors had an average score of 1.46 while that of normal ductal epithelial was 0 ($p<0.05$). Tumors expressing low levels of CLPTM1L as defined by an average score of 1.15 or less were associated with increased survival, while patients with higher expression had shorter survival times. Patients exhibiting low CLPTM1L expression in tumor tissues had a median survival of 24 months, while patients with medium and high CLPTM1L expression in tumor tissues had a median survival of 11 and 14 months respectively (FIG. 5B). Representative tissues stained with CLPTM1L antibody are shown in FIG. 5C.

[0091] We previously demonstrated that targeting CLPTM1L in tumor cells modulates Bcl-xL and Akt survival signaling and sensitizes cells to killing with cisplatin. Polyclonal antibody raised against amino acid residues 18-307 of human CLPTM1L was applied to A549 human lung tumor cells at concentrations of 0, 1, and 5 $\mu\text{g/mL}$. After culture for 48 hours with polyclonal antibody treatment, Western blotting was conducted for Bcl-xL, total Akt1, and phospho-Akt1. Dose dependent decreases in the accumulation of both Bcl-xL and Akt1 with anti-CLPTM1L treatment were observed (FIG. 6A). To determine the effect of treatment with polyclonal antibody on sensitivity to cisplatin, cells were treated with dilutions of anti-CLPTM1L from 0.01 to 1 $\mu\text{g/mL}$ for 24 hours. Cells were then exposed to cisplatin at 0, 10, and 20 μM concentrations for 48 hours. Treatment with anti-CLPTM1L resulted in a decrease in cell viability following cisplatin treatment that was dependent on the dose of antibody (FIG. 6B). These results demonstrate sensitization to cisplatin killing using the antibody.

[0092] Treatment with monoclonal anti-CLP (10-2) increased sensitivity of pancreatic adenocarcinoma cells to killing by gemcitabine (500 μM for 72 hours) (FIG. 6C) and also reduced accumulation of phosphorylated Akt (FIG. 6D).

[0093] In addition, treatment of A549 and H838 human lung tumor cells with our panel of monoclonal antibodies resulted in depletion of Bcl-xL and Akt phosphorylation with a range of efficacies. Western analysis revealed that treatment with antibodies targeting epitopes 6 and 10 resulted in dose dependent decreases in Bcl-xL and phospho-Akt1 accumulation. Likewise, sensitization to cisplatin killing was accomplished by treatment of tumor cells with anti-CLPTM1L monoclonal antibodies. Monoclonal antibody raised against epitope 10 effectively inhibited anchorage independent growth of A549 cells. Among all monoclonal antibodies raised against the extracellular domains of CLPTM1L, only those raised against epitope 10 were effective in inhibiting anchorage independent tumor spheroid growth of H838 cells. A summary of these results for each antibody clone is given in FIG. 10.

[0094] As shown in FIG. 8, we also discovered that treatment with monoclonal anti-CLPTM1L antibodies inhibited colony growth of pancreatic tumor cells. Panc1 pancreatic adenocarcinoma cells were plated at low density and treated with anti-CLPTM1L 6-1, 10-2, or control mouse IgG (FIGS. 8A-B). The number of resulting colonies after 5 days of

growth was inhibited by up to 47% by anti-CLPTM1L treatment.

Example 3 - Effect of CLPTM1L Monoclonal Antibodies on Human Lung Tumor Xenograft Model

[0095] Monoclonal antibodies raised against epitope 6 in ECR1 (SEQ ID NO:6) and epitope 10 in ECR2 (SEQ ID NO:10) were evaluated in xenograft models for inhibition of the growth of established human lung tumor xenografts. A549 cell xenografts were established and mice were separated into groups of equal average tumor size and variance. Mice were treated intraperitoneally with 1 μ L/g (estimated to be equivalent to 2-3 mg/kg of antibody) of the indicated monoclonal antibody clone or control ascites on days 0, 5, and 12 post-separation. One group was similarly treated with 1 mg/kg polyclonal antibody. Polyclonal antibody and monoclonal clone 6-1 modestly but significantly inhibited tumor growth over 17 days posttreatment (FIG. 9). Antibodies raised against epitope 10 (10-2 and 10-3) had a more robust and more significant inhibitory effect on tumor growth. Monoclonal clone 10-2 inhibited tumor growth by 88% ($p < 0.005$). **[0096]** The examples and data set forth above demonstrate that anti-CLPTM1L immunoglobulins are anti-tumor agents *in vivo* and *in vitro*. Furthermore, they describe specific epitopes expected to be efficacious in targeting the extracellular domains of CLPTM1L.

Claims

1. A CLPTM1L-targeting agent, wherein the CLPTM1L-targeting agent is a monoclonal antibody that binds specifically to a peptide having an amino acid sequence of SEQ ID NO:6 or SEQ ID NO:10.
2. The CLPTM1L-targeting agent of claim 1, wherein the monoclonal antibody is a chimeric antibody, human antibody, humanized antibody, recombinant antibody, engineered antibody, conjugated antibody, or a fragment thereof.
3. The CLPTM1L-targeting agent of claim 1, wherein the monoclonal antibody is a CLPTM1L-targeted bispecific monoclonal antibody.
4. The CLPTM1L-targeting agent of claim 1, wherein the monoclonal antibody is an immunoconjugate comprising a therapeutic agent selected from a pharmacologic agent, radioisotope, and toxin; and a linker.
5. A pharmaceutical composition comprising the agent according to any of claims 1-4 and a pharmaceutically acceptable carrier.
6. The agent of any of claims 1-4 or the pharmaceutical composition of claim 5, for use as a medicament.
7. A compound comprising a CLPTM1L-targeting agent for use in a method for treating or preventing a tumor comprising administering to a subject in need thereof a therapeutically effective amount of said compound to a subject, wherein the CLPTM1L-targeting agent is a monoclonal antibody that binds specifically to a peptide having an amino acid sequence of SEQ ID NO:6 or SEQ ID NO:10, whereby the tumor is treated or prevented in the subject.
8. The compound for use in the method of claim 7, wherein:
 - (a) the tumor is a solid tumor selected from the group consisting of glioblastoma, sarcoma, carcinoma, and lymphoma; or
 - (b) the tumor is associated with a cancer or pre-neoplastic lesion selected from the group consisting of lung cancer, pancreatic cancer, prostate cancer, skin cancer, bladder cancer, kidney cancer, ovarian cancer, colon cancer, colorectal cancer, breast cancer, cervical cancer, brain cancer, esophageal cancer, and stomach cancer, or a pre-cancerous lesion thereof, preferably wherein the tumor is associated with lung cancer or pre-cancerous lesion thereof.
9. The compound for use in the method of claim 7, wherein the subject is a human.
10. The compound for use in the method of claim 7, wherein the tumor exhibits resistance to a chemotherapeutic agent; optionally wherein the chemotherapeutic agent is selected from the group consisting of cisplatin and gemcitabine.
11. The compound for use in the method of claim 7, wherein the compound is administered with a pharmaceutically acceptable carrier.

12. A compound comprising a CLPTM1L-targeting agent for use in a method of treating or preventing a disease or condition associated with CLPTM1L protein dysfunction, the method comprising administering said targeting agent to a cell, tissue, or organ of a subject at risk of, diagnosed as having, or exhibiting a symptom of the disease or condition, wherein the CLPTM1L-targeting agent is a monoclonal antibody that binds specifically to a peptide having an amino acid sequence of SEQ ID NO:6 or SEQ ID NO:10.
13. The compound for use in the method of claim 12, wherein the CLPTM1L protein dysfunction comprises one or more of pathogenic protein production, pathogenic protein modification, or pathogenic protein function.
14. The compound for use in the method of claim 12, wherein the disease or condition is a tumor, a cancer, or a pre-neoplastic lesion.
15. The compound for use in the method of claim 14, wherein the tumor is a solid tumor selected from the group consisting of glioblastoma, sarcoma, carcinoma, and lymphoma; optionally wherein the tumor exhibits resistance to a chemotherapeutic agent; optionally wherein the chemotherapeutic agent is selected from the group consisting of cisplatin and gemcitabine.
16. The compound for use in the method of claim 12, wherein the compound is administered with a pharmaceutically acceptable carrier.

Patentansprüche

1. CLPTM1L-Targeting-Mittel, wobei das CLPTM1L-Targeting-Mittel ein monoklonaler Antikörper ist, der spezifisch an ein Peptid mit einer Aminosäuresequenz von SEQ ID NO: 6 oder SEQ ID NO: 10 bindet.
2. CLPTM1L-Targeting-Mittel nach Anspruch 1, wobei der monoklonale Antikörper ein chimärer Antikörper, humaner Antikörper, humanisierter Antikörper, rekombinanter Antikörper, gentechnisch veränderter Antikörper, konjugierter Antikörper oder ein Fragment davon ist.
3. CLPTM1L-Targeting-Mittel nach Anspruch 1, wobei der monoklonale Antikörper ein CLPTM1L-gezielter bispezifischer monoklonaler Antikörper ist.
4. CLPTM1L-Targeting-Mittel nach Anspruch 1, wobei der monoklonale Antikörper ein Immunkonjugat ist, das ein Therapeutikum, das aus einem pharmakologischen Mittel, Radioisotop und Toxin ausgewählt ist, und einen Linker umfasst.
5. Pharmazeutische Zusammensetzung, umfassend das Mittel nach einem der Ansprüche 1-4 und einen pharmazeutisch annehmbaren Träger.
6. Mittel nach einem der Ansprüche 1-4 oder pharmazeutische Zusammensetzung nach Anspruch 5 zur Verwendung als ein Medikament.
7. Verbindung, umfassend ein CLPTM1L-Targeting-Mittel zur Verwendung in einem Verfahren zum Behandeln oder Vorbeugen eines Tumors, umfassend Verabreichen an ein Subjekt, das deren bedarf, einer therapeutisch wirksamen Menge der Verbindung an ein Subjekt, wobei das CLPTM1L-Targeting-Mittel ein monoklonaler Antikörper ist, der spezifisch an ein Peptid mit einer Aminosäuresequenz von SEQ ID NO: 6 oder SEQ ID NO: 10 bindet, wodurch der Tumor in dem Subjekt behandelt oder verhindert wird.
8. Verbindung zur Verwendung in dem Verfahren nach Anspruch 7, wobei:
 - (a) der Tumor ein solider Tumor ist, der aus der Gruppe bestehend aus Glioblastom, Sarkom, Karzinom und Lymphom ausgewählt ist; oder
 - (b) der Tumor mit einem Krebs oder einer präneoplastischen Läsion assoziiert ist, der/die aus der Gruppe bestehend aus Lungenkrebs, Bauchspeicheldrüsenkrebs, Prostatakrebs, Hautkrebs, Blasenkrebs, Nierenkrebs, Eierstockkrebs, Darmkrebs, Kolorektalkrebs, Brustkrebs, Gebärmutterhalskrebs, Hirnkrebs, Speiseröhrenkrebs und Magenkrebs oder einer präkanzerösen Läsion davon ausgewählt ist, bevorzugt wobei der Tumor mit Lungenkrebs oder einer präkanzerösen Läsion davon assoziiert ist.

9. Verbindung zur Verwendung in dem Verfahren nach Anspruch 7, wobei das Subjekt ein Mensch ist.
10. Verbindung zur Verwendung in dem Verfahren nach Anspruch 7, wobei der Tumor Resistenz gegen ein Chemotherapeutikum aufweist; optional wobei das Chemotherapeutikum aus der Gruppe bestehend aus Cisplatin und Gemcitabin ausgewählt ist.
11. Verbindung zur Verwendung in dem Verfahren nach Anspruch 7, wobei die Verbindung mit einem pharmazeutisch annehmbaren Träger verabreicht wird.
12. Verbindung, umfassend ein CLPTM1L-Targeting-Mittel zur Verwendung in einem Verfahren eines Behandeln oder Vorbeugens einer Erkrankung oder eines Zustands, die/der mit CLPTM1L-Protein-Dysfunktion assoziiert ist, wobei das Verfahren ein Verabreichen des Targeting-Mittels an eine Zelle, ein Gewebe oder ein Organ eines Subjekts, bei dem das Risiko eines Symptoms der Erkrankung oder des Zustands besteht, das als dieses habend diagnostiziert wird oder dieses aufweist, umfasst, wobei das CLPTM1L-Targeting-Mittel ein monoklonaler Antikörper ist, der spezifisch an ein Peptid mit einer Aminosäuresequenz von SEQ ID NO: 6 oder SEQ ID NO: 10 bindet.
13. Verbindung zur Verwendung in dem Verfahren nach Anspruch 12, wobei die CLPTM1L-Protein-Dysfunktion eines oder mehrerer von pathogener Proteinproduktion, pathogener Proteinmodifikation oder pathogener Proteinfunktion umfasst.
14. Verbindung zur Verwendung in dem Verfahren nach Anspruch 12, wobei die Erkrankung oder der Zustand ein Tumor, ein Krebs oder eine präneoplastische Läsion ist.
15. Verbindung zur Verwendung in dem Verfahren nach Anspruch 14, wobei der Tumor ein solider Tumor ist, der aus der Gruppe bestehend aus Glioblastom, Sarkom, Karzinom und Lymphom ausgewählt ist; optional wobei der Tumor Resistenz gegen ein Chemotherapeutikum aufweist; optional wobei das Chemotherapeutikum aus der Gruppe bestehend aus Cisplatin und Gemcitabin ausgewählt ist.
16. Verbindung zur Verwendung in dem Verfahren nach Anspruch 12, wobei die Verbindung mit einem pharmazeutisch annehmbaren Träger verabreicht wird.

Revendications

1. Agent de ciblage de CLPTM1L, l'agent de ciblage de CLPTM1L étant un anticorps monoclonal qui se lie spécifiquement à un peptide ayant la séquence d'acides aminés de SEQ ID N° : 6 ou SEQ ID N° : 10.
2. Agent de ciblage de CLPTM1L selon la revendication 1, dans lequel l'anticorps monoclonal est un anticorps chimérique, un anticorps humain, un anticorps recombinant, un anticorps modifié, un anticorps conjugué ou un fragment de ceux-ci.
3. Agent de ciblage de CLPTM1L selon la revendication 1, dans lequel l'anticorps monoclonal est un anticorps monoclonal bispécifique ciblé sur CLPTM1L.
4. Agent de ciblage de CLPTM1L selon la revendication 1, dans lequel l'anticorps monoclonal est un immunoconjugué comprenant un agent thérapeutique choisi parmi un agent pharmacologique, un radio-isotope et une toxine ; et un lieu.
5. Composition pharmaceutique comprenant l'agent selon l'une quelconque des revendications 1 à 4 et un support pharmaceutiquement acceptable.
6. Agent selon l'une quelconque des revendications 1 à 4 ou composition pharmaceutique selon la revendication 5, pour une utilisation en tant que médicament.
7. Composé comprenant un agent de ciblage de CLPTM1L pour une utilisation dans un procédé de traitement ou de prévention d'une tumeur comprenant l'administration à un sujet qui la nécessite d'une quantité thérapeutiquement efficace dudit composé à un sujet, l'agent de cible de CLPTM1L étant un anticorps monoclonal qui se lie spécifiquement à un peptide ayant une séquence d'acides aminés de SEQ ID N° : 6 ou SEQ ID N° : 10, la tumeur étant

ainsi traitée ou prévenue chez le sujet.

8. Composé pour une utilisation dans le procédé selon la revendication 7, dans lequel :

- (a) la tumeur est une tumeur solide choisie dans le groupe constitué par un glioblastome, un sarcome, un carcinome et un lymphome ; ou
- (b) la tumeur est associée à un cancer ou une lésion pré-néoplasique choisi dans le groupe constitué par le cancer du poumon, le cancer pancréatique, le cancer de la prostate, le cancer de la peau, le cancer de la vessie, le cancer du rein, le cancer ovarien, le cancer du côlon, le cancer colorectal, le cancer du sein, le cancer du col de l'utérus, le cancer du cerveau, le cancer de l'œsophage et le cancer de l'estomac, ou une lésion précancéreuse de ceux-ci, la tumeur étant de préférence associée a cancer du poumon ou à une lésion précancéreuse de celui-ci.

9. Composé pour une utilisation dans le procédé selon la revendication 7, dans lequel le sujet est un être humain.

10. Composé pour une utilisation dans le procédé selon la revendication 7, dans lequel la tumeur présente une résistance à un agent chimiothérapeutique ; l'agent chimiothérapeutique étant éventuellement choisi dans le groupe constitué par le cisplatine et la gemcitabine.

11. Composé pour une utilisation dans le procédé selon la revendication 7, le composé étant administré avec un support pharmaceutiquement acceptable.

12. Composé comprenant un agent de ciblage de CLPTM1L pour une utilisation dans un procédé de traitement ou de prévention d'une maladie ou affection associée à un dysfonctionnement de la protéine CLPTM1L, le procédé comprenant l'administration dudit agent de ciblage à une cellule, un tissu ou un organe présentant un risque, un diagnostic ou un symptôme de la maladie ou affection, l'agent de ciblage de CLPTM1L étant un anticorps monoclonal qui se lie spécifiquement à un peptide ayant une séquence d'acides aminés de SEQ ID N° : 6 ou SEQ ID N° : 10.

13. Composé pour une utilisation dans le procédé selon la revendication 12, dans lequel le dysfonctionnement de la protéine CLPTM1L comprend au moins l'une d'une production de protéine pathogène, d'une modification de protéine pathogène ou d'une fonction de protéine pathogène.

14. Composé pour une utilisation dans le procédé selon la revendication 12, dans lequel la maladie ou affection est une tumeur, un cancer ou une lésion pré-néoplasique.

15. Composé pour une utilisation dans le procédé selon la revendication 14, dans lequel la tumeur est une tumeur solide choisie dans le groupe constitué par un glioblastome, un sarcome, un carcinome et un lymphome ; la tumeur présentant éventuellement une résistance à un agent chimiothérapeutique ; l'agent chimiothérapeutique étant éventuellement choisi dans le groupe constitué par le cisplatine et la gemcitabine.

16. Composé pour une utilisation dans le procédé selon la revendication 12, dans lequel le composé est administré avec un support pharmaceutiquement acceptable.

TMHMM posterior probabilities for Sequence

A




FIG. 1

B




FIG. 1 continued




FIG. 2

B.

CLPTM1L -008 Isoform




FIG. 2 continued

Sequence analysis for CLPTH1L







FIG. 3




B



C



D



Summary:

	Mean	SEM	N	t-test
normal	0	0	16	
tumor	1.456784	0.124848	31	$p < 0.05$

Intensity scoring - 0=negative 1=weak 2=moderate 3=strong

FIG. 5

B Overall Survival by cipt tertile




FIG. 5 continued



C

FIG. 5 continued

A




FIG. 6

B




FIG. 6 continued

C

Panc1 PDAC Cells




FIG. 6 continued

D

Monoclonal antibody 10-2 treatment of
PDAC cells




FIG. 6 continued




FIG. 7





FIG. 7 continued




B.

FIG. 8 continued

A.




FIG. 9

B.




FIG. 9 continued

Characterization of in-vitro anti-tumor effects of monoclonal anti-CLPTM1L antibodies

mAb	Targets Both Isoforms	IF Staining Intensity A549	Punctate Surface Staining A549	pAkt Inhibition		Bcl-xL Inhibition		CDDP Sensitization		Tumor Spheroid Growth H838
				A549	H838	A549	H838	A549	H838	
6-1	-	+++	-	++	+/-	+	-	+	+	-
6-2	-	+	-	+	+/-	+	-	-	+/-	-
7-1	-	++	-	+/-	-	+/-	-	-	-	-
7-2	-	+	-	-	-	+/-	-	-	+/-	-
8	+	++	+	+/-	-	-	-	+	+	-
9-1	+	++	+	+	-	+	-	+/-	+	-
9-2	+	+	+	-	+/-	+/-	+/-	+/-	-	-
10-1	+	++	+	+	+/-	-	+/-	-	+/-	+/-
10-2	+	++	+	++	+	+	+	+/-	+/-	+
10-3	+	+++	+	++	+	+	+	+/-	+/-	+

FIG. 10

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5565332 A [0022]
- US 6824780 B [0023]
- US 5567610 A [0023]
- US 5229275 A [0023]
- US 5545806 A [0023]
- US 5569825 A [0023]
- US 5591669 A [0023]
- US 5545807 A [0023]
- WO 9717852 A [0023]
- US 6300064 B [0024]
- US 5969108 A [0024]
- US 4816567 A [0035]
- WO 9616673 A [0035]
- US 5837234 A [0035]
- WO 9802463 A [0035]
- US 5821337 A [0035]
- US 5869046 A [0036]

Non-patent literature cited in the description

- JAMES et al. *PLoS ONE*, 2012, vol. 7 (6), e36116 [0004]
- ZHENHUA et al. *PLoS ONE*, 2012, vol. 7 (12), e52598 [0004]
- OMASITS et al. *Bioinformatics*, 2014, vol. 30 (6), 884-6 [0014]
- SKERRA ; PLUCKTHUN. *Science*, 1988, vol. 240, 1038-41 [0022]
- BIRD et al. *Science*, 1988, vol. 242, 423-26 [0022]
- HUSTON et al. *Proc. Natl. Acad. Sci. USA*, 1988, vol. 85, 5879-83 [0022]
- Antibody Engineering. IRL Press, 1996 [0022][0024]
- JAKOBOVITS et al. *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 2551 [0023]
- JAKOBOVITS et al. *Nature*, 1993, vol. 362, 255-258 [0023]
- BRUGGEMANN et al. *Year in Immuno.*, 1993, vol. 7, 33 [0023]
- Epitope Mapping Protocols in Methods in Molecular Biology. 1996, vol. 66 [0027]
- SCATCHARD et al. *Ann. N.Y. Acad. Sci. USA*, 1949, vol. 51, 660 [0027]
- SMAGLO et al. *Nat. Rev. Clin. Oncol.*, 2014, vol. 11 (11), 637-48 [0032]
- SMAGLO et al. *Nature Reviews Clin. Oncol.*, 2014, vol. 11, 637-48 [0033]
- JONES et al. *Nature*, 1986, vol. 321, 522-525 [0035]
- REICHMANN et al. *Nature*, 1988, vol. 332, 323-327 [0035]
- VERHOEYEN et al. *Science*, 1988, vol. 239, 1534-1536 [0035]
- MILLSTEIN et al. *Nature*, 1983, vol. 305, 537-539 [0035]
- VINCENT ; ZURINI. *Biotechnol. J.*, 2012, vol. 7 (12), 1444-50 [0036]
- KANEKO ; NIWA. *Biodrugs*, 2011, vol. 25 (1), 1-11 [0036]
- CHAMES et al. *mAbs*, 2009, vol. 1 (6), 539-547 [0037]
- TUTT et al. *J. Immunol.*, 1991, vol. 147, 60 [0039]
- GLEASON et al. *Mol. Cancer. Ther.*, 2012, vol. 11 (12), 2674-84 [0039]
- WANG et al. *J Biochem.*, 2004, vol. 135 (4), 555-65 [0040]
- CARON et al. *J. Exp Med.*, 1992, vol. 176, 1191-1195 [0041]
- WOLFF et al. *Cancer Research*, 1993, vol. 53, 2560-2565 [0041]
- STEVENSON et al. *Anti-Cancer Drug Design*, 1989, vol. 3, 219-230 [0041]
- KENDERIAN et al. *Cancer Res.*, 2014, vol. 74 (22), 6383-9 [0042]
- GROVES et al. *J. Immunol. Methods*, 2006, vol. 313, 129-39 [0043]
- BARBAS et al. *Proc. Natl. Acad. Sci. USA*, 1994, vol. 91, 3809-13 [0043]
- ALMAGRO ; STROHL. Antibody Engineering: Humanization. *Affinity Maturation and Selection Methods*, 307-327 [0043]
- Therapeutic Monoclonal Antibodies: From Bench to Clinic. John Wiley & Sons, Inc, 2009 [0043]
- CHOW ; HO. *Sci Transl Med*, 2013, vol. 5, 216rv4 [0045]
- MONTENEGRO et al. *Adv. Drug Delivery Rev.*, 2013, vol. 65, 677-688 [0045]
- ROHLOFF et al. *Molecular Therapy Nucleic Acids*, 2014, vol. 3, e201 [0048]
- ZHU et al. *Theranostics*, 2014, vol. 4 (9), 931-944 [0048]
- ELLINGTON ; SZOSTAK. *Nature*, 1990, vol. 346, 818-822 [0048]

EP 3 094 654 B1

- Remington's Pharmaceutical Sciences. Mack Printing Company, 1990, 1289-1329 [0049]
- **WEINER et al.** *Nature Rev. Immunol.*, 2010, vol. 10, 317-327 [0064]
- **SLANEY et al.** *Cancer Res.*, 2014, vol. 74, 7168-7174 [0066]
- **FREIREICH et al.** *Cancer Chemotherapy Rep.*, 1966, vol. 50 (4), 219-244 [0074]
- Estimating the safe starting dose in clinical trials for therapeutics in adult healthy volunteers. Center for Drug Evaluation and Research, Center for Biologics Evaluation and Research. U.S. Food and Drug Administration, 2002 [0074]
- **THERASSE et al.** *J. Natl. Cancer Inst.*, 2000, vol. 92, 205-16 [0077]
- **CHEN et al.** *J Proteome Res.*, 2009, vol. 8, 651-61 [0085]
- **NI et al.** *PLoS One*, 2012, vol. 7, e52598 [0086]