



(86) Date de dépôt PCT/PCT Filing Date: 2009/06/05

(87) Date publication PCT/PCT Publication Date: 2009/12/10

(45) Date de délivrance/Issue Date: 2019/11/26

(85) Entrée phase nationale/National Entry: 2010/12/02

(86) N° demande PCT/PCT Application No.: US 2009/046421

(87) N° publication PCT/PCT Publication No.: 2009/149359

(30) Priorité/Priority: 2008/06/06 (US61/059,678)

(51) Cl.Int./Int.Cl. *G01N 33/574* (2006.01)

(72) Inventeurs/Inventors:

STAUDT, LOUIS M., US;

RIMSZA, LISA, US;

LISTER, ANDREW T., GB;

WEISENBURGER, DENNIS, US;

DELABIE, JAN, US;

...

(73) Propriétaires/Owners:

THE UNITED STATES OF AMERICA, AS

REPRESENTED BY THE SECRETARY, DEPT. OF
HEALTH AND HUMAN SERVICES, US; ...

(54) Titre : INDICE DE SURVIE POUR UN LYMPHOME A GRANDES CELLULES B DIFFUS

(54) Title: SURVIVAL PREDICTOR FOR DIFFUSE LARGE B CELL LYMPHOMA

(57) Abrégé/Abstract:

The invention provides methods and materials related to a gene expression-based survival predictor for diffuse large B cell lymphoma (DLBCL) patients.

- (72) **Inventeurs(suite)/Inventors(continued)**: SMELAND, ERLEND B., NO; HOLTE, HARALD, NO; KVALOY, STEIN, NO; BRAZIEL, RITA M., US; FISHER, RICHARD I., US; JARES, PEDRO, US; LOPEZ-GUILLERMO, ARMANDO, ES; GUERRI, ELIAS CAMPO, ES; JAFFE, ELAINE S., US; LENZ, GEORGE, DE; WILSON, WYNDHAM H., US; WRIGHT, GEORGE, US; DAVE, SANDEEP S., US; GASCOYNE, RANDY D., CA; CONNORS, JOSEPH M., CA; MULLER-HERMELINK, HANS-KONRAD, DE; ROSENWALD, ANDREAS, DE; OTT, GERMAN, DE; CHAN, WING C., US
- (73) **Propriétaires(suite)/Owners(continued)**:
ARIZONA BOARD OF REGENTS ON BEHALF OF THE UNIVERSITY OF ARIZONA, US;
QUEEN MARY AND WESTFIELD COLLEGE, UNIVERSITY OF LONDON, GB;
BOARD OF REGENTS OF THE UNIVERSITY OF NEBRASKA, US; OSLO UNIVERSITY HOSPITAL HF, NO;
OREGON HEALTH & SCIENCE UNIVERSITY, US; UNIVERSITY OF ROCHESTER, US;
HOSPITAL CLINIC DE BARCELONA, ES; UNIVERSITAT DE BARCELONA, ES;
BRITISH COLUMBIA CANCER AGENCY BRANCH, CA; JULIUS-MAXIMILIANS-UNIVERSITY OF WUERZBURG, DE
- (74) **Agent**: RIDOUT & MAYBEE LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 December 2009 (10.12.2009)

PCT

(10) International Publication Number
WO 2009/149359 A3

(51) International Patent Classification:
G01N 33/574 (2006.01)

(21) International Application Number:
PCT/US2009/046421

(22) International Filing Date:
5 June 2009 (05.06.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/059,678 6 June 2008 (06.06.2008) US

(71) Applicant (for all designated States except US): **THE UNITED STATES OF AMERICA, AS REPRESENTED BY THE SECRETARY, DEPT. OF HEALTH AND HUMAN SERVICES** [US/US]; Office Of Technology Transfer, National Institutes of Health, 6011 Executive Boulevard, Suite 325, MSC 7660, Bethesda, MD 20892-7660 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **STAUDT, Louis, M.** [US/US]; 9309 Greyrock Road, Silver Springs, MD 20910 (US).

(74) Agents: **RIVAS, Marcos, P.** et al.; Leydig, Voit & Mayer, Ltd., Two Prudential Plaza, Suite 4900, 180 North Stetson Avenue, Chicago, IL 60601-6731 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(88) Date of publication of the international search report:
25 February 2010

(54) Title: SURVIVAL PREDICTOR FOR DIFFUSE LARGE B CELL LYMPHOMA

(57) Abstract: The invention provides methods and materials related to a gene expression-based survival predictor for diffuse large B cell lymphoma (DLBCL) patients.

WO 2009/149359 A3

SURVIVAL PREDICTOR FOR DIFFUSE LARGE B CELL LYMPHOMA

CROSS-REFERENCE TO RELATED APPLICATION

STATEMENT REGARDING US FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0001] This invention was made with US Government support under project number Z01 ZIA BC 011006 by the National Institutes of Health, National Cancer Institute. This invention was made with US government support under grant no. U01 CA084967, awarded by the National Institutes of Health. The US Government has certain rights in the invention.

INCORPORATION-BY-REFERENCE OF MATERIAL SUBMITTED ELECTRONICALLY

[0002] Incorporated by reference in its entirety herein is a computer-readable nucleotide sequence listing submitted concurrently herewith and identified as follows: One 1,231,630 Byte ASCII (Text) file named "704777_ST25.TXT," created on June 5, 2009.

BACKGROUND OF THE INVENTION

[0003] The current standard of care for the treatment of diffuse large B cell lymphoma (DLBCL) includes anthracycline-based chemotherapy regimens such as CHOP in combination with the administration of the anti-CD20 monoclonal antibody Rituximab. This combination regimen (R-CHOP) can cure about 60% of patients and has improved the overall survival of DLBCL patients by 10-15% (Coiffier et al., *N. Engl. J. Med.*, 346: 235-42 (2002)). Nonetheless, the molecular basis of response or resistance to this therapy is unknown.

[0004] DLBCL is a molecularly heterogeneous disease (Staudt et al., *Adv. Immunol.*, 87: 163-208 (2005)), and different molecular subtypes of DLBCL can have very different prognoses following treatment. For example, gene expression profiling has identified two molecular subtypes of DLBCL that are biologically and clinically distinct (Rosenwald et al., *N. Engl. J. Med.*, 346: 1937-47 (2002); Alizadeh et al., *Nature*, 403: 503-11 (2000)). The germinal center B cell-like (GCB) DLBCL subtype likely arises from normal germinal center B cells, whereas the activated B cell-like (ABC) DLBCL subtype may arise from a post-germinal center B cell that is blocked during plasmacytic differentiation. Many oncogenic mechanisms distinguish these subtypes: GCB DLBCLs have recurrent t(14,18) translocations, whereas ABC DLBCLs have recurrent trisomy 3 and deletion of the INK4a/ARF locus as well as constitutive activation of the anti-apoptotic NF- κ B signalling pathway (Rosenwald et al., *N. Engl. J. Med.*, 346: 1937-47 (2002); Bea et al., *Blood*, 106: 3183-90 (2005); Tagawa et al., *Blood*, 106: 1770-77 (2005); Davis et al., *J. Exp. Med.*, 194:1861-74 (2001); Ngo et al., *Nature*, 441: 106-10 (2006); Lenz et al., *Science*, 319: 1676-79 (2008)). When treated with CHOP-like chemotherapy, the overall survival rates of patients with GCB DLBCL and ABC DLBCL were 60% and 30%, respectively (Wright et

al., *Proc. Nat'l. Acad. Sci. USA*, 100: 9991-96 (2003)). Thus, the prognosis for different DLBCL can vary widely.

[0005] A separate analytical approach identified four gene expression signatures that reflect distinct DLBCL tumor attributes and that were associated with distinct survival profiles in CHOP-treated DLBCL patients (Rosenwald et al., *N. Engl. J. Med.*, 346: 1937-47 (2002)). A “germinal center B cell” (GCB) signature was associated with a favorable prognosis and paralleled the distinction between ABC and GCB DLBCL. The “proliferation” signature was associated with an adverse prognosis and included MYC and its target genes. The “MHC class II” signature was silenced in the malignant cells in a subset of DLBCL cases, an event that was associated with inferior survival (Rosenwald et al., *N. Engl. J. Med.*, 346: 1937-47 (2002); Rimsza et al., *Blood*, 103: 4251-58 (2004)). A fourth prognostic signature, termed “lymph node” signature was associated with favorable prognosis and included components of the extracellular matrix, suggesting that it reflects the nature of the tumor-infiltrating non-malignant cells. These signatures predicted survival in a statistically independent fashion, indicating that multiple biological variables dictate the response to CHOP chemotherapy in DLBCL.

[0006] Reports have suggested that the benefit of Rituximab immunotherapy might be restricted to certain molecular subtypes of DLBCL. High expression of BCL-2 or low expression of BCL-6 was associated with inferior survival with CHOP therapy. However, this distinction disappeared with R-CHOP therapy (Mounier et al., *Blood*, 101: 4279-84 (2003); Winter et al., *Blood*, 107: 4207-13 (2006)). Immunohistochemistry has also been used to distinguish DLBCLs with a germinal center versus post-germinal center phenotype. Although such immunohistochemical phenotypes were prognostically significant in CHOP-treated cases, they were not prognostic for R-CHOP-treated cases (Nyman et al., *Blood*, 109: 4930-35 (2007)).

[0007] Accordingly, there is a need for new methods of distinguishing among DLBCL subtypes that is prognostically significant for R-CHOP-treated patients.

BRIEF SUMMARY OF THE INVENTION

[0008] The invention provides methods and arrays related to a gene expression-based survival predictor for DLBCL patients, including patients treated with the current standard of care, which includes chemotherapy and the administration of Rituximab.

[0009] The invention provides a method of predicting the survival outcome of a subject suffering from diffuse large B cell lymphoma (DLBCL) that includes obtaining a gene

expression profile from one or more DLBCL biopsy samples from the subject. The gene expression profile, which can be derived from gene expression product isolated from the one or more biopsy samples, includes an expression level for each gene in a germinal center B cell (GCB) gene expression signature and each gene in a stromal-1 gene expression signature. From the gene expression profile, a GCB signature value and a stromal-1 signature value are derived. From these values, a survival predictor score can be calculated using an equation that includes subtracting $[(x) * (\text{the GCB signature value})]$ and subtracting $[(y) * (\text{the stromal-1 signature value})]$. In the equation, (x) and (y) are scale factors. A lower survival predictor score indicates a more favorable survival outcome, and a higher survival predictor score indicates a less favorable survival outcome for the subject.

[0010] The invention also provides a method of generating a survival estimate curve for subjects suffering from DLBCL. Generally the method includes obtaining a gene expression profile from one or more DLBCL biopsy samples taken from each member of a plurality of subjects. Each gene expression profile, which can be derived from gene expression product isolated from the one or more biopsy samples taken from each subject, includes an expression level for each gene in a GCB expression signature, a stromal-1 gene expression signature, and a stromal-2 gene expression signature. For each subject, the GCB signature value, the stromal-1 signature value, and the stromal-2 signature value are determined from the subject's gene expression profile, and, for each subject, a survival predictor score is generated. Each subject's survival outcome following treatment for DLBCL is tracked. A survival estimate curve is generated which correlates the probability of the tracked survival outcome with time following treatment for DLBCL and which also correlates the tracked outcome over time with the survival predictor score for the subjects.

[0011] The invention additionally provides a method of predicting the survival outcome of a subject suffering from DLBCL. Generally, the method includes obtaining a gene expression profile from one or more DLBCL biopsy samples from the subject. The gene expression profile, which can be derived from gene expression product isolated from the one or more biopsy samples, includes an expression level for each gene in a GCB gene expression signature, each gene in a stromal-1 gene expression signature, and each gene in a stromal-2 gene expression signature. The GCB signature value, the stromal-1 signature value, and the stromal-2 signature value are determined from the gene expression profile. The method then includes calculating a survival predictor score using the equation:

survival predictor score = $A - [(x) * (\text{the GCB signature value})] - [(y) * (\text{the stromal-1 signature value})] + [(z) * (\text{the stromal-2 signature value})]$.

In this equation, A is an offset term, and (x), (y), and (z) are scale factors. The method further includes calculating the probability of a survival outcome for the subject beyond an amount of time t following treatment for DLBCL, wherein the subject's probability of the survival outcome P(SO) is calculated using the equation:

$$P(SO) = SO_0(t)^{(\exp((s) * \text{survival predictor score}))}$$

In this equation, $SO_0(t)$ is the probability of the survival outcome, which corresponds to the largest time value smaller than t in a survival outcome curve, and wherein (s) is a scale factor.

[0012] Furthermore, the invention provides a method of evaluating a subject for antiangiogenic therapy of DLBCL. The method includes obtaining a gene expression profile from one or more DLBCL biopsy samples from the subject. The gene expression profile, which can be derived from gene expression product isolated from the one or more biopsy samples, includes an expression level for each gene in a stromal-2 signature. The subject's stromal-2 signature value is then derived from the gene expression profile and evaluated to determine whether the subject's stromal-2 signature value is higher or lower than a standard stromal-2 value. If the subject's stromal-2 signature value is higher than the standard stromal-2 value, then antiangiogenic therapy is indicated, and the subject can be treated with antiangiogenic therapy. If the subject's stromal-2 signature value is not higher than the standard stromal-2 value, then antiangiogenic therapy is not indicated.

[0013] The invention also provides a second method of evaluating a subject for antiangiogenic therapy of DLBCL. The method includes obtaining a gene expression profile from one or more DLBCL biopsy samples from the subject. The gene expression profile, which can be derived from gene expression product isolated from the one or more biopsy samples, includes an expression level for each gene in a stromal-1 signature and in a stromal-2 signature. The subject's stromal-1 signature value and stromal-2 signature value are then derived from the gene expression profile. The stromal-1 signature value is subtracted from the stromal-2 signature value to thereby obtain the subject's stromal score. The subject's stromal score is evaluated to determine whether it is higher or lower than a standard stromal score. If the subject's stromal score is higher than the standard stromal score, then antiangiogenic therapy is indicated, and the subject can be treated with antiangiogenic therapy. If the subject's stromal score is not higher than the standard stromal score, then antiangiogenic therapy is not indicated.

[0014] Additionally, the invention provides a machine-readable medium containing a digitally encoded GCB signature value, a digitally encoded stromal-1 signature value, a digitally encoded stromal-2 signature, or any combination of the foregoing signature values obtained from a subject suffering from DLBCL.

[0015] In another embodiment the invention provides a machine-readable medium containing the digitally encoded survival predictor score obtained using a method disclosed herein for predicting the survival outcome of a subject suffering from diffuse large B cell lymphoma (DLBCL). In yet another embodiment, the invention provides a machine-readable medium containing the survival estimate curve obtained using a method disclosed herein for generating a survival estimate curve for subjects suffering from DLBCL. In still another embodiment, the invention provides a machine-readable medium containing the digitally encoded probability of survival calculated according to a method disclosed herein for predicting the survival outcome (e.g., progression-free survival or overall survival) of a subject suffering from DLBCL. Furthermore, the invention provides a machine-readable medium containing the digitally encoded stromal score generated by a method disclosed herein for evaluating a subject for antiangiogenic therapy of DLBCL.

[0016] The invention also provides a targeted array comprising at least one probe or at least one set of probes for each gene in a germinal center B cell gene (GCB) expression signature, a stromal-1 gene expression signature, and a stromal-2 gene expression signature. The array can include probes for fewer than 20,000 genes or fewer than 10,000 genes.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] Figure 1A is a Kaplan-Meier estimates plot depicting the probability of progression-free-survival versus time (in years) of patients with GCB DLBCL and ABC DLBCL. The plot indicates that GCB patients have a more favorable, i.e., higher probability of progression-free survival rate than ABC patients for at least five years following R-CHOP therapy.

[0018] Figure 1B a Kaplan-Meier estimates plot depicting the probability of overall survival versus time (in years) of patients with GCB DLBCL and ABC DLBCL. The plot indicates that GCB patients have a more favorable, i.e., higher probability, of overall survival than ABC patients for at least five years following R-CHOP therapy.

[0019] Figure 1C is a series of four Kaplan-Meier estimates plots depicting the probabilities of overall survival versus time (in years) in DLBCL patients. Each of the four plots correlates the probability of overall survival with the lymph node/stromal-1, germinal

center B cell, proliferation, or MHC class II gene expression signature, respectively.

Moreover, in each plot, the average expression of the signature genes in each biopsy sample was used to rank cases and divide the cohort into quartile groups as indicated.

[0020] Figure 2A is a pair of Kaplan-Meier estimates plots depicting the probability of progression-free-survival and the probability of overall survival, as indicated, versus time (in years) among DLBCL patients treated with R-CHOP. Patient samples were ranked according to a bivariate model created using the germinal center B cell (GCB) and stromal-1 signatures and divided into quartile groups.

[0021] Figure 2B is a pair of Kaplan-Meier plots depicting the probability of progression-free-survival and the probability of overall survival, as indicated, versus time (in years) among DLBCL patients treated with R-CHOP. Patient samples were ranked according to a survival predictor score derived from a model incorporating the germinal center B cell, stromal-1, and stromal-2 signatures and divided into quartile groups.

[0022] Figure 2C is a series of three Kaplan-Meier estimates plots depicting the probability of overall survival versus time (in years) among R-CHOP treated DLBCL patients in the indicated low, intermediate, or high IPI risk groups. Patient samples were stratified according to the same survival predictor score used in Figure 2B, except that the first and second quartiles were merged, and the third and fourth quartiles were merged.

[0023] Figure 3 depicts the expression levels of the indicated GCB cell, stromal-1, and stromal-2 signature genes in ABC, GCB, and unclassified DLBCL biopsy samples. Relative levels of gene expression are depicted according to the scale shown. Shown at the bottom are the signature averages for each patient. Also shown is the stromal score, which is the component of the survival model contributed by the difference between the stromal-2 and stromal-1 signature averages. The survival predictor score is shown for each patient and was used to order the cases, after grouping into ABC DLBCL, GCB DLBCL, and unclassified categories.

[0024] Figure 4A depicts the relative gene expression of stromal-1, stromal-2, and germinal center B cell signatures in CD19+ malignant and CD19- non-malignant subpopulations of cells isolated from three biopsy specimens from patients with DLBCL. Stromal-1 and stromal-2 signature genes were more highly expressed in the non-malignant cells, whereas the germinal center B cell signature genes were more highly expressed in the malignant cells. The log2 ratios of gene expression levels in the CD19- subpopulation to those in the CD19+ subpopulations are depicted according to the scale shown.

[0025] Figure 4B depicts the results of gene enrichment analysis comparing the stromal-1 gene signature with mesenchyme-1 and mesenchyme-2 signatures (from normal mesenchymal origin cells), with a monocyte signature expressed more highly in normal blood monocytes than in blood B, T, and NK cells, and in a pan-T cell signature expressed more highly in blood T cells than in blood B cells, NK cells, and monocytes. While a relationship was seen between stromal-1 signature and mesenchyme-1, mesenchyme-2, and monocyte signatures, no relationship was observed between the stromal-1 signature and a pan-T cell signature expressed more highly in blood T cells than in blood B cells, NK cells, and monocytes. The relative levels of gene expression are depicted according to the scale shown.

[0026] Figures 5A is a Kaplan-Meier estimates plot depicting the probability of overall survival versus time (in years) in DLBCL cases segregated according to SPARC protein expression levels, as indicated.

[0027] Figure 5B is a pair of images showing the identification of tumor blood vessels by immunohistochemical analysis of CD34+ endothelial cells in representative DLBCL biopsies having low or high blood vessel density (CD34+ objects/ μM^2), as indicated.

[0028] Figure 5C is a plot depicting the correlation between the tumor blood vessel density and the stromal score in analyzed DLBCL biopsies.

[0029] Figures 6A is a Kaplan-Meier estimates plot depicting the probability of overall survival versus time (in years) for "LLMPP CHOP" patients with DLBCL following therapy. The plot indicates that in this cohort, patients with GCB DLBCL show significantly superior overall survival compared to patients with ABC DLBCL following CHOP therapy.

[0030] Figure 6B is a Kaplan-Meier estimates plot depicting the probability of overall survival versus time (in years) for "MMMLNP CHOP" patients with DLBCL following therapy. In this cohort, patients with GCB DLBCL show significantly superior overall survival compared to patients with ABC DLBCL following CHOP therapy.

[0031] Figure 7 is a set of four Kaplan-Meier estimates plots depicting the probability of overall survival versus time (in years) in a "MMMLNP CHOP" cohort. Each of the four plots correlates the probability of overall survival with the lymph node/stromal-1, germinal center B cell, proliferation, or MHC class II gene expression signature, respectively. Moreover, in each plot, the average expression of the signature genes in each biopsy sample was used to rank cases and divide the cohort into quartile groups as indicated.

[0032] Figure 8A is a Kaplan-Meier estimates plot depicting the probability of overall survival versus time (in years) in a "LLMPP CHOP" cohort, which was divided according to

MHC class II signature expression levels. Patients with low MHC class II signature expression have significantly inferior overall survival compared to patients with normal MHC class II expression.

[0033] Figure 8B is a Kaplan-Meier estimates plot depicting the probability of overall survival versus time (in years) in a "MMMLNP CHOP" cohort, which was divided according to MHC class II signature expression levels. Patients with low MHC class II signature expression have significantly inferior overall survival compared to patients with normal MHC class II expression.

[0034] Figure 8C is a Kaplan-Meier estimates plot depicting the probability of overall survival versus time (in years) in a "LLMPP R-CHOP" cohort, which was divided according to MHC class II signature expression levels. There was no significant difference in the overall survival of patients with low MHC class II signature expression as compared to patients with normal MHC class II expression.

[0035] Figure 9A is a pair of Kaplan-Meier estimates plots depicting the probabilities of progression-free survival or overall survival, as indicated, versus time (in years) among patients grouped into quartiles according to a gene expression model consisting of stromal-1 signature, GCB signature, and signature 122 following R-CHOP therapy.

[0036] Figure 9B is a pair Kaplan-Meier estimates plots depicting the probabilities of overall survival versus time (in years) among "MMMLNP CHOP" cohort patients grouped into quartiles according to a gene expression model consisting of either stromal-1 signature and GCB signature or stromal-1, GCB signature, and signature 122, as indicated, following CHOP therapy.

[0037] Figure 9C is a Kaplan-Meier estimates plot depicting the probabilities of overall survival versus time (in years) among "MMMLNP CHOP" cohort patients grouped into quartiles according to a gene expression model consisting of stromal-1 signature, GCB signature, and stromal-2 signature following CHOP therapy.

[0038] Figure 10A is a Kaplan-Meier estimates plot depicting the overall survival among low revised International Prognostic Index (IPI) risk group patients stratified according to the gene expression-based outcome predictor score. After grouping patients into quartiles according to gene expression-based outcome predictor score, quartiles 1 and 2 were merged (Low Model Score), and quartiles 3 and 4 were merged (High Model Score).

[0039] Figure 10B is a Kaplan-Meier estimates plot depicting the overall survival among intermediate revised International Prognostic Index (IPI) risk group patients stratified

according to the gene expression-based outcome predictor. After grouping patients into quartiles according to gene expression-based outcome predictor score, quartiles 1 and 2 were merged (Low Model Score), and quartiles 3 and 4 were merged (High Model Score).

[0040] Figure 10C is a Kaplan-Meier estimates plot depicting the overall survival among high revised International Prognostic Index (IPI) risk group patients stratified according to the gene expression-based outcome predictor. After grouping patients into quartiles according to gene expression-based outcome predictor score, quartiles 1 and 2 were merged (Low Model Score), and quartiles 3 and 4 were merged (High Model Score).

[0041] Figure 11 depicts normal mesenchymal-1 and normal mesenchymal-2 signature gene expression in various normal tissues.

DETAILED DESCRIPTION OF THE INVENTION

[0042] The invention provides a gene expression-based survival predictor for DLBCL patients, including those patients receiving the current standard of care, R-CHOP. The survival predictor can be used to determine the relative probability of a survival outcome in a specific subject. The survival predictor can also be used to predict; i.e., determine the expected probability that a survival outcome will occur by a defined period following treatment for DLBCL. Such prognostic information can be very useful to both the patient and the physician. Patients with survival predictor scores that indicate inferior outcome with R-CHOP therapy could be candidates for a different therapeutic regimen, if, for example, they relapse from R-CHOP treatment. The survival predictor can also be used in the design of clinical studies and analysis of clinical data to provide a quantitative survey of the types of DLBCL patients from which clinical data was gathered. The predictor can be used to improve one or more comparisons between data from different sources (e.g., from different clinical trials), by enabling comparisons with respect to patient characteristics, which are manifested in the gene expression levels that determine and, thus, are embodied in the predictor. Furthermore, the invention provides information that can be very valuable to a DLBCL patient, since the patient may be inclined to order his or her life quite differently, depending on whether the patient has a high or low probability of surviving and/or remaining progression-free for a period of time following treatment.

[0043] The following abbreviations are used herein: ABC, activated B cell-like diffuse large B cell lymphoma; CHOP, cyclophosphamide, doxorubicine, vincristine, and prednisone; CI, confidence interval; COP, cyclophosphamide, vincristine, and prednisone; DLBCL, diffuse large B cell lymphoma; DOD, dead of disease; ECOG, Eastern Cooperative

Oncology Group; FACS, fluorescence-activated cell sorting; FH, follicular hyperplasia; FISH, fluorescence *in situ* hybridization; FL, follicular lymphoma; GC, germinal center; GCB, germinal center B cell-like diffuse large B cell lymphoma; IPI, International Prognostic Index; LPC, lymphoplasmacytic lymphoma; MHC, major histocompatibility complex; NA, not available or not applicable; NK, natural killer; PCR, polymerase chain reaction; RQ-PCR, real-time quantitative PCR; RT-PCR, reverse transcriptase polymerase chain reaction; SAGE, serial analysis of gene expression; WHO, World Health Organization.

[0044] The term "R-CHOP" as used herein refers generally to any therapeutic regimen that includes chemotherapy and the administration of Rituximab. Accordingly, while the term can refer to a Rituximab combination therapy that includes a CHOP regimen of cyclophosphamide, doxorubicine, vincristine, and prednisone, the term R-CHOP can also refer to therapy that includes Rituximab in combination with a chemotherapeutic regimen other than CHOP.

[0045] The phrase "gene expression data" as well as "gene expression level" as used herein refers to information regarding the relative or absolute level of expression of a gene or set of genes in a cell or group of cells. The level of expression of a gene may be determined based on the level of RNA, such as mRNA, encoded by the gene. Alternatively, the level of expression may be determined based on the level of a polypeptide or fragment thereof encoded by the gene. Gene expression data may be acquired for an individual cell, or for a group of cells such as a tumor or biopsy sample. Gene expression data and gene expression levels can be stored on computer readable media, e.g., the computer readable medium used in conjunction with a microarray or chip reading device. Such gene expression data can be manipulated to generate gene expression signatures.

[0046] The term "microarray," "array," or "chip" refers to a plurality of nucleic acid probes coupled to the surface of a substrate in different known locations. The substrate is preferably solid. Microarrays have been generally described in the art in, for example, U.S. Patent Nos. 5,143,854 (Pirrung), 5,424,186 (Fodor), 5,445,934 (Fodor), 5,677,195 (Winkler), 5,744,305 (Fodor), 5,800,992 (Fodor), and 6,040,193 (Winkler), and Fodor et al., *Science*, 251: 767-777 (1991).

[0047] The term "gene expression signature" or "signature" as used herein refers to a group of coordinately expressed genes. The genes making up this signature may be expressed in a specific cell lineage, stage of differentiation, or during a particular biological response. The genes can reflect biological aspects of the tumors in which they are expressed,

such as the cell of origin of the cancer, the nature of the non-malignant cells in the biopsy, and the oncogenic mechanisms responsible for the cancer (Shaffer et al., *Immunity*, 15: 375-385 (2001)). Examples of gene expression signatures include lymph node, proliferation (Rosenwald et al., *New Engl. J. Med.*, 346: 1937-1947 (2002)), MHC class II, ABC DLBCL high, B cell differentiation, T-cell, macrophage, immune response-1, and immune response-2 signatures (U.S. Patent Application Publication No. 2007/0105136 (Staudt)).

[0048] The term “signature value” as used herein corresponds to a mathematical combination of measurements from expression levels of the genes in a gene expression signature. An exemplary signature value is a signature average which corresponds to the average or mean of the individual expression levels in a gene expression signature.

[0049] The phrase “survival predictor score” as used herein refers to a score generated by a multivariate model used to predict survival based on gene expression. A subject with a higher survival predictor score is predicted to have poorer survival than a subject with a lower survival predictor score.

[0050] The term “survival” or “overall survival” as used herein may refer to the probability or likelihood of a subject surviving for a particular period of time. Alternatively, it may refer to the likely term of survival for a subject, such as expected mean or median survival time for a subject with a particular gene expression pattern.

[0051] The term “progression free survival” as used herein can refer to the probability or likelihood of a subject surviving without significant progression or worsening of disease for a particular period of time. Alternatively, it may refer to the likely term for a subject of survival without significant progression or worsening of disease, such as expected mean or median survival time for a subject with a particular gene expression pattern without significant progression or worsening of disease.

[0052] The term “survival outcome” as used herein may refer to survival, overall survival, or progression free survival.

[0053] The phrase “scale factor” as used herein refers to a factor that relates change in gene expression to prognosis. An example of a scale factor is a factor obtained by maximizing the partial likelihoods of the Cox proportional hazards model.

[0054] The gene expression signatures, signature values, survival predictor scores, stromal scores, survival estimate curves, and probabilities of survival disclosed herein may be stored in digitally encoded format on computer readable media, e.g., computer readable media used in conjunction with microarray or chip reading devices or computer readable

media used to store patient data during treatment for DLBCL. Such media and the specialized devices that use them, e.g., for diagnostic and clinical applications, are known in the art.

[0055] The invention provides a method for predicting a survival outcome in a subject diagnosed with DLBCL using gene expression data. Such data may be gathered using any effective method of quantifying gene expression. For example, gene expression data may be measured or estimated using one or more microarrays. The microarrays may be of any effective type, including, but not limited to, nucleic acid based or antibody based. Gene expression may also be measured by a variety of other techniques, including, but not limited to, PCR, quantitative RT-PCR, real-time PCR, RNA amplification, *in situ* hybridization, immunohistochemistry, immunocytochemistry, FACS, serial analysis of gene expression (SAGE) (Velculescu et al., *Science*, 270: 484–87 (1995)), Northern blot hybridization, or western blot hybridization.

[0056] Nucleic acid microarrays generally comprise nucleic acid probes derived from individual genes and placed in an ordered array on a support. This support may be, for example, a glass slide, a nylon membrane, or a silicon wafer. Gene expression patterns in a sample are obtained by hybridizing the microarray with the gene expression product from the sample. This gene expression product may be, for example, total cellular mRNA, rRNA, or cDNA obtained by reverse transcription of total cellular mRNA. The gene expression product from a sample is labeled with a radioactive, fluorescent, or other label to allow for detection. Following hybridization, the microarray is washed, and hybridization of the gene expression product to each nucleic acid probe on the microarray is detected and quantified using a detection device such as a phosphorimager or scanning confocal microscope.

[0057] There are two broad classes of microarrays: cDNA and oligonucleotide arrays. cDNA arrays consist of hundreds or thousands of cDNA probes immobilized on a solid support. These cDNA probes are usually 100 nucleotides or greater in size. There are two commonly used designs for cDNA arrays. The first is the nitrocellulose filter array, which is generally prepared by robotic spotting of purified DNA fragments or lysates of bacteria containing cDNA clones onto a nitrocellulose filter (Southern et al., *Genomics*, 13: 1008–17 (1992); Southern et al., *Nucl Acids Res* 22: 1368–73 (1994); Gress et al., *Oncogene*, 13: 1819–30 (1996); Pietu et al., *Genome Res.*, 6: 492–503 (1996)). The other commonly used cDNA arrays is fabricated by robotic spotting of PCR fragments from cDNA clones onto glass microscope slides (Schena et al., *Science*, 270: 467–70 (1995); DeRisi et al., *Nature*

Genet., 14: 457–60 (1996); Schena et al., *Proc. Nat'l. Acad. Sci. USA*, 93: 10614–19 (1996); Shalon et al., *Genome Res.*, 6: 639–45 (1996); DeRisi et al., *Science*, 278: 680–86 (1997); Heller et al., *Proc. Nat'l. Acad. Sci. USA*, 94: 2150–55 (1997); Lashkari et al., *Proc. Nat'l. Acad. Sci. USA*, 94: 13057–62 (1997)). These cDNA microarrays are simultaneously hybridized with two fluorescent cDNA probes, each labeled with a different fluorescent dye (typically Cy3 or Cy5). In this format, the relative mRNA expression in two samples is directly compared for each gene on the microarray. Oligonucleotide arrays differ from cDNA arrays in that the probes are 20- to 25-mer oligonucleotides. Oligonucleotide arrays are generally produced by *in situ* oligonucleotide synthesis in conjunction with photolithographic masking techniques (Pease et al., *Proc. Nat'l. Acad. Sci. USA*, 91: 5022–26 (1994); Lipshutz et al., *Biotechniques* 19: 442–47 (1995); Chee et al., *Science*, 274: 610–14 (1996); Lockhart et al., *Nature Biotechnol.*, 14: 1675–80 (1996); Wodicka et al., *Nature Biotechnol.*, 15: 1359–6714 (1997)). The solid support for oligonucleotide arrays is typically a glass or silicon surface.

[0058] Methods and techniques applicable to array synthesis and use have been described in, for example, U.S. Patent Nos. 5,143,854 (Pirrung), 5,242,974 (Holmes), 5,252,743 (Barrett), 5,324,633 (Fodor), 5,384,261 (Winkler), 5,424,186 (Fodor), 5,445,934 (Fodor), 5,451,683 (Barrett), 5,482,867 (Barrett), 5,491,074 (Aldwin), 5,527,681 (Holmes), 5,550,215 (Holmes), 5,571,639 (Hubbell), 5,578,832 (Trulson), 5,593,839 (Hubbell), 5,599,695 (Pease), 5,624,711 (Sundberg), 5,631,734 (Stern), 5,795,716 (Chee), 5,831,070 (Pease), 5,837,832 (Chee), 5,856,101 (Hubbell), 5,858,659 (Sapolsky), 5,936,324 (Montagu), 5,968,740 (Fodor), 5,974,164 (Chee), 5,981,185 (Matson), 5,981,956 (Stern), 6,025,601 (Trulson), 6,033,860 (Lockhart), 6,040,193 (Winkler), 6,090,555 (Fiekowsky), and 6,410,229 (Lockhart), and U.S. Patent Application Publication No. 2003/0104411 (Fodor).

[0059] Microarrays may generally be produced using a variety of techniques, such as mechanical or light directed synthesis methods that incorporate a combination of photolithographic methods and solid phase synthesis methods. Techniques for the synthesis of microarrays using mechanical synthesis methods are described in, for example, U.S. Patent Nos. 5,384,261 (Winkler) and 6,040,193 (Winkler). Although a planar array surface is preferred, the microarray may be fabricated on a surface of virtually any shape, or even on a multiplicity of surfaces. Microarrays may be nucleic acids on beads, gels, polymeric surfaces, fibers such as fiber optics, glass, or any other appropriate substrate. See, for

example, U.S. Patent Nos. 5,708,153 (Dower), 5,770,358 (Dower), 5,789,162 (Dower), 5,800,992 (Fodor), and 6,040,193 (Winkler).

[0060] Microarrays can be packaged in such a manner as to allow for diagnostic use, or they can be all-inclusive devices. See, for example, U.S. Patent Nos. 5,856,174 (Lipshutz) and 5,922,591 (Anderson).

[0061] Microarrays directed to a variety of purposes are commercially available from Affymetrix (Santa Clara, CA). For instance, these microarrays may be used for genotyping and gene expression monitoring.

[0062] Gene expression data can be used to identify genes that are coordinately regulated. Genes that encode components of the same multi-subunit protein complex are often coordinately regulated. Coordinate regulation is also observed among genes whose products function in a common differentiation program or in the same physiological response pathway. Recent application of gene expression profiling to the immune system has shown that lymphocyte differentiation and activation are accompanied by parallel changes in expression among hundreds of genes. Gene expression databases may be used to interpret the pathological changes in gene expression that accompany autoimmunity, immune deficiencies, cancers of immune cells and of normal immune responses.

[0063] Scanning and interpreting large bodies of relative gene expression data is a formidable task. This task is greatly facilitated by algorithms designed to organize the data in a way that highlights systematic features, and by visualization tools that represent the differential expression of each gene as varying intensities and hues of color (Eisen et al., *Proc. Nat'l. Acad. Sci. USA*, 95: 14863-68 (1998)). The development of microarrays, which are capable of generating massive amounts of expression data in a single experiment, has greatly increased the need for faster and more efficient methods of analyzing large-scale expression data sets. In order to effectively utilize microarray gene expression data for the prediction of survival in DLBCL patients, there is a need for new algorithms to be developed, which can identify important information and convert it to a more manageable format. In addition, the microarrays used to generate this data can be streamlined to incorporate probe sets that are useful for survival outcome prediction.

[0064] Mathematical analysis of gene expression data is a rapidly evolving science based on a rich mathematics of pattern recognition developed in other contexts (Kohonen, Self-Organizing Maps, Springer Press (Berlin 1997)). Mathematical analysis of gene expression data can be used, for example, to identify groups of genes that are coordinately regulated

within a biological system, to recognize and interpret similarities between biological samples on the basis of similarities in gene expression patterns, and/or to recognize and identify those features of a gene expression pattern that are related to distinct biological processes or phenotypes.

[0065] Mathematical analysis of gene expression data often begins by establishing the expression pattern for each gene on an array across a number (n) of experimental samples. The expression pattern of each gene can be represented by a point in n-dimensional space, with each coordinate specified by an expression measurement in one of the n samples (Eisen et al., *Proc. Nat'l. Acad. Sci. USA*, 95: 14863-68 (1998)). A clustering algorithm that uses distance metrics can then be applied to locate clusters of genes in this n-dimensional space. These clusters indicate genes with similar patterns of variation in expression over a series of experiments. Clustering methods that have been applied to microarray data in the past include hierarchical clustering (Eisen et al., *supra*), self-organizing maps (SOMs) (Tamayo et al., *Proc. Nat'l. Acad. Sci. USA*, 96: 2907-12 (1999)), k-means (Tavazoie et al., *Nature Genet.*, 22: 281-85 (1999)), and deterministic annealing (Alon et al., *Proc. Nat'l. Acad. Sci. USA*, 96: 6745-50 (1999)).

[0066] A variety of different algorithms, each emphasizing distinct orderly features of the data, may be required to glean the maximal biological insight from a set of samples (Alizadeh et al., *J. Clin. Immunol.*, 18: 373-79 (1998)). One such algorithm, hierarchical clustering, begins by determining the gene expression correlation coefficients for each pair of the n genes studied. Genes with similar gene expression correlation coefficients are grouped next to one another in a hierarchical fashion. Generally, genes with similar expression patterns under a particular set of conditions can encode protein products with related roles in the physiological adaptation to those conditions. Novel genes of unknown function that are clustered with a large group of functionally related genes likely participate in similar or related biological process. Likewise, other clustering methods mentioned herein can also group genes together that encode proteins with related biological function.

[0067] In such clustering methods, genes that are clustered together reflect a particular biological function, and are termed gene expression signatures (Shaffer et al., *Immunity* 15: 375-85 (2001)). One general type of gene expression signature includes genes that are characteristically expressed in a particular cell type or at a particular stage of cellular differentiation or activation. Another general type of gene expression signature includes

genes that are regulated in their expression by a particular biological process such as proliferation, or by the activity of a particular transcription factor or signaling pathway.

[0068] The pattern of gene expression in a biological sample can provide a distinctive and accessible molecular picture of its functional state and identity (DeRisi et al., *Science*, 278: 680–86 (1997); Cho et al., *Mol. Cell*, 2: 65–73 (1998); Chu et al., *Science*, 282: 699–705 (1998); Holstege et al., *Cell*, 95: 717–728 (1998); Spellman et al., *Mol. Biol. Cell*, 9: 3273–97 (1998)). Each cell transduces variations in its environment, internal state, and developmental state into readily measured and recognizable variations in its gene expression patterns. Two different samples with related gene expression patterns are therefore likely to be biologically and functionally similar to one another. Thus, a specific gene expression signature in a sample can provide important biological insights into its cellular composition and the function of various intracellular pathways within those cells.

[0069] Databases of gene expression signatures have proven useful in elucidating the complex gene expression patterns of various cancers. For example, the expression pattern of genes in the germinal center B cell signature in a lymphoma biopsy indicates that the lymphoma includes cells derived from the germinal center stage of differentiation. In the same lymphoma biopsy, the expression of genes from the T cell signature can be used to estimate the degree of infiltration of the tumor by host T cells, while the expression of genes from the proliferation signature can be used to quantitate the tumor cell proliferation rate. In this manner, gene expression signatures provide an "executive summary" of the biological properties of a tumor specimen. Gene expression signatures can also be helpful in interpreting the results of a supervised analysis of gene expression data. A supervised analysis generates a list of genes with expression patterns that correlate with survival. Gene expression signatures can be useful in assigning these "predictive" genes to functional categories. In building a multivariate model of survival based on gene expression data, this functional categorization helps to limit the inclusion of multiple genes in the model that measure the same aspect of tumor biology.

[0070] This following approach was utilized to create the survival prediction models for DLBCL of the invention. Gene expression profiles were used to create multivariate models for predicting survival. The methods for creating these models were "supervised" in that they used clinical data to guide the selection of genes to be used in the prognostic classification. The method identified genes with expression patterns that correlated with the length of

overall survival following chemotherapy. Generally the process for identifying the multivariate model for predicting survival included the following steps:

1. Genes were identified having expression patterns univariately associated with a particular clinical outcome using a Cox proportional hazards model. Generally, a univariate p-value of <0.01 is considered the cut-off for significance (however, another criterion can be used). These genes were termed “predictor” genes.
2. Within a set of predictor genes, gene expression signatures were identified.
3. For each gene expression signature significantly associated with survival, the average expression of each component genes within this signature was used to generate a gene expression signature value.
4. A multivariate Cox model of clinical outcome using the gene expression signature values was built.
5. Additional genes were added to the model, which added to the statistical power of the model.

[0071] The model of the invention generates a survival predictor score, with a higher score being associated with worse clinical outcome. The resulting model can be used separately to predict a survival outcome. Alternatively, the model can be used in conjunction with one or more other models, disclosed herein or in other references, to predict a survival outcome.

[0072] The present invention discloses several gene expression signatures related to the clinical outcome of DLBCL patients. The signatures were identified using the clinical data and methods described below in Examples 1 and 2. Three of these gene expression signatures are the germinal center B cell (GCB) signature, the stromal-1 signature, and the stromal-2 signature. Each component gene of these signatures is identified in Table 1 according to its GenBank accession number, its GeneID assigned by Entrez Gene, a common gene symbol, and a descriptive gene title. Table 1 also provides the Affymetrix Probe Set ID, which can be used (e.g., on the Affymetrix U133+ (Affymetrix, Santa Clara, CA) microarray) to determine the gene expression level for the indicated gene. The computer-readable sequence listing filed herewith includes a representative fragment sequence (of about 100 bp or greater) for each genomic target sequence listed in Table 1, followed by the sequence for each probe in the corresponding Affymetrix probe set listed in Table 1.

Table 1

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
GCB	NM_052932	114908	TMEM123	transmembrane protein 123	211967_at
GCB	NM_001014380	84056	KATNAL1	katanin p60 subunit A-like 1	227713_at
GCB	NM_004665	8875	VNN2	vanin 2	205922_at
GCB	NM_004760	9263	STK17A	serine/threonine kinase 17a (apoptosis-inducing)	202693_s_at
GCB	CR590554			Full-length cDNA clone CS0DF007YJ21 of Fetal brain of Homo sapiens (human)	228464_at
GCB	NM_017599	55591	VEZT	vezatin, adherens junctions transmembrane protein	223089_at
GCB	NM_018351	55785	FGD6	FYVE, RhoGEF and PH domain containing 6	1555136_at
GCB	NM_001007075	51088	KLHL5	kelch-like 5 (Drosophila)	226001_at
GCB	NM_004845	9468	PCYT1B	phosphate cytidyltransferase 1, choline, beta	228959_at
GCB	AK026881			CDNA: FLJ23228 fis, clone CAE06654	226799_at
GCB	NM_018440	55824	PAG1	phosphoprotein associated with glycosphingolipid microdomains 1	225626_at
GCB	NM_004965	3150	HMG1	high-mobility group nucleosome binding domain 1	200944_s_at
GCB	NM_001706	604	BCL6	B cell CLL/lymphoma 6 (zinc finger protein 51)	228758_at
GCB	NM_020747	57507	ZNF608	zinc finger protein 608	229817_at
GCB	NM_001001695	400941	FLJ42418	FLJ42418 protein	231455_at
GCB	NM_015055	23075	SWAP70	SWAP-70 protein	209306_s_at
GCB	NM_005607	5747	PTK2	PTK2 protein tyrosine kinase 2	208820_at
GCB	XM_027236	23508	TTC9	tetratricopeptide repeat domain 9	213172_at
GCB	BQ213652	440864	LOC440864	hypothetical gene supported by BC040724	1569034_a_at
GCB	NM_005574	4005	LMO2	LIM domain only 2 (rhombotin-like 1)	204249_s_at
GCB	NM_014667	9686	VGLL4	vestigial like 4 (Drosophila)	212399_s_at
GCB	NM_002221	3707	ITPKB	inositol 1,4,5-trisphosphate 3-kinase B	203723_at
GCB	NM_000902	4311	MME	membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase)	203434_s_at
GCB	NM_012446	23635	SSBP2	single-stranded DNA binding protein 2	203787_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
GCB	NM_024613	79666	PLEKHF2	pleckstrin homology domain containing, family F (with FYVE domain) member 2	222699_s_at
GCB	AV705976			Transcribed locus	204681_s_at
GCB	NM_012108	26228	BRDG1	BCR downstream signaling 1	220059_at
GCB	NM_014397	10783	NEK6	NIMA (never in mitosis gene a)-related kinase 6	223158_s_at
GCB	NM_018981	54431	DNAJC10	DnaJ (Hsp40) homolog, subfamily C, member 10	225174_at
GCB	NM_001379	1786	DNMT1	DNA (cytosine-5)-methyltransferase 1	227684_at
GCB	NM_006152	4033	LRMP	lymphoid-restricted membrane protein	35974_at
GCB	NM_024701	79754	ASB13	ankyrin repeat and SOCS box-containing 13	218862_at
GCB	NM_006085	10380	BPNT1	3'(2'), 5'-bisphosphate nucleotidase 1	232103_at
GCB	NM_023009	65108	MARCKSL1	MARCKS-like 1	200644_at
GCB	NM_033121	88455	ANKRD13A	ankyrin repeat domain 13A	224810_s_at
GCB	NM_015187	23231	KIAA0746	KIAA0746 protein	235353_at
GCB	NM_175739	327657	SERPINA9	serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 9	1553499_s_at
GCB	NM_001012391	400509	RUNDC2B	RUN domain containing 2B	1554413_s_at
GCB	XM_034274	4603	MYBL1	v-myb myeloblastosis viral oncogene homolog (avian)-like 1	213906_at
Stromal-1	NM_024579	79630	C1orf54	chromosome 1 open reading frame 54	219506_at
Stromal-1	NM_001645	341	APOC1	apolipoprotein C-I	213553_x_at
Stromal-1	NM_001562	3606	IL18	interleukin 18 (interferon-gamma-inducing factor)	206295_at
Stromal-1	NM_014479	27299	ADAMDEC1	ADAM-like, decysin 1	206134_at
Stromal-1	NM_003465	1118	CHIT1	chitinase 1 (chitotriosidase)	208168_s_at
Stromal-1	NM_000954	5730	PTGDS	prostaglandin D2 synthase 21kDa (brain)	211748_x_at
Stromal-1	NM_001056	6819	SULT1C1	sulfotransferase family, cytosolic, 1C, member 1	211470_s_at
Stromal-1	NM_018000	55686	MREG	melanoregulin	219648_at
Stromal-1	NM_001018058	22797	TFEC	transcription factor EC	206715_at
Stromal-1	NM_000239	4069	LYZ	lysozyme (renal amyloidosis)	213975_s_at
Stromal-1	NM_006834	10981	RAB32	RAB32, member RAS oncogene family	204214_s_at
Stromal-1	NM_000416	3459	IFNGR1	interferon gamma receptor 1	202727_s_at
Stromal-1	NM_004666	8876	VNN1	vanin 1	205844_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_031491	83758	RBP5	retinol binding protein 5, cellular	223820_at
Stromal-1	NM_001276	1116	CHI3L1	chitinase 3-like 1 (cartilage glycoprotein-39)	209396_s_at
Stromal-1	NM_138434	113763	C7orf29	chromosome 7 open reading frame 29	227598_at
Stromal-1	NM_001005340	10457	GPNUMB	glycoprotein (transmembrane) numb	201141_at
Stromal-1	NM_002294	3920	LAMP2	lysosomal-associated membrane protein 2	203041_s_at
Stromal-1	NM_002888	5918	RARRES1	retinoic acid receptor responder (tazarotene induced) 1	221872_at
Stromal-1	NM_172248	1438	CSF2RA	colony stimulating factor 2 receptor, alpha, low- affinity (granulocyte- macrophage)	210340_s_at
Stromal-1	NM_018344	55315	SLC29A3	solute carrier family 29 (nucleoside transporters), member 3	219344_at
Stromal-1	NM_032413	84419	C15orf48	chromosome 15 open reading frame 48	223484_at
Stromal-1	NM_001001851	80760	ITIH5	inter-alpha (globulin) inhibitor H5	1553243_at
Stromal-1	NM_000211	3689	ITGB2	integrin, beta 2 (complement component 3 receptor 3 and 4 subunit)	1555349_a_at
Stromal-1	NM_005213	1475	CSTA	cystatin A (stefin A)	204971_at
Stromal-1	NM_003874	8832	CD84	CD84 molecule	205988_at
Stromal-1	NM_000228	3914	LAMB3	laminin, beta 3	209270_at
Stromal-1	NM_005651	6999	TDO2	tryptophan 2,3- dioxygenase	205943_at
Stromal-1	NM_001005266	283651	C15orf21	chromosome 15 open reading frame 21	242649_x_at
Stromal-1	AV659177			Transcribed locus	230391_at
Stromal-1	NM_001747	822	CAPG	capping protein (actin filament), gelsolin-like	201850_at
Stromal-1	NM_000784	1593	CYP27A1	cytochrome P450, family 27, subfamily A, polypeptide 1	203979_at
Stromal-1	NM_052998	113451	ADC	arginine decarboxylase	228000_at
Stromal-1	NM_016240	51435	SCARA3	scavenger receptor class A, member 3	219416_at
Stromal-1	Z74615		COL1A1	Collagen, type I, alpha 1	217430_x_at
Stromal-1	NM_052947	115701	ALPK2	alpha-kinase 2	228367_at
Stromal-1	NM_021136	6252	RTN1	reticulum 1	210222_s_at
Stromal-1	AL049370			Full-length cDNA clone CL0BB018ZE07 of Neuroblastoma of Homo sapiens (human)	213100_at
Stromal-1	NM_006042	9955	HS3ST3A1	heparan sulfate (glucosamine) 3-O- sulfotransferase 3A1	219985_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_000041	348	APOE	apolipoprotein E	203382_s_at
Stromal-1	NM_004994	4318	MMP9	matrix metalloproteinase 9 (gelatinase B, 92kDa gelatinase, 92kDa type IV collagenase)	203936_s_at
Stromal-1	NM_001831	1191	CLU	clusterin	222043_at
Stromal-1	NM_002305	3956	LGALS1	lectin, galactoside- binding, soluble, 1 (galectin 1)	201105_at
Stromal-1	NM_032024	83938	C10orf11	chromosome 10 open reading frame 11	223703_at
Stromal-1	NM_001025201	1123	CHN1	chimerin (chimaerin) 1	212624_s_at
Stromal-1	NM_003489	8204	NRIP1	nuclear receptor interacting protein 1	202599_s_at
Stromal-1	NM_032646	94015	TTYH2	tweety homolog 2 (Drosophila)	223741_s_at
Stromal-1	NM_001312	1397	CRIP2	cysteine-rich protein 2	208978_at
Stromal-1	NM_023075	65258	MPPE1	metallophosphoesterase 1	213924_at
Stromal-1	NM_004364	1050	CEBPA	CCAAT/enhancer binding protein (C/EBP), alpha	204039_at
Stromal-1	NM_000248	4286	MITF	microphthalmia- associated transcription factor	207233_s_at
Stromal-1	NM_002185	3575	IL7R	interleukin 7 receptor	226218_at
Stromal-1	NM_021638	60312	AFAP	actin filament associated protein	203563_at
Stromal-1	NM_003786	8714	ABCC3	ATP-binding cassette, sub-family C (CFTR/MRP), member 3	208161_s_at
Stromal-1		730351	LOC730351	hypothetical protein LOC730351	229407_at
Stromal-1	NM_012153	26298	EHF	ets homologous factor	225645_at
Stromal-1	NM_004887	9547	CXCL14	chemokine (C-X-C motif) ligand 14	222484_s_at
Stromal-1	NM_002030	2359	FPRL2	formyl peptide receptor- like 2	230422_at
Stromal-1	NM_001321	1466	CSRP2	cysteine and glycine-rich protein 2	207030_s_at
Stromal-1	NM_001945	1839	HBEGF	heparin-binding EGF-like growth factor	203821_at
Stromal-1	NM_031412	23710	GABARAPL1	GABA(A) receptor- associated protein like 1	208869_s_at
Stromal-1	NM_006022	8848	TSC22D1	TSC22 domain family, member 1	215111_s_at
Stromal-1	NM_016174	51148	CEECAM1	cerebral endothelial cell adhesion molecule 1	224794_s_at
Stromal-1	NM_015103	23129	PLXND1	plexin D1	212235_at
Stromal-1	NM_003270	7105	TSPAN6	tetraspanin 6	209109_s_at
Stromal-1	NM_000887	3687	ITGAX	integrin, alpha X (complement component 3 receptor 4 subunit)	210184_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_001864	1346	COX7A1	cytochrome c oxidase subunit VIIa polypeptide 1 (muscle)	204570_at
Stromal-1	CR599008		GPR157	Full-length cDNA clone CS0DJ007YL22 of T cells (Jurkat cell line) Cot 10-normalized of Homo sapiens (human)	227970_at
Stromal-1	NM_198580	376497	SLC27A1	solute carrier family 27 (fatty acid transporter), member 1	226728_at
Stromal-1	NM_025106	80176	SPSB1	splA/ryanodine receptor domain and SOCS box containing 1	226075_at
Stromal-1	NM_020130	56892	C8orf4	chromosome 8 open reading frame 4	218541_s_at
Stromal-1	NM_173833	286133	SCARA5	scavenger receptor class A, member 5 (putative)	229839_at
Stromal-1	NM_007223	11245	GPR176	G protein-coupled receptor 176	227846_at
Stromal-1	NM_013437	29967	LRP12	low density lipoprotein-related protein 12	219631_at
Stromal-1	NM_007332	8989	TRPA1	transient receptor potential cation channel, subfamily A, member 1	228438_at
Stromal-1	NM_152744	221935	SDK1	sidekick homolog 1 (chicken)	229912_at
Stromal-1	NM_001409	1953	MEGF6	multiple EGF-like-domains 6	226869_at
Stromal-1	NM_012082	23414	ZFPM2	zinc finger protein, multitype 2	219778_at
Stromal-1	NM_080430	140606	SELM	selenoprotein M	226051_at
Stromal-1	NM_030971	81855	SFXN3	sideroflexin 3	217226_s_at
Stromal-1	NM_003246	7057	THBS1	thrombospondin 1	201109_s_at
Stromal-1	NM_003882	8840	WISP1	WNT1 inducible signaling pathway protein 1	235821_at
Stromal-1	NM_005202	1296	COL8A2	collagen, type VIII, alpha 2	221900_at
Stromal-1	NM_003711	8611	PPAP2A	phosphatidic acid phosphatase type 2A	210946_at
Stromal-1	NM_004995	4323	MMP14	matrix metalloproteinase 14 (membrane-inserted)	202828_s_at
Stromal-1	NM_001005336	1759	DNM1	dynammin 1	215116_s_at
Stromal-1	NM_153717	2121	EVC	Ellis van Creveld syndrome	219432_at
Stromal-1	NM_173462	89932	PAPLN	papilin, proteoglycan-like sulfated glycoprotein	226435_at
Stromal-1	XM_496707	441027	FLJ12993	hypothetical LOC441027	229623_at
Stromal-1	NM_001839	1266	CNN3	calponin 3, acidic	228297_at
Stromal-1	NM_015429	25890	ABI3BP	ABI gene family, member 3 (NESH) binding protein	223395_at
Stromal-1	NM_002840	5792	PTPRF	protein tyrosine phosphatase, receptor type, F	200636_s_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_001001522	6876	TAGLN	transgelin	1555724_s_at
Stromal-1	NM_017637	54796	BNC2	basonuclein 2	229942_at
Stromal-1	NM_003391	7472	WNT2	wingless-type MMTV integration site family member 2	205648_at
Stromal-1	NM_015461	25925	ZNF521	zinc finger protein 521	226677_at
Stromal-1	NM_006475	10631	POSTN	periostin, osteoblast specific factor	210809_s_at
Stromal-1	NM_005418	6764	ST5	suppression of tumorigenicity 5	202440_s_at
Stromal-1	NM_005203	1305	COL13A1	collagen, type XIII, alpha 1	211343_s_at
Stromal-1	NM_000681	150	ADRA2A	adrenergic, alpha-2A-, receptor	209869_at
Stromal-1	NM_006622	10769	PLK2	polo-like kinase 2 (Drosophila)	201939_at
Stromal-1	AL528626			Full-length cDNA clone CS0DD001YA12 of Neuroblastoma Cot 50-normalized of Homo sapiens (human)	228573_at
Stromal-1	AF180519	23766	GABARAPL3	GABA(A) receptors associated protein like 3	211458_s_at
Stromal-1	NM_024723	79778	MICAL2	MICAL-like 2	219332_at
Stromal-1	NM_057177	117583	PARD3B	par-3 partitioning defective 3 homolog B (C. elegans)	228411_at
Stromal-1	NM_004949	1824	DSC2	desmocollin 2	226817_at
Stromal-1	NM_032784	84870	RSPO3	R-spondin 3 homolog (Xenopus laevis)	228186_s_at
Stromal-1	NM_007039	11099	PTPN21	protein tyrosine phosphatase, non-receptor type 21	226380_at
Stromal-1	NM_031935	83872	HMCN1	hemicentin 1	235944_at
Stromal-1	AK022877			Clone TUA8 Cri-du-chat region mRNA	213169_at
Stromal-1	AK127644			CDNA FLJ45742 fis, clone KIDNE2016327	236297_at
Stromal-1	AK056963			Full length insert cDNA clone ZE03F06	226282_at
Stromal-1	NM_000899	4254	KITLG	KIT ligand	226534_at
Stromal-1	NM_002387	4163	MCC	mutated in colorectal cancers	226225_at
Stromal-1	NM_198270	4810	NHS	Nance-Horan syndrome (congenital cataracts and dental anomalies)	228933_at
Stromal-1	NM_183376	91947	ARRDC4	arrestin domain containing 4	225283_at
Stromal-1	NM_000216	3730	KAL1	Kallmann syndrome 1 sequence	205206_at
Stromal-1	NM_001008224	55075	UACA	uveal autoantigen with coiled-coil domains and ankyrin repeats	223279_s_at
Stromal-1	NM_133493	135228	CD109	CD109 molecule	226545_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_005545	3671	ISLR	immunoglobulin superfamily containing leucine-rich repeat	207191_s_at
Stromal-1	NM_014365	26353	HSPB8	heat shock 22kDa protein 8	221667_s_at
Stromal-1	NM_014476	27295	PDLIM3	PDZ and LIM domain 3	209621_s_at
Stromal-1	NM_020962	57722	NOPE	likely ortholog of mouse neighbor of Punc E11	227870_at
Stromal-1	NM_018357	55323	LARP6	La ribonucleoprotein domain family, member 6	218651_s_at
Stromal-1	NM_012323	23764	MAFF	v-maf musculoaponeurotic fibrosarcoma oncogene homolog F (avian)	36711_at
Stromal-1	NM_003713	8613	PPAP2B	phosphatidic acid phosphatase type 2B	212230_at
Stromal-1	NM_023016	65124	ANKRD57	ankyrin repeat domain 57	227034_at
Stromal-1	NM_032777	25960	GPR124	G protein-coupled receptor 124	65718_at
Stromal-1	NM_001554	3491	CYR61	cysteine-rich, angiogenic inducer, 61	201289_at
Stromal-1	NM_145117	89797	NAV2	neuron navigator 2	218330_s_at
Stromal-1	NM_001002292	79971	GPR177	G protein-coupled receptor 177	228950_s_at
Stromal-1	NM_001401	1902	EDG2	endothelial differentiation, lysophosphatidic acid G- protein-coupled receptor, 2	204036_at
Stromal-1	NM_198282	340061	TMEM173	transmembrane protein 173	224929_at
Stromal-1	NM_014934	22873	DZIP1	DAZ interacting protein 1	204556_s_at
Stromal-1	NM_001901	1490	CTGF	connective tissue growth factor	209101_at
Stromal-1	NM_024600	79652	C16orf30	chromosome 16 open reading frame 30	219315_s_at
Stromal-1	NM_138370	91461	LOC91461	hypothetical protein BC007901	225380_at
Stromal-1	NM_014632	9645	MICAL2	microtubule associated monooxygenase, calponin and LIM domain containing 2	212472_at
Stromal-1	NM_032866	84952	CGNL1	cingulin-like 1	225817_at
Stromal-1	NM_003687	8572	PDLIM4	PDZ and LIM domain 4	211564_s_at
Stromal-1	BM544548			Transcribed locus	236179_at
Stromal-1	NM_001856	1307	COL16A1	collagen, type XVI, alpha 1	204345_at
Stromal-1	XM_087386	57493	HEG1	HEG homolog 1 (zebrafish)	213069_at
Stromal-1	NM_003887	8853	DDEF2	development and differentiation enhancing factor 2	206414_s_at
Stromal-1	NM_002844	5796	PTPRK	protein tyrosine phosphatase, receptor type, K	203038_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_022138	64094	SMOC2	SPARC related modular calcium binding 2	223235_s_at
Stromal-1	NM_001006624	10630	PDPN	podoplanin	204879_at
Stromal-1	NM_003174	6840	SVIL	supervillin	202565_s_at
Stromal-1	NM_002845	5797	PTPRM	protein tyrosine phosphatase, receptor type, M	155579_s_at
Stromal-1	NM_002889	5919	RARRES2	retinoic acid receptor responder (tazarotene induced) 2	209496_at
Stromal-1	NM_006094	10395	DLC1	deleted in liver cancer 1	210762_s_at
Stromal-1	NM_022463	64359	NXN	nucleoredoxin	219489_s_at
Stromal-1	AK027294			CDNA FLJ14388 fis, clone HEMBA1002716	229802_at
Stromal-1	NM_005711	10085	EDIL3	EGF-like repeats and discoidin I-like domains 3	225275_at
Stromal-1	NM_000177	2934	GSN	gelsolin (amyloidosis, Finnish type)	200696_s_at
Stromal-1	NM_016639	51330	TNFRSF12A	tumor necrosis factor receptor superfamily, member 12A	218368_s_at
Stromal-1	NM_004460	2191	FAP	fibroblast activation protein, alpha	209955_s_at
Stromal-1	NM_000064	718	C3	complement component 3	217767_at
Stromal-1	NM_016206	389136	VGLL3	vestigial like 3 (Drosophila)	227399_at
Stromal-1	NM_004339	754	PTTG1IP	pituitary tumor- transforming 1 interacting protein	200677_at
Stromal-1	NM_003255	7077	TIMP2	TIMP metalloproteinase inhibitor 2	224560_at
Stromal-1	NM_002998	6383	SDC2	syndecan 2 (heparan sulfate proteoglycan 1, cell surface-associated, fibroglycan)	212158_at
Stromal-1	NM_012223	4430	MYO1B	myosin IB	212364_at
Stromal-1	NM_020650	57333	RCN3	reticulocalbin 3, EF-hand calcium binding domain	61734_at
Stromal-1	AL573464			Transcribed locus	229554_at
Stromal-1	AK001903			CDNA FLJ11041 fis, clone PLACE1004405	227140_at
Stromal-1	NM_005928	4240	MFGE8	milk fat globule-EGF factor 8 protein	210605_s_at
Stromal-1	NM_000943	5480	PPIC	peptidylprolyl isomerase C (cyclophilin C)	204518_s_at
Stromal-1	NM_001008397	493869	LOC493869	similar to RIKEN cDNA 2310016C16	227628_at
Stromal-1	AK025431	768211	RELL1	receptor expressed in lymphoid tissues like 1	226430_at
Stromal-1	NM_000297	5311	PKD2	polycystic kidney disease 2 (autosomal dominant)	203688_at
Stromal-1	NM_002975	6320	CLEC11A	C-type lectin domain family 11, member A	211709_s_at
Stromal-1	NM_001920	1634	DCN	decorin	211813_x_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_001723	667	DST	dystonin	215016_x_at
Stromal-1	CR749529			MRNA; cDNA DKFZp686l18116 (from clone DKFZp686l18116)	227554_at
Stromal-1	NM_000165	2697	GJA1	gap junction protein, alpha 1, 43kDa (connexin 43)	201667_at
Stromal-1	NM_012104	23621	BACE1	beta-site APP-cleaving enzyme 1	217904_s_at
Stromal-1	NM_001957	1909	EDNRA	endothelin receptor type A	204464_s_at
Stromal-1	NM_138455	115908	CTHRC1	collagen triple helix repeat containing 1	225681_at
Stromal-1	NM_001331	1500	CTNND1	catenin (cadherin-associated protein), delta 1	208407_s_at
Stromal-1	NM_001613	59	ACTA2	actin, alpha 2, smooth muscle, aorta	200974_at
Stromal-1	NM_002192	3624	INHBA	inhibin, beta A (activin A, activin AB alpha polypeptide)	210511_s_at
Stromal-1	NM_000935	5352	PLOD2	procollagen-lysine, 2-oxoglutarate 5-dioxygenase 2	202620_s_at
Stromal-1	NM_015170	23213	SULF1	sulfatase 1	212354_at
Stromal-1	NM_006039	9902	MRC2	mannose receptor, C type 2	37408_at
Stromal-1	NM_005261	2669	GEM	GTP binding protein overexpressed in skeletal muscle	204472_at
Stromal-1	NM_001008707	2009	EML1	echinoderm microtubule associated protein like 1	204797_s_at
Stromal-1	NM_001031679	253827	MSRB3	methionine sulfoxide reductase B3	225782_at
Stromal-1	NM_001004125	286319	TUSC1	tumor suppressor candidate 1	227388_at
Stromal-1	NM_005965	4638	MYLK	myosin, light chain kinase	202555_s_at
Stromal-1	NM_016205	56034	PDGFC	platelet derived growth factor C	218718_at
Stromal-1	NM_015976	51375	SNX7	sorting nexin 7	205573_s_at
Stromal-1	NM_130830	131578	LRRC15	leucine rich repeat containing 15	213909_at
Stromal-1	NM_002026	2335	FN1	fibronectin 1	212464_s_at
Stromal-1	NM_006855	11015	KDEL3	KDEL (Lys-Asp-Glu-Leu) endoplasmic reticulum protein retention receptor 3	204017_at
Stromal-1	NM_002292	3913	LAMB2	laminin, beta 2 (laminin S)	216264_s_at
Stromal-1	NM_002658	5328	PLAU	plasminogen activator, urokinase	205479_s_at
Stromal-1	NM_005529	3339	HSPG2	heparan sulfate proteoglycan 2 (perlecan)	201655_s_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_001235	871	SERPINH1	serpin peptidase inhibitor, clade H (heat shock protein 47), member 1, (collagen binding protein 1)	207714_s_at
Stromal-1	AJ318805			CDNA FLJ44429 fis, clone UTERU2015653	227061_at
Stromal-1	NM_000396	1513	CTSK	cathepsin K	202450_s_at
Stromal-1	NM_031302	83468	GLT8D2	glycosyltransferase 8 domain containing 2	227070_at
Stromal-1	NM_080821	116151	C20orf108	chromosome 20 open reading frame 108	224690_at
Stromal-1	NM_002345	4060	LUM	lumican	201744_s_at
Stromal-1	NM_005110	9945	GFPT2	glutamine-fructose-6-phosphate transaminase 2	205100_at
Stromal-1	NM_002941	6091	ROBO1	roundabout, axon guidance receptor, homolog 1 (Drosophila)	213194_at
Stromal-1	NM_005429	7424	VEGFC	vascular endothelial growth factor C	209946_at
Stromal-1	NM_002213	3693	ITGB5	integrin, beta 5	201125_s_at
Stromal-1	XM_051017	23363	OBSL1	obscurin-like 1	212775_at
Stromal-1	NM_181724	338773	TMEM119	transmembrane protein 119	227300_at
Stromal-1	NM_003474	8038	ADAM12	ADAM metallopeptidase domain 12 (meltrin alpha)	213790_at
Stromal-1	NM_018222	55742	PARVA	parvin, alpha	217890_s_at
Stromal-1	NM_006478	10634	GAS2L1	growth arrest-specific 2 like 1	31874_at
Stromal-1	NM_000093	1289	COL5A1	collagen, type V, alpha 1	212489_at
Stromal-1	NM_006288	7070	THY1	Thy-1 cell surface antigen	208851_s_at
Stromal-1	CD357685		TIMP2	Transcribed locus, strongly similar to XP_511714.1 similar to Metalloproteinase inhibitor 2 precursor (TIMP-2) (Tissue inhibitor of metalloproteinases-2) (CSC-21K) [Pan troglodytes]	231579_s_at
Stromal-1	NM_003247	7058	THBS2	thrombospondin 2	203083_at
Stromal-1	NM_000088	1277	COL1A1	collagen, type I, alpha 1	1556499_s_at
Stromal-1	NM_006832	10979	PLEKHC1	pleckstrin homology domain containing, family C (with FERM domain) member 1	209210_s_at
Stromal-1	NM_021961	7003	TEAD1	TEA domain family member 1 (SV40 transcriptional enhancer factor)	224955_at
Stromal-1	AK128814			CDNA FLJ25106 fis, clone CBR01467	213675_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_153367	219654	C10orf56	chromosome 10 open reading frame 56	212423_at
Stromal-1	AK092048			MRNA; cDNA DKFZp313C0240 (from clone DKFZp313C0240)	227623_at
Stromal-1	NM_005245	2195	FAT	FAT tumor suppressor homolog 1 (Drosophila)	201579_at
Stromal-1	NM_001129	165	AEBP1	AE binding protein 1	201792_at
Stromal-1	NM_002403	4237	MFAP2	microfibrillar-associated protein 2	203417_at
Stromal-1	NM_004342	800	CALD1	caldesmon 1	201616_s_at
Stromal-1	NM_005576	4016	LOXL1	lysyl oxidase-like 1	203570_at
Stromal-1	NM_199511	151887	CCDC80	coiled-coil domain containing 80	225242_s_at
Stromal-1	NM_012098	23452	ANGPTL2	angiopoietin-like 2	213001_at
Stromal-1	NM_002210	3685	ITGAV	integrin, alpha V (vitronectin receptor, alpha polypeptide, antigen CD51)	202351_at
Stromal-1	NM_000366	7168	TPM1	tropomyosin 1 (alpha)	210986_s_at
Stromal-1	NM_198474	283298	OLFML1	olfactomedin-like 1	217525_at
Stromal-1	NM_001424	2013	EMP2	epithelial membrane protein 2	225078_at
Stromal-1	NM_032575	84662	GLIS2	GLIS family zinc finger 2	223378_at
Stromal-1	NM_007173	11098	PRSS23	protease, serine, 23	226279_at
Stromal-1	NM_001015880	9060	PAPSS2	3'-phosphoadenosine 5'- phosphosulfate synthase 2	203060_s_at
Stromal-1	NM_015645	114902	C1QTNF5	C1q and tumor necrosis factor related protein 5	223499_at
Stromal-1	AK130049			CDNA FLJ26539 fis, clone KDN09310	213429_at
Stromal-1	NM_001849	1292	COL6A2	collagen, type VI, alpha 2	209156_s_at
Stromal-1	NM_001014796	4921	DDR2	discoidin domain receptor family, member 2	225442_at
Stromal-1	NM_015463	25927	C2orf32	chromosome 2 open reading frame 32	226751_at
Stromal-1	AK055628		ADAM12	CDNA FLJ31066 fis, clone HSYRA2001153	226777_at
Stromal-1	NM_014799	9843	HEPH	hephaestin	203903_s_at
Stromal-1	NM_004385	1462	CSPG2	chondroitin sulfate proteoglycan 2 (versican)	221731_x_at
Stromal-1	NM_152330	122786	FRMD6	FERM domain containing 6	225481_at
Stromal-1	BQ917964		PPP4R2	Transcribed locus	235733_at
Stromal-1	NM_002615	5176	SERPINF1	serpin peptidase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1	202283_at
Stromal-1	NM_032348	54587	MXRA8	matrix-remodelling associated 8	213422_s_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_006106	10413	YAP1	Yes-associated protein 1, 65kDa	224894_at
Stromal-1	NM_020182	56937	TMEPAI	transmembrane, prostate androgen induced RNA	222449_at
Stromal-1	CB999028			Transcribed locus	226834_at
Stromal-1	NM_001711	633	BGN	biglycan	201261_x_at
Stromal-1	NM_006902	5396	PRRX1	paired related homeobox 1	226695_at
Stromal-1	NM_000428	4053	LTBP2	latent transforming growth factor beta binding protein 2	204682_at
Stromal-1	NM_004369	1293	COL6A3	collagen, type VI, alpha 3	201438_at
Stromal-1	NM_000393	1290	COL5A2	collagen, type V, alpha 2	221730_at
Stromal-1	NM_015419	25878	MXRA5	matrix-remodelling associated 5	209596_at
Stromal-1	NM_001102	87	ACTN1	actinin, alpha 1	208637_x_at
Stromal-1	NM_000877	3554	IL1R1	interleukin 1 receptor, type I	202948_at
Stromal-1	NM_015927	7041	TGFB1I1	transforming growth factor beta 1 induced transcript 1	209651_at
Stromal-1	NM_032772	84858	ZNF503	zinc finger protein 503	227195_at
Stromal-1	NM_020440	5738	PTGFRN	prostaglandin F2 receptor negative regulator	224937_at
Stromal-1	NM_000138	2200	FBN1	fibrillin 1	202765_s_at
Stromal-1	NM_031442	83604	TMEM47	transmembrane protein 47	209656_s_at
Stromal-1	NM_001734	716	C1S	complement component 1, s subcomponent	208747_s_at
Stromal-1	NM_002290	3910	LAMA4	laminin, alpha 4	202202_s_at
Stromal-1	CN312045		PPP4R2	Transcribed locus, weakly similar to NP_001013658.1 protein LOC387873 [Homo sapiens]	222288_at
Stromal-1	NM_000089	1278	COL1A2	collagen, type I, alpha 2	202403_s_at
Stromal-1	NM_004530	4313	MMP2	matrix metalloproteinase 2 (gelatinase A, 72kDa gelatinase, 72kDa type IV collagenase)	201069_at
Stromal-1	NM_001387	1809	DPYSL3	dihydropyrimidinase-like 3	201431_s_at
Stromal-1	NM_138389	92689	FAM114A1	family with sequence similarity 114, member A1	213455_at
Stromal-1	NM_006670	7162	TPBG	trophoblast glycoprotein	203476_at
Stromal-1	NM_000304	5376	PMP22	peripheral myelin protein 22	210139_s_at
Stromal-1	NM_002775	5654	HTRA1	HtrA serine peptidase 1	201185_at
Stromal-1	NM_002593	5118	PCOLCE	procollagen C- endopeptidase enhancer	202465_at
Stromal-1	NM_003118	6678	SPARC	secreted protein, acidic, cysteine-rich (osteonectin)	212667_at
Stromal-1	NM_007085	11167	FSTL1	folliculin-like 1	208782_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-1	NM_001080393	727936		predicted glycosyl- transferase 8 domain containing 4	235371_at
Stromal-1	NM_018153	84168	ANTXR1	anthrax toxin receptor 1	224694_at
Stromal-1	NM_001733	715	C1R	complement component 1, r subcomponent	212067_s_at
Stromal-1	NM_001797	1009	CDH11	cadherin 11, type 2, OB- cadherin (osteoblast)	207173_x_at
Stromal-1	NM_016938	30008	EFEMP2	EGF-containing fibulin- like extracellular matrix protein 2	209356_x_at
Stromal-2	NM_014601	30846	EHD2	EH-domain containing 2	45297_at
Stromal-2	NM_017789	54910	SEMA4C	sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4C	46665_at
Stromal-2	NM_000484	351	APP	amyloid beta (A4) precursor protein (peptidase nexin-II, Alzheimer disease)	200602_at
Stromal-2	NM_004684	8404	SPARCL1	SPARC-like 1 (mast9, hevin)	200795_at
Stromal-2	NM_002291	3912	LAMB1	laminin, beta 1	201505_at
Stromal-2	NM_000210	3655	ITGA6	integrin, alpha 6	201656_at
Stromal-2	NM_000552	7450	VWF	von Willebrand factor	202112_at
Stromal-2	NM_001233	858	CAV2	caveolin 2	203323_at
Stromal-2	NM_006404	10544	PROCR	protein C receptor, endothelial (EPCR)	203650_at
Stromal-2	NM_000609	6387	CXCL12	chemokine (C-X-C motif) ligand 12 (stromal cell- derived factor 1)	203666_at
Stromal-2	NM_002253	3791	KDR	kinase insert domain receptor (a type III receptor tyrosine kinase)	203934_at
Stromal-2	NM_001442	2167	FABP4	fatty acid binding protein 4, adipocyte	203980_at
Stromal-2	NM_016315	51454	GULP1	GULP, engulfment adaptor PTB domain containing 1	204237_at
Stromal-2	NM_006307	8406	SRPX	sushi-repeat-containing protein, X-linked	204955_at
Stromal-2	NM_000163	2690	GHR	growth hormone receptor	205498_at
Stromal-2	NM_000950	5638	PRRG1	proline rich Gla (G- carboxyglutamic acid) 1	205618_at
Stromal-2	NM_002666	5346	PLIN	perilipin	205913_at
Stromal-2	NM_000459	7010	TEK	TEK tyrosine kinase, endothelial (venous malformations, multiple cutaneous and mucosal)	206702_at
Stromal-2	NM_004797	9370	ADIPOQ	adiponectin, C1Q and collagen domain containing	207175_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-2	NM_000442	5175	PECAM1	platelet/endothelial cell adhesion molecule (CD31 antigen)	208981_at
Stromal-2	NM_198098	358	AQP1	aquaporin 1 (Colton blood group)	209047_at
Stromal-2	NM_021005	7026	NR2F2	nuclear receptor subfamily 2, group F, member 2	209120_at
Stromal-2	NM_014220	4071	TM4SF1	transmembrane 4 L six family member 1	209386_at
Stromal-2	NM_001001549	2887	GRB10	growth factor receptor-bound protein 10	209409_at
Stromal-2	NM_006108	10418	SPON1	spondin 1, extracellular matrix protein	209436_at
Stromal-2	NM_001003679	3953	LEPR	leptin receptor	209894_at
Stromal-2	NM_000599	3488	IGFBP5	insulin-like growth factor binding protein 5	211959_at
Stromal-2	NM_001753	857	CAV1	caveolin 1, caveolae protein, 22kDa	212097_at
Stromal-2	NM_005841	10252	SPRY1	sprouty homolog 1, antagonist of FGF signaling (Drosophila)	212558_at
Stromal-2	NM_015345	23500	DAAM2	dishevelled associated activator of morphogenesis 2	212793_at
Stromal-2	NM_015234	221395	GPR116	G protein-coupled receptor 116	212950_at
Stromal-2	NM_006108	10418	SPON1	spondin 1, extracellular matrix protein	213993_at
Stromal-2	NM_016215	51162	EGFL7	EGF-like-domain, multiple 7	218825_at
Stromal-2	NM_022481	64411	CENTD3	centaurin, delta 3	218950_at
Stromal-2	XM_371262	64123	ELTD1	EGF, latrophilin and seven transmembrane domain containing 1	219134_at
Stromal-2	NM_016563	51285	RASL12	RAS-like, family 12	219167_at
Stromal-2	NM_006094	10395	DLC1	deleted in liver cancer 1	224822_at
Stromal-2	NM_019035	54510	PCDH18	protocadherin 18	225975_at
Stromal-2	NM_019055	54538	ROBO4	roundabout homolog 4, magic roundabout (Drosophila)	226028_at
Stromal-2	NM_002207	3680	ITGA9	integrin, alpha 9	227297_at
Stromal-2	XM_930608	641700	ECSM2	endothelial cell-specific molecule 2	227779_at
Stromal-2	XM_037493	85358	SHANK3	SH3 and multiple ankyrin repeat domains 3	227923_at
Stromal-2	NM_052954	116159	CYYR1	cysteine/tyrosine-rich 1	228665_at
Stromal-2	NM_002837	5787	PTPRB	protein tyrosine phosphatase, receptor type, B	230250_at
Stromal-2	NM_019558	3234	HOXD8	homeobox D8	231906_at
Stromal-2	NM_001442	2167	FABP4	fatty acid binding protein 4, adipocyte	235978_at
Stromal-2	NM_024756	79812	MMRN2	multimerin 2	236262_at

Signature	GenBank Accession No.	Entrez GeneID	Gene Symbol	Gene Title	Affymetrix Probe Set ID
Stromal-2	BQ897248			Transcribed locus	242680_at
Stromal-2	NM_020663	57381	RHOJ	ras homolog gene family, member J	243481_at
Stromal-2	AK091419			CDNA FLJ34100 fis, clone FCBBF3007597	1558397_at
Stromal-2	NM_015719	50509	COL5A3	collagen, type V, alpha 3	52255_s_at
Stromal-2	NM_012072	22918	CD93	CD93 molecule	202878_s_at
Stromal-2	NM_000300	5320	PLA2G2A	phospholipase A2, group IIA (platelets, synovial fluid)	203649_s_at
Stromal-2	NM_019105	7148	TNXB	tenascin XB	206093_x_at
Stromal-2	NM_030754	6289	SAA2	serum amyloid A2	208607_s_at
Stromal-2	NM_019105	7148	TNXB	tenascin XB	208609_s_at
Stromal-2	NM_014220	4071	TM4SF1	transmembrane 4 L six family member 1	209387_s_at
Stromal-2	NM_000668	125	ADH1B	alcohol dehydrogenase IB (class I), beta polypeptide	209612_s_at
Stromal-2	NM_000668	125	ADH1B	alcohol dehydrogenase IB (class I), beta polypeptide	209613_s_at
Stromal-2	NM_001354	1646	AKR1C2	aldo-keto reductase family 1, member C2 (dihydrodiol dehydrogenase 2; bile acid binding protein; 3-alpha hydroxysteroid dehydrogenase, type III)	209699_x_at
Stromal-2	NM_001032281	7035	TFPI	tissue factor pathway inhibitor (lipoprotein-associated coagulation inhibitor)	210664_s_at
Stromal-2	NM_001001924	57509	MTUS1	mitochondrial tumor suppressor 1	212096_s_at
Stromal-2	NM_019105	7148	TNXB	tenascin XB	213451_x_at
Stromal-2	NM_004449	2078	ERG	v-ets erythroblastosis virus E26 oncogene homolog (avian)	213541_s_at
Stromal-2	NM_018407	55353	LAPTM4B	lysosomal associated protein transmembrane 4 beta	214039_s_at
Stromal-2	NM_000331	6288	SAA1	serum amyloid A1	214456_x_at
Stromal-2	NM_019105	7148	TNXB	tenascin XB	216333_x_at
Stromal-2	NM_001034954	10580	SORBS1	sorbin and SH3 domain containing 1	218087_s_at
Stromal-2	NM_017734	54873	PALMD	palmdelphin	218736_s_at
Stromal-2	NM_024756	79812	MMRN2	multimerin 2	219091_s_at
Stromal-2	NM_006744	5950	RBP4	retinol binding protein 4, plasma	219140_s_at
Stromal-2	NM_001034954	10580	SORBS1	sorbin and SH3 domain containing 1	222513_s_at

[0073] The DLBCL survival predictors of the invention were generated using expression data and methods described in Examples 1 and 2, below. The first bivariate survival

predictor incorporates the GCB and stromal-1 gene expression signatures. Fitting the Cox proportional hazards model to the gene expression data obtained from these two signatures resulted in a bivariate model survival predictor score calculated using the following generalized equation:

$$\text{Bivariate DLBCL survival predictor score} = A - [(x) * (\text{GCB signature value})] - [(y) * (\text{stromal-1 signature value})].$$

In this equation, A is an offset term, while (x) and (y) are scale factors. The GCB signature value and the stromal-1 signature value can correspond to the average of the expression levels of all genes in the GCB signature and the stromal-1 signature, respectively. A lower survival predictor score indicates a more favorable survival outcome, and a higher survival predictor score indicates a less favorable survival outcome for the subject.

[0074] The bivariate survival predictor was refined into a multivariate survival predictor that incorporates GCB, stromal-1, and stromal-2 gene expression signatures. Fitting the Cox proportional hazards model to the gene expression data obtained from these three signatures resulted in a multivariate model survival predictor score calculated using the following generalized equation:

$$\text{General multivariate DLBCL survival predictor score} = A - [(x) * (\text{GCB signature value})] - [(y) * (\text{stromal-1 signature value})] + [(z) * (\text{stromal-2 signature value})].$$

In this equation, A is an offset term, while (x), (y), and (z) are scale factors. The GCB signature value, the stromal-1 signature value, and the stromal-2 signature value can correspond to the average of the expression levels of all genes in the GCB signature, the stromal-1 signature, and the stromal-2 signature, respectively. A lower survival predictor score indicates a more favorable survival outcome and a higher survival predictor score indicates a less favorable survival outcome for the subject.

[0075] In one embodiment, the invention provides the following multivariate survival predictor equation:

$$\text{Multivariate DLBCL survival predictor score} = 8.11 - [0.419 * (\text{GCB signature value})] - [1.015 * (\text{stromal-1 signature value})] + [0.675 * (\text{stromal-2 signature value})]$$

In this equation, a lower survival predictor score indicates a more favorable survival outcome, and a higher survival predictor score indicates a poorer survival outcome for the subject.

[0076] In other embodiments of the multivariate DLBCL survival predictor score equation, the offset term (A) or (8.11) can be varied without affecting the equation's

usefulness in predicting clinical outcome. Scale factors (x), (y), and (z) can also be varied, individually or in combination. For example, scale factor (x) can be from about 0.200 or more, from about 0.225 or more, from about 0.250 or more, from about 0.275 or more, from about 0.300, from about 0.325 or more, from about 0.350 or more, from about 0.375 or more, or from about 0.400 or more. Alternatively, or in addition, scale factor (x) can be about 0.625 or less, about 0.600 or less, about 0.575 or less, about 0.550 or less, about 0.525 or less, about 0.500 or less, about 0.475 or less, about 0.450 or less, or about 0.425 or less. Thus, scale factor (z) can be one that is bounded by any two of the previous endpoints. For example scale factor (x) can be a value from 0.200-0.625, from 0.350-0.550, from 0.350-0.475, or from 0.400-0.425. Similarly, scale factor (y) can be from about 0.800 or more, from about 0.825 or more, from about 0.850 or more, from about 0.875 or more, from about 0.900 or more, from about 0.925 or more, from about 0.950 or more, from about 0.975 or more, or from about 1.000 or more. Alternatively, or in addition, scale factor (y) can be, e.g., about 1.250 or less, e.g., about 1.225 or less, about 1.200, about 1.175 or less, about 1.150 or less, about 1.125 or less, about 1.100 or less, about 1.075 or less, about 1.050 or less, or about 1.025 or less. Thus, scale factor (y) can be one that is bounded by any two of the previous endpoints. For example, scale factor (y) can be a value from 0.800-1.250, a value from 0.950-1.1025, a value from 0.950-1.200 or a value from 1.000-1.025. Also similarly, scale factor (z) can be from about 0.450 or more, about 0.475 or more, about 0.500 or more, about 0.525 or more, about 0.550 or more, about 0.575 or more, about 0.600 or more, about 0.625 or more, or about 0.650 or more. Alternatively, or in addition, scale factor (z) can be, e.g., about 0.900 or less, e.g., about 0.875 or less, about 0.850, about 0.825 or less, about 0.800 or less, about 0.775 or less, about 0.750 or less, or about 0.725 or less. Thus, scale factor (z) can be one that is bounded by any two of the previous endpoints. For example, scale factor (z) can be a value from 0.450-0.900, any value from 0.650-0.725, any value from 0.625-0.775 or any value from 0.650-0.700.

[0077] Furthermore, the invention includes any set of scale factors (x), (y), and (z) in conjunction in the general multivariate DLBCL survival predictor score that creates a function that is monotonically related to a multivariate DLBCL survival predictor score equation using any combination of the foregoing specified scale factor (x), (y), and (z) values.

[0078] In some embodiments of the invention, a survival predictor score can be calculated using fewer than all of the gene components of the GCB signature, the stromal-1

signature, and/or the stromal-2 signature listed in Table 1. For example, the survival prediction equations disclosed herein can be calculated using mathematical combinations of the expressions of 98% (38), 95% (37), 93% (36), or 90% (35) of the genes listed in Table 1 for the GCB signature, about 99% (about 280), about 98% (about 277), 97% (about 275), about 96% (about 272), about 95% (about 270), about 94% (about 266), about 93% (about 263), about 92% (about 260), about 91% (about 257), or about 90% (about 255) of the genes listed in Table 1 for the stromal-1 signature, and/or 99% (71), 97% (70), 96% (69), 95% (68), 93% (67), 92% (66), or 90% (65) of the genes listed in Table 1 for the stromal-2 signature (instead of using all of the genes corresponding to a gene signature in Table 1 to calculate the GCB signature value, the stromal-1 signature value, and/or stromal-2 signature value, respectively). In other embodiments, the survival prediction equations disclosed herein can be calculated using mathematical combinations of the expressions of 88% (34 genes), 85% (33 genes), 82% (32 genes), 80% (31 genes) of the genes listed in Table 1 for the GCB signature, about 89% (about 252), about 88% (about 249), about 87% (about 246), about 86% (about 243), about 85% (about 241), about 84% (about 238), about 83% (about 235), about 82% (about 232), about 81% (about 229), or about 80% (about 226) of the genes listed in Table 1 for the stromal-1 signature, and/or 89% (64), 88% (63), 86% (62), 85% (61), 83% (60), 82% (59) or 80% (58) of the genes listed in Table 1 for the stromal-2 signature (instead of using all of the genes corresponding to a gene signature in Table 1 to calculate the GCB signature value, the stromal-1 signature value, and/or stromal-2 signature value, respectively).

[0079] The invention also provides a method of using a DLBCL survival predictor score to predict the probability of a survival outcome beyond an amount of time t following treatment for DLBCL. The method includes calculating the probability of a survival outcome for a subject using the following general equation:

$$P(SO) = SO_0(t)^{(\exp((s) * (\text{survival predictor score})))}$$

In this equation, $P(SO)$ is the subject's probability of the survival outcome beyond time t following treatment for DLBCL, $SO_0(t)$ is the probability of survival outcome, which corresponds to the largest time value smaller than t in a survival outcome curve, and (s) is a scale factor. Treatment for DLBCL can include chemotherapy and the administration of Rituximab. A survival curve can be calculated using statistical methods, such as the Cox Proportional Hazard Model. Additional information regarding survival outcome curves is set forth in Lawless, Statistical Models and Methods for Lifetime Data, John Wiley and Sons (New York 1982) and Kalbfleisch et al., *Biometrika*, 60: 267-79 (1973).

[0080] In one embodiment, the method of the invention includes calculating the probability of overall survival for a subject beyond an amount of time t following treatment for DLBCL. The method includes calculating the probability of a survival outcome for a subject using the following general equation:

$$P(OS) = SO_0(t)^{(\exp(\text{survival predictor score}))}$$

In the equation, $P(OS)$ is the subject's probability of overall survival beyond time t following treatment for DLBCL, $SO_0(t)$ is the curve probability of survival outcome, which corresponds to the largest time value in a survival curve which is smaller than t , and the general equation scale factor (s) = 1. Treatment for DLBCL can include chemotherapy alone or in combination with the administration of Rituximab (R-CHOP).

[0081] In another embodiment, the method of the invention includes calculating the probability of progression-free survival for a subject beyond an amount of time t following treatment for DLBCL. The method includes calculating the probability of a survival outcome for a subject using the following general equation:

$$P(PFS) = SO_0(t)^{(\exp(0.976 * (\text{survival predictor score})))}$$

In this equation, $P(PFS)$ is the subject's probability of progression-free survival beyond time t following treatment for DLBCL, $SO_0(t)$ is the curve probability of progression-free survival, which corresponds to the largest time value in a survival curve which is smaller than t , and the general equation scale factor (s) = 0.976. The treatment for DLBCL can include chemotherapy alone or in combination with the administration of Rituximab (R-CHOP).

[0082] The foregoing equations for $P(OS)$ and $P(PFS)$ were generated by maximizing the partial likelihoods of the Cox proportional hazards model within the LLMPP CHOP data described below in Examples 1 and 2. Separate single variable Cox proportional hazards models were considered for overall survival $P(OS)$ and for progression free survival $P(PFS)$ based on this model score formulation. The single variable scale factor (1.0 for overall survival and 0.997 for progression free survival) were generated for each model by maximization of the partial likelihoods within the R-CHOP patients described below in Examples 1 and 2.

[0083] In other embodiments, the scale factor in the foregoing $P(PFS)$ can be varied such that (instead of 0.976) scale factor (s) is a value between 0.970 and 0.980, e.g. 0.971, 0.972, 0.973, 0.974, 0.975, 0.977, 0.978, and 0.979.

[0084] The invention also provides a method of selecting a subject for antiangiogenic therapy of DLBCL based on the subject's high relative expression of stromal-2 signature

genes. As discussed more fully below in Example 4, the stromal-2 signature includes a number of genes whose expression or gene products are related to angiogenesis. Thus, high relative expression of stromal-2 signature genes in DLBCL can be indicative of high angiogenic activity. Moreover, high relative expression of stromal-2 signature genes can be related to the heavy infiltration of some DLBCL tumors with myeloid lineage cells.

Accordingly, subjects with high relative expression of stromal-2 signature genes are good candidates for treatment with antiangiogenic therapy, either alone or in combination with other anti-oncogenic therapies. Furthermore, as also discussed more fully in Example 4, a stromal score, which was obtained by subtracting the stromal-1 signature value from the stromal-2 signature value, was observed to correlate with high tumor blood vessel density.

[0085] In this regard, the antiangiogenic monoclonal antibody to vascular endothelial growth factor bevacizumab has been clinically tested in patients with DLBCL (Ganjoo et al., *Leuk. Lymphoma*, 47: 998-1005 (2006)). Other antiangiogenic therapies can include small molecule inhibitors of SDF-1 receptor, such as CXCR4 (Petit et al., *Trends Immunol.*, 28: 299-307 (2007)). Still another example of an antiangiogenic therapy can include blocking antibodies to the myeloid lineage cell marker CTGF, which has been implicated in angiogenesis. Moreover, anti-CTGF antibodies have been shown to have anti-cancer activity in pre-clinical models of cancer (Aikawa et al., *Mol. Cancer Ther.*, 5: 1108-16 (2006)).

[0086] In one embodiment, the method of the invention for selecting a subject for antiangiogenic therapy includes obtaining a gene expression profile from a DLBCL biopsy from the subject. The subject's stromal-2 signature value is determined. The subject's stromal-2 signature value is then compared to a standard stromal-2 value. A standard stromal-2 value corresponds to the average of multiple stromal-2 signature values in DLBCL biopsy samples from a plurality of randomly selected subjects with DLBCL, e.g., more than 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, or 250 randomly selected subjects with DLBCL. If the subject's stromal-2 signature value is significantly higher than the standard stromal-2 value, then the subject can be treated with anti-angiogenic therapy.

[0087] In another embodiment, the method of the invention for selecting a subject for anti-angiogenic therapy includes obtaining a gene expression profile from a DLBCL biopsy from the subject. The subject's stromal 1 signature value and stromal-2 signature value are determined. The stromal-1 signature value is then subtracted from the stromal-2 signature value to obtain a stromal score. The subject's stromal score is then compared to a standard stromal score. A standard stromal score corresponds to the average of multiple stromal

scores (each stromal score = [stromal-2 signature value] - [stromal-1 signature value]) derived from DLBCL biopsy samples from a plurality of randomly selected subjects with DLBCL, e.g., more than 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 150, 200, or 250 randomly selected subjects with DLBCL. If the subject's stromal score is significantly higher than the standard stromal score, then the subject can be treated with anti-angiogenic therapy.

[0088] The invention further provides a targeted array that can be used to detect the expression levels of all or most of the genes in a germinal center B cell gene (GCB) expression signature, a stromal-1 gene expression signature, and/or a stromal-2 gene expression signature. A targeted array, as used herein, is an array directed to a limited set of genes and thus differs from a whole genome array. The targeted array of the invention can include probes for fewer than 20,000 genes, fewer than 15,000 genes, fewer than 10,000 genes, fewer than 8,000 genes, fewer than 7,000 genes, fewer than 6,000 genes, fewer than 5,000 genes, or fewer than 4,000 genes. Generally, the targeted array includes probes for at least 80% of the genes in a germinal center B cell gene (GCB) expression signature, a stromal-1 gene expression signature, and/or a stromal-2 gene expression signature. The targeted arrays of the invention can be used, for example, to detect expression levels for use in the methods described herein.

[0089] The invention provides a targeted array that includes probes for all of the genes in the stromal-1 gene expression signature. The invention also provides a targeted array that includes probes for all of the genes in the stromal-2 gene expression signature. Additionally, the invention provides a targeted array that includes probes for all of the genes in the stromal-1 gene expression signature and all of the genes in the stromal-2 gene expression signature. Moreover, the invention provides a targeted array that includes probes for all of the genes in the stromal-1 gene expression signature, all of the genes in the stromal-2 gene expression signature, and all of the genes in the GCB signature.

[0090] In certain embodiments, the arrays of the invention can include 98% (38), 95% (37), 93% (36), or 90% (35) of the genes listed in Table 1 for the GCB signature, about 99% (about 280), about 98% (about 277), 97% (about 275), about 96% (about 272), about 95% (about 270), about 94% (about 266), about 93% (about 263), about 92% (about 260), about 91% (about 257), or about 90% (about 255) of the genes listed in Table 1 for the stromal-1 signature, and/or 99% (71), 97% (70), 96% (69), 95% (68), 93% (67), 92% (66), or 90% (65) of the genes listed in Table 1 for the stromal-2 signature (instead of all of the genes listed in Table 1 for the GCB signature average, the stromal-1 signature average, and/or stromal-2

signature average, respectively). In certain embodiments, the arrays of the invention can include 88% (34 genes), 85% (33 genes), 82% (32 genes), 80% (31 genes) of the genes listed in Table 1 for the GCB signature, about 89% (about 252), about 88% (about 249), about 87% (about 246), about 86% (about 243), about 85% (about 241), about 84% (about 238), about 83% (about 235), about 82% (about 232), about 81% (about 229), or about 80% (about 226) of the genes listed in Table 1 for the stromal-1 signature, and/or 89% (64), 88% (63), 86% (62), 85% (61), 83% (60), 82% (59) or 80% (58) of the genes listed in Table 1 for the stromal-2 signature (instead of all of the genes listed in Table 1 for the GCB signature average, the stromal-1 signature average, and/or stromal-2 signature average, respectively).

[0091] The following examples further illustrate the invention but, of course, should not be construed as in any way limiting its scope.

EXAMPLE 1

[0092] This example demonstrates that significant differences were found between the survival outcomes for R-CHOP treated ABC DLBCL and GCB DLBCL patients and that survival outcome correlated with three prognostic gene expression signatures.

[0093] Pre-treatment tumor biopsy specimens and clinical data were obtained from 414 patients with *de novo* DLBCL treated at 10 institutions in North America and Europe and studied according to a protocol approved by the National Cancer Institute's Institutional Review Board. Patients included in a "LLMP CHOP cohort" of 181 patients were treated with anthracycline-based combinations, most often cyclophosphamide, doxorubicin, vincristine, and prednisone (CHOP) or similar regimens, as previously described (Rosenwald et al., *N. Engl. J. Med.*, 346: 1937-47 (2002)). The remaining 233 patients constituted an R-CHOP cohort that received similar chemotherapy plus Rituximab. The median follow-up in the R-CHOP cohort was 2.1 years (2.8 years for survivors). A panel of expert hematopathologists confirmed the diagnosis of DLBCL using current WHO criteria. Additional clinical patient characteristics for the R-CHOP cohort are described in Table 2. Additional analysis used a second "MMMNLP CHOP" cohort of 177 patients studied by the Molecular Mechanisms of Non-Hodgkin's Lymphoma Network Project (Hummel et al., *N. Engl. J. Med.*, 354: 2419-30 (2006)).

Table 2: Clinical characteristics of DLBCL patients treated with R-CHOP

Characteristic	% Total (N=233)	% Germinal center B cell-like DLBCL (N=107)	% Activated B cell-like DLBCL (N=93)	% Unclassified DLBCL (N=33)	P-value
Age > 60 yr	52	47	63	39	0.02
Ann Arbor stage > II	54	48	62	50	0.06
Lactate Dehydrogenase > 1x Normal	48	43	58	41	0.06
No. of extranodal sites >1	15	14	15	14	0.8
Eastern Cooperative Oncology Group (ECOG) performance status	25	17	33	27	0.02
International Prognostic Index (IPI) Score					<0.001
0 or 1	41	55	21	50	
2 or 3	46	33	63	38	
4 or 5	13	12	15	12	
Revised IPI Score					<0.001
0	19	27	5	28	
1 or 2	56	52	64	48	
3-5	25	21	31	24	

[0094] Gene expression profiling was performed using Affymetrix U133+ 2.0 microarrays. Gene expression profiling data are available through the National Center for Biotechnology Information web site as described in Lenz et al., *New Engl. J. Med.*, 359: 2313-23 (2008), at page 2314. All gene expression array data were normalized using MAS 5.0 software, and were log₂ transformed. To account for technical differences in the microarray processing between the R-CHOP cohort data and the LLMPP CHOP cohort data, the expression values of each gene in the R-CHOP cohort data were adjusted so that its median matched the median of the LLMPP CHOP data.

[0095] Gene expression signature identification and survival predictor model development were based solely on the data from the LLMPP CHOP training set. No prior survival analysis or subgroup analysis was performed with the test sets (MMMLNP CHOP and R-CHOP cohorts). The Cox model was used to identify genes associated with survival in

the CHOP training set and to build multivariate survival models. The models and their associated scaling coefficients were fixed based on the CHOP training set and then evaluated on the test sets. The P-values of survival effects of continuous variables such as gene expression or signature expression were calculated with the Cox likelihood ratio test. The significance of survival effects based on discrete variables such as lymphoma subtype or International Prognostic Index (IPI) was calculated using the log rank test. Validation P-values presented are one-sided in the direction observed in the training set. All other P-values were two sided. Survival curves were estimated using the Kaplan-Meier method.

[0096] All aspects of gene expression signature identification and survival predictor model development were based solely on the data from the CHOP training set. No prior survival analysis or subgroup analysis was performed with the test sets (MMMLNP CHOP and R-CHOP cohorts). The Cox model was used to identify genes associated with survival in the CHOP training set and to build multivariate survival models. The models and their associated scale factors were fixed based on the CHOP training set, and then evaluated on the test sets.

[0097] Since ABC and GCB DLBCL subtypes have distinct overall survival rates with CHOP chemotherapy (Rosenwald et al., *N. Engl. J. Med.*, 346: 1937-47 (2002); Alizadeh et al., *Nature*, 403:503-11(2000); Hummel et al., *N. Engl. J. Med.*, 354:2419-30 (2006); Monti, *Blood*, 105:1851-61(2005)), whether this distinction remains prognostically significant among patients treated with R-CHOP was tested (Coiffier et al., *N. Engl. J. Med.*, 346: 235-42 (2002)). Gene expression profiles were determined for pre-treatment biopsy samples from a “training set” of 181 patients treated with CHOP or CHOP-like chemotherapy alone and from a “test set” of 233 patients treated with R-CHOP. The patients in these two cohorts were comparable with respect to age range and distribution of the clinical prognostic variables that constitute the International Prognostic Index (IPI) (Table 2). In the R-CHOP cohort, patients with GCB DLBCL had better survival rates than those with ABC DLBCL. Specifically, R-CHOP treated GCB DLBCL and ABC DLBCL patients had 3-year overall survival rates of 84% and 56%, respectively, and 3-year progression-free survival rates of 74% and 40%, respectively (Fig. 1A and 1B). In the CHOP training set, and in a second “MMMLMP” CHOP cohort (Hummel et al., *supra*), the overall survival rates for ABC DLBCL and GCB DLBCL were lower than in the R-CHOP cohort (Fig. 6). Multivariate analysis indicated that the relative benefit (i.e., change in survival outcome) due to R-CHOP therapy (as compared to CHOP) was not significantly different between ABC and GCB DLBCL.

[0098] Four gene expression signatures have been previously shown to have prognostic significance in DLBCL patients treated with CHOP (Rosenwald et al., *supra*). Of these, the GCB signature and lymph node signature were associated with favorable survival, and the proliferation signature was associated with inferior survival within the CHOP training set, in the MMMLNP CHOP cohort (see the corresponding signature panels in Figure 7), and in the R-CHOP cohort (see corresponding signature panels in Figure 1C). Thus, the biological differences among DLBCL tumors reflected by these three signatures remain prognostically important in Rituximab treated patients, even though Rituximab treatment generally improved survival in DLBCL.

[0099] The remaining fourth gene expression signature, the MHC class II signature, which was associated with survival in the CHOP training set when treated as a continuous variable, was not associated with survival in the R-CHOP cohort (see MHC class II signature panel in Figure 1C). Moreover, tumors with extremely low “outlier” expression of this signature were associated with inferior survival in both CHOP cohorts (see Figures 8A and 8B), but not in the R-CHOP cohort (see Figure 8C).

[00100] The foregoing results indicate that Rituximab immunotherapy combined with chemotherapy (R-CHOP) benefits both the ABC and GCB subtypes of DLBCL and that gene expression signatures that predicted survival in the context of CHOP chemotherapy retained their prognostic power among R-CHOP-treated patients.

[00101] The foregoing results also indicate that the biological variation among DLBCL tumors, as measured by gene expression signatures, has a consistent relationship to therapeutic response regardless of the treatment regimen used. There is a striking difference in 3-year progression-free survival between ABC DLBCL patients and GCB DLBCL patients treated with R-CHOP (40% vs. 74%). This difference is likely due to genetic and biological differences between these DLBCL subtypes (Staudt et al., *Adv. Immunol.*, 87: 163-208 (2005)).

[00102] Hence, future clinical trials in DLBCL should incorporate quantitative methods to discern these biological differences so that patient cohorts in different trials can be compared and treatment responses can be related to defined tumor phenotypes.

EXAMPLE 2

[0100] This example demonstrates the development of GCB, stromal-1, and stromal-2 survival signatures and a related multivariate model of survival for R-CHOP-treated DLBCL.

[0101] Unless otherwise indicated, patient cohorts and methods of gene expression analysis are as described in Example 1.

[0102] In the LLMPP CHOP cohort data, 936 genes were identified as associated with poor prognosis $p < 0.01$ (1-sided). For genes having multiple array probe sets associated with survival, only the probe set with the strongest association with survival was used. The expression values of the probe sets in the LLMPP CHOP cohort data were then clustered. The largest cluster with an average correlation of >0.6 and containing *myc* was identified as the proliferation survival signature. 1396 genes were identified as associated with favorable outcome. The largest cluster with average correlation of >0.6 and containing *BCL6* was identified as the germinal center B cell (GCB) survival signature. A cluster with average correlation of >0.6 and containing *FN1* was identified as the stromal-I survival signature, whereas another cluster with average correlation of >0.6 containing *HLADRA* was identified as the MHC class II survival signature. The expression levels of genes within each signature were then averaged to create a "signature average" for each biopsy specimen. For the MMMLNP CHOP data set, the average was calculated for those array elements represented on the Affymetrix U133A microarray.

[0103] From the four prognostic clusters or signatures, two signatures, the stromal-I and the GCB signatures were used to create the best two variable survival model. Neither the proliferation nor the MHC class II signatures added to the prognostic value of this two variable model. This bivariate model performed well in the MMMLNP CHOP cohort (Figure 9A) and in the R-CHOP cohort (Figure 2A).

[0104] The CHOP training set was used to discover and refine signatures that added to the prognostic significance of this bivariate model, and the resulting multivariate models were tested in the R-CHOP cohort. 563 genes were identified as adding to the model in the direction of adverse prognosis. These genes were clustered by hierarchical clustering, and three clusters of more than 10 genes with an average correlation of >0.6 were identified. In addition, 542 genes were identified which added to the stromal-I and GCB signature model in the direction of favorable prognosis. These genes were clustered, and two clusters of more than 10 genes with an average correlation of >0.6 were identified. Signature averages were determined for these clusters, and three variable models containing the stromal-I and GCB signature and each of the cluster averages were formed on the MMMLNP CHOP and R-CHOP data sets. Of the five cluster averages, two were found to add statistical significance ($p < 0.02$) in the MMMLNP CHOP data as compared to a model containing the stromal-I and

GCB signatures alone. By contrast, in the R-CHOP data, three of the five cluster averages were found to add significance ($p < 0.02$) to the bivariate model. One of these cluster averages added significantly to the bivariate model in both the MMMLNP CHOP and R-CHOP data. This signature, designated Signature 122, was also found to add to the stromal-1 and GCB signature far more significantly than any of the four other signatures on the LLMPP CHOP data and, thus, was retained for further analysis.

[0105] Signature 122 added significantly to the bivariate model in both the MMMLNP CHOP cohort ($p = 0.011$) and in the R-CHOP cohort ($p = 0.001$) (Figures 9 B and 9C). This Signature 122 positively correlated with the stromal-1 signature, although it was associated with adverse survival when added to the bivariate model. To further refine our model, we identified genes that were more correlated with Signature 122 than with the stromal-1 signature ($p < 0.02$). These genes were organized by hierarchical clustering, and three sets of correlated genes ($r > 0.6$) were observed. One of these clusters, the stromal-2 signature, added to the significance of the bivariate model in both the MMMLNP CHOP cohort ($p = 0.002$) and the R-CHOP cohort ($p < 0.001$) (Figures 2B and 9D).

[0106] A multivariate survival model was formed by fitting a Cox model with the GCB, stromal-1, and stromal-2 signatures to the LLMPP CHOP cohort data shown in Table 3. This final multivariate model with its associated scaling coefficients was then evaluated on the MMLLMPP CHOP and R-CHOP cohort data sets. Survival predictor scores from the final model were used to divide the R-CHOP cohort into quartile groups with 3-year overall survival rates of 89%, 82%, 74%, and 48%, and 3-year progression-free survival rates of 84%, 69%, 61% and 33% (Figure 2B). The survival predictor scores from the final model are illustrated in Figure 3 along with the three component signatures and representative genes of each signature.

TABLE 3

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
2	2.75	0	2.75	0	9.238	8.778	7.475	0.376
3	2.67	0	2.67	0	9.942	8.227	7.102	0.387
5	1.27	1	0.72	1	8.859	9.033	8.716	1.113
21	2.39	0	2.40	0	10.573	8.519	6.959	-0.270
22	2.38	0	2.38	0	8.737	8.686	7.598	0.761
23	2.52	0	2.52	0	10.694	10.322	8.817	-0.897
24	5.11	0	5.11	0	11.376	7.854	7.598	0.500
26	4.01	0	4.01	0	9.829	9.956	8.507	-0.372
28	3.96	0	3.96	0	10.957	9.277	8.248	-0.330
41	0.52	1	0.52	1	9.273	9.437	8.202	0.183
47	1.53	1	0.77	1	9.548	8.802	8.061	0.617
48	0.37	1	0.12	1	8.660	8.279	6.891	0.729
49	2.37	0	2.35	1	10.915	8.988	6.847	-0.965
53	3.89	0	2.23	1	9.530	9.792	9.693	0.721
61	0.90	1	0.46	1	8.649	8.038	8.104	1.798
65	4.04	0	4.04	0	10.744	9.330	7.930	-0.508
66	4.04	0	4.04	0	10.714	10.016	7.536	-1.459
95	0.62	1	0.44	1	9.244	9.197	8.105	0.373
96	5.37	0	5.37	0	10.107	8.723	7.608	0.157
97	5.07	0	5.07	0	9.777	9.192	7.359	-0.349
98	0.94	1	0.59	1	8.794	7.711	7.367	1.571
99	0.40	1	0.40	1	9.024	9.272	9.160	1.101
103	0.03	1	0.02	1	8.883	8.190	7.742	1.301
104	3.76	0	3.76	0	9.785	9.866	7.929	-0.652
106	2.95	0	2.95	0	10.585	7.797	6.824	0.367
107	2.94	0	2.94	0	11.535	8.358	6.660	-0.711

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
108	2.73	0	2.73	0	9.653	8.495	7.550	0.539
109	0.16	1	0.11	1	9.301	9.376	7.994	0.092
110	2.46	0	2.46	0	10.254	8.980	7.324	-0.357
111	2.44	0	2.44	0	10.137	10.691	8.948	-0.949
113	2.12	0	2.12	0	10.746	8.555	6.942	-0.390
114	1.98	0	0.88	1	8.562	8.159	7.120	1.047
115	1.92	0	1.92	0	10.313	9.385	8.157	-0.231
118	1.64	0	1.64	0	10.209	10.194	8.231	-0.959
119	1.60	0	1.60	0	11.059	8.852	7.479	-0.461
1087	0.05	1	0.05	1	8.756	8.491	7.949	1.188
1089	5.12	0	1.27	1	9.863	9.135	8.034	0.129
1091	5.15	0	5.15	0	10.454	9.918	8.742	-0.437
1092	5.06	0	5.07	0	9.452	9.467	8.912	0.556
1093	3.83	1	1.62	1	9.915	9.138	7.747	-0.090
1096	4.02	0	4.02	0	8.887	9.236	7.795	0.274
1097	1.26	1	1.08	1	11.219	9.234	8.321	-0.347
1098	3.53	0	3.53	0	9.117	9.236	7.655	0.082
1099	3.07	0	0.91	1	9.284	8.798	7.741	0.515
1101	5.64	0	5.64	0	9.803	9.466	8.156	-0.101
1108	3.30	0	3.30	0	9.195	10.456	9.065	-0.237
1109	3.78	0	3.78	0	11.008	10.051	8.273	-1.120
1164	0.19	1	0.16	1	9.242	10.307	10.548	0.896
1167	1.49	1	0.45	1	9.809	9.105	8.784	0.687
1168	0.42	1	0.30	1	8.718	8.368	7.149	0.790
1169	1.71	1	1.22	1	11.512	8.108	7.507	0.125
1172	2.82	0	2.82	0	11.137	8.871	8.153	-0.057
1173	0.87	1	0.79	1	11.324	9.914	8.514	-0.950
1175	1.06	1	0.56	1	9.107	10.310	9.063	-0.053

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
1179	2.53	0	2.53	0	9.506	9.437	8.461	0.260
1181	1.72	0	1.72	0	10.688	9.018	7.647	-0.360
1184	4.74	0	2.97	1	10.812	8.979	7.922	-0.187
1185	3.71	0	3.71	0	10.431	8.397	7.317	0.156
1186	3.43	0	3.43	0	8.688	8.944	8.552	1.164
1187	5.23	0	5.23	0	10.072	10.192	8.667	-0.604
1189	5.13	0	5.13	0	10.109	9.212	7.967	-0.097
1190	3.66	0	3.66	0	10.713	10.409	8.910	-0.930
1192	0.16	1	0.16	1	8.825	9.903	8.061	-0.199
1195	4.36	0	4.36	0	11.539	7.567	6.873	0.234
1197	3.13	0	3.13	0	10.287	10.365	9.549	-0.275
1200	0.31	1	0.31	1	9.432	8.950	9.805	1.692
1206	6.51	0	6.51	0	10.410	9.946	8.925	-0.323
1211	6.25	0	6.25	0	11.596	7.908	6.524	-0.372
1215	5.35	0	5.35	0	10.504	9.061	7.550	-0.392
1216	0.46	1	0.29	1	10.017	9.010	7.794	0.028
1219	0.51	1	0.51	1	10.614	10.014	8.619	-0.683
1220	2.24	1	2.25	1	8.850	9.400	8.036	0.286
1221	3.94	0	3.95	0	8.777	7.489	6.672	1.334
1222	3.53	0	3.53	0	10.463	9.310	7.019	-0.986
1224	3.22	0	2.11	1	9.751	9.505	8.453	0.082
1225	2.95	0	2.95	0	8.613	8.313	7.668	1.240
1226	0.08	1	0.08	1	9.229	8.851	7.950	0.625
1228	2.78	0	0.99	1	11.532	8.261	6.932	-0.428
1230	0.59	1	0.54	1	9.369	6.951	6.956	1.825
1231	1.41	0	1.41	0	10.248	8.788	8.011	0.303
1232	2.49	0	0.68	1	10.362	8.528	7.975	0.495
1233	2.50	0	2.50	0	9.239	10.581	8.470	-0.784

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
1236	2.56	0	2.56	0	9.156	10.000	7.805	-0.608
1238	0.16	1	0.16	1	9.488	9.055	8.256	0.517
1239	2.24	0	2.24	0	8.886	8.978	7.838	0.564
1240	1.48	0	1.48	0	10.474	9.073	7.702	-0.288
1241	1.41	1	1.17	1	9.044	9.054	7.451	0.160
1251	2.72	0	2.72	0	8.410	8.687	7.082	0.549
1252	0.01	1	0.01	1	11.167	8.070	7.358	0.206
1255	5.17	0	5.17	0	9.501	9.411	7.887	-0.099
1271	4.72	0	4.73	0	10.718	8.452	7.060	-0.194
1272	5.68	0	5.68	0	9.161	9.080	7.668	0.231
1275	1.89	1	1.48	1	9.257	8.559	8.607	1.354
1277	5.06	0	5.07	0	11.091	9.938	8.274	-1.038
1279	4.87	0	4.87	0	9.309	10.085	9.676	0.504
1281	3.36	0	not available (n/a)	n/a	9.535	9.969	9.090	0.132
1284	3.51	0	3.51	0	10.922	9.680	8.481	-0.567
1288	1.54	0	n/a	n/a	9.430	8.896	8.037	0.554
1289	0.03	1	0.03	1	8.915	9.052	8.002	0.589
1290	5.23	0	5.23	0	10.432	10.426	8.154	-1.340
1291	0.04	1	0.04	1	11.319	8.246	7.323	-0.059
1292	0.10	1	0.10	1	8.667	8.764	8.110	1.058
1293	4.81	0	4.81	0	11.116	9.842	8.083	-1.081
1294	0.53	1	0.53	1	10.138	10.181	8.501	-0.733
1295	5.16	0	5.17	0	9.445	9.694	7.739	-0.463
1296	4.79	0	4.79	0	10.228	9.064	8.852	0.600
1297	4.24	0	4.24	0	9.524	7.990	7.008	0.740
1298	4.56	0	4.56	0	9.022	9.000	7.695	0.389
1331	3.29	0	3.29	0	11.004	9.488	8.289	-0.536
1334	2.87	0	2.87	0	11.434	9.509	8.109	-0.859

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
1335	1.38	1	0.90	1	9.586	8.545	7.423	0.431
1336	2.44	0	2.44	0	10.844	9.704	7.706	-1.082
1337	0.02	1	0.02	1	8.521	7.788	7.860	1.941
1449	1.62	0	1.62	0	9.604	8.463	8.030	0.917
1450	1.30	0	0.53	1	8.571	8.112	7.241	1.173
1451	1.84	0	1.85	0	10.637	9.205	7.759	-0.452
1453	1.71	0	1.71	0	10.964	9.089	8.226	-0.157
1454	0.62	0	0.62	0	11.106	8.514	7.604	-0.052
1553	2.93	0	1.92	1	8.975	9.284	7.475	-0.029
1612	5.37	0	5.37	0	10.526	9.471	7.809	-0.643
1613	5.81	0	n/a	n/a	10.868	9.695	7.730	-1.067
1614	4.36	1	4.36	1	10.358	9.226	8.765	0.322
1617	0.52	0	0.52	0	10.332	8.723	7.180	-0.227
1618	1.70	0	0.98	1	11.233	8.956	7.852	-0.387
1619	0.25	1	0.25	1	8.646	8.028	7.123	1.146
1620	2.17	0	2.17	0	11.647	8.385	7.343	-0.325
1623	2.80	0	2.80	0	9.611	9.484	8.249	0.024
1626	1.76	0	1.76	0	11.236	9.495	8.108	-0.763
1628	3.13	0	1.23	1	8.714	7.972	7.149	1.192
1645	2.85	0	2.85	0	10.146	9.476	8.914	0.258
1647	2.79	0	2.80	0	10.485	10.495	8.707	-1.058
1650	0.75	1	0.75	1	8.830	7.346	6.486	1.333
1651	1.66	0	1.66	0	9.190	7.949	6.829	0.801
1652	1.64	0	n/a	n/a	8.798	8.943	8.331	0.969
1702	1.05	0	1.05	1	9.008	8.217	8.078	1.447
1703	0.70	1	0.70	1	9.499	8.637	7.790	0.621
1704	3.14	0	3.14	0	9.908	9.231	7.503	-0.347
1705	3.94	0	3.94	0	8.933	8.445	8.187	1.321

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
1707	2.80	0	2.80	0	10.610	9.348	7.872	-0.510
1742	3.27	0	n/a	n/a	10.033	8.715	7.412	0.063
1746	1.91	0	1.55	1	9.249	8.705	8.205	0.937
1747	1.48	0	1.48	0	10.162	8.866	7.602	-0.016
1756	3.47	0	3.47	0	10.815	9.638	7.248	-1.312
1761	0.23	1	0.23	1	9.842	10.192	8.664	-0.511
1762	5.20	0	5.20	0	10.583	9.333	7.445	-0.772
1763	5.51	0	5.51	0	8.917	8.925	8.084	0.771
1766	1.59	0	1.59	0	10.919	10.037	8.389	-0.990
1782	1.09	0	1.09	0	10.753	9.600	8.332	-0.516
1788	0.39	1	0.24	1	10.364	8.738	8.914	0.915
1861	0.56	1	0.19	1	9.728	8.604	7.594	0.427
1867	1.17	1	0.38	1	8.903	11.501	10.559	-0.166
1916	1.41	0	n/a	n/a	9.295	11.197	11.508	0.619
1920	1.32	0	1.32	0	10.165	9.630	8.789	0.009
1927	1.53	0	1.53	0	9.195	10.261	9.791	0.451
1928	0.72	0	0.72	0	9.769	8.510	7.330	0.328
1939	0.47	1	0.47	1	9.097	9.363	7.647	-0.043
2002	1.29	0	1.30	0	9.469	9.542	8.600	0.262
2006	1.23	0	1.23	0	10.434	8.223	7.162	0.227
2067	2.18	0	2.18	0	10.244	11.186	9.391	-1.197
2070	0.31	0	0.12	1	10.486	10.680	10.353	-0.135
2162	0.38	1	0.38	1	10.934	10.020	7.960	-1.268
2270	1.59	0	1.59	0	10.117	9.904	8.506	-0.440
2271	1.60	0	1.60	0	8.995	9.349	8.261	0.428
2274	0.41	0	0.41	0	8.863	7.623	7.222	1.533
2283	1.19	0	1.19	0	10.501	8.361	6.741	-0.226

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
2291	0.87	1	0.85	1	10.732	10.184	9.436	-0.353
2299	0.93	0	0.93	0	10.661	9.905	8.189	-0.883
2301	0.61	0	0.61	0	9.852	9.903	8.352	-0.432
2306	0.68	0	0.68	0	8.586	8.759	8.191	1.151
2309	0.43	0	0.43	0	10.839	7.671	6.860	0.413
2311	0.80	0	0.80	0	10.901	7.797	6.912	0.294
2318	0.99	0	0.99	0	10.283	9.403	8.655	0.100
2321	0.82	0	0.82	0	9.691	8.956	7.404	-0.044
2411	0.67	0	0.67	0	8.986	8.383	7.854	1.137
2415	0.62	0	0.62	0	9.296	10.509	9.551	-0.005
2444	3.99	0	3.99	0	10.154	9.871	9.026	-0.071
2445	3.36	0	3.36	0	8.788	8.184	7.964	1.497
2479	0.51	0	0.51	0	11.151	9.023	8.199	-0.186
2482	4.54	0	4.54	0	10.373	9.847	8.208	-0.691
2483	3.89	1	3.89	1	9.241	8.902	7.742	0.428
2484	2.69	1	1.90	1	10.279	9.619	8.312	-0.349
2485	4.43	0	4.43	0	9.957	9.865	8.439	-0.378
2486	4.37	0	n/a	n/a	10.698	10.203	8.041	-1.301
2487	4.34	0	4.34	0	11.227	9.909	8.260	-1.076
2488	4.20	0	4.21	0	9.510	8.709	7.615	0.426
2490	4.02	0	4.02	0	10.510	10.961	8.956	-1.374
2491	0.50	1	0.25	1	9.047	8.554	7.624	0.784
2492	3.96	0	3.96	0	9.904	10.901	9.140	-0.935
2497	3.44	0	3.44	0	9.221	9.438	8.065	0.111
2498	3.37	0	3.37	0	9.318	9.427	8.003	0.040
2500	3.31	0	3.31	0	11.014	9.406	7.375	-1.074
2501	3.28	0	n/a	n/a	8.822	8.551	7.750	0.966
2503	2.99	0	2.99	0	8.301	7.967	6.929	1.222

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
2504	2.78	0	2.78	0	10.145	8.004	7.017	0.472
2505	2.76	0	2.76	0	11.036	8.442	7.136	-0.266
2507	0.86	1	0.54	1	9.737	9.475	8.988	0.480
2508	2.58	0	2.58	0	8.678	9.389	8.230	0.498
2509	0.96	1	0.76	1	8.895	10.441	9.088	-0.081
2511	1.55	1	1.06	1	9.225	9.267	9.191	1.042
2512	2.45	0	2.45	0	11.047	10.465	9.337	-0.838
2513	0.61	1	0.61	1	10.855	10.378	8.395	-1.305
2514	2.18	0	2.18	0	10.477	9.832	7.498	-1.198
2515	2.13	0	2.13	0	9.295	10.519	9.788	0.145
2516	2.07	0	2.07	0	10.575	10.592	8.642	-1.238
2517	2.04	0	0.76	1	9.385	9.163	8.328	0.498
2584	0.68	0	0.68	0	10.759	9.356	8.135	-0.404
2599	4.05	0	4.05	0	10.629	9.158	7.724	-0.425
2600	1.01	1	0.54	1	9.785	8.619	7.291	0.184
2601	1.22	1	0.88	1	9.385	8.044	7.178	0.859
2603	4.43	0	4.43	0	9.582	10.707	9.803	-0.156
2604	0.84	0	0.36	1	9.844	10.511	8.382	-1.026
2609	8.89	0	2.55	1	8.981	8.775	7.506	0.507
2610	0.74	0	0.74	0	10.793	8.964	7.421	-0.502
2611	0.66	0	0.66	0	10.353	10.233	9.032	-0.518
2612	1.17	1	1.13	1	10.290	9.028	8.287	0.230
2613	1.66	0	1.66	0	10.997	9.089	7.749	-0.493
2614	0.21	1	0.21	1	8.768	7.850	7.100	1.261
2615	0.48	0	0.48	0	11.359	9.470	7.647	-1.100
2639	10.29	0	10.30	0	11.085	10.385	8.003	-1.674
2641	1.38	0	1.38	0	9.199	8.818	7.340	0.259
2642	3.67	0	3.67	0	10.731	8.777	7.167	-0.458

Patient	Time to death or last follow up (years)	Status at last follow up (1=dead, 0=alive)	Time to death, or progression, or last follow up (years)	Status at last follow up (1=progressed or died, 0=no progression)	Germinal Center Signature Average	Stromal-1 Signature Average	Stromal-2 Signature Average	Model Score
2643	5.49	0	5.49	0	10.236	10.578	8.473	-1.197
2645	0.19	0	n/a	n/a	11.130	9.997	8.254	-1.129
2646	0.18	1	0.18	1	8.893	7.648	6.871	1.260
2648	0.25	0	0.25	0	8.855	7.745	7.060	1.303
2649	2.13	0	2.13	0	9.688	10.354	9.885	0.214
2650	2.43	0	n/a	n/a	10.007	10.052	8.861	-0.305
2651	1.61	0	n/a	n/a	10.660	9.452	7.831	-0.665
2652	1.84	0	1.84	0	11.378	9.247	7.684	-0.856
2653	1.88	0	1.88	0	11.182	9.638	7.781	-1.106
2654	1.43	0	1.43	0	8.791	9.395	8.905	0.902
2813	3.97	0	3.97	0	10.701	9.366	8.258	-0.306
2814	0.81	1	0.70	1	10.561	9.176	9.275	0.632

[0107] The International Prognostic Index (IPI), which is based on 5 clinical variables, predicts survival in both CHOP-treated and R-CHOP-treated patients (Shipp et al., *N. Engl. J. Med.*, 329:987-94 (1993); Sehn et al., *Blood*, 109: 1857-61 (2007)). The inventive gene expression-based survival model retained its prognostic significance among R-CHOP-treated patients segregated according to IPI into high, intermediate and low IPI risk groups, both as originally defined (Shipp et al., *supra*) ($p < 0.001$) (Figure 2C) and as recently modified for R-CHOP-treated DLBCL (Sehn et al., *supra*) ($p < 0.001$) (Figure 10).

[0108] The foregoing results indicate that the gene expression-based multivariate model can be used to identify large disparities in survival among patients with different DLBCL gene signature profiles. Thus, survival predictor scores were used to divide patients into least and most favorable quartile groups having 3-year progression-free survival rates of 33% and 84%, respectively. Given its statistical independence from the IPI, the gene expression-based survival predictor provides a complementary view of DLBCL variation that can be considered when analyzing data from DLBCL clinical trials. Additionally, the foregoing results indicate that whole-genome gene expression profiles in conjunction with the survival model described herein can be used to provide optimal predictions of expected survival outcomes for subjects suffering from DLBCL.

EXAMPLE 3

[0109] This example demonstrates the use of a survival predictor score to predict the probability of progression free and overall survival outcomes at a period of time t following R-CHOP treatment in accordance with the invention.

[0110] RNA is isolated from a patient's DLBCL biopsy and hybridized to a U133+ array from Affymetrix (Santa Clara, CA). The array is scanned, and MAS 5.0 algorithm is applied to obtain signal values normalized to a target intensity of 500. Signal values are $\log_2$ transformed to intensity values. For genes of interest with multiple probe sets, the intensity value of the multiple probe sets are averaged to obtain a single intensity value for each gene. The single intensity values of genes in the GCB signature are averaged to obtain a GCB signature average of 9.2. The single intensity values of genes in the stromal-1 signature are averaged to obtain a stromal-1 signature average of 8.5. The single intensity values of genes in the stromal-2 signature are averaged to obtain a stromal-2 signature average of 7.2.

[0111] The patient's survival predictor score is calculated using the following equation $8.11 - [0.419 * (\text{GCB signature average})] - [1.015 * (\text{stromal-1 signature average})] +$

$[0.675 * (\text{stromal-2 signature average})]$, such that the survival predictor score = $8.11 - [0.419 * (9.2)] - [1.015 * (8.5)] + [0.675 * (7.2)] = 0.389$

[0112] Table 4 includes values from a progression free survival curve generated using baseline hazard functions calculated from the R-CHOP patient data described in Table 3. The curve was generated in accordance with the methods of Kalbfleisch and Prentice, *Biometrika*, 60: 267-279 (1973), which involves maximizing the full likelihood, under the assumption that the true scaling coefficients were equal to prior estimates. In Table 4, $F_0(t)$ is the probability of progression free survival for each indicated time period following R-CHOP treatment (t-RCHOP).

Table 4

t-RCHOP (years)	$F_0(t)$
0.000	1.000
0.008	0.997
0.016	0.993
0.025	0.990
0.030	0.987
0.036	0.983
0.049	0.980
0.082	0.977
0.096	0.973
0.107	0.970
0.118	0.967
0.120	0.963
0.156	0.960
0.156	0.956
0.159	0.953
0.178	0.950
0.192	0.946
0.211	0.943
0.233	0.939
0.241	0.936
0.246	0.932
0.252	0.928
0.290	0.925
0.298	0.921
0.307	0.918
0.364	0.914
0.381	0.910
0.381	0.907
0.400	0.903
0.441	0.899
0.446	0.895
0.463	0.891
0.468	0.887

t-RCHOP (years)	F ₀ (t)
0.515	0.884
0.517	0.880
0.531	0.876
0.534	0.872
0.537	0.868
0.537	0.864
0.539	0.860
0.561	0.856
0.586	0.852
0.611	0.848
0.679	0.843
0.698	0.839
0.698	0.834
0.720	0.830
0.747	0.826
0.756	0.821
0.761	0.816
0.767	0.812
0.786	0.807
0.849	0.803
0.879	0.798
0.884	0.793
0.898	0.789
0.912	0.784
0.977	0.779
0.986	0.774
1.046	0.770
1.057	0.765
1.076	0.760
1.128	0.755
1.166	0.750
1.216	0.745
1.227	0.740
1.270	0.735
1.481	0.729
1.547	0.724
1.624	0.718
1.900	0.711
1.919	0.705
2.105	0.699
2.231	0.692
2.245	0.685
2.352	0.678
2.546	0.671
2.968	0.662
3.890	0.648
4.364	0.623

[0113] The patient's probability of 2 year progression free survival is calculated using the equation: $P(\text{PFS}) = F_0(t)^{(\exp(0.976 * \text{survival predictor score}))}$, where $F_0(t)$ is the $F_0(t)$ value that corresponds to the largest time value smaller than 2 years in the progression free survival curve. In Table 4, the largest time value smaller than 2 is 1.919, and the corresponding $PF_0(t)$ value is 0.705. Accordingly, the patient's probability of 2 year progression free survival $P(\text{PFS}) = 0.705^{(\exp(0.976 * \text{survival predictor score}))} = 0.705^{1.462} = 0.600$ or about 60%.

[0114] Table 5 includes values from an overall survival curve generated using baseline hazard functions calculated from the R-CHOP patient data described in Table 3. The curve was made according to the method of Kalbfleisch and Prentice, *Biometrika*, 60: 267-279 (1973), which involves maximizing the full likelihood, under the assumption that the true scaling coefficients were equal to our estimates. In Table 5, $OS_0(t)$ is the probability of overall survival for each indicated time period following R-CHOP treatment (t-RCHOP).

Table 5

t-RCHOP (years)	$OS_0(t)$
0.000	1.000
0.008	0.997
0.016	0.994
0.030	0.991
0.033	0.988
0.036	0.984
0.049	0.981
0.082	0.978
0.096	0.975
0.156	0.972
0.156	0.969
0.159	0.965
0.178	0.962
0.192	0.959
0.211	0.956
0.233	0.952
0.246	0.949
0.307	0.946
0.367	0.942
0.380	0.939
0.386	0.935
0.402	0.932
0.416	0.928
0.463	0.925
0.468	0.921
0.504	0.918
0.515	0.914
0.517	0.910

t-RCHOP (years)	OS ₀ (t)
0.531	0.907
0.556	0.903
0.586	0.900
0.610	0.896
0.619	0.892
0.698	0.888
0.747	0.885
0.807	0.881
0.862	0.877
0.868	0.873
0.873	0.869
0.895	0.864
0.944	0.860
0.963	0.856
1.010	0.852
1.057	0.848
1.169	0.843
1.169	0.839
1.215	0.835
1.262	0.830
1.273	0.826
1.382	0.821
1.412	0.817
1.492	0.812
1.527	0.807
1.552	0.802
1.708	0.796
1.889	0.791
2.244	0.784
2.693	0.777
3.826	0.763
3.889	0.749
4.363	0.724

[0115] The patient's probability of 2 year overall survival is calculated using the equation: $P(OS) = OS_0(t)^{(\exp(\text{survival predictor score}))}$, where $OS_0(t)$ is the value that corresponds to the largest time value in the overall survival curve which is smaller than 2 years. In Table 5, the largest time value smaller than 2 is 1.889, and the corresponding $OS_0(t)$ value is 0.791. Accordingly, the patient's probability of 2 year overall survival is $P(PFS) = 0.791^{(\exp(0.389))} = 0.791^{1.4476} = 0.707$ or 70.7%.

EXAMPLE 4

[0116] This example demonstrates the biological basis for DLBCL prognostic signatures.

[0117] Unless otherwise indicated, cohorts and methods of gene expression analysis are described in Examples 1 and 2. Furthermore, cell suspensions from three biopsies were separated by flow cytometry into a CD19+ malignant subpopulation and a CD19- non-malignant subpopulation. Gene expression profiling was performed following two rounds of linear amplification from total RNA (Dave et al., *N. Engl. J. Med.*, 351: 2159-69 (2004)). After MAS5.0 normalization, genes were selected that had a log2 signal value greater than 7 in either the CD19+ or CD19- fractions in at least two of the sorted samples.

[0118] To assess whether the gene expression signatures in the final survival model of Example 2 were derived from the malignant lymphoma cells or from the host microenvironment, three DLBCL biopsy samples were fractionated into CD19+ malignant cells and CD19- non-malignant cells by flow sorting. Most germinal center B cell signature genes were more highly expressed in the malignant fraction, whereas genes from the stromal-1 and stromal-2 signatures were more highly expressed in the non-malignant stromal fraction (Figure 4A), hence their name. Since these two signatures were synergistic in predicting survival, they were combined into a “stromal score” (Figure 3), high values of which were associated with adverse outcome.

[0119] The germinal center B cell signature relates to the distinction between the ABC and GCB DLBCL subtypes (Figure 3). By contrast, the genes defining the stromal-1 signature encodes components of the extracellular matrix, including fibronectin, osteonectin, various collagen and laminin isoforms, and the anti-angiogenic factor thrombospondin (Figure 3 and Table 1). This signature also encodes modifiers of collagen synthesis (*LOXLI*, *SERPINH1*), proteins that remodel the extracellular matrix (*MMP2*, *MMP9*, *MMP14*, *PLAU*, *TIMP2*), and *CTGF*, a secreted protein that can initiate fibrotic responses (Frazier et al., *J. Invest. Dermatol.*, 107(3): 404-11 (1996)). In addition, the stromal-1 signature includes genes characteristically expressed in cells of the monocytic lineage, such as *CEBPA* and *CSF2RA*.

[0120] The stromal-1 signature is significantly related to several previously curated gene expression signatures (Shaffer et al., *Immunol. Rev.*, 210: 67-85 (2006)) based on gene set enrichment analysis (Subramanian et al., *Proc. Nat'l. Acad. Sci. USA*, 102(43): 15545-50 (2005)). Two of these signatures include genes that are coordinately expressed in normal mesenchymal tissues but not in hematopoietic subsets, many of which encode extracellular matrix proteins (false discovery rate (FDR) < 0.001) (Figures 4B and 11) (Su et al., *Proc. Nat'l. Acad. Sci. USA*, 101: 6062-7 (2004)). Also enriched was a “monocyte” signature,

comprised of genes that are more highly expressed in CD14+ blood monocytes than in B cells, T cells, or NK cells (FDR=0.014) (Figure 4B). By contrast, a pan-T cell signature was not related to the stromal-1 signature (Figure 4B). These findings suggest that high expression of the stromal-1 signature identifies tumors with vigorous extracellular matrix deposition and infiltration by cells in the monocytic lineage.

[0121] In this regard, the stromal-1 signature gene product fibronectin was prominently localized by immunohistochemistry to fibrous strands running between the malignant cells in DLBCL biopsy samples, in keeping with its role in extracellular matrix formation. By contrast, the protein products of three other stromal-1 genes – *MMP9*, *SPARC*, and *CTGF* – were localized primarily in histiocytic cells that infiltrated the DLBCL biopsies. By immunofluorescence, *SPARC* and *CTGF* colocalized with CD68, which is a marker for cells in the monocytic lineage. As expected for a stromal-1 gene product, *SPARC* protein levels were associated with favorable overall survival (Figure 5A).

[0122] The stromal-1 signature includes genes that are coordinately expressed in many normal mesenchymal tissues, most of which encode proteins that form or modify the extracellular matrix. The localization of fibronectin to fibrous strands insinuated between the malignant lymphoma cells suggests that the stromal-1 signature reflects the fibrotic nature of many DLBCL tumors. This fibrotic reaction may be related to another stromal-1 signature component, *CTGF*, which participates in many fibrotic responses and diseases, and promotes tumor growth and metastasis of epithelial cancers (Shi-Wen et al., *Cytokine Growth Factor Rev.*, 19: 133-44 (2008)).

[0123] The foregoing results also indicate that the stromal-1 signature reflects a monocyte-rich host reaction to the lymphoma that is associated with the abundant deposition of extracellular matrix. Tumors with high expression of the stromal-1 signature were infiltrated by cells of the myeloid lineage, which include cells that have been implicated in the pathogenesis of epithelial cancers, including tumor-associated macrophages, myeloid-derived suppressor cells, and Tie2-expressing monocytes (reviewed in Wels et al., *Genes Dev.*, 22: 559-74 (2008)). In animal models, these myeloid lineage cells promote tumor cell invasion by secreting matrix metalloproteinases such as *MMP9*, suppress T cell immune responses, and initiate angiogenesis.

[0124] Several stromal-2 signature genes encode well-known markers of endothelial cells. These include von-Willebrand factor (*VWF*) and CD31 (*PECAMI*), as well as other genes specifically expressed in endothelium such as *EGFL7*, *MMRN2*, *GPR116*, and

SPARCL (Table 1). This signature also includes genes encoding key regulators of angiogenesis, such as, for example, KDR (VEGF receptor-2); Grb10, which mediates KDR signaling; integrin alpha 9, which enhances VEGF signaling; TEK, the receptor tyrosine kinase for the cytokine angiopoietin; ROBO4, an endothelial-specific molecular guidance molecule that regulates angiogenesis; and ERG, a transcription factor required for endothelial tube formation. The stromal-2 signature genes *CAVI*, *CAV2*, and *EHD2* encode components of caveolae, which are specialized plasma membrane structures that are abundant in endothelial cells and required for angiogenesis (Frank et al., *Arterioscler. Thromb. Vasc. Biol.*, 23: 1161-8 (2003); Woodman et al., *Am. J. Pathol.*, 162: 2059-68 (2003)). Although the stromal-2 signature includes a large number of genes expressed in endothelial cells, other genes are expressed exclusively in adipocytes, including *ADIPOQ*, *FABP4*, *RBP4*, and *PLIN*.

[0125] Quantitative tests were done to determine whether expression of the stromal-2 signature relative to the stromal-1 signature (i.e., high stromal score) is related to high tumor blood vessel density, given the connection between many stromal-2 signature genes and angiogenesis. More specifically, the stromal-1 signature averages were subtracted from the stromal-2 signature average to thereby obtain a stromal score for each biopsy. Tests showed a quantitative measure of blood vessel density correlated significantly with the stromal score ($r=0.483$, $p=0.019$) (see Figures 5B and 5C), such that higher blood vessel densities correlated with higher stromal scores.

[0126] Thus, the stromal-1 and stromal-2 gene expression signatures reflect the character of the non-malignant cells in DLBCL tumors, and the stromal-2 signature may represent an “angiogenic switch” in which the progression of a hyperplastic lesion to a fully malignant tumor is accompanied by new blood vessel formation (Hanahan et al., *Cell*, 86: 353-64 (1996)). DLBCL tumors with high relative expression of the stromal-2 signature were associated with increased tumor blood vessel density and adverse survival. Significant macrophage infiltration in some DLBCL tumors may predispose to angiogenesis since, in experimental models, tumor-associated macrophages accumulate prior to the angiogenic switch and are required for the switch to occur (Lin et al., *Cancer Res.*, 66: 11238-46 (2006)). Additionally, CXCL12 (SDF-1), a stromal-2 signature component, is a chemokine secreted either by fibroblasts or endothelial cells that can promote angiogenesis by recruiting CXCR4+ endothelial precursor cells from the bone marrow (Orimo et al., *Cell*, 121: 335-48 (2005)). Moreover, an antagonist of angiogenesis, thrombospondin-2 (Kazerounian et al., *Cell Mol. Life Sci.*, 65: 700-12 (2008)), is a stromal-1 signature component, which may

WO 2009/149359

PCT/US2009/046421

62

explain why tumors with low relative expression of this signature had an elevated blood vessel density. Furthermore, the expression of adipocyte-associated genes in DLBCL tumors with high stromal-2 signature expression may play a role in angiogenesis since some cells in adipose tissue may have the potential to differentiate into endothelial cells (Planat-Benard et al., *Circulation*, 109: 656-63 (2004)). Alternatively, the expression of adipose-associated genes may reflect the recruitment of bone marrow-derived mesenchymal stem cells, which home efficiently to tumors (Karnoub et al., *Nature*, 449: 557-63 (2007)) and can stabilize newly formed blood vessels (Au et al., *Blood*, 111: 4551-4558 (2008)).

[0127] The foregoing results indicate that the stromal-1 and stromal-2 gene signatures can be used to generate a stromal score that correlates with increased blood vessel density. Thus, the stromal score can be used to determine if a DLBCL patient is likely to benefit from administration of antiangiogenic therapy (alone, or in conjunction with another DLBCL therapeutic regimen).

[0128] [BLANK]

[0129] The use of the terms “a” and “an” and “the” and similar referents in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms “comprising,” “having,” “including,” and “containing” are to be construed as open-ended terms (i.e., meaning “including, but not limited to,”) unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

[0130] Preferred embodiments of this invention are described herein, including the best mode known to the inventors for carrying out the invention. Variations of those preferred embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the invention to be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the invention unless otherwise indicated herein or otherwise clearly contradicted by context.

WE CLAIM:

1. A method of predicting the survival outcome of a subject suffering from diffuse large B cell lymphoma (DLBCL) and having been treated with chemotherapy and Rituximab (R-CHOP), which method comprises:

a) isolating gene expression product from one or more DLBCL biopsy samples from the subject;

b) obtaining a gene expression profile from the gene expression product, wherein the expression profile comprises an expression level for each gene in a germinal center B cell (GCB) gene expression signature and a stromal-1 gene expression signature;

c) determining a GCB signature value and a stromal-1 signature value from the gene expression profile, wherein the GCB signature value corresponds to the average of the expression levels of the genes in the GCB gene expression signature and the stromal-1 signature value corresponds to the average of the expression levels of the genes in the stromal-1 gene expression signature; and

d) calculating a survival predictor score using an equation that includes subtracting $[(x) * (\text{the GCB signature value})]$ from an offset term and subtracting $[(y) * (\text{the stromal-1 signature value})]$ from the offset term, wherein (x) and (y) are scale factors and given the same offset term and (x) and (y) scale factors a lower survival predictor score compared to another survival predictor score of another subject indicates a more favorable survival outcome and a higher survival predictor score compared to another survival predictor score of another subject indicates a less favorable survival outcome for the subject.

2. The method of claim 1, wherein the method comprises:

b) obtaining a gene expression profile from the gene expression product, wherein the expression profile comprises an expression level for each gene in a GCB expression signature, a stromal-1 gene expression signature, and a stromal-2 gene expression signature;

c) determining a GCB signature value, a stromal-1 signature value, and a stromal-2 gene signature value from the gene expression profile, wherein the stromal-2 signature value corresponds to the average of the expression levels of the genes in the stromal-2 gene expression signature; and

d) calculating a survival predictor score using an equation that includes subtracting $[(x) * (\text{the GCB signature value})]$ from the offset term and subtracting $[(y) * (\text{the stromal-1 signature value})]$ from the offset term, and adding $[(z) * (\text{the stromal-2 signature value})]$ to the offset term, wherein (x), (y), and (z) are scale factors and given the same offset term and (x), (y), and (z) scale factors a lower survival predictor score compared to another survival predictor score of another subject indicates a more favorable survival outcome and a higher survival predictor score compared to another survival predictor score of another subject indicates a less favorable survival outcome for the subject.

3. The method of claim 2, wherein the method comprises:

d) calculating a survival predictor score using the equation: survival predictor score = $8.11 - [0.419 * (\text{the GCB signature value})] - [1.015 * (\text{the stromal-1 signature value})] + [0.675 * (\text{the stromal-2 signature value})]$.

4. A method of evaluating a DLBCL sample for a therapeutic regimen based on survival outcome, wherein the subject has been treated with chemotherapy and Rituximab (R-CHOP), wherein the method comprises:

a) isolating gene expression product from one or more DLBCL biopsy samples from the subject;

b) obtaining a gene expression profile from the gene expression product, wherein the expression profile comprises an expression level for each gene in a germinal center B cell (GCB) gene expression signature and a stromal-1 gene expression signature;

c) determining a GCB signature value and a stromal-1 signature value from the gene expression profile, wherein the GCB signature value corresponds to the average of the expression

levels of the genes in the GCB gene expression signature and the stromal-1 signature value corresponds to the average of the expression levels of the genes in the stromal-1 gene expression signature; and

d) calculating a survival predictor score using an equation that includes subtracting $[(x) * (\text{the GCB signature value})]$ from an offset term and subtracting $[(y) * (\text{the stromal-1 signature value})]$ from the offset term, wherein (x) and (y) are scale factors and given the same offset term and (x) and (y) scale factors (i) a lower survival predictor score compared to another survival predictor score of another subject indicates a more favorable survival outcome and a therapeutic regimen that includes chemotherapy and Rituximab (R-CHOP) and (ii) a higher survival predictor score compared to another survival predictor score of another subject indicates a less favorable survival outcome and a therapeutic regimen other than R-CHOP.

5. The method of claim 4, wherein the DCLBCL sample is from a subject that suffered relapse from R-CHOP treatment.

6. The method of claim 4, wherein the method comprises:

b) obtaining a gene expression profile from the gene expression product, wherein the expression profile comprises an expression level for each gene in a GCB expression signature, a stromal-1 gene expression signature, and a stromal-2 gene expression signature;

c) determining a GCB signature value, a stromal-1 signature value, and a stromal-2 gene signature value from the gene expression profile, wherein the stromal-2 signature value corresponds to the average of the expression levels of the genes in the stromal-2 gene expression signature; and

d) calculating a survival predictor score using an equation that includes subtracting $[(x) * (\text{the GCB signature value})]$ from the offset term and subtracting $[(y) * (\text{the stromal-1 signature value})]$ from the offset term, and adding $[(z) * (\text{the stromal-2 signature value})]$ to the offset term, wherein (x), (y), and (z) are scale factors and given the same offset term and (x), (y), and (z) scale factors (i) a lower survival predictor score compared to another survival predictor score of another subject indicates a more favorable survival outcome and a therapeutic regimen that includes

chemotherapy and Rituximab (R-CHOP) and (ii) a higher survival predictor score compared to another survival predictor score of another subject indicates a less favorable survival outcome and a therapeutic regimen other than R-CHOP.

7. The method of claim 6, wherein the method comprises:

d) calculating a survival predictor score using the equation: survival predictor score = 8.11 - [0.419*(the GCB signature value)] - [1.015*(the stromal-1 signature value)] + [0.675*(the stromal-2 signature value)].

8. A method of predicting the survival outcome of a subject suffering from diffuse large B cell lymphoma (DLBCL) and treated with chemotherapy and Rituximab (R-CHOP), which method comprises:

a) isolating gene expression product from one or more DLBCL biopsy samples from the subject;

b) obtaining a gene expression profile from the gene expression product, wherein the expression profile comprises an expression level for each gene in a germinal center B-cell gene (GCB) expression signature, a stromal-1 gene expression signature, and a stromal-2 gene expression signature;

c) determining a GCB signature value, a stromal-1 signature value, and a stromal-2 signature value from the gene expression profile, wherein the GCB signature value corresponds to the average of the expression levels of the genes in the GCB gene expression signature, the stromal-1 signature value corresponds to the average of the expression levels of the genes in the stromal-1 gene expression signature, and the stromal-2 signature value corresponds to the average of the expression levels of the genes in the stromal-2 gene expression signature; and

d) calculating a survival predictor score using the equation: survival predictor score = A - [(x)*(the GCB signature value)] - [(y)*(the stromal-1 signature value)] + [(z)*(the stromal-2 signature value)], wherein A is an offset term and (x), (y), and (z) are scale factors; and

e) calculating a probability of survival outcome for the subject beyond an amount of time t following treatment for DLBCL, wherein the subject's probability of survival outcome $P(SO)$ is calculated using the equation: $P(SO) = SO_0(t)^{\exp((s) \cdot \text{survival predictor score})}$, wherein $SO_0(t)$ is the probability of the survival outcome, which corresponds to the largest time value smaller than t in a survival outcome curve, and wherein (s) is a scale factor.

9. The method of claim 8, wherein the survival outcome is overall survival and the method comprises:

d) calculating a survival predictor score using the equation: survival predictor score = $[0.419 \cdot (\text{the GCB signature value})] - [1.015 \cdot (\text{the stromal-1 signature value})] + [0.675 \cdot (\text{the stromal-2 signature value})]$; and

e) calculating a probability of overall survival after time t for the subject, wherein the subject's probability of overall survival $P(OS)$ is calculated using the equation: $P(OS) = OS_0(t)^{\exp(\text{survival predictor score})}$, wherein $OS_0(t)$ is the probability of overall survival, which corresponds to the largest time value smaller than t in an overall survival curve, and wherein scale factor $(s) = 1$.

10. The method of claim 8, wherein the survival outcome is progression free survival and the method comprises:



d) calculating a survival predictor score using the equation: survival predictor score = $8.11 - [0.419 \cdot (\text{the GCB signature value})] - [1.015 \cdot (\text{the stromal-1 signature value})] + [0.675 \cdot (\text{the stromal-2 signature value})]$; and

e) calculating a probability of progression free survival after time t for the subject, wherein the subject's probability of progression free survival $P(PFS)$ is calculated using the equation $P(PFS) = F_0(t)^{\exp(0.976 \cdot \text{survival predictor score})}$, wherein $F_0(t)$ is the probability of progression free survival, which corresponds to the largest time smaller than t in a survival curve, and wherein scale factor $(s) = 0.976$.

11. The method of claim 8, wherein the treatment for DLCBL includes chemotherapy and the administration Rituximab.
12. The method of claim 8, wherein the method further includes providing the subject with the calculated probability of the survival outcome after time t.
13. A method of generating a survival estimate curve for a plurality of subjects suffering from diffuse large B cell lymphoma (DLBCL) and having been treated with chemotherapy and Rituximab (R-CHOP), which method comprises:
 - a) obtaining one or more DLBCL biopsy samples from each of the plurality of subjects;
 - b) isolating gene expression product from the one or more DLBCL biopsy samples from each of the plurality of subjects;
 - c) obtaining a gene expression profile from the gene expression product of each subject, wherein the gene expression profile comprises an expression level for each gene in a germinal center B cell gene (GCB) expression signature, a stromal-1 gene expression signature, and a stromal-2 gene expression signature;
 - d) determining for each subject the GCB signature value, the stromal-1 signature value, and the stromal-2 signature value from the subject's gene expression profile, wherein the GCB signature value corresponds to the average of the expression levels of the genes in the GCB gene expression signature, the stromal-1 signature value corresponds to the average of the expression levels of the genes in the stromal-1 gene expression signature, and the stromal-2 signature value corresponds to the average of the expression levels of the genes in the stromal-2 gene expression signature;
 - e) calculating a survival predictor score for each subject;
 - f) tracking a survival outcome for each subject following treatment for DLBCL; and

g) generating a survival estimate curve that correlates the probability of the tracked survival outcome with time following treatment for DLBCL, wherein the survival curve also correlates tracked outcome over time with the survival predictor score for the subjects.

14. The method of claim 13, wherein the survival outcome is overall survival.
15. The method of claim 13, wherein the survival outcome is progression free survival.

**Figure 1A****Figure 1B**

2/22




Figure 1C

3/22




Figure 2A

4/22




Figure 2B





Figure 2C

6/22









Figure 4A



**Figure 4B**

**Figure 5A**

Low Blood-Vessel density
(CD34 + cells per μm^2)



High Blood-Vessel density
(CD34 + cells per μm^2)

**Figure 5B****Figure 5C**

10/22




Figure 6A

11/22





Figure 6B



13/22




Figure 8A

14/22




Figure 8B

15/22




Figure 8C




Figure 9A





Figure 9B

18/22

**Figure 9C**

19/22




Figure 10A

20/22




Figure 10B

21/22




Figure 10C

22/22




Figure 11